

ANALYSIS & PDE

Volume 17

No. 10

2024

PETER HINTZ

SEMICLASSICAL PROPAGATION THROUGH CONE POINTS

SEMICLASSICAL PROPAGATION THROUGH CONE POINTS

PETER HINTZ

We introduce a general framework for the study of the diffraction of waves by cone points at high frequencies. We prove that semiclassical regularity propagates through cone points with an almost sharp loss even when the underlying operator has leading-order terms at the conic singularity which fail to be symmetric. We moreover show improved regularity along strictly diffractive geodesics. Applications include high-energy resolvent estimates for complex- or matrix-valued inverse square potentials and for the Dirac–Coulomb equation. We also prove a sharp propagation estimate for the semiclassical conic Laplacian.

The proofs use the semiclassical cone calculus, introduced recently by the author, and combine radial point estimates with estimates for a scattering problem on an exact cone. A second microlocal refinement of the calculus captures semiclassical conormal regularity at the cone point and thus facilitates a unified treatment of semiclassical cone and b-regularity.

1. Introduction

We present a systematic analysis of the propagation of semiclassical regularity through points which are geometrically singular (cone points), analytically singular (e.g., including inverse square potentials), or both. The novel aspect of our approach is that it handles leading-order singular terms with ease, *regardless of symmetry or sign conditions*.

As a simple application of our main microlocal propagation result, we consider high-energy scattering by complex-valued potentials on \mathbb{R}^n with an inverse square singularity. Denote by $H_0^2(\mathbb{R}^n \setminus \{0\})$ the closure of $C_c^\infty(\mathbb{R}^n \setminus \{0\})$ in the topology of $H^2(\mathbb{R}^n)$; denote further by $\Delta = \sum_{j=1}^n D_{x_j}^2$ (where $D = (1/i)\partial$) the nonnegative Laplacian, and denote polar coordinates on \mathbb{R}^n by $(r, \omega) \in (0, \infty) \times \mathbb{S}^{n-1}$.

Theorem 1.1 (high-energy estimates for potential scattering). *Let $V(x) = V_0(x)/|x|^2$, where $V_0 = V_0(r, \omega) \in C_c^\infty([0, \infty)_r \times \mathbb{S}^{n-1}; \mathbb{C})$ and $V_0(0, \omega) \equiv Z \in \mathbb{C}$. (Thus $V(x) = Z/|x|^2 + \mathcal{O}(|x|^{-1})$ near the origin.) Suppose that $n \geq 5$ and $\operatorname{Re} \sqrt{((n-2)/2)^2 + Z} > 1$. Then there exists $\lambda_0 > 0$ so that for all $\lambda \in \mathbb{C}$ with $\operatorname{Re} \lambda > \lambda_0$ and $0 < \operatorname{Im} \lambda < 1$, the operator*

$$\Delta + V - \lambda : H_0^2(\mathbb{R}^n \setminus \{0\}) \rightarrow L^2(\mathbb{R}^n) \tag{1-1}$$

is invertible, and its inverse obeys the operator norm bound

$$\|\chi(\Delta + V - \lambda)^{-1}\chi\|_{L^2 \rightarrow L^2} \leq C_{\chi, \epsilon} |\lambda|^{-\frac{1}{2} + \epsilon} \tag{1-2}$$

MSC2020: primary 58J47; secondary 35L81, 35P25, 35S05.

Keywords: propagation of singularities, conic singularities, inverse square potentials, semiclassical cone calculus.

for all $\chi \in C_c^\infty(X)$ and $\epsilon > 0$. More generally, $\Delta + V - \lambda : \mathcal{D} \rightarrow L^2(\mathbb{R}^n)$ is invertible for $n \geq 2$ and $Z \in \mathbb{C} \setminus (-\infty, -((n-2)/2)^2]$ for a suitable domain \mathcal{D} (see (5-29) with $l = 1$), and the estimate (1-2) holds in this generality as well.

The point is that we can allow for Z to be nonreal, in which case $\Delta + V$ is not a symmetric operator on $C_c^\infty(\mathbb{R}^n \setminus \{0\})$. (The compact support assumption on V is made only to focus the attention of the reader on a neighborhood of the singularity at $x = 0$. The result holds also for V with sufficient decay at infinity, such as $|\partial_x^\alpha V| \lesssim \langle x \rangle^{-1-\delta}$ for all α where $\delta > 0$.) For a general result for matrix-valued inverse square potentials without symmetry conditions, see Theorem 5.7; Lemma 5.10 verifies the assumptions of Theorem 5.7 for the case considered in Theorem 1.1. Typical applications of high-energy resolvent estimates include decay and local smoothing estimates for solutions to wave and Schrödinger equations; since such applications are orthogonal to the focus of the present paper, we shall not discuss them here.

Burq and Planchon, Stalker, and Tahvildar-Zadeh [Planchon et al. 2003; Burq et al. 2003] proved Strichartz estimates for exact inverse square potentials in the case of real $Z > -((n-2)/2)^2$. Duyckaerts [2006] obtained, by means of estimates for semiclassical defect measures, high-energy resolvent estimates (without the ϵ -loss) in the more general setting of inverse square potentials at a finite collection of points p_j in \mathbb{R}^n , at each of which the coefficient Z_j satisfies $Z_j > -((n-2)/2)^2$. We also mention the work [Baskin and Wunsch 2013] on lossless resolvent estimates in a *geometric* setting, namely in the presence of finitely many conic singularities, and the work by [Hillairet and Wunsch 2020] on resonances in this setting (see also [Galkowski 2017]).

Remark 1.2 (more natural settings). The setting of Theorem 1.1 is chosen here for its simplicity. More natural examples in which leading-order terms without signs or symmetry properties are present arise in particular in the study of PDEs on vector bundles. As an example, motivated by the recent work [Baskin and Wunsch 2023], we prove high-energy resolvent estimates for the Dirac–Coulomb equation in Section 5C; see Theorem 5.14.

The heart of the proof of Theorem 1.1 is the propagation of semiclassical regularity through $r = 0$,¹ which we prove in this paper for a general class of *admissible operators*; see Definition 4.1 and Theorem 4.10. Thus, in addition to inverse square singularities (which may be anisotropic), we allow for the underlying metric g to have a conic singularity at $r = 0$, so $g = dr^2 + r^2k(r, y, dr, dy)$ for some smooth r -dependent tensor k , with $k|_{r=0}$ a Riemannian metric on a closed manifold Y . We moreover allow for further first-order differential operators of the schematic form $r^{-1}D_r, r^{-2}D_y$ to be present. All these singular terms are allowed to be of the same strength at $r = 0$: they are, to leading order at $r = 0$, homogeneous of degree -2 with respect to dilations.

In order to explain the main features of Theorem 1.1, note that the degree -2 homogeneity of the Laplacian and of the potential r^{-2} is reflected also in the Hardy inequality, which demonstrates that any factor of r^{-1} should be regarded as a derivative as far as analysis near the cone point $r = 0$ is concerned. Therefore, when Z in Theorem 1.1 is nonreal, the operator $\Delta + V - \lambda$ is, *even to leading order at the cone*

¹In particular, the choice of the large end of the space, here \mathbb{R}^n , is only made for convenience and allows for simple control of the global structure of the geodesic flow. Thus, we do not discuss the large literature on limiting absorption principles here.

point, not symmetric. Therefore, techniques rooted in the spectral theory of self-adjoint operators do not apply. Furthermore, recall that even for solutions of smooth-coefficient PDEs $Pu = f$ where the principal symbol of P is complex-valued, microlocal regularity of u propagates along the null-bicharacteristics of $\operatorname{Re} P$ *only under a sign condition on $\operatorname{Im} P$* near the boundary of the support of $\operatorname{Im} P$ [Vasy 2018, §4.5]; on a technical level, the term $\operatorname{Im} P$ contributes the leading term in a positive commutator argument for proving the propagation of regularity along null-bicharacteristics of $\operatorname{Re} P$. The absence of sign conditions on $\operatorname{Im} V$ in Theorem 1.1 is thus a significant obstacle for the applicability of existing methods.

In general geometric or analytic settings where one cannot separate variables, propagation estimates through cone points and other types of singularities have so far largely been restricted to self-adjoint settings. Melrose and Wunsch [2004] studied the diffraction of waves by conic singularities by combining microlocal propagation estimates in the edge calculus developed by Mazzeo [1991] with the inversion of a suitable model operator on an exact cone. This point of view is closely related to that adopted in the present paper, see Remark 1.4, though by contrast to the present work, [Melrose and Wunsch 2004] takes full advantage of the self-adjointness of the underlying Laplace operator.

Later works on wave propagation in singular geometries have been based on positive commutator arguments relative to a quadratic form domain (thus still in self-adjoint settings), following the blueprint of [Vasy 2008] on the propagation of singularities on smooth manifolds with corners (see [Lebeau 1997] for the analytic setting). Vasy's work was extended to the setting of manifolds with edge singularities by Melrose, Vasy and Wunsch [Melrose et al. 2008], and the same authors established improved regularity of the strictly diffracted front on manifolds with corners [Melrose et al. 2013]. See [Qian 2009] for the case of inverse square potentials. We remark that in these works, the underlying geometry near the singularity is *not* reflected in the type of singularities which propagate or diffract—for instance, in the case of [Vasy 2008], the geometry is that of a manifold with corners equipped with a smooth (incomplete!) Riemannian metric, but the correct notion of regularity is conormality at the boundary; thus, these works introduce mixed differential-pseudodifferential calculi which are compatible with both structures.

Baskin and Marzuola [2022] combined the techniques of [Vasy 2008] with those developed in [Baskin et al. 2015] to study the long-time behavior of waves on manifolds with conic singularities. An important ingredient in their work is a high-energy estimate for propagation through the conic singularity. In the present paper we give an alternative proof which in particular avoids the use of a mixed calculus; see also Remark 1.5. We also mention that Gannot and Wunsch [2023] analyzed the diffraction by conormal potentials in the semiclassical setting using direct commutator methods involving paired Lagrangian distributions, inspired by [de Hoop et al. 2015].

The recent work [Baskin and Wunsch 2023] on diffraction for the Dirac–Coulomb equation is also rooted in [Melrose and Wunsch 2004; Melrose et al. 2008]. While the (first-order) Dirac–Coulomb operator is self-adjoint for the range of Coulomb charges considered in [Baskin and Wunsch 2023], the wave-type operator obtained by taking an appropriate square has nonsymmetric leading-order terms at the central singularity; thus, the authors work directly with the first-order operator in their proofs of propagation results. We are able to give a direct proof of high-energy estimates for the resolvent associated with the wave-type operator arising in [Baskin and Wunsch 2023]; see Section 5C.

In the high-energy regime under study in the present paper, the strategy for overcoming the issues caused by the absence of symmetry or self-adjointness properties is the following. We distill the contribution of V to the high-frequency propagation of regularity (i.e., in Theorem 1.1, the inverse powers of $|\lambda|$ appearing in uniform estimates of L^2 norms) into a model problem right at the cone point, thus decoupling it from the real-principal-type propagation away from the cone point (where V plays no role due to its subprincipal nature). More precisely, in the setting of Theorem 1.1, set $h := |\lambda|^{-1/2}$ and $z = h^2\lambda = 1 + \mathcal{O}(h)$, and define the semiclassical rescaling

$$P_{h,z} = h^2(\Delta + V - \lambda) = h^2\Delta - z + \frac{h^2}{|x|^2}V_0 = (hD_r)^2 - i(n-1)\frac{h}{r}hD_r + h^2r^{-2}\Delta_{\mathbb{S}^{n-1}} - z + \frac{h^2}{r^2}V_0. \quad (1-3)$$

Recall here that $V_0 \in C_c^\infty([0, \infty)_r \times \mathbb{S}^{n-1}; \mathbb{C})$ is equal to a constant $Z \in \mathbb{C}$ at $r = 0$. The operator $P_{h,z}$ is a semiclassical differential operator in $r > 0$. Its uniform analysis as $h \rightarrow 0$, as far as the novel bit near $r = 0$ is concerned, is based on two ingredients, discussed in more detail in Section 1A:

(1) *Symbolic propagation estimates*: real-principal-type propagation in $r > 0$ in the spirit of [Duistermaat and Hörmander 1972], and radial point estimates down to $r = 0$ in the spirit of [Melrose 1994; Vasy 2013] but taking place in the semiclassical cone algebra introduced by the author in [Hintz 2022]. The advantage of this algebra in the present setting is that $P_{h,z}$ has a smooth and nondegenerate principal symbol in this algebra down to $r = 0$; in this algebra, the proofs of the relevant symbolic estimates are then essentially standard.

(2) *Inversion of a model problem*: Passing to the rescaled variable $\hat{r} = r/h$ and letting $h \rightarrow 0$ for fixed \hat{r} in the resulting expression of $P_{h,z}$ gives

$$N(P) = D_{\hat{r}}^2 - i(n-1)\hat{r}^{-1}D_{\hat{r}} + \hat{r}^{-2}\Delta_{\mathbb{S}^{n-1}} - 1 + \frac{Z}{\hat{r}^2}. \quad (1-4)$$

The inversion of $N(P)$ is a scattering problem on an exact cone at unit frequency and requires the existence of the limiting (outgoing) resolvent. Its analysis is based on b-analysis near the small end of the cone [Melrose 1993] and on the microlocal approach to scattering theory on spaces with conic infinite ends pioneered in [Melrose 1994].

The ϵ -loss in the estimate (1-2) is then due to the analogous loss in the limiting absorption principle for the scattering problem, as one needs to exclude incoming but allow outgoing spherical waves; see Remark 5.3. (We shall in fact deduce the lossy estimates stated in Theorem 1.1 from sharp results — as far as the relationship of domain and codomain of $P_{h,z}$ is concerned — on spaces with variable semiclassical orders.) For general admissible operators, the decay rates of incoming and outgoing solutions of the model problem are typically different, and the semiclassical loss upon propagation through the cone point is equal to their difference (up to an additional ϵ -loss); we give explicit examples in which this loss indeed occurs in Appendix A, demonstrating that our analysis is sharp up to an ϵ -loss. It seems impossible to avoid this ϵ -loss if one proves the propagation estimates in the above step-by-step manner: the microlocal radial point estimates force inequalities on the semiclassical orders (see, however, [Wang 2020] in a Besov space setting), and also on the incoming and outgoing decay orders of the function

spaces on which the model problem is invertible. Thus, avoiding this ϵ -loss requires the development of propagation arguments which provide control near $r = 0$ in one fell swoop. We demonstrate this for operators $P_{h,z} = h^2 \Delta_g - z$, $z = 1 + \mathcal{O}(h)$, on (nonexact) conic manifolds, for which we are able to obtain a *lossless* propagation estimate by means of a positive commutator argument which is global on the level of the normal operator $N(P)$, i.e., which involves the construction of a commutator which is positive *as an operator* on an exact cone, in the spirit of Mourre's construction [1980/81] and Vasy's approach [2000; 2001] to many-body scattering; see Theorem 5.4, in particular the estimate (5-6). It is not clear at present however how to generalize such an argument to more general (in particular nonsymmetric) settings.

The close connection between diffraction by conic singularities and scattering on large ends of cones was recently studied for exact (or "product") cones (i.e., the metric is $g = dr^2 + r^2 k(y, dy)$) by Yang [2022], resulting in a partial improvement of the classical analysis by Cheeger and Taylor [1982a; 1982b] which was based on separation of variables and Bessel function analysis. Recently, Chen Xi [2022] constructed a detailed parametrix for high-frequency diffraction by (nonexact) conic singularities, i.e., for the operator $(h^2 \Delta_g - (1 \pm i0))^{-1}$, with applications to short-time Strichartz estimates for the Schrödinger equation; an important ingredient in his work is the precise resolvent construction by Guillarmou, Hassell and Sikora [Guillarmou et al. 2013], applied on an exact cone which arises similarly to (1-4). (The history of the study of propagation and diffraction phenomena for solutions of wave-type equations on manifolds with singularities is long, starting with Sommerfeld's example [1896] and early developments by Friedlander [1958] and Keller [1985]. The use of geometric and microlocal techniques for the analysis of singularities goes back to [Melrose and Sjöstrand 1978; 1982] on manifolds with boundary using commutator techniques, and [Melrose 1975; Taylor 1976; Melrose and Taylor 2018] using parametrix constructions.) We also mention the recent work by Keeler and Marzuola [2022] who use estimates for the resolvent on exact cones perturbed by a radial (but not homogeneous) potential in order to obtain dispersive estimates for the Schrödinger equation.

Finally, we prove a *diffractive improvement* which gives finer control on the strength of singularities as they propagate through the cone point. Combining our framework with the arguments in [Melrose and Wunsch 2004; Melrose et al. 2008] for the propagation of coisotropic regularity, we show that, under a nonfocusing condition, the strongest singularities propagating towards the cone point only continue along geometric geodesics (limits of geodesics barely missing the cone point), whereas away from those, the diffracted front is smoother; see Section 4F. We do not address here the interesting question of whether one can prove estimates in the presence of multiple scatterers using such diffractive improvements, as done in [Baskin and Wunsch 2013]; in particular, we do not recover Duyckaerts' results [2006] here. See Remark 5.11.

Regarding applications of our high-frequency estimates, we content ourselves with a few conjectural remarks. First, in the context of [Baskin et al. 2015; 2018; Baskin and Marzuola 2022] and Remark 5.12, it should be possible to use our results to justify contour-shifting arguments for obtaining asymptotic expansions (including radiation fields) of solutions to wave-type equations on static conic manifolds with rather general inverse square potentials in the forward cone. Due to the relationship between edge Sobolev spaces and semiclassical cone Sobolev spaces discussed in Remark 3.4, it is likely not necessary to prove b -regularity at the spatial cone point $r = 0$ (unlike in [Baskin and Marzuola 2022, §8.2.2–8.2.3]); instead,




Figure 1. The semiclassical cone single space X_{ch} .

edge propagation results at $r = 0$ (with uniformity as $t \rightarrow \infty$ encoded by b-spaces), which directly apply also to nonsymmetric operators, likely suffice. One does need to show, however, the existence of global forward solutions by means distinct from energy methods since spectral methods are no longer available for defining domains of the spatial operator.² Second, it is not immediately clear how to generalize local smoothing estimates for Schrödinger equations (e.g., as in [Duyckaerts 2006, Corollaire (2)]) if the underlying Hamiltonian is not self-adjoint; it is an interesting question however whether, say in the context of Theorem 1.1, some version of local smoothing (with an ϵ -loss) holds for the evolution defined in terms of the inverse Fourier transform of the resolvent $(\Delta + V - \lambda + i0)^{-1}$, $\lambda \in \mathbb{R}$, assuming it exists. (Estimates in the elliptic regime $\lambda \ll -1$ are discussed in [Hintz 2022].) A similar remark applies to Strichartz estimates.

1A. Sketch of the proof. Consider again the operator $P_{h,z}$ from (1-3); we work locally near $r = 0$, thus on $X = [0, 1)_r \times \mathbb{S}^{n-1}$. In order to achieve a clean separation of the regimes $h \rightarrow 0$, $r > 0$ (corresponding to semiclassical analysis away from the cone point) and $h \sim r \rightarrow 0$ (where the normal operator $N(P)$ in (1-4) enters and semiclassical tools cease to be applicable), we work on a resolution of the total space $[0, 1)_h \times X$ obtained by a real blow-up of $h = r = 0$,³

$$X_{ch} := [[0, 1)_h \times X; \{0\} \times \partial X].$$

See Figure 1. We wish to regard $h/(h+r)$ as the “true” semiclassical parameter; we proceed to make this more precise.

Note first that for $h = 1$, the rescaling $r^2 P_{1,z}$ is a Fuchs-type operator, or b-differential operator in the terminology of [Melrose 1993], namely a differential operator built out of the vector fields $r\partial_r$ and ∂_y (which span the space of *b-vector fields*), where $y \in \mathbb{R}^{n-1}$ denotes local coordinates on ∂X . In this sense, the rescaled operator $r^2 P_{1,z}$ has elliptic principal part given by $(rD_r)^2 + k^{ij} D_{y^i} D_{y^j}$, where k^{ij} is the inverse metric on \mathbb{S}^{n-1} . As h tends to 0, the operator $r^2 P_{h,z}$ is built out of the semiclassical vector fields $hr\partial_r$ and $h\partial_y$ (which span the space of *semiclassical b-vector fields*). In this semiclassical sense (i.e.,

²A closely related setting in which many of these points are addressed, with the exception of the analysis at a spatial inverse square singularity, is described in [Hintz 2023]; see in particular §5.3 of that work.

³Recall here that the real blow-up gives an invariant way of introducing polar coordinates around $\{0\} \times \partial X$. Thus, a neighborhood of $h = r = 0$ in X_{ch} is diffeomorphic to $[0, 1)_\rho \times [0, \frac{\pi}{2}]_\theta \times \partial X$ and equipped with a smooth map (the blow-down map) to $[0, 1) \times X = [0, 1) \times ([0, 1) \times \mathbb{S}^{n-1})$ given by $(\rho, \theta, y) \mapsto (\rho \sin \theta, (\rho \cos \theta, y))$, which is a diffeomorphism away from the front face $\rho^{-1}(0)$. In practice, it is more convenient to work with the smooth functions $h+r$, $h/(h+r)$ on X_{ch} instead of ρ, θ .

ignoring terms with extra powers of h), its principal part is

$$r^2 P_{h,z} \sim (hr D_r)^2 + k^{ij} h D_{y^i} h D_{y^j} - r^2 z.$$

The characteristic set, i.e., the zero set of its principal symbol $\xi_{b\hbar}^2 + |\eta_{b\hbar}|^2 - r^2$, becomes singular at $r = 0$, which is indicative of the inadequacy of the semiclassical b-setting to capture the behavior of $P_{h,z}$ microlocally near $h = r = 0$ (cf. the above discussion regarding the tension between the geometry and the notion of regularity in [Melrose and Wunsch 2004; Vasy 2008] and subsequent works). The way out is to divide by $(h + r)^2$ and thus consider

$$\left(\frac{r}{h+r}\right)^2 P_{h,z} =: p_{h,z}\left(r, y, \frac{h}{h+r} r D_r, \frac{h}{h+r} D_y\right)$$

as a differential operator built out of $(h/(h+r)) r \partial_r$ and $h/((h+r)) \partial_{y^i}$, which are the prototypical *semiclassical cone vector fields* introduced in [Hintz 2022]; see Section 3A. In this sense, the principal part of $(r/(h+r))^2 P_{h,z}$ (i.e., ignoring terms of size $\mathcal{O}(h/(h+r))$) is

$$\left(\frac{r}{h+r}\right)^2 P_{h,z} \sim \left(\frac{h}{h+r} r D_r\right)^2 + k^{ij} \left(\frac{h}{h+r} D_{y^i}\right) \left(\frac{h}{h+r} D_{y^j}\right) - \left(\frac{r}{h+r}\right)^2 z. \tag{1-5}$$

Put differently, we may note that $P_{h,z} \sim (h D_r)^2 + k^{ij} (hr^{-1} D_{y^i})(hr^{-1} D_{y^j}) - z$ is homogeneous of degree 0 with respect to scaling in (h, r) , and approximately homogeneous of degree -2 with respect to scaling in r ; this suggests expressing $P_{h,z}$ in terms of $r/(h+r) = 1 - h/(h+r)$, leading again to (1-5).

Note that in the regime $h/(h+r) \ll 1$, where we are aiming to use semiclassical methods, the operator (1-5) is now nondegenerate in the sense that its principal symbol

$$p_{0,1}(r, y, \xi, \eta) = \xi^2 + |\eta|^2 - 1$$

(recall $z = 1 + \mathcal{O}(h)$) has a smooth zero set on which $p_{0,1}$ vanishes simply. (The microlocal analysis of *semiclassical cone operators* in the semiclassical regime is thus concerned with tracking amplitudes of oscillations

$$r^{\frac{i}{h/(h+r)}} \xi e^{\frac{i}{h/(h+r)} \eta \cdot y}$$

through the phase space over $X_{c\hbar}$ — more precisely, over the “semiclassical face” $h/(h+r) = 0$ — whose fiber variables are (ξ, η) .)

The semiclassical cone calculus $\Psi_{c\hbar}(X)$, introduced in [Hintz 2022] and developed further in Section 3, makes this rigorous. It allows for the symbolic analysis of pseudodifferential operators of the form

$$\text{Op}_{c,h}(p) = “p\left(\frac{h}{h+r}, h+r, y, \frac{h}{h+r} r D_r, \frac{h}{h+r} D_y\right)”$$

using standard methods from microlocal analysis: there is a semiclassical principal symbol $p(0, r, y, \xi, \eta)$, which is a symbol on the aforementioned phase space (defined rigorously after Lemma 3.2). Moreover, as usual, the commutator $i[\text{Op}_{c,h}(p), \text{Op}_{c,h}(q)]$ is given by the quantization of the Poisson bracket of p and q up to operators with an extra factor of $h/(h+r)$. For the operator $P_{h,z}$ in (1-3), the Hamilton vector field of its principal symbol is nondegenerate except at two submanifolds of critical points over $r = 0$;

these critical sets are saddle points for the Hamilton flow, and are end or starting points of geodesics hitting the cone point or emanating from it. (See Figure 10.) One can thus prove quantitative microlocal propagation and radial point estimates on the associated scale of *weighted semiclassical cone Sobolev spaces*, which measure L^2 norms of derivatives along $(h/(h+r))rD_r$, $(h/(h+r))D_y$, and which feature weights which are real powers of $r/(h+r)$ and $h+r$.

Remark 1.3 (semiclassical cone ps.d.o.s as tools). The (large) pseudodifferential calculus $\Psi_{ch}(X)$ was introduced in [Hintz 2022] as the space in which inverses and complex powers of elliptic semiclassical cone operators, such as $h^2\Delta + 1$, live; the goal there was a precise description of their Schwartz kernels. Here, by contrast, we use semiclassical cone pseudodifferential operators (ps.d.o.s) as *tools* to understand propagation phenomena. Correspondingly, we only need to consider the *small* semiclassical cone calculus, as our analysis will be based on proving estimates, rather than on the construction and usage of parametrices. (Parametrices are typically significantly more challenging to construct [Chin 2022] and are very precise tools; on the flipside, they tend to be less convenient when the need for generalizations or for proofs of sharp mapping properties on various function spaces arises.) Thus, in Section 3, we provide a perspective on $\Psi_{ch}(X)$ which makes it easy to work with in nonelliptic settings.

At this point, we control the semiclassical regularity of solutions of $P_{h,z}u = f$ at $h/(h+r) = 0$. This means that we have an estimate of the schematic form

$$\|u\| \lesssim \|P_{h,z}u\| + \|Eu\| + \left\| \frac{h}{h+r}u \right\|, \quad (1-6)$$

where Eu controls u on a transversal to the collection of forward geodesics which encounter $r = 0$; the function spaces here are semiclassical cone Sobolev spaces. That is, control of Eu together with weak control of u at $h/(h+r) = 0$ (finiteness of the final term) gives stronger control of u (finiteness of the left-hand side), provided the forcing term $P_{h,z}u$ has suitable bounds (e.g., equals 0). Notice that the weights 1 and $h/(h+r)$ in the norms $\|u\|$ and $\|hu/(h+r)\|$ are comparable for $h \sim r$, i.e., at the front face in Figure 1; thus, the estimate (1-6) does not provide control of u in this regime.

In order to control u globally, including at $h = r = 0$, one needs to invert the *normal operator* $N(P)$ of $P_{h,z}$, which is the restriction of $P_{h,z}$ to the front face of X_{ch} ; see (1-3)–(1-4) for a concrete example. The function spaces on which one inverts $N(P)$ need to match the function spaces in which the symbolic propagation estimates are obtained. As already observed in [Hintz 2022] (see also the earlier paper [Loya 2002]) and demonstrated in detail on the level of function spaces in Section 3C, the correct function spaces for $N(P)$ are standard Sobolev spaces when $\hat{r} = r/h \gtrsim 1$ (i.e., measuring regularity with respect to $D_{\hat{r}}$ and $\hat{r}^{-1}D_{y^i}$) and b-Sobolev spaces in $\hat{r} \lesssim 1$ (i.e., measuring regularity with respect to $\hat{r}D_{\hat{r}}$ and D_{y^i}). Following [Melrose 1994], we show in Section 4D that the analysis of $N(P)$ on spaces with variable orders of decay as $\hat{r} \rightarrow \infty$ *precisely* matches the above symbolic analysis which involves variable semiclassical orders (powers of $h/(h+r)$) to accommodate the threshold requirements for propagation into/out of the radial sets; see [Dyatlov and Zworski 2019, Appendix E.4].

We stress the *global* (rather than microlocal or symbolic) nature of the requirement that the normal operator $N(P)$ be invertible; while verifying this in concrete situations is nontrivial, one has many standard

techniques at one's disposal (such as boundary-pairing arguments, unique continuation, separation of variables, etc.). We also remark that the necessity to invert model (or "normal") operators for the purpose of controlling solutions of PDE in a singular regime is a typical feature of singular PDE; see for example the role of the invertibility of the Mellin-transformed normal operator family in the asymptotic behavior of waves in [Baskin et al. 2015; 2018; Baskin and Marzuola 2019; 2022].

In combination, the symbolic estimates and the normal operator invertibility provide control of u at both hypersurfaces $h/(h+r) = 0$ and $h+r = 0$ of X_{ch} : schematically, one estimates the final term in (1-6) by $\|(h/(h+r))N(P)u\|$ and then replaces $N(P)$ by $P_{h,z}$, thereby committing an error term which vanishes to leading order at the front face $h+r = 0$; one obtains

$$\|u\| \lesssim \|P_{h,z}u\| + \|Eu\| + \left\| (h+r) \frac{h}{h+r} u \right\| \lesssim \|P_{h,z}u\| + \|Eu\| + h\|u\|.$$

The final term can be absorbed into the left-hand side when h is sufficiently small. Thus, we have uniform control of u as $h \rightarrow 0$. (One can package this into an invertibility statement for a modification of $P_{h,z}$ by placing complex-absorbing potentials away from $r = 0$ in the spirit of [Nonnenmacher and Zworski 2009; Wunsch and Zworski 2011; Datchev and Vasy 2012; Vasy 2013]; see Section 4E.)

Remark 1.4 (relation to edge propagation). The proof of symbolic propagation estimates for wave equations on conic or edge manifolds using the edge calculus [Mazzeo 1991], as done in [Melrose and Wunsch 2004, §8] and [Melrose et al. 2008, §11], is closely related, via the Fourier transform in time, to the semiclassical cone Sobolev spaces associated with $\Psi_{ch}(X)$; see Remark 3.4. Going one step further in the comparison, we note that the fine analysis of diffraction of [Melrose and Wunsch 2004] for waves on a conic manifolds uses a normal operator at the cone point which is defined via a rescaled FBI (Fourier–Bros–Iagolnitzer) transform in time; this normal operator is thus equivalent to the operator $N(P)$ considered here, but used in a different manner.

Remark 1.5 (second microlocalization). Writing

$$hrD_r = (h+r) \frac{h}{h+r} rD_r \quad \text{and} \quad hD_y = (h+r) \frac{h}{h+r} D_y$$

suggests that semiclassical conormal regularity at the cone point (regularity under application of hrD_r and hD_y) can be captured on the scale of semiclassical cone Sobolev spaces as well. We present a systematic second microlocal perspective on this in Section 3D, inspired by recent work of Vasy [2021a; 2021b] on the limiting absorption principle on asymptotically conic manifolds (with the conic nature referring to the *large* end of the manifold). In view of the characterization of the quadratic form domain of $h^2\Delta_g + 1$ as a semiclassical cone Sobolev space in [Hintz 2022, Theorem 6.1], we can thus eliminate the need of working with a mixed differential-pseudodifferential calculus as in [Baskin and Marzuola 2022], and instead work in a single microlocal framework.

1B. Outline of the paper. In Section 2, we review basic notions from b- and scattering analysis, with an eye towards the relationship with semiclassical cone analysis. In Section 3, we describe a hands-on perspective on the semiclassical cone algebra $\Psi_{ch}(X)$ with a focus on its use for symbolic computations.

The heart of the paper is Section 4: we define the general class of operators to which the analysis sketched in Section 1A applies (Section 4A) and analyze in detail their symbolic properties (Section 4B), followed by a general analysis of $N(P)$ (Section 4C). We state and prove the main microlocal result, Theorem 4.10, in Section 4D. We prove the diffractive improvement in Section 4F. Finally, Section 5 contains applications of the general theory: a sharp version of propagation estimates for $h^2\Delta_g - 1$ on conic manifolds in Section 5A, and high-energy resolvent estimates for scattering by inverse square potentials and the Dirac–Coulomb equation in Sections 5B and 5C.

In Appendix B, we provide a brief summary of the Sobolev spaces and pseudodifferential calculi used, to aid the reader in keeping track of the (meanings of the) various orders involved.

2. Review of b- and scattering calculi

We denote by X a smooth n -dimensional compact manifold with nonempty, connected, and embedded boundary ∂X . The Lie algebra $\mathcal{V}_b(X) \subset \mathcal{V}(X) = \mathcal{C}^\infty(X; TX)$ of *b-vector fields* consists of all smooth vector fields on X which are tangent to ∂X . The Lie subalgebra $\mathcal{V}_{sc}(X) \subset \mathcal{V}_b(X)$ of *scattering vector fields* consists of all b-vector fields which vanish, *as b-vector fields*, at ∂X . Thus, if $x \in \mathcal{C}^\infty(X)$ denotes a boundary-defining function (meaning: $\partial X = x^{-1}(0)$, and dx does not vanish on ∂X), then $\mathcal{V}_{sc}(X) = x\mathcal{V}_b(X)$. In local coordinates $(x, y) \in [0, \infty) \times \mathbb{R}^{n-1}$ near a point on ∂X , b-vector fields are of the form

$$a(x, y)x\partial_x + \sum_{j=1}^{n-1} b^j(x, y)\partial_{y^j}, \quad a, b^1, \dots, b^{n-1} \in \mathcal{C}^\infty, \tag{2-1}$$

while scattering vector fields are of the form

$$a(x, y)x^2\partial_x + \sum_{j=1}^{n-1} b^j(x, y)x\partial_{y^j}, \quad a, b^1, \dots, b^{n-1} \in \mathcal{C}^\infty.$$

Correspondingly, there are natural vector bundles

$${}^bTX \rightarrow X, \quad {}^{sc}TX \rightarrow X, \tag{2-2}$$

isomorphic to TX° over X° , but with local frames (in local coordinates as above) given by $x\partial_x, \partial_{y^1}, \dots, \partial_{y^{n-1}}$ and $x^2\partial_x, x\partial_{y^1}, \dots, x\partial_{y^{n-1}}$ respectively, so that $\mathcal{V}_b(X) = \mathcal{C}^\infty(X; {}^bTX)$ and $\mathcal{V}_{sc}(X) = \mathcal{C}^\infty(X; {}^{sc}TX)$. Here, we implicitly use the bundle maps ${}^bTX \rightarrow TX$ and ${}^{sc}TX \rightarrow TX$ (which are isomorphisms over X° but not over ∂X) to identify $\mathcal{C}^\infty(X; {}^bTX)$ and $\mathcal{C}^\infty(X; {}^{sc}TX)$ with subspaces of $\mathcal{C}^\infty(X; TX) = \mathcal{V}(X)$. The dual bundles of (2-2) are the b-cotangent bundle and scattering cotangent bundle, ${}^bT^*X \rightarrow X$ and ${}^{sc}T^*X \rightarrow X$, with local frames $dx/x, dy^1, \dots, dy^{n-1}$ and $dx/x^2, dy^1/x, \dots, dy^{n-1}/x$, respectively. (These 1-forms are thus *smooth, nonzero* sections of ${}^bT^*X$, resp. ${}^{sc}T^*X$, down to ∂X .) Writing the canonical 1-form on T^*X° as

$$\xi_b \frac{dx}{x} + \sum_{j=1}^{n-1} (\eta_b)_j dy^j, \quad \text{resp.} \quad \xi_{sc} \frac{dx}{x^2} + \sum_{j=1}^{n-1} (\eta_{sc})_j \frac{dy^j}{x}, \tag{2-3}$$

thus defines fiber-linear coordinates (ξ_b, η_b) , resp. $(\xi_{sc}, \eta_{sc}) \in \mathbb{R} \times \mathbb{R}^{n-1}$, on ${}^bT^*X$, resp. ${}^{sc}T^*X$. The b-density bundle is denoted by ${}^b\Omega^1 X = |\Lambda^n {}^bT^*X|$; in local coordinates, its smooth sections are of the form $a|(dx/x)dy^1 \cdots dy^{n-1}|$, $a \in \mathcal{C}^\infty$.

The space of finite linear combinations of up to k -fold compositions of elements of $\mathcal{V}_\bullet(X)$, where $\bullet = \text{b, sc}$, is denoted by $\text{Diff}_\bullet^k(X)$, and we put $\text{Diff}_\bullet(X) = \bigoplus_{k \in \mathbb{N}_0} \text{Diff}_\bullet^k(X)$. The space $\text{Diff}_\text{b}(X)$ gives rise to the notion of *conormality* (relative to a fixed function space) of distributions on X° : concretely, the space

$$\mathcal{A}^\alpha(X) \subset x^\alpha L^\infty(X^\circ)$$

consists of all u so that $Au \in x^\alpha L^\infty(X^\circ)$ for all $A \in \text{Diff}_\text{b}(X)$. More generally, for $\delta \leq 1$, one can consider the space

$$\mathcal{A}_{1-\delta}^\alpha(X) \subset x^\alpha L^\infty(X^\circ)$$

of conormal distributions u of type $1 - \delta$, defined by the condition that for any $k \in \mathbb{N}_0$ and $A \in \text{Diff}_\text{b}^k(X)$, one has $Au \in x^{\alpha-k\delta} L^\infty(X^\circ)$. (Thus, $\mathcal{A}^\alpha(X) = \mathcal{A}_1^\alpha(X)$.) A more restrictive class than $\mathcal{A}^\alpha(X)$ is the class of *classical conormal distributions*, $\mathcal{A}_{\text{cl}}^\alpha(X)$, which is defined simply as

$$\mathcal{A}_{\text{cl}}^\alpha(X) = x^\alpha \mathcal{C}^\infty(X) \subset \mathcal{A}^\alpha(X).$$

Given an element $u = x^\alpha u_0 \in \mathcal{A}_{\text{cl}}^\alpha(X)$, the function u_0 is thus not merely conormal (regularity under $x\partial_x, \partial_y$), but smooth (regularity under ∂_x, ∂_y).

As an important example, let $E \rightarrow X$ denote a smooth real vector bundle of rank N , and consider the radial compactification $\bar{E} \rightarrow X$, i.e., the fiber bundle whose fiber \bar{E}_x over $x \in X$ is equal to the radial compactification of $E_x \cong \mathbb{R}^N$ defined by

$$\begin{aligned} \bar{\mathbb{R}}^N &:= (\mathbb{R}^N \sqcup ([0, \infty)_\rho \times \mathbb{S}^{n-1})) / \sim, \\ \mathbb{R}^N \setminus \{0\} \ni \rho^{-1}\omega &\sim (\rho, \omega) \in [0, \infty) \times \mathbb{S}^{n-1}. \end{aligned}$$

Then the total space \bar{E} is a manifold with corners which has two boundary hypersurfaces, $\bar{E}_{\partial X}$ (the radial compactification of $E_{\partial X}$) and $S\bar{E}$ (fiber infinity, locally defined by $\rho = 0$). On \bar{E} , we regard only $S\bar{E}$ as a boundary, in the sense that we declare $\mathcal{V}_\text{b}(\bar{E})$ to consist of all smooth vector fields on \bar{E} which are tangent to $S\bar{E}$ (but not necessarily to $\bar{E}_{\partial X}$). For $s \in \mathbb{R}$, we then put

$$S^s(\bar{E}) := \mathcal{A}^{-s}(\bar{E}).$$

One can of course consider variants of this, e.g., requiring elements of $\mathcal{V}_\text{b}(\bar{E})$ to be tangent to *both* boundary hypersurfaces and defining spaces $S^{s,r}(\bar{E})$ which are conormal of weight $-s, -r$ at $S\bar{E}, \bar{E}_{\partial X}$, respectively; or one may require classicality at one or both of the boundary hypersurfaces.

2A. *b*-pseudodifferential operators. We denote fiber infinity of the radial compactification $\overline{\text{b}T^*X}$ of $\text{b}T^*X$ by $\text{b}S^*X$. Elements of $S^s(\overline{\text{b}T^*X})$ will be symbols of *b*-pseudodifferential operators (of type $(1, 0)$, in Hörmander's (ρ, δ) -terminology [1971, §1.1]). Concretely, consider $a \in S^s(\overline{\text{b}T^*X})$ with support contained in a local coordinate patch near a point on ∂X ; thus, for all $i, j \in \mathbb{N}_0$ and $\alpha, \beta \in \mathbb{N}_0^{n-1}$, there exists a constant $C_{ij\alpha\beta}$ so that

$$|\partial_x^i \partial_y^\alpha \partial_{\xi_\text{b}}^j \partial_{\eta_\text{b}}^\beta a(x, y, \xi_\text{b}, \eta_\text{b})| \leq C_{ij\alpha\beta} (1 + |\xi_\text{b}| + |\eta_\text{b}|)^{s-(j+|\beta|)}.$$

The (left) quantization of a is then defined by

$$\begin{aligned}
 (\text{Op}_b(a)u)(x, y) &:= (2\pi)^{-n} \iiint \exp\left(i\left(\frac{x-x'}{x'}\xi_b + (y-y') \cdot \eta_b\right)\right) \phi\left(\left|\log \frac{x}{x'}\right|\right) \phi(|y-y'|) \\
 &\quad \times a(x, y, \xi_b, \eta_b) u(x', y') \frac{dx'}{x'} dy' d\xi_b d\eta_b,
 \end{aligned}$$

where $\phi \in C_c^\infty((-1, 1))$ is identically 1 near 0. The cutoff ϕ serves to make $C^{-1}x' \leq x \leq Cx'$ and $|y - y'| < c$ on the support of the Schwartz kernel of $\text{Op}_b(a)$ for some $C > 1, c > 0$, i.e., it localizes near the diagonal. We define

$$\Psi_b^s(X) := \text{Op}_b(S^s(\overline{bT^*X})) + \Psi_b^{-\infty}(X).$$

Here, if we write $\pi_{L/R} : X^2 \rightarrow X$ for the left/right projection, the space $\Psi_b^{-\infty}(X)$ of residual operators consists of all operators $\dot{C}^\infty(X) \rightarrow \dot{C}^\infty(X)$ (with $\dot{C}^\infty(X)$ denoting the space of smooth functions on X vanishing to infinite order at ∂X) whose Schwartz kernels $\kappa \in C^{-\infty}(X^2; \pi_R^{*\text{b}}\Omega^1 X)$ (the dual space of $\dot{C}^\infty(X^2; \pi_L^{*\text{b}}\Omega^1 X)$) pull back to smooth right b-densities on the b-double space⁴

$$X_b^2 := [X^2; (\partial X)^2] \tag{2-4}$$

which vanish to infinite order at the left boundary lb_b (the lift of $\partial X \times X$) and the right boundary rb_b (the lift of $X \times \partial X$) but are smooth down to the front face ff_b . (See [Vasy 2018, §6] for more details, and also [Melrose 1993; Grieser 2001].) One often encounters weighted operators as well,

$$\text{Diff}_b^{k,l}(X) := x^l \text{Diff}_b^k(X), \quad \Psi_b^{s,l}(X) := x^l \Psi_b^s(X).$$

More generally still, one can consider quantizations of symbols which are conormal of order s at ${}^bS^*X$ and of order l at ${}^bT_{\partial X}^*X$; this level of generality is occasionally useful; see, e.g., [Vasy 2021b, §5] and Section 3D. Given an operator $A \in \Psi_b^{s,l}(X)$, we denote its Schwartz kernel by K_A .

Elements of $\Psi_b^{s,l}(X)$ define continuous linear operators on $\dot{C}^\infty(X)$, and the composition of two b-ps.d.o.s is again a b-ps.d.o., with orders equal to the sum of the orders of the two factors. The principal symbol ${}^b\sigma_s : \Psi_b^{s,l}(X) \rightarrow (x^l S^s / x^l S^{s-1})(\overline{bT^*X})$ is a $*$ -homomorphism, and maps commutators into Poisson brackets. In local coordinates (and omitting orders for brevity), this means that for two operators $A, B \in \Psi_b(X)$ with principal symbols a, b , we have

$$\begin{aligned}
 {}^b\sigma(i[A, B]) &= \{a, b\} = H_a b, \\
 H_a &= (\partial_{\xi_b} a) x \partial_x + (\partial_{\eta_b} a) \partial_y - (x \partial_x a) \partial_{\xi_b} - (\partial_y a) \partial_{\eta_b}.
 \end{aligned} \tag{2-5}$$

2B. Scattering pseudodifferential operators. It is important to consider more general symbol classes than merely $S^s(\overline{\text{sc}T^*X})$ or $x^{-r} S^s(\overline{\text{sc}T^*X})$. Namely, for $\delta \in [0, \frac{1}{2})$, we shall consider the class

$$S_{1-\delta, \delta}^{s,r}(\overline{\text{sc}T^*X})$$

of symbols which are conormal at ${}^{\text{sc}}S^*X$ with weight $-s$, and conormal of type $1 - \delta$ with weight $-r$ at $\overline{\text{sc}T_{\partial X}^*X}$. (The presence of both $1 - \delta$ and δ as subscripts follows the classical literature on symbol

⁴For a detailed discussion of real blow-ups such as (2-4), we refer the reader to [Melrose 1996]. See [Hintz 2022, Appendix A] for a brief summary which is sufficient for our purposes.

classes; see, e.g., [Hörmander 1971].) This means that $S_{1-\delta,\delta}^{s,r}(\overline{\text{sc}T^*X})$ consists of all smooth functions a on $\text{sc}T^*X$ which over X° are symbols of type $(1, 0)$ and order s , i.e., $a|_{T^*X^\circ} \in S_{1,0}^s(T^*X^\circ) = S^s(\overline{T^*X^\circ})$, and which near ∂X satisfy for all $i, j \in \mathbb{N}_0$ and $\alpha, \beta \in \mathbb{N}_0^{n-1}$ an estimate

$$|(x\partial_x)^i \partial_y^\alpha \partial_{\xi_{\text{sc}}}^j \partial_{\eta_{\text{sc}}}^\beta a(x, y, \xi_{\text{sc}}, \eta_{\text{sc}})| \leq C_{ij\alpha\beta} x^{-r-(i+j+|\alpha+|\beta|)\delta} (1 + |\xi_{\text{sc}}| + |\eta_{\text{sc}}|)^{s-(j+|\beta|)}.$$

In the case $\delta = 0$, we omit the subscript “ $1 - \delta, \delta$ ”. We then define the (left) scattering quantization of a by

$$\begin{aligned} (\text{Op}_{\text{sc}}(a)u)(x, y) &:= (2\pi)^{-n} \iiint \exp\left(i\left[\frac{x-x'}{x^2}\xi_{\text{sc}} + \frac{y-y'}{x}\cdot\eta_{\text{sc}}\right]\right) \phi\left(\left|\log\frac{x}{x'}\right|\right) \phi(|y-y'|) \\ &\quad \times a(x, y, \xi_{\text{sc}}, \eta_{\text{sc}}) u(x', y') \frac{dx'}{x'^2} \frac{dy'}{x'^{n-1}} d\xi_{\text{sc}} d\eta_{\text{sc}}. \end{aligned}$$

(In this generality, scattering ps.d.o.s were introduced by Melrose [1994].) If one were working with global coordinates, one could remove the cutoffs here due to the rapid decay of the partial (in the fiber variables) inverse Fourier transform of a as $|1/x - 1/x'| + |y/x - y'/x'| \rightarrow \infty$.⁵ We then set

$$\Psi_{\text{sc},1-\delta,\delta}^{s,r}(X) := \text{Op}_{\text{sc}}(S_{1-\delta,\delta}^{s,r}(\overline{\text{sc}T^*X})) + \Psi_{\text{sc}}^{-\infty,-\infty}(X),$$

where $\Psi_{\text{sc}}^{-\infty,-\infty}(X)$ consists of all operators whose Schwartz kernels lie in $\dot{C}^\infty(X^2; \pi_R^* \Omega^1 X)$. We shall refer to s as the (scattering) differential order, and to r as the (scattering) decay order.

The principal symbol of scattering operators captures their leading-order behavior for large frequencies as well as at ∂X :

$$\text{sc}\sigma_{s,r} : \Psi_{\text{sc},1-\delta,\delta}^{s,r}(X) \rightarrow (S_{1-\delta,\delta}^{s,r}/S_{1-\delta,\delta}^{s-1,r-1+2\delta})(\overline{\text{sc}T^*X}).$$

This is a *-homomorphism. Thus, for $A_j \in \Psi_{\text{sc},1-\delta,\delta}^{s_j,r_j}(X)$, $j = 1, 2$, we have

$$[A_1, A_2] \in \Psi_{\text{sc},1-\delta,\delta}^{s_1+s_2-1,r_1+r_2-1+2\delta}(X); \tag{2-6}$$

the principal symbol (which captures the commutator modulo $\Psi_{\text{sc},1-\delta,\delta}^{s_1+s_2-2,r_1+r_2-2+4\delta}(X)$) is given in terms of the principal symbols a_1, a_2 of A_1, A_2 by

$$\begin{aligned} \text{sc}\sigma_{s_1+s_2-1,r_1+r_2-1+2\delta}(i[A_1, A_2]) &= H_{a_1} a_2, \\ x^{-1} H_{a_1} &= (\partial_{\xi_{\text{sc}}} a_1)(x\partial_x + \eta_{\text{sc}}\partial_{\eta_{\text{sc}}}) + (\partial_{\eta_{\text{sc}}} a_1)\partial_y - ((x\partial_x + \eta_{\text{sc}}\partial_{\eta_{\text{sc}}})a_1)\partial_{\xi_{\text{sc}}} - (\partial_y a_1)\partial_{\eta_{\text{sc}}}. \end{aligned} \tag{2-7}$$

We refer the reader to [Vasy 2018, §3] for more details in the special case $X = \overline{\mathbb{R}^n}$, in which case the scattering calculus is the same as the standard ps.d.o. calculus on \mathbb{R}^n for amplitudes which are product-type symbols in the base and fiber variables.

A natural setting where one must work with $\delta > 0$ arises when working with operators which have a variable scattering decay order

$$r \in C^\infty(\overline{\text{sc}T^*X}).$$

To wit, for $s \in \mathbb{R}$, we define

$$S^{s,r}(\overline{\text{sc}T^*X})$$

⁵Importantly, one typically does *not* want to localize more sharply to $|1/x - 1/x'| + |y/x - y'/x'| \lesssim 1$ (which is a small neighborhood of the lifted diagonal in the scattering double space; see [Melrose 1994, §21]), as this would thus destroy the leading-order commutativity of the scattering calculus at ∂X .

to consist of all a of the form $a = x^{-r}a_0$, where $a_0 \in \bigcap_{\delta \in (0, 1/2)} S_{1-\delta, \delta}^{s, 0}(\overline{\text{sc}T^*X})$. It is easy to check that $S^{s, r}(\overline{\text{sc}T^*X}) \subset \bigcap_{\delta \in (0, 1/2)} S_{1-\delta, \delta}^{s, r_0}(\overline{\text{sc}T^*X})$ for any $r_0 > \sup r$; in fact, differentiating variable-order symbols produces only logarithmic factors in the boundary-defining function x . Thus, we can quantize such symbols, giving rise to the space

$$\Psi_{\text{sc}}^{s, r}(X) := \text{Op}_{\text{sc}}(S^{s, r}(\overline{\text{sc}T^*X})) + \Psi_{\text{sc}}^{-\infty, -\infty}(X) \subset \bigcap_{\delta \in (0, \frac{1}{2})} \Psi_{\text{sc}, 1-\delta, \delta}^{s, r_0}(X).$$

Principal symbols of elements of $\Psi_{\text{sc}}^{s, r}(X)$ lie in $(S^{s, r} / \bigcap_{\delta > 0} S^{s-1, r-1+2\delta})(\overline{\text{sc}T^*X})$. The (variable) orders are additive under operator composition; this is a consequence of the formula for the full symbol (in local coordinates) of the composition of two ps.d.o.s.

We point out that for fixed $s \in \mathbb{R}$ the space $S^{s, r}(\overline{\text{sc}T^*X})$ (and thus $\Psi_{\text{sc}}^{s, r}(X)$) only depends on the restriction of r to $\overline{\text{sc}T_{\partial X}^*X}$. Indeed, given $r' \in C^\infty(\overline{\text{sc}T^*X})$ with $r' - r = 0$ at $\overline{\text{sc}T_{\partial X}^*X}$, we can write $r' - r = xw$, $w \in C^\infty(\overline{\text{sc}T^*X})$, and therefore $x^{-r'} = x^{-r} \exp(-wx \log x)$; by direct differentiation, one then finds that $\exp(-wx \log x) \in S_{1-\delta, \delta}^{0, 0}(\overline{\text{sc}T^*X})$ for any $\delta > 0$. In view of this, we can define $S^{s, r}(\overline{\text{sc}T^*X})$ and $\Psi_{\text{sc}}^{s, r}(X)$, given a variable order

$$r \in C^\infty(\overline{\text{sc}T_{\partial X}^*X}),$$

to be equal to $S^{s, \tilde{r}}(\overline{\text{sc}T^*X})$ and $\Psi_{\text{sc}}^{s, \tilde{r}}(X)$, respectively, where $\tilde{r} \in C^\infty(\overline{\text{sc}T^*X})$ is any smooth extension of r .

2C. Sobolev spaces. We next recall the corresponding scales of weighted Sobolev spaces. We have some flexibility in the choice of the underlying L^2 -space. Thus, fix any smooth positive b -density $\mu_0 \in C^\infty(X; {}^b\Omega^1 X)$, and fix $a_\mu \in \mathbb{R}$. We then set $\mu := x^{a_\mu} \mu_0$ and

$$H_b^0(X; \mu) \equiv L_b^2(X; \mu) \equiv H_{\text{sc}}^0(X; \mu) \equiv L_{\text{sc}}^2(X; \mu) := L^2(X; \mu). \tag{2-8}$$

These spaces are independent of the choice of μ_0 (but not a_μ), up to equivalence of norms; the same will be true for the spaces defined in the sequel. When the density μ is fixed and clear from the context, we drop it from the notation. Let $\bullet = b, \text{sc}$. For $s \geq 0$, we then let

$$H_\bullet^s(X) := \{u \in H_\bullet^0(X) : Au \in H_\bullet^0(X)\},$$

where $A \in \Psi_\bullet^s(X)$ denotes any fixed elliptic operator. For $s < 0$, we define $H_\bullet^s(X) = (H_\bullet^{-s}(X))^*$ with respect to the $L_\bullet^2(X)$ inner product; an equivalent definition is given by $H_\bullet^s(X) = \{u_1 + Au_2 : u_1, u_2 \in H_\bullet^0(X)\}$, where $A \in \Psi_\bullet^{-s}(X)$ is elliptic. Weighted spaces are defined by

$$H_b^{s, l}(X) = x^l H_b^s(X), \quad H_{\text{sc}}^{s, r}(X) = x^r H_{\text{sc}}^s(X).$$

Finally, we define scattering Sobolev spaces with variable decay orders $r \in C^\infty(\overline{\text{sc}T_{\partial X}^*X})$ by taking $r_0 < \inf r$ and putting

$$H_{\text{sc}}^{s, r}(X) := \{u \in H_{\text{sc}}^{s, r_0}(X) : Au \in H_{\text{sc}}^0(X)\},$$

where $A \in \Psi_{\text{sc}}^{s, r}(X)$ is any fixed elliptic operator.

2D. b -scattering operators and Sobolev spaces. In our application, we shall encounter a compact manifold X whose boundary ∂X has *two* connected components, say H_1, H_2 , both of which are embedded.

We can then consider the space $\mathcal{V}_{b,sc}(X)$ of b-scattering vector fields (which localized to a neighborhood of H_1 , resp. H_2 , lie in \mathcal{V}_b , resp. \mathcal{V}_{sc}), the corresponding b-scattering tangent bundle ${}^{b,sc}TX$ and its dual ${}^{b,sc}T^*X$, as well as weighted b-scattering Sobolev spaces,

$$H_{b,sc}^{s,l,r}(X), \quad s, l \in \mathbb{R}, \quad r \in \mathcal{C}^\infty(\overline{{}^{b,sc}T^*X}).$$

Localized to a neighborhood of H_1 , its elements lie in $H_b^{s,l}$, and localized to a neighborhood of H_2 , they lie in $H_{sc}^{s,r}$.

Let us make this even more concrete in the setting which will arise below,

$$X = [0, \infty]_{\hat{x}} \times Y, \quad H_1 = \hat{x}^{-1}(0), \quad H_2 = \hat{x}^{-1}(\infty), \tag{2-9}$$

where we write $[0, \infty]$ for the closure of $[0, \infty)$ inside of $\overline{\mathbb{R}}$; here Y is a compact $(n-1)$ -dimensional manifold without boundary. Then $\hat{x}/(\hat{x}+1)$ and $(1+\hat{x})^{-1}$ are defining functions of H_1 and H_2 , respectively; hence $\mathcal{V}_{b,sc}(X) = (1+\hat{x})^{-1}\mathcal{V}_b(X)$. Using local coordinates y^1, \dots, y^{n-1} on an open subset $U \subset Y$, the collection of 1-forms

$$(1+\hat{x})\frac{d\hat{x}}{\hat{x}}, \quad (1+\hat{x})dy^1, \quad \dots, \quad (1+\hat{x})dy^{n-1}$$

is a smooth frame of ${}^{b,sc}T^*X$ over $[0, \infty] \times U$. Denoting the corresponding fiber-linear coordinates on ${}^{b,sc}T^*X$ by $(\xi_{b,sc}, \eta_{b,sc}) \in \mathbb{R} \times \mathbb{R}^{n-1}$, we can then quantize a symbol⁶

$$a \in S^{s,l,r}(\overline{{}^{b,sc}T^*X}) = \left(\frac{\hat{x}}{\hat{x}+1}\right)^{-l} (1+\hat{x})^r S^s(\overline{{}^{b,sc}T^*X})$$

by

$$\begin{aligned} (\text{Op}_{b,sc}(a)u)(\hat{x}, y) &:= (2\pi)^{-n} \iiint \exp\left(i\left(\frac{\hat{x}-\hat{x}'}{\hat{x}\frac{1}{1+\hat{x}}}\xi_{b,sc} + \frac{y-y'}{\frac{1}{1+\hat{x}}}\cdot\eta_{b,sc}\right)\right) \\ &\quad \times \phi\left(\left|\log\frac{\hat{x}}{\hat{x}'}\right|\right)\phi(|y-y'|)a(\hat{x}, y, \xi_{b,sc}, \eta_{b,sc})u(\hat{x}', y') \\ &\quad \times \frac{d\hat{x}'}{\hat{x}'\frac{1}{1+\hat{x}'}} \frac{dy'}{\left(\frac{1}{1+\hat{x}'}\right)^{n-1}} d\xi_{b,sc} d\eta_{b,sc}. \end{aligned} \tag{2-10}$$

The space $\Psi_{b,sc}^{s,l,r}(X)$ of b-scattering ps.d.o.s is then the sum

$$\Psi_{b,sc}^{s,l,r}(X) = \text{Op}_{b,sc}(S^{s,l,r}(\overline{{}^{b,sc}T^*X})) + \Psi_{b,sc}^{-\infty,l,-\infty}(X).$$

Here,

$$\Psi_{b,sc}^{-\infty,l,-\infty}(X) = \left(\frac{\hat{x}}{\hat{x}+1}\right)^{-l} \Psi_{b,sc}^{-\infty,0,-\infty}(X)$$

is defined momentarily. First define the double space

$$X_{b,sc}^2 := \left[\overline{[0, \infty)^2} \times Y^2; (\{0\} \times Y)^2; \Delta \cap (\{\infty\} \times Y)^2\right], \tag{2-11}$$

where $\overline{[0, \infty)^2}$ is the radial compactification (equivalently, the closure of $[0, \infty)^2$ inside of $\overline{\mathbb{R}^2}$), and $\Delta \subset \overline{[0, \infty)^2} \times Y^2$ is the diagonal. Then the space $\Psi_{b,sc}^{-\infty,0,-\infty}(X)$ consists of all operators whose Schwartz

⁶We leave the minor, largely notational, changes to accommodate symbols with variable scattering decay orders r to the reader.




Figure 2. The double space $X_{b,sc}^2$ without the factor Y^2 . The dashed line is the lifted diagonal. Indicated are the symbolic orders of Schwartz kernels of elements of $\Psi_{b,sc}^{s,l,r}(X)$.

kernels are smooth right b-densities on $X_{b,sc}^2$ which vanish to infinite order at all boundary hypersurfaces except for the lift of $(\{0\} \times Y)^2$. See Figure 2. Moreover, Schwartz kernels of elements of $\Psi_{b,sc}^{s,0,r}(X)$ are conormal of order s to the lifted diagonal in $X_{b,sc}^2$ smoothly down to the lift of $(\{0\} \times Y)^2$, conormal with weight $-r$ down to the lift of $\Delta \cap (\{\infty\} \times Y)^2$, and vanish to infinite order at all other boundary hypersurfaces.

3. Semiclassical cone calculus

We revisit and generalize the algebra $\Psi_{ch}(X)$ and the associated scale of weighted Sobolev spaces from [Hintz 2022], give a user-friendly treatment of the symbol calculus (including Poisson brackets), and study operators and function spaces with variable (semiclassical) orders and their behavior upon restriction to the transition faces of the semiclassical cone single and double spaces (recalled later in this section). Throughout this section, we denote by X a compact n -dimensional manifold with nonempty, connected, and embedded boundary ∂X . We denote by $x \in C^\infty(X)$ a boundary-defining function.

3A. Vector fields, bundles, Poisson brackets. We recall from Section 1A the *semiclassical cone single space*

$$X_{ch} := [[0, 1)_h \times X; \{0\} \times \partial X],$$

the boundary hypersurfaces of which we denote by cf (conic face, lift of $[0, 1) \times \partial X$), tf (transition face, the front face@), and sf (semiclassical face, lift of $\{0\} \times X$). See Figure 3. Defining functions of these boundary hypersurfaces are $x/(x+h)$, $x+h$, and $h/(h+x)$, respectively. On $X_{ch} \setminus \text{cf}$, it is convenient to use the local defining functions x of tf and h/x of sf.

Definition 3.1 (vector fields). We define the space

$$\mathcal{V}_{ch}(X_{ch})$$

of *semiclassical cone vector fields* to consist of all b-vector fields $V \in \mathcal{V}_b(X_{ch})$ which are horizontal, i.e., tangent to the fibers of $X_{ch} \rightarrow [0, 1)_h$, and whose restriction to sf vanishes.

Lemma 3.2 (spanning set). *Identifying a vector field $V \in \mathcal{V}_b(X)$ with its horizontal lift to X_{ch} along $X_{ch} \rightarrow X$, the space $\mathcal{V}_{ch}(X_{ch})$ is spanned over $C^\infty(X_{ch})$ by $(h/(h+x))\mathcal{V}_b(X)$. Moreover, given $V, W \in \mathcal{V}_{ch}(X_{ch})$, we have $[V, W] \in (h/(h+x))\mathcal{V}_{ch}(X_{ch})$.*




Figure 3. The semiclassical cone single space X_{ch} .

This allows us to define the graded ring

$$\text{Diff}_{ch}(X) = \bigoplus_{k \in \mathbb{N}_0} \text{Diff}_{ch}^k(X)$$

of differential operators in the usual manner.

Proof. Directly from the definition, we have $(h/(h+x))\mathcal{V}_b(X) \subset \mathcal{V}_{ch}(X_{ch})$. Conversely, suppose $V \in \mathcal{V}_{ch}(X_{ch})$. Let us work in local coordinates $(x, y) \in [0, \infty) \times \mathbb{R}^{n-1}$ near a point in ∂X . Near cf, we use the local coordinates (h, \hat{x}, y) with $\hat{x} := x/h$. From the definition, we have

$$V = a(h, \hat{x}, y)\hat{x}\partial_{\hat{x}} + \sum_{j=1}^{n-1} b^j(h, \hat{x}, y)\partial_{y^j}, \tag{3-1}$$

with $a, b^j \in C^\infty$. Since $\hat{x}\partial_{\hat{x}} = x\partial_x$, this expresses V in the desired form.

Near sf on the other hand, we use (\hat{h}, x, y) with $\hat{h} := h/x$. Since $V \in \mathcal{V}_b(X_{ch})$, we can write

$$V = a(\hat{h}, x, y)(x\partial_x - \hat{h}\partial_{\hat{h}}) + \tilde{a}(\hat{h}, x, y)\hat{h}\partial_{\hat{h}} + \sum_{j=1}^{n-1} b^j(\hat{h}, x, y)\partial_{y^j}.$$

The horizontal nature of V means $0 = Vh = V(x\hat{h}) = \tilde{a}x\hat{h}$, which implies $\tilde{a} \equiv 0$ by continuity from $(X_{ch})^\circ = \{x > 0, \hat{h} > 0\}$. The vanishing of V at $\hat{h} = 0$ as a b-vector field implies, in addition, that $a = \hat{h}a'$ and $b^j = \hat{h}b'_j$ with $a', b'_j \in C^\infty$. Since the horizontal lifts of $x\partial_x, \partial_{y^j} \in \mathcal{V}_b(X)$ to X_{ch} are equal to $x\partial_x - \hat{h}\partial_{\hat{h}}, \partial_{y^j}$, the claim follows.

Regarding the Lie algebra structure, we compute, for $V, W \in \mathcal{V}_b(X)$,

$$\left[\frac{h}{h+x}V, \frac{h}{h+x}W \right] = \frac{h}{h+x} \left(\frac{h}{h+x}[V, W] + V \left(\frac{h}{h+x} \right) W - W \left(\frac{h}{h+x} \right) V \right).$$

Since $V, W \in \mathcal{V}_b(X_{ch})$, we have $V(h/(h+x)), W(h/(h+x)) \in (h/(h+x))C^\infty(X_{ch})$. □

There exists a vector bundle

$${}^{ch}TX_{ch} \rightarrow X_{ch}$$

together with a smooth bundle map ${}^{ch}TX_{ch} \rightarrow {}^bTX_{ch}$ so that the space $\mathcal{V}_{ch}(X_{ch})$ is equal to the space of smooth sections of ${}^{ch}TX_{ch}$. In local coordinates on X , a local frame of ${}^{ch}TX_{ch}$ is given by (the horizontal lifts to X_{ch} of)

$$\frac{h}{h+x}x\partial_x, \quad \frac{h}{h+x}\partial_{y^1}, \quad \dots, \quad \frac{h}{h+x}\partial_{y^{n-1}}.$$

We call ${}^{ch}TX_{ch}$ the *ch-tangent bundle* and its dual ${}^{ch}T^*X_{ch}$ the *ch-cotangent bundle*, with local frame

$$\frac{x+h}{h} \frac{dx}{x}, \quad \frac{x+h}{h} dy^1, \quad \dots, \quad \frac{x+h}{h} dy^{n-1}.$$

A choice of local coordinates $(x, y) \in [0, \infty) \times \mathbb{R}^{n-1}$ on an open set $U \subset X$ induces a trivialization of ${}^{ch}T^*X_{ch}$ over the preimage of $[0, 1) \times U$ under $X_{ch} \rightarrow X$, with fiber-linear coordinates $(\xi_{ch}, \eta_{ch}) \in \mathbb{R} \times \mathbb{R}^{n-1}$ defined by the requirement that the canonical 1-form on T^*X° be equal to

$$\xi_{ch} \frac{x+h}{h} \frac{dx}{x} + \sum_{j=1}^{n-1} (\eta_{ch})_j \frac{x+h}{h} dy^j. \tag{3-2}$$

In $X_{ch} \setminus \text{cf}$, where a smooth frame of ${}^{ch}TX_{ch}$ is given by $(h/x)x\partial_x, (h/x)\partial_{y^1}, \dots, (h/x)\partial_{y^{n-1}}$, it is computationally simpler to use the fiber-linear coordinates (ξ, η) in which the canonical 1-form takes the form

$$\xi \frac{x}{h} \frac{dx}{x} + \sum_{j=1}^{n-1} \eta_j \frac{x}{h} dy^j. \tag{3-3}$$

We compute the form of the Hamilton vector field H_a of a smooth function $a \in C^\infty({}^{ch}T^*X_{ch})$ in these fiber coordinates, and using (\hat{h}, x, y) with $\hat{h} = h/x$ as coordinates on the base. In terms of the coordinates on ${}^bT^*X$ used in (2-3), we have $(\xi, \eta) = (h/x)(\xi_b, \eta_b)$ and thus, by changing coordinates in the expression (2-5),

$$H_a = \hat{h}((\partial_\xi a)(x\partial_x - \hat{h}\partial_{\hat{h}} - \eta\partial_\eta) + (\partial_\eta a)\partial_y - ((x\partial_x - \hat{h}\partial_{\hat{h}} - \eta\partial_\eta)a)\partial_\xi a - (\partial_y a)\partial_\eta). \tag{3-4}$$

3B. Symbols, pseudodifferential operators, Sobolev spaces. A simple symbol class for *ch*-operators is $S^s({}^{ch}T^*X_{ch}) = A^{-s}(\overline{{}^{ch}T^*X_{ch}})$, where we only regard fiber infinity ${}^{ch}S^*X_{ch}$ as a boundary, i.e., we require symbols to be smooth down to $\overline{{}^{ch}T^*X_{ch}}$ for $\bullet = \text{cf}, \text{tf}, \text{sf}$. In practice, we need more general symbols: for $\delta \in [0, \frac{1}{2})$ and for $s, l, \alpha, b \in \mathbb{R}$, we define

$$S_{1-\delta, \delta}^{s, l, \alpha, b}(\overline{{}^{ch}T^*X_{ch}}) = \left(\frac{x}{x+h}\right)^{-l} (x+h)^{-\alpha} \left(\frac{h}{h+x}\right)^{-b} S_{1-\delta, \delta}^{s, 0, 0, 0}(\overline{{}^{ch}T^*X_{ch}})$$

to consist of all symbols which are conormal with weight $-s$ at ${}^{ch}S^*X_{ch}$, conormal with weight $-l$ at $\overline{{}^{ch}T_{\text{cf}}^*X_{ch}}$ and with weight $-\alpha$ at $\overline{{}^{ch}T_{\text{tf}}^*X_{ch}}$, and conormal of type $1 - \delta$ at $\overline{{}^{ch}T_{\text{sf}}^*X_{ch}}$ with weight $-b$. In the coordinates (3-3), the membership $a \in S_{1-\delta, \delta}^{s, 0, 0, 0}(\overline{{}^{ch}T^*X_{ch}})$ is equivalent to $a = a(\hat{h}, x, y, \xi, \eta)$ (with $\hat{h} = h/x$) satisfying estimates

$$|(x\partial_x)^i \partial_y^\alpha (\hat{h}\partial_{\hat{h}})^j \partial_\xi^k \partial_\eta^\beta a(\hat{h}, x, y, \xi, \eta)| \leq C_{ijk\alpha\beta} (1 + |\xi| + |\eta|)^{s-(k+|\beta|)} \hat{h}^{-(i+j+k+|\alpha|+|\beta|)\delta}$$

for all $i, j, k \in \mathbb{N}_0$ and $\alpha, \beta \in \mathbb{N}_0^{n-1}$; in coordinates $(h, \hat{x}, y, \xi_b, \eta_b)$ on the *ch*-cotangent bundle over $X_{ch} \setminus \text{sf}$, with $\hat{x} = x/h$ and with the canonical 1-form given by (2-3), a must satisfy

$$|(\hat{x}\partial_{\hat{x}})^i \partial_y^\alpha (h\partial_h)^j \partial_{\xi_b}^k \partial_{\eta_b}^\beta a(h, \hat{x}, y, \xi_b, \eta_b)| \leq C_{ijk\alpha\beta} (1 + |\xi_b| + |\eta_b|)^{s-(k+|\beta|)}.$$

See Figure 4. As usual, we omit the subscript “ $1 - \delta, \delta$ ” when $\delta = 0$.




Figure 4. Illustration of $\overline{chT^*X_{ch}}$ (showing only part of the compactified fibers) and the symbol class $S^{s,l,\alpha,b}(\overline{chT^*X_{ch}})$, indicating the orders at the various boundary hypersurfaces of $\overline{chT^*X_{ch}}$.

It is occasionally useful to restrict attention to symbols which are *classical* conormal down to tf, which amounts to replacing $x\partial_x, h\partial_h$ in the above symbol estimates (which are for symbols of order 0 at tf) by ∂_x, ∂_h . We denote the corresponding symbol classes with a subscript “cl” as in $S_{cl}^{s,l,\alpha,b}(\overline{chT^*X_{ch}})$.

As in Section 2, the main use of $\delta > 0$ is to accommodate symbols with variable orders. Here, we only discuss the case of variable semiclassical orders. Thus, consider $b \in C^\infty(\overline{chT_{sf}^*X_{ch}})$, an arbitrary extension of which to an element of $C^\infty(\overline{chT^*X_{ch}})$ we denote by the same letter; we then put

$$S^{s,l,\alpha,b}(\overline{chT^*X_{ch}}) := \left\{ \left(\frac{h}{h+x} \right)^b a_0 : a_0 \in \bigcap_{\delta \in (0, \frac{1}{2})} S_{1-\delta,\delta}^{s,l,\alpha,0}(\overline{chT^*X_{ch}}) \right\},$$

which is a subset of $\bigcap_{\delta \in (0, 1/2)} S_{1-\delta,\delta}^{s,l,\alpha,b_0}(\overline{chT^*X_{ch}})$ for any $b_0 > \sup b$.

We now proceed to quantize symbols $a = a(h, x, y, \xi_{ch}, \eta_{ch})$, thereby giving meaning to the formal expression

$$\text{“Op}_{c,h}(a) = a\left(h, x, y, \frac{h}{h+x}x D_x, \frac{h}{h+x}D_y\right)\text{”}.$$

Thus, fixing $\phi \in C_c^\infty((-1, 1))$, identically 1 near 0, we define, in local coordinates (x, y) on X ,

$$\begin{aligned} (\text{Op}_{c,h}(a)u)(h, x, y) &:= (2\pi)^{-n} \iiint \exp\left(i\left[\frac{x-x'}{x\frac{h}{h+x}}\xi_{ch} + \frac{y-y'}{\frac{h}{h+x}}\cdot\eta_{ch}\right]\right) \\ &\quad \times \phi\left(\left|\log\frac{x}{x'}\right|\right)\phi(|y-y'|)a(h, x, y, \xi_{ch}, \eta_{ch})u(h, x', y') \\ &\quad \times \frac{dx'}{x'\frac{h}{h+x'}} \frac{dy'}{\left(\frac{h}{h+x'}\right)^{n-1}} d\xi_{ch} d\eta_{ch} \end{aligned} \quad (3-5)$$

for a and u supported in the coordinate chart; for general a, u , one defines $\text{Op}_{c,h}(a)u$ using a partition of unity.

We interpret this in terms of the $c\hbar$ -double space

$$X_{ch}^2 := [0, 1)_h \times X_b^2; \{0\} \times \text{ff}_b; \{0\} \times \Delta_b],$$




Figure 5. The ch -double space X_{ch}^2 .

where we denote by $\Delta_b \subset X_b^2$ the lift of the diagonal in X^2 to X_b^2 ; see (2-4) and the subsequent paragraph for the definition of X_b^2 and its boundary hypersurfaces lb_b, ff_b, rb_b . First, recall from⁷ [Hintz 2022, Definition 3.1] that $lb_2, rb_2, ff_2, tf_2, sf_2$, and df_2 are the lifts of $[0, 1) \times lb_b, [0, 1) \times rb_b, [0, 1) \times ff_b, \{0\} \times ff_b, \{0\} \times X_b^2$, and $\{0\} \times \Delta_b$, respectively; moreover, Δ_{ch} denotes the lift of $[0, 1) \times \Delta_b$. See Figure 5.

Then the Schwartz kernel of $Op_{c,h}(a)$ is a conormal distribution of order $s - \frac{1}{4}$ at Δ_{ch} , conormal down to ff_2, tf_2, df_2 with weights $-l, -\alpha, -b$, and vanishes identically in a neighborhood of lb_2, rb_2, sf_2 .

The composition of two ch -quantizations is almost a ch -quantization itself; one merely has to allow for additional residual terms: Define the space $\Psi_{ch}^{-\infty}(X)$ of residual operators to consist of all operators whose Schwartz kernels are conormal sections of the right b-density bundle on X_{ch}^2 , with weight 0 at ff_2 and tf_2 , and with infinite order vanishing at lb_2, rb_2, df_2, sf_2 . We then put

$$\Psi_{ch}^s(X) := Op_{ch}(S^s(\overline{chT^*X_{ch}})) + \Psi_{ch}^{-\infty}(X),$$

where $Op_{ch} = (Op_{c,h})_{h \in (0,1)}$; this gives the same space as [Hintz 2022, Definition 3.2]. More generally, we define the quantization of symbols $a \in S_{1-\delta,\delta}^{s,l,\alpha,b}(\overline{chT^*X_{ch}})$ by the same formula (3-5); the space of residual operators is now

$$\Psi_{ch}^{-\infty,l,\alpha,-\infty}(X) := \left(\frac{x}{x+h} \right)^{-l} (x+h)^{-\alpha} \Psi_{ch}^{-\infty}(X).$$

Thus, we can now define the spaces

$$\begin{aligned} \Psi_{ch,1-\delta,\delta}^{s,l,\alpha,b}(X) &:= Op_{ch}(S_{1-\delta,\delta}^{s,l,\alpha,b}(\overline{chT^*X_{ch}})) + \Psi_{ch}^{-\infty,l,\alpha,-\infty}(X), \\ \Psi_{ch}^{s,l,\alpha,b}(X) &:= Op_{ch}(S^{s,l,\alpha,b}(\overline{chT^*X_{ch}})) + \Psi_{ch}^{-\infty,l,\alpha,-\infty}(X), \end{aligned}$$

where in the second line $b \in C^\infty(\overline{chT_{sf}^*X_{ch}})$ is a variable-order function. Their Schwartz kernels can be characterized as being conormal distributions (of order $s - \frac{1}{4}$ and type $(1, 0)$) at Δ_{ch} which are conormal at ff_2 (with weight $-l$), tf_2 (with weight $-\alpha$), and conormal of type $1 - \delta$ at df_2 (with weight $-b$), and

⁷We add subscripts “2” here in order to avoid confusion during the frequent changes between X_{ch} and X_{ch}^2 later on.

which vanish to infinite order at lb_2, rb_2, sf_2 . One can also consider subalgebras which are classical at tf , i.e., the symbols are required to be classical conormal at tf , and the residual operators are required to have classical conormal Schwartz kernels at tf_2 ; we denote these algebras by a subscript “cl”, such as

$$\Psi_{ch,cl}^{s,l,\alpha,b}(X).$$

All such ps.d.o.s define h -dependent families of bounded⁸ linear maps on $\dot{C}^\infty(X)$; compositions of two such ps.d.o.s give a ps.d.o. in the same class, with orders given by the sum of the orders of the two factors. The principal symbol map is

$${}^{ch}\sigma_{s,l,\alpha,b} : \Psi_{ch,1-\delta,\delta}^{s,l,\alpha,b}(X) \rightarrow (S^{s,l,\alpha,b}/S^{s-1,l,\alpha,b-1+2\delta})({}^{ch}\overline{T^*X_{ch}}),$$

similarly for the variable-order spaces (with $\delta > 0$ then arbitrary), and it is a $*$ -homomorphism. These facts follow from a minor variation of [Hintz 2022, Proposition 3.9] (using weights instead of index sets), with the statements about principal symbols following by continuity from the corresponding statements for standard semiclassical operators (of type $(1 - \delta, \delta)$) in $x > 0$ and b-ps.d.o.s in $h > 0$; we leave the details to the reader. We moreover have, for $A_j \in \Psi_{ch,1-\delta,\delta}^{s_j,l_j,\alpha_j,b_j}(X)$, $j = 1, 2$, with principal symbols a_j ,

$$\text{Op}_{c,h}(i[A_1, A_2]) - \text{Op}_{c,h}(H_{a_1}a_2) \in \Psi_{ch,1-\delta,\delta}^{s-2,l,\alpha,b-2+4\delta}(X),$$

analogously for variable-order operators. One can evaluate $H_{a_1}a_2$ using the formula (3-4).

Since the principal symbol captures operators to leading order at the union of boundary hypersurfaces ${}^{ch}S^*X_{ch} \cup {}^{ch}\overline{T_{sf}^*X_{ch}}$, the latter set is also the locus of the elliptic and wave front sets of an operator. Thus, for $A \in \Psi_{ch}^{s,l,\alpha,b}(X)$, we define

$$\text{Ell}_{ch}^{s,l,\alpha,b}(A), \text{WF}'_{ch}{}^{l,\alpha}(A) \subset {}^{ch}S^*X_{ch} \cup {}^{ch}\overline{T_{sf}^*X_{ch}}$$

as follows: $\text{Ell}_{ch}^{s,l,\alpha,b}(A)$ is the set of all ζ so that ${}^{ch}\sigma_{s,l,\alpha,b}(A)$ is elliptic in a neighborhood of ζ , and $\text{WF}'_{ch}{}^{l,\alpha}(A)$ is the complement of the set of ζ so that the full symbol of A lies in $S^{-\infty,l,\alpha,-\infty}({}^{ch}\overline{T^*X_{ch}})$ when localized to a sufficiently small neighborhood of ζ . In particular, we have $\text{WF}'_{ch}{}^{l,\alpha}(A) = \emptyset$ if and only if $A \in \Psi_{ch}^{-\infty,l,\alpha,-\infty}(X)$. We omit the orders s, l, α, b and l, α when they are clear from the context. The definitions for type- $(1 - \delta, \delta)$ and variable-order operators are analogous. See Figure 6.

Finally, we define the corresponding weighted Sobolev spaces. As in (2-8), we first fix a weighted b-density $\mu = x^{\alpha_\mu} \mu_0$, where $0 < \mu_0 \in C^\infty(X; {}^b\Omega^1 X)$ and $\alpha_\mu \in \mathbb{R}$, and define

$$H_{c,h}^0(X; \mu) := L^2(X; \mu), \quad H_{c,h}^{0,l,\alpha,b}(X; \mu) := \left(\frac{x}{x+h}\right)^l (x+h)^\alpha \left(\frac{h}{h+x}\right)^b H_{c,h}^0(X; \mu).$$

These spaces depend on α_μ , but are independent of μ_0 (up to equivalence of norms). When the choice of μ is clear from the context, we will omit it from the notation. For $s \geq 0$, we then define $H_{c,h}^{s,l,\alpha,b}(X)$ to consist of all $u \in H_{c,h}^{0,l,\alpha,b}(X)$ so that $Au \in H_{c,h}^0(X)$ for any (thus all) elliptic $A \in \Psi_{ch}^{s,l,\alpha,b}(X)$. We note for $s \in \mathbb{N}_0$ the equivalent characterization

$$H_{c,h}^{s,l,\alpha,b}(X) = \{u \in H_{c,h}^{0,l,\alpha,b}(X) : V_1 \cdots V_j u \in H_{c,h}^{0,l,\alpha,b}(X) \text{ for all } V_i \in \mathcal{V}_{ch}(X_{ch}), 0 \leq i \leq j \leq s\}.$$

⁸Though not uniformly in h unless $b \geq 0$.




Figure 6. The shaded boundary hypersurfaces are the locus of the elliptic set as well as of operator and distributional wave front sets; see also Figure 4.

For $s < 0$, the space $H_{c,h}^{s,l,\alpha,b}(X)$ can be defined either as the dual space $(H_{c,h}^{-s,-l,-\alpha,-b}(X))^*$, or as the space of all $u_1 + Au_2$ where $u_1, u_2 \in H_{c,h}^{0,l,\alpha,b}(X)$ and $A \in \Psi_{ch}^{-s}(X)$. Lastly, for a variable order $b \in C^\infty(\overline{ch T_{sf}^* X_{ch}})$, we pick $b_0 < \inf b$ and put

$$H_{c,h}^{s,l,\alpha,b}(X) := \{u \in H_{c,h}^{s,l,\alpha,b_0}(X) : Au \in H_{c,h}^0(X)\},$$

where $A \in \Psi_{ch}^{s,l,\alpha,b}(X)$ is any elliptic operator; the space $H_{c,h}^{s,l,\alpha,b}(X)$ is independent of the choices of b_0 and A , up to equivalence of norms.

We can define Sobolev wave front sets in the usual manner. Let $l, \alpha \in \mathbb{R}$, and suppose that we are given a distribution $u \in H_{c,h}^{-\infty,l,\alpha,-\infty}(X)$, meaning $u \in H_{c,h}^{-N,l,\alpha,-N}(X)$ for some $N \in \mathbb{R}$. Let $s, b \in \mathbb{R}$. Then

$$WF_{ch}^{s,l,\alpha,b}(u) \subset ch S^* X_{ch} \cup \overline{ch T_{sf}^* X_{ch}}$$

is the complement of all α so that there exists an operator $A \in \Psi_{ch}^{s,l,\alpha,b}(X)$, elliptic at α , so that $Au \in H_{c,h}^0(X)$. (The a priori assumption on u is familiar from the definition of the b -wave front set, see, e.g., [Vasy 2018, Definition 6.2], and ensures that one then also has $Bu \in H_{c,h}^0(X)$ for any $B \in \Psi_{ch}^{s,l,\alpha,b}(X)$ with $WF'_{ch}(B) \subset Ell'_{ch}(A)$.)

Remark 3.3 (operators on vector bundles). If $E, F \rightarrow X$ are smooth vector bundles, one can consider semiclassical cone ps.d.o.s acting between sections of E, F , giving rise to classes $\Psi_{ch}^s(X; E, F)$ and function spaces $H_{c,h}^s(X; E)$ etc. More generally, one can allow E, F to be vector bundles $E, F \rightarrow X_{ch}$ over the semiclassical single space, with Schwartz kernels of elements of $\Psi_{ch}^s(X; E, F)$ defined by taking the tensor product of $\Psi_{ch}^s(X)$ over $C^\infty(X_{ch}^2)$ with $C^\infty(X_{ch}^2; \pi_L^* E \boxtimes \pi_R^* F^*)$, where $\pi_L, \pi_R : X_{ch}^2 \rightarrow X_{ch}$ are the stretched left and right projections. Using such ps.d.o.s, one can define Sobolev spaces $H_{c,h}^s(X; E)$ etc. in this generality.

Remark 3.4 (relationship with edge Sobolev spaces). For the propagation through cone points in the spacetime setting, many authors [Melrose and Wunsch 2004; Melrose et al. 2008] have utilized Mazzeo’s edge algebra [1991]. A typical example is the operator $-D_t^2 + \Delta_g$, where $g = g(x, y, dy)$ is a conic metric on a manifold X with boundary (see (4-2)); upon multiplication by x^2 , this is a second-order differential operator, the principal part of which is a Lorentzian signature quadratic form in the collection

(xD_t, xD_x, D_y) of edge vector fields. The membership $u \in H_e^1(\mathbb{R}_t \times X, |dt \, dg|)$ — meaning that $u, xD_t u, xD_x u, D_y u \in L^2$ — can then be characterized by taking the Fourier transform in t as

$$\hat{u}(\sigma), x|\sigma|\hat{u}(\sigma), xD_x\hat{u}(\sigma), D_y\hat{u}(\sigma) \in L^2(\mathbb{R}_\sigma; L^2(X; |dg|)).$$

Introducing $h = \langle \sigma \rangle^{-1}$, this is equivalent to the $L^2(\mathbb{R}_\sigma; L^2(X))$ membership of $((h+x)/h)\hat{u}, xD_x\hat{u}, D_y\hat{u}$. Upon multiplication by $h/(h+x)$, we thus find

$$u \in H_e^1(\mathbb{R}_t \times X, |dt \, dg|) \iff \hat{u} \in L^2(\mathbb{R}_\sigma; H_{c, \langle \sigma \rangle^{-1}}^{1,0,0,1}(X; |dg|)),$$

and the respective norms of u and \hat{u} are equivalent. (One can show that similar spectral characterizations of edge Sobolev spaces remain valid also for spaces with weights and with variable differential orders; the details will be given elsewhere.)

3C. Restriction to tf. Symbolic arguments for the analysis of semiclassical cone PDEs $Pu = f$ can at best control u microlocally at ${}^{ch}S^*X_{ch} \cup \overline{{}^{ch}T_{sf}^*X_{ch}}$, i.e., modulo errors which are trivial at infinite frequencies and at sf. Crucially however, such errors may well be nontrivial at tf, and thus nontrivial (meaning in particular: not small) as $h \rightarrow 0$. To obtain control at tf, one needs to invert the normal operator $N(P)$, defined in [Hintz 2022, §3.1.2] (denoted by $N_{tf}(P)$ there) and recalled below. The following result, already implicit in the definition of the normal operator in [Hintz 2022, §3.1.2], lays the groundwork for the analysis of $N(P)$.

Lemma 3.5 (restriction to tf: vector fields). *The restriction map*

$$\mathcal{V}_b(X_{ch}) \ni V \mapsto V|_{tf} \in \mathcal{V}_b(tf)$$

restricts to a surjective map

$$N : \mathcal{V}_{ch}(X_{ch}) \rightarrow \mathcal{V}_{b,sc}(tf) \tag{3-6}$$

onto the space $\mathcal{V}_{b,sc}(tf) = (h/(h+x))\mathcal{V}_b(tf)$ of vector fields which are b-vector fields near $tf \cap cf$ and scattering vector fields near $tf \cap sf$. The map (3-6) induces bundle isomorphisms

$${}^{ch}T_{tf}^*X_{ch} \cong {}^{b,sc}T_{tf}, \quad {}^{ch}T_{tf}^*X_{ch} \cong {}^{b,sc}T^*tf. \tag{3-7}$$

Proof. Near $tf \setminus sf$, we write $V \in \mathcal{V}_{ch}(X_{ch})$ in the coordinates (h, \hat{x}, y) , with $\hat{x} = x/h$, as

$$V = a(h, \hat{x}, y)\hat{x}\partial_{\hat{x}} + \sum_{j=1}^{n-1} b^j(h, \hat{x}, y)\partial_{y^j}; \tag{3-8}$$

see (3-1). The restriction to tf, in local coordinates given by $h^{-1}(0) = [0, \infty)_{\hat{x}} \times \mathbb{R}_y^{n-1}$, is the b-vector field

$$V|_{tf} = a(0, \hat{x}, y)\hat{x}\partial_{\hat{x}} + \sum_{j=1}^{n-1} b^j(0, \hat{x}, y)\partial_{y^j}. \tag{3-9}$$

Conversely, every b-vector field W on $[0, \infty) \times \mathbb{R}^{n-1}$ can be written in the form $W = a(\hat{x}, y)\hat{x}\partial_{\hat{x}} + \sum_{j=1}^{n-1} b^j(\hat{x}, y)\partial_{y^j}$, and upon taking $a(h, \hat{x}, y)$ and $b^j(h, \hat{x}, y)$ to be smooth functions which restrict at $h = 0$ to the coefficients $a(\hat{x}, y)$ and $b^j(\hat{x}, y)$ of W defines a ch -vector field V through (3-8) whose

restriction (3-9) to tf is precisely W . We remark moreover that (3-9) vanishes if and only if $a = h\tilde{a}$ and $b^j = h\tilde{b}^j$, where \tilde{a}, \tilde{b}^j are smooth functions of (h, \hat{x}, y) , i.e., if and only if V vanishes at tf as a ch -vector field.

On $\text{tf} \setminus \text{cf}$ on the other hand, and using coordinates (\hat{h}, x, y) with $\hat{h} = h/x$, we can write $V \in \mathcal{V}_{ch}(X_{ch})$ as

$$V = \hat{h}a(\hat{h}, x, y)(x\partial_x - \hat{h}\partial_{\hat{h}}) + \sum_{j=1}^{n-1} \hat{h}b^j(\hat{h}, x, y)\partial_{y^j}, \tag{3-10}$$

with smooth coefficients a, b^1, \dots, b^{n-1} . Restriction to tf , which in these coordinates is given by $x = 0$, produces

$$V|_{\text{tf}} = -a(\hat{h}, 0, y)\hat{h}^2\partial_{\hat{h}} + \sum_{j=1}^{n-1} b^j(\hat{h}, 0, y)\hat{h}\partial_{y^j}, \tag{3-11}$$

which is a scattering vector field on $[0, \infty)_{\hat{h}} \times \mathbb{R}^{n-1}_y$, as claimed. Conversely, every scattering vector field W on $[0, \infty) \times \mathbb{R}^{n-1}$ can be written in the form $W = a(\hat{h}, y)\hat{h}^2\partial_{\hat{h}} + \sum_{j=1}^{n-1} b^j(\hat{h}, y)\hat{h}\partial_{y^j}$, and upon taking $a(\hat{h}, x, y)$ and $b^j(\hat{h}, x, y)$ to be smooth functions which restrict at $x = 0$ to the coefficients $-a(\hat{h}, y)$ and $b^j(\hat{h}, y)$ of W defines a ch -vector field V through (3-10) whose restriction (3-11) to tf is W . Note also that (3-11) vanishes if and only if $a = x\tilde{a}$ and $b^j = x\tilde{b}^j$ for smooth \tilde{a}, \tilde{b}^j , i.e., if and only if V vanishes at tf as a ch -vector field.

The surjectivity of (3-6) follows from these two calculations via a partition of unity subordinate to a cover $X_{ch} = U \cup V$, where $U \cap \text{sf} = \emptyset$ and $V \cap \text{cf} = \emptyset$. Our arguments above also prove that $\ker N = (x + h)\mathcal{V}_{ch}(X_{ch})$. Thus, we have an isomorphism

$$\mathcal{V}_{ch}(X_{ch}) / (x + h)\mathcal{V}_{ch}(X_{ch}) \rightarrow \mathcal{V}_{b,sc}(\text{tf}).$$

This induces the first isomorphism in (3-7) abstractly as follows: if $p \in \text{tf}$, then

$${}^{b,sc}T_p\text{tf} = \mathcal{V}_{b,sc}(\text{tf}) / \mathcal{I}_p\mathcal{V}_{b,sc}(\text{tf}),$$

where $\mathcal{I}_p \subset C^\infty(\text{tf})$ is the ideal of functions vanishing at p . The ideal in $C^\infty(X_{ch}) / (x + h)C^\infty(X_{ch})$ of elements that restrict to an element of \mathcal{I}_p at tf is $\mathcal{J}_p / (x + h)C^\infty(X_{ch})$, where $\mathcal{J}_p \subset C^\infty(X_{ch})$ is the ideal of functions vanishing at p . Since $\mathcal{V}_{ch}(X_{ch}) / \mathcal{J}_p\mathcal{V}_{ch}(X_{ch}) = {}^{ch}T_pX_{ch}$, we obtain (3-7). Concretely, the first isomorphism in (3-7) maps $\hat{x}\partial_{\hat{x}}, \partial_{y^j}$ in the coordinates used in (3-8) to $\hat{x}\partial_{\hat{x}}, \partial_{y^j}$ (cf. (3-9)), and $\hat{h}(x\partial_x - \hat{h}\partial_{\hat{h}}), \hat{h}\partial_{y^j}$ in the coordinates used in (3-10) to $-\hat{h}^2\partial_{\hat{h}}, \hat{h}\partial_{y^j}$ (cf. (3-11)). \square

The map (3-6) induces a surjective map

$$N : \left(\frac{x}{x+h}\right)^{-l} \left(\frac{h}{h+x}\right)^{-b} \text{Diff}_{ch}^k(X) \rightarrow \left(\frac{\hat{x}}{\hat{x}+1}\right)^{-l} (\hat{x}+1)^b \text{Diff}_{b,sc}^k(\text{tf}), \quad \hat{x} := \frac{x}{h}, \tag{3-12}$$

into weighted b -scattering differential operators on tf . More generally:

Lemma 3.6 (restriction to the transition face: ps.d.o.s). *Let $s, l, b \in \mathbb{R}$. Restriction to $\text{tf}_2 \subset X_{ch}^2$ induces a surjective map $N : \Psi_{ch,cl}^{s,l,0,b}(X) \rightarrow \Psi_{b,sc}^{s,l,b}(\text{tf})$.⁹ More generally, if $b \in C^\infty(\overline{{}^{ch}T_{\text{sf}}^*X_{ch}})$, then*

⁹Recall that the subscript “cl” refers to classicality at tf_2 , i.e., smoothness of the Schwartz kernels down to tf_2 .




Figure 7. Illustration of the map π defined in (3-14).

$b' := b|_{\text{tf} \cap \text{sf}} \in C^\infty(\overline{b, \text{sc} T^*_{\text{tf} \cap \text{sf}}(\text{tf})})$, and restriction to $\text{tf}_2 \subset X_{ch}^2$ induces a surjective map

$$N : \Psi_{ch, \text{cl}}^{s, l, 0, b}(X) \rightarrow \Psi_{b, \text{sc}}^{s, l, b'}(\text{tf}). \tag{3-13}$$

Proof. This can be proved entirely on the level of Schwartz kernels, since memberships in Ψ_{ch} or $\Psi_{b, \text{sc}}$ are characterized as conormal distributions with conormal regularity at various boundary hypersurfaces. The point then is that tf_2 is naturally diffeomorphic to the double space $\text{tf}_{b, \text{sc}}^2$ in the notation of (2-11), where we note that $\text{tf} \cong [0, \infty]_{\hat{x}} \times \partial X$ is indeed of the form (2-9). This is the route taken in [Hintz 2022, §3.1.2].

Alternatively, we can proceed explicitly for the symbolically nontrivial part using the quantization map (3-5), and use the Schwartz kernel perspective only to deduce the surjectivity of the restriction map for residual operators, $\Psi_{ch, \text{cl}}^{-\infty, l, 0, -\infty}(X) \rightarrow \Psi_{b, \text{sc}}^{-\infty, l, -\infty}(\text{tf})$. Indeed, on the level of symbols, note that with $\hat{x} = x/h$, we have

$$\begin{aligned} S^{s, l, 0, b}(\overline{ch T^* X_{ch}}) &= \left(\frac{x}{x+h}\right)^{-l} \left(\frac{h}{h+x}\right)^{-b} S^{s, 0, 0, 0}(\overline{ch T^* X_{ch}}) \\ &= \left(\frac{\hat{x}}{\hat{x}+1}\right)^{-l} (\hat{x}+1)^b S^{s, 0, 0, 0}(\overline{ch T^* X_{ch}}), \end{aligned}$$

and hence Lemma 3.5 implies that restriction to $\overline{ch T^*_{\text{tf}} X_{ch}}$ induces a surjective map

$$S_{\text{cl}}^{s, l, 0, b}(\overline{ch T^* X_{ch}}) \rightarrow S^{s, l, b}(\overline{b, \text{sc} T^* \text{tf}}).$$

But changing variables in the ch -quantization (3-5) to $\hat{x} = x/h$, $\hat{x}' = x'/h$ produces precisely the b -scattering quantization (2-10). This proves the lemma for constant orders; the proof in the variable-order case is the same. \square

As a consequence, we can relate semiclassical cone Sobolev spaces to b -scattering Sobolev spaces. In order to state this, we fix a collar neighborhood $\mathcal{U} = [0, x_0)_x \times \partial X$ of ∂X , and define the map

$$\pi : [0, 1)_h \times [0, \infty)_{\hat{x}} \times \partial X \rightarrow [0, 1)_h \times [0, \infty)_x \times \partial X, \quad \pi(h, \hat{x}, y) = (h, h\hat{x}, y). \tag{3-14}$$

Note that $(h\hat{x}, y) \in \mathcal{U}$ if and only if $\hat{x} < x_0 h^{-1}$. Now, since $\text{tf} = [0, \infty]_{\hat{x}} \times \partial X$, the domain of π is $[0, 1)_h \times (\text{tf} \setminus \text{sf})$; moreover, for any fixed $\hat{x} \in [0, \infty)$ and $y \in \partial X$, the point $\pi(h, \hat{x}, y)$ converges, as $h \searrow 0$, to the point (\hat{x}, y) on the transition face of X_{ch} . See Figure 7.

With this setup, we have:

Corollary 3.7 (restriction to tf : Sobolev spaces). *Suppose $\chi \in \mathcal{A}^0([0, 1)_h \times X)$ has compact support in $[0, 1) \times \mathcal{U}$. Let $0 < \mu_0 \in C^\infty(X; {}^b\Omega^1 X)$ and $0 < \hat{\mu}_0 \in C^\infty(\text{tf}; {}^b\Omega^1 X)$, let $\alpha_\mu \in \mathbb{R}$, and put*

$$\mu := x^{\alpha_\mu} \mu_0, \quad \hat{\mu} := \hat{x}^{\alpha_\mu} \hat{\mu}_0.$$

(1) (constant orders) *Let $s, l, \alpha, b \in \mathbb{R}$. Then*

$$\|\chi u\|_{H_{c,h}^{s,l,\alpha,b}(X;\mu)} \sim h^{\frac{\alpha_\mu}{2}-\alpha} \|\pi^*(\chi u)\|_{H_{b,sc}^{s,l,b-\alpha}(\text{tf};\hat{\mu})}, \quad u \in H_{c,h}^{s,l,\alpha,b}(X; \mu), \tag{3-15}$$

in the sense that the left-hand side is bounded by a uniform constant (independent of h and u) times the right-hand side and vice versa.

(2) (variable orders) *Let $b \in C^\infty(\overline{chT_{\text{sf}}^* X_{ch}})$ denote a variable order, and let $b' := b|_{\text{tf} \cap \text{sf}}$. If b is invariant under the lift of the dilation action $(x, y) \mapsto (\lambda x, y)$ in \mathcal{U} , then*

$$\|\chi u\|_{H_{c,h}^{s,l,\alpha,b}(X;\mu)} \sim h^{\frac{\alpha_\mu}{2}-\alpha} \|\pi^*(\chi u)\|_{H_{b,sc}^{s,l,b'-\alpha}(\text{tf};\hat{\mu})}.$$

For general b , and given $\delta > 0$, there exists $x_0(\delta) \in (0, x_0]$ so that for $\chi \in C_c^\infty([0, x_0(\delta)) \times \partial X)$, we have

$$C^{-1} h^{\frac{\alpha_\mu}{2}-\alpha} \|\pi^*(\chi u)\|_{H_{b,sc}^{s,l,b'-\alpha-\delta}(\text{tf};\hat{\mu})} \leq \|\chi u\|_{H_{c,h}^{s,l,\alpha,b}(X;\mu)} \leq C h^{\frac{\alpha_\mu}{2}-\alpha} \|\pi^*(\chi u)\|_{H_{b,sc}^{s,l,b'-\alpha+\delta}(\text{tf};\hat{\mu})}, \tag{3-16}$$

where C does not depend on h, u .

Proof. By factoring out $h^{-\alpha}$, it suffices to consider the case $\alpha = 0$. Consider first the case of constant orders. Factoring out the appropriate powers of $x/(x+h) = \hat{x}/(\hat{x}+1)$ and $h/(h+x) = (\hat{x}+1)^{-1}$, we reduce to the case $l = b = 0$. For $s = 0$, the equivalence of norms (3-15) then follows from

$$\iint |\chi u|^2 x^{\alpha_\mu} \frac{dx}{x} dy = \iint |\pi^*(\chi u)|^2 h^{\alpha_\mu} \hat{x}^{\alpha_\mu} \frac{d\hat{x}}{\hat{x}} dy.$$

For $s \in \mathbb{Z}$, the conclusion follows from (3-12); for general $s \in \mathbb{R}$, use duality and interpolation.

For variable semiclassical orders b (and still with $\alpha = 0$), and under the assumption of dilation-invariance near tf_2 , we first pick an elliptic operator $\hat{A} \in \Psi_{b,sc}^{s,l,b'}(\text{tf})$; we can then extend its Schwartz kernel to a neighborhood of tf_2 to be constant along the orbits of $(h, x) \mapsto (\lambda h, \lambda x)$, and then extend it further to an elliptic operator $A \in \Psi_{ch,cl}^{s,l,0,b}(X)$. In this manner, we obtain a right inverse (with special properties) of the restriction map (3-13). For any fixed $b_0 < \inf b$, we thus have

$$\begin{aligned} \|\chi u\|_{H_{c,h}^{s,l,0,b}(X;\mu)}^2 &\sim \|\chi u\|_{H_{c,h}^{s,l,0,b_0}(X;\mu)}^2 + \|A(\chi u)\|_{H_{c,h}^0(X;\mu)}^2 \\ &\sim h^{\frac{\alpha_\mu}{2}} (\|\pi^*(\chi u)\|_{H_{b,sc}^{s,l,b_0}(\text{tf};\hat{\mu})}^2 + \|\hat{A}(\pi^*(\chi u))\|_{H_{b,sc}^0(\text{tf};\hat{\mu})}^2) \\ &\sim h^{\frac{\alpha_\mu}{2}} \|\chi u\|_{H_{b,sc}^{s,l,b'}(\text{tf};\hat{\mu})}^2. \end{aligned}$$

The lossy estimate (3-16) is a consequence of this, as the dilation-invariant extension of $b' - \delta$, resp. $b' + \delta$ is less, resp. greater than b in a sufficiently small (depending on b and δ) neighborhood of ∂X . □

3D. Relative semiclassical b-regularity. We now make Remark 1.5 precise and demonstrate how to combine the notions of semiclassical cone regularity and semiclassical b- (i.e., conormal) regularity. Recall here that a semiclassical b-vector field is a particular type of h -dependent b-vector field on X ; namely, it is a vector field on $[0, 1)_h \times X$ which is horizontal and which vanishes at $h = 0$. In local coordinates as in (2-1), such a vector fields can be written as

$$a(h, x, y)hx\partial_x + \sum_{j=1}^{n-1} b^j(h, x, y)h\partial_{y^j}. \tag{3-17}$$

The main insight is that the semiclassical b-algebra can be embedded into the semiclassical cone algebra via a phase space resolution; see Lemma 3.8 below. This can alternatively be phrased as a second microlocalization of the semiclassical b-algebra at the zero section over ∂X at $h = 0$; see Remark 3.10.

First, we explain a slightly nonstandard perspective on semiclassical (b-)phase spaces. Let X be an n -dimensional manifold with nonempty embedded boundary ∂X . Thus, paralleling Definition 3.1, we define

$$X_{b\hbar} := [0, 1) \times X, \\ \mathcal{V}_{b\hbar}(X_{b\hbar}) := \{V \in \mathcal{V}_b(X_{b\hbar}) : V \text{ is horizontal, } V|_{h=0} = 0\}.$$

It is then easy to see that $\mathcal{V}_{b\hbar}(X_{b\hbar})$ is spanned over $\mathcal{C}^\infty(X_{b\hbar})$ by hV for $V \in \mathcal{V}_b(X)$ (see (3-17)), where we identify V with an h -independent horizontal vector field on $X_{b\hbar}$. We then have $\mathcal{V}_{b\hbar}(X_{b\hbar}) = \mathcal{C}^\infty(X_{b\hbar}; {}^{b\hbar}TX_{b\hbar})$ for a rank- n vector bundle

$${}^{b\hbar}TX_{b\hbar} \rightarrow X_{b\hbar}.$$

In local coordinates $[0, \infty)_x \times \mathbb{R}_y^{n-1}$, a smooth frame of this bundle is $hx\partial_x, h\partial_{y^1}, \dots, h\partial_{y^{n-1}}$. We can introduce fiber-linear coordinates on the dual bundle ${}^{b\hbar}T^*X_{b\hbar}$ by writing the canonical 1-form as

$$\xi_{b\hbar}h^{-1}\frac{dx}{x} + \sum_{j=1}^{n-1} (\eta_{b\hbar})_j h^{-1}dy^j.$$

Thus, for example, the symbol of the semiclassical b-differential operator $hx D_x$ is $\xi_{b\hbar}$.¹⁰ Denote fiber infinity of the radial compactification $\overline{{}^{b\hbar}T^*X_{b\hbar}}$ by ${}^{b\hbar}S^*X_{b\hbar}$. Given a symbol

$$a \in S^{s,l,b}(\overline{{}^{b\hbar}T^*X_{b\hbar}}) = x^{-l}h^{-b}S^{s,0,0}(\overline{{}^{b\hbar}T^*X_{b\hbar}})$$

(i.e., a is conormal with weight $-s$ at ${}^{b\hbar}S^*X_{b\hbar}$, conormal with weight $-l$ at $\overline{{}^{b\hbar}T^*_{[0,1) \times X}X_{b\hbar}}$, and conormal with weight $-b$ at $h = 0$), we can then define the semiclassical quantization

$$(\text{Op}_{b,h}(a)u)(x, y) := (2\pi)^{-n} \iiint \exp\left(i\left[\frac{x-x'}{hx}\xi_{b\hbar} + \frac{y-y'}{h} \cdot \eta_{b\hbar}\right]\right) \phi\left(\left|\log \frac{x}{x'}\right|\right) \phi(|y-y'|) \\ a(x, y, \xi_{b\hbar}, \eta_{b\hbar})u(x', y') \frac{dx'}{hx'} \frac{dy'}{h^{n-1}} d\xi_{b\hbar} d\eta_{b\hbar}.$$

¹⁰By contrast, the standard convention is to introduce fiber-linear coordinates (ξ_b, η_b) on ${}^bT^*X$ as in (2-3) and declare the principal symbol of $hx D_x$ to be ξ_b ; the translation to the present convention is accomplished by using (the adjoint of) the bundle isomorphism ${}^{b\hbar}TX_{b\hbar} \cong [0, 1)_h \times {}^bTX$ induced by division by h (i.e., induced by the map $\mathcal{V}_{b\hbar}(X_{b\hbar}) \ni V \mapsto (h^{-1}V)_{h \in [0,1)}$).




Figure 8. Left: The resolution $\overline{cb\hbar T^* X_{ch}}$ of the fiber-compactified semiclassical cone phase space at fiber infinity over tf ; see (3-19). (Unlike in Figure 4, we show the full compactified fibers here.) Right: The resolution of the fiber-compactified semiclassical b -phase space at $x = h = 0$ and at the zero section over $x = h = 0$; see (3-20).

If we make the change of variables

$$(\xi_{bh}, \eta_{bh}) = (x + h)(\xi_{ch}, \eta_{ch}), \tag{3-18}$$

see (3-2), this exactly matches the $c\hbar$ -quantization (3-5). The key point is now that this match has a clean interpretation on the level of symbol classes on a joint resolution of the semiclassical cone and b -phase spaces:

Lemma 3.8 (relationship between semiclassical cone and b -phase spaces). *Define the $cb\hbar$ -phase space*

$$\overline{cb\hbar T^* X_{ch}} := [\overline{c\hbar T^* X_{ch}}; {}^{ch}S_{tf}^* X_{ch}]. \tag{3-19}$$

*Denote by $\mathcal{C} := \overline{b\hbar T^*_{\{0\} \times \partial X} X_{bh}}$ the semiclassical b -phase space over the corner $h = x = 0$, and denote by $o_{\mathcal{C}} \subset \mathcal{C}$ the zero section. Then the identity map on $(0, 1)_h \times T^* X^o$ extends by continuity to a diffeomorphism*

$$\overline{cb\hbar T^* X_{ch}} \xrightarrow{\cong} [\overline{b\hbar T^* X_{bh}}; \mathcal{C}; o_{\mathcal{C}}]. \tag{3-20}$$

We refer to the front face of (3-19) as fbf (“finite b -frequencies”). See Figure 8.

Proof of Lemma 3.8. We work in polar coordinates $\rho = x + h, \theta = (x, h)/|(x, h)|$ in the (x, h) -variables. Thus, local coordinates near ${}^{ch}T_{tf}^* X_{ch}$ are $(\rho, y, \theta, \zeta_{ch})$, $\zeta_{ch} := (\xi_{ch}, \eta_{ch})$, while local coordinates near the front face of $[\overline{b\hbar T^* X_{bh}}; \mathcal{C}]$, away from fiber infinity, are $(\rho, y, \theta, \zeta_{bh})$, $\zeta_{bh} = (\xi_{bh}, \eta_{bh})$. Coordinates near the interior of the front face of the final blow-up in (3-20) are then $(\rho, y, \theta, \zeta_{bh}/\rho) = (\rho, y, \theta, \zeta_{ch})$; see (3-18). Near the intersection of the lift of $o_{\mathcal{C}}$ with that of \mathcal{C} , smooth coordinates can be constructed by introducing polar coordinates in the fiber variables, giving $(\rho/|\zeta_{bh}|, y, \theta, |\zeta_{bh}|, \zeta_{bh}/|\zeta_{bh}|)$; this matches, up to a permutation, the local coordinates on $\overline{cb\hbar T^* X_{ch}}$ near the lift of ${}^{ch}T_{tf}^* X_{ch}$ given by $(\rho/|\zeta_{ch}|^{-1}, y, \theta, |\zeta_{ch}|^{-1}, \zeta_{ch}/|\zeta_{ch}|)$. Lastly, near the lift of fiber infinity on the resolved b -phase space, we can use coordinates $(\rho, y, \theta, |\zeta_{bh}|^{-1}, \zeta_{bh}/|\zeta_{bh}|)$, which matches the local coordinates near the lift of ${}^{ch}S^* X_{ch}$ given by $(\rho, y, \theta, |\zeta_{ch}|^{-1}/\rho, \zeta_{ch}/|\zeta_{ch}|)$. \square




Figure 9. Illustration of the orders of (symbols of) $cb\hbar$ -pseudodifferential operators in (3-21) and Definition 3.9.

The blow-up of a boundary face does not enlarge the space of conormal distributions, but allows for more precise accounting of weights. Concretely, define for $s, s', l, \alpha, b \in \mathbb{R}$ the symbol space

$$S^{s,s',l,\alpha,b}(\overline{cb\hbar T^* X_{ch}}), \tag{3-21}$$

where the orders refer, in this order, to fiber infinity, the front face fbf of (3-19), and the phase space over the lifts of cf , tf and sf , see Figure 9. Then we have

$$\begin{aligned} S^{s,l,\alpha,b}(\overline{ch T^* X_{ch}}) &= S^{s,s+\alpha,l,\alpha,b}(\overline{cb\hbar T^* X_{ch}}), \\ S^{s,s',l,\alpha,b}(\overline{cb\hbar T^* X_{ch}}) &\subset S^{\max(s,s'-\alpha),l,\alpha,b}(\overline{ch T^* X_{ch}}). \end{aligned} \tag{3-22}$$

Note that the second inclusion is *false* if we use spaces of *classical* symbols on both sides; after all, blow-ups do enlarge the space of smooth functions (but preserve the space of conormal functions). Since we worked with general conormal symbols and ps.d.o.s in Section 3B, we can immediately quantize symbols on the $cb\hbar$ -phase space:

Definition 3.9 ($cb\hbar$ -pseudodifferential operators). Let $s, s', l, \alpha, b \in \mathbb{R}$. Then we define

$$\Psi_{cb\hbar}^{s,s',l,\alpha,b}(X) := \text{Op}_{ch}(S^{s,s',l,\alpha,b}(\overline{cb\hbar T^* X_{ch}})) + \Psi_{ch}^{-\infty,l,\alpha,-\infty}(X).$$

Operators with variable semiclassical orders $b \in \overline{ch T_{sf}^* X_{ch}}$ are defined similarly.

Remark 3.10 (second microlocalization). In view of Lemma 3.8, one can view $\Psi_{cb\hbar}(X)$ as a second microlocalization of the (conormal) semiclassical b -algebra $\Psi_{b\hbar}(X)$ at the zero section over $h = x = 0$. In terms of symbol classes, we have

$$\begin{aligned} S^{s,l,b}(\overline{b\hbar T^* X_{b\hbar}}) &= S^{s,l+b,l,l+b,b}(\overline{cb\hbar T^* X_{ch}}), \\ S^{s,s',l,\alpha,b}(\overline{cb\hbar T^* X_{ch}}) &\subset S^{s,l,\max(b,s'-l,\alpha-l)}(\overline{b\hbar T^* X_{b\hbar}}), \end{aligned} \tag{3-23}$$

and analogous statements hold for ps.d.o.s. However, similarly to [Vasy 2021b, §5] in the context of b - and scattering algebras, it is analytically advantageous to resolve $\Psi_{ch}(X)$ as in Definition 3.9, as the second microlocal/resolved algebra involves *global* (noncommutative) phenomena at $h = x = 0$ (i.e.,

the lift of tf , associated to which is the normal operator homomorphism into a noncommutative algebra) which are directly inherited from $\Psi_{c\hbar}(X)$, but which are not visible on the level of $\Psi_{b\hbar}(X)$.

For two ps.d.o.s $A_j \in \Psi_{cb\hbar}^{s_j, s'_j, l_j, \alpha_j, b_j}(X)$, one can compute the full symbol, i.e., the symbol modulo

$$S^{-\infty, l_1+l_2, \alpha_1+\alpha_2, -\infty}(\overline{c\hbar T^* X_{c\hbar}}) = S^{-\infty, -\infty, l_1+l_2, \alpha_1+\alpha_2, -\infty}(\overline{cb\hbar T^* X_{c\hbar}}),$$

of the composition $A_1 \circ A_2 \in \Psi_{cb\hbar}^{\max(s_1, s'_1 - \alpha_1) + \max(s_2, s'_2 - \alpha_2), l_1+l_2, \alpha_1+\alpha_2, b_1+b_2}(X)$ in local coordinates using the usual symbol expansion to be the sum of products of derivatives of the full symbols of the two factors along b-vector fields on $\overline{c\hbar T^* X_{c\hbar}}$ which vanish, as b-vector fields, at $c\hbar S^* X_{c\hbar}$ (thus vanishing as b-vector fields at the lift of $c\hbar S^* X_{c\hbar}$ as well as at the front face of (3-19)) and at the lift of $\overline{c\hbar T_{\text{sf}}^* X_{c\hbar}}$. Plugging the $cb\hbar$ -symbols of A_1, A_2 into such an expansion thus shows that, in fact,

$$A_1 \circ A_2 \in \Psi_{cb\hbar}^{s_1+s_2, s'_1+s'_2, l_1+l_2, \alpha_1+\alpha_2, b_1+b_2}(X).$$

Similar arguments show that the principal symbol map

$$cb\hbar \sigma : \Psi_{cb\hbar}^{s, s', l, \alpha, b}(X) \rightarrow (S^{s, s', l, \alpha, b} / S^{s-1, s'-1, l, \alpha, b-1})(\overline{cb\hbar T^* X_{c\hbar}})$$

is well-defined (and a *-homomorphism as usual). One can moreover define an associated scale of Sobolev spaces

$$H_{cb, h}^{s, s', l, \alpha, b}(X) = \{u \in H_{c, h}^{\min(s, s' - \alpha), l, \alpha, b}(X) : Au \in L^2(X) \text{ for all } A \in \Psi_{cb\hbar}^{s, s', l, \alpha, b}(X)\}. \tag{3-24}$$

The relationships (3-22) and (3-23) imply:

Proposition 3.11 (relationships between Sobolev spaces). *Let $s, s', l, \alpha, b \in \mathbb{R}$. Define L^2 using the volume density $\mu = x^{\alpha_\mu} \mu_0$, $0 < \mu_0 \in C^\infty(X; {}^b\Omega^1 X)$, with $\alpha_\mu \in \mathbb{R}$. Then*

$$\begin{aligned} H_{c, h}^{s, l, \alpha, b}(X) &= H_{cb, h}^{s, s+\alpha, l, \alpha, b}(X), \\ H_{b, h}^{s, l, b}(X) &= H_{cb, h}^{s, l+b, l, l+b, b}(X). \end{aligned}$$

One can conversely embed $H_{cb, h}^{s, s', l, \alpha, b}(X)$ into $H_{c, h}^{\tilde{s}, \tilde{l}, \tilde{\alpha}, \tilde{b}}(X)$ and $H_{b, h}^{\tilde{s}, \tilde{l}, \tilde{b}}(X)$ under suitable inequalities (which can be read off from Proposition 3.11) between the orders. In particular, this allows us to give a direct proof of [Hintz 2022, Proposition 3.18] on the relationship between $H_{c, h}(X)$ and $H_{b, h}(X)$; for instance, for $s, l, \alpha \in \mathbb{R}$ (denoted by s, α, τ in the reference), we have

$$H_{c, h}^{s, l, \alpha, 0}(X) = H_{cb, h}^{s, s+\alpha, l, \alpha, 0}(X) \subset H_{b, h}^{s, l, \min(0, \alpha-l, \alpha-l+s)}(X), \tag{3-25}$$

which implies (and is slightly sharper than) the first part of [Hintz 2022, Proposition 3.18]. If one wishes to translate estimates on cone spaces to b-spaces, the advantage of the resolved $cb\hbar$ -Sobolev spaces, compared with $c\hbar$ -Sobolev spaces, is that one can reduce losses in powers of h (or in regularity) in the conversion; as a simple concrete example, we have

$$H_{cb, h}^{s, s'+\alpha, l, \alpha, 0}(X) \subset H_{b, h}^{s, l, \min(0, \alpha-l, \alpha-l+s')}(X),$$

which for $s' \geq -s_-$ gives an improved bound at $h = 0$, and for $s' \geq 0$ a bound which is independent of the differential orders s, s' , unlike (3-25), which gets lossier as s decreases.

Remark 3.12 (variable semiclassical orders). The above discussion applies, mutatis mutandis, to symbols and operators with variable semiclassical orders b as well; here b is a smooth function on the lift of $\overline{c\hbar T_{sf}^* X_{c\hbar}}$ to $\overline{cb\hbar T^* X_{c\hbar}}$.

4. Microlocal propagation estimates at cone points and generalizations

Let $n \geq 1$. We work locally near a cone point; thus on an n -dimensional manifold

$$X = [0, 2x_0)_x \times Y, \quad x_0 > 0, \tag{4-1}$$

where Y is a closed connected $(n-1)$ -dimensional manifold, and where $X^\circ = (0, 2x_0) \times Y$ is equipped with a smooth Riemannian metric g of the form

$$g = dx^2 + x^2 k(x, y, dy), \tag{4-2}$$

where $k \in C^\infty([0, x_0]; C^\infty(Y, S^2 T^* Y))$ is a smooth family of smooth Riemannian metrics on the cross section Y . Any metric which locally near ∂X is of the form $d\tilde{x}^2 + \tilde{x}^2 k(\tilde{x}, y, d\tilde{x}, dy)$, with $k|_{\partial X}$ a Riemannian metric on ∂X , is of the form (4-2) in a suitable smooth collar neighborhood of ∂X , as shown in [Melrose and Wunsch 2004, §1].

While the above X is not compact, all calculations and estimates will take place in the compact subset $[0, x_0] \times Y$ of X ; thus, we shall commit a slight abuse of notation and write $\|u\|_{H_{c,h}^s(X)}$ etc. for norms of functions u on X which will always have support in $x^{-1}([0, x_0])$. We fix the volume density

$$\mu = |dg| = x^{n-1} |dx dk| \in x^n C^\infty(X; {}^b\Omega^1 X) \tag{4-3}$$

on X , and define Sobolev spaces relative to $L^2(X) := L^2(X; \mu)$. We moreover define

$$\begin{aligned} \hat{x} &:= \frac{x}{h}, & \hat{g} &:= d\hat{x}^2 + \hat{x}^2 k(0, y, dy), \\ \hat{h} &:= \hat{x}^{-1} = \frac{h}{x}, & \hat{\mu} &:= |d\hat{g}| = \hat{x}^{n-1} |d\hat{x} dk(0)|. \end{aligned} \tag{4-4}$$

4A. Admissible operators. The class of operators of interest to us is the following.

Definition 4.1 (admissible operators). We call an h -dependent differential operator $P_{h,z}$ on X° *admissible* if it is of the form

$$P_{h,z} = h^2 \Delta_g - z + h^2 x^{-2} Q_{1,z} + hx^{-1} q_{0,z}, \tag{4-5}$$

where $Q_{1,z} \in \text{Diff}_b^1(X)$ and $q_{0,z} \in C^\infty(X)$ depend smoothly on $z \in \mathbb{C}$, $|z-1| < Ch$.

We shall henceforth take $z = z(h)$ to be a smooth function of $h \in [0, 1)$ with $z(0) = 1$.

Remark 4.2 (vector bundles). Our analysis applies also to operators acting on sections of a vector bundle $E \rightarrow X$; we explain the necessary (largely notational) changes in Remark 4.11.

Using local coordinates $y \in \mathbb{R}^{n-1}$ on ∂X , let us write

$$Q_{1,z} = q_{1,z}(x, y, xD_x, D_y), \quad q_{0,z} = q_{0,z}(x, y).$$

The normal operator of $P_{h,z}$ is

$$\begin{aligned} N(P) &:= \Delta_{\hat{g}} - 1 + \hat{x}^{-2}q_{1,1}(0, y, \hat{x}D_{\hat{x}}, D_y) + \hat{x}^{-1}q_{0,1}(0, y) \\ &= D_{\hat{x}}^2 - i(n-1)D_{\hat{x}} + \hat{x}^{-2}\Delta_{k(0)} - 1 + \hat{x}^{-2}q_{1,1}(0, y, \hat{x}D_{\hat{x}}, D_y) + \hat{x}^{-1}q_{0,1}(0, y) \end{aligned} \tag{4-6}$$

on $\text{tf} = [0, \infty]_{\hat{x}} \times \partial X$.¹¹

Lemma 4.3 (structural properties). *We have*

$$P_{h,z} \in \left(\frac{x}{x+h}\right)^{-2} \text{Diff}_{ch}^2(X) \quad \text{and} \quad N(P) \in \left(\frac{\hat{x}}{\hat{x}+1}\right)^{-2} \text{Diff}_{b,sc}^2(\text{tf}).$$

Furthermore, we have

$$P_{h,z} - N(P) \in (x+h) \left(\frac{x}{x+h}\right)^{-2} \text{Diff}_{ch}^2(X),$$

where we abuse notation and write $N(P) \in (x/(x+h))^{-2} \text{Diff}_{ch}^2(X)$ for any operator whose normal operator is equal to $N(P)$.

Proof. In local coordinates y^1, \dots, y^{n-1} on Y , the metric $k(x, y, dy)$ is given by an $(n-1) \times (n-1)$ matrix (k_{ij}) with determinant $|k| > 0$ and inverse (k^{ij}) , and we have

$$\begin{aligned} \Delta_g &= |k|^{-\frac{1}{2}}x^{-n+1}D_x(|k|^{\frac{1}{2}}x^{n-1}D_x) + x^{-2}\Delta_{k(x)} \\ &= D_x^2 - i(n-1+x\gamma)x^{-1}D_x + \sum_{i,j=1}^{n-1} x^{-2}|k|^{-\frac{1}{2}}D_{y^i}(|k|^{\frac{1}{2}}k^{ij}D_{y^j}), \end{aligned}$$

where $\gamma = \frac{1}{2}\partial_x \log |k| \in C^\infty$. Since

$$\begin{aligned} hD_x &= \frac{x+h}{x} \cdot \frac{h}{h+x}x D_x \in \frac{x+h}{x} \mathcal{V}_{ch}(X_{ch}), \\ hx^{-1}D_{y^i} &= \frac{x+h}{x} \cdot \frac{h}{h+x}D_{y^i} \in \frac{x+h}{x} \mathcal{V}_{ch}(X_{ch}), \\ hx^{-1} &= \frac{x+h}{x} \cdot \frac{h}{h+x} \in \frac{x+h}{x} \text{Diff}_{ch}^1(X), \end{aligned} \tag{4-7}$$

we find $h^2\Delta_g \in (x/(x+h))^{-2} \text{Diff}_{ch}^2(X)$, and its normal operator is $D_{\hat{x}}^2 - i(n-1)\hat{x}^{-1}D_{\hat{x}} + \hat{x}^{-2}\Delta_{k(0)} = \Delta_{\hat{g}}$. The remaining terms in (4-5) are analyzed similarly. \square

4B. Characteristic set, Hamilton flow. Using the fiber-linear coordinates (ξ_{ch}, η_{ch}) on ${}^{ch}T^*X_{ch}$ from (3-2), we can read off the principal symbol from (4-7) to be

$$p := \left(\frac{x}{x+h}\right)^2 \cdot {}^{ch}\sigma(P_{h,z}) = \xi_{ch}^2 + |\eta_{ch}|_{k^{-1}}^2 - 1.$$

(Here, we use that $z = 1 + \mathcal{O}(h)$, hence the principal symbol of z is 1.) This is elliptic at fiber infinity ${}^{ch}S^*X_{ch}$, but has a nonempty characteristic set at finite frequencies. Near sf , it is more convenient to use

¹¹This can be defined more invariantly as an operator on the inward pointing normal bundle ${}^+N\partial X$, which is the natural place for the b-normal operators $q_{1,1}(0, y, xD_x, D_y)$ and $q_{0,1}(0, y)$ to live; see [Melrose 1993, §4.15; Hintz 2022, §3] for details.




Figure 10. Illustration of the flow along the rescaled Hamilton vector field H , see (4-9), through the radial sets \mathcal{R}_{in} and \mathcal{R}_{out} . Shown is the characteristic set, the fibers of which over sf are spheres (here 1-spheres); one fiber is drawn as a dotted circle. Also indicated is (in gray) the linearization of H at $\mathcal{R}_{in/out}$ over tf .

the fiber coordinates (ξ, η) from (3-3), and $\hat{h} = h/x$, x, y as coordinates on the base, so that

$$p = \xi^2 + k^{ij}(x, y)\eta_i\eta_j - 1, \tag{4-8}$$

$$\Sigma = p^{-1}(0) \cap {}^{ch}T_{sf}^*X_{ch} = \{\hat{h} = 0, \xi^2 + |\eta|_{k^{-1}}^2 = 1\}.$$

Using (3-4) and writing $|\eta|^2 = k^{ij}\eta_i\eta_j$, we then compute

$$H := \hat{h}^{-1}H_p = 2\xi(x\partial_x - \hat{h}\partial_{\hat{h}} - \eta\partial_{\eta}) + (2|\eta|^2 - x\partial_x k^{ij}\eta_i\eta_j)\partial_{\xi} + 2k^{ij}\eta_i\partial_{y^j} - (\partial_{y^k}k^{ij})\eta_i\eta_j\partial_{\eta_k}. \tag{4-9}$$

Restricted to $x = 0$ as a b-vector field on $\overline{{}^{ch}T^*X_{ch}}$, this is

$$H|_{x=0} = 2\xi(x\partial_x - \hat{h}\partial_{\hat{h}} - \eta\partial_{\eta}) + 2|\eta|^2\partial_{\xi} + (2k^{ij}\eta_i\partial_{y^j} - (\partial_{y^k}k^{ij})\eta_i\eta_j\partial_{\eta_k}). \tag{4-10}$$

This vanishes as a standard vector field on $\hat{h} = x = 0$ if and only if $\eta = 0$. The intersection of $\eta^{-1}(0)$ with $\Sigma \cap x^{-1}(0)$ has two components: the *incoming* and *outgoing radial sets* $\mathcal{R}_{in/out} \subset {}^{ch}T_{sf}^*X_{ch}$,

$$\begin{aligned} \mathcal{R}_{in} &:= \{(\hat{h}, x, y, \xi, \eta) : \hat{h} = 0, x = 0, y \in \partial X, \xi = -1, \eta = 0\}, \\ \mathcal{R}_{out} &:= \{(\hat{h}, x, y, \xi, \eta) : \hat{h} = 0, x = 0, y \in \partial X, \xi = +1, \eta = 0\}. \end{aligned} \tag{4-11}$$

These are saddle points for the rescaled Hamilton vector field H since

$$x^{-1}Hx = \mp 2, \quad \hat{h}^{-1}H\hat{h} = \pm 2, \quad |\eta|^{-2}H|\eta|^2 = \pm 4 \quad \text{at } \mathcal{R}_{in/out}. \tag{4-12}$$

(The top sign is for “in”, the bottom sign for “out”.) See Figure 10.

Over ${}^{ch}T_{tf}^*X_{ch}$, the set \mathcal{R}_{in} is a radial *source* (though this really only makes sense infinitesimally at $tf \cap sf$ since the ch -calculus is not symbolic over $tf \setminus sf$), and \mathcal{R}_{out} is a radial *sink*. This matches precisely the familiar situation of scattering theory on the asymptotically conic space (tf, \hat{g}) , see [Melrose 1994], which we discuss in detail in Section 4C.

In $x > 0$, the flow of H is a reparametrization of the flow of $h^{-1}H_p = x^{-1}H$. Integral curves of H starting over a point in X° never reach ∂X in finite time. Instead, we consider

$$\begin{aligned} H_{sf} &:= h^{-1}H_p|_{\hat{h}=0} \\ &= 2\xi(\partial_x - x^{-1}\eta\partial_{\eta}) + (2x^{-1}|\eta|^2 - \partial_x k^{ij}\eta_i\eta_j)\partial_{\xi} + x^{-1}(2k^{ij}\eta_i\partial_{y^j} - (\partial_{y^k}k^{ij})\eta_i\eta_j\partial_{\eta_k}). \end{aligned} \tag{4-13}$$

Given $y_0 \in \partial X$, the curves

$$\begin{aligned} \gamma_{I,y_0}(s) &:= (-2s, y_0, -1, 0), & s \in (-x_0, 0), \\ \gamma_{O,y_0}(s) &:= (2s, y_0, 1, 0), & s \in (0, x_0), \end{aligned} \tag{4-14}$$

are integral curves of H_{sf} . Here, γ_{I,y_0} strikes ∂X at $s = 0$ at the incoming radial set over point y_0 , whereas γ_{O,y_0} emanates from the outgoing radial set over y_0 at $s = 0$.

Lemma 4.4 (incoming/outgoing null-bicharacteristics). *Let $0 < s_0 < x_0$. Suppose $\gamma : (0, s_0) \rightarrow \Sigma \cap {}^{\text{ch}}T_{\text{sf}\setminus\text{tf}}^* X_{\text{ch}}$ is an integral curve of H_{sf} tending to ∂X as $s \searrow 0$ in the weak sense that $\liminf_{s \searrow 0} x(\gamma(s)) = 0$. Then in the coordinates (x, y, ξ, η) , γ is necessarily of the form $\gamma(s) = \gamma_{O,y_0}(s)$ for some $y_0 \in \partial X$. Similarly, if $\gamma : (-s_0, 0) \rightarrow \Sigma \cap {}^{\text{ch}}T_{\text{sf}\setminus\text{tf}}^* X_{\text{ch}}$ is an integral curve of H_{sf} with $\liminf_{s \nearrow 0} x(\gamma(s)) = 0$, then $\gamma(s) = \gamma_{I,y_0}(s)$ for some $y_0 \in \partial X$.*

Proof. The vector field

$$xH_{\text{sf}} = H|_{x=0} = 2\xi(x\partial_x - \eta\partial_\eta) + (2|\eta|^2 - x\partial_x k^{ij}\eta_i\eta_j)\partial_\xi + (2k^{ij}\eta_i\partial_{y^j} - (\partial_{y^k} k^{ij})\eta_i\eta_j\partial_{\eta_k})$$

vanishes identically at \mathcal{R}_{out} . We study the behavior of xH_{sf} as a vector field on Σ near \mathcal{R}_{out} ; we may use the coordinates $x \geq 0$, $y \in \mathbb{R}^{n-1}$, and $\eta \in \mathbb{R}^{n-1}$, in which ξ is determined by $p = 0$ as the positive square root of $1 - k^{ij}(x, y)\eta_i\eta_j$. The linearization of $(xH_{\text{sf}})|_\Sigma$ in the normal directions at \mathcal{R}_{out} , defined by mapping df to $d(xH_{\text{sf}}f)$ where $f \in C^\infty(\Sigma)$, $f|_{\mathcal{R}_{\text{out}}} = 0$, maps

$$dx \mapsto 2 dx, \quad d\eta \mapsto -2 d\eta,$$

and is thus hyperbolic; the unstable and stable subbundles of $T_{\mathcal{R}_{\text{out}}}\Sigma$ for the $(xH_{\text{sf}})|_\Sigma$ -flow are correspondingly the span of ∂_x and ∂_η . The unstable manifold theorem, in the form given in [Hirsch et al. 1977, Theorem 4.1], thus applies inside Σ at \mathcal{R}_{out} and produces an unstable manifold whose tangent space at a point $\zeta \in \mathcal{R}_{\text{out}}$ is the sum of $T_\zeta(\mathcal{R}_{\text{out}})$ and $\mathbb{R}\partial_x$. (See the proof of [Melrose and Wunsch 2004, Theorem 1.2] for similar, albeit more general, considerations.) Since the manifold $\mathcal{R}_{\text{out}} \cup \{\gamma_{O,y_0}(s) : y_0 \in \partial X, s \in (0, s_0)\}$ is H -invariant with the same tangent space, it must be equal to this unstable manifold. The first part of the lemma follows from this observation; the second part is completely analogous. \square

Definition 4.5 (generalized broken bicharacteristics). Denote by $\dot{\Sigma}$ the topological space defined as the quotient $\Sigma/\partial\Sigma$. Let $I \subset \mathbb{R}$ denote an open interval. We then say that a continuous curve $\gamma : I \rightarrow \dot{\Sigma}$ is a *generalized broken bicharacteristic* (GBB) if either $\gamma(I) \subset \Sigma \setminus \partial\Sigma$ and γ is an integral curve of H_{sf} , or there exist $s_0 \in I$ and $y_I, y_O \in \partial X$ so that $\gamma(s_0 + t) = \gamma_{O,y_O}(t)$ for $t > 0$, $s_0 + t \in I$ and $\gamma(s_0 + t) = \gamma_{I,y_I}(t)$ for $t < 0$, $s_0 + t \in I$.¹² If y_O is at distance π from y_I with respect to the metric $k(0)$ on ∂X , we say that γ is a *geometric GBB*; otherwise γ is a *strictly diffractive GBB*.

See Figure 11. We remark without proof that geometric GBB are uniform limits of H_{sf} -integral curves just barely missing ∂X (see also [Melrose and Wunsch 2004, Lemma 1.5]).

¹²In light of Lemma 4.4, this is equivalent to the condition that γ is an H_{sf} -integral curve outside of $\partial\Sigma$, but may enter and exit the characteristic set over ∂X at different points.




Figure 11. The projection of strictly diffractive (blue) and geometric (red) GBBs to the base X , as well as geodesics (green) just barely missing the cone tip ∂X . Left: the geometric picture, where ∂X is collapsed to a point. Right: the resolved picture.

4C. Scattering theory for the normal operator. Propagation through the “cone point” ∂X will require *global* control of the normal operator, namely the absence of purely outgoing or purely incoming solutions (depending on the direction in which one wants to propagate estimates). Let us define fiber-linear coordinates on the scattering cotangent bundle ${}^{\text{sc}}T^*(\text{tf} \setminus \text{cf})$ via

$$\xi_{\text{sc}} \frac{d\hat{h}}{\hat{h}^2} + \sum_{j=1}^{n-1} (\eta_{\text{sc}})_j \frac{dy^j}{\hat{h}}.$$

Via the identification (3-7), the radial sets $\mathcal{R}_{\text{in/out}}$ defined in (4-11) are then equal to the sets ${}^{\text{sc}}\mathcal{R}_{\text{in/out}} \subset {}^{\text{sc}}T^*(\text{tf} \setminus \text{cf})$, where

$$\begin{aligned} {}^{\text{sc}}\mathcal{R}_{\text{in}} &:= \{(\hat{h}, y, \xi_{\text{sc}}, \eta_{\text{sc}}) : \hat{h} = 0, y \in \partial X, \xi_{\text{sc}} = +1, \eta_{\text{sc}} = 0\}, \\ {}^{\text{sc}}\mathcal{R}_{\text{out}} &:= \{(\hat{h}, y, \xi_{\text{sc}}, \eta_{\text{sc}}) : \hat{h} = 0, y \in \partial X, \xi_{\text{sc}} = -1, \eta_{\text{sc}} = 0\}. \end{aligned}$$

Invariantly, $\mathcal{R}_{\text{in}} = {}^{\text{sc}}\mathcal{R}_{\text{in}}$ is the graph of $-(x/h)(dx/x) = -d(\hat{h}^{-1}) = d\hat{h}/\hat{h}^2$, and likewise for $\mathcal{R}_{\text{out}} = {}^{\text{sc}}\mathcal{R}_{\text{out}}$ but with an overall sign switch.

Definition 4.6 (conditions on the normal operator). Let $l, l' \in \mathbb{R}$, and recall (4-4).

- (1) We say that $N(P)$ is *injective at weight l on outgoing functions* if the only solution u to the equation $N(P)u = 0$ satisfying $u \in \bigcup_{N \in \mathbb{R}} H_{\text{b,sc}}^{\infty, l, -N}(\text{tf}; \hat{\mu})$ and $\text{WF}_{\text{sc}}(u) \subset {}^{\text{sc}}\mathcal{R}_{\text{out}}$ is trivial: $u \equiv 0$.
- (2) We say that $N(P)^*$ (the formal adjoint with respect to $L^2(\text{tf}; \hat{\mu})$) is *injective at weight l' on incoming functions* if the only solution v to the equation $N(P)^*v = 0$ satisfying $v \in \bigcup_{N \in \mathbb{R}} H_{\text{b,sc}}^{\infty, l', -N}(\text{tf}; \hat{\mu})$ and $\text{WF}_{\text{sc}}(v) \subset {}^{\text{sc}}\mathcal{R}_{\text{in}}$ is trivial: $v \equiv 0$.
- (3) If condition (1) and condition (2) with $l' = -l + 2$ hold, we say that $N(P)$ is *invertible at weight l* .

The wave front set assumptions here are the microlocal formulations of outgoing/incoming radiation conditions. In the special case that $N(P) = \Delta_{\hat{g}} - 1$, these assumptions are indeed equivalent to the standard Sommerfeld radiation condition. Our goal is to elevate the qualitative conditions of Definition 4.6 to quantitative estimates; see Lemma 4.8.

Changing variables in the expression (4-6) for $N(P)$ to (\hat{h}, y) gives

$$N(P) = (\hat{h}^2 D_{\hat{h}})^2 + i(n-1)\hat{h}^2 D_{\hat{h}} + \hat{h}^2 \Delta_{k(0)} - 1 + \hat{h}^2 q_{1,1}(0, y, -\hat{h} D_{\hat{h}}, D_y) + \hat{h} q_{0,1}(0, y),$$

with scattering principal symbol at ${}^{\text{sc}}T_{\text{tf} \cap \text{sf}}^* \text{tf}$ given by

$$p_{\text{tf}} = \xi_{\text{sc}}^2 + |\eta_{\text{sc}}|^2 - 1. \tag{4-15a}$$

Its Hamilton vector field is

$$H_{\text{tf}} := \hat{h}^{-1} H_{p_{\text{tf}}} = 2\xi_{\text{sc}}(\hat{h}\partial_{\hat{h}} + \eta_{\text{sc}}\partial_{\eta_{\text{sc}}}) - 2|\eta_{\text{sc}}|^2\partial_{\xi_{\text{sc}}} + \hat{h}^{-1}H_{|\eta_{\text{sc}}|^2} \tag{4-15b}$$

by (2-7), which has a source, resp. sink structure at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$, resp. ${}^{\text{sc}}\mathcal{R}_{\text{out}}$ within the characteristic set $p_{\text{tf}}^{-1}(0)$. Recall then that microlocal propagation estimates near the radial sets ${}^{\text{sc}}\mathcal{R}_{\text{in/out}}$ require suitable orders — here the decay order — of weighted Sobolev spaces to be above or below certain threshold values; see [Melrose 1994, §9], [Vasy 2018, §4.7], and [Dyatlov and Zworski 2019, Appendix E.4].

Definition 4.7 (threshold quantities). Define the functions

$$\begin{aligned} r_1 &:= \text{Im}\left(\left.{}^{\text{b}}\sigma_1(Q_{1,1})\left(-\frac{\text{d}x}{x}\right)\right|_{x=0}\right) \in C^\infty(\partial X), \\ r_0 &:= \text{Im}(q_{0,1}|_{x=0}) \in C^\infty(\partial X). \end{aligned}$$

Then the threshold quantities $r_{\text{in/out}} \in \mathbb{R}$ are defined as

$$r_{\text{in}} := -\frac{1}{2} + \frac{1}{2} \sup_{\partial X}(r_1 + r_0), \quad r_{\text{out}} := -\frac{1}{2} + \frac{1}{2} \inf_{\partial X}(r_1 - r_0).$$

We next recall that at the other end of tf , i.e., the “b-end” $\text{tf} \cap \text{cf}$, the weights l, l' in Definition 4.6 are related to the *boundary spectrum* of $N(P)$. Concretely, from the expression (4-6), we read off

$$\hat{x}^2 N(P) \in (\hat{x}D_{\hat{x}})^2 - i(n-2)\hat{x}D_{\hat{x}} + \Delta_{k(0)} + q_{1,1}(0, y, \hat{x}D_{\hat{x}}, D_y) + \hat{x} \text{Diff}_{\text{b}}(\text{tf} \setminus \text{sf}). \tag{4-16}$$

Its (dilation-invariant in \hat{x}) normal operator at $\hat{x} = 0$ is given by the sum of the first four terms, and the Mellin transformed normal operator family is defined by formally replacing $\hat{x}D_{\hat{x}}$ by multiplication with $\lambda \in \mathbb{C}$, giving

$$\widehat{N}(P)(\lambda, y, D_y) := \lambda^2 - i(n-2)\lambda + \Delta_{k(0)} + q_{1,1}(0, y, \lambda, D_y). \tag{4-17}$$

This is a holomorphic family in λ taking values in elliptic elements of $\text{Diff}^2(\partial X)$. The *boundary spectrum* of $N(P)$ is then

$$\text{spec}_{\text{b}}(N(P)) := \{\lambda \in \mathbb{C} : \widehat{N}(P)(\lambda) : C^\infty(\partial X) \rightarrow C^\infty(\partial X) \text{ is not invertible}\};$$

it is a discrete subset of \mathbb{C} , and its intersection with $|\text{Im } \lambda| < C$ is finite for any fixed value of C [Melrose 1993, §5.3]. Let us now put

$$\Lambda := \{-\text{Im } \lambda : \lambda \in \text{spec}_{\text{b}}(N(P))\}; \tag{4-18}$$

this is a discrete subset of \mathbb{R} .

Lemma 4.8 (estimates for $N(P)$). *Let $s, l \in \mathbb{R}$ and $r \in C^\infty(\overline{{}^{\text{sc}}T_{\text{tf} \cap \text{sf}}^* \text{tf}})$. Suppose that r is constant near ${}^{\text{sc}}\mathcal{R}_{\text{in/out}}$ and satisfies $r > r_{\text{in}}$ at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$, $r < r_{\text{out}}$ at ${}^{\text{sc}}\mathcal{R}_{\text{out}}$. Suppose moreover that $H_{\text{tf}} r \leq 0$ and $\sqrt{-H_{\text{tf}} r} \in C^\infty$ on ${}^{\text{sc}}T_{\text{tf} \cap \text{sf}}^* \text{tf} \cap \{p_{\text{tf}} = 0\}$ in the notation of (4-15a)–(4-15b).*

(1) If $N(P)$ is injective at weight l on outgoing functions and $l - n/2 \notin \Lambda$, then

$$\|u\|_{H_{b,sc}^{s,l,r}(\text{tf}; \hat{\mu})} \leq C \|N(P)u\|_{H_{b,sc}^{s-2,l-2,r+1}(\text{tf}; \hat{\mu})} \tag{4-19}$$

for all u for which both sides are finite.

(2) If $N(P)^*$ is injective at weight $-l + 2$ on incoming functions and $-l + 2 - n/2 \notin \Lambda$, then

$$\|v\|_{H_{b,sc}^{-s+2,-l+2,-r-1}(\text{tf}; \hat{\mu})} \leq C \|N(P)^*v\|_{H_{b,sc}^{-s,-l,-r}(\text{tf}; \hat{\mu})} \tag{4-20}$$

for all v for which both sides are finite.

(3) If $N(P)$ is invertible at weight l and $l - n/2 \notin \Lambda$,¹³ then the operator $N(P)$ is invertible as a map

$$\{u \in H_{b,sc}^{s,l,r}(\text{tf}; \hat{\mu}) : N(P)u \in H_{b,sc}^{s-2,l-2,r+1}(\text{tf}; \hat{\mu})\} \rightarrow H_{b,sc}^{s-2,l-2,r+1}(\text{tf}; \hat{\mu}).$$

Proof. This is a standard application of elliptic b-theory at $\text{tf} \cap \text{cf}$ and radial point estimates at $\text{tf} \cap \text{sf}$ in the scattering calculus as in [Melrose 1994] and [Vasy 2018, §4.8].

We first prove symbolic estimates for $N(P)$ and $N(P)^*$ which do not use the injectivity assumptions. In $\text{tf} \setminus \text{sf}$, the operator $N(P)$ is an elliptic weighted b-differential operator. Let $\phi_j \in \mathcal{C}_c^\infty(\text{tf} \setminus \text{sf})$, $j = 0, 1, 2, 3$, be identically 1 near $\text{tf} \cap \text{cf}$, with $\phi_{j+1} \equiv 1$ on $\text{supp } \phi_j$. Then, only recording the b-regularity and the weight at cf , we have

$$\|\phi_1 u\|_{H_b^{s,l}} \leq C (\|\phi_2 N(P)u\|_{H_b^{s-2,l-2}} + \|\phi_2 u\|_{H_b^{-N,l}}) \tag{4-21}$$

for any fixed N . Now, recalling (4-4), we have

$$H_b^{s,l}([0, \infty)_{\hat{x}} \times \partial X; \hat{\mu}) = H_b^{s,l-\frac{n}{2}}\left([0, \infty) \times \partial X; \left|\frac{d\hat{x}}{\hat{x}} dk(0)\right|\right).$$

Using now that $l - n/2 \notin \Lambda$, we can estimate

$$\|\phi_2 u\|_{H_b^{-N,l}} \leq C \|\hat{x}^{-2} \widehat{N}(P)(\hat{x} D_{\hat{x}}, y, D_y)(\phi_2 u)\|_{H_b^{-N-2,l-2}}$$

by passing to the Mellin transform. Since $N(P) - \hat{x}^{-2} \widehat{N}(P)(\hat{x} D_{\hat{x}}, y, D_y) \in \hat{x}^{-1} \text{Diff}_b$ by (4-16), this can be plugged into (4-21) and yields (putting back the scattering decay orders, which at this point are still arbitrary due to the localizers)

$$\|\phi_1 u\|_{H_{b,sc}^{s,l,r}} \leq C (\|\phi_3 N(P)u\|_{H_{b,sc}^{s-2,l-2,r+1}} + \|\phi_3 u\|_{H_{b,sc}^{-N,l-1,-N}}). \tag{4-22}$$

Turning to the scattering end, and with $\psi_j = 1 - \phi_j$, we claim that (now with the b-decay orders being arbitrary)

$$\|\psi_1 u\|_{H_{b,sc}^{s,l,r}} \leq C (\|\psi_0 N(P)u\|_{H_{b,sc}^{s-2,l-2,r+1}} + \|\psi_0 u\|_{H_{b,sc}^{-N,-N,-N}}). \tag{4-23}$$

This is proved by means of the scattering calculus by a combination of elliptic estimates (controlling $\psi_1 u$ away from $\Sigma_{\text{tf}} := p_{\text{tf}}^{-1}(0)$), radial point estimates at ${}^{\text{sc}}\mathcal{R}_{\text{in/out}}$, and microlocal real-principal-type estimates on $\Sigma_{\text{tf}} \setminus ({}^{\text{sc}}\mathcal{R}_{\text{in}} \cup {}^{\text{sc}}\mathcal{R}_{\text{out}})$. We only sketch the argument for the radial points in order to explain the emergence of the threshold condition on r ; details can be found, e.g., in [Vasy 2018, §4.7].

¹³This condition is automatically satisfied since, for $l - n/2 \notin \Lambda$, the operator $N(P)$ is not even Fredholm; see [Melrose 1993, §6.2].

We work in $[0, 1)_{\hat{h}} \times \partial X \subset \text{tf}$, and consider estimates near ${}^{\text{sc}}\mathcal{R}_{\text{in}}$. Fixing a cutoff function $\chi \in C_c^\infty([0, \frac{1}{2}))$, identically 1 near 0 and with $\chi' \leq 0$ on $[0, \frac{1}{2})$, we consider a commutant

$$a := \hat{h}^{-2r-1} \chi(\hat{h}/\delta) \chi(|\eta_{\text{sc}}|^2) \chi((\xi_{\text{sc}} - 1)^2),$$

$$A := \frac{1}{2}(\text{Op}_{\text{sc}}(a) + \text{Op}_{\text{sc}}(a)^*) \in \Psi_{\text{sc}}^{-\infty, 2r+1},$$

where $\delta > 0$ controls the localization near $\hat{h} = 0$. We compute the commutator

$$2 \text{Im}\langle N(P)u, Au \rangle = \left\langle \left(i[N(P), A] + 2 \frac{N(P) - N(P)^*}{2i} A \right) u, u \right\rangle.$$

(This holds directly for sufficiently decaying u , and for u as in the statement of the lemma can be justified using a regularization argument.) The principal symbol of $i[N(P), A]$ is equal to $\hat{h}H_{\text{tf}}a$. When H_{tf} falls on the cutoff in \hat{h} , the result is supported in the elliptic set of $N(P)$, and hence easily controlled. When H_{tf} falls on either of the second or third cutoff functions, the result is ≤ 0 on $\text{supp } a$ in view of the source character of ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ (or directly using (4-15b)), provided $\delta > 0$ is sufficiently small; at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ then, the principal symbol of

$$i[N(P), A] + 2 \frac{N(P) - N(P)^*}{2i} A$$

has a matching definite sign, i.e., is a *negative* multiple of \hat{h}^{-2r} , provided that

$$2 \cdot (+1) \cdot (-2r - 1) + 2 \cdot {}^{\text{sc}}\sigma \left(\frac{Q - Q^*}{2i} \right) < 0,$$

$$Q := \hat{h}q_{1,1}(0, y, -\hat{h}D_{\hat{h}}, D_y) + q_{0,1}(0, y), \tag{4-24}$$

at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$. But ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ is the graph of the 1-form $d\hat{h}/\hat{h}^2$; hence

$${}^{\text{sc}}\sigma \left(\frac{Q - Q^*}{2i} \right) = \text{Im } {}^{\text{sc}}\sigma(Q)$$

at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ is equal to

$$\text{Im } {}^{\text{b}}\sigma(q_{1,1}(0, y, -\hat{h}D_{\hat{h}}, D_y)) \left(\frac{d\hat{h}}{\hat{h}} \right) + \text{Im } q_{0,1} = r_1 + r_0$$

in the notation of Definition 4.7. The condition (4-24) thus becomes $-2r - 1 + (r_1 + r_0) < 0$, which is satisfied on all of ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ provided that $r > -\frac{1}{2} + \frac{1}{2} \sup(r_1 + r_0) = r_{\text{in}}$ there. Under this assumption, one thus obtains control on u microlocally near ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ in the space $H_{\text{sc}}^{s,r}$ by $N(P)u$ measured in $H_{\text{sc}}^{s-2,r+1}$.

The analysis at ${}^{\text{sc}}\mathcal{R}_{\text{out}}$ is similar, now using the commutant

$$\hat{h}^{-2r-1} \chi(\hat{h}/\delta) \chi(|\eta_{\text{sc}}|^2) \chi((\xi_{\text{sc}} + 1)^2).$$

The derivatives of the latter two cutoffs along H_{tf} are now positive due to the sink character of ${}^{\text{sc}}\mathcal{R}_{\text{out}}$, and the principal symbol of the commutator at the radial set is a negative multiple of \hat{h}^{-2r} (thus allowing us to propagate control from a punctured neighborhood of the radial set into the radial set itself) provided that

$$2 \cdot (-1) \cdot (-2r - 1) + 2 \cdot {}^{\text{sc}}\sigma \left(\frac{Q - Q^*}{2i} \right) < 0 \quad \text{at } {}^{\text{sc}}\mathcal{R}_{\text{out}}. \tag{4-25}$$

In view of ${}^{\text{sc}}\mathcal{R}_{\text{out}}$ being the graph of $-d\hat{h}/\hat{h}^2$ and the calculation

$$\text{Im} {}^{\text{sc}}\sigma(Q)|_{{}^{\text{sc}}\mathcal{R}_{\text{out}}} = \text{Im} {}^{\text{b}}\sigma(q_{1,1}(0, y, -\hat{h}D_{\hat{h}}, D_y))\left(-\frac{d\hat{h}}{\hat{h}}\right) + \text{Im} q_{0,1} = -r_1 + r_0,$$

the condition (4-25) reads $2r + 1 - r_1 + r_0 < 0$, so $r < -\frac{1}{2} + \frac{1}{2} \inf(r_1 - r_0) = r_{\text{out}}$.

Putting (4-22) and (4-23) together, we obtain the estimate

$$\|u\|_{H_{\text{b,sc}}^{s,l,r}} \leq C(\|N(P)u\|_{H_{\text{b,sc}}^{s-2,l-2,r+1}} + \|u\|_{H_{\text{b,sc}}^{-N,l-1,-N}}) \tag{4-26}$$

for any N ; we choose N to satisfy $-N < s$ and $-N < \min r$.

The estimate (4-26) implies that $N(P)$, acting on $H_{\text{b,sc}}^{s,l,r}$, has finite-dimensional kernel; any element u in the kernel automatically lies in $H_{\text{b,sc}}^{\infty,l,r'}$ for any variable-order function r' satisfying $r' < r_{\text{out}}$ at ${}^{\text{sc}}\mathcal{R}_{\text{out}}$. Thus, $\text{WF}_{\text{sc}}(u) \subset {}^{\text{sc}}\mathcal{R}_{\text{out}}$. Under the injectivity assumption on $N(P)$, we thus conclude that $u = 0$. A standard functional analytic argument then allows one to drop the error term in (4-26), which gives the estimate (4-19).

The proof of part (2) is analogous; the direction of propagation in the characteristic set is now reversed, which is precisely matched by the sign switches in the orders in the estimate (4-20). Part (3) is an immediate consequence of the first two parts. □

Remark 4.9 (flexibility in the choice of l). If the assumptions of part (1) of the lemma are satisfied for some value of l , then they continue to hold for all values \tilde{l} with $\tilde{l} - n/2 \notin \Lambda$ for which either $\tilde{l} > l$, or $\tilde{l} \leq l$ but $\tilde{l} - n/2$ and $l - n/2$ lie in the same connected component (a, b) of $\mathbb{R} \setminus \Lambda$. (Indeed, the claim for $\tilde{l} \leq l$ follows from the fact — proved using the Mellin transform upon localizing near $\text{tf} \cap \text{cf}$ — that any element in the kernel of $N(P)$ on $H_{\text{b,sc}}^{s,\tilde{l},r}$ automatically lies in $H_{\text{b,sc}}^{s,b+n/2-\epsilon,r}$ for any $\epsilon > 0$.) A similar statement holds for part (2): we may increase $-l + 2$ (or stay in the same connected component of $(\mathbb{R} \setminus \Lambda) + n/2$), i.e., decrease l . Altogether then, there typically only exists an interval of finite length (possibly empty) of weights l so that the invertibility condition of part (3) is satisfied.

4D. Statement and proof of the microlocal propagation estimate. We are now ready to state the main result of the paper:

Theorem 4.10 (microlocal propagation through the cone point). *Let $P_{h,z}$ denote an admissible operator in the sense of Definition 4.1, and define the threshold quantities $r_{\text{in}}, r_{\text{out}}$ as in Definition 4.7. Let $\Sigma \subset {}^{\text{ch}}T_{\text{sf}}^*X_{\text{ch}}$ denote the characteristic set of $P_{h,z}$ (see (4-8)). Denote by $H = (x/h)H_p \in \mathcal{V}({}^{\text{ch}}T_{\text{sf}}^*X_{\text{ch}})$ the rescaled Hamilton vector field (see (4-9)). Let $s, l, \alpha \in \mathbb{R}$, $\text{b} \in C^\infty(\overline{{}^{\text{ch}}T_{\text{sf}}^*X_{\text{ch}}})$. Assume that b is constant near the radial sets $\mathcal{R}_{\text{in/out}}$ (see (4-11)) and satisfies $\text{b} - \alpha > r_{\text{in}}$ at \mathcal{R}_{in} and $\text{b} - \alpha < r_{\text{out}}$ at \mathcal{R}_{out} ; assume moreover that $H\text{b} \leq 0$ and $\sqrt{-H\text{b}} \in C^\infty$ on Σ . Let $\chi, \tilde{\chi} \in C_c^\infty(X)$ be cutoffs, identically 1 near ∂X , and with $\tilde{\chi} \equiv 1$ on $\text{supp } \chi$. Let $E \in \Psi_{\text{ch}}^0(X)$, with Schwartz kernel supported in $[0, 1)_h \times (\tilde{\chi}^{-1}(1) \times \tilde{\chi}^{-1}(1))$.*

(1) (forward propagation) *Suppose $N(P)$ is injective at weight l on outgoing functions (see Definition 4.6(1)). Suppose that (the preimage in Σ of) all backward GBBs (see Definition 4.5) starting in $\Sigma \cap \text{supp } \chi$ reach $\text{Ell}_{\text{ch}}(E)$ in finite time while remaining inside $\tilde{\chi}^{-1}(1)$. Then for some small $\delta > 0$, we have*

$$\|\chi u\|_{H_{\text{c,h}}^{s,l,\alpha,\text{b}}(X)} \leq C(\|\tilde{\chi} P_{h,z}u\|_{H_{\text{c,h}}^{s-2,l-2,\alpha,\text{b}+1}(X)} + \|Eu\|_{H_{\text{c,h}}^{s,l,\alpha,\text{b}}(X)} + h^\delta \|\tilde{\chi}u\|_{H_{\text{c,h}}^{-N,l,\alpha,\text{b}}(X)}). \tag{4-27}$$

(2) (backward propagation) Suppose $N(P)^*$ is injective at weight $-l + 2$ on incoming functions (see Definition 4.6(2)). Suppose that (the preimage in Σ of) all forward GBBs starting in $\Sigma \cap \text{supp } \chi$ reach $\text{Ell}_{c\hbar}(E)$ in finite time while remaining inside $\tilde{\chi}^{-1}(1)$. Then for some small $\delta > 0$, we have

$$\| \chi u \|_{H_{c,h}^{-s+2, -l+2, -\alpha, -b-1}(X)} \leq C (\| \tilde{\chi} P_{h,z}^* u \|_{H_{c,h}^{-s, -l, -\alpha, -b}(X)} + \| Eu \|_{H_{c,h}^{-s+2, -l+2, -\alpha, -b-1}(X)} + h^\delta \| \tilde{\chi} u \|_{H_{c,h}^{-N, -l+2, -\alpha, -b-1}(X)}). \quad (4-28)$$

Since by Lemma 4.3 and the calculations in Section 4B, the operator $P_{h,z} \in \Psi_{c\hbar}^{2,2,0,0}(X)$ is elliptic at fiber infinity, and is of real-principal-type (except at the radial points) at sf, the estimates (4-27) and (4-28) are sharp as far as the relative orders in the norms on u on the left and $P_{h,z}^{(*)}u$ on the right are concerned. Indeed, it has the well-known real-principal-type loss of one order at sf and is an elliptic estimate in the $c\hbar$ -differentiability sense.

The improvement of the final (error) terms on the right-hand sides in (4-27) and (4-28) relative to the space on the left-hand sides is accomplished at sf by microlocal symbolic means, and at tf using global normal operator estimates. The overall improvement by a positive power of h between error term and left-hand side allows for the inversion of $P_{h,z}$ for small $h > 0$ under suitable assumptions on the global behavior of the null-bicharacteristic flow; see Sections 4E and 5 for examples.

Remark 4.11 (operators on vector bundles). Let $E \rightarrow X$ denote a smooth vector bundle. Theorem 4.10 then holds (with the same proof) also for operators $P_{h,z}$ acting on sections of E , provided $P_{h,z}$ is *admissible* in the sense that

$$P_{h,z} = h^2 x^{-2} Q_{2,z} + h x^{-1} q_{0,z} - z, \\ Q_{2,z} \in \text{Diff}_{\text{b}}^2(X; E), \quad q_{0,z} \in C^\infty(X; \text{End}(E)),$$

where $x^{-2}Q_{2,z}$ (replacing the combination $h^2\Delta_g + h^2x^{-2}Q_{1,z}$ in Definition 4.1) has scalar principal symbol ${}^{\text{b}}\sigma(x^{-2}Q_{2,z}) = {}^{\text{b}}\sigma(x^{-2}\Delta_g)$. That is, ${}^{\text{b}}\sigma(x^{-2}Q_{2,z})(\zeta) = |\zeta|_g^{-1}$ for $\zeta \in {}^{\text{b}}T^*X$, with g the conic metric (4-2). The normal operator is of class

$$N(P) \in \left(\frac{\hat{x}}{\hat{x} + 1} \right)^{-2} \text{Diff}_{\text{b,sc}}^2(\text{tf}; \pi^* E_{\partial X}),$$

where $\pi : \text{tf} = [0, \infty]_{\hat{x}} \times \partial X \rightarrow \partial X$ denotes the projection map. The injectivity conditions of Definition 4.6 are unchanged. The definition of the threshold quantities $r_{\text{in/out}}$ in Definition 4.7 requires a minor change; to wit, with respect to a choice of a positive definite fiber inner product on $E_{\partial X}$, we set (top sign for “in”, bottom sign for “out”)

$$r_{\text{in/out}} := -\frac{1}{2} \pm \frac{1}{2} \sup_{\partial X}^{\text{sc}} \sigma \left(\frac{x^{-2}Q_{2,1} - (x^{-2}Q_{2,1})^* + q_{0,1} - q_{0,1}^*}{2i} \right) \Big|_{\mp \frac{dx}{x^2}}, \quad (4-29)$$

where the sup is defined to be the supremum of the largest eigenvalue of the scattering symbol (which takes values in self-adjoint endomorphisms of E). One may choose different fiber inner products in the calculation of r_{in} and r_{out} , respectively. A (near-)optimal choice of fiber inner products, resulting in (almost) the smallest possible r_{in} and largest possible r_{out} , is typically easy to read off in concrete situations. For example, if $Q_{2,1} = 0$ and $q_{0,1}|_{\partial X}$ is block-diagonal (or more generally lower triangular)

with respect to some bundle splitting of $E|_{\partial X}$, then the supremum in (4-29) can be made to be arbitrarily close to the supremum of $\text{Im } \lambda$, where λ ranges over all eigenvalues of the diagonal entries of $q_{0,1}(y)$, $y \in \partial X$, if one chooses the fiber inner product appropriately.

Remark 4.12 (technical assumptions on the variable order). One can replace the assumptions that b be locally constant near $\mathcal{R}_{\text{in/out}}$ and satisfy $\sqrt{-\text{H}b} \in C^\infty$ on Σ by the simpler assumption that $\text{H}b \leq 0$ on Σ . This would require the use of the sharp Gårding inequality for the $c\hbar$ -calculus, which however we do not prove here.

Proof of Theorem 4.10. We give details for the proof of part (1); the proof of part (2) is completely analogous. If backward GBBs starting in $\text{WF}'_{c\hbar}(B)$ never pass through $\partial\Sigma \subset {}^{c\hbar}T^*_{\text{sf}\cap\text{tf}}X_{c\hbar}$, the orders l and a are irrelevant, and the estimate (4-27) follows from standard elliptic regularity and real-principal-type propagation in the (variable-order) semiclassical calculus on X° . We shall thus work in a small neighborhood of $x = 0$.

Step 1: symbolic positive commutator estimate. We first work near the incoming radial set \mathcal{R}_{in} defined in (4-11); we shall use the coordinates $(\hat{h}, x, y, \xi, \eta)$ near ${}^{c\hbar}T^*_{\text{sf}\cap\text{tf}}X_{c\hbar}$ defined by (3-3) and (4-4). Fix cutoffs $\chi_\partial, \chi_{\text{sf}}, \chi_{\mathcal{R}} \in C_c^\infty([0, 1])$, identically 1 near 0 and satisfying $\chi'_\bullet \leq 0$ and $\sqrt{-\chi_\bullet \chi'_\bullet} \in C^\infty([0, 1])$. Denote a smooth extension of b to ${}^{c\hbar}T^*X_{c\hbar}$ by the same symbol. For small $\delta > 0$, fixed momentarily, we then consider a commutant

$$\check{a} = \hat{h}^{-b-\frac{1}{2}} x^{-\alpha} \chi_\partial \left(\frac{x}{\delta} \right) \chi_{\text{sf}} \left(\frac{\hat{h}}{\delta} \right) \chi_{\mathcal{R}} \left(\frac{\omega}{\delta} \right), \quad \omega := \sqrt{|\eta|^2 + |\xi + 1|^2}. \tag{4-30}$$

Thus, $\text{supp } a$ is contained in any fixed open neighborhood of \mathcal{R}_{in} when $\delta > 0$ is sufficiently small. We have $\check{a} \in S^{-\infty, -\infty, \alpha, b+1/2}({}^{c\hbar}T^*X_{c\hbar})$. Let

$$\check{A} \in \Psi_{c\hbar}^{-\infty, -\infty, \alpha, b+\frac{1}{2}}(X), \quad {}^{c\hbar}\sigma(\check{A}) = \check{a}, \quad A := \check{A}^* \check{A}.$$

Using the $L^2(X; \mu)$ inner product, we then evaluate the commutator

$$\begin{aligned} 2 \text{Im} \langle \check{A} P_{h,z} u, \check{A} u \rangle &= \langle \mathcal{C} u, u \rangle, \\ \mathcal{C} &= i[P_{h,z}, A] + 2 \frac{P_{h,z} - P_{h,z}^*}{2i} A \in \Psi_{c\hbar}^{-\infty, -\infty, 2a, 2b}(X). \end{aligned} \tag{4-31}$$

The principal symbol of \mathcal{C} is

$$2\hat{h}\check{a}\check{H}\check{a} + 2 \cdot {}^{c\hbar}\sigma \left(\frac{P_{h,z} - P_{h,z}^*}{2i\hat{h}} \right) \hat{h}\check{a}^2. \tag{4-32}$$

When H hits χ_∂ , we obtain a nonnegative contribution (in fact, the square e^2 of a smooth function e), while differentiation of $\chi_{\mathcal{R}}$ gives a nonpositive contribution (in fact, a negative square $-b_{\mathcal{R}}^2$), consistently with the saddle point structure of H at \mathcal{R}_{in} . Differentiation of χ_{sf} produces a symbol with semiclassical order $-\infty$.

The main term of $\hat{h}\check{a}\check{H}\check{a}$ near \mathcal{R}_{in} arises from differentiation of the weight $\hat{h}^{-b-1/2} x^{-\alpha}$; since $H_p = \hat{h}\text{H}$ is, modulo $\hat{h}x\mathcal{V}_b$, given by the expression (4-10), we can compute this modulo $xS^{-\infty, -\infty, 2a, 2b}$ by substituting

the expression (4-10) of $H|_{x=0}$ for H . Thus, the main term is

$$\hat{h}^{-2b} x^{-2\alpha} (2\xi(-2\alpha + 2b + 1) + \mathcal{O}(x)) \chi_{\delta}^2 \chi_{\text{sf}}^2 \chi_{\mathcal{R}}^2.$$

A further contribution arises from the skew-adjoint part of $P_{h,z}$ at \mathcal{R}_{in} , which is the same as the skew-adjoint part of $N(P)$ at ${}^{\text{sc}}\mathcal{R}_{\text{in}}$ upon making the identification (3-7); this was already computed in the proof of Lemma 4.8. Overall then, we can write

$${}^{\text{ch}}\sigma(\mathcal{C}) = e^2 - b_{\mathcal{R}}^2 - \epsilon \hat{h} \check{a}^2 - f^2 \hat{h} \check{a}^2, \tag{4-33}$$

where

$$f = \sqrt{-[2(-2(b - \alpha) - 1) + 2(r_1 + r_0)] - \epsilon}$$

is positive (and smooth) at \mathcal{R}_{in} for small $\epsilon > 0$. Denoting ch -quantizations of the lower case symbols by the corresponding upper case letters, we thus have

$$\mathcal{C} = E^* E - B_{\mathcal{R}}^* B_{\mathcal{R}} - \epsilon \|\hat{h}^{\frac{1}{2}} \check{A} u\|^2 - \|\hat{h}^{\frac{1}{2}} F \check{A} u\|^2 + R,$$

where $R \in \Psi_{\text{ch}}^{-\infty, -\infty, 2\alpha, 2b-1}(X)$ has $\text{WF}'_{\text{ch}}(R) \subset \text{supp } \check{a}$ and arises as the remainder term not controlled by the previous symbolic considerations. We will plug this into the right-hand side of (4-31); the left-hand side is bounded from below by

$$-\epsilon \|\hat{h}^{\frac{1}{2}} \check{A} u\|^2 - \epsilon^{-1} \|\hat{h}^{-\frac{1}{2}} \check{A} P_{h,z} u\|^2 \geq -\epsilon \|\hat{h}^{\frac{1}{2}} \check{A} u\|^2 - C \epsilon^{-1} \|G P_{h,z} u\|_{H_{c,h}^{-N, -N, \alpha, b+1}} - C \epsilon^{-1} \|\tilde{\chi} u\|_{H_{c,h}^{-N, -N, \alpha, -N}},$$

where $G \in \Psi_{\text{ch}}^0(X)$ is elliptic on $\text{supp } \check{a}$; here $N \in \mathbb{R}$ is arbitrary. Putting $B_0 := \hat{h}^{1/2} F \check{A} \in \Psi_{\text{ch}}^{-\infty, -\infty, \alpha, b}(X)$ and dropping the contribution of $B_{\mathcal{R}}$, we thus obtain the estimate

$$\|B_0 u\|^2 \lesssim \|G P_{h,z} u\|_{H_{c,h}^{-N, -N, \alpha, b+1}}^2 + \|E u\|^2 + |\langle Ru, u \rangle| + \|\tilde{\chi} u\|_{H_{c,h}^{-N, -N, \alpha, -N}}^2, \tag{4-34}$$

which provides $H_{c,h}^{-N, -N, \alpha, b}$ -control of u microlocally near \mathcal{R}_{in} provided one has microlocal $H_{c,h}^{-N, -N, -N, b}$ -control of u on $\text{WF}'_{\text{ch}}(E) \subset \{0 < x < \delta, |\xi + 1| < \delta\}$, and provided $|\langle Ru, u \rangle|$ is finite; since G is elliptic near $\text{WF}'_{\text{ch}}(R)$, we can insert the estimate

$$|\langle Ru, u \rangle| \lesssim \|G u\|_{H_{c,h}^{-N, -N, \alpha, b-1/2}}^2 + \|\tilde{\chi} u\|_{H_{c,h}^{-N, -N, \alpha, -N}}^2$$

into the right-hand side of (4-34).

Concatenating this radial point estimate with the propagation of regularity from a punctured neighborhood of \mathcal{R}_{in} to a punctured neighborhood of \mathcal{R}_{out} and then a radial point estimate at \mathcal{R}_{out} — proved by the same method, with the commutant \check{a} again given by (4-30) but now with $\omega = (|\eta|^2 + |\xi - 1|^2)^{1/2}$ and using that $2(b - \alpha) + 1 - r_1 + r_0 < 0$ — and using elliptic estimates away from Σ , we obtain the propagation estimate

$$\|B_1 u\|_{H_{c,h}^{s,l,\alpha,b}} \lesssim \|G P_{h,z} u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1}} + \|E u\|_{H_{c,h}^{s,-N,-N,b}} + \|\tilde{\chi} u\|_{H_{c,h}^{-N,l,\alpha,b-1/2}}; \tag{4-35}$$

the operators $B_1, G, E \in \Psi_{\text{ch}}^{0,0,0,0}(X)$ appearing here are subject to the following conditions: $\text{WF}'_{\text{ch}}(B_1) \subset \text{Ell}_{\text{ch}}(G)$, furthermore G is elliptic on $\Sigma \cap x^{-1}(0)$, and all backward null-bicharacteristics from $\text{WF}'_{\text{ch}}(B_1)$

either tend to \mathcal{R}_{out} or enter $\text{Ell}_{ch}(E)$. The orders at cf are unconstrained at this point, but chosen for compatibility with the normal operator argument below.

Fixing a variable-order function b^\flat so that

$$b^\flat - \alpha > r_{\text{in}} \quad \text{at } \mathcal{R}_{\text{in}}, \tag{4-36}$$

we then have the estimate

$$\|\chi u\|_{H_{c,h}^{s,l,\alpha,b}} \lesssim \|\tilde{\chi} P_{h,z} u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1}} + \|Eu\|_{H_{c,h}^{s,-N,-N,b}} + \|\tilde{\chi} u\|_{H_{c,h}^{-N,l,\alpha,b^\flat}} \tag{4-37}$$

under the assumptions on $E, \chi, \tilde{\chi}$ stated in the theorem. For $b^\flat \geq b - \frac{1}{2}$, this follows directly from (4-35) (upon replacing the microlocal cutoff G by the less precise cutoff $\tilde{\chi}$). For general b^\flat , note that as long as (4-36) is satisfied, one can apply this estimate inductively to the error term $\tilde{\chi} u$ provided $\text{supp } \tilde{\chi}$ is sufficiently close to $\text{supp } \chi$ (so that the same operator Eu satisfies the geometric control assumption for $\tilde{\chi}$ in place of χ), increasing supports of the involved cutoff functions by an arbitrarily small but positive amount and gaining half a semiclassical order at each step. Thus, away from \mathcal{R}_{in} , one can ultimately take b^\flat to be arbitrarily negative, while at \mathcal{R}_{in} , one always needs to have (4-36).

Step 2: normal operator estimate. We now work on the error term $\tilde{\chi} u$ in (4-37). We first prove the desired estimate (4-27) under the stronger condition that $b - \alpha > r_{\text{in}} + 1$ at \mathcal{R}_{in} . We split $\tilde{\chi} u = \chi^\flat u + (1 - \chi^\flat) \tilde{\chi} u$, where $\chi^\flat \in C^\infty(X)$ is identically 1 near ∂X and supported in a very small neighborhood of ∂X ; the part $(1 - \chi^\flat) \tilde{\chi} u$ is supported away from $\text{cf} \cup \text{tf}$; hence

$$\|(1 - \chi^\flat) \tilde{\chi} u\|_{H_{c,h}^{-N,l,\alpha,b^\flat}} \lesssim \|(1 - \chi^\flat) \tilde{\chi} u\|_{H_{c,h}^{-N,-N,-N,b^\flat}}$$

for any N . To estimate $\chi^\flat u$, we use the injectivity assumption on $N(P)$ and the resulting estimate (4-19) together with Corollary 3.7(2) (with $\alpha_\mu = n$). For $0 < \delta < 1$ with $b^\flat + 2\delta < b$, and choosing $\text{supp } \chi^\flat$ sufficiently small, we obtain

$$\begin{aligned} \|\chi^\flat u\|_{H_{c,h}^{-N,l,\alpha,b^\flat}} &\lesssim h^{\frac{n}{2}-\alpha} \|\pi^*(\chi^\flat u)\|_{H_{b,\text{sc}}^{-N,l,b^\flat|\text{sf}\cap\text{tf}-\alpha+\delta}} \\ &\lesssim h^{\frac{n}{2}-\alpha} \|N(P)(\pi^*(\chi^\flat u))\|_{H_{b,\text{sc}}^{-N-2,l-2,b^\flat|\text{sf}\cap\text{tf}-\alpha+\delta+1}} \\ &\lesssim \|N(P)(\chi^\flat u)\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\flat+1+2\delta}}. \end{aligned} \tag{4-38}$$

In the final line, we abuse notation and denote by $N(P) \in \Psi_{ch}^{2,2,0,0}(X)$ any operator whose normal operator is equal to $N(P)$. Put $b^\sharp := b^\flat + 1 + 2\delta$. Using Lemma 4.3, which gives $N(P) - P_{h,z} \in \Psi_{ch}^{2,2,-1,0}(X)$, we further estimate

$$\begin{aligned} &\|N(P)(\chi^\flat u)\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} \\ &\leq \|\chi^\flat P_{h,z} u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} + \|\chi^\flat (P_{h,z} - N(P))u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} + \|[N(P), \chi^\flat]u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} \end{aligned} \tag{4-39}$$

$$\lesssim \|\chi^\flat P_{h,z} u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} + \|\chi^\flat u\|_{H_{c,h}^{-N,l,\alpha-1,b^\sharp}} + \|\chi^\sharp u\|_{H_{c,h}^{-N-1,-N,-N,b^\sharp-1}}, \tag{4-40}$$

where $\chi^\sharp \equiv 1$ on $\text{supp } \chi^\flat$. Under the present condition that $b - \alpha > r_{\text{in}} + 1$ at \mathcal{R}_{in} , we can choose b^\flat as in (4-36) so that $b^\sharp < b$ still. Plugging this into (4-37) finishes the proof of part (1) under this condition.

In order to prove the theorem as stated, thus only assuming $b - \alpha > r_{\text{in}}$ at \mathcal{R}_{in} , we note that the norm on second term on the right in (4-40) is one order weaker at tf than the left-hand side of the desired estimate (4-27), but only $b^\sharp - b = b^\flat + 1 + 2\delta - b < 1$ orders stronger at sf . This suggests revisiting the estimates (4-38)–(4-40) using a more precise cutoff which distinguishes between the regimes $\hat{h} \lesssim x$ and $\hat{h} \gtrsim x$. To wit, consider $\psi^b = \tilde{\psi}^b(\hat{h}/x)$, where $\tilde{\psi}^b \equiv 0$ on $[0, 1]$ and $\tilde{\psi}^b \equiv 1$ on $[2, \infty)$. This is a smooth function on $[X_{c\hat{h}}; \text{sf} \cap \text{tf}]$, and thus conormal on $X_{c\hat{h}}$; in fact, we have

$$1 - \psi^b \in \mathcal{A}^{0,\zeta,-\zeta}(X_{c\hat{h}}) = (x + h)^\zeta \left(\frac{h}{h+x} \right)^{-\zeta} \mathcal{A}^0(X_{c\hat{h}}) \subset \Psi_{c\hat{h}}^{0,0,-\zeta,\zeta}(X) \tag{4-41}$$

for any $\zeta \geq 0$, since on $\text{supp}(1 - \psi^b)$ we have $x \lesssim \hat{h}$; thus $x + h \lesssim h/(h+x)$. Taking $\zeta = \delta$, we can therefore estimate

$$\|(1 - \psi^b)\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha,b^\flat}} \lesssim \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha-\delta,b^\flat+\delta}} \leq h^\delta \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha,b}}.$$

Next, the estimate (4-38) holds without change. (Note that Corollary 3.7 applies for merely conormal cutoffs.) Finally, we need to estimate (4-39) more carefully. Note that

$$\psi^b \in \mathcal{A}^{0,-\zeta,\zeta}(X_{c\hat{h}}) \subset \Psi_{c\hat{h}}^{0,0,\zeta,-\zeta}(X)$$

for any $\zeta \geq 0$. Taking $\zeta = 1 - \delta$, this gives $\psi^b(P_{h,z} - N(P)) \in \Psi_{c\hat{h}}^{2,2,-\delta,-1+\delta}(X)$; hence

$$\|\psi^b(P_{h,z} - N(P))u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} \lesssim \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha-\delta,b^\sharp-1+\delta}} \leq h^\delta \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha,b}}.$$

For the final, commutator, term in (4-39), we note that we can replace ψ^b by $1 - \psi^b$ and use (4-41) with $\zeta = \delta$, so $[N(P), \psi^b] \in \Psi_{c\hat{h}}^{1,-\infty,-\delta,-1+\delta}(X)$, which gives

$$\|[N(P), \psi^b]u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} \lesssim \|\tilde{\chi}u\|_{H_{c,h}^{-N-1,-N,\alpha-\delta,b^\sharp-1+\delta}} \leq h^\delta \|\tilde{\chi}u\|_{H_{c,h}^{-N-1,-N,\alpha,b}}.$$

Altogether, we have shown

$$\|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha,b^\flat}} \lesssim \|\tilde{\chi}P_{h,z}u\|_{H_{c,h}^{-N-2,l-2,\alpha,b^\sharp}} + h^\delta \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha,b}}.$$

Plugged into (4-37), we have now established the desired estimate (4-27). □

We can sharpen Theorem 4.10 by working with the resolved Sobolev spaces defined in (3-24). This is straightforward since admissible operators

$$P_{h,z} \in \Psi_{c\hat{h}}^{2,2,0,0}(X) \subset \Psi_{cb\hat{h}}^{2,2,2,0,0}(X)$$

are elliptic at the front face fbf of $\overline{cb\hat{h}T^*X_{c\hat{h}}}$; indeed, this follows from the ellipticity at fiber infinity $c\hat{h}S_{\text{tf}}^*X_{c\hat{h}} \subset \overline{c\hat{h}T^*X_{c\hat{h}}}$ and the classical nature of the principal symbol of $P_{h,z}$. Therefore:

Theorem 4.13 (propagation estimates with relative b -regularity). *In the notation of Theorem 4.10, and for any $s' \in \mathbb{R}$, the forward propagation estimate (4-27) generalizes to an estimate on $cb\hat{h}$ -Sobolev spaces,*

$$\|\chi u\|_{H_{cb,h}^{s,s',l,\alpha,b}(X)} \leq C(\|\tilde{\chi}P_{h,z}u\|_{H_{cb,h}^{s-2,s'-2,l-2,\alpha,b+1}(X)} + \|Eu\|_{H_{cb,h}^{s,s',l,\alpha,b}(X)} + h^\delta \|\tilde{\chi}u\|_{H_{cb,h}^{-N,-N,l,\alpha,b}(X)}).$$

The backward propagation estimate (4-28) generalizes similarly.

4E. Global estimates with complex absorption. We upgrade the microlocal estimate proved above into a quantitative invertibility statement for an operator which effectively localizes the interesting nonelliptic phenomena near the cone point into a small neighborhood of ∂X via complex absorption.

Let us assume first for simplicity that Y is null-cobordant; see Remark 4.16 below on how to remove this assumption. With $X = [0, 2x_0] \times Y$ and g as in (4-1)–(4-2), consider then a compact n -dimensional manifold $X' \supset X$ with boundary $\partial X' = \partial X$, equipped with a smooth metric g' which is equal to g on $X^b := [0, x_0] \times Y$. Given an admissible operator $P_{h,z}$ on X , let $P'_{h,z} \in (x/(x+h))^{-2} \text{Diff}_{ch}^2(X')$ denote an extension of $P_{h,z}$ from $[0, 1) \times X^b$ to $[0, 1) \times X'$ with principal part equal to $h^2 \Delta_{g'}$. For $c \in (0, 1)$, let

$$K_c := X' \setminus ([0, cx_0] \times Y).$$

In order to implement complex absorption, let us take $c \in (0, \frac{1}{2})$ small and fix an operator

$$Q \in \Psi_h^{-\infty}((X')^\circ)$$

whose Schwartz kernel is supported in $K_c \times K_c$, and so that Q is elliptic on T^*K_{2c} with nonnegative principal symbol. We then consider

$$\mathcal{P}_{h,z} := P'_{h,z} - iQ, \tag{4-42}$$

and assume that

$$\text{all backward GBBs of } P'_{h,z} \text{ enter } \text{Ell}_h(Q) \text{ in finite time.} \tag{4-43}$$

By construction, $\mathcal{P}_{h,z}$ is a semiclassically elliptic second-order semiclassical ps.d.o. on $(X')^\circ$ which is elliptic over K_{2c} . Moreover, due to the sign condition on the principal symbol of Q , one can propagate semiclassical regularity for solutions of $\mathcal{P}_{h,z}u = f$ along *forward* null-bicharacteristics of $P'_{h,z}$; see [Vasy 2013, §2.5] and [Dyatlov and Zworski 2019, §5.6.3]. For our fixed metric g on $[0, 2x_0] \times Y$, the control condition (4-43) is satisfied if we choose $c > 0$ sufficiently small. Indeed, from the expression (4-13), one finds that if $H_{\text{sf}}x = 2\xi = 0$ on the characteristic set, then $|\eta|^2 = 1$ and thus

$$H_{\text{sf}}^2x = 2H\xi \geq 2x^{-1}|\eta|^2 - C|\eta|^2 = 2x^{-1} - C > c^{-1} - C > 0$$

in $x < 2c$ when c is sufficiently small; hence the level sets of x are geodesically convex in $x < 2c$, which implies the claim.

Remark 4.14 (relaxed conditions on Q). One can more generally allow Q to be a second-order operator with real-principal symbol; a concrete choice is then $Q = \psi \cdot (h^2 \Delta_{g'} + 1)$, where $\psi \in C_c^\infty(K_c^\circ)$ is identically 1 on K_{2c} .

We then have:

Proposition 4.15 (global estimates with complex absorption). *Let s, l, α, b be as in the statement of Theorem 4.10 (for the operator $P_{h,z}$). Fix the volume density on X' to be the metric density $|\text{dg}'|$. Then for small $h > 0$, the operator $\mathcal{P}_{h,z}$ defined by (4-42) is invertible as a map $H_b^{s,l}(X') \rightarrow H_b^{s-2,l-2}(X')$, and it satisfies the uniform estimate*

$$\|u\|_{H_{c,h}^{s,l,\alpha,b}(X')} \leq C \|\mathcal{P}_{h,z}u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1}(X')} = Ch^{-1} \|(x+h)\mathcal{P}_{h,z}u\|_{H_{c,h}^{s-2,l-2,\alpha,b}}. \tag{4-44}$$

More generally, for any $s' \in \mathbb{R}$, we have

$$\|u\|_{H_{cb,h}^{s',l,\alpha,b}(X')} \leq C \|\mathcal{P}_{h,z}u\|_{H_{cb,h}^{s-2,s'-2,l-2,\alpha,b+1}}.$$

Proof. By our assumptions on the complex-absorbing potential Q , we can apply Theorem 4.10(1) with E and χ supported in $X' \setminus K_c$. We thus have

$$\|\chi u\|_{H_{c,h}^{s,l,\alpha,b}} \leq C(\|\mathcal{P}_{h,z}u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1}} + \|Eu\|_{H_{c,h}^{s,-N,-N,b}} + h^\delta \|\tilde{\chi}u\|_{H_{c,h}^{-N,l,\alpha-1,b^b}}).$$

On the other hand, we can control Eu and $(1 - \chi)u$ in $H_{c,h}^{s,l,\alpha,b}$ (or simply $h^b H_h^s$ if we take E to be localized away from $x = 0$, as we may arrange) by $\mathcal{P}_{h,z}u$ in $H_{c,h}^{s-2,l-2,\alpha,b+1}$ using a combination of elliptic estimates and real-principal-type propagation estimates (with complex absorption), starting either from $\text{Ell}_h(Q)$ or $\{\chi = 1\}$. Altogether, we obtain

$$\|u\|_{H_{c,h}^{s,l,\alpha,b}} \leq C(\|\mathcal{P}_{h,z}u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1}} + h^\delta \|u\|_{H_{c,h}^{-N,l,\alpha,b}}). \tag{4-45}$$

For $h_0 > 0$ with $Ch_0^\delta < \frac{1}{2}$, we can now drop the error term in (4-45) for $0 < h < h_0$. This proves the injectivity of $\mathcal{P}_{h,z}$ (with a quantitative estimate). Analogous arguments prove the dual estimate

$$\|v\|_{H_{c,h}^{-s+2,-l+2,-\alpha,-b-1}} \leq C \|\mathcal{P}_{h,z}^*v\|_{H_{c,h}^{-s,-l,-\alpha,-b}},$$

which implies the surjectivity of $\mathcal{P}_{h,z}$. □

Remark 4.16 (links Y that are not null-cobordant). When Y is not null-cobordant, we cannot choose X' as above. This is a technical issue, independent of the analysis near the cone point $x^{-1}(0)$, which we circumvent here with the following artificial device: we set $X' := [0, 4x_0] \times Y$, and consider h -dependent families of operators on X' which are semiclassical cone operators near $x^{-1}(0)$ and semiclassical scattering operators [Vasy and Zworski 2000] near $x^{-1}(4x_0)$. We then take $P'_{h,z} \in (x/(x+h))^{-2} \text{Diff}_{c\hbar,sc\hbar}^2(X')$ — the second subscript referring to the semiclassical scattering behavior near $x^{-1}(4x_0)$ — to be equal to $P_{h,z}$ on $[0, 1) \times X^b$. We can arrange for $P'_{h,z}$ to be elliptic near $x^{-1}(4x_0)$ in the semiclassical scattering algebra, e.g., by taking it to be equal to $h^2 \Delta_{g'} + 1$ near $x = 4x_0$, where g' is a scattering metric on $(x_0, 4x_0] \times Y$. We choose the complex absorbing operator Q as before, and so that $\mathcal{P}_{h,z} = P'_{h,z} - iQ$ is elliptic in $x > \frac{1}{2}x_0$. Proposition 4.15 then remains valid upon using function spaces for u which near $x^{-1}(4x_0)$ are semiclassical scattering Sobolev spaces with differential order s , semiclassical order b , and arbitrary decay order r , and similarly for $\mathcal{P}_{h,z}u$ with orders $s - 2, b + 1, r$; this uses elliptic estimates in the semiclassical scattering algebra near $x^{-1}(4x_0)$. (This usage of the semiclassical scattering algebra is only one of several possibilities in which the invertibility of $\mathcal{P}_{h,z}$ for small h is easy to obtain despite the presence of a boundary.)

4F. Propagation of Lagrangian regularity: diffractive improvement. By adapting arguments from [Melrose and Wunsch 2004; Melrose et al. 2008], we improve upon Theorem 4.10 by demonstrating that, under a nonfocusing condition, strong singularities can only propagate along geometric GBBs. The key technical result concerns the propagation of Lagrangian regularity with respect to the incoming and outgoing Lagrangian submanifolds, localized near geometric continuations of a GBB striking the

cone point. Using the coordinates $(\hat{h}, x, y, \xi, \eta)$ and the notation of (4-14), the incoming and outgoing Lagrangians are given by

$$\mathcal{F}_\bullet := \bigcup_{y_0 \in \partial X} \mathcal{F}_{\bullet, y_0}, \quad \bullet = I, O,$$

where

$$\begin{aligned} \mathcal{F}_{I, y_0} &:= \gamma_{I, y_0}((-x_0, 0)) = \{(0, x, y_0, -1, 0) : x < 2x_0\}, \\ \mathcal{F}_{O, y_0} &:= \gamma_{O, y_0}((-x_0, 0)) = \{(0, x, y_0, 1, 0) : x < 2x_0\}. \end{aligned} \tag{4-46}$$

(We are making the \hat{h} -coordinate, which was set to 0 in (4-14), explicit here.)

We shall first show that one can control the Lagrangian regularity of a solution u of $P_{h,z}u = f$, with sufficiently regular forcing f , near \mathcal{F}_{O, y_0} by propagating Lagrangian regularity from the union of all $\mathcal{F}_{I, y'}$, with y' at distance π from y_0 , into ∂X and then within ∂X to $\mathcal{F}_{O, y_0} \cap x^{-1}(0)$. Localization within the radial sets $\mathcal{R}_{in/out}$ requires a more careful choice of commutants compared to the symbolic part of the proof of Theorem 4.10, and the extra Lagrangian regularity is captured using test modules, as introduced in [Hassell et al. 2008] and used for this purpose in [Melrose et al. 2008; 2013]; see also [Haber and Vasy 2013]. (Test modules also feature prominently in [Baskin et al. 2015; 2018; Gell-Redman et al. 2020].) Fix $x_0 < x_1 < x_2 < x_3 < 2x_0$ and cutoffs

$$\chi_j \in C_c^\infty([0, x_j]), \quad \chi_j \equiv 1 \quad \text{on } [0, x_{j-1}], \quad j = 1, 2, 3.$$

Mirroring [Melrose et al. 2008, Definition 4.2], we then introduce:

Definition 4.17 (test module). Let $\mathcal{F} = \mathcal{F}_I \cup \mathcal{F}_O$. Define the $\Psi_{ch}^0(X)$ -module¹⁴

$$\mathcal{M} := \left\{ A \in \Psi_{ch}^{0,0,0,1}(X) : \text{supp } K_A \subset [0, 1)_h \times (\text{supp } \chi_2)^2, \left. {}^{ch}\sigma_0\left(\frac{h}{h+x}A\right)\right|_{\mathcal{F}} = 0 \right\}.$$

Denote by $\mathcal{M}^k \subset \Psi_{ch}^{0,0,0,k}(X)$ the set of finite linear combinations of up to k -fold products of elements of \mathcal{M} . If \mathcal{X} is a function space on which $\Psi_{ch}^0(X)$ acts continuously, we say that u has *Lagrangian regularity of order k relative to \mathcal{X}* if $\mathcal{M}^k u \subset \mathcal{X}$. We say that elements of the space $\mathcal{M}^k \mathcal{X}$ satisfy the *nonfocusing condition of degree k relative to \mathcal{X}* .

Since

$$\Psi_{ch}^{0,0,0,1}(X) = \left(\frac{h}{h+x}\right)^{-1} \Psi_{ch}^0(X),$$

regularity with respect to elements of \mathcal{M} means that the semiclassical order improves upon differentiation along suitable elements of $\Psi_{ch}^0(X)$. A concrete example of an element of \mathcal{M} in local coordinates is

$$\frac{h+x}{h} \left(\frac{h}{h+x} D_{y^j}\right) = D_{y^j}.$$

Lemma 4.18 (properties of \mathcal{M} ; see [Melrose et al. 2008, Lemma 4.4]). *The set \mathcal{M} is closed under commutators. Moreover, \mathcal{M} is finitely generated in the sense that there exist $A_1, \dots, A_N \in \Psi_{ch}^{0,0,0,1}(X)$*

¹⁴Recall that K_A denotes the Schwartz kernel of A .

with $\text{supp } K_{A_j} \in [0, 1)_h \times (\text{supp } \chi_3)^2$ so that, with $A_0 := I$, we have

$$\mathcal{M} = \left\{ A \in \Psi_{ch}^{0,0,0,1}(X) : \text{there exists } Q_j \in \Psi_{ch}^0(X) \text{ such that } A = \sum_{j=0}^N Q_j A_j \right\}.$$

Concretely, one can take A_N to have principal symbol

$$\left(\frac{h+x}{h} \right) \cdot {}^{ch}\sigma \left(\left(\frac{x}{x+h} \right)^2 P_{h,z} \right),$$

and one may take A_j , $1 \leq j \leq N-1$, to have principal symbol $((h+x)/h)a_j$, where $a_j \in C^\infty(\overline{{}^{ch}T^*X_{ch}})$ vanishes on \mathcal{F} and has differential da_j which at a point $\zeta \in \mathcal{R}_{in}$, resp. $\zeta \in \mathcal{R}_{out}$ lies in the unstable, resp. stable eigenspace of the linearization of H (as a vector field on ${}^{ch}T_{sf}^*X_{ch}$) at ζ .

Proof. Let $B = ((h+x)/h)B_0$, $C = ((h+x)/h)C_0 \in \mathcal{M}$. Denote the principal symbols of $B_0, C_0 \in \Psi_{ch}^0(X)$ by b_0, c_0 . We then have $[B, C] \in \Psi_{ch}^{-1,0,0,1}(X)$, and

$$d := {}^{ch}\sigma_0 \left(\frac{h}{h+x} i[B, C] \right) = \frac{h}{h+x} H_b \left(\frac{h+x}{h} c_0 \right) = \frac{h}{h+x} H_b \left(\frac{h+x}{h} \right) c_0 + H_b c_0.$$

But by (3-4), $H_b|_{\hat{h}=0}$ is a smooth b-vector field for $b \in S^{0,0,0,1}$; thus $d \in S^0(\overline{{}^{ch}T^*X_{ch}})$. Moreover, since \mathcal{F} is a Lagrangian submanifold, H_b is tangent to \mathcal{F} ; therefore, $H_b c_0 = 0$ on \mathcal{F} since $c_0|_{\mathcal{F}} = 0$, and thus $d|_{\mathcal{F}} = 0$ as well. This proves $[B, C] \in \mathcal{M}$.

Let us now work in local coordinates $(\hat{h}, x, y, \xi, \eta)$ in which the rescaled Hamilton vector field $H = \hat{h}^{-1} H_p$ of $P_{h,z}$ takes the form (4-9). The linearization of H at $\mathcal{R}_{out/in}$ as a vector field on ${}^{ch}T^*X_{ch}$ is (top sign for “in”, bottom sign for “out”)

$$\mp 2(x\partial_x - \hat{h}\partial_{\hat{h}} - \eta\partial_\eta) + 2k^{ij}\eta_i\partial_{y^j}, \tag{4-47}$$

which thus has eigenvalue ∓ 2 (with eigenvector dx), ± 2 (with eigenspace spanned by $d\hat{h}$ and $d\eta_j$), and 0 (with eigenspace spanned by $d\xi$ and $dy^j \pm k^{ij}d\eta_j$). Upon restriction to $\hat{h} = 0$, the same statements remain true except there is no contribution from $d\hat{h}$ anymore. Since \mathcal{F} is locally the joint zero set of $\eta^1, \dots, \eta^{n-1}$, and p , which have linearly independent differentials, every smooth function vanishing on \mathcal{F} can be written as a linear combination (with smooth coefficients) of p and η_j . Thus, we may take quantizations of $\hat{h}^{-1}\eta_j$ for the operators A_j in local coordinates. The full collection of A_j can be defined using a partition of unity. □

The fact that \mathcal{M} is a $\Psi_{ch}^0(X)$ -module and a Lie algebra implies that

$$\mathcal{M}^k = \left\{ \sum_{|\alpha| \leq k} Q_\alpha A^\alpha : Q_\alpha \in \Psi_{ch}^0(X) \right\}, \quad A^\alpha := \prod_{i=1}^N A_i^{\alpha_i}, \tag{4-48}$$

where $\alpha = (\alpha_1, \dots, \alpha_N) \in \mathbb{N}_0^N$. Modulo $\Psi_{ch}^0(X)$, the operator A_N is a multiple of $P_{h,z}$; therefore, regularity of solutions u of $P_{h,z}u = f$, with f having Lagrangian regularity of order k , under application of an element $Q_\alpha \prod_{i=1}^N A_i^{\alpha_i} \in \mathcal{M}^k$ with $\alpha_N > 0$ is automatic once Lagrangian regularity of order $k-1$ has been established. In order to prove regularity of solutions of $P_{h,z}u = f$ under application of A_j , $1 \leq j \leq N-1$, we need to control the commutators of $P_{h,z}$ with the A_j chosen in Lemma 4.18:

Lemma 4.19 (commutators; see [Melrose et al. 2008, Lemma 4.5]). *With the A_j chosen as in Lemma 4.18, we have, for $j = 1, \dots, N - 1$,*

$$i[P_{h,z}, A_j] = \sum_{k=0}^N C_{jk} A_k, \quad C_{jk} \in \Psi_{ch}^{1,2,0,-1}(X),$$

and ${}^{ch}\sigma_{1,2,0,-1}(C_{jk})|_{\mathcal{F} \cap x^{-1}(0)} = 0$ for $k \neq 0$.

Proof. Denote by a_j the principal symbol of $\hat{h}A_j$ for $j = 1, \dots, N - 1$, so ${}^{ch}\sigma(A_j) = \hat{h}^{-1}a_j$. Since $P_{h,z} \in \Psi_{ch}^{2,2,0,0}(X)$, we have $i[P_{h,z}, A_j] \in \Psi_{ch}^{1,2,0,0}(X)$, with principal symbol at sf given by $\hat{h}H(\hat{h}^{-1}a_j)$ in the notation used in (4-9). It thus suffices to prove the existence of $c_{jk} \in S^{1,2,0,0}(\overline{{}^{ch}T^*X_{ch}})$ such that near $\hat{h} = 0$,

$$\hat{h}H(\hat{h}^{-1}a_j) = \sum_{k=1}^N \hat{h}c_{jk}\hat{h}^{-1}a_k, \quad c_{jk}|_{\mathcal{F} \cap x^{-1}(0)} = 0 \quad (k \neq 0); \tag{4-49}$$

indeed, if $C_{jk} \in \Psi_{ch}^{1,2,0,-1}(X)$ is a quantization of $\hat{h}c_{jk}$ times a cutoff to a neighborhood of sf, then (4-49) implies that

$$C_{j0} := i[P_{h,z}, A_j] - \sum_{k=1}^N C_{jk} A_k \in \Psi_{ch}^{0,2,0,-1}(X).$$

In order to verify (4-49), we note that the left-hand side equals $\hat{h}^{-1}(\hat{h}Ha_j - a_jH\hat{h})$; but since at $\mathcal{F} \cap x^{-1}(0)$, the differentials da_j and $d\hat{h}$ are eigenvectors of the linearization of H with the same eigenvalue, as discussed after (4-47), this vanishes quadratically at $\mathcal{F} \cap x^{-1}(0)$, completing the proof. \square

We are now ready to propagate Lagrangian regularity through the radial sets. For $s, l, \alpha, b \in \mathbb{R}$ and $k \in \mathbb{N}_0$, and using the notation (4-48), let

$$H_{c,h}^{s,l,\alpha,b;k}(X) := \{u \in H_{c,h}^{s,l,\alpha,b}(X) : A^\alpha u \in H_{c,h}^{s,l,\alpha,b}(X) \text{ for all } |\alpha| \leq k\}.$$

We recall that we will only encounter distributions on X with compact support, justifying the convenient, albeit slightly imprecise, notation here.

Proposition 4.20 (microlocalized propagation near the radial sets). *Let $s, l, \alpha, b \in \mathbb{R}$. Let $B, E, G \in \Psi_{ch}^0(X)$ denote operators with Schwartz kernels supported in $[0, 1)_h \times (\text{supp } \chi_1)^2$. Recall the quantities $r_{\text{in/out}}$ from Definition 4.7.*

(1) (propagation into \mathcal{R}_{in}) *Suppose that all backward integral curves of H starting in $\Sigma \cap \text{WF}'_{ch}(B)$ either tend to a subset $\mathcal{S} \subset \mathcal{R}_{\text{in}}$ or enter $\text{Ell}'_{ch}(E)$ in finite time while remaining inside $\text{Ell}_{ch}(G)$; and suppose that, for all incoming null-bicharacteristics $\gamma_{I,y_0} : (-x_0, 0) \rightarrow \Sigma$ with $\gamma_{I,y_0}(0) \in \mathcal{S}$, there exists $s \in (-x_0, 0)$ (depending on y_0) such that $\gamma_{I,y_0}((s, 0]) \subset \text{Ell}_{ch}(G)$ and $\gamma_{I,y_0}(s) \in \text{Ell}_{ch}(E)$. Under the condition $b - \alpha > r_{\text{in}}$, we then have*

$$\|Bu\|_{H_{c,h}^{s,l,\alpha,b;k}(X)} \leq C(\|GP_{h,z}u\|_{H_{c,h}^{s-2,l-2,\alpha,b+1;k}(X)} + \|Eu\|_{H_{c,h}^{-N,l,\alpha,b;k}(X)} + \|\chi_2u\|_{H_{c,h}^{-N,l,\alpha,b-1/2}(X)}). \tag{4-50}$$

(2) (propagation out of \mathcal{R}_{out}) *Suppose that all backward integral curves of H starting in $\Sigma \cap \text{WF}'_{ch}(B)$ either tend to a subset $\mathcal{S} \subset \mathcal{R}_{\text{out}}$ or enter $\text{WF}'_{ch}(E)$ in finite time while remaining inside $\text{Ell}_{ch}(G)$. Suppose moreover that $\mathcal{S} \subset \text{Ell}_{ch}(G)$, and that, for every integral curve $\gamma \subset \Sigma \cap x^{-1}(0) \setminus \mathcal{R}_{\text{out}}$ of H*

with $\lim_{s \rightarrow \infty} \gamma(s) \in \mathcal{S}$, there exists s so that $\gamma((s, \infty)) \subset \text{Ell}_{c\hat{h}}(G)$ and $\gamma(s) \in \text{Ell}_{c\hat{h}}(E)$. Then the estimate (4-50) holds under the condition $b - \alpha < r_{\text{out}}$.

Proof. We begin with the proof of part (1). By compactness of $\mathcal{R}_{\text{in/out}}$ and since $\text{Ell}_{c\hat{h}}$ is open, it suffices to prove microlocal estimates near a single point $\zeta_0 \in \mathcal{R}_{\text{in}}$, which in the coordinate system $(\hat{h}, x, y, \xi, \eta)$ used in (4-46) has coordinates $\zeta_0 = (0, 0, y_0, -1, 0)$.

Now, restricted to $x = \hat{h} = 0$ and writing $k = k(y, \eta)$ for the dual metric function of the metric $k(0)$ on ∂X in local coordinates, we have

$$H = -2\xi\eta\partial_\eta + 2|\eta|^2\partial_\xi + (\partial_\eta k)\partial_y - (\partial_y k)\partial_\eta.$$

Following [Melrose and Zworski 1996, Lemma 2], introducing $|\eta|$, $\hat{\eta} = \eta/|\eta|$, one has

$$\partial_s y = (\partial_\eta k)(y, \hat{\eta})|\eta|, \quad \partial_s \hat{\eta} = -(\partial_y k)(y, \hat{\eta})|\eta|$$

along H-integral curves; reparametrizing to $t = t(s)$ satisfying $t' = 2|\eta|$, one thus obtains

$$\partial_t y = \frac{1}{2}(\partial_\eta k)(y, \hat{\eta}), \quad \partial_t \hat{\eta} = -\frac{1}{2}(\partial_y k)(y, \hat{\eta}), \quad \partial_t |\eta| = -\xi, \quad \partial_t \xi = |\eta|.$$

Thus, $\xi(t) = a \cos(t + \varphi_0)$ and $|\eta(t)| = a \sin(t + \varphi_0)$, where $a = \sqrt{\xi^2 + |\eta|^2}$ is constant, and $\varphi_0 \in [0, \pi]$. Therefore, the function Υ assigning to (y, ξ, η) near $(y_0, -1, 0)$ the limiting point along the backward H-integral curve is given by evaluation at $t = -\varphi_0$, so

$$\Upsilon(y, \xi, \eta) = \left(\exp_y \left(\left(-\arccos \frac{\xi}{\sqrt{\xi^2 + |\eta|^2}} \right) \frac{\eta}{|\eta|} \right), -1, 0 \right).$$

In particular, Υ is smooth, and $H\Upsilon = 0$ at $\hat{h} = x = 0$. Extending Υ to a smooth function in a neighborhood of $x = \hat{h} = 0$, with values in $\mathbb{R}^{n-1} \times \mathbb{R} \times \mathbb{R}^{n-1}$, we thus have $H\Upsilon = \mathcal{O}(x)$ at $\hat{h} = 0$. Since $x^{-1}Hx = -2$ at ζ_0 , we can choose C so that in any sufficiently small neighborhood \mathcal{V} of ζ_0 ,

$$H(|\Upsilon - \zeta_0|^2 - Cx) \geq x > 0 \quad \text{in } \mathcal{V}. \tag{4-51}$$

Fix now cutoffs $\chi_{\mathcal{S}}, \chi_\partial, \chi_{\text{sf}}, \chi_{\mathcal{F}}, \chi_\Sigma \in C_c^\infty([0, 1])$, identically 1 near 0, with nonpositive derivative and with $\sqrt{-\chi_\bullet \chi'_\bullet} \in C^\infty$, and consider the commutant

$$\check{a} = \hat{h}^{-b-\frac{1}{2}} x^{-\alpha} \chi_\partial \left(\frac{x}{\delta} \right) \chi_{\text{sf}} \left(\frac{\hat{h}}{\delta} \right) \chi_{\mathcal{F}} \left(\delta^{-1} \sum_{j=1}^{N-1} a_j^2 \right) \chi_\Sigma \left(\frac{p^2}{\delta} \right) \chi_{\mathcal{S}}(\delta^{-1}(|\Upsilon - \zeta_0|^2 - Cx)),$$

where $\delta > 0$ controls the size of $\text{supp } \check{a}$. We now proceed as in the first step of the proof of Theorem 4.10. Thus, in the symbol (4-32) of the commutator appearing in (4-31), and specifically in the term $2\hat{h}\check{a}\check{H}\check{a}$, the main contribution near ζ_0 arises from differentiation of the weights (and then the subprincipal symbol of $P_{h,z}$ enters in the threshold condition on $b - \alpha$ as there), giving a negative multiple of $\hat{h}^{-2b} x^{-2\alpha}$. Differentiation of $\chi_{\mathcal{F}}$ gives a term of the same sign, namely a negative square, since $\sum a_j^2$ is a local quadratic defining function of \mathcal{R}_{in} inside of $\Sigma \cap x^{-1}(0)$. In view of (4-51), differentiation of $\chi_{\mathcal{S}}$ produces x times the negative of a square, thus another term with sign matching that of the main term. Derivatives falling on χ_∂ produce a nonnegative square, corresponding to the a priori control required along $\gamma_{I,y}$ for y near y_0 , at $x \sim \delta$. Finally, differentiation of χ_Σ produces a term vanishing near Σ which thus can

be controlled by elliptic regularity, and differentiation of χ_{sf} produces a semiclassically trivial (namely, vanishing near $\hat{h} = 0$) term. We can then proceed as in (4-33), obtaining the desired propagation estimate.

For $k \geq 1$, we argue as in the proof of [Baskin et al. 2015, Proposition 4.4]: rather than using $\check{A} = \text{Op}_{c,h}(\check{a})$ as the commutant, we use (in the notation (4-48)) the vector of ps.d.o.s $(\check{A}A^\alpha)_{\alpha \in \mathcal{I}}$, where $\mathcal{I} \subset \mathbb{N}_0^N$ consists of all $\alpha \in \mathbb{N}_0^N$ with $|\alpha| = k$ and $\alpha_0 = \alpha_N = 0$. The main term of the commutator arises from \check{A} as before; the new contributions, from commutators of $P_{h,z}$ with a factor A_j , can be expanded as in Lemma 4.18, and those which have the maximal number of module factors A_l , $1 \leq l \leq N - 1$, can be absorbed into this main term due to the vanishing property of the C_{jk} in Lemma 4.18. Thus, one can control k module derivatives of u in a neighborhood of ζ_0 provided one has control of $k - 1$ module derivatives in a slightly bigger neighborhood. Thus, one obtains the estimate (4-50) inductively.

The proof of part (2) is completely analogous; one now takes Υ at $x = \hat{h} = 0$ to be the limiting point along forward H-integral curves. □

Note that for any $\zeta \in \overline{c^h T_{sf}^* X_{ch}} \setminus \mathcal{F}$, there exists an element $A \in \mathcal{M}$ which is elliptic at ζ ; hence microlocally near such ζ , membership in $H_{c,h}^{s,l,\alpha,b;k}$ is equivalent to membership in $H_{c,h}^{s,l,\alpha,b+k}$.¹⁵ In particular, in $\Sigma \cap x^{-1}(0)$ but away from the radial sets, the propagation of $H_{c,h}^{s,l,\alpha,b;k}$ regularity is equivalent to the standard (real-principal-type) propagation of $H_{c,h}^{s,l,\alpha,b+k}$ regularity. One can thus concatenate the radial point estimates of Proposition 4.20 with such real-principal-type estimates. To state this succinctly, we introduce:

Definition 4.21 (integral curves connecting the radial sets). (1) For a point $y \in \partial X$, denote by $\Gamma^\rightarrow(y) \subset C^0([0, \pi]; \Sigma)$ the set of integral curves of H inside $\Sigma \cap x^{-1}(0)$, smoothly reparametrized to uniformly continuous curves $\gamma : (0, \pi) \rightarrow \Sigma \cap x^{-1}(0)$, which satisfy $\gamma(\pi) = (0, 0, y, 1, 0) \in \mathcal{R}_{out}$ and $\gamma(0) \in \mathcal{R}_{in}$. Denoting by $\Pi : \Sigma \cap x^{-1}(0) \rightarrow \partial X$ the projection to the base, define the set of starting points of such curves by

$$\mathcal{Y}^\rightarrow(y) = \{\Pi(\gamma(0)) : \gamma \in \Gamma^\rightarrow(y)\}.$$

(2) We call a continuous curve $\gamma : I \rightarrow \Sigma$ a *resolved GBB* if it is either an integral curve of $h^{-1}H_p$ disjoint from $x^{-1}(0)$, or otherwise if, for some $y \in \partial X$ and $y_0 \in \mathcal{Y}^\rightarrow(y)$, the curve γ is the concatenation of γ_{I,y_0} , an element γ of $\Gamma^\rightarrow(y)$ with $\Pi(\gamma(0)) = y_0$, $\Pi(\gamma(\pi)) = y$, and the curve $\gamma_{O,y}$.

See Figure 12.

Corollary 4.22 (microlocalized propagation of Lagrangian regularity). *Let $s, l, \alpha \in \mathbb{R}$, $b \in C^\infty(\overline{c^h T_{sf}^* X_{ch}})$ and $\chi \in C_c^\infty(X)$ with $\text{supp } \chi \subset \text{supp } \chi_1$ be as in Theorem 4.10, with $b - \alpha$ satisfying the monotonicity and threshold conditions stated there. Let $k \in \mathbb{N}_0$. Let $B, E, G \in \Psi_{ch}^0(X)$, with Schwartz kernels supported in $[0, 1]_h \times (\chi^{-1}(1) \times \chi^{-1}(1))$. Suppose that all backward resolved GBBs starting at a point in $\text{WF}_{ch}^l(B)$ reach $\text{Ell}_{ch}(E)$ in finite time while remaining in $\text{Ell}_{ch}(G)$. Then the estimate (4-50) holds.*

A dualization argument gives the propagation of the nonfocusing condition through ∂X . The simplest setting uses the modification of $P_{h,z}$ via extension to an operator $P'_{h,z}$ on compact manifold $X' \supset X$ and

¹⁵That is, for $B, \tilde{B} \in \Psi_{ch}^0(X)$ with $\text{WF}'_{ch}(B) \subset \text{Ell}_{ch}(\tilde{B}) \setminus \mathcal{F}$, one has $\|Bu\|_{H_{c,h}^{s,l,\alpha,b+k}} \lesssim \|\tilde{B}u\|_{H_{c,h}^{s,l,\alpha,b;k}} + \|u\|_{H_{c,h}^{-N,l,\alpha,-N}}$.




Figure 12. Illustration of a part of the characteristic set, with the Lagrangians \mathcal{F}_I and \mathcal{F}_O in red, and a resolved GBB in blue.

the inclusion of a complex absorbing term $Q \in \Psi_h^{-\infty}((X')^\circ)$ as in Section 4E, resulting in the operator

$$\mathcal{P}_{h,z} = P'_{h,z} - iQ$$

in (4-42). (This requires ∂X to be null-cobordant; if this is not true, one can use the modification described in Remark 4.16.) Recall that the Schwartz kernel of Q has empty intersection with $x^{-1}([0, cx_0]) \times x^{-1}([0, cx_0])$, where $0 < c \ll \frac{1}{2}$. We shall use the notation of Proposition 4.15.

Theorem 4.23 (diffractive improvement). *Let s, l, α, b be as in the statement of Theorem 4.10 (for the operator $P_{h,z}$). Let $E, G \in \Psi_{ch}^0(X')$ be such that all forward resolved GBBs starting at a point in $\text{WF}'_{ch}(E) \subset \text{Ell}_{ch}(G)$ remain in $\text{Ell}_{ch}(G)$ until they enter $\text{Ell}_h(Q)$. Let $f_+ \in H_{c,h}^{s-2, l-2, \alpha, b+1}(X')$, $f_- \in \mathcal{M}^k H_{c,h}^{s-2, l-2, \alpha, b+1}(X')$ be such that $\text{supp } f_\pm \subset x^{-1}([0, cx_0])$. Then the solution u of*

$$\mathcal{P}_{h,z}u = f := f_+ + Ef_-$$

can be written in the form

$$u = u_+ + Gu_-, \quad u_+ \in H_{c,h}^{s, l, \alpha, b}(X'), \quad u_- \in \mathcal{M}^k H_{c,h}^{s, l, \alpha, b}(X').$$

Note that on the scale of semiclassical cone Sobolev spaces, we have $Ef_- \in H_{c,h}^{s-2, l-2, \alpha, b'+1}$ with $b' = b - k$, but typically Ef_- is no better than this. Thus, Theorem 4.23 (for $f_+ = 0$ for concreteness) implies that the strong semiclassical singularities of u resulting from the forcing term Ef_- only propagate along geometric GBBs (resulting in the term Gu_-), whereas microlocally away from these, u has $H_{c,h}^{s, l, \alpha, b}$ -regularity.

In a simple case, a formulation of Theorem 4.23 which highlights regularity rather than singularities reads as follows: Fix $y_0 \in \partial X$, and define the set

$$K := \gamma_{O, y_0} \cup \bigcup_{\gamma \in \Gamma \rightarrow (y_0)} \gamma([0, \pi]) \cup \bigcup_{y \in \mathcal{Y} \rightarrow (y_0)} \gamma_{I, y}.$$

Thus, the quotient $K/(K \cap x^{-1}(0))$ contains the image of all backward geometric GBB continuing γ_{O, y_0} , and K in addition contains all curves inside of $\Sigma \cap x^{-1}(0)$ which connect an incoming base point y (at distance π from y_0) with the outgoing base point y_0 of geometric GBBs. Fixing any $E \in \Psi_{ch}^0(X')$ with $\text{WF}'_{ch}(E) \cap K = \emptyset$, there then exists $G \in \Psi_{ch}^0(X')$ with $\text{WF}'_{ch}(G) \cap K = \emptyset$ which satisfies the

conditions of Theorem 4.23. Thus, if f satisfies the nonfocusing condition (of some degree k) relative to $H_{c,h}^{s-2,l-2,\alpha,b+1}$, and with f microlocally near K lying in $H_{c,h}^{s-2,l-2,\alpha,b+1}$ (thus f in particular does not have strong singularities along the incoming directions $\gamma_{I,y}$), then the semiclassical wave u forced by f lies in $H_{c,h}^{s,l,\alpha,b}$ microlocally near K (thus u in particular does not have a strong singularity along γ_{O,y_0}).

Proof of Theorem 4.23. As follows from Proposition 4.15 by taking adjoints (or directly from the proof of Proposition 4.15), the adjoint $\mathcal{P}_{h,z}^*$ is invertible, and

$$(\mathcal{P}_{h,z}^*)^{-1} : H_{c,h}^{-s,-l,-\alpha,-b} \rightarrow H_{c,h}^{-s+2,-l+2,-\alpha,-b-1}$$

is uniformly bounded. We now apply a backward propagation version of Corollary 4.22 to $P_{h,z}^*$: For E^*, G^* the adjoints of the operators E, G in the statement of the theorem, and for $B^* \in \Psi_{ch}^0(X')$ so that all forward resolved GBBs starting at a point in $WF'_{ch}(E^*)$ remain in $\text{Ell}_{ch}(G^*)$ until they enter $\text{Ell}_{ch}(B^*)$, we have

$$\|E^*v\|_{H_{c,h}^{-s+2,-l+2,-\alpha,-b-1;k}} \leq C(\|G^*P_{h,z}^*v\|_{H_{c,h}^{-s,-l,-\alpha,-b;k}} + \|B^*v\|_{H_{c,h}^{-s+2,-l+2,-\alpha,-b-1;k}} + \|\chi u\|_{H_{c,h}^{-N,-l+2,-\alpha,-b-3/2}})$$

for any $k \in \mathbb{N}_0$. In particular, we may take B^* so that all forward null-bicharacteristics of $P_{h,z}$ starting in $WF'_{ch}(B^*)$ miss the cone point and enter $\text{Ell}_{ch}(Q^*)$ in finite time. The term B^*u is then automatically controlled for solutions of $\mathcal{P}_{h,z}^*v = w$ when $G^*w \in H_{c,h}^{-s,-l,-\alpha,-b;k}$ by elliptic regularity (on $\text{Ell}_{ch}(Q)$) and real-principal-type propagation (along the backward null-bicharacteristic flow) with complex absorption. We conclude that

$$\begin{aligned} (\mathcal{P}_{h,z}^*)^{-1} : \{w \in H_{c,h}^{-s,-l,-\alpha,-b} : G^*w \in H_{c,h}^{-s,-l,-\alpha,-b;k}\} \\ \rightarrow \{v \in H_{c,h}^{-s+2,-l+2,-\alpha,-b-1} : E^*v \in H_{c,h}^{-s+2,-l+2,-\alpha,-b-1;k}\}. \end{aligned}$$

Upon taking adjoints (see also [Melrose et al. 2013, Appendix A]), this implies that

$$\mathcal{P}_{h,z}^{-1} : H_{c,h}^{s-2,l-2,\alpha,b+1} + E(\mathcal{M}^k H_{c,h}^{s-2,l-2,\alpha,b+1}) \rightarrow H_{c,h}^{s,l,\alpha,b} + G(\mathcal{M}^k H_{c,h}^{s,l,\alpha,b})$$

is a bounded map. □

Remark 4.24 (second microlocalization at \mathcal{F}). A sharper approach would be to second microlocalize at \mathcal{F}_I and \mathcal{F}_O , thus cleanly decoupling the semiclassical orders at \mathcal{F}_I and \mathcal{F}_O (subject to threshold conditions at the radial sets) and the semiclassical order away from \mathcal{F} ; this would allow for a unified treatment of Lagrangian and nonfocusing spaces and thus for a direct proof of Theorem 4.23. We leave such refinements for future work. We note that second microlocalization in the semiclassical setting was studied in [Sjöstrand and Zworski 2007; Vasy and Wunsch 2009] following [Bony 1986]; a second microlocal refinement (at the outgoing radial set) for the scattering theory of the corresponding normal operator was recently obtained in [Vasy 2021c].

5. Applications

We now present applications of the propagation estimates proved in Section 4. First, we discuss the familiar geometric case of $h^2\Delta_g - 1$ in Section 5A, where we can moreover prove a result sharpening both Theorem 4.10 and the propagation results of [Baskin and Marzuola 2022]. We discuss high-frequency

scattering by inverse square potentials on Euclidean space in Section 5B, and high-frequency scattering for the Dirac–Coulomb equation in Section 5C.

5A. Propagation estimates for conic Laplacians. For a conic metric g as in (4-2) on the manifold $X = [0, 2x_0)_x \times Y$ of dimension $n = \dim X \geq 3$, we consider

$$P_{h,z} = h^2 \Delta_g - z, \quad |z - 1| < Ch.$$

We fix the volume density $\mu = |dg|$ on X .

Lemma 5.1 (admissibility, thresholds, invertibility). *The operator $P_{h,z}$ is admissible in the sense of Definition 4.1, with threshold quantities $r_{\text{in}} = -\frac{1}{2}$ and $r_{\text{out}} = -\frac{1}{2}$ (see Definition 4.7). Moreover, the normal operator $N(P) = \Delta_{\hat{g}} - 1$, with \hat{g} given in (4-4), is invertible at weight l (in the sense of Definition 4.6(3)) for all*

$$l \in \left(1 - \frac{n-2}{2}, 1 + \frac{n-2}{2} \right). \tag{5-1}$$

Proof. Only the final statement is nontrivial. In the notation (4-17), and passing to a spectral decomposition of $\Delta_{k(0)}$ whose eigenvalues we denote by $0 \leq \lambda_j^2$, $j = 0, 1, 2, \dots$, one finds that $\lambda \in \text{spec}_b(N(P))$ if and only if there exists j with $\lambda^2 - i(n-2)\lambda + \lambda_j^2 = 0$, so

$$\text{spec}_b(N(P)) = \left\{ i \left(\frac{n-2}{2} \pm \sqrt{\left(\frac{n-2}{2} \right)^2 + \lambda_j^2} \right) : j = 0, 1, 2, \dots \right\}.$$

Therefore, the complement of the set Λ defined in (4-18) contains $(-n+2, 0)$. As noted in Remark 4.9, the invertibility of $N(P)$ at weight l is independent of the choice of l inside the shifted interval

$$\frac{n}{2} + (-n+2, 0) = \left(1 - \frac{n-2}{2}, 1 + \frac{n-2}{2} \right).$$

The choice $l = 1$ is particularly natural, as the space $H_{b,\text{sc}}^{1,1,0}(\text{tf}; \hat{\mu})$ is the quadratic form domain of $\Delta_{\hat{g}}$ (as follows from Hardy’s inequality). The invertibility of $N(P)$ at weight $l = 1$ is then equivalent to the limiting absorption principle for the exact conic metric \hat{g} , the proof of which is a standard application of a boundary-pairing argument [Melrose 1995, §2.3] and unique continuation at infinity. See Lemma 5.10 below for a proof in a more general setting. □

As a consequence, we may apply Theorem 4.10 for l in the range (5-1), any value of $s \in \mathbb{R}$, $\alpha = 0$, and variable orders b satisfying in particular $b > -\frac{1}{2}$ at \mathcal{R}_{in} , $b < -\frac{1}{2}$ at \mathcal{R}_{out} , and we may arrange that $|b - (-\frac{1}{2})| < \epsilon$ for any fixed $\epsilon > 0$. Packaged in the form of Proposition 4.15 using complex absorption, we thus have, using the volume density $|dg|$ near ∂X ,

$$\|u\|_{H_{c,h}^{s,l,0,b}} \leq C \| \mathcal{P}_{h,z} u \|_{H_{c,h}^{s-2,l-2,0,b+1}}; \tag{5-2}$$

this estimate is sharp in the sense explained after the statement of Theorem 4.10. Lossy estimates on constant order spaces are given by

$$\begin{aligned} \|(x+h)^{-\frac{1}{2}-\epsilon} u\|_{H_{c,h}^{s,l,0,0}} &\sim h^{-\frac{1}{2}-\epsilon} \|u\|_{H_{c,h}^{s,l,0,-1/2-\epsilon}} \\ &\lesssim h^{-\frac{1}{2}-\epsilon} \| \mathcal{P}_{h,z} u \|_{H_{c,h}^{s-2,l-2,0,1/2+\epsilon}} \sim h^{-1-2\epsilon} \|(x+h)^{\frac{1}{2}+\epsilon} \mathcal{P}_{h,z} u\|_{H_{c,h}^{s-2,l-2,0,0}}. \end{aligned}$$

In the special case $s = l$, and recalling from [Hintz 2022, Theorem 6.3] that the domain

$$\mathcal{D}_h^l = \mathcal{D}((h^2 \Delta_{g'} + 1)^{l/2})$$

of the $(l/2)$ -th power of the Friedrichs extension of the conic Laplacian $h^2 \Delta_{g'} + 1$ is equal to $H_{c,h}^{l,l,0,0}$ in present notation, this gives:

Proposition 5.2 (constant-order estimates). *In the above setting and with l as in (5-1), we have for all $\epsilon > 0$ the estimates*

$$\|(x + h)^{-\frac{1}{2}-\epsilon} u\|_{\mathcal{D}_h^l} \leq C_\epsilon h^{-1-2\epsilon} \|(x + h)^{\frac{1}{2}+\epsilon} \mathcal{P}_{h,z} u\|_{\mathcal{D}_h^{l-2}}, \tag{5-3a}$$

$$\|u\|_{\mathcal{D}_h^l} \leq C_\epsilon h^{-1-2\epsilon} \|\mathcal{P}_{h,z} u\|_{\mathcal{D}_h^{l-2}}, \tag{5-3b}$$

as well as more general estimates with \mathcal{D}_h^l and \mathcal{D}_h^{l-2} replaced by $H_{c,h}^{s,l,0,0}$ and $H_{c,h}^{s-2,l-2,0,0}$.

The estimate (5-3b) is an immediate consequence of (5-3a). We recall that in the case $l = 1$, the (arbitrarily small) 2ϵ -loss in (5-3b) can be removed, as shown in the semiclassical cone setting by Baskin and Marzuola [2022] following arguments by Melrose, Wunsch, and Vasy [Melrose and Wunsch 2004; Melrose et al. 2008]; in the full range of weights l considered here, a lossless estimate was obtained by the author in [Hintz 2022, §6.2] via reduction to the case $l = 1$ via conjugation by $(1 + h^2 \Delta_{g'})^{(l-1)/2}$ and reduction to the case $l = 1$. On the other hand, the estimate (5-3b), even for $\epsilon = 0$, loses a full order at tf compared to the sharper estimate (5-3a).

Remark 5.3 (limiting absorption principle). The $h^{-1-2\epsilon}$ loss in Proposition 5.2 is familiar from (and essentially arises from) the loss of slightly more than one power of $\langle z \rangle$ in the limiting absorption principle

$$(\Delta - 1 \pm i0)^{-1} : \langle z \rangle^{-\frac{1}{2}-\epsilon} L^2(\mathbb{R}^n) \rightarrow \langle z \rangle^{\frac{1}{2}+\epsilon} H^2(\mathbb{R}^n)$$

on Euclidean space, which is a consequence of a sharp variable-order estimate akin to (5-2); see Lemma 4.8.

A natural question is whether one can prove an estimate which removes both the ϵ -loss of (5-3b) while retaining the lossless character of (5-3a) (or (5-2)) at tf . We answer this in the affirmative:

Theorem 5.4 (sharp propagation estimate). *Consider a conic manifold (X, g) as in (4-1)–(4-2) and with $\dim X \geq 3$. Let $P_{h,z} = h^2 \Delta_g - z$, $|z - 1| < Ch$. Denote the characteristic set of $P_{h,z}$ by $\Sigma \subset {}^{ch}T_{\text{sf}}^* X_{ch}$; see (4-8). Let $\chi, \tilde{\chi} \in C_c^\infty(X)$, with $\tilde{\chi} \equiv 1$ near $\text{supp } \chi$, and $E \in \Psi_{ch}^{-\infty}(X)$. Suppose that all backward GBB from $\Sigma \cap \text{supp } \chi$ enter $\text{Ell}_{ch}(E)$ in finite time while remaining inside $\text{supp } \tilde{\chi}$. Then, for any $s, N \in \mathbb{R}$, we have an estimate*

$$\|\chi u\|_{H_{c,h}^{s,1,0,0}(X)} \leq C(\|\tilde{\chi} P_{h,z} u\|_{H_{c,h}^{s-2,-1,0,1}(X)} + \|Eu\|_{H_{c,h}^{-N,-N,-N,0}(X)} + h^{\frac{1}{2}} \|\tilde{\chi} u\|_{H_{c,h}^{-N,-N,0,0}(X)}). \tag{5-4}$$

This holds more generally if the first two norms above are replaced by $\|\chi u\|_{H_{cb,h}^{s,s',1,0,0}}$ and $\|\tilde{\chi} P_{h,z} u\|_{H_{cb,h}^{s-2,s'-2,-1,0,1}}$, with $s' \in \mathbb{R}$ arbitrary. Taking $s = 1, N = 0$ in (5-4) gives

$$\|\chi u\|_{\mathcal{D}_h^1} \lesssim h^{-1} \|(x + h) \tilde{\chi} P_{h,z} u\|_{\mathcal{D}_h^{-1}} + \|Eu\|_{L^2} + h^{\frac{1}{2}} \|\tilde{\chi} u\|_{L^2}, \tag{5-5}$$

and upon adding complex absorption as in Section 4E and (4-42), we have

$$\|u\|_{\mathcal{D}_h^1} = \|u\|_{H_{c,h}^{1,1,0,0}} \lesssim h^{-1} \|(x + h) \mathcal{P}_{h,z} u\|_{\mathcal{D}_h^{-1}} = \|\mathcal{P}_{h,z} u\|_{H_{c,h}^{-1,-1,0,1}}. \tag{5-6}$$

For comparison, the h -lossless version of (5-3b) for $l = 1$ reads

$$\|u\|_{H_{c,h}^{1,1,0,0}} \lesssim h^{-1} \|\mathcal{P}_{h,z}u\|_{H_{c,h}^{-1,-1,0,0}} = \|\mathcal{P}_{h,z}u\|_{H_{c,h}^{-1,-1,1,1}}, \tag{5-7}$$

which is weaker than Theorem 5.4 in that the required control on $\mathcal{P}_{h,z}$ at tf is one order stronger than in the theorem.

As discussed after Theorem 4.10, the estimate (5-4) is sharp in the sense that the relative orders on u on the left and $\mathcal{P}_{h,z}u$ on the right cannot be improved, but here the semiclassical order remains fixed upon propagation through the cone point.

The proof of Theorem 5.4 uses the global positivity (as an operator) of a commutator on tf, reminiscent of proofs of similar lossless results in N -body scattering [Vasy 2000; 2001], as well as a splitting of u , using the functional calculus for $h^2\Delta_g$, into a part localized near the characteristic set and a part where $h^2\Delta_g - 1$ is elliptic and can be inverted by spectral theory. The following technical result is proved at the end of this section.

Lemma 5.5 (functional calculus). *Let $\phi \in C_c^\infty(\mathbb{R})$. Then, for all $\epsilon > 0$,*

$$\phi(h^2\Delta_g) \in \mathcal{A}^{-\epsilon, n-\epsilon, n-\epsilon, 0, 0, \infty}(X_{ch}^2; {}^b\Omega^{\frac{1}{2}}), \tag{5-8}$$

where the orders of the conormal space refer to $\text{lb}_2, \text{ff}_2, \text{rb}_2, \text{tf}_2, \text{df}_2, \text{sf}_2 \subset X_{ch}^2$ in this order.

Remark 5.6 (dimension). We only study the case $\dim X \geq 3$ here. The methods used in [Melrose et al. 2008; Baskin and Marzuola 2022], which are based on quadratic forms, and also the methods in [Melrose and Wunsch 2004], work in the case $\dim X = 2$ as well. However, the identification of the quadratic form domain with a semiclassical cone Sobolev space fails in this case (see [Melrose and Wunsch 2004, equation (3.11)] for $h = 1$), which is why we do not consider it here.

Proof of Theorem 5.4. We present the proof in the case that $\mathcal{P}_{h,z}$ agrees with its normal operator, equivalently $\mathcal{P}_{h,z} = h^2\Delta_g - 1$ with $g = dx^2 + x^2k(y, dy)$ an exact conic metric. In the general case, the error terms arising from $\mathcal{P}_{h,z} - N(P) \in \Psi_{ch}^{2,-2,-1,0}(X)$ are handled easily; we leave the details to the reader. (In particular, since we shall use a global commutator argument which controls u at sf and tf in one fell swoop, there is no need for a delicate argument for the combination of the symbolic estimate at sf and a normal operator estimate at tf as in the end of the proof of Theorem 4.10.) We write $P \equiv \mathcal{P}_{h,z}$ for brevity.

Positive commutator argument: Define the operator

$$A := \frac{h}{2}(xD_x + (xD_x)^*) - 1 = hxD_x - \frac{inh}{2} - 1, \quad a := {}^{ch}\sigma(A) = x\xi - 1, \tag{5-9}$$

where we use the coordinates (3-3). This will be the main piece of the commutant in a positive commutator calculation, and it is in essence the key term both in the commutator argument of [Vasy 2008], as well as in the Mourre commutant [1980/81] in classical scattering theory. Let $\chi = \chi(x)$ be identically 1 near $\partial X = x^{-1}(0)$, with support in any prespecified neighborhood of ∂X , and so that $\chi' \leq 0$, $\sqrt{-\chi\chi'} \in C^\infty$, and so that $a < 0$ has a constant (negative) sign on $\Sigma \cap \text{supp } \chi$; arranging the latter property is what the constant term in (5-9) is for. We then consider the operator

$$\tilde{A} := \chi A \chi,$$

and estimate in two different ways the expression

$$2h^{-1} \operatorname{Im}\langle Pu, \tilde{A}u \rangle = \left\langle \frac{i}{h}[P, \tilde{A}]u, u \right\rangle. \tag{5-10}$$

Consider first the commutator term. Since $h^2\Delta_g$ is homogeneous of degree -2 , we have $(i/h)[P, A] = -[x\partial_x, P] = 2h^2\Delta_g$, which is the crucial global positive commutator. Therefore,

$$\frac{i}{h}[P, \chi A\chi] = 2\chi h^2\Delta_g\chi + \frac{i}{h}(\chi A[P, \chi] + [P, \chi]A\chi).$$

The contribution of the first term to the right-hand side of (5-10) is

$$2\|h\nabla_g(\chi u)\|^2,$$

where we write $\|\cdot\| \equiv \|\cdot\|_{L^2}$. The second term on the other hand consists of operators with coefficients supported strictly away from $x = 0$. It suffices to merely capture its principal symbol, which by (4-9) is $2\chi ah^{-1}H_p\chi = ax^{-1}H(\chi^2) = 4a\xi\chi\chi'$; near incoming directions, where $\xi < 0$, this is negative, whereas near outgoing directions, where $\xi > 0$, this is positive and thus has a sign matching that of the above main commutator term. For a suitable microlocal cutoff $E \in \Psi_{ch}^0(X)$ which is elliptic on Σ in the region $\xi < \epsilon$ for some fixed small $\epsilon \in (0, 1)$, we thus conclude from (5-10) that

$$2\|h\nabla_g(\chi u)\|^2 \leq 2\operatorname{Im}\left\langle \left(hD_x - \frac{in}{2}\frac{h}{x}\right)(\chi u), h^{-1}x\chi Pu \right\rangle - 2h^{-1} \operatorname{Im}\langle \chi u, \chi Pu \rangle + \|Eu\|^2. \tag{5-11}$$

Hardy's inequality gives $\|(h/x)\chi u\| \leq C_n\|hD_x(\chi u)\|$; hence the first term on the right is bounded from above by

$$\epsilon\|hD_x(\chi u)\|^2 + C_\epsilon\|h^{-1}x\chi Pu\|^2.$$

For the second term in (5-11), we rewrite

$$\begin{aligned} \langle \chi u, \chi Pu \rangle &= -\|\chi u\|^2 + \langle h\nabla_g(\chi^2 u), h\nabla_g u \rangle \\ &= -\|\chi u\|^2 + \|\chi h\nabla_g u\|^2 + \langle h(\nabla_g \chi^2)u, h\nabla_g u \rangle; \end{aligned}$$

taking the imaginary part annihilates the first two terms, while for the final term we have

$$\begin{aligned} \langle h(\nabla_g \chi^2)u, h\nabla_g u \rangle - \langle h\nabla_g u, h(\nabla_g \chi^2)u \rangle &= ah^2\nabla_g((\nabla_g \chi^2)u) - h^2(\nabla_g \chi^2) \cdot \nabla_g u, u \\ &= \langle h^2\Delta_g(\chi^2)u, u \rangle, \end{aligned} \tag{5-12}$$

with no derivatives falling on u anymore. Altogether, we obtain from (5-11) the estimate

$$(2 - C\epsilon)\|h\nabla_g(\chi u)\|^2 \leq C_\epsilon\|h^{-1}x\chi Pu\|^2 + Ch\|\tilde{\chi}u\|^2 + \|Eu\|^2, \tag{5-13}$$

where $\tilde{\chi} \equiv 1$ on $\operatorname{supp} \chi$, used to bound the contribution of (5-12).

Control of χu in $H_{c,h}^{1,1,0,0}$: Since the principal symbol of

$$h\nabla_g \in \Psi_{ch}^{1,1,0,0}(X; \mathbb{C}, {}^{ch}T^*X_{ch})$$

(mapping complex-valued functions into sections of ${}^{ch}T^*X_{ch}$, see Remark 3.3) is not injective at the zero section over sf , the estimate (5-13) does not yet give full control of χu in $H_{c,h}^{1,1,0,0}$: an estimate of $\|\chi u\|_{L^2}^2$

is lacking at this point. (Note that the control of $(h/x)\chi u$ via Hardy’s inequality degenerates precisely at sf, i.e., the lift of $h = 0$.) The key observation is that the characteristic set of P and the set where the principal symbol of $h\nabla_g$ fails to be injective are disjoint. Thus, for some $A_1 \in \Psi_{ch}^{-1,-1,0,0}(X; {}^{ch}T^*X_{ch}, \mathbb{C})$ and $A_2 \in \Psi_{ch}^{-2,-2,0,0}(X)$, we have

$$I = A_1 \circ h\nabla_g + A_2 P + R, \quad R \in \Psi_{ch}^{-\infty,0,0,-\infty}(X);$$

this implies

$$\|\chi u\| \leq C(\|h\nabla_g(\chi u)\|_{H_{c,h}^{-1,-1,0,0}(X)} + \|P(\chi u)\|_{H_{c,h}^{-2,-2,0,0}(X)} + \|R(\chi u)\|_{L^2}). \tag{5-14}$$

Using $[P, \chi] \in \Psi_{ch}^{1,-\infty,-\infty,-1}(X)$, we can estimate the second term by

$$\begin{aligned} \|P(\chi u)\|_{H_{c,h}^{-2,-2,0,0}} &\leq \|\chi Pu\|_{H_{c,h}^{-2,-2,0,0}} + \|[P, \chi]u\|_{H_{c,h}^{-2,-2,0,0}} \\ &\lesssim \|h^{-1}x\chi Pu\|_{H_{c,h}^{-2,-1,0,-1}} + \|\tilde{\chi}u\|_{H_{c,h}^{-1,-N,-N,-1}} \end{aligned}$$

for any $N \in \mathbb{R}$. The remainder term in (5-14) is simply estimated by

$$\|R(\chi u)\|_{L^2} \leq C \left\| \frac{h}{h+x} \chi u \right\|_{L^2} \leq C \left\| \frac{h}{x} \chi u \right\|_{L^2}.$$

Applying Hardy’s inequality to this term, the estimate (5-14) then implies, a fortiori,

$$\|\chi u\|_{L^2} \leq C'(\|h\nabla_g(\chi u)\|_{L^2} + \|h^{-1}x\chi Pu\|_{L^2} + h\|\tilde{\chi}u\|_{L^2}).$$

We can now add η times this, with $\eta C' < \frac{1}{2}$, to the estimate (5-13) (in which we fix $\epsilon < C^{-1}$), in order to obtain

$$\|\chi u\|_{\mathcal{D}_h^1}^2 = \|\chi u\|^2 + \|h\nabla_g(\chi u)\|^2 \lesssim \|h^{-1}x\chi Pu\|^2 + \|Eu\|^2 + h\|\tilde{\chi}u\|^2. \tag{5-15}$$

As far as weights in h and x are concerned, this is already the desired estimate. However, the differential order is forced to be 1 here, and in addition the order of differentiability required on Pu in (5-15) is too strong (0 instead of -1) even in this special case.

Sharp improvement at tf: The basic idea is to apply the estimate (5-15) to $\phi(h^2\Delta_g)u$, where $\phi \in \mathcal{C}_c^\infty(\mathbb{R})$ is equal to 1 on $[-4, 4]$; on the remaining piece $(1-\phi(h^2\Delta_g))u$, the operator $h^2\Delta_g - 1$ can be inverted directly using the functional calculus. In order to define $h^2\Delta_g$ as a self-adjoint operator, we need to pass from X to a compact manifold X' . If $Y = \partial X$ is null-cobordant, we may choose X' so that $\partial X' = \partial X$, and we then extend g to a Riemannian metric on X' which we continue to denote by g . The operator $\phi(h^2\Delta_g)$ does depend on the choice of extension, but its structural properties, as used in the following argument, do not. If Y is not null-cobordant, we may set $X' = [0, 6x_0]_x \times Y$ and define a smooth metric on X' which is equal to g on $[0, 2x_0] \times Y$ and equal to the pullback of g along the map $(x, y) \mapsto (6x_0 - x, y)$ on $(4x_0, 6x_0] \times Y$; we denote this metric g again. Thus, we have two identical cone points at $x = 0$ and at $6x_0 - x = 0$.

Concretely then, $\Phi := \phi(h^2\Delta_g)$ is given by Lemma 5.5, the notation of which we shall use here. Now, in order to remain localized near ∂X , we apply the estimate (5-15) to

$$u_1 := \tilde{\chi} \Phi \tilde{\chi} u.$$

Using $[P, \Phi] \equiv 0$ and $\chi[P, \tilde{\chi}] \equiv 0$, we estimate the first term on the right in (5-15) by

$$\|h^{-1}x\chi P\tilde{\chi}\Phi\tilde{\chi}u\| \leq \left\| \frac{x}{h}\chi\Phi\frac{h}{h+x}(h^{-1}(x+h)\tilde{\chi}Pu) \right\| + \|h^{-1}x\chi\Phi[P, \tilde{\chi}]u\|. \tag{5-16}$$

Denoting the lift of x to the left, resp. right factor of X_{ch}^2 by x , resp. x' , we note that

$$\frac{x}{h}\frac{h}{x'+h} \in \mathcal{A}^{1,1,0,0,0,0}(X_{ch}^2) \implies \Psi := \frac{x}{h}\chi\Phi\frac{h}{x'+h} \in \mathcal{A}^{1-\epsilon, n+1-\epsilon, n-\epsilon, 0, 0, \infty}(X_{ch}^2).$$

Passing to a b-density $0 < \mu_0 \in C^\infty(X; {}^b\Omega^1 X)$, we claim that Ψ is continuous as a map

$$H_{c,h}^{-\infty, -1, 0, 0}(X; |dg|) = H_{c,h}^{-\infty, -1-\frac{n}{2}, -\frac{n}{2}, 0}(X; \mu_0) \rightarrow H_{c,h}^{\infty, -\frac{n}{2}, -\frac{n}{2}, 0}(X; \mu_0) = H_{c,h}^{\infty, 0, 0, 0}(X; |dg|),$$

but since Ψ is smoothing in the sense of ch -differentiability, it suffices to show the boundedness on $L^2(X; \mu_0)$ of

$$x^{\frac{n}{2}}\Psi(x')^{-\frac{n}{2}}\left(\frac{x'}{x'+h}\right)^{-1} \in \mathcal{A}^{\frac{n}{2}+1-\epsilon, n-\epsilon, \frac{n}{2}-1-\epsilon, 0, 0, \infty}(X_{ch}^2; {}^b\Omega^{\frac{1}{2}}). \tag{5-17}$$

Since this kernel is bounded section of ${}^b\Omega^{1/2}$ (all indices being ≥ 0), this is a consequence of Schur's lemma.

The operator acting on u in the second term on the right in (5-16) has Schwartz kernel supported in $x' \geq c > 0$ and $|x - x'| > c > 0$ (since $\text{supp } \chi \cap \text{supp } d\tilde{\chi} = \emptyset$), hence lies in $\mathcal{A}^{1-\epsilon, \infty, \infty, \infty, \infty, \infty}(X_{ch}^2)$; therefore, the second term in (5-16) can be bounded by $h^N \|\tilde{\chi}^\sharp u\|$ for any N , where $\tilde{\chi}^\sharp = 1$ on $\text{supp } \tilde{\chi}$. Altogether, forgetting the cutoff $\tilde{\chi}$ and renaming $\tilde{\chi}^\sharp$ as $\tilde{\chi}$, we have proved

$$\|\chi u_1\|_{H_{c,h}^{N, 1, 0, 0}} \lesssim h^{-1} \|(x+h)\tilde{\chi}Pu\|_{H_{c,h}^{-N, -1, 0, 0}} + \|Eu\|_{H_{c,h}^{-N, -N, -N, 0}} + h^{\frac{1}{2}} \|\tilde{\chi}u\|_{H_{c,h}^{-N, 0, 0, 0}} \tag{5-18}$$

for any $s, N \in \mathbb{R}$.

It remains to control χu_2 , where

$$u_2 := u - u_1 = u - \tilde{\chi}\Phi\tilde{\chi}u.$$

Let $\phi^b \in C_c^\infty((-3, 3))$ be identically 1 on $[-2, 2]$, and let $\Phi^b = \phi^b(h^2\Delta_g)$. Then χu_2 is localized near high frequencies, in the sense that its localization to low frequencies

$$\Phi^b(\chi u_2) = \Phi^b(\chi u) + \Phi^b[\Phi, \chi]\tilde{\chi}u - \Phi^b\Phi\chi\tilde{\chi}u = \Phi^b[\Phi, \chi]\tilde{\chi}u \tag{5-19}$$

(using $\Phi^b\Phi = \Phi^b$ and $\chi\tilde{\chi} = \chi$) is $\mathcal{O}(h^\infty)$ (due to the presence of $[\Phi, \chi]$) near $x = 0$ and vanishes to an order h more than u near $\text{supp } d\chi \subset X^\circ$. Moreover, χu_2 satisfies the equation

$$P(\chi u_2) = (\chi - \chi\Phi\tilde{\chi})Pu + ([P, \chi]u - [P, \chi]\Phi\tilde{\chi}u - \chi\Phi[P, \tilde{\chi}]u), \tag{5-20}$$

Altogether, if we put

$$P^\sharp := P + 2\Phi^b,$$

then we have

$$P^\sharp(\chi u_2) = f_2 := (\chi - \chi\Phi\tilde{\chi})Pu + [P, \chi]u - ([P, \chi]\Phi\tilde{\chi}u + \chi\Phi[P, \tilde{\chi}]u - 2\Phi^b[\Phi, \chi]\tilde{\chi}u). \tag{5-21}$$

We moreover have $P^\sharp = p^\sharp(h^2\Delta_g)$, where $p^\sharp(\sigma) := (\sigma - 1) + 2\phi^b \geq \frac{1}{2}(\sigma + 1)$ for $\sigma \geq 0$; hence we can invert P^\sharp using the functional calculus for Δ_g by $(P^\sharp)^{-1} = q^\sharp(h^2\Delta)$ where $q^\sharp(\sigma) = 1/p^\sharp(\sigma)$ is equal

to $(\sigma - 1)^{-1}$ for large σ . One can then show, by a combination of the arguments leading to Lemma 5.5 and [Hintz 2022, Theorem 5.2], that

$$(P^\sharp)^{-1} \in \left(\frac{x}{x+h} \right)^2 \Psi_{ch}^{-2}(X) + \Psi_{ch}^{-\infty, \mathcal{E}}(X)$$

where the collection \mathcal{E} of index sets is equal to $\mathcal{E}(-1)$ in the notation of [Hintz 2022, Theorem 6.1]. Therefore, using (5-21), the mapping properties of elements of $(x/(x+h))^2 \Psi_{ch}^{-2}(X)$, and estimating the smoothing contribution in the space $\Psi_{ch}^{-\infty, \mathcal{E}}(X)$ to $(P^\sharp)^{-1}$ by means of Schur’s lemma, we have

$$\|\chi u\|_{H_{c,h}^{s,1,0,0}} = \|(P^\sharp)^{-1} f_2\|_{H_{c,h}^{s,1,0,0}} \lesssim \|\tilde{\chi} P u\|_{H_{c,h}^{s-2,-1,0,0}} + \|\tilde{\chi} [P, \chi] u\|_{H_{c,h}^{s-2,-1,0,0}} + h \|\tilde{\chi} u\|_{H_{c,h}^{-N,-N,0,0}}. \tag{5-22}$$

Here, the first term on the right comes from the first term in (5-21) and the boundedness of $\chi - \chi \Phi \tilde{\chi}^\sharp$ (with $\tilde{\chi}^\sharp \equiv 1$ on $\text{supp } \tilde{\chi}$) on $H_{c,h}^{s-2,-1,0,0}$; this boundedness follows from the boundedness of the Schwartz kernel of

$$x^{\frac{n}{2}} \frac{x}{x+h} \Phi \left(\frac{x'}{x'+h} \right)^{-1} (x')^{-\frac{n}{2}} \in \mathcal{A}^{\frac{n}{2}+2-\epsilon, n+1-\epsilon, \frac{n}{2}-1-\epsilon, 0, 0, \infty} (X_{ch}^2; {}^b\Omega^{\frac{1}{2}})$$

similarly to the discussion of Ψ in (5-17). The final term in (5-22) comes from the big parenthesis in (5-21), every term of which involves the localizer Φ to low frequencies as well as a commutator with a cutoff χ or $\tilde{\chi}$. But $\tilde{\chi} [P, \chi] \in \Psi_{ch}^{1,-\infty,-\infty,-1}(X)$; hence the second term on the right is bounded from above by $\|\tilde{\chi} u\|_{H_{c,h}^{s-1,-N,-N,-1}}$ for any N . By elliptic regularity at infinite semiclassical cone frequencies, this can be bounded by $C(\|\tilde{\chi}^\sharp P u\|_{H_{c,h}^{s-3,-N-2,-N,-1}} + \|\tilde{\chi}^\sharp u\|_{H_{c,h}^{-N,-N,-N,-1}})$. Combining the resulting estimate with (5-18) proves the theorem for $H_{c,h}$ -spaces. The proof of the more general statement for $H_{cb,h}$ -spaces requires only notational changes which are left to the reader. \square

To complete the proof, it remains to prove Lemma 5.5.

Proof of Lemma 5.5. This can be proved using the Helffer–Sjöstrand formula [1989] similarly to [Vasy 2000, Lemma 10.1 and Proposition 10.2]. Choosing a compactly supported almost analytic extension $\tilde{\phi} \in C_c^\infty(\mathbb{C})$ of ϕ (that is, $\tilde{\phi}|_{\mathbb{R}} = \phi$ and $|\partial_{\bar{z}} \tilde{\phi}| = \mathcal{O}(|\text{Im } z|^N)$ for all N), we have

$$\phi(h^2 \Delta_g) = \frac{1}{2\pi i} \int \partial_{\bar{z}} \tilde{\phi}(z) (h^2 \Delta_g - z)^{-1} d\bar{z} \wedge dz. \tag{5-23}$$

For $z \notin \mathbb{R}$, [Hintz 2022, Theorem 3.10 and §6.1] gives

$$(h^2 \Delta_g - z)^{-1} \in \left(\frac{x}{x+h} \right)^2 \Psi_{ch}^{-2}(X) + \Psi_{ch}^{-\infty, \mathcal{E}}(X), \tag{5-24}$$

where $\mathcal{E} = (\mathcal{E}_{\text{lb}}, \mathcal{E}_{\text{ff}}, \mathcal{E}_{\text{rb}}, \mathcal{E}_{\text{tf}})$, with $\text{Re } z \geq 0$ for $(z, k) \in \mathcal{E}_{\text{lb}}$; $\text{Re } z \geq n$ for $(z, k) \in \mathcal{E}_{\text{rb}}$; $\text{Re } z \geq 2$ for $(z, k) \in \mathcal{E}_{\text{ff}}$; and $\text{Re } z \geq 0$ for $(z, k) \in \mathcal{E}_{\text{tf}}$. Using that the principal symbol of $h^2 \Delta_g$ is real-valued and its normal operator $\Delta_{\hat{g}}$ is self-adjoint, we claim that any fixed seminorm (5-24) is moreover bounded by $|\text{Im } z|^{-k}$ for some k (depending on the seminorm).

To justify this claim, it is instructive to first consider the corresponding statement for $(\Delta - z)^{-1} \in \Psi^{-2}$, $z \in \mathbb{C} \setminus \mathbb{R}$, where Δ is the Laplacian on a closed Riemannian manifold: for any N , the construction of a symbolic parametrix of $\Delta - z$ of order N gives $Q_{N,z}, Q'_{N,z} \in \Psi^{-2}$, with seminorms bounded

by $|\operatorname{Im} z|^{-C}$ for some C depending on N and the seminorm, so that $(\Delta - z)Q_{N,z} = I - R_{N,z}$ and $Q'_{N,z}(\Delta - z) = I - R'_{N,z}$, where $R_{N,z}, R'_{N,z} \in \Psi^{-N}$ obey such bounds as well. But then $(\Delta - z)^{-1} = Q_{N,z} + Q'_{N,z}R_{N,z} + R'_{N,z}(\Delta - z)^{-1}R_{N,z}$, where the first two summands as pseudodifferential operators and the third summand (the “remainder operator”) $R'_{N,z}(\Delta - z)^{-1}R_{N,z}$ as a map $H^{-N} \rightarrow H^N$ obey such bounds; this uses that $\|(\Delta - z)^{-1}\| = |\operatorname{Im} z|^{-1}$ as an operator on L^2 . But any seminorm on Ψ^{-2} is continuous on the space of bounded operators $H^{-N} \rightarrow H^N$ for sufficiently large N . This implies that any Ψ^{-2} -seminorm of $(\Delta - z)^{-1}$ is bounded by $|\operatorname{Im} z|^{-k}$ for some k . (In this simple example, the claim follows also directly from Beals’ Theorem [1977].)

Analogous arguments can be used to control the inverse of the normal operator $\Delta_{\hat{g}} - z$: in addition to carrying out N steps of the symbolic parametrix construction, one uses the inverse of the b-normal operator (which is independent of z) to solve away to order N at the left/right boundary (for the left/right parametrix) and the b-front face of $\operatorname{tf}_{\text{b,sc}}^2$; by virtue of taking only finitely many steps, one ensures the validity of $|\operatorname{Im} z|^{-k}$ bounds on seminorms. The true inverse $(\Delta_{\hat{g}} - z)^{-1}$ then obeys such bounds as well since any seminorm on the space of large b-scattering ps.d.o.s in which $(\Delta_{\hat{g}} - z)^{-1}$ lies (see [Hintz 2022, equation (3.28)] for a general statement) is continuous on the space of remainder operators $H_{\text{b,sc}}^{-N,l_1,-N} \rightarrow H_{\text{b,sc}}^{N,l_2,N}$ (for appropriate orders l_1, l_2 , with $l_2 = -N$, resp. $l_1 = -N$ in the case of the left, resp. right parametrix construction) for sufficiently large N .

The $|\operatorname{Im} z|^{-k}$ bounds for seminorms of (5-24) can then be proved by completely analogous means, namely by constructing a parametrix which is accurate to some finite order, and observing that any fixed seminorm on the space (5-24) is continuous on the relevant space of remainder operators.

Plugging these into (5-23), we conclude that $\phi(h^2\Delta_g)$ is of the same class as the resolvent. To improve the orders, let $m \in \mathbb{N}$ and write $\phi(\sigma) = (\sigma + C)^{-m}\phi_m(\sigma)$ with $C > -\inf \operatorname{supp} \phi$. Then

$$\phi(h^2\Delta_g) = (h^2\Delta_g + C)^{-m}\phi_m(h^2\Delta_g).$$

Applying the previous discussion to ϕ_m and using [Hintz 2022, Theorem 6.3(3)] (with $w = -m$) to control $(h^2\Delta_g + C)^{-m}$ implies, upon letting $m \rightarrow \infty$, that $\phi(h^2\Delta_g) \in \Psi_{\text{ch}}^{-\infty,-\infty,0,0}(X) + \mathcal{A}^{-\epsilon,n-\epsilon,n-\epsilon,0,\infty,\infty}(X_{\text{ch}}^2)$. This gives (5-8). □

5B. Scattering by potentials with inverse square singularities. Complex absorption is a somewhat drastic method for gaining microlocal control along incoming directions. As a more natural setting, let us thus consider scattering by potentials on \mathbb{R}^n , $n \geq 2$, which are singular at the origin $0 \in \mathbb{R}^n$, as in Theorem 1.1. (Working on more general conic manifolds requires only minor modifications.) That is, the underlying spatial manifold is

$$X = [\mathbb{R}_x^n; \{0\}] \cong [0, \infty)_r \times \mathbb{S}^{n-1}, \quad g = dx^2 = dr^2 + r^2g_{\mathbb{S}^{n-1}}. \tag{5-25}$$

We write $\Delta \equiv \Delta_g = \sum D_{x_j}^2$ for the (nonnegative) Laplacian. Let $N \in \mathbb{N}$, and denote by Δ the Laplacian acting componentwise on \mathbb{C}^N -valued functions. We consider scattering by matrix-valued potentials

$$V(x) = |x|^{-2}V_0(x), \quad V_0 \in C_c^\infty(X; \mathbb{C}^{N \times N}).$$

The assumption of compact support of V can of course be relaxed considerably, but since our interest lies in understanding the effect of the singularity at $r = 0$, we shall not concern ourselves with more general conditions on V at infinity here.

We are interested in high-energy estimates for the resolvent of $\Delta + V$; concretely, we shall consider $\Delta + V - \sigma^2$, where $\text{Im } \sigma > 0$ is bounded and $|\text{Re } \sigma| \gg 1$. Upon introducing

$$h = |\sigma|^{-1}, \quad z = (h\sigma)^2 = 1 + \mathcal{O}(h), \tag{5-26}$$

we define

$$P_{h,z} := h^2(\Delta + V - \sigma^2) = h^2\Delta - z + h^2r^{-2}V_0. \tag{5-27}$$

This is admissible in the sense of Definition 4.1, with $Q_{1,z} = V_0$ and $q_{0,z} = 0$. Since $Q_{1,z}$ has differential order 0, the threshold quantities in Definition 4.7 are $r_{\text{in}} = r_{\text{out}} = -\frac{1}{2}$. The normal operator of $P_{h,z}$ is computed by passing to $\hat{r} = \frac{r}{h}$ and setting $h = 0$:

$$\begin{aligned} N(P) &= \Delta_{\hat{g}} - 1 + \hat{r}^{-2}V_{\partial}, \\ \hat{g} &:= d\hat{r}^2 + \hat{r}^2g_{\mathbb{S}^{n-1}}, \quad V_{\partial} := V_0|_{\partial X} \in \mathcal{C}^{\infty}(\partial X; \mathbb{C}^{N \times N}). \end{aligned} \tag{5-28}$$

(Thus $V_{\partial}(\omega) = V_0(0, \omega)$ in the coordinates $(r, \omega) \in [0, \infty) \times \mathbb{S}^2$ on X .)

Theorem 5.7 (potential scattering). *Assume that the operator $N(P)$ defined in (5-28) is invertible at weight $l \in \mathbb{R}$ (in the sense of Definition 4.6(3)). Let $C > 0$, and let $\chi_0 \in \mathcal{C}_c^{\infty}(X)$ be identically 1 near $r = 0$. Then there exists $C' > 0$ so that for $0 < \text{Im } \sigma < C$ and $|\text{Re } \sigma| > C'$, the operator $\Delta + V - \sigma^2$ is invertible as a map*

$$\begin{aligned} \Delta + V - \sigma^2 : \{u \in H_{\text{loc}}^2(X^{\circ}) : \chi_0 u \in r^l H_b^2(X), (1 - \chi_0)u \in H^2(\mathbb{R}^n)\} \\ \rightarrow \{f \in L_{\text{loc}}^2(X^{\circ}) : \chi_0 f \in r^{l-2} L^2(X), (1 - \chi_0)f \in L^2(\mathbb{R}^n)\}. \end{aligned} \tag{5-29}$$

Moreover, in the notation (5-26)–(5-27), the following uniform estimate holds for all $\epsilon, \delta > 0$, a suitable constant $C_{\epsilon,\delta} > 0$, and all $0 < \text{Im } \sigma < C$, $|\text{Re } \sigma| > C'$:

$$\begin{aligned} \|\chi_0 u\|_{H_{c,h}^{s,l,1/2+\epsilon,0}} + \|(1 - \chi_0)u\|_{H_{sc,h}^{s,-1/2-\delta}} \\ \leq C_{\epsilon,\delta} h^{-1-2\epsilon} (\|\chi_0 P_{h,z} u\|_{H_{c,h}^{s-2,l-2,-1/2-\epsilon,0}} + \|(1 - \chi_0)P_{h,z} u\|_{H_{sc,h}^{s-2,1/2+\delta}}); \end{aligned} \tag{5-30}$$

here, $H_{sc}^{s,\gamma} = H_{sc}^{s,\gamma}(\overline{\mathbb{R}^n}) = \langle r \rangle^{-\gamma} \mathcal{F}^{-1}((hD)^{-s} L^2(\mathbb{R}^n))$ is the semiclassical scattering Sobolev space. In particular, for $l_1 \leq \min(l, \frac{1}{2} + \epsilon)$, $l_2 \geq \max(l - 2, -\frac{1}{2} - \epsilon)$, and for any fixed $\chi \in \mathcal{C}_c^{\infty}(X)$ we have

$$\|\chi(\Delta + V - \sigma^2)^{-1} \chi f\|_{r^{l_1} L^2} \leq C_{\epsilon,\chi} |\sigma|^{-1+2\epsilon} \|f\|_{r^{l_2} L^2}. \tag{5-31}$$

We recall that for $V(x) = |x|^{-2}V_0(x)$, with real-valued V_0 satisfying $V_0(0) > -((n - 2)/2)^2$ (and indeed under relaxed regularity requirements on V_0 , and allowing for the presence of several such inverse square singularities), Duyckaerts [2006] obtained cutoff resolvent estimates of the form (5-31) without the 2ϵ -loss. It is an interesting question — which we do not address here — whether in this setting, or perhaps even in the general setting of Theorem 5.7, one can prove a lossless estimate using a global commutator argument similar to the one used in Section 5A.

Remark 5.8 (meromorphic continuation). For V with compact support as above, the resolvent $(\Delta + V - \sigma^2)^{-1}$ can be meromorphically continued to the complex plane when n is odd, and the logarithmic cover of \mathbb{C}^\times when n is even; the estimate (5-31) holds in strips of bounded $\text{Im } \sigma$ for large $|\text{Re } \sigma|$. The construction of this continuation can be accomplished along the lines of black box scattering [Sjöstrand and Zworski 1991] (see also [Dyatlov and Zworski 2019, §4]), with those estimates in the references in $\text{Im } \sigma \gg 1$ relying on self-adjointness replaced by estimates on the off-spectrum resolvent that follow from [Hintz 2022, Theorem 3.10]. For applications of such estimates to expansions of scattered waves for n odd, we refer the reader to [Dyatlov and Zworski 2019, §3.2.2].

Remark 5.9 (vector bundles). One may more generally consider potentials valued in endomorphisms of a vector bundle $E \rightarrow X$, with Δ denoting an operator acting on sections of E with scalar principal symbol given by the dual metric function. The main difference to the case of a trivial bundle is that the threshold quantities $r_{\text{in}}, r_{\text{out}}$ depend on subprincipal terms of Δ (and their calculation requires the choice of a fiber inner product on E ; see Remark 4.11). In the special case of tensor bundles E , and with Δ denoting the tensor Laplacian, the fiber inner product on E induced by g does give $r_{\text{in}} = r_{\text{out}} = -\frac{1}{2}$.

Proof of Theorem 5.7. Semiclassical propagation estimates near infinity of \mathbb{R}^n are standard, see, e.g., [Vasy and Zworski 2000, Theorem 1] (following [Melrose 1994]) in a general geometric setting, and can be combined with the propagation estimates through the singularity at $r = 0$ given in Theorem 4.10 (where we shall take $\alpha = \frac{1}{2} + \epsilon$, $b = 2\epsilon$ near \mathcal{R}_{in} , and $b = 0$ near \mathcal{R}_{out}). Altogether, upon simplifying to constant orders, we obtain, for any $\delta > 0$, and for $0 < h < h_0$ with $h_0 > 0$ sufficiently small,

$$\|\chi_0 u\|_{H_{c,h}^{s,l,1/2+\epsilon,0}} + \|(1 - \chi_0)u\|_{H_{sc,h}^{s,-1/2-\delta}} \lesssim \|\chi_0 P_{h,z} u\|_{H_{c,h}^{s-2,l-2,1/2+\epsilon,1+2\epsilon}} + h^{-1-2\epsilon} \|(1 - \chi_0)P_{h,z} u\|_{H_{sc,h}^{s-2,1/2+\delta}},$$

which is the estimate (5-30). (The loss of $h^{-2\epsilon}$ in the second term on the right is due to the fact that the propagation through $r = 0$ comes with this loss, which then gets propagated out to infinity.) This estimate also gives the injectivity of $P_{h,z}$ for small $h > 0$ and $|z - 1| < Ch$, with surjectivity following from the analogous estimate for the adjoint; this proves the first part of the theorem, albeit on function spaces with weights $\langle r \rangle^{\pm(1/2+\delta)}$ at infinity. But for any fixed σ with $\sigma^2 \notin [0, \infty)$, $\Delta + V - \sigma^2$ is an elliptic scattering operator near infinity; hence these weights can be removed. (It is only in the high-energy limit $|\text{Re } \sigma| \rightarrow \infty$ with $|\text{Im } \sigma / \text{Re } \sigma| \rightarrow 0$ that one loses uniform (in σ) ellipticity.)

The simplified estimate (5-31) follows by setting $s = 2$ in

$$h^{-1-2\epsilon} \|\chi P_{h,z} u\|_{H_{c,h}^{s-2,l-2,-1/2-\epsilon,0}} \leq h^{1-2\epsilon} \|\chi(\Delta + V - \sigma^2)u\|_{H_{c,h}^{s-2,l_2,l_2,0}},$$

$$\|\chi u\|_{H_{c,h}^{s,l,1/2+\epsilon,0}} \leq \|\chi u\|_{H_{c,h}^{s,l_1,l_1,0}}. \quad \square$$

We describe a few scenarios in which the invertibility assumption on $N(P)$ can be verified:

- (1) The operator $N(P)$ is invertible for an open set of $V_\partial \in \mathcal{C}^\infty(\mathbb{S}^{n-1}; \mathbb{C}^{N \times N})$. In particular, it holds when $n \geq 3$, $l \in (1 - (n - 2)/2, 1 + (n - 2)/2)$, and $V_\partial \equiv 0$ by Lemma 5.1, and therefore also when $\|V_\partial\|_{\mathcal{C}^k}$ is sufficiently small (depending on l) for some sufficiently large k .
- (2) Consider V_∂ which depends holomorphically on a parameter $w \in \Omega$, where $\Omega \subset \mathbb{C}$ is open and contains 0. (For example, this is the case when $V_\partial(w) = wV_{\partial,0}$.) Let us write $N(P_w)$ for the w -dependent

normal operator, and assume that $N(P_0)$ is invertible at weight l_0 ; assume moreover that there is a continuous function $l : \Omega \rightarrow \mathbb{R}$ with $l(0) = l_0$ so that $l(w) \notin -\text{Im spec}_b(N(P_w))$. Then there exists a discrete set $D \subset \Omega$ so that $N(P_w)$ is invertible at weight $l(w)$ for $w \in \Omega \setminus D$. This follows from analytic Fredholm theory in w ; we leave the details to the reader.

A very concrete third scenario is the following:

Lemma 5.10 (scalar inverse square potentials). *Let $n \geq 2$, consider the scalar case $N = 1$, and suppose $V_\partial = Z \in \mathbb{C} \setminus (-\infty, -((n-2)/2)^2]$ is a constant (so $V(x) = Z/|x|^2 + \mathcal{O}(|x|^{-1})$). Then $N(P)$ is invertible at weight l if and only if $|l - 1| < \text{Re} \sqrt{((n-2)/2)^2 + Z}$.*

Proof of Lemma 5.10. The boundary spectrum of $N(P)$ can be computed, via expansion into spherical harmonics, as

$$\text{spec}_b(N(P)) = \left\{ i \left(\frac{n-2}{2} \pm \sqrt{\left(\frac{n-2}{2} + \ell \right)^2 + Z} \right) : \ell \in \mathbb{N}_0 \right\}.$$

The condition on Z ensures that

$$\text{Re} \sqrt{\left(\frac{n-2}{2} \right)^2 + Z} > 0,$$

and thus for l as in the statement of the lemma, one has $l - n/2 \notin -\text{Im spec}_b(N(P))$.

Expanding an outgoing solution u of $N(P)u = 0$ at weight l into spherical harmonics, $u(\hat{r}, \omega) = \sum_{|m| \leq \ell} u_{\ell m}(\hat{r}) Y_{\ell m}(\omega)$, the coefficient $u_{\ell m}$ satisfies a Bessel ODE

$$-u''_{\ell m} - \frac{n-1}{\hat{r}} u'_{\ell m} + \frac{\ell(\ell+n-2) + Z}{\hat{r}^2} u_{\ell m} - u_{\ell m} = 0,$$

and hence is a linear combination of $\hat{r}^{-(n-2)/2} H_{\nu_\ell}^{(1)}(\hat{r})$ and $\hat{r}^{-(n-2)/2} H_{\nu_\ell}^{(2)}(\hat{r})$ where we set

$$\nu_\ell = \sqrt{\left(\frac{n-2}{2} + l \right)^2 + Z}.$$

The outgoing condition can only be satisfied if $u_{\ell m}$ is a multiple of $\hat{r}^{-(n-2)/2} H_{\nu_\ell}^{(1)}(\hat{r})$. But for $0 < \hat{r} \ll 1$, one has

$$|\hat{r}^{-\frac{n-2}{2}} H_{\nu_\ell}^{(1)}(\hat{r})| \geq c_\ell \hat{r}^{-\frac{n-2}{2} - \text{Re } \nu_\ell} \geq c_\ell \hat{r}^{-\frac{n-2}{2} - \text{Re } \nu_0},$$

with $c_\ell > 0$, which lies in $\hat{r}^{l'} L^2(\hat{r}^{n-1} d\hat{r})$ if and only if $l' < 1 - \text{Re } \nu_0$, which is violated for $l' = l$. Hence necessarily $u_{\ell m} = 0$. This proves that $N(P)$ is injective at weight l on outgoing functions; the injectivity of $N(P)^*$ at weight $-l + 2$ on incoming functions is proved similarly. \square

Theorem 1.1 follows from Theorem 5.7 and Lemma 5.10 upon taking $\sigma = \sqrt{\lambda}$ and $l = 2$, which allows for the choice $l_1 = l_2 = 0$. Note that for $l = 2$, the target space in (5-29) in $L^2(X) = L^2(\mathbb{R}^n)$, and the domain is $H_0^2(\mathbb{R}^n \setminus \{0\})$ for $n \geq 5$ by Hardy's inequality.

Remark 5.11 (multiple scatterers). By exploiting the diffractive improvement obtained in Section 4F as in [Baskin and Wunsch 2013], it is conceivable that one can generalize (up to ϵ -losses) Duyckaerts' high-energy resolvent estimates [2006] for scattering by a finite number of real-valued inverse square

potentials and analyze the complex-valued case $Z_j \in \mathbb{C} \setminus (-\infty, -((n-2)/2)^2]$ (or more generally the case of finitely many matrix-valued inverse square potentials). However, the study of this problem exceeds the scope of this paper.

Remark 5.12 (operators with inverse square singularities arising in the study of wave equations). Following [Baskin and Marzuola 2022], consider a static metric $g = -dt^2 + dr^2 + r^2k$, where k is a Riemannian metric on a closed manifold Z ; e.g., $(Z, k) = (\mathbb{S}^{n-1}, g_{\mathbb{S}^{n-1}})$, in which case g is the Minkowski metric. In the region $t > 1$, $r/t < \frac{1}{2}$, we introduce $T = 1/t$, $R = r/t$ (in the notation of the reference: ρ, x) and compute

$$t^2 \square_g = -(TD_T + RD_R)^2 + i(TD_T + RD_R) + D_R^2 - \frac{i(n-1)}{R}D_R + R^{-2}\Delta_k.$$

Restricting the coefficients to $T = 0$ (as a b-operator) and formally passing to the Mellin transform by replacing TD_T with multiplication by $\sigma \in \mathbb{C}$ gives

$$P_\sigma = \Delta_G - (RD_R + \sigma)^2 + i(RD_R + \sigma) = \Delta_G - \sigma^2 + Q, \quad G := dR^2 + R^2k,$$

where $Q = i\sigma - (2\sigma - i)RD_R$. When $|\text{Im } \sigma|$ is bounded and $|\text{Re } \sigma| \rightarrow \infty$, we let $h = |\sigma|^{-1}$; then the rescaling

$$P_{h,z} = h^2 P_{h^{-1}z} = h^2 \Delta_G - z^2 + h^2 Q \tag{5-32}$$

is an admissible operator on $[0, \frac{1}{2})_R \times Z$ in the sense of Definition 4.1. More generally, consider the coupling of \square_g with a potential $V = V(r, z)$ which asymptotes to an inverse square potential as $r \rightarrow \infty$, i.e., $V(r, z) = r^{-2}V_0(z) + \mathcal{O}(r^{-2-\delta})$ with $V_0 \in C^\infty(Z)$. Then we have $t^2(\square_g + V) = t^2 \square_g + R^{-2}V_0(z) + \mathcal{O}(T^\delta R^{-2-\delta})$, and therefore the rescaling (5-32) has an additional $h^2 R^{-2}V_0$ term, as studied in the present section; that is, a stationary *asymptotically* inverse square potential on $[0, \infty)_r \times Z$ gives rise to an inverse square *singularity* of $P_{h,z}$ at $R = 0$.¹⁶ (If $V = r^{-2}V_0(z)$ is an exact inverse square potential, then $P_{h,z}$ has the same additional term.) Operators of this type, acting on sections of vector bundles without natural positive definite fiber inner products (and correspondingly without symmetry conditions on V_0), appear in the study of the equations of linearized gravity on stationary and asymptotically flat spacetimes in certain gauges, and indeed this was the author’s original motivation for the investigations in the present paper; the details will appear elsewhere.

5C. Scattering for the Dirac–Coulomb equation. Motivated by [Baskin and Wunsch 2023], we consider the stationary scattering theory for the Dirac–Coulomb equation on Minkowski space at high energies. As discussed in Section 1, our framework allows us to deal directly with the associated matrix-valued Klein–Gordon operator — which has nonsymmetric leading-order terms at the Coulomb singularity — albeit with an arbitrarily small loss upon propagation through the singularity. Moreover, our results include a larger range of Coulomb charges $Z \in \mathbb{R}$ than [Baskin and Wunsch 2023] (which requires $|Z| < \frac{1}{2}$ for technical reasons); we can even allow for Z which $|Z| > \frac{\sqrt{3}}{2}$, in which case the Dirac–Coulomb Hamiltonian is not essentially self-adjoint.

¹⁶After the original version of the present paper appeared, this has been worked out in detail by the author in the preprint [Hintz 2023].

The underlying spatial manifold is again given by (5-25), now with $n = 3$. We recall relevant notation from [Baskin and Wunsch 2023]. Denote the Pauli matrices by

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Put further

$$\beta := \gamma^0 := \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \gamma^j := \begin{pmatrix} 0 & \sigma_j \\ -\sigma_j & 0 \end{pmatrix},$$

$$\alpha^j := \beta \gamma^j = \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix}, \quad \alpha_r := \sum_{j=1}^3 \frac{x_j}{r} \alpha_j.$$

The equation governing a massive Dirac field (with mass $m \in \mathbb{R}$) minimally coupled to an electromagnetic potential $\mathbf{A} = (A_0, A_1, A_2, A_3)$ is

$$(i \not{\partial}_{\mathbf{A}} - m) \psi = 0, \quad \not{\partial}_{\mathbf{A}} := \gamma^\mu (\partial_\mu + i A_\mu),$$

where ψ takes values in \mathbb{C}^4 . We now take

$$A_0 = \frac{Z}{r} + V, \quad V \in C^\infty(X), \quad A_j \in C^\infty(X), \tag{5-33}$$

with $Z \in \mathbb{R}$ the charge of the Coulomb field. As shown in [Baskin and Wunsch 2023, §4.3], the operator $-(i \not{\partial}_{\mathbf{A}} + m)(i \not{\partial}_{\mathbf{A}} - m)$ is then of the form

$$P = -\left(D_t + \frac{Z}{r}\right)^2 + \Delta + m^2 + i \frac{Z}{r^2} \alpha_r + r^{-1} \mathbf{R}, \quad \mathbf{R} \in \text{Diff}_b^1(X; \mathbb{C}^4). \tag{5-34}$$

Let us pass to a fixed temporal frequency $\sigma \in \mathbb{C}$, thus replacing D_t in (5-34) by $-\sigma$, resulting in the operator family $\widehat{P}(\sigma)$. Introducing $h = |\sigma|^{-1}$ and $z = (h\sigma)^2$ and multiplying $\widehat{P}(\sigma) = \widehat{P}(h^{-1} \sqrt{z})$ by h^2 gives

$$P_{h,z} = h^2 \Delta - \left(\sqrt{z} - \frac{hZ}{r}\right)^2 + \frac{h^2}{r^2} (iZ\alpha_r + r^2 m^2) + \frac{h^2}{r^2} r \mathbf{R} = h^2 \Delta - z + h^2 r^{-2} Q_{1,z} + h r^{-1} q_{0,z}, \tag{5-35}$$

$$Q_{1,z} = -Z^2 + iZ\alpha_r + r^2 m^2 + r \mathbf{R}, \quad q_{0,z} = 2\sqrt{z}Z.$$

When $\text{Im } \sigma$ is bounded while $|\text{Re } \sigma| \rightarrow \infty$, one has $z = 1 + \mathcal{O}(h)$; thus, $P_{h,z}$ is an admissible operator in the sense of Definition 4.1. The threshold quantities in Definition 4.7 take the values

$$r_{\text{in}} = r_{\text{out}} = -\frac{1}{2}$$

since $q_{0,1}$ is real and the principal symbol of $Q_{1,z}$ (as a first-order b-differential operator) vanishes at $r = 0$.

The normal operator of $P_{h,z}$ is obtained by passing to $\hat{r} = r/h$ and restricting to $h = 0$, giving in polar coordinates $(\hat{r}, \omega) \in [0, \infty) \times \mathbb{S}^2$

$$N(P) = D_{\hat{r}}^2 - \frac{2i}{\hat{r}} D_{\hat{r}} - \left(1 - \frac{Z}{\hat{r}}\right)^2 + \hat{r}^{-2} (\not{\Delta} + iZ\alpha_r(\omega)). \tag{5-36}$$

For $Z = 0$, the operator $N(P)$ is equal to $(\Delta_{\hat{g}} - 1) \otimes \text{Id}_{\mathbb{C}^4}$, where $\hat{g} = d\hat{r}^2 + \hat{r}^2 g_{\mathbb{S}^2}$, and hence is invertible at weight $l \in (\frac{1}{2}, \frac{3}{2})$ by Lemma 5.1. For fixed l , this will remain true for Z in a small neighborhood of 0.

The determination of the largest set of Z for which $N(P)$ is invertible at some weight requires explicit calculations:

Lemma 5.13 (invertibility of $N(P)$). *Let $Z \in \mathbb{R}$ be such that $|Z| \neq \sqrt{\kappa^2 - \frac{1}{4}}$ for all $\kappa \in \mathbb{N}$. Then the operator $N(P)$ is invertible at weight $l = 1$.*

The conclusion of the Lemma in particular holds in the range $|Z| < \frac{\sqrt{3}}{2}$; this is related to the essential self-adjointness of Dirac operators; see [Weidmann 1971; Lesch 1997; Baskin and Wunsch 2023, §4.1].

Proof of Lemma 5.13. We begin by separating into spinor spherical harmonics following [Baskin and Wunsch 2023, §2.1]: For

$$\kappa \in \mathbb{Z} \setminus \{0\}, \quad \mu \in \left\{-|\kappa| + \frac{1}{2}, \dots, |\kappa| - \frac{1}{2}\right\},$$

define the \mathbb{C}^2 -valued function on \mathbb{S}^2

$$\Omega_{\kappa,\mu}(\omega) = \begin{pmatrix} -\operatorname{sgn}(\kappa)\left(\left(\kappa + \frac{1}{2} - \mu\right)/(2\kappa + 1)\right)^{1/2} Y_{l,\mu-1/2}(\omega) \\ \left(\left(\kappa + \frac{1}{2} + \mu\right)/(2\kappa + 1)\right)^{1/2} Y_{l,\mu+1/2}(\omega) \end{pmatrix}, \quad l = \left|\kappa + \frac{1}{2}\right| - \frac{1}{2}.$$

Thus $\mathbb{A}\Omega_{\kappa,\mu} = \kappa(\kappa + 1)\Omega_{\kappa,\mu}$. Moreover,

$$\alpha_r \begin{pmatrix} a\Omega_{\kappa,\mu} \\ b\Omega_{-\kappa,\mu'} \end{pmatrix} = \begin{pmatrix} -b\Omega_{\kappa,\mu'} \\ -a\Omega_{-\kappa,\mu} \end{pmatrix},$$

as follows from [Baskin and Wunsch 2023, equations (3), (4), (9)] or [Szmytkowski 2007, equation (3.1.3)]. Thus, the action of the spherical operator $\mathbb{A} + iZ\alpha_r \in \operatorname{Diff}^2(\mathbb{S}^2; \mathbb{C}^4)$ appearing in (5-36) on the 2-dimensional space with basis $(\Omega_{\kappa,\mu}, 0)$ and $(0, \Omega_{-\kappa,\mu})$ is given by the matrix

$$\begin{pmatrix} \kappa(\kappa + 1) & -iZ \\ -iZ & \kappa(\kappa - 1) \end{pmatrix}. \tag{5-37}$$

This can be diagonalized when $|Z| \neq |\kappa|$, and it has eigenvalue $\lambda_\kappa^\pm = \kappa^2 \pm \sqrt{\kappa^2 - Z^2}$ on the eigenspace spanned by

$$\mathcal{Y}_{\kappa,\mu}^\pm(\omega) := \begin{pmatrix} iZ\Omega_{\kappa,\mu}(\omega) \\ (\kappa \mp \sqrt{\kappa^2 - Z^2})\Omega_{-\kappa,\mu}(\omega) \end{pmatrix}, \quad \mu \in \left\{-|\kappa| + \frac{1}{2}, \dots, |\kappa| - \frac{1}{2}\right\}.$$

Thus, the action of $N(P)$ on separated functions of the form $u(\hat{r})\mathcal{Y}_{\kappa,\mu}^\pm(\omega)$ is given by the action on u of the differential operator

$$N_\kappa^\pm = D_{\hat{r}}^2 - \frac{2i}{\hat{r}}D_{\hat{r}} - \left(1 - \frac{Z}{\hat{r}}\right)^2 + \hat{r}^{-2}\lambda_\kappa^\pm.$$

The Mellin-transformed normal operator of $\hat{r}^2 N_\kappa^\pm$ at $\hat{r} = 0$ is the polynomial

$$\lambda^2 - i\lambda - Z^2 + \lambda_\kappa^\pm, \quad \lambda \in \mathbb{C};$$

for its roots, we have

$$-(\operatorname{Im} \lambda) + \frac{3}{2} \in \left\{1 - \left(\frac{1}{2} \pm \sqrt{\kappa^2 - Z^2}\right), 1 + \left(\frac{1}{2} \pm \sqrt{\kappa^2 - Z^2}\right)\right\}.$$

Now if $Z^2 > \kappa^2$, then these two roots have real parts $\frac{1}{2}$ and $\frac{3}{2}$, whereas if $Z^2 < \kappa^2$, they are disjoint from an open interval $(1 - \delta, 1 + \delta)$ around 1 due to the assumption that $\kappa^2 - Z^2 \neq \frac{1}{4}$.

An outgoing solution of $N(P)$ at weight $l = 1$, expanded into the spherical eigenfunctions $\mathcal{Y}_{\kappa,\mu}^\pm$, is an outgoing solution of N_κ^\pm ; one easily finds $u = u_\infty \hat{r}^{-1-iZ} e^{i\hat{r}} + \mathcal{O}(\hat{r}^{-2})$ as $\hat{r} \rightarrow \infty$, where $u_\infty \in \mathbb{C}$, and the $\mathcal{O}(\hat{r}^{-2})$ term is conormal at $\hat{r} = 0$; near $\hat{r} = 0$ on the other hand, we have $u = \mathcal{A}^{-1/2+\delta}([0, 1]_{\hat{r}})$. A boundary-pairing argument, i.e., the evaluation of

$$0 = \lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{1/\epsilon} ((N_\kappa^\pm u)\bar{u} - u\overline{N_\kappa^\pm u}) \hat{r}^2 d\hat{r} = 2i \lim_{\epsilon \rightarrow 0} \text{Im}(\hat{r}^2 u \bar{u}'|_{\epsilon}^{1/\epsilon}) = -2i |u_\infty|^2,$$

gives $u_\infty = 0$, and thus $u \equiv 0$ by standard ODE analysis near $\hat{r} = \infty$. This shows that $N(P)$ is injective at weight $l = 1$ on outgoing solutions. Since $(N_\kappa^\pm)^* = N_\kappa^\mp$ with respect to the $L^2(\hat{r}^2 d\hat{r})$ inner product, the injectivity of $N(P)^*$ at weight $-l + 2$ on incoming functions is proved similarly. This completes the proof when Z is not a nonzero integer.

When $Z \in \mathbb{Z} \setminus \{0\}$ and κ satisfies $|Z| = |\kappa|$, then the action of $\mathcal{X} + iZ\alpha_r$ on the span of $(\Omega_{\kappa,\mu}, 0)$ and $(0, \Omega_{-\kappa,\mu'})$ is not diagonalizable anymore. By inspection of (5-37), it still has the eigenvalue κ^2 with eigenspace spanned by $\mathcal{Y}_{\kappa,\mu}^+ = \mathcal{Y}_{\kappa,\mu}^-$. Let $\tilde{\mathcal{Y}}_{\kappa,\mu} = (\Omega_{\kappa,\mu}, 0)$, then an outgoing solution $u = u_1 \mathcal{Y}_{\kappa,\mu}^+ + u_2 \tilde{\mathcal{Y}}_{\kappa,\mu}$ of $N(P)$ satisfies a lower triangular ODE system, with a decoupled equation for u_1 which implies $u_1 \equiv 0$ by the previous arguments, whence u_2 is now an outgoing solution to the same equation as u_1 and must therefore also vanish. □

If we cut \mathbf{A} off via multiplication by a cutoff $\chi \in C_c^\infty(X)$, the operator $P_{h,z}$ is equal to $h^2 \Delta - z$ near infinity and can thus be analyzed as in Section 5B. In this setting, we thus obtain invertibility and quantitative estimates for $P_{h,z}$:

Theorem 5.14 (high-energy estimates for the Dirac–Coulomb equation). *Suppose $\mathbf{A} = (A_0, A_1, A_2, A_3)$ as in (5-33) has compact support. Let $Z \in \mathbb{R}$ be such that $|Z| \neq \sqrt{\kappa^2 - \frac{1}{4}}$ for all $\kappa \in \mathbb{N}$. Then for $l = 1$ (and indeed for l sufficiently close to 1), $0 < \text{Im } z < Ch$, and for all sufficiently small $h > 0$, the operator $P_{h,z} = h^2 \hat{P}(h^{-1}z)$ defined in (5-34)–(5-35) is invertible as a map between the spaces (5-29) and satisfies the uniform bound (5-30) as well as the bound (5-31) (with $\Delta_g + V - \sigma^2$ replaced by $\hat{P}(\sigma)$, where $\sigma = h^{-1}z$) for $l_1 = l_2 = 0$.*

Remark 5.15 (complex charges). One can also analyze the case of nonreal $Z \in \mathbb{C}$, in which case $r_{\text{in}} = -\frac{1}{2} + \text{Im } Z$ and $r_{\text{out}} = -\frac{1}{2} - \text{Im } Z$. The difference $r_{\text{out}} - r_{\text{in}} = -2 \text{Im } Z$ results in an additional $2 \text{Im } Z$ loss of powers of the semiclassical parameter h when propagating through the singularity at $r = 0$. Nonetheless, the invertibility of $N(P)$ automatically holds for values of Z close to those allowed in Theorem 5.14, as discussed prior to Lemma 5.10.

Appendix A: A class of examples with sharp semiclassical loss

Note that the semiclassical order b in Theorem 4.10 must decrease from \mathcal{R}_{in} to \mathcal{R}_{out} by more than

$$D = \max(r_{\text{in}} - r_{\text{out}}, 0); \tag{A-1}$$

thus, the estimate (4-27) controls u in L^2 , say, microlocally near the flow-out of \mathcal{R}_{out} by $h^{-D-\epsilon}$ (for any $\epsilon > 0$) times the L^2 -norm of the microlocalization of u near the flow-in of \mathcal{R}_{in} . While in many natural settings, such as those discussed in Section 5, one has $D = 0$, it is easy to construct examples where $D > 0$. The following example (placed into a general context at the end of this appendix) shows that a loss of h^{-D} typically *does* occur, whence our estimates are sharp up to an ϵ -loss. This ϵ -loss may be avoidable, though we are not able to prove or disprove this here.

Consider $X = [0, 2)_r$, $\mu = |dr|$, and

$$P_{h,z} = h^2 \left(-\partial_r^2 - \frac{2}{r} \partial_r \right) - z + \frac{2ih}{r} q, \quad z = 1 + \mathcal{O}(h),$$

where $q \in \mathbb{C}$ is a parameter. (The term in parentheses is the radial part of the Laplacian on \mathbb{R}^3 in polar coordinates.) The normal operator is

$$N(P) = -\partial_{\hat{r}}^2 - \frac{2}{\hat{r}} \partial_{\hat{r}} - 1 + \frac{2i}{\hat{r}} q.$$

For $q = 0$, the kernel of $N(P)$ is spanned by $\hat{r}^{-1} e^{\pm i\hat{r}}$; since \hat{r}^{-1} barely fails to lie in $\hat{r}^{-1/2} L^2([0, 1)_{\hat{r}}, |d\hat{r}|)$, it is easy to see that $N(P)$ is invertible at weight l in the sense of Definition 4.6 for $l \in (-\frac{1}{2}, \frac{1}{2})$; this persists for small values of $q \in \mathbb{C}$. (The boundary spectrum of $N(P)$ at $\hat{r} = 0$ is independent of q .) In the notation of Definition 4.7, we have

$$r_{\text{in}} = \frac{1}{2} + \text{Re } q, \quad r_{\text{out}} = \frac{1}{2} - \text{Re } q, \tag{A-2}$$

so $D = \max(2 \text{Re } q, 0)$. The quantities (A-2) correspond precisely to the L^2 -decay rates of incoming and outgoing solutions $\hat{u}_{\text{in}}, \hat{u}_{\text{out}} \in \ker N(P)$, which have the asymptotic behavior

$$\hat{u}_{\text{in}} \sim \hat{r}^{-1-\text{Re } q} e^{-i\hat{r}}, \quad \hat{u}_{\text{out}} \sim \hat{r}^{-1+\text{Re } q} e^{i\hat{r}}, \quad \hat{r} \rightarrow \infty. \tag{A-3}$$

(We omit the explicit expressions involving confluent hypergeometric functions.)

We can now construct an element $\hat{u} \in \ker N(P)$ which lies in $\hat{r}^l L^2$, $l \in (-\frac{1}{2}, \frac{1}{2})$, near $\hat{r} = 0$ and which is uniquely specified by requiring its incoming data at $\hat{r} = \infty$ to be given by \hat{u}_{in} . Indeed, \hat{u} is necessarily of the form

$$\hat{u}(\hat{r}) = \hat{u}_{\text{in}} + c \hat{u}_{\text{out}},$$

where $c \in \mathbb{C}$ is the ‘‘scattering matrix’’; necessarily $c \neq 0$ (since \hat{u}_{in} fails to lie in $\hat{r}^l L^2$ near $\hat{r} = 0$). But then

$$P_{h,1} u_h(r) = 0, \quad u_h(r) := \hat{u}(r/h).$$

(This exact formula uses the invariance of $P_{h,1}$ under dilations in (h, r) .) Considering a neighborhood of $r = 1$ then, the asymptotics (A-3) for

$$u_{\bullet,h}(r) := u_{\bullet}(r/h), \quad \bullet = \text{in, out},$$

imply

$$u_h = u_{\text{in},h} + c u_{\text{out},h},$$

$$u_{\text{in},h} \sim h^{1+\text{Re } q} e^{-ir/h}, \quad u_{\text{out},h} \sim h^{1-\text{Re } q} e^{ir/h} \quad (\text{near } r = 1).$$

This demonstrates the loss of $h^{-2\text{Re } q}$ between the amplitudes $h^{1+\text{Re } q}$, resp. $h^{1-\text{Re } q}$ of the incoming, resp. outgoing pieces of u_h . (The fact that there is in fact a gain when $\text{Re } q < 0$ is a peculiar feature of the 1-dimensional situation considered here: the characteristic set of $P_{h,z}$ has two connected components, with the incoming and outgoing radial sets lying in different components, and the monotonicity requirement in Theorem 4.10 does not relate the two components.)

The same idea can be applied to produce many more examples with sharp loss D . Indeed, when $N(P)$ is invertible at weight l , then the solution $\hat{u} = \hat{u}(\hat{r}, y)$ (with y denoting points on ∂X) of $N(P)\hat{u} = 0$, where \hat{u} has prescribed incoming data and lies in $\hat{r}^l L^2$ near $\hat{r} = 0$, gives rise to a solution $u_h(r, y) = \hat{u}(r/h, y)$ of $P_{h,1}u_h(r, y) = 0$, where $P_{h,1} = N(P)$ (upon changing coordinates $\hat{r} = r/h$). The relative decay rates of incoming/outgoing solutions of $N(P)$ are then directly reflected in the relative semiclassical orders of u_h near the flow-in/flow-out of the cone point. (Since the characteristic set of $P_{h,1}$ is connected when $\dim X \geq 2$, the loss is at least 0, see (A-1); after all, even away from the cone point, semiclassical regularity cannot improve under real-principal-type propagation.)

Appendix B: Sobolev spaces and pseudodifferential calculi

This section consists entirely of figures which illustrate the orders of weighted Sobolev spaces as well as of spaces of pseudodifferential operators, with references to the original definitions. Concretely, labeling a boundary hypersurface H by “ l ” means that the order l of some weighted Sobolev space $H^{\dots;l,\dots}$ refers to ρ_H^l decay at H of its elements, where ρ_H is a defining function of H , or l orders of regularity when H is a boundary hypersurface at fiber infinity of some compactified phase space. For spaces of pseudodifferential operators on the other hand, the same label “ l ” refers to a ρ_H^{-l} bound of the full symbol of the operator, or of its Schwartz kernel at the hypersurface of the double space corresponding to H .

See Figure 13 for b- and scattering Sobolev spaces (or operators), Figure 14 for b, sc-Sobolev spaces, Figure 15 for semiclassical cone Sobolev spaces, and Figure 16 for cb \hbar -Sobolev spaces.




Figure 13. X is a manifold with boundary ∂X . Left: the orders of $H_b^{s,l}(X)$; see Section 2A. Right: the orders of $H_{sc}^{s,r}(X)$; see Section 2B.




Figure 14. X is a manifold with two connected and embedded boundary hypersurfaces $\partial X = H_1 \sqcup H_2$. Indicated are the orders for $H_{b,sc}^{s,l,r}(X)$.




Figure 15. X is a manifold with boundary ∂X . Indicated are the orders for $H_{c,h}^{s,l,\alpha,b}(X)$; see Section 3B.




Figure 16. This is essentially a repetition of Figure 9. Indicated are the orders for $H_{cb,h}^{s',l,\alpha,b}(X)$; see Section 3D.

Acknowledgments

I am grateful to András Vasy and Jared Wunsch for helpful conversations, and to two anonymous referees for their careful reading of the paper and for excellent comments and suggestions. I gratefully acknowledge support from a Sloan Research Fellowship and from the NSF under grant no. DMS-1955614.

References

- [Baskin and Marzuola 2019] D. Baskin and J. L. Marzuola, “Locating resonances on hyperbolic cones”, *Math. Res. Lett.* **26**:2 (2019), 365–381. MR Zbl
- [Baskin and Marzuola 2022] D. Baskin and J. L. Marzuola, “The radiation field on product cones”, *Adv. Math.* **408** (2022), art.id. 108589. MR Zbl
- [Baskin and Wunsch 2013] D. Baskin and J. Wunsch, “Resolvent estimates and local decay of waves on conic manifolds”, *J. Differential Geom.* **95**:2 (2013), 183–214. MR Zbl
- [Baskin and Wunsch 2023] D. Baskin and J. Wunsch, “Diffraction for the Dirac–Coulomb propagator”, *Ann. Henri Poincaré* **24**:8 (2023), 2607–2659. MR Zbl
- [Baskin et al. 2015] D. Baskin, A. Vasy, and J. Wunsch, “Asymptotics of radiation fields in asymptotically Minkowski space”, *Amer. J. Math.* **137**:5 (2015), 1293–1364. MR Zbl
- [Baskin et al. 2018] D. Baskin, A. Vasy, and J. Wunsch, “Asymptotics of scalar waves on long-range asymptotically Minkowski spaces”, *Adv. Math.* **328** (2018), 160–216. MR Zbl

- [Beals 1977] R. Beals, “Characterization of pseudodifferential operators and applications”, *Duke Math. J.* **44**:1 (1977), 45–57. MR Zbl
- [Bony 1986] J.-M. Bony, “Second microlocalization and propagation of singularities for semilinear hyperbolic equations”, pp. 11–49 in *Hyperbolic equations and related topics* (Katata/Kyoto, 1984), edited by S. Mizohata, Academic Press, Boston, MA, 1986. MR Zbl
- [Burq et al. 2003] N. Burq, F. Planchon, J. G. Stalker, and A. S. Tahvildar-Zadeh, “Strichartz estimates for the wave and Schrödinger equations with the inverse-square potential”, *J. Funct. Anal.* **203**:2 (2003), 519–549. MR Zbl
- [Cheeger and Taylor 1982a] J. Cheeger and M. Taylor, “On the diffraction of waves by conical singularities, I”, *Comm. Pure Appl. Math.* **35**:3 (1982), 275–331. MR Zbl
- [Cheeger and Taylor 1982b] J. Cheeger and M. Taylor, “On the diffraction of waves by conical singularities, II”, *Comm. Pure Appl. Math.* **35**:4 (1982), 487–529. MR Zbl
- [Chin 2022] X. Chin, “The semiclassical resolvent on conic manifolds and application to Schrödinger equations”, *Commun. Math. Phys.* **390**:2 (2022), 757–826. Zbl
- [Datchev and Vasy 2012] K. Datchev and A. Vasy, “Gluing semiclassical resolvent estimates via propagation of singularities”, *Int. Math. Res. Not.* **2012**:23 (2012), 5409–5443. MR Zbl
- [Duistermaat and Hörmander 1972] J. J. Duistermaat and L. Hörmander, “Fourier integral operators, II”, *Acta Math.* **128**:3–4 (1972), 183–269. MR Zbl
- [Duyckaerts 2006] T. Duyckaerts, “Inégalités de résolvente pour l’opérateur de Schrödinger avec potentiel multipolaire critique”, *Bull. Soc. Math. France* **134**:2 (2006), 201–239. MR Zbl
- [Dyatlov and Zworski 2019] S. Dyatlov and M. Zworski, *Mathematical theory of scattering resonances*, Grad. Stud. in Math. **200**, Amer. Math. Soc., Providence, RI, 2019. MR Zbl
- [Friedlander 1958] F. G. Friedlander, *Sound pulses*, Cambridge Univ. Press, 1958. MR Zbl
- [Galkowski 2017] J. Galkowski, “A quantitative Vainberg method for black box scattering”, *Comm. Math. Phys.* **349**:2 (2017), 527–549. MR Zbl
- [Gannot and Wunsch 2023] O. Gannot and J. Wunsch, “Semiclassical diffraction by conormal potential singularities”, *Ann. Sci. École Norm. Sup. (4)* **56**:3 (2023), 713–800. MR
- [Gell-Redman et al. 2020] J. Gell-Redman, A. Hassell, J. Shapiro, and J. Zhang, “Existence and asymptotics of nonlinear Helmholtz eigenfunctions”, *SIAM J. Math. Anal.* **52**:6 (2020), 6180–6221. MR Zbl
- [Grieser 2001] D. Grieser, “Basics of the b -calculus”, pp. 30–84 in *Approaches to singular analysis* (Berlin, 1999), edited by J. Gil et al., Oper. Theory Adv. Appl. **125**, Birkhäuser, Basel, 2001. MR Zbl
- [Guillarmou et al. 2013] C. Guillarmou, A. Hassell, and A. Sikora, “Resolvent at low energy, III: The spectral measure”, *Trans. Amer. Math. Soc.* **365**:11 (2013), 6103–6148. MR Zbl
- [Haber and Vasy 2013] N. Haber and A. Vasy, “Propagation of singularities around a Lagrangian submanifold of radial points”, pp. 113–116 in *Microlocal methods in mathematical physics and global analysis*, edited by D. Grieser et al., Birkhäuser, Basel, 2013. MR Zbl
- [Hassell et al. 2008] A. Hassell, R. Melrose, and A. Vasy, “Microlocal propagation near radial points and scattering for symbolic potentials of order zero”, *Anal. PDE* **1**:2 (2008), 127–196. MR Zbl
- [Helffer and Sjöstrand 1989] B. Helffer and J. Sjöstrand, “Équation de Schrödinger avec champ magnétique et équation de Harper”, pp. 118–197 in *Schrödinger operators* (Sønderborg, Denmark, 1988), edited by H. Holden and A. Jensen, Lecture Notes in Phys. **345**, Springer, 1989. MR Zbl
- [Hillairet and Wunsch 2020] L. Hillairet and J. Wunsch, “On resonances generated by conic diffraction”, *Ann. Inst. Fourier (Grenoble)* **70**:4 (2020), 1715–1752. MR Zbl
- [Hintz 2022] P. Hintz, “Resolvents and complex powers of semiclassical cone operators”, *Math. Nachr.* **295**:10 (2022), 1990–2035. MR Zbl
- [Hintz 2023] P. Hintz, “Linear waves on asymptotically flat spacetimes, I”, preprint, 2023. arXiv 2302.14647
- [Hirsch et al. 1977] M. W. Hirsch, C. C. Pugh, and M. Shub, *Invariant manifolds*, Lecture Notes in Math. **583**, Springer, 1977. MR Zbl

- [de Hoop et al. 2015] M. de Hoop, G. Uhlmann, and A. Vasy, “Diffraction from conormal singularities”, *Ann. Sci. École Norm. Sup.* (4) **48**:2 (2015), 351–408. MR Zbl
- [Hörmander 1971] L. Hörmander, “Fourier integral operators, I”, *Acta Math.* **127**:1-2 (1971), 79–183. MR Zbl
- [Keeler and Marzuola 2022] B. Keeler and J. L. Marzuola, “Pointwise dispersive estimates for Schrödinger operators on product cones”, *J. Differential Equations* **320** (2022), 419–468. MR Zbl
- [Keller 1985] J. B. Keller, “One hundred years of diffraction theory”, *IEEE Trans. Antennas and Propagation* **33**:2 (1985), 123–126. MR Zbl
- [Lebeau 1997] G. Lebeau, “Propagation des ondes dans les variétés à coins”, *Ann. Sci. École Norm. Sup.* (4) **30**:4 (1997), 429–497. MR Zbl
- [Lesch 1997] M. Lesch, *Operators of Fuchs type, conical singularities, and asymptotic methods*, Teubner-Texte Math. **136**, Teubner, Stuttgart, 1997. MR Zbl
- [Loya 2002] P. Loya, “On the resolvent of differential operators on conic manifolds”, *Comm. Anal. Geom.* **10**:5 (2002), 877–934. MR Zbl
- [Mazzeo 1991] R. Mazzeo, “Elliptic theory of differential edge operators, I”, *Comm. Partial Differential Equations* **16**:10 (1991), 1615–1664. MR Zbl
- [Melrose 1975] R. B. Melrose, “Microlocal parametrices for diffractive boundary value problems”, *Duke Math. J.* **42**:4 (1975), 605–635. MR Zbl
- [Melrose 1993] R. B. Melrose, *The Atiyah–Patodi–Singer index theorem*, Res. Notes in Math. **4**, Peters, Wellesley, MA, 1993. MR Zbl
- [Melrose 1994] R. B. Melrose, “Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces”, pp. 85–130 in *Spectral and scattering theory* (Sanda, Japan, 1992), edited by M. Ikawa, Lecture Notes in Pure Appl. Math. **161**, Dekker, New York, 1994. MR Zbl
- [Melrose 1995] R. B. Melrose, *Geometric scattering theory*, Cambridge Univ. Press, 1995. MR Zbl
- [Melrose 1996] R. B. Melrose, “Differential analysis on manifolds with corners”, book in preparation, 1996, available at <https://math.mit.edu/~rbm/book.html>.
- [Melrose and Sjöstrand 1978] R. B. Melrose and J. Sjöstrand, “Singularities of boundary value problems, I”, *Comm. Pure Appl. Math.* **31**:5 (1978), 593–617. MR Zbl
- [Melrose and Sjöstrand 1982] R. B. Melrose and J. Sjöstrand, “Singularities of boundary value problems, II”, *Comm. Pure Appl. Math.* **35**:2 (1982), 129–168. MR Zbl
- [Melrose and Taylor 2018] R. B. Melrose and M. E. Taylor, “Boundary problems for wave equations with grazing and gliding rays”, preprint, 2018, available at <http://mtaylor.web.unc.edu/wp-content/uploads/sites/16915/2018/04/glide.pdf>.
- [Melrose and Wunsch 2004] R. Melrose and J. Wunsch, “Propagation of singularities for the wave equation on conic manifolds”, *Invent. Math.* **156**:2 (2004), 235–299. MR Zbl
- [Melrose and Zworski 1996] R. Melrose and M. Zworski, “Scattering metrics and geodesic flow at infinity”, *Invent. Math.* **124**:1-3 (1996), 389–436. MR Zbl
- [Melrose et al. 2008] R. Melrose, A. Vasy, and J. Wunsch, “Propagation of singularities for the wave equation on edge manifolds”, *Duke Math. J.* **144**:1 (2008), 109–193. MR Zbl
- [Melrose et al. 2013] R. Melrose, A. Vasy, and J. Wunsch, *Diffraction of singularities for the wave equation on manifolds with corners*, Astérisque **351**, Soc. Math. France, Paris, 2013. MR Zbl
- [Mourre 1980/81] E. Mourre, “Absence of singular continuous spectrum for certain selfadjoint operators”, *Comm. Math. Phys.* **78**:3 (1980/81), 391–408. MR Zbl
- [Nonnenmacher and Zworski 2009] S. Nonnenmacher and M. Zworski, “Quantum decay rates in chaotic scattering”, *Acta Math.* **203**:2 (2009), 149–233. MR Zbl
- [Planchon et al. 2003] F. Planchon, J. G. Stalker, and A. S. Tahvildar-Zadeh, “Dispersive estimate for the wave equation with the inverse-square potential”, *Discrete Contin. Dyn. Syst.* **9**:6 (2003), 1387–1400. MR Zbl
- [Qian 2009] R. Z. Qian, *Diffractive theorems for the wave equation with inverse square potentials*, Ph.D. thesis, Northwestern University, 2009, available at <https://www.proquest.com/docview/304971420>.

- [Sjöstrand and Zworski 1991] J. Sjöstrand and M. Zworski, “Complex scaling and the distribution of scattering poles”, *J. Amer. Math. Soc.* **4**:4 (1991), 729–769. MR Zbl
- [Sjöstrand and Zworski 2007] J. Sjöstrand and M. Zworski, “Fractal upper bounds on the density of semiclassical resonances”, *Duke Math. J.* **137**:3 (2007), 381–459. MR Zbl
- [Sommerfeld 1896] A. Sommerfeld, “Mathematische Theorie der Diffraction”, *Math. Ann.* **47**:2-3 (1896), 317–374. MR Zbl
- [Szmytkowski 2007] R. Szmytkowski, “Recurrence and differential relations for spherical spinors”, *J. Math. Chem.* **42**:3 (2007), 397–413. MR Zbl
- [Taylor 1976] M. E. Taylor, “Grazing rays and reflection of singularities of solutions to wave equations”, *Comm. Pure Appl. Math.* **29**:1 (1976), 1–38. MR Zbl
- [Vasy 2000] A. Vasy, *Propagation of singularities in three-body scattering*, Astérisque **262**, Soc. Math. France, Paris, 2000. MR Zbl
- [Vasy 2001] A. Vasy, “Propagation of singularities in many-body scattering”, *Ann. Sci. École Norm. Sup. (4)* **34**:3 (2001), 313–402. MR Zbl
- [Vasy 2008] A. Vasy, “Propagation of singularities for the wave equation on manifolds with corners”, *Ann. of Math. (2)* **168**:3 (2008), 749–812. MR Zbl
- [Vasy 2013] A. Vasy, “Microlocal analysis of asymptotically hyperbolic and Kerr–de Sitter spaces”, *Invent. Math.* **194**:2 (2013), 381–513. MR Zbl
- [Vasy 2018] A. Vasy, “A minicourse on microlocal analysis for wave propagation”, pp. 219–374 in *Asymptotic analysis in general relativity* (Grenoble, France, 2014), edited by T. Daudé et al., Lond. Math. Soc. Lect. Note Ser. **443**, Cambridge Univ. Press, 2018. MR Zbl
- [Vasy 2021a] A. Vasy, “Limiting absorption principle on Riemannian scattering (asymptotically conic) spaces, a Lagrangian approach”, *Comm. Partial Differential Equations* **46**:5 (2021), 780–822. MR Zbl
- [Vasy 2021b] A. Vasy, “Resolvent near zero energy on Riemannian scattering (asymptotically conic) spaces”, *Pure Appl. Anal.* **3**:1 (2021), 1–74. MR Zbl
- [Vasy 2021c] A. Vasy, “Resolvent near zero energy on Riemannian scattering (asymptotically conic) spaces, a Lagrangian approach”, *Comm. Partial Differential Equations* **46**:5 (2021), 823–863. MR Zbl
- [Vasy and Wunsch 2009] A. Vasy and J. Wunsch, “Semiclassical second microlocal propagation of regularity and integrable systems”, *J. Anal. Math.* **108** (2009), 119–157. Correction in **115** (2011), 389–391. MR Zbl
- [Vasy and Zworski 2000] A. Vasy and M. Zworski, “Semiclassical estimates in asymptotically Euclidean scattering”, *Comm. Math. Phys.* **212**:1 (2000), 205–217. MR Zbl
- [Wang 2020] J. Wang, “Sharp radial estimates in Besov spaces”, preprint, 2020. arXiv 2003.11218
- [Weidmann 1971] J. Weidmann, “Oszillationsmethoden für Systeme gewöhnlicher Differentialgleichungen”, *Math. Z.* **119** (1971), 349–373. MR Zbl
- [Wunsch and Zworski 2011] J. Wunsch and M. Zworski, “Resolvent estimates for normally hyperbolic trapped sets”, *Ann. Henri Poincaré* **12**:7 (2011), 1349–1385. MR Zbl
- [Yang 2022] M. Yang, “Propagation of polyhomogeneity, diffraction, and scattering on product cones”, *J. Spectr. Theory* **12**:2 (2022), 381–415. MR Zbl

Received 28 May 2022. Revised 14 May 2023. Accepted 18 Jul 2023.

PETER HINTZ: phintz@mit.edu, peter.hintz@math.ethz.ch

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States

and

Department of Mathematics, ETH Zürich, Zürich, Switzerland

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 10 2024

Scattering of the three-dimensional cubic nonlinear Schrödinger equation with partial harmonic potentials	3371
XING CHENG, CHANG-YU GUO, ZIHUA GUO, XIAN LIAO and JIA SHEN	
On Gagliardo–Nirenberg inequalities with vanishing symbols	3447
RAINER MANDEL	
Semiclassical propagation through cone points	3477
PETER HINTZ	
Small cap decoupling for the moment curve in \mathbb{R}^3	3551
LARRY GUTH and DOMINIQUE MALDAGUE	
The quermassintegral-preserving mean curvature flow in the sphere	3589
ESTHER CABEZAS-RIVAS and JULIAN SCHEUER	
Upper bound on the number of resonances for even asymptotically hyperbolic manifolds with real-analytic ends	3623
MALO JÉZÉQUEL	