

ANALYSIS & PDE

Volume 17

No. 8

2024

CAMIL MUSCALU AND ITAMAR OLIVEIRA

**A NEW APPROACH TO THE FOURIER EXTENSION PROBLEM
FOR THE PARABOLOID**



A NEW APPROACH TO THE FOURIER EXTENSION PROBLEM FOR THE PARABOLOID

CAMIL MUSCALU AND ITAMAR OLIVEIRA

Dedicated to the memory of Robert S. Strichartz

We propose a new approach to the Fourier restriction conjectures. It is based on a discretization of the Fourier extension operators in terms of quadratically modulated wave packets. Using this new point of view, and by combining natural scalar and mixed norm quantities from appropriate level sets, we prove that all the L^2 -based k -linear extension conjectures are true up to the endpoint for every $1 \leq k \leq d + 1$ if one of the functions involved is a full tensor. We also introduce the concept of *weak transversality*, under which we show that all conjectured L^2 -based multilinear extension estimates are still true up to the endpoint, provided that one of the functions involved has a weaker tensor structure, and we prove that this result is sharp. Under additional tensor hypotheses, we show that one can improve the conjectured threshold of these problems in some cases. In general, the largely unknown multilinear extension theory beyond L^2 inputs remains open even in the bilinear case; with this new point of view, and still under the previous tensor hypothesis, we obtain the near-restriction target for the k -linear extension operator if the inputs are in a certain L^p space for p sufficiently large. The proof of this result is adapted to show that the k -fold product of linear extension operators (no transversality assumed) also “maps near restriction” if one input is a tensor. Finally, we exploit the connection between the geometric features behind the results of this paper and the theory of Brascamp–Lieb inequalities, which allows us to verify a special case of a conjecture by Bennett, Bez, Flock and Lee.

1. Introduction	2842
2. Discrete models	2846
3. Transversality versus weak transversality	2850
4. Our approach and its building blocks	2852
5. Proof of Proposition 4.1: Strichartz/Tomas–Stein for E_d ($k = 1, p = 2$)	2854
6. Proof of Proposition 4.3-Conjecture 1.1 for E_1 ($k = 1, d = 1, p = 4$)	2857
7. Proof of Proposition 4.4-Conjecture 1.2 for $ME_{2,1}$ ($k = 2, d = 1$)	2859
8. Case $k = 1$ of Theorem 1.5	2860
9. Case $2 \leq k \leq d + 1$ of Theorem 1.5	2862
10. The endpoint estimate of the case $k = d + 1$ of Theorem 1.5	2876
11. Improved k -linear bounds for tensors	2878
12. Beyond the L^2 -based k -linear theory with and without transversality	2884
13. Weak transversality, Brascamp–Lieb and an application	2895
14. Further remarks	2903
Appendix A. Sharp examples	2905
Appendix B. Technical results	2913
Acknowledgements	2919
References	2919

MSC2020: 42B10, 42B37.

Keywords: Fourier restriction, extension operator, multilinear restriction, transversality, weak transversality.

1. Introduction

Given a compact submanifold $S \subset \mathbb{R}^{d+1}$ and a function $f : \mathbb{R}^{d+1} \mapsto \mathbb{R}$, the *Fourier restriction problem* asks for which pairs (p, q) one has

$$\|\hat{f}|_S\|_{L^q(S)} \lesssim \|f\|_{L^p(\mathbb{R}^{d+1})},$$

where $\hat{f}|_S$ is the restriction of the Fourier transform \hat{f} to S . This problem arises naturally in the study of certain Fourier summability methods and is known to be connected to questions in geometric measure theory and in nonlinear dispersive PDEs. The interaction between curvature and the Fourier transform has been exploited in a variety of contexts since the works [Hörmander 1973; Fefferman 1971; Stein and Wainger 1978] in the study of oscillatory integrals. For a more detailed description of the restriction problem we refer the reader to the classical survey [Tao 2004]. In this paper we work with the equivalent dual formulation of the question above (known as the *Fourier extension problem*), and specialize to the case where S is the compact piece of the paraboloid parametrized by $\Gamma(x) = (x, |x|^2) \subset \mathbb{R}^{d+1}$ with $x \in [0, 1]^d$. In this setting, the *Fourier extension operator* is initially defined on $C([0, 1]^d)$ by

$$\mathcal{E}_d g(x_1, \dots, x_d, t) = \int_{[0,1]^d} g(\xi_1, \dots, \xi_d) e^{-2\pi i(\xi_1 x_1 + \dots + \xi_d x_d)} e^{-2\pi i t(\xi_1^2 + \dots + \xi_d^2)} d\xi. \tag{1}$$

E. Stein [1993, Chapter IX] proposed the following conjecture:

Conjecture 1.1. *The inequality*

$$\|\mathcal{E}_d g\|_{L^q(\mathbb{R}^{d+1})} \lesssim_{p,q,d} \|g\|_{L^p([0,1]^d)} \tag{2}$$

holds if and only if $q > \frac{2(d+1)}{d}$ and $q \geq \frac{(d+2)}{d} p'$.

Multilinear variants¹ of Conjecture 1.1 arose naturally from the works [Klainerman and Machedon 1993; 1995; 1996] on wellposedness of certain PDEs. Given $2 \leq k \leq d + 1$ compact and connected domains $U_j \subset \mathbb{R}^d$, $1 \leq j \leq k$, define

$$\mathcal{E}_{U_j} g(x, t) := \int_{U_j} g(\xi) e^{-2\pi i x \cdot \xi} e^{-2\pi i t|\xi|^2} d\xi, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R}. \tag{3}$$

Taking the product of all k such operators associated to a set of *transversal* U_j leads to the following conjecture (see Appendix A):

Conjecture 1.2 [Bennett 2014]. *If the caps parametrized by U_j are transversal, then*

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_p \lesssim \prod_{j=1}^k \|g_j\|_2 \quad \text{for all } p \geq \frac{2(d+k+1)}{k(d+k-1)}.$$

Roughly, *transversality* means that any choice of one normal vector per cap is a set of linearly independent vectors, as shown in Figure 1.

Remark 1.3. From now on, we shall refer to Conjecture 1.1 as *the case $k = 1$* . It was settled only for $d = 1$ in [Fefferman 1970; Zygmund 1974]. In higher dimensions we highlight the case $p = 2$ solved in [Strichartz 1977], which is equivalent to the Tomas–Stein theorem [Tomas 1975] in the restriction setting.

¹Multilinear extension estimates also play a fundamental role in Bourgain and Demeter’s decoupling theory [2015].




Figure 1. A choice of normal vectors to the caps parametrized by U_j via $x \mapsto |x|^2$.

Progress beyond these two results was made in many works over the last decades through a diverse set of techniques: localization, bilinear estimates, wave-packet decompositions and more recently polynomial methods. We mention [Bourgain 1991; Tao and Vargas 2000; Tao 2003; Moyua et al. 1996; Wang 2018; Guth 2018; Hickman and Rogers 2019]. Analogous problems for other manifolds were studied in [Wolff 2001; Strichartz 1977; Ou and Wang 2022].

Remark 1.4. Guth [2018] proved a weaker version of Conjecture 1.2 for all $2 \leq k \leq d + 1$ and up to the endpoint, which is known as the k -broad restriction inequality. This estimate plays a central role in his argument in [Guth 2018] to improve the range for which Conjecture 1.1 is known. In Lemma A.3 of [Bourgain and Guth 2011], the authors proved an L^2 -based k -linear estimate for an exponent p slightly larger than the conjectured threshold in Conjecture 1.2.

Only three cases of Conjecture 1.2 are well understood:

- (i) Tao [2003] settled the case $k = 2$ up to the endpoint inspired by [Wolff 2001] for the cone. Lee [2021] obtained the endpoint for $k = 2$.
- (ii) Bennett, Carbery and Tao [Bennett et al. 2006] settled the case $k = d + 1$ up to the endpoint.
- (iii) Bejenaru [2022] settled the case $k = d$ up to the endpoint.

The goal of this paper is to propose a new approach to these problems based on a natural discretization of the operators in terms of scalar products against quadratically modulated wave-packets. Our main theorem reads as follows:

Theorem 1.5. *Conjectures 1.1 and 1.2 hold up to the endpoint if one (any) of the functions involved is a full tensor.²*

Remark 1.6. The endpoint $(p, q) = (\frac{2(d+1)}{d}, \frac{2(d+1)}{d})$ is not included in the range where (2) is supposed to hold; therefore our main theorem implies the case $k = 1$ when g is a full tensor.

Remark 1.7. For $2 \leq k \leq d + 1$, Theorem 1.5 can be proved if the caps are assumed to be *weakly transversal*, which is defined in Section 3. We will prove that transversality implies weak transversality (up to dividing the caps into finitely many pieces), the latter being what is actually exploited in this paper.

²A function g in d variables is a *full tensor* if it can be written as $g(x_1, \dots, x_d) = g_1(x_1) \cdots g_d(x_d)$. We refer the reader to [Igari 1986; Tanaka 2001] for other results related to the restriction problem involving tensors, and we thank Terence Tao for pointing these papers out to us.

Under weak transversality, [Theorem 1.5](#) holds if one (any) of the functions has a weaker tensor structure. This will be made precise in [Section 9](#).

Remark 1.8. For $2 \leq k \leq d + 1$, [Theorem 1.5](#) is sharp under weak transversality in the following sense: if all functions g_1, \dots, g_k are generic, it does not hold if the caps are assumed to be weakly transversal. This is explained in [Appendix A](#).

Remark 1.9. For $2 \leq k \leq d + 1$ we do not use the tensor structure explicitly. It is used in an implicit way when comparing the sizes of natural scalar and mixed norm quantities that appear in the proofs.

Remark 1.10. For $2 \leq k \leq d$, if all functions involved are full tensors, one has more estimates than those predicted by [Conjecture 1.2](#) assuming extra *degrees of transversality*, as proven in [Section 11](#).

It is natural to try to generalize the statement of [Conjecture 1.2](#) for L^p inputs rather than just L^2 . A motivation for that is to deeply understand the role played by transversality; as we will see, the farther our inputs are from L^2 , the less impact the configuration of the caps on the paraboloid has in the best possible estimate (with a single exception to be detailed soon). The general statement of the k -linear extension conjecture for the paraboloid is (as in [\[Bennett 2014\]](#)):

Conjecture 1.11. *Let $k \geq 2$ and suppose that U_1, \dots, U_k parametrize transversal caps of the paraboloid $x \mapsto |x|^2$ in \mathbb{R}^{d+1} . If*

$$\frac{1}{q} < \frac{d}{2(d+1)}, \quad \frac{1}{q} \leq \frac{d+k-1}{d+k+1} \frac{1}{p'} \quad \text{and} \quad \frac{1}{q} \leq \frac{d-k+1}{d+k+1} \frac{1}{p'} + \frac{k-1}{k+d+1},$$

then

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_{L^{q/k}(\mathbb{R}^{d+1})} \lesssim_{p,q} \prod_{j=1}^k \|g_j\|_{L^p(U_j)}.$$

For $2 \leq k < d+1$, to recover the interior of the conjectured range, it is enough³ to prove [Conjecture 1.2](#) and

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^{2(d+1)/d}(U_j)} \tag{4}$$

for all $\varepsilon > 0$.

Remark 1.12. Observe that (4) covers the case $(p, q) = \left(\frac{2(d+1)}{d}, \frac{2(d+1)}{d} + \varepsilon\right)$ of [Conjecture 1.11](#). Notice also that this case would follow from the case $(p, q) = \left(\frac{2(d+1)}{d}, \frac{2(d+1)}{d} + \varepsilon\right)$ of the linear extension of [Conjecture 1.1](#) and Hölder’s inequality. This means that the closer we get to the endpoint extension exponent, the fewer improvements transversality yields in the multilinear theory. The exception to this is the $k = d + 1$ case, for which L^2 functions give the best possible output for the corresponding multilinear operator (rather than $L^{2(d+1)/d}$). Indeed, when one function is a tensor, the best result in this case is obtained in [Section 10](#).

By adapting the argument that shows the case $2 \leq k \leq d + 1$ of [Theorem 1.5](#), we are able to prove the following weaker version of (4):

³The interior of the full range of estimates follows by interpolation between these two cases and the trivial bound $(p, q) = (1, \infty)$.

Theorem 1.13. *Let $2 \leq k < d + 1$. If g_1 is a tensor in addition to the hypotheses of [Conjecture 1.11](#), the following estimate holds:*

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^{p(k,d)}(U_j)} \tag{5}$$

for all $\varepsilon > 0$, where

$$p(k, d) = \begin{cases} \frac{4(d+1)}{d+k+1} & \text{if } 2 \leq k < \frac{d}{2}, \\ \frac{4(d+1)}{2d-k+1} & \text{if } \frac{d}{2} \leq k < d + 1. \end{cases}$$

Remark 1.14. Notice that $\frac{2(d+1)}{d} < p(k, d)$, so [Theorem 1.13](#) is not optimal on the space of the input functions. On the other hand, the output $L^{2(d+1)/(kd)+\varepsilon}$ (for all $\varepsilon > 0$) is the best to which one can hope to map the multilinear operator on the left-hand side. The case $k = d + 1$ of the theorem above coincides with the case $k = d + 1$ of the L^2 -based theory, which is covered in [Section 10](#).

Remark 1.15. Bounds such as the one from [Theorem 1.13](#), i.e., in which one needs p big enough (and not sharp) to map L^p inputs to a fixed L^q , are common in linear extension theory. For example, Wang [\[2018\]](#) showed that \mathcal{E}_2 maps $L^\infty([-1, 1]^2)$ to $L^q(\mathbb{R}^3)$ for $q > 3 + \frac{3}{13}$. As mentioned in [\[Wang 2018\]](#), this implies the (seemingly stronger) bound

$$\|\mathcal{E}_2 g\|_{L^q(\mathbb{R}^3)} \lesssim_q \|g\|_{L^q([-1, 1]^2)}$$

for $q > 3 + \frac{3}{13}$ via the factorization theory of Nikishin and Pisier (see [\[Bourgain 1991\]](#)).

Remark 1.16. The multilinear extension theory for inputs near $L^{2(d+1)/d}$ remains largely unknown in general (except for the almost optimal result in the $k = d + 1$ case in [\[Bennett et al. 2006\]](#)). In fact, it is not fully settled even in the $k = 2, d > 1$ case (whose L^2 -based analogue is known). We refer the reader to [\[Oh 2023\]](#) for partial results in this direction.

Remark 1.17. As the reader may expect, any function can be taken to be the tensor in the statement of [Theorem 1.13](#).

The linear and multilinear theories studied in this paper meet very naturally once more in the context of the techniques we use: the simplest multilinear variant of a linear operator T is given by the product of a certain number of identical copies of it:

$$T_{(k)}(g_1, \dots, g_k) := \prod_{j=1}^k T g_j.$$

Proving that T maps $L^p(U)$ to $L^q(V)$ is equivalent to proving that $T_{(k)}$ maps $L^p(U)$ to $L^{q/k}(V)$, as one can easily check with Hölder’s inequality. Multilinearizing \mathcal{E}_d without any regard to transversality yields the operator

$$\mathcal{E}_{d,(k)}(g_1, \dots, g_k) := \prod_{j=1}^k \mathcal{E}_d g_j. \tag{6}$$

Combining the previous observation with the factorization theory of Nikishin and Pisier, [Conjecture 1.1](#) follows from the bound

$$\left\| \prod_{j=1}^k \mathcal{E}_d g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^\infty([0,1]^d)}. \tag{7}$$

The proof of [Theorem 1.13](#) can be adapted to show the following:

Theorem 1.18. *Let $2 \leq k \leq d + 1$. If g_1 is a tensor, the inequality*

$$\left\| \prod_{j=1}^k \mathcal{E}_d g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^4([0,1]^d)} \tag{8}$$

holds for all $\varepsilon > 0$.

Remark 1.19. Since the inputs g_j are compactly supported, [Theorem 1.18](#) implies (7).

Remark 1.20. Given that the proof of [Theorem 1.18](#) has the L^4 - $L^{4+\varepsilon}$ bound for \mathcal{E}_1 as its main building block, it is not surprising that we have a product of L^4 norms in the right-hand side of the statement above.

We finish this introduction by highlighting the close connection between our results and the theory of linear and nonlinear Brascamp–Lieb inequalities. The concept of *weak transversality* that we introduce can be characterized in terms of certain Brascamp–Lieb data, and by exploiting the geometric features arising from this fact we are able to verify a special case of a conjecture by Bennett, Bez, Flock and Lee.

The paper is organized as follows: in [Section 2](#) we present the linear and multilinear models that we will work with in the proof of [Theorem 1.5](#). We also highlight the main differences between the linearized models that are used in most recent approaches and ours. In [Section 3](#) we define the concepts of transversality and weak transversality, and state in what sense the former implies the latter. [Section 4](#) presents what we refer to as the *building blocks* of our approach. Sections [5](#), [6](#) and [7](#) establish these building blocks: in [Section 5](#) we revisit the case $k = 1$ and $p = 2$ for our model, in [Section 6](#) we revisit Zygmund’s argument and recover the case $k = 1$ for $d = 1$, and in [Section 7](#) we deal with the case $k = 2$ and $d = 1$. In [Section 8](#) we settle the case $k = 1$ of [Theorem 1.5](#), and in [Section 9](#) we show the cases $2 \leq k \leq d + 1$. [Section 10](#) covers the endpoint estimate of the case $k = d + 1$. In [Section 11](#) we discuss how one can improve the bounds of [Conjecture 1.2](#) under extra transversality and tensor hypotheses. [Theorem 1.13](#) (our partial result beyond the L^2 -based k -linear theory) is presented in [Section 12](#) along with its “nontransversal” counterpart [Theorem 1.18](#). In [Section 13](#) we establish a connection between the classical theory of Brascamp–Lieb inequalities and our results, and give an application of this link to a conjecture made in [[Bennett et al. 2018](#)]. In [Section 14](#) we make a few additional remarks. [Appendix A](#) contains examples that show that the range of p in [Conjecture 1.2](#) is sharp, and also that one cannot obtain this range in general under a condition that is strictly weaker than transversality. [Appendix B](#) contains technical results used throughout the paper.

2. Discrete models

A common first step of the earlier works is to *linearize* the contribution of the quadratic phase $x \mapsto |x|^2$. One starts by studying $\mathcal{E}_d g$ on a ball of radius R (hence $|(x, t)| \leq R$) and splits the domain of g into

balls θ_k of radius $R^{-1/2}$. Let us consider $d = 1$ here for simplicity. If

$$g_{\theta_k} := g \cdot \varphi_{\theta_k},$$

where φ_{θ_k} is a bump adapted to $[kR^{-1/2}, (k + 1)R^{-1/2}]$, the quadratic exponential

$$e_{x,t}(\xi) = e^{2\pi i x \xi} e^{2\pi i t \xi^2} \tag{9}$$

behaves in a similar way to a linear exponential $e^{i \# \xi}$ when restricted to this interval. Indeed, the phase-space portrait of $e_{x,t}$ is the (oblique if $t \neq 0$) line

$$u \mapsto x + 2tu,$$

as is explained in more detail in Chapter 1 of [Muscalu and Schlag 2013b]. When we evaluate this line at the endpoints of the support of g_{θ_k} (taking into account that $|t| \leq R$), we see that the phase-space portrait of

$$\varphi_{\theta_k} \cdot e_{x,t}$$

is a parallelogram that essentially coincides with the rectangle

$$I \times J = [kR^{-\frac{1}{2}}, (k + 1)R^{-\frac{1}{2}}] \times [x + 2tkR^{-\frac{1}{2}}, x + 2tkR^{-\frac{1}{2}} + R^{\frac{1}{2}}]. \tag{10}$$

Observe that $I \times J$ has area 1. On the other hand, the phase-space portrait of φ_{θ_k} is a Heisenberg box of sizes $R^{-1/2}$ and $R^{1/2}$, and the linear modulation

$$e^{2\pi i \xi(x+2tkR^{-1/2})} \tag{11}$$

shifts it in frequency to J . The conclusion is that the phase-space portrait of

$$\varphi_{\theta_k} \cdot e^{2\pi i \xi(x+2tkR^{-1/2})}$$

is the Heisenberg box (10); hence the effect of the quadratic modulation $e_{x,t}$ in this setting is essentially the same as the linear one in (11).

Using bumps such as φ_θ to decompose the domain of g and expanding each g_θ into Fourier series allows us to write

$$g(x) = \sum_{\theta \in R^{-1/2}\mathbb{Z}^d \cap [0,1]^d} \overbrace{g(x)\varphi_\theta(x)}^{g_\theta(x)} \tilde{\varphi}_\theta(x) = \sum_{\theta \in R^{-1/2}\mathbb{Z}^d \cap [0,1]^d} \sum_{v \in R^{1/2}\mathbb{Z}^d} \overbrace{c_{v,\theta} e^{2\pi i x \cdot v}}^{g_{\theta,v}(x)} \tilde{\varphi}_\theta(x),$$

where $\tilde{\varphi}_\theta \equiv 1$ on the support of φ_θ and decays very fast away from it. Applying \mathcal{E}_d and using the previous intuition gives rise to the *wave packet decomposition*

$$\mathcal{E}_d g = \sum_{(\theta,v) \in R^{-1/2}\mathbb{Z}^d \cap [0,1]^d \times R^{1/2}\mathbb{Z}^d} \mathcal{E}_d(g_{\theta,v}),$$

where $\mathcal{E}_d(g_{\theta,v})$ is essentially supported on a tube in \mathbb{R}^{d+1} of size $R^{1/2} \times \dots \times R^{1/2} \times R$ whose direction is determined by θ and that is translated by a parameter depending on v . With this linearized model at hand, one can study the interference between these tubes pointing in different directions (both in the

linear and multilinear settings) and take advantage of orthogonality both in space and in frequency. This leads to local estimates of type

$$\|\mathcal{E}_d g\|_{L^q(B(0,R))} \lesssim_\varepsilon R^\varepsilon \|f\|_p \quad \text{for all } \varepsilon > 0$$

and multilinear analogues of it that are later used to obtain global estimates via ε -removal arguments (as in [Tao 1999]). The reader is referred to [Guth 2016] for the details of the decomposition above. This approach has given the current best L^p bounds for \mathcal{E}_d .

In our case, we do not linearize the contribution of the quadratic phase. Instead, we consider a discrete model that keeps the quadratic nature of \mathcal{E}_d intact.

2A. The linear model ($k = 1$). We consider $d = 1$ for simplicity, but the discretization process is analogous for all $d > 1$. Recall that the extension operator for the parabola defined for functions supported on $[0, 1]$ is given by

$$\mathcal{E}_1 g(x, t) = \int_0^1 g(\xi) e^{-2\pi i x \xi} e^{-2\pi i t \xi^2} d\xi. \tag{12}$$

We can insert a bump φ in the integrand that is equal to 1 on $[0, 1]$ and supported in a small neighborhood of this interval. Tiling \mathbb{R}^2 with unit squares with vertices in \mathbb{Z}^2 and rewriting \mathcal{E}_1 ,

$$\mathcal{E}_1 g(x, t) = \sum_{n,m \in \mathbb{Z}} \left[\int g(u) \varphi(u) e^{-2\pi i x u} e^{-2\pi i t u^2} du \right] \chi_n(x) \chi_m(t),$$

where $\chi_n := \chi_{[n, n+1)}$. For a fixed (x, t) , one can write

$$\begin{aligned} e^{-2\pi i x \xi} e^{-2\pi i t \xi^2} \varphi(\xi) &= e^{-2\pi i n \xi} e^{-2\pi i m \xi^2} \cdot e^{-2\pi i (x-n)\xi} e^{-2\pi i (t-m)\xi^2} \varphi(\xi) \\ &= e^{-2\pi i n \xi} e^{-2\pi i m \xi^2} \cdot \sum_{u \in \mathbb{Z}} \langle e^{-2\pi i (x-n)(\cdot)} e^{-2\pi i (t-m)(\cdot)^2}, \varphi_{[0,1]}^u \rangle \cdot \varphi_{[0,1]}^u(\xi) \\ &= e^{-2\pi i n \xi} e^{-2\pi i m \xi^2} \cdot \sum_{u \in \mathbb{Z}} C_u^{n,m,x,t} \cdot \varphi_{[0,1]}^u(\xi), \end{aligned}$$

where we expanded $e^{-2\pi i (x-n)\xi} e^{-2\pi i (t-m)\xi^2}$ as a Fourier series at scale 1,

$$\begin{aligned} C_u^{n,m,x,t} &:= \langle e^{-2\pi i (x-n)(\cdot)} e^{-2\pi i (t-m)(\cdot)^2}, \varphi_{[0,1]}^u \rangle, \\ \varphi_{[0,1]}^u(\xi) &:= \varphi_{[0,1]}(\xi) \cdot e^{-2\pi i u \cdot \xi} \end{aligned}$$

and $\varphi_{[0,1]}$ is a bump adapted to $[0, 1]$ (and compactly supported) just like⁴ φ . Plugging this in (12),

$$\begin{aligned} \mathcal{E}_1 g(x, t) &= \sum_{n,m \in \mathbb{Z}} \left[\int g(\xi) \varphi(\xi) e^{-2\pi i x \xi} e^{-2\pi i t \xi^2} d\xi \right] \chi_n(x) \chi_m(t) \\ &= \sum_{n,m \in \mathbb{Z}} \left[\int g(\xi) \left(e^{-2\pi i n \xi} e^{-2\pi i m \xi^2} \cdot \sum_{u \in \mathbb{Z}} C_u^{n,m,x,t} \cdot \varphi^u(\xi) \right) d\xi \right] \chi_n(x) \chi_m(t) \\ &= \sum_{u \in \mathbb{Z}} \sum_{n,m \in \mathbb{Z}} C_u^{n,m,x,t} \cdot \left[\int g(\xi) e^{-2\pi i n \xi} e^{-2\pi i m \xi^2} \cdot \varphi^u(\xi) d\xi \right] \chi_n(x) \chi_m(t). \end{aligned}$$

⁴We will not distinguish between $\varphi_{[0,1]}$ and φ from now on.




Figure 2. The phase-space portrait of $\varphi_{n,m}$.

For the expression defining \mathcal{E}_1 to be nonzero, (n, m) must satisfy $|x - n| \leq 1$ and $|t - m| \leq 1$; hence the Fourier coefficients $C_u^{n,m,x,t}$ decay like $O(|u|^{-100})$. In addition, the extra factor φ^u in the integral simply shifts the integrand in frequency, and this does not affect in any way the arguments that follow. In order to obtain the final form of our linear model, let us introduce the following notation: if φ is a compactly supported bump (say, in a very small open neighborhood of $[0, 1]^d$) with $\varphi \equiv 1$ on $[0, 1]^d$, we set

$$\varphi_{\vec{n},m}(x) := \varphi(x)e^{2\pi i x \cdot \vec{n}} e^{2\pi i |x|^2 m}. \tag{13}$$

Due to the fast decay of $C_u^{n,x}$ and $C_v^{m,t}$, it is then enough to bound the $u = v = 0$ piece of the sum above, which leads to the discretized model:⁵

$$E_1(g) = \sum_{(n,m) \in \mathbb{Z}^2} \langle g, \varphi_{n,m} \rangle (\chi_n \otimes \chi_m).$$

With the appropriate adaptations, one proceeds in the exact same way in dimension d to reduce matters to the study of the following model operator:

Definition 2.1. Let E_d be defined on $C([0, 1]^d)$ given by

$$E_d(g) = \sum_{\vec{n} \in \mathbb{Z}^d, m \in \mathbb{Z}} \langle g, \varphi_{\vec{n},m} \rangle (\chi_{\vec{n}} \otimes \chi_m),$$

where $\chi_{\vec{n}}$ and χ_m are the characteristic functions of the boxes $[n_1, n_1 + 1) \times \dots \times [n_d, n_d + 1)$ and $[m, m + 1)$, respectively.⁶

The wave packets (13) have a natural phase-space portrait that consist of parallelograms in the phase plane. See Figure 2.

⁵There is a slight abuse of notation here: observe that $\tilde{\chi}_n(x)\tilde{\chi}_m(t) := C_0^{n,m,x,t} \cdot \chi_n(x)\chi_m(t)$ is a smooth function supported in $[n, n + 1) \times [m, m + 1)$, which is all that is needed in the proof. We will continue to call it $\chi_n(x)\chi_m(t)$ to lighten the notation.

⁶Morally speaking, the discrete model and the original operator are “comparable”, but we were not able to prove that rigorously. For that reason we included the proof of known extension estimates for E_d .

By keeping the quadratic nature of E_d intact we take advantage of orthogonality in different ways. For example, for a fixed m the wave packets $\varphi_{n,m}$ are almost orthogonal, as suggested by the fact that the corresponding parallelograms are (almost) disjoint.

2B. The multilinear model ($2 \leq k \leq d + 1$). We recall the definition of the k -linear extension operator:

Definition 2.2. For $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ a transversal set of cubes, the k -linear extension operator is given by

$$\mathcal{ME}_{k,d}(g_1, \dots, g_k) := \prod_{j=1}^k \mathcal{E}_{Q_j} g_j, \tag{14}$$

where

$$\mathcal{E}_{Q_j} g_j(x, t) = \int_{Q_j} g_j(\xi) e^{-2\pi i x \cdot \xi} e^{-2\pi i t |\xi|^2} d\xi, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R}.$$

By an analogous argument to the one we showed in [Section 2A](#), it is enough to prove the corresponding bounds for the following model operator:

Definition 2.3. Let $\text{ME}_{k,d}$ be defined on $C(Q_1) \times \dots \times C(Q_k)$ by

$$\text{ME}_{k,d}(g_1, \dots, g_k) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \langle g_j, \varphi_{\vec{n}, m}^j \rangle (\chi_{\vec{n}} \otimes \chi_m).$$

where

$$\varphi_{\vec{n}, m}^j = \bigotimes_{l=1}^d \varphi_{n_l, m}^{l,j}, \quad \varphi_{n_l, m}^{l,j}(x_l) = \varphi^{l,j}(x_l) e^{2\pi i n_l x_l} e^{2\pi i m x_l^2}$$

and $\varphi^{l,j}(x)$ is $\equiv 1$ on the l -coordinate projection of the domain of g_j defined above and decays fast away from it.

Remark 2.4. It is clear that the discretization process does not depend on whether the collection \mathcal{Q} is made of transversal cubes or not. In particular, it will be of interest in [Section 12B](#) to study the operator given by the right-hand side of (14), but *without* the assumption that the cubes Q_j are transversal. The model for such operator is also given by $\text{ME}_{k,d}$, but without that hypothesis.

3. Transversality versus weak transversality

We recall the following definition from [\[Bennett 2014\]](#):

Definition 3.1. Let $2 \leq k \leq d + 1$ and $c > 0$. A k -tuple S_1, \dots, S_k of smooth codimension-1 submanifolds of \mathbb{R}^{d+1} is c -transversal if

$$|v_1 \wedge \dots \wedge v_k| \geq c$$

for all choices v_1, \dots, v_k of unit normal vectors to S_1, \dots, S_k , respectively. We say that S_1, \dots, S_k are transversal if they are c -transversal for some $c > 0$.

In other words, if the k -dimensional volume of the parallelepiped generated by v_1, \dots, v_k is bounded below by some absolute constant for any choice of normal vectors v_j , the submanifolds are transversal.

From now on, we will say that a collection of k cubes in \mathbb{R}^d is *transversal* if the associated caps defined by them on the paraboloid are transversal in the sense of [Definition 3.1](#).

One can assume without loss of generality that the U_j in the statements of [Conjecture 1.2](#) are cubes that parametrize transversal caps on \mathbb{P}^d via the map $x \mapsto |x|^2$. Even though these conjectures are known to fail in general if one does not assume transversality between the caps (see [Section AB](#)), the theorem that we will prove holds under a weaker condition, since one of the functions is a tensor.

Definition 3.2. Let $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ be a collection of k (open or closed) cubes⁷ in \mathbb{R}^d . The collection \mathcal{Q} is said to be *weakly transversal with pivot Q_j* if there is a set of $k-1$ distinct directions $\mathcal{E}_j = \{e_{i_1}, \dots, e_{i_{k-1}}\}$ (depending on j) of the canonical basis such that

$$\left\{ \begin{array}{l} \overline{\pi_{i_1}(Q_j)} \cap \overline{\pi_{i_1}(Q_1)} = \emptyset, \\ \vdots \\ \overline{\pi_{i_{j-1}}(Q_j)} \cap \overline{\pi_{i_{j-1}}(Q_{j-1})} = \emptyset, \\ \overline{\pi_{i_j}(Q_j)} \cap \overline{\pi_{i_j}(Q_{j+1})} = \emptyset, \\ \vdots \\ \overline{\pi_{i_{k-1}}(Q_j)} \cap \overline{\pi_{i_{k-1}}(Q_k)} = \emptyset, \end{array} \right. \tag{15}$$

where π_l is the projection onto e_l . We say that \mathcal{Q} is *weakly transversal* if it is weakly transversal with pivot Q_j for all $1 \leq j \leq k$.⁸

Remark 3.3. For each $1 \leq j \leq k$, from now on we will refer to a set⁹ \mathcal{E}_j above as *a set of directions associated to Q_j* . Notice that there could be many of such sets for a single j . Also, if $j_1 \neq j_2$, it could be the case that no set of directions associated to Q_{j_1} is associated to Q_{j_2} .

Let us give a few examples to distinguish between [Definitions 3.1](#) and [3.2](#). Consider the case $d = 2$, $k = 3$, $Q_1 = [0, 1]^2$, $Q_2 = [2, 3]^2$, and $Q_3 = [4, 5]^2$. The line $y = x$ intersects Q_1 , Q_2 and Q_3 ; then it follows from [Definition 3.1](#) that they are not transversal. However, observe that

$$\begin{cases} \pi_1(Q_1) \cap \pi_1(Q_2) = \emptyset, \\ \pi_2(Q_1) \cap \pi_2(Q_3) = \emptyset, \end{cases}$$

so $\{e_1, e_2\}$ is a set associated to Q_1 (and similarly one can verify that it is also associated to Q_2 and Q_3). This shows that the collection defined by Q_1 , Q_2 and Q_3 is weakly transversal.

Consider now the cubes $K_1 = [0, 1]^2$, $K_2 = [4, 5] \times [0, 1]$ and $K_3 = [2, 3]^2$. Not only are they transversal in the sense of [Definition 3.1](#), but also weakly transversal.

This is not by chance: a given transversal collection of k cubes can be “decomposed” into finitely many collections of k cubes that are *also* weakly transversal.

⁷The word *cube* will be used throughout the paper to refer to any rectangular box in \mathbb{R}^d , regardless of the sizes of its edges, and they always refer to the supports of the input functions of our linear and multilinear operators. In this paper, it will not be relevant whether the sides of a box have the same length or not; therefore this slight abuse of terminology is harmless.

⁸The estimates that we will prove depend on the separation of the projections in [Definition 3.2](#), just as they depend on the behavior of c from [Definition 3.1](#) in the general case for transversal caps.

⁹The typeface \mathcal{E}_j is being used to distinguish this concept from the previously defined operators \mathcal{E}_d and E_d .




Figure 3. Transversality versus weak transversality.

Claim 3.4. *Given a collection $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ of transversal cubes, each $Q_l \in \mathcal{Q}$ can be partitioned into $O(1)$ many subcubes*

$$Q_l = \bigcup_i Q_{l,i}$$

so that all collections $\tilde{\mathcal{Q}}$ made of picking one subcube $Q_{l,i}$ per Q_l

$$\tilde{\mathcal{Q}} = \{\tilde{Q}_1, \dots, \tilde{Q}_k\}, \quad \tilde{Q}_l \in \{Q_{l,i}\}_i,$$

are weakly transversal.

Proof. See Claim B.4 in Appendix B. □

As a consequence of Claim 3.4, to prove the case $2 \leq k \leq d + 1$ of Theorem 1.5 it suffices to show it for weakly transversal collections. To simplify the exposition, we will present our results for the cubes

$$Q_1 = [0, 1]^d,$$

$$Q_j = [2, 3]^{j-2} \times [4, 5] \times [0, 1]^{d-j+1}, \quad 2 \leq j \leq k.$$

The associated directions to Q_1 are $\{e_1, \dots, e_{k-1}\}$, and we will use it as the pivot. Any other weakly transversal collection of cubes can be dealt with in the same way.

4. Our approach and its building blocks

Notice that the operators \mathcal{E}_d and $\mathcal{ME}_{k,d}$ are pointwise bounded by E_d and $ME_{k,d}$, respectively; therefore we cannot directly conclude any result about the models from the fact that they hold for the original operators. Some of these results will be reproven for the models in this paper, and they will act as *building blocks* in the proof of Theorem 1.5, which is presented in Sections 8 and 9. More precisely, Theorem 1.5 relies on the following:

(1) *Mixed norm Strichartz/Tomas–Stein* ($k = 1, p = 2$). In Section 5 we show the following:

Proposition 4.1. *For all $p > \frac{2(d+2)}{d}$,*

$$\|E_d g\|_p \lesssim_p \|g\|_2.$$

As a consequence, we have:

Corollary 4.2. For all $\varepsilon > 0$,

$$\|E_d(g)\|_{L^{2(d-l+2)/(d-l)+\varepsilon}_{x_{l+1}, \dots, x_d, t} L^2_{x_1, \dots, x_l}} \lesssim_\varepsilon \|g\|_2. \tag{16}$$

Proof. Apply Minkowski’s inequality and Proposition 4.1 in dimension $d - l$. Notice that, after taking L^2 norm in the first l variables, we can use Bessel to bound the left-hand side of (16) by

$$\left[\sum_{n_{l+1}, \dots, n_d, m} \left(\sum_{n_1, \dots, n_l} |\langle g, \varphi_{n_{l+1}, \dots, n_d, m} \rangle, \varphi_{n_1, \dots, n_l, m} \rangle|^2 \right)^{\frac{p_0}{2}} \right]^{\frac{1}{p_0}} \lesssim \left[\sum_{n_{l+1}, \dots, n_d, m} \| \langle g, \varphi_{n_{l+1}, \dots, n_d, m} \rangle \|_2^{p_0} \right]^{\frac{1}{p_0}}, \quad \text{where } p_0 = \frac{2(d-l+2)}{d-l} + \varepsilon.$$

This is how we will use Corollary 4.2 in (56). □

We will use Corollary 4.2 in Conjecture 1.2 to prove Theorem 1.5 for $2 \leq k \leq d + 1$. It will not be needed when $k = d + 1$.

(2) *Extension conjecture for the parabola* ($k = 1, d = 1, p = 4$). In Section 6 we prove the following:

Proposition 4.3. For all $\varepsilon > 0$,

$$\|E_1 g\|_{4+\varepsilon} \lesssim_\varepsilon \|g\|_4. \tag{17}$$

One can show by interpolation that Proposition 4.3 implies Conjecture 1.1 for $d = 1$. We will use it in Section 8 to settle the case $k = 1$ of Theorem 1.5.

(3) *Bilinear extension conjecture for the parabola* ($k = 2, d = 1$). In Section 7 we show that the model $ME_{2,1}$ in Definition 2.3 maps $L^2([0, 1]) \times L^2([4, 5])$ to $L^2(\mathbb{R}^2)$.

Proposition 4.4. The following estimate holds:

$$\|ME_{2,1}(f, g)\|_2 \lesssim \|f\|_2 \cdot \|g\|_2. \tag{18}$$

Transversality will be captured in Section 9 through (18).

By combining scalar and mixed norm stopping times¹⁰ performed simultaneously, we are able to put together the key estimates (16), (17) and (18). In the $2 \leq k \leq d + 1$ case, the tensor structure is used in an implicit way to allow us to better relate these scalar and mixed norm stopping times.

Remark 4.5. The tensor structure $g = g_1 \otimes \dots \otimes g_d$ in the $k = 1$ case allows us to write

$$\langle g, \varphi_{\vec{n}, m} \rangle = \prod_{j=1}^d \langle g_j, \varphi_{n_j, m} \rangle. \tag{19}$$

¹⁰This is not meant in a literal probabilistic sense; strictly speaking, the argument combines the level sets of various scalar and mixed norm quantities that appear naturally in our analysis.

We then obtain the following multilinear form by dualization:

$$\Lambda_d(g_1, \dots, g_d, h) = \langle E_d(g), h \rangle = \sum_{\vec{n} \in \mathbb{Z}^d, m \in \mathbb{Z}} \prod_{j=1}^d \langle g_j, \varphi_{n_j, m} \rangle \cdot \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle, \tag{20}$$

The goal in the $k = 1$ case is to show that

$$|\Lambda_d(g_1, \dots, g_d, h)| \lesssim \|h\|_q \cdot \prod_{j=1}^d \|g_j\|_{p_j}$$

for appropriate exponents p_j and q . Interpolation theory shows that it suffices to obtain

$$|\Lambda_d(g_1, \dots, g_d, h)| \lesssim_\varepsilon |F|^{\gamma_{d+1}} \cdot \prod_{j=1}^d |E_j|^{\gamma_j} \tag{21}$$

for all $\varepsilon > 0$, $|g_j| \leq \chi_{E_j}$, $|h| \leq \chi_F$,¹¹ $E_j \subset [0, 1]$ and $F \subset \mathbb{R}^3$ measurable sets such that γ_j ($1 \leq j \leq d$) and γ_{d+1} are in a small neighborhood of $\frac{d}{2(d+1)}$ and $\frac{d+2}{2(d+1)} + \varepsilon$, respectively.¹² We refer the reader to [Thiele 2006, Chapter 3] for a detailed account of multilinear interpolation theory. To keep the notation simple, all restricted weak-type estimates we will prove in this paper will be for the centers of such neighborhoods. For example, we will show that

$$|\Lambda_d(g_1, \dots, g_d, h)| \lesssim_\varepsilon |F|^{\frac{d+2}{2(d+1)} + \varepsilon} \cdot \prod_{j=1}^d |E_j|^{\frac{d}{2(d+1)}} \tag{22}$$

for all $\varepsilon > 0$, but it will be clear from the arguments that as long as we give this $\varepsilon > 0$ away, a slightly different choice of interpolation parameters yields (21). The restricted weak-type estimates that we will prove in the $2 \leq k \leq d + 1$ case will also be for the centers of the corresponding neighborhoods.

5. Proof of Proposition 4.1: Strichartz/Tomas–Stein for E_d ($k = 1, p = 2$)

Our proof is inspired by the classical TT^* argument. It is possible to prove the endpoint estimate directly for the model E_d by repeating the steps of this argument (see for example [Muscalu and Schlag 2013a, Section 11.2.2]), but we chose the following approach because of its similarity with the one we will use to prove Theorem 1.5. By interpolation with the trivial bound for $q = \infty$, it is enough to prove the bound

$$\|E_d g\|_{\frac{2(d+2)}{d} + \varepsilon} \lesssim_\varepsilon \|g\|_2$$

for all $\varepsilon > 0$.

We start by dualizing E_d to obtain a bilinear form Λ_d :

$$\Lambda_d(g, h) = \langle E_d(g), h \rangle = \sum_{\vec{n} \in \mathbb{Z}^d, m \in \mathbb{Z}} \langle g, \varphi_{\vec{n}, m} \rangle \cdot \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle.$$

¹¹There is an overlap of classical notation here that we hope will not compromise the comprehension of the paper: we chose the typeface E_d to represent the discrete model of the official extension operator \mathcal{E} . On the other hand, the classical theory of restricted weak-type multilinear interpolation usually labels the measurable sets involved in the problems by E_j or F_j . The context will make it clear which object we are referring to.

¹²Rigorously, this only verifies the case $k = 1$ near the endpoint $(\frac{2(d+1)}{d}, \frac{2(d+1)}{d})$, but this is known to imply the desired estimates in the full range. For details, see [Mattila 2015, Theorem 19.8].

Let $E_1 \subset \mathbb{R}^d$ and $E_2 \subset \mathbb{R}^{d+1}$ be measurable sets of finite measure with $|g| \leq \chi_{E_1}$ and $|h| \leq \chi_{E_2}$. Split \mathbb{Z}^{d+1} in two ways:

$$\begin{aligned} \mathbb{Z}^{d+1} &= \bigcup_{l_1 \in \mathbb{Z}} \mathbb{A}^{l_1}, \quad \text{where } (\vec{n}, m) \in \mathbb{A}^{l_1} \iff |\langle g, \varphi_{\vec{n}, m} \rangle| \approx 2^{-l_1}, \\ \mathbb{Z}^{d+1} &= \bigcup_{l_2 \in \mathbb{Z}} \mathbb{B}^{l_2}, \quad \text{where } (\vec{n}, m) \in \mathbb{B}^{l_2} \iff |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-l_2}. \end{aligned}$$

Define $\mathbb{X}^{l_1, l_2} := \mathbb{A}^{l_1} \cap \mathbb{B}^{l_2}$ and observe that

$$|\Lambda_d(g, h)| \lesssim \sum_{l_1, l_2 \in \mathbb{Z}} 2^{-l_1} 2^{-l_2} \#\mathbb{X}^{l_1, l_2}.$$

Notice that, for all $(\vec{n}, m) \in \mathbb{X}^{l_1, l_2}$,

$$\begin{aligned} 2^{-l_1} &\lesssim \int_{\mathbb{R}^d} |g(x)| |\varphi_{\vec{n}, m}(x)| \, dx \leq \min\{|E_1|, 1\}, \\ 2^{-l_2} &\lesssim \int_{\mathbb{R}^d} |h(x)| |\chi_{\vec{n}} \otimes \chi_m(x)| \, dx \leq \min\{|E_2|, 1\}. \end{aligned}$$

In particular, $l_1, l_2 \geq 0$ in the sum above. Now we bound $\#\mathbb{X}^{l_1, l_2}$ in two different ways and interpolate between them:

(a) *L¹-type bound:* Exploit h :

$$\#\mathbb{X}^{l_1, l_2} \leq \#\mathbb{B}^{l_2} \lesssim 2^{l_2} \sum_{(\vec{n}, m) \in \mathbb{B}^{l_2}} |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \lesssim 2^{l_2} \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \int_{Q_{\vec{n}, m}} |h| = 2^{l_2} \|h\|_1 \leq 2^{l_2} |E_2|, \quad (23)$$

where $Q_{\vec{n}, m} := \prod_{i=1}^d [n_i, n_i + 1] \times [m, m + 1]$, $\vec{n} = (n_1, \dots, n_d)$.

(b) *L²-type bound:* Exploit g :

$$\begin{aligned} \#\mathbb{X}^{l_1, l_2} &\lesssim 2^{2l_1} \sum_{(\vec{n}, m) \in \mathbb{X}^{l_1, l_2}} |\langle g, \varphi_{\vec{n}, m} \rangle|^2 \\ &= 2^{2l_1} \left| \left\langle \sum_{(\vec{n}, m) \in \mathbb{X}^{l_1, l_2}} \langle g, \varphi_{\vec{n}, m} \rangle \varphi_{\vec{n}, m}, g \right\rangle \right| \\ &\leq 2^{2l_1} |E_1|^{\frac{1}{2}} \underbrace{\left\| \sum_{(\vec{n}, m) \in \mathbb{X}^{l_1, l_2}} \langle g, \varphi_{\vec{n}, m} \rangle \varphi_{\vec{n}, m} \right\|_2}_{(*)}. \end{aligned} \quad (24)$$

For each set \mathbb{X}^{l_1, l_2} define $\pi_m := \{\vec{n} \in \mathbb{Z}^d; (\vec{n}, m) \in \mathbb{X}^{l_1, l_2}\}$. Observe that

$$(*)^2 = \sum_{m: \pi_m \neq \emptyset} \sum_{\tilde{m}: \pi_{\tilde{m}} \neq \emptyset} \underbrace{\sum_{\vec{n} \in \pi_m} \sum_{\vec{k} \in \pi_{\tilde{m}}} \langle g, \varphi_{\vec{n}, m} \rangle \overline{\langle g, \varphi_{\vec{k}, \tilde{m}} \rangle} \langle \varphi_{\vec{n}, m}, \varphi_{\vec{k}, \tilde{m}} \rangle}_{U(\langle g, \varphi_{\vec{n}, m} \rangle)_{\vec{n} \in \pi_m}, (\langle g, \varphi_{\vec{k}, \tilde{m}} \rangle)_{\vec{k} \in \pi_{\tilde{m}}}}$$

We will estimate U in two ways. Let $a_{\vec{n},m} := \langle g, \varphi_{\vec{n},m} \rangle$. First, by the triangle inequality and the stationary phase [Theorem B.3](#)

$$\begin{aligned} |U((a_{\vec{n},m})_{\vec{n} \in \pi_m}, (a_{\vec{k},\tilde{m}})_{\vec{k} \in \pi_{\tilde{m}}})| &\leq \sum_{\vec{n} \in \pi_m} \sum_{\vec{k} \in \pi_{\tilde{m}}} |\langle g, \varphi_{\vec{n},m} \rangle| \cdot |\langle g, \varphi_{\vec{k},\tilde{m}} \rangle| \frac{1}{\langle m - \tilde{m} \rangle^{\frac{d}{2}}} \\ &= \frac{\| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^1(\pi_m)} \cdot \| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^1(\pi_{\tilde{m}})}}{\langle m - \tilde{m} \rangle^{\frac{d}{2}}}. \end{aligned}$$

Another possibility is

$$\begin{aligned} |U((a_{\vec{n},m})_{\vec{n} \in \pi_m}, (a_{\vec{k},\tilde{m}})_{\vec{k} \in \pi_{\tilde{m}}})| &\leq \left| \int_{\mathbb{R}^d} \left(\sum_{\vec{n} \in \pi_m} \langle g, \varphi_{\vec{n},m} \rangle e^{2\pi i \vec{n} \cdot x} \right) \left(\sum_{\vec{k} \in \pi_{\tilde{m}}} \langle g, \varphi_{\vec{k},\tilde{m}} \rangle e^{2\pi i \vec{k} \cdot x} \right) \varphi(x) \varphi(x) e^{2\pi i(m-\tilde{m})|x|^2} dx \right| \\ &\lesssim \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^2(\pi_m)} \cdot \| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^2(\pi_{\tilde{m}})} \end{aligned}$$

by Cauchy–Schwarz and orthogonality on the sets π_m and $\pi_{\tilde{m}}$ (recall that m and \tilde{m} are fixed). Interpolating between these bounds for $1 \leq p \leq 2$,

$$|U((a_{\vec{n},m})_{\vec{n} \in \pi_m}, (a_{\vec{k},\tilde{m}})_{\vec{k} \in \pi_{\tilde{m}}})| \lesssim \frac{\| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \cdot \| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^p(\pi_{\tilde{m}})}}{\langle m - \tilde{m} \rangle^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}}.$$

Back to (*):

$$\begin{aligned} (*)^2 &\lesssim \sum_{m: \pi_m \neq \emptyset} \sum_{\tilde{m}: \pi_{\tilde{m}} \neq \emptyset} \frac{\| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \cdot \| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^p(\pi_{\tilde{m}})}}{\langle m - \tilde{m} \rangle^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}} \\ &= \sum_{m: \pi_m \neq \emptyset} \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \sum_{\tilde{m}: \pi_{\tilde{m}} \neq \emptyset} \frac{\| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^p(\pi_{\tilde{m}})}}{\langle m - \tilde{m} \rangle^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}} \\ &\leq \| \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \|_{\ell^p(\mathbb{Z})} \left\| \sum_{\tilde{m}: \pi_{\tilde{m}} \neq \emptyset} \frac{\| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^p(\pi_{\tilde{m}})}}{\langle m - \tilde{m} \rangle^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}} \right\|_{\ell^{p'}(\mathbb{Z})} \\ &\leq \| \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \|_{\ell^p(\mathbb{Z})} \cdot \| \| \langle g, \varphi_{\cdot,\tilde{m}} \rangle \|_{\ell^p(\pi_{\tilde{m}})} \|_{\ell^p(\mathbb{Z})} \\ &= \| \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \|_{\ell^p(\mathbb{Z})}^2, \end{aligned}$$

as long as

$$\frac{1}{p} - \frac{1}{p'} = 1 - \frac{d}{2} \left(\frac{1}{p} - \frac{1}{p'} \right) \iff \frac{1}{p} - \frac{1}{p'} = \frac{2}{d+2} \iff \frac{2}{p'} = \frac{d}{d+2} \iff p' = \frac{2d+4}{d},$$

by discrete fractional integration. Plugging this back in (24),

$$\begin{aligned} \#\mathbb{X}^{l_1, l_2} &\lesssim 2^{2l_1} |E_1|^{\frac{1}{2}} \| \| \langle g, \varphi_{\cdot,m} \rangle \|_{\ell^p(\pi_m)} \|_{\ell^p(\mathbb{Z})} \\ &= 2^{2l_1} |E_1|^{\frac{1}{2}} \left(\sum_{(\vec{n},m) \in \mathbb{X}'^{l_1, l_2}} |\langle g, \varphi_{\vec{n},m} \rangle|^p \right)^{\frac{1}{p}} \lesssim 2^{2l_1} |E_1|^{\frac{1}{2}} (2^{-pl_1} \#\mathbb{X}^{l_1, l_2})^{\frac{1}{p}}, \end{aligned}$$

which implies

$$\#\mathbb{X}^{l_1, l_2} \lesssim 2^{(2+\frac{4}{d})l_1} |E_1|^{1+\frac{2}{d}}. \tag{25}$$

Interpolating between (23) and (25):

$$\begin{aligned}
 |\Lambda_d(g, h)| &\lesssim \sum_{l_1, l_2 \geq 0} 2^{-l_1} 2^{-l_2} (2^{(2+\frac{4}{d})l_1} |E_1|^{1+\frac{2}{d}})^{\theta_1} (2^{l_2} |E_2|)^{\theta_2} \\
 &= \left(\sum_{l_1 \geq 0} 2^{-l_1(1-(2+\frac{4}{d})\theta_1)} \right) \left(\sum_{l_2 \geq 0} 2^{-l_2(1-\theta_2)} \right) |E_1|^{(1+\frac{2}{d})\theta_1} |E_2|^{\theta_2} \\
 &\lesssim 2^{-\tilde{l}_1(1-(2+\frac{4}{d})\theta_1)} 2^{-\tilde{l}_2(1-\theta_2)} |E_1|^{(1+\frac{2}{d})\theta_1} |E_2|^{\theta_2} \\
 &\lesssim \min\{|E_1|^{(1-(2+\frac{4}{d})\theta_1)}, 1\} \min\{|E_2|^{1-\theta_2}, 1\} |E_1|^{(1+\frac{2}{d})\theta_1} |E_2|^{\theta_2} \\
 &\lesssim |E_1|^{\alpha_1(1-(2+\frac{4}{d})\theta_1)+(1+\frac{2}{d})\theta_1} |E_2|^{\alpha_2(1-\theta_2)+\theta_2}
 \end{aligned} \tag{26}$$

for all $0 \leq \alpha_1, \alpha_2 \leq 1$, $\theta_1 + \theta_2 = 1$, with $0 \leq (2 + \frac{4}{d})\theta_1 < 1$, $0 \leq \theta_2 < 1$, where \tilde{l}_1 is the smallest possible value of l_1 for which $\mathbb{A}^{l_1} \neq \emptyset$ and \tilde{l}_2 is defined analogously. Picking $\alpha_1 = \frac{1}{2}$, $\alpha_2 = 0$, $\theta_1 = \frac{d}{2(d+2)} - \varepsilon$ and $\theta_2 = \frac{d+4}{2(d+2)} + \varepsilon$ gives

$$|\Lambda_d(g, h)| \lesssim_\varepsilon |E_1|^{\frac{1}{2}} \cdot |E_2|^{\frac{d+4}{2(d+2)} + \varepsilon}$$

for all $\varepsilon > 0$, which proves the proposition by restricted weak-type interpolation.

6. Proof of Proposition 4.3-Conjecture 1.1 for E_1 ($k = 1, d = 1, p = 4$)

The following argument is inspired by Zygmund’s original proof of this case. Define

$$\Phi_{n,m}(s, t) := |t - s|^{\frac{1}{2}} \varphi(s) \varphi(t) e^{2\pi i(s-t)n} e^{2\pi i(s^2-t^2)m}.$$

Claim 6.1. $\langle \Phi_{n,m}, \Phi_{\tilde{n},\tilde{m}} \rangle = O_N \left(\frac{1}{|(n - \tilde{n})(m - \tilde{m})|^N} \right)$

for any natural N if $n \neq \tilde{n}$ and $m \neq \tilde{m}$.

Proof. We have

$$\begin{aligned}
 \langle \Phi_{n,m}, \Phi_{\tilde{n},\tilde{m}} \rangle &= \iint_{[0,1]^2} |t - s| |\varphi(s)|^2 |\varphi(t)|^2 e^{2\pi i(s-t)(n-\tilde{n})} e^{2\pi i(s^2-t^2)(m-\tilde{m})} ds dt \\
 &= \iint_R \frac{|u|}{|u|} \psi(u, v) e^{2\pi i u(n-\tilde{n})} e^{2\pi i v(m-\tilde{m})} du dv,
 \end{aligned}$$

where R is the region that we obtain after making the change of variables $s - t = u$, $s^2 - t^2 = v$, and

$$\psi(u, v) = \varphi \otimes \varphi \left(\frac{v + u^2}{u}, \frac{v - u^2}{u} \right).$$

The claim follows by the nonstationary phase [Theorem B.2](#). □

We now prove the following:

Lemma 6.2. For G smooth supported on $[0, 1] \times [0, 1]$,

$$\left\| \sum_{n,m \in \mathbb{Z}} \langle G, \varphi_{n,m} \otimes \bar{\varphi}_{\tilde{n},\tilde{m}} \rangle (\chi_n \otimes \chi_m) \right\|_2 \lesssim \left(\iint_{[0,1]^2} \frac{|G(s,t)|^2}{|s-t|} ds dt \right)^{\frac{1}{2}}.$$

Proof. Define

$$\tilde{G}(s, t) = \frac{G(s, t)}{|s - t|^{\frac{1}{2}}}$$

on $[0, 1]^2 \setminus \{(x, x); 0 \leq x \leq 1\}$. Observe that

$$\left\| \sum_{n,m \in \mathbb{Z}} \langle G, \varphi_{n,m} \otimes \bar{\varphi}_{n,m} \rangle (\chi_n \otimes \chi_m) \right\|_2^2 = \sum_{n,m \in \mathbb{Z}} |\langle G, \varphi_{n,m} \otimes \bar{\varphi}_{n,m} \rangle|^2 = \sum_{n,m \in \mathbb{Z}} |\langle \tilde{G}, \Phi_{n,m} \rangle|^2 \lesssim \|\tilde{G}\|_2^2,$$

by the almost orthogonality of the $\Phi_{n,m}$ proved in the previous claim. □

Remark 6.3. By the triangle inequality,

$$\left\| \sum_{n,m \in \mathbb{Z}} \langle G, \varphi_{n,m} \otimes \bar{\varphi}_{n,m} \rangle (\chi_n \otimes \chi_m) \right\|_\infty \lesssim \iint_{[0,1]^2} |G(s, t)| \, ds \, dt.$$

Hence by interpolation we obtain

$$\left\| \sum_{n,m \in \mathbb{Z}} \langle G, \varphi_{n,m} \otimes \bar{\varphi}_{n,m} \rangle (\chi_n \otimes \chi_m) \right\|_p \lesssim \left(\iint_{[0,1]^2} \frac{|G(s, t)|^{p'}}{|s - t|^{p'-1}} \, ds \, dt \right)^{\frac{1}{p'}} \tag{27}$$

for $2 \leq p \leq \infty$.

Let $E \subset \mathbb{R}^d$ be a measurable set of finite measure with $|g| \leq \chi_E$. Using [Remark 6.3](#) and [Lemma 6.2](#) for $G = g \otimes \bar{g}$, we have

$$\begin{aligned} \left[\sum_{n,m \in \mathbb{Z}} |\langle g, \varphi_{n,m} \rangle|^{4+\varepsilon} \right]^{\frac{2}{4+\varepsilon}} &= \left[\int_{\mathbb{R}^2} \left(\sum_{n,m \in \mathbb{Z}} |\langle g, \varphi_{n,m} \rangle|^{4+\varepsilon} (\chi_n \otimes \chi_m) \right) \right]^{\frac{2}{4+\varepsilon}} \\ &\leq \left[\int_{\mathbb{R}^2} \left(\sum_{n,m \in \mathbb{Z}} |\langle g, \varphi_{n,m} \rangle|^2 (\chi_n \otimes \chi_m) \right)^{\frac{4+\varepsilon}{2}} \right]^{\frac{2}{4+\varepsilon}} \\ &= \left\| \sum_{n,m \in \mathbb{Z}} |\langle g, \varphi_{n,m} \rangle|^2 (\chi_n \otimes \chi_m) \right\|_{2+\frac{\varepsilon}{2}} \\ &\lesssim \left(\iint_{[0,1]^2} \frac{|g(s)|^{p'} |g(t)|^{p'}}{|s - t|^{p'-1}} \, ds \, dt \right)^{\frac{1}{p'}}, \quad \text{where } p' = \frac{4 + \varepsilon}{2 + \varepsilon}. \end{aligned}$$

To bound this last integral, we proceed as follows:

$$\begin{aligned} \int_0^1 \int_0^1 \frac{|\rho(s)| \cdot |\rho(t)|}{|s - t|^\gamma} \, ds \, dt &= \int_0^1 |\rho(t)| \int_0^1 \frac{|\rho(s)|}{|s - t|^\gamma} \, ds \, dt = \int_0^1 |\rho(t)| \cdot \left(|\rho| * \frac{1}{|s|^\gamma} \right) (t) \, dt \\ &= \left\| |\rho| \left(|\rho| * \frac{1}{|s|^\gamma} \right) \right\|_{L^1(dt)} \leq \|\rho\|_{L^q(dt)} \left\| |\rho| * \frac{1}{|s|^\gamma} \right\|_{L^{p'}(dt)} \lesssim_\varepsilon \|\rho\|_p^2 \end{aligned}$$

if $\frac{1}{p'} = \frac{1}{p} - (1 - \gamma)$, by [Theorem B.1](#). In our case, $\rho = |g|^{p'}$, $\gamma = p' - 1$ and

$$pp' = \frac{(4 + \varepsilon)^2}{2(2 + \varepsilon)} > 4.$$

Then

$$\begin{aligned} \left(\int_0^1 \int_0^1 \frac{|g(s)|^{p'} \cdot |g(t)|^{p'}}{|s-t|^{p'-1}} ds dt \right)^{\frac{1}{p'}} &\lesssim \left(\int_0^1 |g(t)|^{pp'} dt \right)^{\frac{2}{pp'}} = \left(\int_0^1 |g(t)|^{4+\frac{(4+\varepsilon)^2}{2(2+\varepsilon)}-4} dt \right)^{\frac{4(2+\varepsilon)}{(4+\varepsilon)^2}} \\ &\lesssim \left(\int_0^1 |g(t)|^4 dt \right)^{\frac{4(2+\varepsilon)}{(4+\varepsilon)^2}} = |E|^{\frac{4(2+\varepsilon)}{(4+\varepsilon)^2}}. \end{aligned}$$

Observed that in the second line of the chain of inequalities above we used the fact that $|g| \leq 1$. Finally,

$$\|E_1 g\|_{4+\varepsilon} = \left[\sum_{n,m \in \mathbb{Z}} |\langle g, \varphi_{n,m} \rangle|^{4+\varepsilon} \right]^{\frac{1}{4+\varepsilon}} \lesssim |E|^{\frac{2(2+\varepsilon)}{(4+\varepsilon)^2}} \leq |E|^{\frac{1}{4}}.$$

This shows that E_1 maps $L^4([0, 1])$ to $L^q(\mathbb{R}^2)$ for any $q > 4$ by restricted weak-type interpolation.

7. Proof of Proposition 4.4-Conjecture 1.2 for $ME_{2,1}$ ($k = 2, d = 1$)

The model to be treated is

$$ME_{2,1}(f, g) := \sum_{(n,m) \in \mathbb{Z}^2} \langle f, \varphi_{n,m}^1 \rangle \cdot \langle g, \varphi_{n,m}^2 \rangle (\chi_n \otimes \chi_m).$$

Since $d = 1$, we do not have to deal with the multivariable quantity

$$\varphi_{\vec{n},m}^j = \bigotimes_{l=1}^d \varphi_{n_l,m}^{l,j}$$

from Definition 2.3, so we will simplify the notation by taking $\varphi_{n,m}^1 := \varphi_{n,m}^{1,1}$ and $\varphi_{n,m}^2 := \varphi_{n,m}^{1,2}$. We also replaced (g_1, g_2) by (f, g) here to reduce the number of indices carried through the section.

We provide a simple argument involving Bessel’s inequality. After a change of variables to move the domain of φ^2 to be the same as the one of φ^1 , we have

$$\begin{aligned} |ME_{2,1}(f, g)| &\lesssim \sum_{(n,m) \in \mathbb{Z}^2} |\langle f, \varphi_{n,m}^1 \rangle| |\overline{\langle (g)_{-4}, \varphi_{n+8m,m}^1 \rangle}| (\chi_n \otimes \chi_m) \\ &= \sum_{(n,m) \in \mathbb{Z}^2} |\langle f \otimes (g)_{-4}, \varphi_{n,m}^1 \otimes \bar{\varphi}_{n+8m,m}^1 \rangle| (\chi_n \otimes \chi_m), \end{aligned}$$

where¹³ $(g)_{-4}(y) = g(y + 4)$. Observe that

$$\begin{aligned} &\langle f \otimes (g)_{-4}, \varphi_{n,m}^1 \otimes \bar{\varphi}_{n+8m,m}^1 \rangle \\ &= \iint f(x)g(y+4)\varphi^1(x)\varphi^1(y)e^{-2\pi i n x} e^{-2\pi i m x^2} e^{2\pi i(n+8m)y} e^{2\pi i m y^2} dx dy \\ &= \iint f(x)g(y+4)e^{2\pi i n(y-x)} e^{2\pi i m(y-x)(y+x)} e^{16\pi i m y} dx dy \\ &\approx \int \underbrace{\left[\int f\left(\frac{v-u}{2}\right)g\left(\frac{v+u}{2}+4\right)e^{2\pi i m u v} e^{8\pi i m(u+v)} dv \right]}_{H_m(u)} e^{2\pi i n u} du = \hat{H}_m(-n) \end{aligned}$$

¹³This was done to bring the support of $\varphi_{n,m}^2$ to the one of $\varphi_{n+8m,m}^1$. The price to pay is the $+4m$ shift in the linear modulation index of the bump.

Hence

$$\|ME_{2,1}(f, g)\|_2^2 \lesssim \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} |\hat{H}_m(-n)|^2 = \sum_{m \in \mathbb{Z}} \|H_m\|_2^2,$$

by Bessel. On the other hand,

$$\begin{aligned} \|H_m\|_2^2 &= \int \left| \int f\left(\frac{v-u}{2}\right) g\left(\frac{v+u}{2} + 4\right) e^{2\pi i m u v} e^{8\pi i m(u+v)} dv \right|^2 du \\ &= \int \underbrace{\left| \int f\left(\frac{v-u}{2}\right) g\left(\frac{v+u}{2} + 4\right) e^{2\pi i m v(u+4)} dv \right|^2}_{\tilde{H}_u(v)} du = \int |\hat{\tilde{H}}_u(m(u+4))|^2 du. \end{aligned}$$

Transversality enters the picture here through the factor $(u+4)$ above: the $+4$ shift in u comes from the fact that the supports of φ^1 and φ^2 are disjoint and far enough from each other; hence $u+4 \geq c > 0$. This way,

$$\begin{aligned} \|ME_{2,1}(f, g)\|_2^2 &\lesssim \int \left(\sum_{m \in \mathbb{Z}} |\hat{\tilde{H}}_u(m(u+4))|^2 \right) du \\ &\lesssim \iint |\tilde{H}_u(v)|^2 dv du \lesssim \|f\|_2^2 \|g\|_2^2, \end{aligned}$$

by Bessel again.

8. Case $k = 1$ of Theorem 1.5

In this section we start the proof of Theorem 1.5. There are two main ingredients in the argument for the case $k = 1$: Proposition 4.3 and the fact that the wave packets

$$\varphi_{\vec{n},m}(x) := \varphi(x_1) \cdots \varphi(x_d) e^{2\pi i x \cdot \vec{n}} e^{2\pi i |x|^2 m}$$

are almost orthogonal for a fixed m and \vec{n} varying in \mathbb{Z}^d . The latter fact will be exploited through Bessel’s inequality whenever possible. Recall from Remark 4.5 that, since $g = g_1 \otimes \cdots \otimes g_d$, it suffices to study the multilinear form

$$\Lambda_d(g_1, \dots, g_d, h) = \sum_{\vec{n} \in \mathbb{Z}^d, m \in \mathbb{Z}} \prod_{j=1}^d \langle g_j, \varphi_{n_j, m} \rangle \cdot \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle,$$

Now we focus on obtaining (22). Let $E_j \subset [0, 1], 1 \leq j \leq d$, and $F \subset \mathbb{R}^{d+1}$ be measurable sets for which $|g_j| \leq \chi_{E_j}$ and $|h| \leq \chi_F$. Define the sets

$$\begin{aligned} \mathbb{A}_j^{l_j} &:= \{(n_j, m) \in \mathbb{Z}^2 : |\langle g_j, \varphi_{n_j, m} \rangle| \approx 2^{-l_j}\}, \quad 1 \leq j \leq d. \\ \mathbb{B}^{l_{d+1}} &:= \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-l_{d+1}}\}, \\ \mathbb{X}^{l_1, \dots, l_{d+1}} &:= \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_j, m) \in \mathbb{A}_j^{l_j}, 1 \leq j \leq d\} \cap \mathbb{B}^{l_{d+1}}. \end{aligned}$$

Hence,

$$|\Lambda_d(g_1, \dots, g_d, h)| \lesssim \sum_{l_1, \dots, l_{d+1} \in \mathbb{Z}} 2^{-l_1} \cdots 2^{-l_{d+1}} \#\mathbb{X}^{l_1, \dots, l_{d+1}}.$$

As in Section 5, we know that $l_1, \dots, l_{d+1} \geq 0$. We can estimate $\#\mathbb{X}^{l_1, \dots, l_{d+1}}$ using the function h :

$$\#\mathbb{X}^{l_1, \dots, l_{d+1}} \lesssim 2^{l_{d+1}} \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \lesssim 2^{l_{d+1}} |F|. \tag{28}$$

Alternatively, many bounds for $\#\mathbb{X}^{l_1, \dots, l_{d+1}}$ can be obtained using the input functions g_1, \dots, g_d :

$$\begin{aligned} \#\mathbb{X}^{l_1, \dots, l_{d+1}} &\lesssim \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \mathbb{1}_{\mathbb{A}_1^{l_1}}(n_1, m) \cdots \mathbb{1}_{\mathbb{A}_d^{l_d}}(n_d, m) \\ &= \sum_{m \in \mathbb{Z}} \sum_{n_1 \in \mathbb{Z}} \cdots \sum_{n_{d-1} \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_1^{l_1}}(n_1, m) \cdots \mathbb{1}_{\mathbb{A}_{d-1}^{l_{d-1}}}(n_{d-1}, m) \underbrace{\sum_{n_d \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_d^{l_d}}(n_d, m)}_{\alpha_{d,m}} \end{aligned} \tag{29}$$

Observe that $\alpha_{d,m} = \#\{n; (n, m) \in \mathbb{A}_d^{l_d}\}$ and $(n, m) \in \mathbb{A}_d^{l_d} \Rightarrow 1 \lesssim 2^{2l_d} |\langle g_d, \varphi_{n,m} \rangle|^2$. Adding up in n ,

$$\alpha_{d,m} \lesssim 2^{2l_d} \sum_{n: (n,m) \in \mathbb{A}_d^{l_d}} |\langle g_d, \varphi_{n,m} \rangle|^2 \lesssim 2^{2l_d} |E_d|$$

by orthogonality. Notice that this quantity does not depend on m ; therefore we can iterate this argument for $d - 2$ of the remaining $d - 1$ characteristic functions:

$$\begin{aligned} \#\mathbb{X}^{l_1, \dots, l_{d+1}} &\lesssim 2^{2l_d} |E_d| \sum_{m \in \mathbb{Z}} \sum_{n_1 \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_1^{l_1}}(n_1, m) \cdots \mathbb{1}_{\mathbb{A}_{d-2}^{l_{d-2}}}(n_{d-2}, m) \underbrace{\sum_{n_{d-1} \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_{d-1}^{l_{d-1}}}(n_{d-1}, m)}_{\alpha_{d-1,m}} \\ &\lesssim 2^{2l_d} |E_d| 2^{2l_{d-1}} |E_{d-1}| \sum_{m \in \mathbb{Z}} \sum_{n_1 \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_1^{l_1}}(n_1, m) \cdots \mathbb{1}_{\mathbb{A}_{d-3}^{l_{d-3}}}(n_{d-3}, m) \sum_{n_{d-2} \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_{d-2}^{l_{d-2}}}(n_{d-2}, m) \\ &\lesssim 2^{2l_d} 2^{2l_{d-1}} \dots 2^{2l_2} |E_d| \cdots |E_2| \underbrace{\sum_{m \in \mathbb{Z}} \sum_{n_1 \in \mathbb{Z}} \mathbb{1}_{\mathbb{A}_1^{l_1}}(n_1, m)}_{\#\mathbb{A}_1^{l_1}}. \end{aligned} \tag{30}$$

To bound $\#\mathbb{A}_1^{l_1}$ we can use Proposition 4.3. For $\varepsilon > 0$ we have

$$\begin{aligned} (n, m) \in \mathbb{A}_1^{l_1} &\Rightarrow 1 \lesssim 2^{(4+\varepsilon)l_1} |\langle g_1, \varphi_{n,m} \rangle|^{4+\varepsilon} \\ &\Rightarrow \#\mathbb{A}_1^{l_1} \lesssim 2^{(4+\varepsilon)l_1} \sum_{(n,m) \in \mathbb{A}_1^{l_1}} |\langle g_1, \varphi_{n,m} \rangle|^{4+\varepsilon} \lesssim_\varepsilon 2^{(4+\varepsilon)l_1} |E_1|. \end{aligned}$$

Using this above,

$$\#\mathbb{X}^{l_1, \dots, l_{d+1}} \lesssim_\varepsilon 2^{2l_d} 2^{2l_{d-1}} \dots 2^{2l_2} 2^{(4+\varepsilon)l_1} |E_d| \cdots |E_2| |E_1|. \tag{31}$$

We could have used the L^4 - $L^{4+\varepsilon}$ bound for any g_j and a Bessel bound for the remaining ones. More precisely, if $\sigma \in S_d$ is a permutation, we have

$$\#\mathbb{X}^{l_1, \dots, l_{d+1}} \lesssim_\varepsilon 2^{2l_{\sigma(d)}} 2^{2l_{\sigma(d-1)}} \dots 2^{2l_{\sigma(2)}} 2^{(4+\varepsilon)l_{\sigma(1)}} |E_{\sigma(d)}| \cdots |E_{\sigma(2)}| |E_{\sigma(1)}|. \tag{32}$$

This amounts to exactly d different estimates. Interpolating between all of them with equal weight $\frac{1}{d}$, we obtain

$$\begin{aligned} \#\times^{l_1, \dots, l_{d+1}} &\lesssim_\varepsilon 2^{\frac{2(d-1)+4+\varepsilon}{d}l_1} \dots 2^{\frac{2(d-1)+4+\varepsilon}{d}l_d} |E_1| \dots |E_d| \\ &= 2^{(2+\frac{2}{d}+\frac{\varepsilon}{d})l_1} \dots 2^{(2+\frac{2}{d}+\frac{\varepsilon}{d})l_d} |E_1| \dots |E_d|. \end{aligned} \tag{33}$$

Finally, we interpolate between bounds (28) and (33):

$$\begin{aligned} |\Lambda_d(g_1, \dots, g_d, h)| &\lesssim \sum_{l_1, \dots, l_{d+1} \in \mathbb{Z}_+} 2^{-l_1} \dots 2^{-l_{d+1}} \#\times^{l_1, \dots, l_{d+1}} \\ &\lesssim \sum_{l_1, \dots, l_{d+1} \in \mathbb{Z}_+} 2^{-l_1} \dots 2^{-l_{d+1}} (2^{(2+\frac{2}{d}+\frac{\varepsilon}{d})l_1} \dots 2^{(2+\frac{2}{d}+\frac{\varepsilon}{d})l_d} |E_1| \dots |E_d|)^{\theta_1} (2^{l_{d+1}} |F|)^{\theta_2} \\ &\lesssim \left(\sum_{l_{d+1} \geq 0} 2^{-(1-\theta_2)l_{d+1}} |F|^{\theta_2} \right) \prod_{j=1}^d \sum_{l_j \geq 0} 2^{-(1-(2+\frac{2}{d}+\frac{\varepsilon}{d})\theta_1)l_j} |E_j|^{\theta_1} \\ &\lesssim |E_1|^{\alpha(1-(2+\frac{2}{d}+\frac{\varepsilon}{d})\theta_1)+\theta_1} \dots |E_d|^{\alpha(1-(2+\frac{2}{d}+\frac{\varepsilon}{d})\theta_1)+\theta_1} |F|^{\theta_2} \end{aligned}$$

for any $0 \leq \alpha \leq 1$. On the other hand, for several of the series above to converge we need $(2 + \frac{2}{d} + \frac{\varepsilon}{d})\theta_1 > 1$. By choosing the appropriate α and θ_1 close to $(2 + \frac{2}{d})^{-1}$, one concludes this case.

9. Case $2 \leq k \leq d + 1$ of Theorem 1.5

Recall that we fixed a set of weakly transversal cubes $Q = \{Q_1, \dots, Q_k\}$ in Section 3 and let g_j be supported on Q_j . The averaged k -linear extension operator¹⁴ in \mathbb{R}^d is given by

$$\text{ME}_{k,d}^{\frac{1}{k}}(g_1, \dots, g_k) = \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \left(\prod_{j=1}^k |\langle g_j, \varphi_{\vec{n}, m}^j \rangle| \right)^{\frac{1}{k}} (\chi_{\vec{n}} \otimes \chi_m).$$

The conjectured bounds for it are

$$\| \text{ME}_{k,d}^{\frac{1}{k}}(g_1, \dots, g_k) \|_{L^p(\mathbb{R}^{d+1})} \lesssim \prod_{j=1}^k \|g_j\|_{L^2(Q_j)}^{\frac{1}{k}} \quad \text{for all } p \geq \frac{2(d+k+1)}{(d+k-1)}. \tag{34}$$

As done in the case $k = 1$, it's enough to prove certain restricted weak-type bounds for its associated form

$$\tilde{\Lambda}_{k,d}(g, h) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \left(\prod_{i=1}^k |\langle g_i, \varphi_{\vec{n}, m}^i \rangle| \right)^{\frac{1}{k}} \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle, \tag{35}$$

where $g := (g_1, \dots, g_k)$ by a slight abuse of notation.

¹⁴We consider this averaged version of $\text{ME}_{k,d}$ for technical reasons. The conjectured bounds for it have a Banach space as target, as opposed to the quasi-Banach space (for most k and d) $L^{2(d+k+1)/(k(d+k-1))}$ that is the target of Conjecture 1.2. The fact that L^p for $p \geq 2(d+k+1)/(d+k-1)$ is Banach lets us use (49) effectively in the interpolation argument, since it forces the final power γ on $|F|^\gamma$ to be positive.

When $k = d = 2$, Conjecture 1.2 has $L^{5/3}$ as target. We will discuss this case first to help digest the main ideas of the general argument, and since this space is Banach, we can work directly with $\text{ME}_{2,2}$ instead of considering the averaged operator $\text{ME}_{2,2}^{1/2}$.

Remark 9.1. We will prove (34) up to the endpoint assuming that g_1 is a full tensor, but the argument can be repeated if any other g_j is assumed to be of this type. As the reader will notice, the proof depends on the fact that we can find $k - 1$ canonical directions associated to Q_j , which is the defining property of a weakly transversal collection of cubes with pivot Q_j . In what follows, we are taking $\{e_1, \dots, e_{k-1}\}$ to be the set of directions associated to Q_1 .

Remark 9.2. As we mentioned in Remark 1.7, under weak transversality alone we do not need g_1 to be a full tensor to prove the case $2 \leq k \leq d$ of Theorem 1.5. In fact, the following structure is enough in this section:

$$g_1(x_1, \dots, x_d) = g_{1,1}(x_1) \cdot g_{1,2}(x_2) \cdots g_{1,k-1}(x_{k-1}) \cdot g_{1,k}(x_k, \dots, x_d).$$

Notice that we have $k - 1$ single-variable functions and one function in $d - k + 1$ variables. The single-variable ones are defined along $k - 1$ canonical directions $\{e_1, \dots, e_{k-1}\}$ associated to Q_1 , and $g_{1,k}$ is a function in the remaining variables.

In general, if we are given a weakly transversal collection \tilde{Q} , for a fixed $1 \leq j \leq k - 1$ we have a set of associated directions $\mathcal{E}_j = \{e_{i_1}, \dots, e_{i_{k-1}}\}$ (see Definition 3.2). Denote by $x_{\mathcal{E}_j^c}$ the vector of $d - k + 1$ entries obtained after removing $x_{i_1}, \dots, x_{i_{k-1}}$ from (x_1, \dots, x_d) . Assuming that the functions g_l for $l \neq j$ are generic and that g_j has the weaker tensor structure

$$g_j(x_1, \dots, x_d) = g_{j,1}(x_{i_1}) \cdots g_{j,k-1}(x_{i_{k-1}}) \cdot g_{\mathcal{E}_j^c}(x_{\mathcal{E}_j^c}) \tag{36}$$

will suffice to conclude Theorem 1.5 for \tilde{Q} through the argument that we will present in this section.

Remark 9.3. As a consequence of Claim 3.4, a collection $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ of transversal cubes generates finitely many subcollections \tilde{Q} of weakly transversal ones (after partitioning each Q_l into small enough cubes and defining new collections with them). However, for a fixed $1 \leq j \leq k$, the associated $k - 1$ directions in \mathcal{E}_j can potentially change from one such weakly transversal subcollection to another, and this is why we assume g_j to be a full tensor under the transversality assumption.

In this section we will use the following conventions:

- The variables of g_j are x_1, x_2, \dots, x_d , but we will split them in two groups: $k - 1$ blocks of one variable represented by $x_i, 1 \leq i \leq k - 1$, and one block of $d - k + 1$ variables $\vec{x}_k = (x_k, x_{k+1}, \dots, x_{d-1}, x_d)$.
- The index x_i in $\langle \cdot, \cdot \rangle_{x_i}$ indicates that the inner product is an integral in the variable x_i only. For instance,

$$\langle g_j, \varphi \rangle_{x_1} := \int_{\mathbb{R}} g_j(x_1, \dots, x_d) \cdot \bar{\varphi}(x_1, \dots, x_d) dx_1 \tag{37}$$

is now a function of the variables x_2, \dots, x_d . The vector index \vec{x}_k in $\langle \cdot, \cdot \rangle_{\vec{x}_k}$ is understood analogously:

$$\langle g_j, \varphi \rangle_{\vec{x}_k} := \int_{\mathbb{R}^{d-k+1}} g_j(x_1, \dots, x_d) \cdot \bar{\varphi}(x_1, \dots, x_d) d\vec{x}_k \tag{38}$$

- The expression $\|\langle g_j, \cdot \rangle_{x_i}\|_2$ is the L^2 norm of a function in the variables x_l , $1 \leq l \leq k - 1$, $l \neq i$. To illustrate using (37),

$$\|\langle g_j, \varphi \rangle_{x_1}\|_2 = \left[\int_{\mathbb{R}^{d-1}} \left| \int_{\mathbb{R}} g_j(x_1, \dots, x_d) \cdot \bar{\varphi}(x_1, \dots, x_d) dx_1 \right|^2 dx_2 \cdots dx_d \right]^{\frac{1}{2}}.$$

The quantity $\|\langle g_j, \cdot \rangle_{\vec{x}_k}\|_2$ is defined analogously as

$$\|\langle g_j, \varphi \rangle_{\vec{x}_k}\|_2 = \left[\int_{\mathbb{R}^{k-1}} \left| \int_{\mathbb{R}^{d-k+1}} g_j(x_1, \dots, x_d) \cdot \bar{\varphi}(x_1, \dots, x_d) dx_k \right|^2 dx_1 \cdots dx_{k-1} \right]^{\frac{1}{2}}.$$

- For $\vec{n} = (n_1, \dots, n_d)$, define the vector

$$\hat{n}_i := (n_1, \dots, n_{i-1}, n_{i+1}, \dots, n_d).$$

In other words, the hat on \hat{n}_i indicates that n_i was removed from the vector \vec{n} . For $f : \mathbb{Z}^d \rightarrow \mathbb{C}$, define

$$\|f(\vec{n})\|_{\ell^1_{\hat{n}_i}} := \sum_{\hat{n}_i \in \mathbb{Z}^{d-1}} |f(\vec{n})|.$$

That is, $\|f(\vec{n})\|_{\ell^1_{\hat{n}_i}}$ is the ℓ^1 norm of f over all n_1, \dots, n_d , except for n_i . Hence $\|f(\vec{n})\|_{\ell^1_{\hat{n}_i}}$ is a function of the remaining variable n_i . The quantity $\|f(\vec{n})\|_{\ell^1_{\hat{n}_k}}$ is defined analogously as

$$\|f(\vec{n})\|_{\ell^1_{\hat{n}_k}} := \sum_{(n_1, \dots, n_{k-1}) \in \mathbb{Z}^{k-1}} |f(\vec{n})|.$$

Finally, the integral $\int g d\hat{x}_i$ means

$$\int g(x_1, \dots, x_d) d\hat{x}_i := \int g(x_1, \dots, x_d) dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_d.$$

In what follows, let $E_{1,1}, \dots, E_{1,k-1} \subset [0, 1]$, $E_{1,k} \subset [0, 1]^{d-k+1}$, $E_j \subset Q_j$ ($2 \leq j \leq k$) and $F \subset \mathbb{R}^{d+1}$ be measurable sets such that $|g_{1,l}| \leq \chi_{E_{1,l}}$ for $1 \leq l \leq k - 1$, $|g_{1,k}| \leq \chi_{E_{1,k}}$, $|g_j| \leq \chi_{E_j}$ for $2 \leq j \leq k$ and $|h| \leq \chi_F$. Furthermore, $E_1 := E_{1,1} \times \cdots \times E_{1,k-1} \times E_{1,k}$.

A rough description of the argument in one sentence is: *the proof is a combination of Strichartz in some variables and bilinear extension in many pairs of the other variables*. In order to illustrate that, we will first present the simplest case in an informal way, which means that we will avoid the purely technical aspects in this preliminary part. Once this is understood, it will be clear how to rigorously extend the argument in general.

9A. Understanding the core ideas in the $k = d = 2$ case. Consider the model

$$ME_{2,2}(g_1, g_2) = \sum_{(\vec{n}, m) \in \mathbb{Z}^3} \langle g_1, \varphi_{\vec{n}, m}^1 \rangle \langle g_2, \varphi_{\vec{n}, m}^2 \rangle (\chi_{\vec{n}} \otimes \chi_m)$$

and its associated trilinear form¹⁵

$$\tilde{\Lambda}_{2,2}(g_1, g_2, h) = \sum_{(\vec{n}, m) \in \mathbb{Z}^3} \langle g_1, \varphi_{\vec{n}, m}^1 \rangle \langle g_2, \varphi_{\vec{n}, m}^2 \rangle (\chi_{\vec{n}} \otimes \chi_m).$$

Assuming that $g_1 = g_{1,1} \otimes g_{1,2}$, we want to prove that

$$|\tilde{\Lambda}_{2,2}(g_1, g_2)| \lesssim_\varepsilon |E_1|^{\frac{1}{2}} \cdot |E_2|^{\frac{1}{2}} \cdot |F|^{\frac{2}{5} + \varepsilon}$$

for all $\varepsilon > 0$. The $L^2 \times L^2 \mapsto L^{5/3 + \varepsilon}$ bound will then follow by multilinear interpolation and [Remark 4.5](#).

Given the expository character of this subsection, we adopt the informal convention

$$\begin{cases} x^+ \text{ means } x + \delta, \text{ where } \delta > 0 \text{ is arbitrarily small,} \\ x^- \text{ means } x - \delta, \text{ where } \delta > 0 \text{ is arbitrarily small.} \end{cases}$$

We will always be able to control how small the δ above is, so we do not worry about making it precise for now.

The first step is to define the level sets of the scalar products appearing in $\text{ME}_{2,2}$:

$$\begin{aligned} \mathbb{A}_1^{l_1} &= \{(\vec{n}, m) : |\langle g_1, \varphi_{\vec{n}, m}^1 \rangle| \approx 2^{-l_1}\}, \\ \mathbb{A}_2^{l_2} &= \{(\vec{n}, m) : |\langle g_2, \varphi_{\vec{n}, m}^2 \rangle| \approx 2^{-l_2}\}. \end{aligned}$$

Transversality will be captured by exploiting the sizes of “lower-dimensional” information: in fact, we want to make the operator $\text{ME}_{2,1}$ appear, and this will be possible thanks to the interaction between the quantities associated to the level sets

$$\begin{aligned} \mathbb{B}_1^{r_1} &= \{(n_1, m) : \|\langle g_1, \varphi_{n_1, m}^{1,1} \rangle_{x_1}\|_2 \approx 2^{-r_1}\}, \\ \mathbb{C}_1^{s_1} &= \{(n_1, m) : \|\langle g_2, \varphi_{n_1, m}^{1,2} \rangle_{x_1}\|_2 \approx 2^{-s_1}\}. \end{aligned}$$

Since there is only one direction along which one can exploit transversality, we will use the L^2 theory for E_1 (i.e., Strichartz) along the remaining one. In order to do that, the following level sets will be used:

$$\begin{aligned} \mathbb{B}_2^{r_2} &= \{(n_2, m) : \|\langle g_1, \varphi_{n_2, m}^{2,1} \rangle_{x_2}\|_2 \approx 2^{-r_2}\}, \\ \mathbb{C}_2^{s_2} &= \{(n_2, m) : \|\langle g_2, \varphi_{n_2, m}^{2,2} \rangle_{x_2}\|_2 \approx 2^{-s_2}\}. \end{aligned}$$

The size of the scalar product involving h will be captured by the set

$$\mathbb{D}^k = \{(\vec{n}, m) : |\langle H, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-k}\}.$$

We will also need to organize all the information above in appropriate “slices” and in a major set that takes everything into account. The sets that do that are

$$\begin{aligned} \mathbb{X}^{l_2, s_1} &:= \mathbb{A}_2^{l_2} \cap \{(\vec{n}, m) : (n_1, m) \in \mathbb{C}_1^{s_1}\}, \\ \mathbb{X}^{l_2, s_2} &:= \mathbb{A}_2^{l_2} \cap \{(\vec{n}, m) : (n_2, m) \in \mathbb{C}_2^{s_2}\}, \\ \mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k} &= \mathbb{A}_1^{l_1} \cap \mathbb{A}_2^{l_2} \cap \{(\vec{n}, m) : (n_1, m) \in \mathbb{B}_1^{r_1} \cap \mathbb{C}_1^{s_1}, (n_2, m) \in \mathbb{B}_2^{r_2} \cap \mathbb{C}_2^{s_2}\} \cap \mathbb{D}^k, \end{aligned}$$

¹⁵There is a slight abuse of notation here: we are using $\tilde{\Lambda}_{2,2}$ for the form associated to $\text{ME}_{2,2}$ and not for its averaged version $\text{ME}_{2,2}$, as established in the beginning of this section.

where we are using the abbreviations $\vec{l} = (l_1, l_2)$, $\vec{r} = (r_1, r_2)$ and $\vec{s} = (s_1, s_2)$. This gives us

$$|\tilde{\Lambda}_{2,2}(g_1, g_2, h)| \lesssim \sum_{\vec{l}, \vec{r}, \vec{s}, k} 2^{-l_1} 2^{-l_2} 2^{-k} \#\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k}.$$

For the sake of simplicity, let us assume that $g_1 = \mathbb{1}_{E_{1,1}} \otimes \mathbb{1}_{E_{1,2}}$, $g_2 = \mathbb{1}_{E_2}$ and $h = \mathbb{1}_F$.¹⁶ We will need efficient ways of relating the scalar and mixed-norm quantities above. A direct computation (using the definition of $\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k}$) shows that

$$2^{-l_1} = \frac{2^{-r_1} \cdot 2^{-r_2}}{|E_1|^{\frac{1}{2}}}. \tag{39}$$

Using Bessel along a direction, for $(n_1, n_2, m) \in \mathbb{X}^{l_2, s_1}$ we have

$$\begin{aligned} 1 \approx 2^{2l_2} |\langle g_2, \varphi_{\vec{n}, m}^2 \rangle|^2 &\implies \#\mathbb{X}_{(n_1, m)}^{l_2, s_1} \approx 2^{2l_2} \sum_{n_2 \in \mathbb{X}_{(n_1, m)}^{l_2, s_1}} |\langle g_2, \varphi_{\vec{n}, m}^2 \rangle|^2 \\ &\implies \#\mathbb{X}_{(n_1, m)}^{l_2, s_1} \lesssim 2^{2l_2} \|\langle g_2, \varphi_{n_1, m}^{1,2} \rangle\|_2^2 \\ &\implies 2^{-l_2} \lesssim \frac{2^{-s_1}}{(\#\mathbb{X}_{(n_1, m)}^{l_2, s_1})^{\frac{1}{2}}} \implies 2^{-l_2} \lesssim \frac{2^{-s_1}}{\|\mathbb{1}_{\mathbb{X}^{l_2, s_1}}\|_{\ell_{n_1, m}^\infty}^{\frac{1}{2}} \ell_{n_1, m}^1}, \end{aligned} \tag{40}$$

by taking the supremum in (n_1, m) . Analogously,

$$2^{-l_2} \lesssim \frac{2^{-s_2}}{\|\mathbb{1}_{\mathbb{X}^{l_2, s_2}}\|_{\ell_{n_2, m}^\infty}^{\frac{1}{2}} \ell_{n_2, m}^1}. \tag{41}$$

Relations (39), (40) and (41) play a major role in the proof. The last major piece is a way of bounding $\#\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k}$ that allows us to exploit transversality and Strichartz along the right directions, as well as the dual function h . We start with the simplest one of them:

$$\#\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k} \lesssim 2^k \sum_{(\vec{n}, m) \in \mathbb{Z}^3} |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| = 2^k |F|. \tag{42}$$

By dropping most of the indicator functions in the definition of $\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k}$ and using Hölder, we obtain

$$\#\mathbb{X}^{\vec{l}, \vec{r}, \vec{s}, k} \leq \sum_{(\vec{n}, m) \in \mathbb{Z}^3} \mathbb{1}_{\mathbb{X}^{l_2, s_1}}(\vec{n}, m) \cdot \mathbb{1}_{\mathbb{B}_1^{r_1} \cap \mathbb{C}_1^{s_1}}(n_1, m) \leq \|\mathbb{1}_{\mathbb{X}^{l_2, s_1}}\|_{\ell_{n_1, m}^\infty} \ell_{n_2}^1 \cdot \|\mathbb{1}_{\mathbb{B}_1^{r_1} \cap \mathbb{C}_1^{s_1}}\|_{\ell_{n_1, m}^1}.$$

The second factor of the inequality above will be bounded by the one-dimensional bilinear theory:

$$\begin{aligned} \#\mathbb{B}_1^{r_1} \cap \mathbb{C}_1^{s_1} &\lesssim 2^{2r_1+2s_1} \sum_{n_1, m \in \mathbb{Z}} \|\langle g_1, \varphi_{n_1, m}^{1,1} \rangle_{x_1}\|_2^2 \cdot \|\langle g_2, \varphi_{n_1, m}^{1,2} \rangle_{x_1}\|_2^2 \\ &= 2^{2r_1+2s_1} \iint \left(\sum_{n_1, m \in \mathbb{Z}} |\langle g_1, \varphi_{n_1, m}^{1,1} \rangle_{x_1}|^2 \cdot |\langle g_2, \varphi_{n_1, m}^{1,2} \rangle_{x_1}|^2 \right) dx_2 d\tilde{x}_2 \\ &= 2^{2r_1+2s_1} \iint \|g_1\|_{L_{x_1}^2}^2 \cdot \|g_2\|_{L_{x_1}^2}^2 dx_2 d\tilde{x}_2 \leq 2^{2r_1+2s_1} \|g_1\|_2^2 \cdot \|g_2\|_2^2, \end{aligned}$$

¹⁶These indicator functions actually bound g_1 and g_2 , but this does not affect the core of the argument.

by Proposition 4.4 since the supports of $\varphi^{1,1}$ and $\varphi^{1,2}$ are disjoint (this is equivalent to transversality in dimension one). This gives us

$$\#\mathbb{X}^{\vec{l},\vec{r},\vec{s},k} \leq \|\mathbb{1}_{\mathbb{X}^{l_2,s_1}}\|_{\ell_{n_1,m}^\infty \ell_{n_2}^1} \cdot 2^{2r_1+2s_1} \cdot |E_1| \cdot |E_2|. \tag{43}$$

Alternatively,

$$\begin{aligned} \#\mathbb{X}^{\vec{l},\vec{r},\vec{s},k} &\leq \sum_{(n_2,m) \in \mathbb{Z}^2} \mathbb{1}_{\mathbb{B}_2^{r_2} \cap \mathbb{C}_2^{s_2}}(n_2, m) \sum_{n_1 \in \mathbb{Z}} \mathbb{1}_{\mathbb{X}^{l_2,s_2}}(\vec{n}, m) \cdot \mathbb{1}_{\mathbb{B}_1^{r_1}}(n_1, m) \\ &\leq \|\mathbb{1}_{\mathbb{X}^{l_2,s_2}}\|_{\ell_{n_2,m}^\infty \ell_{n_1}^1}^{\frac{1}{2}} \cdot \|\mathbb{1}_{\mathbb{B}_1^{r_1}}\|_{\ell_m^\infty \ell_{n_1}^1}^{\frac{1}{2}} \cdot \|\mathbb{1}_{\mathbb{B}_2^{r_2} \cap \mathbb{C}_2^{s_2}}\|_{\ell_{n_2,m}^1}. \end{aligned}$$

We can treat the last two factors appearing in the right-hand side above as follows: For a fixed $m \in \mathbb{Z}$,

$$\sum_{n_1 \in \mathbb{Z}} \mathbb{1}_{\mathbb{B}_1^{r_1}}(n_1, m) \lesssim 2^{2r_1} \sum_{n_1 \in \mathbb{Z}} \|\langle g_1, \varphi_{n_1,m}^{1,1} \rangle_{x_1}\|_2^2 \leq 2^{2r_1} \cdot \|g_1\|_2^2$$

by Bessel (recall that the modulated bumps $\varphi_{n_1,m}^{1,1}$ are almost-orthogonal if n_1 varies and m is fixed), and then we take the supremum in m . As for the other factor, observe that¹⁷

$$\begin{aligned} \#\mathbb{B}_2^{r_2} \cap \mathbb{C}_2^{s_2} &\lesssim 2^{5r_2+s_2} \sum_{n_2,m \in \mathbb{Z}} \|\langle g_1, \varphi_{n_2,m}^{2,1} \rangle_{x_2}\|_2^5 \cdot \|\langle g_2, \varphi_{n_2,m}^{2,2} \rangle_{x_2}\|_2 \\ &\lesssim 2^{5r_2+s_2} \left(\sum_{n_2,m \in \mathbb{Z}} \|\langle g_1, \varphi_{n_2,m}^{2,1} \rangle_{x_2}\|_2^6 \right)^{\frac{5}{6}} \left(\sum_{n_2,m \in \mathbb{Z}} \|\langle g_2, \varphi_{n_2,m}^{2,2} \rangle_{x_2}\|_2^6 \right)^{\frac{1}{6}} \\ &\leq 2^{5r_2+s_2} \|g_1\|_2^5 \cdot \|g_2\|_2 \end{aligned}$$

by Corollary 4.2. These last two estimates give the following bound on $\#\mathbb{X}^{\vec{l},\vec{r},\vec{s},k}$:

$$\#\mathbb{X}^{\vec{l},\vec{r},\vec{s},k} \lesssim \|\mathbb{1}_{\mathbb{X}^{l_2,s_2}}\|_{\ell_{n_2,m}^\infty \ell_{n_1}^1}^{\frac{1}{2}} \cdot 2^{r_1} \cdot |E_1|^{\frac{1}{2}} \cdot 2^{5r_2+s_2} \cdot |E_1|^{\frac{5}{2}} \cdot |E_2|^{\frac{1}{2}}. \tag{44}$$

In what follows, we interpolate between (43), (44) and (42) with weights $\frac{2}{5}^-$, $\frac{1}{5}^-$ and $\frac{2}{5}^+$, respectively. We also take an appropriate of combination between (40) and (41), and use (39):

$$\begin{aligned} |\tilde{\Lambda}_{2,2}(g_1, g_2, h)| &\lesssim \sum_{\vec{r},\vec{s},k} \frac{2^{-r_1} \cdot 2^{-r_2}}{|E_1|^{\frac{1}{2}}} \cdot \frac{2^{-\frac{4}{5}s_1}}{\|\mathbb{1}_{\mathbb{X}^{l_2,s_1}}\|_{\ell_{n_1,m}^\infty \ell_{n_2}^1}^{\frac{2}{5}}} \cdot \frac{2^{-\frac{1}{5}s_2}}{\|\mathbb{1}_{\mathbb{X}^{l_2,s_2}}\|_{\ell_{n_2,m}^\infty \ell_{n_1}^1}^{\frac{1}{10}}} \cdot 2^{-k} \\ &\quad \cdot \left(\|\mathbb{1}_{\mathbb{X}^{l_2,s_1}}\|_{\ell_{n_1,m}^\infty \ell_{n_2}^1} \cdot 2^{2r_1+2s_1} \cdot |E_1| \cdot |E_2| \right)^{\frac{2}{5}^-} \\ &\quad \cdot \left(\|\mathbb{1}_{\mathbb{X}^{l_2,s_2}}\|_{\ell_{n_2,m}^\infty \ell_{n_1}^1}^{\frac{1}{2}} \cdot 2^{r_1} \cdot |E_1|^{\frac{1}{2}} \cdot 2^{5r_2+s_2} \cdot |E_1|^{\frac{5}{2}} \cdot |E_2|^{\frac{1}{2}} \right)^{\frac{1}{5}^-} \cdot (2^k |F|)^{\frac{2}{5}^-} \\ &\lesssim |E_1|^{\frac{1}{2}} \cdot |E_2|^{\frac{1}{2}} \cdot |F|^{\frac{2}{5}^+}, \end{aligned}$$

which is the estimate that we were looking for.¹⁸

¹⁷Here we are also ignoring the fact that we do not prove the endpoint L^2 - L^6 estimate for the model E_1 . It will not compromise this preliminary exposition.

¹⁸This bound on $\tilde{\Lambda}_{2,2}$ is of course informal, which is why we wrote “ \lesssim ”. Observe that we also removed the sum in \vec{l} ; it contributes with a term that depends on ε in the formal argument. Later in the text we will see why we can assume $\vec{r}, \vec{s}, k \geq 0$ in the sum above.

9B. The general argument. Roughly, this is a one-paragraph outline of the proof: we split the sum in (35) into certain level sets, find good upper bounds for how many points (\vec{n}, m) are in each level set using the weak transversality and Strichartz information, and then average all this data appropriately.

First we will prove the bound

$$\| \text{ME}_{k,d}^{\frac{1}{k}}(g) \|_{L^{2(d+k+1)/(d+k-1)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_{\varepsilon} \prod_{l=1}^k |E_{1,l}|^{\frac{1}{2k}} \cdot \prod_{j=2}^k |E_j|^{\frac{1}{2k}} \tag{45}$$

for every $\varepsilon > 0$. As we remarked at the end of Section 4, this is the restricted weak-type bound that will be proved directly; all the other ones that are necessary for multilinear interpolation can be proved in a similar way, as the reader will notice.

We will define several level sets that encode the sizes of many quantities that will play a role in the proof. We start with the ones involving the scalar products in the multilinear form above:

$$\mathbb{A}_j^{l_j} := \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle g_j, \varphi_{\vec{n},m}^j \rangle| \approx 2^{-l_j}\}, \quad 1 \leq j \leq k.$$

The sizes of the $\langle g_j, \varphi_{\vec{n},m} \rangle$ are not the only information that we will need to control. As in the previous subsection, some mixed-norm quantities appear naturally after using Bessel’s inequality along certain directions, and we will need to capture these as well:

$$\begin{aligned} \mathbb{B}_{i,1}^{r_{i,1}} &:= \{(n_i, m) \in \mathbb{Z}^2 : \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_2 \approx 2^{-r_{i,1}}\}, & 1 \leq i \leq k-1, \\ \mathbb{B}_{i,i+1}^{r_{i,i+1}} &:= \{(n_i, m) \in \mathbb{Z}^2 : \|\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{x_i}\|_2 \approx 2^{-r_{i,i+1}}\}, & 1 \leq i \leq k-1, \\ \mathbb{B}_{k,j}^{r_{k,j}} &:= \{(\vec{n}_k, m) \in \mathbb{Z}^{d-k+2} : \|\langle g_j, \varphi_{\vec{n}_k,m}^{k,j} \rangle_{\vec{x}_k}\|_2 \approx 2^{-r_{k,j}}\}, & 1 \leq j \leq k. \end{aligned}$$

Set $\mathbb{B}_{i,j}^{r_{i,j}} := \emptyset$ for any other pair (i, j) not included in the above definitions. Observe that g_1 (the function that has a tensor structure) has k sets \mathbb{B} associated to it: $k-1$ sets $\mathbb{B}_{i,1}^{r_{i,1}}$ and one set $\mathbb{B}_{k,1}^{r_{k,1}}$. The other functions $g_j, j \neq 1$, have only two: one set $\mathbb{B}_{j-1,j}^{r_{j-1,j}}$ and one set $\mathbb{B}_{k,j}^{r_{k,j}}$ for each $1 \leq j \leq k$. The idea behind the sets $\mathbb{B}_{i,1}^{r_{i,1}}$ and $\mathbb{B}_{i,i+1}^{r_{i,i+1}}$ is to isolate the “piece” of each function that encodes the weak transversality information from the part that captures the Strichartz/Tomas–Stein behavior, which is in the set $\mathbb{B}_{k,j}^{r_{k,j}}$. For each $1 \leq i \leq k-1$, we will pair the information of the sets $\mathbb{B}_{i,1}^{r_{i,1}}$ and $\mathbb{B}_{i,i+1}^{r_{i,i+1}}$ and use Proposition 4.4 to extract the gain yielded by weak transversality. The information contained in the sets $\mathbb{B}_{k,j}^{r_{k,j}}$ will be exploited via Corollary 4.2.

The last quantity we have to control is the one arising from the dualizing function h :

$$\mathbb{C}^t := \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-t}\}.$$

In order to prove some crucial bounds, at some point we will have to isolate the previous information for only one of the functions g_j . This will be done in terms of the following set:¹⁹

$$\mathbb{X}^{l_j;r_{i,j}} = \mathbb{A}_j^{l_j} \cap \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_i, m) \in \mathbb{B}_{i,j}^{r_{i,j}}\}.$$

¹⁹Many of these sets are empty since we set $\mathbb{B}_{i,j}^{r_{i,j}} = \emptyset$ for most (i, j) , but only the nonempty ones will appear in the argument.

In other words, $\mathbb{X}^{l_j; r_{i,j}}$ contains all the (n_1, \dots, n_d, m) whose corresponding scalar product $\langle g_j, \varphi_{\vec{n}, m} \rangle$ has size about 2^{-l_j} and with (n_i, m) being such that $\|\langle g_j, \varphi_{n_i, m}^{i,j} \rangle_{x_i}\|_2$ has size about $2^{-r_{i,j}}$.

Finally, it will also be important to encode all the previous information into one single set. This will be done with

$$\mathbb{X}^{\vec{l}, R, t} := \bigcap_{1 \leq j \leq k} \mathbb{A}_j^{l_j} \cap \left\{ (\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_i, m) \in \bigcap_j \mathbb{B}_{i,j}^{r_{i,j}}, 1 \leq i \leq d \right\} \cap \mathbb{C}^t,$$

where we are using the abbreviations $\vec{l} = (l_1, \dots, l_k)$ and $R := (r_{i,j})_{i,j}$. Hence we can bound the form $\tilde{\Lambda}_{k,d}$ as follows:

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \prod_{j=1}^k 2^{-\frac{l_j}{k}} \#\mathbb{X}^{\vec{l}, R, t}. \tag{46}$$

Observe that we are assuming without loss of generality that $l_j, r_{i,j}, t \geq 0$. Indeed,

$$2^{-l_j} \lesssim |\langle g_j, \varphi_{\vec{n}, m}^j \rangle| \leq \|g_j\|_\infty \cdot \|\varphi\|_1 \lesssim 1,$$

so l_j is at least as big as a universal integer. The argument for the remaining indices is the same.

The following two lemmas play a crucial role in the argument by relating the scalar and mixed-norm quantities involved in the stopping-time above. [Lemma 9.4](#) allows us to do that for the quantities associated to g_1 , the function that has a tensor structure. We remark that this is the only place in the proof where the tensor structure is used.

Lemma 9.4. *If $\mathbb{X}^{\vec{l}, R, t} \neq \emptyset$, then*

$$2^{-l_1} \approx \frac{2^{-r_{1,1}} \dots 2^{-r_{k,1}}}{\|g_1\|_2^{k-1}},$$

Proof. Observe that

$$\begin{aligned} 2^{-r_{1,1}} \dots 2^{-r_{k,1}} &\approx \prod_{i=1}^k \|\langle g_1, \varphi_{n_i, m}^{i,1} \rangle_{x_i}\|_2 = \prod_{i=1}^k \|\langle g_{1,1} \otimes \dots \otimes g_{1,k}, \varphi_{n_i, m}^{i,1} \rangle_{x_i}\|_2 \\ &= \prod_{i=1}^k |\langle g_{1,i}, \varphi_{n_i, m}^{i,1} \rangle_{x_i}| \cdot \|g_{1,1} \otimes \dots \otimes \hat{g}_{1,i} \otimes \dots \otimes g_{1,k}\|_2 \\ &= |\langle g_1, \varphi_{\vec{n}, m}^1 \rangle| \cdot \|g_1\|_2^{k-1} \approx 2^{-l_1} \cdot \|g_1\|_2^{k-1}, \end{aligned}$$

and this proves the lemma. □

[Lemma 9.5](#) gives us an alternative way of relating the quantities previously defined for the generic functions g_2, \dots, g_k .

Lemma 9.5. *If $\mathbb{X}^{\vec{l}, R, t} \neq \emptyset$, then*

$$2^{-l_{i+1}} \lesssim \frac{2^{-r_{i,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}; r_{i,i+1}}}\|_{\ell_{n_i, m}^\infty}^{\frac{1}{2}} \ell_{\hat{n}_i}^1}, \tag{47}$$

$$2^{-l_{i+1}} \lesssim \frac{2^{-r_{k,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}; r_{k,i+1}}}\|_{\ell_{n_k, m}^\infty}^{\frac{1}{2}} \ell_{\hat{n}_k}^1} \tag{48}$$

for all $1 \leq i \leq k - 1$.

Proof. Inequality (47) is a consequence of orthogonality: for a fixed (n_i, m) , define

$$\mathbb{X}_{(n_i, m)}^{l_{i+1}; r_{i, i+1}} := \{\hat{n}_i : (\vec{n}, m) \in \mathbb{X}^{l_{i+1}, r_{i, i+1}}\}.$$

This way,

$$\begin{aligned} \#\mathbb{X}_{(n_i, m)}^{l_{i+1}; r_{i, i+1}} &\approx 2^{2l_{i+1}} \sum_{\hat{n}_i \in \mathbb{X}_{(n_i, m)}^{l_{i+1}; r_{i, i+1}}} |\langle g_{i+1}, \varphi_{\vec{n}, m}^{i+1} \rangle|^2 \\ &\leq 2^{2l_{i+1}} \sum_{\hat{n}_i} \left| \int \langle g_{i+1}, \varphi_{\vec{n}_i, m}^{i, i+1} \rangle_{x_i} \cdot e^{-2\pi i m (\sum_{j \neq i} x_j^2)} \cdot \prod_{j \neq i} e^{-2\pi i n_j x_j} \, d\hat{x}_i \right|^2 \\ &\leq 2^{2l_{i+1}} \int \|\langle g_{i+1}, \varphi_{\vec{n}_i, m}^{i, i+1} \rangle_{x_i}\|^2 \, d\hat{x}_i \\ &\approx 2^{2l_{i+1}} \cdot 2^{-2r_{i, i+1}}, \end{aligned}$$

where we used Bessel’s inequality from the second to the third line. The lemma follows by taking the supremum in (n_i, m) . Equation (48) is proven analogously. \square

The following corollary gives a convex combination of the relations in Lemma 9.5 that will be used in the proof.

Corollary 9.6. *For $1 \leq i \leq k - 1$ we have*

$$2^{-l_{i+1}} \lesssim \frac{2^{-\frac{2k}{d+k+1} \cdot r_{i, i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}; r_{i, i+1}}}\|_{\ell_{\vec{n}_i, m}^\infty \ell_{\vec{n}_i}^1}} \cdot \frac{2^{-\frac{(d-k+1)}{(d+k+1)} \cdot r_{k, i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}; r_{k, i+1}}}\|_{\ell_{\vec{n}_k, m}^\infty \ell_{\vec{n}_k}^1}}.$$

Proof. Interpolate between the bounds of Lemma 9.5 with weights

$$\frac{2k}{d+k+1} \quad \text{and} \quad \frac{d-k+1}{d+k+1},$$

respectively. \square

We now concentrate on estimating the right-hand side of (46) by finding good bounds for $\#\mathbb{X}^{\vec{l}, R, t}$. The following bound follows immediately from the disjointness of the supports of $\chi_{\vec{n}} \otimes \chi_m$:

$$\#\mathbb{X}^{\vec{l}, R, t} \lesssim \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \lesssim 2^t |F|. \tag{49}$$

By definition of the set $\mathbb{X}^{\vec{l}, R, t}$,

$$\#\mathbb{X}^{\vec{l}, R, t} \leq \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \mathbb{1}_{\mathbb{A}_j^{l_j}}(\vec{n}, m) \cdot \prod_{i, j, \mathbb{B}_{i, j}^{r_{i, j}} \neq \emptyset} \mathbb{1}_{\mathbb{B}_{i, j}^{r_{i, j}}}(n_i, m). \tag{50}$$

We will manipulate (50) in k different ways: $k - 1$ of them will exploit orthogonality (through the one-dimensional bilinear theory after combining the sets $\mathbb{B}_{i, 1}^{r_{i, 1}}$ and $\mathbb{B}_{i, i+1}^{r_{i, i+1}}$, $1 \leq i \leq k - 1$) and the last one

will reflect Strichartz/Tomas–Stein in an appropriate dimension. The following lemma gives us estimates for the cardinality of $\mathbb{X}^{\vec{l},R,t}$ based on the sizes of some of its slices along canonical directions.²⁰

Lemma 9.7. *The bounds above imply:*

(a) *The orthogonality-type bounds:*²¹

$$\#\mathbb{X}^{\vec{l},R,t} \lesssim \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{\hat{n}_i,m}^\infty \ell_{\hat{n}_i}^1} \cdot 2^{2r_{i,1}+2r_{i,i+1}} \cdot \|g_1\|_2^2 \cdot \|g_{i+1}\|_2^2, \quad 1 \leq i \leq k-1. \tag{51}$$

(b) *The Strichartz-type bound:*

$$\#\mathbb{X}^{\vec{l},R,t} \lesssim \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{k,j}}}\|_{\ell_{\hat{n}_k,m}^{\frac{1}{k}} \ell_{\hat{n}_k}^1} \cdot 2^{\frac{2}{k} \sum_{i=1}^{k-1} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(k-1)}{k}} \cdot 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \cdot \|g_1\|_2^\alpha \cdot \prod_{l=2}^k \|g_l\|_2^\beta, \tag{52}$$

where

$$\begin{aligned} \alpha &= \frac{2(d+k+1)}{k(d-k+1)} + \delta \cdot \frac{(d+k+1)}{k(d-k+3)}, \\ \beta &= \frac{2}{k} + \tilde{\delta} \cdot \frac{(d-k+1)}{k(d-k+3)}, \end{aligned}$$

with $\delta, \tilde{\delta} > 0$ being arbitrarily small parameters to be chosen later.²²

Proof. For each $1 \leq i \leq k-1$ we bound most of the indicator functions in (50) by 1 and obtain

$$\begin{aligned} \#\mathbb{X}^{\vec{l},R,t} &\leq \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \mathbb{1}_{\mathbb{A}_{i+1}^{l_{i+1}}}(\vec{n},m) \cdot \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \cdot \mathbb{1}_{\mathbb{B}_{i,i+1}^{r_{i,i+1}}}(n_i,m) \\ &= \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}(\vec{n},m) \cdot \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}(n_i,m) \\ &= \sum_{n_i,m} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}(n_i,m) \sum_{\hat{n}_i} \mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}(\vec{n},m) \\ &\leq \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{\hat{n}_i,m}^\infty \ell_{\hat{n}_i}^1} \cdot \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}\|_{\ell_{\hat{n}_i,m}^1}. \end{aligned} \tag{53}$$

Transversality is exploited now: the cube Q_1 with $\{e_1, \dots, e_{k-1}\}$ as associated set of directions satisfies (15), which allows us to apply Proposition 4.4 for each $1 \leq i \leq k-1$ since weak transversality is equivalent to transversality in dimension $d = 1$. By definition of the sets $\mathbb{B}_{i,1}^{r_{i,1}}$ and $\mathbb{B}_{i,i+1}^{r_{i,i+1}}$, Fubini and Proposition 4.4

²⁰The reader may associate this idea to certain discrete Loomis–Whitney or Brascamp–Lieb inequalities. While reducing matters to lower-dimensional theory is at the core of our paper, we do not yet have a genuine “Brascamp–Lieb way” of bounding $\#\mathbb{X}^{\vec{l},R,t}$ for which our methods work. For instance, no “slice” of $\mathbb{X}^{\vec{l},R,t}$ given by fixing a few (or all) n_j and summing over m appears in our estimates, which breaks the Loomis–Whitney symmetry.

²¹Weak transversality enters the picture here.

²²One should think of δ and $\tilde{\delta}$ as being “morally zero”. They will be chosen as a function of the initially given $\varepsilon > 0$, and the only reason we introduce them is to make the appropriate up to the endpoint Strichartz exponent appear in (56). The main terms of α and β are also chosen with that in mind.

we have

$$\begin{aligned}
 \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}\|_{\ell_{n_i,m}^1} &\lesssim 2^{2r_{i,1}+2r_{i,i+1}} \sum_{(n_i,m) \in \mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}} \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_2^2 \cdot \|\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}\|_2^2 \\
 &\leq 2^{2r_{i,1}+2r_{i,i+1}} \iint \left(\sum_{(n_i,m) \in \mathbb{Z}^2} |\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}|^2 \cdot |\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}|^2 \right) d\hat{x}_i d\hat{y}_i \\
 &\leq 2^{2r_{i,1}+2r_{i,i+1}} \int \int \|g_1\|_{L_{\hat{x}_i}^2}^2 \cdot \|g_{i+1}\|_{L_{\hat{y}_i}^2}^2 d\hat{x}_i d\hat{y}_i \\
 &= 2^{2r_{i,1}+2r_{i,i+1}} \cdot \|g_1\|_2^2 \cdot \|g_{i+1}\|_2^2.
 \end{aligned}$$

Using this in (53) gives (a). As for (b), bound $\#\mathbb{X}^{\vec{l},R,t}$ as follows:

$$\begin{aligned}
 \#\mathbb{X}^{\vec{l},R,t} &= \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \mathbb{1}_{\mathbb{X}^{\vec{l},R,t}}(\vec{n},m) \\
 &\leq \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=2}^k \mathbb{1}_{\mathbb{X}^{l_j:r_{k,j}}}(\vec{n},m) \prod_{i=1}^{k-1} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \cdot \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{k,l}^{r_{k,l}}}(\vec{n}_k,m) \\
 &= \sum_{\vec{n}_k,m} \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{k,l}^{r_{k,l}}}(\vec{n}_k,m) \sum_{n_1, \dots, n_{k-1}} \prod_{j=2}^k \mathbb{1}_{\mathbb{X}^{l_j:r_{k,j}}}(\vec{n},m) \prod_{i=1}^{k-1} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \\
 &\leq \sum_{\vec{n}_k,m} \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{k,l}^{r_{k,l}}}(\vec{n}_k,m) \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{k,j}}}(\vec{n},m)\|_{\ell_{\vec{n}_k}^1}^{\frac{1}{k}} \cdot \left\| \prod_{i=1}^{k-1} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \right\|_{\ell_{\vec{n}_k}^1}^{\frac{1}{k}} \\
 &\leq \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{k,j}}}\|_{\ell_{\vec{n}_k,m}^\infty}^{\frac{1}{k}} \ell_{\vec{n}_k}^1 \cdot \prod_{i=1}^{k-1} \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}\|_{\ell_m^\infty}^{\frac{1}{k}} \ell_{n_i}^1 \cdot \left\| \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{k,l}^{r_{k,l}}}\right\|_{\ell_{\vec{n}_k,m}^1}, \tag{54}
 \end{aligned}$$

where we used Hölder’s inequality from the third to fourth line. Next, notice that

$$\begin{aligned}
 \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}\|_{\ell_m^\infty} \ell_{n_i}^1 &\lesssim \sup_m 2^{2r_{i,1}} \sum_{n_i} \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_2^2 \\
 &= \sup_m 2^{2r_{i,1}} \int \sum_{n_i} |\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}|^2 d\hat{x}_i \\
 &\lesssim 2^{2r_{i,1}} \cdot \|g_1\|_2^2 \tag{55}
 \end{aligned}$$

by orthogonality. Now let

$$p_{k,1} := \frac{k(d-k+3)}{(d+k+1)}, \quad p_{k,l} := \frac{k(d-k+3)}{(d-k+1)} \quad \text{for all } 2 \leq l \leq k$$

and notice that

$$\sum_{l=1}^k \frac{1}{p_{k,l}} = 1.$$

This way, by definition of $\mathbb{B}_{k,l}^{r_{k,l}}$ and by Hölder’s inequality with these $p_{k,l}$ we have

$$\begin{aligned}
 & \left\| \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{k,l}^{r_{k,l}}} \right\|_{\ell_{\vec{n}_k, m}^1} \\
 & \lesssim 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \sum_{(\vec{n}_k, m)} \left\| \langle g_1, \varphi_{\vec{n}_k, m}^{k,1} \rangle_{\vec{x}_k} \right\|_2^\alpha \cdot \prod_{l=2}^k \left\| \langle g_l, \varphi_{\vec{n}_k, m}^{k,l} \rangle_{\vec{x}_k} \right\|_2^\beta \\
 & \leq 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \left(\sum_{(\vec{n}_k, m)} \left\| \langle g_1, \varphi_{\vec{n}_k, m}^{k,1} \rangle_{\vec{x}_k} \right\|_2^{\alpha \cdot p_{k,1}} \right)^{\frac{1}{p_{k,1}}} \cdot \prod_{l=2}^k \left(\sum_{(\vec{n}_k, m)} \left\| \langle g_l, \varphi_{\vec{n}_k, m}^{k,l} \rangle_{\vec{x}_k} \right\|_2^{\beta \cdot p_{k,l}} \right)^{\frac{1}{p_{k,l}}} \\
 & = 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \left(\sum_{(\vec{n}_k, m)} \left\| \langle g_1, \varphi_{\vec{n}_k, m}^{k,1} \rangle_{\vec{x}_k} \right\|_2^{\frac{2(d-k+3)}{(d-k+1)} + \delta} \right)^{\frac{1}{p_{k,1}}} \cdot \prod_{l=2}^k \left(\sum_{(\vec{n}_k, m)} \left\| \langle g_l, \varphi_{\vec{n}_k, m}^{k,l} \rangle_{\vec{x}_k} \right\|_2^{\frac{2(d-k+3)}{(d-k+1)} + \delta} \right)^{\frac{1}{p_{k,l}}} \\
 & \leq 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \cdot \|g_1\|_2^\alpha \cdot \prod_{l=2}^k \|g_l\|_2^\beta, \tag{56}
 \end{aligned}$$

by the up to the endpoint mixed-norm Strichartz bound in Corollary 4.2.²³ Using (55) and (56) in (54) yields (b). □

Given $\varepsilon > 0$ small,²⁴ we interpolate between $k + 1$ bounds for $\#\mathbb{X}^{\vec{l}, R, t}$ with the following weights.²⁵

$$\begin{cases} \theta_l = \frac{1}{d+k+1} - \frac{\varepsilon}{k}, & 1 \leq l \leq k-1, & \text{for (51),} \\ \theta_k = \frac{(d-k+1)}{2(d+k+1)} - \frac{\varepsilon}{k} & & \text{for (52),} \\ \theta_{k+1} = \left[1 - \frac{(d+k-1)}{2(d+k+1)} \right] + \varepsilon & & \text{for (49),} \end{cases}$$

which leads to

$$\begin{aligned}
 & |\tilde{\Lambda}_{k,d}(g, h)| \\
 & \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times \prod_{j=1}^k 2^{-\frac{l_j}{k}} \times \prod_{l=1}^{k-1} \left(\|\mathbb{1}_{\mathbb{X}^{l_l+1; r_{l,l}+1}}\|_{\ell_{\vec{n}_l, m}^\infty} \ell_{\vec{n}_l}^1 \cdot 2^{2r_{l,1}+2r_{l,l}+1} \cdot \|g_1\|_2^2 \cdot \|g_{l+1}\|_2^2 \right)^{\frac{1}{d+k+1} - \frac{\varepsilon}{k}} \\
 & \quad \times \left(\prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j; r_{k,j}}}\|_{\ell_{\vec{n}_k, m}^\infty}^{\frac{1}{k}} \ell_{\vec{n}_k}^1 \cdot 2^{\frac{2}{k} \sum_{i=1}^{k-1} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(k-1)}{k}} \cdot 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \cdot \|g_1\|_2^\alpha \cdot \prod_{l=2}^k \|g_l\|_2^\beta \right)^{\frac{(d-k+1)}{2(d+k+1)} - \frac{\varepsilon}{k}} \\
 & \quad \times (2^t |F|)^{\left[1 - \frac{(d+k-1)}{2(d+k+1)} \right] + \varepsilon},
 \end{aligned}$$

²³See the footnote related to Corollary 4.2.

²⁴Perhaps it is helpful for the reader to think of ε, δ and $\tilde{\delta}$ as equal to zero to focus on the important parts of the proof. The presence of these parameters here is a mere technicality, except of course for the fact that $\varepsilon > 0$ makes us lose the endpoint in this case.

²⁵Observe that $\sum_{l=1}^{k+1} \theta_l = 1$. These weights are chosen so that the correct powers of the measures $|E_j|$ and $|F|$ appear in (58).

Using Lemma 9.4 and Corollary 9.6 to bound the 2^{-l_j} in the form $\tilde{\Lambda}_{k,d}$ yields

$$\begin{aligned}
 & |\tilde{\Lambda}_{k,d}(g, h)| \\
 & \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times 2^{-\frac{\varepsilon}{k^2} l_1} \times \left(\frac{1}{\|g_1\|_2^{k-1}} \prod_{j=1}^k 2^{-r_{j,1}} \right)^{\frac{1}{k} - \frac{\varepsilon}{k^2}} \\
 & \times \prod_{i=1}^{k-1} 2^{-\frac{\varepsilon}{k^2} l_{i+1}} \times \prod_{i=1}^{k-1} \left[\frac{2^{-\frac{2}{d+k+1} \cdot r_{i,i+1}} \cdot 2^{-\frac{(d-k+1)}{k(d+k+1)} \cdot r_{k,i+1}}}{\|\mathbb{1}_{\times^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i, m}^\infty} \ell_{\hat{n}_i}^1} \cdot \frac{2^{-\frac{(d-k+1)}{2k(d+k+1)} \cdot r_{k,i+1}}}{\|\mathbb{1}_{\times^{l_{i+1}:r_{k,i+1}}}\|_{\ell_{n_k, m}^\infty} \ell_{\hat{n}_k}^1}} \right]^{1 - \frac{\varepsilon}{k}} \\
 & \times \prod_{l=1}^{k-1} \left(\|\mathbb{1}_{\times^{l+1}:r_{l,l+1}}\|_{\ell_{n_l, m}^\infty} \ell_{\hat{n}_l}^1 \cdot 2^{2r_{l,1} + 2r_{l,l+1}} \cdot \|g_1\|_2^2 \cdot \|g_{l+1}\|_2^2 \right)^{\frac{1}{d+k+1} - \frac{\varepsilon}{k}} \\
 & \times \left(\prod_{j=2}^k \|\mathbb{1}_{\times^{l_j:r_{k,j}}}\|_{\ell_{n_k, m}^\infty} \ell_{\hat{n}_k}^1 \cdot 2^{\frac{2}{k} \sum_{i=1}^{k-1} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(k-1)}{k}} \cdot 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}} \cdot \|g_1\|_2^\alpha \cdot \prod_{l=2}^k \|g_l\|_2^\beta \right)^{\frac{(d-k+1)}{2(d+k+1)} - \frac{\varepsilon}{k}} \\
 & \times (2^t |F|)^{[1 - \frac{(d+k-1)}{2(d+k+1)}] + \varepsilon},
 \end{aligned}$$

Developing the expression above,

$$\begin{aligned}
 |\tilde{\Lambda}_{k,d}(g, h)| & \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times 2^{-\frac{\varepsilon}{k^2} l_1} \times \left(\prod_{j=1}^k 2^{-r_{j,1}} \right)^{\frac{1}{k} - \frac{\varepsilon}{k^2}} \cdot \|g_1\|_2^{\frac{(k-1)}{k^2} \varepsilon - \frac{(k-1)}{k}} \\
 & \times \prod_{i=1}^{k-1} 2^{-\frac{\varepsilon}{k^2} l_{i+1}} \times \prod_{i=1}^{k-1} [2^{-\frac{2}{d+k+1} \cdot r_{i,i+1}} \cdot 2^{-\frac{(d-k+1)}{k(d+k+1)} \cdot r_{k,i+1}}]^{1 - \frac{\varepsilon}{k}} \\
 & \times \prod_{i=1}^{k-1} \left[\|\mathbb{1}_{\times^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i, m}^\infty} \ell_{\hat{n}_i}^1 \right]^{\frac{1}{d+k+1} \cdot (\frac{\varepsilon}{k} - 1)} \cdot \left[\|\mathbb{1}_{\times^{l_{i+1}:r_{k,i+1}}}\|_{\ell_{n_k, m}^\infty} \ell_{\hat{n}_k}^1 \right]^{\frac{(d-k+1)}{2k(d+k+1)} \cdot (\frac{\varepsilon}{k} - 1)} \\
 & \times \left[\prod_{l=1}^{k-1} \|\mathbb{1}_{\times^{l+1}:r_{l,l+1}}\|_{\ell_{n_l, m}^\infty} \ell_{\hat{n}_l}^1 \right]^{\frac{1}{d+k+1} - \frac{\varepsilon}{k}} \times \left[\prod_{l=1}^{k-1} (2^{r_{l,1} + r_{l,l+1}})^{\frac{2}{d+k+1} - \frac{2\varepsilon}{k}} \right] \\
 & \times \|g_1\|_2^{\frac{2(k-1)}{d+k+1} - \frac{2(k-1)\varepsilon}{k}} \cdot \prod_{l=1}^{k-1} \|g_{l+1}\|_2^{\frac{2}{d+k+1} - \frac{2\varepsilon}{k}} \\
 & \times \prod_{j=2}^k \|\mathbb{1}_{\times^{l_j:r_{k,j}}}\|_{\ell_{n_k, m}^\infty} \ell_{\hat{n}_k}^1 \cdot \left[\frac{1}{k} \cdot \left(\frac{d-k+1}{2(d+k+1)} - \frac{\varepsilon}{k} \right) \right] \cdot (2^{\frac{2}{k} \sum_{i=1}^{k-1} r_{i,1}} \cdot 2^{\alpha \cdot r_{k,1} + \sum_{l=2}^k \beta \cdot r_{k,l}})^{\frac{(d-k+1)}{2(d+k+1)} - \frac{\varepsilon}{k}} \\
 & \times \|g_1\|_2^{(\frac{2(k-1)}{k} + \alpha) \cdot (\frac{d-k+1}{2(d+k+1)} - \frac{\varepsilon}{k})} \cdot \prod_{l=2}^k \|g_l\|_2^{\beta \cdot (\frac{d-k+1}{2(d+k+1)} - \frac{\varepsilon}{k})} \\
 & \times (2^t |F|)^{[1 - \frac{(d+k-1)}{2(d+k+1)}] + \varepsilon}.
 \end{aligned}$$

At this point we set the values of δ and $\tilde{\delta}$ (as functions of ε) to be such that²⁶

$$\begin{aligned}
 \delta \cdot \left[\frac{(d-k+1)}{k(d-k+3)} - \frac{(d+k+1)\varepsilon}{k^2(d-k+3)} \right] &= \frac{1}{2} \left[-\frac{\varepsilon}{k^2} + \frac{2(d+k+1)\varepsilon}{k^2(d-k+1)} \right], \\
 \tilde{\delta} \cdot \left[\frac{(d-k+1)^2}{2k(d+k+1)(d-k+3)} - \frac{(d-k+1)\varepsilon}{k^2(d-k+3)} \right] &= \frac{1}{2} \left[\frac{2\varepsilon}{k^2} - \frac{(d-k+1)\varepsilon}{k^2(d+k+1)} \right].
 \end{aligned}$$

²⁶We emphasize that these particular choices are just for computational convenience, and we have not developed the expressions because this is exactly how we use them to simplify the previous calculations.

Simplifying (and using the expressions that define α and β in Lemma 9.7),

$$\begin{aligned}
 |\tilde{\Lambda}_{k,d}(g,h)| &\lesssim \left[\sum_{l_1 \geq 0} 2^{-\frac{\varepsilon}{k^2} l_1} \right] \times \left[\prod_{j=1}^{k-1} \left(\sum_{r_{j,1} \geq 0} 2^{-(\frac{2\varepsilon}{k} + \frac{\varepsilon}{k^2}) r_{j,1}} \right) \right] \cdot \left[\sum_{r_{k,1} \geq 0} 2^{-r_{k,1} \left(-\frac{\varepsilon}{2k^2} + \frac{(d+k+1)}{(d-k+1)} \frac{\varepsilon}{k^2} \right)} \right] \\
 &\times \left[\prod_{i=1}^{k-1} \left(\sum_{l_{i+1} \geq 0} 2^{-\frac{\varepsilon}{k^2} l_{i+1}} \right) \right] \times \left[\sum_{t \geq 0} 2^{-t \left(\frac{(d+k-1)}{2(d+k+1)} - \varepsilon \right)} \right] \\
 &\times \left[\prod_{i=1}^{k-1} \left(\sum_{r_{i,i+1} \geq 0} 2^{-\frac{2\varepsilon}{k} \left(1 - \frac{1}{d+k+1} \right) r_{i,i+1}} \right) \right] \times \left[\prod_{i=1}^{k-1} \left(\sum_{r_{k,i+1} \geq 0} 2^{-\frac{\varepsilon}{k^2} \left(1 - \frac{(d-k+1)}{2(d+k+1)} \right) r_{k,i+1}} \right) \right] \\
 &\times \prod_{i=1}^{k-1} \left[\sup_{l_{i+1}, r_{i,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{i,i+1}}} \|_{\ell_{n_i, m}^\infty \ell_{\hat{n}_i}^1}^{-\frac{\varepsilon}{k} \left(1 - \frac{1}{d+k+1} \right)} \cdot \sup_{l_{i+1}, r_{k,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{k,i+1}}} \|_{\ell_{n_k, m}^\infty \ell_{\hat{n}_k}^1}^{-\frac{\varepsilon}{k^2} \left(1 - \frac{(d-k+1)}{2(d+k+1)} \right)} \right] \\
 &\times \|g_1\|_2^{\frac{1}{k} - \frac{4\varepsilon}{k(d-k+1)} + \frac{\varepsilon}{k} - \frac{\varepsilon}{k^2} - 2\varepsilon + \frac{1}{2} \left(-\frac{\varepsilon}{k^2} + \frac{2(d+k+1)}{(d-k+1)} \frac{\varepsilon}{k^2} \right)} \times \prod_{l=2}^k \|g_l\|_2^{\frac{1}{k} - \frac{2\varepsilon}{k} + \frac{\varepsilon}{k^2} \left(\frac{(d-k+1)}{2(d+k+1)} - 1 \right)} \\
 &\times |F|^{[1 - \frac{(d+k-1)}{2(d+k+1)}] + \varepsilon}.
 \end{aligned}$$

Observe that

$$\sum_{l_1 \geq 0} 2^{-\frac{\varepsilon}{k^2} l_1} \lesssim_\varepsilon 2^{-\frac{\varepsilon}{k^2} \tilde{l}_1},$$

where \tilde{l}_1 is the smallest index l_1 such that $\times^{\tilde{l}_1, R, t} \neq \emptyset$. Hence there exists some (\vec{k}, \vec{m}) such that

$$2^{-\tilde{l}_1} \approx |\langle g_1, \varphi_{\vec{k}, \vec{m}}^1 \rangle| \leq |E_1|.$$

Therefore

$$\sum_{l_1 \geq 0} 2^{-\frac{\varepsilon}{k^2} l_1} \lesssim_\varepsilon |E_1|^{\frac{\varepsilon}{k^2}}.$$

Notice also that

$$\sum_{r_{j,1} \geq 0} 2^{-(\frac{2\varepsilon}{k} + \frac{\varepsilon}{k^2}) r_{j,1}} \lesssim_\varepsilon 2^{-(\frac{2\varepsilon}{k} + \frac{\varepsilon}{k^2}) \tilde{r}_{j,1}},$$

where $\tilde{r}_{j,1}$ is defined analogously. We can then find (n_j, m) such that

$$2^{-r_{j,1}} \lesssim \|\langle g_1, \varphi_{n_j, m}^{j,1} \rangle_{x_j}\|_2 \leq |E_1|^{\frac{1}{2}}.$$

Therefore

$$\sum_{r_{j,1} \geq 0} 2^{-(\frac{2\varepsilon}{k} + \frac{\varepsilon}{k^2}) r_{j,1}} \lesssim_\varepsilon |E_1|^{\frac{\varepsilon}{k} + \frac{\varepsilon}{2k^2}}.$$

We can estimate all other sums in the bound above analogously. Observe that since the cardinalities appearing in

$$\prod_{i=1}^{k-1} \left[\sup_{l_{i+1}, r_{i,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{i,i+1}}} \|_{\ell_{n_i, m}^\infty \ell_{\hat{n}_i}^1}^{-\frac{\varepsilon}{k} \left(1 - \frac{1}{d+k+1} \right)} \cdot \sup_{l_{i+1}, r_{k,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{k,i+1}}} \|_{\ell_{n_k, m}^\infty \ell_{\hat{n}_k}^1}^{-\frac{\varepsilon}{k^2} \left(1 - \frac{(d-k+1)}{2(d+k+1)} \right)} \right] \quad (57)$$

are integers, the whole expression (57) is $O(1)$. Using these observations and the fact that $|E_j| < 1$ gives us

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim_\varepsilon |F|^{1 - \frac{(d+k-1)}{2(d+k+1)} + \varepsilon} \cdot \prod_{j=1}^k |E_j|^{\frac{1}{2k}}. \tag{58}$$

To simplify our notation, set $g := (g_{1,1}, g_{1,2}, \dots, g_{1,k-1}, g_{1,k}, g_2, \dots, g_k)$. To rigorously use multilinear interpolation theory, one can run the argument above for the following averaged multilinearized version of $ME_{k,d}$:

$$\widetilde{ME}_{k,d}^{\frac{1}{k}}(g) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \left(\prod_{l=1}^{k-1} |\langle g_{1,l}, \varphi_{\vec{n}_l, m}^{l,1} \rangle| \right)^{\frac{1}{k}} \cdot |\langle g_{1,k}, \varphi_{\vec{n}_k, m}^{k,1} \rangle|^{\frac{1}{k}} \cdot \left(\prod_{j=2}^k |\langle g_j, \varphi_{\vec{n}, m}^j \rangle| \right)^{\frac{1}{k}} (\chi_{\vec{n}} \otimes \chi_m),$$

with associated dual form²⁷

$$\tilde{\Lambda}_{k,d}(g, h) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \left(\prod_{l=1}^{k-1} |\langle g_{1,l}, \varphi_{\vec{n}_l, m}^{l,1} \rangle| \right)^{\frac{1}{k}} \cdot |\langle g_{1,k}, \varphi_{\vec{n}_k, m}^{k,1} \rangle|^{\frac{1}{k}} \left(\prod_{j=2}^k |\langle g_j, \varphi_{\vec{n}, m}^j \rangle| \right)^{\frac{1}{k}} \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle.$$

Hence (58) gives us

$$\|\widetilde{ME}_{k,d}^{\frac{1}{k}}(g)\|_{L^{2(d+k+1)/(d+k-1)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{l=1}^k |E_{1,l}|^{\frac{1}{2k}} \cdot \prod_{j=2}^k |E_j|^{\frac{1}{2k}}, \tag{59}$$

which is (45) for $\widetilde{ME}_{k,d}$. Finally, observe that

$$\begin{aligned} & \|\widetilde{ME}_{k,d}(g)\|_{L^{2(d+k+1)/(k(d+k-1))+\varepsilon}(\mathbb{R}^{d+1})} \\ & \leq \overbrace{\|\widetilde{ME}_{k,d}(g)^{\frac{1}{k}}\|_{L^{2(d+k+1)/(d+k-1)+k\varepsilon}(\mathbb{R}^{d+1})} \cdots \|\widetilde{ME}_{k,d}(g)^{\frac{1}{k}}\|_{L^{2(d+k+1)/(d+k-1)+k\varepsilon}(\mathbb{R}^{d+1})}}^{k \text{ times}} \\ & \lesssim \left[\prod_{l=1}^k |E_{1,l}|^{\frac{1}{2k}} \cdot \prod_{j=2}^k |E_j|^{\frac{1}{2k}} \right]^k = \prod_{l=1}^k |E_{1,l}|^{\frac{1}{2}} \cdot \prod_{j=2}^k |E_j|^{\frac{1}{2}}, \end{aligned} \tag{60}$$

which finishes the proof of the case $2 \leq k \leq d + 1$ by restricted weak-type interpolation.

10. The endpoint estimate of the case $k = d + 1$ of Theorem 1.5

Let $g_1 : Q_1 \rightarrow \mathbb{R}$, $g_j : Q_j \rightarrow \mathbb{R}$ for $2 \leq j \leq d + 1$ be continuous functions. Recall that the multilinear model for $k = d + 1$ is given in Section 2 by

$$ME_{d+1,d}(g_1, \dots, g_{d+1}) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^{d+1} \langle g_j, \varphi_{\vec{n}, m}^j \rangle (\chi_{\vec{n}} \otimes \chi_m),$$

²⁷There is a slight difference between the forms $\tilde{\Lambda}_{k,d}$ and $\tilde{\Lambda}_{k,d}$: the latter is $2(k-1)$ -linear, whereas the former is k -linear. We cannot apply multilinear interpolation theory with inequality (58) directly, because all we proved is that it holds when g_1 is a tensor. In order to correctly place our estimates in the context of multilinear interpolation, we need to consider a form that has the appropriate level of multilinearity, which is $\tilde{\Lambda}_{k,d}$.

where

$$\varphi_{\vec{n},m}^j = \bigotimes_{l=1}^d \varphi_{n_l,m}^{l,j}, \quad \varphi_{n_l,m}^{l,j}(x_l) = \varphi^{l,j}(x_l) e^{2\pi i n_l x_l} e^{2\pi i m x_l^2},$$

and $\varphi^{l,j}(x)$ was defined in Section 2. From now on, we will assume without loss of generality that g_1 is the full tensor. To simplify our notation, set $g := (g_{1,1}, \dots, g_{1,d}, g_2, \dots, g_{d+1})$. Define

$$\widetilde{ME}_{d+1,d}(g) := \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{l=1}^d \langle g_{1,l}, \varphi_{n_l,m}^{l,1} \rangle \prod_{j=2}^{d+1} \langle g_j, \varphi_{\vec{n},m}^j \rangle (\chi_{\vec{n}} \otimes \chi_m).$$

We will show that $\widetilde{ME}_{d+1,d}$ maps

$$\underbrace{L^2([0, 1]) \times \dots \times L^2([0, 1]) \times L^2(Q_2) \times \dots \times L^2(Q_{d+1})}_{2d \text{ times}}$$

to $L^{2/d}$, which implies the endpoint estimate of the case $k = d + 1$ in Theorem 1.5.

Endpoint estimate of the case $k = d + 1$. Notice that we have d factors in the first product and d factors in the second. We will pair them in the following way:

$$\widetilde{ME}_{d+1,d}(g) := \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=2}^{d+1} \langle g_j, \varphi_{\vec{n},m}^j \rangle \cdot \langle g_{1,j-1}, \varphi_{n_{j-1},m}^{1,j-1} \rangle (\chi_{\vec{n}} \otimes \chi_m)$$

Now observe that

$$\begin{aligned} \|\widetilde{ME}_{d+1,d}(g)\|_{\frac{2}{d}} &= \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=2}^{d+1} |\langle g_j \otimes \bar{g}_{1,j-1}, \varphi_{\vec{n},m}^j \otimes \bar{\varphi}_{n_{j-1},m}^{1,1} \rangle|_{\frac{2}{d}} \\ &\leq \prod_{j=2}^{d+1} \left(\sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} |\langle g_j \otimes \bar{g}_{1,j-1}, \varphi_{\vec{n},m}^j \otimes \bar{\varphi}_{n_{j-1},m}^{1,1} \rangle|^2 \right)^{\frac{1}{d}}. \end{aligned} \tag{61}$$

Let us analyze the $j = 2$ scalar product inside the parentheses (the others are dealt with in a similar way):

$$\begin{aligned} &\langle g_2 \otimes \bar{g}_{1,1}, \varphi_{\vec{n},m}^2 \otimes \bar{\varphi}_{n_1,m}^{1,1} \rangle \\ &= \int_{\mathbb{R}^{d-1}} \langle g_{2,1} \otimes \bar{g}_{1,1}, \varphi_{n_1,m}^{1,2} \otimes \bar{\varphi}_{n_1,m}^{1,1} \rangle \left(\prod_{u \geq 2} \varphi^{u,2}(x_u) \right) e^{-2\pi i m (\sum_{l \geq 2} x_l^2)} e^{-2\pi i (\sum_{l \geq 2} n_l x_l)} \widehat{dx}_1 \\ &= \widehat{H}_{n_1,m}(n_2, \dots, n_d), \end{aligned}$$

where

$$H_{n_1,m}(x_2, \dots, x_d) := \langle g_{2,1} \otimes \bar{g}_{1,1}, \varphi_{n_1,m}^{1,2} \otimes \bar{\varphi}_{n_1,m}^{1,1} \rangle \left(\prod_{u \geq 2} \varphi^{u,2}(x_u) \right) e^{-2\pi i m (\sum_{l \geq 2} x_l^2)}.$$

We can then use Plancherel if we sum over n_2, \dots, n_d first:

$$\begin{aligned} &\sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} |\langle g_j \otimes \bar{g}_{1,j-1}, \varphi_{\vec{n},m}^j \otimes \bar{\varphi}_{n_{j-1},m}^{1,1} \rangle|^2 \\ &= \sum_{n_1,m} \sum_{n_2, \dots, n_d} |\widehat{H}_{n_1,m}(n_2, \dots, n_d)|^2 = \sum_{n_1,m} \|H_{n_1,m}\|_2^2 \\ &= \int_{\mathbb{R}^{d-1}} \left(\prod_{u \geq 2} \varphi^{u,2}(x_u) \right) \left(\sum_{n_1,m} |\langle g_2 \otimes \bar{g}_{1,1}, \varphi_{n_1,m}^{1,2} \otimes \bar{\varphi}_{n_1,m}^{1,1} \rangle|^2 \right) \widehat{dx}_1. \end{aligned}$$

By our initial choice of cubes, $\text{supp}(\varphi_{n_1, m}^{1,1}) \cap \text{supp}(\varphi_{n_1, m}^{1,2}) = \emptyset$, so the sum in (n_1, m) is actually $M_{2,1}$ (we are freezing $d - 1$ variables of g_2 in this sum). Our results from Section 7 imply

$$\sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} |\langle g_j \otimes \bar{g}_{1, j-1}, \varphi_{\vec{n}, m}^j \otimes \bar{\varphi}_{n_{j-1}, m}^{1,1} \rangle|^2 = \|g_2 \otimes \bar{g}_{1,1}\|_2^2.$$

Arguing like that for all $2 \leq j \leq d + 1$, (61) gives us

$$\|\widetilde{ME}_{d+1, d}(g)\|_{\frac{2}{d}}^{\frac{2}{d}} \leq \prod_{j=2}^{d+1} \|g_2 \otimes \bar{g}_{1, j-1}\|_2^{\frac{2}{d}} = \prod_{j=1}^{d+1} \|g_j\|_2^{\frac{2}{d}}$$

and the result follows. □

11. Improved k -linear bounds for tensors

In this section we investigate the following question: *can one obtain better bounds than those of Conjecture 1.2 if one is restricted to the class of tensors?*²⁸ The answer depends on the concept of *degree of transversality*. The extra information that the input functions are supported on cubes that have disjoint projections along many directions leads to new transversality conditions, and we can take advantage of it in the full tensor case. This is the content of Theorem 11.2.

Let $\{e_j\}_{1 \leq j \leq d}$ be the canonical basis of \mathbb{R}^d . If $Q \subset \mathbb{R}^d$ is a cube, $\pi_j(Q)$ represents the projection of Q along the e_j direction.

Definition 11.1. Let $\{Q_1, \dots, Q_k\}$ be a collection of k closed unit cubes in \mathbb{R}^d with vertices in \mathbb{Z}^d . We associate to this collection its *transversality vector*

$$\tau = (\tau_1, \dots, \tau_d),$$

where $\tau_j = 1$ if there are at least two distinct intervals among the projections $\pi_j(Q_l)$, $1 \leq l \leq k$, and $\tau_j = 0$ otherwise. The *total degree of transversality* of the collection $\{Q_1, \dots, Q_k\}$ is

$$|\tau| := \sum_{1 \leq l \leq d} \tau_l.$$

The k -linear extension model for a set of cubes $\{Q_l\}_{1 \leq l \leq k}$ as in Definition 11.1 is initially given on $C(Q_1) \times \dots \times C(Q_k)$ by

$$ME_{k, d}^{Q_1, \dots, Q_k}(g_1, \dots, g_k) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \langle g_j, \varphi_{\vec{n}, m}^j \rangle (\chi_{\vec{n}} \otimes \chi_m), \tag{62}$$

where the bumps $\varphi_{\vec{n}, m}^j$ are analogous to the ones in Section 9, but now adapted to the cubes Q_k .

From now on we will assume that g_j is a full tensor $g_j^1 \otimes \dots \otimes g_j^d$ for $1 \leq j \leq k$ and that the transversality vector of the collection $\{Q_1, \dots, Q_k\}$ is τ . To simplify the notation, we will replace the superscripts Q_j in (62) with τ and define

$$g := (g_1^1, \dots, g_1^d, \dots, g_j^1, \dots, g_j^d, \dots, g_k^1, \dots, g_k^d).$$

²⁸Extension estimates beyond the conjectured range have been verified in [Mandel and Oliveira e Silva 2023] for a certain class of functions when the underlying submanifold is \mathbb{S}^{d-1} ; [Shao 2009] also contains results of this kind for the paraboloid.

We are then led to consider

$$ME_{k,d}^\tau(g) := \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \prod_{l=1}^d \langle g_j^l, \varphi_{n_l,m}^{l,j} \rangle (\chi_{\vec{n}} \otimes \chi_m), \tag{63}$$

where

$$\varphi_{n_l,m}^{l,j}(x) = \varphi^{l,j}(x) e^{2\pi i n_l x} e^{2\pi i m x^2}, \quad \text{supp}(\varphi^{l,j}) \subset \pi_l(Q_j).$$

As was the case in Section 9, we will deal first with an averaged version of $ME_{k,d}^\tau$ for technical reasons. Define

$$\widetilde{ME}_{k,d}^\tau(g) := \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \prod_{l=1}^d |\langle g_j^l, \varphi_{n_l,m}^{l,j} \rangle|^{\frac{1}{k}} (\chi_{\vec{n}} \otimes \chi_m), \tag{64}$$

and consider its dual form

$$\widetilde{\Lambda}_{k,d}^\tau(g, h) := \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \prod_{l=1}^d |\langle g_j^l, \varphi_{n_l,m}^{l,j} \rangle|^{\frac{1}{k}} \cdot \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle.$$

Let $E_{j,l}$, $1 \leq j \leq k$ and $1 \leq l \leq d$, be measurable sets such that $|g_j^l| \leq \chi_{E_{j,l}}$. Let $F \subset \mathbb{R}^{d+1}$ be a measurable set such that $|h| \leq \chi_F$. Under these conditions we have the following result:

Theorem 11.2. $ME_{k,d}^\tau$ satisfies

$$\|ME_{k,d}^\tau(g)\|_{L^p(\mathbb{R}^{d+1})} \lesssim_p \prod_{j=1}^k \prod_{l=1}^d \|g_j^l\|_2 \quad \text{for all } p > p_\tau := \frac{2(d + |\tau| + 2)}{k(d + |\tau|)}.$$

Proof. It is enough to prove that

$$\|\widetilde{ME}_{k,d}^\tau(g)\|_{L^p(\mathbb{R}^{d+1})} \lesssim_p \prod_{j=1}^k \prod_{l=1}^d |E_{j,l}|^{\frac{1}{2k}},$$

holds for every

$$p > \frac{2(d + |\tau| + 2)}{(d + |\tau|)}.$$

Define the level sets

$$\begin{aligned} \mathbb{A}_{j,l}^{r_{j,l}} &:= \{(n_l, m) \in \mathbb{Z}^2 : |\langle g_j^l, \varphi_{n_l,m}^{l,j} \rangle| \approx 2^{-r_{j,l}}\}, \\ \mathbb{B}^t &:= \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-t}\}. \end{aligned}$$

Set $R := (r_{i,j})_{i,j}$ and

$$\mathbb{X}^{R,t} := \left\{ (\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_l, m) \in \bigcap_{j=1}^k \mathbb{A}_{j,l}^{r_{j,l}}, \quad 1 \leq l \leq d \right\} \cap \mathbb{B}^t.$$

We then have

$$|\widetilde{\Lambda}_{k,d}^\tau(g, h)| \lesssim \sum_{R,t \geq 0} 2^{-t} \cdot \prod_{j=1}^k \prod_{l=1}^d 2^{-\frac{r_{j,l}}{k}} \cdot \#\mathbb{X}^{R,t}.$$

As in the previous section, we can assume without loss of generality that $r_{j,l}, t \geq 0$. We can estimate $\#\mathbb{X}^{R,t}$ using the function h :

$$\#\mathbb{X}^{R,t} \lesssim 2^t \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} |(h, \chi_{\vec{n}} \otimes \chi_m)| \lesssim 2^t |F|. \tag{65}$$

Alternatively, by the definition of $\mathbb{X}^{R,t}$,

$$\#\mathbb{X}^{R,t} \leq \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \prod_{l=1}^d \mathbb{1}_{\mathbb{A}_{j,l}^{r_{j,l}}}(n_l, m) \tag{66}$$

There are many ways to estimate the right-hand side above. We will obtain d different bounds for it, each one arising from summing in a different order. Fix $1 \leq l \leq d$ and leave the sum over (n_l, m) for last:

$$\begin{aligned} \#\mathbb{X}^{R,t} &= \sum_{(n_l,m) \in \mathbb{Z}^2} \left[\prod_{j=1}^k \mathbb{1}_{\mathbb{A}_{j,l}^{r_{j,l}}}(n_l, m) \right] \cdot \prod_{\tilde{l}=1, \tilde{l} \neq l}^d \left[\sum_{n_{\tilde{l}}} \prod_{\tilde{j}=1}^k \mathbb{1}_{\mathbb{A}_{\tilde{j},\tilde{l}}^{r_{\tilde{j},\tilde{l}}}}(n_{\tilde{l}}, m) \right] \\ &\leq \sum_{(n_l,m) \in \mathbb{Z}^2} \left[\prod_{j=1}^k \mathbb{1}_{\mathbb{A}_{j,l}^{r_{j,l}}}(n_l, m) \right] \cdot \prod_{\tilde{l}=1, \tilde{l} \neq l}^d \prod_{\tilde{j}=1}^k \left[\sum_{n_{\tilde{l}}} \mathbb{1}_{\mathbb{A}_{\tilde{j},\tilde{l}}^{r_{\tilde{j},\tilde{l}}}}(n_{\tilde{l}}, m) \right]^{\gamma_{l,\tilde{j},\tilde{l}}}, \end{aligned} \tag{67}$$

where we used Hölder's inequality in the last line and $\gamma_{l,\tilde{j},\tilde{l}}$ are generic parameters such that

$$\sum_{\tilde{j}=1}^k \gamma_{l,\tilde{j},\tilde{l}} = 1 \tag{68}$$

for all $1 \leq l, \tilde{l} \leq d$ with $l \neq \tilde{l}$ fixed. Let us briefly explain the labels in these parameters that we just introduced:

$$\gamma_{l,\tilde{j},\tilde{l}} \longrightarrow \begin{cases} l \text{ indicates that the last variables to be summed are } (n_l, m), \\ \tilde{j} \text{ corresponds to the } \tilde{j}\text{-th function } g_{\tilde{j}}, \\ \tilde{l} \neq l \text{ corresponds to the } \tilde{l}\text{-th variable } n_{\tilde{l}}. \end{cases}$$

We will not make any specific choice for the $\gamma_{l,\tilde{j},\tilde{l}}$ since condition (68) will suffice. Now observe that for a fixed $m \in \mathbb{Z}$ we have

$$\sum_{n_{\tilde{l}}} \mathbb{1}_{\mathbb{A}_{\tilde{j},\tilde{l}}^{r_{\tilde{j},\tilde{l}}}}(n_{\tilde{l}}, m) \leq 2^{2r_{\tilde{j},\tilde{l}}} \sum_{n_{\tilde{l}}} |(g_{\tilde{j}}^{\tilde{l},\tilde{j}}, \varphi_{n_{\tilde{l}},m}^{\tilde{l},\tilde{j}})|^2 \leq 2^{2r_{\tilde{j},\tilde{l}}} \cdot |E_{\tilde{j},\tilde{l}}| \tag{69}$$

by Bessel's inequality. Using (69) back in (67):

$$\begin{aligned} \#\mathbb{X}^{R,t} &\leq \prod_{\tilde{l}=1, \tilde{l} \neq l}^d \prod_{\tilde{j}=1}^k 2^{2\gamma_{l,\tilde{j},\tilde{l}} r_{\tilde{j},\tilde{l}}} |E_{\tilde{j},\tilde{l}}|^{\gamma_{l,\tilde{j},\tilde{l}}} \cdot \sum_{(n_l,m) \in \mathbb{Z}^2} \left[\prod_{j=1}^k \mathbb{1}_{\mathbb{A}_{j,l}^{r_{j,l}}}(n_l, m) \right], \\ &= \prod_{\tilde{l}=1, \tilde{l} \neq l}^d \prod_{\tilde{j}=1}^k 2^{2\gamma_{l,\tilde{j},\tilde{l}} r_{\tilde{j},\tilde{l}}} |E_{\tilde{j},\tilde{l}}|^{\gamma_{l,\tilde{j},\tilde{l}}} \cdot \sum_{(n_l,m) \in \mathbb{Z}^2} \left[\prod_{(j_1,j_2), j_1 \neq j_2} \mathbb{1}_{\mathbb{A}_{j_1,l}^{r_{j_1,l}}}(n_l, m) \cdot \mathbb{1}_{\mathbb{A}_{j_2,l}^{r_{j_2,l}}}(n_l, m) \right]. \end{aligned} \tag{70}$$

We simply used the fact that $\mathbb{1}^2 = \mathbb{1}$ in the last line above. Our goal is to pair the scalar products in (63) corresponding to the functions $g_{j_1}^l$ and $g_{j_2}^l$. There are two kinds of such pairs:

- (a) A pair (j_1, j_2) with $j_1 \neq j_2$ is l -transversal if $\text{supp}(\varphi^{l,j_1}) \cap \text{supp}(\varphi^{l,j_2}) = \emptyset$.
- (b) A pair (j_1, j_2) with $j_1 \neq j_2$ is non- l -transversal along the direction e_l if $\text{supp}(\varphi^{l,j_1}) \cap \text{supp}(\varphi^{l,j_2}) \neq \emptyset$.

Thus we have by Hölder’s inequality for generic parameters α_{l,j_1,j_2} and β_{l,j_1,j_2} ,

$$\begin{aligned} \#\mathbb{X}^{R,t} &\leq \prod_{\substack{\tilde{l}=1 \\ \tilde{l} \neq l}}^d \prod_{\substack{\tilde{j}=1 \\ \tilde{j} \neq l}}^k 2^{2\gamma_{l,\tilde{j},\tilde{l}} r_{\tilde{j},\tilde{l}}} \cdot |E_{\tilde{j},\tilde{l}}|^{\gamma_{l,\tilde{j},\tilde{l}}} \cdot \prod_{\substack{(j_1,j_2) \\ l\text{-transversal, } j_1 \neq j_2}} \left(\sum_{(n_l,m) \in \mathbb{Z}^2} \mathbb{1}_{\mathbb{A}_{j_1,l}^{r_{j_1,l}}} (n_l, m) \cdot \mathbb{1}_{\mathbb{A}_{j_2,l}^{r_{j_2,l}}} (n_l, m) \right)^{\alpha_{l,j_1,j_2}} \\ &\times \prod_{\substack{(j_1,j_2) \\ \text{non-}l\text{-transversal, } j_1 \neq j_2}} \left(\sum_{(n_l,m) \in \mathbb{Z}^2} \mathbb{1}_{\mathbb{A}_{j_1,l}^{r_{j_1,l}}} (n_l, m) \cdot \mathbb{1}_{\mathbb{A}_{j_2,l}^{r_{j_2,l}}} (n_l, m) \right)^{\beta_{l,j_1,j_2}}. \end{aligned} \tag{71}$$

Define

$$\begin{aligned} \alpha_{l,j_1,j_2} &= 0 \quad \text{if } (j_1, j_2) \text{ is non-}l\text{-transversal,} \\ \beta_{l,j_1,j_2} &= 0 \quad \text{if } (j_1, j_2) \text{ is } l\text{-transversal.} \end{aligned}$$

Hence Hölder’s condition is

$$\sum_{\substack{(j_1,j_2) \\ 1 \leq j_1, j_2 \leq k \\ j_1 \neq j_2}} \alpha_{l,j_1,j_2} + \beta_{l,j_1,j_2} = 2, \tag{72}$$

since we are counting each α_{l,j_1,j_2} and β_{l,j_1,j_2} twice, for all $1 \leq l \leq d$. The labels in the parameters α and β track the following information:

$$\alpha_{l,j_1,j_2} \text{ and } \beta_{l,j_1,j_2} \longrightarrow \begin{cases} l \text{ indicates that we are summing over } (n_l, m), \\ j_1 \text{ and } j_2 \text{ correspond to two distinct functions } g_{j_1} \text{ and } g_{j_2}. \end{cases}$$

We can then use Proposition 4.4 for the transversal pairs and a combination of one-dimensional Strichartz/Tomas–Stein with Hölder for the nontransversal ones:

$$\begin{aligned} \#\mathbb{X}^{R,t} &\leq \prod_{\substack{\tilde{l}=1 \\ \tilde{l} \neq l}}^d \prod_{\substack{\tilde{j}=1 \\ \tilde{j} \neq l}}^k 2^{2\gamma_{l,\tilde{j},\tilde{l}} r_{\tilde{j},\tilde{l}}} \cdot |E_{\tilde{j},\tilde{l}}|^{\gamma_{l,\tilde{j},\tilde{l}}} \cdot \prod_{\substack{(j_1,j_2) \\ l\text{-transversal, } j_1 \neq j_2}} (2^{2\alpha_{l,j_1,j_2}(r_{j_1,l}+r_{j_2,l})} \cdot |E_{j_1,l}|^{\alpha_{l,j_1,j_2}} \cdot |E_{j_2,l}|^{\alpha_{l,j_1,j_2}}) \\ &\times \prod_{\substack{(j_1,j_2) \\ \text{non-}l\text{-transversal, } j_1 \neq j_2}} (2^{3\beta_{l,j_1,j_2}(r_{j_1,l}+r_{j_2,l})} \cdot |E_{j_1,l}|^{\frac{3}{2}\beta_{l,j_1,j_2}} \cdot |E_{j_2,l}|^{\frac{3}{2}\beta_{l,j_1,j_2}}). \end{aligned} \tag{73}$$

As mentioned earlier in this section, we have d estimates like (73). We will interpolate between them with weights θ_l :

$$\#\mathbb{X}^{R,t} = \prod_{l=1}^d (\#\mathbb{X}^{R,t})^{\theta_l},$$

with

$$\sum_{l=1}^d \theta_l = 1. \tag{74}$$

This yields

$$\#\mathbb{X}^{R,t} \lesssim \prod_{j=1}^k \prod_{l=1}^d 2^{\#j,l r_{j,l}} \cdot |E_{j,l}|^{\frac{\#j,l}{2}}, \tag{75}$$

where

$$\#_{j,l} = \left[\sum_{j_1 \neq j} (2\alpha_{l,j,j_1} + 3\beta_{l,j,j_1}) \right] \cdot \theta_l + \sum_{\tilde{l} \neq l} 2\gamma_{\tilde{l},j,l} \cdot \theta_{\tilde{l}}.$$

In order to prove an estimate like $L^2 \times \dots \times L^2 \mapsto L^p$, we will need all these coefficients $\#_{j,l}$ to be equal. Let us call them all X for now and sum over j :

$$\sum_{j=1}^k X = \left[\sum_{j=1}^k \sum_{j_1 \neq j} (2\alpha_{l,j,j_1} + 3\beta_{l,j,j_1}) \right] \cdot \theta_l + \sum_{\tilde{l} \neq l} 2 \left[\sum_{j=1}^k \gamma_{\tilde{l},j,l} \right] \cdot \theta_{\tilde{l}}$$

By (68) and (72)

$$X = \frac{1}{k} \left[6 - \sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{l,j,j_1} \right] \cdot \theta_l + \sum_{\tilde{l} \neq l} \frac{2}{k} \cdot \theta_{\tilde{l}} \tag{76}$$

for all $1 \leq l \leq d$. Together with (74), (76) gives us a linear system of d equations in the d variables $\theta_1, \dots, \theta_d$. The solution is

$$\theta_l = \left[\sum_{\tilde{l}=1}^d \frac{4 - \sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{l,j,j_1}}{4 - \sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{\tilde{l},j,j_1}} \right]^{-1}. \tag{77}$$

Plugging (77) back in (76) gives us

$$X = \frac{2}{k} \left[1 + \left(\sum_{\tilde{l}=1}^d \frac{1}{[4 - \sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{\tilde{l},j,j_1}]} \right)^{-1} \right]. \tag{78}$$

To minimize X we must maximize

$$\sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{\tilde{l},j,j_1}.$$

This is achieved by choosing $\beta_{l,j_1,j_2} = 0$ for all (j_1, j_2) if there is at least one l -transversal pair (j_1, j_2) . In other words, choose

$$\beta_{l,j_1,j_2} = 0 \quad \text{for all } (j_1, j_2) \text{ if } \tau_l = 1.$$

Hence by (72),

$$\sum_{j=1}^k \sum_{j_1 \neq j} \alpha_{\tilde{l},j,j_1} = \begin{cases} 2 & \text{if } \tau_{\tilde{l}} = 1, \\ 0 & \text{if } \tau_{\tilde{l}} = 0. \end{cases}$$

This choice of parameters gives us

$$X = \frac{2(d + |\tau| + 2)}{k(d + |\tau|)},$$

which implies the following estimate for $\#\mathbb{X}^{R,t}$:

$$\#\mathbb{X}^{R,t} \lesssim \prod_{j=1}^k \prod_{l=1}^d 2^{X \cdot r_{j,l}} \cdot |E_{j,l}|^{\frac{X}{2}}, \tag{79}$$

Finally, we interpolate between (79) with weight $\frac{1}{k \cdot X} - \varepsilon$ and (65) with weight $(1 - \frac{1}{k \cdot X}) + \varepsilon$ to bound the form $\Lambda_{k,d}^\tau$:

$$|\Lambda_{k,d}^\tau(g, h)| \lesssim \sum_{R,t \geq 0} 2^{-t} \cdot \prod_{j=1}^k \prod_{l=1}^d 2^{-\frac{r_{j,l}}{k}} \times \left[\prod_{j=1}^k \prod_{l=1}^d 2^{X \cdot r_{j,l}} \cdot |E_{j,l}|^{\frac{X}{2}} \right]^{\frac{1}{k \cdot X} - \varepsilon} \cdot [2^t |F|]^{(1 - \frac{1}{k \cdot X}) + \varepsilon}.$$

Developing the right-hand side:

$$|\Lambda_{k,d}^\tau(g, h)| \lesssim \left(\sum_{t \geq 0} 2^{-(\frac{1}{k \cdot X} - \varepsilon)t} \right) \prod_{j=1}^k \prod_{l=1}^d \left(\sum_{r_{j,l} \geq 0} 2^{-\varepsilon X \cdot r_{j,l}} \right) \times \left[\prod_{j=1}^k \prod_{l=1}^d |E_{j,l}|^{\frac{1}{2k} - \frac{\varepsilon X}{2}} \right] \cdot |F|^{(1 - \frac{1}{k \cdot X}) + \varepsilon}.$$

As in the previous section, these series are summable. We have

$$\sum_{r_{j,l} \geq 0} 2^{-\varepsilon X \cdot r_{j,l}} \lesssim_\varepsilon |E_{j,l}|^{\varepsilon X}.$$

For the series in t we can just bound it by an absolute constant depending on ε . This leads to

$$|\Lambda_{k,d}^\tau(g, h)| \lesssim_\varepsilon \left[\prod_{j=1}^k \prod_{l=1}^d |E_{j,l}|^{\frac{1}{2k} + \frac{\varepsilon X}{2}} \right] \cdot |F|^{(1 - \frac{1}{k \cdot X}) + \varepsilon} \lesssim \left[\prod_{j=1}^k \prod_{l=1}^d |E_{j,l}|^{\frac{1}{2k}} \right] \cdot |F|^{(1 - \frac{1}{k \cdot X}) + \varepsilon},$$

since $|E_{j,l}| \leq 1$, which finishes the proof by multilinear interpolation. □

Remark 11.3. If $\tau_l = 0$ for $1 \leq l \leq d$, then

$$p_\tau = \frac{2(d + 2)}{kd},$$

which could have been proven in general with Hölder and Strichartz/Tomas–Stein. This is because there is no transversality to exploit; therefore the best bounds we can hope for in the multilinear setting come from the linear one.

Remark 11.4. If there are exactly $k - 1$ indices l such that $\tau_l = 1$, then

$$p_\tau = \frac{2(d + k + 1)}{k(d + k - 1)},$$

which is consistent with Theorem 1.5.

Remark 11.5. Finally, if one has more than $k - 1$ indices l such that $\tau_l = 1$, then

$$p_\tau < \frac{2(d + k + 1)}{k(d + k - 1)},$$

which clearly illustrates the point of this section. The extreme case is when $\tau_l = 1$ for $1 \leq l \leq d$, which gives

$$p_\tau = \frac{2(d + 1)}{kd}.$$

This can be seen as an improvement upon the linear extension conjecture itself in the following sense: if we take the product of k extensions $E_{U_j} g_j$, $1 \leq j \leq k$, and combine the linear extension conjecture with Hölder’s inequality, we obtain an operator that maps $L^{2(d+1)/d} \times \dots \times L^{2(d+1)/d}$ to $L^{2(d+1)/(kd) + \varepsilon}$.

On the other hand, if we are in a situation in which we have as much transversality as possible and all g_j are full tensors, we obtain $L^2 \times \dots \times L^2$ to $L^{2(d+1)/(kd)+\varepsilon}$.

12. Beyond the L^2 -based k -linear theory with and without transversality

Given a collection $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ of cubes, the purpose of this section is to investigate *near-restriction* k -linear estimates associated to \mathcal{Q} . In other words, we study bounds of the form

$$\left\| \prod_{j=1}^k \mathcal{E}_{Q_j} g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^p(Q_j)} \tag{80}$$

for all $\varepsilon > 0$ and for some $p > 1$. There are two cases of interest here:

- \mathcal{Q} is a collection of transversal cubes.
- All cubes in \mathcal{Q} are the same.

It will be clear that all cases in between these two can be studied in the same framework that we now present.

12A. Near-restriction estimates with transversality. We start by restating (4). For $2 \leq k < d + 1$, to recover the whole range of the generalized k -linear extension conjecture, it is enough to prove [Conjecture 1.2](#) and

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^{2(d+1)/d}(U_j)} \tag{81}$$

for all $\varepsilon > 0$.

Let $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ be our initially fixed set of cubes.²⁹ In what follows, we recast the statement of [Theorem 1.13](#) in terms of this set:

Theorem 12.1. *If \mathcal{Q} is a collection of transversal cubes and g_1 is a tensor, the operator $\mathcal{M}\mathcal{E}_{k,d}(g_1, \dots, g_k)$ satisfies*

$$\|\mathcal{M}\mathcal{E}_{k,d}(g_1, \dots, g_k)\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_\varepsilon \prod_{j=1}^k \|g_j\|_{L^{p(k,d)}(Q_j)}, \tag{82}$$

where

$$p(k, d) = \begin{cases} \frac{4(d+1)}{d+k+1} & \text{if } 2 \leq k < \frac{d}{2}, \\ \frac{4(d+1)}{2d-k+1} & \text{if } \frac{d}{2} \leq k < d+1. \end{cases}$$

As anticipated in the [Introduction](#), we prove it by adapting the argument from [Section 9](#).

Remark 12.2. As in [Section 9](#), the theorem above holds under the assumption that the given set of cubes is weakly transversal and any other $g_j, j \neq 1$, can be assumed to be the tensor.

²⁹See [Section 3](#).

Remark 12.3. Roughly speaking, the difference between the proof of [Theorem 12.1](#) and the one done in [Section 9](#) is in the building blocks we use: instead of Strichartz/Tomas–Stein (in the form of [Corollary 4.2](#)), we will use the best extension bound for the parabola (in the form of [Proposition 4.3](#)). One can think of the argument in this section as a rigorous way of replacing the former piece by the latter in our machinery.

Proof of Theorem 12.1. We work in the same setting as in [Section 9](#). Even though there are some slight differences between the level sets from that section and the ones that we will define here, the approach is very similar.

It is convenient to recall a few important points from [Section 9](#):

- The form of interest here is (in its averaged form):

$$\tilde{\Lambda}_{k,d}(g, h) := \sum_{(\vec{n}, m) \in \mathbb{Z}^{d+1}} \left(\prod_{i=1}^k |\langle g_j, \varphi_{\vec{n}, m}^j \rangle| \right)^{\frac{1}{k}} \langle h, \chi_{\vec{n}} \otimes \chi_m \rangle. \tag{83}$$

- The tensor g_1 has the structure $g_1 = g_{1,1} \otimes \dots \otimes g_{1,d}$.
- $E_{1,1}, \dots, E_{1,d} \subset [0, 1]$, $E_j \subset Q_j$ ($2 \leq j \leq k$) and $F \subset \mathbb{R}^{d+1}$ are measurable sets such that $|g_{1,l}| \leq \chi_{E_{1,l}}$ for $1 \leq l \leq d$, $|g_j| \leq \chi_{E_j}$ for $2 \leq j \leq k$ and $|h| \leq \chi_F$. Furthermore, $E_1 := E_{1,1} \times \dots \times E_{1,d}$.

We start by encoding the sizes of the scalar products appearing in [\(83\)](#):

$$\mathbb{A}_j^{l_j} := \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle g_j, \varphi_{\vec{n}, m} \rangle| \approx 2^{-l_j}\}, \quad 1 \leq j \leq k.$$

Now we see the first difference between the argument in this section and the one in [Section 9](#): the mixed-norm quantities here are all of the same kind, in the sense that the inner products inside the L^2 norms are all one-dimensional:

$$\mathbb{B}_{i,j}^{r_{i,j}} := \{(n_i, m) \in \mathbb{Z}^2 : \|\langle g_j, \varphi_{n_i, m}^{i,j} \rangle_{x_i}\|_2 \approx 2^{-r_{i,j}}\}, \quad 1 \leq i \leq d, \quad 1 \leq j \leq k,$$

The remaining sets are defined just as in [Section 9](#), and with the exact same purpose:

$$\begin{aligned} \mathbb{C}^t &:= \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \approx 2^{-t}\}, \\ \mathbb{X}_j^{l_j; r_{i,j}} &= \mathbb{A}_j^{l_j} \cap \{(\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_i, m) \in \mathbb{B}_{i,j}^{r_{i,j}}\}, \\ \mathbb{X}^{\vec{l}, R, t} &:= \bigcap_{1 \leq j \leq k} \mathbb{A}_j^{l_j} \cap \left\{ (\vec{n}, m) \in \mathbb{Z}^{d+1} : (n_i, m) \in \bigcap_{1 \leq i \leq d} \mathbb{B}_{i,j}^{r_{i,j}}, 1 \leq i \leq d \right\} \cap \mathbb{C}^t, \end{aligned}$$

where we are using the abbreviations $\vec{l} = (l_1, \dots, l_k)$ and $R := (r_{i,j})_{i,j}$. Hence,

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim \sum_{\vec{l}, R, t} 2^{-t} \prod_{j=1}^k 2^{-\frac{l_j}{k}} \#\mathbb{X}^{\vec{l}, R, t}.$$

The analogue of [Lemma 9.4](#) is the bound

$$2^{-l_1} \approx \frac{2^{-r_{1,1}} \dots 2^{-r_{d,1}}}{\|g_1\|_2^{d-1}}, \tag{84}$$

which is proven in the same way. By an argument entirely analogous to that of [Lemma 9.5](#), we can show

$$2^{-l_j} \lesssim \frac{2^{-r_{i,j}}}{\|\mathbb{1}_{\mathbb{X}^{l_j:r_{i,j}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1}^{\frac{1}{2}}} \quad \text{for all } 1 \leq i \leq d, 2 \leq j \leq k. \tag{85}$$

The following corollary of the estimates above will give us the appropriate convex combination of such relations.³⁰

Corollary 12.4. *For $1 \leq i \leq k - 1$ we have*

$$2^{-l_{i+1}} \lesssim \frac{2^{-\frac{k}{d+1} \cdot r_{i,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1}^{\frac{1}{2(d+1)}}} \cdot \prod_{u=k}^d \frac{2^{-\frac{1}{d+1} \cdot r_{u,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{u,i+1}}}\|_{\ell_{n_u,m}^\infty \ell_{\hat{n}_u}^1}^{\frac{1}{2k(d+1)}}}.$$

Proof. Interpolate between the bounds in [\(85\)](#) with one weight equal to $\frac{k}{d+1}$ for $(i, j) := (i, i + 1)$ and $d - k + 1$ weights $\frac{1}{d+1}$ for $(i, j) := (u, i + 1)$, $k \leq u \leq d$. □

We can estimate $\#\mathbb{X}^{\vec{l},R,t}$ using the function h :

$$\#\mathbb{X}^{\vec{l},R,t} \lesssim \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} |\langle h, \chi_{\vec{n}} \otimes \chi_m \rangle| \lesssim 2^t |F|. \tag{86}$$

Alternatively,

$$\#\mathbb{X}^{\vec{l},R,t} \leq \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=1}^k \mathbb{1}_{\mathbb{A}_j^{l_j}}(\vec{n}, m) \prod_{i=1}^d \prod_{j=1}^k \mathbb{1}_{\mathbb{B}_{i,j}^{r_{i,j}}}(n_i, m). \tag{87}$$

Similarly to what was done in [Section 9](#), we will manipulate the inequality above in d ways: $k - 1$ of them will exploit orthogonality (from the combination of the sets $\mathbb{B}_{i,1}^{r_{i,1}}$ and $\mathbb{B}_{i,i+1}^{r_{i,i+1}}$, $1 \leq i \leq k - 1$), but now the other $d - k + 1$ ones will reflect the linear extension problem in dimension 1. The following lemma is the appropriate analogue of [Lemma 9.7](#) in this section:

Lemma 12.5. *The bounds above imply:*

(a) *The orthogonality-type bounds: for all $1 \leq i \leq k - 1$,*

$$\#\mathbb{X}^{\vec{l},R,t} \lesssim \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1} \cdot 2^{2r_{i,1} + 2r_{i,i+1}} \cdot \|g_1\|_2^2 \cdot \|g_{i+1}\|_2^2. \tag{88}$$

(b) *The extension-type bounds: for all $k \leq u \leq d$,*

$$\begin{aligned} \#\mathbb{X}^{\vec{l},R,t} &\lesssim \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}\|_{\ell_{n_u,m}^\infty \ell_{\hat{n}_u}^1}^{\frac{1}{k}} \cdot 2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(d-1)}{k}} \\ &\quad \times 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \cdot \left(\prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \cdot \|g_{1,u}\|_4^\alpha \cdot \prod_{l=2}^k \|g_l\|_4^\beta, \end{aligned} \tag{89}$$

where

$$\alpha := \frac{2(k+1)}{k} + \delta \cdot \frac{(k+1)}{2k}, \quad \beta := \frac{2}{k} + \tilde{\delta} \cdot \frac{1}{2k},$$

with $\delta, \tilde{\delta} > 0$ being arbitrarily small parameters to be chosen later.

³⁰Notice that instead of using just two mixed quantities for each scalar one (as in [Corollary 9.6](#)), we are using $d - k + 2$ many of them here.

Proof. Part (a) is the same as in Lemma 9.7(a). As for (b), fix $k \leq u \leq d$ and bound $\#\mathbb{X}^{\vec{l},R,t}$ as follows:

$$\begin{aligned}
 \#\mathbb{X}^{\vec{l},R,t} &= \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \mathbb{1}_{\mathbb{X}^{\vec{l},R,t}}(\vec{n},m) \\
 &\leq \sum_{(\vec{n},m) \in \mathbb{Z}^{d+1}} \prod_{j=2}^k \mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}(\vec{n},m) \prod_{i \neq u} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \cdot \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u,l}^{r_{u,l}}}(n_u,m) \\
 &= \sum_{n_u,m} \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u,l}^{r_{u,l}}}(n_u,m) \sum_{\hat{n}_u} \prod_{j=2}^k \mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}(\vec{n},m) \prod_{i \neq u} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \\
 &\leq \sum_{n_u,m} \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u,l}^{r_{u,l}}}(n_u,m) \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}(\vec{n},m)\|_{\ell_{\hat{n}_u}^1}^{\frac{1}{k}} \cdot \left\| \prod_{i \neq u} \mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}(n_i,m) \right\|_{\ell_{\hat{n}_u}^1}^{\frac{1}{k}} \\
 &\leq \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}\|_{\ell_{n_u,m}^\infty \ell_{\hat{n}_u}^1}^{\frac{1}{k}} \cdot \prod_{i \neq u} \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}\|_{\ell_m^\infty \ell_{n_i}^1}^{\frac{1}{k}} \cdot \left\| \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u,l}^{r_{u,l}}}\right\|_{\ell_{n_u,m}^1}, \tag{90}
 \end{aligned}$$

where we used Hölder’s inequality from the third to fourth line. Next, notice that

$$\begin{aligned}
 \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}}}\|_{\ell_m^\infty \ell_{n_i}^1} &\lesssim \sup_m 2^{2r_{i,1}} \sum_{n_i} \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_2^2 \\
 &= \sup_m 2^{2r_{i,1}} \int \sum_{n_i} |\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}|^2 d\hat{x}_i \lesssim 2^{2r_{i,1}} \cdot \|g_1\|_2^2
 \end{aligned} \tag{91}$$

by orthogonality. Now let

$$\begin{aligned}
 p_{u,1} &:= \frac{2k}{(k+1)}, \\
 p_{u,l} &:= 2k \quad \text{for all } 2 \leq l \leq k
 \end{aligned}$$

and notice that

$$\sum_{l=1}^k \frac{1}{p_{u,l}} = 1.$$

This way, by the definition of $\mathbb{B}_{u,l}^{r_{u,l}}$ and by Hölder’s inequality with these $p_{u,l}$ we have

$$\begin{aligned}
 &\left\| \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u,l}^{r_{u,l}}}\right\|_{\ell_{n_u,m}^1} \\
 &\lesssim 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \sum_{(n_u,m)} \|\langle g_1, \varphi_{n_u,m}^{u,1} \rangle_{x_u}\|_2^\alpha \cdot \prod_{l=2}^k \|\langle g_l, \varphi_{n_u,m}^{u,l} \rangle_{x_u}\|_2^\beta \\
 &\leq 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \left(\sum_{(n_u,m)} \|\langle g_1, \varphi_{n_u,m}^{u,1} \rangle_{x_u}\|_2^{\alpha \cdot p_{u,1}} \right)^{\frac{1}{p_{u,1}}} \cdot \prod_{l=2}^k \left(\sum_{(n_u,m)} \|\langle g_l, \varphi_{n_u,m}^{u,l} \rangle_{x_u}\|_2^{\beta \cdot p_{u,l}} \right)^{\frac{1}{p_{u,l}}} \\
 &= 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \left(\sum_{(n_u,m)} \|\langle g_1, \varphi_{n_u,m}^{u,1} \rangle_{x_u}\|_2^{4+\delta} \right)^{\frac{1}{p_{u,1}}} \cdot \prod_{l=2}^k \left(\sum_{(n_u,m)} \|\langle g_l, \varphi_{n_u,m}^{u,l} \rangle_{x_u}\|_2^{4+\delta} \right)^{\frac{1}{p_{u,l}}}. \tag{92}
 \end{aligned}$$

At this point we see another difference between this proof and the argument in Section 9: We do not obtain a pure L^p norm when using the near- L^4 extension analogue of Corollary 4.2 for $l = d - 1$. Alternatively, we use Hölder in the term involving g_1 once more:

$$\begin{aligned} \|\langle g_1, \varphi_{n_u, m}^{u, 1} \rangle_{x_u}\|_2^{4+\delta} &= \left[\int \left(\prod_{j \neq u} |g_{1, j}|^2(x_j) \right) \cdot |\langle g_{1, u}, \varphi_{n_u, m}^{u, 1} \rangle_{x_u}|^2 \widehat{d}x_u \right]^{\frac{4+\delta}{2}} \\ &\leq \left(\prod_{j \neq u} \|g_{1, j}\|_2 \right)^{4+\delta} \cdot |\langle g_{1, u}, \varphi_{n_u, m}^{u, 1} \rangle_{x_u}|^{4+\delta}. \end{aligned}$$

For the remaining g_l we simply use Hölder and the fact that they are compactly supported:³¹

$$\|\langle g_l, \varphi_{n_u, m}^{u, l} \rangle_{x_u}\|_2^{4+\tilde{\delta}} \lesssim \|\langle g_l, \varphi_{n_u, m}^{u, l} \rangle_{x_u}\|_4^{4+\tilde{\delta}}.$$

These observations imply

$$\begin{aligned} \left\| \prod_{l=1}^k \mathbb{1}_{\mathbb{B}_{u, l}^{r_{u, l}}} \right\|_{\ell_{n_u, m}^1} &\lesssim 2^{\alpha \cdot r_{u, 1} + \sum_{l=2}^k \beta \cdot r_{u, l}} \cdot \left(\prod_{j \neq u} \|g_{1, j}\|_2 \right)^{\frac{4+\delta}{p_{u, 1}}} \cdot \left(\sum_{(n_u, m)} |\langle g_{1, u}, \varphi_{n_u, m}^{u, 1} \rangle_{x_u}|^{4+\delta} \right)^{\frac{1}{p_{u, 1}}} \\ &\quad \cdot \prod_{l=2}^k \left(\sum_{(n_u, m)} \|\langle g_l, \varphi_{n_u, m}^{u, l} \rangle_{x_u}\|_4^{4+\tilde{\delta}} \right)^{\frac{1}{p_{u, l}}} \\ &\leq 2^{\alpha \cdot r_{u, 1} + \sum_{l=2}^k \beta \cdot r_{u, l}} \cdot \left(\prod_{j \neq u} \|g_{1, j}\|_2 \right)^\alpha \cdot \|g_{1, u}\|_4^\alpha \cdot \prod_{l=2}^k \|g_l\|_4^\beta, \end{aligned} \tag{93}$$

where we used Minkowski for norms and the L^4 - $L^{4+\tilde{\delta}}$ one-dimensional extension estimate from the second to third line above. Part (b) follows from applying (91) and (93) to (90). \square

Given $\varepsilon > 0$, we bound the multilinear form $\tilde{\Lambda}_{k, d}$ using the estimates from (84) and Corollary 12.4 (with the appropriate ε -losses for later convenience), and the ones from Lemma 12.5 with the following weights:

$$\begin{cases} \theta_l = \frac{1}{2(d+1)} - \frac{\varepsilon}{d}, & 1 \leq l \leq d \quad \text{for the } d \text{ estimates in (88) and (89),} \\ \theta_{d+1} = 1 - \frac{d}{2(d+1)} + \varepsilon & \text{for (86).} \end{cases}$$

Hence,

$$\begin{aligned} |\tilde{\Lambda}_{k, d}(g, h)| &\lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times 2^{-\frac{(d+1)\varepsilon}{2kd}} l_1 \times \left(\frac{1}{\|g_1\|_2^{d-1}} \prod_{j=1}^d 2^{-r_{j, 1}} \right)^{\frac{1}{k} - \frac{(d+1)\varepsilon}{2kd}} \\ &\quad \times \prod_{i=1}^{k-1} 2^{-\frac{(d+1)\varepsilon}{2kd} l_{i+1}} \times \prod_{i=1}^{k-1} \left[\frac{2^{-\frac{1}{d+1} \cdot r_{i, i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}: r_{i, i+1}}}\|_{\ell_{\hat{n}_i, m}^\infty} \ell_{\hat{n}_i}^1} \cdot \prod_{u=k}^d \frac{2^{-\frac{1}{k(d+1)} \cdot r_{u, i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}: r_{u, i+1}}}\|_{\ell_{n_u, m}^\infty} \ell_{\hat{n}_u}^1} \right]^{1 - \frac{(d+1)\varepsilon}{2d}} \end{aligned}$$

³¹We use this crude estimate for the remaining g_l because they do not have the same structure that allows “pulling out” the one-dimensional functions $g_{1, j}$, like g_1 does. There is a clear loss here and it is reflected in the fact that $p(k, d)$ is not the best exponent for which (82) holds.

$$\begin{aligned}
 & \times \prod_{l=1}^{k-1} \left(\mathbb{1}_{\mathbb{X}^{l+1}:r_{l,l+1}} \|\ell_{\hat{n}_l, m}^\infty \ell_{\hat{n}_l}^1 \cdot 2^{2r_{l,1}+2r_{l,l+1}} \cdot \|g_1\|_2^2 \cdot \|g_{l+1}\|_2 \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\
 & \times \prod_{k \leq u \leq d} \left(\prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j}:r_{u,j}}\| \ell_{\hat{n}_u, m}^{\frac{1}{k}} \ell_{\hat{n}_u}^1 \cdot 2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(d-1)}{k}} \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\
 & \times \prod_{k \leq u \leq d} \left(2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \cdot \left(\prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \cdot \|g_{1,u}\|_4^\alpha \cdot \prod_{l=2}^k \|g_l\|_4^\beta \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\
 & \times (2^t |F|)^{1 - \frac{d}{2(d+1)} + \varepsilon}.
 \end{aligned}$$

Developing the expression above,

$$\begin{aligned}
 & |\tilde{\Lambda}_{k,d}(g, h)| \\
 & \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times 2^{-\frac{(d+1)\varepsilon}{2kd} l_1} \times \left(\prod_{j=1}^d 2^{-r_{j,1}} \right)^{\frac{1}{k} - \frac{(d+1)\varepsilon}{2kd}} \times \|g_1\|_2^{\frac{(d+1)(d-1)\varepsilon}{2kd} - \frac{(d-1)}{k}} \\
 & \times \prod_{i=1}^{k-1} 2^{-\frac{(d+1)\varepsilon}{2kd} l_{i+1}} \times \prod_{i=1}^{k-1} \left[2^{-\frac{1}{d+1} \cdot r_{i,i+1}} \cdot \prod_{u=k}^d 2^{-\frac{1}{k(d+1)} \cdot r_{u,i+1}} \right]^{1 - \frac{(d+1)\varepsilon}{2d}} \\
 & \times \prod_{i=1}^{k-1} \left[\|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\| \ell_{\hat{n}_i, m}^\infty \ell_{\hat{n}_i}^1 \cdot \prod_{u=k}^d \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{u,i+1}}}\| \ell_{\hat{n}_u, m}^\infty \ell_{\hat{n}_u}^1 \right]^{\frac{(d+1)\varepsilon}{2d} - 1} \\
 & \times \left[\prod_{l=1}^{k-1} \|\mathbb{1}_{\mathbb{X}^{l+1}:r_{l,l+1}}\| \ell_{\hat{n}_l, m}^\infty \ell_{\hat{n}_l}^1 \right]^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \times \left[\prod_{l=1}^{k-1} (2^{r_{l,1}+r_{l,l+1}})^{\frac{1}{d+1} - \frac{2\varepsilon}{d}} \right] \times \|g_1\|_2^{\frac{(k-1)}{d+1} - \frac{2(k-1)\varepsilon}{d}} \cdot \prod_{l=1}^{k-1} \|g_{l+1}\|_2^{\frac{1}{d+1} - \frac{2\varepsilon}{d}} \\
 & \times \prod_{u=k}^d \left[\left(\prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j}:r_{u,j}}\| \ell_{\hat{n}_u, m}^{\frac{1}{k}} \ell_{\hat{n}_u}^1 \right)^{\frac{1}{k} \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \cdot \left(2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \right] \\
 & \times \|g_1\|_2^{\frac{2(d-k+1)(d-1)}{k} \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \cdot \prod_{k \leq u \leq d} \left[\left(\|g_{1,u}\|_4 \cdot \prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right) \right] \\
 & \times \prod_{l=2}^k \|g_l\|_4^{\beta(d-k+1) \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \times (2^t |F|)^{[1 - \frac{d}{2(d+1)}] + \varepsilon}. \tag{94}
 \end{aligned}$$

Observe that the product of the blue factors above (for $k \leq u \leq d$) is³²

$$\begin{aligned}
 \prod_{k \leq u \leq d} \left(\|g_{1,u}\|_4 \cdot \prod_{j \neq u} \|g_{1,j}\|_2 \right) &= \left[\prod_{l=1}^{k-1} \|g_{1,l}\|_2^{d-k+1} \right] \cdot \prod_{u=k}^d \left[\|g_{1,u}\|_2^{d-k} \cdot \|g_{1,u}\|_4 \right] \\
 &= \left[\prod_{j=1}^d \|g_{1,j}\|_2^{d-k} \right] \cdot \left[\prod_{l=1}^{k-1} \|g_{1,l}\|_2 \right] \cdot \left[\prod_{u=k}^d \|g_{1,u}\|_4 \right] \leq \|g_1\|_2^{d-k} \cdot |E_1|^{\frac{1}{4}}.
 \end{aligned}$$

³²Recall that $|g_1| = |g_{1,1} \otimes \dots \otimes g_{1,d}| \leq \mathbb{1}_{E_{1,1}} \otimes \dots \otimes \mathbb{1}_{E_{1,d}} \leq \mathbb{1}_{E_1}$.

Notice that the previous step was lossy, which also reflects in the suboptimal final exponent $p(k, d)$. Now we set the values of δ and $\tilde{\delta}$ (as functions of ε) to be such that

$$\begin{aligned} \delta \cdot \frac{(k+1)}{2k} \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right) &= \frac{(d+1)\varepsilon}{kd}, \\ \tilde{\delta} \cdot \frac{1}{2k} \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right) &= \frac{\varepsilon}{kd}. \end{aligned}$$

Simplifying the expression above with this choice of δ and $\tilde{\delta}$,

$$\begin{aligned} &|\tilde{\Lambda}_{k,d}(g, h)| \\ &\lesssim \left[\sum_{l_1 \geq 0} 2^{-\frac{(d+1)\varepsilon}{2kd} \cdot l_1} \right] \times \left[\prod_{j=1}^{k-1} \left(\sum_{r_{j,1} \geq 0} 2^{-\frac{3(d+1)\varepsilon}{2kd} \cdot r_{j,1}} \right) \right] \times \left[\prod_{u=k}^d \left(\sum_{r_{u,1} \geq 0} 2^{-\frac{(d+1)\varepsilon}{2kd} \cdot r_{u,1}} \right) \right] \\ &\times \left[\prod_{i=1}^{k-1} \left(\sum_{l_{i+1} \geq 0} 2^{-\frac{(d+1)\varepsilon}{2kd} \cdot l_{i+1}} \right) \right] \times \left[\sum_{t \geq 0} 2^{-t \left(\frac{d}{2(d+1)} - \varepsilon \right)} \right] \\ &\times \prod_{i=1}^{k-1} \left[\left(\sum_{r_{i,i+1} \geq 0} 2^{-\frac{3\varepsilon}{2d} \cdot r_{i,i+1}} \right) \cdot \prod_{u=k}^d \left(\sum_{r_{u,i+1} \geq 0} 2^{-\frac{\varepsilon}{2kd} \cdot r_{u,i+1}} \right) \right] \\ &\times \prod_{i=1}^{k-1} \left[\sup_{l_{i+1}, r_{i,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{i,i+1}}}\|_{\ell_{n_i, m}^\infty \ell_{\hat{n}_i}^1}^{-\frac{3\varepsilon}{4d}} \cdot \prod_{u=k}^d \sup_{l_{i+1}, r_{u,i+1}} \|\mathbb{1}_{\times^{l_{i+1}; r_{u,i+1}}}\|_{\ell_{n_u, m}^\infty \ell_{\hat{n}_u}^1}^{-\frac{3\varepsilon}{4kd}} \right] \\ &\times \|g_1\|_2^{\frac{(d-k)}{k(d+1)} - \frac{2(d-k)(k+1)\varepsilon}{kd} + \frac{(d-k)(d+1)\varepsilon}{kd} + \frac{(d+1)(d-1)\varepsilon}{2kd} - \frac{2(k-1)\varepsilon}{d} - \frac{2(d-k+1)(d-1)\varepsilon}{kd}} \cdot |E_1|^{\frac{(k+1)}{4k(d+1)} + \frac{(d+1)\varepsilon}{4kd}} \\ &\times \prod_{l=1}^{k-1} |E_{l+1}|^{\frac{(d+k+1)}{4k(d+1)} - \frac{\varepsilon}{d} - \frac{(d-k+1)\varepsilon}{2kd} + \frac{(d-k+1)\varepsilon}{4kd}} \times |F|^{[1 - \frac{d}{2(d+1)}] + \varepsilon}. \end{aligned} \tag{95}$$

By considerations identical to the ones in the end of Section 9, this implies

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim_\varepsilon |F|^{1 - \frac{d}{2(d+1)} + \varepsilon} \cdot |E_1|^{\frac{2d-k+1}{4k(d+1)}} \prod_{l=1}^{k-1} |E_{l+1}|^{\frac{d+k+1}{4k(d+1)}}. \tag{96}$$

To make all exponents of $|E_j|$ ($1 \leq j \leq k$) the same, we have to take

$$\frac{1}{\tilde{p}(k, d)} = \min \left\{ \frac{2d - k + 1}{4k(d + 1)}, \frac{d + k + 1}{4k(d + 1)} \right\}.$$

Again by the same considerations from Section 9, (96) implies³³ Theorem 12.1. □

12B. Near-restriction estimates without transversality. To make the notation lighter, let us omit the index Q and set \mathcal{E}_d be the extension operator associated to a fixed cube $Q \subset \mathbb{R}^d$. Recall the k -product

³³Notice that we obtain something slightly better than Theorem 12.1 if one is looking for *asymmetric estimates*: (96) implies a bound of type $L^{p_1} \times L^{p_2} \times L^{p_2} \times \dots \times L^{p_2} \rightarrow L^{2(d+1)/(kd) + \varepsilon}$, $p_1 \neq p_2$ and $p_1, p_2 \leq p(k, d)$, if g_1 is a tensor.

operator obtained from \mathcal{E}_d defined in (6)

$$\mathcal{E}_{d,(k)}(g_1, \dots, g_k) = \prod_{j=1}^k \mathcal{E}_d g_j.$$

In this subsection we prove [Theorem 1.18](#), which we restate here for the convenience of the reader.

Theorem 12.6. *Let $2 \leq k \leq d + 1$. If g_1 is a tensor, the inequality*

$$\left\| \prod_{j=1}^k \mathcal{E}_d g_j \right\|_{L^{2(d+1)/(kd)+\varepsilon}(\mathbb{R}^{d+1})} \lesssim_{\mathcal{Q},\varepsilon} \prod_{j=1}^k \|g_j\|_{L^4(\mathcal{Q})} \tag{97}$$

holds for all $\varepsilon > 0$.

Remark 12.7. As in the previous subsection, the difference between the proof of [Theorem 12.6](#) and the one done in [Section 9](#) is in the building blocks used: since there is no transversality to be exploited, we only use the best extension bound for the parabola (in the form of [Proposition 4.3](#)).

Proof of Theorem 12.6. The framework is the exact same as in the proof of [Theorem 12.1](#). We have to bound $\#\mathbb{X}^{\vec{l},R,t}$ to effectively estimate³⁴

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim \sum_{\vec{l},R,t} 2^{-t} \prod_{j=1}^k 2^{-\frac{t_j}{k}} \#\mathbb{X}^{\vec{l},R,t}$$

in terms of the measures of the sets $E_{1,\ell}$, $1 \leq \ell \leq d$, E_j , $2 \leq j \leq k$, and F . This will be done by the following analogue of [Lemma 12.5](#):

Lemma 12.8. *The two following extension-type bounds for the cardinality $\#\mathbb{X}^{\vec{l},R,t}$ hold:*

(a) *For all $1 \leq i \leq k - 1$ and all³⁵ $\lambda > 0$,*

$$\#\mathbb{X}^{\vec{l},R,t} \lesssim \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1} \cdot 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \cdot \|g_{1,i}\|_4^{2+\lambda} \cdot \left(\prod_{\ell \neq i} \|g_{1,\ell}\|_{2+\lambda}^{2+\lambda} \right) \cdot \|g_{i+1}\|_4^{2+\lambda}. \tag{98}$$

(b) *If $k < d + 1$, for all $k \leq u \leq d$,*

$$\begin{aligned} \#\mathbb{X}^{\vec{l},R,t} &\lesssim \prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j:r_{u,j}}}\|_{\ell_{n_u,m}^\infty \ell_{\hat{n}_u}^1}^{\frac{1}{k}} \cdot 2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(d-1)}{k}} \\ &\quad \times 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \cdot \left(\prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \cdot \|g_{1,u}\|_4^\alpha \cdot \prod_{l=2}^k \|g_{1,l}\|_4^\beta, \end{aligned} \tag{99}$$

where

$$\alpha := \frac{2(k+1)}{k} + \delta \cdot \frac{(k+1)}{2k}, \quad \beta := \frac{2}{k} + \tilde{\delta} \cdot \frac{1}{2k},$$

with $\delta, \tilde{\delta} > 0$ being arbitrarily small parameters to be chosen later.

Remark 12.9. We highlight that (99) is only going to be used if $k < d + 1$. The argument that follows will make it clear what changes in the case $k = d + 1$ if we only use (98).

³⁴Rigorously, we are dealing with a different operator here, but we will keep the notation unchanged for simplicity.

³⁵The parameter λ will be chosen later. It should be regarded as morally zero, and we only introduce it to be able to use [Proposition 4.3](#) since it does not hold at the endpoint.

Proof. We only prove (98), since (99) is identical to (89). From (53),

$$\#\mathbb{X}^{\vec{l},R,t} \leq \|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1} \cdot \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}\|_{\ell_{n_i,m}^1}.$$

We bound the second factor in the right-hand side above as follows:

$$\begin{aligned} & \|\mathbb{1}_{\mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}}\|_{\ell_{n_i,m}^1} \\ & \lesssim 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \sum_{(n_i,m) \in \mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}} \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_2^{2+\lambda} \cdot \|\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}\|_2^{2+\lambda} \\ & \leq 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \sum_{(n_i,m) \in \mathbb{B}_{i,1}^{r_{i,1}} \cap \mathbb{B}_{i,i+1}^{r_{i,i+1}}} \|\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}\|_{2+\lambda}^{2+\lambda} \cdot \|\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}\|_{2+\lambda}^{2+\lambda} \\ & \leq 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \iint \left(\sum_{(n_i,m) \in \mathbb{Z}^2} |\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}|^{2+\lambda} \cdot |\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}|^{2+\lambda} \right) d\hat{x}_i d\hat{y}_i \\ & \leq 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \iint \left(\sum_{(n_i,m) \in \mathbb{Z}^2} |\langle g_1, \varphi_{n_i,m}^{i,1} \rangle_{x_i}|^{4+2\lambda} \right)^{\frac{1}{2}} \cdot \left(\sum_{(n_i,m) \in \mathbb{Z}^2} |\langle g_{i+1}, \varphi_{n_i,m}^{i,i+1} \rangle_{y_i}|^{4+2\lambda} \right)^{\frac{1}{2}} d\hat{x}_i d\hat{y}_i \\ & \lesssim \lambda^{2(2+\lambda)(r_{i,1}+r_{i,i+1})} \iint \|g_1\|_{L_{x_i}^{4+\lambda}}^{2+\lambda} \cdot \|g_{i+1}\|_{L_{y_i}^{4+\lambda}}^{2+\lambda} d\hat{x}_i d\hat{y}_i \\ & \lesssim 2^{(2+\lambda)(r_{i,1}+r_{i,i+1})} \cdot \|g_{1,i}\|_4^{2+\lambda} \cdot \left(\prod_{\ell \neq i} \|g_{1,\ell}\|_{2+\lambda}^{2+\lambda} \right) \cdot \|g_{i+1}\|_4^{2+\lambda}, \end{aligned}$$

where we used Hölder’s inequality from the second to third lines, Fubini from the third to fourth, Hölder again twice, Proposition 4.3 and the fact that g_1 is a tensor. This finishes the proof of the lemma. \square

As in the previous subsection, given $\varepsilon > 0$, we bound $\tilde{\Lambda}_{k,d}$ using the estimates from (84) and Corollary 12.4, and the ones from Lemma 12.8 with the exact same weights³⁶ we used in the proof of Theorem 12.1:

$$\begin{cases} \theta_l = \frac{1}{2(d+1)} - \frac{\varepsilon}{d}, & 1 \leq l \leq d, \quad \text{for the } d \text{ estimates in (98) and (99),} \\ \theta_{d+1} = 1 - \frac{d}{2(d+1)} + \varepsilon & \text{for (86).} \end{cases}$$

Hence,

$$\begin{aligned} |\tilde{\Lambda}_{k,d}(g,h)| & \lesssim \sum_{\vec{l},R,t \geq 0} 2^{-t} \times 2^{-(\frac{d+1}{2kd})\varepsilon} l_1 \times \left(\frac{1}{\|g_1\|_2^{d-1}} \prod_{j=1}^d 2^{-r_{j,1}} \right)^{\frac{1}{k} - \frac{(d+1)\varepsilon}{2kd}} \\ & \times \prod_{i=1}^{k-1} 2^{-\frac{(d+1)\varepsilon}{2kd} l_{i+1}} \times \prod_{i=1}^{k-1} \left[\frac{2^{-\frac{1}{d+1} \cdot r_{i,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{i,i+1}}}\|_{\ell_{n_i,m}^\infty \ell_{\hat{n}_i}^1}} \cdot \prod_{u=k}^d \frac{2^{-\frac{1}{k(d+1)} \cdot r_{u,i+1}}}{\|\mathbb{1}_{\mathbb{X}^{l_{i+1}:r_{u,i+1}}}\|_{\ell_{n_u,m}^\infty \ell_{\hat{n}_u}^1}} \right]^{1 - \frac{(d+1)\varepsilon}{2d}} \end{aligned}$$

³⁶If $k = d + 1$, we give weight $\frac{1}{2(d+1)} - \frac{\varepsilon}{d}$ to each one of the d estimates in (98) only.

$$\begin{aligned} & \times \prod_{l=1}^{k-1} \left(2^{(2+\lambda)(r_{l,1}+r_{l,l+1})} \cdot \|g_{1,l}\|_4^{2+\lambda} \cdot \left(\prod_{\ell \neq l} \|g_{1,\ell}\|_{2+\lambda}^{2+\lambda} \right) \cdot \|g_{l+1}\|_4^{2+\lambda} \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\ & \times \prod_{k \leq u \leq d} \left(\prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j}; r_{u,j}}\|_{\ell_{\hat{n}_u, m}^{\infty} \ell_{\hat{n}_u}^1}^{\frac{1}{k}} \cdot 2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot \|g_1\|_2^{\frac{2(d-1)}{k}} \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\ & \times \prod_{k \leq u \leq d} \left(2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \cdot \left(\prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \cdot \|g_{1,u}\|_4^\alpha \cdot \prod_{l=2}^k \|g_l\|_4^\beta \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \\ & \times (2^t |F|)^{1 - \frac{d}{2(d+1)} + \varepsilon}. \end{aligned}$$

Developing the expression above³⁷,

$$\begin{aligned} & |\tilde{\tilde{\Lambda}}_{k,d}(g, h)| \\ & \lesssim \sum_{\vec{l}, R, t \geq 0} 2^{-t} \times 2^{-\frac{(d+1)\varepsilon}{2kd}} l_1 \times \left(\prod_{j=1}^d 2^{-r_{j,1}} \right)^{\frac{1}{k} - \frac{(d+1)\varepsilon}{2kd}} \times \|g_1\|_2^{\frac{(d+1)(d-1)\varepsilon}{2kd} - \frac{(d-1)}{k}} \\ & \times \prod_{i=1}^{k-1} 2^{-\frac{(d+1)\varepsilon}{2kd} l_{i+1}} \times \prod_{i=1}^{k-1} \left[2^{-\frac{1}{d+1} \cdot r_{i,i+1}} \cdot \prod_{u=k}^d 2^{-\frac{1}{k(d+1)} \cdot r_{u,i+1}} \right]^{1 - \frac{(d+1)\varepsilon}{2d}} \\ & \times \prod_{i=1}^{k-1} \left[\|\mathbb{1}_{\mathbb{X}^{l_{i+1}}; r_{i,i+1}}\|_{\ell_{\hat{n}_i, m}^{\infty} \ell_{\hat{n}_i}^1}^{\frac{1}{k} \cdot \left(\frac{(d+1)\varepsilon}{2d} - 1 \right)} \cdot \prod_{u=k}^d \|\mathbb{1}_{\mathbb{X}^{l_{i+1}}; r_{u,i+1}}\|_{\ell_{\hat{n}_u, m}^{\infty} \ell_{\hat{n}_u}^1}^{\frac{1}{k(d+1)} \cdot \left(\frac{(d+1)\varepsilon}{2d} - 1 \right)} \right] \\ & \times \left[\prod_{l=1}^{k-1} \|\mathbb{1}_{\mathbb{X}^{l_{l+1}}; r_{l,l+1}}\|_{\ell_{\hat{n}_l, m}^{\infty} \ell_{\hat{n}_l}^1}^{\frac{1}{k} \cdot \left(\frac{(d+1)\varepsilon}{2d} - \frac{\varepsilon}{d} \right)} \right] \times \left[\prod_{l=1}^{k-1} (2^{r_{l,1}+r_{l,l+1}})^{(2+\lambda) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \right] \\ & \times \left[\prod_{l=1}^{k-1} |E_{1,l}|^{\left(\frac{2+\lambda}{4} + (k-2) \right) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \right] \cdot \left[\prod_{u=k}^d |E_{1,u}|^{\left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right) \cdot (k-1)} \right] \cdot \left[\prod_{l=1}^{k-1} |E_{l+1}|^{\left(\frac{2+\lambda}{4} \right) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \right] \\ & \times \prod_{u=k}^d \left[\left(\prod_{j=2}^k \|\mathbb{1}_{\mathbb{X}^{l_j}; r_{u,j}}\|_{\ell_{\hat{n}_u, m}^{\infty} \ell_{\hat{n}_u}^1}^{\frac{1}{k} \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \right) \cdot \left(2^{\frac{2}{k} \sum_{i \neq u} r_{i,1}} \cdot 2^{\alpha \cdot r_{u,1} + \sum_{l=2}^k \beta \cdot r_{u,l}} \right)^{\frac{1}{2(d+1)} - \frac{\varepsilon}{d}} \right] \\ & \times \|g_1\|_2^{\frac{2(d-k+1)(d-1)}{k} \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \cdot \prod_{k \leq u \leq d} \left[\left(\|g_{1,u}\|_4 \cdot \prod_{j \neq u} \|g_{1,j}\|_2 \right)^\alpha \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right) \right] \\ & \times \prod_{l=2}^k \|g_l\|_4^{\beta(d-k+1) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d} \right)} \times (2^t |F|)^{\left[1 - \frac{d}{2(d+1)} \right] + \varepsilon}. \tag{100} \end{aligned}$$

Observe that we highlighted a few factors in red in (100); this is just to compare them to the red terms in (94): the red terms are the only ones that differ in the right-hand sides of (94) and (100). On the other hand, we will bound the product of the blue factors³⁸ in (100) in a slightly better way than we did in the

³⁷The products in the fourth and fifth lines above are void if $k = d + 1$. We can think of them as being 1.

³⁸The seventh and eighth lines are void if $k = d + 1$, hence the blue factors do not contribute at all in this case.

proof of [Theorem 12.1](#):

$$\begin{aligned} \prod_{k \leq u \leq d} \left(\|g_{1,u}\|_4 \cdot \prod_{j \neq u} \|g_{1,j}\|_2 \right) &= \left[\prod_{j=1}^d \|g_{1,j}\|_2^{d-k} \right] \cdot \left[\prod_{l=1}^{k-1} \|g_{1,l}\|_2 \right] \cdot \left[\prod_{u=k}^d \|g_{1,u}\|_4 \right] \\ &\leq \left[\prod_{l=1}^{k-1} |E_{1,l}|^{\frac{d-k+1}{2}} \right] \cdot \left[\prod_{u=k}^d |E_{1,u}|^{\frac{d-k}{2} + \frac{1}{4}} \right]. \end{aligned} \tag{101}$$

Setting δ and $\tilde{\delta}$ exactly as in the previous subsection and using the observations we just made, we conclude that the final bound for $|\tilde{\Lambda}_{k,d}(g, h)|$ compares to [\(96\)](#) exactly as follows:

- The coefficients of the “ $r_{j,1}$ power” is now

$$2^{\left[-\frac{3(d+1)\varepsilon}{2kd} + \lambda\left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d}\right)\right]} r_{j,1},$$

whereas in [\(96\)](#) it was

$$2^{\left(-\frac{3(d+1)\varepsilon}{2kd}\right)} r_{j,1}.$$

- For $1 \leq l \leq k - 1$, [\(101\)](#) gives $|E_{1,l}|$ an extra power of³⁹

$$\left(\frac{1}{2} + \frac{1}{2k}\right) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d}\right) + \frac{(d+1)\varepsilon}{4kd}.$$

On the other hand, still for $1 \leq l \leq k - 1$, the red factors in [\(100\)](#) produce a power of $|E_{1,l}|$ that is exactly

$$\frac{(2-\lambda)}{4} \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d}\right) \tag{102}$$

less than the one produced by the corresponding red factors in [\(94\)](#). If $k < d + 1$, these provide a *net gain* of

$$\left(\frac{1}{2k} - \frac{\lambda}{4}\right) \cdot \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d}\right) + \frac{(d+1)\varepsilon}{4kd}$$

in the final power of $|E_{1,l}|$. If $k = d + 1$, we just lose (compared to [\(96\)](#)) [\(102\)](#) in the final power of $|E_{1,l}|$.

- For $k \leq u \leq d$, the powers of the measures $|E_{1,u}|$ are exactly the same in both [\(94\)](#) and in [\(100\)](#).
- For $2 \leq l \leq k$, the red factors in [\(100\)](#) produce a power of $|E_l|$ that is exactly

$$\frac{(2-\lambda)}{4} \left(\frac{1}{2(d+1)} - \frac{\varepsilon}{d}\right)$$

less than the one produced by the corresponding red factors in [\(94\)](#).

- All other factors are precisely the same.

³⁹Here we are using the explicit choice of δ .

By choosing λ small enough compared to ε and by the same considerations made in the end of [Section 9](#), this implies

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim_\varepsilon |F|^{1-\frac{d}{2(d+1)}+\varepsilon} \cdot |E_1|^{\frac{2d-k+2}{4k(d+1)}} \prod_{l=1}^{k-1} |E_{l+1}|^{\frac{1}{4k}}$$

for $k < d + 1$ and

$$|\tilde{\Lambda}_{k,d}(g, h)| \lesssim_\varepsilon |F|^{1-\frac{d}{2(d+1)}+\varepsilon} \cdot \prod_{l=1}^k |E_l|^{\frac{1}{4k}}$$

for $k = d + 1$. Again by the same considerations from [Section 9](#), these imply [Theorem 12.6](#). □

13. Weak transversality, Brascamp–Lieb and an application

We were recently asked by Jonathan Bennett if there was a link between our results and the theory of Brascamp–Lieb inequalities. The motivation for that comes from the fact that, assuming $g_1 = g_{1,1} \otimes \dots \otimes g_{1,d}$, one can see the operator $\mathcal{ME}_{d+1,d}$ as the $2d$ -linear object

$$T(g_{1,1}, \dots, g_{1,d}, g_2, \dots, g_{d+1}) := \mathcal{ME}_{d+1,d}(g_{1,1} \otimes \dots \otimes g_{1,d}, g_2, \dots, g_{d+1}),$$

and given that such a link exists in the theory of $\mathcal{ME}_{d+1,d}$ (see [\[Bennett 2014\]](#)), it is natural to wonder if boundedness for T is related somehow to the finiteness condition of certain Brascamp–Lieb constants $\text{BL}(\mathbf{L}, \mathbf{p})$.

The purposes of this section are to make this connection clear and to give a modest application of our results to the theory of *restriction-Brascamp–Lieb inequalities*.

13A. A link between weak transversality and Brascamp–Lieb inequalities. We start with some classical background. Let $L_j : \mathbb{R}^n \rightarrow \mathbb{R}^{n_j}$ be linear maps and $p_j \geq 0$, $1 \leq j \leq m$. Inequalities of the form

$$\int_{\mathbb{R}^n} \prod_{j=1}^m (f_j \circ L_j)^{p_j}(v) \, dv \leq C \prod_{j=1}^m \left(\int_{\mathbb{R}^{n_j}} f_j(y_j) \, dy_j \right)^{p_j} \tag{103}$$

are called *Brascamp–Lieb inequalities*. Bennett, Carbery, Christ and Tao [\[Bennett et al. 2008\]](#) established for which *Brascamp–Lieb data* (\mathbf{L}, \mathbf{p}) the inequality above holds, where $\mathbf{L} = (L_1, \dots, L_m)$ and $\mathbf{p} = (p_1, \dots, p_m)$. The best constant for which [\(103\)](#) holds for all nonnegative input functions $f_j \in L^1(\mathbb{R}^{n_j})$ is denoted by $\text{BL}(\mathbf{L}, \mathbf{p})$.

Theorem 13.1 [\[Bennett et al. 2008\]](#). *The constant $\text{BL}(\mathbf{L}, \mathbf{p})$ in [\(103\)](#) is finite if and only if for all subspaces $V \subset \mathbb{R}^n$*

$$\dim(V) \leq \sum_{j=1}^m p_j \dim(L_j V) \tag{104}$$

and

$$\sum_{j=1}^m p_j n_j = n. \tag{105}$$

Remark 13.2. By taking $V = \mathbb{R}^n$ in [\(104\)](#) it follows that each L_j must be surjective for [\(105\)](#) to hold as well.

We will work with explicit maps L_j and use [Theorem 13.1](#) to establish a link between the concept of weak transversality and inequalities such as [\(103\)](#).⁴⁰ These maps will be associated to the submanifolds relevant to the problem at hand: the d -dimensional paraboloid \mathbb{P}^d in \mathbb{R}^{d+1} and some “canonical” two-dimensional parabolas.

In order to define L_j , we fix standard parametrizations for the submanifolds mentioned above. Let

$$\Gamma : \mathbb{R}^d \longrightarrow \mathbb{R}^{d+1}, \tag{106}$$

$$(x_1, \dots, x_d) \longmapsto (x_1, \dots, x_d, \sum_{i=1}^d x_i^2), \tag{107}$$

parametrize \mathbb{P}^d and

$$\gamma_j : \mathbb{R} \longrightarrow \mathbb{R}^{d+1}, \tag{108}$$

$$x \longmapsto (x \cdot \delta_{1j}, \dots, x \cdot \delta_{dj}, x^2), \tag{109}$$

parametrize a parabola in the two-dimensional canonical subspace generated by e_j and e_{d+1} (δ_{ij} is the Kronecker delta). Their differentials are given by

$$d\Gamma : \mathbb{R}^d \longrightarrow M_{(d+1) \times d}, \quad (x_1, \dots, x_d) \longmapsto \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \\ 2x_1 & 2x_2 & \dots & 2x_d \end{bmatrix}$$

and

$$d\gamma_j : \mathbb{R} \longrightarrow M_{(d+1) \times 1}, \quad x \longmapsto [\delta_{1j} \ \delta_{2j} \ \dots \ \delta_{dj} \ 2x]^\top.$$

For $d + 1$ points $x^j = (x_1^j, \dots, x_d^j) \in \mathbb{R}^d$, $1 \leq j \leq d + 1$, define the linear maps⁴¹

$$\begin{aligned} L_\ell^{x_\ell^1} &:= (d\gamma_\ell(x_\ell^1))^* && \text{for all } 1 \leq \ell \leq d, \\ L_{d+\ell}^{x_\ell^{d+1}} &:= (d\Gamma(x_1^{\ell+1}, \dots, x_d^{\ell+1}))^* && \text{for all } 1 \leq \ell \leq d. \end{aligned} \tag{110}$$

It is important to emphasize that $L_{d+\ell}$ depends on $x^{\ell+1}$ (and similarly, L_ℓ depends on x_ℓ^1). The main result of this subsection is:

Theorem 13.3. *Let $\mathcal{Q} = \{Q_1, \dots, Q_{d+1}\}$ be a collection of closed cubes in \mathbb{R}^d . If \mathcal{Q} is weakly transversal with pivot Q_1 , then for any choice of points $x^j = (x_1^j, \dots, x_d^j) \in Q_j$, the linear maps in [\(110\)](#) satisfy*

$$\text{BL}(\mathbf{L}(x), \mathbf{p}) < \infty \quad \text{for } \mathbf{L}(x) = (L_1^{x_1^1}, \dots, L_{2d}^{x_d^{d+1}}) \text{ and } \mathbf{p} = \left(\frac{1}{d}, \dots, \frac{1}{d}\right). \tag{111}$$

Conversely, if [\(111\)](#) is satisfied by the linear maps in [\(110\)](#) for any choice of points $x^j = (x_1^j, \dots, x_d^j) \in Q_j$, then \mathcal{Q} can be decomposed into $O(1)$ weakly transversal collections \mathcal{Q}' of $d + 1$ cubes, each one having a cube $Q'_1 \subset Q_1$ as pivot.

⁴⁰From now on, we will replace n by $d + 1$ when referring to the dimension of the euclidean space.

⁴¹We highlight that the *superscript* j in x^j_i denotes the *point*, whereas the *subscript* i denotes the i -coordinate of the corresponding point. Notice also that we are identifying the adjoint operator T^* with the transpose of the matrix that represents T in the canonical basis.

Remark 13.4. If \mathcal{Q} can be decomposed into $O(1)$ weakly transversal collections \mathcal{Q}' of $d+1$ cubes (in the sense of Claim 3.4), each one having a cube $Q'_1 \subset Q_1$ as pivot, then the conclusion of the first part of the theorem above also holds for \mathcal{Q} . Some important examples to keep in mind are the ones of transversal configurations that are *not* weakly transversal by themselves, but that are decomposable into such: for instance, $\{Q_1, Q_2, Q_3\}$, where $Q_1 = [1, 4] \times [2, 3]$, $Q_2 = [0, 2] \times [0, 1]$ and $Q_3 = [3, 5] \times [0, 1]$ is a transversal collection of cubes in \mathbb{R}^2 , but not weakly transversal with pivot Q_1 since $\pi_1(Q_1)$ intersects both $\pi_1(Q_2)$ and $\pi_1(Q_3)$.

Remark 13.5. We can of course obtain a similar statement if \mathcal{Q} is weakly transversal with any other pivot $Q_j, j \neq 1$. The linear maps L_ℓ and $L_{d+\ell}$ would have to be changed accordingly.

Proof of Theorem 13.3. Suppose that \mathcal{Q} is weakly transversal with pivot Q_1 . We can then assume without loss of generality that

$$\begin{cases} \pi_1(Q_1) \cap \pi_1(Q_2) = \emptyset, \\ \vdots \\ \pi_d(Q_1) \cap \pi_d(Q_{d+1}) = \emptyset. \end{cases} \tag{112}$$

The strategy is to apply Theorem 13.1. Condition (105) is trivially satisfied, so we just have to check (104). Fix the points $x^j = (x_1^j, \dots, x_d^j) \in Q_j, 1 \leq j \leq d$. To avoid heavy notation, we will omit the superscripts x_ℓ^1 and $x^{\ell+1}$ when referring to $L_\ell^{x_\ell^1}$ and $L_{d+\ell}^{x^{\ell+1}}$, respectively, but these points will be referenced whenever they play an important role. We emphasize that the maps $L_\ell, 1 \leq \ell \leq d$, are being identified with the row vector

$$[\delta_{1\ell} \ \delta_{2\ell} \ \dots \ \delta_{d\ell} \ 2x_\ell^1],$$

whereas the maps $L_{d+\ell}, 1 \leq \ell \leq d$, are identified with the $d \times (d+1)$ matrix

$$\begin{bmatrix} 1 & 0 & \dots & 0 & 2x_1^{\ell+1} \\ 0 & 1 & \dots & 0 & 2x_2^{\ell+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 2x_d^{\ell+1} \end{bmatrix}.$$

If $V \subset \mathbb{R}^{d+1}$ is a subspace of dimension k , we have to verify that

$$dk \leq \sum_{j=1}^d \dim(L_j V) + \sum_{\ell=1}^d \dim(L_{d+\ell} V). \tag{113}$$

Suppose that there are exactly $m \geq 0$ indices $j \in \{1, \dots, d\}$ such that $\dim(L_j V) = 0$. If $m = 0$, we must have $L_j V = \mathbb{R}$ for all $1 \leq j \leq d$; hence

$$\sum_{j=1}^d \dim(L_j V) = d. \tag{114}$$

Surjectivity of $L_{d+\ell}, 1 \leq \ell \leq d$, implies $\dim(\ker(L_{d+\ell})) = 1$, which gives the lower bound $\dim(L_{d+\ell} V) \geq k - 1$. We then obtain

$$\sum_{\ell=1}^d \dim(L_{d+\ell} V) \geq d(k - 1). \tag{115}$$

It is clear that (114) and (115) together verify (113) in the $m = 0$ case. If $m \geq 1$, assume without loss of generality that

$$L_1V = \dots = L_mV = 0, \tag{116}$$

$$L_{m+1}V = \dots = L_dV = \mathbb{R}. \tag{117}$$

This gives us

$$\sum_{j=1}^d \dim(L_jV) = d - m. \tag{118}$$

We will show that

$$\sum_{\ell=1}^d \dim(L_{d+\ell}V) \geq (d - m)(k - 1) + mk. \tag{119}$$

Observe that (118) and (119) together verify (113) in the $m \geq 1$ case.

We claim that there are at least m maps L_{ℓ_j} among $L_{\ell+1}, \dots, L_{2d}$ such that $\dim(L_{\ell_j}V) = k$. If not, there are $d - m + 1$ maps $L_{\ell_1}, \dots, L_{\ell_{d-m+1}}$ with $\dim(L_{\ell_j}V) \leq k - 1$. Since $\dim V = k$, the rank-nullity theorem implies the existence of

$$0 \neq v^{\ell_j} \in \ker(L_{\ell_j}) \cap V, \quad 1 \leq j \leq d - m + 1. \tag{120}$$

By (116),

$$L_r v^{\ell_j} = v_r^{\ell_j} + 2x_r^1 v_{d+1}^{\ell_j} = 0, \quad 1 \leq r \leq m, \tag{121}$$

and by (120) we have

$$L_{\ell_j} v^{\ell_j} = \begin{bmatrix} 1 & 0 & \dots & 0 & 2x_1^{\ell_j-d+1} \\ 0 & 1 & \dots & 0 & 2x_2^{\ell_j-d+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 2x_d^{\ell_j-d+1} \end{bmatrix} \cdot \begin{bmatrix} v_1^{\ell_j} \\ v_2^{\ell_j} \\ \vdots \\ v_{d+1}^{\ell_j} \end{bmatrix} = \begin{bmatrix} v_1^{\ell_j} + 2x_1^{\ell_j-d+1} v_{d+1}^{\ell_j} \\ v_2^{\ell_j} + 2x_2^{\ell_j-d+1} v_{d+1}^{\ell_j} \\ \vdots \\ v_d^{\ell_j} + 2x_d^{\ell_j-d+1} v_{d+1}^{\ell_j} \end{bmatrix} = 0 \tag{122}$$

for $1 \leq j \leq d - m + 1$. For each $1 \leq r \leq m$, combining the information from (121) and (122) gives us

$$v_{d+1}^{\ell_j} \cdot (x_r^1 - x_r^{\ell_j-d+1}) = 0.$$

If $v_{d+1}^{\ell_j} = 0$, then (122) also implies $v_n^{\ell_j} = 0$ for all $n \in \{1, \dots, d\}$; thus $v^{\ell_j} = 0$, which contradicts (120). Then we must have

$$x_r^1 = x_r^{\ell_j-d+1}, \quad 1 \leq r \leq m.$$

Let us now see why this cannot happen. We have just shown that there are $d - m + 1$ values of α for which

$$\begin{cases} \pi_1(Q_1) \cap \pi_1(Q_\alpha) \neq \emptyset, \\ \vdots \\ \pi_m(Q_1) \cap \pi_m(Q_\alpha) \neq \emptyset. \end{cases} \tag{123}$$

On the other hand, (112) tells us that $\alpha \notin \{2, 3, \dots, m + 1\}$; hence there are at most $d - m$ possible values for α (we cannot have $\alpha = 1$ either), which is a contradiction.

Hence there are at least m maps L_{ℓ_j} among $L_{\ell+1}, \dots, L_{2d}$ such that $\dim(L_{\ell_j}V) = k$. The remaining $d - m$ maps have kernels of dimension 1, so the image of V through them has dimension at least $k - 1$ (again by surjectivity of L_{ℓ_j} and the rank-nullity theorem). This verifies (119).

For the converse implication, suppose that (111) is satisfied by the linear maps in (110) for any choice of points $(x_1^j, \dots, x_d^j) \in Q_j$. As a consequence of the proof of Claim B.4, each $Q_l \in \mathcal{Q}$ can be partitioned into $O(1)$ subcubes

$$Q_l = \bigcup_i Q_{l,i}$$

so that all collections $\tilde{\mathcal{Q}}$ made of picking one subcube $Q_{l,i}$ per Q_l

$$\tilde{\mathcal{Q}} = \{\tilde{Q}_1, \dots, \tilde{Q}_{d+1}\}, \quad \tilde{Q}_l \in \{Q_{l,i}\}_i,$$

satisfy the following:

- (a) For any two $\tilde{Q}_r, \tilde{Q}_s \in \tilde{\mathcal{Q}}$, either $\pi_j(\tilde{Q}_r) \cap \pi_j(\tilde{Q}_s) = \emptyset$, or $\pi_j(\tilde{Q}_r) = \pi_j(\tilde{Q}_s)$, or $\pi_j(\tilde{Q}_r) \cap \pi_j(\tilde{Q}_s) = \{p_{r,s}\}$, where $p_{r,s}$ is an endpoint of both $\pi_j(\tilde{Q}_r)$ and $\pi_j(\tilde{Q}_s)$.
- (b) All $\pi_j(\tilde{Q}_s)$ that intersect a given $\pi_j(\tilde{Q}_r)$ (but distinct from it) do so at the same endpoint.⁴²

By a slight abuse of notation, let \mathcal{Q} denote one such subcollection that has the two properties above. Suppose, by contradiction, that \mathcal{Q} is not weakly transversal with pivot Q_1 (recall that this is a cube obtained from the original Q_1). The strategy now is to construct a subspace $V \subset \mathbb{R}^{d+1}$ that contradicts (104) for a certain choice of one point per cube in \mathcal{Q} . This construction will exploit a certain feature of a special subset of \mathcal{Q} , which is the content of Claim 13.6.

For simplicity of future references, let us say that a subset $\mathcal{A} \subset \mathcal{Q}$ has the *property (P)* if:

- (1) $Q_1 \in \mathcal{A}$.
- (2) \mathcal{A} is not weakly transversal with pivot Q_1 .

We say that a subset $\mathcal{A} \subset \mathcal{Q}$ is *minimal* if $\mathcal{A}' \subset \mathcal{A}$ has the property (P) if and only if $\mathcal{A}' = \mathcal{A}$. It is clear that, since \mathcal{Q} has the property (P) itself, it must contain a minimal subset of cardinality at least 2.

Claim 13.6. *Let $\mathcal{A} = \{Q_1, K_2, \dots, K_n\}$ be a minimal set of n cubes.⁴³ There is a set D of $d - n + 2$ canonical directions v for which*

$$\pi_v(Q_1) \cap \pi_v(K_j) \neq \emptyset \quad \text{for all } 2 \leq j \leq n. \tag{124}$$

Proof of Claim 13.6. See Claim B.6 in Appendix B. □

We know that \mathcal{Q} has a minimal subset of cardinality $2 \leq n \leq d + 1$. By the previous claim and by conditions (a) and (b) of our initial reductions, if $\mathcal{A}' = \{Q_1, K_2, \dots, K_n\}$ is a minimal subset of \mathcal{Q} , for

⁴²In other words, all $\pi_j(\tilde{Q}_s)$ that intersect a given $\pi_j(\tilde{Q}_r)$ (but distinct from it) do so on the same side. In short notation, let $S_{j,r}$ be the set of s for which $\pi_j(\tilde{Q}_r) \cap \pi_j(\tilde{Q}_s) \neq \emptyset$. The conclusion is that there is some real number γ_j such that $\gamma_j \in \pi_j(Q_r) \cap \bigcap_{s \in S_{j,r}} \pi_j(Q_s)$.

⁴³Observe that Q_1 is the only “ \mathcal{Q} ” cube in this collection. The others are labeled by K_j .

every $v \in D$ there is a number γ_v such that

$$\gamma_v \in \pi_v(Q_1) \cap \bigcap_{j=2}^n \pi_v(K_j).$$

Indeed, $\pi_v(Q_1)$ intersects each $\pi_v(Q_j)$ “on the same side”, so the intersection above must be nonempty (the existence of these γ_v is the only reason why we may need to decompose the initial collection \mathcal{Q} into subcollections that satisfy (a) and (b)).

For simplicity and without loss of generality, assume that $\mathcal{A} = \{Q_1, Q_2, \dots, Q_n\}$ is minimal⁴⁴ and $D = \{e_1, \dots, e_{d-n+2}\}$. Consider the points

$$\begin{aligned} (\gamma_1, \dots, \gamma_{d-n+2}, x_{d-n+3}^j, \dots, x_d^j) &\in Q_j, \quad 1 \leq j \leq n, \\ (x_1^l, \dots, x_d^l) &\in Q_l, \quad n+1 \leq l \leq d+1. \end{aligned}$$

By hypothesis, $\text{BL}(L(x), p) < \infty$ for the following collection of linear maps and exponents:

$$\begin{aligned} L_r^{\gamma_r}(v_1, \dots, v_{d+1}) &= v_r + 2\gamma_r v_{d+1}, \quad 1 \leq r \leq d-n+2, \\ L_s^{x_s^1}(v_1, \dots, v_{d+1}) &= v_s + 2x_s^1 v_{d+1}, \quad d-n+3 \leq s \leq d, \end{aligned}$$

$$L_{d+r}^{(\gamma_1, \dots, \gamma_{d-n+2}, x_{d-n+3}^{r+1}, \dots, x_d^{r+1})}(v_1, \dots, v_{d+1}) = \begin{bmatrix} v_1 + 2\gamma_1 v_{d+1} \\ \vdots \\ v_{d-n+2} + 2\gamma_{d-n+2} v_{d+1} \\ v_{d-n+3} + 2x_{d-n+3}^{r+1} v_{d+1} \\ \vdots \\ v_d + 2x_d^{r+1} v_{d+1} \end{bmatrix}, \quad 1 \leq r \leq n-1,$$

$$L_{d+l}^{x^{l+1}} = \begin{bmatrix} v_1 + 2x_1^{l+1} v_{d+1} \\ \vdots \\ v_d + 2x_d^{l+1} v_{d+1} \end{bmatrix}, \quad n \leq l \leq d, \quad p = \left(\frac{1}{d}, \dots, \frac{1}{d}\right).$$

Define

$$V := \bigcap_{r=1}^{d-n+2} \ker(L_r^{\gamma_r}).$$

Observe that $\dim(V) = n - 1$. Indeed, if we start with a vector $v = (v_1, \dots, v_{d+1})$ of $d + 1$ “free coordinates”, we lose one degree of freedom for each kernel in the intersection above, since $L_r^{\gamma_r}(v) = 0$ gives a relation between v_r and v_{d+1} . We have $d - n + 2$ many of them; hence the total degree of freedom is $(d + 1) - (d - n + 2) = n - 1$, which is the dimension of V . On the other hand, for every $v \in V$ we have by definition

$$L_r^{\gamma_r}(v) = 0, \quad 1 \leq r \leq d - n + 2.$$

Hence

$$\sum_{j=1}^d \dim(L_j V) \leq n - 2.$$

⁴⁴Here we are assuming $K_j = Q_j$, $2 \leq j \leq n$.

Also,

$$L_{d+r}^{(\gamma_1, \dots, \gamma_{d-n+2}, x_{d-n+3}^{r+1}, \dots, x_d^{r+1})}(v) = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ v_{d-n+3} + 2x_{d-n+3}^{r+1}v_{d+1} \\ \vdots \\ v_d + 2x_d^{r+1}v_{d+1} \end{bmatrix}, \quad 1 \leq r \leq n-1.$$

Thus

$$\dim(L_{d+r}V) \leq n-2, \quad 1 \leq r \leq n-1.$$

Since $\dim(V) = n-1$, we have the trivial bound

$$\dim(L_{d+l}V) \leq n-1, \quad n \leq l \leq d.$$

Altogether, these bounds imply

$$\begin{aligned} \frac{1}{d} \left(\sum_{j=1}^d \dim(L_j V) + \sum_{\ell=1}^d \dim(L_{d+\ell} V) \right) &\leq \frac{1}{d} [(n-2) + (n-1)(n-2) + (d-n+1)(n-1)] \\ &= \frac{1}{d} [(n-1)d - 1] < n-1 = \dim(V). \end{aligned}$$

Our initial hypothesis, however, is that $\text{BL}(\mathbf{L}(x), \mathbf{p}) < \infty$; therefore by [Theorem 13.1](#) we must have

$$\dim(V) \leq \frac{1}{d} \left(\sum_{j=1}^d \dim(L_j V) + \sum_{\ell=1}^d \dim(L_{d+\ell} V) \right),$$

which gives a contradiction. We conclude that \mathcal{Q} is weakly transversal with pivot Q_1 . □

13B. An application to Restriction-Brascamp–Lieb inequalities. The following conjecture was proposed in Bennett, Bez, Flock and Lee [\[Bennett et al. 2018\]](#):

Conjecture 13.7. *Suppose that, for each $1 \leq j \leq m$, $\Sigma_j : U_j \mapsto \mathbb{R}^n$ is a smooth parametrization of a n_j -dimensional submanifold S_j of \mathbb{R}^n by a neighborhood U_j of the origin in \mathbb{R}^{n_j} . Let*

$$\mathcal{E}_j g_j(\xi) := \int_{U_j} e^{-2\pi i \xi \cdot \Sigma_j(x)} g_j(x) \, dx$$

be the associated (parametrized) extension operator. If the Brascamp–Lieb constant $\text{BL}(\mathbf{L}, \mathbf{p})$ is finite for the linear maps $L_j := (d\Sigma_j(0))^ : \mathbb{R}^n \mapsto \mathbb{R}^{n_j}$, then provided the neighborhoods U_j of 0 are chosen to be small enough, the inequality*

$$\int_{\mathbb{R}^n} \prod_{j=1}^m |\mathcal{E}_j g_j|^{2p_j} \lesssim \prod_{j=1}^m \|g_j\|_{L^2(U_j)}^{2p_j} \tag{125}$$

holds for all $g_j \in L^2(U_j)$, $1 \leq j \leq m$.

Remark 13.8. The weaker inequality

$$\int_{B(0,R)} \prod_{j=1}^m |\mathcal{E}_j g_j|^{2p_j} \lesssim_\varepsilon R^\varepsilon \prod_{j=1}^m \|g_j\|_{L^2(U_j)}^{2p_j} \tag{126}$$

involving an arbitrary $\varepsilon > 0$ loss was established in [\[Bennett et al. 2018\]](#).

Remark 13.9. Very few cases of [Conjecture 13.7](#) are fully understood.⁴⁵ Recently, Bennett, Nakamura and Shiraki settled the *rank-1 case* $n_1 = \dots = n_m = 1$ as an application of their results on *tomographic Fourier analysis*.⁴⁶

Given their hybrid nature, estimates such as (125) are called *restriction-Brascamp–Lieb inequalities*.

Our goal here is to verify [Conjecture 13.7](#) in a special case. We chose to state the main result of this subsection in a way that does not emphasize the origin in the domains of Σ_j . The reason for this choice is that it brings to light key geometric features of the problem.

We will need a result from [\[Bennett et al. 2018\]](#) on the stability of Brascamp–Lieb constants⁴⁷:

Theorem 13.10 [\[Bennett et al. 2018\]](#). *Suppose that (L^0, \mathbf{p}) is a Brascamp–Lieb datum for which $\text{BL}(L^0, \mathbf{p}) < \infty$. Then there exists $\delta > 0$ and a constant $C < \infty$ such that*

$$\text{BL}(\mathbf{L}, \mathbf{p}) \leq C$$

whenever $\|\mathbf{L} - L^0\| < \delta$.

Now we are ready to state and prove our result:

Theorem 13.11. *Let Γ and γ_j be the parametrizations from (106) and (108), respectively. If, for $x^j = (x_1^j, \dots, x_d^j) \in \mathbb{R}^d$, the linear maps in (110) satisfy*

$$\text{BL}(\mathbf{L}(x), \mathbf{p}) < \infty \quad \text{for } \mathbf{L}(x) = (L_1^{x_1^1}, \dots, L_{2d}^{x_d^{d+1}}) \text{ and } \mathbf{p} = \left(\frac{1}{d}, \dots, \frac{1}{d}\right), \tag{127}$$

then there are small enough cube-neighborhoods $U_i \subset \mathbb{R}$ ($1 \leq i \leq d$) of x_i^1 and $V_\ell \subset \mathbb{R}^d$ of x^ℓ ($2 \leq \ell \leq d + 1$) for which (125) holds.

Remark 13.12. Rephrasing [Theorem 13.11](#) in terms of the original statement, it says that [Conjecture 13.7](#) holds for⁴⁸

$$\begin{aligned} \Sigma_i &= \gamma_i - (\delta_{1i} \cdot x_i^1, \dots, \delta_{di} \cdot x_i^1, 0), \quad 1 \leq i \leq d. \\ \Sigma_\ell &= \Gamma - (x^{\ell-d+1}, 0), \quad d + 1 \leq \ell \leq 2d. \\ m &= 2d, \quad \mathbf{p} = \left(\frac{1}{d}, \dots, \frac{1}{d}\right). \end{aligned}$$

Proof of Theorem 13.11. The argument is just a matter of putting the pieces together. By (127) and [Theorem 13.10](#), there are small enough cube-neighborhoods $U_i \subset \mathbb{R}$ ($1 \leq i \leq d$) of x_i^1 and $V_\ell \subset \mathbb{R}^d$ of x^ℓ ($2 \leq \ell \leq d + 1$) for which (127) still holds⁴⁹. Define

$$Q_1 := \overline{U_1 \times \dots \times U_d}, \quad Q_\ell := \overline{V_\ell}, \quad 2 \leq \ell \leq d + 1.$$

⁴⁵Most of them being very elementary situations, as mentioned in [\[Bennett et al. 2018\]](#).

⁴⁶See [\[Bennett and Nakamura 2021\]](#) for a more detailed exposition of this approach.

⁴⁷[Theorem 13.10](#) says that the map $L \mapsto \text{BL}(\mathbf{L}, \mathbf{p})$ is *locally bounded* for a fixed \mathbf{p} , and this is enough for our purposes. On the other hand, it was shown in [\[Bennett et al. 2017\]](#) that the Brascamp–Lieb constant is *continuous* in \mathbf{L} . It was later shown in [\[Bennett et al. 2020\]](#) that $\text{BL}(\mathbf{L}, \mathbf{p})$ is in fact *locally Hölder continuous* in \mathbf{L} .

⁴⁸Observe that we are just translating the domain of the Σ 's back to the origin.

⁴⁹Our maps L_j are sufficiently smooth for the stability theorem to be applied. The entries of the matrices that represent them are polynomials.




Figure 4. Unveiling the geometric features of the problem when $d = 2$. The cubes we find from [Theorem 13.10](#) are weakly transversal, which gives us access to our earlier results.

Now we apply [Theorem 13.3](#) to conclude that the collection $\mathcal{Q} = \{Q_1, \dots, Q_{d+1}\}$ can be decomposed into $O(1)$ weakly transversal collections \mathcal{Q}' of $d + 1$ cubes, each one having a cube $Q'_1 \subset Q_1$ as pivot.

To each such subcollection we apply the endpoint estimate from [Section 10](#) (all we need to apply it is weak transversality), which finishes the proof. \square

14. Further remarks

Remark 14.1. It was pointed out to us by Jonathan Bennett that the d -dimensional estimates (2) for tensors are equivalent to certain one-dimensional mixed norm bounds. We present this remark in the following proposition:

Proposition 14.2 (Bennett). *For all $p, q \geq 1$, the estimate*

$$\|\mathcal{E}_d g\|_{L_{\xi_1, \dots, \xi_{d+1}}^q} \lesssim \|g\|_p \tag{128}$$

holds for tensors $g(x) = g_1(x_1) \cdots g_d(x_d)$ if and only if

$$\|\mathcal{E}_1 f\|_{L_{\xi_2}^{dq} L_{\xi_1}^q} \lesssim \|f\|_p. \tag{129}$$

holds.

Proof. Assume first that (128) holds for tensors. Then

$$\begin{aligned} \|\mathcal{E}_1 f\|_{L_{\xi_2}^{dq} L_{\xi_1}^q} &= \left[\int \left[\int |\mathcal{E}_1 f(\xi_1, \xi_2)|^q d\xi_1 \right]^d d\xi_2 \right]^{\frac{1}{dq}} \\ &= \left[\int \prod_{j=1}^d \left[\int |\mathcal{E}_1 f(\eta_j, \xi_2)|^q d\eta_j \right] d\xi_2 \right]^{\frac{1}{dq}} \\ &= \left[\int \prod_{j=1}^d \int |\mathcal{E}_d(f \otimes \cdots \otimes f)(\eta_1, \dots, \eta_d)|^q d\vec{\eta} d\xi_2 \right]^{\frac{1}{dq}} \\ &= \|\mathcal{E}_d(f \otimes \cdots \otimes f)\|_q^{\frac{1}{d}} \lesssim \|f \otimes \cdots \otimes f\|_p^{\frac{1}{d}} \lesssim \|f\|_p, \end{aligned}$$

which proves (129). Conversely, assuming that (129) holds for all $f \in L^p([0, 1])$ yields

$$\begin{aligned} \|\mathcal{E}_d(g_1 \otimes \cdots \otimes g_d)\|_q^q &= \int |\mathcal{E}_1 g_1(\xi_1, \xi_{d+1})|^q \cdots |\mathcal{E}_1 g_d(\xi_d, \xi_{d+1})|^q d\xi_1 \cdots d\xi_{d+1} \\ &= \int \prod_{j=1}^d \left[\int |\mathcal{E}_1 g_j(\xi_j, \xi_{d+1})|^q d\xi_j \right] d\xi_{d+1} \\ &\leq \prod_{j=1}^d \left[\int \left[\int |\mathcal{E}_1 g_j(\xi_j, \xi_{d+1})|^q d\xi_j \right]^d d\xi_{d+1} \right]^{\frac{1}{d}} \\ &= \prod_j \|\mathcal{E}_1 g_j\|_{L_{\xi_{d+1}}^{dq} L_{\xi_j}^q}^q \lesssim \prod_{j=1}^d \|g_j\|_p^q = \|g\|_p^q. \quad \square \end{aligned}$$

Estimates such as (129) can be verified directly by interpolation. Taking sup in ξ_2 gives

$$\|\mathcal{E}_1 f\|_{L_{\xi_2}^\infty L_{\xi_1}^2} \lesssim_\varepsilon \|f\|_{L^2([0,1])}. \tag{130}$$

Conjecture 1.1 for $d = 1$ follows from

$$\|\mathcal{E}_1 f\|_{L_{\xi_2, \xi_1}^{4+\varepsilon}} \lesssim_\varepsilon \|f\|_{L^4([0,1])} \tag{131}$$

for all $\varepsilon > 0$. Using mixed-norm Riesz-Thorin interpolation with weights $\approx \frac{d-1}{d+1}$ for (130) and $\approx \frac{2}{d+1}$ for (131), one obtains (129) for $p = \frac{2(d+1)}{d}$ and $q = \frac{2(d+1)}{d} + \varepsilon'$, which shows (128) by the previous claim.

The reader will notice that our proof for the case $k = 1$ of Theorem 1.5 has a similar idea in its core: we interpolate (at the level of the sets $\mathbb{X}^{l_1, \dots, l_d}$) between two estimates similar to (130) and (131). On the other hand, we have not found an extension of Bennett’s remark to the case $2 \leq k \leq d + 1$, in which we still need to interpolate locally instead of globally and assume that only one function has a tensor structure.

Remark 14.3. In [Tao et al. 1998] the authors obtain the following off-diagonal type bounds:

Theorem [Tao et al. 1998]. $\mathcal{ME}_{2,d}$ satisfies

$$\begin{aligned} \|\mathcal{ME}_{2,d}(g_1, g_2)\|_2 &\lesssim \|g_1\|_2 \cdot \|g_2\|_{\frac{d+1}{d}}, \\ \|\mathcal{ME}_{2,d}(g_1, g_2)\|_2 &\lesssim \|f\|_{\frac{d+1}{d}} \cdot \|g\|_2. \end{aligned}$$

In general, under the extra hypothesis that either g_1 or g_2 is a full tensor, one can obtain all k -linear off-diagonal type bounds like $L^{p_1} \times \dots \times L^{p_k} \mapsto L^2$ by a straightforward adaptation of the argument presented in Section 9. We chose not to include them in this manuscript.

Remark 14.4. Under the assumption that g_j are full tensors

$$g_j(x_1, \dots, x_d) = g_{j,1}(x_1) \cdots g_{j,d}(x_d), \quad 1 \leq j \leq k,$$

the methods of this work allow to prove Conjecture 1.11. We will not cover the details of this result here, but the idea is simply to interpolate between the $p = 2$ result and the case $k = 1$ for tensors.

Appendix A: Sharp examples

The goal of this first appendix is to discuss the sharpness of Theorems 1.5 and 11.2. We remark that sharp examples already exist in the literature, notably in the context of the bilinear problem for the sphere in [Foschi and Klainerman 2000], and in the multilinear case for surfaces of any signature in [Hickman and Iliopoulou 2022]. Our examples, however, exploit different ideas than those present in those works in the sense that they are robust enough to address weakly transversal configurations of caps and give sharp results in such cases as well.

The first part of this appendix is about Theorem 11.2, whereas in the second one we prove that, to attain the sharp range of Conjecture 1.2 in general, transversality cannot be replaced by the concept of weak transversality that we introduce.

AA. Range optimality. The main result of this subsection is the following:

Proposition A.1. *The condition*

$$p \geq \frac{2(d + |\tau| + 2)}{k(d + |\tau|)}$$

is necessary for Theorem 11.2 to hold.

Our examples are constructed based on one-dimensional considerations. For the benefit of simplifying the notation, smoothing the exposition to the reader and to establish a clear link with Conjecture 1.2, we present them in the $|\tau| = k - 1$ case, which is the smallest possible value for the corresponding $|\tau|$ of a given collection of transversal cubes (up to decomposing it into weakly transversal collections, see Claim B.4). It will be clear, however, how to work out the general case of arbitrary $|\tau|$, and we will point that out along the proof of Claim A.3.

Consider the caps that project onto the following transversal domains via $x \mapsto |x|^2$:

$$U_1 = [0, 1]^d, \\ U_j = [2, 3]^{j-2} \times [4, 5] \times [0, 1]^{d-j+1}, \quad 2 \leq j \leq k.$$

Observe that these caps are transversal as well;⁵⁰ therefore the following argument for the case $|\tau| = k - 1$ of Proposition A.1 also shows that the range of Conjecture 1.2 is necessary.

⁵⁰For general $|\tau|$ we would have to start with a different collection of cubes with the appropriate total degree of transversality.




Figure 5. Cases $k = 3$ and $k = 4$ when $d = 3$.

We present the examples separately to distinguish their features. For $k = d + 1$ we will take appropriately placed cubes, whereas for $2 \leq k \leq d$ we will take slabs (boxes with edges of two different scales).

Claim A.2. Let $k = d + 1$, $\delta > 0$ small and let A_j^δ be given by

$$A_1^\delta = [0, \delta]^d,$$

$$A_j^\delta = [2, 2 + \delta]^{j-2} \times [4, 4 + \delta] \times [0, \delta]^{d-j+1}, \quad 2 \leq j \leq d + 1.$$

Define $f_j^\delta := \mathbb{1}_{A_j^\delta}$. Then

$$\frac{\|\prod_{j=1}^{d+1} \mathcal{E}_{U_j} f_j^\delta\|_p}{\prod_{j=1}^{d+1} \|f_j^\delta\|_2} \gtrsim \delta^{\frac{d(d+1)}{2} - \frac{1}{p}(d+1)}.$$

Therefore, letting $\delta \rightarrow 0$ implies $p \geq \frac{2}{d}$ is a necessary condition for the $(d + 1)$ -linear extension conjecture to hold for this choice of the U_j and for all f_j that are full tensors.

Claim A.3. Let $2 \leq k < d + 1$, $\delta > 0$ small and let B_j^δ be given by

$$B_1^\delta = [0, \delta^2]^{k-1} \times [0, \delta]^{d-k+1},$$

$$B_j^\delta = [2, 2 + \delta^2]^{j-2} \times [4, 4 + \delta^2] \times [0, \delta^2]^{k-j} \times [0, \delta]^{d-k+1}, \quad 2 \leq j \leq k.$$

Define $g_j^\delta := \mathbb{1}_{B_j^\delta}$. Then

$$\frac{\|\prod_{j=1}^k \mathcal{E}_{U_j} g_j^\delta\|_p}{\prod_{j=1}^k \|g_j^\delta\|_2} \gtrsim \delta^{\frac{k}{2}(d+k-1) - \frac{1}{p}(d+k+1)}.$$

Therefore, letting $\delta \rightarrow 0$ implies

$$p \geq \frac{2(d + k + 1)}{k(d + k - 1)}$$

is a necessary condition for the k -linear extension conjecture to hold for this choice of the U_j and for all g_j that are full tensors.

Before proving the claims, we need the following lemma:

Lemma A.4 (scale-1 phase-space portrait of $e^{2\pi i x^2}$). *There exists a sequence of smooth bumps $(\varphi_n)_{n \in \mathbb{Z}}$ such that*

- (i) $\text{supp}(\varphi_n) \subset [n - 1, n + 1]$, $n \in \mathbb{Z}$,
- (ii) $|\varphi_n^{(\ell)}(x)| \leq C_\ell$ uniformly in $n \in \mathbb{Z}$ and such that

$$e^{2\pi i x^2} = \sum_{n \in \mathbb{Z}} e^{4\pi i n x} \varphi_n(x).$$

Proof. See [Muscalu and Schlag 2013b, Proposition 1.10, page 23]. □

Rescaling with $t > 0$, the corresponding phase space portrait of $e^{2\pi i t x^2}$ is

$$e^{2\pi i t x^2} = e^{2\pi i (\sqrt{t}x)^2} = \sum_{n \in \mathbb{Z}} e^{4\pi i n \sqrt{t}x} \varphi_n(\sqrt{t}x).$$

Observe that $\tilde{\varphi}_t(x) = \varphi_n(\sqrt{t}x)$ is adapted to the Heisenberg box $[\frac{n}{\sqrt{t}}, \frac{n+1}{\sqrt{t}}] \times [0, \sqrt{t}]$, but strictly supported on $[\frac{n-1}{\sqrt{t}}, \frac{n+1}{\sqrt{t}}]$. This way, we can write

$$e^{2\pi i t x^2} = \sum_{n \in \mathbb{Z}} \Phi_{n,t}(x), \tag{132}$$

where $\Phi_{n,t}$ is adapted to the Heisenberg box $[\frac{n}{\sqrt{t}}, \frac{n+1}{\sqrt{t}}] \times [2n\sqrt{t}, (2n+1)\sqrt{t}]$.

Proof of Claim A.2. Motivated by the uncertainty principle, the first step is to analyze the behavior of the extension operator \mathcal{E}_{U_j} applied to f_j^δ on a box whose sizes are reciprocal to the ones of $\text{supp}(f_j^\delta)$. More precisely, we will show that $|\mathcal{E}_{U_j}(f_j^\delta)| \gtrsim \delta^d$ on such boxes.

If $\delta < \frac{1}{\sqrt{t}}$,

$$\begin{aligned} \mathcal{E}_{U_1}(f_1^\delta)(\xi_1, \dots, \xi_d, t) &= \prod_{j=1}^d \left[\int_0^\delta e^{-2\pi i \xi_j x_j} e^{-2\pi i t x_j^2} dx_j \right] \\ &= \prod_{j=1}^d \left[\int_0^\delta e^{-2\pi i \xi_j x_j} \cdot [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right], \end{aligned}$$

since $\text{supp}(\Phi_{n,t}) \cap [0, \delta] = \emptyset$ if $n \in \mathbb{Z} \setminus \{0, 1\}$. If $|\xi_j x_j| < \frac{1}{N}$ (N is a big number to be chosen later), we then have

$$\begin{aligned} |\mathcal{E}_{U_1}(f_1^\delta)(\xi_1, \dots, \xi_d, t)| &= \prod_{j=1}^d \left| \int_0^\delta e^{-2\pi i \xi_j x_j} \cdot [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right|, \\ &\geq \prod_{j=1}^d \left(\left| \int_0^\delta [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right| - \left| \int_0^\delta [e^{-2\pi i \xi_j x_j} - 1] \cdot [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right| \right), \tag{133} \end{aligned}$$

where N is picked so that $[e^{-2\pi i \xi_j x_j} - 1]$ is close enough to zero to make

$$A_j := \left| \int_0^\delta [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right|$$

dominate each factor above. Since $A_j \gtrsim \delta$ (recall that $\Phi_{0,t}$ and $\Phi_{1,t}$ are adapted to Heisenberg boxes of size $\frac{1}{\sqrt{t}} \times \sqrt{t}$ and $\delta < \frac{1}{\sqrt{t}}$), we conclude that if $|\xi_j| \lesssim \frac{1}{\delta}$ for $1 \leq j \leq d$ and $|t| < \frac{1}{\delta^2}$, then

$$|\mathcal{E}_{U_1}(f_1^\delta)(\xi_1, \dots, \xi_d, t)| \geq \delta^d.$$

If ϕ is a bump supported on $[-1, 1]$, we have just proved that

$$|\mathcal{E}_{U_1}(f_1^\delta)(\xi_1, \dots, \xi_d, t)| \gtrsim \delta^d \phi_\delta(\xi_1) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t), \tag{134}$$

where $\phi_\delta(\xi) := \phi(\delta x)$. Analogously, if $\delta < \frac{1}{\sqrt{t}}$,

$$\begin{aligned} \mathcal{E}_{U_2}(f_2^\delta)(\xi_1, \dots, \xi_d, t) &= \left[\int_4^{4+\delta} e^{-2\pi i \xi_1 x_1} e^{-2\pi i t x_1^2} dx_1 \right] \cdot \prod_{j=2}^d \left[\int_0^\delta e^{-2\pi i \xi_j x_j} e^{-2\pi i t x_j^2} dx_j \right] \\ &= \underbrace{\left[\int_4^{4+\delta} e^{-2\pi i \xi_1 x_1} \left(\sum_{n \in \mathbb{Z}} \Phi_{n,t}(x_1) \right) dx_1 \right]}_{I_1} \cdot \prod_{j=2}^d \underbrace{\left[\int_0^\delta e^{-2\pi i \xi_j x_j} \cdot [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right]}_{I_j}. \end{aligned}$$

There are at most $O(1)$ integers n such that $\text{supp}(\Phi_{n,t}) \cap [4, 4 + \delta] \neq \emptyset$, and they cluster around $[4\sqrt{t}]$. Without loss of generality, one can assume that $n = 4\sqrt{t}$ so that the main contribution for I_1 comes from $\Phi_{4\sqrt{t},t}$ whose Heisenberg box is $[4, 4 + \frac{1}{\sqrt{t}}] \times [8t, 8t + \sqrt{t}]$. The modulation $e^{-2\pi i \xi_1 x_1}$ shifts this box vertically by $-\xi_1$, and I_1 is negligible if the boxes $[4, 4 + \frac{1}{\sqrt{t}}] \times [8t - \xi_1, 8t + \sqrt{t} - \xi_1]$ and $[0, \delta] \times [0, \frac{1}{\delta}]$ are disjoint in frequency, so we need $|\xi_1 - 8t| \lesssim \frac{1}{\delta}$ to have a significant contribution to I_1 . In that case,

$$|I_1| \gtrsim \left| \int_4^{4+\delta} e^{-2\pi i \xi_1 x_1} \Phi_{4\sqrt{t},t}(x_1) dx_1 \right| \gtrsim \delta.$$

The analysis of I_j for $j \geq 2$ is the same as the one for the factors of $\mathcal{E}_{U_1}(f_1^\delta)$. We conclude that if $|\xi_1 - 8t| \lesssim \frac{1}{\delta}$, $|\xi_j| \lesssim \frac{1}{\delta}$ for $2 \leq j \leq d$ and $|t| \leq \frac{1}{\delta^2}$, then

$$|\mathcal{E}_{U_2}(f_2^\delta)(\xi_1, \dots, \xi_d, t)| \geq \delta^d.$$

As before,

$$|\mathcal{E}_{U_2}(f_2^\delta)(\xi_1, \dots, \xi_d, t)| \gtrsim \delta^d \phi_\delta(\xi_1 - 8t) \cdot \phi_\delta(\xi_2) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t).$$

The extensions $\mathcal{E}_{U_j}(f_j^\delta)$ for $3 \leq j \leq d + 1$ are treated in the same way we treated $\mathcal{E}_{U_2}(f_2^\delta)$. The conclusion is that

$$\begin{aligned} |\mathcal{E}_{U_j}(f_j^\delta)(\xi_1, \dots, \xi_d, t)| &\gtrsim \delta^d \phi_\delta(\xi_1 - 4t) \cdots \phi_\delta(\xi_{j-2} - 4t) \cdot \phi_\delta(\xi_{j-1} - 8t) \cdot \phi_\delta(\xi_j) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t) \tag{135} \end{aligned}$$

for all $2 \leq j \leq d + 1$.

Let $\xi = (\xi_1, \dots, \xi_d)$. From (134) and (135) we obtain

$$\prod_{j=1}^{d+1} |\mathcal{E}_{U_j}(f_j^\delta)(\xi, t)| \gtrsim \delta^{d(d+1)} \left[\phi_{\delta^2}(t) \prod_{l=1}^d \phi_\delta(\xi_l) \right] \times \left[\prod_{j=2}^d \phi_\delta(\xi_{j-1} - 4t) \cdots \phi_\delta(\xi_{j-2} - 4t) \cdot \phi_\delta(\xi_{j-1} - 8t) \cdot \phi_\delta(\xi_j) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t) \right]. \tag{136}$$

Now we analyze the support of the product of the right-hand side of (136). Notice that we have at least one bump like $\phi_\delta(\xi_j)$ for every $1 \leq j \leq d + 1$, so $|\xi_j| \lesssim \frac{1}{\delta}$ is a necessary condition for the product not to be zero. On the other hand, the conditions

$$|\xi_j| \lesssim \frac{1}{\delta}, \quad |\xi_j - 8t| \lesssim \frac{1}{\delta}$$

together imply $|t| \lesssim \frac{1}{\delta}$, which is much more restrictive than the $|t| \lesssim \frac{1}{\delta^2}$ that comes from the support of the bump $\phi_{\delta^2}(t)$. We conclude that the right-hand side of (136) is supported on the box

$$R_\delta^* = \left\{ (\xi_1, \dots, \xi_d, t) \in \mathbb{R}^{d+1} : |t| \lesssim \frac{1}{\delta}, |\xi_j| \lesssim \frac{1}{\delta}, 1 \leq j \leq d \right\}.$$

Finally,

$$\frac{\|\prod_{j=1}^{d+1} \mathcal{E}_{U_j} f_j^\delta\|_p}{\prod_{j=1}^{d+1} \|f_j^\delta\|_2} \gtrsim \frac{\delta^{d(d+1)} \cdot |R_\delta^*|^{\frac{1}{p}}}{\delta^{\frac{d(d+1)}{2}}} \gtrsim \delta^{\frac{d(d+1)}{2} - \frac{1}{p}(d+1)} \tag{137}$$

and the claim follows. □

Proof of Claim A.3. The outline of the following argument is the same as the one used in previous proof. Let $\xi = (\xi_1, \dots, \xi_d)$. If $\delta^2 < \frac{1}{\sqrt{t}}$,

$$\begin{aligned} \mathcal{E}_{U_1}(g_1^\delta)(\xi, t) &= \prod_{j=1}^{k-1} \left[\int_0^{\delta^2} e^{-2\pi i \xi_j x_j} e^{-2\pi i t x_j^2} dx_j \right] \cdot \prod_{l=k}^d \left[\int_0^\delta e^{-2\pi i \xi_l x_l} e^{-2\pi i t x_l^2} dx_l \right] \\ &= \prod_{j=1}^{k-1} \left[\int_0^{\delta^2} e^{-2\pi i \xi_j x_j} [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right] \cdot \underbrace{\prod_{l=k}^d \left[\int_0^\delta e^{-2\pi i \xi_l x_l} \left(\sum_{n \in \mathbb{Z}} \Phi_{n,t}(x_l) \right) dx_l \right]}_{(*)}, \end{aligned}$$

since $\text{supp}(\Phi_{n,t}) \cap [0, \delta^2] = \emptyset$ if $n \in \mathbb{Z} \setminus \{0, 1\}$. If $\delta < \frac{1}{\sqrt{t}}$ (which is stronger than the previous condition $\delta^2 < \frac{1}{\sqrt{t}}$), we can eliminate most $\Phi_{n,t}$ in (*) as well:

$$\begin{aligned} \mathcal{E}_{U_1}(g_1^\delta)(\xi, t) &= \prod_{j=1}^{k-1} \left[\int_0^{\delta^2} e^{-2\pi i \xi_j x_j} [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right] \cdot \prod_{l=k}^d \left[\int_0^\delta e^{-2\pi i \xi_l x_l} \cdot [\Phi_{0,t}(x_l) + \Phi_{1,t}(x_l)] dx_l \right], \end{aligned}$$

If $|\xi_j x_j| < \frac{1}{N}$ (for N big enough), we then have

$$\begin{aligned}
 & |\mathcal{E}_{U_1}(g_1^\delta)(\xi, t)| \\
 &= \prod_{j=1}^{k-1} \left| \int_0^{\delta^2} e^{-2\pi i \xi_j x_j} [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right| \cdot \prod_{l=k}^d \left| \int_0^\delta e^{-2\pi i \xi_l x_l} \cdot [\Phi_{0,t}(x_l) + \Phi_{1,t}(x_l)] dx_l \right|, \\
 &\gtrsim \delta^{2(k-1)+(d-k+1)} = \delta^{d+k-1},
 \end{aligned}$$

by the same argument presented when we analyzed (133). We conclude that if $|\xi_j| \lesssim \frac{1}{\delta^2}$ for $1 \leq j \leq k-1$, $|\xi_l| \lesssim \frac{1}{\delta}$ for $k \leq l \leq d$ and $|t| < \frac{1}{\delta^2}$, then⁵¹

$$|\mathcal{E}_{U_1}(g_1^\delta)(\xi, t)| \gtrsim \delta^{d+k-1}.$$

Using the same notation from the proof of Claim A.2, we have just proved that

$$|\mathcal{E}_{U_1}(g_1^\delta)(\xi, t)| \gtrsim \delta^d \phi_{\delta^2}(\xi_1) \cdots \phi_{\delta^2}(\xi_{k-1}) \phi_\delta(x_k) \cdots \phi_\delta(x_d) \cdot \phi_{\delta^2}(t), \tag{138}$$

where $\phi_\delta(\xi) := \phi(\delta\xi)$ and ϕ is a bump supported on $[-1, 1]$. Analogously, if $\delta < \frac{1}{\sqrt{t}}$,

$$\begin{aligned}
 & \mathcal{E}_{U_2}(g_2^\delta)(\xi, t) \\
 &= \left[\int_4^{4+\delta^2} e^{-2\pi i \xi_1 x_1} e^{-2\pi i t x_1^2} dx_1 \right] \cdot \prod_{j=2}^{k-1} \left[\int_0^{\delta^2} e^{-2\pi i \xi_j x_j} e^{-2\pi i t x_j^2} dx_j \right] \cdot \prod_{l=k}^d \left[\int_0^\delta e^{-2\pi i \xi_l x_l} e^{-2\pi i t x_l^2} dx_l \right] \\
 &= \underbrace{\left[\int_4^{4+\delta^2} e^{-2\pi i \xi_1 x_1} \left(\sum_{n \in \mathbb{Z}} \Phi_{n,t}(x_1) \right) dx_1 \right]}_{M_1} \cdot \prod_{j=2}^{k-1} \underbrace{\left[\int_0^{\delta^2} e^{-2\pi i \xi_j x_j} \cdot [\Phi_{0,t}(x_j) + \Phi_{1,t}(x_j)] dx_j \right]}_{M_j} \\
 &\quad \times \prod_{l=k}^d \underbrace{\left[\int_0^\delta e^{-2\pi i \xi_l x_l} \cdot [\Phi_{0,t}(x_l) + \Phi_{1,t}(x_l)] dx_l \right]}_{M_l}. \tag{139}
 \end{aligned}$$

As in the proof of Claim A.2, the main contribution for M_1 comes from $\Phi_{4\sqrt{t},t}$, whose Heisenberg box is $[4, 4 + \frac{1}{\sqrt{t}}] \times [8t, 8t + \sqrt{t}]$. The modulation $e^{-2\pi i \xi_1 x_1}$ shifts this box vertically by $-\xi_1$, and M_1 is negligible if the boxes $[4, 4 + \frac{1}{\sqrt{t}}] \times [8t - \xi_1, 8t + \sqrt{t} - \xi_1]$ and $[0, \delta^2] \times [0, \frac{1}{\delta^2}]$ are disjoint in frequency, so we need $|\xi_1 - 8t| \lesssim \frac{1}{\delta^2}$ to have a significant contribution to M_1 . In that case,

$$|M_1| \gtrsim \left| \int_4^{4+\delta^2} e^{-2\pi i \xi_1 x_1} \Phi_{2\sqrt{t},t}(x_1) dx_1 \right| \gtrsim \delta^2.$$

The analyses of M_j for $2 \leq j \leq k-1$ and of M_l for $k \leq l \leq d-k+1$ are the same as the one for the factors of $\mathcal{E}_{U_1}(g_1^\delta)$. We conclude that if $|\xi_1 - 8t| \lesssim \frac{1}{\delta^2}$, $|\xi_j| \lesssim \frac{1}{\delta^2}$ for $2 \leq j \leq k-1$, $|\xi_l| \lesssim \frac{1}{\delta}$ for $k \leq l \leq d$ and $|t| \leq \frac{1}{\delta^2}$, then

$$|\mathcal{E}_{U_2}(g_2^\delta)(\xi, t)| \geq \delta^{d+k-1}.$$

⁵¹For general $|\tau|$, we would have $|\tau|$ conditions of type $|\xi_j| \lesssim \frac{1}{\delta^2}$ and $(d - |\tau|)$ like $|\xi_l| \lesssim \frac{1}{\delta}$.

As before,

$$|\mathcal{E}_{U_2}(g_2^\delta)(\xi, t)| \gtrsim \delta^d \phi_\delta(\xi_1 - 8t) \cdot \phi_{\delta^2}(\xi_2) \cdots \phi_{\delta^2}(\xi_{k-1}) \cdot \phi_\delta(\xi_k) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t).$$

The extensions $\mathcal{E}_{U_j}(g_j^\delta)$ for $3 \leq j \leq k$ are treated in the same way. The conclusion is that

$$|\mathcal{E}_{U_j}(g_j^\delta)(\xi, t)| \gtrsim \delta^d \phi_\delta(\xi_1 - 4t) \cdots \phi_\delta(\xi_{j-2} - 4t) \cdot \phi_\delta(\xi_{j-1} - 8t) \cdot \phi_\delta(\xi_j) \cdots \phi_\delta(\xi_d) \phi_{\delta^2}(t) \quad (140)$$

for all $2 \leq j \leq k$. From (138) and (140) we obtain

$$\begin{aligned} & \prod_{j=1}^k |\mathcal{E}_{U_j}(g_j^\delta)(\xi, t)| \\ & \gtrsim \delta^{k(d+k-1)} \left[\phi_{\delta^2}(t) \prod_{l=1}^{k-1} \phi_{\delta^2}(\xi_l) \cdot \prod_{n=k}^d \phi_\delta(\xi_n) \right] \\ & \times \left[\prod_{j=2}^d \left(\prod_{n=1}^{j-2} \phi_{\delta^2}(\xi_n - 4t) \right) \cdot \phi_{\delta^2}(\xi_{j-1} - 8t) \cdot \left(\prod_{m=j}^{k-1} \phi_{\delta^2}(\xi_m) \right) \cdot \left(\prod_{r=k}^d \phi_\delta(\xi_r) \right) \cdot \phi_{\delta^2}(t) \right]. \quad (141) \end{aligned}$$

Notice that we have at least one bump like $\phi_{\delta^2}(\xi_j)$ for every $1 \leq j \leq k - 1$ and at least one $\phi_\delta(\xi_l)$ for $k \leq l \leq d$, so $|\xi_j| \lesssim \frac{1}{\delta^2}$ and $|\xi_l| \lesssim \frac{1}{\delta}$ are necessary conditions for the product not to be zero. On the other hand, the conditions

$$|\xi_j| \lesssim \frac{1}{\delta^2}, \quad |\xi_j - 8t| \lesssim \frac{1}{\delta^2}$$

together imply $|t| \lesssim \frac{1}{\delta^2}$, which does not add any new information compared to the one coming from the bump $\phi_{\delta^2}(t)$ (this is the main difference between the analysis in Claims A.2 and A.3). We conclude that the right-hand side of (141) is supported on the box

$$S_\delta^* = \left\{ (\xi_1, \dots, \xi_d, t) \in \mathbb{R}^{d+1} : |t| \lesssim \frac{1}{\delta^2}; \quad |\xi_j| \lesssim \frac{1}{\delta^2}, \quad 1 \leq j \leq k - 1; \quad |\xi_l| \lesssim \frac{1}{\delta}, \quad k \leq l \leq d \right\}.$$

Finally,

$$\frac{\|\prod_{j=1}^k \mathcal{E}_{U_j} g_j^\delta\|_p}{\prod_{j=1}^{d+1} \|g_j^\delta\|_2} \gtrsim \frac{\delta^{(d+k-1)k} \cdot |S_\delta^*|^{\frac{1}{p}}}{\delta^{\frac{(d+k-1)k}{2}}} \gtrsim \delta^{\frac{(d+k-1)k}{2} - \frac{(d+k+1)}{p}} \quad (142)$$

and the claim follows. □

AB. Transversality as a necessary condition in general. A natural question is: given k cubes U_j , $1 \leq j \leq k$, is it possible to prove

$$\left\| \prod_{j=1}^k \mathcal{E}_{U_j} g_j \right\|_p \lesssim \prod_{j=1}^k \|g_j\|_2$$

for

$$p \geq \frac{2(d+k+1)}{k(d+k-1)}$$

and all $g_j \in L^2(U_j)$ if the U_j are assumed to be weakly transversal?

The answer is no and we will address it in this second part of the first appendix. As a consequence, we conclude that [Theorem 1.5](#) is sharp under weak transversality, as observed in [Remark 1.8](#).

We will treat the case $k = 3$ and $d = 2$ for simplicity, but a similar construction holds in general. If three boxes $U_1, U_2, U_3 \subset \mathbb{R}^2$ are not transversal, there is a line that crosses them. Assume without loss of generality that $U_1 = [0, 1]^2$, $U_2 = [2, 3]^2$ and $U_3 = [4, 5]^2$. We will show that

$$\|E_{U_1}(h_1) \cdot E_{U_2}(h_2) \cdot E_{U_3}(h_3)\|_p \lesssim \|h_1\|_2 \cdot \|h_2\|_2 \cdot \|h_3\|_2$$

only if $p \geq \frac{10}{9}$. The trilinear extension conjecture for $d = 2$ states that $p \geq 1$ is the sharp range under the transversality hypothesis.

Claim A.5. Define the sets D_j^δ by

$$\begin{aligned} D_1^\delta &= \left[\frac{\sqrt{2} - \delta^2}{2}, \frac{\sqrt{2} + \delta^2}{2} \right] \times \left[-\frac{\delta}{2}, \frac{\delta}{2} \right], \\ D_2^\delta &= \left[\frac{5\sqrt{2} - \delta^2}{2}, \frac{5\sqrt{2} + \delta^2}{2} \right] \times \left[-\frac{\delta}{2}, \frac{\delta}{2} \right], \\ D_3^\delta &= \left[\frac{9\sqrt{2} - \delta^2}{2}, \frac{9\sqrt{2} + \delta^2}{2} \right] \times \left[-\frac{\delta}{2}, \frac{\delta}{2} \right]. \end{aligned}$$

Define $h_j^\delta := \mathbb{1}_{D_j^\delta}$. Then

$$\frac{\|\prod_{j=1}^3 \mathcal{E}_{D_j} h_j^\delta\|_p}{\prod_{j=1}^3 \|h_j^\delta\|_2} \gtrsim \delta^{\frac{9}{2} - \frac{5}{p}}.$$

Proof. The proof is analogous to the ones of [Claims A.2](#) and [A.3](#). □

Let the rhombuses \tilde{D}_j be given by

$$\begin{aligned} \tilde{D}_1 &= \text{Conv} \left((0, 0); \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right); \left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right); (\sqrt{2}, 0) \right), \\ \tilde{D}_2 &= \text{Conv} \left((2\sqrt{2}, 0); \left(\frac{5\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right); \left(\frac{5\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right); (3\sqrt{2}, 0) \right), \\ \tilde{D}_3 &= \text{Conv} \left((4\sqrt{2}, 0); \left(\frac{9\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right); \left(\frac{9\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right); (5\sqrt{2}, 0) \right). \end{aligned}$$

Observe that $D_j^\delta \subset \tilde{D}_j$ for $\delta > 0$ small enough. Extend the domain of h_j^δ to \tilde{D}_j so that it is 0 on $\tilde{D}_j \setminus D_j^\delta$. Let T be a $\frac{\pi}{4}$ counterclockwise rotation and let

$$H_j^\delta(x) := h_j^\delta \circ T^{-1}(x).$$

Notice that T takes \tilde{D}_j to U_j , as shown in the picture below.

Since L^p norms are invariant under rotations, we have

$$\frac{\|\prod_{j=1}^3 \mathcal{E}_{U_j} H_j^\delta\|_p}{\prod_{j=1}^3 \|H_j^\delta\|_2} \gtrsim \delta^{\frac{9}{2} - \frac{5}{p}}$$




Figure 6. Left: The function h_j^δ . Right: H_j^δ is supported on U_j .

from Claim A.5. Letting $\delta \rightarrow 0$ shows that we need $p \geq \frac{10}{9}$, so the sharp range $p \geq 1$ cannot be obtained if the boxes U_1, U_2, U_3 are not transversal.

Remark A.6. As expected, the functions H_j^δ do not have a tensor structure with respect to the canonical basis. If this was the case, our methods would have allowed us to prove that the corresponding trilinear extension operator maps $L^2 \times L^2 \times L^2$ to L^1 .

Appendix B: Technical results

Here we collect a few technical results used throughout the paper.

Theorem B.1. For $0 < \gamma < d$, $1 < p < q < \infty$, and $\frac{1}{q} = \frac{1}{p} - \frac{d-\gamma}{d}$, we have

$$\|f * (|y|^{-\gamma})\|_{L^q(\mathbb{R}^d)} \leq A_{p,q} \cdot \|f\|_{L^p(\mathbb{R}^d)}. \tag{143}$$

Proof. See Proposition 7.8 in [Muscalu and Schlag 2013a]. □

Theorem B.2 (nonstationary phase). Let $a \in C_0^\infty$ and

$$I(\lambda) = \int_{\mathbb{R}^d} e^{2\pi i \lambda \phi(\xi)} a(\xi) \, d\xi.$$

If $\nabla \phi \neq 0$ on $\text{supp}(a)$, then

$$|I(\lambda)| \leq C(N, a, \phi) \lambda^{-N}$$

as $\lambda \rightarrow \infty$ for arbitrary $N \geq 1$.

Proof. See Lemma 4.14 in [Muscalu and Schlag 2013a]. □

Theorem B.3 (stationary phase). If $\nabla \phi(\xi_0) = 0$ for some $\xi_0 \in \text{supp}(a)$, $\nabla \phi \neq 0$ away from ξ_0 and the Hessian of ϕ at the stationary point ξ_0 is nondegenerate, i.e., $\det D^2 \phi(\xi_0) \neq 0$, then for all $\lambda \geq 1$

$$|I(\lambda)| \leq C(N, a, \phi) \lambda^{-\frac{d}{2}}.$$

Proof. See Lemma 4.15 in [Muscalu and Schlag 2013a]. □

We now restate and prove the main claim from [Section 3](#):

Claim B.4. *Given a collection $\mathcal{Q} = \{Q_1, \dots, Q_k\}$ of transversal cubes, each $Q_l \in \mathcal{Q}$ can be partitioned into $O(1)$ many subcubes*

$$Q_l = \bigcup_i Q_{l,i}$$

so that all collections $\tilde{\mathcal{Q}}$ made of picking one subcube $Q_{l,i}$ per Q_l

$$\tilde{\mathcal{Q}} = \{\tilde{Q}_1, \dots, \tilde{Q}_k\}, \quad \tilde{Q}_l \in \{Q_{l,i}\}_i,$$

are weakly transversal.

Proof. For each $1 \leq j \leq d$, consider the set A_j of endpoints of the intervals $\pi_j(Q_1), \dots, \pi_j(Q_k)$. Using these endpoints to partition this collection of intervals, one can assume that there are three cases for two cubes Q_r and Q_s :

- (1) $\pi_j(Q_r) \cap \pi_j(Q_s) = \emptyset$.
- (2) $\pi_j(Q_r) = \pi_j(Q_s)$.
- (3) $\pi_j(Q_r) \cap \pi_j(Q_s) = \{p_{r,s}\}$, where $p_{r,s}$ is an endpoint of both $\pi_j(Q_r)$ and $\pi_j(Q_s)$.

We can go one step further and assume that all $\pi_j(Q_s)$ that intersect a given $\pi_j(Q_r)$ (but distinct from it) do so at the same endpoint. Indeed, if $\pi_j(Q_{s_1}) \cap \pi_j(Q_r) = \{p\}$, $\pi_j(Q_{s_2}) \cap \pi_j(Q_r) = \{q\}$ and $\pi_j(Q_r) = [p, q]$, we can simply split $\pi_j(Q_r)$ in half and obtain intervals that satisfy this property.

Now we choose a point $x_{j,r}$ in every interval $\pi_j(Q_r)$:

- (1) If $\pi_j(Q_r) \cap \pi_j(Q_s) = \emptyset$ for all $s \neq r$, let $x_{j,r}$ be $c_{j,r}$, the center of $\pi_j(Q_r)$.
- (2) If $\pi_j(Q_r)$ intersects some $\pi_j(Q_{s_1})$ at p , any other $\pi_j(Q_{s_2})$ that intersects $\pi_j(Q_r)$ also does it at p . In this case choose $x_{j,r} = x_{j,s} = p$ for all s such that $\pi_j(Q_r) \cap \pi_j(Q_s) \neq \emptyset$.

Let us now show that, after the reductions above, the transversal set of cubes \mathcal{Q} is weakly transversal. More precisely, for a fixed $1 \leq l \leq k$, we will show that there is a set of $k-1$ canonical directions that together with Q_l satisfy (15). Let $\vec{x}_i \in Q_i$ for $1 \leq i \leq k$ be given in coordinates by

$$\vec{x}_i = (x_{1,i}, x_{2,i}, \dots, x_{d,i}).$$

The normal vector to \mathbb{P}^d at \vec{x}_i is

$$\vec{v}_i = (-2x_{1,i}, -2x_{2,i}, \dots, -2x_{d,i}, 1).$$

Then the cubes in \mathcal{Q} are transversal if and only if the matrix

$$\begin{pmatrix} -2x_{1,1} & -2x_{1,2} & \cdots & -2x_{1,k} \\ -2x_{2,1} & -2x_{2,2} & \cdots & -2x_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ -2x_{d,1} & -2x_{d,2} & \cdots & -2x_{d,k} \\ 1 & 1 & \cdots & 1 \end{pmatrix}$$

has rank k for all $x_{j,i} \in \pi_j(Q_i)$, $1 \leq j \leq d$, $1 \leq i \leq k$.

By [Lemma B.5](#) (proven at the end of this appendix), there are $k-1$ rows

$$R_{i_n} = (-2x_{i_n,1}, \dots, -2x_{i_n,k})$$

of the above matrix, $1 \leq n \leq k-1$, such that

$$\begin{cases} x_{i_1,l} \neq x_{i_1,1} \\ \vdots \\ x_{i_{l-1},l} \neq x_{i_{l-1},l-1} \\ x_{i_l,l} \neq x_{i_l,l+1} \\ \vdots \\ x_{i_{k-1},l} \neq x_{i_{k-1},k}. \end{cases}$$

Because of the choices we made, $x_{i_n,l} \neq x_{i_n,r}$ implies

$$\pi_{i_n}(Q_l) \cap \pi_{i_n}(Q_r) = \emptyset,$$

which finishes the proof. □

Finally, we state and prove the auxiliary linear algebra lemma used in the proof of [Claim B.4](#).

Lemma B.5. *Let M be the $(d+1) \times k$ matrix*

$$M = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,k} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{d,1} & a_{d,2} & \cdots & a_{d,k} \\ 1 & 1 & \cdots & 1 \end{pmatrix}$$

and assume that it has rank k . For each column $C_j = (a_{1,j}, \dots, a_{d,j}, 1)$ there are $k-1$ rows $R_{i_l} = (a_{i_l,1}, \dots, a_{i_l,k})$, $1 \leq l \leq k-1$, such that

$$\begin{cases} a_{i_1,j} \neq a_{i_1,l_1} \\ a_{i_2,j} \neq a_{i_2,l_2} \\ \vdots \\ a_{i_{k-1},j} \neq a_{i_{k-1},l_{k-1}}, \end{cases}$$

where $(l_1, l_2, \dots, l_{k-1})$ is some permutation of $(1, 2, \dots, j-1, j+1, \dots, k)$.

Proof. Let us first consider the case $k = d+1$. We have to show that for all columns C_j the first $k-1$ rows satisfy the property of the lemma. Observe that the product

$$MA = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,k} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{k-1,1} & a_{k-1,2} & \cdots & a_{k-1,k} \\ 1 & 1 & \cdots & 1 \end{pmatrix} \cdot \underbrace{\begin{pmatrix} 1 & 1 & \cdots & 1 & 1 & 1 \\ -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & 0 \\ 0 & 0 & \cdots & 0 & -1 & 0 \end{pmatrix}}_{k \times k \text{ matrix } A}$$

is a rank k matrix equal to

$$\begin{pmatrix} (a_{1,1} - a_{1,2}) & (a_{1,1} - a_{1,3}) & \cdots & (a_{1,1} - a_{1,k-1}) & (a_{1,1} - a_{1,k}) & a_{1,1} \\ (a_{2,1} - a_{2,2}) & (a_{2,1} - a_{2,3}) & \cdots & (a_{2,1} - a_{2,k-1}) & (a_{2,1} - a_{2,k}) & a_{2,1} \\ (a_{3,1} - a_{3,2}) & (a_{3,1} - a_{3,3}) & \cdots & (a_{3,1} - a_{3,k-1}) & (a_{3,1} - a_{3,k}) & a_{3,1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ (a_{k-1,1} - a_{k-1,2}) & (a_{k-1,1} - a_{k-1,3}) & \cdots & (a_{k-1,1} - a_{k-1,k-1}) & (a_{k-1,1} - a_{k-1,k}) & a_{k-1,1} \\ 0 & 0 & \cdots & 0 & 0 & 1 \end{pmatrix}.$$

By computing the Laplace expansion with respect to the last row, we conclude that $\det(MA)$ is equal to

$$\det \begin{pmatrix} (a_{1,1} - a_{1,2}) & (a_{1,1} - a_{1,3}) & \cdots & (a_{1,1} - a_{1,k-1}) & (a_{1,1} - a_{1,k}) \\ (a_{2,1} - a_{2,2}) & (a_{2,1} - a_{2,3}) & \cdots & (a_{2,1} - a_{2,k-1}) & (a_{2,1} - a_{2,k}) \\ (a_{3,1} - a_{3,2}) & (a_{3,1} - a_{3,3}) & \cdots & (a_{3,1} - a_{3,k-1}) & (a_{3,1} - a_{3,k}) \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (a_{k-1,1} - a_{k-1,2}) & (a_{k-1,1} - a_{k-1,3}) & \cdots & (a_{k-1,1} - a_{k-1,k-1}) & (a_{k-1,1} - a_{k-1,k}) \end{pmatrix}.$$

The entries of this matrix are

$$x_{i,j} := a_{i,1} - a_{i,j+1}, \quad 1 \leq i, j \leq k - 1.$$

The column C_1 has the property of the lemma if and only if there is some permutation π of $(1, 2, \dots, k - 1)$ such that

$$\begin{cases} x_{1,\pi(1)} = a_{1,1} - a_{1,\pi(1)+1} \neq 0 \\ x_{2,\pi(2)} = a_{2,1} - a_{2,\pi(2)+1} \neq 0 \\ \vdots \\ x_{k-1,\pi(k-1)} = a_{k-1,1} - a_{k-1,\pi(k-1)+1} \neq 0. \end{cases}$$

If this was not the case, for all such permutations π of $(1, 2, \dots, k - 1)$ at least one among $x_{1,\pi(1)}, x_{2,\pi(2)}, \dots, x_{k-1,\pi(k-1)}$ would be zero. Hence

$$\det(MA) = \sum_{\pi \in S_{k-1}} \text{sgn}(\pi) \cdot x_{1,\pi(1)} \cdots x_{k-1,\pi(k-1)} = 0,$$

a contradiction. A similar argument shows that any other column also has this property.

The case $k < d + 1$ can be reduced to the previous one. Indeed, the rank- k condition guarantees that there is a $k \times k$ minor of M that has rank k . There are two possibilities:

(1) *There is a $k \times k$ minor of rank k that has a row of 1s.* This is identical to the case $k = d + 1$ and we conclude that the rows that generate this minor are the ones that satisfy the property of the lemma.

(2) No $k \times k$ minor of rank k has a row of 1's. Here the rows of all nonsingular minors are among the first d ones of M . Let R_{i_l} , $1 \leq l \leq k$, be k rows of M that generate such a minor \tilde{M} :

$$\tilde{M} = \begin{pmatrix} a_{i_1,1} & a_{i_1,2} & \cdots & a_{i_1,k} \\ a_{i_2,1} & a_{i_2,2} & \cdots & a_{i_2,k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{i_{k-1},1} & a_{i_{k-1},2} & \cdots & a_{i_{k-1},k} \\ a_{i_k,1} & a_{i_k,2} & \cdots & a_{i_k,k} \end{pmatrix}.$$

Proceed as in the case $k = d + 1$ and multiply \tilde{M} by the matrix A to obtain

$$\tilde{M}A = \begin{pmatrix} (a_{i_1,1} - a_{i_1,2}) & (a_{i_1,1} - a_{i_1,3}) & \cdots & (a_{i_1,1} - a_{i_1,k}) & a_{i_1,1} \\ (a_{i_2,1} - a_{i_2,2}) & (a_{i_2,1} - a_{i_2,3}) & \cdots & (a_{i_2,1} - a_{i_2,k}) & a_{i_2,1} \\ (a_{i_3,1} - a_{i_3,2}) & (a_{i_3,1} - a_{i_3,3}) & \cdots & (a_{i_3,1} - a_{i_3,k}) & a_{i_3,1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (a_{i_{k-1},1} - a_{i_{k-1},2}) & (a_{i_{k-1},1} - a_{i_{k-1},3}) & \cdots & (a_{i_{k-1},1} - a_{i_{k-1},k}) & a_{i_{k-1},1} \\ (a_{i_k,1} - a_{i_k,2}) & (a_{i_k,1} - a_{i_k,3}) & \cdots & (a_{i_k,1} - a_{i_k,k}) & a_{i_k,1} \end{pmatrix}.$$

By computing the Laplace expansion along the last column of $\tilde{M}A$, we conclude that at least one $(k-1) \times (k-1)$ minor obtained from the first $k-1$ columns of $\tilde{M}A$ is nonsingular. We argue again as in the $k = d + 1$ case to find the $k-1$ rows that satisfy the property of the lemma for the column C_1 . An analogous argument works for any other column of M , but these $k-1$ special rows may vary from column to column. □

Let us recall some of the terminology from the proof of [Theorem 13.3](#) in [Section 13](#). A subset $\mathcal{A} \subset \mathcal{Q}$ has the *property (P)* if:

- (1) $Q_1 \in \mathcal{A}$.
- (2) \mathcal{A} is not weakly transversal with pivot Q_1 .

We say that $\mathcal{A} \subset \mathcal{Q}$ is *minimal* if $\mathcal{A}' \subset \mathcal{A}$ has the property (P) if and only if $\mathcal{A}' = \mathcal{A}$. Since \mathcal{Q} itself has the property (P), it must contain a minimal subset of cardinality at least 2.

Claim B.6. *Let $\mathcal{A} = \{Q_1, K_2, \dots, K_n\}$ be a minimal set of n cubes.⁵² There is a set D of $d-n+2$ canonical directions v for which*

$$\pi_v(Q_1) \cap \pi_v(K_j) \neq \emptyset \quad \text{for all } 2 \leq j \leq n. \tag{144}$$

Proof of Claim B.6. If $n = 2$, then $Q_1 \cap K_2 \neq \emptyset$ and the claim follows directly. If $n > 2$, observe that $\mathcal{A}' = \{Q_1, K_2, \dots, K_{n-1}\}$ is weakly transversal with pivot Q_1 ; otherwise \mathcal{A} would not be minimal. Hence there are $1 \leq j_1, \dots, j_{n-2} \leq d$ distinct such that

$$\begin{cases} \pi_{j_1}(Q_1) \cap \pi_{j_1}(K_2) = \emptyset, \\ \vdots \\ \pi_{j_{n-2}}(Q_1) \cap \pi_{j_{n-2}}(K_{n-1}) = \emptyset. \end{cases} \tag{145}$$

⁵²Observe that Q_1 is the only “ Q ” cube in this collection. The others are labeled by K_j .

Let $D := \{e_1, \dots, e_d\} \setminus \{e_{j_1}, \dots, e_{j_{n-2}}\}$. In what follows, we will show that (144) holds for this set of directions. Notice that if

$$\pi_l(Q_1) \cap \pi_l(K_n) = \emptyset \tag{146}$$

for some $l \in D$, then \mathcal{A} would be weakly transversal with pivot Q_1 (because (145) together with (146) verify the definition of weak transversality), which is false by hypothesis. Hence (144) holds for $j = n$.

Let us argue by induction that, if (144) holds for $1 \leq m < n - 1$ cubes $K_n, K_{\alpha_1}, \dots, K_{\alpha_{m-1}}$, then it's possible to find a new one K_{α_m} for which (144) also holds⁵³ This will be achieved by the following algorithm: consider the set

$$\mathcal{A}'' := \{Q_1, K_n, K_{\alpha_1}, \dots, K_{\alpha_{m-1}}\}.$$

By the minimality of \mathcal{A} , we know \mathcal{A}'' is weakly transversal with pivot Q_1 ; hence there are $1 \leq r_1, \dots, r_m \leq d$ distinct such that

$$\begin{cases} \pi_{r_1}(Q_1) \cap \pi_{r_1}(K_n) = \emptyset, \\ \pi_{r_2}(Q_1) \cap \pi_{r_2}(K_{\alpha_1}) = \emptyset, \\ \vdots \\ \pi_{r_m}(Q_1) \cap \pi_{r_m}(K_{\alpha_{m-1}}) = \emptyset. \end{cases} \tag{147}$$

Property (P) for \mathcal{A} implies $r_1 \in \{j_1, \dots, j_{n-2}\}$.⁵⁴ Then there is j_{β_1} such that $r_1 = j_{\beta_1}$; therefore

$$\begin{cases} \pi_{j_{\beta_1}}(Q_1) \cap \pi_{j_{\beta_1}}(K_{\beta_1+1}) = \emptyset, \\ \pi_{j_{\beta_1}}(Q_1) \cap \pi_{j_{\beta_1}}(K_n) = \emptyset. \end{cases} \tag{148}$$

Since K_{β_1+1} appears in (145), it is one among K_2, \dots, K_{n-1} ; hence $K_{\beta_1+1} \neq K_n$. We are done if $K_{\beta_1+1} \notin \mathcal{A}''$: indeed, if

$$\pi_l(Q_1) \cap \pi_l(K_{\beta_1+1}) = \emptyset \tag{149}$$

for some $l \in D$, then

$$\begin{cases} \pi_{j_1}(Q_1) \cap \pi_{j_1}(K_2) = \emptyset, \\ \vdots \\ \pi_{j_{\beta_1-1}}(Q_1) \cap \pi_{j_{\beta_1-1}}(K_{\beta_1}) = \emptyset, \\ \pi_l(Q_1) \cap \pi_l(K_{\beta_1+1}) = \emptyset, \\ \pi_{j_{\beta_1+1}}(Q_1) \cap \pi_{j_{\beta_1+1}}(K_{\beta_1+2}) = \emptyset, \\ \vdots \\ \pi_{j_{n-2}}(Q_1) \cap \pi_{j_{n-2}}(K_{n-1}) = \emptyset, \\ \pi_{j_{\beta_1}}(Q_1) \cap \pi_{j_{\beta_1}}(K_n) = \emptyset, \end{cases} \tag{150}$$

and \mathcal{A} would be weakly transversal with pivot Q_1 (by definition again), which contradicts property (P). This way, we would find a new (*not in \mathcal{A}''*) cube K_{β_1+1} for which (144) also holds.

⁵³We are done if there are $m = n - 1$ for which (144) holds, therefore we assume the strict inequality $m < n - 1$.

⁵⁴Otherwise we face the same problem that appeared in (146).

On the other hand, if $K_{\beta_1+1} = K_{\alpha_{q_1}}$ for some $K_{\alpha_{q_1}} \in \mathcal{A}'' \setminus \{K_n\}$, then we simply switch the projections $\pi_{j_{\beta_1}}$ and $\pi_{r_{q_1+1}}$ in (145) (they are distinct because $j_{\beta_1} = r_1 \neq r_{q_1+1}$) and consider the conditions

$$\left\{ \begin{array}{l} \pi_{j_1}(Q_1) \cap \pi_{j_1}(K_2) = \emptyset, \\ \vdots \\ \pi_{j_{\beta_1-1}}(Q_1) \cap \pi_{j_{\beta_1-1}}(K_{\beta_1}) = \emptyset, \\ \pi_{r_{q_1+1}}(Q_1) \cap \pi_{r_{q_1+1}}(K_{\alpha_{q_1}}) = \emptyset, \\ \pi_{j_{\beta_1+1}}(Q_1) \cap \pi_{j_{\beta_1+1}}(K_{\beta_1+2}) = \emptyset, \\ \vdots \\ \pi_{j_{n-2}}(Q_1) \cap \pi_{j_{n-2}}(K_{n-1}) = \emptyset \\ \pi_{j_{\beta_1}}(Q_1) \cap \pi_{j_{\beta_1}}(K_n) = \emptyset, \end{array} \right. \tag{151}$$

where the last condition is taken from (148). Since $j_{\beta_1} \neq r_{q_1+1}$, property (P) for \mathcal{A} again implies that $r_{q_1+1} = j_{\beta_2}$. Notice that $\beta_2 \neq \beta_1$ because $r_1 = j_{\beta_1}$ and $r_1 \neq r_{q_1+1}$. This way, from (145),

$$\left\{ \begin{array}{l} \pi_{j_{\beta_2}}(Q_1) \cap \pi_{j_{\beta_2}}(K_{\beta_2+1}) = \emptyset, \\ \pi_{j_{\beta_2}}(Q_1) \cap \pi_{j_{\beta_2}}(K_{\alpha_{q_1}}) = \emptyset. \end{array} \right. \tag{152}$$

The index j_{β_2} is one of the elements in the set $\{j_1, \dots, j_{\beta_1-1}, j_{\beta_1+1}, \dots, j_{n-2}\}$; hence K_{β_2+1} is in the set $\{K_2, \dots, K_{\beta_1}, K_{\beta_1+2}, \dots, K_{n-1}\}$. As before, we are done if $K_{\beta_2+1} \notin \mathcal{A}''$. If not, $K_{\beta_2+1} = K_{\alpha_{q_2}}$ for some $K_{\alpha_{q_2}} \in \mathcal{A}'' \setminus \{K_n, K_{\alpha_{q_1}}\}$ and we switch the projections $\pi_{j_{\beta_2}}$ and $\pi_{r_{q_2+1}}$ in (151) to find some $\beta_3 \notin \{\beta_1, \beta_2\}$ such that

$$\left\{ \begin{array}{l} \pi_{j_{\beta_3}}(Q_1) \cap \pi_{j_{\beta_3}}(K_{\beta_3+1}) = \emptyset, \\ \pi_{j_{\beta_3}}(Q_1) \cap \pi_{j_{\beta_3}}(K_{\alpha_{q_2}}) = \emptyset. \end{array} \right. \tag{153}$$

We keep doing that until we find some $K_{\beta_{\ell+1}} \notin \mathcal{A}''$. This is guaranteed to happen since there are $n - 1$ cubes K_j , but only $m < n - 1$ of them in \mathcal{A}'' . The conclusion is that

$$m < n - 1 \text{ cubes } K_j \text{ satisfy (144)} \implies m + 1 \text{ cubes } K_j \text{ satisfy (144);}$$

therefore (144) holds for $2 \leq j \leq n$. □

Acknowledgements

We thank David Beltran, Jonathan Bennett, Emanuel Carneiro, Andres Fernandez, Jonathan Hickman, Victor Lie, Diogo Oliveira e Silva, Keith Rogers, Mateus Sousa, Terence Tao, Joshua Zahl and the anonymous referees for many important remarks, corrections and for pointing out references in the literature.

References

[Bejenaru 2022] I. Bejenaru, “The almost optimal multilinear restriction estimate for hypersurfaces with curvature: the case of $n - 1$ hypersurfaces in \mathbb{R}^n ”, *Int. Math. Res. Not.* **2022**:20 (2022), 16363–16404. [MR](#) [Zbl](#)

[Bennett 2014] J. Bennett, “Aspects of multilinear harmonic analysis related to transversality”, pp. 1–28 in *Harmonic analysis and partial differential equations* (El Escorial, Spain, 2012), edited by P. Cifuentes et al., *Contemp. Math.* **612**, Amer. Math. Soc., Providence, RI, 2014. [MR](#) [Zbl](#)

- [Bennett and Nakamura 2021] J. Bennett and S. Nakamura, “Tomography bounds for the Fourier extension operator and applications”, *Math. Ann.* **380**:1-2 (2021), 119–159. [MR](#) [Zbl](#)
- [Bennett et al. 2006] J. Bennett, A. Carbery, and T. Tao, “On the multilinear restriction and Kakeya conjectures”, *Acta Math.* **196**:2 (2006), 261–302. [MR](#) [Zbl](#)
- [Bennett et al. 2008] J. Bennett, A. Carbery, M. Christ, and T. Tao, “The Brascamp–Lieb inequalities: finiteness, structure and extremals”, *Geom. Funct. Anal.* **17**:5 (2008), 1343–1415. [MR](#) [Zbl](#)
- [Bennett et al. 2017] J. Bennett, N. Bez, M. G. Cowling, and T. C. Flock, “Behaviour of the Brascamp–Lieb constant”, *Bull. Lond. Math. Soc.* **49**:3 (2017), 512–518. [MR](#) [Zbl](#)
- [Bennett et al. 2018] J. Bennett, N. Bez, T. C. Flock, and S. Lee, “Stability of the Brascamp–Lieb constant and applications”, *Amer. J. Math.* **140**:2 (2018), 543–569. [MR](#) [Zbl](#)
- [Bennett et al. 2020] J. Bennett, N. Bez, S. Buschenhenke, M. G. Cowling, and T. C. Flock, “On the nonlinear Brascamp–Lieb inequality”, *Duke Math. J.* **169**:17 (2020), 3291–3338. [MR](#) [Zbl](#)
- [Bourgain 1991] J. Bourgain, “Besicovitch type maximal operators and applications to Fourier analysis”, *Geom. Funct. Anal.* **1**:2 (1991), 147–187. [MR](#) [Zbl](#)
- [Bourgain and Demeter 2015] J. Bourgain and C. Demeter, “The proof of the l^2 decoupling conjecture”, *Ann. of Math. (2)* **182**:1 (2015), 351–389. [MR](#) [Zbl](#)
- [Bourgain and Guth 2011] J. Bourgain and L. Guth, “Bounds on oscillatory integral operators based on multilinear estimates”, *Geom. Funct. Anal.* **21**:6 (2011), 1239–1295. [MR](#) [Zbl](#)
- [Fefferman 1970] C. Fefferman, “Inequalities for strongly singular convolution operators”, *Acta Math.* **124** (1970), 9–36. [MR](#) [Zbl](#)
- [Fefferman 1971] C. Fefferman, “The multiplier problem for the ball”, *Ann. of Math. (2)* **94** (1971), 330–336. [MR](#) [Zbl](#)
- [Foschi and Klainerman 2000] D. Foschi and S. Klainerman, “Bilinear space-time estimates for homogeneous wave equations”, *Ann. Sci. École Norm. Sup. (4)* **33**:2 (2000), 211–274. [MR](#) [Zbl](#)
- [Guth 2016] L. Guth, “A restriction estimate using polynomial partitioning”, *J. Amer. Math. Soc.* **29**:2 (2016), 371–413. [MR](#) [Zbl](#)
- [Guth 2018] L. Guth, “Restriction estimates using polynomial partitioning, II”, *Acta Math.* **221**:1 (2018), 81–142. [MR](#) [Zbl](#)
- [Hickman and Iliopoulou 2022] J. Hickman and M. Iliopoulou, “Sharp L^p estimates for oscillatory integral operators of arbitrary signature”, *Math. Z.* **301**:1 (2022), 1143–1189. [MR](#) [Zbl](#)
- [Hickman and Rogers 2019] J. Hickman and K. M. Rogers, “Improved Fourier restriction estimates in higher dimensions”, *Camb. J. Math.* **7**:3 (2019), 219–282. [MR](#) [Zbl](#)
- [Hörmander 1973] L. Hörmander, “Oscillatory integrals and multipliers on FL^p ”, *Ark. Mat.* **11** (1973), 1–11. [MR](#) [Zbl](#)
- [Igari 1986] S. Igari, “Interpolation of operators in Lebesgue spaces with mixed norm and its applications to Fourier analysis”, *Tohoku Math. J. (2)* **38**:3 (1986), 469–490. [MR](#) [Zbl](#)
- [Klainerman and Machedon 1993] S. Klainerman and M. Machedon, “Space-time estimates for null forms and the local existence theorem”, *Comm. Pure Appl. Math.* **46**:9 (1993), 1221–1268. [MR](#) [Zbl](#)
- [Klainerman and Machedon 1995] S. Klainerman and M. Machedon, “Finite energy solutions of the Yang–Mills equations in \mathbb{R}^{3+1} ”, *Ann. of Math. (2)* **142**:1 (1995), 39–119. [MR](#) [Zbl](#)
- [Klainerman and Machedon 1996] S. Klainerman and M. Machedon, “Estimates for null forms and the spaces $H_{s,\delta}$ ”, *Int. Math. Res. Not.* **1996**:17 (1996), 853–865. [MR](#) [Zbl](#)
- [Lee 2021] J. Lee, “An endpoint estimate of the bilinear paraboloid restriction operator”, preprint, 2021. [arXiv 2106.15619](#)
- [Mandel and Oliveira e Silva 2023] R. Mandel and D. Oliveira e Silva, “The Stein–Tomas inequality under the effect of symmetries”, *J. Anal. Math.* **150**:2 (2023), 547–582. [MR](#)
- [Mattila 2015] P. Mattila, *Fourier analysis and Hausdorff dimension*, Cambridge Stud. Adv. Math. **150**, Cambridge Univ. Press, 2015. [MR](#) [Zbl](#)
- [Moyua et al. 1996] A. Moyua, A. Vargas, and L. Vega, “Schrödinger maximal function and restriction properties of the Fourier transform”, *Int. Math. Res. Not.* **1996**:16 (1996), 793–815. [MR](#) [Zbl](#)

- [Muscalu and Schlag 2013a] C. Muscalu and W. Schlag, *Classical and multilinear harmonic analysis, I*, Cambridge Stud. Adv. Math. **137**, Cambridge Univ. Press, 2013. [MR](#) [Zbl](#)
- [Muscalu and Schlag 2013b] C. Muscalu and W. Schlag, *Classical and multilinear harmonic analysis, II*, Cambridge Stud. Adv. Math. **138**, Cambridge Univ. Press, 2013. [MR](#) [Zbl](#)
- [Oh 2023] C. Oh, “An improved bilinear restriction estimate for the paraboloid in \mathbb{R}^3 ”, *Math. Z.* **303**:4 (2023), art. id. 88. [MR](#) [Zbl](#)
- [Ou and Wang 2022] Y. Ou and H. Wang, “A cone restriction estimate using polynomial partitioning”, *J. Eur. Math. Soc.* **24**:10 (2022), 3557–3595. [MR](#) [Zbl](#)
- [Shao 2009] S. Shao, “Sharp linear and bilinear restriction estimates for paraboloids in the cylindrically symmetric case”, *Rev. Mat. Iberoam.* **25**:3 (2009), 1127–1168. [MR](#) [Zbl](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Math. Ser. **43**, Princeton Univ. Press, 1993. [MR](#) [Zbl](#)
- [Stein and Wainger 1978] E. M. Stein and S. Wainger, “Problems in harmonic analysis related to curvature”, *Bull. Amer. Math. Soc.* **84**:6 (1978), 1239–1295. [MR](#) [Zbl](#)
- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. [MR](#) [Zbl](#)
- [Tanaka 2001] H. Tanaka, “An estimate for the Bochner–Riesz operator on functions of product type in \mathbb{R}^2 ”, *Tokyo J. Math.* **24**:2 (2001), 567–578. [MR](#) [Zbl](#)
- [Tao 1999] T. Tao, “The Bochner–Riesz conjecture implies the restriction conjecture”, *Duke Math. J.* **96**:2 (1999), 363–375. [MR](#) [Zbl](#)
- [Tao 2003] T. Tao, “A sharp bilinear restriction estimate for paraboloids”, *Geom. Funct. Anal.* **13**:6 (2003), 1359–1384. [MR](#) [Zbl](#)
- [Tao 2004] T. Tao, “Some recent progress on the restriction conjecture”, pp. 217–243 in *Fourier analysis and convexity*, edited by L. Brandolini et al., Birkhäuser, Boston, 2004. [MR](#) [Zbl](#)
- [Tao and Vargas 2000] T. Tao and A. Vargas, “A bilinear approach to cone multipliers, I: Restriction estimates”, *Geom. Funct. Anal.* **10**:1 (2000), 185–215. [MR](#) [Zbl](#)
- [Tao et al. 1998] T. Tao, A. Vargas, and L. Vega, “A bilinear approach to the restriction and Kakeya conjectures”, *J. Amer. Math. Soc.* **11**:4 (1998), 967–1000. [MR](#) [Zbl](#)
- [Thiele 2006] C. Thiele, *Wave packet analysis*, CBMS Region. Conf. Ser. Math. **105**, Amer. Math. Soc., Providence, RI, 2006. [MR](#) [Zbl](#)
- [Tomas 1975] P. A. Tomas, “A restriction theorem for the Fourier transform”, *Bull. Amer. Math. Soc.* **81** (1975), 477–478. [MR](#) [Zbl](#)
- [Wang 2018] H. Wang, “A restriction estimate in \mathbb{R}^3 using brooms”, preprint, 2018. [arXiv 1802.04312](#)
- [Wolff 2001] T. Wolff, “A sharp bilinear cone restriction estimate”, *Ann. of Math. (2)* **153**:3 (2001), 661–698. [MR](#) [Zbl](#)
- [Zygmund 1974] A. Zygmund, “On Fourier coefficients and transforms of functions of two variables”, *Studia Math.* **50** (1974), 189–201. [MR](#) [Zbl](#)

Received 9 Jun 2022. Revised 15 Feb 2023. Accepted 27 Apr 2023.

CAMIL MUSCALU: camil@math.cornell.edu

Department of Mathematics, Cornell University, Ithaca, NY, United States

ITAMAR OLIVEIRA: oliveira.itamar.w@gmail.com

Department of Mathematics, Cornell University, Ithaca, NY, United States

Current address: School of Mathematics, University of Birmingham, Birmingham, United Kingdom

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 8 2024

Dynamical torsion for contact Anosov flows	2619
YANN CHAUBET and NGUYEN VIET DANG	
Measure propagation along a \mathcal{C}^0 -vector field and wave controllability on a rough compact manifold	2683
NICOLAS BURQ, BELHASSEN DEHMAN and JÉRÔME LE ROUSSEAU	
The Dirichlet problem for the Lagrangian mean curvature equation	2719
ARUNIMA BHATTACHARYA	
Local lens rigidity for manifolds of Anosov type	2737
MIHAJLO CEKIĆ, COLIN GUILLARMOU and THIBAUT LEFEUVRE	
The rank-one theorem on RCD spaces	2797
GIOACCHINO ANTONELLI, CAMILLO BRENA and ENRICO PASQUALETTO	
A new approach to the Fourier extension problem for the paraboloid	2841
CAMIL MUSCALU and ITAMAR OLIVEIRA	
A Poincaré–Steklov map for the MIT bag model	2923
BADREDDINE BENHELLAL, VINCENT BRUNEAU and MAHDI ZREIK	
The weak null condition on Kerr backgrounds	2971
HANS LINDBLAD and MIHAI TOHANEANU	