

ANALYSIS & PDE

Volume 17

No. 8

2024

HANS LINDBLAD AND MIHAI TOHANEANU

THE WEAK NULL CONDITION ON KERR BACKGROUNDS

THE WEAK NULL CONDITION ON KERR BACKGROUNDS

HANS LINDBLAD AND MIHAI TOHANEANU

We study a system of semilinear wave equations on Kerr backgrounds that satisfies the weak null condition. Under the assumption of small initial data, we prove global existence and pointwise decay estimates.

1. Introduction	2971
2. The Kerr metric and local energy estimates	2975
3. Pointwise estimates from local energy decay estimates	2980
4. Improved pointwise bounds	2981
5. Pointwise estimates from the Minkowski fundamental solution	2985
6. Setup for pointwise estimates	2987
7. The bootstrap argument for the Einstein model	2988
Acknowledgments	2994
References	2994

1. Introduction

The semilinear system of wave equations in \mathbb{R}^{1+3}

$$\square\phi = Q[\partial\phi, \partial\phi], \quad \phi|_{t=0} = \phi_0, \quad \partial_t\phi|_{t=0} = \phi_1,$$

where Q is a quadratic form, for small initial data has been studied extensively. For the scalar equation, it is known that the solution can blow up in finite time for $\square\phi = (\partial_t\phi)^2$; see [John 1979]. On the other hand, if the nonlinearity satisfies the null condition by Klainerman [1984], e.g., $\square\phi = (\partial_t\phi)^2 - |\partial_x\phi|^2$, it was shown independently in [Christodoulou 1986] and [Klainerman 1986] that the solution exists globally. This result was extended to quasilinear systems with multiple speeds, as well as the case of exterior domains; see, for instance, [Metcalf et al. 2005; Metcalfe and Sogge 2005; 2007; Hidano 2004; Lindblad et al. 2013; Klainerman and Sideris 1996; Alinhac 2003; Lindblad 1992; 2008; Sideris and Tu 2001; Facci and Metcalfe 2022]. There have also been many works for small data in the variable coefficient case. Almost global existence for nontrapping metrics was shown in [Bony and Häfner 2010; Sogge and Wang 2010]. Global existence for stationary, small perturbations of Minkowski was shown in [Wang and Yu 2014], for nonstationary, compactly supported perturbations in [Yang 2013], and for large, asymptotically flat perturbations that satisfy the strong local energy decay estimates in [Looi and Tohaneanu 2022]. In the context of black holes, global existence was shown in [Luk 2013] for Kerr space-times with small angular momentum, and in [Angelopoulos et al. 2020] for the Reissner–Nordström backgrounds.

MSC2020: primary 35L05, 35L71; secondary 83C57.

Keywords: wave equation, Kerr, weak null condition.

Written in harmonic coordinates, the Einstein equations take the form

$$\square_g g_{\mu\nu} = P[\partial_\mu g, \partial_\nu g] + Q_{\mu\nu}[\partial g, \partial g],$$

where \square_g is the wave operator on the background of the Lorentzian metric g , and P and $Q_{\mu\nu}$ are quadratic forms with coefficients depending on the metric. Unfortunately the nonlinear terms do not satisfy the null condition. Yet Christodoulou and Klainerman [1993] were able to prove global existence for Einstein vacuum equations $R_{\mu\nu} \equiv 0$ for small asymptotically flat initial data. Their proof avoids using coordinates since it was believed the metric in harmonic coordinates would blow up for large times. However, later Lindblad and Rodnianski [2003] noticed that Einstein's equations in harmonic coordinates satisfy a weak null condition, and subsequently used it to prove stability of Minkowski in harmonic coordinates [Lindblad and Rodnianski 2005; 2010]. Whereas it is still unknown if general equations satisfying the weak null condition have global existence for small initial data, there have been a number of results in that direction, including detailed asymptotics of the solution; see for example [Alinhac 2003; Lindblad 1992; 2008; 2017; Keir 2018; Deng and Pusateri 2020; Yu 2021a; 2021b].

There has recently been a lot of activity in proving asymptotic stability of black holes. As a first step people have proved decay of solutions to wave equations on Schwarzschild and Kerr background [Blue and Soffer 2003; 2005; Blue and Sterbenz 2006; Marzuola et al. 2010; Dafermos and Rodnianski 2009; Tataru and Tohaneanu 2011; Dafermos et al. 2016; Andersson and Blue 2015]. People have also studied semilinear perturbations [Luk 2013; Ionescu and Klainerman 2015] satisfying the null condition, but apart from our recent papers [Lindblad and Tohaneanu 2018; 2020], and a global existence result for the Maxwell–Born–Infeld system on a Schwarzschild background [Pasqualotto 2019], little is known about quasilinear perturbations or semilinear perturbations satisfying the weak null condition. There has more recently been progress on the nonlinear stability of Schwarzschild and Kerr [Klainerman and Szeftel 2022a; 2022b; 2023; Dafermos et al. 2021; Giorgi et al. 2022]. These proofs are very long, using sophisticated geometric constructions. We hope that studying models of Einstein's equations in wave coordinates will simplify the proofs and lead to a better understanding and extensions as it did for the stability of Minkowski space.

Finally we remark that there are several recent works on the cosmological case. Hintz and Vasy [2018] proved the stability of Kerr–de Sitter with small angular momentum; see also [Fang 2021; 2022] for an alternative proof. More recently there have been works on the wave equation on Kerr–de Sitter background for large angular momentum assuming there are no growing modes [Peterson and Vasy 2021; Mavrogiannis 2022].

1.0.1. *The semilinear Einstein model.* An example of a simple semilinear system satisfying the weak null condition, but not the classical null condition, is the system

$$\square\phi_1 = (\partial_t\phi_2)^2, \quad \square\phi_2 = 0.$$

It is trivial to see that this has global solutions, and moreover that ϕ_1 decays slower than $1/t$. A less trivial example is the semilinear system

$$\square\phi_1 = (\partial_t\phi_2)^2 + Q_1[\partial\phi, \partial\phi], \quad \square\phi_2 = Q_2[\partial\phi, \partial\phi],$$

where Q_j are null forms. These systems have the advantage that the components ϕ_1 and ϕ_2 decouple to highest order. For Einstein's equations there is the additional difficulty that this decoupling can only be seen in a null frame, and contractions with the frame do not commute with the wave operator as far as the L^2 estimate. Hence a more realistic model is the system is

$$\square\phi_{\mu\nu} = P[\partial_\mu\phi, \partial_\nu\phi] + Q_{\mu\nu}[\partial\phi, \partial\phi],$$

where P is assumed to have a certain weak null structure. Contracting with a nullframe this resembles the decoupled systems with $\phi_{\underline{LL}}$ in place of ϕ_1 , where $\underline{L}^\mu\partial_\mu = \partial_t - \partial_r$, and ϕ_2 replaced by the other components ϕ_{TU} , where T is tangential to the outgoing light cones. The only really bad component is $\partial\phi_{\underline{LL}}$ but this one does not show up quadratically in P for Einstein's equations. It shows up linearly but multiplied with a component $\partial\phi_{\underline{LL}}$ that has vanishing radiation field due to the wave coordinate condition.

With the goal of understanding Einstein's equations in (generalized) harmonic coordinates close to Kerr with small angular momentum, we will focus on the following system, which resembles the semilinear part of Einstein's equations:

$$\square_K\phi_{\mu\nu} = P[\partial_\mu\phi, \partial_\nu\phi] + Q_{\mu\nu}[\partial\phi, \partial\phi], \quad \tilde{t} \geq 0, \quad \phi|_{\tilde{t}=0} = \phi_0, \quad \tilde{T}\phi|_{\tilde{t}=0} = \phi_1. \quad (1-1)$$

Here \square_K denotes the d'Alembertian with respect to the Kerr metric, and \tilde{T} is a smooth, everywhere timelike vector field that equals ∂_t away from the black hole. The coordinate \tilde{t} is chosen so that the slices $\tilde{t} = \text{const.}$ are space-like and $\tilde{t} = t$ away from the black hole. For simplicity we will consider compactly supported smooth initial data, but suitably weighted Sobolev spaces of large enough order would suffice. Moreover, $Q_{\mu\nu}$ are null forms and P is a symmetric quadratic form:

$$P[\phi, \psi] = P^{\alpha\beta\gamma\delta}(x/\tilde{t})\phi_{\alpha\beta}\psi_{\gamma\delta},$$

with coefficients with a certain weak null structure. We remove the component $\partial\phi_{\underline{LL}}$ by imposing the condition

$$P^{\underline{LL}\alpha\beta}(x/\tilde{t}) = P^{\alpha\beta\underline{LL}}(x/\tilde{t}) = 0.$$

For this system we cannot have different energy estimates for different components because the null structure is only seen in a null frame and contractions with the frame do not commute with the wave operator. Because of this, one cannot get the decay estimates directly from the L^2 estimates but one has to use the equations again to get improved decay estimates. As a result, the proof is more involved. The method we develop avoids boosts vector fields and combines local energy decay in a compact set with estimates in characteristic coordinates at the light cone. It gives an essentially optimal decay of almost \tilde{t}^{-1} , which is an improvement over $\tilde{t}^{-1/2}$, which can be obtained more easily from energy estimates. The method in particular works close to Minkowski where it gives the optimal decay without using boosts.

Finally we remark that this system can be combined with the quasilinear system that we previously studied [Lindblad and Tohaneanu 2018; 2020] (see also [Looi 2022] for improved pointwise bounds) to resemble also the quasilinear part of Einstein's equations

$$\square_{g[\phi]}\phi_{\mu\nu} = P[\partial_\mu\phi, \partial_\nu\phi] + Q_{\mu\nu}[\partial\phi, \partial\phi],$$

where

$$g^{\alpha\beta}[\phi] = K^{\alpha\beta} + H^{\alpha\beta}[\phi], \quad \text{where } H^{\alpha\beta}[\phi] = H^{\alpha\beta\mu\nu}(x/\tilde{t})\phi_{\mu\nu} \text{ and } H^{LL\mu\nu}(x/\tilde{t}) = 0.$$

1.0.2. Statement of the results. We are now ready to state our main result. We define \tilde{r} to be some function that equals r near the event horizon, and r_K^* away from it; see Section 2 for more details. We also fix r_e satisfying $r_- < r_e < r_+$, and define $\langle x \rangle = (2 + |x|^2)^{1/2}$.

Theorem 1.1. *Let $R_0 > r_e$, and assume that ϕ_0, ϕ_1 are smooth and compactly supported in $\tilde{r} \leq R_0$. Then there exists a global classical solution to the system (1-1) (on a Kerr metric with $|a| \ll M$) provided that, for a certain $\epsilon_0 \ll 1$ and large enough N , we have*

$$\mathcal{E}_N(0) = \|(\phi_0, \phi_1)\|_{H^{N+1} \times H^N} \leq \epsilon_0.$$

Moreover, for some fixed positive integer m , independent of N , we have for any $\delta > 0$

$$\begin{aligned} |\phi_{\leq N-m}| &\lesssim \frac{\mathcal{E}_N(0) \langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{\langle \tilde{t} \rangle}, & |\partial\phi_{\leq N-m}| &\lesssim \frac{\mathcal{E}_N(0) \langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{r \langle \tilde{t} - \tilde{r} \rangle}, \\ |(\partial\phi_{TU})_{\leq N-m}| &\lesssim \frac{\mathcal{E}_N(0)}{r \langle \tilde{t} - \tilde{r} \rangle^{1-\delta}}. \end{aligned}$$

Note that is an improvement of the decay estimates we previously proved essentially by a factor of $\tilde{t}^{-1/2}$. Note also the structure here, that a derivative decreases the homogeneity, but because the homogeneous vector fields we can use together with the wave operator do not span the tangent space at the origin or at the light cone, a derivative only improves by a power of r close to the origin and a power of $\tilde{t} - \tilde{r}$ close to the light cone. Note also that close to the light cone we have a better estimate for the good components which is due to the weak null structure.

1.0.3. Structure of the proof. The starting point is the local energy estimate in Section 2. The local energy scales like the energy, which is consistent with a decay $\tilde{t}^{-1/2}$ of order $-\frac{1}{2}$ for ϕ and $-\frac{3}{2}$ for the derivatives, and this is also the decay we were able to obtain in our previous paper from a bound of the local energy applied to scaling and rotation vector fields; see Section 3. Assuming these decay estimates, one can go back into the equation and get improved decay estimates. In fact from these decay estimates the total decay of the inhomogeneous term would be -3 , which would be consistent with a solution of the wave equation with decay of order -1 . We prove this using L^∞ estimates for the wave operator from Section 5. However the first improved estimates we obtain have the improved decay in r or $\tilde{t} - \tilde{r}$ and we need improved decay in \tilde{t} . For this we have other estimates turn decay in r or $\tilde{t} - \tilde{r}$ into decay in \tilde{t} ; see Section 4. The whole argument is put together in the last section.

The paper is structured as follows. In Section 2 we introduce the Kerr metric, the vector fields we will use, and the local energy estimates which will play a key role in the proof. Sections 3, 4, 5, and 6 contain various estimates that will allow us to extract the necessary pointwise bounds for (vector fields applied to) the solution. Finally, Section 7 contains the bootstrap argument.

2. The Kerr metric and local energy estimates

2.1. The Kerr metric. The Kerr geometry in Boyer–Lindquist coordinates is given by

$$ds^2 = g_{tt}^K dt^2 + g_{t\phi} dt d\phi + g_{rr}^K dr^2 + g_{\phi\phi}^K d\phi^2 + g_{\theta\theta}^K d\theta^2,$$

where $t \in \mathbb{R}$, $r > 0$, (ϕ, θ) are the spherical coordinates on \mathbb{S}^2 and

$$g_{tt}^K = -\frac{\Delta - a^2 \sin^2 \theta}{\rho^2}, \quad g_{t\phi}^K = -2a \frac{2Mr \sin^2 \theta}{\rho^2}, \quad g_{rr}^K = \frac{\rho^2}{\Delta},$$

$$g_{\phi\phi}^K = \frac{(r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta}{\rho^2} \sin^2 \theta, \quad g_{\theta\theta}^K = \rho^2,$$

with

$$\Delta = r^2 - 2Mr + a^2, \quad \rho^2 = r^2 + a^2 \cos^2 \theta.$$

Here M represents the mass of the black hole, and aM its angular momentum.

A straightforward computation gives us the inverse of the metric:

$$g_K^{tt} = -\frac{(r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta}{\rho^2 \Delta}, \quad g_K^{t\phi} = -a \frac{2Mr}{\rho^2 \Delta}, \quad g_K^{rr} = \frac{\Delta}{\rho^2},$$

$$g_K^{\phi\phi} = \frac{\Delta - a^2 \sin^2 \theta}{\rho^2 \Delta \sin^2 \theta}, \quad g_K^{\theta\theta} = \frac{1}{\rho^2}.$$

The case $a = 0$ corresponds to the Schwarzschild space-time. We shall subsequently assume that a is small $0 < a \ll M$, so that the Kerr metric is a small perturbation of the Schwarzschild metric. Note also that the coefficients depend only r and θ but are independent of ϕ and t . We denote the d'Alembertian associated to the Kerr metric by \square_K .

In the above coordinates the Kerr metric has singularities at $r = 0$, on the equator $\theta = \frac{\pi}{2}$, and at the roots of Δ , namely $r_{\pm} = M \pm \sqrt{M^2 - a^2}$. To remove the singularities at $r = r_{\pm}$ we introduce functions $r_K^* = r_K^*(r)$, $v_+ = t + r_K^*$ and $\phi_+ = \phi_+(\phi, r)$ so that (see [Hawking and Ellis 1973])

$$dr_K^* = (r^2 + a^2) \Delta^{-1} dr, \quad dv_+ = dt + dr_K^*, \quad d\phi_+ = d\phi + a \Delta^{-1} dr.$$

Note that when $a = 0$ the r_K^* -coordinate becomes the Schwarzschild Regge–Wheeler coordinate

$$r^* = r + 2M \log(r - 2M).$$

The Kerr metric can be written in the new coordinates (v_+, r, ϕ_+, θ) ,

$$ds^2 = -\left(1 - \frac{2Mr}{\rho^2}\right) dv_+^2 + 2dr dv_+ - 4a\rho^{-2} Mr \sin^2 \theta dv_+ d\phi_+ - 2a \sin^2 \theta dr d\phi_+ + \rho^2 d\theta^2$$

$$+ \rho^{-2} [(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta] \sin^2 \theta d\phi_+^2,$$

which is smooth and nondegenerate across the event horizon up to but not including $r = 0$. We introduce the function

$$\tilde{t} = v_+ - \mu(r),$$

where μ is a smooth function of r . In the $(\tilde{t}, r, \phi_+, \theta)$ -coordinates the metric has the form

$$ds^2 = \left(1 - \frac{2Mr}{\rho^2}\right) d\tilde{t}^2 + 2\left(1 - \left(1 - \frac{2Mr}{\rho^2}\right)\mu'(r)\right) d\tilde{t} dr - 4a\rho^{-2}Mr \sin^2 \theta d\tilde{t} d\phi_+ \\ + \left(2\mu'(r) - \left(1 - \frac{2Mr}{\rho^2}\right)(\mu'(r))^2\right) dr^2 - 2a(1 + 2\rho^{-2}Mr\mu'(r)) \sin^2 \theta dr d\phi_+ \\ + \rho^2 d\theta^2 + \rho^{-2}[(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta] \sin^2 \theta d\phi_+^2.$$

On the function μ we impose the following two conditions:

- (i) $\mu(r) \geq r_K^*$ for $r > 2M$, with equality for $r > \frac{5}{2}M$.
- (ii) The surfaces $\tilde{t} = \text{const.}$ are space-like, i.e.,

$$\mu'(r) > 0, \quad 2 - \left(1 - \frac{2Mr}{\rho^2}\right)\mu'(r) > 0.$$

As long as a is small, we can use the same function μ as in the case of the Schwarzschild space-time in [Marzuola et al. 2010].

We also introduce

$$\tilde{\phi} = \zeta(r)\phi_+ + (1 - \zeta(r))\phi,$$

where ζ is a cutoff function supported near the event horizon.

We fix r_e satisfying $r_- < r_e < r_+$. The choice of r_e is unimportant, and for convenience we may simply use $r_e = M$ for all Kerr metrics with $a/M \ll 1$. Let $\mathcal{M} = \{\tilde{t} \geq 0 : r \geq r_e\}$, $\Sigma(T) = \mathcal{M} \cap \{\tilde{t} = T\}$, and $d\Sigma_K$ be the induced volume element on $\Sigma(T)$.

Let \tilde{r} denote a smooth strictly increasing function (of r) that equals r for $r \leq R$ and r_K^* for $r \geq 2R$ for some large R . We will use the coordinates (\tilde{t}, x^i) , where $x^i = \tilde{r}\omega$. Note that, since $r \approx \tilde{r}$, we can use r^k and \tilde{r}^k interchangeably when defining our spaces of functions in what follows.

2.2. Vector fields and spaces of functions. Our favorite sets of vector fields will be

$$\partial \in \{\partial_{\tilde{t}}, \partial_i\}, \quad \Omega \in \{x^i \partial_j - x^j \partial_i\}, \quad S = \tilde{t} \partial_{\tilde{t}} + \tilde{r} \partial_{\tilde{r}},$$

namely the generators of translations, rotations and scaling. We set $Z = \{\partial, \Omega, S\}$.

We also denote by ∂ the angular derivatives,

$$\partial_j = \frac{x^i}{\tilde{r}} \partial_{\tilde{r}} + \partial_i$$

and let

$$\bar{\partial} \in (\partial_v, \partial), \quad \partial_v = \partial_{\tilde{t}} + \partial_{\tilde{r}}$$

denote the tangential derivatives. We also let $\underline{L} = \partial_{\tilde{t}} - \partial_{\tilde{r}}$.

For a triplet $\alpha = (i, j, k)$, we define $|\alpha| = i + 3j + 3k$ and

$$u_\alpha = \partial^i \Omega^j S^k u, \quad u_{\leq m} = (u_\Lambda)_{|\Lambda| \leq m}.$$

The notation is borrowed from [Lindblad and Tohaneanu 2018], and takes into account the loss of derivatives that occurs when applying weak local energy estimates to vector fields.

Given a norm $\|\cdot\|_X$, we write

$$\|u_{\leq m}\|_X = \sum_{|\Lambda| \leq m} \|u_\Lambda\|_X.$$

We define the classes $S^Z(r^k)$ of functions in $\mathbb{R}^+ \times \mathbb{R}^3$ by

$$f \in S^Z(r^k) \iff |Z^j f(t, x)| \leq c_j (r)^k, \quad j \geq 0.$$

Given a family of functions \mathcal{G} , we will also use the notation

$$f \in S^Z(r^k)\mathcal{G}$$

to mean that

$$f = \sum h_i g_i, \quad h_i \in S^Z(r^k), \quad g_i \in \mathcal{G}.$$

We will also use the notation U for an element of $S^Z(1)Z$, and T for an element of $S^Z(1)\bar{\partial}$. An important observation is that, since

$$\partial_v = \frac{\tilde{t} - \tilde{r}}{\tilde{t}} \partial_{\tilde{r}} + \frac{1}{\tilde{t}} S, \quad \not\partial \phi \in S^Z(r^{-1})\Omega\phi,$$

we have

$$|\bar{\partial} w| \lesssim \frac{\tilde{t} - \tilde{r}}{r} |\partial w| + \frac{1}{r} |\Omega w|. \tag{2-1}$$

Moreover, an easy computation gives

$$\begin{aligned} [\square_K, \partial]\phi &\in S^Z(r^{-2})\partial\partial^{\leq 1}\phi, & [\square_K, \Omega]\phi &\in S^Z(r^{-2})\partial\partial^{\leq 1}\phi, \\ [\square_K, S]\phi &\in S^Z(1)\square_K\phi + S^Z(r^{-2+})\partial\phi + S^Z(r^{-2+})\partial\Omega\phi + S^Z(r^{-2})\partial\partial^{\leq 1}\phi, \end{aligned}$$

and thus by induction we obtain that

$$[\square_K, Z^\alpha]\phi = F_1 + F_2, \quad F_1 \in S^Z(1)(\square_K\phi)_{\leq |\alpha|}, \quad F_2 \in S^Z(r^{-2+})\partial\phi_{\leq |\alpha|}. \tag{2-2}$$

We now claim that

$$[Z, \bar{\partial}] \in S^Z(1)\bar{\partial} + S^Z(r^{-1})\partial. \tag{2-3}$$

Indeed, we compute

$$\begin{aligned} [\partial_{\tilde{r}}, \bar{\partial}] &= 0, & [\partial_i, \partial_v] &= [\not\partial_i, \partial_{\tilde{r}}] \in S^Z(r^{-1})\not\partial, & [\partial_i, \not\partial] &\in S^Z(r^{-1})\partial, \\ [\Omega, \partial_v] &= 0, & [\Omega, \not\partial] &\in S^Z(1)\not\partial, & [S, \partial_v] &= \partial_v, & [S, \not\partial] &\in S^Z(1)\not\partial. \end{aligned}$$

This proves (2-3).

Given vector fields X and Y , we define

$$\phi_{XY} = X^\alpha Y^\beta \phi_{\alpha\beta}.$$

Similarly, we can write the coefficients P with respect to the vector frame $\{\underline{L}, \bar{\partial}\}$ as

$$P^{\alpha\beta\gamma\delta} = P^{\underline{L}\underline{L}\gamma\delta} \underline{L}^\alpha \underline{L}^\beta + \sum P^{TU\gamma\delta} T^\alpha U^\beta,$$

$$P^{\alpha\beta\gamma\delta} = P^{\alpha\beta\underline{L}\underline{L}} \underline{L}^\gamma \underline{L}^\delta + \sum P^{\alpha\beta TU} T^\gamma U^\delta.$$

The assumptions on the coefficients $P^{\alpha\beta\gamma\delta}$ are the following:

$$P^{\alpha\beta\gamma\delta} \in S^Z(1), \tag{2-4}$$

$$P^{\underline{L}\underline{L}\alpha\beta} = P^{\alpha\beta\underline{L}\underline{L}} = 0. \tag{2-5}$$

Equation (2-5) means that terms like $\underline{L}\phi_{\underline{L}\underline{L}} \partial\phi$ do not appear on the right-hand side of (1-1).

The assumption on the null forms $Q_{\mu\nu}$ is that

$$Q_{\mu\nu}[\partial\phi, \partial\phi] \in S^Z(1) \partial\phi \bar{\partial}\phi. \tag{2-6}$$

2.3. Local energy estimates. We consider a partition of \mathbb{R}^3 into the dyadic sets $A_R = \{R \leq \langle \tilde{r} \rangle \leq 2R\}$ for $R \geq 1$.

We now introduce the local energy norm LE

$$\|u\|_{LE} = \sup_R \|\langle r \rangle^{-1/2} u\|_{L^2(\mathcal{M} \cap \mathbb{R} \times A_R)},$$

$$\|u\|_{LE_{[0,1]}} = \sup_R \|\langle r \rangle^{-1/2} u\|_{L^2(\mathcal{M} \cap [0,1] \times \mathbb{A}_R)},$$

its H^1 counterpart

$$\|u\|_{LE^1} = \|\partial u\|_{LE} + \|\langle r \rangle^{-1} u\|_{LE},$$

$$\|u\|_{LE^1_{[0,1]}} = \|\partial u\|_{LE_{[0,1]}} + \|\langle r \rangle^{-1} u\|_{LE_{[0,1]}},$$

as well as the dual norm

$$\|f\|_{LE^*} = \sum_R \|\langle r \rangle^{1/2} f\|_{L^2(\mathcal{M} \cap \mathbb{R} \times A_R)},$$

$$\|f\|_{LE^*_{[0,1]}} = \sum_R \|\langle r \rangle^{1/2} f\|_{L^2(\mathcal{M} \cap [0,1] \times \mathbb{A}_R)}.$$

We also define similar norms for higher Sobolev regularity

$$\|u_{\leq m}\|_{LE^1} = \sum_{|\alpha| \leq m} \|u_\alpha\|_{LE^1},$$

$$\|u_{\leq m}\|_{LE^1_{[0,1]}} = \sum_{|\alpha| \leq m} \|u_\alpha\|_{LE^1_{[0,1]}},$$

$$\|u_{\leq m}\|_{LE_{[0,1]}} = \sum_{|\alpha| \leq m} \|u_\alpha\|_{LE_{[0,1]}},$$

respectively,

$$\|f\|_{LE^{*,k}} = \sum_{|\alpha| \leq k} \|\partial^\alpha f\|_{LE^*},$$

$$\|f\|_{LE^{*,k}_{[0,1]}} = \sum_{|\alpha| \leq k} \|\partial^\alpha f\|_{LE^*_{[0,1]}}.$$

Finally, we introduce a weaker version of the local energy decay norm

$$\begin{aligned} \|u\|_{LE_w^1} &= \|(1 - \chi_{ps}) \partial u\|_{LE} + \|\partial_r u\|_{LE} + \|\langle r \rangle^{-1} u\|_{LE}, \\ \|u\|_{LE_w^1[0,1]} &= \|(1 - \chi_{ps}) \partial u\|_{LE[0,1]} + \|\partial_r u\|_{LE[0,1]} + \|\langle r \rangle^{-1} u\|_{LE[0,1]}. \end{aligned}$$

To measure the inhomogeneous term, we define

$$\begin{aligned} \|f\|_{LE_w^*} &= \inf_{f_1+f_2=f} \|f_1\|_{L^1L^2} + \|(1 - \chi_{ps}) f_2\|_{LE^*}, \\ \|f\|_{LE_w^*[0,1]} &= \inf_{f_1+f_2=f} \|f_1\|_{L^1[0,1]L^2} + \|(1 - \chi_{ps}) f_2\|_{LE^*[0,1]}. \end{aligned}$$

Here χ_{ps} is a smooth, compactly supported spatial cutoff function that equals 1 in a neighborhood of the trapped set. We also define the higher-order weak norms as above.

We define the (nondegenerate) energy

$$E[u](\tilde{t}) = \left(\int_{\Sigma(\tilde{t})} |\partial u|^2 d\Sigma_K \right)^{1/2}.$$

We now fix some $\delta_1 \ll 1$, and define, for a large enough constant R_1 (so that in particular $\chi_{ps} = 0$ when $r > R_1$):

$$\mathcal{E}_N(T) = \sup_{0 \leq \tilde{t} \leq T} E[\phi_{\leq N}](\tilde{t}) + \|\phi_{\leq N}\|_{LE_w^1[0,T]} + \|\langle \tilde{t} - \tilde{r} \rangle^{(-1-\delta_1)/2} \bar{\partial} \phi_{\leq N}\|_{L^2[0,T]L^2(r \geq R_1)}. \tag{2-7}$$

We will need the following local energy estimates for the linear problem:

Lemma 2.1. *Assume that $\square_K \phi = F$, and N is any nonnegative integer. We then have for any $T \geq 0$ that*

$$\mathcal{E}_N(T) \lesssim \mathcal{E}_N(0) + \|F_{\leq N}\|_{LE_w^*[0,T]}, \tag{2-8}$$

where the implicit constant is independent of T .

Proof. We start by proving the base case $N = 0$, that is,

$$\sup_{0 \leq \tilde{t} \leq T} E[\phi](\tilde{t}) + \|\phi\|_{LE_w^1[0,T]} + \|\langle \tilde{t} - \tilde{r} \rangle^{(-1-\delta_1)/2} \bar{\partial} \phi\|_{L^2[0,T]L^2(r \geq R_1)} \lesssim E[\phi](0) + \|F\|_{LE_w^*[0,T]}. \tag{2-9}$$

Theorem 4.1 from [Tataru and Tohaneanu 2011] gives the desired bound for the first two terms on the left-hand side. On the other hand, Lemma 4.3 in [Lindblad and Tohaneanu 2018] and Cauchy–Schwarz yield

$$\|\langle \tilde{t} - \tilde{r} \rangle^{(-1-\delta_1)/2} \bar{\partial} \phi\|_{L^2[0,T]L^2(r \geq R_1)} \lesssim \|\phi\|_{LE_w^1[0,T]} + \|F\|_{LE_w^*[0,T]}.$$

We now commute the equation with the vector fields in Z . Applying the base case estimate (2-9) to ϕ_α for some $|\alpha| = N$ yields

$$\begin{aligned} \sup_{0 \leq \tilde{t} \leq T} E[\phi_\alpha](\tilde{t}) + \|\phi_\alpha\|_{LE_w^1[0,T]} + \|\langle \tilde{t} - \tilde{r} \rangle^{(-1-\delta_1)/2} \bar{\partial} \phi_\alpha\|_{L^2[0,T]L^2(r \geq R_1)} \\ \lesssim \|\phi_\alpha\|_{LE_w^1[0,T]} + \|F_\alpha\|_{LE_w^*[0,T]} + \|[\square_K, Z^\alpha] \phi\|_{LE_w^*[0,T]}. \end{aligned}$$

We are left with bounding the last term on the right-hand side. By (2-2) we have

$$\|[\square_K, Z^\alpha]\phi\|_{LE_w^*[0,T]} \lesssim \|F_{\leq|\alpha|}\|_{LE_w^*[0,T]} + \|r^{-2+} \partial\phi_{\leq|\alpha|}\|_{LE_w^*[0,T]} \lesssim \|F_{\leq|\alpha|}\|_{LE_w^*[0,T]} + \|\phi_{\leq|\alpha|}\|_{LE_w^1[0,T]}.$$

We now sum over all $|\alpha| = N$. □

The first estimate of this kind was obtained by Morawetz for the Klein–Gordon equation [Morawetz 1968]. In the Schwarzschild case, similar estimates were shown in [Blue and Soffer 2003; 2005; Blue and Sterbenz 2006; Dafermos and Rodnianski 2009; 2007; Marzuola et al. 2010]. The estimate for Kerr with small angular momentum was proven in [Tataru and Tohaneanu 2011] (see also [Andersson and Blue 2015; Dafermos and Rodnianski 2013] for related works). For large angular momentum, see [Dafermos et al. 2016] ($|a| < M$) and [Aretakis 2012] ($|a| = M$).

3. Pointwise estimates from local energy decay estimates

The goal of this section is to show how to extract (weak) pointwise estimates from local energy norms. These bounds will serve as the starting point in an iteration that will yield strong enough pointwise bounds to close the bootstrap argument in Section 7.

Let

$$C_T = \{T \leq \tilde{t} \leq 2T : \tilde{r} \leq \tilde{t}\}.$$

We use a double dyadic decomposition of C_T with respect to either the size of $\tilde{t} - \tilde{r}$ or the size of r , depending on whether we are close or far from the cone,

$$C_T = \bigcup_{1 \leq R \leq T/4} C_T^R \bigcup \bigcup_{1 \leq U < T/4} C_T^U,$$

where for $R, U > 1$ we set

$$C_T^R = C_T \cap \{R < r < 2R\}, \quad C_T^U = C_T \cap \{U < \tilde{t} - \tilde{r} < 2U\},$$

while for $R = 1$ and $U = 1$ we have

$$C_T^{R=1} = C_T \cap \{0 < r < 2\}, \quad C_T^{U=1} = C_T \cap \{0 < \tilde{t} - \tilde{r} < 2\}.$$

The sets C_T^R and C_T^U represent the setting in which we apply Sobolev embeddings, which allow us to obtain pointwise bounds from L^2 bounds. Precisely, we have (see Lemma 3.8 from [Metcalf et al. 2012] and Lemma 6.2 in [Lindblad and Tohaneanu 2018]):

Lemma 3.1. *For any function w and all $T \geq 1$ and $1 \leq R, U \leq \frac{1}{4}T$ we have*

$$\|w\|_{L^\infty(C_T^R)} \lesssim \frac{1}{T^{1/2}R^{3/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j w\|_{L^2(C_T^R)} + \frac{1}{T^{1/2}R^{1/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j \partial w\|_{L^2(C_T^R)}, \tag{3-1}$$

respectively,

$$\|w\|_{L^\infty(C_T^U)} \lesssim \frac{1}{T^{3/2}U^{1/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j w\|_{L^2(C_T^U)} + \frac{U^{1/2}}{T^{3/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j \partial w\|_{L^2(C_T^U)}. \tag{3-2}$$

Using the lemma above, we prove the following pointwise bound:

$$\|w\|_{L^\infty(C_T)} \lesssim \langle \tilde{t} \rangle^{-1} \langle \tilde{t} - \tilde{r} \rangle^{1/2} \|w_{\leq 12}\|_{LE^1[T, 2T]}. \tag{3-3}$$

Indeed, in the region C_T^R , this is an immediate application of (3-1). On the other hand, in the region C_T^U this follows from (3-2) and Hardy’s inequality; see (6.7) in [Lindblad and Tohaneanu 2018].

We also need an L^∞ bound on the derivative that is better than (3-3) for large r . This is the content of the following, which is essentially Proposition 3.5 in [Looi and Tohaneanu 2022]

Proposition 3.2. *Let*

$$\mu := \min(\langle \tilde{t} \rangle, \langle \tilde{t} - \tilde{r} \rangle)^{1/2}.$$

Assume that ϕ solve (1-1) for $t \in [T, 2T]$. Then for any dyadic region $C \in \{C_T^R, C_U^R\}$ and $m \geq 0$ we have

$$\|\partial\phi_{\leq m}\|_{L^\infty(C)} \leq \bar{C}_m \frac{1}{\mu} \left(\frac{1}{\langle r \rangle} + \|\partial\phi_{\leq (m+10)/2}\|_{L^\infty(C)} \right) \|\phi_{\leq m+5}\|_{LE^1[T, 2T]}. \tag{3-4}$$

Here the crucial estimate was the following Klainerman–Sideris-type estimate; see Lemma 5.4 in [Lindblad and Tohaneanu 2018] (for Schwarzschild) combined with the remarks after (5.13) in [Lindblad and Tohaneanu 2020]:

Lemma 3.3. *For any w and multiindex Λ we have in the region $r \geq 2R_1$ that*

$$|\partial^2 w_\Lambda| \lesssim \frac{\tilde{t}}{r \langle \tilde{t} - \tilde{r} \rangle} |\partial w_{\leq |\Lambda|+3}| + \frac{\tilde{t}}{\langle \tilde{t} - \tilde{r} \rangle} |(\square_K w)_{\leq |\Lambda|}|.$$

We now apply (3-2) to $\partial\phi_\Lambda$ for any $|\Lambda| \leq m$. We obtain

$$\begin{aligned} \|\partial\phi_\Lambda\|_{L^\infty(C_T^U)} &\lesssim \frac{1}{T^{3/2}U^{1/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j \partial\phi_\Lambda\|_{L^2(C_T^U)} + \frac{U^{1/2}}{T^{3/2}} \sum_{i \leq 1, j \leq 2} \|S^i \Omega^j \partial^2\phi_\Lambda\|_{L^2(C_T^U)} \\ &\lesssim \frac{1}{TU^{1/2}} \|\phi_{\leq |\Lambda|+13}\|_{LE^1[T, 2T]} + \frac{1}{(TU)^{1/2}} \|(\square_K \phi)_{\leq |\Lambda|+10}\|_{L^2(C_T^U)}. \end{aligned}$$

Since

$$|(\square_K \phi)_{\leq |\Lambda|+10}| \lesssim |\partial\phi_{\leq |\Lambda|/2+5}| |\partial\phi_{\leq |\Lambda|+10}|,$$

the conclusion follows in the region C_T^U . A similar computation yields the result in C_T^R .

4. Improved pointwise bounds

We will use three lemmas that will help us improve our pointwise bounds. The first one is Proposition 3.14 from [Metcalf et al. 2012], which will allow us to turn r -decay into t -decay in the region $r \leq \frac{1}{2}t$.

Lemma 4.1. *The following estimate holds for all $m \geq 0$ and some fixed (m -independent) n :*

$$\|u_{\leq m}\|_{LE^1(C_T^{<T/2})} \lesssim T^{-1} \|\langle r \rangle u_{\leq m+n}\|_{LE^1(C_T^{<T/2})} + \|(\square_K u)_{\leq m+n}\|_{LE^*(C_T^{<T/2})}.$$

The second lemma is a slight modification of Lemma 3.11 from [Metcalf et al. 2012], the difference being that we may not enlarge our regions in time. The role of the lemma is to gain a factor of $\tilde{t}/(r(\tilde{t} - \tilde{r}))$ for the derivative.

We let \tilde{C}_T^R and \tilde{C}_T^U denote enlargements of C_T^R and C_T^U in space (but not in time) that contain all the integral curves of the scaling vector field S (i.e., if $(t, x) \in C_T^R$ then $(st, sx) \in \tilde{C}_T^R$ as long as $T \leq st \leq 2T$ and similarly for C_T^U). More precisely, let

$$\begin{aligned} \tilde{C}_T^R &= \left\{ T \leq \tilde{t} \leq 2T : \frac{8}{10} \frac{T}{2R} \leq \frac{\tilde{t}}{\tilde{r}} \leq \frac{12}{10} \frac{2T}{R} \right\}, & \tilde{C}_T^R(\tau) &= \tilde{C}_T^R \cap \{\tilde{t} = \tau\}, \\ \tilde{C}_T^U &= \left\{ T \leq \tilde{t} \leq 2T : \frac{8}{10} \frac{T}{T-2U} \leq \frac{\tilde{t}}{\tilde{r}} \leq \frac{12}{10} \frac{2T}{2T-U} \right\}, & \tilde{C}_T^U(\tau) &= \tilde{C}_T^U \cap \{\tilde{t} = \tau\}. \end{aligned}$$

An important observation here is that $\tilde{r} \approx R$ and $\tilde{t} - \tilde{r} \approx U$ in \tilde{C}_T^R and \tilde{C}_T^U respectively.

Lemma 4.2. For $1 \ll U, R \leq \frac{1}{4}T$ we have

$$\|\partial w\|_{L^2(C_T^R)} \lesssim R^{-1} \|w\|_{L^2(\tilde{C}_T^R)} + T^{-1} (\|Sw\|_{L^2(\tilde{C}_T^R)} + \|S^2w\|_{L^2(\tilde{C}_T^R)}) + R \|\square_K w\|_{L^2(\tilde{C}_T^R)}, \tag{4-1}$$

respectively,

$$\|\partial w\|_{L^2(C_T^U)} \lesssim U^{-1} (\|w\|_{L^2(\tilde{C}_T^U)} + \|Sw\|_{L^2(\tilde{C}_T^U)} + \|S^2w\|_{L^2(\tilde{C}_T^U)}) + T \|\square_K w\|_{L^2(\tilde{C}_T^U)}. \tag{4-2}$$

Proof. The proof is similar to the one in Lemma 3.11 from [Metcalf et al. 2012], except that we need to estimate the boundary terms at $\tilde{t} = T$ and $\tilde{t} = 2T$.

To keep the ideas clear we first prove the lemma with \square_K replaced by \square . We consider a cutoff function χ supported in $[\frac{8}{20}, \frac{22}{10}]$ which equals 1 on $[\frac{9}{20}, \frac{21}{10}]$. Let

$$\beta(\tilde{t}, \tilde{r}) = \chi\left(\frac{\tilde{r} T}{\tilde{t} R}\right).$$

Note that $\beta \equiv 1$ on C_T^R , and that the restriction of β to $T \leq \tilde{t} \leq 2T$ is supported in \tilde{C}_T^R .

Integrating $\frac{1}{2}\beta \square w^2 = \beta(w \square w + m^{\alpha\beta} \partial_\alpha w \partial_\beta w)$ by parts twice gives

$$\begin{aligned} &\int_T^{2T} \int \beta (|\partial_x w|^2 - |\partial_t w|^2) dx dt \\ &= \int_T^{2T} \int \square w \cdot \beta w dx dt - \frac{1}{2} \int_T^{2T} \int (\square \beta) w^2 dx dt - \int \left(\beta w \partial_t w - \frac{1}{2} \beta_t w^2 \right) dx \Big|_T^{2T}. \end{aligned}$$

Since we can write $w_t = (Sw - x^i \partial_i w)/t$ it follows after integration by parts that

$$\int \beta w \partial_t w dx = \frac{1}{\tilde{t}} \int \beta w Sw dx + \frac{1}{2\tilde{t}} \int w^2 \partial_i (x^i \beta) dx.$$

Since $|\partial_i (x^i \beta)| + \tilde{t} |\partial_t \beta| \leq C$ on the support of β , it follows that the boundary terms are bounded by

$$CT^{-1} (\|w(2T, \cdot)\|_{L^2(\tilde{C}_T^R(2T))}^2 + \|Sw(2T, \cdot)\|_{L^2(\tilde{C}_T^R(2T))}^2 + \|w(T, \cdot)\|_{L^2(\tilde{C}_T^R(T))}^2 + \|Sw(T, \cdot)\|_{L^2(\tilde{C}_T^R(T))}^2).$$

Let $\chi(t/T)$ be another smooth cutoff such that $\chi(2) = 1$ and $\chi(1) = 0$. We write

$$\begin{aligned} w(2T, x)^2 &= \int_{1/2}^1 \frac{d}{ds} (\chi w^2)(s2T, sx) ds \\ &= \int_{1/2}^1 S(\chi w^2)(s2T, sx) \frac{ds}{s} = \int_T^{2T} S(\chi w^2)\left(t, t \frac{x}{2T}\right) \frac{dt}{t} \end{aligned}$$

and thus

$$\|w(2T, \cdot)\|_{L^2(\tilde{C}_T^R(2T))}^2 \lesssim \frac{1}{T} \|S(\chi w^2)(t, x)\|_{L^2(\tilde{C}_T^R)}^2 \lesssim \frac{1}{T} (\|w\|_{L^2(\tilde{C}_T^R)}^2 + \|Sw\|_{L^2(\tilde{C}_T^R)}^2).$$

A similar argument holds for $2T$ replaced by T , and for w replaced by Sw . Hence the boundary term can be estimated by

$$\frac{1}{T^2} \sum_{j=0}^2 \|S^j w\|_{L^2(\tilde{C}_T^R)}^2.$$

To estimate ∂w we use the pointwise inequality

$$|\partial w|^2 \leq \tilde{C} \frac{1}{(\tilde{t} - \tilde{r})^2} |Sw|^2 + \frac{\tilde{t}}{\tilde{t} - \tilde{r}} (|\partial_x w|^2 - |\partial_t w|^2), \tag{4-3}$$

which is valid inside the cone C for a fixed large constant \tilde{C} . Hence

$$\int \beta |\partial w|^2 dx dt \lesssim \int \frac{1}{(\tilde{t} - \tilde{r})^2} \beta |Sw|^2 + \frac{\tilde{t}}{\tilde{t} - \tilde{r}} |\square \beta| w^2 + \frac{\tilde{t}}{\tilde{t} - \tilde{r}} \beta |\square w| |w| dx dt, \tag{4-4}$$

where all weights have a fixed size in the support of β . The function β also satisfies $|\square \beta| \lesssim R^{-2}$. Then the conclusion of the lemma follows by applying Cauchy–Schwarz to the last term.

The argument for C_T^U is similar. We now consider

$$\beta(\tilde{t}, \tilde{r}) = \chi\left(\frac{\tilde{t} - \tilde{r}}{\tilde{t}} \frac{T}{U}\right).$$

We multiply by βw and integrate by parts as above. The boundary terms are now controlled by

$$CU^{-1} (\|w(2T, \cdot)\|_{L^2(\tilde{C}_T^U(2T))}^2 + \|Sw(2T, \cdot)\|_{L^2(\tilde{C}_T^U(2T))}^2 + \|w(T, \cdot)\|_{L^2(\tilde{C}_T^U(T))}^2 + \|Sw(T, \cdot)\|_{L^2(\tilde{C}_T^U(T))}^2),$$

which in turn is controlled, by using the scaling S as above, by

$$\frac{1}{TU} \sum_{j=0}^2 \|S^j w\|_{L^2(\tilde{C}_T^R)}^2.$$

The estimate now follows from (4-4), using the fact that $|\square \beta| \lesssim T^{-1}U^{-1}$.

Now consider the above proof but with \square replaced by \square_K . Integrating

$$\frac{1}{2} \beta \square_K w^2 = \beta (w \square_K w + g_K^{\alpha\beta} \partial_\alpha w \partial_\beta w)$$

by parts twice gives

$$\begin{aligned}
 & - \int_T^{2T} \int \beta g_K^{\alpha\beta} \partial_\alpha w \partial_\beta w \sqrt{|g_K|} dx dt \\
 & = \int_T^{2T} \int \left(\beta w \square_K w - \frac{1}{2} (\square_K \beta) w^2 \right) \sqrt{|g_K|} dx dt - \frac{1}{2} \int (\beta g_K^{0\alpha} \partial_\alpha w^2 - g_K^{\alpha 0} w^2 \partial_\alpha \beta) \sqrt{|g_K|} dx \Big|_T^{2T}.
 \end{aligned}$$

First we estimate the boundary term. The terms with $\alpha = 0$ are handled as before and so is the second term with $\alpha > 0$. For the first term with $\alpha > 0$ we integrate by parts and see that it is bounded by a term of the same form as the second term plus a term of the form

$$\frac{1}{2} \int \beta \partial_\alpha (g_K^{0\alpha} \sqrt{|g_K|}) w^2 dx \lesssim \int \beta r^{-2} w^2 dx,$$

which can be estimated as above. To estimate the interior term we just note that

$$\sqrt{|g_K|} g_K^{\alpha\beta} \partial_\alpha w \partial_\beta w = |\partial_x w|^2 - |\partial_t w|^2 + O(r^{-1}) |\partial w|^2,$$

where the error term can be absorbed in the left of (4-3) for large enough R .

This finishes the proof of (4-1), and (4-2) follows in a similar manner. □

Applying Lemma 4.2 to w_α for some multiindex α , and using (2-2), we obtain the higher-order version of the estimates:

$$\|\partial w_\alpha\|_{L^2(C_T^R)} \lesssim R^{-1} \|w_{|\alpha|+n}\|_{L^2(\tilde{C}_T^R)} + R \|(\square_K w)_{|\alpha|+n}\|_{L^2(\tilde{C}_T^R)}, \tag{4-5}$$

$$\|\partial w_\alpha\|_{L^2(C_T^U)} \lesssim U^{-1} \|w_{|\alpha|+n}\|_{L^2(\tilde{C}_T^U)} + T \|(\square_K w)_{|\alpha|+n}\|_{L^2(\tilde{C}_T^U)}. \tag{4-6}$$

Combining the two estimates above (4-5) and (4-6) with the Sobolev embeddings from Lemma 3.1 and the pointwise estimate for second-order derivatives in Lemma 3.3 we obtain:

Corollary 4.3. *For all $T \geq 1$ and $1 \leq R, U \leq \frac{1}{4}T$ we have for some n independent of α*

$$\|\partial w_\alpha\|_{L^\infty(C_T^R)} \lesssim \frac{1}{R} \|w_{\leq|\alpha|+n}\|_{L^\infty(\tilde{C}_T^R)} + R \|(\square_K w)_{|\alpha|+n}\|_{L^\infty(\tilde{C}_T^R)},$$

respectively,

$$\|\partial w_\alpha\|_{L^\infty(C_T^U)} \lesssim \frac{1}{U} \|w_{\leq|\alpha|+n}\|_{L^\infty(\tilde{C}_T^U)} + T \|(\square_K w)_{|\alpha|+n}\|_{L^\infty(\tilde{C}_T^U)}.$$

Finally, we will derive a sharp estimate for the bad first-order derivative, following [Lindblad 1990].

Lemma 4.4. *Let $D_t = \{x : 0 \leq t - |x| \leq \frac{1}{4}t\}$, $C_t^q = \{x : t - |x| = q\}$, and let $\bar{w}(q)$ be any positive continuous function, where $q = t - r$. Suppose that $\square\phi = F$. Then the following holds in D_t , $t \geq 1$:*

$$\begin{aligned}
 t|\partial\phi(t, x)\bar{w}(q)| \lesssim & \sup_{4q \leq \tau \leq t} \left(\|q\partial\phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \sum_{|I| \leq 1} \|Z^I \phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} \right) \\
 & + \int_{4q}^t \left(\langle \tau \rangle \|F(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \sum_{|I|+|J| \leq 2} \langle \tau \rangle^{-1} \|\partial^I \Omega^J \phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} \right) d\tau.
 \end{aligned}$$

Proof. We write

$$\square\phi = -\frac{1}{r} \partial_v \partial_u (r\phi) + \frac{1}{r^2} \Delta_\omega \phi,$$

where $\partial_u = \partial_t - \partial_r$ and $\partial_v = \partial_t + \partial_r$. Hence in D_t

$$|\partial_v \partial_u(r\phi)| \lesssim |r\Box\phi| + \langle r \rangle^{-1} \sum_{|I|+|J|\leq 2} |\partial^I \Omega^J \phi| \lesssim |\langle t \rangle \Box\phi| + \langle t \rangle^{-1} \sum_{|I|+|J|\leq 2} |\partial^I \Omega^J \phi|. \tag{4-7}$$

Integrating this along the flow lines of the vector field ∂_v from the boundary of $D = \bigcup_{\tau \geq 0} D_\tau$ to any point inside D_t for $t \geq 1$. Using that \bar{w} is constant along the flow lines, and (4-7), we obtain

$$\begin{aligned} & |\partial_u(r\phi(t, x))\bar{w}(q)| \\ & \lesssim |\partial_u(r\phi)(4q, 3q)\bar{w}(q)| + \int_{4q}^t \left(\langle \tau \rangle \|F(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \sum_{|I|+|J|\leq 2} \langle \tau \rangle^{-1} \|\partial^I \Omega^J \phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} \right) d\tau. \end{aligned}$$

Moreover

$$t|\partial_u\phi(t, x)\bar{w}(q)| \lesssim |\partial_u(r\phi(t, x))\bar{w}(q)| + |\phi(t, x)\bar{w}(q)|$$

and

$$|\partial_u(r\phi)(4q, 3q)\bar{w}(q)| \lesssim |q\partial_u\phi(4q, 3q)\bar{w}(q)| + |\phi(4q, 3q)\bar{w}(q)|.$$

The last three inequalities yield

$$\begin{aligned} t|\partial_u\phi(t, x)\bar{w}(q)| & \lesssim \sup_{4q \leq \tau \leq t} (\|q\partial\phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \|\phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)}) \\ & \quad + \int_{4q}^t \left(\langle \tau \rangle \|F(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \sum_{|I|+|J|\leq 2} \langle \tau \rangle^{-1} \|\partial^I \Omega^J \phi(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} \right) d\tau. \end{aligned}$$

The lemma follows from also using that $r|\partial\phi| \lesssim |r\partial_q\phi| + |S\phi| + |\Omega\phi|$. □

5. Pointwise estimates from the Minkowski fundamental solution

In this section, we translate pointwise bounds on the inhomogeneous terms into pointwise bounds for the solution by using the fundamental solution of the Minkowski metric.

For any $\beta, \gamma, \eta \in \mathbb{R}$, we define the weighted L^∞ norms

$$\|G\|_{L_{\beta,\gamma,\eta}^\infty} = \|\langle r \rangle^\beta \langle t \rangle^\gamma \langle t-r \rangle^\eta H(t, r)\|_{L_{t,r}^\infty}, \quad H(t, r) = \sum_0^2 \|\Omega^i G(t, r\omega)\|_{L^2(\mathbb{S}^2)}.$$

We use the following lemma (see Section 6 of [Tohaneanu 2022]).

Lemma 5.1. *Let ψ solve*

$$\Box\psi = G, \quad \psi(0) = 0, \quad \partial_t\psi(0) = 0,$$

where G is supported in $\{|x| \leq t + R_0\}$. Assume also that $2 \leq \beta \leq 3$ and $\eta \geq -\frac{1}{2}$. We define, for any arbitrary $\delta > 0$,

$$\tilde{\eta} = \begin{cases} \eta - \delta - 2, & \eta < 1, \\ -1, & \eta > 1. \end{cases}$$

(i) *If $\gamma \geq 0$, we have*

$$r\psi(t, x) \lesssim \frac{1}{\langle t-r \rangle^{\beta+\gamma+\tilde{\eta}-1}} \|G\|_{L_{\beta,\gamma,\eta}^\infty}. \tag{5-1}$$

(ii) If $\gamma < 0$, we have

$$r\psi(t, x) \lesssim \frac{\langle t \rangle^{-\gamma}}{\langle t-r \rangle^{\beta+\tilde{\eta}-1}} \|G\|_{L^\infty_{\beta,\gamma,\eta}}. \tag{5-2}$$

(iii) If $\eta > 1$, we have

$$r\psi(t, x) \lesssim \left\langle \ln \frac{\langle t \rangle}{\langle t-r \rangle} \right\rangle \|G\|_{L^\infty_{2,0,\eta}}. \tag{5-3}$$

Proof. Note first that, after a translation in time, we may assume that $R_0 = 0$.

We use the ideas from [Metcalf et al. 2012]. Define

$$H(t, r) = \sum_0^2 \|\Omega^i G(t, r\omega)\|_{L^2(\mathbb{S}^2)}.$$

By Sobolev embeddings on the sphere, we have $|G| \lesssim H$. Let v be the radial solution to

$$\square v = H, \quad v[0] = 0.$$

By the positivity of the fundamental solution, we have that $|\psi| \lesssim |v|$. On the other hand, we can write v explicitly:

$$rv(t, r) = \frac{1}{2} \int_{D_{tr}} \rho H(s, \rho) ds d\rho,$$

where D_{tr} is the rectangle

$$D_{tr} = \{0 \leq s - \rho \leq t - r, t - r \leq s + \rho \leq t + r\}.$$

We partition the set D_{tr} into a double dyadic manner as

$$D_{tr} = \bigcup_{R \leq t} D_{tr}^R, \quad D_{tr}^R = D_{tr} \cap \{R < r < 2R\}$$

and estimate the corresponding parts of the above integral.

We clearly have

$$\int_{D_{tr}^R} \rho H ds d\rho \lesssim \|G\|_{L^\infty_{\beta,\gamma,\eta}} \int_{D_{tr}^R} \rho^{1-\beta} \langle s \rangle^{-\gamma} \langle s - \rho \rangle^{-\eta} d\rho ds.$$

We now consider two cases:

(i) $R < \frac{1}{8}(t - r)$. Here we have $\rho \sim R$ and $s \approx s - \rho \approx \langle t - r \rangle$; therefore we obtain

$$\int_{D_{tr}^R} \rho^{1-\beta} \langle s \rangle^{-\gamma} \langle s - \rho \rangle^{-\eta} d\rho ds \lesssim R^{3-\beta} \langle t - r \rangle^{-\gamma-\eta},$$

and after summation, using that $\beta \leq 3$, we obtain

$$\sum_{R < (t-r)/8} \int_{D_{tr}^R} \rho H ds d\rho \lesssim \frac{\ln \langle t - r \rangle \langle t - r \rangle^{3-\beta}}{\langle t - r \rangle^{\gamma+\eta}} \lesssim \frac{1}{\langle t - r \rangle^{\beta+\tilde{\eta}}},$$

which is the desired bound in all cases.

(ii) $\frac{1}{8}(t-r) < R < t$. Here we have $\rho \sim R$ and $t \geq s \gtrsim R$. Let $u = s - \rho$.

Assume first that $\gamma \geq 0$; then

$$\int_{D_{tr}^R} \rho^{1-\beta} \langle s \rangle^{-\gamma} \langle s - \rho \rangle^{-\eta} d\rho ds \lesssim R^{2-\beta-\gamma} \int_0^{t-r} \langle u \rangle^{-\eta} du \lesssim R^{2-\beta-\gamma} \langle t-r \rangle^{\mu(\eta)},$$

where

$$\mu(\eta) = \begin{cases} 1 - \eta, & \eta < 1, \\ 0, & \eta > 1. \end{cases}$$

If $\beta + \gamma > 2$, we obtain after summation

$$\sum_{R > (t-r)/8} \int_{D_{tr}^R} \rho H ds d\rho \lesssim \langle t-r \rangle^{2-\beta-\gamma+\mu(\eta)},$$

which is (5-1).

Assume now that $\beta = 2$ and $\gamma = 0$. Equation (5-3) is obvious when $t \leq 1$. When $t \geq 1$, we see that there are $\ln(t/(t-r))$ dyadic regions when $\frac{1}{8}(t-r) < R < t$, so we obtain (5-3) after summation.

Finally, if $\gamma < 0$ we obtain

$$\int_{D_{tr}^R} \rho^{1-\beta} \langle s \rangle^{-\gamma} \langle s - \rho \rangle^{-\eta} d\rho ds \lesssim R^{2-\beta} \langle t \rangle^{-\gamma} \int_0^{t-r} \langle u \rangle^{-\eta} du \lesssim R^{2-\beta} \langle t \rangle^{-\gamma} \langle t-r \rangle^{\mu(\eta)}.$$

Since $\beta \geq 2$, we obtain after summation

$$\sum_{R > (t-r)/8} \int_{D_{tr}^R} \rho H ds d\rho \lesssim \langle t \rangle^{-\gamma} \langle t-r \rangle^{2-\beta+\mu(\eta)},$$

which is (5-2). □

6. Setup for pointwise estimates

In this section, we will slightly adjust \square_K to an operator closer to \square (with respect to the (\tilde{t}, x) -coordinates). Indeed, we let

$$P = |g_K|^{1/4} (-g_K^{\tilde{t}\tilde{t}})^{-1/2} \square_K (-g_K^{\tilde{t}\tilde{t}})^{-1/2} |g_K|^{-1/4}.$$

P is self-adjoint with respect to $d\tilde{t} dx$. More importantly, a quick computation yields that

$$P = \partial_\alpha (g_K^{\alpha\beta} (-g_K^{\tilde{t}\tilde{t}}) \partial_\beta) + V, \quad V = |g_K|^{1/4} (-g_K^{\tilde{t}\tilde{t}})^{-1/2} \square_K ((-g_K^{\tilde{t}\tilde{t}})^{-1/2} |g_K|^{-1/4}).$$

It is easy to see that $V \in S^Z(r^{-3})$.

Let us first consider the Schwarzschild metric. In this case we have that, for large r , $-g_S^{\tilde{t}\tilde{t}} = g_S^{r^*r^*}$ and $\tilde{g}_S^{r^*} = 0$. We thus have

$$P = \square + P_{lr},$$

where the long-range spherically symmetric part P_{lr} has the form

$$P_{lr} = g_{lr}(r) \Delta_\omega + V, \quad g_{lr} \in S^Z(r^{-3}), \quad V \in S^Z(r^{-3}). \tag{6-1}$$

For the Kerr metric, we use the fact that the metric coefficients have the following properties:

$$g_K^{\alpha\beta} - g_S^{\alpha\beta} \in S^Z(r^{-2}), \tag{6-2}$$

$$\partial g_K \in S^Z(r^{-2}), \quad \partial^2 g_K \in S^Z(r^{-3}). \tag{6-3}$$

Using (6-1) and (6-2) we see that we can write

$$P = \square + P_{lr} + P_{sr}, \tag{6-4}$$

where the short-range part P_{sr} has the form

$$P_{sr} = \partial_\alpha g_{sr}^{\alpha\beta} \partial_\beta, \quad g_{sr}^{\alpha\beta} \in S^Z(r^{-2}). \tag{6-5}$$

Using (6-3) we see that for any function ϕ we have

$$P\phi = (-g_K^{\tilde{t}\tilde{t}})\square_K\phi + h_1\phi + h_2\partial\phi, \quad h_1 \in S^Z(r^{-3}), \quad h_2 \in S^Z(r^{-2}). \tag{6-6}$$

Now pick any multiindex α . After commuting with vector fields, using (6-4), (6-1), and (6-5), we obtain

$$P\phi_\alpha \in S^Z(1)(\square_K\phi)_{\leq|\alpha|} + S^Z(r^{-3})\phi_{\leq|\alpha|+6} + S^Z(r^{-2})\partial\phi_{\leq|\alpha|+5},$$

which in turn implies, using (6-4),

$$\square\phi_\alpha \in S^Z(1)(\square_K\phi)_{\leq|\alpha|} + S^Z(r^{-3})\phi_{\leq|\alpha|+6} + S^Z(r^{-2})\partial\phi_{\leq|\alpha|+5}. \tag{6-7}$$

Moreover, by finite speed of propagation, and the assumption on the support of the initial data, the right-hand side is supported in the forward light cone $\{|x| < \tilde{t} + R_0\}$.

We will use (6-7) in the next section to extract more decay for the solution.

7. The bootstrap argument for the Einstein model

We now prove Theorem 1.1 by using a bootstrap argument. We first write

$$\mathcal{E}_N(0) = \mu_N \epsilon,$$

where $\mu_N > 0$ is a fixed, small N -dependent constant to be determined below (see (7-5), (7-6)).

Let $N_1 = \frac{1}{2}N$. We will assume that the following a priori bounds hold for some large constant \tilde{C} independent of ϵ and \tilde{t} , and a fixed small $\delta > 0$:

$$\mathcal{E}_N(\tilde{t}) \leq \tilde{C}\mu_N\epsilon(\tilde{t})^\delta, \tag{7-1}$$

$$|\phi_{\leq N_1+2}| \leq \frac{\epsilon(\ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle))}{\langle \tilde{t} \rangle}, \quad |\partial\phi_{\leq N_1+2}| \leq \frac{\epsilon(\ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle))}{r\langle \tilde{t} - \tilde{r} \rangle}, \tag{7-2}$$

$$|(\partial\phi_{TU})_{\leq N_1+2}| \leq \frac{\epsilon}{\langle \tilde{t} \rangle}. \tag{7-3}$$

Clearly (7-1), (7-2) and (7-3) hold for small times. We assume now that the bounds hold on some time interval $0 \leq \tilde{t} \leq T$, and we improve the constants by $\frac{1}{2}$. By the continuity method this implies that the solution exists globally, and that the bounds also hold globally.

In order to improve (7-1), we show that, for small enough ϵ , there is C_N independent of T so that

$$\mathcal{E}_N(\tilde{t}) \leq C_N \langle \tilde{t} \rangle^{C_N \epsilon} \mathcal{E}_N(0), \quad 0 \leq \tilde{t} \leq T. \tag{7-4}$$

If we now additionally take $\tilde{C} = 2C_N$ and $\epsilon < \delta/C_N$, we thus improve the a priori bound for $\mathcal{E}_N(\tilde{t})$ to

$$\mathcal{E}_N(\tilde{t}) \leq \frac{1}{2} \tilde{C} \mu_N \epsilon \langle \tilde{t} \rangle^\delta.$$

In order to improve the pointwise bounds, we will show that, for some fixed positive integer m , independent of N , we have

$$|\phi_{\leq N-m}| \lesssim \frac{\mathcal{E}_N(0) \langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{\langle \tilde{t} \rangle}, \quad |\partial \phi_{\leq N-m}| \lesssim \frac{\mathcal{E}_N(0) \langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{r \langle \tilde{t} - \tilde{r} \rangle}, \tag{7-5}$$

$$|(\partial \phi_{TU})_{\leq N-m}| \lesssim \frac{\mathcal{E}_N(0)}{r \langle \tilde{t} - \tilde{r} \rangle^{1-\delta}}. \tag{7-6}$$

We can now pick a small μ_N to improve (7-2) and (7-3).

7.1. The energy estimates. We will now use assumptions (7-2) and (7-3) to show (7-4) for small enough ϵ . By Gronwall's inequality and (2-8), it is enough to show that

$$\|(\square_K \phi)_{\leq N}\|_{LE_w^*[0, \tilde{t}]} \lesssim \int_0^{\tilde{t}} \frac{\epsilon}{\langle \tau \rangle} \mathcal{E}_N(\tau) d\tau + \epsilon \mathcal{E}_N(\tilde{t}). \tag{7-7}$$

We can write, using (2-4), (2-5) and (2-6),

$$\square_K \phi \in S^Z(1)(\partial \phi_{TU})^2 + S^Z(1)\partial \phi \bar{\partial} \phi.$$

After commuting with vector fields, and using (2-3), we also get that

$$(\square_K \phi)_{\leq N} \lesssim (\partial \phi_{TU})_{\leq N_1} (\partial \phi_{TU})_{\leq N} + \partial \phi_{\leq N_1} \bar{\partial} \phi_{\leq N} + \bar{\partial} \phi_{\leq N_1} \partial \phi_{\leq N} + r^{-1} \partial \phi_{\leq N_1} \partial \phi_{\leq N-1}. \tag{7-8}$$

The first term is easy. By (7-3) we have

$$\|(\partial \phi_{TU})_{\leq N_1} (\partial \phi_{TU})_{\leq N}\|_{L^1[0, \tilde{t}]L^2} \lesssim \int_0^{\tilde{t}} \frac{\epsilon}{\langle \tau \rangle} \mathcal{E}_N(\tau) d\tau.$$

Similarly, the last term can be estimated in L^1L^2 . Indeed, we note that (7-2) implies

$$|r^{-1} \partial \phi_{\leq N_1}| \lesssim \frac{\epsilon}{\langle \tilde{t} \rangle},$$

and thus

$$\|r^{-1} \partial \phi_{\leq N_1} \partial \phi_{\leq N-1}\|_{L^1[0, \tilde{t}]L^2} \lesssim \int_0^{\tilde{t}} \frac{\epsilon}{\langle \tau \rangle} \mathcal{E}_N(\tau) d\tau.$$

For the second term, we divide it into two parts. When $r < R_1$ we have by (7-2)

$$\|\partial \phi_{\leq N_1} \bar{\partial} \phi_{\leq N}\|_{L^1[0, \tilde{t}]L^2(r < R_1)} \lesssim \int_0^{\tilde{t}} \frac{\epsilon}{\langle \tau \rangle} \mathcal{E}_N(\tau) d\tau.$$

When $r > R_1$, we use (7-2) and the last term in (2-7):

$$\begin{aligned} \|\partial\phi_{\leq N_1} \bar{\partial}\phi_{\leq N}\|_{LE^*[0,\tilde{t}]}^2 &\lesssim \int_0^{\tilde{t}} \int_{r>R_1} \frac{\epsilon^2 \tau^{2\delta}}{r \langle \tau - \tilde{r} \rangle^{2+2\delta}} |\bar{\partial}\phi_{\leq N}|^2 dV \\ &\lesssim \|\epsilon \langle \tau - \tilde{r} \rangle^{(-1-\delta_1)/2} \bar{\partial}\phi_{\leq N}\|_{L^2[0,\tilde{t}]L^2(r\geq R_1)}^2 \lesssim (\epsilon \mathcal{E}_N(\tilde{t}))^2. \end{aligned}$$

For the third term, note that (2-1) and (7-2) imply that

$$|\bar{\partial}\phi_{\leq N_1}| \lesssim \frac{\epsilon}{\langle \tilde{t} \rangle}. \tag{7-9}$$

Using (7-9) gives

$$\|\bar{\partial}\phi_{\leq N_1} \partial\phi_{\leq N}\|_{L^1[0,\tilde{t}]L^2} \lesssim \int_0^{\tilde{t}} \frac{\epsilon}{\langle \tau \rangle} \mathcal{E}_N(\tau) d\tau.$$

Putting all these together we obtain (7-7).

7.2. The decay estimates. We now show that (7-5) and (7-6) hold.

The proof uses an iteration procedure. The most important part here is to obtain pointwise decay rates of \tilde{t}^{-1} near the trapped set for all components. We start with a weak decay rate of $\tilde{t}^{-1/2+C\epsilon}$ given by the slow growth $\tilde{t}^{C\epsilon}$ combined with the results of Section 3. We then use Lemma 5.1 to improve decay in r , followed by Corollary 4.3 to improve the decay of derivatives. Lemma 4.1 then allows us to turn the r -decay into \tilde{t} -decay. This yields an improved global decay rate of $\tilde{t}^{-1+C\epsilon}$, which is barely not enough. We then use Lemma 4.4 to improve the decay of the derivative of the good components $\partial\phi_{TU}$ to \tilde{t}^{-1} near the cone. We can now go back to the iteration procedure, and use the improved bounds combined with Lemma 5.1, Corollary 4.3 and Lemma 4.1 to improve the decay rate of all components to \tilde{t}^{-1} away from the cone. This finishes the proof.

Let $N_2 = N - 13$. We first note that (3-3) and (3-4), combined with the energy bounds (7-4), yield the weak pointwise bounds

$$|\partial\phi_{\leq N_2}| \lesssim \frac{\langle \tilde{t} \rangle^{C\epsilon} \mathcal{E}_N(0)}{r \langle \tilde{t} - \tilde{r} \rangle^{1/2}}, \quad |\phi_{\leq N_2}| \lesssim \frac{\langle \tilde{t} - \tilde{r} \rangle^{1/2} \mathcal{E}_N(0)}{\langle \tilde{t} \rangle^{1-C\epsilon}}. \tag{7-10}$$

We now need to improve the decay of $\phi_{\leq N-m}$ and $\partial\phi_{\leq N-m}$. To that extent, we will use Lemma 5.1, followed by Lemma 4.1 and Corollary 4.3.

We cannot apply Lemma 5.1 directly. On one hand, we have no control on the solution for $r \ll 2M$, and on the other hand, the initial data is not trivial. Instead, let

$$\chi = \chi_1(\tilde{r})\chi_2(\tilde{t}).$$

Here $\chi_1 \equiv 1$ for $\tilde{r} \geq R \gg M$ and supported in $\tilde{r} \geq \frac{1}{2}R$, while $\chi_2 \equiv 1$ for $\tilde{t} \geq 1$ and supported in $\tilde{t} \geq \frac{1}{2}$.

We now consider $\psi_{\alpha\beta} = \chi\phi_{\alpha\beta}$. Using (6-7), we see that ψ satisfies the system

$$\square(\psi_{\leq n}) = G_n, \quad G_n \in S^Z(r^{-2})\partial\phi_{\leq n+5} + S^Z(r^{-3})\phi_{\leq n+6} + S^Z(1)(\partial\phi_{\leq n})^2,$$

with trivial initial data, and G_n supported in the region $r \geq \frac{1}{2}R$. Using (7-10), we see that, for all $n \leq N_3 := N_2 - 12$, we have

$$G_{n+6} \lesssim \mathcal{E}_N(0) \left(\frac{\langle \tilde{t} \rangle^{C\epsilon}}{r^3 \langle \tilde{t} - \tilde{r} \rangle^{1/2}} + \frac{\langle \tilde{t} - \tilde{r} \rangle^{1/2}}{r^3 \langle \tilde{t} \rangle^{1-C\epsilon}} + \frac{\langle \tilde{t} \rangle^{C\epsilon}}{r^2 \langle \tilde{t} - \tilde{r} \rangle} \right).$$

We now apply Lemma 5.1. The first term on the right-hand side is controlled by the other two terms. For the second term we use (5-1) with $\beta = 3$, $\gamma = 1 - C\epsilon$ and $\eta = -\frac{1}{2}$. For the third term, we use (5-1) with $\beta = 2$, $\gamma = -C\epsilon$ and $\eta = 1 - C\epsilon$. We obtain

$$|\phi_{\leq N_3}| \lesssim \frac{\langle \tilde{t} \rangle^{C\epsilon}}{r} \mathcal{E}_N(0). \tag{7-11}$$

We now plug in the bounds (7-11) and (7-10) into Corollary 4.3. We thus obtain for $N_4 = N_3 - n$ with n from Corollary 4.3:

$$\begin{aligned} \|\partial\phi_{N_4}\|_{L^\infty(C_T^R)} &\lesssim \frac{1}{R} \frac{T^{C\epsilon}}{R} \mathcal{E}_N(0) + R \left(\frac{T^{C\epsilon}}{RT^{1/2}} \mathcal{E}_N(0) \right)^2 \lesssim \frac{T^{C\epsilon}}{R^2} \mathcal{E}_N(0), \\ \|\partial\phi_{N_4}\|_{L^\infty(C_T^U)} &\lesssim \frac{1}{U} \frac{T^{C\epsilon}}{R} \mathcal{E}_N(0) + T \left(\frac{T^{C\epsilon}}{RU^{1/2}} \mathcal{E}_N(0) \right)^2 \lesssim \frac{T^{C\epsilon}}{RU} \mathcal{E}_N(0). \end{aligned}$$

The last two inequalities can be written as

$$|\partial\phi_{\leq N_4}| \lesssim \frac{\langle \tilde{t} \rangle^{1+C\epsilon}}{r^2 \langle \tilde{t} - \tilde{r} \rangle} \mathcal{E}_N(0). \tag{7-12}$$

We now use Lemma 4.1. Note that (7-11) and (7-12) yield

$$\|\langle r \rangle \phi_{\leq N_4}\|_{LE^1(C_T^{<T/2})} \lesssim T^{1/2+C\epsilon} \mathcal{E}_N(0).$$

Moreover, (7-10) implies that

$$\|\langle \square_K \phi \rangle_{\leq N_4}\|_{LE^*(C_T^{<T/2})} \lesssim T^{-1/2+C\epsilon} \mathcal{E}_N(0).$$

The two inequalities above and Lemma 4.1 with $N_5 = N_4 - n$ give us

$$\|\phi_{\leq N_5}\|_{LE^1(C_T^{<T/2})} \lesssim T^{-1/2+C\epsilon} \mathcal{E}_N(0),$$

which combined with the Sobolev embeddings from Lemma 3.1 give for $N_6 = N_5 - 13$

$$|\phi_{\leq N_6}| \lesssim \langle \tilde{t} \rangle^{-1+C\epsilon} \mathcal{E}_N(0). \tag{7-13}$$

We now plug in the bounds (7-13) and (7-10) into Corollary 4.3. We thus obtain for $N_7 = N_6 - n$

$$\|\partial\phi_{\leq N_7}\|_{L^\infty(C_T^R)} \lesssim \frac{1}{R} \frac{T^{C\epsilon}}{T} \mathcal{E}_N(0) + R \left(\frac{T^{C\epsilon}}{RT^{1/2}} \mathcal{E}_N(0) \right)^2 \lesssim \frac{T^{C\epsilon}}{RT} \mathcal{E}_N(0).$$

Combined with (7-12), this gives

$$|\partial\phi_{\leq N_7}| \lesssim \frac{\langle \tilde{t} \rangle^{C\epsilon}}{r \langle \tilde{t} - \tilde{r} \rangle} \mathcal{E}_N(0). \tag{7-14}$$

Note also that (2-1), (7-13) and (7-14) give

$$|\bar{\partial}\phi_{\leq N_7-2}| \lesssim \frac{\langle \tilde{t} \rangle^{C\epsilon}}{r \langle \tilde{t} \rangle} \mathcal{E}_N(0). \tag{7-15}$$

Equations (7-13) and (7-14) almost finish the proof of (7-5), except that we need to replace $\langle \tilde{t} \rangle^{C\epsilon}$ by $\langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle$.

We now prove the fact that ψ_{TU} actually satisfies better decay estimates. Indeed, note first that

$$\square(T^\alpha U^\beta \phi_{\alpha\beta}) - T^\alpha U^\beta \square\phi_{\alpha\beta} \in S^Z(r^{-2})\phi_{\leq 1}.$$

Using (6-4) and (6-6) we obtain

$$\square\phi_{TU} \in S^Z(1)(\square_K\phi)_{TU} + S^Z(r^{-2})\phi_{\leq 6}.$$

Moreover,

$$(\square_K\phi)_{TU} \in S^Z(1)\partial\phi\bar{\partial}\phi,$$

and thus

$$\square\phi_{TU} \in S^Z(1)\partial\phi\bar{\partial}\phi + S^Z(r^{-2})\phi_{\leq 6}.$$

After commuting with vector fields (in particular using (2-3)) and applying the cutoff we thus obtain

$$\square(\psi_{TU})_{\leq m} = H_m, \quad H_m \in S^Z(r^{-2})\phi_{\leq m+6} + S^Z(1)\partial\phi_{\leq m}\bar{\partial}\phi_{\leq m} + S^Z(r^{-1})(\partial\phi_{\leq m})^2.$$

Using (7-13), (7-14) and (7-15), we see that

$$|H_m| \lesssim \frac{\mathcal{E}_N(0)}{r^2 \langle \tilde{t} \rangle^{1-C\epsilon}}, \quad m \leq N_7 - 2. \tag{7-16}$$

Let $N_8 = N_7 - 6$. We now apply Lemma 4.4 with $\bar{w}(q) = \langle q \rangle^{1-\delta}$ to $(\psi_{TU})_{\leq N_8}$. Note first that, due to (7-14) and (7-13) we have

$$\sup_{4q \leq \tau \leq \tilde{t}} \left(\|q \partial\phi_{\leq N_8}(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} + \sum_{|I| \leq 1} \|Z^I \phi_{\leq N_8}(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} \right) \lesssim \mathcal{E}_N(0).$$

Moreover, (7-13) implies that

$$\int_{4q}^{\tilde{t}} \sum_{|I| \leq 2} \langle \tau \rangle^{-1} \|\Omega^I \phi_{\leq N_8}(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} d\tau \lesssim \int_{4q}^{\tilde{t}} \langle \tau \rangle^{-1} \frac{\langle q \rangle^{1-\delta} \mathcal{E}_N(0)}{\langle \tau \rangle^{1-C\epsilon}} d\tau \lesssim \mathcal{E}_N(0).$$

Finally, we obtain by (7-16) that

$$\int_{4q}^{\tilde{t}} \langle \tau \rangle \|H_m(\tau, \cdot)\bar{w}\|_{L^\infty(C_\tau^q)} d\tau \lesssim \int_{4q}^{\tilde{t}} \langle \tau \rangle \frac{\langle q \rangle^{1-\delta} \mathcal{E}_N(0)}{\langle \tau \rangle^{3-C\epsilon}} d\tau \lesssim \mathcal{E}_N(0).$$

Lemma 4.4 thus implies, in conjunction with (7-14), that

$$|\partial(\psi_{TU})_{\leq N_8}| \lesssim \frac{\mathcal{E}_N(0)}{r \langle \tilde{t} - \tilde{r} \rangle^{1-\delta}}. \tag{7-17}$$

This finishes the proof of (7-6).

Finally, to obtain a decay rate of $1/\tilde{t}$ in the interior, we see that, using (6-7) and (7-8), we can write our system as

$$\square(\psi_{\leq m}) = J_m,$$

$$J_m \in S^Z(r^{-2})\partial\phi_{\leq m+5} + S^Z(r^{-3})\phi_{\leq m+6} + S^Z(1)(\partial\phi_{TU})_{\leq m}^2 + S^Z(1)\partial\phi_{\leq m}\bar{\partial}\phi_{\leq m} + S^Z(r^{-1})(\partial\phi_{\leq m})^2,$$

and J_m is supported in the region $\{\tilde{t} \geq \frac{1}{2}, \tilde{r} \geq \frac{1}{2}R\}$. Due to the improved bounds (7-13), (7-14) and (7-17) we obtain

$$|J_{m+6}| \lesssim \mathcal{E}_N(0) \left(\frac{\langle \tilde{t} \rangle^{C\epsilon}}{r^3 \langle \tilde{t} - \tilde{r} \rangle} + \frac{1}{r^2 \langle \tilde{t} - \tilde{r} \rangle^{2-2\delta}} \right), \quad m \leq N_9 := N_8 - 8.$$

We now apply Lemma 5.1 and in particular (5-3) to control the last term. We obtain

$$|\psi_{\leq N_9}| \lesssim \frac{\langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{r} \mathcal{E}_N(0). \tag{7-18}$$

Corollary 4.3 thus implies, with $N_{10} = N_9 - n$,

$$|\partial\psi_{\leq N_{10}}| \lesssim \frac{\langle \ln(\langle \tilde{t} \rangle / \langle \tilde{t} - \tilde{r} \rangle) \rangle}{r \langle \tilde{t} - \tilde{r} \rangle} \mathcal{E}_N(0). \tag{7-19}$$

Equations (7-18) and (7-19) finish the proof of (7-5) when $\tilde{r} \geq \frac{1}{2}\tilde{t}$.

All that is left is to replace r by \tilde{t} in the region $\tilde{r} \leq \frac{1}{2}\tilde{t}$. Note first that (7-18) and (7-19), combined with (7-13) and (7-14), yield the (relatively weak) bound

$$|\phi_{\leq N_{10}}| \lesssim \frac{1}{r} \mathcal{E}_N(0), \quad \partial\phi_{\leq N_{10}} \lesssim \frac{1}{r^2} \mathcal{E}_N(0), \quad \tilde{r} < \frac{3\tilde{t}}{4}. \tag{7-20}$$

We now use Lemma 4.1. Note that (7-20) gives

$$\|\langle r \rangle \phi_{\leq N_{10}}\|_{\text{LE}^1(C_T^{<T/2})} \lesssim T^{1/2} \mathcal{E}_N(0).$$

Moreover, (7-14) implies that

$$\|(\square_K \phi)_{\leq N_{10}}\|_{\text{LE}^*(C_T^{<T/2})} \lesssim T^{-1/2} \mathcal{E}_N(0).$$

The two inequalities above and Lemma 4.1 give us, for $N_{11} = N_{10} - n$,

$$\|\phi_{\leq N_{11}}\|_{\text{LE}^1(C_T^{<T/2})} \lesssim T^{-1/2} \mathcal{E}_N(0),$$

which, combined with the Sobolev embeddings from Lemma 3.1 with $N_{12} = N_{11} - 13$, gives

$$|\phi_{\leq N_{12}}| \lesssim \frac{\mathcal{E}_N(0)}{\langle \tilde{t} \rangle}, \quad \tilde{r} \leq \frac{\tilde{t}}{2}.$$

Finally, one last application of Corollary 4.3 with $N_{13} = N_{12} - n$ gives

$$|\partial\phi_{\leq N_{13}}| \lesssim \frac{\mathcal{E}_N(0)}{r \langle \tilde{t} \rangle}, \quad \tilde{r} \leq \frac{\tilde{t}}{2}.$$

This finishes the proof of (7-5) if we pick N large enough so that $N_{13} \geq N_1$.

Acknowledgments

Lindblad was supported in part by NSF grant DMS-1500925 and Simons collaboration grant 638955. Tohaneanu was supported in part by the Simons collaboration grant 586051. We would also like to thank the Mittag Leffler Institute for their hospitality during the Fall 2019 program in geometry and relativity.

References

- [Alinhac 2003] S. Alinhac, “An example of blowup at infinity for a quasilinear wave equation”, pp. 1–91 in *Autour de l’analyse microlocale*, edited by G. Lebeau, Astérisque **284**, Soc. Math. France, Paris, 2003. MR Zbl
- [Andersson and Blue 2015] L. Andersson and P. Blue, “Hidden symmetries and decay for the wave equation on the Kerr spacetime”, *Ann. of Math. (2)* **182**:3 (2015), 787–853. MR Zbl
- [Angelopoulos et al. 2020] Y. Angelopoulos, S. Aretakis, and D. Gajic, “Nonlinear scalar perturbations of extremal Reissner–Nordström spacetimes”, *Ann. PDE* **6**:2 (2020), art. id. 12. MR Zbl
- [Aretakis 2012] S. Aretakis, “Decay of axisymmetric solutions of the wave equation on extreme Kerr backgrounds”, *J. Funct. Anal.* **263**:9 (2012), 2770–2831. MR Zbl
- [Blue and Soffer 2003] P. Blue and A. Soffer, “Semilinear wave equations on the Schwarzschild manifold, I: Local decay estimates”, *Adv. Differential Equations* **8**:5 (2003), 595–614. MR Zbl
- [Blue and Soffer 2005] P. Blue and A. Soffer, “The wave equation on the Schwarzschild metric, II: Local decay for the spin-2 Regge–Wheeler equation”, *J. Math. Phys.* **46**:1 (2005), art. id. 012502. MR Zbl
- [Blue and Sterbenz 2006] P. Blue and J. Sterbenz, “Uniform decay of local energy and the semi-linear wave equation on Schwarzschild space”, *Comm. Math. Phys.* **268**:2 (2006), 481–504. MR Zbl
- [Bony and Häfner 2010] J.-F. Bony and D. Häfner, “The semilinear wave equation on asymptotically Euclidean manifolds”, *Comm. Partial Differential Equations* **35**:1 (2010), 23–67. MR Zbl
- [Christodoulou 1986] D. Christodoulou, “Global solutions of nonlinear hyperbolic equations for small initial data”, *Comm. Pure Appl. Math.* **39**:2 (1986), 267–282. MR Zbl
- [Christodoulou and Klainerman 1993] D. Christodoulou and S. Klainerman, *The global nonlinear stability of the Minkowski space*, Princeton Math. Ser. **41**, Princeton Univ. Press, 1993. MR Zbl
- [Dafermos and Rodnianski 2007] M. Dafermos and I. Rodnianski, “A note on energy currents and decay for the wave equation on a Schwarzschild background”, preprint, 2007. arXiv 0710.0171
- [Dafermos and Rodnianski 2009] M. Dafermos and I. Rodnianski, “The red-shift effect and radiation decay on black hole spacetimes”, *Comm. Pure Appl. Math.* **62**:7 (2009), 859–919. MR Zbl
- [Dafermos and Rodnianski 2013] M. Dafermos and I. Rodnianski, “Lectures on black holes and linear waves”, pp. 97–205 in *Evolution equations* (Zurich, 2008), edited by D. Ellwood et al., Clay Math. Proc. **17**, Amer. Math. Soc., Providence, RI, 2013. MR Zbl
- [Dafermos et al. 2016] M. Dafermos, I. Rodnianski, and Y. Shlapentokh-Rothman, “Decay for solutions of the wave equation on Kerr exterior spacetimes, III: The full subextremal case $|a| < M$ ”, *Ann. of Math. (2)* **183**:3 (2016), 787–913. MR Zbl
- [Dafermos et al. 2021] M. Dafermos, G. Holzegel, I. Rodnianski, and M. Taylor, “The non-linear stability of the Schwarzschild family of black holes”, preprint, 2021. arXiv 2104.08222
- [Deng and Pusateri 2020] Y. Deng and F. Pusateri, “On the global behavior of weak null quasilinear wave equations”, *Comm. Pure Appl. Math.* **73**:5 (2020), 1035–1099. MR Zbl
- [Facci and Metcalfe 2022] M. Facci and J. Metcalfe, “Global existence for quasilinear wave equations satisfying the null condition”, *Houston J. Math.* **48**:3 (2022), 631–653. Zbl
- [Fang 2021] A. J. Fang, “Nonlinear stability of the slowly-rotating Kerr–de Sitter family”, preprint, 2021. arXiv 2112.07183
- [Fang 2022] A. J. Fang, “Linear stability of the slowly-rotating Kerr–de Sitter family”, preprint, 2022. arXiv 2207.07902
- [Giorgi et al. 2022] E. Giorgi, S. Klainerman, and J. Szeftel, “Wave equations estimates and the nonlinear stability of slowly rotating Kerr black holes”, preprint, 2022. arXiv 2205.14808

- [Hawking and Ellis 1973] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time*, Cambridge Monogr. Math. Phys. **1**, Cambridge Univ. Press, 1973. MR Zbl
- [Hidano 2004] K. Hidano, “The global existence theorem for quasi-linear wave equations with multiple speeds”, *Hokkaido Math. J.* **33**:3 (2004), 607–636. MR Zbl
- [Hintz and Vasy 2018] P. Hintz and A. Vasy, “The global non-linear stability of the Kerr–de Sitter family of black holes”, *Acta Math.* **220**:1 (2018), 1–206. MR Zbl
- [Ionescu and Klainerman 2015] A. D. Ionescu and S. Klainerman, “On the global stability of the wave-map equation in Kerr spaces with small angular momentum”, *Ann. PDE* **1**:1 (2015), art. id. 1. MR Zbl
- [John 1979] F. John, “Blow-up of solutions of nonlinear wave equations in three space dimensions”, *Manuscripta Math.* **28**:1-3 (1979), 235–268. MR Zbl
- [Keir 2018] J. Keir, “The weak null condition and global existence using the p -weighted energy method”, preprint, 2018. arXiv 1808.09982
- [Klainerman 1984] S. Klainerman, “Long time behaviour of solutions to nonlinear wave equations”, pp. 1209–1215 in *Proceedings of the International Congress of Mathematicians, II* (Warsaw, 1983), edited by Z. Ciesielski and C. Olech, PWN, Warsaw, 1984. MR Zbl
- [Klainerman 1986] S. Klainerman, “The null condition and global existence to nonlinear wave equations”, pp. 293–326 in *Nonlinear systems of partial differential equations in applied mathematics, I* (Santa Fe, NM, 1984), edited by B. Nicolaenko et al., Lectures in Appl. Math. **23**, Amer. Math. Soc., Providence, RI, 1986. MR Zbl
- [Klainerman and Sideris 1996] S. Klainerman and T. C. Sideris, “On almost global existence for nonrelativistic wave equations in 3D”, *Comm. Pure Appl. Math.* **49**:3 (1996), 307–321. MR Zbl
- [Klainerman and Szeftel 2022a] S. Klainerman and J. Szeftel, “Construction of GCM spheres in perturbations of Kerr”, *Ann. PDE* **8**:2 (2022), art. id. 17. MR Zbl
- [Klainerman and Szeftel 2022b] S. Klainerman and J. Szeftel, “Effective results on uniformization and intrinsic GCM spheres in perturbations of Kerr”, *Ann. PDE* **8**:2 (2022), art. id. 18. MR Zbl
- [Klainerman and Szeftel 2023] S. Klainerman and J. Szeftel, “Kerr stability for small angular momentum”, *Pure Appl. Math. Q.* **19**:3 (2023), 791–1678. MR Zbl
- [Lindblad 1990] H. Lindblad, “On the lifespan of solutions of nonlinear wave equations with small initial data”, *Comm. Pure Appl. Math.* **43**:4 (1990), 445–472. MR Zbl
- [Lindblad 1992] H. Lindblad, “Global solutions of nonlinear wave equations”, *Comm. Pure Appl. Math.* **45**:9 (1992), 1063–1096. MR Zbl
- [Lindblad 2008] H. Lindblad, “Global solutions of quasilinear wave equations”, *Amer. J. Math.* **130**:1 (2008), 115–157. MR Zbl
- [Lindblad 2017] H. Lindblad, “On the asymptotic behavior of solutions to the Einstein vacuum equations in wave coordinates”, *Comm. Math. Phys.* **353**:1 (2017), 135–184. MR Zbl
- [Lindblad and Rodnianski 2003] H. Lindblad and I. Rodnianski, “The weak null condition for Einstein’s equations”, *C. R. Math. Acad. Sci. Paris* **336**:11 (2003), 901–906. MR Zbl
- [Lindblad and Rodnianski 2005] H. Lindblad and I. Rodnianski, “Global existence for the Einstein vacuum equations in wave coordinates”, *Comm. Math. Phys.* **256**:1 (2005), 43–110. MR Zbl
- [Lindblad and Rodnianski 2010] H. Lindblad and I. Rodnianski, “The global stability of Minkowski space-time in harmonic gauge”, *Ann. of Math. (2)* **171**:3 (2010), 1401–1477. MR Zbl
- [Lindblad and Tohaneanu 2018] H. Lindblad and M. Tohaneanu, “Global existence for quasilinear wave equations close to Schwarzschild”, *Comm. Partial Differential Equations* **43**:6 (2018), 893–944. MR Zbl
- [Lindblad and Tohaneanu 2020] H. Lindblad and M. Tohaneanu, “A local energy estimate for wave equations on metrics asymptotically close to Kerr”, *Ann. Henri Poincaré* **21**:11 (2020), 3659–3726. MR Zbl
- [Lindblad et al. 2013] H. Lindblad, M. Nakamura, and C. D. Sogge, “Remarks on global solutions for nonlinear wave equations under the standard null conditions”, *J. Differential Equations* **254**:3 (2013), 1396–1436. MR Zbl
- [Looi 2022] S.-Z. Looi, “Improved decay for quasilinear wave equations close to asymptotically flat spacetimes including black hole spacetimes”, preprint, 2022. arXiv 2208.05439

- [Looi and Tohaneanu 2022] S.-Z. Looi and M. Tohaneanu, “Global existence and pointwise decay for the null condition”, preprint, 2022. arXiv 2204.03626
- [Luk 2013] J. Luk, “The null condition and global existence for nonlinear wave equations on slowly rotating Kerr spacetimes”, *J. Eur. Math. Soc.* **15**:5 (2013), 1629–1700. MR Zbl
- [Marzuola et al. 2010] J. Marzuola, J. Metcalfe, D. Tataru, and M. Tohaneanu, “Strichartz estimates on Schwarzschild black hole backgrounds”, *Comm. Math. Phys.* **293**:1 (2010), 37–83. MR Zbl
- [Mavrogiannis 2022] G. Mavrogiannis, *Decay for quasilinear wave equations on cosmological black hole backgrounds*, Ph.D. thesis, University of Cambridge, 2022.
- [Metcalfe and Sogge 2005] J. Metcalfe and C. D. Sogge, “Hyperbolic trapped rays and global existence of quasilinear wave equations”, *Invent. Math.* **159**:1 (2005), 75–117. MR Zbl
- [Metcalfe and Sogge 2007] J. Metcalfe and C. D. Sogge, “Global existence of null-form wave equations in exterior domains”, *Math. Z.* **256**:3 (2007), 521–549. MR Zbl
- [Metcalfe et al. 2005] J. Metcalfe, M. Nakamura, and C. D. Sogge, “Global existence of solutions to multiple speed systems of quasilinear wave equations in exterior domains”, *Forum Math.* **17**:1 (2005), 133–168. MR Zbl
- [Metcalfe et al. 2012] J. Metcalfe, D. Tataru, and M. Tohaneanu, “Price’s law on nonstationary space-times”, *Adv. Math.* **230**:3 (2012), 995–1028. MR Zbl
- [Morawetz 1968] C. S. Morawetz, “Time decay for the nonlinear Klein–Gordon equations”, *Proc. Roy. Soc. Lond. Ser. A* **306** (1968), 291–296. MR Zbl
- [Pasqualotto 2019] F. Pasqualotto, “Nonlinear stability for the Maxwell–Born–Infeld system on a Schwarzschild background”, *Ann. PDE* **5**:2 (2019), art. id. 19. MR Zbl
- [Peterson and Vasy 2021] O. Peterson and A. Vasy, “Wave equations in the Kerr–de Sitter spacetime: the full extremal range”, 2021. To appear in *J. Eur. Math. Soc.* arXiv 2112.01355
- [Sideris and Tu 2001] T. C. Sideris and S.-Y. Tu, “Global existence for systems of nonlinear wave equations in 3D with multiple speeds”, *SIAM J. Math. Anal.* **33**:2 (2001), 477–488. MR Zbl
- [Sogge and Wang 2010] C. D. Sogge and C. Wang, “Concerning the wave equation on asymptotically Euclidean manifolds”, *J. Anal. Math.* **112** (2010), 1–32. MR Zbl
- [Tataru and Tohaneanu 2011] D. Tataru and M. Tohaneanu, “A local energy estimate on Kerr black hole backgrounds”, *Int. Math. Res. Not.* **2011**:2 (2011), 248–292. MR Zbl
- [Tohaneanu 2022] M. Tohaneanu, “Pointwise decay for semilinear wave equations on Kerr spacetimes”, *Lett. Math. Phys.* **112**:1 (2022), art. id. 6. MR Zbl
- [Wang and Yu 2014] C. Wang and X. Yu, “Global existence of null-form wave equations on small asymptotically Euclidean manifolds”, *J. Funct. Anal.* **266**:9 (2014), 5676–5708. MR Zbl
- [Yang 2013] S. Yang, “Global solutions of nonlinear wave equations in time dependent inhomogeneous media”, *Arch. Ration. Mech. Anal.* **209**:2 (2013), 683–728. MR Zbl
- [Yu 2021a] D. Yu, “Asymptotic completeness for a scalar quasilinear wave equation satisfying the weak null condition”, 2021. To appear in *Mem. Amer. Math. Soc.* arXiv 2105.11573
- [Yu 2021b] D. Yu, “Modified wave operators for a scalar quasilinear wave equation satisfying the weak null condition”, *Comm. Math. Phys.* **382**:3 (2021), 1961–2013. MR Zbl

Received 27 Oct 2022. Revised 25 Apr 2023. Accepted 6 Jun 2023.

HANS LINDBLAD: lindblad@math.jhu.edu

Department of Mathematics, Johns Hopkins University, Baltimore, MD, United States

MIHAI TOHANEANU: mihai.tohaneanu@uky.edu

Department of Mathematics, University of Kentucky, Lexington, KY, United States

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 8 2024

Dynamical torsion for contact Anosov flows	2619
YANN CHAUBET and NGUYEN VIET DANG	
Measure propagation along a \mathcal{C}^0 -vector field and wave controllability on a rough compact manifold	2683
NICOLAS BURQ, BELHASSEN DEHMAN and JÉRÔME LE ROUSSEAU	
The Dirichlet problem for the Lagrangian mean curvature equation	2719
ARUNIMA BHATTACHARYA	
Local lens rigidity for manifolds of Anosov type	2737
MIHAJLO CEKIĆ, COLIN GUILLARMOU and THIBAUT LEFEUVRE	
The rank-one theorem on RCD spaces	2797
GIOACCHINO ANTONELLI, CAMILLO BRENA and ENRICO PASQUALETTO	
A new approach to the Fourier extension problem for the paraboloid	2841
CAMIL MUSCALU and ITAMAR OLIVEIRA	
A Poincaré–Steklov map for the MIT bag model	2923
BADREDDINE BENHELLAL, VINCENT BRUNEAU and MAHDI ZREIK	
The weak null condition on Kerr backgrounds	2971
HANS LINDBLAD and MIHAI TOHANEANU	