

ANALYSIS & PDE

Volume 17

No. 9

2024

IOANN VASILYEV

**A GENERALIZATION OF THE BEURLING–MALLIAVIN
MAJORANT THEOREM**



A GENERALIZATION OF THE BEURLING–MALLIAVIN MAJORANT THEOREM

IOANN VASILYEV

We prove a generalization of the Beurling–Malliavin majorant theorem. In more detail, we establish a new sufficient condition for a function to be a Beurling–Malliavin majorant. Our result is strictly more general than that of the Beurling–Malliavin majorant theorem. We also show that our result is sharp in a number of senses.

1. Introduction

Let $\text{Lip}(\mathbb{R})$ denote the space of Lipschitz functions in \mathbb{R} (i.e., functions f satisfying for all $x, y \in \mathbb{R}$ the inequality $|f(x) - f(y)| \leq C|x - y|$ with $C > 0$ independent of x, y). By $\text{Lip}(\xi, \mathbb{R})$ we shall denote all Lipschitz functions in \mathbb{R} with the Lipschitz constant ξ .

The following theorem was first proved by A. Beurling and P. Malliavin.

Theorem A [Beurling and Malliavin 1962]. *Let $\omega : \mathbb{R} \rightarrow (0, 1]$ be a function such that $\log(1/\omega) \in L^1(\mathbb{R}, dx/(1+x^2))$, and $\log(1/\omega)$ is a Lipschitz function. Then for each $\delta > 0$ there exists a function $f \in L^2(\mathbb{R})$, which is not identically zero and which satisfies $\text{spec}(f) \subset [0, \delta]$ and $|f(x)| \leq \omega(x)$ for all $x \in \mathbb{R}$.*

For a function $f \in L^2(\mathbb{R})$, by $\text{spec}(f)$ we mean the spectrum of f , i.e., the support of its Fourier transformation. Note that the spectrum is defined up to a set of the Lebesgue measure zero. Let us also remark that here the term “not identically zero” means “not zero almost everywhere”. We shall further sometimes write just “nonzero” for brevity.

The Beurling–Malliavin theorems are considered by many experts to be among the most deep and important results of the 20th century harmonic analysis. **Theorem A** above is called the Beurling–Malliavin majorant theorem (or the first Beurling–Malliavin theorem). This result gives conditions for the majorant ω ensuring existence of a nonzero function whose spectrum lies in an arbitrary small interval and whose modulus is majorized by ω . This theorem is a crucial tool in the proof of the second Beurling–Malliavin theorem about the radius of completeness of an exponential system. Moreover, **Theorem A** was recently used by J. Bourgain and S. Dyatlov [2018] in the theory of resonances for hyperbolic surfaces. Deep connections of the first Beurling–Malliavin theorem with nowadays popular gap and type problems are discussed in [Borichev and Sodin 2011; Poltoratski 2012; Makarov and Poltoratski 2010].

Note that **Theorem A** is in a certain sense a contradiction to the following postulate, called the uncertainty principle: “It is impossible for a nonzero function and its Fourier transform to be simultaneously very small, unless the function is zero”. Indeed, **Theorem A** shows that there exist nonzero functions that are “small” and whose Fourier transforms are also “small”. Of course these smallnesses are different from each

MSC2020: primary 42B05, 42B20, 46E35; secondary 26A16.

Keywords: uncertainty principle, Beurling and Malliavin theorems.

other and from the smallness in $L^2(\mathbb{R})$. So in fact [Theorem A](#) does not contradict the most well-known variant of the uncertainty principle, the Heisenberg inequality. For recent violations of the uncertainty principle of a completely different nature, see [\[Kislyakov and Perstneva 2021; Nazarov and Olevskii 2018\]](#).

In addition to the original proof of A. Beurling and P. Malliavin, there are many approaches to the proof of Beurling–Malliavin theorems due to H. Redheffer [\[1977\]](#), L. De Branges [\[1968\]](#), P. Kargaev [\[Koosis\]](#), N. Makarov and A. Poltoratski [\[Makarov and Poltoratski 2010\]](#), to name just some of them. V. Havin, J. Mashregi and F. Nazarov [\[Mashregi et al. 2005\]](#) suggested a new proof of the first Beurling–Malliavin theorem. An essential novelty of their proof was that it was done by (almost) purely real methods and did not use complex analysis except at one place; see [\[Mashregi et al. 2005\]](#) and the remark right after the formulation of [Theorem B](#) below.

Among the goals of the present paper is to give a proof of a new nontrivial generalization of [Theorem A](#). Before stating our main results, we recall some classical definitions and fix some notation.

One of the principle objects of this paper is the class of BM majorants.

Definition 1. Let ω be a bounded nonnegative function on \mathbb{R} . This function is called a Beurling–Malliavin majorant (we shall further write “BM majorant” to save space) if for any $\sigma > 0$ there exists a nonzero function $f \in L^2(\mathbb{R})$ such that

- (a) $|f| \leq \omega$,
- (b) $\text{spec}(f) \subset [0, \sigma]$.

The set of all BM majorants will be further referred to as the BM class. If the conditions (a) and (b) just above are satisfied for a function ω with some fixed $\sigma > 0$, then we call such function ω a σ -admissible majorant. If we replace the condition (a) with a stronger two-sided condition $C\omega \leq |f| \leq \omega$ for some constant $C > 0$, then what we get is the definition of a strictly admissible majorant.

Recall that the *Poisson measure* dP on \mathbb{R} is defined by the formula

$$dP(x) := \frac{dx}{1+x^2}.$$

The corresponding weighted Lebesgue space $L^1(dP)$ is the space of all functions f that satisfy $\int_{\mathbb{R}} |f| dP < \infty$. The expression $\int_{\mathbb{R}} \log(1/\omega) dP$ will be sometimes further referred to as the logarithmic integral of ω .

Note that the condition $\log(1/\omega) \in L^1(dP)$ is necessary for ω to be a BM majorant, but not sufficient; see [\[Mashregi et al. 2005\]](#). What [Theorem A](#) establishes is that some additional regularity suffices for admissibility.

We remind the reader of how one should modify the Cauchy kernel in order to extend the definition of the Hilbert transformation up to the space $L^1(dP)$.

Definition 2. The *Hilbert transformation* of a function $f \in L^1(dP)$ is defined as the principal value integral

$$\mathcal{H}f(x) := \int_{\mathbb{R}} \left(\frac{1}{x-t} + \frac{t}{t^2+1} \right) f(t) dt.$$

It is worth noting that the integral above converges for almost all $x \in \mathbb{R}$.

To avoid ambiguity, we stress that this definition coincides, up to an additive constant, with the classical one for functions in $L^1(\mathbb{R})$.

Let us now introduce function classes that will play an important role in what follows. To this end, we first define an auxiliary system of intervals: $J_0 = [-2, 2)$, and for $j \in \mathbb{N}$,

$$J_j = [2^j, 2^{j+1}), \quad J_{-j} = [-2^{j+1}, -2^j).$$

Definition 3. Let $\beta \in (0, 1]$. If $\beta < 1$, then we shall say that an absolutely continuous function φ belongs to the class V_β if φ is a β -Hölder function on the interval J_j with the constant κ_j and moreover these constants satisfy

$$\left(\sum_{n \in \mathbb{Z}} 2^{-|j|} \kappa_j^{1/(1-\beta)} \right)^{1-\beta} < \infty. \tag{1}$$

In the case when $\beta = 1$, we use the convention $V_\beta = \text{Lip}(\mathbb{R})$.

Note that these classes resemble homogeneous weighted Sobolev spaces. We are going to work with functions that belong to intersections $L^1(dP) \cap V_\beta$. From the functional-analytic point of view, these intersections are Banach spaces with respect to the norms $\| \cdot \|_{L^1(dP)} + \| \cdot \|_{V_\beta}$.

We are now in position to formulate the first main result of this paper to be proved in the next section.

Theorem 1. *Let $\omega : \mathbb{R} \rightarrow (0, 1]$ be a function such that $\log(1/\omega) \in L^1(dP)$, with $\log(1/\omega)$ absolutely continuous and satisfying $\log(1/\omega) \in V_\beta$ for some $\beta \in (0, 1]$. Then for each $\delta > 0$ there exists a function $f \in L^2(\mathbb{R})$, not identically zero, such that $\text{spec}(f) \subset [0, \delta]$ and $|f(x)| \leq \omega(x)$ for all $x \in \mathbb{R}$.*

Remark. We would like to stress that one can replace the intervals J_j in the definition of the spaces V_β with any system of intervals $[\lambda_j, \lambda_{j+1})$, where $\{\lambda_j\}$ is any sequence of reals satisfying $\lambda \leq \lambda_{j+1}/\lambda_j \leq \Lambda$ with $1 < \lambda < \Lambda < \infty$, in a way that the corresponding version of **Theorem 1** holds true.

Remark. Throughout this paper, Ω will mean $\log(1/w)$ for a function $\omega : \mathbb{R} \rightarrow (0, 1]$.

In order to get some intuition of what a “typical” function satisfying $\Omega \in L^1(dP)$ and $\Omega \in V_\beta$ looks like, the reader is welcome to think of a function whose graph consists of an infinite number of “pits” and “hills”; see the pictures of Section 1.5 in [Mashregi et al. 2005] and **Figure 1** below. Of course, the same intuition applies to the functions with Lipschitz logarithm and finite logarithmic integral (i.e., those satisfying the conditions of the first Beurling–Malliavin theorem). However, we shall shortly see that there are drastic differences between these classes of functions.

Indeed, let us compare our sufficient condition of **Theorem 1** with these already known. First, it is obvious that our theorem is a generalization of the first Beurling–Malliavin theorem, since it is a particular case of our result that corresponds to $\beta = 1$.

A “typical” function in classes V_β is visualized at **Figure 1**.

There are many other sufficient “regularity” conditions for the admissibility; see for instance those contained in [Koosis 1988; 1992; Belov and Havin 2015]. However, all these conditions are either imposed on the Hilbert transform of Ω , or they claim that only some regularization or some minorant of ω is admissible. For instance, if the condition $(\log \omega(\cdot))/(1 + (\cdot)^2)^{1/2} \in \dot{W}^{1/2,2}(\mathbb{R})$ is fulfilled for




Figure 1. A “typical” function in V_β .

a function ω that has convergent logarithmic integral, then some regularization of this function is an admissible majorant; see [Beurling and Malliavin 1962].

Note that the following approximation property of the spaces V_β with $\beta \in (0, 1]$ is a direct consequence of our Theorem 1.

Corollary. *By a theorem of A. Baranov and V. Havin [2006, Section 6], we get that, for any $\beta \in (0, 1]$, $\sigma > 0$, and any $\omega \in V_\beta$, the space of all functions in $L^2(\mathbb{R})$ with the spectrum in $\mathbb{R} \setminus [0, \sigma]$ is not dense in the weighted Lebesgue space $L^1(\omega)$.*

We hope that our main results will find other applications in harmonic and complex analysis, in particular for the uncertainty principle and for exponential systems.

The main step of the proof of Theorem 1 is the following lemma.

Lemma 1 (a new variant of the global Nazarov lemma). *Let $0 < \beta \leq 1$. Suppose that $\Omega \in L^1(dP) \cap V_\beta$ is positive. Then, for each $\varepsilon > 0$, there exists a function Ω_1 , satisfying*

- (A) $\Omega(x) \leq \Omega_1(x)$ for all $x \in \mathbb{R}$,
- (B) $\Omega_1 \in L^1(\mathbb{R}, dx/(1+x^2))$,
- (C) $\mathcal{H}\Omega_1 \in \text{Lip}(\varepsilon, \mathbb{R})$, where \mathcal{H} is the Hilbert transform on the real line.

Indeed, Theorem 1 follows from Lemma 1, thanks to the following sufficient condition for a function to be a BM majorant, which is a consequence of a more general result, proved by Mashreghi and Havin.

Theorem B. *If $\omega : \mathbb{R} \rightarrow (0, 1]$, $\log(1/\omega) \in L^1(dP)$ and $\|(\mathcal{H} \log(1/\omega))'\|_\infty < \pi \sigma$, then ω is a σ -admissible majorant.*

Remark. The proof of Theorem B uses a one-dimensional construction coming from the classical (complex) theory of Hardy spaces on the unit circle. Namely, given a nonnegative function on the unit circle with convergent logarithmic integral there exists an analytic function whose modulus coincides with the former function. Such functions are called outer; see [Nikolski 2012] for details.

For necessary conditions for σ -admissible majorants, see [Belov 2007; 2008b; Baranov and Khavin 2006].

We briefly discuss main ideas lying behind our proof of Lemma 1. Our proof is inspired by that of the Nazarov lemma from [Mashregi et al. 2005]. Indeed, we use the beautiful idea of a so-called regularized system of intervals, which was first introduced by F. Nazarov and fruitfully used in [Mashregi et al. 2005]. Another important feature of the proof of the Nazarov lemma in [Mashregi et al. 2005] is a version of the Hadamard–Landau inequality. We have had to modify this result drastically in order for it to fit the conditions of our Lemma 1. This culminated in Lemma 4 of the present paper. On top of that, most estimates from the proof in [Mashregi et al. 2005] become considerably harder under our assumptions, in comparison to the Lipschitz condition of that work.

Note that Nazarov’s lemma is by itself a highly nontrivial and very interesting result in harmonic analysis. To illustrate this, we mention [Stolyarov and Zatitskiy 2021], where the authors have utilized the main object of the Nazarov lemma, the regularized system of intervals, in some special form. For a multidimensional version of the classical Nazarov lemma, see our paper [Vasilyev 2022].

Let us now discuss the second main result of this article. Our Theorem 2, gives an answer to the following question: “How sharp is the result of Theorem 1 ?” The answer to this question is given in the following result.

Theorem 2. *For any $\beta \in (0, 1)$, there are functions $\omega : \mathbb{R} \rightarrow (0, 1]$ satisfying $\log(1/\omega) \in L^1(dP)$ and $2^{-|j|} \kappa_j^{1/(1-\beta)} \asymp 1$ in the notation of Theorem 1, that are not BM majorants.*

We remark that our Theorem 2 shows that the condition $\log(1/\omega) \in V_\beta$ in our Theorem 1 is sharp in a number of senses.

The proof of Theorem 2 builds upon one construction from [Belov and Havin 2015]. This construction says that smallness of a bandlimited function is “contagious”: if such a function is small on an interval, it is also small on a much larger concentric interval. This construction is due to A. Borichev and it works only for majorants that have a growth strictly greater than linear at a sequence tending to infinity. Majorants that appear in the formulation of Theorem 2 have at most linear growth at infinity. Nevertheless, for some of these majorants, we were able to use a combination of Borichev’s construction with an iteration method to prove Theorem 2.

The paper is organized as follows. Theorem 1 is proved in Sections 2 and 3. Section 4 is devoted to the proof of Theorem 2.

We finally mention some open questions concerning Theorems 1 and 2. The first question consists of determining whether the condition $\log(1/\omega) \in V_\beta$ in Theorem 1 can be weakened down to, roughly speaking, a condition of the kind “ ω belongs to some Orlicz-type class, defined in the spirit of V_β classes”. The second question concerns the system of intervals that are used in the definition of the spaces V_β . Namely, we would like to find a necessary and sufficient condition on the system of intervals instead of the dyadic system in Definition 3, for which the first theorem still holds. Yet another question is to find a multidimensional version of Theorem 1 which seems unavailable at the present time, according to [Han and Schlag 2020]. The fourth and the final question reads as follows. It would be also interesting to find

counterparts of the main results of this paper in the context of the so-called model spaces, in spirit of Yu. S. Belov's early papers. The author plans to attack the aforementioned questions in the nearest future.

2. A new local Nazarov lemma

We accumulate here the list of the frequently used technical abbreviations and notation. For an interval $a \subset \mathbb{R}$ its length is denoted by $l(a)$, c_a will stand for the center of a and λa with λ positive will be the interval centered at c_a and whose edge length equals $\lambda l(a)$. Let I be an interval on the real line. We will denote by $T_I(x)$ the distance from $x \in \mathbb{R}$ to $\mathbb{R} \setminus I$. For a dyadic interval b , we will denote by b^\sharp the dyadic parent of b . Throughout this paper, I^* will denote the unit interval $[-\frac{1}{2}, \frac{1}{2}]$. For $\beta \in (0, 1)$, we denote $\text{Hol}_\beta(\kappa, I)$ the class of β -Hölder functions on the interval I , with the constant κ , i.e., all f defined on I such that for all $x \in I$ and $y \in I$ holds $|f(x) - f(y)| \leq \kappa|x - y|^\beta$.

The main step of the proof of our new global Nazarov lemma is its following local variant.

Lemma 2 (a new local Nazarov lemma). *Let $I \subset \mathbb{R}$ be an interval and let $\beta \in (0, 1]$. Suppose that f is a nonnegative absolutely continuous function such that holds $f \in \text{Hol}_\beta(\kappa, I)$ and $\|f\|_{L^\infty(I)} \leq \delta l(I)$ for some $0 < \delta \leq 1$ and $1 \leq \kappa$. Then there exists a nonnegative function $F \in C^\infty(\mathbb{R})$ such that*

- (i) $F = 0$ outside $1.5I$,
- (ii) $f(x) \leq F(x)$ for all $x \in I$,
- (iii) $\|(\mathcal{H}F)'\|_{L^\infty(\mathbb{R})} \lesssim \delta$,
- (iv) $\int_{\mathbb{R}} F(x) dx \lesssim \int_I f + \kappa \delta^{-\beta} l(I)^{1-\beta} (\int_I f)^\beta$.

In the case when $\beta = 1$ in [Lemma 2](#), the corresponding result coincides with [Lemma 2.6](#) from [\[Mashregi et al. 2005\]](#).

In the formulation of [Lemma 2](#) and until the end of the third section, the signs \lesssim and \gtrsim indicate that the left-hand (right-hand) part of an inequality is less than the right-hand (left-hand) part multiplied by a constant independent of δ , f , κ and I .

The rest of this section is entirely devoted to the proof of [Lemma 2](#).

Proof of the new local Nazarov lemma. The following definition is very important.

Definition 4. We say that a dyadic interval $a \subset I$ is essential if $\|f\|_{L^\infty(a)} \geq \delta l(a)/2$. Denote by A the set of essential intervals.

It is straightforward to see that we have

$$\{x \in I : f(x) > 0\} \subseteq \bigcup_{a \in A} a.$$

However, we will not use this fact later on in our estimates.

Consider A^M , the set of maximal by inclusion elements of A . To each interval $a \in A^M$ we associate its tail $t(a)$. Informally, the tail $t(a)$ is a family of dyadic intervals that is composed of a countable number of finite series $t_p(a)$, $p = 0, 1, 2, \dots$, of dyadic intervals. For $p = 0$ we define $t_0(a) := a$ and for a

fixed $p \geq 1$, the intervals of the family $t_p(a)$ all have length equal to $l(a)/2^p$ and their unions form the sets

$$a \cup \bigcup_{1 \leq q \leq p} t_q(a) = \left\{ x \in \mathbb{R} : \frac{l(a)}{2} + l(a) \sum_{q=1}^{p-1} \frac{3^q}{2^q} \leq |x - c_a| < \frac{l(a)}{2} + l(a) \sum_{q=1}^p \frac{3^q}{2^q} \right\}.$$

For a detailed discussion of tails, see [Mashregi et al. 2005, Section 2.6.5]. In fact, after we have added these tails, we will get a regularized system of intervals; see [Mashregi et al. 2005, Sections 2.6 and 2.7]. Next, we define $B := \bigcup_{a \in A^M} t(a)$, and then pose $\tau := \{c \in B^M : c \subseteq I\}$. Here, B^M stands for the set of maximal by inclusion elements of B . Note that the system τ covers I , consists of dyadic intervals and any $c \in \tau$ satisfies $\delta l(c) \geq \|f\|_{L^\infty(c)}$; see [Mashregi et al. 2005].

Define for an interval $a \in \tau$ its neighborhood $N(a)$ by

$$N(a) := \left\{ b \in \tau : d(a, b) \leq 2l(a), \frac{1}{2} \leq \frac{l(a)}{l(b)} \leq 2 \right\}.$$

Note that $\#N(a) \leq 9$. We shall need the following property of the system τ .

Lemma 3. *Suppose that $a \in \tau$ and $b \in \tau \setminus N(a)$. If $l(b) \leq 2l(a)$ then $d(2a, 2b) \geq l(a)/2$, and if $l(b) = 2^k l(a)$ for some natural $k \geq 2$, then $d(2a, 2b) \geq 2 \cdot 3^{k-2} l(a)$.*

Proof. The proof of this lemma is not detailed here, since it can be found in [Mashregi et al. 2005, Section 2.6.6]. □

Define $2\tau := \{2c : c \in \tau\}$. As a direct consequence of the lemma, we deduce that the multiplicity $\#\{b \in 2\tau : x \in b\}$ is uniformly bounded in $x \in \mathbb{R}$. Indeed, if $b \in \tau \setminus N(a)$, then $d(2a, 2b) > 0$ and

$$\sup_{x \in \mathbb{R}} \#\{b \in 2\tau : x \in b\} \leq \sup_{a \in \tau} \#N(a) \lesssim 1.$$

Fix a bump function ϕ , i.e., $\phi \in C^\infty(\mathbb{R})$ satisfying $0 \leq \phi(x) \leq 1$ for all $x \in \mathbb{R}$, $\phi \equiv 0$ outside $1.5I^*$ and $\phi \equiv 1$ on I^* . Second, for an interval $a \in \tau$ define

$$\phi_a(\cdot) := \delta l(a) \phi\left(\frac{(\cdot) - c_a}{l(a)}\right).$$

Simple calculation shows that

$$\mathcal{H}\phi_b(\cdot) = \delta l(b) \mathcal{H}\phi\left(\frac{(\cdot) - c_b}{l(b)}\right).$$

Hence we infer the inequality $\|(\mathcal{H}\phi_b)'\|_{L^\infty(\mathbb{R})} \lesssim \delta$. We finally define F by

$$F := \sum_{a \in \tau} \phi_a.$$

Now we have to check the required properties of the majorant F . The first one follows readily from the definition of F . To prove the second one, note that for all $a \in \tau$ we have $\|f\|_{L^\infty(a)} \leq \delta l(a)$. Indeed, suppose the contrary, i.e., that $\|f\|_{L^\infty(a_0)} > \delta l(a_0)$ for some $a_0 \in \tau$. This means that

$$\|f\|_{L^\infty(a_0^\sharp)} \geq \|f\|_{L^\infty(a_0)} > \delta l(a_0) = \delta \left(\frac{l(a_0^\sharp)}{2}\right),$$

which in turn signifies that a_0^\sharp is an essential interval and hence $a_0^\sharp \in \tau$. This contradicts the definition of τ . From here we deduce that if $x \in a \in \tau$, then

$$F(x) \geq \delta l(a) \geq \|f\|_{L^\infty(a)} \geq f(x).$$

Next we estimate the integral of the function F . To this end, we prove a variant of the Hadamard–Landau inequality which is appropriate for our goals.

Lemma 4. *Let a be an interval such that $a \in A^M$. Then we have*

$$\|f\|_{L^\infty(a)}^2 \lesssim \left(\int_a f\right) \delta + \kappa \left(\int_a f\right)^\beta (\delta l(a))^{1-\beta},$$

where $C(r)$ is a positive constant, depending on r only.

Proof. Let $x_0 \in a$ be a point such that $\|f\|_{L^\infty(a)} = f(x_0)$. Suppose with no loss of generality that $a_+ - x_0 \geq l(a)/2$, where a_+ is the right end of the interval a . Consider a point $x \in (x_0, a_+)$. Since f is Hölder continuous, we hence infer the estimate

$$f(x) \geq f(x_0) - \kappa(x - x_0)^\beta.$$

Let $\nu := (f(x_0)/\kappa)^{1/\beta}$. We shall treat two cases separately, according to the value of ν . First, we suppose that $\nu < l(a)/2$. Observe that in this case the point $x_0 + \nu$ belongs to the interval a . We integrate the estimate just above using this observation and deduce that

$$\begin{aligned} \int_a f &\geq \int_{x_0}^{x_0+\nu/2} f(x) dx \geq \int_{x_0}^{x_0+\nu/2} f(x_0) - \kappa(x - x_0)^\beta dx \\ &= \frac{f(x_0)}{2} \left(\frac{f(x_0)}{\kappa}\right)^{1/\beta} - \frac{\kappa}{2^{1+\beta}(\beta+1)} \left(\frac{f(x_0)}{\kappa}\right)^{(\beta+1)/\beta} \\ &\gtrsim \frac{\|f\|_{L^\infty(a)}^{(\beta+1)/\beta}}{\kappa^{1/\beta}} = \frac{\|f\|_{L^\infty(a)}^{2/\beta} \|f\|_{L^\infty(a)}^{(\beta-1)/\beta}}{\kappa^{1/\beta}} \gtrsim \frac{\|f\|_{L^\infty(a)}^{2/\beta} (\delta l(a))^{(\beta-1)/\beta}}{\kappa^{1/\beta}}, \end{aligned} \tag{2}$$

where the last bound above follows from the fact that $a \in A^M$. Hence we have that

$$\|f\|_{L^\infty(a)}^2 \lesssim \kappa \left(\int_a f\right)^\beta (\delta l(a))^{1-\beta}.$$

Consider now the second case, where $\nu \geq l(a)/2$. In this case we shall use the fact that the point $x_0 + l(a)/2$ belongs to the interval a . Integrating the same inequality as in the first case yields

$$\int_a f \geq \int_{x_0}^{x_0+l(a)/2} f(x) dx \geq \frac{l(a)f(x_0)}{2} - \frac{\kappa l(a)^{\beta+1}}{2^{\beta+1}(\beta+1)} = \frac{l(a)}{2} \left(f(x_0) - \frac{\kappa}{\beta+1} \cdot \left(\frac{l(a)}{2}\right)^\beta\right). \tag{3}$$

Note that since $\nu \geq l(a)/2$, we have also that $f(x_0)/\kappa a \geq (l(a)/2)^\beta$. Let us use this in the following way:

$$\int_a f \geq \frac{l(a)}{2} \left(f(x_0) - \frac{f(x_0)}{\beta+1}\right) \gtrsim \delta^{-1} \|f\|_{L^\infty(a)}^2,$$

thanks to the fact that $a \in A^M$. Hence Lemma 4 is proved. □

So, let us start the estimates of the integral of the function F :

$$\begin{aligned} \int_{\mathbb{R}} F &\leq \sum_{b \in A^M} \int_{\mathbb{R}} \phi_b + \sum_{c \in A^M} \sum_{b \in I(c) \setminus c} \int_{\mathbb{R}} \phi_b \leq \delta \sum_{b \in A^M} l(b)^2 + \delta \sum_{c \in A^M} \sum_{b \in I(c) \setminus c} l(b)^2 \\ &\lesssim \delta \sum_{b \in A^M} l(b)^2 + \delta \sum_{c \in A^M} \sum_{p=1}^{\infty} \sum_{b \in I_p(c)} l(b)^2 \lesssim \delta \sum_{b \in A^M} l(b)^2 + \delta \sum_{c \in A^M} \sum_{p=1}^{\infty} 3^p \left(\frac{l(c)}{2^p}\right)^2 \\ &\lesssim \delta \sum_{c \in A^M} l(c)^2 \lesssim \delta^{-1} \sum_{c \in A^M} \|f\|_{L^\infty(c)}^2. \end{aligned} \tag{4}$$

We further use the result of [Lemma 4](#) to continue the estimates of the integral of the function F :

$$\begin{aligned} \int_{\mathbb{R}} F &\lesssim \sum_{c \in A^M} \int_c f + \delta^{-1} \sum_{c \in A^M} \kappa \left(\int_c f \right)^\beta (\delta l(c))^{1-\beta} \\ &\leq \int_I f + \delta^{-\beta} \kappa \left(\sum_{c \in A^M} \int_c f \right)^\beta \cdot \left(\sum_{c \in A^M} l(c) \right)^{1-\beta} \\ &\lesssim \int_I f + \delta^{-\beta} \kappa l(I)^{1-\beta} \left(\int_I f \right)^\beta. \end{aligned} \tag{5}$$

The last and second-to-last inequalities just above are in need of explanation. The last estimate uses the fact that intervals of A^M are nonoverlapping, whereas the penultimate bound follows from the Hölder inequality.

It remains to derive the inequality on the derivative of the Hilbert transformation of the function F . First, we shall obtain this estimate for $x \in \bigcup_{b \in \tau} 2b$. Let $a (= a(x))$ denote the interval from τ such that $x \in 2a$. We isolate the neighborhood $N(a)$ from its complement in τ and infer the inequality

$$|(\mathcal{H}F)'(x)| \leq \sum_{b \in N(a)} |(\mathcal{H}\phi_b)'(x)| + \sum_{\substack{b \in \tau \setminus N(a) \\ l(b) \leq 2l(a)}} |(\mathcal{H}\phi_b)'(x)| + \sum_{k=2}^{\infty} \sum_{\substack{b \in \tau \setminus N(a) \\ l(b) = 2^k l(a)}} |(\mathcal{H}\phi_b)'(x)| =: S_1 + S_2 + S_3.$$

We shall estimate the terms S_1 , S_2 and S_3 separately. We start with the sum S_1 , whose estimate turns out to be easy:

$$S_1 \leq \#N(a) \sup_{b \in \tau} \|(\mathcal{H}\phi_b)'\|_{L^\infty(\mathbb{R})} \lesssim \delta.$$

We further proceed to the second term. We use a simple estimate on the kernel of the Hilbert transformation, the fact that the system of intervals $\{2b\}_{b \in \tau}$ (by [Lemma 3](#)) has finite multiplicity and [Lemma 3](#) to get

$$\begin{aligned} S_2 &\lesssim \sum_{\substack{b \in \tau \setminus N(a) \\ l(b) \leq 2l(a)}} \int_{\mathbb{R}} \phi_b(t) \frac{\partial}{\partial x} \left(\frac{1}{t-x} \right) dt \lesssim \sum_{\substack{b \in \tau \setminus N(a) \\ l(b) \leq 2l(a)}} \int_{\mathbb{R}} \frac{\phi_b(t)}{(t-x)^2} dt \\ &\lesssim \sum_{\substack{b \in \tau \setminus N(a) \\ l(b) \leq 2l(a)}} \int_{1.5b} \frac{\delta l(a) dt}{(t-x)^2} \lesssim \delta l(a) \int_{\{|u| \geq l(a)/2\}} \frac{du}{|u|^2} \lesssim \delta. \end{aligned}$$

The third term can be estimated as well using [Lemma 3](#):

$$S_3 \lesssim \sum_{k=2}^{\infty} \sum_{\substack{b \in \tau \setminus N(a) \\ l(b)=2^k l(a)}} \int_{2b} \frac{\phi_b(t) dt}{(t-x)^2} \leq \sum_{k=2}^{\infty} 2^k \delta l(a) \sum_{\substack{b \in \tau \setminus N(a) \\ l(b)=2^k l(a)}} \int_{2b} \frac{dt}{(t-x)^2} \lesssim \sum_{k=2}^{\infty} 2^k \delta l(a) \int_{\{|u| \geq 2 \cdot 3^{k-2} l(a)\}} \frac{du}{u^2} \lesssim \delta,$$

and the lemma for $x \in \bigcup_{b \in \tau} 2b$ follows.

Next, if a point $z \in \mathbb{R}$ is situated at a positive distance from the set $\bigcup_{b \in \tau} 2b$, then denote by x the point of this set closest to z , and let $a(= a(x))$ be an interval as above. We infer the estimates

$$|(\mathcal{H}F)'(z)| \leq \sum_{b \in \tau \setminus N(a)} |(\mathcal{H}\phi_b)'(z)| + \sum_{b \in N(a)} |(\mathcal{H}\phi_b)'(z)| \lesssim \sum_{b \in \tau \setminus N(a)} \int_{\mathbb{R}} \frac{\phi_b(t) dt}{(t-x)^2} + \#N(a) \sup_{b \in \tau} \|(\mathcal{H}\phi_b)'\|_{L^\infty(\mathbb{R})}.$$

Thanks to the estimates of the terms S_1, S_2 and S_3 , we conclude that the needed variant of the local Nazarov lemma is proved. □

3. Proof of a new global Nazarov lemma

In this section, we shall derive the global Nazarov lemma from the local one.

Proof. Until the end of the third section, the signs \lesssim and \gtrsim indicate that the left-hand (right-hand) part of an inequality is less than the right-hand (left-hand) part multiplied by a “harmless” positive constant.

Note that we may assume in the global Nazarov lemma that $\Omega(x) = 0$ for $|x| \leq R$, with R being an arbitrary large positive number. Indeed, if it is not the case, then consider the function $\Omega(\cdot) = \max(0, \Omega - \mathcal{M})(\cdot)$, where $\mathcal{M} := \max_{x \in B(0,R)} \Omega(x)$. If Ω_I is a majorant of the function Ω satisfying properties [\(B\)](#) and [\(C\)](#) then the function $\Omega_I + \mathcal{M}$ will be the desired majorant of the function Ω .

Fix $0 < \varepsilon \leq 1$ and choose $1 < R_1$ so big that

$$\int_{\mathbb{R} \setminus (-R_1, R_1)} \Omega dP \leq \varepsilon.$$

Since the series [\(1\)](#) converges, there exists a natural N_1 so big that for all $j > N_1$ it holds that $\kappa_j^{1/(1-\beta)} 2^{-j} \leq \varepsilon^{1/(1-\beta)}$. As a consequence, we infer for all such j the bound

$$\kappa_j 2^{j(\beta+1)} \leq \varepsilon 2^{2j}. \tag{6}$$

By the previous paragraph, we may assume Ω is equal to zero on the interval $(-\max(R_1, 2^{N_1}), \max(R_1, 2^{N_1}))$.

Recall the above-defined system of intervals $J_0 = [-2, 2)$, and, for $j \in \mathbb{N}$,

$$J_j = [2^j, 2^{j+1}), \quad J_{-j} = [-2^{j+1}, -2^j).$$

Next, we shall prove for $x \in \mathbb{R}$ the inequality

$$\Omega(x) \lesssim \varepsilon |x|. \tag{7}$$

With no loss of generality, we suppose that $x > 0$ and we let $n \in \mathbb{N}$ be such that $2^n \leq |x| < 2^{n+1}$. First, note that according to the previous paragraph, the bound [\(7\)](#) is obvious once $|x| \lesssim 1$. Second, for $1 \lesssim |x|$

we will argue as in [Lemma 4](#). We thus find a point $x_0 \in [2^n, 2^{n+1})$ such that $\Omega(x_0) = \|\Omega\|_{L^\infty(J_n)}$. Then, for any $y \in J_n$ we have

$$\Omega(y) \geq \Omega(x_0) - \kappa_n |y - x_0|^\beta.$$

Once again, without loss of generality we suppose that the point $x_0 + 2^n/2$ belongs to the interval J_n . We finally infer the chain of inequalities

$$\begin{aligned} \varepsilon &\geq \int \Omega dP \geq |x|^{-2} \int_{J_n} \Omega(y) dy \geq |x|^{-2} \int_{x_0}^{x_0+2^n/2} (\Omega(x_0) - \kappa_n(y - x_0)^\beta) dy \\ &\geq |x|^{-2} (2^{n-1} \Omega(x_0) - C(\beta) \kappa_n 2^{n(\beta+1)}). \end{aligned} \tag{8}$$

Hence, the bound [\(7\)](#) is proved, by virtue of [\(6\)](#).

Apply the local lemma to each interval J_j and the corresponding restriction $f_j = \Omega \lfloor J_j$. Indeed, [Lemma 2](#) can be applied since these functions satisfy

$$\|f_j\|_\infty \leq \varepsilon 2^j \leq \varepsilon l(J_j)$$

by [\(7\)](#). Thus we obtain functions F_j for $j \in \mathbb{Z}$. The needed majorant Ω_1 is defined by

$$\Omega_1 = \sum_{j \in \mathbb{Z}} F_j.$$

Now, we shall check the required properties of Ω_1 . The first property follows obviously from the local lemma. We proceed to the second one:

$$\begin{aligned} \int_{\mathbb{R}} \Omega_1(t) dP(t) &= \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}} F_j(t) dP(t) \lesssim \sum_{j \in \mathbb{Z}} \int_{1.5J_j} F_j(t) \frac{dt}{2^{2|j|}} \\ &\lesssim \varepsilon^{-\beta} \sum_{j \in \mathbb{Z}} 2^{|j|(\beta-1)} \kappa_j \left(\int_{1.5J_j} \Omega(t) \frac{dt}{2^{2|j|}} \right)^\beta + \sum_{j \in \mathbb{Z}} \int_{1.5J_j} \Omega(t) \frac{dt}{2^{2|j|}} \\ &\lesssim \varepsilon^{-\beta} \left(\sum_{j \in \mathbb{Z}} 2^{-|j|} \kappa_j^{1/(1-\beta)} \right)^{1-\beta} \cdot \left(\sum_{j \in \mathbb{Z}} \int_{1.5J_j} \Omega(t) \frac{dt}{2^{2|j|}} \right)^\beta + \varepsilon \lesssim \varepsilon, \end{aligned} \tag{9}$$

where in the third inequality above we have used the local lemma and in the penultimate bound we have used the Hölder inequality.

So, it remains to check that the third conclusion holds. First, fix a point $x \in \mathbb{R}$. Second, denote by $S(x)$ the interval from the system $\mathcal{F} = \{J_j\}_{j \in \mathbb{Z}}$ such that $x \in S(x)$. Next, denote by $U(x)$ the subset of \mathcal{F} consisting of $S(x)$ and its two neighbor intervals and by $W(x)$ its complement: $W(x) = \mathcal{F} \setminus U(x)$. Finally, write the function Ω_1 as a sum of two functions as follows:

$$\Omega_1 = \sum_{j \in W(x)} F_j + \sum_{j \in U(x)} F_j =: \omega_1 + \omega_2.$$

Since there is only a finite number of intervals in the family $U(x)$, we see that

$$|(\mathcal{H}\omega_2)'(x)| \lesssim \#U(x) \sup_{j \in U(x)} \|(\mathcal{H}F_j)'\|_\infty \lesssim \varepsilon,$$

where we have just used condition (iii) of Lemma 2 in the last estimate. On the other hand, since $\text{supp}(\omega_1) \subseteq \bigcup_{j \in W(x)} 1.5J_j$ we deduce that

$$\text{supp}(\omega_1) \subseteq \left\{ t \in \mathbb{R} : |t - x| \geq \frac{l(S(x))}{4} \right\} \subseteq \left\{ t \in \mathbb{R} : |t - x| \geq \frac{|x|}{16} \right\}.$$

Therefore, we arrive at the chain of inequalities

$$\begin{aligned} |(\mathcal{H}\omega_1)'(x)| &= \left| \left(\int_{\mathbb{R}} \omega_1(t) \frac{1}{t-x} dt \right)' \right| = \left| \int_{\mathbb{R}} \omega_1(t) \frac{\partial}{\partial x} \left(\frac{1}{t-x} \right) dt \right| \\ &= \int_{\mathbb{R}} \omega_1(t) \frac{1}{(t-x)^2} dt \lesssim \int_{\mathbb{R}} \Omega_1(t) dP(t) \lesssim \varepsilon, \end{aligned}$$

thanks to the bound (9).

Hence, the needed variant of the Nazarov lemma is proved. □

Thus, Theorem 1 is also proved, via Theorem B.

4. Sharpness of Lemma 1

Note that the proof of Theorem 2 is a direct consequence of the following proposition.

Proposition 1. *Let $\gamma > \frac{1}{2}$ and define $I_n := [2^n - 2^n/n^\gamma, 2^n + 2^n/n^\gamma]$ for $n \geq 3$. Consider for $x \in \mathbb{R}$ the function*

$$\omega(x) := \begin{cases} \exp(-n^{\gamma-1/2} T_{I_n}(x)) & \text{if } x \in I_n \text{ with } n \geq 3, \\ 1 & \text{otherwise.} \end{cases} \tag{10}$$

We claim that $\log(1/\omega) \in L^1(dP)$ and that $\log(1/\omega)$ satisfies the regularity assumption of Theorem 2, though ω is not a BM majorant.

The graph of the function $\Omega = \log(1/\omega)$ and the main idea of the proof below (i.e., the iteration) is illustrated at Figure 2.

Proof. The first two claims are easy to verify, so we omit their proofs.

Let σ be a positive constant and consider the Bernstein space $\mathcal{E}_{\sigma,1}$, i.e., the space of all entire functions f such that

$$|f(z)| \leq e^{\sigma|z|} \text{ for any } z \in \mathbb{C} \quad \text{and} \quad |f| \leq 1 \text{ on } \mathbb{R}.$$

Recall Lemma 1 from [Belov and Havin 2015].

Lemma A. *For any $\sigma > 0$ there exist a (small) $\alpha(\sigma) \in (0, \frac{1}{2})$ and a (big) $h(\sigma) > 2$ such that for any $h \geq h(\sigma)$, any $f \in \mathcal{E}_{\sigma,1}$ and any compact interval $I \subset \mathbb{R}$*

$$|f| \leq e^{-hT_I} \text{ on } \mathbb{R} \quad \implies \quad |f| \leq e^{-Ch|I|} \text{ on } \tilde{I},$$

where $C > 0$ is an absolute constant and \tilde{I} is the interval centered at $c(I)$ with $|\tilde{I}| = h^{\alpha(\sigma)}|I|$.




Figure 2. The main idea of the proof of [Theorem 2](#).

Let us prove that ω is not in the BM class. Suppose the contrary. Hence, for a fixed $\sigma > 0$ there exists a function, not identically zero, satisfying $f \in L^2(\mathbb{R})$, $\text{spec}(f) \subset [0, \sigma]$ and $|f(x)| \leq \omega(x)$ for all real x . We shall now use [Lemma A](#). Notice that the function f satisfies the conditions of this lemma with $h := n^\vartheta$, where $\vartheta := \gamma - \frac{1}{2} > 0$ and $I := I_n$ for $n \geq n(\gamma)$. We deduce from this lemma that there exists a universal constant C and a power $\alpha \in (0, \frac{1}{2})$, depending only on σ such that

$$|f(x)| \leq \exp(-C2^n)$$

on the interval $I_{n,1} := (n^{\vartheta\alpha}/2)I_n$.

Remark. From now until the end of the present article, the sign $X \asymp Y$ means that $C_1Y \leq X \leq C_2Y$ for some constants C_1 and C_2 depending only on γ, α, σ and C . In this case, we shall say that X is of order Y .

Note that the length of this interval satisfies the bound $|I_{n,1}| \asymp n^{\vartheta(\alpha-1)}2^n$. As a consequence, we infer that the inequality

$$|f(x)| \leq \exp(-Cn^{\vartheta(\alpha-1)}T_{I_{n,1}}(x))$$

is valid for $x \in I_{n,1}$. This means that we can apply [Lemma A](#) once again, now for $h := Cn^{\vartheta(1-\alpha)}$ and $I := I_{n,1}$. This yields the bound

$$|f(x)| \leq e^{-Ch|I_{n,1}|} = e^{-C^22^n},$$

which is true for $x \in I_{n,2} := ((Cn^{\vartheta(1-\alpha)})^\alpha/2)I_{n,1}$. It is not difficult to see that the corresponding interval $I_{n,2}$ has length of order

$$C^\alpha n^{\vartheta(1-\alpha)\alpha + \vartheta(\alpha-1)}2^n = C^\alpha n^{-\vartheta(1-\alpha)^2}2^n.$$

Acting inductively, after m steps, we arrive at the estimate $|f(x)| \leq \exp(-C^m2^n)$, verified by f for $x \in I_{n,m}$ with

$$|I_{n,m}| \asymp n^{-\vartheta(1-\alpha)^m}2^n.$$

Maybe, it is worth noting that $I_{k,m} \cap I_{n,m} = \emptyset$ for any natural m , once $n \neq k$. This results from the fact that we assume, as we can, that $C < 1$.

We are now in position to prove that $f = 0$ identically, which will lead to a contradiction. To this end, we estimate the logarithmic integral of f . For each natural number m it holds that

$$\int_{\mathbb{R}} \log |f(x)| dP(x) \leq - \sum_{n \geq 3} \int_{I_{n,m}} 2^{-2n} C^m 2^n dx \asymp - \sum_{n \geq 3} n^{-\vartheta(1-\alpha)^m}.$$

Choosing m sufficiently large and recalling that $(1 - \alpha) \in (0, 1)$, we arrive at the formula

$$\int_{\mathbb{R}} \log |f(x)| dP(x) = -\infty.$$

Since $f \in L^2(\mathbb{R})$ has the spectrum in the interval $[0, \sigma]$, it hence belongs to the Hardy class $H^2(\mathbb{R})$. From the Jensen inequality, see [Havin and Jörnicke 1994], we deduce that $f = 0$ identically, which contradicts our assumption. Hence, the second theorem is proved. \square

Remark. Alas, our proof above does not work if one replaces in (10) and in the definition of intervals I_n the powers n^γ by θ^n with $\theta \in (1, 2)$.

Remark. It can be seen exactly as above that the function ω_* is not a strictly admissible majorant; recall Definition 1. For a detailed discussion of strictly admissible majorants, see [Belov 2008a].

Acknowledgments

The author is deeply grateful to Anton D. Baranov, Alexander A. Borichev, Evgueni S. Doubtsov, Konstantin M. Dyakonov, Sergei V. Kislyakov, and Yiyu Tang for a number of helpful discussions. The author would also like to thank the anonymous referees for the careful reading of the paper and for a number of helpful suggestions. This work was done under support of the grant no. 075-15-2022-289 for creation and development of Euler International Mathematical Institute.

References

- [Baranov and Khavin 2006] A. D. Baranov and V. P. Khavin, “Admissible majorants for model subspaces and arguments of inner functions”, *Funktsional. Anal. i Prilozhen.* **40**:4 (2006), 3–21. In Russian; translated in *Funct. Anal. Appl.* **40**:4 (2006), 249–263. [MR](#) [Zbl](#)
- [Belov 2007] Y. S. Belov, “Criteria for the admissibility of majorants for model subspaces with a rapidly increasing argument of the generating inner function”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **345** (2007), 55–84. In Russian; translated in *J. Math. Sci. (N.Y.)* **148**:6 (2008), 813–829. [MR](#) [Zbl](#)
- [Belov 2008a] Y. S. Belov, “Model functions with an almost prescribed modulus”, *Algebra i Analiz* **20**:2 (2008), 3–18. In Russian; translated in *St. Petersburg Math. J.* **20**:2 (2009), 163–174. [MR](#) [Zbl](#)
- [Belov 2008b] Y. S. Belov, “Necessary conditions for the admissibility of majorants for some model subspace”, *Algebra i Analiz* **20**:4 (2008), 1–26. In Russian; translated in *St. Petersburg Math. J.* **20**:4 (2009), 507–525. [MR](#) [Zbl](#)
- [Belov and Havin 2015] Y. Belov and V. Havin, “The Beurling–Malliavin multiplier theorem and its analogs for the de Branges spaces”, pp. 581–609 in *Operator theory, I*, edited by D. Alpay, Springer, 2015. [Zbl](#)
- [Beurling and Malliavin 1962] A. Beurling and P. Malliavin, “On Fourier transforms of measures with compact support”, *Acta Math.* **107** (1962), 291–309. [MR](#) [Zbl](#)
- [Borichev and Sodin 2011] A. Borichev and M. Sodin, “Weighted exponential approximation and non-classical orthogonal spectral measures”, *Adv. Math.* **226**:3 (2011), 2503–2545. [MR](#) [Zbl](#)

- [Bourgain and Dyatlov 2018] J. Bourgain and S. Dyatlov, “Spectral gaps without the pressure condition”, *Ann. of Math.* (2) **187**:3 (2018), 825–867. [MR](#) [Zbl](#)
- [de Branges 1968] L. de Branges, *Hilbert spaces of entire functions*, Prentice-Hall, Englewood Cliffs, NJ, 1968. [MR](#) [Zbl](#)
- [Han and Schlag 2020] R. Han and W. Schlag, “A higher-dimensional Bourgain–Dyatlov fractal uncertainty principle”, *Anal. PDE* **13**:3 (2020), 813–863. [MR](#) [Zbl](#)
- [Havin and Jöricke 1994] V. Havin and B. Jöricke, *The uncertainty principle in harmonic analysis*, *Ergebnisse der Math.* (3) **28**, Springer, 1994. [MR](#) [Zbl](#)
- [Kislyakov and Perstneva 2021] S. V. Kislyakov and P. S. Perstneva, “Indicator functions with uniformly bounded Fourier sums and large gaps in the spectrum”, *J. Fourier Anal. Appl.* **27**:2 (2021), art. id. 33. [MR](#) [Zbl](#)
- [Koosis 1988] P. Koosis, *The logarithmic integral, I*, *Cambridge Stud. Adv. Math.* **12**, Cambridge Univ. Press, 1988. [MR](#) [Zbl](#)
- [Koosis 1992] P. Koosis, *The logarithmic integral, II*, *Cambridge Stud. Adv. Math.* **21**, Cambridge Univ. Press, 1992. [MR](#) [Zbl](#)
- [Koosis] P. Koosis, “Kargaev’s proof of Beurling’s lemma”, unpublished manuscript.
- [Makarov and Poltoratski 2010] N. Makarov and A. Poltoratski, “Beurling–Malliavin theory for Toeplitz kernels”, *Invent. Math.* **180**:3 (2010), 443–480. [MR](#) [Zbl](#)
- [Mashregi et al. 2005] D. Mashregi, F. L. Nazarov, and V. P. Khavin, “The Beurling–Malliavin multiplier theorem: the seventh proof”, *Algebra i Analiz* **17**:5 (2005), 3–68. In Russian; translated in *St. Petersburg Math. J.* **17**:5 (2006), 699–744. [MR](#) [Zbl](#)
- [Nazarov and Olevskii 2018] F. Nazarov and A. Olevskii, “A function with support of finite measure and ‘small’ spectrum”, pp. 389–393 in *50 years with Hardy spaces*, edited by A. Baranov et al., *Oper. Theory Adv. Appl.* **261**, Birkhäuser, Cham, 2018. [MR](#) [Zbl](#)
- [Nikolski 2012] N. Nikolski, *Éléments d’analyse avancée, I: Espaces de Hardy*, Belin, Paris, 2012.
- [Poltoratski 2012] A. Poltoratski, “Spectral gaps for sets and measures”, *Acta Math.* **208**:1 (2012), 151–209. [MR](#) [Zbl](#)
- [Redheffer 1977] R. M. Redheffer, “Completeness of sets of complex exponentials”, *Adv. Math.* **24**:1 (1977), 1–62. [MR](#) [Zbl](#)
- [Stolyarov and Zatitskiy 2021] D. Stolyarov and P. Zatitskiy, “Sharp transference principle for BMO and A_p ”, *J. Funct. Anal.* **281**:6 (2021), art. id. 109085. [MR](#) [Zbl](#)
- [Vasilyev 2022] I. Vasilyev, “On the multidimensional Nazarov lemma”, *Proc. Amer. Math. Soc.* **150**:4 (2022), 1601–1611. [MR](#) [Zbl](#)

Received 10 Oct 2022. Revised 21 Jun 2023. Accepted 18 Jul 2023.

IOANN VASILYEV: ivasilyev@pdmi.ras.ru

St. Petersburg Department of Steklov Mathematical Institute Russian Academy of Sciences (PDMI RAS) St. Petersburg, Russia

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 9 2024

Relative heat content asymptotics for sub-Riemannian manifolds ANDREI AGRACHEV, LUCA RIZZI and TOMMASO ROSSI	2997
Minkowski inequality on complete Riemannian manifolds with nonnegative Ricci curvature LUCA BENATTI, MATTIA FOGAGNOLO and LORENZO MAZZIERI	3039
The Willmore flow of tori of revolution ANNA DALL'ACQUA, MARIUS MÜLLER, REINER SCHÄTZLE and ADRIAN SPENER	3079
Optimal Prandtl expansion around a concave boundary layer DAVID GÉRARD-VARET, YASUNORI MAEKAWA and NADER MASMOUDI	3125
Nonlocal operators related to nonsymmetric forms, II: Harnack inequalities MORITZ KASSMANN and MARVIN WEIDNER	3189
Transference of scale-invariant estimates from Lipschitz to nontangentially accessible to uniformly rectifiable domains STEVE HOFMANN, JOSÉ MARÍA MARTELL and SVITLANA MAYBORODA	3251
Optimal regularity and the Liouville property for stable solutions to semilinear elliptic equations in \mathbb{R}^n with $n \geq 10$ FA PENG, YI RU-YA ZHANG and YUAN ZHOU	3335
A generalization of the Beurling–Malliavin majorant theorem IOANN VASILYEV	3355