

ANALYSIS & PDE

Volume 18

No. 3

2025

LÉO BIGORGNE

**GLOBAL EXISTENCE AND MODIFIED SCATTERING
FOR THE SOLUTIONS TO THE VLASOV-MAXWELL SYSTEM
WITH A SMALL DISTRIBUTION FUNCTION**



GLOBAL EXISTENCE AND MODIFIED SCATTERING FOR THE SOLUTIONS TO THE VLASOV–MAXWELL SYSTEM WITH A SMALL DISTRIBUTION FUNCTION

LÉO BIGORGNE

The purpose of this paper is two-fold. In the first part, we provide a new proof of the global existence of the solutions to the Vlasov–Maxwell system with a small initial distribution function. Our approach relies on vector field methods, together with the Glassey–Strauss decomposition of the electromagnetic field, and does not require any support restriction on the initial data or smallness assumption on the Maxwell field. Contrary to previous works on Vlasov systems in dimension 3, we do not modify the linear commutators and avoid then many technical difficulties.

In the second part of this paper, we prove a modified scattering result for these solutions. More precisely, we obtain that the electromagnetic field has a radiation field along future null infinity and approaches, for large time, a smooth solution to the vacuum Maxwell equations. As for the Vlasov–Poisson system, in contrast, the distribution function converges to a new density function f_∞ along *modifications* of the characteristics of the free relativistic transport equation. In order to define these logarithmic corrections, we identify an effective asymptotic Lorentz force. By considering logarithmical modifications of the linear commutators, defined in terms of derivatives of the asymptotic Lorentz force, we finally prove higher-order regularity results for f_∞ .

1. Introduction	629
2. Preliminaries and detailed statement of the main result	633
3. Strategy of the proof and the bootstrap assumptions	650
4. Estimates for the distribution function	652
5. Improvement of the bootstrap assumptions on the electromagnetic field	663
6. Modified scattering for the distribution function	675
7. Scattering result for the electromagnetic field	695
8. Conservation of the total energy of the system	705
Appendix A. Estimates for the gradients of the kernels	707
Appendix B. The radiation field of the derivatives of the Maxwell field	707
Appendix C. Remarks on F^∞ and the modified characteristics	710
Acknowledgements	711
References	711

1. Introduction

The Vlasov–Maxwell system, which is used to describe the dynamics of collisionless plasma, can be written as

MSC2020: 35A01, 35B40, 35L99.

Keywords: Vlasov–Maxwell, scattering, asymptotic, stability.

$$\partial_t f + \hat{v} \cdot \nabla_x f + (E + \hat{v} \times B) \cdot \nabla_v f = 0, \quad (1)$$

$$\nabla_x \cdot E = \int_{\mathbb{R}_v^3} f \, dv, \quad \partial_t E = \nabla_x \times B - \int_{\mathbb{R}_v^3} \hat{v} f \, dv, \quad (2)$$

$$\nabla_x \cdot B = 0, \quad \partial_t B = -\nabla_x \times E, \quad (3)$$

where

- $f : \mathbb{R}_{+,t} \times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}_+$ is the density distribution function of the particles,
- $\hat{v} = v/v^0$, with $v^0 := \sqrt{1 + |v|^2}$, is the relativistic speed of a particle of momentum $v \in \mathbb{R}_v^3$,
- $\int_{\mathbb{R}_v^3} f \, dv$ and $\int_{\mathbb{R}_v^3} \hat{v} f \, dv$ are respectively the total charge density and the total current density,
- $E, B : \mathbb{R}_{+,t} \times \mathbb{R}_x^3 \rightarrow \mathbb{R}^3$ are respectively the electric and the magnetic field.

For simplicity, we assume that the plasma is composed of one species of particles of charge $q = 1$ and mass $m = 1$. Our results can be extended without any additional difficulty to several families of particles of different charges and positive masses.¹ We refer to [Glassey 1996] for a detailed introduction to these equations.

The initial value problem for the Vlasov–Maxwell equations, together with a regular initial data set (f_0, E_0, B_0) composed of a function $f_0 : \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}_+$ and two fields $E_0, B_0 : \mathbb{R}_x^3 \rightarrow \mathbb{R}^3$ satisfying the constraint equations $\nabla_x \cdot E_0 = \int_v f_0 \, dv$ and $\nabla_x \cdot B_0 = 0$, is well-posed [Wollman 1984]. On the other hand, the global existence problem for classical solutions to the Vlasov–Maxwell system is still open² and has only been addressed in some particular cases, such as under certain symmetry assumptions [Glassey and Schaeffer 1990; 1997; 1998; Luk and Strain 2016; Rein 1990; Wang 2022a]. For the general case, since the pioneering work [Glassey and Strauss 1986], several continuation criteria have been obtained [Glassey and Strauss 1987b; 1989; Klainerman and Staffilani 2002; Bouchut et al. 2003; Pallard 2005; 2015; Sospedra-Alfonso and Illner 2010; Luk and Strain 2014; Kunze 2015; Patel 2018].

1.1. Small data solutions to the Vlasov–Maxwell system. Much more is known for this particular perturbative regime, in which global existence holds and the solutions disperse. For small compactly supported initial data Glassey and Strauss [1987a] proved the optimal decay rate $\int_v f \, dv \lesssim t^{-3}$ on the velocity average of the distribution function and obtained estimates for the electromagnetic field and its first-order derivatives. Shortly after, in the multispecies case, the smallness assumptions on the individual particle densities was relaxed by [Glassey and Schaeffer 1988]. Later, Schaeffer [2004] removed the support restriction on the velocity variable. However, his method leads to a loss on the estimate of $\int_v f \, dv$.

It is only recently that all the compact support assumptions on the initial data were removed in two independent results [Bigorgne 2020a; Wang 2022b]. Both of these works are based on vector field methods and the latter used also Fourier analysis. These robust approaches allow for the derivation of sharp pointwise decay estimates on the solutions and their (high-order) derivatives. Moreover, in [Bigorgne 2020a], the initial decay hypothesis in v is optimal and improved estimates on certain *null* components of the electromagnetic field are derived. Finally, using the framework of Glassey and Strauss

¹The case of massless particles requires in fact a different analysis [Bigorgne 2021b].

²In contrast, a global in time existence result for weak solutions was proved in [DiPerna and Lions 1989] and revisited in [Rein 2004].

and without any compact support restriction, Wei and Yang [2021] derived a global existence result which does not require the initial Maxwell field to be small.

In the first part of this article, we provide an alternative but shorter proof of the main results of [Bigorgne 2020a; Wang 2022b], without assuming any smallness assumption on the electromagnetic field. Compared to [Wei and Yang 2021], we require more regularity on the initial data but our method allows us to control the derivatives of the solutions, up to an arbitrary order N . This information is needed for the second part of the paper.

1.2. Modified scattering results for the Vlasov–Poisson system. Sharp decay estimates for the small data solutions to the Vlasov–Poisson system were first derived by [Bardos and Degond 1985] and then, with various improvements, by [Hwang et al. 2011; Smulevici 2016; Duan 2022; Schaeffer 2021] (for the relativistic cases, see [Glasse and Schaeffer 1985; Wang 2023; Bigorgne 2020b]). Modified scattering for these solutions was established in [Choi and Kwon 2016] and then in [Ionescu et al. 2022; Pankavich 2022], where more information was obtained on the asymptotic dynamics governing the modification of the linear characteristics. Furthermore, a scattering map has been constructed by [Flynn et al. 2023] and let us finally mention that similar results hold for perturbations of a point charge [Pausader and Widmayer 2021; Pausader et al. 2024].

In the second part of this paper, we investigate such problems in the context of the Vlasov–Maxwell equations. In particular, as in [Ionescu et al. 2022] for the Vlasov–Poisson system, we prove that

$$\int_{\mathbb{R}^3} f(t, x, v) \, dx \rightarrow Q_\infty(v) \quad \text{as } t \rightarrow +\infty.$$

The scattering charge Q_∞ is deeply related to the leading-order term of the asymptotic expansion of both the charge density $\int_v f \, dv$ and the current density. It allows us to define an asymptotic Lorentz force $v \mapsto \text{Lor}(v)$, from which we deduce the modified scattering statement for f (see Theorem 1.1 and Remark 1.3 for more details). We also prove higher-order regularity properties for the limit distribution f_∞ , which require a more thorough analysis. To our knowledge, there is no such regularity result for the Vlasov–Poisson system.

1.3. Vector field methods for relativistic transport equations. Our analysis of the asymptotic behavior of both the electromagnetic field and the distribution function relies on vector field methods (see Section 2.4 for an overview of the key ideas). This kind of technique was first developed by Klainerman [1985] in order to study solutions to nonlinear wave equations and then adapted in [Christodoulou and Klainerman 1990] to the Maxwell equations. It is only recently that the approach has been adapted to relativistic transport equations by Fajman, Joudioux and Smulevici [Fajman et al. 2017], leading in particular to a proof of the stability of Minkowski spacetime for both the massive and massless Einstein–Vlasov system [Fajman et al. 2021; Bigorgne et al. 2021] (see also [Lindblad and Taylor 2020; Taylor 2017] for alternative proofs). Our work [Bigorgne 2020a] concerning the small data solutions to the Vlasov–Maxwell system relies on such techniques as well. The method has also been successfully used to derive boundedness and decay estimates for the solutions to the massless Vlasov equation on a fixed Kerr black hole [Andersson et al. 2018; Bigorgne 2023]. Finally, even if it concerns the nonrelativistic setting, let us also mention

that such approaches have been applied in the collisional regime [Chaturvedi 2021; 2022; Chaturvedi et al. 2023].

In order to deal with slowly decaying error terms, all the works on the small data solutions to massive relativistic Vlasov systems or the Vlasov–Poisson system [Fajman et al. 2021; Bigorgne 2020a; Smulevici 2016; Duan 2022], based on vector field methods, require dynamically modifying certain linear commutators of the Vlasov operator. One of the main novelties of this article consists in proving that the solutions are global without using these modified vector fields, which considerably simplifies the analysis. For this, even though certain quantities grow logarithmically in time, we are able to close the energy estimates by identifying several hierarchies in the commuted equations (see Section 2.8.2 for more details). It is then important to derive the optimal decay rate t^{-3} for $\int_v f \, dv$ and its derivatives by a method allowing well-chosen weighted $W_{x,v}^{N,\infty}$ norms of the distribution function to grow slowly in time. We believe that this approach could be applied to other systems of equations, in particular for both the Einstein–Vlasov and the Vlasov–Poisson systems.

1.4. The main result. We present here a short version of our main result, stated in Theorems 2.10–2.11 below, where we also describe the behavior of the derivatives of the solutions.

Theorem 1.1. *Any solution (f, E, B) to the Vlasov–Maxwell system (1)–(3) arising from a small initial distribution function and smooth as well as sufficiently decaying initial data is global in time. Moreover:*

(1) *There exists a solution $(E^{\text{vac}}, B^{\text{vac}})$ to the vacuum Maxwell equations³ approaching (E, B) as $t + |x| \rightarrow \infty$,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |E - E^{\text{vac}}|(t, x) + |B - B^{\text{vac}}|(t, x) \leq C_q (1 + t + |x|)^{-1-q}, \quad \frac{1}{2} \leq q < 1.$$

(2) *The Lorentz force has a self-similar asymptotic profile $v \mapsto \text{Lor}(v)$,*

$$\forall (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |t^2(E(t, x + t\hat{v}) + \hat{v} \times B(t, x + t\hat{v})) - \text{Lor}(v)| \lesssim \langle x \rangle^2 |v^0|^8 \frac{\log^n(3+t)}{1+t},$$

where $\langle x \rangle := (1 + |x|^2)^{1/2}$ and, say, $n = 70$. We have modified scattering to a new density function $f_\infty : \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}_+$,

$$\forall t \geq 3, \quad \|f(t, X_\mathcal{C}(t, \cdot, \cdot), \cdot) - f_\infty\|_{L_{x,v}^1 \cap L_{x,v}^\infty} \lesssim t^{-1} \log^n(t),$$

where the Cartesian components $X_\mathcal{C}^k$ of the modified spatial characteristics $X_\mathcal{C} \in \mathbb{R}_x^3$ are defined as

$$X_\mathcal{C}^k(t, x, v) := x^k + t\hat{v}^k - \frac{\log(t)}{v^0} (\text{Lor}^k(v) - \hat{v} \cdot \text{Lor}(v)\hat{v}^k), \quad 1 \leq k \leq 3.$$

Remark 1.2. No modification of the spatial characteristics is in fact required in the exterior of the light cone $\{|x| \geq t\}$ in order to prove such a result (see Section C.2). We already observed in [Bigorgne 2021a] that the small data solutions to the Vlasov–Maxwell system have better behavior in this region.

Similarly, no correction of the linear characteristics should in principle be necessary in order to prove a scattering statement in higher dimensions. This is consistent with the result of [Pankavich 2023]

³The vacuum Maxwell equations are given by (2)–(3) with $f = 0$.

concerning the Vlasov–Poisson system in dimension $d \geq 4$ and our study of the asymptotic behavior of the small data solutions of the Vlasov–Maxwell system in high dimensions [Bigorgne 2022].

The case of massless particles differs from the case of massive particles treated in this paper. Indeed, in view of [Bigorgne 2021b], we expect the small data solutions of the massless Vlasov–Maxwell system to satisfy linear scattering in dimension $d = 3$.

Remark 1.3. The behavior of the Lorentz force along the linear trajectories suggests that the characteristics of the Vlasov–Maxwell system satisfy, for $t \gg 1$,

$$\dot{X} = \widehat{V}, \quad \dot{V} \approx t^{-2} \text{Lor}(V), \quad X(0) = x_0, \quad V(0) = v_0.$$

Hence, we can presume that V converges to v , so that

$$V(t) \approx v - \frac{1}{t} \text{Lor}(v), \quad \dot{X}(t) \approx \widehat{v} - \frac{1}{tv^0} \text{Lor}(v) + \frac{\widehat{v} \cdot \text{Lor}(v)}{tv^0} \widehat{v} + O(t^{-2}),$$

and we can then expect $X(t) \approx X_{\mathcal{G}}(t, x_0, v)$.

Moreover, we could in fact decompose $\text{Lor}(v)$ as $E^\infty(v) + \widehat{v} \times B^\infty(v)$ and observe that, as $v \rightarrow 0$,

$$X_{\mathcal{G}}(t, x, v) = x + tv - \log(t)E^\infty(v) + o(v).$$

In other words, for small velocities, the modified characteristics $X_{\mathcal{G}}$ of the Vlasov–Maxwell system approach the ones constructed in [Ionescu et al. 2022] for the Vlasov–Poisson system.

1.5. Structure of the paper. In Section 2 we introduce the notations and the tools used throughout this article. Then, we state our main results, Theorems 2.10–2.11, and present the key ideas of the proof. In Section 3, we set up the bootstrap assumptions and discuss their immediate consequences. Section 4 concerns the study of the distribution function. In particular, we prove that weighted $L_{x,v}^\infty$ norms of f and its derivatives grow at most logarithmically and we improve the bootstrap assumption on their velocity average. Then, in Section 5, we conclude the proof of the global existence of the small data solutions to (1)–(3) by exploiting the Glassey–Strauss decomposition of the electromagnetic field in order to improve the bounds on (E, B) and their derivatives. Next, in Section 6 we refine our estimates by proving that the particle current density and the electromagnetic field have a self-similar asymptotic profile. This allows us to define the modified trajectories along which the distribution function converges. Section 7 is devoted to the scattering results for the electromagnetic field. A crucial part of the proof consists in constructing a scattering map for the vacuum Maxwell equations. In Section 8, we relate the conserved total energy of the system to the ones of the scattering states. Finally, Appendices A and B contain two useful computations and Appendix C presents alternative expressions for the profile of F and the modified characteristics.

2. Preliminaries and detailed statement of the main result

2.1. Basic notations. In this paper we work on the 1+3-dimensional Minkowski spacetime (\mathbb{R}^{1+3}, η) . We will use two sets of coordinates, the Cartesian $(x^0 = t, x^1, x^2, x^3)$, in which $\eta = \text{diag}(-1, 1, 1, 1)$, and null coordinates $(u, \underline{u}, \theta, \varphi)$, where

$$\underline{u} = t + r, \quad u = t - r, \quad r := |x| = \sqrt{|x^1|^2 + |x^2|^2 + |x^3|^2},$$

and $(\theta, \varphi) \in]0, \pi[\times]0, 2\pi[$ are spherical coordinates on the spheres of constant (t, r) . These coordinates are defined globally on \mathbb{R}^{1+3} apart from the usual degeneration of spherical coordinates and at $r = 0$. Sometimes, for a tensor field T defined on $\mathbb{R}_+ \times \mathbb{R}_x^3$, it will be convenient to write

$$T(u, \underline{u}, \omega) := T\left(\frac{u+u}{2}, \frac{u-u}{2}\omega\right), \quad \underline{u} \geq 0, \quad |u| \leq \underline{u}, \quad \omega \in \mathbb{S}^2.$$

We will work with the null frame $(L, \underline{L}, e_\theta, e_\varphi)$, where $L = 2\partial_u$, $\underline{L} = 2\partial_{\underline{u}}$ are null derivatives and (e_θ, e_φ) is the standard orthonormal basis on the spheres. More precisely,

$$L = \partial_t + \partial_r, \quad \underline{L} = \partial_t - \partial_r, \quad e_\theta = \frac{1}{r}\partial_\theta, \quad e_\varphi = \frac{1}{r \sin \theta}\partial_\varphi.$$

The Einstein summation convention will often be used; for instance $v^\mu \partial_{x^\mu} f = \sum_{\mu=0}^3 v^\mu \partial_{x^\mu} f$. The Latin indices goes from 1 to 3 and the Greek indices from 0 to 3. We will raise and lower indices using the Minkowski metric η , so that $x^i = x_i$ and $x^0 = -x_0$.

The four-momentum vector $\mathbf{v} = (v^\mu)_{0 \leq \mu \leq 3}$ is parametrized by $v = (v^i)_{1 \leq i \leq 3} \in \mathbb{R}_v^3$ and $v^0 = \sqrt{1 + |v|^2}$ since the mass of the particles is equal to 1. Let $(v^L, v^{\underline{L}}, v^{e_1}, v^{e_2})$ be the null components of the momentum vector and $\not{v} = (v^{e_\theta}, v^{e_\varphi})$ its angular part, so that

$$\mathbf{v} = v^L L + v^{\underline{L}} \underline{L} + v^{e_\theta} e_\theta + v^{e_\varphi} e_\varphi, \quad v^L = \frac{v^0 + (x_i/r)v^i}{2}, \quad v^{\underline{L}} = \frac{v^0 - (x_i/r)v^i}{2}, \quad |\not{v}|^2 = |v^{e_\theta}|^2 + |v^{e_\varphi}|^2.$$

The relativistic speed $\hat{v} \in \mathbb{R}^3$ is given by $\hat{v}^i = v^i/v^0$ and, for convenience, we define the quantities

$$\hat{v}^0 := \frac{v^0}{v^0} = 1, \quad \hat{v}^L := \frac{v^L}{v^0}, \quad \hat{v}^{\underline{L}} := \frac{v^{\underline{L}}}{v^0}, \quad \hat{\not{v}} := \frac{\not{v}}{v^0}, \quad \hat{v}^{e_A} := \frac{v^{e_A}}{v^0}, \quad A \in \{\theta, \varphi\}.$$

Sometimes, we will write $(|v^0|^p g)(w)$ to denote $|w^0|^p g(w)$, where $w \in \mathbb{R}_v^3$ and $g : \mathbb{R}_v^3 \rightarrow \mathbb{R}$.

In order to capture the good properties of certain geometric quantities associated to the solutions in the good null directions (L, e_θ, e_φ) , we introduce the Faraday tensor $F_{\mu\nu}$, which is a 2-form, and the four-current density $J(f)_\mu$,

$$F = \begin{pmatrix} 0 & E^1 & E^2 & E^3 \\ -E^1 & 0 & -B^3 & B^2 \\ -E^2 & B^3 & 0 & -B^1 \\ -E^3 & -B^2 & B^1 & 0 \end{pmatrix}, \quad J(f) := \begin{pmatrix} -\int_{\mathbb{R}_v^3} f \, dv \\ \int_{\mathbb{R}_v^3} (v_1/v^0) f \, dv \\ \int_{\mathbb{R}_v^3} (v_2/v^0) f \, dv \\ \int_{\mathbb{R}_v^3} (v_3/v^0) f \, dv \end{pmatrix}. \tag{4}$$

The Cartesian components of F are then either equal to 0 or to a component of $\pm(E, B)$. We will in fact be more interested in its null decomposition $(\alpha(F), \underline{\alpha}(F), \rho(F), \sigma(F))$ defined, for $A \in \{\theta, \varphi\}$, as

$$\alpha(F)_{e_A} := F_{e_A L}, \quad \underline{\alpha}(F)_{e_A} := F_{e_A \underline{L}}, \quad \rho(F) := \frac{1}{2} F_{\underline{L} L}, \quad \sigma(F) := F_{e_\theta e_\varphi}. \tag{5}$$

In particular, $\rho(F) = E \cdot \partial_r$ and $-\sigma(F) = B \cdot \partial_r$ are the radial components of the electric field and the magnetic field. Moreover, the 1-forms $\alpha(F)$ and $\underline{\alpha}(F)$ are tangential to the 2-spheres and we will use the

pointwise norms

$$|\alpha(F)|^2 := |\alpha(F)_{e_\theta}|^2 + |\alpha(F)_{e_\varphi}|^2, \quad |\underline{\alpha}(F)|^2 := |\underline{\alpha}(F)_{e_\theta}|^2 + |\underline{\alpha}(F)_{e_\varphi}|^2, \\ |F|^2 := \sum_{0 \leq \mu < \nu \leq 3} |F_{\mu\nu}|^2 = \frac{1}{2}|\alpha(F)|^2 + \frac{1}{2}|\underline{\alpha}(F)|^2 + |\rho(F)|^2 + |\sigma(F)|^2.$$

The Vlasov equation (1) can be rewritten as

$$\mathbf{T}_F(f) = 0, \quad \text{where } \mathbf{T}_F : f \mapsto \partial_t f + \hat{v} \cdot \nabla_x f + \hat{v}^\mu F_\mu^j \partial_{v^j} f, \tag{6}$$

and the Maxwell equations (2)–(3) take a concise form. The Gauss–Ampère law and the Gauss–Faraday law⁴

$$\nabla^\mu F_{\mu\nu} = J(f)_\nu, \quad \nabla^\mu {}^*F_{\mu\nu} = 0, \tag{7}$$

where ${}^*F_{\mu\nu} = \frac{1}{2}F^{\lambda\sigma} \varepsilon_{\lambda\sigma\mu\nu}$ is the Hodge dual of F and ε is the Levi-Civita symbol. Here ∇ stands for the covariant derivative (or Levi-Civita connection), so that (7) holds in any coordinate system.

The operators ∇_x and ∇_v will denote the standard gradients in x and v respectively. For instance,

$$\nabla_x f = (\partial_{x^1} f, \partial_{x^2} f, \partial_{x^3} f), \quad \nabla_v f = (\partial_{v^1} f, \partial_{v^2} f, \partial_{v^3} f).$$

Given a 2-form G and $0 \leq \lambda \leq 3$, we will denote by $\nabla_{\partial_{x^\lambda}} G$ the covariant derivative of G according to ∂_{x^λ} , where $\partial_{x^0} = \partial_t$. For any multi-index $\kappa \in \{0, 1, 2, 3\}^p$, we define $\nabla_{t,x}^\kappa G := \nabla_{\partial_{x^{\kappa_1}}} \cdots \nabla_{\partial_{x^{\kappa_p}}} G$. In Cartesian coordinates, we then have

$$\nabla_{t,x}^\kappa (G)_{\mu\nu} = \partial_{t,x}^\kappa (G_{\mu\nu}), \quad 0 \leq \mu, \nu \leq 3.$$

Finally, for $x \in \mathbb{R}^3$ we will use the Japanese brackets $\langle x \rangle := (1 + |x|^2)^{1/2}$ and the notation $D_1 \lesssim D_2$ will stand for the statement that there exists $C > 0$ a positive constant independent of the solutions such as $D_1 \leq C D_2$.

2.2. Backward light cones and future null infinity. The scattering state for a smooth electromagnetic field F , which in our case is also called radiation field, will be a tensor field depending on the variables $(u, \omega) \in \mathbb{R} \times \mathbb{S}^2$. It will be obtained as the limit, when $\underline{u} \rightarrow +\infty$, of $rF(u, \underline{u}, \omega)$. For this reason, we introduce the backward light cones $\underline{C}_{\underline{u}}$ and give their induced volume form $d\mu_{\underline{C}_{\underline{u}}}$ in accordance with the choice of the null vector field \underline{L} as their normal vector. Let, for any $\underline{u} \geq 0$,

$$\underline{C}_{\underline{u}} := \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3 \mid t + |x| = \underline{u}\}, \quad d\mu_{\underline{C}_{\underline{u}}} = \frac{1}{2}r^2 du d\mu_{\mathbb{S}^2},$$

where $d\mu_{\mathbb{S}^2} = \sin(\theta) d\theta d\varphi$ is the volume form on \mathbb{S}^2 .

Even if we will not need this formalism, we mention that the radiation field is in fact defined on a part of the conformal boundary of the Minkowski space, called future null infinity \mathcal{I}^+ and corresponding to the future end points of the null geodesics $t - |x| = u$. It can be viewed as $\underline{C}_{+\infty}$. More precisely,

$$(t, r, \omega) \mapsto (T(t, r) = \tan^{-1}(t+r) + \tan^{-1}(t-r), \quad R(t, r) = \tan^{-1}(t+r) - \tan^{-1}(t-r), \omega) \in \mathbb{R} \times \mathbb{S}^3$$

⁴Note that $\nabla^\mu {}^*F_{\mu\nu} = 0$ is equivalent to $\nabla_{[\lambda} F_{\mu\nu]} := \nabla_\lambda F_{\mu\nu} + \nabla_\mu F_{\nu\lambda} + \nabla_\nu F_{\lambda\mu} = 0$.




Figure 1. The set \underline{C}_u and the Penrose diagram of the Minkowski space.

is a conformal diffeomorphism between Minkowski spacetime and the interior of the triangle $0 \leq R \leq \pi$, $|T| = \pi - R$ of the space $\mathbb{R} \times \mathbb{S}^3$, equipped with the metric $-dT^2 + dR^2 + \sin^2(R) d\mu_{\mathbb{S}^2}$. Then

$$\mathcal{I}^+ := \{(T, R, \omega) \in \mathbb{R} \times \mathbb{S}^3 \mid 0 < R < \pi, T = \pi - R\}.$$

Past null infinity \mathcal{I}^- is defined similarly as $\{0 < R < \pi, T = R - \pi\}$ and can be viewed as $t - |x| = -\infty$. See Figure 1.

2.3. Charged electromagnetic field. For our global existence result, it will be sufficient to assume that the electromagnetic field satisfies $|F|(0, \cdot) \lesssim r^{-2}$, whereas our scattering result will require a slightly stronger initial decay hypothesis. However, if the plasma is not neutral, one cannot expect F to decay faster than r^{-2} . Indeed, if (f, F) is a sufficiently regular solution to (6)–(7) on $[0, T]$, we obtain from Gauss’s law that the total charge

$$Q_F(t) := \lim_{r \rightarrow +\infty} \int_{\omega \in \mathbb{S}^2} \rho(F)(t, r\omega) r^2 d\mu_{\mathbb{S}^2} = \int_{x \in \mathbb{R}^3} \int_{v \in \mathbb{R}^3} f(t, x, v) dv dx, \quad t \in [0, T],$$

is a conserved quantity and that $|F| = o(r^{-2})$ implies $Q_F = 0$. In order to avoid such a restrictive assumption, we introduce the pure charge part \bar{F} of F ,

$$\bar{F}(t, x) := \frac{Q_F}{4\pi|x|^2} \frac{x_i}{|x|} dt \wedge dx^i, \quad \rho(\bar{F})(t, x) = \frac{Q_F}{4\pi|x|^2}, \quad \alpha(\bar{F}) = \underline{\alpha}(\bar{F}) = \sigma(\bar{F}) = 0, \quad (8)$$

which corresponds to the electromagnetic field generated by a point charge Q_F at $x = 0$. One can verify that $Q_{\bar{F}} = Q_F$, so that $F - \bar{F}$ is chargeless and it will then be consistent to assume that F has an asymptotic expansion of the form $F = \bar{F} + O(r^{-2-\delta})$, $\delta > 0$. In fact, $E = E^{\text{df}} + E^{\text{cf}}$ and $B = B^{\text{df}} + B^{\text{cf}}$ can be decomposed into their divergence-free and curl-free components. Then, $B^{\text{cf}} = 0$ and $E^{\text{cf},i} = \bar{F}_{0i} + O(r^{-3})$ if $J(f)_0$ is sufficiently regular, so that the stronger initial decay assumption required for the scattering result concerns the divergence-free components of E and B .

2.4. Commutation vector fields. We will derive estimates on both the electromagnetic field and the distribution function using vector field methods. These kinds of approaches are usually based on

- a set of vector fields, which commute with the linear operator of the equation studied,
- energy inequalities, applied in order to prove boundedness for L^2 or L^1 norms of the solutions and their derivatives (for instance, see [Bigorgne 2020a, Section 4.1]),
- weighted Sobolev embeddings, such as [Fajman et al. 2017, Theorem 6], used to obtain decay estimates on the fields.

In this paper, in order to simplify the analysis, we will prove L^∞ estimates and then obtain pointwise decay estimates on the solutions in a different way (see Section 2.8 for more details). We now elaborate on the commutators for the Maxwell equations and the ones for the relativistic transport equation.

Definition 2.1. Let \mathbb{K} be the set composed of the vector fields

$$\partial_t, \quad \partial_{x^i}, \quad \Omega_{0i} := t \partial_{x^i} + x^i \partial_t, \quad \Omega_{jk} := x^j \partial_{x^k} - x^k \partial_{x^j}, \quad S := t \partial_t + x^\ell \partial_{x^\ell} = t \partial_t + r \partial_r,$$

where $1 \leq i \leq 3$ and $1 \leq j < k \leq 3$. The translations ∂_{x^μ} , the Lorentz boosts Ω_{0i} and the rotations Ω_{jk} are Killing vector fields, so that they generate isometries of the Minkowski space. The scaling vector field S is merely conformal Killing.

We will use this set for differentiating the electromagnetic field geometrically. More precisely, for a 2-form F and a vector field $Z = Z^\mu \partial_{x^\mu}$, the Lie derivative $\mathcal{L}_Z(F)$ of F with respect to Z is given, in coordinates, by

$$\mathcal{L}_Z(F)_{\mu\nu} = Z(F_{\mu\nu}) + \partial_\mu(Z^\lambda)F_{\lambda\nu} + \partial_\nu(Z^\lambda)F_{\mu\lambda}.$$

Furthermore, if F is a smooth solution to the vacuum Maxwell equations $\nabla^\mu F_{\mu\nu} = \nabla^{\mu*}F_{\mu\nu} = 0$ and $Z \in \mathbb{K}$, then $\mathcal{L}_Z(F)$ is also a solution to the vacuum Maxwell equations, that is, $\nabla^\mu \mathcal{L}_Z(F)_{\mu\nu} = \nabla^{\mu*} \mathcal{L}_Z(F)_{\mu\nu} = 0$.

Definition 2.2. Let $\widehat{\mathbb{P}}_0$ be the set composed of

$$\partial_t, \quad \partial_{x^i}, \quad \widehat{\Omega}_{0i} := t \partial_{x^i} + x^i \partial_t + v^0 \partial_{v^i}, \quad \widehat{\Omega}_{jk} := x^j \partial_{x^k} - x^k \partial_{x^j} + v^j \partial_{v^k} - v^k \partial_{v^j}, \quad S = t \partial_t + r \partial_r,$$

where $1 \leq i \leq 3$ and $1 \leq j < k \leq 3$. In fact, $\widehat{\partial}_{x^\mu} = \partial_{x^\mu}$, $\widehat{\Omega}_{0i}$ and $\widehat{\Omega}_{jk}$ are obtained as the complete lift, a classical operation in differential geometry,⁵ of the Killing fields ∂_{x^μ} , Ω_{0i} and Ω_{jk} .

These vector fields have good commutation properties with the linear transport operator $T_0 = \partial_t + \widehat{v} \cdot \nabla_x$. Indeed, $[T_0, S] = T_0$ and $[v^0 T_0, \widehat{Z}] = 0$ for all $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{S\}$.

In order to consider higher-order derivatives, we introduce an ordering on $\mathbb{K} = \{Z^i \mid 1 \leq i \leq 11\}$ and on $\widehat{\mathbb{P}}_0 = \{\widehat{Z}^i \mid 1 \leq i \leq 11\}$. It will be convenient to assume that $Z^{11} = \widehat{Z}^{11} = S$ and $\widehat{Z}^i = \widehat{Z}^i$ for any $1 \leq i \leq 10$. Moreover, for a multi-index $\beta \in \llbracket 1, 11 \rrbracket^p$ of length $|\beta| = p$, we denote by \mathcal{L}_{Z^β} the

⁵We refer to [Fajman et al. 2017, Section 2G] for more details about the relations between the Vlasov operator on a Lorentzian manifold and the complete lift of its Killing vector fields.

Lie derivative $\mathcal{L}_{Z^{\beta_1}} \cdots \mathcal{L}_{Z^{\beta_p}}$ of order $|\beta|$. Similarly, we define \widehat{Z}^β as $\widehat{Z}^{\beta_1} \cdots \widehat{Z}^{\beta_p}$. Note the equivalence between the pointwise norms

$$\sum_{|\gamma| \leq N} |\mathcal{L}_{Z^\gamma}(F)| \lesssim \sum_{|\beta| \leq N} \sum_{0 \leq \mu, \nu \leq 3} |Z^\beta(F_{\mu\nu})| \lesssim \sum_{|\gamma| \leq N} |\mathcal{L}_{Z^\gamma}(F)|. \tag{9}$$

Since $\mathcal{L}_{\partial_{x^\mu}}(F)$ and $\partial_{x^\mu} f$ have better behavior than the other derivatives, it will be crucial, in order to identify certain hierarchies in the commuted equations, to count the number of homogeneous vector fields composing Z^β or \widehat{Z}^β . We denote by β_H (respectively β_T) the number of homogeneous vector fields Ω_{0i} , Ω_{jk} and S (respectively translations ∂_{x^μ}) composing Z^β . Note that $\beta_H + \beta_T = |\beta|$ and that \widehat{Z}^β is also composed of β_H homogenous vector fields and β_T translations. If $Z^\beta = \Omega_{01} \partial_t S$, we have $\beta_H = 2$ and $\beta_T = 1$.

The following geometric commutation formula, proved in [Bigorne 2021b, Lemma 2.8], will be fundamental for us.

Lemma 2.3. *Let G be a 2-form and $g : [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a function, both of class C^1 , such that*

$$\nabla^\mu G_{\mu\nu} = J(g)_\nu, \quad \nabla^{\mu*} G_{\mu\nu} = 0.$$

Let further $Z \in \mathbb{K} \setminus \{S\}$ be a Killing vector field and $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{S\}$ be its complete lift. Then,

$$\begin{aligned} \nabla^\mu \mathcal{L}_Z(G)_{\mu\nu} &= J(\widehat{Z}g)_\nu, \quad \nabla^{\mu*} \mathcal{L}_Z(G)_{\mu\nu} = 0, \\ \nabla^\mu \mathcal{L}_S(G)_{\mu\nu} &= J(Sg)_\nu + 3J(g)_\nu, \quad \nabla^{\mu*} \mathcal{L}_S(G)_{\mu\nu} = 0, \\ \widehat{Z}(v^\mu G_\mu^j \partial_{v^j} g) &= v^\mu \mathcal{L}_Z(G)_\mu^j \partial_{v^j} g + v^\mu G_\mu^j \partial_{v^j} \widehat{Z}g, \\ S(v^\mu G_\mu^j \partial_{v^j} g) &= v^\mu \mathcal{L}_S(G)_\mu^j \partial_{v^j} g - 2v^\mu G_\mu^j \partial_{v^j} g + v^\mu G_\mu^j \partial_{v^j} Sg. \end{aligned}$$

Iterating the above, we obtain that the structure of the Vlasov–Maxwell equations (6)–(7) is preserved by commutation.

Proposition 2.4. *Let (f, F) be a sufficiently regular solution to the Vlasov–Maxwell system. For any multi-index β , there exists $C_{\gamma, \kappa}^\beta, C_\xi^\beta \in \mathbb{Z}$ such that*

$$\begin{aligned} \mathbf{T}_F(\widehat{Z}^\beta f) &= \sum_{\substack{|\gamma| + |\kappa| \leq |\beta| \\ |\kappa| \leq |\beta| - 1}} C_{\gamma, \kappa}^\beta \widehat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu^j \partial_{v^j} \widehat{Z}^\kappa f, \\ \nabla^\mu \mathcal{L}_{Z^\beta}(F)_{\mu\nu} &= \sum_{|\xi| \leq |\beta|} C_\xi^\beta J(\widehat{Z}^\xi f), \quad \nabla^{\mu*} \mathcal{L}_{Z^\beta}(F)_{\mu\nu} = 0. \end{aligned}$$

Moreover, the multi-indices $|\gamma| + |\kappa| \leq |\beta|$ satisfy $\gamma_H + \kappa_H \leq \beta_H$ and the equality $\kappa_H = \beta_H$ implies $\gamma_T \geq 1$.

Proof. For the condition on the multi-indices $|\gamma| + |\kappa| \leq |\beta|$, note from Lemma 2.3 that $\gamma_H + \kappa_H \leq \beta_H$ and $\gamma_T + \kappa_T = \beta_T$. Hence, if $\kappa_H = \beta_H$, we necessarily have $\kappa_T < \beta_H$ since $|\kappa| < |\beta|$. This implies $\gamma_T \geq 1$. \square

2.5. Weights preserved along the linear flow. The set \mathbf{k}_1 of weight functions given by

$$z_{0i} := t \widehat{v}^i - x^i, \quad z_{jk} := x^j \widehat{v}^k - x^k \widehat{v}^j, \quad 1 \leq i \leq 3, \quad 1 \leq j < k \leq 3, \tag{10}$$

are conserved along any timelike straight line $t \mapsto (t, x + t\hat{v})$. They are obtained as $|v^0|^{-1}\eta(v, K)$, where K is a Killing vector field⁶ and they are then solutions to the relativistic transport equation, for all $z \in \mathbf{k}_1$, $T_0(z) = 0$. As a consequence, if $T_0(g) = 0$ then the same goes for zg , so that certain weighted norms of g are conserved. In our nonlinear setting these norms will grow logarithmically in time and will then provide useful decay properties on the Vlasov field. For convenience, we will rather work with

$$z := \left(1 + \sum_{z \in \mathbf{k}_1} z^2\right)^{\frac{1}{2}}, \quad T_0(z) = \hat{v}^\mu \partial_{x^\mu}(z) = 0. \tag{11}$$

In particular, as $z_{0i} \in \mathbf{k}_1$, one has

$$1 \leq z \quad \text{and} \quad \forall(t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad \langle x \rangle \leq z(t, x + t\hat{v}, v), \tag{12}$$

which will allow us to obtain space decay for $f(t, x + t\hat{v}, v)$, the particle density evaluated along the linear characteristics. Note also the following properties, which will be particularly useful for us in order to exploit the null structure of the system.

Lemma 2.5. *The four-momentum vector v has good null components, v^L and $\hat{\phi}$. More precisely,*

$$\forall(t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad 0 < \hat{v}^L \lesssim \frac{1 + |t - |x||}{1 + t + |x|} + \frac{z}{1 + t + |x|}, \quad |\hat{\phi}| \lesssim \frac{z}{1 + t + |x|}.$$

In certain circumstances, v^L will be the best component for exploiting decay in $t - r$. We will then use

$$|v^0|^{-2} + |\hat{\phi}|^2 \leq 4\hat{v}^L.$$

Proof. The first two inequalities are proved in [Bigorgne 2020a, Lemma 2.4]; using

$$4v^0 v^L \geq 4v^L v^L = |v^0|^2 - \left|\frac{x^i}{r} v_i\right|^2 = 1 + |v|^2 - |v \cdot \partial_r|^2 = 1 + |v \cdot e_\theta|^2 + |v \cdot e_\phi|^2 = 1 + |\hat{\phi}|^2, \tag{13}$$

the last inequality follows. □

Since the particles are massive and then travel at a speed strictly lower than 1, the speed of light, Vlasov fields enjoy much better decay properties along null rays than along timelike geodesics $t \mapsto x + t\hat{v}$. After a long time, many of the particles should be located in the interior of the light cone. We will capture this property with the following inequality.

Lemma 2.6. *By losing powers of v^0 and z , one can gain decay near the light cone $t = |x|$,*

$$\forall(t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad 1 \lesssim \frac{1 + |t - |x||}{1 + t + |x|} |v^0|^2 + \frac{|v^0|^2 z}{1 + t + |x|}.$$

Moreover, in the exterior of the light cone, for $|x| \geq t$, one has $1 \lesssim (1 + t + |x|)^{-1} |v^0|^2 z$.

Proof. For the first inequality, note that (13) gives $1 \leq 4|v^0|^2 \hat{v}^L$ and apply Lemma 2.5. For the second one, we refer to [Bigorgne 2020a, Remark 2.5]. □

Recall from [Bigorgne 2020a, Lemma 3.2] that z enjoys good commutation properties with the vector fields of $\hat{\mathbb{P}}_0$.

⁶On any smooth Lorentzian manifold (Y, g) , if γ is a timelike geodesic and K a Killing vector field, then $g(\dot{\gamma}, K) = \text{constant}$.

Lemma 2.7. *For any $a \in \mathbb{R}$ and $\widehat{Z} \in \widehat{\mathbb{P}}_0$, we have $|\widehat{Z}(z^a)| \lesssim |a|z^a$.*

Finally, motivated by the fact that any regular solution to the linear relativistic transport equation $T_0(h) = 0$ is constant along the timelike straight lines, $h(t, x + \widehat{v}t, v) = h(0, x, v)$, it will sometimes be useful to work with $g(t, x, v) := f(t, x + t\widehat{v}, v)$, in particular during the study of the asymptotic properties of $\int_v f dv$ and its derivatives. The following result suggests that g enjoys strong space decay and that its v derivatives behave better than the ones of the distribution function f .

Lemma 2.8. *Let $f : [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a sufficiently regular function and $g(t, x, v) := f(t, x + t\widehat{v}, v)$. Then the following properties hold:*

$$(x)^a |g|(t, x, v) \leq |z^a f|(t, x + t\widehat{v}, v), \quad v^0 |\nabla_v g|(t, x, v) \leq \sum_{\widehat{Z} \in \widehat{\mathbb{P}}_0} |z \widehat{Z} f|(t, x + t\widehat{v}, v).$$

Proof. The first property follows from $z^2 \geq 1 + |z_{01}|^2 + |z_{02}|^2 + |z_{03}|^2$ and $|z_{0i}|(t, x + t\widehat{v}, v) = |x^i|$. For the second one, we have, using the Einstein summation convention,

$$v^0 \partial_{v_j} g(t, x, v) = (v^0 \partial_{v_j} f)(t, x + t\widehat{v}, v) + t \partial_{x^j} f(t, x + t\widehat{v}, v) - t \widehat{v}_j \widehat{v}^i \partial_{x^i} f(t, x + t\widehat{v}, v).$$

Then by $v^0 \partial_{v_j} = \widehat{\Omega}_{0j} - t \partial_{x^j} - x^j \partial_t$ and

$$x^j \partial_t + t \widehat{v}^j \widehat{v}^i \partial_{x^i} = (x^j - t \widehat{v}^j) \partial_t + \widehat{v}^j t \partial_t + \widehat{v}^j (t \widehat{v}^i - x^i) \partial_{x^i} + \widehat{v}^j x^i \partial_{x^i} = -z_{0j} \partial_t + \widehat{v}^j S + \sum_{1 \leq i \leq 3} \widehat{v}^j z_{0i} \partial_{x^i}, \quad (14)$$

the result follows. □

2.6. Inverse function of the relativistic speed. In order to perform the change of variables $y = x - \widehat{v}t$ for integrals on the domain \mathbb{R}_v^3 , it will be useful to determine certain properties of the function $v \mapsto \widehat{v}$.

Lemma 2.9. *We define, on the domain $\{y \in \mathbb{R}^3 \mid |y| < 1\}$, the operator $\check{\cdot}$ as*

$$y \mapsto \check{y} = \frac{y}{\sqrt{1 - |y|^2}}, \quad \text{so that } \forall |y| < 1, \quad v \in \mathbb{R}_v^3, \quad \widehat{y} = y, \quad \check{\widehat{v}} = v.$$

Note also that $v^0 = (1 - |\widehat{v}|^2)^{-1/2}$. Moreover, for any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$, the Jacobian determinant of the transformation $v \mapsto x - \widehat{v}t$ is equal to $-t^3/|v^0|^5$.

Proof. The fact that $\check{\cdot}$ is the reciprocal function of $\widehat{\cdot}$ can be obtained by direct computations. Let V be the column vector such that its transpose is $V^T = (v^1/v^0, v^2/v^0, v^3/v^0)$. Then the Jacobian determinant of the transformation $v \mapsto x - \widehat{v}t$ is equal to

$$-\frac{t^3}{|v^0|^3} \det(\mathbf{I}_3 - VV^T) = -\frac{t^3}{|v^0|^3} \det\left(\text{diag}\left(1, 1, 1 - \frac{|v|^2}{1 + |v|^2}\right)\right) = -\frac{t^3}{|v^0|^5}. \quad \square$$

Let us also mention the inequality $2(1 - |\widehat{v}|) \geq (1 - |\widehat{v}|)(1 + |\widehat{v}|) = |v^0|^{-2}$, which will be used several times throughout this paper.

2.7. Complete version of the main result. We are now ready to give a full and detailed version of [Theorem 1.1](#). Recall the alternative geometric form (6)–(7) of the Vlasov–Maxwell equations (1)–(3).

Theorem 2.10. *Let $N \geq 3$ and (f_0, F_0) be an initial data set of class C^N for the Vlasov–Maxwell system. Consider further $\Lambda \geq \epsilon > 0$, two constants $(N_v, N_x) \in \mathbb{R}_+^2$ and assume that*

$$\sum_{|\gamma| \leq N+1} \sup_{x \in \mathbb{R}^3} \langle x \rangle^{2+|\gamma|} |\nabla_x^\gamma F_0|(x) \leq \Lambda, \quad \sum_{|\beta|+|\kappa| \leq N} \sup_{(x,v) \in \mathbb{R}^6} \langle v \rangle^{N_v+|\kappa|} \langle x \rangle^{N_x+|\beta|} |\partial_v^\kappa \partial_x^\beta f_0|(x, v) \leq \epsilon.$$

If $N_v \geq 15$ and $N_x > 7$, there exist $D > 0$ and $\epsilon_0 > 0$, depending only on (N, N_v, N_x) , such that, if $\bar{\epsilon} := \epsilon e^{D\Lambda} \leq \epsilon_0$, then the unique solution (f, F) to (1)–(3) arising from these data is global in time. Moreover:

- *The following pointwise estimates hold for the distribution function:*

$$\forall (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_v^3 \times \mathbb{R}_x^3, \forall |\beta| \leq N, \quad |v^0|^{N_v-3} |z^{N_x-2} \widehat{Z}^\beta f|(t, x, v) \lesssim \bar{\epsilon} \log^{3N_x+3N}(3+t),$$

$$\forall |\kappa| \leq N, \quad |v^0|^{N_v-3} |\partial_{t,x}^\kappa f|(t, x, v) \lesssim \bar{\epsilon}.$$

- *The electromagnetic field and its derivatives $\mathcal{L}_{Z^\gamma} F$, up to order $|\gamma| \leq N - 1$, decay as,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1}.$$

If $|\gamma| \leq N - 2$, the good null components enjoy stronger decay properties near the light cone,

$$\forall (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\alpha(\mathcal{L}_{Z^\gamma} F)|(t, x) + |\rho(\mathcal{L}_{Z^\gamma} F)|(t, x) + |\sigma(\mathcal{L}_{Z^\gamma} F)|(t, x) \lesssim \Lambda \frac{\log(3+t)}{(1+t+|x|)^2}.$$

Let us formulate two remarks.

- (1) More estimates, such as $\int_v f \, dv \lesssim t^{-3}$, are derived during the proof of [Theorem 2.10](#).
- (2) With our method, contrary to our previous work [[Bigorgne 2020a](#)], we cannot reach the optimal assumption $N_v = 3$. We list in [Remark 3.3](#) below the precise parts of the proof where the control of higher spatial and momentum moments of f are required.

We now state our scattering result. For this, recall from (8) the definition of the pure charge part \bar{F} of F .

Theorem 2.11. *Let $0 < \delta \leq 1$ and (f, F) be a smooth solution to the Vlasov–Maxwell system arising from initial data satisfying the assumptions of [Theorem 2.10](#). Suppose further that the initial electromagnetic field has the asymptotic expansion*

$$\sum_{|\gamma| \leq N+1} \sup_{|x| \geq 1} \langle x \rangle^{2+\delta+|\gamma|} |\nabla_{t,x}^\gamma (F - \bar{F})|(0, x) \leq \Lambda. \tag{15}$$

Then, with $n := 7(N_x + N)$, we have the following properties.

- *The spatial average of f converges to a function $Q_\infty \in L^1(\mathbb{R}_v^3) \cap L^\infty(\mathbb{R}_v^3)$ of class C^{N-1} ,*

$$\forall t \in \mathbb{R}_+, \quad \left\| |v^0|^5 \left(\int_{\mathbb{R}_x^3} f(t, x, v) \, dx - Q_\infty(v) \right) \right\|_{L_v^1 \cap L_v^\infty} \lesssim \bar{\epsilon} \frac{\log^n(3+t)}{1+t}.$$

- The four-current density $J(\widehat{Z}^\beta f)_\mu = \int_v (v_\mu/v^0)\widehat{Z}^\beta f \, dv$ has the following self-similar asymptotic profile. For any $|\beta| \leq N - 1$ and $0 \leq \mu \leq 3$,

$$\forall t \in \mathbb{R}_+^*, \quad \sup_{|x| < t} \left| t^3 \int_{\mathbb{R}_v^3} \frac{v^\mu}{v^0} \widehat{Z}^\beta f(t, x, v) \, dv - \frac{x^\mu}{t} (|v^0|^5 Q_\infty^\beta) \left(\frac{\tilde{x}}{t} \right) \right| \lesssim \bar{\epsilon} \frac{\log^n(3+t)}{t}, \quad x^0 = t,$$

where Q_∞^β can be computed in terms of $\partial_v^\kappa Q_\infty$, $|\kappa| \leq |\beta|$. Moreover, $J(\widehat{Z}^\beta f)$ decays much faster in the exterior of the light cone.

- The electromagnetic field and their derivatives up to order $|\gamma| \leq N - 1$ have a self-similar asymptotic profile $v \mapsto \mathcal{L}_{Z^\gamma}(F)^\infty(v)$,

$$\forall (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |t^2 \mathcal{L}_{Z^\gamma}(F)(t, x + \hat{v}t) - \mathcal{L}_{Z^\gamma}(F)^\infty(v)| \lesssim \Lambda \langle x \rangle^2 |v^0|^8 \frac{\log^n(3+t)}{(1+t)^\delta}.$$

F^∞ is of class C^{N-1} and the components of $\mathcal{L}_{Z^\gamma}(F)^\infty$ can be computed in terms of $\partial_v^\kappa F_{\mu\nu}^\infty$, $|\kappa| \leq |\gamma|$.

- We have modified scattering to a state $f_\infty \in L_{x,v}^1 \cap L_{x,v}^\infty$ of class C^{N-2} . For any $|\kappa| + |\beta| \leq N - 2$,

$$\forall t \geq 3, \quad \left\| |v^0|^{N_v-10+|\xi|} \langle x \rangle^{N_x-4-|\xi|} (\partial_v^\xi \partial_x^\kappa f(t, X_{\mathcal{C}}(t, x, v), v) - \partial_v^\xi \partial_x^\kappa f_\infty(x, v)) \right\|_{L_{x,v}^\infty} \lesssim \bar{\epsilon} \frac{\log^n(t)}{t^\delta},$$

where the corrections of the linear spatial characteristics are defined as

$$X_{\mathcal{C}}^j(t, x, v) := x^j + t\hat{v}^j - \frac{\log(t)}{v^0} \hat{v}^\mu (F_{\mu}^{\infty,j}(v) + \hat{v}^j F_{\mu 0}^\infty(v)), \quad 1 \leq j \leq 3. \tag{16}$$

- The modified complete lifts, of the Lorentz boosts $\widehat{\Omega}_{0k}$ and the rotations $\widehat{\Omega}_{jk}$, and the modified scaling,

$$\begin{aligned} \widehat{\Omega}_{\lambda k}^{\text{mod}} &:= \widehat{\Omega}_{\lambda k} - \frac{\log(t)}{v^0} \hat{v}^\mu (\mathcal{L}_{\Omega_{\lambda k}}(F)_\mu^{\infty,j}(v) + \hat{v}^j \mathcal{L}_{\Omega_{\lambda k}}(F)_{\mu 0}^\infty(v)) \partial_{x^j}, \quad 0 \leq \lambda < k \leq 3, \\ S^{\text{mod}} &:= S + \frac{\log(t)}{v^0} \hat{v}^\mu (F_\mu^{\infty,j}(v) + \hat{v}^j F_{\mu 0}^\infty(v)) \partial_{x^j}, \end{aligned}$$

satisfy the improved estimates $\|\widehat{\Omega}_{\lambda k}^{\text{mod}} f(t, \cdot, \cdot)\|_{L_{x,v}^\infty} \lesssim \bar{\epsilon}$ and $\|S^{\text{mod}} f(t, \cdot, \cdot)\|_{L_{x,v}^\infty} \lesssim \bar{\epsilon}$.

- For any $|\gamma| \leq N - 3$, there exists a scattering state $\underline{\alpha}_\gamma^{\mathcal{I}^+}(u, \omega)$ on \mathcal{I}^+ such that,

$$\forall \underline{u} \geq 3, \quad \sup_{|u| \leq \underline{u}, \omega \in \mathbb{S}^2} |r \underline{\alpha}(\mathcal{L}_{Z^\gamma} F)(u, \underline{u}, \omega) - \underline{\alpha}_\gamma^{\mathcal{I}^+}(u, \omega)| \lesssim \Lambda \frac{\log(\underline{u})}{\underline{u}}.$$

Moreover, $\underline{\alpha}^{\mathcal{I}^+}$ is of class C^{N-3} and $\underline{\alpha}_\gamma^{\mathcal{I}^+}$ can be expressed in terms of the derivatives of $\underline{\alpha}^{\mathcal{I}^+}$.

- The conserved energy of the system can be related to the ones of the scattering states. For all $t \in \mathbb{R}_+$,

$$\int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f(t, x, v) \, dv \, dx + \frac{1}{2} \int_{\mathbb{R}_x^3} |F|^2(t, x) \, dx = \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f_\infty(x, v) \, dv \, dx + \frac{1}{4} \int_{\mathbb{R}^u} \int_{\mathbb{S}_\omega^2} |\underline{\alpha}^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du.$$

- If $N \geq 10$, there exists a solution F^{vac} of class C^{N-5} to the vacuum Maxwell equations (19) such that, for any $\frac{1}{2} \leq q < 1$ and $|\gamma| \leq N - 10$,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F) - \mathcal{L}_{Z^\gamma}(F)^{\text{vac}}|(t, x) \leq \Lambda C_q (1+t+|x|)^{-1-q}, \quad C_q > 0.$$

Remark 2.12. As suggested by the scattering result, we could improve the logarithmic powers in the $L^\infty_{x,v}$ estimates for f stated in [Theorem 2.10](#). We could then prove that [Theorem 2.11](#) holds for $n = 3N_x + 3N$. However, such a tiny improvement would require a relatively long and technical proof.

Remark 2.13. We emphasize two main differences with previous works on Vlasov systems in dimension 3 based on vector field methods [[Fajman et al. 2021](#); [Smulevici 2016](#); [Bigogne 2020a](#); [Duan 2022](#)].

- (1) The logarithmic correction of the linear commutators $\widehat{\Omega}_{\lambda,v}$ and S can be geometrically interpreted in terms of the asymptotic dynamic of the Lorentz force $\hat{v}^\mu F_{\mu k}$ and its derivatives (see also [Remark 6.31](#)).
- (2) Our approach does not require modifying the linear commutators in order to prove the global existence of the solutions, so that we avoid many technical difficulties. In these previous works, the analysis of the Vlasov field relied on propagating $L^1_{x,v}$ bounds. The source term of the wave equations (or the Poisson equation) were estimated through weighted Sobolev embeddings as $t^3 |Z^\beta \int_v f \, dv| \leq t^3 \int_v |\widehat{Z}^\beta f| \, dv \lesssim \mathbb{E}(t)$, where $\mathbb{E}(t)$ is a certain $L^1_{x,v}$ norm. However, we know from [Theorems 2.10–2.11](#) that, in general, $\|\widehat{Z} f\|_{L^1_{x,v}} \gtrsim \log(t)$ if $\widehat{Z} \neq \partial_{t,x}$. As a consequence, the optimal decay t^{-3} cannot be obtained in such a way without modifying the linear commutators.

Remark 2.14. The profile F^∞ of F can be explicitly expressed in terms of the limit of the spatial average Q_∞ (see [Remark 6.17](#) and [Appendix C.1](#)). Moreover, the Maxwell field admits the decomposition $F = F^T + F^2$, where

$$\lim_{t \rightarrow +\infty} t^2 F(t, x + t\hat{v}) = \lim_{t \rightarrow +\infty} t^2 F^T(t, x + t\hat{v}) = F^\infty(v), \quad \lim_{\underline{u} \rightarrow +\infty} r F^T(u, \underline{u}, \omega) = 0.$$

In other words, the part of the electromagnetic field which gives rise to F^∞ (respectively $\underline{\alpha}^{\mathcal{I}^+}$) has no impact on $\underline{\alpha}^{\mathcal{I}^+}$ (respectively F^∞).

2.8. Key ingredients of the proof. For the global existence result, our strategy relies on vector field methods and a continuity argument. The proof then essentially consists in improving bootstrap assumptions, which are pointwise decay estimates on the solutions and their derivatives. The scattering statements are then obtained by refining the analysis carried out during of the proof of [Theorem 2.10](#) and by investigating further the asymptotic behavior of the electromagnetic field.

2.8.1. The large Maxwell field. The assumptions of [Theorems 2.10–2.11](#) imply that, initially, the distribution function f is at most of size $\epsilon \ll 1$ and the electromagnetic field F is at most of size Λ . The goal of our bootstrap argument is to prove that these properties are preserved over time. Our proof allows for Λ to be large for the following reasons.

- Since the Maxwell equations are *linear*, we can expect $F(t, \cdot)$ and its derivatives to be at most of size $\Lambda + C\epsilon \sim \Lambda$, provided that ϵ is small enough. Here, the constant C possibly depends on Λ . Indeed, the data are bounded by Λ and we expect the source term $J(f)$ to remains of size ϵ .
- In contrast, the Vlasov equation is nonlinear and we can expect, at first glance, to bound $\|\partial_{t,x}^k f(t, \cdot)\|_{L^\infty_{x,v}}$ by $\epsilon + D\Lambda\epsilon = C(\Lambda)\epsilon$.

In fact, since our argument will rely on Grönwall’s inequality, $C(\Lambda)$ will rather be of the form $e^{D\Lambda}$. The difficulty, if Λ is large, is related to the logarithmic growth of quantities such as $\|\widehat{\Omega}_{01}f\|_{L_{x,v}^\infty}$. More precisely, certain error terms are at the threshold of time-integrability. Consequently a naive application of Grönwall’s inequality would lead to $\|\widehat{\Omega}_{01}f\|_{L_{x,v}^\infty} \lesssim \epsilon(1+t)^{D\Lambda}$. We discuss how to circumvent this obstacle in the next section.

2.8.2. Estimates for the Vlasov field. In order to control sufficiently well the electromagnetic field and close our estimates, we would like to recover the linear decay for $|\int_v \widehat{Z}^\beta f(t, x, v) dv| \lesssim t^{-3}$, with $|\beta| \leq N - 1$, and similar quantities. This is done as follows:

- The main step consists in proving that $|v^0|^{N_v} z^{N_x} \widehat{Z}^\beta f$ grows slowly, and in fact logarithmically, in time.
- Then, by performing the standard change of variables $y = x - t\hat{v}$, we are able to reduce the problem to proving a uniform bound for the spatial averages $|v^0|^5 \int_y \widehat{Z}^\beta f(t, y, v) dy$. This turns out to be a consequence of the first step as well but our argument requires a loss of regularity, which is why we do not attain the optimal decay t^{-3} for the top-order derivatives $|\beta| = N$.

Let us illustrate certain difficulties of the first step, which relies on Duhamel’s formula, by considering the first-order derivatives. If $Z \in \mathbb{K} \setminus \{S\}$ is a Killing vector field, then

$$|\mathbf{T}_F(\widehat{Z}f)| = |\hat{v}^\mu \mathcal{L}_Z(F)_\mu^j \partial_{v^j} f| \lesssim \sum_{1 \leq j \leq 3} \frac{t + |x|}{v^0} |\hat{v}^\mu \mathcal{L}_Z(F)_\mu^j| |\partial_{t,x} f| + \text{better terms.} \tag{17}$$

Since $\mathcal{L}_Z(F)$ is supposed to decay as⁷ $|\mathcal{L}_Z(F)| \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1}$, there are two problems.

- (1) The decay rate degenerates near the light cone $t = |x|$.
- (2) Even far from the light cone, $|\mathbf{T}_F(\widehat{Z}f)| \sim \Lambda t^{-1} |\partial_{t,x} f|$ is not integrable in time, preventing us from proving that $\|\widehat{Z}f\|_{L_{x,v}^\infty}$ grows slowly by a direct application of Grönwall’s inequality if Λ is large.

We deal with the first issue by taking advantage of the null structure of the Lorentz force, which, roughly speaking, allows us to transform decay in $t - r$ into decay in $t + r$. More precisely, $\hat{v}^\mu \mathcal{L}_Z(F)_\mu^j$ can be decomposed as the sum of terms containing either a good null component α , ρ or σ of $\mathcal{L}_Z(F)$ or one of the good null components of \hat{v} . The first group enjoys improved decay estimates near the light cone, whereas the latter allows us to exploit the decay in $t - r$. We refer to Lemmas 4.1 and 4.4 for more details.

We circumvent the second problem by identifying hierarchies in the commuted equations. More precisely, if $Z = \partial_{x^\mu}$ is a translation, one can use that $|\mathcal{L}_{\partial_{x^\mu}}(F)| \lesssim t^{-1}(1+|t-|x||)^{-2}$ in order to prove that $\mathbf{T}_F(\partial_{x^\mu} f)$ is in fact time-integrable. Then, one can observe that the system of the commuted Vlasov equations (17) is in some sense triangular and expect $\|\widehat{Z}f\|_{L_{x,v}^\infty}$ to grow at most logarithmically. A toy model for the system of the first-order commuted equations, once the null structure is well understood, is then

$$\mathbf{T}_F(g) = \Lambda(1+t)^{-2}g + \Lambda(1+t)^{-3}h, \quad \mathbf{T}_F(h) = \Lambda(1+t)^{-1}g + \Lambda(1+t)^{-2}h, \quad g \geq 0, h \geq 0,$$

where g is supposed to capture the behavior of $|\partial_{x^\mu} f|$, $0 \leq \mu \leq 3$, and h that of $|\widehat{Z}f|$, with \widehat{Z} a homogeneous vector field such as $\widehat{\Omega}_{01}$. The source terms having h as a factor represent the strongly

⁷This pointwise decay estimate is consistent with the expected behavior of the source term of the Maxwell equations.

decaying error terms in (17). Using the Duhamel formula and applying Grönwall’s inequality, we have, for $\mathbb{E}(t) := \|g(t, \cdot, \cdot)\|_{L_{x,v}^\infty} + \|h(t, \cdot, \cdot)\|_{L_{x,v}^\infty}$,

$$\mathbb{E}(t) \leq \mathbb{E}(0) + \int_{s=0}^t \frac{\Lambda}{1+s} \mathbb{E}(s) \, ds, \quad \mathbb{E}(t) \leq \mathbb{E}(0)(1+t)^\Lambda.$$

As mentioned earlier, without any smallness assumption on Λ , this estimate is not good enough to derive a satisfying decay estimate for $\int_v f \, dv$. The idea then is to exploit that

$$\mathbf{T}_F(\log^{-1}(3+t)) \leq 0, \quad \mathbf{T}_F(h \log^{-2}(3+t)) \leq \Lambda(1+t)^{-1} \log^{-2}(3+t)g + \Lambda(1+t)^{-2}h \log^{-2}(3+t).$$

By considering the hierarchized norm $\bar{\mathbb{E}}(t) := \|g(t, \cdot, \cdot)\|_{L_{x,v}^\infty} + \|h(t, \cdot, \cdot)\|_{L_{x,v}^\infty} \log^{-2}(3+t)$, we finally get

$$\bar{\mathbb{E}}(t) \leq \bar{\mathbb{E}}(0) + \int_{s=0}^t \frac{2\Lambda}{(1+s) \log^2(3+s)} \bar{\mathbb{E}}(s) \, ds, \quad \bar{\mathbb{E}}(t) \leq \bar{\mathbb{E}}(0)e^{2\Lambda}.$$

More generally, the hierarchies are determined by the number of homogeneous vector fields β_H composing \widehat{Z}^β and the exponent of the weight z .

A new difficulty arises for the higher-order derivatives since we do not have improved estimates at our disposal on the good null components of $\mathcal{L}_{Z^\gamma}(F)$ for $|\gamma| \geq N - 1$. This time, we transform decay in $t - r$ into decay in $t + r$ by losing powers of $|v^0|^2 z$ through Lemma 2.6. For this, it is important to observe that, in the error terms, such a $\mathcal{L}_{Z^\gamma}(F)$ is always multiplied by a low-order derivative of f . We can then close the estimates by propagating weaker $L_{x,v}^\infty$ norms of $\widehat{Z}^\beta f$ when $|\beta| \geq N - 1$.

Remark 2.15. Let us make some comparisons between the decay properties of the electromagnetic F and the ones of the electric field E associated to a solution to the Vlasov–Poisson system arising from small data.

- As $\|E(t, \cdot)\|_{L_x^\infty} \lesssim t^{-2}$ and $|F|(t, x) \lesssim t^{-1}(1 + |t - |x||)^{-1}$, the electromagnetic field has a much weaker decay rate near the light cone $t = r$ than E .
- The difference is even more marked for their derivatives since $|\partial_{t,x}^\kappa E|(t, x) \lesssim t^{-2-|\kappa|}$, whereas we merely have $|\mathcal{L}_{\partial_{t,x}^\kappa} F|(t, x) \lesssim t^{-1}(1 + |t - |x||)^{-1-|\kappa|}$. Thus, in order to exploit the extra decay provided by these derivatives of F , one has to take advantage of the null structure of the system or lose powers of $|v^0|^2 z$.

2.8.3. Estimates for the electromagnetic field. We control the Cartesian components of $\mathcal{L}_{Z^\gamma}(F)$ using the representation formula for the wave equation since, for instance, $\square F_{01} = -\int_v \partial_{x^1} f + \hat{v}^1 \partial_t f \, dv$. However, two difficulties arise for the higher-order derivatives:

- (1) There is a loss of regularity. We need to control $\int_v \hat{v}^\mu \partial_{t,x} \widehat{Z}^\gamma f \, dv$ in order to estimate $\mathcal{L}_{Z^\gamma}(F)$.
- (2) With our method, we do not have the optimal decay rate for $\int_v \widehat{Z}^\gamma f \, dv$, $|\gamma| = N$. Moreover, any logarithmic loss would prevent us from closing our estimates.

We treat the first problem by using the Glassey–Strauss decomposition [1986] of the electromagnetic field, presented in detail in Section 5.1. The idea is to express the derivatives ∂_{x^μ} in terms of derivatives tangential to backward light cones and $\mathbf{T}_0 = \partial_t + \hat{v} \cdot \nabla_x$, which is transverse to light cones. Exploiting then the Vlasov equation $\mathbf{T}_F(f) = 0$, we can perform integration by parts and save one derivative.

We deal with the second issue by estimating $\nabla_{t,x}\mathcal{L}_{Z^\xi}(F)$, for $|\xi| = N - 1$, by the Glassey–Strauss decomposition of the derivatives of the electromagnetic field. Roughly speaking, it allows us to control the inhomogeneous part of $\nabla_{t,x}\mathcal{L}_{Z^\xi}(F)$ by $\int_v |v^0|^3 |\widehat{Z}^\beta f| dv$, where $|\beta| \leq N - 1$ (see [Proposition 5.7](#) and [Corollary 5.8](#) for more details). However, with this process, we get a bad control of the other top-order derivatives near the light cone,

$$|\mathcal{L}_{ZZ^\xi}(F)|(t, x) \lesssim (1+t+|x|)|\nabla_{t,x}\mathcal{L}_{Z^\xi}F|(t, x) + |\mathcal{L}_{Z^\xi}F|(t, x) \lesssim (1+|t-r|)^{-2} \log(3+|t-r|), \quad |\xi| = N - 1.$$

This forces us to lose a power more of $|v^0|^2 \mathbf{z}$ for the estimates of the top-order derivatives of the Vlasov field f .

Once we proved that the solutions are global in time, we use null properties of the Maxwell equations (7) to derive the existence of a scattering state for F and its derivatives. We then address the problem of finding a solution F^{vac} to the vacuum Maxwell equations which approaches F by constructing a scattering map for these equations. For this, we make crucial use of the corresponding result for the homogeneous wave equation [[Lindblad and Schlue 2023](#)]. This is carried out in [Section 7](#).

2.8.4. Modified scattering result. In the context of the Vlasov–Poisson system, except for the trivial solution, the distribution function does not converge along the linear characteristics [[Choi and Ha 2011](#)]. We then do not expect $f(t, x + t\hat{v}, v)$ to converge and the reason is related to the long-range effect of the Lorentz force (recall [Remark 1.3](#)). More precisely, isolating the leading-order term of the source term of the Maxwell equations,

$$\sup_{|x|<t} \left| t^3 \int_{\mathbb{R}_v^3} \frac{v^\mu}{v^0} f(t, x, v) dv - \frac{x^\mu}{t} (|v^0|^5 Q_\infty) \left(\frac{\check{x}}{t} \right) \right| = O(t^{-\frac{\delta}{2}}), \quad Q_\infty(v) := \lim_{t \rightarrow +\infty} \int_{\mathbb{R}_v^3} f(t, x, v) dv,$$

where $x^0 = t$, we are able to prove $t^2 F(t, x + t\hat{v}) = F^\infty(v) + O(t^{-\delta/2})$. Consequently, the slow decay of the electromagnetic field along timelike trajectories implies that the right-hand side of

$$\partial_t(f(t, x + t\hat{v}, v)) = \frac{t}{v^0} \hat{v}^\mu (F_\mu^j(t, x + t\hat{v}) + \hat{v}^j F_{\mu 0}(t, x + t\hat{v})) \partial_{x^j} f(t, x + t\hat{v}, v) + O(t^{-\frac{\delta}{2}})$$

should not be time-integrable, preventing $f(t, x + t\hat{v}, v)$ from converging. Instead, by considering the logarithmic corrections $X_\mathcal{E}$, given in (16), of the timelike straight lines, one can compensate for the worst term in the right-hand side of the previous identity and prove the modified scattering statement $f(t, X_\mathcal{E}, v) \rightarrow f_\infty(x, v)$.

Although the regularity of f_∞ according to x can be obtained in a similar fashion, the regularity in v requires a more thorough analysis. In fact, $v^0 \partial_{v^i}(f(t, X_\mathcal{E}, v))$ can be expressed as terms such as $\widehat{\Omega}_{0i} f(t, X_\mathcal{E}, v)$ which, contrary to $\partial_{t,x} f(t, X_\mathcal{E}, v)$, does not converge. The reason is related to the weak decay of the error term $[T_F, \widehat{\Omega}_{0i}] \sim t^{-1}$. As for the characteristics, the idea consists in considering a logarithmic correction of $\widehat{\Omega}_{0i}$, introduced and studied in [Section 6.4](#), which has improved commutation properties with the Vlasov operator T_F . As stated in [Theorem 2.11](#), these corrections are given in terms of first-order derivatives of the effective electromagnetic field $F^\infty(v)$.

2.9. Null properties of electromagnetic fields. We recall here the classical results which will be used throughout this paper in order to study solutions to the Maxwell equations

$$\nabla^\mu F_{\mu\nu} = J_\nu, \quad \nabla^{\mu*} F_{\mu\nu} = 0, \quad (18)$$

where the source term $J = J_\mu dx^\mu$ is a sufficiently regular 1-form. In particular, solutions to the vacuum Maxwell equations will satisfy

$$\nabla^\mu F_{\mu\nu} = 0, \quad \nabla^{\mu*} F_{\mu\nu} = 0. \quad (19)$$

We point out that some of the estimates presented here could be refined in a general setting. For the purpose of performing energy estimates during the construction of the scattering map for (19), we recall the electromagnetic stress-energy tensor.

Definition 2.16. Let G be a 2-form of class C^1 such that $\nabla^\mu G_{\mu\nu} = J_\nu$ and $\nabla^{\mu*} G_{\mu\nu} = 0$. The energy-momentum tensor $\mathbb{T}[G]_{\mu\nu}$ is defined as

$$\mathbb{T}[G]_{\mu\nu} := G_{\mu\beta} G_\nu{}^\beta - \frac{1}{4} \eta_{\mu\nu} G_{\xi\lambda} G^{\xi\lambda}.$$

Moreover, we have

$$\nabla^\mu T[G]_{\mu\nu} = G_{\nu\lambda} J^\lambda, \quad T[G]_{LL} = |\alpha(G)|^2, \quad T[G]_{\underline{L}\underline{L}} = |\underline{\alpha}(G)|^2, \quad T[G]_{\underline{L}L} = |\rho(G)|^2 + |\sigma(G)|^2.$$

We now present inequalities relying on the relations

$$(t-r)\underline{L} = S - \frac{x^i}{r} \Omega_{0i}, \quad (t+r)L = S + \frac{x^i}{r} \Omega_{0i}, \quad re_\theta = -\cos(\varphi)\Omega_{13} - \sin(\varphi)\Omega_{23}, \quad re_\varphi = \Omega_{12}. \quad (20)$$

Lemma 2.17. Let G be a sufficiently regular solution to the Maxwell equations (18) with a smooth source term J . Then,

$$\forall |x| \geq \frac{1}{2}(1+t), \quad (|\nabla_{\underline{L}}\alpha(G)| + |\nabla_{\underline{L}}\rho(G)| + |\nabla_{\underline{L}}\sigma(G)|)(t, x) \lesssim |J|(t, x) + \sum_{|\gamma| \leq 1} \frac{|\mathcal{L}_{Z^\gamma}(G)|(t, x)}{1+t+|x|}$$

and,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\nabla_L(r\underline{\alpha}(G))|(t, x) \lesssim r|J|(t, x) + \sum_{|\gamma| \leq 1} |\rho(\mathcal{L}_{Z^\gamma}G)|(t, x) + |\sigma(\mathcal{L}_{Z^\gamma}G)|(t, x).$$

Remark 2.18. Compared to $Z \in \mathbb{K}$, $Z \neq \partial_{x^\mu}$, the derivatives tangential to the light cone (L, e_θ, e_φ) provide an extra decay in $t+r$, whereas \underline{L} merely provides an additional decay in $t-r$. The second estimate then reflects that α , $\underline{\alpha}$, ρ and σ are the good null components. The last inequality provides an improved control of $\nabla_L(r\underline{\alpha})$ near the light cone and will be useful in order to prove the existence of scattering states.

Proof. Let us denote by ∇^\flat the intrinsic covariant differentiation on the spheres and by ζ any of the null components α , $\underline{\alpha}$, ρ or σ . Then, according to [Bigorgne 2021b, Lemma D.2], we have, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$(1+t+|x|)|\nabla_L\zeta(G)|(t, x) + (1+|x|)|\nabla^\flat\zeta(G)|(t, x) + (1+|t-|x||)|\nabla_{\underline{L}}\zeta(G)|(t, x) \lesssim \sum_{|\gamma| \leq 1} |\zeta(\mathcal{L}_{Z^\gamma}G)|(t, x).$$

We now express the Maxwell equations in null coordinates. According to [Christodoulou and Klainerman 1990, equations $(M''_1)-(M''_6)$], we have, for any $A \in \{\theta, \varphi\}$,

$$\begin{aligned} \nabla_{\underline{L}}\rho(G) - \frac{2}{r}\rho(G) - \not{\nabla}^{e_B}\underline{\alpha}(G)_{e_B} &= J_{\underline{L}}, & \nabla_{\underline{L}}\underline{\alpha}(G)_{e_A} - \frac{\alpha(G)_{e_A}}{r} + \not{\nabla}_{e_A}\rho(G) - \varepsilon^{AB}\not{\nabla}_{e_B}\sigma(G) &= J_{e_A}, \\ \nabla_{\underline{L}}\sigma(G) - \frac{2}{r}\sigma(G) + \varepsilon^{AB}\not{\nabla}_{e_A}\underline{\alpha}(G)_{e_B} &= 0, & \nabla_{\underline{L}}\underline{\alpha}(G)_{e_A} + \frac{\alpha(G)_{e_A}}{r} - \not{\nabla}_{e_A}\rho(G) - \varepsilon^{AB}\not{\nabla}_{e_B}\sigma(G) &= J_{e_A}. \end{aligned}$$

This allows us to deduce the first estimate. For the last one, use the same arguments and remark further that $\nabla_{\underline{L}}e_A = 0$ implies

$$|\nabla_{\underline{L}}(r\underline{\alpha})| \lesssim \sum_{B \in \{\theta, \varphi\}} |\nabla_{\underline{L}}(r\underline{\alpha})_{e_B}| = \sum_{B \in \{\theta, \varphi\}} |\nabla_{\underline{L}}(r\underline{\alpha}_{e_B})| = \sum_{B \in \{\theta, \varphi\}} |r\nabla_{\underline{L}}\underline{\alpha}(G)_{e_B} + \underline{\alpha}(G)_{e_B}|. \quad \square$$

In the same spirit, we have the following identity which is proved in [Bigorgne 2020a, Proposition 3.7, equation (18)].

Lemma 2.19. *For any sufficiently regular 2-form G and any null component $\zeta \in \{\alpha, \underline{\alpha}, \rho, \sigma\}$,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\zeta(\nabla_{t,x}G)|(t, x) \lesssim \sum_{|\gamma| \leq 1} \frac{|\zeta(\mathcal{L}_{Z^\gamma}G)|(t, x)}{1 + |t - |x||} + \frac{|\mathcal{L}_{Z^\gamma}(G)|(t, x)}{1 + t + |x|}.$$

We now illustrate how the previous lemmas can be used in order to obtain improved estimates for the good null components of the electromagnetic field.

Corollary 2.20. *Consider a 2-form G of class C^1 , a solution to the Maxwell equations (18) with a continuous source term J . Assume that there exist two constants $C[G] > 0$ and $q > 0$ such that,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad (1+t+|x|)|J|(t, x) + \sum_{|\gamma| \leq 1} |\mathcal{L}_{Z^\gamma}(G)|(t, x) \leq \frac{C[G]}{(1+t+|x|)(1+|t-|x||)^q}. \quad (21)$$

Then, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$(|\alpha(G)| + |\rho(G)| + |\sigma(G)|)(t, x) \lesssim C[G] \begin{cases} (1+t+|x|)^{-1-q} & \text{if } 0 < q < 1, \\ \log(3+t)(1+t+|x|)^{-2} & \text{if } q = 1, \\ (1+t+|x|)^{-2}(1+|t-|x||)^{-q+1} & \text{if } q > 1. \end{cases}$$

Moreover, if G is merely defined on $[0, T[\times \mathbb{R}^3$, $T > 0$, we have the weaker estimate, for the case $q > 1$,

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad (|\alpha(G)| + |\rho(G)| + |\sigma(G)|)(t, x) \lesssim C[G](1+t+|x|)^{-2} \quad \text{if } q > 1.$$

Proof. Note first that the assumptions give $|G|(t, x) \lesssim (1+t+|x|)^{-1-q}$ if $1+t \geq 2|x|$ or $|x| \geq 2(1+t)$. We then fix $(t, r\omega) \in \mathbb{R}_+ \times \mathbb{R}^3$ such that $1+t \leq 2r \leq 4(1+t)$, $\omega \in \mathbb{S}^2$, and we denote by ζ any of the null components α, ρ or σ . Consider further

$$\phi(u, \underline{u}) := \zeta(G) \left(\frac{u+u}{2}, \frac{u-u}{2} \omega \right).$$

By Lemma 2.17 and (21), we have

$$|\nabla_{\partial_u} \phi|(u, \underline{u}) = \frac{1}{2} |\nabla_{\underline{L}} \zeta(G)| \left(\frac{u+u}{2}, \frac{u-u}{2} \omega \right) \lesssim \frac{C[G]}{(1+\underline{u})^2(1+|u|)^q}.$$

Now, note that, for $t-r \leq 0$,

$$\begin{aligned} |\zeta(G)|(t, r\omega) &= |\phi|(t-r, t+r) = |\phi|(-t-r, t+r) + \int_{u=-t-r}^{-|t-r|} \nabla_{\partial_u} \phi(u, t+r) \, du \\ &\lesssim |\zeta(G)|(0, t\omega+r\omega) + \frac{C[G]}{(1+t+r)^2} \int_{u=-t-r}^{-|t-r|} \frac{du}{(1+|u|)^q}. \end{aligned}$$

Similarly, if $t-r \geq 0$, we obtain by integrating between $u = t-r$ and $t+r$,

$$|\zeta(G)|(t, r\omega) \lesssim |\zeta(G)|(t+r, 0) + \frac{C[G]}{(1+t+r)^2} \int_{u=|t-r|}^{t+r} \frac{du}{(1+|u|)^q}.$$

By (21),

$$|\zeta(G)|(t+r, 0) + |\zeta(G)|(0, t\omega+r\omega) \lesssim C[G](1+t+r)^{-1-q}$$

and the first part of the result then follows from the computations of the integrals in the previous two estimates. For the case $q = 1$, note that $\log(1+t+r) \leq 3 \log(3+t)$ since $r \leq 2+2t$.

If G is merely defined on $[0, T] \times \mathbb{R}^3$ and $t < T$, then we cannot apply the previous computations in the case $t \geq r$. Instead, we integrate between $u = 0$ and $t-r$ in order to get

$$|\zeta(G)|(t, r\omega) \lesssim |\zeta(G)| \left(\frac{t+r}{2}, \frac{t+r}{2} \omega \right) + \frac{C[G]}{(1+t+r)^2} \int_{u=0}^{|t-r|} \frac{du}{(1+|u|)^q}.$$

It remains to bound $|\zeta(G)|((t+r)/2, (t+r)\omega/2)$ by the estimate obtained in the region $t \leq r$ and to compute the integral in the three different cases. □

Finally, we prove pointwise decay estimates for a solution to the homogeneous wave equation. Since the Cartesian components $F_{\mu\nu}$ of a solution F to the vacuum Maxwell equations satisfy $\square F_{\mu\nu} = 0$, the next result will also allow us to estimate such electromagnetic fields.

Proposition 2.21. *Let ϕ be a C^2 solution to the free wave equation $\square \phi = 0$ such that*

$$\mathcal{E}^q[\phi] := \sup_{x \in \mathbb{R}^3} \langle x \rangle^q |\phi|(0, x) + \sup_{x \in \mathbb{R}^3} \langle x \rangle^{q+1} |\partial_{t,x} \phi|(0, x) < +\infty, \quad q \geq 2.$$

Then, there holds,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\phi|(t, x) \lesssim \frac{\mathcal{E}^q[\phi]}{(1+t+|x|)(1+|t-|x||)^{q-1}}.$$

Proof. By Kirchhoff's formula we have

$$\phi(t, x) = \frac{1}{4\pi t^2} \int_{|y-x|=t} \phi(0, y) \, dy + \frac{1}{4\pi t} \int_{|y-x|=t} \frac{y-x}{|y-x|} \cdot \nabla_y \phi(0, y) + \partial_t \phi(0, y) \, dy.$$

We obtain the result by applying⁸ [Wei and Yang 2021, Lemma 4.1], which gives that, for any $h \in C(\mathbb{R}^3)$ such that $|h|(x) \leq K_0(1 + |x|)^{-p}$,

$$\int_{|y-x|=t} |h|(y) \, dy \leq \begin{cases} 8\pi K_0 t^2 (1+t+|x|)^{-1} (1+|t-|x||)^{-p+1} & \text{if } 2 \leq p < 3, \\ 4\pi K_0 t (1+t+|x|)^{-1} (1+|t-|x||)^{-p+2} & \text{if } p \geq 3, \end{cases} \tag{22}$$

completing the proof. □

3. Strategy of the proof and the bootstrap assumptions

Let $N \geq 3$, $N_v \geq 15$, $N_x > 7$ and consider an initial data set (f_0, F_0) satisfying the hypotheses of [Theorem 2.10](#). By a standard local well-posedness argument, there exists a unique maximal solution (f, F) to the Vlasov–Maxwell system arising from these data. Let $T_{\max} \in \mathbb{R}_+^* \cup \{+\infty\}$ such that the solution is defined on $[0, T_{\max}[$. By continuity, there exists a largest time $0 < T \leq T_{\max}$ and a constant $C_{\text{boot}} > 0$, independent of ϵ , such that the following bootstrap assumptions hold. For all $(t, x) \in [0, T[\times \mathbb{R}^3$,

$$\forall |\gamma| \leq N - 1, \quad |\mathcal{L}_{Z^\gamma}(F)|(t, x) \leq \frac{C_{\text{boot}}\Lambda}{(1+t+|x|)(1+|t-|x||)}, \tag{BA1}$$

$$\forall |\gamma| = N - 1, \quad |\nabla_{t,x}\mathcal{L}_{Z^\gamma}(F)|(t, x) \leq \frac{C_{\text{boot}}\Lambda \log(3+|t-|x||)}{(1+t+|x|)(1+|t-|x||)^2}, \tag{BA2}$$

$$\forall |\beta| \leq N - 2, \quad \left| \int_{\mathbb{R}^3} \frac{v^\mu}{v^0} \widehat{Z}^\beta f(t, x, v) \, dv \right| \leq \frac{C_{\text{boot}}\Lambda}{(1+t+|x|)^3}, \quad 0 \leq \mu \leq 3. \tag{BA3}$$

The goal consists in improving, for C_{boot} chosen large enough and if ϵ is small enough, [\(BA1\)–\(BA3\)](#). We stress that [\(BA3\)](#) will only be used for the proof of [Proposition 3.1](#), where we improve the pointwise decay estimates for the good null components of the electromagnetic field.

3.1. Immediate consequences of the bootstrap assumptions. We start by improving, near the light cone, the estimates for the good null components of the electromagnetic field and its derivatives up to order $N - 2$.

Proposition 3.1. *For any $|\gamma| \leq N - 2$ and all $(t, x) \in [0, T[\times \mathbb{R}^3$, we have*

$$\begin{aligned} (|\alpha(\mathcal{L}_{Z^\gamma} F)| + |\rho(\mathcal{L}_{Z^\gamma} F)| + |\sigma(\mathcal{L}_{Z^\gamma} F)|)(t, x) &\lesssim \frac{\Lambda \log(3+t)}{(1+t+|x|)^2(1+|t-|x||)^{\gamma_T}}, \\ |\underline{\alpha}(\mathcal{L}_{Z^\gamma} F)|(t, x) &\lesssim \frac{\Lambda}{(1+t+|x|)(1+|t-|x||)^{1+\gamma_T}}, \end{aligned}$$

where we recall that γ_T is number of translations ∂_{x^μ} composing Z^γ .

Proof. Consider $|\gamma| \leq N - 2$ and recall from [Proposition 2.4](#) that $\mathcal{L}_{Z^\gamma} F$ is solution to the Maxwell equations [\(18\)](#) with a source term which is a linear combination of $J(\widehat{Z}^\beta f)$, $|\beta| \leq N - 2$, which are bounded by the bootstrap assumption [\(BA3\)](#). Hence, by applying [Corollary 2.20](#) and using the bootstrap

⁸The case $2 < p < 3$, not considered by [Wei and Yang 2021], can be treated as the case $p = 2$ since $\int_b^a \lambda/(1+\lambda)^p \, d\lambda \leq (1+b)^{p-2} \int_b^a \lambda/(1+\lambda)^2 \, d\lambda$.

assumption (BA1), we get

$$\begin{aligned} (|\alpha(\mathcal{L}_{Z^\gamma} F)| + |\rho(\mathcal{L}_{Z^\gamma} F)| + |\sigma(\mathcal{L}_{Z^\gamma} F)|)(t, x) &\lesssim \Lambda \log(3+t)(1+t+|x|)^{-2}, \\ |\underline{\alpha}(\mathcal{L}_{Z^\gamma} F)|(t, x) &\lesssim |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1}. \end{aligned}$$

Now, note that, for any $0 \leq \mu \leq 3$ and $Z \in \mathbb{K}$, we have $[Z, \partial_{x^\mu}] = 0$ or $[Z, \partial_{x^\mu}] = \pm \partial_{x^\lambda}$ for a certain $0 \leq \lambda \leq 3$. As a consequence, and since $\mathcal{L}_{\partial_{x^\mu}} = \nabla_{\partial_{x^\mu}}$, there exists constants $D_{\kappa, \xi}^\gamma \in \mathbb{N}$ such that

$$\mathcal{L}_{Z^\gamma} F = \sum_{|\kappa|=\gamma_T} \sum_{|\xi| \leq |\gamma| - \gamma_T} D_{\kappa, \xi}^\gamma \mathcal{L}_{\partial_{t,x}^\kappa} \mathcal{L}_{Z^\xi} F = \sum_{|\kappa|=\gamma_T} \sum_{|\xi| \leq |\gamma| - \gamma_T} D_{\kappa, \xi}^\gamma \nabla_{t,x}^\kappa \mathcal{L}_{Z^\xi} F. \tag{23}$$

The result then follows from γ_T applications of Lemma 2.19. □

In contrast, we have very bad control of the top-order derivatives near the light cone.

Proposition 3.2. *For any $|\gamma| = N$, there holds,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda \frac{\log(3+|t-|x||)}{(1+|t-|x||)^{2+\gamma_T}}.$$

If $|\gamma| \leq N - 1$, we have the better estimate,

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1-\gamma_T}.$$

Proof. Let $|\gamma| = N$, $(t, x) \in [0, T[\times \mathbb{R}^3$ and note that $|\mathcal{L}_Z G| \lesssim (1+t+r)|\nabla_{t,x} G| + |G|$ for any $Z \in \mathbb{K}$ and any 2-form G . Consequently, we obtain from the bootstrap assumptions (BA1)–(BA2) that,

$$|\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim (1+t+|x|) \frac{\Lambda \log(3+|t-|x||)}{(1+t+|x|)(1+|t-|x||)^2} + \frac{\Lambda}{(1+t+|x|)(1+|t-|x||)} \lesssim \Lambda \frac{\log(3+|t-|x||)}{(1+|t-|x||)^2}.$$

As previously, when $\gamma_T \geq 1$, the extra decay in $t - r$ is given by (23) and Lemma 2.19. The case $|\gamma| \leq N - 1$ is easier and follows from (BA1), (23) and Lemma 2.19. □

3.2. Structure of the proof. The remainder of the paper is divided as follows.

- (1) First, in Section 4, we prove that, for any $|\beta| \leq N$, an $L_{x,v}^\infty$ norm of $\widehat{Z}^\beta f$, weighted by powers of v^0 and z , grows at most logarithmically in time. Next, we control uniformly in time weighted space averages of $\widehat{Z}^\beta f$ for any $|\beta| \leq N - 1$. This will allow us to prove, in Section 4.4, decay estimates for $\int_v \widehat{Z}^\beta f \, dv$ and improve (BA3).
- (2) Then, we introduce the Glassey–Strauss decomposition of the electromagnetic field in Section 5.1. It allows us to improve the bootstrap assumptions (BA1) and (BA2), respectively in Sections 5.3 and 5.4, thus implying the global existence of the solution (f, F) .
- (3) Finally, refining the estimates carried out during the previous steps, we prove our modified scattering result for the distribution function in Section 6. The scattering result for the electromagnetic field is treated in Section 7 and will require an additional step, the construction of a scattering map for the vacuum Maxwell equations.

Remark 3.3. If one is interested in relaxing the assumptions on N_v and N_x , though it would force us to either modify the proof or obtain weaker rate of convergences, we give here the precise results where losses in v^0 and z occur.

- Two powers of z are lost in order to close the $L_{x,v}^\infty$ estimates in Proposition 4.5; $5 + \delta$ powers of z are required in order to apply Lemma 4.7 and prove boundedness for $\int_x f dx$ and its derivatives.
- Three powers of v^0 are lost for closing the $L_{x,v}^\infty$ estimates, and eight for the pointwise decay estimates (see Lemma 4.12 and Proposition 4.13). Finally, the Glassey–Strauss decomposition of the derivative of the Maxwell field requires losing four powers of v^0 , as suggested by Proposition 5.7 and Corollary 5.8.

Note that the various applications of Proposition 4.11 will not require controlling as many moments of f as the results mentioned here.

4. Estimates for the distribution function

4.1. Control of the Lorentz force. In view of the structure of the error terms for the commuted Vlasov equations, given by Proposition 2.4, it is important to obtain precise estimates of the Lorentz force and its derivatives by exploiting its null structure.

Lemma 4.1. *Let $|\gamma| \leq N - 2$ and $j \in \llbracket 1, 3 \rrbracket$. For all $(t, x, v) \in [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$, we have*

$$\frac{1}{v^0} |\hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu^j|(t, x) \lesssim \frac{\Lambda \log(3+t)}{(1+t+|x|)^2} + \frac{\Lambda \hat{v}^{\underline{L}}}{(1+t+|x|)(1+|t-|x||)}.$$

If $\gamma_T \geq 1$, then we have the improved estimate

$$\frac{1}{v^0} |\hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu^j|(t, x) \lesssim \frac{\Lambda}{(1+t+|x|)^{\frac{5}{2}}} + \frac{\Lambda \hat{v}^{\underline{L}}}{(1+t+|x|)(1+|t-|x||)^2}.$$

Proof. Recall the definition of the null components of a 2-form (5) and expand $\hat{v}^\mu F_\mu^j$ according to the null frame $(\underline{L}, L, e_\theta, e_\varphi)$ in order to get

$$\begin{aligned} |\hat{v}^\mu F_\mu^j| &= |\hat{v}^{\underline{L}} F_{\underline{L}}^j + \hat{v}^{\underline{L}} F_{\underline{L}}^j + \hat{v}^{e_\theta} F_{e_\theta}^j + \hat{v}^{e_\varphi} F_{e_\varphi}^j| \\ &\lesssim \hat{v}^{\underline{L}} (|\alpha(F)| + |\rho(F)|) + \hat{v}^{\underline{L}} (|\rho(F)| + |\underline{\alpha}(F)|) + |\hat{v}| (|\sigma(F)| + |\alpha(F)| + |\underline{\alpha}(F)|). \end{aligned} \tag{24}$$

Since $\hat{v}^{\underline{L}}, \hat{v}^{\underline{L}}, |\hat{v}| \leq 1$ and $|\hat{v}| + |v^0|^{-1} \leq 2\sqrt{\hat{v}^{\underline{L}}}$ by Lemma 2.5, we obtain

$$\frac{1}{v^0} |\hat{v}^\mu F_\mu^j| \lesssim \sqrt{\hat{v}^{\underline{L}}} (|\alpha(F)| + |\rho(F)| + |\sigma(F)|) + \hat{v}^{\underline{L}} |\underline{\alpha}(F)|. \tag{25}$$

Note that the same applies to $\mathcal{L}_{Z^\gamma}(F)$, $|\gamma| \leq N - 2$, so that the first estimate follows from Proposition 3.1. Assume now that $\gamma_T \geq 1$ and apply once again (25) to $\mathcal{L}_{Z^\gamma} F$ together with Proposition 3.1. We obtain

$$\begin{aligned} \frac{1}{v^0} |\hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu^j|(t, x) &\lesssim \frac{\Lambda \log(3+t) \sqrt{\hat{v}^{\underline{L}}}}{(1+t+|x|)^2 (1+|t-|x||)} + \frac{\Lambda \hat{v}^{\underline{L}}}{(1+t+|x|)(1+|t-|x||)^2} \\ &\lesssim \Lambda \frac{\log^2(3+t)}{(1+t+|x|)^3} + \frac{\Lambda \hat{v}^{\underline{L}}}{(1+t+|x|)(1+|t-|x||)^2}, \end{aligned}$$

which implies the result. □

If $N - 1 \leq |\gamma| \leq N$, we do not have improved estimates on the null components of the electromagnetic field. Moreover, if $|\gamma| = N$ and $\gamma_T = 0$, we have a very bad control of $\mathcal{L}_{Z^\gamma} F$ near the light cone. The idea then is to transform decay in $t - r$ into decay in $t + r$ at the cost of powers of \mathbf{z} and v^0 .

Lemma 4.2. *Consider $|\gamma| \leq N$ and $j \in \llbracket 1, 3 \rrbracket$. Then, for all $(t, x, v) \in \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,*

$$\frac{1}{v^0} |\hat{v}^\mu \mathcal{L}_{Z^\gamma} (F)_\mu^j|(t, x) \lesssim \frac{1}{v^0} |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda \frac{\log(3 + t + |x|)}{(1 + t + |x|)^2} |v^0|^3 \mathbf{z}^2(t, x, v),$$

and, if $\gamma_T \geq 1$,

$$\frac{1}{v^0} |\hat{v}^\mu \mathcal{L}_{Z^\gamma} (F)_\mu^j|(t, x) \lesssim \frac{1}{v^0} |\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda \frac{\log(3 + t + |x|)}{(1 + t + |x|)^3} |v^0|^3 \mathbf{z}^2(t, x, v).$$

Proof. Recall from Lemma 2.6 that $(1 + t + r)^2 \lesssim (1 + |t - r|)^2 |v^0|^4 \mathbf{z}^2$. The first estimate then follows from Proposition 3.2 and the second one from (BA2) together with (23). \square

Remark 4.3. If $|\gamma| \leq N - 1$, we have $|\mathcal{L}_{Z^\gamma} F|(t, x) \lesssim \Lambda (1 + t + |x|)^{-2} |v^0|^2 \mathbf{z}(t, x, v)$. If $|\gamma| \leq N - 2$, by combining Lemmas 2.5 and 4.1, we could even save a power of $|v^0|^3 \mathbf{z}$ in the first estimate of the Lorentz force and then avoid any loss in v .

4.2. Pointwise bounds for f and its derivatives. As explained in Section 2.8.2, the main difficulties here are related to the weak decay rate of the electromagnetic field. We deal with them by exploiting several hierarchies in the commuted equations and by taking advantage of the null structure of the Lorentz force. Our approach, based on the method of characteristics, will require various applications of the following result.

Lemma 4.4. *Let $g : [0, T] \times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}_+$ and $h : [0, T] \times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}_+$ be two nonnegative sufficiently regular functions such that, for all $(t, x, v) \in [0, T] \times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,*

$$|\mathbf{T}_F(g)|(t, x, v) \leq \frac{C_g}{(1 + t) \log^2(3 + t)} g + \frac{C_g \hat{v}^L}{(1 + |t - |x||) \log^2(3 + |t - |x||)} g + \frac{1}{(1 + t) \log^2(3 + t)} h$$

for some constant $C_g > 0$. Then,

$$\forall (t, x, v) \in [0, T] \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |g|(t, x, v) \leq (\|g(0, \cdot, \cdot)\|_{L_{x,v}^\infty} + 3 \|h\|_{L_{t,x,v}^\infty}) e^{6C_g}.$$

Proof. Fix, for all of this proof, $(x, v) \in \mathbb{R}_x^3 \times \mathbb{R}_v^3$ and denote by $t \mapsto (X_t, V_t)$ the characteristic of the operator $\mathbf{T}_F = \partial_t + \hat{v}^i \partial_{x^i} + \hat{v}^\mu F_\mu^j \partial_{v^j}$ satisfying,

$$\forall 1 \leq j \leq 3, \quad \dot{X}_t^j = \hat{V}_t^j, \quad \dot{V}_t^j = \hat{V}_t^\mu F_\mu^j(t, X_t), \quad X_0 = x, \quad V_0 = v.$$

According to the Duhamel formula, we have,

$$\forall t \in [0, T], \quad g(t, X_t, V_t) = g(0, x, v) + \int_{s=0}^t \mathbf{T}_F(g)(s, X_s, V_s) ds.$$

We are then lead to introduce the two functions

$$\psi_1(s) := (1 + s)^{-1} \log^{-2}(3 + s), \quad \psi_2(s) := \hat{v}^L(X_s) (1 + |s - |X_s||)^{-1} \log^{-2}(3 + |s - |X_s||).$$

In view of the expression of $T_F(g)$, we have, for all $t \in [0, T[$,

$$g(t, X_t, V_t) \leq \|g(0, \cdot, \cdot)\|_{L^\infty_{x,v}} + \|h\|_{L^\infty_{t,x,v}} \int_{s=0}^t \psi_1(s) \, ds + \int_{s=0}^t C_g(\psi_1(s) + \psi_2(s))g(s, X_s, V_s) \, ds.$$

Consequently, Grönwall’s inequality and

$$\int_{s=0}^{+\infty} \psi_1(s) \, ds = \int_{s=0}^{+\infty} \frac{ds}{(1+s)\log^2(3+s)} \leq \int_{s=0}^{+\infty} \frac{3 \, ds}{(3+s)\log^2(3+s)} \leq \frac{3}{\log(3)} \leq 3$$

yield,

$$\forall t \in [0, T[, \quad \sup_{0 \leq s \leq t} g(s, X_s, V_s) \leq (\|g(0, \cdot, \cdot)\|_{L^\infty_{x,v}} + 3\|h\|_{L^\infty_{t,x,v}}) \exp\left(3C_g + C_g \int_{s=0}^t \psi_2(s) \, ds\right).$$

It remains us to estimate the integral of ψ_2 . For this, we will perform a change of variables reflecting that the Vlasov operator reads, in the coordinate system (u, x, v) , where $u = t - |x|$,

$$T_F = \partial_u - \hat{v}^i \frac{X_i}{|x|} \partial_u + \hat{v}^i \partial_{x^i} + \hat{v}^\mu F_\mu^j \partial_{v^j} = 2\hat{v}^L \partial_u + \hat{v}^i \partial_{x^i} + \hat{v}^\mu F_\mu^j \partial_{v^j}.$$

As $\hat{v}^L > 0$ by [Lemma 2.5](#), we can then parametrize $t \mapsto (X_t, V_t)$ by the variable u . Hence, we perform the change of variables $\tilde{u}(s) = s - |X_s|$, so that $\tilde{u}'(s) = 2\widehat{V}^L(X_s) > 0$ and

$$\int_{s=0}^t \psi_2(s) \, ds = \int_{u=t-|x|}^{\tilde{u}(t)} \frac{du}{2(1+|u|)\log^2(3+|u|)} \leq \int_{u \in \mathbb{R}} \frac{du}{2(1+|u|)\log^2(3+|u|)} \leq 3. \quad \square$$

We are now able to prove that quantities such as $z\widehat{Z}^\beta f$ are almost uniformly bounded in phase space. We recall that for a multi-index β , the number of homogeneous vector fields (respectively translations) composing \widehat{Z}^β is denoted by β_H (respectively β_T).

Proposition 4.5. *There exists $D > 0$, depending only on (N, N_v, N_x) , such that the following estimates hold. For all $(t, x, v) \in [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,*

$$\forall 0 \leq q \leq N_x, \quad |\beta| \leq N - 2, \quad |v^0|^{N_v} |z^q \widehat{Z}^\beta f|(t, x, v) \lesssim e^{D\Lambda} \log^{3q+3\beta_H}(3+t), \quad (26)$$

$$\forall 0 \leq q \leq N_x - 2, \quad |\beta| \leq N, \quad |v^0|^{N_v-3} |z^q \widehat{Z}^\beta f|(t, x, v) \lesssim e^{D\Lambda} \log^{3q+3\beta_H}(3+t). \quad (27)$$

Throughout this paper, it will be convenient to work with $\bar{\epsilon} := \epsilon e^{(D+1)\Lambda}$.

Proof. For simplicity, we assume here that $N \geq 4$ and we sketch the proof of the case $N = 3$ in [Remark 4.6](#) below. Note further that, by interpolation, it suffices to deal with the cases $q \in \{0, N_x\}$ for (26) and $q \in \{0, N_x - 2\}$ for (27). Motivated by the analysis of the toy model carried out in [Section 2.8.2](#), we introduce the following hierarchized norms in order to deal with nonintegrable error terms and still obtain satisfying estimates if the electromagnetic field is large. Consider, for $(N_0, p, q) = (N - 2, N_v, N_x)$ or $(N, N_v - 3, N_x - 2)$,

$$\mathbb{E}_{N_0}^{p,q}[f](t, x, v) := \sum_{|\beta| \leq N_0} \frac{|v^0|^p |\widehat{Z}^\beta f|(t, x, v)}{\log^{3\beta_H}(3+t)} + \frac{|v^0|^p |z^q \widehat{Z}^\beta f|(t, x, v)}{\log^{3q+3\beta_H}(3+t)}$$

and let us prove that, for all $(t, x, v) \in [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,

$$\mathbf{T}_F(\mathbb{E}_{N-2}^{N_v, N_x}[f])(t, x, v) \lesssim \frac{\Lambda \mathbb{E}[f]_{N-2}^{N_v, N_x}(t, x, v)}{(1+t) \log^2(3+t)} + \frac{\Lambda \hat{v}^L(x) \mathbb{E}[f]_{N-2}^{N_v, N_x}(t, x, v)}{(1+|t-|x||) \log^2(3+|t-|x||)}, \tag{28}$$

$$\begin{aligned} \mathbf{T}_F(\mathbb{E}_N^{N_v-3, N_x-2}[f])(t, x, v) &\lesssim \frac{\Lambda \mathbb{E}_N^{N_v-3, N_x-2}[f](t, x, v)}{(1+t) \log^2(3+t)} \\ &\quad + \frac{\Lambda \hat{v}^L(x) \mathbb{E}_N^{N_v-3, N_x-2}[f]}{(1+|t-|x||) \log^2(3+|t-|x||)} + \frac{\Lambda \mathbb{E}_{N-2}^{N_v, N_x}[f](t, x, v)}{(1+t) \log^2(3+t)}. \end{aligned} \tag{29}$$

We are able to apply \mathbf{T}_F to these energy norms since $\mathbf{T}_F(|h|) = \mathbf{T}_F(h)(h/|h|)$ almost everywhere for any $h \in W_{\text{loc}}^{1,1}$. The result would then follow from two applications of [Lemma 4.4](#). Fix now $(t, x, v) \in [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$ as well as either $|\beta| \leq N-2$, $p = N_v$ and $a \in \{0, N_x\}$ or $|\beta| \leq N$, $p = N_v - 3$ and $a \in \{0, N_x - 2\}$. Note then, since $\mathbf{T}_F(\log^{-1}(3+t)) < 0$, that

$$\begin{aligned} \mathbf{T}_F\left(\frac{|v^0|^p \mathbf{z}^a |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)}\right) &\leq p \mathbf{T}_F(v^0) \frac{|v^0|^{p-1} \mathbf{z}^a |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)} \\ &\quad + a \mathbf{T}_F(\mathbf{z}) \frac{|v^0|^p \mathbf{z}^{a-1} |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)} + \mathbf{T}_F(\widehat{Z}^\beta f) \frac{\widehat{Z}^\beta f}{|\widehat{Z}^\beta f|} \frac{|v^0|^p \mathbf{z}^a}{\log^{3a+3\beta_H}(3+t)}. \end{aligned} \tag{30}$$

It is important to note that the second term on the right-hand side vanishes if $a = 0$. We start by dealing with the first two terms on the right-hand side since the last one requires a more thorough analysis. As $|\nabla_v v^0| \leq 1$, we obtain, by applying [Lemma 4.1](#),

$$\frac{1}{v^0} |\mathbf{T}_F(v^0)|(t, x, v) = \frac{1}{v^0} |\hat{v}^\mu F_\mu^j \partial_{v^j}(v^0)|(t, x) \lesssim \frac{\Lambda \log(3+t)}{(1+t+|x|)^2} + \frac{\Lambda \hat{v}^L}{(1+t+|x|)(1+|t-|x||)}, \tag{31}$$

so that

$$|\mathbf{T}_F(v^0)| \frac{|v^0|^{p-1} |\mathbf{z}^a \widehat{Z}^\beta f|(t, x, v)}{\log^{3a+3\beta_H}(3+t)} \lesssim \left(\frac{\Lambda}{(1+t)^{\frac{3}{2}}} + \frac{\Lambda \hat{v}^L}{(1+t)(1+|t-|x||)} \right) \frac{|v^0|^p |\mathbf{z}^a \widehat{Z}^\beta f|(t, x, v)}{\log^{3a+3\beta_H}(3+t)}. \tag{32}$$

Next, recall from [\(11\)](#) the identity $\hat{v}^\mu \partial_{x^\mu}(\mathbf{z}) = 0$ and note that $|\nabla_v \mathbf{z}| \lesssim (t+r)/v^0$. We get, using [Lemma 4.1](#),

$$|\mathbf{T}_F(\mathbf{z})|(t, x, v) \lesssim \sum_{1 \leq j \leq 3} \frac{t+|x|}{v^0} |\hat{v}^\mu F_\mu^j(t, x)| \lesssim \frac{\Lambda \log(3+t)}{1+t+|x|} + \frac{\Lambda \hat{v}^L}{1+|t-|x||}.$$

Using Young inequality for products, we obtain, if $a \neq 0$,

$$\frac{\mathbf{z}^{a-1}}{\log^{3a}(3+t)} \leq \frac{a-1}{a \log^3(3+t)} \frac{\mathbf{z}^a}{\log^{3a}(3+t)} + \frac{1}{a \log^3(3+t)}.$$

We then deduce that

$$\begin{aligned} a |\mathbf{T}_F(\mathbf{z})| \frac{|v^0|^p \mathbf{z}^{a-1} |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)} \\ \lesssim \left(\frac{\Lambda}{(1+t) \log^2(3+t)} + \frac{\Lambda \hat{v}^L}{(1+|t-|x||) \log^3(3+t)} \right) \left(\frac{|v^0|^p |\widehat{Z}^\beta f|}{\log^{3\beta_H}(3+t)} + \frac{|v^0|^p \mathbf{z}^a |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)} \right). \end{aligned} \tag{33}$$

We now focus on the last term in (30). The first step consists in applying the commutation formula of Proposition 2.4 and noting that $v^0 \partial_{v^i} = \widehat{\Omega}_{0i} - t \partial_{x^i} - x^i \partial_t$. We can then bound

$$|\mathbf{T}_F(\widehat{Z}^\beta f)| |v^0|^p \mathbf{z}^a \log^{-3a-3\beta_H}(3+t)$$

by a linear combination of terms of the following form. The good ones, which are strongly decaying and can then be easily handled,

$$\mathcal{G}_{\gamma,\kappa}^{p,a} := \frac{1}{v^0} |\widehat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu{}^j| \frac{|v^0|^p \mathbf{z}^a |\widehat{\Omega}_{0j} \widehat{Z}^\kappa f|}{\log^{3a+3\beta_H}(3+t)}, \quad |\gamma| + |\kappa| \leq |\beta|, |\kappa| \leq |\beta| - 1, \tag{34}$$

and the bad ones,

$$\mathcal{B}_{\gamma,\kappa}^{p,a} := (t+r) \sup_{1 \leq j \leq 3} \frac{1}{v^0} |\widehat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_\mu{}^j| \frac{|v^0|^p \mathbf{z}^a |\partial_{t,x} \widehat{Z}^\kappa f|}{\log^{3a+3\beta_H}(3+t)}, \quad \begin{cases} \gamma_H + \kappa_H \leq \beta_H, \\ \kappa_H = \beta_H \implies \gamma_T \geq 1, \end{cases} \tag{35}$$

where, again, $|\gamma| + |\kappa| \leq |\beta|$ and $|\kappa| \leq |\beta| - 1$. We emphasize that $\widehat{Z}^\xi := \partial_{t,x} \widehat{Z}^\kappa$ is composed of the same number of homogeneous vector fields as \widehat{Z}^κ , so that $\xi_H = \kappa_H$. In contrast, $\widehat{Z}^\zeta := \widehat{\Omega}_{0j} \widehat{Z}^\kappa$ satisfies $\zeta_H = \kappa_H + 1$. Moreover, $\widehat{\Omega}_{0j} \widehat{Z}^\kappa$ and $\partial_{t,x} \widehat{Z}^\kappa$ are of order at most $|\beta|$.

Consider first the case $|\beta| \leq N - 2$, so that $p = N_v$ and $a \in \{0, N_x\}$, and fix two multi-indices $|\gamma| \leq |\beta|$, $|\kappa| \leq |\beta| - 1$. Then, according to Lemma 4.1, we have

$$\begin{aligned} \mathcal{G}_{\gamma,\kappa}^{N_v,a} &\lesssim \Lambda \left(\frac{\log(3+t)}{(1+t+|x|)^2} + \frac{\widehat{v}^L}{(1+t+|x|)(1+|t-|x|)} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \widehat{\Omega}_{0j} \widehat{Z}^\kappa f|(t,x,v)}{\log^{3a+3\beta_H}(3+t)} \\ &\lesssim \left(\frac{\Lambda}{(1+t)^{\frac{3}{2}}} + \frac{\Lambda \widehat{v}^L}{(1+t)^{\frac{1}{2}}(1+|t-|x|)} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \widehat{\Omega}_{0j} \widehat{Z}^\kappa f|(t,x,v)}{\log^{3a+3(\kappa_H+1)}(3+t)}. \end{aligned} \tag{36}$$

We now focus on $\mathcal{B}_{\gamma,\kappa}^{N_v,a}$ and we start by treating the case $\kappa_H = \beta_H$ and $\gamma_T \geq 1$. Applying once again Lemma 4.1, we get

$$\begin{aligned} \mathcal{B}_{\gamma,\kappa}^{N_v,a} &\lesssim (t+|x|) \left(\frac{\Lambda}{(1+t+|x|)^{\frac{5}{2}}} + \frac{\Lambda \widehat{v}^L}{(1+t+|x|)(1+|t-|x|)^2} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \partial_{t,x} \widehat{Z}^\kappa f|}{\log^{3a+3\beta_H}(3+t)} \\ &\leq \left(\frac{\Lambda}{(1+t)^{\frac{3}{2}}} + \frac{\Lambda \widehat{v}^L}{(1+|t-|x|)^2} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \partial_{t,x} \widehat{Z}^\kappa f|}{\log^{3a+3\kappa_H}(3+t)}. \end{aligned} \tag{37}$$

Otherwise $\kappa_H \leq \beta_H - 1$, so necessarily $\beta_H \geq 1$, and

$$\begin{aligned} \mathcal{B}_{\gamma,\kappa}^{N_v,a} &\lesssim (t+|x|) \left(\frac{\Lambda \log(3+t)}{(1+t+|x|)^2} + \frac{\Lambda \widehat{v}^L}{(1+t+|x|)(1+|t-|x|)} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \partial_{t,x} \widehat{Z}^\kappa f|}{\log^{3a+3\beta_H}(3+t)} \\ &\leq \left(\frac{\Lambda}{(1+t) \log^2(3+t)} + \frac{\Lambda \widehat{v}^L}{(1+|t-|x|) \log^3(3+t)} \right) \frac{|v^0|^{N_v} |\mathbf{z}^a \partial_{t,x} \widehat{Z}^\kappa f|}{\log^{3a+3\kappa_H}(3+t)}. \end{aligned} \tag{38}$$

We obtain from (30)–(33) and (36)–(38),

$$\mathbf{T}_F \left(\frac{|v^0|^{N_v} \mathbf{z}^a |\widehat{Z}^\beta f|}{\log^{3a+3\beta_H}(3+t)} \right) \lesssim \frac{\Lambda \mathbb{E}_{N-2}^{N_v, N_x}[f](t,x,v)}{(1+t) \log^2(3+t)} + \frac{\Lambda \widehat{v}^L(x) \mathbb{E}_{N-2}^{N_v, N_x}[f](t,x,v)}{(1+|t-|x|) \log^2(3+t)} + \frac{\Lambda \widehat{v}^L(x) \mathbb{E}_{N-2}^{N_v, N_x}[f](t,x,v)}{(1+|t-|x|)^2}.$$

As $|t - r| \gtrsim t$ for $r \geq 2t$ and $t \geq |t - r|$ otherwise, we have

$$(1 + |t - r|)^{-1} \log^{-2}(3 + t) \lesssim (1 + t)^{-1} \log^{-2}(3 + t) + (1 + |t - r|)^{-1} \log^{-2}(3 + |t - r|) \quad (39)$$

and we then deduce that (28) holds. Lemma 4.4 then implies (26).

Assume now that $N - 1 \leq |\beta| \leq N$, $p = N_v - 3$ and $a \in \{0, N_x - 2\}$. We fix two multi-indices γ, κ verifying $|\gamma| + |\kappa| \leq |\beta|$, $|\kappa| \leq |\beta| - 1$ and we consider two cases.

Case 1: $|\gamma| \leq N - 2$. The Lorentz force can still be estimated using Lemma 4.1. One can then follow the analysis carried out in (36)–(39) and obtain

$$\mathcal{G}_{\gamma, \kappa}^{N_v-3, a}, \mathcal{B}_{\gamma, \kappa}^{N_v-3, a} \lesssim \frac{\Lambda \mathbb{E}_N^{N_v-3, N_x-2}[f](t, x, v)}{(1 + t) \log^2(3 + t)} + \frac{\Lambda \hat{v}^L(x) \mathbb{E}_N^{N_v-3, N_x-2}[f](t, x, v)}{(1 + |t - |x||) \log^2(3 + |t - |x||)}, \quad (40)$$

where the term $\mathcal{B}_{\gamma, \kappa}^{N_v-3, a}$ is of course merely defined when γ_T and κ_H satisfy the condition given in (35).

Case 2: $N - 1 \leq |\gamma| \leq N$. Then, as $N \geq 4$, we have $|\kappa| \leq 1$ so that we will be able to control the terms (34)–(35) using (26). In particular, we are allowed to lose two powers of $|v^0|^2 z$ in the upcoming estimates in order to deal with the weak decay rate of $\mathcal{L}_{Z^\gamma} F$ near the light cone. More precisely, using first Lemma 4.2 and then $a + 2 \leq N_x$,

$$\begin{aligned} \mathcal{G}_{\gamma, \kappa}^{N_v-3, a} &\lesssim \frac{\Lambda \log(3 + t + |x|)}{(1 + t + |x|)^2} |v^0|^3 z^2(t, x, v) \frac{|v^0|^{N_v-3} |z^a \widehat{\Omega}_{0j} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3a+3\beta_H}(3 + t)} \\ &\lesssim \frac{\Lambda}{(1 + t)^{\frac{3}{2}}} \frac{|v^0|^{N_v} |z^{N_x} \widehat{\Omega}_{0j} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3N_x+3(\kappa_H+1)}(3 + t)}. \end{aligned}$$

Next, consider the terms (35) and assume first that $\gamma_T \geq 1$. In that case, $\mathcal{B}_{\gamma, \kappa}^{N_v-3, a}$ can be easily handled since it is strongly decaying. Indeed, using again Lemma 4.2, we get

$$\begin{aligned} \mathcal{B}_{\gamma, \kappa}^{N_v-3, a} &\lesssim (t + |x|) \frac{\Lambda \log(3 + t + |x|)}{(1 + t + |x|)^3} |v^0|^3 z^2(t, x, v) \frac{|v^0|^{N_v-3} |z^a \partial_{t,x} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3a+3\beta_H}(3 + t)} \\ &\lesssim \frac{\Lambda}{(1 + t)^{\frac{3}{2}}} \frac{|v^0|^{N_v} |z^{N_x} \partial_{t,x} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3N_x+3\kappa_H}(3 + t)}. \end{aligned}$$

Finally, if $\gamma_T = 0$, we necessarily have $\gamma_H = |\gamma| \geq N - 1 \geq 3$. Since $\beta_H \geq \gamma_H + \kappa_H$, we have $\kappa_H \leq \beta_H - 3$, so that $3a + 3\beta_H \geq 3(a + 2) + 3\kappa_H + 3$. Thus, Lemma 4.2 yields

$$\begin{aligned} \mathcal{B}_{\gamma, \kappa}^{N_v-3, a} &\lesssim (t + |x|) \frac{\Lambda \log(3 + t + |x|)}{(1 + t + |x|)^2} |v^0|^3 z^2(t, x, v) \frac{|v^0|^{N_v-3} |z^a \partial_{t,x} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3a+3\beta_H}(3 + t)} \\ &\lesssim \frac{\Lambda}{(1 + t) \log^2(3 + t)} \frac{|v^0|^{N_v} |z^{a+2} \partial_{t,x} \widehat{Z}^\kappa f|(t, x, v)}{\log^{3(a+2)+3\kappa_H}(3 + t)}. \end{aligned}$$

We then deduce that, in this case,

$$\mathcal{G}_{\gamma, \kappa}^{N_v-3, a}, \mathcal{B}_{\gamma, \kappa}^{N_v-3, a} \lesssim \frac{\Lambda}{(1 + t) \log^2(3 + t)} \mathbb{E}_{N-2}^{N_v, N_x}[f](t, x, v).$$

The estimate (29) ensues from (40) and this last inequality. To conclude the proof, it then remains to apply again the previous Lemma 4.4. □

Remark 4.6. If $N = 3$, the proof of [Proposition 4.5](#) requires an additional step. Once the estimate for $\mathbb{E}_{N-2}^{N_v, N_x}[f]$ is proved, we need to control the intermediary norm $\mathbb{E}_{N-1}^{N_v-1, N_x-1}[f]$. For this, compared to the treatment of $\mathbb{E}_N^{N_v-3, N_x-2}[f]$ carried out during the proof of [Proposition 4.5](#), there are two differences.

- First, we can exploit the much stronger decay estimate satisfied by the derivatives of order $N - 1$ of the electromagnetic field than that on its top-order ones (see [Proposition 3.2](#)). This explains why we can propagate higher moments for the derivatives of order $N - 1$ of f than for the top-order ones.
- Moreover, for controlling sufficiently well $\mathcal{B}_{\gamma, \kappa}^{N_v-1, 0}$ and $\mathcal{B}_{\gamma, \kappa}^{N_v-1, N_x-1}$ in the case $\beta_H = \kappa_H$, we can prove, through a direct application of [Lemma 2.17](#), that the good null components of $\mathcal{L}_{Z^\gamma}(F)$ still satisfy improved estimates when $|\gamma| = N - 1$ and $\gamma_T \geq 1$.

Finally, in order to bound uniformly in time $\mathbb{E}_N^{N_v-3, N_x-2}[f]$, the analysis of the terms [\(34\)–\(35\)](#) is slightly more technical. It is necessary to consider three cases ($|\gamma| \leq N - 2$, $|\gamma| = N - 1$ as well as $|\gamma| = N$) and to use the estimates on the first two energy norms.

4.3. Uniform boundedness of the spatial averages. We start by a preparatory result, which will also be useful later in [Section 6](#). Recall the constant $\bar{\epsilon} := \epsilon e^{(D+1)\Lambda}$ introduced in [Proposition 4.5](#).

Lemma 4.7. *For any $|\beta| \leq N - 1$, we have,*

$$\forall (t, v) \in [0, T[\times \mathbb{R}_v^3, \quad |v^0|^{N_v-6} \left| \partial_t \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) dx \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{(1+t)^2}.$$

Proof. Fix $|\beta| \leq N - 1$, $t \in [0, T[$ and $v \in \mathbb{R}_v^3$. Integrating the commutation formula of [Proposition 2.4](#) for $\widehat{Z}^\beta f$ and performing integration by parts in x gives

$$\begin{aligned} \partial_t \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) dx &= - \int_{\mathbb{R}_x^3} \widehat{v}^\mu F_{\mu}{}^j \partial_{v^j} \widehat{Z}^\beta f(t, x, v) dx \\ &\quad + \sum_{|\gamma|+|\kappa| \leq |\beta|} C_{\gamma, \kappa}^\beta \int_{\mathbb{R}_x^3} \widehat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu}{}^j \partial_{v^j} \widehat{Z}^\kappa f(t, x, v) dx. \end{aligned}$$

Now, we write

$$v^0 \partial_{v^j} = \widehat{\Omega}_{0j} - x^j \partial_t - t \partial_{x^j} = \widehat{\Omega}_{0j} - (x^j - \widehat{v}^j t) \partial_t - v^j S + v^j x^i \partial_{x^i} - t \partial_{x^j}, \quad |x^j - \widehat{v}^j t| \leq z,$$

so that, integrating once again by parts,

$$\begin{aligned} \left| \partial_t \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) dx \right| &\lesssim \sum_{\substack{|\gamma|+|\kappa| \leq |\beta|+1 \\ |\gamma| \leq |\beta|}} \sup_{1 \leq j \leq 3} \int_{\mathbb{R}_x^3} \frac{1}{v^0} |\widehat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu}{}^j(t, x)| |z \widehat{Z}^\kappa f|(t, x, v) dx \\ &\quad + \int_{\mathbb{R}_x^3} \frac{t + |x|}{v^0} |\widehat{v}^\mu \nabla_{t,x} \mathcal{L}_{Z^\gamma}(F)_{\mu}{}^j(t, x)| |\widehat{Z}^\kappa f|(t, x, v) dx. \end{aligned}$$

According to the bootstrap assumptions [\(BA1\)–\(BA2\)](#) and [Lemma 2.6](#), we have

$$\begin{aligned} |\mathcal{L}_{Z^\gamma}(F)_{\mu}{}^j(t, x)| &\lesssim \Lambda (1+t+|x|)^{-2} |v^0|^2 z, \\ |\nabla_{t,x} \mathcal{L}_{Z^\gamma}(F)_{\mu}{}^j(t, x)| &\lesssim \Lambda \log(3+t+|x|) (1+t+|x|)^{-3} |v^0|^4 z^2, \end{aligned}$$

so that

$$\begin{aligned} \left| \partial_t \int_{\mathbb{R}^3_x} \widehat{Z}^\beta f(t, x, v) dx \right| &\lesssim \Lambda \sum_{|\kappa| \leq |\beta|+1} \int_{\mathbb{R}^3_x} \frac{\log(3+t+|x|)}{(1+t+|x|)^2} |v^0|^3 |z^2 \widehat{Z}^\kappa f|(t, x, v) dx \\ &\leq \Lambda \sup_{|\kappa| \leq |\beta|+1} \sup_{x \in \mathbb{R}^3} \left(\frac{\log(3+t+|x|)}{(1+t+|x|)^2} |v^0|^3 |z^{N_x-2} \widehat{Z}^\kappa f|(t, x, v) \right) \int_{\mathbb{R}^3_x} \frac{dx}{z^{N_x-4}(t, x, v)}. \end{aligned}$$

Note then that, in view of (11) and $N_x > 7$,

$$\int_{\mathbb{R}^3_x} \frac{dx}{z^{N_x-4}(t, x, v)} \leq \int_{\mathbb{R}^3_x} \frac{dx}{(1+|x-\hat{v}t|)^{N_x-4}} = \int_{y \in \mathbb{R}^3} \frac{dy}{(1+|y|)^{N_x-4}} < +\infty.$$

Then, multiply both sides of the inequality by $|v^0|^{N_v-6}$ and bound above the right-hand side by applying Proposition 4.5. It remains to use $\Lambda \epsilon e^{D\Lambda} \leq \epsilon e^{(D+1)\Lambda} = \bar{\epsilon}$. □

Remark 4.8. If $|\beta| \leq N - 3$, by using the estimates of the Lorentz force provided by Lemma 4.1, we can even prove $|v^0|^{N_v} |\partial_t \int_x \widehat{Z}^\beta f dv| \lesssim \bar{\epsilon} (1+t)^{-2} \log^{-3N_x-3N} (3+t)$.

Note now that $|\int_x \widehat{Z}^\beta f(0, x, v) dx| \leq 2 \sup_x |z^4 \widehat{Z}^\beta f|(0, x, v) \leq 2\epsilon$. Hence, by integrating in time the inequality of the previous Lemma 4.7, we obtain, for any $|\beta| \leq N - 1$,

$$\forall (t, v) \in [0, T[\times \mathbb{R}^3_v, \quad |v^0|^{N_v-6} \left| \int_{\mathbb{R}^3_x} \widehat{Z}^\beta f(t, x, v) dx \right| \lesssim \epsilon + \bar{\epsilon} \int_{\tau=0}^t \frac{\log^{3N_x+3N} (3+\tau)}{(1+\tau)^2} d\tau \lesssim \bar{\epsilon}.$$

It directly implies the following result.

Corollary 4.9. *Let $|\beta| \leq N - 1$ and $\psi : \mathbb{S}^2_\omega \times \mathbb{R}^3_v \rightarrow \mathbb{R}$ be a function such that $\|\psi(\cdot, v)\|_{L^\infty_\omega} \lesssim |v^0|^{N_v-6}$. Then, for any $\omega \in \mathbb{S}^2$,*

$$\forall (t, v) \in [0, T[\times \mathbb{R}^3_v, \quad \left| \psi(\omega, v) \int_{\mathbb{R}^3_x} \widehat{Z}^\beta f(t, x, v) dx \right| \lesssim \bar{\epsilon}.$$

We allowed the function ψ to depend on a parameter $\omega \in \mathbb{S}^2$ in order to prove optimal decay estimates on certain elements of the Glassey–Strauss decomposition of the electromagnetic field, defined as integral kernels.

4.4. Pointwise decay estimates for velocity averages. We prove here that the decay rate of $\int_v \widehat{Z}^\beta f dv$, for $|\beta| \leq N - 1$, coincides with the one of the linear setting. In particular, we improve the bootstrap assumption (BA3). The starting point consists of performing the change of variables $y = x - t\hat{v}$. For this, recall Lemma 2.9 and that $y \mapsto \check{y}$ is the inverse function of $v \mapsto \hat{v}$.

Lemma 4.10. *Let $g : [0, T[\times \mathbb{R}^3_x \times \mathbb{R}^3_v \rightarrow \mathbb{R}$ be a sufficiently regular function. Then,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad t^3 \int_{\mathbb{R}^3_v} g(t, x - \hat{v}t, v) dv = \int_{|y-x|<t} (|v^0|^5 g) \left(t, y, \frac{\widetilde{x-y}}{t} \right) dy.$$

This change of variables is motivated by the linear case. Any regular solution to the relativistic transport equation $\partial_t h + \hat{v} \cdot \nabla_x h = 0$ is constant along the timelike straight lines, $h(t, x + \hat{v}t, v) = h(0, x, v)$. The previous lemma, applied for $g(t, x, v) = h(0, x, v)$, then leads to $\int_v h(t, x, v) dv \lesssim t^{-3}$.

As a first step, we control $\int_v |\widehat{Z}^\beta f| dv$ for any $|\beta| \leq N$, which has a slightly slower decay rate than in the linear case in the interior of the light cone. These estimates will also be useful on their own.

Proposition 4.11. *Let $|\beta| \leq N$ and $0 \leq a \leq N_x - 6$. Then, the following properties hold.*

- *Almost optimal pointwise decay estimate,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}_x^3, \quad \int_{\mathbb{R}_v^3} |v^0|^{N_v-8} |z^a \widehat{Z}^\beta f|(t, x, v) dv \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{(1+t)^3}.$$

- *Improved decay estimates near and in the exterior of the light cone,*

$$\begin{aligned} \forall |x| \leq t < T, \quad & \int_{\mathbb{R}_v^3} |v^0|^{N_v-8-2a} |\widehat{Z}^\beta f|(t, x, v) dv \lesssim \bar{\epsilon} \log^{3N_x+3N}(3+t) \frac{(1+t-|x|)^a}{(1+t)^{3+a}}, \\ \forall t < \sup(|x|, T), \quad & \int_{\mathbb{R}_v^3} |v^0|^{N_v-8-2a} |\widehat{Z}^\beta f|(t, x, v) dv \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{(1+t+|x|)^{3+a}}. \end{aligned}$$

Proof. Fix $|\beta| \leq N$, $(t, x) \in [0, T[\times \mathbb{R}_x^3$ and $0 \leq a \leq N_x - 6$. If $t \leq 1$, we have by [Proposition 4.5](#),

$$\int_{\mathbb{R}_v^3} |v^0|^{N_v-7} |z^a \widehat{Z}^\beta f|(t, x, v) dv \lesssim \sup_{v \in \mathbb{R}^3} |v^0|^{N_v-3} |z^{N_x-6} \widehat{Z}^\beta f|(t, x, v) \int_{\mathbb{R}_w^3} \frac{dw}{\langle w \rangle^4} \lesssim \bar{\epsilon}.$$

Assume now, unless $T \leq 1$, that $t \geq 1$ and introduce the function $g(t, x, v) := |v^0|^{N_v-8} |z^a \widehat{Z}^\beta f|(t, x+t\hat{v}, v)$.

Applying the previous [Lemma 4.10](#) to g , we get

$$\begin{aligned} t^3 \int_{\mathbb{R}_v^3} |v^0|^{N_v-8} |z^a \widehat{Z}^\beta f|(t, x, v) dv &\leq \int_{|y-x|<t} \sup_{v \in \mathbb{R}^3} |v^0|^5 g(t, y, v) dy \\ &\leq \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |v^0|^5 \langle y \rangle^4 g(t, y, v) \int_{\mathbb{R}_y^3} \frac{dy}{\langle y \rangle^4}. \end{aligned}$$

Using now [Lemma 2.8](#) and then [Proposition 4.5](#), we obtain

$$t^3 \int_{\mathbb{R}_v^3} |v^0|^{N_v-8} |z^a \widehat{Z}^\beta f|(t, x, v) dv \leq \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |v^0|^{N_v-3} |z^{a+4} \widehat{Z}^\beta f|(t, y, v) \lesssim \bar{\epsilon} \log^{3a+12+3N}(3+t).$$

This concludes the proof of the first estimate, which, together with [Lemma 2.6](#), implies the second one as well as the last one in the region $t < |x| \leq 2t$. If $|x| \geq 2t$, note that $z \gtrsim 1 + |x - t\hat{v}| \gtrsim 1 + t + |x|$, so that

$$\int_{\mathbb{R}_v^3} |v^0|^{N_v-7} |\widehat{Z}^\beta f|(t, x, v) dv \lesssim (1+t+|x|)^{-N_x+2} \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |v^0|^{N_v-3} |z^{N_x-2} \widehat{Z}^\beta f|(t, y, v) \int_{\mathbb{R}_w^3} \frac{dw}{\langle w \rangle^4}.$$

It remains to apply [Proposition 4.5](#). □

Our goal now is to remove the logarithmic loss of the estimate of $\int_v \widehat{Z}^\beta f dv$ provided by [Proposition 4.11](#). Since our analysis will rely on the following result, we will not be able to deal with top-order derivatives. We recall that $N_x - 3 > 4$.

Lemma 4.12. *Let $g : [0, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a sufficiently regular function. Then, for all $|x| < t < T$,*

$$\left| t^3 \int_{\mathbb{R}_v^3} g(t, x - \hat{v}t, v) dv - \int_{y \in \mathbb{R}^3} (|v^0|^5 g) \left(t, y, \frac{\tilde{x}}{t} \right) dy \right| \lesssim \frac{1}{t} \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} \langle y \rangle^{N_x-3} (|v^0|^7 |g| + |v^0|^8 |\nabla_v g|)(t, y, v).$$

Proof. According to Lemma 4.10, we have

$$t^3 \int_{\mathbb{R}_v^3} g(t, x - \hat{v}t, v) \, dv - \int_{y \in \mathbb{R}^3} g\left(t, y, \frac{\check{x}}{t}\right) \, dy = \mathcal{I}_1 + \mathcal{I}_2,$$

where

$$\begin{aligned} \mathcal{I}_1 &:= \int_{|x-y|<t} (|v^0|^5 g)\left(t, y, \frac{\widetilde{x-y}}{t}\right) \, dy - \int_{|x-y|<t} (|v^0|^5 g)\left(t, y, \frac{\check{x}}{t}\right) \, dy, \\ \mathcal{I}_2 &:= - \int_{|x-y|\geq t} (|v^0|^5 g)\left(t, y, \frac{\check{x}}{t}\right) \, dy. \end{aligned}$$

Since, by Lemma 2.9, we have $|\nabla_y \check{y}| \lesssim \sqrt{1-|y|^2}^{-3} = \langle \check{y} \rangle^3 = |v^0|^3 \langle \check{y} \rangle$, the mean value theorem gives us

$$\left| (|v^0|^5 g)\left(t, y, \frac{\widetilde{x-y}}{t}\right) - (|v^0|^5 g)\left(t, y, \frac{\check{x}}{t}\right) \right| \lesssim \frac{|y|}{t} \sup_{v \in \mathbb{R}^3} |v^0|^7 |g|(t, y, v) + |v^0|^8 |\nabla_v g|(t, y, v).$$

Consequently,

$$|\mathcal{I}_1| \lesssim \frac{1}{t} \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} \langle y \rangle^{N_x-3} (|v^0|^7 |g|(t, y, v) + |v^0|^8 |\nabla_v g|(t, y, v)) \int_{|x-y|<t} \frac{dy}{\langle y \rangle^{N_x-4}}, \quad N_x - 4 > 3.$$

In order to bound \mathcal{I}_2 recall that $|x| < t$ and note that, for $v = \widetilde{x}/t$ and any $y \in \mathbb{R}$ such that $|y-x| \geq t$,

$$1 = |v^0|^2 \left(1 - \frac{|x|^2}{t^2}\right) \leq |v^0|^2 \frac{|y|(t+|x|)}{t^2} \leq 2 \frac{|y||v^0|^2}{t}.$$

We then finally deduce that

$$|\mathcal{I}_2| \leq \frac{2}{t} \int_{|y-x|\geq t} (|v^0|^7 g)\left(t, y, \frac{\check{x}}{t}\right) \frac{\langle y \rangle^{N_x-3}}{\langle y \rangle^{N_x-4}} \, dy \leq \frac{4}{t} \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |v^0|^7 \langle y \rangle^{N_x-3} |g|(t, y, v). \quad \square$$

We are able to prove that the decay of quantities such as $\int_v \widehat{Z}^\beta f \, dv$ is optimal. We state a general result since we will later have to deal with integral kernels.

Proposition 4.13. *Let $|\beta| \leq N - 1$ and $\Psi : \mathbb{S}_\omega^2 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a sufficiently regular function such that $\|\Psi(\cdot, v)\|_{L^\infty} + \|v^0 \nabla_v \Psi(\cdot, v)\|_{L^\infty} \lesssim |v^0|^{N_v-11}$. Then, for any $\omega \in \mathbb{S}^2$,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}_x^3, \quad \left| \int_{\mathbb{R}_v^3} \Psi(\omega, v) \widehat{Z}^\beta f(t, x, v) \, dv \right| \lesssim \frac{\bar{\epsilon}}{(1+t+|x|)^3}.$$

Proof. Assume first that $|x| \leq t \leq 1$ or $|x| \geq t$. Then, as $|\Psi|(\cdot, v) \lesssim |v^0|^{N_v-9}$, it suffices to use the first or the third estimate of Proposition 4.11, applied for $a = \frac{1}{2}$. Otherwise $t > \max(1, |x|)$ and we introduce, for any $\omega \in \mathbb{S}^2$, $g_\omega(t, x, v) = \Psi(\omega, v) \widehat{Z}^\beta f(t, x + t\hat{v}, v)$. Using first Lemma 2.8 and then Proposition 4.5, we have

$$\begin{aligned} & \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} \langle y \rangle^{N_x-3} (|v^0|^7 |g_\omega| + |v^0|^8 |\nabla_v g_\omega|)(t, y, v) \\ & \lesssim \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |\nabla_v \Psi|(\omega, v) |v^0|^8 |z^{N_x-3} \widehat{Z}^\beta f|(t, y, v) + \sum_{|\kappa| \leq 1} |\Psi|(\omega, v) |v^0|^7 |z^{N_x-2} \widehat{Z}^\kappa \widehat{Z}^\beta f|(t, y, v) \\ & \lesssim \sum_{|\xi| \leq N} \sup_{(y,v) \in \mathbb{R}^3 \times \mathbb{R}^3} |v^0|^{N_v-3} |z^{N_x-2} \widehat{Z}^\xi f|(t, y, v) \lesssim \bar{\epsilon} \log^{3N_x+3N} (3+t). \end{aligned} \tag{41}$$

Now, apply [Lemma 4.12](#) to g_ω in order to get,

$$\forall \omega \in \mathbb{S}^2, \quad t^3 \left| \int_{\mathbb{R}_v^3} \Psi(\omega, v) \widehat{Z}^\beta f(t, x, v) \, dv \right| \lesssim \left| \int_{\mathbb{R}_y^3} (|v^0|^5 g_\omega) \left(t, y, \frac{\check{x}}{t} \right) \, dy \right| + \bar{\epsilon} \frac{\log^{3N_x+3N} (3+t)}{t}.$$

As $t \geq 1$, it remains to bound by $\bar{\epsilon}$ the first term on the right-hand side. For this, perform the change of variables $z = y - t\hat{v}$ and apply [Corollary 4.9](#) with $\psi(\omega, v) = |v^0|^5 \Psi(\omega, v)$. \square

The next result is a direct application of the previous proposition to $\Psi(\omega, v) = v^\mu/v^0$ for any $0 \leq \mu \leq 3$.

Corollary 4.14. *For any $|\beta| \leq N - 1$, the decay of the current density $J(\widehat{Z}^\beta f)$ is optimal. There exists a constant $C > 0$ independent of ϵ such that,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}_x^3, \quad \left| \int_{\mathbb{R}_v^3} \frac{v^\mu}{v^0} \widehat{Z}^\beta f(t, x, v) \, dv \right| \leq \frac{C\bar{\epsilon}}{(1+t+|x|)^3}, \quad 0 \leq \mu \leq 3.$$

If ϵ satisfies $C\bar{\epsilon} = C\epsilon e^{(D+1)\Lambda} < C_{\text{boot}}\Lambda$, it improves the bootstrap assumption [\(BA3\)](#).

4.5. Improved estimates for derivatives of velocity averages. In the linear case, derivatives of averages in v , such as $\partial_{t,x} \int_v f \, dv$, enjoy stronger decay properties. Our study of the top-order derivatives of the electromagnetic field will require the following improved estimates.

Proposition 4.15. *Let $|\beta| \leq N - 1$, $\mu \in \llbracket 0, 3 \rrbracket$ and $\Phi : \mathbb{S}^2 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a sufficiently regular function such that $\|\Phi(\cdot, v)\|_{L^\infty_\omega} + \|v^0 \nabla_v \Phi(\cdot, v)\|_{L^\infty_\omega} \lesssim |v^0|^{N_v-10}$. Then, for any $\omega \in \mathbb{S}^2$,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}_x^3, \quad \left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) \partial_{x^\mu} \widehat{Z}^\beta f(t, x, v) \, dv \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N} (3+t)}{(1+t+|x|)^4}.$$

Proof. Let $(t, x) \in [0, T[\times \mathbb{R}_x^3$ and note that, if $|x| \geq t - 1$, the result is given by [Proposition 4.11](#), applied for $a = 1$. We then consider the case $t - |x| \geq 1$. Using [\(20\)](#) together with $t\Omega_{ij} = (x^i/r)\Omega_{0j} - (x^j/r)\Omega_{0i}$, one has

$$\mathcal{I}_{t,x}^\beta := |t - |x|| \left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) \partial_{x^\mu} \widehat{Z}^\beta f(t, x, v) \, dv \right| \leq \sum_{Z \in \mathbb{K}} \left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) Z \widehat{Z}^\beta f(t, x, v) \, dv \right|.$$

Fix now $Z \in \mathbb{K}$. If Z is a translation ∂_{x^μ} or if $Z = S$, then $Z \in \widehat{\mathbb{P}}_0$. Otherwise, either $Z = \Omega_{ij}$ is a rotation and $Z = \widehat{Z} - v^i \partial_{v_j} + v^j \partial_{v_i}$ or $Z = \Omega_{0k}$ is a Lorentz boost and $Z = \widehat{Z} - v^0 \partial_{v^k}$, so that

$$\mathcal{I}_{t,x}^\beta \leq \sum_{\widehat{Z} \in \widehat{\mathbb{P}}_0} \left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) \widehat{Z} \widehat{Z}^\beta f(t, x, v) \, dv \right| + \sum_{\lambda=0}^3 \sum_{k=1}^3 \left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) v^\lambda \partial_{v^k} \widehat{Z}^\beta f(t, x, v) \, dv \right|.$$

Integration by parts and $|\partial_{v^k}(\Phi(\omega, v)v^\lambda)| \leq v^0 |\nabla_v \Phi|(\omega, v) + |\Phi|(\omega, v) \lesssim |v^0|^{N_v-10}$ yield

$$\left| \int_{\mathbb{R}_v^3} \Phi(\omega, v) \partial_{x^\mu} \widehat{Z}^\beta f(t, x, v) \, dv \right| \lesssim \frac{1}{|t - |x||} \sum_{|\kappa| \leq 1} \int_{\mathbb{R}_v^3} |v^0|^{N_v-10} |\widehat{Z}^\kappa \widehat{Z}^\beta f|(t, x, v) \, dv.$$

As $t - |x| \geq 1$, it remains to apply once again [Proposition 4.11](#) for $a = 1$. \square

5. Improvement of the bootstrap assumptions on the electromagnetic field

We are now able to prove pointwise decay estimates for the Maxwell field and its derivatives. We improve first (BA1), whereas the case of the top-order derivatives (BA2) will require a different strategy since we did not recover the linear decay t^{-3} for $\int_v \widehat{Z}^\beta f(t, x, v) dv$, $|\beta| = N$.

5.1. The Glassey–Strauss decomposition of the electromagnetic field. We separate F as well as its derivatives $\mathcal{L}_{Z^\gamma}(F)$ into three parts according to the Glassey–Strauss decomposition. For this, recall from (4) the relation between the electric field E , the magnetic field B and the Faraday tensor F . We have $E^i = F_{0i}$, $B^1 = F_{32}$, $B^2 = F_{13}$ and $B^3 = F_{21}$. To simplify the statement of the decomposition, we introduce a weight tensor field.

Definition 5.1. Let $w_{\mu\nu}(\omega, v)$ be the antisymmetric tensor defined for all $(\omega, v) \in \mathbb{S}^2 \times \mathbb{R}_v^3$ by

$$w_{0i}(\omega, v) = -w_{i0}(\omega, v) := \omega_i + \hat{v}_i, \quad w_{jk}(\omega, v) := \omega_j \hat{v}_k - \omega_k \hat{v}_j, \quad 1 \leq i, j, k \leq 3,$$

where $\omega_i := x_i/|x|$ if $x \in \mathbb{R}^3$ satisfies $\omega = x/|x|$. We further define

$$W_{\mu\nu}(\omega, v) := \frac{w_{\mu\nu}(\omega, v)}{1 + \omega \cdot \hat{v}}.$$

Remark 5.2. Since w is antisymmetric, $w_{\mu\mu} = 0$ for any $\mu \in \llbracket 0, 3 \rrbracket$. Note also that $1 + \omega \cdot \hat{v} = 2v^L > 0$.

We can now prove an adaptation of [Glassey and Strauss 1986, Theorem 3]. The key idea of their proof consists in rewriting the standard derivatives $\partial_{t,x}$ as combinations of derivatives tangential to a backward light cone, which naturally appears in the representation formula for solutions to wave equations, and $T_0 := \partial_t + \hat{v} \cdot \nabla_x$, the free relativistic transport operator which is transverse to light cones. To avoid any confusion with the scaling vector field, we do not keep the notation S , used by Glassey and Strauss, in order to denote T_0 .

Proposition 5.3. Let $|\gamma| \leq N - 1$. Then, there exist $C_\beta^\gamma, N_{\xi,\kappa}^\gamma \in \mathbb{N}$ such that

$$4\pi \mathcal{L}_{Z^\gamma}(F) = \mathcal{L}_{Z^\gamma}(F)^{\text{data}} + \mathcal{L}_{Z^\gamma}(F)^T + \mathcal{L}_{Z^\gamma}(F)^S,$$

where, for any $0 \leq \mu, \nu \leq 3$ and with $\omega = (y - x)/|y - x|$ in the following integrals:

- $\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{data}}$ can be explicitly computed in terms of the initial data. More precisely,

$$\begin{aligned} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{data}}(t, x) &= 4\pi \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}}(t, x) - \sum_{|\beta| \leq |\gamma|} \frac{C_\beta^\gamma}{t} \int_{|y-x|=t} \int_{\mathbb{R}_v^3} (W_{\mu\nu}(\omega, v) - \delta_\mu^0 \hat{v}^\nu + \delta_\nu^0 \hat{v}^\mu) \widehat{Z}^\beta f(0, y, v) dv dy \end{aligned}$$

and $\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}}$ is the unique solution to the homogeneous wave equation $\square \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}} = 0$ which initially verifies $\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}}(0, \cdot) = \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(0, \cdot)$ and $\partial_t \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}}(0, \cdot) = \partial_t \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(0, \cdot)$.

- The 2-form $\mathcal{L}_{Z^\gamma}(F)^T$ is given by

$$\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^T(t, x) := - \sum_{|\beta| \leq |\gamma|} C_\beta^\gamma [\widehat{Z}^\beta f]_{\mu\nu}^T(t, x),$$

where the field $[\widehat{Z}^\beta f]^T$ generated by $\widehat{Z}^\beta f$ is

$$[\widehat{Z}^\beta f]_{\mu\nu}^T(t, x) := \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \frac{\mathcal{W}_{\mu\nu}(\omega, v)}{|v^0|^2(1 + \omega \cdot \hat{v})} \widehat{Z}^\beta f(t - |y - x|, y, v) \frac{dv dy}{|y - x|^2}.$$

• The 2-form $\mathcal{L}_{Z^\gamma}(F)^S$ is defined by

$$\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^S(t, x) := \sum_{|\xi|+|\kappa| \leq |\gamma|} N_{\xi, \kappa}^\gamma \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} (\widehat{Z}^\kappa f \hat{v}^\lambda \mathcal{L}_{Z^\xi}(F)_{\lambda^j})(t - |y - x|, y, v) \partial_{v^j} \mathcal{W}_{\mu\nu}(\omega, v) \frac{dv dy}{|y - x|}.$$

Proof. Fix $|\gamma| \leq N - 1$ and apply Proposition 2.4 in order to rewrite the Maxwell equations satisfied by $\mathcal{L}_{Z^\gamma}(F)$ as

$$\nabla^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} = \int_{\mathbb{R}_v^3} \frac{v_\nu}{v^0} f_\gamma(t, x, v) dv, \quad \nabla^{\mu*} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} = 0, \quad v \in \llbracket 0, 3 \rrbracket, \quad f_\gamma := \sum_{|\beta| \leq |\gamma|} C_\beta^\gamma \widehat{Z}^\beta f, \quad (42)$$

with $C_\beta^\gamma \in \mathbb{N}$. Introduce further the electric E_γ and magnetic B_γ parts of $\mathcal{L}_{Z^\gamma}(F)$,

$$E_\gamma^i := \mathcal{L}_{Z^\gamma}(F)_{0i}, \quad i \in \llbracket 1, 3 \rrbracket, \quad B_\gamma^1 = \mathcal{L}_{Z^\gamma}(F)_{32}, \quad B_\gamma^2 = \mathcal{L}_{Z^\gamma}(F)_{13}, \quad B_\gamma^3 = \mathcal{L}_{Z^\gamma}(F)_{21}. \quad (43)$$

In such a way, our framework exactly corresponds to the one of Glassey and Strauss. More precisely, one can compute the source terms of the wave equations satisfied by the components of E_γ and B_γ . For any $0 \leq \mu, \nu \leq 3$, we have

$$\square \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} = \int_{\mathbb{R}_v^3} \hat{v}_\mu \partial_{x^\nu} f_\gamma - \hat{v}_\nu \partial_{x^\mu} f_\gamma dv, \quad \text{so, for instance,} \quad \square E_\gamma^i = - \int_{\mathbb{R}_v^3} \partial_{x^i} f_\gamma + \hat{v}_i \partial_t f_\gamma dv.$$

Applying [Glassey and Strauss 1986, Theorem 3] to $(f_\gamma, E_\gamma, B_\gamma)$ provides us, for any $0 \leq \mu, \nu \leq 3$,

$$4\pi \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} = \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{data}} + \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^T - \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \mathcal{W}_{\mu\nu}(\omega, v) (T_0 f_\gamma)(t - |y - x|, y, v) \frac{dv dy}{|y - x|},$$

where we recall that $T_0 = \hat{v}^\lambda \partial_{x^\lambda}$. Note that the constants C_β^γ in the definitions of $\mathcal{L}_{Z^\gamma}(F)^{\text{data}}$, $\mathcal{L}_{Z^\gamma}(F)^T$ and f_γ are the same. Applying the commutation formula of Proposition 2.4 for any $|\beta| \leq |\gamma|$ yields

$$T_0 f_\gamma = - \sum_{|\beta| \leq |\gamma|} C_\beta^\gamma \hat{v}^\mu F_{\mu^j} \partial_{v^j} \widehat{Z}^\beta f + C_\beta^\gamma \sum_{|\xi|+|\kappa| \leq |\beta|} C_{\xi, \kappa}^\beta \hat{v}^\mu \mathcal{L}_{Z^\xi}(F)_{\mu^j} \partial_{v^j} \widehat{Z}^\kappa f, \quad (44)$$

with $C_{\xi, \kappa}^\beta \in \mathbb{N}$. It remains to integrate by parts in v and to recall that $\nabla_{v^j} \cdot \hat{v}^\mu \mathcal{L}_{Z^\xi}(F)_{\mu^j} = \mathcal{L}_{Z^\xi}(F)_{j^j} = 0$. \square

It will then be important to estimate the kernels introduced in the previous proposition.

Lemma 5.4. For all $(\omega, v) \in \mathbb{S}^2 \times \mathbb{R}_v^3$, we have $|\omega + \hat{v}|^2$, $|\omega \wedge \hat{v}|^2 \leq 2(1 + \omega \cdot \hat{v})$ and $(1 + \omega \cdot \hat{v})^{-1} \leq 2|v^0|^2$.

Proof. For the second inequality, simply note that

$$2|v^0|^2(1 + \omega \cdot \hat{v}) \geq 2|v^0|^2(1 - |\hat{v}|) = 2v^0(v^0 - |v|) \geq (v^0 + |v|)(v^0 - |v|) = |v^0|^2 - |v|^2 = 1.$$

For the first ones, since $|\omega| = 1$ and $|\hat{v}| \leq 1$,

$$|\omega + \hat{v}|^2 = |\omega|^2 + |\hat{v}|^2 + 2\omega \cdot \hat{v} \leq 2(1 + \omega \cdot \hat{v}),$$

$$|\omega \wedge \hat{v}|^2 = |\omega|^2 |\hat{v}|^2 - |\omega \cdot \hat{v}|^2 \leq (1 + \omega \cdot \hat{v})(1 - \omega \cdot \hat{v}) \leq 2(1 + \omega \cdot \hat{v}). \quad \square$$

Corollary 5.5. *For any $0 \leq \mu, \nu \leq 3$ and all $(\omega, v) \in \mathbb{S}^2 \times \mathbb{R}_v^3$, there holds*

$$|\mathcal{W}_{\mu\nu}|(\omega, v) \leq 2v^0, \quad \frac{|\mathcal{W}_{\mu\nu}|(\omega, v)}{|v^0|^2(1 + \omega \cdot \hat{v})} \leq 4v^0, \quad |\nabla_v \mathcal{W}_{\mu\nu}|(\omega, v) \leq 6v^0.$$

We have similar bounds for their first-order derivatives,

$$|\nabla_v \mathcal{W}_{\mu\nu}|(\omega, v) \lesssim v^0, \quad \left| \nabla_v \left(\frac{\mathcal{W}_{\mu\nu}(\omega, v)}{|v^0|^2(1 + \omega \cdot \hat{v})} \right) \right| \lesssim v^0, \quad |\nabla_v \nabla_v \mathcal{W}_{\mu\nu}|(\omega, v) \lesssim v^0.$$

Proof. The first two inequalities are a direct consequence of the previous lemma. The other ones ensue from straightforward computations carried out in Lemma A.2. \square

Remark 5.6. These bounds are sharp. Let us focus for instance on the first one, $|\mathcal{W}_{\mu\nu}|(\omega, v) \leq 2v^0$. For this, consider, for any $v \in \mathbb{R}_v^3$, the function $\phi_v : \omega \mapsto 1 + \omega \cdot \hat{v}$ defined on \mathbb{S}^2 . Then,

$$\min_{\omega \in \mathbb{S}^2} \phi_v(\omega) = \frac{v^0 - |v|}{v^0} = \frac{1}{v^0(v^0 + |v|)} \leq \frac{1}{|v^0|^2}, \quad \max_{\omega \in \mathbb{S}^2} \phi_v(\omega) = \frac{v^0 + |v|}{v^0} \geq 1.$$

By continuity, there exists $\omega_v \in \mathbb{S}^2$ such that $1 + \omega_v \cdot \hat{v} = |v^0|^{-2}$. Then, using $|\omega + \hat{v}|^2 = 2(1 + \omega \cdot \hat{v}) - |v^0|^{-2}$, we have

$$\sum_{1 \leq i \leq 3} |\mathcal{W}_{0i}|^2(\omega_v, v) = \frac{|\omega_v + \hat{v}|^2}{|1 + \omega_v \cdot \hat{v}|^2} = \frac{1}{1 + \omega_v \cdot \hat{v}} \left(2 - \frac{1}{|v^0|^2(1 + \omega_v \cdot \hat{v})} \right) = v^0.$$

In order to improve the bootstrap assumption (BA2), we will need to use the Glassey–Strauss decomposition of the spatial derivatives of the electromagnetic field. A similar result holds for the time derivative but we will estimate it by exploiting the Maxwell equations. For instance, one can check that (2)–(3) imply $|\nabla_{\partial_t} F| \lesssim \sum_{1 \leq k \leq 3} |\nabla_{\partial_{x^k}} F| + |J(f)|$. We lighten the notations by denoting the Lorentz force as

$$K^j := \hat{v}^\mu F_{\mu}^j, \quad K_\xi^j := \hat{v}^\mu \mathcal{L}_{Z^\xi}(F)_{\mu}^j, \quad 1 \leq j \leq 3, \quad 1 \leq |\xi| \leq N. \tag{45}$$

Proposition 5.7. *Let $|\gamma| = N - 1$ and $1 \leq k \leq 3$. Then, $\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\gamma}(F)$ can be written as*

$$4\pi \nabla_{\partial_{x^k}} \mathcal{L}_{Z^\gamma}(F) = A_{\gamma,k}^{\text{data}} + A_{\gamma,k}^{\text{ver}} + A_{\gamma,k}^{TT} + A_{\gamma,k}^{TS} + A_{\gamma,k}^{SS},$$

where the five 2-forms satisfy the following properties. We fix $0 \leq \mu, \nu \leq 3$ and we use again the notation $\omega = (y - x)/|y - x|$ in the integrals written below. Moreover, we give the definition of the kernels at the end of the statement.⁹

- $A_{\gamma,k}^{\text{data}}$ can be explicitly computed in terms of the initial data,

$$\begin{aligned} A_{\gamma,k,\mu\nu}^{\text{data}}(t, x) &= 4\pi \partial_{x^k} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{data}}(t, x) - \sum_{|\beta| \leq N-1} C_\beta^\gamma \frac{1}{t^2} \int_{|y-x|=t} \int_{\mathbb{R}_v^3} \mathcal{D}_{\mu\nu}^k(\omega, v) \widehat{Z}^\beta f(0, y, v) \, dv \, dy \\ &\quad - \sum_{|\beta| \leq N-1} C_\beta^\gamma \frac{1}{t} \int_{|y-x|=t} \int_{\mathbb{R}_v^3} C_{\mu\nu}^k(\omega, v) T_0 \widehat{Z}^\beta f(0, y, v) \, dv \, dy. \end{aligned}$$

⁹We point out that we are only interested in the qualitative properties of these kernels.

- $A_{\gamma,k}^{\text{ver}}$ is the vertex term,

$$A_{\gamma,k,\mu\nu}^{\text{ver}}(t, x) := \sum_{|\beta| \leq N-1} C_{\beta}^{\gamma} \int_{\sigma \in \mathbb{S}^2} \int_{\mathbb{R}_v^3} \mathcal{D}_{\mu\nu}^k(\sigma, v) \widehat{Z}^{\beta} f(t, x, v) \, dv \, d\mu_{\mathbb{S}^2}.$$

- $A_{\gamma,k}^{TT}$ is the most singular term,

$$A_{\gamma,k,\mu\nu}^{TT}(t, x) := \sum_{|\beta| \leq N-1} C_{\beta}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \mathcal{A}_{\mu\nu}^k(\omega, v) \widehat{Z}^{\beta} f(t - |y-x|, y, v) \frac{dv \, dy}{|y-x|^3}$$

and the crucial identity $\int_{|\sigma|=1} \mathcal{A}_{\mu\nu}^k(\sigma, \hat{v}) \, d\mu_{\mathbb{S}^2} = 0$ holds for all $v \in \mathbb{R}_v^3$.

- $A_{\gamma,k}^{T,S}$ is given by

$$A_{\gamma,k,\mu\nu}^{T,S}(t, x) := \sum_{|\xi|+|\kappa| \leq N-1} N_{\xi,\kappa}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \nabla_v \mathcal{B}_{\mu\nu}^k(\omega, v) \cdot (\widehat{Z}^{\kappa} f K_{\xi})(t - |y-x|, y, v) \frac{dv \, dy}{|y-x|^2}.$$

- $A_{\gamma,k}^{SS}$ is the sum of the four following quantities, where $N_{\xi,\zeta,\kappa}^{\gamma} \in \mathbb{N}$,

$$A_{\gamma,k,\mu\nu}^{SS,I} := \sum_{|\xi|+|\zeta|+|\kappa| \leq N-1} N_{\xi,\zeta,\kappa}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} [\nabla_v (\nabla_v \mathcal{C}_{\mu\nu}^k(\omega, \cdot) \cdot K_{\xi}) \cdot K_{\zeta} \widehat{Z}^{\kappa} f](t - |y-x|, y, v) \frac{dv \, dy}{|y-x|},$$

$$A_{\gamma,k,\mu\nu}^{SS,II} := \sum_{|\xi|+|\kappa| \leq N-1} N_{\xi,\kappa}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \nabla_v \mathcal{C}_{\mu\nu}^k(\omega, v) \cdot (\mathbf{T}_0(K_{\xi}) \widehat{Z}^{\kappa} f)(t - |y-x|, y, v) \frac{dv \, dy}{|y-x|},$$

$$A_{\gamma,k,\mu\nu}^{SS,III} := \sum_{|\xi|+|\kappa| \leq N-1} N_{\xi,\kappa}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \mathcal{C}_{\mu\nu}^k(\omega, v) \frac{\delta_j^n - \hat{v}_j \hat{v}^n}{v^0} (\partial_{x^n} (K_{\xi}^j) \widehat{Z}^{\kappa} f)(t - |y-x|, y, v) \frac{dv \, dy}{|y-x|},$$

$$A_{\gamma,k,\mu\nu}^{SS,IV} := \sum_{|\xi|+|\kappa| \leq N-1} N_{\xi,\kappa}^{\gamma} \int_{|y-x| \leq t} \int_{\mathbb{R}_v^3} \mathcal{C}_{\mu\nu}^k(\omega, v) \frac{\delta_j^n - \hat{v}_j \hat{v}^n}{v^0} (K_{\xi}^j \partial_{x^n} \widehat{Z}^{\kappa} f)(t - |y-x|, y, v) \frac{dv \, dy}{|y-x|}.$$

- The kernels are smooth functions of $(\omega, v) \in \mathbb{S}^2 \times \mathbb{R}_v^3$ given by

$$\mathcal{A}_{\mu\nu}^k(\omega, v) := -3 \frac{\mathbf{w}_{\mu\nu}(\omega, v) \omega_k}{|v^0|^4 (1 + \omega \cdot \hat{v})^4} - 3 \frac{\mathbf{w}_{\mu\nu}(\omega, v) \hat{v}_k}{|v^0|^2 (1 + \omega \cdot \hat{v})^3} + \frac{\delta_{k\mu} \hat{v}_\nu - \delta_{k\nu} \hat{v}_\mu}{|v^0|^2 (1 + \omega \cdot \hat{v})^2},$$

$$\mathcal{B}_{\mu\nu}^k(\omega, v) := 3 \frac{\mathbf{w}_{\mu\nu}(\omega, v) \omega_k}{|v^0|^2 (1 + \omega \cdot \hat{v})^3} - 2 \frac{\mathbf{w}_{\mu\nu}(\omega, v) \hat{v}_k}{(1 + \omega \cdot \hat{v})^2} - \frac{\delta_{k\mu} \hat{v}_\nu - \delta_{k\nu} \hat{v}_\mu}{1 + \omega \cdot \hat{v}},$$

$$\mathcal{C}_{\mu\nu}^k(\omega, v) := \frac{\omega_k \mathbf{w}_{\mu\nu}(\omega, v)}{(1 + \omega \cdot \hat{v})^2}, \quad \mathcal{D}_{\mu\nu}^k(\omega, v) := \frac{\omega_k \mathbf{w}_{\mu\nu}(\omega, v)}{|v^0|^2 (1 + \omega \cdot \hat{v})^3}.$$

Proof. Let $k \in \llbracket 1, 3 \rrbracket$, $|\gamma| = N - 1$ and recall from (42) the definition of f_{γ} and that $\mathcal{L}_{Z^{\gamma}}(F)$ solves the Maxwell equations with source term $J(f_{\gamma})$. Recall further the electric and magnetic parts (E_{γ}, B_{γ}) of $\mathcal{L}_{Z^{\gamma}}(F)$, introduced in (43). In the same spirit as in the proof of Proposition 5.3, we apply¹⁰ [Glassey 1996, Theorem 5.4.1] to $(f_{\gamma}, E_{\gamma}, B_{\gamma})$. This yields

$$\nabla_{\partial_{x^k}} \mathcal{L}_{Z^{\gamma}}(F)_{\mu\nu} = A_{\gamma,k,\mu\nu}^{\text{data}} + A_{\gamma,k,\mu\nu}^{\text{ver}} + A_{\gamma,k,\mu\nu}^{TT} + \tilde{A}_{\gamma,k,\mu\nu}^{T,S} + \tilde{A}_{\gamma,k,\mu\nu}^{SS},$$

¹⁰See also the original version of the result, [Glassey and Strauss 1986, Theorem 4].

where

$$\begin{aligned} \tilde{A}_{\gamma,k,\mu\nu}^{T,S} &:= \int_{|y-x|\leq t} \int_{\mathbb{R}_v^3} \mathcal{B}_{\mu\nu}^k(\omega, v)(\mathbf{T}_0 f_\gamma)(t - |y - x|, y, v) \frac{dv dy}{|y - x|^2}, \\ \tilde{A}_{\gamma,k,\mu\nu}^{SS} &:= - \int_{|y-x|\leq t} \int_{\mathbb{R}_v^3} \mathcal{C}_{\mu\nu}^k(\omega, v)(\mathbf{T}_0 \mathbf{T}_0 f_\gamma)(t - |y - x|, y, v) \frac{dv dy}{|y - x|}, \end{aligned}$$

as well as $\int_{|\sigma|=1} A_{\mu\nu}^k(\sigma, \hat{v}) d\mu_{\mathbb{S}^2} = 0$. One can then prove that $\tilde{A}_{\gamma,k,\mu\nu}^{T,S} = A_{\gamma,k,\mu\nu}^{T,S}$ by rewriting $\mathbf{T}_0 f_\gamma$ using the (commuted) Vlasov equation. More precisely, we use (44) and we then integrate by parts in v . It remains to deal with $\tilde{A}_{\gamma,k,\mu\nu}^{SS}$ and we recall for this that $\nabla_v \cdot K_\xi = \nabla_{v_j} \cdot \hat{v}^\mu \mathcal{L}_{Z^\xi}(F)_\mu^j = 0$. Hence, using again (44), we get that there exists $N_{\xi,\kappa}^\gamma \in \mathbb{N}$ such that

$$\mathbf{T}_0 \mathbf{T}_0(f_\gamma) = \sum_{|\xi|+|\kappa|\leq|\gamma|} N_{\xi,\kappa}^\gamma \mathbf{T}_0 \partial_{v^j} (K_\xi^j \widehat{Z}^\kappa f).$$

Now, we write $\mathbf{T}_0 \partial_{v^j} = \partial_{v^j} \mathbf{T}_0 - \partial_{v^j}(\hat{v}^n) \partial_{x^n}$ and we apply the commutation formula of Proposition 2.4 to $\widehat{Z}^\kappa f$. We get

$$\mathbf{T}_0 \partial_{v^j} (\hat{v}^\lambda \mathcal{L}_{Z^\xi}(F)_\lambda^j \widehat{Z}^\kappa f) = \partial_{v^j} (\mathbf{T}_0 (K_\xi^j \widehat{Z}^\kappa f)) + \partial_{v^j} (K_\xi^j \mathbf{T}_0 (\widehat{Z}^\kappa f)) - \frac{\delta_j^n - \hat{v}_j \hat{v}^n}{v_0} (\partial_{x^n} (K_\xi^j \widehat{Z}^\kappa f) + K_\xi^j \partial_{x^n} \widehat{Z}^\kappa f),$$

so that, by integration by parts in v for the quantities related to the two first terms on the right-hand side of the previous equality,

$$\begin{aligned} \tilde{A}_{\gamma,k,\mu\nu}^{SS} &= A_{\gamma,k,\mu\nu}^{SS,II} + A_{\gamma,k,\mu\nu}^{SS,III} + A_{\gamma,k,\mu\nu}^{SS,IV} \\ &\quad + \sum_{|\xi|+|\kappa|\leq|\gamma|} N_{\xi,\kappa}^\gamma \int_{|y-x|\leq t} \int_{\mathbb{R}_v^3} \nabla_{v^j} (\mathcal{C}_{\mu\nu}^k(\omega, v))(K_\xi^j \mathbf{T}_0 (\widehat{Z}^\kappa f))(\tau_y, y, v) \frac{dv dy}{|y - x|}, \end{aligned}$$

where $\tau_y := t - |y - x|$. Finally, we deal with the last term by applying first the commutation relation of Proposition 2.4, giving that $\mathbf{T}_0 (\widehat{Z}^\kappa f) = -K \cdot \nabla_v \widehat{Z}^\kappa f + C_{\xi,\beta}^\kappa K_\xi \cdot \nabla_v \widehat{Z}^\beta f$, and then by integrating by parts in v . □

These kernels and their derivatives can be estimated by a direct application of Lemmas 5.4 and A.2.

Corollary 5.8. *For any $1 \leq k, j, n \leq 3$ and for all $v \in \mathbb{R}_v^3$, we have*

$$(|\mathcal{A}^k| + |\nabla_v \mathcal{A}^k| + |\nabla_v \mathcal{B}^k| + |\mathcal{C}^k| + |\nabla_v \mathcal{C}^k| + |\nabla_v \nabla_v \mathcal{C}^k| + |\mathcal{D}^k| + |\nabla_v \mathcal{D}^k|)(\cdot, v) \lesssim |v^0|^3.$$

5.2. Three integral bounds. The estimate of most of the terms listed in Propositions 5.3 and 5.7 will in fact be reduced to the analysis of three different integrals. We will deal with all of them by applying a particular case of [Glassey 1996, Lemma 6.5.2].

Lemma 5.9. *Let $p \in \mathbb{R}$ and $g : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ be a continuous function. Then, for all $(t, x) \in [0, T] \times \mathbb{R}^3 \setminus \{0\}$,*

$$\int_{|y-x|\leq t} g(t - |y - x|, |y|) \frac{dy}{|y - x|^p} = \frac{2\pi}{|x|} \int_{\tau=0}^t \int_{\lambda=||x|-t+\tau}^{|x|+t-\tau} g(\tau, \lambda) \lambda d\lambda \frac{d\tau}{(t - \tau)^{p-1}}.$$

The following result will be useful for controlling $\mathcal{L}_{Z^\gamma}(F)^S$ and $A_{\gamma,k}^{SS}$.

Lemma 5.10. *For any $b \geq 4$ and for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$, there holds*

$$Y_{b,1}^{p=1}(t, x) := \int_{|y-x| \leq t} \frac{1}{(1+t-|y-x|+|y|)^b (1+|t-|y-x|-|y|)|y-x|} dy$$

$$\lesssim \frac{\log(3+|t-|x||)}{(1+t+|x|)(1+|t-|x||)^{b-2}}.$$

Proof. Note first that, on the domain of integration,

$$t - |y - x| + |y| \geq t - |y| - |x| + |y| = t - |x|, \quad t - |y - x| + |y| \geq |y| \geq |x| - |y - x| \geq |x| - t, \quad (46)$$

so that $Y_{b,1}^{p=1}(t, x) \leq (1+|t-|x||)^{-b+4} Y_{4,1}^{p=4}(t, x)$ and it suffices to treat the case $b = 4$. By continuity, we can assume further that $x \neq 0$. According to Lemma 5.9,

$$Y_{4,1}^{p=1}(t, x) \leq \frac{2\pi}{|x|} \int_{\tau=0}^t \int_{\lambda=|x|-t+\tau}^{|x|+t-\tau} \frac{d\lambda d\tau}{(1+\tau+\lambda)^3 (1+|\tau-\lambda|)}.$$

We perform the change of variables $\underline{u} = \tau + \lambda$ and $u = \tau - \lambda$. Then, on the domain of integration $||x| - t| \leq \underline{u} \leq t + |x|$ and $u \leq ||x| - t|$. Moreover, $u \geq -\underline{u}$ since $2\tau \geq 0$. Consequently,

$$Y_{4,1}^{p=1}(t, x) \leq \frac{\pi}{|x|} \int_{\underline{u}=||x|-t|}^{t+|x|} \int_{u=-\underline{u}}^{||x|-t|} \frac{du}{1+|u|} \frac{d\underline{u}}{(1+\underline{u})^3} \leq \frac{2\pi}{|x|} \int_{\underline{u}=||x|-t|}^{t+|x|} \frac{\log(3+\underline{u})}{(1+\underline{u})^3} d\underline{u}.$$

Now, note that

$$Y_{4,1}^{p=1}(t, x) \lesssim \frac{2\pi \log(3+|t-|x||)}{(1+|t-|x||)|x|} \int_{\underline{u}=||x|-t|}^{t+|x|} \frac{d\underline{u}}{(1+\underline{u})^2}$$

$$= \frac{2\pi \log(3+|t-|x||)}{(1+t+|x|)(1+|t-|x||)^2} \frac{t+|x|-|t-|x||}{|x|}$$

and it remains to note that the last factor on the right-hand side is bounded by $2 \min(t, |x|)/|x| \leq 2$. \square

We will apply the next lemma in order to deal with $\mathcal{L}_{Z^{\gamma}}(F)^T$ and $A_{\gamma,k}^{T,S}$.

Lemma 5.11. *Let, for any $b \geq 3$ and all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,*

$$Y_b^{p=2}(t, x) := \int_{|y-x| \leq t} (1+t-|y-x|+|y|)^{-b} \frac{dy}{|y-x|^2}.$$

Then, the following range of estimates holds. For any $0 < \delta \leq 1$,

$$Y_b^{p=2}(t, x) \lesssim \delta^{-1} (1+t+|x|)^{-2+\delta} (1+|t-|x||)^{-b-\delta+3},$$

$$Y_b^{p=2}(t, x) \lesssim (1+t+|x|)^{-2} (1+|t-|x||)^{-b+3} \log(1+t).$$

Proof. In view of (46), we have $Y_b^{p=2}(t, x) \leq (1+|t-|x||)^{-b+3} Y_3^{p=2}(t, x)$ and it suffices to treat the case $b = 3$. Then note that

$$Y_3^{p=2}(t, x) = K_{[0, \frac{t}{2}]} + K_{[\frac{t}{2}, t]}, \quad K_I := \int_{|y-x| \in I} (1+t-|y-x|+|y|)^{-3} \frac{dy}{|y-x|^2}.$$

On the domain of integration of $\mathbf{K}_{[0,t/2]}$, we have $t - |y - x| + |y| \gtrsim t + |x|$. Indeed, $t - |y - x| \geq t/2$ and $|y| \geq |x| - t$ (which controls $|x|/2$ if $|x| \geq 2t$). Consequently,

$$\mathbf{K}_{[0,\frac{t}{2}]} \lesssim (1 + t + |x|)^{-3} \int_{r=0}^{\frac{t}{2}} dr \leq \frac{1}{2}(1 + t + |x|)^{-2}. \tag{47}$$

Applying Lemma 5.9, we have

$$\mathbf{K}_{[\frac{t}{2},t]} \leq \frac{2\pi}{|x|} \int_{\tau=0}^{\frac{t}{2}} \int_{\lambda=||x|-t+\tau|}^{|x|+t-\tau} \frac{d\lambda d\tau}{(1 + \tau + \lambda)^2(t - \tau)}.$$

Now, observe that, for all $0 \leq \tau \leq t/2$,

$$\begin{aligned} \frac{1}{|x|(t - \tau)} \int_{\lambda=||x|-t+\tau|}^{|x|+t-\tau} \frac{d\lambda}{(1 + \tau + \lambda)^2} &= \frac{2 \min(|x|, t - \tau)}{|x|(t - \tau)(1 + t + |x|)(1 + \tau + ||x| - t + \tau|)} \\ &\leq \frac{8}{\max(|x|, t)(1 + t + |x|)(1 + \tau + |t - |x||)}. \end{aligned} \tag{48}$$

Let $0 \leq \delta \leq 1$ and write $(1 + \tau + |t - |x||) \geq (1 + \tau)^{1-\delta}(1 + |t - |x||)^\delta$. It remains to integrate in τ in order to derive the expected range of estimates for $\mathbf{K}_{[t/2,t]}$. \square

Finally, a part of our analysis of $A_{\nu,k}^{TT}$ will rely on the following estimate.

Lemma 5.12. *For all $(t, x) \in [1, +\infty[\times \mathbb{R}^3$, we have*

$$\mathbf{Y}_3^{p=3}(t, x) := \int_{1 \leq |y-x| \leq t} (1 + t - |y - x| + |y|)^{-3} \frac{dy}{|y - x|^3} \lesssim \frac{\log(t)}{(1 + t + |x|)^3}.$$

Proof. The inequality can be easily proved if $t \leq 2$ so we assume $t \geq 2$. Start by writing

$$\mathbf{Y}_3^{p=3}(t, x) = \bar{\mathbf{K}}_{[1,\frac{t}{2}]} + \bar{\mathbf{K}}_{[\frac{t}{2},t]}, \quad \bar{\mathbf{K}}_I := \int_{|y-x| \in I} (1 + t - |y - x| + |y|)^{-3} \frac{dy}{|y - x|^3}.$$

Following (47), we have

$$\bar{\mathbf{K}}_{[1,\frac{t}{2}]} \lesssim (1 + t + |x|)^{-3} \int_{r=1}^{\frac{t}{2}} \frac{dr}{r} \leq \log\left(\frac{t}{2}\right)(1 + t + |x|)^{-3}.$$

Next, we apply Lemma 5.9 to get

$$\bar{\mathbf{K}}_{[\frac{t}{2},t]} \leq \frac{2\pi}{|x|} \int_{\tau=0}^{\frac{t}{2}} \int_{\lambda=||x|-t+\tau|}^{|x|+t-\tau} \frac{d\lambda d\tau}{(1 + \tau + \lambda)^2(t - \tau)^2}.$$

If $2t \geq |x|$, we use (48) and $t - \tau \geq t/2$ in order to derive $\bar{\mathbf{K}}_{[t/2,t]} \lesssim t^{-2}(1 + t + |x|)^{-1} \log(1 + t/2)$, which implies the result. Otherwise, $2t \leq |x|$ and we have, for all $0 \leq \tau \leq t/2$,

$$\begin{aligned} \frac{1}{|x|(t - \tau)^2} \int_{\lambda=||x|-t+\tau|}^{|x|+t-\tau} \frac{d\lambda d\tau}{(1 + \tau + \lambda)^2} &= \frac{2 \min(|x|, t - \tau)}{|x|(t - \tau)^2(1 + t + |x|)(1 + \tau + ||x| - t + \tau|)} \\ &\leq \frac{2}{|x|(1 + t + |x|)(1 + |x| - t)(t - \tau)}. \end{aligned} \tag{49}$$

We get, as $2 \leq 2t \leq |x|$,

$$\mathbf{K}_{[\frac{t}{2},t]} \leq 4\pi \log(2)|x|^{-1}(1 + |x|/2)^{-1}(1 + t + |x|)^{-1} \lesssim (1 + t + |x|)^{-3}. \quad \square$$

5.3. The derivatives of order up to $N - 1$. In this subsection, we prove pointwise decay estimates for each of the elements of the decomposition of $\mathcal{L}_{Z^\gamma}(F)$ provided by [Proposition 5.3](#). We start by dealing with $\mathcal{L}_{Z^\gamma}(F)^{\text{data}}$, which is defined on $\mathbb{R}_+ \times \mathbb{R}^3$.

Proposition 5.13. *There exists $C_{\text{data}} > 0$, depending only on N , such that,*

$$\forall |\gamma| \leq N - 1, \forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^{\text{data}}|(t, x) \leq \Lambda C_{\text{data}}(1 + t + |x|)^{-1}(1 + |t - |x||)^{-1}.$$

Proof. In view of the assumptions on the initial data (see [Theorem 2.10](#)) and applying [Corollary 5.5](#) in order to estimate $\mathcal{W}_{\mu\nu}$, we have, for any $|\beta| \leq N - 1$, $\omega \in \mathbb{S}^2$ and $0 \leq \mu, \nu \leq 3$,

$$\forall y \in \mathbb{R}^3, \quad |\mathcal{L}_{Z^\beta}(F)|(0, y) + \langle y \rangle |\nabla_{t,x} \mathcal{L}_{Z^\beta}(F)|(0, y) \lesssim \sum_{|\kappa| \leq |\beta| + 1} \langle y \rangle^{|\kappa|} |\nabla_{t,x}^\kappa F|(0, y) \lesssim \Lambda \langle y \rangle^{-2},$$

$$\left| \int_{\mathbb{R}_v^3} (\mathcal{W}_{\mu\nu}(\omega, \hat{v}) - \delta_\mu^0 \hat{v}^\nu + \delta_\nu^0 \hat{v}^\mu) \widehat{Z}^\beta f(0, y, v) \, dv \right| \leq 3 \int_{\mathbb{R}_v^3} |v^0|^{N_\nu} |\widehat{Z}^\beta f(0, y, v)| \frac{dv}{\langle v \rangle^{N_\nu - 1}} \lesssim \epsilon \langle y \rangle^{-N_x}.$$

The estimates, at $t = 0$, for the time derivatives of the solutions are obtained by using that (1)–(3) are initially verified. Using (22) for $p = N_x \geq 3$, we then deduce that,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^{\text{data}} - \mathcal{L}_{Z^\gamma}(F)^{\text{hom}}|(t, x) \lesssim \epsilon(1 + t + |x|)^{-1}(1 + |t - |x||)^{-N_x + 1} \quad (50)$$

and it remains to use $\epsilon \leq \Lambda$ and to apply [Proposition 2.21](#) to $\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}}$. □

Next, we consider $\mathcal{L}_{Z^\gamma}(F)^S$, which is strongly decaying far from the light cone.

Proposition 5.14. *For any $|\gamma| \leq N - 1$, there holds,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^S|(t, x) \lesssim \bar{\epsilon} \Lambda \frac{\log(3 + |t - |x||)}{(1 + t + |x|)(1 + |t - |x||)^2}.$$

Proof. Fix $0 \leq \mu, \nu \leq 3$ and recall from [Proposition 5.3](#) the definition of $\mathcal{L}_{Z^\gamma}(F)^S$. We have, with $\omega = (y - x)/|y - x|$,

$$\begin{aligned} & |\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^S|(t, x) \\ & \lesssim \sum_{|\xi| + |\kappa| \leq |\gamma|} \int_{|y-x| \leq t} |\mathcal{L}_{Z^\xi}(F)_{\lambda^j}|(t - |y - x|, y) \left| \int_{\mathbb{R}_v^3} \hat{v}^\lambda \partial_{v^j} \mathcal{W}_{\mu\nu}(\omega, v) \widehat{Z}^\kappa f(t - |y - x|, y, v) \, dv \right| \frac{dy}{|y - x|}. \end{aligned}$$

Fix now $|\xi| + |\kappa| \leq N - 1$, $j \in \llbracket 1, 3 \rrbracket$ and $\lambda \in \llbracket 0, 3 \rrbracket$. In view of [Corollary 5.5](#), $\Psi(\omega, v) := \hat{v}^\lambda \partial_{v^j} \mathcal{W}_{\mu\nu}(\omega, v)$ satisfies $|\Psi|(\cdot, v) + |v^0 \nabla_v \Psi|(\cdot, v) \lesssim |v^0|^2 \leq |v^0|^{N_\nu - 1}$. As $|\kappa| \leq N - 1$, [Proposition 4.13](#) then gives us,

$$\forall (\sigma, \tau, y) \in \mathbb{S}^2 \times [0, T[\times \mathbb{R}^3, \quad \left| \int_{\mathbb{R}_v^3} \hat{v}^\lambda \partial_{v^j} \mathcal{W}_{\mu\nu}(\sigma, v) \widehat{Z}^\kappa f(\tau, y, v) \, dv \right| \lesssim \frac{\bar{\epsilon}}{(1 + \tau + |y|)^3}.$$

Applying this last inequality for $(\sigma, \tau) = (\omega, t - |y - x|)$ and estimating the electromagnetic field using the bootstrap assumption (BA1), we get

$$|\mathcal{L}_{Z^\gamma}(F)^S|(t, x) \lesssim \int_{|y-x| \leq t} \frac{\bar{\epsilon} \Lambda}{(1 + t - |y - x| + |y|)^4 (1 + |t - |y - x| - |y||)} \frac{dy}{|y - x|} = \bar{\epsilon} \Lambda Y_{4,1}^{p=1}(t, x).$$

The result then follows from [Lemma 5.10](#). □

We finally deal with $\mathcal{L}_{Z^\gamma}(F)^T$, which actually enjoys stronger decay properties than $\mathcal{L}_{Z^\gamma}(F)^S$ for $t \sim |x|$ (see [Remark 5.16](#) below).

Proposition 5.15. *For any $|\gamma| \leq N - 1$ and all $(t, x) \in [0, T[\times \mathbb{R}^3$, we have*

$$|\mathcal{L}_{Z^\gamma}(F)^T|(t, x) \lesssim \bar{\epsilon}(1+t+|x|)^{-\frac{7}{4}}(1+|t-|x||)^{-\frac{1}{4}}.$$

Proof. In view of the definition of $\mathcal{L}_{Z^\gamma}(F)^T$, introduced in [Proposition 5.3](#), we have

$$|\mathcal{L}_{Z^\gamma}(F)^T|(t, x) \lesssim \sum_{0 \leq \mu, \nu \leq 3} \sum_{|\beta| \leq |\gamma|} \int_{|y-x| \leq t} \left| \int_{\mathbb{R}_v^3} \frac{\mathcal{W}_{\mu\nu}(\omega, v)}{|v^0|^2(1+\omega \cdot \hat{v})} \widehat{Z}^\beta f(t-|y-x|, y, v) dv \right| \frac{dy}{|y-x|^2}, \quad \omega = \frac{y-x}{|y-x|}.$$

Fix $0 \leq \mu, \nu \leq 3$, $|\beta| \leq |\gamma|$ and recall from [Corollary 5.5](#) that

$$\Psi(\sigma, v) := \frac{\mathcal{W}_{\mu\nu}(\sigma, v)}{|v^0|^2(1+\sigma \cdot \hat{v})}$$

satisfies $|\Psi|(\cdot, v) + |\nabla_v \Psi|(\cdot, v) \lesssim v^0$. We then obtain from [Proposition 4.13](#) that,

$$\forall \sigma \in \mathbb{S}^2, \forall (\tau, z) \in [0, T[\times \mathbb{R}^3, \quad \left| \int_{\mathbb{R}_v^3} \frac{\mathcal{W}_{\mu\nu}(\sigma, v)}{|v^0|^2(1+\sigma \cdot \hat{v})} \widehat{Z}^\beta f(\tau, z, v) dv \right| \lesssim \frac{\bar{\epsilon}}{(1+\tau+|z|)^3}.$$

Applying this estimate for $\sigma = \omega$, $\tau = t - |y - x|$ and $z = y$, we get from [Lemma 5.11](#) that

$$|\mathcal{L}_{Z^\gamma}(F)^T|(t, x) \lesssim \bar{\epsilon} Y_3^{p=2}(t, x) \lesssim \bar{\epsilon}(1+t+|x|)^{-\frac{7}{4}}(1+|t-|x||)^{-\frac{1}{4}}. \quad \square$$

Remark 5.16. In fact, [Lemma 5.11](#) also provides $|\mathcal{L}_{Z^\gamma}(F)^T|(t, x) \lesssim \bar{\epsilon}(1+t+|x|)^{-2} \log(1+t)$. Moreover, the estimate could be significantly improved in the exterior of the light cone, where $|x| \geq t$.

If the constant C_{boot} is chosen such that $C_{\text{boot}} \geq 2C_{\text{data}}$ and if ϵ is small enough, [Propositions 5.13, 5.14](#) and [5.15](#) allow us to improve the bootstrap assumption [\(BA1\)](#).

5.4. The top-order derivatives. In this section, we estimate all the terms listed in [Proposition 5.7](#) in order to improve the bootstrap assumption [\(BA2\)](#). We start by dealing with the ones depending explicitly on the data.

Proposition 5.17. *There exists a constant \bar{C}_{data} , depending only on N , such that, for any $k \in \llbracket 1, 3 \rrbracket$ and $|\gamma| = N - 1$,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |A_{\gamma, k}^{\text{data}}|(t, x) \leq \Lambda \bar{C}_{\text{data}}(1+t+|x|)^{-1}(1+|t-|x||)^{-2}.$$

Proof. Recall from [Propositions 5.3](#) and [5.7](#) the expression of $A_{\gamma, k}^{\text{data}}$ and from [Corollaries 5.5](#) and [5.8](#) the bounds on the kernels. Hence, for $(t, x) \in [0, T[\times \mathbb{R}^3$,

$$|A_{\gamma, k}^{\text{data}}|(t, x) \lesssim |\nabla_{\partial_x k} \mathcal{L}_{Z^\gamma}(F)^{\text{hom}}|(t, x) + \sum_{|\beta| \leq |\gamma|+1} \min(t^{-1}, t^{-2}) \int_{|y-x|=t} \int_{\mathbb{R}_v^3} |v^0|^3 |\widehat{Z}^\beta f|(0, y, v) dv dy.$$

As $[\partial_{x^\mu}, Z] = 0$ or $[\partial_{x^\mu}, Z] = \pm \partial_{x^\nu}$ for any $Z \in \mathbb{K}$, by the equivalence of the pointwise norms (9) and in view of the smallness assumptions on the initial data, there holds

$$|\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)^{\text{hom}}|(0, y) = |\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)|(0, y) \lesssim \sum_{1 \leq |\kappa| \leq N} \langle y \rangle^{|\kappa|-1} |\nabla_{t,x}^\kappa F|(0, y) \lesssim \Lambda \langle y \rangle^{-3},$$

$$|\nabla_{t,x} \nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)^{\text{hom}}|(0, y) = |\nabla_{t,x} \nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)|(0, y) \lesssim \sum_{2 \leq |\kappa| \leq N+1} \langle y \rangle^{|\kappa|-2} |\nabla_{t,x}^\kappa F|(0, y) \lesssim \Lambda \langle y \rangle^{-4}.$$

As $\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)_{\mu\nu}^{\text{hom}}$ is solution to the homogeneous wave equation, Proposition 2.21 gives

$$|\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\nu}(F)^{\text{hom}}|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-2}.$$

Since $|v^0|^{-N_v+3} \in L^1(\mathbb{R}_v^3)$, we have, for any $|\beta| \leq N$,

$$\int_{\mathbb{R}_v^3} |v^0|^3 |\widehat{Z}^\beta f|(0, y, v) \, dv \lesssim \langle y \rangle^{-N_x} \sup_{|\kappa|+|\xi| \leq N} \sup_{(x,v) \in \mathbb{R}^6} |v^0|^{N_v+|\xi|} \langle x \rangle^{N_x+|\kappa|} |\partial_v^\kappa \partial_x^\xi f|(0, x, v) \lesssim \epsilon \langle y \rangle^{-N_x}.$$

Consequently, as $N_x \geq 5$, we have

$$|A_{\gamma,k}^{\text{data}}|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-2} + \epsilon \min(t^{-1}, t^{-2}) \mathcal{Q}_{t,x}, \quad \mathcal{Q}_{t,x} := \int_{|y-x|=t} \frac{d\mu_{\mathbb{S}^2}}{\langle y \rangle^5}.$$

As $\epsilon \leq \Lambda$, it remains to prove $\min(t^{-1}, t^{-2}) \mathcal{Q}_{t,x} \lesssim (1+t+|x|)^{-1}(1+|t-|x||)^{-2}$ and, for this, we consider two cases.

- If $t \leq 1$, then $|y| \geq |x| - 1$ on the domain of integration and $\mathcal{Q}_{t,x} \lesssim 4\pi t^2 \langle x \rangle^{-5}$. It remains to note that $\langle x \rangle \geq 1+t+|x| \geq 1+|t-|x||$ and $t^{-1} \leq t^{-2}$ in the region considered.
- Otherwise $t \geq 1$ and we have $\mathcal{Q}_{t,x} \lesssim t(1+t+|x|)^{-1}(1+|t-|x||)^{-2}$ according to the estimate (22). The result follows from $t^{-2} \leq t^{-1}$ in the domain treated here. \square

Next, we consider the vertex term.

Proposition 5.18. *Let $k \in \llbracket 1, 3 \rrbracket$ and $|\gamma| = N - 1$. We have,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |A_{\gamma,k}^{\text{ver}}|(t, x) \lesssim \bar{\epsilon}(1+t+|x|)^{-3}.$$

Proof. Fix $0 \leq \mu, \nu \leq 3$, $(t, x) \in [0, T[\times \mathbb{R}^3$ and recall $N_v \geq 15$, so that Corollary 5.8 implies $|\mathcal{D}_{\mu\nu}^k(\omega, v) + |v^0 \nabla_v \mathcal{D}_{\mu\nu}^k|(\omega, v) \lesssim |v^0|^{N_v-11}$. Proposition 4.13, applied for $\Psi = \mathcal{D}_{\mu\nu}^k$ and to any $|\beta| \leq N - 1$, then yields

$$\begin{aligned} |A_{\gamma,k,\mu\nu}^{\text{ver}}|(t, x) &\lesssim \sum_{|\beta| \leq |\gamma|} \int_{\sigma \in \mathbb{S}^2} \left| \int_{\mathbb{R}_v^3} \mathcal{D}_{\mu\nu}^k(\sigma, v) \widehat{Z}^\beta f(t, x, v) \, dv \right| d\mu_{\mathbb{S}^2} \\ &\lesssim \frac{\bar{\epsilon}}{(1+t+|x|)^3} \int_{\sigma \in \mathbb{S}^2} d\mu_{\mathbb{S}^2} = \frac{4\pi \bar{\epsilon}}{(1+t+|x|)^3}. \end{aligned} \quad \square$$

We now estimate $A_{\gamma,k}^{T,S}$. Note that the next result could be easily improved but it is more than enough for the purpose of improving the bootstrap assumption (BA2).

Proposition 5.19. *For any $k \in \llbracket 1, 3 \rrbracket$ and $|\gamma| = N - 1$, there holds,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |A_{\gamma,k}^{T,S}|(t, x) \lesssim \bar{\epsilon} \Lambda (1 + t + |x|)^{-\frac{3}{2}} (1 + |t - |x||)^{-2}.$$

Proof. Let $0 \leq \mu, \nu \leq 3$, $(t, x) \in [0, T[\times \mathbb{R}^3$ and recall that $K_\xi^j := \hat{v}^\lambda \mathcal{L}_{Z^\xi}(F)_\lambda^j$. Consequently, $|A_{\gamma,k,\mu\nu}^{T,S}|(t, x)$ is bounded by a linear combination of terms of the form

$$\mathcal{Q}_{t,x}^{\xi,\kappa} := \int_{|y-x| \leq t} |\mathcal{L}_{Z^\xi}(F)_\lambda^j|(t - |y - x|, y) \int_{\mathbb{R}_v^3} |\partial_{v^j} \mathcal{B}_{\mu\nu}^k(\omega, v)| |\widehat{Z}^\kappa f(t - |y - x|, y, v)| \, dv \frac{dy}{|y - x|^2},$$

with $|\xi| + |\kappa| \leq N - 1$ and where we recall that $\omega = (y - x)/|y - x|$. Since $|\partial_{v^j} \mathcal{B}_{\mu\nu}^k(\omega, v)| \lesssim |v^0|^3$ by [Corollary 5.8](#) and $N_v \geq 13$, [Proposition 4.11](#), applied for $a = 1$, provides

$$\int_{\mathbb{R}_v^3} |\partial_{v^j} \mathcal{B}_{\mu\nu}^k(\omega, v)| |\widehat{Z}^\kappa f(t - |y - x|, y, v)| \, dv \lesssim \frac{\bar{\epsilon} (1 + |t - |y - x| - |y||)}{(1 + t - |y - x| + |y|)^{3+\frac{1}{2}}}.$$

Moreover, as $|\xi| \leq N - 1$, the bootstrap assumption [\(BA1\)](#) gives

$$|\mathcal{L}_{Z^\xi}(F)_\lambda^j|(\tau, y) \lesssim \Lambda (1 + t - |y - x| + |y|)^{-1} (1 + |t - |y - x| - |y||)^{-1}.$$

Consequently, the last two estimates yield

$$\mathcal{Q}_{t,x}^{\xi,\kappa} \lesssim \bar{\epsilon} \Lambda \int_{|y-x| \leq t} (1 + t - |y - x|)^{-4-\frac{1}{2}} \frac{dy}{|y - x|^2} = \bar{\epsilon} \Lambda Y_{4+\frac{1}{2}}^{p=2}(t, x)$$

and the result follows from [Lemma 5.11](#). □

We pursue with the analysis of $A_{\gamma,k}^{SS}$. As for the previous term, the estimate could be improved.

Proposition 5.20. *We have, for any $k \in \llbracket 1, 3 \rrbracket$ and $|\gamma| = N - 1$,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |A_{\gamma,k}^{SS}|(t, x) \lesssim \bar{\epsilon} \Lambda \langle \Lambda \rangle (1 + t + |x|)^{-1} (1 + |t - |x||)^{-\frac{5}{2}}.$$

Proof. We fix $(t, x) \in [0, T[\times \mathbb{R}^3$ and we recall that $K_\xi^j := \hat{v}^\lambda \mathcal{L}_{Z^\xi}(F)_\lambda^j$. Recall further from [Proposition 5.7](#) that $A_{\gamma,k}^{SS}$ can be decomposed as the sum of four terms. Bounding the kernel in $A_{\gamma,k}^{SS,I}$ by [Corollary 5.8](#) and estimating the derivatives of the electromagnetic field using [\(BA1\)](#), we have

$$\begin{aligned} & |A_{\gamma,k}^{SS,I}|(t, x) \\ & \lesssim \sum_{|\kappa| \leq N-1} \int_{|y-x| \leq t} \frac{\Lambda^2}{(1 + \tau + |y|)^2 (1 + |\tau - |y||)^2} \int_{\mathbb{R}_v^3} |v^0|^3 |\widehat{Z}^\kappa f|(\tau, y, v) \, dv \frac{dy}{|y - x|}, \quad \tau := t - |y - x|. \end{aligned}$$

For the next two terms, recall that $T_0 = \hat{v}^\lambda \partial_{x^\lambda}$ and the expression of K_ξ . Recall further from [Corollary 5.8](#) that the integral kernels are bounded by $|v^0|^3$. Consequently, we can bound $|A_{\gamma,k}^{SS,II}|(t, x) + |A_{\gamma,k}^{SS,III}|(t, x)$ by a linear combination of terms of the form

$$\mathcal{R}_{t,x}^{\xi,\kappa} := \int_{|y-x| \leq t} |\nabla_{t,x} \mathcal{L}_{Z^\xi}(F)| (t - |y - x|, y) \int_{\mathbb{R}_v^3} |v^0|^3 |\widehat{Z}^\kappa f(t - |y - x|, y, v)| \, dv \frac{dy}{|y - x|},$$

where $|\xi| + |\kappa| \leq N - 1$. We estimate the electromagnetic field through (BA2) if $|\xi| = N - 1$ or by Proposition 3.2 if $|\xi| \leq N - 2$. This leads to the bound

$$|A_{\gamma,k}^{SS,II}(t, x) + |A_{\gamma,k}^{SS,III}(t, x)| \lesssim \sum_{|\kappa| \leq N-1} \int_{|y-x| \leq t} \frac{\Lambda \log(3 + |\tau - |y||)}{(1 + \tau + |y|)(1 + |\tau - |y||)^2} \int_{\mathbb{R}^3} |v^0|^3 |\widehat{Z}^\kappa f|(\tau, y, v) dv \frac{dy}{|y-x|},$$

where $\tau = t - |y - x|$. Controlling the velocity average through the improved estimates of Proposition 4.11 yields, as $N_v \geq 13$,

$$|A_{\gamma,k}^{SS,I}(t, x) + |A_{\gamma,k}^{SS,II}(t, x) + |A_{\gamma,k}^{SS,III}(t, x)| \lesssim \bar{\epsilon} \Lambda \langle \Lambda \rangle \int_{|y-x| \leq t} \frac{\log^{3N_x+3N+1}(3 + t - |y-x| + |y|)}{(1 + t - |y-x| + |y|)^5 (1 + |t - |y-x| - |y||)} \frac{dy}{|y-x|} \lesssim \bar{\epsilon} \Lambda \langle \Lambda \rangle \mathbf{Y}_{4+\frac{3}{4},1}^{p=1}(t, x).$$

Finally, we can bound similarly $|A_{\gamma,k}^{SS,IV}(t, x)$ by a linear combination of terms of the form

$$\bar{\mathcal{R}}_{t,x}^{\xi,\kappa} := \int_{|y-x| \leq t} |\mathcal{L}_{Z^\xi}(F)|(t - |y-x|, y) \left| \int_{\mathbb{R}^3} \mathcal{V}(\omega, v) \partial_{x^n} \widehat{Z}^\kappa f(t - |y-x|, y, v) dv \right| \frac{dy}{|y-x|},$$

where $|\xi| + |\kappa| \leq N - 1$, $1 \leq n \leq 3$ and $\mathcal{V}(\omega, v)$ is of the form $\mathcal{C}_{\mu\nu}^k(\omega, v) \hat{v}^\lambda |v^0|^{-1}$. We get from Corollary 5.8 that $|\mathcal{V}(\omega, v)| + |v^0 \nabla_v \mathcal{V}(\omega, v)| \lesssim |v^0|^3$, so that Proposition 4.15 gives,

$$\forall (\tau, y, \sigma) \in [0, T[\times \mathbb{R}^3 \times \mathbb{S}^2, \quad \left| \int_{\mathbb{R}^3} \mathcal{V}(\sigma, v) \partial_{x^n} \widehat{Z}^\kappa f(\tau, y, v) dv \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3 + \tau)}{(1 + \tau + |y|)^4}.$$

Applying it to $\sigma = \omega$ and $\tau = t - |y - x|$ and estimating the electromagnetic field using (BA1), we get

$$\bar{\mathcal{R}}_{t,x}^{\xi,\kappa} \lesssim \bar{\epsilon} \Lambda \int_{|y-x| \leq t} \frac{\log^{3N_x+3N}(3 + t - |y-x|)}{(1 + t - |y-x| + |y|)^5 (1 + |t - |y-x| - |y||)} \frac{dy}{|y-x|} \lesssim \bar{\epsilon} \Lambda \mathbf{Y}_{4+\frac{3}{4},1}^{p=1}(t, x).$$

Consequently, $|A_{\gamma,k}^{SS}(t, x) \lesssim \bar{\epsilon} \Lambda \langle \Lambda \rangle \mathbf{Y}_{4+\frac{3}{4},1}^{p=1}(t, x)$, so that the result follows from Lemma 5.10. □

Finally, we deal with the most problematic term, the one with an integral kernel carrying the nonintegrable weight $|y - x|^{-3}$.

Proposition 5.21. *Let $k \in \llbracket 1, 3 \rrbracket$ and $|\gamma| = N - 1$. Then,*

$$\forall (t, x) \in [0, T[\times \mathbb{R}^3, \quad |A_{\gamma,k}^{TT}(t, x) \lesssim \bar{\epsilon} \frac{\log(3 + t)}{(1 + t + |x|)^3}.$$

Proof. Let $0 \leq \mu, \nu \leq 3$, $|\beta| \leq N - 1$ and

$$G_\sigma^\beta(\tau, y) := \int_{\mathbb{R}^3} \mathcal{A}_{\mu\nu}^k(\sigma, v) \widehat{Z}^\beta f(\tau, y, v) dv, \quad (\sigma, \tau, y) \in \mathbb{S}^2 \times [0, T[\times \mathbb{R}^3.$$

Recall from Corollary 5.8 the bound on the kernel $\mathcal{A}_{\mu\nu}^k$ and apply Proposition 4.13 for $\Psi = \mathcal{A}_{\mu\nu}^k$. We obtain,

$$\forall (\sigma, \tau, y) \in \mathbb{S}^2 \times [0, T[\times \mathbb{R}^3, \quad |G_\sigma^\beta(\tau, y) \lesssim \bar{\epsilon} (1 + \tau + |y|)^{-3},$$

which, applied for $(\sigma, \tau) = (\omega, t - |y - x|)$, yields

$$|A_{\gamma,k,\mu\nu}^{TT}(t, x)| \lesssim \bar{\epsilon} \mathbf{Y}_3^{p=3}(t, x) + \sum_{|\beta| \leq N-1} \mathcal{U}_{t,x}^\beta, \quad \mathcal{U}_{t,x}^\beta := \left| \int_{|y-x| \leq 1} \int_{\mathbb{R}_v^3} \mathcal{A}_{\mu\nu}^k(\omega, v) \widehat{Z}^\beta f(t - |y - x|, y, v) \frac{dv dy}{|y - x|^3} \right|.$$

Fix $|\beta| \leq N - 1$ and recall from Proposition 5.7 that the average of $\sigma \mapsto \mathcal{A}_{\mu\nu}^k(\sigma, \cdot)$ on \mathbb{S}^2 vanishes. Hence,

$$\begin{aligned} \mathcal{U}_{t,x}^\beta &= \left| \int_{|y-x| \leq 1} \int_{\mathbb{R}_v^3} \mathcal{A}_{\mu\nu}^k(\omega, v) (\widehat{Z}^\beta f(t - |y - x|, y, v) - \widehat{Z}^\beta f(t - |y - x|, x, v)) \frac{dv dy}{|y - x|^3} \right| \\ &\leq \int_{|y-x| \leq 1} |G_\omega^\beta(t - |y - x|, y) - G_\omega^\beta(t - |y - x|, x)| \frac{dy}{|y - x|^3}. \end{aligned}$$

For any $(\sigma, \tau) \in \mathbb{S}^2 \times [0, T[$, we apply the mean value theorem to $s \mapsto G_\sigma^\beta(\tau, x + s(y - x))$ on the interval $[0, 1]$. Then, there exists $x_{\sigma,\tau}$ in the segment $[x, y] \subset \mathbb{R}^3$ such that

$$G_\sigma^\beta(\tau, y) - G_\sigma^\beta(\tau, x) = \omega \cdot \nabla_x G_\sigma^\beta(\tau, x_{\sigma,\tau}) |y - x|, \quad \omega = \frac{y - x}{|y - x|}.$$

Apply now Proposition 4.15 for $\Phi = \mathcal{A}_{\mu\nu}$ in order to get, for any $1 \leq i \leq 3$,

$$\begin{aligned} \forall (\sigma, \tau, z) \in \mathbb{S}^2 \times [0, T[\times \mathbb{R}^3, \quad |\partial_{x^i} G_\sigma^\beta|(\tau, z) &= \left| \int_{\mathbb{R}^3} \mathcal{A}_{\mu\nu}^k(\sigma, v) \partial_{x^i} \widehat{Z}^\beta f(\tau, z, v) dv \right| \\ &\lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3 + \tau)}{(1 + \tau + |z|)^4}. \end{aligned}$$

Applying the last two identities for $\sigma = \omega$, $\tau = t - |y - x|$ and $z = x_{\sigma,\tau}$ yields

$$\mathcal{U}_{t,x}^\beta \lesssim \int_{|y-x| \leq 1} \frac{\bar{\epsilon}}{(1 + t - |y - x| + |x_{\omega,t-|y-x||})^3} \frac{dy}{|y - x|^2}.$$

As $|y - x| \leq 1$ and $x_{\omega,t-|y-x|} \in [x, y]$, we have $1 + t - |y - x| \geq \frac{1}{2}(1 + t)$ and $|x_{\omega,t-|y-x||} \geq |x| - 1$, so that

$$|A_{\gamma,k,\mu\nu}^{TT}(t, x)| \lesssim \bar{\epsilon} \mathbf{Y}_3^{p=3}(t, x) + \bar{\epsilon} (1 + t + |x|)^{-3}.$$

We conclude the proof by applying Lemma 5.12. □

As in the previous subsection, if C_{boot} is chosen such that $C_{\text{boot}} \geq 2\bar{C}_{\text{data}}$ and if ϵ is small enough, we can improve the bootstrap assumption (BA2) for the spatial derivatives $\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\gamma}(F)$, with $1 \leq k \leq 3$ and $|\gamma| = N - 1$, by applying Propositions 5.17–5.21. The time derivative can then be controlled using

$$|\nabla_{\partial_t} \mathcal{L}_{Z^\gamma}(F)| \lesssim \sum_{1 \leq k \leq 3} |\nabla_{\partial_{x^k}} \mathcal{L}_{Z^\gamma}(F)| + \sum_{|\beta| \leq |\gamma|} |J(\widehat{Z}^\beta f)|,$$

which follows from the commuted Maxwell equations (see Proposition 2.4). We stress, for the smallness condition on ϵ , that $\bar{\epsilon} \langle \Lambda \rangle^2 \leq 2\epsilon e^{(D+3)\Lambda}$.

6. Modified scattering for the distribution function

In this section, we determine the precise asymptotic behavior of the particle density and its derivatives under the additional assumption (15) on the initial electromagnetic field. In particular, we determine the

self-similar profile of the current density $J(f)$ as well as the one of the Maxwell field F and we define modified trajectories along which f converges to a new smooth density function.

6.1. Convergence of the spatial averages. Since the solution (f, F) is global in time, all the statements of Sections 3–5 hold true for $T = +\infty$. We can then deduce that $\int_x \widehat{Z}^\beta f \, dx$ converges to a function defined on \mathbb{R}_v^3 .

Proposition 6.1. *Let $|\beta| \leq N - 1$. There exists a continuous function $Q_\infty^\beta \in L_v^1 \cap L_v^\infty$ such that,*

$$\forall t \in \mathbb{R}_+, \quad \left\| |v^0|^{N_v-6} \left(Q_\infty^\beta - \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, \cdot) \, dx \right) \right\|_{L^\infty(\mathbb{R}_v^3)} \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{1+t}.$$

Remark 6.2. This estimate directly implies that $|v^0|^{N_v-10} \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, \cdot) \, dx \rightarrow |v^0|^{N_v-10} Q_\infty^\beta$ in $L^1(\mathbb{R}_v^3)$, as $t \rightarrow +\infty$, with the same rate for convergence.

Proof. Let $v \in \mathbb{R}_v^3$ and apply Lemma 4.7 in order to get, for all $0 \leq t \leq s$,

$$|v^0|^{N_v-6} \left| \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(s, x, v) \, dx - \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) \, dx \right| \lesssim \bar{\epsilon} \int_{\tau=t}^s \frac{\log^{3N_x+3N}(3+\tau)}{(1+\tau)^2} \, d\tau \leq \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{1+t}.$$

Consequently, there exists $Q_\infty^\beta \in L_v^\infty$ such that $\int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(s, x, v) \, dx \rightarrow Q_\infty^\beta$ in L_v^∞ as $s \rightarrow +\infty$. Moreover, letting $s \rightarrow +\infty$ in the previous estimate provides the rate of convergence stated in the proposition. It implies $|v^0|^{N_v-6} Q_\infty^\beta \in L_v^\infty$ and then, as $N_v > 9$, $Q_\infty^\beta \in L_v^1$. \square

It turns out that these functions are differentiable for $|\beta| \leq N - 2$ and that $\partial_{v^i} Q_\infty^\beta$ can be related to other such functions Q_∞^κ . For this reason, if $\widehat{Z}^\beta = \widehat{\Omega}_{0i} \widehat{Z}^\beta$, we will use $Q_\infty^{\widehat{\Omega}_{0i}\beta}$ in order to denote Q_∞^κ .

Proposition 6.3. *For any $|\beta| \leq N - 2$, $Q_\infty^\beta \in C^{N-1-|\beta|}(\mathbb{R}_v^3)$ and its derivatives can be obtained by iterating the relations*

$$v^0 \partial_{v^i} Q_\infty^\beta = Q_\infty^{\widehat{\Omega}_{0i}\beta} - \hat{v}^i Q_\infty^\beta, \quad 1 \leq i \leq 3. \tag{51}$$

Proof. Let $(t, v) \in \mathbb{R}_+ \times \mathbb{R}_v^3$ and note that

$$v^0 \partial_{v^i} \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) \, dx = \int_{\mathbb{R}_x^3} \widehat{\Omega}_{0i} \widehat{Z}^\beta f(t, x, v) \, dx - t \int_{\mathbb{R}_x^3} \partial_{x^i} \widehat{Z}^\beta f(t, x, v) \, dx - \int_{\mathbb{R}_x^3} x^i \partial_t \widehat{Z}^\beta f(t, x, v) \, dx.$$

Writing $\partial_t = -\hat{v} \cdot \nabla_x - \hat{v}^\mu F_\mu^j \partial_{v^j} + \mathbf{T}_F$, we get by performing integration by parts,

$$\begin{aligned} v^0 \partial_{v^i} \int_{\mathbb{R}_x^3} \widehat{Z}^\beta f(t, x, v) \, dx &= \int_{\mathbb{R}_x^3} \widehat{\Omega}_{0i} \widehat{Z}^\beta f(t, x, v) - \hat{v}^i \widehat{Z}^\beta f(t, x, v) \, dx + \int_{\mathbb{R}_x^3} x^i (\hat{v}^\mu F_\mu^j \partial_{v^j} - \mathbf{T}_F)(\widehat{Z}^\beta f)(t, x, v) \, dx. \end{aligned}$$

According to Proposition 6.1, the first term on the right-hand side converges to $Q_\infty^{\widehat{\Omega}_{0i}\beta} - \hat{v}^i Q_\infty^\beta$, as $t \rightarrow +\infty$ and in $L^\infty(\mathbb{R}_v^3)$. Following the proof of Lemma 4.7 and then using Proposition 4.5, one can prove

$$\begin{aligned} \left| \int_{\mathbb{R}^3} x^i (\hat{v}^\mu F_\mu^j \partial_{v^j} - \mathbf{T}_F)(\widehat{Z}^\beta f)(t, x, v) dx \right| &\lesssim \Lambda \frac{\log(3+t)}{1+t} \sup_{|\kappa| \leq |\beta|+1} \sup_{x \in \mathbb{R}^3} |v^0|^3 z^{N_x-2} \widehat{Z}^\kappa f|(t, x, v) \\ &\lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{1+t}. \end{aligned}$$

We then deduce (51) and, by a direct induction, $Q_\infty^\beta \in C^{N-1-|\beta|}(\mathbb{R}_v^3)$. □

Let us mention that any Q_∞^β can be written as a combination of Q_∞ and Q_∞^κ , where \widehat{Z}^κ is only composed of complete lifts of Lorentz boosts $\widehat{\Omega}_{0i}$.

Proposition 6.4. *Let $|\beta| \leq N - 1$. Then:*

- If $\beta_T \geq 1$, which means that \widehat{Z}^β is composed of at least one translation, we have $Q_\infty^\beta = 0$.
- Otherwise there exists $n + |\kappa| \leq |\beta|$ such that $\widehat{Z}^\beta = S^n \widehat{Z}^\kappa$ and $Q_\infty^\beta = (-3)^n Q_\infty^\kappa$.
- Moreover, if $\widehat{Z}^\beta = \widehat{\Omega}_{jk} \widehat{Z}^\kappa$, $1 \leq j < k \leq 3$, then $Q_\infty^\beta = \hat{v}^j Q^{\widehat{\Omega}_{0k\kappa}} - \hat{v}^k Q^{\widehat{\Omega}_{0j\kappa}}$.

Proof. Assume first that $\beta_T \geq 1$. Since $[\widehat{Z}, \partial_{x^\mu}] = 0$ or $\pm \partial_{x^v}$ for any $0 \leq \mu \leq 3$ and $\widehat{Z} \in \widehat{\mathbb{P}}_0$, it suffices to consider the case $\widehat{Z}^\beta = \partial_{x^\mu} \widehat{Z}^\xi$. Then, by either applying Lemma 4.7 or by performing integration by parts,

$$\left| \int_{\mathbb{R}^3} \partial_t \widehat{Z}^\xi f(t, x, v) dx \right| \lesssim \bar{\epsilon} (1+t)^{-\frac{3}{2}} \rightarrow 0, \quad \int_{\mathbb{R}^3} \partial_{x^i} \widehat{Z}^\xi f(t, x, v) dx = 0, \quad 1 \leq i \leq 3.$$

Otherwise $\beta_T = 0$ and since S commutes with $\widehat{\Omega}_{jk}$ and $\widehat{\Omega}_{0i}$, there exists $n + |\kappa| \leq |\beta|$ such that $\widehat{Z}^\beta = S^n \widehat{Z}^\kappa$. The result follows from an easy induction and the following properties, which hold for any $|\xi| \leq N - 2$:

$$\left| \int_{\mathbb{R}^3} t \partial_t \widehat{Z}^\xi f(t, x, v) dx \right| \lesssim \bar{\epsilon} (1+t)^{-\frac{1}{2}} \rightarrow 0, \quad \int_{\mathbb{R}^3} x_i \partial_{x^i} \widehat{Z}^\xi f(t, x, v) dx = - \int_{\mathbb{R}^3} \widehat{Z}^\xi f(t, x, v) dx, \quad 1 \leq i \leq 3.$$

Finally, if $\widehat{Z}^\beta = \widehat{\Omega}_{jk} \widehat{Z}^\kappa$, note that by integration by parts,

$$\int_x \widehat{Z}^\beta f dx = \hat{v}^j \int_x v^0 \partial_{v^k} \widehat{Z}^\kappa f dx - \hat{v}^k \int_x v^0 \partial_{v^j} \widehat{Z}^\kappa f dx$$

and it remains to apply Proposition 6.3. □

We are now able to establish the precise behavior of $J(f)$ in the interior of the light cone. In other words, we improve Corollary 4.14. No such result holds for the exterior region since the decay can be arbitrarily fast (we refer for this to the third estimate of Proposition 4.11). Recall the notation $x^0 = t$.

Proposition 6.5. *For any $|\beta| \leq N - 1$, the components of the electric current density $J(\widehat{Z}^\beta f)$, that is, $J^\mu(\widehat{Z}^\beta f) = \int_{\mathbb{R}^3} (v^\mu/v^0) \widehat{Z}^\beta f dv$, satisfy,*

$$\forall |x| < t, \quad \left| t^3 J^\mu(\widehat{Z}^\beta f)(t, x) - \frac{x^\mu}{t} (|v^0|^5 Q_\infty^\beta) \left(\frac{\check{x}}{t} \right) \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{t}, \quad \mu \in \llbracket 0, 3 \rrbracket.$$

Proof. Let $|\beta| \leq N - 1$, $0 \leq \mu \leq 3$ and $|x| < t$. Apply Lemma 4.12 and the estimate (41) to $g(t, x, v) := \hat{v}^\mu \widehat{Z}^\beta f(t, x + t\hat{v}, v)$. Since the spatial average of $|v^0|^5 g$ is equal to the one of $\hat{v}^\mu |v^0|^5 \widehat{Z}^\beta f$, we get

$$\left| t^3 \int_{\mathbb{R}_v^3} \frac{v^\mu}{v^0} \widehat{Z}^\beta f(t, x, v) dv - \int_{\mathbb{R}_y^3} \left(\frac{v^\mu}{v^0} |v^0|^5 \widehat{Z}^\beta f \right) \left(t, y, \frac{\check{x}}{t} \right) dy \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{t}. \tag{52}$$

As $N_v - 6 \geq 5$, we obtain from Proposition 6.1 that,

$$\forall v \in \mathbb{R}_v^3, \quad \left| \frac{v^\mu}{v^0} |v^0|^5 Q_\infty^\beta(v) - \frac{v^\mu}{v^0} |v^0|^5 \int_{\mathbb{R}_y^3} \widehat{Z}^\beta f(t, y, v) dy \right| \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{1+t}.$$

The result follows from (52) and the last estimate, applied for $v = \widetilde{x/t}$. □

6.2. Self-similar asymptotic profile of the electromagnetic field. To identify the profile of F , we will see that Q_∞ generates an effective electromagnetic field. For this, we study F^T since it is the element of the Glassey–Strauss decomposition of F with the slower decay rate along timelike geodesics $t \mapsto (t, x + t\hat{v})$. If the plasma is not neutral, $Q_F \neq 0$, we will also have to improve the estimate for F^{data} .

6.2.1. Behavior of $\mathcal{L}_{Z^V}(F)^T$ along timelike straight lines. It will be convenient to lighten the notations by denoting the kernel in the integral defining F^T , which was bounded in Corollary 5.5, as

$$\mathcal{W}^T(\omega, v) := \frac{\mathcal{W}(\omega, v)}{|v^0|^2(1 + \omega \cdot \hat{v})}, \quad |\mathcal{W}^T(\cdot, v) + |\nabla_v \mathcal{W}^T|(\cdot, v)| \lesssim v^0. \tag{53}$$

Definition 6.6. Let, for any $|\beta| \leq N - 1$, $[\widehat{Z}^\beta f]^\infty(v)$ be the 2-form defined as,

$$\forall v \in \mathbb{R}_v^3, \quad [\widehat{Z}^\beta f]^\infty(v) := \int_{\substack{|z| \leq 1 \\ |z + \hat{v}| < 1 - |z|}} \mathcal{W}^T\left(\frac{z}{|z|}, \frac{z + \hat{v}}{1 - |z|}\right) (|v^0|^5 Q_\infty^\beta)\left(\frac{z + \hat{v}}{1 - |z|}\right) \frac{dz}{|z|^2(1 - |z|)^3}.$$

Remark 6.7. We recall our convention $(|v^0|^5 Q_\infty^\beta)(w) := |w^0|^5 Q_\infty^\beta(w)$ for any $w \in \mathbb{R}_v^3$.

Remark 6.8. It is crucial to observe that the domain of integration is included in $\{0 \leq |z| \leq (1 + |\hat{v}|)/2\}$. Indeed, if $|z| \geq (1 + |\hat{v}|)/2$, we have

$$|z + \hat{v}| \geq |z| - 1 + 1 - |\hat{v}| \geq \frac{1 - |\hat{v}|}{2} \geq 1 - |z|.$$

Consequently,

$$|z| \leq 1, \quad |z + \hat{v}| < 1 - |z| \quad \implies \quad \frac{1}{4|v^0|^2} \leq \frac{1 - |\hat{v}|}{2} \leq 1 - |z| \leq 1.$$

In order to transform decay in $|t - r|$ into decay in t along timelike trajectories, we will use the next property.

Lemma 6.9. Let $(x, v) \in \mathbb{R}_x^3 \times \mathbb{R}_v^3$. Then,

$$\forall 1 \leq t \leq 4\langle x \rangle |v^0|^2, \quad 1 \leq 4 \frac{\langle x \rangle |v^0|^2}{t}, \quad \forall t \geq 4\langle x \rangle |v^0|^2, \quad t - |x + t\hat{v}| \geq \frac{t}{4|v^0|^2}.$$

Proof. It suffices to observe that,

$$\forall t \geq 4|x| |v^0|^2, \quad t \geq \frac{2|x|}{1 - |\hat{v}|}, \quad \text{so that} \quad t - |x + t\hat{v}| \geq t - \frac{1 - |\hat{v}|}{2} t - |\hat{v}|t = t - \frac{1 + |\hat{v}|}{2} t \geq \frac{t}{4|v^0|^2}. \quad \square$$

We have the following convergence result.

Proposition 6.10. Let $|\beta| \leq N - 1$ and $(x, v) \in \mathbb{R}_x^3 \times \mathbb{R}_v^3$. For all $t \geq 1$, there holds

$$|t^2 [\widehat{Z}^\beta f]^T(t, x + \hat{v}t) - [\widehat{Z}^\beta f]^\infty(v)| \lesssim \bar{\epsilon} \langle x \rangle^2 |v^0|^8 \frac{\log^{3N_x+3N+1}(3+t)}{t}.$$

Proof. Fix $|\beta| \leq N - 1$, $(t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$ and recall from [Proposition 5.3](#) the definition of $[\widehat{Z}^\beta f]^T$. Next, we split the domain of integration of $[\widehat{Z}^\beta f]^T$ into two parts,

$$t^2[\widehat{Z}^\beta f]^T(t, x + \hat{v}t) = t^2 \int_{\substack{|y-x-t\hat{v}| \leq t \\ |y-x| \geq t - |y-x-t\hat{v}|}} \int_{\mathbb{R}_w^3} \mathcal{W}^T\left(\frac{y-x}{|y-x|}, w\right) \widehat{Z}^\beta f(t - |y-x-t\hat{v}|, y, w) \frac{dw dy}{|y-x-t\hat{v}|^2} + \mathcal{J},$$

$$\mathcal{J} := \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \int_{\mathbb{R}_w^3} \mathcal{W}^T\left(\frac{z}{|z|}, w\right) \widehat{Z}^\beta f(t(1-|z|), x + tz + t\hat{v}, w) dw \frac{t^3 dz}{|z|^2},$$

where we performed the change of variables $z = (y - x - t\hat{v})/t$ in order to obtain the second integral \mathcal{J} . As we shall see below, this splitting is useful in order to identify and isolate the asymptotic profile. We start by controlling the first term. For this, note that [\(53\)](#), $N_v \geq 10$ and the last two estimates of [Proposition 4.11](#), applied for $a = 1$, yield, for all $(\omega, \tau, y) \in \mathbb{S}^2 \times \mathbb{R}_+ \times \mathbb{R}^3$,

$$\left| \int_{\mathbb{R}_w^3} \mathcal{W}^T(\omega, w) Z^\beta f(\tau, y, w) dw \right| \lesssim \int_{\mathbb{R}_w^3} w^0 |Z^\beta f|(\tau, y, w) dw \lesssim \bar{\epsilon} \log^{3N_x+3N} (3+\tau) \frac{1+\max(\tau-|y|, 0)}{(1+\tau+|y|)^4}.$$

Note now that $|y - x| \geq t - |y - x - t\hat{v}|$ implies

$$t - |y - x - t\hat{v}| - |y| \leq t - |y - x - t\hat{v}| - |y - x| + |x| \leq |x|.$$

Hence, applying first the previous estimate for $\tau = t - |y - x - t\hat{v}|$ and then [\(46\)](#), we get

$$|t^2[\widehat{Z}^\beta f]^T(t, x + \hat{v}t) - \mathcal{J}| \lesssim \bar{\epsilon}(1+|x|)t^2 \int_{\substack{|y-x-t\hat{v}| \leq t \\ |y-x| \geq t - |y-x-t\hat{v}|}} \frac{\log^{3N_x+3N} (3+t - |y-x-t\hat{v}|)}{(1+t - |y-x-t\hat{v}| + |y|)^4} \frac{dy}{|y-x-t\hat{v}|^2}$$

$$\lesssim \bar{\epsilon} \langle x \rangle \frac{\log^{3N_x+3N} (3+|t-|x+t\hat{v}||)}{1+|t-|x+t\hat{v}||} t^2 Y_3^{p=2}(t, x + t\hat{v}).$$

According to [Lemma 5.11](#), $t^2 Y_3^{p=2}(t, x + t\hat{v}) \lesssim \log(1+t)$. By applying [Lemma 6.9](#), we then deduce

$$|t^2[\widehat{Z}^\beta f]^T(t, x + \hat{v}t) - \mathcal{J}| \lesssim \bar{\epsilon} \langle x \rangle \log(1+t) \left(\frac{\langle x \rangle |v^0|^2}{1+t} + |v^0|^2 \frac{\log^{3N_x+3N} (3+t)}{1+t} \right),$$

so that it remains for us to compare \mathcal{J} with $[\widehat{Z}^\beta f]^\infty(v)$. As in [Section 4.4](#), it is convenient to change the reference frame and work with $g^\beta(\tau, y, w) := \widehat{Z}^\beta f(\tau, y + \tau \hat{w}, w)$. In view of [Lemma 2.9](#), the change of variables $y = x + tz + \hat{v}t - \hat{w}t(1 - |z|)$, for z fixed, leads to

$$\mathcal{J} = \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \int_{|x-y+tz+\hat{v}t| < t(1-|z|)} \mathcal{W}^T\left(\frac{z}{|z|}, w\right) (|v^0|^5 g^\beta)(t(1-|z|), y, w) \frac{dy dz}{|z|^2(1-|z|)^3},$$

where we used w to denote the following function of (y, z) :

$$w = \frac{x - y + \widehat{tz + t\hat{v}}}{t(1-|z|)} \iff \hat{w} = \frac{x - y + tz + t\hat{v}}{t(1-|z|)}.$$

By the triangular inequality, we have $|\mathcal{J} - [\widehat{Z}^\beta f]^\infty| \leq \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3$, where

$$\begin{aligned} \mathcal{J}_1 &:= \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \int_{|x-y+tz+t\hat{v}| < t(1-|z|)} |\Delta_1^\beta| \frac{dy dz}{|z|^2(1-|z|)^3}, \\ \Delta_1^\beta &:= \mathcal{W}^T\left(\frac{z}{|z|}, w\right)(|v^0|^5 g^\beta)(t(1-|z|), y, w) - \mathcal{W}^T\left(\frac{z}{|z|}, \frac{z+\hat{v}}{1-|z|}\right)(|v^0|^5 g^\beta)\left(t(1-|z|), y, \frac{z+\hat{v}}{1-|z|}\right), \\ \mathcal{J}_2 &:= \left| \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \int_{|x-y+tz+t\hat{v}| \geq t(1-|z|)} \mathcal{W}^T\left(\frac{z}{|z|}, \frac{z+\hat{v}}{1-|z|}\right)(|v^0|^5 g^\beta)\left(t(1-|z|), y, \frac{z+\hat{v}}{1-|z|}\right) \frac{dy dz}{|z|^2(1-|z|)^3} \right|, \\ \mathcal{J}_3 &:= \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} |\Delta_3^\beta| \frac{dz}{|z|^2(1-|z|)^3}, \\ \Delta_3^\beta &:= \mathcal{W}^T\left(\frac{z}{|z|}, \frac{z+\hat{v}}{1-|z|}\right) \left[\int_{\mathbb{R}^3} (|v^0|^5 \widehat{Z}^\beta f)\left(t(1-|z|), y, \frac{z+\hat{v}}{1-|z|}\right) dy - (|v^0|^5 Q_\infty^\beta)\left(\frac{z+\hat{v}}{1-|z|}\right) \right], \end{aligned}$$

where, for Δ_3^β , we used that the spatial average of g^β is equal to the one of $\widehat{Z}^\beta f$. In view of [Remark 6.8](#), we will be able to transform time decay for the integrands of \mathcal{J}_i into decay in t , at the cost of powers of v^0 . In particular, [Remark 6.8](#) and $N_x > 7$ imply the following inequality that we will use several times:

$$\int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \int_{\mathbb{R}^3} \frac{dy}{\langle y \rangle^{N_x-4}} \frac{dz}{|z|^2(1-|z|)^n} \lesssim \int_{\substack{|z| \leq 1 \\ |z+\hat{v}| < 1-|z|}} \frac{dz}{|z|^2(1-|z|)^n} \leq 2^{2n+2}\pi |v^0|^{2n}, \quad n \in \mathbb{N}. \quad (54)$$

We start by dealing with \mathcal{J}_1 . Since $|\nabla_V \check{V}| \lesssim (1-|V|^2)^{-3/2} = |\check{V}^0|^3$ for all $|V| < 1$ by [Lemma 2.9](#) and in view of the bounds [\(53\)](#) on \mathcal{W}^T , the mean value theorem yields

$$\begin{aligned} |\Delta_1^\beta| &\leq \frac{|x-y|}{t(1-|z|)} \sup_{V \in \mathbb{R}^3} |V^0|^9 (|g^\beta| + |\nabla_V g^\beta|)(t(1-|z|), y, V) \\ &\leq \frac{1+|x|}{t(1-|z|)\langle y \rangle^{N_x-4}} \sup_{(X,V) \in \mathbb{R}^6} |V^0|^9 \langle X \rangle^{N_x-3} (|g^\beta| + |\nabla_V g^\beta|)(t(1-|z|), X, V). \end{aligned}$$

By applying [Lemma 2.8](#) and then the estimates of [Proposition 4.5](#), we obtain

$$|\Delta_1^\beta| \leq \frac{\langle x \rangle}{t(1-|z|)\langle y \rangle^{N_x-4}} \sum_{|\kappa| \leq N} \sup_{(X,V) \in \mathbb{R}^6} |V^0|^9 |z^{N_x-2} \widehat{Z}^\kappa f|(t(1-|z|), X, V) \lesssim \frac{\bar{\epsilon} \langle x \rangle \log^{3N_x+3N}(3+t)}{t(1-|z|)\langle y \rangle^{N_x-4}},$$

where we used $N_v \geq 12$ and $|\beta| + 1 \leq N$. We then deduce from [\(54\)](#) that

$$\mathcal{J}_1 \lesssim \bar{\epsilon} \langle x \rangle |v^0|^8 \frac{\log^{3N_x+3N}(3+t)}{t}.$$

Next, we control Δ_3^β using $|\mathcal{W}^T(\cdot, V)| \lesssim V^0$, $N_v \geq 12$ and [Proposition 6.1](#). This allows us to bound \mathcal{J}_3 through [\(54\)](#),

$$\Delta_3^\beta \lesssim \bar{\epsilon} \frac{\log^{3N_x+3N}(3+t)}{(1+t)(1-|z|)}, \quad \mathcal{J}_3 \lesssim \bar{\epsilon} |v^0|^8 \frac{\log^{3N_x+3N}(3+t)}{1+t}.$$

Finally, note that on the domain of integration of \mathcal{J}_2 , we have, for $\hat{w} = (z + \hat{v})/(1 - |z|)$,

$$1 = |w^0|^2 \left(1 - \frac{|z + \hat{v}|^2}{(1 - |z|)^2} \right) = |w^0|^2 \frac{(1 - |z| + |z + \hat{v}|)(1 - |z| - |z + \hat{v}|)}{(1 - |z|)^2} \leq |w^0|^2 \frac{2|x - y|}{(1 - |z|)t}.$$

Since $|\mathcal{W}^T|(\cdot, w) \lesssim w^0$, we get

$$\mathcal{J}_2 := \frac{\langle x \rangle}{t} \sup_{\tau \leq t} \sup_{(y, w) \in \mathbb{R}^6} |w^0|^8 \langle y \rangle^{N_x - 3} |g^\beta|(\tau, y, w) \int_{\substack{|z| \leq 1 \\ |z + \hat{v}| < 1 - |z|}} \int_{\mathbb{R}^3} \frac{dy}{\langle y \rangle^{N_x - 4}} \frac{dz}{|z|^2 (1 - |z|)^4}.$$

Using once again [Lemma 2.8](#) together with [Proposition 4.5](#), we get, in view of (54),

$$\mathcal{J}_2 \lesssim \bar{\epsilon} \langle x \rangle t^{-1} \log^{3N_x + 3N} (3 + t) |v^0|^8. \quad \square$$

This directly provides us the asymptotic profile of $\mathcal{L}_{Z^\gamma}(F)^T = -\sum_{|\beta| \leq |\gamma|} C_\beta^\gamma [\widehat{Z}^\beta f]^T$.

Corollary 6.11. *Let $|\gamma| \leq N - 1$ and $\mathcal{L}_{Z^\gamma}(F)^\infty := -\sum_{|\beta| \leq |\gamma|} C_\beta^\gamma [\widehat{Z}^\beta f]^\infty$. Then,*

$$\forall (t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3,$$

$$|t^2 \mathcal{L}_{Z^\gamma}(F)^T(t, x + \hat{v}t) - \mathcal{L}_{Z^\gamma}(F)^\infty(v)| \lesssim \bar{\epsilon} \langle x \rangle^2 |v^0|^8 \frac{\log^{3N_x + 3N + 1}(3 + t)}{t}.$$

Moreover, if Z^γ contains a translation ∂_{x^μ} or the scaling vector field S , then $\mathcal{L}_{Z^\gamma}(F)^\infty = 0$.

Proof. We only have to focus on the second part of the statement. Recall from the proof of [Proposition 6.4](#) that we can reduce the analysis to the cases $Z^\gamma = \partial_{x^\lambda} Z^\kappa$ if $\gamma_T \geq 1$, and $Z^\gamma = SZ^\kappa$ otherwise. Recall further from the commutation formula of [Lemma 2.3](#) and [Proposition 2.4](#) that

$$\nabla^\mu \mathcal{L}_{\partial_{x^\lambda} Z^\kappa}(F)_{\mu\nu} = \sum_{|\xi| \leq |\kappa|} C_\xi^\kappa J(\partial_{x^\lambda} \widehat{Z}^\xi f)_\nu, \quad \nabla^\mu \mathcal{L}_{SZ^\kappa}(F)_{\mu\nu} = \sum_{|\xi| \leq |\kappa|} C_\xi^\kappa J(S\widehat{Z}^\xi f)_\nu + 3C_\xi^\kappa J(\widehat{Z}^\xi f)_\nu.$$

It remains to recall from [Proposition 6.4](#) that $Q_\infty^{\partial_{x^\lambda} \xi} = 0$ and $Q_\infty^{S\xi} = -3Q^\xi$, so that $\mathcal{L}_{Z^\gamma}(F)^\infty = 0$. \square

6.2.2. Behavior of $\mathcal{L}_{Z^\gamma}(F)^{\text{data}}$ along timelike straight lines. Recall from [Proposition 5.3](#) and (50) that F^{data} is the sum of F^{hom} , which verifies $\square F_{\mu\nu}^{\text{hom}} = 0$, and a term which is strongly decaying in the interior of the light cone. If $Q_F \neq 0$, F decays initially as r^{-2} and one cannot expect to prove strong decay estimates for F^{hom} through [Proposition 2.21](#). For this reason, we need to analyse in detail the homogeneous part F^{hom} . It turns out that it decays faster in the interior of the light cone and then along timelike straight lines, so that it will not contribute to the asymptotic Lorentz force.

In order to improve the naive estimate of [Proposition 5.13](#), one can note that the leading-order term $\bar{F}(0, x) = Q_F x_i / (4\pi |x|^3) dt \wedge dx^i$ of the asymptotic expansion of $F^{\text{hom}}(0, \cdot)$ corresponds to the static electromagnetic field generated by a point charge Q_F located at $x = 0$. It is derived from the potential $A = Q(4\pi r)^{-1} dt$ which satisfies the Lorenz gauge, and then $\square A_\mu = 0$ on $\mathbb{R} \times (\mathbb{R}^3 \setminus \{0\})$. To deal with our evolution problem and the singularity of the Newton potential, we introduce

$$\tilde{A}(t, x) := \chi(|x| - t) A(t, x) = \frac{Q_F}{4\pi |x|} \chi(|x| - t) dt, \quad \chi \in C^\infty(\mathbb{R}, [0, 1]), \quad \chi|_{]-\infty, \frac{1}{2}[} = 0, \quad \chi|_{]1, +\infty[} = 1.$$

Then, \tilde{A} is smooth on $\mathbb{R}_+ \times \mathbb{R}^3$ and $\square \tilde{A}_\mu = 0$. It motivates the introduction of

$$\begin{aligned} \tilde{F}(t, x) &:= d\tilde{A}(t, x) = \frac{Q_{F^{X_i}}}{4\pi|x|^3} \chi(|x| - t) dt \wedge dx^i - \frac{Q_{F^{X_i}}}{4\pi|x|^2} \chi'(|x| - t) dt \wedge dx^i \\ &= \chi(|x| - t) \bar{F}(t, x) - \frac{Q_{F^{X_i}}}{4\pi|x|^2} \chi'(|x| - t) dt \wedge dx^i, \end{aligned}$$

which, in view of $[\square, \partial_{x^\lambda}] = 0$ and $\square \tilde{A}_\lambda = 0$, satisfies $\square \tilde{F}_{\mu\nu} = 0$. Since,

- for any $\Gamma \in \mathbb{K} \setminus \{S\}$, $[\square, \Gamma] = 0$ and $[\square, S] = 2\square$,
- for any $Z = Z^\lambda \partial_{x^\lambda} \in \mathbb{K}$ and any 2-form H , we have $\mathcal{L}_Z(H)_{\mu\nu} = Z(H_{\mu\nu}) + \partial_{x^\mu}(Z^\lambda)H_{\lambda\nu} + \partial_{x^\nu}(Z^\lambda)H_{\mu\lambda}$,

we then have $\square \mathcal{L}_{Z^\gamma}(\tilde{F})_{\mu\nu} = 0$ for any $|\gamma| \leq N-1$. The key idea will then be to consider $\mathcal{L}_{Z^\gamma}(F)^{\text{hom}} - \mathcal{L}_{Z^\gamma}(\tilde{F})$. More precisely, the following estimates hold.

Proposition 6.12. *For any $|\gamma| \leq N - 1$, we have,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^{\text{data}}(t, x) - \mathcal{L}_{Z^\gamma}(\tilde{F})(t, x)| \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1-\delta}.$$

Remark 6.13. We will not use it here, but we have

$$|\mathcal{L}_{Z^\gamma}(\tilde{F}) - \chi(|x| - t)\mathcal{L}_{Z^\gamma}(\bar{F})(t, x)| \lesssim Q_F(1+t)^{-1} \mathbb{1}_{0 \leq |x| - t \leq 1}.$$

Moreover, $\mathcal{L}_{\partial_t}(\bar{F}) = \mathcal{L}_{\Omega_{jk}}(\bar{F}) = \mathcal{L}_S(\bar{F}) = 0$ for all $1 \leq j < k \leq 3$. We refer to [Bigorgne 2020a, Section 5] for more information concerning \bar{F} .

This result implies that the leading-order term of $\mathcal{L}_{Z^\gamma}(F)^{\text{data}}(t, x)$ is supported in the exterior of the light cone. Before proving it, let us investigate its direct consequence for the behavior of F^{data} along timelike trajectories.

Proposition 6.14. *For any $|\gamma| \leq N - 1$, we have,*

$$\forall (t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |t^2 \mathcal{L}_{Z^\gamma}(F)^{\text{data}}(t, x + t\hat{v})| \lesssim \Lambda \langle x \rangle^2 |v^0|^4 t^{-\delta}.$$

Proof. Let $(t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$. If $t \leq 4\langle x \rangle |v^0|^2$, it suffices to apply Proposition 5.13, providing

$$|\mathcal{L}_{Z^\gamma}(F)^{\text{data}}(t, x + t\hat{v})| \lesssim \Lambda t^{-1} \leq 16\Lambda \langle x \rangle^2 |v^0|^4 t^{-3}.$$

Otherwise, according to Lemma 6.9, we have $t - |x + t\hat{v}| \geq t/(4|v^0|^2)$, so that $\chi^{(n)}(|x + t\hat{v}| - t) = 0$ for all $n \in \mathbb{N}$. Consequently, we get from Proposition 6.12 that

$$|\mathcal{L}_{Z^\gamma}(F)^{\text{data}}(t, x + t\hat{v})| \lesssim \Lambda t^{-1} (1 + |t - |x + t\hat{v}||)^{-1-\delta} \leq 16\Lambda |v^0|^4 t^{-3}. \quad \square$$

The first step of the proof of Proposition 6.12 consists in controlling the initial data for $\mathcal{L}_{Z^\gamma}(F)^{\text{hom}}$.

Lemma 6.15. *The assumption (15) on the initial electromagnetic field $F(0, \cdot)$ implies,*

$$\forall |\gamma| \leq N - 1, \quad \sup_{|\kappa| \leq 1} \sup_{|x| \geq 1} \langle x \rangle^{2+\delta+|\kappa|} |\nabla_{t,x}^\kappa \mathcal{L}_{Z^\gamma}(F)^{\text{hom}} - \nabla_{t,x}^\kappa \mathcal{L}_{Z^\gamma}(\bar{F})|(0, x) \lesssim \Lambda. \quad (55)$$

Note that $\nabla_{t,x}^\kappa \mathcal{L}_{Z^\gamma}(\bar{F})(0, x) = \nabla_{t,x}^\kappa \mathcal{L}_{Z^\gamma}(\tilde{F})(0, x)$ for all $|x| \geq 1$ since $\chi = 1$ on $[1, +\infty[$.

Proof. As \bar{F} is defined on $\mathbb{R} \times \mathbb{R}^3 \setminus \{0\}$, $\mathcal{L}_{Z^\gamma}(\bar{F})$ is well-defined for $|x| \geq 1$. We point out that $\nabla_t \mathcal{L}_{Z^\gamma}(\bar{F})(0, \cdot)$ does not necessarily vanish (consider for instance the case $Z^\gamma = \Omega_{01}$). Moreover, $\mathcal{L}_{Z^\gamma}(F)^{\text{hom}}(0, \cdot) = \mathcal{L}_{Z^\gamma}(F)(0, \cdot)$ by definition. Hence, the left-hand side of (55) is bounded by

$$\begin{aligned} \sup_{|\kappa| \leq 1} \sup_{|x| \geq 1} \langle x \rangle^{2+\delta+|\kappa|} |\nabla_{t,x}^\kappa \mathcal{L}_{Z^\gamma}(F - \bar{F})|(0, x) &\lesssim \sup_{|\xi| \leq |\gamma|+1} \sup_{|x| \geq 1} \langle x \rangle^{2+\delta+|\xi|} |\nabla_{t,x}^\xi (F - \bar{F})|(0, x) \\ &\leq \Lambda + \sup_{|\beta| \leq |\gamma|} \sup_{|x| \geq 1} \langle x \rangle^{2+\delta+n+|\beta|} |\nabla_{\partial_t} \nabla_{t,x}^\beta F|(0, x), \end{aligned} \tag{56}$$

where, in the last step, we used the assumption (15) and that \bar{F} is independent of t . Now, remark that if $n \geq 1$, the Maxwell equations implies

$$\partial_t(\partial_t^{n-1} \partial_x^\beta B) = -\partial_t^{n-1} \partial_x^\beta (\nabla_x \times E), \quad \partial_t(\partial_t^{n-1} \partial_x^\beta E) = \partial_t^{n-1} \partial_x^\beta (\nabla_x \times B) - \int_{\mathbb{R}^3} \hat{v} \partial_t^{n-1} \partial_x^\beta f \, dv.$$

Let \bar{E} and \bar{B} be the electric and magnetic field associated to \bar{F} according to (4), so that $\bar{E}^i = x^i Q_F / (4\pi r^3)$ and $\bar{B} = 0$. As $\nabla_x \times \bar{E} = \nabla_x \times \bar{B} = 0$, we can bound (56) by Λ by performing an induction and using (15) as well as the initial assumptions on f . \square

We are now able to prove Proposition 6.12 and conclude this subsection. As $\epsilon \leq \Lambda$, (50) implies,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^{\text{data}} - \mathcal{L}_{Z^\gamma}(F)^{\text{hom}}|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1-\delta}.$$

Finally, $\square \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^{\text{hom}} - \square \mathcal{L}_{Z^\gamma}(\tilde{F})_{\mu\nu} = 0$, the decay assumptions on the initial data given by Lemma 6.15 and Proposition 2.21 yield,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\mathcal{L}_{Z^\gamma}(F)^{\text{hom}} - \mathcal{L}_{Z^\gamma}(\tilde{F})|(t, x) \lesssim \Lambda(1+t+|x|)^{-1}(1+|t-|x||)^{-1-\delta}.$$

6.2.3. Self-similar asymptotic profile of $\mathcal{L}_{Z^\gamma}(F)$. We are now able to study the full Maxwell field.

Corollary 6.16. *For any $|\gamma| \leq N - 1$, there exists a 2-form $\mathcal{L}_{Z^\gamma}(F)^\infty$, independent of t , such that,*

$$\forall (t, x, v) \in [1, \infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |t^2 \mathcal{L}_{Z^\gamma}(F)(t, x + \hat{v}t) - \mathcal{L}_{Z^\gamma}(F)^\infty(v)| \lesssim \Lambda \langle x \rangle^2 |v^0|^8 \frac{\log^{3N_x+3N+1}(3+t)}{t^\delta}.$$

Moreover, for any $\eta > 0$, there exists $C_\eta > 0$ such that,

$$\forall (t, x) \in [1, +\infty[\times \mathbb{R}_x^3, \quad \frac{|x|}{t} \leq 1 - \eta, \quad \left| t^2 \mathcal{L}_{Z^\gamma}(F)(t, x) - \mathcal{L}_{Z^\gamma}(F)^\infty\left(\frac{\check{x}}{t}\right) \right| \lesssim \Lambda C_\eta \frac{\log^{3N_x+3N+1}(3+t)}{t^\delta}.$$

Remark 6.17. For the most important case, $|\gamma| = 0$, we have $4\pi F^\infty = -[f]^\infty$, where $[f]^\infty$ is explicitly written in Definition 6.6.

Proof. Fix $|\gamma| \leq N - 1$ and $(t, x, v) \in [1, \infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$. Applying Proposition 5.14 and Lemma 6.9, we have

$$t^2 |\mathcal{L}_{Z^\gamma}(F)^S|(t, x + \hat{v}t) \lesssim \bar{\epsilon} \Lambda \frac{t \log(3 + |t - |x - t\hat{v}||)}{(1 + |t - |x - t\hat{v}||)^2} \lesssim \Lambda \left(\frac{\langle x \rangle^2 |v^0|^4}{t} + |v^0|^4 \frac{\log(3+t)}{t} \right).$$

We then get the first part of the statement using the Glassey–Strauss decomposition given by [Proposition 5.3](#), [Corollary 6.11](#), where $\mathcal{L}_{Z^\gamma}(F)^\infty$ is introduced, and [Proposition 6.14](#). For the second part, it suffices to apply the first estimate, with a slight abuse of notation, for $x = 0$ and $\hat{v} = x/t$. \square

We deduce from the previous result a uniform bound on $\mathcal{L}_{Z^\gamma}(F)^\infty$. Moreover, it turns out that this quantity vanishes in certain cases, providing improved estimates for $\mathcal{L}_{Z^\gamma}(F)$.

Proposition 6.18. *For any $|\gamma| \leq N - 1$, we have $|\mathcal{L}_{Z^\gamma}(F)^\infty|(v) \lesssim \bar{\epsilon} \sqrt{v^0}$. Moreover, if $|\gamma| \geq 1$ and Z^γ contains a translation ∂_{x^μ} or the scaling vector field S , then $\mathcal{L}_{Z^\gamma}(F)^\infty = 0$.*

Proof. According to [Proposition 5.15](#), $t^2 |\mathcal{L}_{Z^\gamma}(F)^T|(t, t\hat{v}) \lesssim \bar{\epsilon} (1 - |\hat{v}|)^{-1/4} \leq 2\bar{\epsilon} \sqrt{v^0}$. All the properties then follow from [Corollary 6.11](#). \square

Finally, we investigate the regularity of $\mathcal{L}_{Z^\gamma}(F)^\infty$.

Proposition 6.19. *For any $|\gamma| \leq N - 2$ and $0 \leq \mu, \nu \leq 3$, $\mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu}$ is of class $C^{N-1-|\gamma|}$. Moreover, for any $1 \leq k \leq 3$, we have*

$$v^0 \partial_{v^k} \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu} = \mathcal{L}_{\Omega_{0k} Z^\gamma}(F)^\infty_{\mu\nu} + 2\hat{v}^k \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu} - \delta_\mu^0 \mathcal{L}_{Z^\gamma}(F)^\infty_{k\nu} - \delta_\mu^k \mathcal{L}_{Z^\gamma}(F)^\infty_{0\nu} - \delta_\nu^0 \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu k} - \delta_\nu^k \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu 0}.$$

The angular derivatives satisfy

$$(v^j \partial_{v^k} - v^k \partial_{v^j}) \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu} = \mathcal{L}_{\Omega_{jk} Z^\gamma}(F)^\infty_{\mu\nu} - \delta_\mu^j \mathcal{L}_{Z^\gamma}(F)^\infty_{k\nu} + \delta_\mu^k \mathcal{L}_{Z^\gamma}(F)^\infty_{j\nu} - \delta_\nu^j \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu k} + \delta_\nu^k \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu j}.$$

Proof. In order to lighten the notations, we introduce $X := x + t\hat{v}$. Then, we compute

$$\begin{aligned} &v^0 \partial_{v^k} (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(t, X)) \\ &= t(\delta_k^i - \hat{v}^k \hat{v}^i) \partial_{x^i} (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) \\ &= (\Omega_{0k} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) - X^k \partial_t (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) + \hat{v}^k (x^i - X^i) \partial_{x^i} (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) \\ &= (\Omega_{0k} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) - \hat{v}^k (S \mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) - x^k \partial_t (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) + \hat{v}^k x^i \partial_{x^i} (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X). \end{aligned}$$

One can already notice that the last two terms enjoy strong decay properties. More precisely, since [Lemma 6.9](#) implies $1 + |t - |X|| \gtrsim (1 + t)/(\langle x \rangle |v^0|^2)$, we have from [Proposition 3.2](#)

$$|t^2 - x^k \partial_t (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) + \hat{v}^k x^i \partial_{x^i} (\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X)| \lesssim \frac{\Lambda \langle x \rangle^3 |v^0|^4}{1 + t}.$$

The result then follows from

$$\mathcal{L}_{S Z^\gamma}(F)_{\mu\nu} = S(\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}) + 2\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}, \quad \mathcal{L}_{S Z^\gamma}(F)^\infty = 0, \tag{57}$$

$$\mathcal{L}_{\Omega_{0k} Z^\gamma}(F)_{\mu\nu} = \Omega_{0k}(\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}) + \delta_\mu^0 \mathcal{L}_{Z^\gamma}(F)_{k\nu} + \delta_\mu^k \mathcal{L}_{Z^\gamma}(F)_{0\nu} + \delta_\nu^0 \mathcal{L}_{Z^\gamma}(F)_{\mu k} + \delta_\nu^k \mathcal{L}_{Z^\gamma}(F)_{\mu 0}$$

and [Corollary 6.16](#), which give us

$$|t^2 v^0 \partial_{v^k} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(t, x + t\hat{v}) - v^0 \partial_{v^k} \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu}(v)| \lesssim \Lambda \langle x \rangle^3 |v^0|^8 \frac{\log^{1+3N_x+3N}(3+t)}{(1+t)^\delta},$$

where $v^0 \partial_{v^k} \mathcal{L}_{Z^\gamma}(F)^\infty_{\mu\nu}(v)$ is given in the statement of the proposition. To get the expression of the angular derivatives, notice that

$$(v^j \partial_{v^k} - v^k \partial_{v^j})(\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(t, X)) = (\Omega_{jk} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X) - (x^j \partial_{x^k} - x^k \partial_{x^j})(\mathcal{L}_{Z^\gamma}(F)_{\mu\nu})(t, X),$$

and

$$\mathcal{L}_{\Omega_{jk}Z^\gamma}(F)_{\mu\nu} = \Omega_{jk}(\mathcal{L}_{Z^\gamma}(F)_{\mu\nu}) + \delta_\mu^j \mathcal{L}_{Z^\gamma}(F)_{k\nu} - \delta_\mu^k \mathcal{L}_{Z^\gamma}(F)_{j\nu} + \delta_\nu^j \mathcal{L}_{Z^\gamma}(F)_{\mu k} - \delta_\nu^k \mathcal{L}_{Z^\gamma}(F)_{\mu j}$$

and apply the same arguments. The $C^{N-1-|\gamma|}$ regularity is obtained by an induction. □

For later use, we prove that the structure of the asymptotic Lorentz force is preserved by differentiation.

Corollary 6.20. *Let $0 \leq \nu \leq 3$ and define*

$$\Delta_{Z^\gamma, \nu}(t, x, v) := t^2 \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(t, x) - \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v), \quad |\gamma| \leq N - 1.$$

For any $|\gamma| \leq N - 2$, there holds

$$\begin{aligned} S(\Delta_{Z^\gamma, \nu}) &= \Delta_{SZ^\gamma, \nu}, \\ \widehat{\Omega}_{jk}(\Delta_{Z^\gamma, \nu}) &= \Delta_{\Omega_{jk}Z^\gamma, \nu} - \delta_\nu^j \Delta_{Z^\gamma, k} + \delta_\nu^k \Delta_{Z^\gamma, j}, \quad 1 \leq j < k \leq 3, \\ \widehat{\Omega}_{0i}(\Delta_{Z^\gamma, \nu}) &= \Delta_{\Omega_{0i}Z^\gamma, \nu} - \delta_\nu^0 \Delta_{Z^\gamma, i} - \delta_\nu^i \Delta_{Z^\gamma, 0} + 2 \frac{t}{v^0} (x^i - t \hat{v}^i) \hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}(t, x), \quad 1 \leq i \leq 3. \end{aligned}$$

Proof. The first identity follows from $S(t^2) = 2t^2$ and (57). For the other ones, start by noticing that, according to Proposition 6.19 and for $1 \leq i \leq 3$,

$$\begin{aligned} \widehat{\Omega}_{0i} \left(\frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v) \right) &= v^0 \partial_{v^i} \left(\frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v) \right) \\ &= \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{\Omega_{0i}Z^\gamma}(F)_{\mu\nu}^\infty(v) - \delta_\nu^0 \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu i}^\infty - \delta_\nu^i \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu 0}^\infty. \end{aligned} \tag{58}$$

Similarly, for $1 \leq j < k \leq 3$,

$$\widehat{\Omega}_{jk} \left(\frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v) \right) = \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{\Omega_{jk}Z^\gamma}(F)_{\mu\nu}^\infty(v) - \delta_\nu^j \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu k}^\infty + \delta_\nu^k \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu j}^\infty. \tag{59}$$

Recall that we denote by \mathbf{v} the 4-vector $(v^\mu)_{0 \leq \mu \leq 4}$, so that

$$\widehat{Z} \left(t^2 \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} \right) = \widehat{Z} \left(\frac{t^2}{|v^0|^2} \right) v^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} + \frac{t^2}{|v^0|^2} \mathcal{L}_{ZZ^\gamma}(F)(\mathbf{v}, \partial_{x^\nu}) + \frac{t^2}{|v^0|^2} \mathcal{L}_{Z^\gamma}(F)(\mathbf{v}, [Z, \partial_{x^\nu}]) \tag{60}$$

$$+ \frac{t^2}{|v^0|^2} \mathcal{L}_{Z^\gamma}(F)([Z, \mathbf{v}], \partial_{x^\nu}) + \frac{t^2}{|v^0|^2} \widehat{Z}(v^\mu) \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}. \tag{61}$$

- If $Z = \Omega_{0i}$, we have $[Z, \mathbf{v}] = -v^i \partial_t - v^0 \partial_{x^i}$ and $\widehat{Z}(v^\mu) = \delta_\mu^0 v^i + \delta_\mu^i v^0$, so that the sum of two terms in (61) vanishes. It remains to remark that $[Z, \partial_{x^\nu}] = -\delta_\nu^i \partial_t - \delta_\nu^0 \partial_i$, $\widehat{Z}(t^2/|v^0|^2) = 2t(x^i - t \hat{v}^i)/|v^0|^2$ and to combine (58) with (60).

- If $Z = \Omega_{jk}$, there holds $[Z, \mathbf{v}] = -v^j \partial_{x^k} + v^k \partial_{x^j}$ and $\widehat{Z}(v^\mu) = \delta_\mu^k v^j - \delta_\mu^j v^k$, so that the sum of the two terms in (61) vanishes once again. The result then ensues from $\widehat{Z}(t^2/|v^0|^2) = 0$, $[Z, \partial_{x^\nu}] = -\delta_\nu^j \partial_{x^k} + \delta_\nu^k \partial_{x^j}$, (59) and (60). □

6.3. Convergence of the distribution function along modified characteristics. Motivated by the discussion in Section 2.8.4 and by Corollary 6.16, we modify the linear spatial characteristics $t \mapsto x + t \hat{v}$ as follows.

Definition 6.21. For $(x, v) \in \mathbb{R}_x^3 \times \mathbb{R}_v^3$, let $X_{\mathcal{C}}(\cdot, x, v) : t \mapsto x + t\hat{v} + \mathcal{C}(t, v)$ be the trajectory¹¹

$$\begin{aligned} X_{\mathcal{C}}^i(t, x, v) &:= x^i + t\hat{v}^i - \log(t)\hat{v}^\mu F_{\mu}^{\infty, j}(v) \frac{\delta_j^i - \hat{v}_j \hat{v}^i}{v^0} \\ &= x^i + t\hat{v}^i - \frac{\log(t)}{v^0} (\hat{v}^\mu F_{\mu i}^{\infty}(v) + \hat{v}^i \hat{v}^\mu F_{\mu 0}^{\infty}(v)), \quad t \in \mathbb{R}_+^*, i \in \llbracket 1, 3 \rrbracket. \end{aligned} \tag{62}$$

For simplicity, we will often write $X_{\mathcal{C}}$ instead of $X_{\mathcal{C}}(t, x, v)$. By Proposition 6.18, the components \mathcal{C}^i of the correction term \mathcal{C} satisfy,

$$\forall t > 0, \quad |\mathcal{C}^i|(t, v) \lesssim \bar{\epsilon} |v^0|^{-\frac{1}{2}} \log(t), \quad i \in \llbracket 1, 3 \rrbracket. \tag{63}$$

We now bound the time derivative of a function evaluated along the modified characteristics.

Proposition 6.22. Let $f : \mathbb{R}_+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3 \rightarrow \mathbb{R}$ be a sufficiently regular function and introduce $h(t, x, v) := f(t, X_{\mathcal{C}}(t, x, v), v)$. Then, for all $(t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,

$$|\partial_t h|(t, x, v) \leq |\mathbf{T}_F(f)|(t, X_{\mathcal{C}}, v) + \Lambda \frac{\log^{3+3N_x+3N}(3+t)}{(1+t)^{1+\delta}} \sum_{\widehat{Z} \in \widehat{\mathbb{P}}_0} \|v^0\|^7 z^2 \widehat{Z} f|(t, X_{\mathcal{C}}, v).$$

Proof. We have, for all $(t, x, v) \in [1, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,

$$\begin{aligned} \partial_t h(t, x, v) &= (\partial_t f + \hat{v}^i \partial_{x^i} f)(t, X_{\mathcal{C}}, v) + \partial_t \mathcal{C}^i(t, v) \partial_{x^i} f(t, X_{\mathcal{C}}, v) \\ &= \mathbf{T}_F(f)(t, X_{\mathcal{C}}) - \hat{v}^\mu F_{\mu}^j(t, X_{\mathcal{C}}) \partial_{v^j} f(t, X_{\mathcal{C}}, v) + \partial_t \mathcal{C}^i(t, v) \partial_{x^i} f(t, X_{\mathcal{C}}, v). \end{aligned} \tag{64}$$

Recall from (14) the relation

$$v^0 \partial_{v^j} = -t(\partial_{x^j} - \hat{v}^j \hat{v}^i \partial_{x^i}) + \widehat{\Omega}_{0j} + z_{0j} \partial_t - \hat{v}^j S - \sum_{1 \leq i \leq 3} \hat{v}^j z_{0i} \partial_{x^i}, \quad 1 \leq j \leq 3, \tag{65}$$

in order to rewrite $\partial_{v^j} f(t, X_{\mathcal{C}}, v)$. As $v^0 \partial_t \mathcal{C}^i(t, v) = -(1/t) \hat{v}^\mu F_{\mu}^{\infty, j}(v) (\delta_j^i - \hat{v}_j \hat{v}^i)$, we get

$$\begin{aligned} |\partial_t h|(t, x, v) &\leq |\mathbf{T}_F(f)|(t, X_{\mathcal{C}}, v) + \sum_{1 \leq j \leq 3} \sum_{\widehat{Z} \in \widehat{\mathbb{P}}_0} |\hat{v}^\mu F_{\mu}^j|(t, X_{\mathcal{C}}) \left| \frac{z}{v^0} \widehat{Z} f \right|(t, X_{\mathcal{C}}, v) \\ &\quad + \frac{1}{t v^0} |t^2 F(t, X_{\mathcal{C}}) - F^\infty(v)| |\partial_{t,x} f|(t, X_{\mathcal{C}}, v). \end{aligned}$$

We deal with the second term on the right-hand side of the previous inequality by controlling the Lorentz force through Remark 4.3, so that $|\hat{v}^\mu F_{\mu}^j|(t, X_{\mathcal{C}}) \lesssim \Lambda(1+t)^{-2} |v^0|^2 z(t, X_{\mathcal{C}}, v)$. Next, by Corollary 6.16 and the mean value theorem,

$$\begin{aligned} |t^2 F(t, X_{\mathcal{C}}) - F^\infty(v)| &\leq |t^2 F(t, x + t\hat{v}) - F^\infty(v)| + t^2 |F(t, X_{\mathcal{C}}) - F(t, x + t\hat{v})| \\ &\lesssim \Lambda \langle x \rangle^2 |v^0|^8 \frac{\log^{3N_x+3N+1}(3+t)}{(1+t)^\delta} + t^2 |\mathcal{C}(t, v)| \sup_{|y - X_{\mathcal{C}}| \leq |\mathcal{C}(t, v)|} |\nabla_{t,x} F|(t, y). \end{aligned}$$

¹¹Recall that F^∞ is a 2-form, so that $\hat{v}^\mu \hat{v}^\nu F_{\mu\nu}^\infty = 0$.

In view of the estimate of $\nabla_{t,x}F$ given by Lemma 4.2 and the bound (63) on \mathcal{C} , we have

$$t^2|\mathcal{C}(t, v)| \sup_{|y-x|\leq|\mathcal{C}(t,v)} |\nabla_{t,x}F|(t, y) \lesssim \frac{\Lambda}{\sqrt{v^0}} t^2 \log(3+t) \frac{\log(3+t)}{(1+t)^3} |v^0|^4 \sup_{|y-X_{\mathcal{C}}|\leq|C|(t,v)} z^2(t, y, v).$$

Since $|\nabla_x z| \lesssim 1$, the mean value theorem yields

$$z(t, x + t\hat{v}, v) \leq \sup_{|y-X_{\mathcal{C}}|\leq|\mathcal{C}(t,v)} z(t, y, v) \leq z(t, X_{\mathcal{C}}, v) + \frac{\bar{\epsilon}}{\sqrt{v^0}} \log(3+t) \lesssim \log(3+t)z(t, X_{\mathcal{C}}, v). \quad (66)$$

Consequently, as $\langle x \rangle \leq z(t, x + t\hat{v}, v)$, we have

$$|t^2 F(t, X_{\mathcal{C}}) - F^\infty(v)| \lesssim \Lambda(1+t)^{-\delta} \log^{3N_x+3N+3}(3+t) |v^0|^8 z^2(t, X_{\mathcal{C}}, v).$$

We then deduce the result from the previous estimates. □

By applying this result to f , we obtain that there exists $f_\infty \in L^\infty_{x,v}$ such that $f(t, X_{\mathcal{C}}, v) \rightarrow f_\infty(x, v)$ as $t \rightarrow 0$ (see Proposition 6.34 for more details). Applying it again to $\partial_x^k f$ we could easily deduce that f_∞ is smooth with respect to the spatial variables. However, obtaining the regularity in the velocity variables requires a more thorough analysis. Indeed, $\partial_{v^i}(f(t, X_{\mathcal{C}}, v))$ is deeply related to $\widehat{\Omega}_{0i} f(t, X_{\mathcal{C}}, v)$, which does not converge.

6.4. Modified commutators. Let $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{\partial_t, \partial_{x^1}, \partial_{x^2}, \partial_{x^3}\}$ be a homogeneous vector field. Contrary to the case of the translations, the error term $[T_F, \widehat{Z}](f)$ does not decay sufficiently fast in order to prove a convergence result for $\widehat{Z}f$, even along the modified characteristics. Indeed, recall from Lemma 2.3 that

$$T_F(\widehat{Z}f) = -\hat{v}^\mu \mathcal{L}_Z(F)_{\mu^j} \partial_{v^j} f + \delta_{\widehat{Z}}^S \hat{v}^\mu F_{\mu^j} \partial_{v^j} f$$

and let us identify the terms with the slowest decay rate. Rewriting ∂_{v^j} by using (65) and estimating the electromagnetic field through Remark 4.3, we have

$$\left| T_F(\widehat{Z}f) - \frac{t}{v^0} (\hat{v}^\mu \mathcal{L}_Z(F)_{\mu^j} - \delta_{\widehat{Z}}^S \hat{v}^\mu F_{\mu^j}) (\delta_j^i - \hat{v}_j \hat{v}^i) \partial_{x^i} f \right| \lesssim \Lambda(1+t)^{-2} \sum_{\widehat{\Gamma} \in \widehat{\mathbb{P}}_0} v^0 |z^2 \widehat{\Gamma} f|. \quad (67)$$

In view of Proposition 4.5, the right-hand side is bounded by $\bar{\epsilon}(1+t)^{-2} \log^9(3+t)$ and then belongs to $L_t^1 L_{x,v}^\infty$. On the other hand, if $\mathcal{L}_Z(F)^\infty$ and F^∞ does not vanish, the decay rate of $t|\mathcal{L}_Z F| + t|F| \lesssim t^{-1}$ along timelike trajectories is at the threshold of time-integrability. For this reason, we modify the linear commutator \widehat{Z} in a way that is similar to how we modify the spatial characteristics. More precisely, motivated by Corollary 6.16 and (67), we introduce the following vector fields.

Definition 6.23. For any $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{\partial_t, \partial_{x^1}, \partial_{x^2}, \partial_{x^3}, S\}$, we define \widehat{Z}^{mod} and S^{mod} as

$$\widehat{Z}^{\text{mod}} := \widehat{Z} - \log(t) \hat{v}^\mu \mathcal{L}_Z(F)_{\mu^j}^{\infty,j}(v) \frac{\delta_j^i - \hat{v}_j \hat{v}^i}{v^0} \partial_{x^i}, \quad S^{\text{mod}} := S + \log(t) \hat{v}^\mu F_{\mu^j}^{\infty,j}(v) \frac{\delta_j^i - \hat{v}_j \hat{v}^i}{v^0} \partial_{x^i}.$$

We further define the correction coefficients $\mathcal{C}_S^i(t, v) = -\mathcal{C}^i(t, v)$ and

$$\mathcal{C}_{\widehat{Z}}^i(t, v) = -\log(t) \hat{v}^\mu \mathcal{L}_Z(F)_{\mu^j}^{\infty,j}(v) \frac{\delta_j^i - \hat{v}_j \hat{v}^i}{v^0} = -\frac{\log(t)}{v^0} (\hat{v}^\mu \mathcal{L}_Z(F)_{\mu^j}^{\infty,i}(v) + \hat{v}^i \hat{v}^\mu \mathcal{L}_Z(F)_{\mu^0}^{\infty}(v)),$$

so that $S^{\text{mod}} = S + \mathcal{C}_S^i(t, v) \partial_{x^i}$ and $\widehat{Z}^{\text{mod}} = \widehat{Z} + \mathcal{C}_{\widehat{Z}}^i(t, v) \partial_{x^i}$.

Remark 6.24. Recall that $t|\mathcal{L}_S(F)| \lesssim (1+t)^{-1-\delta}$ in domains of the form $\{t \geq (1-\delta)r\}$ since $\mathcal{L}_S(F)^\infty = 0$. This is why we do not need to compensate the term related to $\mathcal{L}_S(F)$ in (67).

We have the improved commutation relations.

Proposition 6.25. *Let $Z \in \mathbb{K}$ be a rotational vector field Ω_{jk} or a Lorentz boost Ω_{0i} . Then, for $t > 0$,*

$$[\mathbf{T}_F, \widehat{Z}^{\text{mod}}] = \frac{1}{t} \left(t^2 \widehat{v}^\mu (\mathcal{L}_Z(F)_\mu^j - \mathcal{L}_Z(F)_\mu^{\infty,j}) \frac{\delta_j^i - \widehat{v}_j \widehat{v}^i}{v^0} \right) \partial_{x^i} - \frac{\widehat{v}^\mu}{v^0} \mathcal{L}_Z(F)_\mu^j \left(\widehat{\Omega}_{0j} + z_{0j} \partial_t - \widehat{v}^j S - \sum_{1 \leq i \leq 3} \widehat{v}^j z_{0i} \partial_{x^i} \right) - \mathcal{C}_Z^i \widehat{v}^\mu \mathcal{L}_{\partial_{x^i}}(F)_\mu^j \partial_{v^j} + \widehat{v}^\mu F_\mu^j \partial_{v^j} \mathcal{C}_Z^i \partial_{x^i}.$$

For the scaling vector field, we have

$$[\mathbf{T}_F, S^{\text{mod}}] = -\frac{1}{t} \left(t^2 \widehat{v}^\mu (F_\mu^j - F_\mu^{\infty,j}) \frac{\delta_j^i - \widehat{v}_j \widehat{v}^i}{v^0} \right) \partial_{x^i} + \frac{1}{t} \left(t^2 \widehat{v}^\mu (\mathcal{L}_S(F)_\mu^j - \mathcal{L}_S(F)_\mu^{\infty,j}) \frac{\delta_j^i - \widehat{v}_j \widehat{v}^i}{v^0} \right) \partial_{x^i} + \frac{\widehat{v}^\mu}{v^0} (F_\mu^j - \mathcal{L}_S(F)_\mu^j) \left(\widehat{\Omega}_{0j} + z_{0j} \partial_t - \widehat{v}^j S - \sum_{1 \leq i \leq 3} \widehat{v}^j z_{0i} \partial_{x^i} \right) - \mathcal{C}_S^i \widehat{v}^\mu \mathcal{L}_{\partial_{x^i}}(F)_\mu^j \partial_{v^j} + \widehat{v}^\mu F_\mu^j \partial_{v^j} \mathcal{C}_S^i \partial_{x^i}.$$

Proof. Consider first the case $Z = \Omega_{jk}$ or $Z = \Omega_{0i}$. In view of the commutation relation of Lemma 2.3,

$$[\mathbf{T}_F, \widehat{Z}^{\text{mod}}] = \mathbf{T}_F(\mathcal{C}_Z^i) \partial_{x^i} + [\mathbf{T}_F, \widehat{Z}] + \mathcal{C}_Z^i [\mathbf{T}_F, \partial_{x^i}] = \mathbf{T}_F(\mathcal{C}_Z^i) \partial_{x^i} - \widehat{v}^\mu \mathcal{L}_Z(F)_\mu^j \partial_{v^j} - \mathcal{C}_Z^i \widehat{v}^\mu \mathcal{L}_{\partial_{x^i}}(F)_\mu^j \partial_{v^j}.$$

It then suffices to use (65) in order to rewrite ∂_{v^j} in the second term and to compute

$$\mathbf{T}_F(\mathcal{C}_Z^i) = -\frac{1}{t} \widehat{v}^\mu \mathcal{L}_Z(F)_\mu^{\infty,j} \frac{\delta_j^i - \widehat{v}_j \widehat{v}^i}{v^0} + \widehat{v}^\mu F_\mu^j \partial_{v^j} \mathcal{C}_Z^i.$$

The case of the scaling S can be treated similarly since $\mathcal{L}_S(F)^\infty = 0$ according to Proposition 6.18. \square

Apart from the term involving $\mathcal{L}_S(F)$, already discussed in Remark 6.24, it is clear that any of the error terms decay almost as $t^{-1-\delta}$ for, say, $|x| < t/2$. At this point, we could then prove that f_∞ is C^1 in v . However, since we would like to show $f_\infty \in C^{N-2}(\mathbb{R}_x^3 \times \mathbb{R}_v^3)$, we need to state a higher-order commutator formula for the modified vector fields. For this purpose, we introduce the set

$$\widehat{\mathbb{P}}_0^{\text{mod}} := \{ \partial_t, \partial_{x^i}, \widehat{\Omega}_{0i}^{\text{mod}}, \widehat{\Omega}_{jk}^{\text{mod}}, S^{\text{mod}} \mid 1 \leq i \leq 3, 1 \leq j < k \leq 3 \},$$

and we consider an ordering on it, so that $\widehat{\mathbb{P}}_0^{\text{mod}} = \{ \widehat{Z}^{\text{mod},i} \mid 1 \leq i \leq 11 \}$. Given a multi-index $\beta \in \llbracket 1, 11 \rrbracket^p$, we will then denote $\widehat{Z}^{\text{mod},\beta_1} \dots \widehat{Z}^{\text{mod},\beta_p}$ by $\widehat{Z}^{\text{mod},\beta}$. We will further denote by β_H (respectively β_T) the number of modified vector fields (respectively translations) composing $\widehat{Z}^{\text{mod},\beta}$, so that $|\beta| = \beta_H + \beta_T$. Furthermore, we will use the schematic notation $P_{p,q}(\mathcal{C})$ in order to denote any quantity of the form

$$\prod_{1 \leq k \leq p} \widehat{Z}^{\xi_k} (\mathcal{C}_{\widehat{Z}^k}^{i_k}), \quad (p, q) \in \mathbb{N}^2, \quad 1 \leq i_k \leq 3, \quad \widehat{Z}^k \in \widehat{\mathbb{P}}_0, \quad \sum_{1 \leq k \leq p} |\xi_k| = q, \quad q_T := \sum_{1 \leq k \leq q} \xi_{k,T}, \quad q_H := q - q_T,$$

where $q_T \geq 1$ when at least one translation ∂_{x^μ} is applied to at least one of the correction coefficients. By convention, we set $P_{0,0}(\mathcal{C}) = 1$ for $p = q = 0$. We recall from (10) the weights $z_{\lambda k} \in \mathbf{k}_1$, $0 \leq \lambda < k \leq 3$, which commute with the linear transport operator \mathbf{T}_0 .

Proposition 6.26. *Let $\widehat{Z}^{\text{mod},\beta} \in \widehat{\mathbb{P}}_0^{|\beta|}$. Then, $[\mathbf{T}_F, \widehat{Z}^{\text{mod},\beta}]$ can be written as a linear combination of the following types of terms:*

$$\frac{1}{v^0 t} R\left(\frac{1}{t}, \hat{v}, z\right) P_{p,q}(\mathcal{C}) (t^2 \hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} - \hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v)) \widehat{Z}^\kappa, \tag{T-1}$$

$$\frac{1}{v^0} R\left(\frac{1}{t}, \hat{v}, z\right) P_{p,q}(\mathcal{C}) \mathcal{L}_{Z^\gamma}(F)_{\lambda\nu} \widehat{Z}^\kappa, \tag{T-2}$$

$$\frac{x^\alpha}{v^0} R\left(\frac{1}{t}, \hat{v}, z\right) P_{p,q}(\mathcal{C}) \mathcal{L}_{Z^\gamma}(F)_{\lambda\nu} \widehat{Z}^\kappa, \quad q_T + \gamma_T \geq 1, \tag{T-3}$$

where R is a polynomial in $1/t$, $\hat{v} = (\hat{v}^i)_{1 \leq i \leq 3}$ and $z = (z_{\mu k})_{0 \leq \mu < k \leq 3}$, of degree $\text{deg}_z R$ in z , and

$$q_H + \text{deg}_z R \leq \beta_H, \quad p \leq \beta_H, \quad q + |\gamma| + |\kappa| \leq |\beta| + 1, \quad q, |\gamma|, |\kappa| \leq |\beta|, \quad 0 \leq \alpha, \lambda, \nu \leq 3.$$

Remark 6.27. In fact, we could prove that, as for the first-order commutation formula, most of the error terms satisfy a form of null condition. Since this property is not crucial for our purpose, we chose to demonstrate a result requiring a much simpler analysis.

Proof. Note first that the result holds for any $|\beta| = 1$. One can see it by applying either [Lemma 2.3](#), for the translation, or [Proposition 6.25](#) and by rewriting all the v derivatives as $v^0 \partial_{v_j} = \widehat{\Omega}_{0j} - t \partial_{x_j} - x^j \partial_t$. Let $n \geq 1$ such that the proposition holds for any $|\beta| = n$ and consider a multi-index $|\beta_0| = n + 1$. Consider further $|\beta| = n$ as well as $\widehat{Z}^{\text{mod}} \in \widehat{\mathbb{P}}_0^{\text{mod}}$ such that $\widehat{Z}^{\text{mod},\beta_0} = \widehat{Z}^{\text{mod}} \widehat{Z}^{\text{mod},\beta}$ and note

$$[\mathbf{T}_F, \widehat{Z}^{\text{mod},\beta_0}] = [\mathbf{T}_F, \widehat{Z}^{\text{mod}}] \widehat{Z}^{\text{mod},\beta} + \widehat{Z}^{\text{mod}} [\mathbf{T}_F, \widehat{Z}^{\text{mod},\beta}]. \tag{68}$$

We can deal with the first term on the right-hand side by applying the result for first-order operators and by noticing that $\widehat{Z}^\xi \widehat{Z}^{\text{mod},\beta}$, for $|\xi| \leq 1$, can be written as a linear combination of terms of the form

$$P_{p,q}(\mathcal{C}) \widehat{Z}^\zeta, \quad p \leq \beta_H, \quad q_H \leq \beta_H + \xi_H - 1, \quad q \leq |\beta| + |\xi| - 1, \quad q + |\zeta| \leq |\beta| + |\xi|. \tag{69}$$

For the second term, we apply the induction hypothesis, so that $[\mathbf{T}_F, \widehat{Z}^{\text{mod},\beta}]$ can be written as a linear combination of terms of the form [\(T-1\)](#)–[\(T-3\)](#). In order to deal with them, we will use the following properties:

- $\partial_t(t) = 1$, $\widehat{\Omega}_{0j}^{\text{mod}}(t) = x^j = -z_{0j} - t \hat{v}^j$, $S^{\text{mod}}(t) = t$ and $\widehat{Z}^{\text{mod}}(t) = 0$ otherwise.
- If $\widehat{Z}^{\text{mod}} = \partial_{x^\mu}$, then $\widehat{Z}^{\text{mod}}(x^k) = \delta_\mu^k$. Otherwise, there exists $0 \leq \lambda \leq 3$ such that $\widehat{Z}^{\text{mod}}(x^k) = \pm x^\lambda + \mathcal{C}_Z^k$.
- $\widehat{\Omega}_{0j}^{\text{mod}}(v^0) = v^j$ for any $1 \leq j \leq 3$ and $\widehat{Z}^{\text{mod}}(v^0) = 0$ otherwise.
- There exist four polynomials R_0, \dots, R_3 such that

$$\widehat{Z}^{\text{mod}}(R(1/t, \hat{v}, z)) = R_0(1/t, \hat{v}, z) + \mathcal{C}_Z^i R_i(1/t, \hat{v}, z), \quad \text{deg}_z R_0 \leq \text{deg}_z R + 1, \quad \text{deg}_z R_i \leq \text{deg}_z R,$$

where we set $\mathcal{C}_{\partial_{x^\mu}}^i := 0$. Moreover, if $\widehat{Z}^{\text{mod}} \neq \widehat{\Omega}_{0j}^{\text{mod}}$, then $\text{deg}_z R_0 \leq \text{deg}_z R$. This can be obtained by the first property and [\[Bigorgne 2020a, Lemma 3.2\]](#), giving

$$\forall \widehat{\Gamma} \in \widehat{\mathbb{P}}_0, \quad \forall 1 \leq i \leq 3, \quad \forall z \in \mathbf{k}_1, \quad \widehat{\Gamma}(v^0 z) \in \{0\} \cup \mathbf{k}_1, \quad \partial_{x^i}(z) \in \{0, 1, \hat{v}^k \mid 1 \leq k \leq 3\}.$$

- If $\widehat{Z}^{\text{mod}} = \partial_{x^\mu}$, we schematically have $\widehat{Z}^{\text{mod}}(P_{p,q}(\mathcal{C})) = P_{p,q^0}^0(\mathcal{C})$, with $q^0 = q + 1$ and $q_H^0 = q_H$. Otherwise, $\widehat{Z}^{\text{mod}}(P_{p,q}(\mathcal{C})) = P_{p,q^1}^1(\mathcal{C}) + P_{p+1,q^2}^2(\mathcal{C})$, with $q^1 = q^2 = q + 1$, $q_H^1 = q_H + 1$ and $q_H^2 = q_H$.
- $\widehat{Z}^{\text{mod}}\widehat{Z}^\kappa = \widehat{Z}\widehat{Z}^\kappa + \mathcal{C}_{\widehat{Z}}^i \partial_{x^i} \widehat{Z}^\kappa$ and $\widehat{Z}^{\text{mod}}\mathcal{L}_{Z^\gamma}(F)_{\lambda\nu}$ can be written as a linear combination of

$$\mathcal{L}_{ZZ^\gamma}(F)_{\lambda\nu}, \quad \mathcal{C}_{\widehat{Z}}^i \mathcal{L}_{\partial_{x^i} Z^\gamma}(F)_{\lambda\nu}, \quad \mathcal{L}_{Z^\gamma}(F)_{\mu\xi}, \quad 0 \leq \mu, \xi \leq 3.$$

Hence, we obtain by applying \widehat{Z}^{mod} to any quantity of the form (T-1), (T-2) or (T-3) (corresponding to $|\beta| = n$), a combination of terms of the form (T-1)–(T-3) (corresponding to $|\beta_0| = n + 1$), as well as

$$\mathcal{T}[\widehat{Z}^{\text{mod}}] = \frac{1}{t} R\left(\frac{1}{t}, \hat{v}, z\right) P_{p,q}(\mathcal{C}) \widehat{Z}^{\text{mod}} \left(t^2 \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} - \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\gamma}(F)_{\mu\nu}^\infty(v) \right) \widehat{Z}^\kappa,$$

where $0 \leq \nu \leq 3$, $q + |\gamma| + |\kappa| \leq |\beta| + 1$, $\max(q, |\gamma|, |\kappa|) \leq |\beta|$, $p \leq \beta_H$ and $q_H + \deg_z R \leq \beta_H$. Assume first that \widehat{Z}^{mod} is a translation ∂_{x^λ} . Then,

$$\mathcal{T}[\partial_{x^\lambda}] = \frac{2\delta_\lambda^0}{v^0} R(1/t, \hat{v}, z) P_{p,q}(\mathcal{C}) \hat{v}^\mu \mathcal{L}_{Z^\gamma}(F)_{\mu\nu} \widehat{Z}^\kappa + \frac{t}{v^0} R(1/t, \hat{v}, z) P_{p,q}(\mathcal{C}) \hat{v}^\mu \mathcal{L}_{\partial_{x^\lambda} Z^\gamma}(F)_{\mu\nu} \widehat{Z}^\kappa$$

is the sum of a term of type (T-2) and a term of type (T-3). Otherwise, $\widehat{Z}^{\text{mod}} = \widehat{Z} + \mathcal{C}_{\widehat{Z}}^i \partial_{x^i}$ and, following the previous computations, we have

$$\mathcal{T}[\widehat{Z}^{\text{mod}}] = \mathcal{T}[\widehat{Z}] + \mathcal{C}_{\widehat{Z}}^i \mathcal{T}[\partial_{x^i}] = \mathcal{T}[\widehat{Z}] + \frac{t}{v^0} R(1/t, \hat{v}, z) P_{p,q}(\mathcal{C}) \mathcal{C}_{\widehat{Z}}^i \hat{v}^\mu \mathcal{L}_{\partial_{x^i} Z^\gamma}(F)_{\mu\nu} \widehat{Z}^\kappa,$$

where the last three terms are of type (T-3). According to Corollary 6.20, $\mathcal{T}[\widehat{Z}]$ is a combination of terms of type (T-1) and, in the case $\widehat{Z} = \widehat{\Omega}_{0j}$, (T-2). □

We now control these error terms and then prove a uniform boundedness statement for $\widehat{Z}^{\text{mod},\beta} f$. Because of regularity issues on the coefficients $\mathcal{C}_{\widehat{Z}}$, which are of class C^{N-2} , we are not able to deal with the multi-indices $|\beta| \geq N - 1$.

Proposition 6.28. *Let $|\beta| \leq N - 2$. For all $(t, x, v) \in [3, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$, there holds*

$$|\mathbf{T}_F(\widehat{Z}^{\text{mod},\beta} f)|(t, x, v) \lesssim \Lambda \frac{\log^{3N_x+4N}(t)}{t^{1+\delta}} \sum_{|\kappa| \leq |\beta|} |v^0|^7 |z^{2+\beta_H} \widehat{Z}^\kappa f|(t, x, v).$$

Moreover,

$$|v^0|^{N_v-7} |z^{N_x-2-\beta_H} \mathbf{T}_F(\widehat{Z}^{\text{mod},\beta} f)|(t, x, v) \lesssim \bar{\epsilon} \frac{\log^{6N_x+7N}(t)}{t^{1+\delta}}.$$

Proof. Fix $(t, x, v) \in [3, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$ and let us prove first the following property. Consider $P_{p,q}(\mathcal{C})$ and $R(1/t, \hat{v}, z)$ a polynomial such that $p \leq \beta_H$, $q \leq |\beta|$ and $q_H + \deg_z R \leq \beta_H$. Then,

$$|R(1/t, \hat{v}, z)| |P_{p,q}(\mathcal{C})|(t, x, v) \lesssim \frac{\log^{N-2}(t)}{t^{q_T}} z^{\beta_H}(t, x, v). \tag{70}$$

For this, remark first that, for $|\xi| \leq N - 2$, $i \in \llbracket 1, 3 \rrbracket$ and $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{\partial_t, \partial_{x^1}, \partial_{x^2}, \partial_{x^3}, S\}$,

$$|\widehat{Z}^\xi(\mathcal{C}_{\widehat{Z}}^i)|(t, x, v) \leq \sum_{\substack{|\gamma|+|\kappa| \leq |\xi| \\ \gamma_T = \xi_T}} \mathcal{I}_{\gamma,\kappa}, \quad \mathcal{I}_{\gamma,\kappa} := \sum_{0 \leq \nu \leq 3} |\widehat{Z}^\gamma \log(t)| \left| \widehat{Z}^\kappa \left(\frac{\hat{v}^\mu}{v^0} \mathcal{L}_Z(F)_{\mu\nu}^\infty \right) \right|(v).$$

Note that the case $\widehat{Z} = S$ leads to a similar estimate.

• We have $|\widehat{Z}^\gamma \log(t)| \lesssim t^{-\xi_T} \mathbf{z}^{\xi_H}(t, x, v) \log(t)$. Indeed, $|\widehat{Z}^\gamma \log(t)| \leq |t^{-\gamma_T} P_{\gamma_H}(x/t) \log(t)|$, where P_{γ_H} is a polynomial of degree at most $\gamma_H \leq \xi_H$, and $\gamma_T = \xi_T$. Finally, recall that $|x|/t \leq |x - t\hat{v}|/t + 1 \leq 2z(t, x, v)$.

• To deal with the last factor in $\mathcal{I}_{\gamma, \kappa}$, note first that $|\kappa| + 1 \leq N - 1$ and that this quantity vanishes if κ is composed of at least a translation or the scaling vector field S according to Proposition 6.18. Then, using first the relations (58)–(59) and then Proposition 6.18, we get

$$\left| \widehat{Z}^\kappa \left(\frac{\hat{v}^\mu}{v^0} \mathcal{L}_Z(F)_{\mu\nu}^\infty \right) \right| (v) \lesssim \sum_{|\zeta| \leq |\kappa| + 1} \left| \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{Z^\zeta}(F)_{\mu\nu}^\infty \right| (v) \lesssim \bar{\epsilon}. \tag{71}$$

We then deduce that

$$|R(1/t, \hat{v}, z)| |P_{p,q}(\mathcal{C})|(t, x, v) \lesssim \mathbf{z}^{\deg_z R}(t, x, v) t^{-qT} \mathbf{z}^{qH}(t, x, v) \log^p(t) \bar{\epsilon}^p,$$

which implies (70).

Apply Proposition 6.26 in order to reduce the analysis to the treatment of terms of type (T-1), (T-2) and (T-3). By Corollary 6.16 and (70), we can bound any term of type (T-1) by

$$\Lambda \frac{|v^0|^8 \log^{3N_x + 4N}(t)}{v^0 t^{1+\delta}} \langle x - t\hat{v} \rangle^2 |\mathbf{z}^{\beta_H} \widehat{Z}^\kappa f|(t, x, v) \lesssim \Lambda \frac{\log^{3N_x + 4N}(t)}{t^{1+\delta}} |v^0|^7 |\mathbf{z}^{2+\beta_H} \widehat{Z}^\kappa f|(t, x, v),$$

since $\langle x - t\hat{v} \rangle \leq z(t, x, v)$ and where $|\kappa| \leq N - 2$. We deal with the ones of type (T-2) by using (BA1), (70) and Lemma 2.6. There are bounded above by

$$\frac{\Lambda \log^{N-2}(t)}{(t + |x|)(1 + |t - |x||)v^0} \frac{(1 + |t - |x||)|v^0|^2 z}{t + |x|} |\mathbf{z}^{\beta_H} \widehat{Z}^\kappa f|(t, x, v) \lesssim \Lambda \frac{\log^{N-2}(t)}{t^2} v^0 |\mathbf{z}^{1+\beta_H} \widehat{Z}^\kappa f|(t, x, v).$$

Finally, let \mathcal{T}_3 be a term of type (T-3). Using first (70) together with Proposition 3.2 and then Lemma 2.6,

$$\mathcal{T}_3 \lesssim \frac{\Lambda \log^{N-2}(t)}{v^0 t^{qT} (1 + |t - |x||)^{1+\gamma_T}} |\mathbf{z}^{\beta_H} \widehat{Z}^\kappa f|(t, x, v) \lesssim \Lambda \frac{\log^N(t)}{t^2} |v^0|^3 |\mathbf{z}^{2+\beta_H} \widehat{Z}^\kappa f|(t, x, v).$$

We deduce from that the first estimate of the statement, which, through an application of Proposition 4.5, implies the second one. □

Corollary 6.29. *Let $|\beta| \leq N - 2$. If $\beta_H \leq N_x - 2$, there exists $\bar{D} > 0$ such that,*

$$\forall t \geq 3, \quad \| |v^0|^{N_v - 7} \widehat{Z}^{\text{mod}, \beta} f(t, \cdot, \cdot) \|_{L_{x,v}^\infty} \lesssim \epsilon e^{\bar{D}\Lambda}. \tag{72}$$

Proof. Note first that we can obtain, by a much simpler analysis than in the proof of Proposition 4.5, that $\| |v^0|^{N_v} \mathbf{z}^{N_x} \widehat{Z}^\beta f(3, \cdot, \cdot) \|_{L_{x,v}^\infty} \lesssim \bar{\epsilon}$ for all $|\beta| \leq N$. Consequently, using (69) and (71), we get,

$$\forall |\beta| \leq N - 1, \quad \| |v^0|^{N_v} \mathbf{z}^{N_x} \widehat{Z}^{\text{mod}, \beta} f(3, \cdot, \cdot) \|_{L_{x,v}^\infty} \lesssim \sum_{|\kappa| \leq |\beta|} \| |v^0|^{N_v} \mathbf{z}^{N_x} \widehat{Z}^\kappa f(3, \cdot, \cdot) \|_{L_{x,v}^\infty} \lesssim \bar{\epsilon}.$$

Hence, it suffices to prove, according to Lemma 4.4, that

$$|\mathbf{T}_F(|v^0|^{N_v - 7} \widehat{Z}^{\text{mod}, \beta} f)|(t, x, v) \lesssim \left(\frac{\Lambda |v^0|^{N_v - 7} |\widehat{Z}^{\text{mod}, \beta} f|}{(1+t)^{\frac{3}{2}}} + \frac{\Lambda \hat{v}^L |v^0|^{N_v - 7} |\widehat{Z}^{\text{mod}, \beta} f|}{(1+|t - |x||)^2} \right) + \frac{\bar{\epsilon}}{(1+t) \log^2(3+t)}$$

for all $(t, x, v) \in [3, T[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$ and any $|\beta| \leq N - 2$. For this, we bound $T_F(v^0)$ using (31) and we apply the previous Proposition 6.28 in order to control $T_F(\widehat{Z}^{\text{mod},\beta} f)$. \square

6.5. Regularity of the asymptotic state. In order to prove that f_∞ is differentiable with respect to v , we will need to compute the first-order v -derivatives of the correction terms in the modified spatial characteristics and to bound their higher-order derivatives.

Lemma 6.30. *Let $(i, k) \in \llbracket 1, 3 \rrbracket^2$. Then, for all $(t, x, v) \in [3, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,*

$$v^0 \partial_{v^k} \mathcal{C}^i(t, v) = \mathcal{C}_{\Omega_{0k}}^i(t, v) - \hat{v}^i \mathcal{C}^k(t, v).$$

More generally, for any multi-index $|\kappa| \leq N - 1$,

$$|v^0|^{|\kappa|} |\partial_v^\kappa \mathcal{C}^i|(t, v) \lesssim \bar{\epsilon} |v^0|^{-\frac{1}{2}} \log(t).$$

Proof. According to (58), we have, for any $v \in \llbracket 0, 3 \rrbracket$,

$$v^0 \partial_{v^k} \left(\frac{\hat{v}^\mu}{v^0} F_{\mu v}^\infty \right) = \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{\Omega_{0k}}(F)_{\mu v}^\infty - \delta_v^0 \frac{\hat{v}^\mu}{v^0} F_{\mu k}^\infty - \delta_v^k \frac{\hat{v}^\mu}{v^0} F_{\mu 0}^\infty.$$

This implies in particular that

$$v^0 \partial_{v^k} \left(\frac{\hat{v}^i \hat{v}^\mu}{v^0} F_{\mu 0}^\infty + \frac{\hat{v}^\mu}{v^0} F_{\mu i}^\infty \right) = \frac{\hat{v}^i \hat{v}^\mu}{v^0} \mathcal{L}_{\Omega_{0k}}(F)_{\mu 0}^\infty + \frac{\hat{v}^\mu}{v^0} \mathcal{L}_{\Omega_{0k}}(F)_{\mu i}^\infty - \hat{v}^i \left(\frac{\hat{v}^k \hat{v}^\mu}{v^0} F_{\mu 0}^\infty + \frac{\hat{v}^\mu}{v^0} F_{\mu k}^\infty \right).$$

In view of the definition of the correction coefficients (see Definitions 6.21 and 6.23), we deduce from this last equality the first part of the statement. The second part follows from a direct induction as well as Propositions 6.18–6.19. \square

Remark 6.31. Similarly, we could prove using (59) that $\Omega_{jk}^v \mathcal{C}^i(t, v) = \mathcal{C}_{\Omega_{jk}^i}^i(t, v) - \delta_j^i \mathcal{C}^k(t, v) + \delta_k^i \mathcal{C}^j(t, v)$, where $\Omega_{jk}^v := v^j \partial_{v^k} - v^k \partial_{v^j}$. Consequently, the following quantities, related to the asymptotic Lorentz force,

$$\Gamma(v) := \frac{\hat{v}^\mu}{v^0} (F_{\mu i}^\infty(v) + \hat{v}_i F_{\mu 0}^\infty(v)) dv^i, \quad \Gamma_Z(v) := \frac{\hat{v}^\mu}{v^0} (\mathcal{L}_Z(F)_{\mu i}^\infty(v) + \hat{v}_i \mathcal{L}_Z(F)_{\mu 0}^\infty(v)) dv^i,$$

satisfy $\mathcal{L}_{v^0 \partial_{v^k}}(\Gamma) = \Gamma_{\Omega_{0k}}$ and $\mathcal{L}_{\Omega_{jk}^v}(\Gamma) = \Gamma_{\Omega_{jk}}$.

We now perform a computation, which holds for any sufficiently regular function f . In particular, we will apply it to $f = \partial_{t,x}^k f$. We have

$$v^0 \partial_{v^k} (f(t, X_\mathcal{C}, v)) = t \partial_{x^k} f(t, X_\mathcal{C}, v) - t \hat{v}^k \hat{v}^i \partial_{x^i} f(t, X_\mathcal{C}, v) + v^0 \partial_{v^k} f(t, X_\mathcal{C}, v) + v^0 \partial_{v^k} \mathcal{C}^i(t, v) \partial_{x^i} f(t, X_\mathcal{C}, v).$$

Then, we use (65) in order to rewrite the third term on the right-hand side. We get

$$v^0 \partial_{v^k} (f(t, X_\mathcal{C}, v)) = \left(\widehat{\Omega}_{0k} f + z_{0k} \partial_t f - \hat{v}^k S f - \hat{v}^k \sum_{1 \leq i \leq 3} z_{0i} \partial_{x^i} f \right) (t, X_\mathcal{C}, v) + v^0 \partial_{v^k} \mathcal{C}^i(t, v) \partial_{x^i} f(t, X_\mathcal{C}, v).$$

Hence, as $z_{0i}(t, X_{\mathcal{C}}, v) = -x^i - \mathcal{C}^i(t, v)$,

$$\begin{aligned} v^0 \partial_{v^k}(\mathbf{f}(t, X_{\mathcal{C}}, v)) &= (\widehat{\Omega}_{0k} \mathbf{f})(t, X_{\mathcal{C}}, v) - x^k (\partial_t \mathbf{f})(t, X_{\mathcal{C}}, v) - \frac{\mathcal{C}^k(t, v)}{t} (S \mathbf{f})(t, X_{\mathcal{C}}, v) \\ &\quad + \frac{\mathcal{C}^k(t, v)}{t} X_{\mathcal{C}}^i \partial_{x^i} \mathbf{f}(t, X_{\mathcal{C}}, v) - \hat{v}^k (S \mathbf{f})(t, X_{\mathcal{C}}, v) + \hat{v}^k \mathcal{C}^i(t, v) \partial_{x^i} \mathbf{f}(t, X_{\mathcal{C}}, v) \\ &\quad + \hat{v}^k x^i \partial_{x^i} \mathbf{f}(t, X_{\mathcal{C}}, v) + v^0 \partial_{v^k} \mathcal{C}^i(t, v) \partial_{x^i} \mathbf{f}(t, X_{\mathcal{C}}, v). \end{aligned}$$

Now, according to [Lemma 6.30](#),

$$\widehat{\Omega}_{0k} + v^0 \partial_{v^k} \mathcal{C}^i(t, v) \partial_{x^i} = \widehat{\Omega}_{0k} + \mathcal{C}_{\Omega_{0k}}^i(t, v) \partial_{x^i} - \mathcal{C}^k(t, v) \hat{v}^i \partial_{x^i} = \widehat{\Omega}_{0k}^{\text{mod}} - \mathcal{C}^k(t, v) \hat{v}^i \partial_{x^i},$$

and, in view of the relations $S^{\text{mod}} = S - \mathcal{C}^i(t, v) \partial_{x^i}$ and $X_{\mathcal{C}}^i = x^i + t \hat{v}^i + \mathcal{C}^i(t, v)$,

$$\begin{aligned} v^0 \partial_{v^k}(\mathbf{f}(t, X_{\mathcal{C}}, v)) &= (\widehat{\Omega}_{0k}^{\text{mod}} \mathbf{f})(t, X_{\mathcal{C}}, v) - \left(\hat{v}^k + \frac{\mathcal{C}^k(t, v)}{t} \right) (S^{\text{mod}} \mathbf{f})(t, X_{\mathcal{C}}, v) \\ &\quad - x^k (\partial_t \mathbf{f})(t, X_{\mathcal{C}}, v) + \left(\hat{v}^k + \frac{\mathcal{C}^k(t, v)}{t} \right) x^i \partial_{x^i} \mathbf{f}(t, X_{\mathcal{C}}, v). \quad (73) \end{aligned}$$

Iterating this process to the functions $\mathbf{f} = \partial_{t,x}^{\kappa} f$ yields the following result.

Proposition 6.32. *Let $|\kappa| + |\xi| \leq N - 2$. Then, there exist functions $P_{\beta}^{\kappa, \xi}$ such that,*

$$\forall (t, x, v) \in [3, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \quad |v^0|^{|\xi|} \partial_v^{\xi} ((\partial_{t,x}^{\kappa} f)(t, X_{\mathcal{C}}, v)) = \sum_{|\beta| \leq |\kappa| + |\xi|} P_{\beta}^{\kappa, \xi}(t, x, v) \widehat{Z}^{\text{mod}, \beta} f(t, X_{\mathcal{C}}, v)$$

and $P_{\beta}^{\kappa, \xi}(t, x, v)$ is a linear combination of terms of the form $P(x, \hat{v})M(\mathcal{C})$, where P is a polynomial and

$$M(\mathcal{C}) = \prod_{k=1}^d \frac{1}{t} |v^0|^{|\xi_k|} \partial_v^{\xi_k} \mathcal{C}^{i_k}(t, v), \quad d + \sum_{1 \leq k \leq d} |\xi_k| \leq |\xi|, \quad |\beta| + \sum_{1 \leq k \leq d} |\xi_k| \leq |\xi|, \quad \deg_x(P) + \beta_H \leq |\xi|.$$

The value $d = 0$ is allowed, in which case we set $M(\mathcal{C}) = 1$.

In order to prove, through [Proposition 6.22](#), that the functions considered in the previous statement converge, as $t \rightarrow +\infty$, we will be lead to estimate these polynomials and their time derivative.

Lemma 6.33. *Let $|\kappa| + |\xi| \leq N - 2$ and $|\beta| \leq |\kappa| + |\xi|$. Then, for all $(t, x, v) \in [3, +\infty[\times \mathbb{R}_x^3 \times \mathbb{R}_v^3$,*

$$|P_{\beta}^{\kappa, \xi}|(t, x, v) \lesssim \langle x \rangle^{|\xi| - \beta_H}, \quad |\partial_t P_{\beta}^{\kappa, \xi}|(t, x, v) \lesssim \bar{\epsilon} \langle x \rangle^{|\xi| - \beta_H} \frac{\log(t)}{t^2}.$$

Proof. It is enough to bound terms of the form $P(x, \hat{v})M(\mathcal{C})$ satisfying the conditions given in [Proposition 6.32](#). The first factor satisfies $|P(x, \hat{v})| \lesssim \langle x \rangle^{\deg_x P} \leq \langle x \rangle^{|\xi| - \beta_H}$ and does not depend on t . In view of [Lemma 6.30](#), we have $|M(\mathcal{C})| \lesssim \bar{\epsilon}^d \log^d(t) t^{-d}$, which implies the first estimate. The second one can be obtained similarly. Either $|\partial_t M(\mathcal{C})| = 0$ or $d \geq 1$ and $|\partial_t M(\mathcal{C})| \lesssim \bar{\epsilon}^d \log^d(t) t^{-d-1}$ by [Lemma 6.30](#). \square

We are now able to prove the main result of this paper. For this, let us introduce

$$h : (t, x, v) \mapsto f(t, x + t \hat{v} + \mathcal{C}(t, v), v), \quad h^{\xi, \kappa} := |v^0|^{|\xi|} \partial_v^{\xi} \partial_x^{\kappa} h(t, x, v) = |v^0|^{|\xi|} \partial_v^{\xi} (\partial_x^{\kappa} f(t, X_{\mathcal{C}}(t, x, v), v)).$$

Proposition 6.34. *There exists a function $f_\infty \in C^{N-2}(\mathbb{R}_x^3 \times \mathbb{R}_v^3, \mathbb{R}_+)$ such that, for any $|\kappa| + |\xi| \leq N - 2$,*

$$\forall t \geq 3, \quad \| |v^0|^{N_v-10+|\xi|} \langle x \rangle^{N_x-4-|\xi|} (\partial_v^\xi \partial_x^\kappa h(t, \cdot, \cdot) - \partial_v^\xi \partial_x^\kappa f_\infty) \|_{L_{x,v}^\infty} \lesssim \bar{\epsilon} \frac{\log^{7(N_x+N)}(t)}{t^\delta}.$$

In particular, as $N_v > 13$ and if $N_x > 7 + |\xi|$, we have $\partial_v^\xi \partial_x^\kappa f_\infty \in L_{x,v}^1$.

Proof. Fix $t \geq 3$ and $(x, v) \in \mathbb{R}_x^3 \times \mathbb{R}_v^3$. Applying the previous [Proposition 6.32](#) and [Lemma 6.33](#), we get

$$|\partial_t h^{\xi,\kappa}|(t, x, v) \lesssim \sum_{\substack{|\beta| \leq N-2 \\ \beta_H \leq |\xi|}} \langle x \rangle^{|\xi|-\beta_H} |\partial_t \widehat{Z}^{\text{mod},\beta} f|(t, X_\ell, v) + \bar{\epsilon} \frac{\log(t)}{t^2} \langle x \rangle^{|\xi|-\beta_H} |\widehat{Z}^{\text{mod},\beta} f|(t, X_\ell, v).$$

Next, we recall from (66) the inequality $\langle x \rangle \lesssim \log(t)z(t, X_\ell, v)$ and note, using the same arguments, that $z(t, X_\ell, v) \lesssim \log(t)\langle x \rangle$ holds as well. Bounding $\partial_t \widehat{Z}^{\text{mod},\beta} f$ by [Proposition 6.22](#), we then get

$$\begin{aligned} |v^0|^{N_v-10} \langle x \rangle^{N_x-4-|\xi|} |\partial_t h^{\kappa,\xi}|(t, x, v) &\lesssim \sum_{\substack{|\beta| \leq N-2 \\ \beta_H \leq |\xi|}} \log^{N_x}(t) |v^0|^{N_v-10} |z^{N_x-4-\beta_H} \mathbf{T}_F(\widehat{Z}^{\text{mod},\beta} f)|(t, X_\ell, v) \\ &\quad + \Lambda \frac{\log^{4N_x+3N}(t)}{t^{1+\delta}} \sum_{|\gamma| \leq 1} |v^0|^{N_v-3} |z^{N_x-2-\beta_H} \widehat{Z}^\gamma \widehat{Z}^{\text{mod},\beta} f|(t, X_\ell, v). \end{aligned}$$

We control the first term on the right-hand side by [Proposition 6.28](#) and we claim that the second one is bounded by

$$\Lambda \frac{\log^{4N_x+4N}(t)}{t^{1+\delta}} \sum_{|\kappa| \leq N-1} |v^0|^{N_v-3} |z^{N_x-2} \widehat{Z}^\kappa f|(t, X_\ell, v).$$

Indeed, we rewrite the modified vector fields using (69) and we control $P_{p,q}(\mathcal{C})$ by (70). We then deduce from [Proposition 4.5](#) that

$$|v^0|^{N_v-10} \langle x \rangle^{N_x-4-|\xi|} |\partial_t h^{\kappa,\xi}|(t, x, v) \lesssim \bar{\epsilon} \frac{\log^{7N_x+7N}(t)}{t^{1+\delta}}.$$

We obtain from that,

$$\forall 3 \leq t \leq \tau, \quad \left| |v^0|^{N_v-10} \langle x \rangle^{N_x-4-|\xi|} (h^{\kappa,\xi}(\tau, x, v) - h^{\kappa,\xi}(t, x, v)) \right| \lesssim \bar{\epsilon} \frac{\log^{7(N_x+N)}(t)}{t^\delta}. \tag{74}$$

Consequently, there exists $f_\infty^{\kappa,\xi} \in L_{x,v}^\infty$ such that $h^{\kappa,\xi}(t, \cdot, \cdot) \rightarrow f_\infty^{\kappa,\xi}$ as $t \rightarrow +\infty$, uniformly on any compact subset of $\mathbb{R}_x^3 \times \mathbb{R}_v^3$. By uniqueness of the limit in $\mathcal{D}'(\mathbb{R}_x^3 \times \mathbb{R}_v^3)$ and by continuity of the distributional partial derivatives, we get $f_\infty^{\kappa,\xi} = |v^0|^{|\xi|} \partial_v^\xi \partial_x^\kappa f_\infty$. Letting $\tau \rightarrow +\infty$ in (74) yields the stated rate of convergence and concludes the proof. \square

Remark 6.35. We can improve the result for f_∞ . [Propositions 4.5](#) and [6.22](#) give,

$$\forall t \geq 3, \quad \left\| |v^0|^{N_v-7} \langle x \rangle^{N_x-2} (f(t, X_\ell(t, \cdot, \cdot), \cdot) - f_\infty) \right\|_{L_{x,v}^\infty} \lesssim \bar{\epsilon} \frac{\log^{12+3N_x+3N}(t)}{t^\delta}.$$

Moreover, we could prove that f_∞ is of class C^{N-1} according to the spatial variable x .

Remark 6.36. We could prove that $\partial_v^\xi (\partial_t^n \partial_x^\kappa f(t, X_\ell, v)) \rightarrow \partial_v^\xi (-\hat{v} \cdot \nabla_x)^n \partial_x^\kappa f_\infty$. The idea consists in rewriting the time derivatives using $\partial_t = -\hat{v} \cdot \nabla_x + \mathbf{T}_F - \hat{v}^\mu F_\mu^j \partial_{v_j}$.

7. Scattering result for the electromagnetic field

In this section, we start by defining the scattering state of a sufficiently regular Maxwell field. Then, we construct a scattering map for the vacuum Maxwell equations. Finally, we apply these results together with the estimates derived in Section 3.1 in order to prove that the electromagnetic field F scatters, in the sense that it is approached by a solution to the homogeneous Maxwell equations.

Since the asymptotic states will be functions of the variables (u, θ, φ) , defined on future null infinity \mathcal{I}^+ introduced in Section 2.2, it will be convenient to work in null coordinates. For a function $\psi(t, x)$, in order to simplify the presentation, we will write $\psi(u, \underline{u}, \omega)$ to denote $\psi((\underline{u} + u)/2, (\underline{u} - u)\omega/2)$, where $(u, \underline{u}, \omega)$ are the null coordinates such that $x = r\omega$, $\underline{u} = t + r$ and $u = t - r$.

The scattering state of a smooth electromagnetic field G will give the leading-order term in the asymptotic expansion of rG , as $\underline{u} \rightarrow +\infty$. This motivates the introduction of the following terminology.

Definition 7.1. Let $\phi : \mathbb{R}_+ \times \mathbb{R}^3 \rightarrow \mathbb{R}$ be a function such that the limit

$$\Phi(u, \omega) := \lim_{r \rightarrow +\infty} r\phi(u + r, r\omega) = \lim_{\underline{u} \rightarrow +\infty} (r\phi)(u, \underline{u}, \omega), \quad \Phi(u, \omega) < +\infty,$$

exists and is finite for all $(u, \omega) \in \mathbb{R}_u \times \mathbb{S}^2$. Then, we say that the function Φ , defined on $\mathbb{R}_u \times \mathbb{S}^2$, is the radiation field $\mathcal{R}(\phi)$ of ϕ along future null infinity \mathcal{I}^+ .

Definition 7.2. Similarly, consider β , a 1-form on $\mathbb{R}_+ \times \mathbb{R}^3$ tangential to the 2-spheres¹² such that β_{e_θ} and β_{e_φ} have a radiation field $\beta_{e_\theta}^{\mathcal{I}^+}$ and $\beta_{e_\varphi}^{\mathcal{I}^+}$. Then, $\beta^{\mathcal{I}^+}$, defined on $\mathbb{R}_u \times \mathbb{S}^2$ as the 1-form $\beta_{e_\theta}^{\mathcal{I}^+} d\theta + \beta_{e_\varphi}^{\mathcal{I}^+} d\varphi$ tangential to the 2-spheres, is called the radiation field of β along \mathcal{I}^+ .

If $\beta^{\mathcal{I}^+}$ is of class C^1 , we define

$$\nabla_{\partial_u}(\beta) := \partial_u(\beta_{e_\theta}^{\mathcal{I}^+})d\theta + \partial_u(\beta_{e_\varphi}^{\mathcal{I}^+})d\varphi, \quad \nabla_{e_\theta}(\beta)(u, \cdot, \cdot) := \nabla_{e_\theta}(\beta(u, \cdot, \cdot)), \quad \nabla_{e_\varphi}(\beta)(u, \cdot, \cdot) := \nabla_{e_\varphi}(\beta(u, \cdot, \cdot)),$$

where ∇ denotes the covariant derivative on \mathbb{S}^2 .

We already know from Corollary 2.20 that, given a sufficiently decaying electromagnetic field G , the radiation field of the good null components $\alpha(G)$, $\rho(G)$ and $\sigma(G)$ exist and vanish. Concerning the component $\underline{\alpha}(G)$, we have the following result.

Proposition 7.3. Let G be a C^1 solution to the Maxwell equations (18) with a continuous source term J . Assume that there exist three constants $C[G] > 0$, $p \in \mathbb{N}$ and $q > 0$ such that, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$r|J|(t, x) + \sum_{|\gamma| \leq 1} |\rho(\mathcal{L}_{Z^\gamma} G)|(t, x) + |\sigma(\mathcal{L}_{Z^\gamma} G)|(t, x) \leq \frac{C[G] \log^p(3 + t + |x|)}{(1 + t + |x|)^{1+q}}. \tag{75}$$

Then, $\underline{\alpha}(G)$ has a radiation field along \mathcal{I}^+ . For any $B \in \{\theta, \varphi\}$ and for all $(u, \omega) \in \mathbb{R}_u \times \mathbb{S}^2$, the limit

$$\underline{\alpha}_{e_B}^{\mathcal{I}^+}(u, \omega) := \lim_{r \rightarrow +\infty} r\underline{\alpha}(G)_{e_B}(r + u, r\omega) = \lim_{\underline{u} \rightarrow +\infty} r\underline{\alpha}(G)_{e_B}(u, \underline{u}, \omega)$$

¹²More generally, we could consider tensor fields tangential to the cones \underline{C}_u .

exists and is finite. Moreover,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad \left| r\underline{\alpha}(G)_{e_B}(t, x) - \underline{\alpha}_{e_B}^{\mathcal{I}^+} \left(t - |x|, \frac{x}{|x|} \right) \right| \lesssim C[G] \frac{\log^p(3+t+|x|)}{(1+t+|x|)^q}.$$

Consequently, $\underline{\alpha}^{\mathcal{I}^+}$ is a continuous tensor field, defined on $\mathbb{R}_u \times \mathbb{S}^2$ and tangential to the 2-spheres.

Proof. The last inequality of [Lemma 2.17](#), together with [\(75\)](#), provides,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |\nabla_L(r\underline{\alpha}(G))|(t, x) \lesssim \log^p(3+t+|x|)(1+t+|x|)^{-1-q}. \tag{76}$$

Using the null coordinates $\underline{u} = t + r$ and $u = t - r$, where $x = r\omega$, we get, as $L = 2\partial_{\underline{u}}$ and $\nabla_L e_B = 0$,

$$\forall 0 \leq \underline{u} \leq \underline{z}, \quad |r\underline{\alpha}(F)(u, \underline{z}, \omega) - r\underline{\alpha}(F)(u, \underline{u}, \omega)| \lesssim \int_{s=\underline{u}}^{\underline{z}} \frac{\log^p(3+s) ds}{(1+s)^{1+q}} \lesssim \frac{\log^p(3+\underline{u})}{(1+\underline{u})^q},$$

implying the existence of $\underline{\alpha}_{e_B}^{\mathcal{I}^+}$, for any $B \in \{\theta, \varphi\}$, and the rate of convergence given in the statement. \square

If the electromagnetic field is sufficiently regular, we can relate the radiation fields of the derivatives of G to the ones of $\underline{\alpha}^{\mathcal{I}^+}$. For this, we will use the bounded functions $\omega_i := x^i/|x|$ and $\omega_i^A := \langle \partial_{x^i}, e_A \rangle$, where $1 \leq i \leq 3$ and $A \in \{\theta, \varphi\}$, which depend only on $\omega \in \mathbb{S}^2$ and which are given explicitly in [Appendix B](#).

Proposition 7.4. *Suppose that G satisfies, in addition to the hypotheses of the previous [Proposition 7.3](#), the inequality $|rG|(t, x) \leq C[G]$. Then, for any $Z \in \mathbb{K}$,*

$$\exists \underline{\alpha}_Z^{\mathcal{I}^+} \in \mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2), \quad r\underline{\alpha}(\mathcal{L}_Z G)(\cdot, \underline{u}, \cdot) \xrightarrow{u \rightarrow +\infty} \underline{\alpha}_Z^{\mathcal{I}^+} \quad \text{in } \mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2).$$

Moreover, for any $1 \leq i \leq 3$ and $1 \leq j < k \leq 3$,

$$\begin{aligned} \underline{\alpha}_{\partial_t}^{\mathcal{I}^+} &= \nabla_u \underline{\alpha}^{\mathcal{I}^+}, & \underline{\alpha}_{\partial_{x^i}}^{\mathcal{I}^+} &= -\omega_i \nabla_u \underline{\alpha}^{\mathcal{I}^+}, & \underline{\alpha}_S^{\mathcal{I}^+} &= u \nabla_u \underline{\alpha}^{\mathcal{I}^+} + \underline{\alpha}^{\mathcal{I}^+}, \\ \underline{\alpha}_{\Omega_{jk}}^{\mathcal{I}^+} &= \mathcal{L}_{\Omega_{jk}}(\underline{\alpha}^{\mathcal{I}^+}), & \underline{\alpha}_{\Omega_{0i}}^{\mathcal{I}^+} &= -\omega_i u \nabla_u \underline{\alpha}^{\mathcal{I}^+} - 2\omega_i \underline{\alpha}^{\mathcal{I}^+} + \omega_i^{e_A} \nabla_{e_A} \underline{\alpha}^{\mathcal{I}^+}. \end{aligned}$$

This result is proved in [Appendix B](#).

7.1. Scattering map for the vacuum Maxwell equations. Before starting the construction of the forward map for the homogeneous Maxwell equations, we introduce two functional spaces adapted to our problem. The first one contains the initial electromagnetic fields which are in L^2 and the second one contains the scattering states which belong to L^2 . For a smooth solution F to [\(19\)](#), this state will be the radiation field of $\underline{\alpha}(F)$. Note that the electromagnetic fields considered in this subsection will be denoted by F . Since, we will only consider solutions to the homogeneous Maxwell equations here, there is no risk of confusion with the electromagnetic field of the plasma considered in the remainder of the article.

Definition 7.5. Let $\mathcal{E}_{\{t=0\}}$ be the set containing all the 2-form on \mathbb{R}^{1+3} which does not depend on t and which is in $L^2(\mathbb{R}^3)$. Equipped with the norm

$$\|F_0\|_{\mathcal{E}_{\{t=0\}}}^2 := \int_{\mathbb{R}^3} (|\alpha(F_0)|^2 + |\underline{\alpha}(F_0)|^2 + 2|\rho(F_0)|^2 + 2|\sigma(F_0)|^2)(x) dx,$$

$\mathcal{E}_{\{t=0\}}$ is a Hilbert space.

We define $\mathcal{E}_{\mathcal{I}^+}$ as the set of the 1-forms on $\mathbb{R}_u \times \mathbb{S}^2$ which are tangential to the 2-spheres and in L^2 . For

$$\|\underline{\alpha}^{\mathcal{I}^+}\|_{\mathcal{I}^+}^2 := \int_{\mathbb{R}_u} \int_{\mathbb{S}_\omega^2} |\underline{\alpha}^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du,$$

$(\mathcal{E}_{\mathcal{I}^+}, \|\cdot\|_{\mathcal{I}^+})$ is a Hilbert space.

We now state the two main results of this section.

Theorem 7.6. *The linear map*

$$\mathcal{F}^+ : \mathcal{E}_{\{t=0\}} \cap C_c^\infty \rightarrow \mathcal{E}_{\mathcal{I}^+}, \quad F_0 \mapsto \lim_{u \rightarrow +\infty} r\underline{\alpha}(F)(u, \underline{u}, \omega),$$

where F is the unique solution to the vacuum Maxwell equations (19) such that $F(0, \cdot) = F_0$, is well-defined and preserves the norm $\|F_0\|_{\mathcal{E}_{\{t=0\}}} = \|\mathcal{F}^+(F_0)\|_{\mathcal{I}^+}$.

Moreover, this forward map can be uniquely extended in a bijective isometry $\mathcal{F}^+ : \mathcal{E}_{\{t=0\}} \rightarrow \mathcal{E}_{\mathcal{I}^+}$.

Remark 7.7. When $F_0 \notin C_c^\infty$ but is still sufficiently regular, $\mathcal{F}^+(F_0)$ is also given by the formula written in Theorem 7.6. Otherwise, $\mathcal{F}^+(F_0)$ can still be interpreted, in a weak sense, as the radiation field of $\underline{\alpha}(F)$, with F the solution to (19) arising from the data F_0 (see Lemma 7.9 below).

The proof will in particular rely on the following result, which is also important in itself. It provides precise estimates for solutions arising from the preimage by \mathcal{F}^+ of smooth elements of $\mathcal{E}_{\mathcal{I}^+}$.

Proposition 7.8. *Let $0 < a < \frac{1}{2}$, $N \in \mathbb{N}$ and $\underline{\alpha}^{\mathcal{I}^+} \in \mathcal{E}_{\mathcal{I}^+}$ be a sufficiently regular scattering state. Then, the unique solution F to the vacuum Maxwell equations (19) satisfying $\mathcal{F}^+(F) = \underline{\alpha}^{\mathcal{I}^+}$ satisfies, for any $0 \leq q - \frac{1}{2} < a$,*

$$\begin{aligned} \sum_{|\gamma| \leq N} \|(t-r)^{q-\frac{1}{2}} |\mathcal{L}_{Z^\gamma} F|(t, \cdot)\|_{L_x^2}^2 &\lesssim C[\underline{\alpha}^{\mathcal{I}^+}] \\ &:= \sum_{n_1+n_2+n_3 \leq N+3} \int_{\mathbb{R}_u} \int_{\mathbb{S}_\omega^2} \langle u \rangle^{2a+2n_1} |\nabla_u^{n_1} \not\psi_{e_\theta}^{n_2} \not\psi_{e_\varphi}^{n_3} \underline{\alpha}^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du \end{aligned}$$

for all $t \in \mathbb{R}_+$. In particular, if $N \geq 4$, we have, for any $|\gamma| \leq N - 3$ and $|\xi| \leq N - 4$,

$$\begin{aligned} \forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad (|\alpha(\mathcal{L}_{Z^\gamma} F)| + |\rho(\mathcal{L}_{Z^\gamma} F)| + |\sigma(\mathcal{L}_{Z^\gamma} F)|)(t, x) &\leq \frac{C}{(1+t+|x|)^{1+q}}, \\ \left| r\underline{\alpha}(\mathcal{L}_{Z^\xi} F)(t, x) - \mathcal{F}^+(\mathcal{L}_{Z^\xi} F(0, \cdot))\left(t - |x|, \frac{x}{|x|}\right) \right| &\leq \frac{C}{(1+t+|x|)^q}, \end{aligned}$$

where the constant C depends only on $C[\underline{\alpha}^{\mathcal{I}^+}]$ and q .

We start by proving that \mathcal{F}^+ is well-defined for sufficiently regular electromagnetic field, including those arising from smooth compactly supported data.

Lemma 7.9. *The linear map \mathcal{F}^+ introduced in Theorem 7.6 is well-defined and extends in an injective isometry from $\mathcal{E}_{\{t=0\}}$ to $\mathcal{E}_{\mathcal{I}^+}$. Moreover, if F is a solution to the free Maxwell equations (19) such that*

$$C_F := \sum_{|\gamma| \leq 4} \|\mathcal{L}_{Z^\gamma} F(0, \cdot)\|_{\{t=0\}} < +\infty, \tag{77}$$

then, $\underline{\alpha}(F)$ has a continuous radiation field $\mathcal{F}^+(F(0, \cdot))$ and, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$(|\alpha(F)| + |\rho(F)| + |\sigma(F)|)(t, x) \lesssim C_F(1 + t + |x|)^{-\frac{3}{2}}, \tag{78}$$

$$\left| r\underline{\alpha}(F)(t, x) - \mathcal{F}^+(F(0, \cdot))\left(t - |x|, \frac{x}{|x|}\right) \right| \lesssim C_F(1 + t + |x|)^{-\frac{1}{2}}. \tag{79}$$

This implies that the radiation fields of $\alpha(F)$, $\rho(F)$ and $\sigma(F)$ vanish.

Finally, if F is a mildly regular solution to (19) such that $F(0, \cdot) \in \mathcal{E}_{\{t=0\}}$, then $r\underline{\alpha}(F)$ converges to $\mathcal{F}^+(F(0, \cdot))$, as $\underline{u} \rightarrow +\infty$, in the space of distributions $\mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$.

Proof. Recall from Definition 2.16 the energy momentum tensor $\mathbb{T}[F]_{\mu\nu}$, its principal null components and that $\nabla^\mu \mathbb{T}[F]_{\mu 0} = 0$. For any $t > 0$, the divergence theorem, applied to $\mathbb{T}[F]_{\mu 0}$ in the domain $\{(s, x) \in \mathbb{R}^{1+3} \mid 0 \leq s \leq t\}$, gives

$$\begin{aligned} \|F(0, \cdot)\|_{\{t=0\}} &= 4 \int_{\mathbb{R}_x^3} \mathbb{T}[F]_{00}(0, x) \, dx = 4 \int_{\mathbb{R}_x^3} \mathbb{T}[F]_{00}(t, x) \, dx \\ &= 2 \sum_{0 \leq \mu, \nu \leq 3} \int_{\mathbb{R}_x^3} |F_{\mu\nu}|^2(t, x) \, dx = 2\|F(t, \cdot)\|_{L_x^2}. \end{aligned}$$

This also applies to $\mathcal{L}_{Z^\gamma}(F)$, for any $|\gamma| \leq 4$, since it is a solution to the free Maxwell equations (19) as well. In view of the equivalence of the pointwise norms (9), the standard Klainerman–Sobolev inequality (see for instance Theorem 1.3 of [Sogge 1995, Chapter II]) yields, for any $|\gamma| \leq 2$,

$$\begin{aligned} \forall(t, x) \in \mathbb{R}_+ \times \mathbb{R}_x^3, \quad |\mathcal{L}_{Z^\gamma} F|(t, x) &\lesssim \sum_{|\beta| \leq 2 + |\gamma|} \sum_{0 \leq \mu, \nu \leq 3} |Z^\beta(F_{\mu\nu})|(t, x) \\ &\lesssim \frac{C_F}{(1 + t + |x|)(1 + |t - |x||)^{\frac{1}{2}}}. \end{aligned} \tag{80}$$

Applying Corollary 2.20 to $\mathcal{L}_{Z^\xi} F$, for any $|\xi| \leq 1$ and $q = \frac{1}{2}$, gives,

$$\forall|\xi| \leq 1, \forall(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad (|\alpha(\mathcal{L}_{Z^\xi} F)| + |\rho(\mathcal{L}_{Z^\xi} F)| + |\sigma(\mathcal{L}_{Z^\xi} F)|)(t, x) \leq C_F(1 + t + |x|)^{-\frac{3}{2}}.$$

The existence of the radiation field $\underline{\alpha}^{\mathcal{I}^+}$ of $\underline{\alpha}(F)$ and the rate of convergence given in the statement then follows from Proposition 7.3. Since the convergence is uniform in (u, ω) , $\underline{\alpha}^{\mathcal{I}^+}$ is continuous on $\mathbb{R}_u \times \mathbb{S}^2$.

Before defining \mathcal{F}^+ , we need to bound the L^2 norm of the radiation field. For this, we prove conservation laws which hold for any mildly regular solution G to the free Maxwell equations (19).

Fix $\underline{u} \geq 0$ and apply the divergence theorem to $\mathbb{T}[G]_{\mu 0}$, in the domain $\{t + |x| \leq \underline{u}\}$, in order to get

$$\begin{aligned} \int_{\underline{C}_{\underline{u}}} \mathbb{T}[G]_{\underline{L}0} \, d\mu_{\underline{C}_{\underline{u}}} &= \int_{|x| \leq \underline{u}} \mathbb{T}[G]_{00}(0, x) \, dx \\ &= \frac{1}{4} \int_{|x| \leq \underline{u}} (|\alpha(G)|^2 + |\underline{\alpha}(G)|^2 + 2|\rho(G)|^2 + 2|\sigma(G)|^2)(0, x) \, dx, \end{aligned} \tag{81}$$

where

$$\int_{\underline{C}_{\underline{u}}} \mathbb{T}[G]_{\underline{L}0} \, d\mu_{\underline{C}_{\underline{u}}} = \frac{1}{4} \int_{|u| \leq \underline{u}} \int_{\mathbb{S}_\omega^2} (|\underline{\alpha}(G)|^2 + |\rho(G)|^2 + |\sigma(G)|^2)(u, \underline{u}, \omega) r^2 \, d\mu_{\mathbb{S}^2} \, du. \tag{82}$$

Assume now that $F_{\mu\nu}(0, \cdot) \in C_c^\infty(\mathbb{R}_x^3)$ for all $0 \leq \mu, \nu \leq 3$ and let us apply the previous equality to F . On the one hand, the right-hand side of (81) converges to $\frac{1}{4} \|F(0, \cdot)\|_{\{t=0\}}^2$ as $\underline{u} \rightarrow +\infty$. On the other hand, we know from the Huygens–Fresnel principle that there exists $U > 0$ such that $F(t, x) = 0$ for all $|t - |x|| = |\underline{u}| \geq U$. This implies that the domain of integration of the integrals in (82) is in fact included in $\{|\underline{u}| \leq U\}$ for all $\underline{u} \geq 0$. The triangular inequality in L^2 together with the estimates (78)–(79) then leads to

$$\int_{\underline{C}_{\underline{u}}} \mathbb{T}[F]_{\underline{L}0} \, d\mu_{\underline{C}_{\underline{u}}} \xrightarrow{\underline{u} \rightarrow +\infty} \frac{1}{4} \int_{|\underline{u}| \leq U} \int_{\mathbb{S}_\omega^2} |\mathcal{F}^+(F(0, \cdot))|^2 \, d\mu_{\mathbb{S}^2} \, d\underline{u} = \frac{1}{4} \|\underline{\alpha}^{\mathcal{I}^+}\|_{\mathcal{I}^+}^2.$$

We can then define $\mathcal{F}^+ : \mathcal{E}_{\{t=0\}} \cap C_c^\infty \rightarrow \mathcal{E}_{\mathcal{I}^+}$, with $\mathcal{F}^+(F(0, \cdot)) := \underline{\alpha}^{\mathcal{I}^+}$, and extend it to an injective isometry from $\mathcal{E}_{\{t=0\}}$ to $\mathcal{E}_{\mathcal{I}^+}$.

Consider now a, say, C^1 solution F to (19) such that $F(0, \cdot) \in \mathcal{E}_{\{t=0\}}$. Fix $\psi \in C_c^\infty(\mathbb{R}_u \times \mathbb{S}^2)$ and $R > 0$ satisfying $\text{supp}(\psi) \subset [-R, R] \times \mathbb{S}^2$. Let further $(F_n)_{n \geq 0}$ be a sequence of smooth solutions to the vacuum Maxwell equations such that $F_n(0, \cdot)$ is compactly supported for any $n \in \mathbb{N}$ and $F_n(0, \cdot) \rightarrow F(0, \cdot)$ in $\mathcal{E}_{\{t=0\}}$. Fix $A \in \{\theta, \varphi\}$ and start by observing that

$$\begin{aligned} & |(r\underline{\alpha}(F)_{e_A} - \mathcal{F}^+(F(0, \cdot))_{e_A})\psi| \\ & \lesssim (|r\underline{\alpha}(F) - r\underline{\alpha}(F_n)| + |r\underline{\alpha}(F_n) - \mathcal{F}^+(F_n(0, \cdot))| + |\mathcal{F}^+((F_n - F)(0, \cdot))|) \mathbb{1}_{|\underline{u}| \leq R}. \end{aligned}$$

Then, in order to prove $r\underline{\alpha}_{e_A} \rightarrow \mathcal{F}^+(F(0, \cdot))_{e_A}$ in $\mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$, as $\underline{u} \rightarrow +\infty$, it suffices to prove that the integral on $\mathbb{R}_u \times \mathbb{S}^2$ of each of the three terms on the right-hand side converges to 0 as $\underline{u} \rightarrow +\infty$. For this, consider $\epsilon > 0$ and start by noticing that the energy equality (81)–(82), applied to $F - F_n$, gives,

$$\forall n \geq 0, \forall \underline{u} \geq 0, \quad \int_{\mathbb{R}_u} \int_{\mathbb{S}_\omega^2} |r\underline{\alpha}(F) - r\underline{\alpha}(F_n)|^2(u, \underline{u}, \omega) \, d\mu_{\mathbb{S}^2} \, d\underline{u} \leq \|F(0, \cdot) - F_n(0, \cdot)\|_{\{t=0\}}^2.$$

According to (79), applied to F_n , there exists a constant C_n , such that,

$$\forall n \in \mathbb{N}, \forall \underline{u} \geq 0, \quad \int_{|\underline{u}| \leq R} \int_{\mathbb{S}_\omega^2} |r\underline{\alpha}(F_n)(u, \underline{u}, \omega) - \mathcal{F}^+(F_n(0, \cdot))(u, \omega)| \, d\mu_{\mathbb{S}^2} \, d\underline{u} \leq \frac{C_n}{(1 + \underline{u})^{\frac{1}{2}}}.$$

Moreover, since \mathcal{F}^+ is an isometry, we have $\|\mathcal{F}^+(F_n(0, \cdot)) - \mathcal{F}^+(F(0, \cdot))\|_{\mathcal{I}^+} = \|F(0, \cdot) - F_n(0, \cdot)\|_{\{t=0\}}$. The last four estimates, together with the Cauchy–Schwarz inequality in $L^2([-R, R] \times \mathbb{S}^2)$, yields

$$\left| \int_{\mathbb{R}_u} \int_{\mathbb{S}_\omega^2} (r\underline{\alpha}(F)_{e_A}(u, \underline{u}, \omega) - \mathcal{F}^+(F)_{e_A}(u, \omega))\psi(u, \omega) \, d\mu_{\mathbb{S}^2} \, d\underline{u} \right| \lesssim \|F(0, \cdot) - F_n(0, \cdot)\|_{\{t=0\}} + \frac{C_n}{(1 + \underline{u})^{\frac{1}{2}}}$$

for all $n \in \mathbb{N}$ and $\underline{u} \geq 0$. For a sufficiently large n and \underline{U} , which depends on n , we can bound the right-hand side by ϵ for all $\underline{u} \geq \underline{U}$. This concludes the proof of the last part of the lemma.

It remains to show that for any $F(0, \cdot)$ satisfying (77), we have $\mathcal{F}^+(F(0, \cdot)) = \underline{\alpha}^{\mathcal{I}^+}$. For this, it suffices to recall that we proved $r\underline{\alpha}(F) \rightarrow \underline{\alpha}^{\mathcal{I}^+}$ in $L_{u, \omega}^\infty$. □

Remark 7.10. In fact, assuming more decay on the initial data, we could prove using the equations (M_2'') , (M_5'') and (M_6'') of [Christodoulou and Klainerman 1990] that $|\alpha(F)| = O(\underline{u}^{-2-\delta})$ and that $r^2\rho(F)$ as well as $r^2\sigma(F)$ converge as $\underline{u} \rightarrow +\infty$.

To conclude the proof of [Theorem 7.6](#), it remains us to show that \mathcal{F}^+ is surjective. For this, it suffices to prove [Proposition 7.8](#), which in particular implies that any smooth and compactly supported $\alpha^{\mathcal{I}^+}$ has a preimage by \mathcal{F}^+ . For this, we will make crucial use of [[Lindblad and Schlue 2023](#), Theorem 1.1], which is a similar result for solutions to the homogeneous wave equation, and exploit that $\square F_{\mu\nu} = 0$ for any Cartesian component $F_{\mu\nu}$.

Lemma 7.11. *Let $\Phi \in C(\mathbb{R}_u \times \mathbb{S}^2)$ be a sufficiently regular function, $0 < a < \frac{1}{2}$ and $N \in \mathbb{N}$. Then, there exists a unique solution to wave equation $\square\phi = 0$ on $\mathbb{R}_+ \times \mathbb{R}^3$ satisfying, for any $0 < \delta \leq a$ and all $t \in \mathbb{R}_+$,*

$$\sum_{|\gamma| \leq N} \|\langle t-r \rangle^{a-\delta} Z^\gamma \phi(t, \cdot)\|_{L^2(\mathbb{R}_x^3)}^2 \lesssim \sum_{|k|+|\beta| \leq N+3} \int_{u=-\infty}^{+\infty} \int_{\omega \in \mathbb{S}^2} |(\langle u \rangle \partial_u)^k \partial_\omega^\beta \Phi(u, \omega)|^2 \langle u \rangle^{2a} d\mu_{\mathbb{S}^2} du$$

and such that Φ is the radiation field $\mathcal{R}(\phi)$ of ϕ along \mathcal{I}^+ .

We will also require standard estimates for smooth solutions to the wave equation.

Lemma 7.12. *Let ϕ be a smooth solution to the wave equation $\square\phi = 0$ such that $\|Z^\gamma \phi(0, \cdot)\|_{L_x^2} < +\infty$ for any $|\gamma| \leq 5$. Then, for any $|\beta| \leq 1$, the radiation field $\mathcal{R}(\partial_{t,x}^\beta \phi)$ of $\partial_{t,x}^\beta \phi$ is well-defined and*

$$\forall \underline{u} \geq 1, \forall (u, \omega) \in [-\underline{u}, \underline{u}] \times \mathbb{S}^2, \quad |r \partial_{t,x}^\beta \phi(u, \underline{u}, \omega) - \mathcal{R}(\partial_{t,x}^\beta \phi)(u, \omega)| \lesssim \underline{u}^{-\frac{1}{2}}.$$

Moreover, $\mathcal{R}(\partial_t \phi) = \partial_u \mathcal{R}(\phi)$ and $\mathcal{R}(\partial_{x^i} \phi) = -(x^i/|x|) \partial_u \mathcal{R}(\phi)$ for all $i \in \llbracket 1, 3 \rrbracket$.

Proof. The first part of the result is classical. Indeed, since $\square Z^\gamma \phi = 0$ for any $|\gamma| \leq 4$, we obtain by applying the standard Klainerman–Sobolev inequality and then an energy inequality (for a proof, see for instance Theorem 1.3 and Lemma 3.5 of [[Sogge 1995](#), Chapter II]), that, for all $|\gamma| \leq 2$ and $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$(1+t+|x|)(1+|t-|x||)^{\frac{1}{2}} |Z^\gamma \phi|(t, x) \lesssim \sum_{|\beta| \leq |\gamma|+2} \|Z^\beta \phi(t, \cdot)\|_{L_x^2} \lesssim \sum_{|\beta| \leq 4} \|Z^\beta \phi(0, \cdot)\|_{L_x^2}. \quad (83)$$

Now we claim that,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad |L(r\phi)|(t, x) \lesssim (1+t+|x|)^{-\frac{3}{2}}.$$

Indeed, if $|x| = r \leq (1+t)/2$, we have $1+t+r \lesssim 1+|t-r|$. Moreover, (20) leads to $|L(r\phi)| \leq \sum_{|\beta| \leq 1} |Z^\beta \phi|$, so that the claim is implied by (83). Otherwise, $|x| \gtrsim 1+t+|x| = 1+\underline{u}$ and, by writing the d’Alembertian in spherical coordinates, we obtain from $\square\phi = 0$ that

$$0 = -L\underline{L}\phi + \frac{2L-\underline{L}}{r} \phi + \sum_{1 \leq i < j \leq 3} \frac{\Omega_{ij} \Omega_{ij} \phi}{r^2}, \quad \text{leading to} \quad \underline{L}(L(r\phi)) = \sum_{1 \leq i < j \leq 3} \frac{\Omega_{ij} \Omega_{ij} \phi}{r}. \quad (84)$$

In order to integrate along a null straight line $t+r = \underline{u}$, it will be convenient to work with the null coordinate system. We then write $x = |x|\omega$, with $\omega \in \mathbb{S}^2$. As $\underline{L} = 2\partial_u$ and in view of (83)–(84), we have

$$\begin{aligned} |L(r\phi)|(t, x) &= |L(r\phi)|(t-|x|, \underline{u}, \omega) \leq |L(r\phi)|(-t-|x|, \underline{u}, \omega) + \frac{1}{2} \int_{u=-t-|x|}^{t-|x|} |\underline{L}(L(r\phi))|(u, \underline{u}, \omega) du \\ &\lesssim |L(r\phi)|(0, (t+|x|)\omega) + \int_{u=-t-|x|}^{t-|x|} \frac{du}{(1+\underline{u})^2(1+|u|)^{\frac{1}{2}}} \lesssim (1+\underline{u})^{-\frac{3}{2}}, \end{aligned}$$

which concludes the proof of the claim. As $L = 2\partial_{\underline{u}}$, we directly deduce from it that,

$$\forall \underline{z} \geq \underline{u} \geq 0, \forall |u| \leq \underline{u}, \forall \omega \in \mathbb{S}^2, \quad |r\phi(u, \underline{z}, \omega) - r\phi(u, \underline{u}, \omega)| \lesssim \int_{s=\underline{u}}^{\underline{z}} |L(r\phi)|(u, s, \omega) ds \lesssim (1 + \underline{u})^{-\frac{1}{2}}.$$

This implies the existence of the radiation field $\mathcal{R}(\phi)$ of ϕ as well as the rate of convergence given in the statement of the lemma. Since $\square \partial_{x^\mu} \phi = 0$ and $\|Z^\gamma \partial_{x^\mu} \phi(0, \cdot)\|_{L_x^2} < +\infty$ for any $|\gamma| \leq 4$, the same applies to $\partial_{x^\mu} \phi$. Now, note that

$$2r\partial_t \phi = rL\phi + r\underline{L}\phi, \quad 2r\partial_{x^i} \phi = \frac{x^i}{|x|}rL\phi - \frac{x^i}{|x|}r\underline{L}\phi + 2\langle \partial_{x^i}, e_\theta \rangle r e_\theta \phi + 2\langle \partial_{x^i}, e_\varphi \rangle r e_\varphi \phi, \quad 1 \leq i \leq 3.$$

Combining (83) with (20) yields $r|L\phi| + r|e_\theta \phi| + r|e_\varphi| + |\phi| \lesssim \underline{u}^{-1}$ so that

$$\text{there exists } \phi_\infty^{\underline{L}} \in L^\infty(\mathbb{R}_u \times \mathbb{S}_\omega^2) \text{ such that } \underline{L}(r\phi) \xrightarrow[\underline{u} \rightarrow +\infty]{L_{u,\omega}^\infty} \phi_\infty^{\underline{L}}, \quad \phi_\infty^{\underline{L}} = 2\mathcal{R}(\partial_t \phi), \quad \frac{x^i}{|x|} \phi_\infty^{\underline{L}} = -2\mathcal{R}(\partial_{x^i} \phi).$$

It remains to use that $\underline{L}(r\phi)(\cdot, \underline{u}, \cdot) \rightarrow 2\partial_{\underline{u}}\mathcal{R}(\phi)$ in $\mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$ since $r\phi(\cdot, \underline{u}, \cdot)$ converges to $\mathcal{R}(\phi)$ in $L_{u,\omega}^\infty$. □

We are now ready for the last part of this subsection.

Proof of Proposition 7.8. Fix $0 \leq q - \frac{1}{2} < a < \frac{1}{2}$, $N \in \mathbb{N}$ and $\underline{\alpha}^{\mathcal{I}^+} \in \mathcal{E}_{\mathcal{I}^+}$ such that the norm $C[\underline{\alpha}^{\mathcal{I}^+}]$ is finite. Recall that any sufficiently regular solution F to the vacuum Maxwell equations (19) satisfies $\square F_{\mu\nu} = 0$ for any $0 \leq \mu, \nu \leq 3$. The first step consists in constructing each Cartesian component $F_{\mu\nu}$ of the electromagnetic field by applying Lemma 7.11 to well-chosen radiation fields. This will define a 2-form F which will verify the stated estimate. Then, we will prove that F is indeed a solution to the Maxwell equations and, finally, we will derive the pointwise decay estimates.

Assume first that $N \geq 5$ and let us start by identifying the expected radiation field of $F_{\mu\nu}$. For this, assume that F exists and recall the transfer matrix between the Cartesian and the null frame

$$\partial_t = \frac{1}{2}L + \frac{1}{2}\underline{L}, \quad \partial_{x^i} = \frac{1}{2}\omega_i L - \frac{1}{2}\omega_i \underline{L} + \omega_i^{e_\theta} e_\theta + \omega_i^{e_\varphi} e_\varphi, \quad 1 \leq i \leq 3,$$

where ω_i and $\omega_i^{e_A}$ are bounded functions of the spherical variables and are given explicitly in Appendix B. For convenience, we set $\omega_0 := -1$ and $\omega_0^{e_A} := 0$. Consequently, for any $0 \leq \mu, \nu \leq 3$, there exist smooth functions of $\omega \in \mathbb{S}^2$, $g_{\mu\nu}^{\alpha,\theta}$, $g_{\mu\nu}^{\alpha,\varphi}$, $g_{\mu\nu}^\rho$ and $g_{\mu\nu}^\sigma$, such that

$$rF_{\mu\nu} = -\frac{1}{2}(\omega_\mu^{e_A} \omega_\nu - \omega_\nu \omega_\mu^{e_A})r\underline{\alpha}(F)_{e_A} + g_{\mu\nu}^{\alpha,A}r\alpha(F)_{e_A} + g_{\mu\nu}^\rho r\rho(F) + g_{\mu\nu}^\sigma r\sigma(F).$$

We then obtain by (78)–(79) that

$$\mathcal{R}(F_{\mu\nu}) = -\frac{1}{2}(\omega_\mu^{e_A} \omega_\nu - \omega_\nu \omega_\mu^{e_A})\underline{\alpha}_{e_A}^{\mathcal{I}^+}, \quad 0 \leq \mu, \nu \leq 3. \tag{85}$$

According to Lemma 7.11, we can indeed define a 2-form F satisfying (85) as well as $\square F_{\mu\nu} = 0$ and, for all $t \in \mathbb{R}_+$,

$$\sum_{|\gamma| \leq N} \|(t-r)^{q-\frac{1}{2}}|\mathcal{L}_{Z^\gamma} F|(t, \cdot)\|_{L_x^2} \lesssim \sum_{|\gamma| \leq N} \sum_{0 \leq \mu, \nu \leq 3} \|(t-r)^{q-\frac{1}{2}}Z^\gamma(F_{\mu\nu})(t, \cdot)\|_{L_x^2} \lesssim C[\underline{\alpha}^{\mathcal{I}^+}]. \tag{86}$$

The remainder of the proof of the case $N \geq 5$ essentially consists in performing linear algebra computations. In order to lighten the notations we temporarily denote ∂_{x^λ} by ∂_λ . Our goal now is to prove that F is a solution to the vacuum Maxwell equations (19), which read in Cartesian coordinates

$$\partial^\mu F_{\mu\nu} = 0, \quad \partial^{\mu*} F_{\mu\nu} = \partial_{[\lambda} F_{\mu\nu]} := \partial_\lambda F_{\mu\nu} + \partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} = 0. \tag{87}$$

For a proof of the second identity, see for instance [Bigorgne 2021b, Lemma 2.2]. Since $\square \partial^\mu F_{\mu\nu} = 0$ and $\square \partial^{\mu*} F_{\mu\nu} = 0$, (87) would be implied, according to Lemma 7.11, by

$$\mathcal{R}(\partial^\mu F_{\mu\nu}) = 0, \quad \mathcal{R}(\partial^{\mu*} F_{\mu\nu}) = 0, \quad 0 \leq \nu \leq 3.$$

We compute, using Lemma 7.12, that, for any $0 \leq \lambda \leq 3$,

$$\mathcal{R}(\partial_\lambda F_{\mu\nu}) = -\omega_\lambda \partial_\mu \mathcal{R}(F_{\mu\nu}) = \frac{1}{2} \omega_\lambda (\omega_\mu^{e_A} \omega_\nu - \omega_\mu \omega_\nu^{e_A}) \partial_\mu \underline{\alpha}_{e_A}^{T^+}, \quad 0 \leq \mu, \nu \leq 3.$$

This implies in particular that $\mathcal{R}(\partial_{[\lambda} F_{\mu\nu]}) = 0$. Furthermore, as $\partial^\mu = \eta^{\mu\lambda} \partial_\lambda$, we have

$$\mathcal{R}(\partial^\mu F_{\mu\nu}) = \frac{1}{2} \eta^{\mu\lambda} \omega_\lambda (\omega_\mu^{e_A} \omega_\nu - \omega_\mu \omega_\nu^{e_A}) \partial_\mu \underline{\alpha}_{e_A}^{T^+} = \frac{1}{2} (\eta(e_A, L) \omega_\nu - \eta(L, L) \omega_\nu^{e_A}) \partial_\mu \underline{\alpha}_{e_A}^{T^+} = 0.$$

We then deduce that F is a smooth solution to the vacuum Maxwell equations. Finally, since the Cartesian components of $\underline{L} = \eta_{\underline{L}}^\mu \partial_\mu$ and $e_A = \eta_{e_A}^\mu \partial_\mu$ are bounded functions of $\omega \in \mathbb{S}^2$, we obtain from (85) and Lemmas 7.9, 7.12 that

$$\begin{aligned} \mathcal{F}^+(F(0, \cdot))_{e_A} &= \lim_{\underline{u} \rightarrow \infty} r \underline{\alpha}(F)_{e_A}(\cdot, \underline{u}, \cdot) \\ &= \eta_{e_A}^\mu \eta_{\underline{L}}^\nu \lim_{\underline{u} \rightarrow \infty} r F_{\mu\nu}(\cdot, \underline{u}, \cdot) = \eta_{e_A}^\mu \eta_{\underline{L}}^\nu \mathcal{R}(F_{\mu\nu}) = \underline{\alpha}_{e_A}^{T^+}, \quad A \in \{\theta, \varphi\}. \end{aligned}$$

This concludes the proof of the first part of the proposition for the case $N \geq 5$. Consider now the case $N = 0$ and define similarly $F_{\mu\nu}$, through Lemma 7.11, as the unique solution to $\square F_{\mu\nu} = 0$ such that (85) holds. This directly provides the estimate (86); let us prove that F is a weak solution to (19). For this, consider a sequence $(\underline{\alpha}_n^{I^+}) \in \mathcal{E}_{T^+}^{\mathbb{N}}$ of smooth and compactly supported scattering states such that $C[\underline{\alpha}^{I^+} - \underline{\alpha}_n^{I^+}] \rightarrow 0$ as $n \rightarrow +\infty$. Then, denote by F_n the unique smooth solution to the vacuum Maxwell equations such that $\mathcal{F}^+(F_n(0, \cdot)) = \underline{\alpha}_n^{I^+}$. Applying once again Lemma 7.11 to $\mathcal{R}(F_{\mu\nu} - F_{n,\mu\nu})$ yields

$$\sup_{t \in \mathbb{R}_+} \|F(t, \cdot) - F_n(t, \cdot)\|_{L_x^2} \lesssim C[\underline{\alpha}^{I^+} - \underline{\alpha}_n^{I^+}]. \tag{88}$$

Fix $\psi \in C_c^\infty(\mathbb{R}_+ \times \mathbb{R}_x^3)$ and T_ψ such that $\psi(t, \cdot) = 0$ for all $t \geq T_\psi$. Note, since F_n is a classical and then a weak solution to (19), that for any $0 \leq \nu \leq 3$ and $n \in \mathbb{N}$,

$$\begin{aligned} &\left| \int_{\mathbb{R}_+ \times \mathbb{R}_x^3} F_{\mu\nu}(t, x) \partial^\mu \psi(t, x) \, dx \, dt + \int_{\mathbb{R}_x^3} F_{\mu\nu}(0, x) \psi(0, x) \, dx \right| \\ &= \left| \int_{\mathbb{R}_+ \times \mathbb{R}_x^3} (F - F_n)_{\mu\nu}(t, x) \partial^\mu \psi(t, x) \, dx \, dt + \int_{\mathbb{R}_x^3} (F - F_n)_{\mu\nu}(0, x) \psi(0, x) \, dx \right| \\ &\lesssim (1 + T_\psi) \sup_{t \in \mathbb{R}_+} \|(F - F_n)(t, \cdot)\|_{L_x^2}. \end{aligned} \tag{89}$$

By (88), the right-hand side converges to 0 as $n \rightarrow +\infty$ whereas the left-hand side does not depend on n . This implies that (89) vanishes. The same applies to *F , so that F is a weak solution to the vacuum Maxwell equations (19). Finally, by continuity of \mathcal{F}^+ and (88), $\mathcal{F}^+(F(0, \cdot)) = \underline{\alpha}^{\mathcal{I}^+}$.

We now focus on the second part of Proposition 7.8, which merely concerns the cases $N \geq 4$. We apply [Lindblad and Schlue 2023, Lemma 3.3], a weighted version of the standard Klainerman–Sobolev inequality, to $Z^\beta(F_{\mu\nu})$. Using (9), we obtain, for any $|\gamma| \leq N - 2$ and all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$|\mathcal{L}_{Z^\gamma}(F)|(t, x) \lesssim \sum_{|\beta| \leq N-2} \sum_{0 \leq \mu, \nu \leq 3} |Z^\beta(F_{\mu\nu})|(t, x) \lesssim \sum_{|\beta| \leq N} \frac{\|(t-r)^{q-\frac{1}{2}}|\mathcal{L}_{Z^\beta}(F)|(t, \cdot)\|_{L^2_x}}{(1+t+|x|)(1+|t-|x||)^q}. \tag{90}$$

The numerator in the right-hand side is bounded by $C[\underline{\alpha}^{\mathcal{I}^+}]$. Recall now that $\mathcal{L}_{Z^\gamma}(F)$ is a solution to the vacuum Maxwell equations as well. To conclude the proof, it then suffices to use the previous estimate and to apply Corollary 2.20 to $\mathcal{L}_{Z^\gamma}(F)$ for any $|\gamma| \leq N - 3$, as well as Proposition 7.3, to $\mathcal{L}_{Z^\xi}(F)$ for any $|\xi| \leq N - 4$. □

Remark 7.13. A statement similar to Theorem 7.6 holds for scattering toward past null infinity $\mathcal{I}^- \cong \mathbb{R}_u \times \mathbb{S}^2$. One can construct the past forward evolution bijective isometry $\mathcal{F}^- : \mathcal{E}_{\{t=0\}} \rightarrow \mathcal{E}_{\mathcal{I}^-}$, where, if $F(0, \cdot) \in \mathcal{E}_{\{t=0\}} \cap C_c^\infty$, $\mathcal{F}^-(F)(\underline{u}, \omega) := \lim_{u \rightarrow -\infty} r\alpha(F)(u, \underline{u}, \omega)$ and $\|\cdot\|_{\mathcal{I}^-} := \|\cdot\|_{L^2(\mathbb{R}_u \times \mathbb{S}^2)}$. The scattering map $\mathcal{S} = (\mathcal{F}^-)^{-1} \circ \mathcal{F}^+$ then defines a unitary isomorphism of Hilbert spaces.

Finally, we state a direct consequence of Theorem 7.6, Proposition 7.8 and the commutation properties of the vacuum Maxwell equations with \mathcal{L}_Z , $Z \in \mathbb{K}$.

Definition 7.14. Let $N \geq 0$ and $\mathcal{E}_{\{t=0\}}^N \subset \mathcal{E}_{\{t=0\}}$ be the set of the 2-forms on \mathbb{R}^{1+3} independent of t verifying

$$\|F_0\|_{\mathcal{E}_{\{t=0\}}^N}^2 := \sum_{|\gamma| \leq N} \|\mathcal{L}_{Z^\gamma}(F_0)(0, \cdot)\|_{\mathcal{E}_{\{t=0\}}}^2 < +\infty.$$

Consider $\mathcal{E}_{\mathcal{I}^+}^N \subset \mathcal{E}_{\mathcal{I}^+}$, the set of the 1-forms on $\mathbb{R}_u \times \mathbb{S}^2$ which are tangential to the 2-spheres and such that

$$\|\underline{\alpha}^{\mathcal{I}^+}\|_{N, \mathcal{I}^+}^2 := \sum_{|\gamma| \leq N} \|\underline{\alpha}_{Z^\gamma}^{\mathcal{I}^+}\|_{\mathcal{I}^+}^2 < +\infty,$$

where $\underline{\alpha}_{Z^\gamma}^{\mathcal{I}^+}$ is defined recursively from $\underline{\alpha}^{\mathcal{I}^+}$ through Proposition 7.4. Then, $(\mathcal{E}_{\{t=0\}}^N, \|\cdot\|_{\mathcal{E}_{\{t=0\}}^N}^2)$ and $(\mathcal{E}_{\mathcal{I}^+}^N, \|\cdot\|_{N, \mathcal{I}^+}^2)$ are Hilbert spaces.

Corollary 7.15. For any $N \geq 0$, the restriction of \mathcal{F}^+ to $\mathcal{E}_{\{t=0\}}^N$ is a bijective isometry from $\mathcal{E}_{\{t=0\}}^N$ to $\mathcal{E}_{\mathcal{I}^+}^N$.

7.2. Existence of an asymptotic state for F and its derivatives. In order to avoid any confusion, we make precise that, as in Sections 3–6, F denotes the electromagnetic field of our solution to the Vlasov–Maxwell system (f, F) . The following statement can be easily deduced from previous results.

Proposition 7.16. For any $|\gamma| \leq N - 3$, $\underline{\alpha}(\mathcal{L}_{Z^\gamma}F)$ has a continuous radiation field $\underline{\alpha}_\gamma^{\mathcal{I}^+}$. Moreover, for any $0 \leq \eta < 1$, we have the rate of convergence,

$$\forall \underline{u} \in \mathbb{R}_+, |u| \leq \underline{u}, \omega \in \mathbb{S}^2, \quad \left| \langle u \rangle^\eta (r \underline{\alpha}(\mathcal{L}_{Z^\gamma}F)(\underline{u}, u, \omega) - \underline{\alpha}_\gamma^{\mathcal{I}^+}(u, \omega)) \right| \lesssim \Lambda \frac{\log(3 + \underline{u})}{(1 + \underline{u})^{1-\eta}}.$$

If $|\gamma| = 0$, we simply denote the radiation field of F by $\underline{\alpha}^{\mathcal{I}^+}$.

Proof. Recall from [Proposition 2.4](#) the form of the source term in the commuted Maxwell equations. Hence, according to the estimates of [Proposition 3.1](#) and [Corollary 4.14](#), $\mathcal{L}_{Z^\gamma} F$ satisfies the hypotheses of [Proposition 7.3](#). \square

It turns out that our decomposition of F allows us to improve the estimate on the radiation field.

Proposition 7.17. *For any $|\gamma| \leq N - 3$, we have,*

$$\forall (u, \omega) \in \mathbb{R} \times \mathbb{S}^2, \quad |\underline{\alpha}_\gamma^{\mathcal{I}^+}|(u, \omega) \lesssim \begin{cases} \Lambda \langle u \rangle^{-1-\delta} & \text{if } 0 < \delta < 1, \\ \Lambda \langle u \rangle^{-2} \log(1 + \langle u \rangle) & \text{if } \delta = 1. \end{cases}$$

Proof. Recall the decomposition

$$r \mathcal{L}_{Z^\gamma} F = r \mathcal{L}_{Z^\gamma} (F)^S + r(\mathcal{L}_{Z^\gamma} (F)^{\text{data}}(t, x) - \mathcal{L}_{Z^\gamma}(\tilde{F})) + r \mathcal{L}_{Z^\gamma}(\tilde{F}) + r \mathcal{L}_{Z^\gamma} (F)^T.$$

Then, we use that $u = t - r$ as well as:

- The first term is bounded by $\Lambda \langle t - r \rangle^{-2} \log(1 + \langle t - r \rangle)$ according to [Proposition 5.14](#).
- By [Proposition 6.12](#), the second one is controlled by $\Lambda \langle t - r \rangle^{-1-\delta}$.
- By [Remark 6.13](#), the third term is bounded by $\epsilon(1 + t + r)^{-1} + \epsilon \mathbb{1}_{|t-r| \leq 1}$ and $u = t - r$.
- Finally, the last one is bounded above by $\bar{\epsilon}(1 + t + r)^{-3/4}$ according to [Proposition 5.15](#). \square

The last goal of this section consists in proving, if N is large enough, that F can be approached by a solution to the vacuum Maxwell equations through an application of [Proposition 7.8](#), which requires us to control $\underline{\alpha}^{\mathcal{I}^+}$ and its derivatives up to order at least 3. Note then that by iterating [Proposition 7.4](#), we get that $\underline{\alpha}_\gamma^{\mathcal{I}^+}$ can be computed in terms of derivatives of $\underline{\alpha}^{\mathcal{I}^+}$. Conversely, for any $0 \leq a < \frac{1}{2}$, we have

$$\sum_{n_u+n_\theta+n_\varphi \leq N-3} \int_{\mathbb{R}_u} \int_{\mathbb{S}^2} \langle u \rangle^{2a+2n_u} |\nabla_u^{n_u} \nabla_{e_\theta}^{n_\theta} \nabla_{e_\varphi}^{n_\varphi} \underline{\alpha}^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du \lesssim \sum_{|\gamma| \leq N-3} \int_{\mathbb{R}_u} \int_{\mathbb{S}^2} \langle u \rangle^{2a} |\underline{\alpha}_\gamma^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du.$$

Applying [Proposition 7.16](#) for $\eta = (3 + 2a)/4$ then yields

$$\sum_{n_u+n_\theta+n_\varphi \leq N-3} \int_{\mathbb{R}_u} \int_{\mathbb{S}^2} \langle u \rangle^{2a+2n_u} |\nabla_u^{n_u} \nabla_{e_\theta}^{n_\theta} \nabla_{e_\varphi}^{n_\varphi} \underline{\alpha}^{\mathcal{I}^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du \lesssim \Lambda \int_{\mathbb{R}_u} \langle u \rangle^{a-\frac{3}{2}} \, du \lesssim \frac{\Lambda}{1-2a}. \quad (91)$$

We are now ready to prove the following result.

Proposition 7.18. *If $N \geq 10$, there exists a solution F^{vac} of class C^{N-8} to the vacuum Maxwell equations (19) such that, for any $\frac{1}{2} \leq q < 1$ and $|\gamma| \leq N - 10$,*

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad r|\mathcal{L}_{Z^\gamma}(F) - \mathcal{L}_{Z^\gamma}(F^{\text{vac}})|(t, x) \leq \Lambda C_q (1 + t + |x|)^{-q},$$

where the constant $C_q > 0$ depends on q .

Proof. We fix $0 \leq q - \frac{1}{2} < a < \frac{1}{2}$. Since (91) holds, we get from [Proposition 7.8](#) that there exists a solution F^{vac} of class C^{N-8} to the vacuum Maxwell equations satisfying, for any $|\gamma| \leq N - 9$ and $|\xi| \leq N - 10$,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad (|\alpha(\mathcal{L}_{Z^\gamma} F^{\text{vac}})| + |\rho(\mathcal{L}_{Z^\gamma} F^{\text{vac}})| + |\sigma(\mathcal{L}_{Z^\gamma} F^{\text{vac}})|)(t, x) \lesssim \frac{\Lambda}{(1+t+|x|)^{1+q}}, \quad (92)$$

$$\left| r\alpha(\mathcal{L}_{Z^\xi} F^{\text{vac}})(t, x) - \mathcal{F}^+(\mathcal{L}_{Z^\xi} F^{\text{vac}}(0, \cdot))\left(t - |x|, \frac{x}{|x|}\right) \right| \lesssim \frac{\Lambda}{(1+t+|x|)^q} \quad (93)$$

and $\mathcal{F}^+(F^{\text{vac}}(0, \cdot)) = \alpha^{\mathcal{I}^+}$. Together with Proposition 3.1 and Corollary 4.14, these estimates imply that $\mathcal{L}_{Z^\gamma}(F - F^{\text{vac}})$ satisfies the assumptions of Proposition 7.4 for any $|\gamma| \leq N - 10$. We then deduce, by a straightforward induction, that $\alpha^{\mathcal{I}^+}_\gamma = \mathcal{F}^+(\mathcal{L}_{Z^\gamma} F^{\text{vac}}(0, \cdot))$. Combining (93) with Proposition 7.16 then yields,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad r|\alpha(\mathcal{L}_{Z^\gamma} F) - \alpha(\mathcal{L}_{Z^\gamma} F^{\text{vac}})|(t, x) \lesssim \Lambda(1+t+|x|)^{-q}, \quad |\gamma| \leq N - 10.$$

On the other hand, Proposition 3.1 and (92) give, for any null component $\zeta \in \{\alpha, \rho, \sigma\}$,

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \quad r|\zeta(\mathcal{L}_{Z^\gamma} F) - \zeta(\mathcal{L}_{Z^\gamma} F^{\text{vac}})|(t, x) \lesssim \Lambda(1+t+|x|)^{-q}, \quad |\gamma| \leq N - 9,$$

which concludes the proof. □

Remark 7.19. According to Corollary 7.15 and Lemma 7.9, F^{vac} is in fact of class C^{N-5} . Moreover, if $N \geq 7$, then the statement of Proposition 7.18 still holds for any $|\gamma| \leq N - 7$ and the particular value $q = \frac{1}{2}$.

8. Conservation of the total energy of the system

Since (f, F) is a solution to the Vlasov–Maxwell system, the energy momentum tensor $\mathbb{T}[f, F]$, defined as

$$\mathbb{T}[f, F]_{\mu\nu} := \mathbb{T}[f]_{\mu\nu} + \mathbb{T}[F]_{\mu\nu}, \quad \mathbb{T}[f]_{\mu\nu} := \int_{\mathbb{R}_v^3} f v_\mu v_\nu \frac{dv}{v^0}, \quad \mathbb{T}[F]_{\mu\nu} := F_{\mu\beta} F_{\nu}{}^\beta - \frac{1}{4} \eta_{\mu\nu} F_{\xi\lambda} F^{\xi\lambda},$$

is divergence free. It provides the conservation of the total energy of the system

$$\mathbb{E}_t := \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f(t, x, v) dv dx + \frac{1}{2} \int_{\mathbb{R}_x^3} |F|^2(t, x) dx = \mathbb{E}_0, \quad |F|^2 = \sum_{0 \leq \mu < \nu \leq 3} |F_{\mu\nu}|^2 = |E|^2 + |B|^2.$$

We would like to relate \mathbb{E}_0 to the energy of the scattering states f_∞ and $\alpha^{\mathcal{I}^+}$. More precisely, the goal of this section is to prove

$$\mathbb{E}_\infty := \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f_\infty(x, v) dv dx + \frac{1}{4} \int_{\mathbb{R}_u} \int_{\mathbb{S}_\omega^2} |\alpha^{\mathcal{I}^+}|^2(u, \omega) d\mu_{\mathbb{S}^2} du = \mathbb{E}_0. \quad (94)$$

Note that $\mathbb{E}_\infty < +\infty$ according to Remark 6.35 and Proposition 7.16. The statement (94) is a consequence of $\mathbb{E}_t = \mathbb{E}_0$ and the following two propositions.

Proposition 8.1. *There holds*

$$\lim_{t \rightarrow +\infty} \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f(t, x, v) dv dx = \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f_\infty(x, v) dv dx.$$

Proof. Let $t \geq 3$ and perform the change of variables

$$x^j = y^j + \hat{v}^j t - \log(t) \hat{v}^\mu (F_{\mu j}^\infty(v) + \hat{v}^j F_{\mu 0}^\infty(v))$$

to get

$$\int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f(t, x, v) \, dv \, dx = \int_{\mathbb{R}_y^3} \int_{\mathbb{R}_v^3} v^0 f(t, X_{\mathcal{E}}(t, y, v), v) \, dv \, dy.$$

We then deduce that

$$\left| \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f(t, x, v) \, dv \, dx - \int_{\mathbb{R}_x^3} \int_{\mathbb{R}_v^3} v^0 f_{\infty}(x, v) \, dv \, dx \right| \leq \sup_{(x,v) \in \mathbb{R}^6} \langle x \rangle^{\frac{7}{2}} |v^0|^5 |f(t, X_{\mathcal{E}}(t, x, v), v) - f_{\infty}(x, v)|,$$

which, in view of $N_v \geq 12$, $N_x \geq \frac{11}{2}$ and [Remark 6.35](#), implies the result. □

Proposition 8.2. *We have*

$$\lim_{t \rightarrow +\infty} \frac{1}{2} \int_{\mathbb{R}_x^3} |F|^2(t, x) \, dx = \frac{1}{4} \int_{\mathbb{R}_u} \int_{\mathbb{S}_{\omega}^2} |\underline{\alpha}^{Z^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du.$$

Proof. Consider $\underline{u} \geq \tau \geq 3$ and introduce the domain $\mathcal{D}_{\underline{u}}^{\tau} = \{t + |x| \leq \underline{u}, t \geq \tau\}$, which is bounded by the truncated backward light cone $\underline{C}_{\underline{u}}^{\tau} := \{t + |x| = \underline{u}, t \geq \tau\}$ and $\{t = \tau\} \cap \{|x| \leq \underline{u} - \tau\}$. In the same spirit as [\(81\)](#), the divergence theorem, applied to $\mathbb{T}[F]_{\mu 0}$ in $\mathcal{D}_{\underline{u}}^{\tau}$, yields

$$\int_{\underline{C}_{\underline{u}}^{\tau}} \mathbb{T}[F]_{\underline{L}0} d\mu_{\underline{C}_{\underline{u}}^{\tau}} = \int_{|x| \leq \underline{u} - \tau} \mathbb{T}[F]_{00}(\tau, x) \, dx + \int_{(t,x) \in \mathcal{D}_{\underline{u}}^{\tau}} F_{0\lambda} J(f)^{\lambda} \, dx \, dt. \tag{95}$$

First, we have

$$\lim_{\underline{u} \rightarrow +\infty} \int_{|x| \leq \underline{u} - \tau} \mathbb{T}[F]_{00}(\tau, x) \, dx = \lim_{\underline{u} \rightarrow +\infty} \frac{1}{2} \int_{|x| \leq \underline{u} - \tau} |F|^2(\tau, x) \, dx = \frac{1}{2} \int_{\mathbb{R}_x^3} |F|^2(\tau, x) \, dx.$$

Next, since $|F|(t, x) \lesssim (1 + t + |x|)^{-1} (1 + |t - |x||)^{-1}$ by [\(BA1\)](#) and $|J(f)| \lesssim (1 + t + |x|)^{-3}$ by [Corollary 4.14](#),

$$\int_{(t,x) \in \mathcal{D}_{\underline{u}}^{\tau}} F_{0\lambda} J(f)^{\lambda} \, dx \, dt \lesssim \int_{t=\tau}^{+\infty} \int_{r=0}^{+\infty} \frac{r^2 \, dr \, dt}{(1+t+r)^4 (1+|t-r|)} \lesssim \int_{t=\tau}^{+\infty} \int_{r=0}^{+\infty} \frac{dr \, dt}{(1+t)^{\frac{3}{2}} (1+|t-r|)^{\frac{3}{2}}} \lesssim \tau^{-\frac{1}{2}}.$$

Recall from [Definition 2.16](#) the value of the null components of $\mathbb{T}[F]$. As

$$|\rho|(t, x) + |\sigma|(t, x) \lesssim (1 + t + |x|)^{-\frac{7}{4}}$$

by [Proposition 3.1](#) and in view of [Proposition 7.16](#), applied for $\eta > \frac{1}{2}$,

$$\begin{aligned} \int_{\underline{C}_{\underline{u}}^{\tau}} \mathbb{T}[F]_{\underline{L}0} \, d\mu_{\underline{C}_{\underline{u}}^{\tau}} &= \frac{1}{4} \int_{2\tau - \underline{u} \leq u \leq \underline{u}} \int_{\mathbb{S}_{\omega}^2} (|\underline{\alpha}(F)|^2 + |\rho(F)|^2 + |\sigma(F)|^2)(u, \underline{u}, \omega) r^2 \, d\mu_{\mathbb{S}^2} \, du \\ &= \frac{1}{4} \int_{2\tau - \underline{u} \leq u \leq \underline{u}} \int_{\mathbb{S}_{\omega}^2} r^2 |\underline{\alpha}(F)|^2(u, \underline{u}, \omega) \, d\mu_{\mathbb{S}^2} \, du + O(\underline{u}^{-\frac{1}{2}}) \\ &\xrightarrow{\underline{u} \rightarrow +\infty} \frac{1}{4} \int_{\mathbb{R}_u} \int_{\mathbb{S}_{\omega}^2} |\underline{\alpha}^{Z^+}|^2(u, \omega) \, d\mu_{\mathbb{S}^2} \, du. \end{aligned}$$

Letting $\underline{u} \rightarrow +\infty$ and then $\tau \rightarrow +\infty$ in [\(95\)](#) yields the result. □

Appendix A: Estimates for the gradients of the kernels

In order to estimate the kernels and their derivatives in the integrals of Propositions 5.3 and 5.7, we introduce the following class of terms.

Definition A.1. Let $(p, q, d, d_w) \in \mathbb{N}^4$. We define $\mathbf{S}_{p,q}^{d,d_w}$ as the set of the functions $\mathcal{G} : \mathbb{S}^2 \times \mathbb{R}^3 \rightarrow \mathbb{R}$ of the form

$$\mathcal{G}(\omega, v) = \frac{P(\hat{v}, \omega) Q(\mathbf{w}(\omega, v))}{|v^0|^p (1 + \omega \cdot \hat{v})^q}, \tag{96}$$

where P is a monomial of degree d in $(\hat{v}^1, \hat{v}^2, \hat{v}^3, \omega_1, \omega_2, \omega_3)$ and Q is a monomial of degree d_w in $\mathbf{w}_{\mu\nu}(\omega, v)$, where $0 \leq \mu < \nu \leq 3$.

All the kernels considered in this paper can be written as linear combination of such terms, with $d_w \in \llbracket 0, 3 \rrbracket$. Moreover, if $2q \geq d_w$, by a direct application of Lemma 5.4, one can bound $\mathcal{G}(\omega, v)$ in (96) by $|v^0|^{2q-d_w-p}$. The estimates of Corollaries 5.5 and 5.8 of the derivatives of the kernels then follows from the next result.

Lemma A.2. Let $(p, q, d, d_w) \in \mathbb{N}^4$ and consider $\mathcal{G} \in \mathbf{S}_{p,q}^{d,d_w}$. Then, for any multi-index γ , $\partial_v^\gamma \mathcal{G}(\omega, v)$ can be written as linear combination of terms belonging to certain $\mathbf{S}_{p_0,q_0}^{d_0,d_{w,0}}$, where

$$(p_0, q_0, d_0, d_{w,0}) \in \mathbb{N}^4, \quad 2q_0 - d_{w,0} - p_0 \leq 2q - d_w - p, \quad q - d_w \leq q_0 - d_{w,0}.$$

This implies $|\partial_v^\gamma \mathcal{G}(\cdot, v)| \lesssim |v^0|^{2q-d_w-p}$ if $2q \geq d_w$.

Proof. This follows from a straightforward induction and the following relations. For any $(i, j, k) \in \llbracket 1, 3 \rrbracket^3$,

$$\begin{aligned} \partial_{v^j} \hat{v}^i &= \frac{\delta_i^j - \hat{v}^i \hat{v}^j}{v^0}, & \partial_{v^j} \omega^i &= 0, & \partial_{v^j} |v^0|^{-p} &= -p \frac{\hat{v}^j}{|v^0|^{p+1}}, \\ \partial_{v^j} \mathbf{w}_{0i}(\omega, v) &= \frac{\delta_i^j - \hat{v}^i \hat{v}^j}{v^0}, & \partial_{v^j} \mathbf{w}_{ik}(\omega, v) &= \omega^i \frac{\delta_k^j - \hat{v}^k \hat{v}^j}{v^0} - \omega^k \frac{\delta_i^j - \hat{v}^i \hat{v}^j}{v^0}, \\ \partial_{v^j} \left(\frac{1}{1 + \omega \cdot \hat{v}} \right) &= \frac{\hat{v}^j}{v^0 (1 + \omega \cdot \hat{v})} - \frac{\mathbf{w}_{0j}(\omega, v)}{v^0 (1 + \omega \cdot \hat{v})^2}. \end{aligned} \quad \square$$

Appendix B: The radiation field of the derivatives of the Maxwell field

We fix, for all of this section, a C^1 solution G to the Maxwell equations (18) with a continuous source term J . We assume that there exist $C[G] > 0$ and $q > 0$ such that, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$,

$$|rG|(t, x) \leq C[G], \quad r|J|(t, x) + \sum_{|\gamma| \leq 1} |\rho(\mathcal{L}_{Z^\gamma} G)|(t, x) + |\sigma(\mathcal{L}_{Z^\gamma} G)|(t, x) \leq \frac{C[G]}{(1+t+|x|)^{1+q}}.$$

As a consequence, G verifies the hypotheses (75) of Proposition 7.3 and then has a radiation field $\underline{\alpha}^{\mathcal{I}^+}$. The purpose of this section is to prove that, for any $Z \in \mathbb{K}$, $\mathcal{L}_Z G$ has a radiation field $\underline{\alpha}_Z^{\mathcal{I}^+}$ which can be expressed in terms of the derivatives of $\underline{\alpha}^{\mathcal{I}^+}$. For this, we will use the following bounded functions

depending only on the spherical variables:

$$\begin{aligned} \omega_i &:= \langle \partial_{x^i}, \partial_r \rangle = \frac{x^i}{|x|}, & \omega_i^{e_A} &:= \langle \partial_{x^i}, e_A \rangle, & 1 \leq i \leq 3, & A \in \{\theta, \varphi\}, \\ \omega_1^{e_\theta} &= \cos(\varphi) \cos(\theta), & \omega_2^{e_\theta} &= \sin(\varphi) \cos(\theta), & \omega_3^{e_\theta} &= -\sin(\theta), \\ \omega_1^{e_\varphi} &= -\sin(\varphi), & \omega_2^{e_\varphi} &= \cos(\varphi), & \omega_3^{e_\varphi} &= 0, \end{aligned}$$

and we will work in the space of distributions $\mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$. For simplicity, we will simply write $\psi \rightharpoonup \psi^{\mathcal{I}^+}$ if the weak convergence

$$\psi(u, \underline{u}, \omega) \xrightarrow{u \rightarrow +\infty} \psi^{\mathcal{I}^+}(u, \omega) \quad \text{in } \mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$$

holds. In particular, the following convergences will be crucial for us.

Lemma B.1. *For any $1 \leq i \leq 3$ and $B \in \{\theta, \varphi\}$,*

$$\begin{aligned} |G| &\rightharpoonup 0, & \frac{1}{2}r\underline{L}(\underline{\alpha}(G)_{e_B}) &\rightharpoonup \partial_u(\underline{\alpha}_{e_B}^{\mathcal{I}^+}), & r^2L(\underline{\alpha}(G)_{e_B}) &\rightharpoonup -\underline{\alpha}_{e_B}^{\mathcal{I}^+}, \\ r^2\omega_i^A e_A(\underline{\alpha}(G)_{e_B}) &\rightharpoonup \omega_i^A e_A(\underline{\alpha}_{e_B}^{\mathcal{I}^+}), & r\rho(G) &\rightharpoonup 0, & r\sigma(G) &\rightharpoonup 0. \end{aligned}$$

Since $2r = \underline{u} - u$, we also have

$$rL(\underline{\alpha}(G)_{e_B}) \rightharpoonup 0, \quad r\omega_i^A e_A(\underline{\alpha}(G)_{e_B}) \rightharpoonup 0, \quad \rho(G) \rightharpoonup 0, \quad \sigma(G) \rightharpoonup 0.$$

Proof. The first weak convergence follows from $2|G|(u, \underline{u}, \omega) \leq C[G](\underline{u} - u)^{-1}$, so that $|G|(\cdot, \underline{u}, \cdot) \rightarrow 0$ uniformly on any compact subset of $\mathbb{R}_u \times \mathbb{S}^2$. The others are a direct consequence of the strong uniform convergence $r\underline{\alpha}(G)(u, \underline{u}, \omega) \rightarrow \underline{\alpha}^{\mathcal{I}^+}(u, \omega)$ as $\underline{u} \rightarrow +\infty$, which is given by [Proposition 7.3](#) since G satisfies [\(75\)](#).

- For the second one, use $r\underline{L} = \underline{L}r + 1$, $\underline{L} = 2\partial_u$ and that $\underline{\alpha}(F)_{e_B}(\cdot, \underline{u}, \cdot) \rightarrow 0$ uniformly on compact subsets of $\mathbb{R}_u \times \mathbb{S}^2$.
- The third one is in fact a strong and uniform convergence. Indeed, $r^2L(\underline{\alpha}(G)_{e_B}) = rL(r\underline{\alpha}(G)_{e_B}) - r\underline{\alpha}(G)_{e_B}$ and according to [\(76\)](#), $r|L(r\underline{\alpha}(G)_{e_B})| \lesssim \underline{u}^{-q}$.
- Next, fix $(t, r) \in \mathbb{R}_+^2$, $\psi \in C_c^\infty(\mathbb{R}_u \times \mathbb{S}^2)$ and denote by \vec{v}_i the vector field $\omega_i^{e_A} e_A$, which is the projection on the 2-spheres of ∂_{x^i} . Since $(re_\theta, re_\varphi) = (\partial_\theta, \partial_\varphi / \sin(\theta))$, we have

$$\begin{aligned} \omega_i^A r^2(e_A(\underline{\alpha}(G)_{e_B}))(t, r\omega)\psi(u, \omega) &= r\psi(u, \omega)\vec{v}_i \cdot \nabla(\underline{\alpha}(G)_{e_B}(t, r\omega)), \\ \omega_i^A e_A(\underline{\alpha}_{e_B}^{\mathcal{I}^+})(u, \omega)\psi(u, \omega) &= \psi(u, \omega)\vec{v}_i \cdot \nabla(\underline{\alpha}_{e_B}^{\mathcal{I}^+})(u, \omega), \end{aligned}$$

so that it suffices to apply the divergence theorem on \mathbb{S}^2 and to use $r\underline{\alpha}(G)_{e_B} \rightharpoonup \underline{\alpha}_{e_B}^{\mathcal{I}^+}$.

- Finally, the last two follow from $r|\rho(G)| + r|\sigma(G)| \lesssim \underline{u}^{-q}$. □

We now prove a result which directly implies [Proposition 7.4](#). We consider a more general setting since it does not complicate the proof and so we will be able to apply these properties in different contexts. For this, given a strictly increasing and unbounded sequence of advanced times $s = (\underline{u}_n)_{n \geq 0}$, we will write $\psi \rightharpoonup_s \psi^{\mathcal{I}^+}$ if the following weak convergence holds:

$$\psi(u, \underline{u}_n, \omega) \xrightarrow{n \rightarrow +\infty} \psi^{\mathcal{I}^+}(u, \omega) \quad \text{in } \mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2).$$

Proposition B.2. Consider G an $H^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3)$ 2-form and $\underline{\alpha}^{\mathcal{I}^+}$ an $L^2_{\text{loc}}(\mathbb{R}_u \times \mathbb{S}^2_\omega)$ 2-form tangential to the spheres. Assume that there exists a strictly increasing and unbounded sequence of advanced times $s = (\underline{u}_n)_{n \geq 0}$ such that

- $r\underline{\alpha}(G) \rightharpoonup_s \underline{\alpha}^{\mathcal{I}^+}$,
- all the weak convergences of Lemma B.1 hold, at least for the sequence $s \subset \mathbb{R}_{+,u}$.

Then, for any $Z \in \mathbb{K}$, there exists $\underline{\alpha}^{\mathcal{I}^+}_Z \in L^2_{\text{loc}}(\mathbb{R}_u \times \mathbb{S}^2_\omega)$, a 2-form tangential to the spheres, which satisfies $r\underline{\alpha}(\mathcal{L}_Z G) \rightharpoonup_s \underline{\alpha}^{\mathcal{I}^+}_Z$. Moreover, for any $1 \leq i \leq 3$ and $1 \leq j < k \leq 3$,

$$\begin{aligned} \underline{\alpha}^{\mathcal{I}^+}_{\partial_t} &= \nabla_u \underline{\alpha}^{\mathcal{I}^+}, & \underline{\alpha}^{\mathcal{I}^+}_{\partial_{x^i}} &= -\omega_i \nabla_u \underline{\alpha}^{\mathcal{I}^+}, & \underline{\alpha}^{\mathcal{I}^+}_S &= u \nabla_u \underline{\alpha}^{\mathcal{I}^+} + \underline{\alpha}^{\mathcal{I}^+}, \\ \underline{\alpha}^{\mathcal{I}^+}_{\Omega_{jk}} &= \mathcal{L}_{\Omega_{jk}}(\underline{\alpha}^{\mathcal{I}^+}), & \underline{\alpha}^{\mathcal{I}^+}_{\Omega_{0i}} &= -\omega_i u \nabla_u \underline{\alpha}^{\mathcal{I}^+} - 2\omega_i \underline{\alpha}^{\mathcal{I}^+} + \omega_i^{e_A} \nabla_{e_A} \underline{\alpha}^{\mathcal{I}^+}. \end{aligned}$$

Proof. In order to avoid technical difficulties related to the degeneracies of the spherical coordinate system, we will in fact prove weak convergences in

$$\mathcal{D}'(\mathbb{R}_u \times K), \quad K := \left\{ \omega \in \mathbb{S}^2 \mid \sin \theta > \frac{1}{8} \right\}.$$

The convergences in the full space $\mathcal{D}'(\mathbb{R}_u \times \mathbb{S}^2)$ can then be obtained by applying the upcoming results to another well-chosen spherical coordinate system.

We fix, for all of this proof, $B \in \{\theta, \varphi\}$, $i \in \llbracket 1, 3 \rrbracket$ and we recall that, for any $Z \in \mathbb{K}$,

$$r\underline{\alpha}(\mathcal{L}_Z G)_{e_B} = rZ(\underline{\alpha}(G)_{e_B}) - rG([Z, e_B], \underline{L}) - rG(e_B, [Z, \underline{L}]).$$

Then, we have

$$r\underline{\alpha}(\mathcal{L}_{\partial_t} G)_{e_B} = \frac{r}{2} \underline{L}(\underline{\alpha}(G)_{e_B}) + \frac{r}{2} L(\underline{\alpha}(G)_{e_B}) \rightharpoonup_s \partial_u(\underline{\alpha}^{\mathcal{I}^+}_{e_B}).$$

For the spatial translation $\partial_{x^i} = -\frac{1}{2}\omega_i \underline{L} + \frac{1}{2}\omega_i L + \omega_i^A e_A$, we use that

$$[\partial_{x^i}, \underline{L}] = -\frac{\omega_i^{e_A}}{r} e_A$$

and $[\partial_{x^i}, e_A] = \partial_{x^i}(\omega_j^{e_A}) \partial_{x^j}$, with $\partial_{x^i}(\omega_j^{e_A}) \lesssim r^{-1}$ on K . We get

$$\begin{aligned} r\underline{\alpha}(\mathcal{L}_{\partial_{x^i}} G)_{e_B} &= -\frac{\omega_i r}{2} \underline{L}(\underline{\alpha}(G)_{e_B}) + \frac{\omega_i r}{2} L(\underline{\alpha}(G)_{e_B}) + r\omega_i^A e_A(\underline{\alpha}(G)_{e_B}) - r\partial_{x^i}(\omega_j^{e_A}) G(e_B, \partial_{x^j}) + \omega_i^{e_A} G(e_B, e_A) \\ &\rightharpoonup_s -\omega_i \partial_u(\underline{\alpha}^{\mathcal{I}^+}_{e_B}). \end{aligned}$$

For the scaling, recall that $[S, \underline{L}] = -\underline{L}$ and $[S, e_B] = -e_B$. As $2S = u\underline{L} + (u + 2r)L$, we have

$$r\underline{\alpha}(\mathcal{L}_S G)_{e_B} = \frac{u}{2} r \underline{L}(\underline{\alpha}(G)_{e_B}) + \frac{u + 2r}{2} r L(\underline{\alpha}(G)_{e_B}) + 2rG(e_B, \underline{L}) \rightharpoonup_s u \partial_u(\underline{\alpha}^{\mathcal{I}^+}_{e_B}) + \underline{\alpha}^{\mathcal{I}^+}_{e_B}.$$

Next, for the Lorentz boost Ω_{0i} , we use

$$\Omega_{0i} = \frac{\omega_i}{2} (\underline{u}L - u\underline{L}) + t\omega_i^{e_A} e_A, \quad [\Omega_{0i}, e_B] = \frac{\omega_i^{e_B}}{2r} (\underline{u}L - u\underline{L}) + \frac{t}{r} \omega_i^{e_A} \nabla_{AB}^D e_D, \quad [\Omega_{0i}, \underline{L}] = \omega_i \underline{L} - \frac{u}{r} \omega_i^{e_A} e_A,$$

where \mathcal{V}_{AB}^D are the Christoffel symbols of \mathbb{S}^2 in the nonholonomic basis (e_θ, e_φ) . In particular, \mathcal{V}_{AB}^D is bounded on K . As $\underline{u} = u + 2r$ and $t = u + r$, we obtain

$$\begin{aligned} r\underline{\alpha}(\mathcal{L}_{\Omega_{0i}}G)_{e_B} &= -\frac{\omega_i u}{2} r\underline{L}(\underline{\alpha}(G)_{e_A}) + \frac{\omega_i(u+2r)}{2} r\underline{L}(\underline{\alpha}(G)_{e_B}) + \omega_i^{e_A}(u+r) r e_A(\underline{\alpha}(G)_{e_B}) - \frac{\omega_i^{e_B}}{2} u G(\underline{L}, \underline{L}) \\ &\quad + \frac{\omega_i^{e_B}}{2} \underline{u} G(\underline{L}, \underline{L}) - (u+r) \omega_i^{e_A} \mathcal{V}_{AB}^D G(e_D, \underline{L}) - \omega_i r G(e_B, \underline{L}) + \underline{u} \omega_i^{e_A} G(e_B, e_A). \end{aligned}$$

Since $G(\underline{L}, \underline{L}) = 0$ and $\underline{u}(|G(\underline{L}, \underline{L})| + |G(e_A, e_B)|) = (u+2r)(2|\rho(G)| + |\sigma(G)|) \rightarrow_s 0$, we get

$$\begin{aligned} r\underline{\alpha}(\mathcal{L}_{\Omega_{0i}}G)_{e_B} &\rightarrow_s -\omega_i u \partial_u(\underline{\alpha}_{e_B}^{I^+}) - \omega_i \underline{\alpha}_{e_B}^{I^+} + \omega_i^{e_A} e_A(\underline{\alpha}_{e_B}^{I^+}) - 0 + 0 - \omega_i^{e_A} \mathcal{V}_{AB}^D \underline{\alpha}_{e_D}^{I^+} - \omega_i \underline{\alpha}_{e_B}^{I^+} + 0 \\ &= -\omega_i u \partial_u(\underline{\alpha}_{e_B}^{I^+}) - 2\omega_i \underline{\alpha}_{e_B}^{I^+} + \omega_i^{e_A} \mathcal{V}_{e_A}(\underline{\alpha}^{I^+})_{e_B}. \end{aligned}$$

Finally, we recall the expression of the rotations in the spherical coordinate system (t, r, θ, φ) ,

$$\Omega_{12} = \partial_\varphi, \quad \Omega_{13} = \cos(\varphi)\partial_\theta - \cot(\theta)\sin(\varphi)\partial_\varphi, \quad \Omega_{23} = -\sin(\varphi)\partial_\theta - \cot(\theta)\cos(\varphi)\partial_\varphi.$$

In particular, these vector fields, tangential to the spheres, are well-defined on $\mathcal{I}^+ \simeq \mathbb{R}_u \times \mathbb{S}^2$. Fix now $(j, k) \in \llbracket 1, 3 \rrbracket^2$ and write $\Omega_{jk} = \Omega_{jk}^\theta \partial_\theta + \Omega_{jk}^\varphi \partial_\varphi$. Note, using first $[\Omega_{jk}, \underline{L}] = 0$ and then the expression of the Lie derivative in the spherical coordinate system, that

$$\underline{\alpha}(\mathcal{L}_{\Omega_{jk}}G)_{\partial_B} = \mathcal{L}_{\Omega_{jk}}(\underline{\alpha}(G))_{\partial_B} = \Omega_{jk}(\underline{\alpha}(G)_{\partial_B}) + \partial_B(\Omega_{jk}^A)\underline{\alpha}(G)_{\partial_A}.$$

Recall now that $(re_\theta, re_\varphi) = (\partial_\theta, \partial_\varphi/\sin(\theta))$ on $\mathbb{R}_+ \times \mathbb{R}^3$ and $(e_\theta, e_\varphi) = (\partial_\theta, \partial_\varphi/\sin(\theta))$ on $\mathbb{R}_u \times \mathbb{S}^2$. Hence, using $r\underline{\alpha}(G)_{e_A} \rightarrow_s \underline{\alpha}_{e_A}^{I^+}$ and since any of the quantities considered is smooth and bounded on K ,

$$\begin{aligned} r\underline{\alpha}(\mathcal{L}_{\Omega_{jk}}G)_{e_\theta} &= \Omega_{jk}^\theta \partial_\theta(r\underline{\alpha}(G)_{e_\theta}) + \Omega_{jk}^\varphi \partial_\varphi(r\underline{\alpha}(G)_{e_\theta}) + \partial_\theta(\Omega_{jk}^\theta)r\underline{\alpha}(G)_{e_\theta} + \sin(\theta)\partial_\theta(\Omega_{jk}^\varphi)r\underline{\alpha}(G)_{e_\varphi} \\ &\rightarrow_s \Omega_{jk}(\underline{\alpha}_{e_\theta}^{I^+}) + \partial_\theta(\Omega_{jk}^\theta)\underline{\alpha}_{e_\theta}^{I^+} + \sin(\theta)\partial_\theta(\Omega_{jk}^\varphi)\underline{\alpha}_{e_\varphi}^{I^+} \\ &= \Omega_{jk}(\underline{\alpha}_{\partial_\theta}^{I^+}) + \partial_\theta(\Omega_{jk}^A)\underline{\alpha}_{\partial_A}^{I^+} = \mathcal{L}_{\Omega_{kl}}(\underline{\alpha}^{I^+})_{e_\theta}. \end{aligned}$$

Similarly, we get

$$\begin{aligned} r\underline{\alpha}(\mathcal{L}_{\Omega_{jk}}G)_{e_\varphi} &= \Omega_{jk}(\underline{\alpha}(G)_{e_\varphi}) - \Omega_{kl}\left(\frac{1}{\sin\theta}\right)r\underline{\alpha}(G)_{e_\varphi} + \frac{1}{\sin\theta}\partial_\varphi(\Omega_{jk}^\theta)r\underline{\alpha}(G)_{e_\theta} + \partial_\varphi(\Omega_{jk}^\varphi)r\underline{\alpha}(G)_{e_\varphi} \\ &\rightarrow_s \Omega_{jk}(\underline{\alpha}_{e_\varphi}^{I^+}) - \Omega_{kl}\left(\frac{1}{\sin\theta}\right)\underline{\alpha}_{e_\varphi}^{I^+} + \frac{1}{\sin\theta}\partial_\varphi(\Omega_{jk}^\theta)\underline{\alpha}_{e_\theta}^{I^+} + \partial_\varphi(\Omega_{jk}^\varphi)\underline{\alpha}_{e_\varphi}^{I^+} = \mathcal{L}_{\Omega_{kl}}(\underline{\alpha}^{I^+})_{e_\varphi}. \quad \square \end{aligned}$$

Appendix C: Remarks on F^∞ and the modified characteristics

C.1. Alternative expression for F^∞ . We could define F^∞ in a slightly different way. However, contrary to what we did in [Section 6.2](#), we could not define in such a way $\mathcal{L}_{Z^\nu}(F)^\infty$ for the derivatives of order $|\gamma| = N - 1$. Using the representation formula for the wave equation satisfied by $F_{\mu\nu}$,

$$F_{\mu\nu} = F_{\mu\nu}^{\text{hom}} + [f]_{\mu\nu}^{\text{inh}}, \quad [f]_{\mu\nu}^{\text{inh}}(t, x) := \frac{1}{4\pi} \int_{|y-x| \leq t} \int_{\mathbb{R}^3} (\hat{v}_\mu \partial_{x^\nu} f - \hat{v}_\nu \partial_{x^\mu} f)(t - |y-x|, y, v) dv \frac{dy}{|y-x|}.$$

In order to investigate the asymptotic behavior of $[f]_{\mu\nu}^{\text{inh}}$, it is important to determine the asymptotic profile of the source term of the wave equation. In particular, we need to obtain a better estimate than the one

given by Proposition 4.15 which does not provide the expected time decay t^{-4} . The starting point consists in observing that a kind of null condition holds,

$$\begin{aligned}
 t(\partial_{x^i} + \hat{v}_i \partial_t) &= \Omega_{0i} + z_{0i} \partial_t = \widehat{\Omega}_{0i} - v^0 \partial_{v^i} + \partial_t z_{0i} - \hat{v}_i = \widehat{\Omega}_{0i} - \partial_{v^i} v^0 + \partial_t z_{0i}, \quad 1 \leq i \leq 3, \\
 t(\hat{v}_j \partial_{x^k} - \hat{v}_k \partial_{x^j}) &= \hat{v}_j \widehat{\Omega}_{0k} - \hat{v}_k \widehat{\Omega}_{0j} - \partial_t z_{jk} - \hat{v}_j \partial_{v^k} v^0 + \hat{v}_k \partial_{v^j} v^0, \quad 1 \leq j < k \leq 3.
 \end{aligned}$$

Hence, using the convention $\widehat{\Omega}_{00} = 0$ and performing integration by parts, we obtain, for any $0 \leq \mu < \nu \leq 3$,

$$\int_{\mathbb{R}^3} \hat{v}_\mu \partial_{x^\nu} f - \hat{v}_\nu \partial_{x^\mu} f \, dv = \frac{1}{t} \int_{\mathbb{R}^3} (\hat{v}_\mu \widehat{\Omega}_{0\nu} f - \hat{v}_\nu \widehat{\Omega}_{0\mu} f) \, dv - \frac{1}{t} \int_{\mathbb{R}^3} \partial_t(z_{\mu\nu} f) \, dv.$$

The leading-order term of its asymptotic expansion is the first term on the right-hand side. Its behavior can be easily obtained from Proposition 6.5. Following the proof of Proposition 4.15, one could prove that last term almost decay as t^{-5} . It will then be convenient to use the notation $Q_\infty^{\widehat{\Omega}_{0\ell}}$ in order to denote Q_∞^κ , where $\widehat{Z}^\kappa = \widehat{\Omega}_{0\ell}$, and to set $Q_\infty^{\widehat{\Omega}_{00}} := 0$. Following the proof Proposition 6.10, we could obtain

$$\lim_{t \rightarrow +\infty} t^2 [f]_{\mu\nu}^{\text{inh}}(t, x + t\hat{v}) := \frac{1}{4\pi} \int_{\substack{|z| \leq 1 \\ |z + \hat{v}| < 1 - |z|}} (|v^0|^5 (\hat{v}_\mu Q_\infty^{\widehat{\Omega}_{0\nu}} - \hat{v}_\nu Q_\infty^{\widehat{\Omega}_{0\mu}})) \left(\frac{z + \hat{v}}{1 - |z|} \right) \frac{dz}{|z|(1 - |z|)^4},$$

which is necessarily equal to F^∞ .

C.2. The support of the corrections of the linear characteristics and commutators. We could obtain similar results by modifying the trajectories and the homogeneous vector fields only inside the light cone. More precisely, we could consider, for $\widehat{Z} \in \widehat{\mathbb{P}}_0 \setminus \{\partial_t, \partial_{x^1}, \partial_{x^2}, \partial_{x^3}\}$,

$$\widetilde{X}_\mathcal{C}(t, x, v) := x + t\hat{v} + \mathcal{C}(t, v)\chi(t - |x - t\hat{v}|), \quad \widetilde{Z}^{\text{mod}} := \widehat{Z} + \mathcal{C}_Z^i \chi(t - r) \partial_{x^i},$$

where χ is a cutoff function satisfying $\chi(s) = 0$ for $s \leq 1$ and $\chi(s) = 1$ for $s \geq 2$. It is not surprising that all the results proved for $X_\mathcal{C}$ and \widehat{Z}^{mod} hold as well with these corrections since the Vlasov field enjoys strong decay properties in the exterior of the light cone (see Lemma 2.6). We could even avoid the loss of the weight z^{β_H} in Proposition 6.28 and Corollary 6.29. Indeed, these weights come from the identity $x^i/t = (x^i - t\hat{v}^i)/t + \hat{v}^i$ that we performed during the proof of Proposition 6.26. On the support of χ , we can simply use that $|x|/t \leq 1$. However, we could not save any $\langle x \rangle$ weight in the analogue version of the scattering statement of Proposition 6.34 since we would have to lose a power of z^{β_H} in order to estimate $|v^0|^\xi \partial_v^\xi (\chi(t - |x + t\hat{v}|))$.

Acknowledgements

I would like to thank Jacques Smulevici for suggesting this problem to me. This work was conducted within the France 2030 framework programme, the Centre Henri Lebesgue ANR-11-LABX-0020-01.

References

[Andersson et al. 2018] L. Andersson, P. Blue, and J. Joudioux, “Hidden symmetries and decay for the Vlasov equation on the Kerr spacetime”, *Comm. Partial Differential Equations* **43**:1 (2018), 47–65. MR Zbl

[Bardos and Degond 1985] C. Bardos and P. Degond, “Global existence for the Vlasov–Poisson equation in 3 space variables with small initial data”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **2**:2 (1985), 101–118. MR Zbl

- [Bigorgne 2020a] L. Bigorgne, “Sharp asymptotic behavior of solutions of the 3d Vlasov–Maxwell system with small data”, *Comm. Math. Phys.* **376**:2 (2020), 893–992. [MR](#) [Zbl](#)
- [Bigorgne 2020b] L. Bigorgne, “A vector field method for massless relativistic transport equations and applications”, *J. Funct. Anal.* **278**:4 (2020), art. id. 108365. [MR](#) [Zbl](#)
- [Bigorgne 2021a] L. Bigorgne, “Asymptotic properties of the solutions to the Vlasov–Maxwell system in the exterior of a light cone”, *Int. Math. Res. Not.* **2021**:5 (2021), 3729–3793. [MR](#) [Zbl](#)
- [Bigorgne 2021b] L. Bigorgne, “Sharp asymptotics for the solutions of the three-dimensional massless Vlasov–Maxwell system with small data”, *Ann. Henri Poincaré* **22**:1 (2021), 219–273. [MR](#) [Zbl](#)
- [Bigorgne 2022] L. Bigorgne, *Asymptotic properties of small data solutions of the Vlasov–Maxwell system in high dimensions*, Mém. Soc. Math. France (N.S.) **172**, Soc. Math. France, Paris, 2022. [MR](#) [Zbl](#)
- [Bigorgne 2023] L. Bigorgne, “Decay estimates for the massless Vlasov equation on Schwarzschild spacetimes”, *Ann. Henri Poincaré* **24**:11 (2023), 3763–3836. [MR](#) [Zbl](#)
- [Bigorgne et al. 2021] L. Bigorgne, D. Fajman, J. Joudioux, J. Smulevici, and M. Thaller, “Asymptotic stability of Minkowski space-time with non-compactly supported massless Vlasov matter”, *Arch. Ration. Mech. Anal.* **242**:1 (2021), 1–147. [MR](#) [Zbl](#)
- [Bouchut et al. 2003] F. Bouchut, F. Golse, and C. Pallard, “Classical solutions and the Glassey–Strauss theorem for the 3D Vlasov–Maxwell system”, *Arch. Ration. Mech. Anal.* **170**:1 (2003), 1–15. [MR](#) [Zbl](#)
- [Chaturvedi 2021] S. Chaturvedi, “Stability of vacuum for the Boltzmann equation with moderately soft potentials”, *Ann. PDE* **7**:2 (2021), art. id. 15. [MR](#) [Zbl](#)
- [Chaturvedi 2022] S. Chaturvedi, “Stability of vacuum for the Landau equation with hard potentials”, *Probab. Math. Phys.* **3**:4 (2022), 791–838. [MR](#) [Zbl](#)
- [Chaturvedi et al. 2023] S. Chaturvedi, J. Luk, and T. Nguyen, “The Vlasov–Poisson–Landau system in the weakly collisional regime”, *J. Amer. Math. Soc.* **36**:4 (2023), 1103–1189. [MR](#) [Zbl](#)
- [Choi and Ha 2011] S.-H. Choi and S.-Y. Ha, “Asymptotic behavior of the nonlinear Vlasov equation with a self-consistent force”, *SIAM J. Math. Anal.* **43**:5 (2011), 2050–2077. [MR](#) [Zbl](#)
- [Choi and Kwon 2016] S.-H. Choi and S. Kwon, “Modified scattering for the Vlasov–Poisson system”, *Nonlinearity* **29**:9 (2016), 2755–2774. [MR](#) [Zbl](#)
- [Christodoulou and Klainerman 1990] D. Christodoulou and S. Klainerman, “Asymptotic properties of linear field equations in Minkowski space”, *Comm. Pure Appl. Math.* **43**:2 (1990), 137–199. [MR](#) [Zbl](#)
- [DiPerna and Lions 1989] R. J. DiPerna and P.-L. Lions, “Global weak solutions of Vlasov–Maxwell systems”, *Comm. Pure Appl. Math.* **42**:6 (1989), 729–757. [MR](#) [Zbl](#)
- [Duan 2022] X. Duan, “Sharp decay estimates for the Vlasov–Poisson and Vlasov–Yukawa systems with small data”, *Kinet. Relat. Models* **15**:1 (2022), 119–146. [MR](#) [Zbl](#)
- [Fajman et al. 2017] D. Fajman, J. Joudioux, and J. Smulevici, “A vector field method for relativistic transport equations with applications”, *Anal. PDE* **10**:7 (2017), 1539–1612. [MR](#) [Zbl](#)
- [Fajman et al. 2021] D. Fajman, J. Joudioux, and J. Smulevici, “The stability of the Minkowski space for the Einstein–Vlasov system”, *Anal. PDE* **14**:2 (2021), 425–531. [MR](#) [Zbl](#)
- [Flynn et al. 2023] P. Flynn, Z. Ouyang, B. Pausader, and K. Widmayer, “Scattering map for the Vlasov–Poisson system”, *Peking Math. J.* **6**:2 (2023), 365–392. [MR](#) [Zbl](#)
- [Glassey 1996] R. T. Glassey, *The Cauchy problem in kinetic theory*, SIAM, Philadelphia, PA, 1996. [MR](#) [Zbl](#)
- [Glassey and Schaeffer 1985] R. T. Glassey and J. Schaeffer, “On symmetric solutions of the relativistic Vlasov–Poisson system”, *Comm. Math. Phys.* **101**:4 (1985), 459–473. [MR](#) [Zbl](#)
- [Glassey and Schaeffer 1988] R. T. Glassey and J. W. Schaeffer, “Global existence for the relativistic Vlasov–Maxwell system with nearly neutral initial data”, *Comm. Math. Phys.* **119**:3 (1988), 353–384. [MR](#) [Zbl](#)
- [Glassey and Schaeffer 1990] R. Glassey and J. Schaeffer, “On the ‘one and one-half dimensional’ relativistic Vlasov–Maxwell system”, *Math. Methods Appl. Sci.* **13**:2 (1990), 169–179. [MR](#) [Zbl](#)
- [Glassey and Schaeffer 1997] R. Glassey and J. Schaeffer, “The ‘two and one-half-dimensional’ relativistic Vlasov Maxwell system”, *Comm. Math. Phys.* **185**:2 (1997), 257–284. [MR](#) [Zbl](#)

- [Glasse and Schaeffer 1998] R. T. Glassey and J. Schaeffer, “The relativistic Vlasov–Maxwell system in two space dimensions, I, II”, *Arch. Ration. Mech. Anal.* **141**:4 (1998), 331–374. [MR](#) [Zbl](#)
- [Glasse and Strauss 1986] R. T. Glassey and W. A. Strauss, “Singularity formation in a collisionless plasma could occur only at high velocities”, *Arch. Ration. Mech. Anal.* **92**:1 (1986), 59–90. [MR](#) [Zbl](#)
- [Glasse and Strauss 1987a] R. T. Glassey and W. A. Strauss, “Absence of shocks in an initially dilute collisionless plasma”, *Comm. Math. Phys.* **113**:2 (1987), 191–208. [MR](#) [Zbl](#)
- [Glasse and Strauss 1987b] R. T. Glassey and W. A. Strauss, “High velocity particles in a collisionless plasma”, *Math. Methods Appl. Sci.* **9**:1 (1987), 46–52. [MR](#) [Zbl](#)
- [Glasse and Strauss 1989] R. T. Glassey and W. A. Strauss, “Large velocities in the relativistic Vlasov–Maxwell equations”, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **36**:3 (1989), 615–627. [MR](#) [Zbl](#)
- [Hwang et al. 2011] H. J. Hwang, A. Rendall, and J. J. L. Velázquez, “Optimal gradient estimates and asymptotic behaviour for the Vlasov–Poisson system with small initial data”, *Arch. Ration. Mech. Anal.* **200**:1 (2011), 313–360. [MR](#) [Zbl](#)
- [Ionescu et al. 2022] A. D. Ionescu, B. Pausader, X. Wang, and K. Widmayer, “On the asymptotic behavior of solutions to the Vlasov–Poisson system”, *Int. Math. Res. Not.* **2022**:12 (2022), 8865–8889. [MR](#) [Zbl](#)
- [Klainerman 1985] S. Klainerman, “Uniform decay estimates and the Lorentz invariance of the classical wave equation”, *Comm. Pure Appl. Math.* **38**:3 (1985), 321–332. [MR](#) [Zbl](#)
- [Klainerman and Staffilani 2002] S. Klainerman and G. Staffilani, “A new approach to study the Vlasov–Maxwell system”, *Commun. Pure Appl. Anal.* **1**:1 (2002), 103–125. [MR](#) [Zbl](#)
- [Kunze 2015] M. Kunze, “Yet another criterion for global existence in the 3D relativistic Vlasov–Maxwell system”, *J. Differential Equations* **259**:9 (2015), 4413–4442. [MR](#) [Zbl](#)
- [Lindblad and Schlue 2023] H. Lindblad and V. Schlue, “Scattering from infinity for semilinear wave equations satisfying the null condition or the weak null condition”, *J. Hyperbolic Differ. Equ.* **20**:1 (2023), 155–218. [MR](#) [Zbl](#)
- [Lindblad and Taylor 2020] H. Lindblad and M. Taylor, “Global stability of Minkowski space for the Einstein–Vlasov system in the harmonic gauge”, *Arch. Ration. Mech. Anal.* **235**:1 (2020), 517–633. [MR](#) [Zbl](#)
- [Luk and Strain 2014] J. Luk and R. M. Strain, “A new continuation criterion for the relativistic Vlasov–Maxwell system”, *Comm. Math. Phys.* **331**:3 (2014), 1005–1027. [MR](#) [Zbl](#)
- [Luk and Strain 2016] J. Luk and R. M. Strain, “Strichartz estimates and moment bounds for the relativistic Vlasov–Maxwell system”, *Arch. Ration. Mech. Anal.* **219**:1 (2016), 445–552. [MR](#) [Zbl](#)
- [Pallard 2005] C. Pallard, “On the boundedness of the momentum support of solutions to the relativistic Vlasov–Maxwell system”, *Indiana Univ. Math. J.* **54**:5 (2005), 1395–1409. [MR](#) [Zbl](#)
- [Pallard 2015] C. Pallard, “A refined existence criterion for the relativistic Vlasov–Maxwell system”, *Commun. Math. Sci.* **13**:2 (2015), 347–354. [MR](#) [Zbl](#)
- [Pankavich 2022] S. Pankavich, “Asymptotic dynamics of dispersive, collisionless plasmas”, *Comm. Math. Phys.* **391**:2 (2022), 455–493. [MR](#) [Zbl](#)
- [Pankavich 2023] S. Pankavich, “Scattering and asymptotic behavior of solutions to the Vlasov–Poisson system in high dimension”, *SIAM J. Math. Anal.* **55**:5 (2023), 4727–4750. [MR](#) [Zbl](#)
- [Patel 2018] N. Patel, “Three new results on continuation criteria for the 3D relativistic Vlasov–Maxwell system”, *J. Differential Equations* **264**:3 (2018), 1841–1885. [MR](#) [Zbl](#)
- [Pausader and Widmayer 2021] B. Pausader and K. Widmayer, “Stability of a point charge for the Vlasov–Poisson system: the radial case”, *Comm. Math. Phys.* **385**:3 (2021), 1741–1769. [MR](#) [Zbl](#)
- [Pausader et al. 2024] B. Pausader, K. Widmayer, and J. Yang, “Stability of a point charge for the repulsive Vlasov–Poisson system”, *J. Eur. Math. Soc. (JEMS)* (online publication August 2024).
- [Rein 1990] G. Rein, “Generic global solutions of the relativistic Vlasov–Maxwell system of plasma physics”, *Comm. Math. Phys.* **135**:1 (1990), 41–78. [MR](#) [Zbl](#)
- [Rein 2004] G. Rein, “Global weak solutions to the relativistic Vlasov–Maxwell system revisited”, *Commun. Math. Sci.* **2**:2 (2004), 145–158. [MR](#) [Zbl](#)

- [Schaeffer 2004] J. Schaeffer, “A small data theorem for collisionless plasma that includes high velocity particles”, *Indiana Univ. Math. J.* **53**:1 (2004), 1–34. [MR](#) [Zbl](#)
- [Schaeffer 2021] J. Schaeffer, “An improved small data theorem for the Vlasov–Poisson system”, *Commun. Math. Sci.* **19**:3 (2021), 721–736. [MR](#) [Zbl](#)
- [Smulevici 2016] J. Smulevici, “Small data solutions of the Vlasov–Poisson system and the vector field method”, *Ann. PDE* **2**:2 (2016), art. id. 11. [MR](#) [Zbl](#)
- [Sogge 1995] C. D. Sogge, *Lectures on nonlinear wave equations*, Monogr. Anal. **2**, International Press, Boston, MA, 1995. [MR](#) [Zbl](#)
- [Sospedra-Alfonso and Illner 2010] R. Sospedra-Alfonso and R. Illner, “Classical solvability of the relativistic Vlasov–Maxwell system with bounded spatial density”, *Math. Methods Appl. Sci.* **33**:6 (2010), 751–757. [MR](#) [Zbl](#)
- [Taylor 2017] M. Taylor, “The global nonlinear stability of Minkowski space for the massless Einstein–Vlasov system”, *Ann. PDE* **3**:1 (2017), art. id. 9. [MR](#) [Zbl](#)
- [Wang 2022a] X. Wang, “Global solution of the 3D relativistic Vlasov–Maxwell system for large data with cylindrical symmetry”, preprint, 2022. [arXiv 2203.01199](#)
- [Wang 2022b] X. Wang, “Propagation of regularity and long time behavior of the 3D massive relativistic transport equation, II: Vlasov–Maxwell system”, *Comm. Math. Phys.* **389**:2 (2022), 715–812. [MR](#) [Zbl](#)
- [Wang 2023] X. Wang, “Decay estimates for the 3D relativistic and non-relativistic Vlasov–Poisson systems”, *Kinet. Relat. Models* **16**:1 (2023), 1–19. [MR](#) [Zbl](#)
- [Wei and Yang 2021] D. Wei and S. Yang, “On the 3D relativistic Vlasov–Maxwell system with large Maxwell field”, *Comm. Math. Phys.* **383**:3 (2021), 2275–2307. [MR](#) [Zbl](#)
- [Wollman 1984] S. Wollman, “An existence and uniqueness theorem for the Vlasov–Maxwell system”, *Comm. Pure Appl. Math.* **37**:4 (1984), 457–462. [MR](#) [Zbl](#)

Received 26 Aug 2022. Revised 31 Aug 2023. Accepted 21 Nov 2023.

LÉO BIGORGNE: leo.bigorgne@univ-rennes.fr

Université de Rennes, CNRS, IRMAR - UMR 6625, Rennes, France

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: "Linear Ramp"


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2025 is US \$475/year for the electronic version, and \$735/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 18 No. 3 2025

Rotating spirals in segregated reaction-diffusion systems	549
ARIEL SALORT, SUSANNA TERRACINI, GIANMARIA VERZINI and ALESSANDRO ZILIO	
Stahl–Totik regularity for continuum Schrödinger operators	591
BENJAMIN EICHINGER and MILIVOJE LUKIĆ	
Global existence and modified scattering for the solutions to the Vlasov–Maxwell system with a small distribution function	629
LÉO BIGORGNE	
Strong ill-posedness in L^∞ for the Riesz transform problem	715
TAREK M. ELGINDI and KARIM R. SHIKH KHALIL	
Fractal uncertainty for discrete two-dimensional Cantor sets	743
ALEX COHEN	
Linear potentials and applications in conformal geometry	773
SHIGUANG MA and JIE QING	