

ANALYSIS & PDE

Volume 18

No. 6

2025

GÁBOR SZABÓ AND LISE WOUTERS

**EQUIVARIANT PROPERTY GAMMA
AND THE TRACIAL LOCAL-TO-GLOBAL PRINCIPLE FOR
 C^* -DYNAMICS**



EQUIVARIANT PROPERTY GAMMA AND THE TRACIAL LOCAL-TO-GLOBAL PRINCIPLE FOR C^* -DYNAMICS

GÁBOR SZABÓ AND LISE WOUTERS

We consider the notion of equivariant uniform property Gamma for actions of countable discrete groups on C^* -algebras that admit traces. In case the group is amenable and the C^* -algebra has a compact tracial state space, we prove that this property implies a kind of tracial local-to-global principle for the C^* -dynamical system, generalizing a recent such principle for C^* -algebras exhibited in work of Castillejos et al. For actions on simple nuclear \mathcal{Z} -stable C^* -algebras, we use this to prove that equivariant uniform property Gamma is equivalent to equivariant \mathcal{Z} -stability, generalizing a result of Gardella, Hirshberg, and Vaccaro.

Introduction	1385
1. Preliminaries	1390
2. Equivariant uniform property Gamma	1398
3. Dynamical complemented partitions of unity	1402
4. Dynamical tracial local-to-global principle	1417
5. Equivariant Jiang–Su stability	1425
Acknowledgements	1429
References	1429

Introduction

This article aims to extend the fine structure theory for actions of amenable groups on finite simple C^* -algebras, in particular those covered by the Elliott program. The classification of such C^* -algebras, which mirrors the celebrated Connes–Haagerup classification of injective factors [Connes 1976; Haagerup 1987], has been nearly completed as a culmination of numerous articles by many researchers over the past decade, such as [Elliott et al. 2024; 2020; Gong and Lin 2020; 2022; Gong et al. 2020a; 2020b; Schafhauser 2020; Tikuisis et al. 2017]. Furthermore, Carrion, Gabe, Schafhauser, Tikuisis, and White have announced an eagerly awaited new conceptual proof of the classification theorem [Carrión et al. 2023b], which does not directly rely on the prior works related to tracial approximation. By now it has been recognized that the next natural step is to understand the underlying symmetries of classifiable C^* -algebras, which can be interpreted as the goal to classify group actions on them. This mirrors the work

Szabó was supported by BOF project C14/19/088, funded by the research council of KU Leuven, and research project G085020N, funded by the Research Foundation Flanders (FWO). Wouters was supported by Ph.D. grant 11B6620N, funded by the Research Foundation Flanders (FWO).

MSC2020: 46L35, 46L55.

Keywords: C^* -dynamics, amenability, Jiang–Su algebra.

of Connes, Jones, Ocneanu and others [Connes 1977; Jones 1980; Katayama et al. 1998; Kawahigashi et al. 1992; Masuda 2007; 2013; Ocneanu 1985; Sutherland and Takesaki 1989]. When it comes to classifying group actions on C^* -algebras, a number of researchers have introduced many sophisticated methods over the years to classify specific kinds of group actions utilizing certain Rokhlin-type properties [Evans and Kishimoto 1997; Izumi 2004a; 2004b; Izumi and Matui 2010; 2021a; 2021b; Katsura and Matui 2008; Kishimoto 1995; 1998a; 1998b; Matui 2008; 2010; 2011; Nakamura 2000; Sato 2010; Szabó 2021a]. In direct comparison to the generality achieved for actions on von Neumann algebras, the implementation of the involved methods (in particular the Evans–Kishimoto intertwining argument) for actions on C^* -algebras remained challenging beyond some specific classes of actions or acting groups.

To combat these methodological obstacles, the first author introduced a categorical framework in [Szabó 2021c] to open up the classification of C^* -dynamics up to cocycle conjugacy to methodology directly inspired by [Elliott 2010]. For actions on classifiable C^* -algebras without traces, the so-called Kirchberg algebras [1995], this idea led to the recent breakthrough in [Gabe and Szabó 2022; 2024]. The main result of said work implies that, given any countable amenable group G , any outer G -action on a Kirchberg algebra is uniquely determined by its KK^G -class up to cocycle conjugacy.

Although one might be tempted to guess that similar breakthrough results ought to be in reach for actions on finite classifiable C^* -algebras, one still has a long way to go before such a goal can be achieved. In analogy to the original obstacles to classify all simple nuclear C^* -algebras [Rørdam 2003; Toms 2008; Villadsen 1999], there are basic structural questions to be settled before a classification theory such as in [Gabe and Szabó 2024] can be attempted on finite C^* -algebras. When concerned with just the underlying C^* -algebras, this is already a serious challenge. On the one hand, there is the question whether the C^* -algebras under consideration automatically satisfy certain properties predicted by classification. For the purpose of this article we highlight the property of Jiang–Su stability. If \mathcal{Z} is the so-called Jiang–Su algebra from [Jiang and Su 1999], then a C^* -algebra A is called Jiang–Su stable or \mathcal{Z} -stable when $A \cong A \otimes \mathcal{Z}$. Although this might seem like a technical property at first glance, it becomes natural with more context: Firstly, \mathcal{Z} behaves (as a C^* -algebra) very much like an infinite-dimensional version of the complex numbers \mathbb{C} , for instance at the level of K -theory and traces. Secondly, there is by now a pile of evidence that \mathcal{Z} -stability holds automatically for many C^* -algebras arising from various constructions like the crossed product [Kerr 2020; Kerr and Naryshkin 2021; Kerr and Szabó 2020; Toms and Winter 2013]. The discovery that \mathcal{Z} -stability does in fact *not* hold automatically for all simple nuclear C^* -algebras has, among other things, led to the nearly proven Toms–Winter conjecture, which asserts that \mathcal{Z} -stability can only hold or fail in conjunction with some other, a priori different, regularity conditions.

On the other hand, there is the question about precisely what additional structural consequences (not necessarily equivalent characterizations) are shared by Jiang–Su stable C^* -algebras, a good example of which is the recent breakthrough work [Castillejos et al. 2021b] (which was in turn continuing work from [Bosa et al. 2019; Matui and Sato 2014a; Sato et al. 2015]). The most novel technical achievement therein can be identified as the *tracial local-to-global principle* for C^* -algebras satisfying the so-called uniform property Γ , which is a weaker assumption than Jiang–Su stability. Said principle concerns the behavior of elements in a given C^* -algebra A with respect to the 2-seminorm $\|\cdot\|_{2,\tau}$ induced by

individual tracial states τ on A on the one hand, and the behavior with respect to the uniform tracial 2-norm $\|\cdot\|_u = \sup_{\tau} \|\cdot\|_{2,\tau}$ on the other hand. While the latter is often of interest in the deeper structure and classification theory for C*-algebras, the former can be understood by studying the tracial von Neumann algebra $\pi_{\tau}(A)''$ arising as the weak closure of A under the GNS representation associated to an individual trace τ . In a nutshell, the tracial local-to-global principle asserts that any suitable behavior that can be observed one trace at a time can also be observed uniformly, which often allows one to *transfer*, so to speak, phenomena from von Neumann algebras to the C*-algebraic context. This became in turn the main technical driving force behind the main results in [Castillejos et al. 2021b; 2022], which can be summarized by saying that the Toms–Winter conjecture holds for all simple nuclear C*-algebras having the uniform property Gamma. Whether the latter property automatically holds for simple nuclear nonelementary C*-algebras is presently unknown but is of high interest as it currently represents the main obstacle towards a full solution to the Toms–Winter conjecture.

When we turn our attention to C*-dynamics instead of C*-algebras, we can (and should) study analogous well-behavedness properties as for C*-algebras, one important example of which is equivariant Jiang–Su stability. An action $\alpha : G \curvearrowright A$ on a C*-algebra is called (equivariantly) Jiang–Su stable or \mathcal{Z} -stable, if α is cocycle conjugate to $\alpha \otimes \text{id}_{\mathcal{Z}} : G \curvearrowright A \otimes \mathcal{Z}$. Assuming G is amenable, it is presently open if this happens automatically when A is simple nuclear and \mathcal{Z} -stable; see [Szabó 2021b, Conjecture A]. We note that the analogous question for nonamenable groups is known to have a negative answer [Gardella and Lupini 2021; Jones 1983], although recent insights as in [Gardella et al. 2024; Suzuki 2021] leave some hope for the class of amenable actions of nonamenable groups, which we shall not investigate in this article. If one stresses the point again that \mathcal{Z} essentially looks like an infinite-dimensional version of \mathbb{C} , it should not come as a surprise that we can only expect a classification of G -actions on classifiable C*-algebras by reasonable invariants if they are equivariantly \mathcal{Z} -stable. The existing work in this direction seems to indicate that equivariant Jiang–Su stability may indeed hold automatically whenever one can reasonably expect it [Gardella et al. 2022; Matui and Sato 2012; 2014b; Sato 2019; Szabó 2018a; Wouters 2023]. The other possible line of investigation, namely further structural consequences of equivariant Jiang–Su stability for group actions, was initiated in [Gardella et al. 2022] as a direct adaption of techniques in [Castillejos et al. 2021b], albeit under rather restrictive assumptions on the actions. We recall one of the key concepts from both said paper and the present work but restrict ourselves in this introduction to the case of unital simple C*-algebras for convenience, despite actually investigating the concept in broader generality.¹

Definition A. Let G be a countable discrete group. Let A be a separable unital simple C*-algebra such that all 2-quasitraces on A are traces and $T(A) \neq \emptyset$. Given a free ultrafilter ω on \mathbb{N} , form the uniform tracial ultrapower A^{ω} of A . An action $\alpha : G \curvearrowright A$ is said to have equivariant uniform property Gamma if, for any $k \geq 2$, there exist pairwise orthogonal projections $p_1, \dots, p_k \in (A^{\omega} \cap A')^{\alpha^{\omega}}$ such that

$$\tau(ap_j) = \frac{1}{k}\tau(a) \quad \text{for all } j = 1, \dots, k, a \in A, \tau \in T_{\omega}(A).$$

¹The reader might consult Definitions 1.7, 1.8, and 2.1 and compare with [Gardella et al. 2022, Definition 3.1].

One can observe easily (Remark 2.2) that equivariant uniform property Gamma is always implied by equivariant Jiang–Su stability. This is important to note because it means that all nontrivial consequences of this property will also hold for Jiang–Su stable actions, and may in fact turn out to be useful for subsequent applications. The most important technical consequence relevant to the present work is the tracial local-to-global principle for C^* -dynamical systems over amenable groups. An ad hoc version of this principle appeared in [Gardella et al. 2022, Lemma 4.5] but is only applicable (see [Gardella et al. 2022, Remark 2.2]) for actions that induce an action on tracial states with finite orbits of uniformly bounded size. This assumption appeared not only as a prerequisite for the theory in [Gardella et al. 2022] but is implicitly crucial for the usefulness of the conclusion of this ad hoc principle, which only involves tracial states that are fixed by the action.² In this article we aim to remove any assumptions about how actions $G \curvearrowright A$ are allowed to act on the traces of A , as well as strengthen the conclusion of the tracial local-to-global principle compared to [Gardella et al. 2022], in such a way as to directly generalize and strengthen the known principle for C^* -algebras [Castillejos et al. 2021b, Lemma 4.1]. In addition to formulating our result in the language of $*$ -polynomials as all prior papers did, we would also like to promote the following (formally equivalent) formulation of the tracial local-to-global principle for C^* -dynamics, which becomes our main technical result. We not only restrict ourselves for the moment to actions on unital simple nuclear C^* -algebras (similarly to before) for convenience but give a slightly weaker version here that is easier to state. We treat a stronger version of the statement in broader generality in the main body of the paper; see Theorems 4.2 and 4.6.

Theorem B. *Let G be a countable amenable group and A a separable unital simple nuclear C^* -algebra. Let $\alpha : G \curvearrowright A$ be an action with equivariant uniform property Gamma. Let $\delta : G \curvearrowright D$ be an action on a separable C^* -algebra and $B \subseteq D$ a δ -invariant C^* -subalgebra. Suppose that $\varphi : (B, \delta) \rightarrow (A^\omega, \alpha^\omega)$ is an equivariant $*$ -homomorphism. Then φ extends to an equivariant $*$ -homomorphism $\bar{\varphi} : (D, \delta) \rightarrow (A^\omega, \alpha^\omega)$ if and only if, for every trace $\tau \in \overline{T_\omega(A)}^{\omega^*}$, there exists a $*$ -homomorphism $\varphi^\tau : (D, \delta) \rightarrow (\pi_\tau^{\alpha^\omega}(A^\omega)'', \alpha^\omega)$ with $\varphi^\tau|_B = \pi_\tau^{\alpha^\omega} \circ \varphi$.³ Here π_τ denotes the GNS representation associated to the trace τ , and $\pi_\tau^{\alpha^\omega}$ denotes the direct sum representation $\bigoplus_{g \in G} \pi_\tau \circ \alpha_{g^{-1}}^\omega$.*

Not unlike previous approaches, the proof of this main result factors through a kind of dynamical version of CPoU (the existence of so-called *complemented partitions of unity*) that we establish along the way; see Lemma 3.2. There are two things to note about this, however. Firstly, the present version of dynamical CPoU does not generally match the property suggested for this purpose in [Gardella et al. 2022], and based on our work we are in fact uncertain whether that property can be expected to hold even under the validity of the above theorem. Secondly, we would like to propose a slight perspective shift by viewing the above local-to-global principle as the primary conceptual property to be studied and exploited instead of the dynamical CPoU, which we feel — especially compared to previous iterations — to be rather unwieldy by itself due to its elaborate technical nature.

²This uses that if $\alpha : G \curvearrowright A$ is assumed to induce an action $G \curvearrowright T(A)$ with finite orbits of uniformly bounded cardinality $M > 0$, then one has $\|\cdot\|_{2,u} \leq M \|\cdot\|_{2,T(A)^\alpha}$ as norms on A .

³In actuality one may even allow φ^τ to have range in the tracial ultrapower of this von Neumann algebra, but this requires more cumbersome notation to state rigorously.

In the main body of the paper, we actually prove a stronger version of [Theorem B](#) for a class of actions on much more general C*-algebras. We would like to comment that the starting point of our theory presumes that the underlying C*-algebra satisfies a kind of *weak CPoU*, namely the one shown to hold for nuclear C*-algebras in [\[Castillejos et al. 2021b, Lemma 3.6\]](#). Fortunately, a result in the recent preprint [\[Carrión et al. 2023a\]](#) implies that this kind of weak CPoU in fact holds automatically for all C*-algebras with compact tracial state space, which we can use to our advantage.

As for the rest of the paper, we apply [Theorem B](#) (or rather [Theorems 4.2 and 4.6](#)) to gain insight on equivariant Jiang–Su stability. A famous argument due to Matui and Sato [\[2012\]](#) and the main result of [\[Szabó 2021b\]](#) allows us to argue (as explained in [Section 5](#)) that an action α as above is equivariantly Jiang–Su stable if and only if $A \cong A \otimes \mathcal{Z}$ and α is uniformly McDuff, i.e., there exist unital *-homomorphisms

$$M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega} \quad \text{for all } n \in \mathbb{N}.$$

Once we note that the latter property is known to hold one trace at a time as a consequence of Ocneanu’s theorem [\[1985\]](#) (in the generality we need it, this is imported from [\[Szabó and Wouters 2024\]](#)), the above result can be applied to the α -equivariant inclusion

$$1_n \otimes \text{id}_A : A \rightarrow M_n(A)$$

to deduce the following consequence. As before, we note that we prove this result in greater generality than stated here; see [Theorem 5.7](#).

Corollary C (cf. [\[Gardella et al. 2022, Theorem 7.6\]](#)). *Let $\alpha : G \curvearrowright A$ be an action of a countable amenable group on a separable unital simple nuclear \mathcal{Z} -stable C*-algebra. Then α has equivariant uniform property Gamma if and only if α is equivariantly Jiang–Su stable.*

We expect the main result of this article to have an impact on subsequent applications of equivariant uniform property Gamma or equivariant Jiang–Su stability, in particular in the context of classifying actions on tracial C*-algebras.

As far as potential further research is concerned, let us point out that, for group actions $\alpha : G \curvearrowright A$ that are assumed to be “sufficiently free”,⁴ the theory pursued in this article can be seen as an instance where one studies uniform property Gamma for the inclusion of C*-algebras $A \subseteq A \rtimes_{\alpha,r} G$ in such a way as to strengthen uniform property Gamma for A . It is a tantalizing issue to determine a common framework encompassing all applications of interest regarding uniform property Gamma for more general inclusions of C*-algebras. For instance, it has been hypothesized in past work [\[Kerr and Szabó 2020, Remark 9.6\]](#) that, for a free minimal action $G \curvearrowright X$ of an amenable group on a compact metric space, some desirable dynamical properties ought to follow from a different kind of uniform property Gamma for the inclusion $\mathcal{C}(X) \subseteq \mathcal{C}(X) \rtimes G$, namely the one that strengthens uniform property Gamma for the crossed product; see also [\[Liao and Tikuisis 2022\]](#).

⁴A priori, this may have several different interpretations.

1. Preliminaries

Notation 1.1. Throughout this paper, we will use the following notation and conventions unless specified otherwise:

- By default, ω denotes some free ultrafilter on \mathbb{N} . At times it can make it easier to state a claim using two free ultrafilters, in which case we denote a second one by κ .
- If F is a finite subset inside another set M , we often denote this by $F \Subset M$.
- \mathbb{K} denotes the compact operators on the Hilbert space $\ell^2(\mathbb{N})$.
- Let A be a C^* -algebra. We denote its positive elements by A_+ and its minimal unitization by \tilde{A} . We will also make use of its Pedersen ideal, denoted by $\mathcal{P}(A)$. We assume the reader is familiar with the basic properties of this object. Given a positive element $a \in A$ and $\varepsilon > 0$, we denote by $(a - \varepsilon)_+$ the positive part of the self-adjoint element $a - \varepsilon 1_{\tilde{A}}$.
- The topological cone of lower semicontinuous traces on A_+ will be denoted by $\tilde{T}(A)$; cf. [Elliott et al. 2011]. We call such a trace τ on A *trivial* if it is $\{0, \infty\}$ -valued. It is well known that trivial traces are in one-to-one correspondence with the ideal lattice of A by mapping a trivial trace τ to the linear span of $\tau^{-1}(0)$. The set of nontrivial lower semicontinuous traces on A_+ will be denoted by $T^+(A)$ and the set of tracial states will be denoted by $T(A)$. In this paper, we say that a compact subset $K \subset T^+(A)$ is a *compact generator* for $T^+(A)$ if $\mathbb{R}^{>0}K = T^+(A)$.⁵
- In addition, we denote by $Q\tilde{T}_2(A)$ the set of lower semicontinuous 2-quasitraces (see [Blanchard and Kirchberg 2004, Definition 2.22]) on A , which contains $\tilde{T}(A)$. We usually only mention them to assume in appropriate contexts that there are no genuine quasitraces, i.e., $Q\tilde{T}_2(A) = \tilde{T}(A)$.

We recall the following existence theorem for traces. This follows from a combination of the work of Blackadar and Cuntz [1982, Theorem 1.5] and Haagerup [2014]; see also [Blanchard and Kirchberg 2004, Remark 2.29 (i)].

Theorem 1.2. *Let A be a simple, exact C^* -algebra such that $A \otimes \mathbb{K}$ contains no infinite projections. Then $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$.*

In particular, this implies that each stably finite, simple, separable, nuclear C^* -algebra admits a nontrivial trace.

Definition 1.3 [Kirchberg 2006, Definition 1.1; Kirchberg and Rørdam 2014, Definition 4.3]. Let A be a C^* -algebra with an action $\alpha : G \curvearrowright A$ of a discrete group.

- (1) The *ultrapower* of A is defined as

$$A_\omega := \ell^\infty(A) / \{(a_n)_{n \in \mathbb{N}} \in \ell^\infty(A) : \lim_{n \rightarrow \omega} \|a_n\| = 0\}.$$

- (2) Pointwise application of α on representing sequences induces an action on the ultrapower, which we will denote by $\alpha_\omega : G \curvearrowright A_\omega$.

⁵In case A is simple, an example of such a compact generator is given by $\{\tau \in T^+(A) \mid \tau(a) = 1\}$ for some $a \in \mathcal{P}(A)_+ \setminus \{0\}$.

- (3) There is a natural inclusion $A \subset A_\omega$ by identifying an element of A with its constant sequence. Define

$$A_\omega \cap A' := \{x \in A_\omega \mid [x, A] = 0\} \quad \text{and} \quad A_\omega \cap A^\perp := \{x \in A_\omega \mid xA = Ax = 0\}.$$

The quotient

$$F_\omega(A) := (A_\omega \cap A') / (A_\omega \cap A^\perp)$$

is called the (*corrected*) *central sequence algebra*. If A is σ -unital, then $F_\omega(A)$ is unital, where the unit is represented by a sequential approximate unit $(e_n)_{n \in \mathbb{N}}$.

- (4) Since A is α_ω -invariant, so are $A_\omega \cap A'$ and $A_\omega \cap A^\perp$. Thus, α_ω induces an action on $F_\omega(A)$, which we will denote by $\tilde{\alpha}_\omega : G \curvearrowright F_\omega(A)$.

Definition 1.4. Let A be a C*-algebra. A sequence of tracial states $(\tau_n)_{n \in \mathbb{N}}$ on A defines a trace on A_ω via

$$[(a_n)_{n \in \mathbb{N}}] \mapsto \lim_{n \rightarrow \omega} \tau_n(a_n).$$

A trace of this form is called a *limit trace*. The set of all limit traces on A_ω will be denoted by $T_\omega(A)$. More generally, following [Szabó 2021b, Definition 2.1], a sequence $(\tau_n)_{n \in \mathbb{N}}$ in $\tilde{T}(A)$ defines a lower semicontinuous trace $\tau : \ell^\infty(A)_+ \rightarrow [0, \infty]$ by

$$\tau((a_n)_{n \in \mathbb{N}}) = \sup_{\varepsilon > 0} \lim_{n \rightarrow \omega} \tau_n((a_n - \varepsilon)_+).$$

This trace is the lower semicontinuous regularization of the trace given by $\lim_{n \rightarrow \omega} \tau_n(a_n)$; see [Elliott et al. 2011, Lemma 3.1]. This regularization ensures that $\tau((a_n)_{n \in \mathbb{N}}) = 0$ if $\lim_{n \rightarrow \omega} \|a_n\| = 0$, so τ also induces a lower semicontinuous trace on A_ω . A trace of this form on A_ω is called a *generalized limit trace*. The set of all generalized limit traces is denoted by $\tilde{T}_\omega(A)$.

For the next part, assume A is separable. For any $a \in A_+$ and $\tau \in \tilde{T}_\omega(A)$, we can define a trace

$$\tau_a : (A_\omega \cap A')_+ \rightarrow [0, \infty], \quad x \mapsto \tau(ax).$$

We have that $\tau_a(x) \leq \|x\| \tau(a)$, so this trace is bounded whenever $\tau(a) < \infty$. Note that this trace also induces a trace on $F_\omega(A)$, which by abuse of notation will also be denoted by τ_a . Clearly this yields a tracial state under the assumption $\tau(a) = 1$. Let us say that a given tracial state τ on $F_\omega(A)$ is a *canonical trace* if it belongs to the weak- $*$ -closed convex hull of $\{\tau_a \mid \tau \in \tilde{T}_\omega(A), a \in A_+, \tau(a) = 1\}$.

Remark 1.5. We point out that it is not necessary to consider generalized limit traces in an important subcase that often occurs in the literature. Namely, assume A is a separable simple C*-algebra with $\emptyset \neq T^+(A) = \mathbb{R}^{>0}T(A)$ such that $T(A)$ is compact.⁶ Then it follows by [Castillejos and Evington 2021, Proposition 2.3] that every generalized limit trace $\tau \in \tilde{T}_\omega(A)$ that is finite on some nonzero positive element of A is a multiple of an ordinary limit trace.

Next we recall how various versions of tracial ultrapowers are defined.

⁶For instance, this is automatic when A is separable, simple, nuclear, unital, and stably finite.

Definition 1.6 [Ando and Haagerup 2014, Propositions 3.1 and 3.2]. Suppose \mathcal{M} is a finite von Neumann algebra with faithful normal tracial state τ . Then the *tracial von Neumann algebra ultrapower* is defined as

$$\mathcal{M}^\omega := \ell^\infty(\mathcal{M}) / \{(x_n)_{n \in \mathbb{N}} \in \ell^\infty(\mathcal{M}) \mid \lim_{n \rightarrow \infty} \|x_n\|_{2,\tau} = 0\}. \quad (1-1)$$

This is again a von Neumann algebra with a faithful normal tracial state τ^ω that is defined on representative sequences by $\tau^\omega((x_n)_{n \in \mathbb{N}}) = \lim_{n \rightarrow \infty} \tau(x_n)$.

The notation used for the tracial von Neumann algebra ultrapower is the same as for the uniform tracial ultrapower of a suitable C^* -algebra as defined below. It will be clear from context which of the two notions we use. In the special case that A is a C^* -algebra with unique tracial state τ and no unbounded traces, the uniform tracial ultrapower A^ω is naturally isomorphic, by Kaplansky's density theorem, to the von Neumann tracial ultrapower $(\pi_\tau(A)')^\omega$, where π_τ denotes the GNS representation associated to τ . Note that on a tracial von Neumann algebra (\mathcal{M}, τ) , the topology induced by the $\|\cdot\|_{2,\tau}$ -norm agrees with the $*$ -strong operator topology on bounded subsets. So equivalently, in (1-1) one can quotient out by the sequences that converge to 0 in the $*$ -strong operator topology, which makes the construction equivalent to the Ocneanu ultrapower; cf. [Ando and Haagerup 2014].

Definition 1.7. Let A be a C^* -algebra. Given a constant $p \geq 1$ and $\tau \in T(A)$, we define a seminorm $\|\cdot\|_{p,\tau}$ on A by

$$\|a\|_{p,\tau} = \tau(|a|^p)^{1/p}, \quad a \in A.$$

We will in particular appeal to the cases $p = 1$ or $p = 2$ subsequently. For a nonempty set $X \subset T(A)$, we define a seminorm $\|\cdot\|_{2,X}$ on A by

$$\|a\|_{2,X} := \sup_{\tau \in X} \|a\|_{2,\tau}$$

for all $a \in A$. The seminorm $\|\cdot\|_{2,T(A)}$ is also denoted by $\|\cdot\|_{2,u}$. This is a norm if and only if, for all nonzero $a \in A$, there exists some $\tau \in T(A)$ such that $\tau(a^*a) > 0$, which is in particular the case when A is simple with $T(A)$ nonempty.

We note that in the construction below, we deviate from other sources by making a very explicit distinction in terminology between C^* -algebras that do or do not admit nontrivial unbounded traces.

Definition 1.8 (cf. [Castillejos et al. 2021b, Section 1.3]⁷). Let A be a C^* -algebra with $T(A) \neq \emptyset$. Then the *trace-kernel ideal (with respect to bounded traces)* inside A_ω is defined by

$$J_A^b := \{(a_n)_{n \in \mathbb{N}} \in A_\omega \mid \lim_{n \rightarrow \infty} \|a_n\|_{2,T(A)} = 0\}.$$

The *uniform bounded tracial ultrapower* is defined as the quotient

$$A^{\omega,b} := A_\omega / J_A^b.$$

Whenever $\|\cdot\|_{2,T(A)}$ defines a norm on A , there also exists a canonical embedding of A into $A^{\omega,b}$. Then $A^{\omega,b} \cap A'$ is called the *uniform bounded tracial central sequence algebra*. Whenever we have an action

⁷The cited source assumes separability, but we generalize the definition beyond that case.

$\alpha : G \curvearrowright A$ of a discrete group, the ideal J_A^b is α_ω -invariant. Hence, there is an induced action on the uniform bounded tracial ultrapower, which we will denote by $\alpha^\omega : G \curvearrowright A^{\omega,b}$.

Clearly, every limit trace vanishes on J_A^b and hence also induces a tracial state on $A^{\omega,b}$. We will also use $T_\omega(A)$ to denote the collection of limit traces on $A^{\omega,b}$. Note that

$$J_A^b = \{x \in A_\omega : \|x\|_{2, T_\omega(A)} = 0\},$$

so in particular $\|\cdot\|_{2, T_\omega(A)}$ defines a norm on $A^{\omega,b}$.

Finally, if we assume A is a simple C*-algebra such that $QT_2(A) = \tilde{T}(A)$ and $\emptyset \neq T^+(A) = \mathbb{R}^{>0} \cdot T(A)$ with $T(A)$ compact, then we simply call $J_A = J_A^b$ the trace-kernel ideal, $A^\omega = A^{\omega,b}$ the uniform tracial ultrapower, and $A^\omega \cap A'$ the uniform tracial central sequence algebra.

Remark 1.9. Our choice to add the extra “bounded” in the terminology above and the extra letter “b” in the notation, which is usually not included in other sources such as the ones we cite, is deliberate and has the purpose to not overuse the word “uniform”, in particular in cases where it becomes rather misleading. This is most apparent for nonsimple C*-algebras; if B is any unital simple C*-algebra with $T(B) \neq \emptyset$, then the above construction applied to $A = B \oplus \mathbb{K}$ yields $A^{\omega,b} = B^\omega$ by virtue of the fact that the canonical trace on \mathbb{K} is unbounded. Since one of the two tracial direct summands is entirely forgotten in this construction, this object seems unfit to be called “uniform tracial”. However, even the case of simple C*-algebras is enough to illustrate why one should not equate $A^{\omega,b}$ with the object capturing all “uniform tracial” data. Namely, the range result [Gong and Lin 2022] combined with a little playing around with invariants allows one to see that, given any metrizable Choquet simplex S with $\partial_e S$ admitting some isolated point, there exists a (nonunital) classifiable C*-algebra A such that $T^+(A)$ has a Choquet base affinely homeomorphic to S , yet A has a unique tracial state τ . In this scenario, we have $A^{\omega,b} \cong (\pi_\tau(A))^\omega \cong \mathcal{R}^\omega$ as a consequence of Connes’ theorem. So despite A having a rich tracial structure, the only trace captured by this construction is τ , which compels us to not apply the word “uniform” or the notation “ A^ω ” to such an example.

Note that the phenomenon discussed here is also what motivated us to subsequently revise the definition of (equivariant) uniform property Gamma in the spirit of [Castillejos and Evington 2021], as well as introduce an auxiliary version of it that explicitly only takes into account tracial states, even when the surrounding C*-algebra may have other unbounded traces.

Remark 1.10. Let A be a σ -unital C*-algebra with $T(A) \neq \emptyset$. By [Castillejos et al. 2021b, Proposition 1.11] the uniform bounded tracial ultrapower $A^{\omega,b}$ is unital if and only if $T(A)$ is compact.⁸ Moreover, [Castillejos et al. 2021b, Lemma 1.10] shows that, in that case, the natural map $A_\omega \cap A' \rightarrow A^{\omega,b} \cap A'$ factors through $F_\omega(A)$. In case A is separable, this natural map is surjective by a combination of Propositions 4.5 (iii) and 4.6 in [Kirchberg and Rørdam 2014] (the unitality hypothesis in the second cited proposition is not needed, as it suffices to take a unit in the minimal unitization for the proof).

As we have argued above, there are some issues if one is trying to define the object A^ω for a C*-algebra A that may possess many unbounded traces. In fact, trying to find a viable general definition that has the

⁸The cited statement assumes separability of A , but a closer look at the proof shows that σ -unitality is sufficient.

same level of utility as in the case of unital C^* -algebras has eluded a number of researchers for years. By introducing the next few definitions and observations, however, we wish to promote the viewpoint that there is a rather natural way to define the object $A^\omega \cap A'$ for any separable C^* -algebra A , even if we do not know at present how to properly define the object A^ω itself. Since we are unsure of the viability of this definition when A admits genuine quasitraces, we wish to be cautious and shall only define the concepts below under the assumption that A does not admit them.⁹

Definition 1.11. Let A be a separable C^* -algebra with $QT_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$. The *trace-kernel ideal* \mathcal{J}_A inside $F_\omega(A)$ is defined as the set of elements $x \in F_\omega(A)$ such that, for every generalized limit trace $\tau \in \tilde{T}_\omega(A)$ and $a \in A_+$ with $0 < \tau(a) < \infty$, we have $\tau_a(x^*x) = 0$. With some abuse of notation, we denote the quotient by

$$A^\omega \cap A' = F_\omega(A) / \mathcal{J}_A. \tag{1-2}$$

It is clear from construction that a canonical trace on $F_\omega(A)$ vanishes on \mathcal{J}_A , so it descends to a tracial state on $A^\omega \cap A'$. As before, we call a given tracial state on $A^\omega \cap A'$ a *canonical trace* if it is induced by a canonical trace on $F_\omega(A)$, or equivalently if it belongs to the weak- $*$ -closed convex hull of the tracial states τ_a on $A^\omega \cap A'$, where $\tau \in \tilde{T}_\omega(A)$ and $a \in A_+$ with $\tau(a) = 1$.

If $\alpha : G \curvearrowright A$ is an action of a discrete group with induced action $\tilde{\alpha}_\omega : G \curvearrowright F_\omega(A)$, then clearly \mathcal{J}_A is $\tilde{\alpha}_\omega$ -invariant, so we obtain an induced action $\alpha^\omega : G \curvearrowright A^\omega \cap A'$.

Remark 1.12. In the case that A is simple, and $QT_2(A) = \tilde{T}(A)$ and $\emptyset \neq T^+(A) = \mathbb{R}^{>0}T(A)$ with $T(A)$ compact, it follows from Remarks 1.5 and 1.10 that $F_\omega(A) / \mathcal{J}_A = A^{\omega,b} \cap A'$, so the notation $A^\omega \cap A'$ is consistent with the last part of Definition 1.8.

Remark 1.13 (see remark after [Kirchberg and Rørdam 2014, Definition 4.3]). Let $p \geq 1$ be any constant. Given any element x in a C^* -algebra B with a tracial state θ , one has the inequalities

$$\|x\|_{1,\theta} \leq \|x\|_{p,\theta} \leq \|x\|_{1,\theta}^{1/p} \|x\|^{1-1/p}.$$

This implies that an element in either Definition 1.8 or 1.11 belongs to the trace-kernel ideal if and only if its tracial p -norms vanish with respect to the appropriately chosen (limit) traces. We will frequently use this without further mention for $p = 1$.

Remark 1.14. One of Kirchberg’s initial observations about $F_\omega(A)$, which attests to the naturality of its construction, is that it is a stable invariant. We are about to argue that the same applies to the construction $A \mapsto A^\omega \cap A'$. For this purpose, let $\{e_{k,\ell} \mid k, \ell \geq 1\}$ be a set of matrix units generating \mathbb{K} , and let $1_n \in \mathbb{K}$ be the increasing approximate unit given by $1_n = \sum_{j=1}^n e_{j,j}$. We recall (see [Kirchberg 2006, Proposition 1.9, Corollary 1.10]) that there is a canonical isomorphism $\theta : F_\omega(A) \rightarrow F_\omega(A \otimes \mathbb{K})$ defined as follows: given an element $x \in F_\omega(A)$ represented by a central sequence $(x_n)_{n \in \mathbb{N}}$ in A , it is sent to the element $\theta(x)$ represented by the central sequence $(x_n \otimes 1_n)_{n \in \mathbb{N}}$.

⁹At the same time, we note that the concepts make sense formally anyway, and none of the subsequent arguments hinge on the assumption that A does not admit genuine quasitraces.

Proposition 1.15. *Let A be a separable C*-algebra with $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$. Then the canonical isomorphism $F_\omega(A) \cong F_\omega(A \otimes \mathbb{K})$ preserves the canonical traces on both sides. Consequently, it descends to a canonical isomorphism*

$$A^\omega \cap A' \cong (A \otimes \mathbb{K})^\omega \cap (A \otimes \mathbb{K})'.$$

Proof. As we set up before the proposition, we denote the canonical isomorphism by θ . It is clear that it induces an affine homeomorphism between all tracial states on $F_\omega(A)$ and on $F_\omega(A \otimes \mathbb{K})$ via $\tau \mapsto \tau \circ \theta^{-1}$. The claim amounts to showing that the image of the canonical traces on the left is equal to the canonical traces on the right.

Let Tr be the unique lower semicontinuous trace on \mathbb{K} with $\text{Tr}(e_{1,1}) = 1$. We keep in mind that the assignment $\tilde{T}(A) \rightarrow \tilde{T}(A \otimes \mathbb{K})$ given by $\tau \mapsto \tau \otimes \text{Tr}$ is an affine homeomorphism. Given a generalized limit trace $\tau \in \tilde{T}_\omega(A)$ induced by a sequence $(\tau_n)_{n \in \mathbb{N}}$ in $\tilde{T}(A)$, let us denote by $\tau^s \in \tilde{T}_\omega(A \otimes \mathbb{K})$ the generalized limit trace induced by the sequence $(\tau_n \otimes \text{Tr})_{n \in \mathbb{N}}$ in $\tilde{T}(A \otimes \mathbb{K})$. Clearly the assignment $\tau \mapsto \tau^s$ is also a bijection between generalized limit traces. Let such a generalized limit trace τ be given on A_ω . Given $a \in (A \otimes \mathbb{K})_+$, we can write $a = \sum_{k,\ell=1}^\infty a_{k,\ell} \otimes e_{k,\ell}$ for uniquely determined elements $a_{k,\ell} \in A$. It then follows from [Castillejos and Evington 2021, Proposition 2.9] that we have a norm-convergent sum expression

$$\tau_a^s \circ \theta = \sum_{\ell=1}^\infty \tau_{a_{\ell,\ell}}. \tag{1-3}$$

Applied to $a = b \otimes e_{1,1}$ for some $b \in A_+$ with $\tau(b) = 1$, this gives $\tau_b \circ \theta^{-1} = \tau_a^s$. From this we can infer that canonical traces are mapped to canonical traces. The general expression (1-3) applied to $a \in (A \otimes \mathbb{K})_+$ with $\tau^s(a) = 1$ shows that we have a bijection. □

We give two more technical lemmas that will be useful later on.

Lemma 1.16. *Let A be a C*-algebra with positive element $a \in A_+$. Let $(\varepsilon_n)_{n \in \mathbb{N}}$ be a sequence of positive constants such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$, and let $(b_n)_{n \in \mathbb{N}}$ be a sequence of positive elements such that $\|b_n - a\| < \varepsilon_n$. For each $\tau \in \tilde{T}_\omega(A)$ and $c \in F_\omega(A)$, one has*

$$\lim_{n \rightarrow \infty} \tau_{(b_n - \varepsilon_n)_+}(c) = \tau_a(c).$$

Proof. For each $n \in \mathbb{N}$, there exists a contraction $d_n \in A$ such that $(b_n - \varepsilon_n)_+ = d_n a d_n^*$ by [Kirchberg and Rørdam 2002, Lemma 2.2]. This implies that $\tau_{(b_n - \varepsilon_n)_+}(c) \leq \tau_a(c)$. Since the sequence $(b_n - \varepsilon_n)_+$ converges to a and τ is lower semicontinuous, this leads to the desired result. □

Lemma 1.17. *Let A be a simple C*-algebra with $a \in \mathcal{P}(A)_+ \setminus \{0\}$, and let K be a compact generator for $T^+(A) \neq \emptyset$. Take a generalized limit trace $\tau \in \tilde{T}_\omega(A)$ such that $0 < \tau(a) < \infty$. Then there exists a sequence $(\theta_n)_{n \in \mathbb{N}}$ in K such that the associated generalized limit trace θ on A_ω is a scalar multiple of τ and such that, for each sequence $(b_n)_{n \in \mathbb{N}}$ representing an element of A_ω , we have that*

$$\theta((ab_n)_{n \in \mathbb{N}}) = \lim_{n \rightarrow \omega} \theta_n(ab_n). \tag{1-4}$$

Proof. The fact that there exists a sequence $(\theta_n)_{n \in \mathbb{N}}$ in K such that the associated generalized limit trace is a multiple of τ follows directly from [Szabó 2021b, Lemma 2.10 and Remark 2.11]. Let $B := \overline{a^{1/2} A a^{1/2}}$ denote the hereditary subalgebra generated by $a^{1/2}$. Then $B \subseteq \mathcal{P}(A)$; see for example [Pedersen 1979, Proposition 5.6.2]. Since K is compact, we claim that $\sup_{\sigma \in K} \|\sigma|_B\| < \infty$. Suppose that this would not be the case; then, for all $n \in \mathbb{N}$, we could find a $\sigma_n \in K$ and a positive contraction $d_n \in B$ such that $\sigma_n(d_n) \geq n2^n$. Consider $d := \sum_{n=1}^\infty 2^{-n} d_n \in B$ and $\sigma = \lim_{n \rightarrow \omega} \sigma_n \in K$ (using the compactness of K). Then, for each $n \in \mathbb{N}$, we would get

$$\sigma(d) = \lim_{n \rightarrow \omega} \sigma_n(d) \geq \lim_{n \rightarrow \omega} \sigma_n(2^{-n} d_n) = \infty,$$

but this is a contradiction, since d belongs to the Pedersen ideal. As a consequence we get that, when restricted to the hereditary subalgebra $\overline{a^{1/2} \ell^\infty(A) a^{1/2}} \subseteq \ell^\infty(B)$, the trace formed by $\lim_{n \rightarrow \omega} \theta_n$ is already bounded and hence continuous, so formula (1-4) holds. \square

The following proposition is a useful lifting property in various contexts. The proof relies on the concept of G - σ -ideals; see [Szabó 2018c, Definition 4.1]. Let $\alpha : G \curvearrowright A$ and $\beta : G \curvearrowright B$ be actions on C^* -algebras. As in [Kirchberg 2006], we call an equivariant surjective $*$ -homomorphism $\pi : (A, \alpha) \rightarrow (B, \beta)$ strongly locally semisplit, if for every separable β -invariant C^* -subalgebra $D \subseteq B$, there exists an equivariant c.p.c. order-zero map $\phi : (D, \beta) \rightarrow (A, \alpha)$ such that $\pi \circ \phi = \text{id}_D$.

Proposition 1.18. *Let A be a separable simple C^* -algebra with $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$. Let $\alpha : G \curvearrowright A$ be an action of a countable discrete group. Then the quotient map*

$$(F_\omega(A), \tilde{\alpha}_\omega) \rightarrow (A^\omega \cap A', \alpha^\omega)$$

is strongly locally semisplit.

Proof. By [Szabó 2018c, Proposition 4.5 (ii)], it suffices to prove that $\mathcal{J}_A \subset F_\omega(A)$ is a G - σ -ideal. Fix an element $0 \neq a \in \mathcal{P}(A)_+$ and a compact generator $K \subset T^+(A)$.¹⁰ By [Szabó 2021b, Proposition 2.4] and Lemma 1.16, we can conclude that \mathcal{J}_A coincides with the ideal of those elements $x \in F_\omega(A)$ such that $\tau_a(x^*x) = 0$ for all $\tau \in \tilde{T}_\omega(A)$ induced by any sequence $\tau_n \in K$. Since a belongs to the Pedersen ideal and K is compact, this further implies that an element $x \in F_\omega(A)$ represented by a sequence $(x_n)_{n \in \mathbb{N}}$ in A belongs to \mathcal{J}_A precisely when $\lim_{n \rightarrow \omega} \max_{\tau \in K} \tau(a^{1/2} x_n^* x_n a^{1/2}) = 0$.

We proceed to show that \mathcal{J}_A is a G - σ -ideal. Let $D \subset F_\omega(A)$ be a separable $\tilde{\alpha}_\omega$ -invariant C^* -subalgebra. Let $(d_{k,n})_{n,k \in \mathbb{N}}$ and $(c_{k,n})_{n,k \in \mathbb{N}}$ be two bounded double sequences in A such that, for each $k \in \mathbb{N}$, the sequences $(d_{k,n})_{n \in \mathbb{N}}$ and $(c_{k,n})_{n \in \mathbb{N}}$ are approximately central, the set $\{d^{(k)} = [(d_{k,n})_{n \in \mathbb{N}}] \mid k \in \mathbb{N}\}$ defines a dense subset in the unit ball of D , and the set $\{c^{(k)} = [(c_{k,n})_{n \in \mathbb{N}}] \mid k \in \mathbb{N}\}$ defines a dense subset in the unit ball of $D \cap \mathcal{J}_A$. By Kasparov’s lemma [1988, Lemma 1.4], we can find, for any $\varepsilon > 0$, $F \in G$, and $m \in \mathbb{N}$, a positive element $e \in \mathcal{J}_A$ such that

$$\max_{k \leq m} \|[e, d^{(k)}]\| \leq \varepsilon, \quad \max_{k \leq m} \|(1 - e)c^{(k)}\| \leq \varepsilon, \quad \text{and} \quad \max_{g \in F} \|e - \tilde{\alpha}_\omega(e)\| \leq \varepsilon.$$

¹⁰As pointed out in the footnote after defining compact generators in Notation 1.1, this always exists as a consequences of simplicity.

Let $b \in A$ be a strictly positive contraction. If we represent e by an approximately central sequence $(e_n)_{n \in \mathbb{N}}$ of positive contractions in A , then it follows that

$$\max_{k \leq m} \lim_{n \rightarrow \omega} \|[e_n, d_{k,n}]b\| \leq \varepsilon, \quad \max_{k \leq m} \lim_{n \rightarrow \omega} \|(1 - e_n)c_{k,n}b\| \leq \varepsilon,$$

and

$$\max_{g \in F} \lim_{n \rightarrow \omega} \|(e_n - \alpha_g(e_n))b\| \leq \varepsilon, \quad \lim_{n \rightarrow \omega} \max_{\tau \in K} \tau(a^{1/2}e_n a^{1/2}) = 0.$$

Appealing to Kirchberg’s ε -test [Kirchberg and Rørdam 2014, Lemma 3.1], we can find another approximately central sequence $(e_n)_{n \in \mathbb{N}}$ of positive contractions in A satisfying the stronger property

$$\lim_{n \rightarrow \omega} (\|[e_n, d_{k,n}]b\| + \|(1 - e_n)c_{k,n}b\| + \|(e_n - \alpha_g(e_n))b\|) = 0 \quad \text{and} \quad \lim_{n \rightarrow \omega} \max_{\tau \in K} \tau(a^{1/2}e_n a^{1/2}) = 0$$

for all $k \in \mathbb{N}$ and $g \in G$. This means that this sequence represents a positive contraction $e \in (\mathcal{J}_A \cap D')^{\tilde{\alpha}_\omega}$ such that $ec = c$ for all $c \in \mathcal{J}_A \cap D$. This finishes the proof. \square

To end this preliminary section, we prove the following tracial inequality.

Lemma 1.19. *Let B be a C*-algebra with $a, b \in B_+$ and $\tau \in T(B)$. Then*

$$\|a - b\|_{2,\tau}^2 \leq \|a^2 - b^2\|_{1,\tau}.$$

Proof. If we replace B by its weak closure of the GNS representation $\pi_\tau(B)''$, it is enough to show this in the case that B is a von Neumann algebra with faithful normal tracial state τ .

Historically, this was proved by Powers and Størmer [1970, Lemma 4.1] in the case $B = M_n(\mathbb{C})$ for some $n \in \mathbb{N}$. When B is a von Neumann algebra with faithful normal tracial state τ , this follows from applying [Haagerup 1975, Lemma 2.10], which is formulated for the space $L^2(B, \tau)$, to elements in $B \subset L^2(B, \tau)$. For the reader’s convenience we give here a more direct proof using an idea from [Anantharaman and Popa 2014, Theorem 7.3.7].

Given $a, b \in B_+$, let p and q denote the spectral projections of $a - b$ corresponding to $[0, +\infty)$ and $(-\infty, 0)$, respectively. This means that $a - b = (p - q)|a - b|$ and $p \perp q$. First of all, we have

$$\tau((a^2 - b^2)p) - \tau((a - b)^2 p) = \tau(b(a - b)p) + \tau((a - b)bp) = 2\tau(b^{1/2}(a - b)pb^{1/2}) \geq 0$$

since $(a - b)p \geq 0$. So we get

$$\tau((a - b)^2 p) \leq \tau((a^2 - b^2)p), \tag{1-5}$$

and in a similar way we can obtain that

$$\tau((b - a)^2 q) \leq \tau((b^2 - a^2)q). \tag{1-6}$$

Combining (1-5) and (1-6) gives

$$\tau((a - b)^2) = \tau((a - b)^2(p + q)) \leq \tau((a^2 - b^2)(p - q)).$$

Also,

$$\tau((a^2 - b^2)(p - q)) = \tau(p(a^2 - b^2)p) + \tau(q(b^2 - a^2)q) \leq \tau(|a^2 - b^2|(p + q)) \leq \|a^2 - b^2\|_{1,\tau}$$

since $\|p + q\| \leq 1$. This implies the result. \square

2. Equivariant uniform property Gamma

The notion of uniform property Gamma was introduced in [Castillejos et al. 2021b] and further studied in [Castillejos et al. 2022], where it served as a uniform C^* -algebraic version of property Gamma introduced by Murray and von Neumann for II_1 factors [1943]. Recently, a dynamical version of this property was introduced in the separable unital setting in [Gardella et al. 2022] called the *equivariant uniform property Gamma*. Here we revise the definition to account for separable C^* -algebras with possibly unbounded traces, generalizing the concept called “stabilised property Gamma” by Castillejos and Evington [2021, Definition 2.5]. We choose not to adopt that name because one can argue that uniform property Gamma ought to be a stable property in the first place, just like property Gamma is for von Neumann algebras. In light of recent work by Lin [2023] who proposed a more general framework for C^* -algebras that admit genuine quasitraces, we shall state the definition only in the absence of such.

For separable unital simple exact C^* -algebras, the definition below corresponds to the earlier definition given in [Gardella et al. 2022] (see Proposition 2.4 below) but not in the nonsimple case, as demonstrated by C^* -algebras that arise as extensions of unital classifiable C^* -algebras by the compacts (see the type of example mentioned in Remark 1.9, for instance).

Definition 2.1. Let A be a separable C^* -algebra with $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$, and let $\alpha : G \curvearrowright A$ be an action by a countable discrete group. We say that α has *equivariant uniform property Gamma* (or *equivariant property Gamma* for short) if, for all $n \in \mathbb{N}$, there exist pairwise orthogonal projections $p_1, \dots, p_n \in (A^\omega \cap A')^{\alpha^\omega}$ such that, for all $a \in A_+$ and $\tau \in \tilde{T}_\omega(A)$ with $\tau(a) < \infty$,

$$\tau_a(p_i) = \frac{1}{n} \tau(a).$$

Remark 2.2. We can notice immediately from the naturality of the isomorphism in Proposition 1.15 that equivariant uniform property Gamma is preserved under stable cocycle conjugacy. That is, if A and B are C^* -algebras as above and we have actions $\alpha : G \curvearrowright A$ and $\beta : G \curvearrowright B$ such that $\alpha \otimes \text{id}_{\mathbb{K}}$ is cocycle conjugate to $\beta \otimes \text{id}_{\mathbb{K}}$, then α^ω is conjugate to β^ω via a map preserving the canonical traces. In particular, equivariant uniform property Gamma holds for α if and only if it holds for β .

Next, we observe (cf. [Castillejos et al. 2021b, Proposition 2.3]) that, whenever $\alpha : G \curvearrowright A$ is an equivariantly \mathcal{Z} -stable action on a separable C^* -algebra with $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$, it automatically has equivariant property Gamma. Indeed, a cocycle conjugacy between α and $\alpha \otimes \text{id}_{\mathcal{Z}}$ is easily seen to give rise to a unital $*$ -homomorphism

$$\mathcal{Z}^\omega \cap \mathcal{Z}' \rightarrow (A^\omega \cap A')^{\alpha^\omega}.$$

For this purpose one chooses an approximate unit $e_n \in A$ and considers a sequence of maps $\mathcal{Z} \rightarrow A \otimes \mathcal{Z}$, $x \mapsto e_n \otimes x$, composed with such a cocycle conjugacy, which is seen to induce such a homomorphism. It is well known that $\mathcal{Z}^\omega \cap \mathcal{Z}'$ admits unital embeddings of matrix algebras of arbitrary size $n \geq 2$. So if we fix n and define $p_1, \dots, p_n \in (A^\omega \cap A')^{\alpha^\omega}$ as the image of the canonical rank-one projections inside a matrix algebra under the aforementioned $*$ -homomorphism, then they satisfy the necessary requirements for equivariant property Gamma by uniqueness of the trace on the $n \times n$ matrices.

The following is a version of equivariant property Gamma for possibly nonseparable C*-algebras that exclusively takes into account the bounded traces. This agrees with [Gardella et al. 2022, Definition 3.1] for separable unital C*-algebras but not with the general definition of equivariant property Gamma given above.

Definition 2.3. Let A be a σ -unital C*-algebra with $T(A)$ nonempty and compact, and let $\alpha : G \curvearrowright A$ be an action of a countable discrete group. We say that α has *local equivariant property Gamma with respect to bounded traces* if, for all $n \in \mathbb{N}$ and $\|\cdot\|_{2, T_\omega(A)}$ -separable subsets $S \subset A^{\omega, b}$, there exist pairwise orthogonal projections $p_1, \dots, p_n \in (A^{\omega, b})^{\alpha^\omega} \cap S'$ such that $\tau(ap_i) = \tau(a)/n$ for all $a \in S$ and $\tau \in T_\omega(A)$.

In the unital separable simple setting, Definitions 2.1 and 2.3 are equivalent. We prove this fact in a slightly more general setting in the proposition below.

Proposition 2.4. Let A be a simple separable C*-algebra with $QT_2(A) = \tilde{T}(A)$, and such that $T(A) \neq \emptyset$ is compact and $T^+(A) = \mathbb{R}^{>0}T(A)$.¹¹ Then an action $\alpha : G \curvearrowright A$ of a countable discrete group has equivariant property Gamma if and only if α has local equivariant property Gamma with respect to bounded traces.

Proof. The assumptions on A imply that every generalized limit trace on A that is finite on some nonzero positive element of A is a multiple of an ordinary limit trace (see Remark 1.5) and that $A^\omega \cap A' = A^{\omega, b} \cap A'$ (see Remark 1.12). Therefore, it suffices to show that the existence of pairwise orthogonal projections $p_1, \dots, p_n \in (A^\omega \cap A')^{\alpha^\omega}$ such that $\tau(ap_i) = \tau(a)/n$ for all $a \in A$ and $\tau \in T_\omega(A)$ implies, for any $\|\cdot\|_{2, T_\omega(A)}$ -separable $S \subset A^\omega$, the existence of pairwise orthogonal projections $p'_1, \dots, p'_n \in (A^\omega \cap S')^{\alpha^\omega}$ such that $\tau(ap'_i) = \tau(a)/n$ for all $a \in S$ and $\tau \in T_\omega(A)$. This follows by a standard reindexation argument, which we omit. □

The next part of this section is devoted to proving an equivalence between equivariant property Gamma for an action $\alpha : G \curvearrowright A$ and local equivariant property Gamma with respect to bounded traces for its induced action $\alpha^\omega : G \curvearrowright A^\omega \cap A'$, at least in the setting when A is simple nuclear and has stable rank one.¹² Recall that A is said to have stable rank one if the invertibles of \tilde{A} are dense in \tilde{A} . We start by observing the following description of the tracial state space of $A^\omega \cap A'$.

Proposition 2.5. Let A be a separable, simple, nuclear C*-algebra with uniform property Gamma and stable rank one. Then every tracial state on $A^\omega \cap A'$ is a canonical trace, i.e., one has

$$T(A^\omega \cap A') = \overline{\text{conv}}^{w*} \{ \tau_a \mid \tau \in \tilde{T}_\omega(A), a \in A_+, \tau(a) = 1 \}.$$

Proof. Using exactly the same argument as in the proof of [Castillejos and Evington 2021, Lemma 3.3] and modifying it as hinted in the remark stated before [Castillejos and Evington 2021, Theorem 3.4], we may appeal to [Antoine et al. 2022, Theorem 7.13] (since we assume stable rank one) and pick a nonzero hereditary C*-subalgebra $B \subset A \otimes \mathbb{K}$ with $T^+(B) = \mathbb{R}^{>0}T(B)$ and for which $T(B)$ is nonempty and

¹¹We note that this is automatic if one assumes, e.g., that A has continuous scale (see [Lin 1991, Definition 2.5]), which is a rather common assumption in the context of classification.

¹²Although we use it in the proof, it is likely that stable rank one is not so important for the claim to hold, although we take no guess as to pinning down the correct general assumptions. We note, however, that simple finite \mathcal{Z} -stable C*-algebras have stable rank one; see [Fu et al. 2022; Rørdam 2004].

compact. By Brown’s theorem, it follows that A and B are stably isomorphic. Proposition 1.15 implies that we have an isomorphism $A^\omega \cap A' \cong B^\omega \cap B'$ that induces a bijection between the canonical traces on the left and the right. Hence the claim holds for A if and only if it holds for B .

Now B has uniform property Gamma (see Remark 2.2), so [Castillejos et al. 2021b, Lemma 3.7] implies that B has CPoU. By the “no silly trace” theorem [Castillejos et al. 2021a, Proposition 2.5],¹³ one has that $T(B^\omega)$ is the weak- $*$ -closed convex hull of the limit traces. If B is unital, then the claim follows directly from [Castillejos et al. 2021b, Proposition 4.6]. If B is nonunital, we can extend the inclusion map $B \subset B^\omega$ to a unital inclusion $B^\dagger \subset B^\omega$. From this point of view, we have a trivial equality of algebras

$$B^\omega \cap B' = B^\omega \cap (B^\dagger)' \cap \{1_{B^\omega} - 1_{B^\dagger}\}^\perp.$$

In this case it follows from [Castillejos and Evington 2020, Proposition 5.7] that $T(B^\omega \cap B')$ is the closed convex hull of traces of the form τ_a , where $\tau \in T_\omega(B)$ is a limit trace and $a \in B^\dagger$ is a positive element with $\tau(a) = 1$. If $(e_n)_{n \in \mathbb{N}}$ is an increasing approximate unit in B , then $b_n = e_n a e_n \in B$ converges to a strictly, and hence $\|b_n - a\|_{2,\tau} \rightarrow 0$. This implies the convergence of tracial states $\tau(b_n)^{-1} \tau_{b_n} \rightarrow \tau_a$ in the norm topology, so we observe the equality

$$T(B^\omega \cap B') = \overline{\text{conv}}^{w*} \{ \tau_b \mid \tau \in T_\omega(B), b \in B_+, \tau(b) = 1 \}. \quad \square$$

Theorem 2.6. *Let A be a separable, simple, nuclear C^* -algebra with stable rank one. Then $\alpha : G \curvearrowright A$ has equivariant uniform property Gamma if and only if $\alpha^\omega : G \curvearrowright A^\omega \cap A'$ has local equivariant uniform property Gamma with respect to bounded traces.*

Proof. In order to increase readability in this proof, let us specify another free ultrafilter κ on \mathbb{N} (which may or may not be equal to ω).

We shall show the “if” part first, which actually holds for arbitrary separable simple C^* -algebras with $Q\tilde{T}_2(A) = \tilde{T}(A)$ and $T^+(A) \neq \emptyset$. Let $k \geq 2$. Assuming α^ω has local equivariant property Gamma with respect to bounded traces, we can find pairwise orthogonal projections $p_1, \dots, p_k \in ((A^\omega \cap A')^{\kappa,b})^{(\alpha^\omega)^\kappa}$ such that

$$\tau(ap_j) = \frac{1}{k} \tau(a) \quad \text{for } j = 1, \dots, k, \quad a \in A, \quad \tau \in T_\kappa(A^\omega \cap A').$$

For each $j = 1, \dots, k$, let p_j be represented by a sequence of positive contractions $(p_{j,n})_{n \in \mathbb{N}}$ in $A^\omega \cap A'$. Let in turn each element $p_{j,n}$ be represented by a central sequence $(x_{j,n,\ell})_{\ell \in \mathbb{N}}$ of positive contractions in A . Traces in $T_\kappa(A^\omega \cap A')$ in particular include limit traces associated to sequences of canonical traces. Let $C \subset \mathcal{P}(A)_+ \setminus \{0\}$ be a countable dense subset. Let $K \subset T^+(A)$ be a compact generator. By the conclusion of Lemma 1.17, it follows, for all $a \in C$ and all sequences $(\theta_\ell)_{\ell \in \mathbb{N}}$ in K , that, if τ is the limit trace on A_ω induced by $(\theta_\ell)_{\ell \in \mathbb{N}}$ and τ_a is the induced bounded trace on $A^\omega \cap A'$ that we view in a trivial way as a multiple of a (constant) limit trace on $(A^\omega \cap A')^{\kappa,b}$, then

$$0 = \lim_{n \rightarrow \kappa} \|p_{j,n} - p_{j,n}^2\|_{1,\tau_a} = \lim_{n \rightarrow \kappa} \tau(a|p_{j,n} - p_{j,n}^2|) \stackrel{\text{Lemma 1.17}}{=} \lim_{n \rightarrow \kappa} \lim_{\ell \rightarrow \omega} \theta_\ell(a|x_{j,n,\ell} - x_{j,n,\ell}^2|).$$

¹³Strictly speaking the conclusion is about the reduced tracial product B^∞ in the reference, but this makes no difference to the argument there.

Since the sequence $(\theta_\ell)_{\ell \in \mathbb{N}}$ in K was arbitrary, we may rewrite this as

$$0 = \lim_{n \rightarrow \kappa} \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(a|x_{j,n,\ell} - x_{j,n,\ell}^2|).$$

We may argue in a completely analogous fashion to see that

$$0 = \lim_{n \rightarrow \kappa} \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(a|x_{j,n,\ell} - \alpha_g(x_{j,n,\ell})|), \quad g \in G,$$

as well as

$$0 = \lim_{n \rightarrow \kappa} \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \left| \theta(ax_{j,n,\ell}) - \frac{1}{k} \theta(a) \right| = \lim_{n \rightarrow \kappa} \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(ax_{j,n,\ell} x_{i,n,\ell})$$

for all $i, j = 1, \dots, k$ with $i \neq j$. Lastly, we have by definition that $(x_{j,n,\ell})_{\ell \in \mathbb{N}}$ is a central sequence as $\ell \rightarrow \omega$. Appealing to Kirchberg’s ε -test, we can find central sequences of positive contractions $e_\ell^{(j)}$ in A for $j = 1, \dots, k$ satisfying, for all $a \in C$, the properties

$$0 = \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \left| \theta(ae_\ell^{(j)}) - \frac{1}{k} \theta(a) \right| = \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(a|e_\ell^{(j)} - e_\ell^{(j)2}|)$$

and

$$0 = \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(ae_\ell^{(j)} e_\ell^{(i)}) = \lim_{\ell \rightarrow \omega} \max_{\theta \in K} \theta(a|e_\ell^{(j)} - \alpha_g(e_\ell^{(j)})|), \quad g \in G \text{ and } i \neq j.$$

We consider the resulting elements $e_j \in A^\omega \cap A'$ represented by $(e_\ell^{(j)})_{\ell \in \mathbb{N}}$. Given that C was dense in A_+ , we may conclude that they are pairwise orthogonal projections belonging to $(A^\omega \cap A')^{\alpha^\omega}$ satisfying $\tau_a(e_j) = \tau(a)/k$ for all $\tau \in \tilde{T}_\omega(A)$ and $a \in C$ with $\tau(a) < \infty$. In conclusion, this shows that α has equivariant uniform property Gamma.

For the “only if” part, suppose that α has equivariant property Gamma. Given $k \geq 2$, there exist pairwise orthogonal projections $p_1, \dots, p_k \in (A^\omega \cap A')^{\alpha^\omega}$ such that, for all $a \in A_+$ and $\tilde{\tau} \in \tilde{T}_\omega(A)$ with $\tilde{\tau}(a) < \infty$,

$$\tilde{\tau}_a(p_j) = \frac{1}{k} \tilde{\tau}(a) \quad \text{for } j = 1, \dots, k.$$

As above, choose a compact generator $K \subset T^+(A)$. If we represent each element p_j by a central sequence of positive contractions $(p_{j,n})_{n \in \mathbb{N}}$ in A , then we can argue as before and see that, for all $a \in \mathcal{P}(A)_+ \setminus \{0\}$, $g \in G$, and $i \neq j$, one has the limit properties

$$0 = \lim_{n \rightarrow \omega} \max_{\theta \in K} \left| \theta(ap_{j,n}) - \frac{1}{k} \theta(a) \right| = \lim_{n \rightarrow \omega} \max_{\theta \in K} \theta(a|p_{j,n} - \alpha_g(p_{j,n})|)$$

and

$$0 = \lim_{n \rightarrow \omega} \max_{\theta \in K} \theta(ap_{j,n} p_{i,n}) = \lim_{n \rightarrow \omega} \max_{\theta \in K} \theta(a|p_{j,n} - p_{j,n}^2|).$$

Now take a countable subset $S \subset (A^\omega \cap A')^{\kappa,b}$ whose closure would represent a separable subset as in Definition 2.3. Without loss of generality, let us assume S consists of positive elements. Choose a countable subset $S_0 \subset (A^\omega \cap A')_+$ such that every element of S is represented by a bounded S_0 -valued sequence. Next, choose an increasing sequence of finite sets $F_n \subset \mathcal{P}(A)_+ \setminus \{0\}$ such that their union is dense in A_+ and every element in S_0 has a representing sequence in $\prod_{n \in \mathbb{N}} F_n$. Appealing to the above

stated properties of the sequences $(p_{j,n})_{n \in \mathbb{N}}$ for $j = 1, \dots, n$, we may find an increasing sequence of natural numbers $\ell \mapsto n_\ell$ such that the resulting subsequences satisfy

$$0 = \lim_{\ell \rightarrow \infty} \max_{a \in F_\ell} \|[a, p_{j,n_\ell}]\| = \lim_{\ell \rightarrow \infty} \max_{a \in F_\ell} \max_{\theta \in K} \theta(a|p_{j,n_\ell} - \alpha_g(p_{j,n_\ell})|), \tag{2-1}$$

$$0 = \lim_{\ell \rightarrow \infty} \max_{a \in F_\ell} \max_{\theta \in K} \theta(ap_{j,n_\ell} p_{i,n_\ell}) = \lim_{\ell \rightarrow \infty} \max_{a \in F_\ell} \max_{\theta \in K} \theta(a|p_{j,n_\ell} - p_{j,n_\ell}^2|), \tag{2-2}$$

and

$$0 = \lim_{\ell \rightarrow \infty} \max_{a, b \in F_\ell} \max_{\theta \in K} \left| \theta(abp_{j,n_\ell}) - \frac{1}{k} \theta(ab) \right| \tag{2-3}$$

for all $i, j = 1, \dots, k$ with $i \neq j$. By the choice of the sets F_ℓ , we can see that $(p_{j,n_\ell})_{\ell \in \mathbb{N}}$ defines a central sequence in A , and its induced element $e_j \in A^\omega \cap A'$ commutes with elements in S_0 . We keep in mind the conclusion of [Lemma 1.17](#). Then conditions (2-1) and (2-2) imply that e_1, \dots, e_k are pairwise orthogonal projections in $(A^\omega \cap A')^{\alpha^\omega}$. Condition (2-3) implies that, for all $j = 1, \dots, k$, $\tau \in \tilde{T}_\omega(A)$, every $a \in A_+$ with $\tau(a) = 1$, and every $b \in S_0$, we have

$$\tau_a(be_j) = \tau(abe_j) = \frac{1}{k} \tau(ab) = \frac{1}{k} \tau_a(b).$$

By [Proposition 2.5](#), the weak- $*$ -closed convex hull of such tracial states τ_a yields the whole tracial state space of $A^\omega \cap A'$. In other words, we may conclude

$$\tau(be_j) = \frac{1}{k} \tau(b) \quad \text{for all } j = 1, \dots, k, b \in S_0 \text{ and } \tau \in T(A^\omega \cap A').$$

We may view e_j as constant elements inside $(A^\omega \cap A')^{\kappa, b}$. Since every element in S was represented by a sequence in S_0 , we may conclude that the elements e_1, \dots, e_k satisfy the required property from [Definition 2.3](#) applied to the action

$$\alpha^\omega : G \curvearrowright A^\omega \cap A'.$$

We conclude that α^ω has local equivariant property Gamma with respect to bounded traces. □

3. Dynamical complemented partitions of unity

This section contains the most involved technical arguments of the article, namely the proof that local equivariant property Gamma implies the existence of a dynamical version of complemented partitions of unity [[Castillejos et al. 2021b](#), Definition 3.1], or dynamical CPoU for short. In the case where the induced action on the tracial state space has the property that all orbits are finite with uniformly bounded cardinality, a different iteration of dynamical CPoU was proved in [[Gardella et al. 2022](#), Theorem 4.3]. However, we note that the general statement we prove is a weaker and more intricate version compared to earlier versions but will nevertheless be sufficient to deduce the tracial local-to-global principle.

The starting point for the approach in this section is the following weaker version of CPoU shown in [[Castillejos et al. 2021b](#), Lemma 3.6] for nuclear C^* -algebras, which turns out to hold automatically with the aid of the theory of tracially complete C^* -algebras [[Carrión et al. 2023a](#)].

Proposition 3.1. *Let A be a σ -unital C*-algebra with $T(A)$ nonempty and compact. Then, for every $\|\cdot\|_{2, T_\omega(A)}$ -separable subset $S \subset A^{\omega, b}$, every $k \in \mathbb{N}$, every family $a_1, \dots, a_k \in A_+$, and every*

$$\delta > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i),$$

there exist $e_1, \dots, e_k \in (A^{\omega, b} \cap S'_+)^1$ such that, for all $\tau \in T_\omega(A)$,

- $\tau(\sum_{i=1}^k e_i) = 1$,
- $\tau(a_i e_i) \leq \delta \tau(e_i)$ for $i = 1, \dots, k$.

Proof. Let S , k , a_1, \dots, a_k , and δ be chosen as in the assumption. Set

$$\delta_0 := \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i) < \delta.$$

Since S is $\|\cdot\|_{2, T_\omega(A)}$ -separable, it is first of all clear that one may find a nondegenerate separable C*-subalgebra $A_0 \subseteq A$ containing the tuple a_1, \dots, a_k such that every element of S can be represented by a bounded sequence in A_0 . As every tracial state on A restricts to one on A_0 , the tracial state space of A_0 is still nonempty and compact, and furthermore

$$\sup_{\tau \in T(A_0)} \min_{i=1, \dots, k} \tau(a_i) \geq \delta_0.$$

Let $\eta > 0$. We claim that there exists a finite set $F_\eta \Subset A$ and $\varepsilon_\eta > 0$ such that, if ρ is any state on A with

$$\max_{x \in F_\eta} |\rho(x^*x) - \rho(xx^*)| < \varepsilon_\eta,$$

then $\min_{i=1, \dots, k} \rho(a_i) < \delta_0 + \eta$. If we suppose for a moment that this were false, then it follows that, for every finite set $F \Subset A$ and every $\varepsilon > 0$, there exists a state $\rho_{(F, \varepsilon)}$ on A with

$$\max_{x \in F} |\rho_{(F, \varepsilon)}(x^*x) - \rho_{(F, \varepsilon)}(xx^*)| < \varepsilon \quad \text{and} \quad \min_{i=1, \dots, k} \rho_{(F, \varepsilon)}(a_i) \geq \delta_0 + \eta.$$

We can view $\rho_{(F, \varepsilon)}$ as a net of states by equipping the set of pairs (F, ε) with the obvious order. By the Banach–Anaoğlu theorem, there exists a subset $(\rho_\lambda)_{\lambda \in \Lambda}$ that weak-*converges to a positive functional ρ' with norm at most one on A . By the properties of the net $\rho_{(F, \varepsilon)}$, it is clear that ρ' is tracial. Hence $\min_{i=1, \dots, k} \rho'(a_i) \leq \delta_0$, while at the same time

$$\min_{i=1, \dots, k} \rho'(a_i) = \lim_{(F, \varepsilon)} \min_{i=1, \dots, k} \rho_{(F, \varepsilon)}(a_i) \geq \delta_0 + \eta,$$

which is a contradiction.

Using this intermediate claim, we choose for each $n \geq 1$ a finite set $F_n \Subset A$ and $\varepsilon_n > 0$ satisfying the above conclusion for $\eta = 1/n$. Let $A_1 \subseteq A$ be the C*-algebra generated by A_0 and all the finite sets F_n , which is clearly still separable. Since A_1 contains all the finite sets F_n , it follows that every tracial state τ on A_1 must satisfy

$$\min_{i=1, \dots, k} \tau(a_i) \leq \delta_0 + \frac{1}{n}, \quad n \geq 1,$$

which leads to

$$\sup_{\tau \in T(A_1)} \min_{i=1, \dots, k} \tau(a_i) = \delta_0 < \delta.$$

By all the properties arranged for the subalgebra $A_1 \subseteq A$ so far, it is clear for proving our main claim that we may swap A for the subalgebra A_1 . In other words, we may assume without loss of generality that A is separable.

By [Carrion et al. 2023a, Definition 3.19, Proposition 3.23], the tracial completion $\bar{A}^{T(A)}$ of A yields a factorial tracially complete C^* -algebra. Note that as per the ultraproduct construction of tracially complete C^* -algebras in [Carrion et al. 2023a], the object $(\bar{A}^{T(A)})^\omega$ in that sense becomes canonically isomorphic to the C^* -algebra $A^{\omega, b}$ as considered in Definition 1.8. Because A is separable, $\bar{A}^{T(A)}$ is $\|\cdot\|_{2, T(A)}$ -separable. Thus we may directly apply [Carrion et al. 2023a, Theorem 6.15] (inserting the unit in place of the projection q appearing there) and find the elements $e_1, \dots, e_k \in (A^{\omega, b} \cap S')^1_+$ with the desired properties. \square

The main achievement of this section is the following technical lemma.

Lemma 3.2. *Given $\varepsilon > 0$ and $t \in (0, 1)$, there exists a universal constant $\eta = \eta(\varepsilon, t) > 0$ such that the following holds: Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact. Let G be a countable discrete group, and let $\alpha : G \curvearrowright A$ be an action with local equivariant property Gamma with respect to bounded traces. Suppose that $F, H \in G$ are finite subsets such that*

$$|gH\Delta H| < \eta|H| \quad \text{for all } g \in F.$$

Then, for every $\|\cdot\|_{2, T_\omega(A)}$ -separable subset $S \subset A^{\omega, b}$, every family $a_1, \dots, a_k \in (A^{\omega, b})_+$, and every constant $\delta > 0$ with

$$\frac{\delta}{|H|} > \sup_{\tau \in T_\omega(A)} \min_{i=1, \dots, k} \tau(a_i), \tag{3-1}$$

there exist pairwise orthogonal projections $p_1, \dots, p_k \in A^{\omega, b} \cap S'$ such that, for all $\tau \in T_\omega(A)$, one has

$$\tau(p_1 + \dots + p_k) > t, \tag{3-2}$$

$$\tau(a_i p_i) \leq \delta \tau(p_i) \quad \text{for } i = 1, \dots, k, \tag{3-3}$$

$$\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2 < \varepsilon. \tag{3-4}$$

Remark 3.3. A standard argument shows that the statement in Lemma 3.2 is equivalent to the existence of a universal constant $\eta(\varepsilon, t) > 0$ satisfying the following statement (using approximations instead of the uniform bounded tracial ultrapower):

If $\alpha : G \curvearrowright A$ is an action and $F, H \in G$ are all given as in Lemma 3.2, then, for every finite subset $S \in A$, every $\xi > 0$, every family $a_1, \dots, a_k \in A_+$, and every $\delta > 0$ with

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i), \tag{3-5}$$

there exist pairwise orthogonal contractions $e_1, \dots, e_k \in A_+$ such that

$$\begin{aligned} \|[e_i, x]\|_{2,u} &< \xi \quad \text{for } x \in S, \quad i = 1, \dots, k, \\ \|e_i - e_i^2\|_{2,u} &< \xi \quad \text{for } i = 1, \dots, k, \\ \tau(e_1 + \dots + e_k) &> t - \xi \quad \text{for } \tau \in T(A), \\ \tau(a_i e_i) &< \delta \tau(e_i) + \xi \quad \text{for } \tau \in T(A), \quad i = 1, \dots, k, \\ \max_{g \in F} \sum_{i=1}^k \|\alpha_g(e_i) - e_i\|_{2,\tau} &< \varepsilon + \xi \quad \text{for } \tau \in T(A). \end{aligned}$$

In particular, this means that it suffices to prove [Lemma 3.2](#) for positive elements a_1, \dots, a_k taken in A instead of $A^{\omega,b}$. In this case, (3-1) and (3-5) are equivalent.

The proof of [Lemma 3.2](#) is an adapted version of the proof in the nondynamical setting (cf. [[Castillejos et al. 2021b](#), Section 3]) but also incorporates new ideas related to the dynamical structure. Before we delve into the details, we shall give an overview of the strategy. The construction of the pairwise orthogonal projections p_1, \dots, p_k in the statement of [Lemma 3.2](#) is done in three steps:

(1) Instead of producing pairwise orthogonal projections p_1, \dots, p_k , we start by producing (not yet pairwise orthogonal) positive contractions $e_1, \dots, e_k \in A^{\omega,b} \cap S'$ that satisfy

$$\tau(e_1 + \dots + e_k) = 1, \quad \tau(a_i e_i) \leq \delta \tau(e_i) \quad \text{for } i = 1, \dots, k, \quad \tau \in T_\omega(A),$$

and that are approximately invariant under α^ω in the right sense. This is done in [Lemma 3.4](#) and is the only part of the proof that makes use of the approximate Følner property that appears in the assumption of the lemma.

(2) Next, we use equivariant property Gamma to turn these contractions into orthogonal projections $p'_1, \dots, p'_k \in A^{\omega,b} \cap S'$. As a consequence of this procedure we get that

$$\tau(p'_1 + \dots + p'_k) = \frac{1}{k} \quad \text{for } \tau \in T_\omega(A),$$

but they still satisfy (3-3) and are still approximately invariant in the right sense. This is done in [Lemma 3.6](#).

(3) In order to enlarge the trace of the sum of the projections, we repeat the above steps underneath the projection $1_{A^\omega} - \sum_{i=1}^k p'_i$.¹⁴ We continue this procedure inductively until we end up with orthogonal projections p_1, \dots, p_k whose sum exceeds t in trace and that still satisfy (3-3). If everything is done carefully from the start and $\eta > 0$ is chosen correctly, we can control the error in the invariance of the projections and make sure they satisfy (3-4) in the end. (We note, informally, that this error grows with the number of times this procedure is repeated, which is the ultimate reason why we cannot simply work with $t = 1$ in the statement.)

We shall now implement the above strategy. Combining the contractions arising from [Proposition 3.1](#) with an averaging argument over suitable Følner sets allows us to carry out the first step.

¹⁴For this one actually needs a somewhat stronger version of the second step; see [Lemma 3.7](#).

Lemma 3.4. *Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action by a countable discrete group. Let $\varepsilon > 0$, and let finite subsets $F \subseteq G$ and $H \subseteq G$ be given such that $|gH\Delta H| < \varepsilon|H|$ for all $g \in F$. Then, for all $\|\cdot\|_{2, T_\omega(A)}$ -separable subsets $S \subset A^{\omega, b}$, all $\delta > 0$, and all $a_1, \dots, a_k \in A_+$ with*

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i),$$

there exist $e_1, \dots, e_k \in (A^{\omega, b} \cap S')^1_+$ such that, for $\tau \in T_\omega(A)$,

- $\tau(\sum_{i=1}^k e_i) = 1$,
- $\tau(a_i e_i) \leq \delta \tau(e_i)$ for $i = 1, \dots, k$, and
- $\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau} < \varepsilon$.

Proof. Given $S \subset A^{\omega, b}$, $\delta > 0$, and $a_1, \dots, a_k \in A_+$ as above, we define

$$a'_i := \frac{1}{|H|} \sum_{g \in H} \alpha_{g^{-1}}(a_i), \quad i = 1, \dots, k.$$

Note that, for each $\tau \in T(A)$, the trace $\frac{1}{|H|} \sum_{g \in H} \tau \circ \alpha_{g^{-1}}$ is again an element of $T(A)$, so we see that

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a'_i).$$

By Proposition 3.1, we know that there exist

$$e'_1, \dots, e'_k \in \left(A^\omega \cap \left(\bigcup_{g \in G} \alpha_g^\omega(S) \right)' \right)^1_+$$

such that, for $\tau \in T_\omega(A)$,

$$\tau\left(\sum_{i=1}^k e'_i\right) = 1, \tag{3-6}$$

$$\tau(a'_i e'_i) \leq \frac{\delta}{|H|} \tau(e'_i) \quad \text{for } i = 1, \dots, k.$$

In particular, this last equation implies that

$$\tau(\alpha_{g^{-1}}(a_i) e'_i) \leq \delta \tau(e'_i), \quad g \in H, \quad \tau \in T_\omega(A). \tag{3-7}$$

Now, for $i = 1, \dots, k$, define

$$e_i := |H|^{-1} \sum_{g \in H} \alpha_g^\omega(e'_i).$$

Clearly this still is a positive contraction in $A^\omega \cap S'$. Notice that

$$\tau\left(\sum_{i=1}^k e_i\right) = |H|^{-1} \sum_{g \in H} (\tau \circ \alpha_g^\omega)\left(\sum_{i=1}^k e'_i\right) \stackrel{(3-6)}{=} 1 \quad \text{for } \tau \in T_\omega(A).$$

For $\tau \in T_\omega(A)$ and $i = 1, \dots, k$, we have

$$\begin{aligned} \tau(a_i e_i) &= |H|^{-1} \tau \left(a_i \sum_{g \in H} \alpha_g^\omega(e'_i) \right) = |H|^{-1} \sum_{g \in H} (\tau \circ \alpha_g^\omega)(\alpha_{g^{-1}}(a_i) e'_i) \\ &\stackrel{(3-7)}{\leq} |H|^{-1} \sum_{g \in H} \delta(\tau \circ \alpha_g^\omega)(e'_i) = \delta\tau(e_i). \end{aligned}$$

Lastly, we see that, for $g \in F$ and $\tau \in T_\omega(A)$, we have

$$\begin{aligned} \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1,\tau} &\leq |H|^{-1} \sum_{i=1}^k \sum_{h \in gH\Delta H} \|\alpha_h^\omega(e'_i)\|_{1,\tau} = |H|^{-1} \sum_{h \in gH\Delta H} \sum_{i=1}^k \tau(\alpha_h^\omega(e'_i)) \\ &\stackrel{(3-6)}{=} |H|^{-1} |gH\Delta H| < \varepsilon. \quad \square \end{aligned}$$

Analogously as in the nondynamical setting (cf. [Castillejos et al. 2021b, Lemma 2.4]), (local) equivariant property Gamma allows one to replace positive contractions by projections without changing the tracial values in $A^{\omega,b}$. A different generalization of this lemma was proved in [Gardella et al. 2022, Proposition 3.4], but for the purposes of this paper we need a way to control the (tracially) approximate fixedness of the elements for the action.

Lemma 3.5. *Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact, and let $\alpha : G \curvearrowright A$ be an action of a countable discrete group. Assume that α has local equivariant property Gamma with respect to bounded traces. Let $S \subset A^{\omega,b}$ be a $\|\cdot\|_{2,T_\omega(A)}$ -separable subset, and let $b \in A^{\omega,b} \cap S'$ be a positive contraction. Then there exists a projection $p \in A^{\omega,b} \cap S'$ such that*

$$\tau(ap) = \tau(ab) \quad \text{for } a \in S, \tau \in T_\omega(A), \quad (3-8)$$

and such that, for all $g \in G$ and $\tau \in T_\omega(A)$, one has

$$\|\alpha_g^\omega(p) - p\|_{2,\tau}^2 \leq \|\alpha_g^\omega(b) - b\|_{2,\tau}^2 + \|b\|_{2,\tau}^2. \quad (3-9)$$

Proof. Fix $n \in \mathbb{N}$. By a common reindexation trick, it suffices to find a positive contraction $p \in A^{\omega,b} \cap S'$ satisfying (3-8), (3-9), and $\|p - p^2\|_{2,T_\omega(A)}^2 \leq 1/n$. An element p satisfying all the necessary properties except (3-9) is constructed in the proof of [Castillejos et al. 2021b, Lemma 2.4] with the use of uniform property Gamma. We show that when the construction is carried out using (local) equivariant property Gamma instead, the resulting projection also satisfies the extra condition (3-9).

Just as in [Castillejos et al. 2021b], we define functions $f_1, \dots, f_n \in C([0, 1])$ such that $f_i|_{[0, (i-1)/n]} = 0$, $f_i|_{[(i-1)/n, 1]} = 1$, and f_i is linear on $[(i-1)/n, i/n]$. Note that not only $(1/n) \sum_{i=1}^n f_i = \text{id}_{[0,1]}$, but the monotonicity of each f_i also implies

$$\frac{1}{n} \sum_{i=1}^n |f_i(t_1) - f_i(t_2)| = |t_1 - t_2|, \quad t_1, t_2 \in [0, 1]. \quad (3-10)$$

By local equivariant property Gamma with respect to bounded traces, we can find pairwise orthogonal projections $p_1, \dots, p_n \in (A^{\omega,b})^{\alpha^\omega} \cap S' \cap \{b\}'$ such that $\tau(p_i x) = \tau(x)/n$ for $i = 1, \dots, n$, $\tau \in T_\omega(A)$,

$x \in C^*(S \cup \{b\})$. Define

$$p := \sum_{i=1}^n p_i f_i(b) \in A^{\omega, b} \cap S'.$$

By repeating the arguments in the proof of [Castillejos et al. 2021b, Lemma 2.4] verbatim, we may conclude that $\|p - p^2\|_{2, T_\omega(A)}^2 \leq 1/n$ and $\tau(ap) = \tau(ab)$ for all $a \in S$ and $\tau \in T_\omega(A)$. We need to show that (3-9) holds as well. Fix $\tau \in T_\omega(A)$ and $g \in G$. By [Connes 1976, I.1], there exists a positive Radon measure ν on $[0, 1]^2$ such that, for every pair of functions $h_1, h_2 \in C_0((0, 1])$, the functions $(s, t) \mapsto h_1(s)$ and $(s, t) \mapsto h_2(t)$ are square integrable and

$$\|h_1(\alpha_g^\omega(b)) - h_2(b)\|_{2, \tau}^2 = \int_{[0, 1]^2} |h_1(s) - h_2(t)|^2 d\nu(s, t).$$

Then we get

$$\begin{aligned} \|\alpha_g^\omega(p) - p\|_{2, \tau}^2 &= \left\| \sum_{i=1}^n p_i (f_i(\alpha_g^\omega(b)) - f_i(b)) \right\|_{2, \tau}^2 \\ &= \frac{1}{n} \sum_{i=1}^n \|f_i(\alpha_g^\omega(b)) - f_i(b)\|_{2, \tau}^2 \\ &= \frac{1}{n} \sum_{i=1}^n \int_{[0, 1]^2} |f_i(s) - f_i(t)|^2 d\nu(s, t) \\ &\leq \frac{1}{n} \sum_{i=1}^n \int_{[0, 1]^2} |f_i(s) - f_i(t)| d\nu(s, t) \\ &\stackrel{(3-10)}{=} \int_{[0, 1]^2} |s - t| d\nu(s, t) \\ &\leq \sqrt{\int_{[0, 1]^2} |s^{1/2} - t^{1/2}|^2 d\nu(s, t)} \sqrt{\int_{[0, 1]^2} |s^{1/2} + t^{1/2}|^2 d\nu(s, t)} \\ &= \|\alpha_g^\omega(b)^{1/2} - b^{1/2}\|_{2, \tau} \|\alpha_g^\omega(b)^{1/2} + b^{1/2}\|_{2, \tau}. \quad \square \end{aligned}$$

Using Lemma 3.5, we can construct orthogonal projections that play a similar role to the positive elements in Lemma 3.4.

Lemma 3.6. *Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action by a countable discrete group and assume it has local equivariant property Gamma with respect to bounded traces. Let $\varepsilon > 0$, $F \in G$, and $H \in G$ be such that $|gH\Delta H| < \varepsilon|H|$ for all $g \in F$. Then, for every $\|\cdot\|_{2, T_\omega(A)}$ -separable subset $S \subset A^{\omega, b}$, all $\delta > 0$, and all $a_1, \dots, a_k \in A_+$ with*

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i),$$

there exist pairwise orthogonal projections $p_1, \dots, p_k \in A^{\omega, b} \cap S'$ such that, for all $\tau \in T_\omega(A)$,

- $\tau(\sum_{i=1}^k p_i) = 1/k$,
- $\tau(a_i p_i) \leq \delta \tau(p_i)$ for $i = 1, \dots, k$, and
- $\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2 < 2\sqrt{\varepsilon}/k$.

Proof. By [Lemma 3.4](#), we can find $e_1, \dots, e_k \in (A^{\omega, b} \cap S')_+^1$ such that, for all $\tau \in T_\omega(A)$,

$$\tau\left(\sum_{i=1}^k e_i\right) = 1, \quad (3-11)$$

$$\tau(a_i e_i) \leq \delta \tau(e_i) \quad \text{for } i = 1, \dots, k, \quad (3-12)$$

$$\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau} < \varepsilon. \quad (3-13)$$

Let $S_0 = S \cup \{1_{A^{\omega, b}}, a_1, \dots, a_k\}$. Apply [Lemma 3.5](#) for each $i \in \{1, \dots, k\}$ and find a projection $p_i \in A^{\omega, b} \cap S'_0$ such that, for all $a \in S_0$, $\tau \in T_\omega(A)$, and $g \in G$, we have

$$\tau(ap_i) = \tau(ae_i), \quad (3-14)$$

$$\|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2 \leq \|\alpha_g^\omega(e_i)^{1/2} - e_i^{1/2}\|_{2, \tau} \|\alpha_g^\omega(e_i)^{1/2} + e_i^{1/2}\|_{2, \tau}. \quad (3-15)$$

This already implies the two following facts:

$$\tau\left(\sum_{i=1}^k p_i\right) \stackrel{(3-14)}{=} \tau\left(\sum_{i=1}^k e_i\right) \stackrel{(3-11)}{=} 1 \quad \text{for } \tau \in T_\omega(A), \quad (3-16)$$

$$\tau(a_i p_i) \stackrel{(3-14)}{=} \tau(a_i e_i) \stackrel{(3-12)}{\leq} \delta \tau(e_i) \stackrel{(3-14)}{=} \delta \tau(p_i) \quad \text{for } i = 1, \dots, k, \quad \tau \in T_\omega(A). \quad (3-17)$$

Furthermore, we get, for $g \in F$ and $\tau \in T_\omega(A)$, that

$$\begin{aligned} \sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2 &\stackrel{(3-15)}{\leq} \sum_{i=1}^k \|\alpha_g^\omega(e_i)^{1/2} - e_i^{1/2}\|_{2, \tau} \|\alpha_g^\omega(e_i)^{1/2} + e_i^{1/2}\|_{2, \tau} \\ &\leq \sum_{i=1}^k \|\alpha_g^\omega(e_i)^{1/2} - e_i^{1/2}\|_{2, \tau} (\|\alpha_g^\omega(e_i)^{1/2}\|_{2, \tau} + \|e_i^{1/2}\|_{2, \tau}) \\ &\stackrel{\text{Lemma 1.19}}{\leq} \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau}^{1/2} (\|\alpha_g^\omega(e_i)^{1/2}\|_{2, \tau} + \|e_i^{1/2}\|_{2, \tau}) \\ &= \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau}^{1/2} \tau(\alpha_g^\omega(e_i))^{1/2} + \sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau}^{1/2} \tau(e_i)^{1/2} \\ &\leq \sqrt{\sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau} \sum_{i=1}^k (\tau \circ \alpha_g^\omega)(e_i)} + \sqrt{\sum_{i=1}^k \|\alpha_g^\omega(e_i) - e_i\|_{1, \tau} \sum_{i=1}^k \tau(e_i)} \\ &\stackrel{(3-11), (3-13)}{<} 2\sqrt{\varepsilon}. \end{aligned} \quad (3-18)$$

Set

$$S_1 = S \cup C^*({a_1, \dots, a_k} \cup \{\alpha_g^\omega(p_j) \mid 1 \leq j \leq k, g \in G\}) \subset A^{\omega, b}.$$

Since A has local equivariant property Gamma with respect to bounded traces, we can find pairwise orthogonal projections $r_1, \dots, r_k \in (A^{\omega, b})^{\alpha^\omega} \cap S'_1$ such that

$$\tau(r_i a) = \frac{1}{k} \tau(a) \quad \text{for } \tau \in T_\omega(A), \quad a \in S_1, \quad i = 1, \dots, k. \tag{3-19}$$

Set $p'_i := r_i p_i$. Then clearly $p'_1, \dots, p'_k \in A^{\omega, b} \cap S'$ are pairwise orthogonal projections. We get, for each $\tau \in T_\omega(A)$, that

$$\tau\left(\sum_{i=1}^k p'_i\right) = \tau\left(\sum_{i=1}^k r_i p_i\right) \stackrel{(3-19)}{=} \frac{1}{k} \tau\left(\sum_{i=1}^k p_i\right) \stackrel{(3-16)}{=} \frac{1}{k}.$$

For $i = 1, \dots, k$ and $\tau \in T_\omega(A)$, we get

$$\tau(a_i p'_i) = \tau(a_i r_i p_i) \stackrel{(3-19)}{=} \frac{1}{k} \tau(a_i p_i) \stackrel{(3-17)}{\leq} \frac{\delta}{k} \tau(p_i) \stackrel{(3-19)}{=} \delta \tau(p'_i).$$

Lastly, for $g \in F$ and $\tau \in T_\omega(A)$, we get

$$\sum_{i=1}^k \|\alpha_g^\omega(p'_i) - p'_i\|_{2, \tau}^2 \stackrel{(3-19)}{=} \frac{1}{k} \sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2 \stackrel{(3-18)}{<} \frac{2\sqrt{\varepsilon}}{k}.$$

Hence the projections p'_i satisfy all the required properties. This finishes the proof. □

In order to carry out the inductive argument to enlarge the trace of the sum of the constructed orthogonal projections, we need a stronger version of the previous lemma that allows us to find the orthogonal projections under an arbitrary tracially constant projection q .

Lemma 3.7. *Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action by a countable discrete group and assume it has local equivariant property Gamma with respect to bounded traces. Let $\varepsilon > 0$, $F \subseteq G$, and $H \subseteq G$ be such that $|gH\Delta H| < \varepsilon|H|$ for all $g \in F$. Choose $\delta > 0$ and $a_1, \dots, a_k \in A_+$ such that*

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i).$$

For every $\mu \in (0, 1]$ and $\|\cdot\|_{2, T_\omega(A)}$ -separable subset $S_0 \subset A^{\omega, b}$, there exists a $\|\cdot\|_{2, T_\omega(A)}$ -separable subset $S \subset A^{\omega, b}$ such that, if $q \in A^{\omega, b} \cap S'$ is a projection with $\tau(q) = \mu$ for all $\tau \in T_\omega(A)$, there exist pairwise orthogonal projections $p_1, \dots, p_k \in A^{\omega, b} \cap S'_0 \cap \{\alpha_g^\omega(q) \mid g \in G\}'$ such that, for all $\tau \in T_\omega(A)$,

- $\sum_{i=1}^k \tau(p_i q) = \mu/k$,
- $\tau(a_i p_i q) \leq \delta \tau(p_i q)$ for $i = 1, \dots, k$,
- $\max_{g \in F} \sum_{i=1}^k \|q(\alpha_g^\omega(p_i) - p_i)\|_{2, \tau}^2 \leq 2\mu\sqrt{\varepsilon}/k$, and
- $\sum_{i=1}^k \|\alpha_g^\omega(p_i)(\alpha_g^\omega(q) - q)\|_{2, \tau}^2 \leq (1/k)\|\alpha_g^\omega(q) - q\|_{2, T_\omega(A)}^2$ for all $g \in G$.

Proof. Let $\varepsilon > 0$, $F, H \subseteq G$, $\delta > 0$, and $a_1, \dots, a_k \in A_+$ be as in the statement of the lemma. In order to prove the claim, it suffices to prove the following local statement:

For every $\mu \in (0, 1]$, $\zeta > 0$, $T \in A$, and $E \in G$, there exist $S \in A$ and $\xi > 0$ such that, if $q \in A_+^1$ satisfies

$$\|q - q^2\|_{2,u} < \xi, \quad \sup_{\tau \in T(A)} |\tau(q) - \mu| < \xi, \quad \text{and} \quad \|[q, s]\|_{2,u} < \xi \quad \text{for } s \in S,$$

then there exist pairwise orthogonal projections $p_1, \dots, p_k \in A_+^1$ such that

$$\begin{aligned} \|p_i - p_i^2\|_{2,u} &< \zeta, \\ \|[p_i, t]\|_{2,u} &< \zeta \quad \text{for } t \in T \cup \{\alpha_g(q) \mid g \in E\}, \\ \sup_{\tau \in T(A)} \left| \sum_{i=1}^k \tau(p_i q) - \frac{\mu}{k} \right| &< \zeta, \\ \tau(a_i p_i q) &< \delta \tau(p_i q) + \zeta \quad \text{for } \tau \in T(A), \\ \sup_{\tau \in T(A)} \max_{g \in F} \sum_{i=1}^k \|q(\alpha_g(p_i) - p_i)\|_{2,\tau}^2 &< \frac{2\mu\sqrt{\varepsilon}}{k} + \zeta, \\ \sup_{\tau \in T(A)} \max_{g \in E} \sum_{i=1}^k \|\alpha_g(p_i)^{1/2}(\alpha_g(q) - q)\|_{2,\tau}^2 &< \frac{1}{k} \|\alpha_g(q) - q\|_{2,u}^2 + \zeta. \end{aligned}$$

We prove this local statement by contradiction. Suppose there exist $\mu \in (0, 1]$, $\zeta > 0$, $T \in A$, and $E \in G$ for which the statement does not hold. In other words, this means that, for every $\emptyset \neq S \in A$, we can find a $q_S \in A_+^1$ such that

$$\|q_S - q_S^2\|_{2,u} < 1/|S|, \quad \sup_{\tau \in T(A)} |\tau(q_S) - \mu| < 1/|S|, \quad \text{and} \quad \|[q_S, s]\|_{2,u} < 1/|S| \quad \text{for } s \in S,$$

but there exist no pairwise orthogonal projections $p_1, \dots, p_k \in A_+^1$ satisfying

$$\|p_i - p_i^2\|_{2,u} < \zeta, \tag{3-20}$$

$$\|[p_i, t]\|_{2,u} < \zeta \quad \text{for } t \in T \cup \{\alpha_g(q_S) \mid g \in E\}, \tag{3-21}$$

$$\sup_{\tau \in T(A)} \left| \sum_{i=1}^k \tau(p_i q_S) - \frac{\mu}{k} \right| < \zeta, \tag{3-22}$$

$$\tau(a_i p_i q_S) < \delta \tau(p_i q_S) + \zeta \quad \text{for } \tau \in T(A), \tag{3-23}$$

$$\sup_{\tau \in T(A)} \max_{g \in F} \sum_{i=1}^k \|q_S(\alpha_g(p_i) - p_i)\|_{2,\tau}^2 < \frac{2\mu\sqrt{\varepsilon}}{k} + \zeta, \tag{3-24}$$

$$\sup_{\tau \in T(A)} \max_{g \in E} \sum_{i=1}^k \|\alpha_g(p_i)^{1/2}(\alpha_g(q_S) - q_S)\|_{2,\tau}^2 < \frac{1}{k} \|\alpha_g(q_S) - q_S\|_{2,u}^2 + \zeta. \tag{3-25}$$

In this way we get a net $(q_S)_S$ indexed by the finite subsets of A equipped with inclusion as the natural partial order. We can take a free ultrafilter $\tilde{\omega}$ on this index set of finite subsets of A as follows. For each $I \in A$ consider the set

$$P_I = \{J \in A \mid I \subseteq J\}.$$

As the collection of P_I is closed under finite intersections, they form a filter basis and hence there is a minimal filter on the set of finite subsets of A containing all the sets P_I for $I \in \mathcal{A}$. This filter will be free and can be extended to a free ultrafilter $\tilde{\omega}$ by Zorn’s lemma.

Similarly as in Definitions 1.3 and 1.8, we can define the norm and bounded tracial ultrapower of A over the ultrafilter $\tilde{\omega}$. We also get a set of limit traces $T_{\tilde{\omega}}(A)$ over $\tilde{\omega}$. Then the net $(q_S)_S$ defines a projection $q \in A^{\tilde{\omega},b} \cap A'$ with value μ on all limit traces on $A^{\tilde{\omega},b}$. By Lemma 3.6, we can find pairwise orthogonal projections $p'_1, \dots, p'_k \in A_+^1$ such that

$$\|p'_i - p_i\|^2_{2,u} < \zeta \quad \text{for } i = 1, \dots, k, \tag{3-26}$$

$$\|[p'_i, t]\|_{2,u} < \zeta \quad \text{for } t \in T, \tag{3-27}$$

$$\sup_{\tau \in T(A)} \left| \tau \left(\sum_{i=1}^k p'_i \right) - \frac{1}{k} \right| < \frac{1}{4} \zeta, \tag{3-28}$$

$$\tau(a_i p'_i) < \delta \tau(p'_i) + \zeta \quad \text{for } i = 1, \dots, k, \tau \in T(A), \tag{3-29}$$

$$\max_{g \in F} \sum_{i=1}^k \|\alpha_g(p'_i) - p'_i\|^2_{2,\tau} < \frac{2\sqrt{\varepsilon}}{k} \quad \text{for } \tau \in T(A). \tag{3-30}$$

Since $q \in A^{\tilde{\omega},b} \cap A'$, it follows, for each $\tau \in T_{\tilde{\omega}}(A)$ and $g \in G$, that the assignment

$$A \rightarrow \mathbb{C} : a \mapsto \tau(a\alpha_g^{\tilde{\omega}}(q))/\tau(\alpha_g^{\tilde{\omega}}(q)) = \tau(a\alpha_g^{\tilde{\omega}}(q))/\mu$$

defines a tracial state on A . In particular, we get that, for $\tau \in T_{\tilde{\omega}}(A)$, the following hold:

$$\left| \tau \left(\sum_{i=1}^k p'_i \alpha_g^{\tilde{\omega}}(q) \right) - \frac{\mu}{k} \right| \stackrel{(3-28)}{<} \frac{1}{4} \mu \zeta \leq \frac{1}{4} \zeta \quad \text{for } g \in G, \tag{3-31}$$

$$\tau(a_i p'_i q) \stackrel{(3-29)}{<} \delta \tau(p'_i q) + \mu \zeta \leq \delta \tau(p'_i q) + \zeta, \tag{3-32}$$

$$\max_{g \in F} \sum_{i=1}^k \|q(\alpha_g(p'_i) - p'_i)\|^2_{2,\tau} \stackrel{(3-30)}{<} \frac{2\mu\sqrt{\varepsilon}}{k}. \tag{3-33}$$

Next we show that, for all $\tau \in T_{\tilde{\omega}}(A)$ and $g \in G$,

$$\sum_{i=1}^k \|\alpha_g(p'_i)^{1/2}(\alpha_g^{\tilde{\omega}}(q) - q)\|^2_{2,\tau} < \frac{1}{k} \|\alpha_g^{\tilde{\omega}}(q) - q\|^2_{2,u} + \zeta. \tag{3-34}$$

Fix $\tau \in T_{\tilde{\omega}}(A)$ and $g \in G$. Assume $\tau(q\alpha_g^{\tilde{\omega}}(q)) > 0$. Then the map

$$A \rightarrow \mathbb{C}, \quad a \mapsto \tau(aq\alpha_g^{\tilde{\omega}}(q))/\tau(q\alpha_g^{\tilde{\omega}}(q))$$

defines a tracial state on A . This means that

$$\left| \tau \left(\sum_{i=1}^k \alpha_g(p'_i) q \alpha_g^{\tilde{\omega}}(q) \right) - \frac{1}{k} \tau(q\alpha_g^{\tilde{\omega}}(q)) \right| \stackrel{(3-28)}{<} \frac{1}{4} \zeta \tau(q\alpha_g^{\tilde{\omega}}(q)) \leq \frac{1}{4} \zeta. \tag{3-35}$$

Note that, if $\tau(q\alpha_g^{\tilde{\omega}}(q)) = 0$, then

$$\tau\left(\sum_{i=1}^k \alpha_g(p'_i)q\alpha_g^{\tilde{\omega}}(q)\right) = 0 = \frac{1}{k}\tau(q\alpha_g^{\tilde{\omega}}(q)),$$

and hence (3-35) also holds in this case. Now

$$\begin{aligned} \tau\left(\sum_{i=1}^k \alpha_g(p'_i)(\alpha_g^{\tilde{\omega}}(q) - q)^2\right) &= \tau\left(\sum_{i=1}^k \alpha_g(p'_i)(\alpha_g^{\tilde{\omega}}(q) + q - \alpha_g^{\tilde{\omega}}(q)q - q\alpha_g^{\tilde{\omega}}(q))\right) \\ &\stackrel{(3-31),(3-35)}{<} \frac{1}{k}\tau(\alpha_g^{\tilde{\omega}}(q) + q - \alpha_g^{\tilde{\omega}}(q)q - q\alpha_g^{\tilde{\omega}}(q)) + \zeta \\ &= \frac{1}{k}\tau((\alpha_g^{\tilde{\omega}}(q) - q)^2) + \zeta. \end{aligned}$$

This proves (3-34). If we combine this with (3-26), (3-27), and (3-31)–(3-33), we can conclude that, for some q_S in the net, (3-20)–(3-25) must hold. This gives the desired contradiction. \square

Proof of Lemma 3.2. Given $\varepsilon > 0$ and $t \in (0, 1)$, choose $\eta > 0$ small enough that

$$4\left\lceil \frac{t}{1-t} \right\rceil \sqrt{\eta} < \varepsilon. \tag{3-36}$$

We show that such a constant η satisfies the required properties. Let $\alpha : G \curvearrowright A$, finite sets $F, H \in G$, and $S \subset A^{\omega, b}$ be given as in the statement of the lemma. By Remark 3.3, it suffices to consider $\delta > 0$ and $a_1, \dots, a_k \in A_+$ such that

$$\frac{\delta}{|H|} > \sup_{\tau \in T(A)} \min_{i=1, \dots, k} \tau(a_i).$$

We construct the pairwise orthogonal projections p_1, \dots, p_k in $N := k\lceil t/(1-t) \rceil$ steps. Define a sequence (s_n) in $[0, 1)$ inductively by setting $s_0 = 0$ and setting $s_{i+1} = s_i + (1 - s_i)/k$. Note that, when $s < t$, we have $s + (1 - s)/k > s + (1 - t)/k$. If we assumed for a moment that $s_N < t$, then this sequence is less than t for all of the first N steps, leading to $s_N > N(1 - t)/k \geq t$, which is a contradiction; hence we must have $s_N > t$. Next, we construct separable subsets S_0, \dots, S_N as follows: Set $S_N := S$. Given $i \in \{1, \dots, N\}$ such that S_i is defined, let S_{i-1} be the union of S_i and the set determined by Lemma 3.7 with $1 - s_{i-1}$ in place of μ and S_i in place of S_0 .

In the initial step, we set $p_1^{(0)} = \dots = p_k^{(0)} = 0$. Now suppose that, for some $n \in \mathbb{N}$, we have pairwise orthogonal projections $p_1^{(n)}, \dots, p_k^{(n)} \in A^{\omega, b} \cap S'_n$ such that, for all $\tau \in T_\omega(A)$,

$$\tau(p_1^{(n)} + \dots + p_k^{(n)}) = s_n, \tag{3-37}$$

$$\tau(a_i p_i^{(n)}) \leq \delta \tau(p_i^{(n)}) \quad \text{for } i = 1, \dots, k, \tag{3-38}$$

$$\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n)}) - p_i^{(n)}\|_{2, \tau}^2 \leq \frac{4n\sqrt{\eta}}{k}, \tag{3-39}$$

$$\max_{g \in F} \left\| \sum_{i=1}^k \alpha_g^\omega(p_i^{(n)}) - \sum_{i=1}^k p_i^{(n)} \right\|_{2, T_\omega(A)}^2 \leq 2s_n\sqrt{\eta}. \tag{3-40}$$

Note that the $p_i^{(0)}$ trivially satisfy (3-37)–(3-40). We show that we can construct pairwise orthogonal projections $p_1^{(n+1)}, \dots, p_k^{(n+1)} \in A^{\omega, b} \cap S'_{n+1}$ such that, for all $\tau \in T_\omega(A)$,

$$\tau(p_1^{(n+1)} + \dots + p_k^{(n+1)}) = s_{n+1}, \tag{3-41}$$

$$\tau(a_i p_i^{(n+1)}) \leq \delta \tau(p_i^{(n+1)}) \quad \text{for } i = 1, \dots, k, \tag{3-42}$$

$$\max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n+1)}) - p_i^{(n+1)}\|_{2, \tau}^2 \leq \frac{4(n+1)\sqrt{\eta}}{k}, \tag{3-43}$$

$$\max_{g \in F} \left\| \sum_{i=1}^k \alpha_g^\omega(p_i^{(n+1)}) - \sum_{i=1}^k p_i^{(n+1)} \right\|_{2, T_\omega(A)}^2 \leq 2s_{n+1}\sqrt{\eta}. \tag{3-44}$$

Define

$$q := 1_{A^{\omega, b}} - \sum_{i=1}^k p_i^{(n)}.$$

Note that q is a projection in $A^{\omega, b} \cap S'_n$ with $\tau(q) = 1 - s_n$ for all $\tau \in T_\omega(A)$. By Lemma 3.7 and our choice of S_n , we can find pairwise orthogonal projections

$$r_1, \dots, r_k \in A^{\omega, b} \cap S'_{n+1} \cap \{\alpha_g^\omega(q) \mid g \in G\}'$$

such that, for all $\tau \in T_\omega(A)$,

$$\tau\left(\sum_{i=1}^k r_i q\right) = \frac{1 - s_n}{k}, \tag{3-45}$$

$$\tau(a_i r_i q) \leq \delta \tau(r_i q) \quad \text{for } i = 1, \dots, k, \tag{3-46}$$

$$\max_{g \in F} \sum_{i=1}^k \|q(\alpha_g^\omega(r_i) - r_i)\|_{2, \tau}^2 \leq \frac{2(1 - s_n)\sqrt{\eta}}{k}, \tag{3-47}$$

$$\sum_{i=1}^k \|\alpha_g^\omega(r_i)(\alpha_g^\omega(q) - q)\|_{2, \tau}^2 \leq \frac{1}{k} \|\alpha_g^\omega(q) - q\|_{2, T_\omega(A)}^2 \quad \text{for } g \in G. \tag{3-48}$$

Define

$$p_i^{(n+1)} := p_i^{(n)} + q r_i.$$

By construction (recall that $S_{n+1} \subset S_n$), the elements $p_1^{(n+1)}, \dots, p_k^{(n+1)}$ are pairwise orthogonal projections in $A^{\omega, b} \cap S'_{n+1}$. We show that they satisfy (3-41)–(3-44). Fix $\tau \in T_\omega(A)$. Note first of all that

$$\tau\left(\sum_{i=1}^k p_i^{(n+1)}\right) = \tau\left(\sum_{i=1}^k p_i^{(n)}\right) + \tau\left(\sum_{i=1}^k q r_i\right) \stackrel{(3-37), (3-45)}{=} s_n + \frac{1}{k}(1 - s_n) = s_{n+1}.$$

Moreover, for $i = 1, \dots, k$, we have

$$\tau(a_i p_i^{(n+1)}) = \tau(a_i p_i^{(n)}) + \tau(a_i q r_i) \stackrel{(3-38), (3-46)}{\leq} \delta \tau(p_i^{(n)}) + \delta \tau(q r_i) = \delta \tau(p_i^{(n+1)}).$$

This shows already that the projections satisfy (3-41) and (3-42). Next we prove that they satisfy (3-43). Note that $p_i^{(n)}$ is orthogonal to q for $i = 1, \dots, k$. Hence, we get, for each $g \in G$ and $\tau \in T_\omega(A)$, that

$$\begin{aligned}
& \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n+1)}) - p_i^{(n+1)}\|_{2,\tau}^2 \\
&= \sum_{i=1}^k \tau((\alpha_g^\omega(p_i^{(n+1)}) - p_i^{(n+1)})^2) \\
&= \sum_{i=1}^k (\tau((\alpha_g^\omega(p_i^{(n)}) - p_i^{(n)})^2) + \tau((\alpha_g^\omega(qr_i) - qr_i)^2) - 2\tau(\alpha_g^\omega(p_i^{(n)})qr_i) - 2\tau(p_i^{(n)}\alpha_g^\omega(qr_i))) \\
&\leq \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n)}) - p_i^{(n)}\|_{2,\tau}^2 + \sum_{i=1}^k \|\alpha_g^\omega(qr_i) - qr_i\|_{2,\tau}^2. \tag{3-49}
\end{aligned}$$

For $i = 1, \dots, k$, we have

$$\begin{aligned}
\|\alpha_g^\omega(qr_i) - qr_i\|_{2,\tau}^2 &\leq (\|\alpha_g^\omega(q) - q\|_{2,\tau} \|\alpha_g^\omega(r_i)\|_{2,\tau} + \|q(\alpha_g^\omega(r_i) - r_i)\|_{2,\tau})^2 \\
&\leq 2(\|\alpha_g^\omega(q) - q\|_{2,\tau}^2 + \|q(\alpha_g^\omega(r_i) - r_i)\|_{2,\tau}^2).
\end{aligned}$$

Combining this with (3-49), we find that, for $g \in F$,

$$\begin{aligned}
& \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n+1)}) - p_i^{(n+1)}\|_{2,\tau}^2 \\
&\leq \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(n)}) - p_i^{(n)}\|_{2,\tau}^2 + 2 \sum_{i=1}^k (\|\alpha_g^\omega(q) - q\|_{2,\tau}^2 + \|q(\alpha_g^\omega(r_i) - r_i)\|_{2,\tau}^2) \\
&\stackrel{(3-39),(3-48)}{\leq} \frac{4n\sqrt{\eta}}{k} + \frac{2}{k} \left\| \sum_{i=1}^k \alpha_g^\omega(p_i^{(n)}) - \sum_{i=1}^k p_i^{(n)} \right\|_{2,T_\omega(A)}^2 + 2 \sum_{i=1}^k \|q(\alpha_g^\omega(r_i) - r_i)\|_{2,\tau}^2 \\
&\stackrel{(3-40),(3-47)}{\leq} \frac{4n\sqrt{\eta}}{k} + \frac{4s_n\sqrt{\eta}}{k} + \frac{4(1-s_n)\sqrt{\eta}}{k} = \frac{4(n+1)\sqrt{\eta}}{k}.
\end{aligned}$$

Lastly, we show that the elements $p_i^{(n+1)}$ satisfy (3-44). We get, for each $g \in G$, that

$$\begin{aligned}
\sum_{i=1}^k (\alpha_g^\omega(p_i^{(n+1)}) - p_i^{(n+1)}) &= \sum_{i=1}^k (\alpha_g^\omega(p_i^{(n)}) - p_i^{(n)}) + \alpha_g^\omega\left(q\left(\sum_{j=1}^k r_j\right)\right) - q\left(\sum_{j=1}^k r_j\right) \\
&= q - \alpha_g^\omega(q) + \alpha_g^\omega\left(q\left(\sum_{j=1}^k r_j\right)\right) - q\left(\sum_{j=1}^k r_j\right) \\
&= q\left(1 - \sum_{j=1}^k r_j\right) - \alpha_g^\omega\left(q\left(1 - \sum_{j=1}^k r_j\right)\right) \\
&= q\left(\sum_{j=1}^k (\alpha_g^\omega(r_j) - r_j)\right) + q\left(1 - \sum_{j=1}^k \alpha_g^\omega(r_j)\right) - \alpha_g^\omega\left(q\left(1 - \sum_{j=1}^k r_j\right)\right) \\
&= q\left(\sum_{j=1}^k (\alpha_g^\omega(r_j) - r_j)\right) + (q - \alpha_g^\omega(q))\left(1 - \sum_{j=1}^k \alpha_g^\omega(r_j)\right).
\end{aligned}$$

Keeping in mind that q is a projection and the elements r_j are pairwise orthogonal projections, we have, for all $g \in F$ and $\tau \in T_\omega(A)$, that

$$\begin{aligned} & \left\| \sum_{i=1}^k \alpha_g^\omega(p_i^{(n+1)}) - \sum_{i=1}^k p_i^{(n+1)} \right\|_{2,\tau}^2 \\ &= \tau \left(\left(\sum_{i=1}^k \alpha_g^\omega(p_i^{(n+1)}) - \sum_{i=1}^k p_i^{(n+1)} \right)^2 \right) = \tau \left(\left(\alpha_g^\omega(p_i^{(n)} + qr_i) - \sum_{i=1}^k p_i^{(n)} + qr_i \right)^2 \right) \\ &= \tau \left(\left(\sum_{i=1}^k q(\alpha_g^\omega(r_i) - r_i) + (q - \alpha_g^\omega(q))(1 - \alpha_g^\omega(r_i)) \right)^2 \right) \\ &= \tau \left(q \left(\sum_{i,j=1}^k \alpha_g^\omega(r_i r_j) + r_i r_j - r_i \alpha_g^\omega(r_j) - \alpha_g^\omega(r_i) r_j \right) + (q - \alpha_g^\omega(q))^2 \left(1 - \sum_{j=1}^k \alpha_g^\omega(r_j) \right) \right) \\ & \qquad \qquad \qquad - 2\tau \left(\left(\sum_{j=1}^k r_j \right) q (1 - \alpha_g^\omega(q)) \left(1 - \sum_{j=1}^k \alpha_g^\omega(r_j) \right) \right) \\ &\leq \tau \left(q \left(\sum_{i,j=1}^k \alpha_g^\omega(r_i r_j) + r_i r_j - r_i \alpha_g^\omega(r_j) - \alpha_g^\omega(r_i) r_j \right) + (q - \alpha_g^\omega(q))^2 \right) \\ &\leq \tau \left(q \left(\sum_{i=1}^k \alpha_g^\omega(r_i) + r_i - r_i \alpha_g^\omega(r_i) - \alpha_g^\omega(r_i) r_i \right) + (q - \alpha_g^\omega(q))^2 \right) \\ &= \tau \left(q \left(\sum_{i=1}^k (\alpha_g^\omega(r_i) - r_i)^2 \right) + (q - \alpha_g^\omega(q))^2 \right) = \sum_{i=1}^k \|q(r_i - \alpha_g^\omega(r_i))\|_{2,\tau}^2 + \|q - \alpha_g^\omega(q)\|_{2,\tau}^2 \\ &\stackrel{(3-40),(3-47)}{\leq} 2 \left(s_n + \frac{1}{k}(1 - s_n) \right) \sqrt{\eta} = 2s_{n+1} \sqrt{\eta}. \end{aligned}$$

In the first inequality in the above computation, we used the fact that q commutes with all the elements r_i , thus the term $(\sum_{j=1}^k r_j)q(1 - \alpha_g^\omega(q))(1 - \sum_{j=1}^k \alpha_g^\omega(r_j))$ is a product of two positive elements, whose trace value must be nonnegative. In the second inequality, we used the pairwise orthogonality of the projections r_i , so the mixed terms in the double sum appearing above contribute the trace value of $-(r_i \alpha_g^\omega(r_j) + \alpha_g^\omega(r_i) r_j)$, which likewise is nonpositive.

As explained before, one has $s_N > t$. So if we start the inductive procedure with the projections $p_1^{(0)} = \dots = p_k^{(0)} = 0$, then after N steps we obtain projections $p_1^{(N)}, \dots, p_k^{(N)}$ satisfying

$$\sum_{i=1}^k \tau(p_i^{(N)}) > t \quad \text{for } \tau \in T_\omega(A).$$

Moreover, at that point we have

$$\begin{aligned} \tau(a_i p_i^{(N)}) &\stackrel{(3-42)}{\leq} \delta \tau(p_i^{(N)}) \quad \text{for } i = 1, \dots, k, \tau \in T_\omega(A), \\ \max_{g \in F} \sum_{i=1}^k \|\alpha_g^\omega(p_i^{(N)}) - p_i^{(N)}\|_{2,\tau}^2 &\stackrel{(3-43)}{\leq} \frac{4N\sqrt{\eta}}{k} \stackrel{(3-36)}{<} \varepsilon. \end{aligned}$$

We conclude that $p_1^{(N)}, \dots, p_k^{(N)}$ satisfy all the required properties. □

4. Dynamical tracial local-to-global principle

Here we prove our main technical result, namely that equivariant property Gamma implies a *tracial local-to-global principle* for actions of amenable groups. Roughly, this means that whenever a $*$ -polynomial identity has (local) approximate solutions one tracial presentation at a time, then it has (global) approximate solutions in the uniform tracial 2-norm. We begin by precisely defining these polynomial identities.

Definition 4.1 (cf. [Gardella et al. 2022, Definition 4.4]). Let G be a discrete group, and let X be a countable set of noncommutative variables. A *noncommutative G - $*$ -polynomial* in the variables X is a noncommutative $*$ -polynomial in the variables $\{g \cdot x \mid g \in G, x \in X\}$.

Let A be a C*-algebra with action $\alpha : G \curvearrowright A$. Suppose that $h(x_1, \dots, x_r)$ is a G - $*$ -polynomial in r noncommuting variables. Given a tuple $(a_1, \dots, a_r) \in A^r$, the evaluation $h(a_1, \dots, a_r)$ is computed by interpreting $g \cdot x_i$ as $\alpha_g(a_i)$ for $g \in G$ and $i = 1, \dots, r$.

The main theorem that we prove in this section is the following.

Theorem 4.2. *Let A be a σ -unital C*-algebra with $T(A)$ nonempty and compact, and with weak CPoU. Let $\alpha : G \curvearrowright A$ be an action by an amenable countable discrete group and assume it has local equivariant property Gamma with respect to bounded traces. For each $m \in \mathbb{N}$, let*

$$h_m(x_1, \dots, x_{r_m}, z_1, \dots, z_{s_m})$$

be a G - $$ -polynomial in $r_m + s_m$ noncommuting variables. Let $(a_i)_{i \in \mathbb{N}}$ be a sequence in $A^{\omega, b}$. Suppose, for every $\varepsilon > 0$, $\ell \in \mathbb{N}$, and $\tau \in \overline{T_\omega(A)}^{w*}$, there exist contractions $(y_i^\tau)_{i \in \mathbb{N}}$ in $A^{\omega, b}$ such that*

$$\|h_m(a_1, \dots, a_{r_m}, y_1^\tau, \dots, y_{s_m}^\tau)\|_{2, \tau} < \varepsilon \quad \text{for } m = 1, \dots, \ell.$$

Then there exist contractions $(y_i)_{i \in \mathbb{N}}$ in $A^{\omega, b}$ such that

$$h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m}) = 0 \quad \text{for all } m \in \mathbb{N}. \quad (4-1)$$

We reduce the complexity of the polynomials involved in the proof of this result with the following.

Lemma 4.3. *Let G be a countable discrete group. Consider sets of variables $X = \{x_i \mid i \in \mathbb{N}\}$ and $Z = \{z_i \mid i \in \mathbb{N}\}$. Assume*

$$\mathcal{P} = \{h_m(x_1, \dots, x_{r_m}, z_1, \dots, z_{s_m}) \mid m \in \mathbb{N}\}$$

is a countable set of noncommutative G - $$ -polynomials in the variables $X \cup Z$. Then there exists another set of variables $Z' = \{z'_i \mid i \in \mathbb{N}\}$ and another countable set of noncommutative G - $*$ -polynomials*

$$\mathcal{P}' = \{h'_m(x_1, \dots, x_{r'_m}, z'_1, \dots, z'_{s'_m}) \mid m \in \mathbb{N}\}$$

in the variables $X \cup Z'$ such that every G - $$ -polynomial h'_m satisfies one of the following properties:*

- (1) $h'_m(x_1, \dots, x_{r'_m}, 0, \dots, 0) = 0$ (i.e., no terms in the polynomial with variables only in X) and $h'_m(1, \dots, 1, z'_1, \dots, z'_{s'_m})$ is an ordinary $*$ -polynomial in the variables Z' ;
- (2) $h'_m(x_1, \dots, x_{r'_m}, z'_1, \dots, z'_{s'_m}) = \|h'_m(x_1, \dots, x_{r'_m}, 0, \dots, 0)\| z'_{s'_m} - h'_m(x_1, \dots, x_{r'_m}, 0, \dots, 0)$;
- (3) $h'_m(x_1, \dots, x_{r'_m}, z'_1, \dots, z'_{s'_m}) = g \cdot z'_i - z'_j$ for some $1 \leq i, j \leq s'_m$ and some $g \in G$.

Moreover we have that, for every action $\beta : G \curvearrowright B$ on any C^* -algebra, any sequence $(b_i)_{i \in \mathbb{N}}$ in B , and subset $T \subset T(B)$ the following two statements hold:

(a) There exist contractions $(y_i)_{i \in \mathbb{N}}$ in B such that

$$\|h_m(b_1, \dots, b_{r_m}, y_1, \dots, y_{s_m})\|_{2,T} = 0 \quad \text{for all } m \in \mathbb{N}$$

if and only if there exist contractions $(y'_i)_{i \in \mathbb{N}}$ in B such that

$$\|h'_m(b_1, \dots, b_{r'_m}, y'_1, \dots, y'_{s'_m})\|_{2,T} = 0 \quad \text{for all } m \in \mathbb{N}.$$

(b) For each $\varepsilon > 0$ and each $\ell \in \mathbb{N}$, there exist contractions $(y_i)_{i \in \mathbb{N}}$ in B such that

$$\|h_m(b_1, \dots, b_{r_m}, y_1, \dots, y_{s_m})\|_{2,T} < \varepsilon \quad \text{for } m = 1, \dots, \ell$$

if and only if, for each $\varepsilon' > 0$ and each $\ell' \in \mathbb{N}$, there exist contractions $(y'_i)_{i \in \mathbb{N}}$ in B such that

$$\|h'_m(b_1, \dots, b_{r'_m}, y'_1, \dots, y'_{s'_m})\|_{2,T} < \varepsilon' \quad \text{for } m = 1, \dots, \ell'.$$

Proof. Define

$$Z' = \{z_i \mid i \in \mathbb{N}\} \cup \{z_{i,g} \mid i \in \mathbb{N}, g \in G \setminus \{e\}\} \cup \{w_m \mid m \in \mathbb{N}\}.$$

For each $m \in \mathbb{N}$, we can take the G -*-polynomial h_m in the variables $X \cup Z$ and define h''_m in the variables $X \cup Z'$ by replacing every instance of a variable $g \cdot z_i$ for some $i \in \mathbb{N}$ and $g \in G \setminus \{e\}$ by $z_{i,g}$, e.g., the polynomial $g \cdot z_1 - z_2$ would be transformed into $z_{1,g} - z_2$. Next, we define a new G -*-polynomial h'''_m for each $m \in \mathbb{N}$ by setting

$$h'''_m(X \cup Z') = h''_m(X \cup Z') + \|h_m(x_1, \dots, x_{r_m}, 0, \dots, 0)\|w_m - h_m(x_1, \dots, x_{r_m}, 0, \dots, 0). \quad (4-2)$$

Set $\mathcal{P}'_1 := \{h'''_m(X \cup Z') \mid m \in \mathbb{N}\}$. All the polynomials in this set are of type (1) mentioned in the statement of this lemma. Next, define the sets of G -*-polynomials

$$\mathcal{P}'_2 := \{\|h_m(x_1, \dots, x_{r_m}, 0, \dots, 0)\|w_m - h_m(x_1, \dots, x_{r_m}, 0, \dots, 0) \mid m \in \mathbb{N}\}$$

and

$$\mathcal{P}'_3 := \{g \cdot z_i - z_{i,g} \mid i \in \mathbb{N}, g \in G \setminus \{e\}\}.$$

These sets consist of polynomials of type (2) and (3), respectively.

Consider $\mathcal{P}' = \mathcal{P}'_1 \cup \mathcal{P}'_2 \cup \mathcal{P}'_3$. The G -*-polynomials in this set are all of the right form, and we claim that this does the job. It suffices to show part (a). This is because, for a given C^* -algebra B , action $\beta : G \curvearrowright B$, sequence $(b_i)_{i \in \mathbb{N}} \in B$, and subset $T \subset T(B)$, statement (b) immediately follows from statement (a) when applied to the C^* -algebra B_ω , with action β_ω , sequence $(b_i)_{i \in \mathbb{N}}$ in $B \subset B_\omega$, and the set of limit traces on B_ω arising from sequences of traces in T .

To show part (a), fix a C^* -algebra B , an action $\beta : G \curvearrowright B$, a sequence $(b_i)_{i \in \mathbb{N}}$, and subset $T \subset T(B)$. Let $(y_i)_{i \in \mathbb{N}}$ be any sequence of contractions in B . For notational brevity, we denote these sequences by $\bar{b} = (b_i)_i$ and $\bar{y} = (y_i)_i$. Furthermore we shall also write (exclusively in this proof), for two elements $x, y \in B$, the expression “ $x =_T y$ ” as shorthand for $\|x - y\|_{2,T} = 0$.

We set $y_{i,g} = \beta_g(y_i)$ for $i \in \mathbb{N}$ and $g \in G \setminus \{e\}$ and

$$w_m = \begin{cases} 0, & h_m(b_1, \dots, b_{r_m}, 0, \dots, 0) = 0, \\ \frac{h_m(b_1, \dots, b_{r_m}, 0, \dots, 0)}{\|h_m(b_1, \dots, b_{r_m}, 0, \dots, 0)\|}, & h_m(b_1, \dots, b_{r_m}, 0, \dots, 0) \neq 0. \end{cases}$$

Then the tuple

$$\bar{z} := (y_i)_{i \in \mathbb{N}} \times (y_{i,g})_{i \in \mathbb{N}, g \in G \setminus \{e\}} \times (w_m)_{m \in \mathbb{N}}$$

represents a choice for the free variables of Z' inside B . By definition we have $p(\bar{b}, \bar{z}) = 0$ for all $p \in \mathcal{P}'_2 \cup \mathcal{P}'_3$. By definition of the polynomials h'''_m , we have

$$h'''_m(b_1, \dots, b_{r_m}, \bar{z}) = h''_m(b_1, \dots, b_{r_m}, \bar{z}) + q(\bar{b}, \bar{z}) = h''_m(b_1, \dots, b_{r_m}, \bar{z})$$

for some $*$ -polynomial $q \in \mathcal{P}'_2$. Due to the vanishing of all the $*$ -polynomials of \mathcal{P}'_3 in \bar{z} and given how the polynomial h'''_m arises from the polynomial h_m via substitution of variables, we may finally observe

$$h_m(b_1, \dots, b_{r_m}, y_1, \dots, y_{s_m}) = h''_m(b_1, \dots, b_{r_m}, \bar{z}).$$

This shows immediately that if the sequence $(y_i)_i$ satisfies

$$\|h_m(b_1, \dots, b_{r_m}, y_1, \dots, y_{s_m})\|_{2,T} = 0 \quad \text{for all } m \in \mathbb{N},$$

then we also have $\|p(\bar{b}, \bar{z})\|_{2,T} = 0$ for all $p \in \mathcal{P}'$. In particular, we get the “only if” part in (a).

Conversely, suppose that

$$\bar{z} := (y_i)_{i \in \mathbb{N}} \times (y_{i,g})_{i \in \mathbb{N}, g \in G \setminus \{e\}} \times (w_m)_{m \in \mathbb{N}}$$

is an arbitrary tuple with values in the unit ball of B representing a choice for the free variables in Z' such that $p(\bar{b}, \bar{z}) = 0$ for all $p \in \mathcal{P}'$. By doing the above computations in reverse, we can see that $p(\bar{z}) =_T 0$ for $p \in \mathcal{P}'_3$ forces $y_{i,g} =_T \beta_g(y_i)$ for all $g \in G \setminus \{e\}$. Moreover, the vanishing $p(\bar{b}, \bar{z}) =_T 0$ for $p \in \mathcal{P}'_2$ forces the equation

$$w_m = \frac{h_m(b_1, \dots, b_{r_m}, 0, \dots, 0)}{\|h_m(b_1, \dots, b_{r_m}, 0, \dots, 0)\|}$$

when $h_m(b_1, \dots, b_{r_m}, 0, \dots, 0) \neq 0$. Similar to how we argued above, this implies, for all $m \geq 1$, that

$$h'''_m(b_1, \dots, b_{r_m}, \bar{z}) = h''_m(b_1, \dots, b_{r_m}, \bar{z}) + q(\bar{b}, \bar{z}) =_T h''_m(b_1, \dots, b_{r_m}, \bar{z})$$

for some $*$ -polynomial $q \in \mathcal{P}'_2$. Given how the polynomial h'''_m arises from the polynomial h_m via substitution of variables, we may finally observe

$$h_m(b_1, \dots, b_{r_m}, y_1, \dots, y_{s_m}) =_T h''_m(b_1, \dots, b_{r_m}, \bar{z}), \quad m \geq 1.$$

This shows the “if” part of (a) and finishes the proof. □

Proof of Theorem 4.2. Equation (4-1) is equivalent to

$$\|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})\|_{2,T_{\omega(A)}} = 0 \quad \text{for all } m \in \mathbb{N}.$$

By the previous lemma, we may assume that the $*$ -polynomials h_m are all of one of the following types:

- (1) $h_m(a_1, \dots, a_{r_m}, 0, \dots, 0) = 0$ and $h_m(1_{A^{\omega,b}}, \dots, 1_{A^{\omega,b}}, z_1, \dots, z_{s_m})$ is an ordinary $*$ -polynomial.
- (2) h_m is of the form $\|h_m(x_1, \dots, x_{r_m}, 0, \dots, 0)\|z_i - h_m(x_1, \dots, x_{r_m}, 0, \dots, 0)$ for some $i \in \mathbb{N}$. Equivalently, since this doesn't change the solutions, we may assume that $h_m(a_1, \dots, a_{r_m}, z_1, \dots, z_{s_m})$ is of the form $z_i - a$ for some $i \in \mathbb{N}$ and $a \in C^*(\{\alpha_g^\omega(a_i) \mid i \in \mathbb{N}, g \in G\})$ with $\|a\| = 1$.
- (3) h_m is of the form $g \cdot z_i - z_{i'}$ for some $i, i' \in \mathbb{N}$ and $g \in G$.

By Kirchberg's ε -test, it suffices to find, for each $\varepsilon > 0$ and $\ell \in \mathbb{N}$, contractions $(y_i)_{i \in \mathbb{N}}$ in $A^{\omega,b}$ such that

$$\|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})\|_{2,\tau} \leq \varepsilon \quad \text{for } m = 1, \dots, \ell \text{ and } \tau \in T_\omega(A).$$

Choose $\varepsilon > 0$ and $\ell \in \mathbb{N}$ arbitrarily. Denote by $F \subseteq G$ the set of $g \in G$ appearing in h_m for some $m = 1, \dots, \ell$. Set

$$\delta = \frac{1}{18}\varepsilon^2. \tag{4-3}$$

Choose $t \in (0, 1)$ such that

$$1 - t < \frac{1}{2}\varepsilon^2. \tag{4-4}$$

Let $\eta > 0$ be the universal constant from Lemma 3.2 corresponding to the tuple (δ^2, t) . Since G is amenable, we can find $H \subseteq G$ such that $|gH\Delta H| < \eta|H|$ for each $g \in F$. By assumption, for each $\tau \in \overline{T_\omega(A)}^{w*}$, we can find contractions $(y_i^\tau)_{i \in \mathbb{N}} \in A^{\omega,b}$ such that

$$\|h_m(a_1, \dots, a_{r_m}, y_1^\tau, \dots, y_{s_m}^\tau)\|_{2,\tau}^2 < \frac{\varepsilon^2}{2\ell|H|} \quad \text{for } m = 1, \dots, \ell.$$

Define

$$b^\tau := \sum_{m=1}^{\ell} |h_m(a_1, \dots, a_{r_m}, y_1^\tau, \dots, y_{s_m}^\tau)|^2 \in A^{\omega,b}. \tag{4-5}$$

Then we get

$$\tau(b^\tau) = \sum_{m=1}^{\ell} \|h_m(a_1, \dots, a_{r_m}, y_1^\tau, \dots, y_{s_m}^\tau)\|_{2,\tau}^2 < \frac{\varepsilon^2}{2|H|}.$$

By continuity and compactness of $\overline{T_\omega(A)}^{w*}$, we can find finitely many tracial states $\tau_1, \dots, \tau_k \in \overline{T_\omega(A)}^{w*}$ such that

$$\frac{\varepsilon^2}{2|H|} > \sup_{\tau \in T_\omega(A)} \min_{i=1, \dots, k} \tau(b^{\tau_i}).$$

By Lemma 3.2, it follows that we can find pairwise orthogonal projections

$$p_1, \dots, p_k \in A^{\omega,b} \cap \left(\bigcup_{g \in G} \alpha_g^\omega(\{y_i^{\tau_1}, \dots, y_i^{\tau_n}, a_i \mid i \in \mathbb{N}\}) \right)'$$

such that, for $\tau \in T_\omega(A)$,

$$\tau\left(\sum_{j=1}^k p_j\right) > t, \tag{4-6}$$

$$\tau(b^{\tau_j} p_j) \leq \frac{1}{2}\varepsilon^2 \tau(p_j) \quad \text{for } j = 1, \dots, k, \tag{4-7}$$

$$\max_{g \in F} \sum_{j=1}^k \|\alpha_g(p_j) - p_j\|_{2,\tau}^2 < \delta^2. \tag{4-8}$$

Define $y_i = \sum_{j=1}^k p_j y_i^{\tau_j}$ for $i \in \mathbb{N}$. We show that, for $\tau \in T_\omega(A)$ and $m = 1, \dots, \ell$,

$$\left| \tau(|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2) - \tau\left(\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2\right) \right| < \frac{1}{2}\varepsilon^2. \quad (4-9)$$

We distinguish three cases. First, assume that $h_m(a_1, \dots, a_{r_m}, 0, \dots, 0) = 0$ and that

$$h_m(1_{A^{\omega,b}}, \dots, 1_{A^{\omega,b}}, z_1, \dots, z_{s_m})$$

is an ordinary $*$ -polynomial. In this case

$$|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2 = \sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2$$

since p_1, \dots, p_k are pairwise orthogonal projections. Second, assume h_m is of the form $z_i - a$ for some $i \in \mathbb{N}$ and some element $a \in C^*(\{\alpha_g^\omega(a_i) \mid i \in \mathbb{N}, g \in G\})$ with $\|a\| = 1$. In this case we have

$$\begin{aligned} & \left| \tau(|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2) - \tau\left(\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2\right) \right| \\ &= \left| \tau\left(\left|\sum_{j=1}^k p_j y_i^{\tau_j} - a\right|^2\right) - \tau\left(\sum_{j=1}^k p_j |y_i^{\tau_j} - a|^2\right) \right| = \left| \tau(|a|^2) - \tau\left(\sum_{j=1}^k p_j |a|^2\right) \right| \\ &\leq \tau\left(1 - \sum_{j=1}^k p_j\right) \stackrel{(4-6)}{<} 1 - t \stackrel{(4-4)}{<} \frac{1}{2}\varepsilon^2. \end{aligned}$$

Third, assume h_m is of the form $g \cdot z_i - z_{i'}$ for some $i, i' \in \mathbb{N}$ and $g \in F$. Then we have

$$\begin{aligned} & \left| \tau(|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2) - \tau\left(\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2\right) \right| \\ &= \left| \tau\left(\left|\alpha_g^\omega\left(\sum_{j=1}^k p_j y_i^{\tau_j}\right) - \sum_{j'=1}^k p_{j'} y_{i'}^{\tau_{j'}}\right|^2\right) - \tau\left(\sum_{j=1}^k p_j |\alpha_g^\omega(y_i^{\tau_j}) - y_{i'}^{\tau_j}|^2\right) \right| \\ &= \left| \tau\left(\sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(|y_i^{\tau_j}|^2)\right) - \tau\left(\sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(y_i^{\tau_j})^* \sum_{j'=1}^k p_{j'} y_{i'}^{\tau_{j'}}\right) \right. \\ & \quad \left. - \tau\left(\left(\sum_{j'=1}^k p_{j'} y_{i'}^{\tau_{j'}}\right)^* \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(y_i^{\tau_j})\right) \right| \\ &\leq \left\| \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(|y_i^{\tau_j}|^2) \right\|_{2,\tau} + 2 \left\| \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(y_i^{\tau_j}) \right\|_{2,\tau} \left\| \sum_{j=1}^k p_j y_{i'}^{\tau_j} \right\|_{2,\tau} \\ &\leq \left\| \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(|y_i^{\tau_j}|^2) \right\|_{2,\tau} + 2 \left\| \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(y_i^{\tau_j}) \right\|_{2,\tau}. \end{aligned} \quad (4-10)$$

Note that, for any $g \in G$ and any positive contractions $c_1, \dots, c_k \in A^{\omega, b}$ commuting with the p_j and $\alpha_g^\omega(p_j)$, one has that

$$\begin{aligned} \left\| \sum_{i=1}^k (\alpha_g^\omega(p_i) - p_i) c_i \right\|_{2, \tau}^2 &= \sum_{i=1}^k \tau((\alpha_g^\omega(p_i) - p_i)^2 c_i^2) + \sum_{\substack{i, j=1 \\ i \neq j}}^k \tau(c_i (\alpha_g^\omega(p_i) - p_i) (\alpha_g^\omega(p_j) - p_j) c_j) \\ &\leq \sum_{i=1}^k \left(\tau((\alpha_g^\omega(p_i) - p_i)^2) - \sum_{\substack{j=1, \dots, k \\ j \neq i}} \tau(c_i \alpha_g^\omega(p_i) p_j c_j) - \sum_{\substack{j=1, \dots, k \\ j \neq i}} \tau(c_i p_i \alpha_g^\omega(p_j) c_j) \right) \\ &\leq \sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2, \end{aligned}$$

where in the last inequality we used the tracial property and the fact that the c_i commute with the p_j and $\alpha_g^\omega(p_j)$ to show that the last two terms can be rewritten as the negative of the trace of positive elements.

In particular we have, for $g \in F$, that

$$\left\| \sum_{j=1}^k (\alpha_g^\omega(p_j) - p_j) \alpha_g^\omega(|y_i^{\tau_j}|^2) \right\|_{2, \tau} \leq \sqrt{\sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2} \stackrel{(4-8)}{<} \delta \stackrel{(4-3)}{=} \frac{1}{18} \varepsilon^2. \tag{4-11}$$

For $j = 1, \dots, k$, we have that $\alpha_g^\omega(y_i^{\tau_j})$ is a contraction that can be written as a linear combination of positive contractions commuting with the p_i and $\alpha_g^\omega(p_i)$. An application of the triangle inequality yields

$$\left\| \sum_{i=1}^k (\alpha_g^\omega(p_i) - p_i) \alpha_g^\omega(y_i^{\tau_j}) \right\|_{2, \tau} \leq 4 \sqrt{\sum_{i=1}^k \|\alpha_g^\omega(p_i) - p_i\|_{2, \tau}^2} \stackrel{(4-8)}{<} 4\delta \stackrel{(4-3)}{=} \frac{2}{9} \varepsilon^2. \tag{4-12}$$

Combining (4-10) with (4-11) and (4-12), we get

$$\left| \tau(|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2) - \tau \left(\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2 \right) \right| < \frac{9}{18} \varepsilon^2 = \frac{1}{2} \varepsilon^2.$$

Thus, we have indeed shown that (4-9) holds for all $\tau \in T_\omega(A)$ and $m = 1, \dots, \ell$. From (4-5) we see that

$$\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2 \leq \sum_{j=1}^k p_j b^{\tau_j} \quad \text{for } m = 1, \dots, \ell. \tag{4-13}$$

As a consequence, for $\tau \in T_\omega(A)$ and $m = 1, \dots, \ell$, we get

$$\begin{aligned} \|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})\|_{2, \tau}^2 &= \tau(|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})|^2) \\ &\stackrel{(4-9)}{<} \tau \left(\sum_{j=1}^k p_j |h_m(a_1, \dots, a_{r_m}, y_1^{\tau_j}, \dots, y_{s_m}^{\tau_j})|^2 \right) + \frac{1}{2} \varepsilon^2 \\ &\stackrel{(4-13)}{\leq} \sum_{j=1}^k \tau(p_j b^{\tau_j}) + \frac{1}{2} \varepsilon^2 \stackrel{(4-7)}{\leq} \sum_{j=1}^k \frac{1}{2} \varepsilon^2 \tau(p_j) + \frac{1}{2} \varepsilon^2 \leq \frac{1}{2} \varepsilon^2 + \frac{1}{2} \varepsilon^2 = \varepsilon^2. \quad \square \end{aligned}$$

The next theorem gives an alternative formulation of the tracial local-to-global principle that is convenient to use in certain applications. Before we state it, we introduce some notation.

Notation 4.4. Let A be a C*-algebra with an action $\alpha : G \curvearrowright A$ of a countable discrete group. Given a tracial state $\tau \in T(A)$, denote by $\pi_\tau : A \rightarrow \mathcal{B}(H_\tau)$ the corresponding GNS representation. Then we define the representation

$$\pi_\tau^\alpha : A \rightarrow \mathcal{B}(\ell^2(G, H_\tau)), \quad \pi_\tau^\alpha(x)(\xi)(h) = \pi_\tau(\alpha_h^{-1}(x))\xi(h).$$

The left-regular representation $\lambda : G \rightarrow \mathcal{U}(\ell^2(G, H_\tau))$, defined by $(g \cdot \xi)(h) = \xi(g^{-1}h)$ for $\xi \in \ell^2(G, H_\tau)$ and $g, h \in G$, implements the action α on $\pi_\tau^\alpha(A)$, so we get a continuous extension of the action $\alpha : G \curvearrowright \pi_\tau^\alpha(A)''$ on the weak closure.

Notice that $\pi_\tau^\alpha(A)'' \subseteq \prod_{g \in G} \pi_\tau(A)''$. The trace τ on A extends to a faithful normal trace on $\pi_\tau(A)''$ and, by composition with the natural quotient map $q_g : \prod_{g \in G} \pi_\tau(A)'' \rightarrow \pi_\tau(A)''$ onto the summand with index $g \in G$, also to a normal trace on $\prod_{g \in G} \pi_\tau(A)''$, which we will denote by $\tilde{\tau}_g$. Notice that $\tilde{\tau}_g \circ \pi_\tau^\alpha = \tau \circ \alpha_g^{-1}$. Let $(c_g)_{g \in G}$ be a sequence in $(0, 1)$ such that $\sum_{g \in G} c_g = 1$. Then $\tilde{\tau} := \sum_{g \in G} c_g \tilde{\tau}_g$ defines a faithful normal tracial state on $\prod_{g \in G} \pi_\tau(A)''$ and hence also on the subalgebra $\pi_\tau^\alpha(A)''$. In this way we can form the tracial von Neumann algebra ultrapower $(\pi_\tau^\alpha(A)')^\omega$. Note that, on bounded subsets of $\prod_{g \in G} \pi_\tau(A)''$, the strong operator topology is induced by the norm $\|\cdot\|_{2, \tilde{\tau}}$, or equivalently by the seminorms $\{\|\cdot\|_{2, \tilde{\tau}_g} \mid g \in G\}$. Since $\pi_\tau^\alpha(A)''$ is a von Neumann subalgebra, it follows that on bounded subsets its strong operator topology is also induced by (the restrictions of) these (semi)norms.

Remark 4.5. With the above notation and terminology, the condition in [Theorem 4.2](#) that requires, for every $\varepsilon > 0$, $\ell \in \mathbb{N}$, and $\tau \in \overline{T_\omega(A)}^{w*}$, the existence of contractions $(y_i)_{i \in \mathbb{N}}$ in $A^{\omega, b}$ such that

$$\|h_m(a_1, \dots, a_{r_m}, y_1, \dots, y_{s_m})\|_{2, \tau} < \varepsilon \quad \text{for } m = 1, \dots, \ell$$

is equivalent to the following statement (by Kaplansky's density theorem): for every $\varepsilon > 0$, $\ell \in \mathbb{N}$, and $\tau \in \overline{T_\omega(A)}^{w*}$, there exist contractions $(y'_i)_{i \in \mathbb{N}}$ in $\pi_\tau^{\alpha^\omega}(A^{\omega, b})''$ such that

$$\|h_m(a_1, \dots, a_{r_m}, y'_1, \dots, y'_{s_m})\|_{2, \tilde{\tau}} < \varepsilon \quad \text{for } m = 1, \dots, \ell.$$

Making use of the tracial von Neumann algebra ultrapowers, this is also equivalent to the following statement: for every $\tau \in \overline{T_\omega(A)}^{w*}$, there are contractions $(y''_i)_{i \in \mathbb{N}}$ in $(\pi_\tau^{\alpha^\omega}(A^{\omega, b})'')^\kappa$ such that $h_m(a_1, \dots, a_{r_m}, y''_1, \dots, y''_s) = 0$ for every $m \in \mathbb{N}$.

Theorem 4.6. Let A be a σ -unital C*-algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action by an amenable countable discrete group G and assume it has local equivariant property Gamma with respect to bounded traces. Let ω and κ be two free ultrafilters on \mathbb{N} . Let $\delta : G \curvearrowright D$ be an action on a separable C*-algebra and let $B \subset D$ be a separable, δ -invariant C*-subalgebra. Suppose $\varphi : (B, \delta) \rightarrow (A^{\omega, b}, \alpha^\omega)$ is an equivariant *-homomorphism. Then the following are equivalent:

- (1) For every $\tau \in \overline{T_\omega(A)}^{w*}$, there exists an equivariant *-homomorphism

$$\varphi^\tau : (D, \delta) \rightarrow ((\pi_\tau^{\alpha^\omega}(A^{\omega, b})'')^\kappa, (\alpha^\omega)^\kappa)$$

such that $\varphi^\tau|_B = \pi_\tau^{\alpha^\omega} \circ \varphi$.

- (2) There is an equivariant *-homomorphism $\bar{\varphi} : (D, \delta) \rightarrow (A^{\omega, b}, \alpha^\omega)$ with $\bar{\varphi}|_B = \varphi$.

Proof. It is clear that (2) implies (1). To prove the other implication, take a countable dense $\mathbb{Q}[i]$ -*-subalgebra $C \subset D$ such that it is δ -invariant and such that $C \cap B$ is also dense in B . By inductively enlarging C we may in addition assume that, for each contraction $x \in C$, one has $1 - \sqrt{1 - x^*x} \in C$. Let \mathcal{P} denote the countable family of G -*-polynomials with coefficients in $A^{\omega, b}$ in the variables $\{X_c\}_{c \in C}$ encoding all relations in C :

- $g \cdot X_c - X_{\delta_g(c)}$ for all $c \in C$ and $g \in G$,
- $\lambda X_c + X_{c'} - X_{\lambda c + c'}$ for all $c, c' \in C$ and $\lambda \in \mathbb{Q}[i]$,
- $X_c X_{c'} - X_{cc'}$ for $c, c' \in C$,
- $X_c^* - X_{c^*}$ for $c \in C$,
- $\varphi(b) - X_b$ for $b \in B \cap C$.

It follows from (1) that, for every $\tau \in \overline{T_\omega(A)}^{w^*}$, the equations in \mathcal{P} have exact solutions in $(\pi_\tau^{\alpha^\omega}(A^{\omega, b})'')^\kappa$. By Remark 4.5 this means precisely that all conditions to apply Theorem 4.2 are fulfilled, and we can find exact solutions to all equations in \mathcal{P} in $A^{\omega, b}$. This is equivalent to the existence of a $\mathbb{Q}[i]$ -linear, *-preserving, multiplicative, equivariant map $\bar{\varphi} : C \rightarrow A^{\omega, b}$ with $\bar{\varphi}|_{B \cap C} = \varphi|_{B \cap C}$. We observe that $\bar{\varphi}$ is contractive. Indeed, if $x \in C$ is a contraction, then $y = 1 - \sqrt{1 - x^*x}$ is a self-adjoint element also belonging to C , which satisfies

$$x^*x + y^2 - 2y = x^*x + (y - 1)^2 - 1 = 0.$$

Hence

$$\bar{\varphi}(x)^* \bar{\varphi}(x) + \bar{\varphi}(y)^2 - 2\bar{\varphi}(y) = 0,$$

or equivalently,

$$\bar{\varphi}(x)^* \bar{\varphi}(x) + (1 - \bar{\varphi}(y))^2 = 1.$$

We see that $\bar{\varphi}(x)^* \bar{\varphi}(x)$ is a contraction, and hence $\bar{\varphi}(x)$ is as well. In conclusion, φ extends to an equivariant *-homomorphism $\bar{\varphi} : (D, \delta) \rightarrow (A^{\omega, b}, \alpha^\omega)$ with $\bar{\varphi}_B = \varphi$. \square

In many cases of interest we get the following corollary from Proposition 2.4, which directly generalizes and recovers the technical machinery related to uniform property Gamma from the nondynamical setting; see [Castillejos et al. 2021b, Lemma 4.1]. We note that, upon close inspection of our proof so far, this particular corollary can be obtained based on [Castillejos et al. 2021b, Lemma 3.6] without relying on the preprint [Carrion et al. 2023a].

Corollary 4.7. *Let A be a separable, simple, nuclear C^* -algebra with $T(A)$ nonempty and compact, and such that $T^+(A) = \mathbb{R}^{>0}T(A)$. Let $\alpha : G \curvearrowright A$ be an action by a countable amenable discrete group that has equivariant property Gamma. Then α satisfies the conclusion of Theorems 4.2 and 4.6.*

Remark 4.8. For potential subsequent applications of the theory in this article, let us reflect on how we ended up with the main result of this section. It is worthwhile to note that the amenability of the group G is used (in the proof of Theorem 4.2) through the Følner condition exclusively for the purpose of having access to a finite set $H \Subset G$ that satisfies the conclusion of Lemma 3.4. At no other point in the whole chain of argument is it necessary to know that H is actually a set that is almost invariant with respect

to F , or anything else about H for that matter. This culminates in the following more explicit observation, which we suspect may be, at some point, interesting to consider for certain actions of nonamenable groups.

Let A be a σ -unital C*-algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action of a countable discrete group. Suppose that, for all $\varepsilon > 0$ and $F \in G$, there exists a finite subset $H \in G$ that satisfies the same conclusion as in [Lemma 3.4](#). If α has local equivariant property Gamma with respect to bounded traces, then α also satisfies the conclusion of [Theorems 4.2](#) and [4.6](#).

5. Equivariant Jiang–Su stability

In this section we use the dynamical tracial local-to-global principle derived in the previous section combined with von Neumann algebraic results to conclude that, for actions of countable amenable groups on separable, simple, nuclear, finite, \mathcal{Z} -stable C*-algebras, equivariant property Gamma implies equivariant \mathcal{Z} -stability. Although one can get by with known variations of Ocneanu’s theorem [\[1985\]](#), for many applications treated in this section, our most general results here need a more general McDuff-type theorem for actions of amenable groups on von Neumann algebras, which we import from our recent work [\[Szabó and Wouters 2024\]](#).

We begin by reducing the problem of equivariant \mathcal{Z} -stability to the existence of so-called tracially large c.p.c. order-zero maps $M_n \rightarrow F_\omega(A)^{\tilde{\alpha}_\omega}$ for $n \geq 2$. The argument is well known to experts and traces back to the work of Matui and Sato [\[2012\]](#). It makes use of an equivariant version of their property (SI), for which the general framework needed here was developed in [\[Szabó 2021b\]](#).

Definition 5.1 [\[Szabó 2021b, Definition 2.5; Castillejos et al. 2023, Definition 1.3\]](#). Let A be a separable, simple C*-algebra with $T^+(A) \neq \emptyset$.

- (1) We say that a positive contraction $f \in F_\omega(A)$ is *tracially supported at 1* if the following holds: for every nonzero positive element $a \in \mathcal{P}(A)$, there exists a constant $\kappa = \kappa(f, a) > 0$ such that, for every $\tau \in \tilde{T}_\omega(A)$ with $0 < \tau(a) < \infty$, one has $\inf_{k \in \mathbb{N}} \tau_a(f^k) \geq \kappa \tau(a)$.
- (2) A positive element $e \in F_\omega(A)$ is called *tracially null* if $e \in \mathcal{J}_A$ in the sense of [Definition 1.11](#).
- (3) Let B be a unital C*-algebra. A c.p.c. order-zero map $\phi : B \rightarrow F_\omega(A)$ is called *tracially large* if $\tau_a \circ \phi(1) = \tau(a)$ for all nonzero positive elements $a \in \mathcal{P}(A)$ and $\tau \in \tilde{T}_\omega(A)$ with $\tau(a) < \infty$.

Remark 5.2. It follows from [\[Szabó 2021b, Proposition 2.4\]](#) that any of the conditions above hold for all nonzero positive elements $a \in \mathcal{P}(A)$ if and only if they hold for just a single such element, so in practice it suffices to check them for a single $a \in \mathcal{P}(A)_+ \setminus \{0\}$.

Definition 5.3 [\[Szabó 2021b, Definition 2.7\]](#). Let A be a separable, simple C*-algebra with $T^+(A) \neq \emptyset$ and an action $\alpha : G \curvearrowright A$ of a countable discrete group. We say that α has *equivariant property (SI)* if the following holds:

Whenever $e, f \in F_\omega(A)^{\tilde{\alpha}_\omega}$ are two positive contractions such that f is tracially supported at 1 and e is tracially null, there exists a contraction $s \in F_\omega(A)^{\tilde{\alpha}_\omega}$ such that $fs = s$ and $s^*s = e$.

It follows from [Szabó 2021b, Corollary 4.3] that all actions of amenable groups on nonelementary, separable, simple, nuclear C^* -algebras with strict comparison have property (SI). Combined with the following theorem, it gives a powerful sufficient criterion for equivariant Jiang–Su stability. This is not new to the experts but has never been formally stated in this generality before, so we shall give the proof for the reader’s convenience.

Theorem 5.4. *Let A be a separable, simple C^* -algebra with $T^+(A) \neq \emptyset$, and let $\alpha : G \curvearrowright A$ be an action of a countable discrete group with equivariant property (SI). Then α is equivariantly \mathcal{Z} -stable if and only if, for every $n \in \mathbb{N}$, there exists a unital $*$ -homomorphism $M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega}$.*

Proof. Since the “only if” part can be obtained with the standard argument sketched in Remark 2.2, we prove the “if” part. Given $n \in \mathbb{N}$, let $\phi' : M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega}$ be a unital $*$ -homomorphism. By Proposition 1.18, we can find a tracially large c.p.c. order-zero map $\phi : M_n \rightarrow F_\omega(A)^{\tilde{\alpha}^\omega}$ that lifts ϕ' . Set $e := 1_{F_\omega(A)} - \phi(1)$, and set $f := \phi(e_{1,1})$. Both are positive contractions in $F_\omega(A)^{\tilde{\alpha}^\omega}$. Since ϕ is tracially large, we can conclude immediately that e is tracially null. Since $e_{1,1}$ is a projection, it follows that $\phi(e_{1,1}) - \phi(e_{1,1})^m$ is tracially null for any $m \geq 1$. Moreover, for every $\tau \in \tilde{T}_\omega(A)$ and $a \in \mathcal{P}(A)_+ \setminus \{0\}$ such that $\tau(a) < \infty$, the functional $\tau_a \circ \phi$ is a bounded trace and therefore a multiple of the unique tracial state on M_n . So, for every $k \in \mathbb{N}$, we have

$$\tau_a(f^k) = \tau_a(\phi(e_{1,1})^k) = \tau_a(\phi(e_{1,1})) = \frac{1}{n} \tau(a).$$

This proves that f is tracially supported at 1.

Since α has equivariant property (SI), we can find a contraction $s \in F_\omega(A)^{\tilde{\alpha}^\omega}$ such that $fs = s$ and $s^*s = e$. By [Rørddam and Winter 2010, Theorem 5.1], this implies the existence of a unital $*$ -homomorphism from the dimension drop algebra $Z_{n,n+1}$ into $F_\omega(A)^{\tilde{\alpha}^\omega}$. As \mathcal{Z} is an inductive limit of those algebras, we find a unital $*$ -homomorphism $\mathcal{Z} \rightarrow F_\omega(A)^{\tilde{\alpha}^\omega}$.¹⁵ This implies equivariant \mathcal{Z} -stability by [Szabó 2018b, Corollary 3.8]. □

We shall now prove that, for actions of amenable groups on simple nuclear \mathcal{Z} -stable C^* -algebras, equivariant uniform property Gamma is equivalent to equivariant \mathcal{Z} -stability. We end up giving two separate arguments to prove this result in two cases. Firstly, we prove this result for actions on C^* -algebras that have a compact nonempty tracial state space and no unbounded traces, for which it is sufficient to appeal to Corollary 4.7. Secondly, we prove the result in full generality, but this requires the full power of our theory based on the results from [Carrión et al. 2023a].

Let us proceed in the first case.

Theorem 5.5. *Let A be a separable, nuclear, simple \mathcal{Z} -stable C^* -algebra with $T(A)$ nonempty and compact, and such that $T^+(A) = \mathbb{R}^{>0}T(A)$. Let $\alpha : G \curvearrowright A$ be an action of a countable discrete amenable group. If α has equivariant property Gamma, then α is equivariantly \mathcal{Z} -stable.*

Proof. By Theorem 5.4, it suffices to construct a unital $*$ -homomorphism $M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega}$ for $n \geq 2$. We appeal to Corollary 4.7 and hence know that α obeys the conclusion of Theorem 4.2.

¹⁵This is a standard reindexation trick. Alternatively one can deduce this for example by a combination of Corollary 3.9 and Lemma 4.2 in [Barlak and Szabó 2016].

Let $(a_k)_{k \in \mathbb{N}}$ be a dense sequence in A . Then the existence of such a desired $*$ -homomorphism is equivalent to the existence of elements $e_{1,1}, e_{2,1}, \dots, e_{n,1} \in A^\omega$ satisfying the equations

$$e_{i,1}e_{j,1} = 0, \quad e_{i,1}^*e_{j,1} = \delta_{ij}e_{1,1}, \quad e_{1,1}^2 = e_{1,1}, \quad \alpha_g^\omega(e_{i,1}) = e_{i,1}, \quad \text{and} \quad a_k e_{i,1} - e_{i,1} a_k = 0$$

for all $g \in G$, $i, j = 2, \dots, n$, and $k \in \mathbb{N}$. By [Theorem 4.2](#), it suffices to show that, for each $\varepsilon > 0$, finite subset $F \subseteq G$, $m \in \mathbb{N}$, and every tracial state $\tau \in T(A)$, there exists contractions $f_{1,1}, f_{2,1}, \dots, f_{n,1} \in A$ satisfying

$$\begin{aligned} \|f_{i,1}f_{j,1}\|_{2,\tau} < \varepsilon, \quad \|f_{i,1}^*f_{j,1} - \delta_{ij}f_{1,1}\|_{2,\tau} < \varepsilon, \quad \|f_{1,1}^2 - f_{1,1}\|_{2,\tau} < \varepsilon, \\ \|\alpha_g(f_{i,1}) - f_{i,1}\|_{2,\tau} < \varepsilon, \quad \text{and} \quad \|a_k f_{i,1} - f_{i,1} a_k\|_{2,\tau} < \varepsilon \end{aligned}$$

for all $g \in F$, $i, j = 2, \dots, n$, and $1 \leq k \leq m$. This is the case, however, if and only if, for every $\tau \in T(A)$, there exists a unital equivariant $*$ -homomorphism $M_n \rightarrow ((\pi_\tau^\alpha(A)')^\omega \cap A')^{\alpha^\omega}$. Since A is nuclear, the tracial von Neumann algebra $N_\tau := \pi_\tau^\alpha(A)''$ is injective; see for example [\[Blackadar 2006, Theorem IV.2.2.13\]](#). Since it does not have any direct summand of type I, it follows from Connes' theorem [\[1976\]](#) that $N_\tau \bar{\otimes} \mathcal{R} \cong N_\tau$. Hence the claim follows directly from [\[Szabó and Wouters 2024, Theorem A\]](#). \square

Next we proceed in the second and more general case.

Lemma 5.6. *Let A be a σ -unital C^* -algebra with $T(A)$ nonempty and compact. Let $\alpha : G \curvearrowright A$ be an action by a countable discrete amenable group G and assume it has local equivariant property Gamma with respect to bounded traces. Then the following are equivalent:*

- (1) For all $\tau \in T(A)$, there exists a unital $*$ -homomorphism $\varphi^\tau : M_n \rightarrow ((\pi_\tau^\alpha(A)')^\omega)^{\alpha^\omega}$.
- (2) There exists a unital $*$ -homomorphism $\varphi : M_n \rightarrow (A^{\omega,b})^{\alpha^\omega}$.

Proof. For each $\tau \in T(A)$, the map π_τ^α induces a unital $*$ -homomorphism

$$(A^{\omega,b})^{\alpha^\omega} \rightarrow ((\pi_\tau^\alpha(A)')^\omega)^{\alpha^\omega},$$

so (2) implies (1). We use the fact that α has local equivariant property Gamma to prove the other implication. By [Theorem 4.6](#) the following are equivalent:

- (a) For all $\tau \in \overline{T_\omega(A)}^{w*}$, there exists a unital equivariant $*$ -homomorphism

$$\varphi^\tau : (M_n, \text{id}_{M_n}) \rightarrow ((\pi_\tau^{\alpha^\omega}(A^{\omega,b})')^\kappa, (\alpha^\omega)^\kappa).$$

- (b) There exists a unital equivariant $*$ -homomorphism $\varphi : (M_n, \text{id}_{M_n}) \rightarrow (A^{\omega,b}, \alpha^\omega)$.

Statement (b) is equivalent to statement (2) above. Hence, in order to prove the implication it suffices to prove that statement (1) implies (a). Take $\tau \in \overline{T_\omega(A)}^{w*}$ and denote its restriction to A by σ . The canonical map $A \rightarrow A^{\omega,b}$ induces an equivariant $*$ -homomorphism $(\pi_\sigma^\alpha(A), \alpha) \rightarrow (\pi_\tau^{\alpha^\omega}(A^{\omega,b})', \alpha^\omega)$ that is continuous on the unit ball with respect to the strong operator topology. Hence, it can be extended to a unital equivariant $*$ -homomorphism $\pi_\sigma^\alpha(A)'' \rightarrow \pi_\tau^{\alpha^\omega}(A^{\omega,b})''$. Combining this with (1), this means we can find a unital $*$ -homomorphism $M_n \rightarrow ((\pi_\tau^{\alpha^\omega}(A^{\omega,b})')^\kappa)^{(\alpha^\omega)^\kappa}$. This ends the proof. \square

Theorem 5.7. *Let A be a separable, simple, nuclear \mathcal{Z} -stable C^* -algebra such that $T^+(A) \neq \emptyset$. Let $\alpha : G \curvearrowright A$ be an action of a countable discrete amenable group. If α has equivariant property Gamma, then α is equivariantly \mathcal{Z} -stable.*

Proof. Combining [Szabó 2021b, Corollary 4.3] with Theorem 5.4, we see that, given $n \geq 2$, it suffices to construct a unital $*$ -homomorphism $M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega}$. For convenience, let us specify a (possibly different) free ultrafilter κ on \mathbb{N} . It suffices to show that we can construct a unital $*$ -homomorphism $M_n \rightarrow ((A^\omega \cap A')^{\kappa, b})^{(\alpha^\omega)^\kappa}$, as a reindexation trick will then yield the required unital $*$ -homomorphism $M_n \rightarrow (A^\omega \cap A')^{\alpha^\omega}$. By Theorem 2.6, we conclude that $\alpha^\omega = G \curvearrowright A^\omega \cap A'$ has local equivariant property Gamma with respect to bounded traces. Thus, by Lemma 5.6, it suffices to prove that, for all $\tau \in T(A^\omega \cap A')$, there exists a unital $*$ -homomorphism $\varphi^\tau : M_n \rightarrow ((\pi_\tau^{\alpha^\omega}(A^\omega \cap A')'')^\kappa)^{(\alpha^\omega)^\kappa}$. We note that the C^* -algebra A has uniform property Gamma. Using the same trick as in the proof of Proposition 2.5, we conclude that $A^\omega \cap A' \cong B^\omega \cap B'$ for a hereditary subalgebra $B \subset A \otimes \mathbb{K}$ such that $T^+(B) = \mathbb{R}^{>0}T(B)$ and $T(B)$ is compact. Using [Castillejos et al. 2022, Theorem 4.6], we can hence conclude that there exists a unital $*$ -homomorphism $M_2 \rightarrow A^\omega \cap A'$. By a standard reindexation trick, we can argue that such a $*$ -homomorphism can be chosen to additionally commute with any specified separable subset of $A^\omega \cap A'$.

Fix $\tau \in T(A^\omega \cap A')$. We show first that $N_\tau := \pi_\tau^{\alpha^\omega}(A^\omega \cap A')''$ contains a $\|\cdot\|_{2, \tau}$ -separable, α^ω -invariant von Neumann subalgebra that tensorially absorbs the hyperfinite II_1 -factor. By the aforementioned property of A , we can find a unital embedding $\phi_1 : M_2 \rightarrow \pi_\tau^{\alpha^\omega}(A^\omega \cap A') \subseteq N_\tau$. Set

$$B_1 := C^*\left(\bigcup_{g \in G} \alpha_g^\omega(\phi_1(M_2))\right).$$

Using that B_1 is a separable subquotient of $A^\omega \cap A'$, we again use the aforementioned property of A and find a unital embedding

$$\phi_2 : M_2 \rightarrow \pi_\tau^{\alpha^\omega}(A^\omega \cap A') \cap B_1' \subseteq N_\tau \cap B_1'.$$

Set

$$B_2 := C^*\left(B_1 \cup \bigcup_{g \in G} \alpha_g^\omega(\phi_2(M_2))\right).$$

Carry on with this procedure inductively, i.e., given the C^* -algebra $B_i \subset \pi_\tau^{\alpha^\omega}(A^\omega \cap A')$, find a unital $*$ -homomorphism $\phi_{i+1} : M_2 \rightarrow \pi_\tau^{\alpha^\omega}(A^\omega \cap A') \cap B_i'$ and set

$$B_{i+1} := C^*\left(B_i \cup \bigcup_{g \in G} \alpha_g^\omega(\phi_{i+1}(M_2))\right).$$

Define $\mathcal{B} := \overline{\bigcup_{i \in \mathbb{N}} B_i}^{\|\cdot\|_{2, \tau}} \subset N_\tau$. Then \mathcal{B} is a $\|\cdot\|_{2, \tau}$ -separable, α^ω -invariant von Neumann subalgebra of N_τ such that additionally $\mathcal{B} \cong \mathcal{B} \bar{\otimes} \mathcal{R}$ by [Szabó and Wouters 2024, Corollary 3.8] because it satisfies the McDuff-type criterion (existence of a unital $*$ -homomorphism $\mathcal{R} \rightarrow \mathcal{B}_\omega$) by construction. Denote the restriction of α^ω to \mathcal{B} by β . By [Szabó and Wouters 2024, Theorem A], it follows that β is cocycle conjugate to $\beta \otimes \text{id}_\mathcal{R}$. In particular, we can find a unital $*$ -homomorphism

$$M_n \rightarrow (B^\kappa)^{\beta^\kappa} \subset ((\pi_\tau^{\alpha^\omega}(A^\omega \cap A')'')^\kappa)^{(\alpha^\omega)^\kappa}.$$

□

Acknowledgements

We would like to express our sincere gratitude to the anonymous reviewers for their valuable comments that helped improve this article.

References

- [Anantharaman and Popa 2014] C. Anantharaman and S. Popa, “An introduction to II_1 factors”, preprint, 2014, available at <https://www.math.ucla.edu/~popa/Books/IIunV15.pdf>.
- [Ando and Haagerup 2014] H. Ando and U. Haagerup, “Ultraproducts of von Neumann algebras”, *J. Funct. Anal.* **266**:12 (2014), 6842–6913. [MR](#) [Zbl](#)
- [Antoine et al. 2022] R. Antoine, F. Perera, L. Robert, and H. Thiel, “ C^* -algebras of stable rank one and their Cuntz semigroups”, *Duke Math. J.* **171**:1 (2022), 33–99. [MR](#) [Zbl](#)
- [Barkl and Szabó 2016] S. Barkl and G. Szabó, “Sequentially split $*$ -homomorphisms between C^* -algebras”, *Int. J. Math.* **27**:13 (2016), art. id. 1650105. [MR](#) [Zbl](#)
- [Blackadar 2006] B. Blackadar, *Operator algebras: theory of C^* -algebras and von Neumann algebras*, *Enycl. Math. Sci.* **122**, Springer, 2006. [MR](#) [Zbl](#)
- [Blackadar and Cuntz 1982] B. E. Blackadar and J. Cuntz, “The structure of stable algebraically simple C^* -algebras”, *Amer. J. Math.* **104**:4 (1982), 813–822. [MR](#) [Zbl](#)
- [Blanchard and Kirchberg 2004] E. Blanchard and E. Kirchberg, “Non-simple purely infinite C^* -algebras: the Hausdorff case”, *J. Funct. Anal.* **207**:2 (2004), 461–513. [MR](#) [Zbl](#)
- [Bosa et al. 2019] J. Bosa, N. P. Brown, Y. Sato, A. Tikuisis, S. White, and W. Winter, *Covering dimension of C^* -algebras and 2-coloured classification*, *Mem. Amer. Math. Soc.* **1233**, Amer. Math. Soc., Providence, RI, 2019. [MR](#) [Zbl](#)
- [Carrión et al. 2023a] J. R. Carrión, J. Castillejos, S. Evington, J. Gabe, C. Schafhauser, A. Tikuisis, and S. White, “Tracially complete C^* -algebras”, preprint, 2023. [arXiv 2310.20594](#)
- [Carrión et al. 2023b] J. R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis, and S. White, “Classifying $*$ -homomorphisms, I: Unital simple nuclear C^* -algebras”, preprint, 2023. [arXiv 2307.06480](#)
- [Castillejos and Evington 2020] J. Castillejos and S. Evington, “Nuclear dimension of simple stably projectionless C^* -algebras”, *Anal. PDE* **13**:7 (2020), 2205–2240. [MR](#) [Zbl](#)
- [Castillejos and Evington 2021] J. Castillejos and S. Evington, “Stabilising uniform property Γ ”, *Proc. Amer. Math. Soc.* **149**:11 (2021), 4725–4737. [MR](#) [Zbl](#)
- [Castillejos et al. 2021a] J. Castillejos, S. Evington, A. Tikuisis, and S. White, “Classifying maps into uniform tracial sequence algebras”, *Münster J. Math.* **14**:2 (2021), 265–281. [MR](#) [Zbl](#)
- [Castillejos et al. 2021b] J. Castillejos, S. Evington, A. Tikuisis, S. White, and W. Winter, “Nuclear dimension of simple C^* -algebras”, *Invent. Math.* **224**:1 (2021), 245–290. [MR](#) [Zbl](#)
- [Castillejos et al. 2022] J. Castillejos, S. Evington, A. Tikuisis, and S. White, “Uniform property Γ ”, *Int. Math. Res. Not.* **2022**:13 (2022), 9864–9908. [MR](#) [Zbl](#)
- [Castillejos et al. 2023] J. Castillejos, K. Li, and G. Szabó, “On tracial \mathcal{Z} -stability of simple non-unital C^* -algebras”, *Canad. J. Math.* (online publication April 2023).
- [Connes 1976] A. Connes, “Classification of injective factors: cases II_1 , II_∞ , III_λ , $\lambda \neq 1$ ”, *Ann. of Math. (2)* **104**:1 (1976), 73–115. [MR](#) [Zbl](#)
- [Connes 1977] A. Connes, “Periodic automorphisms of the hyperfinite factor of type II_1 ”, *Acta Sci. Math. (Szeged)* **39**:1-2 (1977), 39–66. [MR](#) [Zbl](#)
- [Elliott 2010] G. A. Elliott, “Towards a theory of classification”, *Adv. Math.* **223**:1 (2010), 30–48. [MR](#) [Zbl](#)
- [Elliott et al. 2011] G. A. Elliott, L. Robert, and L. Santiago, “The cone of lower semicontinuous traces on a C^* -algebra”, *Amer. J. Math.* **133**:4 (2011), 969–1005. [MR](#) [Zbl](#)
- [Elliott et al. 2020] G. A. Elliott, G. Gong, H. Lin, and Z. Niu, “The classification of simple separable KK -contractible C^* -algebras with finite nuclear dimension”, *J. Geom. Phys.* **158** (2020), art. id. 103861. [MR](#) [Zbl](#)

- [Elliott et al. 2024] G. A. Elliott, G. Gong, H. Lin, and Z. Niu, “On the classification of simple C^* -algebras with finite decomposition rank, II”, *J. Noncommut. Geom.* (online publication June 2024).
- [Evans and Kishimoto 1997] D. E. Evans and A. Kishimoto, “Trace scaling automorphisms of certain stable AF algebras”, *Hokkaido Math. J.* **26**:1 (1997), 211–224. [MR](#) [Zbl](#)
- [Fu et al. 2022] X. Fu, K. Li, and H. Lin, “Tracial approximate divisibility and stable rank one”, *J. Lond. Math. Soc.* (2) **106**:4 (2022), 3008–3042. [MR](#) [Zbl](#)
- [Gabe and Szabó 2022] J. Gabe and G. Szabó, “The stable uniqueness theorem for equivariant Kasparov theory”, 2022. To appear in *Amer. J. Math.* [arXiv 2202.09809](#)
- [Gabe and Szabó 2024] J. Gabe and G. Szabó, “The dynamical Kirchberg–Phillips theorem”, *Acta Math.* **232**:1 (2024), 1–77. [MR](#) [Zbl](#)
- [Gardella and Lupini 2021] E. Gardella and M. Lupini, “Group amenability and actions on \mathcal{Z} -stable C^* -algebras”, *Adv. Math.* **389** (2021), art. id. 107931. [MR](#) [Zbl](#)
- [Gardella et al. 2022] E. Gardella, I. Hirshberg, and A. Vaccaro, “Strongly outer actions of amenable groups on \mathcal{Z} -stable nuclear C^* -algebras”, *J. Math. Pures Appl.* (9) **162** (2022), 76–123. [MR](#) [Zbl](#)
- [Gardella et al. 2024] E. Gardella, S. Geffen, J. Kranz, P. Naryshkin, and A. Vaccaro, “Tracially amenable actions and purely infinite crossed products”, *Math. Ann.* (online publication March 2024).
- [Gong and Lin 2020] G. Gong and H. Lin, “On classification of non-unital amenable simple C^* -algebras, II”, *J. Geom. Phys.* **158** (2020), art. id. 103865. [MR](#) [Zbl](#)
- [Gong and Lin 2022] G. Gong and H. Lin, “On classification of nonunital amenable simple C^* -algebras, III : The range and the reduction”, *Ann. K-Theory* **7**:2 (2022), 279–384. [MR](#) [Zbl](#)
- [Gong et al. 2020a] G. Gong, H. Lin, and Z. Niu, “A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, I: C^* -algebras with generalized tracial rank one”, *C. R. Math. Acad. Sci. Soc. R. Can.* **42**:3 (2020), 63–450. [MR](#) [Zbl](#)
- [Gong et al. 2020b] G. Gong, H. Lin, and Z. Niu, “A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, II: C^* -algebras with rational generalized tracial rank one”, *C. R. Math. Acad. Sci. Soc. R. Can.* **42**:4 (2020), 451–539. [MR](#) [Zbl](#)
- [Haagerup 1975] U. Haagerup, “The standard form of von Neumann algebras”, *Math. Scand.* **37**:2 (1975), 271–283. [MR](#) [Zbl](#)
- [Haagerup 1987] U. Haagerup, “Connes’ bicentralizer problem and uniqueness of the injective factor of type III₁”, *Acta Math.* **158**:1-2 (1987), 95–148. [MR](#) [Zbl](#)
- [Haagerup 2014] U. Haagerup, “Quasitraces on exact C^* -algebras are traces”, *C. R. Math. Acad. Sci. Soc. R. Can.* **36**:2-3 (2014), 67–92. [MR](#) [Zbl](#)
- [Izumi 2004a] M. Izumi, “Finite group actions on C^* -algebras with the Rohlin property, I”, *Duke Math. J.* **122**:2 (2004), 233–280. [MR](#) [Zbl](#)
- [Izumi 2004b] M. Izumi, “Finite group actions on C^* -algebras with the Rohlin property, II”, *Adv. Math.* **184**:1 (2004), 119–160. [MR](#) [Zbl](#)
- [Izumi and Matui 2010] M. Izumi and H. Matui, “ \mathbb{Z}^2 -actions on Kirchberg algebras”, *Adv. Math.* **224**:2 (2010), 355–400. [MR](#) [Zbl](#)
- [Izumi and Matui 2021a] M. Izumi and H. Matui, “Poly- \mathbb{Z} group actions on Kirchberg algebras, I”, *Int. Math. Res. Not.* **2021**:16 (2021), 12077–12154. [MR](#) [Zbl](#)
- [Izumi and Matui 2021b] M. Izumi and H. Matui, “Poly- \mathbb{Z} group actions on Kirchberg algebras, II”, *Invent. Math.* **224**:3 (2021), 699–766. [MR](#) [Zbl](#)
- [Jiang and Su 1999] X. Jiang and H. Su, “On a simple unital projectionless C^* -algebra”, *Amer. J. Math.* **121**:2 (1999), 359–413. [MR](#) [Zbl](#)
- [Jones 1980] V. F. R. Jones, *Actions of finite groups on the hyperfinite type II₁ factor*, Mem. Amer. Math. Soc. **237**, Amer. Math. Soc., Providence, RI, 1980. [MR](#) [Zbl](#)
- [Jones 1983] V. F. R. Jones, “A converse to Ocneanu’s theorem”, *J. Operator Theory* **10**:1 (1983), 61–63. [MR](#) [Zbl](#)
- [Kasparov 1988] G. G. Kasparov, “Equivariant KK -theory and the Novikov conjecture”, *Invent. Math.* **91**:1 (1988), 147–201. [MR](#) [Zbl](#)
- [Katayama et al. 1998] Y. Katayama, C. E. Sutherland, and M. Takesaki, “The characteristic square of a factor and the cocycle conjugacy of discrete group actions on factors”, *Invent. Math.* **132**:2 (1998), 331–380. [MR](#) [Zbl](#)

- [Katsura and Matui 2008] T. Katsura and H. Matui, “Classification of uniformly outer actions of \mathbb{Z}^2 on UHF algebras”, *Adv. Math.* **218**:3 (2008), 940–968. [MR](#) [Zbl](#)
- [Kawahigashi et al. 1992] Y. Kawahigashi, C. E. Sutherland, and M. Takesaki, “The structure of the automorphism group of an injective factor and the cocycle conjugacy of discrete abelian group actions”, *Acta Math.* **169**:1-2 (1992), 105–130. [MR](#) [Zbl](#)
- [Kerr 2020] D. Kerr, “Dimension, comparison, and almost finiteness”, *J. Eur. Math. Soc.* **22**:11 (2020), 3697–3745. [MR](#) [Zbl](#)
- [Kerr and Naryshkin 2021] D. Kerr and P. Naryshkin, “Elementary amenability and almost finiteness”, preprint, 2021. [arXiv 2107.05273](#)
- [Kerr and Szabó 2020] D. Kerr and G. Szabó, “Almost finiteness and the small boundary property”, *Comm. Math. Phys.* **374**:1 (2020), 1–31. [MR](#) [Zbl](#)
- [Kirchberg 1995] E. Kirchberg, “Exact C^* -algebras, tensor products, and the classification of purely infinite algebras”, pp. 943–954 in *Proceedings of the International Congress of Mathematicians, II* (Zürich, 1994), edited by S. D. Chatterji, Birkhäuser, Basel, 1995. [MR](#) [Zbl](#)
- [Kirchberg 2006] E. Kirchberg, “Central sequences in C^* -algebras and strongly purely infinite algebras”, pp. 175–231 in *Operator algebras* (Oslo, 2004), edited by O. Brattelli et al., Abel Symp. **1**, Springer, 2006. [MR](#) [Zbl](#)
- [Kirchberg and Rørdam 2002] E. Kirchberg and M. Rørdam, “Infinite non-simple C^* -algebras: absorbing the Cuntz algebra O_∞ ”, *Adv. Math.* **167**:2 (2002), 195–264. [MR](#) [Zbl](#)
- [Kirchberg and Rørdam 2014] E. Kirchberg and M. Rørdam, “Central sequence C^* -algebras and tensorial absorption of the Jiang–Su algebra”, *J. Reine Angew. Math.* **695** (2014), 175–214. [MR](#) [Zbl](#)
- [Kishimoto 1995] A. Kishimoto, “The Rohlin property for automorphisms of UHF algebras”, *J. Reine Angew. Math.* **465** (1995), 183–196. [MR](#) [Zbl](#)
- [Kishimoto 1998a] A. Kishimoto, “Automorphisms of AT algebras with the Rohlin property”, *J. Operator Theory* **40**:2 (1998), 277–294. [MR](#) [Zbl](#)
- [Kishimoto 1998b] A. Kishimoto, “Unbounded derivations in AT algebras”, *J. Funct. Anal.* **160**:1 (1998), 270–311. [MR](#) [Zbl](#)
- [Liao and Tikuisis 2022] H.-C. Liao and A. Tikuisis, “Almost finiteness, comparison, and tracial \mathcal{Z} -stability”, *J. Funct. Anal.* **282**:3 (2022), art. id. 109309. [MR](#) [Zbl](#)
- [Lin 1991] H. X. Lin, “Simple C^* -algebras with continuous scales and simple corona algebras”, *Proc. Amer. Math. Soc.* **112**:3 (1991), 871–880. [MR](#) [Zbl](#)
- [Lin 2023] H. Lin, “Hereditary uniform property Γ ”, *Sci. China Math.* **66**:8 (2023), 1813–1830. [MR](#) [Zbl](#)
- [Masuda 2007] T. Masuda, “Evans–Kishimoto type argument for actions of discrete amenable groups on McDuff factors”, *Math. Scand.* **101**:1 (2007), 48–64. [MR](#) [Zbl](#)
- [Masuda 2013] T. Masuda, “Unified approach to the classification of actions of discrete amenable groups on injective factors”, *J. Reine Angew. Math.* **683** (2013), 1–47. [MR](#) [Zbl](#)
- [Matui 2008] H. Matui, “Classification of outer actions of \mathbb{Z}^N on O_2 ”, *Adv. Math.* **217**:6 (2008), 2872–2896. [MR](#) [Zbl](#)
- [Matui 2010] H. Matui, “ \mathcal{Z} -actions on AH algebras and \mathbb{Z}^2 -actions on AF algebras”, *Comm. Math. Phys.* **297**:2 (2010), 529–551. [MR](#) [Zbl](#)
- [Matui 2011] H. Matui, “ \mathbb{Z}^N -actions on UHF algebras of infinite type”, *J. Reine Angew. Math.* **657** (2011), 225–244. [MR](#) [Zbl](#)
- [Matui and Sato 2012] H. Matui and Y. Sato, “ \mathcal{Z} -stability of crossed products by strongly outer actions”, *Comm. Math. Phys.* **314**:1 (2012), 193–228. [MR](#) [Zbl](#)
- [Matui and Sato 2014a] H. Matui and Y. Sato, “Decomposition rank of UHF-absorbing C^* -algebras”, *Duke Math. J.* **163**:14 (2014), 2687–2708. [MR](#) [Zbl](#)
- [Matui and Sato 2014b] H. Matui and Y. Sato, “ \mathcal{Z} -stability of crossed products by strongly outer actions, II”, *Amer. J. Math.* **136**:6 (2014), 1441–1496. [MR](#) [Zbl](#)
- [Murray and von Neumann 1943] F. J. Murray and J. von Neumann, “On rings of operators, IV”, *Ann. of Math. (2)* **44** (1943), 716–808. [MR](#) [Zbl](#)
- [Nakamura 2000] H. Nakamura, “Aperiodic automorphisms of nuclear purely infinite simple C^* -algebras”, *Ergodic Theory Dynam. Systems* **20**:6 (2000), 1749–1765. [MR](#) [Zbl](#)
- [Ocneanu 1985] A. Ocneanu, *Actions of discrete amenable groups on von Neumann algebras*, Lecture Notes in Math. **1138**, Springer, 1985. [MR](#) [Zbl](#)

- [Pedersen 1979] G. K. Pedersen, *C*-algebras and their automorphism groups*, Lond. Math. Soc. Monogr. **14**, Academic Press, London, 1979. [MR](#) [Zbl](#)
- [Powers and Størmer 1970] R. T. Powers and E. Størmer, “Free states of the canonical anticommutation relations”, *Comm. Math. Phys.* **16** (1970), 1–33. [MR](#) [Zbl](#)
- [Rørdam 2003] M. Rørdam, “A simple C*-algebra with a finite and an infinite projection”, *Acta Math.* **191**:1 (2003), 109–142. [MR](#) [Zbl](#)
- [Rørdam 2004] M. Rørdam, “The stable and the real rank of \mathcal{Z} -absorbing C*-algebras”, *Int. J. Math.* **15**:10 (2004), 1065–1084. [MR](#) [Zbl](#)
- [Rørdam and Winter 2010] M. Rørdam and W. Winter, “The Jiang–Su algebra revisited”, *J. Reine Angew. Math.* **642** (2010), 129–155. [MR](#) [Zbl](#)
- [Sato 2010] Y. Sato, “The Rohlin property for automorphisms of the Jiang–Su algebra”, *J. Funct. Anal.* **259**:2 (2010), 453–476. [MR](#) [Zbl](#)
- [Sato 2019] Y. Sato, “Actions of amenable groups and crossed products of \mathcal{Z} -absorbing C*-algebras”, pp. 189–210 in *Operator algebras and mathematical physics*, edited by M. Izumi et al., Adv. Stud. Pure Math. **80**, Math. Soc. Japan, Tokyo, 2019. [MR](#) [Zbl](#)
- [Sato et al. 2015] Y. Sato, S. White, and W. Winter, “Nuclear dimension and \mathcal{Z} -stability”, *Invent. Math.* **202**:2 (2015), 893–921. [MR](#) [Zbl](#)
- [Schafhauser 2020] C. Schafhauser, “Subalgebras of simple AF-algebras”, *Ann. of Math. (2)* **192**:2 (2020), 309–352. [MR](#) [Zbl](#)
- [Sutherland and Takesaki 1989] C. E. Sutherland and M. Takesaki, “Actions of discrete amenable groups on injective factors of type III $_{\lambda}$, $\lambda \neq 1$ ”, *Pacific J. Math.* **137**:2 (1989), 405–444. [MR](#) [Zbl](#)
- [Suzuki 2021] Y. Suzuki, “Equivariant \mathcal{O}_2 -absorption theorem for exact groups”, *Compos. Math.* **157**:7 (2021), 1492–1506. [MR](#) [Zbl](#)
- [Szabó 2018a] G. Szabó, “Equivariant Kirchberg–Phillips-type absorption for amenable group actions”, *Comm. Math. Phys.* **361**:3 (2018), 1115–1154. [MR](#) [Zbl](#)
- [Szabó 2018b] G. Szabó, “Strongly self-absorbing C*-dynamical systems”, *Trans. Amer. Math. Soc.* **370**:1 (2018), 99–130. [MR](#) [Zbl](#)
- [Szabó 2018c] G. Szabó, “Strongly self-absorbing C*-dynamical systems, II”, *J. Noncommut. Geom.* **12**:1 (2018), 369–406. [MR](#) [Zbl](#)
- [Szabó 2021a] G. Szabó, “The classification of Rokhlin flows on C*-algebras”, *Comm. Math. Phys.* **382**:3 (2021), 2015–2070. [MR](#) [Zbl](#)
- [Szabó 2021b] G. Szabó, “Equivariant property (SI) revisited”, *Anal. PDE* **14**:4 (2021), 1199–1232. [MR](#) [Zbl](#)
- [Szabó 2021c] G. Szabó, “On a categorical framework for classifying C*-dynamics up to cocycle conjugacy”, *J. Funct. Anal.* **280**:8 (2021), art. id. 108927. [MR](#) [Zbl](#)
- [Szabó and Wouters 2024] G. Szabó and L. Wouters, “Dynamical McDuff-type properties for group actions on von Neumann algebras”, *J. Inst. Math. Jussieu* **23**:6 (2024), 2593–2629. [MR](#) [Zbl](#)
- [Tikuisis et al. 2017] A. Tikuisis, S. White, and W. Winter, “Quasidiagonality of nuclear C*-algebras”, *Ann. of Math. (2)* **185**:1 (2017), 229–284. [MR](#) [Zbl](#)
- [Toms 2008] A. S. Toms, “On the classification problem for nuclear C*-algebras”, *Ann. of Math. (2)* **167**:3 (2008), 1029–1044. [MR](#) [Zbl](#)
- [Toms and Winter 2013] A. S. Toms and W. Winter, “Minimal dynamics and K-theoretic rigidity: Elliott’s conjecture”, *Geom. Funct. Anal.* **23**:1 (2013), 467–481. [MR](#) [Zbl](#)
- [Villadsen 1999] J. Villadsen, “On the stable rank of simple C*-algebras”, *J. Amer. Math. Soc.* **12**:4 (1999), 1091–1102. [MR](#) [Zbl](#)
- [Wouters 2023] L. Wouters, “Equivariant \mathcal{Z} -stability for single automorphisms on simple C*-algebras with tractable trace simplices”, *Math. Z.* **304**:1 (2023), art. id. 22. [MR](#) [Zbl](#)

Received 28 Feb 2023. Revised 22 Apr 2024. Accepted 26 Jun 2024.

GÁBOR SZABÓ: gabor.szabo@kuleuven.be
 Department of Mathematics, KU Leuven, Leuven, Belgium

LISE WOUTERS: *Leuven, Belgium*

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: "Linear Ramp"


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2025 is US \$475/year for the electronic version, and \$735/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 18 No. 6 2025

Damped Strichartz estimates and the incompressible Euler–Maxwell system	1309
DIOGO ARSÉNIO and HAROUNE HOUAMED	
Equivariant property Gamma and the tracial local-to-global principle for C^* -dynamics	1385
GÁBOR SZABÓ and LISE WOUTERS	
Multijet bundles and application to the finiteness of moments for zeros of Gaussian fields	1433
MICHELE ANCONA and THOMAS LETENDRE	
Existence of solutions to a fractional semilinear heat equation in uniformly local weak Zygmund-type spaces	1477
NORISUKE IOKU, KAZUHIRO ISHIGE and TATSUKI KAWAKAMI	
A Marcinkiewicz multiplier theory for Schur multipliers	1511
CHIAN YEONG CHUAH, ZHEN-CHUAN LIU and TAO MEI	
Double duals and Hilbert modules	1531
HUAXIN LIN	