

ANALYSIS & PDE

Volume 18

No. 6

2025

MICHELE ANCONA AND THOMAS LETENDRE

**MULTIJET BUNDLES AND APPLICATION TO
THE FINITENESS OF MOMENTS FOR ZEROS OF GAUSSIAN
FIELDS**

MULTIJET BUNDLES AND APPLICATION TO THE FINITENESS OF MOMENTS FOR ZEROS OF GAUSSIAN FIELDS

MICHELE ANCONA AND THOMAS LETENDRE

We define a notion of multijet for functions on \mathbb{R}^n , which extends the classical notion of jets in the sense that the multijet of a function is defined by contact conditions at several points. For all $p \geq 1$ we build a vector bundle of p -multijets, defined over a well-chosen compactification of the configuration space of p distinct points in \mathbb{R}^n . As an application, we prove that the linear statistics associated with the zero set of a centered Gaussian field on a Riemannian manifold have a finite p -th moment as soon as the field is of class C^p and its $(p-1)$ -jet is nowhere degenerate. We prove a similar result for the linear statistics associated with the critical points of a Gaussian field and those associated with the vanishing locus of a holomorphic Gaussian field.

1. Introduction	1433
2. Notation: partitions and function spaces	1440
3. Divided differences and Kergin interpolation	1441
4. Evaluation maps and their kernels	1443
5. Definition of the multijet bundles	1445
6. Application to zeros of Gaussian fields	1454
7. Multijets adapted to a differential operator	1465
8. Multijets of holomorphic maps	1469
Acknowledgments	1474
References	1474

1. Introduction

This paper is concerned with two different but related problems. The first one is to define a natural notion of multijet for a C^k function on \mathbb{R}^n , generalizing the usual notion of k -jet. By multijet we mean that we want to consider a collection of jets at different points in \mathbb{R}^n and patch them together in a relevant way. The second one is to find natural conditions on a Gaussian field $f : \mathbb{R}^n \rightarrow \mathbb{R}^r$ ensuring that the $(n-r)$ -dimensional volume of $f^{-1}(0) \cap \mathbb{B}$ admits finite higher moments, where \mathbb{B} stands for the unit ball in \mathbb{R}^n . One way to tackle this second problem is by considering the multijet of the random field f . In the following, we give more details about our contributions concerning the previous two problems, as well as some variations on these questions.

Letendre was partially supported by the ANR grant UniRandDom (ANR-17-CE40-0008) and a PEPS grant from the CNRS INSMI.

MSC2020: 51M15, 51M25, 55R80, 58A20, 60D05, 60G60.

Keywords: compactification of configuration spaces, Gaussian fields, moments, multijets, random submanifolds.

1.1. Multijet bundles. Let us start by recalling some standard facts about jets. See [Saunders 1989] for background on this matter. Let $x \in \mathbb{R}^n$ and $k \geq 0$. Two smooth functions f and g on \mathbb{R}^n are said to have the same k -jet at x if $f - g$ vanishes at x , as well as all its partial derivatives up to order k . Having the same k -jet at x is an equivalence relation on $C^\infty(\mathbb{R}^n)$, and the space $\mathcal{J}_k(\mathbb{R}^n)_x$ of k -jets at x is the set of equivalence classes for this relation. We denote by $j_k(f, x)$ the k -jet of f at x , that is, its class in $\mathcal{J}_k(\mathbb{R}^n)_x$. The map $j_k(\cdot, x)$ is a linear surjection from $C^\infty(\mathbb{R}^n)$ onto the finite-dimensional vector space $\mathcal{J}_k(\mathbb{R}^n)_x$. Of course, $j_k(f, x)$ makes sense even if f is only C^k and defined on some neighborhood of x .

Considering the family of k -jet spaces for all $x \in \mathbb{R}^n$, the set $\mathcal{J}_k(\mathbb{R}^n) = \bigsqcup_{x \in \mathbb{R}^n} \mathcal{J}_k(\mathbb{R}^n)_x$ is equipped with a natural vector bundle structure over \mathbb{R}^n . Then, if $\Omega \subset \mathbb{R}^n$ is open and $f : \Omega \rightarrow \mathbb{R}$ is C^k , the map $j_k(f, \cdot)$ is a local section of $\mathcal{J}_k(\mathbb{R}^n) \rightarrow \mathbb{R}^n$ over Ω . These definitions are well-behaved with respect to smooth changes of coordinates, so one can define similarly the vector bundle of k -jets of functions on a manifold M . More generally, if $E \rightarrow M$ is a vector bundle over M , there is a corresponding vector bundle $\mathcal{J}_k(M, E) \rightarrow M$ of k -jets of sections of $E \rightarrow M$.

In this paper, we are interested in defining similarly a notion of multijet and the associated vector bundles. That is, we want to consider smooth functions on \mathbb{R}^n up to an equivalence relation defined by the vanishing of some derivatives at several points.

Let us make this more precise. Let $p \geq 1$ and $\Delta_p = \{(x_1, \dots, x_p) \in (\mathbb{R}^n)^p \mid \exists i \neq j \text{ such that } x_i = x_j\}$ denote the diagonal in $(\mathbb{R}^n)^p$. Given $\underline{x} = (x_1, \dots, x_p) \notin \Delta_p$, we say that f and g have the same multijet at \underline{x} if $f(x_i) = g(x_i)$ for all $i \in \llbracket 1, p \rrbracket$ (here we use the notation $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{N}$). This is an equivalence relation on functions, defined by the vanishing of $f - g$ on the set $\{x_1, \dots, x_p\} \subset \mathbb{R}^n$, that is, by p independent linear conditions. Thus the corresponding set of classes is a vector space of dimension p , which we denote by $\mathcal{MJ}_p(\mathbb{R}^n)_{\underline{x}}$. We also denote by $\text{mj}_p(f, \underline{x})$ the class of f in this space, that is, its multijet at \underline{x} .

As will be explained later, this defines a vector bundle $\mathcal{MJ}_p(\mathbb{R}^n)$ of rank p over $(\mathbb{R}^n)^p \setminus \Delta_p$. Moreover, for all $\underline{x} \notin \Delta_p$ the linear map $\text{mj}_p(\cdot, \underline{x}) : C^\infty(\mathbb{R}^n) \rightarrow \mathcal{MJ}_p(\mathbb{R}^n)_{\underline{x}}$ is surjective, and, for all smooth f , its multijet $\text{mj}_p(f, \cdot)$ is smooth. We would like to extend this picture over the whole of $(\mathbb{R}^n)^p$. Note that the surjectivity conditions rule out defining $\mathcal{MJ}_p(\mathbb{R}^n)$ as $\mathcal{J}_0(\mathbb{R}^n)^p$ with $\text{mj}_p(f, \underline{x}) = (j_0(f, x_i))_{1 \leq i \leq p}$. When $\underline{x} \notin \Delta_p$, the previous notion of multijet is defined by p independent linear conditions: vanishing at each of the x_i . The main issue is that, when $\underline{x} \in \Delta_p$, these conditions are no longer independent and we need to replace them by another p -tuple of independent conditions.

A first natural idea is to look at vanishing with multiplicities. In dimension $n = 1$ this works very well. Let $\underline{x} \in \mathbb{R}^p$ be a permutation of $(y_1, \dots, y_1, \dots, y_m, \dots, y_m)$, where $(y_j)_{1 \leq j \leq m} \in \mathbb{R}^m \setminus \Delta_m$ and y_j appears exactly $k_j + 1$ times. We say that f and g have the same multijet at \underline{x} if $(f - g)^{(k_j)}(y_j) = 0$ for all $j \in \llbracket 1, m \rrbracket$ and $k \in \llbracket 0, k_j \rrbracket$. In this sense, having the same multijet is equivalent to having the same Hermite interpolating polynomials at \underline{x} . Thus, we can define $\mathcal{MJ}_p(\mathbb{R})$ as the trivial bundle $\mathbb{R}_{p-1}[X] \times \mathbb{R}^p \rightarrow \mathbb{R}^p$ and $\text{mj}_p(f, \underline{x})$ as the Hermite interpolating polynomial of f at \underline{x} .

If $n > 1$, the previous approach fails already for $p = 2$. Let us consider $x \in \mathbb{R}^n$ and the corresponding $\underline{x} = (x, x) \in \Delta_2$. Asking for the vanishing of $f - g$ and its differential at x gives us $n + 1$ independent conditions, which define the 1-jet space $\mathcal{J}_1(\mathbb{R}^n)_x$. This space has dimension $n + 1 > 2$; hence it is too large to be the $\mathcal{MJ}_2(\mathbb{R}^n)_{\underline{x}}$ we are looking for. The next natural idea is to ask only for the vanishing

at x of $f - g$ and one of its directional derivatives. But whatever choice of directional derivative we make will lead to $\text{mj}_2(f, \cdot)$ not being continuous at x for most $f \in C^\infty(\mathbb{R}^n)$. Actually, one cannot extend $\mathcal{MJ}_2(\mathbb{R}^n)$ nicely over $(\mathbb{R}^n)^2$ if $n > 1$. However, we can extend it nicely over a larger space: the blow-up $\text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$ of $(\mathbb{R}^n)^2$ along Δ_2 . The key idea is that $\text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$ contains $(\mathbb{R}^n)^2 \setminus \Delta_2$ as a dense open subset and that points in the complement of $(\mathbb{R}^n)^2 \setminus \Delta_2$ can be described by the following data: a base point $x \in \mathbb{R}^n$ and a direction $u \in \mathbb{R}\mathbb{P}^{n-1}$. This data tells us exactly which directional derivative to consider at the corresponding point in the exceptional locus of $\text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$. We will come back to this example later; see Example 5.9.

This long discussion shows that there is a natural way to define a multijet bundle $\mathcal{MJ}_p(\mathbb{R}^n)$ over the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p$, but that it does not extend nicely over $(\mathbb{R}^n)^p$ in general. The case $p = 2$ hints that it might however be possible to define a natural multijet bundle over a slightly larger space, containing a copy of $(\mathbb{R}^n)^p \setminus \Delta_p$ as a dense open subset. Our first main contribution is to define such an object. Its main properties are gathered in the following statement, where $\mathcal{C}^k(\mathbb{R}^n, V)$ denotes the space of \mathcal{C}^k functions from \mathbb{R}^n to V .

Theorem 1.1 (existence of multijet bundles). *Let $n \geq 1$ and $p \geq 1$ and let V be a real vector space of finite dimension $r \geq 1$. There exist a smooth manifold $C_p[\mathbb{R}^n]$ of dimension np without boundary and a smooth vector bundle $\mathcal{MJ}_p(\mathbb{R}^n, V) \rightarrow C_p[\mathbb{R}^n]$ of rank rp with the following properties:*

(1) *There exists a smooth proper surjection $\pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ such that $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ is a dense open subset of $C_p[\mathbb{R}^n]$, and π restricted to $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ is a C^∞ -diffeomorphism onto $(\mathbb{R}^n)^p \setminus \Delta_p$.*

(2) *There exists a map $\text{mj}_p : \mathcal{C}^{p-1}(\mathbb{R}^n, V) \times C_p[\mathbb{R}^n] \rightarrow \mathcal{MJ}_p(\mathbb{R}^n, V)$ such that*

- *for all $z \in C_p[\mathbb{R}^n]$, the linear map $\text{mj}_p(\cdot, z) : \mathcal{C}^{p-1}(\mathbb{R}^n, V) \rightarrow \mathcal{MJ}_p(\mathbb{R}^n, V)_z$ is surjective;*
- *for all $f \in \mathcal{C}^{l+p-1}(\mathbb{R}^n, V)$, the section $\text{mj}_p(f, \cdot)$ of $\mathcal{MJ}_p(\mathbb{R}^n, V) \rightarrow C_p[\mathbb{R}^n]$ is \mathcal{C}^l .*

(3) *Let $z \in C_p[\mathbb{R}^n]$ be such that $\pi(z) = (x_1, \dots, x_p) \notin \Delta_p$. Then for all $f \in \mathcal{C}^{p-1}(\mathbb{R}^n, V)$ we have*

$$\text{mj}_p(f, z) = 0 \iff \forall i \in \llbracket 1, p \rrbracket, f(x_i) = 0.$$

(4) *Let $z \in C_p[\mathbb{R}^n]$ be such that $\pi(z)$ is obtained as a permutation of $(y_1, \dots, y_1, \dots, y_m, \dots, y_m)$, where y_j appears exactly $k_j + 1$ times and y_1, \dots, y_m are pairwise distinct vectors in \mathbb{R}^n . Then, there exists a linear surjection $\Theta_z : \prod_{i=1}^m \mathcal{J}_{k_j}(\mathbb{R}^n, V)_{y_j} \rightarrow \mathcal{MJ}_p(\mathbb{R}^n, V)_z$ such that*

$$\forall f \in \mathcal{C}^{p-1}(\mathbb{R}^n, V), \quad \text{mj}_p(f, z) = \Theta_z(j_{k_1}(f, y_1), \dots, j_{k_m}(f, y_m)).$$

Remark 1.2. In Theorem 1.1, the manifold $C_p[\mathbb{R}^n]$ does not depend on V . Part (1) means that we can consider $(\mathbb{R}^n)^p \setminus \Delta_p$ as a dense open subset in $C_p[\mathbb{R}^n]$. Part (2) consists of properties that we expect any reasonable notion of multijet to satisfy. Part (3) means that, as in the previous discussions, if $\pi(z) \notin \Delta_p$ then $\mathcal{MJ}_p(\mathbb{R}^n)_z = \mathcal{C}^{p-1}(\mathbb{R}^n, V) / \sim$, where $f \sim g$ if and only if $f(x_i) = g(x_i)$ for all $i \in \llbracket 1, p \rrbracket$. Part (4) means that, more generally, $\text{mj}_p(f, z)$ only depends on the collection of jets $(j_{k_j}(f, y_j))_{1 \leq j \leq m}$. In particular, $\text{mj}_p(f, z)$ still makes sense if f is only \mathcal{C}^{k_j} on some neighborhood of y_j . This last condition also means that we can think of $\text{mj}_p(f, z)$ intuitively as a family of p independent linear combinations of partial derivatives of f , up to order k_j at y_j . However this family is neither explicit nor unique in general.

Let us now introduce some definitions and notation.

Definition 1.3 (multijets). Let $\Omega \subset \mathbb{R}^n$ be open. We let $C_p[\Omega] = \pi^{-1}(\Omega^p)$ and denote by $\mathcal{MJ}_p(\Omega, V) \rightarrow C_p[\Omega]$ the restriction of $\mathcal{MJ}_p(\mathbb{R}^n, V)$ to $C_p[\Omega]$. We call $\mathcal{MJ}_p(\Omega, V) \rightarrow C_p[\Omega]$ the *bundle of p -multijets* of functions from Ω to V . Its fiber $\mathcal{MJ}_p(\mathbb{R}^n, V)_z$ above $z \in C_p[\Omega]$ is the *space of p -multijets at z* . If $V = \mathbb{R}$, we drop it from the notation and write $\mathcal{MJ}_p(\Omega) \rightarrow C_p[\Omega]$. Let $f : \Omega \rightarrow V$ be of class \mathcal{C}^{p-1} , we call the section $\text{mj}_p(f, \cdot)$ of $\mathcal{MJ}_p(\Omega, V)$ the *p -multijet of f* and its value at $z \in C_p[\Omega]$ the *p -multijet of f at z* .

The manifold $C_p[\mathbb{R}^n]$ is what is called in the literature a “compactification” of the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p$. We will use this terminology, even though it is ill-chosen in our case since $C_p[\mathbb{R}^n]$ is not compact. However $C_p[\mathbb{R}^n]$ contains a diffeomorphic copy of $(\mathbb{R}^n)^p \setminus \Delta_p$ as a dense open subset and it is equipped with a proper surjection onto $(\mathbb{R}^n)^p$ so that, in a sense, it is locally a compactification of $(\mathbb{R}^n)^p \setminus \Delta_p$.

Compactifications of configuration spaces are built to understand how a configuration (ordered or not) of p distinct points can degenerate as these points converge toward one another. They are usually obtained by blowing up various pieces of the diagonal. Points in the exceptional locus then correspond to singular configurations, with some extra data encoding along which paths regular configurations are allowed to degenerate in order to reach this singular configuration. The hope is that the extra data attached to singular configurations is enough to lift the singularities of the problem under consideration. The simplest example of this kind is the blow-up $\text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$ discussed above. More evolved examples are the space defined by Le Barz [1988], the compactification of Fulton and MacPherson [1994] (see also [Axelrod and Singer 1994; Sinha 2004]), Olver’s multispace [2001], the polydiagonal compactification of Ulyanov [2002], the construction of Evain [2005] using Hilbert schemes, and many others.

In dimension $n = 1$, most of the compactifications of configuration spaces that we found in the literature coincide and can be used to define multijets; see for example [Ancona 2021], where Olver’s multispace is used. In higher dimensions they are different and none of them exactly suited our needs. Thus to the best of our knowledge, the manifold $C_p[\mathbb{R}^n]$ in Theorem 1.1 is a new addition to the previous list. We define it by resolving the singularities of some real-algebraic variety, using Hironaka’s theorem [1964a; 1964b]. In particular, $C_p[\mathbb{R}^n]$ is obtained by a sequence of blow-ups along Δ_p . Note that this sequence of blow-ups is neither explicit nor unique. Actually, $C_p[\mathbb{R}^n]$ itself is not uniquely defined, but this is not an issue for the applications we have in mind.

1.2. Finiteness of moments for zeros of Gaussian fields. Let us now describe our contributions concerning zeros of Gaussian fields. Let $n \geq 1$ and let $r \in \llbracket 1, n \rrbracket$. In the following n will always denote the dimension of the ambient space and r the codimension of the random objects we are interested in.

Let $\Omega \subset \mathbb{R}^n$ be open and let $f : \Omega \rightarrow \mathbb{R}^r$ be a centered Gaussian field of class \mathcal{C}^1 . We will always assume that f is nondegenerate, in the sense that $\det \text{Var}(f(x)) > 0$ for all $x \in \Omega$. Under this hypothesis the zero set $Z = f^{-1}(0)$ is almost surely $(n-r)$ -rectifiable; see [Armentano et al. 2023b]. As such, it admits a well-defined $(n-r)$ -dimensional volume measure dVol_Z induced by the Euclidean metric on \mathbb{R}^n .

We denote by ν the random Radon measure on Ω defined by

$$\forall \phi \in \mathcal{C}_c^0(\Omega), \quad \langle \nu, \phi \rangle = \int_Z \phi(x) \, d\text{Vol}_Z(x), \tag{1-1}$$

where $\mathcal{C}_c^0(\Omega)$ denotes the space of continuous functions on Ω with compact support.

Actually, $\langle \nu, \phi \rangle$ makes sense as an almost surely defined random variable as soon as $\phi \in L^\infty(\Omega)$ and has compact support. This kind of test-function includes $\mathcal{C}_c^0(\Omega)$ and indicator functions of bounded Borel subsets, which are the examples we are most interested in. Random variables of the type $\langle \nu, \phi \rangle$ are called the linear statistics of ν (or of f). Understanding the distribution of these linear statistics is one way to understand the distribution of the random measure ν , or equivalently of the random set Z . For example, if $B \subset \Omega$ is a bounded Borel set and $\mathbf{1}_B$ denotes its indicator function, then $\langle \nu, \mathbf{1}_B \rangle$ is the $(n-r)$ -dimensional volume of $Z \cap B$.

In this setting, a classical question is to determine conditions on the field f ensuring that its linear statistics admit finite moments. If $n = r = 1$, such conditions were first obtained in [Belyaev 1966]. More generally see [Azaïš and Wschebor 2009, Theorem 3.6], which holds even if f is not Gaussian. If $n \geq r = 1$, a similar result is proved in [Armentano et al. 2023a]; see also [Armentano et al. 2019, Theorem 4.4]. For a survey of previous results for hypersurfaces (i.e., $r = 1$), we refer to [Azaïš and Wschebor 2009, Chapter 3, Section 2.7] in dimension $n = 1$ and to the introduction of [Armentano et al. 2023a] in dimension $n \geq 1$. Note that [Priya 2020, Theorem 1.2] implies the finiteness of all moments of the nodal length for some Gaussian fields in \mathbb{R}^2 . This problem was also studied in [Malevich and Volodina 1993] for points in \mathbb{R}^2 .

Our second main result gives simple conditions on the field f ensuring the finiteness of the p -th moments of its linear statistics in any dimension and codimension. These conditions are of two kinds: we require the field to be regular enough, and to be nondegenerate in the following sense.

Definition 1.4 (p -nondegeneracy). Let $p \geq 1$ and let $f : \Omega \rightarrow \mathbb{R}^r$ be a \mathcal{C}^p centered Gaussian field. We say that the field f is p -nondegenerate if for all $x \in \Omega$ the centered Gaussian vector

$$(f(x), D_x f, \dots, D_x^p f) \in \bigoplus_{k=0}^p \text{Sym}^k(\mathbb{R}^n) \otimes \mathbb{R}^r$$

is nondegenerate, where $\text{Sym}^k(\mathbb{R}^n)$ denotes the space of symmetric k -linear forms on \mathbb{R}^n and $D_x^k f \in \text{Sym}^k(\mathbb{R}^n) \otimes \mathbb{R}^r$ stands the k -th differential of f at x .

Remark 1.5. If $f = (f_1, \dots, f_r)$, the p -nondegeneracy condition means more concretely that for all $x \in \Omega$ the Gaussian vector $(\partial^\alpha f_i(x))_{1 \leq i \leq r; |\alpha| \leq p}$ is nondegenerate, where we used multi-index notation (see Section 2.2). More abstractly, this condition means that the p -jet $j_p(f, x)$ of f is nondegenerate for all $x \in \Omega$.

Theorem 1.6 (finiteness of moments). Let $n \geq 1$, let $r \in \llbracket 1, n \rrbracket$ and let $p \geq 1$. Let $\Omega \subset \mathbb{R}^n$ be open, let $f : \Omega \rightarrow \mathbb{R}^r$ be a centered Gaussian field and let ν be defined as in (1-1). If f is \mathcal{C}^p and $(p-1)$ -nondegenerate then $\mathbb{E}[|\langle \nu, \phi \rangle|^p] < +\infty$ for all $\phi \in L^\infty(\Omega)$ with compact support.

Example 1.7. Let us give some examples of fields satisfying the assumptions of Theorem 1.6.

- The Bargmann–Fock field, i.e., the smooth stationary Gaussian field on \mathbb{R}^n whose covariance function is $x \mapsto e^{-\|x\|^2/2}$, satisfies the hypotheses of Theorem 1.6.
- Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a stationary \mathcal{C}^p centered Gaussian field. If the support of its spectral measure has nonempty interior then f is $(p-1)$ -nondegenerate.
- In codimension r , if $(f_i)_{1 \leq i \leq r}$ are r independent $(p-1)$ -nondegenerate \mathcal{C}^p Gaussian fields then so is $f = (f_1, \dots, f_r)$.
- The Berry field, i.e., the smooth stationary Gaussian field f on \mathbb{R}^n whose spectral measure is the uniform measure on \mathbb{S}^{n-1} , is 1-nondegenerate but not 2-nondegenerate. Indeed it almost surely satisfies $\Delta f + f = 0$, so that $(f(x), D_x f, D_x^2 f)$ is degenerate for all $x \in \mathbb{R}^n$.

We can consider the same question in a more geometric setting. Let (M, g) be a Riemannian manifold of dimension $n \geq 1$ without boundary and let $E \rightarrow M$ be a smooth vector bundle of rank $r \in \llbracket 1, n \rrbracket$ over M . Let s be a centered Gaussian field on M with values in E , in the sense that s is a random section of $E \rightarrow M$ such that for all $m \geq 1$ and all $x_1, \dots, x_m \in M$ the random vector $(s(x_1), \dots, s(x_m))$ is a centered Gaussian. We assume that s is almost surely \mathcal{C}^1 and that $\det \text{Var}(s(x)) > 0$ for all $x \in M$.

As in the Euclidean setting, $Z = s^{-1}(0)$ is almost surely $(n-r)$ -rectifiable. As before, we denote by ν the random Radon measure on M defined by integrating over Z with respect to the $(n-r)$ -dimensional volume measure $d\text{Vol}_Z$ induced by g . For all $\phi \in L^\infty(M)$ with compact support, we define the linear statistic $\langle \nu, \phi \rangle$ as in (1-1). In this context Definition 1.4 adapts as follows.

Definition 1.8 (p -nondegeneracy for Gaussian sections). Let $p \geq 1$ and let s be a \mathcal{C}^p centered Gaussian field on M with values in E . We say that s is p -nondegenerate if, for all $x \in M$, the centered Gaussian vector $j_p(s, x) \in \mathcal{J}_p(M, E)_x$ is nondegenerate.

Theorem 1.9 (finiteness of moments for zeros of Gaussian sections). Let $p \geq 1$, let s be a centered Gaussian field on M with values in E and let ν be defined as in (1-1). If s is \mathcal{C}^p and $(p-1)$ -nondegenerate then $\mathbb{E}[|\langle \nu, \phi \rangle|^p] < +\infty$ for all $\phi \in L^\infty(M)$ with compact support.

We are aware of the very recent paper [Gass and Stecconi 2024], in which the authors prove a result similar to Theorem 1.6, as well as its analogue for zeros of Gaussian fields on a Riemannian manifold. Their work and ours are independent, and the proofs are different. Their idea is to compare the Kac–Rice densities (see Section 6.3) of the field f with those of a well-chosen Gaussian polynomial P . Then they deduce the result for f from the result for P , which is a consequence of Bézout’s theorem. Our proof follows a different path, as it relies on the multijet bundle that we defined in Theorem 1.1. Our idea is to observe that the zero set of $F : (x_1, \dots, x_p) \mapsto (f(x_1), \dots, f(x_p))$ in the configuration space $\Omega^p \setminus \Delta_p$ is exactly the vanishing locus of the multijet $\text{mj}_p(f, \cdot)$ restricted to $\Omega^p \setminus \Delta_p \subset C_p[\Omega]$. Instead of working with F , which degenerates along Δ_p , we work with the field $\text{mj}_p(f, \cdot)$ that we built to be nondegenerate everywhere. Then, we deduce Theorem 1.6 from the Kac–Rice formula for the expectation (see Proposition 6.17) applied to the p -multijet of f and a compactness argument.

1.3. Higher-order multijets and holomorphic multijets. Let us now discuss two important variations on our main results, Theorems 1.1 and 1.9. In Section 1.1, we said that two functions f and g on \mathbb{R}^n have the same p -multijet at a point $\underline{x} = (x_1, \dots, x_p) \in (\mathbb{R}^n)^p \setminus \Delta_p$ if and only if f and g have the same value, i.e., the same 0-jet, at x_i for all $i \in \llbracket 1, p \rrbracket$. In a sense, the p -multijet of f at \underline{x} is obtained by patching together the 0-jets of f at each of the x_i in a relevant way. A natural generalization is to define a higher-order multijet of f at \underline{x} by patching together the k -jets of f at each of the x_i . We define such a higher-order multijet in Section 7. More generally, we define a multijet bundle adapted to a differential operator \mathcal{D} . The case of higher-order multijets corresponds to $\mathcal{D} = j_k$. The analogue of Theorem 1.1 in this framework is Theorem 7.4 below. We use it to prove an analogue of Theorem 1.9 adapted to \mathcal{D} ; see Theorem 7.8 for a general statement. In the special case where $\mathcal{D} = D$ is the standard differential, the statement is the following.

Theorem 1.10 (finiteness of moments for critical points). *Let M be a smooth manifold without boundary. Let $f : M \rightarrow \mathbb{R}$ be a centered Gaussian field and let ν_D denote the counting measure of its critical locus. Let $p \geq 1$, we assume that f is C^{2p} and $(2p-1)$ -nondegenerate. Then, for all $\phi \in L_c^\infty(M)$, we have $\mathbb{E}[|\langle \nu_D, \phi \rangle|^p] < +\infty$.*

Another variation on Theorem 1.1 is to define holomorphic multijets for holomorphic maps. This is done in Section 8, and more precisely in Theorem 8.2. This is used to prove a holomorphic version of Theorem 1.9. The general statement is given in Theorem 8.13. For a holomorphic Gaussian field on an open subset of \mathbb{C}^n , it takes the following form.

Theorem 1.11 (finiteness of moments for zeros of holomorphic Gaussian fields). *Let $\Omega \subset \mathbb{C}^n$ be open and let $f : \Omega \rightarrow \mathbb{C}^r$ be a centered holomorphic Gaussian field, where $r \in \llbracket 1, n \rrbracket$. Let ν be as in Definition 6.11. Let $p \geq 1$, we assume that, for all $x \in \Omega$, the complex Gaussian vector*

$$(f(x), D_x f, \dots, D_x^{p-1} f) \in \bigoplus_{k=0}^{p-1} \text{Sym}^k(\mathbb{C}^n) \otimes \mathbb{C}^r$$

is nondegenerate. Then, for all $\phi \in L_c^\infty(\Omega)$, we have $\mathbb{E}[|\langle \nu, \phi \rangle|^p] < +\infty$.

Note that Theorems 1.10 and 1.11 are not consequences of Theorem 1.9. Indeed, if $f : M \rightarrow \mathbb{R}$ is a smooth Gaussian field then Df cannot be 1-nondegenerate because $D^2 f$ is symmetric. Similarly, if M is a complex manifold and s is holomorphic, then s is never 1-nondegenerate because it satisfies the Cauchy–Riemann equations.

Gass and Stecconi [2024] proved, independently and by a different method, results analogous to Theorems 1.10 and 1.11. Actually, they prove Theorem 1.10 under the weaker and optimal hypotheses that f is C^{p+1} and p -nondegenerate. The finiteness of the third moment for the number of critical points of a stationary Gaussian field on \mathbb{R}^d was proved in [Beliaev et al. 2024, Theorem 1.6]. For holomorphic Gaussian fields in dimension $n = 1$, see [Nazarov and Sodin 2012].

1.4. Organization of the paper. In Section 2 we gather useful notation that appears in several parts of the paper. In Section 3 we discuss Kergin interpolation, which is a multivariate polynomial interpolation appearing in the definition of multijets. Section 4 is dedicated to evaluations maps on spaces of polynomials,

and more precisely the properties of their kernels. We define our multijet bundles and prove Theorem 1.1 in Section 5. Section 6 is concerned with the application of multijets to the finiteness of moments for the zeros of Gaussian fields and the proofs of Theorems 1.6 and 1.9. Multijets adapted to a differential operator are discussed in Section 7, where we also prove the analogue of Theorem 1.9 for critical points. Finally, holomorphic multijets are defined in Section 8, where we prove the analogue of Theorem 1.9 for holomorphic Gaussian fields.

2. Notation: partitions and function spaces

The goal of this section is to quickly introduce definitions and notation that will appear in different parts of the paper. We gather them here for the reader’s convenience.

2.1. Sets, partitions and diagonals. In this paper, we denote by \mathbb{N} the set of nonnegative integers. Let a and $b \in \mathbb{N}$, we use the following notation for integer intervals $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{N}$.

Let A be a nonempty finite set. For simplicity, in all the notation introduced in this section, if $A = \llbracket 1, p \rrbracket$ we allow ourselves to replace A by p in the indices and exponents. We denote by $|A|$ the cardinality of A . Let M be any set. We denote by M^A the Cartesian product of $|A|$ copies of M indexed by the elements of A . A generic element of M^A is usually denoted by $\underline{x} = (x_a)_{a \in A}$. If $\emptyset \neq B \subset A$, we denote by $\underline{x}_B = (x_a)_{a \in B}$.

Definition 2.1 (large diagonal). We denote by Δ_A the large diagonal in M^A , that is,

$$\Delta_A = \{(x_a)_{a \in A} \in M^A \mid \exists a, b \in A \text{ such that } a \neq b \text{ and } x_a = x_b\}.$$

Definition 2.2 (partitions). Let A be a nonempty and finite set, a *partition* of A is a family $\mathcal{I} = \{I_1, \dots, I_m\}$ of nonempty disjoint subsets of A such that $\bigsqcup_{i=1}^m I_i = A$. The subsets I_1, \dots, I_m are called the *cells* of \mathcal{I} . Given $a \in A$, we denote by $[a]_{\mathcal{I}}$ the only cell of \mathcal{I} that contains a . Finally, we denote by \mathcal{P}_A the set of partitions of A .

Definition 2.3 (clustering partition). Let $\underline{x} = (x_a)_{a \in A} \in M^A$. We denote by $\mathcal{I}(\underline{x}) \in \mathcal{P}_A$ the only partition such that for all a and $b \in A$ we have $x_a = x_b$ if and only if $[a]_{\mathcal{I}(\underline{x})} = [b]_{\mathcal{I}(\underline{x})}$.

Example 2.4. If $\underline{x} = (x, \dots, x)$ then $\mathcal{I}(\underline{x}) = \{A\}$. If $\underline{x} \in M^A \setminus \Delta_A$ then $\mathcal{I}(\underline{x}) = \{\{a\} \mid a \in A\} = \mathcal{I}_0$.

Definition 2.5 (strata of the diagonal). For all $\mathcal{I} \in \mathcal{P}_A$, we set $\Delta_{A, \mathcal{I}} = \{\underline{x} \in M^A \mid \mathcal{I}(\underline{x}) = \mathcal{I}\}$, so that $\Delta_{A, \mathcal{I}_0} = M^A \setminus \Delta_A$ and $\Delta_A = \bigsqcup_{\mathcal{I} \neq \mathcal{I}_0} \Delta_{A, \mathcal{I}}$.

Definition 2.6 (diagonal inclusions). Let $\mathcal{I} \in \mathcal{P}_A$. We denote by $\iota_{\mathcal{I}} : M^{\mathcal{I}} \setminus \Delta_{\mathcal{I}} \rightarrow \Delta_{A, \mathcal{I}}$ the bijection defined by $\iota_{\mathcal{I}}((y_I)_{I \in \mathcal{I}}) = (y_{[a]_{\mathcal{I}}})_{a \in A}$.

2.2. Spaces of functions, sections and jets. We use the following multi-index notation. Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$. We denote its length by $|\alpha| = \alpha_1 + \dots + \alpha_n$. Let ∂_i denote the i -th partial derivative in some product space we denote by $\partial^\alpha = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n}$. Finally, if $X = (X_1, \dots, X_n)$, we let $X^\alpha = X_1^{\alpha_1} \dots X_n^{\alpha_n}$.

Definition 2.7 (polynomials). We denote by $\mathbb{R}_d[X]$ the space of real polynomials in n variables of degree at most d , where $d \in \mathbb{N}$ and $X = (X_1, \dots, X_n)$ is multivariate.

Definition 2.8 (symmetric forms and differentials). Let $k \in \mathbb{N}$. We denote by $\text{Sym}^k(\mathbb{R}^n)$ the space of symmetric k -linear forms on \mathbb{R}^n . Let V be a finite-dimensional real vector space. Then $\text{Sym}^k(\mathbb{R}^n) \otimes V$ is the space of symmetric k -linear maps from \mathbb{R}^n to V . Given a C^k map $f : \mathbb{R}^n \rightarrow V$, we denote by $D_x^k f \in \text{Sym}^k(\mathbb{R}^n) \otimes V$ its k -th differential at $x \in \mathbb{R}^n$.

Let M and N be two manifolds without boundary. For all $k \in \mathbb{N} \cup \{\infty\}$, we denote by $C^k(M, N)$ the space of C^k maps from M to N . If $N = \mathbb{R}$, we drop it from the notation and we simply write $C^k(M)$. We denote by $L^1_{\text{loc}}(M)$ the space of locally integrable functions on M . We denote by $C_c^0(M)$ (resp. $L^\infty_c(M)$) the space of continuous (resp. L^∞) functions on M with compact support. Finally, for any Borel subset $B \subset M$, we denote by $\mathbf{1}_B : M \rightarrow \mathbb{R}$ its indicator function.

Let $E \rightarrow M$ be a vector bundle of finite rank over M , we denote by E_x the fiber above $x \in M$. For all $k \in \mathbb{N} \cup \{\infty\}$, we denote by $\Gamma^k(M, E)$ the space of C^k sections of $E \rightarrow M$.

Definition 2.9 (jets). Let $k \in \mathbb{N}$, we denote by $\mathcal{J}_k(M, E) \rightarrow M$ the vector bundle of k -jets of sections of $E \rightarrow M$. If $E = V \times M$ is trivial with fiber V , we denote its k -jet bundle by $\mathcal{J}_k(M, V) \rightarrow M$. If $V = \mathbb{R}$, we simply write $\mathcal{J}_k(M) \rightarrow M$. Given $s \in \Gamma^k(M, E)$, we denote by $j_k(s, x) \in \mathcal{J}_k(M, E)_x$ its k -jet at $x \in M$.

3. Divided differences and Kergin interpolation

An important step in our construction of a multijet for C^k functions is to reduce the problem to that of defining a multijet for polynomials. This is done by polynomial interpolation. In several variables, polynomial interpolation is rather ill-behaved, at least compared with the one-variable case. However, a multivariate polynomial interpolation suiting our needs was defined by Kergin [1980]. A constructive version of his proof was then given in [Micchelli and Milman 1980], using a multivariate version of the so-called divided differences. In this section, we give the definitions of these objects and recall their relevant properties. We refer to the survey [Lorentz 2000] for more background on polynomial interpolation in \mathbb{R}^n .

3.1. Divided differences. In this section, we recall the definition of multivariate divided differences; see [Micchelli and Milman 1980]. Let $k \in \mathbb{N}$. We denote by σ_k the standard simplex of dimension k , that is,

$$\sigma_k = \left\{ \underline{t} = (t_0, \dots, t_k) \in [0, 1]^{k+1} \mid \sum_{i=0}^k t_i = 1 \right\} \subset \mathbb{R}^{k+1}. \tag{3-1}$$

The simplex σ_k is a subset of $\{ \underline{t} \in \mathbb{R}^{k+1} \mid \sum t_i = 1 \}$, and we denote by ν_k the (k -dimensional) Lebesgue measure on this hyperplane, normalized so that $\nu_k(\sigma_k) = 1/k!$. One can check that its restriction to σ_k satisfies

$$\int_{\sigma_k} \phi(\underline{t}) \, d\nu_k(\underline{t}) = \int_{\substack{t_1, \dots, t_k \geq 0 \\ \sum_{i=1}^k t_i \leq 1}} \phi\left(1 - \sum_{i=1}^k t_i, t_1, \dots, t_k\right) dt_1 \cdots dt_k, \tag{3-2}$$

where $dt_1 \cdots dt_k$ is the Lebesgue measure on \mathbb{R}^k . For any $\underline{x} = (x_0, \dots, x_k) \in (\mathbb{R}^n)^{k+1}$, we denote by $\sigma(\underline{x})$ the convex hull of the x_i and we define $\nu_{\underline{x}} : \underline{t} \mapsto \sum_{i=0}^k t_i x_i$ from σ_k onto $\sigma(\underline{x})$. Recalling Definition 2.8, we have the following.

Definition 3.1 (divided differences). Let $\underline{x} = (x_i)_{0 \leq i \leq k} \in (\mathbb{R}^n)^{k+1}$ and let f be a C^k function defined on some open neighborhood of $\sigma(\underline{x})$ in \mathbb{R}^n . We define the *divided difference* of f at \underline{x} by

$$f[x_0, \dots, x_k] = \int_{\sigma_k} D_{\nu_{\underline{x}}(t)}^k f \, d\nu_k(t) \in \text{Sym}^k(\mathbb{R}^n),$$

that is, as the average of $D^k f$ over $\sigma(\underline{x})$ with respect to the pushed-forward measure $(\nu_{\underline{x}})_*(\nu_k)$.

Remark 3.2. • If $\underline{x} = (x, \dots, x)$ for some $x \in \mathbb{R}^n$ then $f[x, \dots, x] = (1/k!)D_x^k f$.

- Definition 3.1 is invariant under permutation of (x_0, \dots, x_k) .
- When $n = 1$, Definition 3.1 coincides with the classical definition of divided differences, under the canonical isomorphism $\text{Sym}^k(\mathbb{R}) \simeq \mathbb{R}$. This is known as the Hermite–Genocchi formula [Micchelli and Milman 1980].

Lemma 3.3 (regularity of divided differences). *For all $\underline{x} \in (\mathbb{R}^n)^{k+1}$, the map $f \mapsto f[x_0, \dots, x_k]$ is linear. Moreover, if f is of class C^{k+l} then $\underline{x} \mapsto f[x_0, \dots, x_k]$ is of class C^l .*

Proof. The linearity with respect to f is clear. The regularity with respect to \underline{x} is obtained by derivation under the integral, using Definition 3.1 and (3-2). □

3.2. Kergin interpolation. This section is dedicated to Kergin interpolation. In the following, we recall the construction of Kergin interpolation in [Micchelli and Milman 1980], which relies on the divided differences introduced in Definition 3.1. We will use the notation introduced in Definition 2.7.

Proposition 3.4 (Kergin interpolation). *Let $\underline{x} \in (\mathbb{R}^n)^p$ and let f be a function of class C^{p-1} defined on some neighborhood of $\sigma(\underline{x})$ in \mathbb{R}^n . There exists a unique polynomial $K(f, \underline{x}) \in \mathbb{R}_{p-1}[X]$ such that, for all nonempty $I \subset \llbracket 1, p \rrbracket$, we have $f[\underline{x}_I] = (K(f, \underline{x}))[\underline{x}_I]$. Moreover,*

$$K(f, \underline{x}) = \sum_{k=1}^p f[x_1, \dots, x_k](X - x_1, \dots, X - x_{k-1}). \tag{3-3}$$

Proof. This is the content of [Bojanov et al. 1993, Theorem 12.5] for $m = 0$. See also [Micchelli and Milman 1980]. □

Remark 3.5. In particular, Proposition 3.4 implies the following:

- The restriction of $K(\cdot, \underline{x})$ to $\mathbb{R}_{p-1}[X]$ is the identity.
- If x appears with multiplicity at least $k + 1$ in \underline{x} , then

$$D_x^k f = k! \underbrace{f[x, \dots, x]}_{k+1 \text{ times}} = k! (K(f, \underline{x})) \underbrace{[x, \dots, x]}_{k+1 \text{ times}} = D_x^k (K(f, \underline{x})).$$

- The map $P \mapsto (P[x_1, \dots, x_j])_{1 \leq j \leq p}$ is an isomorphism from $\mathbb{R}_{p-1}[X]$ to $\bigoplus_{j=0}^{p-1} \text{Sym}^j(\mathbb{R}^n)$ whose inverse map is given by $(S_j)_{0 \leq j \leq p-1} \mapsto \sum_{j=0}^{p-1} S_j(X - x_1, \dots, X - x_j)$.

Definition 3.6 (Kergin polynomial). The polynomial $K(f, \underline{x})$ from Proposition 3.4 is called the *Kergin interpolating polynomial* of f at \underline{x} .

Example 3.7. If $n = 1$, then $K(f, \underline{x})$ is the Hermite interpolating polynomial of f at $\underline{x} \in \mathbb{R}^p$. If $\underline{x} = (x, \dots, x)$, then $K(f, \underline{x})$ is the Taylor polynomial of order $p - 1$ of f at $x \in \mathbb{R}^n$.

Lemma 3.8 (regularity of the Kergin polynomial). *For all $\underline{x} \in (\mathbb{R}^n)^p$, the map $K(\cdot, \underline{x})$ is linear. Moreover, if f is \mathcal{C}^{l+p-1} then $K(f, \cdot)$ is of class \mathcal{C}^l .*

Proof. This is a consequence of Lemma 3.3 and (3-3). □

We need to prove a form of compatibility in Kergin interpolation, when the set of interpolation points is refined. We will use this fact to prove that the multijet bundle we define below satisfies (4) in Theorem 1.1. The following lemma is stated using the clustering partition $\mathcal{I}(\underline{x})$ from Definition 2.3.

Lemma 3.9 (compatibility in Kergin interpolation). *For all $\underline{x} \in (\mathbb{R}^n)^p$ the linear map from $\mathbb{R}_{p-1}[X]$ to $\prod_{I \in \mathcal{I}(\underline{x})} \mathbb{R}_{|I|-1}[X]$ defined by $(K(\cdot, \underline{x}_I))_{I \in \mathcal{I}(\underline{x})} : P \mapsto (K(P, \underline{x}_I))_{I \in \mathcal{I}(\underline{x})}$ is surjective.*

Proof. Let $\underline{x} \in (\mathbb{R}^n)^p$ and let us write $\mathcal{I} = \mathcal{I}(\underline{x})$ for simplicity. As explained at the end of Section 2.1, there exists a unique $\underline{y} = (y_I)_{I \in \mathcal{I}} \in (\mathbb{R}^n)^{\mathcal{I}} \setminus \Delta_{\mathcal{I}}$ such that $\underline{x} = \iota_{\mathcal{I}}(\underline{y})$. Let $(\chi_I)_{I \in \mathcal{I}}$ be smooth functions on \mathbb{R}^n with pairwise disjoint compact supports and such that χ_I is equal to 1 in a neighborhood of y_I .

Let $(P_I)_{I \in \mathcal{I}} \in \prod_{I \in \mathcal{I}} \mathbb{R}_{|I|-1}[X]$. We consider the function $f = \sum_{I \in \mathcal{I}} \chi_I P_I \in \mathcal{C}^\infty(\mathbb{R}^n)$. Let $P = K(f, \underline{x})$ and let us prove that $K(P, \underline{x}_I) = P_I$ for all $I \in \mathcal{I}$. For all $k \leq |I| - 1$ we have $D_{y_I}^k P = D_{y_I}^k f = D_{y_I}^k P_I$. Indeed y_I appears with multiplicity $|I|$ in \underline{x} (see Remark 3.5) and f is equal to P_I in a neighborhood of y_I . Recalling Example 3.7, we know that $K(P, \underline{x}_I)$ is the Taylor polynomial of order $|I| - 1$ at y_I of P , and hence of P_I . Since $P_I \in \mathbb{R}_{|I|-1}[X]$, we get $K(P, \underline{x}_I) = P_I$. □

4. Evaluation maps and their kernels

The goal of this section is to study evaluation maps on spaces of polynomials and their kernels. Defining multijets is closely related to these objects. Indeed, let $n \geq 1$ and $p \geq 1$ and recall that Δ_p stands for the large diagonal in $(\mathbb{R}^n)^p$; see Definition 2.1. As explained in the Introduction, when $\underline{x} \notin \Delta_p$ we want the multijet of a \mathcal{C}^{p-1} function f at \underline{x} to be the class of f in $\mathcal{C}^{p-1}(\mathbb{R}^n) / \sim$, where $f \sim g$ if and only if $(f(x_i))_{1 \leq i \leq p} = (g(x_i))_{1 \leq i \leq p}$. The Kergin interpolation of Section 3.2 shows that any such class can be represented by a polynomial. Hence, the space of p -multijets at \underline{x} is canonically isomorphic to $\mathbb{R}_{p-1}[X] / \ker \text{ev}_{\underline{x}}$, where $\text{ev}_{\underline{x}} : P \mapsto (P(x_1), \dots, P(x_p))$.

Definition 4.1 (Grassmannian). Let V be a vector space of finite dimension N and $k \in \llbracket 0, N \rrbracket$. We denote by $\text{Gr}_k(V)$ the *Grassmannian* of vector subspaces of V of *codimension* k .

Remark 4.2. Beware that this notation is slightly unusual, since in most textbooks $\text{Gr}_k(V)$ stands for the Grassmannian of subspaces of dimension k .

Let us denote by $\mathcal{L}_{\text{reg}}(V, \mathbb{R}^k) \subset V^* \otimes \mathbb{R}^k$ the open dense subset of linear surjective maps from V to \mathbb{R}^k . The group $\text{GL}_k(\mathbb{R})$ acts on $\mathcal{L}_{\text{reg}}(V, \mathbb{R}^k)$ by multiplication on the left. On the other hand, $L \mapsto \ker(L)$ defines a surjective map from $\mathcal{L}_{\text{reg}}(V, \mathbb{R}^k)$ to $\text{Gr}_k(V)$, and $\ker(L_1) = \ker(L_2)$ if and only if there exists $M \in \text{GL}_k(\mathbb{R})$ such that $L_2 = ML_1$. Thus, one can identify $\text{Gr}_k(V)$ with the orbit space $\mathcal{L}_{\text{reg}}(V, \mathbb{R}^k) / \text{GL}_k(\mathbb{R})$ of the previous action. This is one of the many ways to describe $\text{Gr}_k(V)$ as a smooth real-algebraic manifold.

Definition 4.3 (evaluation map). Let $\underline{x} \in (\mathbb{R}^n)^p$. We set $\text{ev}_{\underline{x}} : f \mapsto (f(x_1), \dots, f(x_p))$ from any space of functions defined at the x_i to \mathbb{R}^p . The source space will always be clear from the context.

Lemma 4.4 (nondegeneracy of $\text{ev}_{\underline{x}}$). Let $\underline{x} \notin \Delta_p$. Then $\text{ev}_{\underline{x}} : \mathbb{R}_{p-1}[X] \rightarrow \mathbb{R}^p$ is surjective.

Proof. Since $\underline{x} \notin \Delta_p$, we have $\mathcal{I}(\underline{x}) = \{\{1\}, \dots, \{p\}\}$ and $\text{ev}_{\underline{x}} = (K(\cdot, x_i))_{1 \leq i \leq p}$ under the canonical identification $\mathbb{R}_0[X] \simeq \mathbb{R}$. Hence this is just a special case of Lemma 3.9. Alternatively, in the right basis, one can extract a Vandermonde matrix from that of $\text{ev}_{\underline{x}}$. □

Lemma 4.4 shows that the following map is well-defined from $(\mathbb{R}^n)^p \setminus \Delta_p$ to $\text{Gr}_p(\mathbb{R}_{p-1}[X])$:

$$\mathcal{G} : \underline{x} \mapsto \ker \text{ev}_{\underline{x}}. \tag{4-1}$$

Lemma 4.5 (algebraicity of \mathcal{G}). The map $\mathcal{G} : (\mathbb{R}^n)^p \setminus \Delta_p \rightarrow \text{Gr}_p(\mathbb{R}_{p-1}[X])$ is algebraic.

Proof. Recalling the previous discussion, we have $\mathcal{L}_{\text{reg}}(\mathbb{R}_{p-1}[X], \mathbb{R}^p) / \text{GL}_p(\mathbb{R}) \simeq \text{Gr}_p(\mathbb{R}_{p-1}[X])$, where the isomorphism is obtained as the quotient map of $\ker : L \mapsto \ker(L)$. In particular,

$$\ker : \mathcal{L}_{\text{reg}}(\mathbb{R}_{p-1}[X], \mathbb{R}^p) \longrightarrow \text{Gr}_p(\mathbb{R}_{p-1}[X]) \simeq \mathcal{L}_{\text{reg}}(\mathbb{R}_{p-1}[X], \mathbb{R}^p) / \text{GL}_p(\mathbb{R})$$

is just the canonical projection, which is algebraic.

Writing $\text{ev} : \underline{x} \mapsto \text{ev}_{\underline{x}}$, we have $\mathcal{G} = \ker \circ \text{ev}$. Thus it is enough to prove that ev is algebraic from $(\mathbb{R}^n)^p \setminus \Delta_p$ to $\mathcal{L}_{\text{reg}}(\mathbb{R}_{p-1}[X], \mathbb{R}^p)$. In the basis of $\mathbb{R}_{p-1}[X]$ formed by the monomials $(X^\alpha)_{|\alpha| < p}$, the matrix of $\text{ev}_{\underline{x}}$ is $(x_i^\alpha)_{1 \leq i \leq p; |\alpha| < p}$, which depends algebraically on \underline{x} . □

Let $\underline{x} \in (\mathbb{R}^n)^p \setminus \Delta_p$, we defined $\mathcal{G}(\underline{x}) \in \text{Gr}_p(\mathbb{R}_{p-1}[X])$ by (4-1). For any nonempty $I \subset \llbracket 1, p \rrbracket$, we define similarly

$$\mathcal{G}_I(\underline{x}) = \ker \text{ev}_{\underline{x}_I} \in \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X]) \quad \text{and} \quad \tilde{\mathcal{G}}_I(\underline{x}) = \ker \text{ev}_{\underline{x}_I} \in \text{Gr}_{|I|}(\mathbb{R}_{p-1}[X]). \tag{4-2}$$

Because of the interpolation properties of the Kergin polynomials (see Remark 3.5), we have that $\text{ev}_{\underline{x}_I} = (\text{ev}_{\underline{x}_I})_{|\mathbb{R}_{|I|-1}[X]} \circ K(\cdot, \underline{x}_I)$ on $\mathbb{R}_{p-1}[X]$. Hence $\tilde{\mathcal{G}}_I(\underline{x}) = K(\cdot, \underline{x}_I)^{-1}(\mathcal{G}_I(\underline{x}))$. Since $K(\cdot, \underline{x}_I)$ is surjective from $\mathbb{R}_{p-1}[X]$ to $\mathbb{R}_{|I|-1}[X]$, this shows that $\tilde{\mathcal{G}}_I(\underline{x})$ has indeed codimension $|I|$, like $\mathcal{G}_I(\underline{x})$.

This collection of subspaces satisfies some incidence relations that will be useful in the following. For all nonempty $I \subset \llbracket 1, p \rrbracket$, we have $\mathcal{G}(\underline{x}) \subset \tilde{\mathcal{G}}_I(\underline{x})$. Actually, we can be more precise: for any $\mathcal{I} \in \mathcal{P}_p$, we have $\mathcal{G}(\underline{x}) = \bigcap_{I \in \mathcal{I}} \tilde{\mathcal{G}}_I(\underline{x})$, and this intersection is transverse by a codimension argument.

Remark 4.6. The map $\mathcal{G} : (\mathbb{R}^n)^p \setminus \Delta_p \rightarrow \text{Gr}_p(\mathbb{R}_{p-1}[X])$ does not admit an extension as a regular map from $(\mathbb{R}^n)^p$ to $\text{Gr}_p(\mathbb{R}_{p-1}[X])$, except if $n = 1$ or $p = 1$, that is, if $\text{Gr}_p(\mathbb{R}_{p-1}[X])$ is a point.

For example, when $n = 2 = p$, the Grassmannian $\text{Gr}_p(\mathbb{R}_{p-1}[X])$ is the set of lines in $\mathbb{R}_1[X_1, X_2]$. Taking $x = R(\cos \theta, \sin \theta) \in \mathbb{R}^2 \setminus \{0\}$, the reader can check that $\mathcal{G}(0, x) = \text{Span}(X_1 \sin \theta - X_2 \cos \theta)$, which does not converge as $R \rightarrow 0$. However, in this case, $\mathcal{G}(0, \cdot)$ extends to the blow-up $\text{Bl}_0(\mathbb{R}^2)$ of \mathbb{R}^2 at 0 and similarly \mathcal{G} extends smoothly to $\text{Bl}_{\Delta_2}((\mathbb{R}^2)^2)$. This suggests that, even though \mathcal{G} does not extend smoothly to $(\mathbb{R}^n)^p$, it might extend to a larger space.

5. Definition of the multijet bundles

In this section we define the vector bundle $\mathcal{MJ}_p(\mathbb{R}^n, V) \rightarrow C_p[\mathbb{R}^n]$ of p -multijets for functions from \mathbb{R}^n to some finite-dimensional vector space V and prove Theorem 1.1. The singularity of \mathcal{G} along Δ_p makes it impossible to define such a bundle over $(\mathbb{R}^n)^p$, which is why we define it over a compactification $C_p[\mathbb{R}^n]$ of the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p$.

The manifold $C_p[\mathbb{R}^n]$ does not depend on V . It is defined in Section 5.1. In the next two sections, we work in the case $V = \mathbb{R}$. All important ideas appear in this case but the notation is slightly simpler. In Section 5.2, we define the bundle $\mathcal{MJ}_p(\mathbb{R}^n)$. In Section 5.3, we prove that p -multijets are local, in the sense of (4) in Theorem 1.1. Finally, we define the bundle $\mathcal{MJ}_p(\mathbb{R}^n, V)$ of multijets for vector-valued maps and prove Theorem 1.1 in Section 5.4.

5.1. Definition of the basis $C_p[\mathbb{R}^n]$. In this section, we define the basis $C_p[\mathbb{R}^n]$ over which our p -multijet bundles are defined. This is a smooth manifold, obtained a compactification of the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p$ such that $(\mathcal{G}_I)_{I \subset \llbracket 1, p \rrbracket}$ extends smoothly to $C_p[\mathbb{R}^n]$. Let us first introduce some notation. We denote by Π_0 the projection from the product space

$$(\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$$

onto the factor $(\mathbb{R}^n)^p$. Similarly, we denote by Π_I the projection onto $\text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$. Then, let

$$\Sigma = \{(\underline{x}, (\mathcal{G}_I(\underline{x}))_{I \subset \llbracket 1, p \rrbracket}) \mid \underline{x} \in (\mathbb{R}^n)^p \setminus \Delta_p\} \subset (\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X]) \tag{5-1}$$

denote the graph of the map $(\mathcal{G}_I)_{I \subset \llbracket 1, p \rrbracket}$. We denote by $\bar{\Sigma}$ the closure of Σ in the product space on the right-hand side of (5-1).

Lemma 5.1 (surjectivity of $(\Pi_0)_{|\bar{\Sigma}}$). *Let $\underline{x} \in (\mathbb{R}^n)^p$. Then there exists $z \in \bar{\Sigma}$ such that $\Pi_0(z) = \underline{x}$.*

Proof. Let $(\underline{x}_n)_{n \in \mathbb{N}}$ be a sequence of points in $(\mathbb{R}^n)^p \setminus \Delta_p$ converging to \underline{x} . Since Grassmannians are compact manifolds, up to extracting subsequences finitely many times, we can assume that for all nonempty $I \subset \llbracket 1, p \rrbracket$ there exists $G_I \in \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$ such that $\mathcal{G}_I(\underline{x}_n) \xrightarrow{n \rightarrow +\infty} G_I$. Then

$$(\underline{x}_n, (\mathcal{G}_I(\underline{x}_n))_{I \subset \llbracket 1, p \rrbracket}) \xrightarrow{n \rightarrow +\infty} (\underline{x}, (G_I)_{I \subset \llbracket 1, p \rrbracket}) = z \in \bar{\Sigma}. \quad \square$$

Lemma 5.2 (location of the new points). *We have $\bar{\Sigma} \setminus \Sigma \subset \Pi_0^{-1}(\Delta_p)$.*

Proof. Since Σ is the graph of a continuous function on $(\mathbb{R}^n)^p \setminus \Delta_p$, it is closed in the open subset $\Pi_0^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$. Hence $\bar{\Sigma} \cap \Pi_0^{-1}((\mathbb{R}^n)^p \setminus \Delta_p) = \Sigma$ and $\bar{\Sigma} \setminus \Sigma \subset \Pi_0^{-1}(\Delta_p)$. \square

Lemma 5.3 (algebraicity of Σ and $\bar{\Sigma}$). *The graph Σ is a smooth real-algebraic manifold and $(\Pi_0)_{|\Sigma} : \Sigma \rightarrow (\mathbb{R}^n)^p \setminus \Delta_p$ is an isomorphism. Moreover, $\bar{\Sigma}$ is a real-algebraic variety whose singular locus is contained in $\bar{\Sigma} \setminus \Sigma$.*

Proof. By Lemma 4.5, the set Σ is the graph of an algebraic map, hence a smooth real-algebraic manifold. Additionally, Π_0 is algebraic and its restriction to Σ is the inverse of $\underline{x} \mapsto (\underline{x}, (\mathcal{G}_I(\underline{x}))_{I \subset \llbracket 1, p \rrbracket})$. Thus

$(\Pi_0)_{|\Sigma}$ is an algebraic isomorphism from Σ onto $(\mathbb{R}^n)^p \setminus \Delta_p$. Since Σ is real-algebraic, so is its closure $\bar{\Sigma}$. By Lemma 5.2, we know that $\bar{\Sigma} \cap \Pi_0^{-1}((\mathbb{R}^n)^p \setminus \Delta_p) = \Sigma$ is smooth. Hence, the singular locus of $\bar{\Sigma}$ is contained in $\bar{\Sigma} \cap \Pi_0^{-1}(\Delta_p) = \bar{\Sigma} \setminus \Sigma$. \square

Example 5.4. In simple cases, we understand very well what $\bar{\Sigma}$ is.

- If $p = 1$ and $n \geq 1$, then $\Delta_p = \emptyset$ and $\text{Gr}_p(\mathbb{R}_{p-1}[X]) = \{\{0\}\}$, so that $\bar{\Sigma} = \Sigma = \mathbb{R}^n$.
- If $n = 1$ and $p \geq 1$, then $\text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X]) = \{\{0\}\}$ for all $I \subset \llbracket 1, p \rrbracket$ and $\bar{\Sigma} = \mathbb{R}^p$.
- If $p = 2$ and $n \geq 2$, then for $x \neq y$ in \mathbb{R}^n we know that $\mathcal{G}(x, y) \subset \mathbb{R}_1[X]$ is the subspace of affine forms on \mathbb{R}^n vanishing at x and y , i.e., on the affine line through x and y . Thus $\mathcal{G}(x, y)$ encodes this line. As $y \rightarrow x$, the accumulation points of $\mathcal{G}(x, y)$ correspond to all the affine lines passing through x , and they encode “the direction from which y converges to x ”. In this case, one can check that $\bar{\Sigma} = \text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$.

In the previous examples the variety $\bar{\Sigma}$ is smooth, hence the following natural question.

Question. *Is $\bar{\Sigma}$ smooth for all $n \geq 1$ and $p \geq 1$?*

Lacking a positive answer to this question, since we want $C_p[\mathbb{R}^n]$ to be a smooth manifold, we will define it by resolving the singularities of $\bar{\Sigma}$. The existence of a resolution of singularities is given by Hironaka’s theorem [1964a; 1964b]. Our references on this matter are [Kollár 2007; Włodarczyk 2005]. See also [Hauser 2003] for a softer introduction to this theory.

Proposition 5.5 (resolution of singularities). *There exists a smooth manifold $C_p[\mathbb{R}^n]$ without boundary of dimension np and a smooth proper $\Pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$ such that*

- (1) $\Pi(C_p[\mathbb{R}^n]) = \bar{\Sigma}$;
- (2) $\Pi^{-1}(\Sigma)$ is an open dense subset of $C_p[\mathbb{R}^n]$;
- (3) $\Pi|_{\Pi^{-1}(\Sigma)}$ is C^∞ -diffeomorphism from $\Pi^{-1}(\Sigma)$ onto Σ .

Proof. We apply Hironaka’s theorem [Kollár 2007, Theorem 3.27] to resolve the singularities of $\bar{\Sigma}$. Since $\bar{\Sigma}$ is algebraic by Lemma 5.3, there exists a smooth real-algebraic manifold $C_p[\mathbb{R}^n]$ and a projective morphism $\Pi : C_p[\mathbb{R}^n] \rightarrow \bar{\Sigma}$ such that Π is an isomorphism over the smooth locus of $\bar{\Sigma}$.

In particular $C_p[\mathbb{R}^n]$ is smooth, the map $\Pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$ is smooth and proper, and $\Pi(C_p[\mathbb{R}^n]) \subset \bar{\Sigma}$. Since Σ is contained in the smooth locus of $\bar{\Sigma}$, the restriction of Π to $\Pi^{-1}(\Sigma)$ is an isomorphism; in particular (3) is satisfied.

According to [Włodarczyk 2005, Theorem 1.0.2], the manifold $C_p[\mathbb{R}^n]$ and the projection Π are obtained by a sequence of blow-ups along smooth submanifolds that do not intersect the regular locus of $\bar{\Sigma}$, and hence Σ . This ensures that conditions (1) and (2) are satisfied. \square

The following corollary proves the existence of the manifold $C_p[\mathbb{R}^n]$ and the proper surjection $\pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ satisfying (1) in Theorem 1.1.

Corollary 5.6 (existence of the basis $C_p[\mathbb{R}^n]$). *There exists a smooth manifold $C_p[\mathbb{R}^n]$ without boundary of dimension np and a smooth proper surjection $\pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ such that*

- (1) *the open subset $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ is dense in $C_p[\mathbb{R}^n]$ and π induces a C^∞ -diffeomorphism from this set onto $(\mathbb{R}^n)^p \setminus \Delta_p$;*
- (2) *for any nonempty $I \subset \llbracket 1, p \rrbracket$, the map $\mathcal{G}_I \circ \pi$ admits a unique smooth extension to $C_p[\mathbb{R}^n]$.*

Proof. We consider $\Pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$ given by Proposition 5.5 and we let $\pi = \Pi_0 \circ \Pi$. Since Grassmannians are compact, Π_0 is proper. Hence π is smooth and proper because Π and Π_0 are. The surjectivity of π is given by Lemma 5.1 and (1) in Proposition 5.5.

Item (1) in Corollary 5.6 is a consequence of Lemmas 5.2 and 5.3 and of conditions (2) and (3) in Proposition 5.5. Let $I \subset \llbracket 1, p \rrbracket$ be nonempty. On the dense open subset $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ we have $\mathcal{G}_I \circ \pi = \Pi_I \circ \Pi$ by definition. In the last equality, the right-hand side is well-defined and smooth on $C_p[\mathbb{R}^n]$, which yields the unique extension we are looking for. □

Since it is defined using Hironaka’s theorem, the manifold $C_p[\mathbb{R}^n]$ is not unique. However, the value of the smooth extension of $\mathcal{G}_I \circ \pi = \Pi_I \circ \Pi$ at $z \in C_p[\mathbb{R}^n]$ only depends on $\Pi(z) \in \bar{\Sigma}$. So this extension does not really depend on the choice of a resolution of singularities. In the following we choose once and for all a realization of $\pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ as in Corollary 5.6. Thanks to (1), we can identify the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p$ with its open dense preimage by π . Under this identification, (2) states that the maps $(\mathcal{G}_I)_{I \subset \llbracket 1, p \rrbracket}$ extend smoothly to $C_p[\mathbb{R}^n]$. So, from now on, we consider \mathcal{G}_I as a smooth map from $C_p[\mathbb{R}^n]$ to $\text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$.

5.2. Definition of the bundle $\mathcal{MJ}_p(\mathbb{R}^n)$. Now that we have defined the base space $C_p[\mathbb{R}^n]$ of our multijet bundle, we can define the bundle itself. The purpose of this section is to construct the vector bundle $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ of multijets for functions from \mathbb{R}^n to \mathbb{R} , and the associated multijet map. The construction for vector-valued maps, explained in Section 5.4, is basically a fiberwise direct sum of this simpler case.

Recall that we defined the projections

$$C_p[\mathbb{R}^n] \xrightarrow{\Pi} \bar{\Sigma} \xrightarrow{\Pi_0} (\mathbb{R}^n)^p$$

and that $\pi = \Pi_0 \circ \Pi$. Thanks to Corollary 5.6, and under the identification discussed above, the map $\mathcal{G} = \mathcal{G}_{\llbracket 1, p \rrbracket}$ defined by (4-1) extends as a smooth map from $C_p[\mathbb{R}^n]$ to $\text{Gr}_p(\mathbb{R}_{p-1}[X])$. Seen as a collection of subspaces of $\mathbb{R}_{p-1}[X]$ indexed by $C_p[\mathbb{R}^n]$, this means that \mathcal{G} defines a smooth vector sub-bundle of corank p in the trivial bundle $\mathbb{R}_{p-1}[X] \times C_p[\mathbb{R}^n] \rightarrow C_p[\mathbb{R}^n]$. We define our multijet bundle as the quotient of this trivial bundle by \mathcal{G} .

Definition 5.7 (vector bundle of multijets). Let $n \geq 1$ and $p \geq 1$. The *vector bundle of multijets of order p on \mathbb{R}^n* is the smooth vector bundle of rank p over $C_p[\mathbb{R}^n]$ defined by

$$\mathcal{MJ}_p(\mathbb{R}^n) = (\mathbb{R}_{p-1}[X] \times C_p[\mathbb{R}^n]) / \mathcal{G}.$$

In particular, for any $z \in C_p[\mathbb{R}^n]$, the fiber $\mathcal{MJ}_p(\mathbb{R}^n)_z = \mathbb{R}_{p-1}[X] / \mathcal{G}(z)$ only depends on $\Pi(z) \in \bar{\Sigma}$.

Recalling the definition of Kergin polynomials given in Section 3.2, we can now define the p -multijet of a C^{p-1} function on \mathbb{R}^n .

Definition 5.8 (multijet of a function). Let $f \in \mathcal{C}^{p-1}(\mathbb{R}^n)$ and $z \in C_p[\mathbb{R}^n]$. The *multijet of f at z* is the element of $\mathcal{MJ}_p(\mathbb{R}^n)_z$ defined as

$$\text{mj}_p(f, z) = K(f, \pi(z)) \bmod \mathcal{G}(z).$$

In particular, as an element of $\mathbb{R}_{p-1}[X]/\mathcal{G}(z)$, the multijet $\text{mj}_p(f, z)$ only depends on $\Pi(z) \in \bar{\Sigma}$.

Example 5.9. In Example 5.4 we saw that in simple cases $\bar{\Sigma}$ is smooth. In these cases we set $C_p[\mathbb{R}^n] = \bar{\Sigma}$ and we can describe the bundle $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ and the map mj_p .

- If $p = 1$, then $C_1[\mathbb{R}^n] = \mathbb{R}^n$ and $\mathcal{G} : x \mapsto \{0\} \subset \mathbb{R}_0[X] \simeq \mathbb{R}$. Thus $\mathcal{MJ}_1(\mathbb{R}^n)$ is the trivial bundle $\mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$. Moreover, if $f \in \mathcal{C}^0(\mathbb{R}^n)$ then $K(f, x) = f(x) \in \mathbb{R}_0[X] \simeq \mathbb{R}$ and $\text{mj}_1(f, x) = f(x)$ for all $x \in \mathbb{R}^n$.
- If $n = 1$, then $C_p[\mathbb{R}] = \mathbb{R}^p$ and $\mathcal{G} : \underline{x} \mapsto \{0\} \subset \mathbb{R}_{p-1}[X]$. Thus $\mathcal{MJ}_p(\mathbb{R})$ is the trivial bundle $\mathbb{R}_{p-1}[X] \times \mathbb{R}^p \rightarrow \mathbb{R}^p$. If $f \in \mathcal{C}^{p-1}(\mathbb{R})$ then $\text{mj}_p(f, \underline{x}) = K(f, \underline{x})$ is the Hermite interpolating polynomial of f at \underline{x} ; see Example 3.7.

Given $\underline{x} = (x_1, \dots, x_p) \notin \Delta_p$, Lemma 4.4 shows that $\text{ev}_{\underline{x}} : \mathbb{R}_{p-1}[X] \rightarrow \mathbb{R}^p$ is an isomorphism. We can then consider the Lagrange basis $(L_i(\underline{x}))_{1 \leq i \leq p}$ of $\mathbb{R}_{p-1}[X]$ which is the preimage by $\text{ev}_{\underline{x}}$ of the canonical basis of \mathbb{R}^p . We then have $\text{mj}_p(f, \underline{x}) = K(f, \underline{x}) = \sum_{i=1}^p f(x_i)L_i(\underline{x})$. Geometrically, this means that the map $(P, \underline{x}) \mapsto (\text{ev}_{\underline{x}}(P), \underline{x})$ defines a local trivialization of $\mathcal{MJ}_p(\mathbb{R}) \rightarrow C_p[\mathbb{R}]$ over $\mathbb{R}^p \setminus \Delta_p$ and that $\underline{x} \mapsto (L_i(\underline{x}))_{1 \leq i \leq p}$ is the corresponding frame. Moreover, it is tautological that $\text{mj}_p(f, \underline{x})$ reads as $(f(x_1), \dots, f(x_p))$ in this trivialization.

In this example, one can also define a global trivialization of $\mathcal{MJ}_p(\mathbb{R})$ by considering the global frame of Newton polynomials $\underline{x} \mapsto (N_k(\underline{x}))_{1 \leq k \leq p}$, where $N_k(\underline{x}) = \prod_{1 \leq i < k} (X - x_i)$. By (3-3) we have $K(f, \underline{x}) = \sum_{k=1}^p f[x_1, \dots, x_k]N_k(\underline{x})$, so that $\text{mj}_p(f, \underline{x})$ reads as $(f[x_1, \dots, x_k])_{1 \leq k \leq p}$ in this trivialization, where the divided differences are the classical ones in dimension 1. In this setting, we used in [Ancona and Letendre 2021] a strategy that can be roughly summarized as replacing $(f(x_i))_{1 \leq i \leq p}$ by $(f[x_1, \dots, x_k])_{1 \leq k \leq p}$. Our present point of view shows that we were actually considering $\text{mj}_p(f, \underline{x})$ all along, but read in different trivializations.

- If $p = 2$, we saw that $C_2[\mathbb{R}^n] = \text{Bl}_{\Delta_2}((\mathbb{R}^n)^2)$. Given $z \in C_2[\mathbb{R}^n]$, if $\pi(z) = (x_1, x_2) \notin \Delta_2$, we know that $\mathcal{G}(z) \subset \mathbb{R}_1[X]$ is the subspace of affine forms vanishing on the line $L_z \subset \mathbb{R}^n$ through x_1 and x_2 . It is then natural to think of the class of P modulo $\mathcal{G}(z)$ as its restriction to L_z . Parametrizing L_z by

$$t \mapsto x_1 + \frac{x_2 - x_1}{\|x_2 - x_1\|}t,$$

one can check that

$$P \mapsto P\left(x_1 + \frac{x_2 - x_1}{\|x_2 - x_1\|}T\right)$$

induces an isomorphism $\mathcal{MJ}_2(\mathbb{R}^n)_z \simeq \mathbb{R}_1[T] \simeq \mathbb{R}^2$, where T is univariate and the second isomorphism is obtained by reading coordinates in the canonical basis $(1, T)$ of $\mathbb{R}_1[T]$.

Recalling Definition 3.1, we have

$$P[x_1, x_2] \cdot \frac{x_2 - x_1}{\|x_2 - x_1\|}T = \left(\int_0^1 D_{x_1+t(x_2-x_1)}P \cdot (x_2 - x_1) dt\right) \frac{T}{\|x_2 - x_1\|} = \frac{P(x_2) - P(x_1)}{\|x_2 - x_1\|}T,$$

and $P = K(P, x_1, x_2)$ is given by (3-3). Letting $\tilde{P}(z) = P(x_2) - P(x_1)/\|x_2 - x_1\|$, we have

$$P\left(x_1 + \frac{x_2 - x_1}{\|x_2 - x_1\|}T\right) = K(P, x_1, x_2)\left(x_1 + \frac{x_2 - x_1}{\|x_2 - x_1\|}T\right) = P(x_1) + \tilde{P}(z)T.$$

Thus, the previous isomorphism $\mathcal{MJ}_2(\mathbb{R}^n)_z \rightarrow \mathbb{R}^2$ is $(P \bmod \mathcal{G}(z)) \mapsto (P(\pi(z)_1), \tilde{P}(z))$.

Let us now consider $z \in \pi^{-1}(\Delta_2)$. This exceptional divisor is the projectivized normal bundle of Δ_2 in $(\mathbb{R}^n)^2$. So we can think of z as a point in the diagonal, say $(x, x) \in \Delta_2$, and a line in $(\mathbb{R}^n)^2$ which is orthogonal to Δ_2 , say spanned by $(u, -u)$ with $u \in \mathbb{S}^{n-1}$. Then $z = \lim_{\epsilon \rightarrow 0}(x + \epsilon u, x - \epsilon u)$ in $C_2[\mathbb{R}^n]$. By continuity, $\mathcal{G}(z)$ is the space of affine forms vanishing on the line $L_z \subset \mathbb{R}^n$ parametrized by $t \mapsto x + tu$. As above, mapping P to the coefficients of $P(x + Tu) = P(x) + (D_x P \cdot u)T \in \mathbb{R}_1[T]$ induces an isomorphism $\mathcal{MJ}_2(\mathbb{R}^n)_z \rightarrow \mathbb{R}^2$. Letting $\tilde{P}(z) = D_x P \cdot u$, this isomorphism is again $(P \bmod \mathcal{G}(z)) \mapsto (P(\pi(z)_1), \tilde{P}(z))$.

Actually, one can check that everything depends smoothly on the base point $z \in C_2[\mathbb{R}^n]$, so that the bundle map $(P \bmod \mathcal{G}(z), z) \mapsto (P(\pi(z)_1), \tilde{P}(z), z)$ defines a global trivialization $\mathcal{MJ}_2(\mathbb{R}^n) \rightarrow \mathbb{R}^2 \times C_2[\mathbb{R}^n]$. If $f \in C^1(\mathbb{R}^n)$, with the same notation as above, $\text{mj}_2(f, z)$ reads in this trivialization as

$$\left(f(x_1), \frac{f(x_2) - f(x_1)}{\|x_2 - x_1\|}\right)$$

if $z \notin \pi^{-1}(\Delta_2)$ and as $(f(x), D_x f \cdot u)$ otherwise.

In these examples, the multijet bundle $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ is trivial. This raises the following.

Question. *Is $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ trivial for all $n \geq 1$ and $p \geq 1$?*

The following two lemmas prove that the bundle map $\text{mj}_p : C^{p-1}(\mathbb{R}^n) \times C_p[\mathbb{R}^n] \rightarrow \mathcal{MJ}_p(\mathbb{R}^n)$ satisfies (2) and (3) in Theorem 1.1.

Lemma 5.10 (regularity of multijets). *The map $\text{mj}_p(\cdot, z) : C^{p-1}(\mathbb{R}^n) \rightarrow \mathcal{MJ}_p(\mathbb{R}^n)_z$ is a linear surjection for all $z \in C_p[\mathbb{R}^n]$. Additionally, let $l \geq 0$ and let $f \in C^{l+p-1}(\mathbb{R}^n)$. Then $\text{mj}_p(f, \cdot)$ is a section of class C^l of $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$.*

Proof. Let $z \in C_p[\mathbb{R}^n]$. The map $K(\cdot, \pi(z)) : C^{p-1}(\mathbb{R}^n) \rightarrow \mathbb{R}_{p-1}[X]$ is linear by Lemma 3.8. It is also surjective since its restriction to $\mathbb{R}_{p-1}[X]$ is the identity. Since $\text{mj}_p(\cdot, z)$ is the composition of $K(\cdot, \pi(z))$ with the canonical projection from $\mathbb{R}_{p-1}[X]$ onto $\mathcal{MJ}_p(\mathbb{R}^n)_z$, it is a linear surjection.

Let $l \geq 0$ and let $f \in C^{l+p-1}(\mathbb{R}^n)$. By Lemma 3.8, we have $K(f, \cdot) \in C^l((\mathbb{R}^n)^p, \mathbb{R}_{p-1}[X])$. Since π is smooth, we get $K(f, \cdot) \circ \pi \in C^l(C_p[\mathbb{R}^n], \mathbb{R}_{p-1}[X])$. In other words, $K(f, \cdot) \circ \pi$ defines a section of class C^l of the trivial bundle $\mathbb{R}_{p-1}[X] \times C_p[\mathbb{R}^n] \rightarrow C_p[\mathbb{R}^n]$. Since \mathcal{G} is a smooth sub-bundle of $\mathbb{R}_{p-1}[X] \times C_p[\mathbb{R}^n]$, projecting onto the quotient bundle $\mathcal{MJ}_p(\mathbb{R}^n)$ does not decrease the regularity. \square

Lemma 5.11 (multijets and evaluation). *Let $z \in C_p[\mathbb{R}^n]$ be such that $\pi(z) = (x_1, \dots, x_p) \notin \Delta_p$. Then for all $f \in C^{p-1}(\mathbb{R}^n)$ we have $\text{mj}_p(f, z) = 0$ if and only if, for all $i \in \llbracket 1, p \rrbracket$, $f(x_i) = 0$.*

Proof. Let us denote by $\underline{x} = (x_1, \dots, x_p) = \pi(z) \notin \Delta_p$. For all $f \in C^{p-1}(\mathbb{R}^n)$, we have

$$\text{mj}_p(f, z) = 0 \iff K(f, \underline{x}) \in \mathcal{G}(\underline{x}) \iff \text{ev}_{\underline{x}}(K(f, \underline{x})) = 0 \iff \text{ev}_{\underline{x}}(f) = 0,$$

since $K(f, \underline{x})$ interpolates the values of f on at x_1, \dots, x_p (see Remark 3.5). \square

Actually, we can describe more precisely the relation between multijets and evaluation outside of the diagonal. This will appear in the proof of Theorem 6.26 below. Thanks to Lemma 4.4, the smooth bundle map $\text{ev} : (P, \underline{x}) \mapsto (\text{ev}_{\underline{x}}(P), \underline{x})$ from $\mathbb{R}_{p-1}[X] \times ((\mathbb{R}^n)^p \setminus \Delta_p)$ to $\mathbb{R}^p \times ((\mathbb{R}^n)^p \setminus \Delta_p)$ is surjective and its kernel is exactly the sub-bundle $\ker \text{ev} = \mathcal{G}$. Thus it induces a smooth bundle map $\tau : \mathcal{MJ}_p(\mathbb{R}^n)_{|(\mathbb{R}^n)^p \setminus \Delta_p} \rightarrow \mathbb{R}^p \times ((\mathbb{R}^n)^p \setminus \Delta_p)$ defined by $\tau(P \bmod \mathcal{G}(\underline{x})) = (\text{ev}_{\underline{x}}(P), \underline{x})$, which is bijective. Thus τ defines a smooth local trivialization of $\mathcal{MJ}_p(\mathbb{R}^n)$ over $(\mathbb{R}^n)^p \setminus \Delta_p$. Moreover, for all $f \in \mathcal{C}^{p-1}(\mathbb{R}^n)$ and $z \in C_p[\mathbb{R}^n]$ such that $\underline{x} = \pi(z) \notin \Delta_p$ we have

$$\tau(\text{mj}_p(f, z)) = \tau(K(f, \underline{x}) \bmod \mathcal{G}(\underline{x})) = (\text{ev}_{\underline{x}}(K(f, \underline{x})), \underline{x}) = (\text{ev}_{\underline{x}}(f), \underline{x}).$$

Hence $\text{mj}_p(f, z)$ simply reads as $(f(x_1), \dots, f(x_p))$ in this trivialization.

5.3. Localness of multijets. The goal of this section is to prove that the multijet bundle $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ defined in the previous section satisfies (4) in Theorem 1.1. Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$ and let $\mathcal{I} = \mathcal{I}(\underline{x})$ be as in Definition 2.3. As explained in Section 2.1, there is a unique $\underline{y} = (y_I)_{I \in \mathcal{I}} \in \mathbb{R}^{\mathcal{I}} \setminus \Delta_{\mathcal{I}}$ such that $\underline{x} = \iota_{\mathcal{I}}(\underline{y})$. Recalling that we dropped $V = \mathbb{R}$ from the notation in the present case, we can restate (4) in Theorem 1.1 as: there exists $\Theta_z : \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \rightarrow \mathcal{MJ}_p(\mathbb{R}^n)_z$ a linear surjection such that $\text{mj}_p(f, z) = \Theta_z((j_{|I|-1}(f, y_I))_{I \in \mathcal{I}})$ for all $f \in \mathcal{C}^{p-1}(\mathbb{R}^n)$.

This property is fundamental. First, it shows that $\text{mj}_p(f, z)$ is obtained by patching together (part of) the jets of order $|I| - 1$ of f at y_I , which justifies the name multijet. Second, it shows that $\text{mj}_p(f, z)$ only depends on the values of f in arbitrarily small neighborhoods of the y_I . This is not obvious at all since the definition of $\text{mj}_p(f, z)$ involves divided differences of f , which are obtained by integrating on the whole convex hull $\sigma(\underline{x})$ of the x_i (see Definition 3.1). In particular, it shows that $\text{mj}_p(f, z)$ makes sense even if f is only $\mathcal{C}^{|I|-1}$ in some neighborhood of y_I for all $I \in \mathcal{I}$. Hence Definition 1.3 makes sense even if Ω is not convex.

In the following, we consider what we think of as the I -th part of a multijet, where $I \subset \llbracket 1, p \rrbracket$. This is just a variation on what we did in Definitions 5.7 and 5.8 and it is defined as follows.

Definition 5.12 (I -multijets). Let $n \geq 1$ and $p \geq 1$ and let $I \subset \llbracket 1, p \rrbracket$ be nonempty, we define the bundle of I -multijets as the following smooth bundle of rank $|I|$ over $C_p[\mathbb{R}^n]$:

$$\mathcal{MJ}_I(\mathbb{R}^n) = (\mathbb{R}_{|I|-1}[X] \times C_p[\mathbb{R}^n]) / \mathcal{G}_I.$$

Let $f \in \mathcal{C}^{|I|-1}(\mathbb{R}^n)$ and $z \in C_p[\mathbb{R}^n]$. We define by $\text{mj}_I(f, z) = K(f, \pi(z)_I) \bmod \mathcal{G}_I(z) \in \mathcal{MJ}_I(\mathbb{R}^n)_z$ the I -multijet of f at z .

As explained in Section 5.1, for all $\emptyset \neq I \subset \llbracket 1, p \rrbracket$ we have a map $\mathcal{G}_I : C_p[\mathbb{R}^n] \rightarrow \text{Gr}_{|I|}(\mathbb{R}_{|I|-1}[X])$ extending the one on $(\mathbb{R}^n)^p \setminus \Delta_p$. Let $z \in C_p[\mathbb{R}^n]$ and $\underline{x} = \pi(z)$. As in Section 4 we define $\tilde{\mathcal{G}}_I(z) = K(\cdot, \underline{x}_I)^{-1}(\mathcal{G}_I(z)) \in \text{Gr}_{|I|}(\mathbb{R}_{p-1}[X])$, where $K(\cdot, \underline{x}_I) : \mathbb{R}_{p-1}[X] \rightarrow \mathbb{R}_{|I|-1}[X]$. Note that $\tilde{\mathcal{G}}_I(z)$ has the same codimension as $\mathcal{G}_I(z)$ since $K(\cdot, \underline{x}_I)$ is surjective.

Lemma 5.13 (compatibility of the \mathcal{G}_I). For all $I \subset \llbracket 1, p \rrbracket$ and $z \in C_p[\mathbb{R}^n]$ we have $\mathcal{G}(z) \subset \tilde{\mathcal{G}}_I(z)$.

Proof. Recall from Section 4 that $\mathcal{G}(z) \subset \tilde{\mathcal{G}}_I(z)$ for any $z \in (\mathbb{R}^n)^p \setminus \Delta_p \subset C_p[\mathbb{R}^n]$, that is, $K(\cdot, \pi(z)_I)(\mathcal{G}(z)) \subset \mathcal{G}_I(z)$. This incidence relation is a closed condition. By construction, the subset $(\mathbb{R}^n)^p \setminus \Delta_p$ is dense

in $C_p[\mathbb{R}^n]$ and both terms in the previous inclusion are continuous with respect to z ; see Lemma 3.8 and Corollary 5.6. Hence the inclusion actually holds for any $z \in C_p[\mathbb{R}^n]$. Thus $\mathcal{G}(z) \subset \tilde{\mathcal{G}}_I(z)$ for all $z \in C_p[\mathbb{R}^n]$. \square

Let $\emptyset \neq I \subset \llbracket 1, p \rrbracket$, let $z \in C_p[\mathbb{R}^n]$ and $\underline{x} = \pi(z)$. We consider $\text{mj}_I(\cdot, z) : \mathbb{R}_{p-1}[X] \rightarrow \mathcal{MJ}_I(\mathbb{R}^n)_z$ from Definition 5.12. This linear map is surjective as the composition of $K(\cdot, \underline{x}_I)$ and the projection modulo $\mathcal{G}_I(z)$. Moreover, $\ker(\text{mj}_I(\cdot, z)) = \tilde{\mathcal{G}}_I(z)$ contains $\mathcal{G}(z)$ by Lemma 5.13. Hence, $\text{mj}_I(\cdot, z)$ induces a surjective linear map from $\mathcal{MJ}_p(\mathbb{R}^n)_z = \mathbb{R}_{p-1}[X]/\mathcal{G}(z)$ onto $\mathcal{MJ}_I(\mathbb{R}^n)_z$ that we still denote by $\text{mj}_I(\cdot, z)$. This is summarized in the following commutative diagram, where the vertical arrows are the canonical projections and all arrows are surjective:

$$\begin{array}{ccc}
 \mathbb{R}_{p-1}[X] & \xrightarrow{K(\cdot, \underline{x}_I)} & \mathbb{R}_{|I|-1}[X] \\
 \downarrow & \searrow \text{mj}_I(\cdot, z) & \downarrow \\
 \mathcal{MJ}_p(\mathbb{R}^n)_z & \xrightarrow{\text{mj}_I(\cdot, z)} & \mathcal{MJ}_I(\mathbb{R}^n)_z
 \end{array} \tag{5-2}$$

Note that $(P, z) \mapsto (K(P, \pi(z)_I), z)$ is a smooth bundle map over $C_p[\mathbb{R}^n]$ from $\mathbb{R}_{p-1}[X] \times C_p[\mathbb{R}^n]$ to $\mathbb{R}_{|I|-1}[X] \times C_p[\mathbb{R}^n]$. Hence, the previous diagram (5-2) defines a smooth surjective bundle map $\text{mj}_I : (P \bmod \mathcal{G}(z)) \mapsto (P \bmod \mathcal{G}_I(z))$ from $\mathcal{MJ}_p(\mathbb{R}^n)$ to $\mathcal{MJ}_I(\mathbb{R}^n)$ over $C_p[\mathbb{R}^n]$.

Definition 5.14 (partitioned multijet). For all $\mathcal{I} \in \mathcal{P}_p$ and $z \in C_p[\mathbb{R}^n]$, we define a linear map from $\mathcal{MJ}_p(\mathbb{R}^n)_z$ to $\prod_{I \in \mathcal{I}} \mathcal{MJ}_I(\mathbb{R}^n)_z$ by $\text{mj}_{\mathcal{I}}(\cdot, z) : \alpha \mapsto (\text{mj}_I(\alpha, z))_{I \in \mathcal{I}}$.

As above, $\text{mj}_{\mathcal{I}} : (\alpha, z) \mapsto \text{mj}_{\mathcal{I}}(\alpha, z)$ defines a smooth bundle map over $C_p[\mathbb{R}^n]$ from $\mathcal{MJ}_p(\mathbb{R}^n)$ to $\bigoplus_{I \in \mathcal{I}} \mathcal{MJ}_I(\mathbb{R}^n)$, which is obtained as the quotient of $(P, z) \mapsto ((K(P, \pi(z)_I))_{I \in \mathcal{I}}, z)$. However, $\text{mj}_{\mathcal{I}}(\cdot, z)$ is not surjective in general. The following lemma proves its surjectivity in some cases.

Lemma 5.15 (splitting of multijets). Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$ and let $\mathcal{I}(\underline{x})$ be defined as in Definition 2.3. Then $\text{mj}_{\mathcal{I}(\underline{x})}(\cdot, z) : \mathcal{MJ}_p(\mathbb{R}^n)_z \rightarrow \prod_{I \in \mathcal{I}(\underline{x})} \mathcal{MJ}_I(\mathbb{R}^n)_z$ is an isomorphism.

Proof. The map $\text{mj}_{\mathcal{I}(\underline{x})}(\cdot, z)$ is linear between two spaces of the same dimension $p = \sum_{I \in \mathcal{I}(\underline{x})} |I|$, so it is enough to prove its surjectivity. Let $(\alpha_I)_{I \in \mathcal{I}(\underline{x})} \in \prod_{I \in \mathcal{I}(\underline{x})} \mathcal{MJ}_I(\mathbb{R}^n)_z$. For each $I \in \mathcal{I}(\underline{x})$ there exists $P_I \in \mathbb{R}_{|I|-1}[X]$ such that $\alpha_I = P_I \bmod \mathcal{G}_I(z)$. By Lemma 3.9, there exists $P \in \mathbb{R}_{p-1}[X]$ such that $K(P, \underline{x}_I) = P_I$ for all $I \in \mathcal{I}(\underline{x})$. Let $\alpha = P \bmod \mathcal{G}(z) \in \mathcal{MJ}_p(\mathbb{R}^n)_z$. Then, for all $I \in \mathcal{I}(\underline{x})$, we have

$$\text{mj}_I(\alpha, z) = \text{mj}_I(P, z) = K(P, \underline{x}_I) \bmod \mathcal{G}_I(z) = P_I \bmod \mathcal{G}_I(z) = \alpha_I.$$

Hence $\text{mj}_{\mathcal{I}(\underline{x})}(\alpha, z) = (\alpha_I)_{I \in \mathcal{I}(\underline{x})}$, and $\text{mj}_{\mathcal{I}(\underline{x})}(\cdot, z)$ is indeed surjective. \square

Let $k \in \mathbb{N}$ and let $x \in \mathbb{R}^n$. By definition, two maps f and $g \in C^k(\mathbb{R}^n)$ have the same k -jet at x if and only if they have the same Taylor polynomial of order k at x . Let $\underline{x} = (x, \dots, x)$. Recalling Example 3.7, the linear map $K(\cdot, \underline{x}) : C^k(\mathbb{R}^n) \rightarrow \mathbb{R}_k[X]$ is surjective, and it induces an isomorphism $\mathcal{J}_k(\mathbb{R}^n)_x \simeq \mathbb{R}_k[X]$.

Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$, let $\mathcal{I} = \mathcal{I}(\underline{x})$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\mathcal{I}}^{-1}(\underline{x})$; see Definitions 2.3 and 2.6. For all $I \in \mathcal{I}$, the canonical isomorphism $\mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \simeq \mathbb{R}_{|I|-1}[X]$ allows us to see the projection from $\mathbb{R}_{|I|-1}[X]$ onto $\mathcal{MJ}_I(\mathbb{R}^n)_z$ as a canonical linear surjection $\varpi_{z,I} : \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \rightarrow \mathcal{MJ}_I(\mathbb{R}^n)_z$.

Definition 5.16 (gluing map). Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\underline{x}}^{-1}(\underline{x})$, where $\mathcal{I} = \mathcal{I}(\underline{x})$. We define $\varpi_z : (\alpha_I)_{I \in \mathcal{I}} \mapsto (\varpi_{z,I}(\alpha_I))_{I \in \mathcal{I}}$ from $\prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I}$ to $\prod_{I \in \mathcal{I}} \mathcal{MJ}_I(\mathbb{R}^n)_z$. We also define $\Theta_z = \text{mj}_{\mathcal{I}}(\cdot, z)^{-1} \circ \varpi_z$.

We can now check that Θ_z satisfies (4) in Theorem 1.1.

Lemma 5.17 (localness of multijets). *For all $z \in C_p[\mathbb{R}^n]$, the map Θ_z is a linear surjection from $\prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I}$ to $\mathcal{MJ}_p(\mathbb{R}^n)_z$. Moreover, it is the only map such that*

$$\forall f \in \mathcal{C}^{p-1}(\mathbb{R}^n), \quad \Theta_z((j_{|I|-1}(f, y_I))_{I \in \mathcal{I}}) = \text{mj}_p(f, z).$$

Proof. With the same notation as in Definition 5.16, for all $I \in \mathcal{I}$ the map $\varpi_{z,I}$ is a linear surjection by definition. Hence so is ϖ_z . Since $\text{mj}_{\mathcal{I}}(\cdot, z)$ is an isomorphism by Lemma 5.15, the map $\Theta_z = \text{mj}_{\mathcal{I}}(\cdot, z)^{-1} \circ \varpi_z$ is also a linear surjection.

Let $f \in \mathcal{C}^{p-1}(\mathbb{R}^n)$. For all $I \in \mathcal{I}$, the image of $j_{|I|-1}(f, y_I)$ under the canonical isomorphism $\mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \simeq \mathbb{R}_{|I|-1}[X]$ is the Taylor polynomial $K(f, \underline{x}_I)$. Hence

$$\varpi_{z,I}(j_{|I|-1}(f, y_I)) = K(f, \underline{x}_I) \bmod \mathcal{G}_I(z) = \text{mj}_I(K(f, \underline{x}_I), z).$$

Thus, we have

$$\varpi_z((j_{|I|-1}(f, y_I))_{I \in \mathcal{I}}) = (\text{mj}_I(K(f, \underline{x}_I), z))_{I \in \mathcal{I}} = \text{mj}_{\mathcal{I}}(\text{mj}_p(f, z), z),$$

and finally

$$\Theta_z((j_{|I|-1}(f, y_I))_{I \in \mathcal{I}}) = \text{mj}_p(f, z).$$

Since the y_I are pairwise distinct, every element of $\prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I}$ can be realized as $(j_{|I|-1}(f, y_I))_{I \in \mathcal{I}}$ for some $f \in \mathcal{C}^\infty(\mathbb{R}^n)$. Hence the previous relation completely defines Θ_z . \square

5.4. Multijets of vector-valued maps. So far we have only defined multijets of real-valued functions. In this section, we extend the previous construction to maps from \mathbb{R}^n to some vector space V of dimension $r \geq 1$. Let $\pi : C_p[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ be given by Corollary 5.6 as before. We define $\mathcal{MJ}_p(\mathbb{R}^n, V) \rightarrow C_p[\mathbb{R}^n]$ as the tensor product of the bundle $\mathcal{MJ}_p(\mathbb{R}^n) \rightarrow C_p[\mathbb{R}^n]$ from Definition 5.7 with the trivial bundle $V \times C_p[\mathbb{R}^n] \rightarrow C_p[\mathbb{R}^n]$.

Definition 5.18 (multijet bundle of vector-valued maps). Let $n \geq 1$ and $p \geq 1$; let V be a real vector space of dimension $r \geq 1$. We define the *bundle of p -multijets of V -valued maps on \mathbb{R}^n* as the following smooth bundle of rank pr over $C_p[\mathbb{R}^n]$:

$$\mathcal{MJ}_p(\mathbb{R}^n, V) = \mathcal{MJ}_p(\mathbb{R}^n) \otimes V.$$

Definition 5.19 (multijet of a map). Let (v_1, \dots, v_r) denote a basis of V . Let $z \in C_p[\mathbb{R}^n]$ and let $f = \sum_{i=1}^r f_i v_i \in \mathcal{C}^{p-1}(\mathbb{R}^n, V)$. We define by $\text{mj}_p(f, z) = \sum_{i=1}^r \text{mj}_p(f_i, z) \otimes v_i \in \mathcal{MJ}_p(\mathbb{R}^n, V)_z$ the p -multijet of f at z .

Lemma 5.20 (independence from the basis). *In Definition 5.19, the vector $\text{mj}_p(f, z)$ does not depend on the choice of the basis (v_1, \dots, v_r) .*

Proof. Let (w_1, \dots, w_r) be another basis of V . There exists a matrix $(a_{ij})_{1 \leq i, j \leq r} \in \text{GL}_r(\mathbb{R})$ such that $v_i = \sum_{j=1}^r a_{ij} w_j$ for all $i \in \llbracket 1, r \rrbracket$. Letting $g_j = \sum_{i=1}^r a_{ij} f_i$ for all $j \in \llbracket 1, r \rrbracket$, we get

$$f = \sum_{i=1}^r f_i v_i = \sum_{1 \leq i, j \leq r} a_{ij} f_i w_j = \sum_{j=1}^r g_j w_j.$$

Then, by linearity of the p -multijet for functions, we have

$$\sum_{j=1}^r \text{mj}_p(g_j, z) \otimes w_j = \sum_{1 \leq i, j \leq r} a_{ij} \text{mj}_p(f_i, z) \otimes w_j = \sum_{i=1}^r \text{mj}_p(f_i, z) \otimes v_i. \quad \square$$

Example 5.21. If $V = \mathbb{R}^r$, then for all $z \in C_p[\mathbb{R}^n]$ and $f = (f_1, \dots, f_r) \in C^{p-1}(\mathbb{R}^n, \mathbb{R}^r)$ we have

$$\mathcal{MJ}_p(\mathbb{R}^n, \mathbb{R}^r)_z = \mathcal{MJ}_p(\mathbb{R}^n)_z \otimes \mathbb{R}^r \simeq (\mathcal{MJ}_p(\mathbb{R}^n)_z)^r,$$

and under this canonical isomorphism $\text{mj}_p(f, z) = (\text{mj}_p(f_i, z))_{1 \leq i \leq r}$, as one would expect.

Let $x \in \mathbb{R}^n$ and $k \in \mathbb{N}$. We have a canonical isomorphism $\mathcal{J}_k(\mathbb{R}^n, V)_x \simeq \mathcal{J}_k(\mathbb{R}^n)_x \otimes V$. If (v_1, \dots, v_r) is a basis of V , this isomorphism is totally determined by the fact that the image of $j_k(f, x)$ is $\sum_{i=1}^r j_k(f_i, x) \otimes v_i$ for all $f = \sum_{i=1}^r f_i v_i \in C^k(\mathbb{R}^n, V)$. As in the proof of Lemma 5.20, this does not depend on the choice of the basis (v_1, \dots, v_r) .

Definition 5.22 (gluing map for vector-valued multijets). Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$, let $\mathcal{I} = \mathcal{I}(\underline{x})$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\underline{x}}^{-1}(\underline{x})$. Using the previous canonical isomorphisms, we have

$$\prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n, V)_{y_I} = \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \otimes V = \left(\prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I} \right) \otimes V.$$

Recalling Definition 5.16, we define a linear map

$$\Theta_z : \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n, V)_{y_I} \rightarrow \mathcal{MJ}_p(\mathbb{R}^n, V)_z$$

by $\Theta_z(\alpha \otimes v) = \Theta_z(\alpha) \otimes v$ for all $\alpha \in \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I}$ and all $v \in V$.

We now have everything we need to prove Theorem 1.1.

Proof of Theorem 1.1. The base space $C_p[\mathbb{R}^n]$ and the projection π are given by Corollary 5.6. In particular, they satisfy (1) in Theorem 1.1. Definition 5.19 and Lemma 5.10 show that mj_p satisfies (2). Similarly, (3) is satisfied thanks to Lemma 5.11 and the definition of mj_p for V -valued maps.

Let us check that the linear maps Θ_z from Definition 5.22 satisfy (4). Let $z \in C_p[\mathbb{R}^n]$, let $\underline{x} = \pi(z)$, let $\mathcal{I} = \mathcal{I}(\underline{x})$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\underline{x}}^{-1}(\underline{x})$. Let us also denote by (v_1, \dots, v_r) a basis of V . Let $\alpha \in \mathcal{MJ}_p(\mathbb{R}^n, V)_z$. There exists $\alpha_1, \dots, \alpha_r \in \mathcal{MJ}_p(\mathbb{R}^n)_z$ such that $\alpha = \sum_{i=1}^r \alpha_i \otimes v_i$. By Lemma 5.17, for each $i \in \llbracket 1, r \rrbracket$, there exists $\beta_i \in \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}(\mathbb{R}^n)_{y_I}$ such that $\alpha_i = \Theta_z(\beta_i)$. Hence,

$$\Theta_z \left(\sum_{i=1}^r \beta_i \otimes v_i \right) = \sum_{i=1}^r \Theta_z(\beta_i) \otimes v_i = \alpha,$$

and Θ_z is indeed surjective.

Finally, let us consider $f = \sum_{i=1}^r f_i v_i \in \mathcal{C}^{p-1}(\mathbb{R}^n, V)$. Then, by Lemma 5.17 once again,

$$\begin{aligned} \Theta_z((j_{|I|-1}(f, y_I))_{I \in \mathcal{I}}) &= \Theta_z\left(\sum_{i=1}^r (j_{|I|-1}(f_i, y_I))_{I \in \mathcal{I}} \otimes v_i\right) = \sum_{i=1}^r \Theta_z((j_{|I|-1}(f_i, y_I))_{I \in \mathcal{I}}) \otimes v_i \\ &= \sum_{i=1}^r \text{mj}_p(f_i, z) \otimes v_i = \text{mj}_p(f, z). \end{aligned} \quad \square$$

Remark 5.23. Another way to define $\mathcal{MJ}_p(\mathbb{R}^n, V)$ and mj_p is the following. If $f : \mathbb{R}^n \rightarrow V$ is regular enough, then the divided differences from Definition 3.1 still make sense, only this time $f[x_0, \dots, x_k] \in \text{Sym}^k(\mathbb{R}^n) \otimes V$. Then, one can still define $K(f, \underline{x})$ as in Proposition 3.4, and it defines an element of $\mathbb{R}_{p-1}[X] \otimes V$ that interpolates the divided differences of f . Similarly, everything we did from Section 3.1 to Section 5.3 can be adapted to the case of V -valued maps, simply by tensoring each vector space by V , and each linear map by Id_V . One can check that we recover the same objects as in Definitions 5.18 and 5.19, up to canonical isomorphisms.

6. Application to zeros of Gaussian fields

This section is concerned with our application of multijet bundles to Gaussian fields. In Section 6.1, we describe the local model for the Gaussian fields with values in a vector bundle that we consider. In Section 6.2, we prove a Bulinskaya-type lemma and a Kac–Rice formula for the zeros of these fields. Section 6.3 is dedicated to the definition of the Kac–Rice densities of order larger than 2. We also relate the properties of these functions with the moments of the linear statistics associated with our field. Finally, we prove Theorems 1.6 and 1.9 in Section 6.4, using the multijet bundles defined in Theorem 1.1.

6.1. Gaussian vectors and Gaussian sections. In this section, we briefly recall some notation and conventions concerning Gaussian vectors. Then we describe the local model for Gaussian fields with values in a vector bundle. We will mostly consider centered random vectors in finite-dimensional vector spaces, so we restrict ourselves to this setting. In the following, V is a finite-dimensional real vector space.

Definition 6.1 (Gaussian vector). We say that a random vector X with values in V is a *centered Gaussian vector* if, for all $\eta \in V^*$, the real random variable $\eta(X)$ is a centered Gaussian in \mathbb{R} .

In particular, a centered Gaussian vector in V has finite moments up to any order. Let us assume that V is endowed with an inner product $\langle \cdot, \cdot \rangle$. Then for all $v \in V$, we define $v^* = \langle v, \cdot \rangle \in V^*$.

Definition 6.2 (variance operator). Let X be a centered Gaussian vector in $(V, \langle \cdot, \cdot \rangle)$. Then its *variance operator* is the nonnegative self-adjoint endomorphism $\text{Var}(X) = \mathbb{E}[X \otimes X^*]$ of V . We say that X is *nondegenerate* if $\text{Var}(X)$ is invertible.

Recall that a centered Gaussian vector in $(V, \langle \cdot, \cdot \rangle)$ is completely determined by its variance. In the following, we denote by $\mathcal{N}(0, \Lambda)$ the centered Gaussian distribution of variance Λ , and by $X \sim \mathcal{N}(0, \Lambda)$ the fact that X follows this distribution.

Definition 6.3 (Gaussian field). Let $E \rightarrow M$ be a vector bundle over some manifold M . We say that a random section s of $E \rightarrow M$ is a *centered Gaussian field* if for all $m \geq 1$ and all x_1, \dots, x_m the

random vector $(s(x_1), \dots, s(x_m))$ is a centered Gaussian. We say that this field is *nondegenerate* if $s(x)$ is nondegenerate for all $x \in M$.

If the centered Gaussian field s is \mathcal{C}^p , then its jet $j_k(s, x)$ is a centered Gaussian for all $x \in M$. Thus, the definition of p -nondegeneracy of the field makes sense; see Definition 1.8. Note that 0-nondegenerate simply means nondegenerate.

Since this will appear in several places later on, let us describe the local model for Gaussian fields in this context. Let $x_0 \in M$. There exists a chart (U, φ) of M around x_0 . That is $\varphi : U \rightarrow \Omega$ is a diffeomorphism between an open neighborhood U of x_0 and an open subset $\Omega \subset \mathbb{R}^n$. Up to reducing U , we can assume that E is trivial over U , i.e., there exists a trivialization $\tau : E|_U \rightarrow \mathbb{R}^r \times U$. Letting $\tau_\varphi = (\text{Id}, \varphi) \circ \tau$, we have the following commutative diagram, where arrows on the top row are bundle maps covering the maps on the bottom row:

$$\begin{array}{ccccc}
 & & \tau_\varphi & & \\
 & \nearrow & & \searrow & \\
 E|_U & \xrightarrow{\tau} & \mathbb{R}^r \times U & \xrightarrow{(\text{Id}, \varphi)} & \mathbb{R}^r \times \Omega \\
 \downarrow & & \downarrow & & \downarrow \\
 U & \xrightarrow{\text{Id}} & U & \xrightarrow{\varphi} & \Omega
 \end{array} \tag{6-1}$$

Let s be a local section of $E|_U$. Then $\tau_\varphi \circ s \circ \varphi^{-1}$ is a section of the trivial bundle on the right-hand side of (6-1). Hence there exists a map $f : \Omega \rightarrow \mathbb{R}^r$ such that $\tau_\varphi \circ s \circ \varphi^{-1} = (f, \text{Id})$. For all $x \in \Omega$, the vector $f(x)$ is the image of $s(\varphi^{-1}(x))$ by a linear bijection. Thus, if $s : M \rightarrow E$ is a centered Gaussian field, its restriction to U corresponds to a centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$. Moreover, f has the same regularity as s .

The local trivializations on the diagram (6-1) induce a similar picture for jet bundles so that we have a local trivialization $\mathcal{J}_p(U, E|_U) \simeq \mathcal{J}_p(\Omega, \mathbb{R}^r)$, under which $j_p(f, x)$ corresponds to $j_p(s, \varphi^{-1}(x))$. Thus, the Gaussian section s is p -nondegenerate in the sense of Definition 1.8 if and only if f is p -nondegenerate in the sense of Definition 1.4. If this is the case, up to replacing Ω by a smaller Ω' such that $\bar{\Omega}' \subset \Omega$ is compact, we can assume that f is uniformly p -nondegenerate, in the sense that $\det \text{Var}(j_p(f, x))$ is bounded from below on Ω . This local picture is summarized in the following lemma.

Lemma 6.4 (local model for Gaussian fields). *Let $s : M \rightarrow E$ be a centered p -nondegenerate Gaussian field. For all $x_0 \in M$, there exist an open neighborhood U of x_0 and a local trivialization of the form (6-1) such that s reads in local coordinates as a centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$ of the same regularity as s and which is uniformly p -nondegenerate on Ω .*

6.2. Bulinskaya lemma and Kac–Rice formula for the expectation. Let (M, g) be a Riemannian manifold of dimension $n \geq 1$ without boundary and let $E \rightarrow M$ be a vector bundle of rank $r \in \llbracket 1, n \rrbracket$. We consider a nondegenerate centered Gaussian field $s : M \rightarrow E$, in the sense of Definition 6.3. The goal of this section is to state a Bulinskaya-type lemma and a Kac–Rice formula for the expectation of the linear statistics of s .

When M is an open subset of \mathbb{R}^n and $E = \mathbb{R}^r \times M$ is trivial, these results are proved by Armentano, Azaïs and Leòn [Armentano et al. 2023b, Proposition 2.1 and Theorem 2.2]. They are extended to fields

on submanifolds of \mathbb{R}^N in [Armentano et al. 2023b, Section 9.1]. In the following we check that the results of that work can be adapted to the case of Gaussian sections.

Remark 6.5. Some readers are most interested in the zeros of Gaussian fields from \mathbb{R}^n to \mathbb{R}^r and the present geometric setting may seem overly complicated to them. Let us stress that, even in the simpler setting of fields from \mathbb{R}^n to \mathbb{R}^r , our proof of Theorem 1.6 uses the Kac–Rice formula in the more general setting we are studying here.

In order to state the Bulinskaya lemma and the Kac–Rice formula, we need the following.

Definition 6.6 (Jacobian determinant). Let $L : V \rightarrow V'$ be a linear map between Euclidean spaces and let L^* denote its adjoint map. The *Jacobian* of L is defined as $\text{Jac}(L) = \det(LL^*)^{1/2}$.

Remark 6.7. We have $\text{Jac}(L) \geq 0$, and $\text{Jac}(L) > 0$ if and only if L is surjective. In particular, the fact that $\text{Jac}(L) = 0$ depends only on L and not on the Euclidean structures on V and V' . Thus the condition $\text{Jac}(L) = 0$ makes sense even if no inner product is specified.

Proposition 6.8 (weak Bulinskaya lemma). *Let ∇ be a connection on $E \rightarrow M$. If the centered Gaussian field $s : M \rightarrow E$ is \mathcal{C}^1 and nondegenerate, the $(n-r)$ -dimensional Hausdorff measure of*

$$\{x \in M \mid s(x) = 0 \text{ and } \text{Jac}(\nabla_x s) = 0\}$$

is almost surely 0.

Remark 6.9. If $s(x) = 0$ then $\nabla_x s$ does not depend on ∇ . Hence the random set we are interested in Proposition 6.8 does not depend on the choice ∇ .

Proof of Proposition 6.8. We can cover M by countably many open trivialization domains of the type described in Lemma 6.4. Then it is enough to prove the result in each of these domains.

Let $U \subset M$ be as Lemma 6.4. In local coordinates, the restriction of s reads as a uniformly nondegenerate \mathcal{C}^1 centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$, where $\Omega \subset \mathbb{R}^n$ is open. Moreover, for any $x \in \Omega$ such that $f(x) = 0$, the covariant derivative of s reads as $D_x f$, independently of the choice of ∇ . Thus, we are left with proving that the $(n-r)$ -dimensional Hausdorff measure of

$$\{x \in \Omega \mid f(x) = 0 \text{ and } \text{Jac}(D_x f) = 0\}$$

is almost surely 0, which is given by [Armentano et al. 2023b, Proposition 2.1]. □

Let us assume from now on that the centered Gaussian field $s : M \rightarrow E$ is \mathcal{C}^1 and nondegenerate. We denote its zero set by $Z = s^{-1}(0)$. Let us define $Z_{\text{sing}} = \{x \in Z \mid \text{Jac}(\nabla_x s) = 0\}$ and $Z_{\text{reg}} = Z \setminus Z_{\text{sing}}$. By Proposition 6.8, the $(n-r)$ -dimensional Hausdorff measure of the singular part Z_{sing} is almost surely 0. On the other hand, the regular part Z_{reg} is the set of points where s vanishes transversally. As such, it is a (possibly empty) \mathcal{C}^1 submanifold of M of codimension r without boundary. Thus, Z is almost surely the union of an open (in Z) regular part Z_{reg} of dimension $n-r$, and a negligible singular part Z_{sing} that we can think of as a set of lower dimension. Let us mention that, under additional assumptions on the field, the singular part is almost surely empty.

Proposition 6.10 (strong Bulinskaya lemma). *If the centered Gaussian field $s : M \rightarrow E$ is \mathcal{C}^2 and 1-nondegenerate, then $Z_{\text{sing}} = \emptyset$ almost surely.*

Proof. As in the proof of Proposition 6.8, it is enough to prove the result in local coordinates given by Lemma 6.4. In these coordinates, s reads as a \mathcal{C}^2 centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$ which is uniformly 1-nondegenerate, that is, $\det \text{Var}(f(x), D_x f)$ is bounded from below on Ω . Then the result follows from [Azaïs and Wschebor 2009, Proposition 6.12]. □

The Riemannian metric g induces a metric on Z_{reg} , which in turn defines an $(n-r)$ -dimensional Riemannian volume measure $d\text{Vol}_Z$. This measure coincides with the $(n-r)$ -dimensional Hausdorff measure on Z . In the following, we consider this measure as a Radon measure on M defined as follows. Recall that the space of Radon measures is the topological dual of $\mathcal{C}_c^0(M)$, and that being a nonnegative Radon is equivalent to being a Borel measure which is finite on compact subsets.

Definition 6.11 (random measure associated with Z). We denote by ν the random nonnegative Radon measure on M defined by

$$\forall \phi \in \mathcal{C}_c^0(M), \quad \langle \nu, \phi \rangle = \int_{Z_{\text{reg}}} \phi(x) d\text{Vol}_Z(x).$$

We define $\langle \nu, \phi \rangle$ similarly if ϕ is nonnegative Borel function (in which case $\langle \nu, \phi \rangle \in [0, +\infty]$) or if ϕ is a Borel function such that $\langle \nu, |\phi| \rangle < +\infty$ almost surely.

Example 6.12. If $n = r$ then Z is almost surely locally finite. In this case $\nu = \sum_{x \in Z} \delta_x$ is the random counting measure of this point process.

Let us go back to the local model of Lemma 6.4. Around any $x_0 \in M$ there exists a chart (U, φ) and a local trivialization of the kind described by (6-1). Since s is \mathcal{C}^1 and nondegenerate, it corresponds in local coordinates to a \mathcal{C}^1 nondegenerate Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$. We still denote by Z (resp. Z_{reg}) the image of Z (resp. Z_{reg}) by φ , which is the zero set of f (resp. its regular part). Similarly, we still denote by g (resp. $d\text{Vol}_Z$) the push-forward to Ω of the metric g (resp. of the measure $d\text{Vol}_Z$), and we identify test-functions on U with test-functions on Ω . Thus, if $\phi \in L_c^\infty(U)$ we have

$$\langle \nu, \phi \rangle = \int_{Z_{\text{reg}}} \phi(x) d\text{Vol}_Z(x),$$

where we think of everything on the right-hand side as defined on Ω . Now, the measure $d\text{Vol}_Z$ is the $(n-r)$ -dimensional Riemannian volume on $Z \subset \Omega \subset \mathbb{R}^n$ induced by g . In the following, we will need to understand how it compares with the Riemannian volume $d\text{Vol}_Z^0$ on Z induced by the Euclidean metric. This is the purpose of what comes next.

Definition 6.13 (Riemannian densities). Let $x \in \Omega$ and let G be a subspace of \mathbb{R}^n . We denote by $\det(g(x)|_G)$ the determinant of the restriction to G of the inner product $g(x)$, in any basis of G which is orthonormal for the Euclidean inner product of \mathbb{R}^n .

For all $r \in \llbracket 0, n \rrbracket$, we denote by $\gamma_r : \Omega \times \text{Gr}_r(\mathbb{R}^n) \rightarrow (0, +\infty)$ the continuous map defined by $\gamma_r : (x, G) \mapsto \det(g(x)|_G)^{1/2}$. We also write $\gamma : x \mapsto \gamma_0(x, \mathbb{R}^n)$ for simplicity.

Lemma 6.14 (comparing volumes). *Let Z be a submanifold of codimension r of Ω and let $d\text{Vol}_Z$ (resp. $d\text{Vol}_Z^0$) denote the $(n-r)$ -dimensional Riemannian volume on Z induced by g (resp. the Euclidean metric). Then $d\text{Vol}_Z$ admits the density $x \mapsto \gamma_r(x, T_x Z)$ with respect to $d\text{Vol}_Z^0$. In particular $d\text{Vol}_\Omega$ admits the density γ with respect to the Lebesgue measure on Ω .*

Proof. This follows directly from the definition of the Riemannian volume measures; see [Lee 2018, Chapter 3] for example. □

We can now state and prove the Kac–Rice formula for the expectation of the linear statistics in our setting of Gaussian fields on M with values in a vector bundle E .

Definition 6.15 (Kac–Rice density). Let $\rho_1 : M \rightarrow [0, +\infty)$ be defined by

$$\rho_1 : x \longmapsto \frac{\mathbb{E}[\text{Jac}(\nabla_x s) \mid s(x) = 0]}{\det(2\pi \text{Var}(s(x)))^{1/2}},$$

where the numerator stands for the conditional expectation of $\text{Jac}(\nabla_x s)$ given that $s(x) = 0$.

Remark 6.16. Since s is nondegenerate and C^1 , the function ρ_1 is well-defined and continuous. Moreover, it does not depend on the choice of ∇ , nor on the choice of a metric on E .

Proposition 6.17 (Kac–Rice formula for the expectation). *Let s be a nondegenerate C^1 centered Gaussian field. Then, for any Borel function $\phi : M \rightarrow \mathbb{R}$ which is nonnegative or such that $\phi\rho_1 \in L^1(M)$, we have*

$$\mathbb{E}[\langle v, \phi \rangle] = \int_M \phi(x)\rho_1(x) d\text{Vol}_M(x),$$

i.e., $\mathbb{E}[v]$ is the Radon measure on M with density ρ_1 with respect to the Riemannian volume $d\text{Vol}_M$.

Remark 6.18. For any Borel maps ϕ_1 and ϕ_2 , we have

$$\mathbb{E}[|\langle v, \phi_1 \rangle - \langle v, \phi_2 \rangle|] \leq \mathbb{E}[\langle v, |\phi_1 - \phi_2| \rangle] = \int_M |\phi_1(x) - \phi_2(x)|\rho_1(x) d\text{Vol}_M(x).$$

Then, if $\phi_1 = \phi_2$ almost everywhere on M , we have $\langle v, \phi_1 \rangle = \langle v, \phi_2 \rangle$ almost surely. Thus, $\langle v, \phi \rangle$ makes sense as a random variable even if ϕ is only defined up to modification on a negligible set.

Proof of Proposition 6.17. By a partition of unity argument, it enough to prove the result if ϕ is compactly supported in an open domain U satisfying the same properties as in Lemma 6.4. In this case, in local coordinates, the field s corresponds to a nondegenerate C^1 centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$ with $\Omega \subset \mathbb{R}^n$ open. Thanks to Remark 6.16, we can assume that ∇ corresponds in this trivialization to the standard derivation for maps from Ω to \mathbb{R}^r and that the metric on E corresponds to the canonical inner product on \mathbb{R}^r . Identifying Z_{reg} , the metric g , the measure $d\text{Vol}_Z$ and the test-function ϕ with their images in the trivialization, we have reduced our problem to proving the result for the vanishing locus of f with the volume measures induced by g .

By Lemma 6.14, we have

$$\mathbb{E}[\langle v, \phi \rangle] = \mathbb{E}\left[\int_{Z_{\text{reg}}} \phi(x) d\text{Vol}_Z(x)\right] = \mathbb{E}\left[\int_{Z_{\text{reg}}} \phi(x)\gamma_r(x, \ker D_x f) d\text{Vol}_Z^0(x)\right].$$

For all $x \in \Omega$ and $\lambda \in \mathcal{C}^0(\Omega, \mathcal{L}(\mathbb{R}^n, \mathbb{R}^r))$ we define $\Psi(x, \lambda) = \phi(x)\gamma_r(x, \ker \lambda(x))\mathbf{1}_O(\lambda(x))$, where $O = \{L \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^r) \mid \text{Jac}(L) > 0\}$. Since O is open and the maps $\ker : O \rightarrow \text{Gr}_r(\mathbb{R}^n)$ and γ_r are continuous, the map Ψ is lower semicontinuous with respect to each variable, where $\mathcal{C}^0(\Omega, \mathcal{L}(\mathbb{R}^n, \mathbb{R}^r))$ is equipped with the weak topology. Thus, we can apply the Euclidean Kac–Rice formula with weight from [Armentano et al. 2023b, Theorem 7.1 and Remark 8] to $\Psi(x, Df)$. This yields

$$\mathbb{E}[\langle \nu, \phi \rangle] = \int_{\Omega} \phi(x) \frac{\mathbb{E}[\gamma_r(x, \ker D_x f) \text{Jac}^0(D_x f) \mid f(x) = 0]}{\det(2\pi \text{Var}(f(x)))^{1/2}} dx,$$

where Jac^0 means that we computed the Jacobian with respect to the Euclidean metric on \mathbb{R}^n .

To conclude, we need to compare Jac^0 with the Jacobian Jac with respect to g . This is the content of Lemma 6.19 below, which yields that $\gamma_r(x, \ker D_x f) \text{Jac}^0(D_x f) = \gamma(x) \text{Jac}(D_x f)$. Since $\gamma(x)$ is deterministic, by Lemma 6.14 we have

$$\mathbb{E}[\langle \nu, \phi \rangle] = \int_{\Omega} \phi(x) \frac{\mathbb{E}[\text{Jac}(D_x f) \mid f(x) = 0]}{\det(2\pi \text{Var}(f(x)))^{1/2}} \gamma(x) dx = \int_{\Omega} \phi(x) \rho_1(x) d\text{Vol}_{\Omega}(x).$$

This proves that the result holds locally, that is, for a field $f : \Omega \rightarrow \mathbb{R}^r$, with the volume measures induced by any Riemannian metric on Ω , which concludes the proof. □

Lemma 6.19 (comparing Jacobians). *Let $x \in \Omega$ and let $L : \mathbb{R}^n \rightarrow \mathbb{R}^r$ be a surjective linear map. With the same notation as above, we have $\gamma_r(x, \ker L) \text{Jac}^0(L) = \gamma(x) \text{Jac}(L)$.*

Proof. We denote by L_g^* (resp. L_0^*) the adjoint of L with respect to the inner product $g(x)$ (resp. the Euclidean inner product). In a Euclidean orthonormal basis adapted to $\ker(L)^\perp \oplus \ker(L)$, the matrix of $g(x)$ is symmetric of the form $\begin{pmatrix} A & {}^t B \\ B & C \end{pmatrix}$ with A and C positive-definite, the matrix of L is $(F \ 0)$, that of L_0^* is $\begin{pmatrix} {}^t F \\ 0 \end{pmatrix}$ and that of L_g^* is $\begin{pmatrix} X \\ Y \end{pmatrix}$. We have $\begin{pmatrix} {}^t F \\ 0 \end{pmatrix} = \begin{pmatrix} A & {}^t B \\ B & C \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}$, which leads to ${}^t F = (A - {}^t B C^{-1} B)X$. Hence,

$$\det(LL_0^*) = \det(F {}^t F) = \det(A - {}^t B C^{-1} B) \det(FX) = \det(A - {}^t B C^{-1} B) \det(LL_g^*),$$

and $\gamma_r(x, G) \text{Jac}^0(L) = \det(C)^{1/2} \det(A - {}^t B C^{-1} B)^{1/2} \text{Jac}(L)$. Since $A - {}^t B C^{-1} B$ is the Schur complement of C in the matrix of $g(x)$, we have $\det(C)^{1/2} \det(A - {}^t B C^{-1} B)^{1/2} = \gamma(x)$. □

6.3. Factorial moment measures and Kac–Rice densities. As in the previous section, we consider a nondegenerate \mathcal{C}^1 centered Gaussian field $s : M \rightarrow E$ which is a random section of some vector bundle $E \rightarrow M$. Recall that $n = \dim(M)$ and $r \in \llbracket 1, n \rrbracket$ is the rank of E . We are interested in the finiteness of the moments of the linear statistics $\langle \nu, \phi \rangle$ with $\phi \in L_c^\infty(M)$; see Definition 6.11. In this section, we introduce the factorial moment measures and Kac–Rice densities of the fields s , and we relate them to higher moments of the linear statistics.

In the following, p will always denote the order of the moment we are considering. Recall that, under our hypothesis on s , the random measure ν is almost surely a nonnegative Radon measure on M ; see Remark 6.18.

Definition 6.20 (product measures). Let $p \geq 1$. We denote by $\nu^{\otimes p}$ the product measure of ν with itself p -times. We also denote by $\nu^{[p]}$ the restriction of $\nu^{\otimes p}$ to $M^p \setminus \Delta_p$. That is, for any test-function Φ ,

$$\langle \nu^{\otimes p}, \Phi \rangle = \int_{Z_{\text{reg}}^p} \Phi(\underline{x}) \, d\text{Vol}_Z^{\otimes p}(\underline{x}) \quad \text{and} \quad \langle \nu^{[p]}, \Phi \rangle = \int_{Z_{\text{reg}}^p \setminus \Delta_p} \Phi(\underline{x}) \, d\text{Vol}_Z^{\otimes p}(\underline{x}).$$

Almost surely, these measures are Radon measures on M^p . More generally, if we want to consider product spaces indexed by a nonempty finite set A instead of $\llbracket 1, p \rrbracket$, we denote by $\nu^{\otimes A}$ the product measure of ν with itself $|A|$ times on M^A and by $\nu^{[A]}$ its restriction to $M^A \setminus \Delta_A$.

The following lemma describes the relation between $\nu^{\otimes p}$ and $\nu^{[p]}$, using the notation introduced in Section 2.1.

Lemma 6.21 (relation between $\nu^{\otimes p}$ and $\nu^{[p]}$). Let $p \geq 1$. If $r < n$ then $\nu^{\otimes p} = \nu^{[p]}$. If $r = n$ then $\nu^{\otimes p} = \sum_{\mathcal{I} \in \mathcal{P}_p} (\iota_{\mathcal{I}})_*(\nu^{[\mathcal{I}]})$.

Proof. If $r < n$ then Z_{reg} is a C^1 submanifold of positive dimension in M . In particular, the large diagonal in $(Z_{\text{reg}})^p$ has positive codimension, and hence is negligible for $d\text{Vol}_Z^{\otimes p}$. Thus $\nu^{\otimes p} = \nu^{[p]}$ in this case.

If $r = n$ then Z_{reg} is a locally finite set and ν is its counting measure. Similarly, $\nu^{\otimes p}$ is the counting measure of the locally finite $(Z_{\text{reg}})^p$, and Δ_p is no longer negligible for this measure. If $r = n = 1$, we proved in [Ancona and Letendre 2021, Lemma 2.7] that $\nu^{\otimes p} = \sum_{\mathcal{I} \in \mathcal{P}_p} (\iota_{\mathcal{I}})_*(\nu^{[\mathcal{I}]})$. The proof is purely combinatorics and it extends immediately to the case $r = n \geq 1$. \square

Our interest in these measures is that, by the Fubini theorem, for all $\phi \in L_c^\infty(M)$ we have $\mathbb{E}[\langle \nu, \phi \rangle^p] = \mathbb{E}[\langle \nu^{\otimes p}, \phi^{\otimes p} \rangle] = \langle \mathbb{E}[\nu^{\otimes p}], \phi^{\otimes p} \rangle$, where $\phi^{\otimes p} : (x_1, \dots, x_p) \mapsto \phi(x_1) \cdots \phi(x_p)$. Thus, the measure $\mathbb{E}[\nu^{\otimes p}]$ is closely related with the computation of moments of linear statistics. For technical reasons, it is more convenient to consider $\mathbb{E}[\nu^{[p]}]$ instead.

Definition 6.22 (moment measures). Let $p \geq 1$. The measure $\mathbb{E}[\nu^{\otimes p}]$ is called the p -th moment measure of the field s and $\mathbb{E}[\nu^{[p]}]$ is called its p -th factorial moment measure.

Definition 6.23 (Kac–Rice density of order p). Let $p \geq 1$ and let us assume that the random vector $(s(x_1), \dots, s(x_p))$ is nondegenerate for all $(x_1, \dots, x_p) \in M^p \setminus \Delta_p$. Then we define

$$\rho_p : (x_1, \dots, x_p) \mapsto \frac{\mathbb{E}[\prod_{i=1}^p \text{Jac}(\nabla_{x_i} s) \mid \forall i \in \llbracket 1, p \rrbracket, s(x_i) = 0]}{\det(2\pi \text{Var}(s(x_1), \dots, s(x_p)))^{1/2}}$$

from $M^p \setminus \Delta_p$ to $[0, +\infty)$, where the numerator is the conditional expectation of $\prod_{i=1}^p \text{Jac}(\nabla_{x_i} s)$ given that $s(x_i) = 0$ for all $i \in \llbracket 1, p \rrbracket$.

Once again, ρ_p is well-defined and continuous on $M^p \setminus \Delta_p$ thanks to our nondegeneracy hypothesis. However, its expression is singular along Δ_p . In particular, ρ_p is in general not bounded, which raises the question of its local integrability near Δ_p . For example, if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a nondegenerate enough stationary Gaussian field and $p = 2$, one can check that as $y \rightarrow x$, the corresponding Kac–Rice density $\rho_2(x, y)$ behaves like $\|y - x\|$ if $n = 1$ and like $1/\|y - x\|$ if $n > 1$.

Proposition 6.24 (Kac–Rice formula for the p -th factorial moment). *Let s be a \mathcal{C}^1 centered Gaussian field such that $(s(x_1), \dots, s(x_p))$ is nondegenerate for all $(x_1, \dots, x_p) \in M^p \setminus \Delta_p$. Then, for any Borel function $\Phi : M^p \rightarrow \mathbb{R}$ which is nonnegative or such that $\Phi \rho_p \in L^1(M^p)$, we have*

$$\mathbb{E}[\langle \nu^{[p]}, \Phi \rangle] = \int_{M^p} \Phi(\underline{x}) \rho_p(\underline{x}) \, d\text{Vol}_M^{\otimes p}(\underline{x}),$$

i.e., $\mathbb{E}[\nu^{[p]}]$ is the measure on M^p with density ρ_p with respect to the Riemannian volume $d\text{Vol}_M^{\otimes p}$.

Proof. Let us consider $S : (x_1, \dots, x_p) \mapsto (s(x_1), \dots, s(x_p))$ on $M^p \setminus \Delta_p$, which is a random section of the restriction over $M^p \setminus \Delta_p$ of the vector bundle $E^p \rightarrow M^p$. This is a nondegenerate \mathcal{C}^1 centered Gaussian field on $M^p \setminus \Delta_p$, and $\nu^{[p]}$ is the measure of integration over its zero set. Bearing in mind that Δ_p is negligible in M^p for $d\text{Vol}_M^{\otimes p}$, the result follows from Proposition 6.17 applied to S . \square

The following proposition relates the properties of the Kac–Rice densities, the moment measures and the moments of linear statistics.

Proposition 6.25 (relation between moments, measures and densities). *Let $p \geq 1$ and let s be a \mathcal{C}^1 centered Gaussian field such that $(s(x_1), \dots, s(x_p))$ is nondegenerate for all $(x_1, \dots, x_p) \notin \Delta_p$. Then the following four properties are equivalent:*

- (1) For all $\phi \in L_c^\infty(M)$, we have $\mathbb{E}[|\langle \nu, \phi \rangle|^p] < +\infty$.
- (2) For all $k \in \llbracket 1, p \rrbracket$, the moment measure $\mathbb{E}[\nu^{\otimes k}]$ is Radon on M^k , i.e., finite on compact sets.
- (3) For all $k \in \llbracket 1, p \rrbracket$, the factorial moment measure $\mathbb{E}[\nu^{[k]}]$ is Radon on M^k .
- (4) For all $k \in \llbracket 1, p \rrbracket$, the Kac–Rice density satisfies $\rho_k \in L_{\text{loc}}^1(M^k)$.

Proof. Let us assume (1). Let $k \in \llbracket 1, p \rrbracket$ and let $K_0 \subset M^k$ be compact. There exists a compact set $K \subset M$ such that $K_0 \subset K^k$. Then,

$$\langle \mathbb{E}[\nu^{\otimes k}], \mathbf{1}_{K_0} \rangle \leq \langle \mathbb{E}[\nu^{\otimes k}], \mathbf{1}_K^{\otimes k} \rangle = \mathbb{E}[\langle \nu, \mathbf{1}_K \rangle^k] = \mathbb{E}[|\langle \nu, \mathbf{1}_K \rangle|^k].$$

Since $\mathbf{1}_K \in L_c^\infty(M)$, the p -th absolute moment of $\langle \nu, \mathbf{1}_K \rangle$ is finite; hence so is its k -th absolute moment. Thus $\langle \mathbb{E}[\nu^{\otimes k}], \mathbf{1}_{K_0} \rangle < +\infty$ for all compact K_0 and (2) is satisfied.

If (2) is satisfied then so is (3). Indeed, for any $k \in \llbracket 1, p \rrbracket$, the measure $\nu^{[k]}$ is the restriction of $\nu^{\otimes k}$ to $M^k \setminus \Delta_k$. Thus, for any compact $K \subset M^k$ we have

$$\langle \mathbb{E}[\nu^{[k]}], \mathbf{1}_K \rangle = \mathbb{E}[\langle \nu^{[k]}, \mathbf{1}_K \rangle] \leq \mathbb{E}[\langle \nu^{\otimes k}, \mathbf{1}_K \rangle] = \langle \mathbb{E}[\nu^{\otimes k}], \mathbf{1}_K \rangle < +\infty.$$

If (3) is satisfied, let $k \in \llbracket 1, p \rrbracket$ and let $K \subset M^k$ be a compact. By Proposition 6.24 we have

$$\int_K \rho_k(\underline{x}) \, d\text{Vol}_M^{\otimes p}(\underline{x}) = \int_M \mathbf{1}_K(\underline{x}) \rho_k(\underline{x}) \, d\text{Vol}_M^{\otimes p}(\underline{x}) = \mathbb{E}[\langle \nu^{[k]}, \mathbf{1}_K \rangle] = \langle \mathbb{E}[\nu^{[k]}], \mathbf{1}_K \rangle < +\infty.$$

Thus ρ_k is integrable on any compact set, that is, $\rho_k \in L_{\text{loc}}^1(M^k)$. This proves (4) in this case.

Finally, let us assume that (4) holds. Let $\phi \in L_c^\infty(M)$, and let us denote by $K \subset M$ its compact support. We have $\mathbb{E}[|\langle \nu, \phi \rangle|^p] \leq \mathbb{E}[\langle \nu, |\phi| \rangle^p] \leq \|\phi\|_\infty^p \mathbb{E}[\langle \nu, \mathbf{1}_K \rangle^p]$, so it is enough to prove that $\mathbb{E}[\langle \nu, \mathbf{1}_K \rangle^p] =$

$\mathbb{E}[\langle v^{\otimes p}, \mathbf{1}_{K^p} \rangle]$ is finite. By Lemma 6.21, whether $r = n$ or not, we have

$$\mathbb{E}[\langle v^{\otimes p}, \mathbf{1}_{K^p} \rangle] \leq \sum_{\mathcal{I} \in \mathcal{P}_p} \mathbb{E}[\langle v^{|\mathcal{I}|}, \mathbf{1}_{K^p \circ \iota_{\mathcal{I}}} \rangle] = \sum_{\mathcal{I} \in \mathcal{P}_p} \langle \mathbb{E}[v^{|\mathcal{I}|}], \mathbf{1}_{K^{\mathcal{I}}} \rangle.$$

Then, the Kac–Rice formula for moments and the local integrability of the $(\rho_k)_{1 \leq k \leq p}$ yields

$$\mathbb{E}[\langle v, \mathbf{1}_K \rangle^p] \leq \sum_{\mathcal{I} \in \mathcal{P}_p} \langle \mathbb{E}[v^{|\mathcal{I}|}], \mathbf{1}_{K^{\mathcal{I}}} \rangle = \sum_{\mathcal{I} \in \mathcal{P}_p} \int_{K^{|\mathcal{I}|}} \rho_{|\mathcal{I}|}(\underline{x}) \, d\text{Vol}_M^{\otimes |\mathcal{I}|}(\underline{x}) < +\infty,$$

which proves (1) and concludes the proof. □

6.4. Proofs of Theorems 1.6 and 1.9: finiteness of moments. The goal of this section is to prove Theorems 1.6 and 1.9, which give simple conditions for the finiteness of the moments of the linear statistics of a Gaussian field. We begin by proving a local version of Theorem 1.6, under a nondegeneracy hypothesis for the multijets of the field. This is Theorem 6.26 below. Then we deduce Theorem 1.9 from Theorem 6.26, in the case of Gaussian fields with value in a vector bundle. Finally, Theorem 1.6 is obtained as a special case of Theorem 1.9.

Let $\Omega \subset \mathbb{R}^n$ be open. Recall that $\mathcal{MJ}_p(\Omega, \mathbb{R}^r) \rightarrow C_p[\Omega]$ is defined in Definition 1.3 as the restriction over $C_p[\Omega] \subset C_p[\mathbb{R}^n]$ of the vector bundle $\mathcal{MJ}_p(\mathbb{R}^n, \mathbb{R}^r) \rightarrow C_p[\mathbb{R}^n]$ from Theorem 1.1.

Theorem 6.26 (finiteness of moments, local version). *Let $f : \Omega \rightarrow \mathbb{R}^r$ be a centered Gaussian field and v be as in Definition 6.11. Let $p \geq 1$. If f is C^p and for all $k \in \llbracket 1, p \rrbracket$ the Gaussian field $\text{mj}_k(f, \cdot) : C_k[\Omega] \rightarrow \mathcal{MJ}_k(\Omega, \mathbb{R}^r)$ is nondegenerate, then the four equivalent statements in Proposition 6.25 hold.*

Proof. Let $f : \Omega \rightarrow \mathbb{R}^r$ be a C^p centered Gaussian field such that $\text{mj}_k(f, \cdot) : C_k[\Omega] \rightarrow \mathcal{MJ}_k(\Omega, \mathbb{R}^r)$ is nondegenerate for all $k \in \llbracket 1, p \rrbracket$.

Step 1: Gaussianity and nondegeneracy of the multijets. Since f is C^p , for all $k \in \llbracket 1, p \rrbracket$ we have $\text{mj}_k(f, \cdot) \in \Gamma^1(C_k[\Omega], \mathcal{MJ}_k(\Omega, \mathbb{R}^r))$ because of (2) in Theorem 1.1. Since f is centered and Gaussian, so is any finite collection of jets of f . Then, for all $m \geq 1$ and all $z_1, \dots, z_m \in C_k[\Omega]$ we have that $(\text{mj}_k(f, z_1), \dots, \text{mj}_k(f, z_m))$ is a centered Gaussian. Indeed, by (4) in Theorem 1.1, this is the image of a centered Gaussian by a linear map. Thus, $\text{mj}_k(f, \cdot)$ is a nondegenerate C^1 centered Gaussian field on $C_k[\Omega]$ with values in $\mathcal{MJ}_k(\Omega, \mathbb{R}^r)$.

Let $z \notin \pi^{-1}(\Delta_p)$ and let $\underline{x} = (x_1, \dots, x_p) = \pi(z)$. By (4) in Theorem 1.1, the map Θ_z is a linear surjection. A dimension argument shows that it is actually a bijection. Thus

$$(f(x_1), \dots, f(x_p)) = (j_0(f, x_1), \dots, j_0(f, x_p)) = \Theta_z^{-1}(\text{mj}_p(f, z)),$$

which proves that $(f(x_1), \dots, f(x_p))$ is nondegenerate. Thus, the hypotheses of Proposition 6.25 are satisfied, and the four statements appearing in this proposition are indeed equivalent.

Step 2: Comparing zeros of f and $\text{mj}_k(f, \cdot)$. Let $k \in \llbracket 1, p \rrbracket$. In the following we are going to prove that $\mathbb{E}[v^{[k]}]$ is a Radon measure on Ω^k , which is enough to conclude the proof. In the following, we say that a subset of $C_k[\Omega]$ (resp. Ω^k) is negligible if its $k(n-r)$ -dimensional Hausdorff measure is 0.

Let us consider the Gaussian field $\text{mj}_k(f, \cdot) : C_k[\Omega] \rightarrow \mathcal{MJ}_k(\Omega, \mathbb{R}^r)$. We have checked above that it satisfies the hypotheses of Proposition 6.8. Let $X \subset C_k[\Omega]$ denote the zero set of $\text{mj}_k(f, \cdot)$. As in Section 6.2, we define $X_{\text{sing}} = \{z \in C_k[\Omega] \mid \text{mj}_k(f, z) = 0 \text{ and } \text{Jac}(\nabla_z \text{mj}_k(f, \cdot)) = 0\}$ and $X_{\text{reg}} = X \setminus X_{\text{sing}}$. Recall that X_{reg} is a C^1 submanifold of codimension kr and that X_{sing} is almost surely negligible by Proposition 6.8. Let $Y = X \cap \pi^{-1}(\Omega^k \setminus \Delta_k)$. We also let $Y_{\text{sing}} = Y \cap X_{\text{sing}}$ and $Y_{\text{reg}} = Y \cap X_{\text{reg}}$.

Recalling that $Z = f^{-1}(0) \subset \Omega$, for all $z \in C_k[\Omega]$ we have $z \in Y$ if and only if $\pi(z) \in Z^k \setminus \Delta_k$; see (3) in Theorem 1.1. By (1) in the same theorem, the restriction of π to $\pi^{-1}(\Omega^k \setminus \Delta_k)$ is a diffeomorphism. Hence $\pi(Y) = Z^k \setminus \Delta_k$, the set $\pi(Y_{\text{reg}})$ is a C^1 submanifold of $\Omega^k \setminus \Delta_k$, and $\pi(Y_{\text{sing}})$ is almost surely negligible. Since $Z^k \setminus (Z_{\text{reg}}^k)^k$ is also almost surely negligible, the submanifolds $Z_{\text{reg}}^k \setminus \Delta_k$ and $\pi(Y_{\text{reg}})$ are almost surely the same, up to a negligible set (actually the reader can check that $\pi(Y_{\text{reg}}) = Z_{\text{reg}}^k \setminus \Delta_k$ using the trivialization τ introduced at the end of Section 5.2). Recalling Definition 6.20, this shows that $\nu^{[k]}$ is the same as the integral over $\pi(Y_{\text{reg}})$ with respect to the Riemannian volume $d\text{Vol}_{\pi(Y)}$ induced by the Euclidean metric on $(\mathbb{R}^n)^k$. At this stage we know that, almost surely,

$$\forall \Phi \in L_c^\infty(\Omega^k), \quad \langle \nu^{[k]}, \Phi \rangle = \int_{\pi(Y_{\text{reg}})} \Phi(\underline{x}) \, d\text{Vol}_{\pi(Y)}(\underline{x}). \tag{6-2}$$

Step 3: Comparing volumes. Let us introduce a Riemannian metric g on $C_k[\Omega]$. It induces a volume measure $d\text{Vol}_X$ on X_{reg} , and hence on Y_{reg} . Additionally, let $z \in C_k[\Omega]$ and let $G \subset T_z(C_k[\Omega])$ be a vector subspace of codimension kr . We define $J(G) = \det((D_z \pi|_G)^* D_z \pi|_G)^{1/2}$, where the adjoint of $D_z \pi|_G$ is computed with respect to g on G and to the Euclidean metric on $(\mathbb{R}^n)^k$. Since π is smooth, this defines a smooth nonnegative function J on the total space of the Grassmannian bundle $\text{Gr}_{kr}(T(C_k[\Omega])) \rightarrow C_k[\Omega]$ of subspaces of codimension kr in the tangent of $C_k[\Omega]$. Our interest in this map is that if $z \in Y_{\text{reg}}$ and $G = T_z Y_{\text{reg}}$ then $J(G)$ is the Jacobian determinant of $D_z(\pi_Y)$, where the C^∞ -diffeomorphism $\pi_Y : Y_{\text{reg}} \rightarrow \pi(Y_{\text{reg}})$ is the restriction of π on both sides.

Let $K \subset \Omega^k$ be compact and let us apply (6-2) to $\mathbf{1}_K$. Using the previous notation, the change of variables π_Y yields

$$\langle \nu^{[k]}, \mathbf{1}_K \rangle = \int_{Y_{\text{reg}}} \mathbf{1}_K(\pi(z)) J(T_z Y_{\text{reg}}) \, d\text{Vol}_X(z) \leq \int_{X_{\text{reg}}} \mathbf{1}_{\pi^{-1}(K)}(z) J(T_z X_{\text{reg}}) \, d\text{Vol}_X(z).$$

By (1) in Theorem 1.1, the projection π is proper; hence $\tilde{K} = \pi^{-1}(K)$ is compact. Since the bundle $\text{Gr}_{kr}(T(C_k[\Omega])) \rightarrow C_k[\Omega]$ has compact fiber, its restriction over $\tilde{K} \subset C_k[\Omega]$ is compact. By continuity, the function J is bounded on this compact set by some constant C_K . Finally, we have proved that, almost surely,

$$\langle \nu^{[k]}, \mathbf{1}_K \rangle \leq C_K \langle \tilde{\nu}, \mathbf{1}_{\tilde{K}} \rangle,$$

where $\tilde{\nu}$ is defined by integrating over X_{reg} with respect to $d\text{Vol}_X$. Taking expectation on both sides we get $\langle \mathbb{E}[\nu^{[k]}], \mathbf{1}_K \rangle \leq C_K \langle \mathbb{E}[\tilde{\nu}], \mathbf{1}_{\tilde{K}} \rangle$.

Step 4: Applying the Kac–Rice formula to multijets. Now, X_{reg} is the regular part of the zero set X of the Gaussian field $\text{mj}_k(f, \cdot)$. We have checked at the beginning of the proof that $\text{mj}_k(f, \cdot)$ satisfies the hypotheses of Proposition 6.17. This proposition yields that $\mathbb{E}[\tilde{\nu}]$ is a Radon measure on $C_k[\Omega]$. Hence $\langle \mathbb{E}[\tilde{\nu}], \mathbf{1}_{\tilde{K}} \rangle$ is finite, and so is $\langle \mathbb{E}[\nu^{[k]}], \mathbf{1}_K \rangle$. Thus $\mathbb{E}[\nu^{[k]}]$ is Radon on Ω^k , which concludes the proof. \square

We can now prove Theorem 1.9, which gives a criterion for the finiteness of the p -th moments of the linear statistics associated with a centered Gaussian field $s : M \rightarrow E$, where $E \rightarrow M$ is some vector bundle of rank r over a Riemannian manifold (M, g) without boundary of dimension $n \geq r$. The idea of the proof is to patch together the local results obtained by applying Theorem 6.26 in nice local trivializations.

Proof of Theorem 1.9. Let $p \geq 1$ and let $s \in \Gamma^p(M, E)$ be a centered Gaussian field which is C^p and $(p-1)$ -nondegenerate.

Step 1: Existence of nice local trivializations. Let $x_0 \in M$. There exists an open neighborhood U of x_0 and a local trivialization of E over U given by Lemma 6.4. In this trivialization, the Gaussian section s corresponds to a centered Gaussian field $f : \Omega \rightarrow \mathbb{R}^r$ which is C^p and $(p-1)$ -nondegenerate. We denote by $x \in \Omega$ the image of x_0 in local coordinates.

Let $k \in \llbracket 1, p \rrbracket$ and let $\underline{x} = (x, \dots, x) \in \Omega^k$. Since $j_{p-1}(f, x)$ is nondegenerate so is $j_{k-1}(f, x)$; see Definition 1.4. Then, by (4) in Theorem 1.1, for all $z \in \pi^{-1}(\{\underline{x}\}) \subset C_k[\Omega]$ the Gaussian vector $\text{mj}_k(f, z) = \Theta_z(j_{k-1}(f, x)) \in \mathcal{MJ}_k(\Omega, \mathbb{R}^r)_z$ is nondegenerate. By (1) in Theorem 1.1, the map π is proper; hence $\pi^{-1}(\{\underline{x}\})$ is compact. On the other hand, since f is C^p , we know $\text{mj}_k(f, \cdot)$ is at least C^1 . Thus $z \mapsto \det \text{Var}(\text{mj}_k(f, z))$ is continuous on $C_k[\Omega]$ and positive on the compact $\pi^{-1}(\{\underline{x}\})$, and hence on some neighborhood V_k of $\pi^{-1}(\{\underline{x}\})$ in $C_k[\Omega]$.

Up to reducing V_k we can assume that $V_k = \pi^{-1}(W_k)$, where W_k is an open neighborhood of \underline{x} in Ω^k . Otherwise, there would exist a sequence $(z_n)_{n \in \mathbb{N}} \in C_k[\Omega] \setminus V_k$ such that $\pi(z_n) \xrightarrow{n \rightarrow +\infty} \underline{x}$. By properness of π , up to extracting a subsequence, we could assume that $z_n \xrightarrow{n \rightarrow +\infty} z$. By continuity $z \in \pi^{-1}(\{\underline{x}\})$, which would be absurd. Since W_k is a neighborhood of \underline{x} in Ω^k , there exists an open neighborhood Υ_k of x in Ω such that $(\Upsilon_k)^k \subset W_k$.

Let us define $\Upsilon = \bigcap_{k=1}^p \Upsilon_k$, which is an open neighborhood of x . For all $k \in \llbracket 1, p \rrbracket$, we have $C_k[\Upsilon] = \pi^{-1}(\Upsilon^k) \subset \pi^{-1}((\Upsilon_k)^k) \subset \pi^{-1}(W_k) = V_k$ and $\text{mj}_k(f, \cdot)$ is nondegenerate on $C_k[\Upsilon]$. Thus, up to replacing Ω by the smaller neighborhood Υ of x in Ω and replacing U by the corresponding neighborhood of x_0 on M , we can assume that the local trivialization given by Lemma 6.4 is such that, for all $k \in \llbracket 1, p \rrbracket$, the Gaussian field $\text{mj}_k(f, \cdot) : C_k[\Omega] \rightarrow \mathcal{MJ}_k(\Omega, \mathbb{R}^r)$ is nondegenerate.

Step 2: Reduction to the local case. Let $\phi \in L_c^\infty(M)$ and let K denote its support. By compactness, there exists a finite family $(U_i)_{i=1}^m$ of open subsets such that $K \subset \bigcup_{i=1}^m U_i$ and each U_i is the domain of nice trivialization of the type we built in the previous paragraph. Letting $U_0 = M \setminus K$, there exists a smooth partition of unity $(\chi_i)_{i=0}^m$ subordinated to the open covering $(U_i)_{i=0}^m$ of M . Then $\phi = \sum_{i=1}^m \chi_i \phi$ by the definition of K .

Recall that ν is the measure from Definition 6.11. We have $|\langle \nu, \phi \rangle| \leq \langle \nu, |\phi| \rangle = \sum_{i=1}^m \langle \nu, \phi_i \rangle$, where $\phi_i = \chi_i |\phi|$ for all $i \in \llbracket 1, m \rrbracket$. Let $p \geq 1$, by Hölder's inequality we get

$$\begin{aligned} \mathbb{E}[|\langle \nu, \phi \rangle|^p] &\leq \mathbb{E}\left[\left(\sum_{i=1}^m \langle \nu, \phi_i \rangle\right)^p\right] = \sum_{1 \leq i_1, \dots, i_p \leq m} \mathbb{E}\left[\prod_{j=1}^p \langle \nu, \phi_{i_j} \rangle\right] \leq \sum_{1 \leq i_1, \dots, i_p \leq m} \prod_{j=1}^p \mathbb{E}[\langle \nu, \phi_{i_j} \rangle^p]^{1/p} \\ &\leq m^p \max_{1 \leq i \leq m} \mathbb{E}[\langle \nu, \phi_i \rangle^p]. \end{aligned}$$

Thus, in order to prove Theorem 1.9, it is enough to prove that $\mathbb{E}[\langle \nu, \phi \rangle^p] < +\infty$ for any nonnegative $\phi \in L_c^\infty(M)$ whose support is included in the domain of a nice trivialization.

Step 3: Local case. Let $U \subset M$ be an open subset over which we have a nice trivialization of E and s . That is, U is as in Lemma 6.4, the section s reads as $f : \Omega \rightarrow \mathbb{R}^r$ in local coordinates, and in addition we can assume that for all $k \in \llbracket 1, p \rrbracket$ the field $\text{mj}_k(f, \cdot)$ is nondegenerate on $C_k[\Omega]$. Identifying objects on U with their image in the local trivialization, we reduced our problem to proving that $\mathbb{E}[\langle \nu, \phi \rangle^p] < +\infty$ for all nonnegative $\phi \in L_c^\infty(\Omega)$. Note that ν is the measure of integration over Z_{reg} with respect to the Riemannian volume measure $d\text{Vol}_Z$ induced by the metric g . In order to apply Theorem 6.26, we need to compare ν with $\tilde{\nu}$, which is the measure of integration over Z_{reg} with respect to the Euclidean volume measure $d\text{Vol}_Z^0$.

Let $\phi \in L_c^\infty(\Omega)$ be nonnegative and let K denote its compact support. Recalling Definition 6.13, Lemma 6.14 shows that

$$\langle \nu, \phi \rangle = \int_{Z_{\text{reg}}} \phi(x) d\text{Vol}_Z(x) = \int_{Z_{\text{reg}} \cap K} \phi(x) \gamma_r(x, \ker D_x f) d\text{Vol}_Z^0(x).$$

Since γ_r is continuous and $K \times \text{Gr}_r(\mathbb{R}^n)$ is compact, the nonnegative function γ_r is bounded by some $C_K > 0$ on this set. Thus $\langle \nu, \phi \rangle \leq C_K \langle \tilde{\nu}, \phi \rangle$. Since $f : \Omega \rightarrow \mathbb{R}^r$ satisfies the hypotheses of Theorem 6.26 and $\tilde{\nu}$ is the measure of integration over its zero set induced by the Euclidean metric, we have $\mathbb{E}[\langle \nu, \phi \rangle^p] \leq C_K^p \mathbb{E}[\langle \tilde{\nu}, \phi \rangle^p] < +\infty$. □

We conclude this section with the proof of Theorem 1.6, which is a corollary of Theorem 1.9.

Proof of Theorem 1.6. Let $f : \Omega \rightarrow \mathbb{R}^r$ be a centered Gaussian field which is C^p and $(p-1)$ -nondegenerate in the sense of Definition 1.4. Then $s = (f, \text{Id})$ is a random section of the trivial bundle $\mathbb{R}^r \times \Omega \rightarrow \Omega$. This s is also a C^p and $(p-1)$ -nondegenerate centered Gaussian field. Its vanishing locus (as a section) is the same as the vanishing locus of f . Hence, the result follows from applying Theorem 1.9 to s . □

7. Multijets adapted to a differential operator

In Theorem 1.1 we defined multijets such that, over the configuration space $(\mathbb{R}^n)^p \setminus \Delta_p \subset C_p[\mathbb{R}^n]$, the p -multijet $\text{mj}_p(f, \underline{x})$ reads as $(f(x_1), \dots, f(x_p))$ in the natural trivialization τ (see the end of Section 5.2). Thus $\text{mj}_p(f, \underline{x})$ is a way to patch together the 0-jets of f at x_i into a smooth object that does not degenerate along Δ_p . In this section, we explain how a similar construction allows us to build a multijet that patches together the k -jets of f at x_i , and more generally the values at x_i of $\mathcal{D}f$, where \mathcal{D} is a differential operator. In Section 7.1 we recall the definition of a differential operator. Then we define a multijet adapted to a given differential operator in Section 7.2. Finally, in Section 7.3, we prove Theorem 1.10.

7.1. Differential operator. In this section, we recall a few fact about differential operators. In the following, we use the multi-index notation introduced in Section 2.2.

Definition 7.1 (differential operator). Let $\Omega \subset \mathbb{R}^n$ be open, let $q, r \geq 1$ and let $d \geq 0$. We say a *differential operator of order at most d* is a linear map $\mathcal{D} : C^d(\Omega, \mathbb{R}^q) \rightarrow C^0(\Omega, \mathbb{R}^r)$ such that there exist continuous

functions $(a_{ij\alpha})_{1 \leq i \leq r; 1 \leq j \leq q; |\alpha| \leq d}$ on Ω such that, for all $f = (f_1, \dots, f_q) \in \mathcal{C}^d(\Omega, \mathbb{R}^q)$,

$$\mathcal{D}(f) : x \longmapsto \left(\sum_{j=1}^q \sum_{|\alpha| \leq d} a_{ij\alpha}(x) \partial^\alpha f_j(x) \right)_{1 \leq i \leq r}. \tag{7-1}$$

More generally, let M be a manifold of dimension n and let $E \rightarrow M$ and $F \rightarrow M$ be two vector bundles of ranks q and r respectively. We say that $\mathcal{D} : \Gamma^d(M, E) \rightarrow \Gamma^0(M, F)$ is a *differential operator of order at most d* if around any point $x \in M$ there exist a chart and local trivializations of E and F such that \mathcal{D} is of the form (7-1) in the corresponding local coordinates. We say that \mathcal{D} is of *order $d \in \mathbb{N}$* if it is of order at most d and not of order at most $d - 1$. If $s \in \Gamma^d(M, E)$ and $x \in M$, we write $\mathcal{D}s = \mathcal{D}(s)$ and $\mathcal{D}_x s = \mathcal{D}(s)(x)$ for simplicity.

Remark 7.2. Let us make some important comments.

- If $\mathcal{D} : \Gamma^d(M, E) \rightarrow \Gamma^0(M, F)$ is a differential operator of order at most d , then it is of the form (7-1) in any set of local coordinates on M, E and F .
- An equivalent definition of a differential operator of order at most d is that it factors linearly through the bundle of d -jets. That is, there exists $L \in \Gamma^0(M, \mathcal{J}_d(M, E)^* \otimes F)$ such that $\mathcal{D}_x s = L(x) j_d(s, x) \in F_x$ for all $s \in \Gamma^d(M, E)$ and $x \in M$.

In the following we always assume that M, E, F and L are smooth. In particular, the functions $(a_{ij\alpha})$ appearing in the local expression (7-1) of \mathcal{D} are smooth. This implies that if $s \in \Gamma^{d+l}(M, E)$ then $\mathcal{D}s \in \Gamma^l(M, F)$.

Example 7.3. The main examples we have in mind are the following.

- The differential $D : \mathcal{C}^1(M) \rightarrow \Gamma^0(M, T^*M)$ is a differential operator of order 1.
- For all $k \in \mathbb{N}$, the jet map $j_k : \Gamma^k(M, E) \rightarrow \Gamma^0(M, \mathcal{J}_k(M, E))$ is a differential operator of order k corresponding to $L(x)$ being the identity of $\mathcal{J}_k(M, E)_x$ for all $x \in M$.
- If M is equipped with a Riemannian metric, the Laplace–Beltrami operator Δ acting on $\mathcal{C}^2(M)$ is a differential operator of order 2.
- If ∇ is a connection on $E \rightarrow M$ then $\nabla : \Gamma^1(M, E) \rightarrow \Gamma^0(M, T^*M \otimes E)$ is a differential operator of order 1. Indeed, in a local frame (e_1, \dots, e_q) of E and local coordinates (x_1, \dots, x_n) on M the covariant derivative of $s = \sum_{j=1}^q f_j e_j \in \Gamma^1(M, E)$ at x is given by

$$\nabla_x s = \sum_{i=1}^n \sum_{j=1}^q \left(\partial_i f_j(x) + \sum_{k=1}^q \mu_{ijk}(x) f_k(x) \right) dx_i \otimes e_j(x),$$

where the (μ_{ijk}) are defined by the relations $\nabla e_k = \sum_{i=1}^n \sum_{j=1}^q \mu_{ijk} dx_i \otimes e_j$ for all $k \in \llbracket 1, q \rrbracket$.

7.2. Multijets adapted to \mathcal{D} . The purpose of this section is to explain how to modify the construction of Section 5 in order to define a multijet bundle adapted to a given differential operator.

Let n, q and $r \geq 1$. We consider a differential operator $\mathcal{D} : \mathcal{C}^d(\mathbb{R}^n, \mathbb{R}^q) \rightarrow \mathcal{C}^0(\mathbb{R}^n, \mathbb{R}^r)$ of order d . As in Remark 7.2, there exists a section L of $\mathcal{J}_d(\mathbb{R}^n, \mathbb{R}^q)^* \otimes \mathbb{R}^r$ such that for any $f \in \mathcal{C}^d(\mathbb{R}^n, \mathbb{R}^q)$ and $x \in \mathbb{R}^n$

we have $\mathcal{D}_x f = L(x) j_d(f, x)$. We assume that for all $x \in \mathbb{R}^n$ the linear map $\mathcal{D}_x : \mathcal{C}^d(\mathbb{R}^n, \mathbb{R}^q) \rightarrow \mathbb{R}^r$ is surjective, which is equivalent to $L(x) : \mathcal{J}_d(\mathbb{R}^n, \mathbb{R}^q)_x \rightarrow \mathbb{R}^r$ being surjective. Moreover, we assume that L is smooth. In this context, we have the following analogue of Theorem 1.1. It holds in particular if $\mathcal{D} = j_k$ or $\mathcal{D} = D$ is the differential.

Theorem 7.4 (existence of multijets adapted to \mathcal{D}). *Let $\mathcal{D} : \mathcal{C}^d(\mathbb{R}^n, \mathbb{R}^q) \rightarrow \mathcal{C}^0(\mathbb{R}^n, \mathbb{R}^r)$ be a differential operator of order d as above. Let $p \geq 1$. There exist a smooth manifold $C_p^{\mathcal{D}}[\mathbb{R}^n]$ of dimension np without boundary and a smooth vector bundle $\mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n) \rightarrow C_p^{\mathcal{D}}[\mathbb{R}^n]$ of rank rp with the following properties:*

- (1) *There exists a smooth proper surjection $\pi : C_p^{\mathcal{D}}[\mathbb{R}^n] \rightarrow (\mathbb{R}^n)^p$ such that $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ is a dense open subset of $C_p^{\mathcal{D}}[\mathbb{R}^n]$, and π restricted to $\pi^{-1}((\mathbb{R}^n)^p \setminus \Delta_p)$ is a \mathcal{C}^∞ -diffeomorphism onto $(\mathbb{R}^n)^p \setminus \Delta_p$.*
- (2) *There exists a map $\text{mj}_p^{\mathcal{D}} : \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^r) \times C_p^{\mathcal{D}}[\mathbb{R}^n] \rightarrow \mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n)$ such that*
 - *for all $z \in C_p^{\mathcal{D}}[\mathbb{R}^n]$, the map $\text{mj}_p^{\mathcal{D}}(\cdot, z) : \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^r) \rightarrow \mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n)_z$ is surjective;*
 - *for all $f \in \mathcal{C}^{l+(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^r)$, the section $\text{mj}_p^{\mathcal{D}}(f, \cdot)$ of $\mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n) \rightarrow C_p^{\mathcal{D}}[\mathbb{R}^n]$ is \mathcal{C}^l .*
- (3) *Let $z \in C_p^{\mathcal{D}}[\mathbb{R}^n]$ be such that $\pi(z) = (x_1, \dots, x_p) \notin \Delta_p$. Then for all $f \in \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^q)$*

$$\text{mj}_p^{\mathcal{D}}(f, z) = 0 \iff \forall i \in \llbracket 1, p \rrbracket, \mathcal{D}_{x_i} f = 0.$$

- (4) *Let $z \in C_p^{\mathcal{D}}[\mathbb{R}^n]$, let $\mathcal{I} = \mathcal{I}(\pi(z))$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\mathcal{I}}^{-1}(\pi(z)) \in (\mathbb{R}^n)^{\mathcal{I}} \setminus \Delta_{\mathcal{I}}$. There exists a linear surjection $\Theta_z^{\mathcal{D}} : \prod_{I \in \mathcal{I}} \mathcal{J}_{(d+1)|I|-1}(\mathbb{R}^n, \mathbb{R}^q)_{y_I} \rightarrow \mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n)_z$ such that*

$$\forall f \in \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^r), \quad \text{mj}_p^{\mathcal{D}}(f, z) = \Theta_z^{\mathcal{D}}((j_{(d+1)|I|-1}(f, y_I))_{I \in \mathcal{I}}).$$

Proof. The proof follows the same strategy as what we did in Sections 4 and 5 in order to prove Theorem 1.1. Let us sketch its main steps.

Let $\underline{x} = (x_1, \dots, x_p) \in (\mathbb{R}^n)^p$ and let $\hat{x} = (\underline{x}, \dots, \underline{x}) \in (\mathbb{R}^n)^{(d+1)p}$. Let $f \in \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^q)$. The polynomial map $K(f, \hat{x}) \in \mathbb{R}_{(d+1)p-1}[X] \otimes \mathbb{R}^q$ is defined as in Definition 3.6. For all $i \in \llbracket 1, p \rrbracket$, since x_i appears with multiplicity $d + 1$ in \hat{x} , the map $K(f, \hat{x})$ has the same d -jet as f at x_i . Hence $\mathcal{D}_{x_i} f = L(x_i) j_d(f, x_i) = L(x_i) j_d(K(f, \hat{x}), x_i) = \mathcal{D}_{x_i}(K(f, \hat{x}))$.

If $\underline{x} \notin \Delta_p$, let us define $\text{ev}_{\underline{x}}^{\mathcal{D}} : P \mapsto (\mathcal{D}_{x_i} P)_{1 \leq i \leq p}$ from $\mathbb{R}_{(d+1)p-1}[X] \otimes \mathbb{R}^q$ to $(\mathbb{R}^r)^p$. Since we assumed that $L(x_i)$ is surjective for all $i \in \llbracket 1, p \rrbracket$, the previous interpolation result proves that $\text{ev}_{\underline{x}}^{\mathcal{D}}$ is surjective. Then, as in (4-2), for all nonempty $I \subset \llbracket 1, p \rrbracket$ we define

$$\mathcal{G}_I^{\mathcal{D}}(\underline{x}) = \ker \text{ev}_{\underline{x}_I}^{\mathcal{D}} \in \text{Gr}_{r|I|}(\mathbb{R}_{(d+1)|I|-1}[X] \otimes \mathbb{R}^q).$$

We also define $\mathcal{G}^{\mathcal{D}}(\underline{x}) = \mathcal{G}_{\llbracket 1, p \rrbracket}^{\mathcal{D}}(\underline{x})$.

Following the same strategy as in Section 5, we denote by $\Sigma_{\mathcal{D}}$ the graph of $(\mathcal{G}_I^{\mathcal{D}})_{I \subset \llbracket 1, p \rrbracket}$ defined on $(\mathbb{R}^n)^p \setminus \Delta_p$. We define $C_p^{\mathcal{D}}[\mathbb{R}^n]$ as a resolution of the singularities of the algebraic variety

$$\bar{\Sigma}_{\mathcal{D}} \subset (\mathbb{R}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{r|I|}(\mathbb{R}_{(d+1)|I|-1}[X] \otimes \mathbb{R}^q).$$

The manifold $C_p^{\mathcal{D}}[\mathbb{R}^n]$ satisfies the analogue of Corollary 5.6. In particular the maps $\mathcal{G}_I^{\mathcal{D}}$ with $I \subset \llbracket 1, p \rrbracket$ extend smoothly to $C_p^{\mathcal{D}}[\mathbb{R}^n]$. Then we define the p -multijet bundle adapted to \mathcal{D} as

$$\mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n) = ((\mathbb{R}_{(d+1)p-1}[X] \otimes \mathbb{R}^q) \times C_p^{\mathcal{D}}[\mathbb{R}^n]) / \mathcal{G}^{\mathcal{D}}$$

over $C_p^{\mathcal{D}}[\mathbb{R}^n]$, similarly to Definition 5.7. Given a function $f \in \mathcal{C}^{(d+1)p-1}(\mathbb{R}^n, \mathbb{R}^q)$, we define its p -multijet adapted to \mathcal{D} at $z \in C_p^{\mathcal{D}}[\mathbb{R}^n]$ as

$$\text{mj}_p^{\mathcal{D}}(f, z) = K(f, \widehat{\pi(z)}) \text{ mod } \mathcal{G}^{\mathcal{D}}(z).$$

Then, following the same steps as in Section 5, one can check that the objects we just defined satisfy the conditions in Theorem 7.4. □

As before, thanks to the localness condition in Theorem 7.4(4), the multijet $\text{mj}_p^{\mathcal{D}}(f, z)$ makes sense even if f is only defined and $\mathcal{C}^{(d+1)|I|-1}$ near y_I for all $I \in \mathcal{I}(\pi(z))$. Hence, the following definition makes sense.

Definition 7.5 (multijets adapted to \mathcal{D}). Let $\Omega \subset \mathbb{R}^n$ be open. We define $C_p^{\mathcal{D}}[\Omega] = \pi^{-1}(\Omega^p)$ and denote by $\mathcal{MJ}_p^{\mathcal{D}}(\Omega)$ the restriction of $\mathcal{MJ}_p^{\mathcal{D}}(\mathbb{R}^n)$ to $C_p^{\mathcal{D}}[\Omega]$. We call $\mathcal{MJ}_p^{\mathcal{D}}(\Omega) \rightarrow C_p^{\mathcal{D}}[\Omega]$ the *bundle of p -multijets adapted to \mathcal{D}* . Let $f : \Omega \rightarrow \mathbb{R}^q$ be of class $\mathcal{C}^{(d+1)p-1}$. We call the section $\text{mj}_p^{\mathcal{D}}(f, \cdot)$ of $\mathcal{MJ}_p^{\mathcal{D}}(\Omega)$ the *p -multijet of f adapted to \mathcal{D}* .

7.3. Finiteness of moments for critical points. The purpose of this section is to prove Theorem 1.10. More generally we prove an analogous result for the zero set of $\mathcal{D}s$, where s is a section of a vector bundle $E \rightarrow M$ and \mathcal{D} is a differential operator; see Theorem 7.8. This is done by adapting what we did in Section 6 to this framework.

Let (M, g) be Riemannian manifold of dimension $n \geq 1$ without boundary. Let $E \rightarrow M$ (resp. $F \rightarrow M$) be a smooth vector bundle of rank $q \geq 1$ (resp. $r \in \llbracket 1, n \rrbracket$). We consider a differential operator $\mathcal{D} : \Gamma^d(M, E) \rightarrow \Gamma^0(M, F)$ of order $d \geq 0$, corresponding to a smooth section $L \in \Gamma^\infty(M, \mathcal{J}_d(M, E)^* \otimes F)$; see Remark 7.2. Thanks to this smoothness assumption we have $\mathcal{D} : \Gamma^{d+l}(M, E) \rightarrow \Gamma^l(M, F)$ for all $l \geq 0$. Finally we assume that $L(x) : \mathcal{J}_d(M, E)_x \rightarrow F_x$ (or equivalently $\mathcal{D}_x : \Gamma^d(M, E) \rightarrow F_x$) is surjective for all $x \in M$.

Let $s : M \rightarrow E$ be a centered Gaussian field on M with values in E in the sense of Definition 6.3. We assume that s is \mathcal{C}^{d+1} and d -nondegenerate, so that $j_d(s, \cdot)$ is a centered Gaussian field with values in $\mathcal{J}_d(M, E)$ which is \mathcal{C}^1 and nondegenerate. Then $\mathcal{D}s \in \Gamma^1(M, F)$ is a centered Gaussian field with values in F which is nondegenerate because of the surjectivity assumption on L . Everything we did in Sections 6.2 and 6.3 applies to $\mathcal{D}s$. In particular, $\mathcal{D}s$ satisfies the weak Bulinskaya lemma (see Proposition 6.8). Hence its vanishing locus is almost surely the union of a codimension- r submanifold of M and a negligible singular set. We denote by $\nu_{\mathcal{D}}$ the random Radon measure on M defined by integrating over the zero set of $\mathcal{D}s$. The formal definition is similar to Definition 6.11.

Example 7.6. Let us assume that $E = \mathbb{R} \times M$ is trivial. Then we can identify $\Gamma^1(M, E)$ with $\mathcal{C}^1(M)$ and consider the differential $D : \mathcal{C}^1(M) \rightarrow \Gamma^0(M, T^*M)$, which is a differential operator of order 1. Let $f : M \rightarrow \mathbb{R}$ be a \mathcal{C}^2 and 1-nondegenerate centered Gaussian field. Then Df is a nondegenerate \mathcal{C}^1

centered Gaussian field on M with values in T^*M , the vanishing locus of Df is the set of critical points of f , and ν_D is the counting measure this random set.

We can now state the analogue of Theorem 6.26 in this context, bearing in mind that Proposition 6.25 applies to ν_D .

Theorem 7.7 (finiteness of moments for ν_D , local version). *Let $\Omega \subset \mathbb{R}^n$ be open and $f : \Omega \rightarrow \mathbb{R}^q$ be a centered Gaussian field. Let $r \in \llbracket 1, n \rrbracket$. Let $\mathcal{D} : \mathcal{C}^d(\Omega, \mathbb{R}^q) \rightarrow \mathcal{C}^0(\Omega, \mathbb{R}^r)$ be a differential operator of order d satisfying the previous hypotheses and ν_D denote the measure of integration over the zero set of $\mathcal{D}f$. Let $p \geq 1$. If f is $\mathcal{C}^{(d+1)p}$ and the Gaussian field $\text{mj}_k^{\mathcal{D}}(f, \cdot) : \mathcal{C}_k^{\mathcal{D}}[\Omega] \rightarrow \mathcal{M}\mathcal{J}_k^{\mathcal{D}}(\Omega)$ is nondegenerate for all $k \in \llbracket 1, p \rrbracket$, then the four equivalent statements in Proposition 6.25 hold for ν_D .*

Proof. Under these hypotheses, for all $k \in \llbracket 1, p \rrbracket$ the Gaussian field $\text{mj}_k^{\mathcal{D}}(f, \cdot)$ is at least \mathcal{C}^1 . Then the proof is the same as that of Theorem 6.26. □

Theorem 7.8 (finiteness of moments for zeros of $\mathcal{D}s$). *In the setting introduced at the beginning of this section, let $s : M \rightarrow E$ be a centered Gaussian field and ν_D denote the measure of integration over the zero set of $\mathcal{D}s$. Let $p \geq 1$. If s is $\mathcal{C}^{(d+1)p}$ and $((d+1)p-1)$ -nondegenerate then $\mathbb{E}[|\langle \nu_D, \phi \rangle|^p] < +\infty$ for all $\phi \in L_c^\infty(M)$.*

Proof. We deduce Theorem 7.8 from Theorem 7.7 in the same way that we deduced Theorem 1.9 from Theorem 6.26; see Section 6.4. □

8. Multijets of holomorphic maps

The purpose of this section is to explain how to adapt what we did in Sections 3 to 6 to the case of holomorphic maps. Theorem 1.6 asks for the $(p-1)$ -nondegeneracy of the field f , that is, $j_{p-1}(f, x)$ needs to be nondegenerate for all x . If $f : \mathbb{C}^n \rightarrow \mathbb{C}$ is a centered holomorphic Gaussian field, then $(f(x), D_x f)$ is always degenerate. Indeed, identifying canonically \mathbb{C} with \mathbb{R}^2 , the differential $D_x f$ takes values in the subspace of $\mathcal{L}(\mathbb{R}^{2n}, \mathbb{R}^2)$ consisting of \mathbb{R} -linear maps that are actually \mathbb{C} -linear. Thus, if we see the holomorphic field f as a smooth field from \mathbb{R}^{2n} to \mathbb{R}^2 , we cannot apply Theorem 1.6. From the point of view of multijets, the multijet $\text{mj}_p(f, \cdot)$ of a holomorphic function $f : \mathbb{C}^n \rightarrow \mathbb{C}$ takes values in a strict sub-bundle of $\mathcal{M}\mathcal{J}_p(\mathbb{R}^{2n}, \mathbb{R}^2)$, which is similar to what happens for jet bundles. Thus, the field $\text{mj}_p(f, \cdot)$ associated with a holomorphic Gaussian field f is necessarily degenerated and Theorem 6.26 does not apply. To remedy this situation, we define in Section 8.1 a multijet bundle adapted to holomorphic maps. Then, in Section 8.2, we use this holomorphic multijet to prove Theorem 1.11.

8.1. Definition of the holomorphic multijet bundles. In this section, we define a multijet bundle for holomorphic maps. Our main result is an equivalent of Theorem 1.1 in this context. Let us first introduce some notation.

Definition 8.1 (spaces of holomorphic maps). We define the following spaces.

- We denote by $\mathbb{C}_d[X]$ the space of complex polynomials of degree at most d in n variables, where $X = (X_1, \dots, X_n)$ is multivariate.

- If M and N are two complex manifolds, we denote by $\mathcal{O}(M, N)$ the space of holomorphic maps from M to N . If $N = \mathbb{C}$, we simply write $\mathcal{O}(M)$.
- If $E \rightarrow M$ is a holomorphic vector bundle, we denote by $\mathcal{J}_k^{\mathbb{C}}(M, E) \rightarrow M$ the holomorphic bundle of k -jets of holomorphic sections of E . If $E = V \times M$ is trivial with fiber V , we denote its holomorphic k -jet bundle by $\mathcal{J}_k^{\mathbb{C}}(M, V) \rightarrow M$. If $V = \mathbb{C}$, we simply write $\mathcal{J}_k^{\mathbb{C}}(M) \rightarrow M$. Given a holomorphic section s of E , we denote by $j_k^{\mathbb{C}}(s, x)$ its holomorphic k -jet at $x \in M$.

Theorem 8.2 (existence of holomorphic multijet bundles). *Let $n \geq 1$ and $p \geq 1$ and let V be a complex vector space of dimension $r \geq 1$. There exist a complex manifold $C_p^{\mathbb{C}}[\mathbb{C}^n]$ of dimension np and a holomorphic vector bundle $\mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V) \rightarrow C_p^{\mathbb{C}}[\mathbb{C}^n]$ of rank rp with the following properties:*

- (1) *There exists a holomorphic proper surjection $\pi : C_p^{\mathbb{C}}[\mathbb{C}^n] \rightarrow (\mathbb{C}^n)^p$ such that $\pi^{-1}((\mathbb{C}^n)^p \setminus \Delta_p)$ is a dense open subset of $C_p^{\mathbb{C}}[\mathbb{C}^n]$, and π restricted to $\pi^{-1}((\mathbb{C}^n)^p \setminus \Delta_p)$ is a biholomorphism onto $(\mathbb{C}^n)^p \setminus \Delta_p$.*
- (2) *There exists a map $\text{mj}_p^{\mathbb{C}} : \mathcal{O}(\mathbb{C}^n, V) \times C_p^{\mathbb{C}}[\mathbb{C}^n] \rightarrow \mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V)$ such that*
 - *for all $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$, the linear map $\text{mj}_p^{\mathbb{C}}(\cdot, z) : \mathcal{O}(\mathbb{C}^n, V) \rightarrow \mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V)_z$ is surjective;*
 - *for all $f \in \mathcal{O}(\mathbb{C}^n, V)$, the section $\text{mj}_p^{\mathbb{C}}(f, \cdot)$ of $\mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V) \rightarrow C_p^{\mathbb{C}}[\mathbb{C}^n]$ is holomorphic.*
- (3) *Let $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$ be such that $\pi(z) = (x_1, \dots, x_p) \notin \Delta_p$. Then for all $f \in \mathcal{O}(\mathbb{C}^n, V)$ we have*

$$\text{mj}_p^{\mathbb{C}}(f, z) = 0 \iff \forall i \in \llbracket 1, p \rrbracket, f(x_i) = 0.$$

- (4) *Let $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$, let $\mathcal{I} = \mathcal{I}(\pi(z))$ and let $(y_I)_{I \in \mathcal{I}} = \iota_{\mathcal{I}}^{-1}(\pi(z)) \in (\mathbb{C}^n)^{\mathcal{I}} \setminus \Delta_{\mathcal{I}}$. There exists a linear surjection $\Theta_z^{\mathbb{C}} : \prod_{I \in \mathcal{I}} \mathcal{J}_{|I|-1}^{\mathbb{C}}(\mathbb{C}^n, V)_{y_I} \rightarrow \mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V)_z$ such that*

$$\forall f \in \mathcal{O}(\mathbb{C}^n, V), \quad \text{mj}_p^{\mathbb{C}}(f, z) = \Theta_z^{\mathbb{C}}((j_{|I|-1}^{\mathbb{C}}(f, y_I))_{I \in \mathcal{I}}).$$

Proof. The proof follows the same steps as what we did in Sections 3, 4 and 5 to prove Theorem 1.1. In the following, we sketch how the proof of Theorem 1.1 adapts to the holomorphic case.

Step 1: Divided differences and Kergin interpolation. Let us consider $f \in \mathcal{O}(\mathbb{C}^n)$ and $\underline{x} = (x_0, \dots, x_k) \in (\mathbb{C}^n)^{k+1}$. The divided difference $f[x_0, \dots, x_k]$ from Definition 3.1 still makes sense. Since f is holomorphic, it is now a symmetric \mathbb{C} -multilinear form on \mathbb{C}^n that depends linearly on f and is holomorphic with respect to \underline{x} . As explained in [Kergin 1980, Proposition 5.1], the Kergin interpolating polynomial is well-behaved with respect to holomorphic maps. Given $f \in \mathcal{O}(\mathbb{C}^n)$ and $\underline{x} \in (\mathbb{C}^n)^p$, (3-3) defines $K(f, \underline{x}) \in \mathbb{C}_{p-1}[X]$ that interpolates the values of $f[\underline{x}_I]$ for all nonempty $I \subset \llbracket 1, p \rrbracket$. The equivalent of Lemma 3.8 is true, in the sense that $K(\cdot, \underline{x})$ is \mathbb{C} -linear and $K(f, \cdot)$ is holomorphic from $(\mathbb{C}^n)^p$ to $\mathbb{C}_{p-1}[X]$.

In this complex framework, the equivalent of Lemma 3.9 holds, that is: for all $\underline{x} \in (\mathbb{C}^n)^p$ the map $P \mapsto (K(P, \underline{x}_I))_{I \in \mathcal{I}(\underline{x})}$ is surjective from $\mathbb{C}_{p-1}[X]$ to $\prod_{I \in \mathcal{I}(\underline{x})} \mathbb{C}_{|I|-1}[X]$. Note however that the proof we gave of Lemma 3.9 does not adapt to the holomorphic setting since it uses bump functions. Here, we deduce the surjectivity of $(K(\cdot, \underline{x}_I))_{I \in \mathcal{I}(\underline{x})}$ from a general amplitude result in algebraic geometry. Let $\mathcal{I} = \mathcal{I}(\underline{x})$ and $(y_I)_{I \in \mathcal{I}} = \iota_{\mathcal{I}}^{-1}(\underline{x})$. For all $I \in \mathcal{I}$, let $P_I \in \mathbb{C}_{|I|-1}[X]$. Multiplying each monomial in P_I by the right power of X_0 yields a homogeneous polynomial $\tilde{P}_I \in \mathbb{C}_{p-1}^{\text{hom}}[X_0, \dots, X_n]$, that is, a global holomorphic section of the line bundle $\mathcal{O}(p-1) \rightarrow \mathbb{C}P^n$. Recall that $\mathcal{O}(p-1)$ is the $(p-1)$ -th tensor

power of the hyperplane line bundle $\mathcal{O}(1) \rightarrow \mathbb{C}P^n$. Since $\mathcal{O}(1)$ is very ample, the bundle $\mathcal{O}(p-1)$ is $(p-1)$ -jet ample; see [Beltrametti and Sommese 1993, Corollary 2.1]. This means that there exists $\tilde{P} \in \mathbb{C}_{p-1}^{\text{hom}}[X_0, \dots, X_n]$ with the same $(|I|-1)$ -jet as \tilde{P}_I at y_I for all $I \in \mathcal{I}$, where we see \mathbb{C}^n as a standard affine chart in $\mathbb{C}P^n$. Then, for all $I \in \mathcal{I}$, the polynomial $P = \tilde{P}(1, X_1, \dots, X_n) \in \mathbb{C}_{p-1}[X]$ has the same $(|I|-1)$ -jet (i.e., the same Taylor polynomial of order $|I|-1$) as P_I at y_I . Thus $(K(P, \underline{x}_I))_{I \in \mathcal{I}} = (P_I)_{I \in \mathcal{I}}$.

Step 2: Kernel of the evaluation and resolution of singularities. As in Definition 4.3, we define a complex evaluation map by $\text{ev}_{\underline{x}}^{\mathbb{C}} : f \mapsto (f(x_1), \dots, f(x_p))$, where $\underline{x} \in (\mathbb{C}^n)^p$. If $\underline{x} \notin \Delta_p$, this map is surjective from $\mathbb{C}_{p-1}[X]$ to \mathbb{C}^p . Hence we can define $\mathcal{G}_I^{\mathbb{C}}(\underline{x}) = \ker \text{ev}_{\underline{x}_I}^{\mathbb{C}} \in \text{Gr}_{|I|}(\mathbb{C}_{|I|-1}[X])$ for all nonempty $I \subset \llbracket 1, p \rrbracket$, where the Grassmannian is now the Grassmannian of complex subspaces of codimension $|I|$. Then, everything we did in Sections 4 and 5 works in the holomorphic setting after replacing \mathbb{R} -linear objects by \mathbb{C} -linear ones.

We define $\Sigma_{\mathbb{C}}$ as the graph of $(\mathcal{G}_I^{\mathbb{C}})_{I \subset \llbracket 1, p \rrbracket}$ from $(\mathbb{C}^n)^p \setminus \Delta_p$ to $\prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{C}_{|I|-1}[X])$ and $C_p^{\mathbb{C}}[\mathbb{C}^n]$ as a resolution of the singularities of its closure $\bar{\Sigma}_{\mathbb{C}}$ in $(\mathbb{C}^n)^p \times \prod_{\emptyset \neq I \subset \llbracket 1, p \rrbracket} \text{Gr}_{|I|}(\mathbb{C}_{|I|-1}[X])$. The resolution of singularities is a result from algebraic geometry which holds over fields of characteristic 0. In particular, in Proposition 5.5 and Corollary 5.6, “smooth” can be replaced by “algebraic” everywhere. The same results hold over \mathbb{C} , in which case algebraic implies holomorphic. Thus, $C_p^{\mathbb{C}}[\mathbb{C}^n]$ is a complex manifold of dimension np , which satisfies the equivalent of Corollary 5.6 with “smooth” replaced by “holomorphic”.

Step 3: Definition of the holomorphic multijet bundles. Everything we did in Sections 5.2, 5.3 and 5.4 adapts to the holomorphic setting. It is enough to replace C^k functions by holomorphic ones and to write the linear arguments over \mathbb{C} instead of \mathbb{R} . We can then define the holomorphic vector bundle of p -multijets of holomorphic functions on \mathbb{C}^n by

$$\mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n) = (\mathbb{C}_{p-1}[X] \times C_p^{\mathbb{C}}[\mathbb{C}^n]) / \mathcal{G}^{\mathbb{C}} \tag{8-1}$$

and the p -multijet of $f \in \mathcal{O}(\mathbb{C}^n)$ by $\text{mj}_p^{\mathbb{C}}(f, z) = K(f, \pi(z)) \bmod \mathcal{G}^{\mathbb{C}}(z)$ for all $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$. If V is a complex vector space of finite dimension, we define as in Definition 5.18

$$\mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V) = \mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n) \otimes V. \tag{8-2}$$

Then we define the p -multijet of $f \in \mathcal{O}(\mathbb{C}^n, V)$ as in Definition 5.19. If (v_1, \dots, v_r) is a basis of V and $f = \sum_{i=1}^r f_i v_i$ is holomorphic, then $\text{mj}_p^{\mathbb{C}}(f, z) = \sum_{i=1}^r \text{mj}_p^{\mathbb{C}}(f_i, z) \otimes v_i$ for all $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$. As in Lemma 5.20, this definition does not depend on a choice of basis. This defines the holomorphic p -multijet that we are looking for. \square

As in the real case, thanks to (4) in Theorem 8.2, the holomorphic multijet $\text{mj}_p^{\mathbb{C}}(f, z)$ of f at $z \in C_p^{\mathbb{C}}[\mathbb{C}^n]$ only depends on the germ of f near the x_i , where $(x_i)_{1 \leq i \leq p} = \pi(z)$. Thus, we can define a holomorphic multijet bundle over any open subset of \mathbb{C}^n .

Definition 8.3 (holomorphic multijets). Let $\Omega \subset \mathbb{C}^n$ be open. We denote by $C_p^{\mathbb{C}}[\Omega] = \pi^{-1}(\Omega^p)$ and by $\mathcal{MJ}_p^{\mathbb{C}}(\Omega, V) \rightarrow C_p^{\mathbb{C}}[\Omega]$ the restriction of $\mathcal{MJ}_p^{\mathbb{C}}(\mathbb{C}^n, V)$ to $C_p^{\mathbb{C}}[\Omega]$. If $V = \mathbb{C}$, we drop it from the notation and write $\mathcal{MJ}_p^{\mathbb{C}}(\Omega) \rightarrow C_p^{\mathbb{C}}[\Omega]$. Let $f \in \mathcal{O}(\Omega, V)$, we call the section $\text{mj}_p^{\mathbb{C}}(f, \cdot)$ of $\mathcal{MJ}_p^{\mathbb{C}}(\Omega, V)$ the *holomorphic p -multijet of f* .

8.2. Application to the zeros of holomorphic Gaussian fields. In this section, we explain how the holomorphic multijets defined in Section 8.1 allow us to prove Theorem 1.11, and the analogue of Theorem 6.26 for holomorphic Gaussian fields. We start by recalling a few facts about complex Gaussian vectors; see [Andersen et al. 1995, Chapter 2].

A random variable $X \in \mathbb{C}$ is called a *centered complex Gaussian* if its real and imaginary parts are independent real centered Gaussian variables of the same variance, i.e., there exists $\lambda \geq 0$ such that $X = X_{\Re} + iX_{\Im}$, with $(X_{\Re}, X_{\Im}) \sim \mathcal{N}(0, \lambda \text{Id})$ in \mathbb{R}^2 .

Definition 8.4 (complex Gaussian vector). We say that a random vector X in a finite-dimensional complex vector space V is a *centered Gaussian* if for all $\eta \in V^*$ the complex variable $\eta(X)$ is a centered complex Gaussian.

If V is equipped with a Hermitian inner product $\langle \cdot, \cdot \rangle$ and we define $v^* = \langle v, \cdot \rangle$ then the *variance* of X is the nonnegative Hermitian operator $\text{Var}_{\mathbb{C}}(X) = \mathbb{E}[X \otimes X^*]$. We say that X is *nondegenerate* if $\text{Var}_{\mathbb{C}}(X)$ is positive-definite.

Remark 8.5. As in the real case, the Gaussianity and nondegeneracy of X do not depend on $\langle \cdot, \cdot \rangle$, but the variance operator does.

A centered complex Gaussian vector X in $(V, \langle \cdot, \cdot \rangle)$ is completely determined by its variance. For example, if $\text{Var}_{\mathbb{C}}(X) = \Lambda$ is positive-definite, then X admits the density $v \mapsto e^{-\langle v, \Lambda^{-1}v \rangle} / \det(\pi \Lambda)$ with respect to the Lebesgue measure on V . We denote by $\mathcal{N}_{\mathbb{C}}(0, \Lambda)$ the centered complex Gaussian distribution of variance Λ . Then $X \sim \mathcal{N}_{\mathbb{C}}(0, \Lambda)$ in \mathbb{C}^n if and only if its real and imaginary part satisfy

$$(X_{\Re}, X_{\Im}) \sim \mathcal{N}\left(0, \frac{1}{2} \begin{pmatrix} \Re(\Lambda) & \Im(\Lambda) \\ -\Im(\Lambda) & \Re(\Lambda) \end{pmatrix}\right)$$

in \mathbb{R}^{2n} .

Let $E \rightarrow M$ be a holomorphic vector bundle over a complex manifold M . We denote by $H^0(M, E)$ the vector space of global holomorphic sections of $E \rightarrow M$.

Definition 8.6 (holomorphic Gaussian field). We say that a random section $s \in H^0(M, E)$ is a *centered holomorphic Gaussian field* if for all $m \geq 1$ and all x_1, \dots, x_m the random vector $(s(x_1), \dots, s(x_m))$ is a centered complex Gaussian. We say that this field is *nondegenerate* if $s(x)$ is nondegenerate for all $x \in M$.

Note that if $s \in H^0(M, E)$ is a centered holomorphic Gaussian field then, for all $k \in \mathbb{N}$, its holomorphic k -jet $j_k^{\mathbb{C}}(s, \cdot)$ defines a centered holomorphic Gaussian field with values in $\mathcal{J}_k^{\mathbb{C}}(M, E)$.

Definition 8.7 (p -nondegeneracy for holomorphic fields). Let $p \geq 1$. We say that the centered holomorphic Gaussian field $s \in H^0(M, E)$ is *p -nondegenerate* if the centered complex Gaussian $j_p^{\mathbb{C}}(s, x) \in \mathcal{J}_p^{\mathbb{C}}(M, E)_x$ is nondegenerate for all $x \in M$.

As in the real framework, we need the following definition.

Definition 8.8 (complex Jacobian). Let $L : V \rightarrow V'$ be a \mathbb{C} -linear map between Hermitian spaces and let L^* denote its adjoint map. The *complex Jacobian* of L is defined as $\text{Jac}_{\mathbb{C}}(L) = \det(LL^*)$.

Remark 8.9. If we see V, V' and L as \mathbb{R} -linear objects and we equip V and V' with the Euclidean structures induced by their Hermitian inner products, then the real and complex Jacobians are related by $\text{Jac}_{\mathbb{C}}(L) = \text{Jac}(L)^2$; see [Andersen et al. 1995, Theorem A.5].

Let us consider a complex manifold M of complex dimension n equipped with a Riemannian metric g and a holomorphic vector bundle $E \rightarrow M$ of complex rank $r \in \llbracket 1, n \rrbracket$. In the following, we denote by ∇ a connection on E which is compatible with the complex structure. As in the real case, the choice of this connection will not matter. Let $s \in H^0(M, E)$ be a centered holomorphic Gaussian field on M with values in E and let $Z = s^{-1}(0)$ denote its vanishing locus. We will always assume that s is nondegenerate. In this setting, the random section s satisfies a strong Bulinskaya-type lemma.

Proposition 8.10 (holomorphic Bulinskaya lemma). *Almost surely the following set is empty:*

$$\{x \in M \mid s(x) = 0 \text{ and } \text{Jac}_{\mathbb{C}}(\nabla_x s) = 0\}.$$

In particular, the zero set Z is almost surely a (possibly empty) complex submanifold of complex codimension r in M .

Proof. It is enough to check the result locally. On an open subset $\Omega \subset M$ over which E is trivial, we can consider s as a nondegenerate smooth centered Gaussian field from Ω to $\mathbb{C}^r \simeq \mathbb{R}^{2r}$. Then, the local result follows from [Lerario and Stecconi 2019, Theorem 7]. □

Let us consider Z as random submanifold of real codimension $2r$ in the Riemannian manifold (M, g) of real dimension $2n$. The metric g induces a $(2n - 2r)$ -dimensional Riemannian volume $d\text{Vol}_Z$ on Z and we can define ν as in Definition 6.11, bearing in mind that $Z = Z_{\text{reg}}$. Then Propositions 6.17, 6.24 and 6.25 hold for the holomorphic field s and the associated linear statistics $\langle \nu, \phi \rangle$ with $\phi \in L_c^\infty(M)$. More generally, everything we did in Sections 6.2, 6.3 and 6.4 adapts to the holomorphic setting.

Remark 8.11. In Definition 6.23, the function ρ_p is defined in terms of real Jacobians and the variance of $(s(x_1), \dots, s(x_p))$ seen as a real Gaussian vector. One can check that another expression of $\rho_p(x_1, \dots, x_p)$ is the following, which is more natural in our holomorphic framework:

$$\forall (x_1, \dots, x_p) \notin \Delta_p, \quad \rho_p(x_1, \dots, x_p) = \frac{\mathbb{E}\left[\prod_{i=1}^p \text{Jac}_{\mathbb{C}}(\nabla_{x_i} s) \mid \forall i \in \llbracket 1, p \rrbracket, s(x_i) = 0\right]}{\det(\pi \text{Var}_{\mathbb{C}}(s(x_1), \dots, s(x_p)))}.$$

We can now state the equivalent of Theorem 6.26 for holomorphic Gaussian fields. Let $\Omega \subset \mathbb{C}^n$ be open. Recall that $\mathcal{M}\mathcal{J}_p^{\mathbb{C}}(\Omega, \mathbb{C}^r) \rightarrow C_p^{\mathbb{C}}[\Omega]$ is defined in Definition 8.3 as the restriction over $C_p^{\mathbb{C}}[\Omega] \subset C_p^{\mathbb{C}}[\mathbb{C}^n]$ of the vector bundle $\mathcal{M}\mathcal{J}_p^{\mathbb{C}}(\mathbb{C}^n, \mathbb{C}^r) \rightarrow C_p^{\mathbb{C}}[\mathbb{C}^n]$ from Theorem 8.2.

Theorem 8.12 (finiteness of moments for holomorphic fields, local version). *Let $f : \Omega \rightarrow \mathbb{C}^r$ be a centered holomorphic Gaussian field and ν be as in Definition 6.11. Let $p \geq 1$. If for all $k \in \llbracket 1, p \rrbracket$ the holomorphic Gaussian field $\text{mj}_k^{\mathbb{C}}(f, \cdot) \in H^0(C_k^{\mathbb{C}}[\Omega], \mathcal{M}\mathcal{J}_k^{\mathbb{C}}(\Omega, \mathbb{C}^r))$ is nondegenerate, then the four equivalent statements in Proposition 6.25 hold.*

Proof. The proof of Theorem 6.26 relies mostly on two facts that are valid for all $k \in \llbracket 1, p \rrbracket$. First, on the open dense subset $\Omega^k \setminus \Delta_k \subset C_k[\Omega]$, the zero set of $\text{mj}_k(f, \cdot)$ is the same as that of $(x_1, \dots, x_k) \mapsto$

$(f(x_1), \dots, f(x_k))$. And second, the field $\text{mj}_k(f, \cdot)$ is nondegenerate on $C_k[\Omega]$, so that we can apply the Kac–Rice formula (Proposition 6.17) to the k -multijet.

These two facts are still true in the present holomorphic setting. Hence, the same proof as that of Theorem 6.26 yields the result. \square

We deduce from this result the equivalent of Theorem 1.9 for a centered holomorphic Gaussian field s on a complex manifold M of dimension n with values in a holomorphic vector bundle E of rank $r \in \llbracket 1, n \rrbracket$. Theorem 1.11 is a special case of the following.

Theorem 8.13 (finiteness of moments for zeros of holomorphic Gaussian sections). *Let $p \geq 1$, let $s \in H^0(M, E)$ be a centered holomorphic Gaussian field and let v be as in Definition 6.11. If s is $(p-1)$ -nondegenerate then $\mathbb{E}[|\langle v, \phi \rangle|^p] < +\infty$ for all $\phi \in L_c^\infty(M)$.*

Proof. We deduce Theorem 8.13 from Theorem 8.12 in the same way that we deduced Theorem 1.9 from Theorem 6.26; see Section 6.4. \square

Remark 8.14. In particular, Theorem 8.13 proves the local integrability of the p -points correlation functions studied in [Bleher et al. 2000] and their scaling limit.

Acknowledgments

The authors thank Vincent Borrelli, Nicolas Vichery and Gabriel Rivière for independently suggesting to take a look at various compactifications of configuration spaces.

References

- [Ancona 2021] M. Ancona, “Random sections of line bundles over real Riemann surfaces”, *Int. Math. Res. Not.* **2021**:9 (2021), 7004–7059. MR Zbl
- [Ancona and Letendre 2021] M. Ancona and T. Letendre, “Zeros of smooth stationary Gaussian processes”, *Electron. J. Probab.* **26** (2021), art. id. 68. MR Zbl
- [Andersen et al. 1995] H. H. Andersen, M. Højbjerg, D. Sørensen, and P. S. Eriksen, *Linear and graphical models for the multivariate complex normal distribution*, Lecture Notes in Stat. **101**, Springer, 1995. MR Zbl
- [Armentano et al. 2019] D. Armentano, J.-M. Azaïs, D. Ginsbourger, and J. R. León, “Conditions for the finiteness of the moments of the volume of level sets”, *Electron. Commun. Probab.* **24** (2019), art. id. 17. MR Zbl
- [Armentano et al. 2023a] D. Armentano, J. M. Azaïs, F. Dalmao, J. R. León, and E. Mordecki, “On the finiteness of the moments of the measure of level sets of random fields”, *Braz. J. Probab. Stat.* **37**:1 (2023), 219–245. MR Zbl
- [Armentano et al. 2023b] D. Armentano, J.-M. Azaïs, and J. R. León, “On a general Kac–Rice formula for the measure of a level set”, preprint, 2023. arXiv 2304.07424
- [Axelrod and Singer 1994] S. Axelrod and I. M. Singer, “Chern–Simons perturbation theory, II”, *J. Differential Geom.* **39**:1 (1994), 173–213. MR Zbl
- [Azaïs and Wschebor 2009] J.-M. Azaïs and M. Wschebor, *Level sets and extrema of random processes and fields*, Wiley, Hoboken, NJ, 2009. MR Zbl
- [Beliaev et al. 2024] D. Beliaev, M. McAuley, and S. Muirhead, “A central limit theorem for the number of excursion set components of Gaussian fields”, *Ann. Probab.* **52**:3 (2024), 882–922. MR Zbl
- [Beltrametti and Sommese 1993] M. C. Beltrametti and A. J. Sommese, “On k -jet ampleness”, pp. 355–376 in *Complex analysis and geometry*, edited by V. Ancona and A. Silva, Springer, 1993. MR Zbl

- [Belyaev 1966] Y. K. Belyaev, “On the number of intersections of a level by a Gaussian stochastic process, I”, *Theory Probab. Appl.* **11**:1 (1966), 106–113. Zbl
- [Bleher et al. 2000] P. Bleher, B. Shiffman, and S. Zelditch, “Universality and scaling of correlations between zeros on complex manifolds”, *Invent. Math.* **142**:2 (2000), 351–395. MR Zbl
- [Bojanov et al. 1993] B. D. Bojanov, H. A. Hakopian, and A. A. Sahakian, *Spline functions and multivariate interpolations*, Math. Appl. **248**, Kluwer, Dordrecht, 1993. MR Zbl
- [Evain 2005] L. Evain, “Compactifications des espaces de configuration dans les schémas de Hilbert”, *Bull. Soc. Math. France* **133**:4 (2005), 497–539. MR Zbl
- [Fulton and MacPherson 1994] W. Fulton and R. MacPherson, “A compactification of configuration spaces”, *Ann. of Math. (2)* **139**:1 (1994), 183–225. MR Zbl
- [Gass and Stecconi 2024] L. Gass and M. Stecconi, “The number of critical points of a Gaussian field: finiteness of moments”, *Probab. Theory Related Fields* **190**:3-4 (2024), 1167–1197. MR Zbl
- [Hauser 2003] H. Hauser, “The Hironaka theorem on resolution of singularities (or: A proof we always wanted to understand)”, *Bull. Amer. Math. Soc. (N.S.)* **40**:3 (2003), 323–403. MR Zbl
- [Hironaka 1964a] H. Hironaka, “Resolution of singularities of an algebraic variety over a field of characteristic zero, I”, *Ann. of Math. (2)* **79**:1 (1964), 109–203. MR Zbl
- [Hironaka 1964b] H. Hironaka, “Resolution of singularities of an algebraic variety over a field of characteristic zero, II”, *Ann. of Math. (2)* **79**:2 (1964), 205–326. MR Zbl
- [Kergin 1980] P. Kergin, “A natural interpolation of C^K functions”, *J. Approx. Theory* **29**:4 (1980), 278–293. MR Zbl
- [Kollár 2007] J. Kollár, *Lectures on resolution of singularities*, Ann. of Math. Stud. **166**, Princeton Univ. Press, 2007. MR Zbl
- [Le Barz 1988] P. Le Barz, “La variété des triplets complets”, *Duke Math. J.* **57**:3 (1988), 925–946. MR Zbl
- [Lee 2018] J. M. Lee, *Introduction to Riemannian manifolds*, 2nd ed., Grad. Texts in Math. **176**, Springer, 2018. MR Zbl
- [Lerario and Stecconi 2019] A. Lerario and M. Stecconi, “Differential topology of Gaussian random fields”, preprint, 2019. arXiv 1902.03805
- [Lorentz 2000] R. A. Lorentz, “Multivariate Hermite interpolation by algebraic polynomials: a survey”, *J. Comput. Appl. Math.* **122**:1-2 (2000), 167–201. MR Zbl
- [Malevich and Volodina 1993] T. L. Malevich and L. N. Volodina, “Some finiteness conditions for factorial moments of the number of zeros of Gaussian field zeros”, *Theory Probab. Appl.* **38**:1 (1993), 27–45. Zbl
- [Micchelli and Milman 1980] C. A. Micchelli and P. Milman, “A formula for Kergin interpolation in \mathbb{R}^k ”, *J. Approx. Theory* **29**:4 (1980), 294–296. MR Zbl
- [Nazarov and Sodin 2012] F. Nazarov and M. Sodin, “Correlation functions for random complex zeroes: strong clustering and local universality”, *Comm. Math. Phys.* **310**:1 (2012), 75–98. MR Zbl
- [Olver 2001] P. J. Olver, “Geometric foundations of numerical algorithms and symmetry”, *Appl. Algebra Engrg. Comm. Comput.* **11**:5 (2001), 417–436. MR Zbl
- [Priya 2020] L. Priya, “Overcrowding estimates for zero count and nodal length of stationary Gaussian processes”, preprint, 2020. arXiv 2012.10857
- [Saunders 1989] D. J. Saunders, *The geometry of jet bundles*, Lond. Math. Soc. Lect. Note Ser. **142**, Cambridge Univ. Press, 1989. MR Zbl
- [Sinha 2004] D. P. Sinha, “Manifold-theoretic compactifications of configuration spaces”, *Selecta Math. (N.S.)* **10**:3 (2004), 391–428. MR Zbl
- [Ulyanov 2002] A. P. Ulyanov, “Polydiagonal compactification of configuration spaces”, *J. Algebraic Geom.* **11**:1 (2002), 129–159. MR Zbl
- [Włodarczyk 2005] J. Włodarczyk, “Simple Hironaka resolution in characteristic zero”, *J. Amer. Math. Soc.* **18**:4 (2005), 779–822. MR Zbl

MICHELE ANCONA: michele.ancona@unice.fr
Université Côte d'Azur, CNRS, LJAD, Nice, France

THOMAS LETENDRE: letendre@math.cnrs.fr
Université Paris-Saclay, CNRS, LMO, Orsay, France

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: "Linear Ramp"

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2025 is US \$475/year for the electronic version, and \$735/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 18 No. 6 2025

Damped Strichartz estimates and the incompressible Euler–Maxwell system DIOGO ARSÉNIO and HAROUNE HOUAMED	1309
Equivariant property Gamma and the tracial local-to-global principle for C^* -dynamics GÁBOR SZABÓ and LISE WOUTERS	1385
Multijet bundles and application to the finiteness of moments for zeros of Gaussian fields MICHELE ANCONA and THOMAS LETENDRE	1433
Existence of solutions to a fractional semilinear heat equation in uniformly local weak Zygmund-type spaces NORISUKE IOKU, KAZUHIRO ISHIGE and TATSUKI KAWAKAMI	1477
A Marcinkiewicz multiplier theory for Schur multipliers CHIAN YEONG CHUAH, ZHEN-CHUAN LIU and TAO MEI	1511
Double duals and Hilbert modules HUAXIN LIN	1531