

# ANALYSIS & PDE

Volume 18

No. 8

2025

QIZHI ZHAO

THE  $L^\infty$  ESTIMATE  
FOR PARABOLIC COMPLEX MONGE-AMPÈRE EQUATIONS



# THE $L^\infty$ ESTIMATE FOR PARABOLIC COMPLEX MONGE–AMPÈRE EQUATIONS

QIZHI ZHAO

Following the recent developments in Chen and Cheng (2023) and Guo et al. (2023), we derive the  $L^\infty$  estimate for Kähler–Ricci flows under certain integral assumptions. The technique also extends to some other parabolic Monge–Ampère equations derived from Kähler geometry and  $G_2$  geometry.

## 1. Introduction

We will derive the  $L^\infty$  estimate for the Kähler–Ricci flow

$$\begin{cases} \partial_t \varphi = \log \left( \frac{\omega_\varphi^n}{e^{nF} \omega_0^n} \right), \\ \varphi(\cdot, 0) = \varphi_0(\cdot) \end{cases} \quad (1-1)$$

under the assumption that the  $p$ -entropy  $\text{Ent}_p(F) = \int_M |F|^p e^{nF} \omega_0^n$  is bounded and  $\int_M F \omega_0^n$  has a lower bound. Here is our main theorem.

**Theorem 1.1.** *Let us consider the flow equation (1-1) on  $M \times [0, T)$ , where  $M$  is an  $n$ -dimensional compact Kähler manifold. Let  $F$  be a space function, i.e.,  $F : M \rightarrow \mathbb{R}$ . Assume, for some  $p > n + 1$ , the  $p$ -entropy  $\text{Ent}_p(F) = \int_M |F|^p e^{nF} \omega_0^n$  is bounded and  $\int_M nF \omega_0^n \geq -K$ . Moreover, suppose  $\varphi$  is a  $C^2$  solution, and let  $\tilde{\varphi} = \varphi - \int \varphi \omega_0^n$  be a normalization which has the zero integral. Then we have the  $L^\infty$  estimate*

$$\|\tilde{\varphi}\|_{L^\infty(M \times [0, T))} \leq C,$$

where  $C$  depends on  $n$ ,  $\omega_0$ ,  $\varphi_0$ ,  $p$ ,  $K$ , and  $\text{Ent}_p(F)$ . Most importantly, such a  $C$  does not depend on  $T$ .

Yau [1978] applied the method of Moser iteration to derive the  $L^\infty$  estimate for Monge–Ampère equations when  $\|e^{nF}\|_{L^p}$  is bounded for some  $p > n$ . Later, Kołodziej [2003] gave another proof by using the pluripotential theory under a weaker assumption that  $\|e^{nF}\|_{L^p}$  is bounded for some  $p > 1$ . More recently, Guo, Phong, and Tong [Guo et al. 2023] recovered Kołodziej’s estimate by a PDE method which was partly motivated by the breakthrough on the cscK metric of Chen and Cheng [2021].

The Kähler–Ricci flow was firstly studied by Cao [1985] when he gave an alternative proof of Calabi’s conjecture for  $c_1(M) = 0$  and  $c_1(M) < 0$ , which investigated the estimates for the Kähler–Ricci flow instead of Monge–Ampère equations. There are abundant results on Kähler–Ricci flow, see [Eyssidieux et al. 2015; 2016; Guedj et al. 2021; Jian and Shi 2024]. Our result requires a weaker regularity on the right-hand side than Cao’s  $L^\infty$  estimate and can be viewed as a parabolic analogue of [Guo et al. 2023; Wang et al. 2021].

MSC2020: 53E30, 58J90.

Keywords: auxiliary equations, energy estimates, parabolic complex Monge–Ampère equations.

There are some technical improvements in our paper compared with previous results in [Chen and Cheng 2023; Guo and Phong 2023; 2024; Guo et al. 2023]. The difficulty for the flow problem is that we want to derive an  $L^\infty$  estimate independent of  $T$ . But the original auxiliary equation may not serve as a good choice. Our approach is to consider a local version of auxiliary flows instead, see Section 2. Compared with the elliptic version of  $L^\infty$  estimates, our theorem requires an extra integral condition. Rewriting (1-1) by  $\omega_\varphi^n = e^{\dot{\varphi}+nF} \omega_0^n$ , we could see that the  $L^\infty$  estimate of  $\varphi$  comes from some  $p$ -entropy bounds on  $e^{\dot{\varphi}+nF}$ . Roughly speaking, we need not only the  $p$ -entropy bound controls on  $e^{nF}$  but also some upper bounds on  $\dot{\varphi}$ . In Kähler–Ricci flow, Cao proved the supremum of  $\dot{\varphi}$  can be controlled by the infimum of  $e^{nF}$  when  $F$  is smooth. Indeed such estimates can be generalized to general parabolic Monge–Ampère flows. However, in our theorem, we can improve the pointwise condition by some integral condition on  $F$ .

There are two directions to generalize Theorem 1.1. As in [Chen and Cheng 2023], we can replace the Monge–Ampère operator on the right-hand side by a more general nonlinear operator  $\mathcal{F}$ . Write  $\omega_0 = \sqrt{-1}g_{j\bar{m}} dz^j \wedge d\bar{z}^m$  in local coordinate; then the corresponding endomorphism  $h_\varphi$ , which is relative to  $\omega_\varphi$ , can be expressed by  $(h_\varphi)_k^j = g^{j\bar{m}}(\omega_\varphi)_{\bar{m}k}$  in local coordinate. Let  $\lambda[h_\varphi]$  be the vector of eigenvalues of  $h_\varphi$ , and consider the nonlinear operator  $\mathcal{F} : \Gamma \subset \mathbb{R}^n \rightarrow \mathbb{R}_+$  with the following four conditions:

- (1) The domain  $\Gamma$  is a symmetric cone with  $\Gamma_n \subset \Gamma \subset \Gamma_1$ , where  $\Gamma_k$  is defined to be the cone of vectors  $\lambda$  with  $\sigma_j(\lambda) > 0$  for  $1 \leq j \leq k$ , where  $\sigma_j$  is the  $j$ -th symmetric polynomial in  $\lambda$ .
- (2)  $\mathcal{F}(\lambda)$  is symmetric in  $\lambda \in \Gamma$  and it is of homogeneous degree  $r$ .
- (3)  $\frac{\partial \mathcal{F}}{\partial \lambda_j} > 0$  for each  $j = 1, \dots, n$  and  $\lambda \in \Gamma$ .
- (4) There is a  $\gamma > 0$  such that

$$\prod_{j=1}^n \frac{\partial \mathcal{F}(\lambda)}{\partial \lambda_j} \geq \gamma \mathcal{F}^{n(1-1/r)} \quad \text{for all } \lambda \in \Gamma. \tag{1-2}$$

The above requirements come from [Guo et al. 2023], and there is a slight modification on the last condition since the homogeneous degree of  $\mathcal{F}$  is  $r$  under our assumption. The complex Hessian operators and  $p$ -Monge–Ampère operators are examples. More examples can be found in [Harvey and Lawson 2023]. Here is our first generalization.

**Theorem 1.2.** *Let  $\varphi$  be a  $C^2$  solution of the flow*

$$\begin{cases} \partial_t \varphi = \log\left(\frac{\mathcal{F}(\lambda[h_\varphi])}{e^{rF}}\right), \\ \varphi(\cdot, 0) = \varphi_0 \end{cases} \tag{1-3}$$

*on  $M \times [0, T)$ , where  $M$  is an  $n$ -dimensional compact Kähler manifold. Let  $F : M \rightarrow \mathbb{R}$  be a space function. Assume, for some  $p > n + 1$ , the  $p$ -entropy  $\text{Ent}_p(F) = \int_M |F|^p e^{nF} \omega_0^n$  is bounded and  $\int_M F \omega_0^n \geq -K$ . Then we have the  $L^\infty$  estimate on the normalization  $\tilde{\varphi}$*

$$\|\tilde{\varphi}\|_{L^\infty(M \times [0, T))} \leq C,$$

*where  $C$  depends on  $n, \omega_0, \varphi_0, p, K, \gamma, r$ , and  $\text{Ent}_p(F)$ .*

Another direction of generalization is motivated by Chen and Cheng [2023], who considered the  $L^\infty$  estimate for the inverse Monge–Ampère flow

$$\begin{cases} (-\partial_t u)\omega_\varphi^n = e^{nF}\omega_0^n, \\ \varphi(\cdot, 0) = \varphi_0. \end{cases} \tag{1-4}$$

Indeed, we can consider the general complex Monge–Ampère flow

$$\begin{cases} \partial_t \varphi = \Theta\left(\frac{\omega_\varphi^n}{e^{nF}\omega_0^n}\right), \\ \varphi(\cdot, 0) = \varphi_0, \end{cases} \tag{1-5}$$

where  $\Theta : \mathbb{R}_+ \rightarrow \mathbb{R}$  is a strictly increasing smooth function. Picard and Zhang [2020] proved the long time existence and convergence of the flow (1-5) under the assumption that  $F \in C^\infty(M, \mathbb{R})$ . It is the Kähler–Ricci flow when  $\Theta(y) = \log y$  and the inverse Monge–Ampère flow (1-4) when  $\Theta(y) = -1/y$ . The general parabolic Monge–Ampère flow (1-5) also arises from many other geometric problems. For example, when  $\Theta(y) = y$ , this is the flow reduced from the anomaly flow with conformal Kähler initial data; see [Phong et al. 2019]. When  $\Theta(y) = y^{1/3}$ , this is the reduction of the  $G_2$ -Laplacian flow over a seven dimensional manifold [Picard and Suan 2024]. We can apply the new technique to prove the  $L^\infty$  estimate of the solution  $\varphi$  to the flow (1-5) under an analogue assumption on  $F$ .

**Theorem 1.3.** *Assume  $\Theta(y) = -1/y$ ,  $y$ , or  $y^{1/3}$ , and  $\text{Ent}_p(F)$  is bounded. Moreover, consider the constant  $K$  equal to  $\max(0, \int_M \Theta(e^{-nF})\omega_0^n)$ . Then, there exists a constant  $C$  depending on  $n$ ,  $\omega_0$ ,  $\varphi_0$ ,  $p$ ,  $K$ , and  $\text{Ent}_p(F)$  such that*

$$\|\tilde{\varphi}\|_{L^\infty(M \times [0, T])} \leq C,$$

where  $\tilde{\varphi}$  is a normalization of a  $C^2$  solution of the flow (1-5).

Since  $\Theta < 0$  in the inverse Monge–Ampère equation, we have  $K \equiv 0$ , which means there is no extra condition for this case.

Going forward, a constant is called *universal* if it depends only on  $n$ ,  $\omega_0$ ,  $\varphi_0$ ,  $p$ ,  $\gamma$ ,  $r$ ,  $K$ , and  $\text{Ent}_p(F)$ .

## 2. Auxiliary equations

In this section, we want to find suitable auxiliary equations as in [Guo et al. 2023] and [Chen and Cheng 2023]. To motivate what a good auxiliary equation is, we first consider the parabolic Monge–Ampère flow (1-5). To have some monotonicity properties of the auxiliary solutions, we prefer a flow with negative time derivative of  $\psi$ . Thus the inverse Monge–Ampère flow, see (1-4), is a good candidate.

Let us consider the inverse Monge–Ampère flow

$$\begin{cases} (-\dot{\psi}_s)\omega_{\psi_s}^n = f_s e^{nF}\omega_0^n, \\ \psi_s(\cdot, 0) = 0, \end{cases} \tag{2-1}$$

where

$$f_s = \frac{(-\varphi - s)_+}{A_s}, \quad A_s = \int_{\Omega_s} (-\varphi - s)e^{nF}\omega_0^n dt, \quad \Omega_s = \{(z, t) \mid -\varphi(z, t) - s > 0\}.$$

But such a flow has singularities, since the factor  $(-\varphi - s)_+/A_s$  is not smooth in a neighborhood of  $\partial\Omega_s$ . Thus we need to consider a sequence of smooth functions  $\tau_k(x)$  which converges uniformly to  $x \cdot \chi_{\mathbb{R}^+}(x)$  and replace  $f_s$  on the right-hand side by

$$\frac{\tau_k(-\varphi - s)}{\int_{\Omega} \tau_k(-\varphi - s)e^{nF}}.$$

By the dominated convergence theorem,  $\psi_{s,k}$  converges to  $\psi_s$  uniformly, which means we can always take a limit in the inequalities to get the desired estimates as in [Guo et al. 2023] and [Chen and Cheng 2023]. To simplify our computations, we will keep using (2-1) as our auxiliary equation.

Another crucial modification of our auxiliary flow is that we must restrict the integration over the time slices. To express our idea more clearly, we need Lemma 2.1 as well as Corollary 2.2 in [Chen and Cheng 2023], which will be stated below. For the reader’s convenience, we will also include the proof from [Chen and Cheng 2023] here.

**Lemma 2.1.** *Consider the inverse Monge–Ampère flow*

$$\begin{cases} (-\dot{\varphi})\omega_{\varphi}^n = e^{nF}\omega_0^n, \\ \varphi|_{t=0} = \varphi_0. \end{cases}$$

Assuming

$$\int_{M \times [0, T]} e^{nF}\omega_0^n dt = C_1 < \infty,$$

we have  $|\sup_M \varphi| \leq C$ , where  $C$  depends on  $n, \omega_0, C_1$ , and  $\|\varphi_0\|_{L^\infty}$ .

*Proof.* Since  $\dot{\varphi} < 0$ , we can get the upper bound by

$$\sup_M \varphi \leq \sup_M \varphi_0 \leq \|\varphi_0\|_{L^\infty}.$$

To estimate the lower bound of  $\sup_M \varphi$ , let us consider the  $I$ -functional

$$I(\varphi) = \frac{1}{n+1} \int_M \varphi \sum_{j=0}^n \omega_0^{n-j} \wedge \omega_{\varphi}^j$$

and its derivative

$$\frac{d}{dt} I(\varphi) = \int_M \partial_t \varphi \omega_{\varphi}^n = - \int_M e^{nF} \omega_0^n.$$

Therefore, for any  $t' \in [0, T]$ , we have

$$\begin{aligned} I(\varphi) - I(\varphi_0) &= \int_0^{t'} \frac{d}{dt} I(\varphi) dt = - \int_{M \times [0, t']} e^{nF} \omega_0^n dt \\ &\geq - \int_{M \times [0, T]} e^{nF} \omega_0^n dt = -C_1, \end{aligned}$$

which implies  $I(\varphi)$  is bounded from below on  $[0, T]$ .

The lower bound estimate of  $\int_M \varphi \omega_0^n$  comes from integration by parts:

$$\begin{aligned} \int_M \varphi \omega_0^n - I(\varphi) &= \int_M \varphi \frac{1}{n+1} \sum_{j=0}^n \omega_0^{n-j} \wedge (\omega_0^j - \omega_\varphi^j) \\ &= \frac{1}{n+1} \int_M \varphi \sum_{j=0}^n \omega_0^{n-j} \wedge \sqrt{-1} \partial \bar{\partial}(-\varphi) \sum_{l=0}^{j-1} \omega_0^{j-1-l} \wedge \omega_\varphi^l \\ &= \frac{1}{n+1} \int_M \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \sum_{j=0}^n \sum_{l=0}^{j-1} \omega_0^{n-1-l} \wedge \omega_\varphi^l \geq 0. \end{aligned}$$

Therefore, we have  $\int_M \varphi \omega_0^n \geq -C_1$  and

$$\sup_M \varphi \geq \frac{1}{\text{Vol}(M, \omega_0)} \int_M \varphi \omega_0^n \geq -\frac{C_1}{\text{Vol}(M, \omega_0)}. \quad \square$$

**Corollary 2.2.** *There exists a constant  $\alpha > 0$  depending only on  $\omega_0$  such that*

$$\sup_{t \in [0, T]} \int_M e^{-\alpha \varphi} \omega_0^n \leq C_2,$$

where  $C_2$  depends on  $M$ ,  $\omega_0$ ,  $C_1$ , and  $\|\varphi_0\|_{L^\infty}$ .

This is a flow version of Hörmander’s result; see [Hörmander 1973, Lemma 4.4] and [Tian 1987] for local and global version of such integral estimate, respectively.

*Proof.* Since  $\varphi$  is a  $\omega_0$ -psh function for every  $t \in [0, T]$ , we have

$$\sup_{t \in [0, T]} \int_M e^{-\alpha(\varphi - \sup_M \varphi)} \omega_0^n \leq C.$$

From Lemma 2.1, the uniform bound of  $\sup_M \varphi$  gives us the desired inequality. □

The above Corollary 2.2 gives us a uniform bound on each time slice. If we apply this corollary on the space time  $M \times [0, T)$ , then a factor  $T$  seems unavoidable on the right-hand side. Therefore it is better to divide the whole space-time into several pieces  $M \times [t_0, t_0 + 1]$  and try to seek an estimate independent of  $t_0$ . This idea inspires us to consider such auxiliary equations involving only local information.

To get the  $L^\infty$  estimate, we need also to consider the normalization in Theorem 1.1, which follows the same normalization in [Picard and Zhang 2020]. In conclusion, we need to use the domain

$$\tilde{\Omega}_s = \{(z, t) \mid -\tilde{\varphi}(z, t) - s > 0\}$$

as a substitute for  $\Omega_s$ .

For any  $t_0 \in [0, T - 1)$ , let us consider a family of regions  $\Omega_{s, t_0} = \tilde{\Omega}_s \cap (M \times [t_0, t_0 + 1])$  and define a family of auxiliary equations

$$\begin{cases} (-\dot{\psi}_{s, t_0}) \omega_{\psi_{s, t_0}}^n = f_{s, t_0} e^{nF} \omega_0^n, \\ \psi_{s, t_0}(\cdot, 0) = 0, \end{cases} \tag{2-2}$$

where  $A_{s, t_0} = \int_{\Omega_{s, t_0}} (-\tilde{\varphi} - s) e^{nF} \omega_0^n dt$  and  $f_{s, t_0} = (-\tilde{\varphi} - s) \cdot \chi_{\Omega_{s, t_0}} / A_{s, t_0}$ .

The benefits appear when we apply Corollary 2.2 to such auxiliary equations. As a result of the choice of  $f_{s,t_0}$ , the integral on the right-hand side of the equation is 1. This implies

$$\int_{\Omega_{s,t_0}} e^{-\alpha\psi_{s,t_0}} \omega_0^n dt \leq C_2, \tag{2-3}$$

where the  $C_2$  is universal now. The above inequality is crucial, because it integrates against  $t$  while it remains independent of  $T$  and  $t_0$ . We will use inequality (2-3) frequently in the following sections.

The family of auxiliary equations (2-2) meets the same problem as (2-1). To be precise, we also need to apply  $\tau_k$  to remove the singularities. For the same reason, we will keep using (2-2) in the following sections.

The extra integral condition was chosen to make the normalization  $\tilde{\varphi}$  satisfy three properties, as follows.

**Lemma 2.3.** *Let  $\tilde{\varphi}$  be given in Theorem 1.1. Then we have*

- (1)  $\sup_{t \in [0, T]} \int_M \dot{\varphi} \omega_0^n \leq C_3,$
- (2)  $\tilde{\varphi} \leq C_3,$
- (3)  $\int_M |\tilde{\varphi}| \omega_0^n \leq C_3,$

where  $C_3$  is universal.

*Proof.* For (1), it comes directly from the estimates

$$\begin{aligned} \int_M \dot{\varphi} \omega_0^n &= \int_M \log\left(\frac{\omega_\varphi^n}{e^{nF} \omega_0^n}\right) \omega_0^n = \int \log\left(\frac{\omega_\varphi^n}{\omega_0^n}\right) \omega_0^n - \int \log(e^{nF}) \omega_0^n \\ &\leq \log\left(\int \omega_\varphi^n\right) - \int \log(e^{nF}) \omega_0^n = - \int \log(e^{nF}) \omega_0^n \leq K. \end{aligned}$$

The first estimate comes from Jensen’s inequality while the second one comes from the assumption on  $F$ . The average integral is chosen with respect to  $V = \int_M \omega_0^n$ .

To prove (2) and (3), let’s consider Green’s formula

$$\tilde{\varphi} = \int_M \tilde{\varphi} \omega_0^n - \int_M G \Delta \tilde{\varphi} \omega_0^n = - \int_M G \Delta \tilde{\varphi} \omega_0^n = - \int_M G \Delta \varphi \omega_0^n,$$

where  $G$  is the Green’s function with respect to  $\omega_0$ .

It is well known that the Green’s function  $G$  could be shifted to be nonnegative and with  $L^1$  norm bound  $C'$ . Combining with  $\text{tr}_{\omega_0} \omega_\varphi = n + \Delta\varphi > 0$  and Green’s formula, we have the universal estimate  $\tilde{\varphi} \leq nC'$ .

Let  $I_+$  and  $I_-$  be the integrals of the positive and negative parts of  $\tilde{\varphi}$ , respectively. Then we have

$$0 = \int_M \tilde{\varphi} = I_+ - I_- \quad \text{and} \quad I_+ \leq nC'V.$$

Thus

$$\int_M |\tilde{\varphi}| = I_+ + I_- = 2I_+ \leq 2nC'V.$$

The lemma follows from choosing  $C_3 = \max(KV, nC', 2nC'V)$ . □

### 3. Entropy bounded by energy

From Section 2, we get a good choice of a family of auxiliary equations (2-2). The following lemma is a key to the proof of Theorem 1.1,

**Lemma 3.1.** *Let  $\varphi$  be as in Theorem 1.1 and  $\psi_{s,t_0}$  be a solution of the auxiliary flow (2-2). Then there are constants  $\beta$ ,  $\epsilon$ , and  $\Lambda$ , with*

$$\beta = \frac{n+1}{n+2}, \quad \epsilon^{n+2} = \left(\frac{n+2}{n+1}\right)^{n+2} \Lambda, \quad \epsilon^{n+2} = \left(\frac{C_4}{(n+1)\beta}\right)^{n+1} A_{s,t_0},$$

where  $C_4$  is a universal constant defined below, such that

$$-\epsilon(-\psi_{s,t_0} + \Lambda)^\beta - \tilde{\varphi} - s \leq 0 \tag{3-1}$$

holds on  $M \times [t_0, t_0 + 1]$ .

In the following estimates we will use  $\psi$  and  $f$  to denote  $\psi_{s,t_0}$  and  $f_{s,t_0}$ , respectively. Let us consider the test function  $H = -\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - s$  and the linearization operator  $L = -\partial/\partial t + \Delta_{\omega_{\varphi_t}}$  of the Kähler-Ricci flow (1-1). The idea of the following argument comes from [Guo et al. 2023].

Let  $\Omega_{s,t_0}^\circ$  denote the interior of  $\Omega_{s,t_0}$ . Suppose the maximum of  $H$  is attained at some point  $x_0 = (z_0, t_0)$  outside  $\Omega_{s,t_0}^\circ$ ; then we have  $H \leq H(x_0) \leq -\tilde{\varphi}(x_0) - s \leq 0$ . To complete the proof, we only need to assume the maximal point  $x_0$  of  $H$  is in  $\Omega_{s,t_0}^\circ$  and then apply the maximum principle.

The scheme of the proof is to estimate  $LH$  at  $x_0$ . The constants are chosen to make  $0 < \beta < 1$  and  $1 - \beta\epsilon\Lambda^{\beta-1} = 0$ , which imply some cancellations. Moreover, such relations among the constants imply that  $LH$  is different from  $H$  by a positive coefficient at  $x_0$ . Roughly speaking, the lemma holds because of the facts that  $H$  is proportional to  $LH$  and  $LH \leq 0$ . Therefore let us firstly apply the operator  $L$  to  $H$  at  $x_0$ :

$$\begin{aligned} 0 \geq LH &= -\beta\epsilon(-\psi + \Lambda)^{\beta-1} \dot{\psi} + \dot{\varphi} - f \dot{\varphi} \omega_0^n \\ &\quad + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \Delta_{\omega_\varphi} \psi + \beta(1 - \beta)\epsilon(-\psi + \Lambda)^{\beta-2} |\partial\varphi|_{\omega_\varphi}^2 - \Delta_{\omega_\varphi} \varphi \\ &\geq \beta\epsilon(-\psi + \Lambda)^{\beta-1} (-\dot{\psi}) - (-\dot{\varphi}) + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \Delta_{\omega_\varphi} \psi - \Delta_{\omega_\varphi} \varphi - C_3 \\ &= \beta\epsilon(-\psi + \Lambda)^{\beta-1} (-\dot{\psi}) - (-\dot{\varphi}) + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \text{tr}_{\omega_\varphi} \omega_\psi \\ &\quad - \text{tr}_{\omega_\varphi} \omega_\varphi + (1 - \beta\epsilon(-\psi + \Lambda)^{\beta-1}) \text{tr}_{\omega_\varphi} \omega_0 - C_3 \\ &\geq \beta\epsilon(-\psi + \Lambda)^{\beta-1} (-\dot{\psi} + \text{tr}_{\omega_\varphi} \omega_\psi) - (-\dot{\varphi} + C_3 + n). \end{aligned}$$

The last estimate comes from the choice of auxiliary equations. Since  $\psi$  solves some inverse Monge-Ampère flows with initial data being identically 0, we have  $\psi \leq 0$  on  $M \times [0, T)$ , and moreover

$$1 - \beta\epsilon(-\psi + \Lambda)^{\beta-1} \geq 1 - \beta\epsilon\Lambda^{\beta-1} = 0.$$

Then we need to deal with the factor  $-\dot{\psi} + \text{tr}_{\omega_\varphi} \omega_\psi$ , which is the main term of the estimate. By the geometric-arithmetic inequality, we have

$$-\dot{\psi} + \text{tr}_{\omega_\varphi} \omega_\psi \geq -\dot{\psi} + n \left(\frac{\omega_\psi^n}{\omega_\varphi^n}\right)^{1/n}. \tag{3-2}$$

Combining (3-2) with the two flow equations (1-1) and (2-2) and using the geometric-arithmetic inequality again, we have

$$\begin{aligned}
 -\dot{\psi} + \text{tr}_{\omega_\psi} \omega_\psi &\geq -\dot{\psi} + n \left( \frac{\omega_\psi^n}{\omega_\phi^n} \right)^{1/n} \\
 &\geq -\dot{\psi} + n f^{1/n} \exp\left(-\frac{1}{n} \dot{\phi}\right) (-\dot{\psi})^{-1/n} \\
 &\geq (n+1) f^{1/(n+1)} \exp\left(-\frac{1}{n+1} \dot{\phi}\right).
 \end{aligned}
 \tag{3-3}$$

Thus replacing  $-\dot{\psi} + \text{tr}_{\omega_\psi} \omega_\psi$  by (3-3), we have the following estimate at  $x_0$ :

$$\begin{aligned}
 0 \geq LH &\geq (n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} \exp\left(-\frac{1}{n+1} \dot{\phi}\right) - (-\dot{\phi} + C_3 + n) \\
 &\geq \left[ (n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} - (-\dot{\phi} + C_3 + n) \exp\left(\frac{1}{n+1} \dot{\phi}\right) \right] \exp\left(-\frac{1}{n+1} \dot{\phi}\right).
 \end{aligned}$$

Since the exponential function is positive, we can simplify it by

$$0 \geq (n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} + (\dot{\phi} - C_3 - n) \exp\left(\frac{1}{n+1} \dot{\phi}\right),
 \tag{3-4}$$

which looks similar to the desired inequality (3-1) in Lemma 3.1. Let us consider a function

$$h(x) = (x - n - C_3) \exp\left(\frac{1}{n+1} x\right).$$

The function  $h : \mathbb{R} \rightarrow \mathbb{R}$  has a universal lower bound  $-C_4$ , where

$$C_4 = (n+1) \exp\left(\frac{C_3-1}{n+1}\right).$$

Thus we have  $h(\dot{\phi}) \geq -C_4$ , and moreover

$$(n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} - C_4 \leq 0.
 \tag{3-5}$$

Since  $f = (-\tilde{\varphi} - s)/A_{s,t_0}$  at the maximal point  $x_0$ , inequality (3-5) is equivalent to

$$\left( \frac{(n+1)\beta}{C_4} \right)^{n+1} \epsilon^{n+1} \frac{-\tilde{\varphi} - s}{A_{s,t_0}} \leq (-\psi + \Lambda)^{(n+1)(1-\beta)},
 \tag{3-6}$$

which is the test function when we chose the constants  $\beta$ ,  $\epsilon$ , and  $\Lambda$  as stated in Lemma 3.1. Thus we have

$$H \leq H(x_0) \leq 0,$$

which completes the proof.

From the above Lemma 3.1, we have

$$\frac{-\tilde{\varphi} - s}{A_{s,t_0}^{1/(n+2)}} \leq (c(-\psi + \Lambda))^{(n+1)/(n+2)},
 \tag{3-7}$$

where

$$c = \left( \frac{(n+1)^2 \epsilon}{(n+2)C_4} \right)^{n+2}.$$

The following estimate comes from (3-7):

$$\int_{\Omega_{s,t_0}} \exp\left[\lambda\left(\frac{-\tilde{\varphi} - s}{A_{s,t_0}^{1/(n+2)}}\right)^{(n+2)/(n+1)}\right] \omega_0^n dt \leq \int_{\Omega_{s,t_0}} \exp\{\lambda c(-\psi_{s,t_0} + \Lambda)\} \omega_0^n dt.$$

If the universal constant  $\lambda$  is chosen to make  $\lambda c = \alpha$ , then, by Corollary 2.2 and inequality (2-3), we have

$$\int_{\Omega_{s,t_0}} e^{-\lambda c \psi_{s,t_0}} \omega_0^n dt \leq \int_{M \times [t_0, t_0+1]} e^{-\alpha \psi_{s,t_0}} \omega_0^n dt \leq C_2,$$

where  $C_2$  is universal. In Lemma 3.1, The constant  $\Lambda$  is chosen to be proportional to  $A_{s,t_0}$ , i.e.,  $\Lambda = c' A_{s,t_0}$  for some universal constant  $c'$ . Then we can bound the right integral by  $C \exp(C A_{s,t_0})$  for some universal constant  $C$ .

Let us define  $E = \sup_{t_0 \in [0, T-1]} \int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3) e^{nF} \omega_0^n dt$ . Then we have

$$\begin{aligned} A_{s,t_0} &= \int_{\Omega_{s,t_0}} (-\tilde{\varphi} - s) e^{nF} \leq \int_{\Omega_{s,t_0}} (-\tilde{\varphi} + C_3) e^{nF} + \int_{\Omega_{s,t_0}} (-C_3 - s) e^{nF} \\ &\leq \int_{\Omega_{s,t_0}} (-\tilde{\varphi} + C_3) e^{nF} = \int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3) e^{nF} + \int_{M \times [t_0, t_0+1] \setminus \Omega_{s,t_0}} (\tilde{\varphi} - C_3) e^{nF} \\ &= E + \int_{M \times [t_0, t_0+1] \setminus \Omega_{s,t_0}} (\tilde{\varphi} - C_3) e^{nF} \leq E, \end{aligned}$$

where the last inequality comes from  $\tilde{\varphi} \leq C_3$  by Lemma 2.3. In summary, we have

$$\int_{\Omega_{s,t_0}} \exp\left[\lambda\left(\frac{-\tilde{\varphi} - s}{A_{s,t_0}^{1/(n+2)}}\right)^{(n+2)/(n+1)}\right] \omega_0^n dt \leq C \exp(CE). \tag{3-8}$$

The  $E$  defined above is called the energy. The  $C_3$  term inside the integral comes purely from a technical consideration that makes the inside function of the integral positive. There is no significant difference from the elliptic case where  $E = \int (-\tilde{\varphi})$  since the extra integral  $0 < \int e^{nF} \leq \text{Vol}(M, \omega_0) + \text{Ent}_p(F)$  is universally bounded from both sides.

To end this section, we will use the De Giorgi iteration method to derive the  $C^0$  estimate by assuming  $E$  is universally bounded. In the next section, we will apply the ABP estimate to get the universal bound on  $E$  which will complete the proof of Theorem 1.1. To prepare for the iteration procedure, we need such an inequality to run the iteration:

$$r\phi_{t_0}(s+r) \leq A_{s,t_0} \leq B_0\phi_{t_0}(s)^{1+\delta_0}, \tag{3-9}$$

where  $\phi_{t_0}(s) = \int_{\Omega_{s,t_0}} e^{nF} \omega_0^n dt$ .

The following lemma is the De Giorgi iteration mentioned above.

**Lemma 3.2.** *Let  $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a decreasing right continuous function with  $\lim_{s \rightarrow \infty} \Phi(s) = 0$ . Moreover, assume  $r\Phi(s+r) \leq B_0\Phi(s)^{1+\delta_0}$  for some constant  $B_0 > 0$  and all  $s > 0$  and  $r \in [0, r]$ . Then there exists a constant  $S_\infty = S_\infty(\delta_0, B_0, s_0) > 0$  such that  $\Phi(s) = 0$  for all  $s \geq S_\infty$ , where  $s_0$  is defined during the proof.*

*Proof.* Fix an  $s_0 > 0$  such that  $\Phi(s_0)^{\delta_0} < 1/(2B_0)$ . Such an  $s_0$  exists since  $\Phi(s) \rightarrow 0$  as  $s \rightarrow \infty$ . Define  $\{s_j\}$  by

$$s_{j+1} = \sup\{s > s_j \mid \phi(s) > \frac{1}{2}\phi(s_j)\}.$$

Thus

$$s_{j+1} - s_j \leq B_0\Phi(s_j)^{1+\delta} / \Phi(s_{j+1}) \leq 2B_0\Phi(s_j)^\delta \leq 2B_02^{-j\delta_0}\Phi^{\delta_0} \leq 2^{-j\delta_0}.$$

Letting

$$S_\infty = s_0 + \sum_{j \geq 0} (s_{j+1} - s_j) \leq s_0 + \frac{1}{1 - 2^{-\delta_0}},$$

we complete the proof. □

In our application,  $\Phi$  is chosen to be  $\phi_{t_0}$ . Let us derive the two sides of (3-9) separately. The left-hand side of (3-9) can be derived by definition which is similar to the proof in [Guo et al. 2023]. The following calculations are direct:

$$\begin{aligned} A_{s,t_0} &= \int_{\Omega_{s,t_0}} (-\tilde{\varphi} - s)e^{nF} \omega_0^n \, dt \geq \int_{\Omega_{s+r,t_0}} (-\tilde{\varphi} - s)e^{nF} \omega_0^n \, dt \\ &\geq \int_{\Omega_{s+r,t_0}} (s+r-s)e^{nF} \omega_0^n \, dt = r\phi(s+r). \end{aligned}$$

To get the right-hand side of (3-9), we need to apply the following inequality coming from Young’s inequality:

$$\int_{\Omega_{s,t_0}} v^p e^{nF} \omega_0^n \, dt \leq \|e^{nF}\|_{L^1(\log L)^p(M \times [t_0, t_0+1])} + C_p \int_{\Omega_{s,t_0}} e^{2v} \omega_0^n \, dt. \tag{3-10}$$

If we choose

$$v = \frac{\lambda}{2} \left( \frac{-\tilde{\varphi} - s}{A_{s,t_0}^{1/(n+2)}} \right)^{(n+2)/(n+1)},$$

then by the above inequality (3-10) we have

$$\int_{\Omega_{s,t_0}} (-\tilde{\varphi} - s)^{(n+2)p/(n+1)} e^{nF} \omega_0^n \, dt \leq C(E)A_{s,t_0}^{p/(n+1)}, \tag{3-11}$$

where the factor  $C(E)$  is a constant dependent on  $n, \omega_0, \varphi_0, p, \gamma, K, \text{Ent}_p(F)$  and additionally on  $E$ . The explicit dependence of the constant  $C(E)$  on  $E$  can be expressed by combining inequalities (3-10) and (3-8).

Thus the right-hand side can be derived by the estimate

$$\begin{aligned} A_{s,t_0} &= \int_{\Omega_{s,t_0}} (-\tilde{\varphi} - s)e^{nF} \omega_0^n \, dt \\ &\leq \left( \int_{\Omega_{s,t_0}} (-\tilde{\varphi} - s)^{(n+2)p/(n+1)} e^{nF} \right)^{(n+1)/((n+2)p)} \cdot \left( \int_{\Omega_{s,t_0}} e^{nF} \right)^{1/q} \\ &\leq C(E)^{(n+1)/((n+2)p)} A_{s,t_0}^{1/(n+2)} \phi_{t_0}^{1-(n+1)/(p(n+2))}, \end{aligned}$$

where the first line is by Hölder’s inequality and  $p$  is the Hölder coefficient

$$\frac{n+1}{p(n+2)} + \frac{1}{q} = 1.$$

In (3-9), we can choose

$$\delta_0 = 1 + \frac{p-n-1}{p(n+1)} \quad \text{and} \quad B_0 = C(E)^{1/p}.$$

To complete this section, we need to get an explicit expression on  $s_0$ . By Chebyshev’s inequality

$$\phi_{t_0}(s) \leq \frac{1}{s} \int_{\Omega_{s,t_0}} (-\tilde{\varphi}) e^{nF} \omega_0^n \leq \frac{E}{s},$$

we can choose  $s_0 = (2B_0)^{1/\delta_0} E$ .

In conclusion, we get the following theorem from the above arguments.

**Theorem 3.3.** *Let  $\tilde{\varphi}$  be as in Theorem 1.1. Then we have*

$$\sup_{M \times [0, T]} |\tilde{\varphi}| \leq C(n, \omega_0, \varphi_0, p, \gamma, K, \text{Ent}_p(F), E).$$

Moreover, if  $E$  can be controlled by a universal constant, then we have Theorem 1.1.

#### 4. Energy bounds by the ABP estimate

In this section we use a parabolic version of the ABP estimate proved by Krylov [1976] and Tso [1985] to give us a uniform energy bound. This approach was introduced in [Chen and Cheng 2023] and is an analogue to the elliptic version in [Guo et al. 2023].

Let  $u$  be a function defined on  $D = \Omega \times [0, T]$ , where  $\Omega$  is a bounded domain in  $\mathbb{R}^n$ . Then the parabolic ABP estimate says that

$$\sup_D u \leq \sup_{\partial_P D} u + C_n (\text{diam } \Omega)^{n/(n+1)} \left( \int_\Gamma |\partial_t u \det D_x^2 u| \, dx \, dt \right)^{1/(n+1)}, \tag{4-1}$$

where  $\partial_P D$  is the parabolic boundary of  $D$  and  $\Gamma = \{(x, t) \mid \partial_t u \geq 0, D_x^2 u \leq 0\}$ .

As mentioned in Section 2, we want to construct a family of local auxiliary equations. The auxiliary equations in this section are chosen to be

$$\begin{cases} (-\partial_t \psi_{t_0}) \omega_{\psi_{t_0}}^n = \frac{(|F|^p + 1) \cdot \chi_{M \times [t_0, t_0+1]}}{\int_{M \times [t_0, t_0+1]} (|F|^p + 1) e^{nF} \omega_0^n \, dt} e^{nF} \omega_0^n, \\ \psi_{t_0}(\cdot, 0) = 0. \end{cases} \tag{4-2}$$

We will use  $\psi$  to denote  $\psi_{t_0}$  for convenience and will skip the computation involved with  $\tau_k$  for the same reason we did in Section 2. Moreover, define a universal constant

$$\Psi = \int_{M \times [t_0, t_0+1]} (|F|^p + 1) e^{nF} \omega_0^n \, dt.$$

Parallel with Lemma 3.1, the following lemma plays a key role in this section.

**Lemma 4.1.** *Let  $\varphi$  be as in Theorem 1.1 and  $\psi$  be a solution of (4-2). For any  $0 < \beta < 1$ , there exists a constant  $C$  which depends on  $n, \omega_0, \varphi_0, p, \gamma, K, \text{Ent}_p(F)$ , and additionally on  $\beta$ , such that the following holds on  $M \times [t_0, t_0 + 1]$ :*

$$-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} \leq C, \tag{4-3}$$

where the constants  $\epsilon > 0$  and  $\Lambda > 0$  are defined as

$$\beta \epsilon \Lambda^{\beta-1} = \frac{1}{4}, \quad \Lambda = \left( \frac{n^p (2n+1)^p 2^p 4^{n+1} 9^{n+1} (n+1)^{n+1} C_4^{n+1} \Psi}{10^{n+1} \alpha^p} \right)^{1/((n+1)(1-\beta))}. \tag{4-4}$$

The constants  $\epsilon$  and  $\Lambda$  depend additionally on  $\beta$ .

Let  $\rho$  be the test function defined by

$$\rho = -\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi}$$

and  $L$  be the linearization as above. To prove Lemma 4.1, we only need to restrict  $\rho$  to its positive part. More precisely, consider  $h_s(x) = x + \sqrt{x^2 + s}$  and use  $h_s(\rho)$  to approximate  $2\rho_+$ . Therefore we have

$$2 \sup \rho \leq 2 \sup \rho_+ \leq \sup h_s(\rho),$$

and in addition the upper bounds of  $h_s$  imply an upper bound of  $\rho$ .

Let us consider  $h_s(\rho)^b$ , where

$$b = 1 + \frac{1}{(2n+2)(2n+1)},$$

and assume  $h_s(\rho)^b$  attains its maximal value  $Q$  at some point  $x_0 \in M \times [0, T]$ . Moreover, we can assume  $Q > 1$ , otherwise there is nothing to prove. Let us apply the parabolic ABP estimate for  $H = h_s(\rho)^b \cdot \eta$ , where  $\eta$  is a cut-off function defined below.

Assume  $r_0 = \min\{1, \text{inj}(M, \omega_0)\}$ , where  $\text{inj}(M, \omega_0)$  is the injectivity radius of  $(M, \omega_0)$ . The cut-off function  $\eta : M \rightarrow \mathbb{R}$  is defined in the following way:

$$\eta \equiv 1 \quad \text{on } B_{\omega_0}(x_0, \frac{1}{2}r_0), \tag{4-5}$$

$$\eta \equiv 1 - \theta \quad \text{on } \{M \setminus B_{\omega_0}(x_0, \frac{3}{4}r_0)\}, \tag{4-6}$$

$$1 - \theta \leq \eta \leq 1 \quad \text{on } \{B_{\omega_0}(x_0, \frac{3}{4}r_0) \setminus B_{\omega_0}(x_0, \frac{1}{2}r_0)\}, \tag{4-7}$$

$$|\nabla \eta|_{\omega_0}^2 \leq 10\theta^2/r_0^2, \tag{4-8}$$

$$|\nabla^2 \eta|_{\omega_0} \leq 10\theta/r_0^2, \tag{4-9}$$

where  $0 < \theta < 1$  is a small constant defined by

$$\theta = \min \left\{ \frac{r_0^2}{100Q^{1/b}}, \frac{1}{2(2n+1)(2n+2)} \right\} \leq \frac{1}{10}.$$

Since  $\eta$  is a space function, it has vanishing time derivative, which reduce our later computations.

*Proof of Lemma 4.1.* The following inequality can be derived directly by applying the operator  $L$  on  $H$ :

$$LH \geq bh'h^{b-1}(-\partial_t \rho)\eta + (\Delta_{\omega_\varphi} h^b)\eta + 2 \operatorname{Re}\langle \nabla h^b, \bar{\nabla} \eta \rangle_{\omega_\varphi} + h^b \Delta_{\omega_\varphi} \eta. \tag{4-10}$$

We will consider each of the terms separately to get good controls. Let us bound the last two terms:

$$h^b \Delta_{\omega_\varphi} \eta \geq -\frac{10\theta}{r_0^2} h^b \operatorname{tr}_{\omega_\varphi} \omega_0, \tag{4-11}$$

$$2 \operatorname{Re}\langle \nabla h^b, \bar{\nabla} \eta \rangle \geq -\frac{b(b-1)}{2} |\nabla h|_{\omega_\varphi}^2 h^{b-2} - \frac{2b}{b-1} h^b |\nabla \eta|_{\omega_\varphi}^2. \tag{4-12}$$

Then, we expand the second term and get

$$(\Delta_{\omega_\varphi} h^b)\eta = b(b-1)|\nabla h|_{\omega_\varphi}^2 h^{b-2}\eta + bh'h^{b-1}(\Delta_{\omega_\varphi} \rho)\eta + b|\nabla \rho|_{\omega_\varphi}^2 h''h^{b-1}\eta. \tag{4-13}$$

Combining (4-10)–(4-13), and noticing that the first term in (4-13) can be absorbed into the first term in (4-12) and the third term of (4-13) is positive, we have

$$\begin{aligned} LH &\geq bh'h^{b-1}(-\partial_t \rho)\eta + bh'h^{b-1}(\Delta_{\omega_\varphi} \rho)\eta - \frac{2b}{b-1} h^b |\nabla \eta|_{\omega_\varphi}^2 - \frac{10\theta}{r_0^2} h^b \operatorname{tr}_{\omega_\varphi} \omega_0 \\ &\geq bh'h^{b-1}(L\rho)\eta - \frac{2b}{b-1} \frac{10\theta^2}{r_0^2} h^b \operatorname{tr}_{\omega_\varphi} \omega_0 - \frac{10\theta}{r_0^2} h^b \operatorname{tr}_{\omega_\varphi} \omega_0. \end{aligned} \tag{4-14}$$

As we mentioned, the derivatives of the cut-off function  $\eta$  will produce  $\operatorname{tr}_{\omega_\varphi} \omega_0$  terms in (4-14) which will be absorbed in the later estimates. The  $L\rho$  term is the main term of (4-14), and it has the same structure as the main term of the test function appearing in Lemma 3.1. This fact motivates the following argument.

Let us compute  $L\rho$  and drop the positive term  $\beta(1-\beta)\epsilon(-\psi + \Lambda)^{\beta-2} |\nabla \psi|^2$ . Then we have

$$\begin{aligned} L\rho &\geq -\beta\epsilon\dot{\psi}(-\psi + \Lambda)^{\beta-1} + \dot{\varphi} + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \Delta_{\omega_\varphi} \psi + \beta(1-\beta)\epsilon(-\psi + \Lambda)^{\beta-2} |\nabla \psi|^2 - \Delta_{\omega_\varphi} \tilde{\varphi} \\ &\geq -\beta\epsilon\dot{\psi}(-\psi + \Lambda)^{\beta-1} + \dot{\varphi} + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \Delta_{\omega_\varphi} \psi - \Delta_{\omega_\varphi} \varphi - C_3. \end{aligned}$$

Since  $\Delta_{\omega_\varphi} \varphi + \operatorname{tr}_{\omega_\varphi} \omega_0 = n$  and  $\Delta_{\omega_\varphi} \psi + \operatorname{tr}_{\omega_\varphi} \omega_0 = \operatorname{tr}_{\omega_\varphi} \omega_\psi$ , we have

$$L\rho \geq \beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi} + \operatorname{tr}_{\omega_\varphi} \omega_\psi) + \dot{\varphi} + (1-\beta\epsilon\Lambda^{\beta-1}) \operatorname{tr}_{\omega_\varphi} \omega_0 - C_3 - n. \tag{4-15}$$

The  $\operatorname{tr}_{\omega_\varphi} \omega_0$  term in (4-15) will serve as a good term to absorb the last two terms in (4-14). The estimate for the rest of the terms in (4-15) follows the same idea in (3-2)–(3-4).

$$\begin{aligned} &\beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi} + \operatorname{tr}_{\omega_\varphi} \omega_\psi) + \dot{\varphi} - C_3 - n \\ &\geq (n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) + \dot{\varphi} - C_3 - n \\ &= \left[ (n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} + (\dot{\varphi} - C_3 - n) \exp\left(\frac{1}{n+1}\dot{\varphi}\right) \right] \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) \\ &\geq [(n+1)\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - C_4] \exp\left(-\frac{1}{n+1}\dot{\varphi}\right), \end{aligned} \tag{4-16}$$

where

$$\tilde{f} = \frac{(|F|^p + 1)\chi_{M \times [t_0, t_0+1]}}{\int_{M \times [t_0, t_0+1]} (|F|^p + 1)e^{nF} \omega_0^n dt}.$$

What we have now is the following inequality, noting that the cut-off function satisfies  $\frac{9}{10} \leq \eta \leq 1$  for any points in the space-time  $M \times [0, T]$ :

$$\begin{aligned} LH \geq (n+1)bh'h^{b-1} & \left[ \frac{9}{10}\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - \frac{C_4}{n+1} \right] \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) \\ & + bh^{b-1} \left[ \frac{9}{10}h'(1 - \beta\epsilon\Lambda^{\beta-1}) - \frac{20\theta^2}{(b-1)r_0^2}h - \frac{10\theta}{br_0^2}h \right] \text{tr}_{\omega_\varphi} \omega_0. \end{aligned} \quad (4-17)$$

Although the core of the lemma is to control  $\rho_+$ , the choice of  $h_s$  makes the negative part of  $\rho$  involved in the above inequality. So we will estimate on sets  $\Omega_+ = \{\rho > 0\}$  and  $\Omega_- = \{\rho \leq 0\}$  separately.

On  $\Omega_-$ , we have

$$0 \leq h_s(\rho) = \rho + \sqrt{\rho^2 + s} = \frac{s}{\sqrt{\rho^2 + s} - \rho} \leq \sqrt{s}$$

and

$$0 \leq h'_s(\rho) = 1 + \frac{\rho}{\sqrt{\rho^2 + s}} \leq 1.$$

Combining the two bounds on  $h$  and  $h'$  and inequality (4-17), we have

$$\begin{aligned} LH & \geq bs^{(b-1)/2} \left[ -C \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) - \frac{20\theta^2}{(b-1)r_0^2}h \text{tr}_{\omega_\varphi} \omega_0 - \frac{10\theta}{br_0^2}h \text{tr}_{\omega_\varphi} \omega_0 \right] \\ & = bs^{(b-1)/2} \left[ -C \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) - c(b, \theta, r_0)h \text{tr}_{\omega_\varphi} \omega_0 \right] \end{aligned}$$

on  $\Omega_-$ , where  $C$  is universal and

$$c(b, \theta, r_0) = \frac{20\theta^2}{(b-1)r_0^2} + \frac{10\theta}{br_0^2}.$$

On the other hand,  $1 \leq h' \leq 2$  on  $\Omega_+$ . By the choice of the constants  $\Lambda$  and  $\epsilon$  in (4-4), the coefficient of the  $\text{tr}_{\omega_\varphi} \omega_0$  term in (4-17) is positive.

Therefore, on the set  $\Omega_+$ , we have

$$LH \geq Cbh^{b-1} \left[ \beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - \frac{10C_4}{9(n+1)} \right] \exp\left(-\frac{1}{n+1}\dot{\varphi}\right).$$

Combining the above two estimates, we obtain

$$\begin{aligned} LH & \geq bs^{(b-1)/2} \left[ -C \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) - c(b, \theta, r_0)h \text{tr}_{\omega_\varphi} \omega_0 \right] \chi_{\Omega_-} \\ & + Cbh^{b-1} \left( \beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - \frac{10C_4}{9(n+1)} \right) \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) \chi_{\Omega_+} =: R. \end{aligned} \quad (4-18)$$

To apply the parabolic ABP estimate for the test function  $H$ , we define the domain

$$\tilde{\Gamma} = \{(z, t) \mid \partial_t H \geq 0, D_z^2 H \leq 0\}.$$

We need also discuss how to control the operator  $L$  on  $\tilde{\Gamma}$ . The estimate can be derived by

$$LH = -\frac{\partial}{\partial t}H + \Delta_{\omega_\varphi}H \leq -(2n+1) \left( \left| \frac{\partial}{\partial t}H \cdot \det D^2H \right| \cdot \left( \frac{\omega_0^n}{\omega_\varphi^n} \right)^2 \right)^{1/(2n+1)},$$

which connects our operator with the parabolic ABP estimate. In conclusion,

$$\left( \left| \frac{\partial}{\partial t}H \cdot \det D^2H \right| \cdot \left( \frac{\omega_0^n}{\omega_\varphi^n} \right)^2 \right)^{1/(2n+1)} \leq -\frac{R}{2n+1} \leq \frac{R_-}{2n+1}.$$

Note that the Hessian matrix  $D^2H$  is the real Hessian of  $H$  instead of the complex Hessian of  $H$ . Therefore we have

$$\left| \frac{\partial}{\partial t}H \cdot \det D^2H \right| \leq c_n R_-^{2n+1} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^2.$$

The factor  $\omega_\varphi^n/\omega_0^n$  is hard to control since it appears as a quadratic term. We can use different strategies to bound such term on  $\Omega_+$  and  $\Omega_-$ . On the domain  $\Omega_+$ , we have

$$\begin{aligned} h^{(2n+1)(b-1)} & \left( \beta \epsilon (-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - \frac{10C_4}{9(n+1)} \right)_-^{2n+1} \exp\left(-\frac{2n+1}{n+1}\dot{\varphi}\right) \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^2 \\ & \leq h^{(2n+1)(b-1)} \left( \frac{10C_4}{9(n+1)} \right)^{2n+1} e^{(n(2n+1)/(n+1))F} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^{2-(2n+1)/(n+1)} \\ & = \left( \frac{10C_4}{9(n+1)} \right)^{2n+1} h^{(2n+1)(b-1)} e^{(n(2n+1)/(n+1))F} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^{1/(n+1)}, \end{aligned} \tag{4-19}$$

while on  $\Omega_-$ , we have

$$\begin{aligned} s^{(2n+1)(b-1)/2} & \left( \exp\left(-\frac{1}{n+1}\dot{\varphi}\right) + c(b, \theta, r_0)h \operatorname{tr}_{\omega_\varphi} \omega_0 \right)^{2n+1} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^2 \\ & \leq C_5 s^{(2n+1)(b-1)/2} = C_5 s^{1/(4n+4)}, \end{aligned} \tag{4-20}$$

where  $C_5$  is not universal since it depends additionally on  $\varphi$ ,  $t$ , and  $t_0$ .

Defining  $D = B_r(x_0) \times [t_0, t_0 + 1]$  and combining (4-19) and (4-20), the parabolic estimate (4-1) tells us that

$$\begin{aligned} & \sup_D(H) - \sup_{\partial_p D}(H) \\ & \leq C_6 \left( \int_{D \cap \Omega_+} h^{(2n+1)(b-1)} e^{(n(2n+1)/(n+1))F} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^{1/(n+1)} \omega_0^n dt + \int_{D \cap \Omega_-} C_5 s^{1/(4n+4)} \omega_0^n dt \right)^{1/(2n+1)} \\ & \leq C_6 \left( \int_{D \cap \Omega_+} \frac{1}{n+1} \left( nh^{(2n+1)(n+1)(b-1)} e^{n(2n+1)F} + \frac{\omega_\varphi^n}{\omega_0^n} \right) \omega_0^n dt + C_5 s^{1/(4n+4)} \right)^{1/(2n+1)} \\ & \leq C_6 \left( \int_{D \cap \Omega_+} h^{1/2} e^{n(2n+1)F} + \int_{M \times [t_0, t_0+1]} \omega_\varphi^n dt + C_5 s^{1/(4n+4)} \right)^{1/(2n+1)} \\ & \leq C_6 \left( \int_{D \cap \Omega_+} h^{1/2} e^{n(2n+1)F} + 1 + C_5 s^{1/(4n+4)} \right)^{1/(2n+1)}, \end{aligned} \tag{4-21}$$

where  $C_6$  is universal. Moreover,  $C_6$  changes line by line as it absorbs all universal coefficients derived from the estimates. We use the volume of  $M$  to absorb the bad factor on  $\Omega_+$  and  $C_5$  to absorb the same bad factor on  $\Omega_-$ .

The integral over the set  $D \cap \Omega_+$  is in fact integrated over the set

$$\{\rho > 0\} \cap \left\{ \beta \epsilon (-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - \frac{10C_4}{9(n+1)} < 0 \right\}.$$

Over this set we have, from the choice of constants,

$$\begin{aligned} n(2n+1)|F| &\leq \left( \frac{10C_4}{9(n+1)} \right)^{(n+1)/p} \Psi^{1/p} (\beta \epsilon)^{-(n+1)/p} (-\psi + \Lambda)^{(1-\beta)(n+1)/p} \\ &= \frac{\alpha}{2} (-\psi + \Lambda)^{(1-\beta)(n+1)/p}. \end{aligned} \tag{4-22}$$

Moreover,  $h(\rho) \leq 2\rho + \sqrt{s}$ . Then by combining with the inequality (4-21), we have

$$\begin{aligned} \sup_D(H) - \sup_{\partial_p D}(H) &\leq C_6 \left( \int_{D \cap \Omega_+} (-\tilde{\varphi} + \sqrt{s})^{1/2} \exp\left(\frac{\alpha}{2}(-\psi + \Lambda)^{(1-\beta)(n+1)/p}\right) \omega_0^n dt + 1 + C_5 s^{1/(4n+4)} \right)^{1/(2n+1)} \\ &\leq C_6 \left( \int_{D \cap \Omega_+} (-\tilde{\varphi} + \exp(\alpha(-\psi + \Lambda)^{(1-\beta)(n+1)/p})) \omega_0^n dt + 1 + s^{1/2} + C_5 s^{1/(4n+4)} \right)^{1/(2n+1)} \\ &\leq C + C_5 s^{1/(4n+2)}, \end{aligned} \tag{4-23}$$

where  $C$  has the same dependencies as  $\Lambda$ . Moreover  $C_5$  changes line by line.

The last inequality is derived based on the following two inequalities: the integral  $\int_M (-\varphi) \omega_0^n dt$  is uniformly bounded by Lemma 2.3, and the fact that  $0 < (1-\beta)(n+1)/p < (n+1)/p < 1$  since  $0 < \beta < 1$  and  $p > n+1$ . Therefore the second integral is bounded by Corollary 2.2 and inequality (2-3).

By (4-23) and the definition of  $\theta$ , we have

$$cQ^{1-1/b} \leq \theta Q \leq \sup_D(H) - \sup_{\partial_p D}(H) \leq C + C_5 s^{1/(4n+2)},$$

where  $c$  is universal. In addition,  $\sup \rho$  can be controlled by

$$2 \sup \rho_+ \leq \sup h_s(\rho) \leq Q^{1/b}.$$

The proof of Lemma 4.1 follows from taking the limit  $s \rightarrow 0^+$ . □

Once we have Lemma 4.1, the following theorem is a direct application of Jensen’s inequality.

**Theorem 4.2.** *Let  $\varphi$  be the  $C^2$  solution defined in Theorem 1.1. For any  $\beta \in (0, 1)$  and  $t_0 \in [0, T - 1)$ , we have the energy estimate*

$$\int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3)^{1/\beta} e^{nF} \omega_0^n dt \leq C,$$

where the constant  $C$  has the same dependencies as  $\Lambda$  defined in Lemma 4.1.

*Proof.* Similar to the application of Lemma 3.1 and (3-7), we have

$$\int_{M \times [t_0, t_0+1]} \exp(c_\beta(-\tilde{\varphi} + C_3)^{1/\beta}) \omega_0^n dt \leq C_\beta,$$

where  $c_\beta$  and  $C_\beta$  both have the same dependencies as  $\Lambda$ .

Let  $\tilde{V} = \int_M e^{nF} \omega_0^n dt$  be the volume on the weighted volume form  $e^{nF} \omega_0^n dt$ . Taking logarithms of both sides and applying Jensen's inequality, we have

$$\begin{aligned} \tilde{V} \log\left(\frac{C_\beta}{\tilde{V}}\right) &\geq \tilde{V} \log\left(\frac{1}{\tilde{V}} \int_{M \times [t_0, t_0+1]} \exp(c_\beta(-\tilde{\varphi} + C_3)^{1/\beta} - nF) e^{nF} \omega_0^n dt\right) \\ &\geq \int_{M \times [t_0, t_0+1]} (c_\beta(-\tilde{\varphi} + C_3)^{1/\beta} - nF) e^{nF} \omega_0^n dt \\ &\geq c_\beta \int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3)^{1/\beta} e^{nF} \omega_0^n dt - \text{Ent}_p(F). \end{aligned}$$

The theorem follows from  $\tilde{V} \leq \text{Ent}_p(F) + V(M, \omega_0)$  and the fact  $y \log(y) > -1/e$  for  $y > 0$ . □

*Proof of Theorem 1.1.* Theorem 3.3 tells us the result follows directly from a uniform control on  $E$ . By Hölder's inequality, we have

$$\int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3) e^{nF} \omega_0^n dt \leq \tilde{V}^{1/n} \left( \int_{M \times [t_0, t_0+1]} (-\tilde{\varphi} + C_3)^{n/(n-1)} e^{nF} \omega_0^n dt \right)^{(n-1)/n}.$$

If we fix  $\beta = 1 - 1/n$  in Theorem 4.2, then the integral estimate is universal and independent of  $t_0$ . Thus we complete the proof of Theorem 1.1. □

### 5. Some generalizations

In this section, we derive some generalizations, Theorems 1.2 and 1.3, of Theorem 1.1.

The idea of Theorem 1.2 comes from the result of Chen and Cheng [2023] for general parabolic Hessian equations. Recall that  $r$  denotes the homogeneous degree of the operator  $\mathcal{F}$  and the linearization of the flow (1-3) is

$$Lu = -\partial_t u + G^{i\bar{j}} u_{i\bar{j}},$$

where

$$G^{i\bar{j}} = \frac{1}{\mathcal{F}} \frac{\partial \mathcal{F}(\lambda[h_\varphi])}{\partial h_{i\bar{j}}}.$$

To prove Theorem 1.2, we will use the family of auxiliary equations (2-2) and follow the same argument as in Sections 3 and 4.

The proof of main estimate (3-1) is tedious, and we will only show the essential differences compared to previous sections. When we apply the operator  $L$  to the test function

$$-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - s,$$

the Laplacian operator will be replaced by the trace operator  $\text{tr}_G v = G^{i\bar{j}} v_{i\bar{j}}$ . More precisely, we have the estimates

$$\begin{aligned} 0 &\geq L(-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - s) \\ &\geq -\beta\epsilon(-\psi + \Lambda)^{\beta-1} \dot{\psi} + \dot{\tilde{\varphi}} + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \text{tr}_G(\sqrt{-1}\partial\bar{\partial}\psi) \\ &\quad + \beta(1 - \beta)\epsilon(-\varphi + \Lambda)^{\beta-2} |\partial\varphi|_G^2 - \text{tr}_G(\sqrt{-1}\partial\bar{\partial}\varphi) \\ &\geq \beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi}) - (-\dot{\varphi}) + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \text{tr}_G(\sqrt{-1}\partial\bar{\partial}\psi) - \text{tr}_G(\sqrt{-1}\partial\bar{\partial}\varphi) - C_3 \\ &\geq \beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi}) - (-\dot{\varphi}) + \beta\epsilon(-\psi + \Lambda)^{\beta-1} \text{tr}_G \omega_\psi - \text{tr}_G \omega_\varphi - C_3 \\ &\geq \beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi} + \text{tr}_G \omega_\psi) - (-\dot{\varphi} + r + C_3). \end{aligned}$$

We also must deal with the factor  $-\dot{\psi} + \text{tr}_G \omega_\psi$  as in inequalities (3-2)–(3-4). The lower bound of the determinant on the condition of  $\mathcal{F}$  will give us the lower bound  $\det G^{i\bar{j}} \geq \gamma \mathcal{F}^{-n/r}$ . By the flows (2-2) and (1-3) and the homogeneous degree  $r$  condition, we have

$$\begin{aligned} -\dot{\psi} + \text{tr}_G \omega_\psi &\geq (n + 1) \sqrt[n+1]{\frac{f e^{nF} \omega_0^n}{\omega_\psi^n} \cdot \omega_\psi^n \det G^{i\bar{j}}} \\ &\geq C_7 f^{1/(n+1)} \sqrt[n+1]{\left(\frac{e^{rF}}{\mathcal{F}}\right)^{n/r}} \geq C_7 f^{1/(n+1)} \exp\left(-\frac{n}{r(n+1)}\dot{\varphi}\right), \end{aligned} \tag{5-1}$$

where  $C_7$  is a universal constant.

Since the function

$$h(x) = (x - r - C_3) \exp\left(\frac{n}{r(n+1)}x\right) > c,$$

where

$$c = -\frac{r(n+1)}{n} \exp\left(\frac{nC_3 - r}{r(n+1)}\right),$$

we have the same estimate

$$0 \geq C_7 \beta \epsilon (-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} - \frac{r(n+1)}{n} \exp\left(\frac{nC_3 - r}{r(n+1)}\right).$$

To derive the ABP estimate and (4-16) for this case, we need to calculate the estimate of the operator  $L$  on  $\tilde{\Gamma}$ . We have

$$LH \leq -(2n + 1) \left( |H_t \cdot \det D^2 H| \cdot \left(\frac{1}{(\det G)^2}\right) \right)^{1/(2n+1)}.$$

Moreover the bad factor in the integration over  $D \cap \Omega_+$  is

$$\begin{aligned} \exp\left(-\frac{n(2n+1)}{r(n+1)}\dot{\varphi}\right) \frac{1}{(\det G)^2} &= e^{(n^2(2n+1)/(r(n+1)))F} \mathcal{F}^{-n(2n+1)/(r(n+1))} \frac{1}{(\det G)^2} \\ &\leq \frac{1}{\gamma^2} e^{(n^2(2n+1)/(r(n+1)))F} \mathcal{F}^{n/(r(n+1))}. \end{aligned}$$

The exponent of  $\mathcal{F}$  is  $n/(r(n+1))$ , which is less than or equal to 1 if we assume the degree satisfies  $1 \leq r \leq n$ , and so we have a similar control as in (4-21). Theorem 1.2 follows from an analogue of Theorem 4.2.

We can also consider the much more general flow equation (1-5). Let us list the linearization operators for different choices of  $\Theta$  firstly:

$$Lu = \begin{cases} -\frac{\partial}{\partial t}u + \dot{\phi} \Delta_\phi u, & \Theta(x) = x, \\ -\frac{\partial}{\partial t}u + \frac{1}{3}\dot{\phi} \Delta_\phi u, & \Theta(x) = x^{1/3}, \\ -\frac{\partial}{\partial t}u - \dot{\phi} \Delta_\phi u, & \Theta(x) = -1/x. \end{cases} \tag{5-2}$$

To get Lemmas 3.1 and 4.1 under the new setting, we need to reprove Lemma 2.3 to get the upper bound of the integral. The following arguments are divided into two cases.

When  $\Theta(y) = -1/y$ , we have

$$\int_M \dot{\phi} = \int_M -e^{nF} \frac{\omega_0^n}{\omega_\phi^n} < 0.$$

When  $\Theta(y) = y^a$  for  $a > 0$ , we have

$$\ddot{\phi} = \frac{d}{dt} \left( \frac{\omega_\phi^n}{e^{nF} \omega_0^n} \right)^a = a \Delta_\phi \dot{\phi} \frac{\omega_\phi^n}{e^{nF} \omega_0^n} \left( \frac{\omega_\phi^n}{e^{nF} \omega_0^n} \right)^{a-1} = a \dot{\phi} \Delta_\phi \dot{\phi}.$$

Consider the first variation of the functional  $\int_M \dot{\phi} \omega_\phi^n$ , given by

$$\begin{aligned} \frac{d}{dt} \int_M \dot{\phi} \omega_\phi^n &= \int_M \ddot{\phi} \omega_\phi^n + \int_M \dot{\phi} \frac{d}{dt} (\omega_\phi^n) \\ &= \int_M \ddot{\phi} \omega_\phi^n + \int_M \dot{\phi} \Delta_\phi \dot{\phi} \omega_\phi^n \\ &= (a+1) \int_M \dot{\phi} \Delta_\phi \dot{\phi} \omega_\phi^n \\ &= -(a+1) \int_M |\nabla \dot{\phi}|_{\omega_\phi}^2 \omega_\phi^n \leq 0. \end{aligned}$$

Then the estimate of  $\int_M \dot{\phi} \omega_0^n$  follows from

$$\begin{aligned} \int_M \dot{\phi} \omega_0^n &\leq \int_M \dot{\phi} \omega_0^n - \int_M \dot{\phi} \omega_\phi^n + \int_M \dot{\phi}(\cdot, 0) \omega_{\phi_0}^n \\ &\leq \int_M \dot{\phi} (\omega_0^n - \omega_\phi^n) + \int_M e^{-anF} \left( \frac{\omega_{\phi_0}^n}{\omega_0^n} \right)^a \omega_{\phi_0}^n \\ &\leq \int_M \dot{\phi} (1 - e^{nF} \dot{\phi}^{1/a}) \omega_0^n + C \int_M e^{-anF} \omega_0^n, \end{aligned}$$

where  $C$  is universal.

Consider a function  $A(y) = y - ly^{1+1/a}$  defined on  $y \in [0, \infty)$ , where  $l$  is positive. Using calculus, we have

$$A(y) \leq A\left(\frac{a^a}{l^a(a+1)^a}\right) = \frac{a^a}{(a+1)^{a+1}} l^{-a}.$$

This yields the estimate

$$\int_M \dot{\phi} \omega_0^n \leq \frac{a^a}{(a+1)^{a+1}} \int_M e^{-anF} \omega_0^n + C \int_M e^{-anF} \omega_0^n \leq CK.$$

Once we have the above results, we need to prove Lemmas 3.1 and 4.1 for the flow equation (1-5). However the operator  $L$  has an extra factor  $-\dot{\varphi}$  in the Laplacian term which requires slightly different calculations. We will only discuss the case when  $\Theta = -1/y$ ; the other two cases can be treated similarly. To start with, we have

$$L(-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - s) \geq \beta\epsilon(-\psi + \Lambda)^{\beta-1}(-\dot{\psi} - \dot{\varphi} \operatorname{tr}_{\omega_\psi} \omega_\varphi) + \dot{\varphi} + n\dot{\varphi}. \tag{5-3}$$

There is no constant  $C_3$  since  $\int \dot{\varphi} < 0$ , and there is a term  $n\dot{\varphi}$  since we have the extra factor when we compute the second derivative. By applying the geometric-arithmetic inequality, we have

$$-\dot{\psi} - \dot{\varphi} \operatorname{tr}_{\omega_\psi} \omega_\varphi \geq (n + 1) \left( \frac{f e^{nF} \omega_0^n}{\omega_\psi^n} (-\dot{\varphi})^n \frac{\omega_\psi^n}{\omega_\varphi^n} \right)^{1/(n+1)} = (n + 1) f^{1/(n+1)} (-\dot{\varphi})$$

and

$$L(-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - s) \geq (n + 1)(\beta\epsilon(-\psi + \Lambda)^{\beta-1} f^{1/(n+1)} - 1)(-\dot{\varphi}).$$

Therefore we can drop the positive factor  $-\dot{\varphi}$  and evaluate the inequality at the maximal point.

To derive an analogue of Lemma 4.1, we need to consider

$$\rho = -\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi} - (n + 1)(t - t_0),$$

where the last two terms  $-(t - t_0)$  do not affect the result since we only estimate locally on  $M \times [t_0, t_0 + 1]$ . If the new  $\rho$  has an upper bound which has the same dependencies as the constants in Lemma 4.1, then  $-\epsilon(-\psi + \Lambda)^\beta - \tilde{\varphi}$  does as well. Following a similar calculation, we have

$$L\rho \geq (n + 1) \left( \beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - 1 + \frac{1}{-\dot{\varphi}} \right) (-\dot{\varphi}),$$

where  $1/-\dot{\varphi}$  comes from  $L(t - t_0) = -1$ . Applying the ABP estimate, we have

$$\left( |\partial_t H \cdot \det D^2 H| \cdot (-\dot{\varphi})^{2n} \left( \frac{\omega_0^n}{\omega_\varphi^n} \right)^2 \right)^{1/(2n+1)} \leq \frac{R_-}{2n + 1}. \tag{5-4}$$

In conclusion, the main term in the ABP estimate is

$$\begin{aligned} & \left( \beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - 1 + \frac{1}{-\dot{\varphi}} \right)_-^{2n+1} (-\dot{\varphi})^{2n+1-2n} \left( \frac{\omega_\varphi^n}{\omega_0^n} \right)^2 \\ & = \left( \beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - 1 + \frac{1}{-\dot{\varphi}} \right)_-^{2n+1} e^{2nF} \frac{1}{-\dot{\varphi}}. \end{aligned} \tag{5-5}$$

The term  $1/-\dot{\varphi}$  can be controlled pointwisely on the domain  $D \cap \Omega_+$ . More specifically,

$$\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - 1 + \frac{1}{-\dot{\varphi}} < 0$$

implies both inequalities

$$\beta\epsilon(-\psi + \Lambda)^{\beta-1} \tilde{f}^{1/(n+1)} - 1 < 0 \quad \text{and} \quad \frac{1}{-\dot{\varphi}} - 1 < 0.$$

The rest of the proof for the  $\Theta = -1/y$  case follows the same procedure.

### Acknowledgements

The author would like to thank Prof. Xiangwen Zhang for helpful discussions and frequent encouragement throughout the project. The author is also grateful for Prof. Duong H. Phong and Prof. Bin Guo’s interest and useful comments. Thanks also to Prof. Freid Tong for some private conversations that were enlightening.

### References

- [Cao 1985] H. D. Cao, “Deformation of Kähler metrics to Kähler–Einstein metrics on compact Kähler manifolds”, *Invent. Math.* **81**:2 (1985), 359–372. MR Zbl
- [Chen and Cheng 2021] X. Chen and J. Cheng, “On the constant scalar curvature Kähler metrics, I: A priori estimates”, *J. Amer. Math. Soc.* **34**:4 (2021), 909–936. MR Zbl
- [Chen and Cheng 2023] X. Chen and J. Cheng, “The  $L^\infty$  estimates for parabolic complex Monge–Ampère and Hessian equations”, *Pure Appl. Math. Q.* **19**:6 (2023), 2869–2913. MR Zbl
- [Eyssidieux et al. 2015] P. Eyssidieux, V. Guedj, and A. Zeriahi, “Weak solutions to degenerate complex Monge–Ampère flows, I”, *Math. Ann.* **362**:3-4 (2015), 931–963. MR Zbl
- [Eyssidieux et al. 2016] P. Eyssidieux, V. Guedj, and A. Zeriahi, “Weak solutions to degenerate complex Monge–Ampère flows, II”, *Adv. Math.* **293** (2016), 37–80. MR Zbl
- [Guedj et al. 2021] V. Guedj, C. H. Lu, and A. Zeriahi, “The pluripotential Cauchy–Dirichlet problem for complex Monge–Ampère flows”, *Ann. Sci. Éc. Norm. Supér. (4)* **54**:4 (2021), 889–944. MR Zbl
- [Guo and Phong 2023] B. Guo and D. H. Phong, “Auxiliary Monge–Ampère equations in geometric analysis”, *ICCM Not.* **11**:1 (2023), 98–135. MR Zbl
- [Guo and Phong 2024] B. Guo and D. H. Phong, “On  $L^\infty$  estimates for fully non-linear partial differential equations”, *Ann. of Math. (2)* **200**:1 (2024), 365–398. MR Zbl
- [Guo et al. 2023] B. Guo, D. H. Phong, and F. Tong, “On  $L^\infty$  estimates for complex Monge–Ampère equations”, *Ann. of Math. (2)* **198**:1 (2023), 393–418. MR Zbl
- [Harvey and Lawson 2023] F. R. Harvey and H. B. Lawson, Jr., “Determinant majorization and the work of Guo–Phong–Tong and Abja–Olive”, *Calc. Var. Partial Differential Equations* **62**:5 (2023), art. id. 153. MR Zbl
- [Hörmander 1973] L. Hörmander, *An introduction to complex analysis in several variables*, North-Holland Mathematical Library **7**, North-Holland, 1973. MR Zbl
- [Jian and Shi 2024] W. Jian and Y. Shi, “ $L^\infty$  estimates for Kähler–Ricci flow on Kähler–Einstein Fano manifolds: a new derivation”, *Proc. Amer. Math. Soc.* **152**:3 (2024), 1279–1286. MR Zbl
- [Kołodziej 2003] S. Kołodziej, “The Monge–Ampère equation on compact Kähler manifolds”, *Indiana Univ. Math. J.* **52**:3 (2003), 667–686. MR Zbl
- [Krylov 1976] N. V. Krylov, “Sequences of convex functions, and estimates of the maximum of the solution of a parabolic equation”, *Sibirsk. Mat. Ž.* **17**:2 (1976), 290–303, 478. In Russian; translated in *Sib. Math. J.* **17**:2 (1976), 226–236. MR Zbl
- [Phong et al. 2019] D. H. Phong, S. Picard, and X. Zhang, “A flow of conformally balanced metrics with Kähler fixed points”, *Math. Ann.* **374**:3-4 (2019), 2005–2040. MR Zbl
- [Picard and Suan 2024] S. Picard and C. Suan, “Flows of  $G_2$ -structures associated to Calabi–Yau manifolds”, *Math. Res. Lett.* **31**:6 (2024), 1837–1877. MR Zbl
- [Picard and Zhang 2020] S. Picard and X. Zhang, “Parabolic complex Monge–Ampère equations on compact Kähler manifolds”, pp. 639–665 in *Proceedings of the International Consortium of Chinese Mathematicians* (Taipei, 2018), edited by L. Ji and S.-T. Yau, International, Boston, MA, 2020. MR Zbl
- [Tian 1987] G. Tian, “On Kähler–Einstein metrics on certain Kähler manifolds with  $C_1(M) > 0$ ”, *Invent. Math.* **89**:2 (1987), 225–246. MR Zbl

- [Tso 1985] K. Tso, “On an Aleksandrov–Bakel’man type maximum principle for second-order parabolic equations”, *Comm. Partial Differential Equations* **10**:5 (1985), 543–553. MR Zbl
- [Wang et al. 2021] J. Wang, X.-J. Wang, and B. Zhou, “A priori estimate for the complex Monge–Ampère equation”, *Peking Math. J.* **4**:1 (2021), 143–157. MR Zbl
- [Yau 1978] S. T. Yau, “On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I”, *Comm. Pure Appl. Math.* **31**:3 (1978), 339–411. MR Zbl

Received 29 Jun 2023. Revised 12 Jul 2024. Accepted 20 Sep 2024.

QIZHI ZHAO: qizhiz3@uci.edu  
University of California, Irvine, Irvine, CA, United States

# Analysis & PDE

msp.org/apde

## EDITORS-IN-CHIEF

Anna L. Mazzucato Penn State University, USA  
alm24@psu.edu  
Clément Mouhot Cambridge University, UK  
c.mouhot@dpms.cam.ac.uk

## BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

## PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: "Linear Ramp"

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

The subscription price for 2025 is US \$475/year for the electronic version, and \$735/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 18 No. 8 2025

---

Uniform contractivity of the Fisher infinitesimal model with strongly convex selection	1835
VINCENT CALVEZ, DAVID POYATO and FILIPPO SANTAMBROGIO	
The $L^\infty$ estimate for parabolic complex Monge–Ampère equations	1875
QIZHI ZHAO	
Spectral asymptotics of the Neumann Laplacian with variable magnetic field on a smooth bounded domain in three dimensions	1897
MAHA AAFARANI, KHALED ABOU ALFA, FRÉDÉRIC HÉRAU and NICOLAS RAYMOND	
Characterization of weighted Hardy spaces on which all composition operators are bounded	1921
PASCAL LEFÈVRE, DANIEL LI, HERVÉ QUEFFÉLEC and LUIS RODRÍGUEZ-PIAZZA	
Long-time behavior of the Stokes-transport system in a channel	1955
ANNE-LAURE DALIBARD, JULIEN GUILLOD and ANTOINE LEBLOND	
Reconstruction for the Calderón problem with Lipschitz conductivities	2033
PEDRO CARO, MARÍA ÁNGELES GARCÍA-FERRERO and KEITH M. ROGERS	
Weakly turbulent solution to the Schrödinger equation on the two-dimensional torus with real potential decaying to zero at infinity	2061
AMBRE CHABERT	