

ANALYSIS & PDE

Volume 19

No. 1

2026

LARRY GUTH, DOMINIQUE MALDAGUE AND CHANGKEUN OH

ℓ^2 DECOUPLING THEOREM FOR SURFACES IN \mathbb{R}^3

ℓ^2 DECOUPLING THEOREM FOR SURFACES IN \mathbb{R}^3

LARRY GUTH, DOMINIQUE MALDAGUE AND CHANGKEUN OH

We identify a new way to divide the δ -neighborhood of surfaces $\mathcal{M} \subset \mathbb{R}^3$ into a finitely overlapping collection of rectangular boxes S . We obtain a sharp (ℓ^2, L^p) decoupling estimate using this decomposition for the sharp range of exponents $2 \leq p \leq 4$. Our decoupling inequality leads to new exponential sum estimates where the frequencies lie on surfaces which do not contain a line.

1. Introduction

Consider the manifold

$$\mathcal{M}_\phi := \{(\xi_1, \xi_2, \phi(\xi_1, \xi_2)) : \xi_1, \xi_2 \in [0, 1]\}, \quad (1-1)$$

where $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a smooth function. Denote by $N_\delta(\mathcal{M}_\phi) \subset \mathbb{R}^3$ the δ -neighborhood of the manifold \mathcal{M}_ϕ . Our main theorem is an ℓ^2 decoupling theorem for \mathcal{M}_ϕ into (ϕ, δ) -flat sets. This type of theorem was introduced by [Bourgain et al. 2020].

Definition 1.1. Let $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be smooth. We say that $S \subset [0, 1]^2$ is (ϕ, δ) -flat if

$$\sup_{u, v \in S} |\phi(u) - \phi(v) - \nabla\phi(u) \cdot (u - v)| \leq \delta. \quad (1-2)$$

For a measurable function $f : \mathbb{R}^3 \rightarrow \mathbb{C}$ and a measurable set $S \subset \mathbb{R}^2$, denote by f_S a smooth Fourier restriction of f on the set $S \times \mathbb{R}$. We refer to Definition 2.1 for the rigorous definition.

Theorem 1.2. Let $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function. Fix $\epsilon > 0$. Then there exists a sufficiently large number A depending on ϵ and ϕ satisfying the following.

For any $\delta > 0$, there exists a collection \mathcal{S}_δ of finitely overlapping sets S such that:

(1) The overlapping number is $O(\delta^{-\epsilon})$ in the sense that

$$\sum_{S \in \mathcal{S}_\delta} \chi_S \leq C_{\epsilon, \phi} \delta^{-\epsilon}. \quad (1-3)$$

(2) S is $(\phi, A\delta)$ -flat and of the form of a parallelogram.

(3) For $2 \leq p \leq 4$, we have

$$\|f\|_{L^p} \leq C_{\epsilon, \phi} \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^p}^2 \right)^{1/2} \quad (1-4)$$

for all functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$.

MSC2020: 42B25.

Keywords: decoupling inequality, exponential sum.

One important aspect of our theorem is that the collection \mathcal{S}_δ is not a strict partition of $[0, 1]^2$ since the sets $S \in \mathcal{S}_\delta$ are $O(\delta^{-\epsilon})$ -overlapping. The decoupling inequality (1-4) for the hyperbolic paraboloid is not true if a family \mathcal{S}_δ is a partition of $[0, 1]^2$. In other words, the $O(\delta^{-\epsilon})$ -overlapping property is necessary for the theorem to hold; see Appendix A. The situation is different for curves $(\xi, \phi(\xi))$ in \mathbb{R}^2 , where Yang [2021] proved a version of Theorem 1.2 using a strict partition. We realized that to prove an ℓ^2 decoupling inequality of the form (1-4), it is crucial to understand the number of possible choices of (ϕ, δ) -flat sets. For a curve $(\xi, \phi(\xi)) \in \mathbb{R}^2$ and a given point $p \in [0, 1]$, there is essentially one choice of a (ϕ, δ) -flat set containing the point p with the maximal size. On the other hand, for the hyperbolic paraboloid in \mathbb{R}^3 and a given point $p \in [0, 1]^2$, there are essentially $O(\log \delta^{-1})$ many choices of (ϕ, δ) -flat sets, which is the form of a rectangle with dyadic sidelengths, containing the point p and with maximal size. This turns out to be the reason why a partition is not enough and we need (1-3). For the hyperbolic paraboloid in \mathbb{R}^n with $n \geq 4$, the possible choices of (ϕ, δ) -flat sets containing a point is $\sim \delta^{-A_n}$ for some $A_n > 0$, which is why we do not have an analogous theorem without any δ^{-1} -power loss; see Appendix B.

Next we describe our main application of Theorem 1.2, which is to prove discrete restriction estimates for manifolds \mathcal{M}_ϕ not containing a line segment. For a function F and a set A , define

$$\|F\|_{L^\#_p(A)}^p := \frac{1}{|A|} \|F\|_{L^p(A)}^p. \tag{1-5}$$

Denote by Λ_δ a collection of δ -separated points in $[0, 1]^2$. Define $e(t) := e^{2\pi it}$.

Corollary 1.3. *Let ϕ be a polynomial of degree d with coefficients bounded by 1. Suppose that the manifold \mathcal{M}_ϕ does not contain a line segment. Then, for $2 \leq p \leq 4$, $\epsilon > 0$, and any sequence $\{a_\xi\}_{\xi \in \Lambda_\delta}$, we have*

$$\left\| \sum_{\xi \in \Lambda_\delta} a_\xi e(x \cdot (\xi, \phi(\xi))) \right\|_{L^\#_p(B_{\delta^{-d}})} \leq C_{\phi, \epsilon} \delta^{-\epsilon} \left(\sum_{\xi \in \Lambda_\delta} |a_\xi|^2 \right)^{1/2}. \tag{1-6}$$

An (ℓ^2, L^p) discrete restriction estimate is an inequality of the form (1-6). Corollary 1.3 is sharp in the following senses: First, the range of p of (1-6) is optimal for the paraboloid, though we expect that a wider range of p is possible for generic polynomials ϕ . Second, the inequality (1-6) is false for any $p > 2$ for some collection Λ_δ if the manifold \mathcal{M}_ϕ contains a line segment.

The (ℓ^2, L^p) discrete restriction estimate (1-6) implies the sharp (ℓ^p, L^p) discrete restriction estimate. Li and Yang [2021] obtained a sharp ℓ^p decoupling inequality for smooth manifolds, but their decoupling does not seem to imply (ℓ^p, L^p) discrete restriction estimates.

Our second application is about the Strichartz estimate for the nonelliptic Schrödinger equation on irrational tori. Let us introduce the conjecture. For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $\alpha < 0$, define $\tilde{\Delta} := \frac{1}{2\pi}(\partial_{11} + \alpha \partial_{22})$. Let v be a solution of the partial differential equation

$$i \partial_t v - \tilde{\Delta} v = 0, \quad v(0, x) = f(x), \tag{1-7}$$

where the initial data f is defined on the torus \mathbb{T}^2 . We let $e^{it\tilde{\Delta}} f(x) := v(x, t)$. The Strichartz estimate for the elliptic Schrödinger equation on irrational tori, which corresponds to the case $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $\alpha > 0$,

was studied in [Deng et al. 2017; 2022]. As in the case of the elliptic Schrödinger equation on irrational tori in [Deng et al. 2017, Conjecture 1.2], we make the following conjecture.

Conjecture 1.4. *Let α be an irrational number. Then*

$$\|e^{it\tilde{\Delta}} f\|_{L^p([0,T]\times\mathbb{T}^2)} \leq C_{\epsilon,\alpha} N^\epsilon \|f\|_{L^2(\mathbb{T}^2)} \begin{cases} T^{1/p} & \text{for } 2 \leq p \leq 4, \\ T^{1/p} + N^{1-4/p} & \text{for } 4 \leq p \leq 6, \\ T^{1/p} N^{1-6/p} + N^{1-4/p} & \text{for } 6 \leq p \end{cases} \quad (1-8)$$

for functions f whose Fourier support is in $[-N, N]^2$.

Corollary 1.5 gives partial progress on the conjecture for the range $2 \leq p \leq 4$.

Corollary 1.5. *Let α be an irrational number. Consider $2 \leq p \leq 4$. For $T \geq N$,*

$$\|e^{it\tilde{\Delta}} f\|_{L^p([0,T]\times\mathbb{T}^2)} \leq C_{\epsilon,\alpha} N^\epsilon T^{1/p} \|f\|_{L^2(\mathbb{T}^2)} \quad (1-9)$$

for functions f whose Fourier support is in $[-N, N]^2$.

A theorem analogous to Corollary 1.5 for the elliptic Schrödinger equation is obtained as a corollary of an ℓ^2 decoupling theorem [Bourgain and Demeter 2015] for the paraboloid, which is observed by [Deng et al. 2017]. Corollary 1.5 is not always true if α is not irrational. We refer to [Sun and Tzvetkov 2020; Mizutani and Tzvetkov 2015] for related topics.

We prove the corollaries in Section 4.

1.1. Comparison to decoupling inequalities in the literature. Let $\mathcal{M} \subset \mathbb{R}^3$ be a manifold in \mathbb{R}^3 with nonvanishing Gaussian curvature, and let $\mathcal{P}_\delta(\mathcal{M})$ be a partition of $N_\delta(\mathcal{M})$ into approximate $\delta^{1/2} \times \delta^{1/2} \times \delta$ caps θ . In their foundational decoupling paper, Bourgain and Demeter [2015] proved that if \mathcal{M} has everywhere positive Gaussian curvature and $2 \leq p \leq 4$, then

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M})} \|f_\theta\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2} \quad (1-10)$$

for all f with $\text{supp } \hat{f} \subset N_\delta(\mathcal{M})$. It is observed in [Bourgain and Demeter 2017a] that (1-10) is false for the hyperbolic paraboloid. The counterexample comes from the geometric fact that the hyperbolic paraboloid contains a line. Based on this observation, one might expect (1-10) to hold for manifolds avoiding a line segment.

Conjecture 1.6. *Let ϕ be a polynomial of degree d . Assume that \mathcal{M}_ϕ does not contain a line segment and has everywhere negative Gaussian curvature. Define*

$$\mathcal{P}_\delta := \{[a, a + \delta^{1/d}] \times [b, b + \delta^{1/d}] : a, b \in \delta^{1/d}\mathbb{Z} \cap [0, 1 - \delta^{1/d}]\}. \quad (1-11)$$

Then, for $2 \leq p \leq 4$,

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_{L^p(\mathbb{R}^3)}^2 \right)^{1/2} \quad (1-12)$$

for functions f whose Fourier supports are in $N_\delta(\mathcal{M})$.

Conjecture 1.6 implies **Corollary 1.3** in the case that the determinant of the Hessian matrix of ϕ is negative. Unfortunately, it is not clear how to prove **Conjecture 1.6** using the current decoupling techniques. Instead of proving **Conjecture 1.6**, we introduce a modified version of the decoupling inequality, which is inspired by (ϕ, δ) -flat sets, and prove that this decoupling still implies **Corollary 1.3**. Both **Conjecture 1.6** and **Theorem 1.2** imply **Corollary 1.3**, but neither of them seems to imply the other.

Let us review decoupling for smooth manifolds to motivate the use of (ϕ, δ) -sets. Bourgain and Demeter [2015; 2017a] proved that if \mathcal{M} has everywhere nonzero Gaussian curvature and $2 \leq p \leq 4$, then

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} (\delta^{-1})^{1/2-1/p} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M})} \|f_\theta\|_{L^p(\mathbb{R}^3)}^p \right)^{1/p} \tag{1-13}$$

for all f with $\text{supp } \hat{f} \subset N_\delta(\mathcal{M})$. As mentioned before, if the Gaussian curvature is positive, then the above decoupling inequality may be refined to an (ℓ^2, L^p) decoupling (see [Bourgain and Demeter 2015]), which has the form

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M})} \|f_\theta\|_{L^p(\mathbb{R}^3)}^2 \right)^{1/2}. \tag{1-14}$$

When \mathcal{M} is the truncated paraboloid $\mathbb{P}^2 = \{(\xi, |\xi|^2) \in \mathbb{R}^3 : |\xi| \leq 1\}$, we have that $\mathcal{P}_\delta(\mathbb{P}^2)$ is the coarsest partition of $N_\delta(\mathbb{P}^2)$ such that each piece θ is essentially flat. Bourgain and Demeter [2015] also proved sharp decoupling estimates for the cylinder $\text{Cyl} = \{(\xi_1, \xi_2, \xi_3) : \xi_1^2 + \xi_2^2 = 1, |\xi_3| \lesssim 1\}$ and the cone $\mathcal{C} = \{(\xi_1, \xi_2, \sqrt{\xi_1^2 + \xi_2^2}) : \frac{1}{4} \leq \xi_1^2 + \xi_2^2 \leq 4\}$, where their δ -neighborhoods are partitioned into planks θ of dimensions about $1 \times \delta^{1/2} \times \delta$. They observed that these are the coarsest (ϕ, δ) -flat sets. Based on this, Bourgain, Demeter, and Kemp [Bourgain et al. 2020] introduced the notion of (ϕ, δ) -flat sets (**Definition 1.1**) for a general manifold \mathcal{M}_ϕ and asked if there is a partition of $N_\delta(\mathcal{M}_\phi)$, denoted by $\mathcal{P}_\delta(\mathcal{M}_\phi)$, such that each element is (ϕ, δ) -flat and for $2 \leq p \leq 4$ the following ℓ^p decoupling is true:

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} |\mathcal{P}_\delta(\mathcal{M}_\phi)|^{1/2-1/p} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M}_\phi)} \|f_\theta\|_{L^p(\mathbb{R}^3)}^p \right)^{1/p} \tag{1-15}$$

for all functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$. This question is answered affirmatively in [Li and Yang 2021].

One may ask if there is a partition of $N_\delta(\mathcal{M}_\phi)$ such that each element is (ϕ, δ) -flat and for $2 \leq p \leq 4$ the following ℓ^2 decoupling is true:

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M}_\phi)} \|f_\theta\|_{L^p(\mathbb{R}^3)}^2 \right)^{1/2} \tag{1-16}$$

for all functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$. Note that (1-16) implies (1-15) by Hölder’s inequality. When S is the truncated hyperbolic paraboloid, Bourgain and Demeter proved that

$$\|f\|_{L^p(\mathbb{R}^3)} \lesssim_\epsilon \delta^{-\epsilon} \delta^{-1/2(1/2-1/p)} \left(\sum_{\theta \in \mathcal{P}_\delta(\mathcal{M}_\phi)} \|f_\theta\|_{L^p(\mathbb{R}^3)}^2 \right)^{1/2}. \tag{1-17}$$

They also proved that the loss of δ on the right-hand side is necessary up to the loss of $\delta^{-\epsilon}$, and the sharp example comes from the geometric fact that the hyperbolic paraboloid contains a line. At first, this might look like a counterexample to the question raised in (1-16). However, since there are multiple choices of (ϕ, δ) -flat sets, it does not give a counterexample. Also, given that the ℓ^2 decoupling theorem for a curve in \mathbb{R}^2 is proved by [Yang 2021] (which is the two-dimensional version of (1-16)), one might still expect that (1-16) is true for some collection of (ϕ, δ) -flat sets. In Appendix A, we prove Theorem A.1, which says that, for the hyperbolic paraboloid, there is no partition, whose elements are (ϕ, δ) -flat, such that (1-16) holds. The proof of Theorem A.1 crucially uses the assumption that a collection of (ϕ, δ) -flat sets is a partition. However, in all of the known applications of decouplings we are aware of, it does not matter if a collection of (ϕ, δ) -flat sets is a partition or $O(\delta^{-\epsilon})$ -finitely overlapping. Theorem 1.2 says the ℓ^2 decoupling theorem is true after allowing the collection \mathcal{S}_δ to be $O(\log \delta^{-1})$ -finitely overlapping; see (1-3). Our decoupling theorem is still useful in the sense that it implies Corollaries 1.3 and 1.5 and Proposition 1.10. Also, by Hölder’s inequality, our decoupling “essentially” implies the ℓ^p decoupling theorem (1-15) of Yang and Li.

Let us finish this subsection by mentioning higher-dimensional manifolds. It is conceivable that the ℓ^p decoupling conjecture is still true in high dimensions. Let us introduce some notation to state the conjecture. For a function $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, we say $S \subset [0, 1]^{n-1}$ is a (ϕ, δ) -set if (1-2) is true. Consider the manifold

$$\mathcal{M}_\phi := \{(\xi, \phi(\xi)) : \xi \in [0, 1]^{n-1}\}. \tag{1-18}$$

For a measurable function $f : \mathbb{R}^n \rightarrow \mathbb{C}$ and a measurable set $S \subset \mathbb{R}^{n-1}$, denote by f_S the Fourier restriction of f on the set $S \times \mathbb{R}$.

Conjecture 1.7. *Let $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ be a smooth function. Fix $\epsilon > 0$. Then there exists a sufficiently large number A depending on ϵ satisfying the following.*

For any $\delta > 0$, there exists a collection \mathcal{S}_δ of finitely overlapping sets $S \subset [0, 1]^{n-1}$ such that:

(1) *The overlapping number is $O(\log \delta^{-1})$ in the sense that*

$$\sum_{S \in \mathcal{S}_\delta} \chi_S \leq C_\epsilon \log(\delta^{-1}). \tag{1-19}$$

(2) *S is $(\phi, A\delta)$ -flat.*

(3) *For $2 \leq p \leq \frac{2(n+1)}{n-1}$, we have*

$$\|f\|_{L^p} \leq C_\epsilon \delta^{-\epsilon} (\#\mathcal{S}_\delta)^{1/2-1/p} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^p}^p \right)^{1/p} \tag{1-20}$$

for all functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$.

Conjecture 1.7 is known for the paraboloid and hyperbolic paraboloid but would be new for general manifolds. The ℓ^p decoupling inequality for manifolds in \mathbb{R}^2 and \mathbb{R}^3 are proved in [Yang 2021] and [Li and Yang 2021]. It remains open for higher dimensions. One main ingredient of ℓ^p decoupling for manifolds in \mathbb{R}^3 is ℓ^2 decoupling for a curve in \mathbb{R}^2 . Since we now have ℓ^2 decoupling for manifolds in \mathbb{R}^3 , it might be possible to prove ℓ^p decoupling for manifolds in \mathbb{R}^4 .

1.2. Example: hyperbolic paraboloid. Our theorem is even new for the hyperbolic paraboloid. Let us state this special case.

Theorem 1.8. Consider $\phi(\xi_1, \xi_2) = \xi_1 \xi_2$. For $2 \leq p \leq 4$,

$$\|f\|_{L^4} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{m=-2^{-1} \log \delta^{-1}}^{2^{-1} \log \delta^{-1}} \sum_{S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}} \|f_S\|_{L^4}^2 \right)^{1/2} \tag{1-21}$$

for all functions f whose Fourier support is in $N_\delta(\mathcal{M}_\phi)$.

The set $\mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}$ is defined in Section 2.1.

While the ℓ^p decoupling for the hyperbolic paraboloid does not imply the Tomas–Stein theorem for the manifold, our theorem implies the Tomas–Stein theorem for the manifold. As an application of Theorem 1.8 to exponential sum estimates, we will prove Corollary 1.5 in Section 4.

Theorem 1.8 itself does not imply ℓ^p decoupling by [Bourgain and Demeter 2017a]. However, in the proof of the theorem, we proved a slightly stronger inequality, which is a mixture of ℓ^2 and ℓ^p norms for all the intermediate scales. We do not state this as a theorem as it is very involved. But this inequality likely implies the ℓ^p decoupling by [Bourgain and Demeter 2017a].

We state the following version of the Stein–Tomas restriction theorem, which is recorded in [Demeter 2020, Theorem 1.16 and Proposition 1.27]. Let \mathbb{H} be the truncated hyperbolic paraboloid.

Theorem 1.9 [Tomas 1975; Stein 1986]. For $4 \leq p \leq \infty$, we have

$$\|\widehat{F}\|_{L^p(B_{\delta^{-1}})} \lesssim \delta^{1/2} \|F\|_{L^2(N_\delta(\mathbb{H}))}$$

for each $0 < \delta < 1$, each ball $B_{\delta^{-1}} \subset \mathbb{R}^3$, and each $F \in L^2(N_\delta(\mathbb{H}))$ supported in $N_\delta(\mathbb{H})$.

Proposition 1.10. Theorem 1.8 implies Theorem 1.9 up to a $\delta^{-\epsilon}$ loss.

Proof of Proposition 1.10. Theorem 1.9 with $p = \infty$ is trivial, so it suffices to prove the $p = 4$ case and invoke interpolation for the remaining exponents $p \in (4, \infty)$. Apply Theorem 1.8 with $p = 4$ to obtain the estimate

$$\|\widehat{F}\|_{L^4(B_{\delta^{-1}})} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{m=-2^{-1} \log \delta^{-1}}^{2^{-1} \log \delta^{-1}} \sum_{S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}} \|(\widehat{F})_S\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2}. \tag{1-22}$$

Note that $(\widehat{F})_S = \widehat{F} \chi_S$. Using Hölder’s inequality, we have, for each $S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}$, that

$$\|\widehat{F} \chi_S\|_{L^4(\mathbb{R}^3)} \leq \|\widehat{F} \chi_S\|_{L^\infty(\mathbb{R}^3)}^{1/2} \|\widehat{F} \chi_S\|_{L^2(\mathbb{R}^3)}^{1/2} \leq \|F\|_{L^1(N_\delta(S))}^{1/2} \|F\|_{L^2(N_\delta(S))}^{1/2} \leq |N_\delta(S)|^{1/4} \|F\|_{L^2(N_\delta(S))}.$$

Using the above inequality to bound the right-hand side of (1-22), we have

$$\begin{aligned} \|\widehat{F}\|_{L^4(B_{\delta^{-1}})} &\lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{m=-2^{-1} \log \delta^{-1}}^{2^{-1} \log \delta^{-1}} \sum_{S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}} |N_\delta(S)|^{1/2} \|F\|_{L^2(N_\delta(S))}^2 \right)^{1/2} \\ &\lesssim_\epsilon \delta^{-\epsilon} \max_{\substack{|m| \lesssim \log \delta^{-1} \\ S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}}} |N_\delta(S)|^{1/4} \|F\|_{L^2(N_\delta(\mathbb{H}))}. \end{aligned}$$

It remains to note that $|N_\delta(S)| \leq \delta^2$ for each $S \in \mathcal{R}_{2^m \delta^{1/2}, 2^{-m} \delta^{1/2}, 0}$ and each $|m| \lesssim \log \delta^{-1}$. □

1.3. Proof strategy and structure of the paper. In Section 2, we prove ℓ^2 decoupling inequalities for perturbed hyperbolic paraboloids using a collection of $O(\log \delta^{-1})$ -overlapping (ϕ, δ) -flat sets. Our result is new even for the hyperbolic paraboloid. The main new idea for the hyperbolic paraboloid is an iterative way of using broad-narrow analysis, which is combined with the induction-on-scale argument by [Bourgain and Demeter 2015; Bourgain et al. 2016]. To prove the theorem for perturbed hyperbolic paraboloids (Theorem 2.2), we combine the aforementioned argument with a restriction theory for perturbed hyperbolic paraboloids developed by [Buschenhenke et al. 2020; 2022; 2023; Guo and Oh 2024]. In Section 3, we use the ℓ^2 decoupling for the perturbed hyperbolic paraboloids as a black box and prove the ℓ^2 decoupling for general polynomials. This argument is essentially the same as that of [Li and Yang 2021]. In Section 4, we prove Corollaries 1.3 and 1.5 from the decoupling theorem. In Appendix A, we prove that there is no ℓ^2 decoupling for the hyperbolic paraboloid in \mathbb{R}^3 using a partition. This explains why it is necessary to introduce $\log(\delta^{-1})$ many partitions in Theorem 1.2. In Appendix B, we prove that there does not exist a collection of (ϕ, δ) -flat rectangles such that ℓ^2 decoupling for the collection is true in high dimensions without any additional power. In Appendix C, we prove the sharpness of Theorem 1.2.

1.4. Notation. For a function F and a set A , define

$$\|F\|_{L^\#_p(A)}^p := \frac{1}{|A|} \|F\|_{L^p(A)}^p. \quad (1-23)$$

For real numbers a_i , define

$$\left| \prod_{i=1}^3 a_i \right| := \prod_{i=1}^3 |a_i|^{1/3}. \quad (1-24)$$

For a ball B_K of radius K centered at $c(B_K)$, define

$$w_{B_K}(x) := \left(1 + \left| \frac{x - c(B_K)}{K} \right|^2 \right)^{-100}. \quad (1-25)$$

For a measurable set A and a function f , define

$$\int_A |f(x)| dx := \frac{1}{|A|} \int_A |f(x)| dx. \quad (1-26)$$

For a ball B , we define

$$\|f\|_{L^\#_p(w_B)}^p := |B|^{-1} \|f\|_{L^p(w_B)}^p. \quad (1-27)$$

For a rectangular box T , we denote by CT the dilation of T by a factor of C with respect to its centroid.

For two nonnegative numbers A_1 and A_2 , we write $A_1 \lesssim A_2$ to mean that there exists a constant C such that $A_1 \leq CA_2$. We write $A_1 \sim A_2$ if $A_1 \lesssim A_2$ and $A_2 \lesssim A_1$. We also write $A_1 \lesssim_\epsilon A_2$ if there exists C_ϵ depending on a parameter ϵ such that $A_1 \leq C_\epsilon A_2$.

2. Perturbed hyperbolic paraboloids

Consider the manifold \mathcal{M}_ϕ associated with

$$\phi(\xi_1, \xi_2) := \xi_1 \xi_2 + a_{2,0} \xi_1^2 + a_{0,2} \xi_2^2 + \sum_{3 \leq j+k \leq d} a_{j,k} \xi_1^j \xi_2^k, \quad (2-1)$$

where the coefficients $a_{j,k}$ satisfy

$$|a_{j,k}| \leq 10^{-10d} \tag{2-2}$$

for $d \geq 3$.

When $d = 2$, the manifold \mathcal{M}_ϕ is associated with

$$\phi(\xi_1, \xi_2) := \xi_1 \xi_2.$$

Note that this is a special case of (2-1) as we allow $a_{j,k}$ to be zero.

Definition 2.1. Let S be a rectangle in \mathbb{R}^2 . Take a smooth function $\Xi_S : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that:

- (1) The support of $\widehat{\Xi}_S$ is contained in $2S$.
- (2) $0 \leq \widehat{\Xi}_S(\xi_1, \xi_2) \leq 1$ for all $(\xi_1, \xi_2) \in \mathbb{R}^2$.
- (3) $\widehat{\Xi}_S$ is greater than $\frac{1}{10}$ on the set S .
- (4) $\|\Xi_S\|_{L^1} \leq 1000$.

Given a function $f : \mathbb{R}^3 \rightarrow \mathbb{C}$, define $f_S : \mathbb{R}^3 \rightarrow \mathbb{C}$ by

$$f_S(x_1, x_2, x_3) := \int_{\mathbb{R}^2} f(x_1 - y_1, x_2 - y_2, x_3) \Xi_S(y_1, y_2) dy_1 dy_2. \tag{2-3}$$

Note that this is a convolution of f and Ξ_S for the first two variables.

Theorem 2.2 (uniform ℓ^2 decoupling for perturbed hyperbolic paraboloids). *Fix $d \geq 2$ and $\epsilon > 0$. Then there exists a sufficiently large number A depending on d and ϵ satisfying the following.*

Let \mathcal{M}_ϕ be a manifold of the form (2-1) satisfying (2-2). Then, for any $\delta > 0$, there exists a family \mathcal{S}_δ of rectangles $S \subset \mathbb{R}^2$ such that:

- (1) *The overlapping number is $O(\log \delta^{-1})$ in the sense that*

$$\sum_{S \in \mathcal{S}_\delta} \chi_S \leq C_{d,\epsilon} \log(\delta^{-1}). \tag{2-4}$$

- (2) *Every S is $(\phi, A\delta)$ -flat and of the form of a parallelogram.*
- (3) *For all f whose Fourier support is in $N_\delta(\mathcal{M}_\phi)$,*

$$\|f\|_{L^4} \leq C_{d,\epsilon} \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2}. \tag{2-5}$$

The constant $C_{d,\epsilon}$ is independent of the choice of ϕ .

The first step of the proof is to construct the family \mathcal{S}_δ . A restriction theory for perturbed hyperbolic paraboloids is developed by [Buschenhenke et al. 2020; 2022; 2023; Guo and Oh 2024]. We use this theory, in particular, the language in [Buschenhenke et al. 2023] to construct the family.

2.1. Construction of the family \mathcal{S}_δ . To construct a family \mathcal{S}_δ , we will first define null vectors of the Hessian matrix of ϕ . Let us recall the notions in Section 3.1.1 of [Buschenhenke et al. 2023]. Consider two vectors at $\xi \in [0, 1]^2$ defined by

$$w_\xi := (-A(\xi), 1) \quad \text{and} \quad v_\xi := (1, -B(\xi)). \quad (2-6)$$

Here A and B are given by

$$A(\xi) = \frac{\phi_{22}(\xi)}{\phi_{12}(\xi) + \sqrt{|H_\phi(\xi)|}} \quad \text{and} \quad B(\xi) = \frac{\phi_{11}(\xi)}{\phi_{12}(\xi) + \sqrt{|H_\phi(\xi)|}}, \quad (2-7)$$

and $H_\phi(\xi)$ is the determinant of the Hessian matrix of ϕ . The functions $\phi_{ij}(\xi)$ are defined by $\partial_i \partial_j \phi(\xi)$. Note that by (2-2) we have

$$|A(\xi)| \leq 10^{-5d} \quad \text{and} \quad |B(\xi)| \leq 10^{-5d}. \quad (2-8)$$

By (3.7) of [Buschenhenke et al. 2023], the vectors w_ξ and v_ξ have the following properties:

$$(w_\xi)H_\phi(\xi)(w_\xi)^T = 0 \quad \text{and} \quad (v_\xi)H_\phi(\xi)(v_\xi)^T = 0. \quad (2-9)$$

Moreover, by (2-2), w_ξ and v_ξ are linearly independent.

We are now ready to define a family \mathcal{S}_δ . Assume that δ^{-1} is a dyadic number. Let us first define \mathcal{R}_δ ; the family \mathcal{S}_δ will be a subset of it:

$$\mathcal{R}_\delta := \bigcup_{\{1 \leq \alpha \leq \delta^{-1/2}: \alpha \in 2^{\mathbb{Z}}\}} \mathcal{R}_{\delta\alpha, \alpha^{-1}}. \quad (2-10)$$

We need to define $\mathcal{R}_{\delta\alpha, \alpha^{-1}}$. Every element in the set will be a rectangle with dimension $\delta\alpha \times \alpha^{-1}$. Introduce a parameter β which will be related to the angle of the long direction of the rectangle and the ξ_1 -axis. Define

$$\mathcal{R}_{\delta\alpha, \alpha^{-1}} := \bigcup_{\{\beta \in \mathbb{Z}: 0 \leq \beta \leq \delta^{-1}\alpha^{-2}\pi\}} \mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}. \quad (2-11)$$

Here $\mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$ is a collection of the translated copies of the rectangle of dimension $\delta\alpha \times \alpha^{-1}$ with the angle $\delta\alpha^2\beta$ with respect to the ξ_1 -axis, so that the elements of $\mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$ are disjoint and their union contains $[0, 1]^2$. So we have defined

$$\mathcal{R}_\delta = \bigcup_{\{1 \leq \alpha \leq \delta^{-1/2}: \alpha \in 2^{\mathbb{Z}}\}} \bigcup_{\{\beta \in \mathbb{Z}: 0 \leq \beta \leq \delta^{-1}\alpha^{-2}\pi\}} \mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}. \quad (2-12)$$

The family \mathcal{S}_δ in Theorem 2.2 will be a subcollection of \mathcal{R}_δ .

Definition 2.3. Let A be a sufficiently large constant, which will be the constant A in Theorem 2.2. We say two rectangles R_1 and R_2 are comparable if

$$R_1 \subset 2AR_2 \quad \text{and} \quad R_2 \subset 2AR_1. \quad (2-13)$$

Remark 2.4. Suppose that R_1 and R_2 are comparable. Let L be an affine transformation such that $L(R_1)$ and $L(R_2)$ are rectangles. Then $L(R_1)$ and $L(R_2)$ are also comparable.

Let us now explain how to choose rectangles. Let A be a large number. Let $R \in \mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$. If the two following conditions are satisfied, we add R to $\mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$:

- (1) R is $(\phi, A\delta)$ -flat.
- (2) For any point $z \in R$, we consider a rectangle centered at z , of dimension $\delta\alpha \times \alpha^{-1}$, with a long direction parallel to the vector w_z . This rectangle is comparable to R .

If the two following conditions are satisfied, we also add R to $\mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$:

- (1) R is $(\phi, A\delta)$ -flat.
- (2) For any point $z \in R$, we consider a rectangle centered at z , of dimension $\delta\alpha \times \alpha^{-1}$, with a long direction parallel to the vector v_z . This rectangle is comparable to R .

Recall that v_z and w_z are defined in (2-6). After this process, we finally have

$$\mathcal{S}_\delta := \bigcup_{\alpha} \bigcup_{\beta} \mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}. \tag{2-14}$$

We define a partition of unity associated with $\mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$. Namely, we take smooth functions $\{\Xi_R\}_{R \in \mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}}$ such that

- (1) the support of $\widehat{\Xi}_R$ is contained in $2R$,
- (2) $0 \leq \widehat{\Xi}_R(\xi_1, \xi_2) \leq 1$ for all $(\xi_1, \xi_2) \in \mathbb{R}^2$,
- (3) $\widehat{\Xi}_R$ is greater than $\frac{1}{10}$ on the set R ,
- (4) for any $\xi \in [0, 1]^2$, we have

$$\sum_{R \in \mathcal{R}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}} \widehat{\Xi}_R(\xi) = 1, \tag{2-15}$$

- (5) $\|\Xi_R\|_{L^1} \leq 1000$.

For a function f , define f_R by

$$f_R(x_1, x_2, x_3) := \int_{\mathbb{R}^2} f(x_1 - y_1, x_2 - y_2, x_3) \Xi_R(y_1, y_2) dy_1 dy_2. \tag{2-16}$$

2.2. Properties of partitions. We have defined partitions \mathcal{S}_δ ; see (2-14). Now let us study some properties of these partitions.

Proposition 2.5. *Let A be a sufficiently large constant. Then*

$$\mathcal{R}_{\delta^{1/2}, \delta^{1/2}, 0} \subset \mathcal{S}_\delta. \tag{2-17}$$

This says that our partitions contain the squares with the canonical scale. As a remark, for the hyperbolic paraboloid, note that the decoupling

$$\|f\|_{L^4} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{S \in \mathcal{R}_{\delta^{1/2}, \delta^{1/2}, 0}} \|f_S\|_{L^4}^2 \right)^{1/2} \tag{2-18}$$

is false for some functions f whose Fourier transforms are supported on the δ -neighborhood of the hyperbolic paraboloid.

Proof. Take $R \in \mathcal{R}_{\delta^{1/2}, \delta^{1/2}, 0}$. Since R is a square, by the definition of \mathcal{S}_δ , it suffices to prove that R is $(\phi, A\delta)$ -flat. By definition, we need to prove that

$$\sup_{u, v \in R} |\phi(u) - \phi(v) - \nabla\phi(u) \cdot (u - v)| \leq A\delta. \quad (2-19)$$

Since R is a square of side length $\delta^{1/2}$ and ϕ has the Hessian matrix whose determinant is bounded by 2 (see (2-1) and (2-2)), (2-19) follows from an application of Taylor's theorem. \square

Proposition 2.6 (finitely overlapping property). *For any $\xi \in [0, 1]^2$,*

$$\#\{S \in \mathcal{S}_\delta : \xi \in S\} \lesssim A \log(\delta^{-1}). \quad (2-20)$$

This proposition gives a proof of the first property of the family \mathcal{S}_δ in [Theorem 2.2](#) (recall that A is a constant depending only on d and ϵ). Let us give a proof.

Proof. Fix $\xi \in [0, 1]^2$. Since the number of dyadic numbers α with $1 \leq \alpha \leq \delta^{-1/2}$ is $O(\log(\delta^{-1}))$, it suffices to show that, given α ,

$$\#\{\beta \in \mathbb{Z} : \text{there exist } S \in \mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta} \text{ s.t. } \xi \in S\} \lesssim A. \quad (2-21)$$

Denote by $S_{1, \xi}$ (resp. $S_{2, \xi}$) the rectangle centered at z , of dimension $\delta\alpha \times \alpha^{-1}$, with a long direction parallel to the vector w_ξ (resp. v_ξ).

Suppose that $\xi \in S$ for some $S \in \mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$. Then S is comparable to either $S_{1, \xi}$ or $S_{2, \xi}$. Without loss of generality, we may assume that S is comparable to $S_{1, \xi}$. Then the angle between the long direction of S and w_ξ is $\lesssim A\delta\alpha^2$. Since $\{\delta\alpha^2\beta\}_{\beta \in \mathbb{Z}}$ is $\delta\alpha^2$ -separated, there are only $\sim A$ many β satisfying the property. This gives the proof. \square

What follows is a technical lemma, which says that the angle between the longest direction of a rectangle S and the ξ_1 -axis (or ξ_2 -axis) is small. If S is a square, it is not clear how to define ‘‘the longest direction of S ’’. So we prove such a statement under the condition (2-22) to guarantee that S looks like a rectangle quantitatively.

Lemma 2.7. *Let A be a sufficiently large number. Let $S \in \mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$. Suppose that*

$$\frac{\text{length of the long direction of } S}{\text{length of the short direction of } S} \geq A^2. \quad (2-22)$$

Then we have

$$\delta\alpha^2\beta \leq \frac{1}{10^{5d}}. \quad (2-23)$$

Proof. Suppose that $S \in \mathcal{S}_{\delta\alpha, \alpha^{-1}, \delta\alpha^2\beta}$. Let us follow the proof of [Proposition 2.6](#). Fix $\xi \in S$. Without loss of generality, we may assume that S is comparable to $S_{1, \xi}$. Then, by Euclidean geometry, the angle between the long direction of S and w_ξ is $\lesssim A\delta\alpha^2$. The condition (2-22) says that $\delta\alpha^2 \leq A^{-2}$, so the angle is bounded by $\lesssim A^{-1}$. On the other hand, by (2-6) and the normalization condition (2-2), the angle between w_ξ and the ξ_1 -axis is bounded by 10^{-9d} . Since A is sufficiently large, the angle between the long direction of S and the ξ_1 -axis is bounded by 10^{-5d} . Therefore, $\delta\alpha^2\beta$, which indicates the angle, is bounded by 10^{-5d} . \square

Definition 2.8. Define $D(\delta)$ to be the smallest constant such that

$$\|f\|_{L^4} \leq D(\delta) \left(\sum_{R \in \mathcal{S}_\delta} \|f_R\|_{L^4}^2 \right)^{1/2} \tag{2-24}$$

for all manifolds \mathcal{M}_ϕ of the form (2-1) satisfying (2-2) and functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$.

One of the key properties of the partitions is the parabolic rescaling lemma. This type of rescaling is very crucial in [Bourgain and Demeter 2015; Li and Yang 2021]. We observe that the same rescaling lemma still holds for our collections. The proof uses a basic property of perturbed hyperbolic paraboloids; for example, see Lemma 3.2 of [Guo and Oh 2024].

Lemma 2.9 (parabolic rescaling). *Let $\delta \leq \sigma \leq 1$. Let $R' \in \mathcal{S}_\sigma$. Then we have*

$$\|f_{R'}\|_{L^4} \leq CD(\sigma^{-1}\delta) \left(\sum_{R \in \mathcal{S}_\delta} \|f_R\|_{L^4}^2 \right)^{1/2} \tag{2-25}$$

for all manifolds \mathcal{M}_ϕ of the form (2-1) satisfying (2-2) and functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$.

Proof. Suppose that R' has dimension $\sigma\alpha \times \alpha^{-1}$ for some $1 \leq \alpha \leq \sigma^{-1}$. Then the Fourier support of $f_{R'}$ is contained in $2R' \times \mathbb{R}$. By using the triangle inequality, we may assume that the Fourier support of $f_{R'}$ is contained in $B \times \mathbb{R}$, where B is a box with dimension $\sigma\alpha \times \alpha^{-1}$. By translation and rotation, we may assume that the Fourier support of $f_{R'}$ is $[0, \alpha^{-1}] \times [0, \sigma\alpha]$. By abusing notation, let us still call this box R' . After the change of variables, our phase function changes. Let us denote the new phase function by

$$\phi_0(\xi_1, \xi_2) := \xi_1 \xi_2 + b_{2,0} \xi_1^2 + b_{0,2} \xi_2^2 + \sum_{3 \leq j+k \leq d} b_{j,k} \xi_1^j \xi_2^k. \tag{2-26}$$

For convenience, we introduce $b_{1,1} = 0$. Note that

$$|b_{j,k}| \lesssim 1, \quad 2 \leq j+k \leq d. \tag{2-27}$$

Since R' is $(\phi_0, A\sigma)$ -flat, by definition,

$$\sup_{u,v \in R'} |\phi_0(u) - \phi_0(v) - \nabla \phi_0(u) \cdot (u - v)| \leq A\sigma. \tag{2-28}$$

Take $u = (0, 0)$ and $v = (t, 0)$. Since $\nabla \phi_0(u) = 0$, this inequality becomes

$$\sup_{0 \leq t \leq \alpha^{-1}} \left| \sum_{j=2}^d b_{j,0} t^j \right| \leq A\sigma. \tag{2-29}$$

By [Li and Yang 2021, Proposition 7.1], this inequality gives

$$|b_{j,0}| \lesssim \sigma \alpha^j, \quad 2 \leq j \leq d. \tag{2-30}$$

Similarly, we take $u = (0, 0)$ and $v = (0, t)$, where $0 \leq t \leq \sigma\alpha$. Then, by the same reasoning, we obtain

$$|b_{0,j}| \lesssim \sigma (\sigma\alpha)^{-j}, \quad 2 \leq j \leq d. \tag{2-31}$$

We next do rescaling. Define $L(\xi, \eta) := (\alpha^{-1}\xi, \sigma\alpha\eta)$ and

$$\tilde{\phi}(\xi, \eta) := \sigma^{-1}\phi_0(L(\xi, \eta)). \quad (2-32)$$

The function $\tilde{\phi}(\xi, \eta)$ can be rewritten as

$$\xi_1\xi_2 + (\sigma^{-1}\alpha^{-2}b_{2,0})\xi_1^2 + (\sigma\alpha^2b_{0,2})\xi_2^2 + \sum_{3 \leq j+k \leq d} (\sigma^{k-1}\alpha^{k-j}b_{j,k})\xi_1^j\xi_2^k. \quad (2-33)$$

To apply the induction hypothesis, we will prove

$$|b_{j,k}| \lesssim \sigma^{-k+1}\alpha^{-k+j}, \quad 2 \leq j+k \leq d. \quad (2-34)$$

The cases for $j=0$ or $k=0$ follow from (2-30) and (2-31). Hence we may assume that $j \geq 1$ and $k \geq 1$.

Let us consider the subcase $k \leq j$. By the inequality $1 \leq \alpha \leq \sigma^{-1}$, we have $1 \leq \alpha^{-k+j}$. So what we need to prove follows from

$$|b_{j,k}| \lesssim \sigma^{-k+1}. \quad (2-35)$$

Since $k \geq 1$ and $\sigma \leq 1$, this follows from (2-27). Let us next consider the subcase $k \geq j$. By the inequality $1 \leq \alpha \leq \sigma^{-1}$, we have $\sigma^{k-j} \leq \alpha^{-k+j}$. So what we need to prove follows from

$$|b_{j,k}| \lesssim \sigma^{-j+1}. \quad (2-36)$$

Since $j \geq 1$ and $\sigma \leq 1$, this follows from (2-27).

We have now proved (2-34). So the function (2-33) can be rewritten as

$$\xi_1\xi_2 + c_{2,0}\xi_1^2 + c_{0,2}\xi_2^2 + \sum_{3 \leq j+k \leq d} c_{j,k}\xi_1^j\xi_2^k, \quad (2-37)$$

where

$$|c_{j,k}| \lesssim 1, \quad 2 \leq j+k \leq d. \quad (2-38)$$

By applying a linear transformation, we may assume that $c_{2,0} = c_{0,2} = 0$. Moreover, by using the triangle inequality, we may assume that $|c_{j,k}| \leq 10^{-10d}$. By abusing the notation, let us still denote by $\tilde{\phi}$ the new phase function. By (2-32) and a change of variables on physical variables, the δ -neighborhood of \mathcal{M}_ϕ becomes the $\sigma^{-1}\delta$ -neighborhood of $\mathcal{M}_{\tilde{\phi}}$. For functions g whose Fourier supports are in $N_{\sigma^{-1}\delta}(\mathcal{M}_{\tilde{\phi}})$, we have

$$\|g\|_{L^4} \leq D(\sigma^{-1}\delta) \left(\sum_{R'' \in \mathcal{S}_{\sigma^{-1}\delta}} \|g_{R''}\|_{L^4}^2 \right)^{1/2}. \quad (2-39)$$

We next rescale back to the original variables. Given $R'' \in \mathcal{S}_{\sigma^{-1}\delta}$, by definition, we have

$$\sup_{u,v \in R''} |\tilde{\phi}(u) - \tilde{\phi}(v) - \nabla\tilde{\phi}(u) \cdot (u-v)| \leq A\sigma^{-1}\delta. \quad (2-40)$$

By (2-32), we can see that $L^{-1}(R'')$ is $(\phi, A\delta)$ -flat. To conclude that $L^{-1}(R'') \in \mathcal{S}_\delta$, it remains to show the following property: For any point $z \in L^{-1}(R'')$, we consider a rectangle centered at z , of dimension $\sigma\alpha \times \alpha^{-1}$, with a long direction parallel to the vector v_z ; let us denote this by R_z . This rectangle is comparable to $L^{-1}(R'')$.

Let us prove the property. Fix $z \in L^{-1}(R'')$, and let v_z be a vector associated with the phase ϕ . By [Remark 2.4](#), it suffices to show that $L(R_z)$ is comparable to R'' . After calculating the Hessian matrix, we see that $L(v_z)$ is comparable to the vector $v_{L(z)}$ associated with the phase function $\tilde{\phi}$. Since R'' belongs to $\mathcal{S}_{\sigma^{-1}\delta}$, this gives the desired property. \square

2.3. The broad-narrow analysis. In this subsection, we introduce a multilinear decoupling and show that a linear decoupling constant is comparable to a multilinear decoupling constant up to epsilon loss ([Theorem 2.12](#)). This framework is introduced by [[Bourgain and Demeter 2015](#)].

Definition 2.10. Let $n(\xi)$ be the unit normal vector of \mathcal{M}_ϕ at the point $(\xi, \phi(\xi))$. We say three points $\xi_{(1)}, \xi_{(2)}, \xi_{(3)} \in [0, 1]^2$ are N^{-1} -transverse if

$$|n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi_{(3)})| \geq N^{-1}. \tag{2-41}$$

Three squares $\tau_1, \tau_2, \tau_3 \subset [0, 1]^2$ are called N^{-1} -transverse if every triple $\xi_{(i)} \in \tau_i$ is N^{-1} -transverse.

Definition 2.11. Define $D_{\text{mul}}(\delta, N^{-1})$ to be the smallest constant satisfying the following: for any N^{-1} -transverse squares $\tau_1, \tau_2, \tau_3 \in \mathcal{R}_{a,a,0}$ for $\delta^{1/2} \leq a \leq 1$,

$$\left\| \prod_{i=1}^3 f_{\tau_i} \right\|_{L^4} \leq D_{\text{mul}}(\delta, N^{-1}) \left(\sum_{R \in \mathcal{S}_\delta} \|f_R\|_{L^4}^2 \right)^{1/2} \tag{2-42}$$

for all manifolds \mathcal{M}_ϕ of the form (2-1) satisfying (2-2) and functions f whose Fourier supports are in $N_\delta(\mathcal{M}_\phi)$.

Recall that the definition of f_{τ_i} is given in the paragraph below (2-14). The main theorem of this subsection follows.

Theorem 2.12. *Given $\epsilon > 0$, there exists a sufficiently large number N depending on ϵ such that, for all $\delta > 0$, we have*

$$D(\delta) \leq C_\epsilon \delta^{-\epsilon} \left(\sup_{\delta \leq \delta' \leq 1} D_{\text{mul}}(\delta', N^{-1}) + 1 \right). \tag{2-43}$$

Proof. We do a broad-narrow analysis by [[Bourgain and Guth 2011](#)]. Let K be a sufficiently large number, which will be determined later. Take a sufficiently large constant K_1 such that K is sufficiently large compared to K_1 . Fix a ball B_K . We write

$$f = \sum_{\tau \in \mathcal{R}_{K^{-1/2}, K^{-1/2}, 0}} f_\tau. \tag{2-44}$$

Recall the definition in (2-11). Each τ is a square of side length $K^{-1/2}$. Consider a collection of squares

$$\mathcal{C} := \{ \tau \in \mathcal{R}_{K^{-1/2}, K^{-1/2}, 0} : \|f_\tau\|_{L^4(B_K)} \geq K^{-100} \|f\|_{L^4(B_K)} \}. \tag{2-45}$$

We will consider several cases. By a pigeonholing argument, we can see that \mathcal{C} is nonempty. Take $\tau_1 \in \mathcal{C}$. Consider the first case that all the squares $\tau \in \mathcal{C}$ are in the $(K_1)^{-1}$ -neighborhood of τ_1 . Denote

by τ'_1 the neighborhood. Then, by the triangle inequality, we have

$$\|f\|_{L^4(B_K)} \leq \left\| \sum_{\{\tau \in \mathcal{R}_{K^{-1/2}, K^{-1/2}, 0} : \tau \cap \tau'_1 \neq \emptyset\}} f_\tau \right\|_{L^4(B_K)} + \sum_{\tau \notin \mathcal{C}} \|f_\tau\|_{L^4(B_K)}. \quad (2-46)$$

By the definition of \mathcal{C} , this gives

$$\|f\|_{L^4(B_K)} \leq 2 \left\| \sum_{\{\tau \in \mathcal{R}_{K^{-1/2}, K^{-1/2}, 0} : \tau \cap \tau'_1 \neq \emptyset\}} f_\tau \right\|_{L^4(B_K)}. \quad (2-47)$$

The set τ'_1 depends on a choice of B_K . Hence, by summing over all possible squares, we have

$$\|f\|_{L^4(B_K)} \lesssim \left(\sum_{\tau' \in \mathcal{R}_{K_1^{-1}, K_1^{-1}, 0}} \|f_{\tau'}\|_{L^4(B_K)}^4 \right)^{1/4}. \quad (2-48)$$

We next consider the case that there exists $\tau_2 \in \mathcal{C}$ outside of the $(K_1)^{-1}$ -neighborhood of τ_1 . There are two subcases. Suppose that there is $\tau_3 \in \mathcal{C}$ such that τ_1, τ_2, τ_3 are $10K^{-1}$ -transverse. Then we have

$$\|f\|_{L^4(B_K)} \lesssim K^{100} \prod_i \|f_{\tau_i}\|_{L^4(B_K)}. \quad (2-49)$$

By an application of a randomization argument and the uncertainty principle, we have

$$\|f\|_{L^4(B_K)} \lesssim K^{200} \left\| \prod_i \tilde{f}_{\tau_i} \right\|_{L^4(w_{B_K})}, \quad (2-50)$$

where \tilde{f}_{τ_i} is a modulation of f_{τ_i} . The definition of w_{B_K} is given in (1-25). We refer to (2.5)–(2.8) of [Guth et al. 2023] for the details of the argument.

Lastly, suppose that there does not exist such a τ_3 . Denote by $\xi_{(i)}$ the center of τ_i for $i = 1, 2$. By Definition 2.10, all the elements $\tau \in \mathcal{C}$ are contained in the set

$$Z_0 := \{\xi \in [0, 1]^2 : |n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi)| < 100K^{-1}\}. \quad (2-51)$$

Here $n(\xi)$ is the unit normal vector of the manifold (2-1) at the point $(\xi, \phi(\xi))$. By the definition of \mathcal{C} , we can obtain

$$\|f\|_{L^4(B_K)} \lesssim \left\| \sum_{\tau \subset Z_0} f_\tau \right\|_{L^4(B_K)}. \quad (2-52)$$

We apply a uniform ℓ^2 decoupling theorem for the set Z_0 of [Li and Yang 2021, Theorem 3.1] and obtain

$$\|f\|_{L^4(B_K)} \lesssim_\epsilon K^\epsilon \left(\sum_S \|f_S\|_{L^4(w_{B_K})}^2 \right)^{1/2} \quad (2-53)$$

for any $\epsilon > 0$. Here $S \subset [0, 1]^2$ is a rectangular box, and the collection $\{S\}_S$ has bounded overlap. Moreover,

$$\bigcup_S S \subset \{\xi \in [0, 1]^2 : |n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi)| \lesssim K^{-1}\}. \quad (2-54)$$

The function f_S is the Fourier restriction of f to the rectangle $S \times \mathbb{R}$.¹ Define

$$Z := \{\xi \in [0, 1]^2 : |n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi)| = 0\}. \tag{2-55}$$

Then, by the separation between τ_1 and τ_2 and by the normalization conditions (2-1) and (2-2), one can see that

$$(2-54) \subset N_{CK_1/K}(Z). \tag{2-56}$$

Let us state this as a lemma.

Lemma 2.13. $\{\xi \in [0, 1]^2 : |n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi)| \lesssim K^{-1}\} \subset N_{CK_1/K}(Z)$.

We postpone the proof of this lemma and continue the proof of Theorem 2.12. By the lemma, each S in (2-54) is contained in a rectangle with dimension $1 \times CK_1K^{-1}$. Let us fix S . Suppose that

$$\frac{\text{length of the long direction of } S}{\text{length of the short direction of } S} \leq K_1. \tag{2-57}$$

Then we decompose S into squares of side length equal to the length of the short direction of S . Then the number of squares is bounded by $O(K_1)$. By Proposition 2.5, we have

$$\|f_S\|_{L^4} \lesssim (K_1)^{1/4} \left(\sum_{\{S' \in \mathcal{S}_X : S' \cap 2S \neq \emptyset\}} \|f_{S'}\|_{L^4}^2 \right)^{1/2}, \tag{2-58}$$

where $K^{-2} \lesssim X \lesssim (K_1K^{-1})^2$. Since K is sufficiently large compared to K_1 , the term $(K_1)^{1/4}$ on the right-hand side will not make any trouble.

Let us consider the case that

$$\frac{\text{length of the long direction of } S}{\text{length of the short direction of } S} \geq K_1. \tag{2-59}$$

Assume that K_1 is large enough that $K_1 > A^2$. By translation and rotation, we may assume that S is contained in $S' := [0, 1] \times [0, CK_1K^{-1}]$. By Lemma 2.7, after some change of variables, we may write the new phase function as

$$\xi_1 \xi_2 + a \xi_1^2 + b \xi_2^2 + \sum_{3 \leq j+k \leq d} b_{j,k} \xi_1^j \xi_2^k, \tag{2-60}$$

where

$$|a| + |b| + \sum_{3 \leq j+k \leq d} |b_{j,k}| \leq 10^{-d}. \tag{2-61}$$

We will apply the following lemma.

Lemma 2.14. *Let K_2 be a number such that $1 \leq K_2 \leq \delta^{-1/2}$. Let $S' = [0, 1] \times [0, K_2^{-1}]$. Suppose that the phase function is*

$$\Phi(\xi_1, \xi_2) := \xi_1 \xi_2 + a \xi_1^2 + b \xi_2^2 + \sum_{3 \leq j+k \leq d} b_{j,k} \xi_1^j \xi_2^k, \tag{2-62}$$

¹There is a minor technical issue. In the work of [Li and Yang 2021], the authors used a characteristic function for the cutoff function. The proof can be modified so that the cutoff is smooth. In our application, we use a smooth cutoff function.

where

$$|a| + |b| + \sum_{3 \leq j+k \leq d} |b_{j,k}| \leq 10^{-d}. \quad (2-63)$$

Then, for any $\epsilon > 0$, we have

$$\|f_{S'}\|_{L^4(B_{K_2})} \leq C_{\epsilon,d,A} K_2^\epsilon \sup_{K_2 \leq L \leq (K_2)^2} \left(\sum_{\{T \in \mathcal{S}_{L^{-1}} : T \subset 2S'\}} \|f_T\|_{L^4(w_{B_{K_2}})}^2 \right)^{1/2} \quad (2-64)$$

for all functions f whose Fourier support is in $N_\delta(\mathcal{M}_\Phi)$. Here $f_{S'}$ is the Fourier restriction of f to the rectangle $S' \times \mathbb{R}$.

Let us assume this lemma and finish the proof. By (2-53) and Lemma 2.14, we have

$$\|f\|_{L^4(B_K)} \leq C_\epsilon (K_1^{-1}K)^\epsilon K^\epsilon \sup_{(K_1^{-1}K) \leq L \leq (K)^2} \left(\sum_{T \in \mathcal{S}_{L^{-1}}} \|f_T\|_{L^4(w_{B_K})}^2 \right)^{1/2}. \quad (2-65)$$

To summarize, by considering all possible cases, we have

$$\|f\|_{L^4} \lesssim_\epsilon \left(\sum_{\tau' \in \mathcal{R}_{K_1^{-1}, K_1^{-1}, 0}} \|f_{\tau'}\|_{L^4}^4 \right)^{1/4} + K^{200} \left\| \prod_{i=1}^3 \tilde{f}_{\tau_i} \right\|_{L^4} + K^\epsilon \sup_{K_1^{-1}K \leq L \leq (K)^2} \left(\sum_{T \in \mathcal{S}_L} \|f_T\|_{L^4}^2 \right)^{1/2} \quad (2-66)$$

for some transverse τ_i . By Definition 2.11 and Lemma 2.9, we have

$$D(\delta) \lesssim_\epsilon D(K_1^2 \delta) + K^{200} D_{\text{mul}}(\delta, 10K^{-1}) + K^\epsilon \sup_{(K_1^{-1}K) \leq L \leq (K)^2} D(L\delta). \quad (2-67)$$

We apply this inequality to the first and third terms on the right-hand side of (2-67) repeatedly and obtain the desired result. \square

Proof of Lemma 2.13. For simplicity, we introduce $F(\xi) := n(\xi_{(1)}) \wedge n(\xi_{(2)}) \wedge n(\xi)$. By the definition of the normal vector $n(\xi)$, this can be rewritten as

$$F(\xi) = c \begin{vmatrix} \partial_1 \phi(\xi_{(1)}) & \partial_2 \phi(\xi_{(1)}) & -1 \\ \partial_1 \phi(\xi_{(2)}) & \partial_2 \phi(\xi_{(2)}) & -1 \\ \partial_1 \phi(\xi) & \partial_2 \phi(\xi) & -1 \end{vmatrix} \quad (2-68)$$

for some $|c| \sim 1$. After routine computations, this is equal to

$$-\partial_1 \phi(\xi) (\partial_2 \phi(\xi_{(1)}) - \partial_2 \phi(\xi_{(2)})) + \partial_2 \phi(\xi) (\partial_1 \phi(\xi_{(1)}) - \partial_1 \phi(\xi_{(2)})) - \partial_1 \phi(\xi_{(1)}) \partial_2 \phi(\xi_{(2)}) + \partial_1 \phi(\xi_{(2)}) \partial_2 \phi(\xi_{(1)}). \quad (2-69)$$

Recall that $\xi_{(1)}$ and $\xi_{(2)}$ are fixed. For simplicity, we write the function as

$$F(\xi) = A \partial_1 \phi(\xi) + B \partial_2 \phi(\xi) + C. \quad (2-70)$$

We first claim that

$$\max(|\partial_2 \phi(\xi_{(1)}) - \partial_2 \phi(\xi_{(2)})|, |\partial_1 \phi(\xi_{(1)}) - \partial_1 \phi(\xi_{(2)})|) \gtrsim \frac{1}{K_1}. \quad (2-71)$$

The claim is equivalent to

$$|\nabla\phi(\xi_{(1)}) - \nabla\phi(\xi_{(2)})| \gtrsim \frac{1}{K_1}. \tag{2-72}$$

By Taylor’s theorem, note that

$$|\nabla^2\phi(\xi_{(1)}) \cdot (\xi_{(1)} - \xi_{(2)})| \sim |\nabla\phi(\xi_{(1)}) - \nabla\phi(\xi_{(2)})|. \tag{2-73}$$

Since the absolute value of eigenvalues of the Hessian matrix of ϕ is comparable to 1 and by the separation $|\xi_{(1)} - \xi_{(2)}| \gtrsim 1/K_1$, we have

$$|\nabla^2\phi(\xi_{(1)}) \cdot (\xi_{(1)} - \xi_{(2)})| \gtrsim \frac{1}{K_1}. \tag{2-74}$$

This completes the proof of the claim.

Let us continue the proof of [Lemma 2.13](#). Recall [\(2-70\)](#). Let us consider the case $|A| \leq |B|$. The case $|A| \geq |B|$ can be dealt with similarly. To prove [Lemma 2.13](#), let us fix $\xi \in [0, 1]^2$ and suppose that

$$F(\xi) = X, \quad |X| \lesssim K^{-1}. \tag{2-75}$$

It suffices to show that there exists $\alpha \in \mathbb{R}$ such that

- $|\alpha| \lesssim K_1 K^{-1}$,
- $F(\xi + (\alpha, 0)) = 0$.

By Taylor’s theorem, since F is a polynomial of degree $d - 1$,

$$F(\xi + (\alpha, 0)) = F(\xi) + \partial_1 F(\xi)\alpha + \sum_{2 \leq j \leq d-1} \frac{\partial_1^j F(\xi)}{j!} \alpha^j. \tag{2-76}$$

By the expression [\(2-70\)](#),

$$\partial_1^j F(\xi) = A \partial_1^{j+1} \phi(\xi) + B \partial_1^j \partial_2 \phi(\xi). \tag{2-77}$$

By the normalization [\(2-1\)](#) and [\(2-2\)](#), we have

$$\begin{aligned} F(\xi + (\alpha, 0)) &= F(\xi) + (\partial_1 F(\xi))\alpha + O(B\alpha^2) \\ &= X + (\partial_1 F(\xi))\alpha + O(B\alpha^2). \end{aligned} \tag{2-78}$$

Note that $|\partial_1 F(\xi)| \sim |B|$. By [\(2-71\)](#), we have $|B| \gtrsim K_1^{-1}$, and we can find $|\alpha| \lesssim K_1 K^{-1}$ such that

$$X + (\partial_1 F(\xi))\alpha + O(B\alpha^2) = 0. \quad \square$$

Proof of [Lemma 2.14](#). We use an induction on K_2 . By taking $C_{\epsilon, d, A}$ sufficiently large, we may assume that [Lemma 2.14](#) is true for sufficiently large K_2 (compared to A).

We write our phase function [\(2-62\)](#) as

$$A(\xi_1) + B(\xi_1, \xi_2) := \left(a\xi_1^2 + \sum_{3 \leq j \leq d} b_{j,0} \xi_1^j \right) + (\xi_1 \xi_2 + b\xi_2^2 + \dots). \tag{2-79}$$

On the set $(\xi_1, \xi_2) \in S'$,

$$A(\xi_1) + B(\xi_1, \xi_2) = A(\xi_1) + O(K_2^{-1}). \tag{2-80}$$

By [Li and Yang 2021, Proposition 7.1],

$$\sup_{0 \leq \xi_1 \leq 1} |A(\xi_1)| \sim_d \max\left(|a|, \max_j |b_{j,0}|\right). \tag{2-81}$$

Let us consider the case that the left-hand side of (2-81) is smaller than AK_2^{-1} for some fixed large number A . We would like to show that $S' \in \mathcal{S}_{(K_2)^{-1}}$. First, note that S' is (ϕ, K_2^{-1}) -flat. Also, for any $z \in S'$, the angle between the ξ_1 -axis and v_z is smaller than $O(AK_2^{-1})$. Hence the rectangle, centered at z of dimension $1 \times K_2^{-1}$ with a long direction parallel to the vector v_z , is comparable to S' . So we see that $S' \in \mathcal{S}_{(K_2)^{-1}}$. This already gives the desired result (2-64).

We next consider the case that the left-hand side of (2-81) is larger than $\frac{1}{2}AK_2^{-1}$. We write our function as

$$f(x) = \int_{\mathbb{R}^3} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi. \tag{2-82}$$

We do a change of variables

$$x_3 \mapsto M^{-1}x_3 \quad \text{and} \quad \xi_3 \mapsto M\xi_3,$$

so that the Fourier support of f is contained in the δM -neighborhood of

$$\{M(A(\xi_1) + B(\xi_1, \xi_2)) : \xi_1 \in [0, 1], \xi_2 \in [0, K_2^{-1}]\} \tag{2-83}$$

for some $M^{-1} \gtrsim AK_2^{-1}$ such that

$$M \sup_{0 \leq \xi_1 \leq 1} |A(\xi_1)| \sim 1.$$

Note that

$$M(A(\xi_1) + B(\xi_1, \xi_2)) = MA(\xi_1) + O(A^{-1}). \tag{2-84}$$

After the change of variables, the ball B_{K_2} on the physical ball becomes a ball of radius K_2/M , which is still larger than A . Then we use a uniform ℓ^2 decoupling for a curve of [Yang 2021, Theorem 1.4] by ignoring the ξ_2 -variable and obtain

$$\|f_{S'}\|_{L^4} \leq C_{\epsilon/2}^{\text{curve}} (A')^{\epsilon/2} \left(\sum_{T'} \|f_{T'}\|_{L^4}^2 \right)^{1/2}. \tag{2-85}$$

Here the sidelength of the longest direction of T' is $(A')^{-1}$, which is smaller than or equal to $A^{-1/d}$. The set T' has dimension $(A')^{-1} \times (K_2)^{-1}$. We next do an isotropic rescaling (with translation), and T' becomes a rectangle with dimension $1 \times A'/K_2$. Since $A'/K_2 > 1/K_2$, by applying the induction hypothesis on K_2 and rescaling back, the left-hand side of (2-85) is bounded by

$$C_{\epsilon/2}^{\text{curve}} C_{\epsilon} (A')^{\epsilon/2} \left(\frac{K_2}{A'} \right)^{\epsilon} \left(\sum_T \|f_T\|_{L^4}^2 \right)^{1/2}. \tag{2-86}$$

Recall that $A' \geq A^{1/d}$. Since A is sufficiently large, we have

$$(A')^{-\epsilon/2} \leq (C_{\epsilon/2}^{\text{curve}})^{-1},$$

and we can close the induction. □

2.4. Bourgain–Demeter-type iteration. We have shown that the linear decoupling constant is bounded by the multilinear decoupling constant (Theorem 2.12). The next step is to bound the multilinear decoupling constant. Here is an ingredient.

Lemma 2.15 (ball inflation lemma). *Let ϕ be a function (2-1) satisfying (2-2). Let $p = 4$, $q = \frac{8}{3}$, and $N > 0$. Let τ_1, τ_2, τ_3 be N^{-1} -transverse squares. For any $\rho > 0$, $\epsilon > 0$, and $x_0 \in \mathbb{R}^3$,*

$$\left(\int_{B(x_0, \rho^{-2})} \left(\prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\rho, \rho, 0} : J \subset \tau_i\}} \|f_J\|_{L_{\#}^q(w_{B_{\rho^{-1}(x)})}}^2 \right)^{1/2} \right)^p dx \right)^{1/p} \leq C_{\epsilon, N} \rho^{-\epsilon} \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\rho, \rho, 0} : J \subset \tau_i\}} \|f_J\|_{L_{\#}^q(w_{B_{\rho^{-2}(x_0)})}}^2 \right)^{1/2} \tag{2-87}$$

for all functions f whose Fourier supports are in $N_{\rho^2}(\mathcal{M}_{\phi})$.

The proof of the ball inflation lemma is standard. We refer to [Bourgain et al. 2016, Theorem 6.6] for the details; see also [Bourgain and Demeter 2017b, Theorem 9.2].

We are ready to give a proof of Theorem 2.2. Let Γ be the smallest constant such that, for every $\epsilon > 0$, we have

$$D(\delta) \leq C_{\epsilon} \delta^{-\Gamma - \epsilon} \quad \text{for every dyadic } \delta < 1, \tag{2-88}$$

where C_{ϵ} is a constant depending on ϵ . Our goal is to prove $\Gamma = 0$.

We introduce some notation. Take $q := \frac{8}{3}$, and note that

$$\frac{1}{q} = \frac{1/2}{2} + \frac{1/2}{4}. \tag{2-89}$$

Define

$$\begin{aligned} \tilde{A}_2(b) &:= \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta b/2, \delta b/2, 0} : J \subset \tau_i\}} \|f_J\|_{L_{\#}^2(w_{B(x, \delta^{-b})})}^2 \right)^{1/2} \right\|_{L_{x \in \mathbb{R}^3}^4}, \\ \tilde{A}_q(b) &:= \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta b/2, \delta b/2, 0} : J \subset \tau_i\}} \|f_J\|_{L_{\#}^q(w_{B(x, \delta^{-b/2})})}^2 \right)^{1/2} \right\|_{L_{x \in \mathbb{R}^3}^4}, \\ \tilde{A}_4(b) &:= \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta b/2, \delta b/2, 0} : J \subset \tau_i\}} \|f_J\|_{L_{\#}^4(w_{B(x, \delta^{-b})})}^2 \right)^{1/2} \right\|_{L_{x \in \mathbb{R}^3}^4}, \end{aligned} \tag{2-90}$$

where $0 < b < 1$. For $0 < b < 1$ and $* = 2, q, 4$, we let $a_*(b)$ be the infimum over all exponents a satisfying

$$\tilde{A}_*(b) \lesssim_{a, N} \delta^{-a} \left(\sum_{R \in \mathcal{S}_{\delta}} \|f_R\|_{L^2(\mathbb{R}^3)}^2 \right)^{1/2} \tag{2-91}$$

for every $\delta > 0$, every N^{-1} -transverse squares τ_i , and every choice of a function f . Lastly, define

$$a_* := \liminf_{b \rightarrow 0} \frac{\Gamma - a_*(b)}{b} \tag{2-92}$$

for $* = 2, q, 4$.

Lemma 2.16. *We have the following inequalities:*

- (1) (setup) $a_* < \infty$ for $* = 2, q, 4$,
- (2) (rescaling) $a_4 \geq \Gamma$,
- (3) (L^2 -orthogonality) $a_2 \geq 2a_q$,
- (4) (ball inflation) $a_q \geq \frac{1}{2}a_4 + \frac{1}{2}a_2$.

Notice that by combining all the inequalities in the lemma, we obtain

$$\Gamma \leq a_4 \leq 2a_q - a_2 \leq 0. \quad (2-93)$$

Hence, in order to prove [Theorem 2.2](#), it suffices to prove [Lemma 2.16](#).

Proof of Lemma 2.16. Let us show inequality (1). It suffices to show

$$\Gamma \leq Cb + a_*(b) \quad (2-94)$$

for some constant C . By the essentially constant property and Bernstein's inequality, we have

$$\left\| \prod_{i=1}^3 f_{\tau_i} \right\|_{L^4} \lesssim \delta^{-Cb} \tilde{A}_*(b) \quad (2-95)$$

for any $* = 2, q, 4$ and $0 < b < 1$. By the definition of $a_*(b)$ (see [\(2-91\)](#)), we have

$$\left\| \prod_{i=1}^3 f_{\tau_i} \right\|_{L^4} \lesssim \delta^{-Cb-a_*(b)} \left(\sum_{R \in \mathcal{S}_\delta} \|f_R\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2}. \quad (2-96)$$

Hence we have

$$D_{\text{mul}}(\delta, N^{-1}) \lesssim \delta^{-Cb-a_*(b)}. \quad (2-97)$$

Combining this with [Theorem 2.12](#) gives

$$D(\delta) \lesssim_\epsilon \delta^{-\epsilon} \delta^{-Cb-a_*(b)}. \quad (2-98)$$

Since $\epsilon > 0$ is arbitrary, by the definition of Γ , we obtain [\(2-94\)](#).

Let us next show inequality (2). By the definition of a_4 , it follows from

$$a_4(b) \leq \Gamma \cdot (1 - b). \quad (2-99)$$

By Hölder's inequality and Minkowski's inequality, we have

$$\tilde{A}_4(b) \lesssim \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta^{b/2}, \delta^{b/2}, 0}: J \subset \tau_i\}} \|f_J\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2}. \quad (2-100)$$

By the parabolic rescaling lemma ([Lemma 2.9](#)), this is further bounded above by

$$D(\delta^{1-b}) \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{S}_\delta: J \subset \tau_i\}} \|f_J\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2} \lesssim D(\delta^{1-b}) \left(\sum_{J \in \mathcal{S}_\delta} \|f_J\|_{L^4(\mathbb{R}^3)}^2 \right)^{1/2}. \quad (2-101)$$

By (2-88) and (2-91), we have

$$\delta^{-a_4(b)} \lesssim \delta^{-\Gamma \cdot (1-b)}. \tag{2-102}$$

This gives (2-99).

Let us move onto inequality (3). By the L^2 -orthogonality and Hölder’s inequality, we obtain

$$\begin{aligned} \tilde{A}_2(b) &\lesssim \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta^b, \delta^b, 0} : J \subset \tau_i\}} \|f_J\|_{L^2_{\#}(w_{B(x, \delta^{-b})})}^2 \right)^{1/2} \right\|_{L^4_{x \in \mathbb{R}^3}} \\ &\lesssim \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta^b, \delta^b, 0} : J \subset \tau_i\}} \|f_J\|_{L^q_{\#}(w_{B(x, \delta^{-b})})}^2 \right)^{1/2} \right\|_{L^4_{x \in \mathbb{R}^3}}. \end{aligned} \tag{2-103}$$

The last expression is $\tilde{A}_q(2b)$, so we have

$$\delta^{-a_2(b)} \lesssim \delta^{-a_q(2b)}, \tag{2-104}$$

or equivalently,

$$a_2(b) \leq a_q(2b). \tag{2-105}$$

After some computations, this gives

$$a_2 \geq 2a_q. \tag{2-106}$$

This completes the proof of inequality (3).

Lastly, let us show inequality (4). By the ball inflation lemma (Lemma 2.15),

$$\tilde{A}_q(b) \lesssim_{\epsilon} \delta^{-\epsilon} \left\| \prod_{i=1}^3 \left(\sum_{\{J \in \mathcal{R}_{\delta^{b/2}, \delta^{b/2}, 0} : J \subset \tau_i\}} \|f_J\|_{L^q_{\#}(w_{B(x, \delta^{-b})})}^2 \right)^{1/2} \right\|_{L^4_{x \in \mathbb{R}^3}}. \tag{2-107}$$

By Hölder’s inequality and (2-89),

$$\tilde{A}_q(b) \lesssim_{\epsilon} \delta^{-\epsilon} \tilde{A}_2(b)^{1/2} \tilde{A}_4(b)^{1/2}. \tag{2-108}$$

By the definition of $a_*(b)$, this gives

$$\delta^{-a_q(b)} \lesssim \delta^{-\epsilon} \delta^{-(a_2(b)+a_4(b))/2}. \tag{2-109}$$

After some computations, this gives

$$a_q \geq \frac{1}{2}a_4 + \frac{1}{2}a_2. \tag{2-110}$$

This completes the proof of inequality (4). □

3. General manifolds

We have proved the ℓ^2 decoupling for perturbed hyperbolic paraboloids. We will use this decoupling as a black box and prove Theorem 3.1. This section does not contain any novelty, and we simply follow the argument of [Li and Yang 2021] in Appendix A.

The proof of Theorem 1.2 can be reduced to that of ℓ^2 decoupling for polynomials (Theorem 3.1). We refer to Section 2.3 of [Li and Yang 2021] for the details.

Theorem 3.1. Fix $d \geq 2$ and $\epsilon > 0$. Then there exists a sufficiently large number A depending on d and ϵ satisfying the following.

Let ϕ be a polynomial of two variables of degree d with coefficients bounded by 1. For any $\delta > 0$, there exists a collection \mathcal{S}_δ of finitely overlapping sets S such that:

(1) The overlapping number is $O(\delta^{-\epsilon})$ in the sense that

$$\sum_{S \in \mathcal{S}_\delta} \chi_S \leq C_{d,\epsilon} \delta^{-\epsilon}. \tag{3-1}$$

(2) S is $(\phi, A\delta)$ -flat and of the form of a parallelogram.

(3) We have

$$\|f\|_{L^4} \leq C_{d,\epsilon} \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2} \tag{3-2}$$

for all f whose Fourier support is in $N_\delta(\mathcal{M}_\phi)$.

The constant $C_{d,\epsilon}$ is independent of the choice of ϕ .

To prove [Theorem 3.1](#), we use [Theorem 2.2](#) as a black box. [Theorem 2.2](#) is a decoupling theorem for a phase function (2-1) satisfying (2-2), but by a simple change of variables, this theorem can be generalized to that for a polynomial function ϕ with bounded coefficients such that

$$|H_\phi(\xi)| \gtrsim 1 \tag{3-3}$$

for all $\xi \in [0, 1]^2$. Here H_ϕ is the determinant of the Hessian matrix of ϕ .

3.1. Sketch of the proof of [Theorem 3.1](#). Let M be a sufficiently large number, which will be determined later; see the end of [Section 3](#). The proof is composed of three steps.

Step 1: Dichotomy: a curved part and a flat part. Decompose $[0, 1]^2$ according to the size of $|H_\phi(\xi)|$:

$$\begin{aligned} S_{\text{curved}} &:= \{(\xi_1, \xi_2) \in [0, 1]^2 : |H_\phi(\xi_1, \xi_2)| > M^{-1}\}, \\ S_{\text{flat}} &:= \{(\xi_1, \xi_2) \in [0, 1]^2 : |H_\phi(\xi_1, \xi_2)| < M^{-1}\}. \end{aligned} \tag{3-4}$$

Introduce a parameter $M \ll M_1$. Decompose $[0, 1]^2$ into squares of side length M_1 . Then on each square, one can see that $|H_\phi| > \frac{1}{2}M^{-1}$ because $M_1^{-1} \ll M^{-1}$ and ∇H_ϕ is bounded. By the triangle inequality,

$$\|f\|_{L^4} \leq \left\| \sum_{\{|\tau|=M_1^{-1}: \tau \cap S_{\text{curved}} \neq \emptyset\}} f_\tau \right\|_{L^4} + \left\| \sum_{\{|\tau|=M_1^{-1}: \tau \subset S_{\text{flat}}\}} f_\tau \right\|_{L^4}. \tag{3-5}$$

After applying the triangle inequality, we apply [Theorem 2.2](#) (see also the discussion below [Theorem 3.1](#)) to the first term on the right-hand side, and this gives

$$\left\| \sum_{\{|\tau|=M_1^{-1}: \tau \cap S_{\text{curved}} \neq \emptyset\}} f_\tau \right\|_{L^4} \lesssim M_1 \left(\sum_{\{|\tau|=M_1^{-1}: \tau \cap S_{\text{curved}} \neq \emptyset\}} \|f_\tau\|_{L^4}^2 \right)^{1/2} \lesssim_\epsilon M_1 \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2}. \tag{3-6}$$

Since M_1 is a fixed number independent of δ , this already gives the desired bound. It remains to bound the second term on the right-hand side of (3-5).

Step 2: Analysis of the flat part. We next bound the term

$$\left\| \sum_{\{|\tau|=M_1^{-1}:\tau\subset S_{\text{flat}}\}} f_\tau \right\|_{L^4}. \quad (3-7)$$

By the generalized two-dimensional uniform ℓ^2 decoupling for polynomials, see Theorem 3.1 in [Li and Yang 2021], we can cover S_{flat} by rectangles T satisfying

$$T \subset \{(\xi_1, \xi_2) \in [0, 1]^2 : |H_\phi(\xi_1, \xi_2)| \lesssim M^{-1}\} \quad (3-8)$$

and

$$\left\| \sum_{\{|\tau|=M_1^{-1}:\tau\subset S_{\text{flat}}\}} f_\tau \right\|_{L^4} \lesssim_\epsilon M^\epsilon \left(\sum_T \|f_T\|_{L^4}^2 \right)^{1/2}. \quad (3-9)$$

Let us fix T . Take an affine transformation L such that it maps $[0, 1]^2$ to T . Take $\phi_1 := \phi \circ L$. Then one can see that

$$|H_{\phi_1}(\xi)| \lesssim M^{-1} \quad (3-10)$$

for all $\xi \in [0, 1]^2$. By [loc. cit., Proposition 7.1], all the coefficients of $H_{\phi_1}(\xi)$ are bounded above by $\sim M^{-1}$. Hence we can apply [loc. cit., Theorem 3.2] and obtain a rotation ρ such that

$$\phi_2(\xi_1, \xi_2) := \phi_1 \circ \rho(\xi_1, \xi_2) = a(\xi_1) + M^{-\alpha} b(\xi_1, \xi_2). \quad (3-11)$$

Here a and b have coefficients bounded above by ~ 1 , and α is a positive number. By the uncertainty principle, the term $M^{-\alpha} b(\xi_1, \xi_2)$ is negligible on the ball of radius M^α on the physical side. Hence we apply the two-dimensional uniform ℓ^2 decoupling of [loc. cit., Theorem 4.4] to the one variable polynomial $a(\xi_1)$ and partition $[0, 1]$ into intervals I such that each rectangle is $(\phi_2, AM^{-\alpha})$ -flat. Combining all the inequalities we have obtained so far, we have

$$\left\| \sum_{\{|\tau|=M_1^{-1}:\tau\subset S_{\text{flat}}\}} f_\tau \right\|_{L^4} \lesssim_\epsilon M^\epsilon \left(\sum_T \|f_T\|_{L^4}^2 \right)^{1/2} \lesssim_\epsilon M^\epsilon \left(\sum_{S' \in \mathcal{S}_{M^{-\alpha}}} \|f_{S'}\|_{L^4}^2 \right)^{1/2}. \quad (3-12)$$

Step 3: Iteration. By (3-5), (3-6), and (3-12), we obtain

$$\|f\|_{L^4} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2} + C_\epsilon (M^\alpha)^\epsilon \left(\sum_{S' \in \mathcal{S}_{M^{-\alpha}}} \|f_{S'}\|_{L^4}^2 \right)^{1/2}. \quad (3-13)$$

The first term is already of the desired form. To bound the second term, we fix S' and do rescaling. Let L be an affine transformation mapping $[0, 1]^2$ to S' . Since S' is $(\phi, AM^{-\alpha})$ -flat, we have

$$\sup_{u, v \in S'} |\phi(u) - \phi(v) - \nabla\phi(u) \cdot (u - v)| \leq AM^{-\alpha}. \quad (3-14)$$

Fix any point $u_0 \in S'$, and define

$$\tilde{\phi}(\xi) := A^{-1} M^\alpha (\phi(Lu_0) - \phi(L\xi) - \nabla\phi(Lu_0) \cdot (Lu_0 - L\xi)). \quad (3-15)$$

Then by (3-14), we have $\sup_{\xi \in [0,1]^2} |\tilde{\phi}(\xi)| \leq 1$. By [Li and Yang 2021, Proposition 7.1], all the coefficients of $\tilde{\phi}$ are bounded by 1. So this phase function satisfies the hypothesis of Theorem 3.1. We apply (3-13) to the function and obtain

$$\|f_{S'}\|_{L^4} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{\{S \in \mathcal{S}_\delta : S \cap S' \neq \emptyset\}} \|f_S\|_{L^4}^2 \right)^{1/2} + C_\epsilon (M^\alpha)^\epsilon \left(\sum_{\{S' \in \mathcal{S}_{M^{-2\alpha}} : S \cap S' \neq \emptyset\}} \|f_{S'}\|_{L^4}^2 \right)^{1/2}. \quad (3-16)$$

By (3-13) and (3-16), we have

$$\|f\|_{L^4} \leq (C_\epsilon + C_\epsilon^2 (M^\alpha)^\epsilon) \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2} + C_\epsilon^2 (M^{2\alpha})^\epsilon \left(\sum_{S' \in \mathcal{S}_{M^{-2\alpha}}} \|f_{S'}\|_{L^4}^2 \right)^{1/2}. \quad (3-17)$$

We repeat this process $O(\log_M \delta^{-1})$ times, and we obtain

$$\|f\|_{L^4} \lesssim_\epsilon (\delta^{-2\epsilon} + \delta^{-\log_M C_\epsilon}) \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2}. \quad (3-18)$$

We take M large enough that $\log_M C_\epsilon \leq \epsilon$. This completes the proof.

4. Proof of Corollaries 1.3 and 1.5

We prove Corollaries 1.3 and 1.5. We first remark that Corollary 1.3 holds under a general condition.

Remark 4.1. Fix $d \geq 2$. Let $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function. Given a straight line l intersecting $[0, 1]^2$, we parametrize the line by $\gamma(t)$ with the unit speed. Assume that

$$\phi(\gamma(t)) = a_0 + a_1 t + a_2 t^2 + \cdots + a_d t^d + E(t), \quad (4-1)$$

where

$$|E(t)| \leq c_\phi t^{d+1} \quad (4-2)$$

and

$$|a_2| + \cdots + |a_d| \geq C_\phi > 0. \quad (4-3)$$

Here, c_ϕ and C_ϕ are independent of the choice of the line l . Then (1-6) is true.

Corollary 1.5 is stated using the language of a partial differential equation. By Fourier series, we can write the operator $e^{it\tilde{\Delta}} f$ as

$$e^{it\tilde{\Delta}} f(x) = \sum_{\xi \in [-N, N]^2 \cap \mathbb{Z}^2} \hat{f}(\xi) e(i(x_1, x_2, x_3) \cdot (\xi_1, \xi_2, \xi_1^2 - \alpha \xi_2^2)). \quad (4-4)$$

Also by Parseval's identity, we have

$$\|f\|_{L^2(\mathbb{T}^2)}^2 \sim \sum_{\xi \in [-N, N]^2 \cap \mathbb{Z}^2} |\hat{f}(\xi)|^2. \quad (4-5)$$

Hence Corollary 1.5 can be rephrased as follows.

Corollary 4.2. *Let α be irrational. Suppose that*

$$\phi(\xi_1, \xi_2) = \xi_1^2 - \xi_2^2 \quad \text{and} \quad \Lambda_\delta = (\delta\mathbb{Z} \times \alpha\delta\mathbb{Z}) \cap [0, 1]^2. \tag{4-6}$$

Then, for $2 \leq p \leq 4$ and $\epsilon > 0$, we have

$$\left\| \sum_{\xi \in \Lambda_\delta} a_\xi e(x \cdot (\xi, \phi(\xi))) \right\|_{L^p_\#(B_{\delta^{-3}})} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{\xi \in \Lambda_\delta} |a_\xi|^2 \right)^{1/2}. \tag{4-7}$$

4.1. Proof of Corollary 1.3. Suppose that the manifold \mathcal{M}_ϕ does not contain a line. Fix $\epsilon > 0$. By Theorem 3.1 with δ replaced by δ^d , we have

$$\|f\|_{L^4} \leq C_{d,\epsilon} \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_{\delta^d}} \|f_S\|_{L^4}^2 \right)^{1/2} \tag{4-8}$$

for all f whose Fourier support is in $N_{\delta^d}(\mathcal{M}_\phi)$. Since ϕ does not contain a line, by a compactness argument, for any l , we can parametrize it as a function of t and write

$$\phi(t, at + b) = a_0 + a_1t + a_2t^2 + \dots + a_d t^d, \tag{4-9}$$

where

$$\max(|a_2|, |a_3|, \dots, |a_d|) \sim 1. \tag{4-10}$$

Suppose that S is a rectangle and is $(\phi, A\delta^d)$ -flat. We claim that the length of a long direction of S is smaller than or equal to $C_\phi\delta$ for some constant C_ϕ depending on the choice of ϕ . We may assume that the angle between the long direction of S and the ξ_1 -axis is smaller than or equal to $\frac{\pi}{4}$. Let l be a line segment passing through the center of S and parallel to the long direction of S but contained in S . For convenience, we introduce a parametrization of the line l : $\gamma(t) = (t, at + b)$, where $t \in [b_0, b_1]$. To prove the claim, it suffices to show that

$$|b_1 - b_0| \leq C_{\phi,A}\delta. \tag{4-11}$$

By the definition of $(\phi, A\delta^d)$ -flat, we have

$$\sup_{t \in [b_0, b_1]} |\phi(\gamma(b_0)) - \phi(\gamma(t)) - \nabla\phi(\gamma(b_0)) \cdot (\gamma(b_0) - \gamma(t))| \leq A\delta^d. \tag{4-12}$$

By (4-9) and (4-10), after some computations, this can be rewritten as

$$\sup_{t \in [b_0, b_1]} |\tilde{a}_1(t - b_0) + \tilde{a}_2(t - b_0)^2 + \dots + \tilde{a}_d(t - b_0)^d| \leq A\delta^d, \tag{4-13}$$

where

$$\max(|\tilde{a}_2|, |\tilde{a}_3|, \dots, |\tilde{a}_d|) \sim 1. \tag{4-14}$$

By [Li and Yang 2021, Proposition 7.1], (4-13) gives

$$\max(|\tilde{a}_2(b_0 - b_1)^2|, |\tilde{a}_3(b_0 - b_1)^3|, \dots, |\tilde{a}_d(b_0 - b_1)^d|) \lesssim A\delta^d. \tag{4-15}$$

The condition (4-14) gives $|b_1 - b_0| \lesssim \delta$, and this finishes the proof of the claim.

We have proved the claim. [Corollary 1.3](#) will follow by counting the number of frequencies in Λ_δ contained in a δ^d -neighborhood of each $S \in \mathcal{S}_\delta$. The claim says that each S contains at most $\lesssim 1$ many frequencies in Λ_δ , and this gives the desired result. Let us give more details. We take f to be a Fourier transform of

$$\sum_{\xi \in \Lambda_\delta} \psi_\xi, \quad (4-16)$$

where ψ_ξ is a smooth bump function supported on the ball of radius δ^d centered at the point $(\xi, \phi(\xi))$. By the claim, we have

$$\left(\sum_{S \in \mathcal{S}_{\delta^d}} \|f_S\|_{L^4}^2 \right)^{1/2} \lesssim \left(\sum_{\xi \in \Lambda_\delta} \|\widehat{\psi}_\xi\|_{L^4}^2 \right)^{1/2}. \quad (4-17)$$

On the other hand, by the definition of f , we have

$$\|f\|_{L^4} \sim \left\| \sum_{\xi} \widehat{\psi}_\xi \right\|_{L^4}. \quad (4-18)$$

Note that

$$\widehat{\psi}_\xi(x) = \delta^{3d} e(i\xi, \phi(\xi)) \psi_{B(0, \delta^{-d})}, \quad (4-19)$$

where ψ_B is a smooth function essentially supported on the ball B . [Corollary 1.3](#) follows from this, [\(4-8\)](#), [\(4-9\)](#), and [\(4-10\)](#). This completes the proof.

4.2. Proof of [Corollary 1.5](#). As we discussed, it suffices to prove [Corollary 4.2](#). For simplicity, we only consider $\alpha = \sqrt{2}$. The general case can be proved identically. For $\phi(\xi_1, \xi_2) = \xi_1^2 - \xi_2^2$ and $\Lambda_\delta = \delta\mathbb{Z} \times \sqrt{2}\delta\mathbb{Z}$, our goal is to show that

$$\left\| \sum_{\xi \in \Lambda_\delta} a_\xi e(x \cdot (\xi, \phi(\xi))) \right\|_{L^\#_p(B_{\delta^{-3}})} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{\xi \in \Lambda_\delta} |a_\xi|^2 \right)^{1/2}. \quad (4-20)$$

This will follow easily from [Theorem 1.8](#) by counting the number of frequencies in Λ_δ contained in a δ^3 -neighborhood of each $S \in \mathcal{S}_\delta$. Suppose that $(m_i\delta, n_i\sqrt{2}\delta) \in \Lambda_\delta \cap N_{\delta^3}(S)$ for $i = 1, 2$. Then using the definition of (ϕ, δ^3) -flat, we have

$$|m_1^2\delta^2 - 2n_1^2\delta^2 - (m_2^2\delta^2 - 2n_2^2\delta^2) - 2(m_1\delta, -\sqrt{2}n_1\delta) \cdot ((m_1 - m_2)\delta, \sqrt{2}(n_1 - n_2)\delta)| \lesssim \delta^3.$$

This simplifies to

$$|(m_1 - m_2)^2 - 2(n_1 - n_2)^2| \lesssim \delta,$$

so

$$|[m_1 - m_2 - \sqrt{2}(n_1 - n_2)][m_1 - m_2 + \sqrt{2}(n_1 - n_2)]| \lesssim \delta.$$

By Lemma 4 of Section 2, Chapter 2 of [[Cassels 1972](#)], we have $1/b^{1+\epsilon'} \lesssim_{\epsilon'} |a + \sqrt{2}b|$ for any $a, b \in \mathbb{Z}$. Therefore, if $(m_1, n_1) \neq (m_2, n_2)$, the above displayed inequality implies that $|n_1 - n_2| \gtrsim_{\epsilon'} \delta^{-1+\epsilon'}$. Since the elements of $\Lambda_\delta \cap N_{\delta^3}(S)$ are $\gtrsim \delta$ -separated, there are fewer than $\lesssim_{\epsilon'} \delta^{-\epsilon'}$ frequencies in $\Lambda_\delta \cap N_{\delta^3}(S)$, as desired.

Appendix A: Partition is not enough

In this appendix, we prove that there is no ℓ^2 decoupling for the hyperbolic paraboloid using a partition. Note that in [Theorem 1.8](#) we introduce $O(\log \delta^{-1})$ many partitions to obtain ℓ^2 decoupling for the hyperbolic paraboloid. The following theorem shows that it is necessary to introduce many partitions.

Theorem A.1. *Let \mathcal{M}_ϕ be the hyperbolic paraboloid given by $\phi(\xi_1, \xi_2) = \xi_1 \xi_2$. Fix $0 < \epsilon < \frac{1}{100}$ and $A \geq 1$. Then the following is false: for any $\delta > 0$, there exists a family \mathcal{S}_δ of rectangles $S \subset \mathbb{R}^2$ such that*

(1) *the interiors of rectangles are disjoint:*

$$\text{int}(S_1) \cap \text{int}(S_2) = \emptyset, \quad S_1, S_2 \in \mathcal{S}_\delta, \quad (\text{A-1})$$

(2) *every S is $(\phi, A\delta)$ -flat,*

(3) *for all f whose Fourier support is in $N_\delta(\mathcal{M}_\phi)$,*

$$\|f\|_{L^4} \leq C_{\epsilon, A} \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2}. \quad (\text{A-2})$$

Proof. Fix $\delta > 0$. For simplicity, assume that δ^{-1} is a dyadic number. Suppose that such a family \mathcal{S}_δ exists for a contradiction. Let us first consider the partition

$$\begin{aligned} [0, 1]^2 &= \bigcup_{a \in \mathbb{Z} \cap [0, \delta^{-1} - 1]} ([0, 1] \times [a\delta, a\delta + \delta]) =: \bigcup_a C_a, \\ [0, 1]^2 &= \bigcup_{b \in \mathbb{Z} \cap [0, \delta^{-1} - 1]} ([b\delta, b\delta + \delta] \times [0, 1]) =: \bigcup_b D_b. \end{aligned} \quad (\text{A-3})$$

Let us fix C_a . Consider a collection of elements of \mathcal{S}_δ which have a large intersection with C_a :

$$\mathcal{S}_{\delta, C_a} := \{S \in \mathcal{S}_\delta : |S \cap C_a| \geq \delta^1 \delta^{1/2 - \epsilon}\}. \quad (\text{A-4})$$

Define $\mathcal{S}_{\delta, C_b}$ similarly. By abusing notation, let us denote by $\mathcal{S}_{\delta, a}$ and $\mathcal{S}_{\delta, b}$ the sets $\mathcal{S}_{\delta, C_a}$ and $\mathcal{S}_{\delta, D_b}$. Recall that elements of the set $\mathcal{S}_{\delta, a}$ are disjoint by the first condition [\(A-1\)](#).

We claim that

$$\left| \bigcup_{S \in \mathcal{S}_{\delta, a}} S \cap C_a \right| \geq \frac{99}{100} \delta \quad \text{and} \quad \left| \bigcup_{S \in \mathcal{S}_{\delta, b}} S \cap D_b \right| \geq \frac{99}{100} \delta. \quad (\text{A-5})$$

Let us assume this claim for a moment and finish the proof of the theorem. By taking the union over a in [\(A-5\)](#), we have

$$\left| \bigcup_{a \in \mathbb{Z} \cap [0, \delta^{-1} - 1]} \bigcup_{S \in \mathcal{S}_{\delta, a}} S \cap C_a \right| \geq \frac{99}{100}. \quad (\text{A-6})$$

By pigeonholing, there exists $D_b \in \mathcal{S}_{\delta, b}$ such that

$$\left| D_b \cap \left(\bigcup_{a \in \mathbb{Z} \cap [0, \delta^{-1} - 1]} \bigcup_{S \in \mathcal{S}_{\delta, a}} S \cap C_a \right) \right| \geq \frac{99}{100} \delta. \quad (\text{A-7})$$

This implies

$$\left| D_b \cap \left(\bigcup_{a \in \mathbb{Z} \cap [0, \delta^{-1} - 1]} \bigcup_{S \in \mathcal{S}_{\delta, a}} S \right) \right| \geq \frac{99}{100} \delta. \quad (\text{A-8})$$

We claim that if $S \in \mathcal{S}_{\delta, a}$ for some a , then

$$|D_b \cap S| \lesssim \delta^1 \delta^{1/2+\epsilon}. \quad (\text{A-9})$$

Note that this means that S does not belong to $\mathcal{S}_{\delta, D_b}$. Since the area of D_b is δ , the first inequality of (A-5) gives

$$\left| \bigcup_{S \in \mathcal{S}_{\delta, D_b}} S \cap D_b \right| \leq \frac{1}{100} \delta. \quad (\text{A-10})$$

This contradicts the second inequality of (A-5). Let us give a proof of (A-9). Suppose that $S \in \mathcal{S}_{\delta, a}$. It suffices to prove that

$$S \subset (\alpha, \beta) + ([0, 1] \times [0, C\delta^{1/2+\epsilon}]) \quad (\text{A-11})$$

for some $\alpha, \beta \in [0, 1]$. To prove this, we use the assumption that S is $(\phi, A\delta)$ -flat. By the definition of $\mathcal{S}_{\delta, a}$, the rectangle S contains a line segment parallel to the ξ_1 -axis with length $\delta^{1/2-\epsilon}$. Since our manifold is translation invariant, for simplicity, assume that the line is $[0, \delta^{1/2-\epsilon}] \times \{0\}$. We will show that if $p = (p_1, p_2)$ is an element of S , then $|p_2| \lesssim \delta^{1/2+\epsilon}$. This gives the proof of (A-9). To show the bound of p_2 , we set $u = (0, 0)$. Then by the definition of $(\phi, A\delta)$ -flat (Definition 1.1),

$$|p_1 p_2| \leq A\delta. \quad (\text{A-12})$$

Similarly, we use $u = (\delta^{1/2-\epsilon}, 0)$, and this gives

$$|p_1 p_2 - \delta^{1/2-\epsilon} p_2| \leq A\delta. \quad (\text{A-13})$$

Combining these two gives $|p_2| \lesssim \delta^{1/2+\epsilon}$.

It remains to prove (A-5). By symmetry, let's show only the first inequality. We will use assumption (A-2). Fix C_a . Let us first show that there exists $S \in \mathcal{S}_{\delta}$ such that

$$|S \cap C_a| \geq (\log \delta^{-1})^{-10} \delta^{1+4\epsilon}. \quad (\text{A-14})$$

Suppose that such an S does not exist for a contradiction. We take f such that

- (1) \hat{f} is supported on the δ -neighborhood of the set $\mathcal{M}_{\phi} \cap (C_a \times \mathbb{R})$,
- (2) \hat{f} is equal to 1 on the $\frac{1}{2}\delta$ -neighborhood of the set $\mathcal{M}_{\phi} \cap (C_a \times \mathbb{R})$.

Here \mathcal{M}_{ϕ} is the hyperbolic paraboloid. Then f is essentially supported on a box with dimension $\delta^{-1} \times 1 \times \delta^{-1}$ and has amplitude $\sim \delta^2$. So we have

$$\|f\|_{L^4} \sim \delta^2 \delta^{-2/4} \sim \delta^{3/2}. \quad (\text{A-15})$$

On the other hand, by pigeonholing, there exists j such that

$$\left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^4}^2 \right)^{1/2} \lesssim (\log \delta^{-1})^{1/2} \left(\sum_{S \in \mathcal{S}_\delta^j} \|f_S\|_{L^4}^2 \right)^{1/2}, \tag{A-16}$$

where the elements S of \mathcal{S}_δ^j satisfy

$$|S \cap C_a| \sim \delta 2^{-j}. \tag{A-17}$$

Here, we used the fact that, for large j , the right-hand side of (A-16) is negligible (which is proved in (A-18)). Since we are assuming that there does not exist S satisfying (A-14), (A-16) is true for $j > 4\epsilon \log_2 \delta^{-1} + 10 \log_2 \log \delta^{-1}$. Note that f_S is essentially constant on a box with volume $(\delta \delta 2^{-j})^{-1}$ and has amplitude $\delta \delta 2^{-j}$. Since \mathcal{S}_δ is a partition (see (A-1)), the cardinality of \mathcal{S}_δ^j is bounded by 2^j . So we have

$$\left(\sum_{S \in \mathcal{S}_\delta^j} \|f_S\|_{L^4}^2 \right)^{1/2} \lesssim (\#\mathcal{S}_\delta^j)^{1/2} \max_S \|f_S\|_{L^4} \lesssim 2^{j/2} (\delta^2 2^{-j}) (\delta^2 2^{-j})^{-1/4} \sim \delta^{3/2} 2^{-j/4}. \tag{A-18}$$

Inequality (A-2), (A-15), and (A-18) give

$$\delta^{3/2} \lesssim (\log \delta^{-1})^{1/2} \delta^\epsilon \delta^{3/2} 2^{-j/4}. \tag{A-19}$$

We get a contradiction from the assumption that $j > 4\epsilon \log_2 \delta^{-1} + 10 \log_2 \log \delta^{-1}$. The same argument works for a general situation. Let $C \subset C_a \subset [0, 1]^2$ be a convex set (polygon). Then we can show that there exists $S \in \mathcal{S}_\delta$ such that

$$|S \cap C| \gtrsim (\log \delta^{-1})^{-10} \delta^{4\epsilon} |C|. \tag{A-20}$$

Let us explain the proof. First, since C is a convex set, we can find a rectangle $\tilde{C} \subset C$ such that $|\tilde{C}| \gtrsim |C|$; for example, by the Kovner–Besicovitch theorem. Since \tilde{C} is a rectangle in C_a , we can repeat the proof of (A-14) and find $S \in \mathcal{S}_\delta$ such that

$$|S \cap C| \geq |S \cap \tilde{C}| \geq (\log \delta^{-1})^{-10} \delta^{4\epsilon} |\tilde{C}| \gtrsim (\log \delta^{-1})^{-10} \delta^{4\epsilon} |C|. \tag{A-21}$$

This gives (A-20).

We have shown that there exists $S_1 \in \mathcal{S}_\delta$ such that (A-14) holds. In particular, $S_1 \in \mathcal{S}_{\delta,a}$; see (A-4) for the definition of $\mathcal{S}_{\delta,a}$ and note that we used the assumption that ϵ is small. If we have $|S_1 \cap C_a| \geq \frac{99}{100} \delta$, then this gives (A-5). So suppose that

$$\delta^{1+4\epsilon} \leq |S_1 \cap C_a| \leq \frac{99}{100} \delta. \tag{A-22}$$

Since S_1 is a rectangle, the set $C_a \setminus (S_1)^c$ is either a convex set or a union of two convex sets. Let us write it as

$$C_a \setminus (S_1)^c = C_{11} \cup C_{12}. \tag{A-23}$$

Here C_{11} and C_{12} are disjoint convex sets and

$$|C_{11}| + |C_{12}| \leq |C_a| (1 - \delta^{4\epsilon}). \tag{A-24}$$

This finishes the first round of the iteration. Let us explain how to proceed. If C_{11} satisfies $|C_{11}| \leq \delta\delta^{1/4}$, then we leave this set. If $|C_{11}| > \delta\delta^{1/4}$, then we apply (A-20) to the convex set C_{11} and obtain the set $S_2 \in \mathcal{S}_\delta$ such that

$$|S_2 \cap C_{11}| \gtrsim (\log \delta^{-1})^{-10} \delta^{4\epsilon} |C_{11}|. \quad (\text{A-25})$$

By the stopping time condition $|C_{11}| > \delta\delta^{1/4}$ and the assumption that ϵ is small, we have $S_2 \in \mathcal{S}_{\delta, C_a}$; see (A-4) for the definition. We write $C_{11} \setminus (S_2)^c = C_{111} \cup C_{112}$. Then we have

$$|C_{111}| + |C_{112}| \leq |C_{11}|(1 - \delta^{4\epsilon}), \quad (\text{A-26})$$

where the C_{11j} are convex sets. Repeat this process for C_{12} . If $|C_{12}| < \delta\delta^{1/4}$, then we have

$$|C_{111}| + |C_{112}| + |C_{12}| \leq |C_a|(1 - \delta^{4\epsilon})^2 + |C_{12}|\delta^{4\epsilon}. \quad (\text{A-27})$$

If $|C_{12}| > \delta\delta^{1/4}$, then we have

$$|C_{111}| + |C_{112}| + |C_{113}| + |C_{114}| \leq |C_a|(1 - \delta^{4\epsilon})^2. \quad (\text{A-28})$$

This finishes the second round of the iteration. We repeat this process M times with $M \gtrsim \delta^{-6\epsilon}$. Denote by $\{C_X\}_X$ and $\{S_Y\}_Y$ a collection of convex sets and a collection of rectangles $S \in \mathcal{S}_\delta$, respectively, that we obtained via this process. Note that

$$\sum_X |C_X| + \sum_Y |S_Y \cap C_a| = |C_a| = \delta. \quad (\text{A-29})$$

We have

$$\begin{aligned} \sum_X |C_X| &= \sum_{\{X: |C_X| \geq \delta\delta^{1/4}\}} |C_X| + \sum_{\{X: |C_X| < \delta\delta^{1/4}\}} |C_X| \\ &\leq |C_a|(1 - \delta^{4\epsilon})^M + \left| \bigcup_{\{X: |C_X| < \delta\delta^{1/4}\}} C_X \right| \delta^{4\epsilon} \leq \frac{1}{100} |C_a|. \end{aligned} \quad (\text{A-30})$$

The last two inequalities follow from the fact that all C_X are disjoint and $C_X \subset C_a$. This completes the proof of the claim (A-5). \square

Appendix B: Higher dimensions

In this appendix, we prove that an analogous result to Theorem 1.2 is false in higher dimensions. For convenience, let us consider only a manifold in \mathbb{R}^4 . Higher-dimensional manifolds can be handled in a similar way. Definitions 1.1 and 2.1 can be naturally generalized to higher dimensions. We will not state them. Recall that the critical exponent p of decoupling for the hyperbolic paraboloid in \mathbb{R}^4 is $\frac{10}{3}$.

Theorem B.1. *Consider $\phi(\xi_1, \xi_2, \xi_3) = \xi_1^2 + \xi_2^2 - \xi_3^2$. Fix $0 < \epsilon < \frac{1}{1000}$. Let A be a constant. Then the following statement is false.*

For any $\delta > 0$, there exists a collection \mathcal{S}_δ of rectangular boxes $S \subset [0, 1]^3$ such that:

(1) *The overlapping number is $O(\log \delta^{-1})$ in the sense that*

$$\sum_{S \in \mathcal{S}_\delta} \chi_S \leq C_\epsilon \log(\delta^{-1}). \quad (\text{B-1})$$

(2) S is $(\phi, A\delta)$ -flat and of the form of a parallelogram.

(3) We have

$$\|f\|_{L^{10/3}} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^{10/3}}^2 \right)^{1/2} \tag{B-2}$$

for all f whose Fourier support is in $N_\delta(\mathcal{M}_\phi)$.

The same conclusion of [Theorem B.1](#) occurs if (B-2) is replaced with

$$\|f\|_{L^p} \leq C_\epsilon \delta^{-\epsilon} \left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^p}^2 \right)^{1/2} \tag{B-3}$$

for $2 < p \leq \frac{10}{3}$. Inequality (B-6) is the only place where (B-2) is used. One may check that (B-3) still implies the desired result.

Proof. Let us prove the result by contradiction. Fix $0 < \epsilon < 10^{-10}$. Define

$$\mathcal{C} := \{(\xi_1, \xi_2, \xi_3) \in [0, 1]^3 : \xi_1^2 + \xi_2^2 - \xi_3^2 = 0\}. \tag{B-4}$$

We cover \mathcal{C} by $\sim \delta^{-1/2}$ many canonical blocks P with dimension $1 \times \delta^{1/2} \times \delta$. Let us denote by \mathcal{P} the collection of blocks. Note that each block is $(\phi, A\delta)$ -flat for some large constant A . For given $P \in \mathcal{P}$, we take translated copies and cover $[0, 1]^3$ so that they are disjoint. Denote by \mathcal{P}_P the collection of translated copies of P . Note that the cardinality of \mathcal{P}_P is comparable to $\delta^{-3/2}$.

Fix $P' \in \mathcal{P}_P$ for some $P \in \mathcal{P}$. We next decompose P' into parallel tubes with radius δ^X , where X is sufficiently large number. Let us denote by $\mathcal{P}_{P,P'}$ the collections of tubes P'' . We claim that, for given $P'' \in \mathcal{P}_{P,P'}$, there is an element S of \mathcal{S}_δ such that the diameter of the set $S \cap P''$ is larger than or equal to $(\log \delta^{-1})\delta^{5\epsilon}$. To prove the claim, let us use the hypothesis (B-2). Take a function f such that

$$\hat{f}(\xi) := \sum_{j=1}^{(\log \delta^{-1})\delta^{-5\epsilon}} f_j(\xi) := \sum_{j=1}^{(\log \delta^{-1})\delta^{-5\epsilon}} 1_{N_{\delta^X}(B_j)}(\xi), \tag{B-5}$$

where $\{B_j\}_{j=1}^{(\log \delta^{-1})\delta^{-5\epsilon}}$ is an arithmetic progression with difference $(\log \delta^{-1})\delta^{5\epsilon}$ and $\bigcup_j B_j \subset P''$. By direct computations, one can see that

$$\|f\|_{L^{10/3}} \gtrsim (\log \delta^{-1})^{1/5} \delta^{-5\epsilon(1/2-1/(10/3))} \left(\sum_j \|f_j\|_{L^{10/3}}^2 \right)^{1/2}. \tag{B-6}$$

Hence inequality (B-2) says that there must be an element S of \mathcal{S}_δ containing at least two B_j . This means that the diameter of $S \cap P''$ is larger than or equal to $(\log \delta^{-1})\delta^{5\epsilon}$. This completes the proof of the claim.

For given $P' \in \mathcal{P}_P$, denote by $\mathcal{S}_{P'}$ the elements of $S \in \mathcal{S}_\delta$ satisfying the following property: for some $P'' \in \mathcal{P}_{P,P'}$ the diameter of $S \cap P''$ is larger than or equal to $(\log \delta^{-1})\delta^{5\epsilon}$. The claim says that $\mathcal{S}_{P'}$ is nonempty.

We next claim that if S is $(\phi, A\delta)$ -flat and $S \in \mathcal{S}_\delta$, then S is contained in a box with dimension $\delta^{1/2-5\epsilon/2} \times \delta^{1-5\epsilon} \times \delta^{5\epsilon}$. Let us give a proof. First of all, by an affine transformation, we may assume that S

contains a line $\{0\} \times \{0\} \times [0, \delta^{5\epsilon}]$, and our new phase function is $\tilde{\phi}(\xi) = \xi_1^2 + \xi_2 \xi_3$. Let $v = (v_1, v_2, v_3) \in S$. We will show that

$$|v_1| \lesssim \delta^{1/2-5\epsilon/2} \quad \text{and} \quad |v_2| \lesssim \delta^{1-5\epsilon}. \quad (\text{B-7})$$

By the definition of a $(\phi, A\delta)$ -flat set with $u = (0, 0, 0)$, we obtain

$$|v_1^2 + v_2 v_3| \leq A\delta. \quad (\text{B-8})$$

We next use $u = (0, 0, \delta^{5\epsilon})$ and obtain

$$|v_1^2 + v_2 v_3 - \delta^{5\epsilon} v_2| \leq A\delta. \quad (\text{B-9})$$

These two inequalities give $|v_2| \lesssim \delta^{1-5\epsilon}$. This bound and (B-8) give $|v_1| \lesssim \delta^{1/2-5\epsilon/2}$. This proves the claim.

By the claim, if $P'_1 \in \mathcal{P}_{P_1}$ and $P'_2 \in \mathcal{P}_{P_2}$ and the angle of the longest directions of P_1 and P_2 is greater than $\delta^{1/2-100\epsilon}$, then any two sets $S_1 \in \mathcal{S}_{P'_1}$ and $S_2 \in \mathcal{S}_{P'_2}$ are distinct. Note also that

$$\left| \bigcup_{S' \in \mathcal{S}_{P'}} S' \cap P' \right| \gtrsim \delta^{5\epsilon} |P'|. \quad (\text{B-10})$$

Let us finish the proof of the theorem. To get a contradiction, we need to prove that there exists $\xi \in [0, 1]^3$ such that the number of $S \in \mathcal{S}_\delta$ containing ξ is greater than the right-hand side of (B-1). To prove this, it suffices to show that there exists $\xi \in [0, 1]^3$ satisfying

$$\sum_{P \in \mathcal{P}} \sum_{P' \in \mathcal{P}_P} \sum_{S' \in \mathcal{S}_{P'}} 1_{S' \cap P'}(\xi) \gtrsim \delta^{-\alpha-100\epsilon} \quad (\text{B-11})$$

for some number $\alpha > 0$. This follows from a pigeonholing argument with

$$\begin{aligned} \int_{[0,1]^3} \sum_{P \in \mathcal{P}} \sum_{P' \in \mathcal{P}_P} \sum_{S' \in \mathcal{S}_{P'}} 1_{S' \cap P'}(\xi) d\xi &= \sum_{P \in \mathcal{P}} \sum_{P' \in \mathcal{P}_P} \int_{[0,1]^3} \sum_{S' \in \mathcal{S}_{P'}} 1_{S' \cap P'}(\xi) d\xi \\ &\gtrsim \delta^{-3/2} \delta^{-1/2} \delta^{5\epsilon} \delta^{3/2} \sim \delta^{-1/2+5\epsilon}. \quad \square \end{aligned}$$

Appendix C: Sharpness of Theorem 1.2

Let us discuss the sharpness of Theorem 1.2. More precisely, we would like to discuss if the range of p for which (1-4) holds is sharp. An example of a manifold where the aforementioned range of p is not sharp is given by

$$\phi(\xi_1, \xi_2) = \xi_1^2. \quad (\text{C-1})$$

The ℓ^2 decoupling for (C-1) is proved for $2 \leq p \leq 6$ by [Bourgain and Demeter 2015], and this range of p is sharp. Motivated by this example, let us introduce the following definition.

Definition C.1. We say $\phi(\xi_1, \xi_2)$ depends only on a variable if there exists an affine transformation L such that

$$\phi(L(\xi_1, \xi_2)) = \psi(\xi_1) + a\xi_2 \quad (\text{C-2})$$

for some function ψ and some $a \in \mathbb{R}$. Second, we say $\phi(\xi_1, \xi_2)$ does not depend on any variable if

$$\phi(\xi_1, \xi_2) = a + b\xi_1 + c\xi_2 \tag{C-3}$$

for some $a, b, c \in \mathbb{R}$. Lastly, we say $\phi(\xi_1, \xi_2)$ depends on two variables if it is not of the form of (C-3) and there does not exist L such that (C-2) holds.

Here is a complete characterization of the ℓ^2 decoupling theorem.

Proposition C.2. *Let $p \geq 2$. Consider a polynomial $\phi := \phi(\xi_1, \xi_2)$. Suppose that \mathcal{S}_δ is a family constructed in Section 2.1 and Section 3.*

- (1) *Let ϕ depend on two variables. Then (1-4) is true for $2 \leq p \leq 4$.*
- (2) *Let ϕ depend only on one variable. Then (1-4) is true for $2 \leq p \leq 6$.*
- (3) *Let ϕ not depend on any variable. Then (1-4) is true for $2 \leq p < \infty$.*

The ranges of p stated in items (1), (2), and (3) are sharp.

Proof. Let us first prove item (1). Theorem 1.2 gives (1-4) for $2 \leq p \leq 4$, so it suffices to prove that the range is sharp. Let $H_\phi(\xi)$ be the Hessian matrix of ϕ . By [de Bondt and van den Essen 2004, Theorem 3.1], $H_\phi(\xi)$ is not identically zero. Consider

$$Z_\phi := \{\xi \in \mathbb{R}^2 : \det(H_\phi(\xi)) = 0\}. \tag{C-4}$$

Since $\det H_\phi(\xi)$ is not identically zero, it is a zero set of a polynomial. So there exists a square τ not intersecting Z_ϕ . Since $H_\phi(\xi)$ is independent of the parameter δ , we may assume that the size of τ is independent of δ . Let us fix such a τ . Assume that δ is smaller than the sidelength of τ . Let \hat{f} be a smooth bump function of the δ -neighborhood of

$$\{(\xi_1, \xi_2, \phi(\xi_1, \xi_2)) : \xi \in \tau\}. \tag{C-5}$$

One can see that if $\xi \in \tau$, then

$$|\det H_\phi(\xi)| \gtrsim 1. \tag{C-6}$$

So the multiplication of eigenvalues of $H_\phi(\xi)$ does not change the sign over $\xi \in \tau$.

Suppose that the eigenvalues of $H_\phi(\xi)$ have the same sign for all $\xi \in \tau$. Then each $S \in \mathcal{S}_\delta$ intersecting τ is a square with length $\delta^{1/2} \times \delta^{1/2}$. So we have

$$f = \sum_S f_S, \quad |f_S(x)| = \delta^2 1_{T_S}, \tag{C-7}$$

where T_S is a tube with dimension $\delta^{-1/2} \times \delta^{-1/2} \times \delta^{-1}$. Then

$$\delta^2 \delta^{-1} \lesssim \|f\|_{L^p(B_1)} \lesssim \|f\|_{L^p(\mathbb{R}^3)}. \tag{C-8}$$

On the other hand,

$$\left(\sum_S \|f_S\|_{L^p}^2\right)^{1/2} \lesssim \delta^{-1/2} \delta^2 |T_S|^{1/p} \sim \delta^{-1/2} \delta^2 \delta^{-2/p}. \tag{C-9}$$

This shows that the decoupling inequality is false for $p > 4$.

Suppose that the eigenvalues have different signs. Then, as in the previous case, we have

$$\delta \lesssim \|f\|_{L^p(\mathbb{R}^3)}. \quad (\text{C-10})$$

We need to calculate

$$\left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^p}^2 \right)^{1/2}. \quad (\text{C-11})$$

By the construction of \mathcal{S}_δ , we have $|f_S| = \delta^2 1_{T_S}$, where T_S has dimension $\delta^A \times \delta^{1-A} \times \delta$ for some $0 \leq A \leq 1$. Here the number A depends on the choice of S . Also, the cardinality of \mathcal{S}_δ is comparable to δ^{-1} up to $\delta^{-\epsilon}$ losses. So we have

$$\left(\sum_{S \in \mathcal{S}_\delta} \|f_S\|_{L^p}^2 \right)^{1/2} \lesssim_{\epsilon} \delta^{-\epsilon} \delta^{-1/2} \delta^2 |T_S|^{1/p} \sim \delta^{-\epsilon} \delta^{-1/2} \delta^2 \delta^{-2/p}. \quad (\text{C-12})$$

This shows that the decoupling inequality is false for $p > 4$, and completes the proof of item (1).

Let us move on to item (2). Since ϕ depends only on one variable, after some change of variables and abusing notation, we may assume that $\phi(\xi_1, \xi_2) = \psi(\xi_1)$. Fix x_2 , and define

$$g(x_1, x_3) := f(x_1, x_2, x_3). \quad (\text{C-13})$$

Then (1-4) simply follows from [Yang 2021, Theorem 1.4]. Let us show the sharpness. Since ψ is a one-variable polynomial, the zeros of ψ are finite. So we can find an open interval $I \subset [0, 1]$ such that, for every $\xi_1 \in I$, we have $|\psi''(\xi_1)| \gtrsim 1$. We write I as a union of intervals J with length $\delta^{1/2}$. Take \hat{f} to be a smooth bump function of the δ -neighborhood of

$$\{(\xi_1, \xi_2, \phi(\xi_1, \xi_2)) : \xi_1 \in I, \xi_2 \in [0, 1]\}. \quad (\text{C-14})$$

Then similar calculations in the proof of item (1) give the sharpness for p . We leave out the details.

Lastly, let us prove item (3). If ϕ does not depend on any variable, then, according to our construction, $f_S = f$. So (1-4) is true for $2 \leq p < \infty$. This completes the proof of Proposition C.2. \square

Acknowledgements

The authors would like to thank Yuqiu Fu for discussion at the early stage of the project. Guth is supported by a Simons Investigator award. Maldague is supported by the National Science Foundation under award no. 2103249. Oh was supported by the NSF grant DMS-1800274 and the POSCO Science Fellowship of POSCO TJ Park Foundation.

References

- [de Bondt and van den Essen 2004] M. de Bondt and A. van den Essen, “Singular Hessians”, *J. Algebra* **282**:1 (2004), 195–204. [MR](#)
- [Bourgain and Demeter 2015] J. Bourgain and C. Demeter, “The proof of the ℓ^2 decoupling conjecture”, *Ann. of Math. (2)* **182**:1 (2015), 351–389. [MR](#)
- [Bourgain and Demeter 2017a] J. Bourgain and C. Demeter, “Decouplings for curves and hypersurfaces with nonzero Gaussian curvature”, *J. Anal. Math.* **133** (2017), 279–311. [MR](#)

- [Bourgain and Demeter 2017b] J. Bourgain and C. Demeter, “A study guide for the l^2 decoupling theorem”, *Chinese Ann. Math. Ser. B* **38**:1 (2017), 173–200. [MR](#)
- [Bourgain and Guth 2011] J. Bourgain and L. Guth, “Bounds on oscillatory integral operators based on multilinear estimates”, *Geom. Funct. Anal.* **21**:6 (2011), 1239–1295. [MR](#)
- [Bourgain et al. 2016] J. Bourgain, C. Demeter, and L. Guth, “Proof of the main conjecture in Vinogradov’s mean value theorem for degrees higher than three”, *Ann. of Math. (2)* **184**:2 (2016), 633–682. [MR](#)
- [Bourgain et al. 2020] J. Bourgain, C. Demeter, and D. Kemp, “Decouplings for real analytic surfaces of revolution”, pp. 113–125 in *Geometric aspects of functional analysis, I*, edited by B. Klartag and E. Milman, Lecture Notes in Math. **2256**, Springer, 2020. [MR](#)
- [Buschenhenke et al. 2020] S. Buschenhenke, D. Müller, and A. Vargas, “A Fourier restriction theorem for a perturbed hyperbolic paraboloid”, *Proc. Lond. Math. Soc. (3)* **120**:1 (2020), 124–154. [MR](#)
- [Buschenhenke et al. 2022] S. Buschenhenke, D. Müller, and A. Vargas, “A Fourier restriction theorem for a perturbed hyperbolic paraboloid: polynomial partitioning”, *Math. Z.* **301**:2 (2022), 1913–1938. [MR](#)
- [Buschenhenke et al. 2023] S. Buschenhenke, D. Müller, and A. Vargas, “Fourier restriction for smooth hyperbolic 2-surfaces”, *Math. Ann.* **387**:1-2 (2023), 17–56. [MR](#)
- [Cassels 1972] J. W. S. Cassels, *An introduction to Diophantine approximation*, Cambridge Tracts in Mathematics and Mathematical Physics **45**, Hafner Publishing Co., New York, 1972. [MR](#)
- [Demeter 2020] C. Demeter, *Fourier restriction, decoupling, and applications*, Cambridge Studies in Advanced Mathematics **184**, Cambridge University Press, 2020. [MR](#)
- [Deng et al. 2017] Y. Deng, P. Germain, and L. Guth, “Strichartz estimates for the Schrödinger equation on irrational tori”, *J. Funct. Anal.* **273**:9 (2017), 2846–2869. [MR](#)
- [Deng et al. 2022] Y. Deng, P. Germain, L. Guth, and S. L. Rydin Myerson, “Strichartz estimates for the Schrödinger equation on non-rectangular two-dimensional tori”, *Amer. J. Math.* **144**:3 (2022), 701–745. [MR](#)
- [Guo and Oh 2024] S. Guo and C. Oh, “A restriction estimate for surfaces with negative Gaussian curvatures”, *Peking Math. J.* **7**:1 (2024), 155–202. [MR](#)
- [Guth et al. 2023] L. Guth, D. Maldague, and C. Oh, “Small cap decoupling for the paraboloid in \mathbb{R}^l ”, preprint, 2023. [arXiv 2307.06445](#)
- [Li and Yang 2021] J. Li and T. Yang, “Decoupling for smooth surfaces in \mathbb{R}^3 ”, preprint, 2021. [arXiv 2110.08441](#)
- [Mizutani and Tzvetkov 2015] H. Mizutani and N. Tzvetkov, “Strichartz estimates for non-elliptic Schrödinger equations on compact manifolds”, *Comm. Partial Differential Equations* **40**:6 (2015), 1182–1195. [MR](#)
- [Stein 1986] E. M. Stein, “Oscillatory integrals in Fourier analysis”, pp. 307–355 in *Beijing lectures in harmonic analysis* (Beijing, 1984), edited by E. M. Stein, Ann. of Math. Stud. **112**, Princeton Univ. Press, 1986. [MR](#)
- [Sun and Tzvetkov 2020] C. Sun and N. Tzvetkov, “Concerning the pathological set in the context of probabilistic well-posedness”, *C. R. Math. Acad. Sci. Paris* **358**:9-10 (2020), 989–999. [MR](#)
- [Tomas 1975] P. A. Tomas, “A restriction theorem for the Fourier transform”, *Bull. Amer. Math. Soc.* **81** (1975), 477–478. [MR](#)
- [Yang 2021] T. Yang, “Uniform l^2 -decoupling in \mathbb{R}^2 for polynomials”, *J. Geom. Anal.* **31**:11 (2021), 10846–10867. [MR](#)

Received 27 Mar 2024. Revised 6 Oct 2024. Accepted 8 Jan 2025.

LARRY GUTH: lguth@math.mit.edu

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States

DOMINIQUE MALDAGUE: dm672@cam.ac.uk

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States

Current address: Department of Mathematics, University of Cambridge, Cambridge, United Kingdom

CHANGKEUN OH: changkeun@snu.ac.kr

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States

Current address: Department of Mathematical Sciences and RIM, Seoul National University, Seoul, South Korea

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Anna L. Mazzucato Penn State University, USA
alm24@psu.edu

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Sc. Intern. Sup. di Studi Avan., Italy berti@sissa.it	Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Yu Deng	University of Chicago, USA yudeng@uchicago.edu	Omar Mohsen	Université Paris-Cité, France omar.mohsen.fr@gmail.com
Thierry Gallay	Université Grenoble Alpes, France Thierry.Gallay@univ-grenoble-alpes.fr	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Xavier Ros Oton	Catalan Inst. for Res. and Adv. Std., Spain xros@icrea.cat
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Nicolas Rougerie	ENS Lyon, France nicolas.rougerie@ens-lyon.fr
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Sebastian Herr	Universität Bielefeld, Germany herr@math.uni-bielefeld.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Jonathan Wing-hong Luk	Stanford University jluk@stanford.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rmb@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: “Linear Ramp”


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2026 is US \$500/year for the electronic version, and \$780/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 19 No. 1 2026

Microlocal analysis near null infinity in asymptotically flat spacetimes	1
PETER HINTZ and ANDRÁS VASY	
Complex Hessian measures with respect to a background Hermitian form	107
SŁAWOMIR KOŁODZIEJ and NGOC CUONG NGUYEN	
ℓ^2 decoupling theorem for surfaces in \mathbb{R}^3	167
LARRY GUTH, DOMINIQUE MALDAGUE and CHANGKEUN OH	