

ANALYSIS & PDE

Volume 19

No. 2

2026

COLETTE DE COSTER, SIMONE DOVETTA, DAMIEN GALANT,
ENRICO SERRA AND CHRISTOPHE TROESTLER

**CONSTANT SIGN AND SIGN CHANGING NLS GROUND STATES
ON NONCOMPACT METRIC GRAPHS**

CONSTANT SIGN AND SIGN CHANGING NLS GROUND STATES ON NONCOMPACT METRIC GRAPHS

COLETTE DE COSTER, SIMONE DOVETTA, DAMIEN GALANT,
ENRICO SERRA AND CHRISTOPHE TROESTLER

We investigate existence and nonexistence of action ground states and nodal action ground states for the nonlinear Schrödinger equation on noncompact metric graphs with mixed homogeneous Kirchhoff and Dirichlet boundary conditions. We first obtain abstract sufficient conditions for existence, typical of problems with lack of compactness, in terms of “levels at infinity” for the action functional associated with the problems. Then we analyze in detail two relevant classes of graphs. For noncompact graphs with at least one half-line, we detect purely topological sharp conditions preventing the existence of ground states or of nodal ground states. We also investigate analogous conditions of metrical nature. The negative results are complemented by several sufficient conditions to ensure existence, either of topological or metrical nature, or a combination of the two. For graphs with infinitely many edges, all bounded, we focus on periodic graphs and infinite trees. In these cases, our results completely describe the phenomenology. Furthermore, we study nodal domains and nodal sets of nodal ground states and we show that the situation on graphs can be totally different from that on domains of \mathbb{R}^N .

1. Introduction

We investigate the existence of constant sign and sign changing solutions of the nonlinear Schrödinger equation

$$u'' + |u|^{p-2}u = \lambda u, \quad (1-1)$$

where $p > 2$ and λ is a real parameter, on noncompact metric graphs under various assumptions.

Throughout this paper we consider the class \mathbf{G} of connected metric graphs $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ where the sets \mathbb{V} and \mathbb{E} are at most countable, every vertex $v \in \mathbb{V}$ has finite degree, and the lengths of the edges $e \in \mathbb{E}$ are bounded away from zero (see Definition 2.1 below). A graph of this type is noncompact if at least one edge is unbounded (i.e., it is a half-line) or if it consists of an infinite number of bounded edges, giving rise to two classes of graphs that behave quite differently and that we will treat separately. Every half-line is considered to end at a “vertex at infinity” of degree 1. The set of all such vertices of \mathbb{V} is denoted by \mathbb{V}_∞ .

The analysis of differential equations on metric graphs experienced a massive growth in recent years, in particular motivated by the potential of graphs to serve as simple models for signal propagation in branched structures. In this context, stationary nonlinear Schrödinger equations such as (1-1) gained

MSC2020: 35Q55, 35R02, 49J40, 58E30.

Keywords: nonlinear Schrödinger, ground states, nodal solutions, least action, constrained minimization.

a prominent interest, as it is well known that to any couple (u, λ) satisfying (1-1) there corresponds a standing wave solution $\psi(t, x) := e^{i\lambda t} u(x)$ of the time-dependent nonlinear Schrödinger equation

$$-i\partial_t \psi(t, x) = \partial_{xx}^2 \psi(t, x) + |\psi(t, x)|^{p-2} \psi(t, x). \quad (1-2)$$

Nonlinear dispersive equations such as (1-2) are largely studied in view of the role they play in many applications, such as, e.g., in the modeling of quantum mechanical systems in Bose–Einstein condensation or in the modeling of optical fibers.

It is clear however that, when considering any differential model on graphs, it is not enough to prescribe a differential equation that describes the behavior of the system in the interior of each edge. The equation must be complemented with suitable matching conditions at the vertices, specifying how the interaction among edges behaves at the junctions. In the case of nonlinear Schrödinger equations, there is a wide class of vertex conditions that can be considered. In the present paper, we couple (1-1) with a specific choice of boundary conditions. Precisely, given a (not necessarily finite) set $Z \subseteq \mathbb{V} \setminus \mathbb{V}_\infty$ of degree-1 vertices, we are interested in solutions to the problem

$$\begin{cases} u'' + |u|^{p-2}u = \lambda u & \text{on every edge of } \mathcal{G}, \\ u \text{ is continuous} & \text{on } \mathcal{G}, \\ \sum_{e \succ v} u'_e(v) = 0 & \text{for every } v \in \mathbb{V} \setminus (Z \cup \mathbb{V}_\infty), \\ u(v) = 0 & \text{for every } v \in Z, \end{cases} \quad (1-3)$$

where $u'_e(v)$ is the outgoing derivative along the edge e incident at the vertex v and $e \succ v$ means that the sum is extended to all such edges. The boundary condition for $v \notin Z$ (together with the continuity of u) is the homogeneous Kirchhoff condition, by far the most used in the literature. It is a natural analogue of the Neumann boundary condition for metric graphs (see, e.g., [Berkolaiko and Kuchment 2013, Section 1.4]). The boundary condition for $v \in Z$ is the homogeneous Dirichlet condition, which by contrast has been discussed only in a few papers (see, for instance, [Esteban 2022]). Here we choose to include mixed conditions to highlight their role in the existence (or nonexistence) of various types of solutions to (1-3). The requirement that all nodes of Z have degree 1 prevents the graph from being disconnected by the Dirichlet conditions, but more general frameworks can easily be treated building on the results of this paper.

Coupling the operator $-d^2/dx^2$ with our boundary conditions guarantees its self-adjointness. This is a natural requirement in the analysis of quantum mechanical problems (see, e.g., [Adami et al. 2020] for an overview of boundary conditions ensuring self-adjointness on metric graphs and [Berkolaiko and Kuchment 2013, Section 1.4] for a thorough discussion).

Solutions to (1-3) can be found by a variational approach that has been employed very frequently to deal with this kind of problem or with its variants. In our setting, the appropriate function space to set problem (1-3) is

$$H_Z^1(\mathcal{G}) := \{u \in H^1(\mathcal{G}) \mid u(v) = 0 \text{ for every } v \in Z\}.$$

Standard arguments show that the $H^1(\mathcal{G})$ solutions of problem (1-3) are exactly the critical points of the action functional $J_\lambda : H_Z^1(\mathcal{G}) \rightarrow \mathbb{R}$ defined by

$$J_\lambda(u) := \frac{1}{2} \|u'\|_{L^2(\mathcal{G})}^2 + \frac{1}{2} \lambda \|u\|_{L^2(\mathcal{G})}^2 - \frac{1}{p} \|u\|_{L^p(\mathcal{G})}^p, \quad (1-4)$$

that is of class C^2 on $H_Z^1(\mathcal{G})$. Hereafter the parameter λ satisfies as usual $\lambda > -\omega_Z(\mathcal{G})$, where

$$\omega_Z(\mathcal{G}) := \inf_{v \in H_Z^1(\mathcal{G}) \setminus \{0\}} \frac{\|v'\|_{L^2(\mathcal{G})}^2}{\|v\|_{L^2(\mathcal{G})}^2}$$

denotes the bottom of the spectrum of the Laplacian on \mathcal{G} associated to the boundary conditions in (1-3). This assumption is standard when working with this problem and is justified by the fact that, under it, the quadratic part in (1-4) provides a norm on $H^1(\mathcal{G})$ equivalent to the usual H^1 -norm.

When looking at solutions to (1-3) from the variational perspective, one has to take into account that the functional J_λ is not bounded from below on $H_Z^1(\mathcal{G})$. A standard way to recover the notion of minimality is to introduce the Nehari manifold associated to J_λ , defined by

$$\begin{aligned} \mathcal{N}_{\lambda,Z}(\mathcal{G}) &:= \{u \in H_Z^1(\mathcal{G}) \mid u \neq 0, J'_\lambda(u)u = 0\} \\ &= \{u \in H_Z^1(\mathcal{G}) \mid u \neq 0, \|u'\|_{L^2(\mathcal{G})}^2 + \lambda\|u\|_{L^2(\mathcal{G})}^2 = \|u\|_{L^p(\mathcal{G})}^p\}. \end{aligned}$$

Clearly, $\mathcal{N}_{\lambda,Z}(\mathcal{G})$ contains all solutions to (1-3). It is a C^1 -manifold diffeomorphic to the unit sphere of $H_Z^1(\mathcal{G})$ and is a natural constraint for J_λ , in the sense that constrained critical points of J_λ are in fact true critical points. Other approaches are possible (for instance, one could constrain J_λ on the unit sphere of $L^p(\mathcal{G})$), but the Nehari approach has the advantage that it works also in cases where the nonlinearity is not homogeneous, thus providing a framework suitable to be generalized to a wider class of nonlinear terms.

Definition 1.1. We say that a function $u \in \mathcal{N}_{\lambda,Z}(\mathcal{G})$ is a *ground state* for problem (1-3) if

$$J_\lambda(u) = \inf_{v \in \mathcal{N}_{\lambda,Z}(\mathcal{G})} J_\lambda(v).$$

Since we want to discuss both one-signed and sign changing solutions, we will also consider *nodal ground states*, roughly the analogue of ground states among sign changing functions. To define rigorously this notion, we let

$$u^+ := \max(u, 0), \quad u^- := \min(u, 0)$$

and define the *nodal Nehari set* as

$$\mathcal{M}_{\lambda,Z}(\mathcal{G}) := \{u \in H_Z^1(\mathcal{G}) \mid u^\pm \in \mathcal{N}_{\lambda,Z}(\mathcal{G})\} = \{u \in H_Z^1(\mathcal{G}) \mid u^\pm \neq 0, J'_\lambda(u)u^\pm = 0\}.$$

The nodal Nehari set contains all nodal solutions of (1-3) but, contrary to $\mathcal{N}_{\lambda,Z}(\mathcal{G})$, in general is not a manifold (see, e.g., [Bartsch and Weth 2003; Castro et al. 1997; Szulkin and Weth 2010]) and is not a natural constraint for J_λ , which causes some extra difficulties when proving existence results.

Definition 1.2. We say that a function $u \in \mathcal{M}_{\lambda,Z}(\mathcal{G})$ is a *nodal ground state* for problem (1-3) if

$$J_\lambda(u) = \inf_{v \in \mathcal{M}_{\lambda,Z}(\mathcal{G})} J_\lambda(v).$$

It is well known that, whenever they exist, ground states (resp. nodal ground states) provide constant sign solutions (resp. sign changing solutions) of (1-1) of minimal action. Actually, to look for one-signed solutions of nonlinear Schrödinger equations as minimizers of suitable functionals is a standard strategy,

that has been widely exploited on graphs in the mass constrained setting, where ground states are defined as minimizers of the energy functional $u \mapsto \frac{1}{2} \|u'\|_{L^2(\mathcal{G})}^2 - \frac{1}{p} \|u\|_{L^p(\mathcal{G})}^p$ restricted to an L^2 -sphere (see, e.g., [Adami et al. 2014a; 2015; 2016; 2019; 2022; Berkolaiko et al. 2021; Boni and Carlone 2023; Besse et al. 2022a; 2022b; Boni and Dovetta 2021; 2022; Dovetta and Tentarelli 2022; Kairzhan et al. 2021; 2022; Pierotti and Soave 2022; Pierotti et al. 2021; Tentarelli 2016]). In particular, these works have shown that the existence of ground states for the prescribed-mass problem on noncompact graphs is a rather unlikely event. Obstructions to existence are provided mostly by the topology of the graph, and sometimes also by its metrical properties. Conversely, the action approach has not received much attention so far (some results in this direction can be found, e.g., in [Adami et al. 2014b; De Coster et al. 2023; Kurata and Shibata 2020; Pankov 2018]). Let us stress that, though clearly critical points of the action on Nehari sets are also critical points of the energy on a suitable L^2 -sphere and vice versa, the general relation between the action and the energy approach is not fully understood. First investigations in this direction started only recently in [Dovetta et al. 2023; Jeanjean and Lu 2022] for NLS equations posed on domains of \mathbb{R}^N , but many of the results obtained therein extend with no difference to metric graphs. Specifically, if on the one hand those analyses proved that mass constrained ground states of the energy are always also ground states of the action, on the other hand they showed that the converse is in general not true. Hence, there may well exist action ground states that are not energy ground states among functions with the same mass, and the actual occurrence of this phenomenon on metric graphs has been proved, e.g., in [Agostinho et al. 2024, Theorem 1.4; Dovetta 2024, Proposition 2.4]. This somehow further motivates the independent study of the action ground state problem even in a context, as that of graphs, in which a well-developed theory of energy ground states is already available. In addition, to the best of our knowledge, nodal ground states, and more generally sign changing solutions, on general metric graphs have never been investigated up to now, neither for the action nor for the energy problem. In the nodal setting, actually, the asymmetry between the action and the energy point of views is more pronounced, as it is not even clear how to define a general variational framework to deal with sign changing solutions with prescribed mass (that is, no analogue of the nodal Nehari set is known for the energy).

The main concern of the present paper is thus to begin a systematic study of action ground states and nodal ground states for (1-3), characterizing the dependence of the problem on topological and metrical properties of the graph. On one side, as it is reasonable to expect, with purely Kirchhoff vertex conditions (i.e., $Z = \emptyset$), we will find that sometimes the same topological conditions that rule out mass constrained ground states do prevent existence of action ground states too (this is the case for graphs with half-lines, as in Theorem 1.6 below), and analogous conditions for nodal ground states will be identified. On the other side, we will also show that there are graphs (periodic ones and trees) for which action ground states exist for every admissible value of λ , whereas existence of mass constrained ground states depends on the value of the mass.

To begin the discussion of our results, we observe that if \mathcal{G} is compact, the existence of a ground state and of a nodal ground state can be proved via standard compactness arguments. Indeed, the compactness of the domain guarantees compactness for Sobolev embeddings $H^1(\mathcal{G}) \hookrightarrow L^p(\mathcal{G})$, which is all that is needed to obtain strong convergence of minimizing sequences for the above variational

problems (see, e.g., [Dovetta 2018, Proposition 3.1] for further details on such compactness results in the context of mass constrained critical points of the energy).

On the contrary, if \mathcal{G} is noncompact, since as we said above the existence of ground states is unlikely, it is not surprising that the analogous eventuality for nodal ground states is even more so. As for ground states, we are going to derive sufficient conditions for both existence and nonexistence of nodal ground states involving topological features, metrical ones and combinations of the two.

Our analysis is based on a rather abstract procedure, typical of problems with lack of compactness, consisting in locating thresholds on the levels of J_λ , involving the so-called level at infinity

$$J_\lambda^\infty(\mathcal{G}; Z) := \inf\left\{\liminf_n J_\lambda(u_n) \mid (u_n)_n \subseteq \mathcal{N}_{\lambda, Z}(\mathcal{G}), \lim_n u_n = 0 \text{ weakly in } H_Z^1(\mathcal{G})\right\}.$$

Theorem 1.3. *Let $\mathcal{G} \in \mathbf{G}$ be a noncompact graph and $\lambda > -\omega_Z(\mathcal{G})$.*

(i) *If*

$$\inf_{v \in \mathcal{N}_{\lambda, Z}(\mathcal{G})} J_\lambda(v) < J_\lambda^\infty(\mathcal{G}; Z), \quad (1-5)$$

then there exists a ground state of J_λ in $\mathcal{N}_{\lambda, Z}(\mathcal{G})$. Moreover, ground states have constant sign on $\mathcal{G} \setminus Z$.

(ii) *If*

$$\inf_{v \in \mathcal{M}_{\lambda, Z}(\mathcal{G})} J_\lambda(v) < J_\lambda^\infty(\mathcal{G}; Z) + \inf_{v \in \mathcal{N}_{\lambda, Z}(\mathcal{G})} J_\lambda(v), \quad (1-6)$$

then there exists a nodal ground state of J_λ in $\mathcal{M}_{\lambda, Z}(\mathcal{G})$.

Theorem 1.4. *For every noncompact graph $\mathcal{G} \in \mathbf{G}$ and $\lambda > -\omega_Z(\mathcal{G})$,*

$$\inf_{v \in \mathcal{M}_{\lambda, Z}(\mathcal{G})} J_\lambda(v) \geq 2 \inf_{v \in \mathcal{N}_{\lambda, Z}(\mathcal{G})} J_\lambda(v). \quad (1-7)$$

If equality holds, then \mathcal{G} admits no nodal ground states of J_λ in $\mathcal{M}_{\lambda, Z}(\mathcal{G})$.

Remark 1.5. Inequality (1-7) also holds when \mathcal{G} is a compact metric graph, and is then strict as nodal ground states always exist in this case.

This abstract strategy, though general, is absolutely insufficient to obtain existence results if one is not able to compute the level $J_\lambda^\infty(\mathcal{G}; Z)$ in concrete cases. Here we will detect specific properties of the graph that permit such computation and make sure that, in certain cases, the ground state level or the nodal ground state level lie below the level at infinity. Since this is where the topology and the metric of the graph become crucial, the analysis of such questions is carried out separately according to the class of graphs under study.

We first discuss the case of graphs with at least one half-line. For every such graph, $\omega_Z(\mathcal{G}) = 0$ and so the following results hold for every $\lambda > 0$.

We identify topological conditions on \mathcal{G} that prevent the existence of ground states and nodal ground states. We describe them here in a concise way, referring to Section 4 for a more detailed discussion. To begin with, recall that the set of *vertices at infinity* of \mathcal{G} is

$$\mathbb{V}_\infty = \{v \in \mathbb{V} \mid v \text{ is the vertex of degree 1 of some half-line}\}$$




Figure 1. Examples of graphs \mathcal{G} with corresponding set $F(\mathcal{G})$ containing 0 (A), 1 (B), and 4 (C) edges, respectively. Here, Kirchhoff conditions are assumed at every vertex, and edges in $F(\mathcal{G})$ are drawn thicker.

and define the set

$$F(\mathcal{G}) = \{e \in \mathbb{E} \mid \text{at least one connected component of } (\mathbb{V}, \mathbb{E} \setminus \{e\}) \text{ has no vertices in } \mathbb{V}_\infty \cup Z\}.$$

The set $F(\mathcal{G})$ is thus the set of edges of \mathcal{G} (if any) whose removal disconnects \mathcal{G} creating a connected component without vertices in $\mathbb{V}_\infty \cup Z$. Basically, if $F(\mathcal{G})$ is nonempty, there exists at least one “bridging” edge in the graph that, once removed, creates at least one connected component separated from all the half-lines and all the vertices with homogeneous Dirichlet condition (see Figure 1 for a concrete illustration of $F(\mathcal{G})$ on different graph structures). The cardinality of $F(\mathcal{G})$ is a purely topological notion and plays a fundamental role in the nonexistence of ground states and nodal ground states. Roughly, we will see that the presence of bridging edges in $F(\mathcal{G})$ may facilitate existence of such states, especially if the connected components without vertices in $\mathbb{V}_\infty \cup Z$ they isolate have a very simple structure. On the contrary, a low cardinality of $F(\mathcal{G})$ somehow corresponds to a too-intricate graph structure not compatible with existence. This is stated rigorously in the next theorem, where for every $\lambda > 0$ we denote by

$$s_\lambda := \inf_{v \in \mathcal{N}_\lambda(\mathbb{R})} J_\lambda(v)$$

the ground state level of J_λ on \mathbb{R} .

Theorem 1.6. *Let $\mathcal{G} \in \mathcal{G}$ be a noncompact graph with at least one half-line and $\lambda > 0$. Then*

$$\inf_{v \in \mathcal{N}_{\lambda,Z}(\mathcal{G})} J_\lambda(v) \leq s_\lambda \quad (1-8)$$

and

$$\inf_{v \in \mathcal{M}_{\lambda,Z}(\mathcal{G})} J_\lambda(v) \leq s_\lambda + \inf_{v \in \mathcal{N}_{\lambda,Z}(\mathcal{G})} J_\lambda(v). \quad (1-9)$$

Moreover,

(i) if $\#F(\mathcal{G}) = 0$, then

$$\inf_{v \in \mathcal{N}_{\lambda,Z}(\mathcal{G})} J_\lambda(v) = s_\lambda \quad (1-10)$$

and it is not achieved, unless \mathcal{G} is isometric to \mathbb{R} or to a “tower of bubbles” depicted in Figure 2;

(ii) if $\#F(\mathcal{G}) \leq 1$, then

$$\inf_{v \in \mathcal{M}_{\lambda,Z}(\mathcal{G})} J_\lambda(v) = s_\lambda + \inf_{v \in \mathcal{N}_{\lambda,Z}(\mathcal{G})} J_\lambda(v) \quad (1-11)$$

and it is never achieved.




Figure 2. An example of a “tower of bubbles” graph. Each of these graphs, identified in Example 2.4 of [Adami et al. 2015], is built of a real line and a finite sequence of two-by-two tangent circles.

The preceding results are the main examples where a purely topological assumption on the graph rules out the existence of ground states or nodal ground states. The families of graphs fulfilling each of the conditions of Theorem 1.6 are rather large and it is not difficult to exhibit examples of structures with these properties (see Figure 1(A)–(B) and Figure 3). In the case of ground states, the condition $\#F(\mathcal{G}) = 0$ was already shown to prevent existence of mass constrained ground states of the energy in [Adami et al. 2015, Theorem 2.5], where it was named assumption (H). In contrast, the analogous condition for nodal ground states is established here for the first time. We underline that both assumptions on the cardinality of $F(\mathcal{G})$ are sharp for nonexistence. Indeed, in Section 4 we will show that there exist graphs satisfying $\#F(\mathcal{G}) \geq 1$ that admit a ground state, and graphs satisfying $\#F(\mathcal{G}) \geq 2$ that admit a nodal ground state.

These nonexistence results are complemented in Section 4 by a number of sufficient conditions to ensure existence. Relying on techniques developed for energy ground states in [Adami et al. 2015, Section 6; 2016, Section 3], it is easy to construct graphs where existence of action ground states is guaranteed by purely *topological* properties whenever $Z = \emptyset$ (see Theorem 4.8 and Figure 6 below). Notably, this turns out to be impossible as soon as $Z \neq \emptyset$. In this case, a necessary condition of *metrical* nature for the existence of ground states arises: the diameter of the set of all bounded edges of the graph must exceed a threshold depending on λ but not on the graph itself (Theorem 4.9). The same constraint holds true for nodal ground states, where it is not even needed to have a nonempty set Z (Theorem 4.11). In addition to providing purely metrical nonexistence results, these theorems imply that the interplay between topology and metric must be further exploited if one hopes to recover existence. We give examples of this fact by describing two general procedures to construct graphs where existence is granted (Theorems 4.12–4.16). The former relies on the metric only, and shows that one and two sufficiently long edges with vertices of degree 1 not in Z are enough to have ground states and nodal ground states, respectively. The latter basically consists in a suitable gluing of graphs hosting ground states to obtain structures supporting nodal ground states (see, e.g., Figure 7).




Figure 3. Further examples of graphs with half-lines satisfying $\#F(\mathcal{G}) = 0$ (A) and $\#F(\mathcal{G}) = 1$ (B).

The previous results highlight how the presence of at least one vertex with homogeneous Dirichlet conditions affects the existence of ground states and nodal ground states. Indeed, the fact that vertices in Z play the same role as those at infinity in the definition of $F(\mathcal{G})$ suggests the idea that an edge ending at a vertex with zero Dirichlet conditions behaves as a half-line. This heuristic comparison makes some sense, since H^1 functions tend to zero at infinity along each half-line. However, Theorem 4.9 unravels that the analogy with half-lines is not complete: edges with endpoints in Z are somehow “worse”, as they make metrical assumptions necessary to obtain existence. We may be tempted to say that existence of ground states simply requires edges ending in Z to be long enough (and thus sufficiently close to half-lines), but this is not true in general, as ground states can exist even on graphs where all edges with vertices in Z are arbitrarily short (see Remark 4.10 below).

Section 5 of the paper deals with noncompact graphs in the class \mathbf{G} with an infinite number of edges, whose length is uniformly bounded from above. Given the huge variety of structures in this class, we confine ourselves to two subclasses of major relevance, that have already been studied extensively in the literature in many contexts (see, e.g., [Adami et al. 2019; Besse et al. 2022a; 2022b; Dovetta and Tentarelli 2022; Dovetta et al. 2020; Gilg et al. 2022; Pankov 2018; Pelinovsky and Schneider 2017] for results related to those we discuss here): periodic graphs and regular trees.

Without entering the details of the definition of periodic graphs (for which we refer to [Berkolaiko and Kuchment 2013, Definition 4.1.1]), let us mention here that we always work assuming that the set Z shares the same type of periodicity as the graph itself. Our main result in this respect completely describes the phenomenology from the point of view of ground states and nodal ground states (results in this direction for ground states on periodic graphs were already given in [Pankov 2018]).

As for graphs with half-lines, if \mathcal{G} is a periodic graph then $\omega_Z(\mathcal{G}) = 0$, so that the next theorem holds again for every $\lambda > 0$.

Theorem 1.7. *Let $\mathcal{G} \in \mathbf{G}$ be a periodic graph and $\lambda > 0$. Then \mathcal{G} admits a ground state. Furthermore,*

$$\inf_{v \in \mathcal{M}_{\lambda, Z}(\mathcal{G})} J_{\lambda}(v) = 2 \inf_{v \in \mathcal{N}_{\lambda, Z}(\mathcal{G})} J_{\lambda}(v)$$

and there are no nodal ground states.

It is interesting to note that the above results are insensitive of the degree of periodicity, i.e., the dimension n of the group \mathbb{Z}^n whose action leaves the graph invariant. This is particularly relevant when put in relation with the available results for prescribed-mass energy ground states (compare, e.g., [Adami et al. 2019, Theorems 1.1–1.2], where $n = 2$, with [Dovetta 2019, Theorem 1.1], where $n = 1$), whose existence has been shown to depend strongly on the value of n .

The last results of Section 5 concern regular trees, i.e., acyclic, noncompact metric graphs with infinitely many bounded edges, all of the same length, and where all the vertices have the same degree, with the possible exception of a unique root vertex of degree 1. If such a vertex with degree 1 is present, we speak of a rooted tree (see Figure 4(A)), otherwise we speak of an unrooted tree (see Figure 4(B)). Regular trees are well-known examples of noncompact graphs satisfying $\omega_Z(\mathcal{G}) > 0$ (see, e.g., [Dovetta et al. 2020]). Hence, in this setting our results involve also negative values of λ .




Figure 4. Examples of a rooted tree (A) and an unrooted tree (B).

Theorem 1.8. *Let \mathcal{G} be a regular tree and $\lambda > -\omega_Z(\mathcal{G})$. Then*

- (i) *if \mathcal{G} is unrooted or if \mathcal{G} is rooted and Z is empty, \mathcal{G} admits a ground state;*
- (ii) *if \mathcal{G} is rooted and Z is not empty, there are no ground states on \mathcal{G} ;*
- (iii) *independently of Z , there are no nodal ground states on \mathcal{G} .*

As in the case of periodic graphs, the above theorem provides a complete description of the problem for regular trees. As one may expect, the role of the set Z is crucial to discriminate between existence and nonexistence on rooted trees. Moreover, as already seen for periodic graphs, observe that Theorem 1.8(i) establishes the existence of ground states of the action on trees for every admissible value of the parameter λ . It is interesting to compare this result with [Dovetta et al. 2020, Theorem 1.2], where it is shown that existence of mass constrained ground states of the energy on trees does depend on the mass. In particular, when $p \in [4, 6)$, energy ground states do not exist if the mass is smaller than a positive threshold. However, this seeming asymmetry is not sufficient to guarantee that, on trees, the action approach is more general than the energy one. Indeed, [Dovetta et al. 2020] does not provide any information on the values of the parameter λ associated to energy ground states, and nothing is known about the mass of action ground states identified in this paper. Hence, it is not clear whether the ground states of the action given by Theorem 1.8(i) for $\lambda > -\omega_Z(\mathcal{G})$ coincide with the mass constrained ground states of the energy found in [Dovetta et al. 2020, Theorem 1.2].

We observe that the discussion developed here requires no restrictions on the nonlinearity power p , so that all our results apply for every $p > 2$. In particular, the existence statements listed above provide constant sign and sign changing solutions to (1-3) also when $p > 6$, the so-called L^2 -supercritical regime, whose analysis is much harder in the context of fixed mass critical points of the energy (first investigations in this direction have been initiated in [Borthwick et al. 2023; Chang et al. 2024]).

To conclude, we study nodal domains (i.e., the connected components of $\mathcal{G} \setminus u^{-1}(0)$) and the nodal set (i.e., the set $u^{-1}(0)$) of nodal ground states u . As one may expect, nodal ground states have exactly

two nodal domains (Theorem 6.1). We also show that the nodal set can have an arbitrary number of components and an arbitrary measure. This is in contrast with the case of open domains of \mathbb{R}^N , where unique continuation principles forbid nonzero solutions to vanish on nonempty open subsets.

Theorem 1.9. *For every $k, m, n \in \mathbb{N}$ with $m \geq 2$, there exists a graph \mathcal{G} and a nodal ground state u on \mathcal{G} such that $u^{-1}(0)$ is the union of k isolated points, m half-lines and n line segments.*

The remainder of the paper is organized as follows. Section 2 collects some preliminary facts useful for the subsequent analysis, while Section 3 provides the proof of the abstract results contained in Theorems 1.3–1.4. Section 4 analyzes the case of graphs with half-lines, while periodic graphs and trees are dealt with in Section 5. Qualitative properties of nodal ground states are studied in Section 6.

Notation. Throughout, we will drop the dependence of $\mathcal{N}_{\lambda,Z}(\mathcal{G})$, $\mathcal{M}_{\lambda,Z}(\mathcal{G})$, J_λ on λ and \mathcal{G} , writing \mathcal{N}_Z , \mathcal{M}_Z and J whenever possible, keeping the complete notation only if necessary. Similarly, when the context permits it, we will not explicitly indicate, in norms, the dependence on the domain of integration. Furthermore, when $Z = \emptyset$, we do not put \emptyset as a subscript and simply write $H^1(\mathcal{G})$, \mathcal{N} , \mathcal{M} , etc.

2. Preliminaries

For the precise notion of metric graphs, we refer to [Berkolaiko and Kuchment 2013]. However, we make precise in the following definition the class of graphs that we consider in this paper.

Definition 2.1. We denote by \mathbf{G} the class of metric graphs $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ such that

- \mathcal{G} is connected and has at most countable number of edges;
- $\deg(v) < \infty$ for every $v \in \mathbb{V}$, where $\deg(v)$ denotes the degree of the vertex v , i.e., the number of edges incident at v ;
- $\ell := \inf_{e \in \mathbb{E}} \ell_e > 0$, where ℓ_e denotes the length of the edge e .

A graph $\mathcal{G} \in \mathbf{G}$ is noncompact as soon as one of the following two eventualities occurs:

- (i) \mathcal{G} has at least one unbounded edge (i.e., a half-line),
- (ii) the number of edges of \mathcal{G} is infinite.

Remark 2.2. One could add in the definition of \mathcal{G} the assumption that every vertex v satisfies $\deg(v) \neq 2$. Indeed, vertices v of degree 2 can a priori be eliminated from any metric graph, by melting the two edges incident at v into a single edge. In some cases however (see Remark 3.2) the possibility of using vertices of degree 2 turns out to be quite handy. Adding or removing vertices of degree 2 from a graph changes it combinatorially, but not as a metric space, and in this paper we will identify graphs that differ only by vertices of degree 2.

As anticipated in the Introduction, we couple (1-1) with mixed Kirchhoff and Dirichlet boundary conditions. Given a noncompact graph $\mathcal{G} \in \mathbf{G}$, we let $Z \subset \mathbb{V}$ denote a set of vertices of degree 1 (possibly empty or infinite) where we impose homogeneous Dirichlet conditions and we set

$$H_Z^1(\mathcal{G}) := \{u \in H^1(\mathcal{G}) \mid u(v) = 0 \text{ for every } v \in Z\}.$$

The Nehari manifold associated to J on $H_Z^1(\mathcal{G})$ is

$$\begin{aligned}\mathcal{N}_Z &:= \{u \in H_Z^1(\mathcal{G}) \mid u \neq 0, J'(u)u = 0\} \\ &= \{u \in H_Z^1(\mathcal{G}) \mid u \neq 0, \|u'\|_2^2 + \lambda\|u\|_2^2 = \|u\|_p^p\},\end{aligned}$$

while the nodal Nehari set is

$$\mathcal{M}_Z := \{u \in H_Z^1(\mathcal{G}) \mid u^\pm \in \mathcal{N}_Z\} = \{u \in H_Z^1(\mathcal{G}) \mid u^\pm \neq 0, J'(u)u^\pm = 0\}.$$

The nodal Nehari set contains all nodal solutions of (1-3), but, generally, it is not a manifold. However, the following fundamental property holds for global minimizers on compact graphs.

Proposition 2.3. *Let $\mathcal{G} \in \mathbf{G}$ be compact and $\lambda > -\omega_Z(\mathcal{G})$. If $u \in \mathcal{M}_Z$ satisfies*

$$J(u) = \inf_{v \in \mathcal{M}_Z} J(v),$$

then $J'(u) = 0$.

Proof. The fact that any function realizing the minimum of the action restricted to its nodal Nehari set is in fact a critical point of the action is a very general property, that holds true for a large class of NLS equations including the one we consider in this paper, and is not specific of graphs. A detailed proof can be found, e.g., in [Bartsch et al. 2005, Proposition 3.1] in the case of NLS equations (with more general nonlinearities than the one of this paper) posed on bounded domains of \mathbb{R}^N with homogeneous Dirichlet conditions at the boundary. The proof reported therein uses only abstract tools of critical point theory (in particular, the deformation lemma). For this reason, that argument extends with no modification to compact graphs. \square

Obviously $\mathcal{M}_Z \subset \mathcal{N}_Z$ and, for $u \in \mathcal{N}_Z$, the functional J defined in (1-4) takes the simple form

$$J(u) = \kappa \|u\|_p^p = \kappa (\|u'\|_2^2 + \lambda \|u\|_2^2), \quad \kappa := \frac{1}{2} - \frac{1}{p}, \quad (2-1)$$

from which we see that J is positive on \mathcal{N}_Z . Actually much more can be said, as stated in the next proposition, which rephrases in the present setting an analogous result of [De Coster et al. 2023, Proposition 2.3].

Proposition 2.4. *For every $\lambda > -\omega_Z(\mathcal{G})$ and $p > 2$, there exists a constant $C > 0$ depending only on λ and p such that, for all noncompact $\mathcal{G} \in \mathbf{G}$,*

$$\inf_{u \in \mathcal{N}_Z} \|u\|_p \geq C > 0.$$

Moreover, if $(u_n)_n \subset \mathcal{N}_Z$ satisfies $\sup_n J(u_n) < \infty$, then $(u_n)_n$ is bounded in $H^1(\mathcal{G})$ and

$$\inf_n \|u_n\|_2 > 0, \quad \inf_n \|u_n\|_\infty > 0.$$

As is well known, there is a natural continuous projection $\pi_\lambda : H_Z^1(\mathcal{G}) \setminus \{0\} \rightarrow \mathcal{N}_Z$, defined by

$$\pi_\lambda(u) = n_\lambda(u)u, \quad n_\lambda(u) = \left(\frac{\|u'\|_2^2 + \lambda \|u\|_2^2}{\|u\|_p^p} \right)^{\frac{1}{p-2}}, \quad (2-2)$$

so that $u \in \mathcal{N}_Z$ if and only if $n_\lambda(u) = 1$. If $u \in H_Z^1(\mathcal{G})$ satisfies $u^\pm \neq 0$, then $\pi_\lambda(u^+) + \pi_\lambda(u^-) \in \mathcal{M}_Z$.

Remark 2.5. For every metric graph \mathcal{G} and every set Z of degree-1 vertices, the map

$$t \mapsto \inf_{v \in \mathcal{N}_{t,Z}(\mathcal{G})} J_t(v)$$

is increasing and continuous on $(-\omega_Z(\mathcal{G}), +\infty)$. This property of the action ground state level is actually general and does not depend on the fact that we are considering the problem on graphs. For this reason, we redirect the interested reader, e.g., to [Dovetta et al. 2023, Lemma 2.4] for a detailed proof (in the context of open subsets of \mathbb{R}^N).

3. Proof of the abstract results

In this section we prove the abstract results stated in Theorems 1.3–1.4. The strategy for the proof of the existence results is to construct special minimizing sequences for J on \mathcal{N}_Z or \mathcal{M}_Z , to avoid problems caused by the noncompactness of the graphs.

Remark 3.1. In this and in the next sections we will freely use the following fact: if $u \not\equiv 0$ solves problem (1-3) and $u \geq 0$ on \mathcal{G} , then $u > 0$ on \mathcal{G} , except of course at vertices in Z . For the convenience of the reader we sketch a proof (full details can be found in [Adami et al. 2015]). If the claim were false, there would exist an edge e of the graph on which u is not identically zero but on which there exists $x_0 \in \mathcal{G} \setminus Z$ such that $u(x_0) = 0$. If x_0 belongs to the interior of e , u' vanishes at that point, by regularity. Otherwise, x_0 is a vertex $v \notin Z$, so that the derivatives $u'_e(v)$ are nonnegative for every $e \succ v$, thus they all vanish by the Kirchhoff condition. By uniqueness in the Cauchy problem, we deduce that u vanishes identically on the edge e , a contradiction.

Remark 3.2. The proof of the next results relies on the following approximation procedure. Given a noncompact graph $\mathcal{G} \in \mathbf{G}$, we construct an increasing sequence $(\mathcal{K}_n)_n \subseteq \mathcal{G}$ of connected compact graphs such that $\bigcup_{n \geq 1} \mathcal{K}_n = \mathcal{G}$, and a sequence $(\chi_n)_n \subseteq H^1(\mathcal{G})$ of cut-off functions such that

$$0 \leq \chi_n \leq 1, \quad \|\chi_n'\|_\infty \leq \frac{1}{\ell}, \quad \chi_n|_{\mathcal{K}_{n-1}} = 1, \quad \text{supp } \chi_n \subseteq \mathcal{K}_n,$$

with ℓ as in Definition 2.1.

To describe the graphs \mathcal{K}_n we begin by performing a preliminary operation on \mathcal{G} as follows. On each half-line of \mathcal{G} (if any) we insert vertices of degree 2 at the points of coordinates $k\ell$, $k = 1, 2, \dots$. Every half-line can now be viewed as a sequence of consecutive edges (each of length ℓ). With some abuse of notation, we still call $\mathcal{G} = (\mathbb{V}, \mathbb{E})$ the new graph obtained in this way (see Remark 2.2). Let now $v_0 \in \mathcal{G}$ be a fixed vertex. For every $n \geq 1$, let \mathbb{V}_n be the set of vertices of \mathbb{V} that can be reached from v_0 traveling on at most n edges. As each node of \mathcal{G} has finite degree, the sets \mathbb{V}_n are finite and, since \mathcal{G} is connected, for each vertex $v \in \mathcal{G}$ (different from v_0) there exists $n_0(v) \geq 1$ such that v belongs to \mathbb{V}_n for every $n \geq n_0(v)$. Then we define the graph \mathcal{K}_n as $(\mathbb{V}_n, \mathbb{E}_n)$, where \mathbb{E}_n is the set of edges of \mathbb{E} whose vertices belong to \mathbb{V}_n . Clearly, each \mathcal{K}_n is connected and compact, and $\bigcup_{n \geq 1} \mathcal{K}_n = \mathcal{G}$. Finally, we define χ_n to be equal to 1 on \mathcal{K}_{n-1} , to 0 on $\mathcal{G} \setminus \mathcal{K}_n$ and affine on every edge of $\mathcal{K}_n \setminus \mathcal{K}_{n-1}$. All the required properties trivially hold (the bound on χ_n' follows from the fact that all edges of \mathcal{G} have length at least ℓ).

Finally, given $u \in H_Z^1(\mathcal{G})$, it is straightforward to check that $\chi_n u \rightarrow u$ in $H^1(\mathcal{G})$ as $n \rightarrow \infty$.

Exploiting Remark 3.2, we now construct suitable minimizing sequences for J on \mathcal{N}_Z and \mathcal{M}_Z .

Proposition 3.3. *Let $\mathcal{G} \in \mathbf{G}$ be noncompact and $\lambda > -\omega_Z(\mathcal{G})$. There exists a minimizing sequence $(u_n)_n \subseteq \mathcal{N}_Z$ for J and $u \in H^1_Z(\mathcal{G})$ such that*

$$u_n \rightharpoonup u \text{ weakly in } H^1(\mathcal{G}), \quad u \geq 0, \quad J'(u) = 0.$$

Proof. Keeping in mind the notation of Remark 3.2, let $(\mathcal{K}_n)_n$ be the sequence of compact graphs approximating \mathcal{G} and let $\partial\mathcal{K}_n = \mathbb{V}_n \setminus \mathbb{V}_{n-1}$. Define the Hilbert space $H_n := H^1_{Z \cup \partial\mathcal{K}_n}(\mathcal{K}_n)$ and the Nehari manifold associated to J on H_n , namely

$$\mathcal{N}_n := \{u \in H_n \mid u \neq 0, J'(u)u = 0\}.$$

If $u \in H_n$, it vanishes on $\partial\mathcal{K}_n$ and, after extending it by 0, it can be viewed as a function in $H^1_Z(\mathcal{G})$, that we still denote by u . Therefore, $\mathcal{N}_n \subseteq \mathcal{N}_Z$ for every $n \geq 1$. Let $u_n \in \mathcal{N}_n$ be a ground state for J restricted to H_n , that is,

$$J(u_n) = \inf_{v \in \mathcal{N}_n} J(v).$$

The existence of u_n is standard by the compactness of the embedding of $H^1(\mathcal{K}_n)$ into $L^p(\mathcal{K}_n)$ observing that, by construction, $\omega_{Z \cup \partial\mathcal{K}_n}(\mathcal{K}_n) \geq \omega_Z(\mathcal{G})$. Also, as $u \in \mathcal{N}_n$ if and only if $|u| \in \mathcal{N}_n$ and $J(u) = J(|u|)$, we can assume that $u_n \geq 0$ on \mathcal{G} .

We claim that $(u_n)_n$ is a minimizing sequence for J on \mathcal{N}_Z . First, since $\mathcal{K}_{n-1} \subset \mathcal{K}_n$ for every n , the sequence $(J(u_n))_n$ is nonincreasing.

Given any $\varepsilon > 0$, let $\bar{u} \in \mathcal{N}_Z$ be such that $J(\bar{u}) \leq \inf_{v \in \mathcal{N}_Z} J(v) + \varepsilon/2$. Let $(\chi_n)_n$ be the sequence of cut-off functions of Remark 3.2. For every n , the function $\tilde{u}_n := \pi_\lambda(\chi_n \bar{u})$ is in \mathcal{N}_Z and $\text{supp } \tilde{u}_n \subseteq \mathcal{K}_n$, which means, in particular, that \tilde{u}_n (restricted to \mathcal{K}_n) is in \mathcal{N}_n . Moreover, by Remark 3.2 and the continuity of π_λ , as soon as n is large enough we have

$$J(\tilde{u}_n) \leq \inf_{v \in \mathcal{N}_Z} J(v) + \varepsilon.$$

Therefore, for all n large,

$$J(u_n) = \inf_{v \in \mathcal{N}_n} J(v) \leq J(\tilde{u}_n) \leq \inf_{v \in \mathcal{N}_Z} J(v) + \varepsilon.$$

Thus $(u_n)_n$ is a minimizing sequence for J on \mathcal{N}_Z , and the claim is proved. Since $(u_n)_n$ is bounded in $H^1_Z(\mathcal{G})$ (like all minimizing sequences), up to a subsequence it converges weakly to some $u \in H^1_Z(\mathcal{G})$ that also satisfies $u \geq 0$. Since u_n minimizes J over \mathcal{N}_n , it follows that $J'(u_n)\phi = 0$ for every $\phi \in H_n$. As $u \mapsto J'(u)\phi$ is weakly continuous on $H^1_Z(\mathcal{G})$, letting $n \rightarrow \infty$ shows that $J'(u)\phi = 0$ for every $\phi \in H_n$ and every n , and thus, by density, that $J'(u) = 0$. \square

Proposition 3.4. *Let $\mathcal{G} \in \mathbf{G}$ be noncompact and $\lambda > -\omega_Z(\mathcal{G})$. There exists a minimizing sequence $(u_n)_n \subseteq \mathcal{M}_Z$ for J and $u \in H^1_Z(\mathcal{G})$ such that*

$$u_n \rightharpoonup u \text{ weakly in } H^1(\mathcal{G}) \quad \text{and} \quad J'(u) = 0.$$

Proof. The proof is very similar to the one of Proposition 3.3, to which we refer for the notation. Let

$$\mathcal{M}_n := \{v \in H_n \mid v^\pm \in \mathcal{N}_n\}$$

and, for each n , let $u_n \in \mathcal{M}_n$ be a nodal ground state for J restricted to H_n , that is,

$$J(u_n) = \inf_{v \in \mathcal{M}_n} J(v).$$

The existence of u_n follows plainly by the compactness of the embedding of $H^1(\mathcal{K}_n)$ into $L^p(\mathcal{K}_n)$ as, for example, in [Szulkin and Weth 2010, Theorem 18] observing again that $\omega_{Z \cup \partial \mathcal{K}_n}(\mathcal{K}_n) \geq \omega_Z(\mathcal{G})$.

We claim that $(u_n)_n$ is a minimizing sequence for J on \mathcal{M}_Z . As above, $(J(u_n))_n$ is nonincreasing. If $u \in \mathcal{M}_Z$, we have $(\chi_n u)^\pm = \chi_n u^\pm$, and both functions are nonzero if n is large enough. By the continuity of π_λ and Remark 3.2, as $n \rightarrow \infty$,

$$\pi_\lambda(\chi_n u^+) + \pi_\lambda(\chi_n u^-) \rightarrow \pi_\lambda(u^+) + \pi_\lambda(u^-) = u \quad \text{in } H_Z^1(\mathcal{G}).$$

Now, given any $\varepsilon > 0$, let $\bar{u} \in \mathcal{M}_Z$ satisfy $J(\bar{u}) \leq \inf_{\mathcal{M}_Z} J + \varepsilon/2$. Define $\tilde{u}_n := \pi_\lambda(\chi_n \bar{u}^+) + \pi_\lambda(\chi_n \bar{u}^-)$, so $\tilde{u}_n \in \mathcal{M}_Z$, $\text{supp } \tilde{u}_n \subseteq \mathcal{K}_n$, whence $\tilde{u}_n \in \mathcal{M}_n$. Then, for every n large enough,

$$J(u_n) = \inf_{v \in \mathcal{M}_n} J(v) \leq J(\tilde{u}_n) \leq \inf_{v \in \mathcal{M}_Z} J(v) + \varepsilon,$$

showing that $(u_n)_n$ is a minimizing sequence for J on \mathcal{M}_Z . Since, by Proposition 2.3, $J'(u_n)\phi = 0$ for every $\phi \in H_n$, we conclude exactly as in the proof of Proposition 3.3. \square

We are now in position to prove Theorems 1.3–1.4.

Proof of Theorem 1.3. Let us prove the two statements separately.

Proof of (i). Let $(u_n)_n \subseteq \mathcal{N}_Z$ be the minimizing sequence for J on \mathcal{N}_Z constructed in Proposition 3.3 and let $u \geq 0$ be its weak limit. We first show that $u \not\equiv 0$. Indeed, if this were the case, then $u_n \rightarrow 0$ in $H_Z^1(\mathcal{G})$, so that

$$\inf_{v \in \mathcal{N}_Z} J(v) = \liminf_{n \rightarrow \infty} J(u_n) \geq J^\infty(\mathcal{G}; Z),$$

which is ruled out by assumption (1-5). Now as $u \not\equiv 0$ and $J'(u) = 0$, u is a nontrivial solution of (1-3). In particular, $u \in \mathcal{N}_Z$ and then, by (2-1) and weak lower semicontinuity,

$$J(u) = \kappa \|u\|_p^p \leq \liminf_{n \rightarrow \infty} \kappa \|u_n\|_p^p = \liminf_{n \rightarrow \infty} J(u_n) = \inf_{v \in \mathcal{N}_Z} J(v),$$

showing that u is a ground state. As such, u is positive on $\mathcal{G} \setminus Z$ by Remark 3.1. \square

Proof of (ii). Consider the minimizing sequence $(u_n)_n$ given by Proposition 3.4 and its weak limit $u \in H_Z^1(\mathcal{G})$ satisfying $J'(u) = 0$. We first show that $u^\pm \not\equiv 0$. For every n ,

$$J(u_n) = J(u_n^+) + J(u_n^-) \geq J(u_n^+) + \inf_{v \in \mathcal{N}_Z} J(v).$$

If, for instance, $u^+ \equiv 0$, then $u_n^+ \rightarrow 0$ in $H_Z^1(\mathcal{G})$, so that

$$\inf_{v \in \mathcal{M}_Z} J(v) = \liminf_{n \rightarrow \infty} J(u_n) \geq \liminf_{n \rightarrow \infty} J(u_n^+) + \inf_{v \in \mathcal{N}_Z} J(v) \geq J^\infty(\mathcal{G}; Z) + \inf_{v \in \mathcal{N}_Z} J(v),$$

by definition of $J^\infty(\mathcal{G}; Z)$, which contradicts (1-6). In the same way one proves that $u^- \neq 0$. As $J'(u) = 0$, it follows that u is a nonzero sign changing solution of (1-3), and hence $u \in \mathcal{M}_Z$. Then by weak lower semicontinuity, we conclude that

$$J(u) = \kappa \|u\|_p^p \leq \kappa \liminf_{n \rightarrow \infty} \|u_n\|_p^p = \liminf_{n \rightarrow \infty} J(u_n) = \inf_{v \in \mathcal{M}_Z} J(v),$$

namely that u is the required minimizer, i.e., a nodal ground state of (1-3). \square

Proof of Theorem 1.4. Let $u \in \mathcal{M}_Z$. Since $u^\pm \in \mathcal{N}_Z$,

$$J(u) = J(u^+) + J(u^-) \geq 2 \inf_{v \in \mathcal{N}_Z} J(v),$$

which is (1-7).

Now assume that $u \in \mathcal{M}_Z$ satisfies

$$J(u) = \inf_{v \in \mathcal{M}_Z} J(v) = 2 \inf_{v \in \mathcal{N}_Z} J(v).$$

Then $J(u^+) = J(u^-) = \inf_{v \in \mathcal{N}_Z} J(v)$, and therefore u^\pm are both ground states of J . As such, by Remark 3.1, they cannot vanish in $\mathcal{G} \setminus Z$, which is a contradiction since $u^\pm \neq 0$. \square

4. Graphs with at least one half-line

In this section we discuss ground states and nodal ground states for noncompact graphs with at least one half-line. These graphs may have infinitely many edges. When a finite number of edges is required, it is explicitly mentioned in the statement of the result. Since for such kind of graphs the bottom of the spectrum of $-\mathrm{d}^2/\mathrm{d}x^2$ on $H_Z^1(\mathcal{G})$ always satisfies

$$\omega_Z(\mathcal{G}) = 0,$$

all the results of this section will hold for every $\lambda \in (0, +\infty)$.

The prototype cases in this context are given by the real line and the half-line, about which everything is known (see, e.g., [Cazenave 2003; Le Coz 2009]). Since the ground states on \mathbb{R} play a very important role in what follows, we recall briefly their main features. On the real line the only nontrivial L^2 solutions to (1-1) are called *solitons* and are unique up to translations and sign. Denoting by ϕ_λ the unique positive and even soliton, for every $\lambda > 0$,

$$s_\lambda := J_\lambda(\phi_\lambda) = \inf_{v \in \mathcal{N}_\lambda(\mathbb{R})} J_\lambda(v),$$

namely the solitons are the ground states on \mathbb{R} (see, e.g., [Le Coz 2009, Proposition 3.12]). Similarly, on the half-line (with $Z = \emptyset$) there is a unique nontrivial L^2 solution (up to sign) to (1-1), given by the so-called *half-soliton* ψ_λ , i.e., the restriction of ϕ_λ to \mathbb{R}^+ . It is the ground state, and

$$J_\lambda(\psi_\lambda) = \inf_{v \in \mathcal{N}_\lambda(\mathbb{R}^+)} J_\lambda(v) = \frac{1}{2} s_\lambda. \quad (4-1)$$

If $Z = \{0\}$ (the vertex of the half-line) there are no nontrivial L^2 solutions to (1-1), as any nontrivial solution of $u'' + |u|^{p-2}u = \lambda u$ on \mathbb{R}^+ that vanishes at the origin corresponds to a periodic orbit in the phase plane associated to the equation and thus is not in $L^2(\mathbb{R}^+)$.

For a general graph with half-lines, a first marker of the importance of the level s_λ is given by the following straightforward property.

Proposition 4.1. *Let $\mathcal{G} \in \mathbf{G}$ contain at least one half-line and $\lambda > 0$. Then*

$$\frac{1}{2}s_\lambda \leq \inf_{v \in \mathcal{N}_Z} J(v) \leq s_\lambda. \quad (4-2)$$

Proof. The inequalities can be easily proved, using rearrangement techniques, arguing exactly as in the proof of [Adami et al. 2015, Theorem 2.2]. \square

In the search for ground states, it is crucial to understand whether one can reverse the second inequality in (4-2) (see, e.g., the discussion in [Adami et al. 2015; 2016] in the context of energy ground states of prescribed mass). In [Adami et al. 2015, Theorems 2.3–2.5] the authors individuated a topological condition on \mathcal{G} under which this can actually be done. To state it we recall that \mathbb{V}_∞ denotes the set of *vertices at infinity* of \mathcal{G} . Every vertex at infinity is a vertex of the graph \mathcal{G} , but is *not* a point of the metric space \mathcal{G} . The assumption introduced in [Adami et al. 2015] is:

$$\begin{aligned} \text{for every } e \in \mathbb{E}, \text{ every connected component of the graph } (\mathbb{V}, \mathbb{E} \setminus \{e\}) \\ \text{contains at least one vertex } v \in \mathbb{V}_\infty. \end{aligned} \quad (\text{H})$$

In [Adami et al. 2015, Theorem 2.3] the authors proved that, if \mathcal{G} satisfies assumption (H), then for every $u \in H^1(\mathcal{G})$ we have $\#u^{-1}(t) \geq 2$ for almost every $t \in (0, \|u\|_\infty)$. The main consequence of this (originally proved in [Adami et al. 2015, Theorem 2.5] for the problem of prescribed-mass ground states) is described in the following result.

Theorem 4.2 [De Coster et al. 2023, Theorem 2.6]. *If $\mathcal{G} \in \mathbf{G}$ satisfies assumption (H) and $\lambda > 0$, then*

$$\inf_{v \in \mathcal{N}} J(v) = s_\lambda$$

and it is never achieved, unless \mathcal{G} is isometric to \mathbb{R} or to a “tower of bubbles” shown in Figure 2.

In this paper the setting is different from that of [Adami et al. 2015; De Coster et al. 2023] for at least two reasons: first, the boundary conditions are more general and the presence of the set Z must be taken into account; second, we are also interested in nodal ground states. For these reasons it is convenient to reformulate and generalize assumption (H) in a form that is more suited to handle the questions under study. As in the Introduction, consider the set

$$F(\mathcal{G}) = \{e \in \mathbb{E} \mid \text{at least one connected component of } (\mathbb{V}, \mathbb{E} \setminus \{e\}) \text{ has no vertices in } \mathbb{V}_\infty \cup Z\}$$

and the assumptions

$$\#F(\mathcal{G}) = 0, \quad (\text{H0})$$

$$\#F(\mathcal{G}) \leq 1. \quad (\text{H1})$$

Note that (H0) and (H1) are, respectively, the assumptions in (i) and (ii) in Theorem 1.6. From now on, with some abuse of notation, we denote the graph $(\mathbb{V}, \mathbb{E} \setminus \{e\})$ simply by $\mathcal{G} \setminus e$.

To investigate the relations between assumptions (H0) and (H), it is convenient to define a new graph $\tilde{\mathcal{G}}$ in the following way. If $Z = \emptyset$, we set $\tilde{\mathcal{G}} = \mathcal{G}$. Otherwise, we replace every (finite) edge e ending at a vertex of Z by a half-line, still called e . We obtain in this way a new graph $\tilde{\mathcal{G}} = (\tilde{\mathbb{V}}, \tilde{\mathbb{E}})$ that has the same number of vertices and edges as \mathcal{G} . The only difference is that edges of \mathcal{G} terminating at vertices of Z are replaced, in $\tilde{\mathcal{G}}$, by half-lines terminating at vertices in $\tilde{\mathbb{V}}_\infty$.

Then it is easily seen that

$$\mathcal{G} \text{ satisfies (H0)} \iff \tilde{\mathcal{G}} \text{ satisfies (H)}. \quad (4-3)$$

Indeed, to say that \mathcal{G} satisfies (H0) means that there are no edges in \mathbb{E} whose removal generates a connected component without vertices in $\mathbb{V}_\infty \cup Z$, namely that for every $e \in \mathbb{E}$, every connected component of $\mathcal{G} \setminus e$ has a vertex in $\mathbb{V}_\infty \cup Z$. But this, read on $\tilde{\mathcal{G}}$, means that every connected component of $\tilde{\mathcal{G}} \setminus e$ has a vertex in $\tilde{\mathbb{V}}_\infty$, which is (H) for $\tilde{\mathcal{G}}$.

Furthermore, to say that $\#F(\mathcal{G}) = 1$, namely that $F(\mathcal{G}) = \{e\}$ for exactly one edge e , means that the graph $\mathcal{G} \setminus e$ decomposes as

$$\mathcal{G} \setminus e = \mathcal{G}_K \cup \mathcal{G}', \quad (4-4)$$

where \mathcal{G}_K is connected and has no vertices in $\mathbb{V}_\infty \cup Z$, while \mathcal{G}' is connected and contains *all* the vertices of $\mathbb{V}_\infty \cup Z$. Also, there are no edges other than e that permit a decomposition like (4-4). As \mathcal{G} has at least one half-line, the unique $e \in F(\mathcal{G})$ can never have a vertex in Z . However, e can be a half-line. In this case, though, $\mathcal{G} \setminus e = \mathcal{G}_K \cup \{v_\infty\}$, where v_∞ is the vertex at infinity of the half-line e . Thus in this case the graph \mathcal{G} is made of a set of bounded edges without vertices in Z and a *single* half-line attached to it.

The next result plays a key role in the proof of some of the subsequent results. Roughly, it states that any graph satisfying $\#F(\mathcal{G}) = 1$ can be turned into a graph satisfying (H0) by attaching to it a suitable half-line.

Lemma 4.3. *Let $\mathcal{G} \in \mathbf{G}$ be a graph with at least one half-line satisfying $\#F(\mathcal{G}) = 1$. Let e be such that $F(\mathcal{G}) = \{e\}$ and \mathcal{G}_K be the connected component of $\mathcal{G} \setminus e$ as in (4-4). Choose a vertex v in \mathcal{G}_K and define a new graph $\tilde{\mathcal{G}}_v$ by¹ $\tilde{\mathcal{G}}_v = \mathcal{G} \cup h$, where h is a half-line attached at v . Then $\tilde{\mathcal{G}}_v$ satisfies (H0).*

Proof. Let v_∞ be the vertex at infinity of h and assume by contradiction that $\#F(\tilde{\mathcal{G}}_v) \geq 1$, namely that there exists $\tilde{e} \in F(\tilde{\mathcal{G}}_v)$. We claim that $\tilde{e} \neq h$. Indeed, removing h from $\tilde{\mathcal{G}}_v$ would leave v_∞ isolated, splitting $\tilde{\mathcal{G}}_v \setminus h$ into the two connected components \mathcal{G} and $\{v_\infty\}$. Since both of them contain vertices in $\tilde{\mathbb{V}}_\infty$, this violates the definition of \tilde{e} . Similarly, it cannot be $\tilde{e} = e$: removing e from $\tilde{\mathcal{G}}_v$, and recalling that h is attached to \mathcal{G}_K , would decompose $\tilde{\mathcal{G}}_v$ into connected components as $(\mathcal{G}_K \cup h) \cup \mathcal{G}'$, violating again the definition of \tilde{e} as before. We are left with the case where \tilde{e} is different from both h and e . In this case we have the decomposition

$$\tilde{\mathcal{G}}_v \setminus \tilde{e} =: \tilde{\mathcal{G}}_K \cup \tilde{\mathcal{G}}',$$

¹Shorthand for $(\mathbb{V} \cup \{v_\infty\}, \mathbb{E} \cup \{h\})$, where v_∞ is the vertex at infinity of h .

with obvious meaning of the symbols. By construction, the half-line h is attached to $\tilde{\mathcal{G}}'$. Removing h and v_∞ from $\tilde{\mathcal{G}}'$ does not disconnect it and, since $\tilde{\mathcal{G}}'$ contains at least another vertex in $\mathbb{V}_\infty \cup Z$, we see that $\tilde{\mathcal{G}}' \setminus (\{v_\infty\}, \{h\})$ is not empty. Therefore $\tilde{\mathcal{G}}_K$ and $\tilde{\mathcal{G}}' \setminus (\{v_\infty\}, \{h\})$ are both nonempty, connected, disjoint and their union is $\mathcal{G} \setminus \tilde{e}$, namely $\tilde{e} \in F(\mathcal{G})$. Since we also have $e \in F(\mathcal{G})$, this shows that $\#F(\mathcal{G}) \geq 2$, violating the assumption. \square

We can now prove that the assumptions (H0) and (H1) are sufficient to rule out the existence of ground states and nodal ground states respectively, as stated in Theorem 1.6.

Proof of Theorem 1.6. We split the proof into two parts.

Part 1: proof of (1-8) and (1-10). Of course it is sufficient to work with nonnegative functions, which we do without further warnings. Since \mathcal{G} contains at least one half-line, Proposition 4.1 guarantees that $\inf_{v \in \mathcal{N}_Z} J(v) \leq s_\lambda$. To prove the reverse inequality under assumption (H0), let $\tilde{\mathcal{G}}$ be the graph defined after the statement of assumptions (H0)–(H1).

Since, as metric spaces, $\mathcal{G} \subseteq \tilde{\mathcal{G}}$, every function $u \in H_Z^1(\mathcal{G})$ extended by 0 on $\tilde{\mathcal{G}} \setminus \mathcal{G}$ can be seen as a function $\tilde{u} \in H^1(\tilde{\mathcal{G}})$. Plainly,

$$\|\tilde{u}\|_{L^q(\tilde{\mathcal{G}})} = \|u\|_{L^q(\mathcal{G})} \quad \text{for every } q \in [1, +\infty], \quad \|\tilde{u}'\|_{L^2(\tilde{\mathcal{G}})} = \|u'\|_{L^2(\mathcal{G})}.$$

This implies that $\tilde{u} \in \mathcal{N}(\tilde{\mathcal{G}})$ and since $\tilde{\mathcal{G}}$ satisfies (H) (because \mathcal{G} satisfies (H0); see (4-3)),

$$J(u) = \kappa \|u\|_{L^p(\mathcal{G})}^p = \kappa \|\tilde{u}\|_{L^p(\tilde{\mathcal{G}})}^p = J(\tilde{u}) \geq \inf_{v \in \mathcal{N}(\tilde{\mathcal{G}})} J(v) = s_\lambda$$

by Theorem 4.2. As this holds for every (nonnegative) $u \in \mathcal{N}_Z$, (1-10) is proved.

Assume now that for some nonnegative $u \in \mathcal{N}_Z$ we have $J(u) = s_\lambda$. Considering, as above, the function $\tilde{u} \in \mathcal{N}(\tilde{\mathcal{G}})$, we see that $J(\tilde{u}) = J(u) = s_\lambda$, namely that \tilde{u} is a ground state for J on $\mathcal{N}(\tilde{\mathcal{G}})$. As such, $\tilde{u}(x) > 0$ for every $x \in \tilde{\mathcal{G}}$, which shows that $Z = \emptyset$, namely that $\tilde{\mathcal{G}} = \mathcal{G}$. We then conclude by Theorem 4.2.

Part 2: proof of (1-9) and (1-11). We first prove (1-9). By density, for every $\varepsilon > 0$ there exists a nonnegative $u_1 \in \mathcal{N}_Z(\mathcal{G})$ with compact support such that

$$J(u_1) \leq \inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v) + \varepsilon.$$

Similarly, there exists a nonnegative $u_2 \in \mathcal{N}(\mathbb{R})$ with compact support such that $J(u_2) \leq s_\lambda + \varepsilon$. By taking a translation of u_2 (if necessary), we can make sure that its support, identified with an interval on some half-line of \mathcal{G} , does not intersect the support of u_1 . We then define $w \in H^1(\mathcal{G})$ by

$$w(x) = \begin{cases} u_1(x) & \text{if } x \in \mathcal{G} \setminus \text{supp}(u_2), \\ -u_2(x) & \text{if } x \in \text{supp}(u_2). \end{cases}$$

Obviously, $w \in \mathcal{M}_Z$ and

$$J(w) = J(u_1) + J(u_2) \leq s_\lambda + \inf_{v \in \mathcal{N}_Z} J(v) + 2\varepsilon.$$

Since ε is arbitrary, we conclude.

We now prove the reverse inequality in (1-11) under assumption (H1). If $\#F(\mathcal{G}) = 0$, then \mathcal{G} satisfies (H0) and Theorem 1.6(i) shows that $\inf_{v \in \mathcal{N}_Z} J(v) = s_\lambda$, so that the inequality to be proved is $\inf_{v \in \mathcal{M}_Z} J(v) \geq 2s_\lambda$. Given any $u \in \mathcal{M}_Z$, applying Theorem 1.6(i) to u^+ and u^- immediately yields

$$J(u) = J(u^+) + J(u^-) > 2s_\lambda,$$

the inequality being strict since both u^+ and u^- vanish somewhere on \mathcal{G} . This ensures that nodal ground states do not exist in this case.

Suppose now that $\#F(\mathcal{G}) = 1$. Let e be the unique element of $F(\mathcal{G})$ and consider the decomposition (4-4):

$$\mathcal{G} \setminus e = \mathcal{G}_K \cup \mathcal{G}'.$$

Given a vertex v of \mathcal{G}_K , for every $u \in \mathcal{M}_Z$, at least one among u^+ and u^- , say u^+ , vanishes at v . Let $\tilde{\mathcal{G}}_v$ be the graph constructed in Lemma 4.3, obtained by attaching to v a half-line h . Since u^+ vanishes at v , it can be extended to a function \tilde{u}^+ simply by defining it to be 0 on h . Clearly, $\tilde{u}^+ \in \mathcal{N}_Z(\tilde{\mathcal{G}}_v)$ and, since $\tilde{\mathcal{G}}_v$ satisfies (H0) by Lemma 4.3, we obtain $J(\tilde{u}^+) \geq s_\lambda$. Then

$$J(u) = J(u^+) + J(u^-) = J(\tilde{u}^+) + J(u^-) \geq s_\lambda + \inf_{v \in \mathcal{N}_Z} J(v),$$

concluding the proof of (1-11).

It remains to show that the infimum is not achieved when $\#F(\mathcal{G}) = 1$. To this end, it suffices to observe that the inequality used above, $J(\tilde{u}^+) \geq s_\lambda$, is in fact strict. Indeed if $J(\tilde{u}^+) = s_\lambda = \inf_{v \in \mathcal{N}(\tilde{\mathcal{G}})} J(v)$, then \tilde{u}^+ is a ground state on $\tilde{\mathcal{G}}$, and hence it cannot vanish anywhere, contrary to the fact that $\tilde{u}^+ \equiv 0$ on h . \square

Remark 4.4. The assumptions of Theorem 1.6 are sharp. Indeed, Theorem 4.8 below shows that there exist graphs \mathcal{G} satisfying $\#F(\mathcal{G}) \geq 1$ that admit ground states, while in Theorem 4.16 and Remark 4.18 we exhibit graphs \mathcal{G} satisfying $\#F(\mathcal{G}) \geq 2$ that admit nodal ground states.

Theorem 1.6 shows that nonexistence of ground states or nodal ground states can be determined by purely topological properties of the graph. The situation for existence is, on the contrary, more involved.

In some cases, existence results for ground states based solely on the topology of the graph can be easily obtained when $Z = \emptyset$, as, for example, if \mathcal{G} has a finite number of edges. In this respect, there is not much to say since the techniques developed in [Adami et al. 2016, Section 3] for the problem of prescribed-mass minimizers of the energy work in the present setting as well, as we now briefly show.

For the reader's convenience, let us first recall with the next lemma a standard relation between the action of a function and the number of preimages of each value it attains. These properties are stated in [De Coster et al. 2023, Propositions 2.2 and 2.8] and are consequences of standard rearrangement techniques on graphs. For this reason, we omit the proof of the lemma.

Lemma 4.5. *Let \mathcal{G} be a noncompact metric graph. Given $\lambda > 0$ and an integer $K \geq 1$, let $u \in \mathcal{N}_{\lambda, Z}$ satisfy $u \geq 0$ on \mathcal{G} and $\#u^{-1}(t) \geq K$ for almost every $t \in (0, \max_{\mathcal{G}} u)$. Then $J_\lambda(u) \geq K \frac{1}{2} s_\lambda$. Furthermore, if equality holds then*

- $\#u^{-1}(t) = K$ for almost every $t \in (0, \max_{\mathcal{G}} u)$;
- the support of u has infinite measure;
- $u^{-1}(t)$ has zero measure for all $t \in (0, \max_{\mathcal{G}} u)$.

Since we use Theorem 1.3 in what follows, we prove the following characterization of $J^\infty(\mathcal{G}; Z)$.

Proposition 4.6. *Let $\mathcal{G} \in \mathbf{G}$ be a noncompact graph with a finite number of edges and $\lambda > 0$. Then*

$$J_\lambda^\infty(\mathcal{G}; Z) = s_\lambda. \quad (4-5)$$

Proof. By density, for every $\varepsilon > 0$ there exists $u = u_\varepsilon \in \mathcal{N}(\mathbb{R})$ with compact support such that $J(u) \leq s_\lambda + \varepsilon$. For every n large enough, the function $u_n(x) = u(x - n)$ is supported in \mathbb{R}^+ and, as such, it can be seen as an element of \mathcal{N}_Z by placing its support on a half-line of \mathcal{G} and then extending it by 0 outside its support. Clearly $u_n \rightharpoonup 0$ in $H^1(\mathcal{G})$ and

$$\liminf_n J(u_n) \leq s_\lambda + \varepsilon.$$

Since ε is arbitrary, the “ \leq ” part is proved. For the reverse inequality, let $(u_n)_n \subset \mathcal{N}_Z$ be a sequence converging weakly to 0 in $H^1(\mathcal{G})$ with $J_\lambda(u_n) \rightarrow J_\lambda^\infty(\mathcal{G}; Z)$. We can assume $u_n \geq 0$ for every n since otherwise we replace it by $|u_n|$ that is still in \mathcal{N}_Z . By weak convergence, we also see that $u_n \rightarrow 0$ in $L_{\text{loc}}^\infty(\mathcal{G})$. Hence, if ε_n denotes the maximum of u_n on the set of all bounded edges of \mathcal{G} , clearly $\varepsilon_n \rightarrow 0$. Therefore, letting $v_n := (u_n - \varepsilon_n)^+$, we see from Proposition 2.4 that $v_n \not\equiv 0$ for every n large enough. Since \mathcal{G} contains at least one half-line, by construction $\#v_n^{-1}(t) \geq 2$ for every $t \in (0, \max v_n)$ and every n , since v_n vanishes on the set of all bounded edges, and the same holds for $\pi_\lambda(v_n)$. So $J(\pi_\lambda(v_n)) \geq s_\lambda$ by Lemma 4.5. Furthermore, as $n \rightarrow \infty$,

$$n_\lambda(v_n)^{p-2} = \frac{\|v_n'\|_2^2 + \lambda \|v_n\|_2^2}{\|v_n\|_p^p} \leq \frac{\|u_n'\|_2^2 + \lambda \|u_n\|_2^2}{\|u_n\|_p^p + o(1)} = 1 + o(1), \quad (4-6)$$

entailing

$$s_\lambda \leq J(\pi_\lambda(v_n)) = J(n_\lambda(v_n)v_n) = \kappa n_\lambda(v_n)^p \|v_n\|_p^p \leq \kappa(1 + o(1))\|u_n\|_p^p = J(u_n) + o(1),$$

from which we obtain $\liminf_n J(u_n) \geq s_\lambda$. Since this holds for every sequence converging weakly to 0, the proof is complete. \square

Remark 4.7. The assumption that the graph has a finite number of edges in Proposition 4.6 cannot be removed. This can be seen considering, for instance, the following example. On a real line we insert, for each integer $k \geq 1$, a node v_k at the point of coordinate k and a terminal edge L_k of length k , by identifying v_k with an endpoint of L_k (Figure 5). By density, for every $\varepsilon > 0$ there exist $k \geq 1$ and $u_k \in \mathcal{N}(\mathbb{R}^+)$ with compact support in $[0, k]$ such that $J(u_k) \leq \frac{1}{2}s_\lambda + \varepsilon$. Since u_k can be considered as a function on the edge L_k , we obtain a sequence $(u_k)_k \in \mathcal{N}(\mathcal{G})$ that converges weakly to 0 and such that $\liminf_k J_\lambda(u_k) \leq \frac{1}{2}s_\lambda + \varepsilon$. This proves that $J_\lambda^\infty(\mathcal{G}) \leq \frac{1}{2}s_\lambda$. By Proposition 4.1, we obtain in fact $J_\lambda^\infty(\mathcal{G}) = \frac{1}{2}s_\lambda$.

Having established (4-5), Theorem 1.3 yields existence of a ground state on a noncompact graph with a finite number of edges as soon as one can prove that $\inf_{v \in \mathcal{N}_Z} J(v) < s_\lambda$. This condition is analogous to the one appearing in the fixed mass case. As we anticipated above, such an inequality can be shown to hold for a number of graphs with $Z = \emptyset$ exploiting only topological properties, by the use of the “graph surgery” techniques developed in [Adami et al. 2016, Section 3].




Figure 5. The graph \mathcal{G} described in Remark 4.7.

Theorem 4.8. For every $\lambda > 0$, every graph \mathcal{G} depicted in Figure 6, for every length of its edges, satisfies

$$\inf_{v \in \mathcal{N}} J(v) < s_\lambda$$

and therefore admits a ground state.

Proof. The inequality can be proved starting with a soliton on \mathbb{R} , via rearrangement techniques, exactly as in the final part of Section 3 in [Adami et al. 2016]. The basic idea is that, on each of these graphs, one can construct explicit functions built from pieces of the positive even soliton ϕ_λ on \mathbb{R} and pieces of its monotone rearrangement on \mathbb{R}^+ , then projected on the Nehari manifold. For instance, on the tadpole such a function coincides on the loop of the graph (of total length L) with the restriction of the soliton to the interval $[-L/2, L/2]$, and on the half-line with the monotone rearrangement on \mathbb{R}^+ of the restriction of the soliton to $\mathbb{R} \setminus [-L/2, L/2]$. The construction on the other graphs in Figure 6 is analogous. Since rearrangements always lower the normalizing factor n_λ defined in (2-2), it is then easy to check that these functions realize action levels strictly smaller than that of the soliton s_λ . Existence of a ground state follows then from Theorem 1.3. \square

When Z is not empty, the existence of a ground state is harder to obtain and further conditions of metrical nature have to be imposed. Indeed, the next theorem shows that, if a graph hosts a ground state,




Figure 6. Some graphs with $Z = \emptyset$ admitting ground states. (A): line with a pendant; (B): signpost; (C): tadpole; (D): 3-fork.

the diameter of the set \mathcal{B} of all bounded edges cannot be arbitrarily small. Recall that $\text{diam}(\mathcal{B})$ is given by the supremum of lengths of the shortest paths between any two points of \mathcal{B} .

Theorem 4.9. *There exists a constant $C > 0$ depending only on $\lambda > 0$ and p such that, for every $\mathcal{G} \in \mathbf{G}$ with at least one half-line and every $Z \neq \emptyset$ such that $\inf_{v \in \mathcal{N}_Z} J(v)$ is achieved, we have*

$$\text{diam}(\mathcal{B}) \geq C,$$

where, as above, \mathcal{B} is the set of all bounded edges of \mathcal{G} .

Proof. Let $u \in \mathcal{N}_Z$ satisfy $J(u) = \inf_{v \in \mathcal{N}_Z} J(v)$. As usual we can assume that $u \geq 0$. Let us show that u attains its L^∞ -norm in \mathcal{B} . Suppose by contradiction that u instead attains its maximum at a point y on a half-line h . Then $\#u^{-1}(t) \geq 2$ for every $0 < t < \max u$. Indeed, t is attained at least once on h and once on a path γ joining y to a point in Z . Thus Lemma 4.5 and Proposition 4.1 imply that $J(u) \geq s_\lambda = \inf_{v \in \mathcal{N}_Z} J(v)$. Let us show that the inequality must be strict, which gives the desired contradiction. If this were not the case, Lemma 4.5 would imply that $\#u^{-1}(t) = 2$ for almost every $t \in (0, \max u)$. Since u already has two preimages on $h \cup \gamma$, this means that u must be constant on $\mathcal{G} \setminus (h \cup \gamma)$. This contradicts the last assertion of Lemma 4.5 unless $\mathcal{G} = h \cup \gamma$, that is \mathcal{G} is a half-line and homogeneous Dirichlet conditions are imposed at its origin. But this is impossible, because, as we already recalled, on the half-line with zero Dirichlet boundary conditions there is no nonzero solution.

Let then $\bar{x} \in \mathcal{B}$ be such that $\|u\|_\infty = u(\bar{x})$. By the Cauchy–Schwarz inequality, (2-1) and Proposition 4.1, letting z be any vertex in Z , we have

$$\|u\|_\infty = |u(\bar{x})| = |u(\bar{x}) - u(z)| \leq \sqrt{\text{diam}(\mathcal{B})} \|u'\|_{L^2(\mathcal{G})} \leq \sqrt{\text{diam}(\mathcal{B})} \sqrt{\frac{s_\lambda}{\kappa}}$$

which, coupled with Proposition 2.4, yields

$$C \leq \|u\|_p^p \leq \|u\|_\infty^{p-2} \|u\|_2^2 \leq \frac{1}{\lambda} \text{diam}(\mathcal{B})^{\frac{p}{2}-1} \left(\frac{s_\lambda}{\kappa}\right)^{\frac{p}{2}}$$

for a suitable constant $C > 0$ depending on λ and p only. \square

Remark 4.10. Comparing Theorems 4.8 and 4.9 highlights the effect of the set Z on the existence of ground states. One may wonder whether Theorem 4.9 can be improved to obtain a universal lower bound involving only the total length of the edges with a vertex in Z , rather than the whole set \mathcal{B} . However, this cannot be done in general: it is easy to exhibit graphs where ground states do exist and the length of the edges with vertices in Z is arbitrarily small. To see this, let \mathcal{G} be any given graph with $Z = \emptyset$ and such that $\inf_{v \in \mathcal{N}(\mathcal{G})} J(v) < s_\lambda$ (e.g., any of the graphs in Figure 6). Exploiting, for instance, the approximation procedure described in Remark 3.2, one can construct a function $u \in \mathcal{N}(\mathcal{G})$ so that $J(u) < s_\lambda$ and the support of u is contained in a suitable neighborhood of \mathcal{B} . In particular, there exists $M > 0$ such that the restriction of u to each half-line of \mathcal{G} satisfies $u \equiv 0$ on $[M, +\infty)$. For every $\ell > 0$, let then \mathcal{G}_ℓ be the graph obtained by attaching a single edge of length ℓ at the point $x = M$ of one of the half-lines of \mathcal{G} , and assume that the vertex of degree 1 of this edge is the only vertex in Z . Clearly, one can think of u as a

function in $\mathcal{N}_Z(\mathcal{G}_\ell)$ for every ℓ , so that $\inf_{v \in \mathcal{N}_Z(\mathcal{G}_\ell)} J(v) \leq J(u) < s_\lambda$, thus implying existence of ground states in $\mathcal{N}_Z(\mathcal{G}_\ell)$ by Theorem 1.3 and Proposition 4.6.

In the case of nodal ground states, it is not even needed to have $Z \neq \emptyset$ to recover the analogue of Theorem 4.9.

Theorem 4.11. *There exists a constant $C > 0$ depending only on $\lambda > 0$ and p such that, for every $\mathcal{G} \in \mathbf{G}$ with at least one half-line and every Z such that $\inf_{v \in \mathcal{M}_Z} J(v)$ is achieved, we have*

$$\text{diam}(\mathcal{B}) \geq C,$$

where, as above, \mathcal{B} is the set of all bounded edges of \mathcal{G} .

Proof. Let u be a nodal ground state. Observe that if u^+ attains its L^∞ -norm on a half-line, as in Theorem 4.9, we prove that $J_\lambda(u^+) > s_\lambda$. Hence $J_\lambda(u) = J_\lambda(u^+) + J_\lambda(u^-) > s_\lambda + \inf_{v \in \mathcal{N}_Z(\mathcal{G})} J_\lambda(v)$ which contradicts (1-9). The same is valid for u^- . Hence both u^+ and u^- attain their L^∞ -norm on \mathcal{B} only. Thus u changes sign in \mathcal{B} and as such, has a zero in \mathcal{B} . We then conclude as in Theorem 4.9 working on u^+ . \square

In view of Theorems 1.6, 4.9 and 4.11, it is clear that a suitable combination of topological and metrical features is needed to guarantee existence of ground states with $Z \neq \emptyset$ and nodal ground states. Towards this direction, we conclude the discussion of this section with two general procedures to construct graphs where ground states and nodal ground states do exist. The first one is genuinely of metrical nature, in that it is completely independent of the topology of the graph. The second one mixes topological and metrical properties.

In the next statement, by *pendant* we mean a finite-length terminal edge whose vertex of degree 1 is not in Z .

Theorem 4.12. *There exists a constant $C > 0$ depending only on $\lambda > 0$ and p such that, for every noncompact graph $\mathcal{G} \in \mathbf{G}$ with a finite number of edges,*

- (i) *if \mathcal{G} has a pendant of length $a \geq C$, then $\inf_{v \in \mathcal{N}_Z} J(v)$ is achieved;*
- (ii) *if \mathcal{G} has two pendants of lengths $a_1, a_2 \geq C$, then $\inf_{v \in \mathcal{M}_Z} J(v)$ is achieved.*

Remark 4.13. The assumption that \mathcal{G} has a finite number of edges cannot be removed. This can be easily seen as follows. For point (1) it is enough to consider the graph \mathcal{G} in Remark 4.7, for which $\inf_{v \in \mathcal{N}_\lambda(\mathcal{G})} J(v) = \frac{1}{2}s_\lambda$. Therefore, if u were a (positive) ground state, by Lemma 4.5, almost every $t \in (0, \max u)$ would be attained only once on \mathcal{G} , which is incompatible with the presence of vertices of degree 3. For point (2) one can simply consider any periodic graph, since such graphs admit no nodal ground state by Theorem 1.7.

Remark 4.14. In Theorem 4.12, C is the same for ground states and for nodal ground states and depends on the presence of the pendants but not on the rest of the graph. This will be of great help in Section 6.

Proof of Theorem 4.12. Let $\psi_\lambda \in H^1(\mathbb{R}^+)$ be the positive half-soliton satisfying, by (4-1), $J(\psi_\lambda) = \frac{1}{2}s_\lambda$. By density, there exists a function $u_1 \in \mathcal{N}(\mathbb{R}^+)$ supported in some interval $[0, C]$ such that

$$J(u_1) < \frac{3}{4}s_\lambda$$

(for example, one may take $(\psi_\lambda - \delta)^+$ with δ small and then project it on $\mathcal{N}_\lambda(\mathbb{R}^+)$).




Figure 7. Example of a graph \mathcal{G}_L as in Theorem 4.16, constructed starting with two graphs in Figure 6. If the vertical edge on the right is sufficiently far from the pendants of the graph, nodal ground states exist.

(1) Let \mathcal{G}_a be a graph with at least one pendant of length a . If a is larger than C , we may consider $[0, C]$ as contained in the pendant, identifying $x = 0$ with its vertex of degree 1. Extending u_1 by 0 on the remaining part of \mathcal{G}_a , we obtain a function $\tilde{u}_1 \in \mathcal{N}_Z(\mathcal{G}_a)$ such that

$$J(\tilde{u}_1) = J(u_1) < \frac{3}{4}s_\lambda.$$

The existence of a ground state follows from Proposition 4.6 and Theorem 1.3.

(2) We denote by \mathcal{G}_{a_1, a_2} a graph with two pendants e_1, e_2 of lengths a_1, a_2 and we show that if $a_1 \geq C$ and $a_2 \geq C$, then

$$\inf_{v \in \mathcal{M}_Z(\mathcal{G}_{a_1, a_2})} J(v) < J^\infty(\mathcal{G}_{a_1, a_2}; Z) + \inf_{v \in \mathcal{N}_Z(\mathcal{G}_{a_1, a_2})} J(v), \tag{4-7}$$

which, via Theorem 1.3, establishes the existence of a nodal ground state. By Propositions 4.1–4.6, for every a_1, a_2 ,

$$J^\infty(\mathcal{G}_{a_1, a_2}; Z) = s_\lambda, \quad \inf_{v \in \mathcal{N}_Z(\mathcal{G}_{a_1, a_2})} J(v) \geq \frac{1}{2}s_\lambda,$$

so that

$$J^\infty(\mathcal{G}_{a_1, a_2}; Z) + \inf_{v \in \mathcal{N}_Z(\mathcal{G}_{a_1, a_2})} J(v) \geq \frac{3}{2}s_\lambda.$$

Now, using the same u_1 as in (1), we define

$$\tilde{u}_1(x) = \begin{cases} u_1(x) & \text{if } x \in [0, a_1] \subset e_1, \\ -u_1(x) & \text{if } x \in [0, a_2] \subset e_2, \\ 0 & \text{elsewhere on } \mathcal{G}_{a_1, a_2}. \end{cases}$$

Clearly $\tilde{u}_1 \in \mathcal{M}_Z(\mathcal{G}_{a_1, a_2})$ and

$$J(\tilde{u}_1) = 2J(u_1) < \frac{3}{2}s_\lambda.$$

This proves (4-7). □

We now discuss the second procedure to find nodal ground states. The idea is to take two graphs admitting ground states and connect them by the addition of a faraway edge. So, let $\mathcal{G}^1, \mathcal{G}^2$ be any two noncompact graphs with a finite number of edges for which $\inf_{v \in \mathcal{N}_{Z^i}(\mathcal{G}^i)} J(v) < s_\lambda$. Given $L > 0$, we glue together \mathcal{G}^1 and \mathcal{G}^2 by taking a new edge of length 1, attaching its first endpoint to the point $x = L$ on a half-line h^1 of \mathcal{G}^1 and its second endpoint to the point $x = L$ on a half-line h^2 of \mathcal{G}^2 . We call \mathcal{G}_L the resulting graph (see Figure 7) and we let the set Z_L of vertices of degree 1 in \mathcal{G}_L with homogeneous Dirichlet conditions be given by the union of the corresponding sets of vertices in \mathcal{G}^1 and \mathcal{G}^2 .

Lemma 4.15. *Let $\lambda > 0$ and $\mathcal{G}^1, \mathcal{G}^2, \mathcal{G}_L$ be the graphs described above. Then*

$$\lim_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) = \min\left(\inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v), \inf_{v \in \mathcal{N}_{Z^2}(\mathcal{G}^2)} J(v)\right).$$

Proof. Without loss of generality, assume that

$$c_1 := \inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v) \leq \inf_{v \in \mathcal{N}_{Z^2}(\mathcal{G}^2)} J(v) =: c_2. \quad (4-8)$$

For every $\varepsilon > 0$ there exists a function $u_\varepsilon \in \mathcal{N}_{Z^1}(\mathcal{G}^1)$ with compact support such that $J(u_\varepsilon) \leq c_1 + \varepsilon$. For every L large enough, we can view u_ε as a function in $\mathcal{N}_{Z_L}(\mathcal{G}_L)$, after extending it to zero on \mathcal{G}_L outside its support. Therefore

$$\limsup_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) \leq \limsup_{L \rightarrow \infty} J(u_\varepsilon) = J(u_\varepsilon) \leq c_1 + \varepsilon$$

and since ε is arbitrary we deduce that

$$\limsup_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) \leq c_1. \quad (4-9)$$

We now prove a complementary inequality. For every L , let $u_L \in \mathcal{N}_{Z_L}(\mathcal{G}_L)$ satisfy

$$J(u_L) \leq \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) + \frac{1}{L} \quad (4-10)$$

and notice that by (2-1), (4-9) and (4-10), u_L is bounded independently of L . Let

$$u_L(x_L) = \min_{h^1 \cap [0, L]} u_L(x), \quad u_L(y_L) = \min_{h^2 \cap [0, L]} u_L(x)$$

and set

$$\delta_L = \max(u_L(x_L), u_L(y_L)).$$

Since u_L is uniformly bounded in $L^2(\mathcal{G}_L)$, $\delta_L \rightarrow 0$ as $L \rightarrow \infty$.

Consider the function $(u_L - \delta_L)^+ \in H_{Z_L}^1(\mathcal{G}_L)$, which does not vanish identically, for all L large, by Proposition 2.4. Exactly as in (4-6), $n_\lambda((u_L - \delta_L)^+) \leq 1 + o(1)$ as $L \rightarrow \infty$.

Set $v_L = \pi_\lambda((u_L - \delta_L)^+)$. Now $v_L \in \mathcal{N}_{Z_L}(\mathcal{G}_L)$, it vanishes at x_L, y_L and

$$J(v_L) = \kappa n_\lambda((u_L - \delta_L)^+)^p \|(u_L - \delta_L)^+\|_p^p \leq \kappa(1 + o(1)) \|u_L\|_p^p = J(u_L) + o(1) \quad (4-11)$$

as $L \rightarrow \infty$.

We now cut \mathcal{G}_L at x_L and y_L , splitting it into three parts $\bar{\mathcal{G}}^1 \subseteq \mathcal{G}^1$, $\bar{\mathcal{G}}^2 \subseteq \mathcal{G}^2$ and $\mathcal{G}^3 = \mathcal{G}_L \setminus (\bar{\mathcal{G}}^1 \cup \bar{\mathcal{G}}^2)$. We call v_i ($i = 1, 2$) the two vertices of \mathcal{G}^3 created on h^i (see Figure 8).

We can use v_L to construct a function $v_L^1 \in H_{Z^1}^1(\mathcal{G}^1)$ by setting

$$v_L^1(x) = \begin{cases} v_L(x) & \text{if } x \in \bar{\mathcal{G}}^1, \\ 0 & \text{elsewhere on } \mathcal{G}^1 \end{cases}$$

and in the same way we construct a function $v_L^2 \in H_{Z^2}^1(\mathcal{G}^2)$. Finally, we call v_L^3 the restriction of v_L to \mathcal{G}^3 . Setting $Z^3 = \{v_1, v_2\}$, by construction we have $v_L^3 \in H_{Z^3}^1(\mathcal{G}^3)$.




Figure 8. The graph \mathcal{G}_L of Figure 7 splits into the three graphs $\bar{\mathcal{G}}^1$, $\bar{\mathcal{G}}^2$ and \mathcal{G}^3 .

If $v_L^i \neq 0$, then there exists $\theta_i \in \mathbb{R}$ so that $v_L^i \in \mathcal{N}_{\theta_i, Z^i}(\mathcal{G}^i)$. Taking $\theta_i = 0$ if $v_L^i = 0$ and recalling that $v_L \in \mathcal{N}_{Z_L}(\mathcal{G}_L)$, we obtain

$$\lambda = \frac{\|v_L^1\|_2^2}{\|v_L\|_2^2} \theta_1 + \frac{\|v_L^2\|_2^2}{\|v_L\|_2^2} \theta_2 + \frac{\|v_L^3\|_2^2}{\|v_L\|_2^2} \theta_3. \quad (4-12)$$

Furthermore,

$$J(v_L) = \kappa(\|v_L^1\|_p^p + \|v_L^2\|_p^p + \|v_L^3\|_p^p) \geq \kappa \max\{\|v_L^1\|_p^p, \|v_L^2\|_p^p, \|v_L^3\|_p^p\}.$$

Since, by (4-12), λ is a convex combination of the θ_i 's, at least one of the three will satisfy $\theta_i \geq \lambda$.

If $\theta_1 \geq \lambda$, by Remark 2.5, we have $\kappa \|v_L^1\|_p^p \geq c_1$. If $\theta_2 \geq \lambda$, by (4-8), we have $\kappa \|v_L^2\|_p^p \geq c_2 \geq c_1$. If $\theta_3 \geq \lambda$, we have $\kappa \|v_L^3\|_p^p \geq \inf_{\mathcal{N}_{\lambda, Z^3}(\mathcal{G}^3)} J \geq s_\lambda \geq c_1$ since \mathcal{G}^3 satisfies (H0) and by the assumptions on \mathcal{G}^1 . In each case we deduce, via (4-11), that

$$J(u_L) \geq J(v_L) + o(1) \geq c_1 + o(1)$$

as $L \rightarrow \infty$, so that by (4-10),

$$\liminf_L \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) \geq \liminf_L \left(J(u_L) - \frac{1}{L} \right) \geq c_1.$$

In view of (4-9), this ends the proof. \square

Theorem 4.16. *Let $\lambda > 0$ and \mathcal{G}^1 , \mathcal{G}^2 and \mathcal{G}_L be the graphs considered above. If L is large enough, then there exist nodal ground states on \mathcal{G}_L .*

Proof. Without loss of generality, we assume that

$$\min \left(\inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v), \inf_{v \in \mathcal{N}_{Z^2}(\mathcal{G}^2)} J(v) \right) = \inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v).$$

Let $\varepsilon := \frac{1}{3}(s_\lambda - \inf_{v \in \mathcal{N}_{Z^2}(\mathcal{G}^2)} J(v))$. By Lemma 4.15, we choose L so large that

$$\inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) \geq \inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v) - \varepsilon \quad (4-13)$$

and that there exist nonnegative $u^i \in \mathcal{N}_{Z^i}(\mathcal{G}^i)$, with compact support satisfying

$$J(u^i) \leq \inf_{v \in \mathcal{N}_{Z^i}(\mathcal{G}^i)} J(v) + \varepsilon.$$




Figure 9. A graph with $\#F(\mathcal{G}) = 2$ where nodal ground states never exist, independently of the length of the edges.

In particular, there is $M > 0$ such that the restriction of u^i to each half-line of \mathcal{G}^i vanishes on $[M, +\infty)$. Hence, for every $L \geq M$, we define $w : \mathcal{G}_L \rightarrow \mathbb{R}$ as

$$w(x) := \begin{cases} u^1(x) & \text{if } x \in \mathcal{G}^1, \\ -u^2(x) & \text{if } x \in \mathcal{G}^2, \\ 0 & \text{elsewhere on } \mathcal{G}_L, \end{cases}$$

where with a slight abuse of notation we still denote by $\mathcal{G}^1, \mathcal{G}^2$ the corresponding subgraphs of \mathcal{G}_L . Clearly, $w \in \mathcal{M}_{Z_L}(\mathcal{G}_L)$ and, by (4-13) and the choice of ε , we have

$$\inf_{v \in \mathcal{M}_{Z_L}(\mathcal{G}_L)} J(v) \leq J(w) = J(u^1) + J(u^2) \leq \inf_{v \in \mathcal{N}_{Z^1}(\mathcal{G}^1)} J(v) + \inf_{v \in \mathcal{N}_{Z^2}(\mathcal{G}^2)} J(v) + 2\varepsilon < \inf_{v \in \mathcal{N}_{Z_L}(\mathcal{G}_L)} J(v) + s_\lambda,$$

in turn implying existence of nodal ground states by Theorem 1.3 and Proposition 4.6. □

Remark 4.17. Concrete examples of graphs fulfilling the hypotheses of Theorem 4.16 can be produced starting, for instance, from any of the graphs in Figure 6 (see, e.g., Figure 7). Since $\inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v) < s_\lambda$ implies $\#F(\mathcal{G}) \geq 1$, by construction we have $\#F(\mathcal{G}_L) \geq 2$.

Remark 4.18. Theorems 1.6(ii) and 4.11 show that, if (H1) holds, or if the set of all bounded edges of \mathcal{G} is too small, nodal ground states never exist, whereas Theorem 4.16 proves that there exist graphs with $\#F(\mathcal{G}) \geq 2$ and a sufficiently large compact core where nodal ground states do exist. However, even though the former provides sufficient conditions for nonexistence, the latter are not sufficient conditions for existence. It is in fact not difficult to produce examples of graphs with $\#F(\mathcal{G}) = 2$, and compact core of arbitrary size, where nodal ground states do not exist. For instance, consider the graph in Figure 9. If u is a nodal ground state on this graph, we know that either u is identically equal to zero on the two half-lines or it has constant sign on them. Assume thus that $u \geq 0$ on the half-lines. Then, since u^+ is not identically zero, u^+ vanishes on the set \mathcal{B} of the bounded edges of \mathcal{G} at least at a point different from the vertex of the half-lines. We then have that u^+ has always at least two preimages for every $t \in (0, \max u^+)$ by Theorem 6.1 and hence $J(u^+) \geq s_\lambda$ by Lemma 4.5. In view of Remark 3.1, as $u^- \in \mathcal{N}_Z(\mathcal{G})$ vanishes somewhere on the graph, we have also that $J(u^-) > \inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v)$. This implies that

$$J(u) = J(u^-) + J(u^+) > \inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v) + s_\lambda,$$

which contradicts (1-9).

5. Graphs with infinitely many bounded edges

In this section we extend our discussion about ground states and nodal ground states to graphs with infinitely many edges whose length is uniformly bounded. In particular, we focus on two subclasses of such graphs that have already been considered in the literature: periodic graphs and regular trees.

5.1. Periodic graphs. Throughout this section, when we speak of a periodic metric graph we mean a graph fulfilling [Berkolaiko and Kuchment 2013, Definition 4.1.1]. We avoid reporting the full details of the definition here. For our purposes, it is enough to recall that, if \mathcal{G} is a periodic graph, then there exists a number $n \in \mathbb{N}$ and a compact subset $W \subset \mathcal{G}$, called a *fundamental domain* of \mathcal{G} , such that

$$\mathcal{G} = \bigcup_{k \in \mathbb{Z}^n} W_k,$$

where W_k is a copy of W for every $k \in \mathbb{Z}^n$, and $W_i \cap W_j$ contains at most finitely many points for every $i \neq j$. In this case, we say that \mathcal{G} is a \mathbb{Z}^n -periodic graph.

Since we are concerned with problems involving homogeneous Dirichlet conditions on a subset Z of the vertices of \mathcal{G} , we specify that when \mathcal{G} is a \mathbb{Z}^n -periodic graph, we only consider here \mathbb{Z}^n -periodic subsets Z (that is, $Z \cap W_k$ is a copy of $Z \cap W$ for every $k \in \mathbb{Z}^n$).

Proof of Theorem 1.7. We address independently the case of ground states and nodal ground states.

Part 1: existence of ground states. Let $(u_n)_n \subset \mathcal{N}_Z$ be such that $\lim_n J(u_n) = \inf_{v \in \mathcal{N}_Z} J(v)$. Exploiting the periodicity of \mathcal{G} and Z , we can assume with no loss of generality that u_n attains its L^∞ -norm on W_0 , for every n . Since $(u_n)_n$ is bounded in $H^1(\mathcal{G})$, up to subsequences $u_n \rightharpoonup u$ in $H^1(\mathcal{G})$ and $u_n \rightarrow u$ in $L^\infty_{\text{loc}}(\mathcal{G})$ as $n \rightarrow \infty$. Furthermore, $u \not\equiv 0$ on \mathcal{G} because, if this were not the case, by the strong convergence of (u_n) to u in $L^\infty(W_0)$ we would have $\|u_n\|_{L^\infty(\mathcal{G})} = \|u_n\|_{L^\infty(W_0)} \rightarrow 0$ as $n \rightarrow \infty$, contradicting Proposition 2.4.

If $u_n \rightarrow u$ in $L^2(\mathcal{G})$, then by standard Gagliardo–Nirenberg inequalities $u_n \rightarrow u$ in $L^p(\mathcal{G})$ and, by weak lower semicontinuity, $n_\lambda(u) \leq 1$, so that

$$\inf_{v \in \mathcal{N}_Z} J(v) \leq J(\pi_\lambda(u)) = \kappa n_\lambda(u)^p \|u\|_p^p \leq \lim_n \kappa \|u_n\|_p^p = \inf_{v \in \mathcal{N}_Z} J(v),$$

i.e., $\pi_\lambda(u)$ is a ground state.

Let us thus show that u_n converges to u strongly in $L^2(\mathcal{G})$. Assume by contradiction that

$$\liminf_n \|u_n - u\|_2^2 > 0.$$

Let $\theta \in \mathbb{R}$ and $(\lambda_n)_n \subset \mathbb{R}$ be such that $u \in \mathcal{N}_{\theta, Z}$, $u_n - u \in \mathcal{N}_{\lambda_n, Z}$ for every n . By the weak convergence of $(u_n)_n$ to u in $H^1(\mathcal{G})$, Brézis–Lieb lemma [1983] and the fact that $u_n \in \mathcal{N}_{\lambda_n, Z}$, we have

$$\begin{aligned} \lambda_n &= \frac{\|u_n - u\|_p^p - \|u'_n - u'\|_2^2}{\|u_n - u\|_2^2} = \frac{\|u_n\|_p^p - \|u'_n\|_2^2 - \|u\|_p^p + \|u'\|_2^2 + o(1)}{\|u_n - u\|_2^2} \\ &= \frac{\lambda \|u_n\|_2^2 - \theta \|u\|_2^2 + o(1)}{\|u_n - u\|_2^2} = \lambda + \frac{(\lambda - \theta) \|u\|_2^2 + o(1)}{\|u_n - u\|_2^2} = \lambda + \frac{\|u\|_2^2}{\|u_n - u\|_2^2} (\lambda - \theta) + o(1) \end{aligned} \quad (5-1)$$

as $n \rightarrow \infty$. Applying again Brézis–Lieb lemma, we obtain

$$\inf_{v \in \mathcal{N}_{\lambda, Z}} J_{\lambda}(v) = \lim_n \kappa \|u_n\|_p^p = \lim_n \kappa \|u_n - u\|_p^p + \kappa \|u\|_p^p. \tag{5-2}$$

Keeping in mind that $\lambda > 0$, we distinguish three cases. If $\theta > \lambda$, then, by Remark 2.5,

$$\lim_n \kappa \|u_n - u\|_p^p + \kappa \|u\|_p^p \geq \kappa \|u\|_p^p \geq \inf_{v \in \mathcal{N}_{\theta, Z}} J_{\theta}(v) > \inf_{v \in \mathcal{N}_{\lambda, Z}} J_{\lambda}(v).$$

If $\theta = \lambda$, we see from (5-1) that $\lambda_n \rightarrow \lambda$ as $n \rightarrow \infty$. Therefore, using again Remark 2.5,

$$\lim_n \kappa \|u_n - u\|_p^p + \kappa \|u\|_p^p \geq \lim_n \inf_{v \in \mathcal{N}_{\lambda_n, Z}} J_{\lambda_n}(v) + \inf_{v \in \mathcal{N}_{\theta, Z}} J_{\theta}(v) = 2 \inf_{v \in \mathcal{N}_{\lambda, Z}} J_{\lambda}(v).$$

If $\theta < \lambda$, we have $\liminf_n \lambda_n > \lambda$ and, similarly,

$$\lim_n \kappa \|u_n - u\|_p^p + \kappa \|u\|_p^p \geq \lim_n \inf_{v \in \mathcal{N}_{\lambda_n, Z}} J_{\lambda_n}(v) > \inf_{v \in \mathcal{N}_{\lambda, Z}} J_{\lambda}(v).$$

In all three cases, (5-2) yields a contradiction.

Part 2: nonexistence of nodal ground states. By Theorem 1.4, it is enough to show that

$$\inf_{v \in \mathcal{M}_Z} J(v) \leq 2 \inf_{v \in \mathcal{N}_Z} J(v).$$

To this end, given $\varepsilon > 0$, let $u \in \mathcal{N}_Z$ be such that $J(u) \leq \inf_{v \in \mathcal{N}_Z} J(v) + \varepsilon$. With no loss of generality, we can take such u to be nonnegative, compactly supported on \mathcal{G} and attaining its L^∞ -norm on W_0 (it is, for instance, enough to apply Remark 3.2 to a suitable ground state of J in \mathcal{N}_Z , that exists by the first part of the proof). Hence, there exists $M > 0$ such that $\text{supp}(u) \subset \bigcup_{|k| \leq M} W_k$. Let then $\bar{u} \in \mathcal{N}_Z$ be a translation of u on \mathcal{G} such that $\text{supp}(\bar{u}) \subset \bigcup_{|k| > M} W_k$ and define $w : \mathcal{G} \rightarrow \mathbb{R}$ as

$$w(x) := \begin{cases} u(x) & \text{if } x \in \text{supp}(u), \\ -\bar{u}(x) & \text{if } x \in \text{supp}(\bar{u}), \\ 0 & \text{elsewhere on } \mathcal{G}. \end{cases}$$

By construction, $w \in \mathcal{M}_Z$ and $J(w) = J(u) + J(\bar{u}) \leq 2 \inf_{v \in \mathcal{N}_Z} J(v) + 2\varepsilon$. Given the arbitrariness of $\varepsilon > 0$, we conclude. □

Remark 5.1. Observe that $J^\infty(\mathcal{G}; Z) = \inf_{v \in \mathcal{N}_Z} J(v)$ for every periodic graph \mathcal{G} and every set Z with the same periodicity. This is the reason why it is not possible to rely directly on the abstract result of Theorem 1.3 to prove Theorem 1.7.

5.2. Regular trees. Recall that a regular tree is an acyclic, noncompact metric graph with edges all of the same length and vertices all of the same degree $d \geq 3$ (unrooted tree), with the possible exception of a single vertex of degree 1 (rooted tree). If \mathcal{G} is an unrooted tree, then necessarily $Z = \emptyset$ since every vertex has degree at least 3, whereas if \mathcal{G} is a rooted tree either $Z = \emptyset$ or it coincides with the root of \mathcal{G} (i.e., the unique vertex of degree 1).

We divide the proof of Theorem 1.8 into two parts, proving first statements (i)–(ii) on ground states and then statement (iii) on nodal ground states.

Proof of Theorem 1.8. We split the proof in several steps.

Step 1: ground states when \mathcal{G} is an unrooted tree. Let $(u_n)_n \subset \mathcal{N}$ be such that $\lim_n J(u_n) = \inf_{v \in \mathcal{N}} J(v)$. Exploiting the symmetry of \mathcal{G} , it is not restrictive to assume that u_n attains its L^∞ -norm in the same fixed edge of \mathcal{G} , for every n . Indeed, the problem is invariant under any isometry of \mathcal{G} and the isometry group of the tree \mathcal{G} acts transitively on the edges of \mathcal{G} . Hence, arguing as in the proof of Theorem 1.7 shows that the weak limit in $H^1(\mathcal{G})$ of $(u_n)_n$ provides a desired ground state for J in \mathcal{N} .

Step 2: ground states when \mathcal{G} is a rooted tree and $Z = \emptyset$. Let r be the root of \mathcal{G} , $d \geq 3$ be the degree of each vertex of \mathcal{G} different from the root, and $\bar{\mathcal{G}}$ be the unrooted tree obtained by gluing together d copies of \mathcal{G} at their roots. We first prove that

$$J^\infty(\mathcal{G}) = \inf_{v \in \mathcal{N}_{\{r\}}(\mathcal{G})} J(v) = \inf_{v \in \mathcal{N}(\bar{\mathcal{G}})} J(v). \quad (5-3)$$

To this aim, given any function $u \in \mathcal{N}_{\{r\}}(\mathcal{G})$, we construct a sequence $(u_n)_n \subset \mathcal{N}_{\{r\}}(\mathcal{G})$ converging weakly to 0 in $H^1(\mathcal{G})$ by translating u along \mathcal{G} and extending it by 0 on the remaining part of the graph. This proves that $J^\infty(\mathcal{G}) \leq \inf_{v \in \mathcal{N}_{\{r\}}(\mathcal{G})} J(v)$. Next, let $(u_n)_n \subset \mathcal{N}(\bar{\mathcal{G}})$ be a sequence converging weakly to 0 in $H^1(\bar{\mathcal{G}})$ and such that $J(u_n) \rightarrow J^\infty(\mathcal{G})$. Since $u_n(r) \rightarrow 0$ by $L^\infty_{\text{loc}}(\bar{\mathcal{G}})$ convergence, we can assume without loss of generality that each u_n satisfies $u_n(r) = 0$, namely that $u_n \in \mathcal{N}_{\{r\}}(\mathcal{G})$. This shows that $J^\infty(\mathcal{G}) \geq \inf_{v \in \mathcal{N}_{\{r\}}(\mathcal{G})} J(v)$, and the first equality is proved.

To prove the second equality, notice that any $u \in \mathcal{N}_{\{r\}}(\mathcal{G})$ can be seen as an element of $\mathcal{N}(\bar{\mathcal{G}})$, after extending it by 0 on $\bar{\mathcal{G}} \setminus \mathcal{G}$. On the other hand, any $u \in \mathcal{N}(\bar{\mathcal{G}})$ with compact support can be considered (when translated in such a way that $r \notin \text{supp}(u)$) as an element of $\mathcal{N}_{\{r\}}(\mathcal{G})$. By density this is enough to conclude the proof of (5-3).

By (5-3) and Theorem 1.3, in order to prove the existence of a ground state, it is sufficient to show that

$$\inf_{v \in \mathcal{N}(\mathcal{G})} J(v) < \inf_{v \in \mathcal{N}(\bar{\mathcal{G}})} J(v). \quad (5-4)$$

To this end, let $u \in \mathcal{N}(\bar{\mathcal{G}})$ be a positive ground state of J in $\mathcal{N}(\bar{\mathcal{G}})$, whose existence is guaranteed by the previous step. Take a vertex v of $\bar{\mathcal{G}}$ and split the graph at v into d disjoint rooted trees \mathcal{G}_i , $i = 1, \dots, d$. For every i , let $u_i > 0$ be the restriction of u to \mathcal{G}_i and $\lambda_i \in \mathbb{R}$ be such that $u_i \in \mathcal{N}_{\lambda_i}(\mathcal{G}_i)$. Since $u \in \mathcal{N}(\bar{\mathcal{G}})$ and $u > 0$ on $\bar{\mathcal{G}}$, we have

$$\lambda = \sum_{i=1}^d \frac{\|u_i\|_{L^2(\mathcal{G}_i)}^2}{\|u\|_{L^2(\bar{\mathcal{G}})}^2} \lambda_i,$$

so that

$$\lambda \leq \left(\max_{1 \leq i \leq d} \lambda_i \right) \sum_{i=1}^d \frac{\|u_i\|_{L^2(\mathcal{G}_i)}^2}{\|u\|_{L^2(\bar{\mathcal{G}})}^2} = \max_{1 \leq i \leq d} \lambda_i.$$

Hence, there exists $j \in \{1, \dots, d\}$ such that $n_\lambda(u_j) \leq 1$. Since each \mathcal{G}_i is a copy of \mathcal{G} , we then have

$$\inf_{v \in \mathcal{N}(\mathcal{G})} J(v) \leq J(\pi_\lambda(u_j)) = \kappa n_\lambda(u_j)^p \|u_j\|_{L^p(\mathcal{G}_j)}^p < \kappa \sum_{i=1}^d \|u_i\|_{L^p(\mathcal{G}_i)}^p = \kappa \|u\|_{L^p(\bar{\mathcal{G}})}^p = \inf_{v \in \mathcal{N}(\bar{\mathcal{G}})} J(v),$$

that is (5-4).

Step 3: ground states when \mathcal{G} is a rooted tree and $Z \neq \emptyset$. Since Z coincides with the root of \mathcal{G} , as in the first part of Step 2 we have

$$\inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v) = \inf_{v \in \mathcal{N}(\bar{\mathcal{G}})} J(v),$$

where $\bar{\mathcal{G}}$ is the unrooted tree corresponding to \mathcal{G} as above. That the problem on \mathcal{G} has no ground state follows by the fact that, if u were a ground state in $\mathcal{N}_Z(\mathcal{G})$, it would be also a ground state in $\mathcal{N}(\bar{\mathcal{G}})$, as any function on \mathcal{G} vanishing at the root can be regarded as a function on $\bar{\mathcal{G}}$ as well after extending it by 0. Since this is impossible because ground states never vanish on $\bar{\mathcal{G}}$, we conclude.

Step 4: nodal ground states when \mathcal{G} is an unrooted tree or \mathcal{G} is a rooted tree and $Z \neq \emptyset$. If \mathcal{G} is an unrooted tree, exploiting again its symmetry, it is easy to adapt the argument developed in the proof of Theorem 1.7 to show again that

$$\inf_{v \in \mathcal{M}(\mathcal{G})} J(v) = 2 \inf_{v \in \mathcal{N}(\mathcal{G})} J(v),$$

and likewise, if \mathcal{G} is a rooted tree with $Z \neq \emptyset$, that

$$\inf_{v \in \mathcal{M}_Z(\mathcal{G})} J(v) = 2 \inf_{v \in \mathcal{N}_Z(\mathcal{G})} J(v).$$

This implies that nodal ground states do not exist by Theorem 1.4.

Step 5: nodal ground states when \mathcal{G} is a rooted tree and $Z = \emptyset$. Since, given any $u \in \mathcal{M}$, at least one between u^+ and u^- vanishes at the root r , it follows that

$$\inf_{v \in \mathcal{M}(\mathcal{G})} J(v) = \inf_{v \in \mathcal{N}(\mathcal{G})} J(v) + \inf_{v \in \mathcal{N}_{\{r\}}(\mathcal{G})} J(v).$$

Arguing as in the proof of Theorem 1.4, this immediately implies that nodal ground states do not exist. \square

6. Qualitative properties of nodal ground states

The first result of this section concerns the *nodal domains* (i.e., the connected components of $\mathcal{G} \setminus u^{-1}(0)$) of any minimizer u in $\mathcal{M}_{\lambda, Z}(\mathcal{G})$.

Theorem 6.1. *Let $\mathcal{G} \in \mathbf{G}$ and $\lambda > -\omega_Z(\mathcal{G})$. Let $u \in \mathcal{M}_Z$ be a nodal ground state. Then u has exactly two nodal domains.*

Proof. Assume for contradiction that there are at least three nodal domains. Up to a change of sign, we can make sure that on at least two of them u is positive, and we call \mathcal{G}_1 one of the two. Since u solves (1-1), multiplying by u and integrating on \mathcal{G}_1 we have

$$\int_{\mathcal{G}_1} (|u'|^2 + \lambda|u|^2 - |u|^p) dx = uu'|_{\partial\mathcal{G}_1} + \int_{\mathcal{G}_1} (-u'' + \lambda u - |u|^{p-2}u)u dx = 0, \tag{6-1}$$

because on $\partial\mathcal{G}_1$ either $u = 0$ or $u' = 0$ (this happens at vertices of degree 1 not in Z).

Now we define $v \in H_Z^1(\mathcal{G})$ by

$$v(x) := \begin{cases} u(x) & \text{if } x \in \mathcal{G} \setminus \mathcal{G}_1, \\ 0 & \text{if } x \in \mathcal{G}_1, \end{cases}$$

and we observe that $v^- = u^- \in \mathcal{N}_Z$ and that v^+ (not identically zero by construction) satisfies

$$\int_{\mathcal{G}} (|(v^+)'|^2 + \lambda|v^+|^2 - |v^+|^p) dx = \int_{\mathcal{G}} (|(u^+)'|^2 + \lambda|u^+|^2 - |u^+|^p) dx - \int_{\mathcal{G}_1} (|u'|^2 + \lambda|u|^2 - |u|^p) dx = 0$$

by (6-1) and because $u^+ \in \mathcal{N}_Z$. Therefore $v \in \mathcal{M}_Z$ and

$$J(v) = \kappa \|v\|_{L^p(\mathcal{G})}^p = \kappa \|u\|_{L^p(\mathcal{G})}^p - \kappa \|u\|_{L^p(\mathcal{G}_1)}^p < \kappa \|u\|_{L^p(\mathcal{G})}^p = J(u),$$

violating the minimality of u . □

To conclude we are left to prove Theorem 1.9. To this end, we will actually prove three independent statements, the full proof of Theorem 1.9 then following by their combination. Each of these statements exhibits a graph supporting a nodal ground state with nodal set respectively given by

- (1) k isolated points;
- (2) $m \geq 2$ half-lines all attached to the same vertex;
- (3) n line segments all attached to the same vertex.

These three constructions, though mutually independent, can all be carried out on the same kind of graph, that we now describe. Given $N \in \mathbb{N}$ and $L > 0$, let v_1, v_2 be two vertices joined by N edges e_1, \dots, e_N , each of length L . Attach to v_1 a pendant and a half-line and do the same to v_2 . In this way we obtain the graph $\mathcal{G}_{N,L}$ depicted in Figure 10. Throughout, we fix $\lambda > 0$ and the length of the two pendants so that nodal ground states in $\mathcal{M}_\lambda(\mathcal{G}_{N,L})$ exist (independently of any other feature of the graph), which is possible by Theorem 4.12.

Proof of (1). Here we show that, for a suitable choice of L , the graph $\mathcal{G}_{k,L}$ admits a nodal ground state u such that $u^{-1}(0)$ consists of k isolated points.

Proposition 6.2. *For every $k \in \mathbb{N}$ there exists $\bar{L} > 0$, depending on λ and k , such that, for every $L \geq \bar{L}$, every nodal ground state u on $\mathcal{G}_{k,L}$ has a nodal set of the form $u^{-1}(0) = \{x_1, \dots, x_k\}$, where x_i belongs to the interior of the edge e_i .*

To prove this proposition we consider also the graph $\bar{\mathcal{G}}_{k+1}$ made of $k+1$ half-lines and a pendant all attached at the same vertex. The length of the pendant of $\bar{\mathcal{G}}_{k+1}$ coincides with that of the two pendants of $\mathcal{G}_{k,L}$. Hence, by Theorem 4.12, ground states exist in $\mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})$.

Lemma 6.3. *For every $k \in \mathbb{N}$,*

$$\lim_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v) = \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v).$$

Proof. The argument is similar to that in the proof of Lemma 4.15. Using suitable compactly supported functions in $\mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})$, one immediately checks that

$$\limsup_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v) \leq \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v).$$

To show that

$$\liminf_{L \rightarrow \infty} \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v) \geq \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v), \tag{6-2}$$




Figure 10. The graph $\mathcal{G}_{N,L}$ with $N = 5$.

it is enough to note that, if $u_L \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})$ satisfies

$$J(u_L) \leq \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v) + \frac{1}{L},$$

then

$$\max_{1 \leq i \leq k} \min_{x \in e_i} |u_L(x)| \rightarrow 0 \quad \text{as } L \rightarrow \infty.$$

This allows one to obtain (6-2) working exactly as in the proof of Lemma 4.15. □

Lemma 6.4. *If $L \rightarrow \infty$, then*

$$\limsup_{L \rightarrow \infty} \inf_{v \in \mathcal{M}_\lambda(\mathcal{G}_{k,L})} J(v) \leq 2 \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v).$$

Proof. The proof follows the same lines as the one of Theorem 4.16. □

Proof of Proposition 6.2. Let u be a nodal ground state in $\mathcal{M}_\lambda(\mathcal{G}_{k,L})$.

Step 1: for L long enough, either $u \equiv 0$ on the pendants or it has no zero on their closure. Assume by contradiction that $u \not\equiv 0$ on a pendant p , but $u(x_0) = 0$ for some x_0 on p . With no loss of generality, let $u > 0$ at the vertex of degree 1 of p . Outside p , $u < 0$ thanks to Theorem 6.1. Denoting as usual by $|p|$ the length of p , since u is a solution to (1-1), we have $u^+ \in \mathcal{N}_\lambda(0, |p|)$ with $u^+(|p|) = 0$, so that

$$J(u^+) \geq \inf_{\substack{v \in \mathcal{N}_\lambda(0, |p|) \\ v(|p|)=0}} J(v) > \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v).$$

This is because the pendant of $\bar{\mathcal{G}}_{k+1}$ can be identified with the interval $[0, |p|]$, but u^+ is not a ground state in $\mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})$ as ground states never vanish.

Letting then

$$\delta := \inf_{\substack{v \in \mathcal{N}_\lambda(0, |p|) \\ v(|p|)=0}} J(v) - \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v) > 0$$

and recalling that $u^- \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})$, it follows that

$$\inf_{v \in \mathcal{M}_\lambda(\mathcal{G}_{k,L})} J(v) = J(u) = J(u^-) + J(u^+) \geq \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v) + \inf_{v \in \mathcal{N}_\lambda(\bar{\mathcal{G}}_{k+1})} J(v) + \delta,$$

contradicting Lemmas 6.3–6.4 for L large enough.

Step 2: $u(v_1)u(v_2) < 0$. Assume that this is not the case. Since u solves (1-1), on any of the two half-lines either $u \equiv 0$ or it never vanishes. Combining with Step 1, this implies that there exist $i \in \{1, \dots, k\}$ and $\bar{x}_1, \bar{x}_2 \in e_i \cup \{v_1, v_2\}$ with $u(\bar{x}_1) = u(\bar{x}_2) = 0$ and, for all $x \in (\bar{x}_1, \bar{x}_2)$, $u \neq 0$. Without loss of generality, let $u > 0$ on (\bar{x}_1, \bar{x}_2) . By Theorem 6.1, we know that $u < 0$ on the remaining part of the graph and $u^- \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})$, while we can think of u^+ as a function in $\mathcal{N}_\lambda(\mathbb{R})$ with compact support. Hence we have

$$\inf_{v \in \mathcal{M}_\lambda(\mathcal{G}_{k,L})} J(v) = J(u^+) + J(u^-) > s_\lambda + \inf_{v \in \mathcal{N}_\lambda(\mathcal{G}_{k,L})} J(v),$$

which contradicts Theorem 1.6.

Step 3: conclusion. The previous steps ensure that $u^{-1}(0) \subset \bigcup_{i=1}^N e_i$ and that it is a finite union of points by uniqueness of the solution of the Cauchy problem for (1-1). The uniqueness of the zero of u on each e_i follows then by Theorem 6.1. \square

Proof of (2). Here we prove the following result.

Proposition 6.5. *Let $m \geq 2$. There exists a graph $\bar{\mathcal{G}}$ that admits a nodal ground state u such that $u^{-1}(0)$ is the union of $m \geq 2$ half-lines attached at the same vertex.*

The graph $\bar{\mathcal{G}}$ is obtained from the graph $\mathcal{G}_{1,L}$ by attaching m half-lines at a suitable point. Before proving Proposition 6.5 we establish the following lemma.

Lemma 6.6. *Let \mathcal{G} be a noncompact graph with a finite number of edges. Let $\tilde{\mathcal{G}}$ be a graph obtained from \mathcal{G} by attaching $m \geq 2$ half-lines h_1, \dots, h_m at one of its points p . If there exists a nodal ground state in $\mathcal{M}_\lambda(\tilde{\mathcal{G}})$, then*

$$\inf_{v \in \mathcal{M}_\lambda(\tilde{\mathcal{G}})} J(v) \geq \inf_{v \in \mathcal{M}_\lambda(\mathcal{G})} J(v). \quad (6-3)$$

Proof. Let \tilde{u} be a nodal ground state on $\tilde{\mathcal{G}}$ and assume without loss of generality that $\tilde{u}(p) \geq 0$. Denote by u the restriction of \tilde{u} on \mathcal{G} and by ϕ_i the restriction of \tilde{u} to the half-line h_i for $i = 1, \dots, m$.

If $\tilde{u}(p) = 0$, since \tilde{u} solves (1-1), each ϕ_i vanishes identically. Hence, $u \in \mathcal{M}_\lambda(\mathcal{G})$, $J(\tilde{u}) = J(u)$ and (6-3) follows.

If $\tilde{u}(p) > 0$, each ϕ_i coincides with a portion of the same soliton ϕ_λ . With a slight abuse of notation we denote by $\phi'_i(p)$ the outgoing derivative of ϕ_i at p along h_i . Note that $\phi'_i(p) < 0$ for every i . Indeed, if on the contrary we had, for instance, $\phi'_1(p) \geq 0$, then the restriction of \tilde{u} to the union of the h_i 's would contain at least one full soliton ϕ_λ , so that $\|\tilde{u}\|_{L^p(\bigcup_i h_i)}^p \geq \|\phi_\lambda\|_p^p$. This would lead to

$$\inf_{v \in \mathcal{M}_\lambda(\tilde{\mathcal{G}})} J(v) = J(\tilde{u}) = \kappa(\|\tilde{u}^+\|_p^p + \|\tilde{u}^-\|_p^p) > \kappa\|\phi_\lambda\|_p^p + \inf_{v \in \mathcal{N}_\lambda(\tilde{\mathcal{G}})} J(v) = s_\lambda + \inf_{v \in \mathcal{N}_\lambda(\tilde{\mathcal{G}})} J(v),$$

which contradicts (1-9).

As, for all $i \in \{1, \dots, m\}$, ϕ_i is a solution to (1-1), we have in particular $\phi_i \in \mathcal{N}_{\theta_i}(h_i)$ with

$$\theta_i = \frac{\int_{h_i} \phi_i^p dx - \int_{h_i} (\phi_i')^2 dx}{\int_{h_i} \phi_i^2 dx} = \frac{\int_{h_i} \phi_i^p dx + \phi_i(p)\phi_i'(p) + \int_{h_i} \phi_i'' \phi_i dx}{\int_{h_i} \phi_i^2 dx} = \frac{\lambda \int_{h_i} \phi_i^2 dx + \phi_i(p)\phi_i'(p)}{\int_{h_i} \phi_i^2 dx} < \lambda$$

since ϕ_i is a portion of ϕ_λ and $\phi'_i(\mathfrak{p}) < 0$. Letting then μ be the number such that $u^+ \in \mathcal{N}_\mu(\mathcal{G})$,

$$\lambda = \sum_{i=1}^m \frac{\|\phi_i\|_{L^2(h_i)}^2}{\|\tilde{u}^+\|_{L^2(\tilde{\mathcal{G}})}^2} \theta_i + \frac{\|u^+\|_{L^2(\mathcal{G})}^2}{\|\tilde{u}^+\|_{L^2(\tilde{\mathcal{G}})}^2} \mu,$$

which, combined with the preceding inequality, yields $\mu > \lambda$.

Since, analogously to Remark 2.5, for a given λ , the map

$$\mu \mapsto \inf_{v \in \mathcal{M}_{\mu,\lambda}(\mathcal{G})} \frac{1}{2} \|v'\|_{L^2(\mathcal{G})}^2 + \frac{1}{2} \mu \|v^+\|_{L^2(\mathcal{G})}^2 + \frac{1}{2} \lambda \|v^-\|_{L^2(\mathcal{G})}^2 - \frac{1}{p} \|v\|_{L^p(\mathcal{G})}^p,$$

where $\mathcal{M}_{\mu,\lambda}(\mathcal{G}) := \{v \in H^1(\mathcal{G}) \mid v^+ \in \mathcal{N}_\mu(\mathcal{G}) \text{ and } v^- \in \mathcal{N}_\lambda(\mathcal{G})\}$, is increasing, we have

$$\inf_{v \in \mathcal{M}_\lambda(\tilde{\mathcal{G}})} J(v) = J(\tilde{u}) = \kappa (\|\tilde{u}^+\|_p^p + \|\tilde{u}^-\|_p^p) \geq \kappa (\|u^+\|_p^p + \|u^-\|_p^p) \geq \inf_{v \in \mathcal{M}_{\mu,\lambda}(\mathcal{G})} J(v) > \inf_{v \in \mathcal{M}_\lambda(\mathcal{G})} J(v). \quad \square$$

Proof of Proposition 6.5. Consider the graph $\mathcal{G}_{1,L}$ with $L \geq \bar{L}$ given by Proposition 6.2. On this graph, by Theorem 4.12 and Proposition 6.2, we have a nodal ground state u with $u^{-1}(0) = \{x_0\}$. Let now $\bar{\mathcal{G}}$ be the graph obtained from $\mathcal{G}_{1,L}$ by attaching m half-lines at the point x_0 and let $\bar{u} \in \mathcal{M}_\lambda(\bar{\mathcal{G}})$ be the function obtained extending u by 0 on each of the additional half-lines.

By Theorem 4.12, nodal ground states exist on $\bar{\mathcal{G}}$ and, by Lemma 6.6,

$$\inf_{v \in \mathcal{M}_\lambda(\mathcal{G}_{1,L})} J(v) = J(u) = J(\bar{u}) \geq \inf_{v \in \mathcal{M}_\lambda(\bar{\mathcal{G}})} J(v) \geq \inf_{v \in \mathcal{M}_\lambda(\mathcal{G}_{1,L})} J(v).$$

This proves that \bar{u} is a nodal ground state on $\bar{\mathcal{G}}$ and hence the existence of a nodal ground state whose nodal set is given by m half-lines attached at the same point. \square

Proof of (3). Here we prove the following statement.

Proposition 6.7. *Let $n \in \mathbb{N}$. There exist a graph $\bar{\mathcal{G}}$, a subset $Z \subseteq \mathbb{V} \setminus \mathbb{V}_\infty$ of its vertices of degree 1 and a nodal ground state $u \in \mathcal{M}_{\lambda,Z}(\bar{\mathcal{G}})$ such that $u^{-1}(0)$ consists of n line segments attached at the same point, each of length smaller than or equal to $\frac{\kappa}{2s_\lambda} \left(\frac{p\lambda}{2}\right)^{2/(p-2)}$.*

Similarly to construction (2), the graph $\bar{\mathcal{G}}$ will be obtained from $\mathcal{G}_{1,L}$ by attaching n line segments at one of its points. To do this we need the next lemma.

Lemma 6.8. *Let \mathcal{G} be a noncompact graph with a finite number of edges. Let $\tilde{\mathcal{G}}$ be a graph obtained from \mathcal{G} by attaching n line segments s_1, \dots, s_n at one of its points \mathfrak{p} . Assume that each line segment has a length smaller than or equal to a number $S > 0$ and ends at a vertex with Dirichlet boundary condition. Suppose also that \tilde{u} and S are such that \tilde{u} is a nodal ground state on $\tilde{\mathcal{G}}$ and that $S \leq \frac{\kappa}{J(\tilde{u})} \left(\frac{p\lambda}{2}\right)^{2/(p-2)}$. Then*

$$\inf_{v \in \mathcal{M}_{\lambda,Z}(\tilde{\mathcal{G}})} J(v) \geq \inf_{v \in \mathcal{M}_{\lambda,Z}(\mathcal{G})} J(v).$$

Proof. We proceed in the same way as in the proof of Lemma 6.6. With no loss of generality, let $\tilde{u}(\mathfrak{p}) \geq 0$. Denote by u the restriction of \tilde{u} to \mathcal{G} and by u_i the restriction of u to s_i for every i . Moreover, let $u'_i(\mathfrak{p})$ be the outward derivative of u_i at \mathfrak{p} along s_i . As \tilde{u} is a nodal ground state, $u_i(\mathfrak{p})u'_i(\mathfrak{p}) \leq 0$. Indeed, if this were not the case, we would have $u'_i(\mathfrak{p}) > 0$ and, since u_i satisfies the Dirichlet condition at the end of s_i ,




Figure 11. Example of a graph as in Theorem 1.9 hosting a nodal ground state whose nodal set (thick on the picture) is made of two isolated points, two groups of three and four line segments respectively, and a group of five half-lines.

by a phase plane analysis we would have $u_i(x_0) := \max u_i \geq \max \phi_\lambda = \left(\frac{1}{2}p\lambda\right)^{1/(p-2)}$. Considering the first zero $x_1 \in s_i$ of u_i , it would then follow that

$$\left(\frac{p\lambda}{2}\right)^{1/(p-2)} \leq u_i(x_0) - u_i(x_1) = \int_{x_1}^{x_0} u_i'(s) ds \leq \sqrt{x_0 - x_1} \|\tilde{u}'\|_2 < \sqrt{\frac{SJ(\tilde{u})}{\kappa}},$$

which contradicts the choice of S . The rest of the proof follows as in that of Lemma 6.6. \square

Proof of Proposition 6.7. The proof is the same as the one of Proposition 6.5, using Lemma 6.8 instead of Lemma 6.6 and observing that, by Theorem 4.12, $J(\tilde{u}) \leq 2s_\lambda$. \square

Remark 6.9. Graphs fulfilling Theorem 1.9 can be obtained combining ad libitum the constructions (1), (2), (3). The general result is a graph as the one depicted in Figure 11.

Acknowledgements

This work was carried out when Galant was a F. R. S.-FNRS Research Fellow. This work has been partially supported by IEA Project “Nonlinear Schrödinger Equations on Metric Graphs”, by GNAMPA Project 2023 “Modelli nonlineari in presenza di interazioni puntuali”, and by the project E53D23005450006 “Nonlinear dispersive equations in presence of singularities” — funded by the European Union — Next Generation EU within the PRIN 2022 program (D. D. 104 -02/02/2022 Ministero dell’Università e della Ricerca). This article reflects only the authors’ views and opinions and the Ministry cannot be considered responsible for them. This work has been carried out in the framework of the Project NQG (ANR-23-CE40-0005-01), funded by the French National Research Agency (ANR). De Coster and Galant thank the ANR for its support.

References

- [Adami et al. 2014a] R. Adami, C. Cacciapuoti, D. Finco, and D. Noja, “Constrained energy minimization and orbital stability for the NLS equation on a star graph”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **31**:6 (2014), 1289–1310. MR
- [Adami et al. 2014b] R. Adami, C. Cacciapuoti, D. Finco, and D. Noja, “Variational properties and orbital stability of standing waves for NLS equation on a star graph”, *J. Differential Equations* **257**:10 (2014), 3738–3777. MR

- [Adami et al. 2015] R. Adami, E. Serra, and P. Tilli, “NLS ground states on graphs”, *Calc. Var. Partial Differential Equations* **54**:1 (2015), 743–761. MR
- [Adami et al. 2016] R. Adami, E. Serra, and P. Tilli, “Threshold phenomena and existence results for NLS ground states on metric graphs”, *J. Funct. Anal.* **271**:1 (2016), 201–223. MR
- [Adami et al. 2019] R. Adami, S. Dovetta, E. Serra, and P. Tilli, “Dimensional crossover with a continuum of critical exponents for NLS on doubly periodic metric graphs”, *Anal. PDE* **12**:6 (2019), 1597–1612. MR
- [Adami et al. 2020] R. Adami, F. Boni, and A. Ruighi, “Non-Kirchhoff vertices and nonlinear Schrödinger ground states on graphs”, *Mathematics* **8**:4 (2020), art. id. 617.
- [Adami et al. 2022] R. Adami, F. Boni, and S. Dovetta, “Competing nonlinearities in NLS equations as source of threshold phenomena on star graphs”, *J. Funct. Anal.* **283**:1 (2022), art. id. 109483. MR
- [Agostinho et al. 2024] F. Agostinho, S. Correia, and H. Tavares, “Classification and stability of positive solutions to the NLS equation on the \mathcal{T} -metric graph”, *Nonlinearity* **37**:2 (2024), art. id. 025005. MR
- [Bartsch and Weth 2003] T. Bartsch and T. Weth, “A note on additional properties of sign changing solutions to superlinear elliptic equations”, *Topol. Methods Nonlinear Anal.* **22**:1 (2003), 1–14. MR
- [Bartsch et al. 2005] T. Bartsch, T. Weth, and M. Willem, “Partial symmetry of least energy nodal solutions to some variational problems”, *J. Anal. Math.* **96** (2005), 1–18. MR
- [Berkolaiko and Kuchment 2013] G. Berkolaiko and P. Kuchment, *Introduction to quantum graphs*, Mathematical Surveys and Monographs **186**, Amer. Math. Soc., Providence, RI, 2013. MR
- [Berkolaiko et al. 2021] G. Berkolaiko, J. L. Marzuola, and D. E. Pelinovsky, “Edge-localized states on quantum graphs in the limit of large mass”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **38**:5 (2021), 1295–1335. MR
- [Besse et al. 2022a] C. Besse, R. Duboscq, and S. Le Coz, “Gradient flow approach to the calculation of stationary states on nonlinear quantum graphs”, *Ann. H. Lebesgue* **5** (2022), 387–428. MR
- [Besse et al. 2022b] C. Besse, R. Duboscq, and S. Le Coz, “Numerical simulations on nonlinear quantum graphs with the GraFiDi library”, *SMAI J. Comput. Math.* **8** (2022), 1–47. MR
- [Boni and Carlone 2023] F. Boni and R. Carlone, “NLS ground states on the half-line with point interactions”, *NoDEA Nonlinear Differential Equations Appl.* **30**:4 (2023), art. id. 51. MR
- [Boni and Dovetta 2021] F. Boni and S. Dovetta, “Prescribed mass ground states for a doubly nonlinear Schrödinger equation in dimension one”, *J. Math. Anal. Appl.* **496**:1 (2021), art. id. 124797. MR
- [Boni and Dovetta 2022] F. Boni and S. Dovetta, “Doubly nonlinear Schrödinger ground states on metric graphs”, *Nonlinearity* **35**:7 (2022), 3283–3323. MR
- [Borthwick et al. 2023] J. Borthwick, X. Chang, L. Jeanjean, and N. Soave, “Normalized solutions of L^2 -supercritical NLS equations on noncompact metric graphs with localized nonlinearities”, *Nonlinearity* **36**:7 (2023), 3776–3795. MR
- [Brézis and Lieb 1983] H. Brézis and E. Lieb, “A relation between pointwise convergence of functions and convergence of functionals”, *Proc. Amer. Math. Soc.* **88**:3 (1983), 486–490. MR
- [Castro et al. 1997] A. Castro, J. Cossio, and J. M. Neuberger, “A sign-changing solution for a superlinear Dirichlet problem”, *Rocky Mountain J. Math.* **27**:4 (1997), 1041–1053. MR
- [Cazenave 2003] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics **10**, Amer. Math. Soc., Providence, RI, 2003. MR
- [Chang et al. 2024] X. Chang, L. Jeanjean, and N. Soave, “Normalized solutions of L^2 -supercritical NLS equations on compact metric graphs”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **41**:4 (2024), 933–959. MR
- [De Coster et al. 2023] C. De Coster, S. Dovetta, D. Galant, and E. Serra, “On the notion of ground state for nonlinear Schrödinger equations on metric graphs”, *Calc. Var. Partial Differential Equations* **62**:5 (2023), art. id. 159. MR
- [Dovetta 2018] S. Dovetta, “Existence of infinitely many stationary solutions of the L^2 -subcritical and critical NLSE on compact metric graphs”, *J. Differential Equations* **264**:7 (2018), 4806–4821. MR
- [Dovetta 2019] S. Dovetta, “Mass-constrained ground states of the stationary NLSE on periodic metric graphs”, *NoDEA Nonlinear Differential Equations Appl.* **26**:5 (2019), art. id. 30. MR
- [Dovetta 2024] S. Dovetta, “Singular limit of periodic metric grids”, *Adv. Math.* **444** (2024), art. id. 109633. MR
- [Dovetta and Tentarelli 2022] S. Dovetta and L. Tentarelli, “Symmetry breaking in two-dimensional square grids: persistence and failure of the dimensional crossover”, *J. Math. Pures Appl.* (9) **160** (2022), 99–157. MR

- [Dovetta et al. 2020] S. Dovetta, E. Serra, and P. Tilli, “NLS ground states on metric trees: existence results and open questions”, *J. Lond. Math. Soc.* (2) **102**:3 (2020), 1223–1240. MR
- [Dovetta et al. 2023] S. Dovetta, E. Serra, and P. Tilli, “Action versus energy ground states in nonlinear Schrödinger equations”, *Math. Ann.* **385**:3-4 (2023), 1545–1576. MR
- [Esteban 2022] M. J. Esteban, “Gagliardo–Nirenberg–Sobolev inequalities on planar graphs”, *Commun. Pure Appl. Anal.* **21**:6 (2022), 2101–2114. MR
- [Gilg et al. 2022] S. Gilg, G. Schneider, and H. Uecker, “Nonlinear dynamics of modulated waves on graphene like quantum graphs”, *Math. Nachr.* **295**:11 (2022), 2147–2170. MR
- [Jeanjean and Lu 2022] L. Jeanjean and S.-S. Lu, “On global minimizers for a mass constrained problem”, *Calc. Var. Partial Differential Equations* **61**:6 (2022), art. id. 214. MR
- [Kairzhan et al. 2021] A. Kairzhan, R. Marangell, D. E. Pelinovsky, and K. L. Xiao, “Standing waves on a flower graph”, *J. Differential Equations* **271** (2021), 719–763. MR
- [Kairzhan et al. 2022] A. Kairzhan, D. Noja, and D. E. Pelinovsky, “Standing waves on quantum graphs”, *J. Phys. A* **55**:24 (2022), art. id. 243001. MR
- [Kurata and Shibata 2020] K. Kurata and M. Shibata, “Least energy solutions to semi-linear elliptic problems on metric graphs”, *J. Math. Anal. Appl.* **491**:1 (2020), art. id. 124297. MR
- [Le Coz 2009] S. Le Coz, “Standing waves in nonlinear Schrödinger equations”, pp. 151–192 in *Analytical and numerical aspects of partial differential equations*, edited by E. Emmrich and P. Wittbold, Walter de Gruyter, Berlin, 2009. MR
- [Pankov 2018] A. Pankov, “Nonlinear Schrödinger equations on periodic metric graphs”, *Discrete Contin. Dyn. Syst.* **38**:2 (2018), 697–714. MR
- [Pelinovsky and Schneider 2017] D. Pelinovsky and G. Schneider, “Bifurcations of standing localized waves on periodic graphs”, *Ann. Henri Poincaré* **18**:4 (2017), 1185–1211. MR
- [Pierotti and Soave 2022] D. Pierotti and N. Soave, “Ground states for the NLS equation with combined nonlinearities on noncompact metric graphs”, *SIAM J. Math. Anal.* **54**:1 (2022), 768–790. MR
- [Pierotti et al. 2021] D. Pierotti, N. Soave, and G. Verzini, “Local minimizers in absence of ground states for the critical NLS energy on metric graphs”, *Proc. Roy. Soc. Edinburgh Sect. A* **151**:2 (2021), 705–733. MR
- [Szulkin and Weth 2010] A. Szulkin and T. Weth, “The method of Nehari manifold”, pp. 597–632 in *Handbook of nonconvex analysis and applications*, edited by D. Y. Gao and D. Motreanu, International Press, Somerville, MA, 2010. MR
- [Tentarelli 2016] L. Tentarelli, “NLS ground states on metric graphs with localized nonlinearities”, *J. Math. Anal. Appl.* **433**:1 (2016), 291–304. MR

Received 25 Sep 2023. Revised 25 Sep 2024. Accepted 1 Feb 2025.

COLETTE DE COSTER: colette.decoster@uphf.fr

Université Polytechnique Hauts-de-France, INSA Hauts-de-France, CERAMATHS - Laboratoire de Matériaux Céramiques et de Mathématiques, F-59313 Valenciennes, France

SIMONE DOVETTA: simone.dovetta@polito.it

Dipartimento di Scienze Matematiche “G. L. Lagrange”, Politecnico di Torino, Torino, Italy

DAMIEN GALANT: damien_galant@brown.edu

Département de Mathématiques, Université de Mons, Mons, Belgium

and

Université Polytechnique Hauts-de-France, INSA Hauts-de-France, CERAMATHS - Laboratoire de Matériaux Céramiques et de Mathématiques, F-59313 Valenciennes, France

Current address: Department of Mathematics, Brown University, Providence, RI, United States

ENRICO SERRA: enrico.serra@polito.it

Dipartimento di Scienze Matematiche “G. L. Lagrange”, Politecnico di Torino, Torino, Italy

CHRISTOPHE TROESTLER: christophe.troestler@umons.ac.be

Département de Mathématiques, Université de Mons, Mons, Belgium

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Anna L. Mazzucato Penn State University, USA
alm24@psu.edu

Clément Mouhot Cambridge University, UK
c.mouhot@dpms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Sc. Intern. Sup. di Studi Avanz., Italy berti@sissa.it	Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Yu Deng	University of Chicago, USA yudeng@uchicago.edu	Omar Mohsen	Université Paris-Cité, France omar.mohsen.fr@gmail.com
Thierry Gallay	Université Grenoble Alpes, France Thierry.Gallay@univ-grenoble-alpes.fr	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Xavier Ros Oton	Catalan Inst. for Res. and Adv. Std., Spain xros@icrea.cat
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Nicolas Rougerie	ENS Lyon, France nicolas.rougerie@ens-lyon.fr
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Sebastian Herr	Universität Bielefeld, Germany herr@math.uni-bielefeld.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Jonathan Wing-hong Luk	Stanford University jluk@stanford.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

Cover image: Eric J. Heller: “Linear Ramp”


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2026 is US \$500/year for the electronic version, and \$780/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 19 No. 2 2026

Constant sign and sign changing NLS ground states on noncompact metric graphs	203
COLETTE DE COSTER, SIMONE DOVETTA, DAMIEN GALANT, ENRICO SERRA and CHRISTOPHE TROESTLER	
Controllability of parabolic equations with inverse square infinite potential wells via global Carleman estimates	241
ALBERTO ENCISO, ARICK SHAO and BRUNO VERGARA	
Focusing dynamics of 2D Bose gases in the instability regime	281
LEA BOSSMANN, CHARLOTTE DIETZE and PHAN THÀNH NAM	
Lower bounds on fibered Yang–Mills functionals: generic nefness and semistability of direct images	317
SIARHEI FINSKI	
Hessian estimates for special Lagrangian equation by doubling	339
RAVI SHANKAR	
Regularity for nonlocal equations with local Neumann boundary conditions	353
XAVIER ROS-OTON and MARVIN WEIDNER	