


Geometry & Topology
Volume 9 (2005) 1915–1951
Published: 17 October 2005



Representations of polygons of finite groups

MICHAEL KAPOVICH

*Department of Mathematics, University of California
1 Shields Ave, Davis, CA 95616, USA*

Email: kapovich@math.ucdavis.edu

Abstract

We construct discrete and faithful representations into the isometry group of a hyperbolic space of the fundamental groups of acute negatively curved even-sided polygons of finite groups.

AMS Classification numbers Primary: 20F69

Secondary: 20G05

Keywords: Hyperbolic groups, polygons of groups

Proposed: Benson Farb
Seconded: Martin Bridson, Walter Neumann

Received: 30 January 2004
Revised: 6 October 2005

1 Introduction

There are only few known obstructions for existence of an isometric properly discontinuous action of a Gromov-hyperbolic group G on the *real-hyperbolic space* \mathbb{H}^p for some p :

(1) If G is a group satisfying Kazhdan property (T) then each isometric action $G \curvearrowright \mathbb{H}^p$ fixes a point in \mathbb{H}^p ; hence no infinite hyperbolic group satisfying property (T) admits an isometric properly discontinuous action $G \curvearrowright \mathbb{H}^p$, for any p .

(2) Suppose that G is the fundamental group of a compact Kähler manifold and $G \curvearrowright \mathbb{H}^p$ is an isometric properly discontinuous action. Then, according to a theorem of Carlson and Toledo [8], this action factors through an epimorphism $G \rightarrow Q$, where Q is commensurable to a surface group. Hence, unless G itself is commensurable to a surface group, it does not admit an isometric properly discontinuous action $G \curvearrowright \mathbb{H}^p$. Examples of Gromov-hyperbolic groups which are Kähler (and are not commensurable to surface groups) are given by the uniform lattices in $PU(m, 1)$, $m \geq 2$, as well as the fundamental groups of compact negatively curved Kähler manifolds (see [20]).

On the positive side, by a theorem of Bonk and Schramm [3], each Gromov-hyperbolic group admits a quasi-isometric embedding to a real-hyperbolic space.

The goal of this paper is to find a better “demarcation line” between hyperbolic groups satisfying property (T) and groups acting discretely on real-hyperbolic spaces. In this paper we will show that a large class of 2-dimensional Gromov-hyperbolic groups admits isometric properly discontinuous *convex-cocompact* actions on real-hyperbolic spaces. We consider a 2-dimensional negatively curved acute polygon \mathcal{P} of finite groups (see section 2.2 for more details). Let $G := \pi_1(\mathcal{P})$ be the fundamental group of this polygon, we refer the reader to [7, Chapter II, section 12] for the precise definitions.

Our main result is:

Theorem 1.1 *Suppose that $n = 2k$ is even. Then the group G admits a discrete, faithful, convex-cocompact action ρ on a constant curvature hyperbolic space \mathbb{H}^p , where $p < \infty$ depends on the polygon \mathcal{P} .*

Our technique in general does not work in the case when n is odd: We were unable to construct a representation. However in section 6 we will construct ρ and prove that it is discrete, faithful, convex-cocompact for a special class of odd-sided n -gons of groups, provided that $n \geq 5$.

Conjecture 1.2 *The assertion of Theorem 1.1 remains valid for all odd $n \geq 5$.*

In contrast, if \mathcal{P} is a *triangle* of finite groups where the vertex links are connected graphs with the 1st positive eigenvalue of the Laplacian $> 1/2$, then the group $G = \pi_1(\mathcal{P})$ satisfies property (T), see [1]. Hence (provided that G is infinite) the group G cannot act properly discontinuously on \mathbb{H}^p for any p . Thus, it appears, that (at least for the polygons of finite groups) the “demarcation line” which we are trying to find, is hidden somewhere among quadrilaterals and triangles of groups. We will address this issue in another paper.

Recall that, by a theorem of Dani Wise [24], G is residually finite (actually, Wise proves that G has separable quasi-convex subgroups, which is used in the proof of our main theorem).

Corollary 1.3 *The group G is linear.*

Remark 1.4 A very different proof of linearity of G was given by Wise and Haglund, who used an embedding of G to a right-angled Coxeter group, [14].

The following problem is open even for right-angled Coxeter groups of virtual cohomological dimension 2.

Problem 1.5 Suppose that G is a Gromov-hyperbolic Coxeter group. Is G isomorphic to a discrete subgroup of $\text{Isom}(\mathbb{H}^n)$ for some n ? Note that if one insists that the Coxeter generators act on \mathbb{H}^n as reflections, then there are examples of Gromov-hyperbolic Coxeter groups which do not admit such actions on \mathbb{H}^n , see [10].

In section 8, we give an example of a nonlinear Gromov-hyperbolic group.

The proof of the main theorem splits in two parts: (1) Construction of ρ , (2) proof of discreteness. To prove discreteness of ρ we show that there exists a ρ -equivariant quasi-isometric embedding $\mu: X \rightarrow \mathbb{H}^p$, where X is the universal cover of the polygon \mathcal{P} . This proves that the action $\rho: G \curvearrowright \mathbb{H}^p$ is properly discontinuous and convex-cocompact. (A priori this action can have finite kernel. In section 7 we explain how to deal with this issue.) The proof that μ is a quasi-isometric embedding is based on the following theorem of independent interest:

Theorem 1.6 *Suppose that X is a 2-dimensional regular cell complex, which is equipped with a $\text{CAT}(-1)$ path-metric so that each face of X is isometric to a right-angled regular n -gon in \mathbb{H}^2 . Let $\mu: X \rightarrow \mathbb{H}^p$ is a continuous map which is a (totally-geodesic) isometric embedding on each face of X . Assume also that for each pair of faces $F', F'' \subset X$ which intersect non-trivially a common face $F \subset X$, we have:*

$$\text{Span}(\mu(F')) \perp \text{Span}(\mu(F'')).$$

Then μ is a quasi-isometric embedding.

Our construction of representations ρ was inspired by the paper of Marc Bourdon, [4], where he proves a theorem which is a special case of Theorem 1.1: In his paper Bourdon considers n -gons of finite groups where the edge groups are cyclic and the vertex groups are direct products of the adjacent edge groups, under the extra assumption that the orders of the edge groups are much smaller than n .

Acknowledgments During the work on this paper I was visiting the Max Plank Institute (Bonn), I was also supported by the NSF grants DMS-02-03045 and DMS-04-05180. I am grateful to Tadeusz Januszkiewicz for inspirational and helpful conversations during the work on this paper. I first tried to prove discreteness of certain representations by verifying that an equivariant map is a quasi-isometric embedding, in a joint project with Bernhard Leeb in 1998. Although our attempt back then was unsuccessful, I am grateful to Bernhard Leeb for that effort. I am also grateful to the referees of this paper for numerous suggestions and to Mark Sapir for discussions of Theorem 8.1.

2 Preliminaries

Notation If Σ is a (finite) set, we define a Euclidean vector space $\text{Vect}(S)$ to be the vector space $L_2(S)$, where S forms an orthonormal basis (we identify each 1-point subset of S with its characteristic function in $L_2(S)$). Suppose that $S \subset \mathbb{H}^q$. Then $\text{Span}(E)$ will denote the smallest totally-geodesic subspace in \mathbb{H}^q which contains S . If $E \subset \mathbb{H}^q$ is a geodesic segment, then $\text{Bis}(E)$ will denote the perpendicular bisector of E .

Suppose that $E_1, E_2 \subset E$ are subspaces of a Euclidean vector space E , whose intersection is E_3 . We say that E_1, E_2 intersect orthogonally if $E_1/E_3, E_2/E_3$ are contained in the orthogonal complements of each other in the Euclidean space E/E_3 .

Suppose that H', H'' are totally-geodesic subspaces in \mathbb{H}^p . We say that H', H'' intersect orthogonally if

$$H' \setminus H'' \neq \emptyset, H'' \setminus H' \neq \emptyset$$

and for some (equivalently, for every) point $x \in H' \cap H''$ we have:

$$T_x(H'), T_x(H'') \subset T_x(\mathbb{H}^p) \text{ intersect orthogonally.}$$

Totally geodesic subspaces $H', H'' \subset \mathbb{H}^p$ are said to be *orthogonal to each other* if either:

- (a) H', H'' intersect orthogonally, or
- (b) H', H'' are within positive distance from each other and for the unique shortest geodesic segment $\sigma := \overline{x'x''} \subset \mathbb{H}^p$ connecting H' to H'' , the totally-geodesic subspaces $H'', \gamma_\sigma(H')$ intersect orthogonally. Here γ_σ is the hyperbolic translation along σ which sends x' to x'' .

We will use the notation $H' \perp H''$ for subspaces H', H'' orthogonal to each other. Clearly, $H' \perp H'' \iff H'' \perp H'$.

2.1 Discrete subgroups of $\text{Isom}(\mathbb{H}^n)$

Recall that a map $f: X \rightarrow Y$ between two metric spaces is called an (L, A) *quasi-isometric embedding* if for all $x, x' \in X$ we have:

$$L^{-1}d(x, x') - A \leq d(f(x), f(x')) \leq Ld(x, x') + A,$$

where $L > 0$. An (L, A) *quasi-isometry* is an (L, A) quasi-isometric embedding $X \xrightarrow{f} Y$ such that each point of Y is within distance $\leq A$ from a point in $\text{Im}(f)$.

A map f is called a *quasi-isometry* (resp. a *quasi-isometric embedding*) if it is an (L, A) quasi-isometry (resp. quasi-isometric embedding) for some L and A .

An (L, A) quasi-geodesic segment in a metric space X is an (L, A) quasi-isometric embedding $f: [0, T] \rightarrow X$, where $[0, T]$ is an interval in \mathbb{R} . By abusing notation we will sometimes refer to the image $\text{Im}(f)$ of an (L, A) quasi-geodesic segment f as an (L, A) quasi-geodesic segment. Recall that by the *Morse lemma* (see for instance [17, Lemma 3.43]), quasi-geodesics in \mathbb{H}^n are *stable*:

There is a function $D = D(L, A)$ such that for each (L, A) quasi-geodesic segment $f: [0, T] \rightarrow \mathbb{H}^n$, the Hausdorff distance between $\text{Im}(f)$ and the geodesic segment

$$\overline{f(0)f(T)} \subset \mathbb{H}^n$$

connecting the end-points of $\text{Im}(f)$ is at most D .

We will use the notation \mathbb{H}^n for the real-hyperbolic n -space; its curvature is normalized to be equal to -1 . The space \mathbb{H}^n has a geometric compactification $\bar{\mathbb{H}}^n = \mathbb{H}^n \cup S^{n-1}$. For a subset $S \subset \mathbb{H}^n$ we let \bar{S} denote its closure in $\bar{\mathbb{H}}^n$.

Discrete subgroups $G \subset \text{Isom}(\mathbb{H}^n)$ are called *Kleinian groups*.

The *convex hull* $C(G)$ of a Kleinian group $G \subset \text{Isom}(\mathbb{H}^n)$ is the smallest non-empty closed convex G -invariant subset $C \subset \mathbb{H}^n$. The convex hull exists for each G whose limit set has cardinality $\neq 1$. The convex hull is unique unless G is finite.

Definition 2.1 A Kleinian group $G \subset \text{Isom}(\mathbb{H}^n)$ is called *convex-cocompact* if $C(G)$ exists and the quotient $C(G)/G$ is compact.

Lemma 2.2 *Suppose we have a representation $\rho: G \rightarrow \text{Isom}(\mathbb{H}^n)$ of a finitely-generated group G . Then the action $\rho: G \curvearrowright \mathbb{H}^n$ is properly discontinuous and convex-cocompact iff there exists a G -equivariant quasi-isometric embedding $f: \Gamma_G \rightarrow \mathbb{H}^n$, where Γ_G is a Cayley graph of G .*

Proof First, suppose that $\rho: G \curvearrowright \mathbb{H}^n$ properly discontinuous and convex-cocompact. Then, because $C(G)$ is a geodesic metric space, there exists a G -equivariant quasi-isometry $f: \Gamma_G \rightarrow C(G)$. Composing this map with the isometric embedding $\iota: C(G) \rightarrow \mathbb{H}^n$, we conclude that $f: \Gamma_G \rightarrow \mathbb{H}^n$ is a quasi-isometric embedding.

Conversely, suppose that $f: \Gamma_G \rightarrow \mathbb{H}^n$ is an equivariant quasi-isometric embedding. In particular, f is a proper map. Hence, if for $1 \in \Gamma_G$ we set $o := f(1)$, then for each compact subset $K \subset \mathbb{H}^n$ there are only finitely many elements $g \in G$ such that $g(o) \in K$. Therefore the action $G \curvearrowright \mathbb{H}^n$ is properly discontinuous. In particular, it has finite kernel.

Observe that *stability* of quasi-geodesics in \mathbb{H}^n implies that $\text{Im}(f)$ is *quasi-convex*, ie, there exists a constant $c < \infty$ such that for any two points $x, y \in \text{Im}(f)$ the geodesic segment \overline{xy} is contained in a c -neighborhood $N_c(\text{Im}(f))$ of $\text{Im}(f)$. On the other hand, by [6, Proposition 2.5.4], there exists $R = R(c)$ such that the convex hull of each c -quasi-convex subset $S \subset \mathbb{H}^n$ is contained in the R -neighborhood $N_R(S)$. Thus, the convex subset $C(\rho(G)) \subset \mathbb{H}^n$ is contained in $N_{R(c)}(\text{Im}(f))$. Since G acts cocompactly on $\text{Im}(f)$ it follows that G acts cocompactly on $C(G)$. Therefore $\rho(G)$ is convex-cocompact. \square

Remark 2.3 (1) Clearly, instead of Γ_G in the above lemma one can use any geodesic metric space on which G acts isometrically, properly discontinuously and cocompactly.

(2) The above lemma shows that existence of a G -equivariant quasi-isometric embedding $f: \Gamma_G \rightarrow \mathbb{H}^n$ implies that ρ has finite kernel. However it does not exclude the possibility that this kernel is nontrivial.

Definition 2.4 Let G be a group with a Cayley graph Γ_G . A subgroup $H \subset G$ is called *quasi-convex* if the orbit $H \cdot 1 \subset \Gamma_G$ is quasi-convex, ie, there exists a number D so that each geodesic segment $\sigma \subset \Gamma_G$ with vertices in $H \cdot 1$ is contained in $N_D(H \cdot 1)$.

If G is Gromov-hyperbolic then quasi-convexity of H is independent of the choice of Cayley graph Γ_G .

2.2 Geometry of polygons of groups

Consider an n -gon P ($n \geq 5$) with vertices x_i and edges e_j , $1 \leq i, j \leq n$. Throughout we will be working mod n , ie, $qn + i$ will be identified with i for $i \in \{1, \dots, n\}$. We will be assuming that each edge e_i has the vertices x_i, x_{i+1} . We will regard P as a (2-dimensional) cell complex and its poset $Pos(P)$ as a (small) category. A *polygon of groups* \mathcal{P} based on P is a covariant functor from $Pos(P)$ to the category of groups and monomorphisms.

In other words, a *polygon of groups* \mathcal{P} based on P is a collection of groups G_{x_i}, G_{e_i}, G_F assigned to the vertices, edges and the 2-face F of P , together with monomorphisms

$$G_F \rightarrow G_e \rightarrow G_x$$

for each edge e containing the vertex x , so that the following diagrams are commutative:

$$\begin{array}{ccc} G_x & \leftarrow & G_e \\ & \swarrow & \uparrow \\ & & G_F \end{array}$$

The direct limit of the above diagrams of monomorphisms is the *fundamental group* $G = \pi(\mathcal{P})$ of the polygon \mathcal{P} . If the vertex, edge and face groups of \mathcal{P} embed naturally into G , the polygon \mathcal{P} is called *developable*. Not every polygon of groups is developable, however under a certain *nonpositive curvature* assumption on \mathcal{P} , the polygon \mathcal{P} is developable, see [7].

Curvature and angles For each vertex $x_i \in P$ define a graph Lk_{x_i} as follows. The vertices of Lk_{x_i} are the right cosets $gG_{e_i}, gG_{e_{i-1}}, g \in G_{x_i}$. The vertices v, w are connected by a (single) edge iff there exists $g \in G_{x_i}$ such that $g(\{v, w\}) = \{G_{e_i}, G_{e_{i-1}}\}$. Thus the group G_{x_i} acts on Lk_{x_i} with the quotient being the edge connecting $G_{e_i}, G_{e_{i-1}}$. We metrize the graph Lk_{x_i} by assigning the same length α_i to each edge, so that the group G_{x_i} acts isometrically. Then the *angle* between the subgroups $G_{e_i}, G_{e_{i-1}}$ is the least number α_i such that the metric graph Lk_{x_i} is a CAT(1) space, ie, the length of the shortest embedded cycle in Lk_{x_i} is at least 2π . Equivalently, the angle between $G_{e_i}, G_{e_{i-1}}$ equals

$$2\pi/\text{girth}(Lk_{x_i}).$$

We will say that the polygon \mathcal{P} is *acute* (or has *acute angles*) if the angle α_i between each pair of edge groups $G_{e_i}, G_{e_{i-1}}$ is at most $\pi/2$.

We refer the reader to [7, Chapter II, section 12] for the precise definitions of the nonpositive/negative curvature of \mathcal{P} ; various examples of negatively curved polygons of groups can be found in [1], [7, Chapter II, section 12] and [22]. Instead, we state the following equivalent definition of negative curvature:

There exists a 2-dimensional simply-connected regular cell complex X (the universal cover of \mathcal{P}) together with a path-metric on X whose restriction to each face of X has constant curvature -1 , so that:

- (1) Each face of X is isometric to an n -gon in \mathbb{H}^2 with angles $\alpha_1, \dots, \alpha_n$.
- (2) Each cell in X is convex.
- (3) There exists an isometric cellular action $G \curvearrowright X$ which is transitive on 2-cells.
- (4) The stabilizer of each 2-face $F \subset X$ is isomorphic to G_F , it fixes F pointwise.
- (5) The stabilizer of each edge e of F is isomorphic to G_e and it fixes e pointwise.
- (6) The stabilizer of each vertex x of F is isomorphic to G_x .
- (7) The inclusion maps $G_F \hookrightarrow G_e \hookrightarrow G_x$ coincide with the monomorphisms $G_F \rightarrow G_e \rightarrow G_x$ in the definition of \mathcal{P} .

Note that the link in X of each vertex $x_i \in F$ is isometric to Lk_{x_i} (where each edge has the length α_i). Thus the above complex X is a CAT(-1) metric space.

Throughout the paper we will be using only the following corollary of negative curvature for *acute* polygons of groups:

Corollary 2.5 *If \mathcal{P} is negatively curved then there is a CAT(-1) complex X where each face is isometric to a regular right-angled polygon in \mathbb{H}^2 , so that the properties (2)–(7) are satisfied.*

In this paper we will consider only the case when the vertex groups are finite, thus the action $G \curvearrowright X$ is properly discontinuous and cocompact, which implies that X is equivariantly quasi-isometric to a Cayley graph of G .

We now return to the original polygon \mathcal{P} assuming that it has even number of sides. Let o denote the center of the face F and let m_j be the midpoint of the edge $e_j \subset F$. We consider two subgraphs $\Gamma_{even}, \Gamma_{odd} \subset F$ which are obtained by conning off from o the sets

$$m_{even} := \{m_{2j}, j = 1, \dots, n/2\}, m_{odd} := \{m_{2j-1}, j = 1, \dots, n/2\}$$

respectively. Let G_{even}, G_{odd} denote the subgroups of G generated by the elements of

$$G_{e_{2j}}, j = 1, \dots, n/2$$

and

$$G_{e_{2j-1}}, j = 1, \dots, n/2$$

respectively. Define subgraphs T_{even} and T_{odd} to be the orbits

$$G_{even} \cdot \Gamma_{even} \quad \text{and} \quad G_{odd} \cdot \Gamma_{odd}.$$

We define a new path-metric τ on the complex X by declaring the closure of each component of $X \setminus (T_{even} \cup T_{odd})$ to be a unit Euclidean square. Clearly, the group G acts on (X, τ) isometrically and (X, τ) is a CAT(0) metric space.

The groups G_{even}, G_{odd} act on the graphs T_{even}, T_{odd} with the fundamental domains $\Gamma_{even}, \Gamma_{odd}$ respectively. It therefore follows that if $g \in G$ and $g(T_{even}) \cap T_{even} \neq \emptyset$ (resp. $g(T_{odd}) \cap T_{odd} \neq \emptyset$) then $g \in G_{even}$ (resp. $g \in G_{odd}$).

Lemma 2.6 *The subgraphs $T_{even}, T_{odd} \subset X$ are convex subsets in X isometric (with respect to the path-metric induced from (X, τ)) to a tree.*

Proof (1) First, let us prove that $T_{even}, T_{odd} \subset X$ are convex. Since X is a CAT(0) space, and T_{even}, T_{odd} are connected, it suffices to test convexity at each vertex of T_{even}, T_{odd} . However, by the definition of the metric τ , the angle between different edges of T_{even} (resp. T_{odd}) at each vertex of T_{even} (resp. T_{odd}) is $\geq \pi$. Therefore convexity follows.

(2) Since, $T_{even}, T_{odd} \subset (X, \tau)$ are convex, it follows that they are contractible. Therefore these graphs are isometric to metric trees. It is clear that T_{even} and T_{odd} are isometric to each other. □

Corollary 2.7 *Each subgroup $G_{\text{even}}, G_{\text{odd}}$ is a quasi-convex subgroup of G .*

Proof Cayley graphs $\Gamma_G, \Gamma_{G_{\text{even}}}, \Gamma_{G_{\text{odd}}}$ are quasi-isometric to $(X, \tau), T_{\text{even}}, T_{\text{odd}}$ respectively. Recall that X is Gromov-hyperbolic. Therefore, by combining stability of quasi-geodesics in Gromov-hyperbolic geodesic metric spaces and convexity of $T_{\text{even}}, T_{\text{odd}} \subset X$, we conclude that $G_{\text{even}}, G_{\text{odd}}$ are quasi-convex subgroups of G . \square

We define functions $\text{odd}(i)$ and $\text{even}(i)$ by

$$\text{odd}(i) = \begin{cases} i, & \text{if } i \text{ is odd} \\ i - 1, & \text{if } i \text{ is even} \end{cases} \quad \text{and} \quad \text{even}(i) = \begin{cases} i, & \text{if } i \text{ is even} \\ i - 1, & \text{if } i \text{ is odd} \end{cases}$$

To motivate the following definition, observe that the group G is generated by the elements g_i of the vertex subgroups G_{x_i} . It will be very important for the later analysis to find out which products of pairs of generators $f = h_j^{-1}g_i, h_j \in G_{x_j}, g_i \in G_{x_i}$, preserve the trees $T_{\text{odd}}, T_{\text{even}}$. Note that the answer is clear for some of these products:

- (a) If g_i preserves $e_{\text{odd}(i)}$ (resp. $e_{\text{even}(i)}$) and h_j^{-1} preserves $e_{\text{odd}(j)}$ (resp. $e_{\text{even}(j)}$), then $g_i, h_j^{-1} \in G_{\text{odd}}$ (resp. G_{even}) and hence f also preserves T_{odd} (resp. T_{even}).
- (b) If $i = j$ and the product $f = h_i^{-1}g_i$ preserves the edge $e_{\text{odd}(i)}$ (resp. $e_{\text{even}(i)}$), then f also preserves T_{odd} (resp. T_{even}).

Accordingly, define finite subsets $\Phi'_{\text{even}}, \Phi_{\text{even}}, \Phi'_{\text{odd}}, \Phi_{\text{odd}} \subset G$ as follows:

- (1) Φ'_{even} consists of products $h_j^{-1}g_i, g_i \in G_{x_i}, h_j \in G_{x_j}$ with either (a) $i \neq j$ and $g_i \in G_{e_{\text{even}(i)}}$ and $h_j \in G_{e_{\text{even}(j)}}$, or (b) $i = j$ and $h_i^{-1}g_i \in G_{e_{\text{even}(i)}}$.
- (2) $\Phi_{\text{even}} := \{h_j^{-1}g_i : g_i \in G_{x_i}, h_j \in G_{x_j}\} \setminus \Phi'_{\text{even}}$.
- (3) Φ'_{odd} consists of products $h_j^{-1}g_i, g_i \in G_{x_i}, h_j \in G_{x_j}$ with either (a) $i \neq j$ and $g_i \in G_{e_{\text{odd}(i)}}$ and $h_j \in G_{e_{\text{odd}(j)}}$, or (b) $i = j$ and $h_i^{-1}g_i \in G_{e_{\text{odd}(i)}}$.
- (4) $\Phi_{\text{odd}} := \{h_j^{-1}g_i : g_i \in G_{x_i}, h_j \in G_{x_j}\} \setminus \Phi'_{\text{odd}}$.

Observe that $\Phi'_{\text{even}}, \Phi'_{\text{odd}}$ are contained in the subgroups $G_{\text{even}}, G_{\text{odd}}$ respectively.

Proposition 2.8 $\Phi_{\text{odd}} \cap G_{\text{odd}} = \emptyset, \Phi_{\text{even}} \cap G_{\text{even}} = \emptyset$. In other words, among the products of the generators, only the “obvious” ones preserve the trees T_{odd} and T_{even} .

Proof We prove that $\Phi_{odd} \cap G_{odd} = \emptyset$, the second assertion is proved by relabeling. We have to show that

$$h_j^{-1}g_i T_{odd} = T_{odd} \Rightarrow h_j^{-1}g_i \in \Phi'_{odd}.$$

Throughout the proof we use the metric τ on X . We begin with the following

Observation 2.9 *Let $g_l \in G_{x_l} \setminus G_{e_{odd(l)}}$ and z be a vertex of $g_l(F)$. Then there is a geodesic segment $\sigma = \overline{zz'}$ $\subset (X, \tau)$ from z to a point $z' \in \Gamma_{odd}$, which intersects T_{odd} orthogonally (at the point z') and which is entirely contained in $F \cup g_l(F)$. For instance, if $g_l(F) \cap F = \{x_l\}$, then the segment σ equals $\overline{zx_l} \cup \overline{x_l m_{odd(l)}}$.*

Moreover, unless $z \in g_l(F) \cap F$, the length of the segment σ is strictly greater than 1. In particular, $d(z, T_{odd}) > 1$.

Let $f := h_j^{-1}g_i$ and assume that $f(T_{odd}) = T_{odd}$. Hence the edge $f(e_{odd(i)}) \subset X$ intersects T_{odd} orthogonally in its midpoint $f(m_{odd(i)})$. The segment $f(e_{odd(i)})$ equals \overline{zw} where $z := h_j^{-1}(x_i)$ and $w := f(x_{i+1})$ (if i is odd), and $w := f(x_{i-1})$ (if i is even). In any case, $d(z, T_{odd}) = 1$.

Suppose that $h_j \notin G_{odd(j)}$. Then, unless $z = x_j$, by applying Observation 2.9 to $l = j$, we obtain a contradiction with $d(z, T_{odd}) > 1$. Therefore $z = x_j$ and $i = j$, $f \in G_{x_i}$. Thus we apply Observation 2.9 to $l = j$, $g_l = f$ and the vertex $w \in f(F)$, and conclude that $w \in F$ as well. Then $f(e_{odd(i)}) = e_{odd(i)}$ and hence $f = h_j^{-1}g_i \in \Phi'_{odd}$.

Hence we conclude that $h_j \in G_{odd(j)}$. We now use the fact that $g_i^{-1}h_j(T_{odd}) = T_{odd}$ and reverse the roles of g_i and h_j . The same argument as above then implies that either

- (a) $g_i \in G_{odd(i)}$, or
- (b) $i = j$, $f^{-1} \in G_{odd(j)}$.

In Case (a), $h_j \in G_{odd(j)}$, $g_i \in G_{odd(i)}$ and thus $f \in \Phi'_{odd}$; in Case (b) $i = j$, $f \in G_{odd(j)}$ and thus $f \in \Phi'_{odd}$. □

Corollary 2.10 *For each $g \in \Phi_{even}, h \in \Phi_{odd}$ we have $gT_{even} \cap T_{even} = \emptyset$, $hT_{odd} \cap T_{odd} = \emptyset$.*

Recall the following definition:

Definition 2.11 A group G is said to satisfy LERF property with respect to a subgroup $H \subset G$ if one of the following equivalent conditions holds:

- (a) For each finite subset $F \subset G \setminus H$ there exists a homomorphism $\phi: G \rightarrow \bar{G}$, where \bar{G} is a finite group and $\phi(H) \cap \phi(F) = \emptyset$.
- (b) For each finite subset $F \subset G \setminus H$ there exists a finite index subgroup $G' \subset G$ so that $H \subset G'$ and $F \cap G' = \emptyset$.

Note that for $H = \{1\}$ the above definition amounts to residual finiteness of G .

Corollary 2.12 *There is an epimorphism $\phi: G \rightarrow \bar{G}$ where \bar{G} is a finite group and $\phi(\Phi_{\text{even}}) \cap \phi(G_{\text{even}}) = \emptyset$, $\phi(\Phi_{\text{odd}}) \cap \phi(G_{\text{odd}}) = \emptyset$.*

Proof According to [24], the group G satisfies the LERF property with respect to each quasi-convex subgroup. Thus there are finite quotients

$$\phi': G \rightarrow \bar{G}', \phi'': G \rightarrow \bar{G}''$$

so that $\phi'(G_{\text{even}}) \cap \phi'(\Phi_{\text{even}}) = \emptyset$, $\phi'(G_{\text{odd}}) \cap \phi'(\Phi_{\text{odd}}) = \emptyset$. Then define the homomorphism $\phi = (\phi', \phi''): G \rightarrow \bar{G}' \times \bar{G}''$ and let the group \bar{G} be the image of ϕ . \square

Let $q: X \rightarrow \bar{X} := X/\text{Ker}(\phi)$ denote the quotient map; the group G acts on the compact complex \bar{X} through the quotient group \bar{G} . We let $\bar{T}_{\text{even}}, \bar{T}_{\text{odd}}$ denote the projections of the trees $T_{\text{even}}, T_{\text{odd}}$ to the complex \bar{X} .

Lemma 2.13 *Suppose that $g_i \in G_{x_i}, h_j \in G_{x_j}$ and $g_i \bar{T}_{\text{even}} \cap h_j \bar{T}_{\text{even}} \neq \emptyset$ (resp. $g_i \bar{T}_{\text{odd}} \cap h_j \bar{T}_{\text{odd}} \neq \emptyset$). Then $h_j^{-1} g_i \in \Phi'_{\text{even}}$ (resp. $h_j^{-1} g_i \in \Phi'_{\text{odd}}$).*

Proof If $g_i \bar{T}_{\text{even}} \cap h_j \bar{T}_{\text{even}} \neq \emptyset$ then $g_i \bar{T}_{\text{even}} = h_j \bar{T}_{\text{even}}$. It follows that $kh_j^{-1} g_i \in G_{\text{even}}$ for some $k \in \text{Ker}(\phi)$; thus $\phi(h_j^{-1} g_i) \in \phi(G_{\text{even}})$ which implies that $h_j^{-1} g_i \in \Phi'_{\text{even}}$. The argument for \bar{T}_{odd} is the same. \square

The graphs $\bar{T}_{\text{even}}, \bar{T}_{\text{odd}}$ determine finite subsets $S_{\text{even}}, S_{\text{odd}}$ of the set $\text{Edges}(\bar{X})$ consisting of those edges in \bar{X} which intersect $\bar{T}_{\text{even}}, \bar{T}_{\text{odd}}$ nontrivially. Let ξ, η denote the characteristic functions of the subsets $S_{\text{even}}, S_{\text{odd}} \subset \text{Edges}(\bar{X})$, normalized to have unit norm in the (finite-dimensional) Hilbert space $H := L^2(\text{Edges}(\bar{X}))$. The group G acts on H by precomposition. We let $V \subset H$ denote the span of the subset $G \cdot \{\xi, \eta\} \subset H$ and let p be the dimension of V .

Corollary 2.14 (1) *The subgroups $G_{\text{even}}, G_{\text{odd}}$ fix the vectors $\xi, \eta \in V$ respectively.*

(2) *The set*

$$\Sigma = \{g^*(\xi), g^*(\eta) : g \in G_{x_1} \cup \dots \cup G_{x_n}\}$$

is an orthonormal system in V .

(3) *For all $g, h \in G_{x_1} \cup \dots \cup G_{x_n}$:*

(a) $g^*(\xi) \neq h^*(\eta)$.

(b) $g^*(\xi) = h^*(\xi)$ (resp. $g^*(\eta) = h^*(\eta)$) iff $h^{-1}g \in \Phi'_{\text{even}}$ (resp. $h^{-1}g \in \Phi'_{\text{odd}}$).

Let $Lk_{x_i}^\circ$ denote the vertex set of the link of x_i in X .

Corollary 2.15 *The representation $G \curvearrowright V$ contains subrepresentations*

$$G_{x_i} \curvearrowright \text{Vect}(Lk_{x_i}^\circ),$$

so that the orthonormal vectors $G_{x_i} \cdot \{\xi, \eta\}$ are identified with the vectors $G_{x_i} \cdot Lk_{x_i}^\circ$.

Proof Let us consider the case when i is odd, since the other case is analogous. Observe that the stabilizer in G_{x_i} of the vector $\xi \in V$ (resp. $\eta \in V$) is the group $G_{x_i} \cap G_{\text{even}} = G_{\text{even}(i)}$ (resp. $G_{x_i} \cap G_{\text{odd}} = G_{\text{odd}(i)}$); the stabilizer in G_{x_i} of the vector f_i^- is $G_{\text{even}(i)}$. Thus we construct an isometric embedding $\text{Vect}(Lk_{x_i}^\circ) \rightarrow V$ by sending f_i^-, f_i^+ to ξ, η respectively, and then extending this map equivariantly to the orthonormal basis $Lk_{x_i}^\circ$. \square

3 Hyperbolic trigonometry

Consider a regular right-angled hyperbolic n -gon $F \subset \mathbb{H}^2$ ($n \geq 5$). Let a_n denote its side-length, ρ_n the radius of the circumscribed circle, r_n the radius of the inscribed circle, b_n the length of the shortest diagonal in F (ie, a diagonal which cuts out a triangle from F); see Figure 1. We then have:

$$\begin{aligned} \cosh(a_n) &= 1 + 2 \cos\left(\frac{2\pi}{n}\right), & \cosh\left(\frac{a_n}{2}\right) &= \sqrt{2} \cos\left(\frac{\pi}{n}\right), & \cosh(b_n) &= \cosh^2(a_n), \\ \cosh(r_n) &= \frac{1}{\sqrt{2} \sin\left(\frac{\pi}{n}\right)}, & \cosh(\rho_n) &= \cosh(r_n) \cosh\left(\frac{a_n}{2}\right). \end{aligned}$$

Note that a_n, b_n, r_n, ρ_n are strictly increasing functions of n .




Figure 1: Geometry of a hyperbolic n -gon

Consider a *Lambert quadrilateral* Q with one zero angle: Q is a quadrilateral in \mathbb{H}^2 with one ideal vertex (where Q has zero angle) and three finite vertices where the angles are $\pi/2$ (Figure 2). Let x, y denote the lengths of the finite sides of Q . Then

$$\sinh(x) \sinh(y) = \cos(0) = 1$$

(see [2, 7.17.1]), or, equivalently

$$\cosh^2(x) \cosh^2(y) = \cosh^2(x) + \cosh^2(y).$$

Thus, if we have two segments $E = \overline{xx'}$, $E' = \overline{x'x''}$ in \mathbb{H}^2 which intersect at the point x' where they meet orthogonally, then the necessary and sufficient condition for $\text{Bis}(E) \cap \text{Bis}(E') = \emptyset$ is

$$\cosh^2(|E|/2) \cosh^2(|E'|/2) \geq \cosh^2(|E|/2) + \cosh^2(|E'|/2),$$

equivalently

$$\sinh(|E|/2) \sinh(|E'|/2) \geq 1.$$

We will refer to these inequalities as the *disjoint bisectors test*.

Lemma 3.1 *Suppose that $E, E' \subset \mathbb{H}^2$ are segments which meet orthogonally at a vertex, where $|E| = 2\rho_n$ and $|E'| = a_n$. Then*

$$\emptyset = \overline{\text{Bis}(E)} \cap \overline{\text{Bis}(E')} \subset \overline{\mathbb{H}^2},$$

provided that $n \geq 7$; in case $n = 6$ we have:

$$\overline{\text{Bis}(E)} \cap \overline{\text{Bis}(E')} \subset \partial_\infty \mathbb{H}^2,$$

is a point at infinity.




Figure 2: A Lambert quadrilateral

Proof Applying the disjoint bisectors test to $|E| = 2\rho_6$, $|E'| = a_6$ we get the equality. Hence the bisectors meet at infinity in case $n = 6$ and are within positive distance from each other if $n \geq 7$. \square

Lemma 3.2 Suppose that $E, E' \subset \mathbb{H}^2$ are segments which meet orthogonally at a vertex, where $|E| = b_n$ and $|E'| = a_n$, $n \geq 7$. Then

$$\emptyset = \overline{\text{Bis}(E)} \cap \overline{\text{Bis}(E')} \subset \overline{\mathbb{H}^2}.$$

Proof Since $b_n \geq b_7, a_n \geq a_7$ it suffices to prove lemma in case $n = 7$. Note that $a_7 > a_6$ and

$$b_7 \approx 2.302366350 > 2\rho_6 = 2.292431670$$

Hence the assertion follows from Lemma 3.1. \square

Below is another application of the *disjoint bisectors test*. Consider three segments s, s', s'' in \mathbb{H}^3 of the length x, y, x respectively, which are mutually orthogonal and so that $s \cap s' = p, s' \cap s'' = q, s' = \overline{pq}$, see Figure 3.

Corollary 3.3 If $x = b_n, y = a_n, n \geq 5$, then $\overline{\text{Bis}(s)} \cap \overline{\text{Bis}(s'')} = \emptyset$.

Proof It suffices to prove the corollary for $n = 5$. We first compute the length $z = 2t$ of the segment s''' coplanar to s and s' such that $\text{Bis}(s) = \text{Bis}(s''')$. By considering the Lambert's quadrilateral with angle ϕ we get:

$$\begin{cases} \cosh(t) \sin(\phi) = \cosh(x/2) \\ \cosh(y) \sin(\phi) = \cosh(h) \\ \sinh(x/2) \sinh(h) = \cos(\phi) \end{cases}$$




Figure 3: Three orthogonal segments

Thus
$$\cosh(t) = \frac{\cosh(y) \cosh(x/2)}{\cosh(h)}.$$

On the other hand, the last two equations in the above system imply that

$$\cosh(h) = \frac{\cosh(x/2)}{\sqrt{\sinh^2(x/2) + 1/\cosh^2(y)}}.$$

Therefore

$$\cosh(z/2) = \sqrt{1 + \sinh^2(x/2) \cosh^2(y)}.$$

By applying the *disjoint bisectors test* to s'' , s''' we get:

$$\text{Bis}(s) \cap \text{Bis}(s'') = \emptyset \iff \sinh^2(x/2) \cosh(y) \geq 1.$$

Lastly, we have:

$$\cosh(a_5) \approx 1.6180, \quad \sinh(b_5/2) \approx 1.85123.$$

Therefore $\sinh^2(b_5/2) \cosh(a_5) > 1$. □

4 Quasi-isometric maps of polygonal complexes

Suppose that X is a simply-connected 2-dimensional regular cell complex which is equipped with a path-metric so that:

1. Each face is isometric to a right-angled regular n -gon in \mathbb{H}^2 (of course, $n \geq 5$).
2. The complex X is *negatively curved*, ie, for each vertex $x \in X$ the length of the shortest embedded loop in $Lk_x(X)$ is at least 2π .

Theorem 4.1 *Suppose that $\mu: X \rightarrow \mathbb{H}^p$ is a continuous map which is a (totally-geodesic) isometric embedding on each face of X . We also assume that for each pair of faces $F', F'' \subset X$ which intersect nontrivially a common face $F \subset X$, we have:*

$$\text{Span}(\mu(F')) \perp \text{Span}(\mu(F'')).$$

Then μ is a quasi-isometric embedding.

Proof Throughout the proof we will be using the notation a_n, b_n, ρ_n for various distances in a regular right-angled hyperbolic n -gon, see Section 3.

Since the inclusion $X^{(1)} \hookrightarrow X$ is a quasi-isometry, it suffices to check that $X^{(1)} \xrightarrow{\mu} \mathbb{H}^p$ is a quasi-isometric embedding. Since μ is 1-Lipschitz, it is enough to show that $d(\mu(z), \mu(w)) \geq C \cdot d(z, w)$ for some $C = C(X) > 0$ and all $z, w \in X^{(0)}$. We first give a proof in case $n \geq 6$ and then explain how to modify it for $n = 5$.

Let $\tilde{\gamma} \subset X^{(1)}$ be an (oriented) geodesic segment connecting z to w . We start by replacing (in case when n is even) each subsegment of $\tilde{\gamma}$ connecting antipodal points in a face F of X with a geodesic segment within F . We will call the resulting (oriented) curve $\gamma \subset X$. Clearly,

$$\text{Length}(\mu(\gamma)) \leq \text{Length}(\mu(\tilde{\gamma})) \leq \frac{n}{2} \text{Length}(\mu(\gamma)),$$

so it suffices to get a lower bound on $\text{Length}(\mu(\gamma))$. We will refer to the edges of γ connecting antipodal points of faces as *diagonals* in γ .

Remark 4.2 Suppose that $\overline{xx'}, \overline{x'x''}$ are (distinct) diagonals in γ , contained in faces F, F' respectively. Then $F \cap F' = \{x'\}$: Otherwise $\tilde{\gamma} \subset X^{(1)}$ would not be a geodesic as there exists a shorter path along the boundaries of F and F' ; see Figure 4. In particular, $\mu(\overline{xx'}) \perp \mu(\overline{x'x''})$ in \mathbb{H}^p .




Figure 4: Diagonals

We will regard γ as concatenation of consecutive segments e_0, e_1, \dots . We define a collection $\mathcal{BE}(\gamma)$ of *bisected edges* E_i in γ inductively as follows:

- (1) Let $E_0 = e_0 \subset \gamma$ be the first edge of γ .
- (2) Suppose that $E_i = e_j \subset \gamma$ was chosen, $i \geq 0$. We will take as $E_{i+1} = e_k$, $k > i$, the first edge on γ following E_i which satisfies two properties:
 - (a) If $e_i \cap e_k \neq \emptyset$ then either e_i or e_k is a diagonal.
 - (b) e_i, e_k do not belong to a common face in X .

Proposition 4.3 *Suppose that $n \geq 6$. Then the edges $E_i, i = 0, 1, \dots$ in γ satisfy the following:*

- (1) *There exists a constant $c = c(X)$ such that $d(E_i, E_{i+1}) \leq c$ in X .*
- (2) *$\overline{\text{Bis}(\mu(E_i))} \cap \overline{\text{Bis}(\mu(E_{i+1}))} \subset \mathbb{H}^p$ is empty unless $n = 6$ and either E_i , or E_{i+1} is not a diagonal. In case $n = 6$ and at least one of these segments is not a diagonal, the bisectors $\text{Bis}(\mu(E_i)), \text{Bis}(\mu(E_{i+1})) \subset \mathbb{H}^p$ are disjoint but have a common ideal point in $\partial_\infty \mathbb{H}^p$.*
- (3) *For all edges $e_j \subset \gamma$ between E_i, E_{i+1} , their images $\mu(e_j)$ are disjoint from $\text{Bis}(\mu(E_i)) \cup \text{Bis}(\mu(E_{i+1}))$.*




Figure 5: Diagonal and an edge

Proof (1) It is clear from the construction, that E_i, E_{i+1} are separated by at most $n/2$ edges on γ . Hence the first assertion follows.

(2) There are several cases we have to consider.

(a) Suppose that either E_i or E_{i+1} is a diagonal (see Figure 5) of the length $2\rho_n$ in the notation of section 3. Then these segments share a common vertex x' and it follows that $\mu(E_i) \perp \mu(E_{i+1})$ (see the Remark above). The worst case occurs when $n = 6$ and one of the segments is an edge of a face of X : The bisectors $\text{Bis}(\mu(E_i)), \text{Bis}(\mu(E_{i+1}))$ are disjoint in \mathbb{H}^p but have a common ideal point (see Lemma 3.1). Since, as n increases, both side-lengths and lengths of diagonals in regular right-angled n -gons in \mathbb{H}^2 strictly increase, it follows that

$$\overline{\text{Bis}(\mu(E_i))} \cap \overline{\text{Bis}(\mu(E_{i+1}))} = \emptyset, \forall n \geq 7.$$

(b) Consider now the case when neither $E_i = e_k$ nor E_{i+1} is a diagonal, E_i is contained in a face F and there exists at least one edge (say, e_{k+1}) between E_i, E_{i+1} which is contained in the face F . Then, by the construction, $E_{i+1} = \overline{x'x''}$ is not contained in F but shares the common point x' with F . Thus $\mu(E_{e+1}) \perp \mu(F)$.

Observe now that *there is a vertex $y \in F$ such that the segments $\overline{yx'}$ and E_i have the same bisector in F* . To find this vertex simply apply the reflection in $\text{Bis}(E_i)$ to the vertex x' : This symmetry preserves F and sends the vertex x'




Figure 6: Bisectors

to a vertex $y \in F$. See Figure 6. Since $x, x' \in F$ are not antipodal, $\overline{yx'}$ is not an edge of F .

Clearly, $\text{Bis}(\mu(\overline{yx'})) = \text{Bis}(\mu(E_i))$. Hence the problem reduces to verifying that the bisectors $\text{Bis}(\mu(\overline{yx'}))$, $\text{Bis}(\mu(E_{i+1}))$ (or their closures in $\overline{\mathbb{H}^P}$) are disjoint. We note that in case $n = 6$ the segment $\overline{yx'}$ connects antipodal points in F ; hence the proof in this case reduces to (a).

Assume now that $n \geq 7$, then, since $\overline{yx'}$ is not an edge of F , $|xy'| \geq b_n \geq b_7$, $|x'x''| = a_n \geq a_7$ and

$$\overline{\text{Bis}(\mu(E_i))} \cap \overline{\text{Bis}(\mu(E_{i+1}))} = \emptyset,$$

follows from Lemma 3.2.

(c) The last case to consider is when E_i, E_{i+1} are not diagonals and they are separated by exactly one edge $e \subset \gamma$ (this edge cannot be a diagonal in this case), which is not contained in a common face with E_i nor with E_{i+1} . Then the edges

$$\mu(E_i), \mu(e), \mu(E_{i+1}) \subset \mathbb{H}^P$$

intersect orthogonally. The lengths of these edges are equal to $a(n) \geq a(6)$. Hence, (as in Case (b)) we replace $\mu(E_i)$ with a segment s of the length $\geq 2\rho_6$ which meets $\mu(E_{i+1})$ orthogonally at the point $\mu(e) \cap \mu(E_{i+1})$. Therefore, by

applying again Lemma 3.1, the bisectors $\text{Bis}(\mu(E_i)), \text{Bis}(\mu(E_{i+1}))$ are disjoint; their closures in \mathbb{H}^P are disjoint provided $n \geq 7$.

This proves the second assertion of the Proposition. The third assertion is clear from the construction: For instance, in Case (b) the edges e_j between E_i, E_{i+1} are all contained in the face F . Therefore they are disjoint from the bisector of E_i within F , which implies the assertion about their images in \mathbb{H}^P . On the other hand, the edge μE_{i+1} is orthogonal to $\mu(F)$, hence $\text{Bis}(\mu E_{i+1})$ is disjoint from μF . \square

Now, let us finish the proof that μ is a quasi-isometric embedding. Suppose that $\gamma \subset X$ has length L , then the subset $\mathcal{BE}(\gamma)$ consists of $\ell \approx L/c$ bisected edges E_i (here c is the constant from Proposition 4.3, Part 1). Hence the geodesic segment $\gamma^* = \overline{\mu(z)\mu(w)}$ in \mathbb{H}^P connecting the end-points of $\mu(\gamma)$ crosses ℓ bisectors $\text{Bis}(\mu E_i)$. In case $n \geq 7$, the bisectors $\text{Bis}(\mu E_i), \text{Bis}(\mu E_{i+1})$ are separated by distance $\delta = \delta(X) > 0$, hence the length of γ^* is at least $\ell\delta$. Since $\ell \approx L/c$, we conclude that $d_{\mathbb{H}^P}(x, y) \geq \text{Const} \cdot L/\delta$. It follows that μ is a quasi-isometry.

Now, consider the exceptional case $n = 6$. We claim that for each i the intersection points $\overline{\text{Bis}(\mu E_i)} \cap \overline{\text{Bis}(\mu E_{i+1})}$ and $\overline{\text{Bis}(\mu E_{i+2})} \cap \overline{\text{Bis}(\mu E_{i+1})}$ are distinct. Given this, instead of the collection $\mathcal{BE}(\gamma)$ we would consider the collection of edges $E_i \in \mathcal{BE}(\gamma)$ for even i , then $\overline{\text{Bis}(\mu E_i)} \cap \overline{\text{Bis}(\mu E_{i+2})} = \emptyset$ for all even i and we are done by the same argument as for $n \geq 7$.

Case I We begin with the case when $E_i \subset F_i, E_{i+2} \subset F_{i+2}$ are diagonals and $E_{i+1} \subset F_i$ is not. (Here F_i are faces of X .) Then $E_i \cap E_{i+1}, E_{i+2} \cap E_{i+1}$ are the end-points of E_{i+1} . Therefore

$$\begin{aligned} \xi_i &= \overline{\text{Bis}(\mu E_i)} \cap \overline{\text{Bis}(\mu E_{i+1})} \in \partial_\infty \text{Span}(\mu(E_i) \cup \mu(E_{i+1})), \\ \xi_{i+1} &= \overline{\text{Bis}(\mu E_{i+1})} \cap \overline{\text{Bis}(\mu E_{i+2})} \in \partial_\infty \text{Span}(\mu(E_{i+1}) \cup \mu(E_{i+2})). \end{aligned}$$

However, by the assumptions on μ ,

$$\text{Span}(\mu(F_{i+2})) \perp \text{Span}(\mu(F_i))$$

Thus

$$\text{Span}(\mu(E_i) \cup \mu(E_{i+1})) \cap \text{Span}(\mu(E_{i+1}) \cup \mu(E_{i+2})) = \text{Span}(\mu(E_{i+1})).$$

Since it is clear that $\xi_i \notin \partial_\infty \text{Span}(\mu(E_{i+1}))$, we conclude that $\xi_i \neq \xi_{i+1}$ and the assertion follows.

We will reduce the case of a general triple of edges E_i, E_{i+1}, E_{i+2} to the Case I discussed above. We consider only one other case, the arguments in the rest of cases are identical to:

Case II Suppose that pairs of edges $E_i \subset F_i, E_{i+1} \subset F_{i+1}, E_{i+2} \subset F_{i+2}$ are as in Figure 7. We take the diagonals $D_i \subset F_i, D_{i+2} \subset F_{i+2}$ so that

$$\text{Bis}(\mu D_i) = \text{Bis}(\mu E_i), \text{Bis}(\mu D_{i+2}) = \text{Bis}(\mu E_{i+2}).$$

Now the proof reduces to the Case I.




Figure 7

Finally, consider the case of pentagons (ie $n = 5$). We define the collection $\mathcal{BE}(\gamma)$ of bisected edges E_i as before. Let E_1, E_2, E_3 be consecutive bisected edges. We will see that $\text{Bis}(\mu E_1) \cap \text{Bis}(\mu E_3) = \emptyset$. Since $n = 5$ we necessarily have: E_2 is separated by a unique edge $e \subset \gamma$ from E_1 and by a unique edge $e' \subset \gamma$ from E_3 , see Figure 8.

Note that it could happen that there is no face F_1 which contains E_1, e , nor a face F_3 which contains E_3, e' . However, in \mathbb{H}^p there exists a unique regular right angled pentagon which contains the edges $\mu(E_1), \mu(e)$ (resp. $\mu(E_3), \mu(e')$) in its boundary. Hence we will give a proof pretending that the corresponding face already exists in X . Observe that, similarly to our discussion above, the diagonals $D_1 \subset F_1, D_3 \subset F_3$ have the property that $\text{Bis}(\mu D_i) = \text{Bis}(\mu E_i)$, $i = 1, 3$. Thus it suffices to consider the triple of pairwise orthogonal segments: $\mu(D_1), \mu(E_2)$ and μD_3 in \mathbb{H}^p . The length of $\mu(E_2)$ equals a_5 , the lengths of




Figure 8

D_1, D_3 are equal to b_5 , hence

$$\overline{\text{Bis}(\mu D_1)} \cap \overline{\text{Bis}(\mu D_3)} = \emptyset$$

by Corollary 3.3. □

5 Proof of the main theorem

In this section establish

Theorem 5.1 *Suppose that $n = 2k$ is even. Then the group G admits an isometric properly discontinuous convex-cocompact action $\rho: G \curvearrowright \mathbb{H}^p$, where $p < \infty$ depends on the polygon \mathcal{P} .*

Proof Let X be the universal cover of the polygon of groups \mathcal{P} . We first construct a representation $\rho: G \rightarrow \text{Isom}(\mathbb{H}^p)$ for a certain p . We then produce a quasi-isometric ρ -equivariant embedding $\mu: X \rightarrow \mathbb{H}^p$. From this, via Lemma

2.2, it will follow that $\rho: G \curvearrowright \mathbb{H}^p$ is an isometric properly discontinuous convex-cocompact action.

Let Lk_x denote the link (in X) of the vertex x , similarly, let Lk_e denote the link of the edge e . Recall that Lk° denotes the vertex set of a link Lk . The set $Lk_{x_i}^\circ$ contains two distinguished elements: f_i^+, f_i^- which correspond to the directions from x_i toward x_{i+1} and toward x_{i-1} respectively. We define the subsets $Lk_{x_i}^\bullet := Lk_{x_i}^\circ \setminus \{f_i^+, f_i^-\}$.

Observe that the directions $\eta_i \in Lk_{x_i}^\circ, \eta_{i+1} \in Lk_{x_{i+1}}^\circ$ belong to the boundary of a common face in X if and only if there exists $g \in G_{e_i}$ so that

$$\eta_i = g(f_i^\pm), \eta_{i+1} = g(f_{i+1}^\pm).$$

Step 1: Construction of ρ

It is clear that to construct a representation $\rho: G \rightarrow \text{Isom}(\mathbb{H}^p)$ (for some p) we have to produce a collection of faithful representations

$$\rho_i: G_{x_i} \rightarrow \text{Isom}(\mathbb{H}^p)$$

so that the following diagram commutes:

$$\begin{array}{ccccc} G_{x_i} & \longleftarrow & G_{e_i} & \longrightarrow & G_{x_{i+1}} \\ & \searrow \rho_i & & \rho_{i+1} \swarrow & \\ & & \text{Isom}(\mathbb{H}^p) & & \end{array}$$

Embed F isometrically (as a convex, regular, right-angled polygon) in the hyperbolic plane \mathbb{H}^2 . Via this embedding we will identify the directions $f_i^\pm \in Lk_{x_i}$ with the unit vectors $\vec{f}_i^\pm \in T_{x_i}\mathbb{H}^2$ which are tangent to the sides of F .

In what follows we will adopt the following convention. Given a number p and a totally-geodesic embedding $\mathbb{H}^2 \subset \mathbb{H}^p$, we observe that the normal bundle $N(\mathbb{H}^2)$ of \mathbb{H}^2 in \mathbb{H}^p admits a canonical flat orthogonal connection (invariant under the stabilizer of \mathbb{H}^2 in $\text{Isom}(\mathbb{H}^p)$). Thus, given normal vectors $\nu \in N_x(\mathbb{H}^2), \nu' \in N_{x'}(\mathbb{H}^2)$, we have well-defined scalar product $\nu \cdot \nu'$ and hence the notion of orthogonality $\nu \perp \nu'$. Set $V_i := T_{x_i}\mathbb{H}^p$ and let $N_i \subset V_i$ denote the orthogonal complement to $T_{x_i}\mathbb{H}^2$. We define $R_i \in \text{Isom}(\mathbb{H}^p)$ to be the isometric reflection in the bisector of the edge e_i of $F \subset \mathbb{H}^p$. Set $J_i := R_{i-1} \circ \dots \circ R_1$, for $i = 2, \dots, n + 1$; observe that $J_{n+1} = Id$.

Remark 5.2 The fact that the identity $J_{n+1} = Id$ fails if n is odd is one of the reasons why our construction requires n to be an even number. An attempt to apply the constructions below to odd n lead to a representation of a certain extension of the group G rather than of G itself.

Proposition 5.3 *Suppose that $n = 2k$ is even. Then there exists a natural number p and a collection of faithful isometric linear actions $d\rho_i: G_{x_i} \curvearrowright V_i$, $i = 1, \dots, n$, so that the following hold:*

(1) *Each representation $d\rho_i$ contains a subrepresentaion $G_{x_i} \curvearrowright Vect(Lk_{x_i}^\circ)$, so that the unit basis vectors $f_i^\pm \in Lk_{x_i}^\circ \subset Vect(Lk_{x_i}^\circ)$ are identified with the vectors $\vec{f}_i^\pm \in V_i$.*

(2) *Each reflection $R_i: \mathbb{H}^p \rightarrow \mathbb{H}^p$ induces an isomorphism of $\mathbb{R}G_{e_i}$ -modules*

$$(V_i, d\rho_i(G_{e_i})) \rightarrow (V_{i+1}, d\rho_{i+1}(G_{e_i})).$$

(3) *“Orthogonality”: The spaces $Vect(Lk_{x_i}^\bullet) \subset N_i, Vect(Lk_{x_j}^\bullet) \subset N_j$ are mutually orthogonal, $|i - j| \geq 2, i, j \in \{1, \dots, n\}$. If $j = i + 1$ then we require orthogonality of the subspaces*

$$Vect(Lk_{x_i}^\circ \setminus G_{e_i} \cdot \{f_i^+, f_i^-\}) \subset N_i, Vect(Lk_{x_{i+1}}^\circ \setminus G_{e_i} \cdot \{f_{i+1}^+, f_{i+1}^-\}) \subset N_{i+1}.$$

Remark 5.4 (1) The assumption that the number of sides of F is even is used only in this part of the proof of the main theorem and very likely is just a technicality.

(2) The “orthogonality” property will be used to prove that the action $G \curvearrowright \mathbb{H}^p$ that we are about to construct, is discrete, faithful and convex-cocompact.

Before beginning the proof of the proposition we first make some observations (where we ignore the *orthogonality* issue). Suppose that we have constructed representations $d\rho_i$. We then “fold” these representations into a single orthogonal representation $G \curvearrowright V_1$ by composing each $d\rho_i$ with the composition of reflections $(R_1)_* \circ \dots \circ (R_{i-1})_*$, where $(R_j)_*$ is the isomorphism $O(V_{j+1}) \rightarrow O(V_j)$ which is induced by $dR_j|_{x_{j+1}}$. Note that under the action $G \curvearrowright V_1$ the vectors \vec{f}_1^+, \vec{f}_1^- are fixed by the “odd” and “even” subgroups G_{odd}, G_{even} , respectively. Moreover, the representation $G \curvearrowright V_1$ contains subrepresentations $G_{x_i} \curvearrowright Vect(Lk_{x_i}^\circ)$.

Recall that in Corollary 2.14 we have constructed a finite-dimensional orthogonal representation $G \curvearrowright V$ which satisfies the same properties as above: It contains unit vectors ξ, η fixed by G_{even}, G_{odd} respectively, and it contains subrepresentations $G_{x_i} \curvearrowright Vect(Lk_{x_i}^\circ)$. Therefore, to construct the representations $d\rho_i$ we begin with the action $G \curvearrowright V$ (which we identify with an action $G \curvearrowright V_1$) and then “unfold” it (using compositions of reflections R_i) to a collection of representations $d\rho_i$. This is the idea of the proof of Proposition 5.3.

The reader familiar with *bending* deformations of representations of groups into $\text{Isom}(\mathbb{H}^p)$ will notice that the “folding” and “unfolding” of representations discussed above is nothing but π -*bending*.

Proof We let $G \curvearrowright V$ denote the orthogonal representation constructed in Corollary 2.14. According to Corollary 2.15, the representation $G \curvearrowright V$ contains subrepresentations $G_{x_i} \curvearrowright \text{Vect}(Lk_{x_i}^\circ)$. Let p denote the dimension of V .

Our goal is to construct isometries $\phi_i: V \rightarrow V_i := T_{x_i}\mathbb{H}^p$. The actions $d\rho_i$ will be obtained by the conjugation:

$$d\rho_i: G_{x_i} \curvearrowright V_i := \phi_i \circ (G_{x_i} \curvearrowright V) \circ \phi_i^{-1}.$$

First take an arbitrary isometry $\phi_1: V \rightarrow V_1$ sending the unit vectors $\xi, \eta \in V$ to the vectors \vec{f}_1^-, \vec{f}_1^+ respectively. Now define isometries $\phi_j, j = 2, \dots, n+1$ by

$$\phi_i = J_i \circ \phi_1,$$

ie

$$\phi_{i+1} = R_i \circ \phi_i.$$

Note that $\phi_{n+1} = \phi_1$. Define the action $d\rho_i: G_{x_i} \curvearrowright V_i$ by conjugating via ϕ_i the action $G_{x_i} \curvearrowright V$.

The group G_{e_1} fixes the vector $\eta \in V$, hence G_{e_1} also fixes the vector $\phi_1(\eta) = \vec{f}_1^+$. Thus

$$dR_1 \circ d\rho_1|_{G_{e_1}} = d\rho_2|_{G_{e_1}}.$$

The same argument shows that

$$dR_i \circ d\rho_i|_{G_{e_i}} = d\rho_{i+1}|_{G_{e_i}}.$$

for all i . This proves (1) and (2). In what follows we will identify the spaces $\text{Vect}(Lk_{x_i}^\circ)$ with their images in $V_i, i = 1, \dots, n$.

We will check that the sets

$$Lk_{x_i}^\circ \setminus G_{e_i} \cdot \{f_i^+, f_i^-\} \subset N_i, Lk_{x_{i+1}}^\circ \setminus G_{e_i} \cdot \{f_{i+1}^+, f_{i+1}^-\} \subset N_{i+1}$$

are orthogonal to each other and will leave the remaining orthogonality assertion to the reader. Let

$$v \in Lk_{x_i}^\circ \setminus G_{e_i} \cdot \{f_i^+, f_i^-\} \subset N_i, w \in Lk_{x_{i+1}}^\circ \setminus G_{e_i} \cdot \{f_{i+1}^+, f_{i+1}^-\} \subset N_{i+1}.$$

In order to show that $v \perp w$ it suffices to verify that the corresponding vectors

$$v, w \in \Sigma \subset V$$

are distinct (recall that Σ is an orthonormal system in V). If, say, $v \in G \cdot \xi, w \in G \cdot \eta$ then $v \neq w$. Hence we will consider the case

$$v = g^* \eta \in G_{x_i} \cdot \eta, w = h^* \eta \in G_{x_{i+1}} \cdot \eta.$$

According to Corollary 2.14, if $g^*(\eta) = h^*(\eta)$ then $h^{-1}g \in \Phi'_{odd}$. In our case, $g \in G_{x_i}, h \in G_{x_{i+1}}$. Then $h^{-1}g \in \Phi'_{odd}$ means that either

(a) g, h do not have a common fixed vertex of F and $g \in G_{odd(i)}, h \in G_{odd(i+1)}$,

or

(b) g, h fix the same vertex of F and $g^*(\eta) = h^*(\eta)$.

In case (a), if i is odd then $g, h \in G_{e_i}$, and hence $g^* \eta = h^* \eta$ corresponds to the vectors $f_i^+ \in N_i, f_i^- \in N_{i+1}$. Therefore the equality $v = w$ implies that

$$\phi_i(v) \in G_{e_i} \cdot \vec{f}_i^+, \phi_{i+1}(w) \in G_{e_i} \cdot \vec{f}_{i+1}^-.$$

If i is even then both $g \in G_{e_{i-1}}, h \in G_{e_{i+1}}$ fix the vector η . Therefore the equality $v = w$ implies that

$$\phi_i(v) = \vec{f}_i^- \in G_{e_i} \cdot \vec{f}_i^-, \phi_{i+1}(w) = \vec{f}_{i+1}^+ \in G_{e_i} \cdot \vec{f}_{i+1}^+.$$

In case (b), we can assume that, say, $h \in G_{x_i} \cap G_{x_{i+1}} = G_{e_i}$. If i is odd then $h^* \eta = \eta$ and therefore $g^* \eta = \eta$. Therefore the equality $v = w$ implies that

$$\phi_i(v) = f_i^+ \in G_{e_i} \cdot f_i^+, \quad \phi_{i+1}(w) = f_{i+1}^- \in G_{e_i} \cdot f_{i+1}^-.$$

Lastly, assume that i is even. Then the vector $v = g^*(\eta) = w = h^*(\eta)$ corresponds to the vectors

$$\phi_i(w) = h(\vec{f}_i^-) \in G_{e_i} \cdot \vec{f}_i^- \subset N_i, \quad \phi_{i+1}(w) = h(\vec{f}_{i+1}^+) \in G_{e_i} \cdot \vec{f}_{i+1}^+ \subset N_{i+1}.$$

This proves the *orthogonality* assertion. □

Now, once we have constructed linear orthogonal representations $d\rho_i: G_{x_i} \curvearrowright V_i$, we extend them (by exponentiation) to isometric actions $\rho_i: G_{x_i} \curvearrowright \mathbb{H}^p$, which fix the points $x_i, i \in 1, \dots, n$. Observe that for each i the group $\rho_i(G_{e_i})$, resp. $\rho_i(G_{e_{i-1}})$, fixes the edge e_i , resp. e_{i-1} , of the polygon $F \subset \mathbb{H}^2$, since $d\rho_i(G_{e_i}), d\rho_i(G_{e_{i-1}})$, fix the vectors \vec{f}_i^+, \vec{f}_i^- respectively. Hence the reflection R_i commutes with the groups $\rho_i(G_{e_i})$ and $\rho_{i+1}(G_{e_i})$. Thus the representations $\rho_i, \rho_{i+1}: G_{e_i} \rightarrow \text{Isom}(\mathbb{H}^p)$ are the same. Therefore the representations ρ_i determine an isometric action $\rho: G \curvearrowright \mathbb{H}^p$.

Step 2: Discreteness of ρ

We will construct a ρ -equivariant continuous mapping $\mu: X \rightarrow \mathbb{H}^p$ satisfying the assumptions of Theorem 4.1. Since such μ is necessarily a quasi-isometric embedding (Theorem 4.1), by applying Lemma 2.2, we will conclude that $\rho: G \curvearrowright \mathbb{H}^p$ is properly discontinuous and convex-cocompact.

Recall that we have identified the face F of X with a regular right-angled hyperbolic polygon F in \mathbb{H}^2 , this defines the (identity) embedding $\mu: F \rightarrow \mathbb{H}^2$. Now, for each $g \in G$ we set

$$\mu|_{gF} := \rho(g) \circ \mu|_F.$$

Let us check that this mapping is well-defined:

- (1) If $g \in G_F$ then, by construction of ρ , $\rho(g)$ fixes the polygon $\mu(F)$ pointwise, hence $\mu \circ g|_F = \rho(g) \circ \mu|_F$ for $g \in G_F$.
- (2) If $g \in G_e$, where e is an edge of \mathcal{P} , then, by the construction, $\rho(g)$ fixes the edge $\mu(e)$ pointwise, hence $\mu \circ g|_e = \rho(g) \circ \mu|_e$ for $g \in G_e$.
- (3) The same argument applies to $g \in G_x$, for the vertices $x \in F$.

Hence $\mu: X \rightarrow \mathbb{H}^p$ is well-defined and thus it is a ρ -equivariant, continuous mapping which is an isometric totally-geodesic embedding on each face of X .

Lastly, we check the *orthogonality* condition required by Theorem 4.1. By equivariance, it is clear that we only need to verify orthogonality for the faces $F', F'' \subset X$ which are adjacent to the face $F \subset X$. We will see that this orthogonality condition will follow from the Assertion 3 of Proposition 5.3. There are several cases which may occur, we will check one of them and will leave the rest to the reader.

Suppose that $F' = g_i(F)$, $F'' = g_j(F)$, $g_i \in G_{x_i}$, $G_j \in G_{x_j}$ and

$$F' \cap F = \{x_i\}, \quad F'' \cap F = \{x_j\},$$

where $|i - j| \geq 2$. Then

$$T_{x_i}(\mu F') \subset N_i, \quad T_{x_j}(\mu F'') \subset N_j$$

and the vectors

$$dg_i(\vec{f}_i^\pm) \neq \vec{f}_i^\pm, \quad dg_j(\vec{f}_j^\pm) \neq \vec{f}_j^\pm$$

span $T_{x_i}(\mu F')$ and $T_{x_j}(\mu F'')$ respectively. According to the Assertion 3 of Proposition 5.3, we have:

$$dg_i(\vec{f}_i^\pm) \perp dg_j(\vec{f}_j^\pm).$$

(Recall that here the orthogonality is defined modulo the parallel translation along curves in $\text{Span}(F)$.) Since both $\text{Span}(\mu F')$, $\text{Span}(\mu F'')$ intersect $\text{Span}(F)$ orthogonally, the geodesic segment $\overline{x_i x_j} \subset \text{Span}(F)$ is orthogonal to both $\text{Span}(\mu F')$ and $\text{Span}(\mu F'')$. Therefore $\mu(F') \perp \mu(F'')$. \square

6 The odd case

In this section we will construct examples of negatively curved right-angled polygons of groups and their actions on \mathbb{H}^p in the case of the odd number of sides.

We define the following polygon of groups. Suppose that we are given finite groups $\Gamma_1, \dots, \Gamma_n$. Let F be an n -gon ($n \geq 3$). Below the indices i are taken modulo n . We assign the group $G_{x_i} = \Gamma_i \times \Gamma_{i+1}$ to each vertex x_i of F . We label each edge e_i of F by the group Γ_{i+1} . The homomorphisms $\Gamma_{i+1} \rightarrow \Gamma_i \times \Gamma_{i+1}$, $\Gamma_{i+1} \rightarrow \Gamma_{i+1} \times \Gamma_{i+2}$ are the natural isomorphisms to the second and the first factor respectively. We set $G_F := \{1\}$. In what follows, let \mathcal{P} denote the resulting polygon of groups (see Figure 9) and set $G := \pi_1(\mathcal{P})$.




Figure 9

Remark 6.1 Note that G is isomorphic to the *cyclic graph-product* of the groups $\Gamma_i, i = 1, \dots, n$ (see [7] for detailed definition). Indeed, the group G is

generated by the elements of $\Gamma_1, \dots, \Gamma_n$ subject to the relations:

$$[g_i, h_{i+1}] = 1, \forall g_i \in \Gamma_i, h_{i+1} \in \Gamma_{i+1}.$$

The polygon of groups \mathcal{P} is negatively curved provided that $n \geq 5$: For each vertex $x = x_i \in \mathcal{P}$, the link of x in the universal cover X of \mathcal{P} is the complete bipartite graph K_{t_{i-1}, t_i} , where $t_j := |G_{e_j}|, j = 1, \dots, n$.

Theorem 6.2 *Suppose that $n \geq 5$, and the polygon of groups \mathcal{P} is as above. Then G admits a properly discontinuous convex-cocompact action on \mathbb{H}^p for certain $p = p(\mathcal{P})$.*

Proof We will assume that $n = 2k - 1$ is odd. Let e_1, \dots, e_n denote the edges of the polygon \mathcal{P} . First, we construct a homomorphism $\rho: G \rightarrow \text{Isom}(\mathbb{H}^p)$ for some p , which is faithful on each vertex group. Let T denote the disjoint union

$$\bigcup_{i=1}^n Lk_{e_i},$$

where the link of e_i is taken in X . One can think of this set as the set of all flags: (e, f) , where e is an edge in F and f is a face of X containing e . Each group G_{e_i} acts naturally on Lk_{e_i} (since G_{e_i} fixes the edge $e_i \subset X$). We extend this action to the *trivial action* on the rest of T . Thus we get an action

$$\prod_{i=1}^n G_{e_i} \curvearrowright T.$$

Observe that there is a tautological epimorphism

$$G \rightarrow \prod_{i=1}^n G_{e_i}$$

which sends each subgroup $G_{e_i} \subset G$ to the subgroup G_{e_i} of the direct product. Hence G acts on T through the quotient group $\prod_{i=1}^n G_{e_i}$.

Let W denote the Euclidean vector space $\text{Vect}(T)$ with the orthonormal basis T , and set $p := \dim(W)$, ie,

$$p = t_1 + \dots + t_n.$$

The set T contains distinguished elements f_1, \dots, f_n consisting of the flags (e_i, F) . The dihedral group D_n acts on F and therefore on $\{f_1, \dots, f_n\}$. We extend this action to the rest of T (and hence to W) by the identity on $T \setminus \{f_1, \dots, f_n\}$. The reflections $R_j \in D$ yield isometric involutions I_j of W . Note

that (since $n = 2k - 1$ is odd) the involution R_{i+k} fixes x_{i+1} , hence I_{i+k} permutes the vectors $f_i, f_{i+1} \in W$.

Embed the polygon F to \mathbb{H}^2 as a right-angled regular polygon $\mu(F) \subset \mathbb{H}^2$ as in the proof of Theorem 1.1. As before, we will identify F and $\mu(F)$. We embed \mathbb{H}^2 into \mathbb{H}^p as a totally-geodesic subspace. Let m_i denote the midpoint of the edge $e_i \subset F$. The tangent space $W_i := T_{m_i}\mathbb{H}^p$ contains a distinguished vector \vec{f}_i which is the unit vector orthogonal to e_i and directed inward F . Note that this vector is parallel (under the parallel transport along e_i) to the vectors $\vec{f}_i^+, \vec{f}_{i+1}^-$ used in the proof of Theorem 1.1. Pick an arbitrary linear isometry $\psi_1: W \rightarrow W_1$ which sends f_1 to \vec{f}_1 . By conjugating via ψ_1 we transport the linear action $G \curvearrowright W$ to a linear action $G \curvearrowright W_1$, exponentiating the latter action we get an isometric action $\rho_1: G \curvearrowright \mathbb{H}^p$.

We now proceed analogously to the proof of Theorem 1.1: Define linear maps $\psi_i: W \rightarrow W_i$ so that we have a commutative diagram:

$$\begin{array}{ccccccc}
 W & \xrightarrow{I_{k+1}} & W & \xrightarrow{I_{k+2}} & W & \dots & \xrightarrow{I_{n+k+1}} & W \\
 \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 & & & \downarrow \psi_{n+1} \\
 W_1 & \xrightarrow{R_{k+1}} & W_2 & \xrightarrow{R_{k+2}} & V_3 & \dots & \xrightarrow{R_{n+k+1}} & W_n
 \end{array}$$

Note that $\psi_i(f_i) = \vec{f}_i$ for all i . Indeed, $\psi_1(f_1) = \vec{f}_1$ by construction. Suppose that $\psi_i(f_i) = \vec{f}_i$. Then

$$\psi_{i+1}(f_{i+1}) = R_j \circ \psi_i \circ I_j(f_{i+1}) = R_j \circ \psi_i(f_i) = R_j(\vec{f}_i) = \vec{f}_{i+1}$$

where $j = i + k$.

Observe that $\psi_{n+1} \neq \psi_1$. However, $I_{n+k+1} \circ \dots \circ I_{k+1} = I_1$ commutes with $G_{e_1} \curvearrowright W$ and $R_{n+k+1} \circ \dots \circ R_{k+1} = R_1$ commutes with $\rho_1(G_{e_1}) \curvearrowright \mathbb{H}^p$. Hence

$$\rho_1 = \rho_{n+1}: G_{e_1} \rightarrow \text{Isom}(\mathbb{H}^p).$$

It remains to verify that for each i the groups $\rho_i(G_{e_i}), \rho_{i+1}(G_{e_{i+1}})$ commute. It is elementary to verify that for all $g \in G_{e_i}, g' \in G_{e_{i+1}}$ the vectors $\psi_i(g(f_i))$ and $\psi_{i+1}(g'(f_{i+1}))$ are mutually orthogonal (after being translated to $T_{x_{i+1}}\mathbb{H}^p$ along e_i, e_{i+1}). The group action $\rho_i(G_{e_i}) \curvearrowright T_{x_{i+1}}\mathbb{H}^p$ permutes the vectors

$$\{\psi_i(g(f_i)), g \in G_{e_i}\}$$

and fixes the orthogonal complement to these vectors; same is true for the action of $G_{e_{i+1}}$ and the vectors

$$\{\psi_{i+1}(g'(f_{i+1})), g' \in G_{e_{i+1}}\}$$

Hence the groups $\rho_i(G_{e_i}), \rho_{i+1}(G_{e_{i+1}})$ commute. Therefore we have constructed a homomorphism $\rho: G \rightarrow \text{Isom}(\mathbb{H}^p)$, $\rho|_{G_{e_i}} = \rho_i|_{G_{e_i}}$.

This action has the same ‘‘orthogonality’’ properties as the homomorphism ρ in the proof of Theorem 1.1, ie, if $i \neq j$ then for all $g \in G_{e_i} \setminus \{1\}, g' \in G_{e_j} \setminus \{1\}$, the hyperbolic planes $\mathbb{H}^2, \rho(g)\mathbb{H}^2$ and $\rho(g')\mathbb{H}^2$ are mutually orthogonal. Thus, the arguments of the second part of the proof of Theorem 1.1 still work and, by applying Theorem 4.1, we conclude that ρ is discrete, faithful, convex-cocompact. \square

Suppose now that X is a (locally finite) right-angled 2-dimensional hyperbolic building whose fundamental chamber F has $n \geq 6$ vertices. Recall that X is uniquely determined by the *thickness* t_i of the edges e_i of F , ie, the number of 2-faces in X containing e_i . Thus every such building is the universal cover of an n -gon \mathcal{P} of finite groups corresponding to a cyclic graph-product. Thickness of the edge e_i is the order of the edge group G_{e_i} in \mathcal{P} .

According to a recent theorem of F. Haglund, [13], all uniform lattices in the building X are commensurable. Hence, as an application of Theorem 6.2, we obtain

Corollary 6.3 *Let H be a group acting discretely, cocompactly and isometrically on X . Then H contains a finite index subgroup which admits a properly discontinuous convex-cocompact action on \mathbb{H}^p for some $p = p(X)$.*

7 Extension of discrete representations

In this section we discuss the following question:

Suppose that $G \curvearrowright \mathbb{H}^n$ is a properly discontinuous isometric action. Is it true that G is isomorphic to a Kleinian group?

Note that the kernel F of the action $G \curvearrowright \mathbb{H}^n$ is necessarily finite, therefore we have a short exact sequence

$$1 \rightarrow F \rightarrow G \rightarrow \bar{G} \rightarrow 1,$$

where \bar{G} is Kleinian. What we are interested in is whether the group G is itself isomorphic to a Kleinian group. Of course, a necessary condition for this is that G is residually finite. Finding a non-residually finite extension G of a Kleinian group \bar{G} is a very difficult task, and presently such extensions are not known. Nevertheless we have:

Theorem 7.1 *Suppose that G is a residually finite group which fits into a short exact sequence*

$$1 \rightarrow F \rightarrow G \rightarrow \bar{G} \rightarrow 1,$$

where \bar{G} admits a discrete and faithful representation $\bar{\rho}$ into $\text{Isom}(\mathbb{H}^n)$. Then G also admits a discrete and faithful representation ρ into $\text{Isom}(\mathbb{H}^m)$ for some m . Moreover, if $\bar{\rho}$ is convex-cocompact (resp. geometrically finite) then ρ can be taken convex-cocompact (resp. geometrically finite).

Proof The proof of this theorem is modeled on the proof of the well-known fact that a finite extension of a residually finite linear group is again linear, but we present it here for the sake of completeness.

We first lift $\bar{\rho}$ to a homomorphism $\bar{\rho}: G \rightarrow \text{Isom}(\mathbb{H}^n)$, so that $\text{Ker}(\bar{\rho}) = F$. Since G is residually finite, there exists a homomorphism

$$\phi: G \rightarrow Q$$

where Q is a finite group, so that $\phi|_F$ is injective. Embed Q in $SO(k)$ for some k . The product group $\text{Isom}(\mathbb{H}^n) \times SO(k)$ embeds in $\text{Isom}(\mathbb{H}^{n+k})$ as the stabilizer of \mathbb{H}^n embedded in \mathbb{H}^{n+k} as a totally-geodesic subspace. Therefore, for $m = n + k$ we get a homomorphism

$$\rho: G \rightarrow \text{Isom}(\mathbb{H}^n) \times SO(k) \subset \text{Isom}(\mathbb{H}^m)$$

given by

$$\rho(g) = (\bar{\rho}(g), \phi(g)).$$

It is clear that $\rho(g)|_{\mathbb{H}^n} = \bar{\rho}(g)$ and therefore ρ is faithful and $\rho(G) \subset \text{Isom}(\mathbb{H}^m)$ is discrete. Moreover, $\Lambda(\rho(G)) = \Lambda(\bar{\rho}(\bar{G}))$. Recall that geometrically finite and convex-cocompact actions can be detected by considering the dynamics of a discrete group on its limit set (see [5]). Therefore, if $\bar{\rho}$ is convex-cocompact (resp. geometrically finite) then ρ is also convex-cocompact (resp. geometrically finite). □

Combining Theorem 7.1 with Theorem 5.1 we get Theorem 1.1.

8 Example of a nonlinear Gromov-hyperbolic group

Theorem 8.1 *There exists an infinite hyperbolic group G such that each representation of G to $GL(m, \mathbf{k})$ factors through a finite group. In particular, G is nonlinear. Here \mathbf{k} is an arbitrary field.*

Proof Let Γ be a uniform lattice in a quaternionic hyperbolic space $\mathbb{H}\mathbb{H}^n$, $n \geq 2$. Since $\mathbb{H}\mathbb{H}^n$ is negatively curved, the group Γ is hyperbolic. Clearly, the group Γ is a nonelementary hyperbolic group; hence Γ admits an infinite proper quotient $\Gamma \rightarrow G$ where G is a hyperbolic group (see [11] or [16]). We first consider the case when \mathbf{k} has zero characteristic. Then without loss of generality we can assume that we are given a linear representation $\rho: G \rightarrow GL(N, \mathbb{R})$. We will show that $\rho(G)$ is finite by using the standard “adelic” trick. The reader can find similar applications of this argument in Margulis’ proof of arithmeticity of higher rank lattices (see [19, 25]), and in Tits’ proof of the *Tits alternative*, [23].

The representation ρ lifts to a linear representation $\tilde{\rho}: \Gamma \rightarrow GL(N, \mathbb{R})$. Let L denote the Zariski closure of $\tilde{\rho}(\Gamma)$ in $GL(N, \mathbb{R})$. Let S denote the solvable radical of L . We first consider the case when $L' := L/S$ is a reductive group with nontrivial noncompact factor H . Then the projection $\Gamma \rightarrow G \rightarrow L \rightarrow H$ has Zariski dense image. Hence, according to Corlette’s Archimedean superrigidity theorem [9], the representation $\Gamma \rightarrow H$ extends to $\text{Isom}(\mathbb{H}\mathbb{H}^n)$. This however contradicts the assumption that the projection $\Gamma \rightarrow G$ is not 1–1. Therefore the group L' is a compact algebraic group.

Suppose that the projection $\rho(G) \subset L'$ is infinite. As a compact Lie group, L' is isomorphic to a subgroup of $O(M)$. Since G satisfies property (T),

$$H^1(G, o(M)_{\text{Ad}(\rho)}) = 0,$$

where $o(M)$ is the Lie algebra of $O(M)$. Vanishing of the above cohomology group implies that the space $\text{Hom}(G, O(M))/O(M)$ is finite. Hence, analogously to the proof of Theorem 7.67 in [21], ρ is conjugate to a representation $\rho': G \rightarrow O(M)$ for which

$$\rho'(G) \subset K(F) \subset O(M, F) \subset GL(M, F),$$

where F is a number field and $K(\mathbb{R})$ is the Zariski closure of $\rho'(G)$.

One would like to replace the representation ρ with another representation ϕ of the group G , whose image is Zariski dense in a certain noncompact algebraic group and so that $\text{Ker}(\phi) = \text{Ker}(\rho)$. The most obvious thing to try is to find an element σ of the Galois group $\text{Gal}(\mathbb{C}/\mathbb{Q})$, so that the image of $\phi = \sigma(\rho)$ is not relatively compact. This does not necessarily work. Note however, that the restriction of the norm on \mathbb{C} to $\sigma(F)$ gives rise to an *Archimedean valuation* on F . The idea of the *adelic trick* is to use *non-Archimedean valuations* v of F together with Archimedean ones. This is done by introducing the *ring of adèles* of F , which is a certain subset of the product

$$\prod_{v \in \text{Val}(F)} F_v,$$

where F_v is the completion of F with respect to the valuation v .

Let $\mathbb{A}(F)$ denote the ring of adèles of F ; then the diagonal embedding $F \hookrightarrow \mathbb{A}(F)$ has *discrete image* (see for example [18]). Hence the diagonal embedding

$$\rho'(G) \hookrightarrow GL(M, \mathbb{A}(F))$$

also has discrete image. If the projection of $\rho'(G)$ to each factor $GL(M, F_v)$ were relatively compact, the image of $\rho'(G)$ in $GL(M, \mathbb{A}(F))$ would be compact as well. However a discrete subset of a compact is finite, which contradicts the assumption that $\rho(G)$ is infinite.

Thus there exists a valuation v of F so that the image of the projection

$$\rho'(G) \rightarrow K(F_v) \subset GL(M, F_v)$$

is not relatively compact. In case when v is an Archimedean valuation, we can again apply Corlette's Archimedean superrigidity [9] to get a contradiction. Hence such v has to be nonarchimedean. Therefore the representation $\Gamma \rightarrow \rho'(G) \rightarrow GL(M, F_v)$ corresponds to an isometric action of Γ on a locally finite Euclidean building X . However, by the non-Archimedean superrigidity theorem of Gromov and Schoen [12], Γ fixes a point in X . Therefore the image of $\rho'(G)$ in $GL(M, F_v)$ is relatively compact, which is a contradiction. Hence $\rho'(G)$ is finite. It follows that the group L is commensurable to its solvable radical S ; hence $\rho(G)$ is a virtually solvable group. By applying property (T) again, we conclude that $\rho(G)$ is finite.

We now consider the case when \mathbf{k} has positive characteristic; since the argument is similar to the zero characteristic case, we give only a sketch. Under the above assumptions, $\rho(G) \subset GL(m, F)$, where F is a finitely generated field (of positive characteristic). The field F is an extension

$$F_q \subset E \subset F$$

where F_q is a finite field, $F_q \subset E$ is an purely transcendental extension and $E \subset F$ is an algebraic extension (see [15, Chapter VI.1]). Since F is finitely generated, F/E is finite-dimensional and therefore, by passing to a bigger matrix group, we reduce the problem to the case when $F = E$ is a purely transcendental extension, which necessarily has finite transcendence degree. Therefore we reduced to the case of $\rho: G \rightarrow GL(N, F)$, where $F = F_q(t_1, \dots, t_m)$ is the field of rational functions with coefficients in F_q .

Then we associate with each variable $t_j^{\pm 1}$ a discrete valuation $v_{\pm j}$ and an action $G \curvearrowright X_{\pm j}$ on the corresponding Euclidean building. The non-Archimedean superrigidity theorem of Gromov and Schoen [12] shows that for each $\pm j$ the

action $G \curvearrowright X_{\pm j}$ has a fixed point. Therefore the matrix coefficients of $\rho(G)$ have bounded degree with respect to all the variables $t_1^{\pm 1}, \dots, t_m^{\pm 1}$. Hence, since F_q is finite, the matrix coefficients of $\rho(G)$ belongs to a finite subset of F and thus $\rho(G)$ is finite. \square

Remark 8.2 After this paper was written, I was informed by Alain Valette that he also knew how to prove Theorem 8.1. I am sure that other people were also aware of this proof since all the arguments here are quite standard.

References

- [1] **W Ballmann, J Świątkowski**, *On L^2 -cohomology and property (T) for automorphism groups of polyhedral cell complexes*, *Geom. Funct. Anal.* 7 (1997) 615–645
- [2] **A F Beardon**, *The geometry of discrete groups*, Graduate Texts in Mathematics 91, Springer–Verlag, New York (1983)
- [3] **M Bonk, O Schramm**, *Embeddings of Gromov hyperbolic spaces*, *Geom. Funct. Anal.* 10 (2000) 266–306
- [4] **M Bourdon**, *Sur la dimension de Hausdorff de l'ensemble limite d'une famille de sous-groupes convexes co-compacts*, *C. R. Acad. Sci. Paris Sér. I Math.* 325 (1997) 1097–1100
- [5] **B H Bowditch**, *Geometrical finiteness for hyperbolic groups*, *J. Funct. Anal.* 113 (1993) 245–317
- [6] **B H Bowditch**, *Geometrical finiteness with variable negative curvature*, *Duke Math. J.* 77 (1995) 229–274
- [7] **M R Bridson, A Haefliger**, *Metric spaces of non-positive curvature*, Grundlehren series 319, Springer–Verlag, Berlin (1999)
- [8] **J A Carlson, D Toledo**, *Harmonic mappings of Kähler manifolds to locally symmetric spaces*, *Inst. Hautes Études Sci. Publ. Math.* (1989) 173–201
- [9] **K Corlette**, *Archimedean superrigidity and hyperbolic geometry*, *Ann. of Math.* (2) 135 (1992) 165–182
- [10] **A Felikson, P Tumarkin**, *A series of word-hyperbolic Coxeter groups*, [arXiv:math.GR/0507389](https://arxiv.org/abs/math/0507389)
- [11] **M Gromov**, *Hyperbolic groups*, from: “Essays in group theory”, *Math. Sci. Res. Inst. Publ.* 8, Springer, New York (1987) 75–263
- [12] **M Gromov, R Schoen**, *Harmonic maps into singular spaces and p -adic superrigidity for lattices in groups of rank one*, *Inst. Hautes Études Sci. Publ. Math.* (1992) 165–246

- [13] **F Haglund**, *Commensurabilité de réseaux uniformes dans les immeubles à angles droits*, preprint (2003)
- [14] **F Haglund**, **D Wise**, in preparation
- [15] **T W Hungerford**, *Algebra*, Graduate Texts in Mathematics 73, Springer-Verlag, New York (1980)
- [16] **S V Ivanov**, **A Y Ol'shanskiĭ**, *Hyperbolic groups and their quotients of bounded exponents*, Trans. Amer. Math. Soc. 348 (1996) 2091–2138
- [17] **M Kapovich**, *Hyperbolic manifolds and discrete groups*, Progress in Mathematics 183, Birkhäuser Boston Inc. Boston, MA (2001)
- [18] **S Lang**, *Algebraic numbers*, Addison-Wesley Publishing Co. Inc. Reading, Mass.–Palo Alto–London (1964)
- [19] **G A Margulis**, *Discrete subgroups of semisimple Lie groups*, Ergebnisse series 17, Springer-Verlag, Berlin (1991)
- [20] **G D Mostow**, **Y T Siu**, *A compact Kähler surface of negative curvature not covered by the ball*, Ann. of Math. (2) 112 (1980) 321–360
- [21] **M S Raghunathan**, *Discrete subgroups of Lie groups*, Springer-Verlag, New York (1972)
- [22] **J Świątkowski**, *A class of automorphism groups of polygonal complexes*, Q. J. Math. 52 (2001) 231–247
- [23] **J Tits**, *Free subgroups in linear groups*, J. Algebra 20 (1972) 250–270
- [24] **D T Wise**, *The residual finiteness of negatively curved polygons of finite groups*, Invent. Math. 149 (2002) 579–617
- [25] **R J Zimmer**, *Ergodic theory and semisimple groups*, Monographs in Mathematics 81, Birkhäuser Verlag, Basel (1984)