

Homogeneous coordinate rings and mirror symmetry for toric varieties

MOHAMMED ABOUZAID

Given a smooth toric variety X and an ample line bundle $\mathcal{O}(1)$, we construct a sequence of Lagrangian submanifolds of $(\mathbb{C}^*)^n$ with boundary on a level set of the Landau–Ginzburg mirror of X . The corresponding Floer homology groups form a graded algebra under the cup product which is canonically isomorphic to the homogeneous coordinate ring of X .

14J32; 53D40

1 Introduction

In this paper we give some evidence for M Kontsevich’s homological mirror symmetry conjecture [13] in the context of toric varieties. Recall that a smooth complete toric variety is given by a simplicial rational polyhedral fan Δ such that $|\Delta| = \mathbb{R}^n$ and all maximal cones are non-singular (Fulton [10, Section 2.1]). The convex hull of the primitive vertices of the 1–cones of Δ is a convex polytope which we denote by P , containing the origin as an interior point, and may be thought of as the Newton polytope of a Laurent polynomial $W: (\mathbb{C}^*)^n \rightarrow \mathbb{C}$. This Laurent polynomial is the Landau–Ginzburg mirror of X .

Our construction will rely on choosing an ample line bundle $\mathcal{O}(1)$ on X . On the complex side, we know that sections of this line bundle are given by lattice points of a polytope Q determined by $\mathcal{O}(1)$. Further, we can recover the original toric variety as

$$(1-1) \quad X = \text{Proj} \left(\bigoplus_{j \geq 0} \mathbb{C}^j Q \cap \mathbb{Z}^n \right),$$

with product given by linearly extending the formula

$$x_k \otimes y_l \rightarrow x_k + y_l.$$

For the bulk of the paper, it will be more convenient to identify the integral points of jQ with the $\frac{1}{j}$ integral points of Q , and re-write the product as

$$(1-2) \quad x_k \otimes y_l \rightarrow \frac{kx_k + ly_l}{k+l}.$$

On the symplectic side, we study the Floer homology of Lagrangian submanifolds of $(\mathbb{C}^*)^n$ with boundary on $W^{-1}(0)$. $\mathcal{O}(1)$ will induce a subdivision of P , and hence a tropical degeneration of $W^{-1}(0)$ in the sense of Mikhalkin. In order to simplify our computations, we will have to replace $\mathcal{O}(1)$ by a sufficiently high power (see Remark 3.10). The combinatorial data which determines $\mathcal{O}(1)$ will then allow us to construct Lagrangians which we suggestively call $\mathcal{L}(j)$ for every integer j . We will write \mathcal{L} for $\mathcal{L}(0)$. Our main result is the following:

Theorem 1.1 *The Floer cohomology groups $HF^0(\mathcal{L}, \mathcal{L}(j))$ for $j \geq 0$ form an associative algebra under the cup product which is canonically isomorphic to the homogeneous coordinate ring of X equipped with the line bundle $\mathcal{O}(1)$.*

We now summarize the contents of this paper. In Section 2, we will introduce the notion of admissible Lagrangian (originally due to Kontsevich), and explain why their Floer theory is well defined. In Section 3, we will review the results of Mikhalkin [15] which we will need, and explain how the datum of a line bundle on a toric variety gives rise to a tropical degeneration of its mirror. Ignoring boundary problems, we will also introduce some flat Lagrangian submanifolds of $T^*\mathbb{R}^n/\mathbb{Z}^n$ whose Floer cohomology groups correspond to holomorphic sections of the line bundles $\mathcal{O}(j)$. In Section 4, we will give an explicit construction of a smooth symplectic submanifold which interpolates between the complex hypersurface and its “tropical” counterpart, as well as a construction of the Lagrangian submanifold \mathcal{L} which corresponds to the structure sheaf of X . In Section 5, we will use the Lagrangians introduced in Section 3 to construct the admissible Lagrangians $\mathcal{L}(j)$ that appear in the main theorem, and complete its proof. There is a minor change of notation which occurs in the middle of the paper as explained in Remark 4.18.

We would like to comment on some related work. We do not discuss any homological algebra in this paper, but the result we prove establishes the existence of a functor from the Fukaya category of the mirror of X_Δ to the category of coherent sheaves on X_Δ . Let us briefly sketch the construction. The functor takes every Lagrangian to the graded vector space $\bigoplus_{j \geq 0} HF^0(\Lambda, \mathcal{L}(j))$. By Theorem 1.1 this is naturally a graded module over the homogeneous coordinate ring of X . A classical result of Serre implies that the category of coherent sheaves is a quotient of the category of graded modules.

In order to prove that this functor yields an equivalence of derived categories, one would need to understand how the A_∞ structures on both sides affect the construction. For $\mathbb{C}\mathbb{P}^2$, this can be done explicitly using essentially the methods of this paper and Beilinson's description of the category of coherent sheaves on projective spaces [1], and we expect the computation to extend to $\mathbb{C}\mathbb{P}^n$. However, for a general toric variety, it is not clear how to obtain such an explicit description of the category of coherent sheaves. In an upcoming paper, we will establish such an equivalence of categories, but the methods are unfortunately much less explicit than those used here, and pass through Morse theory. Also, if we consider the Lagrangians $\partial L(j) \subset M$, we can use the ideas of this paper to give evidence for the mirror conjecture, this time in the case of Calabi–Yau hypersurfaces in toric varieties.

Acknowledgments

I would first like to thank my advisor Paul Seidel for originally suggesting the use of tropical geometry as an approach to Kontsevich's conjecture, and for helpful discussions throughout my work on this project. I also thank Gabriel Kerr for teaching me many things about toric varieties, Kevin Costello for useful comments on an early draft, and Ivan Smith for an important conversation about this work while it was still in maturation. I finally want to acknowledge the referee whose careful reading of an earlier version unearthed many errors, not all of them minor, and whose suggestions for addressing the more significant issues have been particularly helpful.

2 Floer (co)-homology for Lagrangians with boundary along a complex hypersurface

2.1 Admissible Lagrangians

We begin by observing that many properties of closed Lagrangian manifolds can be extended to the case where the Lagrangians have boundary. For example, the cotangent bundle of any manifold with boundary is itself a manifold with boundary which carries the usual symplectic structure.

Recall that the restriction of the cotangent bundle to the boundary carries a canonical oriented rank–1 trivial sub-bundle $E_{\partial L}$ which annihilates the tangent space to the boundary and induces the appropriate co-orientation on ∂L . Note that we can think of this sub-bundle as lying in the restriction of the tangent space of T^*L to the boundary of the zero section. The proof of Weinstein's neighbourhood theorem extends to this setting to show that Lagrangian submanifolds with boundary, with a choice of a rank–1

oriented sub-bundle at the boundary, are locally modeled after the cotangent bundle with its canonical sub-bundle at the boundary.

Lemma 2.1 *Let L be a Lagrangian submanifold of a symplectic manifold N , and let E be an oriented rank-1 sub-bundle of the symplectic orthogonal complement of $T\partial L$ such that the pairing*

$$TL|_{\partial L} \otimes E \rightarrow \mathbb{R}$$

induced by the symplectic form is non-degenerate and yields the appropriate co-orientation on ∂L .

Inside a sufficiently small neighbourhood of L in N , there exists a full dimensional submanifold with boundary $(V_L, \partial V_L)$, such that the inclusion $(L, \partial L) \subset (V_L, \partial V_L)$ satisfies the following properties:

- *The restriction of $T\partial V_L$ to ∂L contains the sub-bundle E .*
- *There exists a symplectomorphism $(V_L, \partial V_L) \rightarrow (T^*L, T^*L|_{\partial L})$ identifying L with the zero section of its cotangent bundle and E with the canonical sub-bundle $E_{\partial L}$ and a projection to $(L, \partial L)$ such that the following diagram commutes:*

$$\begin{array}{ccc} (V_L, \partial V_L) & \xrightarrow{\quad\quad\quad} & (T^*L, T^*L|_{\partial L}) \\ & \searrow & \swarrow \\ & (L, \partial L) & \end{array}$$

□

As usual, this allows us to reduce problems about the topology of nearby Lagrangian submanifolds to questions about closed forms on L . We will find the following lemma particularly useful:

Lemma 2.2 *Let L' and L be two Lagrangian submanifolds of N which have the same boundary. Let V_L be a submanifold of N which satisfies the conditions of Lemma 2.1. If L' is transverse to ∂V_L and there is a neighbourhood of $\partial L'$ in L' which is contained in V_L , then there exists a Lagrangian submanifold L'' which satisfies the following conditions:*

- *L'' is Hamiltonian isotopic to L .*
- *L'' agrees with L' in a sufficiently small neighbourhood of $\partial L'$.*
- *L'' agrees with L away from a larger neighbourhood of the boundary.*

Moreover, L'' is independent, up to Hamiltonian isotopy, of the choices which will be made in its construction.

Proof Consider a point p in $\partial L'$, and let \vec{n} be a tangent vector in TL'_p which points towards the interior. Since L' and ∂V_L are transverse, \vec{n} projects to a vector in TL_p which is transverse to the boundary. Since there is a neighbourhood of $\partial L'$ in L' which is contained in V_L , the image of \vec{n} must point towards the interior of L . Therefore, the restriction of the projection $V_L \rightarrow L$ to L' is a submersion in a neighbourhood of the boundary. In particular, passing to the cotangent bundle of L , we can identify L' locally as the graph of a closed 1-form which vanishes on ∂L . Since the inclusion of ∂L in a neighbourhood induces an isomorphism on cohomology this 1-form is exact, so we may write it as the differential of a function H' .

By choosing a cutoff function with appropriately bounded derivatives, we can construct a function $H'': L \rightarrow \mathbb{R}$ with support in a neighbourhood of ∂L which agrees with H' in a smaller neighbourhood of the boundary, and such that the graph of dH'' lies in a neighbourhood of L which is identified with V_L . Linear interpolation yields an isotopy between any two choices for H'' . Pulling back the graph of the exact 1-form dH'' to V_L yields the desired Lagrangian L'' . \square

In the situation considered in this paper, N will be a Stein manifold and all the Lagrangians we will consider will have their boundary lying on M , the zero level set of a holomorphic map $f: N \rightarrow \mathbb{C}$ with 0 as a regular value. We can equip this Stein manifold with the structure of an exact symplectic manifold by choosing an embedding into \mathbb{C}^R and restricting the usual symplectic form on \mathbb{C}^R and its primitive. We will denote the symplectic form by ω , and the primitive by θ . As usual, the complex and symplectic structures induce a metric g . Given a Lagrangian L , the existence of the complex structure determines ∂V_L to first order since we can let the line bundle E be spanned by $J\vec{n}$, with \vec{n} the normal vector of the inclusion $\partial L \subset L$.

Note that θ restricts to a closed 1-form on every Lagrangian submanifold. The following definition is standard.

Definition 2.3 A Lagrangian submanifold L is *exact* if there is a function h on L such that $dh = \theta|_L$.

In essence, this exactness conditions provides a priori bounds for the energy of pseudo-holomorphic discs. Since our Lagrangian submanifolds may have boundary, we will require an additional condition as a mean to guarantee some compactness results for such discs (see Lemma 2.8). Recall that the symplectic orthogonal complement to the tangent space of the fibre defines a distribution on N . Every tangent vector in \mathbb{C} has a unique lift to this orthogonal complement, so we may associate a connection to the map f . Given a Lagrangian in the symplectic hypersurface $f^{-1}(0)$ and a curve γ




Figure 1: The pair (L_1, L_2) on the left is a positively oriented pair. The one on the right picture is negatively oriented.

in \mathbb{C} with an endpoint at the origin, parallel transport with respect to this connection determines a unique Lagrangian submanifold of N whose image under f is exactly γ .

Definition 2.4 A compact oriented exact Lagrangian submanifold $(L, \partial L)$ of N which has boundary on M is *admissible* if there exists a curve γ in \mathbb{C} such that $\gamma(0) = 0$ and L agrees with the parallel transport of ∂L along γ in some neighbourhood of the origin.

Note that the condition of admissibility is of course vacuous if $\partial L = \emptyset$, which is a possibility that we do not exclude. As far as the author knows, the idea of studying such Lagrangians is due to Kontsevich [12, page 30].

As our goal is to define and compute Floer homology for admissible Lagrangians, we will have to understand the possible behaviours at the boundary. If L is admissible, then there exists a non-zero vector $\gamma'(0) = v \in \mathbb{R}^2$ such that the image of a transverse vector to $T\partial L$ in $TL|_{\partial L}$ lies in the ray \mathbb{R}^+v .

Definition 2.5 A pair of admissible Lagrangians (L_1, L_2) , whose tangent spaces at the boundary project to vectors (v_1, v_2) , is *positively oriented* if the angle from v_1 to v_2 lies in the interval $(0, \frac{\pi}{2})$. If this angle is between $-\frac{\pi}{2}$ and 0 , we say that the pair is *negatively oriented*.

Assume the pair (L_1, L_2) is negatively oriented. We may choose a vector v'_2 such that (v_1, v'_2) is a positively oriented pair and the angle between v_2 and v'_2 is less than $\frac{\pi}{2}$. The latter condition is enough to guarantee that the parallel transport of ∂L_2 along a the straight half ray γ'_2 with tangent vector v'_2 at the origin lies in a neighbourhood of L_2 in which we can apply Lemma 2.2 to obtain a Lagrangian submanifold L''_2 which interpolates between L_2 and the parallel transport of its boundary along γ'_2 .

This process takes the configuration on the right in Figure 1 to the one on the left. We conclude the following:

Lemma 2.6 *If the pair (L_1, L_2) is negatively oriented and $H^1(L_2, \partial L_2) = 0$, then L_2 is Hamiltonian isotopic to an admissible Lagrangian L_2'' such that the pair (L_1, L_2'') is positively oriented. Furthermore, L_2'' is independent of the choices that are made in the construction up to a Hamiltonian isotopy which preserves the admissibility and positivity of the pair (L_1, L_2'') .*

Proof The only parts which we have not checked are the exactness and uniqueness of L_2'' . But the restriction of θ to L_2 , and therefore to ∂L_2 is exact. Since L_2 and L_2'' are homeomorphic, the vanishing of $H^1(L_2, \partial L_2)$ guarantees that $\theta|_{L_2''}$ is also exact.

We already proved uniqueness in Lemma 2.2 by using a linear isotopy. But such an isotopy does not necessarily preserve admissibility. However, any two candidates $L_{2,0}''$ and $L_{2,1}''$ can be made the endpoints of a 1-parameter family of admissible Lagrangians $L_{2,t}''$ such that $(L_1, L_{2,t}'')$ is a positively oriented pair for every time t . Since the submanifolds $L_{2,t}''$ agree away from a tubular neighbourhood of the boundary, this 1-parameter family is generated by a time dependent 1-form which vanishes away from a neighbourhood of the boundary. On the other hand, it also vanishes at the boundary since all Lagrangians $L_{2,t}''$ have the same boundary. Therefore the 1-form is exact, and the 1-parameter family $L_{2,t}''$ is generated by a time-dependent Hamiltonian function. \square

2.2 Compactness for pseudo-holomorphic discs

We will define a Floer theory for admissible Lagrangians in which the possible boundary intersection points are ignored. To this effect, let L_1 and L_2 be admissible Lagrangians which intersect transversely away from M , and choose p and q , a pair of transverse intersection points between these two Lagrangians. In analogy with Morse theory, where we count gradient trajectories connecting critical points, we will count pseudo-holomorphic maps connecting p and q .

Formally, we equip the strip

$$S = \mathbb{R} \times [0, 1] = \{t, s \mid -\infty < t < +\infty, 0 \leq s \leq 1\}$$

with the usual complex structure

$$J_S \frac{\partial}{\partial t} = \frac{\partial}{\partial s} \quad J_S \frac{\partial}{\partial s} = -\frac{\partial}{\partial t}.$$

Just as Morse theory relies on choosing a function whose gradient flow with respect to an underlying metric satisfies appropriate transversality conditions, we will have to perturb the complex structure to an almost complex structure in order to achieve transversality, (these almost complex structures are called regular, Floer–Hofer–Salamon [7]). Perturb the complex structure on N to such a regular almost complex structure J while maintaining the condition that the map $f: N \rightarrow \mathbb{C}$ remains holomorphic in a neighbourhood of M . In the closed case, [7, Remark 5.2] proves that regularity may be achieved by perturbing the almost complex structure on any open subset where every pseudo-holomorphic curve passes. The case of curves with Lagrangian boundary conditions is entirely analogous, and we can therefore choose a perturbation which is supported in a small neighbourhood of p and q , and hence occurs far away from M .

Even though the perturbed almost complex structure on N is not integrable, we may still consider holomorphic maps

$$u: S \rightarrow N,$$

such that $u(t, 0) \in L_1$, $u(t, 1) \in L_2$. We say that such a map has finite energy if the integral

$$\int_S u^*(\omega)$$

is finite.

The classical theory studies $\mathcal{M}(p, q)$, the moduli space of finite energy maps such that for every $s \in [0, 1]$,

$$\begin{aligned} \lim_{t \rightarrow +\infty} u(t, s) &= p \\ \lim_{t \rightarrow -\infty} u(t, s) &= q. \end{aligned}$$

We will need to add an additional restriction.

Definition 2.7 Let L_1 and L_2 be admissible Lagrangians with transverse intersection points p and q which occur away from the boundary. An *admissible strip* u is an element of $\mathcal{M}(p, q)$ whose image does not intersect M .

Note that $\mathcal{M}(p, q)$ admits a free \mathbb{R} action corresponding to translation in the t direction, and the regularity of the almost complex structure implies that the quotient is a smooth manifold. However, unlike the case for closed Lagrangians, $\mathcal{M}(p, q)/\mathbb{R}$ does not admit a compactification to a manifold with boundary because of those pseudo-holomorphic strips whose boundary intersects the boundaries of L_1 or L_2 . However, for admissible strips, we have the following:




Figure 2: The image of $\partial S'$ is contained in the thick lines.

Lemma 2.8 *Let L_1 and L_2 be admissible Lagrangians, and let p and q be transverse intersection points. If u_τ is a 1-parameter family of holomorphic strips in $\mathcal{M}(p, q)$ such that u_0 is admissible, then there exists an $\epsilon > 0$ such that the image of $f \circ u_\tau$ does not intersect the closed ϵ neighbourhood of the origin for all $0 \leq \tau \leq 1$. In particular u_1 is admissible.*

Proof The reader may find Figure 2 useful in what follows.

Let ϵ be such that $f(L_1)$ and $f(L_2)$ do not intersect in the punctured 2ϵ neighbourhood of the origin. Assume that τ is the smallest time at which the lemma does not hold. Consider $(u_\tau \circ f)^{-1}(B_{\epsilon'}(0)) = S'$ for some ϵ' between ϵ and 2ϵ . Since the critical points of holomorphic maps are isolated, we may choose ϵ' such that S' is a submanifold of S with boundary. Notice that the boundary of S' must be mapped to the union of the set $f(L_1) \cup f(L_2)$ with the circle of radius ϵ' .

We can compute the degree of $u_\tau \circ f$ by choosing a generic point p in the image of S' , and counting the number of preimages. However, one may find a path from p to the interior of $B_\epsilon(0)$ which does not pass through the image of $\partial S'$ or through any critical point. Since the number of preimages can only change at a critical point or on the boundary, and the image of S' does not intersect the interior of the ball of radius ϵ , we conclude that p has no preimages. But this proves that $u(S)$ does not intersect the ball of radius ϵ . □

Note that the inclusion,

$$\mathcal{M}^{\text{adm}}(p, q) \subset \mathcal{M}(p, q)$$

is clearly open. The above lemma shows that it is also closed, so $\mathcal{M}^{\text{adm}}(p, q)$ consists of components of the moduli space of all finite energy strips connecting p and

q . Moreover, the lemma also guarantees that the images of all admissible pseudo-holomorphic strips lie in a compact subset of N that does not intersect the boundary of the Lagrangians L_i . In particular we can conclude that the closure of $\mathcal{M}^{\text{adm}}(p, q)/\mathbb{R}$ in the Gromov compactification $\overline{\mathcal{M}(p, q)}/\mathbb{R}$ is a compact manifold with corners as long as we can prove the existence of a bound on the energy of pseudo-holomorphic strips. This is where the exactness conditions are used. Indeed, since ω is exact, we have

$$\int_S u^*(\omega) = \int_{\partial S} \theta$$

by Stokes's theorem. But ∂S consists of two segments, one on L_1 and the other on L_2 , with endpoints at p and q . Since the restriction of θ to L_i is exact, the right hand-side is independent of the paths between p and q . This implies that all pseudo-holomorphic strips in $\mathcal{M}(p, q)$ have the same energy.

2.3 Floer homology

Having proved the necessary compactness result, we can now define relatively graded Floer homology groups over a field of characteristic 2 for admissible Lagrangians. We will follow the construction of these groups for closed Lagrangians which is due to Floer, [6]. We assume (L_1, L_2) is a positively oriented pair of Lagrangians which intersect transversely away from the boundary, and define a chain complex

$$CF_*(L_1, L_2) = \bigoplus_{p \in (L_1 \cap L_2) - M} \mathbb{Z}_2 \cdot [p]$$

with differential

$$d[p] = \sum_{[q]} |\mathcal{M}^{\text{adm}}(p, q)/\mathbb{R}| \cdot [q]$$

where the sum is taken over all points q such that $\mathcal{M}^{\text{adm}}(p, q)$ is 1-dimensional, and $|\mathcal{M}^{\text{adm}}(p, q)/\mathbb{R}|$ is the cardinality of the space of unparametrized strips connecting p to q . The following result is classical.

Lemma 2.9 *If L_1 and L_2 are exact, then $d^2 = 0$.*

Proof The proof that $d^2 = 0$ relies on interpreting the terms in the expression for $d^2[p]$ as the boundaries of 1-dimensional moduli spaces of unparametrized strips. So we must show the boundary of such moduli spaces consists only of pairs of holomorphic strips, ie, that no bubbling of pseudo-holomorphic discs occurs. But given a pseudo-holomorphic disc u with boundary on one of the Lagrangians L_i we can use Stokes's

theorem to compute that

$$0 \neq \int_D u^*(\omega) = \int_{S^1} u^*(\theta).$$

This contradicts the assumption that the restriction of θ to L_i is exact. \square

This construction produces a \mathbb{Z}_2 graded theory which cannot be lifted, in general, to the usual \mathbb{Z} grading which we expect in a homology theory. However, if N admits a complex volume form (ie, if $c_1(N) = 0$), then there is a special class of Lagrangians for which such a theory exists, see Seidel [18]. Choosing a complex volume form Ω , we define a phase map

$$\begin{aligned} L &\rightarrow S^1 \\ p &\mapsto \frac{\Omega(e_1 \wedge \cdots \wedge e_n)}{|\Omega(e_1 \wedge \cdots \wedge e_n)|} \end{aligned}$$

where $\{e_i\}_{i=1}^n$ is an oriented frame for the tangent space of L at p .

Definition 2.10 A *graded Lagrangian* submanifold of N is an oriented Lagrangian submanifold of N together with a lift of its phase map from S^1 to \mathbb{R} .

Note, in particular, that the obstruction to the existence of a lift lies in $H^1(L)$, so that all simply connected Lagrangians are gradable. We refer to [18] for details on the construction of \mathbb{Z} -graded Floer homology. We also need to lift these homology groups to \mathbb{C} -valued invariants in order to compare them to cohomology groups of line bundles on the mirror. The need for appropriate orientations accounts for the additional restrictions in

Definition 2.11 An *admissible Lagrangian brane* is an admissible graded Lagrangian L which is spin, together with a choice of a spin structure.

Remark 2.12 The term brane is borrowed from string theory, where the ‘‘Lagrangian branes’’ that we’re considering can be thought of as boundary conditions for open strings in the A -model. Other than the fact that string theory motivates the homological mirror symmetry conjecture, physical considerations are completely irrelevant to our arguments.

The relevance of spin structures to the orientation of moduli spaces of discs was observed by de Silva in [2] and by Fukaya, Oh, Ohta, and Ono in [9]. We will give a short description of the signed differential. The reader should keep Figure 3 in mind during the next few paragraphs.




Figure 3: Gluing H_q to S

Consider an intersection point p between L_1 and L_2 . The tangent spaces $T_p L_1$ and $T_p L_2$ are linear Lagrangians in a symplectic vector space V . The grading determines a unique path up to homotopy λ_p from $T_p L_1$ to $T_p L_2$. We may therefore define a Cauchy–Riemann operator $\bar{\partial}_{H_p}$ on a copy of the upper half-plane H_p with Lagrangian boundary conditions, given by the path λ_p , that converge to L_1 along the negative real axis and to L_2 along the positive real axis.

More precisely, choose a map $h: \mathbb{R} \rightarrow [0, 1]$ which takes a neighbourhood of $-\infty$ to 0 and of ∞ to 1, and consider maps from H_p to the vector space V that have values in the Lagrangian $\lambda_p(h(t))$ subspace at a point t of the boundary. We denote this space by $C_{\lambda_p}^\infty(H_p, V)$. We can now define the operator

$$\bar{\partial}_{H_p}: C_{\lambda_p}^\infty(H_p, V) \rightarrow C^\infty(H_p, V \otimes \Omega^{0,1}(H_p))$$

to be the usual $\bar{\partial}$ operator. We can perform the same construction at any other intersection point q to produce an operator $\bar{\partial}_{H_q}$ with boundary conditions λ_q .

Let S be an element of $\mathcal{M}^{\text{adm}}(p, q)$. Since the upper half-plane is biholomorphic to a 1-sided strip, we may glue H_q to the negative end of S to yield a surface $S\#H_q$ which is again bi-holomorphic to a one-sided strip. Strictly speaking, this means that we should choose an identification of a neighbourhood of infinity in H_q with $(-\infty, 0] \times [0, 1]$. For any $R > 0$ we can remove $(-\infty, -R) \times [0, 1]$ from S and from H_q , then identify the two copies of $[-R, 0] \times [0, 1]$ in S and in H_q using the biholomorphism

$$(t, s) \mapsto (-R - t, 1 - s).$$

Further, for R sufficiently large, since the asymptotic boundary conditions agree, we can glue the Cauchy–Riemann operators $\bar{\partial}_S$ and $\bar{\partial}_{H_q}$ to get a Cauchy–Riemann operator $\bar{\partial}_{S\#\bar{\partial}_{H_q}}$ on the strip $S\#H_q$. We do not keep track of R in the notation because the construction is essentially independent of R if it is sufficiently large.

To make sense of the boundary conditions, we trivialize the restriction of TN to S , so that the boundary conditions $u(t, 0) \in L_1$ and $u(t, 1) \in L_2$ yield two paths

$$\begin{aligned} \lambda_1: \mathbb{R} &\rightarrow V \text{ such that } \lambda_1(-\infty) = T_q L_1 \text{ and } \lambda_1(+\infty) = T_p L_1 \\ \lambda_2: \mathbb{R} &\rightarrow V \text{ such that } \lambda_2(-\infty) = T_q L_2 \text{ and } \lambda_2(+\infty) = T_p L_2. \end{aligned}$$

The boundary condition $\lambda_{S\#H_q}$ for $\bar{\partial}_{S\#\bar{\partial}_{H_q}}$ is just given by concatenating the paths $-\lambda_1$, λ_q , and λ_2 , where $-\lambda_1$ corresponds to traversing the path “backwards”. In particular, we have the asymptotic conditions

$$\lambda_{S\#H_q}(-\infty) = T_p L_1 \text{ and } \lambda_{S\#H_q}(+\infty) = T_p L_2$$

which are the same conditions as those of λ_p . Since the space of graded Lagrangians is simply connected, the paths λ_p and $\lambda_{S\#H_q}$ are homotopic. Moreover, the choice of spin structures determines, up to homotopy, a unique such homotopy between λ_p and $\lambda_{S\#H_q}$. If we consider the extensions of our Cauchy–Riemann operators to appropriate Hilbert space completions of spaces of smooth functions, the above path will therefore yield an isomorphism

$$\det(\bar{\partial}_{S\#H_q}) \cong \det(\bar{\partial}_{H_p}).$$

On the other hand, the gluing theorem yields the following isomorphism of determinant bundles

$$\det(\bar{\partial}_{S\#H_q}) \cong \det(\bar{\partial}_S) \otimes \det(\bar{\partial}_{H_q}).$$

If S is a strip with 1–dimensional parametrized moduli space, then $\det(\bar{\partial}_S)$ is canonically trivialized by the translation operator $\frac{\partial}{\partial t}$. We therefore obtain an isomorphism

$$\det(\bar{\partial}_{H_p}) \cong \det(\bar{\partial}_{H_q}).$$

The orientations of L_1 and L_2 determine orientations of these two vector spaces. The contribution of S to $d[p]$ will be positive if the above isomorphism preserves orientations, and negative otherwise.

2.4 Stability of Floer homology under perturbations

While we are interested in the symplectic topology of a complex hypersurface $M \subset (\mathbb{C}^*)^n$, our constructions will rely on deforming M to symplectic submanifolds of $(\mathbb{C}^*)^n$ that are not necessarily complex. We will therefore have to generalize the above discussion to the non-integrable situation. There is a more general setting of exact symplectic manifolds with properly embedded hypersurfaces to which this discussion can be extended, but our aims here are more modest. We remark that our construction of Floer homology for admissible Lagrangians is still valid if we perturb both the complex

structure and the fibration near the 0–level set so long as the following conditions are preserved:

- The perturbed almost complex structure is compatible with the symplectic form and J –convex at infinity.
- The map to \mathbb{C} near the 0–level set remains holomorphic.

The first condition is familiar from the study of the symplectic topology of Stein manifolds Eliashberg–Gromov [5]. The second condition is necessary for the validity of Lemma 2.8.

Note that admissibility is not stable under exact Hamiltonian perturbations (even those that preserve M). However, given two positively oriented admissible Lagrangians L_1 and L_2 one may still consider exact Hamiltonian deformations which preserve the admissibility of L_1 . As long as the tangent vectors v_1 and v_2 are appropriately oriented, Lemma 2.8 will apply for some ϵ . In particular, the usual proofs of invariance of Floer homology are valid for those Hamiltonian isotopies that preserve the admissibility of L_1 , and the positivity of the pair (L_1, L_2) . As in the classical situation, this allows us to compute Floer homology for a pair of admissible Lagrangian branes that do not intersect transversely away from M by choosing an appropriate Hamiltonian deformation of one of them.

In particular, if (L_1, L_2) is a pair of admissible Lagrangian branes which is negatively oriented, we can use Lemma 2.6 in order to unambiguously define

$$CF_*(L_1, L_2) \equiv CF_*(L_1, L_2'')$$

where the pair (L_1, L_2'') is positively oriented.

We also observe that our conventions for “positivity” are designed to guarantee that Floer’s old result that expresses Floer homology of nearby Lagrangians in terms of ordinary homology extends to this setting,

$$HF_*(L, L) \cong H_*(L, \partial L).$$

We now consider what happens when we vary the holomorphic map f in a family.

Lemma 2.13 *Let f_t be a 1–parameter family of symplectic fibrations which are holomorphic near the origin, and assume that $(f_t)^{-1}(0) = M$. There is a bijection between Hamiltonian isotopy classes of admissible Lagrangians for f_0 and f_1 . For Lagrangians satisfying $H^1(L, \partial L) = 0$, this bijection respects the Floer homology groups.*

Proof Assume that L is an admissible Lagrangian with respect to f_0 . Since ∂L is compact, we can uniformly bound the derivatives $\frac{\partial f}{\partial t}$ in a neighbourhood of ∂L . We can therefore choose an ϵ such that the parallel transport of ∂L with respect to f_t lies in an arbitrarily small neighbourhood of the parallel transport of ∂L with respect to $f_{t+\epsilon}$. If this neighbourhood is small enough, we can apply Lemma 2.2. In particular, subdividing the interval $[0, 1]$ into sufficiently many subintervals, we obtain a Hamiltonian isotopy between L and a Lagrangian which is admissible with respect to f_1 .

Let L_1 and L_2 be two admissible Lagrangians with respect to f_0 . We can choose the Hamiltonian isotopies which we used in the previous paragraph to be supported in an arbitrarily small neighbourhood of M , and such that no new intersection points are created. In particular, these isotopies will equal the identity near the images of the elements of $\mathcal{M}(p, q)$ for all pairs of intersection p and q between L_1 and L_2 . This yields the desired invariance of Floer homology groups. \square

2.5 Cohomology and cup product

With Poincaré duality in mind, we can now define Floer cohomology by simply re-indexing the Floer complex as

$$CF^*(L_1, L_2) \equiv CF_{n-*}(L_1, L_2)$$

which on cohomology yields

$$HF^*(L_1, L_2) \equiv HF_{n-*}(L_1, L_2).$$

Remark 2.14 Note that Floer homology and cohomology are Poincaré dual with these conventions, but there is no degree preserving duality between them. Whereas the classical analogue of our Floer homology groups is the homology of L relative its boundary, the classical analogue of our Floer cohomology group is the ordinary cohomology of L . These classical groups are indeed Poincaré dual. One can resolve this unfortunate state of affairs by introducing Floer homology in two flavours, with positive and negative orientations at the boundary. In this language, our Floer cohomology between L_1 and L_2 would indeed be the (ordinary) dual of Floer homology between L_2 and L_1 with the opposite convention to the one we have chosen. Since the main goal of this paper is to perform a computation, we will not discuss these issues further, and simply use the above definition.




Figure 4: A marked disc

As in the case of closed Lagrangians the Floer co-chain complex admits a cup product

$$CF^*(L_1, L_2) \otimes CF^*(L_2, L_3) \rightarrow CF^*(L_1, L_3).$$

In order to describe the cup product, we need some preliminary definitions. There are essentially no differences between the case we are studying, and that of closed Lagrangians which is explained in Fukaya and Oh [8].

Definition 2.15 A Riemann surface with *strip-like ends* is an open Riemann surface with a choice of biholomorphisms between its ends (ie, complements of sufficiently large compact subsets) and the strip $[0, 1] \times \mathbb{R}^+$.

We let D be the unit disc in \mathbb{C} , and $\xi = e^{\frac{2\pi i}{3}}$. Note that $T = D - \{\xi, \xi^2, 1\}$ admits the structure of a Riemann surface with strip-like ends. Assuming all Lagrangians intersect transversely and the pairs (L_1, L_2) , (L_2, L_3) and (L_1, L_3) are positively oriented, we consider finite energy pseudo-holomorphic maps $u: T \rightarrow N$ which satisfy the following conditions (see Figure 4):

- u maps the arcs $(1, \xi)$ to L_1 , (ξ, ξ^2) to L_2 , and $(\xi^2, 1)$ to L_3 .
- Along the strip-like ends associated to the punctures $(\xi, \xi^2, 1)$, the image of u converges uniformly to interior intersection points (p, q, r) among the Lagrangians L_j .
- The image of T under u does not intersect M .

We denote the above moduli space by $\mathcal{M}^{\text{adm}}(p, q, r)$. We can now define the cup product over \mathbb{Z}_2 by the formula

$$[p] \otimes [q] \mapsto \sum_r |\mathcal{M}^{\text{adm}}(p, q, r)| \cdot [r],$$

where the sum is taken over 0-dimensional moduli spaces. Just as we proved that $d^2 = 0$, the usual proof that this cup product descends to an associative product on cohomology applies in our situation.

To obtain a product in Floer cohomology over \mathbb{C} , we follow the same strategy as for obtaining signs in the differential. Given a holomorphic triangle T with strip like ends associated to three points p, q and r , we attach copies of the upper half-plane at the incoming ends to form a 1-sided strip $T \# H_p \# H_q$. As before, we obtain an operator $\bar{\partial}_{T \# H_p \# H_q}$. The gluing theorem yields a canonical isomorphism

$$\det(\bar{\partial}_{T \# H_p \# H_q}) \cong \det(\bar{\partial}_T) \otimes \det(\bar{\partial}_{H_p}) \otimes \det(\bar{\partial}_{H_q}).$$

There is a unique path (up to homotopy) between the boundary conditions for this operator and those for the operator $\bar{\partial}_{H_r}$, yielding a canonical isomorphism

$$\det(\bar{\partial}_{T \# H_p \# H_q}) \cong \det(\bar{\partial}_{H_r}).$$

Since u only contributes to the cup product when its associated $\bar{\partial}$ operator is invertible, we have a canonical trivialization of $\det(\bar{\partial}_T)$. So we obtain an isomorphism

$$\det(\bar{\partial}_{H_p}) \otimes \det(\bar{\partial}_{H_q}) \cong \det(\bar{\partial}_{H_r}),$$

whose compatibility with orientations determines the signed contribution of T to the image of $[p] \otimes [q]$ under the cup product.

3 Tropical geometry

3.1 Background

Let f be a Laurent polynomial in n variables over \mathbb{C} . In [11], Gel'fand, Kapranov, and Zelevinsky introduced the amoeba of f as the projection of $f^{-1}(0)$ to \mathbb{R}^n under the logarithm map

$$\text{Log}(z_1, \dots, z_n) = (\log |z_1|, \dots, \log |z_n|).$$

One may also define amoebas for varieties over fields other than \mathbb{C} . Amongst other results on varieties over non-Archimedean fields, Kapranov gave a description of their amoebas in term of polyhedral complexes in \mathbb{R}^n [4]. Mikhalkin then used this

description in order to obtain new results about the topology of complex hypersurfaces [15]. We will follow his point of view with small modifications.

In coordinates, we will write

$$f = \sum_{\alpha \in \mathbb{Z}^n} c_{\alpha} z^{\alpha}$$

with $c_{\alpha} \in \mathbb{C}$ and $z^{\alpha} = z_1^{\alpha_1} z_2^{\alpha_2} \cdots z_n^{\alpha_n}$. Since f is a polynomial, only finitely many monomials have non-zero coefficients, and we let A be the set of vectors which label these monomials. Let P be the convex hull of A thought of as a subset of \mathbb{R}^n (Newton polytope of f).

Definition 3.1 The *convex support* of a function $v: A \rightarrow \mathbb{R}$ is the largest convex function $\hat{v}: P \rightarrow \mathbb{R}$ such that $\hat{v}(\alpha) \leq v(\alpha)$.

The function \hat{v} is in fact piece-wise linear on P , so we may decompose $P = \bigcup P_{v_i}$, where each v_i is a linear function, and P_{v_i} is the domain where v_i and \hat{v} agree. We say that the decomposition $P = \bigcup P_{v_i}$ is the *coherent subdivision* of P induced by v . The following result is well known.

Lemma 3.2 Each polytope of the coherent subdivision induced by a function $v: A \rightarrow \mathbb{R}$ is a lattice polytope whose vertices lie in A . \square

We will be particularly interested in subdivisions which satisfy the following additional condition (See Theorem 3.11).

Definition 3.3 A subdivision of P is *maximal* if each polytope of the subdivision is equivalent under the action of $ASL(n, \mathbb{Z})$ to the standard n -dimensional simplex.

The function v also determines a piecewise linear function $L_v: \mathbb{R}^n \rightarrow \mathbb{R}$, its Legendre transform, which is defined by,

$$(3-1) \quad L_v(u) = \max_{\alpha \in A} (\langle \alpha, u \rangle - v(\alpha)).$$

Since it is defined as the maximum of finitely many linear functions, L_v is smooth away from a subset of zero measure.

Definition 3.4 Given any function $v: A \rightarrow \mathbb{R}$, the locus of non-smoothness of L_v is a *tropical hypersurface* or *tropical amoeba* of f .

Note that this tropical hypersurface of f is the set where two or more linear functions agree. It will be sufficient for our purposes to think of the tropical amoeba Π as an $n - 1$ dimensional polyhedral complex whose k skeleton corresponds to the set where $n - k + 1$ functions agree. In particular, each k -face is dual to a unique $(n - k)$ -polytope of the polyhedral subdivision of A induced by ν . We will denote the dual of τ by $\check{\tau}$. Given a face σ of Π , we will also use $\check{\sigma}$ for its dual polytope in P . Our conventions are that all polytopes and faces are closed.

When $k = 0$, we obtain a component of $\mathbb{R}^n - \Pi$ rather than a face of Π . Concretely, every component of $\mathbb{R}^n - \Pi$ may be labeled by the unique $\alpha \in A$ which, on the given component, achieves the maximum in the equation defining L_ν . We can therefore write

$$\mathbb{R}^n - \Pi = \bigsqcup_{\alpha \in A} C_\alpha - \partial C_\alpha.$$

In general, some of these components may be empty.

Going back to the complex numbers, we may use the function ν to define the patch-working polynomials

$$f_t(z) = \sum_{\alpha \in A} c_\alpha t^{-\nu(\alpha)} z^\alpha.$$

For every t , we consider \mathcal{A}_t the Log-amoeba of f_t . One expression of the connection between tropical and complex amoebas is given by the following theorem which is due to Mikhalkin [15, Theorem 5] and Rullgård [17, Theorem 9].

Theorem 3.5 *In the Gromov–Hausdorff topology, the sets $\mathcal{A}_t / \log(t)$ converge to Π as t goes to infinity.* □

We will be studying the hypersurface $f^{-1}(0) \subset (\mathbb{C}^*)^n$ from the symplectic point of view. In order to be precise, we must choose a symplectic structure on $(\mathbb{C}^*)^n$. We will use

$$\omega = \sum_{j=1}^n \frac{dz_j \wedge d\bar{z}_j}{2i |z_j|^2},$$

where $\{z_j\}_{j=1}^n$ are the standard coordinates on $(\mathbb{C}^*)^n$. With respect to the usual $(\mathbb{C}^*)^n$ action, this is an invariant Kähler form on $(\mathbb{C}^*)^n$. The vectors $\{\frac{\partial}{\partial z_j}\}_{j=1}^n$ and their complex conjugates form an orthogonal basis for the Kähler metric. In particular,

$$\left| \frac{\partial}{\partial z_j} \right| = \frac{1}{|z_j|}$$

and hence

$$|dz^j| = |z_j|.$$

It will be convenient to have a different description of $(\mathbb{C}^*)^n$ as a symplectic manifold. Consider the cotangent bundle of \mathbb{R}^n with standard coordinates $\{u_j\}_{j=1}^n$ on \mathbb{R}^n and $\{\theta_j = du_j\}_{j=1}^n$ on the fibre of the cotangent bundle. The coordinates $\{u_j\}_{j=1}^n$ induce an affine structure on the base, which determines a lattice in the fibre. We will choose this lattice to be spanned by $\{2\pi\theta_j\}_{j=1}^n$.

Lemma 3.6 *The quotient of $T^*\mathbb{R}^n$ by the lattice $2\pi\mathbb{Z}^n$ in each cotangent fibre is symplectomorphic to $(\mathbb{C}^*)^n$ with the identification given by the exponential map*

$$(u_j, \theta_j) \rightarrow e^{u_j + i\theta_j}. \quad \square$$

Note that the inverse of the above exponential map is the logarithmic map whose first component arises in the definition of amoebas,

$$z_j \rightarrow (\log |z_j|, \arg(z_j)).$$

We will omit all notation for the above symplectomorphisms, and will use either coordinate system at our convenience. Note, however, that these symplectomorphisms identify the projection of the cotangent bundle onto its base manifold with the Log projection of $(\mathbb{C}^*)^n$ onto \mathbb{R}^n , hence identify the cotangent fibres with the set of points in $(\mathbb{C}^*)^n$ whose components have fixed norms. We will also be using the standard Euclidean metric on \mathbb{R}^n . As usual this metric identifies the tangent and cotangent bundles, and carries the natural complex structure of the tangent bundle to a complex structure on the cotangent bundle which is compatible with the symplectic form.

The advantage of the cotangent bundle point of view is that many constructions can now be performed on the base \mathbb{R}^n , and some of them reduce to linear algebra. For example, recall that a diffeomorphism of a manifold induces a symplectomorphism of its cotangent bundle. If, in addition, this diffeomorphism is an isomorphism of the affine structure, it will induce a symplectomorphism of the associated Lagrangian torus bundle. Thus, every affine transformation in $ASL(n, \mathbb{Z})$ induces a symplectomorphism of $(\mathbb{C}^*)^n$. Assume, for simplicity, that we have a linear transformation. If we represent it by a matrix A , then in the standard coordinates of $T^*\mathbb{R}^n$ this symplectomorphism is given by multiplying by A in the coordinates corresponding to the base, and by $(A^T)^{-1}$ in the coordinates corresponding to the fibre. In general such a transformation does not preserve the standard Euclidean metric on the base, hence does not preserve the complex structure on the fibre.

Indeed, given an element of $ASL(n, \mathbb{Z})$, there is a different automorphism of the torus fibration which does preserve the complex structure. Assuming again that we have a linear transformation represented by an integral matrix A , this complex automorphism,

in the standard coordinates of $T^*\mathbb{R}^n$, is given by using A in both the base and the fibre directions. In the standard coordinates of $(\mathbb{C}^*)^n$ as a complex manifold, it can be thought of as the multiplicative change of variables

$$(z_1, \dots, z_n) \rightarrow (z^{\alpha_1}, \dots, z^{\alpha_n})$$

where α_i are the rows of A .

3.2 The tropical model

Let X be a complete toric variety. Let v_i be the primitive vertices of the 1-cones of the fan Δ defining X . We consider $A = \{v_i\}_{i=1}^m \cup \{0\}$ and its convex hull P . To each maximal cone $\tau \in \Delta(n)$ we assign the convex polytope $P(\tau)$ which is the convex hull of the set $\{0\} \cup \{v_i | v_i \in \tau\}$.

We will assume that every polytope $P(\tau)$ is a minimal simplex; this is equivalent to the requirement that the set $\{v_i | v_i \in \tau\}$ forms a basis for \mathbb{Z}^n . Fans with this property correspond to smooth toric varieties, [10, page 34]. Let W be the Laurent polynomial

$$W(z) = -1 + \sum_{0 \neq \alpha \in A} z^\alpha$$

and let $M = W^{-1}(0)$.

Remark 3.7 The choice of coefficients on the monomials which appear in W is only done for convenience. After passing to the tropical limit, any other choice of (non-zero) coefficients is related to the one we picked by rescaling followed by a symplectomorphism.

Lemma 3.8 *Every ample line bundle on X induces a coherent subdivision of P which near 0 is given by the polytopes $P(\tau)$. In particular, near 0, we have a maximal triangulation.*

Proof Every ample line bundle on X is determined up to an integral linear function by a strictly convex function ϕ on $|\Delta|$ which is integral linear on each cone. Consider the coherent subdivision of P induced by $\phi|_A$. To prove the lemma, it will suffice to prove that 0 is a vertex of this subdivision, and that each full-dimensional polytope of the subdivision abutting on the origin is equal to $P(\tau)$ for some cone $\tau \in \Delta(n)$.

To prove this we observe that on each polytope $P(\tau)$ the convex support ψ of the function $\phi|_A$ agrees with ϕ since the latter is convex. Since ϕ is in fact strictly convex, we conclude that each $P(\tau)$ is contained in a distinct polytope of the subdivision. It

remains to show that $P(\tau)$ is equal to a polytope of the subdivision. Indeed, if this were not the case, this polytope would have a vertex not belonging to τ . But coherent subdivisions do not create new vertices, so this vertex is in fact the minimal vertex $v_{\tau'}$ of some 1–cone τ' . This contradicts the strict convexity of ϕ . \square

The choice of an ample line bundle therefore induces a choice of patchworking polynomials W_t , whose amoebas converge (after rescaling) to a tropical hypersurface Π as in the previous section. Since the origin is a vertex of the subdivision induced by ϕ , we conclude that there must be a non-empty component, Q , of the complement of Π which is dual to the origin.

Corollary 3.9 *Q is the convex polytope which, in toric geometry, describes X as a toric variety with ample line bundle $\mathcal{O}(1)$.*

Proof Observe that Equation (3–1) shows that Q is the polytope consisting of points y which satisfy

$$\langle v_i, y \rangle \leq \phi(v_i)$$

for every primitive vertex v_i . After adjusting for different sign conventions, this is therefore the polytope whose integral points form a basis for the space of sections of the ample line bundle $\mathcal{O}(1)$ [10, Section 3.4]. \square

We do not get a global triangulation of our Newton polytope, but we certainly have a maximal triangulation near the origin. Since we will only be studying Π in a neighbourhood of Q , we may appeal to the results of Mikhalkin about maximal tropical degenerations.

Remark 3.10 While we could use any ample line bundle, it will be convenient to have a lattice point in the interior of Q . Note that this can be achieved by replacing $\mathcal{O}(1)$ by a sufficiently high power, and we may assume that the origin is an interior point.

3.3 Twisting the tropical zero-section

We begin by producing a “tropical” version of our construction. In particular, the boundary of this version will not be smooth, hence it will not be clear how the results that we obtain would be invariant under a natural class of Hamiltonian isotopies as discussed in the previous section. Nonetheless, as the complete version requires many choices that are necessary to produce a meaningful answer, but that obscure the simple nature of the construction, we will prefer to discuss the tropical case first.

We now consider the situation where our toric variety X is smooth. In this situation, we may restate a result of Mikhalkin.

Theorem 3.11 [15, Lemma 6.2] *Given a maximal triangulation of P , there exists a natural choice of a piecewise smooth symplectic hypersurface M_∞ of $(\mathbb{C}^*)^n$ which projects to the tropical amoeba Π . \square*

In fact, we only need this theorem as motivation, since we will be using Proposition 4.2 to prove the precise results. We will therefore not give a complete description of M_∞ which may be thought of as a limit of $W_t^{-1}(0)$ after rescaling, but we will use the following results that follow from the proof of Theorem 3.11 or of Proposition 4.2.

- The preimage of a point on a k -face σ of Π contains a subtorus of the fibre $\mathbb{R}^n/2\pi\mathbb{Z}^n$ which is parallel to the tangent space of σ (thought of modulo \mathbb{Z}^n of course).
- The preimage a point on the interior of an $n - 1$ dimensional facet σ is equal to a torus in the fibre which is parallel to the tangent space of σ .

Remark 3.12 We have chosen the coefficients of the monomials that appear in W exactly in such a way as to ensure that the intersection of M_∞ with the zero section contains $\partial Q \subset \Pi$, the boundary of the component of $\mathbb{R}^n - \Pi$ which corresponds to the origin. This guarantees that the preimage of an $n - 1$ facet of Q is exactly equal to the tangent space of this facet. Note that the tangent space to a k face is equal to the intersection of the tangent spaces to all the maximal cells that contain it, so that the statement extends to lower dimensional strata of the boundary of Q .

Since the zero section is a Lagrangian submanifold of $T\mathbb{R}^n$, the polytope Q may be thought of as a Lagrangian ball with boundary on M_∞ ; we denote this ball by \mathcal{L}_∞ .

Consider the Hamiltonian function

$$H_\infty(u_1, \dots, u_i) = -\pi \sum_{i=1}^n u_i^2$$

and let ϕ_∞^1 be its time-1 Hamiltonian flow. In the universal cover, an explicit formula for ϕ_∞^1 is given by

$$\phi_\infty^1(u_1, \theta_1, \dots, u_n, \theta_n) = (u_1, \theta_1 - 2\pi u_1, \dots, u_n, \theta_n - 2\pi u_n)$$

which we can write more conveniently as

$$(3-2) \quad \phi_\infty^1(u, \theta) = (u, \theta - 2\pi u).$$

Lemma 3.13 $\phi_\infty^1(\partial\mathcal{L}_\infty) \subset M_\infty$

Proof Note that Equation (3–2) implies that if u is an integral point of \mathbb{R}^n , then ϕ_∞^1 pointwise fixes the inverse image of u under the projection map. In particular, the inverse image of every vertex of Q is fixed.

Consider a top dimensional cell of Q . After a suitable translation by an element of \mathbb{Z}^n , we may assume that one of its vertices is the origin. But the restriction of ϕ_∞^1 to the zero section of a linear subspace is just the map

$$(v, 0) \rightarrow (v, -2\pi v).$$

So the subset of M_∞ which lies over a top dimensional cell is preserved by ϕ_∞^1 . Since $\partial\mathcal{L}_\infty \subset M_\infty$, this establishes the result for the restriction of $\partial\mathcal{L}_\infty$ to the top dimensional cells. The lemma follows from Remark 3.12. \square

We will denote $\phi_\infty^1(\mathcal{L}_\infty)$ by $\mathcal{L}_\infty(1)$. The previous lemma in fact shows that the time l flow satisfies

$$\phi_\infty^l(\partial\mathcal{L}_\infty) \subset M_\infty$$

for every integer l , so we have well defined Lagrangian balls $\mathcal{L}_\infty(l)$ with boundaries on M_∞ .

3.4 A preliminary computation of Floer groups

Even though we don't have smooth boundary conditions yet, we will set out to compute the Lagrangian Floer homology groups of the pairs $(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2))$. We are missing the usual compactness and transversality results that guarantee that our answers will be invariant under Hamiltonian perturbations but we will proceed regardless with the construction. We will justify the use of small Hamiltonian perturbations of the interior points by constructing admissible Lagrangians in Corollaries 5.9 and 5.10, which will also resolve the issue of boundary intersection points.

Note that Lagrangian sections of the cotangent bundle have a natural grading coming from the Morse index [18, Example 2.10]. In particular, whenever two such sections \mathcal{L} , \mathcal{L}' intersect transversely at a point p , we may write them locally as the graphs of df and df' for smooth real valued functions f and f' such that p is a non-degenerate critical point of $f' - f$. As an element of $CF_*(\mathcal{L}, \mathcal{L}')$, the class of p will have degree equal to its Morse index as a critical point of $f' - f$. We write $\mu_p(\mathcal{L}, \mathcal{L}')$ for this index. These choices carry naturally to the quotient of the cotangent bundle by a lattice coming from an affine structure on the base [14] to give a canonical choice of grading on sections of Lagrangian torus bundles.

We also need to resolve the issue of the status of intersection points that occur on the boundary. Our temporary ad-hoc prescription, which we will justify in Section 5.4, is

that boundary intersection points are included in $CF_*(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2))$ if and only if $l_1 < l_2$. We will also stipulate that the group $CF_*(\mathcal{L}_\infty(l), \mathcal{L}_\infty(l))$ is isomorphic to \mathbb{C} concentrated in degree n .

Lemma 3.14 *The Floer homology groups between the Lagrangians $\mathcal{L}_\infty(j)$ are given by*

$$HF_n(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2)) = \bigoplus_{p \in Q \cap \frac{1}{l_2 - l_1} \mathbb{Z}^n} \mathbb{C} \cdot [p]$$

with all other groups zero if $l_1 < l_2$, and

$$HF_0(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2)) = \bigoplus_{p \in (Q - \partial Q) \cap \frac{1}{l_2 - l_1} \mathbb{Z}^n} \mathbb{C} \cdot [p]$$

with all other groups zero if $l_1 > l_2$.

Proof First we reduce to the case where the first Lagrangian is the zero section by applying an appropriate twist. In other words, we have a natural graded isomorphism

$$CF_*(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2)) \cong CF_*(\mathcal{L}_\infty(l_1 + i), \mathcal{L}_\infty(l_2 + i))$$

for any integer i . Considering the case $i = -l_1$ reduces the computation to one we've already done.

Indeed, we have already remarked in our proof of Lemma 3.13 that the intersection points of \mathcal{L}_∞ and $\mathcal{L}_\infty(l)$ correspond to the $\frac{1}{l} \mathbb{Z}^n$ points of Q . For simplicity, we first discuss the case where $l > 0$. Note that the lift of $\mathcal{L}_\infty(l)$ to $T^*\mathbb{R}^n$ which intersects the zero section at a point $p = (p_1, \dots, p_n)$ is given by the differential of the function $-l\pi \sum_{i=1}^n (u_i - p_i)^2$. Therefore, since the Morse index of this function at p is n if $0 < l$, the Floer complex is concentrated in degree n , where it is given by

$$CF_n(\mathcal{L}_\infty, \mathcal{L}_\infty(l)) = \bigoplus_{p \in Q \cap \frac{1}{l} \mathbb{Z}^n} \mathbb{C} \cdot [p].$$

The same computation yields that if $l < 0$ the Floer complex is concentrated in degree 0, ie,

$$CF_0(\mathcal{L}_\infty, \mathcal{L}_\infty(l)) = \bigoplus_{p \in (Q - \partial Q) \cap \frac{1}{l} \mathbb{Z}^n} \mathbb{C} \cdot [p].$$

Since all these complexes are concentrated in one degree, the differential is necessarily trivial. □

We will now pass to cohomology in order to compute the cup product. Recall that the degree of a transverse intersection point $p \in \mathfrak{L} \cap \mathfrak{L}'$ in cohomology is

$$I_p(\mathfrak{L}, \mathfrak{L}') = n - \mu_p(\mathfrak{L}, \mathfrak{L}').$$

We will compute the cup product

$$HF^*(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2)) \otimes HF^*(\mathcal{L}_\infty(l_2), \mathcal{L}_\infty(l_3)) \rightarrow HF^*(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_3)),$$

by counting holomorphic triangles connecting three intersection points. Assume the integers l_1 , l_2 , and l_3 are all distinct.

Lemma 3.15 *If $p = (p_1, \dots, p_n)$ is an intersection point of $\mathcal{L}_\infty(l_1)$ and $\mathcal{L}_\infty(l_2)$, and $q = (q_1, \dots, q_n)$ is an intersection point of $\mathcal{L}_\infty(l_2)$ and $\mathcal{L}_\infty(l_3)$, then there exists at most one intersection point r of $\mathcal{L}_\infty(l_1)$ and $\mathcal{L}_\infty(l_3)$ such that*

$$\mathcal{M}(p, q, r) \neq \emptyset.$$

Furthermore, if the intersection point r exists, it is given by the formula

$$(3-3) \quad r = \frac{(l_2 - l_1)p + (l_3 - l_2)q}{l_3 - l_1}.$$

Remark 3.16 The reader should note the similarity with Equation (1–2).

Proof There is a topological obstruction to the existence of a holomorphic triangle with the appropriate boundary conditions which we now describe.

After choosing a lift \tilde{p} of p to $T^*\mathbb{R}^n$, there are uniquely determined lifts of each Lagrangian. In particular, the lifts of $\mathcal{L}_\infty(l_2)$ and $\mathcal{L}_\infty(l_3)$ intersect at a lift \tilde{q} of q , while the lifts of $\mathcal{L}_\infty(l_1)$ and $\mathcal{L}_\infty(l_3)$ intersect at most in a unique points which we call \tilde{r} . For the next few paragraphs, we will compute everything in term of these lifts.

Using the metric to identify the cotangent fibre with the tangent space, and the vector space structure on the base to identify each tangent space with that of the origin, we may write $\tilde{p} = (p, \tilde{p}_1, \dots, \tilde{p}_n)$. The lift of $\mathcal{L}_\infty(l_1)$ can be identified with the graph of the affine transformation

$$(x_1, \dots, x_n) \rightarrow (-l_1(x_1 - p_1) + \tilde{p}_1, \dots, -l_1(x_n - p_n) + \tilde{p}_n)$$

and similarly for the lift of $\mathcal{L}_\infty(l_2)$. Since q lies on $\mathcal{L}_\infty(l_2)$, its lift is therefore

$$\tilde{q} = (q, -l_2(q_1 - p_1) + \tilde{p}_1, \dots, -l_2(q_n - p_n) + \tilde{p}_n).$$

This allows us to conclude that the lift of $\mathcal{L}_\infty(l_3)$ that we're considering is the graph of

$$\begin{aligned} x_i &\rightarrow -l_3(x_i - q_i) + \tilde{q}_i \\ &= -l_3(x_i - q_i) - l_2(q_i - p_i) + \tilde{p}_i. \end{aligned}$$

Note that our sought after intersection point r is given by the solution to the system of linear equations

$$-l_1(r_i - p_i) + \tilde{p}_i = -l_3(r_i - q_i) - l_2(q_i - p_i) + \tilde{p}_i, \quad 1 \leq i \leq n,$$

which is clearly given by Equation (3-3). □

3.5 Cup product

We must now count holomorphic triangles with appropriate boundary conditions in order to compute the cup product. As in the previous section, any statements that are not justified here are handled in future sections, in particular Section 5.4. As we're working with products of Lagrangians in different copies of \mathbb{C}^* , we appeal to the fact that the holomorphic triangle admits no deformations, so proving regularity for a pseudo-holomorphic map

$$u: (T, \partial T) \rightarrow (M, \cup_i L_i)$$

amounts to proving the surjectivity of the $\bar{\partial}$ operator

$$W^{1,p}((T, \partial T), (u^*TM, \cup_i u^*TL_i)) \rightarrow L^p(T, \Omega^{0,1}(T) \otimes u^*TM).$$

If $L_i \hookrightarrow M$ are given by products of Lagrangians in different factors of the symplectic manifold M , then the above map splits into direct summands, and surjectivity amounts to surjectivity for each of the summands of u .

It is well known that non-constant holomorphic polygons on Riemann surfaces are regular, though we will only give a proof in this specific situation.

Lemma 3.17 *Consider three lines L_1, L_2 and L_3 in $\mathbb{R}^2 = \mathbb{C}$ with rational slope which intersect at three distinct points (p, q, r) . If there is a holomorphic triangle in $\mathcal{M}(p, q, r)$, then it is necessarily regular.*

Proof Note that the usual index theoretic argument shows that such a holomorphic triangle has index 0. In particular, it suffices to show the operator

$$W^{1,p}((T, \partial T), (\mathbb{C}, \cup_i TL_i)) \rightarrow L^p(T, \mathbb{C})$$

is injective.

Assume that F is an element of the kernel, and choose an integer K such that the images of the tangent lines TL_i under the map $z \rightarrow z^K$ is the real axis. Note that F^K will therefore be a holomorphic map from T to \mathbb{C} which takes ∂T to \mathbb{R} . By the maximum principle, all such maps are constant. However, since F has finite $W^{1,p}$ norm, and hence finite L^p norm with respect to an infinite measure (because of the strip-like ends), it must therefore be identically 0. \square

Lemma 3.18 *Assume p and q are such that $r = \frac{(l_2-l_1)p+(l_3-l_2)q}{l_3-l_1}$ lies in the polytope Q and that the integers l_1, l_2 , and l_3 are distinct. If $l_1 < l_2$ then*

$$\mathcal{M}(p, q, r) \neq \emptyset$$

if and only if

$$(3-4) \quad l_3 < l_1 < l_2 \text{ or } l_1 < l_2 < l_3.$$

If $l_2 < l_1$, then the moduli space is non-empty if and only if

$$(3-5) \quad l_2 < l_3 < l_1.$$

Proof By the preceding discussion, the existence of a holomorphic triangle in the total space with boundary on our given affine subspaces is equivalent to the existence of n holomorphic triangles with boundary on a configuration of straight lines in \mathbb{R}^2 . But Riemann's mapping theorem says that such a triangle exists if and only if orientations are preserved. Moreover, since in each factor such a triangle is unique if it exists, there can be at most one holomorphic disc with appropriate boundary conditions. One might also be concerned that the holomorphic triangles we're finding may go outside the the inverse image of Q even when all the intersection points between the different Lagrangians lie within Q . But this doesn't happen because Q is convex. Indeed, representing the triangle with three marked points as a geodesic triangle T in the plane, we see that each component u_i of the holomorphic map $u: T \rightarrow \mathbb{C}^n$ is in fact given by an affine transformation of the plane. In particular, the image of T is the flat triangle determined by the points $(\tilde{p}, \tilde{q}, \tilde{r})$, hence projects to the flat triangle determined by (p, q, r) . Convexity implies that this triangle is contained in Q .

We have therefore reduced the problem to establishing whether the triples $(\xi, \xi^2, 1)$ and $((p_i, \tilde{p}_i), (q_i, \tilde{q}_i), (r_i, \tilde{r}_i))$ have the same orientation, assuming the latter consists of distinct points. It follows from Formula (3-3) (see also, Figure 5) that if $l_1 < l_2$, the orientations are the same if and only if

$$l_3 < l_1 < l_2 \text{ or } l_1 < l_2 < l_3$$




Figure 5: Holomorphic discs with $l_1 < l_2 < l_3$ (1), and $l_2 < l_3 < l_1$ (2)

while if $l_2 < l_1$, then the orientations are the same if and only if

$$l_2 < l_3 < l_1.$$

In particular, this condition is independent of which factor we’re considering, so the above is the necessary and sufficient condition for the existence of a holomorphic disc with marked points on $(\tilde{p}, \tilde{q}, \tilde{r})$ if these points are distinct.

We must now deal with the case where the marked points are not distinct. Observe that since all the slopes l_i are assumed to be different, this can only occur if all three points are equal. We must appeal to a yet unproven invariance under Hamiltonian deformations to interpret the product among Lagrangian submanifolds in non-generic positions in terms of the product for nearby Lagrangians. Note the Conditions (3–4) or (3–5) are clearly invariant under small Hamiltonian deformations. In particular, given a triple intersection point between $\mathcal{L}_\infty(l_1)$, $\mathcal{L}_\infty(l_2)$, and $\mathcal{L}_\infty(l_3)$, we may locally deform one of the Lagrangians, through Lagrangians which are also given as $\mathcal{L}_\infty(l)$ for some real number l , preserving Conditions (3–4) or (3–5). But the argument we gave when l_i are distinct integers did not depend on their integrality, so we reach the desired conclusion that Conditions (3–4) or (3–5) suffice to describe the existence of a regular holomorphic triangle. \square

Remark 3.19 The perturbation argument that we’re using breaks down whenever $p = q = r$ is a point on the boundary. In Section 5, we will “push intersection points to the interior,” and thereby legitimate the perturbation argument.

If two of the Lagrangian sections are equal, recall that we have made an as yet unjustified claim that the corresponding Floer cohomology group is concentrated in degree 0 and is isomorphic to a copy of \mathbb{C} . We will justify in Section 5.4 that under the cup product, this group acts by scalar multiplication on the Floer cohomology group of a pair of Lagrangian manifolds.

To complete our analysis of the presence of holomorphic triangles, we must find appropriate conditions for \tilde{r} to lie within the prescribed region. We will only do this for a special case.

Lemma 3.20 *Assume that $l_1 \leq l_2 \leq l_3$. The product in Floer cohomology is given by*

$$(3-6) \quad HF^0(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_2)) \otimes HF^0(\mathcal{L}_\infty(l_2), \mathcal{L}_\infty(l_3)) \rightarrow HF^0(\mathcal{L}_\infty(l_1), \mathcal{L}_\infty(l_3))$$

$$[p] \otimes [q] \mapsto \pm \left[\frac{(l_2 - l_1)p + (l_3 - l_2)q}{l_3 - l_1} \right].$$

Proof Since Q is convex, and all the coefficients in Equation (3-3) are positive, we conclude that \tilde{r} projects to an intersection point of $\mathcal{L}_\infty(l_1)$ and $\mathcal{L}_\infty(l_3)$. In the case of strict inequalities among the l_j , Equation (3-4) implies that there is necessarily a holomorphic triangle connecting p , q and r . The case where two of the Lagrangians may be equal follows from the previous discussion. \square

We also know from Equation (3-5) that there might be other non-trivial cup products if the condition $l_1 \leq l_2 \leq l_3$ does not hold. The point is that these are the same products that show up on the algebraic side. We will take this up again in Section 5.5.

We will now prove that we may choose all signs in Equation (3-6) to be positive.

Lemma 3.21 *There is a choice of generators for $CF^*(\mathcal{L}(l_i), \mathcal{L}(l_j))$ such that all the signs that arise in the cup product are positive.*

Proof We already know that

$$[p] \otimes [q] \mapsto \pm[r]$$

and it remains to prove that all signs may be chosen to be positive. However, since any configuration of three Lagrangian sections with $l_1 < l_2 < l_3$ is isotopic through Lagrangian sections to any other configuration and the cup product is invariant under such Lagrangian isotopies, we conclude that all signs are either positive or negative. If necessary, we complete the argument by replacing every generator by its negative. \square

In fact, this argument extends to products in which the triple of Lagrangians does not necessarily satisfy the condition $l_1 < l_2 < l_3$. Indeed, one can then simply dualize the operation that induces the cup product to reduce every other possible configuration to the case $l_1 < l_2 < l_3$. For example, if $l_3 < l_1 < l_2$, then, restricting to the interior intersection points, the map

$$CF^*(\mathcal{L}(l_1), \mathcal{L}(l_2)) \otimes CF^*(\mathcal{L}(l_2), \mathcal{L}(l_3)) \rightarrow CF^*(\mathcal{L}(l_1), \mathcal{L}(l_3))$$

can be obtained by dualizing the map

$$CF^*(\mathcal{L}(l_3), \mathcal{L}(l_1)) \otimes CF^*(\mathcal{L}(l_1), \mathcal{L}(l_2)) \rightarrow CF^*(\mathcal{L}(l_3), \mathcal{L}(l_2)).$$

But we know from the previous lemma that with our choice of generators, the sign that appears in this last product is positive, hence so it is for the first one. Once this is established for the interior intersection points, it immediately follows that it is true when boundary intersection points are involved as well because the signs are determined by local considerations which are insensitive to the distinction between the interior and the boundary.

4 Tropical localization

Mikhalkin used Theorem 3.5 to study the topology of smooth complex hypersurfaces M of $(\mathbb{C}^*)^n$ by producing a subset of \mathbb{R}^n which interpolates between the polyhedral complex Π (thought of as the tropical amoeba of a non-Archimedean variety) and the Log-amoeba of M . He observed that the tropical amoeba has a complex analogue to which the hypersurface M degenerates. We will give a slight modification of his proof which will allow us to obtain a stronger result about the symplectic structures.

4.1 Construction

Recall from Section 3 that each component of the complement of $\log(t)\Pi = \Pi_t$ corresponds to a vertex α of a coherent subdivision. We defined C_α in terms of the linear function attached to α , but one might equivalently define $\log(t)C_\alpha = C_{\alpha,t}$ as the image of the open subset of $(\mathbb{C}^*)^n$ where the monomial $t^{-\nu(\alpha)}z^\alpha$ dominates all others. For the purposes of this section, we will need some bounds on polyhedral geometry of P . For simplicity, we will assume that $c_\alpha = 1$.

Let N be the maximum of the ℓ^1 norms of all vectors $\alpha - \beta$ for α, β neighbouring vertices of the subdivision, and of all vectors $\alpha \in A$. Note that N is also an upper bound for the Euclidean norm of such vectors. Assume that ν induces a triangulation, and that the affine map from the standard k -dimensional vector space to \mathbb{R}^n induced

by every simplex of the subdivision of A distorts length by an amount bounded by $\rho > 1$. Note that such a bound exists simply because these affine maps are all injective.

If α and β are distinct elements of A , let $\mathcal{H}(\alpha, \beta)$ denote the hyperplane defined by the equation

$$\langle \alpha, u \rangle - v(\alpha) = \langle \beta, u \rangle - v(\beta).$$

Note that every face of Π is supported by such a hyperplane.

Lemma 4.1 *There exists a constant $c > 0$ such that for ϵ sufficiently small*

$$d(p, C_\alpha) \geq \epsilon \Rightarrow d(p, \mathcal{H}(\alpha, \beta)) \geq 2c\epsilon$$

for every $p \in C_\beta$ and for all pairs $\alpha \neq \beta$.

Proof Since A only has finitely many elements, it suffices to find a bound for each of them and then take the minimum. Accordingly, fix a component C_β .

Choose a constant δ such that 2δ neighbourhood of C_α can only intersect C_β if their boundaries share a face. Since Π is a finite cell complex and $\mathcal{H}(\alpha, \beta)$ cannot asymptotically approach C_β at infinity, there exists a constant K such that $d(p, \mathcal{H}(\alpha, \beta)) \geq K$ whenever p does not lie in the δ neighbourhood of C_α .

Let us assume for simplicity that the face σ shared by C_α and C_β is bounded. In this case, the boundary of the δ neighbourhood of σ intersects C_β in a compact subset which we will denote S . Note that convexity of C_β implies that $C_\beta \cap \mathcal{H}(\alpha, \beta) = \sigma$. In particular, the distances to $\mathcal{H}(\alpha, \beta)$ and σ are bounded above and below on S by non-zero constants, so an appropriate ratio yields the desired constant c . Consider $p \in C_\beta - C_\alpha$ such that $d(p, \sigma) = \epsilon < \delta$. There are points q on σ and r on $\mathcal{H}(\alpha, \beta)$ which realize the distance from p to these respective sets. The ratio between these distances can be computed from the angles of the triangle with corners p, q and r . By extending the segment from p to q , we eventually reach a point p' on S . Note that q is the point of σ closest to p' . If r' is the orthogonal projection of p' on $\mathcal{H}(\alpha, \beta)$, then the right triangles with corners (p, q, r) and (p', q, r') are similar. In particular, the ratios of their sides are equal. Since the ratio of the sides of the triangle (p', q, r') is bounded by $2c$, the same bound works for the triangle (p, q, r) . Further, since

$$d(p, q) \geq d(p, C_\alpha) = \epsilon,$$

we conclude that

$$d(p, \mathcal{H}(\alpha, \beta)) = d(p, r) \geq 2cd(p, q) \geq 2c\epsilon.$$

If the intersection between C_α and C_β is not compact, we choose a ball B that for every point q_1 of $(C_\alpha \cap C_\beta) - B$ there is a point q_2 in $C_\alpha \cap C_\beta \cap B$ such that there is

an isometry of the 2δ neighbourhoods of q_1 and q_2 which takes the intersection of Π with one neighbourhood to its intersection with the other. The existence of such a ball is a consequence of the fact that Π is a finite polyhedral complex. If p lies in the δ neighbourhood of q , then the point of $\mathcal{H}(\alpha, \beta)$ nearest to p lies in the 2δ neighbourhood of q . So the problem is entirely local, and the bound obtained by considering the intersection of B with the boundary of the δ neighbourhood of C_α in C_β yields a bound which is also valid for the complement of B . The rest of the argument proceeds as in the compact case. \square

Now pick ϵ such that $\delta > \epsilon > 0$. By Theorem 3.5, there exists a T such that $d(\mathcal{A}_t, \Pi_t) < \log(t)\epsilon$ for all $t > T$. Choose such ϵ and $T > 1$ such that the following additional conditions are satisfied for all $t > T$

$$(4-1) \quad \frac{e^{-c\epsilon \log(t)}}{\epsilon \log(t)} < \frac{1}{40 |A| \rho}$$

$$(4-2) \quad e^{-c\epsilon \log(t)} < \frac{1}{5 |A|^2 \rho N}.$$

In addition, choose nowhere negative C^∞ functions ϕ_α on \mathbb{R}^n such that the following properties hold

$$(4-3) \quad d(p, C_{\alpha,t}) \leq \frac{\epsilon \log(t)}{2} \Leftrightarrow \phi_\alpha(p) = 0$$

$$(4-4) \quad d(p, C_{\alpha,t}) \geq \epsilon \log(t) \Leftrightarrow \phi_\alpha(p) = 1$$

$$(4-5) \quad \sum_{i=1}^n \left| \frac{\partial \phi_\alpha(p)}{\partial u_i} \right| < \frac{4}{\epsilon \log(t)}.$$

We will also abuse notation and write $\phi_\alpha(z)$ for $\phi_\alpha(\text{Log}(z))$. Further, we will assume that ϕ_α and the norm of its derivatives are C^0 close to functions which depend only on the distance to C_α .

We now consider the family of maps

$$f_{t,s} = \sum_{\alpha \in A} t^{-\nu(\alpha)} (1 - s\phi_\alpha(z)) z^\alpha.$$

Our goal is to prove the following:

Proposition 4.2 *For large enough t , $f_{t,s}^{-1}(0) = M_{t,s}$ is a family of symplectic hypersurfaces parametrized by s .*




Figure 6: The amoeba of the standard hyperplane in $(\mathbb{C}^*)^2$, its “tropical localization,” and its tropical amoeba

We will call $f_{t,1}$ a “tropical localization” of the Laurent polynomial f . The choice of terminology should be clear from Figure 6.

It follows from the definition of $M_{t,1}$ that near a face σ_t of Π_t , the equations that define M depend only on those coordinates which are orthogonal to σ_t . In order to be more precise, we consider a fixed cover of \mathbb{R}^n .

Definition 4.3 Given any polytope τ of the subdivision of P , define $O_{\check{\tau}}$ to be the set of points $p \in \mathbb{R}^n$ such that $\phi_\alpha(p) \neq 1$ for all $\alpha \in \tau$, and $\phi_\alpha(p) = 1$ for all $\alpha \notin \tau$.

Our choices of functions ϕ_α ensure that

$$\bigcup_{\tau} O_{\check{\tau}} = \mathbb{R}^n.$$

One way to see this is to observe that, although it is not an open set, we can still think of each $O_{\check{\tau}}$ as a neighbourhood of an open subset of $\check{\tau}_t = \log(t)\check{\tau}$, the face of Π_t dual to τ . Indeed, it is not hard to show that a point p that is distance more than $\epsilon \log(t)$ away from $\check{\tau}_t$ must be distance at least $\epsilon \log(t)$ from some $C_{\alpha,t}$ with $\alpha \in \tau$, so $\phi_\alpha(p) = 1$ by construction. This proves that $O_{\check{\tau}}$ is contained in an $\epsilon \log(t)$ neighbourhood of $\check{\tau}_t$.

However, since every face of $\partial\check{\tau}$ is dual to a polytope $\tau' \subset Q$ such that $\tau \subset \partial\tau'$, we know that near every face $\check{\tau}'_t$ of the boundary of $\check{\tau}_t$, there exists at least one vertex $\beta \notin \tau$ such that ϕ_β is not equal to 1. This implies that $\partial\check{\tau}_t$ is not contained in $O_{\check{\tau}}$. But if we restrict to the complement of an $\epsilon \log(t)$ neighbourhood of $\partial\check{\tau}_t$ in $\check{\tau}_t$, then $\phi_\beta = 1$ for every $\beta \notin \tau$.

In addition, note that since we’re working with maximal subdivisions, every polytope τ of the subdivision of P induces a product decomposition

$$(\mathbb{C}^*)^n = (\mathbb{C}^*)^\tau \times (\mathbb{C}^*)^{\check{\tau}}$$

where $\check{\tau}$ is the face of Π which is dual to τ . We therefore have the following description.

Lemma 4.4 *In $O_{\check{\tau}}$, $M_{t,1}$ is a product*

$$M_{t,1} = M_{t,1}^{\check{\tau}} \times U,$$

with $M_{t,1}^{\check{\tau}}$ a hypersurface in $(\mathbb{C}^*)^{\check{\tau}}$ and U a subset of $(\mathbb{C}^*)^{\check{\tau}}$. The set U may itself be decomposed at $\mathbb{T}^{\check{\tau}} \times U'$ with $\mathbb{T}^{\check{\tau}}$ the real subtorus of $(\mathbb{C}^*)^{\check{\tau}}$. \square

Away from its boundary, U' is an open set of $(\mathbb{R}^+)^k$ and may be identified with the open subset of $\check{\tau}$ consisting of points which lie sufficiently far away from the boundary. Furthermore, in the identification of $(\mathbb{C}^*)^n$ with $T\mathbb{R}^n/\mathbb{Z}^n$, the torus $\mathbb{T}^{\check{\tau}}$ corresponds to the tangent space of $\check{\tau}$.

To prove Proposition 4.2, we will use a simple observation which was used by Donaldson in [3]: In order to show that $f^{-1}(0)$ is symplectic it suffices to prove that on the preimage of 0, $|\bar{\partial}f| < |\partial f|$, where the norms are computed with respect to the unique metric determined by the complex and symplectic structures. We compute that

$$\partial f_{t,s} = \sum_{\alpha \in A} t^{-\nu(\alpha)} \partial z^\alpha - s \sum_{\alpha \in A} t^{-\nu(\alpha)} (\phi_\alpha(z) \partial z^\alpha + z^\alpha \partial \phi_\alpha(z))$$

while

$$\bar{\partial} f_{t,s} = s \sum_{\alpha \in A} t^{-\nu(\alpha)} z^\alpha \bar{\partial} \phi_\alpha(z).$$

Since our norm comes from a hermitian metric, and ϕ_α is only a function of the norm of z , it suffices to show that

$$(4-6) \quad \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \partial z^\alpha \right| > s \left(\left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \phi_\alpha(z) \partial z^\alpha \right| + 2 \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} z^\alpha \partial \phi_\alpha(z) \right| \right)$$

hence it suffices to show the above inequality for $s = 1$.

The idea behind our choice of cut-off functions ϕ_α is that ϕ_α is only non-zero whenever the corresponding monomial contributes negligibly to $f_{t,s}$. Formally we have the following:

Lemma 4.5 *If $\text{Log}(z) = p$ is in $C_{\beta,t}$ and $\phi_\alpha(p) \neq 0$ then*

$$\left| \frac{t^{-\nu(\alpha)} z^\alpha}{t^{-\nu(\beta)} z^\beta} \right| < e^{-c\epsilon \log(t)|\alpha-\beta|}.$$

Proof Consider the shortest segment with one endpoint at p and the other at q , the closest point in $\mathcal{H}(\alpha, \beta)$. We have

$$\langle q, \beta \rangle - \log(t)v(\beta) = \langle q, \alpha \rangle - \log(t)v(\alpha).$$

We compute

$$\begin{aligned} \left| \frac{t^{-v(\alpha)} z^\alpha}{t^{-v(\beta)} z^\beta} \right| &= \frac{t^{-v(\alpha)} |e^{\langle p, \alpha \rangle}|}{t^{-v(\beta)} |e^{\langle p, \beta \rangle}|} \\ &= e^{\langle p, \alpha - \beta \rangle + \log(t)(-v(\alpha) + v(\beta))} \\ &= e^{\langle q, \alpha - \beta \rangle + \log(t)(-v(\alpha) + v(\beta))} e^{\langle p - q, \alpha - \beta \rangle} \\ &= e^{\langle p - q, \alpha - \beta \rangle}. \end{aligned}$$

Since the segment from p to q is normal to σ_t , the vector from $p - q$ is parallel to $\alpha - \beta$, so

$$\begin{aligned} \langle p - q, \alpha - \beta \rangle &= -|p - q||\alpha - \beta| \\ &\leq -c\epsilon \log(t)|\alpha - \beta| \end{aligned}$$

where the last inequality follows from $\phi_\alpha(p) \neq 0$, and Lemma 4.1. □

We will only be considering points at which $f_{t,s}$ vanishes. The next two lemmata establish constraints that such points must satisfy.

Lemma 4.6 *If $\mathcal{A}_{t,s}$ is the amoeba of the hypersurface $M_{t,s}$, then $d(\mathcal{A}_{t,s}, \Pi_t) < \log(t)\epsilon$ for all s .*

Proof Note that if $d(p, \Pi_t) \geq \log(t)\epsilon$, then there exists a unique monomial $t^{-v(\beta)} z^\beta$ such that $\phi_\beta = 0$. By Lemma 4.5 and Condition (4-2), we have

$$\begin{aligned} \left| (1 - s\phi_\alpha(z))t^{-v(\alpha)} z^\alpha \right| &\leq \left| t^{-v(\alpha)} z^\alpha \right| \\ &< \frac{1}{5|A|^2 \rho N} \left| t^{-v(\beta)} z^\beta \right| \\ &< \frac{1}{|A|} \left| t^{-v(\beta)} z^\beta \right| \end{aligned}$$

for every $\alpha \neq \beta$. Since there are exactly $|A|$ such terms in our expression for $f_{t,s}$ for which $\alpha \neq \beta$, we conclude that $f_{t,s}(z)$ cannot vanish. □

Lemma 4.7 *If s and z are such that $f_{t,s}(z) = 0$, and $\text{Log}(z) \in C_\beta$, then*

$$|f_{t,r}(z)| < \frac{|t^{-\nu(\beta)} z^\beta|}{5|A|\rho N}$$

for every $0 \leq r \leq 1$. The same bound holds for:

$$\left| \sum_{\phi_\alpha(z) \neq 1} t^{-\nu(\alpha)} z^\alpha \right|$$

Proof It suffices to observe that:

$$\begin{aligned} |f_{t,r}(z)| &= |f_{t,r}(z) - f_{t,s}(z)| \\ &= \left| (r-s) \sum_{\phi_\alpha(z) \neq 0} t^{-\nu(\alpha)} \phi_\alpha(z) z^\alpha \right| \\ &\leq \sum_{\phi_\alpha(z) \neq 0} |t^{-\nu(\alpha)} z^\alpha| \\ &\leq |A| e^{-c \in \log(t)} |t^{-\nu(\beta)} z^\beta| \\ &\leq \frac{|t^{-\nu(\beta)} z^\beta|}{5\rho |A| N} \end{aligned}$$

The same computation yields the second part of the lemma:

$$\begin{aligned} \left| \sum_{\phi_\alpha(z) \neq 1} t^{-\nu(\alpha)} z^\alpha \right| &= \left| s \sum_{0 < \phi_\alpha(z) < 1} \phi_\alpha(z) t^{-\nu(\alpha)} z^\alpha - \sum_{\phi_\alpha(z) = 1} t^{-\nu(\alpha)} (1 - s\phi_\alpha(z)) z^\alpha \right| \\ &\leq \sum_{\phi_\alpha(z) \neq 0} |t^{-\nu(\alpha)} z^\alpha| \\ &\leq \frac{|t^{-\nu(\beta)} z^\beta|}{5\rho |A| N} \end{aligned}$$

□

We now complete the proof of this section’s main result.

Proof of Proposition 4.2 Assume that $f_{t,s}(z) = 0$, and $\text{Log}(z) \in O_z$. By Lemma 4.6, τ consists of at least two vectors. Since the subdivision induced by ν is a triangulation, each subpolytope is in fact a simplex. We choose δ in τ such that $\text{Log}(z) \in C_{\delta,t}$,

which implies that $|t^{-\nu(\delta)} z^\delta| \geq |t^{-\nu(\beta)} z^\beta|$ for all β . In addition, choose a vertex γ in τ which differs from δ , and re-write

$$\sum_{\alpha \in A} t^{-\nu(\alpha)} \partial z^\alpha = \sum_{\beta \in \tau} t^{-\nu(\beta)} z^\gamma \partial z^{\beta-\gamma} + \sum_{\beta \in \tau} t^{-\nu(\beta)} z^{\beta-\gamma} \partial z^\gamma + \sum_{\alpha \in A-\tau} t^{-\nu(\alpha)} \partial z^\alpha.$$

In order to establish Equation (4–6), we decompose its left hand-side as above, so it will suffice to prove the inequality:

$$\begin{aligned} \left| z^\gamma \sum_{\beta \in \tau} t^{-\nu(\beta)} \partial z^{\beta-\gamma} \right| &> \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \phi_\alpha(z) \partial z^\alpha \right| + 2 \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \partial \phi_\alpha(z) z^\alpha \right| \\ &+ \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} z^{\beta-\gamma} \partial z^\gamma \right| + \left| \sum_{\alpha \notin \tau} t^{-\nu(\alpha)} \partial z^\alpha \right| \end{aligned}$$

We start by bounding the left hand-side from below.

$$\begin{aligned} \left| z^\gamma \sum_{\beta \in \tau} t^{-\nu(\beta)} \partial z^{\beta-\gamma} \right| &= \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} z^\gamma \sum_{i=1}^n (\beta_i - \gamma_i) z^{\beta-\gamma-e_i} dz_i \right| \\ &= \sum_{i=1}^n \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} (\beta_i - \gamma_i) z^{\beta-e_i} dz_i \right| \\ &= \sum_{i=1}^n \left| \sum_{\beta \in \tau} (\beta_i - \gamma_i) t^{-\nu(\beta)} z^\beta \right| \\ &\geq \frac{|t^{-\nu(\delta)} z^\delta|}{\rho} \end{aligned}$$

To see the last step, we interpret the penultimate line as the norm of a vector in \mathbb{C}^n which is the image of the k -vector $(t^{-\nu(\beta)} z^\beta)_{\beta \neq \gamma}$ under the linear map which takes the standard basis vectors to the vectors $(\beta - \gamma)_{\beta \in \tau}$. The constant ρ was chosen to be a bound for the length distortion of this linear map. It suffices therefore to bound the norm of $(t^{-\nu(\beta)} z^\beta)_{\beta \neq \gamma}$. We have simply used the size of the δ component as the lower bound for the norm of this vector.

We now successively bound all the terms in the right hand-side. We begin with the first term using Lemma 4.5.

$$\begin{aligned} \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \phi_\alpha(z) \partial z^\alpha \right| &\leq \sum_{\phi_\alpha(z) \neq 0} t^{-\nu(\alpha)} \phi_\alpha(z) |\partial z^\alpha| \\ &\leq \sum_{\phi_\alpha(z) \neq 0} t^{-\nu(\alpha)} \sum_{i=1}^n |\alpha_i| |z^\alpha| \\ &\leq N |A| e^{-c\epsilon \log(t)} \left| t^{-\nu(\delta)} z^\delta \right| \\ &< \frac{\left| t^{-\nu(\delta)} z^\delta \right|}{5\rho} \end{aligned}$$

The same bound works for the last term. We use Lemma 4.7 to bound the third term.

$$\begin{aligned} \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} z^{\beta-\gamma} \partial z^\gamma \right| &= \sum_{i=1}^n \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} z^{\beta-\gamma} \gamma_i z^{\gamma-e_i} dz_i \right| \\ &= \sum_{i=1}^n |\gamma_i| \left| \sum_{\beta \in \tau} t^{-\nu(\beta)} z^\beta \right| \\ &< \frac{\left| t^{-\nu(\delta)} z^\delta \right|}{5\rho} \end{aligned}$$

Finally, we bound the second term.

$$\begin{aligned} \left| \sum_{\alpha \in A} t^{-\nu(\alpha)} \partial \phi_\alpha(z) z^\alpha \right| &\leq \sum_{\phi_\alpha(z) \neq 0} \left| t^{-\nu(\alpha)} z^\alpha \right| |\partial \phi_\alpha(z)| \\ &\leq \sum_{\phi_\alpha(z) \neq 0} \left| t^{-\nu(\alpha)} z^\alpha \right| \sum_{i=1}^n \left| \frac{\partial \phi_\alpha(z)}{\partial \log |z_i|} \frac{dz^i}{\bar{z}_i} \right| \\ &= \sum_{\phi_\alpha(z) \neq 0} \left| t^{-\nu(\alpha)} z^\alpha \right| \sum_{i=1}^n \left| \frac{\partial \phi_\alpha(z)}{\partial u_i} \right| \\ &\leq \sum_{\phi_\alpha(z) \neq 0} \left| t^{-\nu(\delta)} z^\delta \right| e^{-c\epsilon \log(t)} \frac{4}{\epsilon \log(t)} \\ &\leq \frac{\left| t^{-\nu(\delta)} z^\delta \right|}{10\rho} \end{aligned}$$

□

Note that we can use the above bounds to conclude the following:

Corollary 4.8 *Under the hypotheses used in this section we have the bound $|\partial f_{t,s}| > \frac{|t^{-\nu(\delta)} z^\delta|}{10\rho}$.* \square

4.2 Moser's argument

Our goal in this section is to prove the following:

Proposition 4.9 *After rescaling the symplectic form, the pairs $((\mathbb{C}^*)^n, M_{t,s})$ are symplectomorphic for all t and all s for which they are defined.*

In the data of a pair (N, M) of symplectic manifolds, we remember the inclusion of M as a symplectic submanifold of N . In particular, a symplectomorphism of pairs is a symplectomorphism between the total spaces which carries the submanifolds to each other. If we fix t , we will define a map

$$\psi_s: ((\mathbb{C}^*)^n, M_{t,0}) \rightarrow ((\mathbb{C}^*)^n, M_{t,s}).$$

We will only prove the result for varying s , as the result for varying t is entirely analogous, although it requires rescaling the symplectic form by $\log(t)$. We begin by showing that the hypersurfaces $M_{t,s}$ are symplectomorphic using the fact that our proof that $M_{t,s}$ is a symplectic hypersurface in fact proves that $f_{t,s}$ is a symplectic fibration in a neighbourhood of each $M_{t,s}$. In particular we obtain a symplectic connection. Let $Y_{t,s}$ be the horizontal lift of $-\frac{\partial f_{t,s}}{\partial s}$. We need to bound $Y_{t,s}$ in order to show that it integrates to a flow. But first, we would like to obtain an explicit formula for it. This is a problem in linear algebra, for which we need a little notation.

Let (V, ω) be a symplectic vector space. The symplectic form ω determines an isomorphism between V and V^* which we denote by $\omega^\#$ and define by the formula

$$\langle \omega^\#(v), w \rangle = \omega(v, w).$$

The inverse isomorphism will be written $\omega_\#$. We will also use the same symbols when we extend these maps i -linearly to the complexification of V and its dual. We equip \mathbb{C} with the standard symplectic form

$$dx \wedge dy = \frac{i}{2} dz \wedge d\bar{z}.$$

Lemma 4.10 *Let f_* be a linear map between the symplectic vector spaces (V, ω) and (\mathbb{C}, ω_0) with symplectic kernel and with dual f^* . The horizontal lift of the vector $a \in \mathbb{C}$ is given by*

$$f^!(a) = \frac{2i}{\omega(f^*dz, f^*d\bar{z})} \omega_{\#}(f^* \omega_0^{\#} a).$$

Proof We will work with the complexification of the vector space throughout. First, note that the image of \mathbb{C} under the composition $\omega_{\#} \circ f^* \circ \omega_0^{\#}$ is precisely the orthogonal complement of the kernel of f_* . Hence it suffices to show that the right hand side of our formula maps to a under the linear map f_* . By linearity, it suffices to check this for $\frac{\partial}{\partial z}$ and $\frac{\partial}{\partial \bar{z}}$, where, keeping in mind our application, we've used a suggestive notation for the standard basis of the complexification of \mathbb{C} . We check it for $\frac{\partial}{\partial z}$ by computing the pairing with dz and $d\bar{z}$.

$$\begin{aligned} \left\langle f_* \left(\frac{2i}{\omega(f^*dz, f^*d\bar{z})} \omega_{\#} f^* \left(\omega_0^{\#} \frac{\partial}{\partial z} \right) \right), dz \right\rangle &= \frac{2i \langle f_* \left(\omega_{\#} f^* \left(\frac{id\bar{z}}{2} \right) \right), dz \rangle}{\omega(f^*dz, f^*d\bar{z})} \\ &= \frac{-\langle \omega_{\#} f^* d\bar{z}, f^* dz \rangle}{\omega(f^*dz, f^*d\bar{z})} \\ &= 1 \\ \left\langle f_* \left(\frac{2i}{\omega(f^*dz, f^*d\bar{z})} \omega_{\#} f^* \left(\omega_0^{\#} \frac{\partial}{\partial z} \right) \right), d\bar{z} \right\rangle &= \frac{\langle \omega_{\#} f^* d\bar{z}, f^* d\bar{z} \rangle}{\omega(f^*dz, f^*d\bar{z})} \\ &= 0. \end{aligned}$$

Of course, the same computation works for $\frac{\partial}{\partial \bar{z}}$. □

Lemma 4.11 *$Y_{t,s}$ is a bounded vector field, hence integrates to a local diffeomorphism near the fibre $M_{t,s}$.*

Proof By the previous lemma, we have the expression

$$f_{t,s}^! \left(\frac{\partial f_{t,s}}{\partial s} \right) = \frac{2i}{\omega(f_{t,s}^* dz, f_{t,s}^* d\bar{z})} \omega_{\#} f_{t,s}^* \left(\omega_0^{\#} \frac{\partial f_{t,s}}{\partial s} \right).$$

Since $f_{t,s}$ is almost holomorphic, this vector field has norm less than or equal to

$$\frac{4 \left| \frac{\partial f_{t,s}}{\partial s} \right| \cdot |f_{t,s}^* dz|}{|f_{t,s}^* dz|^2}.$$

By Corollary 4.8

$$|f_{t,s}^* dz| \geq \frac{|\partial f_{t,s}|}{2} \geq \frac{|t^{-\nu(\delta)} z^{\delta}|}{20\rho}$$

with δ as in the proof of Proposition 4.2. On the other hand

$$\begin{aligned} \left| \frac{\partial f_{t,s}}{\partial s} \right| &= \left| \sum_{\phi_\alpha(z) \neq 0} \phi_\alpha(z) t^{-\nu(\alpha)} z^\alpha \right| \\ &\leq \frac{|t^{-\nu(\delta)} z^\delta|}{5 |A| \rho N} \end{aligned}$$

as in Lemma 4.7. □

Lemma 4.12 *The flow $\bar{\psi}_s$ of $Y_{t,s}$ restricts to a symplectomorphism between $M_{t,0}$ and $M_{t,s}$.*

Proof First, we observe that by construction of $Y_{t,s}$

$$\begin{aligned} \frac{\partial}{\partial s} f_{t,s} (\bar{\psi}_s(z)) &= f_{t,s*} (Y_{t,s}|_{\bar{\psi}_s(z)}) + \frac{\partial f_{t,s}}{\partial s} \Big|_{\bar{\psi}_s(z)} \\ &= 0. \end{aligned}$$

So the flow of $Y_{t,s}$ preserves the level sets of the function $f_{t,s}$, and hence maps the zero fibres to each other. To check that this is, indeed, a symplectomorphism, it suffices to compute

$$\begin{aligned} \frac{\partial}{\partial s} \bar{\psi}_s^* (\omega_{M_{t,s}}) &= \frac{\partial}{\partial s} \bar{\psi}_s^* (\omega|_{M_{t,s}}) \\ &= \bar{\psi}_s^* (\mathcal{L}_{Y_{t,s}} \omega|_{M_{t,s}}) \\ &= \bar{\psi}_s^* (d(i_{Y_{t,s}} \omega)|_{M_{t,s}}) \\ &= 0 \end{aligned}$$

using Cartan’s formula, the fact that ω is closed, and that $Y_{t,s}$ is orthogonal to $M_{t,s}$. In particular, $i_{Y_{t,s}} \omega$ vanishes on $M_{t,s}$ and hence also its exterior derivative. □

We must now check whether this symplectomorphism can be embedded in an appropriate symplectomorphism of pairs. Similar results are well known in the compact case [16] and the proof extends to the open case provided one can bound the vector fields.

Lemma 4.13 *There exists a Hamiltonian time-dependent vector field $Y'_{t,s}$ on $(\mathbb{C}^*)^n$ which is supported in a neighbourhood of $M_{t,s}$ and which integrates to a symplectic flow ψ_s that maps $M_{t,0}$ to $M_{t,s}$.*

Proof There exists a function H_s which vanishes on $M_{t,s}$, and such that $dH_s|_{M_{t,s}} = i_{Y_{t,s}}\omega$. This can be seen locally by taking any trivialization of a neighbourhood of $M_{t,s}$ and thinking of $i_{Y_{t,s}}\omega$ as an initial value problem for a function, then using a partition of unity to produce a global function. We can always choose H_s to be supported in a neighbourhood of $M_{t,s}$. The form dH_s gives the closed form which corresponds to our vector field $Y'_{t,s}$. Since our initial values are bounded, an appropriate choice of cutoff function ensures that the vector field $Y'_{t,s}$ is also bounded, hence integrates to a Hamiltonian isotopy. \square

Remark 4.14 Note that we can choose the germs of the vector fields $Y'_{t,s}$ and $Y_{t,s}$ to be C^∞ close at $M_{t,s}$. In particular, this means we can assume that ψ_s respects the fibration to arbitrarily high order at $M_{t,s}$.

Note that if J is any compatible complex structure J on V such that f_* is a map of complex vector spaces, then the symplectic horizontal lift agrees with the orthogonal lift which for a unit length vector v is given by

$$f^!(v) = \frac{1}{g(f^*g_0^\#v, f^*g_0^\#v)} g_\# f^* g_0^\# v.$$

In particular, if f_* is the derivative at the origin of a symplectic fibration $f: N \rightarrow \mathbb{C}$, we consider the real valued function $f_{\mathbb{R}}$, which is the restriction of f to the inverse image of the real axis. Since $df_{\mathbb{R}} = f^*g_0^\# \frac{\partial}{\partial x}$, we conclude

$$\text{grad } f_{\mathbb{R}} = |df_{\mathbb{R}}|^2 f^! \frac{\partial}{\partial x}.$$

In future section, we will let Λ be a compact Lagrangian submanifold of the zero fibre of f , which we denote M , and consider parallel transports of Λ along straight half-rays originating at the origin, and moving in the left half-plane. We write such a line as

$$\gamma_c(t) = -t(1 + 2ci).$$

Denote the parallel transport along this line by L_c , and the distance to Λ by r_Λ . On L_0 , the gradient of r_Λ and the lift of $\frac{\partial}{\partial x}$ are both orthogonal to Λ , and satisfy

$$\text{grad } r_\Lambda = |df_{\mathbb{R}}| f^! \frac{\partial}{\partial x}.$$

Lemma 4.15 *Let J be an almost complex structure in a neighbourhood of M for which f is holomorphic and M is an almost complex submanifold. If we identify a neighbourhood of L_0 with its cotangent bundle as in Lemma 2.1, then L_c is given as the graph of a 1-form which agrees with the differential of $cr_\Lambda^2(u)$ to first order in u .*

Proof Let L'_c be the graph of the differential of cr_Λ^2 . Since the differential of $cr_\Lambda^2(u)$ vanishes on Λ , L_c and L'_c have the same boundary. To show that the 1-forms that define them agree to first order in u , it suffices to show that their tangent spaces agree on the boundary. However, TN_p decomposes as an orthogonal direct sum $TM \oplus \mathbb{C}$, and it is clear that

$$TL_c = T\Lambda \oplus \mathbb{R} \cdot f^! \left(\frac{\partial}{\partial x} + 2c \frac{\partial}{\partial y} \right);$$

so it remains to compute TL'_c . To do this, we observe that r_Λ vanishes on Λ , and that since TM is closed under J , and J is compatible with ω , its orthogonal complement is also closed under J . We may therefore ignore the directions corresponding to $T\Lambda$ in our computation. Now the orthogonal complement of $T\Lambda$ in TL_0 is spanned by $f^! \frac{\partial}{\partial x}$, so the component of the tangent space of TL'_c which lies in the orthogonal complement to $T\Lambda$ is spanned by

$$(4-7) \quad f^! \frac{\partial}{\partial x} + \nabla_{f^! \frac{\partial}{\partial x}} (2cr_\Lambda(u)J \operatorname{grad} r_\Lambda(u)).$$

Consider the vector field

$$Z = J \left(\operatorname{grad} r_\Lambda - |df_{\mathbb{R}}| f^! \frac{\partial}{\partial x} \right).$$

We know that this vector field vanishes at Λ . In addition, we know that the vector field $Jf^! \frac{\partial}{\partial x} = f^! \frac{\partial}{\partial y}$ is the lift of a geodesic flow in the base. We conclude that on Λ the second term can be computed as follows:

$$\begin{aligned} \nabla_{f^! \frac{\partial}{\partial x}} (r_\Lambda(u)J \operatorname{grad} r_\Lambda(u)) &= \nabla_{f^! \frac{\partial}{\partial x}} \left(r_\Lambda(u)|df_{\mathbb{R}}|Jf^! \frac{\partial}{\partial x} + r_\Lambda(u)Z \right) \\ &= \nabla_{f^! \frac{\partial}{\partial x}} \left(r_\Lambda(u)|df_{\mathbb{R}}|f^! \frac{\partial}{\partial y} \right) + r_\Lambda(u)\nabla_{f^! \frac{\partial}{\partial x}} Z \\ &\quad + f^! \frac{\partial}{\partial x} (r_\Lambda(u))Z \end{aligned}$$

Since r_Λ and Z vanish on Λ , it remains to compute that:

$$\begin{aligned} \nabla_{f^! \frac{\partial}{\partial x}} \left(r_\Lambda(u)|df_{\mathbb{R}}|f^! \frac{\partial}{\partial y} \right) &= f^! \frac{\partial}{\partial x} (|df_{\mathbb{R}}|r_\Lambda(u)) \frac{\partial}{\partial y} \\ &= \left(r_\Lambda(u)f^! \frac{\partial}{\partial x} (|df_{\mathbb{R}}|) + |df_{\mathbb{R}}|f^! \frac{\partial}{\partial x} (r_\Lambda(u)) \right) f^! \frac{\partial}{\partial y} \\ &= \operatorname{grad} r_\Lambda(u)(r_\Lambda(u))f^! \frac{\partial}{\partial y} \\ &= f^! \frac{\partial}{\partial y} \end{aligned}$$

In the last line, we've used the fact that the restriction gradient of r_Λ to Λ has unit norm. Plugging this back in Equation (4–7) completes the proof of the lemma. \square

4.3 The zero-section

In this section, we construct an admissible Lagrangian submanifold of $(\mathbb{C}^*)^n$ with boundary on $M_{t,1}$, which will be the first step of future constructions. The construction can be done for any Laurent polynomial f . However, we will specialize to the situation where we're considering the mirror W of a smooth toric variety. In particular,

$$W_{t,1} = -1 + \sum_{0 \neq \alpha \in A} t^{-\nu(\alpha)} (1 - \phi_\alpha(z)) z^\alpha.$$

Consider the $\epsilon \log(t)$ neighbourhood of the $n - 2$ skeleton of the polytope $Q_t = C_{0,t}$. The intersection of the amoeba of $W_{t,1}$ with a neighbourhood of Q_t agrees with the boundary of Q_t outside this neighbourhood of the $n - 2$ skeleton.

Recall that by thinking of $(\mathbb{C}^*)^n$ as the cotangent bundle of \mathbb{R}^n modulo the lattice \mathbb{Z}^n , we identified a natural zero-section. One might equivalently think of the inclusion of $(\mathbb{R}^+)^n$ into $(\mathbb{C}^*)^n$.

Lemma 4.16 *Near the component of the complement of the amoeba corresponding to the origin, the zero section intersects $M_{t,1}$ in a smooth Lagrangian sphere of $M_{t,1}$. This Lagrangian sphere is the boundary of a compact subset of the zero section which is diffeomorphic to the n -ball.*

Remark 4.17 It is easy to see that the zero section does intersect the zero level set of the linear function

$$-1 + \sum_{i=1}^n t^{-\nu(i)} z_i$$

in a smooth Lagrangian submanifold. Hence, by the multiplicative change of coordinates, the same holds for

$$g(z) = -1 + \sum_{0 \neq \alpha \in \tau} t^{-\nu(\alpha)} z^\alpha$$

whenever τ is a minimal simplex.

Proof We use the polyhedral decomposition provided by the tropical limit, and prove the result for $M_{t,1}$. Consider a point z such that $\text{Log}(z) \in O_\tau$. If τ is a k -simplex, we know that all but $k + 1$ terms in

$$W_{t,1} = -1 + \sum_{0 \neq \alpha \in A} t^{-\nu(\alpha)} (1 - \phi_\alpha(z)) z^\alpha$$

are zero. It suffices therefore to check the result for a simplex as in the previous remark. So consider

$$g_{k,s}(z) = -1 + \sum_{0 \neq \alpha \in \tau} t^{-\nu(\alpha)} (1 - s\phi_\alpha(z))z^\alpha$$

which we think of as a 1-parameter family of functions defined on a domain where $g_{k,1}$ agrees with $W_{t,1}$. By the previous remark, it will be sufficient to show that reality is preserved by our flow, which reduces to showing that the vector field which defines the flow is tangent to $(\mathbb{R}^+)^n$ on the intersection of $(\mathbb{R}^+)^n$ with the zero level set. Using the result of Lemma 4.10, and the expression of the symplectic form on $(\mathbb{C}^*)^n$ in standard coordinates, we are reduced to showing that

$$g_{k,s}^*(\omega_0^\# \frac{\partial g_{k,s}}{\partial s}) = \sum_{i=1}^n h_i(z) dy_i$$

for some real functions h_i . Since $g_{k,s}$ restricts to a real function on the intersection of $(\mathbb{R}^+)^n$ with the zero level set, the restriction of $\frac{\partial g_{k,s}}{\partial s}$ to this intersection is some real multiple of $\frac{\partial}{\partial x}$, hence $\omega_0^\# \frac{\partial g_{k,s}}{\partial s}$ is some real multiple of dy . The result would therefore follow from showing that for all $1 \leq i \leq n$, the following derivative of the imaginary part is zero:

$$\frac{\partial \text{Im}(g_{k,s})}{\partial x_i} = 0$$

But in fact, the function $\text{Im}(g_{k,s})$ itself vanishes on the real locus, so this establishes that the flow maps the positive real locus of $g_{k,0}$ into the positive real locus of $g_{k,1}$. Since we can also use the reverse flow, this map gives a diffeomorphism between the two sets.

To see that the result is simply an n -ball with boundary, we note that the boundary of the component of the complement of the amoeba corresponding to the origin is a polytope which is itself homeomorphic to a sphere which bounds \mathcal{L}_∞ . It is clear from the local model that \mathcal{L}_∞ is star-shaped about the origin, so it must be diffeomorphic to the standard ball. □

We will denote this Lagrangian ball with boundary on $M_{t,1}$ by $\log(t)\mathcal{L}$.

Remark 4.18 Note that the results of the previous section allow us to conclude that any construction which is performed for admissible Lagrangians with respect to $W_{t,1}$ can be “pulled back” to the original complex hypersurface M . Hereafter, we will change our notation and write M for the image of $M_{t,1}$ under the diffeomorphism of $(\mathbb{C}^*)^n$ induced by the diffeomorphism of the base

$$u \rightarrow \frac{u}{\log(t)}.$$

This induces the rescaling of the symplectic form which appears in the statement of Proposition 4.9. But rescaling preserves symplectic and Lagrangian submanifolds, so our constructions are insensitive to it. In particular, we produce a Lagrangian ball \mathcal{L} with boundary on M .

We will also use W for the pull back of $W_{t,1}$ under the above “conformal” symplectomorphism. We will still perform most computations using the coordinates z_i on $(\mathbb{C}^*)^n$ as a subset of \mathbb{C}^n . Strictly speaking, these differ by a “rescaling” from the coordinates used in the previous section. These coordinates come with a natural complex structure which will use.

There are two reasons that allow us to use this integrable almost complex structure. The first reason is that we proved in Section 3.5 that the holomorphic triangles that we will be considering are already regular. This means that there is no issue in the interior. More problematic is the fact that we will also be considering the distance function with respect to the usual metric in order to approximate parallel transport as in Lemma 4.15. However, it is easy to see from the remark following Lemma 4.13 that we can choose the usual complex structure to be C^1 -close near M to an admissible almost complex structure J . This implies that the distance function with respect to the usual metric yields a C^1 -close approximation to parallel transport with respect to J , which is all we will need.

5 The admissible Lagrangians $\mathcal{L}(j)$

5.1 Desiderata

Consider the symplectic hypersurface M with amoeba \mathcal{A} . By Lemma 4.6, \mathcal{A} is ϵ -close to the polyhedral complex Π . If we focus only on Q , we know that the amoeba agrees with the polyhedral complex away from an ϵ neighbourhood of its $n - 2$ skeleton. In the next two sections, we will construct a Hamiltonian function H which should be thought of as a modification of H_∞ and whose time-1 flow preserves M . But first, we would like to describe the properties that H must satisfy, and show that all possible choices of H , up to admissible Hamiltonian isotopy, are equivalent.

Continuing with the change in notation introduced in the last section, we have a cover O_σ of \mathbb{R}^n indexed by the faces σ of Π such that each O_σ is a neighbourhood of a large open subset of σ . Recall that we have been using the Euclidean metric on \mathbb{R}^n to identify each cotangent fibre with a copy of \mathbb{R}^n ; let π denote this cotangent fibration.

Definition 5.1 A Lagrangian submanifold Λ of M is a *boundary for $\mathcal{L}(1)$* if Λ lifts to a submanifold $\tilde{\Lambda} \subset T\mathbb{R}^n$ such the following properties are satisfied:

- $\tilde{\Lambda}$ is a section of $T\mathbb{R}^n|_{\partial\mathcal{L}}$, and
- if $z \in \tilde{\Lambda}$ and $\pi(z) \in \partial\mathcal{L} \cap O_\sigma$ then z lies in the affine subspace supporting $-2\pi\sigma$, and
- $\tilde{\Lambda}$ is the restriction to \mathcal{L} of an exact section of $T^*\mathbb{R}^n$.

Note that the last condition is vacuous if $n > 2$ since $H^1(\partial\mathcal{L})$ vanishes in this case.

Lemma 5.2 *Given Λ , a boundary for $\mathcal{L}(1)$, and a vector v in the left half plane, there exists an admissible Lagrangian $\mathcal{L}(1)$ which is an exact section of $T\mathbb{R}^n|_{\mathcal{L}}$ with boundary equal to Λ and whose image under W agrees near the origin with a curve γ whose tangent vector at the origin is v . Furthermore, any two such Lagrangians, for (possibly) different tangent vectors at the origin, are isotopic through admissible Lagrangians.*

Proof To prove the first statement, we prove that the parallel transport of Λ along a short curve γ with tangent vector v at the origin yields a section of $T\mathbb{R}^n|_U$ where U is a 1-sided neighbourhood of $\partial\mathcal{L}$ in \mathcal{L} . As usual, we consider the problem on each element of the cover O_σ . Note that the condition that $\tilde{\Lambda}$ lies in the hyperplane that supports $-2\pi\sigma$ implies that $\tilde{\Lambda}$ is locally the product of a section of $T\sigma$ with the zero section of a transverse symplectic slice. By Lemma 4.4 we know that in O_σ the function W is independent of the directions corresponding to σ , so parallel transport only depends on the directions which lie in the transverse slice, which we may locally identify with a lower dimensional torus.

Claim If \mathcal{L}_v is the parallel transport of the zero-section along a sufficiently short curve γ starting at 0 with tangent vector v in the left half-plane, then π projects \mathcal{L}_v homeomorphically onto a neighbourhood of the boundary of \mathcal{L} .

Proof of Claim In the proof of Lemma 4.16 we noticed that $W_{t,1}$ takes negative real values on \mathcal{L} . In particular, this implies that \mathcal{L} is the parallel transport of its boundary along the negative real axis. The result now follows immediately from the proof of Lemma 4.15. \square

Constructing a Lagrangian $\mathcal{L}_v(1)$ as the parallel transport of Λ along such a curve γ yields locally a section $T\mathbb{R}^n|_{\mathcal{L}}$. Our third condition on Λ guarantees that this section is exact, hence any extension of the function that defines it will yield the desired section of $T^*\mathbb{R}^n|_{\mathcal{L}}$.

To prove uniqueness, we observe that any Lagrangian $\mathcal{L}(1)$ is the graph of the differential of a function H with prescribed derivatives at the boundary. The linear interpolation

between two such functions H and H' defines an isotopy which may not preserve admissibility. But choosing a sufficiently fine cover of the interval $[0, 1]$ as in Lemma 2.13 allows us to modify this isotopy to one in which admissibility is preserved. \square

This means that we can unambiguously define $\mathcal{L}(1)_\Lambda$ to be an admissible exact section of $T^*\mathbb{R}^n|_{\mathcal{L}}$ whose boundary is Λ . We would like to eliminate the dependence on Λ . Recall from Lemma 4.4 that W is independent of the directions corresponding to the tangent space of σ . In particular, the proof of the previous lemma together with the convexity of the faces of σ imply that we can use interpolation as in Lemma 2.13 to prove the following:

Lemma 5.3 *If Λ and Λ' are boundaries for $\mathcal{O}(1)$, then $\mathcal{L}(1)_\Lambda$ and $\mathcal{L}(1)_{\Lambda'}$ are isotopic through admissible Lagrangians.* \square

So we can define admissible Lagrangian submanifolds $\mathcal{L}(1)$ which are well defined up to Hamiltonian isotopy. In the next sections, we will produce an explicit model for $\mathcal{L}(1)$ that will allow us to compute its Floer cohomology groups. We also note that this construction naturally extends to the construction of Lagrangians $\mathcal{L}(j)$, for any integer j , whose lifts lie in $-2\pi j\sigma$ in every set O_σ .

5.2 A boundary for $\mathcal{L}(1)$

In this section, we will construct a boundary for $\mathcal{L}(1)$. First, we must choose an appropriate cover of a neighbourhood of Π . Given σ a facet of Π , Let U_σ be its ϵ -neighbourhood and V_σ its 2ϵ neighbourhood. If we then consider the $n-2$ skeleton, there exists a constant, c_{n-2} such that every point in the 2ϵ neighbourhood of Π , but which is distance more than $c_{n-2}\epsilon$ from the $n-2$ skeleton is closest to a unique facet of Π . The value of c_{n-2} is independent of ϵ , and is related to the ‘‘angles’’ between the facets near their intersections. Given a face σ in the $n-2$ skeleton, let U_σ be the $c_{n-2}\epsilon$ neighbourhood of σ , and V_σ the $2c_{n-2}\epsilon$ neighbourhood of σ .

Repeating this process inductively, we obtain constants $c_i \geq 1$ for each $i \leq n-1$ independent of ϵ , and define U_σ to be the $c_i\epsilon$ neighbourhood of σ , and V_σ its $2c_i\epsilon$ neighbourhood for $i = \dim(\sigma)$. Let

$$U_i = \bigcup_{\dim(\sigma)=i} U_\sigma$$

$$V_i = \bigcup_{\dim(\sigma)=i} V_\sigma.$$

The main property satisfied by this cover is the following:




Figure 7: The sets U_0 , V_0 , U_1 and V_1 for the hyperplane in $(\mathbb{C}^*)^2$ shown in progressively darker shades of gray

Lemma 5.4 *Every point of $V_i - U_{i-1}$ is closest to a unique face of dimension i . \square*

In the interior of each i -face σ , we identify the normal bundle of σ with a neighbourhood, and define a self map ψ_σ of this bundle which is radial, collapses U_σ to σ , and is the identity away from V_σ . We would like to take the composition of all ψ_σ over the $n-2$ skeleton, to obtain a self map of \mathbb{R}^n . There are two issues with this:

- ψ_σ does not extend continuously to \mathbb{R}^n because σ has non-empty boundary (if $\dim(\sigma) \neq 0$).
- The supports of ψ_σ for different faces are not disjoint.

We resolve both of these issues by defining ψ_i to be the composition of ψ_σ for σ a face of dimension i in an arbitrary order, and defining ψ to be the composition

$$\psi = \psi_0 \circ \psi_1 \circ \cdots \circ \psi_{n-1}.$$

The key point is that the set where ψ_i is not continuous, or depends on the order chosen, lies entirely within U_{i-1} , which is collapsed by the next map (ψ_{i-1}) to the $i-1$ skeleton. It should therefore be clear from the construction that ψ is a well defined smooth self map of \mathbb{R}^n which does not depend on the order of compositions chosen for ψ_σ . Further, we have the following:

Lemma 5.5 *The map ψ can be chosen to satisfy the following additional properties:*

- ψ preserves the i -skeleton of Π . In particular, it maps every face to itself.
- ψ maps every point $p \in U_i$ to the nearest i -face of Π . If $p \in U_i - V_{i-1}$ then $\psi(p)$ is the nearest point on the nearest i -face.
- ψ maps the amoeba \mathcal{A} to the tropical amoeba Π .

- ψ can be chosen to satisfy the condition that its derivative is bounded independently of ϵ . Specifically, given any tangent vector v at u ,

$$|\psi_*(u)(v)| < 4^n |v|.$$

Proof The first two properties are immediate from the construction, so we only prove the other two. Note that we have $O_\sigma \subset U_\sigma$. In particular, since the localization of the tropical amoeba lies entirely in $\cup_\sigma O_\sigma$, ψ maps the localization of the tropical amoeba to the tropical amoeba. Finally, since each ψ_σ collapses a neighbourhood of radius $c_i\epsilon$, and is the identity outside a neighbourhood of radius $2c_i\epsilon$, its differential can be chosen to expand lengths of vectors by a factor less than 4. The result follows by observing that in a neighbourhood of any point p , ψ can be written as the composition of at most n maps corresponding to the faces of each dimension which are closest to p . \square

Consider the Hamiltonian function given by

$$H^\partial(u) = -2\pi \left\langle u - \frac{\psi(u)}{2}, \psi(u) \right\rangle - \pi a \langle \psi(u), \psi(u) \rangle$$

on the set $\mathcal{L} \cap \cup_i U_i$, and define Λ to be the graph of $dH^\partial|_{\partial\mathcal{L}}$. To keep the notation simpler, we do not make the dependence on a explicit.

Lemma 5.6 Λ is a boundary for $\mathcal{L}(1)$.

Proof Let us first set $a = 0$. We compute that for any tangent vector v ,

$$\begin{aligned} dH^\partial(u)(v) &= -2\pi \left\langle v - \frac{\psi_*(u)(v)}{2}, \psi(u) \right\rangle - 2\pi \left\langle u - \frac{\psi(u)}{2}, \psi_*(u)(v) \right\rangle \\ &= \langle v, -2\pi \psi(u) \rangle - 2\pi \langle u - \psi(u), \psi_*(u)(v) \rangle. \end{aligned}$$

The only condition we need to check is that if $u \in O_\sigma$, this co-vector lies in the hyperplane supporting $-2\pi\sigma$ after passing to the tangent space. By the previous lemma, $\psi(u)$ lies in σ , so this is already true of the first term. We must prove that the second term lies in the tangent space of σ . In particular, it suffices to show that the corresponding 1-form annihilates every tangent vector normal to σ . But in U_σ , ψ_σ collapses the directions orthogonal to σ so this is indeed true.

To address the case $a \neq 0$, we observe that the term depending on a has differential

$$v \mapsto -2\pi a \langle \psi_*(u)(v), \psi(u) \rangle$$

which, again, corresponds to a term in the tangent space of σ . \square

5.3 Parallel transport of Λ

As in the proof of Lemma 5.2, we begin the construction of $\mathcal{L}(1)$ by parallel transport of the boundary along a curve $\gamma \subset \mathbb{C}$ starting at the origin and moving in the left half plane. We proved that this parallel transport can be expressed locally as an exact section over \mathcal{L} . Recall that in order to compute

$$HF^*(\mathcal{L}, \mathcal{L}(1))$$

we need to ensure that $W(\mathcal{L}(1))$ agrees with a curve γ near the origin, and that the angle between the negative x -axis and γ is between 0 and $\frac{\pi}{2}$. In particular, we would be considering $c > 0$ in the notation of Lemma 4.15. Let us write $\mathcal{L}_c(1)$ for a Lagrangian that satisfies this condition of being a parallel transport of Λ along a line of slope $2c$ at the origin.

Lemma 5.7 *The graph of dH^∂ agrees with $\mathcal{L}_0(1)$ near Λ .*

Proof We only need to prove that the image under W of the graph of dH^∂ near ∂L is the negative real axis. By construction, dH^∂ lies in the hyperplane supporting $-2\pi\sigma$, so we may think of it as the product of the zero section of $(\mathbb{C}^*)^{\check{\sigma}}$ with a non-trivial section of $T\sigma$. Since in O_σ , W only depends on the directions transverse to σ , it suffices to prove that the zero-section \mathcal{L} is the parallel transport of its restriction to the boundary along the negative real axis. We already observed this in the proof of Lemma 5.2. \square

The graph of the differential of

$$H_c^\partial(u) = -2\pi \left(\left\langle u - \frac{\psi(u)}{2}, \psi(u) \right\rangle + \frac{a}{2} \langle \psi(u), \psi(u) \rangle \right) + cr_\Lambda^2(u)$$

is therefore C^1 -close to an admissible Lagrangian by Lemma 4.15. In particular, using Lemma 2.2, we can construct an admissible Lagrangian $\mathcal{L}_c(1)$ that is C^1 -close to this graph near Λ , and that agrees with it away from an arbitrarily small neighbourhood of Λ . In practice, this means that we can think of the graph of dH_c^∂ as being admissible.

We will interpolate between $\mathcal{L}_c(1)$ and the linear Lagrangians $\mathcal{L}_\infty(1)$ constructed in Section 3.3. We will only do the harder case where $c > 0$ which corresponds to choosing a model of $\mathcal{L}(1)$ such that \mathcal{L} and $\mathcal{L}(1)$ are a positively oriented pair.

Let δ be an arbitrarily small positive number. We assume that the origin is contained in the interior of Q . Choose $\epsilon' > 0$ such that $(1 + \epsilon')Q$ is contained in a 2δ neighbourhood of Q . Note that $(1 + \epsilon')\partial Q$ lies entirely outside of Q . Further, given any face σ of ∂Q , $(1 + \epsilon')\sigma$ lies in the interior of $2\sigma - Q$. Choose $\epsilon > 0$ and go through the

constructions of the previous sections so that $(1 + \epsilon')U_\sigma$ also lies in the interior of $2\sigma - Q$. Note that we can choose $\epsilon < \epsilon' < \delta$.

There exists a number $0 < b < \frac{1}{2}$ such that r_Λ has no critical points in the $b\epsilon$ neighbourhood of Λ . Let g be a non-negative cutoff function satisfying the following conditions:

$$\begin{aligned} g(u) = 0 &\Leftrightarrow r_\Lambda(u) \leq b\epsilon \\ g(u) = 1 &\Leftrightarrow r_\Lambda(u) \geq \epsilon \\ |dg(u)| &\leq \frac{2}{\epsilon} \end{aligned}$$

Consider the Hamiltonian H_c given by the formula

$$\begin{aligned} &-(1 - g(u)) \left(\pi \langle 2u - \psi(u), \psi(u) \rangle + (\epsilon' + 5\epsilon\epsilon')\pi \langle \psi(u), \psi(u) \rangle - cr_\Lambda^2(u) \right) \\ &- g(u)\pi(1 + \epsilon')\langle u, u \rangle. \end{aligned}$$

Note that for $r_\Lambda(u) < b\epsilon$, this function is just H_c^∂ for $a = \epsilon' + 5\epsilon\epsilon'$. In particular, the corresponding Lagrangian is C^1 -close to $\mathcal{L}_c(1)$. Since the images of $\mathcal{L}_c(1)$ and \mathcal{L} under W do not intersect, it is clear that the Lagrangian corresponding to H_c does not intersect \mathcal{L} in this region. Let us abuse notation and write $\mathcal{L}_c(1)$ for the graph of the function H_c . When $r_\Lambda(u) \geq \epsilon$, the Hamiltonian is

$$-\pi(1 + \epsilon')\langle u, u \rangle$$

and the intersection points between \mathcal{L} and $\mathcal{L}(1)_c$ in this region are exactly the $\frac{1}{1+\epsilon'}$ integral points of Q .

Lemma 5.8 *dH_c takes no 2π -integral values in the region where $b\epsilon \leq r_\Lambda(u) \leq \epsilon$.*

Proof Since it suffices to prove non-integrality of some coefficient, we use the cover O_σ and consider only the factor transverse to σ . In other words we may assume, after possibly reducing to a factor, that u lies in the ϵ neighbourhood of a vertex u_0 of Q . Note that this means that $\psi(u) = u_0$, and that the differential of ψ vanishes. We compute that in this case, dH_c equals

$$\begin{aligned} &-2\pi u_0 + g(u) (2\pi u_0 - 2\pi(1 + \epsilon')u) + (1 - g(u))2cr_\Lambda(u)dr_\Lambda(u) \\ &+ \left(\pi \langle 2u - u_0, u_0 \rangle + (\epsilon' + 5\epsilon\epsilon')\pi \langle u_0, u_0 \rangle - cr_\Lambda^2(u) - (1 + \epsilon')\pi \langle u, u \rangle \right) dg(u). \end{aligned}$$

Since $-2\pi u_0$ is an integral vector it suffices to show that the remaining terms do not add up to 0, and have norm much less than 1. The main observation is the following:

Claim The inner product of any of the vectors dg , dr_Λ or $u_0 - (1 + \epsilon')u$ with u_0 is non-positive.

Proof For the last vector, this follows from the choices made about ϵ' and ϵ at the beginning of this section. In particular, we can choose ϵ much smaller than ϵ' , so this vector is well approximated by $-\epsilon'u_0$. Since dg can be chosen approximately parallel to dr_Λ , we will simply prove the claim for this last vector. But the gradient of dr_Λ is a scalar multiple of the normal vector at the nearest point in Λ . On the other hand, it is easy to compute that the negative of the gradient of the Laurent polynomial

$$W_t(z) = -1 + \sum_t t^{-\nu(\alpha)} z^\alpha$$

satisfies this condition on Λ (this is simply the statement that W decreases as we move towards the interior of \mathcal{L}). Since the cutoff functions were chosen C^1 -small with respect to W_t , this result is unaffected by the presence of ϕ_α . We can then rescale by $\log(t)$ to work with our new conventions. \square

Each of these vectors appears in the expression of dH_c . Since the terms that they correspond to cannot all vanish at the same time, it suffices to show that the coefficients of each of these term is non-negative and bounded. It is clear that the first two coefficients are indeed non-negative, and we have the bounds

$$\begin{aligned} |g(u)(2\pi u_0 - 2\pi(1 + \epsilon')u)| &\leq 2\pi\delta \\ |(1 - g(u))2cr_\Lambda(u)dr_\Lambda(u)| &\leq 2c\epsilon. \end{aligned}$$

In the second bound, we used the fact that $|dr_\Lambda| \leq 1$. It remains, therefore, to consider the coefficient of the last term. We re-write it as

$$\epsilon'\pi((1 + 5\epsilon)\langle u_0, u_0 \rangle - \langle u, u \rangle) - \pi(\langle u, u \rangle - 2\langle u, u_0 \rangle + \langle u_0, u_0 \rangle) - cr_\Lambda^2(u),$$

which simplifies to

$$5\epsilon'\pi\langle u_0, u_0 \rangle + \epsilon'\pi\langle u_0 - u, u_0 + u \rangle - \pi\langle u - u_0, u - u_0 \rangle - cr_\Lambda^2(u).$$

Note that the norm of the first term is $5\pi\epsilon'\langle u_0, u_0 \rangle$. The other terms are bounded above by $3\pi\epsilon'\langle u_0, u_0 \rangle$, $\pi\epsilon^2$, and $4c\epsilon^2$. In particular, since u_0 is an integral vector, and $\epsilon < \epsilon'$, the sum of the terms is bounded above by $10\pi\epsilon'\langle u_0, u_0 \rangle$ so long that c is not too large. It is also strictly positive since the first term dominates, so we conclude that the differential of our function does not attain the value $-2\pi u_0$ in the desired neighbourhood of u_0 .

Since the norm of dg is by assumption less than $\frac{2}{\epsilon}$, we conclude that the term corresponds to dg has norm bounded by $20\pi\epsilon'\langle u_0, u_0 \rangle$. We can choose ϵ' small enough to

ensure that this bound is much less than 1, so that dH_c is a bounded distance away from $-2\pi u_0$ as desired. \square

The reason for the careful choices that we have made in the above paragraphs is as follows: in order to interpolate between the differentials of two functions using cutoff functions without creating new critical points, we must know not only that the values of the differentials are close to each other, but also that the values of the functions are close. This is exactly the reason why we need the conical choice of

$$(\epsilon' + 5\epsilon'\epsilon)\langle\psi(u), \psi(u)\rangle$$

as a term in our Hamiltonian.

Corollary 5.9 *For sufficiently small $0 < c$, $\mathcal{L}_c(1)$ is an admissible Lagrangian that intersects the interior of \mathcal{L} in the set*

$$Q \cap \frac{1}{1 + \epsilon'}\mathbb{Z}^n \cong Q \cap \mathbb{Z}^n.$$

Further, $\mathcal{L}_c(1)$ agrees with a linear Lagrangian away from an ϵ neighbourhood of its boundary.

If we want a model for $\mathcal{L}(1)$ where the pair $(\mathcal{L}, \mathcal{L}(1))$ is negatively oriented, then we have to consider $\mathcal{L}_{-c}(1)$. However, following the above argument, we will find that the coefficient of dr_Λ is negative in this case. This means that we may be creating new intersection points if we try to interpolate between $\mathcal{L}_{-c}(1)$ and the graph of $(1 + \epsilon')\langle u, u \rangle$. We can easily fix this by using $(1 - \epsilon')\langle u, u \rangle$ instead. However, this means that we must also use $-(\epsilon' + 5\epsilon'\epsilon)$ in our expression for H_c^∂ . The rest of the argument then carries through as desired and yields

Corollary 5.10 *For sufficiently small $0 < c$, $\mathcal{L}_{-c}(1)$ is an admissible Lagrangian that intersects the interior of \mathcal{L} in the set*

$$(Q - \partial Q) \cap \mathbb{Z}^n.$$

Further, $\mathcal{L}_{-c}(1)$ agrees with a linear Lagrangian away from an ϵ neighbourhood of its boundary.

Since the Lagrangians $\mathcal{L}_{\pm c}(1)$ are equivalent up to admissible Hamiltonian isotopy for any value of c by Lemma 5.2, we revert to our old notation and refer to either of them as $\mathcal{L}(1)$.

We can also use the same method to construct $\mathcal{L}(j)$ for all j . Recall that the first choice we made is that of a constant $0 < \delta$, which allowed us to prove that no intersection

point occurred in a neighbourhood of the boundary by bounding the norm of a certain vector by constants dominated by δ . It was sufficient, in order to construct $\mathcal{L}(1)$, for δ to be much smaller than 1, since we simply had to avoid any lattice points that are not in Q . However, in order to construct $\mathcal{L}(j)$, we will have to avoid all elements of the lattice $\frac{1}{j}\mathbb{Z}^n$. This means that we have to choose δ much smaller than $\frac{1}{j}$. We can then use this to construct the Lagrangian boundary of $\mathcal{L}(j)$ using the function jH^∂ , then carry through the rest of the steps of the preceding construction. If j is negative, we must work with $-\epsilon'$ as explained above. We conclude the following:

Proposition 5.11 *For every integer $j \in \mathbb{Z}$, there exists an admissible Lagrangian $\mathcal{L}(j)$ such that the pair $(\mathcal{L}, \mathcal{L}(j))$ is positively oriented and their interior intersection points are in bijective correspondence with*

$$\frac{1}{j}\mathbb{Z}^N \cap Q$$

if $j > 0$ and with

$$\frac{1}{j}\mathbb{Z}^N \cap (Q - \partial Q)$$

if $j < 0$. Further, all interior intersection points occur on the complement of a small neighbourhood of the boundary. In this open set where the intersection points occur, $\mathcal{L}(j)$ is given by the Hamiltonian

$$-j\pi(1 + \epsilon')\langle u, u \rangle$$

for some small ϵ' .

5.4 Computing Floer groups and products

First we settle the issue of the Floer homology of \mathcal{L} with itself.

Lemma 5.12 *The only non-trivial Floer homology group of \mathcal{L} with itself is*

$$HF_n(\mathcal{L}, \mathcal{L}) = \mathbb{C}.$$

Proof We take a small Hamiltonian isotopy and compute the Floer homology of \mathcal{L} with its image \mathcal{L}' under the isotopy. Of course, this Hamiltonian isotopy must be chosen so that \mathcal{L}' is admissible and the pair $(\mathcal{L}, \mathcal{L}')$ is positively oriented. To achieve this, we simply reproduce the arguments of the previous section and interpolate between the graph of the differential of

$$-\langle u, \eta u \rangle$$

with η positive and the parallel transport of $\partial\mathcal{L}$ along some curve γ' which goes in the third quadrant. The point is that η can be chosen so small that the differential of this function does not take integral values on Q . Further, the positivity of η will guarantee that we can interpolate between the graph of this Hamiltonian and the parallel transport of the boundary in the direction of a “positive” curve γ without creating any new intersection points with \mathcal{L} . The above quadratic function has a critical point of index n at the origin, so we reach the desired conclusion. Note that this is consistent with $H_*(D^n, S^{n-1})$. \square

We can reinterpret Proposition 5.11 as follows:

Lemma 5.13 *The only non-vanishing Floer cohomology groups between \mathcal{L} and $\mathcal{L}(j)$ are given by*

$$HF^0(\mathcal{L}, \mathcal{L}(j)) \cong \bigoplus_{p \in Q \cap \mathbb{Z}^n} \mathbb{C} \cdot [p] \quad \text{if } 0 \leq j \quad \text{and,}$$

$$HF^n(\mathcal{L}, \mathcal{L}(j)) \cong \bigoplus_{p \in (Q - \partial Q) \cap \mathbb{Z}^n} \mathbb{C} \cdot [p] \quad \text{if } j < 0.$$

Proof The fact that all intersections between \mathcal{L} and $\mathcal{L}(j)$ occur in an open region where $\mathcal{L}(j)$ is given by a linear Lagrangian allows us to compute the Floer cohomology as in Section 3.4. \square

We can now discuss the cup product.

Lemma 5.14 *Identifying the generators of Floer homology groups with lattice points as in Lemma 5.13, the product*

$$HF^*(\mathcal{L}(l_1), \mathcal{L}(l_2)) \otimes HF^*(\mathcal{L}(l_2), \mathcal{L}(l_3)) \rightarrow HF^*(\mathcal{L}(l_1), \mathcal{L}(l_3))$$

is the same as the product for the Floer homology groups of the Lagrangians $\mathcal{L}_\infty(l_i)$ which we used in Section 3.4. Concretely, for $l_1 \leq l_2 \leq l_3$, we have

$$[p] \otimes [q] \mapsto \left[\frac{(l_2 - l_1)p + (l_3 - l_2)q}{l_3 - l_1} \right]$$

as in Lemma 3.20.

Proof First, we observe that we can choose the constant δ in the previous section so that our models for $\mathcal{L}(l_i)$ have all their interior intersection points in the open region where these Lagrangians are linear. This means that all the computations of Section 3.4 hold. We now justify the perturbation arguments used in that section.

It should be clear now that the Maslov index computation for triple intersections that we performed is legitimate, even when the intersection point occurs at the boundary of $\mathcal{L}_\infty(l_i)$, since in $\mathcal{L}(l_i)$ these are now interior intersection points, and can be perturbed to achieve transversality without affecting the admissibility condition. It remains to address the situation where two of the three Lagrangians are equal. The most important thing is to check that our generator of $HF^*(\mathcal{L}, \mathcal{L})$ acts as the identity. Say we are trying to compute

$$HF^*(\mathcal{L}, \mathcal{L}) \otimes HF^*(\mathcal{L}, \mathcal{L}(j)) \rightarrow HF^*(\mathcal{L}, \mathcal{L}(j)).$$

Now choose an intersection point $[q]$ between \mathcal{L} and $\mathcal{L}(j)$, and perturb \mathcal{L} to a Lagrangian \mathcal{L}' as in Lemma 5.12. Note that we can ensure that the intersection point p between \mathcal{L} and \mathcal{L}' occurs near q . We claim that the cup product is given by

$$[p] \otimes [q'] \rightarrow [q]$$

where $[q']$ is the corresponding nearby intersection point between \mathcal{L}' and $\mathcal{L}(j)$. In effect, we have reduced everything to a Maslov index computation. The point is that we can choose all these intersections to occur at the same point, and \mathcal{L}' to have slope $-\eta$. For sufficiently small η , the numbers $(0, \eta, j)$ satisfy Conditions (3–4) or (3–5) regardless of what the sign of j is. The usual obstruction in the first homology of the torus fibre allows us to conclude that there can be no other holomorphic triangle. \square

5.5 Completing the argument

Let X be the toric variety of the Introduction with an ample line bundle $\mathcal{O}(1)$. The last section, together with Equation (1–1), allows us to conclude that if $j \leq l$, then

$$\mathrm{Hom}_*(\mathcal{O}(j), \mathcal{O}(l)) \cong HF^*(\mathcal{L}(j), \mathcal{L}(l)),$$

and that this isomorphism is compatible with cup product. This proves Theorem 1.1.

We now use Serre duality to compute the case $j > l$. To simplify the notation, we let $j = 0$, so $l < 0$ and

$$\mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(l)) \cong (\mathrm{Hom}_{n-*}(\mathcal{O}, \mathcal{O}(-l) \otimes \kappa))^\vee.$$

Where κ is the canonical bundle. Since the piecewise linear function which defines κ takes the value -1 on each primitive vertex of a 1–cone of Δ , [10, Section 4.3], we see that our requirement that the interior of Q have an integral point (See Remark 3.10), is equivalent to the fact that the piecewise linear function defining $\mathcal{O}(-l) \otimes \kappa$ is convex. Its sections are in fact given by integral points which satisfy

$$\langle v_i, y \rangle \leq \phi(v_i) - 1$$

for every primitive vertex v_i of the 1–cones of Δ . But these are exactly the interior lattice points of $-lQ$. We have therefore proved that

$$\mathrm{Hom}_*(\mathcal{O}(j), \mathcal{O}(l)) \cong HF^*(\mathcal{L}(j), \mathcal{L}(l))$$

for all values of j and l . One may check that this isomorphism is indeed compatible with the cup product (which, for negative powers is interpreted through Serre duality). We check this for the case $l > 0$, $j < -l$ by computing

$$\mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(l)) \otimes \mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(j)) \rightarrow \mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(l+j)).$$

Note that Serre duality suggests that we should reduce this computation to that of

$$\mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(l)) \otimes \mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(-l-j) \otimes \kappa) \rightarrow \mathrm{Hom}_*(\mathcal{O}, \mathcal{O}(-j) \otimes \kappa),$$

in which all Hom's are concentrated in degree 0. The formula for the product is

$$[p] \otimes [q] \rightarrow [p+q],$$

which then dualizes to

$$[p] \otimes [r] = \sum_{q \in (l+j)(Q-\partial Q) \cap \mathbb{Z}^n \ni p+q=r} [q],$$

where the summation in the right hand side is either empty or consists of a unique term. The usual change of perspective from lattice points to $(l+j)Q$ to $\frac{1}{l+j}Q$ lattice points of Q yields the isomorphism with the product on Floer cohomology.

References

- [1] **A A Beilinson**, *Coherent sheaves on \mathbf{P}^n and problems in linear algebra*, Funktsional. Anal. i Prilozhen. 12 (1978) 68–69 MR509388
- [2] **V De Silva**, *Products in the symplectic Floer homology of Lagrangian intersections*, PhD thesis, Oxford University (1998)
- [3] **S K Donaldson**, *Symplectic submanifolds and almost-complex geometry*, J. Differential Geom. 44 (1996) 666–705 MR1438190
- [4] **M Einsiedler, M Kapranov, D Lind**, *Non-archimedean amoebas and tropical varieties* arXiv:math.AG/0408311
- [5] **Y Eliashberg, M Gromov**, *Convex symplectic manifolds*, from: “Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989)”, Proc. Sympos. Pure Math. 52, Amer. Math. Soc., Providence, RI (1991) 135–162 MR1128541
- [6] **A Floer**, *Morse theory for Lagrangian intersections*, J. Differential Geom. 28 (1988) 513–547 MR965228

- [7] **A Floer, H Hofer, D Salamon**, *Transversality in elliptic Morse theory for the symplectic action*, Duke Math. J. 80 (1995) 251–292 MR1360618
- [8] **K Fukaya, Y-G Oh**, *Zero-loop open strings in the cotangent bundle and Morse homotopy*, Asian J. Math. 1 (1997) 96–180 MR1480992
- [9] **K Fukaya, Y-G Oh, H Ohta, K Ono**, *Lagrangian intersection Floer theory - anomaly and obstruction*, unpublished manuscript available at <http://www.math.kyoto-u.ac.jp/~fukaya/> (2000)
- [10] **W Fulton**, *Introduction to toric varieties*, Annals of Mathematics Studies 131, Princeton University Press, Princeton, NJ (1993) MR1234037 The William H. Roever Lectures in Geometry
- [11] **IM Gel'fand, MM Kapranov, A V Zelevinsky**, *Discriminants, resultants, and multi-dimensional determinants*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA (1994) MR1264417
- [12] **M Kontsevich**, *Lectures at ENS Paris, Spring 1998* Notes by J Bellaïche, J-F Dat, I Marin, G Racinet and H Randriambololona
- [13] **M Kontsevich**, *Homological algebra of mirror symmetry*, from: “Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)”, Birkhäuser, Basel (1995) 120–139 MR1403918
- [14] **N C Leung**, *Mirror symmetry without corrections* arXiv:math.DG/0009235
- [15] **G Mikhalkin**, *Decomposition into pairs-of-pants for complex algebraic hypersurfaces*, Topology 43 (2004) 1035–1065 MR2079993
- [16] **W-D Ruan**, *Lagrangian torus fibration of quintic Calabi-Yau hypersurfaces. II. Technical results on gradient flow construction*, J. Symplectic Geom. 1 (2002) 435–521 MR1959057
- [17] **H Rullgård**, *Polynomial amoebas and convexity*, preprint, Stockholm University (2001)
- [18] **P Seidel**, *Graded Lagrangian submanifolds*, Bull. Soc. Math. France 128 (2000) 103–149 MR1765826

Department of Mathematics, University of Chicago
Chicago, IL 60637, USA

mabouzai@math.uchicago.edu

Proposed: Jim Bryan
Seconded: Lothar Göttsche, Simon Donaldson

Received: 26 November 2005
Revised: 3 May 2006