

Blanchfield and Seifert algebra in high-dimensional boundary link theory I: Algebraic K -theory

ANDREW RANICKI
DESMOND SHEIHAM

The classification of high-dimensional μ -component boundary links motivates decomposition theorems for the algebraic K -groups of the group ring $A[F_\mu]$ and the noncommutative Cohn localization $\Sigma^{-1}A[F_\mu]$, for any $\mu \geq 1$ and an arbitrary ring A , with F_μ the free group on μ generators and Σ the set of matrices over $A[F_\mu]$ which become invertible over A under the augmentation $A[F_\mu] \rightarrow A$. Blanchfield $A[F_\mu]$ -modules and Seifert A -modules are abstract algebraic analogues of the exteriors and Seifert surfaces of boundary links. Algebraic transversality for $A[F_\mu]$ -module chain complexes is used to establish a long exact sequence relating the algebraic K -groups of the Blanchfield and Seifert modules, and to obtain the decompositions of $K_*(A[F_\mu])$ and $K_*(\Sigma^{-1}A[F_\mu])$ subject to a stable flatness condition on $\Sigma^{-1}A[F_\mu]$ for the higher K -groups.

19D50, 57Q45; 20E05

Desmond Sheiham died 25 March 2005.

This paper is dedicated to the memory of Paul Cohn and Jerry Levine.

Introduction

For any integer $\mu \geq 1$ let F_μ be the free group on μ generators z_1, z_2, \dots, z_μ . The classification theory of high-dimensional μ -component boundary links involves ‘Seifert \mathbb{Z} -modules’ and ‘Blanchfield $\mathbb{Z}[F_\mu]$ -modules’, corresponding to the algebraic invariants obtained from μ -component Seifert surfaces and the boundary link exterior. This paper concerns the algebraic relationship between f.g. projective Seifert A -modules and h.d. 1 Blanchfield $A[F_\mu]$ -modules for any ring A , extending the work of Sheiham [41]. Part I deals with the algebraic K -theory of the Seifert and Blanchfield modules. Part II will deal with the algebraic L -theory of the Seifert and Blanchfield forms, such as arises in the computation of the cobordism groups of boundary links. The algebraic K - and L -theory in the knot case $\mu = 1$ have already been done by Ranicki [34].

Combinatorial transversality

Section 1 develops a combinatorial construction of fundamental domains for F_μ -covers of CW complexes which will serve as a role model for the algebraic transversality of $A[F_\mu]$ -module chain complexes in the subsequent sections. The F_μ -covers $p: \widetilde{W} \rightarrow W$ of a space W are classified by the homotopy classes of maps

$$c: W \longrightarrow BF_\mu = \bigvee_{\mu} S^1$$

with $\widetilde{W} = c^*EF_\mu$ the pullback to W of the universal cover EF_μ of BF_μ . Let $0 \in BF_\mu$ be the point at which the circles S^1 are joined, and choose points $1, 2, \dots, \mu \in BF_\mu \setminus \{0\}$, one in each circle. If W is a compact manifold then c is homotopic to a map which is transverse regular at $\{1, 2, \dots, \mu\} \subset BF_\mu$, so that

$$V = c^{-1}(\{1, 2, \dots, \mu\}) = V_1 \sqcup V_2 \sqcup \dots \sqcup V_\mu \subset W$$

is a disjoint union of μ codimension-1 submanifolds $V_i = c^{-1}(\{i\}) \subset W$ (which may be empty) and cutting W at V there is obtained a fundamental domain $U \subset \widetilde{W}$, a compact manifold with boundary

$$\partial U = \bigsqcup_{i=1}^{\mu} (V_i \sqcup z_i V_i).$$

If W is connected and $c_*: \pi_1(W) \rightarrow F_\mu$ is surjective then U is connected and V_1, V_2, \dots, V_μ are non-empty, and may be chosen to be connected. In the combinatorial version of transversality it is only required that W be a finite CW complex, and W may be replaced by a simple homotopy equivalent finite CW complex also denoted by W , with disjoint subcomplexes $V_1, V_2, \dots, V_\mu \subset W$ and a fundamental domain $U \subset \widetilde{W}$ which is a finite subcomplex with a subcomplex

$$\partial U = \bigsqcup_{i=1}^{\mu} (V_i \sqcup z_i V_i) \subset U, \quad V_i = U \cap z_i^{-1} U$$

such that

$$\bigcup_{g \in F_\mu} gU = \widetilde{W}, \quad gU \cap hU = \emptyset \text{ unless } g^{-1}h \in \{1, z_1, z_1^{-1}, \dots, z_\mu, z_\mu^{-1}\}.$$

Ranicki [35] developed combinatorial transversality at $Y \subset X$ for maps of finite CW complexes

$$W \rightarrow X = X_1 \cup_Y X_2$$

with X, X_1, X_2, Y connected and $\pi_1(Y) \rightarrow \pi_1(X_1), \pi_1(Y) \rightarrow \pi_1(X_2)$ injective. The essential difference from [35] is that we are here using the Cayley tree $EF_\mu = G_\mu$ of F_μ rather than the Bass–Serre tree of the amalgamated free product given by the Seifert–van Kampen Theorem

$$\pi_1(X) = \pi_1(X_1) *_{\pi_1(Y)} \pi_1(X_2)$$

for bookkeeping purposes. We show that W can be replaced by a simple homotopy equivalent finite CW complex \tilde{W} with disjoint subcomplexes $V_1, V_2, \dots, V_\mu \subset \tilde{W}$, such that the F_μ -cover \tilde{W} can be constructed from a fundamental domain finite subcomplex $U \subset \tilde{W}$ obtained by cutting W at $V = V_1 \sqcup V_2 \sqcup \dots \sqcup V_\mu \subset W$.

Algebraic transversality

Let A be an associative ring with 1. All A -modules will be understood to be left A -modules, unless a right A -module structure is specified.

Section 2 develops an ‘algebraic transversality’ technique for cutting $A[F_\mu]$ -modules along A -modules, which mimics the geometric transversality method of Section 1. In Section 2 we shall prove:

Theorem A Every $A[F_\mu]$ -module chain complex E admits a ‘Mayer–Vietoris presentation’

$$0 \longrightarrow \bigoplus_{i=1}^{\mu} C^{(i)}[F_\mu] \xrightarrow{f} D[F_\mu] \longrightarrow E \longrightarrow 0$$

with $C^{(i)}, D$ A -module chain complexes, and $f = (f_1^+ z_1 - f_1^- \dots f_\mu^+ z_\mu - f_\mu^-)$ defined using A -module chain maps $f_i^+, f_i^-: C^{(i)} \rightarrow D$. If E is a f.g. free $A[F_\mu]$ -module chain complex then $C^{(i)}, D$ can be chosen to be f.g. free A -module chain complexes, with $D \subset E$ and $f_i^+, f_i^-: C^{(i)} = D \cap z_i^{-1} D \rightarrow D$ given by $f_i^+(x) = x, f_i^-(x) = z_i x$.

Remark For $\mu = 1$ Theorem A was first proved by Waldhausen [47], being the chain complex version of the Higman linearization trick for matrices with entries in the Laurent polynomial extension $A[F_1] = A[z, z^{-1}]$. The algebraic transversality theory of [47] applies to chain complexes over the group rings $A[G_1 *_H G_2]$ of injective amalgamated free products $G_1 *_H G_2$, using the Bass–Serre theory of groups acting on trees. In principle, Theorem A for $\mu \geq 2$ could be proved by applying [47] to the successive free products in

$$F_\mu = F_1 * F_{\mu-1} = F_1 * (F_1 * F_{\mu-2}) = \dots = F_1 * (F_1 * (F_1 * \dots * (F_1)))$$

but this would be quite awkward in practice. In view of both the geometric motivation and the algebraic applications it is better to prove [Theorem A](#) (as will be done in [Section 2](#)) using the Cayley tree of F_μ with respect to the generator set $\{z_1, z_2, \dots, z_\mu\}$.

Boundary links

A μ -component link is a (locally flat, oriented) embedding

$$\ell: \bigsqcup_{\mu} S^n \subset S^{n+2}.$$

Every link admits a Seifert surface $V^{n+1} \subset S^{n+2}$, a codimension-1 submanifold with boundary

$$\partial V = \ell\left(\bigsqcup_{\mu} S^n\right) \subset S^{n+2}.$$

By definition, ℓ is a μ -component boundary link if there exists a μ -component Seifert surface

$$V^{n+1} = V_1 \sqcup V_2 \sqcup \dots \sqcup V_\mu \subset S^{n+2}.$$

The exterior of a link ℓ is the $(n+2)$ -dimensional manifold with boundary

$$(W^{n+2}, \partial W) = \left(\text{cl}\left(S^{n+2} - \left(\ell\left(\bigsqcup_{\mu} S^n\right) \times D^2\right)\right), \ell\left(\bigsqcup_{\mu} S^n\right) \times S^1\right).$$

In particular, a knot $S^n \subset S^{n+2}$ is a 1-component boundary link.

The trivial μ -component boundary link

$$\ell_0: \bigsqcup_{\mu} S^n \subset S^{n+2}$$

is defined by the connected sum of μ copies of the trivial knot

$$S^n \subset (S^n \times D^2) \cup (D^{n+1} \times S^1) = S^{n+2},$$

so that

$$\ell_0: \bigsqcup_{\mu} S^n \subset \#_{\mu} S^{n+2} = S^{n+2} = \left(\bigsqcup_{\mu} S^n \times D^2\right) \cup W_0$$

has Seifert surface and exterior

$$V_0 = \bigsqcup_{\mu} D^{n+1}, \quad W_0 = \#_{\mu} (D^{n+1} \times S^1) \subset S^{n+2}.$$

The exterior W_0 has the homotopy type of $\bigvee_{\mu} S^1 \vee \bigvee_{\mu-1} S^{n+1}$, with $\pi_1(W_0) = F_\mu$.

We shall make much use of the fact that the universal cover of $BF_\mu = \bigvee_\mu S^1$ is the contractible space with free F_μ -action defined by the Cayley tree $EF_\mu = G_\mu$ of F_μ , with vertices $g \in F_\mu$ and edges (g, gz_i) ($g \in F_\mu, 1 \leq i \leq \mu$). The cellular chain complex $C(EF_\mu) = C(G_\mu)$ is the standard 1-dimensional f.g. free $\mathbb{Z}[F_\mu]$ -module resolution of \mathbb{Z}

$$0 \longrightarrow C_1(G_\mu) = \bigoplus_{i=1}^{\mu} \mathbb{Z}[F_\mu] \xrightarrow{d} C_0(G_\mu) = \mathbb{Z}[F_\mu] \longrightarrow \mathbb{Z} \longrightarrow 0,$$

the Mayer-Vietoris presentation with $d = (z_1 - 1 \ z_2 - 1 \ \dots \ z_\mu - 1)$.

The exterior W of an n -dimensional link $\ell: \bigsqcup_\mu S^n \subset S^{n+2}$ is homotopy equivalent to the complement $S^{n+2} \setminus \ell(\bigsqcup_\mu S^n)$, so that

$$\begin{aligned} H_*(W) &= H_*\left(S^{n+2} \setminus \ell\left(\bigsqcup_\mu S^n\right)\right) \\ &= H^{n+2-*}\left(S^{n+2}, \ell\left(\bigsqcup_\mu S^n\right)\right) = H^{n+1-*}\left(\bigsqcup_\mu S^n\right) \quad (* \neq 0, n+2) \end{aligned}$$

by Alexander duality. The homology groups $H_*(W), H_*(W_0)$ are thus the same:

$$H_r(W) = H_r(W_0) = \begin{cases} \mathbb{Z} & \text{if } r = 0 \\ \bigoplus_{\mu} \mathbb{Z} & \text{if } r = 1 \\ \bigoplus_{\mu-1} \mathbb{Z} & \text{if } r = n + 1 \\ 0 & \text{otherwise.} \end{cases}$$

The homotopy groups $\pi_*(W), \pi_*(W_0)$ are in general not the same, on account of linking. By Smythe [43] and Gutierrez [22] ℓ is a boundary link if and only if there exists a surjection $\pi_1(W) \rightarrow \pi_1(W_0) = F_\mu$ sending the meridians $m_1, m_2, \dots, m_\mu: S^1 \subset W$ around the μ components $\ell_1, \ell_2, \dots, \ell_\mu: S^n \subset S^{n+2}$ of ℓ to the generators $z_1, z_2, \dots, z_\mu \in F_\mu$. We shall only be considering boundary links ℓ with a particular choice of such a surjection $\pi_1(W) \rightarrow F_\mu$, the F_μ -links of Cappell and Shaneson [9]. For any such ℓ there exists a map $c: W \rightarrow W_0$ which induces a surjection $c_*: \pi_1(W) \rightarrow \pi_1(W_0)$ and isomorphisms $c_*: H_*(W) \cong H_*(W_0)$. Let $\tilde{W} = c^* \tilde{W}_0$ be the pullback F_μ -cover of W , with a f.g. free $\mathbb{Z}[F_\mu]$ -module cellular chain complex $C(\tilde{W})$. An F_μ -equivariant lift $\tilde{c}: \tilde{W} \rightarrow \tilde{W}_0$ of c induces a $\mathbb{Z}[F_\mu]$ -module chain map $\tilde{c}: C(\tilde{W}) \rightarrow C(\tilde{W}_0)$ and a \mathbb{Z} -module chain equivalence $c: C(W) \rightarrow C(W_0)$. A μ -component Seifert surface $V = V_1 \sqcup V_2 \sqcup \dots \sqcup V_\mu \subset S^{n+2}$ for ℓ has a neighbourhood $V \times [-1, 1] \subset S^{n+2}$, with $V = V \times \{0\}$. The F_μ -cover \tilde{W} can be constructed from F_μ copies of $S^{n+2} \setminus V$, glued together using the inclusions $f_i^+, f_i^-: V_i \rightarrow S^{n+2} \setminus V$

defined by

$$f_i^\pm(v_i) = (v_i, \pm 1) \in V \times [-1, 1] \subset S^{n+2}.$$

It follows that $C(\tilde{W})$ has a f.g. free $\mathbb{Z}[F_\mu]$ -module Mayer–Vietoris presentation

$$0 \longrightarrow \bigoplus_{i=1}^{\mu} C(V_i)[F_\mu] \xrightarrow{f} C(S^{n+2} \setminus V)[F_\mu] \longrightarrow C(\tilde{W}) \longrightarrow 0$$

with $f = f^+z - f^- = (f_1^+z_1 - f_1^- \dots f_\mu^+z_\mu - f_\mu^-)$.

Seifert and Blanchfield modules

There are four fundamental notions in our abstract version for any ring A of the Seifert and Blanchfield modules of μ -component boundary links:

- (i) A *Seifert A -module* is a triple

$$(P, e, \{\pi_i\}) = (A\text{-module, endomorphism, } \{\pi_i\})$$

where $\{\pi_i: P \rightarrow P\}$ is a system of idempotents expressing P as a μ -fold direct sum, with

$$\pi_i: P = P_1 \oplus P_2 \oplus \dots \oplus P_\mu \rightarrow P;$$

$$(x_1, x_2, \dots, x_\mu) \mapsto (0, \dots, 0, x_i, 0, \dots, 0).$$

Let $Sei_\infty(A)$ be the category of Seifert A -modules. A Seifert A -module $(P, e, \{\pi_i\})$ is *f.g. projective* if P is a f.g. projective A -module. Let $Sei(A) \subset Sei_\infty(A)$ be the full subcategory of the f.g. projective Seifert A -modules.

- (ii) A *Blanchfield $A[F_\mu]$ -module* M is an $A[F_\mu]$ -module such that

$$\text{Tor}_*^{A[F_\mu]}(A, M) = 0,$$

regarding A as a right $A[F_\mu]$ -module via the augmentation map

$$\epsilon: A[F_\mu] \rightarrow A; z_i \mapsto 1.$$

Let $Bla_\infty(A)$ be the category of Blanchfield $A[F_\mu]$ -modules. In [Section 3.2](#) Blanchfield $A[F_\mu]$ -modules will be identified with the F_μ -link modules in the sense of Sheiham [\[41\]](#), that is $A[F_\mu]$ -modules M which admit an $A[F_\mu]$ -module presentation

$$0 \longrightarrow P[F_\mu] \xrightarrow{d} Q[F_\mu] \longrightarrow M \longrightarrow 0$$

for A -modules P, Q with the augmentation $\epsilon(d): P \rightarrow Q$ an A -module isomorphism. Thus $\mathcal{B}la_\infty(A)$ is just the F_μ -link module category $\mathcal{F}lk_\infty(A)$ of [41]. A Blanchfield $A[F_\mu]$ -module M has *homological dimension 1* (or *h.d. 1* for short) if it has a 1-dimensional f.g. projective $A[F_\mu]$ -module resolution

$$0 \longrightarrow K \xrightarrow{d} L \longrightarrow M \longrightarrow 0$$

with (necessarily) $\epsilon(d) = 1 \otimes d: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$ an A -module isomorphism. Let $\mathcal{B}la(A) \subset \mathcal{B}la_\infty(A)$ be the full subcategory of the h.d. 1 Blanchfield $A[F_\mu]$ -modules. Let $\mathcal{F}lk(A) \subset \mathcal{B}la(A)$ be the full subcategory of the h.d. 1 Blanchfield modules M which admit a 1-dimensional induced f.g. projective $A[F_\mu]$ -module resolution

$$0 \longrightarrow P[F_\mu] \xrightarrow{d} Q[F_\mu] \longrightarrow M \longrightarrow 0$$

with P, Q f.g. projective A -modules. As in [41] the objects of $\mathcal{F}lk(A)$ will be called *h.d. 1 F_μ -link modules*.

- (iii) The *covering* of a Seifert A -module $(P, e, \{\pi_i\})$ is the Blanchfield $A[F_\mu]$ -module

$$B(P, e, \{\pi_i\}) = \text{coker}(1 - e + ez: P[F_\mu] \rightarrow P[F_\mu])$$

with $z = \sum_{i=1}^{\mu} \pi_i z_i: P[F_\mu] \rightarrow P[F_\mu]$, defining functors

$$B_\infty: \mathcal{S}ei_\infty(A) \rightarrow \mathcal{B}la_\infty(A), \quad B: \mathcal{S}ei(A) \rightarrow \mathcal{F}lk(A).$$

- (iv) A Seifert A -module $(P, e, \{\pi_i\})$ is *primitive* if $B(P, e, \{\pi_i\}) = 0$. Let

$$\mathcal{P}rim_\infty(A) = \ker(B_\infty: \mathcal{S}ei_\infty(A) \rightarrow \mathcal{B}la_\infty(A))$$

be the full subcategory of $\mathcal{S}ei_\infty(A)$ with objects the primitive Seifert A -modules, and let

$$\mathcal{P}rim(A) = \ker(B: \mathcal{S}ei(A) \rightarrow \mathcal{F}lk(A)) \subset \mathcal{S}ei(A)$$

be the full subcategory of $\mathcal{S}ei(A)$ with objects the primitive f.g. projective Seifert A -modules.

Simple boundary links

The motivational examples of f.g. projective Seifert \mathbb{Z} -modules and h.d. 1 F_μ -link $\mathbb{Z}[F_\mu]$ -modules come from the $(2q-1)$ -dimensional μ -component boundary links

$\ell: \bigsqcup_{\mu} S^{2q-1} \subset S^{2q+1}$ which are *simple*, meaning that the exterior W has homotopy groups

$$\pi_r(W) = \begin{cases} F_{\mu} & \text{if } r = 1 \\ 0 & \text{if } 2 \leq r \leq q-1, \end{cases}$$

so that the universal cover \widetilde{W} is $(q-1)$ -connected. These conditions are equivalent to the existence of a μ -component Seifert surface $V = V_1 \sqcup V_2 \sqcup \cdots \sqcup V_{\mu}$ with each component V_i $(q-1)$ -connected:

$$\pi_r(V_i) = 0 \quad (1 \leq i \leq \mu, 1 \leq r \leq q-1).$$

The homology of the Seifert surface defines a f.g. projective (actually f.g. free) Seifert \mathbb{Z} -module $(P, e, \{\pi_i\})$, with

$$\pi_i = 0 \oplus \cdots \oplus 0 \oplus 1 \oplus 0 \oplus \cdots \oplus 0: P = \bigoplus_{i=1}^{\mu} H_q(V_i) \rightarrow P = \bigoplus_{i=1}^{\mu} H_q(V_i)$$

and

$$e = (f_1^+ \ f_2^+ \ \dots \ f_{\mu}^+): P = H_q(V) = \bigoplus_{i=1}^{\mu} H_q(V_i) \longrightarrow H_q(S^{2q+1} \setminus V) = H^q(V) = H_q(V) = P$$

the endomorphism induced by the inclusions $f_i^+: V_i \rightarrow S^{2q+1} \setminus V$, identifying

$$H_q(S^{2q+1} \setminus V) = H^q(V)$$

by Alexander duality and $H^q(V) = H_q(V)$ by Poincaré duality. The covering of $(P, e, \{\pi_i\})$ is the h.d. 1 F_{μ} -link $\mathbb{Z}[F_{\mu}]$ -module

$$B(P, e, \{\pi_i\}) = H_q(\widetilde{W})$$

defined by the homology of the F_{μ} -cover \widetilde{W} of the exterior W . The f.g. projective Seifert \mathbb{Z} -module $(P, e, \{\pi_i\})$ is primitive if and only if $H_q(\widetilde{W}) = 0$; for $q \geq 2$ this is the case if and only if ℓ is unlinked (Gutierrez [22]).

Blanchfield = Seifert/primitive

Section 3 uses algebraic transversality to prove that every h.d. 1 F_{μ} -link module M is isomorphic to the covering $B(P, e, \{\pi_i\})$ of a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$, and that morphisms of h.d. 1 F_{μ} -link modules can be expressed as fractions of morphisms of f.g. projective Seifert A -modules.

The algebraic relation between Seifert A -modules and Blanchfield $A[F_{\mu}]$ -modules for $A = \mathbb{Z}$ was first investigated systematically in the knot case $\mu = 1$, by Levine [25];

[26] and Trotter [46], and for the link case $\mu \geq 1$ by Farber [13; 14] and Sheiham [41]. In particular, [41] expressed the Blanchfield module category $\mathcal{B}la_\infty(A) = \mathcal{F}lk_\infty(A)$ as the quotient of the Seifert A -module category $\mathcal{S}ei_\infty(A)$ by the primitive Seifert A -module subcategory $\mathcal{P}rim_\infty(A)$, as we now recall.

Let \mathcal{A} be an abelian category. By definition, a *Serre subcategory* $\mathcal{C} \subset \mathcal{A}$ is a non-empty full subcategory such that for every exact sequence in \mathcal{A}

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

M is an object in \mathcal{C} if and only if M', M'' are objects in \mathcal{C} . Gabriel [17] defined the quotient abelian category \mathcal{A}/\mathcal{C} with the same objects as \mathcal{A} but different morphisms: if M, N are objects in \mathcal{A} then

$$\text{Hom}_{\mathcal{A}/\mathcal{C}}(M, N) = \varinjlim \text{Hom}_{\mathcal{A}}(M', N'')$$

with the direct limit taken over all the exact sequences in \mathcal{A}

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0, 0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$$

with M'', N' objects in \mathcal{C} . The canonical functor $F: \mathcal{A} \rightarrow \mathcal{A}/\mathcal{C}; A \mapsto A$ sends each object C in \mathcal{C} to $F(C) = 0$, and has the universal property that for any exact functor $G: \mathcal{A} \rightarrow \mathcal{B}$ such that $G(C) = 0$ for all objects in \mathcal{C} there exists a unique functor $\bar{G}: \mathcal{A}/\mathcal{C} \rightarrow \mathcal{B}$ such that $\bar{G}F = G$. In particular, if \mathcal{B} is an exact category and $G: \mathcal{A} \rightarrow \mathcal{B}$ is an exact functor then the full subcategory $\mathcal{C} \subset \mathcal{A}$ with objects C such that $G(C) = 0$ is a Serre subcategory, and there is induced a functor $\bar{G}: \mathcal{A}/\mathcal{C} \rightarrow \mathcal{B}; A \mapsto G(A)$ such that $G = \bar{G}F$.

By definition, a category is *small* if the class of morphisms is a set. In order to avoid set-theoretic difficulties we shall only be dealing with categories which are *essentially small*, ie equivalent to a small category.

Let \mathcal{A} be an essentially small category, and let Σ be a set of morphisms in \mathcal{A} , e.g. the morphisms of a subcategory. A *category of fractions* $\Sigma^{-1}\mathcal{A}$ is a category with a universally Σ -inverting functor $F: \mathcal{A} \rightarrow \Sigma^{-1}\mathcal{A}$, meaning that:

- (i) F sends each $f \in \Sigma$ to an isomorphism $F(f)$ in $\Sigma^{-1}\mathcal{A}$,
- (ii) for any functor $G: \mathcal{A} \rightarrow \mathcal{B}$ which sends each $f \in \Sigma$ to an isomorphism $G(f)$ there exists a unique functor $\bar{G}: \Sigma^{-1}\mathcal{A} \rightarrow \mathcal{B}$ such that $\bar{G}F = G$.

An essentially small category of fractions $\Sigma^{-1}\mathcal{A}$ exists, with the same objects as \mathcal{A} , and such a category is unique up to isomorphism (Gabriel and Zisman [18], Borceux

[7, 5.2.2]). For example, if \mathcal{A} is an abelian category and $\mathcal{C} \subset \mathcal{A}$ is a Serre subcategory, then

$$\mathcal{A}/\mathcal{C} = \Sigma^{-1}\mathcal{A}$$

is a category of fractions inverting the set Σ of morphisms f in \mathcal{A} with $\ker(f)$ and $\text{coker}(f)$ in \mathcal{C} .

An $A[F_\mu]$ -module M is Blanchfield if and only if the A -module morphism

$$\gamma: \bigoplus_{\mu} M \rightarrow M; (m_1, m_2, \dots, m_{\mu}) \mapsto \sum_{i=1}^{\mu} (z_i - 1)m_i$$

is an isomorphism, called the *Sato isomorphism* (after [37], the case $A = \mathbb{Z}$). As in Sheiham [41], for any Blanchfield $A[F_\mu]$ -module M use the A -module morphisms

$$\begin{aligned} p_i: \bigoplus_{\mu} M &\rightarrow M; (m_1, m_2, \dots, m_{\mu}) \mapsto m_i, \\ \omega: \bigoplus_{\mu} M &\rightarrow M; (m_1, m_2, \dots, m_{\mu}) \mapsto \sum_{i=1}^{\mu} m_i, \\ \pi_i &= \gamma p_i \gamma^{-1}: M \rightarrow M, \\ e &= \omega \gamma^{-1}: M \rightarrow M \end{aligned}$$

to define a Seifert A -module $U(M) = (M, e, \{\pi_i\})$.

The categories $\mathcal{P}rim_{\infty}(A)$, $\mathcal{S}ei_{\infty}(A)$ are abelian, while $\mathcal{B}la_{\infty}(A)$ is in general only exact. The covering functor $B_{\infty}: \mathcal{S}ei_{\infty}(A) \rightarrow \mathcal{B}la_{\infty}(A)$ was shown in [41, 5.2] to be exact, so that $\mathcal{P}rim_{\infty}(A) \subset \mathcal{S}ei_{\infty}(A)$ is a Serre subcategory. Thus if Ξ_{∞} is the set of morphisms f in $\mathcal{S}ei_{\infty}(A)$ such that $B(f)$ is an isomorphism in $\mathcal{B}la_{\infty}(A)$, or equivalently $\ker(f)$ and $\text{coker}(f)$ are in $\mathcal{P}rim_{\infty}(A)$, then

$$\mathcal{S}ei_{\infty}(A)/\mathcal{P}rim_{\infty}(A) = \Xi_{\infty}^{-1}\mathcal{S}ei_{\infty}(A).$$

The induced exact functor $\bar{B}_{\infty}: \mathcal{S}ei_{\infty}(A)/\mathcal{P}rim_{\infty}(A) \rightarrow \mathcal{B}la_{\infty}(A)$ is such that

$$B_{\infty}: \mathcal{S}ei_{\infty}(A) \rightarrow \mathcal{S}ei_{\infty}(A)/\mathcal{P}rim_{\infty}(A) \xrightarrow{\bar{B}_{\infty}} \mathcal{B}la_{\infty}(A)$$

and has the universal property of inverting Ξ_{∞} . The functor \bar{B}_{∞} was shown to be an equivalence in [41, 5.15] using the fact that the functor

$$U_{\infty}: \mathcal{B}la_{\infty}(A) \rightarrow \mathcal{S}ei_{\infty}(A); M \mapsto U(M)$$

is right adjoint to B : for any Seifert A -module V there is a natural isomorphism

$$\text{Hom}_{\mathcal{B}la_{\infty}(A)}(B(V), M) \cong \text{Hom}_{\mathcal{S}ei_{\infty}(A)}(V, U(M)).$$

The functor U_∞ is fully faithful, allowing $Bla_\infty(A)$ to be regarded as a full subcategory of $Sei_\infty(A)$. By [41, 5.15] U_∞ induces a functor

$$\bar{U}_\infty: Bla_\infty(A) \rightarrow Sei_\infty(A)/\mathcal{P}rim_\infty(A)$$

which is an equivalence inverse to \bar{B}_∞ . Thus up to equivalence

$$Sei_\infty(A)/\mathcal{P}rim_\infty(A) = \Xi_\infty^{-1}Sei_\infty(A) = Bla_\infty(A).$$

The categories $\mathcal{P}rim(A)$, $Sei(A)$, $\mathcal{F}lk(A)$, $Bla(A)$ are exact but not in general abelian. As in [41] let $Sei(A)/\mathcal{P}rim_\infty(A) \subset Sei_\infty(A)/\mathcal{P}rim_\infty(A)$ be the full subcategory with objects in $Sei(A)$. The equivalence

$$\bar{B}_\infty: Sei_\infty(A)/\mathcal{P}rim_\infty(A) \xrightarrow{\approx} Bla_\infty(A)$$

was shown in [41, 5.17] to restrict to an equivalence of exact sequences

$$\bar{B}: Sei(A)/\mathcal{P}rim_\infty(A) \xrightarrow{\approx} \mathcal{F}lk(A)$$

with

$$B: Sei(A) \rightarrow Sei(A)/\mathcal{P}rim_\infty(A) \xrightarrow[\approx]{\bar{B}} \mathcal{F}lk(A).$$

From the construction of $Sei_\infty(A)/\mathcal{P}rim_\infty(A)$ a morphism in $Sei(A)/\mathcal{P}rim_\infty(A)$ may involve objects in $Sei_\infty(A)$ which are not in $Sei(A)$, so that the equivalence \bar{B} cannot be used to relate the algebraic K -theories of $Sei(A)$ and $\mathcal{F}lk(A)$.

A category of fractions $\Sigma^{-1}\mathcal{A}$ has a *left calculus of fractions* if:

- (i) $(1: A \rightarrow A) \in \Sigma$ for every object A in \mathcal{A} ,
- (ii) if $(s: A \rightarrow B), (t: B \rightarrow C) \in \Sigma$ then $(ts: A \rightarrow C) \in \Sigma$,
- (iii) for any $f: A \rightarrow B$ in \mathcal{A} and $s: A \rightarrow D$ in Σ there exist $g: D \rightarrow C$ in \mathcal{A} and $t: B \rightarrow C$ in Σ such that $tf = gs: A \rightarrow C$,
- (iv) for any $f, g: A \rightarrow B$ in \mathcal{A} and $s: D \rightarrow A$ in Σ with $fs = gs: D \rightarrow B$ there exists $(t: B \rightarrow C) \in \Sigma$ with $tf = tg: A \rightarrow C$.

It then follows that a morphism $A \rightarrow B$ in $\Sigma^{-1}\mathcal{A}$ can be regarded as an equivalence class $s^{-1}f$ of pairs $(f: A \rightarrow C, s: B \rightarrow C)$ of morphisms in \mathcal{A} with $s \in \Sigma$, where

$$(f, s) \sim (f', s') \text{ if there exist morphisms } g: C \rightarrow D, g': C' \rightarrow D \text{ in } \mathcal{A} \\ \text{with } (gs = g's': B \rightarrow D) \in \Sigma \text{ and } gf = g'f': A \rightarrow D$$

so that

$$s^{-1}f = (gs)^{-1}(gf) = (g's')^{-1}(g'f') = s'^{-1}f': A \rightarrow B \text{ in } \Sigma^{-1}\mathcal{A}.$$

Let Ξ be the set of morphisms f in $\mathcal{S}ei(A)$ such that $B(f)$ is an isomorphism in $\mathcal{F}lk(A)$, or equivalently such that $\ker(f)$ and $\operatorname{coker}(f)$ are in $\mathcal{P}rim_\infty(A)$. In [Section 3](#) we shall prove:

Theorem B (i) *The category of fractions $\Xi^{-1}\mathcal{S}ei(A)$ has a left calculus of fractions, and the covering functor $B: \mathcal{S}ei(A) \rightarrow \mathcal{F}lk(A)$ induces an equivalence of exact categories*

$$\bar{B}: \Xi^{-1}\mathcal{S}ei(A) \xrightarrow{\approx} \mathcal{F}lk(A).$$

(ii) *The h.d. 1 Blanchfield $A[F_\mu]$ -module category $\mathcal{B}la(A)$ is the idempotent completion of the h.d. 1 F_μ -link module category $\mathcal{F}lk(A)$.*

The key step in the proof of [Theorem B](#) (i) is the use of the algebraic transversality [Theorem A](#) to verify that for any h.d. 1 F_μ -link module M the Seifert A -module $U(M)$ is a direct limit of morphisms in Ξ .

Primitive = near-projection

[Section 4](#) gives an intrinsic characterization of the primitive f.g. projective Seifert A -modules $(P, e, \{\pi_i\})$ as generalized near-projections.

An endomorphism $e: P \rightarrow P$ of an A -module P is *nilpotent* if $e^N = 0$ for some $N \geq 0$.

An endomorphism $e: P \rightarrow P$ is a *near-projection* if $e(1-e): P \rightarrow P$ is nilpotent ([Lück and Ranicki \[28\]](#)).

In [Section 4](#) we shall prove:

Theorem C *A f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ is primitive if and only if it can be expressed as*

$$(P, e, \{\pi_i\}) = \left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & e^{+-} \\ e^{-+} & e^{--} \end{pmatrix}, \{\pi_i^+\} \oplus \{\pi_i^-\} \right)$$

and the 2μ -component Seifert A -module

$$(P', e', \pi') = \left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix}, \{\pi_i^+\} \oplus \{\pi_i^-\} \right)$$

is such that $e'z': P'[F_{2\mu}] \rightarrow P'[F_{2\mu}]$ is nilpotent, with $F_{2\mu}$ the free group on 2μ generators $z'_1, \dots, z'_{2\mu}$.

For $\mu = 1$ the condition for a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ to be primitive is just that e be a near-projection. For $\mu = 1$ [Theorem C](#) is just the result of Bass, Heller and Swan [\[5\]](#) that $1 - e + ez: P[z, z^{-1}] \rightarrow P[z, z^{-1}]$ is an $A[z, z^{-1}]$ -module isomorphism if and only if e is a near-projection, if and only if $(P, e) = (P^+, e^{++}) \oplus (P^-, e^{--})$ with $e^{++}: P^+ \rightarrow P^+$ and $1 - e^{--}: P^- \rightarrow P^-$ nilpotent.

Algebraic K -theory

[Section 5](#) obtains results on the algebraic K -theory of $A[F_\mu]$, $\mathcal{P}r\imath m(A)$, $\mathcal{S}e\imath(A)$, $\mathcal{F}l\kappa(A)$ and $\mathcal{B}l\alpha(A)$, using the algebraic K -theory noncommutative localization exact sequences of Schofield [\[39\]](#) and Neeman–Ranicki [\[30; 31\]](#).

The class group $K_0(\mathcal{E})$ of an exact category \mathcal{E} is the Grothendieck group with one generator $[M]$ for each object M in \mathcal{E} , and one relation $[K] - [L] + [M] = 0$ for each exact sequence in \mathcal{E}

$$0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0.$$

The algebraic K -groups $K_n(\mathcal{E})$ are defined by Quillen [\[32\]](#) for $n \geq 1$ and by Schlichting [\[38\]](#) for $n \leq -1$. Write

$$\begin{aligned} \mathcal{P}r\imath m_* (A) &= K_*(\mathcal{P}r\imath m(A)), & \mathcal{S}e\imath_* (A) &= K_*(\mathcal{S}e\imath(A)), \\ \mathcal{B}l\alpha_* (A) &= K_*(\mathcal{B}l\alpha(A)), & \mathcal{F}l\kappa_* (A) &= K_*(\mathcal{F}l\kappa(A)), \end{aligned}$$

noting that $\mathcal{B}l\alpha_n(A) = \mathcal{F}l\kappa_n(A)$ for $n \neq 0$.

Theorem D (i) *The algebraic K -groups of $A[F_\mu]$ split as*

$$K_*(A[F_\mu]) = K_*(A) \oplus \bigoplus_{\mu} K_{*-1}(A) \oplus \widetilde{\mathcal{P}r\imath m}_{*-1}(A).$$

(ii) *The sequence of functors*

$$\mathcal{P}r\imath m(A) \longrightarrow \mathcal{S}e\imath(A) \xrightarrow{B} \mathcal{B}l\alpha(A)$$

induces a long exact sequence of algebraic K -groups

$$\dots \rightarrow \mathcal{P}r\imath m_n(A) \rightarrow \mathcal{S}e\imath_n(A) \xrightarrow{B} \mathcal{B}l\alpha_n(A) \rightarrow \mathcal{P}r\imath m_{n-1}(A) \rightarrow \dots$$

with

$$\text{im}(B: \mathcal{S}e\imath_0(A) \rightarrow \mathcal{B}l\alpha_0(A)) = \mathcal{F}l\kappa_0(A) \subseteq \mathcal{B}l\alpha_0(A).$$

(iii) The exact sequence in (ii) splits as a direct sum of exact sequences

$$\begin{aligned} \cdots \rightarrow \bigoplus_{2\mu} K_n(A) \rightarrow \bigoplus_{\mu} K_n(A) \xrightarrow{0} \bigoplus_{\mu} K_{n-1}(A) \rightarrow \bigoplus_{2\mu} K_{n-1}(A) \rightarrow \cdots, \\ \cdots \rightarrow \widetilde{\text{Prim}}_n(A) \rightarrow \widetilde{\text{Sei}}_n(A) \rightarrow \widetilde{\text{Bla}}_n(A) \rightarrow \widetilde{\text{Prim}}_{n-1}(A) \rightarrow \cdots. \end{aligned}$$

For $\mu = 1$ $\mathcal{P}r\text{im}(A)$ is the exact category of f.g. projective A -modules P with a near-projection $e: P \rightarrow P$, which is equivalent to the product $\mathcal{N}il(A) \times \mathcal{N}il(A)$ of two copies of the exact category $\mathcal{N}il(A)$ of f.g. projective A -modules P with a nilpotent endomorphism $e: P \rightarrow P$, and

$$\begin{aligned} \text{Prim}_*(A) &= K_*(\mathcal{P}r\text{im}(A)) = \text{Nil}_*(A) \oplus \text{Nil}_*(A), \\ \text{Nil}_*(A) &= K_*(\mathcal{N}il(A)) = K_*(A) \oplus \widetilde{\text{Nil}}_*(A), \\ \widetilde{\text{Prim}}_*(A) &= \widetilde{\text{Nil}}_*(A) \oplus \widetilde{\text{Nil}}_*(A). \end{aligned}$$

Thus for $\mu = 1$ **Theorem D** (i) is just the splitting theorem of Bass, Heller and Swan [5], [4] for $K_1(A[z, z^{-1}])$ and its generalization to the higher K -groups

$$K_*(A[z, z^{-1}]) = K_*(A) \oplus K_{*-1}(A) \oplus \widetilde{\text{Nil}}_{*-1}(A) \oplus \widetilde{\text{Nil}}_{*-1}(A).$$

Theorem D (ii)–(iii) is new even in the case $\mu = 1$.

Let $\Sigma^{-1}A[F_\mu]$ be the noncommutative Cohn (ie universal) localization of $A[F_\mu]$ inverting the set Σ of the morphisms of f.g. projective $A[F_\mu]$ -modules which induce isomorphisms of f.g. projective A -modules under the augmentation $\epsilon: A[F_\mu] \rightarrow A$. The exact category $H(A[F_\mu], \Sigma)$ of h.d. 1 Σ -torsion $A[F_\mu]$ -modules is such that

$$H(A[F_\mu], \Sigma) = \text{Bla}(A), \quad K_*(H(A[F_\mu], \Sigma)) = \text{Bla}_*(A).$$

Theorem E (i) The localization exact sequence

$$K_1(A[F_\mu]) \rightarrow K_1(\Sigma^{-1}A[F_\mu]) \rightarrow K_0(H(A[F_\mu], \Sigma)) \rightarrow K_0(A[F_\mu]) \rightarrow \cdots$$

splits as a direct sum of the exact sequences

$$\begin{aligned} K_1(A) \oplus \bigoplus_{\mu} K_0(A) \rightarrow K_1(A) \xrightarrow{0} \bigoplus_{\mu} K_{-1}(A) \rightarrow K_0(A) \oplus \bigoplus_{\mu} K_{-1}(A) \rightarrow \cdots, \\ \widetilde{\text{Prim}}_0(A) \rightarrow \widetilde{\text{Sei}}_0(A) \rightarrow \widetilde{\text{Bla}}_0(A) \rightarrow \widetilde{\text{Prim}}_{-1}(A) \rightarrow \cdots. \end{aligned}$$

(ii) If $\Sigma^{-1}A[F_\mu]$ is stably flat (ie if $\text{Tor}_*^{A[F_\mu]}(\Sigma^{-1}A[F_\mu], \Sigma^{-1}A[F_\mu]) = 0$ for $* \geq 1$) the exact sequences and the splitting in (i) extend to the left, involving the algebraic K -groups K_n for $n \geq 2$, with

$$K_*(\Sigma^{-1}A[F_\mu]) = K_*(A) \oplus \widetilde{\text{Sei}}_{*-1}(A).$$

For $\mu = 1$ $Sei(A)$ is the exact category $\mathcal{E}nd(A)$ of f.g. projective A -modules P with an endomorphism $e: P \rightarrow P$, and

$$Sei_*(A) = K_*(\mathcal{E}nd(A)) = \text{End}_*(A) = K_*(A) \oplus \widetilde{\text{End}}_*(A).$$

The special case of [Theorem E](#) (i)

$$K_1(\Sigma^{-1}A[z, z^{-1}]) = K_1(A) \oplus \widetilde{\text{End}}_0(A)$$

is the splitting theorem of Ranicki [\[33, 10.21\]](#).

We are grateful to Pere Ara, Warren Dicks, Marco Schlichting and the referee for helpful comments on the preprint version of the paper, which have led to various improvements. In particular, it was Pere Ara who pointed out that the Blanchfield $A[F_\mu]$ -module category $\mathcal{B}la_\infty(A)$ is the same as the F_μ -link module category $\mathcal{F}lk_\infty(A)$ of [\[41\]](#).

1 Combinatorial transversality for F_μ -covers

For $\mu \geq 1$ let $F_\mu = \langle z_1, z_2, \dots, z_\mu \rangle$ be the free group with generators z_1, z_2, \dots, z_μ .

1.1 F_μ -covers

Definition 1.1 An F_μ -cover of a space W is a regular covering $p: \widetilde{W} \rightarrow W$ with group of covering translations F_μ .

A classifying space BF_μ for F_μ -covers is a connected space such that

$$\pi_j(BF_\mu) = \begin{cases} F_\mu & \text{if } j = 1 \\ 0 & \text{if } j \geq 2. \end{cases}$$

The universal cover of BF_μ is an F_μ -cover

$$p_\mu: EF_\mu = \widetilde{BF}_\mu \rightarrow BF_\mu$$

with EF_μ a contractible space with a free F_μ -action.

Proposition 1.2 (i) Given an F_μ -cover $p: \widetilde{W} \rightarrow W$ and a map $f: V \rightarrow W$ there is defined a pullback square

$$\begin{array}{ccc} \widetilde{V} & \xrightarrow{\tilde{f}} & \widetilde{W} \\ f^*p \downarrow & & \downarrow p \\ V & \xrightarrow{f} & W \end{array}$$

with

$$\begin{aligned} \tilde{V} &= f^* \tilde{W} = \{(x, y) \in V \times \tilde{W} \mid f(x) = p(y) \in W\}, \\ f^* p: \tilde{V} &\rightarrow V; (x, y) \mapsto x, \quad \tilde{f}: \tilde{V} \rightarrow \tilde{W}; (x, y) \mapsto y \end{aligned}$$

such that $f^* p: \tilde{V} \rightarrow V$ is the pullback F_μ -cover.

(ii) The F_μ -covers $p: \tilde{W} \rightarrow W$ of a space W are classified by the homotopy classes of maps $c: W \rightarrow BF_\mu$ with

$$\begin{aligned} \tilde{W} &= c^* EF_\mu = \{(x, y) \in W \times EF_\mu \mid c(x) = [y] \in BF_\mu\}, \\ p(x, y) &= c^* p_\mu(x, y) = x. \end{aligned}$$

For a connected space W the homotopy classes of maps $c: W \rightarrow BF_\mu$ are in one-one correspondence with the morphisms $c_*: \pi_1(W) \rightarrow F_\mu$; the connected F_μ -covers \tilde{W} correspond to surjections $c_*: \pi_1(W) \rightarrow F_\mu$.

Proof Standard. □

1.2 The Cayley tree G_μ


We shall be working with the following explicit constructions of BF_μ and EF_μ , as well as the Cayley tree of F_μ :

Definition 1.3 The Cayley tree G_μ is the tree with vertex set

$$G_\mu^{(0)} = F_\mu$$

and edge set

$$G_\mu^{(1)} = \{(g, gz_i) \mid g \in F_\mu, 1 \leq i \leq \mu\} \subset G_\mu^{(0)} \times G_\mu^{(0)}.$$



Define a transitive F_μ -action on G_μ

$$F_\mu \times G_\mu \rightarrow G_\mu; (g, x) \mapsto gx$$

with quotient the one-point union of μ circles

$$G_\mu / F_\mu = BF_\mu = \bigvee_{\mu} S^1.$$

Let

$$I_\mu = \bigcup_{i=1}^{\mu} [e_i^-, e_i^+] \subset \mathbb{R}^\mu$$

with

$$e_i^+ = (0, \dots, 0, 1, 0, \dots, 0), \quad e_i^- = (0, \dots, 0, -1, 0, \dots, 0) \in \mathbb{R}^\mu,$$

$$[e_i^-, e_i^+] = \{(0, \dots, 0, t, 0, \dots, 0) \mid -1 \leq t \leq 1\} \subset \mathbb{R}^\mu.$$

Thus I_μ is the one-point union of μ copies of the interval $[-1, 1] \subset \mathbb{R}$, identifying the μ copies of $0 \in [-1, 1]$.

We regard BF_μ as the quotient space of I_μ

$$BF_\mu = I_\mu / \{e_i^+ \sim e_i^- \mid 1 \leq i \leq \mu\} = \bigvee_{\mu} S^1,$$

the one-point union of μ copies of the circle $S^1 = [-1, 1]/(-1 \sim 1)$ in which the μ copies of $[0] \in S^1$ are identified, with

$$e_i = [e_i^+] = [e_i^-] \neq [0] \in BF_\mu$$

a point in the i^{th} circle. The universal cover EF_μ of BF_μ is

$$EF_\mu = (F_\mu \times I_\mu) / \{(g, e_i^+) \sim (gz_i, e_i^-) \mid g \in F_\mu, 1 \leq i \leq \mu\},$$

a contractible space with a free F_μ -action

$$F_\mu \times EF_\mu \rightarrow EF_\mu; (g, (h, x)) \mapsto (gh, x)$$

and covering projection

$$p_\mu: EF_\mu \rightarrow BF_\mu; [g, x] \mapsto [x].$$

Define an F_μ -equivariant homeomorphism $G_\mu \xrightarrow{\cong} EF_\mu$ by sending the vertex $g \in G_\mu^{(0)} = F_\mu$ to the point $(g, 0) \in EF_\mu$, and the edge $(g, gz_i) \in G_\mu^{(1)}$ to the line segment

$$\{(g, te_i^+) \mid 0 \leq t \leq 1\} \cup \{(gz_i, te_i^-) \mid 0 \leq t \leq 1\} \subset EF_\mu$$

with endpoints $(g, 0), (gz_i, 0) \in EF_\mu$. The projection $G_\mu \rightarrow G_\mu/F_\mu$ can thus be identified with the universal cover $p_\mu: EF_\mu \rightarrow BF_\mu$.


1.3 Fundamental domains

Definition 1.4 A *fundamental domain* of an F_μ -cover $p: \tilde{W} \rightarrow W$ is a closed subspace $U \subset \tilde{W}$ such that

- (a) $F_\mu U = \tilde{W}$, or equivalently $p(U) = W$,
- (b) for any $g, h \in F_\mu$

$$gU \cap hU = \begin{cases} gV_i & \text{if } g = hz_i \\ hV_i & \text{if } g = hz_i^{-1} \\ gU & \text{if } g = h \\ \emptyset & \text{otherwise} \end{cases}$$

with $V_i = U \cap z_i^{-1}U$.



Thus $U \subset \tilde{W}$ is sufficiently large for the translates $gU \subset \tilde{W}$ ($g \in F_\mu$) to cover \tilde{W} , but sufficiently small for the overlaps $gU \cap hU$ to be non-empty only if $g^{-1}h = 1$ or z_i or z_i^{-1} .

Example 1.5 (i) The subspace $(1, I_\mu) \subset EF_\mu$ is a fundamental domain of the universal cover $p_\mu: EF_\mu \rightarrow BF_\mu$.

(ii) Let G'_μ be the barycentric subdivision of the Cayley tree G_μ , the tree with

$$(G'_\mu)^{(0)} = G_\mu^{(0)} \cup G_\mu^{(1)},$$


$$(G'_\mu)^{(1)} = \{(h, (g, gz_i)) \mid h = g \text{ or } gz_i\} \subset (G'_\mu)^{(0)} \times (G'_\mu)^{(0)}.$$

The F_μ -equivariant homeomorphism $G_\mu = G'_\mu \cong EF_\mu$ sends the vertex $(g, gz_i) \in (G'_\mu)^{(0)}$ to $(g, e_i^+) \in EF_\mu$. The subgraph $U_\mu \subset G'_\mu$ defined by

$$U_\mu^{(0)} = \{1\} \cup \{(1, z_i)\} \cup \{(z_i^{-1}, 1)\}$$

$$U_\mu^{(1)} = \{(1, (1, z_i))\} \cup \{(z_i^{-1}, (z_i^{-1}, 1))\}$$

is the fundamental domain of the cover $G_\mu \rightarrow G_\mu/F_\mu$ corresponding to $(1, I_\mu) \subset EF_\mu$ under the G_μ -equivariant homeomorphism $G_\mu \cong EF_\mu$.



Proposition 1.6 (i) Given an F_μ -cover $p: \tilde{W} \rightarrow W$ and a map $f: V \rightarrow W$ let $f^*p: \tilde{V} = f^*\tilde{W} \rightarrow V$ be the pullback F_μ -cover. If $U \subset \tilde{W}$ is a fundamental domain of p then

$$\tilde{f}^{-1}(U) = \{(x, y) \mid x \in V, y \in U, f(x) = p(y) \in W\} \subset \tilde{V}$$

is a fundamental domain of $f^* p$.

(ii) Every F_μ -cover $p: \tilde{W} \rightarrow W$ has fundamental domains.

Proof (i) By construction.

(ii) Apply (i), using the fundamental domain $U_\mu \subset G'_\mu = EF_\mu$ for the cover

$$p_\mu: EF_\mu \rightarrow EF_\mu/F_\mu = BF_\mu$$

given by [Example 1.5](#), noting that

$$p = c^* p_\mu: \tilde{W} = c^* EF_\mu \rightarrow W$$

is the pullback of the universal F_μ -cover $p_\mu: EF_\mu \rightarrow BF_\mu$ along a classifying map $c: W \rightarrow BF_\mu$

$$\begin{array}{ccc} \tilde{W} & \xrightarrow{\tilde{c}} & EF_\mu \\ p \downarrow & & \downarrow p_\mu \\ W & \xrightarrow{c} & BF_\mu \end{array}$$

The inverse image of $U_\mu \subset EF_\mu$

$$U = \tilde{c}^{-1}(U_\mu) \subset \tilde{W}$$

is a fundamental domain of $c: \tilde{W} \rightarrow W$. □

1.4 Combinatorial transversality

If $p: \tilde{W} \rightarrow W$ is an F_μ -cover of a space W with an additional structure such as a manifold or finite CW complex, we should like to have fundamental domains $U \subset \tilde{W}$ with the additional structure. For manifolds this is achieved by choosing a classifying map $c: W \rightarrow BF_\mu$ transverse at $\{e_1, e_2, \dots, e_\mu\} \subset BF_\mu$ – see [Example 1.11](#) below for a more detailed discussion. For a finite CW complex W we shall develop a combinatorial version of transversality, constructing finite subcomplexes $X \subset X(\infty)$ of the Borel construction $X(\infty) = \tilde{W} \times_{F_\mu} G_\mu$, such that the projection $f(\infty): W(\infty) \rightarrow W$ restricts to a simple homotopy equivalence $f: X \rightarrow W$ such that the pullback F_μ -cover $\tilde{X} = f^* \tilde{W} \rightarrow X$ has a fundamental domain $U \subset \tilde{X}$ which is a finite subcomplex.

Proposition 1.7 For any F_μ -cover $p: \tilde{W} \rightarrow W$ let F_μ act diagonally on $\tilde{W} \times G_\mu$

$$F_\mu \times (\tilde{W} \times G_\mu) \rightarrow (\tilde{W} \times G_\mu); (g, (x, y)) \mapsto (gx, gy).$$

(i) The map

$$\pi: X = \widetilde{W} \times_{F_\mu} G_\mu \rightarrow W; [x, g] \mapsto p(x)$$

is the projection of a fibration

$$G_\mu \longrightarrow X \xrightarrow{\pi} W$$

with contractible point inverses; for each $x \in \widetilde{W}$ there is defined a homeomorphism

$$G_\mu \rightarrow \pi^{-1} p(x); g \mapsto [x, g].$$

In particular, π is a homotopy equivalence.

(ii) The pullback F_μ -cover of X

$$\pi^* p: \widetilde{X} = p^* \widetilde{W} = \widetilde{W} \times G_\mu \rightarrow X = \widetilde{W} \times_{F_\mu} G_\mu$$

has fundamental domain $\widetilde{W} \times U \subset \widetilde{X} = \widetilde{W} \times G_\mu$, with $U \subset G_\mu$ any fundamental domain.

Proof Standard. □

Definition 1.8 (i) An F_μ -splitting (X, Y, Z, h) of a space W is a homeomorphism $h: X \rightarrow W$ from a space with a decomposition

$$X = Y \times [-1, 1] \cup_{Y \times \{-1, 1\}} Z$$

with $Y = Y_1 \sqcup Y_2 \sqcup \dots \sqcup Y_\mu$ the disjoint union of spaces Y_1, Y_2, \dots, Y_μ and $Y \times [-1, 1]$ attached to Z along maps

$$\alpha_i^-: Y_i \times \{-1\} \rightarrow Z, \alpha_i^+: Y_i \times \{1\} \rightarrow Z.$$

(ii) An F_μ -splitting (X, Y, Z, h) of a connected space W is *connected* if each of $Y_1, Y_2, \dots, Y_\mu, Z$ is non-empty and connected.

Proposition 1.9 Let W be a space with an F_μ -splitting (X, Y, Z, h) .

(i) The F_μ -splitting determines an F_μ -cover $p: \widetilde{W} \rightarrow W$ with

$$\begin{aligned} \widetilde{W} &= (F_\mu \times (Y \times [-1, 1] \sqcup Z)) / \sim, \\ (g, y_i, 1) &\sim (z_i g, \alpha_i^+(y_i, 1)), \\ (g, y_i, -1) &\sim (g, \alpha_i^-(y_i, -1)) \quad (g \in F_\mu, y_i \in Y_i, 1 \leq i \leq \mu), \\ p: \widetilde{W} &\rightarrow W; (g, x) \mapsto [h(x)]. \end{aligned}$$

The subspace

$$Z' = (1, Y \times [0, 1]) \cup (1, Z) \cup \bigcup_{i=1}^{\mu} (z_i, Y_i \times [-1, 0]) \subset \widetilde{W}$$

is a fundamental domain of $p: \widetilde{W} \rightarrow W$.

(ii) If there exists a homeomorphism $\phi: Z' \rightarrow Z$ such that

$$\phi(1, y_i, 0) = \alpha_i^-(y_i, -1), \phi(z_i, y_i, 0) = \alpha_i^+(y_i, 1) \quad (y_i \in Y_i, 1 \leq i \leq \mu)$$

the identification space

$$\widetilde{W}' = (F_{\mu} \times Z) / (g, \alpha_i^-(y_i)) \sim (z_i g, \alpha_i^+(y_i))$$

is such that there is defined a homeomorphism

$$(1, \phi): \widetilde{W} \rightarrow \widetilde{W}'; (g, x) \mapsto (g, \phi(x))$$

so that

$$p' = p(1, \phi)^{-1}: \widetilde{W}' \rightarrow W; (g, x) \mapsto p\phi^{-1}(x)$$

is an F_{μ} -cover of W which is isomorphic to $p: \widetilde{W} \rightarrow W$, with fundamental domain

$$(1, \phi)(Z') = (1, Z) \subset \widetilde{W}'.$$

(iii) The fundamental group of a connected space W with a connected F_{μ} -splitting (X, Y, Z, h) is an amalgamated free product

$$\pi_1(W) = \pi_1(Z) * F_{\mu} / \{ \alpha_i^+(g_i) z_i = z_i \alpha_i^-(g_i) \mid g_i \in \pi_1(Y_i), 1 \leq i \leq \mu \}.$$

The surjection $\pi_1(W) \rightarrow F_{\mu}$ is induced by a map $c: W \rightarrow BF_{\mu}$ sending $h(Y_i \times \{0\}) \subset W$ to $\{e_i\} \subset BF_{\mu}$. The surjection $\pi_1(W) \rightarrow F_{\mu}$ classifies the connected F_{μ} -cover $p: \widetilde{W} \rightarrow W$ in (i).

Proof (i) and (ii) follow by construction.

(iii) follows from the Seifert–van Kampen theorem and obstruction theory. □

Example 1.10 Define an F_{μ} -splitting $(H_{\mu}, \{1, 2, \dots, \mu\}, I_{\mu}, f)$ of BF_{μ} by

$$H_{\mu} = \{1, 2, \dots, \mu\} \times [-1, 1] \cup_{(i,1) \leftrightarrow e_i^+, (i,-1) \leftrightarrow e_i^-} I_{\mu},$$

$$f: H_{\mu} \rightarrow BF_{\mu}; \begin{cases} (i, t) \mapsto [(1 - t/2)e_i^+] & \text{for } 0 \leq t \leq 1 \\ (i, t) \mapsto [(1 + t/2)e_i^-] & \text{for } -1 \leq t \leq 0 \\ u \mapsto u/2 & \text{for } u \in I_{\mu} \end{cases}$$

with

$$f(i, 0) = e_i, \quad f(i, 1) = e_i^+/2, \quad f(i, -1) = e_i^-/2.$$

The corresponding F_μ -cover of BF_μ is the universal F_μ -cover $\widetilde{BF}_\mu = G_\mu \rightarrow BF_\mu$, with fundamental domain $I_\mu = (1, I_\mu) \subset G_\mu$. Note that $f(I_\mu) = J_\mu$, with $J_\mu \subset I_\mu$ the homeomorphic copy of I_μ defined by

$$J_\mu = \{(0, \dots, 0, t, 0, \dots, 0) \in I_\mu \mid -1/2 \leq t \leq 1/2\}.$$

A subspace $Y \subset X$ is *collared* if the inclusion $i: Y \rightarrow X$ extends to an embedding $j: Y \times [0, 1] \rightarrow X$, with $i(y) = j(y, 0) \in X$ for $y \in Y$. In particular, $\partial Z \subset Z$ is collared, for any manifold with boundary $(Z, \partial Z)$.

Example 1.11 Use the F_μ -splitting $(H_\mu, \{1, 2, \dots, \mu\}, I_\mu, f)$ of BF_μ given by Example 1.10 to identify

$$BF_\mu = H_\mu = \{1, 2, \dots, \mu\} \times [-1, 1] \cup_{\{1, 2, \dots, \mu\} \times \{-1, 1\}} I_\mu.$$

If $p: \widetilde{X} \rightarrow X$ is an F_μ -cover of a manifold X it is possible to choose a classifying map

$$c: X \rightarrow BF_\mu = \{1, 2, \dots, \mu\} \times [-1, 1] \cup_{\{1, 2, \dots, \mu\} \times \{-1, 1\}} I_\mu$$

which is transverse regular at $\{e_1, e_2, \dots, e_\mu\} \subset BF_\mu$, with the inverse images of $e_i = (i, 0) \in BF_\mu$ disjoint framed codimension-1 submanifolds

$$Y_i = c^{-1}(e_i) \subset X \quad (1 \leq i \leq \mu).$$

Cutting X along

$$Y = c^{-1}\{e_1, e_2, \dots, e_\mu\} = Y_1 \sqcup Y_2 \sqcup \dots \sqcup Y_\mu \subset X$$

there is obtained an F_μ -splitting $(X, Y, Z, \text{id.})$ of X , so that

$$X = Y \times [-1, 1] \cup_{Y \times \{-1, 1\}} Z$$

with $Y = Y \times \{0\} \subset X$ a framed codimension-1 submanifold, and $Z = c^{-1}(I_\mu) \subset X$ a codimension-0 submanifold with

$$\alpha_i^+: Y_i \times \{1\} \rightarrow Z, \quad \alpha_i^-: Y_i \times \{-1\} \rightarrow Z$$

components of the inclusion of the boundary $\partial Z = Y \times \{-1, 1\} \subset Z$. Since $\partial Z \subset Z$ is collared the fundamental domain of the F_μ -cover $\widetilde{X} = c^*G_\mu$

$$Z' = (1, Y \times [0, 1]) \cup (1, Z) \cup \bigcup_{i=1}^{\mu} (z_i, Y_i \times [-1, 0]) \subset \widetilde{X}$$

is such that there exists a homeomorphism $\phi: Z' \rightarrow Z$ with

$$\phi(1, y_i, 0) = \alpha_i^-(y_i, -1), \quad \phi(z_i, y_i, 0) = \alpha_i^+(y_i, 1) \quad (y_i \in Y_i, 1 \leq i \leq \mu).$$

Thus by Proposition 1.9 (ii) $p: \tilde{X} \rightarrow X$ is isomorphic to the F_μ -cover $p': \tilde{X}' \rightarrow X$ with

$$\begin{aligned} \tilde{X}' &= (F_\mu \times Z)/(g, \alpha_i^-(y_i)) \sim (z_i g, \alpha_i^+(y_i)), \\ p' &= p(1, \phi)^{-1}: \tilde{X}' \rightarrow X; (g, x) \mapsto p\phi^{-1}(x). \end{aligned}$$

If X and \tilde{X} are connected it is possible to choose c such that each $Y_i = p^{-1}(e_i)$ is connected, with

$$p_* = p(Y, Z)_*: \pi_1(X) \rightarrow F_\mu.$$

Definition 1.12 (i) A homotopy F_μ -splitting (X, Y, Z, h) of a space W is a homotopy equivalence $h: X \rightarrow W$ from a space with an F_μ -splitting $(X, Y, Z, 1)$, so that

$$X = Y \times [-1, 1] \cup_{Y \times \{-1, 1\}} Z, \quad Y = Y_1 \sqcup Y_2 \sqcup \dots \sqcup Y_\mu.$$

(ii) A homotopy F_μ -splitting (X, Y, Z, h) of a finite CW complex W is simple if X is a finite CW complex, $Y_1, Y_2, \dots, Y_\mu, Z \subset X$ are subcomplexes and $h: W \rightarrow X$ is a simple homotopy equivalence.

Example 1.13 Any finite CW complex W with an F_μ -cover $\tilde{W} \rightarrow W$ admits simple homotopy F_μ -splittings (X, Y, Z, h) : embed $W \subset S^N$ (N large) with closed regular neighbourhood $(X, \partial X)$ and apply the manifold transversality of Example 1.11 to the F_μ -cover $\tilde{X} \simeq \tilde{W} \rightarrow W \simeq X$.

Working as in Ranicki [35] we shall now develop a combinatorial transversality construction of simple homotopy F_μ -splittings of W using finite subcomplexes of the Borel construction (Proposition 1.7) $\tilde{W} \times_{F_\mu} G_\mu$, as follows.

Definition 1.14 The canonical homotopy F_μ -splitting $(X(\infty), Y(\infty), Z(\infty), h(\infty))$ of a space W with an F_μ -cover $p: \tilde{W} \rightarrow W$ is given by

$$X(\infty) = Y(\infty) \times [-1, 1] \cup_{Y(\infty) \times \{-1, 1\}} Z(\infty)$$

with

$$\begin{aligned} \alpha(\infty)_i^+ &: Y(\infty)_i = \tilde{W} \rightarrow Z(\infty) = \tilde{W} \times I_\mu; x \mapsto (z_i x, e_i^+), \\ \alpha(\infty)_i^- &: Y(\infty)_i = \tilde{W} \rightarrow Z(\infty) = \tilde{W} \times I_\mu; x \mapsto (x, e_i^-), \\ h(\infty) &: X(\infty) \rightarrow W; (x, y) \mapsto p(x). \end{aligned}$$

The map $h(\infty)$ is a homotopy equivalence since it is the composite

$$h(\infty) = \pi \circ f: X(\infty) \xrightarrow{f} \widetilde{W} \times_{F_\mu} G_\mu \xrightarrow{\pi} W$$

of the homeomorphism

$$f: X(\infty) \rightarrow \widetilde{W} \times_{F_\mu} G_\mu; \begin{cases} (x, i, t) \mapsto (x, (1-t/2)e_i^+) & \text{for } 0 \leq t \leq 1 \\ (x, i, t) \mapsto (z_i x, (1+t/2)e_i^-) & \text{for } -1 \leq t \leq 0 \\ (x, u) \mapsto (x, u/2) & \text{for } u \in I_\mu \end{cases}$$

and the homotopy equivalence

$$\pi: X = \widetilde{W} \times_{F_\mu} G_\mu \rightarrow W$$

given by [Proposition 1.7](#). For every $y \in G_\mu$ there is a unique $g \in F_\mu$ such that $gy \in I_\mu \setminus \{e_1^+, e_2^+, \dots, e_\mu^+\}$, so that either $gy = te_i^+$ with $0 \leq t < 1$, or $gy = te_i^-$ with $0 \leq t \leq 1$, and

$$f^{-1}: \widetilde{W} \times_{F_\mu} G_\mu \rightarrow X(\infty) : \begin{cases} (gx, i, 2(1-t)) & \text{if } gy = te_i^+ \text{ with } 1/2 \leq t < 1 \\ (z_i^{-1}gx, i, 2(t-1)) & \text{if } gy = te_i^- \text{ with } 1/2 \leq t \leq 1 \\ (gx, gy) & \text{if } 2gy \in I_\mu \text{ (ie if } -1/2 \leq t \leq 1/2). \end{cases}$$

Proposition 1.15 Given a space W with F_μ -cover $p: \widetilde{W} \rightarrow W$ and a subspace $V \subseteq \widetilde{W}$ let

$$X(V) = Y(V) \times [-1, 1] \cup_{Y(V) \times \{-1, 1\}} Z(V) \subseteq X(\infty)$$

with

$$\alpha(V)_i^+: Y(V)_i = V \cap z_i^{-1}V \rightarrow Z(V) = V \times I_\mu; x \mapsto (z_i x, e_i^+),$$

$$\alpha(V)_i^-: Y(V)_i = V \cap z_i^{-1}V \rightarrow Z(V) = V \times I_\mu; x \mapsto (x, e_i^-),$$

and set

$$h(V) = h(\infty)|: X(V) \rightarrow W; (x, t) \mapsto p(x).$$

(i) For any $x \in V$

$$\begin{aligned} h(V)^{-1}(p(x)) &= \{(x, y) \in \widetilde{W} \times_{F_\mu} G_\mu \mid y \in G_\mu(V, x)\} \\ &= \{x\} \times G_\mu(V, x) \subseteq X(V) \subseteq X(\infty) = \widetilde{W} \times_{F_\mu} G_\mu \end{aligned}$$

with $G_\mu(V, x) \subseteq G_\mu$ the subgraph defined by

$$G_\mu(V, x)^{(0)} = \{g \in F_\mu \mid gx \in V\} \subseteq G_\mu^{(0)} = F_\mu,$$

$$G_\mu(V, x)^{(1)} = \{(i, g) \mid gx, gz_i x \in V\} \subseteq G_\mu^{(1)} = \{1, 2, \dots, \mu\} \times F_\mu.$$

(ii) The image of $h(V)$ is

$$h(V)(X(V)) = p(V) \subseteq W,$$

so that $h(V)$ is surjective if and only if $p(V) = W$, if and only if $\bigcup_{g \in F_\mu} gV = \widetilde{W}$.

Proof By construction. □

In particular, if $V = \widetilde{W}$ then

$$(X(V), Y(V), Z(V), h(V)) = (X(\infty), Y(\infty), Z(\infty), h(\infty))$$

and $h(V): X(V) = X(\infty) \rightarrow W$ is a homotopy equivalence (since it has contractible point inverses).

Theorem 1.16 (Combinatorial transversality) *Let W be a connected finite CW complex with a connected F_μ -cover $p: \widetilde{W} \rightarrow W$. The canonical homotopy F_μ -splitting $(X(\infty), Y(\infty), Z(\infty), h(\infty))$ of W is a union*

$$(X(\infty), Y(\infty), Z(\infty), h(\infty)) = \bigcup_{\{V\}} (X(V), Y(V), Z(V), h(V))$$

of simple homotopy F_μ -splittings $(X(V), Y(V), Z(V), h(V))$ of W , with $\{V\}$ a collection of finite subcomplexes $V \subset \widetilde{W}$ such that

$$p(V) = W, \quad \bigcup_{\{V\}} V = \widetilde{W}.$$

In particular, there exist simple homotopy F_μ -splittings of W .

Proof Let

$$W = \bigcup D^0 \cup \bigcup D^1 \cup \dots \cup \bigcup D^n$$

be the given cell structure of W , with skeleta

$$W^{(r)} = \bigcup D^0 \cup \bigcup D^1 \cup \dots \cup \bigcup D^r.$$

The characteristic maps $D^r \rightarrow W$ of the r -cells restrict to embeddings $D^r \setminus S^{r-1} \subset W$ on the interiors, and as a set W is the disjoint union of the interiors

$$W = \bigsqcup D^0 \sqcup \bigsqcup (D^1 \setminus S^0) \sqcup \dots \sqcup \bigsqcup (D^n \setminus S^{n-1}).$$

Choose a lift of each r -cell D^r in W to an r -cell \tilde{D}^r in \tilde{W} , so that

$$\tilde{W} = \bigcup_{g \in F_\mu} \bigcup_g \tilde{D}^0 \cup \bigcup_{g \in F_\mu} \bigcup_g \tilde{D}^1 \cup \dots \cup \bigcup_{g \in F_\mu} \bigcup_g \tilde{D}^n.$$

Write $\phi: S^r \rightarrow W^{(r)}$ for the attaching maps of the $(r+1)$ -cells in W , and let $\tilde{\phi}: S^r \rightarrow \tilde{W}^{(r)}$ be the attaching maps of the chosen lifted $(r+1)$ -cells in \tilde{W} . For any subtree $T_n \subseteq G_\mu$ there exists a sequence of subtrees $T_r \subseteq G_\mu$ for $r = n-1, n-2, \dots, 0$ such that

$$\tilde{\phi}(S^r) \subseteq \tilde{W}^{(r-1)} \cup \bigcup_{g_r \in T_r^{(0)}} g_r \tilde{D}^r. \tag{*}$$

The sequence $T = (T_n, T_{n-1}, \dots, T_0)$ determines a subcomplex

$$V\langle T \rangle = \bigcup_{g_0 \in T_0^{(0)}} \bigcup_{g_0} \tilde{D}^0 \cup \bigcup_{g_1 \in T_1^{(0)}} \bigcup_{g_1} \tilde{D}^1 \cup \dots \cup \bigcup_{g_n \in T_n^{(0)}} \bigcup_{g_n} \tilde{D}^n \subseteq \tilde{W}$$

such that $p(V\langle T \rangle) = W$. The map $h(V\langle T \rangle): X(V\langle T \rangle) \rightarrow W$ constructed in [Proposition 1.15](#) is surjective, with contractible point inverses

$$h(V\langle T \rangle)^{-1}(p(x)) = G_\mu(V, x) = T_r \quad (p(x) \in D^r \setminus S^{r-1} \subset W),$$

so that it is a homotopy equivalence and $(X(V\langle T \rangle), Y(V\langle T \rangle), Z(V\langle T \rangle), h(V\langle T \rangle))$ is a homotopy F_μ -splitting of W . For the maximal sequence $T = (G_\mu, G_\mu, \dots, G_\mu)$ $V\langle T \rangle = \tilde{W}$ and we have the canonical homotopy F_μ -splitting $(X(\infty), Y(\infty), Z(\infty), h(\infty))$ of W . Any finite subtree $T_n \subset G_\mu$ can be used to start a sequence $T = (T_n, T_{n-1}, \dots, T_0)$ of finite subtrees $T_r \subset G_\mu$ satisfying $(*)$, since for each $r = n, n-1, \dots, 1$ the r -cells $\tilde{D}^r \rightarrow \tilde{W}$ are attached to a finite subcomplex of the $(r-1)$ -skeleton $\tilde{W}^{(r-1)}$. For a sequence T of finite subtrees $(X(V\langle T \rangle), Y(V\langle T \rangle), Z(V\langle T \rangle), h(V\langle T \rangle))$ is a simple homotopy F_μ -splitting of W . Finally, note that G_μ is a union of finite subtrees $T_n \subset G_\mu$, so that $(F_\mu, F_\mu, \dots, F_\mu)$ is a union of sequences $T = (T_n, T_{n-1}, \dots, T_0)$ of finite subtrees $T_r \subset G_\mu$ satisfying $(*)$, with corresponding expressions

$$\begin{aligned} \tilde{W} &= \bigcup_T V\langle T \rangle, \\ (X(\infty), Y(\infty), Z(\infty), h(\infty)) &= \bigcup_T (X(V\langle T \rangle), Y(V\langle T \rangle), Z(V\langle T \rangle), h(V\langle T \rangle)). \end{aligned}$$

This completes the proof. □

2 Algebraic transversality for $A[F_\mu]$ -module complexes

Algebraic transversality for $A[F_\mu]$ -module chain complexes is modelled on the combinatorial transversality for F_μ -covers of Section 1. The procedure replaces matrices with entries in $A[F_\mu]$ by (in general larger) matrices with entries of the linear type

$$a_1 + \sum_{i=1}^{\mu} a_{z_i} z_i \in A[F_\mu] \quad (a_1, a_{z_1}, \dots, a_{z_\mu} \in A).$$

Algebraic transversality can be traced back to the work of Higman, Bass–Heller–Swan, Stallings, Casson and Waldhausen on the algebraic K -theory of polynomial extensions and more general amalgamated free products. See of Ranicki [33, Chapter 7] for a treatment of algebraic transversality in the case $\mu = 1$ when $A[F_\mu] = A[z, z^{-1}]$ is the Laurent polynomial extension of A .

Definition 2.1 Given an A -module P and a set F let

$$P[F] = \bigoplus_{x \in F} xP$$

be the direct sum of copies xP of P , consisting of the formal A -linear combinations $\sum_{x \in F} xa_x$ ($a_x \in P$) with $\{x \in F \mid a_x \neq 0\}$ finite.

In particular, if F is a semigroup with 1 then $A[F]$ is a ring.

We shall be particularly concerned with the case of a free group $F = F_\mu$ or the free semigroup F_μ^+ on μ generators z_1, z_2, \dots, z_μ . Thus $F_\mu^+ \subset F_\mu$ consists of all the products $z_{i_1}^{n_1} z_{i_2}^{n_2} \dots z_{i_k}^{n_k}$ with $n_1, n_2, \dots, n_k \geq 0$. The rings $A[F_\mu]$, $A[F_\mu^+]$ are free products

$$\begin{aligned} A[F_\mu] &= A[z_1, z_1^{-1}] *_A A[z_2, z_2^{-1}] *_A \dots *_A A[z_\mu, z_\mu^{-1}], \\ A[F_\mu^+] &= A[z_1] *_A A[z_2] *_A \dots *_A A[z_\mu]. \end{aligned}$$

For any ring morphism $k: A \rightarrow B$ induction and restriction define functors

$$\begin{aligned} k_! : \{A\text{-modules}\} &\rightarrow \{B\text{-modules}\}; L \mapsto k_!L = B \otimes_A L, \\ k^! : \{B\text{-modules}\} &\rightarrow \{A\text{-modules}\}; M \mapsto k^!M = M \end{aligned}$$

such that $k_!$ is left adjoint to $k^!$, with a natural isomorphism

$$\text{Hom}_A(L, k^!M) \rightarrow \text{Hom}_B(k_!L, M); f \mapsto (b \otimes x \mapsto bf(x)).$$

Definition 2.2 An $A[F]$ -module is *induced* if it is of the form

$$P[F] = k_! P = A[F] \otimes_A P$$

for an A -module P , with $k: A \rightarrow A[F]$ the inclusion.

Proposition 2.3 Let P, Q be A -modules.

(i) There is defined a natural isomorphism of additive groups

$$\text{Hom}_A(P, Q[F]) \rightarrow \text{Hom}_{A[F]}(P[F], Q[F]); f \mapsto \left(\sum_{y \in F} yg_y \mapsto \sum_{y \in F} yf(g_y) \right).$$

(ii) There is defined a natural injection of additive groups

$$\text{Hom}_A(P, Q)[F] \rightarrow \text{Hom}_A(P, Q[F]); \sum_{x \in F} xf_x \mapsto \left(y \mapsto \sum_{x \in F} xf_x(y) \right).$$

(iii) If P is a f.g. projective A -module the injection in (ii) is also a surjection, so that the composite with the isomorphism in (i) is a natural isomorphism allowing the identification

$$\text{Hom}_A(P, Q)[F] = \text{Hom}_{A[F]}(P[F], Q[F]).$$

Proof (i) This is just the adjointness of $k_!$ and $k^!$, with $k: A \rightarrow A[F]$ the inclusion.

(ii) Obvious.

(iii) It is sufficient to consider the case $P = A$. □

Definition 2.4 Let P be an A -module which is given as a μ -fold direct sum

$$P = P_1 \oplus P_2 \oplus \cdots \oplus P_\mu$$

with idempotents $\pi_i: P \rightarrow P_i \rightarrow P$.

(i) Define the $A[F]$ -module endomorphism

$$z = \sum_{i=1}^{\mu} \pi_i z_i = \begin{pmatrix} z_1 & 0 & \cdots & 0 \\ 0 & z_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & z_\mu \end{pmatrix} : P[F] = P_1[F] \oplus P_2[F] \oplus \cdots \oplus P_\mu[F]$$

$$\longrightarrow P[F] = P_1[F] \oplus P_2[F] \oplus \cdots \oplus P_\mu[F].$$

For $F = F_\mu$ this is an automorphism, with inverse

$$z^{-1} = \sum_{i=1}^{\mu} \pi_i z_i^{-1} = \begin{pmatrix} z_1^{-1} & 0 & \cdots & 0 \\ 0 & z_2^{-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & z_\mu^{-1} \end{pmatrix} : P[F_\mu] = P_1[F_\mu] \oplus P_2[F_\mu] \oplus \cdots \oplus P_\mu[F_\mu] \\ \longrightarrow P[F_\mu] = P_1[F_\mu] \oplus P_2[F_\mu] \oplus \cdots \oplus P_\mu[F_\mu].$$

(ii) Given a collection of A -module morphisms

$$e = \{e_i \in \text{Hom}_A(P_i, Q) \mid 1 \leq i \leq \mu\}$$

define the $A[F]$ -module morphism

$$ez = \sum_{i=1}^{\mu} e \pi_i z_i = (e_1 z_1 \ e_2 z_2 \ \cdots \ e_\mu z_\mu) : P[F] = P_1[F] \oplus P_2[F] \oplus \cdots \oplus P_\mu[F] \rightarrow Q[F].$$

(iii) An $A[F]$ -module morphism $f : P[F] \rightarrow Q[F]$ is *linear* if

$$f = f^+ z - f^- = (f^{+,1} z_1 - f^{-,1} \ \cdots \ f^{+,\mu} z_\mu - f^{-,\mu}) : \\ P[F] = P_1[F] \oplus P_2[F] \oplus \cdots \oplus P_\mu[F] \rightarrow Q[F]$$

for some A -module morphisms $f^{+,i}, f^{-,i} : P_i \rightarrow Q$.

Definition 2.5 (i) A *Mayer-Vietoris presentation* of an $A[F]$ -module E is an exact sequence of the type

$$0 \longrightarrow \bigoplus_{i=1}^{\mu} C^{(i)}[F] \xrightarrow{f} D[F] \longrightarrow E \longrightarrow 0$$

with $C^{(i)}, D$ A -modules and $f = f^+ z - f^-$ a linear $A[F]$ -module morphism.

(ii) A *Mayer-Vietoris presentation* of an $A[F]$ -module morphism $\phi : E \rightarrow E'$ is a morphism of Mayer-Vietoris presentations

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C^{(i)}[F] & \xrightarrow{f} & D[F] & \longrightarrow & E \longrightarrow 0 \\ & & \downarrow \bigoplus g^{(i)} & & \downarrow h & & \downarrow \phi \\ 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C'^{(i)}[F] & \xrightarrow{f'} & D'[F] & \longrightarrow & E' \longrightarrow 0 \end{array}$$

where $g^{(i)}: C^{(i)} \rightarrow C'^{(i)}$ and $h: D \rightarrow D'$ A -module morphisms.

(iii) A Mayer–Vietoris presentation of an $A[F]$ -module chain complex E is an exact sequence as in (i), with $C^{(i)}, D$ A -module chain complexes and $f^{\pm, i}: C^{(i)} \rightarrow D$ A -module chain maps. Similarly for an $A[F]$ -module chain map $\phi: E \rightarrow E'$, with a morphism of exact sequences as in (ii) in which $g^{(i)}, h$ are A -module chain maps.

(iv) A Mayer–Vietoris presentation of a finite induced f.g. projective $A[F_\mu]$ -module chain complex E is finite if $C^{(i)}, D$ are finite f.g. projective A -module chain complexes.

Example 2.6 Let X be the CW complex

$$X = Z / \{x \sim \beta_i(x) \mid x \in Y_i^+, 1 \leq i \leq \mu\}$$

which is obtained from a CW complex Z and disjoint collared subcomplexes

$$Y_1^+, Y_2^+, \dots, Y_\mu^+, Y_1^-, Y_2^-, \dots, Y_\mu^- \subset Z$$

using cellular homeomorphisms $\beta_i: Y_i^+ \rightarrow Y_i^-$ as identifications. As in [Definition 1.8](#) there is an F_μ -splitting (X, Y, Z, h) , where $Y = Y_1^+ \sqcup Y_2^+ \sqcup \dots \sqcup Y_\mu^+$ and

$$\begin{aligned} \alpha_i^+ &= \text{inclusion}_{Y_i^+ \subset Z}: Y_i = Y_i^+ \rightarrow Z, \\ \alpha_i^- &= (\text{inclusion}_{Y_i^- \subset Z})\beta_i: Y_i = Y_i^+ \rightarrow Z. \end{aligned}$$

The cellular free $\mathbb{Z}[F_\mu]$ -module chain complex $C(\tilde{X})$ of the F_μ -cover \tilde{X} of X given by [Proposition 1.9](#) (i) has a Mayer–Vietoris presentation

$$0 \longrightarrow C(Y)[F_\mu] \xrightarrow{\alpha} C(Z)[F_\mu] \longrightarrow C(\tilde{X}) \longrightarrow 0$$

with $C(Y)^{(i)} = C(Y_i)$, $C(Z)$ free \mathbb{Z} -module chain complexes, and $\alpha = \alpha^+ z - \alpha^-$ a linear $\mathbb{Z}[F_\mu]$ -module chain map. If Z is a finite CW complex the Mayer–Vietoris presentation is finite.

We shall construct Mayer–Vietoris presentations of free $A[F_\mu]$ -module chain complexes using the Cayley tree G_μ ([Definition 1.3](#)) and the subtree $G_\mu^+ \subset G_\mu$ corresponding to $F_\mu^+ \subset F_\mu$.

Definition 2.7 (i) Let $G_\mu^+ \subset G_\mu$ be the subtree with

$$(G_\mu^+)^{(0)} = F_\mu^+, \quad (G_\mu^+)^{(1)} = \{(g, gz_i) \mid g \in F_\mu^+, 1 \leq i \leq \mu\}.$$

(ii) For any subtree $T \subseteq G_\mu$ and $i = 1, 2, \dots, \mu$ let $T^{(i,1)} \subseteq T^{(1)}$ be the set of edges of type (g, gz_i) with $g \in F_\mu$, such that

$$T^{(1)} = \coprod_{i=1}^{\mu} T^{(i,1)},$$

and let

$$T^+ = T \cap G_\mu^+ \subseteq T.$$

(iii) For $F = F_\mu$ (resp. F_μ^+) let $G = G_\mu$ (resp. G_μ^+).

We shall only be considering subtrees $T \subseteq G$ containing the vertex $1 \in G^{(0)}$.

Proposition 2.8 Given an A -module P let $E = P[F]$ be the induced $A[F]$ -module, regarded as a 0-dimensional $A[F]$ -module chain complex.

(i) For any subtree $T \subseteq G$ there is defined a Mayer-Vietoris presentation of E

$$E\langle T \rangle: 0 \longrightarrow \bigoplus_{i=1}^{\mu} C^{(i)}[F] \xrightarrow{f} D[F] \longrightarrow E \longrightarrow 0$$

with

$$D = P[T^{(0)}], \quad C^{(i)} = D \cap z_i^{-1}D = P[T^{(i,1)}] \subseteq E, \\ f^{+,i}: C^{(i)} \rightarrow D; xp \mapsto xp, \quad f^{-,i}: C^{(i)} \rightarrow D; xp \mapsto z_i xp.$$

(ii) The Mayer-Vietoris presentations $E\langle T \rangle$ are such that

$$E\langle T \cap T' \rangle = E\langle T \rangle \cap E\langle T' \rangle, \quad E\langle T \cup T' \rangle = E\langle T \rangle + E\langle T' \rangle \quad (T, T' \subseteq G).$$

If P is f.g. projective and T is finite then $C^{(i)}, D$ are f.g. projective A -modules.

(iii) Given a morphism of induced $A[F]$ -modules

$$\phi: E = P[F] \rightarrow E' = P'[F]$$

and a subtree $T \subseteq G$ let $\phi_*T \subseteq G$ be the smallest subtree such that

$$\phi(P) \subseteq P'[\phi_*T^{(0)}] \subseteq E'.$$

For any subtree $T' \subseteq G$ such that $\phi_*T \subseteq T'$ there is defined a morphism of Mayer-Vietoris presentations

$$\begin{array}{ccccccc} E\langle T \rangle : 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C^{(i)}[F] & \xrightarrow{f} & D[F] & \longrightarrow & E \longrightarrow 0 \\ & & \downarrow \bigoplus g^{(i)} & & \downarrow h & & \downarrow \phi \\ E'\langle T' \rangle : 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C'^{(i)}[F] & \xrightarrow{f'} & D'[F] & \longrightarrow & E' \longrightarrow 0 \end{array}$$

with

$$g^{(i)} = \phi|: C^{(i)} \rightarrow C'^{(i)}, \quad h = \phi|: D \rightarrow D'.$$

If P is a f.g. A -module and $T \subset G$ is finite, then so is $\phi_*T \subset G$.

Proof By construction. □

Example 2.9 The Mayer–Vietoris presentation of E associated to the minimal subtree $T = \{1\} \subset G$ is

$$0 \longrightarrow 0 \longrightarrow P[F] \xrightarrow{\text{id.}} E \longrightarrow 0.$$

Definition 2.10 The canonical Mayer–Vietoris presentation of an $A[F]$ -module chain complex E with each $E_r = P_r[F]$ an induced $A[F]$ -module

$$E\langle\infty\rangle : 0 \longrightarrow \bigoplus_{i=1}^{\mu} C^{(i)}[F] \xrightarrow{f} D[F] \longrightarrow E \longrightarrow 0$$

is the Mayer–Vietoris presentation with $E_r\langle\infty\rangle = E_r\langle T \rangle$ the Mayer–Vietoris presentation of E_r associated to the maximal subtree $T = G \subseteq G$, where

$$f^{+,i} = \text{id.}, \quad f^- = z_i : C^{(i)} = k^!E \rightarrow D = k^!E$$

with $k: A \rightarrow A[F]$ the inclusion.

Remark 2.11 (i) The canonical Mayer–Vietoris presentation can be written in terms of induction and restriction

$$E\langle\infty\rangle : 0 \longrightarrow \bigoplus_{\mu} k_!k^!E \xrightarrow{f} k_!k^!E \longrightarrow E \longrightarrow 0$$

with

$$\begin{aligned} f: \bigoplus_{\mu} k_!k^!E &\rightarrow k_!k^!E; \quad x_i \otimes y \mapsto x_i z_i \otimes y - x_i \otimes z_i y \quad (x_i \in A[F], y \in E), \\ k_!k^!E &\rightarrow E; \quad x \otimes y \mapsto xy \quad (x \in A[F], y \in E). \end{aligned}$$

(ii) The canonical Mayer–Vietoris presentation for $F = F_{\mu}$ is the algebraic analogue of the canonical homotopy F_{μ} -splitting of a space W with an F_{μ} -cover \widetilde{W} in [Definition 1.14](#).

Theorem 2.12 (Algebraic transversality for chain complexes) *Let E be an n -dimensional $A[F]$ -module chain complex*

$$E: E_n \xrightarrow{d_n} E_{n-1} \longrightarrow \cdots \longrightarrow E_1 \xrightarrow{d_1} E_0$$

with each $E_r = P_r[F]$ induced from an A -module P_r .

(i) For any sequence $T = (T_n, T_{n-1}, \dots, T_0)$ of subtrees $T_r \subseteq G$ such that

$$(d_r)_*(T_r) \subseteq T_{r-1} \quad (r = n, n-1, \dots, 1) \tag{*}$$

there is defined a Mayer–Vietoris presentation

$$E\langle T \rangle: 0 \longrightarrow \bigoplus_{i=1}^{\mu} C^{(i)}[F] \xrightarrow{f^+ z - f^-} D[F] \longrightarrow E \longrightarrow 0$$

with

$$E\langle T \rangle_r = E_r\langle T_r \rangle \quad (0 \leq r \leq n), \quad E\langle T \rangle \subseteq E\langle \infty \rangle.$$

(ii) If the A -modules P_r are f.g. projective then for any finite subtree $T_n \subseteq G$ there exists a sequence $T = (T_n, T_{n-1}, \dots, T_0)$ of finite subtrees $T_r \subseteq G$ satisfying (*), so that $E\langle T \rangle$ is a finite Mayer–Vietoris presentation of E . Thus

$$E\langle \infty \rangle = \bigcup_T E\langle T \rangle$$

with the union taken over all such sequences T . In particular, E admits a finite Mayer–Vietoris presentation.

Proof By repeated applications of Proposition 2.8, with the sequences $T = (T_n, T_{n-1}, \dots, T_0)$ the chain complex analogues of the sequences used to construct the homotopy F_μ -splittings of CW complexes in the proof of Theorem 1.16. □

This completes the proof of Theorem A of the Introduction.

3 Blanchfield and Seifert modules

3.1 The Magnus–Fox embedding

This section obtains some technical results on the Magnus–Fox embedding which we shall need to characterize Blanchfield $A[F_\mu]$ -modules, and to approximate h.d. 1 F_μ -link modules by f.g. projective Seifert A -modules.

Let $A\langle\langle x_1, x_2, \dots, x_\mu \rangle\rangle$ be the ring of A -coefficient formal power series in non-commuting indeterminates x_1, x_2, \dots, x_μ . The *Magnus–Fox embedding* is defined by

$$i: A[F_\mu] \rightarrow \widehat{A[F_\mu]} = A\langle\langle x_1, x_2, \dots, x_\mu \rangle\rangle; \quad z_j \mapsto 1 + x_j.$$

See the paper of Ara and Dicks [1] for a recent account of the Magnus–Fox embedding, including the relationship with noncommutative Cohn localization.

The augmentations $\epsilon(z_j) = 1, \hat{\epsilon}(x_j) = 0$ give rise to a commutative triangle of rings

$$\begin{array}{ccc} A[F_\mu] & \xrightarrow{i} & \widehat{A[F_\mu]} \\ & \searrow \epsilon & \swarrow \hat{\epsilon} \\ & A & \end{array}$$

Proposition 3.1 (i) For projective $\widehat{A[F_\mu]}$ -modules \widehat{K}, \widehat{L} the augmentation map

$$\hat{\epsilon}: \text{Hom}_{\widehat{A[F_\mu]}}(\widehat{K}, \widehat{L}) \rightarrow \text{Hom}_A(A \otimes_{\widehat{A[F_\mu]}} \widehat{K}, A \otimes_{\widehat{A[F_\mu]}} \widehat{L}); \hat{f} \mapsto 1 \otimes \hat{f}$$

is surjective.

(ii) A morphism $\hat{f}: \widehat{K} \rightarrow \widehat{L}$ of projective $\widehat{A[F_\mu]}$ -modules is an isomorphism if and only if the A -module morphism

$$1 \otimes \hat{f}: A \otimes_{\widehat{A[F_\mu]}} \widehat{K} \rightarrow A \otimes_{\widehat{A[F_\mu]}} \widehat{L}$$

is an isomorphism.

(iii) A morphism $f: K \rightarrow L$ of projective $A[F_\mu]$ -modules induces an $\widehat{A[F_\mu]}$ -module isomorphism

$$1 \otimes f: \widehat{A[F_\mu]} \otimes_{A[F_\mu]} K \rightarrow \widehat{A[F_\mu]} \otimes_{A[F_\mu]} L$$

if and only if the A -module morphism

$$1 \otimes f: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$$

is an isomorphism.

Proof (i) By additivity this reduces to the special case $\widehat{K} = \widehat{L} = \widehat{A[F_\mu]}$, which is just the fact that $\hat{\epsilon}: \widehat{A[F_\mu]} \rightarrow A$ is surjective.

(ii) It suffices to prove that if $1 \otimes \hat{f}$ is an A -module isomorphism then \hat{f} is an $\widehat{A[F_\mu]}$ -module isomorphism.

Consider first the special case when \widehat{K}, \widehat{L} are free $\widehat{A[F_\mu]}$ -modules, say $\widehat{A[F_\mu]}^k, \widehat{A[F_\mu]}^\ell$ for some sets k, ℓ . The augmentation map

$$\hat{\epsilon}: \text{Hom}_{\widehat{A[F_\mu]}}(\widehat{A[F_\mu]}^k, \widehat{A[F_\mu]}^\ell) \rightarrow \text{Hom}_A(A^k, A^\ell); \hat{f} \mapsto 1 \otimes \hat{f}$$

has a canonical splitting. If $1 \otimes \hat{f}$ is an isomorphism then all the entries in the matrix of the $\widehat{A[F_\mu]}$ -module morphism

$$g = 1 - (1 \otimes \hat{f})^{-1} \hat{f}: \widehat{A[F_\mu]}^k \rightarrow \widehat{A[F_\mu]}^k$$

have constant term 0, so that $1 - g = (1 \otimes \hat{f})^{-1} \hat{f}$ is an $\widehat{A[F_\mu]}$ -module isomorphism with inverse

$$(1 - g)^{-1} = 1 + g + g^2 + g^3 + g^4 + \dots: \widehat{A[F_\mu]}^k \rightarrow \widehat{A[F_\mu]}^k,$$

and $\hat{f} = (1 \otimes \hat{f})(1 - g)$ is an isomorphism.

For the general projective case apply (i) to lift $(1 \otimes \hat{f})^{-1}$ to an $\widehat{A[F_\mu]}$ -module morphism $\hat{e}: \hat{L} \rightarrow \hat{K}$. Choose a projective $\widehat{A[F_\mu]}$ -module \hat{J} such that $\hat{J} \oplus \hat{K} \oplus \hat{L}$ is a free $\widehat{A[F_\mu]}$ -module, and apply the special case to the $\widehat{A[F_\mu]}$ -module morphism

$$1 \oplus \begin{pmatrix} 0 & \hat{e} \\ \hat{f} & 0 \end{pmatrix}: \hat{J} \oplus \hat{K} \oplus \hat{L} \rightarrow \hat{J} \oplus \hat{K} \oplus \hat{L}.$$

(iii) This is a special case of (ii). □

For $j = 1, 2, \dots, \mu$ let y_j be a formal square root of z_j , so that $(y_j)^2 = z_j$. Let $F_\mu(y)$ be the free group generated by y_1, y_2, \dots, y_μ , so that $F_\mu \subset F_\mu(y)$ is the free subgroup generated by z_1, z_2, \dots, z_μ . We can identify $G_\mu^{(1,j)}$ with the subset $F_\mu y_j \subset F_\mu(y)$: the edge $(g, gz_j) \in G_\mu^{(1,j)}$ ($g \in F_\mu$) is identified with the element $gy_j^{-1} \in F_\mu(y)$.

Lemma 3.2 *If $T \subset G_\mu$ is a finite subtree then*

$$A[T^{(0)}] = A[\{1\}] \oplus \left(\bigoplus_{j=1}^{\mu} A[T^{(1,j)}](y_j^{-1} - y_j) \right) \subset A[F_\mu]. \tag{*}$$

Proof If $w \in T^{(1,j)}$ then certainly $w(y_j^{-1} - y_j) \in A[T^{(0)}]$. Let us check linear independence of the generators on the right hand side of (*). Assuming the contrary, let

$$a_1 + \sum_{gy_j^{-1} \in U} a_g g(y_j^{-1} - y_j) = 0 \in A[F_\mu]$$

be a non-trivial relation with $U \subset T^{(1)}$ non-empty and minimal. We reach a contradiction by observing that if $g(y_j)^{-1} \in U$ is a word of maximal length (in reduced form) then $a_g = 0$.

We must also show that every $v \in T^{(0)}$ is an element of the right-hand side of (*). Indeed there is a (unique) path in the tree from 1 to v defined by a sequence of edges $w_1, w_2, \dots, w_n \in T^{(1)}$ and we have

$$v = 1 + \sum_{i=1}^n w_i(y_{j(i)}^{-1} - y_{j(i)})\eta_i \in A[F_\mu]$$

if the signs $\eta_i \in \{\pm 1\}$ are chosen appropriately and $j(i)$ is such that $w_i \in T^{(1,j(i))}$. \square

Proposition 3.3 For any finite subset $S \subset F_\mu$ the inclusion $i|: A[S] \rightarrow \widehat{A[F_\mu]}$ is a split A -module injection.

Proof Since every finite S is contained in the vertex set of some finite tree we may assume that $S = T^{(0)}$ for some finite subtree $T \subset G_\mu$. We proceed by induction on $|T^{(0)}|$.

If the tree T has only one vertex then $T^{(0)} = \{1\}$ with $i(1) = 1 \in \widehat{A[F_\mu]}$ and

$$\widehat{A[F_\mu]} = A[\{1\}] \oplus \bigoplus_{i=1}^\mu \widehat{A[F_\mu]}x_i = A \oplus \bigoplus_{i=1}^\mu \widehat{A[F_\mu]}(1 - z_i^\eta) \quad (**)$$

for any $\eta \in \{\pm 1\}$, and $i|: A[\{1\}] \rightarrow \widehat{A[F_\mu]}$ is a split injection.

Suppose now that $|T^{(0)}| \geq 2$. Let $v_0 \in T^{(0)}$ be a leaf, ie a vertex to which only one edge is incident. Let $T \setminus \{v_0\}$ denote the tree obtained by removing the vertex v_0 and the incident edge. By the inductive hypothesis, $i|: A[T^{(0)} \setminus \{v_0\}] \rightarrow \widehat{A[F_\mu]}$ is a split injection; we denote the image by P .

Since v_0 is incident to precisely one edge then $v_0 = w_0 y_k^\eta$ for unique $\eta \in \{\pm 1\}$, $k \in \{1, \dots, \mu\}$ and $w_0 \in T^{(1,k)}$. Now for every j we have $T^{(1,j)} y_j^{-\eta} \subset T^{(0)} \setminus \{v_0\}$. Thus

$$\begin{aligned} T^{(1,j)}(y_j^{-1} - y_j) &= T^{(1,j)} y_j^{-\eta} (1 - y_j^{2\eta}) \eta \\ &= T^{(1,j)} y_j^{-\eta} (1 - z_j^\eta) \eta \subset (T^{(0)} \setminus \{v_0\})(1 - z_j^\eta) \eta. \end{aligned}$$

It follows from (*) that $i(A[T^{(0)}])$ is a direct summand of

$$Ai(v_0) \oplus \bigoplus_{j=1}^\mu P(1 - z_j^\eta)$$

and hence, by the following [Lemma 3.4](#), a direct summand of $\widehat{A[F_\mu]}$. \square

Lemma 3.4 Suppose P is an A -module which is a direct summand of $\widehat{A[F_\mu]}$. If $\theta \in \widehat{A[F_\mu]}$ is an element such that $\widehat{\epsilon}(\theta) = 1 \in A$ and $\eta = 1$ or -1 then

$$A\theta \oplus \left(\bigoplus_{j=1}^{\mu} P(1 - z_j^\eta) \right) \subset \widehat{A[F_\mu]}$$

is again a direct summand.

Proof We may write $\widehat{A[F_\mu]} = P \oplus Q$ for some A -module Q . Let $\eta = 1$ or -1 . Now it follows easily from (**) that

$$\begin{aligned} \widehat{A[F_\mu]} &= A\theta \oplus \left(\bigoplus_{j=1}^{\mu} \widehat{A[F_\mu]}(1 - z_j^\eta) \right) \\ &= A\theta \oplus \left(\bigoplus_{j=1}^{\mu} P(1 - z_j^\eta) \right) \oplus \left(\bigoplus_{j=1}^{\mu} Q(1 - z_j^\eta) \right) \end{aligned}$$

which completes the proof. \square

3.2 Blanchfield modules

Definition 3.5 (i) A *Blanchfield $A[F_\mu]$ -module* M is an $A[F_\mu]$ -module such that

$$\mathrm{Tor}_*^{A[F_\mu]}(A, M) = 0.$$

(ii) (Sheiham [41]) An F_μ -link module M is an $A[F_\mu]$ -module which has a 1-dimensional induced $A[F_\mu]$ -module resolution

$$0 \longrightarrow P[F_\mu] \xrightarrow{d} P[F_\mu] \longrightarrow M \longrightarrow 0$$

with P an A -module and d an $A[F_\mu]$ -module morphism such that the augmentation A -module morphism $\epsilon(d): P \rightarrow P$ is an isomorphism.

As before, let $k: A \rightarrow A[F_\mu]$ be the inclusion.

Proposition 3.6 The following conditions on an $A[F_\mu]$ -module M are equivalent:

- (i) M is a Blanchfield module,
- (ii) M is an F_μ -link module,

(iii) the A -module morphism

$$\gamma_M: \bigoplus_{\mu} k^1 M \rightarrow k^1 M; (m_1, m_2, \dots, m_{\mu}) \mapsto \sum_{i=1}^{\mu} (z_i - 1)m_i$$

is an isomorphism.

Proof The canonical Mayer–Vietoris presentation (Definition 2.10) of any $A[F_{\mu}]$ -module M is defined by

$$0 \longrightarrow \bigoplus_{\mu} k_1 k^1 M \xrightarrow{d} k_1 k^1 M \longrightarrow M \longrightarrow 0$$

with

$$d: \bigoplus_{\mu} k_1 k^1 M \rightarrow k_1 k^1 M; x_i \otimes y \mapsto x_i z_i \otimes y - x_i \otimes z_i y \quad (x_i \in A[F_{\mu}], y \in M),$$

$$k_1 k^1 M = k^1 M[F_{\mu}] \rightarrow M; x \otimes y \mapsto xy \quad (x \in A[F_{\mu}], y \in M),$$

such that d has augmentation A -module morphism

$$\epsilon(d) = -\gamma_M: \bigoplus_{\mu} k^1 M \rightarrow k^1 M.$$

Regarded as a right $A[F_{\mu}]$ -module A has a 1-dimensional f.g. free resolution

$$0 \longrightarrow \bigoplus_{i=1}^{\mu} A[F_{\mu}] \xrightarrow{\oplus(z_i - 1)} A[F_{\mu}] \xrightarrow{\epsilon} A \longrightarrow 0,$$

so that for any $A[F_{\mu}]$ -module M

$$\text{Tor}_n^{A[F_{\mu}]}(A, M) = \begin{cases} A \otimes_{A[F_{\mu}]} M = \text{coker}(\gamma_M) & \text{if } n = 0, \\ \ker(\gamma_M) & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases}$$

The equivalences (i) \iff (ii) \iff (iii) are now clear. □

Definition 3.7 (i) Let $\mathcal{B}la_{\infty}(A)$ be the category of Blanchfield $A[F_{\mu}]$ -modules, and let $\mathcal{B}la(A) \subset \mathcal{B}la_{\infty}(A)$ be the full subcategory of the h.d. 1 Blanchfield $A[F_{\mu}]$ -modules. (In view of Proposition 3.6 $\mathcal{B}la_{\infty}(A)$ is the same as the F_{μ} -link module category $\mathcal{F}lk_{\infty}(A)$ of Sheiham [41]).

(ii) Let $\mathcal{F}lk(A) \subset \mathcal{B}la(A)$ be the full subcategory of the h.d. 1 Blanchfield $A[F_{\mu}]$ -modules M such that there exists a 1-dimensional induced $A[F_{\mu}]$ -module resolution

$$0 \longrightarrow P[F_{\mu}] \xrightarrow{d} P[F_{\mu}] \longrightarrow M \longrightarrow 0$$

with P a f.g. projective A -module.

Example 3.8 (i) For a principal ideal domain A

$$K_0(A[F_\mu]) = K_0(A) = \mathbb{Z}$$

(see Bass [3]) so that

$$\text{Bla}(A) = \text{Flk}(A).$$

(ii) A finitely presented Blanchfield $\mathbb{Z}[F_\mu]$ -module is a ‘type L ’ $\mathbb{Z}[F_\mu]$ -module in the sense of Sato [37].

(iii) Given a μ -component boundary link $\ell: \bigsqcup_\mu S^n \subset S^{n+2}$ let $c: W \rightarrow W_0$ be a \mathbb{Z} -homology equivalence from the exterior W to the exterior W_0 of the trivial μ -component boundary link $\ell_0: \bigsqcup_\mu S^n \subset S^{n+2}$, with F_μ -equivariant lift $\tilde{c}: \tilde{W} \rightarrow \tilde{W}_0$ to the F_μ -covers. The homology groups $\dot{H}_*(\tilde{W}) = H_{*+1}(\tilde{c}: \tilde{W} \rightarrow \tilde{W}_0)$ are Blanchfield $\mathbb{Z}[F_\mu]$ -modules of homological dimension ≤ 2 . Each $\dot{H}_r(\tilde{W})$ has a \mathbb{Z} -contractible f.g. free $\mathbb{Z}[F_\mu]$ -module resolution of the type

$$0 \rightarrow \mathbb{Z}[F_\mu]^{a_r} \rightarrow \mathbb{Z}[F_\mu]^{b_r} \rightarrow \mathbb{Z}[F_\mu]^{c_r} \rightarrow \dot{H}_r(\tilde{W}) \rightarrow 0 \quad (0 \leq r \leq n+1)$$

with $a_r - b_r + c_r = 0$, and $\dot{H}_r(\tilde{W})/\mathbb{Z}$ -torsion is an h.d. 1 F_μ -link module (Levine [26, 3.5] for $\mu = 1$, Sato [37, 3.1] and Duval [12, 4.1] for $\mu \geq 2$). See Example 3.13 below for the construction of an $(n+1)$ -dimensional chain complex C in $\text{Sei}(\mathbb{Z})$ such that the covering $B(C)$ is an $(n+1)$ -dimensional chain complex in $\text{Flk}(\mathbb{Z})$ with $H_*(B(C)) = \dot{H}_*(\tilde{W})$.

The following Proposition 3.9 characterizes Blanchfield $A[F_\mu]$ -modules in terms of $A[F_\mu]$ -modules K such that

$$\text{Tor}_1^{A[F_\mu]}(A, K) = 0.$$

If K is a flat $A[F_\mu]$ -module then $\text{Tor}_1^{A[F_\mu]}(B, K) = 0$ for any right $A[F_\mu]$ -module B , and in particular $B = A$. If $K = P[F_\mu]$ is induced from an A -module P then

$$\text{Tor}_1^{A[F_\mu]}(A, P[F_\mu]) = \text{Tor}_1^A(A, P) = 0.$$

Proposition 3.9 (i) If M is an $A[F_\mu]$ -module with a resolution

$$0 \longrightarrow K \xrightarrow{d} L \longrightarrow M \longrightarrow 0$$

such that

$$\text{Tor}_1^{A[F_\mu]}(A, K) = \text{Tor}_1^{A[F_\mu]}(A, L) = 0$$

(e.g. the canonical Mayer–Vietoris presentation of [Definition 2.10](#)) then M is Blanchfield if and only if the A –module morphism $1 \otimes d: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$ is an isomorphism.

(ii) A morphism $d: K \rightarrow L$ of projective $A[F_\mu]$ –modules is injective and $M = \text{coker}(d)$ is a Blanchfield $A[F_\mu]$ –module if and only if the A –module morphism $1 \otimes d: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$ is an isomorphism.

Proof (i) It follows from [Proposition 3.6](#) and the commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 \text{Tor}_1^{A[F_\mu]}(A, K) = 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} K & \xrightarrow{\gamma_K} & K & \longrightarrow & A \otimes_{A[F_\mu]} K \longrightarrow 0 \\
 & & \downarrow \oplus d & & \downarrow d & & \downarrow 1 \otimes d \\
 \text{Tor}_1^{A[F_\mu]}(A, L) = 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} L & \xrightarrow{\gamma_L} & L & \longrightarrow & A \otimes_{A[F_\mu]} L \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \bigoplus_{i=1}^{\mu} M & \xrightarrow{\gamma_M} & M & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

that M is Blanchfield if and only if $1 \otimes d$ is an isomorphism.

(ii) If d is injective and M is Blanchfield then $1 \otimes d$ is an isomorphism by (i), since projective $A[F_\mu]$ –modules are flat. Conversely, if $1 \otimes d: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$ is an isomorphism then $1 \otimes d: \widehat{A[F_\mu]} \otimes_{A[F_\mu]} K \rightarrow \widehat{A[F_\mu]} \otimes_{A[F_\mu]} L$ is an isomorphism by [Proposition 3.1](#) (iii), and it follows from the injectivity of $K \rightarrow \widehat{A[F_\mu]} \otimes_{A[F_\mu]} K$, $L \rightarrow \widehat{A[F_\mu]} \otimes_{A[F_\mu]} L$ and the commutative diagram

$$\begin{array}{ccc}
 K & \xrightarrow{d} & L \\
 \downarrow & & \downarrow \\
 \widehat{A[F_\mu]} \otimes_{A[F_\mu]} K & \xrightarrow{1 \otimes d} & \widehat{A[F_\mu]} \otimes_{A[F_\mu]} L
 \end{array}$$

that $d: K \rightarrow L$ is injective. □

The *idempotent completion* $\mathcal{P}(\mathcal{E})$ of an additive category \mathcal{E} is the additive category with objects pairs $(M, p = p^2: M \rightarrow M)$ defined by projections p of objects M in \mathcal{E} , and morphisms $f: (M, p) \rightarrow (N, q)$ defined by morphisms $f: M \rightarrow N$ in \mathcal{E} such that $qfp = f: M \rightarrow N$. As usual, \mathcal{E} is *idempotent complete* if the functor $\mathcal{E} \rightarrow \mathcal{P}(\mathcal{E}); M \mapsto (M, 1)$ is an equivalence, or equivalently if for every idempotent $p = p^2: M \rightarrow M$ in \mathcal{E} there exists a direct sum decomposition $M = P \oplus Q$ with

$$p = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} : M = P \oplus Q \rightarrow M = P \oplus Q.$$

For any exact category \mathcal{E} there exists a full embedding $\mathcal{E} \subset \mathcal{A}$ in an abelian category \mathcal{A} (Gabriel–Quillen), and the idempotent completion $\mathcal{P}(\mathcal{E})$ is equivalent to the full exact subcategory of \mathcal{A} with objects $\text{im}(p)$ for objects (M, p) in $\mathcal{P}(\mathcal{E})$. For $\mathcal{E} = \mathcal{F}lk(A) \subset \mathcal{A} = \mathcal{B}la_\infty(A)$ we have that $\mathcal{P}(\mathcal{F}lk(A)) \subset \mathcal{B}la_\infty(A)$. In fact, we have:

Proposition 3.10 (i) *The exact categories $\mathcal{P}rim(A)$, $\mathcal{S}ei(A)$, $\mathcal{B}la(A)$ are idempotent complete.*

(ii) *The idempotent completion of $\mathcal{F}lk(A)$ is equivalent to $\mathcal{B}la(A)$*

$$\mathcal{P}(\mathcal{F}lk(A)) \approx \mathcal{B}la(A).$$

Proof (i) The exact categories $\mathcal{P}rim(A)$, $\mathcal{S}ei(A)$, $\mathcal{B}la(A)$ are closed under direct summands.

(ii) For any f.g. projective $A[F_\mu]$ -modules K, L the augmentation map

$$\epsilon: \text{Hom}_{A[F_\mu]}(K, L) \rightarrow \text{Hom}_A(A \otimes_{A[F_\mu]} K, A \otimes_{A[F_\mu]} L); d \mapsto 1 \otimes d$$

is surjective, by the following argument: choose f.g. projective $A[F_\mu]$ -modules K', L' such that

$$K \oplus K' = A[F_\mu]^k, L \oplus L' = A[F_\mu]^\ell$$

for some $k, \ell \geq 0$, and note that the augmentation map

$$\begin{aligned} \epsilon: \text{Hom}_{A[F_\mu]}(K \oplus K', L \oplus L') &= \text{Hom}_{A[F_\mu]}(A[F_\mu]^k, A[F_\mu]^\ell) \\ &\longrightarrow \text{Hom}_A(A \otimes_{A[F_\mu]} (K \oplus K'), A \otimes_{A[F_\mu]} (L \oplus L')) = \text{Hom}_A(A^k, A^\ell) \end{aligned}$$

is surjective. Given an h.d. 1 Blanchfield $A[F_\mu]$ -module M with a f.g. projective $A[F_\mu]$ -module resolution

$$0 \longrightarrow K \xrightarrow{d} L \longrightarrow M \longrightarrow 0$$

we know from Proposition 3.9 (i) that $1 \otimes d: A \otimes_{A[F_\mu]} K \rightarrow A \otimes_{A[F_\mu]} L$ is an A -module isomorphism. By Proposition 3.1 (i) it is possible to lift $(1 \otimes d)^{-1}$ to an $A[F_\mu]$ -module morphism $e: L \rightarrow K$, so that by Proposition 3.9 (i) e is an injection with

$$N = \text{coker}(e)$$

an h.d. 1 Blanchfield $A[F_\mu]$ -module. Let J be a f.g. projective $A[F_\mu]$ -module such that $J \oplus K \oplus L$ is f.g. free, say $A[F_\mu]^m$. The $A[F_\mu]$ -module morphism

$$f = 1 \oplus \begin{pmatrix} 0 & e \\ d & 0 \end{pmatrix} : J \oplus K \oplus L = A[F_\mu]^m \rightarrow J \oplus K \oplus L = A[F_\mu]^m$$

is such that $1 \otimes f: A^m \rightarrow A^m$ is an isomorphism, so that $\text{coker}(f) = M \oplus N$ is an h.d. 1 F_μ -link module. The functor

$$\mathcal{Flk}(A) \rightarrow \mathcal{Bla}(A); M \mapsto M$$

is a full embedding such that every object in $\mathcal{Bla}(A)$ is a direct summand of an object in $\mathcal{Flk}(A)$, so that $\mathcal{Bla}(A)$ is (equivalent to) the idempotent completion $\mathcal{P}(\mathcal{Flk}(A))$. \square

3.3 Seifert modules

Let Q_μ be the complete quiver which has μ vertices and μ^2 arrows, one arrow between each ordered pair of vertices. The path ring is given by

$$Q_\mu = \mathbb{Z}[e] * \mathbb{Z}[\pi_1, \pi_2, \dots, \pi_\mu \mid \pi_i \pi_j = \delta_{ij} \pi_i, \sum_{i=1}^\mu \pi_i = 1]$$

where $\pi_i e \pi_j$ corresponds to the unique path of length 1 from the i th vertex to the j th vertex. An A -module P together with a ring morphism $\rho: Q_\mu \rightarrow \text{End}_A(P)$ is essentially the same as a triple $(P, e, \{\pi_i\})$ with $e: P \rightarrow P$ an endomorphism, and $\{\pi_i: P \rightarrow P\}$ a complete system of μ idempotents. (Such representations of Q_μ were first considered by Farber [14] for particular A .)

Definition 3.11 (i) A *Seifert A -module* $(P, e, \{\pi_i\})$ is an A -module P together with an endomorphism $e: P \rightarrow P$, and a system $\{\pi_i: P \rightarrow P\}$ of idempotents expressing P as a μ -fold direct sum, with

$$\pi_i: P = P_1 \oplus P_2 \oplus \dots \oplus P_\mu \rightarrow P; (x_1, x_2, \dots, x_\mu) \mapsto (0, \dots, 0, x_i, 0, \dots, 0).$$

(ii) A *morphism* of Seifert A -modules

$$g: (P, e, \{\pi_i\}) \rightarrow (P', e', \{\pi'_i\})$$

is an A -module morphism such that

$$ge = e'g, \quad g\pi_i = \pi'_i g: P \rightarrow P'.$$

The conditions $g\pi_i = \pi'_i g$ are equivalent to g preserving the direct sum decompositions, so that

$$g = \begin{pmatrix} g_1 & 0 & \dots & 0 \\ 0 & g_2 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & g_\mu \end{pmatrix} : P = P_1 \oplus P_2 \oplus \dots \oplus P_\mu \rightarrow P' = P'_1 \oplus P'_2 \oplus \dots \oplus P'_\mu$$

with $g_i: P_i \rightarrow P'_i$.

(iii) The Seifert A -module category $Sei_\infty(A)$ has objects Seifert A -modules and morphisms as in (ii). Let $Sei(A) \subseteq Sei_\infty(A)$ be the full subcategory of the Seifert A -modules $(P, e, \{\pi_i\})$ with P f.g. projective.

3.4 The covering functor B

Seifert modules determine F_μ -link modules by:

Definition 3.12 (i) The covering of a Seifert A -module $(P, e, \{\pi_i\})$ is the F_μ -link module

$$B(P, e, \{\pi_i\}) = \text{coker}(1 - e + ez: P[F_\mu] \rightarrow P[F_\mu])$$

with Mayer-Vietoris presentation

$$0 \longrightarrow \bigoplus_{i=1}^{\mu} P_i[F_\mu] \xrightarrow{d} P[F_\mu] \longrightarrow B(P, e, \{\pi_i\}) \longrightarrow 0,$$

where $d = 1 - e + ez$.

(ii) The covering of a Seifert A -module morphism $g: (P, e, \{\pi_i\}) \rightarrow (P', e', \{\pi'_i\})$ is the F_μ -link module morphism

$$B(g): B(P, e, \{\pi_i\}) \rightarrow B(P', e', \{\pi'_i\}); x \mapsto g(x)$$

resolved by

$$\begin{array}{ccccccc} 0 & \longrightarrow & P[F_\mu] & \xrightarrow{d} & P[F_\mu] & \longrightarrow & B(P, e, \{\pi_i\}) \longrightarrow 0 \\ & & \downarrow g & & \downarrow g & & \downarrow B(g) \\ 0 & \longrightarrow & P'[F_\mu] & \xrightarrow{d'} & P'[F_\mu] & \longrightarrow & B(P', e', \{\pi'_i\}) \longrightarrow 0 \end{array}$$

with $d = 1 - e + ez$, $d' = 1 - e' + e'z$.

Example 3.13 Let $\ell: \bigsqcup_{\mu} S^n \subset S^{n+2}$ be a μ -component boundary link with exterior W , so that there exists a \mathbb{Z} -homology equivalence $c: W \rightarrow W_0$ to the exterior W_0 of the trivial μ -component boundary link $\ell_0: \bigsqcup_{\mu} S^n \subset S^{n+2}$. The $(n+2)$ -dimensional f.g. free $\mathbb{Z}[F_{\mu}]$ -module chain complex

$$\dot{C}(\tilde{W}) = \mathcal{C}(\tilde{c}: C(\tilde{W}) \rightarrow C(\tilde{W}_0))_{*+1}$$

is \mathbb{Z} -contractible. For any μ -component Seifert surface $V = V_1 \sqcup V_2 \sqcup \dots \sqcup V_{\mu} \subset S^{n+2}$ for ℓ there exists a degree 1 map $V \rightarrow V_0$ to the μ -component Seifert surface $V_0 = \bigsqcup_{\mu} D^{n+1} \subset S^{n+2}$ for ℓ_0 . Let

$$\dot{C}(V_i) = \mathcal{C}(C(V_i) \rightarrow C(D^{n+1}))_{*+1}, \quad \dot{C}(V) = \sum_{i=1}^{\mu} \dot{C}(V_i).$$

The map $V \rightarrow S^{n+2} \setminus V$ pushing V off itself in the positive normal direction combines with chain level Alexander duality to induce a \mathbb{Z} -module chain map

$$e: \dot{C}(V) \rightarrow C(S^{n+2} \setminus V, \bigsqcup_{\mu} \{\text{pt.}\}) \simeq \dot{C}(V)^{n+1-*},$$

so that there is defined an $(n+1)$ -dimensional chain complex $(\dot{C}(V), e, \{\pi_i\})$ in $Sei(\mathbb{Z})$. The covering $B(\dot{C}(V), e, \{\pi_i\})$ is an $(n+1)$ -dimensional chain complex in $\mathcal{F}lk(\mathbb{Z})$, with the projection

$$\begin{aligned} \mathcal{C}(1 - e + ez: \dot{C}(V)[F_{\mu}] \rightarrow \dot{C}(V)[F_{\mu}]) &= \dot{C}(\tilde{W}) \\ \longrightarrow B(\dot{C}(V), e, \{\pi_i\}) &= \text{coker}(1 - e + ez: \dot{C}(V)[F_{\mu}] \rightarrow \dot{C}(V)[F_{\mu}]) \end{aligned}$$

a homology equivalence.

The covering construction defines a functor of exact categories

$$B_{\infty}: Sei_{\infty}(A) \rightarrow Bla_{\infty}(A); (P, e, \{\pi_i\}) \mapsto B(P, e, \{\pi_i\})$$

which restricts to a functor $B: Sei(A) \rightarrow \mathcal{F}lk(A)$.

Definition 3.14 A morphism f in $Sei_{\infty}(A)$ is a B -isomorphism if $B(f)$ is an isomorphism in $Bla_{\infty}(A)$. Let Ξ_{∞} denote the set of B -isomorphisms in $Sei_{\infty}(A)$, and let Ξ denote the set of B -isomorphisms in $Sei(A)$.

3.5 Blanchfield/Seifert algebraic transversality

We shall now use the algebraic transversality of Section 2 to establish that every h.d. 1 F_{μ} -link module M is isomorphic to the covering $B(P, e, \{\pi_i\})$ of a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$, uniquely up to morphisms in Ξ .

We refer to Sheiham [41] for the proof that $B_\infty: Sei_\infty(A) \rightarrow Flk_\infty(A)$ induces an equivalence of exact categories $\bar{B}_\infty: \Xi_\infty^{-1}Sei_\infty(A) \approx Flk_\infty(A)$. Algebraic transversality will be used to prove that the universal localization $Sei(A) \rightarrow \Xi^{-1}Sei(A)$ has a calculus of fractions, and that the covering functor $B: Sei(A) \rightarrow Flk(A)$ induces an equivalence of exact categories $\bar{B}: \Xi^{-1}Sei(A) \approx Flk(A)$.

Given an F_μ -link module M let $U(M) = (M, e_M, \{\pi_i\})$ be the Seifert A -module defined in [41] – the definition is recalled in the Introduction of this paper, along with the fact proved in [41] that B_∞ is a left adjoint of

$$U_\infty: Bla_\infty(A) \rightarrow Sei_\infty(A); M \mapsto U(M).$$

The natural isomorphism of the adjointness

$$\begin{aligned} \text{Hom}_{Bla_\infty(A)}(B(Q, f, \{\rho_i\}), M) &\xrightarrow{\cong} \text{Hom}_{Sei_\infty(A)}((Q, f, \{\rho_i\}), U(M)); \\ g &\mapsto \text{adj}(g) = U(g)h \end{aligned}$$

is defined for any Seifert A -module $(Q, f, \{\rho_i\})$, with

$$h: Q \subset Q[F_\mu] \rightarrow UB(Q, f, \{\rho_i\})$$

the restriction of the canonical surjection $Q[F_\mu] \rightarrow B(Q, f, \{\rho_i\})$. If M is h.d. 1 and $(Q, f, \{\rho_i\})$ is f.g. projective the natural isomorphism can be written as

$$\text{Hom}_{Flk(A)}(B(Q, f, \{\rho_i\}), M) \cong \text{Hom}_{Sei_\infty(A)}((Q, f, \{\rho_i\}), U(M))$$

but note that in general $U(M)$ is not a f.g. projective Seifert A -module.

The following result establishes that for an h.d. 1 F_μ -link module M the Seifert A -module $U(M)$ is the direct limit of a directed system of f.g. projective Seifert A -modules $(P, e, \{\pi_i\})$ and morphisms in Ξ , with isomorphisms $B(P, e, \{\pi_i\}) \cong M$.

Theorem 3.15 (Blanchfield/Seifert algebraic transversality) *Let M be an h.d. 1 F_μ -link module, with a 1-dimensional induced f.g. projective $A[F_\mu]$ -module resolution*

$$0 \longrightarrow P[F_\mu] \xrightarrow{d} P[F_\mu] \longrightarrow M \longrightarrow 0$$

such that $\epsilon(d): P \rightarrow P$ is an A -module isomorphism.

(i) *Let I_∞ be the set of ordered pairs $T = (T_0, T_1)$ of subtrees $T_0, T_1 \subseteq G_\mu$ such that $d_*(T_1) \subseteq T_0$. The set I_∞ is partially ordered by inclusion, with maximal element*

$$T_{\max} = \bigcup_{T \in I_\infty} T = (G_\mu, G_\mu) \in I_\infty.$$

There is defined a directed system of Seifert A -modules $(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$ and morphisms in Ξ_∞

$$\phi\langle T, T' \rangle: (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \longrightarrow (P\langle T' \rangle, e\langle T' \rangle, \{\pi_i\langle T' \rangle\}) \quad (T \subseteq T' \in I_\infty)$$

with direct limit

$$\varinjlim_{T \in I_\infty} (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) = (P\langle T_{\max} \rangle, e\langle T_{\max} \rangle, \{\pi_i\langle T_{\max} \rangle\}) = U(M).$$

For any $T = (T_0, T_1) \in I_\infty$ the morphism $\phi\langle T, T_{\max} \rangle: (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \rightarrow U(M)$ is the adjoint $\phi\langle T, T_{\max} \rangle = \text{adj}(\phi\langle T \rangle)$ of an isomorphism in $\mathcal{Flk}(A)$

$$\phi\langle T \rangle: B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \xrightarrow{\cong} M$$

such that for any $T \subseteq T' \in I_\infty$ there is defined a commutative triangle of isomorphisms in $\mathcal{Flk}_\infty(A)$

$$\begin{array}{ccc} B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) & \xrightarrow[\cong]{B(\phi\langle T, T' \rangle)} & B(P\langle T' \rangle, e\langle T' \rangle, \{\pi_i\langle T' \rangle\}) \\ & \searrow \phi\langle T \rangle \cong & \swarrow \phi\langle T' \rangle \cong \\ & & M \end{array}$$

In particular, $\phi\langle T, T_{\max} \rangle \in \Xi_\infty$.

(ii) Let $I \subset I_\infty$ be the subset of the ordered pairs $T = (T_0, T_1)$ of finite subtrees $T_0, T_1 \subset G_\mu$ such that $d_*T_1 \subseteq T_0$. For $T \in I$ $(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$ is a f.g. projective Seifert A -module, and

$$\varinjlim_{T \in I} (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) = U(M)$$

with $\phi\langle T, T' \rangle \in \Xi$ ($T \subseteq T' \in I$).

(iii) For any f.g. projective Seifert A -module $(Q, f, \{\rho_i\})$ every morphism

$$g: B(Q, f, \{\rho_i\}) \rightarrow M$$

in $\mathcal{Flk}(A)$ factors as

$$g: B(Q, f, \{\rho_i\}) \xrightarrow{B(g\langle T \rangle)} B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \xrightarrow[\cong]{\phi\langle T \rangle} M$$

for some $T \in I$, with $g\langle T \rangle: (Q, f, \{\rho_i\}) \rightarrow (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$ a morphism in $\text{Sei}(A)$.

Proof (i) The induced f.g. projective $A[F_\mu]$ -module chain complex

$$E: E_1 = P[F_\mu] \xrightarrow{d} E_0 = P[F_\mu]$$

is such that $H_0(E) = M$, $H_1(E) = 0$. By [Theorem 2.12](#) for any subtree $T_1 \subseteq G_\mu$ there exists a subtree $d_*(T_1) \subseteq G_\mu$ such that for any subtree $T_0 \subseteq G_\mu$ with $d_*(T_1) \subseteq T_0$ E admits a Mayer–Vietoris presentation

$$\begin{array}{ccccccc} E_1\langle T_1 \rangle: 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C_1^{(i)}[F_\mu] & \xrightarrow{f_1^+ z - f_1^-} & D_1[F_\mu] & \longrightarrow & E_1 \longrightarrow 0 \\ & & \downarrow d_C & & \downarrow d_D & & \downarrow d \\ E_0\langle T_0 \rangle: 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C_0^{(i)}[F_\mu] & \xrightarrow{f_0^+ z - f_0^-} & D_0[F_\mu] & \longrightarrow & E_0 \longrightarrow 0 \end{array}$$

with $C_j^{(i)} = P[T_j^{(i,1)}]$, $D_j = P[T_j^{(0)}] \subseteq E_j = P[F_\mu]$ ($j = 0, 1$),
and $d_C = \bigoplus_{i=1}^{\mu} d|: \bigoplus_{i=1}^{\mu} C_1^{(i)} \rightarrow \bigoplus_{i=1}^{\mu} C_0^{(i)}$, $d_D = d|: D_1 \rightarrow D_0$.

The A -modules defined by

$$\begin{aligned} P_i\langle T \rangle &= \text{coker}(d|: C_1^{(i)} \rightarrow C_0^{(i)}), \\ P\langle T \rangle &= \text{coker}(d_C) = \bigoplus_{i=1}^{\mu} P_i\langle T \rangle, \\ Q\langle T \rangle &= \text{coker}(d_D) \end{aligned}$$

fit into a commutative diagram of $A[F_\mu]$ -modules with exact rows and columns

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C_1^{(i)}[F_\mu] & \xrightarrow{f_1^+ z - f_1^-} & D_1[F_\mu] & \longrightarrow & P[F_\mu] \longrightarrow 0 \\ & & \downarrow d_C & & \downarrow d_D & & \downarrow d \\ 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} C_0^{(i)}[F_\mu] & \xrightarrow{f_0^+ z - f_0^-} & D_0[F_\mu] & \longrightarrow & P[F_\mu] \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & P\langle T \rangle[F_\mu] & \xrightarrow{f^+ z - f^-} & Q\langle T \rangle[F_\mu] & \longrightarrow & M \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

with $f^+, f^-: P\langle T \rangle \rightarrow Q\langle T \rangle$ the A -module morphisms induced by

$$f_0^+, f_0^-: \bigoplus_{i=1}^{\mu} C_0^{(i)} \rightarrow D_0.$$

It follows from $\text{Tor}_1^{A[F_\mu]}(A, M) = 0$ that $f^+ - f^-: P\langle T \rangle \rightarrow Q\langle T \rangle$ is an A -module isomorphism. The Seifert A -module $(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$ defined by

$$e\langle T \rangle = (f^+ - f^-)^{-1} f^+: P\langle T \rangle \rightarrow P\langle T \rangle, \quad \pi_i\langle T \rangle: P\langle T \rangle \rightarrow P_i\langle T \rangle \rightarrow P\langle T \rangle$$

is such that $P\langle T \rangle[F_\mu] \cong Q\langle T \rangle[F_\mu] \rightarrow M$ induces the isomorphism of Blanchfield $A[F_\mu]$ -modules $\phi\langle T \rangle: B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \cong M$ adjoint to the natural map $(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \rightarrow U(M)$. (In particular, $(P\langle T_{\max} \rangle, e\langle T_{\max} \rangle, \{\pi_i\langle T_{\max} \rangle\}) = U(M)$ and $\phi\langle T_{\max} \rangle: BU(M) \cong M$ is the natural isomorphism ψ_M defined in [41, 5.10].) For $T \subseteq T' \in I$ the B -isomorphism $\phi\langle T, T' \rangle$ is induced by the inclusion $T \subseteq T'$.

(ii) The augmentation of the $A[F_\mu]$ -module morphism $d: P[F_\mu] \rightarrow P[F_\mu]$ is an A -module isomorphism $\epsilon(d): P \rightarrow P$, so that the induced $\widehat{A[F_\mu]}$ -module morphism $\widehat{d}: \widehat{P[F_\mu]} \rightarrow \widehat{P[F_\mu]}$ is an isomorphism, by Proposition 3.1. For any $T = (T_0, T_1) \in I$ the inclusion $P[T_1^{(0)}] \rightarrow \widehat{P[F_\mu]}$ is a split A -module injection by Proposition 3.3. Let $s: \widehat{P[F_\mu]} \rightarrow P[T_1^{(0)}]$ be a splitting A -module surjection. The anticlockwise composition of the morphisms (inverting \widehat{d}) in the diagram

$$\begin{array}{ccccc} & & s & & \\ & & \curvearrowright & & \\ D_1 = P[T_1^{(0)}] & \longrightarrow & P[F_\mu] & \longrightarrow & \widehat{P[F_\mu]} \\ d_D = d| & \downarrow & \downarrow d & \cong & \downarrow \widehat{d} \\ D_0 = P[T_0^{(0)}] & \longrightarrow & P[F_\mu] & \longrightarrow & \widehat{P[F_\mu]} \end{array}$$

defines an A -module surjection $P[T_0^{(0)}] \rightarrow P[T_1^{(0)}]$ splitting $d|: P[T_1^{(0)}] \rightarrow P[T_0^{(0)}]$. Thus $d|$ is a split injection of f.g. projective A -modules and $P\langle T \rangle = \text{coker}(d|)$ is a f.g. projective A -module.

(iii) The morphism $g: B(Q, f, \{\rho_i\}) \rightarrow M$ in $\mathcal{Flk}(A)$ has a canonical resolution

$$\begin{array}{ccccccc} 0 & \longrightarrow & Q[F_\mu] & \xrightarrow{1-f+fz} & Q[F_\mu] & \longrightarrow & B(Q, f, \{\rho_i\}) \longrightarrow 0 \\ & & \downarrow \text{adj}(g) & & \downarrow \text{adj}(g) & & \downarrow g \\ 0 & \longrightarrow & P\langle T_{\max} \rangle[F_\mu] & \longrightarrow & P\langle T_{\max} \rangle[F_\mu] & \longrightarrow & M \longrightarrow 0 \end{array}$$


with

$$\text{adj}(g): (Q, f, \{\rho_i\}) \rightarrow (P\langle T_{\max} \rangle, e\langle T_{\max} \rangle, \{\pi_i\langle T_{\max} \rangle\}) = U(M)$$

the adjoint morphism in $\text{Sei}_\infty(A)$. Since Q is f.g. projective there exists $T \in I$ such that

$$\text{im}(g: B(Q, f, \{\rho_i\}) \rightarrow M) \subseteq \text{im}(B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) \rightarrow M)$$

with a lift of g to an A -module morphism $g\langle T \rangle: Q \rightarrow P\langle T \rangle$ which preserves the direct sum structures. The diagram of A -modules and morphisms



commutes except possibly in $(*)$, and $(*)$ commutes if and only if

$$g\langle T \rangle: (Q, f, \{\rho_i\}) \rightarrow (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$$

is a morphism of Seifert A -modules. Since Q is f.g. projective and the composite

$$Q \xrightarrow{g\langle T \rangle f - e\langle T \rangle g\langle T \rangle} P\langle T \rangle \xrightarrow{\phi\langle T \rangle} U(M) = \varinjlim_{T' \in I} P\langle T' \rangle$$

is 0 there exists $T' \in I$ such that $T \subseteq T'$ and the composite

$$g\langle T' \rangle f - e\langle T' \rangle g\langle T' \rangle: Q \xrightarrow{g\langle T' \rangle f - e\langle T' \rangle g\langle T' \rangle} P\langle T \rangle \longrightarrow P\langle T' \rangle$$

is 0, so that

$$g\langle T' \rangle: (Q, f, \{\rho_i\}) \rightarrow (P\langle T' \rangle, e\langle T' \rangle, \{\pi_i\langle T' \rangle\})$$

is a morphism of Seifert A -modules as required (except that T' has to be called T). \square

Definition 3.16 Let $M = B(P, e, \{\pi_i\})$ for a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$.

(i) For any $T \in I_\infty$ let

$$s\langle T \rangle: (P, e, \{\pi_i\}) \rightarrow (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$$

be the B -isomorphism determined by the inclusion $P = P[\{1\}] \subseteq P[T_0^{(0)}]$.

(ii) For $T = T_{\max} \in I_\infty$ write

$$s_M = s\langle T_{\max} \rangle: (P, e, \{\pi_i\}) \rightarrow (P\langle T_{\max} \rangle, e\langle T_{\max} \rangle, \{\pi_i\langle T_{\max} \rangle\}) = U(M).$$

This is the B -isomorphism adjoint of $1: M \rightarrow M$, such that

$$s_M: (P, e, \{\pi_i\}) \xrightarrow{s\langle T \rangle} (P\langle T \rangle, e, \{\pi_i\}) \xrightarrow{\phi\langle T \rangle} U(M)$$

for any $T \in I_\infty$.

Putting everything together:

Theorem 3.17 (i) Every h.d. 1 F_μ -link module M is isomorphic to the covering $B(P, e, \{\pi_i\})$ of a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$.

(ii) For any f.g. projective Seifert A -modules $(P, e, \{\pi_i\})$, $(Q, f, \{\rho_i\})$ every morphism $g: B(Q, f, \{\rho_i\}) \rightarrow B(P, e, \{\pi_i\})$ in $\mathcal{Flk}(A)$ is of the form $g = B(s)^{-1}B(t)$ for some morphisms

$$s: (P, e, \{\pi_i\}) \rightarrow (P', e', \{\pi'_i\}), t: (Q, f, \{\rho_i\}) \rightarrow (P', e', \{\pi'_i\})$$

in $\text{Sei}(A)$ with $s \in \Xi$.

(iii) If $u: (Q, f, \{\rho_i\}) \rightarrow (P, e, \{\pi_i\})$ is a morphism of f.g. projective Seifert A -modules such that $B(u) = 0$ there exists an element $v: (P, e, \{\pi_i\}) \rightarrow (P', e', \{\pi'_i\})$ in Ξ such that $vu = 0$.

(iv) The localization $\Xi^{-1}\text{Sei}(A)$ has a left calculus of fractions, and the covering construction defines an equivalence of exact categories

$$\bar{B}: \Xi^{-1}\text{Sei}(A) \xrightarrow{\cong} \mathcal{Flk}(A); (P, e, \{\pi_i\}) \mapsto B(P, e, \{\pi_i\}).$$

Proof (i) By Theorem 3.15 (i)–(ii) M is isomorphic to $B(P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$ for any $T \in I$, e.g. for the minimal element $T_{\min} = (d_*\{1\}, \{1\}) \in I$.

(ii) By Theorem 3.15 (iii) the adjoint of g factors in $\text{Sei}_\infty(A)$ as

$$\begin{array}{ccc} (Q, f, \{\rho_i\}) & \xrightarrow{\text{adj}(g)} & UB(P, e, \{\pi_i\}) \\ & \searrow g\langle T \rangle & \nearrow \text{adj}(\phi\langle T \rangle) \\ & (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\}) & \end{array}$$

for some $T \in I$. The morphisms in $Sei(A)$ defined by

$$s = s\langle T \rangle: (P, e, \{\pi_i\}) \rightarrow (P', e', \{\pi'_i\}) = (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$$

$$t = g\langle T \rangle: (Q, f, \{\rho_i\}) \rightarrow (P', e', \{\pi'_i\}) = (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$$

are such that s is a B -isomorphism (ie $s \in \Xi$) and $g = B(s)^{-1}B(t)$.

(iii) Let $M = B(P, e, \{\pi_i\})$. We have a commutative diagram in $Sei_\infty(A)$

$$\begin{array}{ccc}
 (Q, f, \{\rho_i\}) & \xrightarrow{\text{adj}(B(u)) = 0} & U(M) \\
 & \searrow u & \nearrow \theta_M \\
 & & (P, e, \{\pi_i\})
 \end{array}$$

Since Q is f.g. projective there exists $T \in I$ such that

$$v = s\langle T \rangle: (P, e, \{\pi_i\}) \rightarrow (P\langle T \rangle, e\langle T \rangle, \{\pi_i\langle T \rangle\})$$

is a B -isomorphism in $Sei(A)$ (ie $v \in \Xi$) with $vu = 0$.

(iv) Immediate from (i)–(iii). □

This completes the proof of [Theorem B](#) of the Introduction.

4 Primitive Seifert modules

This section is devoted to the kernel of the covering functor $B: Sei(A) \rightarrow Flk(A)$. Following the terminology of Sheiham [41]:

Definition 4.1 (i) A Seifert A -module $(P, e, \{\pi_i\})$ is *primitive* if

$$B(P, e, \{\pi_i\}) = 0$$

or equivalently $1 - e + ez: P[F_\mu] \rightarrow P[F_\mu]$ is an $A[F_\mu]$ -module isomorphism.

(ii) Let $Prim(A) \subset Sei(A)$ be the full subcategory with objects the primitive f.g. projective Seifert A -modules.

We shall now obtain an intrinsic characterization of the objects in $Prim(A)$, generalizing the results for $\mu = 1$ recalled below.

Definition 4.2 (Lück and Ranicki [28, Section 5]) A *near-projection* (P, e) is an A -module P together with an endomorphism $e \in \text{End}_A(P)$ such that $e(1 - e) \in \text{End}_A(P)$ is nilpotent.

Proposition 4.3 (Bass, Heller and Swan [5], Lück and Ranicki [28])

(i) A linear morphism of induced f.g. projective $A[z]$ -modules

$$f_0 + f_1 z: P[z] \rightarrow Q[z]$$

is an isomorphism if and only if $f_0 + f_1: P \rightarrow Q$ is an isomorphism and

$$e = (f_0 + f_1)^{-1} f_1: P \rightarrow P$$

is nilpotent.

(ii) A linear morphism of induced f.g. projective $A[z, z^{-1}]$ -modules

$$f_0 + f_1 z: P[z, z^{-1}] \rightarrow Q[z, z^{-1}]$$

is an isomorphism if and only if $f_0 + f_1: P \rightarrow Q$ is an isomorphism and

$$e = (f_0 + f_1)^{-1} f_1: P \rightarrow P$$

is a near-projection.

(iii) Suppose that (P, e) is a near-projection, or equivalently that

$$1 - e + ze: P[z, z^{-1}] \rightarrow P[z, z^{-1}]$$

is an $A[z, z^{-1}]$ -module automorphism. If $N \geq 0$ is so large that $(e(1-e))^N = 0$ then

$$e^N + (1-e)^N: P \rightarrow P$$

is an A -module automorphism, and the endomorphism

$$e_\omega = (e^N + (1-e)^N)^{-1} e^N: P \rightarrow P$$

is a projection, with $e_\omega(1-e_\omega) = 0$. The submodules of P

$$P^+ = (1-e_\omega)(P) = (1-e)^N(P) = \{x \in P \mid (1-e+ez)^{-1}e(x) \in P[z]\},$$

$$P^- = e_\omega(P) = e^N(P) = \{x \in P \mid (1-e+ez)^{-1}(1-e)(x) \in z^{-1}P[z^{-1}]\}$$

are such that

$$(P, e) = (P^+, e^+) \oplus (P^-, e^-)$$

with $e^+: P^+ \rightarrow P^+$ and $1-e^-: P^- \rightarrow P^-$ nilpotent.

Definition 4.4 A f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ is *strongly nilpotent* if the $A[F_\mu^+]$ -module endomorphism

$$ez = \sum_{i=1}^{\mu} e\pi_i z_i: P[F_\mu^+] \rightarrow P[F_\mu^+]$$

is nilpotent, ie $(ez)^N = 0$ for some $N \geq 1$.

The condition for strong nilpotence is equivalent to the $A[F_\mu]$ -module endomorphism

$$ez = \sum_{i=1}^{\mu} e\pi_i z_i: P[F_\mu] \rightarrow P[F_\mu]$$

being nilpotent.

Expressed as a representation of the complete quiver Q_μ , a Seifert module $(P, \rho: Q_\mu \rightarrow \text{End}_A P)$ is strongly nilpotent if and only if there exists $N \geq 1$ such that $\rho(p) = 0$ for every path $p \in Q_\mu$ of length $\geq N$.

Proposition 4.5 *The following conditions on a f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ are equivalent:*

- (i) $(P, e, \{\pi_i\})$ is strongly nilpotent,
- (ii) the $A[F_\mu^+]$ -module endomorphism

$$1 - ez: P[F_\mu^+] \rightarrow P[F_\mu^+]$$

is an automorphism,

- (iii) the $A[F_\mu^+]$ -module endomorphism

$$1 - e + ez: P[F_\mu^+] \rightarrow P[F_\mu^+]$$

is an automorphism.

Proof (i) \implies (ii) If $(ez)^N = 0$ then $1 - ez$ has inverse

$$(1 - ez)^{-1} = 1 + ez + (ez)^2 + \dots + (ez)^{N-1} \in \text{Hom}_{A[F_\mu^+]}(P[F_\mu^+], P[F_\mu^+]) = \text{Hom}_A(P, P)[F_\mu^+].$$

(ii) \implies (i) The inverse of $1 - ez$ is of the form

$$(1 - ez)^{-1} = \sum_{\substack{1 \leq i_1, i_2, \dots, i_k \leq \mu \\ n_1, n_2, \dots, n_k \geq 0 \\ n_1 + n_2 + \dots + n_k < N}} f_{i_1 i_2 \dots i_k} z_{i_1}^{n_1} z_{i_2}^{n_2} \dots z_{i_k}^{n_k}: P[F_\mu^+] \rightarrow P[F_\mu^+]$$

for some $N \geq 1$. We have the identity

$$(1 - ez)^{-1} - (1 + ez + (ez)^2 + \dots + (ez)^{N-1}) = (1 - ez)^{-1}(ez)^N \in \text{Hom}_{A[F_\mu^+]}(P[F_\mu^+], P[F_\mu^+]) = \text{Hom}_A(P, P)[F_\mu^+]$$

in which the left hand side is a sum of monomials in $z_{i_1} z_{i_2}^{n_2} \dots z_{i_k}^{n_k}$ of degree $n_1 + n_2 + \dots + n_k < N$ and the right hand side is a sum of monomials of degree $\geq N$. Both sides of the identity are thus 0,

$$(ez)^N = 0: P[F_\mu^+] \rightarrow P[F_\mu^+]$$

and $(P, e, \{\pi_i\})$ is strongly nilpotent.

(ii) \iff (iii) Immediate from the identity

$$1 - e + ez = 1 - e(1 - z): P[F_\mu^+] \rightarrow P[F_\mu^+]$$

and the change of variables $z_i \mapsto 1 - z_i$. □

Definition 4.6 A μ -component Seifert A -module $(P, e, \{\pi_i\})$ is a *near-projection* if it can be expressed as

$$(P, e, \{\pi_i\}) = \left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & e^{+-} \\ e^{-+} & e^{--} \end{pmatrix}, \{\pi_i^+\} \oplus \{\pi_i^-\} \right)$$

and the 2μ -component Seifert A -module

$$(P', e', \pi') = \left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix}, \{\pi_i^+\} \oplus \{\pi_i^-\} \right)$$

is strongly nilpotent.

Lemma 4.7 For a near-projection $(P, e, \{\pi_i\})$ the pairs (P, e) , (P, e') are near-projections.

Proof We have a decomposition $P = P^+ \oplus P^-$ with respect to which e' is strongly nilpotent. Now

$$\begin{aligned} e(1 - e) &= \begin{pmatrix} e^{++} & e^{+-} \\ e^{-+} & e^{--} \end{pmatrix} \begin{pmatrix} 1 - e^{++} & -e^{+-} \\ -e^{-+} & 1 - e^{--} \end{pmatrix} \\ &= \begin{pmatrix} e^{++} - (e^{++})^2 - e^{+-}e^{-+} & -e^{++}e^{+-} + e^{+-}(1 - e^{--}) \\ e^{-+} - e^{-+}e^{++} - e^{--}e^{-+} & -e^{-+}e^{+-} + e^{--}(1 - e^{--}) \end{pmatrix} \\ &= \begin{pmatrix} e^{++} - (e^{++})^2 - e^{+-}e^{-+} & -e^{++}e^{+-} + e^{+-}(1 - e^{--}) \\ e^{-+} - e^{-+}e^{++} - e^{--}e^{-+} & -e^{-+}e^{+-} + (1 - e^{--}) - (1 - e^{--})^2 \end{pmatrix}. \end{aligned}$$

The matrix

$$e' = \left(\begin{array}{c|c} e^{++} & -e^{+-} \\ \hline e^{-+} & 1 - e^{--} \end{array} \right)$$

denotes a strongly nilpotent representation of the complete quiver $Q_{2\mu}$ on 2μ vertices. In the following illustration $\mu = 1$:



Now each entry in the $2\mu \times 2\mu$ matrix $e(1 - e)$ above is (the image of) a linear combination of paths of length at least one in the quiver. Hence each entry of $(e(1 - e))^N$ is the image of a sum of paths of length at least N . It follows that $(e(1 - e))^N = 0$ for some $N \geq 1$.

The pair (P, e') is a near-projection since $e': P \rightarrow P$ is nilpotent. □

For $\mu = 1$ there is no difference between a near-projection $(P, e, \{\pi_i\})$ and a near-projection (P, e) . For $\mu \geq 2$ a near-projection $(P, e, \{\pi_i\})$ has (P, e) a near-projection (Lemma 4.7) but the splitting $(P, e) = (P^+, e^+) \oplus (P^-, e^-)$ given by Proposition 4.3 does not in general extend to a direct sum decomposition of Seifert A -modules

$$(P, e, \{\pi_i\}) = (P^+, e^+, \{\pi_i^+\}) \oplus (P^-, e^-, \{\pi_i^-\}).$$

This is illustrated by the following example.

Example 4.8 Let A be a field, and consider the 2-component Seifert A -module $(P, e, \{\pi_1, \pi_2\})$ given by

$$P = A^4, \quad e = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \quad \pi_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \pi_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

In this case $e: P \rightarrow P$ is a projection, with $e(1 - e) = 0$. This f.g. projective Seifert A -module has just one submodule

$$(\bar{P}, \bar{e}, \{\bar{\pi}_1, \bar{\pi}_2\}) \subseteq (P, e, \{\pi_1, \pi_2\})$$

namely

$$\bar{P} = e(P) = \{(0, x, 0, y) \in P \mid (x, y) \in A^2\}.$$

It is not possible to decompose $(P, e, \{\pi_1, \pi_2\})$ as a direct sum, since $(\bar{P}, \bar{e}, \{\bar{\pi}_1, \bar{\pi}_2\})$ is not a summand. Neither e nor $1 - e$ is nilpotent but

$$1 - e + ez = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & z_1 & z_2 - 1 & 0 \\ 0 & 0 & 1 & 0 \\ z_1 - 1 & 0 & 0 & z_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & z_2 - 1 & 0 \\ 0 & 0 & 1 & 0 \\ z_1 - 1 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & z_1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & z_2 \end{pmatrix}$$

and

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & z_2 - 1 & 0 \\ 0 & 0 & 0 & 0 \\ z_1 - 1 & 0 & 0 & 0 \end{pmatrix}^2 = 0$$

so $1 - e + ez$ is invertible. Moreover, $(P, e, \{\pi_1, \pi_2\})$ is a near-projection, with

$$P_1^+ = A \oplus 0 \oplus 0 \oplus 0, P_1^- = 0 \oplus A \oplus 0 \oplus 0, P_2^+ = 0 \oplus 0 \oplus A \oplus 0, P_2^- = 0 \oplus 0 \oplus 0 \oplus A$$

such that

$$e' = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} : P = P_1^+ \oplus P_1^- \oplus P_2^+ \oplus P_2^- \rightarrow P = P_1^+ \oplus P_1^- \oplus P_2^+ \oplus P_2^-$$

is strongly nilpotent.

The main result of this section is:

Theorem 4.9 *A f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ is primitive if and only if it is a near-projection.*

Proof Suppose that $(P, e, \{\pi_i\})$ is a near-projection, with $e' = \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix}$ strongly nilpotent. We have

$$\begin{aligned} 1 - e + ez &= 1 - e(1 - z) \\ &= \begin{pmatrix} 1 - e^{++}(1 - z) & -e^{+-}(1 - z) \\ -e^{-+}(1 - z) & 1 - e^{--}(1 - z) \end{pmatrix} \\ &= \begin{pmatrix} 1 - e^{++}(1 - z) & e^{+-}(1 - z^{-1}) \\ -e^{-+}(1 - z) & 1 - (1 - e^{--})(1 - z^{-1}) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix} \\ &= \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix} \begin{pmatrix} 1 - z & 0 \\ 0 & 1 - z^{-1} \end{pmatrix} \right) \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix} : (P^+ \oplus P^-)[F_\mu] \\ &\qquad \qquad \qquad \longrightarrow (P^+ \oplus P^-)[F_\mu]. \end{aligned}$$

It follows from the strong nilpotence of e' that $e'((1 - z) \oplus (1 - z^{-1}))$ is nilpotent, and hence that

$$1 - e(1 - z) = (1 - e'((1 - z) \oplus (1 - z^{-1}))) (1 \oplus z) : (P^+ \oplus P^-)[F_\mu] \rightarrow (P^+ \oplus P^-)[F_\mu]$$

is an isomorphism, so that $B(P, e, \{\pi_i\}) = 0$ and $(P, e, \{\pi_i\})$ is primitive.

Conversely, suppose that $(P, e, \{\pi_i\})$ is a primitive f.g. projective Seifert A -module, ie such that the $A[F_\mu]$ -module morphism

$$1 - e + ez: P[F_\mu] \rightarrow P[F_\mu]$$

is an isomorphism. We shall use a variant \bar{G}_μ of the Cayley tree G_μ (Definition 1.3) to prove that $1 - e + ez: P[F_\mu] \rightarrow P[F_\mu]$ is a near-projection. Define

$$\bar{G}_\mu^{(0)} = F_\mu, \quad \bar{G}_\mu^{(1)} = \{(w, z_i w) \mid w \in F_\mu, i \in \{1, 2, \dots, \mu\}\}$$

so that there is defined a right F_μ -action

$$\bar{G}_\mu \times F_\mu \rightarrow \bar{G}_\mu; (w, g) \mapsto wg.$$

For each $i = 1, 2, \dots, \mu$ partition F_μ as

$$F_\mu = F_\mu^{+,i} \sqcup F_\mu^{-,i} \sqcup \{1\}$$


with $F_\mu^{+,i}$ (resp. $F_\mu^{-,i}$) consisting of the reduced words in z_1, z_2, \dots, z_μ which start (resp. do not start) with z_i . Removing the edge $(w, z_i w)$ disconnects \bar{G}_μ , and the complement is a disjoint union of trees

$$\bar{G}_\mu - \{(w, z_i w)\} = \bar{G}_\mu^+(w, z_i w) \sqcup \bar{G}_\mu^-(w, z_i w)$$

with

$$\bar{G}_\mu^+(w, z_i w)^{(0)} = F_\mu^{+,i} w, \quad \bar{G}_\mu^-(w, z_i w)^{(0)} = (F_\mu^{-,i} \cup \{1\})w.$$

In the diagram



we are placing the components of the range (resp. domain) $P[F_\mu]$ at the vertices (resp. edges) of \bar{G}_μ , with the A -module wP at $w \in \bar{G}_\mu^{(0)}$, and the A -module wP_i at

$(w, z_i w) \in \overline{G}_\mu^{(1)}$. An element

$$x \in P[F_\mu] = \sum_{(w, z_i w) \in \overline{G}_\mu^{(1)}} w P_i$$

is sent to

$$(1 - e)(x) + ez(x) \in P[F_\mu] = \sum_{w \in \overline{G}_\mu^{(0)}} w P,$$

as indicated by the arrows in the diagram. For $i = 1, 2, \dots, \mu$ define the A -modules

$$P_i^+ = \left\{ x \in P_i \mid (1 - e + ez)^{-1} ez(x) \in \sum_{w \in F_\mu^{+,i}} w P \right\},$$

$$P_i^- = \left\{ x \in P_i \mid (1 - e + ez)^{-1} (1 - e)(x) \in \sum_{j \neq i} P_j \oplus \sum_{w \in F_\mu^{-,i}} w P \right\}.$$

An element $x^+ \in P_i$ belongs to P_i^+ if and only if there exist elements $y^+(w) \in P$ ($w \in F_\mu^{+,i}$) such that

$$ez(x^+) = (1 - e + ez) \left(\sum_{w \in F_\mu^{+,i}} w y^+(w) \right) \in \sum_{w \in F_\mu^{+,i}} w P. \quad (*)$$

There is one component $y^+(w)$ for each edge in $\overline{G}_\mu^+(1, z_i)^{(1)}$, and one equation for each vertex in $\overline{G}_\mu^+(1, z_i)^{(0)}$. Similarly, an element $x^- \in P_i$ belongs to P_i^- if and only if there exist elements $y_j \in P_j$ ($j \neq i$) and $y^-(w) \in P$ ($w \in F_\mu^{-,i}$) such that

$$((1 - e)(x^-), 0) = (1 - e + ez) \left(\sum_{j \neq i} y_j + \sum_{w \in F_\mu^{-,i}} w y^-(w) \right) \in P \oplus \sum_{w \in F_\mu^{-,i}} w P. \quad (**)$$

There is one component y_j ($j \neq i$) or $y^-(w)$ for each edge $\overline{G}_\mu^-(1, z_i)^{(1)}$, and one equation for each vertex in $\overline{G}_\mu^-(1, z_i)^{(0)}$. For $i = 1, 2, \dots, \mu$ partition

$$F_\mu^{+,i} = F_\mu^{++ ,i} \sqcup F_\mu^{-+,i}, \quad F_\mu^{-,i} = F_\mu^{+-,i} \sqcup F_\mu^{-- ,i}$$

with $F_\mu^{\alpha+,i}$ consisting of the words $w = z_{i_0}^{\epsilon_0} \dots z_{i_k}^{\epsilon_k} \in F_\mu$ with $(i_0, \epsilon_0) = (i, +)$, $\epsilon_k = \alpha$, and $F_\mu^{\alpha-,i}$ consisting of the words $w = z_{i_0}^{\epsilon_0} \dots z_{i_k}^{\epsilon_k} \in F_\mu$ with $(i_0, \epsilon_0) \neq (i, +)$, $\epsilon_k = \alpha$. For any $x^+ \in P_i^+$ and $w \in F_\mu^{\alpha+,i}$ we have that $y^+(w) \in P_j^\alpha$, as given by all the terms in (*) involving $\overline{G}^+(w, z_i w)$. Similarly, for any $x^- \in P_i^-$ and $w \in F_\mu^{\alpha-,i}$ we have that $y^-(w) \in P_j^\alpha$, as given by all the terms in (**) involving $\overline{G}^-(w, z_i w)$.

Regarded as an A -module isomorphism $1-e+ez: P[F_\mu] \rightarrow P[F_\mu]$ can be expressed as

$$1-e+ez = \begin{pmatrix} ez| & (1-e+ez)| & 0 \\ (1-e)| & 0 & (1-e+ez)| \end{pmatrix}: P_i \oplus \left(\sum_{w \in F_\mu^{+,i}} wP \right) \oplus \left(\sum_{j \neq i} P_j \oplus \sum_{w \in F_\mu^{-,i}} wP \right) \rightarrow \left(\sum_{w \in F_\mu^{+,i}} wP \right) \oplus \left(P \oplus \sum_{w \in F_\mu^{-,i}} wP \right)$$

so that there is induced an A -module isomorphism

$$\begin{aligned} \left[\begin{array}{c} ez| \\ (1-e)| \end{array} \right]: P_i &\rightarrow \left(\text{coker} \left((1-e+ez): \sum_{w \in F_\mu^{+,i}} wP \rightarrow \sum_{w \in F_\mu^{+,i}} wP \right) \right) \\ &\oplus \left(\text{coker} \left((1-e+ez): \sum_{j \neq i} P_j \oplus \sum_{w \in F_\mu^{-,i}} wP \rightarrow P \oplus \sum_{w \in F_\mu^{-,i}} wP \right) \right) \end{aligned}$$

and

$$P_i = P_i^+ \oplus P_i^-,$$

with

$$\begin{aligned} (1-e+ez)^{-1}ez(P_i^+) &\subseteq \sum_{j=1}^\mu \sum_{w \in F_\mu^{+,i}} wP_j^+ \oplus \sum_{j=1}^\mu \sum_{w \in F_\mu^{-,i}} wP_j^-, \\ (1-e+ez)^{-1}(1-e)(P_i^-) &\subseteq \sum_{j=1}^\mu \sum_{w \in F_\mu^{+,i}} wP_j^+ \oplus \sum_{j=1}^\mu \sum_{w \in F_\mu^{-,i}} wP_j^-. \end{aligned}$$

For $\alpha, \beta \in \{\pm\}$ let

$$e_{ji}^{\beta\alpha}: P_i^\alpha \rightarrow P_j^\beta$$

be the A -module morphisms such that

$$e = \begin{pmatrix} e_{ji}^{++} & e_{ji}^{+-} \\ e_{ji}^{-+} & e_{ji}^{--} \end{pmatrix}: P = \sum_{i=1}^\mu (P_i^+ \oplus P_i^-) \rightarrow P = \sum_{j=1}^\mu (P_j^+ \oplus P_j^-).$$

Let

$$v_{ji}^{\beta\alpha}(w): P_i^\alpha \rightarrow P_j^\beta$$

be the A -module morphisms such that

$$\begin{aligned}
 -(1-e+ez)^{-1}ez| &= \left(\begin{array}{c} \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{++},i} wv_{ji}^{++}(w) \\ \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{-+},i} wv_{ji}^{-+}(w) \end{array} \right) : P_i^+ \\
 &\rightarrow \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{++},i} wP_j^+ \oplus \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{-+},i} wP_j^-, \\
 -(1-e+ez)^{-1}(1-e)| &= \left(\begin{array}{c} \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{+-},i} wv_{ji}^{+-}(w) \\ \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{--},i} wv_{ji}^{--}(w) \end{array} \right) : P_i^- \\
 &\rightarrow \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{+-},i} wP_j^+ \oplus \sum_{j=1}^{\mu} \sum_{w \in F_{\mu}^{--},i} wP_j^-.
 \end{aligned}$$

Composing with $1 - e + ez$ gives

(*)

$$\begin{aligned}
 -e_{ji}^{++}z_i &= \sum_{k=1}^{\mu} \left(\sum_{w \in F_{\mu}^{++},i} w(\delta_{jk} - e_{jk}^{++})v_{ki}^{++}(w) + \sum_{w \in F_{\mu}^{-+},i} wz_k(e_{jk}^{+-})v_{ki}^{-+}(w) \right) : \\
 &P_i^+ \rightarrow \sum_{w \in F_{\mu}^{++},i} wP_j^+, \\
 -e_{ji}^{-+}z_i &= \sum_{k=1}^{\mu} \left(\sum_{w \in F_{\mu}^{++},i} we_{jk}^{-+}v_{ki}^{++}(w) + \sum_{w \in F_{\mu}^{-+},i} wz_k e_{jk}^{--}v_{ki}^{-+}(w) \right) : \\
 &P_i^+ \rightarrow \sum_{w \in F_{\mu}^{-+},i} wP_j^-, \\
 -(-e_{ji}^{+-}) &= \sum_{k=1}^{\mu} \left(\sum_{w \in F_{\mu}^{+-},i} w(\delta_{jk} - e_{jk}^{+-})v_{ki}^{+-}(w) + \sum_{w \in F_{\mu}^{--},i} wz_k e_{jk}^{+-}v_{ki}^{--}(w) \right) : \\
 &P_i^- \rightarrow \sum_{w \in F_{\mu}^{+-},i} wP_j^+, \\
 -(\delta_{ji} - e_{ji}^{--}) &= \sum_{k=1}^{\mu} \left(\sum_{w \in F_{\mu}^{+-},i} we_{jk}^{-+}v_{ki}^{+-}(w) + \sum_{w \in F_{\mu}^{--},i} wz_k e_{jk}^{--}v_{ki}^{--}(w) \right) : \\
 &P_i^- \rightarrow \sum_{w \in F_{\mu}^{--},i} wP_j^-.
 \end{aligned}$$

Comparing the coefficients of z_i and 1 gives

$$\begin{aligned}
 - \begin{pmatrix} e_{ji}^{++} \\ e_{ji}^{-+} \end{pmatrix} &= \sum_{k=1}^{\mu} \left(\begin{pmatrix} \delta_{jk} - e_{jk}^{++} \\ -e_{jk}^{-+} \end{pmatrix} v_{ki}^{++}(z_i) + \begin{pmatrix} e_{jk}^{+-} \\ e_{jk}^{--} \end{pmatrix} v_{ki}^{-+}(z_i z_k^{-1}) \right): \\
 & P_i^+ \rightarrow P_j^+ \oplus P_j^-, \\
 - \begin{pmatrix} -e_{ji}^{+-} \\ \delta_{ji} - e_{ji}^{--} \end{pmatrix} &= \sum_{k=1}^{\mu} \left(\begin{pmatrix} \delta_{jk} - e_{jk}^{++} \\ -e_{jk}^{-+} \end{pmatrix} v_{ki}^{+-}(1) + \begin{pmatrix} e_{jk}^{+-} \\ e_{jk}^{--} \end{pmatrix} v_{ki}^{--}(z_k^{-1}) \right): \\
 & P_i^- \rightarrow P_j^+ \oplus P_j^-.
 \end{aligned}$$

Writing

$$\begin{pmatrix} v^{++} & v^{+-} \\ v^{-+} & v^{--} \end{pmatrix} = \begin{pmatrix} v_{ki}^{++}(z_i) & v_{ki}^{+-}(1) \\ v_{ki}^{-+}(z_i z_k^{-1}) & v_{ki}^{--}(z_k^{-1}) \end{pmatrix}: P^+ \oplus P^- \rightarrow P^+ \oplus P^-,$$


we thus have

$$- \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix} = \begin{pmatrix} 1 - e^{++} & e^{+-} \\ -e^{-+} & e^{--} \end{pmatrix} \begin{pmatrix} v^{++} & v^{+-} \\ v^{-+} & v^{--} \end{pmatrix}.$$

Let $Q_{\mu}^{+,-}$ be the quiver with 2μ vertices $(i, \pm)_{1 \leq i \leq \mu}$ and one edge $(i_0, \epsilon_0) \rightarrow (i_1, \epsilon_1)$ for each pair of vertices with $(i_0, \epsilon_0) \neq (i_1, -\epsilon_1)$. (The path ring is given by

$$\begin{aligned}
 Q_{\mu}^{+,-} = \\
 \mathbb{Z}[s] * \mathbb{Z} \left[\pi_1^+, \dots, \pi_{\mu}^+, \pi_1^-, \dots, \pi_{\mu}^- \mid \pi_i^{\alpha} \pi_j^{\beta} = \delta_{\alpha\beta} \delta_{ij} \pi_i^{\alpha}, \sum_{i=1}^{\mu} (\pi_i^+ + \pi_i^-) = 1 \right] \\
 / \{ \pi_i^{\alpha} s \pi_i^{-\alpha} \}
 \end{aligned}$$

where $\pi_i^{\alpha} s \pi_j^{\beta}$ ($(i, \alpha) \neq (j, -\beta)$) corresponds to the unique path of length 1 from (i, α) to (j, β) .) In the illustration $\mu = 2$:



Regard a word $w = z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1} \dots z_{i_k}^{\epsilon_k} \in F_{\mu}$ as a path of length $|w| = k$ in $Q_{\mu}^{+,-}$

$$(i_0, \epsilon_0) \rightarrow (i_1, \epsilon_1) \rightarrow \dots \rightarrow (i_k, \epsilon_k)$$

and for $k \geq 1$ define an A -module morphism $v(w): P_{i_0}^{\epsilon_0} \rightarrow P_{i_1}^{\epsilon_1}$ as follows. Define

$$[w] = [z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1}] [z_{i_1}^{\epsilon_1} z_{i_2}^{\epsilon_2}] \dots [z_{i_{k-1}}^{\epsilon_{k-1}} z_{i_k}^{\epsilon_k}] \in F_{\mu}$$

with

$$[z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1}] = \begin{cases} z_{i_0}^{\epsilon_0} & \text{if } (\epsilon_0, \epsilon_1) = (+, +) \\ z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1} & \text{if } (\epsilon_0, \epsilon_1) = (+, -) \\ z_{i_1}^{\epsilon_1} & \text{if } (\epsilon_0, \epsilon_1) = (-, -) \\ 1 & \text{if } (\epsilon_0, \epsilon_1) = (-, +). \end{cases}$$

For $k = 1$ set

$$v(z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1}) = v_{i_1 i_0}^{\epsilon_1 \epsilon_0}([z_{i_1}^{\epsilon_1} z_{i_0}^{\epsilon_0}])$$

and for $k \geq 2$ set

$$v(z_{i_0}^{\epsilon_0} z_{i_1}^{\epsilon_1} \dots z_{i_k}^{\epsilon_k}) = v(z_{i_{k-1} i_k}^{\epsilon_{k-1} \epsilon_k}) \dots v(z_{i_1 i_2}^{\epsilon_1 \epsilon_2}) v(z_{i_0 i_1}^{\epsilon_0 \epsilon_1}).$$

The identities

$$v(w) = v_{i_k i_0}^{\epsilon_k \epsilon_0}([w]): P_{i_0}^{\epsilon_0} \rightarrow P_{i_k}^{\epsilon_k}$$

may be verified by induction on k , since both sides satisfy the equations (*) and so

$$\begin{aligned} -(1 - e + ez)^{-1} ez| &= \sum_{w \in F_\mu^{+,i}} w v(w): P_i^+ \rightarrow \sum_{w \in F_\mu^{+,i}} w P, \\ -(1 - e + ez)^{-1} (1 - e)| &= \sum_{j \neq i} v_{ji}^{+-} + \sum_{w \in F_\mu^{-,i}} w v(w): P_i^- \rightarrow \sum_{j \neq i} P_j \oplus \sum_{w \in F_\mu^{-,i}} w P. \end{aligned}$$

For $\alpha, \beta \in \{\pm\}$ let $F_\mu^{\beta\alpha}$ be the set of paths

$$(i_0, \epsilon_0) \rightarrow (i_1, \epsilon_1) \rightarrow \dots \rightarrow (i_k, \epsilon_k)$$

in $Q_\mu^{+,-}$ with $\epsilon_0 = \alpha, \epsilon_k = \beta$. The $A[F_\mu]$ -module endomorphism

$$v' = \begin{pmatrix} z & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v^{++} & v^{+-} \\ v^{-+} & v^{--} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & z^{-1} \end{pmatrix}: (P^+ \oplus P^-)[F_\mu] \rightarrow (P^+ \oplus P^-)[F_\mu]$$

is such that for any $N \geq 1$

$$(v')^N = \begin{pmatrix} \sum_{w \in F_\mu^{++}, |w|=N} w v^{++}(w) & \sum_{w \in F_\mu^{+-}, |w|=N} w v^{+-}(w) \\ \sum_{w \in F_\mu^{-+}, |w|=N} w v^{-+}(w) & \sum_{w \in F_\mu^{--}, |w|=N} w v^{--}(w) \end{pmatrix}: (P^+ \oplus P^-)[F_\mu] \rightarrow (P^+ \oplus P^-)[F_\mu].$$

If $N \geq 1$ is so large that

$$(1 - e + ez)^{-1} = \sum_{w \in F_\mu, |w| < N} a_w w: P[F_\mu] \rightarrow P[F_\mu] \quad (a_w \in \text{Hom}_A(P_{i_0}, P_{i_k})),$$

then for any word $w \in F_\mu$ of length $|w| = k > N$

$$v(w) = 0: P_{i_0}^{\epsilon_0} \rightarrow P_{i_k}^{\epsilon_k}.$$

The 2μ -component Seifert module

$$(P', v', \pi') = \left(P^+ \oplus P^-, \begin{pmatrix} v^{++} & v^{+-} \\ v^{-+} & v^{--} \end{pmatrix}, \{\pi_i^+ \oplus \pi_i^-\} \right)$$

is strongly nilpotent, with $(v'z')^N = 0$, regarding $F_{2\mu}$ as free group on 2μ generators $z'_1, z'_2, \dots, z'_{2\mu}$ and letting

$$z' = \begin{pmatrix} z'_1 & 0 & \dots & 0 \\ 0 & z'_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & z'_{2\mu} \end{pmatrix}: P'[F_{2\mu}] \rightarrow P'[F_{2\mu}].$$

Define the 2μ -component Seifert module

$$(P', e', \pi') = \left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & -e^{+-} \\ e^{-+} & 1 - e^{--} \end{pmatrix}, \{\pi_i^+ \oplus \pi_i^-\} \right).$$

Applying the augmentation $\epsilon: z_i \mapsto 1$ to the $A[F_\mu]$ -module morphisms

$$\begin{aligned} & - \sum_{w \in F_\mu^{+,i}} wv(w): P_i^+[F_\mu] \xrightarrow{ez} P[F_\mu] \xrightarrow{(1-e+ez)^{-1}} P[F_\mu], \\ & - \left(\sum_{j \neq i} v_{ji}^{+-} + \sum_{w \in F_\mu^{-,i}} wv(w) \right): P_i^-[F_\mu] \xrightarrow{1-e} P[F_\mu] \xrightarrow{(1-e+ez)^{-1}} P[F_\mu] \end{aligned}$$

shows that the components of e' are given by linear combinations of paths of length ≥ 1

$$\begin{aligned} e^{++} &= - \sum_{w \in F_\mu^{++}} v(w): P^+ \rightarrow P^+, \\ e^{-+} &= - \sum_{w \in F_\mu^{+-}} v(w): P^+ \rightarrow P^-, \\ -e^{+-} &= - \sum_{w \in F_\mu^{-+}} v(w): P^- \rightarrow P^+, \\ 1 - e^{--} &= - \sum_{w \in F_\mu^{--}} v(w): P^- \rightarrow P^-. \end{aligned}$$

The $A[F_{2\mu}]$ -module endomorphism $e'z': P'[F_{2\mu}] \rightarrow P'[F_{2\mu}]$ is nilpotent, with

$$(e'z')^N = 0,$$

so that (P, e, π) is strongly nilpotent. \square

This completes the proof of [Theorem C](#) of the Introduction.

5 Algebraic K -theory

We shall obtain our results on the algebraic K -theory of $A[F_\mu]$ and Blanchfield and Seifert modules using the Waldhausen [50] algebraic K -theory of categories with cofibrations and weak equivalences, and the noncommutative localization algebraic K -theory exact sequence of Neeman and Ranicki [30; 31].

5.1 The algebraic K -theory of exact categories

The higher algebraic K -groups $K_n(\mathcal{E})$ of an exact category \mathcal{E} are defined by Quillen [32] to be the homotopy groups of a connective spectrum $K(\mathcal{E})$

$$\pi_n(K(\mathcal{E})) = K_n(\mathcal{E}) \quad (n \geq 0)$$

with $K_0(\mathcal{E})$ the Grothendieck class group. The idempotent completion $\mathcal{E} \rightarrow \mathcal{P}(\mathcal{E})$ induces an injection $K_0(\mathcal{E}) \rightarrow K_0(\mathcal{P}(\mathcal{E}))$ and isomorphisms $K_n(\mathcal{E}) \rightarrow K_n(\mathcal{P}(\mathcal{E}))$ for $n \geq 1$, by the cofinality theorem of Grayson [21]. The lower K -groups $K_n(\mathcal{E})$ ($n \leq -1$) are defined by Schlichting [38] (following on from the definitions of Karoubi

and Pedersen–Weibel for the lower K –groups of filtered additive categories) to be the lower homotopy groups of a nonconnective spectrum $K\mathcal{P}(\mathcal{E})$ such that

$$\pi_n(K\mathcal{P}(\mathcal{E})) = K_n(\mathcal{P}(\mathcal{E})) \quad (n \in \mathbb{Z}),$$

with $K_n(\mathcal{E}) = K_n(\mathcal{P}(\mathcal{E}))$ for $n \neq 0$.

The algebraic K –groups of a ring R are the algebraic K –groups of the idempotent complete exact category $\mathcal{E} = \mathcal{P}roj(R)$ of f.g. projective R –modules

$$K_n(R) = K_n(\mathcal{P}roj(R)) \quad (n \in \mathbb{Z}),$$

as defined for $-\infty < n \leq 1$ in Bass [4], and for $2 \leq n < \infty$ in Quillen [32]. The nonconnective spectrum defined by $K(R) = K\mathcal{P}(\mathcal{P}roj(R))$ has homotopy groups $\pi_*(K(R)) = K_*(R)$.

A *Waldhausen category* (\mathcal{C}, w) is a small category \mathcal{C} with cofibrations together with a subcategory $w \subset \mathcal{C}$ of weak equivalences satisfying the axioms of [50]. As usual, there is defined a connective algebraic K –theory spectrum

$$K(\mathcal{C}, w) = \Omega|wS_\bullet\mathcal{C}|$$

with homotopy groups the algebraic K –theory groups

$$K_n(\mathcal{C}, w) = \pi_n(K(\mathcal{C}, w)) \quad (n \geq 0).$$

A functor $F: (\mathcal{C}, w) \rightarrow (\mathcal{C}', w')$ of Waldhausen categories induces a long exact sequence of algebraic K –groups

$$\cdots \rightarrow K_{n+1}(F) \rightarrow K_n(\mathcal{C}, w) \xrightarrow{F} K_n(\mathcal{C}', w') \rightarrow \cdots \rightarrow K_0(F) \rightarrow 0$$

with $K_n(F) = \pi_n(F: K(\mathcal{C}, w) \rightarrow K(\mathcal{C}', w'))$ ($n \geq 0$).

As in Thomason and Trobaugh [45, 1.9] we shall only be considering Waldhausen categories (\mathcal{C}, w) which are ‘complicial biWaldhausen’, so that in particular \mathcal{C} is a full subcategory of the category of chain complexes in an abelian category \mathcal{A} , the cofibrations are chain maps which are split injections in each degree, w contains the quasi-isomorphisms (= the chain maps inducing isomorphisms in homology), and which in addition are closed under the formation of canonical homotopy pushouts and pullbacks.

The *homotopy* (or *derived*) *category* [45, page 269] of a Waldhausen category (\mathcal{C}, w) is the category of fractions

$$D(\mathcal{C}, w) = w^{-1}\mathcal{C},$$

which is a triangulated category under the above hypotheses. The idempotent completion $\mathcal{P}D(\mathcal{C}, w)$ is then also triangulated (Balmer and Schlichting [2]), and the class groups $K_0(D(\mathcal{C}, w))$, $K_0(\mathcal{P}D(\mathcal{C}, w))$ are defined, with $K_0(D(\mathcal{C}, w)) = K_0(\mathcal{C}, w)$. Schlichting [38] defined the lower K -groups $K_n(\mathcal{P}D(\mathcal{C}, w))$ for $n \leq -1$ for Waldhausen categories as above, and constructed a nonconnective spectrum $K\mathcal{P}(\mathcal{C}, w)$ with homotopy groups

$$\pi_n(K\mathcal{P}(\mathcal{C}, w)) = K\mathcal{P}_n(\mathcal{C}, w) = \begin{cases} K_n(\mathcal{C}, w) & \text{for } n \geq 1 \\ K_0(\mathcal{P}D(\mathcal{C}, w)) & \text{for } n = 0 \\ K\mathcal{P}_n(\mathcal{C}, w) & \text{for } n \leq -1. \end{cases}$$

A functor $F: (\mathcal{C}, w) \rightarrow (\mathcal{C}', w')$ of Waldhausen categories induces a long exact sequence of algebraic K -groups

$$\cdots \rightarrow K\mathcal{P}_{n+1}(F) \rightarrow K\mathcal{P}_n(\mathcal{C}, w) \xrightarrow{F} K\mathcal{P}_n(\mathcal{C}', w') \rightarrow K\mathcal{P}_n(F) \rightarrow \cdots,$$

with $K\mathcal{P}_n(F) = \pi_n(F: K\mathcal{P}(\mathcal{C}, w) \rightarrow K\mathcal{P}(\mathcal{C}', w'))$ ($n \in \mathbb{Z}$).

Given an exact category \mathcal{E} let $C^b(\mathcal{E})$ be the category of bounded chain complexes in \mathcal{E} and chain maps. An object C in $C^b(\mathcal{E})$ is *acyclic* (in the sense of Keller [24, Chapter 11]) if each differential $d: C_r \rightarrow C_{r-1}$ factors as $C_r \rightarrow Z_r \rightarrow C_{r-1}$ with

$$0 \rightarrow Z_{r+1} \rightarrow C_r \rightarrow Z_r \rightarrow 0$$

exact. A morphism $f: C \rightarrow D$ in $C^b(\mathcal{E})$ is a *quasi-isomorphism* if the mapping cone $\mathcal{C}(f)$ is chain equivalent to an acyclic complex. If \mathcal{E} is fully embedded in an abelian category \mathcal{A} with the embedding closed under extensions and the idempotent completion $\mathcal{P}(\mathcal{E})$ is closed under taking kernels of surjections then a quasi-isomorphism is the same as a chain map inducing isomorphisms in homology in the ambient abelian category \mathcal{A} [45, Appendix A].

Let $(C^b(\mathcal{E}), w_{\mathcal{E}})$ be the Waldhausen category with cofibrations the chain maps which are degreewise split injections, and $w_{\mathcal{E}} \subset C^b(\mathcal{E})$ the subcategory of quasi-isomorphisms. The derived category

$$D^b(\mathcal{E}) = D(C^b(\mathcal{E}), w_{\mathcal{E}})$$

is the category of bounded chain complexes in \mathcal{E} and fractions of chain homotopy classes of chain maps, with denominators quasi-isomorphisms. As usual, let $K^b(\mathcal{E})$ be the category of bounded chain complexes in \mathcal{E} and chain homotopy classes of chain maps, and let $wK_{\mathcal{E}} \subset K^b(\mathcal{E})$ be the subcategory of quasi-isomorphisms: the localization

$$D^b(\mathcal{E}) = (wK_{\mathcal{E}})^{-1} K^b(\mathcal{E})$$

has both a left and a right calculus of fractions. The derived category $D^b(\mathcal{E})$ is a triangulated category [45, 1.9.6]. Balmer and Schlichting [2, 2.12] prove that the idempotent completion of the derived category is the derived category of the idempotent completion

$$\mathcal{P}D^b(\mathcal{E}) = D^b(\mathcal{P}(\mathcal{E}))$$

and the algebraic K -groups are such that

$$\begin{cases} K_n(\mathcal{C}^b(\mathcal{E}), w_{\mathcal{E}}) = K_n(\mathcal{E}) & \text{for } n \geq 0 \text{ (Gillet [20])} \\ K\mathcal{P}_n(\mathcal{C}^b(\mathcal{E}), w_{\mathcal{E}}) = K_n(\mathcal{P}(\mathcal{E})) & \text{for } n \in \mathbb{Z} \text{ (Schlichting [38]).} \end{cases}$$

By [45, 1.9.2] the Waldhausen category defined in the same way but with cofibrations the chain maps which are degreewise admissible monomorphisms has the same algebraic K -theory.

Definition 5.1 Let $F: \mathcal{E} \rightarrow \mathcal{D}$ be a functor of exact categories.

(i) The algebraic K -groups $K\mathcal{P}_*(\mathcal{E}, \mathcal{D})$ are

$$K\mathcal{P}_n(\mathcal{E}, \mathcal{D}) = K\mathcal{P}_n(\mathcal{C}^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}) \quad (n \in \mathbb{Z})$$

with $(\mathcal{C}^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}) \subset (\mathcal{C}^b(\mathcal{E}), w_{\mathcal{E}})$ the Waldhausen subcategory with $\mathcal{C}^b(\mathcal{E}, \mathcal{D}) \subset \mathcal{C}^b(\mathcal{E})$ the full subcategory with objects the bounded chain complexes C in \mathcal{E} which are chain equivalent in \mathcal{D} to acyclic complexes, and

$$w_{(\mathcal{E}, \mathcal{D})} = w_{\mathcal{E}} \cap \mathcal{C}^b(\mathcal{E}, \mathcal{D}) \subset \mathcal{C}^b(\mathcal{E}, \mathcal{D})$$

the subcategory of the quasi-isomorphisms.

(ii) The algebraic ΓK -groups of F are


$$\Gamma K_n(F) = K\mathcal{P}_n(\mathcal{C}^b(\mathcal{E}), w_{\mathcal{D}}) \quad (n \in \mathbb{Z})$$

with $w_{\mathcal{D}} \subset \mathcal{C}^b(\mathcal{E})$ the subcategory with morphisms the chain maps in \mathcal{E} which become quasi-isomorphisms in \mathcal{D} , or equivalently such that the mapping cones are in $\mathcal{C}^b(\mathcal{E}, \mathcal{D})$.

The groups $\Gamma K_*(F)$ are the algebraic K -theory analogues of the algebraic L -theory groups $\Gamma_*(F)$ of Cappell and Shaneson [8].

Theorem 5.2 Let $F: \mathcal{E} \rightarrow \mathcal{D}$ be a functor of exact categories.

(i) The algebraic K -groups fit into a commutative braid of exact sequences



with $\Gamma F: (C^b(\mathcal{E}), w_{\mathcal{D}}) \rightarrow (C^b(\mathcal{D}), w_{\mathcal{D}})$ induced by F .

(ii) If $\Gamma F: \mathcal{P}D(C^b(\mathcal{E}), w_{\mathcal{D}}) \rightarrow \mathcal{P}D(C^b(\mathcal{D}), w_{\mathcal{D}})$ is an equivalence of categories then

$$KP_*(\Gamma F) = 0, KP_{*+1}(F) \cong KP_*(\mathcal{E}, \mathcal{D}), \Gamma K_*(F) \cong KP_*(\mathcal{D})$$

and the braid of (i) collapses to the exact sequence

$$\cdots \longrightarrow KP_{n+1}(\mathcal{D}) \longrightarrow KP_n(\mathcal{E}, \mathcal{D}) \longrightarrow KP_n(\mathcal{E}) \xrightarrow{F} KP_n(\mathcal{D}) \longrightarrow \cdots$$

(iii) The hypothesis of (ii) is satisfied if $F: \mathcal{E} \rightarrow \mathcal{D} = \Sigma^{-1}\mathcal{E}$ is the canonical functor to a category of fractions and \mathcal{D} has a calculus of left fractions.

Proof (i) The cases $n \geq 0$ are a direct application of the version of the localization theorem of [50, 1.6.4] stated in Theorem 2.3 and Lemma 2.5 of Neeman and Ranicki [31], with

$$\mathcal{R}^c = D(C^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}) \subset \mathcal{S}^c = D(C^b(\mathcal{E}), w_{\mathcal{E}}), \quad \mathcal{S}^c / \mathcal{R}^c \approx D(C^b(\mathcal{E}), w_{\mathcal{D}}),$$

$$\mathbf{R} = (C^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}), \quad \mathbf{S} = (C^b(\mathcal{E}), w_{\mathcal{E}}), \quad \mathbf{T} = \mathbf{S}_{\mathbf{R}} = (C^b(\mathcal{E}), w_{\mathcal{D}})$$

giving a fibration sequence of connective spectra

$$K(C^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}) \rightarrow K(C^b(\mathcal{E}), w_{\mathcal{E}}) \rightarrow K(C^b(\mathcal{E}), w_{\mathcal{D}}).$$

The cases $n < 0$ follow from Theorems 2.4, 3.7 of [31] and Schlichting [38, Theorems 1,9], which give a fibration sequence of nonconnective spectra

$$KP(C^b(\mathcal{E}, \mathcal{D}), w_{(\mathcal{E}, \mathcal{D})}) \rightarrow KP(C^b(\mathcal{E}), w_{\mathcal{E}}) \rightarrow KP(C^b(\mathcal{E}), w_{\mathcal{D}}).$$

(ii) This is a direct application of the Approximation Theorem of Waldhausen [50, Theorem 1.6.7]: if $F: (\mathcal{C}, w) \rightarrow (\mathcal{C}', w')$ is a functor which induces an equivalence of

the homotopy categories $F: D(\mathcal{C}, w) \rightarrow D(\mathcal{C}', w')$ then $F: K(\mathcal{C}, w) \rightarrow K(\mathcal{C}', w')$ is a homotopy equivalence inducing isomorphisms $F: K_*(\mathcal{C}, w) \cong K_*(\mathcal{C}', w')$. Similarly, if $F: \mathcal{P}D(\mathcal{C}, w) \rightarrow \mathcal{P}D(\mathcal{C}', w')$ is an equivalence there are induced isomorphisms $F: K\mathcal{P}_*(\mathcal{C}, w) \cong K\mathcal{P}_*(\mathcal{C}', w')$ ([38]).

(iii) Every object D in $\mathcal{C}^b(\mathcal{D})$ is chain equivalent to $F(E)$ for an object E in $\mathcal{C}^b(\mathcal{E})$, and the functors $F: \mathcal{C}^b(\mathcal{E}) \rightarrow \mathcal{C}^b(\mathcal{D})$, $F: D(\mathcal{C}^b(\mathcal{E}), w_{\mathcal{D}}) \rightarrow D(\mathcal{C}^b(\mathcal{D}), w_{\mathcal{D}})$ are localizations. □

Definition 5.3 (i) Write the algebraic K -groups of the exact categories $\mathcal{P}rim(A)$, $\mathcal{S}ei(A)$, $\mathcal{B}la(A)$, $\mathcal{F}lk(A)$ as

$$\begin{aligned} \text{Prim}_*(A) &= K_*(\mathcal{P}rim(A)), \text{Sei}_*(A) = K_*(\mathcal{S}ei(A)), \\ \text{Bla}_*(A) &= K_*(\mathcal{B}la(A)), \text{Flk}_*(A) = K_*(\mathcal{F}lk(A)). \end{aligned}$$

(ii) Write the algebraic K -groups of the idempotent completion of the homotopy category of $(\mathcal{C}^b(\mathcal{S}ei(A), \mathcal{B}la(A)), w_{(\mathcal{S}ei(A), \mathcal{B}la(A))})$ as

$$(\text{Sei}, \text{Bla})_*(A) = K\mathcal{P}_*(\mathcal{C}^b(\mathcal{S}ei(A), \mathcal{B}la(A)), w_{(\mathcal{S}ei(A), \mathcal{B}la(A))}).$$

Proposition 5.4 The covering functor $B: \mathcal{S}ei(A) \rightarrow \mathcal{B}la(A)$ induces morphisms $B: \text{Sei}_*(A) \rightarrow \text{Bla}_*(A)$ which fit into a long exact sequence

$$\cdots \longrightarrow (\text{Sei}, \text{Bla})_n(A) \longrightarrow \text{Sei}_n(A) \xrightarrow{B} \text{Bla}_n(A) \longrightarrow (\text{Sei}, \text{Bla})_{n-1}(A) \longrightarrow \cdots$$

with

$$\text{im}(B: \text{Sei}_0(A) \rightarrow \text{Bla}_0(A)) = \text{Flk}_0(A) \subseteq \text{Bla}_0(A).$$

Proof Apply [Theorem 5.2](#) (iii) with

$$F: \mathcal{E} = \mathcal{S}ei(A) \rightarrow \mathcal{D} = \Xi^{-1}\mathcal{S}ei(A) \approx \mathcal{F}lk(A),$$

noting that $\mathcal{S}ei(A)$ is idempotent complete ([Proposition 3.10](#) (i)), that $\Xi^{-1}\mathcal{S}ei(A) \approx \mathcal{F}lk(A)$ has a left calculus of fractions by [Theorem 3.17](#), and that $\mathcal{B}la(A) \approx \mathcal{P}(\mathcal{F}lk(A))$ ([Proposition 3.10](#)(ii)). □

In the next section it will be shown that the functor

$$\mathcal{P}rim(A) \rightarrow \mathcal{C}^b(\mathcal{S}ei(A), \mathcal{B}la(A)); (P, e, \{\pi_i\}) \mapsto (\cdots \rightarrow 0 \rightarrow (P, e, \{\pi_i\}))$$

induces isomorphisms of algebraic K -groups $\text{Prim}_*(A) \cong (\text{Sei}, \text{Bla})_*(A)$.

5.2 The algebraic K -theory of noncommutative localizations

Given a ring R let $\text{Mod}(R)$ be the abelian category of R -modules, so that $\text{Proj}(R) \subset \text{Mod}(R)$ is an exact subcategory. Write the Waldhausen category of $\text{Proj}(R)$ as

$$(\mathcal{C}^b(R), w_R) = (\mathcal{C}^b(\text{Proj}(R)), w_{\text{Proj}(R)}).$$

An object in $\mathcal{C}^b(R)$ is a bounded chain complex C of f.g. projective R -modules; C is acyclic if and only if $H_*(C) = 0$. A morphism $f: C \rightarrow D$ in $\mathcal{C}^b(R)$ is a chain map; f is in w_R if and only if $f_*: H_*(C) \rightarrow H_*(D)$ is an isomorphism. The algebraic K -groups of R are given by

$$K_*(R) = K_*(\text{Proj}(R)) = K\mathcal{P}_*(\mathcal{C}^b(R), w_R).$$

A ring morphism $\mathcal{F}: R \rightarrow S$ induces a functor of abelian categories

$$\mathcal{F} = S \otimes_R -: \text{Mod}(R) \rightarrow \text{Mod}(S); P \mapsto S \otimes_R P$$

which restricts to an exact functor $F: \text{Proj}(R) \rightarrow \text{Proj}(S)$. There is also induced a functor of Waldhausen categories

$$\mathcal{F}: (\mathcal{C}^b(R), w_R) \rightarrow (\mathcal{C}^b(S), w_S); C \mapsto S \otimes_R C.$$

The relative homotopy groups of $\mathcal{F}: K(R) \rightarrow K(S)$ are the relative K -groups $K_*(\mathcal{F})$ in the long exact sequence

$$\cdots \longrightarrow K_n(R) \xrightarrow{\mathcal{F}} K_n(S) \longrightarrow K_n(\mathcal{F}) \longrightarrow K_{n-1}(R) \longrightarrow \cdots.$$

Let R be a ring, and let Σ be a set of morphisms of f.g. projective R -modules. A ring morphism $R \rightarrow T$ is Σ -inverting if each $(s: P \rightarrow Q) \in \Sigma$ induces a T -module isomorphism $1 \otimes s: T \otimes_R P \rightarrow T \otimes_R Q$. By Cohn [10] there exists a *universal Σ -inverting localization* ring morphism

$$\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$$

such that any Σ -inverting ring morphism $R \rightarrow T$ has a unique factorization

$$R \xrightarrow{\mathcal{F}} S \longrightarrow T.$$

The category of fractions $\Sigma^{-1}\text{Proj}(R)$ is equivalent to the full subcategory

$$\text{Proj}_R(S) \subseteq \text{Proj}(S)$$

with objects isomorphic to the f.g. projective S -modules $\Sigma^{-1}P = S \otimes_R P$ induced from f.g. projective R -modules P , and $\text{Proj}(S) = \mathcal{P}(\text{Proj}_R(S))$ is the idempotent completion.

Definition 5.5 (i) For any ring morphism $\mathcal{F}: R \rightarrow S$ write the Waldhausen categories defined in Definition 5.1 as

$$\begin{aligned} (\mathcal{C}^b(\mathcal{P}roj(R), \mathcal{P}roj(S)), w_{(\mathcal{P}roj(R), \mathcal{P}roj(S))}) &= (\mathcal{C}^b(R, S), w_{(R, S)}), \\ (\mathcal{C}^b(\mathcal{P}roj(R)), w_S) &= (\mathcal{C}^b(R), w_S) \end{aligned}$$

with corresponding nonconnective algebraic K -theory spectra

$$K\mathcal{P}(\mathcal{C}^b(R, S), w_{(R, S)}) = K(R, S), \quad K\mathcal{P}(\mathcal{C}^b(R), w_S) = \Gamma K(\mathcal{F})$$

and algebraic K -groups $K_*(R, S), \Gamma K_*(\mathcal{F})$. An object in $\mathcal{C}^b(R, S)$ is a bounded chain complex C of f.g. projective R -modules such that $H_*(S \otimes_R C) = 0$. A morphism $f: C \rightarrow D$ in $\mathcal{C}^b(R, S)$ is a chain map; f is in $w_{(R, S)}$ if and only if $f_*: H_*(C) \rightarrow H_*(D)$ is an isomorphism. A morphism $f: C \rightarrow D$ in $\mathcal{C}^b(R)$ is in w_S if and only if $1 \otimes f: H_*(S \otimes_R C) \rightarrow H_*(S \otimes_R D)$ is an isomorphism.

(ii) For an injective universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$ let $H(R, \Sigma)$ be the exact category of *h.d. 1* Σ -torsion R -modules, ie the cokernels of injective morphisms $s: P \rightarrow Q$ of f.g. projective R -modules which induce an S -module isomorphism $1 \otimes s: S \otimes_R P \rightarrow S \otimes_R Q$ (eg if $s \in \Sigma$).

(iii) (Neeman and Ranicki [30; 31]) A universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$ is *stably flat* if

$$\text{Tor}_i^R(S, S) = 0 \quad (i \geq 1).$$


In particular, a universal localization $\mathcal{F}: R \rightarrow S$ is stably flat if S has flat dimension ≤ 1 as an R -module, ie if there exists a 1-dimensional flat R -module resolution

$$0 \rightarrow F_1 \rightarrow F_0 \rightarrow S \rightarrow 0.$$

Proposition 5.6 (i) For any ring morphism $\mathcal{F}: R \rightarrow S$ the functor

$$\Gamma \mathcal{F}: (\mathcal{C}^b(R), w_S) \rightarrow (\mathcal{C}^b(S), w_S); \quad C \mapsto S \otimes_R C$$

induces morphisms of algebraic K -groups $\Gamma \mathcal{F}: \Gamma K_*(\mathcal{F}) \rightarrow K_*(S)$ which fit into a commutative braid of exact sequences



(ii) For any universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$

$$\Gamma K_n(\mathcal{F}) = K_n(S), K_n(\mathcal{F}) = K_{n-1}(R, S), K_n(\Gamma\mathcal{F}) = 0 \quad (n \leq 1).$$

(iii) For a stably flat universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$

$$\Gamma K_*(\mathcal{F}) = K_*(S), K_{*+1}(\mathcal{F}) = K_*(R, S), K_*(\Gamma\mathcal{F}) = 0,$$

and there is induced a localization exact sequence in the algebraic K -groups

$$\cdots \longrightarrow K_n(R, S) \longrightarrow K_n(R) \xrightarrow{\mathcal{F}} K_n(S) \longrightarrow K_{n-1}(R, S) \longrightarrow \cdots$$

(iv) For an injective universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$ there is defined an equivalence of homotopy categories

$$D(\mathcal{C}^b(R, S), w_{(R, S)}) \approx D(\mathcal{C}^b(H(R, \Sigma)), w_{H(R, \Sigma)})$$

inducing isomorphisms

$$K_*(R, S) \cong K_*(H(R, \Sigma)).$$

(v) For an injective stably flat universal localization $\mathcal{F}: R \rightarrow S = \Sigma^{-1}R$ there is defined a localization exact sequence in the algebraic K -groups

$$\cdots \longrightarrow K_n(H(R, \Sigma)) \longrightarrow K_n(R) \xrightarrow{\mathcal{F}} K_n(\Sigma^{-1}R) \longrightarrow K_{n-1}(H(R, \Sigma)) \longrightarrow \cdots$$

Proof (i) Immediate from [Theorem 5.2](#) (i) and (ii) applied to $\mathcal{F}: \mathcal{C}^b(R) \rightarrow \mathcal{C}^b(S)$.

(ii)–(v) See Neeman and Ranicki [\[30; 31\]](#). □

5.3 Triangular matrix rings

We refer to Haghany and Varadarajan [\[23\]](#) for the general theory of modules over triangular matrix rings, and to Schofield [\[39\]](#), Ranicki [\[36\]](#) and Sheiham [\[42\]](#) for previous accounts of the universal localization of triangular matrix rings.

Proposition 5.7 *Let*

$$A = \begin{pmatrix} A_1 & B \\ 0 & A_2 \end{pmatrix}$$

be the triangular 2×2 matrix ring defined by rings A_1, A_2 and an (A_1, A_2) -bimodule B .

(i) An A -module $L = (L_1, L_2, \lambda)$ is defined by an A_1 -module L_1 , an A_2 -module L_2 and an A_1 -module morphism $\lambda: B \otimes_{A_2} L_2 \rightarrow L_1$. As an additive group $L = L_1 \oplus L_2$, written $\begin{pmatrix} L_1 \\ L_2 \end{pmatrix}$, with

$$\begin{pmatrix} A_1 & B \\ 0 & A_2 \end{pmatrix} \times \begin{pmatrix} L_1 \\ L_2 \end{pmatrix} \rightarrow \begin{pmatrix} L_1 \\ L_2 \end{pmatrix}: \left(\begin{pmatrix} a_1 & b \\ 0 & a_2 \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) \rightarrow \begin{pmatrix} a_1 x_1 + \lambda(b \otimes x_2) \\ a_2 x_2 \end{pmatrix}.$$

(ii) An A -module morphism $(f_1, f_2): (L_1, L_2, \lambda) \rightarrow (L'_1, L'_2, \lambda')$ is defined by an A_1 -module morphism $f_1: L_1 \rightarrow L'_1$, and an A_2 -module morphism $f_2: L_2 \rightarrow L'_2$ such that the diagram

$$\begin{array}{ccc} B \otimes_{A_2} L_2 & \xrightarrow{\lambda} & L_1 \\ 1 \otimes f_2 \downarrow & & \downarrow f_1 \\ B \otimes_{A_2} L'_2 & \xrightarrow{\lambda'} & L'_1 \end{array}$$

commutes.

(iii) An A -module $L = (L_1, L_2, \lambda)$ is f.g. projective if and only if λ is injective, $\text{coker}(\lambda)$ is a f.g. projective A_1 -module, and L_2 is a f.g. projective A_2 -module.

(iv) The projection

$$\text{Proj}(A) \rightarrow \text{Proj}(A_1) \times \text{Proj}(A_2); L = (L_1, L_2, \lambda) \mapsto (\text{coker}(\lambda), L_2)$$

induces isomorphisms

$$K_*(A) \cong K_*(A_1) \oplus K_*(A_2).$$

(v) If an A -module $L = (L_1, L_2, \lambda)$ is h.d. 1 then

(1) the 1-dimensional A_1 -module chain complex

$$K: \dots \longrightarrow 0 \longrightarrow B \otimes_{A_2} L_2 \xrightarrow{\lambda} L_1$$

is such that there exists a quasi-isomorphism (= homology equivalence) $J \rightarrow K$ for a 1-dimensional f.g. projective A_1 -module chain complex J , and

(2) L_2 is an h.d. 1 A_2 -module.

If B is a flat right A_2 -module the converse also holds: an A -module L is h.d. 1 if and only if conditions 1. and 2. are satisfied.

(vi) The columns of A are f.g. projective A -modules

$$S_1 = (A_1, 0, 0), S_2 = (B, A_2, 1)$$

with

$$S_1 \oplus S_2 = A, \text{End}(S_1) = A_1, \text{End}(S_2) = A_2.$$

The universal localization of A inverting a non-empty subset $\Sigma \subseteq \text{Hom}_A(S_1, S_2)$ is a morphism of 2×2 matrix rings

$$A = \begin{pmatrix} A_1 & B \\ 0 & A_2 \end{pmatrix} \rightarrow \Sigma^{-1}A = M_2(C) = \begin{pmatrix} C & C \\ C & C \end{pmatrix}$$

with C the endomorphism ring of the induced f.g. projective $\Sigma^{-1}A$ -module

$$\Sigma^{-1}S_1 \cong \Sigma^{-1}S_2.$$

The composite of the functor

$$\Sigma^{-1}: \text{Mod}(A) \rightarrow \text{Mod}(\Sigma^{-1}A); P \mapsto \Sigma^{-1}P = \Sigma^{-1}A \otimes_A P$$

and the Morita equivalence of categories

$$\begin{aligned} (C \ C) \otimes_{\Sigma^{-1}A} - : \text{Mod}(\Sigma^{-1}A) &\xrightarrow{\cong} \text{Mod}(C); \\ L = (L_1, L_2, \lambda) &\mapsto (C \ C) \otimes_{\Sigma^{-1}A} L \end{aligned}$$

is the assembly functor

$$\begin{aligned} \text{Mod}(A) &\longrightarrow \text{Mod}(C); \\ L = (L_1, L_2, \lambda) &\mapsto (C \ C) \otimes_{\Sigma^{-1}A} \Sigma^{-1}L = (C \ C) \otimes_A L \\ &= \text{coker} \left(\begin{pmatrix} 1 \otimes \lambda \\ \kappa \otimes 1 \end{pmatrix} : C \otimes_{A_1} B \otimes_{A_2} L_2 \rightarrow C \otimes_{A_1} L_1 \oplus C \otimes_{A_2} L_2 \right) \end{aligned}$$

with

$$\kappa: C \otimes_{A_1} B \rightarrow C; x \otimes y \mapsto xy$$

the (C, A_2) -bimodule morphism defined by multiplication in C , using the A_1 -module morphism $B \rightarrow C$. The assembly functor $\text{Proj}(A) \rightarrow \text{Proj}(C)$ induces the morphisms

$$\Sigma^{-1}: K_*(A) = K_*(A_1) \oplus K_*(A_2) \rightarrow K_*(\Sigma^{-1}A) = K_*(C).$$

(vii) If B and C are flat A_1 -modules and C is a flat A_2 -module then the A -module $\begin{pmatrix} C \\ C \end{pmatrix}$ has a 1-dimensional flat A -module resolution

$$0 \rightarrow \begin{pmatrix} B \\ 0 \end{pmatrix} \otimes_{A_2} C \rightarrow \begin{pmatrix} A_1 \\ 0 \end{pmatrix} \otimes_{A_1} C \oplus \begin{pmatrix} B \\ A_2 \end{pmatrix} \otimes_{A_2} C \rightarrow \begin{pmatrix} C \\ C \end{pmatrix} \rightarrow 0$$

so that $\Sigma^{-1}A = \begin{pmatrix} C \\ C \end{pmatrix} \oplus \begin{pmatrix} C \\ C \end{pmatrix}$ is stably flat. An h.d. 1 A -module $L = (L_1, L_2, \lambda)$ is Σ -torsion if and only if the C -module morphism

$$\begin{pmatrix} 1 \otimes \lambda \\ \kappa \otimes 1 \end{pmatrix} : C \otimes_{A_1} B \otimes_{A_2} L_2 \rightarrow C \otimes_{A_1} L_1 \oplus C \otimes_{A_2} L_2$$

is an isomorphism.

Proof (i) and (ii) Standard.

(iii) For any A -module $L = (L_1, L_2, \lambda)$ there is defined an exact sequence

$$0 \rightarrow (\ker(\lambda), 0, 0) \rightarrow (B \otimes_{A_2} L_2, L_2, 1) \xrightarrow{(\lambda, 1)} (L_1, L_2, \lambda) \rightarrow (\operatorname{coker}(\lambda), 0, 0) \rightarrow 0.$$

Now $(A_1, 0, 0) = \begin{pmatrix} A_1 \\ 0 \end{pmatrix}$ and $(B, A_2, 1) = \begin{pmatrix} B \\ A_2 \end{pmatrix}$ are f.g. projective A -modules, since

$$(A_1, 0, 0) \oplus (B, A_2, 1) = \left(A_1 \oplus B, A_2, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) = A.$$

If $\ker(\lambda) = 0$ and $\operatorname{coker}(\lambda)$ is a f.g. projective A_1 -module then $(\operatorname{coker}(\lambda), 0, 0) = (A_1, 0, 0) \otimes_{A_1} \operatorname{coker}(\lambda)$ is a f.g. projective A -module. If L_2 is a f.g. projective A_2 -module then

$$(B \otimes_{A_2} L_2, L_2, 1) = \begin{pmatrix} B \\ A_2 \end{pmatrix} \otimes_{A_2} L_2$$

is a f.g. projective A -module. Thus if these two conditions are satisfied then the exact sequence splits and L is a f.g. projective A -module.

Conversely, suppose that (L_1, L_2, λ) is a f.g. projective A -module, so that there exists an A -module (L'_1, L'_2, λ') with an A -module isomorphism

$$(L_1, L_2, \lambda) \oplus (L'_1, L'_2, \lambda') \cong \left((A_1)^k \oplus B^k, (A_2)^k, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) = A^k$$

for some $k \geq 0$. It follows from $\ker(\lambda \oplus \lambda') = 0$ that $\ker(\lambda) = 0$, and from $\operatorname{coker}(\lambda \oplus \lambda') \cong (A_1)^k$ that $\operatorname{coker}(\lambda)$ is a f.g. projective A_1 -module. Also, $L_2 \oplus L'_2 \cong (A_2)^k$, so that L_2 is a f.g. projective A_2 -module.

(iv) The result that the inclusion and projection

$$i = A_1 \times A_2 \rightarrow A, \quad j: A \rightarrow A_1 \times A_2$$

induce inverse isomorphisms

$$K_*(A_1 \times A_2) = K_*(A_1) \oplus K_*(A_2) \begin{matrix} \xleftarrow{i_*} \\ \xrightarrow{j_*} \end{matrix} K_*(A)$$

was first obtained by Berrick and Keating [6]. Here is a proof using Waldhausen K -theory. It is immediate from $ji = 1$ that

$$j_*i_* = 1: K_*(A_1 \times A_2) \rightarrow K_*(A) \rightarrow K_*(A_1 \times A_2).$$

Every f.g. projective A -module $L = (L_1, L_2, \lambda: B \otimes_{A_2} L_2 \rightarrow L_1)$ fits into a natural short exact sequence of f.g. projective A -modules

$$0 \rightarrow (B \otimes_{A_2} L_2, L_2, 1) \xrightarrow{(\lambda, 1)} (L_1, L_2, \lambda) \rightarrow (\text{coker}(\lambda), 0) \rightarrow 0.$$

The functors

$$F_1: \mathcal{P}roj(A) \rightarrow \mathcal{P}roj(A); L \mapsto A \otimes_{A_1} A_1 \otimes_A L = (\text{coker}(\lambda), 0),$$

$$F_2: \mathcal{P}roj(A) \rightarrow \mathcal{P}roj(A); L \mapsto A \otimes_{A_2} A_2 \otimes_A L = (B \otimes_{A_2} L_2, L_2, 1)$$

fit into a cofibration sequence

$$F_2 \rightarrow 1_{\mathcal{P}roj(A)} \rightarrow F_1,$$

and are such that

$$F_k: K_*(A) \rightarrow K_*(A_k) \rightarrow K_*(A) \quad (k = 1, 2).$$

Now apply the additivity theorem for Quillen K -theory [50, Proposition 1.3.2 (4)] to identify

$$i_*j_* = F_1 + F_2 = 1: K_*(A) \rightarrow K_*(A),$$

so that i_* , j_* are inverse isomorphisms.

(v) If $L = (L_1, L_2, \lambda)$ is an h.d. 1 A -module there exists a 1-dimensional f.g. projective A -module resolution

$$0 \longrightarrow (P_1, P_2, f) \xrightarrow{(h_1, h_2)} (Q_1, Q_2, g) \longrightarrow (L_1, L_2, \lambda) \longrightarrow 0,$$

so that $\text{coker}(f)$, $\text{coker}(g)$ are f.g. projective A_1 -modules and P_2 , Q_2 are f.g. projective A_2 -modules. The 1-dimensional A_1 -module chain complex

$$K: \dots \longrightarrow 0 \longrightarrow B \otimes_{A_2} L_2 \xrightarrow{\lambda} L_1$$

and the 1-dimensional f.g. projective A_1 -module chain complex

$$J: J_1 = \text{coker}(f) \xrightarrow{h_1} J_0 = \text{coker}(g)$$

are related by a homology equivalence $J \rightarrow K$. Furthermore, $L_2 = \text{coker}(h_2)$ is an h.d. 1 A_2 -module. Thus both conditions 1. and 2. are satisfied.

Conversely, suppose that B is a flat right A_2 -module and that $L = (L_1, L_2, \lambda)$ is an A -module such that there exists a homology equivalence $J \rightarrow K$ with J a 1-dimensional f.g. projective A_1 -module chain complex and that L_2 is an h.d. 1 A_2 -module with a 1-dimensional f.g. projective A_2 -module resolution

$$0 \rightarrow P_2 \rightarrow Q_2 \rightarrow L_2 \rightarrow 0.$$

There is induced a short exact sequence of A_1 -modules

$$0 \rightarrow B \otimes_{A_2} P_2 \rightarrow B \otimes_{A_2} Q_2 \rightarrow B \otimes_{A_2} L_2 \rightarrow 0$$

and it follows from the 1-dimensional f.g. projective A -module resolution of L

$$0 \rightarrow (B \otimes_{A_2} P_2, P_2, 1) \oplus (J_1, 0, 0) \rightarrow (B \otimes_{A_2} Q_2, Q_2, 1) \oplus (J_0, 0, 0) \rightarrow L \rightarrow 0$$

that L is an h.d. 1 A -module.

(vi) and (vii) See [36, 2.2]. □

We shall actually be working with $(\mu + 1) \times (\mu + 1)$ -matrix rings:

Definition 5.8 For any ring A and $\mu \geq 1$ define the triangular $(\mu + 1) \times (\mu + 1)$ -matrix ring

$$T_\mu(A) = \begin{pmatrix} A & A \oplus A & A \oplus A & \dots & A \oplus A \\ 0 & A & 0 & \dots & 0 \\ 0 & 0 & A & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & A \end{pmatrix}.$$

The ring $T_\mu(A)$ is the A -coefficient path algebra of the quiver with vertices $0, 1, \dots, \mu$ and two arrows $s_i^+, s_i^-: i \rightarrow 0$ for $i = 1, 2, \dots, \mu$. A $T_\mu(A)$ -module $L = (L_i, f_i^+, f_i^-)$ consists of A -modules L_0, L_1, \dots, L_μ and A -module morphisms $f_i^+, f_i^-: L_i \rightarrow L_0$ ($1 \leq i \leq \mu$).

Let S_0, S_1, \dots, S_μ be the $T_\mu(A)$ -modules defined by the columns of $T_\mu(A)$, so that

$$S_0 = (A, 0, \dots, 0; 0, \dots, 0),$$

$$S_i = (A \oplus A, 0, \dots, 0, A, 0, \dots, 0; 0, \dots, 0, \text{id.}, 0, \dots, 0) \quad (1 \leq i \leq \mu).$$

It follows from

$$S_0 \oplus S_1 \oplus \dots \oplus S_\mu = T_\mu(A)$$

that each S_i is a f.g. projective $T_\mu(A)$ -module. Let $\sigma = \{s_i^+, s_i^-\}$ be the set of f.g. projective $T_\mu(A)$ -module morphisms

$$s_i^+ = ((1 \ 0), 0, \dots, 0), \quad s_i^- = ((0 \ 1), 0, \dots, 0): S_i \rightarrow S_0 \quad (1 \leq i \leq \mu).$$

Proposition 5.9 (i) The universal σ -inverting localization of $T_\mu(A)$ is given by the inclusion

$$\mathcal{F}: T_\mu(A) \rightarrow \sigma^{-1}T_\mu(A) = M_{\mu+1}(A[F_\mu])$$

with $M_{\mu+1}(A[F_\mu])$ the ring of all $(\mu + 1) \times (\mu + 1)$ -matrices with entries in $A[F_\mu]$. The universal localization \mathcal{F} is both injective and stably flat.

(ii) The composite

$$\text{Mod}(T_\mu(A)) \xrightarrow{\mathcal{F}} \text{Mod}(M_{\mu+1}(A[F_\mu])) \xrightarrow[\approx]{\text{Morita}} \text{Mod}(A[F_\mu])$$

sends a $T_\mu(A)$ -module $L = (L_i, f_i^+, f_i^-)$ to the assembly $A[F_\mu]$ -module

$$M = \text{coker}\left((f_1^+ z_1 - f_1^- \dots f_\mu^+ z_\mu - f_\mu^-): \bigoplus_{i=1}^{\mu} L_i[F_\mu] \rightarrow L_0[F_\mu]\right).$$

(iii) A $T_\mu(A)$ -module $L = (L_i, f_i^+, f_i^-)$ is f.g. projective if and only if L_0, \dots, L_μ are f.g. projective A -modules and the A -module morphism

$$(f_1^+ \ f_1^- \ f_2^+ \ f_2^- \ \dots \ f_\mu^+ \ f_\mu^-): \bigoplus_{i=1}^{\mu} L_i \oplus L_i \rightarrow L_0$$

is a split injection. The projection

$$\text{Proj}(T_\mu(A)) \rightarrow \prod_{\mu+1} \text{Proj}(A); (L_i, f_i^+, f_i^-) \mapsto (L_0, L_1, L_2, \dots, L_\mu)$$

induces isomorphisms in algebraic K -theory

$$K_*(T_\mu(A)) \cong \bigoplus_{\mu+1} K_*(A).$$

(iv) A $T_\mu(A)$ -module $L = (L_i, f_i^+, f_i^-)$ is h.d. 1 σ -torsion if and only if L_0, \dots, L_μ are f.g. projective A -modules and the $A[F_\mu]$ -module morphism

$$(f_1^+ z_1 - f_1^- \ f_2^+ z_2 - f_2^- \ \dots \ f_\mu^+ z_\mu - f_\mu^-): \bigoplus_{i=1}^{\mu} L_i[F_\mu] \rightarrow L_0[F_\mu]$$

is an isomorphism. A f.g. projective Seifert A -module $(P, e, \{\pi_i\})$ is primitive if and only if (P, P_i, f_i^+, f_i^-) is an h.d. 1 σ -torsion $T_\mu(A)$ -module. The functor

$$\text{Prim}(A) \rightarrow H(T_\mu(A), \sigma); (P, e, \{\pi_i\}) \mapsto (P, P_i, e\pi_i, (e - 1)\pi_i)$$

is an equivalence of exact categories, so that

$$\text{Prim}_*(A) = K_*(H(T_\mu(A), \sigma)).$$

The forgetful functor

$$\begin{aligned} \mathcal{P}rim(A) &\rightarrow \prod_{2\mu} \mathcal{P}roj(A); \\ &\left(P^+ \oplus P^-, \begin{pmatrix} e^{++} & e^{+-} \\ e^{-+} & e^{--} \end{pmatrix}, \{\pi_i^+\} \oplus \{\pi_i^-\} \right) \mapsto (P_1^+, P_1^-, \dots, P_\mu^+, P_\mu^-) \end{aligned}$$

(defined using [Theorem 4.9](#)) is split by

$$\prod_{2\mu} \mathcal{P}roj(A) \rightarrow \mathcal{P}rim(A); \quad (P_1^+, P_1^-, \dots, P_\mu^+, P_\mu^-) \mapsto (P^+ \oplus P^-, 0, \{\pi_i^+\} \oplus \{\pi_i^-\}).$$

The reduced K -groups defined by

$$\widetilde{\text{Prim}}_*(A) = \ker(\text{Prim}_*(A) \rightarrow \bigoplus_{2\mu} K_*(A))$$

are such that

$$K_*(H(T_\mu(A), \sigma)) = \text{Prim}_*(A) = \bigoplus_{2\mu} K_*(A) \oplus \widetilde{\text{Prim}}_*(A).$$

Proof The universal localization $\sigma^{-1}T_\mu(A)$ is the $(\mu + 1) \times (\mu + 1)$ -matrix ring $M_{\mu+1}(R)$ with R the endomorphism ring of the induced f.g. projective $\sigma^{-1}T_\mu(A)$ -module $\sigma^{-1}S_0$, and there is defined an isomorphism

$$A[F_\mu] \rightarrow R; \quad z_i \mapsto s_i^+(s_i^-)^{-1}.$$

The remaining parts are given by [Proposition 5.7](#), viewing the $(\mu + 1) \times (\mu + 1)$ matrix ring $T_\mu(A)$ as a triangular 2×2 matrix ring

$$T_\mu(A) = \begin{pmatrix} A_1 & B \\ 0 & A_2 \end{pmatrix}$$

with

$$A_1 = A, \quad A_2 = \begin{pmatrix} A & 0 & \dots & 0 \\ 0 & A & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & A \end{pmatrix}, \quad B = (A \oplus A \ \dots \ A \oplus A)$$

such that

$$\text{Mod}(A_2) = \prod_{\mu} \text{Mod}(A).$$

An A_2 -module is just a μ -tuple $(L_1, L_2, \dots, L_{\mu})$ of A -modules. By the 2×2 theory a $T_{\mu}(A)$ -module L just a $(\mu + 1)$ -tuple $(L_0, L_1, \dots, L_{\mu})$ of A -modules, together with A -module morphisms $f_i^+, f_i^-: L_i \rightarrow L_0$ ($1 \leq i \leq \mu$). Note that B is a flat right A_2 -module, and that for an h.d. 1 σ -torsion $T_{\mu}(A)$ -module $L = (L_i, f_i^+, f_i^-)$ each L_i ($0 \leq i \leq \mu$) is a f.g. projective A -module, by the following argument. The necessary and sufficient conditions of Proposition 5.7 (v) and (vii) for a $T_{\mu}(A)$ -module L to be h.d. 1 σ -torsion are:

- (i) there exists a 1-dimensional f.g. projective A -module chain complex $J: J_1 \rightarrow J_0$ with a homology equivalence

$$\begin{array}{ccc} J_1 & \longrightarrow & J_0 \\ \downarrow & & \downarrow \\ \bigoplus_{i=1}^{\mu} L_i \oplus L_i & \xrightarrow{(f_i^+ \ f_i^-)} & L_0, \end{array}$$

- (ii) each L_i ($1 \leq i \leq \mu$) is an h.d. 1 A -module,
- (iii) the $A[F_{\mu}]$ -module morphism

$$(f_i^+ z_i - f_i^-): \bigoplus_{i=1}^{\mu} L_i[F_{\mu}] \rightarrow L_0[F_{\mu}]$$

is an isomorphism.

If L satisfies these conditions there is defined a commutative diagram of A -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{i=1}^{\mu} L_i & \xrightarrow{\begin{pmatrix} 1 \\ -1 \end{pmatrix}} & \bigoplus_{i=1}^{\mu} L_i \oplus L_i & \xrightarrow{(1 \ 1)} & \bigoplus_{i=1}^{\mu} L_i & \longrightarrow & 0 \\ & & \cong \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & L_0 & \xrightarrow{1} & L_0 & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

with exact rows and with $(f_i^+ - f_i^-)$ an isomorphism. There are defined A -module isomorphisms

$$J_0 \oplus L_0 \cong J_0 \oplus \bigoplus_{i=1}^{\mu} L_i \cong J_1,$$

so that each L_i ($0 \leq i \leq \mu$) is a f.g. projective A -module. □

Example 5.10 The assembly of $A[F_\mu]$ -modules is an algebraic analogue of the geometric construction of an F_μ -cover \widetilde{W} of a space W from a fundamental domain $U \subset \widetilde{W}$. The subspaces

$$V_i = U \cap z_i^{-1}U, \quad z_i V_i = z_i U \cap U \subset U \quad (1 \leq i \leq \mu)$$

are disjoint, with embeddings

$$f_i^+ : V_i \rightarrow U; x \mapsto x, \quad f_i^- : V_i \rightarrow U; x \mapsto z_i x \quad (1 \leq i \leq \mu),$$

and \widetilde{W} can be constructed by glueing together F_μ copies of U

$$\begin{aligned} \widetilde{W} &= (F_\mu \times U) / \{(g, f_i^+(x)) \sim (gz_i, f_i^-(x)) \mid g \in F_\mu, x \in V_i, 1 \leq i \leq \mu\} \\ &= \bigcup_{g \in F_\mu} gU \text{ with } U \cap z_i^{-1}U = V_i. \end{aligned}$$

Such a situation arises if W is a manifold with a surjection $\pi_1(W) \rightarrow F_\mu$, eg a boundary link exterior. The surjection is induced by a map

$$c : W \rightarrow BF_\mu = \bigvee_{\mu} S^1$$

which is transverse regular at $\{1, 2, \dots, \mu\} \subset BF_\mu$. Cutting W open at the inverse image codimension-1 submanifolds $V_i = c^{-1}(\{i\}) \subset W$ there is obtained a fundamental domain $U \subset \widetilde{W}$ for the pullback $\widetilde{W} = c^*EF_\mu$ to W of the universal cover EF_μ of BF_μ . More generally, suppose that W is a finite CW complex with an F_μ -cover \widetilde{W} , and that $U \subset \widetilde{W}$ is a fundamental domain which is a subcomplex. The embeddings $f_i^+, f_i^- : V_i \rightarrow U$ induce inclusions of the cellular f.g. free \mathbb{Z} -module chain complexes $f_i^+, f_i^- : C(V_i) \rightarrow C(U)$. The f.g. projective $T_\mu(\mathbb{Z})$ -module chain complex $C = (C(U), C(V_i), f_i^+, f_i^-)$ has assembly the cellular f.g. free $\mathbb{Z}[F_\mu]$ -module chain complex of \widetilde{W}

$$\text{coker} \left((f_1^+ z_1 - f_1^- \dots f_\mu^+ z_\mu - f_\mu^-) : \bigoplus_{i=1}^{\mu} C(V_i)[F_\mu] \rightarrow C(U)[F_\mu] \right) = C(\widetilde{W}),$$

such that

$$C(\widetilde{W})_r = \text{coker} \left((f_1^+ \dots f_\mu^+) : \bigoplus_{i=1}^{\mu} C(V_i)_r \rightarrow C(U)_r \right) [F_\mu].$$

Theorem 5.11 The algebraic K -groups of $A[F_\mu]$ split as

$$K_*(A[F_\mu]) = K_*(A) \oplus \bigoplus_{\mu} K_{*-1}(A) \oplus \widetilde{\text{Prim}}_{*-1}(A).$$

Proof By Proposition 5.9 the universal localization

$$\mathcal{F}: A[F_\mu] \rightarrow \sigma^{-1}T_\mu(A) = M_{\mu+1}(A)$$

is injective and stably flat. The noncommutative localization exact sequence of Neeman and Ranicki [30; 31]

$$\begin{aligned} \cdots \longrightarrow K_{n+1}(\sigma^{-1}T_\mu(A)) &\longrightarrow K_n(H(T_\mu(A), \sigma)) \\ &\longrightarrow K_n(T_\mu(A)) \longrightarrow K_n(\sigma^{-1}T_\mu(A)) \longrightarrow \cdots \end{aligned}$$

is given by

$$\cdots \rightarrow K_{n+1}(A[F_\mu]) \rightarrow \text{Prim}_n(A) \rightarrow \bigoplus_{\mu+1} K_n(A) \rightarrow K_n(A[F_\mu]) \rightarrow \cdots$$

with $\text{Prim}_n(A) \rightarrow K_n(T_\mu(A)) = \bigoplus_{\mu+1} K_n(A)$ induced by

$$\text{Prim}(A) \rightarrow \prod_{\mu+1} \mathcal{P}\text{roj}(A); (P, e, \{\pi_i\}) \mapsto (P, P_1, P_2, \dots, P_\mu),$$

so that

$$\begin{aligned} \text{Prim}_n(A) &= \bigoplus_{2\mu} K_n(A) \oplus \widetilde{\text{Prim}}_n(A) \rightarrow \bigoplus_{\mu+1} K_n(A); \\ &(x_1^+, x_1^-, x_2^+, x_2^-, \dots, x_\mu^+, x_\mu^-, \tilde{x}) \\ &\mapsto \left(\sum_{i=1}^{\mu} (x_i^+ + x_i^-), x_1^+ + x_1^-, x_2^+ + x_2^-, \dots, x_\mu^+ + x_\mu^- \right). \end{aligned}$$

This completes the proof. □

Definition 5.12 Let $\mathcal{G}: A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu]$ be the universal localization inverting the set Σ of morphisms of f.g. projective $A[F_\mu]$ -modules which induce an isomorphism of f.g. projective A -modules under the augmentation $\epsilon: A[F_\mu] \rightarrow A; z_i \mapsto 1$.

Proposition 5.13 (i) *The universal localization $\mathcal{G}: A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu]$ is injective. The h.d. 1 Σ -torsion $A[F_\mu]$ -module category is*

$$H(A[F_\mu], \Sigma) = \text{Bla}(A).$$

(ii) *The composite*

$$\mathcal{GF}: T_\mu(A) \xrightarrow{\mathcal{F}} \sigma^{-1}T_\mu(A) = M_{\mu+1}(A[F_\mu]) \xrightarrow{\mathcal{G}} \tau^{-1}T_\mu(A) = M_{\mu+1}(\Sigma^{-1}A[F_\mu])$$

is the universal localization inverting the set τ of morphisms of f.g. projective $T_\mu(A)$ -modules which become isomorphisms under the composite

$$\epsilon\mathcal{F}: T_\mu(A) \xrightarrow{\mathcal{F}} \sigma^{-1}T_\mu(A) = M_{\mu+1}(A[F_\mu]) \xrightarrow{\epsilon} M_{\mu+1}(A).$$

(iii) A $T_\mu(A)$ -module $L = (L_i, f_i^+, f_i^-)$ is h.d. 1 τ -torsion if and only if L_0, \dots, L_μ are f.g. projective A -modules and the A -module morphism

$$f = (f_1^+ - f_1^-, f_2^+ - f_2^-, \dots, f_\mu^+ - f_\mu^-): L_1 \oplus L_2 \oplus \dots \oplus L_\mu \rightarrow L_0$$

is an isomorphism, if and only if

$$(P, e, \{\pi_i\}) = \left(\bigoplus_{i=1}^{\mu} L_i, f^{-1}(f_1^+ \ f_2^+ \ \dots \ f_\mu^+), \{\pi_i\} \right)$$

is a f.g. projective Seifert A -module. The functor

$$\mathcal{S}ei(A) \rightarrow H(T_\mu(A), \tau); (P, e, \{\pi_i\}) \mapsto (P, P_i, e\pi_i, (e-1)\pi_i)$$

is an equivalence of exact categories. The assembly of (L_i, f_i^+, f_i^-) is the covering Blanchfield $A[F_\mu]$ -module of $(P, e, \{\pi_i\})$

$$\begin{aligned} \text{coker} \left((f_1^+ z_1 - f_1^- \ \dots \ f_\mu^+ z_\mu - f_\mu^-): \bigoplus_{i=1}^{\mu} L_i[F_\mu] \rightarrow L_0[F_\mu] \right) \\ = \text{coker} (1 - e + ze: P[F_\mu] \rightarrow P[F_\mu]) = B(P, e, \{\pi_i\}), \end{aligned}$$

so that up to equivalence

$$\mathcal{F} = B: H(T_\mu(A), \tau) = \mathcal{S}ei(A) \rightarrow H(M_{\mu+1}(A[F_\mu]), \tau) = \mathcal{B}la(A).$$

(iv) The forgetful functor

$$\mathcal{S}ei(A) \rightarrow \prod_{\mu} \mathcal{P}roj(A); (P, e, \{\pi_i\}) \mapsto (P_1, P_2, \dots, P_\mu)$$

is split by

$$\prod_{\mu} \mathcal{P}roj(A) \rightarrow \mathcal{P}rim(A); (P_1, P_2, \dots, P_\mu) \mapsto \left(\bigoplus_{i=1}^{\mu} P_i, 0, \{\pi_i\} \right)$$

The reduced K -groups defined by

$$\widetilde{\mathcal{S}ei}_*(A) = \ker \left(\mathcal{S}ei_*(A) \rightarrow \bigoplus_{\mu} K_*(A) \right)$$

are such that

$$K_*(H(T_\mu(A), \tau)) = \text{Sei}_*(A) = \bigoplus_{\mu} K_*(A) \oplus \widetilde{\text{Sei}}_*(A).$$


Proof (i) The Magnus–Fox embedding $A[F_\mu] \rightarrow A\langle\langle x_1, \dots, x_\mu \rangle\rangle$ is Σ -inverting, so that there is a unique factorization

$$A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu] \rightarrow A\langle\langle x_1, x_2, \dots, x_\mu \rangle\rangle.$$

The identification $H(A[F_\mu], \Sigma) = \text{Bla}(A)$ is a formality, as is the identification $\mathcal{P}roj(A[F_\mu]) = \mathcal{P}(\mathcal{P}roj_A(A[F_\mu]))$ with $\mathcal{P}roj_A(A[F_\mu]) \subseteq \mathcal{P}roj(A[F_\mu])$ the full subcategory with objects isomorphic to the f.g. projective $A[F_\mu]$ -modules $P[F_\mu]$ induced from f.g. projective A -modules P .

(ii)–(iv) By construction, working as in the proof of Proposition 5.9 (iv) to show that if $L = (L_i, f_i^+, f_i^-)$ is an h.d. 1 τ -torsion $T_\mu(A)$ -module then L_0, L_1, \dots, L_μ are f.g. projective A -modules. \square

Theorem 5.14 (i) The algebraic K -groups of $\mathcal{P}rim(A)$, $\text{Sei}(A)$ and $\text{Bla}(A)$ fit into a commutative braid of exact sequences



for $n \in \mathbb{Z}$, with $\mathcal{G}: A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu]$ the universal localization and

$$\begin{aligned}
 K_*(T_\mu(A)) &= \bigoplus_{\mu+1} K_*(A), \\
 K_*(H(T_\mu(A), \sigma)) &= (\text{Sei}, \text{Bla})_*(A) = \text{Prim}_*(A) = \bigoplus_{2\mu} K_*(A) \oplus \widetilde{\text{Prim}}_*(A), \\
 K_*(H(T_\mu(A), \tau)) &= \text{Sei}_*(A) = \bigoplus_{\mu} K_*(A) \oplus \widetilde{\text{Sei}}_*(A), \\
 K_*(H(A[F_\mu], \Sigma)) &= \text{Bla}_*(A) = \bigoplus_{\mu} K_{*-1}(A) \oplus \widetilde{\text{Bla}}_*(A), \\
 \Gamma K_*(\mathcal{G}) &= K_*(A) \oplus \widetilde{\text{Sei}}_{*-1}^{\mu}(A) (= K_*(\Sigma^{-1}A[F_\mu]) \text{ for } * \leq 1)
 \end{aligned}$$

The reduced K -groups fit into a long exact sequence

$$\cdots \rightarrow \widetilde{\text{Prim}}_n(A) \rightarrow \widetilde{\text{Sei}}_n(A) \rightarrow \widetilde{\text{Bla}}_n(A) \rightarrow \widetilde{\text{Prim}}_{n-1}(A) \rightarrow \cdots .$$

(ii) If $\mathcal{G}: A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu]$ is stably flat then

$$\Gamma K_n(\mathcal{G}) = K_n(\Sigma^{-1}A[F_\mu]) = K_n(A) \oplus \widetilde{\text{Sei}}_{n-1}(A)$$

for all $n \in \mathbb{Z}$.

Proof (i) Consider the commutative square of Waldhausen categories

$$\begin{array}{ccc} (\mathcal{C}^b(T_\mu(A)), w_{T_\mu(A)}) & \longrightarrow & (\mathcal{C}^b(T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) \\ \mathcal{F} \downarrow & & \downarrow \\ (\mathcal{C}^b(T_\mu(A)), w_{\sigma^{-1}T_\mu(A)}) & \xrightarrow{\mathcal{G}} & (\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) . \end{array}$$

Since $\mathcal{F}: T_\mu(A) \rightarrow \sigma^{-1}T_\mu(A) = M_{\mu+1}(A[F_\mu])$ is stably flat there are defined equivalences

$$(\mathcal{C}^b(T_\mu(A)), w_{\sigma^{-1}T_\mu(A)}) \approx (\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\sigma^{-1}T_\mu(A)}) \approx (\mathcal{C}^b(A[F_\mu]), w_{A[F_\mu]})$$

which induce homotopy equivalences

$$K\mathcal{P}(\mathcal{C}^b(T_\mu(A)), w_{\sigma^{-1}T_\mu(A)}) \simeq K(\sigma^{-1}T_\mu(A)) \simeq K(A[F_\mu]) .$$

Also, since $\tau^{-1}T_\mu(A) = M_{\mu+1}(\Sigma^{-1}A[F_\mu])$ the functor

$$(\mathcal{C}^b(T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) \rightarrow (\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\tau^{-1}T_\mu(A)})$$

induces an equivalence of the homotopy categories

$$D(\mathcal{C}^b(T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) \approx D(\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) .$$


The composite of this equivalence and the Morita equivalence

$$D(\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) \approx D(\mathcal{C}^b(A[F_\mu]), w_{\Sigma^{-1}A[F_\mu]})$$


induces a homotopy equivalence

$$\begin{aligned} K\mathcal{P}(\mathcal{C}^b(\sigma^{-1}T_\mu(A)), w_{\tau^{-1}T_\mu(A)}) &\simeq K\mathcal{P}(\mathcal{C}^b(A[F_\mu]), w_{\Sigma^{-1}A[F_\mu]}) \\ &= \Gamma K(\mathcal{G}: A[F_\mu] \rightarrow \Sigma^{-1}A[F_\mu]) . \end{aligned}$$

Thus Propositions 5.6, 5.9 and 5.13 give a braid of Waldhausen categories



inducing a commutative braid of exact sequences



Split off the reduced K -groups in

$$\text{Prim}_*(A) = \bigoplus_{2\mu} K_*(A) \oplus \widetilde{\text{Prim}}_*(A),$$

$$\text{Sei}_*(A) = \bigoplus_{\mu} K_*(A) \oplus \widetilde{\text{Sei}}_*(A)$$

from the long exact sequence

$$\dots \rightarrow \text{Prim}_n(A) \rightarrow \text{Sei}_n(A) \rightarrow \text{Bla}_n(A) \rightarrow \text{Prim}_{n-1}(A) \rightarrow \dots$$

to define the reduced K -groups in

$$\text{Bla}_*(A) = \bigoplus_{\mu} K_{*-1}(A) \oplus \widetilde{\text{Bla}}_*(A)$$

and to obtain the long exact sequence

$$\dots \rightarrow \widetilde{\text{Prim}}_n(A) \rightarrow \widetilde{\text{Sei}}_n(A) \rightarrow \widetilde{\text{Bla}}_n(A) \rightarrow \widetilde{\text{Prim}}_{n-1}(A) \rightarrow \dots$$

(ii) This is a special case of Proposition 5.6 (ii). □

This completes the proofs of Theorems D and E of the Introduction.

Remark 5.15 Unfortunately, we do not know if the universal localization $\Sigma^{-1}A[F_\mu]$ is stably flat in general. See Dicks and Sontag [11], Farber and Vogel [16] for proofs that $\Sigma^{-1}A[F_\mu]$ is stably flat when A is a principal ideal domain, and Ara and Dicks [1, Theorem 4.4] when A is a von Neumann regular ring or a commutative Bezout domain.

Remark 5.16 Sheiham [40] computed

$$K_1(\Sigma^{-1}A[F_\mu]) = K_1(A) \oplus \epsilon_\Sigma^{-1}(1)/C$$

with $\epsilon_\Sigma: \Sigma^{-1}A[F_\mu] \rightarrow A$ the factorization of the augmentation map $\epsilon: A[F_\mu] \rightarrow A$ and $C \subseteq \epsilon_\Sigma^{-1}(1)$ the subgroup generated by the commutators

$$(1 + ab)(1 + ba)^{-1} \quad (a, b \in \Sigma^{-1}A[F_\mu], \epsilon(ab) = \epsilon(ba) = 0).$$

It follows from the splitting given by Theorem 5.14 (i)

$$K_1(\Sigma^{-1}A[F_\mu]) = K_1(A) \oplus \widetilde{\text{Sei}}_0(A)$$

that there is defined an isomorphism

$$\widetilde{\text{Sei}}_0(A) \xrightarrow{\cong} \epsilon_\Sigma^{-1}(1)/C; (P, e, \{\pi_i\}) \mapsto D(1 - e + ez: P[F_\mu] \rightarrow P[F_\mu])$$

with D the generalized Dieudonné noncommutative determinant of [40, 4.3].

Example 5.17 (i) The algebraic K -groups of $\mathbb{Z}[F_\mu]$ are such that

$$K_*(\mathbb{Z}[F_\mu]) = K_*(\mathbb{Z}) \oplus \bigoplus_{\mu} K_{*-1}(\mathbb{Z}),$$

$$K_n(\mathbb{Z}[F_\mu]) = K_n(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } n = 0 \\ 0 & \text{if } n \leq -1 \end{cases}$$

by Stallings [44], Gersten [19], Bass [4, XII] and Waldhausen [49; 48], so that

$$\begin{aligned} \text{Flk}(\mathbb{Z}) &= \text{Bla}(\mathbb{Z}), & \widetilde{\text{Prim}}_*(\mathbb{Z}) &= 0, \\ K_{*+1}(\Sigma^{-1}\mathbb{Z}[F_\mu])/K_{*+1}(\mathbb{Z}) &= \widetilde{\text{Sei}}_*(\mathbb{Z}) = \text{Flk}_*(\mathbb{Z}) = \text{Bla}_*(\mathbb{Z}), \\ K_*(H(\mathbb{Z}[F_\mu], \Sigma)) &= \bigoplus_{\mu} K_{*-1}(\mathbb{Z}) \oplus \widetilde{\text{Sei}}_*(\mathbb{Z}), \\ K_n(H(\mathbb{Z}[F_\mu], \Sigma)) &= \widetilde{\text{Sei}}_n(\mathbb{Z}) \quad (n \leq 0). \end{aligned}$$

(ii) Given a μ -component boundary link $\ell: \bigsqcup_{\mu} S^n \subset S^{n+2}$ with exterior W and given a μ -component Seifert surface $V = V_1 \sqcup V_2 \sqcup \dots \sqcup V_{\mu} \subset S^{n+2}$ for ℓ let

$\dot{C}(\tilde{W})$, $(\dot{C}(V), e, \{\pi_i\})$ be the chain complexes defined in [Example 3.13](#). Thus $\dot{C}(\tilde{W})$ is a $\Sigma^{-1}\mathbb{Z}[F_\mu]$ -acyclic $(n+2)$ -dimensional f.g. free $\mathbb{Z}[F_\mu]$ -module chain complex, $(\dot{C}(V), e, \{\pi_i\})$ is an $(n+1)$ -dimensional chain complex in $\mathcal{Sei}(\mathbb{Z})$, and $B(\dot{C}(V), e, \{\pi_i\})$ is an $(n+1)$ -dimensional chain complex in $\mathcal{Flk}(\mathbb{Z})$ with a homology equivalence $\dot{C}(\tilde{W}) \rightarrow B(\dot{C}(V), e, \{\pi_i\})$. The torsion

$$\begin{aligned} \tau(\ell) &= \tau(\Sigma^{-1}\dot{C}(\tilde{W})) \\ &= (\dot{C}(V), e, \{\pi_i\}) = \sum_{r=0}^{n+1} (-)^r (\dot{C}_r(V), e, \{\pi_i\}) = [\dot{C}(\tilde{W})] \\ &\in K_1(\Sigma^{-1}\mathbb{Z}[F_\mu])/K_1(\mathbb{Z}) = K_0(H(\mathbb{Z}[F_\mu], \Sigma)) = \widetilde{\text{Sei}}_0(\mathbb{Z}) = \text{Bla}_0(\mathbb{Z}) \end{aligned}$$

is an isotopy invariant of ℓ , given by Sheiham [\[40\]](#) to be the generalized Dieudonné determinant

$$\tau(\ell) = \sum_{r=0}^{n+1} (-)^r D(1 - e + ez: \dot{C}_r(V)[F_\mu] \rightarrow \dot{C}_r(V)[F_\mu]) \in \widetilde{\text{Sei}}_0(\mathbb{Z}) = \epsilon_{\Sigma}^{-1}(1)/C$$

with $\epsilon_{\Sigma}: \Sigma^{-1}\mathbb{Z}[F_\mu] \rightarrow \mathbb{Z}$ and $C \subseteq \epsilon_{\Sigma}^{-1}(1)$ as recalled in [Remark 5.16](#). The $\mathbb{Z}[F_\mu]$ -modules $\dot{H}_r(\tilde{W})/\mathbb{Z}$ -torsion ($0 \leq r \leq n+1$) are h.d. 1 F_μ -link modules, and

$$\begin{aligned} \tau(\ell) &= \sum_{r=0}^{n+1} (-)^r D(1 - e + ez: \dot{H}_r(V)[F_\mu] \rightarrow \dot{H}_r(V)[F_\mu]) \\ &= \sum_{r=0}^{n+1} (-)^r [\dot{H}_r(\tilde{W})/\mathbb{Z}\text{-torsion}] \\ &\in K_1(\Sigma^{-1}\mathbb{Z}[F_\mu])/K_1(\mathbb{Z}) = K_0(H(\mathbb{Z}[F_\mu], \Sigma)) \\ &= \widetilde{\text{Sei}}_0(\mathbb{Z}) = \text{Bla}_0(\mathbb{Z}) = (\Sigma^{-1}\mathbb{Z}[F_\mu])^\bullet / \{\pm 1\}. \end{aligned}$$

For $\mu = 1$ this is just the Reidemeister torsion of a knot $\ell: S^n \subset S^{n+2}$, which is the alternating product of the Alexander polynomials

$$\begin{aligned} \tau(\ell) &= \sum_{r=0}^{n+1} (-)^r \det(1 - e + ez: \dot{H}_r(V)[z, z^{-1}] \rightarrow \dot{H}_r(V)[z, z^{-1}]) \\ &= \sum_{r=0}^{n+1} (-)^r [\dot{H}_r(\tilde{W})/\mathbb{Z}\text{-torsion}] \\ &\in K_1(\Sigma^{-1}\mathbb{Z}[z, z^{-1}])/K_1(\mathbb{Z}) = K_0(H(\mathbb{Z}[z, z^{-1}], \Sigma)) \\ &= \widetilde{\text{Sei}}_0(\mathbb{Z}) = \text{Bla}_0(\mathbb{Z}) = \widetilde{\text{End}}_0(\mathbb{Z}) = (\Sigma^{-1}\mathbb{Z}[z, z^{-1}])^\bullet / \{\pm 1\} \end{aligned}$$

(Milnor [\[29\]](#), cf [\[33, Example 17.11\]](#)).

(iii) The isotopy classes of simple μ -component boundary links $\ell: \bigsqcup_{\mu} S^{2q-1} \subset S^{2q+1}$ for $q \geq 3$ are in one-one correspondence with the ‘ l -equivalence classes of Seifert matrices’ (Liang [27], generalizing the case $\mu = 1$ due to Levine [25]), and also with the ‘ R -equivalence classes of $(-)^q$ -symmetric isometry structures of multiplicity μ ’ (Farber [15, 4.7]). For simple ℓ $H_q(\widetilde{W})$ is an h.d. 1 F_{μ} -link module, and the torsion

$$\tau(\ell) = (-)^q [H_q(\widetilde{W})] \in \widetilde{\text{Sei}}_0(\mathbb{Z}) = \text{Flk}_0(\mathbb{Z}) = \text{Bla}_0(\mathbb{Z})$$

is just the K -theory part of these complete isotopy invariants for $q \geq 3$.

References

- [1] **P Ara, W Dicks**, *Universal localizations embedded in power-series rings*, Forum Mathematicum (to appear)
- [2] **P Balmer, M Schlichting**, *Idempotent completion of triangulated categories*, J. Algebra 236 (2001) 819–834 [MR1813503](#)
- [3] **H Bass**, *Projective modules over free groups are free*, J. Algebra 1 (1964) 367–373 [MR0178032](#)
- [4] **H Bass**, *Algebraic K-theory*, W A Benjamin (1968) [MR0249491](#) Downloadable from: <http://www.math.uni-bielefeld.de/~rehmann/DML/>
- [5] **H Bass, A Heller, R G Swan**, *The Whitehead group of a polynomial extension*, Inst. Hautes Études Sci. Publ. Math. (1964) 61–79 [MR0174605](#)
- [6] **A J Berrick, M E Keating**, *The K-theory of triangular matrix rings*, from: “Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983)”, Contemp. Math. 55, Amer. Math. Soc., Providence, RI (1986) 69–74 [MR862629](#)
- [7] **F Borceux**, *Handbook of categorical algebra. 1*, Encyclopedia of Mathematics and its Applications 50, Cambridge University Press, Cambridge (1994) [MR1291599](#)
- [8] **S E Cappell, J L Shaneson**, *The codimension two placement problem and homology equivalent manifolds*, Ann. of Math. (2) 99 (1974) 277–348 [MR0339216](#)
- [9] **S E Cappell, J L Shaneson**, *Link cobordism*, Comment. Math. Helv. 55 (1980) 20–49 [MR569244](#)
- [10] **P M Cohn**, *Free rings and their relations*, Academic Press (1971) [MR0371938](#)
- [11] **W Dicks, E D Sontag**, *Sylvester domains*, J. Pure Appl. Algebra 13 (1978) 243–275 [MR509164](#)
- [12] **J Duval**, *Forme de Blanchfield et cobordisme d’entrelacs bords*, Comment. Math. Helv. 61 (1986) 617–635 [MR870709](#)

- [13] **M Farber**, *Hermitian forms on link modules*, Comment. Math. Helv. 66 (1991) 189–236 [MR1107839](#)
- [14] **M Farber**, *Noncommutative rational functions and boundary links*, Math. Ann. 293 (1992) 543–568 [MR1170526](#)
- [15] **M Farber**, *Stable-homotopy and homology invariants of boundary links*, Trans. Amer. Math. Soc. 334 (1992) 455–477 [MR1079054](#)
- [16] **M Farber**, **P Vogel**, *The Cohn localization of the free group ring*, Math. Proc. Cambridge Philos. Soc. 111 (1992) 433–443 [MR1151322](#)
- [17] **P Gabriel**, *Des catégories abéliennes*, Bull. Soc. Math. France 90 (1962) 323–448 [MR0232821](#)
- [18] **P Gabriel**, **M Zisman**, *Calculus of fractions and homotopy theory*, Ergebnisse series 35, Springer, New York (1967) [MR0210125](#)
- [19] **S M Gersten**, *On class groups of free products*, Ann. of Math. (2) 87 (1968) 392–398 [MR0224655](#)
- [20] **H Gillet**, *Riemann–Roch theorems for higher algebraic K -theory*, Adv. in Math. 40 (1981) 203–289 [MR624666](#)
- [21] **D R Grayson**, *Exact sequences in algebraic K -theory*, Illinois J. Math. 31 (1987) 598–617 [MR909785](#)
- [22] **M A Gutiérrez**, *Boundary links and an unlinking theorem*, Trans. Amer. Math. Soc. 171 (1972) 491–499 [MR0310902](#)
- [23] **A Haghany**, **K Varadarajan**, *Study of modules over formal triangular matrix rings*, J. Pure Appl. Algebra 147 (2000) 41–58 [MR1744654](#)
- [24] **B Keller**, *Derived categories and their uses*, from: “Handbook of algebra, Vol. 1”, North-Holland, Amsterdam (1996) 671–701 [MR1421815](#)
- [25] **J Levine**, *An algebraic classification of some knots of codimension two*, Comment. Math. Helv. 45 (1970) 185–198 [MR0266226](#)
- [26] **J Levine**, *Knot modules I*, Trans. Amer. Math. Soc. 229 (1977) 1–50 [MR0461518](#)
- [27] **C C Liang**, *An algebraic classification of some links of codimension two*, Proc. Amer. Math. Soc. 67 (1977) 147–151 [MR0458439](#)
- [28] **W Lück**, **A Ranicki**, *Chain homotopy projections*, J. Algebra 120 (1989) 361–391 [MR989903](#)
- [29] **J W Milnor**, *Infinite cyclic coverings*, from: “Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, 1967)”, Prindle, Weber & Schmidt, Boston (1968) 115–133 [MR0242163](#)
- [30] **A Neeman**, **A Ranicki**, *Noncommutative localisation and chain complexes I: Algebraic K - and L -theory* [arXiv:math.RA/0109118](#)

- [31] **A Neeman, A Ranicki**, *Noncommutative localisation in algebraic K -theory I*, *Geom. Topol.* 8 (2004) 1385–1425 [MR2119300](#)
- [32] **D Quillen**, *Higher algebraic K -theory I*, from: “Algebraic K -theory I: Higher K -theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)”, Springer, Berlin (1973) 85–147. *Lecture Notes in Math.*, Vol. 341 [MR0338129](#)
- [33] **A Ranicki**, *High-dimensional knot theory*, Springer Monographs in Mathematics, Springer, New York (1998) [MR1713074](#)
- [34] **A Ranicki**, *Blanchfield and Seifert algebra in high-dimensional knot theory*, *Mosc. Math. J.* 3 (2003) 1333–1367 [MR2058802](#)
- [35] **A Ranicki**, *Algebraic and combinatorial codimension-1 transversality*, from: “Proceedings of the Casson Fest”, (C Gordon, R Y, editors), *Geom. Topol. Monogr.* 7 (2004) 145–180 [MR2172482](#)
- [36] **A Ranicki**, *Noncommutative localization in topology*, from: “Non-commutative localization in algebra and topology”, *London Math. Soc. Lecture Note Ser.* 330, Cambridge Univ. Press, Cambridge (2006) 81–102 [MR2222483](#)
- [37] **N Sato**, *Free coverings and modules of boundary links*, *Trans. Amer. Math. Soc.* 264 (1981) 499–505 [MR603777](#)
- [38] **M Schlichting**, *Negative K -theory of derived categories*, *Math. Z.* 253 (2006) 97–134 [MR2206639](#)
- [39] **A H Schofield**, *Representation of rings over skew fields*, *London Mathematical Society Lecture Note Series* 92, Cambridge University Press, Cambridge (1985) [MR800853](#)
- [40] **D Sheiham**, *Whitehead groups of localizations and the endomorphism class group*, *J. Algebra* 270 (2003) 261–280 [MR2016661](#)
- [41] **D Sheiham**, *Invariants of boundary link cobordism II: The Blanchfield–Duval form*, from: “Non-commutative localization in algebra and topology (Edinburgh 2002)”, *LMS Lecture Note Ser.* 330, Cambridge Univ. Press, Cambridge (2006) 143–219 [MR2222485](#)
- [42] **D Sheiham**, *Universal localization of triangular matrix rings*, *Proc. Amer. Math. Soc.* 134 (2006) 3465–3474
- [43] **N Smythe**, *Boundary links*, from: “Wisconsin Topology Seminar”, *Ann. of Maths. Studies* 60, Princeton UP (1966) 69–72
- [44] **J Stallings**, *Whitehead torsion of free products*, *Ann. of Math.* (2) 82 (1965) 354–363 [MR0179270](#)
- [45] **R W Thomason, T Trobaugh**, *Higher algebraic K -theory of schemes and of derived categories*, from: “The Grothendieck Festschrift, Vol. III”, *Progr. Math.* 88, Birkhäuser, Boston, MA (1990) 247–435 [MR1106918](#)

- [46] **H F Trotter**, *Knot modules and Seifert matrices*, from: “Knot theory (Proc. Sem., Plans-sur-Bex, 1977)”, Lecture Notes in Math. 685, Springer, Berlin (1978) 291–299 [MR521739](#)
- [47] **F Waldhausen**, *Whitehead groups of generalized free products*, from: “Algebraic K -theory II: “Classical” algebraic K -theory and connections with arithmetic (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)”, Springer, Berlin (1973) 155–179. Lecture Notes in Math., Vol. 342 [MR0370576](#)
- [48] **F Waldhausen**, *Algebraic K -theory of generalized free products, Part 1*, Ann. of Math. (2) 108 (1978) 135–204 [MR0498807](#)
- [49] **F Waldhausen**, *Algebraic K -theory of generalized free products, Part 2*, Ann. of Math. (2) 108 (1978) 205–256 [MR0498808](#)
- [50] **F Waldhausen**, *Algebraic K -theory of spaces*, from: “Algebraic and geometric topology (New Brunswick, N.J., 1983)”, Lecture Notes in Math. 1126, Springer, Berlin (1985) 318–419 [MR802796](#)

School of Mathematics, University of Edinburgh
Edinburgh EH9 3JZ, Scotland, UK

a.ranicki@ed.ac.uk

Proposed: Wolfgang Lück
Seconded: Peter Teichner, Steve Ferry

Received: 6 October 2005
Revised: 14 July 2006