

Erratum for the paper ‘On the chain-level intersection pairing for PL manifolds’

J E MCCLURE

Greg Friedman has pointed out that there are sign errors in [3], and in particular Lemma 10.5(b) (which is a key step in the proof of the main theorem) is not correct as stated.

The purpose of this note is to provide a correction.

[18D50](#); [57Q65](#)

In Section 5 of [3], the umkehr map

$$H_p(A, B) \rightarrow H_{p+n-m}(A', B')$$

should have a sign $(-1)^{(m-p)(n-m)}$ as in [1, pages 314–315] (see [2] for an explanation of where this sign comes from). Also, it’s convenient to let the symbol $f_!$ stand for the desuspension

$$\Sigma^{-m} H_*(A, B) \rightarrow \Sigma^{-n} H_*(A', B').$$

Note that this map preserves degrees.

With these changes, Lemma 5.1 says that the diagram

$$\begin{array}{ccc} \Sigma^{-m} H_*(A, B) & \xrightarrow{f_!} & \Sigma^{-n} H_*(A', B') \\ \partial \downarrow & & \downarrow \partial \\ \Sigma^{-m} H_*(B) & \xrightarrow{f_!} & \Sigma^{-n} H_*(B') \end{array}$$

commutes.

In Section 8, observe that if C_* and D_* are chain complexes and $m, n \in \mathbb{Z}$ then $(\Sigma^{-m} C_*) \otimes (\Sigma^{-n} D_*)$ and $\Sigma^{-(m+n)}(C_* \otimes D_*)$ will be *different* chain complexes when n is odd: they are the same as graded abelian groups but have different differentials. However, there is an isomorphism

$$\Theta : (\Sigma^{-m} C_*) \otimes (\Sigma^{-n} D_*) \rightarrow \Sigma^{-(m+n)}(C_* \otimes D_*)$$

which takes $\Sigma^{-m} x \otimes \Sigma^{-n} y$ to $(-1)^{n|x|} \Sigma^{-(m+n)}(x \otimes y)$, and similarly for any number of tensor factors.

Given manifolds M_1 and M_2 of dimensions m_1, m_2 , define

$$\bar{\varepsilon} : (\Sigma^{-m_1} C_* M_1) \otimes (\Sigma^{-m_2} C_* M_2) \rightarrow \Sigma^{-(m_1+m_2)} C_*(M_1 \times M_2)$$

to be

$$(-1)^{m_1 m_2} (\Sigma^{-(m_1+m_2)} \varepsilon) \circ \Theta.$$

The motivation for this is that $\bar{\varepsilon}$ is Poincaré dual to the exterior product in cohomology (see [2] for details). Similarly, given M_1, \dots, M_k define

$$\bar{\varepsilon} : \bigotimes \Sigma^{-m_i} C_* M_i \rightarrow \Sigma^{-\sum m_i} C_*(\prod M_i)$$

to be

$$(-1)^{e_2(m_1, \dots, m_k)} (\Sigma^{-\sum m_i} \varepsilon) \circ \Theta$$

where e_2 is the second elementary symmetric function (so that, for example, $e_2(m_1, m_2, m_3) = m_1 m_2 + m_1 m_3 + m_2 m_3$).

Now define

$$G_2 \subset (\Sigma^{-m} C_* M) \otimes (\Sigma^{-m} C_* M)$$

to be $\bar{\varepsilon}^{-1}(\Sigma^{-2m} C_*^\Delta(M \times M))$ and define μ_2 to be $\Delta_! \circ \bar{\varepsilon}$.

With these changes, Remark 8.1 becomes correct (it wasn't before); see [2] for the proof.

In Section 10, Definition 10.1 should be restated: define G_k to be the subcomplex of $(\Sigma^{-m} C_* M)^{\otimes k}$ consisting of elements x for which both $\Sigma^{mk} \bar{\varepsilon}_k(x)$ and $\Sigma^{mk} \bar{\varepsilon}_k(\partial x)$ are in general position with respect to all generalized diagonal maps.

The diagram in Lemma 10.5(b) should be replaced by

$$\begin{array}{ccc} \Sigma^{-m_1} C_*^f M_1 \otimes \Sigma^{-m_2} C_*^f M_2 & \xrightarrow{\bar{\varepsilon}} & \Sigma^{-(m_1+m_2)} C_*^{f \times g}(M_1 \times M_2) \\ f_! \otimes g_! \downarrow & & \downarrow (f \times g)_! \\ \Sigma^{-n_1} C_*^f N_1 \otimes \Sigma^{-n_2} C_*^f N_2 & \xrightarrow{\bar{\varepsilon}} & \Sigma^{-(n_1+n_2)} C_*^{f \times g}(N_1 \times N_2) \end{array}$$

At the end of Section 10, the definition of G_R should be

$$\bar{\varepsilon}_{k'}^{-1} \circ (R^*)_! \circ \bar{\varepsilon}_k.$$

In Section 11, Lemma 11.1 should say that the inclusion

$$G_{k+l} \hookrightarrow (\Sigma^{-m} C_* M)^{k+l} \cong (\Sigma^{-m} C_* M)^k \otimes (\Sigma^{-m} C_* M)^l$$

has its image in $G_k \otimes G_l$. Now define

$$\xi_{k,l} : G_{k+l} \rightarrow G_k \otimes G_l$$

to be the inclusion provided by Lemma 11.1.

References

- [1] **A Dold**, *Lectures on algebraic topology*, Grundlehrenseries 200, Springer, New York (1972) [MR0415602](#)
- [2] **G Friedman**, *On the chain-level intersection pairing for PL manifolds*, to appear [arXiv:0808.1749](#)
- [3] **J E McClure**, *On the chain-level intersection pairing for PL manifolds*, *Geom. Topol.* 10 (2006) 1391–1424 [MR2255502](#)

*Department of Mathematics, Purdue University
150 N University Street, West Lafayette, IN 47907-2067, USA*

mcclure@math.purdue.edu

Received: 18 July 2007