

Width is not additive

RYAN BLAIR
MAGGY TOMOVA

We develop the construction suggested by Scharlemann and Thompson in [15] to obtain an infinite family of pairs of knots K_α and K'_α so that $w(K_\alpha \# K'_\alpha) = \max\{w(K_\alpha), w(K'_\alpha)\}$. This is the first known example of a pair of knots such that $w(K \# K') < w(K) + w(K') - 2$ and it establishes that the lower bound $w(K \# K') \geq \max\{w(K), w(K')\}$ obtained in Scharlemann and Schultens [14] is best possible. Furthermore, the knots K_α provide an example of knots where the number of critical points for the knot in thin position is greater than the number of critical points for the knot in bridge position.

57M25, 57M27, 57M50

1 Introduction

Thin position for knots was first defined by Gabai [5] in his proof of property R . The idea of width has had important applications in 3–manifold topology. In particular, width played an integral role in three celebrated results: the solution to the knot complement problem by Gordon and Luecke [7], the leveling of unknotting tunnels by Goda, Scharlemann and Thompson [6] and Thompson’s solution [19] to the recognition problem for S^3 , which was originally achieved by Rubinstein [12]. However, surprisingly little is known about its intrinsic properties. Most strikingly, the behavior of knot width under connected sums has remained one of the most interesting and difficult problems to elucidate. In an attempt to shed light on this question, width has been compared to bridge number: the least number of maxima over all projections of the knot. Just like bridge number, width depends on the number of critical points of a projection, but it also takes into account their relative heights. The behavior of bridge number under connected sum was first established by Schubert [16]. Later, Schultens [18] gave a considerably more elegant proof of the result. Stacking the two knots vertically and connecting a minimum of the top one to a maximum of the bottom one shows that

$$(1) \quad b(K \# K') \leq b(K) + b(K') - 1.$$

Schubert's result affirms that inequality (1) is in fact an equality.

This construction also gives an easy inequality for the width of a connected sum, namely

$$(2) \quad w(K \# K') \leq w(K) + w(K') - 2.$$

In general, it is unknown for which knots K and K' the inequality (2) is also an equality. Partial results and special cases have been solved. Most notably, Scharlemann and Schultens showed in [14] that

$$(3) \quad w(K \# K') \geq \max\{w(K), w(K')\}$$

and the work of Rieck and Sedgwick in [11] implies that the equality $w(K \# K') = w(K) + w(K') - 2$ holds for meridionally small knots. The main result in this paper is that inequality (2) is strict for some knots and that inequality (3) is best possible if no restrictions are placed on the knots.

Theorem 1.1 *There exists an infinite family of knots K_α and K'_α with $w(K_\alpha \# K'_\alpha) = \max\{w(K_\alpha), w(K'_\alpha)\}$.*

Moreover, the construction also yields examples of another interesting phenomenon.

Theorem 1.2 *There exists an infinite family of knots K_α so that the minimal bridge position for K_α has fewer critical points than any thin position of K_α .*

Our construction is based on ideas proposed by Scharlemann and Thompson in [15]. In their paper, the authors give a large family of pairs of knots K_α and K'_α for which it appears that $w(K_\alpha \# K'_\alpha) = \max\{w(K_\alpha), w(K'_\alpha)\}$. This equality holds if the projections of K_α and K'_α considered by Scharlemann and Thompson have minimal width amongst all possible projections of the knots, ie that K_α and K'_α are in thin position. One of the knots in each pair is quite simple and its width is easily established. However, the authors could not verify the width of the second knot in any of their pairs. In [2], the current authors established that, for most of these pairs, the second knot is not in thin position. Therefore, most of these pairs do not provide the desired counterexample to the conjectured equality $w(K \# K') = w(K) + w(K') - 2$.

In this paper, we construct a family of pairs of knots K_α and K'_α that satisfy the properties required for the pairs presented in [15] and we establish that both K_α and K'_α are in thin position. For such a pair, it follows that $w(K_\alpha \# K'_\alpha) = \max\{w(K_\alpha), w(K'_\alpha)\}$. Figure 1 depicts such a pair where one knot is the trefoil. The figure demonstrates a projection of $K \# \text{trefoil}$ that has the same width as the given projection of K . We will show that this projection of K is of minimal width.




Figure 1

The paper is organized as follows. After some preliminary definitions in Section 3, we show that if a knot is in thin position and P is a thin level sphere, then cut-disks for P that do not intersect any other thin spheres can be isotoped to be vertical. This allows us to use the results about vertical cut-disks developed by the second named author in [22]. In Section 4, we review results found in Johnson and Tomova [9] about the behavior of a second bridge surface for a tangle that has a high distance bridge sphere. In Section 5, we use a theorem of Schubert to construct tangles with high distance properties so that the numerator closures of certain subtangles yield nontrivial knots. In Section 6, we construct a three strand tangle for which we can classify all essential meridional surfaces of Euler characteristic greater than -12 . In Section 7, we construct the candidate knots K_α . Each of these knots has the general schematic introduced in Scharlemann and Thompson [15]; see Figure 1. In Section 8, we establish some of the properties of the knots. The proofs of these properties depend on the results in the previous sections. In Section 9, we classify all essential meridional spheres in the complement of K_α that have fewer than 14 punctures. In Section 10, we determine that bridge position and thin position for K_α do not coincide. In Section 11, we introduce additional constraints on the thin and thick levels of a width minimizing embedding of K_α . Finally, in Section 12, we show that the projection given in Figure 1 of each of the knots K_α is thin by checking the widths of a relatively small number of possible thin positions.

Acknowledgement We would like to thank Tsuyoshi Kobayashi, Yo'av Rieck and Martin Scharlemann for many helpful conversations.

Our research was partially supported by NSF grant number 1054450.

2 Definitions and preliminaries

Let K be a knot embedded in S^3 . We will denote a regular neighborhood of K by $\eta(K)$. Unless otherwise stated, all surfaces under consideration are connected. An *essential meridional surface* in the knot complement is a surface with nonempty, meridional boundary that does not have any compressing disks in the knot complement and that is not boundary parallel in $S^3 - \eta(K)$. Let F be a meridional surface embedded in the complement of K . A *cut-disk* for F is a disk $D^c \subset S^3$ such that $D^c \cap F = \partial D^c$, $|D^c \cap K| = 1$ and the annulus $D^c - \eta(K)$ is not parallel in the knot complement to a subset of $F - \eta(K)$ via an isotopy during which the boundary component of $D^c - \eta(K)$ on the boundary of the knot may move, but has to stay on the boundary of the knot. In particular, if K is prime, the curve ∂D^c is not parallel to a boundary component of $F - \eta(K)$. We use the term *c-disk* to refer to either a compressing disk or a cut-disk.

In the following we will often consider height functions on S^3 , B^3 and $S^2 \times I$. In each of these cases, we only consider height functions with a minimal number of nondegenerate critical points. Let $h: S^3 \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ be the standard height function on S^3 and suppose K is in general position with respect to h . If t is a regular value of $h|_K$, $h^{-1}(t)$ is called a *level sphere* with width $w(h^{-1}(t)) = |K \cap h^{-1}(t)|$. If $c_0 < c_1 < \dots < c_n$ are all the critical values of $h|_K$, choose regular values r_1, r_2, \dots, r_n such that $c_{i-1} < r_i < c_i$. Then the *width of K with respect to h* is defined by $w(K, h) = \sum w(h^{-1}(r_i))$. The *width of K* , $w(K)$, is the minimum of $w(K', h)$ over all knots K' isotopic to K . We say that K is in *thin position* if $w(K, h) = w(K)$. Note that by removing a neighborhood of the north and south pole, we can assume $K \subset S^2 \times I$ and define width there. We will switch between these two ambient spaces freely during this discussion. More details about thin position and basic results can be found in Scharlemann [13].

A level sphere $h^{-1}(t)$ is called *thin* if the highest critical point for K below it is a maximum and the lowest critical point above it is a minimum. If the highest critical point for K below $h^{-1}(t)$ is a minimum and the lowest critical point above it is a maximum, the level sphere is called *thick*. As the lowest critical point of K is a minimum and the highest is a maximum, a thick level sphere can always be found. It is possible that the knot does not have any thin spheres with respect to some height function. When this occurs the unique thick sphere is called a *bridge sphere* and the knot is said to be in *bridge position*.

We will use the following result found in Scharlemann and Schultens [14] to simplify our computations.

Lemma 2.1 [14, Lemma 6.2] *Let K be an embedding of a knot in S^3 and let $h: S^3 \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ be the standard height function on S^3 . If $\{a_i\}$, $i = 0, \dots, n$, and $\{b_j\}$, $j = 0, \dots, n + 1$, are the widths of all thin and all thick spheres respectively, then*

$$w(K) = \frac{\sum_{j=0}^{n+1} b_j^2 - \sum_{i=0}^n a_i^2}{2}.$$

Unless otherwise stated, we will always consider all level spheres to lie in the knot complement, ie they are always meridional surfaces. A key ingredient to our proofs is the behavior of c -disks for the thin level spheres. We review some already known results here and then develop some new results in the next section.

Theorem 2.2 [23] *Suppose K is a prime knot in thin position and let P be the thin sphere of lowest width. Then P is incompressible.*

Theorem 2.3 [20, Theorem 8.1] *Suppose K is a prime knot in thin position and P is a minimal width thin sphere. If P' is a thin sphere so that $w(P') = w(P) + 2$, then P' is incompressible.*

In many of our proofs we will consider various 3-balls containing arcs of the knot. Therefore we need the following definition.

Definition 2.4 *A tangle, \mathcal{R} , is a tuple, (B_R, R) , where B_R is a 3-ball or $S^2 \times I$ and R is a properly embedded compact 1-manifold in B_R . \mathcal{R} is an n -strand tangle if R contains no loops and exactly n arcs.*

Given any tangle \mathcal{T} in a ball or in $S^2 \times I$ and a height function h , the critical points of the tangle can be organized into braid boxes as follows: suppose t_1 and t_2 are adjacent thin levels. Then a *braid box* $B_{[t_1, t_2]} \subset h^{-1}[t_1, t_2]$ is a ball containing $T \cap h^{-1}[t_1, t_2]$. In this ball, the tangle has some number of minima of T followed by some number of maxima of T . Given a tangle \mathcal{S} in $S^2 \times I$ and a c -disk D^c for $S^2 \times \{0\}$, then D^c naturally decomposes \mathcal{S} into two subtangles \mathcal{S}_α and \mathcal{S}_β . These subtangles can be decomposed into braid boxes $\{B_{[a_i^-, a_i^+]}\}$ and $\{B_{[b_j^-, b_j^+]}\}$. A *horizontal isotopy* is an isotopy that preserves the height of each point. For more details see Tomova [22].

Theorem 2.5 [22, Lemma 9.1] *Let $\mathcal{L} = (B_L, L)$ be a prime tangle where B_L is homeomorphic to $S^2 \times I$, let P be a level sphere for \mathcal{L} and let D^* be a c -disk for P that does not have any saddles with respect to the usual height function on $S^2 \times I$. Then there exists a horizontal isotopy ν which keeps D^* fixed such that if $\{B_{[a_i^-, a_i^+]}\}$ and $\{B_{[b_j^-, b_j^+]}\}$ are the collections of braid boxes for the tangles \mathcal{S}_α and \mathcal{S}_β respectively, for any i and j the intervals $[a_i^-, a_i^+]$ and $[b_j^-, b_j^+]$ are disjoint.*

Remark 2.6 As ν is a horizontal isotopy it does not change the total number or the heights of the thin spheres for L .

Corollary 2.7 Let L be a prime knot embedded in $S^2 \times I$, let P be a thin sphere for L , let D^* be a c -disk for P above it that intersects each level sphere in at most one simple closed curve and let P' be the thin sphere directly above D^* . Then either there are some thin spheres between P and P' or all critical points for L between P and P' are on the same side of D^* . In particular, in the latter case c -compressing P along D^* results in a component parallel to P' .

Proof Suppose P and P' are adjacent thin spheres and suppose L has critical points on both sides of D^* . By Theorem 2.5, we may assume that the braid boxes for the two sides are disjoint. However, a level sphere that is disjoint from all braid boxes is necessarily thin and therefore P and P' are not adjacent. \square

3 Vertical cut-disks

Let K be a knot in S^3 and let $h: S^3 \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ be the standard height function on S^3 . Suppose that K is in thin position with respect to h . Let $P = h^{-1}(r)$ be a level sphere and suppose P has a c -disk, C , that lies above it.

We first introduce some notation and definitions. Figure 3 illustrates all of the terminology outlined below. Let F_C be the singular foliation on the c -disk C induced by the level sets of $h|_C$. Although conventionally saddles are points, in this paper a *saddle* of a surface F is any leaf of the foliation induced by level sets of $h|_F$ that is homeomorphic to the wedge of two circles. By standard position, we can assume that all saddles of F_C are disjoint from K .

Given a saddle $\sigma = s_1^\sigma \vee s_2^\sigma$ in a level sphere $S_\sigma = (h^{-1} \circ h)(\sigma)$, let D_1^σ be the closure of the component of $S_\sigma - s_1^\sigma$ that is disjoint from s_2^σ and D_2^σ be the closure of the component of $S_\sigma - s_2^\sigma$ that is disjoint from s_1^σ .

A subdisk D in F_C is *monotone* if its boundary is entirely contained in a leaf of F_C and the interior of D is disjoint from every saddle in F_C . In practice, we will use the term subdisk in a slightly broader sense, allowing ∂D to be immersed in C , where if ∂D is immersed, then ∂D is a saddle. We say a monotone disk is *outermost* if its boundary is s_i^σ for some saddle σ and label the disk D_σ . Similarly, if some s_i^σ bounds an outermost disk D_σ , we say σ is an outermost saddle. It will usually be the case that at most one of s_1^σ and s_2^σ is the boundary of an outermost disk, so, our convention is to relabel so that $\partial D_\sigma = s_1^\sigma$.

Suppose σ is an outermost saddle. The level sphere S_σ cuts S^3 into two 3–balls. The ball that contains D_σ is again cut by D_σ into two 3–balls B_σ and B'_σ . We choose the labeling of B_σ and B'_σ so that $\partial B_\sigma = D_1^\sigma \cup D_\sigma$.

We say σ is an *inessential saddle* if σ is an outermost saddle and D_σ is disjoint from K . An n –*punctured disk* denotes a disk embedded in S^3 that meets K transversely in exactly n points. An embedded simple closed curve in a c –disk C is c –*inessential* if it bounds a 1–punctured disk in C . Similarly, σ is a c –*inessential saddle* if σ is an outermost saddle and D_σ meets K exactly once. We say σ is a *removable saddle* if σ is an outermost saddle where D_σ has a unique maximum (minimum) and $h|_{K \cap B_\sigma}$ has a maximum (minimum) at every point of $K \cap D_\sigma$. See Figure 2.




Figure 2

We say a saddle σ in F_C is *standard* if there is a monotone disk E_σ such that $\partial(E_\sigma) = \sigma$. If σ is a standard saddle, A_σ is the 3–ball with boundary $E_\sigma \cup D_1^\sigma \cup D_2^\sigma$ such that $A_\sigma \cap S_\sigma = D_1^\sigma \cup D_2^\sigma$.

By general position arguments, we can assume every saddle σ in F_C has a bicollared neighborhood in C that is disjoint from K and all other singular leaves of F_C . The boundary of this bicollared neighborhood consists of three circles c_1^σ , c_2^σ , and c_3^σ , where c_1^σ and c_2^σ are parallel to s_1^σ and s_2^σ respectively. We can assume c_1^σ , c_2^σ , and c_3^σ are level with respect to h and that c_1^σ and c_2^σ lie in the same level surface. The terminology for this section is summarized in Figure 3.

Definition 3.1 A c –disk for a level sphere is *vertical* if it does not have any saddles with respect to h , ie if it intersects each level sphere in at most one simple closed curve.

In [23], Wu shows that any compressing disk for a level sphere is isotopic to a vertical compressing disk by an isotopy that does not change the width of the knot. This is generally not true of cut-disks, but we will now show that under certain conditions cut-disks can also be isotoped to be vertical. Many of the arguments in this section are extensions of techniques developed by Schultens in [18] and extended by Blair in [1]. We will need the following definitions.




Figure 3

Definition 3.2 A c -disk, C , for a level surface P is taut with respect to h if the number of saddles in F_C is minimal subject to the condition that K is a minimal width embedding and P is a level surface.

Definition 3.3 Following Heath and Kobayashi [8], a sphere P in S^3 is called *bowl-like* with respect to a height function h if it can be decomposed into two disks, E_1 and E_2 , glued along their boundary such that E_1 is contained in a level surface for h and E_2 is a monotone disk disjoint from K .

Two thin spheres P and P' are *adjacent* if there is no third thin sphere P'' so that P and P' are contained in distinct complementary components of P'' .

Lemma 3.4 Assume P and P' are adjacent thin spheres with P' above P and C is a cut-disk for P above it but disjoint from P' . We allow the special case where P is the highest thin sphere and P' is a level sphere above it disjoint from K . If F_C contains an inessential saddle, then C is not taut.

Proof Suppose σ is an inessential saddle in F_C .

Case 1 Suppose D_σ contains a unique minimum. Call this point a .

Case 1a Additionally, suppose B_σ does not contain $-\infty$. Use the “Pop out Lemma” [18, Lemma 2] to eliminate σ ; see Figure 4. This isotopy fixes all of S^3 below D_σ and above P' . However, D_σ is contained strictly above the level sphere P . Hence, this isotopy eliminates σ while fixing K below P and above P' and not creating any new critical points for $h|_K$. Since all maxima of $h|_K$ are above all minima of $h|_K$ in the region between P and P' , altering the relative heights of the critical points

without creating any new critical points can only decrease the width of K . Since we have decreased the number of saddles of C without increasing the width of K , then C is not taut.




Figure 4




Figure 5

Case 1b Additionally, suppose B_σ contains $-\infty$. We will describe a sequence of isotopies that allows us to decrease the number of saddles; see Figure 6.

Let α be a monotone arc with endpoints a and $-\infty$ that misses K and intersects C only at local minima. Label the points of $\alpha \cap C$ in order of decreasing height with $a = a_1, \dots, a_n$. Since C lies above P , α meets P in a single point b where $h(a_n) > h(b)$. See Figure 5. Again by general position, we can assume none of the a_i or b lie on K . The following isotopy is a modification of the isotopy presented in [18, Lemma 1].

Let S_- be a level sphere contained in a small neighborhood of $-\infty$ such that S_- does not meet K or P . Let α_b be a subarc of α with endpoints b and $-\infty$. Enlarge α_b slightly to be a vertical solid cylinder V such that ∂V consists of a small neighborhood

of b in P , a small disk in S_- and a vertical annulus, A . Replacing P with the isotopic surface $(P - V) \cup A \cup (S_- - V)$ represents an isotopy of P in $S^3 - K$ that fixes K and results in P being bowl-like.

Toward the goal of isotopying C to be disjoint from α , perform the following isotopy. Let α_n be a subarc of α with endpoints a_n and $-\infty$. Enlarge α_n slightly to be a vertical solid cylinder V such that ∂V consists of a small neighborhood of a_n in C , a small disk in S_- and a vertical annulus, A . Replacing C with the cut-disk $(C - V) \cup A \cup (S_- - V)$ represents an isotopy of C in $S^3 - K$ that fixes K , does not change the number of saddles of F_C and preserves P as bowl-like. The current arrangement is represented in the third illustration in Figure 6.

By induction on n , we can assume α is disjoint from C and P except at the point a . By isotopying D_σ to a new disk D_σ^* in the manner described above, we have enlarged B'_σ to contain $-\infty$ and shrunk B_σ so that it is disjoint from $-\infty$. After a small tilt so that h again restricts to a Morse function on D_σ^* , $F_{D_\sigma^*}$ is a collection of circles and one minimum. The resulting cut-disk C^* is isotopic to C via an isotopy that leaves σ and K fixed and does not change the number of saddles of F_C .

By the ‘‘Pop out Lemma’’ [18, Lemma 2], we can eliminate σ without introducing any new maxima to h_K or new saddles to F_C and while preserving P as bowl-like. The current arrangement is represented in the fourth illustration in Figure 6.

Since P is now bowl-like, it can be decomposed into two disks E_1 and E_2 as in the definition of bowl-like. Let a be the unique minimum on E_2 . Again choose a monotone arc α with endpoints a and $-\infty$ that misses K and intersects C only at local minima. The arc α is disjoint from P except at a . Label the points of $\alpha \cap C$ in order of decreasing height with a_1, \dots, a_n . Again by general position, we can assume none of the a_i lie in K . Repeat the above argument to produce an isotopic copy of C with the same number of saddles that is disjoint from α . The current arrangement is represented in the fifth illustration in Figure 6.

Horizontally shrink and vertically lower P until it is strictly below all of C . Let S_- be 2-sphere boundary of a regular neighborhood of $-\infty$ so that S_- is disjoint from K , P and C . After lowering P into the neighborhood of $-\infty$ and expanding P to fill the neighborhood, we have isotoped P to S_- while preserving the width of K below P and above P' . The current arrangement is represented in the sixth illustration in Figure 6.

Since we have produced an isotopy that decreases the number of saddles of F_C while not introducing any new maxima to $h|_K$ and fixing K below P and above P' , then C is not taut.

Case 2 Suppose D_σ contains a unique maximum. The argument is symmetric to the one in Case 1 above. If necessary, isotope P' to be bowl-like to guarantee that B_σ does not contain $+\infty$, then apply the ‘‘Pop out Lemma’’ to reduce the number of saddles for C . Finally, restore P' to be level. As in Case 1, these isotopies do not affect the width of K below P and above P' and do not introduce new critical points for K , so they do not increase the width of the knot. Since we have decreased the number of saddles of C without increasing the width of K , C is not taut. \square




Figure 6

Corollary 3.5 Assume P and P' are adjacent thin spheres with P' above P and C is a cut-disk for P above it but disjoint from P' . If F_C contains a nonstandard saddle σ , then C is not taut.

Proof Suppose F_C contains a nonstandard saddle σ . By definition of nonstandard, c_3^σ does not bound a monotone disk in C . Since c_1^σ , c_2^σ and c_3^σ are the boundary components of an embedded pair of pants in C , then two of these curves, assume first c_1^σ and c_3^σ , bound, possibly punctured, disks in C denoted E_1^* and E_3^* respectively

such that both E_1^* and E_3^* are disjoint from $c_1^\sigma \cup c_2^\sigma \cup c_3^\sigma$ in their interior. Either σ is outermost or a saddle in $F_{E_1^*}$ is outermost. By hypothesis, E_3^* contains a saddle. Hence, E_3^* contains an outermost saddle. Since F_C contains two outermost saddles and C meets K exactly once, then one of these outermost saddles is inessential. By Lemma 3.4, C is not taut. A similar argument establishes that result if c_1^σ and c_2^σ or c_2^σ and c_3^σ bound disks in C . \square

Given a cut-disk C for a level sphere P we require a notion of *nestedness* that we develop here. Without loss of generality, suppose the C lies above P . C decomposes the 3–ball above P into two 3–balls B_1 and B_2 . Each saddle of C is nested with respect to B_1 or is nested with respect to B_2 , but not both. We determine a saddle's nestedness in the following way. Let σ be a saddle in F_C and Q be the level sphere either just above or just below σ that contains c_1^σ and c_2^σ . The surface $Q - (c_1^\sigma \cup c_2^\sigma)$ is composed of two disks and an annulus, A . If a collar of ∂A in A is contained in B_1 , then we say σ is *unnested* with respect to B_1 . If not, we say σ is *nested* with respect to B_1 . We define nested and unnested with respect to B_2 similarly. Note that nested with respect to B_1 is the same as unnested with respect to B_2 and nested with respect to B_2 is unnested with respect to B_1 . As an example, the two saddles depicted in the first illustration in Figure 7 are nested with respect to different 3–balls.

Two saddles $\sigma = s_1^\sigma \vee s_2^\sigma$ and $\tau = s_1^\tau \vee s_2^\tau$ in F_C are *adjacent* if, up to subscript labels, s_1^σ and s_1^τ cobound an annulus in C that is disjoint from s_2^σ, s_2^τ , all other saddles, and K . Recall that, if σ is a standard saddle, E_σ is the monotone disk in C with boundary σ .

Lemma 3.6 *Assume P and P' are adjacent thin spheres with P' above P and C is a cut-disk for P above it but disjoint from P' . If σ and τ are adjacent saddles in F_C such that σ and τ are nested with respect to different 3–balls, then C is not taut.*

Proof Assume σ and τ are adjacent saddles in F_C such that σ and τ are nested with respect to different 3–balls. By Corollary 3.5, we can assume both c_3^σ and c_3^τ bound monotone disks E_σ and E_τ respectively.

Let A be the monotone annulus in C with boundary $s_1^\sigma \cup s_1^\tau$. If K meets $A \cup E_\sigma \cup E_\tau$ (the annulus in C with boundary $s_2^\sigma \cup s_2^\tau$), then one of s_2^σ or s_2^τ bounds a disk E^* in C that is disjoint from K and an outermost saddle of F_{E^*} is inessential. By Lemma 3.4, C is not taut. Hence, we can assume K is disjoint from $A \cup E_\sigma \cup E_\tau$.

Without loss of generality, suppose σ lies above τ . Let B be the 3–ball in S^3 with boundary $D_1^\tau \cup A \cup E_\sigma \cup D_2^\sigma$. If B does not contain $+\infty$, then use the isotopy constructed in [18, Lemma 3] to eliminate τ without introducing any new saddles to

F_C and without introducing any new maxima to $h|_K$. See Figure 7. If B does contain $+\infty$, then use the isotopy in Case 1 of Lemma 3.4 to isotope P' to be bowl-like to guarantee that B_σ does not contain $+\infty$, then apply [18, Lemma 3] to reduce the number of saddles for C . Finally, restore P' to be level. As in Case 1 of Lemma 3.4, these isotopies do not affect the width of K below P and above P' and do not introduce new critical points for K , so they do not increase the width of the knot. Thus, C is not taut. \square




Figure 7

Lemma 3.7 Assume P and P' are adjacent thin spheres with P' above P and C is a cut-disk for P above it but disjoint from P' . If σ is an outermost saddle in F_C such that $B_\sigma \cap C \neq \emptyset$, then C is not taut.

Proof Assume to form a contradiction that C is taut. With out loss of generality, assume σ is nested with respect to B_1 . Since all saddles in F_C are standard, by Corollary 3.5, and no saddles in F_C are inessential, by Lemma 3.4, then there is a labeling of the saddles of F_C as $\sigma = \sigma_1, \dots, \sigma_n$ such that σ is an outermost c -inessential saddle and σ_i is adjacent to σ_{i+1} for each $i \in \{1, \dots, n-1\}$. Inductively, by Lemma 3.6, σ being nested with respect to B_1 implies all saddles in F_C are nested with respect to B_1 . Hence, B_2 can be decomposed into B_σ together with a collection of vertical solid cylinders and solid elbows. Thus, B_σ is disjoint from C , contradicting the hypothesis. \square

Lemma 3.8 Assume P and P' are adjacent thin spheres with P' above P and C is a cut-disk for P above it and disjoint from P' . If F_C contains a c -inessential saddle, then C is not taut.

Proof Suppose σ is a c -inessential saddle in F_C .

The isotopy utilized in the following claim was originally described in [18, page 5].

Claim *If D_σ has a minimum and is punctured by K , we may assume that $h|_{K \cap B_\sigma}$ also has a local minimum at $K \cap D_\sigma$. Symmetrically, if D_σ has a maximum and is punctured by K , we may assume that $h|_{K \cap B_\sigma}$ also has a local maximum at $K \cap D_\sigma$.*

Proof of claim Suppose D_σ has a minimum and $h|_{K \cap B_\sigma}$ has a local maximum at $p_\sigma = K \cap D_\sigma$. Let x be the minimum of K that is nearest p_σ and inside B_σ . Let α be the monotone subarc of K inside B_σ with boundary points p_σ and x . Let β be a monotone arc in D_σ with endpoints p_σ and y such that $h(y) = h(x)$. Let δ be a level arc contained in B_σ connecting x to y . Let E^* be the vertical disk with boundary $\alpha \cup \beta \cup \delta$ that is embedded in B_σ . We can assume the interior of E^* meets K transversely in a collection of points k_1, \dots, k_n where $h(k_1) > h(k_2) > \dots > h(k_n)$. It is important to note that if C meets the interior of E^* then C is not taut, by Lemma 3.6. Hence, we can assume C is disjoint from E^* . Let μ_i be the arc corresponding to a small neighborhood of k_i in $K \cap B_\sigma$ for each i .

Replace μ_n with a monotone arc which starts at an end point of μ_n , runs parallel to E^* until it nearly reaches D_σ , travels along D_σ until it returns to the other side of E^* , travels parallel to E^* (now on the opposite side) and connects to the other end point of μ_n . The result is isotopic to K , does not change the number of maxima of $h|_K$ and reduces n . By induction on n , we may assume that $K \cap E^* = \emptyset$. Isotope α along E^* until it lies just outside of D_σ except where it intersects D_σ exactly at the point y . After a small tilt of K , $h|_{K \cap B_\sigma}$ now has a local minimum at p_σ . \square

After applying the isotopy given by the claim, we can repeat the arguments in Lemma 3.4 to remove this saddle. We give a very brief summary here.

Case 1 Suppose D_σ contains a unique maximum. If necessary, isotope C , P' and all level spheres above P' so that B_σ is disjoint from $+\infty$. This isotopy replaces P' and all level spheres above it with bowl-like spheres. Next, use the isotopy in [1, Lemma 3] to eliminate σ ; see Figure 8. Finally isotope C , P' and all the other spheres that used to be level, to be level again. This can be done without introducing any new saddles or critical points for K . This isotopy fixes all of S^3 below $h(\sigma)$ and above P' . Additionally, this isotopy removes at least one saddle of C and does not create any new critical points for $h|_K$ in the thick region between P and P' . Since all maxima of $h|_K$ are above all minima of $h|_K$ in the region between P and P' , altering the relative heights of the critical points without creating any new critical points can only decrease the width of K . Hence, C is not taut.

Case 2 Suppose D_σ contains a unique minimum.




Figure 8

Note that if B_σ contains $-\infty$, then B_σ contains P and $B_\sigma \cap C \neq \emptyset$. By Lemma 3.7, C is not taut. Therefore, we may assume that B_σ does not contain $-\infty$. By the claim, we may also assume that $h|_{K \cap B_\sigma}$ has a local minimum at $p_\sigma = K \cap D_\sigma$. We can then use the isotopy from [1, Lemma 3] to eliminate σ . As in the previous case this isotopy can only decrease the width of K and decreases the number of saddles. Hence, C is not taut. \square

Theorem 3.9 *Let K be a knot in S^3 in thin position and let P and P' be two adjacent thin spheres or let P be the highest thin sphere for K . Suppose P is cut-compressible with the cut-disk C above P such that $C \cap P' = \emptyset$. Then there is an isotopy of K supported between P and P' that does not change the width of K after which C is vertical.*

Proof We can assume that we have isotoped C to be taut. By Lemma 3.4 and Lemma 3.8, F_C has no inessential and no c -inessential saddles. Hence, C is vertical. \square

The above theorem allows us use the following previously known result.

Theorem 3.10 [22] *Let K be a prime knot in thin position and suppose P is a thin sphere. Let D^* be a compressing disk or a vertical cut-disk for P , say above it. Then there is a thin sphere above D^* and if S_0 is the lowest such thin level sphere, then $w(S_0) < w(P)$.*

Combining Theorem 3.9 and Theorem 3.10 we obtain the following corollary.

Corollary 3.11 *Let K be a prime knot in thin position and suppose P is a thin sphere. Let D^* be a c -disk for P , say above it. Then there is a thin sphere above P and if P' is the lowest such thin sphere, then either $D^* \cap P' \neq \emptyset$ or $w(P') < w(P)$.*

4 Bridge surfaces

In the previous section, we focused on results pertaining to thin position for knots. Here we review results about a tangle in bridge position. We begin with a brief review of the definition of a bridge surface and its distance. For more details see [21].

Suppose M is a 3-manifold homeomorphic either to S^3 , to a ball, or to $S^2 \times I$ and containing a properly embedded compact 1-manifold, T . A sphere Σ is a *bridge sphere* for M if $M - \Sigma$ has two components each of which is either a ball or is homeomorphic to $S^2 \times I$ and each component of $T - \Sigma$ is boundary parallel in $M - \Sigma$ to Σ or it is a vertical arc in $S^2 \times I$. Let H^+ and H^- be closure of the components of $M - \Sigma$ and let $\tau^\pm = T \cap H^\pm$. We say that $(\Sigma, (H^-, \tau^-), (H^+, \tau^+))$ is a *bridge splitting* for (M, T) .

If B is a ball in (M, T) , under certain conditions $B \cap \Sigma$ induces a bridge sphere for $(B, B \cap T)$ as described in the following lemma.

Lemma 4.1 *Suppose K is a knot in thin position with two adjacent thin spheres P and P' such that P' is above P . Suppose D^* is a vertical c -disk for P lying above it and disjoint from P' . Let B be the ball cobounded by D^* and P disjoint from P' and let $T = K \cap B$. Then there exists a thick sphere Σ between P and P' such that the disk $\Delta = B \cap \Sigma$ together with the possibly once punctured disk that $\partial\Delta$ bounds in D^* is a bridge sphere for (B, T) .*

Proof Since D^* is vertical we can lower all of the minima of K in the region bounded by P and P' until each of the minima is below $K \cap D^*$. This isotopy fixes D^* and alters K only in a neighborhood of its minima. After this isotopy, a level sphere just above the highest minimum of K in the region bounded by P and P' is a thick sphere. Denote one such thick sphere by Σ . Since D^* is vertical, $\beta = D^* \cap \Sigma$ is a single essential simple closed curve. Note that β separates D^* into a disk that meets K in at most one point and an annulus that is disjoint from K .

Let E be a bridge disk for some bridge contained in T . By redefining E , we may assume that $E \cap D^*$ consists only of arcs. Let α be an outermost arc of intersection that has both of its endpoints in β so that the disk F that α cobounds with a segment of β in D^* is disjoint from E and does not contain the puncture of D^* . We can redefine E by replacing the subdisk α cuts off in E with the disk F . This reduces $|D^* \cap E|$. Therefore every bridge disk for an arc in B is either disjoint from D^* or intersects it in a single arc that has one endpoint in β and one on the puncture of D^* . Moreover there is at most one bridge disk that does the latter. Thus Δ together with a parallel copy of the possibly once punctured disk that $\partial\Delta$ bounds in D^* is a bridge sphere for (B, T) . \square

Suppose $(\Sigma, (H^-, \tau^-), (H^+, \tau^+))$ is a bridge splitting for (M, T) . The *curve complex*, $\mathcal{C}(\Sigma, T)$, is a graph with vertices corresponding to isotopy classes of essential simple closed curves in $\Sigma - \eta(T)$. Two vertices are adjacent in $\mathcal{C}(\Sigma, T)$ if their corresponding classes of curves have disjoint representatives.

Let \mathcal{V}^+ (respectively \mathcal{V}^-) be the set of all essential simple closed curves in $\Sigma - \eta(T)$ that bound disks in $H^+ - \eta(T)$ (respectively $H^- - \eta(T)$). Then the *distance of the bridge splitting*, $d(\Sigma, T)$, is defined to be the minimum distance between a vertex in \mathcal{V}^+ and a vertex in \mathcal{V}^- measured in $\mathcal{C}(\Sigma, T)$ with the path metric.

We will need the following special case of a result proven in [9].

Theorem 4.2 [9, Theorem 4.4] *Suppose N is a 3–sphere, a 3–ball or $S^2 \times I$ containing a properly embedded compact 1–manifold, K . Let M be a submanifold homeomorphic to a 3–ball or to $S^2 \times I$ such that $T = K \cap M$ is a collection of loops and arcs properly embedded in M . Let Σ be a bridge sphere of (M, T) and let Σ' be a bridge sphere of (N, K) . Then one of the following holds:*

- *There is an isotopy of Σ'_K followed by some number of compressions and cut-compressions of $\Sigma'_K \cap M$ in M giving a compressed surface Σ''_K such that $\Sigma''_K \cap M$ is parallel to Σ_K .*
- *$d(\Sigma, T) \leq 2 - \chi(\Sigma'_K)$.*
- *$\chi(\Sigma_K) \geq -3$.*

The following result can be easily obtained by a simplified version of the proof of Theorem 4.2 given in [9] so we will not prove it here.

Theorem 4.3 *Suppose N is a manifold containing a properly embedded compact 1–manifold, K . Let M be a submanifold homeomorphic to a ball or to $S^2 \times I$ such that $T = K \cap M$ is a compact 1–manifold properly embedded in M . Let Σ be a bridge surface for (M, T) and F be an essential separating surface in N . Then one of the following holds:*

- *$d(\Sigma, T) \leq 2 - \chi(F_K)$.*
- *$\chi(\Sigma_K) \geq -3$.*
- *Each component of $F \cap (M - \eta(T))$ is boundary parallel in $M - \eta(T)$.*

Finally, on multiple occasions in our construction we will need tangles that satisfy certain distance conditions. We will make use of the following two results.

Corollary 4.4 [3, Corollary 4.10] *Let B be a ball containing a b -strand tangle T with $b \geq 3$. Let p_1, \dots, p_k be a collection of points $T \cap \partial B$ such that $1 < k < 2b - 1$ and let n be any positive integer. Then there is a curve γ that bounds a disk in ∂B containing exactly the punctures p_1, \dots, p_k such that for every compressing disk D for $\partial B - \eta(K)$ contained in B , $\text{dist}(\partial D, \gamma) \geq n$.*

Corollary 4.5 [3, Corollary 5.3] *Given positive integers b, c, d , and g with $c \leq b$ such that if $g = 0$, then $b \geq 3$, and if $g = 1$, then $b \geq 1$, there exists a closed orientable 3-manifold M containing a c -component link L and a bridge surface Σ of genus g for (M, L) so that L is b -bridge with respect to Σ and $d(\Sigma, L) \geq d$.*

Letting $g = 0$ and $c = 1$ allows us to choose the above M to be the three sphere and the link to be a knot. Furthermore by removing any 3-ball disjoint from the bridge surface that intersects the link in exactly $k < b$ trivial arcs we obtain the following:

Corollary 4.6 *Given positive integers b, d and k with $k < b$ and $b \geq 3$, there exists a 3-ball containing a k -component tangle T with no closed components and a bridge sphere Σ so that $|T \cap \Sigma| = 2b$ and $d(\Sigma, T) \geq d$.*

5 Knotting

The goal of this section is to produce 3-strand tangles so that the strands of the tangles are knotted in some sense while also controlling the distance between certain curves in the curve complex of the 6-punctured sphere. We will later show how these tangles can be inserted in the boxes in Figure 1 to construct a knot that is in thin position. We begin by reviewing some definitions.

Definition 5.1 An n -strand tangle, \mathcal{R} , is *rational* if all arcs of R can be simultaneously isotoped into $\partial(B_R)$. A tangle \mathcal{T} is an *induced subtangle* of a tangle \mathcal{R} if $B_T = B_R$ and $T \subseteq R$.

Definition 5.2 Given a rational tangle \mathcal{R} and a simple closed curve ϵ in $\partial(B_R)$, (\mathcal{R}, ϵ) is an *equatorial pair* if ϵ is disjoint from R and no arc in R has both of its endpoints on the same component of $\partial(B_R) - \epsilon$. (\mathcal{T}, ϵ) is an *equatorial subpair* of an equatorial pair (\mathcal{R}, ϵ) if (\mathcal{T}, ϵ) is an equatorial pair and \mathcal{T} is an induced subtangle of \mathcal{R} .

Definition 5.3 Given an n -strand tangle, \mathcal{R} , and an equatorial pair, (\mathcal{R}, ϵ) , embed B_R as the unit ball in R^3 such that ϵ is mapped to the unit circle in the xy -plane and

all points of $R \cap \partial(B_R)$ are mapped to the unit circle in the xz -plane. A *projection* of (\mathcal{R}, ϵ) is a projection of such an embedding of R into the xz -plane. If \mathcal{R} is a 2-strand tangle, a *numerator closure* of (\mathcal{R}, ϵ) is a knot obtained by connecting the endpoints of R via two arcs in the unit circle in the xz -plane so that each of the arcs is disjoint from the unit circle in the xy -plane.

In the remainder of the section, we will heavily rely on standard results about rational tangles. In particular, recall that each proper isotopy class of 2-strand rational tangle can be represented by a fraction p/q , where $(p, q) = 1$. See [10], or [4] for a detailed treatment of the subject.

Theorem 5.4 [4] *Two 2-strand rational tangles are properly isotopic if and only if they have the same fraction.*

Theorem 5.5 [17] *Consider two 2-strand rational tangles with fractions p/q and p'/q' . If $K(p/q)$ and $K(p'/q')$ denote the corresponding rational knots obtained by taking the numerator closures of these tangles, then $K(p/q)$ and $K(p'/q')$ are isotopic if and only if*

- $p = p'$ and
- either $q \equiv q' \pmod{p}$ or $qq' \equiv 1 \pmod{p}$.

By Theorem 5.5, if \mathcal{R} is a rational 2-strand tangle, (\mathcal{R}, ϵ) has numerator closure the unknot if and only if (\mathcal{R}, ϵ) has a projection as a $1/q$ rational tangle. We will call such an equatorial pair the *unpair*.

Theorem 5.6 *If \mathcal{R} is a rational 3-strand tangle and (\mathcal{R}, ϵ) is an equatorial pair, then there exists an equatorial pair (\mathcal{R}, δ) such that no 2-strand equatorial subpair is the unpair and $d(\epsilon, \delta) \leq 4$ (where d is the distance function for the curve complex of the 6-punctured sphere).*

Proof Suppose R consists of three arcs α , β and γ . The equatorial pair (\mathcal{R}, ϵ) has three 2-strand equatorial subpairs $(\mathcal{R}_1, \epsilon)$, $(\mathcal{R}_2, \epsilon)$ and $(\mathcal{R}_3, \epsilon)$, where R_1 contains α and β , R_2 contains β and γ , and R_3 contains γ and α . Let p_1/q_1 , p_2/q_2 and p_3/q_3 be the fractions corresponding to the proper isotopy classes of $(\mathcal{R}_1, \epsilon)$, $(\mathcal{R}_2, \epsilon)$ and $(\mathcal{R}_3, \epsilon)$ respectively. In Definition 5.3, we are free to choose how the points of $R \cap \partial(B_R)$ are mapped to the unit circle in the xz -plane. In particular, we are free to twist pairs of points of ∂R that get mapped to the southern hemisphere of the unit sphere. This twisting changes the proper isotopy class of each of $(\mathcal{R}_1, \epsilon)$, $(\mathcal{R}_2, \epsilon)$ and $(\mathcal{R}_3, \epsilon)$. In Conway's notation, this twisting corresponds to multiplying each fraction

p_1/q_1 , p_2/q_2 and p_3/q_3 by $1/m$, where m corresponds to the number of twists. Since m can be chosen to be arbitrarily large, we can map B_R to the unit ball so that each of the rational numbers p_1/q_1 , p_2/q_2 and p_3/q_3 lie strictly between -1 and 1 .

If none of $(\mathcal{R}_1, \epsilon)$, $(\mathcal{R}_2, \epsilon)$ or $(\mathcal{R}_3, \epsilon)$ are the unpair, then we are done.

Suppose $(\mathcal{R}_1, \epsilon)$ is the unpair. By Theorem 5.5, $(\mathcal{R}_1, \epsilon)$ has fraction $1/n$. Note that, by inserting a sufficiently large number of twists between the boundary points of α and β in the southern hemisphere, we can assume n , the denominator for the fraction representation of $(\mathcal{R}_1, \epsilon)$, is positive. Isotope \mathcal{R} as in Figure 9. This isotopy alters α but fixes β and γ . After this isotopy, (\mathcal{R}, δ_1) is an equatorial pair such that the numerator closure of $(\mathcal{R}_1, \delta_1)$ is the twisted Whitehead double of the unknot.

Since ϵ is isotopic to δ_1 in $\partial(B_R) - \partial(R_2)$, then $(\mathcal{R}_2, \epsilon) = (\mathcal{R}_2, \delta_1)$. However, $(\mathcal{R}_3, \epsilon) \neq (\mathcal{R}_3, \delta_1)$. Since $(\mathcal{R}_3, \epsilon)$ has fraction p_3/q_3 and the isotopy in Figure 9 corresponds to Conway sum of the projection of $(\mathcal{R}_3, \epsilon)$ and a tangle with fraction $2/1$, then $(\mathcal{R}_3, \delta_1)$ is a tangle with fraction $(p_3 + 2q_2)/q_2$. However, p_3/q_3 was assumed to be strictly between -1 and 1 . Hence, $(p_3 + 2q_2)/q_2$ is strictly between 1 and 3 and cannot be of the form $1/r$ for any integer r . Thus, by Theorem 5.5, $(\mathcal{R}_3, \delta_1)$ is not the unpair.




Figure 9




Figure 10

As illustrated in Figure 10, there is an essential simple closed curve in $\partial(B_R) - \partial(R)$ that is disjoint from both ϵ and δ_1 . Thus, $d(\epsilon, \delta_1) \leq 2$. If $(\mathcal{R}_2, \delta_1)$ is not an unpair, then we are done.

Suppose $(\mathcal{R}_2, \delta_1)$ is an unpair, we repeat the previous process. Let p_1^1/q_1^1 , p_2^1/q_2^1 and p_3^1/q_3^1 be the fractions corresponding to the proper isotopy classes of $(\mathcal{R}_1, \delta_1)$, $(\mathcal{R}_2, \delta_1)$ and $(\mathcal{R}_3, \delta_1)$ respectively. By Theorem 5.5, $(\mathcal{R}_2, \delta_1)$ has fraction $1/r$. By inserting a sufficiently large number of twists between the boundary points of β and γ in the southern hemisphere, we can assume r , the denominator for the fraction representation of $(\mathcal{R}_1, \epsilon)$, is positive.

Isotope \mathcal{R} in a fashion similar to that of Figure 9, except with δ_1 replacing ϵ , δ_2 replacing δ_1 . We choose this isotopy so that it alters γ but fixes α and β . After this isotopy, (\mathcal{R}, δ_2) is an equatorial pair such that the numerator closure of $(\mathcal{R}_2, \delta_2)$ is a twisted Whitehead double of the unknot. Since this isotopy fixes α and β , the numerator closure of $(\mathcal{R}_1, \delta_2)$ remains a twisted Whitehead double of the unknot. Since p_3^1/q_3^1 is strictly between 1 and 3, then $(p_3^1 + 2q_2^1)/q_2^1$ is strictly between 3 and 5 and cannot be of the form $1/r$ for any integer r . Thus, by Theorem 5.5, $(\mathcal{R}_3, \delta_2)$ is not the unpair. As before, there is an essential simple closed curve in $\partial(B_R) - \partial(R)$ that is disjoint from both δ_1 and δ_2 . Thus, $d(\epsilon, \delta_2) \leq d(\epsilon, \delta_1) + d(\delta_1, \delta_2) = 2 + 2 = 4$. □

6 Essential surfaces in high distance tangles

In this section, we lay the foundations that will eventually allow us to construct a tangle with only one essential surface with Euler characteristic near zero. We begin with some definitions which build on the definitions introduced in the last section.

Definition 6.1 Let \mathcal{R} be a rational n -strand tangles and \mathcal{Q} be a rational m -strand. Let \mathcal{C}_l be the curve complex for the $2l$ -punctured sphere. Let \mathcal{V}_R be the set of all isotopy classes of essential simple closed curves in $\partial(B_R) - R$ that bound disks in $B_R - R$. Define \mathcal{V}_Q analogously.

Let γ_R (respectively γ_Q) be an essential curve on $\partial(B_R) - R$ (respectively $\partial(B_Q) - Q$) such that γ_R (respectively γ_Q) bounds a k -punctured disk D_R (respectively D_Q) in $\partial(B_R)$ (respectively $\partial(B_Q)$). Create a new $(n + m - k)$ -strand tangle \mathcal{T} by identifying D_R and D_Q via a homeomorphism ψ of the k -punctured disk. The resulting tangle \mathcal{T} depends on D_R , D_Q and ψ . Let B_T be the 3-ball obtained by gluing B_R to B_Q in this way. Let D be the image of D_R and D_Q in B_T .

Definition 6.2 Let F be a properly embedded surface or a submanifold in a 3-manifold M and let K be a properly embedded compact 1-manifold in M . If $\eta(K)$ is a regular open neighborhood of K in M , define F_K to be $F - \eta(K)$.

Definition 6.3 Given an n -strand rational tangle \mathcal{R} , let B be a small ball in the interior of B_R disjoint from R . Then $B_R - B$ is homeomorphic to $S^2 \times I$. We may isotope R so each strand of R has exactly one critical point with respect to the height function obtained from the natural projection of $S^2 \times I$ onto its I factor. Connect each of these n critical points to ∂B via n vertical arcs, one from each critical point to ∂B . Let $\Gamma_R = B \cup (\bigcup_1^k \tau_i)$. Then Γ_R is the *spine* of \mathcal{R} . Note that the tangle obtained from (B_R, R) by removing a regular open neighborhood of Γ_R is homeomorphic to $(S^2 \times I, R^-)$, where R^- is a collection of properly embedded vertical arcs.

Lemma 6.4 [21, Lemma 2.9] *If F_K is a connected incompressible surface in a rational tangle \mathcal{R} , then one of the following holds:*

- (1) F_K is a sphere bounding a ball.
- (2) F_K is a twice punctured sphere bounding a ball containing an unknotted arc.
- (3) $F_K \cap \partial(B_R) \neq \emptyset$.

Definition 6.5 A tangle \mathcal{T} is *prime* if every embedded 2-punctured sphere in B_T bounds a 3-ball containing an unknotted arc.

Theorem 6.6 *Let \mathcal{T} be an irreducible, prime 3-strand tangle as in Definition 6.1. Suppose F_K is a properly embedded connected c -incompressible surface in \mathcal{T} with ∂F_K a possibly empty collection of curves isotopic in $\partial(B_R) - R$ to ∂D . If F_K can be isotoped to be disjoint from a spine of \mathcal{R} and a spine of \mathcal{Q} , then F_K is one of the following:*

- (1) F_K is a sphere bounding a ball.
- (2) F_K is a twice punctured sphere bounding a ball containing a unknotted arc.
- (3) F_K is isotopic to $\partial(B_R)_K - \text{int}(D)$.
- (4) F_K is isotopic to $\partial(B_Q)_K - \text{int}(D)$.
- (5) F_K is isotopic to $\partial(B_T)_K$.
- (6) F_K is isotopic to D_K .
- (7) F_K is an annulus isotopic into $\partial(B_T)_K$.

Proof Let $F^R = F \cap B_R$ and $F^Q = F \cap B_Q$, and let Γ_R be the spine of the rational tangle \mathcal{R} such that $F \cap \Gamma_R = \emptyset$. Let M_R be the complement of an open neighborhood of Γ_R in B_R . Then M_R is homeomorphic to $(S^2 \times I)$ with $\partial(M_R) = \partial_+(M_R) \sqcup \partial_-(M_R)$ so that $\partial_+(M_R)$ is the boundary of a regular neighborhood of Γ_R and $\partial_-(M_R) = \partial(B_R)$. Hence, F^R is properly embedded in M_R , is disjoint from

$\partial_+(M_R)$ and meets $\partial_-(M_R)$ only in D . It will be convenient to refer to the height function, h_R , on M_R obtained from the natural projection of $S^2 \times I$ onto its I factor.

As \mathcal{R} is a rational tangle, $K \cap M_R$ is isotopic to a collection of arcs, $\{x_1, \dots, x_6\}$ that are monotone with respect to h_R and where x_4, x_5 and x_6 are the unique arcs of $K \cap M_R$ that meet D . Let E be an embedded vertical rectangle in M_R that is disjoint from D and contains x_1, x_2 and x_3 such that ∂E is the end point union of x_1, γ, x_3 and γ' , where γ is an arc in $\partial_-(M_R)$ and γ' is the image of γ under the natural projection from M_R onto $\partial_+(M_R)$. By assumption, after isotopy, any arc of $E \cap F_K$ is disjoint from both γ and γ' . Suppose $\alpha \in F_K \cap E$ is any arc or simple closed curve.

Claim 1 *If α meets x_i more than once, then we have conclusion (2).*

Proof If α meets some x_i more than once, then for some x_j there is a bigon H' in E cobounded by a subarc of x_j and a subarc of α such that the interior of H' is disjoint from α and from x_1, x_2 and x_3 . F meets the boundary of a closed regular neighborhood of H' , $\partial(\eta(H'))$, in a single simple closed curve. Since H' is a disk, $\partial(\eta(H'))$ is a 2–sphere. The closure of each component of $\partial(\eta(H')) - F$ is a disk. One of these disks, together with a 2–punctured subdisk of F_K cobound a 3–ball containing an unknotted subarc of K . As F_K is incompressible, we must have conclusion (2). \square

Suppose that α is a simple closed curve. If α is disjoint from x_2 , it can be eliminated by using an innermost disk argument that appeals to the incompressibility of F_K and the irreducibility of $M_R - K$. If α intersects x_2 , then it must intersect it at least twice and, therefore, by Claim 1, we have conclusion (2). Hence, we can assume $F_K \cap E$ contains no simple closed curve

Suppose that $\alpha \in F_K \cap E$ is an arc. By Claim 1, one endpoint of α must be in x_1 and the other in x_3 . Furthermore, by Claim 1, we can assume $E \cap F_K$ consists of arcs that meet each of the strands x_1, x_2 and x_3 in exactly one point each. After an isotopy, we can assume that each curve in $E \cap F_K$ is level with respect to h_R . If $\eta(E)$ is a regular open neighborhood of E , then let $N_R = M_R - \eta(E)$. Hence, we can assume that F_K^R meets M_R outside of N_R in a, possibly empty, collection of level disks each meeting K in three points. By repeating the symmetric argument with an embedded vertical rectangle E^* in M_Q , we can assume that F_K^Q meets M_Q outside of N_Q in a, possibly empty, collection of level disks each meeting K in three points.

Let N be the union of N_R and N_Q in B_T , see Figure 11. Outside of N , F_K is a collection of 3–punctured disks that are level with respect to h_R or h_Q . Call the boundary curves of these two collections \mathcal{D}_R and \mathcal{D}_Q respectively. Since F_K meets




Figure 11

$\partial(B_T)$ in a collection of curves, \mathcal{D} , parallel to ∂D , then $F_K \cap \partial N = \mathcal{D} \cup \mathcal{D}_R \cup \mathcal{D}_Q$. Additionally, any two curves in $\mathcal{D} \cup \mathcal{D}_R \cup \mathcal{D}_Q$ are isotopic in $\partial(N)_K$. N is homeomorphic to $D^2 \times I$, where $D^2 \times \{1\} = \partial_+(M_R) - \eta(E)$ and $D^2 \times \{0\} = \partial_+(M_Q) - \eta(E^*)$. Hence, N has a natural height function, h_N , induced by projection onto the I factor. In particular, h_N can be chosen so that every curve in $\mathcal{D} \cup \mathcal{D}_R \cup \mathcal{D}_Q$ is level and each arc of $K \cap N$ is monotone. Let H be a properly embedded vertical disk in N that contains all three strands of $N \cap K$. Note that ∂H meets every curve in $\mathcal{D} \cup \mathcal{D}_R \cup \mathcal{D}_Q$ in exactly two points and each component of $N - H$ is a 3-ball disjoint from K . See Figure 12.

Suppose $H \cap F = \emptyset$. Since every curve in $F \cap \partial N$ meets H , then $F \cap \partial N = \emptyset$ and F is contained in $N - H$. Hence, we have conclusion (1).

Suppose $H \cap F \neq \emptyset$. If any component of $F \cap H$ is a closed curve disjoint from K , then, by the incompressibility of F_K and the irreducibility of $N - \eta(K)$, we can remove it via an isotopy of F_K supported in N_K . If any component of $F \cap H$ is a closed curve not disjoint from K , then, by appealing to the argument in Claim 1, we have conclusion (2). Hence, every component of $F_K \cap H$ is an arc. Label the endpoints of an outermost such arc as in Figure 12, where the a_i^\pm lie on \mathcal{D}_R , the b_i^\pm lie on \mathcal{D}_Q , and the c_i^\pm lie on \mathcal{D} . There is an outermost arc in $F_K \cap H$ with one of the following endpoint labels:

- (1) a_1^- and a_1^+ (similarly, b_1^- and b_1^+)
- (2) a_i^- and a_{i+1}^- (similarly, b_i^- and b_{i+1}^- , a_i^+ and a_{i+1}^+ , or b_i^+ and b_{i+1}^+)
- (3) c_i^- and c_{i+1}^- (similarly, c_i^+ and c_{i+1}^+)
- (4) a_r^+ and c_1^+ (similarly, a_r^- and c_1^- , b_1^+ and c_s^+ , or b_1^- and c_s^-)

- (5) c_1^- and c_1^+ (similarly, c_s^- and c_s^+)
- (6) a_r^+ and b_1^+ (similarly, a_r^- and b_1^-)




Figure 12

Let y_1, y_2 and y_3 be the three strands of K in N . Let α be an outermost arc of $F \cap H$ in H . If α meets one of y_1, y_2 and y_3 in more than one point, then as in Claim 1, we have conclusion (2). Hence, we can assume that α meets each of y_1, y_2 and y_3 in at most one point.

Case 1 Suppose α is an outer most arc of $F_K \cap H$ with endpoints a_1^- and a_1^+ . Since α meets each of y_1, y_2 and y_3 in at most one point, $\alpha \cap K$ consists of exactly three points. Let L be the disk in $F^R - N$ that contains a_1^- and a_1^+ . The disk L together with a neighborhood of α in F_{KR} is a 6-punctured annulus in M_R . Both boundary components of this annulus are contained in the interior of M_R and bound disks in M_R disjoint from both F and K . See Figure 13. By incompressibility of F_K and irreducibility of $M_R - K$, both boundary components of this annulus bound disks in F_K . Hence, F_R is a 6-punctured sphere in M_R isotopic to $\partial(B_R)_K$. This is a contradiction to F_K being incompressible, so such an outermost arc must not exist.

Case 2 Suppose α is an outermost arc of $F_K \cap H$ with endpoints a_i^- and a_{i+1}^- . Since α meets each of y_1, y_2 and y_3 in at most one point, $\alpha \cap K = \emptyset$. Let L_1 and L_2 be the two 3-punctured disks in $F_K^R - N$ that contain a_i^- and a_{i+1}^- in their respective boundaries. Since α is outermost, we can isotope it to be monotone with respect to h_N . Let x_i be one of the strands of K in $M_R - N$. As in Figure 14, there is a disk G in $M_R - R$ that is vertical with respect to h_R and illustrates a parallelism between a subarc of $x_i, i \in \{1, 2, 3\}$, and an arc in F_K . As in the proof of Claim 1, we have conclusion (2).

Case 3 Suppose α is an outermost arc of $F_K \cap H$ with endpoints c_i^- and c_{i+1}^- . Since α meets each of y_1, y_2 and y_3 in at most one point, $\alpha \cap K = \emptyset$. Let γ_1 and γ_2




Figure 13




Figure 14

be the curves in \mathcal{D} that contain c_i^- and c_{i+1}^- respectively. Let C_1 and C_2 be small closed neighborhoods of γ_1 and γ_2 in F respectively. A neighborhood of α in F_K together with C_1 and C_2 form a connected subsurface of F_K with three boundary components γ_1 , γ_2 and β . The curve β bounds a disk in N disjoint from F and K . By incompressibility of F_K and irreducibility of $N - K$, β bounds a disk in F_K . Hence, we have conclusion (7).

Case 4 Suppose α is an outermost arc of $F_K \cap H$ with endpoints a_r^+ and c_1^+ . Since α meets each of y_1 , y_2 and y_3 in at most one point, $\alpha \cap K = \emptyset$. Let L be the disk in $F_K^R - N$ with boundary in \mathcal{D}_R that contains a_r^+ . Let γ be the curve in \mathcal{D} that contains c_1^+ and let C_1 be a small closed neighborhood of γ in F . A neighborhood of α in F_K together with E and C_1 forms a 3-punctured annulus subsurface of F_K with boundary components γ and β . The curve β bounds a disk in N disjoint from

F and K . By incompressibility of F_K and irreducibility of $N - K$, β bounds a disk in F_K . Hence, we have conclusion (3). By a similar argument, if α has endpoints b_1^+ and c_s^+ or b_1^- and c_s^- we have conclusion (4).

Case 5 Suppose α is an outermost arc of $F_K \cap H$ with endpoints c_1^- and c_1^+ . This case can only occur if there are no a_i^\pm (ie \mathcal{D}_R is empty). Since α meets each of y_1 , y_2 and y_3 in at most one point, $\alpha \cap K$ consists of exactly three points. Let γ be the curve in \mathcal{D} that contains both c_1^- and c_1^+ . Let C be a small closed neighborhood of γ in F . C together with a regular neighborhood of α in F_K is a 3-punctured subsurface of F_K with three boundary components γ , β_1 and β_2 . However, β_1 and β_2 each bound disks in $N - K$ disjoint from F and K . By incompressibility of F_K and irreducibility of $N - K$, both β_1 and β_2 bound disks in F_K . Hence, we have conclusion (6).

Case 6 Suppose α is an outermost arc of $F_K \cap H$ with endpoints a_r^+ and b_1^+ . This case can only occur if there are no c_i^\pm (ie \mathcal{D} is empty). Since α meets each of y_1 , y_2 and y_3 in at most one point, $\alpha \cap K = \emptyset$. Let L_R be the disk in $F^R - N$ with boundary in \mathcal{D}_R such that $a_r^+ \in \partial(L_R)$ and let L_Q be the disk in $F^Q - N$ with boundary in \mathcal{D}_Q such that $b_1^+ \in \partial(L_Q)$. A neighborhood of α in F_K together with L_R and L_Q is a 6-punctured disk subsurface of F_K . The boundary of this disk bounds a disk in $N - K$ disjoint from F and K . By incompressibility of F_K and irreducibility of $N - K$, the boundary of this 6-punctured disk bounds a disk in F_K . Hence, we have conclusion (5). □

Lemma 6.7 *If D_K is compressible in $B_R - R$ then $d(\partial D, \mathcal{V}_R) \leq 1$. If D_K is compressible in $B_Q - Q$ then $d(\partial D, \mathcal{V}_Q) \leq 1$*

Proof Let $E \subseteq B_R - R$ be a compressing disk for D_K . Since ∂E is essential in D_K and disjoint from $\partial(D_K)$, then $\partial E \in \mathcal{V}_R$ and $d(\partial D, \mathcal{V}_R) \leq 1$. □

Lemma 6.8 *If D_K is compressible in $B_T - T$ then $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$.*

Proof If D_K is compressible in $B_T - T$, then D_K is compressible in $B_R - R$ or D_K is compressible in $B_Q - Q$. By Lemma 6.7, $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$. □

Lemma 6.9 [21, Proposition 4.1] *If D_K is cut-compressible in $B_R - R$ then $d(\partial D, \mathcal{V}_R) \leq 2$. Similarly, if D_K is cut-compressible in $B_Q - Q$ then $d(\partial D, \mathcal{V}_Q) \leq 2$.*

Proof Let α be the arc of K in B_R that punctures the cut-compressing disk, C , for D_K . Let B be the disk in B_R illustrating the boundary parallelism of α . After perhaps an isotopy of B , $B \cap C$ is a single arc β that separates B into two disks B_1 and B_2 . Consider a regular neighborhood of $C \cup B_1$. Its boundary contains a disk that intersects $\partial(B_R)_K$ in an essential curve γ and does not intersect ∂C . Hence, $d(\partial D, \mathcal{V}_R) \leq d(\partial D, \partial C) + d(\partial C, \gamma) \leq 2$. \square

Lemma 6.10 *If D_K is c -compressible in $B_T - T$ then either $d(\partial D, \mathcal{V}_R) \leq 2$ or $d(\partial D, \mathcal{V}_Q) \leq 2$.*

Proof If D_K is compressible in $B_T - T$ then, by Lemma 6.8, $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$. If D_K is cut-compressible, then D_K is cut-compressible in $B_R - R$ or D_K is cut-compressible in $B_Q - Q$. By Lemma 6.9, $d(\partial D, \mathcal{V}_R) \leq 2$ or $d(\partial D, \mathcal{V}_Q) \leq 2$. \square

Lemma 6.11 *If $B_T - T$ is reducible then $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$.*

Proof Let S be a reducing sphere for $B_T - T$. Since $B_R - R$ and $B_Q - Q$ are irreducible, S cannot be isotoped to be disjoint from D . Isotope S so that $|S \cap D|$ is minimal. If an innermost curve of $D \cap S$ in S is essential in D_K , then D_K is compressible and, by Lemma 6.8 $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$.

Suppose α is a curve in $D \cap S$ that is innermost in S and bounds a subdisk D' in D . Since α is innermost in S , α bounds a subdisk S' of S that is disjoint from D except in its boundary. After pushing D' slightly off of D toward S' , $D' \cup S'$ is a 2-sphere embedded in either $B_R - R$ or $B_Q - Q$. Since both $B_R - R$ and $B_Q - Q$ are irreducible, $D' \cup S'$ bounds a 3-ball disjoint from K . We can use this 3-ball to construct an isotopy of S that reduces $|S \cap D|$. However, this contradicts the assumption that $|S \cap D|$ is minimal. Hence, α must be essential in D_K . \square

Lemma 6.12 *If $B_T - T$ contains an essential 2-punctured sphere then $d(\partial D, \mathcal{V}_R) \leq 2$ or $d(\partial D, \mathcal{V}_Q) \leq 2$.*

Proof Let S_K be an essential 2-punctured sphere in $B_T - T$. Since $B_R - R$ and $B_Q - Q$ are prime, S_K cannot be isotoped to be disjoint from D . Isotope S_K so that $|S_K \cap D|$ is minimal. If an innermost curve of $D \cap S_K$ in S_K bounds a disk in S_K and is essential in D_K , then D_K is compressible and, by Lemma 6.8, $d(\partial D, \mathcal{V}_R) \leq 1$ or $d(\partial D, \mathcal{V}_Q) \leq 1$.

If an innermost curve of $D \cap S_K$ in S_K bounds a cut-disk in S_K and is essential in D_K , then D_K is cut-compressible. By Lemma 6.10, $d(\partial D, \mathcal{V}_R) \leq 2$ or $d(\partial D, \mathcal{V}_Q) \leq 2$.

Suppose α is an innermost curve of $D \cap S_K$ in S_K that bounds a c -disk S' in S_K and is inessential in D_K . Since α is inessential in D_K , α bounds a c -disk D' in D . After pushing D' slightly off of D toward S' , $D' \cup S'$ is a sphere or a 2-punctured sphere embedded in either $B_R - R$ or $B_Q - Q$. Since both $B_R - R$ and $B_Q - Q$ are irreducible and prime, $D' \cup S'$ bounds a 3-ball or a 3-ball containing an unknotted arc of K . We can use this 3-ball to construct an isotopy of S_K that reduces $|S \cap D|$. However, this contradicts the assumption that $|S \cap D|$ is minimal. Hence, no such α exists. \square

Lemma 6.13 *If $\partial(B_T) - T$ is compressible in $B_T - T$, then $d(\partial D, \mathcal{V}_R) \leq 2$ or $d(\partial D, \mathcal{V}_Q) \leq 2$.*

Proof Assume Δ is a compressing disk for $\partial(B_T) - T$ in $B_T - T$. Isotope Δ so that $|\Delta \cap D_K|$ is minimal. By Lemma 6.8, we can assume that D_K is incompressible. By Lemma 6.11, we can assume that $B_T - T$ is irreducible. Since D_K is incompressible and $B_T - T$ is irreducible we can assume $\Delta \cap D_K$ consists only of arcs and no simple closed curves. Let α be an outermost arc of $\Delta \cap D_K$ in Δ and let F be the subdisk of Δ that α cobounds with an arc in $\partial\Delta$ such that the interior of F is disjoint from D_K . Assume F is properly embedded in B_R . If ∂F is inessential in $\partial(B_R) - R$, then F is boundary parallel in $B_R - R$ and this boundary parallelism can be used to construct an isotopy of Δ that decreases $|\Delta \cap D_K|$, a contradiction. Hence, F is a compressing disk for $B_R - R$ that intersects ∂D in exactly two points. It follows that $d(\partial D, \mathcal{V}_R) \leq 2$. If F is properly embedded in B_Q , then $d(\partial D, \mathcal{V}_Q) \leq 2$. \square

Lemma 6.14 *There exists a tangle \mathcal{T} satisfying Definition 6.1 with $d(\partial(D_R), \mathcal{V}_R) \geq 3$ and $d(\partial(D_Q), \mathcal{V}_Q) \geq 3$.*

Proof This follows immediately from Corollary 4.4. \square

Lemma 6.15 *If F_K is a c -incompressible surface in $B_T - T$ with ∂F consisting of a (possibly empty) collection of simple closed curves isotopic to ∂D in $\partial(B_T) - T$, $d(\partial D, \mathcal{V}_R) \geq 3$ and $d(\partial D, \mathcal{V}_Q) \geq 3$, then, after an isotopy of F_K , $F \cap D$ is a (possibly empty) collection of simple closed curves all of which are essential in both F_K and D_K .*

Proof Since all components of ∂F are isotopic to ∂D in $\partial(B_T) - T$, there is an isotopy of F_K supported in a small neighborhood of $\partial(B_T)$ in B_T resulting in $\partial F \cap \partial D = \emptyset$. Suppose the interior of F_K has been isotoped to minimize $|F \cap D|$ and suppose α is a curve in $D \cap F$ which is inessential in D_K . By appealing to an innermost such α ,

we can assume that the disk or 1-punctured disk, D' , that α bounds in D is disjoint from F except in its boundary. Since F_K is c -incompressible, α bounds a disk or 1-punctured disk, F' , in F . By Lemma 6.11 and Lemma 6.12, $D' \cup F'$ cobound a 3-ball or a 3-ball containing an unknotted arc that gives rise to an isotopy which reduces the number of components of $D \cap F$. Hence, if α is inessential in D_K , then we contradict the minimality of $|D \cap F|$. By Lemma 6.10, D_K is c -incompressible so a similar argument implies that α is also essential in F_K . \square

Theorem 6.16 [21, Proposition 4.3] *Suppose $(F, \partial F) \subset (B_R, \partial(B_R))$ is a properly embedded surface transverse to K that satisfies all of the following conditions:*

- (1) F_K has no disk components.
- (2) F_K is c -incompressible.
- (3) F_K intersects every spine Γ_R of B_R .
- (4) All curves of $F \cap \partial(B_R)$ are essential on $\partial(B_R) - R$.

Then there is at least one curve $f \in F \cap \partial(B_R)$ that is essential in $\partial(B_R) - R$ such that $d(\mathcal{V}_R, f) \leq 1 - \chi(F_K)$ and every $g \in F \cap \partial(B_R)$ that is essential on $\partial(B_R) - R$ for which the inequality does not hold lies in the boundary of a $(\partial(B_R) - R)$ -parallel annulus component of F_K .

Theorem 6.17 *Let \mathcal{T} be a tangle as described in Definition 6.1. In addition, choose D_R and D_Q such that $d(\partial(D_R), \mathcal{V}_R) \geq 3$ and $d(\partial(D_Q), \mathcal{V}_Q) \geq 3$. If F_K is a properly embedded connected c -incompressible surface in $B_T - T$ with ∂F a (possibly empty) collection of curves isotopic to ∂D in $\partial(B_T) - K$, then one of the following holds:*

- (1) F_K is a sphere bounding a ball.
- (2) F_K is a twice punctured sphere bounding a ball containing a unknotted arc.
- (3) F_K is isotopic to $\partial(B_R)_K - \text{int}(D)$.
- (4) F_K is isotopic to $\partial(B_Q)_K - \text{int}(D)$.
- (5) F_K is isotopic to $\partial(B_T)_K$.
- (6) F_K is isotopic to D_K .
- (7) F_K is a $\partial(B_T)_K$ -parallel annulus.
- (8) $d(\partial D, \mathcal{V}_R) \leq 2 - \chi(F_K)$.
- (9) $d(\partial D, \mathcal{V}_Q) \leq 2 - \chi(F_K)$.

Proof There are two cases to consider.

Case 1 Suppose F_K can be isotoped to be disjoint from D . Then F is properly embedded in one of B_R or B_Q with boundary (if nonempty) isotopic to parallel copies of ∂D . Without loss of generality, assume F is contained in B_R . Note that since every component of ∂F is essential in $B_R - R$, no component of F_K is a boundary parallel disk in $B_R - R$. If F_K is a disk, then $d(\partial D, \mathcal{V}_R) = 0$ contradicting the hypothesis of the theorem. If ∂F is empty, then conclusions (1) or (2) hold, by Lemma 6.4. Hence, we can assume that F_K is a c -incompressible properly embedded surface with no disk components and nontrivial boundary. If, in addition, F_K intersects every spine Γ_R , then the hypotheses of Theorem 6.16 are satisfied and there is at least one curve $f \in F_K \cap (\partial(B_R) - R)$ that is essential in $\partial(B_R) - R$ such that $d(\mathcal{V}_R, f) \leq 1 - \chi(F_K)$. Since all curves in $F_K \cap \partial(B_R) - R$ are parallel to ∂D , then conclusion (9) holds. If F_K is disjoint from some spine Γ_R , then the hypotheses of Theorem 6.6 are satisfied and one of conclusions (1) to (7) holds.

Case 2 Suppose F_K cannot be isotoped to be disjoint from D . If F_K can be isotoped to be disjoint from some spine Γ_R and some spine of Γ_Q then, by Theorem 6.6, one of conclusions (1) to (7) holds.

Suppose F_K cannot be isotoped to be disjoint from any spine of \mathcal{R} . (The case where F_K cannot be isotoped to be disjoint from any spine of \mathcal{Q} is proven analogously.) In particular, $F_K^R = F_K \cap B_R$ can not be isotoped to be disjoint from any spine of \mathcal{R} . To apply Theorem 6.16 to F_K^R we need to verify the remaining three hypotheses.

(1) By Lemma 6.10, $d(\partial(D_R), \mathcal{V}_R) \geq 3$ and $d(\partial(D_Q), \mathcal{V}_Q) \geq 3$ imply that D_K is c -incompressible. By Lemma 6.11, $d(\partial(D_R), \mathcal{V}_R) \geq 3$ and $d(\partial(D_Q), \mathcal{V}_Q) \geq 3$ imply that $B_T - T$ is irreducible. By c -incompressibility of D and irreducibility of $B_T - T$ we can assume that $F_K - D$ contains no disk components. Hence, F_K^R contains no disk components.

(2) F_K is assumed to be c -incompressible in $B_T - T$. Hence F_K^R is c -incompressible in $B_R - R$.

(3) We have assumed that F_K intersects every spine of \mathcal{R} .

(4) By Lemma 6.15, D_K and F_K can be isotoped to intersect in a nonempty collection of closed curves that are essential in each surface. In particular, every component of $F_K^R \cap D_K$ is essential in D_K .

Since the hypotheses for Theorem 6.16 are satisfied, there exists a curve $f \in F_K^R \cap D_K$ that is essential on $\partial(B_R) - R$ and such that $d(\mathcal{V}_R, f) \leq 1 - \chi(F_K^R)$. Since both F_K^R and F_K^Q are planar surfaces containing no disk or sphere components, $\chi(F_K^R) \leq 0$ and

$\chi(F_K^Q) \leq 0$. As $\chi(F_K) = \chi(F_K^R) + \chi(F_K^Q)$, $\chi(F_K^R) \leq 0$ and $\chi(F_K^Q) \leq 0$, it follows that $1 - \chi(F_K) \geq 1 - \chi(F_K^R)$ and $d(\mathcal{V}_R, f) \leq 1 - \chi(F_K)$. Since $d(f, \partial D) \leq 1$, we conclude that $d(\partial D, \mathcal{V}_R) \leq 2 - \chi(F_K)$. \square

7 Constructing the example

We will now construct a knot K as in Figure 15. We begin with the schematic in that figure and we substitute each of the balls B_1, \dots, B_4 with tangles satisfying particular properties. In the schematic S_1, \dots, S_4 are punctured spheres, B_1 and B_2 are the disjoint balls bounded by S_1 and S_2 containing tangles $\mathcal{T}_1 = (B_1, T_1)$ and $\mathcal{T}_2 = (B_2, T_2)$ respectively. The unique strand of K that connects S_1 and S_2 but is disjoint from S_4 will be labeled ϵ . The sphere S_4 bounds a ball B_T disjoint from B_1 and B_2 containing a 3–strand tangle $\mathcal{T} = (B_T, T)$. Thus, the knot K is naturally decomposed into three tangles: the two 2–strand tangles \mathcal{T}_1 and \mathcal{T}_2 and the 3–strand tangle \mathcal{T} .




Figure 15

By Corollary 4.6, we may assume that $d(\Sigma_1, T_1)$ is arbitrarily high and \mathcal{T}_1 does not have any closed components. In our construction, we will require that $d(\Sigma_1, T_1) \geq 25$. The tangle (B_2, T_2) will be identical to the tangle (B_1, T_1) .

The tangle \mathcal{T} is constructed as follows. Let each of \mathcal{R} and \mathcal{Q} be a 3–strand rational tangle. Let \mathcal{V}_R (respectively \mathcal{V}_Q) be the set of all essential simple closed curves in $\partial(B_R)$ (respectively $\partial(B_Q)$) that bound disks in $B_R - K$ (respectively $B_Q - K$). By Corollary 4.4, there exist equatorial curves γ_R and γ_Q in $(\partial B_R)_K$ and $(\partial B_Q)_K$

respectively such that $d(\gamma_{\mathcal{R}}, \nu_{\mathcal{R}}) \geq 25$ and $d(\gamma_{\mathcal{Q}}, \nu_{\mathcal{Q}}) \geq 25$. By Theorem 5.6, we may simultaneously require that no 2–strand equatorial subpair of $(\mathcal{R}, \nu_{\mathcal{R}})$ or $(\mathcal{Q}, \nu_{\mathcal{Q}})$ is the unpair. Let \mathcal{T} be the tangle obtained from \mathcal{R} and \mathcal{Q} as in Definition 6.1. We may assume that \mathcal{T} is symmetric with respect to the disk D .

Theorem 7.1 *We may assume that K is a knot.*

Proof Consider K to be constructed from the two 2–strand tangles contained in the two balls bounded by S_1 . By our construction, neither of these tangles has any closed components. Note that we may use any automorphism of the 4–punctured sphere to create K . In particular, we can choose a homomorphism that ensures that for each of the strands in B_1 the two endpoints are glued to endpoints of different strands in $S^3 - B_1$, thus, creating a knot. \square

8 Properties of K

In this section, we establish some of the properties of K . For the rest of this paper all surfaces will be punctured spheres so to avoid multiple subscript we will drop the subscript K , ie a surface F will always be punctured unless otherwise specified and will not be denoted by F_K .

The first property of K is based on an easy computation for the schematic in Figure 15 using the equation provided in Lemma 2.1.

Property 8.1 $w(K) \leq 134$.

Property 8.2 *Consider the 4–strand tangle contained between S_1 and S_3 . If T is any sphere separating S_1 from S_3 so that any maxima of the tangle are above T and any minima are below it, then T intersects the tangle in at least 8 points.*

Proof The tangle contained between S_1 and S_3 consists of the tangle \mathcal{R} together with an additional unknotted strand. By construction no equatorial subpair of \mathcal{R} is the unpair. Thus at least 2 of the three strands in \mathcal{R} must intersect T in at least 3 points each. The other two strands have to intersect T in at least one point each as T separates the endpoints of these strands. Thus T intersects the tangle in at least 8 points as required. \square

Property 8.3 *The spheres S_1 , S_2 and S_3 are all distinct.*

Proof Suppose first that S_1 is isotopic to S_2 . Since S_1 and S_2 are disjoint embedded 2–spheres in S^3 , the complementary components of $S_1 \cup S_2$ consist of two open 3–balls and one S^2 cross open interval. Let N be the closure of the complementary component homeomorphic to S^2 cross open interval. Thus, N has boundary $S_1 \cup S_2$, $N \cap K$ is a collection of vertical strands and $S_4 \subset N$. Hence, S_4 is visibly compressible into B_T . However, this is not possible, by Lemma 6.13, since we have assumed $d(\partial D, \mathcal{V}_R) \geq 25$ and $d(\partial D, \mathcal{V}_Q) \geq 25$.

Suppose then that S_1 is isotopic to S_3 . Then a copy of S_1 will intersect the tangle contained between S_1 and S_3 in only 4 points contradicting Property 8.2. \square

Property 8.4 *Any bridge surface for the tangle (B_1, T_1) meets T_1 in at least 10 points, and similarly for (B_2, T_2) .*

Proof By construction, (B_1, T_1) has a bridge surface Σ_1 of distance 25. Suppose Σ' is another bridge surface for (B_1, T_1) . By Theorem 4.2, with B_1 playing the roles of both M and N , it follows that either $\chi(\Sigma') \leq -23$ or, after some c –compressions, Σ' is parallel to Σ_1 . In either case, T_1 intersects Σ' at least as many times as it intersects Σ_1 , ie at least 10 times. \square

Property 8.5 *Both S_1 and S_2 are c –incompressible in $S^3 - K$.*

Proof In search of contradiction, suppose that one of S_1 or S_2 is c –compressible with c –disk Δ' . By taking Δ to be the disk bounded by an innermost curve of $\Delta' \cap (S_1 \cup S_2)$ in Δ' , we can assume that Δ is a c –disk for S_1 and the interior of Δ is disjoint from both S_1 and S_2 . The curve $\partial\Delta$ separates S_1 into two disks D_1 and D_2 .

If Δ is contained in B_1 , then, up to relabeling D_1 and D_2 , $\Delta \cup D_1$ is a 2–punctured sphere in B_1 . If $\Delta \cup D_1$ is an inessential 2–punctured sphere, then Δ is boundary parallel, contradicting the fact that Δ is a c –disk, or (B_1, T_1) is a rational tangle, a contradiction to Property 8.4. Hence, we can assume that $\Delta \cup D_1$ is an essential 2–punctured sphere. Maximally compress and cut-compress $\Delta \cup D_1$ in B_1 and let F be one of the resulting components. Note that F is an essential 2–punctured sphere. By Theorem 4.3, either F can be isotoped to be disjoint from Σ_1 , Σ_1 has four punctures or $25 \leq d(\Sigma_1) \leq 2 - \chi(F) = 2 - \chi(\Delta \cup D_1)$. It is easy to show that there are no essential surfaces in the complement of a tangle that are disjoint from its bridge surface. By the construction of B_1 , Σ_1 has 10 punctures and therefore we conclude that $\chi(\Delta \cup D_1) = \chi(F) \leq -23$. This contradicts the fact that F is an essential 2–punctured sphere.

Suppose Δ is not contained in B_1 or B_2 . Let ϵ be the arc in $K - (S_1 \cup S_2)$ that is disjoint from B_T and connects S_1 to S_2 . As S_4 is isotopic to the boundary of the

neighborhood of $S_1 \cup S_2 \cup \epsilon$, it follows that ∂D is isotopic to a meridional curve for ϵ . Additionally, if Δ is contained strictly between S_1 and S_2 and is disjoint from ϵ , then Δ is a c -disk for S_4 with boundary disjoint from ∂D . By Lemma 6.10, D is c -incompressible, so we can assume that Δ is disjoint from D . Since Δ is disjoint from D then Δ is a c -disk for $\partial(B_R) - R$. If Δ is a compressing disk, then $d(\partial D, \mathcal{V}_R) \leq 1$, a contradiction to the construction of the tangle \mathcal{T} . If Δ is a cut-disk and E is the bridge disk for the strand of R that meets Δ , then, possibly after an isotopy of E , we can assume that Δ meets E in a single arc. The boundary of a regular neighborhood of $E \cup \Delta$ contains a compressing disk, Δ'' , for $\partial(B_R) - R$ that is disjoint from Δ . Since $\partial\Delta$ is disjoint from ∂D and $\partial(\Delta'')$ is disjoint from $\partial\Delta$ then $d(\partial D, \mathcal{V}_R) \leq 2$. This is a contradiction to the construction of \mathcal{T} .

If Δ is contained strictly between S_1 and S_2 and meets ϵ , then we can assume that D_1 meets K in one point and D_2 meets K in three points. If ϵ has an endpoint in D_1 , then $D_1 \cup \Delta$ bounds a 3-ball containing a unknotted arc, since ϵ is unknotted. Hence, Δ is boundary parallel, a contradiction. Therefore, we can assume that D_1 is disjoint from ϵ . The twice punctured sphere $\Delta \cup D_1$ is disjoint from $S_1 \cup S_2$ and meets ϵ in a single point. Therefore, after isotoping S_4 to be the boundary of a regular neighborhood of $S_1 \cup S_2 \cup \epsilon$, $(\Delta \cup D_1) \cap B_{\mathcal{T}}$ is a cut-disk for $\partial(B_{\mathcal{T}}) - T$ with boundary parallel to a meridional curve of ϵ . As noted before, such a meridional curve is isotopic to ∂D . As argued above, this implies $d(\partial D, \mathcal{V}_R) \leq 2$, a contradiction to the construction of \mathcal{T} . □

Property 8.6 *Let F be a connected, planar, meridional, nonboundary parallel, c -incompressible, planar surface in $S^3 - \eta(K)$. Then one of the following holds:*

- (1) F can be isotoped to be disjoint from $B_{\mathcal{T}}$.
- (2) F is isotopic to S_3 .
- (3) F has at least 14 punctures.

Proof Isotope F so that $F \cap (S_1 \cup S_2)$ is minimal. Suppose $F \cap S_1$ is nonempty. Since S_1 is c -incompressible, by Property 8.5, and K is a knot, then minimality of $F \cap (S_1 \cup S_2)$ implies $F - S_1$ contains no disk components. Let F^1 be a component of $B_1 \cap F$. By minimality of $|F \cap (S_1 \cup S_2)|$, F^1 is not isotopic to a subsurface of S_1 . Since F is c -incompressible, so is F^1 . So, F^1 can not be isotoped to be disjoint from Σ_1 . Since F^1 is c -incompressible and F^1 is not isotopic to a subsurface of S_1 , then F^1 is not boundary parallel in $B_1 - \eta(T_1)$. By Theorem 4.3, F^1 , and thus F , has at least 14 punctures. Hence, we can assume that F is disjoint from S_1 and S_2 .

Let M be the $S^2 \times I$ region in S^3 with boundary S_1 and S_2 . Since F is disjoint from S_1 and S_2 , F is contained in the interior of M . Recall that ϵ is the strand of

$K \cap M$ that is disjoint from B_T . Let η be a small open neighborhood of $S_1 \cup S_2 \cup \epsilon$ in M . By transversality, F meets η in a possibly empty collection of parallel, disjoint, 1-punctured disks. Recall D from Definition 6.1. Since B_T is isotopic to $M - \eta$ then F can be isotoped to meet $\partial(B_T)$ in a collection of curves parallel to ∂D . Since F is planar and $F \cap \eta(\epsilon)$ is a collection of once-punctured disks, $F^T = F \cap B_T$ is connected. If F^T is a disk, then F is a 1-punctured sphere in S^3 , a contradiction. Since F is c -incompressible, so is F^T . By Theorem 6.17, F^T is isotopic to one of ten surfaces. Conclusion (1) and (2) cannot occur since F was assumed to be essential. If conclusion (3), (4), (5) or (7) holds, then we can isotope F to be disjoint from B_T . If conclusion (6) holds, F is the boundary union of D and a 1-punctured disk that meets ϵ . In this case, F is isotopic to S_3 . If conclusion (8) or (9) holds then, since $d(\partial D, \mathcal{V}_R) \geq 25$ and $d(\partial D, \mathcal{V}_Q) \geq 25$, we can conclude that F has at least 14 punctures. \square

We will be particularly interested in surfaces obtained by tubing two spheres with a tube that runs along an arc of the knot connecting these spheres. The following definition describes this construction.

Definition 8.7 Let F and G be disjoint embedded spheres in S^3 with the property that $F \cap K \neq \emptyset$ and $G \cap K \neq \emptyset$ and let α be the closure of a component of $K - (F \cup G)$ with an endpoint in each of F and G . Then the boundary of a regular neighborhood of $F \cup \alpha \cup G$ has three components. Let $F \#_\alpha G$ be the component that is not parallel to F or to G in the complement of the other two components.

Equivalently, $F \#_\alpha G$ is the embedded connected sum of F and G obtained by replacing a neighborhood of $\partial(\alpha)$ in F and G with an annulus that runs parallel to α .

Property 8.8 Let $B_{i,j}$ be the ball bounded by $S_i \#_\alpha S_j$ with $i, j \in \{1, 2, 3\}$ and $i \neq j$ that is disjoint from B_1 and B_2 and let $T_{i,j} = K \cap B_{i,j}$. If $S_i \#_\alpha S_j$ is incompressible, then any bridge sphere $\Sigma_{i,j}$ for $(B_{i,j}, T_{i,j})$ has at least 10 punctures.

In the special case where $i = 1, j = 2$ and S_1 is tubed to S_2 along a strand α that intersects S_4 , then the bridge sphere $\Sigma_{1,2}$ for $(B_{1,2}, T_{1,2})$ has at least 14 punctures.

Proof There are three cases to consider.

Case 1 Suppose $S = S_1 \#_\epsilon S_2$. In this case, S is isotopic to S_4 and the tangle under consideration is the tangle \mathcal{T} . This tangle contains three arcs α, β and γ . By construction, ∂D is a simple closed curve in S_4 and each arc of T has an endpoint in each of the two components of $S_4 - \partial D$. For any two arcs in T , say α and β , define $K_{\alpha,\beta}$ to be the link obtained by connecting the endpoints of α and β via two arcs in

S_4 so that each of these arcs is disjoint from ∂D . Under such restrictions, the link type of $K_{\alpha,\beta}$ is well defined. As illustrated in Figure 16, $K_{\alpha,\beta}$ can also be constructed by taking the connected sum of some numerator closure of an equatorial subpair of $(\mathcal{R}, \partial D)$ with some numerator closure of an equatorial subpair of $(\mathcal{Q}, \partial D)$ neither of which is the unknot by construction. Since $K_{\alpha,\beta}$ is the connected sum of two links, neither of which is the unknot, then the bridge number of $K_{\alpha,\beta}$ is at least 3, by [16]. Hence, one of α or β meets the bridge sphere of T in at least 4 points, the other meets the sphere in at least 2 points. By examining $K_{\alpha,\gamma}$ and $K_{\beta,\gamma}$ we conclude that one of α or γ meets the bridge sphere in at least 4 points and one of β or γ meets the bridge sphere in at least 4 points. Hence, two of the arcs α , β , or γ meet the bridge sphere in at least 4 points and the third meets it in at least 2 points. Thus, the bridge sphere for \mathcal{T} has at least 10 punctures.




Figure 16

Case 2 Suppose S is isotopic to $S_1 \#_{\alpha} S_2$ where α is one of the three strands connecting S_1 to S_2 that intersects S_4 and therefore passes through both B_R and B_Q . Let γ , δ be the other two strands of T and, thus, ϵ , γ and δ are the three strands of $T_{1,2}$. By connecting the endpoints of γ with an arc contained in S , we create a knot, K_{γ} . The knot-type of K_{γ} is well-defined and independent of how we connect the points in $\partial\gamma$. We define K_{δ} and K_{ϵ} similarly. As illustrated in Figure 17, K_{γ} can also be constructed by taking the connected sum of some numerator closure of an equatorial subpair of $(\mathcal{R}, \partial D)$ with some numerator closure of an equatorial subpair of $(\mathcal{Q}, \partial D)$. As before, both of these are knotted by construction. Since K_{γ} is the connected sum of two knots, then the bridge number of K_{γ} is at least 3, by [16]. Hence γ meets the bridge sphere in at least 6 points. A similar argument reveals that δ meets the bridge sphere in at least 6 points. Additionally, ϵ must meet this bridge sphere in at least 2 points. Hence, the bridge sphere for $\mathcal{T}_{1,2}$ must have at least 14 punctures.




Figure 17

Case 3 Suppose S is isotopic to $S_1 \#_\alpha S_3$ or $S_2 \#_\alpha S_3$. Without loss of generality, we may assume that S is isotopic to $S_1 \#_\alpha S_3$. There are four components of $K - (S_1 \cup S_3)$ that have a boundary point on each of S_1 and S_3 . One of these components is contained in ϵ , so we denote it by ϵ . Call the other three components β, γ, δ . If $\alpha = \epsilon$, then $S_1 \#_\alpha S_3$ is isotopic to $\partial(B_R)$, and therefore compressible. Hence, we can assume that α passes through B_R . Without loss of generality, assume $\alpha = \beta$. In this case, the strands γ, δ and ϵ become the three strands of $T_{1,3}$. By connecting the endpoints of γ with an arc contained in S , we create a knot, K_γ . The knot-type of K_γ is well-defined independent of how we connect the points in $\partial\gamma$. We define K_δ and K_ϵ similarly. As illustrated in Figure 18, K_γ can also be constructed by taking a numerator closure of an equatorial subpair of $(\mathcal{R}, \partial D)$, which is knotted by construction. Since K_γ is knotted, the bridge number of K_γ is at least 2. Hence, γ meets any bridge sphere in at least 4 points. A similar argument reveals that δ meets any bridge sphere in at least 4 points. Additionally, ϵ must meet any bridge sphere in at least 2 points. Hence, the bridge sphere for $\mathcal{T}_{1,3}$ must have at least 10 punctures. \square

Property 8.9 Let S be the tangle $(S^3 - (B_1 \cup B_2), K \cap (S^3 - (B_1 \cup B_2)))$. Then any bridge sphere for S must have at least 10 punctures.

Proof Let M be the embedded copy of $S^2 \times I$ in S^3 with boundary $S_1 \cup S_2$. The knot K meets M in four arcs α, β, γ and ϵ , where ϵ is the unique arc disjoint from S_4 . For any two arcs in M , say α and β , define $K_{\alpha,\beta}$ to be the knot obtained by connecting the endpoints of α and β via an arc in S_1 and an arc in S_2 . The knot




Figure 18

type of $K_{\alpha,\beta}$ is well defined under this construction. As illustrated in Figure 19, $K_{\alpha,\beta}$ can also be constructed by taking the connected sum of some numerator closure of an equatorial subpair of $(\mathcal{R}, \partial D)$ with some numerator closure of an equatorial subpair of $(\mathcal{Q}, \partial D)$, both of which are knotted by construction. Since $K_{\alpha,\beta}$ is the connected sum of two knots, then the bridge number of $K_{\alpha,\beta}$ is at least 3, by [16]. Similarly, each of $K_{\alpha,\gamma}$ and $K_{\beta,\gamma}$ has bridge number at least 3. Since $K_{\alpha,\beta} \cup K_{\alpha,\gamma} \cup K_{\beta,\gamma}$ meets any bridge sphere in at least 18 points, any bridge sphere for \mathcal{S} meets the union of α , β and γ in at least 9 points. Additionally, ϵ must meet any bridge sphere for \mathcal{S} in at least one point. Thus, any bridge sphere for \mathcal{S} must have at least 10 punctures. \square




Figure 19

Property 8.10 *Let B' and B'' be the two balls bounded by S_3 and let $T' = K \cap B'$ and $T'' = K \cap B''$. Then any bridge surface for each of the tangles (B', T') and (B'', T'') has at least 12 punctures.*

Proof Let Σ' be a bridge surface for the tangle (B', T') . Without loss of generality, suppose B_1 is contained in B' . By Theorem 4.2, either $d(\Sigma_1) \leq 2 - \chi(\Sigma')$, or $\Sigma' \cap B_1$ is a sphere parallel to Σ_1 with tubes attached. In the first case, Σ' has at least 26 punctures. In the second case, if no tubes are attached, then Σ' is isotopic to Σ_1 and so, by Property 8.5, S_1 is an essential surface contained in B' but disjoint from Σ' which is not possible since S_1 is not isotopic to S_3 . Therefore, to recover Σ' from Σ_1 at least one tube must be attached. Since each additional tube adds at least two punctures, Σ' has at least 12 punctures. An analogous argument with S_2 playing the role of S_1 shows the result for (B'', T'') . \square

9 Meridional essential spheres in the complement of K

In this section, we will classify all essential spheres in the complement of K with fewer than 14 punctures.

Proposition 9.1 *Any meridional essential sphere that is embedded in one of B_1 , B_2 or B_T has at least 14 punctures.*

Proof Suppose F is an essential meridional sphere and $F \subset B_1$. Maximally cut-compress F in B_1 and let F' be one of the resulting components. Note that F' is incompressible, $\chi(F') \geq \chi(F)$ and the number of punctures of F' is less than or equal to that of F since F is a meridional essential sphere in a knot complement. By Theorem 4.3, it follows that either F' can be isotoped to be disjoint from Σ_1 , Σ_1 has four punctures or $25 \leq d(\Sigma_1) \leq 2 - \chi(F') \leq 2 - \chi(F)$. It is easy to show that there are no essential surfaces in the complement of a tangle that are disjoint from its bridge sphere. By the construction of B_1 , Σ_1 has 10 punctures and therefore we conclude that $\chi(F) = \chi(F') \leq -23$. It follows that F must have at least 25 punctures. A symmetric argument produces the desired result if $F \subset B_2$.

Suppose F is an essential sphere in $B_T - T$. After maximally cut-compressing F in $B_T - T$, each component is a meridional, c -incompressible sphere in $B_T - T$. Let F' be one such component. By Theorem 6.17, we conclude that $d(\partial(D), \mathcal{V}_R) \leq 2 - \chi(F')$, or $d(\partial(D), \mathcal{V}_Q) \leq 2 - \chi(F')$, or F' is isotopic to $\partial(B_T) - T$. In the first two cases, since $d(\partial(D), \mathcal{V}_R) \geq 25$ and $d(\partial(D), \mathcal{V}_Q) \geq 25$, we conclude that F' , and thus F , has at least 14 punctures. Hence, we can assume that, after maximally cut-compressing F in B_T , every component is isotopic to $\partial(B_T) - T$. If there is more than one such component, then reversing the final cut-compression by attaching a tube between two parallel copies of $\partial(B_T) - T$ results in a compressible surface, a contradiction to the incompressibility of F . Hence, F is isotopic to $\partial(B_T) - T$, a contradiction to F being essential. \square

We associate to the given projection of K a graph Γ embedded in S^2 , the projection sphere of Figure 15, where each vertex corresponds to one of the four 3–balls, B_1, \dots, B_4 , and the edges correspond to strands of the knot connecting these 3–balls. We may assume that K lies in an arbitrarily small neighborhood of S^2 . Note that a simple closed curve in S^2 that is disjoint from Γ bounds a disk in S^2 that is disjoint from Γ . We will use this graph on several occasions.

Lemma 9.2 *Suppose F is an essential meridional sphere disjoint from B_i for $i = 1, \dots, 4$, that is embedded so that $|F \cap S^2|$ is minimal. Then $F \cap S^2$ is a single simple closed curve τ . Moreover there are no bigons in S^2 whose boundary is the endpoint union of a segment of an edge of Γ and a segment of τ .*

Proof As F is disjoint from all B_i , it intersects the 2–sphere containing Γ in circles disjoint from the vertices of the graph. If any such circle bounds a disk in the 2–sphere disjoint from the graph, it can be removed by an isotopy using the incompressibility of F , contradicting the assumption that $|F \cap S^2|$ is minimal.

Suppose α and β are two simple closed curves in $F \cap S^2$. By the above, both curves contain points of intersection between F and K . Consider a product neighborhood $S^2 \times I$ of S^2 so that $\partial(S^2 \times I)$ is disjoint from K and $F \cap (S^2 \times I)$ is a union of vertical fibers. Then α is parallel to two curves in the boundary of the product, one lying in $F \cap (S^2 \times \{0\})$ and the other in $F \cap (S^2 \times \{1\})$. Since F is a sphere, one of these curves, α' , separates α and β in F . Let D be the disk in $\partial(S^2 \times I)$ that α' bounds and note that D is disjoint from K . As F is incompressible, after possibly an isotopy of F , we may assume that $F \cap \text{int}(D) = \emptyset$. As $\partial D = \alpha'$ separates the punctures of F lying in α from the punctures lying in β , the disk D is a compressing disk for F , contradicting the hypothesis. Hence, we can assume $F \cap S^2$ is a single simple closed curve, τ .

Suppose a subarc of $\tau = S^2 \cap F$ cobounds with some edge of Γ a bigon. Without loss of generality, we may assume that the interior of the bigon is disjoint from Γ and F . A regular neighborhood of this bigon contains a compressing disk for F unless F is a twice punctured sphere parallel to a segment of the edge. As F is essential, no such bigons can exist. \square

Proposition 9.3 *Any essential meridional sphere, F , that is disjoint from B_i for $i = 1, \dots, 4$ is parallel to one of S_1, S_2, S_3 or S_4 . In particular, any such surface has at most 6 punctures.*

Proof We will continue using the terminology developed in the proof of Lemma 9.2. By that lemma, it follows that if $\tau = S^2 \cap F$ intersects any edge in a triple of parallel

edges of Γ , it must intersect all three of them. Thus, we can replace these triples with a single edge and assign it a weight of 3. The projection of K is then modeled by a cycle graph on 4 vertices, C_4 , where three of the edges have weight 3 and one has weight 1. The number of punctures of F lying in τ is the sum of the weights of the edges τ intersects. By the Jordan–Brouwer theorem, τ has an even number of intersections with the edges of Γ and, by Lemma 9.2, consecutive edges that τ intersects must be distinct.

Let E be one of the two disks C_4 bounds in S^2 and let l be an outermost arc of intersection of τ and E . Let p_1 and p_2 be the two endpoints of l . Choose p'_1 and p'_2 to be points in τ that are close to p_1 and p_2 respectively but not in l so that the segment of $l' = \tau - (p'_1 \cup p'_2)$ containing l only intersects C_4 in p_1 and p_2 . Let g be any embedded path in $S^2 - (\tau \cup C_4)$ from p'_1 to p'_2 . Consider a small regular neighborhood $S^2 \times I$ of S^2 . Then there is a disk with boundary $(l' \times \{0\}) \cup (p'_1 \times I) \cup (p'_2 \times I) \cup (l' \times \{1\})$ that contains two copies of g . This disk would be a compressing disk for F unless $p_1 \cup p_2 = \tau \cap C_4$. Therefore, we conclude that τ intersects exactly two of the edges of C_4 and intersects each of those exactly once. The result follows by considering all possible pairs of edges. \square

Proposition 9.4 *There are no essential twice-punctured spheres that are disjoint from all of S_1 , S_2 and S_3 .*

Proof Let F be an essential twice punctured sphere in the complement of K disjoint from S_1 , S_2 and S_3 . By Proposition 9.1, F is not embedded in B_1 or B_2 . Without loss of generality, we may assume that F is on the same side of S_3 as B_1 but outside of B_1 . In this case, it is easy to see that F can be isotoped to lie entirely in B_T , contradicting Lemma 6.12. \square

Proposition 9.5 *The spheres S_1 , S_2 and S_3 are c -essential.*

Proof The spheres S_1 and S_2 are c -essential by Property 8.5. Suppose S_3 has a c -disk D^c . We may assume that this c -disk is disjoint from $S_1 \cup S_2$. Let Δ be one of the two punctured disks ∂D^c bounds on S_3 . Then $F = D^c \cup \Delta$ is a sphere that is disjoint from all S_i , $i = 1, 2, 3$. Note that we may choose Δ so it has at most 2-punctures. As D^c has at most one puncture and every sphere has an even number of punctures, F has at most two punctures. As K is not a split link, F is an essential twice punctured sphere disjoint from S_1 , S_2 and S_3 , contradicting Proposition 9.4. \square

Corollary 9.6 *K is a prime knot.*

Proof Suppose F is a decomposing sphere for K . As S_1 , S_2 and S_3 are c -essential by Proposition 9.5, we can find such a sphere that is disjoint from $S_1 \cup S_2 \cup S_3$, contradicting Proposition 9.4 \square

Corollary 9.7 *Every essential 4-punctured sphere is c -essential.*

Property 9.8 *Let $B' = S^3 - B_1$ and $B'' = S^3 - B_2$. Any bridge surface for the tangles $\mathcal{T}' = (B', K - T_1)$ and $\mathcal{T}'' = (B'', K - T_2)$ has at least 14 punctures.*

Proof We show the result for \mathcal{T}' , but note that a completely analogous argument proves the result for \mathcal{T}'' . Let Σ be a bridge surface for \mathcal{T}' . By Theorem 4.2, either $d(\Sigma_2) \leq 2 - \chi(\Sigma)$, or $\Sigma \cap B_2$ is a sphere parallel to Σ_2 with tubes attached. In the first case, Σ has at least 26 punctures, so suppose we are in the second case. If no tubes are attached, ie if Σ is parallel to Σ_2 , then S_2 is an essential sphere that is disjoint from the bridge surface of the tangle T' which is not possible since S_2 is not isotopic to S_3 . If more than one tube is attached, then $w(\Sigma) \geq 14$. Similarly, if some tube corresponds to a compressing disk that decomposes Σ into a component parallel to Σ_2 and a sphere with more than two punctures or if some tube corresponds to a cut-disk that decomposes Σ into a component parallel to Σ_2 and a sphere with more than four punctures, then $w(\Sigma) \geq 14$. Therefore, we may assume that Σ is parallel to Σ_2 with exactly one tube attached. If this tube corresponds to a compressing disk, then it connects Σ_2 to a twice punctured sphere and if it corresponds to a cut-disk, then it connects Σ_2 to a four times punctured sphere.

Claim *We may assume the tube corresponds to a compressing disk.*

Proof of claim Suppose the tube corresponds to a cut-disk D^c . Then ∂D^c bounds a three punctured disk Δ in Σ . Consider the arc κ that has both of its endpoints in Δ . A bridge disk for κ can be isotoped to be disjoint from D^c using an innermost disk/outermost arc argument. Let E be the compressing disk for Σ obtained by taking the boundary of a regular neighborhood of the bridge disk for κ . Note that E is disjoint from D^c . Additionally, D^c , E and a once punctured annulus A in Σ cobound a twice punctured sphere. By Corollary 9.6, K and, thus, \mathcal{T}' is prime. As \mathcal{T}' is prime, it follows that $E \cup A$ is isotopic to D^c . Therefore, the surface F obtained by cut-compressing Σ along D^c that does not contain Δ is isotopic to the surface F' obtained from Σ by compressing it along E that contains A . As F is parallel to Σ_2 , so is F' . Therefore, we may replace the tube corresponding to D^c with a tube corresponding to E . \square

By the claim, we may assume that Σ can be recovered by tubing Σ_2 to a twice punctured sphere Q along a tube that is disjoint from the knot. There are 4 strands of K that have one endpoint in S_1 and the other in Σ_2 . By Property 8.5, S_1 and S_2 are c -essential, so Q can be isotoped to be disjoint from both S_1 and S_2 . Hence, the sphere Q intersects at most one of these strands, so there are at least three strands that have one endpoint in S_1 and one point in Σ . At least two of these strands, α and β , intersect B_T . As Σ is a bridge surface, α and β are both vertical and, therefore, parallel to each other. Let R be the rectangle they cobound. Connecting $\alpha \cap B_T$ and $\beta \cap B_T$ along the two arcs $R \cap S_4$ results in the unknot. However, this is the knot $K_{\alpha,\beta}$ described in Case 1 of the proof of Property 8.8. By the argument there, the bridge number of $K_{\alpha,\beta}$ is at least three, leading to a contradiction. \square

Proposition 9.9 *Any c -essential meridional sphere with fewer than 14 punctures has exactly 4 punctures and is parallel to one of S_1, S_2 or S_3 .*

Proof Let F be a c -essential sphere with fewer than 14 punctures. Isotope it to intersect $B_1 \cup B_2 \cup B_T$ minimally and suppose first that $F \cap B_1 \neq \emptyset$. Note that no component of $F - S_1$ can be a c -disk as S_1 is c -incompressible. It follows that, if F' is any component of $F \cap B_1$, $\chi(F') \geq \chi(F)$. Furthermore, F' cannot be parallel to $\partial(B_1 - \eta(K))$ as in this case either $|F \cap B_1|$ can be reduced or F has a cut-disk (this situation occurs if $F \cap B_1$ has a component that is an annulus parallel to a segment of $B_1 \cap K$). Recall that Σ_1 has 10 punctures and there are no essential surfaces in the complement of the tangle that are disjoint from the bridge sphere of the tangle. By Theorem 4.3, it follows that $d(\Sigma_1, K \cap B_1) \leq 2 - \chi(F')$. As $d(\Sigma_1, K \cap B_1) \geq 25$, this implies that $\chi(F') \leq -23$ and, therefore, $\chi(F) \leq -23$. This contradicts the hypothesis that F has at most 14 punctures and, therefore, it follows that $F \cap B_1 = \emptyset$. Similarly, $F \cap B_2$ must also be empty.

By Property 8.6, it follows that if $F \cap B_T \neq \emptyset$, then F is S_3 or has at least 14 punctures. Hence, we can assume that F is disjoint from B_i for $i = 1, \dots, 4$. By Proposition 9.3, F is parallel to one of S_1, S_2, S_3 or S_4 . However, S_4 is not c -essential. Thus, F is parallel to one of S_1, S_2 or S_3 . \square

Proposition 9.10 *The only incompressible 6-punctured spheres are the ones obtained by tubing two essential 4-punctured spheres that are not mutually parallel along a strand of K . In particular, if P is an incompressible 6-punctured sphere, then its cut-disk, B_1 , and B_2 are on the same side of P .*

Proof Suppose G is an incompressible 6-punctured sphere. By Proposition 9.9, it follows that G is cut-compressible. As K is prime, cut-compressing a 6-punctured sphere

results in a pair of 4-punctured spheres. If the original sphere was incompressible, so are the two 4-punctured ones.

By Corollary 9.7, all essential 4-punctured spheres are c -essential, and all c -essential 4-punctured spheres in the complement of K are parallel to one of S_1 , S_2 or S_3 by Proposition 9.9. If two parallel copies of some 4-punctured sphere are tubed together along a single strand of the knot, the resulting 6-punctured sphere is always compressible. Therefore, any incompressible 6-punctured sphere is the result of tubing together two of S_1 , S_2 or S_3 . \square

Note that S_4 can be obtained by tubing S_1 to S_2 .

10 Bridge number

In this section, we will show that thin and bridge position for K do not coincide.

Lemma 10.1 *Suppose Σ is a minimal bridge sphere for K , and C_1 and C_2 are c -disks for Σ on opposite sides of Σ so that $\partial(C_1) \cap \partial(C_2) = \emptyset$. Then bridge position for K is not thin position.*

Proof Let Δ_1 and Δ_2 be the disjoint disks ∂C_1 and ∂C_2 bound in Σ . The disk Δ_1 must have at least two punctures as ∂C_1 is essential in Σ . In particular, there is a strand κ_1 above Σ that has both of its endpoint in Δ_1 and is disjoint from C_1 . As Σ is a bridge sphere, κ_1 has a bridge disk E_1 . Using the fact that there are no spheres that intersect the knot in exactly one point and an innermost disk argument, we can choose E_1 so that it intersects C_1 only in arcs. If there are two distinct outermost arcs of $E_1 \cap C_1$ in C_1 , then one of these arcs bounds a disk in C_1 that is disjoint from E_1 and does not contain the puncture. If there is a unique outermost arc of $E_1 \cap C_1$ in C_1 , then one of the two disks this arc bounds in C_1 is disjoint from E_1 does not contain the puncture. In either case, we can boundary compress E_1 along the disk we just found in C_1 to find a new bridge disk for κ_1 that meets C_1 in strictly fewer arcs. Repeat this process to produce a bridge disk E_1^* for κ_1 that is disjoint from C_1 . In particular, $E_1^* \cap \Sigma \subset \Delta_1$. Similarly, there is a bridge disk E_2 for some strand κ_2 below Σ so that $E_2 \cap \Sigma \subset \Delta_2$. This pair of disks allows us to push the maximum of κ_1 below the minimum of κ_2 , thus decreasing the width of K . Therefore, bridge position of K is not thin position. \square

We will rely heavily on the terminology introduced in Section 3 and illustrated in Figure 3. In addition, we need the following definition first introduced in [1].

Definition 10.2 Let S be a 2–sphere embedded in S^3 so that S meets K transversely in exactly 4 points. S is *worm-like* if, for every saddle $\sigma = s_1^\sigma \vee s_2^\sigma$, each of s_1^σ and s_2^σ cuts S into two twice punctured disks and every saddle in S is nested with respect to the same side of S .

Theorem 10.3 [1, Theorem A] *If S is a c –incompressible 4–punctured sphere and bridge position for K is thin position, then there is an isotopy of S and K resulting in $h|_K$ having $b(K)$ maxima and S being worm-like.*

Theorem 10.4 *Suppose S is a c –incompressible 4–punctured sphere in S^3 bounding 3–balls B_1 and B_2 on opposite sides. Additionally, suppose bridge position for K is thin position. Then, up to relabeling B_1 and B_2 , there is a minimal bridge sphere, Σ , such $B_1 \cap \Sigma$ is a collection of punctured disks and $S - \Sigma$ is a collection of annuli and 2–punctured disks.*

Proof By Theorem 10.3, we can assume that S is worm-like and $h|_K$ has $b(K)$ maxima. In particular, we can assume that all saddles in S are nested with respect to B_1 . Let E_1 and E_2 be the unique outermost disks of S . By definition of worm-like, E_1 and E_2 are 2–punctured disks. Since all saddles in S are nested with respect to B_1 , the interior of A_σ is disjoint from S for every saddle σ . In particular, A_σ is disjoint from A_τ whenever $\sigma \neq \tau$. If E_σ has a unique maximum, we can horizontally shrink and vertically lower A_σ so that, after this isotopy, A_σ is contained in an arbitrarily small neighborhood of the level sphere containing σ . If E_σ has a unique minimum, we can horizontally shrink and vertically raise A_σ so that after this isotopy A_σ is contained in an arbitrarily small neighborhood of the level sphere containing σ . By general position, we can assume that all saddles occur at distinct heights. Since A_σ is disjoint from A_τ whenever $\sigma \neq \tau$ and all saddles occur at distinct heights, we can isotope S and K so that $h(A_\sigma) \cap h(A_\tau) = \emptyset$ whenever $\sigma \neq \tau$. This isotopy does not change the number of critical points of $h|_K$, fixes the saddles of S and leaves invariant the number of saddles of S .

Suppose that, after isotopying all the A_σ to occur at distinct heights, there exists an A_σ with a unique minimum above an A_τ with a unique maximum. Each of $D_1^\sigma, D_2^\sigma, D_1^\tau, D_2^\tau$ meets K as otherwise S is compressible. Since each of E_σ and E_τ are disjoint from K , A_σ contains a minimum of K and A_τ contains a maximum of K . Since A_σ is contained completely above A_τ , there is a minimum of K above a maximum of K , a contradiction to the assumption that bridge position is thin position. Hence, we can assume that all A_σ with unique maxima lie above all A_σ with unique minima.

Fix σ and τ as the two unique outermost saddles of S . Suppose $E_1 = D_\sigma$ and suppose E_1 has a unique maximum. Let $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$.

The following claim is very similar to the claim in the proof of Lemma 3.8.

Claim 1 *After an isotopy that fixes S and preserves that number of maxima of h_K , we can assume that $h_{K \cap B_\sigma}$ has a local maximum at x_1 .*

Proof Suppose $h_{K \cap B_\sigma}$ has a local minimum at x_1 . Let y be the maximum of K that is nearest x_1 and inside B_σ . Such a y must exist since K does not meet E_1 above x_1 . Let α be the monotone subarc of K with boundary $x_1 \cup y$. The arc α is completely contained in B_σ . Let β be a monotone arc in E_1 with endpoints x_1 and z such that $h(z) = h(y)$. Let δ be a level arc disjoint from K and contained in B_σ connecting x to y . Let E be a vertical disk with boundary $\alpha \cup \beta \cup \delta$ that is embedded in B_σ . We can assume the interior of E meets K transversely in a collection of points k_1, \dots, k_n where $h(k_1) > h(k_2) > \dots > h(k_n)$. Let η_i be the arc corresponding to a small neighborhood of k_i in K for each i .

Replace η_1 with a monotone arc which starts at an end point of η_1 , runs parallel to E until it nearly reaches E_1 , travels along E_1 until it returns to the other side of E , travels parallel to E (now on the opposite side) and connects to the other end point of η_1 . The result is isotopic to K , does not change the number of maxima of $h|_K$ and reduces n . By induction on n , we may assume that $K \cap E = \emptyset$. Isotope α along E until it lies just outside of E_1 except where it intersects E_1 exactly at the point y . After a small tilt of K , we see that x_1 is now a local maximum of $h_{K \cap B_\sigma}$. \square

By a symmetric argument, we conclude that if E_1 has a unique minimum and $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$, then, after an isotopy that fixes S and preserves the number of maxima of h_K , we can assume that $h_{K \cap B_\sigma}$ has a local minimum at x_2 .

Suppose E_1 has a unique maximum, $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$ and there exists an A_ζ such that A_ζ has a unique minimum and $h(\zeta) > h(x_2)$. If x_2 is a maximum of $h_{B_\sigma \cap K}$ then σ is a removable saddle. By [1, Lemma 3.5], we can eliminate σ while preserving the number of maxima of h_K . Hence, we can assume x_2 is a minimum of $h_{B_\sigma \cap K}$. Since x_2 is a minimum of $h_{B_\sigma \cap K}$, then either there is a maximum of K inside B_σ or x_1 is connected to x_2 via a monotone subarc of K . In the later case, examine a level disk in B_σ with boundary in E_1 such that this disk is just below x_2 . If K meets this disk, then there is a maximum of K in B_σ . If K does not meet this disk then S is compressible, a contradiction. Hence, we can assume that there is a maximum of K in B_σ . As previously noted, A_ζ contains a minimum of K .

Since x_1 is a local maximum of $h_{B_\sigma \cap K}$, we can horizontally shrink and vertically lower the portion of B_σ above x_2 to within an arbitrarily small neighborhood of the

level sphere containing x_2 . Similarly, we can horizontally shrink and vertically raise A_ζ to within an arbitrarily small neighborhood of the level sphere containing ζ . These isotopies do not change the number of critical points of h_K , however they do raise a minimum above a maximum, a contradiction to the assumption that bridge position coincides with thin position.

By a similar argument, we can eliminate the following possibilities.

- (1) E_1 has a unique minimum, $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$ and there exists an A_κ such that A_κ has a unique maximum and $h(\kappa) < h(x_1)$.
- (2) E_2 has a unique maximum, $\{x_1, x_2\} = K \cap E_2$ such that $h(x_2) < h(x_1)$ and there exists an A_ζ such that A_ζ has a unique minimum and $h(\zeta) > h(x_2)$.
- (3) E_2 has a unique minimum, $\{x_1, x_2\} = K \cap E_2$ such that $h(x_2) < h(x_1)$ and there exists an A_κ such that A_κ has a unique maximum and $h(\kappa) < h(x_1)$.
- (4) E_1 has a unique minimum where $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$ and E_2 has a unique maximum where $\{y_1, y_2\} = K \cap E_2$ such that $h(y_2) < h(y_1)$ and $h(x_1) > h(y_2)$.

Suppose E_1 has a unique minimum where $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$ and E_2 has a unique maximum where $\{y_1, y_2\} = K \cap E_2$ such that $h(y_2) < h(y_1)$. Let $\{\kappa_1, \dots, \kappa_k\}$ be the set of all saddles such that A_{κ_i} contains a unique maximum and $\{\zeta_1, \dots, \zeta_s\}$ be the set of all saddle such that A_{ζ_i} contains a unique minimum. By the above eliminations, $\min(h(y_2), h(\kappa_1), \dots, h(\kappa_k)) > \max(h(x_1), h(\zeta_1), \dots, h(\zeta_s))$ and any level sphere with height strictly between these two values is a bridge sphere satisfying the conclusions of the theorem.

Suppose E_1 has a unique maximum where $\{x_1, x_2\} = K \cap E_1$ such that $h(x_2) < h(x_1)$ and E_2 has a unique maximum where $\{y_1, y_2\} = K \cap E_2$ such that $h(y_2) < h(y_1)$. Let $\{\kappa_1, \dots, \kappa_k\}$ be the set of all saddles such that A_{κ_i} contains a unique maximum and $\{\zeta_1, \dots, \zeta_s\}$ be the set of all saddle such that A_{ζ_i} contains a unique minimum. By the above eliminations, $\min(h(y_2), h(x_2), h(\kappa_1), \dots, h(\kappa_k)) > \max(h(\zeta_1), \dots, h(\zeta_s))$ and any level sphere with height strictly between these two values is a bridge sphere satisfying the conclusions of the theorem. The case when both E_1 and E_2 have unique minima follows similarly. \square

Remark 10.5 In addition, we have shown in the above proof that the number of components in $B_1 \cap \Sigma$ is one more than the number of saddles of S when S is taut.

Theorem 10.6 *The bridge position and the thin position for K are distinct.*

Proof Assume, towards a contradiction, that bridge position and thin position for K coincide. Recall that $w(K) \leq 134$, so it is enough to show that the bridge number of K is at least 9, or equivalently, that K intersects any bridge sphere in at least 18 points. Let Σ be a minimal bridge sphere for K and consider how S_1 intersects Σ . By Theorem 10.4, we may assume that one of the 3-balls that S_1 bounds intersects Σ in a collection of disks and $S_1 - \Sigma$ is a collection of annuli and 2-punctured disks. Suppose the number of intersection curves $|S_1 \cap \Sigma|$ is minimal subject to these constraints. Let c_1, \dots, c_n be the curves of $S_1 \cap \Sigma$ and let D_1, \dots, D_n be the disjoint disks c_1, \dots, c_n bound in Σ . As S_1 is c -incompressible each D_i has at least 2 punctures.

Case 1 $n \geq 2$.

Claim 1 *Either $\Sigma - \bigcup\{D_i\}$ is c -compressible both above and below or bridge position for K is not thin position.*

Proof The bridge sphere Σ separates S^3 into two 3-balls H_1 and H_2 . The planar surface $S_1 \cap H_1$ consists of a, possibly empty, collection of c -incompressible annuli, A_1, \dots, A_n and a, possibly empty, collection of at most two c -incompressible 2-punctured disks E_1 and E_2 . We will use the convention that $\partial E_1 = \partial D_1$ and $\partial E_2 = \partial D_n$. Since $n \geq 2$, $S_1 \cap H_i$ consists of at least one c -incompressible annulus or $S_1 \cap H_i = E_1 \cup E_2$. Since every properly embedded meridional surface in $H_1 - K$ with nonempty boundary is boundary compressible in $H_1 - K$, we can choose F to be a boundary compressing disk for $S_1 \cap H_1$ in H_1 . In particular, we can assume that $F \cap S_1$ is a single essential arc, α , in $S_1 \cap H_1$. Let $\partial F = \alpha \cup \beta$ where β is contained in Σ . In particular, β is disjoint from S_1 except in its boundary. If α is contained in some A_i and β is contained in $\Sigma - \bigcup\{D_i\}$, then boundary compressing A_i along F produces a compressing disk for $\Sigma - \bigcup\{D_i\}$ above Σ . If α is contained in some A_i and β is contained in some D_i , then α is not essential in A_i , a contradiction. If α is contained in some E_i and β is contained in $\Sigma - \bigcup\{D_i\}$, then boundary compressing E_i along F produces a cut-disk for $\Sigma - \bigcup\{D_i\}$ above Σ . If α is contained in some E_i and β is contained in $\Sigma - \bigcup\{D_i\}$, then boundary compressing E_i along F produces a cut-disk for D_1 or D_n above Σ . We can conclude that one of $\Sigma - \bigcup\{D_i\}$, D_1 or D_n is c -compressible above. Similarly, we conclude one of $\Sigma - \bigcup\{D_i\}$, D_1 or D_n is c -compressible below. In particular, if D_1 is c -compressible above we can always find a c -disk below Σ that is disjoint from D_1 since $n \geq 2$. By examining all remaining possibilities for c -disks above and below Σ and noting that D_1 and D_n are distinct, we concluded that either $\Sigma - \bigcup\{D_i\}$ is c -compressible both above and below or Σ has c -disks that satisfy the hypotheses of Lemma 10.1. Hence, either $\Sigma - \bigcup\{D_i\}$ is c -compressible both above and below or bridge position of K is not thin position. \square

Suppose some D_j is c -compressible. As S_1 is c -incompressible, we may assume that the c -disk is disjoint from it. By taking an innermost curve of intersection of this c -disk with all other disks D_i , we can find a disk D_k that is c -compressible in the complement of the others. Let Δ be this c -disk. Without loss of generality, assume Δ is above Σ . By Claim 1, either $\Sigma - \bigcup\{D_i\}$ is c -compressible both above and below or bridge position of K is not thin position. However, we have assumed that bridge position coincides with thin position, so we may assume $\Sigma - \bigcup\{D_i\}$ is c -compressible both above and below. Therefore, Δ and a c -disk for $\Sigma - \bigcup\{D_i\}$ below Σ give a pair of c -disks for Σ on opposite sides with disjoint boundaries. By Lemma 10.1, bridge position and thin position for K are distinct, a contradiction.

By the previous argument, we may assume all D_i are c -incompressible. If B_1 is contained on the same side of S_1 as the D_i , then each D_i must have at least 20 punctures, by Theorem 4.3. It follows that Σ has at least 40 punctures so the bridge number of K is at least 20.

It remains to consider the case when all D_i are c -incompressible and are contained in $S^3 - B_1$. As S_2 is essential, it must intersect Σ and, therefore, it must intersect some D_i , say D_1 . Since S_2 is c -incompressible, then, after isotoping $|S_2 \cap D_1|$ to be minimal, an innermost curve of $S_2 \cap D_1$ in D_1 must bound a subdisk Δ_1 of D_1 containing at least 2 punctures. This subdisk must be c -incompressible as a c -disk for it would be a c -disk for D_1 . If Δ_1 is contained in B_2 , it must have at least 20 punctures, by Theorem 4.3, so Σ has at least 22 punctures and the bridge number of K is at least 11. We conclude that either bridge and thin position for K do not coincide or all innermost curves of $D_1 \cap S_2$ bound disks in D_1 that have at least 2 punctures and are outside of B_2 . If there are at least 8 such innermost curves, then D_1 has at least 16 punctures. As each of D_2, \dots, D_n has at least 2 punctures and $n \geq 2$, it follows that Σ has at least 18 punctures as desired. If there are fewer than 8 innermost curves, then a second innermost curve cobounds with some of the innermost curves a planar surface $F \subset B_2 \cap D_1$ with at most 8 boundary components. A c -disk for F would also be a c -disk for D_1 so F is c -incompressible. Let b be the number of boundary components of F and p be the number of points of intersection between F and K . By Theorem 4.3, it follows that $2 - b - p = \chi(F) \leq -23$. However, D_1 meets K in at least $2(b - 1)$ points outside of F . Therefore, D_1 has at least $25 - b + 2(b - 1) = 23 + b$ punctures. Since $b \geq 2$ and $n \geq 2$, we conclude that Σ has at least 27 punctures in total and the bridge number of K is at least 14.

Case 2 $n = 1$.

Claim 2 *If $n = 1$, there is an isotopy taking S_1 to a level sphere and adding at most one additional maximum to h_K .*

Proof Since $n = 1$, S_1 is a standard round 2–sphere with no saddles, by Remark 10.5. Label each point of $\{x_1, x_2, x_3, x_4\} = K \cap S_1$ with an m if it is a local minimum of $h_{K \cap B_1}$ and label it with an M if it is a local maximum of $h_{K \cap B_1}$. If all points of $K \cap S_1$ receive a common label, then S_1 is isotopic to a level sphere via an isotopy that preserves the number of maxima of h_K , as in Figure 20.




Figure 20

Order the points in $K \cap S_1$ in terms of increasing height so that $h(x_1) < h(x_2) < h(x_3) < h(x_4)$.

By Claim 1 of Theorem 10.4, there is an isotopy of K fixing S_1 and the number of maxima of h_K so that after this isotopy x_4 receives a label of M . By a symmetric argument, we can assume that x_1 receives the label m .




Figure 21

Suppose x_2 is labeled m . Poke a neighborhood of x_4 in S_1 along K toward and just past the nearest maximum of K causing the highest point of $K \cap S_1$ to now be labeled m . See Figure 21. After this isotopy, S_1 contains a single inessential saddle which can be removed as in the proof of Lemma 3.4. If x_3 is labeled m , then S_1 is isotopic to a level sphere via an isotopy that preserves the number of maxima of h_K , as in Figure 20. If x_3 is labeled M , then we can preform an isotopy of K supported in a neighborhood of x_3 that introduces exactly 1 additional maximum to h_K and results in x_3 receiving a label of m . Now that all points in $K \cap S_1$ receive the same label, S_1 is isotopic to a level sphere via an isotopy that preserves the number of maxima of h_K , as in Figure 20. By a symmetric argument, if x_2 receives a label of M , then there is

an isotopy taking S_1 to a level sphere and adding at most one additional maximum to h_K . \square

Use Claim 2 to isotope S_1 to be level at the cost of introducing at most one additional maximum to h_K . Without loss of generality, suppose B_1 is contained below S_1 and the complement of B_1 is contained above S_1 . By Property 9.8, the tangle below S_1 has at least 7 minima and, therefore, at least 5 maxima. By Property 8.4, the tangle above S_1 has at least 5 maxima and, therefore, in this position K has at least 10 maxima. As at most one maximum was added, K has bridge number at least 9. \square

11 Some useful lemmas

Recall that K' denotes an embedding of K that minimizes width. In this section, we establish additional restrictions on thin and thick spheres for K' .

Lemma 11.1 *If G is an essential, 6-punctured, thin sphere for K' , then there are thick spheres of width 10 below and above G . These spheres are not necessarily adjacent to G .*

Proof We will show that there is a thick sphere of width 10 above G . If there is a thin sphere of width 8 or more above G , the result is clear, so suppose all thin spheres above G , if there are any, have width 4 or 6. Let P be the highest thin sphere above G , possibly $P = G$. If $w(P) = 4$, then P must be one of S_1 , S_2 or S_3 , by Corollary 9.7 and Proposition 9.9. The result then follows by Property 8.4, Property 8.10 or Property 9.8. If $w(P) = 6$, by Theorem 2.3, P is incompressible and, by Corollary 3.11, P is c -incompressible above. Therefore, by Proposition 9.10, P is cut-compressible below. Again by Proposition 9.10, it follows that both B_1 and B_2 are disjoint from and below P . In this case, the thick sphere above P has width at least 10 by Property 8.8. \square

Lemma 11.2 *Suppose that P and P' are two adjacent thin spheres for K' so that $4 \leq w(P)$, $w(P') \leq 10$. Suppose D^c is a c -disk for P lying between them so that ∂D^c bounds a three or four punctured disk Δ in P . Then the sphere $F = D^c \cup \Delta$ is essential. Furthermore, if Σ is the thick sphere between P and P' and F does not separate P and P' then:*

- (1) *If $w(P) \geq 6$ and $w(P') \geq 6$, then $w(\Sigma) \geq 14$.*
- (2) *If $w(P) \geq 6$ and $w(P') = 4$, then $w(\Sigma) \geq 12$.*
- (3) *If $w(P) = 8$ and $w(P') = 4$ and D^c is a compressing disk, then $w(\Sigma) \geq 14$.*

Proof Suppose P and P' are two adjacent thin spheres for K' and D^c is a c -disk for P lying between them so that ∂D^c bounds a three or four punctured disk Δ in P . Consider the four punctured sphere $F = \Delta \cup D^c$ and suppose F has a compressing disk E . By using an outermost arc argument, we may assume that $\partial E \subset \Delta$. As F has only 4 punctures, ∂E bounds a twice punctured disk $\delta \subset \Delta$. Then $\delta \cup E$ is a twice punctured sphere. As K is prime, the strand of the knot with both endpoints in δ can be isotoped to lie in δ . It follows that this strand can be isotoped to lie just past the thin sphere P . This isotopy either eliminates a maximum and an adjacent minimum or slides a maximum below a minimum thus decreasing $w(K')$. As K' is in thin position, this is a contradiction.

By Theorem 3.9, we may assume that D^c is vertical. Let B be the ball bounded by F that is disjoint from P' and let $E = B \cap \Sigma$. By Lemma 4.1 E together with the possibly once punctured disk ∂E bounds in D^c is a bridge sphere for the tangle $K' \cap B$. As F is an incompressible 4-punctured sphere, it must be parallel to one of S_1, S_2 or S_3 . By one of Property 8.4, Property 8.10 or Property 9.8, the width of any bridge sphere for $K' \cap B$ is at least 10 and, therefore, E has at least 9 punctures. In addition, let τ_1, \dots, τ_n be the strands of K between P and P' that are disjoint from B . Then $|\Sigma \cap \tau_i| \geq 1$ if τ_i has one of its endpoints in P and one in P' and $|\Sigma \cap \tau_i| \geq 2$ if τ_i has both of its endpoints on the same sphere. It is easy to check that the conclusions of the lemma are satisfied in all three cases, see Figure 22. \square




Figure 22

Remark 11.3 Note that if P is a six-punctured sphere with a cut-disk disjoint from all other thin spheres or if P is an eight-punctured sphere with a compressing disk disjoint from all other thin spheres, then we can always choose Δ so that the sphere F which is the union of the c -disk and Δ is 4-punctured and does not separate P and its adjacent thin sphere.

Lemma 11.4 Suppose P and P' are two adjacent thin spheres for K' with P' above P and P is incompressible but has a cut-disk D^c above it and disjoint from P' . Suppose furthermore that ∂D^c bounds a 5-punctured disk in P and the 6-punctured

sphere G which is the union of D^c and this disk does not separate P and P' . Then the thick sphere Σ between P and P' has width at least $w(P) + 4$.

In the special case where $G = S_1 \#_{\alpha} S_2$, α is a strand that intersects B_T and B_1 is not contained between P and P' , then Σ has at least $w(P) + 8$ punctures.

Proof Let B_G be the ball bounded by G and disjoint from P and let $T = K' \cap B_G$. By Theorem 3.9, we may assume that $G \cap \Sigma = \beta$, where β is a single essential simple closed curve in Σ . By Lemma 4.1, the sphere R obtained by cut-compressing Σ along the cut-disk that β bounds in D^c is a bridge sphere for T . A compressing disk for G would result in a compressing disk for P , so we conclude that G is incompressible and, therefore, it is one of the spheres in Proposition 9.10. In particular either B_1 and B_2 are contained in B_G or they are both disjoint from it.

There are two cases to consider. If B_1 and B_2 are disjoint from B_G , then by Property 8.8 it follows that R has at least 10 punctures. If B_1 and B_2 are contained in B_G , we can apply Theorem 4.2 with $N = B_G$ with the bridge sphere R and $M = B_1$ with the bridge sphere Σ_1 . It again follows that R has at least 10 punctures.

As each strand that has an endpoint in P' must intersect Σ at least once, it follows that $|\Sigma \cap K| \geq |R \cap K| - 1 + |P \cap K| - 5 \geq w(P) + 4$ as desired.

If $G = S_1 \#_{\alpha} S_2$ and B_1 is not contained between P and P' , then, by Property 8.8, $|R \cap K| \geq 13$, so $|\Sigma \cap K| \geq |R \cap K| - 1 + |P \cap K| - 5 \geq w(P) + 8$. \square

12 $w(K \# \text{trefoil}) = w(K)$

In this section, we show that the width of K is 134. This completes the proof that $w(K \# \text{trefoil}) = w(K) < w(K) + w(\text{trefoil}) - 2$. Let K' be a knot isotopic to K that is in thin position. The argument is separated into two parts depending on whether the minimal width thin sphere for K' is cut-compressible or not.

Theorem 12.1 *If a thinnest thin sphere for K' is cut-compressible, then $w(K') \geq 138$.*

Proof As all essential 4-punctured spheres in the complement of K are c -incompressible, it follows that the thin sphere of lowest width has at least 6 punctures. In particular, if P is a thin sphere for K' , $w(P) \geq 6$.

Case A Suppose first that K' has a compressible thin sphere.

Claim *There exist adjacent thin spheres P and P' so that P has a compressing disk on the same side as P' but disjoint from it.*

Proof Let P'' be a compressible thin sphere for K' with compressing disk D' . Consider the intersection of D' with the collection of all thin spheres for K' . Assume that D' has been isotoped so this intersection is minimal. In particular, every curve of intersection is essential in the corresponding thin sphere. Let σ be an innermost curve of intersection in D' . Let P be the thin sphere containing σ . Then P has a compressing disk $D \subset D'$ disjoint from all other thin spheres. By Corollary 3.11, there are other thin spheres on the same side of P as D . In particular, we can choose P' to be such a sphere so that P and P' are adjacent thin spheres. \square

Let P and P' be the two spheres guaranteed by the claim. Without loss of generality, assume P is below P' . By Theorem 2.3, it follows that $w(P) \geq 10$. If $w(P) \geq 12$, then, by Lemma 2.1, $w(K') \geq \frac{1}{2}(14^2 + 14^2 + 8^2 - 12^2 - 6^2) = 138$, so suppose $w(P) = 10$. By Corollary 2.7, compressing P along D yields a copy of P' . As $w(P') \geq 6$ and K is prime, it follows that $w(P') = 6$ and ∂D bounds a 4-punctured disk $\Delta \subset P$ so that $D \cup \Delta$ is a sphere that does not separate P and P' . By Lemma 11.2 part (1), the thick sphere between P and P' has width at least 14. By Lemma 11.1, there is a thick sphere of width at least 10 above P' and one of width at least 12 below P . It follows that $w(K') \geq 152$.

Case B All thin spheres for K' are incompressible.

Let P be a thinnest thin sphere for K' and suppose it is cut-compressible above. By Corollary 3.11, K' must have at least one other thin sphere, P' , above P . If $w(P') \geq 12$, then, by Lemma 2.1, $w(K') \geq \frac{1}{2}(14^2 + 14^2 + 8^2 - 12^2 - 6^2) = 138$, so suppose every thin sphere for K' has width at most 10.

Let D^c be the cut-disk for P . By taking an innermost curve of intersection of D^c with the union of all other thin spheres for K' , we can find a thin sphere that has a cut-disk which is disjoint from all other thin spheres. Let P' be the thin sphere of lowest width amongst all thin spheres for K' that has this property and let D'^c be its cut-disk. Without loss of generality, we will assume D'^c is above P' . Let P'' be the thin sphere adjacent to P' above it. By Corollary 3.11, this sphere exists and $w(P'') < w(P')$.

Case 1 $w(P') = 6$. In this case, $w(P'') \leq 4$, by Corollary 3.11. Hence, this case does not satisfy the hypotheses of the theorem at hand.

Case 2 $w(P') = 8$. By Corollary 2.7, cut-compressing P' along D'^c yields a copy of P'' . As $w(P'')$ in this case must be 6, it follows that $\partial D'^c$ bounds a three punctured disk in P' so that the union of D'^c and this disk is a sphere that does not separate P' and P'' . By Lemma 11.2 part (1), the thick sphere between P' and P'' has width at

least 14. By Lemma 11.1, there is a thick sphere of width at least 10 above P'' and so $w(K') \geq 148$.

Case 3 $w(P') = 10$. There are three subcases to consider:

Case 3a Suppose that ∂D^c bounds a three-punctured disk in P' so that the union of D^c and this disk is a sphere that does not separate P' and P'' . By Lemma 11.2 part (1), the thick sphere between P' and P'' has width at least 14. If $w(P'') = 6$, then, by Lemma 11.1, there is a thick sphere of width at least 10 above it and so $w(K') \geq 152$. If $w(P'') = 8$, then $w(K') \geq 138$.

Case 3b Suppose that ∂D^c bounds two five-punctured disks in P' . As compressing P' along D^c yields a copy of P'' , $w(P'') = 6$. By Lemma 11.4, the thick sphere between P' and P'' has width at least 14. By Lemma 11.1, the thick sphere above P'' has width at least 10. It follows that $w(K') \geq 152$.

Case 3c Suppose that ∂D^c bounds a seven-punctured disk in P' so that the union of D^c and this disk is a sphere that does not separate P' and P'' . By Theorem 3.9, we may assume that D^c is vertical. By Corollary 2.7, cut-compressing P' along D^c gives a copy of P'' . Hence, $w(P'') = 4$, contradicting the assumption. \square

Theorem 12.2 *If the thinnest thin sphere for K is cut-incompressible, then $w(K') \geq 134$. Moreover, if $w(K') = 134$, any thin position for K' has exactly three thick levels of width 10 and exactly two thin levels of width 4.*

Proof Let P be the thinnest thin sphere for K' and note that, by Theorem 2.2, P is incompressible. As P is cut-incompressible, by Proposition 9.9, P either has 4 punctures or at least 14 punctures. In the second case, $w(K') \geq 158$ so we may assume that P is one of S_1 , S_2 or S_3 .

Case 1 K' has exactly one thin sphere, P .

Suppose P is isotopic to S_1 and, without loss of generality, suppose that B_1 is below it. By Property 8.4, the thick sphere below P has width at least 10 and, by Property 9.8, the thick sphere above it has width at least 14. It follows that $w(K') \geq 140$. Similarly, the same width bound follows if P is isotopic to S_2 .

Suppose then that P is isotopic to S_3 . By Property 8.10, the thick surfaces above and below P have width at least 12. In this case, $w(K') \geq 136$ as desired.

Case 2 Neither S_1 nor S_2 is a thin sphere and there are at least 2 thin spheres.

In this case, we may assume $P = S_3$ and any other thin spheres have width at least 6 and, therefore, are c -compressible. The sphere S_3 splits K into two tangles, \mathcal{T}_a

and \mathcal{T}_b , lying above and below S^3 respectively. Let $w(\mathcal{T}_a)$ (respectively $w(\mathcal{T}_b)$) be the sum of the widths of all level spheres lying above (respectively below) S_3 . Thus $w(K') = w(\mathcal{T}_a) + w(\mathcal{T}_b) + 4$.

By hypothesis, there is at least one other thin sphere say above P . Since all thin spheres other than P are c -compressible, let D'_a be c -disk for some thin sphere above P . As P is c -incompressible, $D'_a \cap P = \emptyset$. By taking an innermost curve of intersection of D'_a with the collection of all other thin spheres for K' , we can find a cut or compressing disk D_a for some thin sphere P_a above P that is disjoint from all other thin spheres for K' . In addition, we can assume that either D_a is a compressing disk or P_a is incompressible. Let T_a be the thick sphere that intersects D_a .

If $w(P_a) \geq 10$, then direct computation shows that $w(\mathcal{T}_a) \geq 88$.

If $w(P_a) = 8$, and D_a is a compressing disk, then, by Lemma 11.2, $w(T_a) \geq 12$ so $w(\mathcal{T}_a) \geq 84$.

If $w(P_a) = 8$, and D_a is a cut-disk (in particular as noted above this allows us to assume P_a is incompressible) then either by Lemma 11.2 or by Lemma 11.4, $w(T_a) \geq 12$ so again $w(\mathcal{T}_a) \geq 84$.

If $w(P_a) = 6$, then, by Lemma 11.2 part (2), the thick sphere T_a that intersects D_a has width at least 12. In this case, $w(\mathcal{T}_a) \geq 80$.

We can apply all of the above arguments also to \mathcal{T}_b but in addition we must consider the case when \mathcal{T}_b doesn't have any thin spheres. In that case by Property 8.10, the thick surface below P has width at least 12 so $w(\mathcal{T}_b) \geq 66$.

Therefore in this case $w(K') = w(\mathcal{T}_a) + w(\mathcal{T}_b) + 4 \geq 80 + 66 + 4 = 150$.

Case 3 Exactly one of S_1 or S_2 is a thin sphere and there are at least two thin spheres.

Without loss of generality, we may assume that $P = S_1$ is a thin sphere and B_1 is below P .

Claim 1 *There is a thick sphere of width at least 10 below P .*

Proof If there aren't any thin spheres below P , there is a thick sphere of width at least 10, by Property 8.4. If there is a thin sphere of width 8 or more, the result follows immediately. Suppose all thin spheres below P have width at most 6. However, all such spheres are incompressible, by Theorem 2.3, and there are no such spheres in B_1 , by Proposition 9.1. □

Claim 2 *If there is a compressible thin sphere P' above P and $w(P') \leq 8$, then $w(K') \geq 154$.*

Proof Suppose that K' has a thin sphere P' that is compressible. By Theorem 2.3, we may assume that $w(P') = 8$. Let D be its compressing disk. Then D is disjoint from all thin spheres of width less than 8 as all such spheres are incompressible. Therefore, by taking the intersection of D with all thin spheres for K' and rechoosing P' we may assume that D is disjoint from all other thin spheres. If this new P' has the property that $w(P') \geq 10$, then there are two distinct thin spheres above P and $w(K') \geq 154$. Hence, we can assume that $w(P') = 8$ and D is a compressing disk for P' contained between consecutive thin spheres P' and P'' . If ∂D bounds a 2-punctured disk in P' , then this disk together with D cobound a 3-ball containing an unknotted arc. Such a 3-ball gives rise to an isotopy that thins K' . Hence, we can assume that ∂D bounds a 4-punctured disk to each side in P' . By Corollary 2.7, compressing P' along D gives rise to a copy of P'' . Since ∂D bounds 4-punctured disks to each side in P' , then P'' is a 4-punctured sphere. By Lemma 11.2 part (3), the thick sphere intersecting D has width at least 14. Then $w(K') \geq 158$ as desired. \square

Subcase 3A Suppose first that K' has no thin spheres above P . By Property 9.8, there is thick sphere of width at least 14 above P . By hypothesis, there is a thin sphere below P . By Proposition 9.1, this sphere cannot have width 4 or 6 as all such spheres are incompressible. Therefore, the sphere must have width at least 8 and so $w(K') \geq 158$.

Subcase 3B Suppose that K' has exactly one other thin sphere P' above P .

If $w(P') = 4$, then P' must be S_3 . By Property 8.10, the thick surface above P' has width at least 12. Consider the thick surface T between S_1 and S_3 . By Property 8.2, $w(T) \geq 8$ and so $w(K') \geq 138$.

Suppose then that $w(P') \geq 6$. By Claim 2, we may assume that P' is incompressible or $w(P') \geq 10$. If $w(P') \geq 10$, then $w(K') \geq 136$ as desired. If P' is c -incompressible, then P' meets K in at least 14 points, by Proposition 9.9. Hence, we can assume that P' is cut-compressible. The cut-disk for P' is disjoint from P and lies below P' , by Corollary 3.11.

If $w(P') = 6$, then the thick sphere between P and P' has width at least 12, by Lemma 11.2 part (2). By Lemma 11.1, it follows that the thick sphere above P' has width at least 10 and, therefore, $w(K') \geq 146$.

Suppose then that $w(P') = 8$, then P' is incompressible, by Claim 2, and P' has a cut-disk D^c , by Proposition 9.9. As P' is the only thin sphere above P , by Corollary 3.11, D^c must be below it and, by Theorem 3.9, we can assume that D^c is vertical. By Corollary 2.7, cut-compressing P' along D^c results in a copy of P and, therefore,

∂D^c bounds a 5-punctured disk Δ in P' so that the 6-punctured sphere $G = D^c \cup \Delta$ does not separate P and P' .

A compressing disk for G would result in a compressing disk for P' and $w(K') \geq 154$, by Claim 2. Hence, we may assume that G is incompressible and, therefore, it is one of the spheres classified in Proposition 9.10. In particular, G does not separate B_1 and B_2 . Let B_G be the ball bounded by G disjoint from P . As B_1 is disjoint from B_G , so is B_2 . Therefore, there are three possibilities to consider: B_2 is below P' but outside of B_G , B_2 is above P' or B_2 intersects P' .

It is clear that B_2 cannot be contained below P' and be disjoint from B_G as that would imply that the essential surface S_2 is completely contained in the product region between P and the 4-punctured sphere resulting from cut-compressing P' along D^c .

If B_2 is completely contained above P' , then let Σ be the thick surface for K above P' . By Theorem 4.2, it follows that either Σ has at least 26 punctures or Σ is isotopic to Σ_2 with possibly some tubes attached. If no tubes are attached, then Σ is parallel to Σ_2 and S_2 is an essential sphere completely disjoint from the bridge sphere of the tangle lying above P' , which is not possible. Therefore, at least one tube is attached. We conclude that Σ has at least 12 punctures. By Lemma 11.4, the thick sphere below P' also has width at least 12. Hence, $w(K) \geq 154$.

Suppose that B_2 intersects P' . We have already assumed that P is isotopic to S_1 and we have established that cut-compressing P' along D^c produces P and an incompressible 6-punctured sphere, G . Since tubing along a strand of K is the inverse operation to cut-compressing, P' is isotopic to $G \#_{\beta} S_1$. However, both S_1 and G can be isotoped to be disjoint from B_2 . Hence, β intersects B_2 , as otherwise B_2 could be isotoped to be disjoint from P' .

There are several cases to consider.

If $G = S_1 \#_{\epsilon} S_3$ or $G = S_2 \#_{\epsilon} S_3$, where ϵ is the strand not passing through B_T , then G is compressible which is not possible.

Suppose $G = S_1 \#_{\gamma} S_3$, where γ is a strand passing through B_T ; see the first schematic of Figure 23. Then $P' = G \#_{\alpha} S_1$. The strand α may contain ϵ or it may not. In either case, the tangle S above P' can be obtained from the tangle T' contained on one side of S_3 by replacing one of the strands with three parallel strands as in Figure 23. By Property 8.10, every bridge surface for T' has at least 12 punctures. Since each of the new strands must intersect the bridge surface in at least two points, the thick surface above P' has at least 16 punctures. By Lemma 11.4, the thick sphere directly below P' has at least 12 punctures, so $w(K) \geq 210$.




Figure 23

Suppose $G = S_1 \#_{\gamma} S_2$, where γ is a strand passing through B_T . In this case, by Lemma 11.4, the thick sphere directly below P' has width at least 16, so $w(K) \geq 188$.

Suppose $G = S_1 \#_{\epsilon} S_2$, where ϵ is the strand not passing through B_T , ie $G = S_4$. Then the tangle S above P' can be obtained from the tangle \mathcal{T}_2 by replacing one of the strands with three parallel strands; see Figure 24. As any bridge surface for \mathcal{T}_2 has at least 10 punctures, by Property 8.4, and each of the two additional strands has to intersect the thick surface for S at least twice, it follows that the thick surface above P' has at least 14 punctures. By Lemma 11.4, the thick sphere directly below P' also has at least 12 punctures, so $w(K) \geq 180$.

Finally, suppose that $G = S_2 \#_{\gamma} S_3$, where γ is a strand passing through B_T , see Figure 25. Let S be the 4 strand tangle above P' . By deleting two of the four strands of S , we can obtain the tangle \mathcal{T}_2 . Therefore, by Property 8.4 the thick surface above P' intersects two of the strands in S in at least 10 points and each of the other two strands in at least 2 points each. Hence, this thick sphere meets K in at least 14 points. As before, the thick sphere below P' has width at least 12, so $w(K) \geq 180$.

Subcase 3C Suppose that there are at least 2 thin spheres above P .

If at least one of these spheres has width at least 10, then $w(K') \geq 146$. Hence, we may assume that all thin spheres above P have width at most 8. By Claim 2, we may assume that all thin spheres for K' above P are incompressible. If one of the thin spheres above P is c -incompressible, then, by Proposition 9.9, that thin sphere is isotopic to S_2 or S_3 , a contradiction to the assumption that S_2 is not a thin sphere. Hence, one of the thin spheres above P is cut-compressible. By taking the intersection




Figure 24




Figure 25

of such a cut-disk with the union of all thin spheres above P , we can find a cut-disk D^c for some thin sphere P' that is disjoint from all other thin spheres. Let P'' be the thin sphere adjacent to P' on the same side of P' as D^c . This sphere exists, by Corollary 3.11. If $w(P') = 6$, then, by Lemma 11.2 part (2), the thick sphere between P' and P'' has width at least 12. It follows that $w(K') \geq 142$. If $w(P') = 8$, then, by Lemma 11.2 or Lemma 11.4, the thick sphere between P' and P'' has width at least 12 and, so $w(K') \geq 142$.

Case 4 S_1 and S_2 are both thin spheres.

By Claim 1 of Case 3, it follows that there are thick spheres of width at least 10 below S_1 and above S_2 . If there is a thin sphere between S_1 and S_2 , it must either have width 4 or it must have width at least 8 as all 6-punctured thin spheres do not separate B_1 from B_2 . If there is a thin sphere of width 8, then $w(K) \geq 152$. Suppose there is a thin sphere of width 4 between S_1 and S_2 . There can be only one such sphere and it is necessarily S_3 . Let T_a and T_b be the thick spheres directly above and below S_3 . Both of these thick spheres must have width at least 6. If $w(T_a) = w(T_b) = 6$, then K has exactly one minimum and one maximum between S_1 and S_3 and similarly between S_3 and S_2 . This implies that the tangle T between S_1 and S_3 has a bridge sphere of width 8, contradicting Property 8.9. Therefore at least one of these spheres has width at least 8. But in this case $w(K) \geq 138$ as desired.

Suppose that S_1 and S_2 are adjacent thin spheres. By Property 8.9, the width of the thick sphere between them is at least 10 and so $w(K') \geq 134$ as desired. \square




Figure 26

Proof of Theorem 1.1 Let K'_α be any two-bridge knot and let K_α be any of the knots constructed in Section 7. By Theorems 12.1 and 12.2, it follows that $w(K_\alpha) = 134$. It is easy to see that the width of any two-bridge knot is eight. Figure 1 demonstrates that $w(K_\alpha \# K'_\alpha) \leq w(K_\alpha) = 134$. By [14], $w(K_\alpha \# K'_\alpha) \geq w(K_\alpha)$ and, therefore, $w(K_\alpha \# K'_\alpha) = w(K_\alpha)$. \square

Proof of Theorem 1.2 By Theorems 12.1 and 12.2, it follows that $w(K_\alpha) = 134$ and K_α has a thin position with exactly three thick spheres of width 10 and exactly two thin spheres of width 4. In this position, K_α has 11 maxima. However, Figure 26 demonstrates that $b(K'_\alpha) \leq 10$. \square

References

- [1] **R C Blair**, *Bridge number and Conway products*, *Algebr. Geom. Topol.* 10 (2010) 789–823 MR2629764
- [2] **R Blair, M Tomova**, *Companions of the unknot and width additivity*, *J. Knot Theory Ramifications* 20 (2011) 497–511 MR2796224
- [3] **R Blair, M Tomova, M Yoshizawa**, *High distance bridge surfaces* arXiv: 1203.4294
- [4] **J H Conway**, *An enumeration of knots and links, and some of their algebraic properties*, from: “Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)”, (J Leech, editor), Pergamon, Oxford (1970) 329–358 MR0258014
- [5] **D Gabai**, *Foliations and the topology of 3–manifolds. III*, *J. Differential Geom.* 26 (1987) 479–536 MR910018
- [6] **H Goda, M Scharlemann, A Thompson**, *Levelling an unknotting tunnel*, *Geom. Topol.* 4 (2000) 243–275 MR1778174
- [7] **C M Gordon, J Luecke**, *Knots are determined by their complements*, *J. Amer. Math. Soc.* 2 (1989) 371–415 MR965210
- [8] **D J Heath, T Kobayashi**, *A search method for thin positions of links*, *Algebr. Geom. Topol.* 5 (2005) 1027–1050 MR2171802
- [9] **J Johnson, M Tomova**, *Flipping bridge surfaces and bounds on the stable bridge number* arXiv:0908.3690
- [10] **L H Kauffman, S Lambropoulou**, *Classifying and applying rational knots and rational tangles*, from: “Physical knots: knotting, linking, and folding geometric objects in \mathbb{R}^3 (Las Vegas, NV, 2001)”, (J A Calvo, K C Millett, E J Rawdon, editors), *Contemp. Math.* 304, Amer. Math. Soc. (2002) 223–259 MR1953344
- [11] **Y Rieck, E Sedgwick**, *Thin position for a connected sum of small knots*, *Algebr. Geom. Topol.* 2 (2002) 297–309 MR1917054
- [12] **J H Rubinstein**, *An algorithm to recognize the 3–sphere*, from: “Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)”, (S D Chatterji, editor), Birkhäuser, Basel (1995) 601–611 MR1403961
- [13] **M Scharlemann**, *Thin position in the theory of classical knots*, from: “Handbook of knot theory”, (W Menasco, M Thistlethwaite, editors), Elsevier, Amsterdam (2005) 429–459 MR2179267
- [14] **M Scharlemann, J Schultens**, *3–manifolds with planar presentations and the width of satellite knots*, *Trans. Amer. Math. Soc.* 358 (2006) 3781–3805 MR2218999
- [15] **M Scharlemann, A Thompson**, *On the additivity of knot width*, from: “Proceedings of the Casson Fest”, (C Gordon, Y Rieck, editors), *Geom. Topol. Monogr.* 7 (2004) 135–144 MR2172481

- [16] **H Schubert**, *Über eine numerische Knoteninvariante*, Math. Z. 61 (1954) 245–288 MR0072483
- [17] **H Schubert**, *Knoten mit zwei Brücken*, Math. Z. 65 (1956) 133–170 MR0082104
- [18] **J Schultens**, *Additivity of bridge numbers of knots*, Math. Proc. Cambridge Philos. Soc. 135 (2003) 539–544 MR2018265
- [19] **A Thompson**, *Thin position and the recognition problem for S^3* , Math. Res. Lett. 1 (1994) 613–630 MR1295555
- [20] **M Tomova**, *Compressing thin spheres in the complement of a link*, Topology Appl. 153 (2006) 2987–2999 MR2248402
- [21] **M Tomova**, *Multiple bridge surfaces restrict knot distance*, Algebr. Geom. Topol. 7 (2007) 957–1006 MR2336246
- [22] **M Tomova**, *Cut-disks for level spheres in link and tangle complements*, Topology Appl. 156 (2009) 783–794 MR2492962
- [23] **Y-Q Wu**, *Thin position and essential planar surfaces*, Proc. Amer. Math. Soc. 132 (2004) 3417–3421 MR2073319

Department of Mathematics, University of Pennsylvania

David Rittenhouse Lab, 709 South 33rd Street, Philadelphia, PA 19104-6395, United States

Department of Mathematics, University of Iowa

14 MacLean Hall, Iowa City, IA 52242-1419, United States

ryblair@math.upenn.edu, maggy-tomova@uiowa.edu

<http://www.math.upenn.edu/~ryblair>, <http://www.math.uiowa.edu/~mtomova>

Proposed: Cameron Gordon

Received: 18 June 2010

Seconded: David Gabai, Ronald Stern

Revised: 25 March 2012