

Loose Legendrians and the plastikstufe

EMMY MURPHY
KLAUS NIEDERKRÜGER
OLGA PLAMENEVSKAYA
ANDRÁS I STIPSICZ

We show that the presence of a plastikstufe induces a certain degree of flexibility in contact manifolds of dimension $2n + 1 > 3$. More precisely, we prove that every Legendrian knot whose complement contains a “nice” plastikstufe can be destabilized (and, as a consequence, is loose). As an application, it follows in certain situations that two nonisomorphic contact structures become isomorphic after connect-summing with a manifold containing a plastikstufe.

57R17

1 Introduction

In dimension 3, it has been known for a long time that contact structures containing a topological object called an *overtwisted* disk are “flexible”, in the sense that two overtwisted contact structures which are homotopic as oriented 2–plane fields are also isotopic; see Eliashberg [5]. Often this property is phrased as saying that overtwisted contact structures on 3–manifolds satisfy an h –principle. A second important property of all contact structures containing overtwisted disks is that such contact 3–manifolds are not symplectically fillable.

In higher dimensions, the quest for the right definition of “overtwisted” contact structures has been going on for a while, but we are far from having a definitive answer. On the one hand, there is a variety of special submanifolds, plastikstufes and bLobs (see the second author alone [22] and with Massot and Wendl [17]), whose presence in a contact manifold is known to obstruct fillability. On the other hand, certain related objects possess flexibility properties desired for “overtwisted” contact structures. More specifically, a class of Legendrian knots, called “loose” Legendrians, satisfies a version of h –principle (see the first author [20]), and Weinstein cobordisms obtained by attaching symplectic handles along loose knots are governed by a “symplectic h –cobordism theorem” (see Cieliebak and Eliashberg [2]). (Roughly speaking, loose Legendrian knots in higher-dimensional contact manifolds are those that contain a

sufficiently wide “kink” whose projection to a 3–dimensional subspace is a stabilized Legendrian arc. See [Section 6](#) for a precise definition.) In this paper, we hope to shed some light on flexibility in high dimensions by studying loose Legendrians in contact manifolds containing a plastikstufe. (The latter is a foliated submanifold of maximal dimension that is a product of an overtwisted disk and a closed manifold.) Our main result is the following:

Theorem 1.1 *Let (M^{2n+1}, ξ) be any contact manifold containing a small plastikstufe \mathcal{P}_B with spherical core and trivial rotation. Then any Legendrian in (M, ξ) which is disjoint from \mathcal{P}_B is loose.*

(The definitions of the various terms in the above theorem are deferred to [Sections 3](#) and [6](#).)

Etnyre and Pancholi [\[10\]](#) have constructed examples of PS –overtwisted contact manifolds satisfying the hypotheses of [Theorem 1.1](#). The proof of [Theorem 1.1](#) is based on the proof of the corresponding statement concerning overtwisted contact 3–manifolds, together with a certain isotopy gained from a classical h –principle (see Gromov [\[14\]](#)) to bring a part of the given Legendrian into product position with respect to the plastikstufe.

As a consequence of the above theorem, we establish a flexibility result motivated by a familiar 3–dimensional fact: if two contact structures on a 3–manifold are homotopic as oriented 2–plane fields, they become isotopic as contact structures after connect-summing with any overtwisted contact manifold. The following theorem is a corollary of [Theorem 1.1](#) and of the results of Cieliebak and Eliashberg [\[2\]](#) on flexible Weinstein cobordisms. Different versions of the symplectic h –cobordism theorem [\[2\]](#) imply several versions of our result below.

Theorem 1.2 *Consider two contact structures ξ_0, ξ_1 on a manifold Y of dimension $2n+1 > 3$. Let (M, ξ_{PS}) be a simply connected manifold containing a small plastikstufe with spherical core and trivial rotation. Assume that one of the following conditions holds:*

- (1) *There exists a manifold W of dimension $2n+2$ with $\partial W = Y$, and W carries two Stein structures J_0, J_1 such that (W, J_0) is a filling for ξ_0 , (W, J_1) is a filling for ξ_1 , and J_0, J_1 are homotopic through almost complex structures.*
- (2) *There exists a Stein cobordism (W, J) from (Y, ξ_0) to (Y, ξ_1) such that W is smoothly the product cobordism $Y \times [0, 1]$.*

Then $(M, \xi_{PS}) \# (Y, \xi_0)$ is contactomorphic to $(M, \xi_{PS}) \# (Y, \xi_1)$.

According to [2], the requirement about the existence of Stein structures on W can be exchanged to Weinstein structures. (For the definition of a Weinstein structure, see Section 6.) Indeed, a Stein structure induces a Weinstein structure, and any Weinstein structure (after possibly a homotopy through Weinstein structures) can be induced by a Stein structure. In addition, two Stein structures are homotopic (through Stein structures) if and only if the induced Weinstein structures are homotopic (through Weinstein structures). In conclusion, in the above theorem the two notions are interchangeable. In fact, in accordance with the results we quote from [2], the proofs in Section 6 will be phrased in the language of Weinstein manifolds, and we appeal to the above interchangeability principle. Unlike in the 3–dimensional situation, we are unable to prove any flexibility result in the absence of Stein/Weinstein fillings or cobordisms.

Seidel and Smith [24], and later McLean [19] constructed infinite families of examples of exotic Stein structures of finite type on \mathbb{R}^{2n+2} for $n > 1$. (Yet further such examples are given by Abouzaid and Seidel [1].) A spherical level set that encloses all the critical points of the plurisubharmonic Morse function in any of the above exotic Stein \mathbb{R}^{2n+2} carries a contact structure ξ_{exo} filled by a nonstandard Stein ball. By calculating symplectic homologies, it is shown [1; 19; 24] that $(S^{2n+1}, \xi_{\text{exo}})$ is not contactomorphic to $(S^{2n+1}, \xi_{\text{std}})$. (Indeed, it is expected [18] that infinitely many of these exotic contact spheres are pairwise noncontactomorphic.) Now, let $(S^{2n+1}, \xi_{\text{PS}})$ be a sphere containing a plastikstufe as in Theorem 1.2 (the existence of which, with the right rotation class in every dimension, will be checked in Section 5).

Corollary 1.3 *With the notations as above, $(S^{2n+1}, \xi_{\text{exo}} \# \xi_{\text{PS}})$ is contactomorphic to $(S^{2n+1}, \xi_{\text{std}} \# \xi_{\text{PS}}) \cong (S^{2n+1}, \xi_{\text{PS}})$.*

Proof This is an immediate consequence of the previous theorem: one could use either condition (1), applied to fillings by the standard and nonstandard Stein ball, or condition (2), applied to the cobordism obtained by puncturing the nonstandard Stein ball. \square

The paper is organized as follows. In Section 2, we quickly review the 3–dimensional case: if M^3 is overtwisted, then every Legendrian knot $L \subset M$ disjoint from an overtwisted disk can be destabilized. The destabilization is realized by taking a connected sum with the boundary of an overtwisted disk. (While this seems to be a “folklore” statement, explicit proofs are absent from the literature.) The definition and properties of plastikstufes are reviewed in Section 3. In Section 4 we recall the definition of loose knots and in Section 5, via the Etnyre–Pancholi construction, we find plastikstufes with trivial rotation. In Section 6 we give the proof of Theorem 1.2. We close the paper with a short list of some open problems in Section 7.

Acknowledgments The main results of the present work were found when the authors visited the American Institute of Mathematics (AIM), as participants of the “*Contact topology in higher dimensions*” workshop. We would like to thank AIM for their hospitality, the ESF and the NSF for the support and other workshop participants, in particular to Atsuhide Mori, for stimulating conversations. We are grateful to John Etnyre for useful conversations and e-mail correspondence. The third author would also like to thank Yasha Eliashberg, Ko Honda and Thomas Vogel for answering some questions. EM was partially supported by NSF grant DMS-0943787. KN was partially supported by Grant ANR-10-JCJC 0102 of the *Agence Nationale de la Recherche*. OP was partially supported by NSF grant DMS-1105674. AS was partially supported by ERC Grant LDTBud, by *ADT Lendület* and by OTKA NK81203.

2 The 3–dimensional case: Sliding over an overtwisted disk

In this section we prove that a Legendrian knot in the complement of an overtwisted disk can be destabilized. (For related results and constructions, see also Huang [16].) More precisely, we need a local statement: assuming that the knot is close to the overtwisted disk, we would like to find a destabilization by modifying the knot *only* in the neighborhood of the overtwisted disk. For this local modification, we restrict to the case of a “simple” overtwisted disk, that is, an embedded disk whose characteristic foliation is isomorphic to the one shown in Figure 1: the boundary of the disk is the closed leaf of the foliation, and there is a unique (elliptic) singularity inside the disk. (Recall that an overtwisted disk is any embedded disk with Legendrian boundary with $tb = 0$. Standard arguments resting on Giroux Flexibility imply that a manifold containing an overtwisted disk also contains a simple one.) Although it is a known fact that any Legendrian knot in the complement of an overtwisted disk can be destabilized (cf Dymara [3]), we were unable to find the above “local” version in the literature, and therefore (for the sake of completeness) we include the arguments below.

Before proceeding any further, we recall the definition of the stabilization for Legendrian knots in \mathbb{R}^3 . (For a general discussion on Legendrian knots, see, eg Etnyre [9].) Suppose that in a Darboux chart containing a strand of the Legendrian knot L , the front projection of this strand has a cusp. To stabilize L , remove this cusp in the projection and replace it by a kink, as shown in Figure 2. It is not hard to check that, up to Legendrian isotopy, the stabilized knot L_{stab} is independent of the choice of the Darboux coordinates and of the point where the stabilization is performed. A more invariant definition can be given by using convex surfaces: L_{stab} is a stabilization of L if $L \cup L_{\text{stab}}$ is the boundary of a convex annulus whose dividing set consists of a single arc with endpoints on L_{stab} .




Figure 1: A simple overtwisted disk




Figure 2: Stabilization of a Legendrian knot in dimension 3

Theorem 2.1 *Let (M^3, ξ) be an overtwisted contact manifold. Suppose that L is a Legendrian knot in the complement of a simple overtwisted disk \mathcal{D}_{ot} . Then L can be destabilized, that is, it is the stabilization of another Legendrian knot. A destabilization of L is given by the Legendrian connected sum $L \# \partial\mathcal{D}_{\text{ot}}$ of L and the boundary of the overtwisted disk.*

Corollary 2.2 *If (M^3, ξ) is an overtwisted contact manifold, then any Legendrian knot in the complement of any overtwisted disk can be destabilized. \square*

The above corollary follows immediately from [Theorem 2.1](#): as we already remarked, any overtwisted contact structure on the knot complement contains a simple overtwisted disk. (Alternatively, the corollary can be derived from results of Eliashberg and Fraser [\[6\]](#) or Etnyre [\[8\]](#).) It is useful, however, to have an explicit destabilization procedure given by [Theorem 2.1](#).

Our proof of [Theorem 2.1](#), or at least its main idea, is essentially borrowed from Vogel [\[26, Proposition 3.22\]](#), although the statement of [\[26\]](#) is different from ours. (We are also indebted to John Etnyre, who pointed out Vogel's lemma to us.) We modify the proof from [\[26\]](#) to adapt it to our present purposes.

Proof of [Theorem 2.1](#) We will use convex surface theory [\[12; 15\]](#) to see the destabilization. First, notice that the overtwisted disk from [Figure 1](#) is convex, and its dividing

set consists of a unique closed curve. By Giroux's Flexibility Principle, we can isotope the disk (while keeping its Legendrian boundary fixed) to obtain a convex disk with any characteristic foliation divided by the same curve. Consider the foliation shown in [Figure 3](#). It is clear that this foliation can be carried by a convex surface and is divided by the same closed curve. Thus, after an isotopy, we can assume that the foliation on \mathcal{D}_{ot} looks like [Figure 3](#) and has the following features:

- (i) A family of straight Legendrian arcs (vertical lines on [Figure 3](#)) separate \mathcal{D}_{ot} into two half-disks.
- (ii) There are smooth Legendrian closed curves that run close to the boundary of these half-disks and are formed by the union of the leaves of the foliation.

Now, consider a (piecewise smooth) Legendrian curve formed by the left semicircle of $\partial\mathcal{D}_{\text{ot}}$ and the leftmost vertical Legendrian arc. Let D_l denote the half-disk bounded by this curve in \mathcal{D}_{ot} . A disk D_r is defined similarly on the right side of \mathcal{D}_{ot} .

Recall that the Thurston–Bennequin number of a Legendrian knot K can be computed [\[15\]](#) from the dividing set Γ_Σ on its convex Seifert surface Σ :

$$(1) \quad \text{tb}(K) = -\frac{1}{2}|\Gamma \cap K|.$$

From this formula, $\text{tb}(\partial D_l) = \text{tb}(\partial D_r) = -1$. Note that the Thurston–Bennequin number is defined even for piecewise smooth curves: indeed, linking with the transverse push-off is still well-defined. The formula of [Equation \(1\)](#) also holds in this context, which can be seen, for example, via approximation by smooth Legendrian curves. We will be attaching the overtwisted disk in two steps, first one half, that is, D_l , then the other, ie the Legendrian arcs and D_r . The idea is that half of \mathcal{D}_{ot} is easier to control: indeed, half of an overtwisted disk represents a bypass, and bypasses can be found in tight contact manifolds [\[15\]](#).

Now we are ready to prove the theorem. Since the statement is local, it suffices to establish the claim for a small Legendrian unknot L_0 with $\text{tb}(L_0) = -1$, located near the overtwisted disk \mathcal{D}_{ot} (and disjoint from it). Let L be another standard Legendrian unknot with $\text{tb}(L) = -1$, such that L_0 and L together bound a thin annulus A whose framing gives the Seifert framing on both knots, ie the contact framings of the knots are one less than the framings provided by the annulus. This property guarantees that after a small perturbation, we can assume A to be convex. We will show that $L \# \partial\mathcal{D}_{\text{ot}}$ is a destabilization of L_0 . To begin, notice that the dividing set on A consists of two parallel arcs, each connecting L_0 and L . To establish the theorem, we will show that the annulus $A \# \mathcal{D}_{\text{ot}}$ bounded by L_0 and $L \# \partial\mathcal{D}_{\text{ot}}$ can be perturbed into a convex surface with dividing set given by one boundary parallel arc with endpoints on L_0 (see [Figure 4](#)). This will be done in two steps.




Figure 3: The overtwisted disk \mathcal{D}_{ot} can be endowed with another characteristic foliation divided by the same dividing set (the thick circle). The foliation in the figure has a collection of singular points at the top and bottom of the boundary circle and on the horizontal diameter (near the endpoints and near the center). A family of vertical leaves separates the right and the left half-disks. The vertical leaves go from singular points at the top (respectively bottom) of the boundary circle to singular points near the center. The right and left halves of the disk are foliated by arcs connecting singularities near the endpoints of the horizontal diameter and those near the center of \mathcal{D}_{ot} . The arcs located near $\partial\mathcal{D}_{\text{ot}}$ have vertical tangencies, thus the union of an arc on the top and the corresponding arc on the bottom (together with the singular points they connect) is a smooth Legendrian knot. The half-disk D_I on the left side of \mathcal{D}_{ot} is cut off by the leftmost vertical Legendrian arc.

First, assume that the part of $\partial\mathcal{D}_{\text{ot}}$ where the connected sum $L \# \partial\mathcal{D}_{\text{ot}}$ was formed lies in ∂D_I , and consider the Legendrian connected sum $L \# \partial D_I$. Make the annulus $A \# D_I$ convex by a small isotopy fixing the boundary (we will keep the same notation for the perturbed surface). This is possible because tb of each boundary component is negative: $\text{tb}(L_0) = -1$ and $\text{tb}(L \# \partial D_I) = \text{tb}(L) + \text{tb}(\partial D_I) - 1 = -1$. Here, we are using [15, Proposition 3.1], where the convex perturbation is described as a two-stage process. First, a C^0 -small isotopy is done near the Legendrian boundary of the surface to put a collar neighborhood of the boundary into standard form. Second, a C^∞ -small isotopy, supported away from the boundary, makes the surface convex. The first perturbation uses a model neighborhood of the Legendrian boundary, and introduces a




Figure 4: Connected sum with the boundary of an overtwisted disk destabilizes a Legendrian knot

certain number of singularities with uniform rotation of the contact structure between them. If a collar neighborhood for part of the boundary is already in standard form, this part can be kept fixed during the first isotopy. (A close examination of Honda's proof shows that the same arguments go through in our case even though the boundary is only piecewise smooth. The key observation here is that the smooth Legendrian knots approximating the boundary of D_l form an annulus which is convex and in standard form in the terminology of [15, Proposition 3.1].) Therefore, we can assume that the isotopy that makes $A \# D_l$ convex fixes a neighborhood of the diameter of D_{ot} separating D_l and D_r , more precisely, that all the straight Legendrian arcs between the two half-disks are kept fixed, as well as the half-disk D_r .

The convex surface $A \# D_l$ (or rather, a surface with the same characteristic foliation) can be found in a tight contact manifold. Indeed, the dividing set on D_l is the same as that on a bypass. Bypasses exist in tight contact manifolds, thus by Giroux Flexibility we can find a surface with foliation isomorphic to that on D_l . Furthermore, in a tight neighborhood of a bypass we can find both an annulus bounded by two small unknots and the strip needed to form the connected sum. Now, consider the dividing set Γ on $A \# D_l$. Tightness of a neighborhood of $A \# D_l$ implies that Γ can have no homotopically trivial closed components. By (1), Γ intersects both components of the boundary of $A \# D_l$ at two points. This gives the following possibilities for the dividing set: either (i) Γ consists of two arcs, each connects a point on L_0 to a point on $L \# \partial D_l$, or (ii) Γ consists of two boundary-parallel arcs, plus possibly a number of closed curves running along the core of the annulus.

To rule out the second possibility, we argue as follows. Since L_0 is a small unknot, we can find a disk U bounded by L_0 , such that U is contained in the tight neighborhood of the bypass D_l . We can assume that U is convex; then its dividing set is given by a single arc connecting two points on L_0 . Moreover, we can assume that $U \cup A \# D_l$ is

a convex surface (with a tight neighborhood). But then, if there is a boundary-parallel dividing curve on $A \# D_l$ connecting two points of L_0 , the surface $U \cup A \# D_l$ would have a homotopically trivial closed component of the dividing set, which contradicts the tightness of the neighborhood.

Once we know that on $A \# D_l$ the dividing set Γ consists of two parallel arcs running from L_0 to $L \# \partial D_l$, it is clear that after we attach the other half of the overtwisted disk, the convex surface $A \# \mathcal{D}_{ot}$ will have the dividing set as in Figure 4. In conclusion, the knot $L \# \mathcal{D}_{ot}$ is a destabilization of L . □

3 Higher dimensions: Definition of the plastikstufe

Many obstructions of fillability of higher-dimensional contact manifolds have been found in the recent past; see [22] and [17]. Many of these obstructions are modeled on the overtwisted disk and take the shape of particular submanifolds in the given contact manifold. The initial incarnation of this type of obstruction is the plastikstufe [22]:

Definition 3.1 Let (M, ξ) be a contact manifold of dimension $2n + 1$, and let B be a closed $(n - 1)$ -manifold. A plastikstufe \mathcal{P}_B with core B is a submanifold

$$\mathcal{P}_B = D^2 \times B \hookrightarrow M$$

such that $\xi \cap T\mathcal{P}_B$ is a singular foliation that is tangent to the fibers $\{z\} \times B$ for every $z \in D^2$, and that restricts on every slice $D^2 \times \{b\}$ to the foliation of the overtwisted disk sketched in Figure 1. A plastikstufe \mathcal{P}_B is called a *small plastikstufe* if there is an embedded open ball in (M, ξ) containing \mathcal{P}_B . A contact manifold that contains a plastikstufe is called *PS-overtwisted*.




Figure 5: The core of a plastikstufe in a contact 5-manifold is always a circle

A plastikstufe determines the germ of the contact structure on its neighborhood [21]. More precisely, let \mathcal{D}_{ot} be an overtwisted disk in $(\mathbb{R}^3_{ot}, \alpha_{ot})$ with $\alpha_{ot} = \cos rdz +$

$r \sin r d\varphi$ written in cylindrical coordinates. Then a plastikstufe \mathcal{P}_B has a neighborhood U_{PS} contactomorphic to a neighborhood of $\mathcal{D}_{ot} \times \{0\text{-section}\}$ in

$$(\mathbb{R}^3_{ot} \times T^*B, \alpha_{PS} = \alpha_{ot} + \lambda_{can}),$$

with $\lambda_{can} = -p dq$ the canonical 1-form on T^*B . (For purposes of the present paper, we could even assume the existence of a standard neighborhood of the plastikstufe as part of definition of a PS -overtwisted manifold.) In the next section, we will prove that Legendrians become flexible in the presence of a plastikstufe, at least if the latter satisfies some technical conditions.

4 h -principles, Legendrians and plastikstufe

In this section, we review the definition of loose Legendrians and prove [Theorem 1.1](#). We begin with some background on Gromov’s h -principle for subcritical isotropic embeddings; see [\[14\]](#) and [\[7\]](#).

4.1 Formal isotropic embeddings and immersions

Note that an isotropic embedding $f: S \rightarrow (M, \xi)$ induces a monomorphism $df: TS \rightarrow TM$ such that $d_x f(T_x S)$ is a isotropic plane in the symplectic space $(\xi_{f(x)}, d\alpha)$ for every $x \in S$. If we choose a compatible almost complex structure J on (M, ξ) , then $d_x f(T_x S)$ is a totally real subspace in $(\xi_{f(x)}, J)$. Thus, we can think of $df: TS \rightarrow \xi$ as a isotropic monomorphism, or as a totally real monomorphism if an almost complex structure is chosen. (Note that in the latter case, the complexification $df^{\mathbb{C}}: T_{\mathbb{C}}S \rightarrow \xi_{f(x)}$ with $T_{\mathbb{C}}S = TS \otimes \mathbb{C}$ gives a complex monomorphism in every fiber.)

Suppose that $g, h: S \rightarrow M$ are two Legendrian immersions. If there is a regular homotopy f_t of Legendrian immersions that connects g and h , then the maps $dg: TS \rightarrow \xi$ and $dh: TS \rightarrow \xi$ are homotopic through $F_t = df_t$.

As a generalization of isotropic isotopies, we recall the definition of formal isotopy of isotropic embeddings. We say that isotropic embeddings $f_0, f_1: S \rightarrow M$ are *formally isotopic* if

- (a') there is a smooth isotopy $f_t: S \rightarrow M$ of embeddings connecting f_0 and f_1 ,
- (b') there is a homotopy F_t of isotropic monomorphisms over f_t so that $df_0 = F_0, df_1 = F_1: TS \rightarrow \xi$, and
- (c') the path of isotropic monomorphisms F_t is homotopic to df_t through paths of monomorphisms with fixed endpoints df_0, df_1 .

By a result of Gromov [14], formally isotopic *subcritical* isotropic embeddings are isotropically isotopic. In general, an h -principle fails for Legendrian embeddings, but it holds for the special class of *loose* Legendrians [20], which we review in the next subsection.

We make a similar definition for Legendrian immersions. Given two Legendrian immersions $f_0, f_1: \Lambda \rightarrow M$, assume that

- (a) there is a homotopy of continuous maps $f_t: \Lambda \rightarrow M$ connecting f_0 and f_1 , and
- (b) a homotopy F_t of Lagrangian monomorphisms over f_t so that $df_0 = F_0, df_1 = F_1: T\Lambda \rightarrow \xi$.

In this case, we say that the Legendrian immersions f_0 and f_1 are *formally homotopic*.

It is also shown in [14] that Legendrian immersions satisfy an h -principle, that is, any formally homotopic Legendrian immersions are homotopic through Legendrian immersions. We do not make use of this h -principle, but formal homotopies of Legendrian immersions will be needed throughout the paper. In particular, formal homotopies are used below in the definition of relative rotation class.

Remark 4.1 The notion of formal isotopy carries over verbatim to the case where ξ is an almost contact structure. Moreover, consider a path of almost contact structures ξ_t that starts and ends at honest contact structures ξ_0 and ξ_1 , respectively. Suppose that $g, h: \Lambda \rightarrow M$ are embeddings that are Legendrian for both ξ_0 and ξ_1 , and the bundle maps dg and dh are Lagrangian for every almost contact structure ξ_t . Then g and h are formally homotopic with respect to ξ_0 if and only if they are formally homotopic with respect to ξ_1 . In this case, Gromov's h -principle implies that g and h are homotopic through immersions that are Legendrian with respect to ξ_0 if and only if they are homotopic through Legendrian immersions with respect to ξ_1 .

Remark 4.2 It is useful to reinterpret condition (b) in the definition of formal homotopy in different terms for the case where the image of the Legendrian immersions $g, h: \Lambda \rightarrow M$ is contained in some open ball $U \subset M$. Fix a complex structure J on ξ that is tamed by the conformal symplectic structure given by $d\alpha|_\xi$, and choose a J -complex trivialization of (ξ, J) over the ball U ; both J and the trivialization will be unique up to homotopy. In particular, these choices allow us to identify $(\xi|_U, J)$ with the trivial bundle $\mathbb{C}^n \times U \rightarrow U$.

If $g: \Lambda \rightarrow (U, \xi)$ is a Legendrian immersion in U , we can write the differential dg with respect to the trivialization chosen above, as a map $dg: T\Lambda \rightarrow \mathbb{C}^n$, and furthermore since $dg(T_x\Lambda)$ is a totally real subspace in $(\xi_{g(x)}, J)$, the complexification

$$dg^{\mathbb{C}}: T_{\mathbb{C}}\Lambda \longrightarrow \mathbb{C}^n$$

gives a complex isomorphism in every fiber.

Given now a second Legendrian immersion $h: \Lambda \rightarrow U$, we can relate the two maps $dg^{\mathbb{C}}$ and $dh^{\mathbb{C}}$ by a map $\varphi: \Lambda \rightarrow \text{GL}(n, \mathbb{C})$ defined by the identity

$$(2) \quad dh_x^{\mathbb{C}} = \varphi(x) \cdot dg_x^{\mathbb{C}}$$

for every $x \in \Lambda$. The homotopy class of φ is called the *relative rotation class* of the Legendrian immersion h relative to g . The relative rotation class is an element of $[\Lambda, \text{GL}(n, \mathbb{C})] \cong [\Lambda, U(n)]$. It is clear that the maps $dg^{\mathbb{C}}: T\Lambda \otimes \mathbb{C} \rightarrow \xi$ and $dh^{\mathbb{C}}: T\Lambda \otimes \mathbb{C} \rightarrow \xi$ are homotopic through fiberwise complex isomorphisms if and only if their relative rotation class vanishes.

4.2 Loose Legendrians

We define loose Legendrian embeddings [20] (or loose Legendrians for short) by requiring that they possess the following model chart.

Definition 4.3 Let $n > 1$. In $(\mathbb{R}^3, \xi_{\text{std}} = \ker(dz - y dx))$, let L_0 be the Legendrian curve

$$t \mapsto (t^2, \frac{15}{4}(t^3 - t), \frac{3}{2}t^5 - \frac{5}{2}t^3),$$

where t is in an open interval containing $[-\sqrt{5}/\sqrt{3}, \sqrt{5}/\sqrt{3}]$. Let $U \subseteq \mathbb{R}^3_{\text{std}}$ be some convex open set containing L_0 (see the right diagram of Figure 2). In $T^*\mathbb{R}^{n-1}$ let Z be the Lagrangian zero section $\{\mathbf{p} = 0\}$, and let V_ρ be the open set $\{|\mathbf{p}| < \rho, |\mathbf{q}| < \rho\}$. Then $U \times V_\rho$ is canonically identified with an open set in $(\mathbb{R}^{2n+1}, \xi_{\text{std}})$, and $L_0 \times Z$ is Legendrian in this contact Darboux chart. If $\rho > 1$, we call the relative pair $(U \times V_\rho, L_0 \times Z)$ a *loose chart*. A connected Legendrian manifold $\Lambda \subseteq (M, \xi)$ with $\dim \Lambda > 1$ is called *loose* if there is a Darboux chart $U \subseteq M$ so that $(U, \Lambda \cap U)$ is a loose chart.

Note that the above definition contains a size restriction $\rho > 1$ on the chart parameter ρ . This is a key condition: indeed, any Legendrian can be shown to have a model chart with sufficiently small ρ .

Remark 4.4 The above definition is slightly different than the one given in [20], because here we are fixing a concrete stabilization with action 2. There is no loss of generality here, due to the existence of the contactomorphism $(x_i, y_i, z) \mapsto (cx_i, cy_i, c^2z)$ for any constant c .

It is not a priori clear why a chart with a small $\rho < 1$ is not isomorphic to a chart with large $\rho > 1$, but this must be true because loose Legendrians satisfy the following *h*-principle:

Theorem 4.5 [20] *If two loose Legendrians Λ_0, Λ_1 in a contact manifold of dimension $2n + 1 > 3$ are formally isotopic, then they are isotopic through Legendrian embeddings.* \square

In other words, loose Legendrians are flexible, ie they are classified up to Legendrian isotopy by easy to calculate invariants coming from smooth topology and bundle theory. Recall that for general Legendrian embeddings, a similar h -principle does not hold in any dimension. Indeed, it is known that holomorphic curve invariants detect Legendrian rigidity in many examples where two knots are formally isotopic but Legendrian nonisotopic [4]. (This also tells us that the requirement $\rho > 1$ imposes a nontrivial restriction.)

4.3 The proof of Theorem 1.1

Our strategy for proving Theorem 1.1 is quite simple. The motivation comes from Theorem 2.1: in every overtwisted \mathbb{R}^3 -slice we can isotope a given Legendrian embedding of a knot to an embedding where its front projection has a “kink”. We can think of a plastikstufe as a “product” family of overtwisted disks and perform a family of the above isotopies for slices $\Lambda \cap \mathbb{R}^3 \times \{b\}$ of a given Legendrian Λ (near the plastikstufe). This product isotopy would produce a chart required by Definition 4.3 and verify that the given Legendrian is, indeed, loose. For us to carry out this plan, the Legendrian Λ , perhaps after an isotopy, must contain a codimension 0 submanifold Λ_0 diffeomorphic to $B \times [0, 1]$, where B is the core of the plastikstufe \mathcal{P}_B . Moreover, Λ_0 must be in a “product position” in some standard neighborhood $U_{PS} = N_\epsilon \mathcal{D}_{ot} \times T_\epsilon^* B$ of \mathcal{P}_B :

$$\begin{aligned}\mathcal{P}_B &= \mathcal{D}_{ot} \times \{0\text{-section}\} \subset U_{PS}, \\ \Lambda_0 &= K \times \{0\text{-section}\} \subset U_{PS},\end{aligned}$$

where $K \subset N_\epsilon \mathcal{D}_{ot}$ is a Legendrian arc. In other words, we must be able to move the given Legendrian Λ by an ambient contact isotopy so that a product strip of the form $K \times \{0\text{-section}\}$ would be contained in Λ . In Lemma 4.8, we will show that this is possible if the plastikstufe has a spherical core and *trivial rotation*, a notion we now define.

A *leaf ribbon* in a plastikstufe \mathcal{P}_B with core B is a thin ribbon diffeomorphic to $B \times (0, 1)$, obtained by shrinking a leaf of \mathcal{P}_B within itself into the standard neighborhood of the core B . Notice that all such ribbons are Legendrian isotopic: ribbons contained in the same leaf are all deformations of the same leaf and ribbons in different leaves can be related by rotating around the core.

From now on, we restrict to plastikstufes with core B diffeomorphic to S^{n-1} (see Remark 5.4 for a brief discussion of the general case). We will compare the formal

Legendrian homotopy class of their leaf ribbons to the class represented by a punctured Legendrian disk.

Definition 4.6 For a small plastikstufe \mathcal{P} with spherical core, we define the rotation class of \mathcal{P} to be the relative rotation class between a leaf ribbon of \mathcal{P} and a punctured Legendrian disk. We say that \mathcal{P} has *trivial rotation* if this class vanishes.

It is not hard to show that any two Legendrian disks in (M, ξ) are Legendrian isotopic. Therefore, the rotation class of a small plastikstufe with spherical core is well-defined.

Remark 4.7 The rotation class of a plastikstufe with core $B = S^{n-1}$ is an element of

$$[B \times [0, 1], \text{GL}(n, \mathbb{C})] \cong [B, U(n)] = \pi_{n-1}(U(n)) = \begin{cases} \mathbb{Z} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Thus, in contact manifolds of dimension $2n + 1 = 4m + 3$, every small plastikstufe with spherical core has trivial rotation. We will see in Section 5 that in the appropriate sense, the rotation of a plastikstufe in dimensions $2n + 1 = 4m + 1 > 5$ is determined by the rotation of its core, while in dimension 5 it is the rotation of the leaf direction that determines the rotation of \mathcal{P} .

Given a plastikstufe \mathcal{P} with trivial rotation, and a Legendrian Λ , we now isotope Λ towards \mathcal{P} .

Lemma 4.8 *Suppose (M^{2n+1}, ξ) contains a small plastikstufe $\mathcal{P} \subset (M, \xi)$ with spherical core $B = S^{n-1}$, and \mathcal{P} has trivial rotation. Let $\Lambda \subset M$ be a Legendrian disjoint from \mathcal{P} . Then there exists an ambient contact isotopy of M , fixed near \mathcal{P} , that takes a submanifold Λ_0 of Λ diffeomorphic to $S^{n-1} \times [0, 1]$ to a product strip $\Lambda_1 = [0, 1] \times \{0\text{-section}\}$ near \mathcal{P} .*

Proof We can assume that the plastikstufe \mathcal{P} and at least some part of Λ are contained in an open ball U . Fix a strip $\Lambda_1 = S^{n-1} \times I$ which is isotopic to a leaf ribbon for \mathcal{P} but lies in the complement of \mathcal{P} . (Here and below, I denotes a closed interval.) Choose a small Legendrian disk in $\Lambda \cap U$, and let $\Lambda_0 = S^{n-1} \times I$ be an annular Legendrian strip in this disk, so that the interior of Λ_0 is isotopic to the punctured disk.

Now we have two Legendrian embeddings $f_0, f_1: S^{n-1} \times I \rightarrow M$, such that Λ_i is the image of f_i , $i = 0, 1$. We would like to connect these by a path of Legendrian embeddings. To this end, we will apply Gromov’s h -principle for subcritical isotropic embeddings; see [14] and [7]. (Alternatively, we could have used the h -principle for open Legendrian embeddings [14]).

To use the subcritical isotropic h -principle, restrict attention to the core spheres $S_i = f_i(S^{n-1} \times \{c\})$, $i = 0, 1$, where c is a point inside I . (In what follows, we will be assuming that c is close to an endpoint of I . It will be clear from the context where we want the core spheres to be; a particular choice is unimportant, since any two are isotropically isotopic.) Let $f_i|_{\text{core}}: S^{n-1} \rightarrow M$ denote the isotropic embeddings given by the corresponding restriction maps. We first construct a formal isotropic isotopy between these embeddings.

Reparameterizing I , we can think of f_0 and f_1 as maps

$$f_0: S^{n-1} \times [0, \delta] \longrightarrow M, \quad f_1: S^{n-1} \times [1 - \delta, 1] \longrightarrow M,$$

for some small $\delta > 0$. Choosing a trivialization of ξ in U as in [Remark 4.2](#), we consider the maps

$$df_0^{\mathbb{C}}: T_{\mathbb{C}}(S^{n-1} \times [0, \delta]) \longrightarrow \mathbb{C}^n, \quad df_1^{\mathbb{C}}: T_{\mathbb{C}}(S^{n-1} \times [1 - \delta, 1]) \longrightarrow \mathbb{C}^n.$$

The map $df_0^{\mathbb{C}}$ is homotopic to the map

$$G_0^{\mathbb{C}}: T(S^{n-1} \times [0, \delta]) \longrightarrow \mathbb{C}^n, \quad G_0^{\mathbb{C}}(x, t) = df_0^{\mathbb{C}}(x, \delta), \quad 0 \leq t \leq \delta.$$

Similarly, $df_1^{\mathbb{C}}$ is homotopic to the map

$$G_1^{\mathbb{C}}: T(S^{n-1} \times [1 - \delta, 1]) \longrightarrow \mathbb{C}^n, \quad G_1^{\mathbb{C}}(x, t) = df_1^{\mathbb{C}}(x, 1 - \delta), \quad 1 - \delta \leq t \leq 1.$$

Then, we can write

$$(3) \quad df_0^{\mathbb{C}}(x, \delta) = \psi(x)df_1^{\mathbb{C}}(x, 1 - \delta)$$

for a map $\psi: S^{n-1} \rightarrow \text{GL}(n, \mathbb{C})$; cf [Remark 4.2](#).

Because \mathcal{P} has trivial rotation and Λ_1 is isotopic to a leaf ribbon of \mathcal{P} , it follows that Λ_0 and Λ_1 are formally Legendrian homotopic. Then, the map ψ is homotopic to the map sending each point of S^{n-1} to the identity transformation in $\text{GL}(n, \mathbb{C})$. Using this homotopy, we can construct a map

$$G^{\mathbb{C}}: T_{\mathbb{C}}(S^{n-1} \times [0, 1]) \longrightarrow \mathbb{C}^n$$

which gives a complex isomorphism in every fiber and coincides with $df_0^{\mathbb{C}}$ near $S^{n-1} \times \{0\}$ and with $df_1^{\mathbb{C}}$ near $S^{n-1} \times \{1\}$. The real part G of $G^{\mathbb{C}}$ is a Lagrangian monomorphism $T(S^{n-1} \times [0, 1]) \rightarrow \xi$, restricting to df_0 and df_1 near the ends of the cylinder $S^{n-1} \times [0, 1]$.

The Smale–Hirsch Immersion Theorem (see eg [\[7\]](#)) implies that G is homotopic to the differential dg of an immersion $g: S^{n-1} \times [0, 1] \rightarrow M$ such that $g = f_0$ near $S^{n-1} \times \{0\}$ and $g = f_1$ near $S^{n-1} \times \{1\}$.

But $S^{n-1} \times [0, 1]$ has dimension n , and M has dimension $2n + 1$, so we can perturb g into general position by homotoping it (through immersions) to an embedding \tilde{g} . We can assume that this homotopy fixes the cylinder near the ends, and that the core spheres we took are close enough to the ends of the cylinder.

Clearly, the maps $\tilde{g}_t = \tilde{g}(\cdot, t)$ are smooth embeddings connecting $f_0|_{\text{core}}$ and $f_1|_{\text{core}}$. The maps $G_t = G(\cdot, t)|_{T(S^{n-1} \times \{t\})}$ are isotropic monomorphisms covering \tilde{g}_t ; moreover, G_t coincides with $df_0|_{TS^{n-1}}$ respectively $df_1|_{TS^{n-1}}$ near 0 respectively 1, and the path G_t is homotopic to $d\tilde{g}$ rel endpoints. In short, we have a formal isotopy of subcritical isotropic embeddings. By Gromov’s h -principle, it follows that $f_0|_{\text{core}}$ and $f_1|_{\text{core}}$ can be connected by a path \tilde{f}_t of isotropic embeddings.

The next step is to upgrade \tilde{f}_t to a path of *framed* isotropic embeddings. For an $(n - 1)$ -dimensional isotropic submanifold S of (M^{2n+1}, ξ) , we consider a framing by a nonvanishing vector field $X: S \rightarrow \xi$, such that $X(s) \in (T_s S)^\perp \setminus T_s S$ for every point $s \in S$. (Here, $(T_s S)^\perp$ stands for the symplectic orthogonal complement of $T_s S$ in $(\xi, d\alpha)$; this subspace contains $T_s S$ and has dimension $n + 1$.) In other words, we look at framings such that at every point of S the direct sum $TS \oplus \text{Span}(X)$ is a Lagrangian subspace of ξ . For an isotropic submanifold contained in a ball, we can use a trivialization to find two vector fields v, v' such that at every point $(T_s S)^\perp = T_s S \oplus \text{Span}(v, v')$, and think of the vector field X as living in $\text{Span}(v, v')$. Since X does not vanish, after normalizing we can think of the framing as a map $X: S \rightarrow S^1$.

Notice that the original embeddings f_0, f_1 of $S^{n-1} \times I$ endow the restrictions $f_i|_{\text{core}}$, $i = 0, 1$ with a framing X_i given by $X_i = df_i$ (the I -direction). We would like to find framings X_t for the isotropic embeddings \tilde{f}_t , so that the path (\tilde{f}_t, X_t) connects $(f_0|_{\text{core}}, X_0)$ and $(f_1|_{\text{core}}, X_1)$. If $n > 2$, then the maps $X_0: f_0(S^{n-1}) \rightarrow S^1$ and $X_1: f_1(S^{n-1}) \rightarrow S^1$ are necessarily homotopic, so we can find the desired path X_t . In the case $n = 2$, recall that the differentials df_0 and df_1 of the embeddings of $S^{n-1} \times I$ were connected by a path of Lagrangian monomorphisms F_t . Then, F_t (the I -direction) is a homotopy between X_0 and X_1 .

The framed isotropic isotopy of the core spheres $f_0(S^{n-1})$ and $f_1(S^{n-1})$ immediately gives a Legendrian isotopy of the original annuli Λ_0 and Λ_1 . Indeed, Λ_0 and Λ_1 can be shrunk and flattened (by a Legendrian isotopy) to thin annuli around their core spheres, linear in the direction of X_0 , respectively X_1 . These can be connected by a family of thin Legendrian annuli built on the spheres $\tilde{f}_t(S^{n-1})$, spanning a linear strip in the X_t direction. By the ambient isotopy theorem, (see eg [11, Theorem 2.6.2]), the resulting isotopy of Legendrian annuli can be realized by a compactly supported contact isotopy of (M, ξ) , concluding the proof. □

Lemma 4.9 Suppose that Λ is a Legendrian in (M, ξ) which lies in the complement of a small plastikstufe with spherical core and trivial rotation. Then there is an open subset V of M with topology $S^{n-1} \times D^{n+2}$ such that the pair $(V, V \cap \Lambda)$ is contactomorphic to $(D_{\text{std}}^3 \times D^*S^{n-1}, \Lambda_0)$, where Λ_0 is the Cartesian product of a stabilization in D_{std}^3 and the zero section in D^*S^{n-1} .

Proof By Lemma 4.8, after an appropriate Legendrian isotopy, in the standard neighborhood of the plastikstufe $\mathbb{R}_{OT}^3 \times D^*S^{n-1}$ the Legendrian Λ is given by $K \times S^{n-1}$. Our entire picture is S^{n-1} -symmetric, so we apply Theorem 2.1 simultaneously for each point in S^{n-1} . □

Lemma 4.10 Suppose that $\Lambda \subset (M, \xi)$ is Legendrian, and there exists an open set $V \subset M$ such that $(V, V \cap \Lambda)$ is contactomorphic to $(D_{\text{std}}^3 \times D^*S^{n-1}, \Lambda_0)$, where Λ_0 is the Cartesian product of a stabilization in D_{std}^3 and the zero section in D^*S^{n-1} . Then Λ is loose.

Proof For any metric on S^{n-1} , D^*S^{n-1} contains a subset symplectomorphic to $U_\rho = \{|q| < \rho, |p| < \rho\} \subset T^*\mathbb{R}^{n-1}$, for some ρ . Fix $\epsilon > 0$ so that $\epsilon < \rho$. In D_{std}^3 , we can isotope any stabilization by compactly supported isotopy to the stabilization L_ϵ given by

$$t \mapsto (\epsilon t^2, \frac{15\epsilon}{4}(t^3 - t), \frac{\epsilon^2}{2}(3t^5 - 5t^3)).$$

We apply this isotopy to Λ_0 simultaneously for each point in S^{n-1} , showing that $(D_{\text{std}}^3 \times D^*S^{n-1}, \Lambda_0)$ is contactomorphic to $(D_{\text{std}}^3 \times D^*S^{n-1}, L_\epsilon \times Z)$, where Z is the zero section in D^*S^{n-1} . Then the subset $D_{\text{std}}^3 \times U_\rho$ is a loose chart, which can be seen by reparametrizing our coordinates by the contactomorphism $(x_i, y_i, z) \mapsto (x_i/\epsilon, y_i/\epsilon, z/\epsilon^2)$. □

5 Small plastikstufes with trivial rotation

For Theorem 1.1 to be useful, we need a supply of contact manifolds containing small plastikstufes satisfying the hypotheses of Lemma 4.8, ie having spherical core and trivial rotation. We will use the following result of Etnyre–Pancholi [10] to find suitable examples.

Theorem 5.1 [10] Let (M, ξ) be a contact manifold of dimension $2n + 1$ and let B be an $(n - 1)$ -dimensional isotropic submanifold with trivial conformal symplectic normal bundle. Then we may alter ξ to a contact structure ξ' that contains a small

plastikstufe with core B such that ξ and ξ' are homotopic through almost contact structures.

In fact, choosing any neighborhood U of B , we can find a smaller neighborhood U' of B such that both the modification and the homotopy only affect ξ in $U \setminus \overline{U'}$. \square

In our application we need a refinement of this result, in which one also determines the rotation of the plastikstufe.

Proposition 5.2 *The modification of Theorem 5.1 can be applied in any contact manifold to produce a small plastikstufe with spherical core and trivial rotation.*

Proof The only part of the statement which needs justification is that we can achieve triviality of the rotation class. Indeed, choose a Legendrian disk D in (M^{2n+1}, ξ) and apply the construction from the proof of Theorem 5.1 to any cooriented codimension 1 submanifold B of D . This B is automatically isotropic, and its symplectic normal bundle is spanned by (X_n, JX_n) where X_n denotes a vector field in D that is transverse to B , and J is a compatible almost complex structure on ξ . Since we can do this construction inside a small ball around the Legendrian disk D , the resulting contact structure ξ' will contain a small plastikstufe \mathcal{P}_B .

Assume for the rest of the proof that B is a small sphere S^{n-1} that bounds a Legendrian disk in L . (See Remark 5.4 below for discussion of the more general case.) By Remark 4.7, the rotation class of a small plastikstufe with spherical core necessarily vanishes in contact manifolds of dimension $4m + 3$, but we still need to study the remaining dimensions $4m + 1$. Below, we will first show that the rotation for any plastikstufe constructed by Theorem 5.1 (on a core $B = S^{n-1}$ that bounds a Legendrian disk before creating the plastikstufe) is trivial if $\dim M > 5$. (This phenomenon already manifested itself in the proof of Theorem 1.1, where any two framings on isotropic spheres were automatically homotopic if $\dim M > 5$.) Finally, we will show that using the Etnyre–Pancholi Theorem 5.1 in dimension 5 with more care, we can produce plastikstufes with arbitrary rotation class, including trivial rotation.

We can assume that the modification to ξ' has been done in a sufficiently small neighborhood of S^{n-1} so that $\xi' = \xi$ both close to the center of the auxiliary Legendrian disk D chosen above, and outside of an embedded ball. Moreover, the Etnyre–Pancholi modification does not change the contact structure near the chosen isotropic submanifold B . Thus, close to B the leaves of the resulting plastikstufe are also Legendrian with respect to the *old* contact structure ξ . Since ξ and ξ' are homotopic, by Remark 4.1 it suffices to examine the rotation of a leaf with respect to ξ .

Choose a leaf ribbon near B and compare its rotation to the punctured Legendrian disk $D - \{0\} \cong B \times I$. Recall that by Remark 4.2 we can trivialize ξ over a ball containing the Legendrians, and then obtain the rotation class as the homotopy class of the map $\varphi: B \times I \rightarrow \text{GL}(n, \mathbb{C})$ that satisfies Equation (2), where h and g denote the two embeddings we wish to compare. In fact, because $B \times I$ deformation retracts to B , it suffices to study the restriction of this map to $B \times \{\frac{1}{2}\}$.

Then the two embeddings are the same over B and only differ in the remaining direction. Over B the tangent bundle to the leaf ribbon splits as $TB \oplus \langle X_l \rangle$, where X_l is the leaf direction, and the tangent bundle to the flat annulus splits as $TB \oplus \langle X_n \rangle$, where X_n is the normal to B in the disk D . Therefore, for any $q \in B$ we have

$$\varphi(q) = \begin{pmatrix} & * & \\ \text{Id} & \vdots & \\ & * & \\ 0 \cdots 0 & \kappa(q) & \end{pmatrix},$$

where $\kappa: B \rightarrow \mathbb{C}^*$ is a continuous function. This helps us understand the homotopy class of φ in $[B, \text{GL}(n, \mathbb{C})]$. We consider two cases.

Case 1 Suppose $2n + 1 > 5$. Then the function $\kappa: B \rightarrow \mathbb{C}^*$ is homotopic to 1, since by assumption $B = S^{n-1}$ with $n - 1 \geq 2$. It follows that φ represents the trivial class in $[B, \text{GL}(n, \mathbb{C})]$, so the plastikstufe \mathcal{P}_B has trivial rotation.

Case 2 Suppose $2n + 1 = 5$. Now, the function κ , and therefore φ , may represent a nontrivial homotopy class. It turns out that this class depends on the choice of coordinates in the Etnyre–Pancholi construction. A reparametrization will allow us to adjust the rotation class. To see this, we first review the setup of the construction of [10]. Given an isotropic manifold B with trivial symplectic conormal bundle, we use the isotropic neighborhood theorem to identify a neighborhood of B in (M, ξ) with $T^*B \times \mathbb{R}^3$, equipped with the contact form $pdq + dz + r^2d\theta$. Choosing a transverse curve $\gamma(t)$ in \mathbb{R}^3 , we can further identify a neighborhood of $B \times \gamma$ in $T^*B \times \mathbb{R}^3$ with

$$(T^*B \times S^1 \times D^2, \ker(pdq + dt + r^2d\theta)).$$

Here, B is the 0–section of T^*B , pdq is the canonical 1–form on T^*B , S^1 is the transverse curve $\gamma(t)$, and D^2 is an open disk with polar coordinates (r, θ) . Since $\dim M = 5$, we have $B = S^1$, and the coordinate q varies in the unit circle. The generalized Lutz twist from the Etnyre–Pancholi construction produces in each of the slices $T^*B \times \{t_0\} \times D^2$ a plastikstufe \mathcal{P}_{t_0} whose core is $B \times \{t_0\}$, where B is the zero section as before. Near the core, the leaves of \mathcal{P}_{t_0} are given by $\{\theta = \text{const}\}$.

We can define for any $a \in \mathbb{Z}$ a contactomorphism

$$F_a: T^*B \times S^1 \times \mathbb{R}^2 \longrightarrow T^*B \times S^1 \times \mathbb{R}^2$$

by $F_a(p, q, t, r e^{i\theta}) = (p - ar^2, q, t, r e^{i(\theta+aq)})$ that preserves $p dq + dz + r^2 d\theta$. Note that the zero section $\{p = 0, r = 0\}$ is preserved by F_a .

Instead of applying now the generalized Lutz twist of [10] to the initial model neighborhood of B we have chosen in the Legendrian disk D above, we use first a restriction of the map F_a to modify the coordinates of the model neighborhood. This means that for a fixed t_0 , the construction will produce a plastikstufe with the same core for either choice of coordinates. The leaf direction near the core, however, is different: we get $\{\theta = \text{const}\}$ for the plastikstufe \mathcal{P} constructed with the old coordinates, and $\theta' = \theta + aq = \text{const}$ for \mathcal{P}' constructed in the new coordinates. Intuitively, the leaf direction of \mathcal{P}' rotates a times with respect to the leaf direction of \mathcal{P} as we move around the circle $S^1 = B$.

We now compute the relative rotation class between \mathcal{P} and \mathcal{P}' . For this, we compare two trivialisations of the bundle $\xi \rightarrow B = S^1$ corresponding to the (Legendrian) leaves of our plastikstufes. With a choice of a compatible almost complex structure J on ξ , each plastikstufe yields a trivialization of $\xi|_B$ given by

$$(\text{core } S^1\text{-direction}, J(\text{core } S^1\text{-direction}), \text{leaf direction}, J(\text{leaf direction})).$$

Since the S^1 -direction for \mathcal{P} and \mathcal{P}' is the same and the leaf directions differ as described above, the two trivialisations are related by the map $\varphi: S^1 \rightarrow U(2)$ given by

$$\varphi(e^{iq}) = \begin{pmatrix} 1 & 0 \\ 0 & e^{iaq} \end{pmatrix}.$$

Different values of a give maps lying in different homotopy classes in $[S^1, U(2)] = \pi_1(U(2)) = \mathbb{Z}$. More precisely, we can realize any given element in $\pi_1(U(2))$ by an appropriate choice of a . In particular, we can realize the class given by a flat Legendrian annulus (ie the one contained in the Legendrian disk D). It follows that by an appropriate choice of coordinates, we can construct a plastikstufe with trivial rotation class. □

Remark 5.3 The previous proposition essentially shows that the Etnyre–Pancholi construction can be used to produce a plastikstufe with an arbitrary given rotation class. Indeed, any rotation class of the core can be realized by Gromov’s h -principle for subcritical isotropic embeddings. In the leaf direction, rotation is always trivial in dimensions higher than 5; in dimension 5, any given rotation can be realized by an

appropriate choice of coordinates in the Etnyre–Pancholi construction. We leave the details to the reader as we have no use for plastikstufes with nontrivial rotation.

Remark 5.4 The construction above could be generalized in a relatively easy way for a small plastikstufe \mathcal{P}_B with core B^{n-1} , as long as B^{n-1} can be (smoothly) embedded into \mathbb{R}^n . To simplify the presentation, we have chosen to restrict our attention to the spherical case; cf [Question 7.2](#).

6 Flexible Weinstein manifolds

Let $(W^{2n+2}, d\lambda)$ be an exact symplectic manifold. Given an exhausting Morse function $\varphi: W \rightarrow \mathbb{R}$, we say the triple (W, λ, φ) is a *Weinstein manifold* if the vector field V determined by $V \lrcorner d\lambda = \lambda$ is gradient-like for the Morse function φ . (If φ has finitely many critical points, we say that W has finite type.) It follows that λ is a contact form on any noncritical level set $\varphi^{-1}(c)$. Similarly, (W, λ, φ) is a *Weinstein cobordism* from (Y_0, ξ_0) to (Y_1, ξ_1) if $\partial W = Y_1 \cup -Y_0$, where Y_0 and Y_1 are level sets for φ , and V points inwards along Y_0 and outwards along Y_1 . If $Y_0 = \emptyset$, W is called a *Weinstein domain*.

Observe that because $\mathcal{L}_V \lambda = \lambda$, the flow of the vector field V expands λ exponentially. If $p \in W$ is a critical point of φ , the *descending manifold* C of p consists of all the points in W which converge to p by the flow of V . Since any vector in TC converges to the zero vector at the point p under the flow of V , but this flow also expands λ exponentially, we conclude that $\lambda|_C = 0$. Thus any descending manifold is an isotropic submanifold of W , and furthermore it intersects every level set in a contact isotropic submanifold. In particular, every critical point of φ must have index less than or equal to $n + 1$.

A Weinstein cobordism of dimension $2n + 2 > 4$ is called *flexible* if every descending manifold of index $n + 1$ intersects a lower level set along a *loose Legendrian sphere* (in other words, attaching spheres for all handles of index $n + 1$ are loose Legendrians). Here, we will assume for simplicity that all critical values of φ are distinct. Using the h -principle result from [\[20\]](#), it is shown in [\[2\]](#) that flexible Weinstein cobordisms can be completely classified by their topology. There are several related theorems proved in [\[2, Chapter 14\]](#); we state two results (in the form relevant to our case).

Theorem 6.1 [\[2\]](#) *Let (W, λ, φ) be a flexible Weinstein cobordism from $(Y_0, \ker \alpha_0)$ to $(Y_1, \ker \alpha_1)$, such that W is smoothly a product cobordism. Then after a homotopy through Weinstein structures, the cobordism (W, λ, φ) is isomorphic to $(Y_0 \times \mathbb{R}, e^t \alpha_0, t)$. In particular, $(Y_0, \ker \alpha_0)$ is contactomorphic to $(Y_1, \ker \alpha_1)$. \square*

More generally:

Theorem 6.2 [2] *Let $(W, \lambda_0, \varphi_0)$ and $(W, \lambda_1, \varphi_1)$ be two flexible Weinstein structures on a fixed cobordism W . Suppose that $\partial_-(W, \lambda_0, \varphi_0)$ is contactomorphic to $\partial_-(W, \lambda_1, \varphi_1)$, and further that $d\lambda_0$ and $d\lambda_1$ are homotopic through nondegenerate 2-forms. Then there exists a homotopy of Weinstein structures connecting $(W, \lambda_0, \varphi_0)$ and $(W, \lambda_1, \varphi_1)$. \square*

Now we are ready to prove [Theorem 1.2](#) from the introduction:

Proof of Theorem 1.2 Suppose hypothesis (1) is satisfied, ie (Y, ξ_0) and (Y, ξ_1) are the respective boundaries of the Weinstein manifolds $(W, \lambda_0, \varphi_0)$ and $(W, \lambda_1, \varphi_1)$. We then puncture each manifold, so that $(W, \lambda_j, \varphi_j)$ is a Weinstein cobordism from S_{std}^{2n+1} to (Y, ξ_j) . Let $(M^{2n+1}, \xi_{\text{PS}})$ be a PS -overtwisted manifold containing a plastikstufe \mathcal{P} satisfying the hypotheses of [Theorem 1.1](#). By connect summing this manifold onto each level set of $(W, \lambda_0, \varphi_0)$ and $(W, \lambda_1, \varphi_1)$ (away from all descending manifolds) we obtain two new Weinstein cobordisms \tilde{W}_j from (M, ξ_{PS}) to $(Y_j, \xi_j) \# (M, \xi_{\text{PS}})$. The cobordism \tilde{W}_j has the same critical points as W , and each descending manifold is disjoint from M (and in particular, from the plastikstufe \mathcal{P}) at every level set. By [Theorem 1.1](#) this implies that all descending manifolds of index $n + 1$ intersect all level sets in loose Legendrians, therefore \tilde{W}_j is a flexible Weinstein cobordism. We then complete the proof by applying [Theorem 6.2](#). The proof of the theorem under hypothesis (2) is similar; in this case we connect sum (M, ξ_{PS}) to each level set of the cobordism between (Y, ξ_0) and (Y, ξ_1) , and apply [Theorem 6.1](#). \square

7 Open questions

The questions in this section are not new, but with the results of this paper they may be formulated in a more precise way. Above we have showed that certain PS -overtwisted manifolds are flexible within the class of PS -overtwisted manifolds being Stein cobordant by a *topologically* trivial cobordism.

Question 7.1 Even if the general flexibility question seems to be out of reach, it would be interesting to tackle more special cases. For example:

(a) Is it true that the connected sum of an Ustilovsky sphere (cf [\[25\]](#)) with a suitable PS -overtwisted manifold is contactomorphic to the PS -overtwisted manifold? Note that in this case, the two contact manifolds are cobordant via a topologically nontrivial Stein cobordism.

(b) Take a model PS -overtwisted sphere $(S^{2n+1}, \xi_{\text{PS}})$. Is it true that $(S^{2n+1}, \xi_{\text{PS}} \# \xi_{\text{PS}})$ is contactomorphic to $(S^{2n+1}, \xi_{\text{PS}})$?

Question 7.2 Can we remove the technical conditions in [Theorem 1.1](#) and prove similar statements for arbitrary plastikstufes, or, more generally, for arbitrary bLobs? (Bordered Legendrian open books, or bLobs for short, were introduced in [\[17\]](#) to generalize the notion of PS -overtwistedness.)

It is not hard to see that the arguments of this paper go through in the presence of a “large” open ball in $\mathbb{R}_{\text{ot}}^3 \times \mathbb{R}^{2n}$. Moreover, some properties of such ball [\[23\]](#) resemble those of a loose chart. Is it true that a neighborhood of an arbitrary plastikstufe or bLob contains such an open ball?

A further question is about finding the correct definition of overtwistedness in higher dimensions. A very useful approach to studying contact structures in all dimensions is provided by open book decompositions [\[13\]](#).

Question 7.3 Can our results be interpreted in terms of open books to obtain some type of flexibility in that context?

References

- [1] **M Abouzaid, P Seidel**, *Altering symplectic manifolds by homologous recombination* [arXiv:1007.3281](#)
- [2] **K Cieliebak, Y Eliashberg**, *From Stein to Weinstein and back: Symplectic geometry of affine complex manifolds*, AMS Colloq. Publ. 59 (2012) [MR3012475](#)
- [3] **K Dymara**, *Legendrian knots in overtwisted contact structures on S^3* , Ann. Global Anal. Geom. 19 (2001) 293–305 [MR1828083](#)
- [4] **T Ekholm, J Etnyre, M Sullivan**, *Non-isotopic Legendrian submanifolds in \mathbb{R}^{2n+1}* , J. Differential Geom. 71 (2005) 85–128 [MR2191769](#)
- [5] **Y Eliashberg**, *Classification of overtwisted contact structures on 3-manifolds*, Invent. Math. 98 (1989) 623–637 [MR1022310](#)
- [6] **Y Eliashberg, M Fraser**, *Topologically trivial Legendrian knots*, J. Symplectic Geom. 7 (2009) 77–127 [MR2496415](#)
- [7] **Y Eliashberg, N Mishachev**, *Introduction to the h -principle*, Graduate Studies in Mathematics 48, Amer. Math. Soc. (2002) [MR1909245](#)
- [8] **J B Etnyre**, *On knots in overtwisted contact structures* [arXiv:1012.3745](#)
- [9] **J B Etnyre**, *Legendrian and transversal knots*, from: “Handbook of knot theory”, (W Menasco, M Thistlethwaite, editors), Elsevier B. V., Amsterdam (2005) 105–185 [MR2179261](#)
- [10] **J B Etnyre, DM Pancholi**, *On generalizing Lutz twists*, J. Lond. Math. Soc. 84 (2011) 670–688 [MR2855796](#)
- [11] **H Geiges**, *An introduction to contact topology*, Cambridge Studies in Advanced Mathematics 109, Cambridge Univ. Press (2008) [MR2397738](#)

- [12] **E Giroux**, *Convexité en topologie de contact*, Comment. Math. Helv. 66 (1991) 637–677 [MR1129802](#)
- [13] **E Giroux**, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, from: “Proceedings of the ICM, Vol. II”, (T Li, editor), Higher Ed. Press, Beijing (2002) 405–414 [MR1957051](#)
- [14] **M Gromov**, *Partial differential relations*, Ergeb. Math. Grenzgeb. 9, Springer, Berlin (1986) [MR864505](#)
- [15] **K Honda**, *On the classification of tight contact structures. I*, Geom. Topol. 4 (2000) 309–368 [MR1786111](#)
- [16] **Y Huang**, *A proof of the classification theorem of overtwisted contact structures via convex surface theory* [arXiv:1102.5398](#)
- [17] **P Massot, K Niederkrüger, C Wendl**, *Weak and strong fillability of higher dimensional contact manifolds*, Invent. Math. 192 (2013) 287–373 [MR3044125](#)
- [18] **M McLean**, *Personal communication*
- [19] **M McLean**, *Lefschetz fibrations and symplectic homology*, Geom. Topol. 13 (2009) 1877–1944 [MR2497314](#)
- [20] **E Murphy**, *Loose Legendrian embeddings in high dimensional contact manifolds* [arXiv:1201.2245](#)
- [21] **K Niederkrüger**, *Habilitation*, Toulouse, in preparation
- [22] **K Niederkrüger**, *The plastikstufe — a generalization of the overtwisted disk to higher dimensions*, Algebr. Geom. Topol. 6 (2006) 2473–2508 [MR2286033](#)
- [23] **K Niederkrüger, F Presas**, *Some remarks on the size of tubular neighborhoods in contact topology and fillability*, Geom. Topol. 14 (2010) 719–754 [MR2602849](#)
- [24] **P Seidel, I Smith**, *The symplectic topology of Ramanujam’s surface*, Comment. Math. Helv. 80 (2005) 859–881 [MR2182703](#)
- [25] **I Ustilovsky**, *Infinitely many contact structures on S^{4m+1}* , Internat. Math. Res. Notices (1999) 781–791 [MR1704176](#)
- [26] **T Vogel**, *Existence of Engel structures*, Ann. of Math. 169 (2009) 79–137 [MR2480602](#)

Department of Mathematics, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

Institut de math., Université Paul Sabatier – Toulouse III, 31062 Toulouse, Cedex 9, France

Department of Mathematics, Stony Brook University, Stony Brook, NY 11790, USA

Rényi Institute of Mathematics, Hungarian Academy of Sciences, 1053 Budapest, Hungary

e_murphy@mit.edu, niederkr@math.univ-toulouse.fr,

olga@math.sunysb.edu, stipsicz@renyi.hu

<http://www.math.univ-toulouse.fr/~niederkr/>,

<http://www.renyi.hu/~stipsicz>

Proposed: Yasha Eliashberg

Received: 18 December 2012

Seconded: Leonid Polterovich, Jean-Pierre Otal

Revised: 3 March 2013