

Uniform hyperbolicity of the graphs of curves

TARIK AOUGAB

Let $\mathcal{C}(S_{g,p})$ denote the curve complex of the closed orientable surface of genus g with p punctures. Masur and Minsky and subsequently Bowditch showed that $\mathcal{C}(S_{g,p})$ is δ -hyperbolic for some $\delta = \delta(g, p)$. In this paper, we show that there exists some $\delta > 0$ independent of g, p such that the curve graph $\mathcal{C}_1(S_{g,p})$ is δ -hyperbolic. Furthermore, we use the main tool in the proof of this theorem to show uniform boundedness of two other quantities which a priori grow with g and p : the curve complex distance between two vertex cycles of the same train track, and the Lipschitz constants of the map from Teichmüller space to $\mathcal{C}(S)$ sending a Riemann surface to the curve(s) of shortest extremal length.

05C12, 20F65, 57M07, 57M15, 57M20

1 Introduction

Let $S_{g,p}$ denote the orientable surface of genus g with p punctures. The *curve complex* of $S_{g,p}$, denoted $\mathcal{C}(S_{g,p})$, is the simplicial complex whose vertices are in one-to-one correspondence with isotopy classes of nonperipheral simple closed curves on $S_{g,p}$, and such that $k + 1$ vertices span a k -simplex if and only if the corresponding $k + 1$ isotopy classes can be realized disjointly on $S_{g,p}$. We make $\mathcal{C}(S)$ into a metric space by identifying each k -simplex with the standard simplex in \mathbb{R}^k with unit length edges (see Masur and Minsky [13] for more details).

Masur and Minsky showed [13] that there is some $\delta = \delta(S_{g,p})$ such that $\mathcal{C}(S_{g,p})$ is δ -hyperbolic, meaning that geodesic triangles in $\mathcal{C}(S)$ are δ -thin: any edge is contained in the δ -neighborhood of the union of the other two edges. Bowditch reproved this result in [2], showing that the hyperbolicity constant δ grows no faster than logarithmically in g and p . Let $\mathcal{C}_n(S)$ denote the n -skeleton of $\mathcal{C}(S)$. The main result of this paper is that for $\mathcal{C}_1(S)$, the *curve graph*, one may take δ to be independent of g and p .

Theorem 1.1 *There exists $k > 0$ so that for any admissible choice of g, p , $\mathcal{C}_1(S_{g,p})$ is k -hyperbolic.*

Note that $\mathcal{C}(S)$ is quasi-isometric to $\mathcal{C}_1(S)$, however the quasi-isometry constants depend on the underlying surface. Therefore Theorem 1.1 does not imply the uniform hyperbolicity of $\mathcal{C}(S)$. We also note that it has recently come to the author's attention that Bowditch has independently obtained the same result [1], as have Clay, Rafi and Schleimer using different methods [4], and Hensel, Przytycki and Webb [9], who have subsequently shown that it suffices to choose $\delta = 17$.

Let $d_{\mathcal{C}(S_{g,p})}(\cdot, \cdot)$ denote distance in $\mathcal{C}_1(S_{g,p})$; when there is no ambiguity, the reference to $S_{g,p}$ will be omitted in this notation. Let $\xi(S_{g,p}) = 3g + p - 4$; $\xi(S)$ is called the *complexity* of S . In the case that S is a disconnected surface, define $\xi(S)$ to be the sum of the complexities of its connected components. If $\xi(S) \leq 0$, S is called *sporadic* and each component of S possesses one of finitely many well understood topological types. In truth, the definition of $\mathcal{C}(S)$ needs to be modified when S is sporadic and connected (see [13] for details), because it is exactly these surfaces for which no two simple closed curves are disjoint.

In what follows, $i(\alpha, \beta)$ is the *geometric intersection number* of α and β , defined to be the minimum value of $|x \cap y|$, where x and y are representatives of the homotopy classes of α and β , respectively. The main tool in the proof of Theorem 1.1 is the following theorem.

Theorem 1.2 *For each $\lambda \in (0, 1)$, there is some $N = N(\lambda) \in \mathbb{N}$ such that if $\alpha, \beta \in \mathcal{C}_0(S_{g,p})$, whenever $\xi(S_{g,p}) > N$ and $d_{\mathcal{C}}(\alpha, \beta) \geq k$,*

$$i(\alpha, \beta) \geq \left(\frac{\xi(S)^\lambda}{f(\xi(S))} \right)^{k-2},$$

where $f(\xi) = O(\log_2(\xi))$.

In order to emphasize how Theorem 1.2 is used in the proof of Theorem 1.1, we define the numbers $q_{k,g,p}$ as follows:

$$q_{k,g,p} := \min\{i(\alpha, \beta) \mid \alpha, \beta \in \mathcal{C}_0(S_{g,p}) \text{ with } d_{\mathcal{C}}(\alpha, \beta) = k\}.$$

Irrespective of ξ , curve complex distance is bounded above by a logarithmic function of intersection number, and therefore if we fix g and p , $q_{k,g,p}$ grows exponentially as a function of k . But one may also study $q_{k,g,p}$ as a function of g or p .

Question 1 How does $q_{k,g,p}$ grow as a function of ξ ?

The content of Theorem 1.2 is that $q_{k,g,p}$ eventually grows faster than $\xi^{\lambda(k-2)}$ as a function of ξ for any $\lambda \in (0, 1)$.

We conclude the paper with two further applications of Theorem 1.2, the first being the resolution of a question posed by Masur and Minsky in [13] regarding vertex cycles of train tracks on surfaces (see Section 5 for relevant definitions). Specifically they ask the following.

Question 2 As a consequence of the fact that there are only finitely many train tracks on a surface S up to combinatorial equivalence, there is a bound $B = B(S)$ depending only on the topology of S such that

$$d_{\mathcal{C}(S)}(\alpha, \beta) < B$$

if $\alpha, \beta \in \mathcal{C}_0(S)$ are two vertex cycles of the same train track τ . Can B be made independent of S ? That is, is there some $B > 0$ such that, for any choice of S , the curve complex distance between two vertex cycles of the same train track on S is no more than B ?

In Section 5, we show the following.

Theorem 1.3 Let $\tau \subset S_{g,p}$ be a train track, and let α, β be vertex cycles of τ . Then if $\xi(S_{g,p})$ is sufficiently large,

$$d_{\mathcal{C}(S)}(\alpha, \beta) \leq 3.$$

In what follows, let $\text{Teich}(S)$ denote the Teichmüller space of S , the space of marked Riemann surfaces homeomorphic to S , modulo conformal equivalence isotopic to the identity. We will assume that $\text{Teich}(S)$ is equipped with the metric topology coming from the *Teichmüller metric*, denoted by $d_{\text{Teich}}(\cdot, \cdot)$. In this metric, the distance between two marked Riemann surfaces x and y is determined by the minimal dilatation constant associated to a quasiconformal map isotopic to the identity between x and y (see Farb and Margalit [6] for more details).

In the final section, we consider the map $\Phi: \text{Teich}(S) \rightarrow \mathcal{C}(S)$ introduced by Masur and Minsky in [13]; here $\Phi(x)$ is the set of isotopy classes of simple closed curves minimizing the extremal length in the conformal class of x . Note that Φ is technically a map into $\mathcal{P}(\mathcal{C}_0(S))$ (the power set of \mathcal{C}_0), but $\text{diam}_{\mathcal{C}}(\Phi(x))$ is uniformly bounded, and furthermore, there exists some $c = c(S)$ so that if x, y are Riemann surfaces within one of each other in $\text{Teich}(S)$, then

$$\text{diam}_{\mathcal{C}}(\Phi(x) \cup \Phi(y)) \leq c.$$

A map F from $\text{Teich}(S)$ to $\mathcal{C}(S)$ is then constructed by defining $F(x)$ to be any element of $\Phi(x)$; an immediate consequence of the existence of c is that F is coarsely

Lipschitz: for any $x, y \in \text{Teich}(S)$,

$$d_{\mathcal{C}}(F(x), F(y)) \leq c(s) \cdot d_{\text{Teich}}(x, y) + c(s).$$

In the final section, we show that c can be taken to be independent of S .

Theorem 1.4 *There exists $c > 0$ so that for any choice of g, p ,*

$$d_{\mathcal{C}(S_{g,p})}(F(x), F(y)) \leq c \cdot d_{\text{Teich}(S_{g,p})}(x, y) + c$$

for any $x, y \in \text{Teich}(S_{g,p})$. In other words, the map sending a Riemann surface to any curve in $\mathcal{C}(S)$ with minimal extremal length is coarsely Lipschitz, with Lipschitz constants independent of the choice of S .

How uniform hyperbolicity follows from Theorem 1.2 In both proofs of hyperbolicity of \mathcal{C} by Masur and Minsky and Bowditch, the method of proof is to construct a family of quasigeodesics satisfying certain properties.

In [13], Masur and Minsky show the existence of a coarsely transitive family of quasigeodesics $\{g_i\}_{i \in I}$ (images of Teichmüller geodesics under the above-mentioned map) equipped with projection maps $\phi_i: \mathcal{C}(S) \rightarrow g_i$, essentially having the property that the diameter of $\phi_i(\mathcal{C}(S) \setminus N_\epsilon(g_i))$ is bounded, with the bound depending only on S and $\epsilon > 0$. Here $N_\epsilon(\cdot)$ denotes the tubular neighborhood of radius ϵ . It is then demonstrated that this is a sufficient condition for δ -hyperbolicity. In [2], Bowditch constructs a similar family of uniform quasigeodesics and uses them to show that $\mathcal{C}(S)$ satisfies a subquadratic isoperimetric inequality, also implying δ -hyperbolicity.

Both Bowditch and Masur and Minsky rely on a key lemma, that every unit area singular Euclidean surface homeomorphic to S contains an annulus of definite width $W = W(S)$. Masur and Minsky prove this lemma using a limiting argument, while Bowditch's proof is more effective and yields some quantitative control on the growth of W as a function of S . Uniform hyperbolicity of \mathcal{C}_1 follows if one plugs the result of Theorem 1.2 into Bowditch's more effective setup, as is demonstrated in Section 4.

We note that, while the conclusion of Theorem 1.2 suffices to prove Theorem 1.1, we conjecture that this lower bound can be improved.

Conjecture 1.5 *Let k, α, β be as in the statement of Theorem 1.1. Then there exists a polynomial $f_k: \mathbb{N} \rightarrow \mathbb{N}$ of degree $k - 2$ such that*

$$d_{\mathcal{C}}(\alpha, \beta) \geq k \Rightarrow i(\alpha, \beta) \geq f_k(\xi(S)).$$

Organization of the paper In Section 2, we establish Theorem 1.2 for $k = 3$ which is used in Section 3 as the base case of an induction argument on the curve graph distance. In Section 3, we complete the proof of Theorem 1.2, and in Section 4 we show how Theorem 1.1 immediately follows from Theorem 1.2, together with the extensive quantitative control that Bowditch obtains on the growth of the hyperbolicity constant in his original proof. In Section 5, we use Theorem 1.2 to prove Theorem 1.3, and in Section 6, we show how to derive Theorem 1.4 as a corollary of Theorem 1.2.

Acknowledgements The author would primarily like to thank his adviser, Yair Minsky, for invaluable guidance and introducing him to the problem of obtaining quantitative control on the hyperbolicity of \mathcal{C} as a function of complexity. The author would also like to thank Ian Biringer, Asaf Hadari, and Thomas Koberda for reading a draft of this paper and offering many helpful comments and suggestions. Finally, the author thanks Yael Algom-Kfir, Spencer Dowdall, Johanna Mangahas and Babak Modami for many helpful and insightful conversations.

2 Lower bounds on intersection numbers for filling pairs

Let $\Gamma = \{\gamma_i\}_{i \in I}$ be a collection of curves on $S_{g,p}$ in pairwise *minimal position*, meaning that for each $k \neq j$,

$$|\gamma_j \cap \gamma_k| = i(\gamma_j, \gamma_k).$$

We say that such a collection *fills* the surface if the complement $S_{g,p} \setminus \Gamma$ is a union of topological disks and once-punctured disks. Equivalently, Γ fills if every homotopically nontrivial, nonperipheral (ie, not homotopic into a neighborhood of a puncture) simple isotopy class has a nonzero geometric intersection number with at least one member of Γ . Henceforth, we will use the word *essential* to refer to any curve which is nonperipheral and homotopically nontrivial. We begin the study of Question 1 with the following simpler question.

Question 3 On $S_{g,p}$, how many times does a pair of simple closed curves need to intersect in order to fill?

Note that two simple closed curves α and β fill if and only if $d_{\mathcal{C}}(\alpha, \beta) \geq 3$, for this precisely means that there is no essential simple closed curve which is simultaneously disjoint from both α and β . In light of the notation used in the introduction, Question 3 can therefore be restated as a question about $q_{3,g,p}$.

Lemma 2.1 $q_{3,g,p} \geq 2g + p - 2$

Proof Suppose α and β fill $S_{g,p}$. Then $\alpha \cup \beta$ can be considered as the 1-skeleton of a decomposition of $S_{g,p}$ into disks and once-punctured disks. Letting D denote the number of disks in this decomposition, we obtain

$$\chi(S_{g,p}) = 2 - 2g - p = -i(\alpha, \beta) + D.$$

The right hand side comes from the fact that there are twice as many edges as there are vertices in this decomposition, and the vertices are precisely the intersections between α and β . Then since $D \geq 0$, $q_{3,g,p} \geq 2g + p - 2$. \square

3 Proof of Theorem 1.2

In this section, we prove Theorem 1.2.

Theorem 1.2 For each $\lambda \in (0, 1)$, there is some $N = N(\lambda) \in \mathbb{N}$ such that if $\alpha, \beta \in \mathcal{C}_0(S_{g,p})$, whenever $\xi(S_{g,p}) > N$ and $d_C(\alpha, \beta) \geq k$,

$$i(\alpha, \beta) \geq \left(\frac{\xi(S)^\lambda}{f(\xi(S))} \right)^{k-2},$$

where $f(\xi) = O(\log_2(\xi))$.

We will show that if $\xi > N(\lambda)$, then

$$i(\alpha, \beta) < \left(\frac{\xi^\lambda}{f(\xi)} \right)^{k-2} \Rightarrow d_C(\alpha, \beta) < k.$$

We induct on the curve graph distance k ; the base case $k = 3$ was established in Section 2. Both $N(\lambda)$ and $f(\xi)$ will be established in the course of the proof.

Thus we begin by assuming that $\alpha, \beta \in \mathcal{C}_0(S_{g,p})$ are such that

$$i(\alpha, \beta) < \left(\frac{\xi^\lambda}{f(\xi)} \right)^{k-2}.$$

Assume that α and β are in minimal position. Cutting along α yields $S' := S \setminus \alpha$, which is topologically either a genus $g - 1$ surface with $p + 2$ punctures, or S' has two connected components. After cutting, β becomes a set B of disjoint arcs $\{b_1, \dots, b_n\}$ with endpoints at the two punctures corresponding to α . Consider a maximal subcollection

$$B' = \{b_{i_1}, \dots, b_{i_q}\} \subseteq B$$

of pairwise nonisotopic arcs; note that B' must fill S' and therefore there is some linear function $g(\xi)$ which bounds $q = |B'|$ from below. Furthermore, we have

$q = |B'| \leq 6g + 3(p + 2) - 4$ (see Korkmaz and Papadopoulos [11, Lemma 2.1]). Choose $h(\xi) = O(\xi)$, and $h(\xi) \gg 6g + 3(p + 2) - 4$, ($h(\xi) = 50\xi$, for instance). Note that the number of complementary regions of the arcs in B' is also no larger than $h(\xi)$. The main point is that $q = |B'|$ is bounded above and below by linear functions of ξ :

$$g(\xi) \leq |B'| \leq 50\xi.$$

Case 1: The original surface S is closed, so that $p = 0$ In this case, since B' fills S' , the collection of complementary components of B' in S' consists of a disjoint union of polygons, each having at least 3 sides. The collection of complementary regions defines a dual graph $G_{S'} = (V(G_{S'}), E(G_{S'}))$, whose vertices $V(G_{S'})$ correspond to complementary regions, and such that two vertices are connected by an edge if and only if they represent adjacent (across an arc in B') complementary regions in S' . For $v \in V(G_{S'})$, let $d(v)$ denote the degree of v , and let $\bar{d}(G)$ denote the average degree of G . Note that

$$\bar{d}(G_{S'}) \geq 3.$$

For each $j, 1 \leq j \leq k$, define the mass of b_{i_j} , denoted $m(j)$, to be the number of arcs in the original collection B that were isotopic to b_{i_j} . If j is such that

$$m(j) > \frac{i(\alpha, \beta)}{\xi^\lambda},$$

b_{i_j} is said to have *large mass*; otherwise b_{i_j} has *small mass*. Note that

$$\sum_{1 \leq j \leq q} m(j) = i(\alpha, \beta),$$

and therefore there can be no more than ξ^λ arcs of large mass. Assume ξ is sufficiently large so that

$$\xi \gg \xi^\lambda.$$

(This will be made more precise below.) Then for all such surfaces, cutting along all large mass arcs yields a possibly disconnected, nonsimply connected (indeed, nonsporadic) surface S'' . This is because cutting along any arc b_{i_j} in B' decreases the complexity by at most 3; to see this, it suffices to analyze the three possible cases:

- (1) b_{i_j} starts and ends at the same puncture and is nonseparating.
- (2) b_{i_j} starts and ends at the same puncture and is separating.
- (3) The terminal punctures of b_{i_j} are distinct.

The complexity of the surface obtained by cutting along b_{i_j} depends only on which of the three cases we are in; the details of this are left to the reader, because the nonsimply connectedness of S'' is implied by the conclusion of the following lemma.

Proposition 3.1 *There exists $N_1 \in \mathbb{N}$ such that if $\xi > N_1$, there exists a homotopically nontrivial simple closed curve $\gamma \subset S''$ intersecting no more than $f(\xi)$ arcs of B' .*

In the statement of Proposition 3.1, $f(\xi)$ is the same function f from the statement of Theorem 1.2, and it will be determined in the course of the proof. As will be proven below, Proposition 3.1 asserts that γ is essential when viewed as a simple closed curve on the original surface S , not just as a curve on S'' .

Proof of Proposition 3.1 The arcs in B'' must fill S'' and therefore, as was the case with B' , these arcs cut the surface into finitely many simply connected regions. Denote by $G_{S''}$ the corresponding dual graph. Then

$$\begin{aligned} |V(G_{S''})| &= |V(G_{S'})|, \\ |E(G_{S''})| &\geq |E(G_{S'})| - \xi^\lambda \geq g(\xi) - \xi^\lambda. \end{aligned}$$

Note that some of the arcs in B'' may be isotopic on S'' ; indeed, if T is a triangular complementary region in S' , and exactly one of the arcs of ∂T has large mass, then the remaining two arcs will form a bigon in S'' . It is also possible for all but one arc on the boundary of a complementary region in S' to have large mass; if this happens, the remaining arc will bound a disk in S'' .

Choose N_1 large enough so that for all $\xi > N_1$,

$$\bar{d}(G_{S''}) \geq \frac{2(g(\xi) - \xi^\lambda)}{|V(G_{S'})|} > 2.02.$$

We will need the following fact (see Fiorini, Joret, Theis and Wood [7, Lemma 3.2]; see also Diestel [5] for more details).

Lemma 3.2 *Let $\epsilon > 0$. There exists a decreasing function $g: (0, 1) \rightarrow \mathbb{R}_+$ so that if $G = (V, E)$ is any graph with $\bar{d}(G) > 2 + \epsilon$, then G has girth no larger than $g(\epsilon) \cdot \log_2(|V|)$.*

We recall that the *girth* of a graph G is the length of the shortest cycle on G .

Thus assuming $\xi(S) > N_1$, by Lemma 3.2, $G_{S''}$ has a simple cycle $\tilde{\gamma}$ no longer than

$$g\left(\frac{1}{50}\right) \cdot \log_2(2h(\xi)) := f(\xi).$$

(Recall that $h(\xi) = 50\xi$, and was chosen to be linear in ξ , and to bound the number of nonisotopic arcs in the collection B' from above.)

By construction, $\tilde{\gamma}$ corresponds to a simple closed loop γ intersecting no more than $f(\xi)$ small mass arcs; it remains to show that γ is homotopically nontrivial in S .

By construction, γ intersects at least one arc of B'' , and if γ intersects $b \in B''$, it only does so once since $\tilde{\gamma}$ is a simple cycle. We will show that all such intersections between γ and β are essential, which immediately implies that γ is essential.

Arguing by contradiction, assume that γ and β are not in minimal position. Then a closed arc j_γ of γ forms a bigon with a closed arc j_β of β . Since γ only intersects each arc in B'' once, it follows that j_β must contain pieces of two distinct arcs e_x, e_y of B'' as subarcs, each of which contains one of the two points of the set $\{x, y\} = j_\gamma \cap j_\beta$. Therefore, j_β must contain an element of $\beta \cap \alpha$.

Let $\tilde{\beta}$ represent the curve obtained from β by replacing j_β with j_γ . Then $\tilde{\beta}$ is isotopic to β but intersects α fewer times than β does, which contradicts the initial assumption that α and β began in minimal position.

This concludes the proof of Proposition 3.1. □

Hence γ is an essential, simple closed curve on S , which only intersects small mass arcs of β , and only at most $f(\xi)$ of them. Therefore,

$$i(\gamma, \beta) \leq f(\xi) \cdot \frac{i(\alpha, \beta)}{\xi^\lambda} < \frac{f(\xi)}{\xi^\lambda} \cdot \left(\frac{\xi^\lambda}{f(\xi)}\right)^{k-2} = \left(\frac{\xi^\lambda}{f(\xi)}\right)^{k-3},$$

where the strict inequality is due to our initial assumption. Then by the induction hypothesis,

$$d_C(\gamma, \beta) < k - 1,$$

and thus by the triangle inequality,

$$d_C(\alpha, \beta) \leq d_C(\alpha, \gamma) + d_C(\beta, \gamma) < 1 + (k - 1) = k.$$

This concludes the proof of Theorem 1.2 in the case that S is closed.

Case 2: $p > 0$ When $p > 0$, the complementary regions of B' in S' may now be once-punctured, and could have a single arc of B' in its boundary. In this case, we modify the definition of $G_{S'} = (V(G_{S'}), E(G_{S'}))$ as follows. There are two edges in the edge set $E(G_{S'})$ for every arc in B' . The vertex set $V(G_{S'})$ of $G_{S'}$ consists of two vertices for every nonpunctured complementary region, and one vertex for each punctured complementary region.

Note that this does not completely determine the graph $G_{S'}$. Given a nonpunctured complementary region, there are choices to be made as to which edges of $G_{S'}$ connect

to which of the two vertices corresponding to that region. However, the conclusion we are seeking will not depend on any of these choices, and therefore we assume they have been made arbitrarily.

The abstract graph $G_{S'}$ comes equipped with a preferred immersion into S' , (whose image we also refer to as $G_{S'}$), satisfying the following property:

Let e_1, e_2 be two edges corresponding to the same arc $e \in B'$ on the boundary of a punctured region R in S' . Then the region bounded $e_1 \cup e_2 \cup e$ contains the puncture of R (see Figure 1).




Figure 1: A possible portion of the graph $G_{S'}$ in the punctured case, immersed in S' . Punctures are represented by white circles, vertices of $G_{S'}$ by black circles.

The goal, as in the case that $p = 0$, will be to show the existence of a simple closed curve γ , intersecting no more than $f(\xi)$ small mass arcs of β , counting multiplicity.

Denote by $P = \{R_1, \dots, R_s\}$ the components of $S' \setminus B'$. As above, let $|R_i|$ denote the number of sides in the boundary of R_i . Let $U_1 \subseteq P$ be the set of punctured regions,

$U_2 \subseteq P$ the set of nonpunctured regions. Then by the Gauss–Bonnet Theorem,

$$\begin{aligned} 2\pi|\chi(S')| &= \pi\left(\sum_{R_i \in U_1} |R_i| + \sum_{R_j \in U_2} (|R_j| - 2)\right), \\ \Rightarrow 2(2g + p - 2) &= \sum_{R_i \in U_1} |R_i| + \sum_{R_j \in U_2} (|R_j| - 2), \\ \Rightarrow 2(2g + |U_1| - 2) &= \sum_{R_i \in U_1 \cup U_2} |R_i| - 2|U_2|. \end{aligned}$$

Note also that

$$|V(G_{S'})| = |U_1| + 2|U_2|, \quad |E(G_{S'})| = \sum_{R_i \in U_1 \cup U_2} |R_i|.$$

Since

$$\bar{d}(G_{S'}) = \frac{2|E(G_{S'})|}{|V(G_{S'})|},$$

we have

$$\begin{aligned} \bar{d}(G_{S'}) &= \frac{2\sum_i |R_i|}{|U_1| + 2|U_2|} = \frac{2(2(2g + |U_1| - 2) + 2|U_2|)}{|U_1| + 2|U_2|} \\ &= 2 \cdot \frac{4g - 4 + 2|U_1| + 2|U_2|}{|U_1| + 2|U_2|} \geq 2 \cdot \frac{\xi(S) + p + 2|U_2|}{p + 2|U_2|}. \end{aligned}$$

Note that the number $p + |U_2|$ of complementary regions R_i is at most twice the number of the arcs in B' , since every such region is bounded by at least one arc, and each arc is adjacent to two regions. Therefore,

$$p + 2|U_2| \leq 2(p + |U_2|) \leq 4(6g + 3(p + 2) - 4) < h(\xi),$$

and hence

$$\begin{aligned} 2 \cdot \frac{\xi(S) + p + 2|U_2|}{p + 2|U_2|} &= 2\left(1 + \frac{\xi}{p + 2|U_2|}\right) \\ &> 2\left(1 + \frac{\xi}{h(\xi)}\right) = 2.04. \end{aligned}$$

We obtain $G_{S''}$ from $G_{S'}$ as in the closed case; any edge in $E(G_{S'})$ corresponding to a large mass arc of β is deleted, and $|V(G_{S''})| = |V(G_{S'})|$. Thus there exists some $N_2 \in \mathbb{N}$ so that for $\xi(S) > N_2$,

$$\bar{d}(G_{S''}) > 2.02,$$

and therefore assuming $\xi > N_2$, $G_{S''}$ has a simple cycle $\tilde{\gamma}$ of length no more than

$$g\left(\frac{1}{50}\right) \log_2(|V(G_{S''})|) \leq f(\xi).$$

Unlike in the closed case, $\tilde{\gamma}$ does not automatically correspond to a simple closed curve on S'' , because $G_{S''}$ is only immersed and not embedded. Keeping this in mind, let γ be a curve in the homotopy class of $\tilde{\gamma}$; we will first show that γ is essential in S . We will again show this by proving that γ is in minimal position with β ; as above, assume that an arc j_γ of γ and an arc j_β of β bound a bigon in S . We claim that this bigon cannot be completely contained in S'' , and therefore j_β intersects α . Then as in the closed case, homotoping j_β across the bigon reduces the number of intersections between α and β , contradicting the assumption that α and β were chosen to be in minimal position.

Thus we must show that the bigon is not completely contained in S'' . Note that, unlike in the closed case, it is now possible for γ to cross a single arc of β in S'' more than once. However, out of all of the arcs of β entering the bigon, there must be some innermost one, characterized by the property that together with a piece of j_γ , it bounds a bigon containing no arc of β in its interior. This piece of j_γ must then correspond to a segment of $\tilde{\gamma}$ contained in one complementary region of S'' , and whose endpoints lie on the same arc on the boundary of this region. Thus this complementary region is punctured, and its puncture is contained in the interior of the bigon in question, a contradiction. Thus γ is essential.

Now, suppose γ intersects itself once. Let γ_1 be one of the two simple closed curves obtained from γ by starting and ending at the self-intersection point x . If γ_1 is nonperipheral in S , replace γ with γ_1 . Otherwise, let γ_2 be the other side of γ , so that $\gamma_2 = (\gamma \setminus \gamma_1) \cup \{x\}$.

If γ_2 is nonperipheral in S , replace γ with γ_2 and stop. We have reduced to the case where both γ_1 and γ_2 are peripheral, and therefore there exists punctures y_1, y_2 of S so that γ_j is homotopic into a neighborhood of y_j , $j = 1, 2$. Note that since γ is nonperipheral, $y_1 \neq y_2$. Let $N(\gamma)$ be a small regular neighborhood of γ . Then the component r of $\partial N(\gamma)$ encompassing both y_1 and y_2 is a simple curve, intersecting no more arcs of β than does $\tilde{\gamma}$. It remains to show that r is homotopically nontrivial and nonperipheral.

To see that this is the case, note that there must be an arc of β separating y_1 from y_2 that γ crosses, and therefore r crosses as well (see Figure 2). By the same argument applied to γ above, this intersection must be essential, and therefore r is essential. Thus replace γ with r .

Now suppose γ has $k > 1$ self-intersections, and let γ_1 be a simple closed curve obtained from γ by starting and ending at some intersection point x as above, and let γ_2 be the other side of γ . Then if γ_i is nonperipheral for either $i = 1, 2$, replace γ with γ_i . This reduces the number of self intersections that γ possesses, so we are done by induction. If γ_1 and γ_2 are both peripheral, then we can homotope γ into a neighborhood of the type pictured in Figure 2, and as above, one of the boundary components of this neighborhood will be simple and essential, by the same argument used above.




Figure 2: A schematic picture of γ in the case of one self-intersection when γ_1, γ_2 are both peripheral. White circles represent punctures, and r is the simple curve encompassing γ . The transverse arc belongs to β , and it must intersect r essentially.

Thus choose $N(\lambda) = \max\{N_1, N_2\}$. This completes the proof of Theorem 1.2. □

4 Independence of the hyperbolicity constant on ξ

We briefly summarize Bowditch’s proof of hyperbolicity of $\mathcal{C}(S)$ as seen in [2], and then demonstrate how Theorem 1.1 follows from his setup, with the use of Theorem 1.2. To avoid confusion, when possible we will use the same notation he introduces in his original article.

Define WX to be the set of *weighted curves*; an element of WX is a pair

$$(\lambda, \alpha), \quad \alpha \in \mathcal{C}_0(S_{g,p}), \lambda \in \mathbb{R}_+.$$

Then WMX , the set of *weighted multicurves* is the set of all finite formal sums of elements of WX with pairwise disjoint summands. Note that any element of $\mathcal{C}_0(S_{g,p})$ is naturally considered as an element of either WX or WMX by assigning unit weight. Given $(p, \lambda_1), (q, \lambda_2) \in WX$, we can extend the notion of geometric intersection number for elements of \mathcal{C}_0 to WX linearly

$$i(\lambda_1 p, \lambda_2 q) = \lambda_1 \lambda_2 i(p, q),$$

and we can again extend linearly to WMX . Then for $\alpha, \beta, \delta \in WMX$ with $i(\alpha, \beta) = 1$, define

$$l_{\alpha\beta}(\delta) := \max[i(\alpha, \delta), i(\beta, \delta)],$$

$$m_{\alpha\beta}(\delta) := \sup \left[\{l_{\alpha\beta}(\delta)\} \cup \left\{ \frac{i(\gamma, \delta)}{l(\gamma)} \mid \gamma \in \mathcal{C}_0 \right\} \right].$$

Then for any $r \geq 0$, we define

$$L(\alpha, \beta, r) = \{\delta \in \mathcal{C}_0 \mid l_{\alpha\beta}(\delta) \leq r\},$$

$$M(\alpha, \beta, r) = \{\delta \in \mathcal{C}_0 \mid m_{\alpha\beta}(\delta) \leq r\}.$$

Note that $M(\alpha, \beta, r) \subseteq L(\alpha, \beta, r)$. In Section 5, Bowditch shows that there exists an essential annulus of width at least $\eta = \eta(\xi)$ in any unit area singular Euclidean surface of complexity ξ , and uses this to show the following.

Lemma 4.1 *There exists $R = R(\xi)$ such that $M(\alpha, \beta, R) \neq \emptyset$. Furthermore, one can choose R so that $R = O(\xi^{5/2})$.*

As a consequence of this, we obtain the following.

Lemma 4.2 *There exists $D = D(\xi)$ so that for any $\alpha, \beta \in \mathcal{C}_0(S_{g,p})$,*

$$\text{diam}(L(\alpha, \beta, R)) < D.$$

Indeed, Lemma 4.1 follows immediately from the definitions and Lemma 4.2.

There is some $\delta \in M(\alpha, \beta, R)$ since it is nonempty. Then for any $\gamma \in L(\alpha, \beta, R)$, we have

$$d(\gamma, \delta) \leq i(\gamma, \delta) + 1 \leq l_{\alpha\beta}(\gamma)m_{\alpha\beta}(\delta) + 1 \leq R^2 + 1.$$

Here, the first inequality is always true, independent of ξ or the choice of γ and δ . The important inequalities in the chain above are the second and third ones, which in particular imply the existence of some $\delta \in L(\alpha, \beta, R)$ so that for any other $\gamma \in L(\alpha, \beta, R)$, $i(\gamma, \delta) \leq R^2 + 1$ and $R^2 + 1 = O(\xi^5)$.

Thus letting $f(\xi)$ be as in the statement of Theorem 1.2, there is some $\eta \in \mathbb{N}$ such that for $\xi > \eta$,

$$R^2(\xi) + 1 < \frac{\xi^6}{(f(\xi))^7} = \left(\frac{\xi^{6/7}}{f(\xi)}\right)^{(9-2)}.$$

Therefore, for $N(6/7)$ the natural number in the statement of Theorem 1.2, for all $\xi > \max(\eta, N(6/7))$, one has

$$d_{C(S)}(\gamma, \delta) < 9.$$

Thus, for all such values of ξ we can use $D = 18$ as the diameter bound for $L(\alpha, \beta, R)$.

Bowditch considers a metric space X having the property that to each pair of points $\alpha, \beta \in X$, there is a subset $\Lambda_{\alpha\beta}$, together with a *coarse ordering* $\leq_{\alpha\beta}$ on $\Lambda_{\alpha\beta}$. By a coarse ordering, we mean a transitive relation satisfying the property that for any $x, y \in \Lambda_{\alpha\beta}$, either $x \leq_{\alpha\beta} y$ or $y \leq_{\alpha\beta} x$.

Moreover, associated to X is a function $\phi: X \times X \times X \rightarrow X$; given three points $x, y, z \in X$, $\phi(x, y, z)$ plays the role of the center of a triangle with vertices x, y, z . We require ϕ to satisfy the relations

$$\phi(x, y, z) = \phi(y, z, x) = \phi(y, x, z), \quad \phi(x, x, y) = x,$$

and define $\Lambda_{\alpha\beta}[xy] \subseteq \Lambda_{\alpha\beta}$ by

$$\Lambda_{\alpha\beta}[xy] = \{z \in \Lambda_{\alpha\beta} \mid x \leq_{\alpha\beta} z \leq_{\alpha\beta} y\}.$$

He then shows the following.

Theorem 4.3 [2] *Suppose that $(X, \{(\Lambda_{\alpha\beta})\}_{(\alpha,\beta) \in X \times X}, \phi)$ satisfies $\phi(\alpha, \beta, \gamma) \in \Lambda_{\alpha\beta} \cap \Lambda_{\beta\gamma} \cap \Lambda_{\gamma\alpha}$, and suppose there exists $K \geq 0$ satisfying*

- (1) $d_{\mathcal{H}}(\Lambda_{\alpha\beta}[\alpha, \phi(\alpha, \beta, \gamma)], \Lambda_{\alpha\gamma}[\alpha, \phi(\alpha, \beta, \gamma)]) \leq K$,
- (2) *given $x, y \in X$ with $d(x, y) \leq 1$, $\text{diam}(\Lambda_{\alpha\beta}[\phi(\alpha, \beta, x), \phi(\alpha, \beta, y)]) \leq K$,*
- (3) *for $c \in \Lambda_{\alpha\beta}$, $\text{diam}(\Lambda_{\alpha\beta}[c, \phi(\alpha, \beta, \gamma)]) \leq K$.*

Then X is δ -hyperbolic, with hyperbolicity constant depending only on K .

In Theorem 4.3, $d_{\mathcal{H}}(\cdot, \cdot)$ denotes the Hausdorff distance.

Bowditch then goes on to define the sets $\Lambda_{\alpha\beta}$ to essentially be the curve obtained by choosing an element of $L(\lambda \cdot \alpha, \mu \cdot \beta, R)$ for each pair $\lambda, \mu \in \mathbb{R}_+$ satisfying $i(\lambda \cdot \alpha, \mu \cdot \beta) = 1$. Hyperbolicity is proved by showing that for this choice of $\Lambda_{\alpha\beta}$ (together with the choice of centers $\phi(\alpha, \beta, \gamma)$ whose definition is not summarized here; see [2]), the conditions of Lemma 4.1 are satisfied, with K depending only on D . This proves Theorem 1.1. □

5 Bounded diameter of vertex cycle sets of birecurrent train tracks

In this section, we prove Theorem 1.3, but before doing so we recall some of the basic terminology of train tracks on surfaces (refer to [13], Penner and Harer [17] and Papadopoulos [16] for more details). Recall that a *train track* $\tau \subset S$ is an embedded 1-complex; edges are called *branches* and vertices *switches*. Each edge is a smooth parameterized path with well-defined tangent vectors at the endpoints. Furthermore, at each switch v , there is a single line $L \subset T_v S$ such that the tangent vector of any branch incident at v lies on L . For each switch v , we choose a preferred direction of L ; a branch b incident at v is called *incoming* if its tangent vector at v is aligned with the chosen direction, and *outgoing* otherwise. We require that the valence of each switch be at least 3, unless τ has a simple closed curve component c ; in this case we have a single bivalent switch on c .

Any component Q of $S \setminus \tau$ is a surface with boundary consisting of smooth arcs running between cusps. We define the *generalized Euler characteristic* of Q to be

$$\chi(Q) - \frac{1}{2}V(Q),$$

where $V(Q)$ is the number of cusps on ∂Q . We require that the generalized Euler characteristic of each component of $S \setminus \tau$ be negative.

A *train path* is a smooth subpath of τ which traverses a switch only by entering via an incoming branch and exiting via an outgoing branch. Given a train track τ , let \mathcal{B} denote the set of branches. A nonnegative, real-valued function $\mu: \mathcal{B} \rightarrow \mathbb{R}$ is called a *transverse measure* on τ if for each switch v , it satisfies

$$\sum_{b \in i(v)} \mu(b) = \sum_{b' \in o(v)} \mu(b'),$$

where $i(v)$ is the set of incoming branches at v , $o(v)$ the set of outgoing branches. These are called the *switch conditions*.

We call τ *recurrent* if it admits a transverse measure with all positive weights, and *transversely recurrent* if, for each branch b , there exists a simple closed curve $c = c(b)$ intersecting b , which intersects τ transversely and is such that $S \setminus (\tau \cup c)$ has no bigons. τ is called *birecurrent* if it is both recurrent, and transversely recurrent, and *generic* if all switches are at most trivalent.

The set of transverse measures is invariant under scaling, thus it follows that the set of all transverse measures, viewed as a subset of $\mathbb{R}^{|\mathcal{B}|}$, is a cone over a compact polyhedron

in projective space. Let $P(\tau)$ denote the projective polyhedron of transverse measures; the class $[\mu] \in P(\tau)$ is called a *vertex cycle* of τ if it is an extreme point of $P(\tau)$.

A geodesic lamination λ is *carried* by τ if there is a C^1 map $\phi: \lambda \rightarrow \tau$ which is isotopic to the identity, and such that the restriction of the differential $d\phi$ to any tangent line of λ is nonsingular; this amounts to saying that $F(\lambda)$ is a train path on τ . Suppose c is a simple closed curve carried by τ . Then c induces a transverse measure called the *counting measure*: each branch b of τ is assigned the integer corresponding to the number of times c traverses b .

In general, if $[v]$ is a vertex cycle of τ , $[v]$ has a unique representative that is the counting measure on a simple closed curve c carried by τ (see [13]). Thus if $[v_1], [v_2]$ are two vertex cycles of a train track τ , we can define $d_C([v_1], [v_2])$ to be the curve graph distance between their simple closed curve representatives.

As a consequence of the fact that there are only finitely many train tracks on S up to homeomorphism, there is a bound $B = B(S)$ such that any two vertex cycles of the same train track are a distance of at most B from one another. However, Masur and Minsky [13] conjecture that B can be made independent of S , and conjecture further that $B = 3$ suffices.

Using Theorem 1.2, we will show the following.

Theorem 1.3 *Let $\tau \subset S_{g,p}$ be a train track, and let α, β be vertex cycles of τ . Then if $C(S_{g,p})$ is sufficiently large,*

$$d_{C(S)}(\alpha, \beta) \leq 3.$$

We note that this constant B also occurs in the proof of the quasiconvexity of the disk set by Masur and Minsky in [14].

Proof We can assume that τ is generic and birecurrent since for any train track τ , there exists a generic birecurrent track τ' such that $P(\tau) = P(\tau')$ (see [17]). Assume $d_C(\alpha, \beta) = 4$, and assume further that $\xi(S_{g,p}) > N(3/4)$. Then by Theorem 1.2,

$$i(\alpha, \beta) \geq \left(\frac{\xi^{3/4}}{f(\xi)}\right)^2 = \frac{\xi^{3/2}}{(f(\xi))^2}.$$

We will need the following fact about vertex cycles (see Hamenstädt [8] for a proof).

Lemma 5.1 *If α is a simple closed curve representative of a vertex cycle $[v]$ of τ , then if $\phi: \alpha \rightarrow \tau$ is the associated carrying map, $\phi(\alpha)$ traverses each branch of τ at most twice, and never twice in the same direction.*

Since τ is birecurrent, for any $\epsilon > 0$, there exists a hyperbolic metric σ on S so that τ has geodesic curvature less than ϵ with respect to σ [17]. Let α_σ (resp. β_σ) denote the unique geodesic representative of α (resp. β) in the metric σ . By choosing ϵ sufficiently small, we can assume α_σ and β_σ both lie within a small embedded tubular neighborhood $N_{\epsilon'}(\tau)$ foliated by transverse ties. Collapsing these ties to points yields a train track isotopic to τ within some small bounded distance of τ .

Let $\text{Br}(\tau)$ denote the number of branches of τ . We show the following.

Lemma 5.2 *If α and β are two vertex cycles of τ , then*

$$i(\alpha, \beta) \leq 4 \text{Br}(\tau).$$

Proof Let b be any branch of τ , and let $N_{\epsilon'}(b)$ denote the restriction of ϵ' tie-neighborhood of τ in the metric σ to b . By Lemma 5.1, α_σ and β_σ can each enter $N_{\epsilon'}(b)$ at most twice. Since α_σ and β_σ are both geodesics, they are in minimal position and therefore any arc of α_σ can intersect a given arc of β_σ at most once within $N_{\epsilon'}(b)$. □

For any train track $\omega \subset S$, one has the bound [17]

$$\text{Br}(\omega) \leq -9\chi(S) - 3p = 18g + 6p - 18.$$

Thus there is some $k \in \mathbb{N}$ so that for all $\xi > k$,

$$\frac{\xi^{3/2}}{(f(\xi))^2} = i(\alpha, \beta) \gg 4 \cdot \text{Br}(\tau),$$

contradicting Lemma 5.2. Thus for all surfaces S satisfying $\xi(S) > \max(k, N(3/4))$, α and β can not both be vertex cycles of the same train track on S , given that their curve graph distance is at least 4. □

6 Uniformity of the Lipschitz constants for the Teichmüller projection map

In this final section, we prove Theorem 1.4.

Theorem 1.4 *There exists $c > 0$ so that for any choice of g, p ,*

$$d_{\mathcal{C}(S_{g,p})}(F(x), F(y)) \leq c \cdot d_{\text{Teich}(S_{g,p})}(x, y) + c$$

for any $x, y \in \text{Teich}(S_{g,p})$. In other words, the map sending a Riemann surface to any curve in $\mathcal{C}(S)$ with minimal extremal length is coarsely Lipschitz, with Lipschitz constants independent of the choice of S .

Proof We follow Masur and Minsky’s proof of the complexity-dependent version of this statement, as seen in [13, Section 2]. Recall that

$$\Phi: \text{Teich}(S) \rightarrow \mathcal{P}(\mathcal{C}_0(S))$$

is the map that associates to each Riemann surface x the set of isotopy classes of curves with smallest extremal length. By subdividing the Teichmüller geodesic segment connecting x to y into unit length subsegments, It suffices to show that there exists some constant c such that for any choice of S , given $x, y \in \text{Teich}(S)$ with $d_{\text{Teich}}(x, y) \leq 1$,

$$\text{diam}_{\mathcal{C}(S)}(\Phi(x) \cup \Phi(y)) \leq c.$$

For each complete, finite volume hyperbolic metric on S , there is an essential simple closed curve with hyperbolic length less than some constant $b = b(S)$, the *Bers constant*; it is known that $b(S) = O(\log(\xi(S)))$ (see Buser [3]).

Furthermore, the extremal length is bounded above by an exponential function of hyperbolic length (see Maskit [12]): concretely, fixing a Riemann surface x and an essential simple closed curve γ , let $\text{ext}_x(\gamma)$ denote extremal length, and $\text{hyp}_x(\gamma)$ the length of the geodesic representative of γ in the unique complete finite volume hyperbolic metric in the conformal class of x . Then

$$\text{ext}_x(\gamma) \leq \frac{\text{hyp}_x(\gamma)}{2} e^{\text{hyp}_x(\gamma)/2}.$$

Therefore, there is some constant $E = E(S)$ such that any Riemann surface in $\text{Teich}(S)$ has a curve with extremal length no more than $E(S)$, and $E(S) = O(\xi \log(\xi))$.

Recall also Kerckhoff’s characterization of d_{Teich} in terms of extremal lengths (see Kerckhoff [10]):

$$d_{\text{Teich}}(x, y) = \frac{1}{2} \log \left(\sup_{\gamma \in \mathcal{C}_0(S)} \frac{\text{ext}_y(\gamma)}{\text{ext}_x(\gamma)} \right).$$

Now, let $\alpha \in \Phi(x)$; since α minimizes extremal length in x , $\text{ext}_x(\alpha) \leq E(S)$. Since $d_{\text{Teich}}(x, y) \leq 1$, by Kerckhoff’s formula for the Teichmüller distance, given any $\beta \in \Phi(y)$, it follows that $\text{ext}_x(\beta) \leq e^2 E(S)$. As seen in both [13] and Minsky [15],

$$\text{ext}_x(\alpha) \text{ext}_x(\beta) \geq i(\alpha, \beta)^2$$

and therefore

$$i(\alpha, \beta) \leq e E(S) = o(\xi(S)^2).$$

Thus Theorem 1.2 implies that there is some $k \in \mathbb{N}$ so that for $\xi(S) > k$,

$$d_{\mathcal{C}(S)}(\alpha, \beta) \leq 4.$$

Independent of the choice of complexity, curve complex distance is bounded above by a logarithmic function of intersection number (see [2]):

$$d_C(\alpha, \beta) \leq f(i(\alpha, \beta)), f: \mathbb{N} \rightarrow \mathbb{N}, \quad f = O(\log(n)).$$

Therefore it suffices to choose

$$c = \max \left(4, \max_{\xi(S) < k} f(\lceil eE(S) \rceil) \right).$$

Here, $\lceil x \rceil$ denotes the smallest integer larger than x . This completes the proof of Theorem 1.4. \square

References

- [1] **B H Bowditch**, *Uniform hyperbolicity of the curve graphs*, To appear in Pacific J. Math. Available at <http://homepages.warwick.ac.uk/~masgak/papers/uniformhyp.pdf>
- [2] **B H Bowditch**, *Intersection numbers and the hyperbolicity of the curve complex*, J. Reine Angew. Math. 598 (2006) 105–129 MR2270568
- [3] **P Buser**, *Geometry and spectra of compact Riemann surfaces*, Birkhäuser, Boston, MA (2010) MR2742784
- [4] **M T Clay, K Rafi, S Schleimer**, *Uniform hyperbolicity and geometric topology*, In preparation
- [5] **R Diestel**, *Graph theory*, 3rd edition, Graduate Texts in Mathematics 173, Springer, Berlin (2005) MR2159259
- [6] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Mathematical Series 49, Princeton Univ. Press (2012) MR2850125
- [7] **S Fiorini, G Joret, D O Theis, D R Wood**, *Small minors in dense graphs*, European J. Combin. 33 (2012) 1226–1245 MR2904987
- [8] **U Hamenstädt**, *Geometry of the complex of curves and of Teichmüller space*, from: “Handbook of Teichmüller theory, Vol. I”, (A Papadopoulos, editor), IRMA Lect. Math. Theor. Phys. 11, Eur. Math. Soc., Zürich (2007) 447–467 MR2349677
- [9] **S Hensel, P Przytycki, R Webb**, *Slim unicorns and uniform hyperbolicity for arc graphs and curve graphs* arXiv:1301.5577
- [10] **S P Kerckhoff**, *The asymptotic geometry of Teichmüller space*, Topology 19 (1980) 23–41 MR559474
- [11] **M Korkmaz, A Papadopoulos**, *On the arc and curve complex of a surface*, Math. Proc. Cambridge Philos. Soc. 148 (2010) 473–483 MR2609303

- [12] **B Maskit**, *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. 10 (1985) 381–386 MR802500
- [13] **H A Masur**, **Y N Minsky**, *Geometry of the complex of curves, I: Hyperbolicity*, Invent. Math. 138 (1999) 103–149 MR1714338
- [14] **H A Masur**, **Y N Minsky**, *Quasiconvexity in the curve complex*, from: “In the tradition of Ahlfors and Bers, III”, (W Abikoff, A Haas, editors), Contemp. Math. 355, Amer. Math. Soc. (2004) 309–320 MR2145071
- [15] **Y N Minsky**, *Teichmüller geodesics and ends of hyperbolic 3-manifolds*, Topology 32 (1993) 625–647 MR1231968
- [16] **A Papadopoulos**, *Introduction to Teichmüller theory, old and new*, from: “Handbook of Teichmüller theory, Vol. I”, IRMA Lect. Math. Theor. Phys. 11, Eur. Math. Soc., Zürich (2007) 1–30 MR2349667
- [17] **R C Penner**, **J L Harer**, *Combinatorics of train tracks*, Annals of Mathematics Studies 125, Princeton Univ. Press (1992) MR1144770

Department of Mathematics, Yale University
10 Hillhouse Avenue, New Haven, CT 06510, USA

tarik.aougab@yale.edu

Proposed: Danny Calegari

Seconded: Jean-Pierre Otal, Walter Neumann

Received: 16 January 2013

Revised: 27 May 2013

