

Surface group representations to $SL(2, \mathbb{C})$ and Higgs bundles with smooth spectral data

RICHARD A WENTWORTH
MICHAEL WOLF

We show that for every nonelementary representation of a surface group into $SL(2, \mathbb{C})$ there is a Riemann surface structure such that the Higgs bundle associated to the representation lies outside the discriminant locus of the Hitchin fibration.

30F60, 32G15, 53C07, 70S15; 30F40, 53C43

1 Introduction

Let Σ be a closed, oriented surface of genus $g \geq 2$. In this short note we answer a special case of the following question posed by Nigel Hitchin: which representations $\rho: \pi_1(\Sigma) \rightarrow SL(n, \mathbb{C})$ correspond to Higgs bundles which lie outside the discriminant locus of the Hitchin fibration for some Riemann surface structure on Σ ? For example, the Higgs field for a unitary representation (ie one whose image lies in a conjugate of $SU(n)$) is identically zero, and a reducible representation (ie one whose image preserves a proper subspace of \mathbb{C}^n for the standard action) necessarily has a Higgs field whose characteristic polynomial is reducible. As a consequence, these representations always lie in fibers over the discriminant locus for any choice of Riemann surface structure. The goal of this paper is to show that for $n = 2$ these examples present essentially the only restrictions. To state the result, recall that a representation $\rho: \pi(\Sigma) \rightarrow SL(2, \mathbb{C})$ is called *elementary* if it is either unitary, reducible, or maps to the subgroup generated by an embedding

$$\mathbb{C}^* \hookrightarrow SL(2, \mathbb{C}) : \lambda \mapsto \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$$

and the element $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. We shall prove the following:

Theorem 1 *A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow SL(2, \mathbb{C})$ defines a point in the fiber of the Hitchin fibration over the discriminant locus for every Riemann surface structure on Σ if and only if ρ is elementary.*

The natural approach to the above statement is to prove that if ρ is nonelementary, one can find a Riemann surface structure X on Σ so that the Higgs bundle on X corresponding to ρ defines a point in the fiber of the Hitchin fibration away from the discriminant locus for X . We shall prove this by combining the powerful result of Gallo, Kapovich and Marden [7] with the method of harmonic maps to trees (see [22; 24] by Wolf).

Let us first review a bit of the background and terminology for this problem. Let

$$(1) \quad \mathcal{X}(\Sigma) = \text{Hom}(\pi_1(\Sigma), \text{SL}(2, \mathbb{C})) // \text{SL}(2, \mathbb{C})$$

denote the $\text{SL}(2, \mathbb{C})$ -character variety of Σ parametrizing semisimple representations (see Culler and Shalen [4] and Lubotzky and Magid [16]). For a (marked) Riemann surface structure X on Σ , let $\mathcal{M}(X)$ denote the moduli space of rank 2 Higgs bundles on X with fixed trivial determinant (see Hitchin [9]). The nonabelian Hodge theorem asserts the existence of a homeomorphism $\mathcal{X}(\Sigma) \simeq \mathcal{M}(X)$ for each X . One direction of the homeomorphism is a consequence of the following result of Corlette [2] and Donaldson [6]: given a semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}(2, \mathbb{C})$ and a Fuchsian representation

$$\sigma: \pi_1(\Sigma) \xrightarrow{\sim} \Gamma \subset \text{PSL}(2, \mathbb{R}),$$

with $X = \Gamma \backslash \mathbb{H}^2$, there exists a smooth harmonic map $v: \mathbb{H}^2 \rightarrow \mathbb{H}^3$ that is equivariant for the action of $\pi_1(\Sigma)$ via σ on the upper half plane $\mathbb{H}^2 \subset \mathbb{C}$ and ρ on the hyperbolic 3-space \mathbb{H}^3 , on which $\text{SL}(2, \mathbb{C})$ acts by isometries. Moreover, v minimizes the energy among all such equivariant maps. We shall refer to v as an *equivariant harmonic map*. If $Q(X)$ denotes the space of holomorphic quadratic differentials on X , then there is a (singular) holomorphic fibration $h: \mathcal{M}(X) \rightarrow Q(X)$ which is a smooth fibration of abelian varieties over the locus of nonzero differentials with simple zeros. The image by h of a Higgs bundle corresponding to a semisimple representation is simply the Hopf differential of any equivariant harmonic map, as described above. The divisor $\Delta(X) \subset Q(X)$ consisting of those quadratic differentials having some zero with multiplicity is called the *discriminant locus*. Points in $\mathcal{M}(X)$ in the fiber over $q \in Q(X) \setminus \Delta(X)$ correspond to certain line bundles on a branched double cover of X called the *spectral curve*. The line bundle and the spectral curve together form the *spectral data*, which completely determine the Higgs bundle, and hence via the other direction of the nonabelian Hodge theorem, the corresponding representation ρ . The spectral data for points in $\mathcal{M}(X)$ lying over the discriminant locus are more difficult to describe; hence, the interest in the question posed by Hitchin. For more on this structure, see [10].

With this understood, [Theorem 1](#) is a direct consequence of the following equivalent statement:

Theorem 2 Let $\rho: \pi_1(\Sigma) \rightarrow SL(2, \mathbb{C})$ be a semisimple representation. Then there exists a Riemann surface structure $X = \Gamma \backslash \mathbb{H}^2$ on Σ such that the Hopf differential of the ρ -equivariant harmonic map $\mathbb{H}^2 \rightarrow \mathbb{H}^3$ has only simple zeros if and only if ρ is nonelementary.

Remark 3 (i) A unitary representation fixes a point in \mathbb{H}^3 , and so the constant map is equivariant and clearly energy-minimizing. Hence, the Hopf differential vanishes. A semisimple elementary representation that is not unitary fixes a geodesic in \mathbb{H}^3 , which then necessarily coincides with the image of any equivariant harmonic map. The Hopf differential is therefore the square of an abelian differential. In particular, since we assume $g \geq 2$, the differential has zeros with multiplicity. Therefore, the “only if” parts of Theorems 1 and 2 are clear.

- (ii) We shall actually prove a slightly stronger statement; namely, for nonelementary representations we can find a Riemann surface structure such that the vertical foliation of the Hopf differential has no saddle connections.
- (iii) Note that there are obviously sections of the bundle of holomorphic quadratic differentials over Teichmüller space $\text{Teich}(\Sigma)$ which at every point have zeros with multiplicity; one class of examples are the squares of abelian differentials just mentioned. Hence, Theorem 2 does not seem to follow from a simple dimension count.
- (iv) As pointed out by Hitchin, there will be other obstructions in any generalization of Theorem 1 for $n \geq 3$. In particular, some of these will come from other real forms of $SL(n, \mathbb{C})$. Representations to $SU(p, q)$, $p \neq q$, for example, will always lie in the discriminant locus (see Schaposnik [19]). Finding a suitable replacement in higher rank for the result of Gallo, Kapovich and Marden remains a challenge.
- (v) Theorem 2 states that if a nonelementary representation ρ is in the discriminant locus relative to one Riemann surface X , there is another Riemann surface Y for which ρ is not in the discriminant locus relative to Y . Neither the statement nor the proof suggests any conclusion about the frontier of the closure of these discriminant loci in any of the natural compactifications of the moduli space $\mathcal{X}(\Sigma)$ (see Daskalopoulos, Dostoglou, and Wentworth [5]).

We briefly outline the strategy for proving Theorem 2. The Hopf differential Φ of a locally defined harmonic map $w: \Omega \rightarrow (N, d)$ from a domain $\Omega \subset \mathbb{C}$ to metric space (N, d) is defined to be $\Phi = 4(u^* d_T)^2$.⁰ In the case of an equivariant harmonic map $v: \mathbb{H}^2 \rightarrow \mathbb{H}^3$, this differential Φ descends to a holomorphic quadratic differential

on the Riemann surface $X = \Gamma \backslash \mathbb{H}^2$, which for convenience we continue to denote by Φ . Projecting in \mathbb{H}^2 along the leaf space of Φ yields a $\pi_1(\Sigma)$ -equivariant harmonic map from \mathbb{H}^2 to a metric tree T_Φ , called the *dual tree* to Φ . There are two relevant observations: First, if the vertices of T_Φ all have valence three, then we recognize that Φ must have had only simple zeroes. Second, we notice that the $\pi_1(\Sigma)$ -equivariant product map $\mathbb{H}^2 \rightarrow \mathbb{H}^3 \times T_\Phi$ is a *conformal harmonic map*, ie it is both harmonic and has vanishing Hopf differential.

The idea then is to reverse this construction: we seek a tree T all of whose vertices are trivalent and a $\pi_1(\Sigma)$ -equivariant conformal harmonic map $\mathbb{H}^2 \rightarrow \mathbb{H}^3 \times T$. In that case, the resulting Hopf differential for the harmonic map to \mathbb{H}^3 will be the negative of the Hopf differential for projection to the tree, which necessarily has only simple zeroes.

If the function on Teichmüller space which records, for each domain Riemann surface, the equivariant energy of the harmonic map to $\mathbb{H}^3 \times T$ is proper, then there exists a conformal harmonic map. Now given a representation

$$\rho: \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{C}),$$

then unless ρ is quasi-Fuchsian one expects there to be certain divergent sequences in Teichmüller space along which the energy of the equivariant harmonic map $\mathbb{H}^2 \rightarrow \mathbb{H}^3$ is uniformly bounded, while for other divergent sequences the energy tends to $+\infty$. A similar statement holds for harmonic maps to trees. Therefore, the challenge is to associate a tree T to a given ρ such that the sum of the energies to T and \mathbb{H}^3 diverges along every choice of proper path in Teichmüller space. We are rescued in this quest by the main result of [7], which by realizing $\mathrm{SL}(2, \mathbb{C})$ -representations of surface groups as holonomies of complex projective structures, provides a measured lamination on the surface with image of at least moderate (quotient) length in \mathbb{H}^3 : that measured lamination can be adjusted so that its dual tree suffices for our needs.

Acknowledgements The authors warmly thank Nigel Hitchin for introducing us to the problem, and Shinpei Baba for discussions about projective structures which led to the formulation of Lemma 6. We also appreciate the helpful comments of the referees.

The authors were supported by National Science Foundation grants DMS-1406513 and DMS-1007383, respectively. Both authors are grateful to the Mathematical Sciences Research Institute at Berkeley, where some of this research was conducted, to the Institute for Mathematical Sciences at the National University of Singapore where the work was begun and to NSF grants DMS-1107452, 1107263, 1107367 “RNMS: Geometric structures And Representation varieties” (the GEAR Network) which supported collaborative travel for the authors.

2 Trees, measured foliations, and harmonic maps

In this section, we prove a lemma that motivates the strategy of the proof of [Theorem 2](#). The basic constructions in the statement of the lemma below were first exploited in [\[24\]](#). Namely, we will find the desired Riemann surface structure as a critical point for an energy function on Teichmüller space. To define this energy function, first choose a measured foliation, say (\mathcal{F}, λ) on the differentiable surface Σ , lift that measured foliation to a $\pi_1(\Sigma)$ -equivariant measured foliation on the universal cover $\tilde{\Sigma}$, and then project the transverse measure λ along the leaves to obtain an \mathbb{R} -tree $T = T_\lambda$ with an isometric action (relative to the metric defined by the projected measure) of $\pi_1(\Sigma)$. For concreteness, we will express the isometric action of the fundamental group on T by a representation $\rho_T: \pi_1(\Sigma) \rightarrow \text{Iso}(T)$. For any $\gamma \in \pi_1(\Sigma)$ whose free homotopy class is represented by a simple closed curve, the intersection $i(\gamma, \lambda)$ with the foliation is equal to the translation length γ as it acts on T :

$$(2) \quad i(\gamma, \lambda) = |\rho_T(\gamma)|_T := \min_{x \in T} d_T(x, \gamma x).$$

Recall that the action of an isometry on an \mathbb{R} -tree is always semisimple (see Culler and Morgan [\[3\]](#)); hence the “min” instead of an “inf” in [\(2\)](#).

We focus initially on two features of this construction. First, given a Fuchsian representation $\sigma: \pi_1(\Sigma) \xrightarrow{\sim} \Gamma \subset \text{PSL}(2, \mathbb{R})$, a Riemann surface $X = \Gamma \backslash \mathbb{H}^2$, and an \mathbb{R} -tree T with an isometric action $\rho_T: \pi_1(\Sigma) \rightarrow \text{Iso}(T)$, a map $u: \mathbb{H}^2 \rightarrow T$ is called $\pi_1(\Sigma)$ -equivariant if $u(\sigma(\gamma)z) = \rho_T(\gamma)u(z)$ for all $\gamma \in \pi_1(\Sigma)$ and all $z \in \mathbb{H}^2$ (when the Riemann surface structure is assumed, we sometimes say that u is ρ_T -equivariant to emphasize the action on the target). We define the ρ_T -energy $E_{\rho_T}(X)$ of X to be the infimum of the energies of locally finite energy $\pi_1(\Sigma)$ -equivariant maps $\mathbb{H}^2 \rightarrow T$ (see [\[22\]](#) for the case of maps to \mathbb{R} -trees and [\[14; 12\]](#) for the general setting of nonpositively curved metric space targets). Here we note that the energy density for such maps is a locally integrable form on \mathbb{H}^2 that is invariant with respect to the action of $\pi_1(\Sigma)$ via σ . It therefore descends to X , and its integral gives a well defined (finite) energy. Moreover, any energy minimizer (or harmonic map) $u: \mathbb{H}^2 \rightarrow T$ has the following properties (see [\[11; 22; 23; 5\]](#)):

- There is a nonzero holomorphic quadratic (Hopf) differential $\Phi \in Q(X)$ whose vertical measured foliation (on \mathbb{H}^2) defines a metric tree T_Φ with an isometric action of $\pi_1(\Sigma)$.
- There is a $\pi_1(\Sigma)$ -equivariant map $\psi: T_\Phi \rightarrow T$ which is a *folding*; in the case when $T = T_\lambda$ is dual to a measured foliation (the only case we will consider here), then ψ is an isometry.

- Finally, $u = \psi \circ \pi$, where $\pi: \mathbb{H}^2 \rightarrow T_\Phi$ is the projection onto the vertical leaf space of Φ .

Moreover, the energy of u is given by

$$(3) \quad E_{\rho_T}(X) := E(u) = 2 \int_X |\Phi|.$$

The energy only depends on the marked isomorphism class of X . Hence, $E_{\rho_T}(X)$ is a well-defined function $E_{\rho_T}: \text{Teich}(\Sigma) \rightarrow \mathbb{R}_{\geq 0}$.

Second, some features of the (Hopf) quadratic differential Φ are reflected in the tree: in particular, if each vertex of the tree has valence three, then Φ can have only simple zeros, as any higher order zeros—or indeed any collection of zeros connected by subarcs of a leaf—would create higher order branching of the leaf space, which is the tree $T = T_\Phi$ in this setting. As it is a generic condition that the zeros of a holomorphic quadratic differential should be simple with no connecting leaves between them, we see that the generic tree dual to a measured foliation should have all vertices of valence three.

The hyperbolic 3–ball $\mathbb{H}^3 = \text{SL}(2, \mathbb{C})/\text{SU}(2)$ has a left action of $\text{SL}(2, \mathbb{C})$ by isometries. Fix a semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}(2, \mathbb{C})$. Then \mathbb{H}^3 inherits a left action of $\pi_1(\Sigma)$ by isometries. Given a Riemann surface structure $X = \Gamma \backslash \mathbb{H}^2$ on Σ , the theorem of Corlette and Donaldson mentioned in the introduction asserts the existence of a harmonic map $v: \mathbb{H}^2 \rightarrow \mathbb{H}^3$ that is equivariant with respect to ρ ; this map is unique if and only if ρ is irreducible. Thus, in analogy with what we did with the target tree T in defining the ρ_T –energy, we may define the ρ –energy $E_\rho(X)$ of a Riemann surface to be the energy $E(v)$ of v . As before, the function E_ρ is well-defined on the Teichmüller space $\text{Teich}(\Sigma)$.

Finally, consider the nonpositively curved metric space $N = T \times \mathbb{H}^3$ with product metric d_N and with the diagonal isometric action $\pi_1(\Sigma) \rightarrow \text{Iso}(N)$ given by $\rho_N(\gamma) = (\rho_T(\gamma), \rho(\gamma))$. The energy of equivariant maps $\mathbb{H}^2 \rightarrow N$ is simply the sum of the energies of the maps to T and \mathbb{H}^3 . This defines our setting well enough to state the following lemma:

Lemma 4 *Let $T = T_\lambda$ be a tree which is both dual to a measured foliation on the surface Σ and has all vertices of valence three, and let $\rho: \pi_1(\Sigma) \rightarrow \text{SL}(2, \mathbb{C})$ be irreducible. Suppose that the function $E_{\rho_N} = E_{\rho_T} + E_\rho$ is proper on $\text{Teich}(\Sigma)$. Then there exists a Riemann surface structure on Σ such that the Hopf differential of the ρ –equivariant harmonic map $\mathbb{H}^2 \rightarrow \mathbb{H}^3$ has only simple zeros.*

Remark 5 By our comments above on the generic nature of such trees, we see that the first sentence is not a vacuous condition.

Proof By a classical result (see [20; 18], and for the case of general nonpositively curved metric target spaces, [21, Corollary 1.3]), the energy function

$$E_{\rho_T} + E_{\rho}: \text{Teich}(\Sigma) \rightarrow \mathbb{R}$$

is differentiable on $\text{Teich}(\Sigma)$, and so, being proper, achieves its minimum at a point $X = \Gamma \backslash \mathbb{H}^2$; moreover, the gradient of that energy function vanishes at X . On the other hand, the classical expression for the gradient as a multiple of the Hopf differential of the ρ_N -equivariant harmonic map from \mathbb{H}^2 to $T \times \mathbb{H}^3$ holds in this case (see [21, Theorem 1.2]), and so the Hopf differential of that harmonic map vanishes. Because the target metric is a product, we may express the harmonic map $f: \mathbb{H}^2 \rightarrow T \times \mathbb{H}^3$ as a product $f = (u, v)$, where u is the unique ρ_T -equivariant harmonic map $\mathbb{H}^2 \rightarrow T$, and v is the unique ρ -equivariant harmonic map $\mathbb{H}^2 \rightarrow \mathbb{H}^3$. The Hopf differential of f is the sum of the Hopf differentials Φ_u and Φ_v of u and v , respectively; and since it vanishes, we have $\Phi_v = -\Phi_u$. However, as explained in the opening of this section, the vertical measured foliation of Φ_u has leaf space which projects to a tree T_{Φ_u} that is equivariantly isometric to T . In particular, since T has all vertices of valence three, the differential Φ_u has simple zeros. The same is therefore true of $\Phi_v = -\Phi_u$. \square

3 Complex projective structures and bending laminations

Let us introduce some more notation. For a hyperbolic surface S and simple closed curve $\gamma \subset S$, let $\ell_S(\gamma)$ denote the length of the geodesic in the free homotopy class of γ as measured on S . For $g \in \text{Iso}(\mathbb{H}^3)$, define the translation length $|g|_{\mathbb{H}^3}$ as in (2):

$$|g|_{\mathbb{H}^3} := \inf_{x \in \mathbb{H}^3} d_{\mathbb{H}^3}(g \cdot x, x).$$

The goal now is to find a tree for which the hypotheses of Lemma 4 are satisfied. To that end, let $\rho: \pi_1(\Sigma) \rightarrow SL(2, \mathbb{C})$ be nonelementary. The foundational result in [7] implies that ρ is the holonomy of a complex projective structure, say (X, \wp) , and hence is the holonomy of a developing map $\text{dev}_{\rho}: \tilde{\Sigma} \rightarrow \mathbb{C}P^1$. (The reader may find it useful to keep in mind that this complex projective structure is not necessarily unique, and in general, the developing map, while a local homeomorphism, is neither necessarily injective nor a covering.) We exploit the rich synthetic hyperbolic geometry of complex projective structures in the following lemma; in that setting, because of hyperbolic geometric constructions, it is more convenient to replace measured foliations with measured laminations in the discussion. As the natural homeomorphism between the space of measured foliations and measured laminations respects the passage to dual trees, there is no loss of content in this change of perspective. A *maximal lamination* is a measured lamination all of whose complementary regions are ideal triangles; for more background on properties of geodesic laminations used below, see [1].

Lemma 6 *Let (X, \wp) be a complex projective structure on Σ with holonomy ρ . Then there is a hyperbolic structure S on Σ , a maximal measured geodesic lamination λ on S , and constants $\varepsilon_1, A > 0$, depending only on (S, λ) , such that the following hold:*

- (i) *If γ is a simple closed curve on Σ with intersection number $i(\gamma, \lambda) < \varepsilon_1$, then $|\rho(\gamma)|_{\mathbb{H}^3} \geq A\ell_S(\gamma)$.*
- (ii) *More generally, for any constant $I > 0$, there is $L > 0$ so that if γ is a simple closed curve on Σ with $i(\gamma, \lambda) < I$ and $\ell_S(\gamma) > L$, then $|\rho(\gamma)|_{\mathbb{H}^3} \geq A\ell_S(\gamma)$.*

Proof We begin by recalling the key property of complex projective structures we will need. Good references for this material, due almost entirely to Thurston, are [13, Section 2] and [15, Theorem 8.6]. Given a complex projective structure (X, \wp) on Σ with holonomy ρ , there is a hyperbolic surface structure S on Σ , a measured geodesic lamination λ_0 and a (pleated surface) map $F: \tilde{S} \rightarrow \mathbb{H}^3$ from the universal cover \tilde{S} to \mathbb{H}^3 , which has image a surface $F(\tilde{S}) \subset \mathbb{H}^3$ and for which $F|_{\tilde{\lambda}_0}$ is an isometry. Here, $\tilde{\lambda}_0$ is the lift to \tilde{S} of the lamination $\lambda_0 \subset S$.

Choose a point $p \in \lambda_0$ and a small neighborhood $U \subset S$ containing p . Some of the leaves, say α_i , of λ_0 that meet U later recur to U , and the images of those arcs α_i determine F -images, say $F(\hat{U}_i) = V_i \subset \mathbb{H}^3$, of lifts \hat{U}_i of U that are separated by (fixed portions of) isometric images of the arcs α_i . In particular, the images V_i of those lifts are at some minimum distance A from each other, depending only on the geometry of S and $\lambda_0 \subset S$.

Note that if γ is a closed curve which lies C^1 -close to a lamination, we can choose such a neighborhood U so that γ meets U some number k times before closing up. Thus, if the image $F(\tilde{q})$ of a lift \tilde{q} of a point $q \in \gamma \cap U$ were to lie in a neighborhood $V_0 \subset \mathbb{H}^3$, then the image $\rho(\gamma)(F(\tilde{q}))$ by the isometry $\rho(\gamma)$ of $F(\tilde{q})$ would have to lie in some lift $V_k \subset \mathbb{H}^3$, with a single lift $\hat{\gamma}$ connecting the neighborhoods V_0 and V_k and meeting other lifts V_1, \dots, V_{k-1} along its path. We conclude that such an isometry $\rho(\gamma)$ has translation length $|\rho(\gamma)|_{\mathbb{H}^3}$ comparable to that of its length $\ell_S(\gamma)$ on S : the construction shows that this comparability constant $|\rho(\gamma)|_{\mathbb{H}^3}/\ell_S(\gamma)$ may be taken to depend only on λ_0 and S , but to be independent of γ , so long as γ is sufficiently close in C^1 to λ_0 .

Therefore, if λ_0 is also a maximal lamination, set $\lambda = \lambda_0$ and our construction of λ is complete. It is of course possible that the lamination λ_0 is not maximal. For example, the lamination λ_0 might consist only of a single simple closed curve, so that the complement in Σ of λ_0 could be a surface of large Euler characteristic. In that case, we may perturb λ_0 into a maximal lamination λ : measured laminations which are maximal in this sense are dense, for example by using [11] and the density of

holomorphic quadratic differentials on a Riemann surface with corresponding properties or the theory of train tracks [17]. This new measured lamination λ will meet the old lamination λ_0 at a maximum angle $\delta > 0$, which we may choose to be as small as we wish. In particular, the perturbation of λ_0 to λ has only a mild effect on our constructions and estimates: by choosing δ small enough, and restricting ourselves to curves γ which are both very long and very close in C^1 to leaves in λ , we find that since λ is close to λ_0 in C^1 , we have already focused on curves which are sufficiently close to λ_0 in C^1 for the previous estimates to hold: for curve classes γ whose S -geodesic representatives are sufficiently close to the S -measured geodesic lamination λ , we have that $|\rho(\gamma)|_{\mathbb{H}^3} \geq A\ell_S(\gamma)$.

With these observations in mind, consider part (ii) of the lemma. Fix a number $I > 0$. It suffices to show that there is a bound $L > 0$ such that for any simple closed curve $\gamma \subset \Sigma$ with intersection number $i(\gamma, \lambda) < I$ and length $\ell_S(\gamma) > L$, the S -geodesic representative of γ lies C^1 -close to the S -geodesic measured lamination λ . For suppose that it is not the case, ie that there is some I and a sequence of curves γ_k for which $i(\gamma_k, \lambda) < I$, while $\ell_S(\gamma_k) \rightarrow \infty$ and the C^1 -distance between γ_k and λ is bounded away from zero. Consider the measured geodesic laminations μ_k whose measure is given, for a transverse arc C , by $\mu_k(C) = i(C, \gamma_k)/\ell_S(\gamma_k)$, ie normalized counting measure. Of course, as $k \rightarrow \infty$, the intersection numbers satisfy

$$i(\mu_k, \lambda) = i(\gamma_k/\ell_S(\gamma_k), \lambda) < \frac{I}{\ell_S(\gamma_k)} \rightarrow 0.$$

Allowing μ to be an accumulation point of μ_k , we see that $i(\mu, \lambda) = 0$. Moreover, μ is nontrivial (for example, a subsequence μ_k can all be carried on a single train track, but then one of the finitely many branches of that track admits an intersection number with a transverse arc that is bounded away from zero). But as λ is maximal and $i(\mu, \lambda) = 0$, we have that μ is a sublamination of λ , and hence the support of μ_k — that is, the curve γ_k — may be taken to approximate λ in the Hausdorff sense. This in turn implies, by the geometry of nearby hyperbolic geodesics, that γ_k lies arbitrarily closely to λ in C^1 , contradicting the assumption.

Similarly, for part (i), if no such constants ε_1, A exist, we may find γ_k for which $i(\gamma_k, \lambda) \rightarrow 0$ and $|\rho(\gamma_k)|_{\mathbb{H}^3}/\ell_S(\gamma_k) \rightarrow 0$, and we derive a contradiction as above. This completes the proof of the lemma. □

4 Proof of the main result

Let $\rho: \pi_1(\Sigma) \rightarrow SL(2, \mathbb{C})$ be nonelementary. The theorem of Gallo, Kapovich and Marden guarantees that ρ is the holonomy of a complex projective structure (X, \wp)

on Σ . Let $T = T_\lambda$ be the dual tree to the measured lamination, and S the hyperbolic structure on Σ , obtained in Lemma 6. Let $N = T \times \mathbb{H}^3$ and ρ_N be as in Section 2. We will need a preliminary result about N : by Lemma 6(i) and equation (2), we immediately have

Lemma 7 *There exists $\varepsilon_2 > 0$, which depends only on ρ , S , and λ , such that for all $1 \neq \gamma \in \pi_1(\Sigma)$, the translation length $|\rho_N(\gamma)|_N \geq \varepsilon_2$.*

We can now give the following proof:

Proof of Theorem 2 By Lemma 4, it suffices to show that the energy function $E_{\rho_N} = E_{\rho_T} + E_\rho$ is proper on $\text{Teich}(\Sigma)$. Let us remark that in case ρ is quasi-Fuchsian, it was shown in [8, Section 5] (see also [24, Prop. 3.6]) that E_ρ is proper, and therefore so is E_{ρ_N} for any choice of T . For general ρ , however, properties of the lamination λ and the associated tree $T = T_\lambda$ play a key role, and the argument is necessarily different from the one used in [8, Section 5]. With the intent of arriving at a contradiction, we therefore suppose to the contrary that E_{ρ_N} is not proper. Under the assumption we can find a sequence $\sigma_i: \pi_1(\Sigma) \xrightarrow{\sim} \Gamma_i \subset \text{PSL}(2, \mathbb{R})$ of Fuchsian representations such that the set of isomorphism classes of marked Riemann surfaces $\{X_i\}_{i \in \mathbb{N}}$, $X_i = \Gamma_i \backslash \mathbb{H}^2$, contains no limit points in $\text{Teich}(\Sigma)$. We suppose furthermore that we have a constant K and unique harmonic maps

$$u_i: \mathbb{H}^2 \rightarrow T \quad \text{and} \quad v_i: \mathbb{H}^2 \rightarrow \mathbb{H}^3$$

that are equivariant with respect to the action of $\pi_1(\Sigma)$, via σ_i on the left, and ρ_T and ρ on the right, with $E(u_i) + E(v_i) \leq K$.

Step 1 By a standard argument (see [20; 18]), the energy bound plus Lemma 7 imply that there is a uniform positive lower bound on the lengths of the shortest geodesics for the hyperbolic surfaces X_i . By the Mumford–Mahler compactness theorem, it follows that we can find quasiconformal homeomorphisms $g_i: \mathbb{H}^2 \rightarrow \mathbb{H}^2$ and a Fuchsian representation $\sigma_\infty: \pi_1(S) \xrightarrow{\sim} \Gamma_\infty$, such that $g_i \circ \Gamma_i \circ g_i^{-1} = \Gamma_i$, and (after passing to a subsequence) $\hat{\sigma}_i = g_i \circ \sigma_i \circ g_i^{-1} \rightarrow \sigma_\infty$, in the Chabauty topology. Introduce the following notation: for any $\gamma \in \pi_1(\Sigma)$, define

$$(4) \quad \hat{\gamma}_i := \sigma_i^{-1} \circ \hat{\sigma}_i(\gamma).$$

Step 2 Let us first focus on the maps u_i to the tree. By [14] and the convergence of the $\hat{\sigma}_i$, the maps u_i are uniformly Lipschitz with a constant proportional to $\sqrt{E(u_i)}$. In particular, since the energy is uniformly bounded, so is the Lipschitz constant. Therefore, we may assume the Hopf differentials Φ_i of u_i , regarded as Γ_i -automorphic holomorphic quadratic differentials on \mathbb{H}^2 , converge $\Phi_i \rightarrow \Phi_\infty$ uniformly to a holomorphic

differential Φ_∞ . It is possible that $\Phi_\infty \equiv 0$; we will deal with this contingency in **Step 6** below. In the intervening steps below, assume $\Phi_\infty \not\equiv 0$.

Step 3 As discussed previously, the leaf space T_{Φ_i} of the vertical measured foliation of Φ_i has the structure of an \mathbb{R} -tree with an isometric action of $\pi_1(\Sigma)$ (via $\hat{\sigma}_i$) that is $\pi_1(\Sigma)$ -equivariantly isometric to T . Denote this isometry by $\psi_i: T_{\Phi_i} \rightarrow T$. If we let $\pi_i: \mathbb{H}^2 \rightarrow T_{\Phi_i}$ be the projection onto the leaf space of the vertical foliation, then as in **Section 2** we have that u_i is given by $u_i = \psi_i \circ \pi_i$.

Step 4 Fix $\gamma \in \pi_1(\Sigma)$. We choose a representative curve α_∞ in \mathbb{H}^2 from 0 to $\sigma_\infty(\gamma) \cdot 0$ that is quasitransverse to the vertical measured foliation of Φ_∞ . Let $\alpha_i: [0, 1] \rightarrow \mathbb{H}^2$ be a path from 0 to $\hat{\sigma}_i(\gamma) \cdot 0$, that is quasitransverse to the vertical foliation of Φ_i . Then since the $\hat{\sigma}_i$ and Φ_i converge, α_i may furthermore be chosen ε -close to α_∞ for i sufficiently large.

Step 5 By **Step 4**, it follows that there exists I (depending on γ) such that for i sufficiently large,

$$d_{T_{\Phi_i}}(\pi_i\alpha_i(1), \pi_i\alpha_i(0)) < I.$$

On the other hand,

$$\begin{aligned} d_{T_{\Phi_i}}(\pi_i\alpha_i(1), \pi_i\alpha_i(0)) &= d_T(\psi_i \circ \pi_i\alpha_i(1), \psi_i \circ \pi_i\alpha_i(0)), \\ &= d_T(u_i(\hat{\sigma}_i(\gamma)\alpha_i(0)), u_i(\alpha_i(0))), \\ &= d_T(u_i(\sigma_i(\hat{\gamma}_i)\alpha_i(0)), u_i(\alpha_i(0))), \\ &= d_T(\rho_T(\hat{\gamma}_i)u_i(\alpha_i(0)), u_i(\alpha_i(0))), \end{aligned}$$

where $\hat{\gamma}_i$ is defined by (4). Hence, in particular,

$$(5) \quad i(\hat{\gamma}_i, \lambda) = |\rho_T(\hat{\gamma}_i)|_T < I,$$

for i sufficiently large.

Step 6 In the case where $\Phi_\infty \equiv 0$, it follows from (3) that $E(u_i) \rightarrow 0$. Hence, by the assertion in **Step 2**, the Lipschitz constants for u_i also tend to zero uniformly. Therefore, for any given $\gamma \in \pi_1(\Sigma)$, since $\hat{\sigma}_i(\gamma) \cdot 0 \rightarrow \sigma_\infty(\gamma) \cdot 0$ remains bounded,

$$\begin{aligned} |\rho_T(\hat{\gamma}_i)|_T &\leq d_T(\rho_T(\hat{\gamma}_i)u_i(0), u_i(0)) \\ &= d_T(u_i(\sigma_i(\hat{\gamma}_i) \cdot 0), u_i(0)) \\ &= d_T(u_i(\hat{\sigma}_i(\gamma) \cdot 0), u_i(0)) \\ &\rightarrow 0 \end{aligned}$$

by the decay of the Lipschitz constants for u_i and the convergence of $\hat{\sigma}_i(\gamma) \cdot 0$. Thus $|\rho_T(\hat{\gamma}_i)|_T < I$ for i sufficiently large so that (5) holds in this case as well.

Step 7 We apply a similar argument to the sequence of harmonic maps v_i . Since the energy $E(v_i)$ is uniformly bounded, and the groups $g_i \circ \Gamma_i \circ g_i^{-1}$ converge, the v_i are uniformly Lipschitz. In particular, for any $\gamma \in \pi_1(\Sigma)$ there is B (depending on γ), such that

$$d_{\mathbb{H}^3}(v_i(\hat{\sigma}_i(\gamma) \cdot 0), v_i(0)) \leq B.$$

In that case,

$$d_{\mathbb{H}^3}(v_i(\hat{\sigma}_i(\gamma) \cdot 0), v_i(0)) = d_{\mathbb{H}^3}(v_i(\sigma_i(\hat{\gamma}_i) \cdot 0), v_i(0)) = d_{\mathbb{H}^3}(\rho(\hat{\gamma}_i)v_i(0), v_i(0))$$

and hence

$$(6) \quad |\rho(\hat{\gamma}_i)|_{\mathbb{H}^3} \leq B.$$

Of course, in this last term, the quantity B still depends on γ but is bounded independently of the index i .

Step 8 We now relate the estimates of the previous three steps to arrive at the following crucial conclusion. Combining (5) and (6) with Lemma 6, we find that the lengths $\ell_S(\hat{\gamma}_i)$ must be uniformly bounded in i . This implies that there are only finitely many homotopy classes among the $\hat{\gamma}_i$. Hence, after passing to a subsequence we may assume there exists a fixed $\hat{\gamma}$ such that $\hat{\gamma}_i = \hat{\gamma}$, for all i .

Step 9 Now apply the argument in Steps 4–8 to a set of generators $\gamma^{(1)}, \dots, \gamma^{(2g)}$ of $\pi_1(\Sigma)$. We conclude that, along some subsequence,

$$\hat{\gamma}^{(j)} = \sigma_i^{-1} \circ \hat{\sigma}_i(\gamma^{(j)}), \quad j = 1, \dots, 2g$$

(see (4)). But then the automorphisms $\sigma_i^{-1} \circ \hat{\sigma}_i$ are constant on all of $\pi_1(\Sigma)$. Since $\hat{\sigma}_i$ converges, so does σ_i , contradicting the hypothesis of no limit points for the sequence $\{X_i\}$.

This contradiction completes the proof. □

References

- [1] **F Bonahon**, *Bouts des variétés hyperboliques de dimension 3*, Ann. of Math. 124 (1986) 71–158 [MR847953](#)
- [2] **K Corlette**, *Flat G -bundles with canonical metrics*, J. Differential Geom. 28 (1988) 361–382 [MR965220](#)

- [3] **M Culler, J W Morgan**, *Group actions on \mathbb{R} -trees*, Proc. London Math. Soc. 55 (1987) 571–604 [MR907233](#)
- [4] **M Culler, P B Shalen**, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. 117 (1983) 109–146 [MR683804](#)
- [5] **G Daskalopoulos, S Dostoglou, R Wentworth**, *On the Morgan–Shalen compactification of the $SL(2, \mathbb{C})$ character varieties of surface groups*, Duke Math. J. 101 (2000) 189–207 [MR1738182](#)
- [6] **S K Donaldson**, *Twisted harmonic maps and the self-duality equations*, Proc. London Math. Soc. 55 (1987) 127–131 [MR887285](#)
- [7] **D Gallo, M Kapovich, A Marden**, *The monodromy groups of Schwarzian equations on closed Riemann surfaces*, Ann. of Math. 151 (2000) 625–704 [MR1765706](#)
- [8] **W M Goldman, R A Wentworth**, *Energy of twisted harmonic maps of Riemann surfaces*, from: “In the tradition of Ahlfors–Bers, IV”, (D Canary, J Gilman, J Heinonen, H Masur, editors), Contemp. Math. 432, Amer. Math. Soc., Providence, RI (2007) 45–61 [MR2342806](#)
- [9] **N J Hitchin**, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. 55 (1987) 59–126 [MR887284](#)
- [10] **N Hitchin**, *Stable bundles and integrable systems*, Duke Math. J. 54 (1987) 91–114 [MR885778](#)
- [11] **J Hubbard, H Masur**, *Quadratic differentials and foliations*, Acta Math. 142 (1979) 221–274 [MR523212](#)
- [12] **J Jost**, *Equilibrium maps between metric spaces*, Calc. Var. Partial Differential Equations 2 (1994) 173–204 [MR1385525](#)
- [13] **Y Kamishima, S P Tan**, *Deformation spaces on geometric structures*, from: “Aspects of low-dimensional manifolds”, (Y Matsumoto, S Morita, editors), Adv. Stud. Pure Math. 20, Kinokuniya, Tokyo (1992) 263–299 [MR1208313](#)
- [14] **N J Korevaar, R M Schoen**, *Sobolev spaces and harmonic maps for metric space targets*, Comm. Anal. Geom. 1 (1993) 561–659 [MR1266480](#)
- [15] **R S Kulkarni, U Pinkall**, *A canonical metric for Möbius structures and its applications*, Math. Z. 216 (1994) 89–129 [MR1273468](#)
- [16] **A Lubotzky, A R Magid**, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. 336, Amer. Math. Soc, Providence, RI (1985) [MR818915](#)
- [17] **R C Penner, J L Harer**, *Combinatorics of train tracks*, Annals of Mathematics Studies 125, Princeton Univ. Press, NJ (1992) [MR1144770](#)
- [18] **J Sacks, K Uhlenbeck**, *Minimal immersions of closed Riemann surfaces*, Trans. Amer. Math. Soc. 271 (1982) 639–652 [MR654854](#)

- [19] **L P Schaposnik**, *Spectral data for G -Higgs bundles*, PhD thesis, University of Oxford (2013) [MR3389247](#) Available at <http://search.proquest.com/docview/1683347472>
- [20] **R Schoen, S T Yau**, *Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature*, Ann. of Math. 110 (1979) 127–142 [MR541332](#)
- [21] **R A Wentworth**, *Energy of harmonic maps and Gardiner's formula*, from: “In the tradition of Ahlfors–Bers, IV”, (D Canary, J Gilman, J Heinonen, H Masur, editors), Contemp. Math. 432, Amer. Math. Soc., Providence, RI (2007) 221–229 [MR2342819](#)
- [22] **M Wolf**, *Harmonic maps from surfaces to \mathbb{R} -trees*, Math. Z. 218 (1995) 577–593 [MR1326987](#)
- [23] **M Wolf**, *On realizing measured foliations via quadratic differentials of harmonic maps to \mathbb{R} -trees*, J. Anal. Math. 68 (1996) 107–120 [MR1403253](#)
- [24] **M Wolf**, *Measured foliations and harmonic maps of surfaces*, J. Differential Geom. 49 (1998) 437–467 [MR1669653](#)

Department of Mathematics, University of Maryland
College Park, MD 20742, United States

Department of Mathematics, Rice University
MS 136, Houston, TX 77005-1892, United States

raw@math.umd.edu, mwolf@rice.edu

Proposed: Benson Farb
Seconded: Bruce Kleiner, John Lott

Received: 3 August 2015
Accepted: 2 January 2016