

On the second homology group of the Torelli subgroup of $\text{Aut}(F_n)$

MATTHEW DAY
ANDREW PUTMAN

Let IA_n be the Torelli subgroup of $\text{Aut}(F_n)$. We give an explicit finite set of generators for $H_2(\text{IA}_n)$ as a $\text{GL}_n(\mathbb{Z})$ -module. Corollaries include a version of surjective representation stability for $H_2(\text{IA}_n)$, the vanishing of the $\text{GL}_n(\mathbb{Z})$ -coinvariants of $H_2(\text{IA}_n)$, and the vanishing of the second rational homology group of the level ℓ congruence subgroup of $\text{Aut}(F_n)$. Our generating set is derived from a new group presentation for IA_n which is infinite but which has a simple recursive form.

20E05, 20E36, 20F05, 20J06

1 Introduction

The *Torelli subgroup* of the automorphism group of a free group F_n on n letters, denoted by IA_n , is the kernel of the action of $\text{Aut}(F_n)$ on $F_n^{\text{ab}} \cong \mathbb{Z}^n$. The group of automorphisms of \mathbb{Z}^n is $\text{GL}_n(\mathbb{Z})$ and the resulting map $\text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$ is easily seen to be surjective, so we have a short exact sequence

$$1 \rightarrow \text{IA}_n \rightarrow \text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z}) \rightarrow 1.$$

Though it has a large literature, the cohomology and combinatorial group theory of IA_n remain quite mysterious. Magnus [26] proved that IA_n is finitely generated, and thus that $H_1(\text{IA}_n)$ has finite rank. Krstić and McCool [24] later showed that IA_3 is not finitely presentable. This was improved by Bestvina, Bux and Margalit [4], who showed that $H_2(\text{IA}_3)$ has infinite rank. However, for $n \geq 4$ it is not known whether or not IA_n is finitely presentable, or whether or not $H_2(\text{IA}_n)$ has finite rank.

Representation-theoretic finiteness It seems to be very difficult to determine whether or not $H_2(\text{IA}_n)$ has finite rank, so it is natural to investigate weaker finiteness properties. Since inner automorphisms act trivially on homology, the conjugation action of $\text{Aut}(F_n)$ on IA_n induces an action of $\text{GL}_n(\mathbb{Z})$ on $H_k(\text{IA}_n)$. Church and Farb [12, Conjecture 6.7] conjectured that $H_k(\text{IA}_n)$ is finitely generated as a $\text{GL}_n(\mathbb{Z})$ -module. In other words, they conjectured that there exists a finite subset of $H_k(\text{IA}_n)$ whose $\text{GL}_n(\mathbb{Z})$ -orbit spans $H_k(\text{IA}_n)$. Our first main theorem verifies their conjecture for $k = 2$.

Theorem A (generators for $H_2(\text{IA}_n)$) For all $n \geq 2$, there exists a finite subset of $H_2(\text{IA}_n)$ whose $\text{GL}_n(\mathbb{Z})$ -orbit spans $H_2(\text{IA}_n)$.

Each element of our finite subset corresponds to a map of a surface into a classifying space for IA_n ; the genera of these surfaces range from 1 to 3. Table 1 below lists our finite set of $\text{GL}_n(\mathbb{Z})$ -generators for $H_2(\text{IA}_n)$. This table expresses these generators using specific “commutator relators” in IA_n ; see below for how to translate these into elements of $H_2(\text{IA}_n)$.

Remark 1.1 The special case $n = 3$ of Theorem A was proven in the unpublished thesis of Owen Baker [2]. His proof uses a “Jacobian” map on outer space and is quite different from our proof. It seems difficult to generalize his proof to higher n .

Surjective representation stability The generators for $H_2(\text{IA}_n)$ given in Theorem A are explicit enough that they can be used to perform a number of interesting calculations. The first verifies part of a conjecture of Church and Farb, which asserts that the homology groups of IA_n are “representation stable”. We begin with some background. An increasing sequence

$$G_1 \subset G_2 \subset G_3 \subset \dots$$

of groups is *homologically stable* if for all $k \geq 1$, the k^{th} homology group of G_n is independent of n for $n \gg 0$. Many sequences of groups are homologically stable; see Hatcher and Wahl [21] for a bibliography. In particular, Hatcher and Vogtmann [20] proved this for $\text{Aut}(F_n)$. However, it is known that IA_n is *not* homologically stable; indeed, even $H_1(\text{IA}_n)$ does not stabilize (see below).

Church and Farb [12] introduced a new form of homological stability for groups like IA_n whose homology groups possess natural group actions. For IA_n , they conjectured that for all $k \geq 1$, there exists some $n_k \geq 1$ such that the following two properties hold for all $n \geq n_k$:

- **Injective stability** The map $H_k(\text{IA}_n) \rightarrow H_k(\text{IA}_{n+1})$ is injective.
- **Surjective representation stability** The map $H_k(\text{IA}_n) \rightarrow H_k(\text{IA}_{n+1})$ is surjective “up to the action of $\text{GL}_{n+1}(\mathbb{Z})$ ”; more precisely, the $\text{GL}_{n+1}(\mathbb{Z})$ -orbit of its image spans $H_k(\text{IA}_{n+1})$.

Remark 1.2 In fact, they made this conjecture in [11] for the Torelli subgroup of the mapping class group; however, they have informed us that they also conjecture it for IA_n .

Our generators for $H_2(\text{IA}_n)$ are “the same in each dimension” starting at $n = 6$, so we are able to derive the following special case of Church and Farb’s conjecture:

Theorem B (surjective representation stability for $H_2(\text{IA}_n)$) *The $\text{GL}_{n+1}(\mathbb{Z})$ -orbit of the image of the natural map $H_2(\text{IA}_n) \rightarrow H_2(\text{IA}_{n+1})$ spans $H_2(\text{IA}_{n+1})$ for $n \geq 6$.*

Remark 1.3 Boldsen and Hauge Døllerup [5] proved a theorem similar to **Theorem B** for the *rational* second homology group of the Torelli subgroup of the mapping class group. Their proof is different from ours; in particular, they were not able to prove an analogue of **Theorem A**. It seems hard to use their techniques to prove **Theorem B**. Similarly, our proof uses special properties of IA_n and does not work for the Torelli subgroup of the mapping class group.

Coinvariants Our tools do not allow us to easily distinguish different homology classes; indeed, for all we know our generators for $H_2(\text{IA}_n)$ might be redundant. This prevents us from proving injective stability for $H_2(\text{IA}_n)$. However, we still can prove some interesting vanishing results. If G is a group and M is a G -module, then the *coinvariants* of G acting on M , denoted by M_G , are the largest quotient of M on which G acts trivially. More precisely, $M_G = M/K$ with $K = \langle m - g \cdot m \mid g \in G, m \in M \rangle$. We then have the following.

Theorem C (vanishing coinvariants) *For $n \geq 6$, we have $(H_2(\text{IA}_n))_{\text{GL}_n(\mathbb{Z})} = 0$.*

Remark 1.4 Church and Farb [12, Conjecture 6.5] conjectured that the $\text{GL}_n(\mathbb{Z})$ -invariants in $H^k(\text{IA}_n; \mathbb{Q})$ are 0. For $k = 1$, this follows from the known computation of $H^1(\text{IA}_n; \mathbb{Q})$; see below. **Theorem C** implies that this also holds for $k = 2$.

Linear congruence subgroups For $\ell \geq 2$, the *level ℓ congruence subgroup* of $\text{Aut}(F_n)$, denoted by $\text{Aut}(F_n, \ell)$, is the kernel of the natural map $\text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z}/\ell)$; one should think of it as a “mod- ℓ ” version of IA_n . It is natural to conjecture that for all $k \geq 1$, there exists some $n_k \geq 1$ such that $H_k(\text{Aut}(F_n, \ell); \mathbb{Q}) \cong H_k(\text{Aut}(F_n); \mathbb{Q})$ for $n \geq n_k$; an analogous theorem for congruence subgroups of $\text{GL}_n(\mathbb{Z})$ is due to Borel [7]. Galatius [19] proved that $H_k(\text{Aut}(F_n); \mathbb{Q}) = 0$ for $n \gg 0$, so this conjecture really asserts that $H_k(\text{Aut}(F_n, \ell); \mathbb{Q}) = 0$ for $n \gg 0$. The case $k = 1$ of this is known. Indeed, Satoh [31] calculated the abelianization of $\text{Aut}(F_n, \ell)$ for $n \geq 3$ and the answer consisted entirely of torsion, so $H_1(\text{Aut}(F_n, \ell); \mathbb{Q}) = 0$ for $n \geq 3$. Using **Theorem A**, we will prove the case $k = 2$.

Theorem D (second homology of congruence subgroups) *For $\ell \geq 2$ and $n \geq 6$, we have $H_2(\text{Aut}(F_n, \ell); \mathbb{Q}) = 0$.*

The key to our proof is that **Theorem A** allows us to show that the image of $H_2(\text{IA}_n; \mathbb{Q})$ in $H_2(\text{Aut}(F_n, \ell); \mathbb{Q})$ vanishes; this allows us to derive **Theorem D** using standard techniques.

Remark 1.5 The second author proved an analogue of [Theorem D](#) for congruence subgroups of the mapping class group in [30]. The techniques in [30] are different from those in the present paper and it seems difficult to prove [Theorem D](#) via those techniques.

Basic elements of Torelli We now wish to describe our generating set for $H_2(\text{IA}_n)$. This requires introducing some basic elements of IA_n . Let $\{x_1, \dots, x_n\}$ be a free basis for F_n . We then make the following definitions:

- For distinct $1 \leq i, j \leq n$, let $C_{x_i, x_j} \in \text{IA}_n$ be defined via the formulas

$$C_{x_i, x_j}(x_i) = x_j x_i x_j^{-1} \quad \text{and} \quad C_{x_i, x_j}(x_\ell) = x_\ell \quad \text{if } \ell \neq i.$$

- For $\alpha, \beta, \gamma \in \{\pm 1\}$ and distinct $1 \leq i, j, k \leq n$, let $M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]} \in \text{IA}_n$ be defined via the formulas

$$M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]}(x_i^\alpha) = [x_j^\beta, x_k^\gamma] x_i^\alpha \quad \text{and} \quad M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]}(x_\ell) = x_\ell \quad \text{if } \ell \neq i.$$

Observe that by definition

$$M_{x_i^{-1}, [x_j^\beta, x_k^\gamma]}(x_i^{-1}) = [x_j^\beta, x_k^\gamma] x_i^{-1} \quad \text{and} \quad M_{x_i^{-1}, [x_j^\beta, x_k^\gamma]}(x_i) = x_i [x_j^\beta, x_k^\gamma]^{-1}.$$

We call C_{x_i, x_j} a *conjugation move* and $M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]}$ a *commutator transvection*.

Surfaces in a classifying space: our generators A *commutator relator* in IA_n is a formula of the form $[a_1, b_1] \cdots [a_g, b_g] = 1$ with $a_i, b_i \in \text{IA}_n$. Given such a commutator relator r , let Σ_g be a genus g surface. There is a continuous map $\zeta: \Sigma_g \rightarrow K(\text{IA}_n, 1)$ that takes the standard basis for $\pi_1(\Sigma_g)$ to $a_1, b_1, \dots, a_g, b_g \in \text{IA}_n$. We obtain an element $\mathfrak{h}_r = \zeta_*([\Sigma_g]) \in H_2(\text{IA}_n)$. With this notation, the generators for $H_2(\text{IA}_n)$ given by [Theorem A](#) are the elements \mathfrak{h}_r where r is one of the relators in [Table 1](#).

The Johnson homomorphism To motivate our proof of [Theorem A](#), we first recall the computation of $H_1(\text{IA}_n)$, which is due independently to Farb [18], Kawazumi [23] and Cohen and Pakianathan [14]. The basic tool is the Johnson homomorphism [22], which was introduced in the context of the Torelli subgroup of the mapping class group (though it also appears in earlier work of Andreadakis [1]). See Satoh [32] for a survey of the IA_n version of it. The Johnson homomorphism is a homomorphism

$$\tau: \text{IA}_n \rightarrow \text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$$

that arises from studying the action of IA_n on the second nilpotent truncation of F_n . It can be defined as follows. For $z \in F_n$, let $[z] \in \mathbb{Z}^n$ be the associated element of the abelianization of F_n . Consider $f \in \text{IA}_n$. For $x \in F_n$, we have $f(x) \cdot x^{-1} \in [F_n, F_n]$.

- (H1) $[C_{x_a, x_b}, C_{x_c, x_d}] = 1$, possibly with $b = d$.
- (H2) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}] = 1$, possibly with $\{b, c\} \cap \{e, f\} \neq \emptyset$, or with $x_a^\alpha = x_d^{-\delta}$ as long as $x_a^\alpha \neq x_d^\delta$, $a \notin \{e, f\}$ and $d \notin \{b, c\}$.
- (H3) $[C_{x_a, x_b}, M_{x_c^\gamma, [x_d^\delta, x_e^\epsilon]}] = 1$,
possibly with $b \in \{d, e\}$ if $c \notin \{a, b\}$ and $a \notin \{c, d, e\}$.
- (H4) $[C_{x_c, x_b}^\beta C_{x_a, x_b}^\beta, C_{x_c, x_a}^\alpha] = 1$.
- (H5) $[C_{x_a, x_c}^{-\gamma}, C_{x_a, x_d}^{-\delta}][C_{x_a, x_b}^{-\beta}, M_{x_b^\beta, [x_c^\gamma, x_d^\delta]}] = 1$.
- (H6) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}][M_{x_d^\delta, [x_a^\alpha, x_e^\epsilon]}, M_{x_d^\delta, [x_c^\gamma, x_b^\beta]}] \times$
 $[M_{x_d^\delta, [x_c^\gamma, x_b^\beta]}, C_{x_d, x_e}^{-\epsilon}] = 1$, possibly with $b = e$ or $c = e$.
- (H7) $[M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]}, C_{x_a, x_b}^\beta][C_{x_c, x_d}^{-\delta}, M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]}][M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]}, M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]}] = 1$,
possibly with $b = d$.
- (H8) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, C_{x_a, x_d}^\delta C_{x_b, x_d}^\delta C_{x_c, x_d}^\delta] = 1$.
- (H9) $[C_{x_a, x_c}^\gamma C_{x_b, x_c}^\gamma, C_{x_a, x_b}^\beta C_{x_c, x_b}^\beta][M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, C_{x_b, x_a}^\alpha C_{x_c, x_a}^\alpha] = 1$.

Table 1: The set of commutator relators whose associated elements of $H_2(\text{IA}_n)$ generate it as a $\text{GL}_n(\mathbb{Z})$ -module. Distinct letters represent distinct indices unless stated otherwise.

There is a natural surjection $\rho: [F_n, F_n] \rightarrow \wedge^2 \mathbb{Z}^n$ satisfying $\rho([a, b]) = [a] \wedge [b]$; the kernel of ρ is $[F_n, [F_n, F_n]]$. We can then define a map $\tilde{\tau}_f: F_n \rightarrow \wedge^2 \mathbb{Z}^n$ via the formula $\tilde{\tau}_f(x) = \rho(f(x) \cdot x^{-1})$. One can check that $\tilde{\tau}_f$ is a homomorphism. It factors through a homomorphism $\tau_f: \mathbb{Z}^n \rightarrow \wedge^2 \mathbb{Z}^n$. We can then define $\tau: \text{IA}_n \rightarrow \text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$ via the formula $\tau(f) = \tau_f$. One can check that τ is a homomorphism.

Generators and their images Define

$$S_{\text{MA}}(n) = \{C_{x_i, x_j} \mid 1 \leq i, j \leq n \text{ distinct}\} \cup \{M_{x_i, [x_j, x_k]} \mid 1 \leq i, j, k \leq n \text{ distinct, } j < k\}.$$

Magnus [26] proved that IA_n is generated by $S_{\text{MA}}(n)$; see Day and Putman [16] and Bestvina, Bux and Margalit [4] for modern proofs. For distinct $1 \leq i, j \leq n$, the image $\tau(C_{x_i, x_j}) \in \text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$ is the homomorphism defined via the formulas

$$[x_i] \mapsto [x_j] \wedge [x_i] \quad \text{and} \quad [x_\ell] \mapsto 0 \quad \text{if } \ell \neq i.$$

Similarly, for distinct $1 \leq i, j, k \leq n$ with $j < k$, the image $\tau(M_{x_i, [x_j, x_k]})$ in $\text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$ is the homomorphism defined via the formulas

$$[x_i] \mapsto [x_j] \wedge [x_k] \quad \text{and} \quad [x_\ell] \mapsto 0 \quad \text{if } \ell \neq i.$$

The key observation is that these form a *basis* for $\text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$.

The abelianization Let $F(S_{MA}(n))$ be the free group on $S_{MA}(n)$ and let $R_{MA}(n) \subset F(S_{MA}(n))$ be a set of relations for IA_n , so $IA_n = \langle S_{MA}(n) \mid R_{MA}(n) \rangle$. Since τ takes $S_{MA}(n)$ bijectively to a basis for the free abelian group $\text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$, we must have $R_{MA}(n) \subset [F(S_{MA}(n)), F(S_{MA}(n))]$. This immediately implies that $H_1(IA_n) \cong \text{Hom}(\mathbb{Z}^n, \wedge^2 \mathbb{Z}^n)$.

Hopf’s formula But even more is true. Recall that Hopf’s formula (see Brown [10]) says that if G is a group with a presentation $G = \langle S \mid R \rangle$, then

$$H_2(G) \cong \frac{\langle\langle R \rangle\rangle \cap [F(S), F(S)]}{[F(S), \langle\langle R \rangle\rangle]},$$

here $\langle\langle R \rangle\rangle$ is the normal closure of R . The intersection in the numerator of this is usually hard to calculate, so Hopf’s formula is not often useful for computation. However, by what we have said it simplifies for IA_n to

$$(1) \quad H_2(IA_n) \cong \frac{\langle\langle R_{MA}(n) \rangle\rangle}{[F(S_{MA}(n)), \langle\langle R_{MA}(n) \rangle\rangle]}.$$

This isomorphism is very concrete: an element $r \in \langle\langle R_{MA}(n) \rangle\rangle$ is a commutator relator, and the associated element of $H_2(IA_n)$ is the homology class \mathfrak{h}_r discussed above.

Summary and trouble For $r \in \langle\langle R_{MA}(n) \rangle\rangle$ and $z \in F(S_{MA}(n))$, the element $zrz^{-1}r^{-1}$ lies in the denominator of (1), $[F(S_{MA}(n)), \langle\langle R_{MA}(n) \rangle\rangle]$. Hence $\mathfrak{h}_{zrz^{-1}} = \mathfrak{h}_r$. It follows that $H_2(IA_n)$ is generated by the set $\{\mathfrak{h}_r \mid r \in R_{MA}(n)\}$. In other words, to calculate generators for $H_2(IA_n)$, it is enough to find a presentation for IA_n with $S_{MA}(n)$ as its generating set. However, this seems like a difficult problem (especially if, as we suspect, IA_n is not finitely presentable). Moreover, the $\text{GL}_n(\mathbb{Z})$ -action on $H_2(IA_n)$ has not yet appeared.

L-presentations To incorporate the $\text{GL}_n(\mathbb{Z})$ -action on $H_2(IA_n)$ into our presentation for IA_n , we use the notion of an L-presentation, which was introduced by Bartholdi [3] (we use a slight simplification of his definition). An L-presentation for a group G is a triple $\langle S \mid R^0 \mid E \rangle$, where S and R^0 and E are as follows:

- S is a generating set for G .
- $R^0 \subset F(S)$ is a set consisting of relations for G (not necessarily complete).
- E is a subset of $\text{End}(F(S))$.

This data must satisfy the following condition: Let $M \subset \text{End}(F(S))$ be the monoid generated by E . Define $R = \{f(r) \mid f \in M, r \in R^0\}$. Then we require $G = \langle S \mid R \rangle$. Each element of E descends to an element of $\text{End}(G)$; we call the resulting subset $\tilde{E} \subset \text{End}(G)$ the *induced endomorphisms* of our L-presentation. We say that our L-presentation is *finite* if the sets S and R^0 and E are all finite.

In our examples, the induced endomorphisms of our L–presentations will actually be automorphisms. Thus in our context one should think of an L–presentation as a group presentation incorporating certain symmetries of a group. Here is an easy example:

Example Let $S = \{z_i \mid i \in \mathbb{Z}/p\}$ and $R^0 = \{z_0^2\}$. Let $\psi: F(S) \rightarrow F(S)$ be the homomorphism defined via the formula $\psi(z_i) = z_{i+1}$. Then $\langle S \mid R^0 \mid \{\psi\} \rangle$ is an L–presentation for the free product of p copies of $\mathbb{Z}/2$.

A finite L–presentation for Torelli The conjugation action of $\text{Aut}(F_n)$ on IA_n gives an injection $\text{Aut}(F_n) \hookrightarrow \text{Aut}(\text{IA}_n)$. If we could somehow construct a finite L–presentation $\langle S_{\text{MA}}(n) \mid R_{\text{MA}}^0(n) \mid E_{\text{MA}}(n) \rangle$ for IA_n whose set of induced endomorphisms generated

$$\text{Aut}(F_n) \subset \text{Aut}(\text{IA}_n) \subset \text{End}(\text{IA}_n),$$

then **Theorem A** would immediately follow. Indeed, since the $\text{GL}_n(\mathbb{Z})$ –action on $H_2(\text{IA}_n)$ is induced by the conjugation action of $\text{Aut}(F_n)$ on IA_n , it would follow that the $\text{GL}_n(\mathbb{Z})$ –orbit of the set $\{\mathfrak{h}_r \mid r \in R_{\text{MA}}^0(n)\} \subset H_2(\text{IA}_n)$ spanned $H_2(\text{IA}_n)$.

Although we find the idea in the previous paragraph illuminating, we do not follow it strictly. To make our L–presentation for IA_n easier to comprehend, we will use the following generating set, which is larger than S_{MA} :

$$S_{\text{IA}}(n) = \{C_{x_i, x_j} \mid 1 \leq i, j \leq n \text{ distinct}\} \\ \cup \{M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]} \mid 1 \leq i, j, k \leq n \text{ distinct}, \alpha, \beta, \gamma \in \{\pm 1\}\}.$$

This has the advantage of making our relations and rewriting rules shorter, and making their meaning easier to understand. It has the disadvantage of making the proof of **Theorem A** less direct. Our theorem giving an L–presentation for IA_n is as follows:

Theorem E (finite L–presentation for Torelli) *For all $n \geq 2$, there exists a finite L–presentation $\text{IA}_n = \langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle$ whose set of induced endomorphisms generates $\text{Aut}(F_n) \subset \text{Aut}(\text{IA}_n) \subset \text{End}(\text{IA}_n)$.*

We note that our presentation is not a presentation in which all relators are commutators. The formulas for the $R_{\text{IA}}^0(n)$ and $E_{\text{IA}}(n)$ in our finite L–presentation are a little complicated, so we postpone them until **Section 2**. The formulas in that section make it clear that $R_{\text{IA}}(n)$ does not lie in $[F(S_{\text{IA}}(n)), F(S_{\text{IA}}(n))]$. Therefore we cannot prove **Theorem A** simply by interpreting the relators as homology classes. We must do something more complicated to deduce that theorem from our presentation.

Remark 1.6 The relations in **Table 1** are not sufficient for our L–presentation. Indeed, they all lie in the commutator subgroup, but the generators $S_{\text{IA}}(n)$ do not map to linearly independent elements of the abelianization.

Sketch of proof We close this introduction by briefly discussing how we prove [Theorem E](#). In particular, we explain why it is easier to verify an L -presentation than a standard presentation. We remark that our proof is inspired by a recent paper [9] of the second author together with Brendle and Margalit, which constructed generators for the kernel of the Burau representation evaluated at -1 .

Assume that we have guessed a finite L -presentation $\langle S_{IA}(n) \mid R_{IA}^0(n) \mid E_{IA}(n) \rangle$ for IA_n as in [Theorem E](#) (we found the one that we use by first throwing in all the relations we could think of and then attempting the proof below; each time it failed it revealed a relation that we had missed). Let Q_n be the group presented by the purported L -presentation. There is thus a surjection $\pi: Q_n \rightarrow IA_n$, and the goal of our proof will be to construct an inverse map $\phi: IA_n \rightarrow Q_n$ satisfying $\phi \circ \pi = \text{id}$. This will involve several steps.

Step 1 We decompose IA_n in terms of stabilizers of conjugacy classes of primitive elements of F_n .

For $z \in F_n$, let $[[z]]$ denote the union of the conjugacy classes of z and z^{-1} . A *primitive element* of F_n is an element that forms part of a free basis. Let $\mathcal{C} = \{[[z]] \mid z \in F_n \text{ primitive}\}$. The set \mathcal{C} forms the set of vertices of a simplicial complex called the *complex of partial bases*, which is analogous to the complex of curves for the mapping class group. Applying a theorem of the second author [29] to the action of IA_n on the complex of partial bases, we will obtain a decomposition

$$(2) \quad IA_n = \ast_{c \in \mathcal{C}} (IA_n)_c / (\text{some relations});$$

here $(IA_n)_c$ denotes the stabilizer in IA_n of c . The unlisted relations play only a small role in our proof and can be ignored at this point.

Step 2 We use induction to construct a partial inverse.

Fix some $c_0 \in \mathcal{C}$. The stabilizer $(IA_n)_{c_0}$ is very similar to IA_{n-1} ; in fact, it is connected to IA_{n-1} by an exact sequence that is analogous to the Birman exact sequence for the mapping class group. We construct this exact sequence in the companion paper [17], which builds on our previous paper [15]. By analyzing this exact sequence and using induction, we will construct a “partial inverse” $\phi_{c_0}: (IA_n)_{c_0} \rightarrow Q_n$. We remark that this step is where most of our relations arise — they actually are relations in the kernel of the Birman exact sequence we construct in [15].

Step 3 We use the L -presentation to lift the conjugation action of $\text{Aut}(F_n)$ on IA_n to Q_n .

Let \tilde{E} be the induced endomorphisms of our L -presentation. We directly prove that these endomorphisms actually give an action of $\text{Aut}(F_n)$ on Q_n such that the projection

map $\pi: \mathcal{Q}_n \rightarrow \text{IA}_n$ is equivariant. This is the key place where we use properties of L -presentations; in general, it is difficult to construct group actions on groups given by generators and relations.

Step 4 We use our group action to construct the inverse.

The conjugation action of $\text{Aut}(F_n)$ on IA_n transitively permutes the terms of (2). Using our lifted action of $\text{Aut}(F_n)$ on \mathcal{Q}_n as a “guide”, we then “move” the partially defined inverse ϕ_{c_0} around and construct ϕ on the rest of IA_n , completing the proof.

Remark 1.7 In [28], the second author constructed an infinite presentation of the Torelli subgroup of the mapping class group. Though this used the same result [29] that we quoted above, the details are quite different. One source of this difference is that instead of an L -presentation with a finite generating set, the paper [28] constructed an ordinary presentation with an infinite generating set.

Computer calculations At several places in this paper, we will need to verify large numbers of equations in group presentations. Rather than displaying these equations in the paper or leaving them as exercises, we use the GAP System to store and check our equations mechanically. The code to verify these equations is in the file `h2ia.g`, which is in an [online supplement](#). We found the equations in this file by hand, and our proof does not rely on a computer search. We will say more about this in [Section 5](#), where said calculations begin.

This is a good place to note that our results rely strongly on the authors’ earlier paper [17] and the computer calculations from that paper. There we use a similar approach to automatically verify identities and prove the existence of certain homomorphisms between groups given by presentations. This is used in more than one place in the present paper, but most crucially in [Proposition 3.14](#). The computations from our earlier paper are in a file `iabes.g`, which is available on the authors’ websites and on arXiv.

Outline We begin in [Section 2](#) by giving a precise statement of the L -presentation whose existence is asserted in [Theorem E](#). Next, in [Section 3](#) we discuss several tools that are needed for the proof of [Theorem E](#). The proof of [Theorem E](#) is in [Section 4](#). This proof depends on some combinatorial group theory calculations that are stated in [Sections 2](#) and [3](#) but whose proofs are postponed until [Section 5](#). In [Section 6](#), we prove [Theorem A](#). That section also shows how to derive [Theorem B](#) from [Theorem A](#). Finally, [Theorems C](#) and [D](#) are proven in [Section 7](#).

Acknowledgements Day was supported in part by NSF grant DMS-1206981 and Putman was supported in part by NSF grant DMS-1255350 and the Alfred P Sloan Foundation.

- (R0) $M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}^{-1} = M_{x_a^\alpha, [x_c^\gamma, x_b^\beta]}$.
- (R1) $[C_{x_a, x_b}, C_{x_c, x_d}] = 1$, possibly with $b = d$.
- (R2) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}] = 1$, possibly with $\{b, c\} \cap \{e, f\} \neq \emptyset$, or with $x_a^\alpha = x_d^{-\delta}$ as long as $x_a^\alpha \neq x_d^\delta$, $a \notin \{e, f\}$ and $d \notin \{b, c\}$.
- (R3) $[C_{x_a, x_b}, M_{x_c^\gamma, [x_d^\delta, x_e^\epsilon]}] = 1$,
possibly with $b \in \{d, e\}$ if $c \notin \{a, b\}$ and $a \notin \{c, d, e\}$.
- (R4) $[C_{x_a, x_b} C_{x_c, x_b}, C_{x_c, x_a}] = 1$.
- (R5) $C_{x_a, x_b}^\beta M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]} C_{x_a, x_b}^{-\beta} = M_{x_a^\alpha, [x_c^\gamma, x_b^{-\beta}]}$.
- (R6) $M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]} M_{x_a^{-\alpha}, [x_b^\beta, x_c^\gamma]} = [C_{x_a, x_c}^{-\gamma}, C_{x_a, x_b}^{-\beta}]$.
- (R7) $[C_{x_a, x_b}^{-\beta}, M_{x_b^\beta, [x_c^\gamma, x_d^\delta]}] = [C_{x_a, x_d}^{-\delta}, C_{x_a, x_c}^{-\gamma}]$.
- (R8) $M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]} M_{x_d^\delta, [x_a^\alpha, x_e^\epsilon]} M_{x_a^\alpha, [x_c^\gamma, x_b^\beta]} =$
 $C_{x_d, x_e}^{-\epsilon} M_{x_d^\delta, [x_c^\gamma, x_b^\beta]} C_{x_d, x_e}^\epsilon M_{x_d^\delta, [x_a^\alpha, x_e^\epsilon]} M_{x_d^\delta, [x_b^\beta, x_c^\gamma]}$, possibly with $e \in \{b, c\}$.
- (R9) $C_{x_a, x_b}^\beta M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]} C_{x_a, x_b}^{-\beta} =$
 $C_{x_c, x_d}^{-\delta} M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]} C_{x_c, x_d}^\delta M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]} M_{x_c^\gamma, [x_b^\beta, x_a^\alpha]}$, possibly with $b = d$.

Table 2: Basic relations for the L–presentation of IA_n . Distinct letters are assumed to represent distinct indices unless stated otherwise. Let $R_{IA}(n)$ denote the finite set of all relations from the above ten classes.

2 Our finite L–presentation

We now discuss the relations $R_{IA}^0(n)$ and endomorphisms $E_{IA}(n)$ of our L–presentation. Two calculations (Propositions 2.1 and 2.3) are postponed until Section 5.

Relations Our set $R_{IA}^0(n)$ of relations consists of the relations in Table 2. It is easy to verify that these are all indeed relations:

Proposition 2.1 *The relations $R_{IA}^0(n)$ all hold when interpreted in IA_n .*

The proof is computational and is postponed until Section 5. We remark that unlike many of our computer calculations, it is not particularly difficult to verify by hand.

Since the relations are rather complicated we suggest to the reader that they not pay too close attention to them on their first pass through the paper. The overall structure of our proof (and, in fact, the majority of its details) can be understood without much knowledge of our relations.

Remark 2.2 The relations in $R_{IA}^0(n)$ have reasonable intuitive interpretations. Relations (R1)–(R3) state that generators acting only in different places commute with

each other. Relation (R4) is a generalization of the fact that for $n = 3$, the conjugation move C_{x_3, x_1} conjugates the inner automorphism $C_{x_1, x_2}C_{x_3, x_2}$ back to itself (since it fixes the conjugating element x_2). Relation (R5) makes sense by looking at either side of x_a : on the right of x_a^α , instances of $x_b^{\pm\beta}$ cancel, but on the left side of x_a^α , we get a conjugate of a basic commutator that is itself a basic commutator. Relation (R6) states that conjugation by a commutator is the same as acting by a commutator of conjugation moves. Relations (R7)–(R9) allow us to rewrite a conjugate of a generator acting on a given element as a product of generators acting only on that same element (x_a , x_d or x_c as stated here, respectively). In this sense, these relations are like the Steinberg relations from the presentation of $\text{GL}_n(\mathbb{Z})$ in algebraic K–theory.

Generators for the automorphism group of a free group Before discussing our endomorphisms $E_{\text{IA}}(n)$, we first introduce a generating set for $\text{Aut}(F_n)$ that goes back to work of Nielsen. For $\alpha = \pm 1$ and distinct $1 \leq i, j \leq n$, let $M_{x_i^\alpha, x_j} \in \text{Aut}(F_n)$ be the *transvection* that takes x_i^α to $x_j x_i^\alpha$ and fixes x_ℓ for $\ell \neq i$. Just like before, we have

$$M_{x_i^{-1}, x_j}(x_i^{-1}) = x_j x_i^{-1} \quad \text{and} \quad M_{x_i^{-1}, x_j}(x_i) = x_i x_j^{-1}.$$

Next, for distinct $1 \leq i, j \leq n$ let $P_{i, j} \in \text{Aut}(F_n)$ be the *swap automorphism* that exchanges x_i and x_j while fixing x_ℓ for $\ell \neq i, j$. Finally, for $1 \leq i \leq n$ let $I_i \in \text{Aut}(F_n)$ be the *inversion automorphism* that takes x_i to x_i^{-1} and fixes x_ℓ for $\ell \neq i$. Define

$$S_{\text{Aut}}(n) = \{M_{x_i^\alpha, x_j}^\beta \mid 1 \leq i, j \leq n \text{ distinct}, \alpha, \beta \in \{\pm 1\}\} \\ \cup \{P_{i, j} \mid 1 \leq i, j \leq n \text{ distinct}\} \cup \{I_i \mid 1 \leq i \leq n\}.$$

Observe that the set $S_{\text{Aut}}(n) \subset \text{Aut}(F_n)$ is closed under inversion.

Endomorphisms Below we will define a function $\theta: S_{\text{Aut}}(n) \rightarrow \text{End}(F(S_{\text{IA}}(n)))$ with this key property: Let $\pi: F(S_{\text{IA}}(n)) \rightarrow \text{IA}_n$ and $\rho: F(S_{\text{Aut}}(n)) \rightarrow \text{Aut}(F_n)$ be the projections. Then for $s \in S_{\text{Aut}}(n)$ and $w \in F(S_{\text{IA}}(n))$, we have

$$(3) \quad \pi(\theta(s)(w)) = \rho(s)\pi(w)\rho(s)^{-1} \in \text{IA}_n.$$

Our set of endomorphisms will then be

$$E_{\text{IA}}(n) = \{\theta(s) \mid s \in S_{\text{Aut}}(n)\}.$$

The relevance of the formula (3) is that we want the induced endomorphisms of our IA–presentation of IA_n to generate the image of $\text{Aut}(F_n)$ in $\text{Aut}(\text{IA}_n) \subset \text{End}(\text{IA}_n)$ arising from the conjugation action of $\text{Aut}(F_n)$ on IA_n .

Defining θ To define an endomorphism $\theta(s): F(S_{\text{IA}}(n)) \rightarrow F(S_{\text{IA}}(n))$ for $s \in S_{\text{Aut}}(n)$, it is enough to say what $\theta(s)$ does to each element of $S_{\text{IA}}(n)$. There are two cases:

$s \in S_I$	$\theta(M_{x_a^\alpha, x_b^\beta}^\beta)(s)$
C_{x_c, x_a}	$(C_{x_c, x_a}^\alpha C_{x_c, x_b}^\beta)^\alpha$
C_{x_a, x_c}	$C_{x_a, x_c} M_{x_a^\alpha, [x_b^{-\beta}, x_c]}^\beta$
C_{x_b, x_c}	$C_{x_b, x_c} M_{x_a^\alpha, [x_b^{-\beta}, x_c^{-1}]}^\beta$
C_{x_b, x_a}	$(C_{x_a, x_b}^\beta C_{x_b, x_a}^\alpha)^\alpha$
$M_{x_a^\alpha, [x_c^\gamma, x_d^\delta]}$	$C_{x_a, x_b}^\beta M_{x_a^\alpha, [x_c^\gamma, x_d^\delta]}^\beta C_{x_a, x_b}^{-\beta}$
$M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]}$	$M_{x_c^\gamma, [x_b^\beta, x_d^\delta]} C_{x_c, x_b}^{-\beta} M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]} C_{x_c, x_b}^\beta$
$M_{x_c^\gamma, [x_a^{-\alpha}, x_d^\delta]}$	$M_{x_c^\gamma, [x_a^{-\alpha}, x_d^\delta]} C_{x_c, x_a}^\alpha M_{x_c^\gamma, [x_b^{-\beta}, x_d^\delta]} C_{x_c, x_a}^{-\alpha}$
$M_{x_b^\beta, [x_c^\gamma, x_d^\delta]}$	$C_{x_a, x_b}^\beta M_{x_a^{-\alpha}, [x_c^\gamma, x_d^\delta]} M_{x_b^\beta, [x_c^\gamma, x_d^\delta]} C_{x_a, x_b}^{-\beta}$
$M_{x_b^{-\beta}, [x_c^\gamma, x_d^\delta]}$	$M_{x_a^\alpha, [x_c^\gamma, x_d^\delta]} M_{x_b^{-\beta}, [x_c^\gamma, x_d^\delta]}$
$M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}$	$C_{x_a, x_b}^\beta M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]} C_{x_a, x_b}^{-\beta}$
$M_{x_a^\alpha, [x_b^{-\beta}, x_c^\gamma]}$	$C_{x_a, x_b}^\beta M_{x_a^\alpha, [x_b^{-\beta}, x_c^\gamma]} C_{x_a, x_b}^{-\beta}$
$M_{x_b^\beta, [x_a^\alpha, x_c^\gamma]}$	$C_{x_a, x_c}^\gamma M_{x_a^\alpha, [x_b^{-\beta}, x_c^\gamma]} C_{x_b, x_c}^{-\gamma} M_{x_b^{-\beta}, [x_a^\alpha, x_c^{-\gamma}]}$
$M_{x_b^\beta, [x_a^{-\alpha}, x_c^\gamma]}$	$C_{x_c, x_b}^{-\beta} C_{x_c, x_a}^{-\alpha} M_{x_b^{-\beta}, [x_c^{-\gamma}, x_a^\alpha]} C_{x_b, x_c}^\beta M_{x_a^\alpha, [x_c^\gamma, x_b^{-\beta}]} C_{x_a, x_c}^{-\gamma} C_{x_c, x_a}^\alpha C_{x_c, x_a}^\beta$
$M_{x_b^{-\beta}, [x_a^\alpha, x_c^\gamma]}$	$C_{x_a, x_c}^{-\gamma} C_{x_c, x_a}^\alpha M_{x_b^{-\beta}, [x_c^\gamma, x_a^{-\alpha}]} C_{x_c, x_b}^\beta M_{x_a^\alpha, [x_c^\gamma, x_b^{-\beta}]} C_{x_b, x_c}^\gamma C_{x_c, x_b}^{-\beta} C_{x_c, x_a}^{-\alpha}$
$M_{x_b^{-\beta}, [x_a^{-\alpha}, x_c^\gamma]}$	$C_{x_b, x_c}^{-\gamma} M_{x_a^\alpha, [x_b^{-\beta}, x_c^\gamma]} C_{x_c, x_b}^{-\beta} M_{x_b^{-\beta}, [x_a^{-\alpha}, x_c^\gamma]} C_{x_c, x_a}^{-\alpha} C_{x_a, x_c}^\gamma C_{x_c, x_a}^\alpha C_{x_c, x_a}^\beta$
$M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]}$	$C_{x_a, x_b}^\beta M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]} C_{x_a, x_b}^{-\beta}$
$M_{x_c^\gamma, [x_a^\alpha, x_b^{-\beta}]}$	$M_{x_c^\gamma, [x_b^\beta, x_a^\alpha]}$

Table 3: Definition of $\theta(M_{x_a^\alpha, x_b^\beta}^\beta)$ on the generators $S_{IA}(n)$. All indices in each entry are assumed to be distinct. If no entry is listed for $t \in S_{IA}(n)$ or for the generator representing t^{-1} (as in relation (R0)) then $\theta(M_{x_a^\alpha, x_b^\beta}^\beta)(t) = t$.

- $s = P_{i,j}$ or $s = I_i$ We then define $\theta(s)$ using the action of s on F_n via $\theta(s)(C_{x_a, x_b}) = C_{s(x_a), s(x_b)}$ and $\theta(s)(M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}) = M_{s(x_a^\alpha), [s(x_b^\beta), s(x_c^\gamma)]}$. Here one should interpret $C_{x_e^{-1}, x_f}$ as C_{x_e, x_f} , $C_{x_e, x_f^{-1}}$ as C_{x_e, x_f}^{-1} and $C_{x_e^{-1}, x_f^{-1}}$ as C_{x_e, x_f}^{-1} .
- $s = M_{x_a^\alpha, x_b^\beta}$ In this case, we define $\theta(s)$ via the formulas in Table 3. These list the cases where $\theta(s)$ does not fix a generator, except that, to avoid redundancy, we do not always list both a commutator transvection and its inverse. Specifically, if $t = M_{x_c^\gamma, [x_d^\delta, x_e^\epsilon]}$, possibly with $\{c, d, e\} \cap \{a, b\} \neq \emptyset$, and a formula is listed for $t' = M_{x_c^\gamma, [x_e^\epsilon, x_d^\delta]}$ but not for t , then we define $\theta(s)(t) = \theta(s)(t')^{-1}$.

If Table 3 lists no entry for t or t' , or the table lists no entry for t and t is a conjugation move, then we define $\theta(s)(t) = t$.

These formulas were chosen to be as simple as possible, among formulas realizing (3). Just like for the relations, we recommend not dwelling on these formulas during one's first read through this paper.

Proposition 2.3 *The definition of θ satisfies (3).*

This proof uses a computer verification and is postponed until Section 5. Propositions 2.1 and 2.3 together imply that all of the extended relations from our L–presentation are trivial in IA_n . This means that the obvious map on generators (sending each generator to the automorphism it names) extends to a well-defined homomorphism

$$\langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle \rightarrow \text{IA}_n .$$

3 Tools for the proof

In this section, we assemble the tools we will need to prove Theorem E. In Section 3.1, we discuss a theorem of the second author that gives a sort of infinite presentation for a group acting on a simplicial complex. In Section 3.2, we introduce the complex of partial bases. In Section 3.3, we give generators for the IA_n –stabilizers of simplices in the complex of partial bases. In Section 3.4, we introduce an action of $\text{Aut}(F_n)$ on the group given by our purported L–presentation for IA_n . Finally, in Section 3.5 we introduce a certain morphism between groups given by L–presentations.

Two results in this sections have computer-aided proofs which are postponed until Section 5: Proposition 3.13 from Section 3.4 and Proposition 3.14 from Section 3.5.

3.1 Presentations from group actions

Consider a group G acting on a simplicial complex X . We say that G acts *without rotations* if for all simplices σ of X , the setwise and pointwise stabilizers of σ coincide. For a simplex σ , denote by G_σ the stabilizer of σ . Letting $X^{(0)}$ denote the vertex set of X , there is a homomorphism from the free product of vertex stabilizers

$$\psi: \bigast_{v \in X^{(0)}} G_v \rightarrow G .$$

As notation, if $g \in G$ stabilizes a vertex v of X , then denote by g_v the associated element of

$$G_v < \bigast_{v \in X^{(0)}} G_v .$$

The map ψ is rarely injective. Two families of elements in its kernel are as follows:

- If e is an edge of X joining vertices v and v' and if $g \in G_e$, then $g_v g_{v'}^{-1} \in \ker(\psi)$. We call these the *edge relators*.
- If $v, w \in X^{(0)}$ and $g \in G_v$ and $h \in G_w$, then $h_w g_v h_w^{-1} (h g h^{-1})_{h(v)}^{-1} \in \ker(\psi)$. We call these the *conjugation relators*.

The second author gave hypotheses under which these generate $\ker(\psi)$:

Theorem 3.1 [29] *Consider a group G acting without rotations on a 1–connected simplicial complex X . Assume that X/G is 2–connected. Then the kernel of the map ψ described above is normally generated by the edge and conjugation relators.*

3.2 The complex of partial bases

We now introduce the simplicial complex to which we will apply [Theorem 3.1](#). For $z \in F_n$, let $\llbracket z \rrbracket$ denote the union of the conjugacy classes of z and z^{-1} .

Definition 3.2 A *partial basis* for F_n is a set $\{z_1, \dots, z_k\} \subset F_n$ such that there exist $z_{k+1}, \dots, z_n \in F_n$ with $\{z_1, \dots, z_n\}$ a free basis for F_n . The *complex of partial bases* for F_n , denoted by \mathcal{B}_n , is the simplicial complex whose $(k-1)$ –simplices are sets $\{\llbracket z_1 \rrbracket, \dots, \llbracket z_k \rrbracket\}$, where $\{z_1, \dots, z_k\}$ is a partial basis for F_n .

The group $\text{Aut}(F_n)$ acts on \mathcal{B}_n , and we wish to apply [Theorem 3.1](#) to the restriction of this action to IA_n . It is clear that IA_n acts on \mathcal{B}_n without rotations, so we must check that \mathcal{B}_n is 1–connected and that $\mathcal{B}_n/\text{IA}_n$ is 2–connected.

We start by verifying that \mathcal{B}_n is 1–connected.

Proposition 3.3 *The simplicial complex \mathcal{B}_n is 1–connected for $n \geq 3$.*

Proof For $z \in F_n$, let $\llbracket z \rrbracket'$ be the conjugacy class of z . Define \mathcal{B}'_n to be the simplicial complex whose $(k-1)$ –simplices are sets $\{\llbracket z_1 \rrbracket', \dots, \llbracket z_k \rrbracket'\}$, where $\{z_1, \dots, z_k\}$ is a partial basis for F_n . In [16], the authors proved that \mathcal{B}'_n is 1–connected for $n \geq 3$. There is a natural simplicial map $\rho: \mathcal{B}'_n \rightarrow \mathcal{B}_n$. Letting $\psi^0: (\mathcal{B}_n)^{(0)} \rightarrow (\mathcal{B}'_n)^{(0)}$ be an arbitrary map satisfying $\rho \circ \psi^0 = \text{id}$, it is clear that ψ^0 extends to a simplicial map $\psi: \mathcal{B}_n \rightarrow \mathcal{B}'_n$ satisfying $\rho \circ \psi = \text{id}$. This implies that ρ induces a surjection on all homotopy groups, so \mathcal{B}_n is 1–connected for $n \geq 3$. □

It also follows from [16] that $\mathcal{B}_n/\text{IA}_n$ is $(n-2)$ –connected. In particular, it is 2–connected for $n \geq 4$, and thus satisfies the conditions of [Theorem 3.1](#) for $n \geq 4$. However, we will need a complex that satisfies the conditions of [Theorem 3.1](#) for $n = 3$ as well. We therefore attach cells to increase the connectivity.

Definition 3.4 The *augmented complex of partial bases* for F_n , denoted by $\widehat{\mathcal{B}}_n$, is the simplicial complex whose $(k-1)$ -simplices are as follows:

- Sets of the form $\{\llbracket z_1 \rrbracket, \dots, \llbracket z_k \rrbracket\}$, where $\{z_1, \dots, z_k\}$ is a partial basis for F_n . These will be called the *standard simplices*.
- Sets of the form $\{\llbracket z_1 z_2 \rrbracket, \llbracket z_1 \rrbracket, \llbracket z_2 \rrbracket, \dots, \llbracket z_{k-1} \rrbracket\}$, where $\{z_1, \dots, z_{k-1}\}$ is a partial basis for F_n . These will be called the *additive simplices*.

Remark 3.5 Since $z_1 z_2$ and $z_2 z_1$ are conjugate, the two additive simplices

$$\{\llbracket z_1 z_2 \rrbracket, \llbracket z_1 \rrbracket, \llbracket z_2 \rrbracket, \dots, \llbracket z_{k-1} \rrbracket\} \quad \text{and} \quad \{\llbracket z_2 z_1 \rrbracket, \llbracket z_1 \rrbracket, \llbracket z_2 \rrbracket, \dots, \llbracket z_{k-1} \rrbracket\}$$

of $\widehat{\mathcal{B}}_n$ are the same.

The group $\text{Aut}(F_n)$ (and hence IA_n) still acts on $\widehat{\mathcal{B}}_n$. Since $\widehat{\mathcal{B}}_n$ is obtained from \mathcal{B}_n by adding simplices of dimension at least 2, it inherits the 1-connectivity of \mathcal{B}_n for $n \geq 3$ asserted in Proposition 3.3.

Proposition 3.6 The complex $\widehat{\mathcal{B}}_n$ is 1-connected for $n \geq 3$.

To help us understand the connectivity of $\widehat{\mathcal{B}}_n / \text{IA}_n$, we introduce the following complex. For $\vec{v} \in \mathbb{Z}^n$, let $(\vec{v})_{\pm}$ denote the set $\{\vec{v}, -\vec{v}\}$.

Definition 3.7 A *partial basis* for \mathbb{Z}^n is a set $\{\vec{v}_1, \dots, \vec{v}_k\} \subset \mathbb{Z}^n$ such that there exist $\vec{v}_{k+1}, \dots, \vec{v}_n \in \mathbb{Z}^n$ with $\{\vec{v}_1, \dots, \vec{v}_n\}$ a basis for \mathbb{Z}^n . The *augmented complex of lax partial bases* for \mathbb{Z}^n , denoted by $\widehat{\mathcal{B}}_n(\mathbb{Z})$, is the simplicial complex whose $(k-1)$ -simplices are as follows:

- Sets of the form $\{(\vec{v}_1)_{\pm}, \dots, (\vec{v}_k)_{\pm}\}$, where $\{\vec{v}_1, \dots, \vec{v}_k\}$ is a partial basis for \mathbb{Z}^n . These will be called the *standard simplices*.
- Sets of the form $\{(\vec{v}_1 + \vec{v}_2)_{\pm}, (\vec{v}_1)_{\pm}, (\vec{v}_2)_{\pm}, \dots, (\vec{v}_{k-1})_{\pm}\}$, where $\{\vec{v}_1, \dots, \vec{v}_{k-1}\}$ is a partial basis for \mathbb{Z}^n . These will be called the *additive simplices*.

We then have the following lemma:

Lemma 3.8 We have $\widehat{\mathcal{B}}_n / \text{IA}_n \cong \widehat{\mathcal{B}}_n(\mathbb{Z})$ for $n \geq 1$.

For the proof of Lemma 3.8, we will need the following result of the authors. For $z \in F_n$, let $[z] \in \mathbb{Z}^n$ be the associated element of the abelianization of F_n .

Lemma 3.9 [16, Lemma 5.3] Let $\{\vec{v}_1, \dots, \vec{v}_n\}$ be a basis for \mathbb{Z}^n and let $\{z_1, \dots, z_k\}$ be a partial basis for F_n such that $[z_i] = \vec{v}_i$ for $1 \leq i \leq k$. Then there exist $z_{k+1}, \dots, z_n \in F_n$ with $[z_i] = \vec{v}_i$ for $k + 1 \leq i \leq n$ such that $\{z_1, \dots, z_n\}$ is a basis for F_n .

Proof of Lemma 3.8 The map $(\widehat{\mathcal{B}}_n)^{(0)} \rightarrow (\widehat{\mathcal{B}}_n(\mathbb{Z}))^{(0)}$ that takes $\llbracket z \rrbracket$ to $[z]$ extends to a simplicial map $\rho: \widehat{\mathcal{B}}_n \rightarrow \widehat{\mathcal{B}}_n(\mathbb{Z})$. Since IA_n acts without rotations on $\widehat{\mathcal{B}}_n$, the quotient $\widehat{\mathcal{B}}_n/\text{IA}_n$ has a natural CW-complex structure whose k -cells are the IA_n -orbits of the k -cells of $\widehat{\mathcal{B}}_n$ (warning: though it will turn out that in this case it is, this CW-complex structure need not be a simplicial complex structure; consider, for example, the action of \mathbb{Z} by translations on the standard triangulation of \mathbb{R} whose vertices are \mathbb{Z}). Since ρ is IA_n -invariant, it factors through a map $\bar{\rho}: \widehat{\mathcal{B}}_n/\text{IA}_n \rightarrow \widehat{\mathcal{B}}_n(\mathbb{Z})$. We will prove that $\bar{\rho}$ is an isomorphism of CW-complexes.

This requires checking two things. The first is that every simplex of $\widehat{\mathcal{B}}_n(\mathbb{Z})$ is in the image of ρ , which is an immediate consequence of Lemma 3.9. The second is that if σ and σ' are simplices of $\widehat{\mathcal{B}}_n$ such that $\rho(\sigma) = \rho(\sigma')$, then there exists some $f \in \text{IA}_n$ such that $f(\sigma) = \sigma'$. It is clear that σ and σ' are either both standard simplices or both additive simplices. Assume first that they are both standard simplices. We can then write

$$\sigma = \{\llbracket z_1 \rrbracket, \dots, \llbracket z_k \rrbracket\} \quad \text{and} \quad \sigma' = \{\llbracket z'_1 \rrbracket, \dots, \llbracket z'_k \rrbracket\}$$

as in the definition of standard simplices, with $[z_i] = [z'_i]$ for $1 \leq i \leq k$. Set $\vec{v}_i = [z_i] = [z'_i]$ for $1 \leq i \leq k$. The set $\{\vec{v}_1, \dots, \vec{v}_k\}$ is a partial basis for \mathbb{Z}^n , so we can extend it to a basis $\{\vec{v}_1, \dots, \vec{v}_n\}$. Applying Lemma 3.9 twice, we can find $z_{k+1}, \dots, z_n \in F_n$ and $z'_{k+1}, \dots, z'_n \in F_n$ such that $[z_i] = [z'_i] = \vec{v}_i$ for $k+1 \leq i \leq n$ and such that both $\{z_1, \dots, z_n\}$ and $\{z'_1, \dots, z'_n\}$ are free bases for F_n . There then exists $f \in \text{Aut}(F_n)$ such that $f(z_i) = z'_i$ for $1 \leq i \leq n$. By construction, we have $f \in \text{IA}_n$ and $f(\sigma) = \sigma'$.

It remains to deal with the case where σ and σ' are both simplices of additive type. Write

$$\sigma = \{\llbracket z_1 z_2 \rrbracket, \llbracket z_1 \rrbracket, \llbracket z_2 \rrbracket, \dots, \llbracket z_{k-1} \rrbracket\} \quad \text{and} \quad \sigma' = \{\llbracket z'_1 z'_2 \rrbracket, \llbracket z'_1 \rrbracket, \dots, \llbracket z'_{k-1} \rrbracket\}$$

as in the definition of additive simplices. The unordered sets

$$\{([z_1] + [z_2])_{\pm}, ([z_1])_{\pm}, ([z_2])_{\pm}\} \quad \text{and} \quad \{([z'_1] + [z'_2])_{\pm}, ([z'_1])_{\pm}, ([z'_2])_{\pm}\}$$

are minimal nonempty subsets of $\rho(\sigma) = \rho(\sigma')$ such that the defining elements of \mathbb{Z}^n are not linearly independent. It follows that as unordered sets we have

$$\begin{aligned} \rho(\{\llbracket z_1 z_2 \rrbracket, \llbracket z_1 \rrbracket, \llbracket z_2 \rrbracket\}) &= \rho(\{\llbracket z'_1 z'_2 \rrbracket, \llbracket z'_1 \rrbracket, \llbracket z'_2 \rrbracket\}), \\ \rho(\{\llbracket z_3 \rrbracket, \dots, \llbracket z_{k-1} \rrbracket\}) &= \rho(\{\llbracket z'_3 \rrbracket, \dots, \llbracket z'_{k-1} \rrbracket\}). \end{aligned}$$

Reordering the z_i and possibly replacing some of the z_i by z_i^{-1} (which does not change $\llbracket z_i \rrbracket$), we can assume that $[z_i] = [z'_i]$ for $3 \leq i \leq k-1$.

The next observation is that all of the following sets define the same additive simplex (but with the vertices in a different order; all six possible orderings occur):

$$\begin{aligned} & \{[[z_1 z_2]], [[z_1]], [[z_2]]\}, \quad \{[[z_2 z_1]], [[z_2]], [[z_1]]\}, \quad \{[[z_1]], [[z_1 z_2]], [[z_2^{-1}]]\}, \\ & \{[[z_1^{-1}]], [[z_2]], [[z_2^{-1} z_1^{-1}]]\}, \quad \{[[z_2]], [[z_2 z_1]], [[z_1^{-1}]]\}, \quad \{[[z_2^{-1}]], [[z_1]], [[z_1^{-1} z_2^{-1}]]\}. \end{aligned}$$

By reordering σ and possibly changing some of our expressions for the elements in it again, we can assume that

$$([z_1] + [z_2])_{\pm} = ([z'_1] + [z'_2])_{\pm}, \quad ([z_1])_{\pm} = ([z'_1])_{\pm}, \quad ([z_2])_{\pm} = ([z'_2])_{\pm}$$

and that $[z_i] = [z'_i]$ for $3 \leq i \leq k - 1$.

The final observation is that either

$$([z_1], [z_2]) = ([z'_1], [z'_2]) \quad \text{or} \quad ([z_1], [z_2]) = (-[z'_1], -[z'_2]);$$

the key point here is that changing the sign of one of $\{[z_1], [z_2]\}$ but not the other changes $([z_1] + [z_2])_{\pm}$. If the second possibility occurs, then replace z_1 and z_2 with z_1^{-1} and z_2^{-1} , respectively; this does not change σ . The upshot is that we now have arranged for $[z_i] = [z'_i]$ for all $1 \leq i \leq k - 1$. By the same argument we used to deal with standard simplices, there exists some $f \in \text{IA}_n$ such that $f(z_i) = z'_i$ for $1 \leq i \leq k - 1$. Since $f(z_1 z_2) = z'_1 z'_2$, we see that $f(\sigma) = \sigma'$, as desired. \square

The second author together with Church proved in [13] that $\widehat{\mathcal{B}}_n(\mathbb{Z})$ is $(n-1)$ -connected for $n \geq 1$. We therefore deduce the following:

Proposition 3.10 *The complex $\widehat{\mathcal{B}}_n / \text{IA}_n$ is $(n-1)$ -connected for $n \geq 1$.*

3.3 Generators for simplex stabilizers

This section is devoted to the following proposition, which gives generators for the stabilizers in IA_n of simplices of \mathcal{B}_n . Recall that $S_{\text{MA}}(n)$ is Magnus’s generating set for IA_n discussed in the introduction.

Proposition 3.11 *Fix $1 \leq k \leq n$ and define $\Gamma = (\text{IA}_n)_{[[x_{n-k+1}], [[x_{n-k+2}], \dots, [[x_n]]}$. Then Γ is generated by*

$$\begin{aligned} S_{\text{MA}}(n) \cap \Gamma &= \{C_{x_a, x_b} \mid 1 \leq a, b \leq n \text{ distinct}\} \\ &\cup \{M_{x_a, [x_b, x_c]} \mid 1 \leq a \leq n - k, 1 \leq b, c \leq n \text{ distinct}\}. \end{aligned}$$

Proof The map $F_n \rightarrow F_{n-k}$ that quotients by the normal closure of $\{x_{n-k+1}, \dots, x_n\}$ induces a split surjection $\rho: \Gamma \rightarrow \text{IA}_{n-k}$. Define $\mathcal{K}_{n-k, k} = \ker(\rho)$, so we have

$\Gamma = \mathcal{K}_{n-k,k} \rtimes \text{IA}_{n-k}$. As we said in the introduction, Magnus [26] proved that IA_{n-k} is generated by

$$(4) \{C_{x_a, x_b} \mid 1 \leq a, b \leq n-k \text{ distinct}\} \cup \{M_{x_a, [x_b, x_c]} \mid 1 \leq a, b, c \leq n-k \text{ distinct}\}.$$

The authors proved in [17, Theorem A] that $\mathcal{K}_{n-k,k}$ is generated by

$$(5) \{C_{x_a, x_b} \mid n-k+1 \leq a \leq n, 1 \leq b \leq n \text{ distinct}\} \\ \cup \{C_{x_a, x_b} \mid 1 \leq a \leq n, n-k+1 \leq b \leq n \text{ distinct}\} \\ \cup \{M_{x_a, [x_b, x_c]} \mid 1 \leq a \leq n-k, n-k-1 \leq b \leq n, 1 \leq c \leq n \text{ distinct}\}.$$

The union of (4) and (5) is the claimed generating set for Γ . □

Remark 3.12 For $z \in F_n$, define $[[z]]'$ to be the conjugacy class of z . In [17] we deal with $(\text{IA}_n)_{[[x_{n-k+1}]]', [[x_{n-k+2}]]', \dots, [[x_n]]'$ instead of $(\text{IA}_n)_{[[x_{n-k+1}]], [[x_{n-k+2}]], \dots, [[x_n]]}$; however, since x_i and x_i^{-1} have different images in F_n^{ab} , these two stabilizer subgroups are actually equal. There are also notational differences: the group denoted by $\mathcal{K}_{n-k,k}$ here is denoted by $\mathcal{K}_{n-k,k}^{\text{IA}}$ in that paper.

3.4 The action of $\text{Aut}(F_n)$

Let \mathcal{Q}_n be the group with the L–presentation $\langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle$ discussed in Section 2. By Propositions 2.1 and 2.3, there is a map $\pi: \mathcal{Q}_n \rightarrow \text{IA}_n$. The group $\text{Aut}(F_n)$ acts on IA_n by conjugation. The goal of this section is to state Proposition 3.13 below, which asserts that this action can be lifted to \mathcal{Q}_n .

To state some important properties of this lifted action, we must introduce some notation. First, let $S_{\text{Aut}}(n) \subset \text{Aut}(F_n)$ be the generating set discussed in Section 2. Recall that $E_{\text{IA}}(n) \subset \text{End}(F(S_{\text{IA}}(n)))$ is the image of a map $\theta: S_{\text{Aut}}(n) \rightarrow \text{End}(F(S_{\text{IA}}(n)))$. There is thus a map $e: S_{\text{Aut}}(n) \rightarrow \text{End}(\mathcal{Q}_n)$ whose image is the set of induced endomorphisms of our L–presentation. It will turn out that the image of e consists of automorphisms, and these automorphisms generate the action of $\text{Aut}(F_n)$ on \mathcal{Q}_n .

Second, recall that $\{x_1, \dots, x_n\}$ is a fixed free basis for F_n . Let

$$(S_{\text{IA}}(n))_{[[x_n]]} = \{C_{x_a, x_b} \mid 1 \leq a, b \leq n \text{ distinct}\} \\ \cup \{M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]} \mid 1 \leq a, b, c \leq n \text{ distinct}, \alpha, \beta, \gamma \in \{\pm 1\}, a \neq n\}.$$

This is exactly the subset of $S_{\text{IA}}(n) \subset \text{IA}_n$ consisting of automorphisms that fix $[[x_n]]$; Proposition 3.11 (with $k = 1$) implies that it generates the stabilizer subgroup $(\text{IA}_n)_{[[x_n]]}$. Define $(\mathcal{Q}_n)_{[[x_n]]}$ be the subgroup of \mathcal{Q}_n generated by $(S_{\text{IA}}(n))_{[[x_n]]}$. We will then require the stabilizer subgroup $(\text{Aut}(F_n))_{[[x]]}$ to preserve the subgroup $(\mathcal{Q}_n)_{[[x_n]]}$.

Our proposition is as follows:

Proposition 3.13 *For all $n \geq 2$, there is an action of $\text{Aut}(F_n)$ on \mathcal{Q}_n that satisfies the following three properties:*

- (1) *The action comes from the induced endomorphisms in the sense that, for $s \in S_{\text{IA}}(n) \subset \text{Aut}(F_n)$ and $q \in \mathcal{Q}_n$, we have $s \cdot q = e(s) \cdot q$.*
- (2) *The restriction of the action to IA_n induces the conjugation action of \mathcal{Q}_n on itself in the sense that, for $q, r \in \mathcal{Q}_n$, we have $\pi(r) \cdot q = rqr^{-1}$.*
- (3) *For $\eta \in (\text{Aut}(F_n))_{[[x_n]]}$ and $q \in (\mathcal{Q}_n)_{[[x_n]]}$, we have $\eta \cdot q \in (\mathcal{Q}_n)_{[[x_n]]}$.*

The proof of [Proposition 3.13](#) is a computation with generators and relations (mostly done by computer), so we have postponed it until [Section 5](#).

3.5 A homomorphism between L-presentations

There is a natural split surjection $\rho: (\text{IA}_n)_{[[x_n]]} \rightarrow \text{IA}_{n-1}$ arising from the quotient map $F_n \rightarrow F_{n-1}$ whose kernel is the normal closure of x_n . Let $\mathcal{K}_{n-1,1} = \ker(\rho)$; so we have a decomposition $(\text{IA}_n)_{[[x_n]]} = \mathcal{K}_{n-1,1} \rtimes \text{IA}_{n-1}$. Building on the Birman exact sequence for $\text{Aut}(F_n)$ we constructed in [\[15\]](#), we constructed an L-presentation for $\mathcal{K}_{n-1,1}$ in [\[17, Theorem D\]](#) ($\mathcal{K}_{n-1,1}$ is denoted by $\mathcal{K}_{n-1,1}^{\text{IA}}$ in that paper). This L-presentation plays a crucial role in the inductive step of our proof, because it allows us to obtain the following proposition:

Proposition 3.14 *There is a homomorphism $\mathcal{K}_{n-1,1} \rightarrow \langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle$ fitting into the following commuting triangle:*

$$\begin{array}{ccc}
 \mathcal{K}_{n-1,1} & \longrightarrow & \langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle \\
 & \searrow & \downarrow \\
 & & \text{IA}_n
 \end{array}$$

Usually finding a homomorphism between groups given by presentations is simple: one checks that the relations map to products of conjugates of relations. This is the spirit of the proof of [Proposition 3.14](#), but the substitution rules and extended relations complicate the picture. Our proof of [Proposition 3.14](#) is computer-assisted and is postponed until [Section 5](#).

4 Verification of our L-presentation

In this section, we prove [Theorem E](#), which says that IA_n has the finite L-presentation $\langle S_{\text{IA}}(n) \mid R_{\text{IA}}^0(n) \mid E_{\text{IA}}(n) \rangle$ discussed in [Section 2](#). Our proof is inspired by the proof of the main theorem of [\[9\]](#). We will make use of [Propositions 2.1, 2.3, 3.13 and 3.14](#),

which were all stated in previous sections and which will be proved (with the aid of a computer) in Section 5.

Proof of Theorem E Let \mathcal{Q}_n be the group given by $\langle S_{IA}(n) \mid R_{IA}^0(n) \mid E_{IA}(n) \rangle$. Elements of $S_{IA}(n)$ play dual roles as elements of \mathcal{Q}_n and as elements of IA_n , and during our proof it will be important to distinguish them. Therefore, throughout this proof elements C_{x_a, x_b} and $M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}$ will always lie in IA_n ; the associated elements of \mathcal{Q}_n will be denoted by \mathfrak{C}_{x_a, x_b} and $\mathfrak{M}_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}$.

There is a natural projection map $\pi: \mathcal{Q}_n \rightarrow IA_n$. We will prove that π is an isomorphism by induction on n . The base cases are $n = 1$ and $n = 2$. For $n = 1$, both IA_n and \mathcal{Q}_n are the trivial group, so there is nothing to prove. For $n = 2$, it is a classical theorem of Nielsen [27] (see also [25, Proposition 4.5]) that IA_2 is the group of inner automorphisms of F_2 , so IA_2 is a free group on the generators C_{x_1, x_2} and C_{x_2, x_1} . Our generating set for \mathcal{Q}_2 is $\{\mathfrak{C}_{x_1, x_2}, \mathfrak{C}_{x_2, x_1}\}$, and for $n = 2$ the set of basic relations $R_{IA}^0(2)$ is empty. Even though our set of substitution rules $E_{IA}(2)$ is nonempty, it follows that our full set of relations for \mathcal{Q}_2 is empty. So our presentation for \mathcal{Q}_2 is $\langle \mathfrak{C}_{x_1, x_2}, \mathfrak{C}_{x_2, x_1} \mid \emptyset \rangle$, and the result is also true in this case.

Assume now that $n \geq 3$ and that the projection map $\mathcal{Q}_{n'} \rightarrow IA_{n'}$ is an isomorphism for all $1 \leq n' < n$. Since π is a surjection, to prove that π is an isomorphism it is enough to construct a homomorphism $\phi: IA_n \rightarrow \mathcal{Q}_n$ such that $\phi \circ \pi = \text{id}$. Propositions 3.6 and 3.10 show that the action of IA_n on \hat{B}_n satisfies the conditions of Theorem 3.1, so

$$IA_n \cong \left(\prod_{[[z]] \in (\hat{B}_n)^{(0)}} * (IA_n)_{[[z]]} \right) / R,$$

where R is the normal closure of the edge and conjugation relators. The construction of ϕ will have two steps. First, we will use the action of $\text{Aut}(F_n)$ on \mathcal{Q}_n provided by Proposition 3.13 to construct a map

$$\tilde{\phi}: \prod_{[[z]] \in (\hat{B}_n)^{(0)}} * (IA_n)_{[[z]]} \rightarrow \mathcal{Q}_n.$$

Second, we will show that $\tilde{\phi}$ takes the edge and conjugation relators to 1, and thus induces a map $\phi: IA_n \rightarrow \mathcal{Q}_n$. We will close by verifying that $\phi \circ \pi = \text{id}$.

Construction of $\tilde{\phi}$ To construct $\tilde{\phi}$, we must construct a map

$$\tilde{\phi}_{[[z]]}: (IA_n)_{[[z]]} \rightarrow \mathcal{Q}_n$$

for each vertex $[[z]]$ of \hat{B}_n . Recalling that $\{x_1, \dots, x_n\}$ is our fixed free basis for F_n , we begin with the vertex $[[x_n]]$. In the following claim, we will use the notation $(S_{IA}(n))_{[[x_n]]}$ and $(\mathcal{Q}_n)_{[[x_n]]}$ introduced in Section 3.4.

Claim 1 *The restriction of π to $(\mathcal{Q}_n)_{[[x_n]]}$ is an isomorphism onto $(IA_n)_{[[x_n]]}$.*

Proof Proposition 3.11 implies that natural map $\pi|_{(\mathcal{Q}_n)_{[[x_n]]}}: (\mathcal{Q}_n)_{[[x_n]]} \rightarrow (\text{IA}_n)_{[[x_n]]}$ is surjective, since the generators from that proposition (with $k = 1$) are in the image.

Our inductive hypothesis says that the map $\pi|_{\mathcal{Q}_{n-1}}: \mathcal{Q}_{n-1} \rightarrow \text{IA}_{n-1}$ is an isomorphism. Recall from Section 3.5 that $(\text{IA}_n)_{[[x_n]]} = \mathcal{K}_{n-1,1} \rtimes \text{IA}_{n-1}$, where the projection $(\text{IA}_n)_{[[x_n]]} \rightarrow \text{IA}_{n-1}$ is the one induced by the map $F_n \rightarrow F_{n-1}$ that quotients out by the normal closure of x_n , and $\mathcal{K}_{n-1,1}$ is the kernel of this projection. The composition

$$(\mathcal{Q}_n)_{[[x_n]]} \rightarrow (\text{IA}_n)_{[[x_n]]} \rightarrow \text{IA}_{n-1} \xrightarrow[\cong]{\pi|_{\mathcal{Q}_{n-1}}^{-1}} \mathcal{Q}_{n-1}$$

is a well-defined homomorphism. It is a composition of surjective maps, and is therefore surjective. We define $\mathfrak{K}_{n-1,1}$ to be the kernel of this composition of maps.

The restriction of π to $\mathfrak{K}_{n-1,1}$ has image in $\mathcal{K}_{n-1,1}$ since the map $(\mathcal{Q}_n)_{[[x_n]]} \rightarrow \mathcal{Q}_{n-1}$ factors through $(\text{IA}_n)_{[[x_n]]} \rightarrow \text{IA}_{n-1}$. Proposition 3.11 says that $\mathcal{K}_{n-1,1}$ is generated by the set

$$S_{\mathcal{K}}(n) := \{C_{x_n, x_a}, C_{x_a, x_n} \mid 1 \leq a < n\} \\ \cup \{M_{x_a^\alpha, [x_b^\beta, x_n^\gamma]}, M_{x_a^\alpha, [x_n^\gamma, x_b^\beta]} \mid 1 \leq a, b < n \text{ distinct, } \alpha, \beta, \gamma \in \{\pm 1\}\}.$$

Since these generators are contained in $\mathfrak{K}_{n-1,1}$, the map $\mathfrak{K}_{n-1,1} \rightarrow \mathcal{K}_{n-1,1}$ is surjective. Further, Proposition 3.14 gives us a left inverse to $\pi|_{\mathfrak{K}_{n-1,1}}$. We conclude that $\pi|_{\mathfrak{K}_{n-1,1}}$ is an isomorphism $\mathfrak{K}_{n-1,1} \cong \mathcal{K}_{n-1,1}$. We note that existence of this isomorphism is a deceptively difficult part of the proof, and it is the main consequence that we draw from [17].

Summing up, we have a commutative diagram of short exact sequences as follows:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathfrak{K}_{n-1,1} & \longrightarrow & (\mathcal{Q}_n)_{[[x_n]]} & \longrightarrow & \mathcal{Q}_{n-1} \longrightarrow 1 \\ & & \downarrow \cong & & \downarrow & & \downarrow \cong \\ 1 & \longrightarrow & \mathcal{K}_{n-1,1} & \longrightarrow & (\text{IA}_n)_{[[x_n]]} & \longrightarrow & \text{IA}_{n-1} \longrightarrow 1 \end{array}$$

The five lemma therefore says that the projection map $(\mathcal{Q}_n)_{[[x_n]]} \rightarrow (\text{IA}_n)_{[[x_n]]}$ is an isomorphism, as desired. \square

Claim 1 implies that we can define a map $\tilde{\phi}_{[[x_n]]}: (\text{IA}_n)_{[[x_n]]} \rightarrow \mathcal{Q}_n$ via the formula $\tilde{\phi}_{[[x_n]]} = (\pi|_{(\mathcal{Q}_n)_{[[x_n]]}})^{-1}$.

Now consider a general vertex $[[z]]$ of $\hat{\mathcal{B}}_n$. Here we will use the action of $\text{Aut}(F_n)$ on \mathcal{Q}_n provided by Proposition 3.13. The group $\text{Aut}(F_n)$ acts transitively on the set of primitive elements of F_n , so there exists some $\nu \in \text{Aut}(F_n)$ such that $\nu(x_n) = z$. We

then define a map $\tilde{\phi}_{[[z]]}: (\mathbf{IA}_n)_{[[z]]} \rightarrow \mathcal{Q}_n$ via the formula

$$\tilde{\phi}_{[[z]]}(\eta) = \nu \cdot \tilde{\phi}_{[[x_n]]}(\nu^{-1}\eta\nu) \quad (\eta \in (\mathbf{IA}_n)_{[[z]])$$

This appears to depend on the choice of ν , but the following claim says that this choice does not matter.

Claim 2 *The map $\tilde{\phi}_{[[z]]}(\eta)$ does not depend on the choice of ν .*

Proof Assume that $\nu_1, \nu_2 \in \text{Aut}(F_n)$ both satisfy $\nu_i(x_n) = z$, and consider some $\eta \in (\mathbf{IA}_n)_{[[z]]}$. Our goal is to prove that

$$(6) \quad \nu_1 \cdot \tilde{\phi}_{[[x_n]]}(\nu_1^{-1}\eta\nu_1) = \nu_2 \cdot \tilde{\phi}_{[[x_n]]}(\nu_2^{-1}\eta\nu_2).$$

Define $\mu = \nu_1^{-1}\nu_2$ and $\omega = \nu_2^{-1}\eta\nu_2$, so $\mu \in (\text{Aut}(F_n))_{[[x_n]]}$ and $\omega \in (\mathbf{IA}_n)_{[[x_n]]}$. We will first prove that

$$(7) \quad \tilde{\phi}_{[[x_n]]}(\mu\omega\mu^{-1}) = \mu \cdot \tilde{\phi}_{[[x_n]]}(\omega).$$

To see this, observe first that by construction both $\tilde{\phi}_{[[x_n]]}(\mu\omega\mu^{-1})$ and $\tilde{\phi}_{[[x_n]]}(\omega)$ lie in $(\mathcal{Q}_n)_{[[x_n]]}$. The third part of [Proposition 3.13](#) implies that $\mu \cdot \tilde{\phi}_{[[x_n]]}(\omega)$ also lies in $(\mathcal{Q}_n)_{[[x_n]]}$. [Claim 1](#) says that $\pi|_{(\mathcal{Q}_n)_{[[x_n]]}}$ is injective, so to prove (7), it is thus enough to prove that $\tilde{\phi}_{[[x_n]]}(\mu\omega\mu^{-1})$ and $\mu \cdot \tilde{\phi}_{[[x_n]]}(\omega)$ have the same image under π . This follows from the calculation

$$\pi(\tilde{\phi}_{[[x_n]]}(\mu\omega\mu^{-1})) = \mu\omega\mu^{-1} = \mu\pi(\tilde{\phi}_{[[x_n]]}(\omega))\mu^{-1} = \pi(\mu \cdot \tilde{\phi}_{[[x_n]]}(\omega)),$$

where the first two equalities follow from the fact that $\pi \circ \tilde{\phi}_{[[x_n]]} = \text{id}$ and the third follows from the first conclusion of [Proposition 3.13](#).

We now verify (6) as follows:

$$\nu_1 \cdot \tilde{\phi}_{[[x_n]]}(\nu_1^{-1}\eta\nu_1) = \nu_1 \cdot \tilde{\phi}_{[[x_n]]}(\mu\omega\mu^{-1}) = \nu_1\mu \cdot \tilde{\phi}_{[[x_n]]}(\omega) = \nu_2 \cdot \tilde{\phi}_{[[x_n]]}(\nu_2^{-1}\eta\nu_2). \quad \square$$

This completes the construction of $\tilde{\phi}$.

Some naturality properties Before we study the edge and conjugation relators, we first need to verify the following two naturality properties of $\tilde{\phi}$. Starting now we will use the notation which was introduced in [Section 2](#): for a vertex $[[z]]$ of $\hat{\mathcal{B}}_n$ and $\eta \in \mathbf{IA}_n$ satisfying $\eta([[z]]) = [[z]]$, we will denote η , considered as an element of

$$(\mathbf{IA}_n)_{[[z]]} < \underset{[[z]] \in (\hat{\mathcal{B}}_n)^{(0)}}{*} (\mathbf{IA}_n)_{[[z]]},$$

by $\eta_{[[z]]}$.

Claim 3 *The following two identities hold:*

- Let $1 \leq a, b \leq n$ be distinct and $1 \leq i \leq n$ be arbitrary. Then $\tilde{\phi}((C_{x_a, x_b})_{\llbracket x_i \rrbracket}) = \mathfrak{C}_{x_a, x_b}$.
- Let $1 \leq a, b, c \leq n$ be distinct, let $\alpha, \beta, \gamma \in \{\pm 1\}$ be arbitrary, and let $1 \leq i \leq n$ be such that $i \neq a$. Then $\tilde{\phi}((M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]})_{\llbracket x_i \rrbracket}) = \mathfrak{M}_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}$.

Proof The proofs of the two identities are similar; we will deal with the first and leave the second to the reader. It is clear from the construction that $\tilde{\phi}((C_{x_a, x_b})_{\llbracket x_n \rrbracket}) = \mathfrak{C}_{x_a, x_b}$. For $1 \leq i < n$, we have $P_{i,n}(x_n) = x_i$, and thus by definition we have

$$\begin{aligned} \tilde{\phi}((C_{x_a, x_b})_{\llbracket x_i \rrbracket}) &= P_{i,n} \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(P_{i,n}^{-1} C_{x_a, x_b} P_{i,n}) = P_{i,n} \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(C_{P_{i,n}^{-1}(x_a), P_{i,n}^{-1}(x_b)}) \\ &= P_{i,n} \cdot \mathfrak{C}_{P_{i,n}^{-1}(x_a), P_{i,n}^{-1}(x_b)} = \mathfrak{C}_{x_a, x_b}; \end{aligned}$$

here the last equality follows from the first part of [Proposition 3.13](#) and the definition of the endomorphisms in [Section 2](#). □

Claim 4 Let $\llbracket z \rrbracket$ be a vertex of $\hat{\mathcal{B}}_n$. Then for $\eta \in (\text{IA}_n)_{\llbracket z \rrbracket}$ we have $\pi(\tilde{\phi}_{\llbracket z \rrbracket}(\eta)) = \eta$.

Proof Pick $v \in \text{Aut}(F_n)$ such that $v(x_n) = z$. Then

$$\pi(\tilde{\phi}_{\llbracket z \rrbracket}(\eta)) = \pi(v \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(v^{-1}\eta v)) = v\pi(\tilde{\phi}_{\llbracket x_n \rrbracket}(v^{-1}\eta v))v^{-1} = v v^{-1} \eta v v^{-1} = \eta;$$

here the second equality uses the first part of [Proposition 3.13](#). □

The edge and conjugation relators We now check that $\tilde{\phi}$ takes the edge and conjugation relators to 1.

Claim 5 (edge relators) If e is an edge of $\hat{\mathcal{B}}_n$ with endpoints $\llbracket z \rrbracket$ and $\llbracket z' \rrbracket$ and $\eta \in (\text{IA}_n)_e$, then $\tilde{\phi}(\eta_{\llbracket z \rrbracket} \eta_{\llbracket z' \rrbracket}^{-1}) = 1$.

Proof We first consider the special case where $z = x_n$ and $z' = x_{n-1}$. [Proposition 3.11](#), with $k = 2$, states that $(\text{IA}_n)_{\llbracket x_{n-1} \rrbracket, \llbracket x_n \rrbracket}$ is generated by

$$(8) \quad \{C_{x_a, x_b} \mid 1 \leq a, b \leq n \text{ distinct}\} \cup \{M_{x_a, [x_b, x_c]} \mid 1 \leq a, b, c \leq n \text{ distinct, } a \neq n-1, n\}.$$

[Claim 3](#) implies that for all elements ω in (8), we have $\tilde{\phi}(\omega_{\llbracket x_{n-1} \rrbracket}) = \tilde{\phi}(\omega_{\llbracket x_n \rrbracket})$. It follows that for all $\eta \in (\text{IA}_n)_{\llbracket x_{n-1} \rrbracket, \llbracket x_n \rrbracket}$ we have $\tilde{\phi}(\eta_{\llbracket x_{n-1} \rrbracket}) = \tilde{\phi}(\eta_{\llbracket x_n \rrbracket})$, as desired.

We now turn to general edges e with endpoints $\llbracket z \rrbracket$ and $\llbracket z' \rrbracket$ and $\eta \in (\text{IA}_n)_e$. There exists some $v \in \text{Aut}(F_n)$ such that $v(x_n) = z$ and $v(x_{n-1}) = z'$, and hence $vP_{n-1,n}(x_n) = z'$. Setting $\eta' = v^{-1}\eta v \in (\text{IA}_n)_{\llbracket x_{n-1} \rrbracket, \llbracket x_n \rrbracket}$, we have

$$\begin{aligned} \tilde{\phi}(\eta_{\llbracket z \rrbracket}) &= v \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(v^{-1}\eta v) = v \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(\eta'), \\ \tilde{\phi}(\eta_{\llbracket z' \rrbracket}) &= vP_{n-1,n} \tilde{\phi}_{\llbracket x_n \rrbracket}(P_{n-1,n}^{-1} v^{-1} \eta v P_{n-1,n}) = v \cdot \tilde{\phi}_{\llbracket x_{n-1} \rrbracket}(\eta'). \end{aligned}$$

By the previous paragraph, we have $\tilde{\phi}_{\llbracket x_n \rrbracket}(\eta') = \tilde{\phi}_{\llbracket x_{n-1} \rrbracket}(\eta')$, so we conclude that $\tilde{\phi}(\eta_{\llbracket z \rrbracket}) = \tilde{\phi}(\eta_{\llbracket z' \rrbracket})$, as desired. \square

Claim 6 (conjugation relators) *If $\llbracket z \rrbracket$ and $\llbracket z' \rrbracket$ are vertices of $\hat{\mathcal{B}}_n$ and $\eta \in (\text{IA}_n)_{\llbracket z \rrbracket}$ and $\omega \in (\text{IA}_n)_{\llbracket z' \rrbracket}$, then $\tilde{\phi}(\omega_{\llbracket z' \rrbracket} \eta_{\llbracket z \rrbracket} \omega_{\llbracket z' \rrbracket}^{-1} (\omega \eta \omega^{-1})_{\llbracket \omega(z) \rrbracket}) = 1$.*

Proof Choose $\nu \in \text{Aut}(F_n)$ such that $\nu(x_n) = z$. We then have

$$\begin{aligned} \tilde{\phi}(\omega_{\llbracket z' \rrbracket} \eta_{\llbracket z \rrbracket} \omega_{\llbracket z' \rrbracket}^{-1}) &= \tilde{\phi}_{\llbracket z' \rrbracket}(\omega) \tilde{\phi}_{\llbracket z \rrbracket}(\eta) \tilde{\phi}_{\llbracket z' \rrbracket}(\omega)^{-1} = \pi(\tilde{\phi}_{\llbracket z' \rrbracket}(\omega)) \cdot \tilde{\phi}_{\llbracket z \rrbracket}(\eta) \\ &= \omega \cdot \tilde{\phi}_{\llbracket z \rrbracket}(\eta) = \omega \nu \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}(\nu^{-1} \eta \nu) \\ &= \omega \nu \cdot \tilde{\phi}_{\llbracket x_n \rrbracket}((\omega \nu)^{-1} \omega \eta \omega^{-1} (\omega \nu)) = \tilde{\phi}((\omega \eta \omega^{-1})_{\llbracket \omega(z) \rrbracket}), \end{aligned}$$

as desired. The second equality follows from the third part of [Proposition 3.13](#), the third equality follows from [Claim 4](#), and the remainder of the equalities are straightforward applications of the definitions. \square

Claims 5 and 6 imply that $\tilde{\phi}$ descends to a homomorphism $\phi: \text{IA}_n \rightarrow \mathcal{Q}_n$.

We have an inverse To complete the proof, it remains to prove the following:

Claim 7 *We have $\phi \circ \pi = \text{id}$.*

Proof [Claim 3](#) implies that this holds for the generators of \mathcal{Q}_n . \square

This completes the proof of [Theorem E](#). \square

5 Computations for the L–presentation

This section contains the postponed proofs of [Propositions 2.1, 2.3, 3.13 and 3.14](#). These proofs are done with the aid of a computer. We will discuss our computational framework in [Section 5.1](#) and then prove the propositions in [Sections 5.2–5.4](#).

We will use the following notation throughout the rest of the paper. Let $S_{\text{Aut}}(n)^*$ denote the free monoid on the set $S_{\text{Aut}}(n)$. In [Section 2](#), we defined a function $\theta: S_{\text{Aut}}(n) \rightarrow \text{End}(F(S_{\text{IA}}(n)))$. This naturally extends to a function

$$\theta: S_{\text{Aut}}(n)^* \rightarrow \text{End}(F(S_{\text{IA}}(n))).$$

5.1 Computational framework

As we discussed in the introduction, we use the GAP system to mechanically verify the large number of equations we have to check. These verifications are in the file `h2ia.g`, available as an [online supplement](#).

We use GAP's built-in functionality to model F_n as a free group on the eight generators $x_a, x_b, x_c, x_d, x_e, x_f, x_g$ and y . Since our computations never involve more than 8 variables, computations in this group suffice to show that our computations hold in general.

Elements of the sets $S_{\text{Aut}}(n)$ and $S_{\text{IA}}(n)$ are parametrized over basis elements from F_n and their inverses, so we model these sets using lists. For example, we model the generator M_{x_a, x_b} as the list ["M", x_a, x_b], C_{y, x_a} as ["C", y, x_a], and $M_{x_a^{-1}, [y, x_c]}$ as ["Mc", x_a^{-1}, y, x_c]. We model $P_{a, b}$ as ["P", x_a, x_b] and I_a as ["I", x_a]. The examples should make clear: the first entry in the list is a string key "M", "C", "Mc", "P" or "I", indicating whether the list represents a transvection, conjugation move, commutator transvection, swap or inversion. The parameters given as subscripts in the generator are then the remaining elements of the list, in the same order.

GAP's built-in free group functionality expects the basis elements to be variables, not lists, so we do not use it to model $S_{\text{Aut}}(n)^*$ and $F(S_{\text{IA}}(n))$. We model inverses of generators as follows: the inverse of ["M", x_a, x_b] is ["M", x_a, x_b^{-1}] and the inverse of ["C", x_a, x_b] is ["C", x_a, x_b^{-1}], but the inverse of ["Mc", x_a, x_b, x_c] is ["Mc", x_a, x_c, x_b]. Swaps and inversions are their own inverses. Technically, this means that we are not really modeling $S_{\text{Aut}}(n)^*$ and $F(S_{\text{IA}}(n))$; instead we model structures where the order relations for swaps and inversions and the relation (R0) for inverting commutator transvections are built in. This is not a problem because our verifications always show that certain formulas are trivial modulo our relations, and we can always apply the (R0) and order relations as needed.

We model words in $S_{\text{Aut}}(n)^*$ and $F(S_{\text{IA}}(n))$ as lists of generators and inverse generators. The empty word [] represents the trivial element. We wrote several functions in `h2ia.g` that perform common tasks on words. The function `pw` takes any number of words (reduced or not) as arguments and returns the freely reduced product of those words in the given order, as a single word. The function `iw` inverts its input word and the function `cyw` cyclically permutes its input word.

The function `iarel` outputs the relations $R_{\text{IA}}^0(n)$. We introduce some extra relations for convenience. The function `exiarel` outputs these extra relations and the code generating the list `exiarelchecklist` derives the extra relations from the basic relations. The function `theta` takes in a word w in $S_{\text{Aut}}(n)$ and a word v in $S_{\text{IA}}(n)$, and returns $\theta(w)(v)$. In addition to the functions described here, we often define simple macros for carrying out the verifications.

The function `applyrels` is particularly useful, because it inserts multiple relations into a word. It takes two inputs: a starting word and a list of words with placement indicators. The function recursively inserts the first word from the list in the starting

word at the given position, reduces the word, and then calls itself with the new word as the starting word and with the same list of insertions, with the first dropped.

For example, the following command appears in the justification of [Proposition 3.13](#):

```
applyrels(
  pw(
    theta([[ "M", xa, xb ], [ "M", xa, xb^-1 ] ], [[ "Mc", xb, xa, xe ] ]),
    [[ "Mc", xb, xe, xa ] ]
  ),
  [
    [6, iarel(5, [xa, xb, xe])],
    [6, iw(exiarel(1, [xb^-1, xe, xa]))],
    [2, iw(iarel(4, [xe, xa, xb]))],
    [2, iarel(6, [xb^-1, xe^-1, xa])],
    [2, iw(iarel(5, [xb, xe, xa]))]
  ]
)
```

This tells the GAP system to compute the effect of $\theta(M_{x_a, x_b} M_{x_a, x_b}^{-1})$ on $M_{x_b, [x_a, x_e]}$. Then it multiplies this by $M_{x_b, [x_e, x_a]}$, the inverse of $M_{x_b, [x_a, x_e]}$. It then freely reduces this word. The system inserts a version of (R5) after the sixth letter in this word, and reduces the result to a new word. Then it inserts the inverse of one of the extra relations after the sixth letter in the new word and reduces it. It continues with inserting relations and reducing the resulting expressions, inserting instances of (R4), (R5) and (R6). Since the entire expression evaluates to [], we have expressed

$$\theta(M_{x_a, x_b} M_{x_a, x_b}^{-1})(M_{x_b, [x_a, x_e]}) \cdot M_{x_b, [x_e, x_a]}$$

as a product of relations in \mathcal{Q}_n . In any example like this, an interested reader can reproduce our reduction process by removing all the list entries from the second input of the `applyrels` call, and then adding them back in one at a time, evaluating after each one.

5.2 Verifying the map to IA_n

First we prove [Proposition 2.1](#), which states that our relations $R_{IA}^0(n)$ hold in IA_n .

Proof of Proposition 2.1 The code generating the list `verifyiarel` generates examples of all the relations in $R_{IA}^0(n)$, with all allowable configurations of coincidences between the subscripts on the generators. It converts each of these relations into automorphisms of F_n and evaluates them on a basis for F_n , returning `true` if all basis

elements are unchanged. We evaluate on a fixed finite-rank free group, but since the basic relations involve at most six generators, those evaluations suffice to show the result in general. Since `verifyare1` evaluates to a list of `true`, this means that all these relations are true. \square

Next we prove [Proposition 2.3](#), which states that θ acts by conjugation when evaluated on generators (see (3)).

Proof of Proposition 2.3 The code generating the list `thetavsconjaut` goes through all possible configurations for a pair of generators s from $S_{\text{Aut}}(n)$ and t from $S_{\text{IA}}(n)$, evaluates $\theta(s)(t)$ as a product of generators, and then evaluates both $\theta(s)(t)$ and sts^{-1} on a basis for F_n . It returns `true` when both have the same effect on all basis elements. Since `thetavsconjaut` evaluates to a list of `true`, the proposition holds. \square

5.3 Verifying Proposition 3.13

Proof of Proposition 3.13 The action of $\text{Aut}(F_n)$ on \mathcal{Q}_n is given by our substitution rule endomorphism map

$$\theta: S_{\text{Aut}}(n) \rightarrow \text{End}(F(S_{\text{IA}}(n))).$$

First of all, it is clear that for each $s \in S_{\text{Aut}}(n)$, the element $\theta(s)$ defines an endomorphism of \mathcal{Q}_n . This is because the subgroup of $F(S_{\text{IA}}(n))$ normally generated by the relations of \mathcal{Q}_n is invariant under $\theta(s)$ by the definition of \mathcal{Q}_n .

Next, we verify that $\theta(s)$ is an automorphism of \mathcal{Q}_n . If s is a swap or an inversion, then it is clear from the definition of θ that this is the case. In the code generating the list `thetainverselist`, we compute $\theta(s)(\theta(s^{-1})(t))t^{-1}$ for $s = M_{x_a, x_b}$ and for all possible configurations of t relative to s . In each case, we reduce it to the trivial word using relations for \mathcal{Q}_n . It is not hard to deduce that $\theta(s)(\theta(s^{-1})(t)) = t$ in \mathcal{Q}_n for the remaining choices of $s = M_{x_a, x_b}^{-1}$, $M_{x_a^{-1}, x_b}$ and $M_{x_a^{-1}, x_b}^{-1}$, using the fact that it is true for $s = M_{x_a, x_b}$.

So this shows that θ defines an action

$$F(S_{\text{Aut}}(n)) \rightarrow \text{Aut}(\mathcal{Q}_n).$$

Now we need to verify that this action descends to an action of $\text{Aut}(F_n)$. To show this, it is enough to show that for every relation r in a presentation for $\text{Aut}(F_n)$, we have

$$(9) \quad \theta(r)(t) = t \quad \text{in } \mathcal{Q}_n,$$

for t taken from a generating set for \mathcal{Q}_n . To check this, we use the same version of Nielsen's presentation for $\text{Aut}(F_n)$ that we used in [17, Theorem 5.5]. The generators

are the same set $S_{\text{Aut}}(n)$ we use here, and the relations fall into five classes (N1)–(N5). Relations (N1) are sufficient for the subgroup generated by swaps and inversions, and (N2) are relations indicating how to conjugate transvections by swaps and inversions. It is an exercise to see that (9) holds for relations of class (N1) and (N2). Relations (N3)–(N5) are more complicated relations. For each of these, we compute $\theta(r)(t)t^{-1}$ on generators t (for t with enough configurations of subscripts to include a generating set) and reduce the resulting expressions to 1 using relations from \mathcal{Q}_n . These computations are given in the code generating the lists `thetaN3list`, `thetaN4list` and `thetaN5list`. Since these evaluate to lists of the trivial word, this verifies (9). We have shown that the action of an element of $\text{Aut}(F_n)$ on \mathcal{Q}_n does not depend on the word in $F(S_{\text{Aut}}(n))$ we use to represent it.

Since we have shown that θ defines an action, now we can check the three properties asserted in Proposition 3.13. We have already verified the first point (we took the definition of the action to agree with it). To verify the second point, we need to check that for $w, s \in S_{\text{IA}}(n)$, there is $\tilde{w} \in F(S_{\text{Aut}}(n))$ representing w with

$$(10) \quad \theta(\tilde{w})(s) = ws w^{-1} \quad \text{in } \mathcal{Q}_n.$$

In fact, it is enough to verify this for w and s in a smaller generating set, and the generating set that s is taken from may depend on w . In the code generating `thetaconjrellist`, for each choice of w from $S_{\text{MA}}(n)$, we lift w to $\tilde{w} \in F(S_{\text{Aut}}(n))$, and for several configurations of subscripts in the generator s , we reduce the element $\theta(\tilde{w})(s)ws^{-1}w^{-1}$ to the identity using relations from \mathcal{Q}_n . We use enough configurations of subscripts in s to cover all cases for s in a generating set (a conjugate of $S_{\text{MA}}(n)$).

To check the third point, we use the generating set $(S_{\text{Aut}}(n))_{[[x_n]]}$ for $(\text{Aut}(F_n))_{[[x_n]]}$ mentioned in the proof of Proposition 3.11 above, namely

$$\begin{aligned} & \{M_{x_a^\alpha, x_b} \mid 1 \leq a \leq n-1, 1 \leq b \leq n, \alpha = \pm 1, a \neq b\} \\ & \cup \{P_{a,b} \mid 1 \leq a, b \leq n, a \neq b, a \neq n\} \\ & \cup \{I_a \mid 1 \leq a \leq n\} \cup \{C_{x_n, x_a} \mid 1 \leq a \leq n-1\}. \end{aligned}$$

We need to check that for each of these generators, there is $w \in F(S_{\text{Aut}}(n))$ representing it with $\theta(w)(s)$ in $(\mathcal{Q}_n)_{[[x_n]]}$ (really, that $\theta(w)(s)$ is equal in \mathcal{Q}_n to an element of $(\mathcal{Q}_n)_{[[x_n]]}$). This is clear from the definition of θ for w a swap or inversion. It can be verified for $w = M_{x_a^\alpha, x_b}$ by inspecting Table 3. For w representing C_{x_n, x_a} , the fact that $\theta(w)(s) \in (\mathcal{Q}_n)_{[[x_n]]}$ follows from the second point in this proposition, since $C_{x_n, x_a} \in \text{IA}_n$. □

5.4 Verifying Proposition 3.14

Here we prove Proposition 3.14. We recall the statement: $\mathcal{K}_{n-1,1}$ is the kernel of the natural map $(\text{IA}_n)_{[[x_n]]} \rightarrow \text{IA}_{n-1}$, and the proposition asserts that the inclusion $\mathcal{K}_{n-1,1} \hookrightarrow \text{IA}_n$ factors as the composition of a map $\mathcal{K}_{n-1,1} \rightarrow \mathcal{Q}_n$ with the projection $\mathcal{Q}_n \rightarrow \text{IA}_n$.

The proof uses the finite L–presentation for $\mathcal{K}_{n-1,1}$ from [17]. We note that [17, Theorem D] asserts the existence of such a presentation, and [17, Theorem 6.2] gives the precise statement that we use in the computations. Since this L–presentation is in fact a presentation for $\mathcal{K}_{n-1,1}$, we use the same notation for $\mathcal{K}_{n-1,1}$ as a subset of IA_n and $\mathcal{K}_{n-1,1}$ as the group given by this presentation.

We do not reproduce the L–presentation here, but instead we describe some of its features. Its finite generating set is

$$S_{\mathcal{K}}(n) = \{M_{x_a^\alpha, [x_n^\epsilon, x_b^\beta]} \mid 1 \leq a, b \leq n-1, a \neq b, \alpha, \beta, \epsilon \in \{1, -1\}\} \\ \cup \{C_{x_n, x_a} \mid 1 \leq a \leq n-1\} \cup \{C_{x_a, x_n} \mid 1 \leq a \leq n-1\}.$$

The substitution endomorphisms of the L–presentation for $\mathcal{K}_{n-1,1}$ are indexed by a finite generating set $(S_{\text{Aut}}(n))_{[[x_n]]}$ for $(\text{Aut}(F_n))_{[[x_n]]}$. The endomorphisms themselves are the image of a map

$$\phi: (S_{\text{Aut}}(n))_{[[x_n]]} \rightarrow \text{End}(F(S_{\mathcal{K}}(n))).$$

Proof of Proposition 3.14 Since $S_{\mathcal{K}}(n)$ is a subset of $S_{\text{IA}}(n)$, we map $\mathcal{K}_{n-1,1}$ to \mathcal{Q}_n by sending each generator to the generator of the same name. To verify that this map on generators extends to a well-defined map of groups, we need to check that each defining relation from $\mathcal{K}_{n-1,1}$ maps to the trivial element of \mathcal{Q}_n . Since $\mathcal{K}_{n-1,1}$ is given by a L–presentation, we proceed as follows:

- (1) We check that each of the basic relations from $\mathcal{K}_{n-1,1}$ maps to the trivial element of \mathcal{Q}_n .
- (2) We check that for $s \in (S_{\text{Aut}}(n))_{[[x_n]]}$ and $t \in S_{\mathcal{K}}(n)$, we have

$$\phi(s)(t) = \theta(s)(t) \quad \text{in } \mathcal{Q}_n,$$

where we use $(S_{\text{Aut}}(n))_{[[x_n]]} \subset S_{\text{Aut}}(n)$ to plug s into θ , and we interpret both expressions in \mathcal{Q}_n using $F(S_{\mathcal{K}}(n)) \subset F(S_{\text{IA}}(n))$.

The first point is verified in the code generating the list `kfromialist`. The function `krel` produces the basic relations from $\mathcal{K}_{n-1,1}$, and we reduce each relation to the identity by applying relations from \mathcal{Q}_n . The second point is verified in the code generating the list `thetavsphlist`. For each choice of pairs of generators, we reduce the difference of θ and ϕ using relations from \mathcal{Q}_n .

With these two points verified, one can easily check by induction that every extended relation (starting with a basic relation, and applying any sequence of rewriting rules) maps to the identity element in \mathcal{Q}_n . \square

6 Generators for $H_2(\mathbf{IA}_n)$

In this section, we prove [Theorem A](#), which asserts that there exists a finite subset of $H_2(\mathbf{IA}_n)$ whose $GL_n(\mathbb{Z})$ -orbit spans $H_2(\mathbf{IA}_n)$. In fact, we gave an explicit list of generators in [Table 1](#); each generator is of the form h_r for a commutator relation r . This list is reproduced in [Table 4](#), which also introduces the notation $h_i(\cdot) \in H_2(\mathbf{IA}_n)$ for the associated elements of homology (this notation will be used during the calculations in [Section 7](#), though we will not use it in this section). The following theorem asserts that this list is complete; it is a more precise form of [Theorem A](#) and will be the main result of this section.

Theorem 6.1 *Fix $n \geq 2$. Let $S_H(n)$ be the set of commutator relators in [Table 4](#). Then the $GL_n(\mathbb{Z})$ -orbit of the set $\{h_r \mid r \in S_H(n)\}$ spans $H_2(\mathbf{IA}_n)$.*

Before proving [Theorem 6.1](#), we will use it to derive [Theorem B](#).

Proof of Theorem B Recall that this theorem asserts that for $n \geq 6$, the $GL_{n+1}(\mathbb{Z})$ -orbit of the image of the natural map $H_2(\mathbf{IA}_n) \rightarrow H_2(\mathbf{IA}_{n+1})$ spans $H_2(\mathbf{IA}_{n+1})$. Let $S_{n+1} \subset GL_{n+1}(\mathbb{Z})$ be the subgroup consisting of permutation matrices. By inspecting [Table 4](#), it is clear that the S_{n+1} -orbit of the image of $\{h_r \mid r \in S_H(n)\} \subset H_2(\mathbf{IA}_n)$ in $H_2(\mathbf{IA}_{n+1})$ is $\{h_r \mid r \in S_H(n+1)\}$. This uses the fact that $n \geq 6$, since the commutator relations in $S_H(n)$ use generators involving at most six basis elements. \square

We now turn to the proof of [Theorem 6.1](#). We start by introducing some notation. Let $F = F(S_{\mathbf{IA}}(n))$ and let $R \subset F$ denote the full set of relations of \mathbf{IA}_n , so $\mathbf{IA}_n = F/R$. Define $\widetilde{H_2(\mathbf{IA}_n)} = R/[F, R]$, and for $r \in R$ denote by $\|r\|$ the associated element of $\widetilde{H_2(\mathbf{IA}_n)}$. There is a natural map $\widetilde{H_2(\mathbf{IA}_n)} \rightarrow F^{ab}$, and the starting point for our proof is the following lemma. In it, recall from the beginning of [Section 5](#) that $S_{\text{Aut}}(n)^*$ is the free monoid on the set $S_{\text{Aut}}(n)$.

Lemma 6.2 *The group $\widetilde{H_2(\mathbf{IA}_n)}$ is an abelian group which is generated by*

$$\{\|\theta(w)(r)\| \mid w \in S_{\text{Aut}}(n)^* \text{ and } r \text{ is one of the relations (R0)–(R9) from [Table 2](#)\}.$$

Also, we have $H_2(\mathbf{IA}_n) = \ker(\widetilde{H_2(\mathbf{IA}_n)} \rightarrow F^{ab})$.

Proof The group $\widetilde{H_2(\mathbf{IA}_n)}$ is abelian since $[R, R] \subset [F, R]$. For $v \in F$ and $r \in R$, we have $[v, r] \in [F, R]$, so $\|vr v^{-1}\| = \|r\|$. The indicated generating set for $\widetilde{H_2(\mathbf{IA}_n)}$ thus follows from [Theorem E](#). As for the final statement of the lemma, we follow one

of the standard proofs of Hopf's formula [10]. The 5-term exact sequence in group homology associated to the short exact sequence

$$0 \rightarrow R \rightarrow F \rightarrow \text{IA}_n \rightarrow 0$$

is

$$H_2(F) \rightarrow H_2(\text{IA}_n) \rightarrow R/[F, R] \rightarrow H_1(F) \rightarrow H_1(\text{IA}_n) \rightarrow 0.$$

Since F is free, we have $H_2(F) = 0$, and the claim follows. \square

Our goal will be to take an element of $\widetilde{H_2(\text{IA}_n)}$ that happens to lie in $H_2(\text{IA}_n)$ and rewrite it as a sum of elements of the form

$$(11) \quad \{ \|\theta(w)(r)\| \mid w \in S_{\text{Aut}}(n)^* \text{ and } r \text{ is one of the relations (H1)–(H9) of Table 4} \}.$$

The relations (H1)–(H4) are the same as (R1)–(R4), and the relations (H5)–(H7) are the same as (R6)–(R9). The troublesome relations are (R0), (R5) and (R6), none of which lie in $H_2(\text{IA}_n)$. For $r \in R$, we have $\|r\| \in H_2(\text{IA}_n)$ if and only if the exponent-sum of each generator in $S_{\text{IA}}(n)$ appearing in it is 0. For our problematic relations (R0), (R5) and (R6), the exponent-sum of all the conjugations moves is already 0, so we will only need to study the exponent-sums of the commutator transvections.

We begin with the following lemma, which will allow us to mostly ignore our rewriting rules $\theta(\cdot)$.

Lemma 6.3 *Consider $w \in S_{\text{Aut}}(n)^*$, and let r be a relation of the form (R0), (R5) or (R6). Then $\|\theta(w)(r)\| = h + h'$, where h and h' are as follows:*

- $h \in H_2(\text{IA}_n)$ is a sum of elements from (11).
- h' is a sum of elements of the form

$$\{ \|r\| \mid r \text{ is one of the relations (R0), (R5) and (R6) from Table 2} \}.$$

Proof We can use induction to reduce to the case where $w = s \in S_{\text{Aut}}(n)$. The proof now is a combinatorial group-theoretic calculation: we will show how to rewrite $\theta(s)(r)$ as a product of relations of the desired form.

We start by dealing with the case where r is of the form (R0). Observe that

$$\theta(s)(M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}) \quad \text{and} \quad \theta(s)(M_{x_a^\alpha, [x_c^\gamma, x_b^\beta]})^{-1}$$

agree up to (R0), except in two cases. These are

$$\begin{aligned} \theta(M_{x_a^\alpha, x_b^\beta})(M_{x_b^{-\beta}, [x_c^\gamma, x_d^\delta]}) &= M_{x_a^\alpha, [x_c^\gamma, x_d^\delta]} M_{x_b^{-\beta}, [x_c^\gamma, x_d^\delta]} \\ \theta(M_{x_a^\alpha, x_b^\beta})(M_{x_b^\beta, [x_c^\gamma, x_d^\delta]}) &= C_{x_a, x_b}^\beta M_{x_a^{-\alpha}, [x_c^\gamma, x_d^\delta]} M_{x_b^\beta, [x_c^\gamma, x_d^\delta]} C_{x_a, x_b}^{-\beta}. \end{aligned}$$

- (H1) $[C_{x_a, x_b}^\beta, C_{x_c, x_d}^\delta] = 1$, possibly with $b = d$.
- (H2) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}] = 1$, possibly with $\{b, c\} \cap \{e, f\} \neq \emptyset$ or $x_a^\alpha = x_d^{-\delta}$ as long as $x_a^\alpha \neq x_d^\delta$, $a \notin \{e, f\}$ and $d \notin \{b, c\}$.
- (H3) $[C_{x_a, x_b}^\beta, M_{x_c^\gamma, [x_d^\delta, x_e^\epsilon]}] = 1$, possibly with $b \in \{d, e\}$ if $c \notin \{a, b\}$ and $a \notin \{c, d, e\}$.
- (H4) $[C_{x_c, x_b}^\beta C_{x_a, x_b}^\beta, C_{x_c, x_a}^\alpha] = 1$.
- (H5) $[C_{x_a, x_c}^{-\gamma}, C_{x_a, x_d}^{-\delta}][C_{x_a, x_b}^{-\beta}, M_{x_b^\beta, [x_c^\gamma, x_d^\delta]}] = 1$.
- (H6) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}][M_{x_d^\delta, [x_e^\epsilon, x_f^\zeta]}, M_{x_d^\delta, [x_c^\gamma, x_b^\beta]}] \times [M_{x_d^\delta, [x_c^\gamma, x_b^\beta]}, C_{x_d, x_e}^{-\epsilon}] = 1$, possibly with $b = e$ or $c = e$.
- (H7) $[M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]}, C_{x_a, x_b}^\beta][C_{x_c, x_d}^{-\delta}, M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]}][M_{x_c^\gamma, [x_a^\alpha, x_b^\beta]}, M_{x_c^\gamma, [x_a^\alpha, x_d^\delta]}] = 1$, possibly with $b = d$.
- (H8) $[M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, C_{x_a, x_d}^\delta C_{x_b, x_d}^\delta C_{x_c, x_d}^\delta] = 1$.
- (H9) $[C_{x_a, x_c}^\gamma C_{x_b, x_c}^\beta, C_{x_a, x_b}^\beta C_{x_c, x_b}^\beta][M_{x_a^\alpha, [x_b^\beta, x_c^\gamma]}, C_{x_b, x_a}^\alpha C_{x_c, x_a}^\alpha] = 1$.

Table 4: The set $S_H(n)$ of basic commutator relators such that the $GL_n(\mathbb{Z})$ -orbit of $\{h_r \mid r \in S_H(n)\}$ spans $H_2(\text{IA}_n)$. Distinct letters are assumed to represent distinct indices unless stated otherwise. We use the notation $h_i(\cdot)$, with inputs the appropriate $x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon, x_f^\zeta$, for the elements in $H_2(\text{IA}_n)$, which we use later.

In the first case,

$$\theta(M_{x_a^\alpha, x_b}^\beta)(M_{x_b^{-\beta}, [x_d^\delta, x_c^\gamma]}^{-1}) = M_{x_b^{-\beta}, [x_c^\gamma, x_d^\delta]} M_{x_a^\alpha, [x_c^\gamma, x_d^\delta]}.$$

In the second case,

$$\theta(M_{x_a^\alpha, x_b}^\beta)(M_{x_b^\beta, [x_d^\delta, x_c^\gamma]}^{-1}) = C_{x_a, x_b}^\beta M_{x_b^\beta, [x_c^\gamma, x_d^\delta]} M_{x_a^{-\alpha}, [x_c^\gamma, x_d^\delta]} C_{x_a, x_b}^{-\beta}.$$

In both cases, the two expressions differ by an application of (R2). This means that $\theta(s)$ of an (R0) relation can always be written using (R0) and (R2) relations.

Next we explain the computations that prove the lemma for (R5) and (R6) relations. These are in the list `rewritetheta(R5) (R6)`. In these computations we reduce $\theta(s)(r)$ to the trivial word where $s \in S_A^{\pm 1}$ and r is an (R5) or (R6) relation. These reductions may use any of the basic relations for IA_n , including (R5) and (R6) themselves, but notably may not use the images of (R5) and (R6) relations under θ . We may use the images of (R1)–(R4), (R7)–(R9), (H8) and (H9) under θ .

Despite these restrictions, we may use the extra relations from `exiare1` in these computations. Our relation `exiare1(3, [xa,xb,xc,xd])` is (H8), and the relation

$\text{exiare1}(4, [\text{xa}, \text{xb}, \text{xc}, \text{xd}])$ is equivalent to (H8) modulo the basic relations. Relation $\text{exiare1}(7, [\text{xa}, \text{xb}, \text{xc}])$ is (H9), and relation $\text{exiare1}(6, [\text{xa}, \text{xb}, \text{xc}])$ and relation $\text{exiare1}(8, [\text{xa}, \text{xb}, \text{xc}, \text{xd}])$ are equivalent to (H9) modulo the basic relations. All the other exiare1 relations can be derived without using images of (R5) or (R6) under θ . These facts can be verified by inspecting exiare1checklist .

If s is a swap or an inversion move, then acting by $\theta(s)$ is always the same as acting on the parameters of the relation by s in the obvious way. Therefore the list $\text{rewritetheta}(R5)(R6)$ only contains cases where s is a transvection.

We use several redundancies between different forms of the relations (R5) and (R6) to reduce the number of computations. Inverting the parameter x_b^β in (R5) (as it appears in Table 2) is the same as cyclically permuting the relation. Inverting the parameter x_a^α in (R6) is the same as applying a relation from (R2) to the original (R6) relation. Swapping the roles of x_b^β and x_c^γ in (R6) is the same as inverting and cyclically permuting the original (R6) relation and applying a relation from (R2).

We use the identity

$$C_{x_a, x_b}^{-\beta} \theta(M_{x_a^\alpha, x_b}^\beta)(t) C_{x_a, x_b}^\beta = \theta(M_{x_a^{-\alpha}, x_b}^{-\beta})(t),$$

which holds in IA_n for any $t \in \text{IA}_n$. This is a consequence of Proposition 3.13. In particular, this means that we only need to consider one of $\theta(M_{x_a^\alpha, x_b^\beta})(r)$ and $\theta(M_{x_a^{-\alpha}, x_b^{-\beta}})(r)$; if one is trivial then so is the other.

Since the computations in $\text{rewritetheta}(R5)(R6)$ rewrite all configurations of $\theta(s)(r)$ for r an (R5) or (R6) relation, up to these reductions, this proves the lemma. \square

The next lemma allows us to deal with certain combinations of (R5) and (R6) relations. The ordered triple of generators of F_n involved in a commutator transvection $M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]}$ is (x_i, x_j, x_j) . There are eight commutator transvections involving a given triple of generators.

Lemma 6.4 *Fix distinct $1 \leq a, b, c \leq n$, and let $w \in R \cap [F, F]$ be a product of (R5) and (R6) relations whose commutator transvections involve only (x_a, x_b, x_c) , in order. Then $\|w\|$ can be written as a sum of elements of the form $\|v\|$ with v an (H2) relation.*

Proof Let F' be the subgroup of F generated by the eight commutator transvections involving (x_a, x_b, x_c) and the two conjugation moves $\{C_{x_a, x_b}, C_{x_a, x_c}\}$, and let $R' \subset F'$ be the normal closure in F' of the (R5) and (R6) relations that can be written as products of elements of F' . We thus have $w \in R' \cap [F', F']$. Our first goal is to better understand $(R' \cap [F', F'])/[F', R']$ and $R'/[F', R']$. Consider the exact

sequence of abelian groups

$$0 \rightarrow \frac{R' \cap [F', F']}{[F', R']} \rightarrow \frac{R'}{[F', R']} \rightarrow \frac{R'}{R' \cap [F', F']} \rightarrow 0.$$

We find generators for the first group in the sequence by considering a related exact sequence of free abelian groups.

Let v_1, \dots, v_8 be the eight commutator transvections in $S_{IA}(n)$ that only involve (x_a, x_b, x_c) , enumerated as in Table 5. Similarly, let r_1, \dots, r_8 be the eight (R5) relations in F' and let r_9, \dots, r_{16} denote the eight (R6) relations lying in F' , enumerated as in Table 5. Let A be the free abelian group freely generated by r_1, \dots, r_{16} , and let B be the free abelian group freely generated by v_1, \dots, v_8 . We consider the map $A \rightarrow B$ that counts the exponent-sum of each commutator transvection generator. Let C denote the kernel of this map and let B' denote the image.

		name	relation
		r_1	$C_{x_a, x_b} M_{x_a, [x_b, x_c]} C_{x_a, x_b}^{-1} M_{x_a, [x_b^{-1}, x_c]}$
		r_2	$C_{x_a, x_b} M_{x_a^{-1}, [x_b, x_c]} C_{x_a, x_b}^{-1} M_{x_a^{-1}, [x_b^{-1}, x_c]}$
		r_3	$C_{x_a, x_c}^{-1} M_{x_a, [x_b, x_c]} C_{x_a, x_c} M_{x_a, [x_b, x_c^{-1}]}$
		r_4	$C_{x_a, x_c}^{-1} M_{x_a^{-1}, [x_b, x_c]} C_{x_a, x_c} M_{x_a^{-1}, [x_b, x_c^{-1}]}$
name	generator	r_5	$C_{x_a, x_c}^{-1} M_{x_a, [x_b^{-1}, x_c]} C_{x_a, x_c} M_{x_a, [x_b^{-1}, x_c^{-1}]}$
v_1	$M_{x_a, [x_b, x_c]}$	r_6	$C_{x_a, x_c}^{-1} M_{x_a^{-1}, [x_b^{-1}, x_c]} C_{x_a, x_c} M_{x_a^{-1}, [x_b^{-1}, x_c^{-1}]}$
v_2	$M_{x_a^{-1}, [x_b, x_c]}$	r_7	$C_{x_a, x_b} M_{x_a, [x_b, x_c^{-1}]} C_{x_a, x_b}^{-1} M_{x_a, [x_b^{-1}, x_c^{-1}]}$
v_3	$M_{x_a, [x_b^{-1}, x_c]}$	r_8	$C_{x_a, x_b} M_{x_a^{-1}, [x_b, x_c^{-1}]} C_{x_a, x_b}^{-1} M_{x_a^{-1}, [x_b^{-1}, x_c^{-1}]}$
v_4	$M_{x_a^{-1}, [x_b^{-1}, x_c]}$	r_9	$M_{x_a, [x_b, x_c]} M_{x_a^{-1}, [x_b, x_c]} [C_{x_a, x_b}^{-1}, C_{x_a, x_c}^{-1}]$
v_5	$M_{x_a, [x_b, x_c^{-1}]}$	r_{10}	$M_{x_a, [x_b^{-1}, x_c]} M_{x_a^{-1}, [x_b^{-1}, x_c]} [C_{x_a, x_b}, C_{x_a, x_c}^{-1}]$
v_6	$M_{x_a^{-1}, [x_b, x_c^{-1}]}$	r_{11}	$M_{x_a, [x_b, x_c^{-1}]} M_{x_a^{-1}, [x_b, x_c^{-1}]} [C_{x_a, x_b}^{-1}, C_{x_a, x_c}]$
v_7	$M_{x_a, [x_b^{-1}, x_c^{-1}]}$	r_{12}	$M_{x_a, [x_b^{-1}, x_c^{-1}]} M_{x_a^{-1}, [x_b^{-1}, x_c^{-1}]} [C_{x_a, x_b}, C_{x_a, x_c}]$
v_8	$M_{x_a^{-1}, [x_b^{-1}, x_c^{-1}]}$	r_{13}	$M_{x_a^{-1}, [x_b, x_c]} M_{x_a, [x_b, x_c]} [C_{x_a, x_b}^{-1}, C_{x_a, x_c}^{-1}]$
		r_{14}	$M_{x_a^{-1}, [x_b^{-1}, x_c]} M_{x_a, [x_b^{-1}, x_c]} [C_{x_a, x_b}, C_{x_a, x_c}^{-1}]$
		r_{15}	$M_{x_a^{-1}, [x_b, x_c^{-1}]} M_{x_a, [x_b, x_c^{-1}]} [C_{x_a, x_b}^{-1}, C_{x_a, x_c}]$
		r_{16}	$M_{x_a^{-1}, [x_b^{-1}, x_c^{-1}]} M_{x_a, [x_b^{-1}, x_c^{-1}]} [C_{x_a, x_b}, C_{x_a, x_c}]$

Table 5: Labels for the eight commutator transvections using x_a, x_b and x_c in order, and for the sixteen (R5) and (R6) relations using these commutator transvections.

Since R' is normally generated by relations r_1, \dots, r_{16} , we know that $R'/[F', R']$ is generated by the images of these relations. Thus there is a surjection $A \rightarrow R'/[F', R']$ that sends each basis element to the image of the relation with the same name. The group B is a subgroup of $F'/[F', F']$. The natural map $R'/[F', R'] \rightarrow F'/[F', F']$ counts exponent-sums of generators. Since the generators r_1, \dots, r_{16} all have zero exponent-sum for conjugation move generators, we do not lose any information by counting only commutator transvection generators in B . This means we have a commuting square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ R'/[F', R'] & \longrightarrow & F'/[F', F'] \end{array}$$

The subgroup B' thus maps surjectively onto $(R'[F', F'])/[F', F']$, which is isomorphic to $R'/(R' \cap [F', F'])$. Therefore we have a commuting diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C & \longrightarrow & A & \longrightarrow & B' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & (R' \cap [F', F'])/[F', R'] & \longrightarrow & R'/[F', R'] & \longrightarrow & R'/(R' \cap [F', F']) & \longrightarrow & 0 \end{array}$$

The map $B' \rightarrow F'/[F', F']$ is injective, so $B' \rightarrow R'/(R' \cap [F', F'])$ is also injective. By construction, $A \rightarrow R'/[F', R']$ is surjective. It follows from a simple diagram chase that $C \rightarrow (R' \cap [F', F'])/[F', R']$ is surjective.

The map $A \rightarrow B$ is given by the 8×16 matrix

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{pmatrix}$$

A straightforward linear algebra computation shows that C , the kernel of this map, is generated by the nine vectors

$$\begin{aligned} r_1 - r_3 - r_5 + r_7, & \quad r_2 - r_4 - r_6 + r_8, & \quad -r_1 - r_2 + r_{13} + r_{14}, \\ -r_3 - r_4 + r_{13} + r_{15}, & \quad r_1 + r_2 - r_5 - r_6 - r_{13} + r_{12}, & \\ r_9 - r_{13}, & \quad r_{10} - r_{14}, & \quad r_{11} - r_{15}, r_{12} - r_{16}. \end{aligned}$$

Since C surjects on $(R' \cap [F', F'])/[F', R']$, we know that $(R' \cap [F', F'])/[F', R']$ is generated by the images of these nine elements.

We will now describe calculations that show that each of the generators above is equivalent modulo $[F', R']$ to an (H2) relation. In each case, we find representatives in $R' \cap [F', F']$ of the image of the given element of C . Since we are working modulo $[F', R']$ we may conjugate any r_i in computing the representative. In reducing to an (H2) relation, we may also apply any relation at all, as long as we apply its inverse somewhere else.

The last four generators are easily equivalent to (H2) relations. We skip the second kernel generator because its image is equal to the first after inverting x_a , and we skip the fourth because its image is equal to the third after swapping x_b and x_c . The three computations in the list `kernelList` finish the lemma by showing that the first, third and fifth generators are equivalent to (H2) relations. □

Proof of Theorem 6.1 We must show that every element of $\widehat{H_2(\text{IA}_n)}$ that happens to lie in $H_2(\text{IA}_n)$ can be written as a sum of elements of

$$\{\|\theta(w)(r)\| \mid w \in S_{\text{Aut}}(n)^* \text{ and } r \text{ is one of the relations (H1)–(H9) from Table 4}\}.$$

Combining Lemmas 6.2 and 6.3 with the fact that (R0) and (R5) and (R6) are the only relations in our L–presentation for IA_n that do not appear as one of the commutator relations in Table 4, we see that it enough to deal with sums of elements of the set

$$\{\|r\| \mid r \text{ is one of the relations (R0), (R5) and (R6)}\}.$$

So consider $\|w\| \in H_2(\text{IA}_n)$ that can be written as

$$\|w\| = \sum_{i=1}^m \|r_i\|$$

with each r_i either an (R0), (R5) or (R6) relation.

For any choice of distinct $1 \leq a, b, c \leq n$, we consider the commutator transvection generators involving (x_a, x_b, x_c) or (x_a, x_c, x_b) , and the (R5), (R6) and (R0) relations involving only these commutator transvections. We write

$$\|w\| = \sum_{i=1}^{n+\binom{n-1}{2}} \|w_i\|,$$

where each $\|w_i\|$ is a sum of (R5), (R6) and (R0) relations involving only a single choice of $(x_a, \{x_b, x_c\})$. To prove the theorem, it is enough to show that we can write

each of the $\|w_i\|$ as a sum of our generators for $H_2(\text{IA}_n)$. So we assume $\|w\| = \|w_i\|$ for some i ; this amounts to fixing a choice of $(x_a, \{x_b, x_c\})$ and assuming $\|w\|$ is a sum of (R0), (R5) and (R6) generators using only commutator transvections involving this triple.

We call a commutator transvection $M_{x_i^\alpha, [x_j^\beta, x_k^\gamma]}$ *positive* if $j < k$, and *negative* otherwise. Each (R5) or (R6) relation contains two positive commutator transvections (and no negative ones), or two negative ones (and no positive ones). Suppose $\|r_i\|$ is an (R5) or (R6) relation with negative generators, appearing in the sum defining $\|w\|$. By inserting an (R0) relation and its inverse into r_i , we replace both of the negative generators with positive ones. Let r'_i denote the word we get by doing this to r_i . Since we have added and subtracted the same element in $\overline{H_2(\text{IA}_n)}$, we have $\|r'_i\| = \|r_i\|$. Modifying an (R5) or (R6) relation in this way gives us the inverse of an (R5) or (R6) relation involving the same $(x_a, \{x_b, x_c\})$, up to cyclic permutation of the relation. So we interpret this move as rewriting the sum defining $\|w\|$: we replace the relation $\|r_i\|$ with the new relation $\|r'_i\|$, which is an (R5) or (R6) relation without negative commutator transvections. We proceed to eliminate all the negative commutator transvections in (R5) and (R6) relations in $\|w\|$ this way.

Having done this, the only negative commutator transvections the sum defining $\|w\|$ appear in (R0) relations. Since $\|w\| \in H_2(\text{IA}_n)$, the negative generators appear with exponent-sum zero; so the (R0) relations appear in inverse pairs. This means that we can simply rewrite the sum without any (R0) relations. So $\|w\|$ is a sum of (R5) and (R6) relations whose only commutator transvections are positive ones involving $(x_a, \{x_b, x_c\})$. Then $\|w\|$ satisfies the hypotheses of [Lemma 6.4](#) and therefore is a sum of (H2) generators. □

7 Coinvariants and congruence subgroups

This section contains the proofs of [Theorems C and D](#), which can be found in [Section 7.2](#) and [7.3](#), respectively. Both of these proofs depend on calculations that are contained in [Section 7.1](#).

7.1 The action of $\text{GL}_n(\mathbb{Z})$ on $H_2(\text{IA}_n)$

This section is devoted to understanding the action of $\text{GL}_n(\mathbb{Z})$ on our generators for $H_2(\text{IA}_n)$. The results in this section consist of long lists of equations that are verified by a computer, so on their first pass a reader might want to skip to the next two sections to see how they are used. For $i = 1, \dots, 9$, we use the notation $h_i(x_{a_1}^{\alpha_1}, \dots, x_{a_{k_i}}^{\alpha_{k_i}}) \in H_2(\text{IA}_n)$ for the image in $H_2(\text{IA}_n)$ of the i^{th} relation from $S_H(n)$,

with the given parameters, as specified in Table 4. Since the action of $GL_n(\mathbb{Z})$ is induced from the action of $Aut(F_n)$, we record the action of various $Aut(F_n)$ generators on these generators.

The computations justifying Lemmas 7.2–7.6 are in the file h2ia.g. We use the Hopf isomorphism $H_2(IA_n) \cong (R \cap [F, F])/[F, R]$, where $F = F(S_{IA}(n))$ and $R < F$ is the group of relations of IA_n . We justify these equations by performing computations in $R \cap [F, F] \subset F$. In each computation, we start with a word representing one side of the equation and reduce to the trivial word using words representing the other side. Since $[F, R]$ is trivial, we may use any relations in inverse pairs, we may apply relations from in any order, and we may cyclically permute relations.

We note some identities, which we leave as an exercise.

Lemma 7.1 *The following identities hold in $H_2(IA_n)$. The letters in subscripts are assumed distinct unless otherwise noted.*

- (a) $h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\delta) = -h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^{-\delta})$, even if $b = d$.
- (b) $h_3(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\delta, x_e^\epsilon) = -h_3(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\delta, x_e^{-\epsilon})$, even if $b = e$ or $c = e$.
- (c) $h_3(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\delta, x_e^\epsilon) = -h_3(x_a^\alpha, x_c^\gamma, x_b^\beta, x_b^\delta, x_e^\epsilon)$, even if $b = e$ or $c = e$.

We also need the following, which is not obvious.

Lemma 7.2 *The following identities hold in $H_2(IA_n)$. The letters in subscripts are assumed distinct unless otherwise noted.*

- (a) $h_6(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta, x_e^\epsilon) = h_6(x_a^\alpha, x_c^\gamma, x_e^{-\epsilon}, x_d^\delta, x_e^\epsilon) - h_7(x_a^\alpha, x_e^\epsilon, x_d^\delta, x_e^\epsilon) - h_7(x_a^\alpha, x_e^{-\epsilon}, x_d^\delta, x_e^\epsilon)$.
- (b) $h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon) = -h_6(x_a^\alpha, x_c^\gamma, x_b^\beta, x_d^\delta, x_e^\epsilon)$, even if $b = e$ or $c = e$.

Proof Computations justifying these equations appear in the list lemma7pt2. □

We proceed by expressing the action of many elementary matrices from $GL_n(\mathbb{Z})$ on our generators.

Lemma 7.3 *The following identities hold in $H_2(IA_n)$. The letters in subscripts are assumed distinct unless otherwise noted.*

- (a) $M_{x_b^\beta, x_e}^\epsilon \cdot h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) = h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_1(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta)$.
- (b) $M_{x_b^\beta, x_e}^\epsilon \cdot h_1(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta) = h_1(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta)$.

- (c) $M_{x_b^\beta, x_d}^\delta \cdot h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) = h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_1(x_a^\alpha, x_d^\delta, x_c^\gamma, x_b^\beta).$
- (d) $M_{x_a^\alpha, x_e}^{-\epsilon} \cdot h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) = h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) - h_3(x_a^\alpha, x_e^\epsilon, x_b^\beta, x_c^\gamma, x_d^\delta),$
even if $x_e^\epsilon = x_d^\delta.$
- (e) $M_{x_a^\alpha, x_e}^{-\epsilon} \cdot h_3(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_f^\zeta) =$
 $h_3(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_f^\zeta) - h_2(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon, x_f^\zeta),$ even if $\{b, c\} \cap \{e, f\} \neq \emptyset.$
- (f) $M_{x_a^\alpha, x_d}^\delta \cdot h_2(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^{-\delta}, x_f^\zeta, x_e^\epsilon) =$
 $h_2(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^{-\delta}, x_f^\zeta, x_e^\epsilon) - h_2(x_a^\alpha, x_b^\beta, x_c^\gamma, x_a^{-\alpha}, x_e^\epsilon, x_f^\zeta),$
even if $\{b, c\} \cap \{e, f\} \neq \emptyset.$

Proof These computations appear in lemma7pt3. The equations where coincidences are allowed are justified in several different computations. □

Lemma 7.4 *The following identities hold in $H_2(\text{IA}_n)$. The letters in subscripts are assumed distinct.*

- (a) $M_{x_d^\delta, x_b}^\beta \cdot h_4(x_a^\alpha, x_d^\delta, x_c^\gamma) = h_4(x_a^\alpha, x_d^\delta, x_c^\gamma) + h_4(x_a^\alpha, x_b^\beta, x_c^\gamma).$
- (b) $M_{x_d^\delta, x_b}^\beta \cdot h_4(x_a^\alpha, x_b^\beta, x_c^\gamma) = h_4(x_a^\alpha, x_b^\beta, x_c^\gamma).$
- (c) $M_{x_b^\beta, x_c}^{-\gamma} \cdot h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) =$
 $h_1(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) - h_3(x_b^\beta, x_d^\delta, x_c^\gamma, x_a^\alpha, x_c^\gamma) +$
 $h_5(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_4(x_c^\gamma, x_d^\delta, x_a^\alpha).$
- (d) $M_{x_d^\delta, x_e}^\epsilon \cdot h_8(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) =$
 $h_8(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_8(x_a^\alpha, x_b^\beta, x_c^\gamma, x_e^\epsilon) + ((H1) \text{ generators}).$
- (e) $M_{x_d^\delta, x_e}^\epsilon \cdot h_8(x_a^\alpha, x_b^\beta, x_c^\gamma, x_e^\epsilon) = h_8(x_a^\alpha, x_b^\beta, x_c^\gamma, x_e^\epsilon).$

Proof These computations appear in lemma7pt4. □

Lemma 7.5 *The following identities hold in $H_2(\text{IA}_n)$. The letters in subscripts are assumed distinct.*

- (a) $M_{x_f^\zeta, x_e}^\epsilon \cdot h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_f^\zeta) =$
 $h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_f^\zeta) + h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon).$
- (b) $M_{x_f^\zeta, x_e}^\epsilon \cdot h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon) = h_6(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta, x_e^\epsilon).$

- (c) $M_{x_c^\gamma, x_e}^{-\epsilon} \cdot h_7(x_a^\alpha, x_b^\beta, x_e^\epsilon, x_d^\delta) = h_7(x_a^\alpha, x_b^\beta, x_e^\epsilon, x_d^\delta) + h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_6(x_e^\epsilon, x_b^\beta, x_a^\alpha, x_c^\gamma, x_d^\delta) + ((H1)–(H3) \text{ generators}).$
- (d) $M_{x_c^\gamma, x_e}^{-\epsilon} \cdot h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) = h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_2(x_c^{-\gamma}, x_e^{-\epsilon}, x_d^{-\delta}, x_c^\gamma, x_b^\beta, x_a^\alpha).$
- (e) $M_{x_d^\delta, x_b}^\beta \cdot h_7(x_a^\alpha, x_d^\delta, x_c^\gamma, x_d^\delta) = h_7(x_a^\alpha, x_d^\delta, x_c^\gamma, x_d^\delta) + h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\beta) + h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_d^\delta) + h_7(x_a^\alpha, x_d^\delta, x_c^\gamma, x_b^\beta) + ((H1) \text{ generators}).$
- (f) $M_{x_d^\delta, x_b}^\beta \cdot h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\beta) = h_7(x_a^\alpha, x_b^\beta, x_c^\gamma, x_b^\beta).$
- (g) $M_{x_d^\delta, x_b}^\beta \cdot h_7(x_a^\alpha, x_d^{-\delta}, x_c^\gamma, x_d^\delta) = h_7(x_a^\alpha, x_d^{-\delta}, x_c^\gamma, x_d^\delta) + h_7(x_a^\alpha, x_b^{-\beta}, x_c^\gamma, x_b^\beta) + h_7(x_a^\alpha, x_d^{-\delta}, x_c^\gamma, x_b^\beta) - h_7(x_a^\alpha, x_b^{-\beta}, x_c^\gamma, x_d^{-\delta}).$
- (h) $M_{x_d^\delta, x_b}^\beta \cdot h_7(x_a^\alpha, x_b^{-\beta}, x_c^\gamma, x_b^\beta) = h_7(x_a^\alpha, x_b^{-\beta}, x_c^\gamma, x_b^\beta).$
- (i) $M_{x_f^\zeta, x_e}^\epsilon \cdot h_6(x_a^\alpha, x_f^\zeta, x_c^\gamma, x_d^\delta, x_f^\zeta) = h_6(x_a^\alpha, x_f^\zeta, x_c^\gamma, x_d^\delta, x_f^\zeta) - h_7(x_a^\alpha, x_e^\epsilon, x_d^\delta, x_f^\zeta) - h_7(x_a^\alpha, x_e^\epsilon, x_d^\delta, x_e^\epsilon) + h_6(x_a^\alpha, x_f^\zeta, x_c^\gamma, x_d^\delta, x_e^\epsilon) - h_7(x_a^\alpha, x_e^{-\epsilon}, x_d^\delta, x_f^\zeta) - h_7(x_a^\alpha, x_e^{-\epsilon}, x_d^\delta, x_e^\epsilon) + h_6(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta, x_f^\zeta) + h_6(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta, x_e^\epsilon).$
- (j) $M_{x_f^\zeta, x_e}^\epsilon \cdot h_6(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta, x_e^\epsilon) = h_6(x_a^\alpha, x_e^\epsilon, x_c^\gamma, x_d^\delta, x_e^\epsilon).$

Proof These computations appear in lemma7pt5. □

Lemma 7.6 *The following identities hold in $H_2(\text{IA}_n)$. The letters in subscripts are assumed distinct.*

- (a) $M_{x_b^\beta, x_d}^\delta \cdot h_9(x_a^\alpha, x_b^\beta, x_c^\gamma) = h_9(x_a^\alpha, x_b^\beta, x_c^\gamma) + h_9(x_a^\alpha, x_d^\delta, x_c^\gamma) + ((H1)–(H5) \text{ generators}).$
- (b) $M_{x_b^\beta, x_d}^\delta \cdot h_9(x_a^\alpha, x_d^\delta, x_c^\gamma) = h_9(x_a^\alpha, x_d^\delta, x_c^\gamma) + ((H1)–(H6) \text{ generators}).$

Proof These computations are contained in the list lemma7pt6, but they take some explanation. We use the relations `exiare1(5, [...])` frequently in these computations; these relations are always combinations of (H1) and (H4) relations. We

use `exiarel(7, [xa,xb,xc])` to represent $h_9(x_a^\alpha, x_b^\beta, x_c^\gamma)$, but we also use other relations in this computation. The relation `exiarel(8, [xa,xb,xc,xd])` is an expanded version of this relation that behaves better under this action. Its derivation in `exiarelchecklist` shows that it differs from $-h_9(x_a^\alpha, x_b^\beta, x_c^\gamma)$ only by (H1), (H4) and (H5) relations. We also use `exiarel(6, [xa,xb,xc,xd])`; this differs from $h_9(x_a^\alpha, x_b^\beta, x_c^\gamma)$ by (H1) and (H4) relations, and an (R6) relation. This is apparent in the derivation of `exiarel(7, [xa,xb,xc])` in `exiarelchecklist`.

The computation justifying [Lemma 7.6\(a\)](#) shows directly that the image of the relation `exiarel(8, [xa,xb,xc,xd])` under `[["M",xb,xd]]` can be reduced to the trivial word by applying

- `exiarel(6, [xa,xd,xc])`,
- `iw(exiarel(7, [xa,xb,xc]))`,
- (H1)–(H5) relations (some in `exiarel(5, [...])` relations),
- (R5) and (R6) relations, and
- elements from $[F, R]$, including inverse pairs of instances of `exiarel(1, [...])` and `exiarel(3, [...])`.

Since we start and end with elements of $R \cap [F, F]$ in this computation, the use of (R5) and (R6) (in one case inside `exiarel(6, [xa,xb,xc,xd])`) is inconsequential; by [Lemma 6.4](#) this can only change the outcome by (H2) relations. So this proves [Lemma 7.6\(a\)](#).

The computation justifying [Lemma 7.6\(b\)](#) is similar, but uses an instance of the relation `exiarel(2, [...])`. The derivation in `exiarelchecklist` shows that this relation is a combination of (H2), (H5) and (H6) relations, and elements of $[F, R]$. □

7.2 Coinvariants of $H_2(\mathbf{IA}_n)$

In this section, we prove [Theorem C](#), which asserts that for the $\text{GL}_n(\mathbb{Z})$ -coinvariants of $H_2(\mathbf{IA}_n)$ vanish for $n \geq 6$.

Proof of Theorem C We use the generators (H1)–(H9) from [Table 4](#). To show that the coinvariants $H_2(\mathbf{IA}_n)_{\text{GL}_n(\mathbb{Z})}$ are trivial, we show that the coinvariance class of each of these generators is trivial. The coinvariants are defined to be the largest quotient of $H_2(\mathbf{IA}_n)$ with a trivial induced action of $\text{GL}(n, \mathbb{Z})$. Since the action of $\text{GL}(n, \mathbb{Z})$ on $H_*(\mathbf{IA}_n)$ is induced by the action of $\text{Aut}(F_n)$ on \mathbf{IA}_n , this means that $H_2(\mathbf{IA}_n)_{\text{GL}_n(\mathbb{Z})}$ is the quotient of $H_2(\mathbf{IA}_n)$ by the subgroup generated by classes of the form $f \cdot r - r$, where $f \in \text{Aut}(F_n)$ and $r \in H_2(\mathbf{IA}_n)$. Elements of the form $f \cdot r - r$ are called *coboundaries*.

In fact, we have already shown in Lemmas 7.1–7.6 that the subgroup of $H_2(\mathrm{IA}_n)$ generated by coboundaries contains our generators from Theorem 6.1. We show this for each generator in the equations above as follows:

- (H1) in equations (a) and (c) of Lemma 7.3, also using an observation from Lemma 7.1,
- (H2) in equations (e) and (f) of Lemma 7.3,
- (H3) in Lemma 7.3(d), also using an observation from Lemma 7.1,
- (H4) in Lemma 7.4(a),
- (H5) in Lemma 7.4(c),
- generic (H6) in Lemma 7.5(a),
- (H7) in equations (c), (e) and (g) of Lemma 7.5,
- the special cases of (H6) in Lemma 7.5(i), also using equations (a) and (b) of Lemma 7.2,
- (H8) in Lemma 7.4(d) and
- (H9) in Lemma 7.6(a).

Each equation shows how to express the given generator as a sum of coboundaries and generators previously expressed in terms of coboundaries: \square

Remark 7.7 The equations above assume that distinct subscripts label distinct elements. This means that Lemma 7.5(a) requires six basis elements. We do not know if the generic (H6) generator (a five-parameter generator) can be expressed as a sum of coboundaries without using a sixth basis element. Therefore we require $n \geq 6$ in the statement and we do not know if the theorem holds for smaller n .

7.3 Second homology of congruence subgroups

In this section, we prove Theorem D, which asserts that $H_2(\mathrm{Aut}(F_n, \ell); \mathbb{Q}) = 0$ for $n \geq 6$ and $\ell \geq 2$. The key to this is the following lemma. Let $\mathrm{GL}_n(\mathbb{Z}, \ell)$ be the level- ℓ congruence subgroup of $\mathrm{GL}_n(\mathbb{Z})$, that is, the kernel of the natural map $\mathrm{GL}_n(\mathbb{Z}) \rightarrow \mathrm{GL}_n(\mathbb{Z}/\ell)$.

Like in Theorem C, we require $n \geq 6$ because of Lemma 7.5(a). We do not know if the result holds for smaller n .

Lemma 7.8 For $n \geq 6$ and $\ell \geq 2$ we have $(H_2(\mathrm{IA}_n; \mathbb{Q}))_{\mathrm{GL}_n(\mathbb{Z}, \ell)} = 0$.

Proof Again we use our generators from Theorem 6.1. The universal coefficient theorem implies that $H_2(\mathrm{IA}_n; \mathbb{Q})$ is generated by the images of our generators from $H_2(\mathrm{IA}_n)$.

We have two approaches for showing that a generator has trivial image. The first is the following: if $f \in \text{Aut}(F_n)$ and $r, s \in H_2(\text{IA}_n)$ with $f \cdot r - r = s$ and $f \cdot s = s$ in $(H_2(\text{IA}_n; \mathbb{Q}))_{\text{GL}_n(\mathbb{Z}, \ell)}$, then

$$f^\ell \cdot r - r = \ell s \quad \text{in } (H_2(\text{IA}_n; \mathbb{Q}))_{\text{GL}_n(\mathbb{Z}, \ell)}.$$

If further f^ℓ lies in $\text{Aut}(F_n, \ell)$ then this shows that s is trivial in $(H_2(\text{IA}_n; \mathbb{Q}))_{\text{GL}_n(\mathbb{Z}, \ell)}$.

The second approach is simpler: if $f \in \text{Aut}(F_n)$ and $r, s \in H_2(\text{IA}_n)$ with $f \cdot r - r = s$ and $r = 0$ in $(H_2(\text{IA}_n; \mathbb{Q}))_{\text{GL}_n(\mathbb{Z}, \ell)}$, then of course, $s = 0$ in $(H_2(\text{IA}_n; \mathbb{Q}))_{\text{GL}_n(\mathbb{Z}, \ell)}$.

We show the generators have trivial images as follows:

- generic (H1) using equations (a) and (b) of Lemma 7.3 by the first approach,
- special cases of (H1) using Lemma 7.3(c) by the second approach, and using Lemma 7.1,
- (H3) by Lemma 7.3(d) using the second approach, and using Lemma 7.1,
- (H2) by equations (e) and (f) of Lemma 7.3, using the second approach,
- (H4) by equations (a) and (b) of Lemma 7.4, using the first approach,
- (H5) by Lemma 7.4(c), by the second approach,
- generic (H6) by equations (a) and (b) of Lemma 7.5, using the first approach,
- generic (H7) by equations (c) and (d) of Lemma 7.5, by the first approach,
- special cases of (H7) by equations (e) and (f), and by equations (g) and (h), of Lemma 7.5, both by the first approach,
- one special case of (H6) by equations (i) and (j) of Lemma 7.5, by the first approach,
- other special cases of (H6) using the first case and equations (a) and (b) of Lemma 7.2,
- (H8) by equations (d) and (e) of Lemma 7.4 by the first approach and
- (H9) by equations (a) and (b) of Lemma 7.6 by the first approach. □

Proof of Theorem D We examine the Hochschild–Serre spectral sequence associated to the short exact sequence

$$(12) \quad 1 \rightarrow \text{IA}_n \rightarrow \text{Aut}(F_n, \ell) \rightarrow \text{GL}_n(\mathbb{Z}, \ell) \rightarrow 1.$$

First, the Borel stability theorem [7] implies that $H_2(\text{GL}_n(\mathbb{Z}, \ell); \mathbb{Q}) = 0$. Next, recall from the introduction that $H_1(\text{IA}_n; \mathbb{Q}) \cong \text{Hom}(\mathbb{Q}^n, \wedge^2 \mathbb{Q}^n)$; the group $\text{GL}_n(\mathbb{Z})$ acts on this in the obvious way. Since $\text{Hom}(\mathbb{R}^n, \wedge^2 \mathbb{R}^n)$ is an irreducible representation of

the algebraic group $\mathrm{SL}_n(\mathbb{R})$, it follows from work of Borel [6, Proposition 3.2] that $H_1(\mathrm{IA}_n; \mathbb{Q})$ is an irreducible representation of $\mathrm{GL}_n(\mathbb{Z}, \ell)$ (we remark that the above reference is one of the steps in the original proof of the Borel density theorem; the result can also be derived directly from the Borel density theorem). It thus follows from the extension of the Borel stability theorem to nontrivial coefficient systems [8] that

$$H_1(\mathrm{GL}_n(\mathbb{Z}, \ell); H_1(\mathrm{IA}_n; \mathbb{Q})) = 0.$$

Lemma 7.8 says that

$$H_0(\mathrm{GL}_n(\mathbb{Z}, \ell); H_2(\mathrm{IA}_n; \mathbb{Q})) \cong (H_2(\mathrm{IA}_n; \mathbb{Q}))_{\mathrm{GL}_n(\mathbb{Z}, \ell)} = 0.$$

The $p + q = 2$ terms of the Hochschild–Serre spectral sequence associated to (12) thus all vanish, so $H_2(\mathrm{Aut}(F_n, \ell); \mathbb{Q}) = 0$, as desired. \square

References

- [1] **S Andreadakis**, *On the automorphisms of free groups and free nilpotent groups*, Proc. London Math. Soc. 15 (1965) 239–268 [MR](#)
- [2] **O Baker**, *The Jacobian map on outer space*, PhD thesis, Cornell University (2011) Available at <https://ecommons.cornell.edu/handle/1813/30769>
- [3] **L Bartholdi**, *Endomorphic presentations of branch groups*, J. Algebra 268 (2003) 419–443 [MR](#)
- [4] **M Bestvina**, **K-U Bux**, **D Margalit**, *Dimension of the Torelli group for $\mathrm{Out}(F_n)$* , Invent. Math. 170 (2007) 1–32 [MR](#)
- [5] **SK Boldsen**, **M Hauge Døllerup**, *Towards representation stability for the second homology of the Torelli group*, Geom. Topol. 16 (2012) 1725–1765 [MR](#)
- [6] **A Borel**, *Density properties for certain subgroups of semi-simple groups without compact components*, Ann. of Math. 72 (1960) 179–188 [MR](#)
- [7] **A Borel**, *Stable real cohomology of arithmetic groups*, Ann. Sci. École Norm. Sup. 7 (1974) 235–272 [MR](#)
- [8] **A Borel**, *Stable real cohomology of arithmetic groups, II*, from “Manifolds and Lie groups” (S Murakami, J Hano, K Okamoto, A Morimoto, H Ozeki, editors), Progr. Math. 14, Birkhäuser, Boston (1981) 21–55 [MR](#)
- [9] **T Brendle**, **D Margalit**, **A Putman**, *Generators for the hyperelliptic Torelli group and the kernel of the Burau representation at $t = -1$* , Invent. Math. 200 (2015) 263–310 [MR](#)
- [10] **KS Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer (1994) [MR](#)

- [11] **T Church, B Farb**, *Parameterized Abel–Jacobi maps and abelian cycles in the Torelli group*, *J. Topol.* 5 (2012) 15–38 [MR](#)
- [12] **T Church, B Farb**, *Representation theory and homological stability*, *Adv. Math.* 245 (2013) 250–314 [MR](#)
- [13] **T Church, A Putman**, *The codimension-one cohomology of $\text{SL}_n\mathbb{Z}$* , *Geom. Topol.* 21 (2017) 999–1032 [MR](#)
- [14] **F Cohen, J Pakianathan**, *On automorphism groups of free groups and their nilpotent quotients*, in preparation
- [15] **M Day, A Putman**, *A Birman exact sequence for $\text{Aut}(F_n)$* , *Adv. Math.* 231 (2012) 243–275 [MR](#)
- [16] **M Day, A Putman**, *The complex of partial bases for F_n and finite generation of the Torelli subgroup of $\text{Aut}(F_n)$* , *Geom. Dedicata* 164 (2013) 139–153 [MR](#)
- [17] **M Day, A Putman**, *A Birman exact sequence for the Torelli subgroup of $\text{Aut}(F_n)$* , *Internat. J. Algebra Comput.* 26 (2016) 585–617 [MR](#)
- [18] **B Farb**, *Automorphisms of F_n which act trivially on homology*, in preparation
- [19] **S Galatius**, *Stable homology of automorphism groups of free groups*, *Ann. of Math.* 173 (2011) 705–768 [MR](#)
- [20] **A Hatcher, K Vogtmann**, *Cerf theory for graphs*, *J. London Math. Soc.* 58 (1998) 633–655 [MR](#)
- [21] **A Hatcher, N Wahl**, *Stabilization for mapping class groups of 3–manifolds*, *Duke Math. J.* 155 (2010) 205–269 [MR](#)
- [22] **D Johnson**, *An abelian quotient of the mapping class group \mathcal{I}_g* , *Math. Ann.* 249 (1980) 225–242 [MR](#)
- [23] **N Kawazumi**, *Cohomological aspects of Magnus expansions*, preprint (2005) [arXiv](#)
- [24] **S Krstić, J McCool**, *The non-finite presentability of $\text{IA}(F_3)$ and $\text{GL}_2(\mathbb{Z}[t, t^{-1}])$* , *Invent. Math.* 129 (1997) 595–606 [MR](#)
- [25] **R C Lyndon, P E Schupp**, *Combinatorial group theory*, *Ergeb. Math. Grenzgeb.* 89, Springer (1977) [MR](#)
- [26] **W Magnus**, *Über n –dimensionale Gittertransformationen*, *Acta Math.* 64 (1935) 353–367 [MR](#)
- [27] **J Nielsen**, *Die Isomorphismen der allgemeinen, unendlichen Gruppe mit zwei Erzeugenden*, *Math. Ann.* 78 (1917) 385–397 [MR](#)
- [28] **A Putman**, *An infinite presentation of the Torelli group*, *Geom. Funct. Anal.* 19 (2009) 591–643 [MR](#)
- [29] **A Putman**, *Obtaining presentations from group actions without making choices*, *Algebr. Geom. Topol.* 11 (2011) 1737–1766 [MR](#)

- [30] **A Putman**, *The second rational homology group of the moduli space of curves with level structures*, Adv. Math. 229 (2012) 1205–1234 [MR](#)
- [31] **T Satoh**, *The abelianization of the congruence IA -automorphism group of a free group*, Math. Proc. Cambridge Philos. Soc. 142 (2007) 239–248 [MR](#)
- [32] **T Satoh**, *A survey of the Johnson homomorphisms of the automorphism groups of free groups and related topics*, preprint (2012) [arXiv](#)

*Department of Mathematical Sciences, University of Arkansas
Fayetteville, AR, United States*

*Department of Mathematics, University of Notre Dame
Notre Dame, IN, United States*

matthewd@uark.edu, andyp@nd.edu

Proposed: Shigeyuki Morita

Received: 22 October 2015

Seconded: Benson Farb, Mark Behrens

Revised: 8 November 2016