

Dynamics on flag manifolds: domains of proper discontinuity and cocompactness

MICHAEL KAPOVICH
BERNHARD LEEB
JOAN PORTI

For noncompact semisimple Lie groups G with finite center, we study the dynamics of the actions of their discrete subgroups $\Gamma < G$ on the associated partial flag manifolds G/P . Our study is based on the observation, already made in the deep work of Benoist, that they exhibit also in higher rank a certain form of convergence-type dynamics. We identify geometrically domains of proper discontinuity in all partial flag manifolds. Under certain dynamical assumptions equivalent to the Anosov subgroup condition, we establish the cocompactness of the Γ -action on various domains of proper discontinuity, in particular on domains in the full flag manifold G/B . In the regular case (eg of B -Anosov subgroups), we prove the nonemptiness of such domains if G has (locally) at least one noncompact simple factor not of the type A_1 , B_2 or G_2 by showing the nonexistence of certain ball packings of the visual boundary.

53C35; 22E40, 37B05, 51E24

To Guiomar

1 Introduction

Let G be a noncompact semisimple Lie group with finite center. In this paper, we study the natural actions

$$\Gamma \curvearrowright G/P$$

of discrete subgroups $\Gamma < G$ on the (partial) flag manifolds G/P associated to G . (Here, $P < G$ denotes a parabolic subgroup.) We are interested in aspects of the topological dynamics of the action of Γ in the higher-rank case, notably in domains of proper discontinuity and criteria for the cocompactness of the Γ -action on these.

The study of such questions has a long history. Kulkarni [27] introduced several notions of limit sets for discrete group actions on general topological spaces and proved

proper discontinuity away from the largest of these limit sets. Guivarc'h [12] was the first to define limit sets for discrete group actions on the flag manifolds of $SL(n, \mathbb{R})$. Benoist's [1] foundational work included a definition of limit sets in flag manifolds of arbitrary reductive Lie groups and a notion of contraction dynamics equivalent to the one used in this paper; it was of major influence on further developments in the field and basic for our paper. The work of Karlsson [25] and Papasoglu and Swenson [34] introduced a notion of higher-rank convergence behavior, similar to the one used in this paper, for discrete group actions on visual boundaries in the general setting of CAT(0) spaces. In his pioneering paper [28], Labourie introduced Anosov subgroups and studied their dynamics on flag manifolds of $SL(n, \mathbb{R})$. Guichard and Wienhard [11] constructed cocompact domains of proper discontinuity for Anosov subgroups of various semisimple Lie groups acting on various flag manifolds. Cano, Navarrete and Seade [6] extensively studied Kulkarni's limit sets in the case of discrete subgroups of $SL(n, \mathbb{C})$ acting on $\mathbb{C}P^{n-1}$.

Our approach to studying the topological dynamics of Γ relies on the geometry of the associated symmetric space $X = G/K$ of noncompact type. The connection is established by the fact that the flag manifolds occur as the G -orbits in the visual boundary $\partial_\infty X$, that is, the boundary at infinity of the visual compactification $\bar{X} = X \sqcup \partial_\infty X$ of X .

If $\text{rank}(X) = 1$, or equivalently, if X has strictly negative sectional curvature, then the only flag manifold is $\partial_\infty X$ itself, and the transitive action $G \curvearrowright \partial_\infty X$ has *convergence dynamics*. This means that divergent sequences in G exhibit a certain attraction-repulsion behavior, namely they subconverge on the complement of one point in $\partial_\infty X$ locally uniformly to a constant map. More precisely, for a sequence $g_n \rightarrow \infty$ in G , there exist a subsequence (g_{n_k}) and (not necessarily distinct) points $\xi_\pm \in \partial_\infty X$ such that

$$g_{n_k} |_{\partial_\infty X - \{\xi_-\}} \rightarrow \xi_+$$

uniformly on compacta.

As a consequence of convergence dynamics, for a discrete subgroup $\Gamma < G$, there is a clean Γ -invariant *dynamical decomposition*

$$\partial_\infty X = \Omega_{\text{disc}} \sqcup \Lambda$$

into the open *domain of discontinuity* or *wandering set* Ω_{disc} and the compact *limit set* Λ . The latter consists of all points occurring as limits ξ_+ as above for sequences $\gamma_n \rightarrow \infty$ in Γ , and the Γ -action on Ω_{disc} is even *properly discontinuous*. In order for

this action to be cocompact, one needs to impose further conditions on the group. The action $\Gamma \curvearrowright \Omega_{\text{disc}}$ is *cocompact* if (but not only if) Γ is *convex-cocompact*.

If $\text{rank}(X) \geq 2$, then the action $G \curvearrowright \partial_\infty X$ is no longer transitive. The G -orbits are compact and, as G -spaces, copies of flag manifolds. They are parametrized by the spherical Weyl chamber σ_{mod} associated to G ; a G -orbit for an interior point of a face $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ (which we will also refer to as a *face type*) is naturally identified with the flag manifold $\text{Flag}_{\tau_{\text{mod}}} \cong G/P_\tau$, the conjugacy class of the parabolic subgroup P_τ corresponding to the face τ_{mod} . In particular, the *regular* G -orbits in $\partial_\infty X$, ie those corresponding to interior points of σ_{mod} itself, are identified with the full flag manifold, $\partial_{\text{Fü}} X \cong \text{Flag}_{\sigma_{\text{mod}}} \cong G/B_\sigma$, the space of Weyl chambers at infinity, also called the Furstenberg boundary; here, B_σ denotes a minimal parabolic subgroup of G .

Our study is based on the observation that a *weak* form of *convergence dynamics* persists for the action $G \curvearrowright \partial_\infty X$ in higher rank (see Sections 5.2 and 6): sequences $g_n \rightarrow \infty$ in G *subaccumulate* outside a compact exceptional subset locally uniformly at another compact subset, meaning that there exist a subsequence (g_{n_k}) and compact subsets $A_\pm \subset \partial_\infty X$ such that

$$g_{n_k} |_{\partial_\infty X - A_-} \text{ accumulates at } A_+$$

uniformly on compacta. We briefly say that (g_{n_k}) is (A_-, A_+) -*accumulating* in this case. There is a certain flexibility in the choice of the pair of compact subsets (A_-, A_+) and a trade-off (“uncertainty relation”): if one shrinks one of the subsets A_\pm , one must enlarge the other.

For instance, one can make the following *metric* choice for the pair of compact subsets: for a suitable subsequence (g_{n_k}) , there exist points $\xi_\pm \in \partial_\infty X$ such that (g_{n_k}) is $(\bar{B}(\xi_-, \pi - r), \bar{B}(\xi_+, r))$ -accumulating for all radii $r \in (0, \pi)$, where $B(\xi, r)$ denotes a ball in $\partial_\infty X$ with respect to the Tits angle metric. This kind of convergence type dynamical behavior had been observed, in the general setting of proper CAT(0) spaces, by Karlsson [25, Theorem 1] and Papasoglu and Swenson [34, Theorem 4]; see also the first version of this paper [18, Theorem 1.1 and Section 6.1].

In our setting of CAT(0) *model* spaces with their rich geometric structure, one can make more flexible “combinatorial” choices for the pair of compact subsets which can be described in terms of the (partial) Bruhat order $<$ on the Weyl group W . These will enable us to construct *larger* domains of proper discontinuity for discrete group actions than those obtained from metric choices. To explain the combinatorial choices, we need some preliminary considerations.

It will be useful for us to interpret the Bruhat order geometrically, and we give a geometric description of it and its generalization as the *folding order*; see Section 3.2.

Since any two Weyl chambers at infinity $\sigma, \sigma' \subset \partial_\infty X$ are contained in an apartment, that is, the visual boundary $\partial_\infty F \subset \partial_\infty X$ of a maximal flat $F \subset X$, we can define a combinatorial *relative position*

$$\text{pos}(\sigma', \sigma) \in W$$

of σ' with respect to σ ; see Section 3.3. The larger the position is with respect to \prec , the more generic it is. We note that the sublevels of $\text{pos}(\cdot, \sigma)$ in $\partial_{\text{Fü}} X$ are precisely the Schubert cycles with respect to σ , that is, the B_σ -orbit closures.

We define a *thickening*

$$\text{Th} \subset W$$

of the neutral element inside the Weyl group as a union of sublevels for the Bruhat order; see Section 3.4. Each thickening $\text{Th} \subset W$ gives rise to corresponding thickenings of chambers inside the Furstenberg boundary,

$$\text{Th}_{\text{Fü}}(\sigma) := \{\text{pos}(\cdot, \sigma) \in \text{Th}\} \subset \partial_{\text{Fü}} X,$$

which can also be regarded as thickenings $\text{Th}(\sigma) \subset \partial_\infty X$ inside the visual boundary by taking the union of the chambers contained in them. The thickenings of chambers in $\partial_{\text{Fü}} X$ are finite unions of Schubert cycles and hence projective subvarieties. Thickenings $\text{Th}_{\text{Fü}}(A) \subset \partial_{\text{Fü}} X$ and $\text{Th}(A) \subset \partial_\infty X$ of compact subsets $A \subset \partial_{\text{Fü}} X$ are defined as the union of the thickenings of the individual chambers $\sigma \in A$; they are again compact.

This discussion generalizes: There is a well-defined $W_{\tau_{\text{mod}}} \backslash W$ -valued position $\text{pos}(\cdot, \tau)$ of chambers relative to a *simplex* $\tau \in \text{Flag}_{\tau_{\text{mod}}}$. Here, $W_{\tau_{\text{mod}}} < W$ denotes the stabilizer of the face $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$. If the thickening $\text{Th} \subset W$ is $W_{\tau_{\text{mod}}}$ -left invariant, then it yields well-defined compact thickenings $\text{Th}_{\text{Fü}}(A) \subset \partial_{\text{Fü}} X$ and $\text{Th}(A) \subset \partial_\infty X$ of compact subsets $A \subset \text{Flag}_{\tau_{\text{mod}}}$. Even more generally, there is a well-defined $W_{\tau_{\text{mod}}} \backslash W / W_{\nu_{\text{mod}}}$ -valued position $\text{pos}(\nu, \tau)$ of *simplices* $\nu \in \text{Flag}_{\nu_{\text{mod}}}$ relative to simplices $\tau \in \text{Flag}_{\tau_{\text{mod}}}$, and a $W_{\tau_{\text{mod}}}$ -left and $W_{\nu_{\text{mod}}}$ -right invariant thickening inside W yields thickenings of subsets of $\text{Flag}_{\tau_{\text{mod}}}$ inside $\text{Flag}_{\nu_{\text{mod}}}$.

For each thickening $\text{Th} \subset W$, there is the *complementary* thickening $\text{Th}^c \subset W$ defined by

$$W = \text{Th} \sqcup w_0 \text{Th}^c,$$

where w_0 denotes the longest element of the Weyl group. (This partition of the Weyl group generalizes the decomposition $\pi = r + (\pi - r)$ of the maximal distance in the unit sphere; compare the metric choices above.) We call a thickening Th *slim*, *fat* or *balanced* if $\text{Th} \subseteq \text{Th}^c$, $\text{Th} \supseteq \text{Th}^c$, or $\text{Th} = \text{Th}^c$, respectively. Existence results for balanced thickenings with different invariance properties are stated in Proposition 1.9 below; examples are given in Section 3.4.

Returning to the dynamics of sequences $g_n \rightarrow \infty$ in G on $\partial_\infty X$, we show (see Lemma 4.5 and Proposition 6.9) that there always exist a subsequence (g_{n_k}) , a face $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ and a pair of simplices $\tau_\pm \in \text{Flag}_{\pm\tau_{\text{mod}}}$ such that

$$(1) \quad g_{n_k} |_{C(\tau_-)} \rightarrow \tau_+$$

uniformly on compacta in the open Schubert cell $C(\tau_-) \subset \text{Flag}_{\tau_{\text{mod}}}$ of simplices opposite to τ_- . Here $-\tau_{\text{mod}} := \iota\tau_{\text{mod}}$ for the canonical involution $\iota = -w_0$ of σ_{mod} . This locally uniform convergence property implies (see Corollary 6.3) that, more generally,

$$(2) \quad (g_{n_k}) \text{ is } (\text{Th}^c(\tau_-), \text{Th}(\tau_+))\text{-accumulating on } \partial_\infty X$$

for all $W_{\tau_{\text{mod}}}$ -left invariant thickenings $\text{Th} \subset W$. Note that (1) is equivalent to (2) for the slimmest nonempty choice $\text{Th} = \{e\}$. We call sequences satisfying (1) τ_{mod} -contracting; see Definition 6.1. An equivalent notion had already been introduced by Benoist in his fundamental work [1]; see in particular part (5) of his Lemma 3.5, and compare also Remark 1.3.

We now turn to discussing the *dynamics of discrete subgroups* $\Gamma < G$ on $\partial_\infty X$.

For a face $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we define the “small” τ_{mod} -limit set

$$\Lambda_{\tau_{\text{mod}}} \subset \text{Flag}_{\tau_{\text{mod}}}$$

as the set of all simplices in $\text{Flag}_{\tau_{\text{mod}}}$ which occur as limits τ_+ as in (1) for sequences $\gamma_n \rightarrow \infty$ in Γ ; see Definition 6.4.

Remark 1.1 In his influential paper, Benoist [1, Section 3.6] introduced a notion of limit set Λ_Γ for Zariski dense subgroups Γ of reductive algebraic groups over local fields which, in the case of real semisimple Lie groups, is equivalent to our concept of σ_{mod} -limit set $\Lambda_{\sigma_{\text{mod}}}$.¹ What we call the τ_{mod} -limit set $\Lambda_{\tau_{\text{mod}}}$ for other face types $\tau_{\text{mod}} \subsetneq \sigma_{\text{mod}}$ is mentioned in his Remark 3.6(3), and his work implies that, in the Zariski

¹Benoist’s limit set Λ_Γ is contained in the flag manifold Y_Γ , which, in the case of real Lie groups, is the full flag manifold G/B ; see the beginning of Section 3 of his paper. It consists of the limit points of sequences contracting on G/B ; see his Definitions 3.5 and 3.6.

dense case, $\Lambda_{\tau_{\text{mod}}}$ is the image of $\Lambda_{\sigma_{\text{mod}}}$ under the natural projection $\text{Flag}_{\sigma_{\text{mod}}} \rightarrow \text{Flag}_{\tau_{\text{mod}}}$ of flag manifolds.

By choosing a $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, we obtain from these small limit sets the “large” *thickened* limit sets $\text{Th}(\Lambda_{\tau_{\text{mod}}}) \subset \partial_{\infty} X$. Our main results concern the proper discontinuity and cocompactness of the Γ -action on the complements

$$\Omega_{\text{Th}} := \partial_{\infty} X - \text{Th}(\Lambda_{\tau_{\text{mod}}}),$$

respectively, on their intersections with the G -orbits $G\eta \subset \partial_{\infty} X$. We obtain the strongest results for the dynamics on the Furstenberg boundary. This is reasonable because the latter fibers with compact fiber over all partial flag manifolds, and cocompact domains of proper discontinuity in any flag manifold pull back to such domains in $\partial_{\text{Fü}} X$. Our results are of the kind, in the spirit of Mumford’s *geometric invariant theory*, that the Γ -actions become properly discontinuous when removing a sufficiently “fat” thickening of $\Lambda_{\tau_{\text{mod}}}$, and remain cocompact when removing a sufficiently “slim” one. (See Example 3.30 and the discussion in Section 7.4 for a concrete connection with configuration spaces and GIT.)

The accumulation property (2) is the key step in constructing *domains of proper discontinuity* for all discrete subgroups; see Propositions 6.15 and 6.16 in Section 6.4. We obtain the most useful results for subgroups which satisfy a certain generalization of the convergence property (see Definition 6.5):

Definition 1.2 (weak convergence subgroup) We call a discrete subgroup $\Gamma < G$ a τ_{mod} -convergence subgroup with respect to a face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, if every sequence $\gamma_n \rightarrow \infty$ in Γ has a subsequence satisfying (1) with this particular face type τ_{mod} , or equivalently, has a subsequence satisfying (2) for any choice of $W_{\tau_{\text{mod}}}$ -left invariant thickenings $\text{Th} \subset W$.

Remark 1.3 (convergence dynamics versus regularity) We note that the τ_{mod} -convergence property of a subgroup $\Gamma < G$, formulated in terms of the dynamics of the action on the visual boundary $\partial_{\infty} X$, can be equivalently described in terms of the asymptotic behavior of Γ -orbits in the symmetric space X . Namely, Γ is a τ_{mod} -convergence subgroup if and only if it is a τ_{mod} -regular subgroup of G ; see Section 6.3. The notion of τ_{mod} -regularity was introduced in our earlier paper [19], where also the equivalence of the two notions was established. In the present paper, we only need (and verify) that τ_{mod} -convergence implies τ_{mod} -regularity.

The notions of regularity and contraction for sequences and their essential equivalence can be found already in the fundamental work of Benoist [1, Section 3]. For the sake of completeness, we give independent proofs in our setting of discrete subgroups of semisimple Lie groups. Also our methods are rather different. We give a geometric treatment and present the material in a form suitable for the further development of our theory of discrete isometry groups acting on Riemannian symmetric spaces and euclidean buildings, such as in our papers [16; 20].

Our main result on proper discontinuity includes (compare Theorem 6.8):

Theorem 1.4 (proper discontinuity outside fat thickenings) *Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be ι -invariant. If $\Gamma < G$ is a τ_{mod} -convergence subgroup, then for any fat $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_{\infty} X - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous.

In order to obtain *cocompactness* for actions of τ_{mod} -convergence subgroups, we must impose further conditions, as it is the case for convergence actions; compare the situation in rank one. Our main requirement is that the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ should be *expanding at $\Lambda_{\tau_{\text{mod}}}$* in the sense of Sullivan [37, Section 9]; see Definition 5.19. Moreover, if τ_{mod} is ι -invariant, we call the limit set $\Lambda_{\tau_{\text{mod}}}$ *antipodal* if the simplices in it are pairwise opposite; see Definition 2.3(ii).

Definition 1.5 (CEA subgroup) For an ι -invariant face $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we call a τ_{mod} -convergence subgroup $\Gamma < G$ a τ_{mod} -CEA subgroup (convergence, expanding, antipodal) if $\Lambda_{\tau_{\text{mod}}}$ is antipodal and if the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ is expanding at $\Lambda_{\tau_{\text{mod}}}$.

The restricted action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}$ is then a convergence action in the traditional sense. Such subgroups are higher-rank generalizations of convex-cocompact subgroups of rank-one Lie groups. In fact, the CEA condition is only one of various equivalent dynamical and (coarse) geometric conditions which can be used to characterize this class of discrete subgroups; see our papers [15; 19; 21; 22] for a detailed study of these conditions and their equivalence. In particular:

Remark 1.6 (CEA versus Anosov) The class of τ_{mod} -CEA subgroups coincides with the class of $P_{\tau_{\text{mod}}}$ -Anosov subgroups; see [19, Section 6.5]. Here, $P_{\tau_{\text{mod}}}$ refers to

the conjugacy class of parabolic subgroups of G corresponding to the face τ_{mod} of the spherical Weyl chamber σ_{mod} . We recall that the notion of Anosov subgroup had first been introduced in [28] using the language of geodesic flows, and further extended in [11]. We gave the first flow-free definitions in [19]. We note that Labourie's original definition did not require τ_{mod} to be ι -invariant; instead, he worked with $(P_{\tau_{\text{mod}}}, P_{\iota\tau_{\text{mod}}})$ -Anosov subgroups. However, as already observed in [11], the general case readily reduces to the ι -invariant one.

Our main result regarding cocompactness includes the following (compare Theorem 7.7 and Corollary 7.8):

Theorem 1.7 (cocompactness outside slim thickenings) *Suppose $\Gamma < G$ is a τ_{mod} -CEA subgroup. Then for each slim $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\tau_{\text{mod}}})$$

is cocompact.

More generally, suppose that $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$ is another face type and that the thickening Th is also $W_{\nu_{\text{mod}}}$ -right invariant. Then for any G -orbit $G\eta \subset \partial_{\infty} X$ corresponding to an interior point of ν_{mod} , the action

$$\Gamma \curvearrowright G\eta - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is cocompact.

By combining the two theorems, we obtain the central result of this paper:

Theorem 1.8 (cocompact domains of proper discontinuity) *Let $\Gamma < G$ and the data $\tau_{\text{mod}}, \nu_{\text{mod}}, \text{Th}$ be as in the previous theorem with the additional requirement that the thickening Th be balanced. Then the respective actions are properly discontinuous and cocompact.*

We note that topology of the quotient space $(\partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\tau_{\text{mod}}})) / \Gamma$, in general, depends on the balanced thickening; see Example 7.14.

Balanced thickenings do not exist for all invariance requirements, but for many, they do. For instance, one can impose arbitrary left invariance, and as a consequence, one has balanced thickenings of τ_{mod} -limits sets inside $\partial_{\text{Fü}} X$ for all ι -invariant τ_{mod} , as the first part of the next result shows (see Section 3.4 for more general results):

Proposition 1.9 (existence of balanced thickenings) *For every ι -invariant face type τ_{mod} , there exists a $W_{\tau_{\text{mod}}}$ -left invariant balanced thickening $\text{Th} \subset W$.*

For an arbitrary face type ν_{mod} , a $W_{\nu_{\text{mod}}}$ -right invariant balanced thickening exists if and only if left multiplication by w_0 has no fixed point on $W/W_{\nu_{\text{mod}}}$. This is the case, for instance, if $w_0 = -\text{id}$, or equivalently, if all irreducible factors of the symmetric space are of type $A_1, B_{n \geq 2}, D_{2k \geq 4}, E_{7,8}, F_4$ or G_2 .

The *nonemptiness* of the domains found in Theorem 1.8 (and Theorem 1.4) is an issue. For instance, uniform lattices in rank-one Lie groups have empty domains of discontinuity at infinity (and such lattices are CEA). See also Example 7.14 for empty domains in the reducible case. If, for a τ_{mod} -convergence subgroup Γ with antipodal τ_{mod} -limit set, all domains given by Theorem 1.4 were empty, it would follow that the visual boundary of X admits a *packing* by a compact family (with respect to the visual topology) of $\frac{\pi}{2}$ -balls (with respect to the Tits metric); see Proposition 8.12. However, the existence of such packings can be ruled out for most Weyl groups (see Theorem 8.8), and we conclude (see Theorem 8.13):

Theorem 1.10 (nonemptiness of domains of proper discontinuity) *Suppose that X has at least one de Rham factor not of the type A_1, B_2 or G_2 , and let $\Gamma < G$ be a σ_{mod} -convergence subgroup with antipodal limit set $\Lambda_{\sigma_{\text{mod}}}$. Then for some balanced thickening $\text{Th} \subset W$, the domain of proper discontinuity $\partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\sigma_{\text{mod}}})$ for the Γ -action provided by Theorem 1.4 is nonempty.*

Note that the theorem covers the case of CEA subgroups, but is more general.

The possible balanced thickenings can be described more precisely; see Theorem 8.13. In the B_2 -case, we have partial nonemptiness results for the groups $G = O(2k + 1, 2)$ with $k \geq 1$ (see Addendum 8.14). The G_2 -case is not discussed in this paper.

The above results yield the following for the dynamics of CEA subgroups on the Furstenberg boundary:

Corollary 1.11 (dynamics on the Furstenberg boundary) *Suppose that $\Gamma < G$ is a τ_{mod} -CEA subgroup. There exist $W_{\tau_{\text{mod}}}$ -left invariant balanced thickenings $\text{Th} \subset W$, and for every such thickening, the action*

$$\Gamma \curvearrowright \partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous and cocompact.

If $\Gamma < G$ is a σ_{mod} -CEA subgroup, and if X has at least one de Rham factor not of the type A_1 , B_2 or G_2 , then for some balanced thickening $\text{Th} \subset W$, the cocompact domain of proper discontinuity $\partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\sigma_{\text{mod}}})$ is nonempty.

Again, the possible thickenings occurring in the σ_{mod} -case can be described more precisely.

Remark 1.12 (dynamics on Finsler compactifications) Our results regarding domains of proper discontinuity and cocompactness for discrete group actions on flag manifolds have analogs for the actions of the same classes of subgroups on a *Finsler compactification* \bar{X}^{Fins} of X . This is done in our paper [16]. The compactification \bar{X}^{Fins} is obtained from X geometrically by applying the horoboundary construction to suitable G -invariant *regular polyhedral Finsler* metrics on X rather than to G -invariant Riemannian metrics (which yields the visual compactification $\bar{X} = X \sqcup \partial_{\infty} X$), and it coincides with the maximal Satake compactification from algebraic group theory; see also Parreau [35]. Note that the Furstenberg boundary $\partial_{\text{Fü}} X$ naturally embeds into \bar{X}^{Fins} as a G -orbit, namely as the only compact one. Some of the results become easier in the Finsler setting; for instance, the nonemptiness of domains of proper discontinuity at infinity is no longer an issue. Each τ_{mod} -convergence subgroup with antipodal limit set $\Lambda_{\tau_{\text{mod}}}$ has a nonempty domain of proper discontinuity in the Finsler ideal boundary (defined using an arbitrary $W_{\tau_{\text{mod}}}$ -left invariant balanced thickening), once $\text{rank}(X) \geq 2$; see [16, Lemma 9.19].

Remark 1.13 There is overlap of our results with [11]. There, cocompact domains of proper discontinuity are constructed for Anosov subgroups of various semisimple Lie groups acting on various partial flag manifolds. However, in the general case of arbitrary semisimple Lie groups G , such domains are constructed only in G -homogeneous spaces fibering over $\partial_{\text{Fü}} X \cong G/B$ with compact fiber [11, Theorem 1.9]. Nonemptiness of these domains is proven for P -Anosov subgroups of small cohomological dimension [11, Theorems 1.11, 1.12 and 9.10], while our nonemptiness results apply to σ_{mod} -convergence subgroups with antipodal σ_{mod} -limit set (this class of subgroups includes B -Anosov subgroups), without restriction on the cohomological dimension.

Observe also that our treatment is intrinsic, while in [11], a theory for Anosov subgroups of Lie groups of the type $\text{Aut}(F)$ is first developed (where the F 's are certain bilinear and hermitian forms), and then generalized to other semisimple Lie groups by embedding these into the groups of type $\text{Aut}(F)$. The intrinsic approach is more uniform and seems to provide better control; eg it allows us to get the domains, for

general semisimple Lie groups, in flag manifolds instead of only in bundles over these as in [11]. While in some low-rank cases the outcomes of the two constructions of thickened limit sets are the same, our construction appears to be more general.

The earlier version [18] of this paper written in 2013 covered only the σ_{mod} -regular case. Some of the material of [18], dealing with equivalent characterizations of τ_{mod} -CEA actions, was later moved to our paper [19]. Most of the rest of the material of [18] was generalized and moved into this paper.

Acknowledgements We thank the referee for deepening our understanding of Benoist’s foundational work [1]. Kapovich was supported by the NSF grants DMS-09-05802, DMS-12-05312 and DMS-16-04241. Porti was supported by the grants Mineco MTM2012-34834, AGAUR SGR2009-1207 and the Icrea Acadèmia Award 2008. The authors are also grateful to the GEAR grant which partially supported the IHP trimester in Winter of 2012 (DMS 1107452, 1107263, 1107367 “RNMS: Geometric structures and representation varieties” (the GEAR Network)), and the Max Planck Institute for Mathematics in Bonn, where some of this work was done.

2 Geometric preliminaries

In this section, we collect some standard material on Coxeter complexes, the geometry of nonpositively curved symmetric spaces and associated spherical Tits buildings; we refer the reader to [26; 29] for more detailed discussion of symmetric spaces and buildings.

2.1 General notation

We will use the notation $B(a, r)$ and $\bar{B}(a, r)$ for the open and closed r -balls, respectively, centered at a in a metric space Z . We will denote the nearest point distance of a point $z \in Z$ to a subset $A \subset Z$ by $d(z, A) := \inf d(z, \cdot)|_A$. The Hausdorff distance between two subsets $A, B \subset Z$ will be denoted by $d_H(A, B)$. A *geodesic* in a metric space is an isometric embedding from a (possibly infinite) interval $I \subset \mathbb{R}$.

2.2 Coxeter complexes

A *spherical Coxeter complex* a_{mod} is a pair (S, W) consisting of a unit sphere S in a euclidean vector space V and a finite group W which acts isometrically on S and is generated by reflections at hyperplanes. A Coxeter complex is *reducible* if W splits

as a (nontrivial) direct product $W_1 \times W_2$ and V admits a W -invariant (nontrivial) orthogonal direct sum decomposition $V = V_1 \oplus V_2$ such that W_i fixes V_{3-i} for $i = 1, 2$. In this case, we obtain two induced Coxeter complexes (S_i, W_i) on the unit spheres $S_i \subset V_i$. A Coxeter complex which is not reducible is called *irreducible*.

We will use the notation \angle for the angular metric on S . Throughout the paper, we assume that W does not fix a point in S and is associated with a root system R . Spherical Coxeter complexes will occur as *model apartments* of spherical buildings, mostly of Tits boundaries of symmetric spaces, and will in this context usually be denoted by a_{mod} .

A wall m_ρ in S is the fixed point set of a hyperplane reflection ρ in W . A *half-apartment* in S is a closed hemisphere bounded by a wall. A point $\xi \in S$ is called *singular* if it belongs to a wall and *regular* otherwise.

The action $W \curvearrowright S$ determines on S a structure as a simplicial complex whose facets, called *chambers*, are the closures of the connected components of

$$S - \bigcup_{\rho} m_{\rho},$$

where the union is taken over all reflections ρ in W . We will refer to the simplices in this complex as *faces*. (If one allows fixed points for W on S , then S carries only a structure as a cell complex.) Codimension-one faces of this complex are called *panels*. The *interior* $\text{int}(\tau)$ of a face τ is the complement in τ to the union of walls not containing τ . The interiors $\text{int}(\tau)$ are called *open simplices*. A geodesic sphere in S is called *singular* if it is simplicial, equivalently, if it equals an intersection of walls.

Each chamber is a fundamental domain for the action $W \curvearrowright S$. We define the *spherical model Weyl chamber* as the quotient $\sigma_{\text{mod}} = S/W$. The natural projection $\theta: S \rightarrow \sigma_{\text{mod}}$ restricts to an isometry on every chamber. An important elementary property of the chamber σ_{mod} is that its diameter (with respect to the spherical metric) is $\leq \frac{\pi}{2}$.

For a face τ_{mod} of σ_{mod} , we define the subgroup $W_{\tau_{\text{mod}}} \subset W$ as the stabilizer of τ_{mod} in W . Accordingly, for a point $\bar{\xi} \in \sigma_{\text{mod}}$, we define $W_{\bar{\xi}} \subset W$ as the stabilizer of $\bar{\xi}$ in W . Then $W_{\bar{\xi}} = W_{\tau_{\text{mod}}}$, where τ_{mod} is the face of σ_{mod} spanned by $\bar{\xi}$, ie which contains $\bar{\xi}$ as an interior point. Note that $W_{\sigma_{\text{mod}}} = 1$ and $W_{\bar{\xi}} = 1$ for $\bar{\xi} \in \text{int}(\sigma_{\text{mod}})$.

It is convenient, and we will frequently do so, to identify σ_{mod} with a chamber $\sigma \subset S$ (traditionally called the *positive chamber*). Such an identification determines a generating set of W , namely the reflections at the walls bounding σ_{mod} , and hence a word

metric on W ; the longest element with respect to this metric is denoted by w_0 . This element sends σ_{mod} to the opposite chamber in S . We say that two points $\xi, \hat{\xi} \in S$ are *Weyl antipodes* if $\hat{\xi} = w_0\xi$. We define the *standard* or *opposition involution*

$$\iota = \iota_S: S \rightarrow S$$

as the composition $-w_0$. This involution preserves σ_{mod} and equals the identity if and only if $-\text{id}_S \in W$, because then $w_0 = -\text{id}_S$.

A point ξ in S is called a *root* if the hemisphere centered at ξ is simplicial, equivalently, is bounded by a wall. If (S, W) is associated with a root system R , then $\xi \in S$ is a root if and only if it has the direction of a *coroot*. Note that irreducible root systems correspond to irreducible Coxeter complexes and vice versa.

Remark 2.1 We will be assuming in what follows that (S, W) is associated with a root system R which spans V^* . Equivalently, W is isomorphic to the linear part of an affine *crystallographic* Coxeter group, ie one acting cocompactly on the affine space underlying the vector space V . The root system R in this situation can be assumed to be *reduced*; ie if roots α, β have the same kernel, then $\alpha = \pm\beta$. In what follows, we will be assuming that R is reduced.

Note that each root type $\bar{\zeta} \in \sigma_{\text{mod}}$ is ι -invariant, since the reflection $w \in W$ corresponding to the root $\bar{\zeta}$ sends $\bar{\zeta}$ to $-\bar{\zeta}$.

Each *irreducible* root system R has one or two distinct root types; ie W acts on R with one or two orbits. Geometrically speaking, this means that W acts on the set of walls with one or two orbits. We refer the reader to [4] for details.

Suppose that S is identified with the sphere at infinity of a euclidean space F , that is, $S \cong \partial_\infty F$, where $\partial_\infty F$ is equipped with the angular metric. For a closed subset $A \subset S$ and a point $x \in F$, we define $V(x, A) \subset F$ as the complete cone over A with tip x , that is, as the union of rays emanating from x and asymptotic to A . If $\tau \subset S$ is a face, we call the cone $V(x, \tau)$ a *Weyl sector*, and if $\sigma \subset S$ is a chamber, we call $V(x, \sigma)$ a *euclidean Weyl chamber*.

After fixing an origin $o \in F$, the group W lifts to a group of isometries of F fixing o . The euclidean Weyl chambers $V(o, \sigma)$ are then fundamental domains for the action of $W \curvearrowright F$.

We define the *euclidean model Weyl chamber* as the quotient $V_{\text{mod}} = F/W$; we will also denote it by Δ or Δ_{euc} . It is canonically isometric to the complete euclidean cone over σ_{mod} . The natural projection

$$\text{proj}: F \rightarrow V_{\text{mod}} = \Delta_{\text{euc}} = \Delta$$

restricts to an isometry on every euclidean Weyl chamber $V(o, \sigma)$.

For a closed subset $\bar{A} \subset \sigma_{\text{mod}}$, we define $V(0, \bar{A}) \subset V_{\text{mod}}$ as the complete cone over \bar{A} with tip 0. In particular, a face τ_{mod} of σ_{mod} corresponds to a face $V(0, \tau_{\text{mod}})$ of V_{mod} .

We define the Δ -valued distance function or Δ -distance d_{Δ} on F by

$$d_{\Delta}(x, y) = \text{proj}(y - x) \in \Delta.$$

Note the symmetry property

$$(3) \quad d_{\Delta}(x, y) = \iota d_{\Delta}(y, x).$$

The Weyl group is precisely the group of isometries for the Δ -valued distance on F_{mod} which fix the origin.

Lemma 2.2 *Suppose that the Coxeter complex (S, W) is irreducible. Then $\iota = \text{id}$ if and only if the root system of (S, W) is of type $A_1, B_{\ell}, C_{\ell}, D_{2k}, E_{7,8}, F_4$ or G_2 . If $\iota \neq \text{id}$, then σ_{mod} contains exactly one root which, therefore, is ι -invariant.*

Proof The proof is by examination of the irreducible root systems (see eg [4]): $w_0 = -\text{id}$ if and only if the root system is of type $A_1, B_{\ell}, C_{\ell}, D_{2k}, E_{7,8}, F_4$ or G_2 . All the remaining irreducible root systems are simply laced; equivalently, W acts transitively on roots. \square

2.3 Hadamard manifolds

In this section only, X denotes a Hadamard manifold, ie a simply connected complete Riemannian manifold with nonpositive sectional curvature. We will use the notation $\text{Isom}(X)$ for the full isometry group of X .

Any two points in X are connected by a unique geodesic segment. We will use the notation xy for the oriented geodesic segment connecting x to y . We will often regard geodesic segments, geodesic rays and complete geodesics as parametrized with unit speed and treat them as isometric maps of intervals to X .

We will denote by $\angle_x(y, z)$ the angle between the geodesic segments xy and xz at the point x . For $x \in X$, we let $\Sigma_x X$ denote the *space of directions* of X at x , ie the unit sphere in the tangent space $T_x X$, equipped with the angle metric.

The *ideal* or *visual boundary* of X , denoted by $\partial_\infty X$, is the set of asymptote classes of geodesic rays in X , where two rays are *asymptotic* if and only if they have finite Hausdorff distance. Points in $\partial_\infty X$ are called *ideal points*. For $\xi \in \partial_\infty X$ and $x \in X$, we denote by $x\xi$ the geodesic ray emanating from x and asymptotic to ξ , ie representing the ideal point ξ . For $x \in X$, we have a natural map

$$\log_x: \partial_\infty X \rightarrow \Sigma_x X$$

sending $\xi \in \partial_\infty X$ to the velocity vector at x of the geodesic ray $x\xi$. The *cone* or *visual topology* on $\partial_\infty X$ is characterized by the property that all the maps \log_x are homeomorphisms; with respect to this topology, $\partial_\infty X$ is homeomorphic to the sphere of dimension $\dim(X) - 1$. The visual topology extends to $\bar{X} = X \cup \partial_\infty X$ as follows: a sequence (x_n) converges to an ideal point $\xi \in \partial_\infty X$ if the sequence of geodesic segments xx_n emanating from some (any) base point x converges to the ray $x\xi$ pointwise (equivalently, uniformly on compacta in \mathbb{R}). This topology makes \bar{X} into a closed ball. We define the visual boundary of a subset $A \subset X$ as the set $\partial_\infty A = \bar{A} \cap \partial_\infty X$ of its accumulation points at infinity.

The visual boundary $\partial_\infty X$ carries the natural *Tits (angle) metric* \angle_{Tits} , defined as

$$\angle_{\text{Tits}}(\xi, \eta) = \sup_{x \in X} \angle_x(\xi, \eta),$$

where $\angle_x(\xi, \eta)$ is the angle between the geodesic rays $x\xi$ and $x\eta$. The *Tits boundary* $\partial_{\text{Tits}} X$ is the metric space $(\partial_\infty X, \angle_{\text{Tits}})$. The Tits metric is lower semicontinuous with respect to the visual topology, and accordingly, the *Tits topology* induced by the Tits metric is finer than the visual topology. It is discrete if there is an upper negative curvature bound, and becomes nontrivial if flat directions occur. For instance, the Tits boundary of flat r -space is the unit $(r-1)$ -sphere, $\partial_{\text{Tits}} \mathbb{R}^r \cong S^{r-1}(1)$. An isometric embedding $X \rightarrow Y$ of Hadamard spaces induces an isometric embedding $\partial_{\text{Tits}} X \rightarrow \partial_{\text{Tits}} Y$ of Tits boundaries.

A subset A of $\partial_{\text{Tits}} X$ is called *convex* if for any two points $\xi, \eta \in A$ with $\angle_{\text{Tits}}(\xi, \eta) < \pi$, the (unique) geodesic $\xi\eta$ connecting ξ and η in $\partial_{\text{Tits}} X$ is entirely contained in A .

2.4 Symmetric spaces of noncompact type

The standard references for this and the following section are [8; 13]. Our treatment of this standard material is more geometric than the one presented in these books.

A symmetric space, denoted by X throughout this paper, is said to be of *noncompact type* if it is nonpositively curved, simply connected and has no euclidean factor. In particular, it is a Hadamard manifold. We will identify X with the quotient G/K where G is a semisimple Lie group with finite center acting isometrically and transitively on X , and K is a maximal compact subgroup of G . We will assume that G is commensurable with the isometry group $\text{Isom}(X)$ in the sense that we allow compact kernel and finite cokernel for the natural map $G \rightarrow \text{Isom}(X)$. In particular, the image of G in $\text{Isom}(X)$ contains the identity component $\text{Isom}(X)_o$. The Lie group G carries a natural structure as a real algebraic group.

A *point reflection* (also known as a *Cartan involution*) at a point $x \in X$ is an isometry σ_x which fixes x and has differential $-\text{id}_{T_x X}$ in x . In a symmetric space, point reflections exist in all points (by definition). A *transvection* of X is an isometry which is the product $\sigma_x \sigma_{x'}$ of two point reflections; it preserves the oriented geodesic through x and x' and the parallel vector fields along it. The transvections preserving a unit speed geodesic $c(t)$ form a one parameter subgroup (T_t^c) of $\text{Isom}(X)_o$ where T_t^c denotes the transvection mapping $c(s) \mapsto c(s+t)$. A nontrivial isometry ϕ of X is called *axial* if it preserves a geodesic l and shifts along it. (It does not have to be a transvection.) The geodesic l is called an *axis* of ϕ . Axes are in general not unique. They are parallel to each other.

A *flat* in X is a totally geodesic flat submanifold, equivalently, a convex subset isometric to a euclidean space. A maximal flat in X is a flat which is not contained in any larger flat; we will use the notation F for maximal flats. The group $\text{Isom}(X)_o$ acts transitively on the set of maximal flats; the common dimension of maximal flats is called the *rank* of X . The space X has rank one if and only if it has strictly negative sectional curvature.

A maximal flat F is preserved by all transvections along geodesic lines contained in it. In general, there exist nontrivial isometries of X fixing F pointwise. The subgroup of isometries of F which are induced by elements of G is isomorphic to a semidirect product $\mathbb{R}^r \rtimes W$, where r is the rank of X . The subgroup \mathbb{R}^r acts simply transitively on F by translations. The linear part W is a finite reflection group, called the *Weyl group* of G and X . Since maximal flats are equivalent modulo G , the action $W \curvearrowright F$ is well defined up to isometric conjugacy.

We will think of the Weyl group as acting on a *model flat* $F_{\text{mod}} \cong \mathbb{R}^r$ and on its visual boundary sphere at infinity, the *model apartment* $a_{\text{mod}} = \partial_{\text{Tits}} F_{\text{mod}} \cong S^{r-1}$. The pair (a_{mod}, W) is the *spherical Coxeter complex* associated with X . We identify the

spherical model Weyl chamber σ_{mod} with a (fundamental) chamber in the model apartment, $\sigma_{\text{mod}} \subset a_{\text{mod}}$. Accordingly, we identify the *euclidean model Weyl chamber* V_{mod} with the sector in F_{mod} with tip in the origin and visual boundary σ_{mod} , so $V_{\text{mod}} \subset F_{\text{mod}}$.

The Δ -valued distance naturally extends from F_{mod} to X because every pair of points lies in a maximal flat. In order to define the Δ -distance $d_{\Delta}(x, y)$ of two points $x, y \in X$, one chooses a maximal flat F containing x, y and identifies it isometrically with F_{mod} so that the Weyl group actions correspond. The resulting quantity $d_{\Delta}(x, y)$ is independent of the choices. We refer the reader to [17] for the detailed discussion of *metric properties* of d_{Δ} .

For every maximal flat $F \subset X$, we have a Tits isometric embedding $\partial_{\infty}F \subset \partial_{\infty}X$ of its visual boundary sphere. There is an identification $\partial_{\infty}F \cong a_{\text{mod}}$ with the model apartment, unique up to composition with elements in W . The Coxeter complex structure on a_{mod} induces a simplicial structure on $\partial_{\infty}F$. The visual boundaries of maximal flats cover $\partial_{\infty}X$ because every geodesic ray in X is contained in a maximal flat. Moreover, their intersections are simplicial. One thus obtains a G -invariant piecewise spherical *simplicial structure* on $\partial_{\infty}X$ which makes $\partial_{\infty}X$ into a *spherical building* and, also taking into account the visual topology, into a topological spherical building. It is called the *spherical* or *Tits building* associated to X . The Tits metric is the path metric with respect to the piecewise spherical structure. We will refer to the simplices as *faces*.

The visual boundaries $\partial_{\infty}F \subset \partial_{\infty}X$ of the maximal flats $F \subset X$ are precisely the *apartments* with respect to the spherical building structure at infinity, which in turn are precisely the convex subsets isometric to the unit $(r-1)$ -sphere with respect to the Tits metric. Any two points in $\partial_{\infty}X$ lie in a common apartment.

The action $G \curvearrowright \partial_{\infty}X$ on ideal points is not transitive if X has rank ≥ 2 . Every G -orbit meets every chamber exactly once. The quotient can be identified with the spherical model chamber, $\partial_{\infty}X/G \cong \sigma_{\text{mod}}$. We call the projection

$$\theta: \partial_{\infty}X \rightarrow \partial_{\infty}X/G \cong \sigma_{\text{mod}}$$

the *type map*. It restricts to an isometry on every chamber $\sigma \subset \partial_{\infty}X$. We call the inverse $\kappa_{\sigma} = (\theta|_{\sigma})^{-1}: \sigma_{\text{mod}} \rightarrow \sigma$ the (*chamber*) *chart* for σ . Consequently, θ restricts to an isometry on every face $\tau \subset \partial_{\infty}X$. We call $\theta(\tau) \subset \sigma_{\text{mod}}$ the *type* of the face τ and $\kappa_{\tau} = (\theta|_{\tau})^{-1}: \theta(\tau) \rightarrow \tau$ its *chart*. We define the *type* of an ideal point $\xi \in \partial_{\infty}X$ as its image $\theta(\xi) \in \sigma_{\text{mod}}$. A point $\xi \in \partial_{\infty}X$ is called *regular* if its type is an interior

point of σ_{mod} , and *singular* otherwise. We denote by $\partial_{\infty}^{\text{reg}} X \subset \partial_{\infty} X$ the set of regular ideal boundary points. A point $\rho \in \partial_{\text{Tits}} X$ is said to be of *root type* if $\theta(\rho)$ is a root in $\sigma_{\text{mod}} \subset S$. Equivalently, the closed $\frac{\pi}{2}$ -ball centered at ρ (with respect to the Tits metric) is simplicial, ie is a simplicial subcomplex of $\partial_{\text{Tits}} X$. If $a \subset \partial_{\infty} X$ is an apartment, we call a type preserving isometry $\kappa_a: a_{\text{mod}} \rightarrow a$ an *apartment chart* for a .

A geodesic segment xy in X is called *regular* if $x \neq y$ and for the unique geodesic ray $x\xi$ extending xy , the point $\xi \in \partial_{\text{Tits}} X$ is regular. Equivalently, the vector $d_{\Delta}(x, y)$ belongs to the interior of V_{mod} .

Definition 2.3 (antipodal) (i) Two ideal points $\xi, \eta \in \partial_{\infty} X$ are called *antipodal* if $\angle_{\text{Tits}}(\xi, \eta) = \pi$. A subset of $\partial_{\infty} X$ is called *antipodal* if the points in it are pairwise antipodal.

(ii) Two simplices $\tau_1, \tau_2 \subset \partial_{\infty} X$ are *opposite* (or *antipodal*) with respect to a point $x \in X$ if $\tau_2 = \sigma_x \tau_1$, where σ_x denotes the reflection at the point x . Two simplices $\tau_1, \tau_2 \subset \partial_{\infty} X$ are *opposite* (or *antipodal*) if they are opposite simplices in the apartments containing both of them.

Note that the last property holds if and only if some (every) interior point of τ_1 has an antipode in the interior of τ_2 , equivalently, if and only if τ_1 and τ_2 are opposite with respect to some point $x \in X$. Their types are then related by $\theta(\tau_2) = \iota(\theta(\tau_1))$. We will frequently use the notation $\tau, \hat{\tau}$ and τ_+, τ_- for pairs of antipodal simplices.

A pair of opposite chambers $\sigma_+, \sigma_- \subset \partial_{\infty} X$ is contained in a unique apartment, which we will denote by $a(\sigma_+, \sigma_-)$; the apartment $a(\sigma_+, \sigma_-)$ is the visual boundary of a unique maximal flat $F(\sigma_+, \sigma_-)$ in X .

For a point $x \in X$ and a simplex $\tau \subset \partial_{\infty} X$, define the (Weyl) *sector* $V = V(x, \tau) \subset X$ as the union of rays $x\xi$ for all ideal points $\xi \in \tau$. Weyl sectors are contained in flats. They are isometric images of faces $V(0, \tau_{\text{mod}}) \subset V_{\text{mod}}$ of the euclidean model Weyl chamber under isometric embeddings $F_{\text{mod}} \rightarrow X$ which are type preserving at infinity. More generally, for a point $x \in X$ and a closed subset $A \subset \partial_{\infty} X$, we define the *Weyl cone* $V(x, A)$ as the union of all rays $x\xi$ for $\xi \in A$. Weyl cones are in general not flat.

The stabilizers $B_{\sigma} \subset G$ of the chambers $\sigma \subset \partial_{\infty} X$ are the *minimal parabolic subgroups* of G . After choosing a reference chamber $\sigma_0 \subset \partial_{\infty} X$, we call $B = B_{\sigma_0}$ the *positive* minimal parabolic subgroup. The group G acts transitively on the set of chambers in $\partial_{\infty} X$, which we will then identify with G/B , the *full flag manifold*

of G . The minimal parabolic subgroups are algebraic subgroups of G , and G/B is a real projective variety. The set $\partial_{\text{Fü}}X \cong G/B$ of chambers in $\partial_\infty X$ is called the *Furstenberg boundary* of X ; we will equip it with the visual topology (as opposed to the Zariski topology coming from G/B) which coincides with its manifold topology as a compact homogeneous G -space. Every regular G -orbit $G\xi \subset \partial_\infty X$, for $\xi \in \text{int}(\sigma_0)$, is G -equivariantly and homeomorphically identified with $\partial_{\text{Fü}}X$ by assigning to the (regular) ideal point $g\xi$ the unique chamber $g\sigma_0$ containing it.

The stabilizers $P_\tau \subset G$ of simplices $\tau \subset \partial_\infty X$ are the *parabolic subgroups* of G . The group G acts transitively on simplices of the same type. The set $\text{Flag}_{\tau_{\text{mod}}} \cong G/P_{\tau_{\text{mod}}}$ of the simplices τ of type $\theta(\tau) = \tau_{\text{mod}} \subset \sigma_{\text{mod}}$ is called the *partial flag manifold* of type τ_{mod} . In particular, $\text{Flag}_{\sigma_{\text{mod}}} = \partial_{\text{Fü}}X$. Again, we equip the flag manifolds with the visual topology; it agrees with their topology as compact homogeneous G -spaces. Every G -orbit $G\xi \subset \partial_\infty X$ of type $\theta(\xi) \in \text{int}(\tau_{\text{mod}})$ is G -equivariantly homeomorphic to $\text{Flag}_{\tau_{\text{mod}}}$.

For a flag manifold $\text{Flag}_{\tau_{\text{mod}}}$ and a simplex $\hat{\tau}$ of type $\iota\tau_{\text{mod}}$, we define the *open Schubert stratum* $C(\hat{\tau}) \subset \text{Flag}_{\tau_{\text{mod}}}$ as the subset of simplices opposite to $\hat{\tau}$ in the sense of Definition 2.3. It follows from semicontinuity of the Tits distance that the subset $C(\hat{\tau}) \subset \text{Flag}_{\tau_{\text{mod}}}$ is indeed open. Furthermore, this subset is also dense in $\text{Flag}_{\tau_{\text{mod}}}$. We note that for rank-1 symmetric spaces, the only flag manifold associated to G is $\partial_\infty X$ and the open Schubert strata are the complements of points.

If τ_{mod} is ι -invariant, we say as in Definition 2.3 that a subset of $\text{Flag}_{\tau_{\text{mod}}}$ is *antipodal* if the simplices in it are pairwise opposite.

3 Geometry of visual boundaries

In this section, we introduce definitions and prove some properties of symmetric spaces of noncompact type and their visual boundaries of more specific nature which are needed for our study of discrete group actions at infinity.

3.1 Stars at infinity and regular points

For a simplex $\tau \subset \partial_\infty X$, the *star* $\text{st}(\tau) \subset \partial_\infty X$ is the union of all closed chambers $\sigma \supseteq \tau$. It is proven in [19, Proposition 2.14] that for each face $\tau \subset \partial_\infty X$, the *Weyl cone* $V(x, \text{st}(\tau))$ is a closed convex subset of X .

For a face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we define the *open star*

$$\text{ost}(\tau_{\text{mod}}) \subset \sigma_{\text{mod}}$$

as the union of all open faces of σ_{mod} whose closure contains τ_{mod} . Its complement

$$\partial \text{st}(\tau_{\text{mod}}) := \sigma_{\text{mod}} - \text{ost}(\tau_{\text{mod}})$$

is the union of all (closed) faces of σ_{mod} which do not contain τ_{mod} .

For a simplex $\tau \subset \partial_{\infty} X$, we define the *open star*

$$\text{ost}(\tau) \subset \text{st}(\tau) \subset \partial_{\infty} X$$

as the union of all open simplices in $\partial_{\infty} X$ whose closure contains τ . Then

$$\partial \text{st}(\tau) := \text{st}(\tau) - \text{ost}(\tau)$$

is the union of all (closed) simplices in $\text{st}(\tau)$ which do not contain τ ,

Definition 3.1 An ideal point $\xi \in \partial_{\infty} X$ is said to be τ_{mod} -regular if $\theta(\xi) \in \text{ost}(\tau_{\text{mod}})$, and τ_{mod} -singular if $\theta(\xi) \in \partial \text{st}(\tau_{\text{mod}})$.

We will call σ_{mod} -regular points simply *regular*. Note that $\text{ost}(\sigma_{\text{mod}}) = \text{int}(\sigma_{\text{mod}})$, and the regular points in $\partial_{\infty} X$ are precisely the interior points of chambers.

Note that $\text{ost}(\tau)$ is the subset of $\theta(\tau)$ -regular points in $\text{st}(\tau)$ and $\partial \text{st}(\tau)$ is subset of $\theta(\tau)$ -singular points. The τ_{mod} -regular part

$$\partial_{\infty}^{\tau_{\text{mod}}\text{-reg}} X = \theta^{-1}(\text{ost}(\tau_{\text{mod}})) \subset \partial_{\infty} X$$

of the visual boundary contains all open chambers and is in particular dense in $\partial_{\infty} X$ (also with respect to the Tits topology). For a τ_{mod} -regular point $\xi \in \partial_{\infty} X$, there is a unique closest (with respect to the Tits metric) simplex $\tau \subset \partial_{\infty} X$ of type τ_{mod} , namely the one with $\xi \in \text{ost}(\tau)$.

The notion of regularity extends to oriented geodesic segments, rays and lines in X : A geodesic ray $x\eta \subset X$ is τ_{mod} -regular if its ideal endpoint $\eta \in \partial_{\infty} X$ is. An oriented geodesic segment $xy \subset X$ is τ_{mod} -regular if the geodesic ray $x\eta$ extending it is.

The geometric significance of σ_{mod} -regularity of geodesic segments comes from the fact that a geodesic segment (or ray) in X is σ_{mod} -regular if and only if it is contained in a unique maximal flat.

3.2 Folding order

In this section, we discuss natural partial orders on Weyl orbits in the model apartment and give different equivalent geometric definitions for them.

By a *folding map* $a_{\text{mod}} \rightarrow a_{\text{mod}}$, we mean a type preserving continuous map which sends chambers isometrically onto chambers.

We will be working with folding maps which fix some reference face and think of them as moving points “closer” towards this face. For instance, for a simplicial hemisphere $h \subset a_{\text{mod}}$ (containing the reference face) there is the folding map fixing h and reflecting the complementary hemisphere onto it, see the discussion of special foldings below.

Definition 3.2 (folding order) For a face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we define the τ_{mod} -*folding order* $\prec_{\tau_{\text{mod}}}$ on a_{mod} as follows: for distinct points $\bar{\xi}_1, \bar{\xi}_2 \in a_{\text{mod}}$, we say that $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$ if and only if there exists a folding map $f: a_{\text{mod}} \rightarrow a_{\text{mod}}$ such that $f|_{\tau_{\text{mod}}} = \text{id}_{\tau_{\text{mod}}}$ and $f(\bar{\xi}_2) = \bar{\xi}_1$.

We will use the notation \prec for $\prec_{\sigma_{\text{mod}}}$.

The relations $\prec_{\tau_{\text{mod}}}$ are *transitive* since compositions of folding maps are folding maps.

Remark 3.3 (o) Our folding order inequalities are nonstrict, ie allowing equality.

(i) The relation $\prec_{\tau_{\text{mod}}}$ is closed. It compares only points which lie in the same Weyl orbit; ie if $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$ then $W\bar{\xi}_1 = W\bar{\xi}_2$.

(ii) The relation $\prec_{\tau_{\text{mod}}}$ on singular Weyl orbits is the closure of the relation $\prec_{\tau_{\text{mod}}}$ on regular ones: it holds that $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$ if and only if there exist sequences of regular points $\bar{\xi}_i^n \rightarrow \bar{\xi}_i$ with $\bar{\xi}_1^n \prec_{\tau_{\text{mod}}} \bar{\xi}_2^n$.

(iii) Any isometry of a_{mod} preserving τ_{mod} as a set preserves the relation $\prec_{\tau_{\text{mod}}}$.

(iv) The relations $\prec_{\tau_{\text{mod}}}$ and \prec are closely related. Clearly, \prec is stronger than $\prec_{\tau_{\text{mod}}}$; ie if $\bar{\xi}_1 \prec \bar{\xi}_2$ then $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$. More precisely, note that a folding map f fixing τ_{mod} is the composition $w \circ f'$ of a folding map f' fixing σ_{mod} with an element $w \in W_{\tau_{\text{mod}}}$. Thus $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$ if and only if there exist $\bar{\xi}'_i \in W_{\tau_{\text{mod}}}\bar{\xi}_i$ such that $\bar{\xi}'_1 \prec \bar{\xi}'_2$.

(v) If $\bar{\xi}_1 \prec_{\tau_{\text{mod}}} \bar{\xi}_2$ then $w_1\bar{\xi}_1 \prec_{\tau_{\text{mod}}} w_2\bar{\xi}_2$ for all $w_1, w_2 \in W_{\tau_{\text{mod}}}$, because $w_1fw_2^{-1}$ is again a folding map fixing τ_{mod} . Hence $\prec_{\tau_{\text{mod}}}$ descends to a relation on the quotient $W_{\tau_{\text{mod}}}\backslash a_{\text{mod}}$, which we also denote by $\prec_{\tau_{\text{mod}}}$.

There is a *metric estimate* for the folding order, because folding maps are 1–Lipschitz:

Lemma 3.4 *If $\bar{\vartheta} \in \tau_{\text{mod}}$, then we have the implication*

$$W_{\tau_{\text{mod}}}\bar{\xi}_1 \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}}\bar{\xi}_2 \implies \angle_{\text{Tits}}(\bar{\xi}_1, \bar{\vartheta}) \leq \angle_{\text{Tits}}(\bar{\xi}_2, \bar{\vartheta}).$$

Also, if $\bar{\vartheta} \in \text{int}(\tau_{\text{mod}})$, equality holds on the right-hand side only if $W_{\tau_{\text{mod}}}\bar{\xi}_1 = W_{\tau_{\text{mod}}}\bar{\xi}_2$.

Proof Suppose $W_{\tau_{\text{mod}}}\bar{\xi}_1 \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}}\bar{\xi}_2$. Then there exists a folding map $a_{\text{mod}} \rightarrow a_{\text{mod}}$ fixing τ_{mod} with $f(\bar{\xi}_2) = \bar{\xi}_1$. It maps the geodesic segment $\bar{\vartheta}\bar{\xi}_2$ to a broken geodesic segment β from $\bar{\vartheta}$ to $\bar{\xi}_1$ of the same length, whence the implication of inequalities.

Suppose now in addition that $\bar{\vartheta} \in \text{int}(\tau_{\text{mod}})$ and $W_{\tau_{\text{mod}}}\bar{\xi}_1 \neq W_{\tau_{\text{mod}}}\bar{\xi}_2$. The initial segments of β and $\bar{\vartheta}\bar{\xi}_2$ have the same type. Therefore, there exists $w \in W_{\bar{\vartheta}} = W_{\tau_{\text{mod}}}$ such that $w\beta$ and $\bar{\vartheta}\bar{\xi}_2$ have a common initial segment. Since $w\bar{\xi}_1 \neq \bar{\xi}_2$, the broken geodesic segment β cannot be a true geodesic segment, and we obtain the strict metric inequality $\angle_{\text{Tits}}(\bar{\xi}_1, \bar{\vartheta}) < \angle_{\text{Tits}}(\bar{\xi}_2, \bar{\vartheta})$. □

As a consequence, we can justify our terminology of “partial order”:

Corollary 3.5 *The relation $\prec_{\tau_{\text{mod}}}$ is a partial order on $W_{\tau_{\text{mod}}}\backslash a_{\text{mod}}$.*

Proof We must verify antisymmetry. Suppose $W_{\tau_{\text{mod}}}\bar{\xi}_1 \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}}\bar{\xi}_2 \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}}\bar{\xi}_1$. By Lemma 3.4, we have $\angle_{\text{Tits}}(\bar{\xi}_1, \bar{\vartheta}) = \angle_{\text{Tits}}(\bar{\xi}_2, \bar{\vartheta})$ for all $\bar{\vartheta} \in \tau_{\text{mod}}$. The equality part implies that $W_{\tau_{\text{mod}}}\bar{\xi}_1 = W_{\tau_{\text{mod}}}\bar{\xi}_2$. □

We discuss next the structure of folding maps and decompositions into simple ones.

Each wall m splits a_{mod} into two hemispheres, the inner hemisphere h^+ containing σ_{mod} and the outer hemisphere h^- . This decomposition gives rise to the folding map which fixes h^+ and reflects h^- onto it. We call a composition of such folding maps at walls m_i a *special folding*. The intersection $\bigcap_i h_i^+$ of inner hemispheres is fixed by the special folding. In particular, special foldings fix the model chamber σ_{mod} .

In general, there are folding maps fixing σ_{mod} which are not special. However, this makes no difference for the folding order \prec :

Lemma 3.6 (compare [24, page 441, Theorem 4.9]) *If for points $\bar{\xi}_1, \bar{\xi}_2 \in a_{\text{mod}}$ there exists a folding map fixing σ_{mod} and mapping $\bar{\xi}_2 \mapsto \bar{\xi}_1$, then there exists a special folding with this property.*

Proof We may assume that $\bar{\xi}_1$ and $\bar{\xi}_2$ are regular and different. We connect a point $\bar{\eta}$ in the interior of σ_{mod} to $\bar{\xi}_2$ by a geodesic segment $\bar{\gamma}$ which avoids faces of codimension at least two. Let f be a folding map fixing σ_{mod} with $f(\bar{\xi}_2) = \bar{\xi}_1$. Then $\bar{\beta} = f \circ \bar{\gamma}$ is a broken geodesic path which connects $\bar{\eta}$ to $\bar{\xi}_1$ and has the same length and initial direction as $\bar{\gamma}$. Its bending points are interior points of panels and $\bar{\beta}$ is locally “reflected” at the walls containing these panels. The assertion follows if we can replace $\bar{\beta}$ by a broken geodesic path from $\bar{\eta}$ to $\bar{\xi}_1$, which is the image of $\bar{\gamma}$ under a *special* folding.

Let $\bar{\eta}_1$ denote the first bending point of $\bar{\beta}$ starting from $\bar{\eta}$. It lies in a wall m_1 . If $\bar{\beta}$ crosses m_1 again in some point $\bar{\eta}_2$, then we replace the subpath $\bar{\eta}_1\bar{\eta}_2$ by its reflection at m_1 . The modified broken path $\bar{\beta}'$ has again reflection folds, the same initial direction and the same endpoint. Moreover, its initial segment is strictly longer. After finitely many such modifications, we may assume that $\bar{\beta}'$ stays inside h_1^+ . (The wall m_1 has changed in the process.) We then can obtain $\bar{\beta}'$ as the image of another broken path $\bar{\beta}''$ under the special folding s_1 at m_1 , ie $\bar{\beta}' = s_1 \circ \bar{\beta}''$, such that $\bar{\beta}''$ has a strictly longer initial segment than $\bar{\beta}'$.

Thus, we can replace $\bar{\beta}$ by another broken path $\bar{\beta}''$ with reflection bends, with the same length and initial direction as $\bar{\gamma}$, with a strictly longer initial segment than $\bar{\beta}$, and such that some special folding s_1 maps the endpoint of $\bar{\beta}''$ to the endpoint of $\bar{\beta}$. It follows by induction that $\bar{\beta}$ can be replaced by another broken path with the same endpoint and which is the image of $\bar{\gamma}$ under a special folding. □

Corollary 3.7 (alternative definition of σ_{mod} -folding order) *We have $\bar{\xi}_1 < \bar{\xi}_2$ if and only if there exists a special folding which maps $\bar{\xi}_2 \mapsto \bar{\xi}_1$.*

We note that the partial order $<$ was defined exactly in this way by Littellmann; see [30, page 509].

The folding orders on the Weyl orbits in the model apartment correspond to orders on the Weyl group and its (double) coset spaces, as we explain now.

We can regard the regular Weyl orbits as copies of W by identifying the orbit point in the chamber $w\sigma_{\text{mod}}$ with the element $w \in W$. Under this identification, it holds that

$$w_1 < w_2$$

for elements $w_1, w_2 \in W$ if and only if there exists a folding map $a_{\text{mod}} \rightarrow a_{\text{mod}}$ fixing σ_{mod} and mapping $w_2\sigma_{\text{mod}} \mapsto w_1\sigma_{\text{mod}}$, and

$$w_1 <_{\tau_{\text{mod}}} w_2$$

if and only if there exists such a folding map fixing only τ_{mod} . Again, $\prec_{\tau_{\text{mod}}}$ descends to an order on $W_{\tau_{\text{mod}}} \setminus W$, also denoted by $\prec_{\tau_{\text{mod}}}$.

Remark 3.8 (Bruhat order) The corollary shows that the folding order \prec on W coincides with the *Bruhat order*; see [14, Chapter 5.9] or [2, Chapter 2] for a definition. Hence, the folding order gives a *geometric interpretation* of the Bruhat order. To verify this, one observes that if the chambers $w\sigma_{\text{mod}}$ and $w'\sigma_{\text{mod}}$ are symmetric with respect to a wall and if $w\sigma_{\text{mod}}$ lies in the inner hemisphere, then we have the inequality $l(w) < l(w')$ for word lengths. Here the word length on W is defined using as generators the reflections at the walls of σ_{mod} .

More generally, if $\bar{\xi}$ is an interior point of the face $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$, ie $\bar{\xi} \in \text{int}(\nu_{\text{mod}})$, then $W_{\bar{\xi}} = W_{\nu_{\text{mod}}}$ and $W\bar{\xi} \cong W/W_{\nu_{\text{mod}}}$. Under this identification, the order $\prec_{\tau_{\text{mod}}}$ on the Weyl orbit quotient $W_{\tau_{\text{mod}}} \setminus W\bar{\xi} \subset W_{\tau_{\text{mod}}} \setminus a_{\text{mod}}$ becomes a partial order on the double quotient $W_{\tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}}$; compare Remark 3.3(v). It holds that

$$W_{\tau_{\text{mod}}} w_1 W_{\nu_{\text{mod}}} \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}} w_2 W_{\nu_{\text{mod}}}$$

if and only if there exist $w'_i \in W_{\tau_{\text{mod}}} w_i W_{\nu_{\text{mod}}}$ such that $w'_1 < w'_2$; see [32] for a slightly different description of this order.

We next describe the effect of the longest element $w_0 \in W$ on the folding order. Recall that w_0 is the involution sending σ_{mod} to the opposite chamber $\hat{\sigma}_{\text{mod}}$ in a_{mod} .

Lemma 3.9 *Left multiplication with w_0 reverses the σ_{mod} -folding order.*

Proof Suppose that the special folding s_m at the wall m maps $\bar{\xi}_2$ to $\bar{\xi}_1$; ie $s_m \bar{\xi}_2 = \bar{\xi}_1$. When applying w_0 , the inner hemisphere bounded by m becomes the outer hemisphere bounded by $w_0 m$ and vice versa: $w_0 h_m^\pm = h_{w_0 m}^\mp$. Hence $s_{w_0 m} w_0 \bar{\xi}_1 = w_0 \bar{\xi}_2$. The assertion follows by applying Corollary 3.7 and induction. \square

Regarding the analogous fact for the orders $\prec_{\tau_{\text{mod}}}$, note that $w_0 W_{\tau_{\text{mod}}} w_0^{-1} = W_{w_0 \tau_{\text{mod}}} = W_{\iota \tau_{\text{mod}}}$, and w_0 maps $W_{\tau_{\text{mod}}}$ -orbits to $W_{\iota \tau_{\text{mod}}}$ -orbits. The action of w_0 therefore induces a natural map

$$(4) \quad W_{\tau_{\text{mod}}} \setminus a_{\text{mod}} \xrightarrow{w_0} W_{\iota \tau_{\text{mod}}} \setminus a_{\text{mod}}, \quad W_{\tau_{\text{mod}}} \bar{\xi} \mapsto w_0 W_{\tau_{\text{mod}}} \bar{\xi} = W_{\iota \tau_{\text{mod}}} w_0 \bar{\xi},$$

and correspondingly,

$$(5) \quad \begin{aligned} W_{\tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}} &\xrightarrow{w_0} W_{\iota \tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}}, \\ W_{\tau_{\text{mod}}} w W_{\nu_{\text{mod}}} &\mapsto w_0 W_{\tau_{\text{mod}}} w W_{\nu_{\text{mod}}} = W_{\iota \tau_{\text{mod}}} w_0 w W_{\nu_{\text{mod}}}. \end{aligned}$$

The lemma implies that these maps are order reversing:

$$(6) \quad W_{\tau_{\text{mod}}} \bar{\xi}_1 \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}} \bar{\xi}_2 \iff W_{\iota\tau_{\text{mod}}} w_0 \bar{\xi}_1 \succ_{\iota\tau_{\text{mod}}} W_{\iota\tau_{\text{mod}}} w_0 \bar{\xi}_2,$$

and, respectively,

$$W_{\tau_{\text{mod}}} w W_{\nu_{\text{mod}}} \prec_{\tau_{\text{mod}}} W_{\tau_{\text{mod}}} w' W_{\nu_{\text{mod}}} \iff W_{\iota\tau_{\text{mod}}} w_0 w W_{\nu_{\text{mod}}} \succ_{\iota\tau_{\text{mod}}} W_{\iota\tau_{\text{mod}}} w_0 w' W_{\nu_{\text{mod}}}.$$

3.3 Relative position at infinity

Let $\sigma_0, \sigma \subset \partial_\infty X$ be chambers. There exists an (in general nonunique) apartment $a \subset \partial_\infty X$ containing these chambers, ie $\sigma_0, \sigma \subset a$, and a unique apartment chart $\alpha: a_{\text{mod}} \rightarrow a$ such that $\sigma_0 = \alpha(\sigma_{\text{mod}})$. We define the *position of σ relative to σ_0* as the chamber

$$\text{pos}(\sigma, \sigma_0) := \alpha^{-1}(\sigma) \subset a_{\text{mod}}.$$

Abusing notation, it can be regarded algebraically as the unique element

$$\text{pos}(\sigma, \sigma_0) \in W$$

such that

$$\sigma = \alpha(\text{pos}(\sigma, \sigma_0)\sigma_{\text{mod}}).$$

The relative position does not depend on the choice of the apartment a . To see this, choose regular points $\xi_0 \in \text{int}(\sigma_0)$ and $\xi \in \text{int}(\sigma)$ which are not antipodal, ie $\angle_{\text{Tits}}(\xi, \xi_0) < \pi$. Then the segment $\xi_0\xi$ is contained in a by convexity, and its image $\alpha^{-1}(\xi_0\xi)$ in a_{mod} is independent of the chart α because its initial portion $\alpha^{-1}(\xi_0\xi \cap \sigma_0)$ in σ_{mod} is.

More generally, we define the position of a chamber σ relative to an arbitrary simplex τ_0 of type τ_{mod} as follows. Let again $a \subset \partial_\infty X$ be an apartment containing τ_0 and σ , and let $\alpha: a_{\text{mod}} \rightarrow a$ be a chart such that $\tau_0 = \alpha(\tau_{\text{mod}})$. It is unique up to precomposition with an element in $W_{\tau_{\text{mod}}}$. We define the *position of σ relative to τ_0* as the $W_{\tau_{\text{mod}}}$ -orbit of the chamber $\alpha^{-1}(\sigma) \subset a_{\text{mod}}$. It can be interpreted algebraically as a coset

$$\text{pos}(\sigma, \tau_0) \in W_{\tau_{\text{mod}}} \backslash W.$$

Even more generally, we define the position of a simplex $\nu \subset \partial_\infty X$ relative τ_0 . Let $a \subset \partial_\infty X$ be an apartment containing τ_0 and ν , and let $\alpha: a_{\text{mod}} \rightarrow a$ be a chart such that $\tau_0 = \alpha(\tau_{\text{mod}})$. We define the *position of ν relative to τ_0* as the $W_{\tau_{\text{mod}}}$ -orbit of the simplex $\alpha^{-1}(\nu) \subset a_{\text{mod}}$. It can be interpreted algebraically as a double coset

$$\text{pos}(\nu, \tau_0) \in W_{\tau_{\text{mod}}} \backslash W / W_{\nu_{\text{mod}}},$$

where $\nu_{\text{mod}} = \theta(\nu)$ is the type of ν . Finally, we define the position of an ideal point $\xi \in \partial_\infty X$ relative τ_0 as the relative position of the simplex $\nu_\xi \subset \partial_\infty X$ spanned by ξ (ie containing ξ as an interior point):

$$\text{pos}(\xi, \tau_0) := \text{pos}(\nu_\xi, \tau_0) \in W_{\tau_{\text{mod}}} \setminus W / W_{\nu_{\text{mod}}},$$

where $\nu_{\text{mod}} = \theta(\nu_\xi)$. In particular, $\text{pos}(\xi, \tau_0) \in W_{\tau_{\text{mod}}} \setminus W$ if ξ is regular.

Lemma 3.10 *Two ideal points ξ_1, ξ_2 in the same G -orbit $G\xi \subset \partial_\infty X$ have the same position relative to a simplex $\tau \subset \partial_\infty X$ if and only if they lie in the same orbit of the parabolic subgroup $P_\tau < G$:*

$$\text{pos}(\xi_1, \tau) = \text{pos}(\xi_2, \tau) \iff P_\tau \xi_1 = P_\tau \xi_2.$$

Proof The implication “ \Leftarrow ” is clear. For “ \Rightarrow ”, let $a_i \subset \partial_\infty X$ be apartments containing τ and ξ_i . There exists $p \in P_\tau$ such that $a_1 = pa_2$. Then $\text{pos}(\xi_1, \tau) = \text{pos}(\xi_2, \tau) = \text{pos}(p\xi_2, \tau)$ if and only if ξ_1 and $p\xi_2$ span the same simplex in a_1 . In view of $\theta(\xi_1) = \theta(\xi_2)$, the latter is equivalent to $\xi_1 = \xi_2$. \square

The positions relative to τ thus correspond to the orbits of P_τ , and we have the identification

$$P_\tau \setminus G\xi \cong P_\tau \setminus \text{Flag}_{\nu_{\text{mod}}} \cong P_\tau \setminus G/P_{\nu_\xi} \cong W_{\tau_{\text{mod}}} \setminus W / W_{\nu_{\text{mod}}}$$

with $\tau_{\text{mod}} = \theta(\tau)$ and $\nu_{\text{mod}} = \theta(\nu_\xi)$.

In particular, for regular orbits, which are copies of the Furstenberg boundary, we obtain the identification

$$P_\tau \setminus \partial_{\text{Fü}} X \cong W_{\tau_{\text{mod}}} \setminus W.$$

The positions relative to a chamber σ correspond to the orbits of the minimal parabolic subgroup B_σ , and we have

$$B_\sigma \setminus G\xi \cong B_\sigma \setminus \text{Flag}_{\nu_{\text{mod}}} \cong B_\sigma \setminus G/P_{\nu_\xi} \cong W / W_{\nu_{\text{mod}}} \quad \text{and} \quad B_\sigma \setminus \partial_{\text{Fü}} X \cong W.$$

The G -orbits $G\xi$, respectively, the flag manifolds $\text{Flag}_{\nu_{\text{mod}}}$ thus decompose into finitely many P_τ -orbits which we call *Schubert strata* relative τ or τ -*Schubert strata*, and their closures (*generalized*) *Schubert cycles*. (We will see below that the cycles are unions of strata.) The level sets of $\text{pos}(\cdot, \sigma)$, ie the B_σ -orbits, are called *Schubert cells* relative σ .

Note that the Schubert cycles in the flag manifolds are *projective subvarieties*.

We will use the following notation. For a simplex $\tau_- \in \text{Flag}_{\tau_{\text{mod}}}$, we denote by

$$C_{\tau_{\text{mod}}}(\tau_-) := \{\tau : \tau \text{ opposite to } \tau_-\} \subset \text{Flag}_{\tau_{\text{mod}}}$$

the *open Schubert stratum* associated with τ_- in $\text{Flag}_{\tau_{\text{mod}}}$,

We can now use the folding order to compare the positions of points in a G -orbit $G\xi \subset \partial_\infty X$, respectively, a flag manifold $\text{Flag}_{\nu_{\text{mod}}}$ relative to simplices τ of a fixed type τ_{mod} .

We begin by proving a useful monotonicity property for the folding order under folding maps. It is a direct consequence of the definition of the folding order that folding maps $a_{\text{mod}} \rightarrow a_{\text{mod}}$ decrease the relative positions of pairs of simplices. We will need the same fact for *folding maps* $a_{\text{mod}} \rightarrow \partial_\infty X$ and $\partial_\infty X \rightarrow \partial_\infty X$ by which we mean, as before, type preserving continuous maps sending chambers isometrically onto chambers.

Lemma 3.11 (monotonicity) (i) *For a folding map $f: \partial_\infty X \rightarrow \partial_\infty X$ and simplices $\tau, \nu \subset \partial_\infty X$, it holds that*

$$\text{pos}(f(\nu), f(\tau)) \prec_{\theta(\tau)} \text{pos}(\nu, \tau).$$

(ii) *For a folding map $\alpha: a_{\text{mod}} \rightarrow \partial_\infty X$, a face type τ_{mod} and a simplex $\bar{\nu} \subset a_{\text{mod}}$, it holds that*

$$\text{pos}(\alpha(\bar{\nu}), \alpha(\tau_{\text{mod}})) \prec_{\tau_{\text{mod}}} \text{pos}(\bar{\nu}, \tau_{\text{mod}}).$$

Proof Part (i) reduces to part (ii) by choosing an apartment $a \supset \tau \cup \nu$ and a chart $\kappa: a_{\text{mod}} \rightarrow a$ with $\kappa(\tau_{\text{mod}}) = \tau$ for the face type $\tau_{\text{mod}} = \theta(\tau)$. Then apply (ii) to $\alpha = f \circ \kappa$ and $\bar{\nu} = \kappa^{-1}(\nu)$.

To verify (ii), consider the composition

$$a_{\text{mod}} \xrightarrow{\alpha} \partial_\infty X \rightarrow \partial_\infty X / B_\alpha(\sigma_{\text{mod}}) \cong a_{\text{mod}},$$

where the second map is the natural projection. It is a folding map $\bar{\alpha}: a_{\text{mod}} \rightarrow a_{\text{mod}}$ fixing σ_{mod} , and therefore

$$\text{pos}(\bar{\alpha}(\bar{\nu}), \tau_{\text{mod}}) \prec_{\tau_{\text{mod}}} \text{pos}(\bar{\nu}, \tau_{\text{mod}}).$$

Since the $B_{\alpha(\sigma_{\text{mod}})}$ -action on $\partial_\infty X$ preserves positions relative to faces of $\alpha(\sigma_{\text{mod}})$, we also have

$$\text{pos}(\alpha(\bar{\nu}), \alpha(\tau_{\text{mod}})) = \text{pos}(\bar{\alpha}(\bar{\nu}), \tau_{\text{mod}}).$$

The assertion follows. □

Lemma 3.12 (semicontinuity of relative position) *If $\xi_n \rightarrow \xi$ in $G\xi \subset \partial_\infty X$, and $\tau_n \rightarrow \tau$ in $\text{Flag}_{\tau_{\text{mod}}}$ are sequences such that the sequence of relative positions $\text{pos}(\xi_n, \tau_n)$ is constant, ie $\text{pos}(\xi_n, \tau_n) = p \in W_{\tau_{\text{mod}}} \setminus W/W_{\theta(v_\xi)}$ for all n , then $\text{pos}(\xi, \tau) \prec_{\tau_{\text{mod}}} p$.*

In particular, the sublevels of $\text{pos}(\cdot, \tau)$ in $G\xi$ are closed.

Proof There are apartment charts $\alpha_n: a_{\text{mod}} \rightarrow \partial_\infty X$ with $\alpha_n|_{\tau_{\text{mod}}} = \kappa_{\tau_n}$ and $\alpha_n(\bar{\xi}) = \xi_n$. The charts subconverge to a folding map α with $\alpha|_{\tau_{\text{mod}}} = \kappa_\tau$ and $\alpha(\bar{\xi}) = \xi$. The assertion follows from monotonicity; see Lemma 3.11(ii). \square

It follows that the *suplevels* $\{\text{pos}(\cdot, \tau) \succ_{\tau_{\text{mod}}} p\}$ in $G\xi$ are *open*, because their complements are finite unions of sublevels $\{\text{pos}(\cdot, \tau) \prec_{\tau_{\text{mod}}} p'\}$.

We show now that the folding order coincides with the inclusion order on Schubert cycles.

We start with the chamber case, where the relation between closures and the Bruhat order is well known. In the case of complex Lie groups, it goes back to the work of Chevalley in the 1950s [7]; for the proofs in the general case (including reductive groups over local fields), see [3] and [31]. (We are grateful to James Humphreys and Shrawan Kumar for the references.)

Proposition 3.13 *For a chamber $\sigma \subset \partial_\infty X$ and ideal points ξ_1, ξ_2 in the same G -orbit $G\xi \subset \partial_\infty X$, we have*

$$\text{pos}(\xi_1, \sigma) \prec \text{pos}(\xi_2, \sigma) \iff B_\sigma \xi_1 \subseteq \overline{B_\sigma \xi_2}.$$

Proof We denote by $\bar{\xi}_i \in a_{\text{mod}}$ the point of type $\theta(\xi)$ with $\text{pos}(\bar{\xi}_i, \sigma_{\text{mod}}) = \text{pos}(\xi_i, \sigma)$.

Suppose first that $\xi_1 \in \overline{B_\sigma \xi_2}$. Then there exists a sequence (b_n) in B_σ such that $b_n \xi_2 \rightarrow \xi_1$. Let a_n be apartments containing σ and $b_n \xi_2$, and let $\alpha_n: a_{\text{mod}} \rightarrow a_n$ be the apartment charts which restrict to the chamber chart of σ , ie $\alpha_n|_{\sigma_{\text{mod}}} = \kappa_\sigma: \sigma_{\text{mod}} \rightarrow \sigma$. Then $\alpha_n(\bar{\xi}_2) = b_n \xi_2$. The Tits isometric embeddings α_n subconverge (with respect to the visual topology) to a limit map $\alpha: a_{\text{mod}} \rightarrow \partial_\infty X$. The map α is, in general, not an isometric embedding (chart), but only a folding map extending κ_σ . It satisfies $\alpha(\bar{\xi}_2) = \xi_1$. Monotonicity yields $\text{pos}(\xi_1, \sigma) \prec \text{pos}(\bar{\xi}_2, \sigma_{\text{mod}}) = \text{pos}(\xi_2, \sigma)$; see Lemma 3.11(ii).

Vice versa, suppose now that $\bar{\xi}_1 \prec \bar{\xi}_2$. By definition of the partial order \prec there exists a folding map of a_{mod} fixing σ_{mod} and carrying $\bar{\xi}_2 \mapsto \bar{\xi}_1$. Furthermore there is an isometric embedding $a_{\text{mod}} \rightarrow \partial_\infty X$ which extends the chamber chart κ_σ and maps $\bar{\xi}_1 \mapsto \xi_1$.

By composition we obtain a folding map $\alpha: a_{\text{mod}} \rightarrow \partial_\infty X$ which extends κ_σ and maps $\alpha(\bar{\xi}_2) = \xi_1$. We want to find a sequence of isometric embeddings $\alpha_n: a_{\text{mod}} \rightarrow \partial_\infty X$ extending κ_σ such that $\alpha_n(\bar{\xi}_2) \rightarrow \alpha(\bar{\xi}_2) = \xi_1$. This will then imply that $\xi_1 \in \overline{B_\sigma \bar{\xi}_2}$. (Note that in general folding maps are not limits of isometric embeddings.)

We may assume that the relative positions $\bar{\xi}_i$ are regular. (Otherwise, we may perturb them keeping the inequality $\bar{\xi}_1 < \bar{\xi}_2$ and perturb the ξ_i accordingly.) We choose in a_{mod} a geodesic $\bar{\gamma}$ of length π starting in an interior point $\bar{\eta}_0$ of σ_{mod} and passing through $\bar{\xi}_2$ while avoiding simplices of codimension ≥ 2 . It crosses successively a sequence (gallery) of chambers $\bar{\sigma}_0 = \sigma_{\text{mod}}, \bar{\sigma}_1, \dots, \bar{\sigma}_k = \widehat{\sigma_{\text{mod}}}$ and intersects the intermediate panels $\bar{\tau}_i = \bar{\sigma}_i \cap \bar{\sigma}_{i-1}$ transversally in interior points $\bar{\eta}_i$. When applying the folding map α , it may happen that successive chambers of the folded gallery coincide, ie that $\alpha(\bar{\sigma}_i) = \alpha(\bar{\sigma}_{i-1})$ for some i . (This happens if and only if α is not an isometric embedding.) One can arbitrarily well approximate (in the visual topology) the folded gallery by an embedded gallery with the same initial chamber σ . To obtain such approximations it is convenient to use the G -action as follows. If $\alpha(\bar{\sigma}_i) = \alpha(\bar{\sigma}_{i-1})$ then one may pick an element $g \in G$ close to the identity, which fixes $\alpha(\bar{\tau}_i)$ and moves $\alpha(\bar{\sigma}_i) = \alpha(\bar{\sigma}_{i-1})$ away from itself, and apply it to the “tail” $\alpha(\bar{\sigma}_i), \dots, \alpha(\bar{\sigma}_k)$ of the gallery. Doing this inductively along the gallery, one obtains an arbitrarily good approximation of the folded gallery $\alpha(\bar{\sigma}_0) = \sigma, \dots, \alpha(\bar{\sigma}_k)$ by an embedded gallery $\sigma_0 = \sigma, \dots, \sigma_k$, that is a sequence of chambers such that $\sigma_i \cap \sigma_{i-1}$ is precisely a panel for all i . This yields at the same time an approximation of the broken geodesic $\alpha(\bar{\gamma})$ in $\partial_\infty X$ by a true geodesic γ such that $\gamma \cap \sigma_i$ and $\bar{\gamma} \cap \bar{\sigma}_i$ are corresponding subsegments of the same type. Now we use the path $\bar{\gamma}$ as a “guiding line” to extend the correspondence $\bar{\sigma}_i \mapsto \sigma_i$ of galleries to an isometric embedding $\alpha': a_{\text{mod}} \rightarrow \partial_\infty X$ extending κ_σ : Since γ connects two antipodal regular points there exists a unique such α' extending the isometry $\bar{\gamma} \rightarrow \gamma$ and hence mapping $\bar{\sigma}_i$ to σ_i . By construction, $\alpha'(\bar{\xi}_2)$ approximates $\alpha(\bar{\xi}_2) = \xi_1$ arbitrarily well. So we find a sequence of apartment charts α_n with the desired properties. \square

The proposition readily generalizes to the simplex case (in the case when G is a complex semisimple Lie group, a proof of the following proposition can be found in [32, Propage 3.13]):

Proposition 3.14 *For simplices $\tau \subset \partial_\infty X$ and ideal points ξ_1, ξ_2 in the same G -orbit $G\xi \subset \partial_\infty X$, we have*

$$\text{pos}(\xi_1, \tau) <_{\theta(\tau)} \text{pos}(\xi_2, \tau) \iff P_\tau \xi_1 \subseteq \overline{P_\tau \xi_2}.$$

Proof Let $\sigma \supset \tau$ be a chamber. Since the quotient space P_τ/B_σ is compact (it is the space of chambers containing τ as a face), the condition $\xi_1 \in \overline{P_\tau \xi_2}$ is equivalent to the existence of an element $p \in P_\tau$ such that $p\xi_1 \subseteq \overline{B_\sigma \xi_2}$. According to Proposition 3.13, this is equivalent to the existence of $p \in P_\tau$ such that $\text{pos}(p\xi_1, \sigma) \prec \text{pos}(\xi_2, \sigma)$. Since P_τ acts transitively on chambers containing τ , we have that $\bigcup_{p \in P_\tau} \text{pos}(p\xi_1, \sigma) = W_{\theta(\tau)} \text{pos}(\xi_1, \sigma) = \text{pos}(\xi_1, \tau)$. This completes the proof. \square

In other words, the proposition says that the τ -Schubert cycles in G/P_ν correspond to the sublevels of the folding order $\prec_{\tau_{\text{mod}}}$ on $W_{\tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}}$, where $\tau_{\text{mod}} = \theta(\tau)$ and $\nu_{\text{mod}} = \theta(\nu)$.

Recall that the simplices opposite to simplices of type τ_{mod} have type $\iota\tau_{\text{mod}}$, and that the action of w_0 induces the natural maps

$$W_{\tau_{\text{mod}}} \setminus a_{\text{mod}} \xrightarrow{w_0} W_{\iota\tau_{\text{mod}}} \setminus a_{\text{mod}},$$

and hence the maps

$$W_{\tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}} \xrightarrow{w_0} W_{\iota\tau_{\text{mod}}} \setminus W/W_{\nu_{\text{mod}}}$$

of relative positions; compare (4) and (5).

Definition 3.15 (complementary position) We define the *complementary position* by

$$\text{c-pos} := w_0 \text{ pos}.$$

This terminology is justified by the following result. (See Definition 2.3(ii) for the notion of antipodality.)

Lemma 3.16 Let $\tau, \hat{\tau}, \nu \subset \partial_\infty X$ be simplices contained in an apartment a , and suppose that τ and $\hat{\tau}$ are antipodal. Then $\text{pos}(\nu, \hat{\tau}) = \text{c-pos}(\nu, \tau)$.

Proof Let $\alpha: a_{\text{mod}} \rightarrow a$ be a chart such that $\alpha|_{\tau_{\text{mod}}} = \kappa_\tau$. Then $\hat{\tau} = \alpha(\hat{\tau}_{\text{mod}}) = (\alpha \circ w_0)(\iota\tau_{\text{mod}})$ with the simplex $\hat{\tau}_{\text{mod}} = w_0\iota\tau_{\text{mod}} \subset a_{\text{mod}}$ opposite to τ_{mod} . Using the reparametrized chart $\alpha \circ w_0$, we obtain $\text{pos}(\nu, \hat{\tau}) = (\alpha \circ w_0)^{-1}(\nu) = w_0\alpha^{-1}(\nu) = w_0 \text{ pos}(\nu, \tau)$. \square

The relation of ‘‘complementarity’’ is clearly symmetric: $\text{c-c-pos} = \text{pos}$. Passing to complementary relative position reverses the partial order (see Lemma 3.9):

$$(7) \quad \text{pos}(\xi_1, \tau) \prec_{\theta(\tau)} \text{pos}(\xi_2, \tau) \iff \text{c-pos}(\xi_1, \tau) \succ_{\iota\theta(\tau)} \text{c-pos}(\xi_2, \tau).$$

Points with smaller position relative to a simplex are closer to it in a *metric* sense. Namely, according to Lemma 3.4, we have the inequality of Tits distances:

$$(8) \quad \text{pos}(\xi_1, \tau) \prec_{\theta(\tau)} \text{pos}(\xi_2, \tau) \implies \angle_{\text{Tits}}(\xi_1, \cdot)|_{\tau} \leq \angle_{\text{Tits}}(\xi_2, \cdot)|_{\tau},$$

and respectively, for simplices τ_1 and τ_2 of the same type τ_{mod} ,

$$(9) \quad \text{pos}(\xi_1, \tau_1) \prec_{\tau_{\text{mod}}} \text{pos}(\xi_2, \tau_2) \implies \angle_{\text{Tits}}(\xi_1, \cdot) \circ \kappa_{\tau_1} \leq \angle_{\text{Tits}}(\xi_2, \cdot) \circ \kappa_{\tau_2}.$$

If τ_1 and τ_2 are simplices of opposite types, $\theta(\tau_1) = \iota\theta(\tau_2)$, and if the relative positions $\text{pos}(\xi_1, \tau_1)$ and $\text{pos}(\xi_2, \tau_2)$ are complementary, then

$$(10) \quad \angle_{\text{Tits}}(\xi_1, \cdot) \circ \kappa_{\tau_1} + \angle_{\text{Tits}}(\xi_2, \cdot) \circ \kappa_{\tau_2} \circ \iota \equiv \pi$$

on $\theta(\tau_1) = \iota\theta(\tau_2)$. To see this, note that the formula reduces to the case when the simplices τ_1 and τ_2 are opposite to each other, and $\xi_1 = \xi_2$ lies in an apartment containing them.

The following triangle inequality extends Lemma 3.16:

Lemma 3.17 *Let $\tau, \hat{\tau} \subset \partial_{\infty} X$ be a pair of antipodal simplices and let $\nu \subset \partial_{\infty} X$ be an arbitrary simplex. Then*

$$\text{pos}(\nu, \hat{\tau}) \succ_{\iota\theta(\tau)} \text{c-pos}(\nu, \tau),$$

*with equality if and only if $\tau, \hat{\tau}$ and ν are contained in an apartment.*²

Proof Let $a \subset \partial_{\infty} X$ be an apartment containing $\tau, \hat{\tau}$, and let

$$\partial_{\infty} X \xrightarrow{r} a$$

be a folding retraction, ie a folding map such that $r|_a = \text{id}_a$. Such a retraction is given eg by the natural projection $\partial_{\infty} X \rightarrow \partial_{\infty} X/B_{\sigma} \cong a$ for a chamber $\sigma \subset a$. By monotonicity (see Lemma 3.11(i)), we have

$$\text{pos}(\nu, \tau) \succ_{\theta(\tau)} \text{pos}(r\nu, \tau)$$

and

$$\text{pos}(\nu, \hat{\tau}) \succ_{\iota\theta(\tau)} \text{pos}(r\nu, \hat{\tau}) = \text{c-pos}(r\nu, \tau);$$

see Lemma 3.16. Since complementing position reverses the folding order (see (7)), we obtain the desired inequality.

²Equivalently, ν lies in the spherical sub-building $B(\tau, \hat{\tau}) \subset \partial_{\infty} X$ consisting of all apartments containing $\tau, \hat{\tau}$.

Suppose that equality holds, $\text{pos}(\nu, \widehat{\tau}) = \text{c-pos}(\nu, \tau)$. Let $\xi, \widehat{\xi}, \eta$ be interior points of $\tau, \widehat{\tau}, \nu$ such that $\xi, \widehat{\xi}$ are antipodal. Then (see Lemma 3.16 again)

$$\angle_{\text{Tits}}(\xi, \eta) + \angle_{\text{Tits}}(\eta, \widehat{\xi}) = \pi;$$

ie η lies on a geodesic segment $\widehat{\xi}\widehat{\xi}$. It follows that there exists an apartment containing $\xi, \widehat{\xi}, \eta$ and hence also the simplices spanned by these points. □

3.4 Thickenings

3.4.1 Thickenings in the Weyl group A *thickening* (of the neutral element) in W is a subset

$$\text{Th} \subset W$$

which is a union of sublevels for the folding order, ie which contains with every element w also every element w' satisfying $w' < w$. In the theory of posets, such subsets are called *ideals*.

Unions and intersections of thickenings are again thickenings, and removing a maximal element from a thickening yields a thickening. Furthermore, note that

$$\text{Th}^c := w_0(W - \text{Th}) = W - w_0\text{Th}$$

is again a thickening, because left multiplication with w_0 reverses the folding order; see Lemma 3.9. It holds that

$$W = \text{Th} \sqcup w_0\text{Th}^c,$$

and we therefore call Th^c the thickening *complementary* to Th .

Definition 3.18 (fat and slim) The thickening $\text{Th} \subset W$ is called *fat* if $\text{Th} \cup w_0\text{Th} = W$, equivalently, $\text{Th} \supseteq \text{Th}^c$. It is called *slim* if $\text{Th} \cap w_0\text{Th} = \emptyset$, equivalently, $\text{Th} \subseteq \text{Th}^c$. It is called *balanced* if it is both fat and slim, equivalently, $\text{Th} = \text{Th}^c$.

For types $\overline{\vartheta}_0, \overline{\vartheta} \in \sigma_{\text{mod}}$ and a radius $r \in [0, \pi]$, we define the *metric thickening*

$$(11) \quad \text{Th}_{\overline{\vartheta}_0, \overline{\vartheta}, r} := \{w \in W : d(w\overline{\vartheta}, \overline{\vartheta}_0) \leq r\},$$

using the natural W -invariant spherical metric d on a_{mod} . It is indeed a thickening by Lemma 3.4.

Recall that for a face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we denote by $W_{\tau_{\text{mod}}}$ its stabilizer in W . Furthermore, $\iota = -w_0: \sigma_{\text{mod}} \rightarrow \sigma_{\text{mod}}$ denotes the canonical involution of the model spherical Weyl chamber.

Lemma 3.19 (i) If $\bar{\vartheta}_0 \in \tau_{\text{mod}}$, then $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}$ is $W_{\tau_{\text{mod}}}$ -left invariant, that is,

$$W_{\tau_{\text{mod}}} \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r} = \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}.$$

(ii) If also $\iota \bar{\vartheta}_0 = \bar{\vartheta}_0$, then $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}$ is fat for $r \geq \frac{\pi}{2}$ and slim for $r < \frac{\pi}{2}$.

(iii) If, in addition, $d(w\bar{\vartheta}, \bar{\vartheta}_0) \neq \frac{\pi}{2}$ for all $w \in W$, then $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}$ is balanced.

Proof (i) For $w' \in W_{\tau_{\text{mod}}}$, we have that $w'\bar{\vartheta}_0 = \bar{\vartheta}_0$, and hence

$$d(w'w\bar{\vartheta}, \bar{\vartheta}_0) = d(w\bar{\vartheta}, \underbrace{w'^{-1}\bar{\vartheta}_0}_{\bar{\vartheta}_0}).$$

(ii) Since $w_0\bar{\vartheta}_0 = -\iota\bar{\vartheta}_0 = -\bar{\vartheta}_0$, we have

$$d(w_0w\bar{\vartheta}, -\bar{\vartheta}_0) = d(w\bar{\vartheta}, \underbrace{-w_0\bar{\vartheta}_0}_{\bar{\vartheta}_0})$$

and

$$d(w_0w\bar{\vartheta}, \bar{\vartheta}_0) = \pi - d(w\bar{\vartheta}, \bar{\vartheta}_0).$$

Hence

$$\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}^c = W - w_0\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r} := \{w \in W : d(w\bar{\vartheta}, \bar{\vartheta}_0) < \pi - r\},$$

which yields the assertion.

(iii) Slimness holds because $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2} = \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}$ for radii r slightly below $\frac{\pi}{2}$. \square

The metric examples provide balanced thickenings with arbitrary left invariance:

Corollary 3.20 (existence of balanced thickenings, I) For every ι -invariant face type τ_{mod} , there exists a $W_{\tau_{\text{mod}}}$ -left invariant balanced thickening $\text{Th} \subset W$.

Proof Since $\iota\tau_{\text{mod}} = \tau_{\text{mod}}$, there exists $\bar{\vartheta}_0 \in \tau_{\text{mod}}$ such that $\iota\bar{\vartheta}_0 = \bar{\vartheta}_0$. Moreover, the set of types $\bar{\vartheta} \in \sigma_{\text{mod}}$ such that $d(\cdot\bar{\vartheta}, \bar{\vartheta}_0) \neq \frac{\pi}{2}$ on W is the complement of a finite union of great spheres in a_{mod} , and hence open and dense. \square

In order to obtain balanced thickenings with additional right invariance, we modify the metric thickenings (11) at their “boundaries”. The rigidity part of Lemma 3.4 implies that the elements of

$$\partial \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r} := \{w \in W : d(w\bar{\vartheta}, \bar{\vartheta}_0) = r\}$$

are pairwise \prec -incomparable and maximal in $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}$. Thus every subset $\text{Th} \subset W$ with

$$\{w \in W : d(w\bar{\vartheta}, \bar{\vartheta}_0) < r\} \subseteq \text{Th} \subseteq \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, r}$$

is a thickening.

Using these modified metric thickenings, we can generalize our last existence result:

Proposition 3.21 (existence of balanced thickenings, II) *Let $\tau_{\text{mod}}, \nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be face types and suppose that τ_{mod} is ι -invariant. Then a $W_{\tau_{\text{mod}}}$ -left invariant and $W_{\nu_{\text{mod}}}$ -right invariant balanced thickening $\text{Th} \subset W$ exists if and only if (left multiplication by) w_0 has no fixed point on $W_{\tau_{\text{mod}}} \backslash W / W_{\nu_{\text{mod}}}$; see (5).*

Proof If a balanced thickening exists, then w_0 cannot have a fixed point as a consequence of the definition of balancedness.

Vice versa, let us assume that w_0 has no fixed point. We choose $\bar{\vartheta}_0 \in \tau_{\text{mod}}$ with $\iota \bar{\vartheta}_0 = \bar{\vartheta}_0$ and $\bar{\nu} \in \nu_{\text{mod}}$. Then the fat thickening $\text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$ is $W_{\tau_{\text{mod}}}$ -left and $W_{\nu_{\text{mod}}}$ -right invariant, and so is the “great sphere” $\partial \text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$. The latter is moreover preserved by the involution w_0 while the “hemispheres bounded by it”, $\text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2} - \partial \text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$ and $W - \text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$, are exchanged; see the proof of Lemma 3.19(ii). Since w_0 has no fixed point, $\partial \text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$ decomposes as a union of pairs of double cosets $W_{\tau_{\text{mod}}} w W_{\nu_{\text{mod}}}$ which are swapped by w_0 . By removing from $\text{Th}_{\bar{\vartheta}_0, \bar{\nu}, \pi/2}$ one double coset of each pair, we therefore obtain a balanced thickening as desired. \square

Corollary 3.22 *If $w_0 = -\text{id}_{a_{\text{mod}}}$, then a $W_{\nu_{\text{mod}}}$ -right invariant balanced thickening exists for every face type ν_{mod} .*

Proof We equivariantly identify the coset space $W / W_{\nu_{\text{mod}}}$ with an orbit $W\bar{\nu} \subset a_{\text{mod}}$ for some $\bar{\nu} \in \text{int}(\nu_{\text{mod}})$. By assumption, w_0 has no fixed point on a_{mod} , and hence none on $W / W_{\nu_{\text{mod}}}$. \square

Remark 3.23 (i) Note that $w_0 = -\text{id}_{a_{\text{mod}}}$ if and only if all irreducible factors of W are of type $A_1, B_{n \geq 2}, D_{2k \geq 4}, E_{7,8}, F_4$ or G_2 ; see [4].

(ii) $W_{\tau_{\text{mod}}}$ -left and $W_{\nu_{\text{mod}}}$ -right invariant balanced thickenings do not always exist. For instance, in the B_2 -case there are no $W_{\tau_{\text{mod}}}$ -biinvariant thickenings for τ_{mod} a vertex type.

In rank two, the balanced thickenings are easy to describe:

Example 3.24 (balanced thickenings in rank 2) (i) If $W = W_{A_2}$, then σ_{mod} is an arc of length $\frac{\pi}{3}$. There is a unique balanced thickening $\text{Th} \subset W$ described by the property that $\text{Th} \cdot \sigma_{\text{mod}} \subset a_{\text{mod}}$ is the $\frac{\pi}{2}$ -ball centered at the midpoint of σ_{mod} .

(ii) If $W = W_{B_2}$ or W_{G_2} , then σ_{mod} is an arc of length $\frac{\pi}{4}$ or $\frac{\pi}{6}$. In these cases, there are two balanced thickenings. Namely, for each vertex $\bar{\xi}$ of σ_{mod} we have the $W_{\bar{\xi}}$ -left invariant thickening $\text{Th} \subset W$ for which $\text{Th} \cdot \sigma_{\text{mod}} = \bar{B}(\bar{\xi}, \frac{\pi}{2})$.

Below, we give two examples in higher rank. First we address the irreducible case:

Example 3.25 (some balanced thickenings of type A_n) The spherical Coxeter complex a_{mod} can be modeled as the unit sphere in the hyperplane

$$x_0 + \dots + x_n = 0$$

in \mathbb{R}^{n+1} . The Weyl group $W \cong S_{n+1}$ acts by permuting the coordinates, and we choose the fundamental chamber $\sigma_{\text{mod}} \subset a_{\text{mod}}$ as given by the inequalities $x_0 \geq \dots \geq x_n$. It holds that

$$(x_0, \dots, x_n) \xrightarrow{w_0 = -\iota} (x_n, \dots, x_0).$$

There are the ι -invariant edge midpoints $\bar{\vartheta}_k \in \sigma_{\text{mod}}$ for $1 \leq k \leq \frac{1}{2}n$ represented by the vectors

$$\underbrace{(1, \dots, 1, 0, \dots, 0)}_k, \underbrace{(-1, \dots, -1)}_k$$

and the unique ι -invariant vertex $\bar{\vartheta}_{(n+1)/2} \in \sigma_{\text{mod}}$ if n is odd. The type $\bar{\vartheta}_1$ is the *unique root type*.

In incidence geometric terms, the Coxeter complex a_{mod} is the spherical building associated to the finite projective n -space $\mathbb{P}_{\text{mod}}^n = \{e_0, \dots, e_n\}$ consisting of $n + 1$ points. Every subset is a projective subspace (of dimension one less than the number of points in it) and corresponds to a vertex of a_{mod} . Vertices are adjacent if and only if the corresponding subspaces are incident, ie one contains the other. The element $w_\pi \in W$ corresponding to the permutation $\pi \in S_{n+1}$ acts by $e_i \xrightarrow{w_\pi} e_{\pi(i)}$. We let the fundamental chamber $\sigma_{\text{mod}} \subset a_{\text{mod}}$ correspond to the full flag

$$\{e_0\} \subset \dots \subset \{e_0, \dots, e_i\} \subset \dots \subset \{e_0, \dots, e_{n-1}\}.$$

The edge spanned by $\bar{\vartheta}_k$ then has the vertices $\{e_0, \dots, e_{k-1}\}$ and $\{e_0, \dots, e_{n-k}\}$.

We determine the metric thickenings $\text{Th}_{\bar{\vartheta}_1, \bar{\vartheta}, \pi/2}$ for regular types $\bar{\vartheta} \in \text{int}(\sigma_{\text{mod}})$: The type $\bar{\vartheta}$ is represented by a vector (t_i) with $t_0 > \dots > t_n$. The element $w_\pi \in W$ carries (t_i) to the vector $(t_{\pi^{-1}(i)})$. Thus, $\angle_{\text{Tits}}(w_\pi \bar{\vartheta}, \bar{\vartheta}_1) < \frac{\pi}{2}$ if and only if $t_{\pi^{-1}(0)} > t_{\pi^{-1}(n)}$ if and only if $\pi^{-1}(0) < \pi^{-1}(n)$, and we obtain the balanced thickening

$$(12) \quad \text{Th}_{\bar{\vartheta}_1, \bar{\vartheta}, \pi/2} = \{w_\pi \in W : \pi^{-1}(0) < \pi^{-1}(n)\}.$$

Similarly, one can describe the thickenings $\text{Th}_{\bar{\vartheta}_k, \bar{\vartheta}, \pi/2}$ for $k \geq 2$. They depend on the type $\bar{\vartheta}$ and are balanced for a dense open set of values.

To give an incidence geometric description of the thickening (12), note that $\pi^{-1}(i)$ is the dimension of the smallest subspace in the flag $w_\pi \sigma_{\text{mod}}$ which contains e_i . Hence $w_\pi \in \text{Th}_{\bar{\vartheta}_1, \bar{\vartheta}, \pi/2}$ if and only if to the flag $w_\pi \sigma_{\text{mod}}$ belongs a subspace U which contains e_0 but not e_n , equivalently,

$$(13) \quad \{e_0\} \subseteq U \subseteq \{e_0, \dots, e_{n-1}\}.$$

Another interesting example is the $W_{\bar{\vartheta}_m}$ -biinvariant thickening $\text{Th}_{\bar{\vartheta}_m, \bar{\vartheta}_m, \pi/2}$ for $n = 2m - 1$. Observe that $\angle_{\text{Tits}}(w_\pi \bar{\vartheta}_m, \bar{\vartheta}_m) \leq \frac{\pi}{2}$ if and only if

$$(14) \quad |w_\pi \{e_0, \dots, e_{m-1}\} \cap \{e_0, \dots, e_{m-1}\}| \geq \frac{1}{2}m,$$

with the equality cases corresponding to each other. Equality cannot occur if m is odd, and in this case the thickening $\text{Th}_{\bar{\vartheta}_m, \bar{\vartheta}_m, \pi/2}$ is balanced.

Remark 3.26 With a bit more work one can classify all balanced thickenings in the A_3 case: There are 10 balanced thickenings. Two of them are $W_{\bar{\vartheta}_2}$ -left invariant for the unique ι -invariant vertex $\bar{\vartheta}_2 \in \sigma_{\text{mod}}$, and one is $W_{\epsilon_{\text{mod}}}$ -left invariant for the unique ι -invariant edge $\epsilon_{\text{mod}} \subset \sigma_{\text{mod}}$.

The next example concerns the reducible case:

Example 3.27 (some balanced thickenings of type A_1^n) The spherical Coxeter complex a_{mod} can be modeled as the unit sphere in \mathbb{R}^n . The Weyl group $W \cong \mathbb{Z}_2^n \cong \{\pm 1\}^n$ acts by changing the signs of the coordinates x_i ; ie its canonical generators act by reflections at the coordinate hyperplanes. We choose the fundamental chamber $\sigma_{\text{mod}} \subset a_{\text{mod}}$ as given by the inequalities $x_1, \dots, x_n \geq 0$.

The longest element $w_0 = (-1, \dots, -1)$ acts as $-\text{id}$. The Bruhat order on W is given by

$$w_\epsilon \prec w_{\epsilon'} \iff \epsilon_i \geq \epsilon'_i \text{ for all } i,$$

where we denote the elements in W by w_ϵ with $\epsilon = (\epsilon_i)$.

The $(k-1)$ -simplices of the spherical Coxeter complex a_{mod} correspond to the $\{\pm 1\}$ -valued maps defined on subsets of $\{1, \dots, n\}$ of cardinality k . In particular, the chambers can be interpreted as the ordered n -point configurations on $\{\pm 1\}$.

Let $\bar{\zeta} \in \sigma_{\text{mod}}$ be the *central* type represented by the vector $(1, \dots, 1)$. We determine the metric thickenings of the form $\text{Th}_{\bar{\zeta}, \bar{\vartheta}, \pi/2}$ for the regular types $\bar{\vartheta} \in \text{int}(\sigma_{\text{mod}})$: The type $\bar{\vartheta}$ is represented by a vector $t = (t_i)$ with $t_i > 0$. The element $w_\epsilon \in W$ carries t to the vector $(\epsilon_i t_i)$. Thus $\angle_{\text{Tits}}(w_\epsilon \bar{\vartheta}, \bar{\zeta}) < \frac{\pi}{2}$ if and only if $\epsilon \cdot t = \epsilon_1 t_1 + \dots + \epsilon_n t_n > 0$, and

$$\text{Th}_t = \{w_\epsilon \in W : \epsilon \cdot t > 0\}, \quad \overline{\text{Th}}_t = \{w_\epsilon \in W : \epsilon \cdot t \geq 0\}$$

are *metric thickenings*. The thickening Th_t is slim, while $\overline{\text{Th}}_t$ is fat. We have that $\overline{\text{Th}}_t = \text{Th}_t$ is balanced if and only if $\epsilon \cdot t \neq 0$ for all sign choices ϵ , which is the case for “generic” values of t .

To phrase it in terms of configurations, we consider *weighted n -point configurations* on $\{\pm 1\}$ with weights t_i . Then the thickenings $\overline{\text{Th}}_t$ and Th_t correspond to the sets of configurations with at least, respectively, strictly more than half of the total mass placed on $+1$.

3.4.2 Thickenings at infinity From thickenings in the Weyl group, we derive *thickenings at infinity* as follows.

Given a $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the induced thickening of a simplex $\tau \in \text{Flag}_{\tau_{\text{mod}}}$ inside the Furstenberg boundary

$$\text{Th}_{\text{Fü}}(\tau) := \{\text{pos}(\cdot, \tau) \in W_{\tau_{\text{mod}}} \setminus \text{Th}\} \subset \partial_{\text{Fü}} X$$

is well defined. Furthermore, we define the thickening of τ inside the visual boundary as the union of the corresponding (closed) chambers

$$(15) \quad \text{Th}(\tau) := \bigcup_{\sigma \in \text{Th}_{\text{Fü}}(\tau)} \sigma \subset \partial_\infty X.$$

Due to the semicontinuity of relative position (see Lemma 3.12), the thickenings $\text{Th}_{\text{Fü}}(\tau)$ and $\text{Th}(\tau)$ are compact. The intersections $\text{Th}(\tau) \cap G\xi$ with the G -orbits $G\xi \subset \partial_\infty X$ are finite unions of Schubert cycles and hence projective subvarieties. For regular G -orbits $G\xi$, the intersection $\text{Th}(\tau) \cap G\xi$ is naturally identified with $\text{Th}_{\text{Fü}}(\tau)$.

Note that if the thickening $\text{Th} \subset W$ is $W_{\nu_{\text{mod}}}$ -right invariant for a face type $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$, then the thickenings of simplices are *unions of stars* of simplices of type ν_{mod} .

For a subset $A \subset \text{Flag}_{\tau_{\text{mod}}}$, we define the induced thickenings

$$\text{Th}_{\text{Fü}}(A) = \bigcup_{\tau \in A} \text{Th}_{\text{Fü}}(\tau) \quad \text{and} \quad \text{Th}(A) = \bigcup_{\tau \in A} \text{Th}(\tau).$$

If A is compact, then its thickenings are compact as well.

Below are several examples of thickenings, based on the examples in the previous section.

Example 3.28 (rank 2) We continue with Example 3.24.

(i) If $W = W_{A_2}$, then chambers in the visual boundary are arcs of Tits length $\frac{\pi}{3}$. For the unique balanced thickening $\text{Th} \subset W$, the associated thickening $\text{Th}(\sigma) \subset \partial_\infty X$ of a chamber $\sigma \subset \partial_\infty X$ with midpoint ζ is the ball $\bar{B}(\zeta, \frac{\pi}{2})$. In incidence geometric terms, regarding $\partial_{\text{Tits}} X$ as the spherical building associated to a projective plane Π , the chamber σ corresponds to a flag (l, p) consisting of a line $l \subset \Pi$ and a point $p \in l$. The thickening $\text{Th}_{\text{Fü}}(\sigma) \subset \partial_{\text{Fü}} X$ inside the Furstenberg boundary consists of all flags (l', p') such that $l' = l$ or $p' = p$.

(ii) If $W = W_{B_2}$ or W_{G_2} , then chambers have length $\frac{\pi}{4}$ or $\frac{\pi}{6}$. For a vertex type $\bar{\xi} \in \sigma_{\text{mod}}$ and the unique $W_{\bar{\xi}}$ -left invariant balanced thickening $\text{Th} \subset W$, the associated thickening of a vertex $\xi \in \partial_\infty X$ of type $\theta(\xi) = \bar{\xi}$ inside $\partial_\infty X$ is given by $\text{Th}(\xi) = \bar{B}(\xi, \frac{\pi}{2})$. The thickening of a chamber $\sigma \subset \partial_\infty X$ equals $\text{Th}(\sigma) = \bar{B}(\xi_\sigma, \frac{\pi}{2})$ where ξ_σ is the vertex of σ with type $\bar{\xi}$.

For instance, if $G = O(n, 2)$ with $n \geq 2$ and hence X has type B_2 , then $\partial_{\text{Tits}} X$ can be regarded from the incidence geometry perspective as the spherical building arising from isotropic flags in $\mathbb{R}^{n,2} = \mathbb{R}^n \oplus \mathbb{R}^2$. A chamber corresponds to a flag (L, P) consisting of an isotropic plane $P \subset \mathbb{R}^{n,2}$ and a(n isotropic) line $L \subset P$. If $\bar{\xi}$ is the vertex type corresponding to isotropic *lines*, then the thickening $\text{Th}_{\text{Fü}}(L) \subset \partial_{\text{Fü}} X$ of an isotropic line $L \in \text{Flag}_{\bar{\xi}}$ consists of all flags (L', P') such that $P' \supset L$. On the other hand, if $\bar{\xi}$ is the vertex type corresponding to isotropic *planes*, then the thickening $\text{Th}_{\text{Fü}}(P)$ of an isotropic plane $P \in \text{Flag}_{\bar{\xi}}$ consists of all flags $(L', P') \in \partial_{\text{Fü}} X$ such that $L' \subset P$.

Example 3.29 (type A_n) We continue with Example 3.25.

Let $G = \text{SL}(n + 1, \mathbb{F})$. We regard $\partial_{\text{Tits}} X$ as the spherical building associated to the projective n -space $\mathbb{F}P^n$.

Let $\tau_{\text{mod}}(\bar{\vartheta}_1) \in \sigma_{\text{mod}}$ denote the edge type with midpoint $\bar{\vartheta}_1$. Then $\text{Flag}_{\tau_{\text{mod}}(\bar{\vartheta}_1)} = \mathcal{F}_{1,n}$, the manifold of 2-flags (L, H) consisting of a hyperplane $H \subset \mathbb{F}^{n+1}$ and a line $L \subset H$. For the $W_{\tau_{\text{mod}}(\bar{\vartheta}_1)}$ -left invariant balanced thickening $\text{Th}_{\bar{\vartheta}_1, \bar{\vartheta}, \pi/2} \subset W$ given by (12), the thickening $\text{Th}_{\bar{\vartheta}_1, \bar{\vartheta}, \pi/2}^{\text{Fü}}((L, H))$ of the flag $(L, H) \in \mathcal{F}_{1,n}$ in the full flag manifold $\partial_{\text{Fü}} X$ consists of all flags $U_1 \subset \dots \subset U_i \subset \dots \subset U_n$ in \mathbb{F}^{n+1} such that (compare (13))

$$L \subseteq U_i \subseteq H \quad \text{for some } i.$$

If $n = 4l + 1$, then $\text{Flag}_{\bar{\vartheta}_{2l+1}} = \mathcal{F}_{2l+1}$ is the middle Grassmannian of $(2l+2)$ -dimensional linear subspaces of \mathbb{F}^{n+1} . The balanced $W_{\bar{\vartheta}_{2l+1}}$ -biinvariant thickening $\text{Th}_{\bar{\vartheta}_{2l+1}, \bar{\vartheta}_{2l+1}, \pi/2}^{\text{Fü}}(U)$ of a subspace $U \in \mathcal{F}_{2l+1}$, inside \mathcal{F}_{2l+1} , consists of all subspaces $U' \in \mathcal{F}_{2l+1}$ such that (compare (14))

$$\dim(U' \cap U) \geq l + 1.$$

Example 3.30 (type A_1^n , configuration spaces and stability in the sense of geometric invariant theory) We continue with Example 3.27.

Let $X = Y^n$ be the n -fold product of a rank-one symmetric space Y , eg $Y = \mathbb{H}^2$. Then $\partial_{\text{Fü}} X \cong (\partial_\infty Y)^n$, and we will view chambers as ordered n -point configurations $\xi = (\eta_i)$ on $\partial_\infty Y$. The relative position of two configurations $\xi, \xi' \in \partial_{\text{Fü}} X$ is given by

$$\text{pos}(\xi', \xi) = w_\epsilon \quad \text{with} \quad \epsilon_i = +1 \iff \eta'_i = \eta_i.$$

Thus, it records the entries i where the configurations agree.

We fix a regular vector $t = (t_i) \in \text{int}(\Delta) \cong \mathbb{R}_+^n$ and assign the weight $t_i > 0$ to the i^{th} point of a configuration. Then a chamber ξ , regarded now as a *weighted configuration* on $\partial_\infty Y$, defines the finite measure

$$\mu_\xi = t_1 \delta_{\eta_1} + \dots + t_n \delta_{\eta_n}$$

on $\partial_\infty Y$, where δ_{η_i} denotes the Dirac measure concentrated in the point η_i . (Masses add when points η_i “collide”). The total mass $|\mu_\xi|$ of μ_ξ equals

$$M = t_1 + \dots + t_n.$$

In the language of geometric invariant theory, the finite measure μ_ξ (and the corresponding weighted configuration ξ) is called *stable* if $\mu_\xi(\eta) < \frac{1}{2}M$ for all points $\eta \in \partial_\infty Y$, and *semistable* if $\mu_\xi(\eta) \leq \frac{1}{2}M$ for all η .

Let $(\eta) := (\eta, \dots, \eta)$ denote the configuration concentrated in the point η . According to Example 3.27, the thickenings $(\overline{\text{Th}}_t)_{\text{Fü}}((\eta))$ and $(\text{Th}_t)_{\text{Fü}}((\eta))$ of (η) consist of the weighted configurations where at least, respectively, strictly more than half of the total mass is concentrated in the point η .

Choose now $A \subset \partial_{\text{Fü}} X$ as the “diagonal”, that is, as the compact antipodal subset of all configurations (η) concentrated in one point. Then the thickenings $(\text{Th}_t)_{\text{Fü}}(A)$ and $(\overline{\text{Th}}_t)_{\text{Fü}}(A)$ of A inside $\partial_{\text{Fü}} X$ equal the subsets of weighted configurations which are *not semistable*, respectively, *not stable*. In the case when Th_t is balanced, both notions agree: “stable=semistable”.

The sets of stable and semistable configurations depend on the weights t . For instance, if $t_i > \frac{1}{2}M$ for some i , then there are no semistable weighted configurations, equivalently,

$$(\text{Th}_t)_{\text{Fü}}(A) = \partial_{\text{Fü}}X.$$

For instance, for $n = 2$ and any $t_1 \neq t_2$, there are no semistable configurations.

In contrast, if $n \geq 3$ and $t_i < \frac{1}{2}M$ for all i , then there are always stable configurations, for instance, the configurations where no two point coincide. Equivalently, in this case,

$$(\text{Th}_t)_{\text{Fü}}(A) \neq \partial_{\text{Fü}}X.$$

We return to the general discussion of thickenings in $\partial_\infty X$. Our motivation for introducing the notion of slimness is the following observation:

Lemma 3.31 (disjointness of slim thickenings) (i) *Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be an ι -invariant face type, and let $\text{Th} \subset W$ be a slim $W_{\tau_{\text{mod}}}$ -left invariant thickening. Then for any two antipodal simplices $\tau, \hat{\tau} \in \text{Flag}_{\tau_{\text{mod}}}$, it holds that*

$$\text{Th}_{\text{Fü}}(\tau) \cap \text{Th}_{\text{Fü}}(\hat{\tau}) = \emptyset.$$

(ii) *More generally, suppose that $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$ is another face type and that the slim thickening Th is also $W_{\nu_{\text{mod}}}$ -right invariant. Then for any G -orbit $G\xi \subset \partial_\infty X$ of type $\bar{\xi} = \theta(\xi) \in \text{int}(\nu_{\text{mod}})$ and any two antipodal simplices $\tau, \hat{\tau} \in \text{Flag}_{\tau_{\text{mod}}}$, it holds that*

$$\text{Th}(\tau) \cap \text{Th}(\hat{\tau}) \cap G\xi = \emptyset.$$

Proof Part (i) follows from Lemma 3.17 and the definition of slimness. Indeed, suppose that $\text{Th}_{\text{Fü}}(\tau) \cap \text{Th}_{\text{Fü}}(\hat{\tau})$ contains a chamber σ . Then $\text{pos}(\sigma, \tau), \text{pos}(\sigma, \hat{\tau}) \in W_{\tau_{\text{mod}}} \setminus \text{Th}$. By Lemma 3.17, $\text{pos}(\sigma, \hat{\tau}) \succ_{\tau_{\text{mod}}} \text{c-pos}(\sigma, \tau)$. Hence also $\text{c-pos}(\sigma, \tau) \in W_{\tau_{\text{mod}}} \setminus \text{Th}$, equivalently, $\text{pos}(\sigma, \tau) \in W_{\tau_{\text{mod}}} \setminus w_0\text{Th}$. It follows that $\text{Th} \cap w_0\text{Th} \neq \emptyset$, contradicting slimness.

Part (ii) follows because the thickenings are unions of stars of simplices of type ν_{mod} . Indeed, by (i), the intersection $\text{Th}(\tau) \cap \text{Th}(\hat{\tau})$ contains no chamber, and hence it cannot contain the star of a simplex of type ν_{mod} . □

4 Asymptotic geometric notions in symmetric spaces

4.1 Shadows at infinity and strong asymptoticity of Weyl cones

For a simplex $\tau_- \subset \partial_\infty X$ of type $\iota\tau_{\text{mod}}$ and a point $x \in X$, we consider the function

$$(16) \quad \tau \mapsto d(x, P(\tau_-, \tau))$$

on the open Schubert stratum $C(\tau_-) \subset \text{Flag}_{\tau_{\text{mod}}}$. We denote by $\tau_+ \in C(\tau_-)$ the chamber x -opposite to τ_- .

Lemma 4.1 *The function (16) is continuous and proper.*

Proof This follows from the fact that $C(\tau_-)$ and X are homogeneous spaces for the parabolic subgroup P_{τ_-} . Indeed, continuity follows from the continuity of the function

$$g \mapsto d(x, P(\tau_-, g\tau_+)) = d(g^{-1}x, P(\tau_-, \tau_+))$$

on P_{τ_-} which factors through the orbit map $P_{\tau_-} \rightarrow C(\tau_-)$, $g \mapsto g\tau_+$.

Regarding properness, note that a simplex $\tau \in C(\tau_-)$ is determined by any point y contained in the parallel set $P(\tau_-, \tau)$, namely as the simplex y -opposite to τ_- . Thus, if $P(\tau_-, \tau) \cap B(x, R) \neq \emptyset$ for some fixed $R > 0$, then there exists $g \in P_{\tau_-}$ such that $\tau = g\tau_+$ and $d(x, gx) < R$. In particular, g is bounded. This implies properness. \square

Moreover, the function (16) has a unique minimum zero in τ_+ .

We define the following open subsets of $C(\tau_-)$ which can be regarded as *shadows* of balls in X with respect to τ_- . For $x \in X$ and $r > 0$, we put

$$U_{\tau_-,x,r} := \{\tau \in C(\tau_-) : d(x, P(\tau_-, \tau)) < r\}.$$

The next fact expresses the uniform strong asymptoticity of asymptotic Weyl cones.

Lemma 4.2 *For $r, R > 0$, there exists $d = d(r, R) > 0$ such that if $y \in V(x, \text{st}(\tau_-))$ with $d(y, \partial V(x, \text{st}(\tau_-))) \geq d(r, R)$, then $U_{\tau_-,x,R} \subset U_{\tau_-,y,r}$.*

Proof If $U_{\tau_-,x,R} \not\subset U_{\tau_-,y,r}$ there exists $x' \in B(x, R)$ such that $d(y, V(x', \text{st}(\tau_-))) \geq r$. Thus, if the assertion is wrong, there exist a sequence $x_n \rightarrow x_\infty$ in $\bar{B}(x, R)$ and a sequence $y_n \rightarrow \infty$ in $V(x, \text{st}(\tau_-))$ such that $d(y_n, \partial V(x, \text{st}(\tau_-))) \rightarrow +\infty$ and $d(y_n, V(x_n, \text{st}(\tau_-))) \geq r$.

Let $\rho: [0, +\infty) \rightarrow V(x, \tau_-)$ be a geodesic ray with initial point x and asymptotic to an interior point of τ_- . Then the sequence (y_n) eventually enters every Weyl cone $V(\rho(t), \text{st}(\tau_-))$. Since the distance function $d(\cdot, V(x_n, \text{st}(\tau_-)))$ is convex and bounded, and hence nonincreasing along rays asymptotic to $\text{st}(\tau_-)$, we have that

$$R \geq d(x, V(x_n, \text{st}(\tau_-))) \geq d(\rho(t), V(x_n, \text{st}(\tau_-))) \geq d(y_n, V(x_n, \text{st}(\tau_-))) \geq r$$

for $n \geq n(t)$. It follows that, for all t ,

$$R \geq d(\rho(t), V(x_\infty, \text{st}(\tau_-))) \geq r.$$

However, the ray ρ is strongly asymptotic to $V(x_\infty, \text{st}(\tau_-))$, a contradiction. □

4.2 Asymptotic properties of sequences and subgroups

We first consider sequences in the model euclidean Weyl chamber Δ .

Definition 4.3 We say that a sequence (δ_n) in Δ is

(i) τ_{mod} -pure if it is contained in a tubular neighborhood of the sector $V(0, \tau_{\text{mod}})$ and drifts away from its boundary $\partial V(0, \tau_{\text{mod}}) = V(0, \partial\tau_{\text{mod}})$, ie

$$d(\delta_n, V(0, \partial\tau_{\text{mod}})) \rightarrow +\infty;$$

(ii) τ_{mod} -regular if

$$d(\delta_n, V(0, \partial \text{st}(\tau_{\text{mod}}))) \rightarrow +\infty.$$

These properties are stable under bounded perturbation of the sequence, due to the triangle inequality $|d_\Delta(x, y) - d_\Delta(x', y')| \leq d(x, x') + d(y, y')$. Therefore, the following definitions for sequences in X and G are sensible:

Definition 4.4 (pure and weakly regular) (i) We say that a sequence (x_n) in X is τ_{mod} -pure, respectively, τ_{mod} -regular if for some (any) base point $o \in X$ the sequence of Δ -distances $d_\Delta(o, x_n)$ has this property.

(ii) We say that a sequence (g_n) in G is τ_{mod} -pure, respectively, τ_{mod} -regular if for some (any) point $x \in X$ the orbit sequence $(g_n x)$ in X has this property.

(iii) We say that a subgroup $\Gamma < G$ is τ_{mod} -regular if all sequences of pairwise distinct elements in Γ have this property.

The face type of a pure sequence is uniquely determined. Moreover, a τ_{mod} -regular sequence is τ'_{mod} -regular for every face type $\tau'_{\text{mod}} \subseteq \tau_{\text{mod}}$, because $\text{ost}(\tau'_{\text{mod}}) \supseteq \text{ost}(\tau_{\text{mod}})$.

Note that τ_{mod} -regular subgroups are in particular discrete. If $\text{rank}(X) = 1$, then discreteness is equivalent to (σ_{mod}) -regularity. In higher rank, regularity can be considered as a strengthening of discreteness: A discrete subgroup $\Gamma < G$ may not be τ_{mod} -regular for any face type τ_{mod} ; this can happen eg for free abelian subgroups of transvections of rank ≥ 2 .

Lemma 4.5 (i) τ_{mod} -pure sequences are τ_{mod} -regular.

(ii) Every sequence which diverges to infinity contains a τ_{mod} -pure subsequence for some face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$.

Proof Assertion (i) is a direct consequence of the definitions, and (ii) follows by induction on face types. □

Note also that a sequence which diverges to infinity is τ_{mod} -regular if and only if it contains ν_{mod} -pure subsequences only for face types $\nu_{\text{mod}} \supseteq \tau_{\text{mod}}$. (We will not use this fact.)

Remark 4.6 (relation to Finsler compactifications) There is a close relation between the regularity of sequences and the asymptotic geometry of certain G -invariant *Finsler metrics* on X ; see [16, Section 8.1.2]. For instance, a sequence in X is (σ_{mod}) -regular if and only if it accumulates at the Furstenberg boundary inside the regular Finsler compactification.

5 Some topological dynamics

5.1 (Proper) discontinuity and dynamical relation

Let Z be a compact metrizable space, and let $\Gamma \subset \text{Homeo}(Z)$ be a countably infinite subgroup. We consider the action $\Gamma \curvearrowright Z$.

Definition 5.1 (discontinuous) A point $z \in Z$ is called *wandering* with respect to the Γ -action if the action is *discontinuous* at z , ie if z has a neighborhood U such that $U \cap \gamma U \neq \emptyset$ for at most finitely many $\gamma \in \Gamma$.

Nonwandering points are called *recurrent*.

Definition 5.2 (domain of discontinuity) We call the set

$$\Omega_{\text{disc}} \subset Z$$

of wandering points the *wandering set* or *domain of discontinuity* for the action $\Gamma \curvearrowright Z$.

Note that Ω_{disc} is open and Γ -invariant.

Definition 5.3 (proper) The action of Γ on an open subset $U \subset Z$ is called *proper* if for every compact subset $K \subset U$, it holds that $K \cap \gamma K \neq \emptyset$ for at most finitely many $\gamma \in \Gamma$.

In particular, the action of Γ on U is then discontinuous, ie $U \subseteq \Omega_{\text{disc}}$, and is therefore called *properly discontinuous*.

Definition 5.4 (domain of proper discontinuity) If Γ is a group, we call a Γ -invariant open subset $\Omega \subseteq \Omega_{\text{disc}}$ on which Γ acts properly a *domain of proper discontinuity* for Γ .

The orbit space Ω/Γ is then Hausdorff. Note that in general there is *no unique* maximal *proper* domain of discontinuity.

Example 5.5 (nonunique maximal domain of proper discontinuity) Consider the infinite cyclic group $\Gamma \cong \mathbb{Z}$ acting projectively on $Z = \mathbb{R}P^2$, so that a generator γ of Γ acts as the projectivization of a diagonal matrix with distinct positive eigenvalues $\lambda_1 > \lambda_2 > \lambda_3$. Let $e_1, e_2, e_3 \in Z$ be the three fixed points of γ (eigenspaces for $\lambda_1, \lambda_2, \lambda_3$, respectively). Let $E_{ij} \subset Z$ denote the projective lines spanned by e_i and e_j for $i < j$. Then $\Omega_{\text{disc}} = Z - \{e_1, e_2, e_3\}$, and both sets $U_1 = Z - (E_{23} \cup \{e_1\})$ and $U_3 = Z - (E_{12} \cup \{e_3\})$ are maximal domains of proper discontinuity for Γ . (The maximality follows from the fact that the points on E_{12} are dynamically related to the points on E_{23} .) Observe also that in this example both U_1/Γ and U_3/Γ are compact.

Discontinuity and proper discontinuity can be nicely expressed using the notion of dynamical relation. The following definition is due to Frances [10, Definition 1]:

Definition 5.6 (dynamically related) Two points $z, z' \in Z$ are called *dynamically related* with respect to a sequence (h_n) in $\text{Homeo}(Z)$, written

$$z \stackrel{(h_n)}{\sim} z',$$

if there exists a sequence $z_n \rightarrow z$ in Z such that $h_n z_n \rightarrow z'$.

The points z, z' are called *dynamically related* with respect to the Γ -action, written

$$z \stackrel{\Gamma}{\sim} z',$$

if there exists a sequence $\gamma_n \rightarrow \infty$ in Γ such that $z \stackrel{(\gamma_n)}{\sim} z'$.

Here, for a sequence (γ_n) in Γ , we write $\gamma_n \rightarrow \infty$ if every element of Γ occurs at most finitely many times in the sequence.

One verifies (see eg [15]) that

- (i) dynamical relation is a closed relation in $Z \times Z$,
- (ii) points in different Γ -orbits are dynamically related if and only if their orbits cannot be separated by disjoint Γ -invariant open subsets.

The concept of dynamical relation is useful for our discussion of discontinuity because

- (i) a point is nonwandering if and only if it is dynamically related to itself,
- (ii) the action is proper on an open subset $U \subset Z$ if and only if no two points in U are dynamically related.

5.2 Accumulation and proper discontinuity

In this paper, we derive proper discontinuity of actions from a certain accumulation behavior which is a relaxation of *convergence dynamics*.

Let Z and Γ be as above. Let (Y_n) be a sequence of subsets of Z . We denote by $\text{Acc}((Y_n)) \subset Z$ the closed subset consisting of the accumulation points of all sequences (y_n) of points $y_n \in Y_n$.

Definition 5.7 (accumulation) We say that the sequence of subsets $Y_n \subset Z$ *accumulates* at a subset $S \subset Z$, written

$$Y_n \text{ acc } S,$$

if $\text{Acc}((Y_n)) \subseteq S$.

If $S \subset Z$ is closed, then the sequence (Y_n) accumulates at S if and only if every neighborhood U of S contains all but finitely many of the subsets Y_n .

We first consider the dynamics of a sequence (h_n) in $\text{Homeo}(Z)$.

Definition 5.8 (accumulating sequence) For compact subsets $A_{\pm} \subset Z$, we say that the sequence (h_n) *accumulates* at A_+ outside A_- , briefly, $(A_-, A_+)\text{-accumulates}$, if

$$(17) \quad h_n K \text{ acc } A_+$$

for all compacta K disjoint from A_- .

Property (17) is a statement about the locally uniform accumulation of the (h_n) -orbits initiating outside the exceptional subset A_- and can be rephrased in terms of dynamical relations between points in Z with respect to the (h_n) -action. Namely, equivalently, for all points $z, z' \in Z$ it holds that

$$(18) \quad z \stackrel{(h_{n_k})}{\sim} z' \text{ for some subsequence } (h_{n_k}) \implies z \in A_- \text{ or } z' \in A_+.$$

Note that the dynamical relation condition $z \stackrel{(h_{n_k})}{\sim} z'$ and the dual condition $z' \stackrel{(h_{n_k}^{-1})}{\sim} z$ are equivalent, and consequently we have the symmetry

$$(h_n) \text{ is } (A_-, A_+)\text{-accumulating} \iff (h_n^{-1}) \text{ is } (A_+, A_-)\text{-accumulating.}$$

Note that if $A_{\pm} \subset A'_{\pm}$, then $(A_-, A_+)\text{-accumulation}$ implies $(A'_-, A'_+)\text{-accumulation}$.

Now we consider the action $\Gamma \curvearrowright Z$.

Definition 5.9 (accumulating action, I) We say that the action $\Gamma \curvearrowright Z$ is $(A_-, A_+)\text{-accumulating}$ if every sequence $\gamma_n \rightarrow \infty$ in Γ has an $(A_-, A_+)\text{-accumulating}$ subsequence.

According to (18), we obtain for dynamical relations:

Lemma 5.10 (dynamical relations, I) *If the action $\Gamma \curvearrowright Z$ is $(A_-, A_+)\text{-accumulating}$, then for any two points $z, z' \in Z$, it holds that*

$$(19) \quad z \stackrel{\Gamma}{\sim} z' \implies z \in A_- \text{ or } z' \in A_+.$$

Proof This is a direct consequence of (18). □

We conclude:

Proposition 5.11 (proper discontinuity, I) *If the subsets A_{\pm} are Γ -invariant and if the action $\Gamma \curvearrowright Z$ is $(A_-, A_+)\text{-accumulating}$, then the action*

$$\Gamma \curvearrowright Z - (A_- \cup A_+)$$

is properly discontinuous.

Proof By Lemma 5.10, there are no dynamical relations between points outside $A_- \cup A_+$. □

Suppose that A_{\pm} are Γ -invariant compact (with respect to the Hausdorff topology) families of compact subsets $A_{\pm} \subset Z$.

Definition 5.12 (limit family) The *forward limit family* of Γ with respect to (A_-, A_+) is the family $\mathcal{L}_+ \subset \mathcal{A}_+$ consisting of all subsets $A_+ \in \mathcal{A}_+$ for which there exists a sequence $\gamma_n \rightarrow \infty$ in Γ which is $(A_-, A_+)\text{-accumulating}$ for some subset $A_- \in \mathcal{A}_-$. Similarly, we define the *backward limit family* $\mathcal{L}_- \subset \mathcal{A}_-$.

The limit families \mathcal{L}_\pm are Γ -invariant. Due to the compactness of the families \mathcal{A}_\pm , they are closed and hence compact themselves:

Lemma 5.13 \mathcal{L}_\pm is closed.

Proof Suppose, for instance, that (A_+^k) is a sequence in \mathcal{L}_+ such that $A_+^k \rightarrow A_+ \in \mathcal{A}_+$. There exist sequences $\gamma_n^k \rightarrow \infty$ in Γ which (A_-^k, A_+^k) -accumulate for some $A_-^k \in \mathcal{A}_-$ (in fact $\in \mathcal{L}_-$). After passing to a subsequence, we may assume that $A_-^k \rightarrow A_- \in \mathcal{A}_-$. A diagonal argument yields an (A_-, A_+) -accumulating sequence $(\gamma_n^{k(m)})_m$ in Γ . Hence $A_+ \in \mathcal{L}_+$ and \mathcal{L}_+ is closed. □

As a consequence of the lemma, the Γ -invariant subsets

$$T_\pm := \bigcup_{A_\pm \in \mathcal{L}_\pm} A_\pm \subset Z$$

are compact.

Definition 5.14 (accumulating action, II) We say that the action $\Gamma \curvearrowright Z$ is $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating if every sequence $\gamma_n \rightarrow \infty$ in Γ has a subsequence which is $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating for some subsets $A_\pm \in \mathcal{A}_\pm$.

For such accumulating actions, the limit families are closely related to their dynamics:

Lemma 5.15 (dynamical relations, II) *If the action $\Gamma \curvearrowright Z$ is $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating, then for any two points $z, z' \in Z$, it holds that*

$$z \overset{\Gamma}{\sim} z' \implies z \in T_- \text{ or } z' \in T_+.$$

Proof If two points are dynamically related with respect to the Γ -action, then they are dynamically related with respect to an $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating sequence in Γ with $A_\pm \in \mathcal{L}_\pm$ and, hence, $A_\pm \subset T_\pm$. The assertion therefore follows from (18). □

We conclude as before:

Proposition 5.16 (proper discontinuity, II) *If the action $\Gamma \curvearrowright Z$ is $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating, then the action*

$$\Gamma \curvearrowright Z - (T_- \cup T_+)$$

is properly discontinuous.

Remark 5.17 (convergence actions) The action $\Gamma \curvearrowright Z$ is a *convergence action* (see eg [5]) if and only if it is $(\mathcal{A}_-, \mathcal{A}_+)$ -accumulating with \mathcal{A}_\pm the family of one point subsets. The limit families \mathcal{L}_\pm then become the *limit set* $\Lambda \subset Z$ of the action. The action on its complement is properly discontinuous, compare Proposition 5.16. We recover the *dynamical decomposition*

$$Z = \Omega_{\text{disc}} \sqcup \Lambda$$

and that the action on the domain of discontinuity is proper. Hence, for convergence actions there exists a unique maximal domain of proper discontinuity.

The main example of convergence actions comes from the following fact: Every discrete group Γ of isometries of a proper Gromov hyperbolic geodesic metric space Y acts as a convergence group on the visual compactification $\bar{Y} = Y \cup \partial_\infty Y$, and in particular on the Gromov boundary $\partial_\infty Y$ of Y .

Remark 5.18 (accumulation phenomena in nonpositive curvature) Convergence type behavior in the sense of accumulation has been studied by Karlsson, Papasoglu and Swenson in the general context of nonpositive curvature. They showed that for proper isometric actions $\Gamma \curvearrowright Y$ on CAT(0) spaces the induced action $\Gamma \curvearrowright \partial_\infty Y$ on the visual boundary is $(\mathcal{B}(\theta), \mathcal{B}(\pi-\theta))$ -accumulating for $0 < \theta < \pi$, where $\mathcal{B}(\theta)$ is the family of closed balls of Tits radius θ in $\partial_\infty Y$; see [25, Theorem 1] and [34, Theorem 4]. Some of our results can be viewed as combinatorial versions of this (Tits) metric result for actions on CAT(0) *model* spaces of higher rank; see eg Corollary 6.3 and Lemma 6.14 below.

5.3 Expansion and cocompactness

In this section, let (Z, d) be a compact metric space and let $\Gamma \curvearrowright Z$ be a continuous action of a discrete group.

The following notion is due to Sullivan [37, Section 9]:

Definition 5.19 (expanding action) We say that the action $\Gamma \curvearrowright Z$ is *expanding* at the *point* $z \in Z$ if there exists an element $\gamma \in \Gamma$ which is *uniformly expanding* on a neighborhood U of z ; ie for some constant $c > 1$ and all points $z_1, z_2 \in U$ we have

$$d(\gamma z_1, \gamma z_2) \geq c \cdot d(z_1, z_2).$$

We say that the action of Γ is *expanding* at a compact Γ -invariant *subset* $E \subset Z$ if it is expanding at all points $z \in E$.

Remark 5.20 If the action $\Gamma \curvearrowright Z$ is expanding at E , then it is *arbitrarily strongly* expanding there, ie for every point $z \in E$ exist a sequence (γ_n) in Γ and a sequence of (shrinking) neighborhoods U_n of z such that the $\gamma_n|_{U_n}$ are uniformly expanding with expansion factors $c_n \rightarrow +\infty$. This follows directly from the definition by iterating locally expanding elements. Note that, as a consequence, the action is expanding at E also with respect to any bilipschitz equivalent metric on Z .

We will need the following more general notion of *partial expansion*. We suppose that the action $\Gamma \curvearrowright Z$ has the following structure: There is a Γ -invariant compact subset $E \subset Z$ and a continuous map $\pi: E \rightarrow \Lambda$ onto a compact topological space Λ (eg a fiber bundle), such that the restricted action $\Gamma \curvearrowright E$ is fiber preserving, ie it descends to a continuous action $\Gamma \curvearrowright \Lambda$. We set $E_\lambda := \pi^{-1}(\lambda)$.

Definition 5.21 (transversely expanding action, I) We say that the action $\Gamma \curvearrowright Z$ is *expanding transversely to π at the fiber E_λ* if there exist an element $\gamma \in \Gamma$ and a neighborhood $U \subset Z$ of E_λ such that for some constant $c > 1$ we have

$$(20) \quad d(\gamma z, E_{\gamma\lambda'}) \geq c \cdot d(z, E_{\lambda'})$$

for all points $z \in U$ and fibers $E_{\lambda'} \subset U$.

We say that the action $\Gamma \curvearrowright Z$ is *expanding at E transversely to π* if it is expanding at all fibers E_λ .

The action $\Gamma \curvearrowright Z$ is expanding at E if and only if it is expanding at E transversely to id_E .

The concept of expansion is important to us due to the following observation:

Proposition 5.22 (transversely expanding implies cocompact on the complement, I) *If the action $\Gamma \curvearrowright Z$ is expanding at E transversely to π , then the action $\Gamma \curvearrowright Z - E$ is cocompact.*

Proof We claim that for some constant $c > 1$,

$$(21) \quad \sup d(\cdot, E)|_{\Gamma z} > c \cdot d(z, E)$$

for all $z \in Z - E$ sufficiently close to E . Otherwise, there would exist a sequence (z_n) in $Z - E$ accumulating at E and a sequence of constants $c_n \rightarrow 1$ such that

$$d(\gamma z_n, E) \leq c_n \cdot d(z_n, E)$$

for all $n \in \mathbb{N}$ and $\gamma \in \Gamma$. Since E is compact, we may assume, after passing to a subsequence, that (z_n) accumulates at a fiber, $z_n \rightarrow E_\lambda$. Due to expansion, there exists an element $\gamma_\lambda \in \Gamma$ which satisfies the expansion property (20) on a neighborhood $U_\lambda \subset Z$ of E_λ with some expansion factor $c_\lambda > 1$. Let $E_{\gamma_\lambda \lambda_n}$ be the fiber closest to $\gamma_\lambda z_n$, $d(\gamma_\lambda z_n, E_{\gamma_\lambda \lambda_n}) = d(\gamma_\lambda z_n, E)$. Then $\lambda_n \rightarrow \lambda$. Since $z_n \in U_\lambda$ and $E_{\lambda_n} \subset U_\lambda$ for large n , it follows that

$$c_\lambda \cdot d(z_n, E) \leq c_\lambda \cdot d(z_n, E_{\lambda_n}) \leq d(\gamma_\lambda z_n, E_{\gamma_\lambda \lambda_n}) = d(\gamma_\lambda z_n, E) \leq c_n \cdot d(z_n, E),$$

a contradiction confirming our claim.

Let $U \subset Z$ be an open tubular neighborhood of E where (21) holds. Thus, no Γ -orbit is entirely contained in $U - E$ and, therefore, every Γ -orbit in $Z - E$ meets the compact subset $Z - U \subset Z - E$. □

The above argument (from [18, Section 2.2]) leads actually to a more general result.

Let us suppose, more generally, that the action $\Gamma \curvearrowright Z$ has the following structure: There is a Γ -invariant compact subset $E \subset Z$ which is represented as the (not necessarily disjoint) union of a Γ -invariant collection $\mathcal{E} = \{E_\lambda : \lambda \in \Lambda\}$ of compact subsets $E_\lambda \subset Z$ parametrized by some set Λ .

Definition 5.23 (transversely expanding action, II) We say that the action $\Gamma \curvearrowright Z$ is *expanding transversely to \mathcal{E} at a point z* if there exist an element $\gamma \in \Gamma$, a neighborhood $U \subset Z$ of z and a constant $c > 1$ such that we have

$$(22) \quad d(\gamma u, \gamma E_\lambda) \geq c \cdot d(u, E_\lambda)$$

for all points $u \in U - E$ and all E_λ which have nonempty intersection with U . We say that the action $\Gamma \curvearrowright Z$ is *expanding transversely to \mathcal{E}* if it is expanding at all points $z \in E$.

Proposition 5.24 (transversely expanding implies cocompact on the complement, II) *If the action $\Gamma \curvearrowright Z$ is expanding transversely to \mathcal{E} , then the action $\Gamma \curvearrowright Z - E$ is cocompact.*

Proof We claim that for some constant $c > 1$,

$$(23) \quad \sup d(\cdot, E)|_{\Gamma u} > c \cdot d(u, E)$$

for all $u \in Z - E$ sufficiently close to E . Otherwise, there would exist a sequence (u_n) in $Z - E$ accumulating at E and a sequence of constants $c_n \rightarrow 1$ such that

$$d(\gamma u_n, E) \leq c_n \cdot d(u_n, E)$$

for all $n \in \mathbb{N}$ and $\gamma \in \Gamma$. Since E is compact, we may assume, after passing to a subsequence, that (u_n) converges to some point $z \in E_\lambda$ for some $\lambda \in \Lambda$. Due to expansion, there exists an element $\gamma_z \in \Gamma$ which satisfies the expansion property (22) on a neighborhood $U \subset Z$ of z with some expansion factor $c > 1$. Let $\gamma_z E_{\lambda_n} \in \mathcal{E}$ be the set in the collection \mathcal{E} closest to $\gamma_z u_n$:

$$d(\gamma_z u_n, E) = d(\gamma_z u_n, \gamma_z E_{\lambda_n}) = d(\gamma_z u_n, \gamma_z z_n)$$

with $z_n \in E_{\lambda_n}$. Then $z_n \rightarrow z$ because $\gamma_z u_n \rightarrow \gamma_z z \in E$, which implies that $E_{\lambda_n} \cap U \neq \emptyset$ for all sufficiently large n . It follows that (for large n),

$$c \cdot d(u_n, E) \leq c \cdot d(u_n, E_{\lambda_n}) \leq d(\gamma_z u_n, \gamma_z E_{\lambda_n}) = d(\gamma_z u_n, E) \leq c_n \cdot d(u_n, E),$$

a contradiction confirming our claim.

Let $U \subset Z$ be an open tubular neighborhood of E where (23) holds. Thus, no Γ -orbit is entirely contained in $U - E$ and, therefore, every Γ -orbit in $Z - E$ meets the compact subset $Z - U \subset Z - E$. □

6 Accumulation dynamics on flag manifolds and proper discontinuity

We now study the dynamics of G and its discrete subgroups $\Gamma < G$ on its associated flag manifolds, equivalently, on (the G -orbits in) the visual boundary $\partial_\infty X$. In this section, we will discuss a certain dynamical behavior, which is a relaxed version of convergence dynamics, and use it to construct domains of proper discontinuity for discrete subgroups.

6.1 Weakly contracting sequences

Let (g_n) be a sequence in G , and let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be a face type.

We consider the following contraction property for the dynamics of (g_n) on $\text{Flag}_{\tau_{\text{mod}}}$. An equivalent notion had been studied by Benoist in his foundational work [1]; see Section 3.5 there.

Definition 6.1 (τ_{mod} -contracting sequence) We say that the sequence (g_n) is τ_{mod} -contracting if there exist simplices τ_{\pm} of type $\pm\tau_{\text{mod}}$ such that

$$(24) \quad g_n|_{C(\tau_-)} \rightarrow \tau_+$$

uniformly on compacta as $n \rightarrow +\infty$.

We recall that $C(\tau_-)$ is a dense open subset of $\text{Flag}_{\tau_{\text{mod}}}$.

Property (24) means (g_n) is $(\text{Flag}_{\tau_{\text{mod}}}-C(\tau_-), \tau_+)$ -accumulating; see Definition 5.8. It can be rephrased in terms of dynamical relations between points in $\text{Flag}_{\tau_{\text{mod}}}$ with respect to the (g_n) -action. Namely, equivalently, for all simplices $\tau, \tau' \in \text{Flag}_{\tau_{\text{mod}}}$, it holds (see (18)) that

$$(25) \quad \tau \stackrel{(g_{n_k})}{\rightsquigarrow} \tau' \text{ for some subsequence } (g_{n_k}) \implies \tau \notin C(\tau_-) \text{ or } \tau' = \tau_+.$$

The conclusion of the last implication can be expressed in terms of relative positions:

$$\text{pos}(\tau, \tau_-) \text{ maximal} \implies \text{pos}(\tau', \tau_+) \text{ minimal}.$$

We observe that the last implication follows from the combinatorial inequality

$$(26) \quad \text{pos}(\tau', \tau_+) < \text{c-pos}(\tau, \tau_-).$$

The next result shows that this inequality holds for dynamically related points on all flag manifolds, thought of as G -orbits in $\partial_{\infty} X$. It is the key step in our study of proper discontinuity.

Proposition 6.2 (dynamical relation inequality) *The following are equivalent:*

- (i) property (24);
- (ii) for any two points $\xi, \xi' \in \partial_{\infty} X$, it holds that

$$(27) \quad \xi \stackrel{(g_{n_k})}{\rightsquigarrow} \xi' \text{ for some subsequence } (g_{n_k}) \implies \text{pos}(\xi', \tau_+) < \text{c-pos}(\xi, \tau_-).$$

Proof Suppose first that property (24) holds and that $\xi \stackrel{(g_n)}{\rightsquigarrow} \xi'$. Then ξ and ξ' lie in the same G -orbit, ie $G\xi = G\xi'$, and there exists a sequence (ξ_n) in this G -orbit such that $\xi_n \rightarrow \xi$ and $g_n\xi_n \rightarrow \xi'$. Let $a \subset \partial_{\infty} X$ be an apartment containing τ_- and ξ . Nearby apartments a_n containing ξ_n can be obtained by using isometries $h_n \rightarrow e$ in G with $\xi_n = h_n\xi$ and putting $a_n = h_n a$. Let $\hat{\tau}_- \subset a$ be the simplex opposite to τ_- ,

and let $\tau_n = h_n \hat{\tau}_- \subset a_n$. Then $\tau_n \rightarrow \hat{\tau}_-$. Since $\hat{\tau}_- \in C(\tau_-)$, the locally uniform convergence (24) implies that $g_n \tau_n \rightarrow \tau_+$. We obtain

$$\begin{aligned} \text{pos}(\xi', \tau_+) &< \text{pos}(g_n \xi_n, g_n \tau_n) \\ &= \text{pos}(\xi_n, \tau_n) = \text{pos}(h_n \xi, h_n \hat{\tau}_-) \\ &= \text{pos}(\xi, \hat{\tau}_-) = \text{c-pos}(\xi, \tau_-), \end{aligned}$$

where the inequality follows from the semicontinuity of relative position; see Lemma 3.12.

Conversely, suppose that (ii) holds. Since inequality (26) is a special case of the inequality in the implication of (27), it follows that (25), and thus (24), holds. \square

We observe a symmetry: condition (27) is equivalent to the dual condition

$$(28) \quad \xi' \stackrel{(g_{n_k}^{-1})}{\sim} \xi \text{ for some subsequence } (g_{n_k}^{-1}) \text{ of } (g_n^{-1}) \implies \text{pos}(\xi, \tau_-) < \text{c-pos}(\xi', \tau_+),$$

because both dynamical relation hypotheses are equivalent, as are the combinatorial inequality conclusions. Therefore, the proposition implies that (24) is equivalent to the dual property on $\text{Flag}_{-\tau_{\text{mod}}}$ that

$$g_n^{-1}|_{C(\tau_+)} \rightarrow \tau_-$$

uniformly on compacta as $n \rightarrow +\infty$.

Note that the simplices τ_{\pm} in (24) are well defined, because this is clear for τ_+ and follows for τ_- by symmetry.

Inequality (27) can be (re)converted into a statement about the asymptotic behavior of arbitrary (g_n) -orbits in $\partial_{\infty} X$. We can in general not expect that these orbits converge, but we obtain information where they accumulate. For individual orbits, it follows that for a point $\xi \in \partial_{\infty} X$ the orbit $(g_n \xi)$ accumulates in $G\xi \subset \partial_{\infty} X$ at the Schubert cycle

$$\{\text{pos}(\cdot, \tau_+) < \text{c-pos}(\xi, \tau_-)\}.$$

A locally uniform statement can be conveniently formulated using the language of thickenings:

Corollary 6.3 (orbit accumulation) *If property (24) holds, and if $\text{Th} \subset W$ is a $W_{\tau_{\text{mod}}}$ -left invariant thickening, then the sequence (g_n) is $(\text{Th}^c(\tau_-), \text{Th}(\tau_+))$ -accumulating (see Definition 5.8).*

Proof Otherwise, there is a dynamical relation $\xi \stackrel{(g_{n_k})}{\sim} \xi'$ with $\xi \notin \text{Th}^c(\tau_-)$ and $\xi' \notin \text{Th}(\tau_+)$; compare (18). That is, $\text{pos}(\xi, \tau_-) \notin \text{Th}^c$ and $\text{pos}(\xi', \tau_+) \notin \text{Th}$. Moreover, (24) implies (27), and hence the inequality $\text{pos}(\xi', \tau_+) < \text{c-pos}(\xi, \tau_-)$. It follows that

$$\text{pos}(\xi', \tau_+) < \text{c-pos}(\xi, \tau_-) = w_0 \text{pos}(\xi, \tau_-) \in \text{Th},$$

and hence $\text{pos}(\xi', \tau_+) \in \text{Th}$, a contradiction. □

6.2 Weak convergence subgroups

Let $\Gamma < G$ be a discrete subgroup.

Definition 6.4 (τ_{mod} -limit set) We define the *forward/backward τ_{mod} -limit set* of Γ as the set

$$\Lambda_{\tau_{\text{mod}}}^{\pm} = \Lambda_{\tau_{\text{mod}}}^{\pm}(\Gamma) \subset \text{Flag}_{\pm\tau_{\text{mod}}}$$

of all simplices τ_{\pm} as in (24) for all τ_{mod} -contracting sequences $\gamma_n \rightarrow \infty$ in Γ .

An equivalent notion had already been introduced by Benoist in his profound work; see [1, Section 3.6].

Note that passing to a finite index subgroup does not change the limit sets.

The limit sets $\Lambda_{\tau_{\text{mod}}}^{\pm}$ are Γ -invariant and compact; see Lemma 5.13. Moreover, one has the symmetry

$$\Lambda_{-\tau_{\text{mod}}}^{\pm}(\Gamma) = \Lambda_{\tau_{\text{mod}}}^{\mp}(\Gamma).$$

In particular, if τ_{mod} is ι -invariant we can define the τ_{mod} -limit set

$$\Lambda_{\tau_{\text{mod}}}(\Gamma) := \Lambda_{\tau_{\text{mod}}}^{\pm}(\Gamma).$$

To any $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, we associate the Γ -invariant compact families of compact subsets

$$\mathcal{A}_{\tau_{\text{mod}}, \text{Th}}^{-} := \{\text{Th}^c(\tau_-) : \tau_- \in \Lambda_{\tau_{\text{mod}}}^{-}\} \quad \text{and} \quad \mathcal{A}_{\tau_{\text{mod}}, \text{Th}}^{+} := \{\text{Th}(\tau_+) : \tau_+ \in \Lambda_{\tau_{\text{mod}}}^{+}\}.$$

The structure of the dynamics of the action $\Gamma \curvearrowright \partial_{\infty} X$ is closely related to the limit sets if it enjoys contraction behavior in the following sense:

Definition 6.5 (τ_{mod} -convergence action) The action $\Gamma \curvearrowright \partial_{\infty} X$ is called a τ_{mod} -convergence action if every sequence $\gamma_n \rightarrow \infty$ in Γ has a τ_{mod} -contracting subsequence. The subgroup $\Gamma < G$ is then called a τ_{mod} -convergence subgroup.

Remark 6.6 (rank one) If $\text{rank}(X) = 1$, this property is equivalent to being a *convergence action* and is satisfied for all discrete subgroups $\Gamma < G$, compare Remark 5.17.

Corollary 6.3 implies:

Proposition 6.7 (weak contraction implies accumulation) *If $\Gamma < G$ is a τ_{mod} -convergence subgroup and if $\text{Th} \subset W$ is a $W_{\tau_{\text{mod}}}$ -left invariant thickening, then the action $\Gamma \curvearrowright \partial_{\infty} X$ is $(\mathcal{A}_{\tau_{\text{mod}}, \text{Th}}^-, \mathcal{A}_{\tau_{\text{mod}}, \text{Th}}^+)$ -accumulating.*

We obtain our main result for proper discontinuity:

Theorem 6.8 (domains of proper discontinuity for τ_{mod} -convergence subgroups) *If $\Gamma < G$ is a τ_{mod} -convergence subgroup, then for any $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_{\infty} X - (\text{Th}^c(\Lambda_{\tau_{\text{mod}}}^-) \cup \text{Th}(\Lambda_{\tau_{\text{mod}}}^+))$$

is properly discontinuous. In particular, if τ_{mod} is ι -invariant and Th is fat, the action

$$\Gamma \curvearrowright \partial_{\infty} X - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous.

Proof The first assertion follows from the last proposition by applying Proposition 5.16 with $\mathcal{L}_{\pm} = \mathcal{A}_{\tau_{\text{mod}}, \text{Th}}^{\pm}$. The second assertion follows because $\text{Th}^c \subseteq \text{Th}$ due to fatness. \square

Note that the thickenings of limit sets $\text{Th}(\Lambda_{\tau_{\text{mod}}}^{\pm}(\Gamma))$ are Γ -invariant and compact.

For examples of thickenings, we refer to Section 3.4.2.

6.3 Weakly regular subgroups

The properties of contraction, defined via the dynamics at infinity (see Definition 6.1), and regularity, defined via the asymptotics of orbits in X (see Definition 4.3), are equivalent in a suitable sense; compare the discussion in [19, Section 5.2]. The most relevant aspect for the purposes of this paper is that regularity implies contraction in a suitable sense.

We first consider sequences of isometries:

Proposition 6.9 *Every τ_{mod} -regular sequence in G contains a τ_{mod} -contracting subsequence.*

Proof Compare the proof of [19, Proposition 5.14].

Suppose that the sequence (g_n) in G is τ_{mod} -regular. Let $x \in X$. There exist simplices $\tau_n^\pm \in \text{Flag}_{\pm\tau_{\text{mod}}}$ (unique for large n) such that

$$g_n^{\pm 1}x \in V(x, \text{st}(\tau_n^\pm)).$$

After passing to a subsequence, we may assume convergence

$$\tau_n^\pm \rightarrow \tau_\pm$$

in $\text{Flag}_{\pm\tau_{\text{mod}}}$, because the flag manifolds are compact.

Since $x \in g_n V(x, \text{st}(\tau_n^-)) = V(g_n x, \text{st}(g_n \tau_n^-))$ and $g_n x \in V(x, \text{st}(\tau_n^+))$, it follows that the Weyl cones $V(g_n x, \text{st}(g_n \tau_n^-))$ and $V(x, \text{st}(\tau_n^+))$ lie in the same parallel set, namely in $P(g_n \tau_n^-, \tau_n^+)$, and face in opposite directions. In particular, the simplices $g_n \tau_n^-$ and τ_n^+ are x -opposite, and thus $g_n \tau_n^-$ converges to the simplex $\hat{\tau}_+$ which is x -opposite to τ_+ :

$$g_n \tau_n^- \rightarrow \hat{\tau}_+.$$

Since the sequence $(g_n x)$ is τ_{mod} -regular, it holds that

$$d(g_n^{-1}x_n, \partial V(x, \text{st}(\tau_n^-))) \rightarrow +\infty.$$

According to Lemma 4.2, for any $r, R > 0$, the inclusion of shadows

$$U_{\tau_n^-, x, R} \subset U_{\tau_n^-, g_n^{-1}x, r}$$

holds for $n \geq n(r, R)$. Therefore, there exist positive numbers $R_n \rightarrow +\infty$ and $r_n \rightarrow 0$ such that

$$U_{\tau_n^-, x, R_n} \subset U_{\tau_n^-, g_n^{-1}x, r_n}$$

for large n , equivalently,

$$(29) \quad g_n U_{\tau_n^-, x, R_n} \subset U_{g_n \tau_n^-, x, r_n}.$$

Since $\tau_n^- \rightarrow \tau_-$ and $R_n \rightarrow +\infty$, the sequence of shadows $U_{\tau_n^-, x, R_n} \subset C(\tau_n^-) \subset \text{Flag}_{\tau_{\text{mod}}}$ exhausts $C(\tau_-)$ in the sense that every compactum in $C(\tau_-)$ is contained in $U_{\tau_n^-, x, R_n}$ for large n . Indeed, for fixed $R > 0$ we have Hausdorff convergence $U_{\tau_n^-, x, R} \rightarrow U_{\tau_-, x, R}$ in $\text{Flag}_{\tau_{\text{mod}}}$, which immediately follows eg using symmetry, ie from the transitivity of the action $K_x \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ of the maximal compact stabilizer $K_x < G$ of x . Furthermore, the shadows $U_{\tau_-, x, R}$ exhaust $C(\tau_-)$ as $R \rightarrow +\infty$; see the continuity part of Lemma 4.1.

On the other hand, since $g_n \tau_n^- \rightarrow \hat{\tau}_+$ and $r_n \rightarrow 0$, the shadows $U_{g_n \tau_n^-, x, r_n}$ shrink, ie Hausdorff converge to the point τ_+ . Indeed, $U_{g_n \tau_n^-, x, r} \rightarrow U_{\hat{\tau}_+, x, r}$ in $\text{Flag}_{\tau_{\text{mod}}}$ for fixed $r > 0$, and $U_{\hat{\tau}_+, x, r} \rightarrow \tau_+$ as $r \rightarrow 0$, using again the continuity part of Lemma 4.1 and the fact that the function (16) assumes the value zero only in τ_+ .

Together with these observations on exhaustion and shrinking of shadows, (29) shows

$$g_n|_{C(\tau_-)} \rightarrow \tau_+$$

uniformly on compacta; ie the (sub)sequence (g_n) is τ_{mod} -contracting. □

Remark 6.10 The converse, that τ_{mod} -contracting sequences in G are τ_{mod} -regular, was shown in [19, Theorem 5.23].

We conclude for groups of isometries:

Corollary 6.11 *τ_{mod} -regular subgroups are τ_{mod} -convergence subgroups.*

Remark 6.12 For τ_{mod} -regular subgroups, the notion of τ_{mod} -limit set introduced in Definition 6.4 is equivalent to the notion of τ_{mod} -limit set introduced in Definition 5.32 of [19]; see Proposition 5.29 of [19].

Based on the corollary, we can translate our proper discontinuity result for convergence subgroups (Theorem 6.8) into one for regular subgroups:

Theorem 6.13 (domains of proper discontinuity for τ_{mod} -regular subgroups) *Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be an arbitrary face type. If $\Gamma < G$ is a τ_{mod} -regular subgroup, then for every $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_\infty X - (\text{Th}^c(\Lambda_{\tau_{\text{mod}}}^-) \cup \text{Th}(\Lambda_{\tau_{\text{mod}}}^+))$$

is properly discontinuous. In particular, if τ_{mod} is ι -invariant and Th is fat, the action

$$\Gamma \curvearrowright \partial_\infty X - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous.

6.4 Discrete subgroups

The general construction of domains of proper discontinuity in Section 5.2 applies equally to arbitrary discrete subgroups $\Gamma < G$. There are several ways to proceed. The most immediate possibility is the following.

Choose for every face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ a $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th}_{\tau_{\text{mod}}}$, and define the Γ -invariant compact families

$$\mathcal{A}_{\pm} := \bigcup_{\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}} \mathcal{A}_{\tau_{\text{mod}}, \text{Th}_{\tau_{\text{mod}}}}^{\pm}.$$

Lemma 6.14 *The action $\Gamma \curvearrowright \partial_{\infty} X$ is $(\mathcal{A}_{-}, \mathcal{A}_{+})$ -accumulating.*

Proof According to Lemma 4.5, every sequence $\gamma_n \rightarrow \infty$ in Γ contains a τ_{mod} -regular (even τ_{mod} -pure) subsequence, and hence a τ_{mod} -contracting subsequence for some face type τ_{mod} . The assertion follows therefore from Corollary 6.3. \square

Thus Proposition 5.16 yields in this case:

Proposition 6.15 (domains of proper discontinuity for discrete subgroups, I) *If $\Gamma < G$ is a discrete subgroup, then the action*

$$(30) \quad \Gamma \curvearrowright \partial_{\infty} X - \bigcup_{\tau_{\text{mod}}} (\text{Th}_{\tau_{\text{mod}}}^c(\Lambda_{\tau_{\text{mod}}}^{-}) \cup \text{Th}_{\tau_{\text{mod}}}(\Lambda_{\tau_{\text{mod}}}^{+}))$$

is properly discontinuous.

In general, this domain of proper discontinuity can be further enlarged by only removing the thickenings of the limit simplices arising from *pure* sequences in the group: define the *pure forward/backward τ_{mod} -limit set*

$$\Lambda_{\tau_{\text{mod}}}^{\text{pure}, \pm} \subseteq \Lambda_{\tau_{\text{mod}}}^{\pm}$$

as the closure of the set of all simplices τ_{\pm} as in (24) for all τ_{mod} -pure τ_{mod} -contracting sequences (γ_n) in Γ . As above, we conclude:

Proposition 6.16 (domains of proper discontinuity for discrete subgroups, II) *If $\Gamma < G$ is a discrete subgroup, then the action*

$$(31) \quad \Gamma \curvearrowright \partial_{\infty} X - \bigcup_{\tau_{\text{mod}}} (\text{Th}_{\tau_{\text{mod}}}^c(\Lambda_{\tau_{\text{mod}}}^{\text{pure}, -}) \cup \text{Th}_{\tau_{\text{mod}}}(\Lambda_{\tau_{\text{mod}}}^{\text{pure}, +}))$$

is properly discontinuous.

Since the domain in (30) is in general smaller than the domain in (31), one cannot expect the Γ -action on it to be cocompact.

On the other hand, if Γ is τ_{mod} -regular, then it contains ν_{mod} -pure sequences only for the face types $\nu_{\text{mod}} \supseteq \tau_{\text{mod}}$, and hence only these limit sets $\Lambda_{\nu_{\text{mod}}}^{\pm}$ can be nonempty. Since $W_{\nu_{\text{mod}}} \leq W_{\tau_{\text{mod}}}$, we may choose $\text{Th}_{\nu_{\text{mod}}} = \text{Th}_{\tau_{\text{mod}}}$ for these face types, and then the domain in (31) coincides with the domain in Theorem 6.13.

7 Cocompactness

7.1 Nearby simplex thickenings

For incident faces $\nu \subset \tau \subset \partial_\infty X$, the parabolic subgroups fixing them are contained in each other, $P_\nu \supset P_\tau$. Correspondingly, for incident face types $\nu_{\text{mod}} \subset \tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ there is the natural forgetful map

$$\pi_{\nu_{\text{mod}}\tau_{\text{mod}}} : \text{Flag}_{\tau_{\text{mod}}} \rightarrow \text{Flag}_{\nu_{\text{mod}}}$$

assigning to a face τ of type τ_{mod} its face ν of type ν_{mod} . It is a G -equivariant smooth fibration with compact base and fiber.

We fix auxiliary Riemannian metrics on all partial flag manifolds $\text{Flag}_{\tau_{\text{mod}}}$. Thereby also the G -orbits $G\xi \subset \partial_\infty X$ are equipped with Riemannian metrics by equivariantly identifying them with the appropriate flag manifolds.

The fibrations $\pi_{\nu_{\text{mod}}\tau_{\text{mod}}}$ are then Lipschitz continuous by compactness. Vice versa, we have:

Lemma 7.1 (controlled lifts) *Let τ and ν' be simplices of types τ_{mod} and ν_{mod} with $\nu_{\text{mod}} \subset \tau_{\text{mod}}$, and let $\nu \subset \tau$ be the face of type ν_{mod} . Then there exists a simplex $\tau' \supset \nu'$ of type τ_{mod} such that*

$$d(\tau', \tau) \leq C_0 \cdot d(\nu', \nu)$$

with a uniform constant $C_0 \geq 1$ only depending on the chosen Riemannian metrics.

Proof The Riemannian metrics on $\text{Flag}_{\tau_{\text{mod}}}$ and $\text{Flag}_{\nu_{\text{mod}}}$ can be chosen so that $\pi_{\nu_{\text{mod}}\tau_{\text{mod}}}$ becomes a Riemannian submersion. With respect to these metrics, there exists τ' such that $d(\tau, \tau') = d(\nu, \nu')$. For other choices of the metrics, a multiplicative constant enters. \square

The lemma generalizes (by induction along galleries) to:

Lemma 7.2 *Let τ, τ' be simplices of type τ_{mod} and let $\tilde{\tau}$ be a simplex of type $\tilde{\tau}_{\text{mod}}$. Then there exists another simplex $\tilde{\tau}'$ of type $\tilde{\tau}_{\text{mod}}$ with relative position $\text{pos}(\tilde{\tau}', \tau') = \text{pos}(\tilde{\tau}, \tau)$ such that*

$$d(\tilde{\tau}', \tilde{\tau}) \leq C_1 \cdot d(\tau', \tau)$$

with a uniform constant $C_1 \geq 1$ only depending on the chosen Riemannian metrics.

Let now $G\xi \subset \partial_\infty X$ be a G -orbit at infinity, which we think of as identified with the appropriate flag manifold. We fix a $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$. Then the distance between simplices τ in $\text{Flag}_{\tau_{\text{mod}}}$ and the Hausdorff distance between their thickenings $\text{Th}(\tau) \cap G\xi$ in $G\xi$ control each other, and through an ideal point in $G\xi$ outside a simplex thickening exists a simplex thickening at controlled distance:

Lemma 7.3 (nearby simplex thickenings) *The following assertions hold with a uniform constant $C \geq 1$ only depending on the chosen Riemannian metrics:*

- (i) *The Hausdorff distance between the thickenings of any two simplices $\tau', \tau \in \text{Flag}_{\tau_{\text{mod}}}$ is controlled by*

$$d_H(\text{Th}(\tau') \cap G\xi, \text{Th}(\tau) \cap G\xi) \leq C \cdot d(\tau', \tau).$$

- (ii) *For a point $\xi' \in G\xi$ and a simplex $\tau \in \text{Flag}_{\tau_{\text{mod}}}$, there exists a simplex $\tau' \in \text{Flag}_{\tau_{\text{mod}}}$ such that $\xi' \in \text{Th}(\tau')$ and*

$$d(\tau', \tau) \leq C \cdot d(\xi', \text{Th}(\tau) \cap G\xi).$$

Proof (i) If $\eta \in \text{Th}(\tau) \cap G\xi$ is arbitrary, then applying Lemma 7.2 (to the flag manifold identified with $G\xi$) yields a point $\eta' \in G\xi$ with $\text{pos}(\eta', \tau') = \text{pos}(\eta, \tau) \in \text{Th}(\tau_{\text{mod}})$, ie $\eta' \in \text{Th}(\tau')$, and controlled distance $d(\eta', \eta) \leq C \cdot d(\tau, \tau')$.

(ii) Suppose that $\xi \in \text{Th}(\tau) \cap G\xi$ is the point closest to ξ' , that is, $d(\xi', \xi) = d(\xi', \text{Th}(\tau) \cap G\xi)$. Lemma 7.2 yields a simplex $\tau' \in \text{Flag}_{\tau_{\text{mod}}}$ with $\text{pos}(\xi', \tau') = \text{pos}(\xi, \tau) \in \text{Th}$ and controlled distance $d(\tau', \tau) \leq C \cdot d(\xi', \xi)$, whence the second inequality. □

7.2 From expansion to transverse expansion

Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be a face type, and let $\text{Th} \subset W$ be a $W_{\tau_{\text{mod}}}$ -left invariant thickening. In this section, we work on a fixed but arbitrary G -orbit $G\eta \subset \partial_\infty X$.

We start with an observation concerning the topology of thickenings in flag manifolds.

Lemma 7.4 (fibration of thickenings) *Let $A \subset \text{Flag}_{\tau_{\text{mod}}}$ be compact, and suppose that the thickenings $\text{Th}(\tau) \cap G\eta$ of the simplices $\tau \in A$ are pairwise disjoint. Then the natural map*

$$\text{Th}(A) \cap G\eta \xrightarrow{\pi} A$$

is a continuous fibration with compact fiber.

Proof Suppose that $\xi_n \rightarrow \xi$ in $\text{Th}(A) \cap G\eta$ and $\tau_n \rightarrow \tau$ in A with $\xi_n \in \text{Th}(\tau_n)$. Then $\xi \in \text{Th}(\tau)$ by semicontinuity of relative position; see Lemma 3.12. The assumption on the disjointness of fibers implies that $\pi(\xi) = \tau$. Thus, π is continuous.

In order to show that π is a fiber bundle, we need to construct local trivializations. Fix $\tau_0 \in A$. There exists a compact subset $S \subset G$ which is mapped by $s \mapsto s\tau_0$ homeomorphically onto a compact neighborhood of τ_0 in A . (Such a subset can be found in a slice through e transverse to P_{τ_0} .) Restricting the action $G \curvearrowright \partial_\infty X$, we obtain a topological embedding

$$S \times (\text{Th}(\tau_0) \cap G\eta) \rightarrow \text{Th}(A) \cap G\eta$$

and a local trivialization of π over a neighborhood of τ_0 in A . □

Now we turn to dynamics.

Let (g_n) be a sequence of isometries in G which preserve A , ie $g_n A = A$. We consider the action of (g_n) on $G\eta$ and derive transverse expansion from expansion on $\text{Flag}_{\tau_{\text{mod}}}$:

Lemma 7.5 (expansion implies transverse expansion) *Suppose (g_n) is on $\text{Flag}_{\tau_{\text{mod}}}$ arbitrarily expanding at $\tau_+ \in A$; ie there exist neighborhoods V_n of τ_+ in $\text{Flag}_{\tau_{\text{mod}}}$ and constants $c_n \rightarrow +\infty$ such that $g_n|_{V_n}$ is expanding with expansion factor c_n .*

Then there exist neighborhoods W_n of $\text{Th}(\tau_+) \cap G\eta$ and constants $C_n \rightarrow +\infty$ such that

$$(32) \quad d(g_n \xi, g_n \text{Th}(\tau) \cap G\eta) \geq C_n \cdot d(\xi, \text{Th}(\tau) \cap G\eta)$$

for all $\xi \in W_n$ and $\tau \in A$ with $\text{Th}(\tau) \cap G\eta \subset W_n$; compare inequality (20).

Proof To simplify notation, we write (only in this proof) $\text{Th}(\cdot)$ instead of $\text{Th}(\cdot) \cap G\eta$.

Let W_n be some (small) neighborhood of the compact subset $\text{Th}(\tau_+)$, and let ξ, τ be as in inequality (32). We work near $\text{Th}(g_n \tau_+)$. According to Lemma 7.3(ii), we can choose $g_n \tau' \in \text{Flag}_{\tau_{\text{mod}}}$ such that $g_n \xi \in \text{Th}(g_n \tau')$, and $g_n \tau'$ has controlled distance from $g_n \tau$:

$$d(g_n \tau', g_n \tau) \leq C \cdot d(g_n \xi, \text{Th}(g_n \tau))$$

with a uniform constant.

After shrinking the neighborhood $g_n W_n$ of $\text{Th}(g_n \tau_+)$, we may assume that $g_n \tau$ is close to $g_n \tau_+$, using that $\text{Th}(A)$ fibers over A ; see Lemma 7.4. Moreover, that $g_n \xi$ is close to $\text{Th}(g_n \tau)$. Thus, after shrinking W_n sufficiently, we may assume that $\tau', \tau \in V_n$.

Then

$$d(g_n\tau', g_n\tau) \geq c_n \cdot d(\tau', \tau).$$

Since we also have uniform control, ie

$$d(\xi, \text{Th}(\tau)) \leq d_H(\text{Th}(\tau'), \text{Th}(\tau)) \leq C \cdot d(\tau', \tau),$$

by Lemma 7.3(ii), it follows that

$$d(g_n\xi, \text{Th}(g_n\tau)) \geq C^{-2}c_n \cdot d(\xi, \text{Th}(\tau)),$$

that is, our assertion with $C_n = C^{-2}c_n$. □

We apply the above discussion to discrete group actions on flag manifolds.

Proposition 7.6 (transverse expansion at slim thickenings) *Let $\Gamma < G$ be a discrete subgroup, and suppose that*

- (i) *the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ is expanding at $\Lambda_{\tau_{\text{mod}}}^+$,*
- (ii) *the thickenings $\text{Th}(\tau) \cap G\eta$ of the simplices $\tau \in \Lambda_{\tau_{\text{mod}}}^+$ are pairwise disjoint.*

Then the action $\Gamma \curvearrowright G\eta$ is expanding at $\text{Th}(\Lambda_{\tau_{\text{mod}}}^+) \cap G\eta$ transversely to the natural fibration $\text{Th}(\Lambda_{\tau_{\text{mod}}}^+) \cap G\eta \rightarrow \Lambda_{\tau_{\text{mod}}}^+$ given by Lemma 7.4.

Proof Since the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ is expanding at $\Lambda_{\tau_{\text{mod}}}^+$, it is arbitrarily strongly expanding there; see Remark 5.20. That is, for every limit simplex $\tau_+ \in \Lambda_{\tau_{\text{mod}}}^+$ there exists a sequence (γ_n) in Γ and a sequence of neighborhoods V_n of τ_+ such that the $\gamma_n|_{V_n}$ are uniformly expanding with expansion factors $c_n \rightarrow +\infty$. Lemma 7.5 then implies that the action $\Gamma \curvearrowright G\eta$ is (arbitrarily) expanding at $\text{Th}(\Lambda_{\tau_{\text{mod}}}^+) \cap G\eta$ transversely to π . □

Using that transverse expansion implies cocompactness on the complement (see Proposition 5.22), we derive our main cocompactness result:

Theorem 7.7 (cocompact domains) *Let Γ and Th be as in the previous proposition. Then the action*

$$\Gamma \curvearrowright G\eta - \text{Th}(\Lambda_{\tau_{\text{mod}}}^+)$$

is cocompact.

The following is a special case of the theorem. Here, for ι -invariant τ_{mod} , a subset of $\text{Flag}_{\tau_{\text{mod}}}$ is called *antipodal* if the simplices in it are pairwise opposite; see Definition 2.3(ii).

Corollary 7.8 (cocompactness outside slim thickenings) *Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be an ι -invariant face type. Suppose that $\Gamma < G$ is a discrete subgroup such that $\Lambda_{\tau_{\text{mod}}}$ is antipodal and the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ is expanding at $\Lambda_{\tau_{\text{mod}}}$.*

(i) *Then for any slim $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\tau_{\text{mod}}})$$

is cocompact.

(ii) *More generally, suppose that $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$ is another face type and that the thickening Th is also $W_{\nu_{\text{mod}}}$ -right invariant. Then for any G -orbit $G\eta \subset \partial_{\infty} X$ of type $\bar{\eta} = \theta(\eta) \in \text{int}(\nu_{\text{mod}})$, the action*

$$\Gamma \curvearrowright G\eta - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is cocompact.

Proof We have that $\Lambda_{\tau_{\text{mod}}}^{\pm} = \Lambda_{\tau_{\text{mod}}}$, because τ_{mod} is ι -invariant. Since Th is slim and the simplices τ in $\Lambda_{\tau_{\text{mod}}}$ are pairwise antipodal, their thickenings $\text{Th}_{\text{Fü}}(\tau)$ in $\partial_{\text{Fü}} X$ are pairwise disjoint; see Lemma 3.31. Thus, the hypotheses of the theorem are satisfied. \square

Remark 7.9 (rank one) If $\text{rank}(X) = 1$, this follows from part of a basic result for Kleinian groups characterizing convex-cocompactness. Namely, the following properties are equivalent for a discrete subgroup $\Gamma < G$:

- (i) Γ is *convex-cocompact*.
- (ii) The action $\Gamma \curvearrowright \bar{X}$, equivalently, the action $\Gamma \curvearrowright \partial_{\infty} X$, is expanding at Λ .
- (iii) The (properly discontinuous) action $\Gamma \curvearrowright \bar{X} - \Lambda$ is cocompact.

In particular, the action $\Gamma \curvearrowright \partial_{\infty} X - \Lambda$ is then cocompact.

7.3 Cocompact domains of proper discontinuity

We consider the following class of discrete subgroups (see Definition 1.5 in the introduction):

Definition 7.10 (CEA subgroup) For a ι -invariant face type $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$, we call a τ_{mod} -convergence subgroup $\Gamma < G$ a τ_{mod} -CEA subgroup (convergence, expanding, antipodal) if $\Lambda_{\tau_{\text{mod}}}$ is antipodal and if the action $\Gamma \curvearrowright \text{Flag}_{\tau_{\text{mod}}}$ is expanding at $\Lambda_{\tau_{\text{mod}}}$.

Remark 7.11 (CEA versus Anosov) The class of τ_{mod} -CEA subgroups coincides with the class of $P_{\tau_{\text{mod}}}$ -Anosov subgroups, see [19, Section 6.5]. Here, $P_{\tau_{\text{mod}}}$ refers to the conjugacy class of parabolic subgroups of G corresponding to the face τ_{mod} of the spherical Weyl chamber σ_{mod} .

Combining our main results on proper discontinuity (see Theorem 6.8) and cocompactness (see Corollary 7.8), we obtain:

Theorem 7.12 (cocompact domains of proper discontinuity) *Suppose that $\Gamma < G$ is a τ_{mod} -CEA subgroup.*

- (i) *Then for any balanced $W_{\tau_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W$, the action*

$$\Gamma \curvearrowright \partial_{\text{F}\ddot{\text{u}}} X - \text{Th}_{\text{F}\ddot{\text{u}}}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous and cocompact.

- (ii) *More generally, suppose that $\nu_{\text{mod}} \subseteq \sigma_{\text{mod}}$ is another face type and that the thickening Th is also $W_{\nu_{\text{mod}}}$ -right invariant. Then for every G -orbit $G\eta \subset \partial_{\infty} X$ of type $\bar{\eta} = \theta(\eta) \in \text{int}(\nu_{\text{mod}})$, the action*

$$\Gamma \curvearrowright G\eta - \text{Th}(\Lambda_{\tau_{\text{mod}}})$$

is properly discontinuous and cocompact.

Remark 7.13 According to Corollary 3.20, balanced $W_{\tau_{\text{mod}}}$ -left invariant thickenings always exist, and Theorem 7.12 therefore provides cocompact domains of discontinuity at least in the Furstenberg boundary $\partial_{\text{F}\ddot{\text{u}}} X$.

The question whether these domains are nonempty will be addressed in Section 8.

7.4 A relation with Mumford’s geometric invariant theory

We continue the discussion in Example 3.30, now looking at actions (of Lie subgroups) on configuration spaces. (See [23; 17] for a more detailed discussion of geometric invariant theory in the context of weighted configurations.)

Let $H = \text{Isom}_o(Y)$. We consider the diagonal action $H \curvearrowright \partial_{\text{F}\ddot{\text{u}}} X$ on configurations. As we discussed in Example 3.30, the choice of a regular vector $t = (t_i) \in \text{int}(\Delta)$ determines subsets

$$(\partial_{\text{F}\ddot{\text{u}}} X)_{\text{st},t} = \partial_{\text{F}\ddot{\text{u}}} X - (\overline{\text{Th}}_t)_{\text{F}\ddot{\text{u}}}(A) \quad \text{and} \quad (\partial_{\text{F}\ddot{\text{u}}} X)_{\text{sst},t} = \partial_{\text{F}\ddot{\text{u}}} X - (\text{Th}_t)_{\text{F}\ddot{\text{u}}}(A)$$

of stable, respectively, semistable weighted configurations in $\partial_\infty Y$. Mumford’s GIT [33] defines the *Mumford quotient*

$$\partial_{\text{Fü}} X //_t H = (\partial_{\text{Fü}} X)_{\text{sst},t} // H$$

by suitably extended orbit equivalence. In the case when the thickening Th_t is balanced, all semistable points are even stable, and one has

$$\partial_{\text{Fü}} X //_t H = (\partial_{\text{Fü}} X)_{\text{sst},t} // H = (\partial_{\text{Fü}} X)_{\text{st},t} / H,$$

the latter being a quotient in the usual sense.

A nice exercise is to prove directly that the space $\partial_{\text{Fü}} X //_t H$ is compact and Hausdorff in this case. For instance, if $H = \text{PSL}(2, \mathbb{R})$, $Y = \mathbb{H}^2$, $n = 3$ and $t = (1, 1, 1)$, then $\partial_{\text{Fü}} X //_t H$ consists of exactly two points represented by configurations of three distinct points on the circle with different cyclic orders. Continuing with $Y = \mathbb{H}^2$ and letting $n = 4$, one verifies that for $t = (2, 1, 1, 1)$, the Mumford quotient is homeomorphic to S^1 , while for $t = (5, 4, 3, 1)$ the Mumford quotient is homeomorphic to the disjoint union of two circles. Taking $n = 5$, one obtains that for $t = (1, 1, 1, 1, 1)$ the Mumford quotient is the genus-4 oriented surface, while for $t = (5, 4, 1, 1, 1)$ the quotient is the disjoint union of two 2–spheres. Thus, we see that quotients are not homeomorphic for distinct choices of t .

More generally, one can describe dependence of the topology of the Mumford quotient $\partial_{\text{Fü}} X //_t H$ on the parameter t as follows.

The hyperplanes $\sum_{i \in I} t_i = \sum_{j \notin I} t_j$ (also called *interior walls*), where I runs over subsets of $\{1, \dots, n\}$, partition the chamber

$$\Delta = \{(t_1, \dots, t_n) : t_i > 0\}$$

into open convex subsets, also called *chambers*. The topology of $\partial_{\text{Fü}} X //_t H$ does not change as long as t varies in a single chamber; permuting the chambers does not change the topology either; however, *crossing through a wall* amounts to a certain Morse surgery on the manifold. This can be seen by identifying the quotients $\partial_{\text{Fü}} X //_t H$ with certain moduli spaces of polygons with fixed side-length: In the case when $H = \text{PSL}(2, \mathbb{R})$, these are polygons in the euclidean plane; see [23].

It was conjectured by Kevin Walker that if t, t' belong to chambers in distinct S_n –orbits then the Mumford quotients are not homeomorphic. This conjecture was proven 20 years later in most cases by Farber, Hausmann and Schütz [9] and in full generality by

Schütz [36]. Similar results hold when the circle is replaced by a k -sphere. In fact, different quotients are distinguished by their cohomology rings.

We will now see how the dependence of the topology of $\partial_{\text{Fü}}X //_t H$ on the parameter t described above leads to the change of the topology of quotients by *discrete* group actions.

Example 7.14 We continue with the notation of Example 3.30. For concreteness, we assume that $Y = \mathbb{H}^2$ and $H = \text{PSL}(2, \mathbb{R})$, and that $\Gamma < H$ is a torsion-free uniform lattice (a closed hyperbolic surface subgroup). The embedding $H < G = H \times \cdots \times H$ is diagonal, and we view Γ as a subgroup of G . Then Γ preserves the diagonally embedded totally geodesic hyperbolic plane $\mathbb{H}^2 \subset X$ and acts cocompactly on it. Thus, $\Lambda_{\sigma_{\text{mod}}} = \partial_{\infty}\mathbb{H}^2 \subset \partial_{\text{Fü}}X$ is the diagonally embedded circle, and $\Lambda = \partial_{\infty}\mathbb{H}^2 \subset \partial_{\infty}X$ is the visual limit set. The ideal boundary points in $\partial_{\infty}\mathbb{H}^2 \subset \partial_{\infty}X$ are contained in the central regular G -orbit $\theta^{-1}(\bar{\zeta}) \subset \partial_{\infty}X$ of type $\bar{\zeta} \in \text{int}(\sigma_{\text{mod}})$ represented by the vector $(1, \dots, 1) \in \text{int}(\Delta)$. It follows that the subgroup $\Gamma < G$ is uniformly σ_{mod} -regular (see [19] for the precise definition). More precisely, it is $\{\bar{\zeta}\}$ -regular. Moreover, the group Γ is obviously quasi-isometrically embedded in H , and hence also in G . We conclude that $\Gamma < G$ is a σ_{mod} -CEA subgroup (eg as a consequence of [20, Theorem 1.5]).

Given a balanced metric thickening $\text{Th} = \text{Th}_t \subset W$, we have that the domain $\Omega_{\text{Th}} = \partial_{\text{Fü}}X - \text{Th}_{\text{Fü}}(\Lambda_{\sigma_{\text{mod}}})$ considered in Theorem 7.12(i) equals the set $(\partial_{\text{Fü}}X)_{\text{st},t}$ of stable weighted n -point configurations on $\partial_{\infty}Y \cong S^1$ (stability being defined with respect to the weights t). The group H acts on $(\partial_{\text{Fü}}X)_{\text{st},t}$ freely, and we have a principal H -bundle

$$H \rightarrow (\partial_{\text{Fü}}X)_{\text{st},t} \rightarrow (\partial_{\text{Fü}}X)_{\text{st},t}/H = \partial_{\text{Fü}}X //_t H.$$

Dividing $(\partial_{\text{Fü}}X)_{\text{st},t}$ by Γ instead of H , we obtain a fiber bundle

$$H/\Gamma \rightarrow (\partial_{\text{Fü}}X)_{\text{st},t}/\Gamma \rightarrow (\partial_{\text{Fü}}X)_{\text{st},t}/H.$$

In particular, by taking nonhomeomorphic Mumford quotients, we obtain nonhomeomorphic quotients $\Omega_{\text{Th}}/\Gamma$. For instance, for $n = 4$ we obtain three distinct topological types of quotients: The empty quotient, a connected nonempty quotient (a bundle over the circle with the fiber H/Γ) and a disconnected quotient which is the disjoint union of two copies of an H/Γ -bundle over S^1 .

8 Nonemptiness

8.1 Thickenings and packings

We will use the following notion of ball packing for the visual boundary (using its structure as a topological spherical building).

Definition 8.1 (packing) A *packing* of $\partial_\infty X$ by $\frac{\pi}{2}$ -balls is a family \mathcal{B} of disjoint open $\frac{\pi}{2}$ -balls (with respect to the Tits metric) the union of whose closures equals $\partial_\infty X$. We call the packing *compact* if the set of centers of these balls is compact with respect to the visual topology.

Note that the set of centers of the balls is necessarily antipodal, see Definition 2.3(i). Hence the centers must have the same ι -invariant type. We call it the *type* of the packing. We call the packing *simplicial* if the balls are simplicial subcomplexes of $\partial_\infty X$. The simplicial $\frac{\pi}{2}$ -balls are precisely the $\frac{\pi}{2}$ -balls centered at points of root type, and hence a packing is simplicial if and only if it is of root type.

We will show that compact packings often do not exist.

Definition 8.2 (nonpacking type) We say that the symmetric space X is of

- (i) *nonpacking type* if $\partial_\infty X$ admits no compact packing by $\frac{\pi}{2}$ -balls,
- (ii) *non- $\bar{\vartheta}$ -packing type* for an ι -invariant type $\bar{\vartheta} \in \sigma_{\text{mod}}$ if $\partial_\infty X$ admits no compact packing by $\frac{\pi}{2}$ -balls of type $\bar{\vartheta}$,
- (iii) *non-root-packing type* if it is of non- $\bar{\vartheta}$ -packing type for *some* root type $\bar{\vartheta} \in \sigma_{\text{mod}}$.

Our motivation for proving the nonexistence of packings is that it implies via the nonfullness of thickenings, as is made precise by the next result, the nonemptiness of domains of proper discontinuity, see Proposition 8.12 below.

Let $\tau_{\text{mod}} \subseteq \sigma_{\text{mod}}$ be an ι -invariant face type. Suppose that

$$A \subset \text{Flag}_{\tau_{\text{mod}}}$$

is an antipodal compact subset. It determines for every ι -invariant type $\bar{\vartheta}_0 \in \tau_{\text{mod}}$ the antipodal compact subset $C \subset \partial_\infty X$ consisting of the points $\zeta_{\tau, \bar{\vartheta}_0} = \tau \cap \theta^{-1}(\bar{\vartheta}_0)$ of type $\bar{\vartheta}_0$ in the simplices $\tau \in A$, and hence the family of disjoint open $\frac{\pi}{2}$ -balls

$$\mathcal{B}(A, \bar{\vartheta}_0) = \left\{ B\left(\zeta_{\tau, \bar{\vartheta}_0}, \frac{\pi}{2}\right) : \tau \in A \right\}.$$

Note that the union of the closed balls is a compact subset of $\partial_\infty X$.

Proposition 8.3 (full thickenings yield packings) *Let τ_{mod} and $\bar{\vartheta}_0 \in \tau_{\text{mod}}$ be ι -invariant, and let $A \subset \text{Flag}_{\tau_{\text{mod}}}$ be an antipodal compact subset. Suppose that*

$$\text{Th}_{\text{Fü}}(A) = \partial_{\text{Fü}} X$$

for all balanced $W_{\tau_{\text{mod}}}$ -left invariant thickenings $\text{Th} \subset W$ of the form $\text{Th} = \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}$ (as defined by (11)). Then the family of balls $\mathcal{B}(A, \bar{\vartheta}_0)$ is a packing of $\partial_{\infty} X$.

Proof Suppose that $\mathcal{B}(A, \bar{\vartheta}_0)$ is not a packing. The union of the corresponding closed balls is compact in $\partial_{\infty} X$ (with respect to the visual topology), and its complement therefore open. Let ξ be a point in the complement, and let $\theta(\xi) = \bar{\vartheta}$. After perturbing ξ , we may assume that ξ is regular and that the (always fat) $W_{\tau_{\text{mod}}}$ -left invariant metric thickening $\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2} \subset W$ is balanced; see Lemma 3.19 and the proof of Corollary 3.20. By the construction of metric thickenings (see (11) and (15)),

$$\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}(\tau) \cap G\xi = \bar{B}\left(\zeta_{\tau, \bar{\vartheta}_0}, \frac{\pi}{2}\right) \cap G\xi$$

for $\tau \in A$. It follows that $\xi \notin \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}(A)$ and hence $(\text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2})_{\text{Fü}}(A) \neq \partial_{\text{Fü}} X$. \square

Remark 8.4 (full thickenings yield fibrations) Note that the hypothesis of the proposition implies in particular the existence of the following kind of fibrations of the Furstenberg boundary: In view of Lemma 7.4, it follows from $\text{Th}_{\text{Fü}}(A) = \partial_{\text{Fü}} X$ that there is a fiber bundle

$$\partial_{\text{Fü}} X \rightarrow A$$

whose fibers are finite unions of Schubert cycles (namely the thickenings $\text{Th}(\tau)$ for $\tau \in A$). Any two fibers are equivalent modulo the G -action on $\partial_{\text{Fü}} X$. If $A = \Lambda_{\tau_{\text{mod}}}$ for a subgroup $\Gamma < G$, then the fibration is Γ -equivariant.

8.2 Nonexistence of packings

We show in this section that compact packings of the visual boundary by $\frac{\pi}{2}$ -balls do not exist for most Weyl groups. Note that the discussion applies more generally to packings of *compact topological spherical buildings*.

8.2.1 Type A_2 Suppose that the symmetric space X has type A_2 . The spherical model chamber σ_{mod} is then an arc $\bar{\xi}\bar{\eta}$ of length $\frac{\pi}{3}$, with $\bar{\xi}, \bar{\eta} \in \sigma_{\text{mod}}$ the two vertex types. The involution ι of σ_{mod} is the reflection at the midpoint $\bar{\zeta}$, which is therefore the only ι -invariant type. We denote by

$$\text{Flag}_{\sigma_{\text{mod}}} \xrightarrow{\pi_{\bar{\xi}}} \text{Flag}_{\bar{\xi}} = \theta^{-1}(\bar{\xi}) \quad \text{and} \quad \text{Flag}_{\sigma_{\text{mod}}} \xrightarrow{\pi_{\bar{\eta}}} \text{Flag}_{\bar{\eta}} = \theta^{-1}(\bar{\eta})$$

the canonical projections from the full flag manifold (of chambers) to the partial flag manifolds (of vertices of fixed type).

A packing \mathcal{B} of $\partial_\infty X$ by $\frac{\pi}{2}$ -balls is necessarily of type $\bar{\xi}$ and hence simplicial. A $\frac{\pi}{2}$ -ball in $\partial_\infty X$ with center of type $\bar{\xi}$ consists of a central chamber and all chambers adjacent to it; ie it is the $\frac{\pi}{3}$ -neighborhood of its central chamber. Thus, the packing corresponds to a set $C \subset \text{Flag}_{\sigma_{\text{mod}}}$ of pairwise opposite chambers such that every other chamber is adjacent to a chamber in C .

We denote by $C_{\bar{\xi}} = \pi_{\bar{\xi}}(C)$ and $C_{\bar{\eta}} = \pi_{\bar{\eta}}(C)$ the sets of vertices of the chambers in C , and by $O_{\bar{\xi}} = \text{Flag}_{\bar{\xi}} - C_{\bar{\xi}}$ and $O_{\bar{\eta}} = \text{Flag}_{\bar{\eta}} - C_{\bar{\eta}}$ their complements. The complement of the union of the chambers in C is the union of the open $\frac{\pi}{3}$ -balls centered at the points in $O_{\bar{\xi}} \cup O_{\bar{\eta}}$; ie the chambers not in C are the chambers with a vertex in $O_{\bar{\xi}}$ or $O_{\bar{\eta}}$. We therefore have the disjoint decomposition

$$\text{Flag}_{\sigma_{\text{mod}}} = C \sqcup \pi_{\bar{\xi}}^{-1}(O_{\bar{\xi}}) \sqcup \pi_{\bar{\eta}}^{-1}(O_{\bar{\eta}}).$$

We observe that, if a chamber has a vertex in $O_{\bar{\xi}}$, then its other vertex lies in $C_{\bar{\eta}}$. Vice versa, every vertex in $C_{\bar{\eta}}$ belongs to a chamber whose other vertex lies in $O_{\bar{\xi}}$. This means that

$$(33) \quad C_{\bar{\eta}} = \pi_{\bar{\eta}}(\pi_{\bar{\xi}}^{-1}(O_{\bar{\xi}})).$$

So far, our discussion applies to packings of arbitrary spherical buildings of type A_2 . Now we take into account the visual topology.

Theorem 8.5 *If X has type A_2 , then it is of nonpacking type.*

Proof We keep the notation from the previous discussion. Suppose that \mathcal{B} is a compact packing of $\partial_\infty X$; ie C is compact and therefore also its images $C_{\bar{\xi}}$ and $C_{\bar{\eta}}$ under the projections $\pi_{\bar{\xi}}$ and $\pi_{\bar{\eta}}$. Then $O_{\bar{\xi}}$ is open. Since the projection $\pi_{\bar{\eta}}$ is open, (33) implies that $C_{\bar{\eta}}$ is also open; ie it is clopen. Since it is a nonempty proper subset, it follows that $\text{Flag}_{\bar{\eta}}$ is disconnected, and consequently also $\text{Flag}_{\sigma_{\text{mod}}}$. This is absurd, because $\text{Flag}_{\sigma_{\text{mod}}}$ is a homogeneous space of $\text{Isom}_o(X)$ and therefore connected. \square

8.2.2 Irreducible case of rank ≥ 3 For most irreducible Weyl groups, the question of the nonexistence of *simplicial* packings can be reduced to the A_2 -case.

Theorem 8.6 (nonexistence of simplicial packings in rank ≥ 3) *If X is irreducible of rank(X) ≥ 3 , then it is of non-root-packing type.*

Proof We make use of the spherical building geometry of $\partial_{\text{Tits}} X$, see [26] for a detailed discussion. The question of nonexistence can be reduced to lower rank by observing that packings of spherical buildings by $\frac{\pi}{2}$ -balls induce such packings of their spaces of directions.

The space of directions $\Sigma_\xi \partial_{\text{Tits}} X$ of a point $\xi \in \partial_\infty X$ carries again a natural spherical building structure. We will use the notation (S_ξ, W_ξ) for the associated Coxeter complex. More precisely, $\Sigma_\xi \partial_{\text{Tits}} X$ is naturally identified with the Tits building of the symmetric subspace $X' \subset X$, which appears in the decomposition $P(l) = X' \times l$ of the parallel set of a geodesic $l \subset X$ asymptotic to ξ .

Furthermore, the spaces of directions of closed $\frac{\pi}{2}$ -balls $\bar{B}(\zeta, \frac{\pi}{2})$ at boundary points $\xi \in \partial \bar{B}(\zeta, \frac{\pi}{2})$ are again $\frac{\pi}{2}$ -balls:

$$\Sigma_\xi \bar{B}(\zeta, \frac{\pi}{2}) = \bar{B}(\vec{\xi}\zeta, \frac{\pi}{2}).$$

This follows from the first variation formula in S^2 , because for any point η sufficiently close to ξ the three points ξ, η, ζ are the vertices of an embedded spherical triangle. If two open balls $B(\zeta_i, \frac{\pi}{2})$ are disjoint and if ξ is a point in the intersection of their boundaries, then the spaces of directions $\Sigma_\xi \bar{B}(\zeta_i, \frac{\pi}{2})$ have disjoint interiors.

Let now \mathcal{B} be a compact packing of $\partial_\infty X$ by $\frac{\pi}{2}$ -balls. Then \mathcal{B} induces packings \mathcal{B}_ξ by $\frac{\pi}{2}$ -balls of the spaces of directions $\Sigma_\xi \partial_{\text{Tits}} X$ for all boundary points ξ of the packing balls; the family \mathcal{B}_ξ consists of the balls $B(\vec{\xi}\zeta, \frac{\pi}{2})$ for which $B(\zeta, \frac{\pi}{2}) \in \mathcal{B}$ and $\xi \in \partial B(\zeta, \frac{\pi}{2})$. If the packing \mathcal{B} is simplicial, then so are the packings \mathcal{B}_ξ .

To see that the compactness of \mathcal{B} implies the compactness of the induced families \mathcal{B}_ξ , consider a convergent sequence $\zeta_n \rightarrow \zeta$ of centers of packing balls in \mathcal{B} such that $\angle_{\text{Tits}}(\xi, \zeta_n) = \frac{\pi}{2}$ for all n . Then $\angle_{\text{Tits}}(\xi, \zeta) \leq \frac{\pi}{2}$ by the semicontinuity of Tits distance. However, strict inequality is impossible, because then ξ would be an interior point of the packing ball $B(\zeta, \frac{\pi}{2})$, which is absurd. Thus also $\angle_{\text{Tits}}(\xi, \zeta) = \frac{\pi}{2}$ and it follows that $\vec{\xi}\zeta_n \rightarrow \vec{\xi}\zeta$. Hence the centers $\vec{\xi}\zeta_n$ of packing balls in \mathcal{B}_ξ converge to the center of such a ball. Thus the families \mathcal{B}_ξ are compact.

Let us now focus on simplicial packings. Suppose that $\partial_\infty X$ admits compact simplicial packings by $\frac{\pi}{2}$ -balls for all (at most two) root types. Consider for all such packings of $\partial_\infty X$ the induced simplicial packings by $\frac{\pi}{2}$ -balls of the spaces of directions $\Sigma_\xi \partial_{\text{Tits}} X$ for all vertices ξ in the boundaries of packing balls. Then for these, the type $\theta(\xi) \in \sigma_{\text{mod}}$ runs through all possible vertex types, and for every fixed vertex type $\theta(\xi)$, the type of the packing \mathcal{B}_ξ runs through all possible root types. (The vertex type $\theta(\xi)$

and the root type of the packing \mathcal{B}_ξ uniquely determine the root type of the packing \mathcal{B} which has to be used.)

The type $\theta(\xi)$ of a vertex ξ corresponds to a wall of the fundamental Weyl chamber, and the Dynkin diagram for the link $\Sigma_\xi \partial_{\text{Tits}} X$ is obtained from the Dynkin diagram for $\partial_{\text{Tits}} X$ by removing the corresponding node. By examining Dynkin diagrams of irreducible root systems, we note that every irreducible root system of rank ≥ 3 has a simple edge and can hence be reduced to the A_2 root system by successively removing nodes without disconnecting it. Thus, if $\text{rank}(X) \geq 3$, it follows that there exists a symmetric space X' of type A_2 whose visual boundary admits a compact packing by $\frac{\pi}{2}$ -balls. This contradicts Theorem 8.5. □

Note that root types are ι -invariant.

Regarding the irreducible case, Theorems 8.5 and 8.6 leave open the cases of type B_2 and G_2 in rank 2. We will prove some partial results for the B_2 -case in Section 8.2.4.

8.2.3 Reducible case We reduce to the irreducible case using the observation:

Lemma 8.7 *Suppose that X decomposes as the product $X = X_1 \times X_2$ of symmetric spaces. If X_1 is of non-root-packing type, then so is X .*

Proof The model chamber σ_{mod} of X splits as the spherical join

$$\sigma_{\text{mod}} = \sigma_{\text{mod}}^1 \circ \sigma_{\text{mod}}^2$$

of the model chambers of the factors. The same applies to the visual boundaries:

$$\partial_\infty X = \partial_\infty X_1 \circ \partial_\infty X_2.$$

A root type $\bar{\vartheta}_1 \in \sigma_{\text{mod}}^1$ remains a root type in σ_{mod} under the inclusion $\sigma_{\text{mod}}^1 \subset \sigma_{\text{mod}}$, and a $\frac{\pi}{2}$ -ball $\bar{B}^{\partial_\infty X}(\zeta_1, \frac{\pi}{2}) \subset \partial_\infty X$ centered at a point $\zeta_1 \in \partial_\infty X_1 \subset \partial_\infty X$ of type $\bar{\vartheta}_1$ splits as the spherical join

$$\bar{B}^{\partial_\infty X}(\zeta_1, \frac{\pi}{2}) = \bar{B}^{\partial_\infty X_1}(\zeta_1, \frac{\pi}{2}) \circ \partial_\infty X_2$$

of the ball $\bar{B}^{\partial_\infty X_1}(\zeta_1, \frac{\pi}{2}) \subset \partial_\infty X_1$ with the full visual boundary $\partial_\infty X_2$. Hence, $\partial_\infty X$ admits a compact packing by $\frac{\pi}{2}$ -balls of type $\bar{\vartheta}_1$ if and only if $\partial_\infty X_1$ does. The assertion follows. □

Combining Theorems 8.5 and 8.6 with Lemma 8.7, we obtain:

Theorem 8.8 (nonexistence of simplicial packings) *If X has at least one de Rham factor not of the type A_1 , B_2 or G_2 , then it is of non-root-packing type.*

Proof The assumptions imply that X has a de Rham factor either of type A_2 or with rank ≥ 3 . □

8.2.4 Type B_2 Suppose now that the symmetric space X has type B_2 . We obtain only partial results on the nonexistence of packings.

The model spherical chamber σ_{mod} is an arc $\bar{\xi}\bar{\eta}$ of length $\frac{\pi}{4}$, and $\bar{\xi}, \bar{\eta} \in \sigma_{\text{mod}}$ are the two vertex types. Moreover, $\iota = \text{id}_{\sigma_{\text{mod}}}$ and all types in σ_{mod} are ι -invariant. We again denote by

$$\text{Flag}_{\sigma_{\text{mod}}} \xrightarrow{\pi_{\bar{\xi}}} \text{Flag}_{\bar{\xi}} = \theta^{-1}(\bar{\xi}) \quad \text{and} \quad \text{Flag}_{\sigma_{\text{mod}}} \xrightarrow{\pi_{\bar{\eta}}} \text{Flag}_{\bar{\eta}} = \theta^{-1}(\bar{\eta})$$

the canonical projections from the full flag manifold (of chambers) to the partial flag manifolds (of vertices of fixed type).

Simplicial $\frac{\pi}{2}$ -balls in $\partial_{\text{Tits}} X$ are centered at vertices, and hence simplicial packings by $\frac{\pi}{2}$ -balls are of vertex type.

A packing $\mathcal{B}_{\bar{\vartheta}_0}$ of regular type $\bar{\vartheta}_0 \in \text{int}(\sigma_{\text{mod}})$ gives rise to a continuous family of packings $\mathcal{B}_{\bar{\vartheta}}$ with $\bar{\vartheta} \in \sigma_{\text{mod}}$ by simultaneously “sliding” its centers along the chambers containing them. Namely, we choose as the centers of $\mathcal{B}_{\bar{\vartheta}}$ the points of type $\bar{\vartheta}$ in those chambers which contain the centers of $\mathcal{B}_{\bar{\vartheta}_0}$. A regular packing thus gives rise to singular packings of *both* vertex types.

Consider now a packing \mathcal{B} of $\partial_{\infty} X$ by $\frac{\pi}{2}$ -balls of type $\bar{\xi}$ with set of centers $C_{\bar{\xi}} \subset \text{Flag}_{\bar{\xi}}$. Note that $\text{Flag}_{\bar{\eta}}$ is *partitioned* by the $\frac{\pi}{4}$ -spheres around the points in $C_{\bar{\xi}}$, and the map

$$\pi_{\bar{\xi}}^{-1} C_{\bar{\xi}} \xrightarrow{\pi_{\bar{\eta}}} \text{Flag}_{\bar{\eta}}$$

is bijective. If $C_{\bar{\xi}}$ is compact, then this map is a homeomorphism, and its inverse is a *section* of $\pi_{\bar{\eta}}$ whose image is $\pi_{\bar{\xi}}$ -saturated, ie is a union of $\pi_{\bar{\xi}}$ -fibers. Conversely, each section of $\pi_{\bar{\eta}}$ whose image is $\pi_{\bar{\xi}}$ -saturated yields a compact packing of $\partial_{\infty} X$ by $\frac{\pi}{2}$ -balls of type $\bar{\xi}$.

Question 8.9 For which groups G of type B_2 the projection $\pi_{\bar{\xi}}$ (resp. $\pi_{\bar{\eta}}$) admits a section whose image is $\pi_{\bar{\eta}}$ -saturated (resp. $\pi_{\bar{\xi}}$ -saturated)?

Example 8.10 Let $G = \text{SO}(n, 2)$ with $n \geq 2$. The partial flag manifolds in this case are the Grassmannian \mathcal{L} of isotropic lines L and the Grassmannian \mathcal{P} of isotropic planes P , and the full flag manifold is the manifold \mathcal{F} of isotropic flags (L, P) .

Fix an orthogonal splitting $\mathbb{R}^{n,2} = \mathbb{R}^n \oplus \mathbb{R}^2$ so that the quadratic form

$$q = x_1^2 + \cdots + x_n^2 - x_{n+1}^2 - x_{n+2}^2$$

is definite on each factor. Then the isotropic planes in $\mathbb{R}^{n,2}$ are the graphs of isometries $\Phi: (\mathbb{R}^2, -q|_{\mathbb{R}^2}) \rightarrow (\mathbb{R}^n, q|_{\mathbb{R}^n})$. The isotropic lines in $\mathbb{R}^{n,2}$ are the graphs of isometries $\phi: (l, -q|_l) \rightarrow (\mathbb{R}^n, q|_{\mathbb{R}^n})$ defined on lines $l \subset \mathbb{R}^2$. A full isotropic flag corresponds to a pair (Φ, l) , its isotropic line corresponding to the restriction $\Phi|_l$. Thus, we have the product splitting $\mathcal{F} \cong \mathcal{P} \times \mathbb{R}P^1$.

The projection $\pi_{\mathcal{P}}: \mathcal{F} \rightarrow \mathcal{P}$ is the projection to the first factor. It admits “constant” sections s_l by fixing l . Their images are $\pi_{\mathcal{L}}$ -saturated, namely $s_l(\mathcal{P}) = \pi_{\mathcal{L}}^{-1}\mathcal{L}_l$, where \mathcal{L}_l denotes the set of isotropic lines contained in the hyperplane $\mathbb{R}^n \oplus l \cong \mathbb{R}^{n,1} \subset \mathbb{R}^{n,2}$. The subset $\mathcal{L}_l \subset \mathcal{L}$ is compact and antipodal, and hence constitutes the set of centers of a packing of $\partial_{\infty}X$ by $\frac{\pi}{2}$ -balls. In incidence geometric terms, the antipodality corresponds to the fact that the hyperplane $\mathbb{R}^n \oplus l$ contains no isotropic plane, and the packing to the (equivalent) fact that every isotropic plane intersects $\mathbb{R}^n \oplus l$ in a line. The hyperplane $\mathbb{R}^n \oplus l$ is the orthogonal complement, in $\mathbb{R}^{n,2}$, of the line $l^{\perp} \subset \mathbb{R}^2$ orthogonal to l . Accordingly, $\mathcal{L}_l \subset \mathcal{L}$ is the orbit of a subgroup of $SO(n, 2)$ isomorphic to $SO(n, 1)$, namely of the one which fixes l^{\perp} .

The projection $\pi_{\mathcal{L}}: \mathcal{F} \rightarrow \mathcal{L}$ is given by $(\Phi, l) \mapsto \Phi|_l$. Let $\mathcal{L}_1 \subset \mathcal{L}$ denote the subset of isotropic lines which project to $l_1 = \mathbb{R}e_1 \subset \mathbb{R}^2$, ie for which the isometry ϕ is defined on l_1 . A section of $\pi_{\mathcal{L}}$ over \mathcal{L}_1 would associate with each unit vector $\phi(e_1)$ in \mathbb{R}^n a unit vector orthogonal to it, namely $\Phi(e_2)$ for the extension Φ of ϕ determined by the section. It would thus yield a unit vector field on $S^{n-1} \subset \mathbb{R}^n$. Such a vector field does not exist if n is odd.

On the other hand, if n is even, then we can use the standard identification $\mathbb{R}^{n,2} \cong \mathbb{C}^{n/2,1}$ and consider the subset $\mathcal{P}_c \subset \mathcal{P}$ of isotropic planes which are invariant under the complex structure, ie which are complex lines. Every isotropic line is contained in a unique such isotropic plane by complexification, which means that

$$\pi_{\mathcal{P}}^{-1}\mathcal{P}_c \xrightarrow{\pi_{\mathcal{L}}} \mathcal{L}$$

is a homeomorphism and, accordingly, \mathcal{P}_c is the set of centers of a packing. It is the orbit of the subgroup $SU(\frac{1}{2}n, 1) \subset SO(n, 2)$.

Our discussion and the example imply:

Theorem 8.11 (i) *If $G = O(2k + 1, 2)$ with $k \geq 1$, then $\partial_{\infty}X$ admits no compact packing by $\frac{\pi}{2}$ -balls of regular type, neither of the singular type corresponding to isotropic planes.*

- (ii) If $G = O(2k, 2)$ with $k \geq 1$, then $\partial_\infty X$ admits a compact packing by $\frac{\pi}{2}$ -balls of the singular type corresponding to isotropic planes whose set of centers is an orbit of $U(k, 1) < G$.
- (iii) If $G = O(n, 2)$ with $n \geq 2$, then $\partial_\infty X$ admits a compact packing by $\frac{\pi}{2}$ -balls of the singular type corresponding to isotropic lines whose set of centers is an orbit of $O(n, 1) < G$.

Proof (i) According to our above discussion, a packing of regular type gives rise to packings of both singular types. We assume therefore that there exists a packing of the singular type corresponding to isotropic planes, ie with centers in $\mathcal{P} \subset \partial_\infty X$. It yields a section of the fiber bundle $\pi_{\mathcal{L}}: \mathcal{F} \rightarrow \mathcal{L}$. However, such a section does not exist (see the previous example), contradiction.

(ii)–(iii) See the previous example. □

The theorem leaves open the question whether packings of regular type exist if n is even.

8.3 Nonemptiness of domains of proper discontinuity

We now apply our results on packings to discrete subgroups. Proposition 8.3 yields:

Proposition 8.12 (empty domains yield packings) *Let τ_{mod} and $\bar{\vartheta}_0 \in \tau_{\text{mod}}$ be ι -invariant. Suppose that $\Gamma < G$ is a discrete subgroup such that $\Lambda_{\tau_{\text{mod}}}$ is antipodal, and*

$$\text{Th}_{\text{F}\ddot{\text{u}}}(\Lambda_{\tau_{\text{mod}}}) = \partial_{\text{F}\ddot{\text{u}}} X$$

for all balanced $W_{\tau_{\text{mod}}}$ -left invariant thickenings $\text{Th} \subset W$ of the form $\text{Th} = \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}$ (as defined by (11)). Then the family of balls $\mathcal{B}(\Lambda_{\tau_{\text{mod}}}, \bar{\vartheta}_0)$ is a packing of $\partial_\infty X$.

Applying our nonexistence results for packings (see Theorem 8.8), we conclude that some of the domains of proper discontinuity constructed earlier (see Theorem 6.8) are nonempty. For instance, in the regular case $\tau_{\text{mod}} = \sigma_{\text{mod}}$, we obtain:

Theorem 8.13 (nonemptiness of domains of proper discontinuity) *Suppose that X has at least one de Rham factor not of the type A_1, B_2 or G_2 , and let $\Gamma < G$ be a σ_{mod} -convergence subgroup with antipodal limit set $\Lambda_{\sigma_{\text{mod}}}$. Then for some balanced thickening $\text{Th} \subset W$, the domain of proper discontinuity $\partial_{\text{F}\ddot{\text{u}}} X - \text{Th}_{\text{F}\ddot{\text{u}}}(\Lambda_{\sigma_{\text{mod}}})$ for the Γ -action (provided by Theorem 6.8) is nonempty. Moreover, the thickening can be chosen of the form $\text{Th} = \text{Th}_{\bar{\vartheta}_0, \bar{\vartheta}, \pi/2}$ (as defined by (11)) with $\bar{\vartheta}_0 \in \sigma_{\text{mod}}$ a root type.*

Proof Otherwise, by the proposition, $\partial_\infty X$ admits compact packings by $\frac{\pi}{2}$ -balls for all (of the at most two) root types. However, this contradicts Theorem 8.5, respectively, Theorem 8.6. (Note that root types are t -invariant.) \square

In the B_2 case, we can only treat a family of examples:

Addendum 8.14 (nonemptiness of domains of proper discontinuity, B_2 case) *Suppose that $G = O(2k+1, 2)$ with $k \geq 1$, and let $\Gamma < G$ be a π_{mod} -convergence subgroup with antipodal limit set $\Lambda_{\pi_{\text{mod}}}$ for the vertex type $\pi_{\text{mod}} \in \sigma_{\text{mod}}$ corresponding to isotropic planes. Then for the balanced $W_{\pi_{\text{mod}}}$ -left invariant thickening $\text{Th} \subset W = W_{B_2}$, the domain of proper discontinuity $\partial_{\text{Fü}} X - \text{Th}_{\text{Fü}}(\Lambda_{\pi_{\text{mod}}})$ for the Γ -action (provided by Theorem 6.8) is nonempty.*

Proof Otherwise, the proposition yields a compact packing of $\partial_\infty X$ by $\frac{\pi}{2}$ -balls of type π_{mod} , contradicting Theorem 8.11(i). \square

This leaves open the question whether, in the case of $G = O(2k, 2)$ for $k \geq 2$, there are σ_{mod} -convergence subgroups with antipodal limit sets, which have empty domains of proper discontinuity for *arbitrary* balanced thickenings Th . We note that Example 7.14 provides examples of σ_{mod} -CEA subgroups with empty domains of proper discontinuity in $\partial_{\text{Fü}} X$ for *some* choices of balanced thickenings.

Remark 8.15 Theorem 8.13 is both weaker and stronger than the nonemptiness results in [11, Theorems 1.11, 1.12 and 9.10]. It is stronger in the sense that it applies to hyperbolic groups Γ without assumptions on their cohomological dimension, unlike the results in [11] which require small cohomological dimension; furthermore, it applies to domains of discontinuity in various partial flag manifolds (always including $G/B = \partial_{\text{Fü}} X$), unlike the results in [11] which work (in general) only for domains of discontinuity in G/AN (which is a certain fiber bundle over G/B). On the other hand, it is weaker in the sense that it addresses only the regular case ($\tau_{\text{mod}} = \sigma_{\text{mod}}$). We also note that some examples of Anosov subgroups for which some discontinuity domains are empty are given in [11, Remark 8.5].

References

- [1] **Y Benoist**, *Propriétés asymptotiques des groupes linéaires*, *Geom. Funct. Anal.* 7 (1997) 1–47 MR
- [2] **A Björner, F Brenti**, *Combinatorics of Coxeter groups*, *Graduate Texts in Mathematics* 231, Springer (2005) MR

- [3] **A Borel, J Tits**, *Compléments à l'article: "Groupes réductifs"*, Inst. Hautes Études Sci. Publ. Math. 41 (1972) 253–276 MR
- [4] **N Bourbaki**, *Éléments de mathématique, XXXIV: Groupes et algèbres de Lie, Chapitres 4–6*, Actualités Scientifiques et Industrielles 1337, Hermann, Paris (1968) MR
- [5] **B H Bowditch**, *Convergence groups and configuration spaces*, from "Geometric group theory down under" (J Cossey, W D Neumann, M Shapiro, editors), de Gruyter, Berlin (1999) 23–54 MR
- [6] **A Cano, J P Navarrete, J Seade**, *Complex Kleinian groups*, Progress in Mathematics 303, Springer (2013) MR
- [7] **C Chevalley**, *Sur les décompositions cellulaires des espaces G/B* , from "Algebraic groups and their generalizations: classical methods" (W J Haboush, B J Parshall, editors), Proc. Sympos. Pure Math. 56, Amer. Math. Soc., Providence, RI (1994) 1–23 MR
- [8] **P B Eberlein**, *Geometry of nonpositively curved manifolds*, Univ. of Chicago Press (1996) MR
- [9] **M Farber, J-C Hausmann, D Schütz**, *On the conjecture of Kevin Walker*, J. Topol. Anal. 1 (2009) 65–86 MR
- [10] **C Frances**, *Lorentzian Kleinian groups*, Comment. Math. Helv. 80 (2005) 883–910 MR
- [11] **O Guichard, A Wienhard**, *Anosov representations: domains of discontinuity and applications*, Invent. Math. 190 (2012) 357–438 MR
- [12] **Y Guivarc'h**, *Produits de matrices aléatoires et applications aux propriétés géométriques des sous-groupes du groupe linéaire*, Ergodic Theory Dynam. Systems 10 (1990) 483–512 MR
- [13] **S Helgason**, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics 34, Amer. Math. Soc., Providence, RI (2001) MR
- [14] **J E Humphreys**, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics 29, Cambridge Univ. Press (1990) MR
- [15] **M Kapovich, B Leeb**, *Discrete isometry groups of symmetric spaces*, MSRI lecture notes (2015) arXiv To appear in *Handbook of Group Actions, IV* (L Ji, A Papadopoulos, S-T Yau, editors) International Press, Boston
- [16] **M Kapovich, B Leeb**, *Finsler bordifications of symmetric and certain locally symmetric spaces*, preprint (2015) arXiv
- [17] **M Kapovich, B Leeb, J Millson**, *Convex functions on symmetric spaces, side lengths of polygons and the stability inequalities for weighted configurations at infinity*, J. Differential Geom. 81 (2009) 297–354 MR
- [18] **M Kapovich, B Leeb, J Porti**, *Dynamics at infinity of regular discrete subgroups of isometries of higher rank symmetric spaces*, preprint (2013) arXiv

- [19] **M Kapovich, B Leeb, J Porti**, *Morse actions of discrete groups on symmetric space*, preprint (2014) arXiv
- [20] **M Kapovich, B Leeb, J Porti**, *A Morse lemma for quasigeodesics in symmetric spaces and euclidean buildings*, preprint (2014) arXiv
- [21] **M Kapovich, B Leeb, J Porti**, *Anosov subgroups: dynamical and geometric characterizations* (2016) arXiv To appear in Eur. J. Math.
- [22] **M Kapovich, B Leeb, J Porti**, *Some recent results on Anosov representations*, Transform. Groups 21 (2016) 1105–1121 MR
- [23] **M Kapovich, J Millson**, *On the moduli space of polygons in the Euclidean plane*, J. Differential Geom. 42 (1995) 133–164 MR
- [24] **M Kapovich, J J Millson**, *A path model for geodesics in Euclidean buildings and its applications to representation theory*, Groups Geom. Dyn. 2 (2008) 405–480 MR
- [25] **A Karlsson**, *On the dynamics of isometries*, Geom. Topol. 9 (2005) 2359–2394 MR
- [26] **B Kleiner, B Leeb**, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, Inst. Hautes Études Sci. Publ. Math. 86 (1997) 115–197 MR
- [27] **R S Kulkarni**, *Groups with domains of discontinuity*, Math. Ann. 237 (1978) 253–272 MR
- [28] **F Labourie**, *Anosov flows, surface groups and curves in projective space*, Invent. Math. 165 (2006) 51–114 MR
- [29] **B Leeb**, *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, Bonner Mathematische Schriften 326, Univ. Bonn, Math. Inst., Bonn (2000) MR
- [30] **P Littelmann**, *Paths and root operators in representation theory*, Ann. of Math. 142 (1995) 499–525 MR
- [31] **S A Mitchell**, *Quillen’s theorem on buildings and the loops on a symmetric space*, Enseign. Math. 34 (1988) 123–166 MR
- [32] **S A Mitchell**, *Parabolic orbits in flag varieties*, preprint (2008)
- [33] **D Mumford, J Fogarty, F Kirwan**, *Geometric invariant theory*, 3rd edition, Ergeb. Math. Grenzgeb. 34, Springer (1994) MR
- [34] **P Papasoglu, E Swenson**, *Boundaries and JSJ decompositions of $CAT(0)$ -groups*, Geom. Funct. Anal. 19 (2009) 559–590 MR
- [35] **A Parreau**, *La distance vectorielle dans les immeubles affines et les espaces symétriques*, in preparation
- [36] **D Schütz**, *The isomorphism problem for planar polygon spaces*, J. Topol. 3 (2010) 713–742 MR
- [37] **D Sullivan**, *Quasiconformal homeomorphisms and dynamics, II: Structural stability implies hyperbolicity for Kleinian groups*, Acta Math. 155 (1985) 243–260 MR

*Department of Mathematics, University of California
Davis, CA, United States*

*Mathematisches Institut, Universität München
München, Germany*

*Departament de Matemàtiques, Universitat Autònoma de Barcelona
Barcelona, Spain*

kapovich@math.ucdavis.edu, leeb@mathematik.uni-muenchen.de,
porti@mat.uab.cat

Proposed: Bruce Kleiner

Seconded: Anna Wienhard, Dmitri Burago

Received: 11 October 2015

Revised: 11 May 2017