

Topology of closed hypersurfaces of small entropy

JACOB BERNSTEIN

LU WANG

We use a weak mean curvature flow together with a surgery procedure to show that all closed hypersurfaces in \mathbb{R}^4 with entropy less than or equal to that of $\mathbb{S}^2 \times \mathbb{R}$, the round cylinder in \mathbb{R}^4 , are diffeomorphic to \mathbb{S}^3 .

53C44; 35K55, 57R65

1 Introduction

If Σ is a hypersurface, that is, a smooth properly embedded codimension-one submanifold of \mathbb{R}^{n+1} , then the *Gaussian surface area* of Σ is

$$(1-1) \quad F[\Sigma] = \int_{\Sigma} \Phi d\mathcal{H}^n = (4\pi)^{-\frac{n}{2}} \int_{\Sigma} e^{-\frac{|x|^2}{4}} d\mathcal{H}^n,$$

where \mathcal{H}^n is n -dimensional Hausdorff measure. Following Colding–Minicozzi [9], define the *entropy* of Σ to be

$$\lambda[\Sigma] = \sup_{\substack{y \in \mathbb{R}^{n+1} \\ \rho \in \mathbb{R}^+}} F[\rho\Sigma + y].$$

That is, the entropy of Σ is the supremum of the Gaussian surface area over all translations and dilations of Σ . Observe that the entropy of a hyperplane is 1. In [2], we show that for $2 \leq n \leq 6$, the entropy of a closed (ie compact and without boundary) hypersurface in \mathbb{R}^{n+1} is uniquely (modulo translations and dilations) minimized by \mathbb{S}^n , the unit sphere centered at the origin. This verifies a conjecture of Colding–Ilmanen–Minicozzi–White [8, Conjecture 0.9] (compare Ketover–Zhou [24]). We further show, in [3, Corollary 1.3], that surfaces in \mathbb{R}^3 of small entropy are topologically rigid. That is, if Σ is a closed surface in \mathbb{R}^3 and $\lambda[\Sigma] \leq \lambda[\mathbb{S}^1 \times \mathbb{R}]$, then Σ is diffeomorphic to \mathbb{S}^2 .

In this article, we use a weak mean curvature flow (see Evans–Spruck [10; 11; 12; 13] and Chen–Giga–Goto [6]) to obtain new topological rigidity for closed hypersurfaces in \mathbb{R}^4 of small entropy. This generalizes a result in [8] for closed self-shrinkers to arbitrary closed hypersurfaces and contrasts with the methods of both [8] and [3, Corollary 1.3], which both use only the classical mean curvature flow.

Theorem 1.1 *If $\Sigma \subset \mathbb{R}^4$ is a closed hypersurface with $\lambda[\Sigma] \leq \lambda[\mathbb{S}^2 \times \mathbb{R}]$, then Σ is diffeomorphic to \mathbb{S}^3 .*

A key ingredient in the proof of Theorem 1.1 is a refinement of [3, Theorem 1.1] about the topology of asymptotically conical self-shrinkers of small entropy. Recall, a hypersurface Σ is said to be *asymptotically conical* if it is smoothly asymptotic to a regular cone; ie $\lim_{\rho \rightarrow 0} \rho \Sigma = \mathcal{C}(\Sigma)$ in $C_{\text{loc}}^\infty(\mathbb{R}^{n+1} \setminus \{\mathbf{0}\})$ for $\mathcal{C}(\Sigma)$ a regular cone. A *self-shrinker* is a hypersurface Σ that satisfies

$$(1-2) \quad H_\Sigma + \frac{1}{2} \mathbf{x}^\perp = \mathbf{0},$$

where $H_\Sigma = -H_\Sigma \mathbf{n}_\Sigma = \Delta_\Sigma \mathbf{x}$ is the mean curvature vector of Σ , and \mathbf{x}^\perp is the normal component of the position vector. Let us denote the set of self-shrinkers in \mathbb{R}^{n+1} by \mathcal{S}_n and the set of asymptotically conical self-shrinkers by \mathcal{ACS}_n . Self-shrinkers generate solutions to the mean curvature flow that move self-similarly by scaling. That is, if $\Sigma \in \mathcal{S}_n$, then

$$\{\Sigma_t\}_{t \in (-\infty, 0)} = \{\sqrt{-t} \Sigma\}_{t \in (-\infty, 0)}$$

moves by mean curvature. Important examples are the maximally symmetric self-shrinking cylinders with k -dimensional spine,

$$\mathbb{S}_*^{n-k} \times \mathbb{R}^k = \{(x, y) \in \mathbb{R}^{n-k+1} \times \mathbb{R}^k = \mathbb{R}^{n+1} : |x|^2 = 2(n-k)\},$$

where $0 \leq k \leq n$. As $\mathbb{S}_*^{n-k} \times \mathbb{R}^k$ are self-shrinkers, their Gaussian surface area and entropy agree (compare [9, Lemma 7.20]). That is,

$$\lambda_n = \lambda[\mathbb{S}^n] = F[\mathbb{S}_*^n] = F[\mathbb{S}_*^n \times \mathbb{R}^l] = \lambda[\mathbb{S}^n \times \mathbb{R}^l].$$

Hence, a computation of Stone [31] gives that

$$2 > \lambda_1 > \frac{3}{2} > \lambda_2 > \dots > \lambda_n > \dots \rightarrow \sqrt{2}.$$

Theorem 1.2 *Let $\Sigma \in \mathcal{ACS}_n$ for $n \geq 2$. If $\lambda[\Sigma] \leq \lambda_{n-1}$, then Σ is contractible and $\mathcal{L}(\Sigma)$, the link of the asymptotic cone $\mathcal{C}(\Sigma)$, is a homology $(n-1)$ -sphere.*

Remark 1.3 We always consider homology with integer coefficients.

For $n = 3$, the classification of surfaces and Alexander's theorem [1] gives:

Corollary 1.4 *Let $\Sigma \in \mathcal{ACS}_3$. If $\lambda[\Sigma] \leq \lambda_2$, then Σ is diffeomorphic to \mathbb{R}^3 .*

To prove Theorem 1.1, we first establish a topological decomposition, ie Theorem 4.5, constructed from the weak mean curvature flow associated to Σ . Together with Corollary 1.4, this allows one to perform a surgery procedure which immediately gives the result. Both these steps require $n = 3$. For $n \geq 4$, one can use Theorem 1.2 and this surgery procedure to show a (strictly weaker) extension of Theorem 1.1 valid in any dimension where the two hypotheses below are satisfied. These hypotheses ensure the existence of the topological decomposition. Specifically, they ensure that if the entropy of an initial hypersurface is small enough, then tangent flows at all singularities are modeled by self-shrinkers that are either closed or asymptotically conical.

In order to state these hypotheses, first let \mathcal{S}_n^* denote the set of nonflat elements of \mathcal{S}_n , and for any $\Lambda > 0$, let

$$\mathcal{S}_n(\Lambda) = \{\Sigma \in \mathcal{S}_n : \lambda[\Sigma] < \Lambda\} \quad \text{and} \quad \mathcal{S}_n^*(\Lambda) = \mathcal{S}_n^* \cap \mathcal{S}_n(\Lambda).$$

Next let \mathcal{RMC}_n denote the space of *regular minimal cones* in \mathbb{R}^{n+1} ; that is, $\mathcal{C} \in \mathcal{RMC}_n$ if and only if it is a proper subset of \mathbb{R}^{n+1} and $\mathcal{C} \setminus \{0\}$ is a hypersurface in $\mathbb{R}^{n+1} \setminus \{0\}$ that is invariant under dilation about 0 and with vanishing mean curvature. Let \mathcal{RMC}_n^* denote the set of nonflat elements of \mathcal{RMC}_n , ie cones whose second fundamental forms do not identically vanish. For any $\Lambda > 0$, let

$$\mathcal{RMC}_n(\Lambda) = \{\mathcal{C} \in \mathcal{RMC}_n : \lambda[\mathcal{C}] < \Lambda\} \quad \text{and} \quad \mathcal{RMC}_n^*(\Lambda) = \mathcal{RMC}_n^* \cap \mathcal{RMC}_n(\Lambda).$$

Let us now fix a dimension $n \geq 3$ and a value $\Lambda > 1$. The first hypothesis is that

$$(\star_{n,\Lambda}) \quad \mathcal{RMC}_k^*(\Lambda) = \emptyset \quad \text{for all } 3 \leq k \leq n.$$

Observe that all regular minimal cones in \mathbb{R}^2 consist of unions of rays, and so $\mathcal{RMC}_1^* = \emptyset$. Likewise, $\mathcal{RMC}_2^* = \emptyset$, since great circles are the only geodesics in \mathbb{S}^2 . The second hypothesis is that

$$(\star\star_{n,\Lambda}) \quad \mathcal{S}_{n-1}^*(\Lambda) = \emptyset.$$

Obviously this holds only if $\Lambda \leq \lambda_{n-1}$. We then show the following conditional result.

Theorem 1.5 *Fix $n \geq 4$ and $\Lambda \in (\lambda_n, \lambda_{n-1}]$. If $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ both hold and Σ is a closed hypersurface in \mathbb{R}^{n+1} with $\lambda[\Sigma] \leq \Lambda$, then Σ is a homology n -sphere.*

Remark 1.6 By the results of [2] and Zhu [34], there does *not* exist a closed hypersurface Σ such that $\lambda[\Sigma] \leq \lambda_n$ unless Σ is a round sphere. Thus, we require $\Lambda > \lambda_n$ in order to make Theorem 1.5 nontrivial.

For general n and $\Lambda \in (\lambda_n, \lambda_{n-1}]$, neither the validity of $(\star_{n,\Lambda})$ nor that of $(\star\star_{n,\Lambda})$ is known. However, both can be established for $n = 3$ and $\Lambda = \lambda_2$. First, as part of their

proof of the Willmore conjecture, Marques–Neves gave a lower bound on the density of nontrivial regular minimal cones in \mathbb{R}^4 . In particular, it follows from their [25, Theorem B] that if $C \in \mathcal{RMC}_3^*$, then $\lambda[C] > \lambda_2$, and so (\star_{3,λ_2}) holds. Furthermore, it follows from [3, Corollary 1.2] that $\mathcal{S}_2^*(\lambda_2) = \emptyset$, and so $(\star\star_{3,\lambda_2})$ holds.

For $n \geq 4$, some partial results suggest that $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold for $\Lambda = \lambda_{n-1}$. For instance, Ilmanen–White [23, Theorem 1*] have shown that if $C \in \mathcal{RMC}_n^*$ is area-minimizing and topologically nontrivial, then $\lambda[C] \geq \lambda_{n-1}$. Additionally, [8, Theorem 0.1] says that the self-shrinking sphere has the lowest entropy among all compact self-shrinkers, and [8, Conjecture 0.10] posits that $(\star\star_{n,\lambda_{n-1}})$ holds for $n \leq 7$. It is important to note that there exist many topologically trivial elements of \mathcal{RMC}_n^* . Indeed, the work of Hsiang [15; 16] and Hsiang–Sterling [17] shows that there exist topologically trivial elements of \mathcal{RMC}_n^* for $n = 5, 7$ and for all even $n \geq 4$.

The paper is organized as follows. In Section 2, we introduce notation and recall some basic facts about the mean curvature flow. In Section 3, we show regularity of self-shrinking measures of low entropy. In Section 4, we study the structure of the singular set for weak mean curvature flows of small entropy. Importantly, we give a topological decomposition, Theorem 4.5, of the regular part of the flow which is the basis of the surgery procedure. In Section 5, we prove Theorem 1.2 and Corollary 1.4. Finally, in Section 6, we carry out the surgery procedure and prove Theorems 1.1 and 1.5.

Acknowledgements Bernstein was partially supported by the NSF Grants DMS-1307953 and DMS-1609340. Wang was partially supported by the Chapman Fellowship of Imperial College London, an Alfred P. Sloan Research Fellowship, and the NSF Grant DMS-1406240. Support for Wang’s research was also provided by the Vice Chancellor for the Research and Graduation Education at the University of Wisconsin–Madison with funding from the Wisconsin Alumni Research Foundation.

2 Notation and background

In this section, we fix notation for the rest of the paper and recall some background on mean curvature flow. Experts should feel free to consult this section only as needed.

2.1 Singular hypersurfaces

We will use results from [20] on weak mean curvature flows. For this reason, we follow the notation of [20] as closely as possible.

We use the notation:

- $\mathcal{M}(\mathbb{R}^{n+1}) = \{\mu : \mu \text{ is a Radon measure on } \mathbb{R}^{n+1}\}$ (see [29, Section 4]);
- $\mathcal{IM}_k(\mathbb{R}^{n+1}) = \{\mu : \mu \text{ is an integer } k\text{-rectifiable Radon measure on } \mathbb{R}^{n+1}\}$ (see [20, Section 1]);
- $\mathbf{IV}_k(\mathbb{R}^{n+1}) = \{V : V \text{ is an integer rectifiable } k\text{-varifold on } \mathbb{R}^{n+1}\}$ (see [20, Section 1] or [29, Chapter 8]).

The space $\mathcal{M}(\mathbb{R}^{n+1})$ is given the weak* topology. That is,

$$\mu_i \rightarrow \mu \iff \int f d\mu_i \rightarrow \int f d\mu \text{ for all } f \in C_c^0(\mathbb{R}^{n+1}).$$

And the topology on $\mathcal{IM}_k(\mathbb{R}^{n+1})$ is the subspace topology induced by the natural inclusion into $\mathcal{M}(\mathbb{R}^{n+1})$. For the details of the topologies considered on $\mathbf{IV}_k(\mathbb{R}^{n+1})$, we refer to [20, Section 1] or [29, Chapter 8]. There are natural bijective maps

$$V : \mathcal{IM}_k(\mathbb{R}^{n+1}) \rightarrow \mathbf{IV}_k(\mathbb{R}^{n+1}) \quad \text{and} \quad \mu : \mathbf{IV}_k(\mathbb{R}^{n+1}) \rightarrow \mathcal{IM}_k(\mathbb{R}^{n+1}).$$

The second map is continuous, but the first is not. Henceforth, we write $V(\mu) = V_\mu$ and $\mu(V) = \mu_V$.

If $\Sigma \subset \mathbb{R}^{n+1}$ is a k -dimensional smooth properly embedded submanifold, we set $\mu_\Sigma = \mathcal{H}^k \llcorner \Sigma \in \mathcal{IM}_k(\mathbb{R}^{n+1})$. Given $(y, \rho) \in \mathbb{R}^{n+1} \times \mathbb{R}^+$ and $\mu \in \mathcal{IM}_k(\mathbb{R}^{n+1})$, we define the rescaled measure $\mu^{y,\rho} \in \mathcal{IM}_k(\mathbb{R}^{n+1})$ by

$$\mu^{y,\rho}(\Omega) = \rho^k \mu(\rho^{-1}\Omega + y).$$

This is defined so that if Σ is a k -dimensional smooth properly embedded submanifold,

$$\mu_\Sigma^{y,\rho} = \mu_{\rho(\Sigma - y)}.$$

One of the defining properties of $\mu \in \mathcal{IM}_k(\mathbb{R}^{n+1})$ is that for μ -ae $x \in \mathbb{R}^{n+1}$, there is an integer $\theta_\mu(x)$ such that

$$\lim_{\rho \rightarrow \infty} \mu^{x,\rho} = \theta_\mu(x) \mu_P,$$

where P is a k -dimensional plane through the origin. When such P exists, we denote it by $T_x \mu$ and call it the *approximate tangent plane at x*. The value $\theta_\mu(x)$ is the *multiplicity of μ at x* , and by definition, $\theta_\mu(x) \in \mathbb{N}$ for μ -ae x . Notice that if $\mu = \mu_\Sigma$, then $T_x \mu = T_x \Sigma$ and $\theta_\mu(x) = 1$. Given a $\mu \in \mathcal{IM}_n(\mathbb{R}^{n+1})$, set

$$\text{reg}(\text{spt}(\mu)) = \{x \in \text{spt}(\mu) : B_\rho(x) \cap \text{spt}(\mu) \text{ is a hypersurface for some } \rho > 0\},$$

and $\text{sing}(\text{spt}(\mu)) = \text{spt}(\mu) \setminus \text{reg}(\text{spt}(\mu))$. Here $B_\rho(x)$ is the open ball in \mathbb{R}^{n+1} centered at x with radius ρ . Likewise,

$$\text{reg}(\mu) = \{x \in \text{reg}(\text{spt}(\mu)) : \theta_\mu(x) = 1\} \quad \text{and} \quad \text{sing}(\mu) = \text{spt}(\mu) \setminus \text{reg}(\mu).$$

For $\mu \in \mathcal{IM}_n(\mathbb{R}^{n+1})$, we extend the definitions of F and λ in the obvious manner:

$$F[\mu] = F[V_\mu] = \int \Phi d\mu \quad \text{and} \quad \lambda[\mu] = \lambda[V_\mu] = \sup_{\substack{\mathbf{y} \in \mathbb{R}^{n+1} \\ \rho \in \mathbb{R}^+}} F[\mu^{\mathbf{y}, \rho}].$$

2.2 Gaussian densities and tangent flows

Historically, the first weak mean curvature flow was the measure-theoretic flow introduced by Brakke [4]. This flow is called a *Brakke flow*. Brakke’s original definition considered the flow of varifolds. We use the (slightly stronger) notion introduced by Ilmanen [20, Definition 6.3]. For our purposes, the Brakke flow has two important roles. The first is the fact that Huisken’s monotonicity formula [19] holds also for Brakke flows; see [21, Lemma 7]. The second is the powerful regularity theory of Brakke [4] for such flows. In particular, we will often refer to White’s version of Brakke’s local regularity theorem [33]. We emphasize that White’s argument is valid only for a special class of Brakke flows, but that all Brakke flows considered in this paper are within this class.

A consequence of Huisken’s monotonicity formula is that if a Brakke flow $\mathcal{K} = \{\mu_t\}_{t \geq t_0}$ has bounded area ratios, then \mathcal{K} has a well-defined *Gaussian density* at every point $(\mathbf{y}, s) \in \mathbb{R}^{n+1} \times (t_0, \infty)$ given by

$$\Theta_{(\mathbf{y}, s)}(\mathcal{K}) = \lim_{t \rightarrow s^-} \int \Phi_{(\mathbf{y}, s)}(\mathbf{x}, t) d\mu_t(\mathbf{x}),$$

where

$$\Phi_{(\mathbf{y}, s)}(\mathbf{x}, t) = (4\pi)^{-\frac{n}{2}} e^{-\frac{|\mathbf{x}-\mathbf{y}|^2}{4(t-s)}}.$$

Furthermore, the Gaussian density is upper semicontinuous.

Combining the compactness of Brakke flows (compare [20, Theorem 7.1]) with the monotonicity formula, one establishes the existence of tangent flows. For a Brakke flow $\mathcal{K} = \{\mu_t\}_{t \geq t_0}$ and a point $(\mathbf{y}, s) \in \mathbb{R}^{n+1} \times (t_0, \infty)$, define a new Brakke flow

$$\mathcal{K}^{(\mathbf{y}, s), \rho} = \{\mu_t^{(\mathbf{y}, s), \rho}\}_{t \geq \rho^2(t_0 - s)},$$

where

$$\mu_t^{(\mathbf{y}, s), \rho} = \mu_{s + \rho^{-2}t}^{\mathbf{y}, \rho}.$$

Definition 2.1 Let $\mathcal{K} = \{\mu_t\}_{t \geq t_0}$ be an integral Brakke flow with bounded area ratios. A nontrivial Brakke flow $\mathcal{T} = \{\nu_t\}_{t \in \mathbb{R}}$ is a *tangent flow* to \mathcal{K} at $(\mathbf{y}, s) \in \mathbb{R}^{n+1} \times (t_0, \infty)$ if there is a sequence $\rho_i \rightarrow \infty$ such that $\mathcal{K}^{(\mathbf{y}, s), \rho_i} \rightarrow \mathcal{T}$. Denote by $\text{Tan}_{(\mathbf{y}, s)}\mathcal{K}$ the set of tangent flows to \mathcal{K} at (\mathbf{y}, s) .

The monotonicity formula implies that any tangent flow is backwardly self-similar.

Theorem 2.2 (Ilmanen [21, Lemma 8]) *Given an integral Brakke flow $\mathcal{K} = \{\mu_t\}_{t \geq t_0}$ with bounded area ratios, a point $(y, s) \in \mathbb{R}^{n+1} \times (t_0, \infty)$ with $\Theta_{(y,s)}(\mathcal{K}) \geq 1$, and a sequence $\rho_i \rightarrow \infty$, there exists a subsequence ρ_{i_j} and a $\mathcal{T} \in \text{Tan}_{(y,s)}\mathcal{K}$ such that $\mathcal{K}^{(y,s),\rho_{i_j}} \rightarrow \mathcal{T}$.*

Furthermore, $\mathcal{T} = \{v_t\}_{t \in \mathbb{R}}$ is backwardly self-similar with respect to parabolic rescaling about $(\mathbf{0}, 0)$. That is, for all $t < 0$ and $\rho > 0$,

$$v_t = v_t^{(\mathbf{0},0),\rho}.$$

Moreover, $V_{v_{-1}}$ is a stationary point of the F functional, and

$$\Theta_{(y,s)}(\mathcal{K}) = F[v_{-1}].$$

2.3 Level set flows and boundary motions

We will also need a set-theoretic weak mean curvature flow called the level-set flow. This flow was first studied in the context of numerical analysis by Osher and Sethian [27]. The mathematical theory was developed by Evans and Spruck [10; 11; 12; 13] and Chen, Giga and Goto [6]. For our purposes, it has the important advantages of being uniquely defined and satisfying a maximum principle.

A technical feature of the level-set flow is that the level sets $L(\Gamma_0) = \{\Gamma_t\}_{t \geq 0}$ may develop nonempty interiors for positive times. This phenomena is called fattening and is unavoidable for certain initial sets Γ_0 and is closely related to nonuniqueness phenomena of weak solutions of the flow. We say $L(\Gamma_0)$ is *nonfattening* if each Γ_t has no interior. It is relatively straightforward to see that the nonfattening condition is generic; see for instance [20, Theorem 11.3].

In [20], Ilmanen synthesized both notions of weak flow. In particular, he showed that for a large class of initial sets, there is a canonical way to associate a Brakke flow to the level-set flow, and observed that this allows, among other things, for the application of Brakke’s partial regularity theorem. For our purposes, it is important that the Brakke flow constructed does not vanish gratuitously. A similar synthesis may be found in [13]. The result we need is the following.

Theorem 2.3 (Ilmanen [20, Theorem 11.4]) *If Σ_0 is a closed hypersurface in \mathbb{R}^{n+1} and the level-set flow $L(\Sigma_0)$ is nonfattening, then there is a set $E \subset \mathbb{R}^{n+1} \times \mathbb{R}$ and a Brakke flow $\mathcal{K} = \{\mu_t\}_{t \geq 0}$ such that:*

- (1) $E = \{(x, t) : u(x, t) > 0\}$, where u solves the level set flow equation with initial data u_0 that satisfies $E_0 = \{x : u_0(x) > 0\}$ and $\partial E_0 = \{x : u_0(x) = 0\} = \Sigma_0$;
- (2) each $E_t = \{(x, t) \in E\}$ is of finite perimeter and $\mu_t = \mathcal{H}^n \llcorner \partial^* E_t$, where $\partial^* E_t$ is the reduced boundary of E_t .

3 Regularity of self-shrinking measures of small entropy

We establish some regularity properties of self-shrinking measures of small entropy when $n \geq 3$. We restrict to $n \geq 3$ in order to avoid certain technical complications coming from the fact that $\lambda_1 > \frac{3}{2}$.

3.1 Self-shrinking measures

We will need a singular analog of S_n . To that end, we define the set of self-shrinking measures on \mathbb{R}^{n+1} by

$$SM_n = \{\mu \in \mathcal{IM}_n(\mathbb{R}^{n+1}) : V_\mu \text{ is stationary for the } F \text{ functional, } \text{spt}(\mu) \neq \emptyset\}.$$

Clearly, if $\Sigma \in S_n$, then $\mu_\Sigma \in SM_n$. There are many examples of singular self-shrinkers. For instance, any element of $\mathcal{C} \in \mathcal{RMC}_n$ satisfies $\mu_{\mathcal{C}} = \mathcal{H}^n \llcorner \mathcal{C} \in SM_n$. For $\mu \in SM_n$, we define the associated Brakke flow $\mathcal{K} = \{\mu_t\}_{t \in \mathbb{R}}$ by

$$\mu_t = \begin{cases} 0 & \text{if } t \geq 0, \\ \mu^{0, \sqrt{-t}} & \text{if } t < 0. \end{cases}$$

One can verify that this is a Brakke flow. Given $\Lambda > 0$, set

$$SM_n(\Lambda) = \{\mu \in SM_n : \lambda[\mu] < \Lambda\} \quad \text{and} \quad SM_n[\Lambda] = \{\mu \in SM_n : \lambda[\mu] \leq \Lambda\}.$$

3.2 Regularity and asymptotic properties of self-shrinking measures of small entropy

A $\mu \in \mathcal{IM}_n(\mathbb{R}^{n+1})$ is a cone if $\mu^{0, \rho} = \mu$. Likewise, $\mu \in \mathcal{IM}_n(\mathbb{R}^{n+1})$ splits off a line if, up to an ambient rotation of \mathbb{R}^{n+1} , we have $\mu = \hat{\mu} \times \mu_{\mathbb{R}}$ for $\hat{\mu} \in \mathcal{IM}_{n-1}(\mathbb{R}^n)$. Observe that if $\mu \in SM_n$ is a cone, then V_μ is stationary (for area). Similarly, if $\mu \in SM_n$ splits off a line, then $\hat{\mu} \in SM_{n-1}$ and $\lambda[\mu] = \lambda[\hat{\mu}]$.

Standard dimension reduction arguments give the following.

Lemma 3.1 Fix $n \geq 3$ and $\Lambda \leq \frac{3}{2}$, and suppose that $(\star_{n, \Lambda})$ holds. If $\mu \in SM_n(\Lambda)$ is a cone, then $\mu = \mu_P$ for some hyperplane P .

Proof We will prove this by showing that if $(\star_{n, \Lambda})$ holds, then for all $3 \leq m \leq n$, if $\mu \in SM_m(\Lambda)$ is a cone, then $\mu = \mu_P$ for a hyperplane P in \mathbb{R}^{m+1} .

We proceed by induction on m . When $m = 3$, note that $\Lambda \leq \frac{3}{2}$ and so we have that $\mu = \mu_{\mathcal{C}}$ for some $\mathcal{C} \in \mathcal{RMC}_3$ by [2, Proposition 4.2]. Hence, by the assumption that $\mathcal{RMC}_3^*(\Lambda) = \emptyset$, we must have that \mathcal{C} is a hyperplane through the origin. To complete

the induction argument, we observe that it suffices to show that if $\mu \in \mathcal{SM}_m(\Lambda)$ is a cone, then $\mu = \mu_C$ for some $C \in \mathcal{RMC}_m(\Lambda)$. Indeed, such a C must be a hyperplane because $(\star_{n,\Lambda})$ holds and so, by definition, $\mathcal{RMC}_m^*(\Lambda) = \emptyset$ for $3 \leq m \leq n$.

To complete the proof, we argue by contradiction. Suppose that $\text{spt}(\mu)$ is not a regular cone. Then there is a point $y \in \text{sing}(\mu) \setminus \{0\}$. As V_μ is stationary, and $\mu \in \mathcal{IM}_m$ with $\lambda[\mu] < \Lambda$, we may apply Allard's integral compactness theorem (see [29, Theorem 42.7 and Remark 42.8]) to conclude that there exists a sequence $\rho_i \rightarrow \infty$ such that $\mu^{y,\rho_i} \rightarrow \nu$ and V_ν is a stationary integral varifold. Moreover, it follows from the monotonicity formula [29, Theorem 17.6] that ν is a cone; see also [29, Theorem 19.3].

As μ is a cone, ν splits off a line. That is, $\nu = \hat{v} \times \mu_{\mathbb{R}}$, where $\hat{v} \in \mathcal{IM}_{m-1}$ and $V_{\hat{v}}$ is a stationary cone, and so $\hat{v} \in \mathcal{SM}_{m-1}$. Also, by the lower semicontinuity of entropy,

$$\lambda[\hat{v}] = \lambda[\hat{v} \times \mu_{\mathbb{R}}] \leq \lambda[\mu] < \Lambda.$$

Thus, it follows from the induction hypotheses that $\hat{v} = \mu_{\hat{P}}$, where \hat{P} is a hyperplane in \mathbb{R}^m and so V_ν is a multiplicity-one hyperplane. Hence, by Allard's regularity theorem (see [29, Theorem 24.2]), $y \in \text{reg}(\mu)$, giving a contradiction. Therefore, $\mu = \mu_C$ for a $C \in \mathcal{RMC}_m(\Lambda)$. □

As a consequence, we obtain regularity for elements of $\mathcal{SM}_n(\Lambda)$ under the hypothesis that $(\star_{n,\Lambda})$ holds.

Proposition 3.2 *Fix $n \geq 3$ and $\Lambda \leq \frac{3}{2}$ and suppose that $(\star_{n,\Lambda})$ holds. If $\mu \in \mathcal{SM}_n(\Lambda)$, then $\mu = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n(\Lambda)$.*

Proof Observe that for $\mu \in \mathcal{SM}_n(\Lambda)$, the mean curvature of V_μ is locally bounded by (1-2). Following the same reasoning in the proof of Lemma 3.1, given $y \in \text{sing}(\mu)$, there exists a sequence $\rho_i \rightarrow \infty$ such that $\mu^{y,\rho_i} \rightarrow \nu$ and V_ν is a stationary cone and so $\nu \in \mathcal{SM}_n$. By the lower semicontinuity of entropy, $\lambda[\nu] \leq \lambda[\mu] < \Lambda$. Hence, together with Lemma 3.1, it follows that $\text{sing}(\mu) = \emptyset$. That is, $\text{spt}(\mu)$ is a smooth submanifold of \mathbb{R}^{n+1} that, moreover, satisfies (1-2). Finally, the entropy bound on μ implies that $\mu(B_R) \leq CR^n$ for some $C > 0$, (where B_R denotes the open ball in \mathbb{R}^{n+1} centered at the origin with radius R) and so, by [7, Theorem 1.3], $\text{spt}(\mu)$ is proper. That is, $\mu = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n$. □

If, in addition, $(\star\star_{n,\Lambda})$ holds, then we have the following.

Proposition 3.3 *Fix $n \geq 3$ and $\Lambda \leq \Lambda_{n-1}$ and suppose that both $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold. If $\mu \in \mathcal{SM}_n(\Lambda)$, then $\mu = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n(\Lambda)$, and either Σ is diffeomorphic to \mathbb{S}^n or $\Sigma \in \mathcal{ACS}_n$.*

Proof First observe that, by Proposition 3.2, $\mu = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n(\Lambda)$. If Σ is closed, then it follows from [8, Theorem 0.7] that Σ is diffeomorphic to \mathbb{S}^n . On the other hand, if Σ is not closed, then it is noncompact.

Let $\mathcal{K} = \{\mu_t\}_{t \in \mathbb{R}}$ be the Brakke flow associated to μ . Note that $\mu_t = \mu_{\sqrt{-t}\Sigma}$ for $t < 0$. Let $\mathcal{X} = \{y : y \neq \mathbf{0}, \Theta_{(y,0)}(\mathcal{K}) \geq 1\} \subset \mathbb{R}^{n+1} \setminus \{\mathbf{0}\}$. As Σ is noncompact, \mathcal{X} is nonempty. Indeed, pick any sequence of points $y_i \in \Sigma$ with $|y_i| \rightarrow \infty$. The points $\hat{y}_i = |y_i|^{-1}y_i \in |y_i|^{-1}\Sigma$. Hence, $\Theta_{(\hat{y}_i, -|y_i|^{-2})}(\mathcal{K}) \geq 1$. As the \hat{y}_i are in a compact subset, up to passing to a subsequence and relabeling, $\hat{y}_i \rightarrow \hat{y}$, and so the upper semicontinuity of Gaussian density implies that $\Theta_{(\hat{y},0)}(\mathcal{K}) \geq 1$.

We next show that \mathcal{X} is a regular cone. The fact that \mathcal{X} is a cone readily follows from the fact that \mathcal{K} is invariant under parabolic scalings. To see that $\text{sing}(\mathcal{X}) \subset \{\mathbf{0}\}$, we note that, by [2, Lemma 4.4], for any $y \in \mathcal{X}$ and $\mathcal{T} \in \text{Tan}_{(y,0)}\mathcal{K}$, $\mathcal{T} = \{\nu_t\}_{t \in \mathbb{R}}$ splits off a line. That is, up to an ambient rotation, $\nu_t = \hat{\nu}_t \times \mu_{\mathbb{R}}$ with $\{\hat{\nu}_t\}_{t \in \mathbb{R}}$ the Brakke flow associated to $\hat{\nu}_{-1} \in \mathcal{SM}_{n-1}(\Lambda)$. Here we use the lower semicontinuity of entropy. Note that $\Lambda \leq \lambda_{n-1} < \frac{3}{2}$. Thus, by Proposition 3.2 and the hypothesis that $(\star_{n,\Lambda})$ holds, $\hat{\nu}_{-1} = \mu_\Gamma$ for $\Gamma \in \mathcal{S}_{n-1}(\Lambda)$. Hence, as we assume that $(\star\star_{n,\Lambda})$ holds, Γ is a hyperplane through the origin. Therefore, it follows from Brakke’s regularity theorem that, for $t < 0$ close to 0, $\text{spt}(\mu_t)$ has uniformly bounded curvature near y and so $\sqrt{-t}\Sigma \rightarrow \mathcal{X}$ in $C_{\text{loc}}^\infty(\mathbb{R}^{n+1} \setminus \{\mathbf{0}\})$, concluding the proof. \square

As a consequence, we establish the following compactness theorem for asymptotically conical self-shrinkers of small entropy.

Corollary 3.4 *Fix $n \geq 3$, $\Lambda \leq \Lambda_{n-1}$, and $\epsilon_0 > 0$. If both $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold, then the set*

$$ACS_n[\Lambda - \epsilon_0] = \{\Sigma : \Sigma \in ACS_n, \lambda[\Sigma] \leq \Lambda - \epsilon_0\}$$

is compact in the $C_{\text{loc}}^\infty(\mathbb{R}^{n+1})$ topology.

Proof Consider a sequence $\Sigma_i \in ACS_n[\Lambda - \epsilon_0]$ and let $\mu_i = \mu_{\Sigma_i} \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. By the integral compactness theorem for F -stationary varifolds, up to passing to a subsequence, $\mu_i \rightarrow \mu$ in the sense of Radon measures. Moreover, by the lower semicontinuity of the entropy, $\mu \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. Hence, by Proposition 3.2, $\mu = \mu_\Sigma$ for $\Sigma \in \mathcal{S}_n[\Lambda - \epsilon_0]$ and so, by Allard’s regularity theorem, $\Sigma_i \rightarrow \Sigma$ in $C_{\text{loc}}^\infty(\mathbb{R}^{n+1})$. Finally, as each Σ_i is noncompact and connected, so is Σ and so, by Proposition 3.3, $\Sigma \in ACS_n[\Lambda - \epsilon_0]$, proving the claim. \square

Recall that $\mathcal{C}(\Sigma)$ denotes the asymptotic cone of any $\Sigma \in ACS_n$. Denote the link of the asymptotic cone by $\mathcal{L}(\Sigma) = \mathcal{C}(\Sigma) \cap \mathbb{S}^n$.

Proposition 3.5 Fix $n \geq 3$, $\Lambda \leq \lambda_{n-1}$, and $\epsilon_0 > 0$. If both $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold, then the set

$$\mathcal{L}_n[\Lambda - \epsilon_0] = \{\mathcal{L}(\Sigma) : \Sigma \in \mathcal{ACS}_n[\Lambda - \epsilon_0]\}$$

is compact in the $C^\infty(\mathbb{S}^n)$ topology.

Proof Consider a sequence $L_i \in \mathcal{L}_n[\Lambda - \epsilon_0]$ and let $\Sigma_i \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$ be chosen so that $\mathcal{L}(\Sigma_i) = L_i$ (observe that the Σ_i are uniquely determined by [32, Theorem 1.3]). By Corollary 3.4, up to passing to a subsequence, $\Sigma_i \rightarrow \Sigma \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$. We claim that $L_i \rightarrow L = \mathcal{L}(\Sigma)$ in $C^\infty(\mathbb{S}^n)$.

To see this, let $\mu_i = \mu_{\Sigma_i}$ and $\mu = \mu_\Sigma$ be the corresponding elements of $SM_n[\Lambda - \epsilon_0]$, and let \mathcal{K}_i and \mathcal{K} be the associated Brakke flows. Clearly, $\mu_i \rightarrow \mu$ in the sense of measures. Hence, by construction, the \mathcal{K}_i converge in the sense of Brakke flows to \mathcal{K} . Since

$$\mathcal{C}(\Sigma) = \{\mathbf{x} \in \mathbb{R}^{n+1} : \Theta_{(\mathbf{x},0)}(\mathcal{K}) \geq 1\}$$

and likewise for $\mathcal{C}(\Sigma_i)$, we have by Brakke’s regularity theorem that $\mathcal{C}(\Sigma_i) \rightarrow \mathcal{C}(\Sigma)$ in $C^\infty_{\text{loc}}(\mathbb{R}^{n+1} \setminus \{\mathbf{0}\})$; that is, $\mathcal{L}(\Sigma_i) \rightarrow \mathcal{L}(\Sigma)$ in $C^\infty(\mathbb{S}^n)$ as claimed. \square

Let B_R denote the open ball in \mathbb{R}^{n+1} centered at the origin with radius R . Combining Corollary 3.4 and Proposition 3.5 gives the following.

Corollary 3.6 Fix $n \geq 3$, $\Lambda \leq \lambda_{n-1}$, and $\epsilon_0 > 0$. Suppose that $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold. If $\Sigma \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$, then

- (1) given $\delta > 0$, there is a $\kappa \in (0, 1)$ and $\mathcal{R} > 1$ depending only on n, Λ, ϵ_0 and δ such that if $p \in \Sigma \setminus B_{\mathcal{R}}$ and $r = \kappa|\mathbf{x}(p)|$, then $\Sigma \cap B_r(p)$ can be written as a connected graph of a function v over a subset of $T_p\Sigma$ with $|Dv| \leq \delta$;
- (2) there are $R_0 = R_0(n, \Lambda, \epsilon_0)$ and $C_0 = C_0(n, \Lambda, \epsilon_0)$ such that $\Sigma \setminus \bar{B}_{R_0}$ is given by the normal graph of a smooth function u over $\mathcal{C}(\Sigma) \setminus \Omega$, where Ω is a compact set, satisfying that for $p \in \mathcal{C}(\Sigma) \setminus \Omega$,

$$|\mathbf{x}(p)||u(p)| + |\mathbf{x}(p)|^2|\nabla_{\mathcal{C}(\Sigma)}u(p)| + |\mathbf{x}(p)|^3|\nabla_{\mathcal{C}(\Sigma)}^2u(p)| \leq C_0.$$

Hence, $\Sigma \setminus B_R$ is diffeomorphic to $\mathcal{L}(\Sigma) \times [0, \infty)$ for any $R \geq R_0$.

Proof For any sequence $\Sigma_i \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$, by Corollary 3.4 and Proposition 3.5, up to passing to a subsequence, $\Sigma_i \rightarrow \Sigma$ in $C^\infty_{\text{loc}}(\mathbb{R}^{n+1})$ for some $\Sigma \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$, and $\mathcal{L}(\Sigma_i) \rightarrow \mathcal{L}(\Sigma)$ in $C^\infty(\mathbb{S}^n)$. Let \mathcal{K}_i and \mathcal{K} be the associated Brakke flows to Σ_i and Σ , respectively. As $\Sigma \in \mathcal{ACS}_n$, we have that $\mathcal{K} \lfloor (B_2 \setminus \bar{B}_1) \times [-1, 0]$ is a smooth mean curvature flow. Furthermore, since $\mathcal{K}_i \rightarrow \mathcal{K}$, it follows from Brakke’s local

regularity theorem that Σ_i have uniform curvature decay, more precisely, there exist $R, C > 0$ such that for all i and $p \in \Sigma_i \setminus B_R$,

$$\sum_{k=0}^2 |\mathbf{x}(p)|^{k+1} |\nabla_{\Sigma_i}^k A_{\Sigma_i}(p)| \leq C,$$

where A_{Σ_i} is the second fundamental form of Σ_i . As the $\mathcal{C}(\Sigma_i) \rightarrow \mathcal{C}(\Sigma)$, by [32, Lemma 2.2] and [3, Proposition 4.2], there exist $R', C' > 0$ such that items (1) and (2) in the statement hold for all Σ_i . This establishes the corollary by the arbitrariness of the Σ_i . \square

Finally, we need the fact that closed self-shrinkers of small entropy have an upper bound on their extrinsic diameter.

Proposition 3.7 *Fix $n \geq 3$, $\Lambda \leq \lambda_{n-1}$, and $\epsilon_0 > 0$. Suppose that both $(\star_{n,\Lambda})$ and $(\star\star_{n,\Lambda})$ hold. Then there is an $R_D = R_D(n, \Lambda, \epsilon_0)$ such that if $\Sigma \in \mathcal{S}_n[\Lambda - \epsilon_0]$ is closed, then $\Sigma \subset \bar{B}_{R_D}$.*

Proof We argue by contradiction. If this was not true, then there would be a sequence of $\Sigma_i \in \mathcal{S}_n[\Lambda - \epsilon_0]$ with the property that the Σ_i are closed and there are points $p_i \in \Sigma_i$ with $|p_i| \rightarrow \infty$. In particular, for each $R > \sqrt{2n}$, there is an $i_0 = i_0(R)$ such that if $i > i_0(R)$, then $\Sigma_i \cap \partial B_R \neq \emptyset$. Indeed, if this was not the case, then the mean curvature flows $\{\sqrt{-t} \Sigma\}_{t \in [-1,0)}$ and $\{\partial B_{\sqrt{R^2 - 2n(t+1)}}\}_{t \in [-1,0)}$ would violate the avoidance principle.

Let $\mu_i = \mu_{\Sigma_i} \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. By the integral compactness theorem for F -stationary varifolds, up to passing to a subsequence the μ_i converge to a $\mu \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. By Proposition 3.2, $\mu = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n[\Lambda - \epsilon_0]$. Furthermore, up to passing to a further subspace, $\Sigma_i \rightarrow \Sigma$ in $C_{\text{loc}}^\infty(\mathbb{R}^{n+1})$. It follows that $\Sigma \cap \partial B_R \neq \emptyset$ for all $R > \sqrt{2n}$. In other words, Σ is noncompact and so, by Proposition 3.3, $\Sigma \in \mathcal{ACS}_n$. However, this implies that Σ is noncollapsed (see [2, Definition 4.6]), while the Σ_i are closed and thus collapsed by [2, Lemma 4.8]. This contradicts [2, Proposition 4.10] and completes the proof. \square

4 Singularities of flows with small entropy

Given a Brakke flow $\mathcal{K} = \{\mu_t\}_{t \in I}$ and a point $(x_0, t_0) \in \text{sing}(\mathcal{K})$ with $t_0 \in \overset{\circ}{I}$, a tangent flow $\mathcal{T} \in \text{Tan}_{(x_0, t_0)} \mathcal{K}$ is of *compact type* if $\mathcal{T} = \{v_t\}_{t \in (-\infty, \infty)}$ and $\text{spt}(v_{-1})$ is compact. Otherwise, the tangent flow is of *noncompact type*. If every element of $\text{Tan}_{(x_0, t_0)} \mathcal{K}$ is of compact type, then (x_0, t_0) is a *compact singularity*. Likewise, if

every element of $\text{Tan}_{(x_0, t_0)}\mathcal{K}$ is of noncompact type, then (x_0, t_0) is a *noncompact singularity*.

For the remainder of this section, we fix a dimension $n \geq 3$ and constants $\Lambda \in (\lambda_n, \lambda_{n-1}]^1$ and $\epsilon_0 > 0$, and suppose that both $(\star_{n, \Lambda})$ and $(\star\star_{n, \Lambda})$ hold. We further assume that $\Sigma_0 \subset \mathbb{R}^{n+1}$ is a closed connected hypersurface with $\lambda[\Sigma_0] \leq \Lambda - \epsilon_0$ and with the property that the level set flow $L(\Sigma_0)$ is nonfattening and that (E, \mathcal{K}) is the pair given by Theorem 2.3.

Proposition 4.1 *Let $(x_0, t_0) \in \text{sing}(\mathcal{K})$ and $\mathcal{T} \in \text{Tan}_{(x_0, t_0)}\mathcal{K}$. If $\mathcal{T} = \{v_t\}_{t \in (-\infty, \infty)}$ is of noncompact type, then $v_{-1} = \mu_\Sigma$ for some $\Sigma \in \mathcal{ACS}_n$. Moreover, there is a constant $R_1 = R_1(n, \Lambda, \epsilon_0)$ such that for all $R \geq R_1$,*

$$\mathcal{T} \lfloor (B_{16R} \setminus \bar{B}_R) \times (-1, 1)$$

is a smooth mean curvature flow. Moreover, for all $\rho \in (R, 16R)$ and $t \in (-1, 1)$, ∂B_ρ meets $\text{spt}(v_t)$ transversally and $\partial B_\rho \cap \text{spt}(v_t)$ is connected.

Proof First, invoking Theorem 2.2 and the monotonicity formula, \mathcal{T} is backwardly self-similar with respect to parabolic scalings about $(\mathbf{0}, 0)$ and $v_{-1} \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. Furthermore, by Proposition 3.3, we have $v_{-1} = \mu_\Sigma$ for some $\Sigma \in \mathcal{ACS}_n[\Lambda - \epsilon_0]$. Finally, by Corollary 3.6, the pseudolocality property of mean curvature flow [22, Theorem 1.5]² and Brakke’s local regularity theorem, there is an $R_1 > 0$ depending only on n, Λ , and ϵ_0 such that for $R > R_1$,

$$\mathcal{T} \lfloor (B_{16R} \setminus \bar{B}_R) \times (-1, 1)$$

is a smooth mean curvature flow. Indeed, for all $t \in (-1, 1)$, the intersection $\text{spt}(v_t) \cap (B_{16R} \setminus \bar{B}_R)$ is the graph of a function over a subset of $\mathcal{C}(\Sigma)$, the asymptotic cone of Σ with small C^2 norm. As such, for all $\rho \in (R, 16R)$ and $t \in (-1, 1)$, the boundary ∂B_ρ meets $\text{spt}(v_t)$ transversally. As $\lambda[\Sigma] \leq \lambda[\Sigma_0] < \lambda_{n-1}$, it follows from [3, Theorem 1.1] that $\mathcal{L}(\Sigma)$, the link of $\mathcal{C}(\Sigma)$, is connected, and hence so is $\partial B_\rho \cap \text{spt}(v_t)$. \square

Lemma 4.2 *Each $(x_0, t_0) \in \text{sing}(\mathcal{K})$ is either a compact or a noncompact singularity.*

Proof Suppose that (x_0, t_0) is not a noncompact singularity. Then there is a $\mathcal{T} = \{v_t\}_{t \in \mathbb{R}} \in \text{Tan}_{(x_0, t_0)}\mathcal{K}$ of compact type. By the monotonicity formula and Theorem 2.2, $v_{-1} \in \mathcal{SM}_n[\Lambda - \epsilon_0]$. It follows from Proposition 3.3 that $v_{-1} = \mu_\Sigma$ for some $\Sigma \in \mathcal{S}_n[\Lambda - \epsilon_0]$ and Σ is closed. Hence, by [28, Corollary 1.2], \mathcal{T} is the only element of $\text{Tan}_{(x_0, t_0)}\mathcal{K}$ and so (x_0, t_0) is a compact singularity, proving the claim. \square

¹The reader may refer to Remark 1.6 for the reason that we restrict to $\Lambda > \lambda_n$.

²The proof of [22, Theorem 1.5] uses the local regularity theorem of White, which is also applicable to the Brakke flows in Theorem 2.3 and their tangent flows; see [33, pages 1487–1488].

Theorem 4.3 Given $(\mathbf{x}_0, t_0) \in \text{sing}(\mathcal{K})$, there exist $\rho_0 = \rho_0(\mathbf{x}_0, t_0, \mathcal{K}) > 0$ and $\alpha = \alpha(n, \Lambda, \epsilon_0) > 1$ such that the following hold:

(1) If (\mathbf{x}_0, t_0) is a compact singularity and $\rho < \rho_0$, then

$$\mathcal{K} \lfloor (B_{2\alpha\rho}(\mathbf{x}_0) \times (t_0 - 4\alpha^2\rho^2, t_0 + 4\alpha^2\rho^2) \setminus \{(\mathbf{x}_0, t_0)\})$$

is a smooth mean curvature flow. Furthermore, $\text{spt}(\mu_t) \cap \partial B_R(\mathbf{x}_0) = \emptyset$ for all $R \in (\frac{1}{2}\alpha\rho, 2\alpha\rho)$ and $t \in (t_0 - \rho^2, t_0 + \rho^2)$.

(2) If (\mathbf{x}_0, t_0) is a noncompact singularity and $\rho < \rho_0$, then

$$\mathcal{K} \lfloor (B_{2\alpha\rho}(\mathbf{x}_0) \times (t_0 - 4\alpha^2\rho^2, t_0] \setminus \{(\mathbf{x}_0, t_0)\})$$

and

$$\mathcal{K} \lfloor (B_{2\alpha\rho}(\mathbf{x}_0) \setminus \bar{B}_{\frac{1}{2}\alpha\rho}(\mathbf{x}_0)) \times (t_0 - \rho^2, t_0 + \rho^2)$$

are both smooth mean curvature flows. Furthermore, for all $R \in (\frac{1}{2}\alpha\rho, 2\alpha\rho)$ and $t \in (t_0 - \rho^2, t_0 + \rho^2)$, the boundary $\partial B_R(\mathbf{x}_0)$ meets $\text{spt}(\mu_t)$ transversally, and the intersection is connected.

Finally, for all $t \in (t_0 - \rho^2, t_0)$, we have that $\text{spt}(\mu_t) \cap \bar{B}_{\alpha\rho}(\mathbf{x}_0)$ is diffeomorphic (possibly as a manifold with boundary) to $\Gamma \cap \bar{B}_\alpha$, where $\Gamma \in \mathcal{S}_n^*[\Lambda - \epsilon_0]$, and if $\Gamma \in \mathcal{ACS}_n$, then $\Gamma \setminus B_\alpha$ is diffeomorphic to $\mathcal{L}(\Gamma) \times [0, \infty)$.

Proof Set $\alpha = 4 \max\{R_1, R_D, 1\}$, where R_1 is given by Proposition 4.1 and R_D is given by Proposition 3.7. Without loss of generality, we may assume that $(\mathbf{x}_0, t_0) = (\mathbf{0}, 0)$.

We establish the regularity near (but not at) $(\mathbf{0}, 0)$ by contradiction. To that end, suppose that there was a sequence of points $(\mathbf{x}_i, t_i) \in \text{sing}(\mathcal{K}) \setminus \{(\mathbf{0}, 0)\}$ such that $(\mathbf{x}_i, t_i) \rightarrow (\mathbf{0}, 0)$. If $(\mathbf{0}, 0)$ is a noncompact singularity, we further assume $t_i \leq 0$. Let $r_i^2 = |\mathbf{x}_i|^2 + |t_i|$. Then up to passing to a subsequence, it follows from Theorem 2.2 that $\mathcal{K}^{(\mathbf{0}, 0), r_i} \rightarrow \mathcal{T}$ in the sense of Brakke flows and $\mathcal{T} = \{v_t\}_{t \in \mathbb{R}} \in \text{Tan}_{(\mathbf{0}, 0)}\mathcal{K}$. Let $\tilde{\mathbf{x}}_i = r_i^{-1}\mathbf{x}_i$ and $\tilde{t}_i = r_i^{-2}t_i$. Then $|\tilde{\mathbf{x}}_i|^2 + |\tilde{t}_i| = 1$; that is, $(\tilde{\mathbf{x}}_i, \tilde{t}_i)$ lies on the unit parabolic sphere in spacetime. Thus, up to passing to a subsequence, $(\tilde{\mathbf{x}}_i, \tilde{t}_i) \rightarrow (\tilde{\mathbf{x}}_0, \tilde{t}_0)$, where $|\tilde{\mathbf{x}}_0|^2 + |\tilde{t}_0| = 1$. Moreover, the upper semicontinuity of Gaussian density implies that $\Theta_{(\tilde{\mathbf{x}}_0, \tilde{t}_0)}(\mathcal{T}) \geq 1$.

As $v_{-1} \in \mathcal{SM}_n[\Lambda - \epsilon_0]$, Proposition 3.3 implies that $\text{sing}(v_t) = \emptyset$ for $t < 0$. That is, $(\tilde{\mathbf{x}}_0, \tilde{t}_0)$ is a regular point of \mathcal{T} if $\tilde{t}_0 < 0$. If $(\mathbf{0}, 0)$ is a noncompact singularity, then \mathcal{T} is of noncompact type and $\tilde{t}_0 \leq 0$. Hence, either $(\tilde{\mathbf{x}}_0, \tilde{t}_0)$ is a regular point or $\tilde{t}_0 = 0$ and $|\tilde{\mathbf{x}}_0| = 1$. However in the later case, Proposition 4.1 applied to $\mathcal{T}^{(\mathbf{0}, 0), \alpha} \in \text{Tan}_{(\mathbf{0}, 0)}\mathcal{K}$ implies that $(\tilde{\mathbf{x}}_0, \tilde{t}_0)$ is also a regular point of \mathcal{T} . If $(\mathbf{0}, 0)$ is a compact singularity, then \mathcal{T} is of compact type and $v_{-1} = \mu_\Gamma$ for some $\Gamma \in \mathcal{S}_n(\Lambda)$ by Proposition 3.3. This

implies that \mathcal{T} is extinct at time 0 and $\text{sing}(\mathcal{T}) = \{(\mathbf{0}, 0)\}$, again implying that $\tilde{t}_0 \leq 0$ and $(\tilde{\mathbf{x}}_0, \tilde{t}_0)$ is a regular point of \mathcal{T} . Hence, it follows from Brakke’s local regularity theorem that for all i sufficiently large, $(\tilde{\mathbf{x}}_i, \tilde{t}_i) \notin \text{sing}(\mathcal{K}^{(\mathbf{0},0),r_i})$, or equivalently, $(\mathbf{x}_i, t_i) \notin \text{sing}(\mathcal{K})$. This is the desired contradiction. Therefore, for $\rho'_0 > 0$ sufficiently small, if $\rho < \rho'_0$ and $(\mathbf{0}, 0)$ is a noncompact singularity, then

$$\mathcal{K} \lfloor (B_{2\alpha\rho} \times (-4\alpha^2\rho^2, 0] \setminus \{(\mathbf{0}, 0)\})$$

is a smooth mean curvature flow, while if $\rho < \rho'_0$ and $(\mathbf{0}, 0)$ is a compact singularity,

$$\mathcal{K} \lfloor (B_{2\alpha\rho} \times (-4\alpha^2\rho^2, 4\alpha^2\rho^2) \setminus \{(\mathbf{0}, 0)\})$$

is a smooth mean curvature flow.

We continue arguing by contradiction and again consider a sequence ρ_i of positive numbers with $\rho_i \rightarrow 0$ and $\rho_i < \rho'_0$. Up to passing to a subsequence, $\mathcal{K}^{(\mathbf{0},0),\rho_i}$ converges, in the sense of Brakke flows, to some $\mathcal{T} = \{v_t\}_{t \in \mathbb{R}} \in \text{Tan}_{(\mathbf{0},0)}\mathcal{K}$. If $(\mathbf{0}, 0)$ is a compact singularity, then as $\alpha \geq 4R_D$, we have $\partial B_R \cap \text{spt}(v_t) = \emptyset$ for $R \geq \frac{1}{2}\alpha$ and $t \in (-1, 1)$ by Proposition 3.7 and the avoidance principle. Hence, the nature of the convergence implies that, for ρ_i sufficiently large, $\partial B_R \cap \text{spt}(\mu_t) = \emptyset$ for $t \in (-\rho_i^2, \rho_i^2)$ and $R \in (\frac{1}{2}\alpha\rho_i, 2\alpha\rho_i)$. If $(\mathbf{0}, 0)$ is a noncompact singularity, then Proposition 4.1 implies that

$$\mathcal{T} \lfloor (B_{4\alpha} \setminus \bar{B}_{\frac{1}{4}\alpha}) \times (-1, 1)$$

is a smooth mean curvature flow and that ∂B_R meets $\text{spt}(v_t)$ transversally and as a connected set for all $R \in (\frac{1}{4}\alpha, 4\alpha)$ and $t \in (-1, 1)$. Thus, by Brakke’s local regularity theorem, for all i sufficiently large,

$$\mathcal{K}^{(\mathbf{0},0),\rho_i} \lfloor (B_{2\alpha} \setminus \bar{B}_{\frac{1}{2}\alpha}) \times (-1, 1)$$

is a smooth mean curvature flow, and hence so is

$$\mathcal{K} \lfloor (B_{2\alpha\rho_i} \setminus \bar{B}_{\frac{1}{2}\alpha\rho_i}) \times (-\rho_i^2, \rho_i^2).$$

Moreover, for all $R \in (\frac{1}{2}\alpha\rho_i, 2\alpha\rho_i)$ and $t \in (-\rho_i^2, \rho_i^2)$, the boundary ∂B_R meets μ_t transversally and as a connected set. Hence, as the sequence ρ_i was arbitrary, there is a $\rho''_0 < \rho'_0$ such that items (1) and (2) hold for $\rho < \rho''_0$.

To complete the proof, we observe that again arguing by contradiction, there is a $\rho_0 < \rho''_0$ such that if $\rho < \rho_0$, then $B_{2\alpha} \cap \rho^{-1} \text{spt}(\mu_{-\rho^2})$ is a normal graph over a smooth domain³ Ω in Γ with small C^2 norm for some $\Gamma \in \mathcal{S}_n[\Lambda - \epsilon_0]$. In particular, as $\alpha \geq 4R_1 \geq 4R_0$, by Corollary 3.6, $\partial\Omega$ is a small normal graph over $\partial B_{2\alpha} \cap \Gamma$,

³Given a manifold M we say a subset $U \subset M$ is a *smooth domain* if U is open and ∂U is a smooth submanifold.

so $\bar{B}_{\alpha\rho} \cap \text{spt}(\mu_{-\rho^2})$ is diffeomorphic to $\bar{B}_\alpha \cap \Gamma$. Furthermore, the choice of α ensures that if $\Gamma \in \mathcal{ACS}_n$, then $\Gamma \setminus B_\alpha$ is diffeomorphic to $\mathcal{L}(\Sigma) \times [0, \infty)$. It remains only to show that $\bar{B}_{\alpha\rho} \cap \text{spt}(\mu_t)$ is diffeomorphic to $\bar{B}_\alpha \cap \Gamma$ for $t \in (-\rho^2, 0)$. This follows from the fact that, as already established, the flow is smooth in $\bar{B}_{2\alpha\rho} \times [-2\rho^2, 0)$, and for all $t \in [-\rho^2, 0)$, either $\partial B_{\alpha\rho} \cap \text{spt}(\mu_t) = \emptyset$ (if the singularity is compact) or the intersection is transverse (if the singularity is noncompact). As such, the flow provides a diffeomorphism between $\bar{B}_{\alpha\rho} \cap \text{spt}(\mu_t)$ and $\bar{B}_{\alpha\rho} \cap \text{spt}(\mu_{-\rho^2})$; see the appendix. \square

Corollary 4.4 *For each $t_0 > 0$, the set $\text{sing}_{t_0}(\mathcal{K}) = \{\mathbf{x} : (\mathbf{x}, t_0) \in \text{sing}(\mathcal{K})\}$ is finite.*

Theorem 4.5 *There is an $N = N(\Sigma_0) \in \mathbb{N}$ and a sequence of closed connected hypersurfaces $\Sigma^1, \dots, \Sigma^N$ such that the following hold:*

- (1) $\Sigma^1 = \Sigma_0$.
- (2) Σ^N is diffeomorphic to S^n .
- (3) *For each i with $1 \leq i \leq N - 1$, there is an $m = m(i) \in \mathbb{N}$ and open connected pairwise disjoint smooth domains $U_1^i, \dots, U_{m(i)}^i \subset \Sigma^i$ and $V_1^i, \dots, V_{m(i)}^i \subset \Sigma^{i+1}$ such that*
 - *there are orientation-preserving diffeomorphisms*

$$\hat{\Phi}^i : \Sigma^{i+1} \setminus \bigcup_{j=1}^{m(i)} V_j^i \rightarrow \Sigma^i \setminus \bigcup_{j=1}^{m(i)} U_j^i;$$

- *each \bar{U}_j^i is diffeomorphic to $\bar{B}_{R_j^i} \cap \Gamma_j^i$, where $\Gamma_j^i \in \mathcal{ACS}_n^*(\Lambda)$ and $\Gamma_j^i \setminus B_{R_j^i}$ is diffeomorphic to $\mathcal{L}(\Gamma_j^i) \times [0, \infty)$.*

Proof Let us denote the set of compact singularities of \mathcal{K} by $\text{sing}^C(\mathcal{K})$ and the set of noncompact singularities by $\text{sing}^{NC}(\mathcal{K})$. By Lemma 4.2, $\text{sing}(\mathcal{K}) = \text{sing}^{NC}(\mathcal{K}) \cup \text{sing}^C(\mathcal{K})$. We note that if $X \in \text{sing}^{NC}(\mathcal{K})$, then by Proposition 3.3, every element of $\text{Tan}_X \mathcal{K}$ is the flow of an element of \mathcal{ACS}_n and so the tangent flows are noncollapsed at time 0 in the sense of [2, Definition 4.9]. Hence, by [2, Lemma 5.1], $\text{sing}^C(\mathcal{K}) \neq \emptyset$. In fact, if we define the extinction time of \mathcal{K} to be

$$T(\mathcal{K}) = \sup \{t : \text{spt}(\mu_t) \neq \emptyset\},$$

then

$$\emptyset \neq \{\mathbf{x} \in \mathbb{R}^{n+1} : \Theta_{(\mathbf{x}_0, T(\mathcal{K}))}(\mathcal{K}) \geq 1\} = \{\mathbf{x} \in \mathbb{R}^{n+1} : (\mathbf{x}, T(\mathcal{K})) \in \text{sing}^C(\mathcal{K})\}.$$

It follows from Theorem 4.3 that $\text{sing}^C(\mathcal{K})$ consists of at most a finite number of points.

Observe that if $\text{sing}(\mathcal{K})$ consists of exactly one point X_0 , then we can take $N = 1$. Indeed, by the above discussion, this singularity must be compact and hence, by Proposition 3.3, there is a $\Gamma \in S_n(\Lambda)$ diffeomorphic to S^n such that one of the tangent

flows at X_0 is the flow associated to μ_Γ . In this case we may write $\mathcal{K} = \{\mu_{\Sigma_t}\}_{t \in [0, T(\mathcal{K})]}$ where $\{\Sigma_t\}_{t \in [0, T(\mathcal{K})]}$ is a smooth mean curvature flow. By Brakke's regularity theorem, there is a t near $T(\mathcal{K})$ such that Σ_t is a small normal graph over Γ and hence $\Sigma^1 = \Sigma_0$ is diffeomorphic to Γ , verifying the claim.

Now let $ST(\mathcal{K}) = \{t \in \mathbb{R} : (\mathbf{x}, t) \in \text{sing}(\mathcal{K})\}$ be the set of singular times. Notice that by Corollary 4.4 there are at most a finite number of singular points associated to each singular time. We observe that as $\Sigma^1 = \Sigma_0$ is smooth, there is a $\delta > 0$ such that $ST(\mathcal{K}) \subset [\delta, T(\mathcal{K})]$. Furthermore, as $\text{sing}(\mathcal{K})$ is a closed set, so is $ST(\mathcal{K})$.

For each $t \in ST(\mathcal{K})$, let

$$\rho(t) = \min \{\rho_0(\mathbf{x}, t, \mathcal{K}) : \mathbf{x} \in \text{sing}_t(\mathcal{K})\} > 0,$$

where $\rho_0(\mathbf{x}, t, \mathcal{K})$ is the constant given by Theorem 4.3. This minimum is positive as $\text{sing}_t(\mathcal{K})$ is a finite set. Observe that by Theorem 4.3,

$$(4-1) \quad B_{\alpha\rho(t)}(\mathbf{x}) \cap B_{\alpha\rho(t)}(\mathbf{x}') = \emptyset$$

when \mathbf{x} and \mathbf{x}' are distinct elements of $\text{sing}_t(\mathcal{K})$, and $\alpha = \alpha(n, \Lambda, \epsilon_0)$ is given by Theorem 4.3. Next, choose $\tau(t) \in (0, \rho^2(t))$ so that

$$\mathcal{K} \lfloor \left(\mathbb{R}^{n+1} \setminus \bigcup_{\mathbf{x} \in \text{sing}_t(\mathcal{K})} \bar{B}_{\alpha\rho(t)}(\mathbf{x}) \right) \times (t - \tau(t), t + \tau(t))$$

is a smooth mean curvature flow. Such a τ exists as $\text{sing}(\mathcal{K})$ is a closed set.

As $ST(\mathcal{K})$ is a closed subset of $[0, T(\mathcal{K})]$, it is a compact set and so the open cover

$$\{(t - \tau(t), t + \tau(t)) : t \in ST(\mathcal{K})\}$$

of $ST(\mathcal{K})$ has a finite subcover. That is, there are a finite number of times $t_1, \dots, t_{N'} \in ST(\mathcal{K})$, labeled so that $t_i < t_{i+1}$ and chosen so that

$$ST(\mathcal{K}) \subset \bigcup_{i=1}^{N'} (t_i - \tau(t_i), t_i + \tau(t_i)).$$

Furthermore, we can assume that for each i ,

- (1) $t_i - \tau(t_i) < t_j - \tau(t_j)$ for all $j > i$,
- (2) $t_i + \tau(t_i) > t_j + \tau(t_j)$ for all $j < i$, and
- (3) $t_j + \tau(t_j) < t_{j'} - \tau(t_{j'})$ for all $j < i < j'$,

as otherwise we could delete $(t_i - \tau(t_i), t_i + \tau(t_i))$ and still have an open cover. Note that, by the definition of $\tau(t)$, one must have $t_{N'} = T(\mathcal{K})$.

By Theorem 4.3 we may choose a sequence of points $s_1^\pm, \dots, s_{N'}^\pm$, with $t_i \in (s_i^-, s_i^+)$ $|s_i^\pm - t_i| < \tau(t_i)$, $s_i^+ \leq s_{i+1}^-$ and so that

$$\left([0, s_1^-] \cup \bigcup_{i=1}^{N'-1} [s_i^+, s_{i+1}^-] \right) \cap \text{ST}(\mathcal{K}) = \emptyset.$$

More concretely, first take $s_1^- \in (t_1 - \tau(t_1), t_1)$ with $s_1^- > 0$ and $s_{N'}^+ = t_{N'} + \frac{1}{2}\tau(t_{N'})$. For $1 \leq i \leq N' - 1$, let

$$\tilde{s}_i^+ = \sup(\text{ST}(\mathcal{K}) \cap (t_i - \tau(t_i), t_i + \tau(t_i))),$$

and for $2 \leq i \leq N'$, let

$$\tilde{s}_i^- = \inf(\text{ST}(\mathcal{K}) \cap (t_i - \tau(t_i), t_i + \tau(t_i))).$$

The definition of $\tau(t_i)$ and Theorem 4.3 imply that $\tilde{s}_i^- = t_i$. As the set of singular times is closed and $t_i \in \text{ST}(\mathcal{K})$, we have $\tilde{s}_i^+ \in \text{ST}(\mathcal{K})$ and $t_i \leq \tilde{s}_i^+$. We treat two cases. In the first case we suppose that $t_{i+1} - \tau(t_{i+1}) < t_i + \tau(t_i)$. As $\tilde{s}_{i+1}^- = t_{i+1}$, there are then no singular times in the interval $(t_{i+1} - \tau(t_{i+1}), t_i + \tau(t_i))$ and so we may take $s_i^+ = s_{i+1}^-$ to be the same point in this interval. In the second case, we suppose that $t_i + \tau(t_i) \leq t_{i+1} - \tau(t_{i+1})$ and observe that $\tilde{s}_i^+ \leq t_i + \tau(t_i) \leq t_{i+1} - \tau(t_{i+1})$. In fact, $\tilde{s}_i^+ < t_i + \tau(t_i)$, as otherwise, in order to cover $\text{ST}(\mathcal{K})$, assumption (3) from above would not hold. Pick s_i^+ as some point in $(\tilde{s}_i^+, t_i + \tau(t_i))$ and s_{i+1}^- as some point in $(t_{i+1} - \tau(t_{i+1}), t_i)$. The lack of singular times in $[0, s_1^-]$ and in each $[s_i^+, s_{i+1}^-]$ follows by our choices and assumptions (1) and (3) above.

For $1 \leq i \leq N'$, set $\Sigma_\pm^i = \text{spt}(\mu_{s_i^\pm})$. By the choice of s_i^\pm , each Σ_\pm^i is a closed hypersurface, and as there are no singular times between s_i^+ and s_{i+1}^- , we have for $1 \leq i \leq N' - 1$ diffeomorphisms $\Phi^i: \Sigma_+^i \rightarrow \Sigma_-^{i+1}$ coming from the flow, and for the same reason, a diffeomorphism $\Phi^0: \Sigma^1 \rightarrow \Sigma_-^1$. Observe that, *a priori*, the Σ_\pm^i need not consist of one component (indeed, $\Sigma_+^{N'}$ is empty). By Corollary 4.4, $\text{sing}_{t_i}(\mathcal{K})$ is finite for each $1 \leq i \leq N'$ and we write

$$\{\mathbf{x}_i^1, \dots, \mathbf{x}_i^{M(i)}\} = \text{sing}_{t_i}(\mathcal{K});$$

ie the (\mathbf{x}_i^j, t_i) are the singular points of the flow at time t_i . Up to relabeling, there is an $0 \leq m(i) \leq M(i)$ such that $(\mathbf{x}_i^j, t_i) \in \text{sing}^{\text{NC}}(\mathcal{K})$ for $1 \leq j \leq m(i)$, while $(\mathbf{x}_i^j, t_i) \in \text{sing}^{\text{C}}(\mathcal{K})$ for $m(i) < j \leq M(i)$. Set $R^i = \alpha\rho(t_i)$, and for each \mathbf{x}_i^j , let $U_{j,\pm}^i \subset \Sigma_\pm^i$ be the sets $B_{R^i}(\mathbf{x}_i^j) \cap \Sigma_\pm^i$. By (4-1) for fixed j , these are pairwise disjoint sets, and by Theorem 4.3, these intersections are transverse and so the $\sigma_{j,\pm}^i = \partial U_{j,\pm}^i$ are submanifolds of Σ_\pm^i . Hence, the $U_{j,\pm}^i$ are smooth pairwise disjoint domains.

Furthermore, by Theorem 4.3 and fact that $\tau(t) < \rho(t)$, each $\bar{U}_{j,-}^i$ is diffeomorphic to $\bar{B}_\alpha \cap \Gamma_j^i$ for some $\Gamma_j^i \in \mathcal{S}_n$. In particular, for $j > m(i)$ we have that $\bar{U}_{j,-}^i$ is a closed connected hypersurface, while for $1 \leq j \leq m(i)$, we have that $\partial \bar{U}_{j,-}^i$ is nonempty and connected. Hence $\bar{U}_{j,+}^i = \emptyset$ for $j > m(i)$, while $\partial \bar{U}_{j,+}^i$ is nonempty and connected for $1 \leq j \leq m(i)$. Furthermore, Theorem 4.3 implies that there are diffeomorphisms (see the appendix)

$$\Psi^i: \Sigma_-^i \setminus \bigcup_{j=1}^{M(i)} U_{j,-}^i \rightarrow \Sigma_+^i \setminus \bigcup_{j=1}^{M(i)} U_{j,+}^i.$$

As Σ^1 is connected and $\Phi^0(\Sigma^1) = \Sigma_-^1$, Σ_-^1 is also connected. As each $\sigma_{j,-}^1$ is connected, we obtain that

$$\hat{\Sigma}_-^1 = \Sigma_-^1 \setminus \bigcup_{j=1}^{M(1)} U_{j,-}^1$$

is connected. Let $\tilde{\Sigma}_+^1$ be the connected component of Σ_+^1 that contains $\Psi^1(\hat{\Sigma}_-^1)$. Inductively, let $\tilde{\Sigma}_-^{i+1} = \Phi^i(\tilde{\Sigma}_+^i)$ and

$$\hat{\Sigma}_-^{i+1} = \tilde{\Sigma}_-^{i+1} \setminus \bigcup_{j=1}^{M(i+1)} U_{j,-}^{i+1}$$

and define $\tilde{\Sigma}_+^{i+1}$ to be the connected component of Σ_+^{i+1} that contains $\Psi^{i+1}(\hat{\Sigma}_-^{i+1})$. Here we adopt the convention that if $\hat{\Sigma}_-^{i+1} = \emptyset$, then $\tilde{\Sigma}_+^{i+1} = \emptyset$. It follows inductively that each $\tilde{\Sigma}_\pm^i$ is connected. Let $\tilde{\Phi}^i: \tilde{\Sigma}_+^i \rightarrow \tilde{\Sigma}_-^{i+1}$ be the diffeomorphisms given by restricting the Φ^i . To be consistent, we also set $\tilde{\Sigma}_-^1 = \Sigma_-^1$ and $\tilde{\Phi}^0 = \Phi^0$.

Finally let

$$N = \max\{1 \leq i \leq N' : \tilde{\Sigma}_-^k \neq \emptyset \text{ for all } 1 \leq k \leq i\}.$$

If $N < N'$, then by constructions, $\hat{\Sigma}_-^N = \emptyset$ and $\tilde{\Sigma}_-^N = U_{j,-}^N$ for some $j > m(N)$. If $N = N'$, then $t_N = T(\mathcal{K})$ at which all singularities are compact. Thus it follows from [8, Theorem 0.7] that $\tilde{\Sigma}_-^N$ is diffeomorphic to S^n . The theorem now follows by taking $\Sigma^i = \tilde{\Sigma}_-^i$ for $2 \leq i \leq N$ and $\hat{\Phi}^i$ are the diffeomorphisms given by $(\tilde{\Phi}^i \circ \Psi^i)^{-1}$. \square

5 A sharpening of [3, Theorem 1.1]

In order to prove Theorem 1.2, we begin with an elementary lemma.

Lemma 5.1 *If $\mathbf{x}_1, \dots, \mathbf{x}_{m+1} \in \mathbb{R}^{n+1}$ is a sequence of points such that*

$$(5-1) \quad |\mathbf{x}_i - \mathbf{x}_{i+1}| \leq \hat{K}(1 + |\mathbf{x}_i|)^{-1}$$

for $1 \leq i \leq m$ and some $\hat{K} \geq 0$, then

$$(5-2) \quad |\mathbf{x}_1 - \mathbf{x}_{m+1}| \leq K(m)(1 + |\mathbf{x}_1|)^{-1},$$

where $K(m) = (\hat{K} + 1)^m - 1$.

Proof We proceed by induction on m . The lemma is obviously true when $m = 1$. Suppose (5-2) holds for $m = m'$. Using this induction hypothesis with (5-1) implies that

$$\begin{aligned} |\mathbf{x}_1 - \mathbf{x}_{m'+2}| &\leq |\mathbf{x}_1 - \mathbf{x}_{m'+1}| + |\mathbf{x}_{m'+1} - \mathbf{x}_{m'+2}| \\ &\leq K(m')(1 + |\mathbf{x}_1|)^{-1} + \widehat{K}(1 + |\mathbf{x}_{m'+1}|)^{-1}. \end{aligned}$$

Furthermore, by the induction hypothesis and triangle inequality,

$$|\mathbf{x}_1| \leq K(m')(1 + |\mathbf{x}_1|)^{-1} + |\mathbf{x}_{m'+1}|.$$

As $K(m') \geq 0$ and $(1 + |\mathbf{x}_1|)^{-1} \leq 1$, this implies that

$$1 + |\mathbf{x}_1| \leq 1 + K(m') + |\mathbf{x}_{m'+1}| \leq (1 + K(m'))(1 + |\mathbf{x}_{m'+1}|).$$

That is,

$$(1 + |\mathbf{x}_{m'+1}|)^{-1} \leq (1 + K(m'))(1 + |\mathbf{x}_1|)^{-1}.$$

Hence,

$$|\mathbf{x}_1 - \mathbf{x}_{m'+2}| \leq (K(m') + \widehat{K}(1 + K(m')))(1 + |\mathbf{x}_1|)^{-1},$$

and by the induction hypothesis, $K(m') = (\widehat{K} + 1)^{m'} - 1$, so setting

$$K(m' + 1) = K(m') + \widehat{K}(1 + K(m')) = (\widehat{K} + 1)^{m'+1} - 1$$

verifies that (5-2) holds for $m = m' + 1$ and finishes the proof. \square

In [3, Theorem 1.1] we showed that for any asymptotically conical self-shrinker with entropy less than or equal to that of a round cylinder, the link of the asymptotic cone separates the unit sphere into exactly two connected components, both diffeomorphic to the self-shrinker. We next observe that the proof of [3, Theorem 1.1] actually allows us to make the following more refined conclusion.

Proposition 5.2 *Fix $n \geq 2$. If $\Sigma \in \mathcal{ACS}_n[\lambda_{n-1}]$, then there is a homeomorphic involution $\phi : \mathbb{S}^n \rightarrow \mathbb{S}^n$ which fixes $\mathcal{L}(\Sigma)$, the link of the asymptotic cone $\mathcal{C}(\Sigma)$ of Σ , and swaps the two components of $\mathbb{S}^n \setminus \mathcal{L}(\Sigma)$.*

Proof By [3, Theorem 1.1], the link $\mathcal{L}(\Sigma)$ is connected and separates \mathbb{S}^n into two components Ω_+ and Ω_- . In particular, $\mathcal{L}(\Sigma) = \partial\bar{\Omega}_+ = \partial\bar{\Omega}_-$. In order to construct ϕ , it is enough to show the existence of a homeomorphism $\psi : \bar{\Omega}_+ \rightarrow \bar{\Omega}_-$ such that $\psi|_{\mathcal{L}(\Sigma)} : \mathcal{L}(\Sigma) \rightarrow \mathcal{L}(\Sigma)$ is the identity map. Indeed, if such a ψ exists, one defines ϕ by

$$\phi(p) = \begin{cases} \psi(p) & \text{if } p \in \bar{\Omega}_+, \\ \psi^{-1}(p) & \text{if } p \in \Omega_-. \end{cases}$$

To explain the construction of ψ , let us first summarize the main objects used in the proof of [3, Theorem 1.1]. First, recall that it is shown there that associated to Σ are two smooth mean curvature flows $\{\Gamma_t^\pm\}_{t \in [-1,0]}$ with Γ_{-1}^+ the normal exponential graph over Σ of a small positive multiple of the lowest eigenfunction of the self-shrinker stability operator of Σ (normalized to be positive) and Γ_{-1}^- to be a small negative multiple of this function. In particular, by choosing the multiple small enough, one can ensure both that Γ_{-1}^+ is the exponential normal graph of some function on Γ_{-1}^- and that Γ_{-1}^- is the exponential normal graph of some function on Γ_{-1}^+ . Furthermore, up to relabeling, each $\Gamma^\pm = \Gamma_0^\pm$ is diffeomorphic to Ω^\pm the components of $S^n \setminus \mathcal{L}(\Sigma)$. Moreover, these diffeomorphisms, which we denote by Π^\pm , are given by restricting, to Γ^\pm , the map

$$\Pi(p) = \frac{\mathbf{x}(p)}{|\mathbf{x}(p)|}.$$

We next use the flow $\{\Gamma_t^\pm\}_{t \in [-1,0]}$ to construct a natural diffeomorphism $\Psi: \Gamma^+ \rightarrow \Gamma^-$ which has the property that there is a constant $K > 0$ such that

$$(5-3) \quad |\mathbf{x}(p) - \mathbf{x}(\Psi(p))| \leq \frac{K}{1 + |\mathbf{x}(p)|}.$$

We do so iteratively. Specifically, by [3, Proposition 4.4(1) and Proposition 4.5] there is a constant $\tilde{C}_0 > 0$ such that

$$(5-4) \quad \sup_{t \in [-1,0]} \sup_{\Gamma_t^\pm} (|A_{\Gamma_t^\pm}| + |\nabla_{\Gamma_t^\pm} A_{\Gamma_t^\pm}|) < \tilde{C}_0.$$

This, together with [3, Proposition 4.4(3)], implies that there is a $\rho > 0$ such that $\mathcal{T}_\rho(\Gamma_t^\pm)$ is a regular tubular neighborhood of Γ_t^\pm for each $t \in [-1, 0]$. It follows from this and (5-4) that there is a $\delta > 0^4$ such that if $t_1, t_2 \in [-1, 0]$ and $|t_1 - t_2| < \delta$, then $\Gamma_{t_1}^\pm$ is a normal exponential graph over $\Gamma_{t_2}^\pm$ and vice versa. As such, for all $t_1, t_2 \in [-1, 0]$ with $|t_1 - t_2| < \delta$, there is a diffeomorphism

$$\Psi_{t_2, t_1}^\pm: \Gamma_{t_1}^\pm \rightarrow \Gamma_{t_2}^\pm$$

defined by nearest-point projection from $\Gamma_{t_1}^\pm$ to $\Gamma_{t_2}^\pm$. Pick $M \in \mathbb{N}$ so $M\delta > 1$, choose $0 = s_0 > s_1 > \dots > s_M = -1$ so that $|s_i - s_{i+1}| < \delta$, and define a diffeomorphism $\Psi^-: \Gamma_{-1}^- \rightarrow \Gamma^-$ by

$$\Psi^- = \Psi_{s_0, s_1}^- \circ \Psi_{s_1, s_2}^- \circ \dots \circ \Psi_{s_{M-1}, s_M}^-.$$

⁴By the mean curvature flow equation, [18, Lemma 3.3], and (5-4), the velocity of a point moving along the flow and the rate of change of the unit normal are bounded by $10\tilde{C}_0$. Thus one may choose $\delta = 10^{-3}\tilde{C}_0^{-1} \min\{\rho, 1\}$ for instance. In particular, this ensures that if $|t_1 - t_2| < \delta$, then $\Gamma_{t_1}^\pm$ is contained in $\mathcal{T}_\rho(\Gamma_{t_2}^\pm)$ and vice versa.

Likewise, define a diffeomorphism $\Psi^+ : \Gamma^+ \rightarrow \Gamma_{-1}^+$ by

$$\Psi^+ = \Psi_{s_M, s_{M-1}}^+ \circ \Psi_{s_{M-1}, s_{M-2}}^+ \circ \dots \circ \Psi_{s_1, s_0}^+,$$

and let $\Psi^{+,-} : \Gamma_{-1}^+ \rightarrow \Gamma_{-1}^-$ be given by nearest-point projection. By construction, this is also a diffeomorphism and so the map

$$\Psi = \Psi^- \circ \Psi^{+,-} \circ \Psi^+$$

is a diffeomorphism $\Psi : \Gamma^+ \rightarrow \Gamma^-$.

By construction, if $t_1, t_2 \in [-1, 0]$ and $|t_1 - t_2| < \delta$, then for all $p \in \Gamma_{t_1}^\pm$,

$$(5-5) \quad |\mathbf{x}(p) - \mathbf{x}(\Psi_{t_2, t_1}^\pm(p))| < \rho.$$

Furthermore, [3, Proposition 4.4(1)] implies that for $t \in [-1, 0]$ each Γ_t^\pm is smoothly asymptotic to $\mathcal{C}(\Sigma)$. In particular, there is a $R > 0$ and functions u_t^\pm on $\mathcal{C}(\Sigma) \setminus B_R$ whose normal exponential graph over $\mathcal{C}(\Sigma)$ sits inside of Γ_t^\pm and contains $\Gamma_t^\pm \setminus B_{2R}$. Moreover, by [3, Proposition 4.2(2)] and [3, Lemma 4.3] there is a constant $K' > 0$ such that for $p \in \mathcal{C}(\Sigma) \setminus B_R$,

$$|u_t^\pm(p)| \leq K'(1 + |\mathbf{x}(p)|)^{-1}.$$

Hence, for any $t_1, t_2 \in [-1, 0]$, if $p \in \Gamma_{t_1}^\pm \setminus B_{2R}$, then there is a point $p' \in \mathcal{C}(\Sigma) \setminus B_R$ such that

$$(5-6) \quad |\mathbf{x}(p) - \mathbf{x}(p')| \leq K'(1 + |\mathbf{x}(p')|)^{-1}$$

and also a point $p'' \in \Gamma_{t_2}^\pm$ such that

$$(5-7) \quad |\mathbf{x}(p') - \mathbf{x}(p'')| \leq K'(1 + |\mathbf{x}(p')|)^{-1}.$$

Hence if $|t_1 - t_2| < \delta$, then as Ψ_{t_2, t_1}^\pm is given by nearest-point projection,

$$\begin{aligned} |\mathbf{x}(p) - \mathbf{x}(\Psi_{t_2, t_1}^\pm(p))| &\leq |\mathbf{x}(p) - \mathbf{x}(p'')| \\ &\leq |\mathbf{x}(p) - \mathbf{x}(p')| + |\mathbf{x}(p') - \mathbf{x}(p'')| \\ &\leq 2K'(1 + |\mathbf{x}(p')|)^{-1}. \end{aligned}$$

As $K' > 0$ and $1 + |\mathbf{x}(p')| \geq 1$, (5-6) implies that

$$(1 + |\mathbf{x}(p')|)^{-1} \leq (1 + K')(1 + |\mathbf{x}(p)|)^{-1},$$

and so

$$|\mathbf{x}(p) - \mathbf{x}(\Psi_{t_2, t_1}^\pm(p))| \leq 2K'(1 + K')(1 + |\mathbf{x}(p)|)^{-1}.$$

Combining this with (5-5) one obtains that for all $p \in \Gamma_{t_1}^\pm$,

$$|\mathbf{x}(p) - \mathbf{x}(\Psi_{t_2, t_1}^\pm(p))| \leq \widehat{K}(1 + |\mathbf{x}(p)|)^{-1},$$

where $\widehat{K} = 2K'(1 + K') + \rho(1 + 2R)$. By the same arguments, for all $p \in \Gamma_{-1}^+$,

$$|\mathbf{x}(p) - \mathbf{x}(\Psi^{+, -}(p))| \leq \widehat{K}(1 + |\mathbf{x}(p)|)^{-1}.$$

Hence, it follows from Lemma 5.1, that

$$|\mathbf{x}(p) - \mathbf{x}(\Psi(p))| \leq K(1 + |\mathbf{x}(p)|)^{-1},$$

where $K = (1 + \widehat{K})^{2M+2} - 1$.

To complete the proof, set

$$\psi(p) = \begin{cases} \Pi^-(\Psi((\Pi^+)^{-1}(p))) & \text{if } p \in \Omega_+, \\ p & \text{if } p \in \partial\Omega_+. \end{cases}$$

We claim that ψ is a homeomorphism. First, note that, by [3, Proposition 4.4(3)], there is an $R > 1$ and $\widetilde{C}_1 > 1$ such that if $p \in \Gamma^\pm \setminus B_R$, then

$$\widetilde{C}_1^{-1} |\mathbf{x}(p)|^{2\mu} < \text{dist}_{\mathbb{R}^{n+1}}(p, \mathcal{L}(\Sigma)) < \widetilde{C}_1 |\mathbf{x}(p)|^{-1},$$

where $\mu < -1$. Hence,

$$(5-8) \quad C^{-1} |\mathbf{x}(p)|^{2\mu-1} < \text{dist}_{\mathbb{S}^n}(\Pi^\pm(p), \mathcal{L}(\Sigma)) < C |\mathbf{x}(p)|^{-2},$$

where $C \geq \widetilde{C}_1$. Hence, for $q \in \Omega^+$ with $\text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))$ sufficiently small, if we set $q' = (\Pi^+)^{-1}(q) \in \Gamma^+$, then

$$|\mathbf{x}(q')| \geq C^{\frac{1}{2\mu-1}} \text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))^{\frac{1}{2\mu-1}}.$$

By (5-3),

$$\begin{aligned} \left| |\mathbf{x}(\Psi(q'))| - |\mathbf{x}(q')| \right| &\leq |\mathbf{x}(\Psi(q')) - \mathbf{x}(q')| \\ &\leq KC^{-\frac{1}{2\mu-1}} \text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))^{-\frac{1}{2\mu-1}}. \end{aligned}$$

Hence, for $\text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))$ sufficiently small,

$$\text{dist}_{\mathbb{S}^n}(q, \psi(q)) \leq 4KC^{-\frac{1}{2\mu-1}} \text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))^{-\frac{1}{2\mu-1}} |\mathbf{x}(q')|^{-1}.$$

Using (5-8), again gives

$$\text{dist}_{\mathbb{S}^n}(q, \psi(q)) \leq 4KC^{-\frac{2}{2\mu-1}} \text{dist}_{\mathbb{S}^n}(q, \mathcal{L}(\Sigma))^{-\frac{2}{2\mu-1}}.$$

As $\mu < -1$, for any $q_0 \in \mathcal{L}(\Sigma)$, the right-hand side goes to 0 as $q \rightarrow q_0$. By the triangle inequality,

$$\text{dist}_{\mathbb{S}^n}(q_0, \psi(q)) \leq \text{dist}_{\mathbb{S}^n}(q, \psi(q)) + \text{dist}_{\mathbb{S}^n}(q, q_0),$$

and so the right-hand side goes to 0 as $q \rightarrow q_0$. Hence, ψ is continuous. Finally, as $\bar{\Omega}_+$ is compact and $\bar{\Omega}_-$ is Hausdorff, ψ is a closed map and hence, as ψ is a bijection, it is a homeomorphism. \square

Theorem 1.2 is a standard topological consequence of Proposition 5.2.

Proof of Theorem 1.2 Recall that [3, Theorem 1.1] states that $\mathcal{L}(\Sigma)$ separates \mathbb{S}^n into exactly two components, which we denote by U^\pm , both diffeomorphic to Σ . As a consequence, $\mathcal{L}(\Sigma)$ is connected. Let $\phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ be the homeomorphism given by Proposition 5.2 so $\phi(U^-) = U^+$. Pick a regular tubular neighborhood $T \subset \mathbb{S}^n$ of $\mathcal{L}(\Sigma)$. We let $V^\pm = U^\pm \cup T$ and observe that \bar{U}^\pm , the closure of U^\pm , is a retract of V^\pm and that $\mathcal{L}(\Sigma)$ is a retraction of $T = V^- \cap V^+$.

As \bar{U}^\pm is a retraction of V^\pm and $\mathcal{L}(\Sigma)$ is a retraction of T , the natural inclusion maps induce isomorphisms between the reduced homology groups $\tilde{H}_k(\bar{U}^\pm)$ and $\tilde{H}_k(V^\pm)$ and between $\tilde{H}_k(\mathcal{L}(\Sigma))$ and $\tilde{H}_k(T)$. As such, there is a natural map $\Phi: \tilde{H}_k(V^-) \rightarrow \tilde{H}_k(V^+)$ defined by the diagram

$$\begin{array}{ccccc}
 & & \tilde{H}_k(T) & \xrightarrow{j_*^-} & \tilde{H}_k(V^-) \\
 & \swarrow \cong & & \searrow j_*^+ & \downarrow \Phi \\
 \tilde{H}_k(\mathcal{L}(\Sigma)) & \xrightarrow{i_*^-} & \tilde{H}_k(\bar{U}^-) & & \tilde{H}_k(V^+) \\
 & \searrow i_*^+ & \downarrow \phi_* & \swarrow \cong & \\
 & & \tilde{H}_k(\bar{U}^+) & &
 \end{array}$$

where $i^\pm: \mathcal{L}(\Sigma) \rightarrow \bar{U}^\pm$ and $j^\pm: T \rightarrow V^\pm$ denote the natural inclusion maps, and we used that $\phi \circ i^- = i^+$. As ϕ is a homeomorphism, both ϕ_* and Φ are isomorphisms. This implies that the map

$$J = (j_*^-, -j_*^+): \tilde{H}_k(T) \rightarrow \tilde{H}_k(V^-) \oplus \tilde{H}_k(V^+)$$

is surjective if and only if $\tilde{H}_k(V^-) = \tilde{H}_k(V^+) = \{0\}$. Indeed, if the map is surjective, then for any element $\alpha \in \tilde{H}_k(V^-)$ there is an element $\beta \in \tilde{H}_k(T)$ such that $J(\beta) = (\alpha, 0)$. That is, $j_*^-(\beta) = \alpha$ and $j_*^+(\beta) = 0$. Hence, $0 = j_*^+(\beta) = \Phi(j_*^-(\beta)) = \Phi(\alpha)$. In other words, as Φ is an isomorphism, $\alpha \in \ker(\Phi) = \{0\}$ and so $\tilde{H}_k(V^-) = \{0\}$. The proof that $\tilde{H}_k(V^+) = \{0\}$ is the same. The converse is immediate.

We next recall several standard facts about the reduced homology of manifolds and of manifolds with boundary. First, as $\mathcal{L}(\Sigma)$ is a connected, oriented $(n-1)$ -dimensional manifold, $\tilde{H}_k(\mathcal{L}(\Sigma)) = \tilde{H}_k(T) = \{0\}$ for $k = 0$ and $k \geq n$, and $\tilde{H}_{n-1}(\mathcal{L}(\Sigma)) = \tilde{H}_{n-1}(T) = \mathbb{Z}$. Likewise, as the \bar{U}^\pm are connected, oriented n -manifolds with boundary, $\tilde{H}_k(\bar{U}^\pm) = \tilde{H}_k(V^\pm) = 0$ for $k = 0$ and $k \geq n$.

In order to compute the remaining reduced homology groups, we use the Mayer–Vietoris long exact sequence for the reduced homology of (V^-, V^+, \mathbb{S}^n) . This gives the following exact sequences for $k \geq 0$:

$$(5-9) \quad \tilde{H}_{k+1}(\mathbb{S}^n) \rightarrow \tilde{H}_k(T) \xrightarrow{J} \tilde{H}_k(V^-) \oplus \tilde{H}_k(V^+) \rightarrow \tilde{H}_k(\mathbb{S}^n).$$

As $\tilde{H}_k(\mathbb{S}^n) = \mathbb{Z}$ for $k = n$ and is otherwise $\{0\}$, (5-9) implies that J is surjective for $0 \leq k \leq n - 1$. Hence $\tilde{H}_k(\bar{U}^\pm) = \tilde{H}_k(V^\pm) = \{0\}$ for these k . As such, (5-9) further implies that $\tilde{H}_k(\mathcal{L}(\Sigma)) = \tilde{H}_k(T) = \{0\}$ for $0 \leq k \leq n - 2$, completing the verification that $\mathcal{L}(\Sigma)$ is a homology $(n-1)$ -sphere.

To conclude the proof, it is enough, by the Hurewicz theorem and the Whitehead theorem, to show that $\pi_1(U^\pm) = \pi_1(\bar{U}^\pm) = \{1\}$. To that end, first observe that the maps $F^\pm: \mathbb{S}^n \rightarrow \bar{U}^\pm$ defined by

$$F^\pm(p) = \begin{cases} p & \text{if } p \in \bar{U}^\pm, \\ \phi(p) & \text{if } p \in U^\mp \end{cases}$$

are continuous. Now suppose γ is a closed loop in \bar{U}^\pm . As $\pi_1(\mathbb{S}^n) = \{1\}$, there is a homotopy $H: \mathbb{S}^1 \times [0, 1] \rightarrow \mathbb{S}^n$ taking γ to a point. Clearly, $F^\pm \circ H: \mathbb{S}^1 \times [0, 1] \rightarrow \bar{U}^\pm$ is also a homotopy taking γ to a point. That is, $\pi_1(\bar{U}^\pm) = \{1\}$. □

Proof of Corollary 1.4 By Theorem 1.2, $\mathcal{L}(\Sigma)$ is a homology 2-sphere. By the classification of surfaces this means that $\mathcal{L}(\Sigma)$ is diffeomorphic to \mathbb{S}^2 and so Alexander’s theorem [1] implies that both components of $\mathbb{S}^3 \setminus \mathcal{L}(\Sigma)$ are diffeomorphic to \mathbb{R}^3 . This together with [3, Theorem 1.1] proves the claim. □

6 Surgery procedure

We prove Theorem 1.1 using Corollary 1.4 and Theorem 4.5.

Proof of Theorem 1.1 We first observe that (\star_{3,λ_2}) holds by [25, Theorem B] and that $(\star\star_{3,\lambda_2})$ holds by [3, Corollary 1.2]. If Σ is (after a translation and dilation) a self-shrinker, then by [8, Theorem 0.7], Σ is diffeomorphic to \mathbb{S}^3 , proving the theorem. Otherwise, flow Σ for a small amount of time by the mean curvature flow (using short time existence of for smooth closed initial hypersurfaces) to obtain a hypersurface Σ' diffeomorphic to Σ and, by Huisken’s monotonicity formula, with $\lambda[\Sigma'] < \lambda[\Sigma]$. On the one hand, if the level set flow of Σ' is nonfattening, then we set $\Sigma_0 = \Sigma'$. On the other hand, if the level set flow of Σ' is fattening, then we can take Σ_0 to be a small normal graph over Σ' so that $\lambda[\Sigma_0] < \lambda[\Sigma]$, and because the nonfattening condition is generic, the level set flow of Σ_0 is nonfattening.

Hence, the hypotheses of Section 4 hold, and we may apply Theorem 4.5 unconditionally to obtain a family of hypersurfaces $\Sigma^1, \dots, \Sigma^N$ in \mathbb{R}^4 . As Σ^N is diffeomorphic to \mathbb{S}^3 , if $N = 1$, then there is nothing more to show and so we may assume that $N > 1$. We will now show that Σ^{N-1} is diffeomorphic to Σ^N and hence to \mathbb{S}^3 .

Let $V = \bigcup_{j=1}^{m(N-1)} V_j^{N-1}$ and $\widehat{\Sigma}^N = \Sigma^N \setminus V$, and let $U = \bigcup_{j=1}^{m(N-1)} U_j^{N-1}$ and $\widehat{\Sigma}^{N-1} = \Sigma^{N-1} \setminus U$, so $\widehat{\Phi}^{N-1}: \widehat{\Sigma}^N \rightarrow \widehat{\Sigma}^{N-1}$ is the orientation-preserving diffeomorphism given by Theorem 4.5. By Corollary 1.4 and [3, Theorem 1.1], each component of \bar{U} is diffeomorphic to a closed 3-ball \bar{B}^3 . Hence, each component of $\partial\widehat{\Sigma}^{N-1}$ and $\partial\widehat{\Sigma}^N$ is diffeomorphic to \mathbb{S}^2 . That is, ∂V_j^{N-1} is diffeomorphic to \mathbb{S}^2 for $1 \leq j \leq m(N-1)$, and so, as Σ^N is diffeomorphic to the 3-sphere, Alexander’s theorem [1] implies that each \bar{V}_j^{N-1} is diffeomorphic to the closed 3-ball. Hence, there are orientation-preserving diffeomorphisms $\Psi_j^{N-1}: \bar{V}_j^{N-1} \rightarrow \bar{U}_j^{N-1}$.

Denote by

$$\widehat{\phi}_j^{N-1}: \partial V_j^{N-1} \rightarrow \partial U_j^{N-1} \quad \text{and} \quad \psi_j^{N-1}: \partial V_j^{N-1} \rightarrow \partial U_j^{N-1}$$

the diffeomorphisms given by restricting $\widehat{\Phi}^{N-1}$ and Ψ_j^{N-1} , respectively. Observe, that the orientation of $\widehat{\Sigma}^N$ and the orientation on \bar{V} induce opposite orientations on $\partial\bar{V}$. Likewise, the orientation of $\widehat{\Sigma}^{N-1}$ and that of \bar{U} induce opposite orientations on $\partial\bar{U}$. By construction, the $\widehat{\phi}_j^{N-1}$ preserve the orientations induced from $\widehat{\Sigma}^N$ and $\widehat{\Sigma}^{N-1}$. Hence, as the orientations induced by \bar{V}_j^{N-1} and \bar{U}_j^{N-1} are opposite to those induced by $\widehat{\Sigma}^N$ and $\widehat{\Sigma}^{N-1}$, the $\widehat{\phi}_j^{N-1}$ also preserve these orientations. The same is true of the ψ_j^{N-1} . As such,

$$\xi_j^{N-1} = (\psi_j^{N-1})^{-1} \circ \widehat{\phi}_j^{N-1} \in \text{Diff}_+(\partial V_j^{N-1}),$$

where $\text{Diff}_+(M)$ is the space of orientation-preserving self-diffeomorphisms of an oriented manifold M (here we may use the orientation on ∂V_j^{N-1} induced by either \bar{V} or $\widehat{\Sigma}^N$). By [26]—see also [30] and [5]—the space $\text{Diff}_+(\mathbb{S}^2)$ is path-connected and so any element of $\text{Diff}_+(\mathbb{S}^2)$ extends to an element of $\text{Diff}_+(\bar{B}^3)$. That is, there are diffeomorphism $\Xi_j^{N-1} \in \text{Diff}_+(\bar{V}_j^{N-1})$ that restrict to ξ_j^{N-1} on ∂V_j^{N-1} . Thus, the maps $\widehat{\Psi}_j^{N-1} = \Psi_j^{N-1} \circ \Xi_j^{N-1}: \bar{V}_j^{N-1} \rightarrow \bar{U}_j^{N-1}$ are diffeomorphisms that agree with $\widehat{\Phi}^{N-1}$ on the common boundary.

Define $\Phi^{N-1}: \Sigma^N \rightarrow \Sigma^{N-1}$ by

$$\Phi^{N-1}(p) = \begin{cases} \widehat{\Phi}^{N-1}(p) & \text{if } p \in \widehat{\Sigma}^N, \\ \widehat{\Psi}_j^{N-1}(p) & \text{if } p \in V_j^{N-1}. \end{cases}$$

By construction, this map is a homeomorphism. However, it is a standard procedure to construct a diffeomorphism between Σ^N and Σ^{N-1} by smoothing this map out (see

for instance [14, Theorem 8.1.9]). Hence, Σ^{N-1} is diffeomorphic to \mathbb{S}^3 and iterating this argument shows that $\Sigma = \Sigma^1$ is diffeomorphic to \mathbb{S}^3 as claimed. \square

Theorem 1.5 follows from Theorem 1.2, Theorem 4.5 and the Mayer–Vietoris long exact sequence for reduced homology. For completeness, we include a proof of the following standard topological fact which we will need to use.

Lemma 6.1 *Let M be a closed n -dimensional manifold and $\Sigma \subset M$ a closed hypersurface. If M is a homology n -sphere and Σ is a homology $(n-1)$ -sphere, then each component of $M \setminus \Sigma$ is a homology n -ball.*

Proof Our hypotheses ensure that both M and Σ are connected and oriented. Hence, Σ is two-sided and there is an open $U^+ \subset M$ such that $\Sigma = \partial U^+$. Let $U^- = M \setminus \bar{U}^+$. To prove the lemma, we will need to compute the Mayer–Vietoris long exact sequence for $(\bar{U}^-, \bar{U}^+, M)$. Strictly speaking, we should “thicken” \bar{U}^+ and \bar{U}^- up with a regular tubular neighborhood of $\Sigma = \partial \bar{U}^\pm$ as in the proof of Theorem 1.2, but we leave the details of this to the reader.

The Mayer–Vietoris long exact sequence and the fact that M is a homology n -sphere and Σ is a homology $(n-1)$ -sphere gives the sequences:

$$\begin{array}{ccccccc}
 \tilde{H}_{k+1}(M) & \xrightarrow{\partial} & \tilde{H}_k(\Sigma) & \longrightarrow & \tilde{H}_k(\bar{U}^-) \oplus \tilde{H}_k(\bar{U}^+) & \longrightarrow & \tilde{H}_k(M) \\
 \uparrow = & & \uparrow = & & \uparrow = & & \uparrow = \\
 \tilde{H}_{k+1}(\mathbb{S}^n) & \xrightarrow{\partial} & \tilde{H}_k(\mathbb{S}^{n-1}) & \longrightarrow & \tilde{H}_k(\bar{U}^-) \oplus \tilde{H}_k(\bar{U}^+) & \longrightarrow & \tilde{H}_k(\mathbb{S}^n)
 \end{array}$$

For $0 \leq k \leq n-2$ and $k \geq n+1$, this immediately gives that $\tilde{H}_k(\bar{U}_\pm) = \{0\}$. When $k = n-1$, the map ∂ is necessarily generated by $[M] \mapsto [\Sigma]$ where $[M]$ is the fundamental class of M and $[\Sigma]$ is the fundamental class of Σ . In particular, this map is an isomorphism and so we conclude that $\tilde{H}_{n-1}(\bar{U}^\pm) = \{0\}$. For the same reason, $\tilde{H}_n(\bar{U}^\pm) = \{0\}$, which verifies the claim. \square

Proof of Theorem 1.5 Arguing as in the first paragraph of the proof of Theorem 1.1, we obtain $\Sigma^1, \dots, \Sigma^N$ the hypersurfaces given by Theorem 4.5. As Σ^N is diffeomorphic to \mathbb{S}^n , it is a homology n -sphere. In particular, if $N = 1$, then there is nothing further to show. As such, we may assume that $N > 1$.

Let us show that Σ^{N-1} is a homology n -sphere. First, set $V = \bigcup_{j=1}^{m(N-1)} V_j^{N-1}$ and $\hat{\Sigma}^N = \Sigma^N \setminus V$, and let $U = \bigcup_{j=1}^{m(N-1)} U_j^{N-1}$ and $\hat{\Sigma}^{N-1} = \Sigma^{N-1} \setminus U$. Next observe that, as $\partial U_j^{N-1} = \mathcal{L}(\Gamma_j^{N-1})$ for some $\Gamma_j^{N-1} \in \mathcal{ACS}_n^*(\Lambda)$, Theorem 1.2 implies that each component of $\partial \hat{\Sigma}^{N-1}$ is a homology $(n-1)$ -sphere. Hence, as $\partial U = \partial \hat{\Sigma}^{N-1}$ is

diffeomorphic to $\partial\widehat{\Sigma}^N = \partial V$, we see that each component of $\partial V = \partial\widehat{\Sigma}^N$ is a homology $(n-1)$ -sphere and so Lemma 6.1 implies that each component of \bar{V} is a homology n -ball.

We may now use the Mayer–Vietoris long exact sequence to show $\tilde{H}_k(\widehat{\Sigma}^N) = \{0\}$ for $k \neq n-1$ and $\tilde{H}_{n-1}(\widehat{\Sigma}^N) = \mathbb{Z}^{m(N-1)-1}$. To see this, consider the Mayer–Vietoris long exact sequence of $(\bar{V}, \widehat{\Sigma}^N, \Sigma^N)$. This long exact sequence and the fact that \bar{V} is the union of homology n -balls gives, for $k > 0$, the exact sequences:

$$\begin{array}{ccccccc} \tilde{H}_{k+1}(\Sigma^N) & \xrightarrow{\partial} & \tilde{H}_k(\partial V) & \longrightarrow & \tilde{H}_k(\bar{V}) \oplus \tilde{H}_k(\widehat{\Sigma}^N) & \longrightarrow & \tilde{H}_k(\Sigma^N) \\ \uparrow = & & \uparrow = & & \uparrow = & & \uparrow = \\ \tilde{H}_{k+1}(\mathbb{S}^n) & \xrightarrow{\partial} & \bigoplus_{j=1}^{m(N-1)} \tilde{H}_k(\mathbb{S}^{n-1}) & \longrightarrow & \tilde{H}_k(\widehat{\Sigma}^N) & \longrightarrow & \tilde{H}_k(\mathbb{S}^n) \end{array}$$

Hence $\tilde{H}_k(\widehat{\Sigma}^N) = \{0\}$ for $1 \leq k \leq n-2$ and $k \geq n+1$. When $k = n-1$, the map ∂ is generated by $[\Sigma^N] \mapsto ([\partial V_1^{N-1}], \dots, [\partial V_{m(N-1)}^{N-1}])$ where $[\Sigma^N]$ is the fundamental class of Σ^N and $[\partial V_j^{N-1}]$ is the fundamental class of ∂V_j^{N-1} . It follows that $\tilde{H}_{n-1}(\widehat{\Sigma}^N) = \mathbb{Z}^{m(N-1)-1}$ and, as this map is injective, that $\tilde{H}_n(\widehat{\Sigma}^N) = \{0\}$. Finally, as $\widehat{\Sigma}^N$ is connected, $\tilde{H}_0(\widehat{\Sigma}^N) = \{0\}$, which completes the computation.

By Theorem 4.5, $\widehat{\Sigma}^N$ is diffeomorphic to $\widehat{\Sigma}^{N-1}$, and so $\tilde{H}_k(\widehat{\Sigma}^{N-1}) = 0$ for $k \neq n-1$ and $\tilde{H}_{n-1}(\widehat{\Sigma}^{N-1}) = \mathbb{Z}^{m(N-1)-1}$. Furthermore, Theorem 1.2 implies that each component of \bar{U} is contractible. Hence, applying the Mayer–Vietoris long exact sequence to $(\widehat{\Sigma}^{N-1}, \bar{U}, \Sigma^{N-1})$ gives the following diagram for $k > 0$:

$$\begin{array}{ccccccc} \tilde{H}_k(\partial\bar{U}) & \longrightarrow & \tilde{H}_k(\bar{U}) \oplus \tilde{H}_k(\widehat{\Sigma}^{N-1}) & \longrightarrow & \tilde{H}_k(\Sigma^{N-1}) & \longrightarrow & \tilde{H}_{k-1}(\partial\bar{U}) \\ \uparrow = & & \uparrow = & & \uparrow = & & \uparrow = \\ \bigoplus_{j=1}^{m(N-1)} \tilde{H}_k(\mathbb{S}^{n-1}) & \longrightarrow & \tilde{H}_k(\widehat{\Sigma}^{N-1}) & \longrightarrow & \tilde{H}_k(\Sigma^{N-1}) & \longrightarrow & \bigoplus_{j=1}^{m(N-1)} \tilde{H}_{k-1}(\mathbb{S}^{n-1}) \end{array}$$

In particular, for $1 \leq k \leq n-2$ and $k \geq n+1$, we obtain that $\tilde{H}_k(\Sigma^{N-1}) = \{0\}$. The Mayer–Vietoris long exact sequence further gives the exact sequences:

$$\begin{array}{ccccccc} \tilde{H}_{n-1}(\partial\bar{U}) & \xrightarrow{\delta} & \tilde{H}_{n-1}(\bar{U}) \oplus \tilde{H}_{n-1}(\widehat{\Sigma}^{N-1}) & \longrightarrow & \tilde{H}_{n-1}(\Sigma^{N-1}) & \longrightarrow & \tilde{H}_{n-2}(\partial\bar{U}) \\ \uparrow = & & \uparrow = & & \uparrow = & & \uparrow = \\ \mathbb{Z}^{m(N-1)} & \xrightarrow{\delta} & \mathbb{Z}^{m(N-1)-1} & \longrightarrow & \tilde{H}_{n-1}(\Sigma^{N-1}) & \longrightarrow & \{0\} \end{array}$$

Here δ is given by $(l_1, \dots, l_{m(N-1)}) \mapsto (l_1 - l_{m(N-1)}, \dots, l_{m(N-1)-1} - l_{m(N-1)})$. As δ is surjective, it follows that $\tilde{H}_{n-1}(\Sigma^{N-1}) = \{0\}$. Finally, as Σ^{N-1} is an oriented, connected n -dimensional manifold $\tilde{H}_n(\Sigma^{N-1}) = \mathbb{Z}$ and $\tilde{H}_0(\Sigma^{N-1}) = \{0\}$. Hence, Σ^{N-1} is a homology n -sphere.

As our argument only used that Σ^N was a homology n -sphere, we may repeat it to see that each of the Σ^i is a homology n -sphere and so conclude that Σ is one as well. \square

Appendix

Fix an open subset $U \subset \mathbb{R}^{n+1}$. A hypersurface Σ in U is a proper, codimension-one submanifold of U . A smooth mean curvature flow S in U is a collection of hypersurfaces $\{\Sigma_t\}_{t \in I}$ in U , for I an interval, such that

- (1) for all $t_0 \in I$ and $p_0 \in \Sigma_{t_0}$, there is a $r_0 = r_0(p_0, t_0)$ and an interval $I_0 = I_0(p_0, t_0)$ with $(p_0, t_0) \in B_{r_0}^{n+1}(p_0) \times I_0 \subset U \times I$;
- (2) there is a smooth map $F: B_1^n \times I_0 \rightarrow \mathbb{R}^{n+1}$ such that $F_t(p) = F(p, t): B_1^n \rightarrow \mathbb{R}^{n+1}$ is a parametrization of $B_{r_0}^{n+1}(p_0) \cap \Sigma_t$; and
- (3) $(\frac{\partial}{\partial t} F(p, t))^\perp = \mathbf{H}_{\Sigma_t}(F(p, t))$.

It is convenient to consider the spacetime track of S (also denoted by S):

$$(A-1) \quad S = \{(x(p), t) \in \mathbb{R}^{n+1} \times \mathbb{R} : p \in \Sigma_t\} \subset U \times I.$$

This is a smooth submanifold of spacetime and is transverse to each constant time hyperplane $\mathbb{R}^{n+1} \times \{t_0\}$. Along the spacetime track S , let d/dt be the smooth vector field

$$(A-2) \quad \frac{d}{dt} \Big|_{(p,t)} = \frac{\partial}{\partial t} + \mathbf{H}_{\Sigma_t}(p).$$

It is not hard to see that this vector field is tangent to S and the position vector satisfies

$$(A-3) \quad \frac{d}{dt} \mathbf{x}(p, t) = \mathbf{H}_{\Sigma_t}(p).$$

It is a standard fact that if each Σ_t in S is closed, ie is compact and without boundary, then there is a smooth map

$$F: M \times I \rightarrow \mathbb{R}^{n+1}$$

such that each $F_t = F(\cdot, t): M \rightarrow \mathbb{R}^{n+1}$ is a parametrization of Σ_t a closed n -dimensional manifold M . As a consequence, each Σ_t is diffeomorphic to M .

We will need the following generalization of this last fact to manifolds with boundary.

Proposition A.1 Fix $R \in (0, \infty]$ and let $\{\bar{B}_{2r_1}(\mathbf{x}_1), \dots, \bar{B}_{2r_m}(\mathbf{x}_m)\}$ be a collection of pairwise disjoint balls in $B_R \subset \mathbb{R}^{n+1}$ and let $U = B_{2R} \setminus \bigcup_{i=1}^m \bar{B}_{r_i}(\mathbf{x}_i)$. If $\{\Sigma_t\}_{t \in (-\tau, \tau)}$ is a smooth mean curvature flow in U with the property that

- (1) each $\hat{\Sigma}_t = \Sigma_t \cap (\bar{B}_R \setminus \bigcup_{i=1}^m B_{2r_i}(\mathbf{x}_i))$ is compact,
- (2) each $\partial B_{2r_i}(\mathbf{x}_i)$, for $1 \leq i \leq m$, intersects Σ_t transversally and nontrivially for all $t \in (-\tau, \tau)$,
- (3) if $R < \infty$, then ∂B_R intersects Σ_t transversally and nontrivially for all $t \in (-\tau, \tau)$,

then $\hat{\Sigma}_{t_1}$ and $\hat{\Sigma}_{t_2}$ are diffeomorphic as compact manifolds with boundary for any $t_1, t_2 \in (-\tau, \tau)$.

Proof For simplicity, we consider only $R = \infty$, $m = 1$, $\mathbf{x}_1 = \mathbf{0}$ and $r_1 = \frac{1}{2}$. It is straightforward to extend the argument to the general case. Let S be the spacetime track of the flow, so S is a smooth hypersurface in $(\mathbb{R}^{n+1} \setminus \bar{B}_{1/2}) \times (-\tau, \tau)$. As each Σ_t intersects ∂B_1 transversally, it is clear that S meets $\partial B_1 \times (-\tau, \tau)$ transversally. In particular, $\tilde{S} = S \setminus (B_1 \times (-\tau, \tau))$ is a smooth hypersurface with boundary. Let $\tilde{B} = \partial \tilde{S} = \{(p, t) : p \in \partial B_1 \cap \Sigma_t, t \in (-\tau, \tau)\}$.

Without loss of generality we may assume that the given t_1, t_2 satisfy $t_1 < t_2$. Let $\hat{S} = \tilde{S} \cap (\mathbb{R}^{n+1} \times [t_1, t_2])$ and $\hat{B} = \tilde{B} \cap (\mathbb{R}^{n+1} \times [t_1, t_2])$. Observe that \hat{S} is a compact manifold with corners and \hat{B} is one of its boundary strata. The other two boundary strata are $\hat{\Sigma}_{t_1} \times \{t_1\}$ and $\hat{\Sigma}_{t_2} \times \{t_2\}$.

As ∂B_1 meets each Σ_t transversally and \hat{B} is compact, there is an $\epsilon > 0$ such that $|\mathbf{x}^\top(p, t)| \geq 2\epsilon$ for $(p, t) \in \hat{B}$, where \mathbf{x}^\top is the tangential component of the position vector. By continuity there is a $\frac{1}{2} > \delta > 0$ such that, for any $t \in [t_1, t_2]$ and $p \in (\bar{B}_{1+\delta} \setminus B_{1-\delta}) \cap \Sigma_t$, we have $|\mathbf{x}^\top(p, t)| \geq \epsilon$. Now let $\eta \in C_0^\infty(\mathbb{R}^{n+1})$ be a smooth function with $0 \leq \eta \leq 1$, $\eta = 1$ on ∂B_1 and $\text{spt}(\eta) \subset \bar{B}_{1+\delta} \setminus B_{1-\delta}$. For $(p, t) \in \hat{S}$, consider the vector

$$V(p, t) = -\eta(\mathbf{x}(p, t)) \frac{(\mathbf{x}(p, t) \cdot \mathbf{H}_{\Sigma_t}(p))}{|\mathbf{x}^\top(p, t)|^2} \mathbf{x}^\top(p, t),$$

and observe this gives a smooth vector field on S that restricts to a smooth compactly supported vector field on each Σ_t . Let $W = d/dt + V$ which is a smooth vector field on S .

We claim that W is tangent to \hat{B} and transverse to $\hat{\Sigma}_{t_1} \times \{t_1\} \cup \hat{\Sigma}_{t_2} \times \{t_2\}$. As V is tangent to $\Sigma_t \times \{t\}$, the transversality of W follows from the transversality of d/dt . This transversality follows immediately from the definition of d/dt . To see the

tangency, note that by construction, $\widehat{B} = \{(p, t) \in \widehat{S} : |\mathbf{x}(p, t)|^2 = 1\}$. For $(p, t) \in \widehat{B}$, one computes

$$\begin{aligned} \partial_{\mathbf{W}} |\mathbf{x}(p, t)|^2 &= 2\mathbf{x}(p, t) \cdot \nabla_{\mathbf{W}} \mathbf{x}(p, t) \\ &= 2\mathbf{x}(p, t) \cdot \mathbf{H}_{\Sigma_t}(p) - 2\eta(\mathbf{x}(p, t)) \frac{(\mathbf{x}(p, t) \cdot \mathbf{H}_{\Sigma_t}(p))}{|\mathbf{x}^\top(p, t)|^2} \mathbf{x}(p, t) \cdot \mathbf{x}^\top(p, t) \\ &= 0, \end{aligned}$$

where the last equality used that $(p, t) \in \widehat{B}$ so $\eta(\mathbf{x}(p, t)) = 1$. This verifies the claim.

To conclude the proof, observe that as \widehat{S} is compact and \mathbf{W} is tangent to \widehat{B} and transverse to $\widehat{\Sigma}_{t_1} \times \{t_1\} \cup \widehat{\Sigma}_{t_2} \times \{t_2\}$, standard ODE theory gives that for any $P_0 = (p_0, t_0) \in \widehat{S}$ the initial value problem

$$\begin{cases} \dot{\gamma}_{P_0}(s) = \mathbf{W}(\gamma_{P_0}(s)), \\ \gamma_{P_0}(0) = P_0 \end{cases}$$

has a unique smooth solution $\gamma_{P_0}: [t_1 - t_0, t_2 - t_0] \rightarrow \widehat{S}$ which depends smoothly on P_0 . These solutions satisfy $t(\gamma_{P_0}(s)) = s + t_0$ and so there is a diffeomorphism $\phi: \Sigma_{t_1} \rightarrow \Sigma_{t_2}$ given by $(\phi(p), t_2) = \gamma_{(p, t_1)}(t_2 - t_1)$. \square

References

- [1] **J W Alexander**, *On the subdivision of 3-space by a polyhedron*, Nat. Acad. Proc. 10 (1924) 6–8
- [2] **J Bernstein, L Wang**, *A sharp lower bound for the entropy of closed hypersurfaces up to dimension six*, Invent. Math. 206 (2016) 601–627 MR
- [3] **J Bernstein, L Wang**, *A topological property of asymptotically conical self-shrinkers of small entropy*, Duke Math. J. 166 (2017) 403–435 MR
- [4] **K A Brakke**, *The motion of a surface by its mean curvature*, Mathematical Notes 20, Princeton Univ. Press (1978) MR
- [5] **J Cerf**, *Sur les difféomorphismes de la sphère de dimension trois ($\Gamma_4 = 0$)*, Lecture Notes in Mathematics 53, Springer (1968) MR
- [6] **Y G Chen, Y Giga, S Goto**, *Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations*, J. Differential Geom. 33 (1991) 749–786 MR
- [7] **X Cheng, D Zhou**, *Volume estimate about shrinkers*, Proc. Amer. Math. Soc. 141 (2013) 687–696 MR
- [8] **T H Colding, T Ilmanen, W P Minicozzi II, B White**, *The round sphere minimizes entropy among closed self-shrinkers*, J. Differential Geom. 95 (2013) 53–69 MR

- [9] **T H Colding, W P Minicozzi II**, *Generic mean curvature flow, I: Generic singularities*, Ann. of Math. 175 (2012) 755–833 MR
- [10] **L C Evans, J Spruck**, *Motion of level sets by mean curvature, I*, J. Differential Geom. 33 (1991) 635–681 MR
- [11] **L C Evans, J Spruck**, *Motion of level sets by mean curvature, II*, Trans. Amer. Math. Soc. 330 (1992) 321–332 MR
- [12] **L C Evans, J Spruck**, *Motion of level sets by mean curvature, III*, J. Geom. Anal. 2 (1992) 121–150 MR
- [13] **L C Evans, J Spruck**, *Motion of level sets by mean curvature, IV*, J. Geom. Anal. 5 (1995) 77–114 MR
- [14] **M W Hirsch**, *Differential topology*, Graduate Texts in Mathematics 33, Springer (1976) MR
- [15] **W-Y Hsiang**, *Minimal cones and the spherical Bernstein problem, I*, Ann. of Math. 118 (1983) 61–73 MR
- [16] **W-Y Hsiang**, *Minimal cones and the spherical Bernstein problem, II*, Invent. Math. 74 (1983) 351–369 MR
- [17] **W-Y Hsiang, I Sterling**, *Minimal cones and the spherical Bernstein problem, III*, Invent. Math. 85 (1986) 223–247 MR
- [18] **G Huisken**, *Flow by mean curvature of convex surfaces into spheres*, J. Differential Geom. 20 (1984) 237–266 MR
- [19] **G Huisken**, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom. 31 (1990) 285–299 MR
- [20] **T Ilmanen**, *Elliptic regularization and partial regularity for motion by mean curvature*, Mem. Amer. Math. Soc. 520, Amer. Math. Soc., Providence, RI (1994) MR
- [21] **T Ilmanen**, *Singularities of mean curvature flow of surfaces*, preprint (1995) Available at <https://people.math.ethz.ch/~ilmanen/papers/sing.ps>
- [22] **T Ilmanen, A Neves, F Schulze**, *On short time existence for the planar network flow*, preprint (2014) arXiv
- [23] **T Ilmanen, B White**, *Sharp lower bounds on density for area-minimizing cones*, Camb. J. Math. 3 (2015) 1–18 MR
- [24] **D Ketover, X Zhou**, *Entropy of closed surfaces and min-max theory*, preprint (2015) arXiv
- [25] **F C Marques, A Neves**, *Min-max theory and the Willmore conjecture*, Ann. of Math. 179 (2014) 683–782 MR
- [26] **J Munkres**, *Differentiable isotopies on the 2-sphere*, Michigan Math. J. 7 (1960) 193–197 MR

- [27] **S Osher, J A Sethian**, *Fronts propagating with curvature-dependent speed: algorithms based on Hamilton–Jacobi formulations*, J. Comput. Phys. 79 (1988) 12–49 MR
- [28] **F Schulze**, *Uniqueness of compact tangent flows in mean curvature flow*, J. Reine Angew. Math. 690 (2014) 163–172 MR
- [29] **L Simon**, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University 3, Aust. Nat. Univ., Centre for Math. Anal., Canberra (1983) MR
- [30] **S Smale**, *Diffeomorphisms of the 2–sphere*, Proc. Amer. Math. Soc. 10 (1959) 621–626 MR
- [31] **A Stone**, *A density function and the structure of singularities of the mean curvature flow*, Calc. Var. Partial Differential Equations 2 (1994) 443–480 MR
- [32] **L Wang**, *Uniqueness of self-similar shrinkers with asymptotically conical ends*, J. Amer. Math. Soc. 27 (2014) 613–638 MR
- [33] **B White**, *A local regularity theorem for mean curvature flow*, Ann. of Math. 161 (2005) 1487–1519 MR
- [34] **J J Zhu**, *On the entropy of closed hypersurfaces and singular self-shrinkers*, preprint (2016) arXiv

*Department Of Mathematics, Johns Hopkins University
Baltimore, MD, United States*

*Department of Mathematics, University of Wisconsin
Madison, WI, United States*

bernstein@math.jhu.edu, luwang@math.wisc.edu

Proposed: Bruce Kleiner

Seconded: Tobias H. Colding, John Lott

Received: 28 June 2016

Revised: 3 February 2017

