

Primes and fields in stable motivic homotopy theory

JEREMIAH HELLER

KYLE M ORMSBY

Let F be a field of characteristic different from 2. We establish surjectivity of Balmer’s comparison map

$$\rho^\bullet: \mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c) \rightarrow \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$$

from the tensor triangular spectrum of the homotopy category of compact motivic spectra to the homogeneous Zariski spectrum of Milnor–Witt K -theory. We also comment on the tensor triangular geometry of compact cellular motivic spectra, producing in particular novel field spectra in this category. We conclude with a list of questions about the structure of the tensor triangular spectrum of the stable motivic homotopy category.

14F42; 19D45, 55P42, 18E30

1 Introduction

The chromatic approach to stable homotopy theory is a powerful organizational and computational tool. Underlying this approach is a good understanding of the tensor triangular geometry of the stable homotopy category, as provided by the celebrated nilpotence and periodicity theorems of Hopkins and Smith [16]. Recent computational work of Andrews [1] suggests a rich and vast chromatic picture in the motivic setting. However, a good (or any) understanding of the tensor triangular geometry of the stable motivic homotopy category is still lacking.

To set the stage for our results, recall that given a (small) tensor triangulated category \mathcal{H} , the set underlying Balmer’s spectrum $\mathrm{Spc}(\mathcal{H})$ is the set of tensor triangular prime ideals; see Balmer [5]. This simple invariant is surprisingly powerful; for example, under mild hypotheses it completely determines the thick tensor ideals of \mathcal{H} [5, Theorem 4.10]. Computing the spectrum of \mathcal{H} is no easy task, but Balmer’s comparison maps [7, Theorem 5.3, Corollary 5.6] make it possible to organize $\mathrm{Spc}(\mathcal{H})$ fiberwise over the classical spectra of (homogenous) prime ideals of the (graded) endomorphism ring of the unit.

This plays out as follows in the motivic setting over a field F whose characteristic is not 2. The tensor triangulated categories of interest here are the homotopy category of compact motivic \mathbb{P}^1 -spectra $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ and the homotopy category of compact cellular motivic \mathbb{P}^1 -spectra $\mathcal{C}(F)$. We write $\mathcal{H}(F)$ for either one of these two categories. Of course the graded endomorphism ring (relative to $\mathbb{A}^1 \setminus 0$) is the same in either case, and by a theorem of Morel [27], this ring is the Milnor–Witt K -theory ring $K_*^{\mathrm{MW}}(F)$ of F . In degree zero, $K_0^{\mathrm{MW}}(F) = \mathrm{GW}(F)$, the Grothendieck–Witt ring of quadratic forms. Balmer’s comparison maps thus take the form

$$\begin{array}{ccc} \mathrm{Spc}(\mathcal{H}(F)) & \xrightarrow{\rho^\bullet} & \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F)) \\ & \searrow \rho & \downarrow \\ & & \mathrm{Spec}(\mathrm{GW}(F)) \end{array}$$

Moreover, Balmer [7, Theorem 7.13] applies to show that ρ is surjective. Complete knowledge of $\mathrm{Spec}(\mathrm{GW}(F))$ can be extracted from classical work of Lorenz and Leicht [24]; see also [7, Remark 10.2]. Thus ρ provides a very explicit lower bound for $\mathrm{Spc}(\mathcal{H}(F))$.

Recent work of Thornton [32] shows that $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$ has precisely $|X_F| + 1$ more elements than $\mathrm{Spec}(\mathrm{GW}(F))$, where X_F denotes the set of orderings on the field F . Thus ρ^\bullet could potentially detect up to $|X_F| + 1$ more primes of $\mathcal{H}(F)$ than ρ can. Our main theorem is that ρ^\bullet is surjective and thus ρ^\bullet does yield a full $(|X_F| + 1)$ -fold improvement on the previously known lower bounds for $\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c)$. This meets the challenge posed in [7, Remark 10.5].

Theorem 1.1 *Suppose F is a field whose characteristic is not 2. Then the maps*

$$\begin{aligned} \rho^\bullet: \mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c) &\rightarrow \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F)), \\ \rho^\bullet: \mathrm{Spc}(\mathcal{C}(F)) &\rightarrow \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F)), \end{aligned}$$

are both surjective.

Remark 1.2 This theorem tells us that we have a commutative diagram as follows, in which both vertical arrows are surjective:

$$\begin{array}{ccc} \mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c) & \longrightarrow & \mathrm{Spc}(\mathcal{C}(F)) \\ \rho^\bullet \downarrow & & \downarrow \rho^\bullet \\ \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F)) & \longleftarrow & \mathrm{Spec}^h(K_*^{\mathrm{MW}}(F)) \end{array}$$

Balmer has informed us of a general criterion [9, Theorem 1.3] for when a tensor triangulated functor $f: \mathcal{K} \rightarrow \mathcal{L}$ between essentially small tensor triangulated categories induces a surjection $\mathrm{Spc}(f): \mathrm{Spc}(\mathcal{L}) \rightarrow \mathrm{Spc}(\mathcal{K})$; namely, if f detects \otimes -nilpotence and \mathcal{K} is rigid, then $\mathrm{Spc}(f)$ is surjective. Since this condition is clearly met by $f: \mathcal{C}(F) \subset \mathrm{SH}^{\mathbb{A}^1}(F)^c$, the top horizontal arrow in the above diagram is surjective as well.

Our argument for Theorem 1.1 is indirect, and does not produce any explicit prime ideals in the stable motivic homotopy categories. After establishing our main theorem we address the question of producing explicit primes for $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ and $\mathcal{C}(F)$ by analyzing realization functors (Section 5) and field spectra for $\mathcal{C}(F)$ (Section 4). Ultimately, when F is isomorphic to a subfield of \mathbb{C} we find infinite towers of tensor primes above all but one positive-characteristic prime in $K_*^{\mathrm{MW}}(F)$; see Figure 4.

Both Joachimi [21, Chapter 7] and Kelly [22] have studied the tensor triangular geometry of $\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c)$. Joachimi's study is primarily concerned with Betti realization functors, and should be compared with our results in Section 5. Kelly proves that ρ^\bullet is surjective when F is a finite field; we adapt his argument to prove Proposition 3.1.

Outline We recall the basic background of tensor triangular geometry, stable motivic homotopy theory, and Milnor–Witt K -theory in Section 2. In particular we recall there Thornton's theorem on the structure of $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$. In Section 3 we establish our main result, Theorem 1.1, on the surjectivity of ρ^\bullet . In Section 4, we find explicit field spectra for $\mathcal{C}(F)$, whose acyclics are tensor triangular primes living over particular elements of $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$; see Theorem 4.10. In Section 5, we completely determine the effect of ρ^\bullet on the injective image of $\mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$ and $\mathrm{Spc}(\mathrm{SH}(C_2)^c)$ under the maps induced by Betti realization; see Propositions 5.3 and 5.5. In Section 6, we devote our attention to field spectra for $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ and elucidate the manner in which singular and realization functors interact with tensor triangular primes. We conclude with a list of questions in Section 7, amounting to an optimist's sketch of how to determine the complete structure of $\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c)$. (Caveat emptor: some of these questions are likely quite difficult.)

Notation We use the following notation throughout:

- F is a field, always of characteristic $\neq 2$.
- Sm/F is the category of smooth finite-type schemes over F .

- $\mathrm{SH}^{\mathbb{A}^1}(F)$, $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ and $\mathcal{C}(F)$ are, respectively, the stable motivic homotopy category over $\mathrm{Spec} F$, the full subcategory of compact motivic spectra, and the full subcategory of compact cellular motivic spectra.
- $\mathcal{K}(F)$ refers to either of $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ or $\mathcal{C}(F)$.
- $K_*^{\mathrm{MW}}(F)$ is the Milnor–Witt K –theory ring of F .
- S^0 is the motivic sphere spectrum over $\mathrm{Spec} F$.
- S^m and $S^{m\alpha}$ for $m \in \mathbb{Z}$ are the “simplicial” and the “geometric” m –sphere, respectively: S^m is the m –fold suspension of S^0 , and $S^{m\alpha} = (\mathbb{A}^1 \setminus 0)^{\wedge m}$.
- $[X, Y]_{\mathcal{D}}$ is the set of maps between objects X, Y of a category \mathcal{D} .
- $\underline{\pi}_m(X)_n$, where $X \in \mathrm{SH}^{\mathbb{A}^1}(F)$ and $m, n \in \mathbb{Z}$, is the Nisnevich sheaf on Sm/F associated with the assignment

$$U \mapsto [S^m \wedge U_+, S^{n\alpha} \wedge X]_{\mathrm{SH}^{\mathbb{A}^1}(F)}.$$

Note that other sources may write $\underline{\pi}_{m-n\alpha}(X)$ or $\underline{\pi}_{m-n, -n}(X)$ for this sheaf. We have chosen our notation to match Morel’s in [26; 27].

- $\pi_m(X)_n$, where $X \in \mathrm{SH}^{\mathbb{A}^1}(F)$ and $m, n \in \mathbb{Z}$, is the set of sections of $\underline{\pi}_m(X)_n$ on $\mathrm{Spec} F$.
- $\mathrm{Spec}(R)$ is the Zariski spectrum of prime ideals in a commutative ring R .
- $\mathrm{Spec}^h(R_*)$ is the Zariski spectrum of homogeneous prime ideals in an ε –commutative \mathbb{Z} –graded ring R_* (which means that $\varepsilon \in R_0$, $\varepsilon^2 = 1$, and $ab = \varepsilon^{mn}ba$ for $a \in R_m, b \in R_n$).
- $\mathrm{Spc}(\mathcal{C})$, for \mathcal{C} a tensor triangulated category, is Balmer’s tensor triangular spectrum of \mathcal{C} .

Acknowledgements It is our pleasure to thank Shane Kelly and Paul Balmer for helpful discussions and for sharing with us drafts of [22] and [9], respectively. We also thank the referee for helpful comments and improvements to the exposition. Ormsby was partially supported by NSF award DMS-1406327.

2 Preliminaries

In this section we recall background notions and material from tensor triangular geometry and stable motivic homotopy theory used in the rest of the paper.

2.1 Tensor triangular geometry

We begin by recalling a few definitions and facts about the spectrum of a tensor triangulated category. We refer the reader to [7; 8] for full details.

A *tensor triangulated category* $(\mathcal{K}, \otimes, 1)$ is a triangulated category \mathcal{K} together with a symmetric monoidal product \otimes which is bi-exact and a unit $1 \in \mathcal{K}$. For \mathcal{K} essentially small, Balmer associates its *tensor triangular spectrum* $\mathrm{Spc}(\mathcal{K})$, which is a topological space defined as follows.

A *tensor triangular prime ideal* (or simply a prime ideal) of \mathcal{K} is a proper thick subcategory $\mathcal{P} \subsetneq \mathcal{K}$ such that:

- \mathcal{P} is a *tensor ideal*, ie if $a \in \mathcal{K}$ and $b \in \mathcal{P}$, then $a \otimes b \in \mathcal{P}$.
- \mathcal{P} is *prime*, ie if for some $a, b \in \mathcal{K}$ we have $a \otimes b \in \mathcal{P}$, then a or b is in \mathcal{P} .

The prime spectrum of \mathcal{K} is the set of primes

$$\mathrm{Spc}(\mathcal{K}) := \{\mathcal{P} \subsetneq \mathcal{K} \mid \mathcal{P} \text{ is prime}\}.$$

Given an object $a \in \mathcal{K}$, its support is

$$\mathrm{supp}(a) := \{\mathcal{P} \in \mathrm{Spc}(\mathcal{K}) \mid a \notin \mathcal{P}\}.$$

Write $U(a) = \mathrm{Spc}(\mathcal{K}) \setminus \mathrm{supp}(a)$. Equip $\mathrm{Spc}(\mathcal{K})$ with the topology defined by letting the sets $U(a)$ for $a \in \mathcal{K}$ be a basis for the open sets. In particular, the support $\mathrm{supp}(a)$ of an object is always closed.

The prime spectrum is contravariantly functorial in \mathcal{K} . That is, given a tensor exact functor $\phi: \mathcal{K} \rightarrow \mathcal{L}$ there is an induced continuous map $\mathrm{Spc}(\phi): \mathrm{Spc}(\mathcal{L}) \rightarrow \mathrm{Spc}(\mathcal{K})$ defined by $\mathcal{P} \mapsto \phi^{-1}(\mathcal{P})$.

Of central interest in this paper are the continuous maps constructed by Balmer [7, Theorem 5.3, Corollary 5.6] which relate the spectrum of a tensor triangulated category to the spectrum of graded and ungraded endomorphism rings. Let $u \in \mathrm{Pic}(\mathcal{K})$ be a \otimes -invertible object and let $R_{\mathcal{K}}^*$ be the graded endomorphism ring of the unit: that is, $R_{\mathcal{K}}^i = [1, u^{\otimes i}]_{\mathcal{K}}$ for $i \in \mathbb{Z}$. We write $R_{\mathcal{K}} = R_{\mathcal{K}}^0$ for ungraded ring of ordinary endomorphisms of the unit. Write $\mathrm{Spec}^h(R_{\mathcal{K}}^*)$ for the Zariski spectrum of homogeneous

prime ideals in this graded ring. We have continuous maps

$$\begin{array}{ccc}
 \mathrm{Spc}(\mathcal{H}) & \xrightarrow{\rho^\bullet} & \mathrm{Spec}^h(R_{\mathcal{H}}^*) \\
 & \searrow \rho & \downarrow (\)^0 \\
 & & \mathrm{Spec}(R_{\mathcal{H}})
 \end{array}$$

where ρ^\bullet is defined by

$$\rho^\bullet(\mathcal{P}) := \{f \in R_{\mathcal{H}}^* \mid \mathcal{P} \in \mathrm{supp}(\mathrm{cone}(f))\}$$

and $(\)^0$ is defined by sending a homogenous prime \mathfrak{p} to the prime ideal $\mathfrak{p}^0 = \mathfrak{p} \cap R_{\mathcal{H}}^0$ of homogeneous degree zero elements. Of course, the map ρ^\bullet depends on the choice of invertible element u ; in general, we will omit u from our notation, but will write ρ_u^\bullet when specification is necessary.

2.2 Stable motivic homotopy theory and Milnor–Witt K –theory

We fix throughout a base field F , which will always be assumed to be of characteristic different from two. The tensor triangulated categories of interest in this paper stem from the homotopy category $\mathrm{SH}^{\mathbb{A}^1}(F)$ of motivic spectra over F , developed in [33], [20] and [26]. This category itself is too large for tensor triangular geometry and, as is standard in this situation, we focus on the homotopy category $\mathrm{SH}^{\mathbb{A}^1}(F)^c$ of compact motivic spectra.¹ In this case, the symmetric monoidal product is the smash product \wedge , and the unit object is the sphere spectrum S^0 .

There are many invertible objects in $\mathrm{SH}^{\mathbb{A}^1}(F)^c$, two of which are singled out. The first is the ordinary circle S^1 and the second is the geometric (or Tate) circle $S^\alpha := \mathbb{A}^1 \setminus 0$. We write $S^{m+n\alpha} = (S^1)^{\wedge m} \wedge (S^\alpha)^{\wedge n}$. The homotopy category of *compact cellular spectra* $\mathcal{C}(F) \subseteq \mathrm{SH}^{\mathbb{A}^1}(F)^c$ is defined to be the thick subcategory generated by the set $\{S^{m+n\alpha} \mid n, m \in \mathbb{Z}\}$. We write $\mathcal{H}(F)$ for either of these categories.

Milnor–Witt K –theory $K_*^{\mathrm{MW}}(F)$ plays a distinguished role in stable motivic homotopy theory because it is isomorphic to a graded ring of endomorphisms of the unit object in $\mathcal{H}(F)$ (at least when $\mathrm{char} F \neq 2$). Specifically, Morel [27] proves that

$$K_*^{\mathrm{MW}}(F) \cong \pi_0(S^0)_* \cong [S^0, S^{*\alpha}]_{\mathcal{H}(F)}.$$

¹Recall that an object X in a triangulated category \mathcal{H} is *compact* if $[X, \bigoplus_I Y_i]_{\mathcal{H}} \cong \bigoplus_I [X, Y_i]_{\mathcal{H}}$ for any set I .

Milnor–Witt K –theory is defined to be a quotient of the free associative \mathbb{Z} –graded ring on the set of symbols $[F^\times] := \{[u] \mid u \in F^\times\}$ in degree 1 and η in degree -1 ; the quotient is by the homogeneous ideal enforcing the following relations:

- **Twisted logarithm** $[uv] = [u] + [v] + \eta[u][v]$.
- **Steinberg relation** $[u][v] = 0$ for $u + v = 1$.
- **Commutativity** $[u]\eta = \eta[u]$.
- **Witt relation** $(2 + [-1]\eta)\eta = 0$.

Milnor–Witt K –theory is ε –commutative for $\varepsilon = -(1 + [-1]\eta)$. Moreover, it is related as follows to several important classical invariants:

- $K_0^{\text{MW}}(F) \cong \text{GW}(F)$, the Grothendieck–Witt ring of quadratic forms.
- $\eta^{-1}K_*^{\text{MW}}(F) \cong W(F)[\eta, \eta^{-1}]$, the ring of Laurent polynomials over the Witt ring.
- $K_*^{\text{MW}}(F)/\eta = K_*^M(F)$, the Milnor K –theory of F .

As observed in [7, Corollary 10.1], the map $\rho: \text{Spc}(\mathcal{K}(F)) \rightarrow \text{Spec}(\text{GW}(F))$ is surjective. (This follows from a general criterion [7, Theorem 7.13] for surjectivity of ρ .) Our results concern the map $\rho^\bullet: \text{Spc}(\mathcal{K}(F)) \rightarrow \text{Spec}^h(K_*^{\text{MW}}(F))$ and rely crucially on the following theorem of Thornton. Recall that X_F is the set of orderings on F equipped with the Harrison topology.²

Theorem 2.1 [32, Theorem 3.12] *Each $\mathfrak{p} \in \text{Spec}^h(K_*^{\text{MW}}(F))$ is of exactly one of the following forms, where p ranges over rational primes, α ranges over orderings of F , and $P_\alpha \subseteq F^\times$ denotes the positive cone of α :*

- (1) $([F^\times], \eta)$,
- (2) $([F^\times], \eta, p)$,
- (3) $([F^\times], 2)$,
- (4) $([P_\alpha], h)$,
- (5) $([P_\alpha], \eta, 2)$, and
- (6) $([P_\alpha], h, p)$ for $p \neq 2$.

Additionally, the topology on $\text{Spec}^h(K_*^{\text{MW}}(F))$ is specified by the following facts:

²A subbasis for the Harrison topology is given by the sets $H(a) = \{\alpha \in X_F \mid a >_\alpha 0\}$ for $a \in F^\times$. The Harrison space X_F is a Stone space; in particular, X_F is discrete if and only if $|X_F| < \infty$.

- (a) The subspace of minimal prime ideals is homeomorphic to $X_F^* = X_F \sqcup \{\infty\}$, where X_F has the Harrison topology and ∞ is an isolated point.
- (b) The Hasse diagram (ie inclusion poset) of $\text{Spec}^h(K_*^{\text{MW}}(F))$ is shown in Figure 1 and the closure operator takes a subset $S \subseteq \text{Spec}^h(K_*^{\text{MW}}(F))$ to everything in or above S in the Hasse diagram.




Figure 1: The Hasse diagram of $\text{Spec}^h(K_*^{\text{MW}}(F))$. Inclusion of tensor triangular primes and closure both go upwards in the diagram. In the picture, we suppose that $\alpha, \beta \in X_F$, and the central ellipses represent the rest of X_F .

3 Surjectivity

Let $\mathcal{H}(F)$ denote either $\text{SH}^{\mathbb{A}^1}(F)^c$ or $\mathcal{C}(F)$. In this section we prove our main theorem, Theorem 1.1: the continuous map $\rho^\bullet: \text{Spc}(\mathcal{H}(F)) \rightarrow \text{Spec}^h(K_*^{\text{MW}}(F))$ hits all the primes enumerated in Theorem 2.1. We warm up by showing that this is true when F is nonreal, ie when -1 is a sum of squares in F .

Proposition 3.1 *Suppose F is a nonreal field whose characteristic is not 2. Then*

$$\rho^\bullet: \text{Spc}(\mathcal{H}(F)) \rightarrow \text{Spec}^h(K_*^{\text{MW}}(F))$$

is surjective.

Proof Our proof is an adaptation of Kelly’s argument for the surjectivity of ρ^\bullet in the case when F is a finite field [22]. We first show that $\rho^\bullet(U(\mathbb{P}^2)) = D_\eta = \{([F^\times], 2)\}$, where $U(\mathbb{P}^2) \subseteq \text{Spc}(\mathcal{H}(F))$ is the complement of $\text{supp}(\mathbb{P}^2)$ and D_η is notation for the basic open set of homogeneous prime ideals not containing η . Indeed, since $\Sigma\mathbb{P}^2 \simeq C\eta$,

the cone of η , we have that the inclusion $U(\mathbb{P}^2) \cong \mathrm{Spc}(\mathcal{K}(F)/\langle \mathbb{P}^2 \rangle) \rightarrow \mathrm{Spc}(\mathcal{K}(F))$ factors through $\mathrm{Spc}(\eta^{-1}\mathcal{K}(F))$. Recall from [28, Corollary 3.11] that $\eta^{-1}K_*^{\mathrm{MW}}(F) \cong W(F)[\eta, \eta^{-1}]$, the ring of Laurent polynomials in a variable η over the Witt ring $W(F)$. We therefore have that

$$(\)^0: \mathrm{Spec}^h(\eta^{-1}K_*^{\mathrm{MW}}(F)) \cong \mathrm{Spec}(W(F)) \cong \mathrm{Spec}(\mathbb{F}_2),$$

where the final identification holds via [24] because F has no orderings. Since the nongraded analogue ρ of ρ^\bullet is surjective (by [7, Theorem 7.13] since $\eta^{-1}\mathcal{K}(F)$ is connective), we conclude that $\rho^\bullet: \mathrm{Spc}(\eta^{-1}\mathcal{K}(F)) \rightarrow \mathrm{Spec}^h(\eta^{-1}K_*^{\mathrm{MW}}(F))$ is surjective. The image of $\mathrm{Spec}^h(\eta^{-1}K_*^{\mathrm{MW}}(F))$ in $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$ is precisely D_η . We conclude that $\rho^\bullet(U(\mathbb{P}^2)) = D_\eta$.

We now check that $\rho^\bullet(\mathrm{supp}(\mathbb{P}^2))$ is precisely the (closed) complement $V_\eta = \{\mathfrak{p} \mid \eta \in \mathfrak{p}\}$ of D_η . We have that $\rho^\bullet(\mathrm{supp}(\mathbb{P}^2)) \subseteq V_\eta$ by definition of ρ^\bullet and supp . Note that by [7, Remark 10.2], $\mathrm{Spec} \mathbb{Z} \cong \mathrm{Spec} \mathrm{GW}(F)$ via the assignment $p\mathbb{Z} \mapsto \dim^{-1}p\mathbb{Z}$ for $\dim: \mathrm{GW}(F) \rightarrow \mathbb{Z}$ the dimension homomorphism. Meanwhile,

$$V_\eta = \{([F^\times], p, \eta) \mid p \text{ prime or } 0\},$$

and it is elementary to observe that $([F^\times], p, \eta)^0 = \dim^{-1}p\mathbb{Z}$. Thus we get that $(\)^0: V_\eta \cong \mathrm{Spec} \mathrm{GW}(F) \cong \mathrm{Spec} \mathbb{Z}$, and the composite $\rho: \mathrm{Spc}(\mathcal{K}(F)) \rightarrow \mathrm{Spec} \mathbb{Z}$ is surjective by [7, Theorem 7.13]. It follows that $\rho^\bullet(\mathrm{supp}(\mathbb{P}^2)) \subseteq V_\eta$ consists of at least every prime in V_η except possibly $([F^\times], 2, \eta)$. Thus it remains to show that $([F^\times], 2, \eta) \in \rho^\bullet(\mathrm{supp}(\mathbb{P}^2))$. For this it suffices to show that the closure of $U(\mathbb{P}^2)$ intersects $\mathrm{supp}(\mathbb{P}^2)$ nontrivially. Indeed, by continuity, $\rho^\bullet(\overline{U(\mathbb{P}^2)} \cap \mathrm{supp}(\mathbb{P}^2))$ is contained in $\rho^\bullet(\overline{U(\mathbb{P}^2)}) \cap V_\eta = \{([F^\times], 2, \eta)\}$.

Suppose for the sake of contradiction that $\mathrm{Spc}(\mathcal{K}(F))$ is separated by $\overline{U(\mathbb{P}^2)}$ and $\mathrm{supp}(\mathbb{P}^2)$. Let p denote the exponential characteristic of F and consider the functor $i: \mathcal{K}(F) \rightarrow \mathcal{K}(F)[1/p]$ and the induced continuous map

$$\phi := \mathrm{Spc}(i): \mathrm{Spc}(\mathcal{K}(F)[1/p]) \rightarrow \mathrm{Spc}(\mathcal{K}(F)).$$

Now we have that

$$\phi^{-1}(\mathrm{supp}(\mathbb{P}^2)) = \mathrm{supp}(\mathbb{P}^2[1/p]), \quad \phi^{-1}(U(\mathbb{P}^2)) = U(\mathbb{P}^2[1/p]).$$

It follows that $\mathrm{Spc}(\mathcal{K}(F)[1/p])$ is separated by $\overline{U(\mathbb{P}^2[1/p])}$ and $\mathrm{supp}(\mathbb{P}^2[1/p])$. By [23, Corollary B.2] every smooth F -scheme is dualizable in $\mathrm{SH}^{\mathbb{A}^1}(F)[1/p]$ and so by [18, Theorem 2.1.3], every object in $\mathcal{K}(F)[1/p]$ is dualizable. We may thus apply [6,

Theorem 2.11] to conclude that the unit of $\mathcal{K}(F)[1/p]$ decomposes as a nontrivial sum of two objects. It follows that $\text{GW}(F)[1/p]$, the endomorphism ring of the unit in $\mathcal{K}(F)$, decomposes as the product of two nontrivial rings. Since $\text{Spec}(\text{GW}(F)[1/p]) \cong \text{Spec}(\mathbb{Z}[1/p])$ has a unique minimal element, $\text{GW}(F)[1/p]$ is irreducible, and we have reached a contradiction. We conclude that

$$\overline{U(\mathbb{P}^2)} \cap \text{supp}(\mathbb{P}^2) \neq \emptyset,$$

and hence $\rho^\bullet(\text{supp}(\mathbb{P}^2)) = V_\eta$. It follows that ρ^\bullet is surjective. □

We can now prove our main theorem.

Proof of Theorem 1.1 The proof proceeds in four steps.

Step 1 We first show that it suffices to check surjectivity of ρ^\bullet when F is algebraically or real closed. For an arbitrary F , let $i: F \hookrightarrow F_\alpha$ be a real or algebraic closure of F . Consider the induced functor $i^*: \mathcal{K}(F) \rightarrow \mathcal{K}(F_\alpha)$, which is tensor triangulated such that $i^*((\mathbb{A}^1 \setminus 0)_F) = (\mathbb{A}^1 \setminus 0)_{F_\alpha}$. Hence we get a commutative square:

$$\begin{CD} \text{Spc}(\mathcal{K}(F_\alpha)) @>\text{Spc}(i^*)>> \text{Spc}(\mathcal{K}(F)) \\ @V\rho^\bullet VV @VV\rho^\bullet V \\ \text{Spec}^h(K_*^{\text{MW}}(F_\alpha)) @>>> \text{Spec}^h(K_*^{\text{MW}}(F)) \end{CD}$$

By Theorem 2.1, the sum of the bottom horizontal maps over all $\alpha \in X_F^* = X_F \sqcup \{\infty\}$, where $F_\infty = \bar{F}$ and F_α is the real closure of F corresponding to $\alpha \in X_F$, is surjective. Thus if ρ^\bullet is surjective for all algebraically or real closed fields, we may conclude that ρ^\bullet is surjective in general.

Step 2 We now check that ρ^\bullet is surjective when F is algebraically closed. This follows from Proposition 3.1 because algebraically closed fields are nonreal.

Step 3 In this step, we will compute $[-1]^{-1}K_*^{\text{MW}}(F)$ for F real closed. We use this computation in Step 4 as part of our proof that ρ^\bullet is surjective when F is real closed.

For the moment, take F to be any field of characteristic different from 2. Via the twisted logarithm relation, in $K_*^{\text{MW}}(F)$ we have

$$0 = [1] = 2[-1] + [-1]^2\eta.$$

Hence $h = 2 + [-1]\eta = 0$ in $[-1]^{-1}K_*^{\text{MW}}(F)$. It follows that

$$[-1]^{-1}K_*^{\text{MW}}(F) = [-1]^{-1}(K_*^{\text{MW}}(F)/h).$$

The ring $K_*^W(F) := K_*^{\text{MW}}(F)/h$ also goes by the name Witt K -theory. By [26, Theorem 6.4.7], we have $K_*^W(F) \cong W(F)[I\eta^{-1}, \eta]$, the extended Rees ring of the Witt ring of F with respect to the fundamental ideal I . This is a graded ring in which $W(F)[I\eta^{-1}, \eta]_n = I^n \cdot \{\eta^{-n}\}$, for I^n the n^{th} power of the fundamental ideal in $W(F)$ (where $I^n = W(F)$ for $n \leq 0$). The image of $[-1]$ in $W[I\eta^{-1}, \eta]$ is $2 \cdot \eta^{-1}$.

Now suppose that F is real closed, in which case $W(F) \cong \mathbb{Z}$ via the signature homomorphism and

$$K_*^W(F) \cong \mathbb{Z}[\eta, [-1]]/(2 + [-1]\eta).$$

Hence

$$[-1]^{-1}K_*^{\text{MW}}(F) \cong [-1]^{-1}K_*^W(F) \cong \mathbb{Z}[[[-1], [-1]^{-1}],$$

ie Laurent polynomials in the variable $[-1]$ over \mathbb{Z} .

Step 4 We now show that ρ^\bullet is surjective when F is real closed, completing our proof. Write

$$\text{Spec}^h(K_*^{\text{MW}}(F)) = V_{[-1]} \cup D_{[-1]},$$

where $V_{[-1]}$ is the subspace of primes of the form (1), (2) or (3), and $D_{[-1]}$ is its complement. Note that $V_{[-1]}$ is the closed set of primes containing $[-1]$, and $D_{[-1]}$ is the basic open of primes not containing $[-1]$. Base change to \bar{F} and Step 2 imply that $V_{[-1]}$ is hit by ρ^\bullet . To show that $D_{[-1]}$ is hit by ρ^\bullet , consider the following diagram:

$$\begin{array}{ccccc} \text{Spc}(\mathcal{H}(F)/\langle C[-1] \rangle) & \longrightarrow & \text{Spc}([-1]^{-1}\mathcal{H}(F)) & \longrightarrow & \text{Spc}(\mathcal{H}(F)) \\ \downarrow & & \downarrow & & \downarrow \\ D_{[-1]} & \longrightarrow & \text{Spec}^h([-1]^{-1}K_*^{\text{MW}}(F)) & \longrightarrow & \text{Spec}^h(K_*^{\text{MW}}(F)) \\ & & \downarrow (\)^0 & & \\ & & \text{Spec}([-1]^{-1}K_0^{\text{MW}}(F)) & & \end{array}$$

Here $C[-1]$ is the cone of $[-1]: S^0 \rightarrow S^\alpha$ and $\mathcal{H}(F)/\langle C[-1] \rangle$ is the Verdier quotient by the thick subcategory $\langle C[-1] \rangle$ generated by $C[-1]$, whence $\text{Spc}(\mathcal{H}(F)/\langle C[-1] \rangle)$ is naturally homeomorphic to $U(C[-1])$. Thus the restriction of ρ^\bullet to $\text{Spc}(\mathcal{H}(F)/\langle C[-1] \rangle)$ has image inside of $D_{[-1]}$, and this specifies the left-hand vertical map in the diagram. Since $[-1]$ is invertible in $\mathcal{H}(F)/\langle C[-1] \rangle$, the quotient functor $\mathcal{H}(F) \rightarrow \mathcal{H}(F)/\langle C[-1] \rangle$

factors through $[-1]^{-1}\mathcal{H}(F)$. The top row of the diagram is given by applying Spc to this factorization. Naturality of ρ^\bullet produces the rest of the diagram except for the map $(\)^0: \mathrm{Spec}^h([-1]^{-1}K_*^{\mathrm{MW}}(F)) \rightarrow \mathrm{Spec}([-1]^{-1}K_0^{\mathrm{MW}}(F))$, which is the usual map of intersection with the 0th graded piece.

By Step 3 the map $(\)^0$ is a homeomorphism. Additionally, the vertical composite is the comparison map ρ , which is surjective by connectivity of the category $[-1]^{-1}\mathcal{H}(F)$ and [7, Theorem 7.13]. We conclude that the middle upper vertical ρ^\bullet is surjective. Furthermore, the bottom row in our diagram is a homeomorphism followed by an embedding so that the composite is the inclusion of $D_{[-1]}$ in $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$. It follows that the image of the right-hand vertical ρ^\bullet contains all of $D_{[-1]}$. We conclude that ρ^\bullet is surjective, as desired. \square

4 Explicit primes

The methods of the previous section show that ρ^\bullet is surjective without explicitly identifying any tensor triangular primes living over elements of $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$. In this section, we construct elements of $\mathrm{Spc}(\mathcal{C}(F))$ living over elements of $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$. We focus on $\mathrm{Spc}(\mathcal{C}(F))$ because we are able to construct a rich class of fields for $\mathcal{C}(F)$. Using a different approach in Section 6, we construct fields for $\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c)$.

Before constructing field spectra, we will need some preliminary facts about fields in tensor triangulated categories, Milnor–Witt K –theory, Morel’s homotopy t –structure, and the Dugger–Isaksen cellularization functor. We gather these facts in the next four subsections. Readers familiar with this content should focus on Definition 4.1, Proposition 4.2, Definition 4.4, and Proposition 4.5, acquaint themselves with our notation, and then move on to Section 4.5.

4.1 Fields in tensor triangulated categories

Let $(\mathcal{C}, \otimes, 1)$ be a tensor triangulated category. Recall that a *ring* in \mathcal{C} is an object R equipped with a unit $\eta: 1 \rightarrow R$ and multiplication $\mu: R \otimes R \rightarrow R$ making R a monoid in \mathcal{C} .³ An R –*module* in \mathcal{C} is an object M equipped with an action map $\alpha: R \otimes M \rightarrow M$ satisfying the standard axioms. Let $\mathrm{Pic}(\mathcal{C})$ denote the Picard group of \otimes –invertible objects in \mathcal{C} . We call an R –module *free* if it is module-isomorphic to a coproduct of modules of the form $P \otimes R$ where each P is in $\mathrm{Pic}(\mathcal{C})$.

³The monoid and ring nomenclature are somewhat at odds. Of course, a monoid in \mathbb{Z} –modules is a ring, and this is the inspiration for the terminology.

Definition 4.1 A ring $K \in \mathcal{C}$ is called a *skew field* if every K -module in \mathcal{C} is free.

We will abuse terminology and call a skew field K a *field*, even when K is not commutative. Field objects are important in tensor triangular geometry because they are a rich source of prime ideals.

Proposition 4.2 Suppose that $K \in \mathcal{C}$ is a field. Let \mathcal{D} be a tensor triangulated subcategory of \mathcal{C} . Then

$$\ker_{\mathcal{D}}(K) := \{X \in \mathcal{D} \mid K \otimes X \cong 0\}$$

is a tensor triangular prime in \mathcal{D} .

Proof It is easy to check that $\ker_{\mathcal{D}} K$ is a tensor triangular ideal, so all that remains to show is primality. Suppose that $X \otimes Y \in \ker_{\mathcal{D}} K$ for some X, Y in \mathcal{D} . Since $K \otimes X$ is a K -module and K is a field, we have

$$K \otimes X \cong \bigoplus_{i \in I} P_i \otimes K,$$

where $P_i \in \text{Pic}(\mathcal{C})$ for $i \in I$. Thus

$$0 \cong K \otimes X \otimes Y \cong \bigoplus_{i \in I} P_i \otimes K \otimes Y.$$

Since each P_i is \otimes -invertible, this is only possible if $I = \emptyset$ (in which case $X \in \ker_{\mathcal{D}} K$) or $K \otimes Y \cong 0$ (in which case $Y \in \ker_{\mathcal{D}} K$). Thus $\ker_{\mathcal{D}} K \in \text{Spc}(\mathcal{D})$, as desired. \square

Remark 4.3 We are considering $P \otimes R$ to be a free R -module for any $P \in \text{Pic}(\mathcal{C})$. In specific cases, one might wish to restrict this notion. For example in the motivic setting, where the structure of $\text{Pic}(\text{SH}^{\mathbb{A}^1}(F))$ is complicated and not completely determined (see [3; 19]), it could be reasonable to only allow smashing with $S^{m+n\alpha}$ rather than arbitrary invertible spectra. (Indeed, the fields we produce are of this form.) However, it will not make a difference in our work below and so we work with the general notion.

4.2 The residue fields of Milnor–Witt K -theory

In order to produce fields for $\mathcal{C}(F)$, we will need to understand the residue fields of Milnor–Witt K -theory, as defined below.

Definition 4.4 For \mathfrak{p} a homogeneous prime ideal of $K_*^{\text{MW}}(F)$, we define the *residue field* of $K_*^{\text{MW}}(F)$ at \mathfrak{p} ,

$$L(\mathfrak{p}) := \text{GrFrac}(K_*^{\text{MW}}(F)/\mathfrak{p}),$$

to be the graded field of fractions of the \mathbb{Z} -graded integral domain $K_*^{\text{MW}}(F)/\mathfrak{p}$; this is formed by inverting all nonzero homogeneous classes in $K_*^{\text{MW}}(F)/\mathfrak{p}$.

By elementary computation and Theorem 2.1, we arrive at the following list of residue fields for $K_*^{\text{MW}}(F)$.

Proposition 4.5 *The residue fields of $K_*^{\text{MW}}(F)$ take the following forms (with the obvious $K_*^{\text{MW}}(F)$ -module structure):*

- (1) $L([F^\times], \eta) \cong \mathbb{Q}$,
- (2) $L([F^\times], \eta, p) \cong \mathbb{F}_p$,
- (3) $L([F^\times], 2) \cong \mathbb{F}_2[\eta^{\pm 1}]$,
- (4) $L([P_\alpha], h) \cong \mathbb{Q}[\eta^{\pm 1}]$,
- (5) $L([P_\alpha], \eta, 2) \cong \mathbb{F}_2[[-1]^{\pm 1}]$, and
- (6) $L([P_\alpha], h, p) \cong \mathbb{F}_p[\eta^{\pm 1}]$ for $p \neq 2$. □

4.3 The homotopy t -structure

Morel [26] has introduced a t -structure on $\text{SH}^{\mathbb{A}^1}(F)$ defined in the following way. Let $\text{SH}_{\geq 0}^{\mathbb{A}^1}(F)$ denote the full subcategory of *connective motivic spectra*, those X such that $\pi_m(X)_n = 0$ for $m < 0$ and $n \in \mathbb{Z}$. Similarly, let $\text{SH}_{\leq 0}^{\mathbb{A}^1}(F)$ consist of all X such that $\pi_m(X)_n = 0$ for $m > 0$ and $n \in \mathbb{Z}$. By [26, Theorem 5.2.3], the triple $(\text{SH}^{\mathbb{A}^1}(F), \text{SH}_{\geq 0}^{\mathbb{A}^1}(F), \text{SH}_{\leq 0}^{\mathbb{A}^1}(F))$ is a nondegenerate t -structure, called the *homotopy t -structure*.

In the usual fashion, we can define $\text{SH}_{\leq m}^{\mathbb{A}^1}(F)$ and $\text{SH}_{\geq m}^{\mathbb{A}^1}(F)$ for $m \in \mathbb{Z}$, and the adjoints to inclusion $\tau_{\leq m}: \text{SH}^{\mathbb{A}^1}(F) \rightarrow \text{SH}_{\leq m}^{\mathbb{A}^1}(F)$ and $\tau_{\geq m}: \text{SH}^{\mathbb{A}^1}(F) \rightarrow \text{SH}_{\geq m}^{\mathbb{A}^1}(F)$. For $X \in \text{SH}^{\mathbb{A}^1}(F)$ we define

$$\mathbb{H}_m X := \tau_{\leq m} \tau_{\geq m} X,$$

and call $\mathbb{H}_0 X$ the *heart* of X . Morel identifies the heart $\mathrm{SH}_{\leq 0}^{\mathbb{A}^1}(F) \cap \mathrm{SH}_{\geq 0}^{\mathbb{A}^1}(F)$ of the homotopy t -structure with the so-called category of *homotopy modules*⁴ but we will not use this result here.

The next lemma follows from [14, Section 2.3]. We include a proof for completeness.

Lemma 4.6 *Let E be a motivic ring spectrum. Then $\mathbb{H}_0 E$ is a motivic ring spectrum as well.*

Proof The multiplication on $\mathbb{H}_0 E$ arises as a composite

$$\mathbb{H}_0 E \wedge \mathbb{H}_0 E \rightarrow \mathbb{H}_0(\mathbb{H}_0 E \wedge \mathbb{H}_0 E) \rightarrow \mathbb{H}_0 E.$$

We begin by constructing the second map. First observe that $\mathbb{H}_0(\mathbb{H}_0 E \wedge \mathbb{H}_0 E) \simeq \mathbb{H}_0(\tau_{\geq 0} E \wedge \tau_{\geq 0} E)$. Then $\mathbb{H}_0(\mathbb{H}_0 E \wedge \mathbb{H}_0 E) \rightarrow \mathbb{H}_0 E$ arises by applying \mathbb{H}_0 to the natural map $\tau_{\geq 0} E \wedge \tau_{\geq 0} E \rightarrow \tau_{\geq 0} E$.

To construct $\mathbb{H}_0 E \wedge \mathbb{H}_0 E \rightarrow \mathbb{H}_0(\mathbb{H}_0 E \wedge \mathbb{H}_0 E)$, simply observe that whenever A is connective, there is a natural map $A \rightarrow \mathbb{H}_0 A$ (which is just the natural map $A \rightarrow \tau_{\leq 0} A$).

It is formal to check that the above map $\mathbb{H}_0 E \wedge \mathbb{H}_0 E \rightarrow \mathbb{H}_0 E$ and the unit

$$S^0 \rightarrow \mathbb{H}_0 S^0 \rightarrow \mathbb{H}_0 E$$

make $\mathbb{H}_0 E$ a monoid in $\mathrm{SH}^{\mathbb{A}^1}(F)$, as desired. □

4.4 Cellularization

Write $i: \mathrm{SH}^{\mathbb{A}^1}(F)^{\mathrm{cell}} \subseteq \mathrm{SH}^{\mathbb{A}^1}(F)$ for the homotopy category of motivic cell spectra, ie the localizing subcategory generated by $S^{m+n\alpha}$, for $m, n \in \mathbb{Z}$. Recall from [12, Proposition 7.3] that, on the level of homotopy categories, there is a functor

$$\mathrm{Cell}: \mathrm{SH}^{\mathbb{A}^1}(F) \rightarrow \mathrm{SH}^{\mathbb{A}^1}(F)^{\mathrm{cell}}$$

which is right adjoint to the inclusion. As is standard, we will abuse notation and write again $\mathrm{Cell}(X)$ for $i \mathrm{Cell}(X)$. In particular, we have a natural map $\mathrm{Cell}(X) \rightarrow X$. Note also that Cell is a lax symmetric monoidal functor, as it is right adjoint to a symmetric monoidal functor. We immediately conclude the following.

Lemma 4.7 *Let E be a motivic ring spectrum. Then $\mathrm{Cell}(E)$ is a motivic ring spectrum as well.* □

⁴These are $\underline{K}_*^{\mathrm{MW}}(F)$ -modules \underline{M}_* equipped with *contraction isomorphisms* $(\underline{M}_{n+1})_{-1} \cong \underline{M}_n$, where $()_{-1}$ is Morel’s contraction construction [26, Definition 4.3.10].

4.5 Field spectra and explicit primes in $\text{Spc}(\mathcal{C}(F))$

We now build some cellular field spectra and realize some explicit tensor triangular primes in $\mathcal{C}(F)$. We write

$$\mathbb{H}_0^{\text{cell}}(E) := \text{Cell}(\mathbb{H}_0(E)).$$

Lemma 4.8 Fix $\mathfrak{p} \in \text{Spec}^h(K_*^{\text{MW}}(F))$ and suppose there exists a (homotopy associative) ring spectrum E such that $\pi_0(E)_* \cong L(\mathfrak{p})$ as $K_*^{\text{MW}}(F)$ –modules. Then $\mathbb{H}_0^{\text{cell}}(E)$ is a field in $\text{SH}^{\mathbb{A}^1}(F)^{\text{cell}}$ and $\rho^\bullet(\ker_{\mathcal{C}(F)}(\mathbb{H}_0^{\text{cell}}(E))) = \mathfrak{p}$.

Proof First, note that Lemmas 4.6 and 4.7 imply that $\mathbb{H}_0^{\text{cell}}(E)$ is a cellular ring spectrum.

Let M be a $\mathbb{H}_0^{\text{cell}}(E)$ –module and note that $\pi_*(M)_*$ is an $L(\mathfrak{p})$ –module. Since $L(\mathfrak{p})$ is a graded field, we may choose a homogeneous basis for $\pi_*(M)_*$ as a free $\pi_*(\mathbb{H}_0^{\text{cell}}(E))_*$ –module. This basis induces a map

$$\bigvee_{\lambda \in \Lambda} \Sigma^\lambda \mathbb{H}_0^{\text{cell}}(E) \rightarrow M,$$

where each λ is in $\mathbb{Z} \oplus \mathbb{Z}\{\alpha\}$, which induces an isomorphism on homotopy groups. This map is thus a weak equivalence since homotopy groups detect weak equivalences between cellular motivic spectra. Hence M is free, as desired, so $\mathbb{H}_0^{\text{cell}}(E)$ is a cellular motivic field spectrum.

By Proposition 4.2, $\ker_{\mathcal{C}(F)}(\mathbb{H}_0^{\text{cell}} E)$ is in $\text{Spc}(\mathcal{C}(F))$. Moreover,

$$\begin{aligned} &\rho^\bullet(\ker_{\mathcal{C}(F)}(\mathbb{H}_0^{\text{cell}} E)) \\ &= \{f \in K_*^{\text{MW}}(F) \mid \text{cone}(f) \notin \ker_{\mathcal{C}(F)}(\mathbb{H}_0^{\text{cell}}(E))\} \\ &= \{f \in K_*^{\text{MW}}(F) \mid \mathbb{H}_0^{\text{cell}}(E) \wedge \text{cone}(f) \not\cong * \} \\ &= \{f \in K_*^{\text{MW}}(F) \mid \mathbb{H}_0^{\text{cell}}(E) \xrightarrow{f} S^{*\alpha} \wedge \mathbb{H}_0^{\text{cell}}(E) \text{ is not a weak equivalence}\} \\ &= \{f \in K_*^{\text{MW}}(F) \mid L(\mathfrak{p}) \xrightarrow{f} L(\mathfrak{p}) \text{ is not an isomorphism}\} \\ &= \mathfrak{p}. \end{aligned}$$

□

Proposition 4.9 If $\mathfrak{p} \in \text{Spec}^h(K_*^{\text{MW}}(F))$ is one of types (3)–(6) in Theorem 2.1, then there exists a motivic ring spectrum $E \in \text{SH}^{\mathbb{A}^1}(F)$ such that $\pi_0(E)_* \cong L(\mathfrak{p})$.

Proof First suppose $\mathfrak{p} = ([F^\times], 2)$ is of type (3), so $L(\mathfrak{p}) \cong \mathbb{F}_2[\eta^{\pm 1}]$. Fix an embedding $i: F \hookrightarrow \bar{F}$ of F into an algebraic closure \bar{F} . The functor $i^*: \text{SH}^{\mathbb{A}^1}(F) \rightarrow \text{SH}^{\mathbb{A}^1}(\bar{F})$

has a right adjoint, i_* , which is lax-symmetric monoidal. In particular, i_* preserves motivic ring spectra. Let $KT_{\bar{F}}$ denote Witt K -theory [17] over \bar{F} . Then we may take $E = i_*KT_{\bar{F}}$. Indeed, we may compute

$$\begin{aligned} \pi_0(E)_* &\cong [S^0, S^{*\alpha} \wedge i_*KT_{\bar{F}}]_{\text{SH}^{\Delta^1}(F)} \\ &\cong [S^{-*\alpha}, i_*KT_{\bar{F}}]_{\text{SH}^{\Delta^1}(F)} \\ &\cong [S^{-*\alpha}, KT_{\bar{F}}]_{\text{SH}^{\Delta^1}(\bar{F})} \\ &\cong W(\bar{F})[\eta^{\pm 1}] \\ &\cong \mathbb{F}_2[\eta^{\pm 1}], \end{aligned}$$

since the Witt ring of an algebraically closed field is always isomorphic to \mathbb{F}_2 via the rank homomorphism.

Now suppose that F is formally real with ordering α . (This is necessary for primes of types (4)–(6) to exist.) Fix an embedding $j: F \hookrightarrow F_\alpha$. Suppose that \mathfrak{p} is of type (4), so that $L(\mathfrak{p}) \cong \mathbb{Q}[\eta^{\pm 1}]$. Then we may take $E = j_*KT_{F_\alpha} \otimes \mathbb{Q}$, and a computation similar to that above gives

$$\pi_0(E)_* \cong \mathbb{Q}[\eta^{\pm 1}],$$

since $W(F_\alpha) \cong \mathbb{Z}$.

Suppose that \mathfrak{p} is of type (6), so $L(\mathfrak{p}) \cong \mathbb{F}_p[\eta^{\pm 1}]$. Then we may take $E = j_*KT_{F_\alpha}/p$, and we again get the desired isomorphism

$$\pi_0(E)_* \cong \mathbb{F}_p[\eta^{\pm 1}].$$

Now suppose $\mathfrak{p} = ([P_\alpha], \eta, 2)$ is of type (5), so that $L(\mathfrak{p}) \cong \mathbb{F}_2[[-1]^{\pm 1}]$. Let $(H\mathbb{F}_2)_{F_\alpha}$ denote the mod-2 motivic cohomology spectrum over F_α . Let $[-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}$ denote the colimit of iterated multiplication by $[-1]$ on $(H\mathbb{F}_2)_{F_\alpha}$. Then we may take $E = j_*[-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}$. Indeed, we compute

$$\begin{aligned} \pi_0(E)_* &\cong [S^0, S^{*\alpha} \wedge j_*[-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}]_{\text{SH}^{\Delta^1}(F)} \\ &\cong [S^{-*\alpha}, j_*[-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}]_{\text{SH}^{\Delta^1}(F)} \\ &\cong [S^{-*\alpha}, [-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}]_{\text{SH}^{\Delta^1}(F_\alpha)} \\ &\cong [-1]^{-1}K_*^M(F_\alpha) \\ &\cong \mathbb{F}_2[[-1]^{\pm 1}]. \end{aligned}$$

The penultimate isomorphism utilizes the Milnor conjecture [34], while the final isomorphism is an invocation of [25, Example 1.6]. □

Combining Lemma 4.8 and Proposition 4.9, we arrive at the following list of explicit primes in $\mathcal{C}(F)$ and their images in $\text{Spec}^h(K_*^{\text{MW}}(F))$.

Theorem 4.10 *Fix an embedding $i: F \hookrightarrow \bar{F}$ of F into its algebraic closure. If $\alpha \in X_F \neq \emptyset$, fix an embedding $j = j_\alpha: F \hookrightarrow F_\alpha$ of F into its real closure with respect to α . Let*

$$E_3 = i_*KT_{\bar{F}}, \quad E_4 = j_*KT_{F_\alpha} \otimes \mathbb{Q}, \quad E_5 = j_*[-1]^{-1}(H\mathbb{F}_2)_{F_\alpha}, \quad E_6 = j_*KT_{F_\alpha}/p.$$

Writing

$$\mathcal{P}_i = \ker_{\mathcal{C}(F)}(\mathbb{H}_0^{\text{cell}}(E_i)),$$

we have

$$\begin{aligned} \rho^\bullet(\mathcal{P}_3) &= ([F^\times], 2), & \rho^\bullet(\mathcal{P}_4) &= ([P_\alpha], h), \\ \rho^\bullet(\mathcal{P}_5) &= ([P_\alpha], \eta, 2), & \rho^\bullet(\mathcal{P}_6) &= ([P_\alpha], h, p). \end{aligned} \quad \square$$

Remark 4.11 In Proposition 5.3 we will determine explicit tensor triangular primes living over primes in $K_*^{\text{MW}}(F)$ of types (1) and (2). These are realized as acyclics for field spectra in Remark 6.4.

5 Betti realization and tensor triangular primes

In this section, we study the role of Betti realization functors in determining tensor triangular primes in $\text{SH}^{\mathbb{A}^1}(F)^c$. We consider both the complex Betti realization functor

$$\text{Re}_{B,i}: \text{SH}^{\mathbb{A}^1}(F)^c \rightarrow \text{SH}^{\text{fin}}$$

associated with a complex embedding $i: F \hookrightarrow \mathbb{C}$ and, for each $\alpha \in X_F$, the C_2 -equivariant Betti realization functor

$$\text{Re}_{B,j}^{C_2}: \text{SH}^{\mathbb{A}^1}(F)^c \rightarrow \text{SH}(C_2)^c$$

associated with an order-preserving embedding $j: F \hookrightarrow \mathbb{R}$. See [15, Section 4.4] for a review of these constructions. We will frequently leave the embedding into \mathbb{C} or \mathbb{R} implicit and call these functors Re_B and $\text{Re}_B^{C_2}$, respectively.

In order to study $\text{Spc}(\text{Re}_B)$ and $\text{Spc}(\text{Re}_B^{C_2})$, we will need to know the structure of the spaces $\text{Spc}(\text{SH}^{\text{fin}})$ and $\text{Spc}(\text{SH}(C_2)^c)$, respectively. We recall these spaces in the next two subsections, following Balmer [7] and Balmer and Sanders [10], respectively. In the final subsection, we present our results on $\text{Spc}(\text{Re}_B)$ and $\text{Spc}(\text{Re}_B^{C_2})$.

5.1 Tensor triangular primes in $\mathrm{SH}^{\mathrm{fin}}$

Building on [16], Balmer determines the structure of $\mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$ in [7, Section 9]. We briefly recall the result here.

Let p be a rational prime and let n be a positive integer. Let $K(p, n - 1)$ denote $(n - 1)^{\mathrm{th}}$ Morava K -theory at the prime p , with the convention that $K(p, 0) = H\mathbb{Q}$, the rational Eilenberg–MacLane spectrum. Let $\mathcal{C}_{p,n}$ denote the kernel of $K(p, n - 1)$ -homology.⁵ In particular, $\mathcal{C}_{p,1} = \mathcal{C}_{q,1}$ for all p, q , and we denote this set by $\mathcal{C}_{0,1}$. Let $\mathcal{C}_{p,\infty}$ denote the set of p -locally trivial spectra.

Theorem 5.1 [7, Corollary 9.5] *Every element of $\mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$ is of the form $\mathcal{C}_{p,n}$ for some rational prime p and n a positive integer or ∞ . The only duplication occurs when $n = 1$, in which case $\mathcal{C}_{p,1} = \mathcal{C}_{0,1}$, the set of finite-torsion spectra. The Hasse diagram for $\mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$ takes the form shown in Figure 2, and the closure of $S \subseteq \mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$ consists of all the elements in or above S in the Hasse diagram.*




Figure 2: The Hasse diagram for $\mathrm{Spc}(\mathrm{SH}^{\mathrm{fin}})$. Inclusion of tensor triangular primes is downwards, and closure is upwards in the diagram.

⁵Balmer sets $\mathcal{C}_{p,n}$ equal to the kernel of $K(p, n - 1) \wedge ()_{(p)}$. Since $K(p, n - 1)$ is already p -local and p -localization is smashing, we have $\ker K(p, n - 1) = \ker K(p, n - 1) \wedge ()_{(p)}$.

5.2 Tensor triangular primes in $\text{SH}(C_2)^c$

In [10], Balmer and Sanders completely determine the tensor triangular spectrum $\text{Spc}(\text{SH}(G)^c)$ for G a finite group of square-free order. Here we recall their results, specializing to the group $G = C_2$.

Let p be a rational prime or 0, let n be a positive integer or ∞ , let H be a subgroup of C_2 , and let $\Phi^H: \text{SH}(C_2) \rightarrow \text{SH}$ denote the geometric fixed points functor for H . Define

$$\mathcal{P}(H, p, n) = (\Phi^H)^{-1}(\mathcal{L}_{p,n}).$$

Theorem 5.2 [10, Theorem 8.12 and Diagram (1.3)] *Every element of $\text{Spc}(\text{SH}(C_2)^c)$ is of the form $\mathcal{P}(H, p, n)$ for precisely one H , p and n . The Hasse diagram for $\text{Spc}(\text{SH}(C_2)^c)$ takes the form shown in Figure 3 and the closure of $S \subseteq \text{Spc}(\text{SH}(C_2)^c)$ consists of all the elements in or above S in the Hasse diagram.*




Figure 3: The Hasse diagram for $\text{Spc}(\text{SH}(C_2)^c)$. Inclusion of tensor triangular primes is downwards, and closure is upwards in the diagram.

5.3 The maps $\text{Spc}(\text{Re}_B)$ and $\text{Spc}(\text{Re}_B^{C_2})$

Proposition 5.3 *Suppose that F embeds into \mathbb{C} with associated Betti realization functor Re_B . Then the composite map $\rho^\bullet \circ \text{Spc}(\text{Re}_B): \text{Spc}(\text{SH}^{\text{fin}}) \rightarrow \text{Spec}^h(K_*^{\text{MW}}(F))$ takes the values*

$$\begin{aligned} \mathcal{C}_{0,1} &\mapsto ([F^\times], \eta), \\ \mathcal{C}_{p,n} &\mapsto ([F^\times], p, \eta), \end{aligned}$$

for all p and $n \geq 2$. In particular, the image of $\rho^\bullet \circ \text{Spc}(\text{Re}_B)$ is

$$V_\eta \setminus \{([P_\alpha], 2, \eta) \mid \alpha \in X_F\},$$

the collection of primes of type (1) or (2).

Proof Let $\pi^*(S^0)$ denote the cohomotopy groups of the sphere, so that $\pi^n(S^0) = [S^0, S^n] \cong \pi_{-n}(S^0)$. Betti realization induces a commutative diagram

$$\begin{array}{ccc} \text{Spc}(\text{SH}^{\text{fin}}) & \xrightarrow{\text{Spc}(\text{Re}_B)} & \text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c) \\ \rho_{S^1}^\bullet \downarrow & & \downarrow \rho_{S^0}^\bullet \\ \text{Spec}^h(\pi^*(S^0)) & \longrightarrow & \text{Spec}^h(K_*^{\text{MW}}(F)) \end{array}$$

wherein the bottom horizontal arrow takes $\pi^{>0}(S^0)$ to $([F^\times], \eta)$ and $(p, \pi^{>0}(S^0))$ to $([F^\times], p, \eta)$. The rest of the computation is easy. \square

For completeness, we include the following slight enhancement of [7, Proposition 10.4].

Corollary 5.4 *When $F = \mathbb{C}$, the map $\text{Spc}(\text{Re}_B)$ is monic with image contained in $\text{supp}(\mathbb{P}^2)$.*

Proof Let $c^*: \text{SH}^{\text{fin}} \rightarrow \text{SH}^{\mathbb{A}^1}(F)^c$ be the functor induced by the constant presheaf functor. The map $\text{Spc}(\text{Re}_B)$ is monic because $\text{Re}_B \circ c^* = \text{id}$. Since $X_F = \emptyset$, the image of the map $\rho^\bullet \circ \text{Spc}(\text{Re}_B)$ is exactly V_η , and $(\rho^\bullet)^{-1}(V_\eta) = \text{supp}(C\eta) = \text{supp}(\mathbb{P}^2)$. \square

We now turn to the study of $\text{Spc}(\text{Re}_B^{C_2})$ when F has a real embedding.

Proposition 5.5 *Suppose F has an embedding into \mathbb{R} compatible with $\alpha \in X_F$, and let $\text{Re}_{B,\alpha}^{C_2}$ denote the corresponding C_2 -equivariant Betti realization functor. Then*

the composite map $\rho^\bullet \circ \text{Spc}(\text{Re}_{B,\alpha}^{C_2}): \text{Spc}(\text{SH}(C_2)^c) \rightarrow \text{Spec}^h(K_*^{\text{MW}}(F))$ takes the following values:

$$\begin{aligned} \mathcal{P}(e, 0, 1) &\mapsto ([F^\times], \eta), \\ \mathcal{P}(C_2, 0, 1) &\mapsto ([P_\alpha], h), \\ \mathcal{P}(e, p, n) &\mapsto ([F^\times], p, \eta) \quad \text{for all } p \text{ and } n \geq 2, \\ \mathcal{P}(C_2, p, n) &\mapsto ([P_\alpha], p, h) \quad \text{for } p \neq 2 \text{ and } n \geq 2, \\ \mathcal{P}(C_2, 2, n) &\mapsto ([P_\alpha], 2, \eta) \quad \text{for } n \geq 2. \end{aligned}$$

Remark 5.6 If F admits an embedding into \mathbb{R} then for each $\alpha \in X_F$ there is an embedding compatible with the ordering α . The proposition implies that the union of the images of all of the resulting $\rho^\bullet \circ \text{Spc}(\text{Re}_{B,\alpha}^{C_2})$ is

$$\text{Spec}^h(K_*^{\text{MW}}(F)) \setminus \{([F^\times], 2)\}.$$

Proof Let $g = \rho^\bullet \circ \text{Spc}(\text{Re}_{B,\alpha}^{C_2})$. It is easy to check that $g(\mathcal{P}(e, 0, 1)) = ([F^\times], \eta)$ and $g(C_2, 0, 1) = ([P_\alpha], h)$. Since g is continuous, we know that $g(\overline{\mathcal{P}(e, 0, 1)}) \subseteq (\overline{[F^\times], \eta})$, where $(\overline{})$ denotes the closure operator. Observing the form of $\text{Spc}(\text{SH}(C_2)^c)$ in Figure 3 and of $\text{Spec}^h(K_*^{\text{MW}}(F))$ in Figure 1, we see that $g(\mathcal{P}(e, p, n)) = ([F^\times], p, \eta)$ for all p and $n \geq 2$.

We now check that $g(\mathcal{P}(C_2, 2, n)) = ([P_\alpha], 2, \eta)$ for $n \geq 2$. Since $g(\mathcal{P}(C_2, 0, 1)) = ([P_\alpha], h)$, it follows from the closure continuity condition that $g(\mathcal{P}(C_2, 2, n))$ is either $([F^\times], 2, \eta)$ or $([P_\alpha], 2, \eta)$. Observe, though, that $\Phi^{C_2} \text{Re}_{B,\alpha}^{C_2} C[-1] \simeq *$. (Simply apply $\Phi^{C_2} \text{Re}_{B,\alpha}^{C_2}$ to the defining cofiber sequence for $C[-1]$.) Thus

$$\begin{aligned} [-1] &\notin \rho^\bullet(\text{Spc}(\text{Re}_{B,\alpha}^{C_2})(\mathcal{P}(C_2, 2, n))) \\ &= \{f \in K_*^{\text{MW}}(F) \mid (\Phi^{C_2} \text{Re}_{B,\alpha}^{C_2} Cf)_{(2)} \wedge K(2, n-1) \not\simeq *\}. \end{aligned}$$

Also $g(\mathcal{P}(C_2, 2, n)) \neq ([P_\alpha], h)$ since, for example, $\Phi^{C_2}(C\eta) \wedge K(2, n-1) \not\simeq *$ so $\eta \in g(\mathcal{P}(C_2, 2, n))$. We conclude that $g(\mathcal{P}(C_2, 2, n)) = ([P_\alpha], 2, \eta)$.

A similar argument shows that $g(C_2, p, n) = ([P_\alpha], p, h)$ for p odd and $n \geq 2$. \square

Remark 5.7 By Theorem 1.1 and the fact that $C\eta \simeq \Sigma\mathbb{P}^2$, we know that $([F^\times], 2)$ belongs to $\rho^\bullet(\text{supp}(\mathbb{P}^2))$. This and Proposition 5.5 imply that $\text{Spc}(\text{Re}_B^{C_2})$ does not hit all of $\text{supp}(\mathbb{P}^2)$, indicating that the tensor triangular spectrum $\text{Spc}(\text{SH}^{\mathbb{A}^1}(\mathbb{R})^c)$ is richer than $\text{Spc}(\text{SH}(C_2)^c)$. Compare with Corollary 5.4. The moral seems to

be that significant new ideas are necessary to unravel the structure of $\text{supp}(\mathbb{P}^2) \subseteq \text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c)$.

Corollary 5.8 *When $F \subseteq \mathbb{R}$, the map $\text{Spc}(\text{Re}_B^{C_2})$ is monic with image contained in*

$$(\rho^\bullet)^{-1}(\text{Spec}^h(K_*^{\text{MW}}(F)) \setminus \{([F^\times], 2)\}) \subsetneq \text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c).$$

Proof Denote the functor of [15, Theorem 4.6] by $c_{F[i]/F}^*$. The map $\text{Spc}(\text{Re}_B^{C_2})$ is monic since $\text{Re}_B^{C_2} \circ c_{F[i]/F}^* = \text{id}$. The restriction on the image of $\text{Spc}(\text{Re}_B^{C_2})$ follows immediately from Proposition 5.5. □

Remark 5.9 This answers a question posed in [21, Remark 7.2.11]. In particular, $\text{Spc}(\text{Re}_B^{C_2})$ is never surjective.

We conclude this section by providing a sketch of known structures in $\text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c)$ when F is a subfield of \mathbb{R} with precisely one ordering. (For example, F could be \mathbb{R} itself.) See Figure 4 below. These structures include the images of the injective maps $\text{Spc}(\text{Re}_B)$ and $\text{Spc}(\text{Re}_B^{C_2})$ along with a mysterious (but nonempty) subspace $(\rho^\bullet)^{-1}([F^\times], 2)$. We have also drawn the map ρ^\bullet with target $\text{Spec}^h(K_*^{\text{MW}}(F))$ so that known parts of fibers are aligned vertically over their images. Note that we have drawn $\text{Spec}^h(K_*^{\text{MW}}(F))$ in a fashion slightly different from that in Figure 1 so as to highlight similarities with $\text{Spc}(\text{SH}(C_2)^c)$.

The interested reader can easily elaborate upon Figure 4 in the case where F is a subfield of \mathbb{R} with more than one ordering. The result would be an “ X_F fan” over primes in $K_*^{\text{MW}}(F)$ of types (4)–(6). To see what is known for F a nonreal subfield of \mathbb{C} , simply delete everything of these types.

Remark 5.10 Since the first version of this paper was written, work of Bachmann [4] appeared, which makes it possible to completely specify $U(\text{cone}([-1]))$, the portion of the spectrum $\text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c)$ living over primes in $K_*^{\text{MW}}(F)$ of types (4)–(6). Bachmann proves that $\text{SH}^{\mathbb{A}^1}(F)[[-1]^{-1}]$ is equivalent to $\text{SH}(X_F)$, the homotopy category of sheaves of spectra on X_F . It follows that $U(\text{cone}([-1]))$ is homeomorphic to $\text{Spc}(\text{SH}(X_F))$. Thus $U(\text{cone}([-1]))$ is the product of X_F with the topological space specified by the towers on the left-hand side of Figure 4.




Figure 4: Known structures in $\text{Sp}(\text{SH}^{\mathbb{A}^1}(F)^c)$ for F a subfield of \mathbb{R} with one ordering α , arranged fiberwise over $\text{Spec}^h(K_*^{\text{MW}}(F))$ with respect to ρ^\bullet .

6 Field spectra for $\text{SH}^{\mathbb{A}^1}(F)^c$

In Section 4, we constructed fields in the homotopy category of cellular spectra. Acyclics for these fields gave explicit tensor triangular primes in $\mathcal{C}(F)$. In this section we further explore motivic fields. We begin by showing that fields from the topological stable homotopy category can be imported into $\text{SH}^{\mathbb{A}^1}(F)$, when F admits a complex embedding.

If F is a subfield of \mathbb{C} then we have an adjoint pair $\text{Re}_B: \text{SH}^{\mathbb{A}^1}(F) \rightleftarrows \text{SH} : \mathcal{S}_B$, where Re_B extends the functor sending a smooth complex variety X to the space $X(\mathbb{C})$. Similarly, if F is a subfield of \mathbb{R} , we have an adjoint pair $\text{Re}_B^{C_2}: \text{SH}^{\mathbb{A}^1}(F) \rightleftarrows \text{SH}(C_2) : \mathcal{S}_B^{C_2}$.

In order to treat both of these cases simultaneously, we write

$$\text{Re}: \text{SH}^{\mathbb{A}^1}(F) \rightleftarrows \text{SH}(G) : \mathcal{S}$$

for either of these pairs, where G is either $\{e\}$ or C_2 .

Lemma 6.1 *The functor $\mathcal{S}: \text{SH}(G) \rightarrow \text{SH}^{\mathbb{A}^1}(F)$ commutes with all coproducts.*

Proof For any X in Sm/F and any $n \in \mathbb{Z}$, we have that $\Sigma_T^n X_+$ is dualizable by [30, Theorem 4.9], and since Re is symmetric monoidal, $\text{Re}(\Sigma_T^n X_+)$ is dualizable (see eg [13, Proposition 3.10]), hence compact. Using adjunction, it is straightforward to verify that

$$\left[\Sigma_T^n X_+, \bigvee_{\alpha} \mathcal{S}(E_{\alpha}) \right]_{\text{SH}^{\mathbb{A}^1}(F)} \cong \left[\Sigma_T^n X_+, \mathcal{S}\left(\bigvee_{\alpha} E_{\alpha}\right) \right]_{\text{SH}^{\mathbb{A}^1}(F)},$$

which implies that the map $\bigvee_{\alpha} \mathcal{S}(E_{\alpha}) \rightarrow \mathcal{S}(\bigvee_{\alpha} E_{\alpha})$ is an equivalence. □

Lemma 6.2 *The canonical map*

$$\mathcal{S}(K) \wedge X \xrightarrow{\cong} \mathcal{S}(K \wedge \text{Re}(X))$$

is an equivalence for any K in $\text{SH}(G)$ and any X in $\text{SH}^{\mathbb{A}^1}(F)$.

Proof The map of the lemma is the composite of maps

$$\mathcal{S}(K) \wedge X \rightarrow \mathcal{S}(K) \wedge \mathcal{S} \text{Re}(X) \rightarrow \mathcal{S}(K \wedge \text{Re}(X)).$$

Consider the full subcategory $\mathcal{A} \subseteq \text{SH}^{\mathbb{A}^1}(F)$ whose objects satisfy the condition that $\mathcal{S}(K) \wedge X \simeq \mathcal{S}(K \wedge \text{Re}(X))$. The category \mathcal{A} is triangulated and by the previous lemma it is closed under arbitrary coproducts, ie it is a localizing subcategory. It therefore suffices to show that \mathcal{A} contains every compact spectrum X .

Since F has characteristic zero, every smooth quasiprojective F -scheme is dualizable by [30, Theorem 4.9]. By [18, Theorem 2.1.3], this implies that $\text{SH}^{\mathbb{A}^1}(F)^c$ is equal to the category of dualizable objects. Since Re is strong symmetric monoidal, $\text{Re}(DX) \simeq D(\text{Re } X)$ and $F(Y, \mathcal{S}(K)) \simeq \mathcal{S}F(\text{Re}(Y), K)$; see eg [13, Proposition 3.10 and Equation (3.4)]. We thus have equivalences

$$\begin{aligned} \mathcal{S}(K) \wedge X &\simeq F(DX, \mathcal{S}(K)) \\ &\simeq \mathcal{S}F(\text{Re}(DX), K) \\ &\simeq \mathcal{S}F(D(\text{Re } X), K) \\ &\simeq \mathcal{S}(K \wedge \text{Re } X). \end{aligned} \quad \square$$

The image of $\text{Spc}(\text{Re}): \text{Spc}(\text{SH}(G)^c) \rightarrow \text{Spc}(\text{SH}^{\mathbb{A}^1}(F)^c)$ is characterized as follows.

Theorem 6.3 *Let K be a field in $\text{SH}(G)$. Then $\mathcal{S}(K)$ is a field in $\text{SH}^{\mathbb{A}^1}(F)$. Moreover, letting $\ker K \in \text{Spc}(\text{SH}(G)^{\text{fin}})$ be the prime ideal of K -acyclics, we have*

$$\text{Spc}(\text{Re})(\ker K) = \ker(\mathcal{S}(K)).$$

Proof The functor \mathcal{S} is lax monoidal as it is right adjoint to a monoidal functor. In particular, if K is a ring in $\text{SH}(G)$, then $\mathcal{S}(K)$ is a ring in $\text{SH}^{\mathbb{A}^1}(F)$.

Let c^* denote the functor from [15, Theorem 4.6]. We first observe that if X is an $\mathcal{S}(K)$ -module then $\text{Re}(X)$ is a K -module. To see this, first note that the adjoint of the canonical isomorphism $\text{Re } c^*(K) \cong K$ is a ring map $c^*(K) \rightarrow \mathcal{S}(K)$. Therefore X is a $c^*(K)$ -module and hence $\text{Re}(X)$ is a K -module. Since K is a field, we have that $\text{Re}(X) = \bigvee_{\alpha} K$. It follows from Lemma 6.1 that $\mathcal{S} \text{Re}(X) \simeq \bigvee_{\alpha} \mathcal{S}(K)$ and so $\mathcal{S} \text{Re}(X)$ is a free $\mathcal{S}(K)$ -module.

Now consider the following comparison of retracts:

$$\begin{array}{ccccc} X & \longrightarrow & \mathcal{S}(K) \wedge X & \longrightarrow & X \\ \downarrow & & \downarrow \simeq & & \downarrow \\ \mathcal{S} \text{Re}(X) & \longrightarrow & \mathcal{S}(K \wedge \text{Re}(X)) & \longrightarrow & \mathcal{S} \text{Re}(X) \end{array}$$

By Lemma 6.2 the middle arrow is an equivalence and therefore so is the outer arrow and so X is a free $\mathcal{S}(K)$ -module as well. This establishes the first statement.

To establish the second statement, we need to see that $\mathcal{S}(K) \wedge X \simeq *$ if and only if $K \wedge \text{Re } X \simeq *$. This follows from Lemma 6.2 and the observation that if $\mathcal{S}(Z) \simeq *$ then $Z \simeq *$ (since $\pi_n(Z) = [S^n, \mathcal{S}(Z)]_{\text{SH}^{\mathbb{A}^1}(\mathbb{C})}$). □

Remark 6.4 In particular, Theorem 6.3 and Proposition 5.3 imply that $\ker(\mathcal{S}_B(H\mathbb{Q}))$ and $\ker(\mathcal{S}_B(K(p, n-1)))$ are in $\text{Spc}(\text{SH}^{\mathbb{A}^1}(F))$, mapping to $([F^\times], \eta)$ and $([F^\times], p, \eta)$, respectively, under ρ^\bullet .

Similarly, one can show that

$$\ker(\mathcal{S}_B^{C_2}(EC_{2+} \wedge K(p, n-1))) \quad \text{and} \quad \ker(\mathcal{S}_B^{C_2}(\tilde{E}C_2 \wedge K(p, n-1)))$$

belong to $\text{Spc}(\text{SH}^{\mathbb{A}^1}(F))$, with images under ρ^\bullet specified by Proposition 5.5.

Lastly we note the following restriction on fields in $\text{SH}^{\mathbb{A}^1}(F)$.

Proposition 6.5 *Let F be a subfield of \mathbb{C} . If E is a field in $\mathrm{SH}^{\mathbb{A}^1}(F)$ then $\mathrm{Re}_B(E)$ is either a sum of suspensions of Morava K -theories or it is contractible. Moreover in the first case, $\mathrm{id}_E \wedge \eta \simeq *$.*

Proof Since $\mathrm{Re}_B(E) \wedge X \simeq \mathrm{Re}_B(E \wedge c^*X)$ and E is a field, $\mathrm{Re}_B(E) \wedge X$ is free over $\mathrm{Re}_B(E)$ for any X in SH . Thus if $\mathrm{Re}_B(E)$ is not contractible, it is a sum of suspensions of Morava K -theories by [16, Proposition 1.9].

Assume the E is not contractible. Then the E -algebra $S_B \mathrm{Re}_B(E)$ is also not contractible. Since it is also free, $P \wedge E$ is a retract of $S_B \mathrm{Re}_B(E)$ for some invertible motivic spectrum P .

Now, via the equivalence of Lemma 6.2, we have a commutative diagram of maps:

$$\begin{array}{ccc}
 S_B \mathrm{Re}_B(E) & \xrightarrow{\mathrm{id} \wedge \eta} & S_B \mathrm{Re}_B(E) \wedge S^{-\alpha} \\
 \searrow S(\mathrm{id} \wedge \eta^{\mathrm{top}}) & & \downarrow \simeq \\
 & & S_B(\mathrm{Re}_B(E) \wedge S^{-1})
 \end{array}$$

Since η^{top} acts by zero on Morava K -theory, it follows that $\mathrm{id}_{S_B \mathrm{Re}_B(E)} \wedge \eta \simeq *$ and hence $\mathrm{id}_{P \wedge E} \wedge \eta \simeq *$ and since P is invertible, $\mathrm{id}_E \wedge \eta \simeq *$. □

Remark 6.6 If F is real closed, then $X_F = *$ and [4] imply $\mathrm{SH}^{\mathbb{A}^1}(F)[[-1]^{-1}] \simeq \mathrm{SH}$, the classical Spanier–Whitehead category. Let $\mathrm{EM}(\underline{W}[\eta^{\pm 1}] \otimes \mathbb{Q})$ denote the Eilenberg–MacLane object associated with the homotopy module given by the rationalization of the Witt sheaf, and note that this object lives in $\mathrm{SH}^{\mathbb{A}^1}(F)[[-1]^{-1}]$ since $(2 + [-1]\eta)\eta = 0$. Under Bachmann’s equivalence, this object is sent to the rational Eilenberg–MacLane spectrum $H\mathbb{Q}$, which is a field in SH . Since every $\mathrm{EM}(\underline{W}[\eta^{\pm 1}] \otimes \mathbb{Q})$ -module is $[-1]$ -periodic, we may conclude that $\mathrm{EM}(\underline{W}[\eta^{\pm 1}] \otimes \mathbb{Q})$ is a motivic field spectrum. Similarly, $\mathrm{EM}(\underline{W}[\eta^{\pm 1}]/p)$ is a motivic field spectrum for all odd primes p . Of course, the complex Betti realizations of such spectra are trivial, but Bachmann’s functor is a real Betti realization, ie the composition of C_2 -equivariant Betti realization with geometric fixed points.

7 Questions

This section contains a list of questions and commentary thereon. It is the authors’ hope that interested readers will take up these problems and advance our collective

understanding of the tensor triangular geometry of stable motivic homotopy theory. In order to speak simultaneously about both compact cellular and all compact spectra, as before, we will write $\mathcal{H}(F)$ to refer to either of $\mathcal{C}(F)$ and $\mathrm{SH}^{\mathbb{A}^1}(F)^c$.

Question 7.1 Is $\mathrm{Spc}(\mathcal{H}(F))$ rigid in the sense that $\mathrm{Spc}(i^*)$ is a homeomorphism whenever $i: F \hookrightarrow F'$ is an extension of algebraically closed or real closed fields?

This question is motivated by the observations that the ring $\mathrm{GW}(F)$ and the space $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$ are rigid amongst these types of extensions. Additionally, the p -complete cellular categories are rigid for algebraically closed fields. Let us write $\mathcal{C}(F)_p^\wedge \subseteq \mathrm{SH}^{\mathbb{A}^1}(F)$ for the thick subcategory generated by $(S^{m\alpha})_p^\wedge$ for $m \in \mathbb{Z}$. Here $E_p^\wedge := L_{S^0/p}(E)$ is the Bousfield localization with respect to the mod- p Moore spectrum.

Proposition 7.2 *Let $i: F \hookrightarrow F'$ be an extension of algebraically closed fields and p a prime different from $\mathrm{char}(F)$. Then $\mathrm{Spc}(\mathcal{C}(F)_p^\wedge) \rightarrow \mathrm{Spc}(\mathcal{C}(F')_p^\wedge)$ is a homeomorphism.*

Proof By [31, Theorem 1.1], $i^*: \mathrm{SH}^{\mathbb{A}^1}(F)_p^\wedge \rightarrow \mathrm{SH}^{\mathbb{A}^1}(F')_p^\wedge$ is full and faithful. On cellular categories it is also essentially surjective and so $\mathcal{C}(F)_p^\wedge \rightarrow \mathcal{C}(F')_p^\wedge$ is an equivalence of categories. □

Write $\mathcal{C}(F)_\mathbb{Q} \subseteq \mathrm{SH}^{\mathbb{A}^1}(F)$ for the thick subcategory generated by $(S^{m\alpha})_\mathbb{Q}$ for $m \in \mathbb{Z}$, and $\mathrm{SH}^{\mathbb{A}^1}(F)_\mathbb{Q}^c$ for the thick subcategory generated by $(X_+)_\mathbb{Q}$, where X is a smooth F -scheme. Work of Cisinski and Déglise [11, Corollary 16.2.14] implies that when F is nonreal, $\mathrm{SH}^{\mathbb{A}^1}(F)_\mathbb{Q}^c$ is equivalent to the triangulated category of rational geometric motives $\mathrm{DM}_{\mathrm{gm}}(F)_\mathbb{Q}$, and similarly that $\mathcal{C}(F)_\mathbb{Q}$ is equivalent to the triangulated category of rational mixed Tate motives, $\mathrm{DMT}(F)_\mathbb{Q}$. Write $\mathcal{H}(F)_\mathbb{Q}$ for either of $\mathcal{C}(F)_\mathbb{Q}$ or $\mathrm{SH}^{\mathbb{A}^1}(F)_\mathbb{Q}^c$.

Question 7.3 Is $\mathrm{Spc}(\mathcal{H}(F)_\mathbb{Q}) = *$ when F is algebraically closed?

Remark 7.4 Peter’s study [29] of the tensor triangular geometry of $\mathrm{DMT}(F)_\mathbb{Q}$ for F an algebraic extension of \mathbb{Q} implies that $\mathrm{Spc}(\mathcal{C}(\overline{\mathbb{Q}})_\mathbb{Q}) = *$, so if Question 7.1 has an affirmative answer, then $\mathrm{Spc}(\mathcal{C}(F)_\mathbb{Q}) = *$ for all algebraically closed fields of characteristic 0. Also note that, over a finite field, assuming the Beilinson–Parshin conjecture and that rational and numerical equivalence on algebraic cycles agree, Kelly [22] has shown that $\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(\mathbb{F}_q)_\mathbb{Q}^c) = *$.

Remark 7.5 Question 7.3 is hard: as remarked to us by Shane Kelly, it implies the conservativity conjecture, which states that Betti realization $DM_{\text{gm}}(\mathbb{C})_{\mathbb{Q}} \rightarrow D(\mathbb{Q})$, on the category of rationalized geometric motives, is a conservative functor (see [2, Conjecture 2.1]), or equivalently, $\text{Re}: SH^{\mathbb{A}^1}(\mathbb{C})_{\mathbb{Q}}^c \rightarrow SH_{\mathbb{Q}}^{\text{fin}}$ is conservative. Indeed, if $\text{Spc}(SH^{\mathbb{A}^1}(\mathbb{C})_{\mathbb{Q}}^c) = *$, then by the classification of thick ideals [5, Theorem 4.10] there is a unique proper thick ideal in $SH^{\mathbb{A}^1}(\mathbb{C})_{\mathbb{Q}}^c$. In particular, the proper thick ideals $\{X \in SH^{\mathbb{A}^1}(\mathbb{C})_{\mathbb{Q}}^c \mid \text{Re}(X) \simeq *\}$ and $\{0\}$ are equal.

In the other direction, we have the following proposition. Recall that a topological space is *local* if it has a unique closed point.

Proposition 7.6 *If the conservativity conjecture holds, then $\text{Spc}(\mathcal{K}(\mathbb{C})_{\mathbb{Q}})$ is local.*

Proof The tensor triangulated category $\mathcal{K}(\mathbb{C})_{\mathbb{Q}}$ is rigid in the sense that every object is dualizable. By [7, Proposition 4.2] these spaces are local if and only if $X \wedge Y \simeq *$ implies $X \simeq *$ or $Y \simeq *$. If $X \wedge Y \simeq *$, then $\text{Re}(X \wedge Y) \simeq \text{Re}(X) \wedge \text{Re}(Y) \simeq *$. But since these are rational spectra, it follows that either $\text{Re}(X) \simeq *$ or $\text{Re}(Y) \simeq *$. Since $SH^{\mathbb{A}^1}(\mathbb{C})_{\mathbb{Q}}^c$ is equivalent to $DM_{\text{gm}}(\mathbb{C})_{\mathbb{Q}}$ (see [11, Corollary 16.2.14]), the Conservativity Conjecture implies that either $X \simeq *$ or $Y \simeq *$. □

Remark 7.7 More generally, Balmer has shown in [9, Theorem 1.2] that if $f: \mathcal{K} \rightarrow \mathcal{L}$ is a tensor triangulated functor between essentially small, tensor triangulated categories, where \mathcal{K} is rigid, then f is conservative if and only if the induced map $\text{Spc}(f): \text{Spc}(\mathcal{L}) \rightarrow \text{Spc}(\mathcal{K})$ is surjective on closed points. This is consistent with Remark 7.5 and Proposition 7.6 because $\text{Spc}(SH_{\mathbb{Q}}^{\text{fin}}) = *$.

Remark 7.8 When F is perfect of finite 2-étale cohomological dimension, Bachmann has shown in [3, Theorem 15] that the functor $M: SH^{\mathbb{A}^1}(F)^c \rightarrow DM_{\text{gm}}(F)$ is conservative. By [9, Theorem 1.2], we see that this is equivalent to $\text{Spc}(M)$ being surjective on closed points. Note, though, that $\text{Spc}(M)$ is not surjective, as its image does not intersect $(\rho^{\bullet})^{-1}([F^{\times}], 2)$. Further study of $\text{Spc}(DM_{\text{gm}}(F))$ and its relationship with $\text{Spc}(SH^{\mathbb{A}^1}(F)^c)$ would be quite interesting.

The following question speculates on the fashion in which $\text{Spc}(\mathcal{K}(F))$ might be reconstructed from the tensor triangular spectra of the stable motivic homotopy categories of the real and algebraic closures of F .

Question 7.9 Let X_F denote the set of orderings on F , and let $X_F^* = X_F \sqcup \{\infty\}$. For $\alpha \in X_F$ let F_α denote the real closure of F with respect to α ; let $F_\infty = \bar{F}$ denote the algebraic closure of F . For $\alpha \in X_F^*$ let $i_\alpha: F \hookrightarrow F_\alpha$ be a chosen embedding. Each of these induces a map $\mathrm{Spc}(i_\alpha^*): \mathrm{Spc}(\mathcal{H}(F_\alpha)) \rightarrow \mathrm{Spc}(\mathcal{H}(F))$. Is the map

$$\coprod_{\alpha \in X_F^*} \mathrm{Spc}(\mathcal{H}(F_\alpha)) \rightarrow \mathrm{Spc}(\mathcal{H}(F))$$

given by the sum of these maps surjective? Can we explicitly describe $\mathrm{Spc}(\mathcal{H}(F))$ as a quotient of $\coprod_{\alpha \in X_F^*} \mathrm{Spc}(\mathcal{H}(F_\alpha))$?

Note that the analogue of the surjectivity question on $\mathrm{Spec}^h(K_*^{\mathrm{MW}}(F))$ is true by Thornton's theorem. If we could produce the quotient description and answer Question 7.1 in the affirmative, this would give a robust description of $\mathrm{Spc}(\mathcal{H}(F))$.

Question 7.10 What is the structure of $\mathrm{Spc}(\mathcal{H}(\mathbb{C}))$? Of $\mathrm{Spc}(\mathcal{H}(\mathbb{R}))$?

While simply stated, the authors believe that this may be the hardest of the questions. (See Remark 5.7.) Were one to successfully tackle this problem and affirmatively answer Questions 7.1 and 7.9, one would completely describe $\mathrm{Spc}(\mathcal{H}(F))$, at least for characteristic 0 fields.

References

- [1] **M Andrews**, *A non-nilpotent self map of S^0/η over \mathbb{C}* , preprint (2014) Available at <http://www.math.ucla.edu/~mjandr/>
- [2] **J Ayoub**, *Motives and algebraic cycles: a selection of conjectures and open questions*, from “Hodge theory and L^2 -analysis” (L Ji, editor), International Press, Somerville, MA (2017) 87–125
- [3] **T Bachmann**, *On the conservativity of the functor assigning to a motivic spectrum its motive*, preprint (2015) arXiv
- [4] **T Bachmann**, *Motivic and real étale stable homotopy theory*, preprint (2016) arXiv
- [5] **P Balmer**, *The spectrum of prime ideals in tensor triangulated categories*, J. Reine Angew. Math. 588 (2005) 149–168 MR
- [6] **P Balmer**, *Supports and filtrations in algebraic geometry and modular representation theory*, Amer. J. Math. 129 (2007) 1227–1250 MR
- [7] **P Balmer**, *Spectra, spectra, spectra — Tensor triangular spectra versus Zariski spectra of endomorphism rings*, Algebr. Geom. Topol. 10 (2010) 1521–1563 MR

- [8] **P Balmer**, *Tensor triangular geometry*, from “Proceedings of the International Congress of Mathematicians, II” (R Bhatia, A Pal, G Rangarajan, V Srinivas, M Vanninathan, editors), Hindustan Book Agency, New Delhi (2010) 85–112 MR
- [9] **P Balmer**, *On the surjectivity of the map of spectra associated to a tensor-triangulated functor*, preprint (2017) arXiv
- [10] **P Balmer**, **B Sanders**, *The spectrum of the equivariant stable homotopy category of a finite group*, *Invent. Math.* 208 (2017) 283–326 MR
- [11] **D-C Cisinski**, **F Déglise**, *Triangulated categories of mixed motives*, preprint (2009) arXiv
- [12] **D Dugger**, **D C Isaksen**, *Motivic cell structures*, *Algebr. Geom. Topol.* 5 (2005) 615–652 MR
- [13] **H Fausk**, **P Hu**, **J P May**, *Isomorphisms between left and right adjoints*, *Theory Appl. Categ.* 11 (2003) No. 4, 107–131 MR
- [14] **J J Gutiérrez**, **O Röndigs**, **M Spitzweck**, **P A Østvær**, *Motivic slices and coloured operads*, *J. Topol.* 5 (2012) 727–755 MR
- [15] **J Heller**, **K Ormsby**, *Galois equivariance and stable motivic homotopy theory*, *Trans. Amer. Math. Soc.* 368 (2016) 8047–8077 MR
- [16] **M J Hopkins**, **J H Smith**, *Nilpotence and stable homotopy theory, II*, *Ann. of Math.* 148 (1998) 1–49 MR
- [17] **J Hornbostel**, *A^1 -representability of hermitian K -theory and Witt groups*, *Topology* 44 (2005) 661–687 MR
- [18] **M Hovey**, **J H Palmieri**, **N P Strickland**, *Axiomatic stable homotopy theory*, *Mem. Amer. Math. Soc.* 610, Amer. Math. Soc., Providence, RI (1997) MR
- [19] **P Hu**, *On the Picard group of the stable \mathbb{A}^1 -homotopy category*, *Topology* 44 (2005) 609–640 MR
- [20] **J F Jardine**, *Motivic symmetric spectra*, *Doc. Math.* 5 (2000) 445–552 MR
- [21] **R Joachimi**, *Thick ideals in equivariant and motivic stable homotopy categories*, preprint (2015) arXiv
- [22] **S Kelly**, *Some observations about motivic tensor triangulated geometry over a finite field*, preprint (2016) arXiv
- [23] **M Levine**, **Y Yang**, **G Zhao**, *Algebraic elliptic cohomology theory and flops, I*, preprint (2013) arXiv
- [24] **F Lorenz**, **J Leicht**, *Die Primideale des Wittschen Ringes*, *Invent. Math.* 10 (1970) 82–88 MR
- [25] **J Milnor**, *Algebraic K -theory and quadratic forms*, *Invent. Math.* 9 (1969/1970) 318–344 MR

- [26] **F Morel**, *An introduction to \mathbb{A}^1 -homotopy theory*, from “Contemporary developments in algebraic K -theory” (M Karoubi, A O Kuku, C Pedrini, editors), ICTP Lect. Notes 15, Abdus Salam Int. Cent. Theoret. Phys., Trieste (2004) 357–441 MR
- [27] **F Morel**, *On the motivic π_0 of the sphere spectrum*, from “Axiomatic, enriched and motivic homotopy theory” (J P C Greenlees, editor), NATO Sci. Ser. II Math. Phys. Chem. 131, Kluwer, Dordrecht (2004) 219–260 MR
- [28] **F Morel**, *\mathbb{A}^1 -algebraic topology over a field*, Lecture Notes in Mathematics 2052, Springer (2012) MR
- [29] **T J Peter**, *Prime ideals of mixed Artin–Tate motives*, J. K-Theory 11 (2013) 331–349 MR
- [30] **O Röndigs, P A Østvær**, *Modules over motivic cohomology*, Adv. Math. 219 (2008) 689–727 MR
- [31] **O Röndigs, P A Østvær**, *Rigidity in motivic homotopy theory*, Math. Ann. 341 (2008) 651–675 MR
- [32] **R Thornton**, *The homogeneous spectrum of Milnor–Witt K -theory*, J. Algebra 459 (2016) 376–388 MR
- [33] **V Voevodsky**, *\mathbb{A}^1 -homotopy theory*, Doc. Math. Extra Vol. ICM I (1998) 579–604 MR
- [34] **V Voevodsky**, *Motivic cohomology with $\mathbf{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. 98 (2003) 59–104 MR

*Department of Mathematics, University of Illinois at Urbana-Champaign
Urbana, IL, United States*

*Department of Mathematics, Reed College
Portland, OR, United States*

jbheller@illinois.edu, ormsbyk@reed.edu

Proposed: Stefan Schwede
Seconded: Mark Behrens, Haynes R Miller

Received: 19 August 2016
Revised: 19 July 2017