

An application of the Duistermaat–Heckman theorem and its extensions in Sasaki geometry

CHARLES P BOYER
HONGNIAN HUANG
EVELINE LEGENDRE

Building on an idea laid out by Martelli, Sparks and Yau (2008), we use the Duistermaat–Heckman localization formula and an extension of it to give rational and explicit expressions of the volume, the total transversal scalar curvature and the Einstein–Hilbert functional, seen as functionals on the Sasaki cone (Reeb cone). Studying the leading terms, we prove they are all proper. Among consequences thereof we get that the Einstein–Hilbert functional attains its minimal value and each Sasaki cone possesses at least one Reeb vector field with vanishing transverse Futaki invariant.

53B99, 53CXX, 53DXX

1 Introduction

The general problem motivating our work is: given a polarized Sasaki type manifold (N^{2n+1}, ξ) , does there exist a compatible constant scalar curvature Sasaki (cscS for short) metric? This is a hard problem and the answer is conjecturally related to some notion of K–stability — see Collins and Székelyhidi [12] — and is closely related to the analogous problem in (compact) Kähler geometry; see eg Donaldson [14; 15], Ross and Thomas [30] and Stoppa [31]. Well-known obstructions are the K–stability — see [12] — the transversal Futaki invariant — see Futaki, Ono and Wang [17] — and the Einstein–Hilbert functional — see Legendre [23] and Boyer, Huang, Legendre and Tønnesen-Friedman [9]. In this paper we study the latter using the Duistermaat–Heckman localization formula, developing an idea of Martelli, Sparks and Yau [27]. More precisely, given a Sasaki manifold $(N, \mathcal{D}, J, \xi_o)$ with maximal torus of automorphisms $T \subset \text{CR}$, the (reduced) Sasaki cone (or Reeb cone), denoted by \mathfrak{t}^+ , is an open polyhedral cone in $\mathfrak{t} = \text{Lie}(T)$ and contains all the T –invariant Reeb vector fields on (N, \mathcal{D}, J) . Picking a quasiregular Reeb vector field $\xi_o \in \mathfrak{t}^+$, that is, it induces a circle action $S_o^1 \subset T$, we build an orbundle $\pi: \mathcal{L} \rightarrow W$ polarizing

the quotient symplectic orbifold $(W := N/S^1_0, \sigma)$ so that there is a biholomorphism between $\mathcal{L} \setminus W$ and the Kähler cone $Y_0 := N \times \mathbb{R}$ over N . The action of T is defined on L and the fixed-point set $Z = \bigsqcup Z$ consists in a disjoint union of symplectic suborbifolds of (W, σ) . Moreover, the normal bundle of Z in M is T -invariant and splits with respect to the action into (real) rank 2 symplectic bundles $\bigoplus_{i=0}^{n-n_Z} E_i^Z$ on which T acts with weight $\kappa_i^Z \in \mathfrak{t}^*$ (where $\dim Z = 2n_Z$). We recall and explain these facts in Section 4.1 and we then we use it to prove that

$$(1) \quad V_\xi = \text{vol}(N, \xi) = \frac{(2\pi)^{n+1}}{n!} \sum_Z \frac{1}{d_Z} \int_Z \frac{1}{\prod_{j=0}^{n-n_Z} (c_1(E_j^Z) - \langle \kappa_j^Z, \xi \rangle)}.$$

In [27], Martelli, Sparks and Yau proved that the volume functional V is a rational function of $\xi \in \mathfrak{t}^+$ when the Kähler cone is Gorenstein. This hypothesis is not needed to get (1) and, moreover, by studying the leading term in (1) we get the following corollary:

Theorem 1.1 *Let (N, \mathcal{D}) be a compact contact manifold of Sasaki type of dimension $2n + 1$ and \mathfrak{t}^+ a compatible Sasaki cone. The volume functional $V: \mathfrak{t}^+ \rightarrow \mathbb{R}$ is a rational function homogenous of order $-(n + 1)$. Moreover, V tends to $+\infty$ when ξ approaches the boundary of \mathfrak{t}^+ .*

We also derive a similar formula for the *total transversal scalar curvature* S_ξ using the same orbundle $\mathcal{L} \rightarrow W$ and an extension — see Theorem 3.1 — of the Duistermaat–Heckman localisation formula. The formula is

$$(2) \quad S_\xi = \frac{2(2\pi)^{n+1}}{(n-1)!} \sum_Z \frac{1}{d_Z} \int_Z \frac{(\iota_Z^* c_1(W) + \sum_{i=1}^{n-n_Z} \langle \kappa_i^Z, \xi \rangle)}{\prod_{j=0}^{n-n_Z} (c_1(E_j^Z) - \langle \kappa_j^Z, \xi \rangle)}.$$

As a direct consequence we get the following theorem:

Theorem 1.2 *Let (N, \mathcal{D}) be a compact contact manifold of Sasaki type of dimension $2n + 1$ with a fixed class of compatible CR-structures $[J]$. Let \mathfrak{t}^+ be a compatible Sasaki cone. The total transversal scalar curvature functional $S: \mathfrak{t}^+ \rightarrow \mathbb{R}$ is a rational function homogenous of order $-n$.*

Remark 1.3 Tian used the Duistermaat–Heckman localization formula to compute the Futaki invariant of some Kähler manifolds in [32]. Using the relation found in [23; 9], Tian’s formula should be related to the derivative of (2) in the regular case.

Remark 1.4 In the toric case the fixed-point set consists in a set of isolated points. The formula (2) becomes

$$(3) \quad S_\xi = \frac{2(2\pi)^{n+1}}{(n-1)!} \sum_p \frac{1}{d_Z} \frac{\sum_{i=1}^n \langle \kappa_i^p, \xi \rangle}{\prod_{j=0}^n \langle \kappa_j^p, \xi \rangle},$$

which coincides with the one found in [23] using integration by parts.

The claims of Theorems 1.1 and 1.2 may also be obtained by studying the Hilbert series of the polarized Kähler cone associated to (N, \mathcal{D}, J, ξ) — see [12] — as pointed out to us by Tristan Collins.

Now, by analyzing the weights of the action of T on \mathcal{L} , using the construction and Morse’s lemma we get that the leading term, when $\xi \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ , of the right-hand side of (2) is positive. From this we derive the following result:

Corollary 1.5 *When $\xi \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ , S_ξ tends to $+\infty$.*

The Einstein–Hilbert functional is defined here to be the homogenous functional

$$(4) \quad H(\xi) = \frac{S_\xi^{n+1}}{V_\xi^n}$$

on the (reduced) Sasaki cone \mathfrak{t}^+ .

In [23; 9] it is shown that this functional detects the Reeb vector field whose transversal Futaki invariant vanishes. That is, fixing the isotopy class of (\mathcal{D}, J) we have

$$\{\xi \in \mathfrak{t} \mid \exists \text{ compatible cscS} \} \subset \text{crit}(H).$$

This is our main motivation to study this functional and, in particular, its critical points. Nonuniqueness is now known, indeed H is not convex and may possess many critical points; see Boyer, Huang, Legendre and Tønnesen-Friedman [9; 11; 23]. In this paper we solve the question of existence by proving that H is proper on the convex (relatively compact) set of rays of the Sasaki cone.

Corollary 1.6 *The Einstein–Hilbert functional is an homogenous rational function on \mathfrak{t}^+ , tending to $+\infty$ when $\xi \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ . In particular, it attains its minimal value along a ray in the interior of the cone \mathfrak{t}^+ .*

Corollary 1.7 *Any Sasaki cone contains rays of Reeb vector fields with vanishing transversal Futaki invariant. Moreover, any ray of Reeb vector fields with vanishing transversal Futaki invariant, in particular a ray of cscS structures, is isolated in any 2–dimensional subcone of the Sasaki cone.*

To obtain formula (2) we relate the total scalar curvature of the metric associated to ξ to the metric associated to ξ_o on W in order to get something independent. The relation we found is quite explicit — see (32) — and we get the following corollary:

Corollary 1.8 *Within a Sasaki cone there is at most one ray of vanishing transverse scalar curvature. Moreover, if there is one Sasaki structure with nonnegative transverse scalar curvature then the total transverse scalar curvature is nonnegative on the whole Sasaki cone.*

The next section gathers facts and conventions we used in this paper. We present in Section 3 the localization formula of Duistermaat–Heckman and the extension we need. In Section 4.1 we build the orbiresolution of the cone we will use in the subsequent sections to prove Theorem 1.1 in Section 4.2 and Theorem 1.2 in Section 4.3. The final section first gathers the proof of the Corollaries 1.6–1.8. Then we discuss how the positivity (or nonnegativity) of \mathcal{S}_ξ affects the transverse geometry, namely that in a certain sense the transverse structure is dominated by rational curves. Although the nonnegativity of \mathcal{S}_ξ is sufficient for this, it is not necessary.

Acknowledgements Legendre would like to thank her colleagues J-F Barraud, P Gauduchon and R Leclercq for illuminating discussions on symplectic manifolds and equivariant cohomology. We also thank T Collins, J Sparks and S Sun for their comments on a previous version of this paper and their interest in this work.

Boyer was partially supported by grant #245002 from the Simons Foundation. Legendre is partially supported by France ANR project EMARKS No. ANR-14-CE25-0010.

2 Some facts and conventions in Sasaki geometry

2.1 Basics of Sasaki geometry

We give in this section the basic definitions and facts we need for our purposes. We refer to [7] for an extensive study of Sasakian geometry.

2.1.1 Sasaki cone A Sasakian manifold of dimension $2n + 1$ is a smooth manifold N endowed with the following structures:

- a cooriented rank $2n$ contact distribution \mathcal{D} ;
- a CR–structure $J \in \text{End}(\mathcal{D})$ (ie $J^2 = -\text{id}_{\mathcal{D}}$ + integrability condition);
- a nowhere-vanishing contact form $\eta \in \Gamma(\mathcal{D}^0)$, where $\mathcal{D}^0 := \text{Annihilator}(\mathcal{D}) \subset T^*N$;
- a Riemannian metric $g = \eta^2 + \frac{1}{2}d\eta(\cdot, J\cdot)$;
- a Reeb vector field ξ lying in the Lie algebra of CR–diffeomorphisms $\mathfrak{cr}(\mathcal{D}, J)$ and satisfying the conditions $\eta(\xi) = 1$ and $\mathcal{L}_{\xi}\eta = 0$.

Any three of these structures determine the remaining ones. For example, given (\mathcal{D}, J, η) we get ξ by solving the equations involved, then $\Phi \in \Gamma(\text{End}(\text{TN}))$, defined as $\Phi(\xi) = 0$ and $\Phi|_{\mathcal{D}} = J$ and finally

$$g = \frac{1}{2}d\eta(\cdot, \Phi(\cdot)) + \eta \otimes \eta.$$

Therefore, the space of Sasaki structures sharing the same CR–structure, denoted by $\text{Sas}(\mathcal{D}, J)$, is in bijection with the cone of Reeb vector fields

$$(5) \quad \mathfrak{cr}^+(\mathcal{D}, J) = \{X \in \mathfrak{cr}(\mathcal{D}, J) \mid \eta(X) > 0\},$$

where η is any fixed, nowhere-vanishing, section of \mathcal{D}^0 . The map $\mathfrak{cr}^+(\mathcal{D}, J) \rightarrow \text{Sas}(\mathcal{D}, J)$ is given by

$$(6) \quad \xi' \mapsto \left(\frac{\eta}{\eta(\xi')}, \mathcal{D}, J \right).$$

To avoid confusion, we sometimes refer to a Sasakian structure redundantly by specifying all the structures involved like $(N, \mathcal{D}, J, \eta, g, \xi)$. From [7] we know that $\mathfrak{cr}^+(\mathcal{D}, J)$ is an open convex cone in $\mathfrak{cr}(\mathcal{D}, J)$, invariant under the adjoint action of $\mathfrak{CR}(\mathcal{D}, J)$. Moreover, the following result will be useful for our study:

Theorem 2.1 [8] *Let N be a compact manifold of dimension $2n + 1$ with a CR–structure (\mathcal{D}, J) of Sasaki type. Then the Lie algebra $\mathfrak{cr}(\mathcal{D}, J)$ decomposes as $\mathfrak{cr}(\mathcal{D}, J) = \mathfrak{t} + \mathfrak{p}$, where \mathfrak{t} is the Lie algebra of a maximal torus \underline{T} of dimension k with $1 \leq k \leq n + 1$ and \mathfrak{p} is a completely reducible \mathfrak{t} –module. Furthermore, every $X \in \mathfrak{cr}^+(\mathcal{D}, J)$ is conjugate to a positive element in the Lie algebra \mathfrak{t} .*

The (reduced) *Sasaki cone* is the set $\underline{t}^+ = \underline{t} \cap \text{cr}^+(\mathcal{D}, J)$. When seeking extremal or csc Sasaki metrics one deforms the contact structure by $\eta \mapsto \eta + d^c\varphi$, where the function φ is invariant under the maximal torus \underline{T} . Thus, the Sasaki cone is associated with an isotopy class of contact structures of Sasaki type that is invariant under \underline{T} . This fact has given rise to several equivalent definitions of \underline{t}^+ — see [20; 12; 6] — and is also called the *Reeb cone*.

2.1.2 Kähler cones Recall — see eg [24] — that cooriented contact manifolds are in one-to-one correspondence with symplectic cones. Given a cooriented contact manifold (N, \mathcal{D}) the symplectic cone over N is given by $(\mathcal{D}_+^0, \omega)$, where \mathcal{D}_+^0 is a connected component of the annihilator of \mathcal{D}_+^0 devoid of the 0–section and ω is the pullback of the Liouville symplectic form of T^*N via the inclusion $\mathcal{D}_+^0 \subset T^*N$. There is a natural projection $\pi: \mathcal{D}_+^0 \rightarrow N$, so that \mathcal{D}_+^0 is an \mathbb{R}^+ –bundle over N . The fibers coincide with the orbits of the vector field, $\tau \in \Gamma(T\mathcal{D}_+^0)$, induced by the \mathbb{R}^+ –action (ie multiplication) along the fibers, so that $(\mathcal{D}_+^0, \omega)$ is a symplectic cone in the sense that $\mathcal{L}_\tau\omega = \omega$. Conversely, given a symplectic cone $(M, \omega, \mathbb{R}^+, \tau)$, then $N := M/\mathbb{R}^+$ and $\mathcal{D} := \ker(\tau \lrcorner \omega)$. Note that any contact 1–form η is a section of \mathcal{D}_+^0 trivializing it as $\mathcal{D}_+^0 = N \times \mathbb{R}^+$.

Similarly, Sasakian manifolds are in one-to-one correspondence with Kähler cones. Any Sasakian structure (g, J, Φ, η, ξ) on (N, \mathcal{D}) provides a Kählerian structure (\hat{g}, \hat{J}) on $(\mathcal{D}_+^0, \omega)$ as follows:

- the metric \hat{g} is the unique Riemannian metric which is conic (that is $\mathcal{L}_\tau\hat{g} = \hat{g}$) and restricts to g on the submanifold $\eta(N) \subset \mathcal{D}_+^0$;
- the complex structure \hat{J} is the unique complex structure which is homogenous of degree 0 with respect to τ (ie $\mathcal{L}_\tau\hat{J} = 0$) and restricts to Φ on $\eta(N)$.

Convention 2.2 Given a manifold N we will let $Y := N \times \mathbb{R}^+$ be the topological cone, $\iota_N: N \rightarrow N \times \{1\}$ the inclusion and $r: Y \rightarrow \mathbb{R}^+$ the projection on the second factor. Any contact 1–form η on N gives rise to a symplectic structure

$$\omega := \frac{1}{2}d(r^2\eta)$$

(so that $\iota_N^*\omega = \frac{1}{2}d\eta$) and any compatible Sasaki structure (J, ξ) gives a compatible Kähler structure on Y as

$$\hat{g} := dr \otimes dr + r^2g, \quad \hat{J} := J - \xi \otimes \frac{dr}{r} + r \frac{\partial}{\partial r} \otimes \eta.$$

Then, by uniqueness, $(Y, \omega, \hat{g}, \hat{J})$ is the Kähler cone over $(N, \mathcal{D}, J, g, \eta, \xi)$. The compatible Kähler structure on Y is then determined by a choice Reeb vector field $\xi \in \mathfrak{t}^+$, giving rise to a polarized affine variety (Y, ξ) .

2.2 The transverse geometry

Most of the important structure of Sasakian geometry arises from its transverse Kählerian structure.

2.2.1 The transverse structure One can consider the class $\text{Sas}(\xi)$ of Sasakian structures having the same Reeb vector field. Let L_ξ be the line bundle having ξ as a section. The inclusion $L_\xi \hookrightarrow TN$ induces a sequence of bundle morphisms

$$0 \rightarrow L_\xi \hookrightarrow TN \rightarrow Q_\xi \rightarrow 0.$$

For any CR–structure (\mathcal{D}, J) in $\text{Sas}(\xi)$, the restriction followed by projection $\mathcal{D} \rightarrow Q_\xi$ is an isomorphism and provides a complex structure \bar{J} on Q_ξ . One can consider the subclass of structures $\text{Sas}(\xi, \bar{J})$ making the following diagram commute:

$$(7) \quad \begin{array}{ccc} TN & \longrightarrow & Q_\xi \\ \Phi \downarrow & & \bar{J} \downarrow \\ TN & \longrightarrow & Q_\xi \end{array}$$

With that comes a natural notion of transversal holomorphic vector fields $\mathfrak{h}(\xi, \bar{J})$; see [8].

For a given Sasakian manifold, $(N, g, \xi, \mathcal{D}, J)$, the transverse Kähler geometry refers to the geometry of $(\mathcal{D}, J, g|_{\mathcal{D}})$. More precisely, N is foliated by the Reeb flow. So there are local submersions $\pi_\alpha: U_\alpha \rightarrow V_\alpha$, where U_α and V_α are open subsets of N and \mathbb{C}^n , respectively, such that $\pi_\alpha^* I = \Phi$, where I is the standard complex structure on \mathbb{C}^n . In particular, $d\pi_\alpha: (\mathcal{D}|_{U_\alpha}, J) \rightarrow (TV_\alpha, I)$ is an isomorphism and the Sasaki metric is sent to a Kähler structure $(g_\alpha, \sigma_\alpha, I)$ on V_α . In the case where ξ is quasiregular — equivalently, it is induced by the action of a closed 1–dimensional subtorus $\underline{S}_\xi^1 \subset \underline{T}$ — the open sets and maps $\pi_\alpha: U_\alpha \rightarrow V_\alpha$ patch up as a global orbifold quotient

$$(8) \quad \pi: N \rightarrow W := N/\underline{S}_\xi^1$$

endowed with the unique symplectic structure σ satisfying $\pi^* \sigma = \frac{1}{2} d\eta$.

2.2.2 Transverse automorphisms

Convention 2.3 We will adopt the following point of view, brought from toric geometry. Instead of working with $\underline{T} \subset \text{CR}(\mathcal{D}, J)$ a Lie group of diffeomorphism of N and \mathfrak{t} its Lie algebra of vector fields on N , we consider a torus $T = \mathfrak{t}/\Lambda$, where \mathfrak{t} is the Lie algebra and $\Lambda \subset \mathfrak{t}$ a lattice and an injective morphism $\phi: T \hookrightarrow \text{CR}(\mathcal{D}, J)$. Given $a \in \mathfrak{t}$, the vector field induced on N is $X_a = \phi_*(a)$ or, equivalently, is given at $p \in N$ by

$$X_a(p) = \left(\frac{d}{dt} \exp(ta) \cdot p \right)_{t=0},$$

where $\exp: \mathfrak{t} \rightarrow T$ is the quotient map. By maximality, a Reeb vector field ξ is necessarily induced from the action of T and so there exists $b \in \mathfrak{t}$ such that $\xi = X_b$. We write $\mathfrak{t}^+ = \phi^{-1}(\mathfrak{t}^+)$ and call it the (reduced) Sasaki cone as well.

Let T be a maximal torus embedded in $\text{CR}(\mathcal{D}, J)$ with Lie algebra \mathfrak{t} . For any T -invariant compatible pair (ξ, η) (ie such that $(N, g, \xi, \mathcal{D}, J)$ is a Sasakian manifold with $T \xrightarrow{\phi} \text{Isom}(N, g)$) there is a η -momentum map

$$\mu_\eta: N \rightarrow \mathfrak{t}^*$$

in the sense that for any $a \in \mathfrak{t}$, we have

$$-\frac{1}{2}d\eta(X_a, \cdot) = d\langle \mu_\eta, a \rangle.$$

Such a map always exists since one can take $\langle \mu_\eta, a \rangle = \frac{1}{2}\eta(X_a)$ (this is the one we pick in this paper). The image of μ_η lies in the affine subspace

$$\mathcal{H}_b = \{x \in \mathfrak{t}^* \mid \langle x, b \rangle = \frac{1}{2}\},$$

where $b \in \mathfrak{t}^+ \subset \mathfrak{t}$ is such that $X_b = \xi$ is the Reeb vector field of η . Such an element b exists by the maximality of T .

We can consider the infinitesimal \mathfrak{t} -action on the open subset U_α . Since ξ is induced by the action of T , this action descends as an infinitesimal action of $\mathfrak{t}/\mathbb{R}b$ on V_α . It is easy to see that the resulting action should be Hamiltonian. However, the $(\mathfrak{t}/\mathbb{R}b)$ -momentum map is not the map $\bar{\mu}_\eta$ making the following diagram commute:

$$(9) \quad \begin{array}{ccc} U_\alpha & \xrightarrow{\mu_\eta} & \mathfrak{t}^* \\ \pi_\alpha \downarrow & & \downarrow \text{id} \\ V_\alpha & \xrightarrow{\bar{\mu}_\eta} & \mathfrak{t}^* \end{array}$$

Indeed, the image of this one is not in $(\mathfrak{t}/\mathbb{R}b)^*$ (ie $\langle \bar{\mu}_\eta, a + tb \rangle \neq \langle \bar{\mu}_\eta, a \rangle$ in general). The momentum map of the $(\mathfrak{t}/\mathbb{R}b)$ -action on $(V_\alpha, \sigma_\alpha)$ can be taken to be

$$(10) \quad \check{\mu}_\eta = \bar{\mu}_\eta - x_o$$

for any $x_o \in \mathcal{H}_b$. Therefore, for any class $[a] \in \mathfrak{t}/\mathbb{R}b$,

$$\langle \check{\mu}_\eta, [a] \rangle = \frac{1}{2}\eta(X_a) - \langle x_o, a \rangle.$$

Remark 2.4 As a direct consequence of this discussion, for any $a \in \mathfrak{t}^*$, we can see $\eta(X_a)$ as a function on V_α and, more precisely, we can identify \mathfrak{t} with the set of affine-linear functions on the linear space $\mathcal{H}_b - \{x_o\} \simeq (\mathfrak{t}/\mathbb{R}b)^*$, in the sense that $\frac{1}{2}\eta(X_a) = \langle \check{\mu}_\eta, [a] \rangle + \langle x_o, a \rangle$, where the second term is a constant.

Remark 2.5 If ξ is quasiregular, it induces an S^1 -action, corresponding to a subgroup $S^1_\xi \subset T$, and we have a global symplectic orbifold quotient $(W = N/S^1_\xi, \sigma)$. The image of the momentum map $\Delta_b = \text{Im}\check{\mu}_\eta \subset \mathcal{H}_b - \{x_o\}$ is a compact polytope. Then, the Sasaki cone can be identified with the affine-linear functions on $\mathcal{H}_b - \{x_o\} \simeq (\text{Lie}(T/S^1_\xi))^*$ which are positive on Δ_b ; see [23].

2.2.3 Transverse curvatures The Kähler structure $(g_\alpha, \sigma_\alpha, I)$ on V_α has a connection ∇^T_α and curvatures $R^T_\alpha, \text{Ric}^T_\alpha, \rho^T_\alpha, s^T_\alpha, \dots$. Since $\pi_\alpha^*\nabla^T_\alpha$ and $\pi_\beta^*\nabla^T_\beta$ coincide on $U_\alpha \cap U_\beta$, these objects patch together to define global objects on N , the *transversal* connection and curvatures $\nabla^T, R^T, \text{Ric}^T, \rho^T, s^T, \dots$. See [7; 17] for more details. These tensors are basic, notably the transversal Ricci form ρ^T satisfies

$$\rho^T(\xi, \cdot) = 0, \quad \mathcal{L}_\xi \rho^T = 0$$

and lies in the basic first Chern class $2\pi c^B_1(\mathcal{F}_\xi)$.

Since the exterior derivative preserves this condition, the graded algebra of basic forms is a subcomplex of the de Rham complex. Moreover, one can define the basic exterior derivative d_B as the restriction of the differential to these forms; its adjoint is δ_B and the basic Laplacian is $\Delta_B = d_B\delta_B + \delta_B d_B$. The Hodge theorem holds for the basic cohomology in this context; for a Sasaki metric g there exists a unique basic function ψ_g (of mean value 0) such that

$$\rho^T = \rho^T_H + i\partial\bar{\partial}\psi_g,$$

where ρ_H^T is Δ_B -harmonic. Note that

$$\Delta_B \psi_g = s^T - \frac{S_\xi}{V_\xi} = s^T - \bar{s}^T,$$

where V_ξ is the volume of (N, g) with volume form

$$dv_\xi = \frac{\eta \wedge (d\eta)^n}{n!},$$

S_ξ is the total transversal scalar curvature, that is

$$(11) \quad S_\xi := \int_N s^T dv_\xi,$$

and \bar{s}^T is the average transverse scalar curvature. The volume of the Sasakian manifold (N, \mathcal{D}, J, ξ) does not depend on the chosen structure in $\text{Sas}(\xi)$ — see [7] — and the total transversal scalar curvature does not depend on the chosen structure in $\text{Sas}(\xi, \bar{J})$; see [17].

3 The Duistermaat–Heckman localisation formula and extensions

Let (M, ω) be a symplectic compact manifold admitting a Hamiltonian action of S^1 generating the (Hamiltonian) vector field $X \in \Gamma(TM)$ associated to $H: M \rightarrow \mathbb{R}$. In this section, to fit with [28], we take the convention that

$$(12) \quad X \lrcorner \omega = dH.$$

Let $Z := \text{Fix } S^1$ be the fixed-point set of S^1 . It is known that Z consists in a union of smooth symplectic submanifolds $Z = \bigsqcup_l Z_l$, so the normal bundle E_Z of any component, say $Z \subset Z$, bears a symplectic structure as well and splits into a sum of rank 2 bundles $E_Z = \bigoplus_j E_j^Z$ according to the action of S^1 , acting on E_j^Z with weight κ_j^Z ; here and after j runs from 1 to $n - m_Z$, where $2m_Z = \dim Z$. The Duistermaat–Heckman theorem [16] says that, under these conditions, we have

$$(13) \quad \int_M e^{\omega - H} = \sum_Z e^{-H(Z)} \int_Z \prod_j \frac{e^{\iota_Z^* \omega}}{2\pi c_1(E_j^Z) + \kappa_j^Z},$$

where, in the right-hand side, Z denotes a generic connected component of Z , $H(Z)$ the (single) value H is taking on that component and $\iota_Z: Z \hookrightarrow M$ the inclusion. For any form $\psi \in \Omega(M)$, we write $e^\psi = \sum_{k \geq 0} \psi^k / k!$.

In the simpler case where H is Morse — equivalently, Z is a collection of isolated points — the formula reads

$$(14) \quad \int_M e^{\omega-H} = \sum_p \frac{e^{-H(p)}}{e(p)},$$

where $e(p)$ is the product of the weights at $p \in Z$.

Here, we will use the following extension:

Theorem 3.1 *Let $\phi \in \Omega_{S^1}^2(M)$ be a closed form and $f \in C^\infty(M)$ be such that $X \lrcorner \phi = df$; then*

$$(15) \quad \int_M (\phi - f)e^{\omega-H} = \sum_Z \int_Z \frac{(i_Z^* \phi - f(Z))e^{i_Z^* \omega - H(Z)}}{\prod_{j=1}^{n-m_Z} (c_1(E_j^Z) + \kappa_j^Z / 2\pi)}.$$

Remark 3.2 The Duistermaat–Heckman formula (13)–(14) has been explained and generalized in term of equivariant cohomology by Atiyah and Bott [1] and Berline and Vergne [4]. The claim (15) is a particular case of Theorem 7.13 of [3, page 216]. A mixture of the equivariant cohomology approach and the original proof of Duistermaat and Heckman is given in [28] when H is a Morse function.

Remark 3.3 Whenever a torus T acts on (M, ω) in a Hamiltonian fashion and with a momentum map

$$\mu: M \rightarrow \mathfrak{t}^*,$$

it might be interesting to use formula (15) when varying $X = -X_a$, where $a \in \mathfrak{t}$ lies in the lattice Λ of circle subgroups. Then $H = \langle \mu, a \rangle$ and we have to replace $\kappa_j^Z = -\langle \underline{\kappa}_j^Z, a \rangle$ for some (fixed) weights $\underline{\kappa}_1^Z, \dots, \underline{\kappa}_{n-m_Z}^Z \in \Lambda^*$. By continuity, formula (15) is valid for any $a \in \mathfrak{t}$ such that $\langle \underline{\kappa}_j^Z, a \rangle \neq 0$ for $j = 1, \dots, n - m_Z$.

Remark 3.4 As explained in [27], the Duistermaat–Heckman formula (13) holds on a noncompact manifold assuming that Z lies in the interior of M and the measure tends to 0 on the boundary of M . Indeed, the proof is essentially of local nature, up to the final integration. Moreover, the power series involved in formulas (13) and (15) are proved to coincide term by term. Therefore, it holds on a noncompact manifold when understood as “if the left-hand side of (15) converges, then it coincides with the right-hand side of (15), which, then, converges”.

Remark 3.5 Formula (13) also holds for orbifolds, as pointed out in [27], with the slight modification

$$(16) \quad \int_M e^{\omega-H} = \sum_Z \frac{1}{d_Z} e^{-H(Z)} \int_Z \prod_j \frac{e^{t_Z^* \omega}}{(2\pi c_1(E_j^Z) + \kappa_j^Z)/2\pi},$$

where d_Z is the order of the orbifold group of the generic points of Z . Again this orbifold extension is also straightforward for formula (15), which becomes

$$(17) \quad \int_M (\phi - f)e^{\omega-H} = \sum_Z \frac{1}{d_Z} \int_Z \frac{(i_Z^* \phi - f(Z))e^{t_Z^* \omega - H(Z)}}{\prod_{j=1}^{n-m_Z} (c_1(E_j^Z) + \kappa_j^Z/2\pi)}.$$

Remark 3.6 Another case of extension is when ω is degenerated. Indeed, the only property of ω which is actually important is the relation $X \lrcorner \omega = dH$ — see [3] — together with the nice properties of Hamiltonian group actions, which may be ensured by other means than the nondegeneracy of ω .

4 Applications in Sasakian geometry

4.1 An orbiresolution of the cone

Pick $\xi_o \in \mathfrak{t}^+$, a quasiregular Reeb vector field on (N, \mathcal{D}) with symplectic orbifold quotient (W, σ) . Denote by $\mathcal{S}_o = (N, \mathcal{D}, J, \xi_o, \eta_o, g_o)$ the quasiregular Sasakian structure on N associated to ξ_o . Let $b_o \in \mathfrak{t}$ be the lattice element giving the Reeb vector field $\xi_o = X_{b_o}$ and $S_o^1 \subset T$ the closed subgroup (a 1–dimensional torus) induced by b_o . We obtain an orbifold resolution of the Kähler cone with the cone point included, $Y = Y_0 \cup \{0\}$, by constructing the line orbibundle \mathcal{L} over W . We construct it as a Kähler reduction as follows. The Kähler cone (Y_0, ω_o, J_o) — see Section 2.1.2 — comes equipped with the Hamiltonian action of T and the (homogenous) momentum map

$$\hat{\mu}: Y_0 \rightarrow \mathfrak{t}^*,$$

called the universal momentum map by Lerman [24], so that, with respect to Convention 2.2, we have

$$\frac{1}{2}r^2 = \langle \hat{\mu}, b_o \rangle.$$

Recall also that we have a natural smooth projection $p: Y_0 \rightarrow N$.

On the symplectic product $(Y_0 \times \mathbb{C}, \omega_o + \omega_{\text{std}})$ we consider the Hamiltonian function

$$v(y, z) = \langle \hat{\mu}_y, b_o \rangle - \frac{1}{2}|z|^2$$

with Hamiltonian vector field $\xi_o - \partial_\theta$ inducing an S^1 -action, which we denote by $S^1_{\mathcal{A}}$. One can check that

$$v^{-1}\left(\frac{1}{2}\right) = \{(y, z) \mid \langle \hat{\mu}_y, b_o \rangle = \frac{1}{2} + \frac{1}{2}|z|^2\} \simeq N \times \mathbb{C}$$

and that $\xi_o - \partial_\theta$ is nowhere zero on $v^{-1}\left(\frac{1}{2}\right)$. Therefore, $1 \in \mathbb{R}$ is a regular value of v and we can perform the symplectic reduction

$$\mathcal{L} = v^{-1}\left(\frac{1}{2}\right)/S^1_{\mathcal{A}},$$

which is an orbifold with a natural symplectic form Ω defined by $q^*\Omega = i^*(\omega_o + \omega_{\text{std}})$, where $q: v^{-1}\left(\frac{1}{2}\right) \rightarrow \mathcal{L}$ is the quotient map and $i: v^{-1}\left(\frac{1}{2}\right) \hookrightarrow Y_0 \times \mathbb{C}$ is the inclusion.

A point of \mathcal{L} is an orbit $[(y, z)]_{\mathcal{A}} = S^1_{\mathcal{A}} \cdot (y, z)$. On the level set $z = 0$ the action of $S^1_{\mathcal{A}}$ coincides with the one of S^1_o and we get $\mathcal{L} \cap \{[(y, z)]_{\mathcal{A}} \mid z = 0\} \simeq W := \{[p(y)]_o = S^1_o \cdot p(y)\}$. The space \mathcal{L} is a line bundle over W with bundle map $\pi([(y, z)]_{\mathcal{A}}) = [p(y)]_o$. Moreover, there is a smooth bijective map

$$(18) \quad f: \mathcal{L}_o \rightarrow Y_0, \quad [(y, z)]_{\mathcal{A}} \mapsto \left(\frac{z}{|z|} \cdot p(y), |z|\right),$$

where \cdot denotes the action of S^1 induced by ξ_o (that is $e^{2\pi i\theta} \cdot x = \exp_T(\theta b_o) \cdot x$) and $\mathcal{L}_o = \{[(y, z)]_{\mathcal{A}} \in \mathcal{L} \mid z \neq 0\} = \mathcal{L} \setminus (0\text{-section})$. Note that the pullback $q^*f: Y_0 \times \mathbb{C} \rightarrow Y_0$ is \mathbb{C}^* -invariant and one can check by direct computation that $(q^*f)_* \circ (J_o + i) = J_o \circ (q^*f)_*$. Hence, $f: \mathcal{L}_o \rightarrow Y_0$ is biholomorphic and we get a birational map $f: \mathcal{L} \dashrightarrow Y_0$.

Note that the total space of \mathcal{L} inherits the Hamiltonian action of $G = (T \times S^1)/S^1_{\mathcal{A}}$ with momentum map

$$\check{\nu}: \mathcal{L} \rightarrow \mathfrak{g}^* \simeq \{(\alpha, t) \in \mathfrak{t}^* \oplus \mathbb{R} \mid \langle \alpha, b_o \rangle = \frac{1}{2} + \frac{1}{2}t\}.$$

Given $u \in T$ we may consider the element $[(u, 1)]_{S^1_{\mathcal{A}}} \in G$; this gives an isomorphism making f an equivariant map. In what follows we identify the action of G and T using that latter map. The image of $\check{\nu}$ is identified with the truncated polyhedral cone $\{x \in \text{Im } \hat{\mu} \subset \mathfrak{t}^* \mid \langle x, b_o \rangle \geq \frac{1}{2}\}$.

The zero section $W \hookrightarrow \mathcal{L}$, with its induced symplectic structure, is naturally identified with the symplectic quotient (W, σ) — see (8) — associated to the quasiregular Reeb

vector field $\xi_o = X_{b_o}$. Also the action of G coincides with the action of T/S_o^1 with momentum map $\check{\mu}_{\eta_o}: W \rightarrow (\mathfrak{t}/\mathbb{R}b_o)^*$; see Section 2.2.2.

4.1.1 The weights of the torus action on the orbiresolution Let $Z \subset \mathcal{L}$ be a connected component of the fixed-point set of T . In particular, $Z \subset W$ is seen inside \mathcal{L} as the zero section. For $p \in Z$, the action of T gives a decomposition into equivariant bundles

$$(19) \quad T_p \mathcal{L} = T_p W \oplus \mathcal{L}_p = T_p Z \oplus \left(\bigoplus_{j=0}^{n-m_Z} E_{j,p}^Z \right).$$

Here, and in what follows, \mathcal{L}_p is canonically identified with its tangent in $T\mathcal{L}$.

Since $Z \subset W$, the fiber \mathcal{L}_p is one of the summands in the right-hand side of (19) and we put

$$\mathcal{L}_p = E_{0,p}^Z.$$

We can easily compute the weight of the action of T on the fiber \mathcal{L}_p . Indeed, p is fixed by T if and only if for any $y \in Y_o$ such that $[p(y)]_o = p$ we have $\{X_a(y) \mid a \in \mathfrak{t}\} = \mathbb{R}\xi_o(y)$. Then, for such a point $p = [(y, 0)]_{\mathcal{A}}$ and $b \in \mathfrak{t}$ we can define $\delta_b \in \mathbb{R}$ by $\delta_b X_{b_o}(y) = X_b(y)$, so that

$$[(\exp_T(tb)y, z)]_{\mathcal{A}} = [(\exp_T(tb - t\delta_b b_o) \cdot y, e^{t\delta_b z})]_{\mathcal{A}} = [(y, e^{t\delta_b z})]_{\mathcal{A}}.$$

Therefore, the weight $\underline{\kappa}_0^Z \in \mathfrak{t}^*$ is

$$(20) \quad \langle \underline{\kappa}_0^Z, b \rangle = \delta_b = \eta_o(X_b)_q$$

for any $q \in N \cap \pi^{-1}(p)$.

Note that the other weights $\underline{\kappa}_1^Z, \dots, \underline{\kappa}_{n-m_Z}^Z \in \mathfrak{t}^*$ all lie in $(\mathfrak{t}/\mathbb{R}b_o)^* \simeq b_o^0 \subset \mathfrak{t}^*$, the annihilator of b_o , since $\bigoplus_{j=0}^{n-m_Z} E_{j,p}^Z \subset T_p W$ and $W \subset \text{Fix}_{\mathcal{L}} S_o^1$.

4.2 The volume functional

To prove Theorem 1.1 when the Kähler cone is Gorenstein, Martelli, Sparks and Yau [27] suggest to use the Duistermaat–Heckman formula (13), proved in [16], on the crepant resolution of the cone (with its cone point), assuming the existence of an approximation of the pulled-back cone metric on the resolution. Further, they produce an algebrogeometric proof of their theorem. Our proof follows the lines of the first suggested approach in [27], but we get rid of the assumption on the existence of the approximation, use an orbiline bundle, and we don't need the assumption that the Kähler cone is Gorenstein.

Let $(N, \mathcal{D}, Jg, \xi, \eta)$ be a Sasakian manifold of dimension $2n + 1$ and (Y_0, ω, J) its Kähler cone; see Convention 2.2. We denote by $Y = Y_0 \cup \{0\}$ the union of Y_0 with its cone point. Note that, with respect to the notation of Section 2.1, the Hamiltonian function of ξ is $\frac{1}{2}r^2$, in the sense that $\omega(\xi, \cdot) = -d(\frac{1}{2}r^2)$, since $\omega = \frac{1}{2}d(r^2\eta)$. Moreover, using Fubini’s theorem and integration by parts, we have

$$(21) \quad V_\xi := \int_N \frac{\eta \wedge (d\eta)^n}{n!} = \frac{a^{n+1}}{n!} \int_{Y_0} e^{-ar^2/2} \frac{\omega^{n+1}}{(n+1)!}$$

for any $a > 0$. The natural idea is to use the Duistermaat–Heckman formula to expand the right-hand side term as a rational function of $\xi \in \mathfrak{t}^+$; however, ξ has no fixed point. An idea of Martelli, Sparks and Yau is then to use a resolution of Y and approximate the Kähler structure of (Y_0, ω, J) on it.

We will use the Duistermaat–Heckman formula (16), Remark 3.3 of Martelli, Sparks and Yau [27] and the observation (21) on the total space of \mathcal{L} constructed in Section 4.1. Observe that in these formulas ω is not required to be symplectic and η does not need to be defined everywhere. Namely we note that

- (i) equation (21) holds if and only if $Y_0 = N \times \mathbb{R}^+$ and

$$r^2 \in C^\infty(Y_0), \quad \omega = \frac{1}{2}d(r^2\eta) \in \Omega^2(Y_0), \quad \eta \text{ is invariant by dilation;}$$

- (ii) equation (16) holds if and only if ω is closed and $\omega(X, \cdot) = dH$;

- (iii) (16) coincides with (21) (for $a = 1$) if and only $-\frac{1}{2}r^2 = -H$.

Observe that the restriction of the function $z \rightarrow |z|$ on $\nu^{-1}(\frac{1}{2}) \subset Y_0 \times \mathbb{C}$ (see Section 4.1) is well defined on the quotient \mathcal{L} and that $|z|^2$ is smooth. Moreover, $|z| = f^*r$, where the map f is defined by (18) and r is the projection on the second factor of Y_0 ; see Convention 2.2.

The vector field $\xi = X_b$, as a vector field on Y_0 , commutes with the action of ξ_o , thus of \mathcal{A} , and descends as a well-defined vector field, still denoted by $\xi = X_b$, on \mathcal{L} . The contact form η_ξ on N , associated to ξ , is not basic with respect to ξ_o and thus not well defined on \mathcal{L} a priori. However, we can define $\tilde{\eta}$ on $\mathcal{L} \setminus W$ by imposing $\tilde{\eta}(\xi) = 1$, $d\tilde{\eta}(\xi, \cdot) = 0$ and that $\tilde{\eta}$ coincides with $f^*\eta$ on $r^{-1}(1)$. It does not define $\tilde{\eta}$ uniquely but we only need these conditions to make our argument work. Note that ξ commutes with the vector field $r \frac{\partial}{\partial r}$ induced by the dilation on the second factor of $N \times \mathbb{C}$. Hence, $\tilde{\eta}$ is invariant by the dilation, that is homogenous of degree 0 with respect to $r \frac{\partial}{\partial r}$. Then, using observation (i) above, (21) holds on $Y_0 \simeq \mathcal{L} \setminus W$ for $\omega := \frac{1}{2}d(r^2\tilde{\eta})$.

Now we can forget $\tilde{\eta}$ and work with the right-hand side of (21) for $a = 1$. Indeed, the 1-form $\tilde{\eta}$ cannot be extended continuously on \mathcal{L} but $r^2\tilde{\eta}$ can and then ω is a smooth closed 2-form on \mathcal{L} . So, the integrand of the right-hand side of (21) makes sense on \mathcal{L} and then, because W has measure zero, we have

$$(22) \quad \int_N \frac{\eta \wedge (d\eta)^n}{n!} = \frac{1}{n!} \int_{\mathcal{L}} e^{-r^2/2} \frac{\omega^{n+1}}{(n+1)!}.$$

Here, we identify N with $r^{-1}(1) \subset \mathcal{L}$.

We can use (16) with ω , $X = -\xi$ and $H = \frac{1}{2}r^2$, which are defined on the whole \mathcal{L} . The equation $\omega(X, \cdot) = \frac{1}{2}dr^2$ holds on W , because both sides vanish, and on $\mathcal{L} \setminus W$, thanks to the fact that $\tilde{\eta}(X) = -1$, $d\tilde{\eta}(X, \cdot) = 0$ and $\omega = \frac{1}{2}d(r^2\tilde{\eta})$. Hence, using observations (ii) and (iii) above we get that (16) coincides with (22). That is,

$$(23) \quad \int_{\mathcal{L}} e^{-r^2/2} \frac{\omega^{n+1}}{(n+1)!} = \sum_Z \frac{1}{d_Z} \int_Z \prod_j \frac{e^{t_Z^* \omega}}{c_1(E_j^Z) + \kappa_j^Z / 2\pi} \\ = \sum_Z \frac{1}{d_Z} \int_Z \prod_j \frac{e^{t_Z^* \omega}}{c_1(E_j^Z) - \langle \kappa_j^Z, b \rangle / 2\pi},$$

where the last line comes from Remark 3.3 to highlight the dependence on $b \in \mathfrak{t}^+$.

In the formula (23), the fixed-point set $Z = \bigcup Z$ of X lies in W , on which ω vanishes identically. Thus, in the integrand of the right-hand side of (23), only the first term of the power series in $t_Z^* \omega$ does not vanish a priori. In conclusion, we have

$$(24) \quad V_{X_b} = \frac{1}{n!} \sum_Z \frac{(2\pi)^{n-m_Z}}{d_Z} \int_Z \prod_{j=0}^{n-m_Z} \frac{1}{2\pi c_1(E_j^Z) - \langle \kappa_j^Z, b \rangle},$$

which can be written as a rational function of the $\langle \kappa_j^Z, b \rangle$ using the relation $(1-x)^{-1} = \sum_{s \geq 0} x^s$. Since the integration over Z picks up only the term of degree $2m_Z$, the factor $(2\pi)^{m_Z}$ goes out and we get a rational function of order $-(n+1)$ up to the overall factor $(2\pi)^{n+1}$,

$$(25) \quad V_{X_b} = \frac{(2\pi)^{n+1}}{n!} \sum_Z \frac{1}{d_Z} \int_Z \prod_{j=0}^{n-m_Z} \frac{1}{\langle \kappa_j^Z, b \rangle} \left(\sum_{s \geq 0} \left(\frac{c_1(E_j^Z)}{\langle \kappa_j^Z, b \rangle} \right)^s \right).$$

This concludes the proof of Theorem 1.1 □

Remark 4.1 If we start with a generic $b \in \mathfrak{t}^+$, (25) holds for the dense open subset of \mathfrak{t}^+ of elements sharing the same fixed-point set $\{Z\} \subset W$ which coincides with

the fixed points of $T/S_{\xi_o}^1$, the maximal torus acting on W . For example, in the toric case starting with a generic $b \in \mathfrak{t}^+$, then $\{Z\}$ is the fixed-point set of the torus and so consists of a finite number of isolated points. In this case, formula (25) gives back the formulas in [26; 23] as expected.

Corollary 4.2 *When $b \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ , V_{X_b} tends to $+\infty$.*

Proof When $b \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ (but away from $0 \in \partial\mathfrak{t}^+$), there is at least one point $p \in N$ such that $\eta_o(X_b)_p \rightarrow 0$ by definition of \mathfrak{t}^+ ; see (5). Therefore, p is a minimum of the function $\eta_o(X_b) = \langle \mu_{\eta_o}, b \rangle$ and $\mu_{\eta_o}(p)$ lies in the boundary of the polytope $\Delta_o = \mu_{\eta_o}(N)$, which, up to translation is the moment polytope of $(W, \sigma, T/S_o^1)$.

Recall from Section 2.2.2 that the function $\eta_o(X_b)$ is the pullback to N of an affine-linear function on Δ_o , so that it is a Morse–Bott function on N of even index. By a well-known argument — see [19] — $\eta_o(X_b)$ has a unique minimum in the sense that there is a unique connected component $\tilde{Z} \subset \text{crit}(\eta_o(X_b)) \subset N$ such that \tilde{Z} is a local minimum, thus it is a global minimum and there is no other part of N on which $\eta_o(X_b)$ tends to 0 faster than it does on \tilde{Z} .

Moreover, near $Z = \pi(\tilde{Z}) \subset W$ and seen as a function on W it can be locally written as $\eta_o(X_b) =_{\text{loc}} \sum_{i=1}^{n-m_Z} |v_i|^2 \langle \underline{\kappa}_i^Z, b \rangle + c_b$, where c_b is a constant and $v_i \in E_i^Z$ in the decomposition (19) corresponding to the action of T/S_o^1 endowed with an invariant and compatible metric $|\cdot|$; see eg [24]. Since p is a minimum we have

$$\langle \underline{\kappa}_i^Z, b \rangle > 0$$

for $i = 1, \dots, n-m_Z$. Therefore, combining this with the observations of Section 4.1.1, in particular that $c_1(E_0^Z) = \iota_Z^* c_1(\mathcal{L}) = \iota_Z^* [\sigma/2\pi] > 0$ and $\langle \underline{\kappa}_0^Z, b \rangle = \eta(X_b)_p$, we get that the leading term of V_{X_b} , which is

$$(26) \quad \frac{(2\pi)^{n+1}}{n! d_Z} \left(\prod_{j=0}^{n-m_Z} \frac{1}{\langle \underline{\kappa}_j^Z, b \rangle} \right) \int_Z \left(\frac{c_1(E_0^Z)}{\langle \underline{\kappa}_0^Z, b \rangle} \right)^{m_Z},$$

is positive and tends to $+\infty$ when $\langle \underline{\kappa}_0^Z, b \rangle = \eta_o(X_b)_p \rightarrow 0$. □

4.3 The total transversal scalar curvature functional

4.3.1 Variation of the transverse scalar curvature The goal is to understand better how the function $\xi \mapsto \mathbf{S}_\xi$ varies with ξ when \mathcal{D} is a fixed (cooriented) contact structure with a compatible CR–structure (\mathcal{D}, J) on N ; see (6).

Given a Sasaki structure $(N, \mathcal{D}, J, \xi, \eta_\xi)$ on N , we recall from [17] that

$$(27) \quad {}_sT \frac{\eta_\xi \wedge (d\eta_\xi)^n}{n!} = 2 \frac{\rho_\xi^T \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!}$$

and ρ_ξ^T is defined on each patch $U_\alpha \subset N$ as the pullback of the Ricci form $\rho_{\xi,\alpha}$ of the Kähler structure $(g_{\xi,\alpha}, \omega_{\xi,\alpha}, I)$ induced on V_α (see the notation from the second paragraph of Section 2.2, taken from [17]; here we add the index ξ to emphasize the dependence in the structure).

Consider another Sasaki structure $(N, \mathcal{D}, J, \xi_o, \eta_o)$ compatible with the same CR–structure (\mathcal{D}, J) and $\xi_o \in \mathfrak{t}^+$. The associated transversal Ricci form $\rho_o^T \in \Gamma(N, \wedge^2 \mathcal{D})$ is not basic with respect to ξ . Indeed, $\mathcal{L}_\xi \rho_o^T \equiv 0$ by T –invariance of the Sasaki structure $(N, \mathcal{D}, J, \xi_o, \eta_o)$, but there is no reason for $\rho_o^T(\xi, \cdot)$ to vanish identically. Actually, ξ_o commutes with ξ by hypothesis and descends as a Killing vector field on $(V_\alpha, g_{\xi,\alpha}, \omega_{\xi,\alpha}, I)$. Therefore, by the well-known formula — see [18, Lemma 1.23.4] — we have that

$$(28) \quad \rho_{\xi,\alpha}(\xi_o, \cdot) = -\frac{1}{2} d\Delta^{g_{\xi,\alpha}} \langle \check{\mu}_\xi, [b_o] \rangle,$$

where $\check{\mu}_\xi: V_\alpha \rightarrow (\mathfrak{t}/\mathbb{R}b)^*$ is the momentum map and b_o is such that $\xi_o = X_{b_o}$ (see Section 2.2.2).

Up to a constant, the pullback by $\pi_{\xi,\alpha}$ of the function $\langle \check{\mu}_\xi, [b_o] \rangle$ coincides with $\langle \mu_\xi, b_o \rangle = \frac{1}{2} \eta_\xi(\xi_o)$, which is globally defined on N . For any function $f \in C^2(V_\alpha)$ we have $\pi_{\xi,\alpha}^* \Delta^{g_{\xi,\alpha}} f = \Delta^{g_\xi} \pi_{\xi,\alpha}^* f$ because π is a Riemannian fibration with totally geodesic fibers [33]. Therefore, on N , we have the global formula

$$\rho_\xi^T(\xi_o, \cdot) = -\frac{1}{2} d\Delta^{g_\xi} \langle \mu_\xi, b_o \rangle,$$

or, similarly,

$$(29) \quad \rho_o^T(\xi, \cdot) = -\frac{1}{2} d\Delta^{g_o} \langle \mu_{\xi_o}, b \rangle.$$

In particular, putting $f_{o,\xi} := \frac{1}{2} \Delta^{g_o} \langle \mu_{\xi_o}, b \rangle$ we get that the 2–form

$$\beta = \rho_o^T + \eta_\xi \wedge df_{o,\xi} - f_{o,\xi} d\eta_\xi = \rho_o^T - d(f_{o,\xi} \eta_\xi)$$

is closed and ξ –basic (ie basic with respect to ξ). So the two closed 2–forms ρ_ξ^T and β lie in the same de Rham cohomology class, namely $c_1(\mathcal{D})$, and are both ξ –basic and J –invariant. Therefore, there exists a ξ –basic function $f \in C_B^\infty(N)$ such that

$$(30) \quad \rho_\xi^T - \beta = d_B d_B^c f.$$

Note that any basic function $f \in C_B^2(N)$ defines a function, still denoted by f , on each chart V_α and, applying a classical Kähler formula on $(V_\alpha, g_{\xi,\alpha}, \omega_{\xi,\alpha}, I)$ (see eg [18]), we have

$$\frac{dd^c f \wedge (\omega_\xi^T)^{n-1}}{(n-1)!} = -(\Delta^{g_{\xi,\alpha}} f) \frac{(\omega_\xi^T)^n}{n!},$$

where $\Delta^{g_{\xi,\alpha}}$ is the Laplacian associated to the metric $g_{\xi,\alpha}$. Therefore, we have, for any basic function $f \in C_B^2(N)$,

$$\frac{d_B d_B^c f \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} = -(\Delta_B^{g_\xi} f) \frac{\eta_\xi \wedge (d\eta_\xi)^n}{n!},$$

where $\Delta_B^{g_\xi} f = d_B^* d_B f$ is the basic Laplacian.

Putting this, (30) and (27) in (11) and using the fact that $\eta_\xi \wedge \eta_\xi \equiv 0$, we get

$$\begin{aligned} (31) \quad \mathbf{S}_\xi &= 2 \int_N \frac{\rho_\xi^T \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} \\ &= 2 \int_N \frac{\beta \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} + \int_N \frac{(d_B d_B^c f) \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} \\ &= 2 \int_N \frac{\beta \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} - \int_N (\Delta_B^{g_\xi} f) \frac{\eta_\xi \wedge (d\eta_\xi)^n}{n!} \\ &= 2 \int_N \frac{\beta \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} \\ &= 2 \int_N \frac{(\rho_o^T - d(f_{o,\xi} \eta_\xi)) \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} \\ &= 2 \int_N \frac{\rho_o^T \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1}}{(n-1)!} - 2 \int_N f_{o,\xi} \frac{\eta_\xi \wedge (d\eta_\xi)^n}{(n-1)!}. \end{aligned}$$

Note that $f_{o,\xi} = \frac{1}{2} \Delta^{g_o} \langle \mu_{\xi_o}, b \rangle = \frac{1}{4} \Delta^{g_o} \eta_o(\xi)$ is linear with respect to $\xi \in \mathfrak{t}^+$. Putting $dv_o = (\eta_o \wedge (d\eta_o)^n)/n!$, another way to write (31) is

$$\begin{aligned} (32) \quad \mathbf{S}_\xi &= \int_N \frac{s_o^T}{\eta_o(\xi)^n} dv_o - 2 \int_N \frac{f_{o,\xi}}{\eta_o(\xi)^{n+1}} dv_o \\ &= \int_N \frac{s_o^T}{\eta_o(\xi)^n} dv_o + \frac{n+1}{2} \int_N \frac{|d\eta_o(\xi)|_{g_o}^2}{\eta_o(\xi)^{n+2}} dv_o. \end{aligned}$$

Corollary 4.3 *Within a Sasaki cone there is at most one ray of vanishing transverse scalar curvature. Moreover, if there is one Sasaki structure with nonnegative transverse*

scalar curvature then the total transverse scalar curvature is nonnegative on the whole Sasaki cone.

4.3.2 A localization formula for the total transverse scalar curvature Using polar coordinates on $Y_0 = N \times \mathbb{R}_{>0}$, Fubini’s theorem and integration by parts, we have, for any $\eta \in \Omega^1(N)$, $\phi \in \Omega^2(N)$ and $a > 0$,

$$(33) \quad \int_N \frac{\eta \wedge \phi \wedge (d\eta)^{n-1}}{(n-1)!} = \frac{a^n}{(n-1)!} \int_{Y_0} e^{-ar^2/2} \frac{\phi \wedge \omega^n}{n!},$$

where $\omega = \frac{1}{2}dr^2\eta$.

Given a Sasakian manifold $(N, \mathcal{D}, J, g, \xi, \eta)$ of dimension $2n + 1$ as above, we use the notation of Sections 4.2 and 2.1. We consider the resolution $f: \mathcal{L} \dashrightarrow Y$ of Section 4.1 associated with a quasiregular Reeb vector field $\xi_o \in \mathfrak{t}^+$, inducing the isometric action of a circle $S^1_{\xi_o} \subset T$ whose quotient Kähler orbifold is $(W, \sigma_o, \check{J}_o, \check{g}_o)$. Recall that \mathcal{L} is obtained as the quotient of $v^{-1}(\frac{1}{2}) = N \times \mathbb{C}$ by the action of S^1 induced by $\xi_o - \partial_\theta$. In particular, a ξ_o basic k -form on N descends as a well-defined k -form on \mathcal{L} ; for example, ρ_o^T and $f_{o,\xi}$ are both defined on the total space of \mathcal{L} . Then, we can use the observations (33) and (21) to modify respectively the first and the second term of the last line of (31) as follows:

$$(34) \quad \begin{aligned} \frac{(n-1)!}{2} \mathbf{S}_\xi &= \int_N \rho_o^T \wedge \eta_\xi \wedge (d\eta_\xi)^{n-1} - \int_N f_{o,\xi} \eta_\xi \wedge (d\eta_\xi)^n \\ &= \int_{Y_0} e^{-r^2/2} \frac{\rho_o^T \wedge \omega^n}{n!} - \int_{Y_0} f_{o,\xi} e^{-r^2/2} \frac{\omega^{n+1}}{(n+1)!} \\ &= \int_Y e^{-r^2/2} (\rho_o^T - f_{o,\xi}) e^\omega \\ &= \int_Y (\rho_o^T - f_{o,\xi}) e^{\omega - r^2/2}, \end{aligned}$$

where $\omega = \frac{1}{2}dr^2\eta_\xi$.

Using Theorem 3.1 in the orbifold case (see Remark 3.5) with $X = -\xi$ and $H = \frac{1}{2}r^2$, we get

$$(35) \quad \begin{aligned} \frac{(n-1)!}{2} \mathbf{S}_\xi &= \sum_Z \frac{1}{d_Z} \int_Z \frac{(i_Z^* \rho_o^T - f_{o,\xi}(Z)) e^{i_Z^* \omega}}{\prod_{j=0}^{n-m_Z} (c_1(E_j^Z) + \kappa_j^Z / 2\pi)} \\ &= \sum_Z \frac{1}{d_Z} \int_Z \frac{(i_Z^* \rho_o^T - f_{o,\xi}(Z)) e^{i_Z^* \omega}}{\prod_{j=0}^{n-m_Z} (c_1(E_j^Z) - \langle \kappa_j^Z, b \rangle / 2\pi)}. \end{aligned}$$

Again, the fixed-point set $Z = \bigcup Z$ of X lies in W , on which ω vanishes identically. Thus, in the integrand of the right-hand side of (35), only the first term of the power series in $\iota_Z^* \omega$ does not vanish a priori. In conclusion, we have

$$\begin{aligned}
 (36) \quad \mathcal{S}_\xi &= \frac{2}{(n-1)!} \sum_Z \frac{1}{d_Z} \int_Z \frac{(\iota_Z^* \rho_o^T - f_{o,\xi}(Z))}{\prod_{j=0}^{n-m_Z} (c_1(E_j^Z) - \langle \underline{\kappa}_j^Z, b \rangle / 2\pi)} \\
 &= \frac{2(2\pi)^{n+1}}{(n-1)!} \sum_Z \frac{1}{d_Z} \left[\int_Z (\iota_Z^* \rho_o^T / 2\pi) \wedge \left(\prod_{j=0}^{n-m_Z} \frac{1}{\langle \underline{\kappa}_j^Z, b \rangle} \sum_{s \geq 0} \left(\frac{c_1(E_j^Z)}{\langle \underline{\kappa}_j^Z, b \rangle} \right)^s \right) \right. \\
 &\quad \left. - f_{o,\xi}(Z) \int_Z \prod_{j=0}^{n-m_Z} \frac{1}{\langle \underline{\kappa}_j^Z, b \rangle} \sum_{s \geq 0} \left(\frac{c_1(E_j^Z)}{\langle \underline{\kappa}_j^Z, b \rangle} \right)^s \right].
 \end{aligned}$$

Remark 4.4 Whenever the fixed-point set Z of X consists in a set of isolated points $Z = \{p\}$, formula (36) simplifies as

$$(37) \quad \mathcal{S}_\xi = \frac{2(2\pi)^{n+1}}{(n-1)!} \sum_p \frac{1}{d_p} \frac{-f_{o,\xi}(p)}{\prod_{j=1}^{n+1} \langle \underline{\kappa}_j^p, b \rangle}.$$

4.3.3 Proof of Theorem 1.2 Using (36) and the facts that $\iota_Z^* \rho_o^T / 2\pi, c_1(E_j^Z) \in H_{\text{dR}}^2(Z, \mathbb{Q})$ and $\lambda_j^Z \in \Lambda^*$ (see Remark 3.3), to prove Theorem 1.2 it remains to show that $f_{o,\xi}(Z) \in \mathbb{Q}$.

Lemma 4.5 Let (M^n, ω, J, g) be a Kähler orbifold endowed with an effective Hamiltonian isometric action of a torus T^m and a momentum map $\mu: M \rightarrow \mathfrak{t}^*$ (where $\mathfrak{t} := \text{Lie}(T)$). For any Z , a connected component of $\text{Fix}_M T$ with $\dim Z = 2m_Z$, denoting the equivariant decomposition by $T_z M = T_z Z \oplus \bigoplus_{j=1}^{n-m_Z} E_j^Z$ with weights $\underline{\kappa}_1^Z, \dots, \underline{\kappa}_{n-m_Z}^Z$, we have

$$(\Delta^g \mu)_z = -2 \sum_{j=1}^{n-m_Z} \underline{\kappa}_j^Z$$

at any $z \in Z$.

Proof We prove the lemma for M a manifold. Since the result is local this assumption is not restrictive.

Let first recall some consequences of the symplectic slice theorem; see eg [2; 25]. We choose $z \in M$, a fixed point of T , and call $\phi: T \rightarrow \text{Symp}(T_x M, \omega)$ the representation such that $\phi_\theta = d_z \theta$, where $\theta \in T$ is viewed as a diffeomorphism of M ; we write

the corresponding representation as $\phi_i: T \rightarrow \text{Symp}(E_i^Z, \omega_i)$. With this notation, $\kappa_i^Z = d\phi_i \in \mathfrak{t}^*$ are the weights of the action ϕ . Note that W_0 is the tangent space of the connected component Z of the fixed-point set of T .

Let $b \in \mathfrak{t}$ and put $\mu_b = \langle \mu, b \rangle$. The Laplacian $\Delta^g \mu_b$ is also $-\text{tr Hess}^g \mu_b$, where we have at z , for vector fields X and Y defined near z ,

$$\begin{aligned}
 (38) \quad \text{Hess}^g \mu_b(X, Y) &= \nabla d\mu_b(X, Y) \\
 &= X(d\mu_b(Y)) \\
 &= -X(\omega(X_b, Y)) \\
 &= -\omega(\nabla_X X_b, Y)
 \end{aligned}$$

by using twice the fact that z is a critical point of μ_b and the Kähler condition $\nabla J = 0$. Hence, since X_b is a real holomorphic vector field, that is, $\nabla_{JY} X_b = J\nabla_Y X_b$, and $\mathcal{L}_X \omega(X_b, Y) = 0$ for any $X \in \Gamma(TZ)$, we obtain

$$\Delta^g \mu_b = \sum_{i=1}^n \omega(\nabla_{w_i} X_b, w_i) + \omega(\nabla_{Jw_i} X_b, Jw_i),$$

where $\{w_i, Jw_i\}$ is a normal symplectic basis of E_i for $i = 1, \dots, n - m_Z$ and $\{w_i, Jw_i\}_{i=n-m+1}^n$ is a normal symplectic basis of $T_z Z$. Thanks to the fact that X_b is real holomorphic, it reduces to

$$\Delta^g \mu_b = 2 \sum_{i=1}^n \omega(\nabla_{w_i} X_b, w_i).$$

Consider the path $\theta_t = \exp tb$ in the torus T , so $(d\phi)(b) = \frac{d}{dt} \Big|_{t=0} \phi_{\theta_t} \in \mathfrak{symp}(T_z M, \omega)$ and $v \in T_z M$ with the flow $\gamma_s \subset M$ (ie $\frac{d}{ds} \Big|_{s=0} \gamma_s = v$). With this notation,

$$\begin{aligned}
 (39) \quad d\phi_b(v) &= \frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} (\theta_t \circ \gamma_s) \\
 &= \frac{d}{ds} \Big|_{s=0} \frac{d}{dt} \Big|_{t=0} (\theta_t \circ \gamma_s) \\
 &= \frac{d}{ds} \Big|_{s=0} X_b(\gamma_s) \\
 &= dX_b(v),
 \end{aligned}$$

seen as a section $X_b: M \rightarrow TM$. Recall that g induces a Riemannian metric on TM , denoted by g as well, so the g -orthogonal decomposition into the vertical and horizontal parts of $T_{(z,v)}(TM) = T_v(T_z M) \oplus \mathcal{H}^\nabla$ gives $dX_b(v) = \nabla_v X_b + \tilde{v}$. Hence,

identifying $T_v(T_Z M) = T_Z M$, we obtain

$$\Delta^g \mu_b = 2 \sum_{i=1}^n \omega(\nabla_{w_i} X_b, w_i) = 2 \sum_{i=1}^n \omega(d\phi_b(w_i), w_i) = -2 \sum_{i=1}^{n-m} \langle d\phi_i, b \rangle. \quad \square$$

A direct consequence of this lemma, applied on a Kähler orbifold $(W, \sigma_o, \check{J}_o, \check{g}_o)$, is that for $\xi = X_b$, the function

$$-2f_{o,\xi}(z) = -(\Delta^{g_o} \langle \check{\mu}_o, [b] \rangle)_z = 2 \left\langle \sum_{j=1}^{n-m_Z} \underline{\kappa}_j^Z, [b] \right\rangle = 2 \left\langle \sum_{j=1}^{n-m_Z} \underline{\kappa}_j^Z, b \right\rangle,$$

when restricted on Z , is a rational polynomial. Actually it is a linear function with rational coefficients. Recall from Section 4.1.1 that $\underline{\kappa}_1^Z, \dots, \underline{\kappa}_{n-m_Z}^Z$ lie in the annihilator of $b_o \in \mathfrak{t}$ and thus are defined on $\mathfrak{t}/\mathbb{R}b_o$.

This completes the proof of Theorem 1.2.

Corollary 4.6 *When $b \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ , S_{X_b} tends to $+\infty$.*

Proof This goes essentially as the proof of Corollary 4.2. When $b \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ (but away from $0 \in \partial\mathfrak{t}^+$), there is at least one point $p \in N$ such that $\eta_o(X_b)_p \rightarrow 0$, by definition of \mathfrak{t}^+ ; see (5). Therefore, p is a minimum of the function $\eta_o(X_b) = \langle \mu_{\eta_o}, b \rangle$ and, arguing as in the proof of Corollary 4.2, we get that there is a unique connected component $\tilde{Z} \subset \text{crit}(\eta_o(X_b)) \subset N$ such that \tilde{Z} is a local minimum, thus it is a global minimum and there is no other part of N on which $\eta_o(X_b)$ tends to 0 faster than it does on \tilde{Z} . Moreover,

$$\langle \underline{\kappa}_i^Z, b \rangle > 0$$

for $i = 1, \dots, n - m_Z$ and — see Section 4.1.1 — we have $\langle \underline{\kappa}_0^Z, b \rangle = \eta_o(X_b)_p$. Hence, the leading term of S_{X_b} in (36) is the one where $\langle \underline{\kappa}_0^Z, b \rangle$ appears with the greater exponent. This term is

$$(40) \quad \frac{-2f_{o,\xi}(Z)}{d_Z(n-1)!} \left(\prod_{j=0}^{n-m_Z} \frac{1}{\langle \underline{\kappa}_j^Z, b \rangle} \right) \int_Z \left(\frac{c_1(E_0^Z)}{\langle \underline{\kappa}_0^Z, b \rangle} \right)^{m_Z}.$$

It is positive by the observations above, Lemma 4.5 and since $c_1(E_0^Z) = \iota_Z^* c_1(\mathcal{L}) = \iota_Z^* [\sigma/2\pi] > 0$ (see Section 4.1.1) and tends to $+\infty$ when $\langle \underline{\kappa}_0^Z, b \rangle = \eta_o(X_b)_p \rightarrow 0$. \square

5 The Einstein–Hilbert functional and scalar curvature

Here, by scalar curvature we mean the transverse scalar curvature s_ξ^T of a Sasakian structure $\mathcal{S} = (\mathcal{D}, J, \xi, \eta, g)$. Of course, s_ξ^T is related to the scalar curvature s_g of the Sasaki metric g by $s_g = s_\xi^T - 2n$ [7]. Recall the Einstein–Hilbert functional (4), which can be written in terms of the average scalar curvature \bar{s}_ξ^T as

$$H(\xi) = \frac{S_\xi^{n+1}}{V_\xi^n} = (\bar{s}_\xi^T)^{n+1} V_\xi.$$

Since it is invariant under the transverse homothety operation $\xi \mapsto a^{-1}\xi$ with $a \in \mathbb{R}^+$, it descends to a function on the space of rays of the Sasaki cone \mathfrak{t}^+ . We have:

Lemma 5.1 *When $b \in \mathfrak{t}^+$ tends to the boundary of \mathfrak{t}^+ , $H(X_b)$ tends to $+\infty$.*

Proof Both S and V tend to $+\infty$ when b tends to the boundary of \mathfrak{t}^+ . Moreover, considering the leading terms (26) and (40) in equation (4) and putting $\langle \underline{\kappa}_0^Z, b \rangle = \eta_o(X_b)_p$ as a factor, we see that $H(X_b)$ behaves as

$$\frac{C(b)}{\langle \underline{\kappa}_0^Z, b \rangle^{m_Z+1}},$$

where $C(b)$ is a rational function tending to

$$f_{o,\xi}(Z)^{n+1} \left(\prod_{j=1}^{n-m_Z} \frac{1}{\langle \underline{\kappa}_j^Z, b \rangle} \right) \left(\int_Z c_1(E_0^Z)^{m_Z} \right) > 0$$

when $\langle \underline{\kappa}_0^Z, b \rangle = \eta_o(X_b)_p \rightarrow 0$. □

Equation (32) has several interesting consequences. Recall the definition of the *type* of a Sasakian structure. A Sasakian structure (N, ξ, η, Φ, g) is said to be of *positive (negative) type* if $c_1(\mathcal{F}_\xi)$ can be represented by a positive (negative) definite basic $(1, 1)$ form. If $c_1(\mathcal{F}_\xi)$ vanishes, it is of *null type*. When none of these hold it is of *indefinite type*. If $\dim \mathfrak{t}^+ \geq 2$, then the Sasakian structure is either of positive or indefinite type.

Lemma 5.2 *Fix a Sasaki structure $S_0 = (\mathcal{D}, J, \xi_0)$ such that its Sasaki automorphism group has dimension at least two.*

- (1) *There is no more than one ray of Reeb vector fields $\xi \in \mathfrak{t}^+$ having vanishing transverse scalar curvature s_0^T and if there is one such ray τ_0 then for all $\xi \in \mathfrak{t}^+ \setminus \{\tau_0\}$ we have $S_\xi > 0$.*

- (2) If S_0 satisfies $s_0^T > 0$ almost everywhere, then $S_\xi > 0$ for all $\xi \in \mathfrak{t}^+$. In particular, if S_0 is of positive type, then $S_\xi > 0$ for all $\xi \in \mathfrak{t}^+$.

Note that if Lemma 5.2(1) holds, then the ray \mathfrak{r}_0 is the unique global minimum of the Einstein–Hilbert functional H and $H(\xi) = 0$ for all $\xi \in \mathfrak{r}_0$. Moreover, no $\xi \in \mathfrak{t}^+$ has strictly positive scalar curvature. Somewhat more generally, we have:

Proposition 5.3 *If the dimension of \mathfrak{t}^+ is at least two and there exists $\xi \in \mathfrak{t}^+$ such that $S_\xi \leq 0$, then all elements of the Sasaki cone are of indefinite type.*

Proof Since $\dim \mathfrak{t}^+ > 1$, a Sasakian structure is either positive or indefinite. Suppose to the contrary that there is a Sasakian structure in \mathfrak{t}^+ of positive type; then taking this to be ξ_o in (32) gives a contradiction. \square

5.1 A global minimum

We now show that H always attains a global minimum.

Theorem 5.4 *Let (N, \mathcal{D}) be a contact manifold of Sasaki type with a T action of Reeb type. Then there exists $\xi_{\min} \in \mathfrak{t}^+$ that minimizes H . Moreover, if $H(\xi_{\min}) \neq 0$ and the corresponding Sasaki metric is extremal, it must have constant scalar curvature.*

Proof Recall [9] that a set Σ is a transversal subset of \mathfrak{t}^+ if $\Sigma \subset \mathfrak{t}_k^+$ and that Σ meets each ray passing through \mathfrak{t}^+ in a single point and is a codimension-one relatively compact subset of \mathfrak{t}^+ whose closure does not contain 0. Let $\bar{\mathfrak{t}}^+$ denote the closure of \mathfrak{t}^+ . Since $H(\xi)$ is scale-invariant, $H|_\Sigma$ is independent of Σ . We choose Σ to be any intersection of $\bar{\mathfrak{t}}^+$ with a transverse hyperplane. Then $P = \bar{\mathfrak{t}}^+ \cap \Sigma$ is a simple compact convex polytope. Now, by Lemma 5.1, $H(\xi)$ tends to $+\infty$ as ξ tends to the boundary ∂P . Thus, since H is a continuous function on the interior $P^\circ = P \setminus \partial P$, there exists ξ_{\min} such that $H(\xi) \geq H(\xi_{\min})$. This proves the first statement.

For the second statement we assume that $S_{\xi_{\min}} \neq 0$. Then it follows from [9] that the Sasaki–Futaki invariant F_ξ vanishes at $\xi = \xi_{\min}$. So, if S_{\min} is extremal, it has constant scalar curvature. \square

Note that the ray \mathfrak{r}_{\min} of ξ_{\min} is generally not necessarily unique. However, by Lemma 5.2(1), it is unique if there exists a Sasakian structure in \mathfrak{t}^+ with vanishing transverse scalar curvature, in which case $H(\xi) \geq 0$ for all $\xi \in \mathfrak{t}^+$ and equality holds if and only if $\xi = a\xi_{\min}$, where $a \in \mathbb{R}^+$. In this case we get a unique global minimum. Explicit examples are given by Proposition 5.17 of [10].

There are counterexamples to the second statement of Theorem 5.4 if we drop the hypothesis that \mathcal{S}_{\min} is extremal. See [6] for many examples where the entire Sasaki cone is relatively K -unstable and admits no extremal Sasaki metrics.

Another functional of interest to handle the case when $\mathcal{S}_\xi < 0$ and n is odd is defined by

$$(41) \quad H_1(\xi) = \text{sign}(\mathcal{S}_\xi) |H(\xi)|.$$

It follows as in the proof of Theorem 5.4 that H_1 also has a global minimum $\xi_{1 \min}$ giving

$$H_1(\xi) \geq H_1(\xi_{1 \min})$$

for all $\xi \in \mathfrak{t}^+$.

5.2 Positivity and rational curves

Recall that a projective algebraic variety X of complex dimension n is said to be *uniruled* if there exists a dominant rational map $X' \times \mathbb{C}P^1 \dashrightarrow X$, where X' is a variety of dimension $n - 1$. It is well known that a uniruled variety has Kodaira dimension $-\infty$ and that there is a rational curve through each point. Its relation with Sasakian geometry was mentioned briefly in Theorem 7.5.33 of [7], which states that a compact quasiregular Sasakian structure of positive type has a uniruled quotient variety. This follows directly by a result of Miyaoka and Mori [29]. In the simply connected case there is a topological classification of positive Sasakian 5-manifolds due to Kollar [22], which shows how the existence of irrational curves determines the torsion in $H_2(N^5, \mathbb{Z})$ (see also Section 10.2.1 of [7]). However, a recent result of Heier and Wong [21] shows that the hypothesis that the Sasaki manifold be of positive type can be weakened considerably.

Theorem 5.5 (Heier and Wong) *Let N be a regular compact Sasaki manifold with Reeb field ξ .*

- (1) *If $\mathcal{S}_\xi > 0$, then its quotient variety is uniruled.*
- (2) *If $\mathcal{S}_\xi = 0$, then either its quotient is uniruled or its canonical line bundle is torsion. In particular, if the dimension of $\mathfrak{t}^+ > 1$, the quotient is uniruled.*

Remark 5.6 It appears quite likely that this result can be generalized to the quasiregular case, although one would need to work with the orbifold canonical bundle. This seemingly entails generalizing the methods of Boucksom, Demailly, Păun and Peternell [5] to the case when the variety X is a normal projective variety with cyclic quotient

singularities. Nevertheless, below we are able to generalize Theorem 5.5 to a certain class of orbifolds.

The nonnegativity of a particular S_ξ is, however, far from the last word. Indeed, Example 5.16 of [10] gives a quasiregular Sasakian structure on an S^3 bundle over a Riemann surface of genus 23 with constant transverse scalar curvature equal to -16π . This is not only uniruled, but a ruled manifold.

Here we are content to consider a special case.

Theorem 5.7 *Let N be a compact manifold of dimension $2n + 1$ with a CR–structure (\mathcal{D}, J) of Sasaki type with Sasaki cone \mathfrak{t}^+ of dimension at least two. Suppose further that all quasiregular $\xi \in \mathfrak{t}^+$ have S^1 quotients of the form (W_ξ, Δ_ξ) , where W_ξ is a smooth projective variety and Δ_ξ is a branch divisor depending on ξ . If $S_\xi \geq 0$ then W_ξ is uniruled.*

Proof Recall [7] that a branch divisor has the form

$$\Delta = \sum_i \left(1 - \frac{1}{m_i}\right) D_i,$$

where D_i is an irreducible hypersurface contained in the orbifold singular locus and m_i is the ramification index of D_i and the sum is finite. The orbifold first Chern class and the first Chern class of X are related by

$$(42) \quad c_1^{\text{orb}}(W, \Delta) = c_1(W) - \sum_i \left(1 - \frac{1}{m_i}\right) c_1(L_i),$$

where L_i is the line bundle associated to D_i . Now $2\pi c_1(W)$ is represented by the Ricci form ρ_W , and $2\pi c_1^{\text{orb}}(W, \Delta)$ is represented by the orbifold Ricci form ρ^{orb} which is the pushforward of the transverse Ricci form ρ^T , which is well defined since it is basic. Let ω be any Kähler form on X ; then, by the work of Demailly [13], $2\pi c_1(L_i)$ is represented by the curvature current $c(L_i)$ of a singular Hermitian metric on L_i . Now equation (42) implies

$$(43) \quad \int_W \rho \wedge \omega^{n-1} = \int_W \rho^{\text{orb}} \wedge \omega^{n-1} + \sum_i \left(1 - \frac{1}{m_i}\right) \int_W c(L_i) \wedge \omega^{n-1}.$$

Note that the scalar curvature of the orbifold Kähler metric ω_0 is just the transverse scalar curvature of the corresponding Sasakian structure. The cohomology class $[\omega_0] \in H^2(W, \mathbb{Q})$ defines a positive orbiline bundle \mathcal{L} on (W, Δ) , so it is orbiample.

It follows from orbifold theory that \mathcal{L}^l , where $l = \text{lcm}(m_i)$, is an ample line bundle on X . Thus, $l[\omega_0] \in H^2(W, \mathbb{Z})$ and, since \mathcal{L}^l is ample, $l[\omega_0]$ can be represented by a Kähler form, which we choose to be ω . This implies

$$\int_W \rho^{\text{orb}} \wedge \omega^{n-1} = l^{n-1} \int_W \rho^{\text{orb}} \wedge \omega_0^{n-1} = \frac{l^{n-1}}{2n} \int_W s^T \omega_0^n = C \mathcal{S}_\xi$$

for some $C > 0$. If we let \mathcal{S}_W denote the total scalar curvature of X , this and equation (43) implies

$$\mathcal{S}_W = C \mathcal{S}_\xi + \sum_i \left(1 - \frac{1}{m_i}\right) \int_W c(L_i) \wedge \omega^{n-1}.$$

Since the D_i are effective, it follows from Proposition 4.2(b) of [13] that $c(L_i)$ satisfies $c(L_i) \geq \epsilon \omega$ for some $\epsilon > 0$, so this equation implies

$$(44) \quad \mathcal{S}_W \geq C \mathcal{S}_\xi + \epsilon n! \sum_i \left(1 - \frac{1}{m_i}\right) V_W.$$

So $\mathcal{S}_\xi \geq 0$ implies $\mathcal{S}_W > 0$, and the result follows by the Heier–Wong theorem, Theorem 5.5. \square

Actually, by looking at sequences of quasiregular Reeb fields that approach the boundary of \mathfrak{t}^+ , we can do better.

Corollary 5.8 *Assuming the hypothesis of Theorem 5.7, if there exists a sequence $\{\xi_k\}$ of quasiregular Reeb fields tending to the boundary in \mathfrak{t}^+ and such that $W_\xi = W_{\xi_k}$ is independent of k , then W_ξ is uniruled.*

Proof Since the dimension of \mathfrak{t}^+ is at least 2, we know by Corollary 4.6 that $\mathcal{S}_\xi \rightarrow +\infty$ near the boundary. So there exists a quasiregular $\xi \in \{\xi_k\} \subset \mathfrak{t}^+$ such that $\mathcal{S}_\xi > 0$. \square

Our assumption that all quasiregular $\xi \in \mathfrak{t}^+$ have a codimension-one orbifold singular set is restrictive. It is, however, realized by a fairly large collection of Sasaki families, for example those coming from the S_w^3 -join studied in [11]. For these one can also easily construct sequences satisfying the conditions of Corollary 5.8.

References

- [1] **MF Atiyah, R Bott**, *The moment map and equivariant cohomology*, *Topology* 23 (1984) 1–28 MR
- [2] **M Audin**, *The topology of torus actions on symplectic manifolds*, *Progress in Mathematics* 93, Birkhäuser, Basel (1991) MR

- [3] **N Berline, E Getzler, M Vergne**, *Heat kernels and Dirac operators*, Grundle Math. Wissen. 298, Springer (1992) MR
- [4] **N Berline, M Vergne**, *Zéros d'un champ de vecteurs et classes caractéristiques équivariantes*, Duke Math. J. 50 (1983) 539–549 MR
- [5] **S Boucksom, J-P Demailly, M Păun, T Peternell**, *The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension*, J. Algebraic Geom. 22 (2013) 201–248 MR
- [6] **C P Boyer, C van Coevering**, *Relative K-stability and extremal Sasaki metrics*, Math. Res. Lett. 25 (2018) 1–19 MR
- [7] **C P Boyer, K Galicki**, *Sasakian geometry*, Oxford Univ. Press (2008) MR
- [8] **C P Boyer, K Galicki, S R Simanca**, *Canonical Sasakian metrics*, Comm. Math. Phys. 279 (2008) 705–733 MR
- [9] **C P Boyer, H Huang, E Legendre, C W Tønnesen-Friedman**, *The Einstein–Hilbert functional and the Sasaki–Futaki invariant*, Int. Math. Res. Not. 2017 (2017) 1942–1974 MR
- [10] **C P Boyer, C W Tønnesen-Friedman**, *Extremal Sasakian geometry on S^3 -bundles over Riemann surfaces*, Int. Math. Res. Not. 2014 (2014) 5510–5562 MR
- [11] **C P Boyer, C W Tønnesen-Friedman**, *The Sasaki join, Hamiltonian 2-forms, and constant scalar curvature*, J. Geom. Anal. 26 (2016) 1023–1060 MR
- [12] **T C Collins, G Székelyhidi**, *K-semistability for irregular Sasakian manifolds*, J. Differential Geom. 109 (2018) 81–109 MR
- [13] **J-P Demailly**, *Singular Hermitian metrics on positive line bundles*, from “Complex algebraic varieties” (K Hulek, T Peternell, M Schneider, F-O Schreyer, editors), Lecture Notes in Math. 1507, Springer (1992) 87–104 MR
- [14] **S K Donaldson**, *Scalar curvature and stability of toric varieties*, J. Differential Geom. 62 (2002) 289–349 MR
- [15] **S K Donaldson**, *Constant scalar curvature metrics on toric surfaces*, Geom. Funct. Anal. 19 (2009) 83–136 MR
- [16] **J J Duistermaat, G J Heckman**, *On the variation in the cohomology of the symplectic form of the reduced phase space*, Invent. Math. 69 (1982) 259–268 MR Addendum in 72 (1983) 153–158 MR
- [17] **A Futaki, H Ono, G Wang**, *Transverse Kähler geometry of Sasaki manifolds and toric Sasaki–Einstein manifolds*, J. Differential Geom. 83 (2009) 585–635 MR
- [18] **P Gauduchon**, *Calabi’s extremal Kähler metrics: an elementary introduction*, book project (2015) Available at <http://germanio.math.unifi.it/wp-content/uploads/2015/03/dercalabi.pdf>
- [19] **V Guillemin, S Sternberg**, *Convexity properties of the moment mapping*, Invent. Math. 67 (1982) 491–513 MR

- [20] **W He, S Sun**, *The generalized Frankel conjecture in Sasaki geometry*, Int. Math. Res. Not. 2015 (2015) 99–118 MR
- [21] **G Heier, B Wong**, *Scalar curvature and uniruledness on projective manifolds*, Comm. Anal. Geom. 20 (2012) 751–764 MR
- [22] **J Kollár**, *Einstein metrics on five-dimensional Seifert bundles*, J. Geom. Anal. 15 (2005) 445–476 MR
- [23] **E Legendre**, *Existence and non-uniqueness of constant scalar curvature toric Sasaki metrics*, Compos. Math. 147 (2011) 1613–1634 MR
- [24] **E Lerman**, *Contact toric manifolds*, J. Symplectic Geom. 1 (2003) 785–828 MR
- [25] **E Lerman, S Tolman**, *Hamiltonian torus actions on symplectic orbifolds and toric varieties*, Trans. Amer. Math. Soc. 349 (1997) 4201–4230 MR
- [26] **D Martelli, J Sparks, S-T Yau**, *The geometric dual of α -maximisation for toric Sasaki–Einstein manifolds*, Comm. Math. Phys. 268 (2006) 39–65 MR
- [27] **D Martelli, J Sparks, S-T Yau**, *Sasaki–Einstein manifolds and volume minimisation*, Comm. Math. Phys. 280 (2008) 611–673 MR
- [28] **D McDuff, D Salamon**, *Introduction to symplectic topology*, Clarendon, New York (1995) MR
- [29] **Y Miyaoka, S Mori**, *A numerical criterion for uniruledness*, Ann. of Math. 124 (1986) 65–69 MR
- [30] **J Ross, R Thomas**, *Weighted projective embeddings, stability of orbifolds, and constant scalar curvature Kähler metrics*, J. Differential Geom. 88 (2011) 109–159 MR
- [31] **J Stoppa**, *K-stability of constant scalar curvature Kähler manifolds*, Adv. Math. 221 (2009) 1397–1408 MR
- [32] **G Tian**, *Kähler–Einstein metrics on algebraic manifolds*, from “Transcendental methods in algebraic geometry” (F Catanese, C Ciliberto, editors), Lecture Notes in Math. 1646, Springer (1996) 143–185 MR
- [33] **B Watson**, *Manifold maps commuting with the Laplacian*, J. Differential Geometry 8 (1973) 85–94 MR

*Department of Mathematics and Statistics, University of New Mexico
Albuquerque, NM, United States*

*Department of Mathematics and Statistics, University of New Mexico
Albuquerque, NM, United States*

*Institut de Mathématiques de Toulouse, Université Paul Sabatier
Toulouse, France*

cboyer@unm.edu, hnhuang@unm.edu, eveline.legendre@math.univ-toulouse.fr

Proposed: Simon Donaldson

Received: 28 August 2017

Seconded: Tobias H Colding, Jean-Pierre Otal

Accepted: 28 June 2018