

# The resolution of paracanonical curves of odd genus

GAVRIL FARKAS  
MICHAEL KEMENY

We prove the Prym–Green conjecture on minimal free resolutions of paracanonical curves of odd genus. The proof proceeds via curves lying on ruled surfaces over an elliptic curve.

14H10

## 0 Introduction

The study of torsion points on Jacobians of algebraic curves has a long history in algebraic geometry and number theory. On the one hand, torsion points of Jacobians have been used to rigidify moduli problems for curves, while on the other hand, such a torsion point determines an unramified cyclic cover over the curve in question, which gives rise to a (generalized) Prym variety; see Birkenhake and Lange [5, Chapter 12] for an introduction to this circle of ideas.

Pairs  $[C, \tau]$ , where  $C$  is a smooth curve of genus  $g \geq 2$  and  $\tau \in \text{Pic}^0(C)$  is a nontrivial torsion line bundle of order  $\ell \geq 2$ , form an irreducible moduli space  $\mathcal{R}_{g,\ell}$ . One may view this moduli space as a higher-genus analogue of the level- $\ell$  modular curve  $X_1(\ell)$ . There is a finite cover

$$\mathcal{R}_{g,\ell} \rightarrow \mathcal{M}_g$$

given by forgetting the  $\ell$ -torsion point. Following ideas going back to Mumford, Tyurin and many others, linearizing the Abel–Prym embedding of the curve in its Prym variety leads to the study of the properties of  $[C, \tau]$  in terms of the projective geometry of the level- $\ell$  *paracanonical* curve

$$\varphi_{K_C \otimes \tau}: C \hookrightarrow \mathbb{P}^{g-2}$$

induced by the line bundle  $K_C \otimes \tau$ . In practice, this amounts to a qualitative study of the equations and the syzygies of the paracanonical curve in question. For instance, in

the case  $\ell = 2$ , there is a close relationship between the study of these syzygies and the Prym map

$$\mathcal{R}_{g,2} \rightarrow \mathcal{A}_{g-1}$$

to the moduli space of principally polarized abelian varieties of dimension  $g - 1$ , which has been exploited fruitfully for some time; see for instance Beauville [3]. For higher levels, the study of these syzygies has significant applications to the study of the birational geometry of  $\mathcal{R}_{g,\ell}$ ; see Chiodo, Eisenbud, Farkas and Schreyer [7].

Denoting by  $\Gamma_C(K_C \otimes \tau) := \bigoplus_{q \geq 0} H^0(C, (K_C \otimes \tau)^{\otimes q})$  the homogeneous coordinate ring of the paracanonical curve, for integers  $p, q \geq 0$ , let

$$K_{p,q}(C, K_C \otimes \tau) := \text{Tor}^p(\Gamma_C(K_C \otimes \tau), \mathbb{C})_{p+q}$$

be the Koszul cohomology group of  $p^{\text{th}}$  syzygies of weight  $q$  of the paracanonical curve, and denote by  $b_{p,q} := \dim K_{p,q}(C, K_C \otimes \tau)$  the corresponding Betti number.

The *Prym–Green conjecture* formulated by Chiodo, Eisenbud, Farkas and Schreyer [7] predicts that the minimal free resolution of the paracanonical curve corresponding to a general level- $\ell$  curve  $[C, \tau] \in \mathcal{R}_{g,\ell}$  of genus  $g \geq 5$  is *natural*, that is, in each diagonal of its Betti table, at most one entry is nonzero. The naturality of the resolution amounts to the vanishing statements  $b_{p,2} \cdot b_{p+1,1} = 0$  for all  $p$ . As explained in [7], for odd genus  $g = 2n + 1$ , this is equivalent to the vanishing statements

$$(1) \quad K_{n-1,1}(C, K_C \otimes \tau) = 0 \quad \text{and} \quad K_{n-3,2}(C, K_C \otimes \tau) = 0.$$

Since the differences  $b_{p,2} - b_{p+1,1}$  are known, naturality entirely determines the resolution of the general level- $\ell$  paracanonical curves and shows that its Betti numbers are as small as the geometry (that is, the Hilbert function) allows. We refer the reader to Farkas and Ludwig [11] and to [7] for background on this conjecture and its important implications on the global geometry of  $\mathcal{R}_{g,\ell}$ .

In particular, a positive solution to the Prym–Green conjecture for bounded genus  $g < 23$  has been shown to be instrumental in determining the Kodaira dimension of  $\mathcal{R}_{g,\ell}$  for small values of  $\ell$ . The Prym–Green conjecture is obviously inspired by the classical Green’s conjecture for syzygies of canonical curves stating that the minimal resolution of a general canonical curve  $C \subseteq \mathbb{P}^{g-1}$  is natural. The main result of this paper is a complete solution to this conjecture in odd genus:

**Theorem 0.1** *The Prym–Green conjecture holds for any odd genus  $g$  and any level  $\ell$ .*

1	2	⋯	$n - 3$	$n - 2$	$n - 1$	$n$	⋯	$2n - 2$
$b_{1,1}$	$b_{2,1}$	⋯	$b_{n-3,1}$	$b_{n-2,1}$	0	0	⋯	0
0	0	⋯	0	$b_{n-2,2}$	$b_{n-1,2}$	$b_{n,2}$	⋯	$b_{2n-2,2}$

Table 1

**Theorem 0.1** implies the general level- $\ell$  paracanonical curve of genus  $g = 2n + 1 \geq 5$  has the minimal resolution given in **Table 1**, where

$$\begin{aligned}
 b_{p,1} &= \frac{p(2n - 2p - 3)}{2n - 1} \binom{2n}{p+1} && \text{if } p \leq n - 2, \\
 b_{p,2} &= \frac{(p + 1)(2p - 2n + 5)}{2n - 1} \binom{2n}{p+2} && \text{if } p \geq n - 2.
 \end{aligned}$$

In odd genus, we have established the conjecture before for level 2 in [9] (using Nikulin surfaces) and for high level  $\ell \geq \sqrt{(g + 2)/2}$  in [10] (using Barth–Verra surfaces). **Theorem 0.1** therefore removes any restriction on the level  $\ell$ . Apart from that, we feel that the rational elliptic surfaces used in this paper are substantially simpler objects than the  $K3$  surfaces used in [9] and [10] and should have further applications to syzygy problems. The Prym–Green conjecture in even genus, amounting to the single vanishing statement

$$(2) \quad K_{\frac{1}{2}g-2,1}(C, K_C \otimes \tau) = 0$$

(or equivalently,  $K_{\frac{1}{2}g-3,2}(C, K_C \otimes \tau) = 0$ ), is still mysterious. It is expected to hold for any genus and level  $\ell > 2$ . For level 2, it has been shown to fail in genus 8 by Colombo, Farkas, Verra and Voisin [8]; a *Macaulay* calculation carried out by Chiodo, Eisenbud, Farkas and Schreyer [7] indicates that the conjecture very likely fails in genus 16 as well. This strongly suggests that for level 2 the Prym–Green conjecture fails for general *Prym canonical* curves of genera having high divisibility properties by 2 and in these cases there should be genuinely new methods of constructing syzygies. At the moment the vanishing (2) is not even known to hold for arbitrary even genus  $g$  in the case when  $\tau$  is a general line bundle in  $\text{Pic}^0(C)$ .

By semicontinuity and the irreducibility of  $\mathcal{R}_{g,\ell}$ , it suffices to establish the vanishing (1) for one particular example of a paracanonical curve of odd genus. In our previous partial results on the Prym–Green conjecture, we constructed suitable examples  $[C, \tau]$  in terms of curves lying on various kinds of lattice-polarized  $K3$  surfaces, namely the Nikulin and Barth–Verra surfaces. In each case, the challenge lies in realizing the  $\ell$ -torsion bundle  $\tau$  as the restriction of a line bundle on the surface, so that the

geometry of the surface can be used to prove the vanishing of the corresponding Koszul cohomology groups, while making sure that the curve  $C$  in question remains general, for instance, from the point of view of Brill–Noether theory. In contrast, in this paper we use the elliptic ruled surfaces recently introduced by Farkas and Tarasca [13] (closely related to the very interesting earlier work of Treibich [18]), in order to provide explicit examples of pointed Brill–Noether general curves defined over  $\mathbb{Q}$ . These surfaces also arise when one degenerates a projectively embedded  $K3$  surface to a surface with isolated, elliptic singularities. They have been studied in detail by Arbarello, Bruno and Sernesi in their important work [2] on the classification of curves lying on  $K3$  surfaces in terms of their Wahl map.

Whereas our previous results required a different  $K3$  surface for each torsion order  $\ell$  for which the construction worked, in the current paper we deal with *all* orders  $\ell$  using a single surface. This is possible because on the elliptic ruled surface in question, a general genus- $g$  curve admits a canonical degeneration within its linear system to a singular curve consisting of a curve of genus  $g - 1$  and an elliptic tail. This leads to an inductive structure involving curves of every genus and makes possible inductive arguments, while working on the same surface all along.

We introduce the elliptic ruled surface central to this paper. For an elliptic curve  $E$ , set

$$\phi: X := \mathbb{P}(\mathcal{O}_E \oplus \eta) \rightarrow E,$$

where  $\eta \in \text{Pic}^0(E)$  is neither trivial nor torsion. We fix an origin  $a \in E$  and let  $b \in E$  be such that  $\eta = \mathcal{O}_E(a - b)$ . Furthermore, choose a point  $r \in E \setminus \{b\}$  such that  $\zeta := \mathcal{O}_E(b - r)$  is torsion of order precisely  $\ell$ . The scroll  $X \rightarrow E$  has two sections  $J_0$  and  $J_1$  corresponding respectively to the quotients  $\mathcal{O}_E \oplus \eta \twoheadrightarrow \eta$  and  $\mathcal{O}_E \oplus \eta \twoheadrightarrow \mathcal{O}_E$ . We have

$$J_1 \cong J_0 - \phi^* \eta, \quad N_{J_0/X} \cong \mathcal{O}_{J_0}(\phi^* \eta), \quad N_{J_1/X} \cong \mathcal{O}_{J_1}(\phi^* \eta^\vee),$$

where we freely mix notation for divisors and line bundles. For any point  $x \in E$  we denote by  $f_x$  the fiber  $\phi^{-1}(x)$ . We let

$$C \in |gJ_0 + f_r|$$

be a general element; this is a smooth curve of genus  $g$ . We further set

$$L := \mathcal{O}_X((g - 2)J_0 + f_a).$$

Using that  $K_X = -J_0 - J_1$ , the adjunction formula shows that the restriction  $L_C$  is a level- $\ell$  paracanonical bundle on  $C$ , that is,  $[C, \tau] \in \mathcal{R}_{g, \ell}$ , where  $\tau := \phi_C^*(\zeta) \cong L_C \otimes K_C^\vee$ .

with  $\phi_C: C \rightarrow E$  being the restriction of  $\phi$  to the curve  $C$ . In this paper we verify the Prym–Green conjecture for this particular paracanonical curve of genus  $g = 2n + 1$ .

Denoting by  $\tilde{X}$  the blow-up of  $X$  at the two basepoints of  $|L|$  and by  $\tilde{L} \in \text{Pic}(\tilde{X})$  the proper transform of  $L$ , one begins by showing that the first vanishing required in the Prym–Green conjecture,  $K_{n-1,1}(C, K_C \otimes \tau) = 0$ , is a consequence of the vanishing of  $K_{n-1,1}(\tilde{X}, \tilde{L})$  and that of the mixed Koszul cohomology group  $K_{n-2,2}(\tilde{X}, -C, \tilde{L})$  (see Section 1 for details). By the Lefschetz hyperplane principle in Koszul cohomology, the vanishing of  $K_{n-1,1}(\tilde{X}, \tilde{L})$  is a consequence of Green’s conjecture for a general curve  $D$  in the linear system  $|L|$  on  $X$ . Since  $D$  has been proven by Farkas and Tarasca [13] to be Brill–Noether general, Green’s conjecture holds for  $D$ . We then show (see (8)) that a sufficient condition for the second vanishing appearing in (1) is that

$$K_{n-2,2}(D, \mathcal{O}_D(-C), K_D) = 0 \quad \text{and} \quad K_{n-1,2}(D, \mathcal{O}_D(-C), K_D) = 0.$$

Via results from Farkas, Mustața and Popa [12] coupled with the usual description of Koszul cohomology in terms of kernel bundles, we prove that these vanishings are both consequences of the following transversality statement between difference varieties in the Jacobian  $\text{Pic}^2(D)$ :

$$(3) \quad \mathcal{O}_D(C) - K_D - D_2 \not\subseteq D_n - D_{n-2},$$

where, as usual,  $D_m$  denotes the  $m^{\text{th}}$  symmetric product of  $D$  (see Lemma 1.7). This last statement is proved inductively, using the canonical degeneration of  $D$  inside its linear system to a curve of lower genus with elliptic tails. It is precisely this feature of the elliptic surface  $X$ , of containing Brill–Noether general curves of every genus (something which is not shared by a  $K3$  surface), which makes the proof possible. To sum up this part of the proof, we point that by using the geometry of  $X$ , we reduce the first half of the Prym–Green conjecture, that is, the statement  $K_{n-1,1}(C, K_C \otimes \tau) = 0$  on the curve  $C$  of genus  $g$ , to the geometric condition (3) on the curve  $D$  of genus  $g - 2$ .

The second vanishing required by the Prym–Green conjecture,  $K_{n-3,2}(C, K_C \otimes \tau) = 0$ , falls in the range covered by the secant conjecture of Green and Lazarsfeld [16]. This feature appears only in odd genus; for even genus the Prym–Green conjecture is beyond the range in which the secant conjecture applies (see Section 2 for details). For a curve  $C$  of genus  $g = 2n + 1$  and maximal Clifford index  $\text{Cliff}(C) = n$ , the secant conjecture predicts that for a nonspecial line bundle  $L \in \text{Pic}^{2g-2}(C)$ , one has the equivalence

$$K_{n-3,2}(C, L) = 0 \iff L - K_C \notin C_{n-1} - C_{n-1}.$$

Despite significant progress, the secant conjecture is not known for arbitrary  $L$ , but in [9, Theorem 1.7], we provided a sufficient condition for the vanishing to hold. Precisely, whenever

$$(4) \quad \tau + C_2 \not\subseteq C_{n+1} - C_{n-1},$$

we have  $K_{n-3,2}(C, K_C \otimes \tau) = 0$ . Thus the second half of the Prym–Green conjecture has been reduced to a transversality statement of difference varieties very similar to (3), but this time on the same curve  $C$ . Using the already mentioned elliptic tail degeneration inside the linear system  $|C|$  on  $X$ , we establish inductively in Section 2 that (4) holds for a general curve  $C \subseteq X$  in its linear system. This completes the proof of the Prym–Green conjecture.

**Acknowledgements** Farkas is supported by DFG Priority Program 1489 *Algorithmische und experimentelle Methoden in Algebra, Geometrie und Zahlentheorie*. Kemeny is supported by NSF grant DMS-1701245 *Syzygies, moduli spaces, and Brill–Noether theory*.

## 1 Elliptic surfaces and paracanonical curves

We fix a level  $\ell \geq 2$  and recall that pairs  $[C, \tau]$ , where  $C$  is a smooth curve of genus  $g$  and  $\tau \in \text{Pic}^0(C)$  is an  $\ell$ -torsion point, form an irreducible moduli space  $\mathcal{R}_{g,\ell}$ . We refer the reader to [7] for a detailed description of the Deligne–Mumford compactification  $\overline{\mathcal{R}}_{g,\ell}$  of  $\mathcal{R}_{g,\ell}$ .

Normally we prefer multiplicative notation for line bundles, but occasionally, in order to simplify calculations, we switch to additive notation and identify divisors and line bundles. If  $V$  is a vector space and  $S := \text{Sym } V$ , for a graded  $S$ -module  $M$  of finite type, we denote by  $K_{p,q}(M, V)$  the Koszul cohomology group of  $p^{\text{th}}$  syzygies of weight  $q$  of  $M$ . If  $X$  is a projective variety,  $L$  is a line bundle and  $\mathcal{F}$  is a sheaf on  $X$ , we set as usual  $K_{p,q}(X, \mathcal{F}, L) := K_{p,q}(\Gamma_X(\mathcal{F}, L), H^0(X, L))$ , where  $\Gamma_X(\mathcal{F}, L) := \bigoplus_{q \in \mathbb{Z}} H^0(X, \mathcal{F} \otimes L^{\otimes q})$  is viewed as a graded  $\text{Sym } H^0(X, L)$ -module. For background questions on Koszul cohomology, we refer the reader to the book [1].

Assume now that  $g := 2n + 1$  is odd and let us consider the decomposable elliptic ruled surface  $\phi: X \rightarrow E$  defined in the introduction. Retaining all the notation, our first aim is to establish the vanishing of the linear syzygy group  $K_{n-1,1}(C, K_C \otimes \tau)$ . Before proceeding, we confirm that  $\tau := \phi_C^*(\zeta)$  is nontrivial of order precisely  $\ell$ , so that  $[C, \tau]$  is indeed a point of  $\mathcal{R}_{g,\ell}$ .

**Lemma 1.1** For any  $1 \leq m \leq \ell - 1$ , the line bundle  $\tau^{\otimes m} \in \text{Pic}^0(C)$  is not effective.

**Proof** Since the order of  $\zeta$  is precisely  $\ell$ , we have

$$H^0(X, \phi^*(\zeta^{\otimes m})) \cong H^0(E, \zeta^{\otimes m}) = 0 \quad \text{for } 1 \leq m \leq \ell - 1.$$

So it suffices to show  $H^1(X, \phi^*(\zeta^{\otimes m})(-C)) = 0$ . By Serre duality, this is equivalent to  $H^1(X, \phi^*(r + \eta - m\zeta)((g - 2)J_0)) = 0$ . Applying the Leray spectral sequence this amounts to

$$H^1(E, \mathcal{O}_E(a + (m + 1)r - (m + 1)b) \otimes \text{Sym}^{g-2}(\mathcal{O}_E \oplus \eta)) = 0,$$

which is clear for degree reasons. □

Any linear system which is a sum of a positive multiple of  $J_0$  and a fiber of  $\phi$  has two basepoints; see [13, Lemma 2]. In particular, the linear system  $|L|$  on  $X$  has two basepoints  $p \in J_1$  and  $q^{(g-2)} \in J_0$ . Here

$$\{p\} := f_a \cdot J_1 \quad \text{and} \quad \{q^{(g-2)}\} := f_{s^{(g-2)}} \cdot J_0,$$

where the point  $s^{(g-2)} \in E$  is determined by the condition  $\eta^{\otimes(g-2)} \cong \mathcal{O}_E(s^{(g-2)} - a)$ .

Let  $\pi: \tilde{X} \rightarrow X$  be the blow-up of  $X$  at these two basepoints, with exceptional divisors  $E_1$  respectively  $E_2$  over  $p$  respectively  $q^{(g-2)}$ . We denote by  $\tilde{L} := \pi^*L - E_1 - E_2$  the proper transform of  $L$ . Note that  $K_{\tilde{X}} = -\tilde{J}_0 - \tilde{J}_1$ , where  $\tilde{J}_0 = J_0 - E_2$  and  $\tilde{J}_1 = J_1 - E_1$  are the proper transforms of  $J_0$  and  $J_1$ . We now observe that the basepoints of the two linear systems  $|L|$  and  $|C|$  on  $X$  are disjoint.

**Lemma 1.2** Let  $x_0 \in J_0$  and  $x_1 \in J_1$  be the two basepoints of  $|C|$ . Then we have  $x_0, x_1 \notin \{p, q^{(g-2)}\}$ .

**Proof** First, since  $r \neq a$ , we obtain  $J_1 \cap f_a \neq J_1 \cap f_r$ , therefore  $p \neq x_1$ . Next, recall that  $\{q^{(g-2)}\} = J_0 \cap f_{s^{(g-2)}}$ , where  $\mathcal{O}_E(s^{(g-2)} - a) = \eta^{\otimes(g-2)}$  and  $\{x_0\} = J_0 \cap f_{t^{(g)}}$ , where the point  $t^{(g)} \in E$  is determined by the equation  $\mathcal{O}_E(t^{(g)} - r) = \eta^{\otimes g}$ . We need to show  $\eta^{\otimes(g-2)}(a) \neq \eta^{\otimes g}(r)$ . Else, since  $\mathcal{O}_E(a - r) = \eta \otimes \zeta$ , it would imply  $\zeta = \eta$ , which is impossible, for  $\zeta$  is a torsion class, whereas  $\eta$  is not. □

Since the curve  $C$  does not pass through the points  $p$  and  $q^{(g-2)}$  which are blown-up, we shall abuse notation by writing  $C$  for  $\pi^*(C)$ . We set  $S := \text{Sym } H^0(\tilde{X}, \tilde{L})$  and consider the short exact sequence of graded  $S$ -modules

$$0 \rightarrow \bigoplus_{q \in \mathbb{Z}} H^0(\tilde{X}, q\tilde{L} - C) \rightarrow \bigoplus_{q \in \mathbb{Z}} H^0(\tilde{X}, q\tilde{L}) \rightarrow M \rightarrow 0,$$

where the first map is defined by multiplication with the section defining  $C$  and the module  $M$  is defined by this exact sequence. By the corresponding long exact sequence in Koszul cohomology [15, Corollary 1.d.4], that is,

$$\cdots \rightarrow K_{p,1}(\tilde{X}, \tilde{L}) \rightarrow K_{p,1}(M, H^0(\tilde{X}, \tilde{L})) \rightarrow K_{p-1,2}(\tilde{X}, -C, \tilde{L}) \rightarrow \cdots,$$

the vanishing of the Koszul cohomology group  $K_{p,1}(M, H^0(\tilde{X}, \tilde{L}))$  follows from  $K_{p,1}(\tilde{X}, \tilde{L}) = 0$  and  $K_{p-1,2}(\tilde{X}, -C, \tilde{L}) = 0$ . The reason we are interested in the Koszul cohomology of  $M$  becomes apparent in the following lemma:

**Lemma 1.3** *The equality  $K_{p,1}(M, H^0(\tilde{X}, \tilde{L})) \cong K_{p,1}(C, K_C \otimes \tau)$  holds for any  $p \geq 0$ .*

**Proof** The restriction map induces an isomorphism  $H^0(\tilde{X}, \tilde{L}) \cong H^0(C, K_C \otimes \tau)$ . First, the restriction map is injective, since  $\tilde{L} - C = \pi^*(-2J_0 + f_a - f_r) - E_1 - E_2$  is not effective (as it has negative intersection with the nef class  $\pi^*(f_r)$ ). Next,  $h^0(\tilde{X}, \tilde{L}) = h^0(X, L) = g - 1$  by a direct computation using the projection formula; see also [13, Lemma 2]. As  $h^0(C, K_C \otimes \tau) = g - 1$ , the restriction to  $C \subseteq \tilde{X}$  induces the claimed isomorphism.

Let  $M_q$  denote the  $q^{\text{th}}$  graded piece of  $M$ . We have an isomorphism  $M_0 \cong H^0(\tilde{X}, \mathcal{O}_{\tilde{X}})$  and have already seen that  $H^0(\tilde{X}, \tilde{L} - C) = 0$ , so  $M_1 \cong H^0(\tilde{X}, \tilde{L}) \cong H^0(C, K_C \otimes \tau)$ . So we have the commutative diagram

$$\begin{array}{ccc} \wedge^{p+1} H^0(\tilde{L}) \otimes M_0 & \longrightarrow & \wedge^{p+1} H^0(K_C + \tau) \\ \downarrow & & \downarrow \\ \wedge^p H^0(\tilde{L}) \otimes M_1 & \longrightarrow & \wedge^p H^0(K_C + \tau) \otimes H^0(K_C + \tau) \\ \downarrow \delta_1 & & \downarrow \delta'_1 \\ \wedge^{p-1} H^0(\tilde{L}) \otimes M_2 & \longrightarrow & \wedge^{p-1} H^0(K_C + \tau) \otimes H^0(2K_C + 2\tau) \end{array}$$

where the two topmost horizontal maps are isomorphisms and the bottommost horizontal map is injective. Thus the middle cohomology of the left-hand column is isomorphic to that of the right, so that we have the equality  $K_{p,1}(M, H^0(\tilde{X}, \tilde{L})) \cong K_{p,1}(C, K_C \otimes \tau)$ , for any  $p \geq 0$ . □

### 1.1 The vanishing of the Koszul cohomology group $K_{n-1,1}(C, K_C \otimes \tau)$

We can summarize the discussion so far. In order to establish the first vanishing required by the Prym–Green conjecture for the pair  $[C, \tau]$ , that is,  $K_{n-1,1}(C, K_C \otimes \tau) = 0$ , it



suffices to prove that

$$(5) \quad K_{n-1,1}(\tilde{X}, \tilde{L}) = 0,$$

$$(6) \quad K_{n-2,2}(\tilde{X}, -C, \tilde{L}) = 0.$$

Vanishing (5) is a consequence of Green’s conjecture on syzygies of canonical curves.

**Proposition 1.4** *We have  $K_{n-1,1}(\tilde{X}, \tilde{L}) = 0$ .*

**Proof** Let  $D \in |\tilde{L}|$  be a general element. Thus  $D$  is a smooth curve of genus  $2n - 1$ . We have an isomorphism  $K_{n-1,1}(\tilde{X}, \tilde{L}) \cong K_{n-1,1}(D, K_D)$ , as  $K_{\tilde{X}|_D} \cong \mathcal{O}_D$  and by applying [1, Theorem 2.20] (note that one only needs that the restriction  $H^0(\tilde{X}, \tilde{L}) \rightarrow H^0(D, K_D)$  is surjective, and not  $H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) = 0$ , for this result). As  $D$  is a smooth curve of genus  $2n - 1$ , the vanishing in question is a consequence of Green’s conjecture, which is known to hold for curves of maximal gonality; see [19; 17]. Hence it suffices to show that  $D$  has maximum gonality  $n + 1$ . But  $D$  is the strict transform of a smooth curve in  $|L|$  and is a Brill–Noether general curve by [13, Remark 2]; in particular, it has maximal gonality.  $\square$

We now turn to the vanishing of the second Koszul group  $K_{n-2,2}(\tilde{X}, -C, \tilde{L})$ . The following argument is inspired by [15, Theorem 3.b.7].

**Proposition 1.5** *Let  $D \in |\tilde{L}|$  be general and let  $p \geq 0$ . Assume*

$$K_{m,2}(D, \mathcal{O}_D(-C), K_D) = 0 \quad \text{for } m \in \{p, p + 1\}.$$

Then

$$K_{p,2}(\tilde{X}, -C, \tilde{L}) = 0.$$

**Proof** Set as before  $S := \text{Sym } H^0(\tilde{X}, \tilde{L})$  and consider the exact sequence of graded  $S$ -modules

$$0 \rightarrow \bigoplus_{q \in \mathbb{Z}} H^0(\tilde{X}, (q - 1)\tilde{L} - C) \rightarrow \bigoplus_{q \in \mathbb{Z}} H^0(\tilde{X}, q\tilde{L} - C) \rightarrow B \rightarrow 0,$$

serving as a definition for  $B$ , and where the first map is given by multiplication by a general section  $s \in H^0(\tilde{X}, \tilde{L})$ . We now argue along the lines of [9, Lemma 2.2]. Taking the long exact sequence in Koszul cohomology and using that multiplication by a section  $s \in H^0(\tilde{X}, \tilde{L})$  induces the zero map on Koszul cohomology, we get

$$K_{p,q}(B, H^0(\tilde{X}, \tilde{L})) \cong K_{p,q}(\tilde{X}, -C, \tilde{L}) \oplus K_{p-1,q}(\tilde{X}, -C, \tilde{L}),$$

for all  $p, q \in \mathbb{Z}$ .

Let  $D = Z(s)$  be the divisor defined by  $s \in H^0(\tilde{X}, \tilde{L})$ , and consider the graded  $S$ -module

$$N := \bigoplus_{q \in \mathbb{Z}} H^0(D, qK_D - C_D).$$

We have the inclusion  $B \subseteq N$  of graded  $S$ -modules. We claim  $B_1 = N_1 = 0$ . By intersecting with the nef class  $f_r$ , we see  $H^0(\tilde{X}, \tilde{L} - C) = 0$ , implying  $B_1 = 0$ . As  $\deg(K_D - C_D) = -4$ , we have  $N_1 = 0$ . Upon taking Koszul cohomology, this immediately gives the inclusion

$$K_{p,2}(B, H^0(\tilde{X}, \tilde{L})) \subseteq K_{p,2}(N, H^0(\tilde{X}, \tilde{L})).$$

In particular,  $K_{p,2}(\tilde{X}, -C, \tilde{L}) \subseteq K_{p+1,2}(B, H^0(\tilde{X}, \tilde{L})) \subseteq K_{p+1,2}(N, H^0(\tilde{X}, \tilde{L}))$ .

To finish the proof, it will suffice to show

$$(7) \quad K_{p,2}(N, H^0(\tilde{X}, \tilde{L})) \cong K_{p,2}(D, \mathcal{O}_D(-C), K_D) \oplus K_{p-1,2}(D, \mathcal{O}_D(-C), K_D).$$

Since  $\tilde{L} \cdot \tilde{J}_0 = 0$  and  $\tilde{L} \cdot \tilde{J}_1 = 0$ , it follows that  $\mathcal{O}_D(K_{\tilde{X}}) \cong \mathcal{O}_D$ . We now closely follow the proof of [9, Lemma 2.2]. The section  $s$  induces a splitting  $H^0(\tilde{X}, \tilde{L}) \cong \mathbb{C}\{s\} \oplus H^0(D, K_D)$ , giving rise, for every  $p$ , to isomorphisms

$$\wedge^p H^0(\tilde{X}, \tilde{L}) \cong \wedge^{p-1} H^0(D, K_D) \oplus \wedge^p H^0(D, K_D).$$

The desired isomorphism (7) follows from a calculation which is identical to the one carried out in the second part of the proof of [9, Lemma 2.2]. There one works with a  $K3$  surface, but the only thing needed for the argument to work is that  $\mathcal{O}_D(K_{\tilde{X}}) \cong \mathcal{O}_D$ .  $\square$

To establish that  $K_{n-1,1}(C, K_C \otimes \tau) = 0$ , it thus suffices to show

$$(8) \quad K_{n-2,2}(D, \mathcal{O}_D(-C), K_D) = 0 \quad \text{and} \quad K_{n-1,2}(D, \mathcal{O}_D(-C), K_D) = 0.$$

Via a well-known description of Koszul cohomology using kernel bundles (see [1, Proposition 2.5]), taking into account that  $H^0(D, K_D - C_D) = 0$ , these two statements are equivalent to

$$(9) \quad \begin{aligned} H^0(D, \wedge^{n-2} M_{K_D} \otimes (2K_D - C_D)) &= 0, \\ H^0(D, \wedge^{n-1} M_{K_D} \otimes (2K_D - C_D)) &= 0, \end{aligned}$$

where we recall that  $M_{K_D}$  is the kernel bundle, defined by the short exact sequence

$$0 \rightarrow M_{K_D} \rightarrow H^0(D, K_D) \otimes \mathcal{O}_D \rightarrow K_D \rightarrow 0.$$

Both statements (9) will be reduced to general position statements with respect to divisorial difference varieties of the various curves on  $X$ .

### 1.2 Containment between difference varieties on curves

If  $C$  is a smooth curve of genus  $g$ , we denote by  $C_a - C_b \subseteq \text{Pic}^{a-b}(C)$  the image of the difference map  $v: C_a \times C_b \rightarrow \text{Pic}^{a-b}(C)$ . We occasionally make use of the realization given in [12] of the *divisorial* difference varieties as nonabelian theta divisors associated to exterior powers of the kernel bundle of  $K_C$ . Precisely, for  $i = 0, \dots, \lfloor (g-1)/2 \rfloor$ , one has the following equality of divisors on  $\text{Pic}^{g-2i-1}(C)$ :

$$(10) \quad C_{g-i-1} - C_i = \{ \xi \in \text{Pic}^{g-2i-1}(C) : H^0(C, \wedge^i M_{K_C} \otimes K_C \otimes \xi^\vee) \neq 0 \}.$$

We now make an observation concerning a containment relation between difference varieties.

**Lemma 1.6** *Let  $C$  be a smooth curve, let  $a \geq 2, b \geq 0, c > 0$  be integers, and let  $A \in \text{Pic}^{a+b-c}(C)$ . Assume  $A - C_a \subseteq C_b - C_c$ . Then  $A - C_{a-2} \subseteq C_{b+1} - C_{c-1}$ .*

**Proof** Let  $B$  be an arbitrary effective divisor of degree  $a - 2$ , and let  $y_0 \in C$  be a fixed point. Since  $A - C_a \subseteq C_b - C_c$ , we have a well-defined morphism

$$f: C \rightarrow C_b - C_c \subseteq \text{Pic}^{b-c}(C), \quad x \mapsto A - (B + x + y).$$

We further have the difference map  $v: C_b \times C_c \rightarrow \text{Pic}^{b-c}(C)$  given by  $v(F_1, F_2) := \mathcal{O}_C(F_1 - F_2)$ , where  $F_1$  and  $F_2$  are effective divisors of degrees  $b$  and  $c$  respectively, as well as the projection  $p_2: C_b \times C_c \rightarrow C_c$ .

Suppose first that  $\dim p_2(v^{-1}(\text{Im}(f))) \geq 1$ . As the divisor  $y_0 + C_{c-1} \subseteq C_c$  is ample — see [14, Lemma 2.7] —  $p_2(v^{-1}(\text{Im}(f)))$  must meet  $y_0 + C_{c-1}$ . This means that there exists a point  $x \in C$  such that  $A - (B + x + y) \equiv F_1 - F_2$ , with  $F_1 \in C_b$  and  $F_2 \in C_c$  being effective divisors such that  $F_2 = y_0 + F'_2$ , where  $F'_2 \in C_{c-1}$  is effective. But then

$$A - B = \mathcal{O}_C(F_1 + x - F'_2) \in C_{b+1} - C_{c-1}.$$

Assume now  $p_2(v^{-1}(\text{Im}(f))) \subseteq C_c$  is finite. Then one can find a divisor  $F_2 \in C_c$ , such that for every  $x \in C$ , there is a divisor  $F_x \in C_b$  with  $A - B - x - y = F_x - F_2$ . Picking  $x \in \text{supp}(F_2)$ , we write  $F_2 = x + F'_2$ , where  $F'_2 \in C_{c-1}$ . Then  $A - B = \mathcal{O}_C(F_x + y_0 - F'_2) \in C_{b+1} - C_{c-1}$ . □

We may now restate the vanishing conditions (8) in terms of difference varieties. From now on we revert to the elliptic surface  $\phi: X \rightarrow E$  and recall that  $C \in |gJ_0 + f_r|$ .

**Lemma 1.7** Set  $g = 2n + 1$  with  $n \geq 2$  and choose a general curve  $D \in |(g-2)J_0 + f_a|$ . Suppose

$$C_D - K_D - D_2 \not\subseteq D_n - D_{n-2}.$$

Then  $K_{n-1,1}(C, K_C \otimes \tau) = 0$ , for a general level- $\ell$  curve  $[C, \eta] \in \mathcal{R}_{g,\ell}$ .

**Proof** By assumption, there exist points  $x, y \in D$  such that  $C_D - K_D - x - y$  is not in  $D_n - D_{n-2}$ . It follows from (10) that this is equivalent to

$$H^0(D, \wedge^{n-2} M_{K_D} \otimes (2K_D - C_D + x + y)) = 0,$$

implying

$$H^0(D, \wedge^{n-2} M_{K_D} \otimes (2K_D - C_D)) = 0.$$

This is equivalent to

$$K_{n-2,2}(D, \mathcal{O}_D(-C), K_D) = 0.$$

Next, by Lemma 1.6, our assumption implies  $C_D - K_D - D_4 \not\subseteq D_{n-1} - D_{n-1}$ . Thus  $H^0(D, \wedge^{n-1} M_{K_D} \otimes (2K_D - C_D + T)) = 0$ , for some effective divisor  $T \in D_4$ , and therefore  $H^0(D, \wedge^{n-1} M_{K_D} \otimes (2K_D - C_D)) = 0$  as well, amounting to

$$K_{n-1,2}(D, \mathcal{O}_D(-C), K_D) = 0. \quad \square$$

Any smooth divisor  $D \in |L|$  carries two distinguished points, namely  $p$  and  $q^{(g-2)}$ . We will prove that if  $D \in |L|$  is general, then

$$(11) \quad C_D - K_D - p - q^{(g-2)} \notin D_n - D_{n-2}.$$

Let us first introduce some notation. For an integer  $m \geq 1$ , we define the line bundle

$$L_m := \mathcal{O}_X(mJ_0 + f_a) \in \text{Pic}(X).$$

A general element  $D \in |L_m|$  is a smooth curve of genus  $m$ , having two distinguished points  $p \in J_1$  and  $q^{(m)} \in J_0$ , which as already explained, are the basepoints of  $|L_m|$ . Recall that for each  $j = 0, \dots, m - 1$ , we introduced the divisorial difference variety

$$D_j - D_{m-1-j} \subseteq \text{Pic}^{2j+1-m}(D).$$

This difference variety is empty for  $j < 0$  or  $j > m - 1$ . We shall prove (11) inductively by contradiction, using the fact that in any family of curves on the surface  $X$ , there is a canonical degeneration to a curve with an elliptic tail.

### 1.3 The induction step

Assume that for a general curve  $D \in |L_{g-2-j}|$  one has, for some  $0 \leq i \leq j$ ,

$$C_D - K_D - p - (2i + 1)q^{(g-2-j)} \in D_{n-i} - D_{n-2-j+i}.$$

Then for a general curve  $Z \in |L_{g-3-j}|$  one has, for some  $0 \leq i' \leq j + 1$ ,

$$(12) \quad C_Z - K_Z - p - (2i' + 1)q^{(g-3-j)} \in Z_{n-i'} - Z_{n-3-j+i'}.$$

Notice that the assumption  $D_{n-i} - D_{n-2-j+i} \neq \emptyset$  for a curve  $D \in |L_{g-2-j}|$  implies

$$(13) \quad 0 \leq n - i \leq g - 3 - j.$$

Let  $D \in |L_{g-2-j}|$  be general. In order to prove the induction step, we degenerate  $D$  within its linear system to the curve of compact type

$$Y := J_0 + Z,$$

for a general  $Z \in |L_{g-3-j}|$ . Notice that  $J_0 \cdot Z = q^{(g-3-j)} =: q$  and the marked point  $p$  lies on  $Z \setminus \{q\}$ . On  $Y$ , in the spirit of limit linear series, we choose the twist of bidegree  $(0, 2g - 2j - 6)$  of its dualizing sheaf, that is, the line bundle

$$\tilde{K} \in \text{Pic}(Y)$$

characterized by  $\tilde{K} \otimes \mathcal{O}_{J_0} \cong \mathcal{O}_{J_0}$  and  $\tilde{K} \otimes \mathcal{O}_Z \cong K_Z(2q)$ . We establish a few technical statements to be used later in the proofs.

**Lemma 1.8** *Assume the bounds (13). Then, for any  $0 \leq i \leq j \leq g - 4$ , we have:*

- (i)  $h^0(Y, \tilde{K}) = h^0(D, K_D) = g - 2 - j$ .
- (ii)  $H^0(Y, \mathcal{O}_Y(C - J_1 - (2i + 1)J_0) \otimes \tilde{K}^\vee) = 0$ .
- (iii)  $h^0(Y, \mathcal{O}_Y(C - J_1 - (2i + 1)J_0)) = h^0(D, C_D(-p - (2i + 1)q^{(g-2-j)}))$ .
- (iv)  $h^0(Y, \mathcal{O}_Y(C - J_1 - (2i + 1)J_0) \otimes \tilde{K}) = h^0(D, C_D \otimes K_D(-p - (2i + 1)q^{(g-2-j)}))$ .

**Proof** (i) Because  $\tilde{K}$  is a limit of canonical bundles on smooth curves, we have  $h^0(Y, \tilde{K}) \geq g - 2 - j = h^0(D, K_D)$ . So it suffices to show  $h^0(Y, \tilde{K}) \leq h^0(D, K_D)$ . Twisting by  $\tilde{K}$  the short exact sequence

$$(14) \quad 0 \rightarrow \mathcal{O}_{J_0}(-q) \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Z \rightarrow 0$$

and taking cohomology, we get  $h^0(Y, \tilde{K}) \leq h^0(Z, K_Z(2q)) = g - 2 - j$ , as required.

(ii) Set  $A_d := \mathcal{O}_Y(C - J_1 - (2i + 1)J_0) \otimes \tilde{K}^{\otimes d} \in \text{Pic}(Y)$ . One needs to show  $H^0(Y, A_{-1}) = 0$ . Via the projection  $\phi: X \rightarrow E$  we identify the section  $J_0$  with the elliptic curve  $E$ . We have  $\mathcal{O}_{J_0}(A_{-1}) \cong \eta^{\otimes(g-2i-1)}(r)$ . Furthermore  $\mathcal{O}_{J_0}(q) \cong \eta^{\otimes(g-3-j)}(a)$ , hence

$$\mathcal{O}_{J_0}(A_{-1}(-q)) \cong \eta^{\otimes(j-2i+2)}(r - a).$$

We have  $H^0(E, \eta^{\otimes(j-2i+2)}(r - a)) = H^0(E, \zeta^\vee \otimes \eta^{\otimes(j-2i+1)}) = 0$ , for  $\zeta$  is  $\ell$ -torsion, whereas  $\eta$  is not a torsion bundle. From the short exact sequence (14) twisted by  $A_{-1}$ , in order to conclude it suffices to show that the restricted line bundle

$$\begin{aligned} \mathcal{O}_Z(A_{-1}) &\cong \mathcal{O}_Z((g - 2i - 3)J_0 - J_1 + f_r) \otimes K_Z^\vee \\ &\cong \mathcal{O}_Z((j + 1 - 2i)J_0 + f_r - f_a) \end{aligned}$$

is not effective. We will first show  $H^0(X, (j + 1 - 2i)J_0 + f_r - f_a) = 0$ . If  $j + 1 - 2i < 0$ , this is immediate since then  $((j + 1 - 2i)J_0 + f_r - f_a) \cdot f_r < 0$  and the curve  $f_r$  is nef. If  $j + 1 - 2i \geq 0$ , we use the isomorphism

$$H^0(X, (j + 1 - 2i)J_0 + f_r - f_a) \cong H^0(E, \mathcal{O}_E(r - a) \otimes \text{Sym}^{j+1-2i}(\mathcal{O}_E \oplus \eta)) = 0.$$

In order to conclude, it is enough to show  $H^1(X, (j + 1 - 2i)J_0 + f_r - f_a - Z) = 0$ . By Serre duality, this is equivalent to

$$H^1(X, K_X + Z + f_a - f_r - (j + 1 - 2i)J_0) = 0.$$

We compute

$$K_X + Z + f_a - f_r - (j + 1 - 2i)J_0 = (g - 6 + 2i - 2j)J_0 + \phi^*\eta + 2f_a - f_r,$$

where  $g - 6 + 2i - 2j \geq -1$  by (13). If  $g - 6 + 2i - 2j \geq 0$ , then

$$\begin{aligned} H^1(X, (g - 6 + 2i - 2j)J_0 + \phi^*\eta + 2f_a - f_r) \\ = H^1(E, \mathcal{O}_E(2a - r + \eta) \otimes \text{Sym}^{g-6+2i-2j}(\mathcal{O}_E \oplus \eta)), \end{aligned}$$

which vanishes for degree reasons. Finally, if  $g - 6 + 2i - 2j = -1$ , an application of the Leray spectral sequence implies  $H^1(X, -J_0 + \phi^*\eta + 2f_a - f_r) = 0$  as well. This completes the proof of (ii).

(iii) By Riemann–Roch and semicontinuity, it suffices to show  $H^1(Y, A_0) = 0$ , that is,

$$H^1(Y, \mathcal{O}_Y((g - 2i - 1)J_0 - J_1 + f_r)) = 0.$$

If so, then the bundle  $\mathcal{O}_X(C - J_1 - (2i + 1)J_0)$  has the same number of sections, when restricted to a general element  $D \in |L_{g-2-j}|$  or to its codimension 1 degen-

eration  $Y$  in its linear system. By (13), we have  $g - 2i - 1 \geq 0$ . First, starting from  $H^1(X, -J_1 + f_r) = 0$ , which is an easy consequence of the Leray spectral sequence, one shows inductively that  $H^1(X, mJ_0 - J_1 + f_r) = 0$  for all  $m \geq 0$ , in particular also  $H^1(X, (g - 2i - 1)J_0 - J_1 + f_r) = 0$ .

To conclude, it is enough to show  $H^2(X, (g - 2i - 1)J_0 - J_1 + f_r - Y) = 0$ . By Serre duality,

$$H^2(X, (g - 2i - 1)J_0 - J_1 + f_r - Y) \cong H^0(X, (2i - 2 - j)J_0 + f_a - f_r)^\vee.$$

If  $2i - 2 - j < 0$ , then the class  $(2i - 2 - j)J_0 + f_a - f_r$  is not effective on  $X$  as it has negative intersection with  $f_r$ . If  $2i - 2 - j \geq 0$  then this class is not effective by projecting to  $E$ .

(iv) It suffices to show  $H^1(Y, A_1) = 0$ . We use the exact sequence on  $Y$

$$0 \rightarrow \mathcal{O}_Z(-q) \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_{J_0} \rightarrow 0.$$

As  $\text{deg } \mathcal{O}_{J_1}(A_1) = 1$ , it is enough to show  $H^1(Z, \mathcal{O}_Z(A_1)(-q)) = 0$ . By direct computation,

$$\text{deg } \mathcal{O}_Z(A_1(-q)) = \text{deg } K_Z + 2n - 2i + g - 3 - j.$$

From (13),  $n - i \geq 0$ , whereas  $j \leq g - 4$  by assumption, so  $g - 3 - j > 0$  and the required vanishing follows for degree reasons.  $\square$

We now have all the pieces needed to prove the induction step. The transversality statement (3) that the first half of the Prym–Green conjecture has been reduced to is proved inductively, by being part of a system of conditions involving difference varieties of curves of every genus on the surface  $X$ .

**Proposition 1.9** Fix  $0 \leq j \leq g - 3$  and assume that for a general curve  $D \in |L_{g-2-j}|$ ,

$$C_D - K_D - p - (2i + 1)q^{(g-2-j)} \in D_{n-i} - D_{n-2-i+j} \quad \text{for some } 0 \leq i \leq j.$$

Then for a general curve  $Z \in |L_{g-3-j}|$ , the following holds:

$$C_Z - K_Z - p - (2i' + 1)q^{(g-3-j)} \in Z_{n-i'} - Z_{n-3-j+i'} \quad \text{for some } 0 \leq i' \leq j + 1.$$

**Proof** Using the determinantal realization of divisorial varieties (10) coming from [12], the assumption may be rewritten as

$$H^0(D, \wedge^{n-2-j+i} M_{K_D}^\vee \otimes K_D^\vee \otimes \mathcal{O}_D(C - p - (2i + 1)q^{(g-2-j)})) \neq 0,$$

or, equivalently,

$$H^0(D, \wedge^{n-i} M_{K_D} \otimes \mathcal{O}_D(C - J_1 - (2i + 1)J_0)) \neq 0.$$

By Lemma 1.8(ii),  $H^0(D, \mathcal{O}_D(C - J_1 - (2i + 1)J_0) \otimes K_D^\vee) = 0$ , so this amounts to

$$K_{n-i,0}(D, \mathcal{O}_D(C - J_1 - (2i + 1)J_0), K_D) \neq 0.$$

We now let  $D$  degenerate inside its linear system to the curve  $Y = J_0 + Z$ , where  $Z \in |L_{g-3-j}|$  and  $J_0 \cdot Z = q^{(g-3-j)} =: q$ . This implies, by semicontinuity for Koszul cohomology [6], together with Lemma 1.8, that

$$K_{n-i,0}(Y, \mathcal{O}_Y(C - J_1 - (2i + 1)J_0), \tilde{K}) \neq 0,$$

where  $\tilde{K}$  is the twist of the dualizing sheaf of  $Y$  introduced just before Lemma 1.8. This is the same as saying that the map

$$\wedge^{n-i} H^0(Y, \tilde{K}) \otimes H^0(Y, A_0) \rightarrow \wedge^{n-i-1} H^0(Y, \tilde{K}) \otimes H^0(Y, A_1)$$

is not injective, where  $A_d := \mathcal{O}_Y(C - J_1 - (2i + 1)J_0) \otimes \tilde{K}^{\otimes d}$  are the line bundles we defined earlier. As seen in the proof of Lemma 1.8(i), restriction induces an isomorphism

$$H^0(Y, \tilde{K}) \cong H^0(Z, K_Z(2q)).$$

Using the identification between  $J_0$  and  $E$ , we have seen in the proof of Lemma 1.8(ii) that  $\mathcal{O}_{J_0}(A_d)(-q) \cong \eta^{\otimes(j-2i+2)}(r-a)$  is a nontrivial line bundle of degree 0 on  $E$ , therefore  $H^i(J_0, \mathcal{O}_{J_0}A_d(-q)) = 0$  for  $i = 0, 1$ . Thus, restriction to  $Z$  induces an isomorphism

$$H^0(Y, A_d) \cong H^0(Z, \mathcal{O}_Z(A_d)).$$

This also gives that the map

$$\begin{aligned} \wedge^{n-i} H^0(Z, K_Z(2q)) \otimes H^0(Z, \mathcal{O}_Z(A_0)) \\ \rightarrow \wedge^{n-i-1} H^0(Z, K_Z(2q)) \otimes H^0(Z, \mathcal{O}_Z(A_1)), \end{aligned}$$

fails to be injective. As one has

$$H^0(Z, \mathcal{O}_Z(A_1)) \subseteq H^0(Z, \mathcal{O}_Z(A_1 + 2q)) \quad \text{and} \quad H^0(Z, \mathcal{O}_Z(A_{-1} - 2q)) = 0,$$

we obtain  $K_{n-i,0}(Z, \mathcal{O}_Z(A_0), K_Z(2q)) \neq 0$ , which can be rewritten as

$$(15) \quad H^0(Z, \wedge^{n-i} M_{K_Z(2q)} \otimes \mathcal{O}_Z(A_0)) \neq 0.$$



We compute that the slope  $\mu$  satisfies  $\mu(\wedge^{n-i} M_{K_Z(2q)} \otimes \mathcal{O}_Z(A_0)) = g(Z) - 1$ , where  $\mu(M_{K_Z(2q)}) = -2$ . By Serre duality, condition (15) can then be rewritten as

$$H^0(Z, \wedge^{n-i} M_{K_Z(2q)}^\vee \otimes K_Z \otimes \mathcal{O}_Z(-A_0)) \neq 0.$$

We now use that Beauville [4, Proposition 2] has described the theta divisors of vector bundles of the form  $\wedge^{n-i} M_{K_Z(2q)}$  as above. Using [4], from (15) it follows that either

$$\mathcal{O}_Z(A_0) - K_Z \in Z_{n-i} - Z_{n-3+i-j},$$

or else

$$\mathcal{O}_Z(A_0) - K_Z - 2q \in Z_{n-i-1} - Z_{n-2+i-j}.$$

Taking into account that  $\mathcal{O}_Z(A_0) = \mathcal{O}_Z(C - p - (2i + 1)q)$ , the desired conclusion now follows. As a final remark, we note that, whilst in [4] it is assumed that  $Z$  is nonhyperelliptic (which, using the Brill–Noether genericity of  $Z$ , happens whenever  $g - 3 - j \geq 3$ ), the above statement is a triviality for  $g - 3 - j = 1$ , whereas in the remaining case  $g - 3 - j = 2$  it follows directly from the argument in [4, Proposition 2]. Indeed, in this case we have a short exact sequence

$$0 \rightarrow \wedge^{n-i-1} M_{K_Z(2q)}^\vee \rightarrow \wedge^{n-i} M_{K_Z(2q)}^\vee \rightarrow \wedge^{n-i} M_{K_Z}^\vee \rightarrow 0.$$

The claim now follows immediately from [12, Section 3]. This completes the proof.  $\square$

By Proposition 1.9 and induction, we now reduce the proof of (8) to a single statement on elliptic curves on the ruled surface  $X$ :

**Theorem 1.10** *Set  $g = 2n + 1$  and  $\ell \geq 2$ . Then for a general element  $[C, \tau] \in \mathcal{R}_{g,\ell}$ , one has  $K_{n-1,1}(C, K_C \otimes \tau) = 0$ .*

**Proof** We apply Lemma 1.7 and the sufficient condition (11). By the inductive step described above, reasoning by contradiction, it suffices to show that if  $D \in |L_1|$  is general, then

$$C_D - K_D - p - (2i + 1)q^{(1)} \notin D_{n-i} - D_{n-i}$$

for each  $1 \leq i \leq g - 3$ . Assume this is not the case. The bounds (13) force  $n = i$  and the difference variety on the right consists of  $\{\mathcal{O}_D\}$ . One needs to prove that  $\mathcal{O}_D(C - p - (2n + 1)q^{(1)}) \cong \mathcal{O}_D(f_r - J_1)$  is not effective. As  $H^0(X, f_r - J_1) = 0$  and  $D \in |J_0 + f_a|$ , it suffices to prove

$$H^1(X, f_r - J_1 - J_0 - f_a) = H^1(X, f_r - f_a + K_X) = 0,$$

or equivalently by Serre duality, that  $H^1(X, f_a - f_r) = 0$ . This follows immediately from the Leray spectral sequence.  $\square$

## 2 The Green–Lazarsfeld secant conjecture for paracanonical curves

We recall the statement of the Green–Lazarsfeld secant conjecture [16]. Let  $p$  be a positive integer,  $C$  a smooth curve of genus  $g$  and  $L$  a nonspecial line bundle of degree

$$(16) \quad d \geq 2g + p + 1 - \text{Cliff}(C).$$

The secant conjecture predicts that if  $L$  is  $(p+1)$ -very ample then  $K_{p,2}(C, L) = 0$  (the converse implication is easy, so one has an equivalence). The secant conjecture has been proved in many cases in [9], in particular for a general curve  $C$  and a general line bundle  $L$ . In the extremal case  $d = 2g + p + 1 - \text{Cliff}(C)$ , Theorem 1.7 in [9] says that whenever

$$L - K_C + C_{d-g-2p-3} \not\subseteq C_{d-g-p-1} - C_{2g-d+p}$$

(the left-hand side being a divisorial difference variety),  $K_{p,2}(C, L) = 0$ . Theorem 1.7 in [9] requires  $C$  to be Brill–Noether–Petri general, but the proof given there shows that for curves of odd genus the only requirement is that  $C$  have maximum gonality  $(g + 3)/2$ .

In the case at hand, we choose a general curve on the decomposable elliptic surface  $X$

$$C \in |gJ_0 + f_r|$$

of genus  $g = 2n + 1$  and Clifford index  $\text{Cliff}(C) = n$ . We apply the above result to  $L_C = K_C \otimes \tau$ , with  $\tau = \mathcal{O}_C(\zeta)$ . In order to conclude that  $K_{n-3,2}(C, K_C \otimes \tau) = 0$ , it suffices to show

$$\tau + C_2 \not\subseteq C_{n+1} - C_{n-1}.$$

We have two natural points on  $C$ , namely those cut out by intersection with  $J_0$  and  $J_1$ , and for those points it suffices to show

$$(17) \quad \phi^* \zeta \otimes \mathcal{O}_C(J_0 + J_1) \notin C_{n+1} - C_{n-1}.$$

We first establish a technical result similar to Lemma 1.8 and because of this analogy we use similar notations. For  $0 \leq i \leq g - 1$ , let  $Y \in |(g - i)J_0 + f_r|$  be the union of  $J_0$  and a general curve  $Z \in |(g - i - 1)J_0 + f_r|$ . We set  $x_0 := Z \cdot J_0 = f_{t^{g-i-1}} \cdot J_0$ , where  $t^{(g-i-1)} \in E$  satisfies  $\mathcal{O}_E(t^{(g-i-1)} - r) = \eta^{\otimes(g-i-1)}$ . We denote by  $\tilde{K} \in \text{Pic}(Y)$

the twist at the node of the dualizing sheaf of  $Y$  such that  $\mathcal{O}_{J_0}(\tilde{K}) \cong \mathcal{O}_{J_0}$  and  $\mathcal{O}_Z(\tilde{K}) \cong K_Z(2x_0)$ . We recall that  $Z$  has a second distinguished point, namely  $x_1 = Z \cdot J_1$ .

**Lemma 2.1** *Let  $Y \in |(g-i)J_0 + f_r|$  for  $0 \leq i \leq g-1$  be as above and assume  $j$  is an integer satisfying  $0 \leq j \leq n-1$  and  $0 \leq i-j \leq n+1$ . For a general  $D \in |(g-i)J_0 + f_r|$ , we have:*

- (i)  $h^0(Y, \tilde{K}) = h^0(D, K_D)$ .
- (ii)  $H^0(Y, \mathcal{O}_Y(\phi^*\zeta^\vee - (2j + 1 - i)J_0 - J_1)) = 0$ .
- (iii)  $h^0(Y, \tilde{K}^{\otimes m} \otimes \phi^*\zeta^\vee(- (2j + 1 - i)J_0 - J_1)) = h^0(D, K_D^{\otimes m} \otimes \phi^*\zeta^\vee(- (2j + 1 - i)J_0 - J_1))$  for  $m \in \{1, 2\}$ .

**Proof** (i) This is similar to Lemma 1.8(i) and we skip the details.

(ii) Set  $k = -(2j + 1 - i)$ . If  $k \leq 0$ , the statement is clear for degree reasons, so assume  $k \geq 1$ . Then

$$H^0(X, \phi^*\zeta^\vee \otimes \mathcal{O}_X(kJ_0 - J_1)) \cong H^0(E, (\eta \otimes \zeta^\vee) \otimes \text{Sym}^{k-1}(\mathcal{O}_E \oplus \eta)) = 0,$$

so it suffices to show that  $H^1(X, \phi^*\zeta^\vee \otimes \mathcal{O}_X(kJ_0 - J_1 - Y)) = 0$ . By Riemann–Roch, this is equivalent to

$$H^1(X, \phi^*\zeta \otimes \mathcal{O}_X((g - k - i - 1)J_0 + f_r)) = 0.$$

Using the given bounds, we have that  $g - k - i - 1 \geq -1$ . It suffices to show  $H^1(X, \phi^*\zeta \otimes \mathcal{O}_X(mJ_0 + f_r)) = 0$ , for  $m \geq -1$ . This follows along the lines of the proof of Lemma 1.8.

(iii) By Riemann–Roch and semicontinuity, it suffices to show that for  $m = 1, 2$ , one has

$$H^1(Y, \tilde{K}^{\otimes m} \otimes \mathcal{O}_Y(\phi^*\zeta^\vee - (2j + 1 - i)J_0 - J_1)) = 0.$$

Consider the exact sequence

$$(18) \quad 0 \rightarrow \mathcal{O}_Z(-x_0) \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_{J_0} \rightarrow 0.$$

Then  $\mathcal{O}_{J_0}(\tilde{K}^{\otimes m} \otimes \phi^*\zeta^\vee(- (2j + 1 - i)J_0 - J_1)) \cong \zeta^\vee \otimes \eta^{\otimes(i-2j-1)} \neq 0 \in \text{Pic}^0(E)$ . So it suffices to show

$$\begin{aligned} H^1(Z, K_Z \otimes \phi^*\zeta^\vee(- (2j - i)x_0 - x_1)) &= 0, \\ H^1(Z, K_Z^{\otimes 2} \otimes \phi^*\zeta^\vee(- (2j - i - 2)x_0 - x_1)) &= 0. \end{aligned}$$

The second vanishing is automatic for degree reasons (using the bounds on  $i$  and  $j$ ), so we just need to establish the first one. By Serre duality, this is equivalent to

$$H^0(Z, \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_Z((2j - i + 1)x_0)) = 0.$$

This is obvious if  $2j - i + 1 < 0$ , so assume  $2j - i + 1 \geq 0$ . Using once more the Leray spectral sequence, it follows  $H^0(X, \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X((2j - i + 1)J_0)) = 0$ , so it suffices to prove

$$H^1(X, \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X((2j - i + 1)J_0 - Z)) = 0.$$

Using Serre duality and the bound  $g - 2j - 4 \geq -1$ , this goes through as in the proof of Lemma 2.1(i). □

The following proposition provides the induction step, to be proved in order to establish the second half of the Prym–Green conjecture:

**Proposition 2.2** *Let  $0 \leq i \leq g - 2$ . Suppose for a general curve  $D \in |(g - i)J_0 + f_r|$  there exists an integer  $0 \leq j \leq i$  such that*

$$\mathcal{O}_D(\phi^*\zeta + (2j - i + 1)J_0 + J_1) \in D_{n+1-i+j} - D_{n-1-j}.$$

*Then for a general curve  $Z \in |(g - i - 1)J_0 + f_r|$ , there exists  $0 \leq j' \leq i + 1$  such that*

$$\mathcal{O}_Z(\phi^*\zeta + (2j' - i)J_0 + J_1) \in Z_{n-i+j'} - Z_{n-1-j'}.$$

**Proof** By assumption,  $j \leq n - 1$  and  $i - j \leq n + 1$ . Applying again the determinantal description of divisorial difference varieties from [12, Section 3] and with Serre duality, the hypothesis turns into

$$H^0(D, \wedge^{n-1-j} M_{K_D} \otimes \mathcal{O}_D(\phi^*\zeta^\vee - (2j - i + 1)J_0 - J_1)) \neq 0.$$

By Lemma 2.1 and semicontinuity,  $H^0(D, \mathcal{O}_D(\phi^*\zeta^\vee - (2j + 1 - i)J_0 - J_1)) = 0$ , so the above is equivalent to

$$K_{n-1-j,1}(D, \mathcal{O}_D(\phi^*\zeta^\vee - (2j - i + 1)J_0 - J_1), K_D) \neq 0.$$

By Lemma 2.1 and semicontinuity for Koszul cohomology, we then also have

$$K_{n-1-j,1}(Y, \mathcal{O}_Y(\phi^*\zeta^\vee - (2j - i + 1)J_0 - J_1), \tilde{K}) \neq 0,$$

where recall that  $Y = Z \cup J_0$ , with  $Z \cdot J_0 = x_0$ . Consider again the exact sequence (18),

and since  $H^0(J_0, \mathcal{O}_{J_0}(mJ_0 + \phi^*\zeta^\vee)) = 0$  for any  $m$ , the inclusion map yields isomorphisms

$$H^0(Z, K_Z^{\otimes m} \otimes \mathcal{O}_Y(\phi^*\zeta^\vee + (2m - 2j + i - 2)x_0 - x_1)) \cong H^0(Y, \tilde{K}^{\otimes m} \otimes \mathcal{O}_Y(\phi^*\zeta^\vee - (2j - i + 1)J_0 - J_1))$$

that are valid for all positive integers  $m$ . Recall the isomorphism  $H^0(Y, \tilde{K}) \cong H^0(Z, K_Z(2x_0))$  given by restriction. We define the graded  $\text{Sym } H^0(Z, K_Z(2x_0))$ -module

$$A := \bigoplus_{q \in \mathbb{Z}} H^0(Z, \mathcal{O}_Z(\tilde{K}^{\otimes q} + \phi^*\zeta^\vee - (2j - i + 1)x_0 - x_1)),$$

as well as the graded  $\text{Sym } H^0(Y, \tilde{K})$ -module

$$B := \bigoplus_{q \in \mathbb{Z}} H^0(Y, \tilde{K}^{\otimes q} \otimes \mathcal{O}_Y(\phi^*\zeta^\vee - (2j - i + 1)J_0 - J_1)).$$

We then have the following commutative diagram, where the vertical arrows are isomorphisms induced by tensoring the exact sequence (18):

$$\begin{array}{ccc} \wedge^{n-1-j} H^0(Z, K_Z(2x_0)) \otimes A_1 & \longrightarrow & \wedge^{n-2-j} H^0(Z, K_Z(2x_0)) \otimes A_2 \\ \downarrow & & \downarrow \\ \wedge^{n-1-j} H^0(Y, \tilde{K}) \otimes B_1 & \longrightarrow & \wedge^{n-2-j} H^0(Y, \tilde{K}) \otimes B_2 \end{array}$$

Thus it follows  $K_{n-1-j,1}(Z, \mathcal{O}_Z(\phi^*\zeta^\vee + (i - 2j - 2)x_0 - x_1), K_Z(2x_0)) \neq 0$ , or, equivalently,

$$H^0(Z, \wedge^{n-1-j} M_{K_Z(2x_0)} \otimes K_Z \otimes \mathcal{O}_Z(\phi^*\zeta^\vee + (i - 2j - 2)x_0 - x_1)) \neq 0.$$

We now compute the slope  $\chi$ :

$$\chi(\wedge^{n-1-j} M_{K_Z(2x_0)} \otimes K_Z \otimes \mathcal{O}_Z(\phi^*\zeta^\vee + (i - 2j - 2)x_0 - x_1)) = 0.$$

Applying once more the description given in [4, Proposition 2] for the theta divisors of the exterior powers of the vector bundle  $M_{K_Z(2x_0)}$ , we obtain that either

$$\mathcal{O}_Z(\phi^*\zeta + (2j - i)x_0 + x_1) \in Z_{n-i+j} - Z_{n-1-j},$$

or

$$\mathcal{O}_Z(\phi^*\zeta + (2j + 2 - i)x_0 + x_1) \in Z_{n+1-i+j} - Z_{n-2-j},$$

which establishes the claim. □

We now complete the proof of the Prym–Green conjecture for odd genus.

**Theorem 2.3** *Set  $g = 2n + 1$  and  $\ell \geq 2$ . Then for a general element  $[C, \tau] \in \mathcal{R}_{g,\ell}$  one has  $K_{n-3,2}(C, K_C \otimes \tau) = 0$ .*

**Proof** Using the inductive argument from [Lemma 2.1](#), it suffices to prove the base case of the induction, that is, show that if  $D \in |J_0 + f_r|$  is general and  $0 \leq j \leq g - 1$ , then

$$\mathcal{O}_D(\phi^*\zeta + (2j - g + 2)J_0 + J_1) \notin D_{n-1-j} - D_{n-1-j},$$

for any  $0 \leq j \leq g - 1$ . Suppose this is not the case, which forces  $j = n - 1$  and then

$$\mathcal{O}_D(\phi^*(\zeta \otimes \eta^\vee)) \cong \mathcal{O}_D(\phi^*\zeta - J_0 + J_1) \cong \mathcal{O}_D.$$

Since  $H^0(X, \phi^*(\zeta \otimes \eta^\vee)) = 0$ , this implies  $H^1(X, \phi^*(\zeta \otimes \eta^\vee \otimes \mathcal{O}_X(-J_0 - f_r))) \neq 0$ . Observe  $H^2(X, \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X(-J_0 - f_r)) = 0$  by Serre duality. Taking the cohomology exact sequence associated to

$$0 \rightarrow \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X(-J_0 - f_r) \rightarrow \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X(-f_r) \rightarrow \mathcal{O}_{J_0}(\phi^*(\zeta \otimes \eta^\vee) - f_r) \rightarrow 0$$

and using the Leray spectral sequence, we immediately get

$$H^1(X, \phi^*(\zeta \otimes \eta^\vee) \otimes \mathcal{O}_X(-J_0 - f_r)) = 0,$$

which is a contradiction. □

## References

- [1] **M Aprodu, J Nagel**, *Koszul cohomology and algebraic geometry*, University Lecture Series 52, Amer. Math. Soc., Providence, RI (2010) [MR](#)
- [2] **E Arbarello, A Bruno, E Sernesi**, *On hyperplane sections of K3 surfaces*, *Algebr. Geom.* 4 (2017) 562–596 [MR](#)
- [3] **A Beauville**, *Prym varieties and the Schottky problem*, *Invent. Math.* 41 (1977) 149–196 [MR](#)
- [4] **A Beauville**, *Some stable vector bundles with reducible theta divisor*, *Manuscripta Math.* 110 (2003) 343–349 [MR](#)
- [5] **C Birkenhake, H Lange**, *Complex abelian varieties*, 2nd edition, *Grundle. Math. Wissen.* 302, Springer (2004) [MR](#)
- [6] **M Boratyński, S Greco**, *Hilbert functions and Betti numbers in a flat family*, *Ann. Mat. Pura Appl.* 142 (1985) 277–292 [MR](#)

- [7] **A Chiodo, D Eisenbud, G Farkas, F-O Schreyer**, *Syzygies of torsion bundles and the geometry of the level  $\ell$  modular variety over  $\overline{\mathcal{M}}_g$* , *Invent. Math.* 194 (2013) 73–118 [MR](#)
- [8] **E Colombo, G Farkas, A Verra, C Voisin**, *Syzygies of Prym and paracanonical curves of genus 8*, *Épjournal Geom. Algébrique* 1 (2017) art. id. 7 [MR](#)
- [9] **G Farkas, M Kemeny**, *The generic Green–Lazarsfeld secant conjecture*, *Invent. Math.* 203 (2016) 265–301 [MR](#)
- [10] **G Farkas, M Kemeny**, *The Prym–Green conjecture for torsion line bundles of high order*, *Duke Math. J.* 166 (2017) 1103–1124 [MR](#)
- [11] **G Farkas, K Ludwig**, *The Kodaira dimension of the moduli space of Prym varieties*, *J. Eur. Math. Soc.* 12 (2010) 755–795 [MR](#)
- [12] **G Farkas, M Mustață, M Popa**, *Divisors on  $\mathcal{M}_{g,g+1}$  and the minimal resolution conjecture for points on canonical curves*, *Ann. Sci. École Norm. Sup.* 36 (2003) 553–581 [MR](#)
- [13] **G Farkas, N Tarasca**, *Du Val curves and the pointed Brill–Noether theorem*, *Selecta Math.* 23 (2017) 2243–2259 [MR](#)
- [14] **W Fulton, R Lazarsfeld**, *On the connectedness of degeneracy loci and special divisors*, *Acta Math.* 146 (1981) 271–283 [MR](#)
- [15] **ML Green**, *Koszul cohomology and the geometry of projective varieties*, *J. Differential Geom.* 19 (1984) 125–171 [MR](#)
- [16] **M Green, R Lazarsfeld**, *On the projective normality of complete linear series on an algebraic curve*, *Invent. Math.* 83 (1986) 73–90 [MR](#)
- [17] **A Hirschowitz, S Ramanan**, *New evidence for Green’s conjecture on syzygies of canonical curves*, *Ann. Sci. École Norm. Sup.* 31 (1998) 145–152 [MR](#)
- [18] **A Treibich**, *Revêtements tangentiels et condition de Brill–Noether*, *C. R. Acad. Sci. Paris Sér. I Math.* 316 (1993) 815–817 [MR](#)
- [19] **C Voisin**, *Green’s canonical syzygy conjecture for generic curves of odd genus*, *Compos. Math.* 141 (2005) 1163–1190 [MR](#)

*Institut für Mathematik, Humboldt Universität zu Berlin  
Berlin, Germany*

*Department of Mathematics, Stanford University  
Palo Alto, CA, United States*

[farkas@math.hu-berlin.de](mailto:farkas@math.hu-berlin.de), [michael.kemeny@gmail.com](mailto:michael.kemeny@gmail.com)

Proposed: Dan Abramovich  
Seconded: Jim Bryan, Richard Thomas

Received: 7 September 2017  
Revised: 23 February 2018

