

The simplicial EHP sequence in \mathbb{A}^1 -algebraic topology

KIRSTEN WICKELGREN

BEN WILLIAMS

We give a tool for understanding simplicial desuspension in \mathbb{A}^1 -algebraic topology: we show that $X \rightarrow \Omega(S^1 \wedge X) \rightarrow \Omega(S^1 \wedge X \wedge X)$ is a fiber sequence up to homotopy in 2-localized \mathbb{A}^1 algebraic topology for $X = (S^1)^m \wedge \mathbb{G}_m^{\wedge q}$ with $m > 1$. It follows that there is an EHP spectral sequence

$$\mathbb{Z}_{(2)} \otimes \pi_{n+1+i}^{\mathbb{A}^1}(S^{2n+2m+1} \wedge (\mathbb{G}_m)^{\wedge 2q}) \Rightarrow \mathbb{Z}_{(2)} \otimes \pi_i^{\mathbb{A}^1, s}(S^m \wedge (\mathbb{G}_m)^{\wedge q}).$$

14F42; 19G12, 55Q40

1. Introduction	1691
2. Overview of \mathbb{A}^1 homotopy theory	1700
3. Localization	1715
4. The Grothendieck–Witt group	1727
5. The Hilton–Milnor splitting	1731
6. The stable isomorphism	1737
7. Fiber of the James–Hopf map	1751
8. \mathbb{A}^1 simplicial EHP fiber sequence	1770
References	1775

1 Introduction

Let Σ denote the suspension functor from pointed simplicial sets (or topological spaces) to itself, defined as $\Sigma X := S^1 \wedge X$. For some maps $f: \Sigma Y \rightarrow \Sigma X$, there is a $g: Y \rightarrow X$ such that $f = \Sigma g$. In this case, f is said to desuspend and g is called a desuspension of f . Under certain conditions, the obstruction to desuspending f is a generalized Hopf invariant, as is proven by the existence of the EHP sequence

$$(1) \quad X \rightarrow \Omega \Sigma X \rightarrow \Omega \Sigma X^{\wedge 2}$$

of I James [19; 20; 21; 22] and Toda [37; 38], which induces a long exact sequence in homotopy groups in a range; see for example Toda [36] or Whitehead [39, Chapter XII, Theorem 2.2]. Namely, f desuspends if and only if the generalized Hopf invariant

$$H(f): Y \rightarrow \Omega \Sigma Y \xrightarrow{\Omega f} \Omega \Sigma X \rightarrow \Omega \Sigma X^{\wedge 2}$$

is null. Because calculations can become easier after applying suspension, it is useful to have such a systematic tool for studying desuspension.

By work of James [20; 21], it is known that when X is an odd-dimensional sphere, (1) is a fiber sequence, and when X is an even-dimensional sphere, (1) is a fiber sequence after localizing at 2. In particular, for any sphere, (1) is a 2-local fiber sequence. Since the suspension of a sphere is again a sphere, the corresponding fiber sequences for all spheres form an exact couple, thereby defining the EHP spectral sequence; see Mahowald [30]. The EHP spectral sequence is a tool for calculating unstable homotopy groups of spheres. See, for example, the extensive calculations of Toda in [38].

We provide the analogous tools for $\Sigma X = S^1 \wedge X$ in \mathbb{A}^1 -algebraic topology, identifying the obstruction to S^1 -desuspension of a map whose codomain is any sphere with a generalized Hopf invariant, and relating S^1 -stable homotopy groups of spheres to unstable homotopy groups, after 2-localization, by the corresponding EHP spectral sequence. We leave the p -localized sequence for future work.

Place ourselves in the setting of \mathbb{A}^1 -algebraic topology over a field; see Morel and Voevodsky [34; 33]. Let \mathbf{Sm}_k denote the category of smooth schemes over a perfect field k , and consider the simplicial model category $\mathbf{sPre}(\mathbf{Sm}_k)$ of simplicial presheaves on \mathbf{Sm}_k with the \mathbb{A}^1 injective local model structure, which will be recalled in Section 2. This model structure can be localized at a set of primes P (see Hornbostel [14] and Section 3) giving rise to the notation of a P -local fiber sequence up to homotopy. See Definition 8.1. Define the notation

$$S^{n+q\alpha} = (S^1)^{\wedge n} \wedge (\mathbb{G}_m)^{\wedge q}.$$

Let $\Omega(-)$ denote the pointed \mathbb{A}^1 -mapping space $\mathrm{Map}_{\mathbf{sPre}(\mathbf{Sm}_k)_*}(S^1, L_{\mathbb{A}^1}-)$, where $L_{\mathbb{A}^1}$ denotes \mathbb{A}^1 -fibrant replacement.

Theorem 1.1 *Let $X = S^{n+q\alpha}$ with $n > 1$. There is a 2-local \mathbb{A}^1 -fiber sequence up to homotopy,*

$$X \rightarrow \Omega \Sigma X \rightarrow \Omega \Sigma X^{\wedge 2}.$$

Let $\pi_i^{\mathbb{A}^1}$ denote the i^{th} \mathbb{A}^1 homotopy sheaf, and more generally define $\pi_{i+v\alpha}^{\mathbb{A}^1}(X)$ to be the sheaf associated to the presheaf taking a smooth k -scheme U to the \mathbb{A}^1 -homotopy classes of maps from $S^{i+j\alpha} \wedge U_+$ to X . The stable \mathbb{A}^1 homotopy groups are defined as the colimit $\pi_{i+v\alpha}^{s,\mathbb{A}^1}(X) = \text{colim}_{r \rightarrow \infty} \pi_{i+r+v\alpha}^{\mathbb{A}^1}(\Sigma^r X)$.

Theorem 1.2 (simplicial EHP sequence) *Choose n, q and v in $\mathbb{Z}_{\geq 0}$ with $n \geq 2$.*

- *There is a spectral sequence*

$$(E_{i,j}^r, d_r: E_{i,j}^r \rightarrow E_{i-1,j-r}^r) \Rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i-n+(v-q)\alpha}^{s,\mathbb{A}^1} = \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1,s} S^{n+q\alpha}$$

with $E_{i,j}^1 = \mathbb{Z}_{(2)} \otimes \pi_{j+1+i+v\alpha}^{\mathbb{A}^1}(S^{2j+2n+1+2q\alpha})$ if $i \geq 2n-1+j$ and otherwise $E_{i,j}^1 = 0$.

- *Choose $n' > n$. There is a spectral sequence*

$$(E_{i,j}^r, d_r: E_{i,j}^r \rightarrow E_{i-1,j-r}^r) \Rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1} S^{n'+q\alpha}$$

with $E_{i,j}^1 = \mathbb{Z}_{(2)} \otimes \pi_{j+1+i+v\alpha}^{\mathbb{A}^1}(S^{2j+2n+1+2q\alpha})$ if $i \geq 2n-1+j$ and $j < n'-n$, and $E_{i,j}^1 = 0$ otherwise.

Theorem 1.2 follows directly from Theorem 1.1. Theorem 1.1 is a summary of a more refined theorem, giving conditions under which (1) is a fiber sequence without 2-localization. To state this theorem, let $\text{GW}(k)$ denote the Grothendieck–Witt group of k , and consider the element of $\text{GW}(k)$ given by $-\langle -1 \rangle = -(1 + \rho\eta)$, where η is the motivic Hopf map and $\rho = [-1]$ in the notation of [33, Definition 3.1]. Let K^{MW} denote Milnor–Witt K -theory as defined in [33, Definition 3.1]. For a set of primes P , write \mathbb{Z}_P for the ring \mathbb{Z} with formal multiplicative inverses adjoined for all primes not in P .

Theorem 1.3 *Let $X = S^{n+q\alpha}$ with $n > 1$, and let $e = (-1)^{n+q} \langle -1 \rangle^q$. Let P be a set of primes. The sequence*

$$X \rightarrow \Omega \Sigma X \rightarrow \Omega \Sigma X^{\wedge 2}$$

is a P -local \mathbb{A}^1 -fiber sequence up to homotopy if $1+m(1+e)$ are units in $\text{GW}(k) \otimes \mathbb{Z}_P$ for all positive integers m .

Corollary 1.4 *In the setting of Theorem 1.3, the sequence*

- *is always a 2-local \mathbb{A}^1 -fiber sequence up to homotopy;*
- *is an \mathbb{A}^1 -fiber sequence up to homotopy when $e = -1$ or when $n + q$ is odd and the field k is not formally real.*

In particular, the sequence is an \mathbb{A}^1 -fiber sequence up to homotopy

- when n is odd and q is even;
- when $n + q$ is odd and $k = \mathbb{C}$, or more generally, when $n + q$ is odd and k is any field such that $2\eta = 0$ in K_*^{MW} .

Although the statement of Theorem 1.3 is a direct analogue of the corresponding theorem in algebraic topology, the proof given here is not a straightforward generalization of a proof in algebraic topology. The difficulty is that \mathbb{A}^1 -fiber sequences are problematic and \mathbb{A}^1 -homotopy groups are not necessarily finitely generated. Standard tools like the Serre spectral sequence are not currently available.

If a theorem holds for every simplicial set in a functorial manner, it may globalize in the following sense. First, one may be able to obtain in **sPre** a naïve analogue by starting with simplicial presheaves instead of simplicial sets, performing corresponding operations, producing corresponding maps in **sPre**. If the theorem in algebraic topology says that some map is always a weak equivalence (resp. weak equivalence through a range), it may be immediate that the corresponding map is a global weak equivalences (resp. global weak equivalence through a range). If the \mathbb{A}^1 -invariant analogues of the operations considered in the theorem are obtained by applying $L_{\mathbb{A}^1}$ to the naïve analogue (defined by applying the operation in simplicial set to the sections over each $U \in \mathbf{Sm}_k$), then the theorem holds in \mathbb{A}^1 -algebraic topology.

This is the case of the Hilton–Milnor splitting shown below:

Theorem 1.5 *There is a natural isomorphism*

$$\Sigma \Omega \Sigma X \rightarrow \Sigma \vee \bigvee_{n=1}^{\infty} X^{\wedge n}$$

in the \mathbb{A}^1 -homotopy category.

This is also the case for the statement that for any simplicial presheaf X , the sequence (1) is a fiber sequence up to homotopy in the range $i \leq 3n - 2$, meaning

$$\begin{aligned} \pi_{3n-2}^{\mathbb{A}^1} X \rightarrow \pi_{3n-1}^{\mathbb{A}^1} \Sigma X \rightarrow \pi_{3n-1}^{\mathbb{A}^1} \Sigma(X^{\wedge 2}) \rightarrow \\ \dots \rightarrow \pi_i^{\mathbb{A}^1} X \rightarrow \pi_{i+1}^{\mathbb{A}^1} \Sigma X \rightarrow \pi_{i+1}^{\mathbb{A}^1} \Sigma(X^{\wedge 2}) \rightarrow \pi_{i-1}^{\mathbb{A}^1} X \rightarrow \dots \end{aligned}$$

is exact. This fact is shown in joint work with A Asok and J Fasel [4].

This is not the case for Theorems 1.1 and 1.3, ie these theorems are not proven by globalizing a corresponding result in algebraic topology, where the sequence (1) fails to be exact for $X = S^n \vee S^n$. See Example 6.20.

Here is a sketch of the proof of Theorem 1.1; its purpose is to help the reader understand the proof given in this paper, and also to explain the similarities with, and differences from, the situation in classical algebraic topology. Let $J(X)$ denote the free monoid on a pointed object X in simplicial presheaves on \mathbf{Sm}_k , where \mathbf{Sm}_k denotes smooth schemes over k .

In algebraic topology, the free monoid on a pointed object is canonically homotopy equivalent to the loops of the suspension. It was understood by Fabien Morel that the same result holds in \mathbb{A}^1 -algebraic topology. Indeed, a result of Morel implies that $L_{\mathbb{A}^1} J(X)$ is simplicially equivalent to $\Omega L_{\mathbb{A}^1} \Sigma X$, for X pointed, fibrant and connected. (The phrase “simplicially equivalent” means weakly equivalent in the injective Nisnevich local model structure. Here, “fibrant” means with respect to this model structure as well.) We show the versions of this result that we need in Section 5. By globalizing a construction from algebraic topology [39, Chapter VII, Section 2], there is a sequence

$$X \rightarrow J(X) \rightarrow J(X^{\wedge 2}),$$

where $X \rightarrow J(X)$ is the canonical map induced from the adjunction between Σ and Ω , and $J(X) \rightarrow J(X^{\wedge 2})$ is the James–Hopf map, ie the above maps exist in \mathbb{A}^1 -algebraic topology and the composite map $X \rightarrow J(X^{\wedge 2})$ is nullhomotopic (simplicially). Thus, there is an induced map in the homotopy category from X to the P -localized \mathbb{A}^1 -homotopy fiber of $J(X) \rightarrow J(X^{\wedge 2})$, where P is a set of primes. Use the notation $h: X \rightarrow F$ for this map. Theorems 1.1 and 1.3 say that for X a sphere, h is a P -localized \mathbb{A}^1 -homotopy equivalence for appropriate P , and it is proved as follows.

By Theorem 1.5, there is a map of S^1 -spectra $b: \Sigma^\infty J(X) \rightarrow \Sigma^\infty X$. Using the tensor structure of spectra over spaces, it follows that there is a map of S^1 -spectra

$$c: \Sigma^\infty J(X) \rightarrow \Sigma^\infty (X \times J(X^{\wedge 2}))$$

which fits into the commutative diagram

$$\begin{array}{ccc} \Sigma^\infty J(X) & \xrightarrow{\quad} & \Sigma^\infty X \times J(X^{\wedge 2}) \\ & \searrow & \swarrow \\ & \Sigma^\infty J(X^{\wedge 2}) & \end{array}$$

(See Section 6.4, and, for general X , see Section 7.2, in particular the discussion following Construction 7.10.)

The two spaces $J(X)$ and $X \times J(X^{\wedge 2})$ are the same size in the sense that stably they are both weakly equivalent to $\Sigma^\infty \vee \bigvee_{n=1}^\infty X^{\wedge n}$. To see this, note that $\Sigma^\infty J(X) \cong \Sigma^\infty \vee \bigvee_{n=1}^\infty X^{\wedge n}$ by Theorem 1.5;

$$\Sigma^\infty(X \times J(X^{\wedge 2})) \cong \Sigma^\infty X \vee \Sigma^\infty J(X^{\wedge 2}) \vee \Sigma^\infty(X \wedge J(X^{\wedge 2}))$$

because the product of two spaces is stably equivalent to the wedge of their smash with their wedge, ie $\Sigma^\infty(X \times Y) \cong \Sigma^\infty(X \vee Y \vee X \wedge Y)$. By Theorem 1.5, we have stable weak equivalences $J(X^{\wedge 2}) \cong \bigvee_{n=1}^\infty X^{\wedge 2n}$ and $X \wedge J(X^{\wedge 2}) \cong \bigvee_{n=1}^\infty X^{\wedge 2n+1}$. These equivalences, when combined with the previous, show that stably $X \times J(X^{\wedge 2}) \cong \bigvee_{n=1}^\infty X^{\wedge n}$. It is not always the case, however, that the stable map $c: \Sigma^\infty J(X) \rightarrow \Sigma^\infty(X \times J(X^{\wedge 2}))$ constructed above is a weak equivalence; see Example 6.20. In algebraic topology, this map is a weak equivalence for X an odd sphere, and an equivalence after inverting 2 for X an even sphere. We show an analogous fact in \mathbb{A}^1 -algebraic topology, in the following way. By the Hilton–Milnor theorem, the map c can be viewed as a “matrix”, which itself is the product of matrices corresponding to the diagonal of $J(X)$ and a combination of b with the James–Hopf map $J(X) \rightarrow J(X^{\wedge 2})$. Nick Kuhn’s calculations of the stable decomposition of the diagonal of $J(X)$ (see [26]) and the stable decomposition of the James–Hopf map (see [25, Section 6]) in algebraic topology globalize to give the matrix entries of c in terms of sums of permutations of smash powers of X . Morel computes that the swap map $X \wedge X \rightarrow X \wedge X$ is e , and more importantly, any permutation σ on $X^{\wedge m}$ is equivalent to $e^{\text{sign}(\sigma)}$ in the homotopy category (see [33, Lemma 3.43]). Since X is a co- H -space, Kuhn’s results imply that the matrix entries of c are diagonal, and when combined with Morel’s result, we calculate the n^{th} such entry to be

- $1 + \frac{1}{2}((2n)!/(2^n n!) - 1)(e + 1)$ for n even;
- $(1 + \frac{1}{2}((2(n - 1))!/(2^{n-1}(n - 1)!) - 1)(e + 1))(\frac{1}{2}(n + 1) + \frac{1}{2}(n - 1)e)$ for n odd.

Note that $(2n)!/(2^n n!) = 1(3)(5)\cdots(2n - 1)$ is an odd integer, so the n^{th} diagonal term of this matrix is of the form $1 + m(e + 1)$ with m an integer for n even, and a product of two such terms for n odd. Note that $e^2 = 1$ in the homotopy category, because e is the class of the swap. It follows that the product of two terms of the form $1 + m(e + 1)$ is also of this form because $(e + 1)^2 = 2(1 + e)$. Also note that for any positive integer m , we have that

$$((m + 1) + me)((m + 1) - me) = 2m + 1,$$

whence $(m + 1) + me$ is a unit after localizing at 2. It follows that c is a weak equivalence after 2-localization. More generally, c is a weak equivalence after P -localization whenever all the terms $(m + 1) + me$ are units in $\text{GW}(k) \otimes \mathbb{Z}_P$. See Proposition 6.17. This produces the corresponding hypothesis in Theorem 1.3. We can furthermore characterize exactly when $(m + 1) + me$ is a unit in $\text{GW}(k)$ for all m : either $e = -1$ or the field is not formally real and $e = -\langle -1 \rangle$. See Corollary 4.8.

We are then in the situation where we have two P -localized \mathbb{A}^1 -fiber sequences

$$(2) \quad F \rightarrow J(X) \rightarrow J(X^{\wedge 2}),$$

$$(3) \quad X \rightarrow X \times J(X^{\wedge 2}) \rightarrow J(X^{\wedge 2}),$$

and a stable equivalence between the total spaces which respects the maps to the base. We would like to “cancel off” the base $J(X^{\wedge 2})$ to conclude that there is an equivalence between the fibers. This is indeed what we do, however, there are two major obstacles to overcome with this approach.

The first is that the standard tool to measure the size of a fiber of a fibration in terms of the base and total space is the Serre spectral sequence, and at present there is no Serre spectral sequence for \mathbb{A}^1 -fiber sequences. The desired such sequence would use a homology theory like $H_*^{\mathbb{A}^1}$ (see [33, Definition 6.29]) because of the need for analogues of the Hurewicz theorem as in [33, Chapter 6.3] to conclude a weak equivalence between the fibers. We use S^1 -stable \mathbb{A}^1 homotopy groups on the obvious analogue of the Serre spectral sequence defined by lifting the skeletal filtration on the base to express the total space as a filtered limit of cofibrations, and then making an exact couple by applying π_i^{s,P,\mathbb{A}^1} . This gives a spectral sequence even for a global fibration, but it is not clear that it can be controlled. We provide some of the desired control in Section 7.2. Assume for simplicity that the base is reduced in the sense that its 0-skeleton is a single point, as is the case for $J(X^{\wedge 2})$. The E^1 -page can be then identified with π_i^{s,P,\mathbb{A}^1} applied to a wedge indexed by the nondegenerate simplices of the base of the fibration. This wedge construction takes P -local \mathbb{A}^1 weak equivalences of the fiber (resp. P -local \mathbb{A}^1 weak equivalences in a range) to P - \mathbb{A}^1 weak equivalences (resp. in a range). See Lemmas 7.12 and 7.15. We then show that this identification of the E^1 -page is natural with respect to maps, and even natural with respect to the stable map c discussed above. See Lemma 7.13. This identification of the E^1 -page does not behave well with respect to weak equivalences of the base, as it involves the specific simplices of the base. It is sufficient here because the map on the base is the identity. We do not understand the E^2 -page.

We then have a map of spectral sequences from the spectral sequence associated to (2) to the spectral sequence associated to (3). We wish to use this map of spectral sequences to show that the stable weak equivalences of the base and total space imply a stable \mathbb{A}^1 equivalence of the fibers, after appropriately localizing.

Then comes the second difficulty. There are infinitely many nonvanishing stable homotopy groups of the fibers in question, and these groups themselves are not necessarily finitely generated abelian groups. We need to show that there is an isomorphism of these E^1 -pages, but to do this, we need to allow for the possibility that all terms of both spectral sequences are nonzero, nonfinitely generated groups. We give an inductive argument to do this in Proposition 7.18, and immediately following the proposition there is a verbal description of what happened.

The strategy of this proof of the motivic EHP sequence is modeled on the proof of the EHP sequence given in Michael Hopkins's stable homotopy course at Harvard University in the fall of 2012. Hopkins credits this proof to James [19; 21] together with some ideas of Ganea [10]. In this argument, the original Serre spectral sequence is used; there is no need to work in spectra, as calculations in (co)homology suffice. Since the (co)homology of spheres in algebraic topology is concentrated in two degrees, there is no analogue of Proposition 7.18.

It is also possible to compute the first differential in the EHP sequence of Theorem 1.2, and this computation will be made available in a joint paper by Asok and the present authors [4].

Computations of unstable motivic homotopy groups of spheres can be applied to classical problems in the theory of projective modules, for example to the problem of determining when algebraic vector bundles decompose as a direct sum of algebraic vector bundles of smaller rank. See [33, Chapter 8] and Asok and Fasel [1; 2].

In a different direction, it can be shown that there is an \mathbb{A}^1 weak equivalence

$$\Sigma(\mathbb{P}^1 - \{0, 1, \infty\}) \cong \Sigma(\mathbb{G}_m \vee \mathbb{G}_m)$$

between the S^1 suspensions of $\mathbb{P}^1 - \{0, 1, \infty\}$ and $\mathbb{G}_m \vee \mathbb{G}_m$. By comparing the actions of the absolute Galois group on geometric étale fundamental groups, it can be shown that this weak equivalence does not desuspend; see Wickelgren [40]. Because the action of the absolute Galois group on $\pi_1^{\text{ét}}(\mathbb{P}_{\mathbb{Q}}^1 - \{0, 1, \infty\})$ is both tied to interesting mathematics — see Ihara [17] — and obstructs desuspension, it is potentially also of

interest to have systematic tools like those provided by the EHP sequence to study the obstructions to desuspension.

1.1 Remark on the field

Throughout this paper, we will use k for a field such that the unstable connectivity results of Morel apply in the \mathbb{A}^1 homotopy theory over k . Specifically, we rely on [33, Theorems 5.46 and 6.1]. At present, these results require k to be perfect, but this may well be unnecessary. The requirement that k should be infinite that arises in the use of [33, Lemma 1.15] can be circumvented by use of Hogadi and G Kulkarni [13].

Organization

The organization of this paper is as follows: Theorem 1.2 is proven in Section 8 as Theorems 8.5 and 8.6. Theorem 1.1 is proven in Section 8 as Theorem 8.3. The core of these arguments is the cancellation property of Section 7.3. The substitute for the Serre spectral sequence is developed in Section 7. In Section 6, the motivic James–Hopf map and the diagonal of the James construction are computed stably as matrices with entries in $\mathrm{GW}(k)$. Section 5 proves the Hilton–Milnor splitting. Section 4 gives results on the Grothendieck–Witt group that are needed to understand when the matrices computed in Section 6 are invertible. Section 3 provides needed results on localizations of $\mathbf{sPre}(\mathbf{Sm}_k)$ and $\mathbf{Spt}(\mathbf{Sm}_k)$, and Section 2 introduces the needed notation and background on \mathbb{A}^1 -homotopy theory.

Acknowledgements

We wish to strongly thank Aravind Asok and Jean Fasel for their generosity in sharing with us unpublished notes about the James construction in \mathbb{A}^1 algebraic topology. Wickelgren wishes to thank Michael Hopkins for his stable homotopy course in the Fall of 2012, and the entire homotopy theory community in Cambridge, Massachusetts, for their energy, enthusiasm and perspicacity. We are also pleased to thank Emily Riehl for help using simplicial model categories, and Aravind Asok, Jean Fasel, David Gepner, Daniel Isaksen and Kyle Ormsby for useful discussions. We also thank an anonymous referee for useful comments, in particular for pointing out a gap in a previous version of the proof of Proposition 7.18.

Wickelgren was supported by an American Institute of Mathematics five-year fellowship and NSF grants DMS-1406380 and DMS-1552730. Some of this work was done while

Wickelgren was in residence at MSRI during the Spring 2014 algebraic topology semester, supported by NSF grant 0932078 000. Further work was done while both authors were in residence at the Institut Mittag-Leffler during the special program *Algebrogeometric and homotopical methods*. We thank MSRI and Institut Mittag-Leffler for the pleasant and productive visits we enjoyed with them.

2 Overview of \mathbb{A}^1 homotopy theory

In the sequel, we will have to draw on many results regarding \mathbb{A}^1 homotopy. We collect those results in this section for ease of reference. We make no claim that any of these results are original.

As stated in the introduction, we assume k is a field so that the unstable connectivity theorem of Morel applies over k .

Let \mathbf{Sm}_k denote a small category equivalent to the category of smooth, finite-type k -schemes. The category $\mathbf{sPre}(\mathbf{Sm}_k)$ is the category of simplicial presheaves on \mathbf{Sm}_k , and $\mathbf{sPre}(\mathbf{Sm}_k)_*$ the category of pointed simplicial presheaves. The category \mathbf{Sm}_k is considered embedded in $\mathbf{sPre}(\mathbf{Sm}_k)$ via the Yoneda embedding. The terminal object of \mathbf{Sm}_k and $\mathbf{sPre}(\mathbf{Sm}_k)$ is therefore $\mathrm{Spec} k$, which is also denoted by k and $*$ depending on the context.

The notation $\mathrm{Map}(X, Y)$ denotes the internal mapping object where it appears, generally in $\mathbf{sPre}(\mathbf{Sm}_k)$. Many categories appearing in the sequel are simplicially enriched, and in them $\mathrm{SMap}(X, Y)$ will denote a simplicial mapping object. Where there is a model structure \mathfrak{a} present, we will use the notation $[X, Y]_{\mathfrak{a}}$ to denote the set of maps in the homotopy category from X to Y . The notation $[X, Y]$ will be used when \mathfrak{a} is clear from the context.

If K is a simplicial set, then we write K_i for the set of i -simplices in K .

2.1 Model structures

This paper makes use of two families of model structure on the category $\mathbf{sPre}(\mathbf{Sm}_k)$ and its descendants: the local injective model structure of [23] — introduced there as the “global” model structure — and the local flasque model structure of [18]. Our use of these terms follows [18]. These model structures are Quillen equivalent. Each gives rise to descendent model structures by \mathbb{A}^1 - or P -localization or by stabilization. The flasque

model structures are employed only to prove technical results regarding spectra; when “flasque” is not specified, it is to be understood that the injective structures are meant. The weak equivalences in the injective local and the flasque local model structures are the local weak equivalences — those maps that induce isomorphisms on homotopy sheaves, properly defined [23]. In the seminal work [34], these maps are called “simplicial weak equivalences” in order to emphasize their nonalgebraic character.

Both the injective and the flasque local model structures are left Bousfield localizations of global model structures on $\mathbf{sPre}(\mathbf{Sm}_k)$, a global model structure being one where the weak equivalences are those maps $\phi: X \rightarrow Y$ that induce weak equivalences $\phi(U): X(U) \rightarrow Y(U)$ for all objects U of \mathbf{Sm}_k . Both the global injective and the global flasque model structures are left proper, simplicial, cellular — see [14] — and combinatorial, so left Bousfield localizations of either at any set of morphisms exist and are again left proper, simplicial and cellular. In the injective model structures all objects are cofibrant, and therefore these model structures are *tractable* in the sense of [5].

We shall need a cartesian model category structure on $\mathbf{sPre}(\mathbf{Sm}_k)$ from time to time. The category $\mathbf{sPre}(\mathbf{Sm}_k)$ is cartesian closed as a category in its own right, and it is well known — and proved in [5, Application IV] — that the injective local model structure is a symmetric monoidal model category in the sense of [15, Chapter 4] with the cartesian product providing the monoidal operation. One then may wish to establish that some model structure \mathfrak{a} obtained as a left Bousfield localization of this structure inherits the structure of a monoidal model category.

Proposition 2.1 *Suppose \mathfrak{a} is a localization of the injective model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$ such that \mathfrak{a} is a simplicial monoidal model category with respect to the cartesian product. Let A denote a set of morphisms in $\mathbf{sPre}(\mathbf{Sm}_k)$ such that for all objects U of \mathbf{Sm}_k , if $f: X \rightarrow Y$ is in A , there is a morphism in A isomorphic to $f \times \text{id}_U: X \times U \rightarrow Y \times U$. Then the localization of \mathfrak{a} at A inherits the monoidal model category structure of \mathfrak{a} .*


Proof This is an application of [5, Proposition 4.47]. Here we assume that we are working within some universe \mathbf{X} . The role of \mathbf{V} is played by \mathbf{sSet} . The model category \mathfrak{a} is left proper because all objects are cofibrant. Then the hypotheses of [5, Proposition 4.47] are that \mathfrak{a} is symmetric monoidal, and that there exists a set of homotopy generators of \mathfrak{a} , here taken to be the representable objects U , such that if Z is A -local and U is representable, the internal mapping object $\text{Map}(U, Z)$ is again A -local. By adjunction, this follows from our hypotheses. \square

Corollary 2.2 *Let \mathfrak{a} be a symmetric monoidal model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$, where the monoidal operation is given by the cartesian product. Let \mathfrak{a}_* denote the pointed analogue. If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$, then the functor $X \times \cdot$ preserves \mathfrak{a} weak equivalences. If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, then the functor $X \wedge \cdot$ preserves \mathfrak{a}_* weak equivalences.*

Proof Let $f: Z \rightarrow Y$ be an \mathfrak{a} weak equivalence. Because \mathfrak{a}_* is a simplicial model category in which all objects are cofibrant and monomorphisms are cofibrations, it follows from [35, Corollary 14.3.2] that $\text{id}_X \vee f: X \vee Z \rightarrow X \vee Y$ is an \mathfrak{a} weak equivalence.

By Proposition 2.1, for any object X , the functor $X \times \cdot$ preserves trivial cofibrations; by Ken Brown’s lemma, it therefore preserves weak equivalences between cofibrant objects, but all objects are cofibrant.

Note that $\text{id}_X \vee f$, id_* , $X \times f$ and $X \wedge f$ determine a map of pushout squares as in the commutative diagram



Furthermore, $X \vee Z \rightarrow X \times Z$ and $X \vee Y \rightarrow X \times Y$ are cofibrations because they are monomorphisms. It now follows from [35, Corollary 14.3.2] that $X \wedge f: X \wedge Z \rightarrow X \wedge Y$ is an \mathfrak{a} weak equivalence, as claimed. □

2.2 Homotopy sheaves

If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$ or $\mathbf{sPre}(\mathbf{Sm}_k)_*$, we write $L_{\text{Nis}}X$ for a functorial fibrant replacement in the local injective model structure; this also serves as a fibrant replacement in the flasque model structure. We write $L_{\mathbb{A}^1}$ for a functorial fibrant replacement in the \mathbb{A}^1 model structures.

Since the purpose of this paper is to establish some identities regarding \mathbb{A}^1 homotopy sheaves, it behoves us to define what a homotopy sheaf means in the sequel.

The following definitions date at least to [23].

Definition 2.3 If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$, then we define $\pi_0^{\text{pre}}(X)$ as the presheaf

$$U \mapsto \pi_0(|X(U)|),$$

where U is an object of \mathbf{Sm}_k and $|X(U)|$ indicates a geometric realization of $X(U)$. We define $\pi_0(X)$ as the associated Nisnevich sheaf to $\pi_0^{\text{pre}}(X)$.

Proposition 2.4 If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$, then $\pi_0(X)$ is the sheaf associated to the presheaf coequalizer:

$$U \mapsto \text{coeq}(X(U)_1 \begin{array}{c} \xrightarrow{d_1} \\ \xrightarrow{d_0} \end{array} X(U)_0).$$

Definition 2.5 If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ with basepoint $x_0 \rightarrow X$, then we define $\pi_i^{\text{pre}}(X, x_0)$ for $i \geq 1$ as the presheaf

$$U \mapsto \pi_i(|X(U)|, x_0),$$

where U is an object of \mathbf{Sm}_k and x_0 , in an abuse of notation, indicates the basepoint of $|X(U)|$ induced by $x_0 \rightarrow X$. We define $\pi_i(X, x_0)$ as the associated Nisnevich sheaf. The basepoint x_0 will generally be understood and omitted.

The reader is reminded that $X(U)$ may have connected components that do not appear in the global sections, $X(*)$. In this case, the sheaves of groups $\pi_i(X, x_0)$ as defined above are insufficient to describe the homotopy theory of X .

It is the case that the functor $\pi_i(\cdot)$ takes simplicial weak equivalences to isomorphisms, and $\pi_0^{\mathbb{A}^1}$ takes \mathbb{A}^1 weak equivalences to isomorphisms.

If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, we reserve the notation $\Omega^i X$ for the derived loop space $\text{Map}_*(S^n, L_{\text{Nis}}X)$. In particular, $\Omega^0 X \cong L_{\text{Nis}}X$.

We rely on the following result throughout:

Proposition 2.6 Equip $\mathbf{sPre}(\mathbf{Sm}_k)_*$ with the Nisnevich local model structure. If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and if $i \geq 0$, then $\pi_i(X)$ is the sheaf associated to the presheaf

$$U \mapsto [\Sigma^i(U_+), X]_*.$$

Proof By applying a functorial fibrant replacement functor if necessary, we may assume that $X(U)$ is a fibrant simplicial set for all objects U of \mathbf{Sm}_k . In the following

sequence of isomorphisms, all homotopy groups considered are simplicial homotopy groups of fibrant simplicial sets:

$$\begin{aligned}
 [S^i \wedge U_+, X]_* &\cong \pi_0(\mathbf{SMap}_*(S^i \wedge U_+, X)) \\
 &\cong \pi_0(\mathbf{SMap}_*(S^i, \mathbf{Map}_*(U_+, X))) \\
 &\cong \pi_0(\mathbf{SMap}_*(S^i, \mathbf{Map}(U, X))) \\
 &\cong \pi_0(\mathbf{sSet}_*(S^i, \mathbf{Map}(U, X)(*))) \\
 &\cong \pi_0(\mathbf{sSet}_*(S^i, X(U))) \\
 &\cong \pi_i(X(U)),
 \end{aligned}$$

as required. □

Corollary 2.7 *If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and $i \geq 0$, then $\pi_i(X) \cong \pi_0(\Omega^i X)$.*

Proof The result follows from the proposition and the adjunction

$$[\Sigma^i(U_+), X]_* \cong [U_+, \Omega^i X]_* \quad \square$$

Since taking global sections, $X \mapsto X(*)$, is taking a stalk, we also have the following corollary:

Corollary 2.8 *If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and if $i \geq 0$, then*

$$\pi_i(X)(*) \cong [S^i, X].$$

If $i, j \geq 0$, we define

$$S^{i+j\alpha} = S^i \wedge \mathbb{G}_m^{\wedge j},$$

where \mathbb{G}_m is pointed at the rational point 1. If $j \geq 0$ and X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_+$, we define $\pi_{i+j\alpha}^{\mathbb{A}^1}(X)$ as $\pi_i(\mathbf{Map}_*(\mathbb{G}_m^{\wedge j}, L_{\mathbb{A}^1} X))$; it is isomorphic to the sheaf associated to the presheaf $U \mapsto [S^{i+j\alpha} \wedge U_+, X]_{\mathbb{A}^1}$. Taking global sections, we have

$$\pi_{i+j\alpha}^{\mathbb{A}^1}(X)(k) \cong [S^{i+j\alpha}, X]_{\mathbb{A}^1}.$$

2.3 Compact objects and flasque model structures

We say that an object X of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is *compact* if

$$\operatorname{colim}_i \mathbf{Map}_*(X, F_i) \cong \mathbf{Map}_*(X, \operatorname{colim}_i F_i)$$

whenever F_i is a filtered system in $\mathbf{sPre}(\mathbf{Sm}_k)$, and similarly for $\mathbf{sPre}(\mathbf{Sm}_k)_*$. An argument similar to that of [9, Lemma 9.13] shows that pointed smooth schemes are compact, and it is easy to see that finite constant simplicial presheaves are compact. If A and B are pointed compact objects, then $A \wedge B$ is compact, and all finite colimits of compact objects are again compact.

We shall frequently make use of the following:

Proposition 2.9 *Let I be a filtered small category and X an I -shaped diagram in $\mathbf{sPre}(\mathbf{Sm}_k)$ (resp. $\mathbf{sPre}(\mathbf{Sm}_k)_*$). Then the natural map $\mathrm{hocolim} X \rightarrow \mathrm{colim} X$ is a global weak equivalence.*

Proof We describe the case of $\mathbf{sPre}(\mathbf{Sm}_k)$; that of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is similar.

By construction — see [12, Chapter 18] — $(\mathrm{hocolim} X)(U) \cong \mathrm{hocolim}(X(U))$ for all $U \in \mathbf{Sm}_k$, and similarly for colim . The result then follows from the classical fact that the natural map $\mathrm{hocolim} X(U) \rightarrow \mathrm{colim} X(U)$ is a weak equivalence [6, XII.3.5]. \square

Proposition 2.10 *If $X_0 \rightarrow X_1 \rightarrow \dots$ is a sequential diagram in $\mathbf{sPre}(\mathbf{Sm}_k)$, then the natural map of sheaves*

$$\mathrm{colim}_i \pi_0(X_i) \rightarrow \pi_0(\mathrm{colim}_i X_i)$$

is an isomorphism.

Proof By Proposition 2.9, we may replace $\{X_i\}$ by a naturally weakly equivalent diagram without changing the homotopy type of $\mathrm{colim}_i X_i$. The group $\mathrm{colim}_i \pi_0(X_i)$ is also unchanged by such a procedure. We can therefore assume that $X_i(U)$ is a fibrant simplicial set for all objects U of \mathbf{Sm}_k .

Since, according to Proposition 2.4, $\pi_0(Y)$ is the sheaf associated to a coequalizer of presheaves provided Y takes values in fibrant simplicial sets, the result follows by commuting colimits. \square

The injective local model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$ suffers from a technical drawback when one wishes to calculate with filtered colimits, which is that filtered colimits of fibrant objects are not necessarily fibrant themselves. This is the problem that motivates the construction of the flasque model structures of [18], and one can see the presence of flasque or flasque-like conditions appearing often throughout the literature when calculations with filtered colimits are being carried out; see [24; 9; 32].

We therefore consider two flasque model structures on $\mathbf{sPre}(\mathbf{Sm}_k)$: the local flasque structure in which the weak equivalences are the simplicial weak equivalences, and the \mathbb{A}^1 flasque structure in which the weak equivalences are the \mathbb{A}^1 weak equivalences. These model structures apply also to $\mathbf{sPre}(\mathbf{Sm}_k)_*$. These model structures are simplicial, proper and cellular, and the \mathbb{A}^1 structures are left Bousfield localizations of the local model structure. There is a square of Quillen adjunctions

$$(4) \quad \begin{array}{ccc} \text{injective local} & \xrightarrow{\simeq} & \text{flasque local} \\ \downarrow & & \downarrow \\ \text{injective } \mathbb{A}^1 & \xrightarrow{\simeq} & \text{flasque } \mathbb{A}^1 \end{array}$$

where the arrows indicate the left adjoints, and each arrow is the identity functor on $\mathbf{sPre}(\mathbf{Sm}_k)$. The horizontal arrows represent Quillen equivalences. A similar diagram obtains in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

Not all objects are cofibrant in $\mathbf{sPre}(\mathbf{Sm}_k)$ or $\mathbf{sPre}(\mathbf{Sm}_k)_*$ in the flasque model structures, in contrast to the case of the injective structures. Since the \mathbb{A}^1 flasque structures are left Bousfield localizations of the local flasque structures, the cofibrant objects in one model structure agree with the cofibrant objects in the other. The results of [18], specifically Lemmas 3.13 and 6.2, show that all pointed simplicial sets and all quotients X/Y of monomorphisms $Y \rightarrow X$ in \mathbf{Sm}_k are flasque cofibrant in $\mathbf{sPre}(\mathbf{Sm}_k)_*$. This includes all smooth schemes pointed at a rational point. Lemma 3.14 of [18] shows that finite smash products of flasque cofibrant objects are again flasque cofibrant in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

Proposition 2.11 *If F_i is a filtered diagram of objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and if X is a compact and flasque cofibrant object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, then there is a zigzag of local (resp. \mathbb{A}^1) weak equivalences*

$$(5) \quad \text{colim}_i \text{Map}_*(X, RF_i) \rightarrow \text{Map}_*(X, R \text{colim}_i RF_i) \leftarrow \text{Map}_*(X, R \text{colim}_i F_i),$$

where R denotes an injective local (resp. injective \mathbb{A}^1) functorial fibrant replacement.

Proof By Proposition 2.9, the local (resp. \mathbb{A}^1) homotopy type of a filtered colimit is invariant under termwise replacement by locally (resp. \mathbb{A}^1) equivalent objects.

Filtered colimits of flasque fibrant objects are again flasque fibrant; see [18].

The objects RF_i are flasque fibrant, so the colimit $\text{colim}_i RF_i$ is flasque fibrant, as is $R \text{colim}_i RF_i$. There is a global weak equivalence $\text{colim}_i RF_i \simeq R \text{colim}_i RF_i$. Since

R preserves weak equivalences, we also have $R \operatorname{colim}_i R F_i \simeq R \operatorname{colim}_i F_i$. Since the object X is flasque cofibrant, the functor $\operatorname{Map}_*(X, \cdot)$ preserves trivial flasque fibrations, and by Ken Brown’s lemma, weak equivalences between flasque fibrant objects. The map $\operatorname{Map}_*(X, \operatorname{colim}_i R F_i) \rightarrow \operatorname{Map}_*(X, R \operatorname{colim}_i F_i)$ is therefore a weak equivalence. The result now follows from the compactness of X . \square

Corollary 2.12 *If F_r is a filtered system of objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and if $i, j \geq 0$ are integers, then there are natural isomorphisms of sheaves*

$$\pi_i(\operatorname{colim}_r F_r) \cong \operatorname{colim}_r \pi_i(F_r)$$

and

$$\pi_{i+j\alpha}^{\mathbb{A}^1}(\operatorname{colim}_r F_r) \cong \operatorname{colim}_r \pi_{i+j\alpha}^{\mathbb{A}^1}(F_r).$$

Proof Combine Corollary 2.7 and Propositions 2.10 and 2.11, noting that the objects $S^{i+j\alpha}$ are compact and flasque cofibrant. \square

We warn the reader that $\pi_{i+j\alpha}^{\mathbb{A}^1}(\Omega^r X)$ differs from $\pi_{i+r+j\alpha}^{\mathbb{A}^1}(X)$ in general; see [33, Theorem 6.46].

2.4 Spectra

We take [16] as our main reference for the theory of spectra in model structures such as those we consider here. We shall require only naïve spectra, rather than symmetric spectra. For us a spectrum, E , shall be an S^1 -spectrum, consisting of a sequence $\{E_i\}_{i=0}^\infty$ of objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, equipped with bonding maps $\sigma: \Sigma E_i \rightarrow E_{i+1}$. The maps of spectra $E \rightarrow E'$ being defined as levelwise maps $E_i \rightarrow E'_i$ which furthermore commute with the bonding maps, we have a category of presheaves of spectra, which we denote by $\mathbf{Spt}(\mathbf{Sm}_k)$.

Just as we have two notions of weak equivalence on $\mathbf{sPre}(\mathbf{Sm}_k)_*$, the local and the \mathbb{A}^1 , we shall have two kinds of weak equivalence between objects of $\mathbf{Spt}(\mathbf{Sm}_k)$, the stable and the \mathbb{A}^1 .

There is a set, I in the notation of [18], of generating cofibrations for which the domains and codomains all possess the property that we call “compact”, which [18] calls “ ω -small” and which is stronger than the property that [16] calls “finitely presented”. Moreover, both model structures are localizations of an objectwise flasque model structure having a set, J in the notation of [18], which again consists of maps having

finitely presented domains and codomains. By the arguments of [16, Section 4], these model structures are *almost finitely generated*.

The theory of [16, Section 3] establishes a stable model structure on $\mathbf{Spt}(\mathbf{Sm}_k)$ based on any cellular, left proper model structure, \mathfrak{a} , on $\mathbf{sPre}(\mathbf{Sm}_k)_*$. In particular, this applies when \mathfrak{a} is a left Bousfield localization of the global injective or global flasque model structure, and therefore when it is one of the four structures of (4). The results of [16, Section 5] ensure that we have Quillen adjunctions and equivalences between these model structures

$$(6) \quad \begin{array}{ccc} \text{stable injective local} & \xrightarrow{\cong} & \text{stable flasque local} \\ \downarrow & & \downarrow \\ \text{stable injective } \mathbb{A}^1 & \xrightarrow{\cong} & \text{stable flasque } \mathbb{A}^1 \end{array}$$

Since the functors of (4) are the identity functors, the same is true of the functors of (6); only the model structure varies.

We write *stable weak equivalence* for the weak equivalences of the stable injective local and stable flasque local model structures, and *stable \mathbb{A}^1 equivalence* for the weak equivalences of the stable injective \mathbb{A}^1 and the stable flasque \mathbb{A}^1 model structures. In keeping with our convention, we write $L_{\mathbb{A}^1} E$ to denote a fibrant replacement of E in the stable \mathbb{A}^1 model structures.

Since the underlying unstable model structures are proper, we may apply fibrant-replacement functors levelwise to objects in $\mathbf{Spt}(\mathbf{Sm}_k)$ to obtain maps of spectra $E \rightarrow RE$ given by $E_i \rightarrow RE_i$, the fibrant replacement in any one of the four unstable model structures under consideration. There is also a spectrum-level infinite loop space functor, Θ^∞ , that takes a spectrum E to the spectrum having i^{th} space

$$(\Theta^\infty E)_i = \operatorname{colim}_{k \rightarrow \infty} \operatorname{Map}_*(S^k, E_{i+k}).$$

Proposition 2.13 *A map $f: E \rightarrow E'$ of $\mathbf{Spt}(\mathbf{Sm}_k)$ is a stable weak equivalence (resp. a stable \mathbb{A}^1 equivalence) if and only if*

$$\Theta^\infty(Rf)_i: (\Theta^\infty RE)_i \rightarrow (\Theta^\infty RE')_i$$

is a weak equivalence for all i , where R represents the flasque local fibrant replacement functor (resp. flasque \mathbb{A}^1 -fibrant replacement functor).

Proof This is a special case of [16, Theorem 4.12]. The ancillary hypotheses given there, that sequential colimits in commute with finite products and that $\text{Map}_*(S^1, \cdot)$ commutes with sequential limits, are satisfied in $\mathbf{sPre}(\mathbf{Sm}_k)_*$. \square

One can verify that a spectrum E is weakly equivalent to the spectrum one obtains from E by replacing each space E_i by the connected component of the basepoint in E_i . We may therefore assume that E_i is connected, meaning we do not have to worry about the problem of components that are not globally defined.

For any integer i , there is an adjunction of categories

$$(7) \quad \Sigma^{\infty-i}: \mathbf{sPre}(\mathbf{Sm}_k)_* \rightleftarrows \mathbf{Spt}(\mathbf{Sm}_k) : \text{Ev}_i,$$

where the spectrum $\Sigma^{\infty-i} X$ is the spectrum the j^{th} space of which is $\Sigma^{j-i} X$ if $j \geq i$, and $*$ otherwise, and where the bonding maps are the evident ones. The right adjoint Ev_i takes E to E_i .

Proposition 2.14 *Suppose α is a left Bousfield localization of either the global injective or the global flasque model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$. Then the adjoint functors of (7) form a Quillen pair between the pointed model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and the stable model structure on $\mathbf{Spt}(\mathbf{Sm}_k)$ induced by α .*

Proof This follows from Definition 1.2, Proposition 1.15 and Definition 3.3 of [16]. \square

The left-derived functor of Ev_0 in the flasque model structures are the functors

$$\Omega^\infty: \{E_n\}_n \mapsto \text{colim}_k \Omega^k R E_{n+k},$$

where R is either $L_{\text{Nis}}^{\text{fl}}$ or $L_{\mathbb{A}^1}^{\text{fl}}$, as appropriate. The smash product on $\mathbf{sPre}(\mathbf{Sm}_k)_*$ extends to an action of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ on $\mathbf{Spt}(\mathbf{Sm}_k)$, and the functors $\Sigma^{\infty-i}$ preserve this structure, and in particular are simplicial functors [16, Section 6].

For an object E of $\mathbf{Spt}(\mathbf{Sm}_k)$, we define the stable homotopy sheaves $\pi_i^s(E)$ as the colimit

$$\pi_i^s(E) = \text{colim}_{r \rightarrow \infty} \pi_{i+r}(E_r).$$

Since $\pi_{i+r}(E_r)$ is the sheaf associated to the presheaf $U \mapsto \pi_{i+r}(E_r(U))$, and sheafification commutes with colimits, it follows that $\pi_i^s(E)$ may also be described as the sheaf associated to the presheaf

$$U \mapsto \pi_i^s(E(U)),$$

where the stable homotopy group is the ordinary stable homotopy group of simplicial spectra. This definition of the sheaf π_i^s is used in [32].

We similarly define the stable \mathbb{A}^1 homotopy sheaves $\pi_{i+j\alpha}^{s,\mathbb{A}^1}(E)$ as the colimit

$$\pi_{i+j\alpha}^{s,\mathbb{A}^1}(E) = \operatorname{colim}_{r \rightarrow \infty} \pi_{i+r+j\alpha}^{\mathbb{A}^1}(E_r).$$

When X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, will use the notation $\pi_*^s(X)$ and $\pi_*^{s,\mathbb{A}^1}(X)$ for $\pi_*^s(\Sigma^\infty X)$ and $\pi_*^{s,\mathbb{A}^1}(\Sigma^\infty X)$, respectively.

Proposition 2.15 *Let $f: E \rightarrow E'$ be a map in $\mathbf{Spt}(\mathbf{Sm}_k)$. Then*

- (i) *f is a stable weak equivalence if and only if $\pi_i^s(f)$ is an isomorphism for all i ;*
- (ii) *f is an \mathbb{A}^1 -stable weak equivalence if and only if $\pi_i^{s,\mathbb{A}^1}(f)$ is an isomorphism for all i .*

Proof The map $f: E \rightarrow E'$ is a stable weak equivalence if and only if the maps $(\Theta^\infty L_{\mathbf{Nis}}^{\mathbb{f}} E)_i \rightarrow (\Theta^\infty L_{\mathbf{Nis}}^{\mathbb{f}} E')_i$ are simplicial weak equivalences for all i . The space $(\Theta^\infty L_{\mathbf{Nis}}^{\mathbb{f}} E)_j$ is

$$\operatorname{colim}_{r \rightarrow \infty} \Omega^r(L_{\mathbf{Nis}}^{\mathbb{f}} E_{j+r})$$

and its $(i+j)^{\text{th}}$ homotopy sheaf is, by Corollary 2.12,

$$\pi_{i+j}(\operatorname{colim}_{r \rightarrow \infty} \Omega^r(L_{\mathbf{Nis}}^{\mathbb{f}} E_{j+r})) \cong \operatorname{colim}_{j+r \rightarrow \infty} \pi_{i+j+r}(E_{j+r}) \cong \pi_i^s(E).$$

The result for π_i^s follows.

For \mathbb{A}^1 equivalence, the same argument applies mutatis mutandis. Writing $L_{\mathbb{A}^1}^{\mathbb{f}}$ for the flasque \mathbb{A}^1 -fibrant replacement functor, we see that the $(i+j)^{\text{th}}$ homotopy sheaf of the j^{th} level of the \mathbb{A}^1 -stable fibrant replacement $\Theta^\infty(L_{\mathbb{A}^1}^{\mathbb{f}} E)$ is

$$\pi_{i+j}(\operatorname{colim}_{r \rightarrow \infty} \Omega^r(L_{\mathbb{A}^1}^{\mathbb{f}} E_{j+r})),$$

which simplifies to $\pi_i^s(L_{\mathbb{A}^1}^{\mathbb{f}} E) = \pi_i^{s,\mathbb{A}^1}(E)$. □

This proposition says that the definition of stable weak equivalence used in this paper agrees with that of [32].

Proposition 2.16 For any object E of $\mathbf{Spt}(\mathbf{Sm}_k)$ and any nonnegative integers i, i' and j ,

(i) the sheaf associated to the presheaf

$$U \mapsto [\Sigma^{\infty-i'}(S^i \wedge U_+), E]$$

is $\pi_{i-i'}^s(E)$;

(ii) the sheaf associated to the presheaf

$$U \mapsto [\Sigma^{\infty-i'}(S^{i+j\alpha} \wedge U_+), E]_{\mathbb{A}^1}$$

is $\pi_{i-i'+j\alpha}^{s, \mathbb{A}^1}(E)$.

Proof We prove the first statement.

The given presheaf, by adjunction, is

$$U \mapsto [S^i \wedge U_+, \text{Ev}_{i'} E],$$

where the functor $\text{Ev}_{i'}$ is a derived functor in the flasque stable model structure. By reference to [16], we write this presheaf more explicitly as

$$U \mapsto [S^i \wedge U_+, \text{colim}_{r \rightarrow \infty} \Omega^r L_{\text{Nis}}^{\text{fl}} E_{i'+r}],$$

which is associated to the sheaf

$$\pi_i(\text{colim}_{r \rightarrow \infty} \Omega^r L_{\text{Nis}}^{\text{fl}} E_{i'+r}) \cong \text{colim}_{r \rightarrow \infty} \pi_{i-i'+(i'+r)}(E_{i'+r}) \cong \pi_{i-i'}^s(E),$$

as asserted.

The proof of the second statement is similar, with the proviso that $L_{\text{Nis}}^{\text{fl}}$ is replaced by $L_{\mathbb{A}^1}^{\text{fl}}$, and one concludes that the sheaf being represented is

$$\pi_{i-i'}^s(\text{Map}_*(\mathbb{G}_m^{\wedge j}, L_{\mathbb{A}^1}^{\text{fl}} E)) \cong \pi_{i-i'+j\alpha}^{s, \mathbb{A}^1}(E). \quad \square$$

Corollary 2.17 For any $i, j \geq 0$, and any object E of $\mathbf{Spt}(\mathbf{Sm}_k)$, taking global sections gives

$$\pi_i^s(E)(*) = [\Sigma^\infty S^i, E]$$

and

$$\pi_{i+j\alpha}^{s, \mathbb{A}^1}(E)(*) = [\Sigma^\infty S^{i+j\alpha}, E]_{\mathbb{A}^1}.$$

Proposition 2.18 *Suppose $\{E_n\}$ is a filtered system of objects in $\mathbf{Spt}(\mathbf{Sm}_k)$. Then the natural maps*

$$\operatorname{colim}_n \pi_i^s(E_i) \rightarrow \pi_i^s(\operatorname{colim}_n E_i)$$

and

$$\operatorname{colim}_n \pi_{i+j\alpha}^{s, \mathbb{A}^1}(E_i) \rightarrow \pi_{i+j\alpha}^{s, \mathbb{A}^1}(\operatorname{colim}_n E_i)$$

are isomorphisms.

Proof These follow from Corollary 2.12 and the observations that taking colimits commute and that colimits of spectra are calculated termwise. \square

2.5 Long exact sequences of homotopy sheaves

We will use the term *cofiber sequence* only in a limited sense: a cofiber sequence in a pointed model category \mathbf{M} is a sequence of maps $X \rightarrow Y \rightarrow Z$ such that $X \rightarrow Y$ is a cofibration of cofibrant spaces and Z is a categorical pushout of $* \leftarrow X \rightarrow Y$. A *fiber sequence* is dual.

The image of a cofiber sequence in $\operatorname{ho} \mathbf{M}$ may also be called a cofiber sequence, as in [15, Chapter 6]. The notion of *fiber sequence* is dual.

The derived functors of left-Quillen functors preserve cofiber sequences, and dually the derived functors of right-Quillen functors preserve fiber sequences.

Suppose α is a model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$, obtained as a left Bousfield localization of the flasque- or injective-local model structure. Consider $\mathbf{Spt}(\mathbf{Sm}_k)$, endowed with the stable model structure derived from α [16, Section 3]. By [16, Theorem 3.9], the homotopy category $\operatorname{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$ is a triangulated category in the sense of [15, Chapter 7.1]. Write $\pi_{i+j\alpha}^{s, \alpha}(E)$ for the sheaf associated to the presheaf $U \mapsto [S^i \wedge \mathbb{G}_m^j \wedge U_+, E]_{s\alpha}$, where the set of maps is calculated in $\operatorname{ho}_\alpha(\mathbf{Spt}(\mathbf{Sm}_k))$. Thus we have the following result by applying the associated-sheaf functor to Lemma 7.1.10 of [15]:

Proposition 2.19 *If $X \rightarrow Y \rightarrow Z$ is a cofiber sequence in $\operatorname{ho}_\alpha(\mathbf{Spt}(\mathbf{Sm}_k))$, then the induced sequence of homotopy sheaves*

$$\cdots \rightarrow \pi_{i+j\alpha}^{s, \alpha}(X) \rightarrow \pi_{i+j\alpha}^{s, \alpha}(Y) \rightarrow \pi_{i+j\alpha}^{s, \alpha}(Z) \rightarrow \pi_{i-1+j\alpha}^{s, \alpha}(X) \rightarrow \cdots$$

is an exact sequence of sheaves of abelian groups.

Examples include π_i^s and $\pi_{i+j\alpha}^{s, \mathbb{A}^1}$, as well as $\pi_i^{s, P}$ and $\pi_{i+j\alpha}^{s, P, \mathbb{A}^1}$ of Section 3.

2.6 \mathbb{A}^1 -unstable and -stable

We say that a spectrum E is \mathbb{A}^1 - n -connected if $\pi_i^{s, \mathbb{A}^1}(E) = 0$ for all $i \leq n$. From the above definition of π_i^{s, \mathbb{A}^1} , combined with [33, Theorem 6.38] saying that $L_{\mathbb{A}^1}$ does not decrease the connectivity of connected objects, and that $L_{\mathbb{A}^1}$ commutes with Ω for simply connected objects, we deduce the following lemma:

Lemma 2.20 *If X is an \mathbb{A}^1 - n -connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, then $\Sigma^\infty X$ is \mathbb{A}^1 - n -connected.*

Recall that a map $f: X \rightarrow Y$ of connected objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is said to be n -connected if the homotopy fiber is $(n-1)$ -connected, and \mathbb{A}^1 - n -connected if the \mathbb{A}^1 -homotopy fiber is \mathbb{A}^1 - $(n-1)$ -connected.

By use of [33, Theorem 6.53 and Lemma 6.54] and the \mathbb{A}^1 -connectivity theorem, we deduce that if $X \rightarrow Y$ is n -connected with $n \geq 1$ and if moreover $\pi_1(Y)$ is strongly \mathbb{A}^1 -invariant, then $X \rightarrow Y$ is \mathbb{A}^1 - n -connected. These conditions hold when X is simply connected, or when $n \geq 2$ and X is \mathbb{A}^1 -local.

The following result is due to Asok and Fasel [3]:

Proposition 2.21 (Blakers–Massey theorem of Asok and Fasel) *Suppose $f: X \rightarrow Y$ is an \mathbb{A}^1 - n -connected map of connected objects in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and X is \mathbb{A}^1 - m -connected, with $n, m \geq 1$; then the morphism $\mathrm{hofib}_{\mathbb{A}^1} f \rightarrow \Omega L_{\mathbb{A}^1} \mathrm{hocofib} f$ is $(m+n)$ -connected.*

Proof We rely on a homotopy excision result, a consequence of the Blakers–Massey theorem, that says that the result of this proposition holds in the setting of classical topology [39, Chapter VII, Theorem 7.12].

We may replace $f: X \rightarrow Y$ by an equivalent \mathbb{A}^1 -fibration of \mathbb{A}^1 -fibrant objects without changing the \mathbb{A}^1 homotopy type of $\mathrm{hofib}_{\mathbb{A}^1} f$ or of $\mathrm{hocofib} f$.

The \mathbb{A}^1 homotopy fiber of f therefore agrees with the ordinary fiber and therefore also with the simplicial homotopy fiber.

The classical homotopy excision result, applied at points, now says that the map $\mathrm{hofib} f \rightarrow \Omega \mathrm{hocofib} f$ is simplicially $(m+n)$ -connected. Since $m+n \geq 2$, and

$\text{hofib } f = \text{hofib}_{\mathbb{A}^1} f$ is \mathbb{A}^1 -local, it follows that $\pi_1(\Omega \text{ hofib } f)$ is strongly \mathbb{A}^1 -invariant and then by [33, Theorem 6.56] it follows that

$$\text{hofib}_{\mathbb{A}^1} f \simeq L_{\mathbb{A}^1} \text{ hofib}_{\mathbb{A}^1} f \rightarrow L_{\mathbb{A}^1} \Omega \text{ hofib } f$$

is $(m+n)$ -connected.

The connectivity hypotheses imply that $\pi_1(Y) \cong \pi_1^{\mathbb{A}^1}(Y)$ is trivial, and therefore by the van Kampen theorem that $\text{hofib } f$ is simply connected. This implies by [33, Theorem 6.46] that $L_{\mathbb{A}^1} \Omega \text{ hofib } f \simeq \Omega L_{\mathbb{A}^1} \text{ hofib}$. This completes the proof. \square

Corollary 2.22 *Suppose $f: X \rightarrow Y$ is a map of \mathbb{A}^1 simply connected objects in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that the homotopy cofiber $\text{hofib } f$ is \mathbb{A}^1 contractible. Then f is an \mathbb{A}^1 weak equivalence.*

Proof We show by induction that $\text{hofib}_{\mathbb{A}^1} f$ is arbitrarily highly connected. Since X and Y are simply connected, $\text{hofib}_{\mathbb{A}^1} f$ is 0-connected, so f is 1-connected.

Suppose we know that $\text{hofib}_{\mathbb{A}^1} f$ is d -connected; then applying Proposition 2.21 with $n = d + 1$ and $m = 1$, we deduce that $\text{hofib}_{\mathbb{A}^1} f \rightarrow \Omega \text{ hofib } f \simeq *$ is $\mathbb{A}^1-(d+2)$ -connected, so that $\pi_{d+1}(\text{hofib}_{\mathbb{A}^1} f)$ is trivial. \square

Corollary 2.23 *Suppose $f: X \rightarrow Y$ is a map of \mathbb{A}^1 simply connected objects in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $\Sigma^\infty f: \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is an \mathbb{A}^1 weak equivalence; then f is an \mathbb{A}^1 weak equivalence.*

Proof We may replace f by a fibration of \mathbb{A}^1 -fibrant objects.

The map f is necessarily 1-connected, and from the proposition we deduce that $\pi_1(\text{hofib } f) \simeq \pi_0(\text{hofib } f)$, which is trivial. Since Σ^∞ is a left Quillen functor, it preserves cofiber sequences in the derived category, and we deduce that $\Sigma^\infty \text{ hofib } f$ is \mathbb{A}^1 contractible. Since $\text{hofib } f$ is simply connected, the \mathbb{A}^1 Hurewicz theorem implies that $\text{hofib } f$ is \mathbb{A}^1 contractible.

An appeal to Corollary 2.22 now completes the argument. \square

2.7 Points

The site $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$ is well known to have enough points. Let Q be a conservative set of points of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$. For each element $q \in Q$, there is an adjunction of categories

$$q^*: \mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k) \rightleftarrows \mathbf{Set} : q_*$$

where q^* , as well as preserving all colimits, preserves finite limits.

There is a Quillen adjunction

$$q^*: \mathbf{sPre}(\mathbf{Sm}_k) \rightleftarrows \mathbf{sSet} : q_*$$

from the injective local model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$ to the usual model structure on \mathbf{sSet} . This extends in the obvious way to the pointed model categories, and to the categories of spectra

$$q^*: \mathbf{Spt}(\mathbf{Sm}_k) \rightleftarrows \mathbf{Spt} : q_*$$

For an object X of $\mathbf{sPre}(\mathbf{Sm}_k)$, there is, by reference to Proposition 2.4, an isomorphism $q^*\pi_0(X) \cong \pi_0(q^*X)$. It is also the case that $p^*\Omega^i(\mathbf{L}_{\mathbf{Nis}}X) \simeq \Omega^i(\mathbf{Ex}^\infty p^*X)$. This gives us the following proposition:

Proposition 2.24 *If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and i is a positive integer and q a point of $\mathbf{Sh}_{\mathbf{Nis}}(\mathbf{Sm}_k)$, then there is an isomorphism of groups $\pi_i(|q^*X|) \cong q^*\pi_i(X)$.*

Corollary 2.25 *If X is an object of $\mathbf{Spt}(\mathbf{Sm}_k)$, and if i is an integer, then there is an isomorphism of abelian groups $\pi_i^s(q^*X) \cong q^*\pi_i^s(X)$.*

These facts are special cases of results concerning ∞ -topoi [29, 6.5.1.4]. They are well known — see for instance [32, Example 2.2.4] — but seldom stated.

3 Localization

Let P denote a nonempty set of prime ideals of \mathbb{Z} , and $P' = \bigcap_{(p) \in P} (\mathbb{Z} \setminus (p))$ the set of integers not lying in any of these ideals. We write \mathbb{Z}_P for the localization $(P')^{-1}\mathbb{Z}$, and $\mathbb{Z}_{(p)}$ in the case where $P = \{(p)\}$ consists of a single ideal. Following [8], where the following is carried out in the category of CW complexes, we define $S^1_\tau = S^1$, a Kan complex equivalent to $\Delta^1 / \partial\Delta^1$. For any integer n , define $\rho_n: S^1_\tau \rightarrow S^1_\tau$ to be a degree- n self-map of S^1 , and let T_P denote the set of all such ρ_n as n ranges over P' .

The local injective and flasque model structures on $\mathbf{sPre}(\mathbf{Sm}_k)$ are cellular in the sense of Hirschhorn [12]; a proof for the injective case appears in [14, Lemma 1.5] and the flasque case is treated in [18]. We may therefore apply the general machinery of [12] and left Bousfield localize $\mathbf{sPre}(\mathbf{Sm}_k)$ at the set T_P . We call the resulting model structures P -local, and if $P = \{(p)\}$, we call the resulting model structures p -local. Write L_P for the functorial fibrant replacement functor in each model category. In the case where $P = \{(p)\}$, we may write $L_{(p)}$.

The localization of the usual model structure on \mathbf{sSet} with respect to the set T_P of maps is a form of P -local model structure on \mathbf{sSet} , we refer the reader to [8], especially Section 8, for the comparisons between different P -localizations in classical topology and for a discussion of nonnilpotent objects. For nilpotent simplicial sets, the various P -localization functors agree up to weak equivalence; see [8, Proposition 8.1]. In particular, they agree for simply connected spaces, and by extension to simply connected simplicial presheaves. For instance, in the case of simply connected simplicial presheaves, the P -localization defined here agrees with $H_*(\cdot, \mathbb{Z}_P)$ -localization.

Lemma 3.1 *With notation as above, if s is a point of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$, the adjunctions*

$$s^*: \mathbf{sPre}(\mathbf{Sm}_k) \rightleftarrows \mathbf{sSet} : s_*$$

and

$$s^*: \mathbf{sPre}(\mathbf{Sm}_k)_* \rightleftarrows \mathbf{sSet}_* : s_*$$

are monoidal Quillen adjunctions between the P -local model categories, where $\mathbf{sPre}(\mathbf{Sm}_k)$ and $\mathbf{sPre}(\mathbf{Sm}_k)_*$ may be given either the flasque or the injective model structure.

Proof It is sufficient to prove the unpointed cases; the pointed follow immediately. The proofs in the flasque and injective cases are the same.

Following [12, Theorem 3.3.20], the adjoint pair

$$s^*: \mathbf{sPre}(\mathbf{Sm}_k) \rightleftarrows \mathbf{sSet} : s_*$$

is a Quillen adjunction between the P -local model structure on the left and the model structure on \mathbf{sSet} obtained by localization at the set of maps

$$s^*(\rho_n^k \times \text{id}_U): s^*(S_\tau^k \times U) \rightarrow s^*(S_\tau^k \times U),$$

where $\rho_n^k \in T_P$. Denote this set of maps by $s^*T'_P$. It will suffice to show that localization of \mathbf{sSet} at $s^*T'_P$ agrees with localization of \mathbf{sSet} at T_P .

Since evaluation at s^* commutes with fiber products, the maps of $s^*T'_P$ maps are of the form $\rho_n^k \times \text{id}_{s^*U}$, and setting $U = *$, we see that $T_P \subset s^*T'_P$. The maps of $s^*T'_P$ are, moreover, weak equivalences in the localization of \mathbf{sSet} at T_P . It follows that the localization of \mathbf{sSet} at $s^*T'_P$ is simply the ordinary P -localization of \mathbf{sSet} . \square

We note in addition that the model categories appearing above are simplicial model categories, and the adjunctions appearing are adjunctions of simplicial model categories in the sense of [15, Chapter 4.2].

We continue to work principally in the injective local not-localized-at- P model structures, but write $A \simeq_P B$ to indicate that A is weakly equivalent to B in the P -local structure, or equivalently that $L_P A \simeq L_P B$. The notation $A \simeq_{(P)} B$ will be used where appropriate. We will use the flasque model structures only when dealing with spectra.

In this section we will occasionally write groups $\pi_i(X)$ in multiplicative notation even when the groups are abelian. The n^{th} power map of a group G will be the map $x \mapsto x^n$, which is necessarily a homomorphism if G is abelian, and is preserved by group homomorphisms in any case. If P is a set of primes, then a group G is said to be P -local if the n^{th} power map is a bijection on G whenever n is not divisible by any of the primes in P . We will say that a presheaf of groups is P -local if all groups of sections are P -local, and a sheaf of groups is P -local if the appropriate n^{th} power maps are isomorphisms of sheaves of sets.

Proposition 3.2 *If X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and P is a set of primes, then the sheaves $\pi_i(L_P X)$ are P -local sheaves of groups.*

Proof It suffices to show that the presheaves

$$U \mapsto \pi_i(L_P X(U))$$

are P -local; the result for the associated sheaves is then an exercise in sheafification.

Let n be an integer not divisible by any of the primes of P , let $i \geq 1$ and let U be an object of \mathbf{Sm}_k . We wish to show that the n^{th} power map on $\pi_i(L_P X(U))$ is a bijection, but this is the map induced by $\rho_n^i \times \text{id}_U$ on $\pi_0(\text{SMap}_*(S_\tau^i \vee U_+, L_P X))$. Since $L_P X$ is P -local and $\rho_n^i \times \text{id}_U$ is in T'_P , this map is a bijection. \square

Lemma 3.3 *Let X be an object of $\mathbf{sPre}(\mathbf{Sm}_k)$, let s be a point of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$ and let P be a set of primes. Then $s^*L_P X \simeq L_P s^* X$.*

Proof We first claim that $s^*L_P X$ is P -local. Since it is fibrant, it suffices to show that if ρ_n^k is an element of T_P , then the induced map

$$(\rho_n^k)_*: \text{SMap}(S_\tau^k, s^*L_P X) \rightarrow \text{SMap}(S_\tau^k, s^*L_P X)$$

is a weak equivalence. If $\{U_i\}$ is a system of neighborhoods for s^* then there is a succession of natural isomorphisms

$$\begin{aligned} \text{SMap}(S_\tau^k, s^*L_P X) &\cong \text{SMap}(S_\tau^k, \text{colim}_U(L_P X)(U)) \\ &\cong \text{colim}_U \text{SMap}(S_\tau^k, (L_P X)(U)) \quad (\text{since } S_\tau^k \text{ is compact}) \\ &\cong \text{colim}_U \text{SMap}(S_\tau^k \times U, L_P X) \end{aligned}$$

and ρ_n^k induces a weak equivalence on the spaces $\text{SMap}(S_\tau^k \times U, L_P X)$ since $L_P X$ is P -local.

The functor s^* preserves trivial cofibrations, and therefore the map $s^*X \rightarrow s^*L_P X$ is a trivial cofibration, the target of which is fibrant in the P -local model structure on \mathbf{sSet} . Therefore $s^*L_P X$ is weakly equivalent in the ordinary model structure on \mathbf{sSet} to any other P -fibrant-replacement for s^*X , notably to $L_P s^*X$, which is what was claimed. □

Proposition 3.4 *Let X be a fibrant object of $\mathbf{sPre}(\mathbf{Sm}_k)$, let S be a conservative set of points of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$ and let P be a set of primes. Then X is P -local if and only if s^*X is P -local for all $s^* \in S$.*

Proof The space X is P -local if and only if X is fibrant and $X \rightarrow L_P X$ is a local weak equivalence. This is the case if and only if $s^*X \rightarrow s^*L_P X$ is a weak equivalence for all $s^* \in S$, which, by Lemma 3.3, is the case if and only if $s^*X \rightarrow L_P s^*X$ is a weak equivalence for all $s^* \in S$, and since s^*X is fibrant, this is the same as saying that s^*X is P -local in \mathbf{sSet} . □

Definition 3.5 An object X of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is said to be *simple* if the action of $\pi_1(X)$ on $\pi_i(X)$ is trivial for all $i \geq 1$.

In particular, if X is simple then the sheaf $\pi_1(X)$ is a sheaf of abelian groups which acts trivially on $\pi_i(X)$ for all $i \geq 2$. A simply connected object is simple, as is an object with an H -space structure.

Proposition 3.6 *Let X be a connected, simple object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$; then the natural maps $\mathbb{Z}_P \otimes_{\mathbb{Z}} \pi_i(X) \rightarrow \pi_i(L_P X)$ are isomorphisms.*

Proof Fix a point s of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$. By Lemma 3.3, there are isomorphisms

$$s^* \pi_i(L_P X) \cong \pi_i(s^*L_P X) \cong \pi_i(L_P s^*X).$$

As remarked above for the case of simply connected spaces, Proposition 8.1 of [8] implies that the P -localization of simplicial sets considered here agrees with the P -localization of [6, Section V] in the case of simple simplicial sets. By reference to [6, V.4.1–V.4.2], the group $\pi_i(L_P s^* X)$ is isomorphic to

$$\pi_i(s^* X) \otimes_{\mathbb{Z}} \mathbb{Z}_P \cong s^*(\pi_i(X) \otimes_{\mathbb{Z}} \mathbb{Z}_P),$$

which proves the proposition. □

Lemma 3.7 *Suppose X is a simply connected object of $\mathbf{sPre}(\mathbf{Sm}_k)$ and P a set of prime numbers; then $L_P(S^1 \wedge X) \simeq S^1 \wedge L_P X$ and $\Omega L_P X \simeq L_P \Omega L_{\text{Nis}} X$.*

Proof For a pointed simplicial set X , there is a map $S^1 \wedge X \rightarrow S^1 \wedge L_P X$ which induces P -localization on homology, and therefore there is a weak equivalence $L_P(S^1 \wedge X) \simeq S^1 \wedge L_P X$. This is promoted to the setting of simply connected objects in $\mathbf{sPre}(\mathbf{Sm}_k)_+$ by arguing at points.

A similar argument applies to ΩX using homotopy in place of homology. □

Proposition 3.8 *If X and Y are objects in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and P is a set of primes, then $L_P(X \times Y) \simeq L_P X \times L_P Y$.*

Proof The object $L_P X \times L_P Y$ is P -locally weakly equivalent to $X \times Y$, and therefore to $L_P(X \times Y)$, by Corollary 2.2. Since $L_P X \times L_P Y$ is P -locally fibrant, the result follows. □

Proposition 3.9 *Suppose X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$ and P is a set of prime numbers; then $X \rightarrow L_P X$ induces an isomorphism on π_0 .*

Proof For any simplicial set K , the set of maps

$$\text{SMap}(S^1 \times K, S^0) \rightarrow \text{SMap}(S^1 \times K, S^0)$$

induced by multiplication by n in S^1 is a bijection, the simplicial set $\text{SMap}(S^1 \times K, S^0)$ depending only on the components of K .

Then the map $X \rightarrow L_P X$ induces an equivalence of mapping objects $\text{Map}(L_P X, S^0) \rightarrow \text{Map}(X, S^0)$, from which the result follows. □

3.1 P and \mathbb{A}^1 localization

Proposition 3.10 *If X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)$ such that X is \mathbb{A}^1 -local and $\pi_1(X)$ is abelian, then $L_P X$ is again \mathbb{A}^1 -local.*

Proof By Proposition 3.9, $L_P X$ is again connected.

Under the hypotheses, it suffices to check that the sheaves $\pi_i(X) \otimes_{\mathbb{Z}} \mathbb{Z}_P$ are strictly \mathbb{A}^1 -invariant [33, Chapter 6], but this follows immediately since the functor $\cdot \otimes_{\mathbb{Z}} \mathbb{Z}_P$ is exact. \square

In the sequel, we consider only the composite localization $L_P L_{\mathbb{A}^1} X$, and not the reverse. The proposition says that, under connectivity hypotheses, $L_P L_{\mathbb{A}^1} X$ is both \mathbb{A}^1 - and P -local.

If X is a connected H -space in $\mathbf{sPre}(\mathbf{Sm}_k)_*$, then it is possible to define self maps

$$\times n: X \xrightarrow{\Delta} X^n \xrightarrow{\mu(\mu(\dots\mu))} X$$

by composing the n -fold diagonal and an iterated multiplication map. The map $\times n$ represents a class in $[X, X]$, which we also denote by $\times n$ in an abuse of notation.

Proposition 3.11 *If X is a connected H -space in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and P is a set of primes, then $L_P X$ is again a connected H -space, and the map $X \rightarrow L_P X$ is a weak equivalence if and only if $\times n \in [X, X]$ is invertible for all n not divisible by the primes of P .*

Proof The object $L_P X$ carries an H -space structure since $L_P(X \times X) \simeq L_P X \times L_P X$; see Proposition 3.8.

An Eckmann–Hilton argument implies that X is simple, that is, the action of $\pi_1(X)$ on $\pi_i(X)$ is trivial for all i , and moreover $\times n$ induces multiplication by n on all homotopy sheaves. The result follows. \square

Proposition 3.12 *Suppose X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and further that X is equipped with an H -space structure. Then $L_{\mathbb{A}^1} L_P X \simeq L_P L_{\mathbb{A}^1} X$, where the localizations are carried out with respect to either the local or the flasque model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$.*

Proof We give the proof in the local case; the flasque is the same mutatis mutandis.

Starting with the Quillen adjunction from the injective local model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$ to the \mathbb{A}^1 -local, we obtain a commutative diagram of model structures,

where the maps indicated are left Quillen adjoints,

$$(8) \quad \begin{array}{ccc} \text{local} & \longrightarrow & \mathbb{A}^1 \\ \downarrow & & \downarrow \\ P\text{-local} & \longrightarrow & P\text{-}\mathbb{A}^1 \end{array}$$

where the $P\text{-}\mathbb{A}^1$ model structure is the P -localization in the evident sense of the \mathbb{A}^1 model structure.

We claim that for a connected H -space object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, the maps $X \rightarrow L_{\mathbb{A}^1} L_P X$ and $X \rightarrow L_P L_{\mathbb{A}^1} X$ are both fibrant replacements in the $P\text{-}\mathbb{A}^1$ model structure, and therefore that $L_{\mathbb{A}^1} L_P X \simeq L_P L_{\mathbb{A}^1} X$ in the original model structure.

The lynchpin of the following argument is the observation, by reference to [12, Proposition 3.4.1], an object W of $\mathbf{sPre}(\mathbf{Sm}_k)$ is $P\text{-}\mathbb{A}^1$ -local if it satisfies the following three conditions:

- (i) W is fibrant in the injective model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$.
- (ii) For any object U of \mathbf{Sm}_k , the maps

$$\text{SMap}(U, W) \rightarrow \text{SMap}(U \times \mathbb{A}^1, W)$$

of simplicial mapping objects are weak equivalences.

- (iii) For any ρ_n^k where $k \geq 1$ and n is not divisible by a prime in P , the maps induced by ρ_n^k ,

$$\text{SMap}(S_\tau^k, W) \rightarrow \text{SMap}(S_\tau^k, W),$$

are weak equivalences.

The object $L_P L_{\mathbb{A}^1} X$ is both \mathbb{A}^1 -fibrant, by Proposition 3.10, and P -locally fibrant, and it is therefore a P -local object in the \mathbb{A}^1 model structure. By reference to [12, Proposition 3.4.1], it is fibrant in the $P\text{-}\mathbb{A}^1$ model structure. Since $X \rightarrow L_{\mathbb{A}^1} X$ is an \mathbb{A}^1 weak equivalence, it is a fortiori a $P\text{-}\mathbb{A}^1$ weak equivalence, and therefore $X \rightarrow L_{\mathbb{A}^1} X \rightarrow L_P L_{\mathbb{A}^1} X$ is a $P\text{-}\mathbb{A}^1$ weak equivalence, and therefore a fibrant replacement.

In the other case, we argue similarly. The object $L_{\mathbb{A}^1} L_P X$ is \mathbb{A}^1 -fibrant. Moreover, $L_P X$ is an H -space for which $\times n \in [L_P X, L_P X]$ is an isomorphism. Since the natural transformation $\text{id} \rightarrow L_{\mathbb{A}^1}$ induces a morphism of H -spaces, it follows that $\times n \in [L_{\mathbb{A}^1} L_P X, L_{\mathbb{A}^1} L_P X]$ is invertible as well, so by Proposition 3.11, it follows that $L_{\mathbb{A}^1} L_P X$ is P -fibrant. Moreover, $X \rightarrow L_P X$ is a P -local weak equivalence, and therefore a $P\text{-}\mathbb{A}^1$ weak equivalence, and consequently $X \rightarrow L_P X \rightarrow L_{\mathbb{A}^1} L_P X$ is a $P\text{-}\mathbb{A}^1$ -fibrant replacement. \square

We recall from [33, Chapter 2] that there is a construction on presheaves of groups, \mathcal{G} , given by

$$\mathcal{G}_{-1}: U \mapsto \ker(\mathcal{G}(\mathbb{G}_m \times X) \xrightarrow{\text{ev}(1)} \mathcal{G}(X)),$$

where $\text{ev}(1)$ is evaluation at 1 in \mathbb{G}_m . Equivalently, \mathcal{G}_{-1} is the kernel of the map of group sheaves $\text{Map}(\mathbb{G}_m, \mathcal{G}) \rightarrow \text{Map}(*, \mathcal{G}) \cong \mathcal{G}$. The assignment $\mathcal{G} \mapsto \mathcal{G}_{-1}$ is functorial, and sends sheaves to sheaves. The j -fold iterate of the “ -1 ” functor applied to \mathcal{G} is denoted by \mathcal{G}_{-j} .

Theorem 6.13 of [33] says that if X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, then $\pi_{i+j\alpha}^{\mathbb{A}^1}(X) = \pi_i^{\mathbb{A}^1}(X)_{-j}$. Recall that $\pi_{i+j\alpha}^{\mathbb{A}^1}(X)$ is notation for $\pi_i(\text{Map}(\mathbb{G}_m^{\wedge j}, L_{\mathbb{A}^1} X))$.

Proposition 3.13 *If \mathcal{G} is an abelian sheaf of groups, then \mathcal{G}_{-1} is also abelian and there is a natural isomorphism $(R \otimes \mathcal{G})_{-1} \cong R \otimes \mathcal{G}_{-1}$.*

Proof The abelian property of \mathcal{G}_{-1} follows immediately from the definition.

For any object U of \mathbf{Sm}_k , we have a natural commutative diagram of left-exact sequences

$$\begin{array}{ccccccc} 1 & \longrightarrow & (R \otimes \mathcal{G})_{-1}(U) & \longrightarrow & (R \otimes \mathcal{G})(\mathbb{G}_m \times U) & \longrightarrow & (R \otimes \mathcal{G})(U) \\ & & \downarrow f & & \parallel & & \parallel \\ 1 & \longrightarrow & R \otimes \mathcal{G}_{-1}(U) & \longrightarrow & R \otimes (\mathcal{G}(\mathbb{G}_m \times U)) & \longrightarrow & R \otimes (\mathcal{G}(U)) \end{array}$$

from which the natural isomorphism $(R \otimes \mathcal{G})_{-1} \cong R \otimes \mathcal{G}_{-1}$ follows. □

Proposition 3.14 *If j is a nonnegative integer, X is a simply connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and P is a set of primes, then there is a natural isomorphism*

$$\text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P L_{\mathbb{A}^1} X) \rightarrow L_P \text{Map}_*(\mathbb{G}_m^{\wedge j}, L_{\mathbb{A}^1} X)$$

in $\text{ho}_{\text{Nis}} \mathbf{sPre}(\mathbf{Sm}_k)_*$.

Proof Each of the two spaces in question is equipped with a natural map to

$$L_P \text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P L_{\mathbb{A}^1} X).$$

It suffices to show that each of these maps is a simplicial weak equivalence.

By Proposition 3.10, the space $L_P L_{\mathbb{A}^1} X$ is \mathbb{A}^1 -local. By the unstable \mathbb{A}^1 -connectivity theorem [33, Theorem 6.38], it is also connected. As is shown in the proof of [33,

Theorem 6.13], the functor $\text{Map}_*(\mathbb{G}_m^{\wedge j}, \cdot)$ preserves the subcategory of connected, \mathbb{A}^1 -local objects in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

Let Y denote either $L_{\mathbb{A}^1} X$ or $L_P L_{\mathbb{A}^1} X$, both of which are \mathbb{A}^1 -local and connected. Then, for any $i \geq 1$, the homotopy sheaf $\pi_i(\text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P Y))$ is naturally isomorphic to

$$\begin{aligned} \pi_i(L_P Y)_{-j} &\cong \pi_i^{P, \mathbb{A}^1}(Y)_{-j} \cong \pi_i^{\mathbb{A}^1}(Y)_{-j} \otimes_{\mathbb{Z}} \mathbb{Z}_P \cong \pi_i(\text{Map}_*(\mathbb{G}_m^{\wedge j}, Y)) \otimes_{\mathbb{Z}} \mathbb{Z}_P \\ &\cong \pi_i(L_P \text{Map}_*(\mathbb{G}_m^{\wedge j}, Y)). \end{aligned}$$

In particular, the spaces $\text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P L_{\mathbb{A}^1} X)$ and $L_P \text{Map}_*(\mathbb{G}_m^{\wedge j}, L_{\mathbb{A}^1} X)$ are both weakly equivalent to the space $L_P \text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P L_{\mathbb{A}^1} X)$, as required. \square

Definition 3.15 For an object X of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and nonnegative integers i and j , the notation $\pi_{i+j\alpha}^{P, \mathbb{A}^1}(X)$ is used to denote $\pi_i(\text{Map}_*(\mathbb{G}_m^{\wedge j}, L_P L_{\mathbb{A}^1} X))$.

Proposition 3.16 *If i and j are nonnegative integers, $L_{\mathbb{A}^1} X$ is a simply connected, \mathbb{A}^1 -local object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and P is a set of primes, then there are natural isomorphisms*

$$\pi_{i+j\alpha}^{P, \mathbb{A}^1}(X) \cong \pi_i^{P, \mathbb{A}^1}(X)_{-j} \cong \pi_i^{\mathbb{A}^1}(X)_{-j} \otimes_{\mathbb{Z}} \mathbb{Z}_P \cong \pi_{i+j\alpha}^{\mathbb{A}^1}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_P.$$

Proof The sheaf $\pi_{i+j\alpha}^{P, \mathbb{A}^1}(X)$ is isomorphic to $\pi_{i+j\alpha}^{\mathbb{A}^1}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_P$ by Proposition 3.14. This is isomorphic to

$$\pi_i^{\mathbb{A}^1}(X)_{-j} \otimes_{\mathbb{Z}} \mathbb{Z}_P \cong \pi_i^{P, \mathbb{A}^1}(X)_{-j},$$

as required. \square

Proposition 3.17 *If P is a set of primes and if X, Y and Z are simply connected objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $X \rightarrow Y \rightarrow Z$ is a P - \mathbb{A}^1 -fiber sequence up to homotopy, and if j is a nonnegative integer, then there is a natural long exact sequence*

$$\cdots \rightarrow \pi_{i+j\alpha}^{P, \mathbb{A}^1}(X) \rightarrow \pi_{i+j\alpha}^{P, \mathbb{A}^1}(Y) \rightarrow \pi_{i+j\alpha}^{P, \mathbb{A}^1}(Z) \rightarrow \pi_{i-1+j\alpha}^{P, \mathbb{A}^1}(X) \rightarrow \cdots .$$

Proof This is an immediate consequence of Definition 3.15. \square

3.2 P -localization of spectra

Throughout this section, the underlying model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is taken to be the flasque, rather than the injective.

One can construct a P -local model category of presheaves of spectra, following [16], as the S^1 -stable model category on the P -local flasque model structure on $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

Lemma 3.18 *The adjunction*

$$\Sigma^\infty : \mathbf{sPre}(\mathbf{Sm}_k)_* \rightleftarrows \mathbf{Spt}(\mathbf{Sm}_k) : \text{Ev}_0$$

is a Quillen adjunction between the P -local model categories.

Proof This is implicit in [16], being the combination of Proposition 1.16 and the definition of the stable model structure as a localization of the level model structure on spectra. □

Explicitly, the fibrant-replacement functor, L_P , in $\mathbf{Spt}(\mathbf{Sm}_k)$ with the P -local model structure is given by

$$(L_P E)_i = \text{colim}_{k \rightarrow \infty} \text{Map}_*(S^k, L_P E_{i+k}).$$

With the P -local model structures and the smash product, the category $\mathbf{Spt}(\mathbf{Sm}_k)$ is a $\mathbf{sPre}(\mathbf{Sm}_k)_*$ model category, in the sense of [15, Chapter 4.2].

Lemma 3.19 *If s is a point of $\mathbf{Sh}_{\text{Nis}}(\mathbf{Sm}_k)$, the adjunction*

$$s^* : \mathbf{Spt}(\mathbf{Sm}_k) \rightleftarrows \mathbf{Spt} : s_*$$

is a Quillen adjunction between the P -local model categories.

Corollary 3.20 *For any object X of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and any set P of primes, there is a stable weak equivalence*

$$\Sigma^\infty L_P X \rightarrow L_P \Sigma^\infty X.$$

A spectrum E is said to be P -local if it is fibrant and the map $E \rightarrow L_P E$ is a stable weak equivalence. Since it is possible to check stable weak equivalence of spectra at points, we deduce the following by arguing at points:

Proposition 3.21 *A spectrum E is P -local if and only if it is fibrant and the maps*

$$E \xrightarrow{n} E$$

are weak equivalences for all n not divisible by the primes in P .

Proposition 3.22 *A spectrum E is P -local if and only if it is fibrant and the localization maps $\pi_i^s(E) \rightarrow \pi_i^s(E) \otimes_{\mathbb{Z}} \mathbb{Z}_P$ are isomorphisms for all i .*

3.3 P - and \mathbb{A}^1 -localization of spectra

We begin with a commutative diagram of model structures on the category $\mathbf{Spt}(\mathbf{Sm}_k)$, which is the application of [16] to the flasque version of diagram (8):

$$\begin{array}{ccc} \text{stable} & \longrightarrow & \mathbb{A}^1 \\ \downarrow & & \downarrow \\ P\text{-stable} & \longrightarrow & P\text{-}\mathbb{A}^1\text{-stable} \end{array}$$

Fibrant replacements in the \mathbb{A}^1 - or P -local model structures are effected by replacing E by the spectrum that has level i given by

$$\operatorname{colim}_k \Omega^k L_{\mathbb{A}^1} E_{i+k}$$

or

$$\operatorname{colim}_k \Omega^k L_P E_{i+k},$$

respectively, L_P and $L_{\mathbb{A}^1}$ being taken in the flasque model structures. The stable fibrant replacements are denoted by $L_P E$ and $L_{\mathbb{A}^1} E$.

Lemma 3.23 *The classes of P -locally flasque fibrant and $P\text{-}\mathbb{A}^1$ -locally flasque fibrant objects in $\mathbf{sPre}(\mathbf{Sm}_k)$ and $\mathbf{sPre}(\mathbf{Sm}_k)_*$ are closed under filtered colimits.*

Proof Since an object is $P\text{-}\mathbb{A}^1$ -locally flasque fibrant if and only if it is both P - and \mathbb{A}^1 -locally flasque fibrant, it suffices to prove the case of P -locally flasque fibrant objects.

Suppose X_k is a filtered diagram of P -locally flasque fibrant objects; then $\operatorname{colim}_k X_k$ is flasque fibrant, by [18]. We wish to show that for any $\rho_n^n \times \operatorname{id}_U: S_\tau^k \times U \rightarrow S_\tau^k \times U$ in T_P , the induced map

$$\operatorname{SMap}(S_\tau^k \times U, \operatorname{colim}_k X_k) \rightarrow \operatorname{SMap}(S_\tau^k \times U, \operatorname{colim}_k X_k)$$

is a weak equivalence. Since $S_\tau^k \times U$ is equivalent to a compact object, $S^k \times U$, of $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and since the X_k and $\operatorname{colim}_k X_k$ are all fibrant, and $S_\tau^k \times U$ and $S^k \times U$ are all cofibrant, we wish to show that the induced map

$$\operatorname{colim}_k \operatorname{SMap}(S_\tau^k \times U, X_k) \rightarrow \operatorname{colim}_k \operatorname{SMap}(S_\tau^k \times U, X_k)$$

is a weak equivalence of simplicial sets, but since the X_k are themselves P -locally fibrant, this is immediate. \square

Proposition 3.24 *For any object E of $\mathbf{Spt}(\mathbf{Sm}_k)$ there is a stable weak equivalence $L_P L_{\mathbb{A}^1} E \simeq L_{\mathbb{A}^1} L_P E$.*

Proof The objects in question are levelwise fibrant for the flasque model structure. It suffices therefore to show that they are levelwise weakly equivalent for the flasque model structure. Since we are working the flasque model structure, filtered colimits of fibrant objects are again fibrant, and so we deduce the existence of weak equivalences

$$L_{\mathbb{A}^1} \operatorname{colim}_k X_k \simeq L_{\mathbb{A}^1} \operatorname{colim}_k L_{\mathbb{A}^1} X_k \simeq \operatorname{colim}_k L_{\mathbb{A}^1} X_k,$$

and similarly for L_P .

We may assume that the spaces E_i appearing are all simply connected H -spaces, and therefore $L_P L_{\mathbb{A}^1} E_i \simeq L_{\mathbb{A}^1} L_P E_i$. We then have

$$\operatorname{colim}_k \Omega^k L_{\mathbb{A}^1} (\operatorname{colim}_{k'} \Omega^{k'} L_P E_{k+k'+i}) \simeq \operatorname{colim}_k \Omega^k (\operatorname{colim}_{k'} \Omega^{k'} L_{\mathbb{A}^1} L_P E_{k+k'+i}),$$

which is symmetric in $L_{\mathbb{A}^1}$ and L_P , up to weak equivalence, whence the result. \square

We therefore conflate $L_P L_{\mathbb{A}^1} E$ and $L_{\mathbb{A}^1} L_P E$, calling either the P - \mathbb{A}^1 -localization of E . We say that a map of spectra $f: E \rightarrow E'$ is a P - \mathbb{A}^1 weak equivalence if $L_P L_{\mathbb{A}^1} f$ is a stable weak equivalence of spectra, or equivalently if $L_P f$ is an \mathbb{A}^1 weak equivalence of spectra, or equivalently again if $L_{\mathbb{A}^1} f$ is a P -local equivalence of spectra. We write $\pi_i^{P, \mathbb{A}^1, s}(E)$ for the homotopy sheaves $\pi_i^s(L_P L_{\mathbb{A}^1} E)$.

Proposition 3.25 *If E is an object in $\mathbf{Spt}(\mathbf{Sm}_k)$ and P is a set of prime numbers, then there is a natural isomorphism*

$$\pi_i^{s, P, \mathbb{A}^1}(E) \cong \pi_i^{s, \mathbb{A}^1}(E) \otimes_{\mathbb{Z}} \mathbb{Z}_P.$$

Proof This is immediate from the above. \square

Proposition 3.26 *If $f: E \rightarrow E'$ is a map in $\mathbf{Spt}(\mathbf{Sm}_k)$ and P is a set of prime numbers, then f is a P - \mathbb{A}^1 weak equivalence if and only if $\pi_i^{s, \mathbb{A}^1}(f) \otimes_{\mathbb{Z}} \mathbb{Z}_P$ is an isomorphism of abelian groups for all i .*

Proof This is immediate from the above. \square

Proposition 3.27 Suppose $\{E_n\}$ is a filtered system of objects in $\mathbf{Spt}(\mathbf{Sm}_k)$ and P is a set of prime numbers; then the natural maps

$$\operatorname{colim}_n \pi_i^{s,P}(E_n) \rightarrow \pi_i^{s,P}(\operatorname{colim}_n E_n)$$

and

$$\operatorname{colim}_n \pi_{i+j\alpha}^{s,P,\mathbb{A}^1}(E_n) \rightarrow \pi_{i+j\alpha}^{s,P,\mathbb{A}^1}(\operatorname{colim}_n E_n)$$

are isomorphisms.

Proof This is immediate from the above and Proposition 2.18. □

Proposition 3.28 Suppose that $f: X \rightarrow Y$ is a map of simply connected objects $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $\Sigma^\infty f$ is a $P-\mathbb{A}^1$ -stable weak equivalence. Then f is a $P-\mathbb{A}^1$ weak equivalence.

Proof The map $L_P L_{\mathbb{A}^1} f: L_P L_{\mathbb{A}^1} \Sigma^\infty X \rightarrow L_P L_{\mathbb{A}^1} \Sigma^\infty Y$ is a weak equivalence, and this map agrees in the stable homotopy category with the map $L_P L_{\mathbb{A}^1} \Sigma^\infty L_{\mathbb{A}^1} X \rightarrow L_P L_{\mathbb{A}^1} \Sigma^\infty L_{\mathbb{A}^1} Y$. We may commute L_P past $L_{\mathbb{A}^1}$ and past Σ^∞ , so we conclude that $L_{\mathbb{A}^1} \Sigma^\infty L_P L_{\mathbb{A}^1} X \rightarrow L_{\mathbb{A}^1} \Sigma^\infty L_P L_{\mathbb{A}^1} Y$ is a weak equivalence. By Corollary 2.23, since $L_P L_{\mathbb{A}^1} X$ and $L_P L_{\mathbb{A}^1} Y$ are simply connected, we deduce that $L_{\mathbb{A}^1} L_P L_{\mathbb{A}^1} X \rightarrow L_{\mathbb{A}^1} L_P L_{\mathbb{A}^1} Y$ is a weak equivalence, and since $L_P L_{\mathbb{A}^1} X, L_P L_{\mathbb{A}^1} Y$ are already \mathbb{A}^1 -local by Proposition 3.10, the result follows. □

4 The Grothendieck–Witt group

4.1 The homotopy of spheres

Consider a motivic sphere $X = S^{n+q\alpha} = S^n \wedge \mathbb{G}_m^{\wedge q}$.

We make frequent use of the following result, which is a paraphrase of some results of [33, Section 6.3]:

Lemma 4.1 (Morel) *If (n, q) and (n', q') are pairs of nonnegative integers and if $n \geq 2$, then*

$$\pi_{n+q\alpha}^{\mathbb{A}^1}(S^{n'+q'\alpha}) = \begin{cases} 0 & \text{if } n < n', \\ \mathbb{K}_{q'-q}^{\text{MW}} & \text{if } n = n' \text{ and } q' > 0, \\ 0 & \text{if } n = n', q' = 0 \text{ and } q > 0, \\ \mathbb{Z} & \text{if } n = n' \text{ and } q = q' = 0. \end{cases}$$

The stable version of this result was known earlier, but may be deduced from the unstable.

Corollary 4.2 *If (n, q) and (n', q') are pairs of integers with q and q' nonnegative, then the sets of maps between $S^{n+q\alpha}$ and $S^{n'+q'\alpha}$ in the \mathbb{A}^1 homotopy category of S^1 -spectra take the form*

$$\pi_{n+q\alpha}^{s, \mathbb{A}^1}(S^{n'+q'\alpha}) = \begin{cases} 0 & \text{if } n < n', \\ \mathbb{K}_{q'-q}^{\text{MW}} & \text{if } n = n' \text{ and } q' > 0, \\ 0 & \text{if } n = n', q' = 0 \text{ and } q > 0, \\ \mathbb{Z} & \text{if } n = n' \text{ and } q = q' = 0. \end{cases}$$

We remark that \mathbb{K}_0^{MW} is the sheaf of Grothendieck–Witt groups, also denoted by GW . We observe that if $q > 0$, then, by Corollary 2.17,

$$[\Sigma^\infty S^{n+q\alpha}, \Sigma^\infty S^{n'+q'\alpha}]_{\mathbb{A}^1} = \text{GW}(*).$$

Proposition 4.3 *Suppose n, n', q and q' are integers such that n, q and q' are nonnegative and $q' \geq 1$. We have an identification*

$$\pi_{n+q\alpha}^{s, P, \mathbb{A}^1}(S^{n'+q'\alpha}) = \begin{cases} 0 & \text{if } n < n', \\ \mathbb{K}_{q'-q}^{\text{MW}} \otimes_{\mathbb{Z}} \mathbb{Z}_P & \text{if } n = n'. \end{cases}$$

Proof This follows immediately from Corollary 4.2 and Proposition 3.25. □

Remark 4.4 Since \mathbb{Z} is a subring of $\mathbb{K}_0^{\text{MW}}(k) = \text{GW}(k)$, it follows that \mathbb{Z}_P is a subring of $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_P$.

4.2 Twist classes

For $a \in k^*$, following [33, Chapter 3], we define $\langle a \rangle: S^{(1,0)} \wedge \mathbb{G}_m \rightarrow S^{(1,0)} \wedge \mathbb{G}_m$ to be the map induced by multiplication $a: \mathbb{G}_m \rightarrow \mathbb{G}_m$ by forming $a_+: (\mathbb{G}_m)_+ \rightarrow \mathbb{G}_m$, suspending

$$S^{(1,0)} \wedge (a_+): S^{(1,0)} \wedge (\mathbb{G}_m)_+ \cong S^{1,0} \vee S^{(1,0)} \wedge \mathbb{G}_m \rightarrow S^{(1,0)} \wedge \mathbb{G}_m$$

and letting $\langle a \rangle$ denote the restriction of this map to the $S^{(1,0)} \wedge \mathbb{G}_m$ summand.

Remark 4.5 The interchange of any two terms in

$$(S^{n+q\alpha})^{\wedge r} = S^{n+q\alpha} \wedge S^{n+q\alpha} \wedge \dots \wedge S^{n+q\alpha} \cong S^{rn+r q\alpha}$$

represents the element

$$e_{n,q} = (-1)^{n+q} \langle -1 \rangle^q \in \pi_{rn+rq\alpha}^{\mathbb{A}^1}(S^{rn+rq\alpha}),$$

by [33, Lemma 3.43]. Observe that $e_{n,q}^2 = 1$.

Much of the following work depends on showing a class $A + B\langle -1 \rangle$, where A and B are integers, is a unit in the ring $\text{GW}(k)$ or $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_{(2)}$.

We remind the reader that a field k is said to be *formally real* if -1 cannot be written as a sum of squares in k [27, Chapter VIII].

Proposition 4.6 *Suppose A and B are integers, and let R be a localization of \mathbb{Z} . Then $A + B\langle -1 \rangle$ is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} R$ if and only if one of the following conditions is met:*

- (i) $A + B$ and $A - B$ are units in R and k is formally real.
- (ii) $A + B$ is a unit in R and the field k is not formally real.

Proof We remark that the dimension homomorphism makes R into a split subring of $\text{GW}(k) \otimes_{\mathbb{Z}} R$.

We first handle the case where k is formally real.

Since $(A + B\langle -1 \rangle)(A + B\langle -1 \rangle) = A^2 - B^2 = (A + B)(A - B)$, the condition in (i) is sufficient.

We may embed k in a real closure $\phi: k \rightarrow k^r$. This embedding induces a ring homomorphism $\phi: \text{GW}(k) \otimes_{\mathbb{Z}} R \rightarrow \text{GW}(k^r) \otimes_{\mathbb{Z}} R = R \oplus R\langle -1 \rangle$ [27, Proposition II.3.2]. Abstractly, the ring $\text{GW}(k^r) \otimes_{\mathbb{Z}} R$ is endowed with an automorphism $\langle -1 \rangle \mapsto -\langle -1 \rangle$, and if $\phi(A + B\langle -1 \rangle) = A + B\langle -1 \rangle$ is a unit, then so too is $A - B\langle -1 \rangle$, from which we deduce that their product, $(A + B\langle -1 \rangle)(A - B\langle -1 \rangle) = A^2 - B^2$, is a unit as well. But $A^2 - B^2$ is a unit if and only if $A + B$ and $A - B$ are units, showing that this condition is necessary and sufficient if k is formally real.

Suppose now that k is not formally real.

We may employ the dimension map $\text{GW}(k) \otimes_{\mathbb{Z}} R \rightarrow R$ to show that if $A + B\langle -1 \rangle$ is a unit, then necessarily $A + B$ is a unit, u .

We wish to show that this condition is also sufficient to imply $A + B\langle -1 \rangle$ is a unit. If the characteristic of k is 2, then $\langle -1 \rangle = 1$ and the result is trivial. We may therefore assume the characteristic of k is not 2. Write $A = u - B$.

Suppose first the characteristic of k is not 2. In the not formally real case, the Witt group $W(k) = \text{GW}(k)/(1 + \langle -1 \rangle)$ is 2-primary torsion [27, Theorem VIII.3.6]. Moreover, the class $1 - \langle -1 \rangle$ lies in the fundamental ideal of $\text{GW}(k)$, which is isomorphic to its image in $W(k)$, so it follows that $1 - \langle -1 \rangle$ is 2-primary torsion. Since $(1 - \langle -1 \rangle)^N = 2^{N-1}(1 - \langle -1 \rangle)$ by an easy induction, the element $1 - \langle -1 \rangle$ is nilpotent in $\text{GW}(k)$ and therefore also in $\text{GW}(k) \otimes_{\mathbb{Z}} R$. We deduce $A + B\langle -1 \rangle = u - B(1 - \langle -1 \rangle)$ is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} R$, as required. \square

We owe the argument in the not formally real case to the anonymous referee.

We employ Proposition 4.6 via the following two corollaries:

Corollary 4.7 *Suppose m is a nonnegative integer and $e_{n,q} = (-1)^{n+q}\langle -1 \rangle^q$ is the twist class of the sphere $S^{n+q\alpha}$. Then the class $1 + m + me_{n,q}$ is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_{(2)}$.*

Proof There are, in general, four cases, $e_{n,q} = \pm 1$ and $e_{n,q} = \pm \langle -1 \rangle$, although it is possible that $\langle -1 \rangle = 1$ in $\text{GW}(k)$. The two cases $e_{n,q} = \pm 1$ are immediate.

For the other two cases, by the proposition, it suffices to check that one or both of $m + 1 + m = 2m + 1$ and $m + 1 - m = 1$ are units in $\mathbb{Z}_{(2)}$, which they both are. \square

Corollary 4.8 *Suppose m is a positive integer and $e_{n,q} = (-1)^{n+q}\langle -1 \rangle^q$ is the twist class of the sphere $S^{n+q\alpha}$. Then the class $1 + m + me_{n,q}$ is a unit in $\text{GW}(k)$ if and only if one of the following conditions holds:*

- n is odd and q is even.
- $n + q$ is odd and k is not formally real.

Proof The cases where q is even, so $e_{n,q} = \pm 1$, are easily dealt with and do not depend on the field. Assume therefore that q is odd.

Suppose k is formally real, then the proposition says that $1 + m \pm m\langle -1 \rangle$ is a unit if and only if 1 and $1 + 2m$ are units in \mathbb{Z} . Therefore there are no cases where the class is a unit, q is odd and k is formally real.

Suppose k is not formally real and q is odd. Then, by the proposition, $1 + m + me_{n,q}$ is a unit if and only if $1 + m - (-1)^n m$ is a unit, whereupon it is necessary and sufficient for n to be even. \square

5 The Hilton–Milnor splitting

The James construction on a pointed simplicial set was introduced by James in [19]. The idea of applying it in \mathbb{A}^1 homotopy theory, and thereby obtaining a weak equivalence $J(X) \simeq \Omega \Sigma X$ as in Proposition 5.2, is not original to us. We learned of it from Asok and Fasel, who attribute it to Morel.

Our presentation is based on that of [39, Chapter VII, Section 2], and we also refer frequently to [41]. Suppose X is a pointed simplicial set. An injection $\alpha: (1, 2, \dots, n) \rightarrow (1, 2, \dots, m)$ induces a map $\alpha_*: X^n \rightarrow X^m$. Let \sim denote the equivalence relation on $\coprod_{n=0}^\infty X^n$ generated by $x \sim \alpha_*(x)$ for all injections α . The James construction on X is $J(X) = \coprod_{n=0}^\infty X^n / \sim$. The construction $J(X)$ is the free monoid on the pointed simplicial set X . The k -simplices $J(X)_k$ of $J(X)$ are the free monoids on the pointed sets X_k , that is, $J(X)_k = \coprod_n X_k^n / \sim$. The James construction is filtered by pointed simplicial sets $J_n(X)$, defined by $J_n(X) = \coprod_{m=0}^n X^m / \sim$.

Define spaces $D_n(X)$ as the cofibers of sequences

$$J_{n-1}(X) \rightarrow J_n(X) \rightarrow D_n(X).$$

There are canonical weak equivalences $D_n(X) \rightarrow X^{\wedge n}$. Define $D(X) = \bigvee_{n=0}^\infty D_n(X)$.

Definition 5.1 For a pointed simplicial presheaf X , define $J(X)$, $J_n(X)$, $D_n(X)$ and $D(X)$ in \mathbf{sPre} by

$$\begin{aligned} J(X)(U) &= J(X(U)), & J_n(X)(U) &= J_n(X(U)), \\ D_n(X)(U) &= D_n(X(U)), & D(X)(U) &= D(X(U)). \end{aligned}$$

Let $\ell: X = J_1(X) \rightarrow J(X)$ denote the map induced by the canonical maps $X(U) \rightarrow J(X(U))$ for $U \in \mathbf{Sm}$.

The *James construction* is then defined to be $L_{\mathbb{A}^1} J(X)$.

We learned the following result from Asok and Fasel. It is a functorial version of a result due to James [19, Theorem 5.6], and is presented in more recent terminology in [39, Chapter VII, Theorem 2.6].

Proposition 5.2 *Suppose X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$. There is a natural isomorphism $J(X) \rightarrow \Omega \Sigma X$ in $\mathbf{hoNis} \mathbf{sPre}(\mathbf{Sm}_k)_*$.*

Proof For any object U in \mathbf{Sm}_k , there is a functorial map $j: J(X(U)) \rightarrow F(X(U))$, where $F(X(U))$ is Milnor’s construction—the free abelian group on the pointed space $X(U)$ —as laid out in [41, Sections 3.3.2 and 3.3.3]. This map is a weak equivalence when $X(U)$ is connected [41, Theorem 3.3.5]. There is a natural isomorphism $F(X(U)) \cong G\Sigma X(U)$ [11, Chapter V, Theorem 6.15]. Here, $G\Sigma X(U)$ is a fibrant model for $\Omega\Sigma(X(U))$. In particular, there is a zigzag of maps of simplicial presheaves

$$J(X) \leftarrow F(X) \xrightarrow{\cong} G\Sigma(X).$$

The formation of $J(X(U))$ and $F[X(U)]$ commute with colimits, since the free abelian monoid and the free abelian monoid functors do. Therefore, both J and F commute with taking stalks at a point p^* . Since p^*X is a connected simplicial set, it follows that $p^*J(X) \cong J(p^*X) \rightarrow F(p^*X) \cong p^*F(X)$ is a weak equivalence, whence the result. \square

Note that this natural isomorphism induces a natural isomorphism $J(X) \rightarrow \Omega\Sigma X$ in $\mathbf{ho}_{\mathbb{A}^1} \mathbf{sPre}(\mathbf{Sm}_k)_*$. It also follows immediately from this result that if $X \rightarrow Y$ is a local weak equivalence, then the functorial map $J(X) \rightarrow J(Y)$ is a local weak equivalence.

Corollary 5.3 *Suppose X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$. Then there is a natural isomorphism*

$$\pi_i^{\mathbb{A}^1} J(X) \cong \pi_{i+1}^{\mathbb{A}^1} \Sigma X.$$

Proof By Proposition 5.2, there is a natural isomorphism $\pi_i^{\mathbb{A}^1} J(X) \cong \pi_i^{\mathbb{A}^1} \Omega\Sigma X$. By definition, $\pi_i^{\mathbb{A}^1} \Omega\Sigma X = \pi_i L_{\mathbb{A}^1} \Omega\Sigma X$. Since ΣX is simplicially simply connected, $\Omega\Sigma X$ is simplicially connected. By Morel’s connectivity theorem [33, Theorem 6.38], $L_{\mathbb{A}^1} \Omega\Sigma X$ is also simplicially connected. Thus, $\pi_0 L_{\mathbb{A}^1} \Omega\Sigma X \cong *$ is strongly \mathbb{A}^1 -invariant. By [33, Theorem 6.46], it follows that the canonical morphism $L_{\mathbb{A}^1} \Omega\Sigma X \rightarrow \Omega L_{\mathbb{A}^1} \Sigma X$ is a simplicial weak equivalence. Thus, there is a natural isomorphism $\pi_i L_{\mathbb{A}^1} \Omega\Sigma X \cong \pi_{i+1} L_{\mathbb{A}^1} \Sigma X$. Combining with the previous gives the claimed natural isomorphism $\pi_i^{\mathbb{A}^1} J(X) \cong \pi_{i+1}^{\mathbb{A}^1} \Sigma X$. \square

Corollary 5.4 *Suppose $X \rightarrow Y$ is an \mathbb{A}^1 weak equivalence of connected objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$; then the functorial map $J(X) \rightarrow J(Y)$ is an \mathbb{A}^1 weak equivalence.*

Proof We will show that if $X \rightarrow Y$ is an \mathbb{A}^1 weak equivalence of connected objects, then $\Omega\Sigma X \rightarrow \Omega\Sigma Y$ is an \mathbb{A}^1 weak equivalence. Since there is a natural isomorphism

$\Omega \Sigma X \cong J(X)$ in $\text{ho } \mathbf{sPre}(\mathbf{Sm}_k)_*$, and therefore in $\text{ho}_{\mathbb{A}^1} \mathbf{sPre}(\mathbf{Sm}_k)_*$, it will follow that $J(X) \rightarrow J(Y)$ is an isomorphism in $\text{ho}_{\mathbb{A}^1} \mathbf{sPre}(\mathbf{Sm}_k)_*$, as claimed.

The following is a sequence of local weak equivalences:

$$\begin{aligned} L_{\mathbb{A}^1} \Sigma X &\xrightarrow{\cong} L_{\mathbb{A}^1} \Sigma Y && \text{(since } \mathbb{A}^1\text{-localization is simplicial),} \\ \Omega L_{\mathbb{A}^1} \Sigma X &\xrightarrow{\cong} \Omega L_{\mathbb{A}^1} \Sigma Y, \\ L_{\mathbb{A}^1} \Omega \Sigma X &\xrightarrow{\cong} L_{\mathbb{A}^1} \Omega \Sigma Y && \text{(by [33, Theorem 6.46]),} \end{aligned}$$

but this is precisely what was to be shown. □

Given $W, X \in \mathbf{sSet}_*$ and a map $f: (J_n W, J_{n-1} W) \rightarrow X$, we define the *combinatorial extension* of f ,

$$h(f): J(W) \rightarrow J(X),$$

by following the procedure of [19, Theorem 1.4 and Section 2] (cf [39, Chapter VII, Section 2]).

We first define the restriction of $h(f)$ to $J_m(W)$. For $m < n$, the restriction of $h(f)$ is the constant map. Suppose $m \geq n$. To an injection $(1, 2, \dots, n) \rightarrow (1, 2, \dots, m)$, we may associate a map $W^m \rightarrow W^n$ and therefore a map $W^m \rightarrow W^n \rightarrow J_n(W)$.

Consider the set of all $\binom{m}{n}$ increasing, n -term subsequences of $(1, 2, \dots, m)$. Order these by lexicographic ordering, reading from the right. Each sequence is an injective map $\{1, \dots, n\} \rightarrow \{1, \dots, m\}$. Taking the ordered product over all injections, we obtain a total map

$$W^m \rightarrow J_n(W)^{\binom{m}{n}}.$$

The $\binom{m}{n}$ -fold product of the map f is the map

$$J_n(W)^{\binom{m}{n}} \rightarrow X^{\binom{m}{n}}.$$

We set the restriction of $h(f)$ to $J_m(W)$ to be

$$W^m \rightarrow J_n(W)^{\binom{m}{n}} \rightarrow X^{\binom{m}{n}} \rightarrow J_{\binom{m}{n}}(X) \rightarrow J(X).$$

One checks that this is well defined.

This definition is functorial, and extends immediately to presheaves:

Definition 5.5 Given $W, X \in \mathbf{sPre}_*$ and a map $f: (J_n(W), J_{n-1}(W)) \rightarrow X$, we may define the *combinatorial extension* of f ,

$$h(f): J(W) \rightarrow J(X), \quad h(f)(U) = h(f(U)).$$

For $X \in \mathbf{sSet}_*$, the cofiber sequences $J_{n-1}(X) \rightarrow J_n(X) \rightarrow D_n(X)$ induce natural maps $(J_n(X), J_{n-1}(X)) \rightarrow D_n(X)$. For $X \in \mathbf{sPre}$, we thereby obtain maps $(J_n(X), J_{n-1}(X)) \rightarrow D_n(X)$, and consequently maps

$$j_n: J(X) \rightarrow J(D_n(X))$$

by combinatorial extension.

Let $i_n: J(D_n(X)) \rightarrow J(D(X))$ be the map induced by the canonical inclusions $D_n(X(U)) \rightarrow D(X(U))$. The monoid structure on $J(D(X(U)))$ induces multiplication maps $\mu_n: J(D(X))^n \rightarrow J(D(X))$. Consider the maps

$$\mu_{n+1} \prod_{m=0}^n i_m j_m: J(X) \rightarrow J(D(X))$$

for $n = 0, 1, 2, \dots$. It is important here that the product be ordered, and we declare it to be ordered by increasing values of m .

The composition of $i_n j_n$ with $J_{n-1}(X) \rightarrow J(X)$ is the constant based map. We'll say that the restriction of $i_n j_n$ to $J_{n-1}(X)$ is the constant map. It follows that $\mu_{n+1} \prod_{m=0}^n i_m j_m$ restricted to $J_{n-1}(X)$ equals the restriction of $\mu_{N+1} \prod_{m=0}^N i_m j_m$ to $J_{n-1}(X)$ for all $N \geq n$. Note that $J(X) = \text{colim } J_n(X)$. Thus, we may define $f: J(X) \rightarrow J(D(X))$ by

$$f = \text{colim}_n \mu_{n+1} \prod_{m=0}^n i_m j_m.$$

For convenience, extend f to $f_+: J(X)_+ \rightarrow J(D(X))$ by mapping the disjoint point via $* = X(U)^{\wedge 0} \rightarrow DX \rightarrow J(D(X))$.

Taking the simplicial suspension of f_+ , we obtain

$$\Sigma f_+: \Sigma(J(X)_+) \rightarrow \Sigma J(D(X)).$$

Let $|\cdot|: \mathbf{sSet} \rightarrow \mathbf{K}$ denote the geometric realization functor from simplicial sets to Kelly spaces, and let $\text{Simp}: \mathbf{K} \rightarrow \mathbf{sSet}$ be the right adjoint functor, which is the functor of singular simplices.

We claim that for any simplicial presheaf Y , for example $Y = D(X)$, there is an evaluation map

$$(9) \quad \Sigma J(Y) \rightarrow \text{Simp} |\Sigma Y|.$$

To see this, let $\Omega^M: \mathbf{Top} \rightarrow \mathbf{Top}$ denote the Moore loops functor [7]. There is a strictly associative multiplication $\Omega^M \times \Omega^M \rightarrow \Omega^M$. Since taking Simp commutes with finite products, there is a strictly associative multiplication on $\text{Simp } \Omega^M$, and therefore an induced commutative diagram

$$\begin{array}{ccc} Y & \xrightarrow{\quad} & \text{Simp } \Omega^M | \Sigma Y | \\ & \searrow & \nearrow \\ & J(Y) & \end{array}$$

Applying Σ , we obtain a map $\Sigma J(Y) \rightarrow \Sigma \text{Simp } \Omega^M | \Sigma Y |$. There is a natural transformation of functors $\Sigma \text{Simp} \rightarrow \text{Simp } \Sigma$ and so we have a composite

$$(10) \quad \Sigma J(Y) \rightarrow \Sigma \text{Simp } \Omega^M | \Sigma Y | \rightarrow \text{Simp } \Sigma \Omega^M | \Sigma Y |.$$

The counit of the adjunction between loops and suspension produces a natural transformation $\Sigma \Omega^M \rightarrow \text{id}$. Composing with (10) produces a map

$$\Sigma J(Y) \rightarrow \Sigma \text{Simp } \Omega^M | \Sigma Y | \rightarrow \text{Simp } \Sigma \Omega^M | \Sigma Y | \rightarrow \text{Simp } | \Sigma Y |,$$

which is what we claimed in (9).

Composing Σf_+ with (9) for $Y = D(X)$ produces a map

$$(11) \quad \Sigma J(X)_+ \rightarrow \text{Simp } | \Sigma D(X) |.$$

For each U , this map (11) evaluated at U is a weak equivalence; see [39, Chapter VII, Theorem 2.10]. Thus, (11) is a weak equivalence in the simplicial model structures on $\mathbf{sPre}(\mathbf{Sm}_k)_*$. Combining with the injective weak equivalence $\Sigma D(X) \rightarrow \text{Simp } | \Sigma D(X) |$, we have the zigzag of injective weak equivalences

$$(12) \quad \Sigma J(X)_+ \rightarrow \text{Simp } | \Sigma D(X) | \leftarrow \Sigma D(X).$$

We have shown:

Proposition 5.6 *Suppose X is a connected object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$. There is a canonical isomorphism $\Sigma J(X)_+ \rightarrow \Sigma D(X)$ in $\text{ho } \mathbf{sPre}(\mathbf{Sm}_k)_*$.*

Corollary 5.7 *There is a canonical isomorphism*

$$\sigma: J(X)_+ \rightarrow D(X)$$

in $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$.

Remark 5.8 Here and subsequently we write $J(X)_+$ and $D(X)$ in place of the stable $\Sigma^\infty J(X)_+$ and $\Sigma^\infty D(X)$ whenever the context demands S^1 -stable objects.

5.1 The low-dimensional simplices of the J construction

Definition 5.9 We say a simplicial presheaf X is n -reduced if the unique map $X \rightarrow *$ induces an isomorphism $X_i \rightarrow *_i$ for $i \leq n$. The term “0-reduced” may be abbreviated to “reduced”.

Equivalently, $|X_i(U)| = 1$ for all smooth schemes U and all $i \leq n$.

Example 5.10 The constant simplicial presheaf representing the simplicial n -sphere, $\Delta^n / \partial \Delta^n$, is $(n-1)$ -reduced.

Example 5.11 Suppose X and Y are n -reduced. Then $X \times Y$ is n -reduced.

Example 5.12 Suppose X is a n -reduced simplicial presheaf, given the unique pointed structure, and Y is a simplicial presheaf pointed by a map $s_0: * \rightarrow S$. Then $X \wedge Y$ is n -reduced. To see this, fix a smooth scheme U and consider the construction of $(X \wedge Y)(U)_i$ for $i \leq n$. It is given by the pushout

$$(13) \quad \begin{array}{ccc} (X \vee Y)(U)_i & \longrightarrow & (X \times Y)(U)_i \\ \downarrow & & \downarrow \\ * & \longrightarrow & (X \wedge Y)(U)_i \end{array}$$

but since X is n -reduced,

$$\begin{aligned} (X \vee Y)(U)_i &= X(U)_i \vee Y(U)_i = Y(U)_i, \\ (X(U) \times Y(U))_i &= X(U)_i \times Y(U)_i = Y(U)_i. \end{aligned}$$

In particular, the top horizontal arrow of diagram (13) is a bijection, whence so too is the bottom arrow.

Example 5.13 As a special case of the above, our models for the motivic spheres, $S^n \wedge \mathbb{G}_m^m = S^{n+m\alpha}$ are $(n-1)$ -reduced. Note that the space $S^{n+m\alpha} \wedge S^{n+m\alpha}$ is $(n-1)$ -reduced but not n -reduced, whereas the weakly equivalent space $S^{2n+2m\alpha}$ is $(2n-1)$ -reduced. This holds even when $m = 0$, that is, in the case of classical homotopy theory.

Proposition 5.14 Let X be an n -reduced simplicial presheaf. Then $J(X)$ is n -reduced.

Proof Let $i \leq n$. We can calculate $J(X)(U)_i$ directly. Since the category of simplicial presheaves on \mathbf{Sm}_k is really the category of presheaves of sets on $\mathbf{Sm}_k \times \Delta$, where Δ is the standard simplex category, both evaluation at a smooth scheme U and taking i^{th} simplices (which is “evaluation at Δ^i ”) commute with all limits and colimits.

Therefore, we may calculate

$$J(X)(U)_i = \left(\prod_{m=0}^{\infty} X(U)^m / \sim \right)_i = \prod_{m=0}^{\infty} (X(U)_i)^m / \sim$$

but this last is simply $\prod_{m=0}^{\infty} * / \sim$. The \sim relation identifies two i -simplices in $(X(U)_i)^m$ and $(X(U)_i)^{m'}$ if one is obtained from the other by means of an order-preserving injection of the indexing set. But all such injections induce the identity map $* = (X(U)_i)^m \rightarrow (X(U)_i)^{m'} = *$, so it follows that $J(X)(U)_i$ collapses to a singleton set, as required. \square

6 The stable isomorphism

6.1 The diagonal

Let $\sigma: J(X)_+ \rightarrow D(X)$ denote the stable isomorphism in $\text{ho } \mathbf{Spt}(\mathbf{Sm}_k)$ of Corollary 5.7. The category $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$ is equipped with localization functors to $\text{ho}_{\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$, $\text{ho}_p \mathbf{Spt}(\mathbf{Sm}_k)$ and $\text{ho}_{p, \mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$, this being the upshot of Section 3. We will denote the images of objects and morphisms under the various localization functors by the same notation as we use in the category $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$, and in order to avoid confusion we will specify the category in which we are working.

Let $\Delta^q: J(X) \rightarrow J(X)^q$ denote the order- q diagonal of $J(X)$.

The study of Δ^q has been extensively carried out by Kuhn in [25; 26] and elsewhere. We describe some of that study here. Fix a natural number i and a finite sequence $A = (a_1, a_2, \dots, a_q)$ of natural numbers. Following [26], we define a *partial A cover* to be a collection of subsets $T = (T_1, \dots, T_q)^1$ of $\{1, \dots, i\}$ with the property that $|T_t| = a_t$. We will say this partial cover is of *type (a_1, \dots, a_q)* . The partial cover T is called a *cover* if $\bigcup_{t=1}^q T_t = \{1, \dots, i\}$. In [26, Section 2], a natural stable map of spaces is constructed, associated to a cover T ,

$$\Psi_T: D_i X \rightarrow D_{a_1}(X) \wedge D_{a_2}(X) \wedge \dots \wedge D_{a_q}(X).$$

¹This is denoted by S in [26], but we have reserved S for the symmetric group because we use Σ for the reduced suspension functor

We should mention that the definition of $D_i X$ in [26] is not the same as ours, merely homotopy equivalent. Since he is considering a vastly more general context than we are, Kuhn defines $D_i X$ in terms of a coefficient system, whereas we consider, in the language of coefficient systems, only the little-intervals operad. The little-intervals operad has $\mathcal{C}(n)$ homotopy equivalent to the space of n distinct marked points on a unit interval, which is homotopy equivalent to the symmetric group S_n . This equivalence allows us our much more elementary definition of $D_i X$, homotopy equivalent to that of Kuhn. Specifically, where we have $X^{\wedge i}$, Kuhn takes the space of i points on an interval, each labeled with an element of X , and then contracts the subspace where any label is the basepoint of X to a point.

Using our construction of $D_i X$, it is possible to give an elementary description of $\Psi_T: D_i X \rightarrow D_{a_1}(X) \wedge D_{a_2}(X) \wedge \cdots \wedge D_{a_q}(X)$. The set $\{1, \dots, i\}$ is covered by subsets A_1, \dots, A_q of cardinalities a_1, \dots, a_q . To each we associate the order-preserving inclusion $\alpha_{T,t}: \{1, \dots, a_t\} \rightarrow \{1, \dots, i\}$, and then define a map

$$\psi_T: X^{\times i} \rightarrow X^{\times a_1} \times \cdots \times X^{\times a_t}, \quad (x_1, \dots, x_i) \mapsto \prod_{t=1}^q \left(\prod_{k=1}^{a_t} x_{\alpha_{T,t}(k)} \right),$$

where the products are taken in the usual order.

For any $\sigma \in S_i$, we may act on $X^{\times i}$ by permuting the factors, and thereby obtain

$$\psi_T \circ \sigma: X^{\times i} \rightarrow X^{\times a_1} \times \cdots \times X^{\times a_t}.$$

We act on the codomain by the unique element of $S_{a_1} \times \cdots \times S_{a_t}$ so that for each $j \in \{1, \dots, t\}$, the composite map $X^{\times i} \rightarrow X^{\times a_1} \times \cdots \times X^{\times a_t} \rightarrow X^{\times a_j}$ is order-preserving. Write $[\sigma]\psi_T$ for the resulting map

$$[\sigma]\psi_T: X^{\times i} \rightarrow X^{\times a_1} \times \cdots \times X^{\times a_t}.$$

It depends only on the coset of σ modulo S_T , the stabilizer of the cover T .

If T is a cover, and if the basepoint of X appears in the i -tuple on the left, then it will appear on the right, and so the maps $[\sigma]\psi_T$ descend to maps $X^{\wedge i} = D_i X \rightarrow D_{a_1} X \wedge \cdots \wedge D_{a_q} X$, which we also write as $[\sigma]\psi_T$ in an abuse of notation. Finally, the stable map

$$\Psi_T: D_i X \rightarrow D_{a_1}(X) \wedge D_{a_2}(X) \wedge \cdots \wedge D_{a_q}(X)$$

is produced as a sum $\sum_{\sigma \in S_i/S_T} [\sigma]\psi_T$.

For example, given the cover $T = (\{1, 2\}, \{3, 4\})$, the function Ψ_T can be represented as the sum of the functions sending a reduced word $x_1x_2x_3x_4$ to each of the following:

$$\begin{aligned} &((x_1x_2), (x_3x_4)), \quad ((x_1x_3), (x_2x_4)), \quad ((x_1x_4), (x_2x_3)), \\ &((x_2x_3), (x_1x_4)), \quad ((x_2x_4), (x_1x_3)), \quad ((x_3x_4), (x_1x_2)). \end{aligned}$$

The construction of the map Ψ_T is natural, and therefore it induces a natural stable map of simplicial presheaves, which we also denote by Ψ_T .

Two covers are said to be *equivalent* if they lie in the same orbit of the symmetric group action on $\{1, \dots, i\}$. The proof in [25, Lemma 2.6] that equivalent covers induce homotopic maps is entirely formal and carries over to the setting of simplicial presheaves.

Definition 6.1 Let $\Delta_{i,(a_1,a_2,\dots,a_q)}^q(X)$ denote the composition in $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$,

$$\begin{aligned} D_i X \rightarrow D(X) \xrightarrow{\sigma^{-1}} J(X)_+ \xrightarrow{\Delta^q} (J(X)^q)_+ \xrightarrow{\wedge^q \sigma} \wedge^q D(X) \\ \rightarrow D_{a_1}(X) \wedge D_{a_2}(X) \wedge \dots \wedge D_{a_q}(X). \end{aligned}$$

The following result is essentially [26, Theorem 2.4], but we assert it in the category of simplicial presheaves.

Proposition 6.2 *There is an equality in the stable homotopy category of presheaves,*

$$\Delta_{i,(a_1,a_2,\dots,a_q)}^q = \sum_T \Psi_T: D_i X \rightarrow D_{a_1}(X) \wedge D_{a_2}(X) \wedge \dots \wedge D_{a_q}(X),$$

where the sum runs over equivalence classes of covers T of type (a_1, \dots, a_q) .

Proof We outline the argument of Kuhn, noting that at all points all constructions are natural, and therefore the proof carries over essentially without modification to the context of simplicial presheaves.

The problem is reduced from that of DX to $D(X_+)$, that is, to the case of a space where the basepoint is disjoint. The device that allows this reduction is that the map in the stable homotopy category $D(X_+) \rightarrow DX$ is a split epimorphism. This follows from the fact that in general the map $X \times Y \rightarrow X \wedge Y$ is split after application of a single suspension functor, and applies equally well in the case of presheaves as in the case of spaces.

Kuhn observes immediately after Lemma 3.1 of [26] that $C(X_+)_+$ is naturally homeomorphic to $D(X_+)$, denoting the natural homeomorphism by s' . We work with a different model for $D(X_+)$ and the $C(X_+)$ of [26] corresponds to our $J(X_+)$, and the homeomorphism s' in our context amounts to the observation that the reduced free monoid on a simplicial set K_+ is naturally isomorphic to $S^0 \vee K_+ \vee (K \times K)_+ \vee \dots$. The homeomorphism $(s'): J(X_+)_+ \rightarrow D(X_+)$ does not stabilize to give the Hilton–Milnor splitting $\sigma: J(X_+)_+ \rightarrow D(X_+)$ of Corollary 5.7. For this reason, Definition 3.2 of [26] defines $\Theta: D(X_+) \rightarrow D(X_+)$ to be the stable map $\sigma \circ (s')^{-1}$, and defines $\Theta_{n,m}: D_n(X_+) \rightarrow D_m(X_+)$ to be the stable maps given by the $(n, m)^{\text{th}}$ component of Θ , ie $\Theta_{n,m}$ is a stable map $D_n(X_+) \rightarrow D_m(X_+)$. The proofs of Propositions 3.3 (the proof of which is the proof of [25, Proposition 4.5]), 3.4 and Lemmas 3.5 and 3.6 of [26] are entirely formal, relying on the behavior of the stable transfer maps associated to subgroups of the symmetric group. They apply without modification to simplicial presheaves, and therefore suffice to establish our proposition. \square

Proposition 6.3 (Kuhn [26]) *If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ equipped with a co- H -structure, then $\Delta_{i,(a_1,\dots,a_q)}^q(X) \cong *$ in $\mathbf{hoNis} \mathbf{Spt}(\mathbf{Sm}_k)$, unless $i = \sum_{j=1}^q a_j$.*

Proof If $i < \sum_{j=1}^q a_j$, then this is true even without the co- H -structure. In Proposition 6.2, there are no covers in this case, and therefore the sum is empty.

If X is equipped with a co- H -structure, then the diagonal $X \rightarrow X \times X$ factors up to homotopy through a co- H -map $X \rightarrow X \vee X$. In this case, the diagonal map $X \rightarrow X \wedge X$ is nullhomotopic. The argument in the first part of [26, Appendix A] applies directly, amounting to the claim that if $i > \sum_{j=1}^q a_j$, then each term in the sum of Proposition 6.2 is also null. \square

6.2 Combinatorics

Our first aim is to prove that a specific stable map, namely $\Delta_{2m+r,(2m,r)}^2: X^{\wedge 2m+r} \rightarrow X^{\wedge 2m} \wedge X^{\wedge r}$, is as an equivalence, at least after localizing at 2, when $X \simeq S^{2n+q\alpha}$ is a motivic sphere and when m is a natural number and r is either 0 or 1. This is a corollary of Propositions 6.2 and 6.3, but the proof requires us to consider some elementary combinatorics. The same combinatorics will prove useful later, when we turn to studying the James–Hopf map.

For a positive integer a and an m -tuple of positive integers (a_1, a_2, \dots, a_m) , let $\binom{a}{a_1, a_2, \dots, a_m}$ denote the set of functions $\sigma: \{1, 2, \dots, a\} \rightarrow \{1, 2, \dots, m\}$ such that $\sigma^{-1}(i)$ has cardinality a_i . Note that $\binom{a}{a_1, a_2, \dots, a_m}$ is nonempty if and only if $a = \sum a_i$.

Given an element $\sigma \in \binom{a}{a_1, a_2, \dots, a_m}$ and a natural number $i \leq m$, write $\sigma^{-1}(i)$ as $\{\sigma^{-1}(i)_1, \sigma^{-1}(i)_2, \dots, \sigma^{-1}(i)_{a_i}\}$ in such a way that $\sigma^{-1}(i)_j < \sigma^{-1}(i)_{j+1}$ for all $j \leq a_i - 1$. Define $\tilde{\sigma}$ to be the permutation on a letters sending $(1, 2, \dots, a)$ to $(\sigma^{-1}(1)_1, \sigma^{-1}(1)_2, \dots, \sigma^{-1}(1)_{a_1}, \sigma^{-1}(2)_1, \dots, \sigma^{-1}(2)_{a_2}, \dots, \sigma^{-1}(m)_{a_m})$.

For instance, if σ is the element of $\binom{5}{1, 2, 2}$ given by sending $2 \mapsto 1$ and $1, 5 \mapsto 2$ and $3, 4 \mapsto 3$, then $\tilde{\sigma}$ is the permutation taking $(1, 2, 3, 4, 5)$ to $(2, 1, 5, 3, 4)$.

Suppose X is an objects of $\mathbf{sPre}(\mathbf{Sm}_k)_*$. For $\sigma \in \binom{a}{a_1, a_2, \dots, a_m}$, define $e(\sigma): X^{\wedge a} \rightarrow X^{\wedge a}$ to be the map induced by $\tilde{\sigma}$, and define $\text{sign}(\sigma)$ to be the number of pairs $r < k$ in $\{1, 2, \dots, a\}$ such that $\tilde{\sigma}(r) > \tilde{\sigma}(k)$. In the example given, $\text{sign}(\sigma)$ is the cardinality of $\{(1, 2), (3, 4), (3, 5)\}$, ie 3.

Remark 6.4 Suppose X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$. Let i and a_1, a_2, \dots, a_q be positive integers such that $i = \sum a_k$. Then

$$\Delta_{i, (a_1, a_2, \dots, a_q)}^q(X) = \sum_{\sigma \in \binom{i}{a_1, a_2, \dots, a_q}} e(\sigma)$$

in $\text{hoNis } \mathbf{Spt}(\mathbf{Sm}_k)$ — this is merely a restatement of a special case of Proposition 6.2 in different notation.

In the case where $X = S^{n+q\alpha}$, Remark 4.5 says that $e(\sigma)$ is $e_{n,q}^{\text{sign } \sigma}$, where $e_{n,q} = (-1)^{n+q} \langle -1 \rangle^q$ in $\text{GW}(k)$. Using the remark above then gives:

Corollary 6.5 Suppose $X = S^{n+q\alpha}$. Let i, a_1, a_2, \dots, a_w be nonnegative integers such that $i = \sum a_k$. Then

$$\Delta_{i, (a_1, a_2, \dots, a_w)}^w(X) = \sum_{\sigma \in \binom{i}{a_1, a_2, \dots, a_w}} e_{n,q}^{\text{sign } \sigma}$$

in $\text{hoNis } \mathbf{Spt}(\mathbf{Sm}_k)$.

We will have occasion to use an involution

$$\gamma: \binom{a_1 + a_2 + \dots + a_m}{a_1, a_2, \dots, a_m} \rightarrow \binom{a_1 + a_2 + \dots + a_m}{a_1, a_2, \dots, a_m}$$

which is defined as follows:

Take $\sigma \in \binom{a_1 + a_2 + \dots + a_m}{a_1, a_2, \dots, a_m}$. There are two possibilities:

- (i) $\sigma(2i - 1) = \sigma(2i)$ for all applicable i . In this case, we say $\gamma(\sigma) = \sigma$, so σ is fixed under the involution. We write $F_\gamma(a_1 + a_2 + \dots + a_m; a_1, a_2, \dots, a_m)$ for the set of fixed points, or F_γ when the coefficients are clear from the context.
- (ii) Otherwise, there exists a least integer i such that $\sigma(2i - 1) \neq \sigma(2i)$. We then let $\gamma(\sigma)$ be the function that agrees with σ except that $\gamma(\sigma)(2i - 1) = \sigma(2i)$ and $\gamma(\sigma)(2i) = \sigma(2i - 1)$.

If σ is not a fixed point of γ , then $\text{sign}(\sigma) + \text{sign}(\gamma(\sigma)) \equiv 1 \pmod{2}$, so that the number of elements in $\binom{a_1+a_2+\dots+a_m}{a_1, a_2, \dots, a_m}$ of even sign is given by the formula

$$(14) \quad \begin{aligned} \text{number of elements of even sign} &= \frac{1}{2} \left(\left| \binom{a_1+a_2+\dots+a_m}{a_1, a_2, \dots, a_m} \right| - |F_\gamma| \right) + |F_\gamma| \\ &= \frac{1}{2} \left(\left| \binom{a_1+a_2+\dots+a_m}{a_1, a_2, \dots, a_m} \right| + |F_\gamma| \right). \end{aligned}$$

We can often find ways to calculate $\left| \binom{a_1+a_2+\dots+a_m}{a_1, a_2, \dots, a_m} \right|$ and $|F_\gamma|$.

The cardinality of $\binom{x+y}{x, y}$ is the binomial coefficient $(x + y)! / (x! y!)$. Let $\lfloor \frac{1}{2}x \rfloor$ denote the greatest integer less than or equal to $\frac{1}{2}x$.

Proposition 6.6 *Among the elements of $\binom{x+y}{x, y}$, the number having even sign is*

$$\frac{1}{2} \left(\left| \binom{x+y}{x, y} \right| + \left| \binom{\lfloor \frac{1}{2}(x+y) \rfloor}{\lfloor \frac{1}{2}x \rfloor, \lfloor \frac{1}{2}y \rfloor} \right| \right).$$

By virtue of our definitions, the second summand is 0 in the case where x and y are both odd.

Proof We rely on the involution γ and (14). Since $\left| \binom{x+y}{x, y} \right|$ is known, it remains to calculate $|F_\gamma|$.

There are several cases to consider:

- (i) If x and y are both odd, then every number in $\{1, \dots, x + y\}$ forms part of a pair $(2i - 1, 2i)$, and there must be at least one pair for which $\sigma(2i - 1) \neq \sigma(2i)$, since $\sigma^{-1}(1)$ is odd. There are therefore no fixed points of the involution.
- (ii) If x and y are both even, then every number in $\{1, \dots, x + y\}$ forms part of a pair $(2i - 1, 2i)$. In order for σ to be fixed by γ , it must be the case that $\sigma(2i - 1) = \sigma(2i)$ for all i . Defining $\tau \in \binom{(x+y)/2}{x/2, y/2}$ by the formula $\tau(i) = \sigma(2i)$, we see that there is a bijection between F_γ and $\binom{(x+y)/2}{x/2, y/2}$.

- (iii) If one of x and y is even and the other odd — say for specificity that x is even and y odd — then σ is fixed under γ if $\sigma(x + y) = 2$ and $\sigma(2i - 1) = \sigma(2i)$ for all $i \leq \lfloor \frac{1}{2}(x + y) \rfloor$. Similarly to the previous case, there is a bijection in this case between F_γ and $\binom{(x+y-1)/2}{x/2, (y-1)/2}$.

Since $|F_\gamma|$ agrees in all cases with

$$\left| \binom{\lfloor \frac{1}{2}(x + y) \rfloor}{\lfloor \frac{1}{2}x \rfloor, \lfloor \frac{1}{2}y \rfloor} \right|,$$

the proposition is proved. □

Proposition 6.7 *Let $X \simeq S^{n+q\alpha}$ be a motivic sphere, suppose m is a nonnegative integer and $r \in \{0, 1\}$. Let P be a set of primes, and let m be a natural number such that $m + 1 + me_{n,q}$ is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_P$; then the map $\Delta_{2m+r, (2m,r)}^2: X^{\wedge 2m+r} \rightarrow X^{\wedge 2m} \wedge X^{\wedge r}$ is an isomorphism in $\text{ho}_{P, \mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$.*

In particular, $\Delta_{2m+r, (2m,r)}^2$ is an isomorphism in $\text{ho}_{2, \mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$.

Proof We have calculated $\Delta_{2m+r, (2m,r)}^2$ in Corollary 6.5 and Proposition 6.6. If $r = 0$, we find $\Delta_{2m, (2m,0)}^2 = 1$ in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_{(2)}$.

When $r = 1$, we find $\Delta_{2m+1, (2m,1)}^2 = m + 1 + me_{n,q}$, from which the first claim follows immediately.

The element $m + 1 + me_{n,q}$ is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_{(2)}$ by Corollary 4.7. □

The same calculations, referring to Corollary 4.8, show the following:

Proposition 6.8 *Let $X \simeq S^{n+q\alpha}$ be a motivic sphere. Assume one of the following two conditions holds:*

- (i) n is odd and q is even.
- (ii) $n + q$ is odd and the ground field k is not formally real.

Suppose m is a nonnegative integer and $r \in \{0, 1\}$. The diagonal map

$$\Delta_{2m+r, (2m,r)}^2: X^{\wedge 2m+r} \rightarrow X^{\wedge 2m} \wedge X^{\wedge r}$$

is an isomorphism in $\text{ho}_{\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$.

Definition 6.9 We impose a total order on the elements of $\binom{a_1 + \dots + a_m}{a_1, a_2, \dots, a_m}$ by declaring $\sigma < \sigma'$ if $\sigma(j) = \sigma'(j)$ for all $j \leq k$ and $\sigma(k) < \sigma'(k)$.

Following [25], define a regular (r, s) -set of size m to be a set, $\{S_1, \dots, S_s\}$, of subsets of $\{1, \dots, m\}$ satisfying

- (i) $|S_i| = r$ for all i ,
- (ii) $\bigcup_{i=1}^s S_i = \{1, \dots, m\}$.

Let $L(r, s, m)$ denote the set of all regular (r, s) sets of size m . This goes by the name $B(B(m, s), r)$ in [25], and the discussion that follows here is a much reduced version of the discussion to be found there. In particular, we concentrate on the case where $r = 2$ and $m = 2s$.

There is a cover $\phi: \binom{2s}{2, \dots, 2} \rightarrow L(2, s, 2s)$, the source being the set of ordered partitions of $\{1, \dots, 2s\}$ into s disjoint subsets of cardinality 2, and the latter the set of unordered partitions. There is an S_s -action on functions $\sigma: \{1, \dots, 2s\} \rightarrow \{1, \dots, s\}$ induced from the action on the target, and the orbits of this action are in bijective correspondence with $L(2, s, 2s)$. For any $\lambda \in L(2, s, 2s)$, define $\text{sign}(\lambda)$ to be $\text{sign}(\sigma)$, where σ is the least element, in the order of Definition 6.9, of $\binom{2s}{2, \dots, 2}$ that maps to λ under ϕ .

Write $E(2, s)$ and $O(2, s)$ for the number of elements in $L(2, s, 2s)$ having even and odd sign, respectively. Trivially, $E(2, 1) = 1$ and $O(2, 1) = 0$.

Proposition 6.10 *The quantities $E(2, s)$ and $O(2, s)$ satisfy $E(2, s) = O(2, s) + 1$.*

Proof An element $\lambda \in E(2, s)$ is a partition of $\{1, \dots, 2s\}$ into s disjoint subsets $\{S_1, \dots, S_s\}$. One orders these subsets in ascending order of their least members. The quantity $\text{sign}(\lambda)$ is the number of pairs of numbers $j_1 < j_2$ such that $j_1 \in S_{\ell_1}$ and $j_2 \in S_{\ell_2}$ with $S_{\ell_1} > S_{\ell_2}$. We can set up an involution $\bar{\gamma}$ on $L(2, s, 2s)$ by observing that γ descends from $\binom{2s}{2, \dots, 2}$.

Explicitly, if $\lambda = \{S_1, \dots, S_s\} \in L(2, s, 2s)$ is not the partition

$$\lambda_0 = \{\{1, 2\}, \{3, 4\}, \dots, \{2s-1, 2s\}\}$$

then there is a least pair of integers $(2i-1, 2i)$ such that $2i-1$ and $2i$ lie in different sets S_i and S_j with $i \neq j$. Let $\bar{\gamma}(\lambda)$ be the partition obtained from λ by interchanging $2i-1$ and $2i$. The exceptional partition, λ_0 , is the unique fixed point of $\bar{\gamma}$.

Observe that if $\lambda \neq \lambda_0$, then $\text{sign}(\lambda) + \text{sign}(\bar{\gamma}(\lambda)) \equiv 1 \pmod{2}$. Since $\text{sign}(\lambda_0) = 0$ is even, it follows that $E(2, s) - 1 = O(2, s)$, as asserted. \square

Remark 6.11 The cardinality of $\binom{2s}{2, \dots, 2}$ is $(2s)!/2^s$ and the cardinality of $L(2, s, 2s)$ is $(2s)!/(s!2^s)$. Explicitly, therefore,

$$E(2, s) = \frac{1}{2} \left(\frac{(2s)!}{s!2^s} - 1 \right) + 1 = \frac{(2s)!}{s!2^{s+1}} + \frac{1}{2}$$

and

$$O(2, s) = \frac{1}{2} \left(\frac{(2s)!}{s!2^s} - 1 \right) = \frac{(2s)!}{s!2^{s+1}} - \frac{1}{2}.$$

The quantity $E(2, s) + O(2, s) = (2s)!/(s!2^s)$ is the product of the first s odd integers, $(2s-1)(2s-3) \cdots (5)(3)(1)$. The fact that this is a unit in $\mathbb{Z}_{(2)}$ appears in the classical study of j_2 .

6.3 Decomposing the second James–Hopf map

Definition 6.12 Let

$$a_{i,s}^2: D_i(X) \rightarrow D_s(X^{\wedge 2})$$

denote the composition in $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$,

$$D_i X \rightarrow D(X) \xrightarrow{\sigma^{-1}} J(X)_+ \xrightarrow{j_2} J(X^{\wedge 2})_+ \xrightarrow{\sigma} D(X^{\wedge 2}) \rightarrow D_s(X^{\wedge 2}).$$

For example, we have

$$(15) \quad a_{2,1}^2 = \text{id}_{D_2(X)}$$

by the commutative diagram

$$\begin{array}{ccc} J(X) & \xrightarrow{j_2} & J(X^{\wedge 2}) \\ \uparrow & & \uparrow \\ J_2(X) & \longrightarrow & X^{\wedge 2} \end{array}$$

where the lower horizontal map is the composite

$$J_2(X) \rightarrow J_2(X)/J_1(X) = D_2(X) \cong X^{\wedge 2} \xrightarrow{\text{id}} X^{\wedge 2} \cong D_1(X^{\wedge 2}).$$

Proposition 6.13 Let $X \simeq S^{n+q\alpha}$ be a motivic sphere with $n \geq 1$. Let $i \geq 2$ be an integer. The maps in $\text{ho}_{\text{Nis}} \mathbf{Spt}(\mathbf{Sm}_k)$,

$$D_i(X) \xrightarrow{\quad \xrightarrow{\quad} \quad \xrightarrow{\quad} \quad \xrightarrow{\quad} \quad} J(X) \xrightarrow{j_2} J(X^{\wedge 2}) \xrightarrow{\quad} D_s(X^{\wedge 2}),$$

$\xrightarrow{\quad a_{i,s}^2 \quad}$

agree in $\text{ho}_{\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$ with

$$a_{i,s}^2 = \begin{cases} E(2,s) + O(2,s)e_{n,q} & \text{if } i = 2s, \\ * & \text{otherwise.} \end{cases}$$

Proof The case where $i \neq 2s$ follows from Corollary 6.3(1) and (3) of [25]. As with the results of [26], the arguments here all yield natural homotopies of maps, and therefore the results carry over from the case of spaces to the case of simplicial presheaves.

When $i = 2s$, then, by Theorem 6.2 of [25], the class $a_{i,s}^2$ is equal to the sum of the classes of permutations of $X^{\wedge 2s}$ associated to regular $(2,s)$ sets of size $2s$. Of these, $E(2,s)$ are even permutations, and therefore equivalent to the identity, and $O(2,s)$ are odd, and therefore equivalent to the single interchange $e_{n,q}$. \square

Corollary 6.14 *The map $a_{2s,s}^2$ is an isomorphism in $\text{ho}_{2,\mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k))$.*

Proof The map in question is $E(2,s) + O(2,s)e_{n,r}$. Since $E(2,s) = O(2,s) + 1$ by Proposition 6.10, it follows from Corollary 4.7 that it is a unit in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_{(2)}$. \square

Similarly, we have the following corollary:

Corollary 6.15 *If $X = S^{n+q\alpha}$ in $\text{GW}(k)$ is a motivic sphere and one of*

- (i) n is even and q is odd, or
- (ii) $n + q$ is odd and the field k is not formally real,

then $a_{2s,s}^2$ represents an isomorphism in $\text{ho}_{\mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k))$.

Proof This follows from Proposition 6.10 and Corollary 4.8. \square

6.4 The stable weak equivalence

Let X be of the form $S^{n+q\alpha}$ for $n \geq 1$ and $q \geq 2$. Let $e_{n,q}$ be the class $(-1)^{n+q}(-1)^q$ in $\text{GW}(k)$.

Fix a set of primes, P . All objects and maps in this section belong to the category $\text{ho}_{P,\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$, unless otherwise stated. If P is the set of all primes, then $\text{ho}_{P,\mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k)) = \text{ho}_{\mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k))$. This case and the case $P = \{(2)\}$ are the two cases that are applied in subsequent sections of this paper.

Write $b_+ : J(X)_+ \cong \bigvee_{i=0}^\infty X^{\wedge i} \rightarrow X_+$ for the projection map.

We will need the following construction again in the sequel, so we present it here for later reference.

Construction 6.16 Suppose given an unstable map $j : J \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and a stable map $b : J \rightarrow X$ in $\mathbf{ho}_{P, \mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k))_*$, that is to say a homotopy class of maps $b : \Sigma^\infty J \rightarrow \Sigma^\infty X$. We produce a stable map $(j \wedge b_+) \circ \Delta_+ : J_+ \rightarrow (X \times Y)_+$ as follows.

We may extend the maps $b : J \rightarrow X$ and $j : J \rightarrow Y$ to maps $b_+ : J_+ \rightarrow X_+$ in $\mathbf{ho}_{P, \mathbb{A}^1}(\mathbf{Spt}(\mathbf{Sm}_k))_*$ and $j_+ : J_+ \rightarrow Y_+$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$. Then we take the smash product of these two maps. For convenience, we note that $j \vee \text{id}$ is a map in the unstable homotopy category, so this may be carried out in an elementary way without recourse to a smash product of spectra. This gives a map $b_+ \wedge j_+ : J_+ \wedge J_+ \rightarrow X_+ \wedge Y_+$. But the source and target of this map may be identified with $(J \times J)_+$ and $(X \times Y)_+$, respectively. Then precomposing with the diagonal map $J \rightarrow (J \times J)_+$ gives the result.

By means of the above, we construct a map in stable homotopy, denoted by c :

$$J(X)_+ \xrightarrow{\Delta} (J(X) \times J(X))_+ \xrightarrow{\quad} (J(X) \times X)_+ \xrightarrow{\quad} (J(X^{\wedge 2}) \times X)_+.$$

c

Here Δ is the image in $\mathbf{ho}_{2, \mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$ of the diagonal map

$$J(X)_+ \rightarrow (J(X) \times J(X))_+$$

in $\mathbf{sPre}(\mathbf{Sm}_k)_*$, and j is the James–Hopf map $j : J(X) \rightarrow J(X^{\wedge 2})$ in $\mathbf{sPre}(\mathbf{Sm}_k)$. Since j_+ is a map in $\mathbf{sPre}(\mathbf{Sm}_k)_*$, we can form the product map $(J(X) \times X)_+ \rightarrow J(X^{\wedge 2} \times X)_+$ in $\mathbf{ho}_{\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)(\mathbf{Sm}_k)$ by means of the action of $\mathbf{sPre}(\mathbf{Sm}_k)_*$ on $\mathbf{Spt}(\mathbf{Sm}_k)$.

Both spaces $J(X)_+$ and $(J(X^{\wedge 2}) \times X)_+$ are isomorphic in the homotopy category $\mathbf{ho}_{2, \mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$ to the spectrum $\bigvee_{i=0}^\infty X^{\wedge i}$. To see the latter, decompose

$$(16) \quad \begin{aligned} (J(X^{\wedge 2}) \times X)_+ &\cong S^0 \vee J(X^{\wedge 2}) \vee X \vee (J(X^{\wedge 2}) \wedge X) \\ &\cong S^0 \vee (\bigvee_{i=1}^\infty X^{\wedge 2i}) \vee X \vee (\bigvee_{i=1}^\infty X^{\wedge 2i+1}). \end{aligned}$$

Use the above to fix a standard isomorphism $\bigvee_{i=0}^\infty X^{\wedge i} \cong (J(X^{\wedge 2}) \times X)_+$ in the homotopy category $\text{ho}_{\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)$, an isomorphism $J(X)_+ \cong \bigvee_{i=0}^\infty X^{\wedge i}$ already having been fixed in the form of the stable map s .

Proposition 6.17 *Fix a sphere $X = S^{n+q\alpha}$. If elements $m + 1 + me_{n,q}$, where m is an integer, are units in $\text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}_p$, then the map c is a weak equivalence.*

Proof Consider the ring

$$R = \text{End}_{\text{ho}_{2,\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)}(\bigvee_{i=0}^\infty X^{\wedge i}).$$

We wish to show that $(j_+ \times b_+) \circ \Delta$ is a unit of this ring.

We may write

$$R = \prod_{i=0}^\infty \text{Hom}_{\text{ho}_{2,\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)}(X^{\wedge i}, \bigvee_{l=0}^\infty X^{\wedge l})$$

and

$$\bigvee_{l=0}^\infty X^{\wedge l} \simeq \bigvee_{l=0}^i X^{\wedge l} \vee \bigvee_{l=i+1}^\infty X^{\wedge l}.$$

It follows from the Hurewicz theorem that $[X^{\wedge i}, \bigvee_{l=i+1}^\infty X^{\wedge l}] = 0$, and so

$$\text{Hom}_{\text{ho}_{2,\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)}(X^{\wedge i}, \bigvee_{l=0}^\infty X^{\wedge l}) = \bigoplus_{l=0}^i \text{Hom}_{\text{ho}_{2,\mathbb{A}^1} \mathbf{Spt}(\mathbf{Sm}_k)}(X^{\wedge i}, X^{\wedge l}),$$

so that $R = \prod_{i=0}^\infty \bigoplus_{l=0}^i \pi_{in+iq\alpha}(S^{ln+lq\alpha})$. We may represent elements of R as infinite, upper-triangular matrices $(d_{i,l})$ such that $d_{i,l} \in \pi_{in+iq\alpha}(S^{ln+lq\alpha})$ by decreeing $d_{i,l} = 0$ whenever $i < l$. It follows from the usual algebra of matrix multiplication that an element of R is a unit if and only if the terms $d_{i,i} \in \pi_{in+iq\alpha}(S^{in+iq\alpha})$ are units for all i .

The invertibility of c in R may be deduced from the classes $d_{i,i}$ appearing in the diagram

$$(17) \quad \begin{array}{ccccccc} & & & c & & & \\ & & & \curvearrowright & & & \\ J(X)_+ & \xrightarrow{\Delta} & (J(X) \times J(X))_+ & \longrightarrow & (J(X) \times X)_+ & \longrightarrow & (J(X^{\wedge 2}) \times X)_+ \\ & \uparrow & & & & & \downarrow \\ X^{\wedge i} & & & d_{i,i} & & & X^{\wedge i} \end{array}$$

where the unmarked arrows are inclusion and projection maps.

We can factor $d_{i,i}$ in diagram (17) as

$$(18) \quad \begin{array}{ccccccc} & & & & c & & \\ & & & & \curvearrowright & & \\ J(X)_+ & \xrightarrow{\Delta} & (J(X) \times J(X))_+ & \longrightarrow & (J(X) \times X)_+ & \longrightarrow & (J(X^{\wedge 2}) \times X)_+ \\ \uparrow & & \uparrow & & & & \downarrow \\ X^{\wedge i} & \xrightarrow{f} & \bigvee_{n=0}^i X^{\wedge i-n} \wedge X^{\wedge n} & \longrightarrow & & \longrightarrow & X^{\wedge i} \\ & & & & d_{i,i} & & \end{array}$$

where f is the wedge sum of maps $\Delta_{i,(i-n,n)}^2: X^{\wedge i} \rightarrow X^{\wedge i-n} \wedge X^{\wedge n}$ as n varies. This factorization follows from Proposition 6.3.

We can further factorize $d_{i,i}$ because the map $\Sigma^\infty(J(X) \times J(X))_+ \rightarrow \Sigma^\infty(J(X) \times X)_+$ is the identity on the first and projection on the second factor:

$$(19) \quad \begin{array}{ccccccc} & & & & c & & \\ & & & & \curvearrowright & & \\ J(X)_+ & \xrightarrow{\Delta} & (J(X) \times J(X))_+ & \longrightarrow & (J(X) \times X)_+ & \longrightarrow & (J(X^{\wedge 2}) \times X)_+ \\ \uparrow & & \uparrow & & \uparrow & & \downarrow \\ X^{\wedge i} & \xrightarrow{f} & \bigvee_{n=0}^i X^{\wedge i-n} \wedge X^{\wedge n} & \longrightarrow & (X^{\wedge i} \wedge X^{\wedge 0}) \vee (X^{\wedge i-1} \wedge X) & \longrightarrow & X^{\wedge i} \\ & & & & d_{i,i} & & \end{array}$$

Write $i = 2m + s$ where $s \in \{0, 1\}$. By use of Proposition 6.13, we deduce that the bottom row can be further factored as

$$(20) \quad \begin{array}{ccccccc} & & & & d_{i,i} & & \\ & & & & \curvearrowright & & \\ X^{\wedge i} & \xrightarrow{f} & \bigvee_{n=0}^i X^{\wedge i-n} \wedge X^{\wedge n} & \longrightarrow & (X^{\wedge i} \wedge X^{\wedge 0}) \vee (X^{\wedge i-1} \wedge X) & \longrightarrow & X^{\wedge i} \\ & & & & \downarrow & \nearrow a_{2m,m}^2 \wedge \text{id} & \\ & & & & X^{\wedge 2m} \wedge X^{\wedge s} & & \end{array}$$

It follows that $d_{i,i}$ factors as $(a_{2m,m}^2 \wedge \text{id}) \circ \Delta_{i,(2m,s)}^2$, and since both these maps are isomorphisms by virtue of Propositions 6.7 and 6.13, so too is $d_{i,i}$, and therefore so too is $c = (j_+ \times b_+) \circ \Delta$. □

Remark 6.18 The hypothesis of the proposition that elements of the form

$$(m + 1) + me_{n,q} \in \text{GW}(k) \otimes_{\mathbb{Z}} \mathbb{Z}P$$

be units holds in particular in the following cases:

- (i) The ring \mathbb{Z}_P is $\mathbb{Z}_{(2)}$ or \mathbb{Q} . In this case, the hypothesis holds by Corollary 4.7.
- (ii) The integer n is odd and the integer q is even. In this case, $e_{n,q} = -1$, and the hypothesis holds by Corollary 4.8.
- (iii) The integer $n + q$ is odd, and the field k is not formally real. Again, the hypothesis holds in this case by Corollary 4.8.

Remark 6.19 If X is an object in $\mathbf{sPre}(\mathbf{Sm}_k)_*$, there is an action of the symmetric group S_n on $X^{\wedge n}$. In the case where X is a motivic sphere, this action factors through the sign representation of S_n . The fact that c is a 2–local weak equivalence depends on this fact, as we can see in the following example.

Example 6.20 Let X be the simplicial set $X = S^2 \vee S^2$. The map $S_n \rightarrow [X^{\wedge n}, X^{\wedge n}]$ is injective because the action of S^n on $H^{2n}(X^{\wedge n}, \mathbb{Q}) \cong H^2(X, \mathbb{Q})^{\otimes n} \cong \mathbb{Q}^{2^n}$ contains a direct sum of two copies of the permutation representation of S_n over \mathbb{Q} as summands. These two copies can be described as follows. The wedge product $X^{\wedge n}$ is the direct sum of copies of S^{2n} indexed by n –tuples of elements of $\{1, 2\}$. The n –tuples which have a single 1 and the rest 2’s form one of the summands, and the other is obtained by switching the roles of 1 and 2.

The objects $J(X)_+$ and $J(X^{\wedge 2})_+$ split stably as $\bigvee_{i=0}^{\infty} X^{\wedge i}$ and $\bigvee_{i=0}^{\infty} X^{\wedge 2i}$, respectively. The second James–Hopf map

$$j_2: \bigvee_{i=0}^{\infty} X^{\wedge i} \rightarrow \bigvee_{i=0}^{\infty} X^{\wedge 2i}$$

restricts to a map $a_{4,2}^2: X^{\wedge 4} \rightarrow X^{\wedge 4}$. The paper [25] calculates this map explicitly as the sum of permutations

$$a_{4,2}^2 = \sum_{\sigma \in \binom{4}{2,2}} e(\sigma).$$

Note that $\binom{4}{2,2}$ is in bijection with $\{((1, 2)(3, 4)), ((1, 3)(2, 4)), ((1, 4)(2, 3))\}$ under the bijection sending $((a, b), (c, d))$ to the map sending a and b to 1 and sending c and d to 2. Using cycle notation for permutations, and representing the identity by e , this sum is

$$a_{4,2}^2 = e + (2\ 3) + (2\ 4\ 3).$$

The induced map on the singular cohomology $H^8(X^{\wedge 4}, \mathbb{Q}) \cong \mathbb{Q}^{16}$ is not of full rank, since $e + (2\ 3) + (2\ 4\ 3)$ is not an isomorphism on the permutation representation,




Figure 1: The first four rows, nine columns of the E_2 -page of the Serre spectral sequence for $H^*(\cdot, \mathbb{Q})$ associated to $\text{hofib}(j_2) \rightarrow J(X) \xrightarrow{j_2} J(X^{\wedge 2})$.

namely on either of the submodules mentioned above $a_{4,2}^2$ acts by the matrix

$$\begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix},$$

which has determinant 0.

The map induced by j_2 on rational cohomology $H^8(J(X^{\wedge 2}), \mathbb{Q}) \rightarrow H^8(J(X), \mathbb{Q})$ is not an isomorphism in this case, and is in particular not injective, and so the analogue of Proposition 6.17 fails in this case, even \mathbb{Q} -locally.

Moreover, associated to the fiber sequence

$$\text{hofib}(j_2) \rightarrow J(X) \xrightarrow{j_2} J(X^{\wedge 2})$$

there is a Serre spectral sequence for rational cohomology, part of which is shown in Figure 1. We have shown that the edge map $H^8(J(X^{\wedge 2}), \mathbb{Q}) \rightarrow H^8(J(X), \mathbb{Q})$ is not injective. Since the edge map is not injective, it is not the case that the spectral sequence collapses at the E_2 -page, and since $H^*(J(X^{\wedge 2}), \mathbb{Q})$ is concentrated in even degrees, it follows that $H^*(\text{hofib}(j_2), \mathbb{Q})$ is not also concentrated in even degrees. In particular, $\text{hofib}(j_2)$ does not have the same rational cohomology as $X = S^2 \vee S^2$, showing that even the \mathbb{Q} -local version of the EHP sequence does not hold for a general space X .

7 Fiber of the James–Hopf map

In Section 7.3, we will have two fiber sequences $F \rightarrow E \rightarrow Y$ and $X \rightarrow X \times Y \rightarrow Y$ with the same base Y and a stable weak equivalence between the total spaces E and $X \times Y$, which is compatible with the map to the base. We will show that in fact the fibers are stably weakly equivalent as well (Proposition 7.18). For this, it is natural to

ask for a Serre spectral sequence, as the Serre spectral sequence gives a good way to measure the size of the total space of a fibration in terms of the size of the base and the fiber. Since the base spaces of the fibrations f and p are the same and their total spaces are the same size, a Serre spectral sequence would give us a tool with which to attempt to “cancel off the base space” and conclude that the fibers have the same size. The purpose of the first part of this section is to show that enough of these ideas remain available in \mathbb{A}^1 –homotopy theory. In Section 7.1, we construct a spectral sequence to substitute for the Serre spectral sequence. We develop needed properties in Section 7.2, and in Section 7.3, we show that the desired cancellation is possible.

7.1 A spectral sequence

Let \mathfrak{a} be a left Bousfield localization of the global model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$. There is an associated stable model structure on the category of S^1 –spectra, $\mathbf{Spt}(\mathbf{Sm}_k)$. See Section 2.4. Let $\mathcal{H}_i: \mathbf{Spt}(\mathbf{Sm}_k) \rightarrow \mathbf{Sh}_{\text{Nis}}$ be an \mathfrak{a} –corepresentable functor, given by a spectrum E , such that $\mathcal{H}_i(F)$ is the Nisnevich sheaf associated to the presheaf

$$U \mapsto [\Sigma^\infty S^i \wedge E \wedge \Sigma^\infty U_+, F]_{\mathfrak{a},s}.$$

We write $\mathcal{H}_i(X)$ for $\mathcal{H}_i(\Sigma^\infty X)$ when X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

Since left Bousfield localization does not change which maps are cofibrations, the notions of global cofibration, Nisnevich local cofibration, \mathbb{A}^1 –cofibration and \mathfrak{a} –cofibration for $\mathbf{sPre}(\mathbf{Sm}_k)$ are the same. For $X_1 \rightarrow X_2$ a cofibration with respect to these model structures, the cofiber C is the pushout

$$\begin{array}{ccc} C = \operatorname{colim} X_1 & \longrightarrow & X_2 \\ & & \downarrow \\ & & * \end{array}$$

A sequence is said to be a cofiber (resp. fiber) sequence up to homotopy if the sequence is isomorphic in the homotopy category to a cofiber (resp. fiber) sequence.

Proposition 7.1 *The homology theory \mathcal{H}_* has the following properties:*

- (i) \mathcal{H}_i takes \mathfrak{a} weak equivalences to isomorphisms.
- (ii) Given a cofibration $X_1 \rightarrow X_2$ with cofiber C in \mathfrak{a} , there is a natural long exact sequence of sheaves of abelian groups

$$\cdots \rightarrow \mathcal{H}_i X_1 \rightarrow \mathcal{H}_i X_2 \rightarrow \mathcal{H}_i C \rightarrow \mathcal{H}_{i-1} X_1 \rightarrow \cdots .$$

- (iii) As a special case of (ii), we see that $\mathcal{H}_i(\Sigma X) \simeq \mathcal{H}_{i-1}(X)$ for $X \in \mathbf{sPre}(\mathbf{Sm}_k)_*$.

Assumption 7.2 We assume that \mathcal{H}_* satisfies two further axioms:

- (i) **Boundedness** $\mathcal{H}_i(X) = 0$ for $i < 0$ for all objects X in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.
- (ii) **Compactness** $\operatorname{colim} \mathcal{H}_i(X_j) = \mathcal{H}_i(\operatorname{colim} X_j)$ for all filtered diagrams $\{X_j\}$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.

These axioms are satisfied by $\mathcal{H}_i = \pi_{i+j\alpha}^{s, \mathbb{A}^1}$ and $\mathcal{H}_i = \pi_{i+j\alpha}^{s, P, \mathbb{A}^1}$: we take α to be the \mathbb{A}^1 injective structure or the P - \mathbb{A}^1 injective structure, respectively. In each case, $E = \Sigma^\infty \mathbb{G}_m^{\wedge j}$. The boundedness axiom follows from Lemma 2.20, the compactness from Proposition 2.18 or Proposition 3.27.

For $f: X \rightarrow Y$ a global fibration, we construct a spectral sequence $E_{ij}^r \Rightarrow \mathcal{H}_{i+j} X$.

This spectral sequence for $\alpha = \mathbb{A}^1$ or its P -localizations will have the property that it relates α -homotopy-invariant information about the total space with α -homotopy-invariant information about the fiber and more delicate information about the base.

Let $\Delta_{\leq n}$ be the full subcategory of Δ on the objects $\{0, 1, \dots, n\}$. The pointed n -skeleton $\operatorname{sk}_n: \mathbf{sSet}_* \rightarrow \mathbf{sSet}_*$ can be defined as the composite of the n -truncation functor $\mathbf{sSet}_* \rightarrow \operatorname{Fun}(\Delta_{\leq n}^{\text{op}}, \mathbf{Set}_*)$ with its left adjoint. Given a simplicial presheaf X , define $\operatorname{sk}_n X \in \mathbf{sPre}(\mathbf{Sm}_k)$ by $U \mapsto \operatorname{sk}_n X(U)$. For $n < 0$, the definition gives $\operatorname{sk}_n X = *$.

Definition 7.3 Let $f: X \rightarrow Y$ be a fibration in the global model structure between pointed simplicial presheaves. We define the spectral sequence

$$(E_{ij}^r, d^r: E_{i+r, j}^r \rightarrow E_{i, j+r-1}^r):$$

The cofibrations

$$\operatorname{sk}_0 Y \rightarrow \operatorname{sk}_1 Y \rightarrow \dots \rightarrow \operatorname{sk}_n Y \rightarrow \dots Y$$

pull back to cofibrations

$$\operatorname{sk}_0 Y \times_Y X \rightarrow \operatorname{sk}_1 Y \times_Y X \rightarrow \dots \rightarrow \operatorname{sk}_n Y \times_Y X \rightarrow \dots Y \times_Y X = X.$$

The cofiber sequences

$$\operatorname{sk}_{n-1} Y \times_Y X \rightarrow \operatorname{sk}_n Y \times_Y X \rightarrow C_n$$

for $n \geq 0$ give rise to the long exact sequences of Proposition 7.1(ii), which form an exact couple:

$$\begin{array}{ccc}
 \bigoplus_{i,n=0}^{\infty} \mathcal{H}_i(\text{sk}_n Y \times_Y X) & \xrightarrow{\quad\quad\quad} & \bigoplus_{i,n=0}^{\infty} \mathcal{H}_i(\text{sk}_n Y \times_Y X) \\
 & \swarrow \text{dotted} & \nwarrow \\
 & \bigoplus_{i,n=0}^{\infty} \mathcal{H}_i C_n &
 \end{array}$$

This exact couple gives rise to the spectral sequence $E_{i,j}^r$ with $E_{i,j}^1 = \mathcal{H}_{i+j} C_i$.

We now relate C_i to the fiber of f . Assume for simplicity that Y is reduced in the sense that $Y_0 = *$, and let F denote the fiber of f over Y_0 .

For $U \in \mathbf{Sm}_k$, let $L_n Y(U) \in \mathbf{sSet}_*$ denote the n^{th} latching object, defined as $L_n Y(U) = (\text{sk}_{n-1} Y(U))_n$, and let $N_n Y(U)$ be the set of nondegenerate n -simplices of $Y(U)$, defined as $N_n Y(U) = Y_n(U) - L_n Y(U)$.

Despite the fact that $N_n Y(-)$ does not necessarily define a presheaf,

$$\bigvee_{N_n Y(F_+ \wedge (\Delta^n / \partial \Delta^n))} = U \mapsto \bigvee_{y \in N_n Y(U)} (F(U)_+ \wedge (\Delta^n / \partial \Delta^n))$$

is a presheaf because it could equally well be written as

$$\frac{\bigvee_{y \in Y_n(U)} (F(U)_+ \wedge (\Delta^n / \partial \Delta^n))}{\bigvee_{y \in L_n Y(U)} (F(U)_+ \wedge (\Delta^n / \partial \Delta^n))},$$

and both Y_n and $L_n Y$ are presheaves. This presheaf is weakly equivalent in the global model structure to the cofiber C_n , as shown by the following lemma:

Lemma 7.4 *There is a global weak equivalence $C_n \simeq \bigvee_{N_n Y(F_+ \wedge (\Delta^n / \partial \Delta^n))}$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$.*

Proof Let $\partial \Delta^n$ denote the boundary of Δ^n . By [11, Chapter VII, Proposition 1.7, page 355], there is a pushout

$$\begin{array}{ccc}
 (Y_n \times \partial \Delta^n) \cup_{L_n Y \times \partial \Delta^n} (L_n Y \times \Delta^n) & \longrightarrow & \text{sk}_{n-1} Y \\
 \downarrow & & \downarrow \\
 Y_n \times \Delta^n & \longrightarrow & \text{sk}_n Y
 \end{array}
 \tag{21}$$

The pullback of a pushout square of simplicial sets is a pushout square because small colimits are pullback stable in the topos of simplicial sets. Since limits and colimits in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ commute with taking the sections above $U \in \mathbf{Sm}$, it follows that the pullback of a pushout square in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is also a pushout square. Thus, applying

the functor $(-)\times_Y X$ to (21) produces a pushout square, which then produces a global weak equivalence between C_n and the cofiber of

$$(22) ((Y_n \times \partial\Delta^n) \times_Y X) \cup_{(L_n Y \times \partial\Delta^n) \times_Y X} ((L_n Y \times \Delta^n) \times_Y X) \rightarrow (Y_n \times \Delta^n) \times_Y X.$$

The composition $Y_n \times F \rightarrow F \rightarrow X$ of the projection with the inclusion determines a map $Y_n \times F \rightarrow X$. The product of the identify map on $Y_n \times F$ with inclusion of the 0th vertex into Δ^n gives a map $Y_n \times F \rightarrow Y_n \times F \times \Delta^n$. The canonical map $Y_n \times \Delta^n \rightarrow Y$ factors as $Y_n \times \Delta^n \rightarrow \text{sk}_n Y \rightarrow Y$, and precomposing with the projection $Y_n \times F \times \Delta^n \rightarrow Y_n \times \Delta^n$, we obtain maps $Y_n \times F \times \Delta^n \rightarrow Y_n \times \Delta^n \rightarrow \text{sk}_n Y \rightarrow Y$. These maps fit into the commutative diagram

$$\begin{array}{ccccccc} Y_n \times F & \longrightarrow & (Y_n \times \Delta^n) \times_Y X & \longrightarrow & \text{sk}_n Y \times_Y X & \longrightarrow & X \\ \downarrow & \nearrow \text{dotted} & \downarrow & & \downarrow & & \downarrow \\ Y_n \times F \times \Delta^n & \longrightarrow & Y_n \times \Delta^n & \longrightarrow & \text{sk}_n Y & \longrightarrow & Y \end{array}$$

formed by the solid arrows.

Since $Y_n \times F \rightarrow Y_n \times F \times \Delta^n$ is a global trivial cofibration, this commutative diagram extends to include a map denoted by the dotted arrow, by the lifting property of global trivial cofibrations and global fibrations. The dotted arrow is a global weak equivalence because it is a map of global fibrations over $Y_n \times \Delta$ such that the induced map on the fibers is a global weak equivalence

$$(23) \begin{array}{ccc} & (Y_n \times \Delta^n) \times_Y X & \\ & \nearrow \text{dotted} & \downarrow \\ Y_n \times F \times \Delta^n & \longrightarrow & Y_n \times \Delta^n \end{array}$$

Pulling back the diagram (23) by a map $A \rightarrow Y_n \times \Delta^n$ produces a map of global fibrations over A such that the induced map on fibers is a global weak equivalence, and it follows that the pullback of the dotted arrow remains a global weak equivalence as well. We apply this to the canonical maps from $A = Y_n \times \partial\Delta^n$, $A = L_n Y \times \partial\Delta^n$ and $A = L_n Y \times \Delta^n$. Since the union in the domain of the map (22) is a homotopy pushout as well as a pushout, we obtain a diagram

$$\begin{array}{ccc} ((Y_n \times \partial\Delta^n) \times_Y X) \cup_{(L_n Y \times \partial\Delta^n) \times_Y X} ((L_n Y \times \Delta^n) \times_Y X) & \longrightarrow & (Y_n \times \Delta^n) \times_Y X \\ \downarrow & & \downarrow \\ (Y_n \times \partial\Delta^n \times F) \cup_{L_n Y \times \partial\Delta^n \times F} (L_n Y \times \Delta^n \times F) & \longrightarrow & Y_n \times \Delta^n \times F \end{array}$$

where the vertical arrows are weak equivalences. Since the horizontal arrows are monomorphisms and therefore global cofibrations, we obtain an induced weak equivalence between the cofibers, proving the lemma. \square

It is convenient to introduce notation for the presheaf $\bigvee_{N_i Y} F_+$, so we do that now.

Definition 7.5 For a global fibration $f: X \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $Y_0 = *$, define K_i in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ to be

$$K_i = \bigvee_{N_i Y} F_+,$$

where F is the fiber of f .

Lemma 7.4 shows that there is a global weak equivalence $C_i \simeq S^i \wedge K_i$.

Proposition 7.6 Let $\{E_{ij}^r, d_{ij}^r\}$ denote the spectral sequence of Definition 7.3 associated to a global fibration $f: X \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $Y_0 = *$.

- (i) There is a canonical isomorphism $E_{i,j}^1 \cong \mathcal{H}_j K_i$
- (ii) $E_{ij}^r = 0$ for i or j less than 0.
- (iii) This spectral sequence converges to the values of the functors \mathcal{H}_* on X

$$(E_{ij}^r, d^r: E_{i+r,j}^r \rightarrow E_{i,j+r-1}^r) \Rightarrow \mathcal{H}_{i+j} X.$$

Proof We prove the claims in order.

- (i) Since there is a global weak equivalence $C_i \simeq S^i \wedge K_i$ (Lemma 7.4), it follows that

$$E_{i,j}^1 = \mathcal{H}_{i+j} C_i \cong \mathcal{H}_{i+j} S^i \wedge K_i \cong \mathcal{H}_j K_i$$

by Proposition 7.1(iii).

- (ii) The claim is immediate for $i < 0$. We show that $E_{ij}^1 = 0$ for $j < 0$, which is sufficient because E_{ij}^r is a subquotient of $E_{i,j}^1$. By (i), $E_{i,j}^1 \cong \mathcal{H}_j K_i$. For $j < 0$, we have $\mathcal{H}_j K_i = 0$ by Assumption 7.2(i).
- (iii) The convergence follows from (ii). Since $\text{colim}_n \text{sk}_n Y = Y$ and finite limits commute with filtered colimits, it follows that $\text{colim}_n \text{sk}_n Y \times_Y X \cong X$. Since \mathcal{H}_{j+i} preserves filtered colimits (Assumption 7.2(ii)), $\text{colim}_n \mathcal{H}_{j+i}(\text{sk}_n Y \times_Y X) \cong \mathcal{H}_{j+i}(X)$, proving the claim. \square

We can summarize this subsection as follows: For a map $f: X \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ such that $Y_0 = *$, we have a spectral sequence

$$(E_{ij}^r, d^r: E_{i+r,j}^r \rightarrow E_{i,j+r-1}^r) \Rightarrow \mathcal{H}_{i+j} X$$

satisfying the properties of Proposition 7.6, where $F = \text{hofib}_{\text{global}} f$ is the homotopy fiber in the global model structure of f , ie factor f as the composition $f = \iota \circ f'$, with $\iota: X \rightarrow Z$ a global trivial cofibration and $f': Z \rightarrow Y$ a global fibration. Let F be the fiber of f' over the basepoint. The spectral sequence is that of Definition 7.3 applied to f' .

Remark 7.7 The construction of this spectral sequence as well as those of its properties given in this section only require the lifting properties of global fibrations. The subtler lifting properties for α -fibrations have not been exploited.

7.2 A functoriality property

We will use a functoriality property of the spectral sequence constructed in Section 7.1 with respect to a particular sort of stable map in the homotopy category.

Recall that α is a left Bousfield localization of the global injective model structure.

Proposition 7.8 *The model structure α on $\mathbf{sPre}(\mathbf{Sm}_k)$ satisfies the following properties:*

- (i) *All monomorphisms in $\mathbf{sPre}(\mathbf{Sm}_k)$ are α cofibrations, and in particular, objects of $\mathbf{sPre}(\mathbf{Sm}_k)$ are α cofibrant.*
- (ii) *α is compatible with the tensor, cotensor and simplicial enrichment in the sense that this structure makes $\mathbf{sPre}(\mathbf{Sm}_k)$ into a simplicial model category [35, Definition 11.4.4].*
- (iii) *$\Sigma: \mathbf{sPre}(\mathbf{Sm}_k) \rightarrow \mathbf{sPre}(\mathbf{Sm}_k)$ takes α weak equivalences to α weak equivalences.*
- (iv) *There is a left Quillen functor $\Sigma^\infty: \mathbf{sPre}(\mathbf{Sm}_k)_+ \rightarrow \mathbf{Spt}(\mathbf{Sm}_k)$, where $\mathbf{Spt}(\mathbf{Sm}_k)$ is endowed with a stable model structure, which we also call α , in an abuse of notation.*

Proof Property (i) is immediate: α is a left Bousfield localization of the global injective model structure, and the same property holds there. Property (ii) follows from [12, Theorem 4.1.1(4)] because the global injective model structure is left proper, simplicial, and cellular (see [14]). Property (iii) is a special case of (ii). Property (iv) is Proposition 2.14. □

We furthermore make the following assumption:

Assumption 7.9 If X is an object of $\mathbf{sPre}(\mathbf{Sm}_k)$, then $X \times \cdot$ preserves weak equivalences.

To construct the EHP fiber sequence, we will set \mathfrak{a} to be the P -localized \mathbb{A}^1 model structure for P a set of primes. By Proposition 2.1 and Corollary 2.2, this choice is valid.

We will employ the following construction, which is a version of Construction 6.16 that applies to maps b that exist after one suspension, rather than simply stably; we use the notation $\Sigma(-)_+$ to mean $\Sigma((-)_+)$.

Construction 7.10 Suppose given a map $j: J \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and a map $b: \Sigma J \rightarrow \Sigma X$ in $\text{ho}_{\mathfrak{a}} \mathbf{sPre}(\mathbf{Sm}_k)_*$. We produce a map $(b \wedge j_+) \circ \Sigma \Delta_+ : \Sigma J_+ \rightarrow \Sigma(X \times Y)_+$ as follows:

We may extend $b: \Sigma J \rightarrow \Sigma X$ to a map in $\text{ho}_{\mathfrak{a}} \mathbf{sPre}(\mathbf{Sm}_k)_*$

$$b: \Sigma J_+ \rightarrow \Sigma X_+$$

since $\Sigma J_+ \simeq \Sigma J \vee S^1$, and similarly for X . We take the smash product of $b: \Sigma J_+ \rightarrow \Sigma X_+$ and $j_+: J_+ \rightarrow Y_+$, obtaining a map $\Sigma J_+ \wedge J_+ \rightarrow \Sigma X_+ \wedge Y_+$. The left-hand side is identified with $\Sigma(J \times J)_+$ and the right with $\Sigma(X \times Y)_+$. Then we precompose with the diagonal map $\Sigma \Delta_+ : \Sigma J_+ \rightarrow \Sigma(J \times J)_+$ to obtain a map in $\text{ho}_{\mathfrak{a}} \mathbf{sPre}(\mathbf{Sm}_k)_*$,

$$(24) \quad (b \wedge j_+) \circ \Sigma \Delta_+ : \Sigma J_+ \rightarrow \Sigma(X \times Y)_+.$$

This construction is functorial in the map j as follows. Suppose given a commutative square

$$\begin{array}{ccc} J' & \xrightarrow{p} & J \\ \downarrow j' & & \downarrow j \\ Y' & \xrightarrow{q} & Y \end{array}$$

Then there is a map $b \circ \Sigma p: \Sigma J' \rightarrow \Sigma X$, and the evident square

$$\begin{array}{ccc} \Sigma J' & \xrightarrow{((b \circ \Sigma p) \wedge j_+) \circ \Sigma \Delta_+} & \Sigma(X \times Y)_+ \\ \downarrow & & \downarrow \\ \Sigma J & \xrightarrow{(b \wedge j_+) \circ \Sigma \Delta_+} & \Sigma(X \times Y)_+ \end{array}$$

is commutative.

We remark that the map $(b \wedge j_+) \circ \Sigma \Delta_+ : \Sigma J \rightarrow \Sigma(X \times Y)$ constructed here agrees in the stable category with Construction 6.16.

We assume the following setup: a map $j : J \rightarrow Y$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ and a map $b : \Sigma J \rightarrow \Sigma X$ in $\mathbf{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$ with X fibrant. We replace $j : J \rightarrow Y$ by a fibration by means of a canonical factorization $J \xrightarrow{\iota} E \xrightarrow{f} Y$, where ι is a trivial cofibration and f is a fibration. There is a well-defined map in the homotopy category,

$$(25) \quad b_E = b \circ \Sigma \iota^{-1} : \Sigma E \rightarrow \Sigma X.$$

Let p denote the projection $X \times Y \rightarrow Y$. Associated to each of the fibrations

$$f : E \rightarrow Y, \quad p : X \times Y \rightarrow Y$$

there are spectral sequences as in Definition 7.3. We denote these by E'_{ij} and $(E')^r_{ij}$, respectively. Let

$$\mathfrak{b} : \Sigma E \rightarrow \Sigma(X \times Y)$$

denote the map formed from $(b_E \wedge f_+) \circ \Sigma \Delta_+ : \Sigma E_+ \rightarrow \Sigma(X \times Y)_+$ (Construction 7.10) by precomposing with a map $\Sigma E \rightarrow \Sigma E \vee S^1 \simeq \Sigma E_+$ and postcomposing with $\Sigma(X \times Y)_+ \rightarrow \Sigma(X \times Y)$. Since \mathcal{H}_* is a stable theory, we obtain a map

$$\mathcal{H}_i \Sigma^{-1} \mathfrak{b} : \mathcal{H}_*(J) \rightarrow \mathcal{H}_*(X \times Y).$$

Lemma 7.11 *With notation as above, \mathfrak{b} induces a map of spectral sequences $E'_{ij} \rightarrow (E')^r_{ij}$. The induced map $E^\infty_{ij} \rightarrow (E')^\infty_{ij}$, which by Proposition 7.6(iii) is a map from the associated graded of a filtration of $\mathcal{H}_{i+j}(E)$ to the associated graded of a filtration of $\mathcal{H}_{i+j}(X \times Y)$, is compatible with*

$$\mathcal{H}_{i+j} \Sigma^{-1} \mathfrak{b} : \mathcal{H}_{i+j}(E) \rightarrow \mathcal{H}_{i+j}(X \times Y).$$

Proof In order to construct the map of spectral sequences, we start with the skeletal filtration $\text{sk}_\bullet Y$ of Y . For all n , there are maps $f_n : \text{sk}_n Y \times_Y E \rightarrow \text{sk}_n Y$. We remark in passing that the fiber product $\text{sk}_n Y \times_Y E$ is a homotopy fiber product by virtue of $f : E \rightarrow Y$ being a fibration. Moreover, there are composite maps $b_n : \Sigma(\text{sk}_n Y \times_Y E) \rightarrow \Sigma E \rightarrow \Sigma X$, where the first map is induced by the inclusion of the skeleton and the second is b .

By means of Construction 7.10, we obtain maps $\Sigma(\text{sk}_n Y \times_Y E)_+ \rightarrow \Sigma(\text{sk}_n Y \times X)_+$ in the homotopy category, and by functoriality, these maps are compatible in that the

diagram

$$(26) \quad \begin{array}{ccccc} \Sigma(\mathrm{sk}_{n-1} Y \times_Y E)_+ & \longrightarrow & \Sigma(\mathrm{sk}_n Y \times_Y E)_+ & \longrightarrow & \Sigma E_+ \\ \downarrow & & \downarrow & & \downarrow \\ \Sigma(\mathrm{sk}_{n-1} Y \times X)_+ & \longrightarrow & \Sigma(\mathrm{sk}_n Y \times X)_+ & \longrightarrow & \Sigma(Y \times X)_+ \end{array}$$

commutes.

The commutative diagram (26) induces a commutative diagram

$$(27) \quad \begin{array}{ccccc} \Sigma(\mathrm{sk}_{n-1} Y \times_Y E) & \longrightarrow & \Sigma(\mathrm{sk}_n Y \times_Y E) & \longrightarrow & \Sigma C_n \\ \downarrow & & \downarrow & & \downarrow \\ \Sigma(\mathrm{sk}_{n-1} Y \times X) & \longrightarrow & \Sigma(\mathrm{sk}_n Y \times X) & \longrightarrow & \Sigma C'_n \end{array}$$

in $\mathrm{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$, where the horizontal rows are cofiber sequences and suspensions of cofiber sequences. Applying $\mathcal{H}_* \Sigma^{-1}$ to the entire diagram then defines a morphism of long exact sequences, and thus a morphism of exact couples, and therefore a morphism of spectral sequences. The compatibility with the induced map on E^∞ -pages follows from applying $\mathcal{H}_* \Sigma^{-1}$ to (26). \square

We can compute the map $E_{ij}^1 \rightarrow (E')_{ij}^1$ of E^1 -pages of the map of spectral sequences of Lemma 7.11. Suppose again that $Y \in \mathbf{sPre}(\mathbf{Sm}_k)_*$ is such that $Y_0 = *$. Let $a: F \rightarrow E$ denote the canonical map of simplicial presheaves given by the definition $F = \mathrm{hofib}_{\mathrm{global}} f = * \times_Y E \rightarrow E$. Composing Σa_+ with b_E yields a map

$$b_E \circ \Sigma a_+: \Sigma F_+ \rightarrow \Sigma X_+$$

in $\mathrm{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$.

Recall that $K_i \in \mathbf{sPre}(\mathbf{Sm}_k)_*$ is defined by $K_i(U) = \bigvee_{N_i Y(U)} (F(U)_+)$. The analogous definition for the global fibration p is then $K'_i = \bigvee_{N_i Y(U)} (X(U)_+)$. We claim that the map $b_E \circ \Sigma a_+: \Sigma F_+ \rightarrow \Sigma X_+$ in $\mathrm{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$ defines a map $\Sigma K_i \rightarrow \Sigma K'_i$ in $\mathrm{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$. This claim is established by the following lemma.

Lemma 7.12 *Suppose I is a presheaf of sets on \mathbf{Sm}_k .*

- (i) *If $g: A \rightarrow B$ is an α weak equivalence in pointed spaces $\mathbf{sPre}(\mathbf{Sm}_k)_*(\mathbf{Sm}_k)$, then $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ is an α weak equivalence.*
- (ii) *If $g: A \rightarrow B$ is an α weak equivalence in spectra $\mathbf{Spt}(\mathbf{Sm}_k)$, then $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ is an α weak equivalence.*

Furthermore, suppose that $Y \in \mathbf{sPre}(\mathbf{Sm}_k)_*$ is pointed. Then we may replace the presheaf I of sets on \mathbf{Sm}_k by $U \mapsto N_n Y(U)$.

Proof (i) $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ is canonically identified with $I_+ \wedge g: I_+ \wedge A \rightarrow I_+ \wedge B$, so $\bigvee_I g$ is a weak equivalence by Corollary 2.2.

(ii) We again have a canonical identification of the map $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ with $I_+ \wedge g: I_+ \wedge A \rightarrow I_+ \wedge B$. For any pointed simplicial sheaf \mathcal{X} , the functor

$$\mathbf{Spt}(\mathbf{Sm}_k) \rightarrow \mathbf{Spt}(\mathbf{Sm}_k), \quad E \mapsto E \wedge \mathcal{X},$$

preserves stable \mathbb{A}^1 weak equivalences, as in [32, Section 4, page 27].

Furthermore, note that we have a canonical bijection $Y_n(U)/L_n Y(U) \cong N_n Y(U) \amalg *$ and that $U \mapsto Y_n(U)/L_n Y(U)$ is a presheaf of (pointed) sets, which we will denote by I . There is a monomorphism $A \rightarrow \bigvee_I A$, and similarly for B . The map $U \mapsto \bigvee_{N_n Y(U)} g(U)$ is the map induced by taking cofibers in the diagram

$$\begin{array}{ccc} A & \longrightarrow & \bigvee_I A \\ g \downarrow & & \bigvee_I g \downarrow \\ B & \longrightarrow & \bigvee_I B \end{array}$$

Since the top and bottom horizontal arrows are cofibrations, the cofibers of these maps are also homotopy cofibers, so it follows that $\bigvee_{N_n Y(U)} g$ is an a weak equivalence. \square

It follows that $b_E \circ \Sigma a_+$ in $\mathbf{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$ induces a map

$$\bigvee_{N_n Y}(b_E \circ \Sigma a_+): \Sigma K_i \rightarrow \Sigma K'_i$$

in $\mathbf{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$. As before, we may apply $\mathcal{H}_j \Sigma^{-1}$ to any map in $\mathbf{ho}_a \mathbf{sPre}(\mathbf{Sm}_k)_*$. Applying $\mathcal{H}_j \Sigma^{-1}$ to $\bigvee_{N_n Y}(b \circ \Sigma a_+)$ gives our identification of the map of E^1 -pages in the map of spectral sequences $E^r_{ij} \rightarrow (E')^r_{ij}$ in Lemma 7.11.

Lemma 7.13 *Suppose that $Y_0 = *$. The identification of $E^1_{i,j}$ with $\mathcal{H}_j(K_i)$ in Proposition 7.6(i) is functorial with respect to the map*

$$E^r_{ij} \rightarrow (E')^r_{ij}$$

in the sense that the induced map for $r = 1$ is $\mathcal{H}_j \Sigma^{-1}(\bigvee_{N_n Y}(b_E \circ \Sigma a_+)) : \mathcal{H}_j(K_i) \rightarrow \mathcal{H}_j(K'_i)$.

Proof By construction, the lemma is equivalent to the claim that the map $\mathcal{H}_{n+j}C_n \rightarrow \mathcal{H}_{n+j}C'_n$ given by applying $\mathcal{H}_{n+j}\Sigma^{-1}$ to the commutative diagram (27) is identified with $\mathcal{H}_{n+j}\Sigma^{-1}$ applied to

$$\bigvee_{\text{id}_{N_n Y}}(b_E \circ \Sigma a_+) \wedge S^n: \Sigma \vee \bigvee_{N_n Y} F_+ \wedge S^n \rightarrow \Sigma \vee \bigvee_{N_n Y} X_+ \wedge S^n$$

via the equivalences given in Lemma 7.4.

The equivalences of Lemma 7.4 are constructed by choosing a trivialization

$$\begin{array}{ccc} Y_n \times F & \longrightarrow & E \\ \downarrow & \nearrow \phi & \downarrow f \\ Y_n \times F \times \Delta^n & \longrightarrow & Y \end{array}$$

and any two are homotopic, as either is equivalent in the homotopy category to the composition

$$Y_n \times F \times \Delta^n \xrightarrow{\cong} Y_n \times F \rightarrow E.$$

The product $X \times Y$ admits a canonical trivialization

$$\begin{array}{ccc} Y_n \times X & \longrightarrow & X \times Y \\ \downarrow & \nearrow & \downarrow f \\ Y_n \times X \times \Delta^n & \longrightarrow & Y \end{array}$$

Applying $\Sigma(-)_+$ to $Y_n \times F \rightarrow E$ and composing with the map

$$(b_E \wedge f_+) \circ \Sigma \Delta_+: \Sigma E_+ \rightarrow \Sigma(X \times Y)_+$$

from Construction 7.10 produces a map

$$\Sigma(Y_n \times F)_+ \rightarrow \Sigma E_+ \rightarrow \Sigma(Y \times X)_+,$$

which factors through the inclusion

$$\Sigma(Y_n \times X)_+ \rightarrow \Sigma(Y \times X)_+.$$

The resulting map

$$\Sigma(Y_n \times F)_+ \rightarrow \Sigma E_+ \rightarrow \Sigma(Y_n \times X)_+$$

is the smash product of the identity $1_{Y_n}: Y_n \rightarrow Y_n$ on Y_n and $b_E \circ \Sigma a_+$. It follows that the map $\Sigma C_n \rightarrow \Sigma C'_n$ of (27) is identified with $\bigvee_{\text{id}_{N_n Y}}(b_E \circ \Sigma a_+) \wedge 1_{S^n}$, proving the lemma. □

The description of the map of E^1 -pages given in Lemma 7.13 means that understanding the construction taking a map $g: A \rightarrow B$ in $\text{ho}_\alpha \mathbf{sPre}(\mathbf{Sm}_k)_*$ to $\bigvee_{N_n Y} g: \bigvee_{N_n Y} A \rightarrow \bigvee_{N_n Y} B$ in $\text{ho}_\alpha \mathbf{sPre}(\mathbf{Sm}_k)_*$ implies understanding the map of E^1 -pages. The next lemma and corollary give some understanding of this construction $g \mapsto \bigvee_{N_n Y} g$ in the case where (α, \mathcal{H}_*) is $(\mathbb{A}^1, \pi_*^{s, \mathbb{A}^1})$ or $(P - \mathbb{A}^1, \pi_*^{s, P, \mathbb{A}^1})$.

To show Lemma 7.15, it is useful to note the following:

Remark 7.14 If X is a simplicially n -connected spectra in the sense that $X \in \mathbf{Spt}(\mathbf{Sm}_k)$ satisfies $\pi_i^s X = 0$ for $i \leq n$, then Morel’s stable connectivity theorem [32] implies that $\pi_i^{s, \mathbb{A}^1} X = \pi_i^s L_{\mathbb{A}^1} X = 0$ for $i \leq n$. Because $\pi_i^{s, P, \mathbb{A}^1}$ is naturally isomorphic to $\mathbb{Z}_P \otimes \pi_i^{s, \mathbb{A}^1}$ by Proposition 3.25, it also follows that $\pi_i^{s, P, \mathbb{A}^1}(X) = 0$ for $i \leq n$, whence $L_P L_{\mathbb{A}^1} X$ is simplicially n -connected.

Lemma 7.15 Let I be a presheaf of sets on \mathbf{Sm}_k and $g: A \rightarrow B$ be a map in $\mathbf{sPre}(\mathbf{Sm}_k)_*$. Let (α, \mathcal{H}_i) be either $(\mathbb{A}^1, \pi_i^{s, \mathbb{A}^1})$ or $(P - \mathbb{A}^1, \pi_i^{s, P, \mathbb{A}^1})$.

- (i) If g induces an isomorphism on \mathcal{H}_i for $i < n - 1$ and surjection for $i = n - 1$, then $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ induces an isomorphism on \mathcal{H}_i for $i < n - 1$ and a surjection for $i = n - 1$.
- (ii) If g induces an isomorphism on \mathcal{H} for $i < n$ and there is a map $h: B \rightarrow A$ in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$ such that $g \circ h = \text{id}_B$ in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$, then $\bigvee_I g: \bigvee_I A \rightarrow \bigvee_I B$ induces an isomorphism on \mathcal{H}_i for $i < n$ and a surjection for $i = n$.

Proof The second statement follows from the first, and the first is proven as follows.

Let C denote the \mathbb{A}^1 -homotopy cofiber of g (ie factor g as $g_1 \circ g_2$ with $g_2: A \rightarrow B'$ a cofibration and $g_1: B' \rightarrow B$ an \mathbb{A}^1 fibration and an \mathbb{A}^1 weak equivalence and let C be the cofiber of the cofibration g_2). Since g_2 is also a $P - \mathbb{A}^1$ cofibration and $B' \rightarrow B$ is also a $P - \mathbb{A}^1$ weak equivalence, we have a long exact sequence

$$(28) \quad \dots \rightarrow \mathcal{H}_{i+1} C \rightarrow \mathcal{H}_i A \xrightarrow{\mathcal{H}_i(g)} \mathcal{H}_i B \rightarrow \mathcal{H}_i C \rightarrow \dots$$

Since $\mathcal{H}_i(g)$ is an isomorphism for $i < n - 1$, we have that $\text{Image}(\mathcal{H}_{i+1} C \rightarrow \mathcal{H}_i A) = 0$. Thus, $\mathcal{H}_{i+1} C = \ker(\mathcal{H}_{i+1} C \rightarrow \mathcal{H}_i A)$. Since (28) is exact, $\ker(\mathcal{H}_{i+1} C \rightarrow \mathcal{H}_i A) = \text{Image}(\mathcal{H}_{i+1} B \rightarrow \mathcal{H}_{i+1} C)$.

Since $\mathcal{H}_i(g)$ is a surjection for $i < n$, we have that $\mathcal{H}_i B \rightarrow \mathcal{H}_i C$ is the zero map. Thus, for $i < n - 1$, we have that $\text{Image}(\mathcal{H}_{i+1} B \rightarrow \mathcal{H}_{i+1} C) = 0$, from which it follows that $\mathcal{H}_{i+1} C = 0$. In other words, $\mathcal{H}_i C = 0$ for $i < n$.

For any point, q^* , the spectrum $q^*L_\alpha\Sigma^\infty C$ satisfies the condition that $\pi_i^s q^*L_\alpha\Sigma^\infty C = 0$ for $i < n$ because $\pi_i^s L_\alpha\Sigma^\infty C = \mathcal{H}_i C = 0$, and $q^* \pi_i^s L_\alpha\Sigma^\infty C = \pi_i^s q^*L_\alpha\Sigma^\infty C$ [32, Section 2.2, page 12].

Thus, $\bigvee_{q^*I} q^*L_\alpha\Sigma^\infty C$ satisfies that condition that $\pi_i^s \bigvee_{q^*I} q^*L_\alpha\Sigma^\infty C = 0$ for $i < n$. Note that $\bigvee_{q^*I} q^*L_\alpha\Sigma^\infty C = q^*(\bigvee_I L_\alpha\Sigma^\infty C)$.

Thus, $q^* \pi_i^s \bigvee_I L_\alpha\Sigma^\infty C = \pi_i^s q^*(\bigvee_I L_\alpha\Sigma^\infty C) = 0$. Since q was arbitrary, we conclude that $\pi_i^s \bigvee_I L_\alpha\Sigma^\infty C = \pi_i^s q^*(\bigvee_I L_\alpha\Sigma^\infty C) = 0$.

By Remark 7.14, we conclude that

$$\mathcal{H}_i \bigvee_I L_\alpha\Sigma^\infty C = 0 \quad \text{for } i < n.$$

By Lemma 7.12, the map $\bigvee_I C \rightarrow \bigvee_I L_\alpha\Sigma^\infty C$ is an ms weak equivalence, whence

$$(29) \quad \mathcal{H}_i \bigvee_I C = 0 \quad \text{for } i < n.$$

By definition, we have that $A \xrightarrow{g_2} B' \rightarrow C$ is such that g_2 is a cofibration, and C is the cofiber of g_2 . By the definition of the global, injective local, \mathbb{A}^1 or $P\text{-}\mathbb{A}^1$ model structures, $\bigvee_I A \xrightarrow{g_2} \bigvee_I B'$ is a cofibration with cofiber $\bigvee_I C$. The sequence

$$\bigvee_I A \xrightarrow{g_2} \bigvee_I B' \rightarrow \bigvee_I C$$

therefore gives rise to a long exact sequence in \mathcal{H}_i . By Lemma 7.12, this long exact sequence can be written as

$$(30) \quad \cdots \rightarrow \mathcal{H}_{i+1} \bigvee_I C \rightarrow \mathcal{H}_i \bigvee_I A \xrightarrow{\mathcal{H}_i(g)} \mathcal{H}_i \bigvee_I B' \rightarrow \mathcal{H}_i \bigvee_I C \rightarrow \cdots .$$

The lemma is proven by combining (30) and (29). □

Corollary 7.16 *Suppose that $Y \in \mathbf{sPre}(\mathbf{Sm}_k)_*$ is pointed. Then we may replace the presheaf I of sets on \mathbf{Sm}_k by $U \mapsto N_n Y(U)$ in Lemma 7.15.*

Proof There is a functorial cofiber sequence

$$\bigvee_{L_n Y(U)} A \rightarrow \bigvee_{Y_n(U)} A \rightarrow \bigvee_{N_n Y(U)} A,$$

whence a functorial long exact sequence

$$\cdots \rightarrow \mathcal{H}_i(\bigvee_{L_n Y(U)} A) \rightarrow \mathcal{H}_i(\bigvee_{Y_n(U)} A) \rightarrow \mathcal{H}_i(\bigvee_{N_n Y(U)} A) \rightarrow \mathcal{H}_{i-1}(\bigvee_{L_n Y(U)} A) \rightarrow \cdots .$$

Applying Lemma 7.15 with $I = L_n Y$ and $I = Y_n$, the claim follows by the five lemma. □

7.3 A cancellation property

Say that a spectral sequence $E_{i,j}^r$ is a *first quadrant spectral sequence* if the differential on the r^{th} page is of bidegree $(-r, r - 1)$, ie $d_{i,j}^r: E_{i,j}^r \rightarrow E_{i-r,j+(r-1)}^r$, and if $E_{i,j}^r$ satisfies the condition that $E_{i,j}^r = 0$ when i or j is less than 0. The following lemma is a straightforward consequence of degree considerations, but we include the proof for completeness.

Lemma 7.17 *Suppose $\theta_{i,j}^r: E_{i,j}^r \rightarrow (E')_{i,j}^r$ is a map of first quadrant spectral sequences such that $\theta_{i,j}^1$ is an isomorphism for $j < q$. Then:*

- (i) $\theta_{i,j}^r$ is injective when $j < q$.
- (ii) $\theta_{i,j}^r$ is an isomorphism when $j + (r - 1) - 1 < q$ and $r \geq 2$.
- (iii) $\theta_{i,j}^r$ is an isomorphism when $j < q$ and $j + i \leq q$.

Proof We prove the claim by induction on q . For $q = 0$, there is nothing to show. Suppose the claim holds for $q - 1$. Now induct on r , which we assume ≥ 2 . Suppose that the claim holds for $r - 1$.

(i) Choose i and j with $j < q$. By the inductive hypothesis on r , we have that $\theta_{i,j}^{r-1}$ is injective. Since

$$(d')_{i,j}^{r-1} \theta_{i,j}^{r-1} = \theta_{i-(r-1),j+(r-1)-1}^{r-1} d_{i,j}^{r-1},$$

it follows that $\ker d_{i,j}^{r-1} \rightarrow \ker (d')_{i,j}^{r-1}$ is injective. It thus suffices to show that $\text{Image } d_{i+(r-1),j-((r-1)-1)}^r \rightarrow \text{Image } (d')_{i+(r-1),j-((r-1)-1)}^{r-1}$ is surjective (which is equivalent to being an isomorphism because $\theta_{i,j}^{r-1}$ is injective). Let $j' = j - ((r - 1) - 1)$. Note that $j' + ((r - 1) - 1) - 1 = j - 1 < q$. Thus, by the inductive hypothesis (ii) on r , $\theta_{i+(r-1),j-((r-1)-1)}^{r-1}$ is an isomorphism, from which the desired surjectivity follows.

(ii) Note that (ii) holds for $r = 2$. Choose i and j such that $j + (r - 1) - 1 < q$. Since $j + ((r - 1) - 1) - 1 < q$, we have by induction that $\theta_{i,j}^{r-1}$ is an isomorphism. We show that $\theta_{i,j}^r$ is an isomorphism. For this, it suffices to show that the inclusions $\ker d_{i,j}^{r-1} \subseteq \ker (d')_{i,j}^{r-1}$ and $\text{Image } d_{i+(r-1),j-((r-1)-1)}^r \subseteq \text{Image } (d')_{i+(r-1),j-((r-1)-1)}^{r-1}$ are isomorphisms. Since $j - ((r - 1) - 1) + ((r - 1) - 1) - 1 = j - 1 < q$, by the inductive hypothesis, we have that $\theta_{i+(r-1),j-((r-1)-1)}^{r-1}$ is an isomorphism. Thus, $\text{Image } d_{i+(r-1),j-((r-1)-1)}^r \subseteq \text{Image } (d')_{i+(r-1),j-((r-1)-1)}^{r-1}$ is an isomorphism. Note that $d_{i,j}^{r-1}$ is a map $E_{i,j}^{r-1} \rightarrow E_{i-(r-1),j+(r-1)-1}^{r-1}$ and similarly for $(d')^{r-1}$. Thus, to show that $\ker d_{i,j}^{r-1} \subseteq \ker (d')_{i,j}^{r-1}$ is an isomorphism, it suffices to show that

$\theta_{i-(r-1),j+(r-1)-1}^{r-1}$ is injective. Since $j + (r - 1) - 1 < q$, $\theta_{i-(r-1),j+(r-1)-1}^{r-1}$ is injective by (i).

(iii) Choose i and j such that $j < q$ and $j + i \leq q$. By (i), we have that $\theta_{i,j}^r$ is injective. We show surjectivity. By the inductive hypothesis, $\theta_{i,j}^{r-1}$ is an isomorphism. Thus, it suffices to see that $\ker d_{i,j}^{r-1} \subseteq \ker (d')_{i,j}^{r-1}$ is an isomorphism. Note that if $r = 2$, then $\theta_{i,j}^1$ and $\theta_{i-1,j}^1$ are isomorphisms and $\theta_{i-1,j}^1 d_{i,j}^1 = (d')_{i,j}^1 \theta_{i,j}^1$, whence $\ker d_{i,j}^1 \subseteq \ker (d')_{i,j}^1$ is an isomorphism. So, we may assume that $r > 2$. Since $(d')_{i,j}^{r-1} \theta_{i,j}^{r-1} = \theta_{i-(r-1),j+(r-1)-1}^{r-1} d_{i,j}^{r-1}$, it suffices to see that $\theta_{i-(r-1),j+(r-1)-1}^{r-1}$ is injective. For $i - (r - 1) < 0$, we have that $E_{i-(r-1),j+(r-1)-1}^{r-1} = 0$ so $\theta_{i-(r-1),j+(r-1)-1}^{r-1}$ is injective. Thus, we may assume $i - (r - 1) \geq 0$ or equivalently $r \leq i + 1$. Thus, $j + (r - 1) - 1 \leq j + (i + 1 - 1) - 1 = j + i - 1 \leq q - 1 < q$. Thus, $\theta_{i-(r-1),j+(r-1)-1}^{r-1}$ is injective by (i). \square

Let (α, \mathcal{H}_i) be either $(\mathbb{A}^1, \pi_i^{s, \mathbb{A}^1})$ or $(P - \mathbb{A}^1, \pi_i^{s, P, \mathbb{A}^1})$.

As in Section 7.2, let $j: J \rightarrow Y$ be a map of pointed simplicial presheaves, and let $b: \Sigma J \rightarrow \Sigma X$ be a map in $\text{ho}_\alpha \mathbf{sPre}(\mathbf{Sm}_k)_*$ between the suspensions of J and X , for X a pointed simplicial presheaf. Factor j as $j = \iota \circ f$ with $\iota: J \rightarrow E$ an α weak equivalence and cofibration, and $f: E \rightarrow Y$ an α fibration. Assume as above that X is fibrant. Then $p: X \times Y \rightarrow Y$ is also an α fibration.

Let $\mathfrak{b}: \Sigma E \rightarrow \Sigma(X \times Y)$ in $\text{ho}_\alpha \mathbf{sPre}(\mathbf{Sm}_k)_*$ be as in Construction 7.10. Suppose that Y is 1-reduced, that is to say, $Y_0 = Y_1 = *$. Let $a: F \rightarrow E$ denote the canonical map of simplicial presheaves $F = * \times_Y E \rightarrow E$. We suppose that the resulting map $\Sigma F \rightarrow \Sigma X$ induces a surjection on \mathcal{H}_i for all i . (Indeed, we will later use this construction when $\Sigma F \rightarrow \Sigma X$ has a section up to homotopy.)

Assume that $\mathcal{H}_0(X)$, $\mathcal{H}_0(F)$, $\mathcal{H}_0(Y)$ and $\mathcal{H}_0(E)$ are all 0. For example, this is satisfied if X, F, Y and E are \mathbb{A}^1 -connected, because $\pi_i^{s, P, \mathbb{A}^1}(-) \cong \pi_i^{s, \mathbb{A}^1}(-) \otimes_{\mathbb{Z}} \mathbb{Z}_P$; see Proposition 3.25.

Proposition 7.18 *If $\mathcal{H}_i(\Sigma^{-1}\mathfrak{b})$ is an isomorphism for all i , then*

$$\mathcal{H}_i(\Sigma^{-1}b \circ a): \mathcal{H}_i F \rightarrow \mathcal{H}_i X$$

is an isomorphism for all i , and $\Sigma^{-1}b \circ a: \Sigma^\infty F \rightarrow \Sigma^\infty X$ is an α weak equivalence.

Proof Since $(\mathfrak{a}, \mathcal{H}_i)$ is either $(\mathbb{A}^1, \pi_i^{s, \mathbb{A}^1})$ or $(P-\mathbb{A}^1, \pi_i^{s, P, \mathbb{A}^1})$, the functors \mathcal{H}_i detect stable \mathfrak{a} weak equivalences by Proposition 2.15 or Propositions 3.25 and 3.26, respectively. Thus, it suffices to prove that $\mathcal{H}_i(\Sigma^{-1}b \circ a): \mathcal{H}_i F \rightarrow \mathcal{H}_i X$ is an isomorphism for all i .

By Lemma 7.11, we have an induced morphism of spectral sequences $\theta_{i,j}^r: (E_{i,j}^r, d^r) \rightarrow ((E')_{i,j}^r, (d')^r)$ from the spectral sequence of Definition 7.3 induced by f to the one induced by p .

Suppose for the sake of contradiction that $\mathcal{H}_i(\Sigma^{-1}b \circ a): \mathcal{H}_i F \rightarrow \mathcal{H}_i X$ is not an isomorphism for all i . Then there exists a minimal q such that $\mathcal{H}_q F \rightarrow \mathcal{H}_q X$ is not an isomorphism, and $q > 0$ by the assumption that both $\mathcal{H}_0 F$ and $\mathcal{H}_0 X$ are both 0. By Lemma 7.15 and Corollary 7.16, it follows that

$$(31) \quad \theta_{i,j}^1: E_{i,j}^1 \xrightarrow{\cong} (E')_{i,j}^1$$

is an isomorphism for $j < q$.

By Proposition 7.6(ii) and Lemma 7.17, $\theta_{i,j}^r$ is an isomorphism for $i + j = q$ and $i > 0$ and all r . For r sufficiently large, $\theta_{i,j}^r = \theta_{i,j}^\infty$ (Proposition 7.6(ii)), whence $\theta_{i,j}^\infty$ is an isomorphism for $i + j = q$ and $i > 0$.

Introduce the notation

$$0 \subseteq R_n^0 \subseteq R_n^1 \subseteq R_n^2 \subseteq \dots \subseteq R_n^n = \mathcal{H}_n(E)$$

for the filtration associated to the spectral sequence E_{ij}^r . Let the corresponding filtration associated to the spectral sequence $(E')_{i,j}^r$ be denoted by $T_n^i \subseteq \mathcal{H}_n(X \times Y)$. Note that we have the commutative diagram

$$(32) \quad \begin{array}{ccccccc} 0 & \longrightarrow & T_q^i & \longrightarrow & T_q^{i+1} & \longrightarrow & (E')_{i+1, q-(i+1)}^\infty \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \theta_{i+1, q-(i+1)}^\infty \\ 0 & \longrightarrow & R_q^i & \longrightarrow & R_q^{i+1} & \longrightarrow & E_{i+1, q-(i+1)}^\infty \longrightarrow 0 \end{array}$$

for $i = -1, \dots, q-1$, where by convention $R_n^{-1} = 0$ and $T_n^{-1} = 0$ for all n .

Since $\Sigma^{-1}b$ induces an isomorphism on \mathcal{H}_q , we have that $R_q^q \rightarrow T_q^q$ is an isomorphism. Applying the five lemma and (32) for $i = q-1, q-2, \dots, 0$, we conclude that $R_q^0 \rightarrow T_q^0$ is an isomorphism. By definition, $R_q^0 = E_{0,q}^\infty$ and $T_q^0 = (E')_{0,q}^\infty$, so we have that

$$\theta_{0,q}^\infty: E_{0,q}^\infty \rightarrow (E')_{0,q}^\infty$$

is an isomorphism.

Since $Y_0 = *$, the map $\theta_{0,1}^1: E_{0,q}^1 \rightarrow (E')_{0,q}^1$ is identified with $\mathcal{H}_q F \rightarrow \mathcal{H}_q X$ by Proposition 7.6(i) and Lemma 7.13. Since we have assumed that $\mathcal{H}_q F \rightarrow \mathcal{H}_q X$ is not an isomorphism, it follows that

$$\theta_{0,q}^1: E_{0,q}^1 \rightarrow (E')_{0,q}^1$$

is not an isomorphism. Since we have assumed that $Y_1 = *$, Proposition 7.6(i) implies that the domains of $d_{1,q}^1$ and $(d')_{1,q}^1$ are zero. Thus, $d_{1,q}^1$ and $(d')_{1,q}^1$ are zero. Thus, $\theta_{0,q}^2$ is not an isomorphism. Let n be maximal such that $E_{0,q}^n \rightarrow (E')_{0,q}^n$ is not an isomorphism. Since $d_{0,q}^n = 0$, $(d')_{0,q}^n = 0$, and $E_{0,q}^{n+1} \rightarrow (E')_{0,q}^{n+1}$ is an isomorphism, we conclude that

$$\text{Image } d_{n,q-(n-1)}^n \rightarrow \text{Image } (d')_{n,q-(n-1)}^n$$

is not an isomorphism.

Note that $q + 1 - i + (r - 1) - 1 < q$ when $r < i + 1$. Thus, by Lemma 7.17(ii), we have that

$$\theta_{i,q+1-i}^r: E_{i,q+1-i}^r \rightarrow (E')_{i,q+1-i}^r$$

is an isomorphism for $i > 1$ and $r = 1, 2, \dots, i$. For $r = i + 1$,

$$\begin{aligned} E_{i,q+1-i}^{i+1} &\cong \ker d_{i,q+1-i}^i / \text{Image } d_{2i,q+1-i-(i-1)}^i, \\ (E')_{i,q+1-i}^{i+1} &\cong \ker (d')_{i,q+1-i}^i / \text{Image } (d')_{2i,q+1-i-(i-1)}^i. \end{aligned}$$

Furthermore, we have the isomorphism

$$\text{Image } d_{2i,q+1-i-(i-1)}^i \cong \text{Image } (d')_{2i,q+1-i-(i-1)}^i$$

by Lemma 7.17(ii). Thus, $E_{i,q+1-i}^{i+1} \xrightarrow{\cong} (E')_{i,q+1-i}^{i+1}$ is an injection, and is an isomorphism if and only if

$$\ker d_{i,q+1-i}^i \rightarrow \ker (d')_{i,q+1-i}^i$$

is an isomorphism, which happens if and only if $\text{Image } d_{i,q+1-i}^i \rightarrow \text{Image } (d')_{i,q+1-i}^i$ is an isomorphism.

By the above, we thus conclude that $E_{n,q+1-n}^{n+1} \xrightarrow{\cong} (E')_{n,q+1-n}^{n+1}$ is not surjective.

Note that by degree reasons, $d_{n,q+1-n}^m = 0$ and $(d')_{n,q+1-n}^m = 0$ for $m \geq n + 1$. Since by Lemma 7.17(i)–(ii) we have $\text{Image } d_{n+m,q+1-n-(m-1)}^m \cong \text{Image } (d')_{n+m,q+1-n-(m-1)}^m$, we conclude that

$$E_{n,q+1-n}^\infty \xrightarrow{\cong} (E')_{n,q+1-n}^\infty$$

is not surjective.

The same reasoning shows that $E_{i,q+1-i}^\infty \xrightarrow{\subseteq} (E')_{i,q+1-i}^\infty$ is an isomorphism for $i > n \geq 2$. Explicitly, for $i > n$, we have that $E_{0,q}^i \rightarrow (E')_{0,q}^i$ is an isomorphism by the choice of n . As above, note that since $d_{0,q}^i = 0$, $(d')_{0,q}^i = 0$ and $E_{0,q}^{i+1} \rightarrow (E')_{0,q}^{i+1}$ is an isomorphism, we conclude that the map $\text{Image } d_{i,q-(i-1)}^i \rightarrow \text{Image}(d')_{i,q-(i-1)}^i$ is an isomorphism. Continuing with the same reasoning, note that $\theta_{i,q+1-i}^i: E_{i,q+1-i}^i \rightarrow (E')_{i,q+1-i}^i$ is an isomorphism by Lemma 7.17(ii), whence $\ker d_{i,q-(i-1)}^i \rightarrow \ker(d')_{i,q-(i-1)}^i$ is an isomorphism. Since

$$\text{Image } d_{2i,q+1-i-(i-1)}^i \cong \text{Image}(d')_{2i,q+1-i-(i-1)}^i$$

by Lemma 7.17(i)–(ii), we have that $\theta_{i,q+1-i}^{i+1}: E_{i,q+1-i}^{i+1} \rightarrow (E')_{i,q+1-i}^{i+1}$ is an isomorphism. By degree reasons $d_{i,q+1-i}^m = 0$ and $(d')_{i,q+1-i}^m = 0$ for $m \geq i + 1$. Since by Lemma 7.17(i)–(ii) we have $\text{Image } d_{i+m,q+1-i-(m-1)}^m \cong \text{Image}(d')_{i+m,q+1-i-(m-1)}^m$, we have that $E_{i,q+1-i}^\infty \xrightarrow{\subseteq} (E')_{i,q+1-i}^\infty$ is an isomorphism for $i > n$, as claimed.

Since $\Sigma^{-1}\mathfrak{b}$ induces an isomorphism on \mathcal{H}_{q+1} , we have that $R_{q+1}^{q+1} \rightarrow T_{q+1}^{q+1}$ is an isomorphism. Combining this with the commutative diagram

$$(33) \quad \begin{array}{ccccccc} 0 & \longrightarrow & T_{q+1}^i & \longrightarrow & T_{q+1}^{i+1} & \longrightarrow & (E')_{i+1,q+1-(i+1)}^\infty & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \theta_{i+1,q+1-(i+1)}^\infty & \uparrow & \\ 0 & \longrightarrow & R_{q+1}^i & \longrightarrow & R_{q+1}^{i+1} & \longrightarrow & E_{i+1,q+1-(i+1)}^\infty & \longrightarrow & 0 \end{array}$$

for $i = q, q-1, \dots, n$ and the five lemma, we see that $R_{q+1}^i \rightarrow T_{q+1}^i$ is an isomorphism for $i = q, q-1, \dots, n$.

In particular, $R_{n+1}^n \rightarrow T_{n+1}^n$ is surjective, which by (33) with $i = n-1$ implies that $E_{n,q+1-n}^\infty \xrightarrow{\subseteq} (E')_{n,q+1-n}^\infty$ is surjective, giving the desired contradiction. \square

Here is a verbal description of the proof of Proposition 7.18. Choose q minimal such that $\pi_q^{s,\mathbb{A}^1}(F) \rightarrow \pi_q^{s,\mathbb{A}^1}(X)$ is not an isomorphism. The failure to be an isomorphism is necessarily a failure of injectivity. In terms of the map of spectral sequences, this implies that $E_{0,1}^1 \rightarrow (E')_{0,q}^1$ is not injective. Since q is minimal, degree arguments with first quadrant spectral sequences imply that $E_{i,q-i}^\infty \rightarrow (E')_{i,q-i}^\infty$ are isomorphisms for $i > 0$. Since $\pi_q^{s,\mathbb{A}^1}((b \wedge \Sigma^\infty f_+) \circ \Sigma^\infty \Delta_+)$ is an isomorphism, and since $\bigoplus_i E_{i,q-i}^\infty$ is the associated graded of $\pi_q^{\mathbb{A}^1,s}(E)$ and the analogous statement for E' and $X \times Y$ holds, it follows that $E_{0,q}^\infty \rightarrow (E')_{i,q-i}^\infty$ is also an isomorphism. Thus, we can choose a maximal $n \geq 2$ for which $E_{0,q}^n \rightarrow (E')_{0,q}^n$ is not an isomorphism. For degree reasons,

the failure of this map to be an isomorphism is a failure of injectivity. Thus, the image of a d must be larger than the image of a d' . Since the domains of these differentials have smaller second index, these domains actually have to be isomorphic. This then implies that the kernel of the d is smaller than the kernel of the (d') . This leads to a contradiction with the surjectivity of $\pi_{q+1}^{s, \mathbb{A}^1}(F) \rightarrow \pi_{q+1}^{s, \mathbb{A}^1}(X)$.

8 \mathbb{A}^1 simplicial EHP fiber sequence

Definition 8.1 Say that the sequence $X \rightarrow Y \rightarrow Z$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is an a fiber sequence up to homotopy if there is a diagram

$$\begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ F & \longrightarrow & E & \xrightarrow{f} & B \end{array}$$

which commutes up to homotopy with B fibrant, f a fibration with fiber $F = * \times_B E$, and all the vertical maps weak equivalences.

Remark 8.2 Morel defines $X \rightarrow Y \rightarrow Z$ to be a simplicial fibration sequence if the composition $X \rightarrow Z$ is the constant map and the induced map from X to the homotopy fiber of $Y \rightarrow Z$ in the injective local model structure is a simplicial weak equivalence.

He then defines $X \rightarrow Y \rightarrow Z$ to be an \mathbb{A}^1 -fibration sequence if $L_{\mathbb{A}^1} X \rightarrow L_{\mathbb{A}^1} Y \rightarrow L_{\mathbb{A}^1} Z$ is a simplicial fibration sequence. See [33, Definition 6.44].

In this vein, it is natural to define $X \rightarrow Y \rightarrow Z$ to be a $P\text{-}\mathbb{A}^1$ fibration sequence if

$$L_P L_{\mathbb{A}^1} X \rightarrow L_P L_{\mathbb{A}^1} Y \rightarrow L_P L_{\mathbb{A}^1} Z$$

is a simplicial fibration sequence.

Note that if $X \rightarrow Y \rightarrow Z$ in $\mathbf{sPre}(\mathbf{Sm}_k)_*$ is a $P\text{-}\mathbb{A}^1$ fiber sequence up to homotopy as in 8.1, then B, E and F are $P\text{-}\mathbb{A}^1$ -local, from which it follows that they can be identified with $L_P L_{\mathbb{A}^1} X, L_P L_{\mathbb{A}^1} E$ and $L_P L_{\mathbb{A}^1} F$, respectively. Since $P\text{-}\mathbb{A}^1$ fibrations are simplicial fibrations, we have that a $P\text{-}\mathbb{A}^1$ fiber sequence up to homotopy is a $P\text{-}\mathbb{A}^1$ fibration sequence.

Note that for P the set of all primes, the $P\text{-}\mathbb{A}^1$ injective model structures on $\mathbf{sPre}(\mathbf{Sm}_k)$ and $\mathbf{Spt}(\mathbf{Sm}_k)$ are the \mathbb{A}^1 injective model structures, and $X \rightarrow L_P X$ is the identity map.

For $X \in \mathbf{sPre}(\mathbf{Sm}_k)_*$, let $j: J(X) \rightarrow J(X^{\wedge 2})$ be j_2 from Definition 5.5. Recall from Section 4.1 and before the notation $S^{n+q\alpha} = S^n \wedge \mathbb{G}_m^{\wedge q}$. Recall from Sections 1 and 6 the notation $\langle -1 \rangle$ in $\mathrm{GW}(k)$. Recall from Section 3 that for a set of primes P , \mathbb{Z}_P denotes \mathbb{Z} with all primes not in P inverted.

Theorem 8.3 *Let $X = S^{n+q\alpha}$ with $n > 1$, and let $e = (-1)^{n+q} \langle -1 \rangle^q$. Let P be a set of primes, and suppose that for all $m \in \mathbb{Z}_{>0}$, the element $(m + 1) + me$ is a unit in $\mathrm{GW}(k) \otimes \mathbb{Z}_P$. Then*

$$(34) \quad X \rightarrow J(X) \xrightarrow{j} J(X^{\wedge 2})$$

is a fiber sequence up to homotopy in the P - \mathbb{A}^1 injective model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$.

By Proposition 5.2, Theorem 8.3 proves Theorem 1.3.

Proof Let \mathfrak{a} denote the P - \mathbb{A}^1 injective model structure on $\mathbf{sPre}(\mathbf{Sm}_k)$ and $\mathbf{Spt}(\mathbf{Sm}_k)$.

Recall the notation $D(X) = \bigvee_{j=0}^{\infty} X^{\wedge j}$ from Definition 5.1. From Section 5, we have the zigzag (12),

$$\Sigma J(X)_+ \rightarrow \mathrm{Simp} |\Sigma D(X)| \leftarrow \Sigma D(X),$$

of weak equivalences in the global model structure.

Let $b_1: D(X) \rightarrow L_P L_{\mathbb{A}^1} X$ denote the composition of the map $D(X) \rightarrow X$ which for $j \neq 1$ crushes the summands $X^{\wedge j}$ to the basepoint with the canonical map $X \rightarrow L_P L_{\mathbb{A}^1} X$.

Replace $J(X^{\wedge 2})$ by a fibrant simplicial presheaf $J(X^{\wedge 2}) \rightarrow L_P L_{\mathbb{A}^1} J(X^{\wedge 2})$. Let $f': E' \rightarrow L_P L_{\mathbb{A}^1} J(X^{\wedge 2})$ be a fibrant replacement of

$$J(X) \xrightarrow{j} J(X^{\wedge 2}) \rightarrow L_P L_{\mathbb{A}^1} J(X^{\wedge 2})$$

is the \mathfrak{a} model structure. Let

$$E = J(X^{\wedge 2}) \times_{L_P L_{\mathbb{A}^1} J(X^{\wedge 2})} E'$$

be the pullback of E' to $J(X^{\wedge 2})$, and let $f: E \rightarrow J(X^{\wedge 2})$ be the canonical projection. (The motivation for defining E' is to obtain the diagram (35) below, which fits precisely into Definition 8.1. If this is overly pedantic, the reader may consider a fibrant replacement of j .) Note that f is an \mathfrak{a} fibration, and that there is a canonical map $J(X) \rightarrow E$. Since \mathfrak{a} is a proper model structure and $J(X^{\wedge 2}) \rightarrow L_P L_{\mathbb{A}^1} J(X^{\wedge 2})$ is an \mathfrak{a} weak equivalence, we have that $E \rightarrow E'$ is a weak equivalence. Since $J(X) \rightarrow E'$

is an α weak equivalence by construction, by the 2-out-of-3 property it follows that the canonical map $J(X) \rightarrow E$ is an α weak equivalence. We thus have a commutative diagram

$$\begin{array}{ccc} J(X) & \xrightarrow{\cong_\alpha} & E \\ & \searrow j & \downarrow f \\ & & J(X^{\wedge 2}) \end{array}$$

with the map $J(X) \xrightarrow{\cong} E$ an α weak equivalence.

It follows that we have the zigzag

$$\Sigma E \xleftarrow{\cong} \Sigma J(X)_+ \xrightarrow{\cong} \text{Simp}|\Sigma D(X)| \xleftarrow{\cong} \Sigma D(X) \xrightarrow{b_1} \Sigma L_P L_{\mathbb{A}^1} X,$$

which determines a map $b: \Sigma^\infty E \rightarrow \Sigma^\infty L_P L_{\mathbb{A}^1} X$ in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$.

As in Sections 6.4 and 7.2, we obtain a map

$$(b_+ \wedge \Sigma^\infty f_+) \circ \Sigma^\infty \Delta_+ : \Sigma^\infty E_+ \rightarrow \Sigma^\infty (L_P L_{\mathbb{A}^1} X \times J(X^{\wedge 2}))_+$$

in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$. The hypothesis that for all $m \in \mathbb{Z}_{>0}$ the element $(m + 1) + me$ is a unit in $\text{GW}(k) \otimes \mathbb{Z}_P$ implies that $(b_+ \wedge \Sigma^\infty f_+) \circ \Sigma^\infty \Delta_+$ is an isomorphism by Proposition 6.17. By Section 5.1, $J(X^{\wedge 2})$ is 1-reduced and $L_P L_{\mathbb{A}^1} X$ is fibrant, so we may apply Proposition 7.18. It follows that $b \circ \Sigma^\infty a: \Sigma^\infty F \rightarrow \Sigma^\infty X$ is an isomorphism in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$, where $a: F \rightarrow E$ is the inclusion of the fiber of f into E .

Since E is the pullback of E' we have the diagram in $\mathbf{sPre}(\mathbf{Sm}_k)$,

$$\begin{array}{ccccc} F & \longrightarrow & E' & \longrightarrow & L_P L_{\mathbb{A}^1} J(X^{\wedge 2}) \\ \uparrow & & \uparrow \cong_\alpha & & \uparrow \cong_\alpha \\ F & \longrightarrow & E & \longrightarrow & J(X^{\wedge 2}) \end{array}$$

We conclude that $b \circ \Sigma^\infty a': \Sigma^\infty F \rightarrow \Sigma^\infty X$ is an isomorphism in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$, where a' is the composition in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$ corresponding to the zigzag $F \rightarrow E' \leftarrow E$.

Since the composition of the two maps in the sequence (34) is constant, the composition

$$X \rightarrow J(X) \rightarrow L_P L_{\mathbb{A}^1} J(X^{\wedge 2})$$

is also constant. We therefore have an induced map

$$h: X \rightarrow F.$$

By construction of b_1 , the composition $\Sigma^\infty X \rightarrow \Sigma^\infty E \xrightarrow{b} \Sigma^\infty L_P L_{\mathbb{A}^1} X$ is the canonical map associated to the identity in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$, and in particular is an isomorphism. Thus, the composition $\Sigma^\infty X \xrightarrow{\Sigma^\infty h} \Sigma^\infty F \xrightarrow{\Sigma^\infty a'} \Sigma^\infty E \xrightarrow{b} \Sigma^\infty X$ is an isomorphism. Since $b \circ \Sigma^\infty a'$ is an isomorphism in $\text{ho}_\alpha \mathbf{Spt}(\mathbf{Sm}_k)$, so is $\Sigma^\infty h$.

Note that $L_P h: L_P X \rightarrow L_P F$ is a map of \mathbb{A}^1 -simply connected objects. $\Sigma^\infty L_P h$ is an \mathbb{A}^1 weak equivalence as follows. By Corollary 3.20, we have that $\Sigma^\infty L_P h \simeq L_P \Sigma^\infty h$. Since $\Sigma^\infty h$ is a P - \mathbb{A}^1 weak equivalence, we have that $L_P L_{\mathbb{A}^1} \Sigma^\infty h$ is a simplicial weak equivalence. By Proposition 3.24, we have $L_P L_{\mathbb{A}^1} \Sigma^\infty h \simeq L_{\mathbb{A}^1} L_P \Sigma^\infty h$, whence $L_{\mathbb{A}^1} L_P \Sigma^\infty h$ is a simplicial weak equivalence, so $L_P \Sigma^\infty h$ is an \mathbb{A}^1 weak equivalence, as claimed. We now apply Corollary 2.23 to conclude that $L_P h$ is an \mathbb{A}^1 weak equivalence, whence h is an α weak equivalence.

The diagram

$$(35) \quad \begin{array}{ccccc} X & \longrightarrow & J(X) & \xrightarrow{j} & J(X^{\wedge 2}) \\ \downarrow & & \downarrow & & \downarrow \\ F & \longrightarrow & E' & \longrightarrow & L_P L_{\mathbb{A}^1} J(X^{\wedge 2}) \end{array}$$

shows that (34) is an α fiber sequence up to homotopy. □

Proof of Corollary 1.4 • By Theorem 8.3, it is sufficient to show that $(m + 1) + me$ is a unit in $\text{GW}(k) \otimes \mathbb{Z}_{(2)}$ for all positive integers m . This was shown in Corollary 4.7.

- Apply Theorem 8.3 and Corollary 4.8.
- When n is odd and q is even, we have $e = -1$, whence $(m + 1) + me = 1$ is a unit in $\text{GW}(k)$.
- When $2\eta = 0$, we have $2\eta\rho = 0$, whence $\eta\rho$ is torsion in $\text{GW}(k)$. All torsion elements of $\text{GW}(k)$ are nilpotent [28, Chapter VIII, Section 8.1] (or because the Grothendieck–Witt ring is a λ -ring, and all torsion elements of λ rings are nilpotent by a result of Graeme Segal), so $\eta\rho$ is nilpotent. When $n + q$ is odd,

$$e = (-1)^n (-1)^q (1 + \rho\eta)^q = -(1 + \rho\eta)^q = \begin{cases} -1 - \rho\eta & \text{if } q \text{ is odd,} \\ -1 & \text{if } q \text{ is even.} \end{cases}$$

Thus, $1 + e$ is nilpotent, from which it follows that $(m + 1) + me = 1 + (e + 1)m$ is a unit in $\text{GW}(k)$.

The fact that $2\eta = 0$ when $k = \mathbb{C}$ is shown [31, Remark 6.3.5 and Lemma 6.3.7]. □

Corollary 8.4 *Let $X = S^{n+q\alpha}$ with $n > 1$. Choose $v \in \mathbb{Z}$. There is a functorial long exact sequence*

$$\begin{aligned} \cdots \rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1} X &\rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+1+v\alpha}^{\mathbb{A}^1} \Sigma X \rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+1+v\alpha}^{\mathbb{A}^1} \Sigma(X \wedge X) \\ &\rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i-1+v\alpha}^{\mathbb{A}^1} X \rightarrow \cdots \end{aligned}$$

Proof Combining Theorem 8.3 and Corollary 1.4, we have that

$$X \rightarrow J(X) \rightarrow J(X^{\wedge 2})$$

is a $2-\mathbb{A}^1$ fiber sequence up to homotopy. It follows that there is an associated long exact sequence in $\pi_{*+v\alpha}^{2,\mathbb{A}^1}$. See Proposition 3.17. By Proposition 3.16 and [33, Theorem 6.13], we may replace $\pi_{*+v\alpha}^{2,\mathbb{A}^1}$ with $\mathbb{Z}_{(2)} \otimes \pi_{*+v\alpha}^{\mathbb{A}^1}$. By Corollary 5.3, we can identify the homotopy groups $\pi_{*+v\alpha}^{\mathbb{A}^1} J(X)$ with $\pi_{*+1+v\alpha}^{\mathbb{A}^1} \Sigma X$. This yields the claimed long exact sequence. \square

The long exact sequences of Corollary 8.4 form an exact couple, which in turn gives rise to an \mathbb{A}^1 simplicial EHP spectral sequence. Here the adjective ‘‘simplicial’’ refers to the suspension with respect to the simplicial circle S^1 .

Theorem 8.5 *Choose $q, v \in \mathbb{Z}_{\geq 0}$ and $n \in \mathbb{Z}$ such that $n \geq 2$. There is a spectral sequence*

$$(E_{i,m}^r, d_r: E_{i,m}^r \rightarrow E_{i-1,m-r}^r) \Rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i-n,v-q}^{s,\mathbb{A}^1} = \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{s,\mathbb{A}^1} S^{n+q\alpha}$$

with $E_{i,m}^1 = \mathbb{Z}_{(2)} \otimes \pi_{m+1+i+v\alpha}^{\mathbb{A}^1} (S^{2m+2n+1+2q\alpha})$ if $i \geq 2n - 1 + m$ and otherwise $E_{i,m}^1 = 0$.

Proof The E^1 -page is as claimed by the construction of the exact couple and by the fact that $\pi_{i+v\alpha}^{\mathbb{A}^1} S^{n+q\alpha} = 0$ for $i < n$. The latter fact follows from Morel’s connectivity theorem [33, Theorem 6.38].

The spectral sequence converges for degree reasons: for all (i, m) there are only finitely many r with a nonzero differential leaving or entering $E_{i,m}^r$.

Since by definition we have $\text{colim}_m \pi_{m+i+v\alpha}^{\mathbb{A}^1} \Sigma^m S^{n+q\alpha} = \pi_{i+v\alpha}^{s,\mathbb{A}^1} S^{n+q\alpha}$ (see Section 2.4), we also have

$$\text{colim}_m (\mathbb{Z}_{(2)} \otimes \pi_{i+m+v\alpha}^{\mathbb{A}^1} \Sigma^m S^{n+q\alpha}) = \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{s,\mathbb{A}^1} S^{n+q\alpha},$$

and it follows that the spectral sequence converges to $\mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{s,\mathbb{A}^1} S^{n+q\alpha}$. \square

As in the setting of algebraic topology, one obtains truncated EHP spectral sequences converging to unstable homotopy groups of spheres.

Theorem 8.6 Choose $q, v \in \mathbb{Z}_{\geq 0}$ and $n_1, n_2 \in \mathbb{Z}$ such that $n_2 \geq n_1 \geq 2$. There is a spectral sequence $(E_{i,m}^r, d_r: E_{i,m}^r \rightarrow E_{i-1,m-r}^r) \Rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1} S^{n_2+q\alpha}$ with

$$E_{i,m}^1 = \begin{cases} \mathbb{Z}_{(2)} \otimes \pi_{m+1+i+v\alpha}^{\mathbb{A}^1} (S^{2m+2n_1+1+2q\alpha}) & \text{if } i \geq 2n-1+m \text{ and } 0 \leq m < n_2-n_1, \\ 0 & \text{otherwise.} \end{cases}$$

Proof The long exact sequences of Corollary 8.4 for $X = S^{n_1+m+q\alpha}$ with $0 \leq m < n_2 - n_1$ can be combined with the long exact sequence

$$\dots \rightarrow 0 \rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1} S^{n_2+q\alpha} \rightarrow \mathbb{Z}_{(2)} \otimes \pi_{i+v\alpha}^{\mathbb{A}^1} S^{n_2+q\alpha} \rightarrow 0 \rightarrow \dots$$

associated to the $2-\mathbb{A}^1$ fiber sequence

$$S^{n_2+q\alpha} \rightarrow S^{n_2+q\alpha} \rightarrow *$$

(which replaces in Theorem 8.5 the long exact sequences of Corollary 8.4 for $X = S^{n_1+m+q\alpha}$ with $n_2 - n_1 \leq m$) to form an exact couple. The E^1 -page equals the E^1 -page of the EHP sequence constructed in Theorem 8.5 for $m < n_2 - n_1$, and $E_{i,m}^1 = 0$ for $m \geq n_2 - n_1$. The convergence is clear. \square

References

- [1] **A Asok, J Fasel**, *A cohomological classification of vector bundles on smooth affine threefolds*, Duke Math. J. 163 (2014) 2561–2601 MR
- [2] **A Asok, J Fasel**, *Splitting vector bundles outside the stable range and \mathbb{A}^1 -homotopy sheaves of punctured affine spaces*, J. Amer. Math. Soc. 28 (2015) 1031–1062 MR
- [3] **A Asok, J Fasel**, *Comparing Euler classes*, Q. J. Math. 67 (2016) 603–635 MR
- [4] **A Asok, K Wickelgren, B Williams**, *The simplicial suspension sequence in \mathbb{A}^1 -homotopy*, Geom. Topol. 21 (2017) 2093–2160 MR
- [5] **C Barwick**, *On left and right model categories and left and right Bousfield localizations*, Homology Homotopy Appl. 12 (2010) 245–320 MR
- [6] **A K Bousfield, DM Kan**, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics 304, Springer (1972) MR
- [7] **G Carlsson, R J Milgram**, *Stable homotopy and iterated loop spaces*, from “Handbook of algebraic topology” (I M James, editor), North-Holland, Amsterdam (1995) 505–583 MR

- [8] **C Casacuberta, G Peschke**, *Localizing with respect to self-maps of the circle*, Trans. Amer. Math. Soc. 339 (1993) 117–140 MR
- [9] **D Dugger, D C Isaksen**, *Motivic cell structures*, Algebr. Geom. Topol. 5 (2005) 615–652 MR
- [10] **T Ganea**, *On the homotopy suspension*, Comment. Math. Helv. 43 (1968) 225–234 MR
- [11] **P G Goerss, J F Jardine**, *Simplicial homotopy theory*, Progress in Mathematics 174, Birkhäuser, Basel (1999) MR
- [12] **P S Hirschhorn**, *Model categories and their localizations*, Mathematical Surveys and Monographs 99, Amer. Math. Soc., Providence, RI (2003) MR
- [13] **A Hogadi, G Kulkarni**, *Gabber’s presentation lemma for finite fields*, J. Reine Angew. Math. (online publication January 2018)
- [14] **J Hornbostel**, *Localizations in motivic homotopy theory*, Math. Proc. Cambridge Philos. Soc. 140 (2006) 95–114 MR
- [15] **M Hovey**, *Model categories*, Mathematical Surveys and Monographs 63, Amer. Math. Soc., Providence, RI (1999) MR
- [16] **M Hovey**, *Spectra and symmetric spectra in general model categories*, J. Pure Appl. Algebra 165 (2001) 63–127 MR
- [17] **Y Ihara**, *Braids, Galois groups, and some arithmetic functions*, from “Proceedings of the International Congress of Mathematicians” (I Satake, editor), volume I, Math. Soc. Japan, Tokyo (1991) 99–120 MR
- [18] **D C Isaksen**, *Flasque model structures for simplicial presheaves*, *K–Theory* 36 (2005) 371–395 MR
- [19] **I M James**, *Reduced product spaces*, Ann. of Math. 62 (1955) 170–197 MR
- [20] **I M James**, *On the suspension triad*, Ann. of Math. 63 (1956) 191–247 MR
- [21] **I M James**, *The suspension triad of a sphere*, Ann. of Math. 63 (1956) 407–429 MR
- [22] **I M James**, *On the suspension sequence*, Ann. of Math. 65 (1957) 74–107 MR
- [23] **J F Jardine**, *Simplicial presheaves*, J. Pure Appl. Algebra 47 (1987) 35–87 MR
- [24] **J F Jardine**, *Motivic symmetric spectra*, Doc. Math. 5 (2000) 445–552 MR
- [25] **N J Kuhn**, *The transfer and James–Hopf invariants*, Math. Z. 196 (1987) 391–405 MR
- [26] **N J Kuhn**, *Stable splittings and the diagonal*, from “Homotopy methods in algebraic topology” (JPC Greenlees, RR Bruner, N Kuhn, editors), Contemp. Math. 271, Amer. Math. Soc., Providence, RI (2001) 169–181 MR
- [27] **T Y Lam**, *The algebraic theory of quadratic forms*, W A Benjamin, Reading, MA (1973) MR

- [28] **T Y Lam**, *Introduction to quadratic forms over fields*, Graduate Studies in Mathematics 67, Amer. Math. Soc., Providence, RI (2005) MR
- [29] **J Lurie**, *Higher topos theory*, Annals of Mathematics Studies 170, Princeton Univ. Press (2009) MR
- [30] **M Mahowald**, *The image of J in the EHP sequence*, Ann. of Math. 116 (1982) 65–112 MR
- [31] **F Morel**, *An introduction to \mathbb{A}^1 -homotopy theory*, from “Contemporary developments in algebraic K -theory” (M Karoubi, A O Kuku, C Pedrini, editors), ICTP Lect. Notes XV, Abdus Salam Int. Cent. Theoret. Phys., Trieste (2004) 357–441 MR
- [32] **F Morel**, *The stable \mathbb{A}^1 -connectivity theorems*, K -Theory 35 (2005) 1–68 MR
- [33] **F Morel**, *\mathbb{A}^1 -algebraic topology over a field*, Lecture Notes in Mathematics 2052, Springer (2012) MR
- [34] **F Morel, V Voevodsky**, *\mathbb{A}^1 -homotopy theory of schemes*, Inst. Hautes Études Sci. Publ. Math. 90 (1999) 45–143 MR
- [35] **E Riehl**, *Categorical homotopy theory*, New Mathematical Monographs 24, Cambridge Univ. Press (2014) MR
- [36] **H Toda**, *Generalized Whitehead products and homotopy groups of spheres*, J. Inst. Polytech. Osaka City Univ. Ser. A. Math. 3 (1952) 43–82 MR
- [37] **H Toda**, *On the double suspension E^2* , J. Inst. Polytech. Osaka City Univ. Ser. A. 7 (1956) 103–145 MR
- [38] **H Toda**, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies 49, Princeton Univ. Press (1962) MR
- [39] **G W Whitehead**, *Elements of homotopy theory*, Graduate Texts in Mathematics 61, Springer (1978) MR
- [40] **K Wickelgren**, *Desuspensions of $S^1 \wedge (\mathbb{P}_{\mathbb{Q}}^1 - \{0, 1, \infty\})$* , Internat. J. Math. 27 (2016) art. id. 1640010 MR
- [41] **J Wu**, *Simplicial objects and homotopy groups*, from “Braids” (A J Berrick, FR Cohen, E Hanbury, Y-L Wong, J Wu, editors), Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap. 19, World Sci., Hackensack, NJ (2010) 31–181 MR

School of Mathematics, Georgia Institute of Technology
Atlanta, GA, United States

Department of Mathematics, University of British Columbia
Vancouver, BC, Canada

kwickelgren3@math.gatech.edu, tbjw@math.ubc.ca

Proposed: Jesper Grodal
Seconded: Haynes R Miller, Mark Behrens

Received: 14 December 2015
Revised: 5 June 2018

