

Nonnegative Ricci curvature, stability at infinity and finite generation of fundamental groups

JIAYIN PAN

We study the fundamental group of an open n -manifold M of nonnegative Ricci curvature. We show that if there is an integer k such that any tangent cone at infinity of the Riemannian universal cover of M is a metric cone whose maximal Euclidean factor has dimension k , then $\pi_1(M)$ is finitely generated. In particular, this confirms the Milnor conjecture for a manifold whose universal cover has Euclidean volume growth and a unique tangent cone at infinity.

53C20, 53C23; 53C21, 57S30

1 Introduction

A longstanding problem in Riemannian geometry is the Milnor conjecture, which was proposed in 1968 [15]. It states that any open n -manifold of nonnegative Ricci curvature has a finitely generated fundamental group. This conjecture remains open.

The Milnor conjecture has been verified under various additional assumptions. For a manifold with Euclidean volume growth, Anderson [1] and Li [12] independently proved that the fundamental group is finite. Sormani showed that the Milnor conjecture holds if the manifold has small linear diameter growth, or linear volume growth [18]. Liu classified open 3-manifolds of $\text{Ric} \geq 0$, which confirms the Milnor conjecture in dimension 3 [13]. Recently, the author has presented a different approach to the Milnor conjecture in dimension 3 [17] with Cheeger–Colding–Naber theory [2; 3; 6].

In this paper, we verify the Milnor conjecture for manifolds with additional conditions on the Riemannian universal cover at infinity. Recall that for any open n -manifold (M, x) of $\text{Ric} \geq 0$, and any sequence $r_i \rightarrow \infty$, passing to a subsequence if necessary, we can consider a tangent cone of M at infinity, which is the Gromov–Hausdorff limit [10] of

$$(r_i^{-1}M, x) \xrightarrow{\text{GH}} (Y, y).$$

In general, a tangent cone of M at infinity may not be unique [3]. By a splitting theorem [2], Y is a metric product $\mathbb{R}^k \times Y'$, where Y' has no lines. Cheeger and Colding showed that when M has Euclidean volume growth, any tangent cone of M at infinity (Y, y) is a metric cone $(\mathbb{R}^k \times C(Z), (0, z))$ of dimension n [2], where $C(Z)$ has $\text{diam}(Z) < \pi$ and the vertex z . However, k may not be unique among all tangent cones of M at infinity [7].

Definition 1.1 Let M be an open n -manifold with $\text{Ric} \geq 0$, and let k be an integer. We say that M is k -Euclidean at infinity if any tangent cone (Y, y) of M at infinity is a metric cone, whose maximal Euclidean factor has dimension k , that is, (Y, y) splits as $(\mathbb{R}^k \times C(Z), (0, z))$, where $C(Z)$ is a metric cone with $\text{diam}(Z) < \pi$ and the vertex z .

We point out that Definition 1.1 indeed is equivalent to a uniform control on the diameter of Z : for any tangent cone $(\mathbb{R}^k \times C(Z), (0, z))$ of M at infinity, $\text{diam}(Z) \leq \pi - \eta(M)$ for some positive constant $\eta(M)$. This follows from the fact that the space of all tangent cones at infinity is compact in the pointed Gromov–Hausdorff topology (see Lemma 3.1 and Remark 3.2).

We state our main result:

Theorem 1.2 Let M be an open n -manifold of $\text{Ric} \geq 0$. If the Riemannian universal cover of M is k -Euclidean at infinity, then $\pi_1(M)$ is finitely generated.

Note that in Theorem 1.2, the k -Euclidean at infinity assumption is on the universal cover. If $k = 0$, then in fact $\pi_1(M)$ is finite (Proposition 2.9).

Under the assumption in Theorem 1.2, the Riemannian universal cover \tilde{M} of M may have different tangent cones at infinity, even with different dimensions. We mention that the condition in Theorem 1.2 can be further weakened; see Remark 5.14.

Corollary 1.3 Let M be an open n -manifold of $\text{Ric} \geq 0$. If the Riemannian universal cover of M has Euclidean volume growth and a unique tangent cone at infinity, then $\pi_1(M)$ is finitely generated.

We explain our approach to Theorem 1.2 as follows. Let \tilde{M} be the Riemannian cover of M . Given any tangent cone of \tilde{M} at infinity, we consider the associated equivariant Gromov–Hausdorff convergence with $\pi_1(M, x)$ -action — see Fukaya and Yamaguchi [8] —

$$(r_i^{-1} \tilde{M}, \tilde{x}, \pi_1(M, x)) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G),$$

where G is a closed subgroup of $\text{Isom}(\tilde{Y})$. We call such a limit space $(\tilde{Y}, \tilde{y}, G)$ an equivariant tangent cone of $(\tilde{M}, \pi_1(M, x))$ at infinity. It is known that G is always a Lie group [4; 6]. If \tilde{Y} is a metric cone $\mathbb{R}^k \times C(Z)$, where $\text{diam}(Z) < \pi$, then $\text{Isom}(\tilde{Y})$ is a product $\text{Isom}(\mathbb{R}^k) \times \text{Isom}(Z)$. Thus the G -action on \tilde{Y} can be naturally projected to the \mathbb{R}^k -factor through $p: G \rightarrow \text{Isom}(\mathbb{R}^k)$ (see Propositions 2.6 and 2.7).

Our main discovery is that when \tilde{M} is k -Euclidean at infinity, there is certain equivariant stability among all the equivariant tangent cones of $(\tilde{M}, \pi_1(M, x))$ at infinity:

Theorem 1.4 *Let M be an open n -manifold of $\text{Ric} \geq 0$. Suppose that $\pi_1(M)$ is abelian and the universal cover \tilde{M} is k -Euclidean at infinity. Then there exist a closed abelian subgroup K of $O(k)$ and an integer $l \in [0, k]$ such that for any equivariant tangent cone $(\tilde{Y}, \tilde{y}, G) = (\mathbb{R}^k \times C(Z), (0, z), G)$ of $(\tilde{M}, \pi_1(M, x))$ at infinity, the projected G -action on the \mathbb{R}^k -factor $(\mathbb{R}^k, 0, p(G))$ satisfies that $p(G) = K \times \mathbb{R}^l$, with K fixing 0 and the subgroup $\{e\} \times \mathbb{R}^l$ acting as translations in \mathbb{R}^k .*

Remark 1.5 Theorem 1.4 can be generalized to nilpotent fundamental groups; see Remark 5.12. We do not pursue the most general statement here, since we do not need this in the present paper.

Theorem 1.2 follows from Theorem 1.4. Suppose that $\pi_1(M, x)$ is not finitely generated; then, without loss of generality, we can assume that $\pi_1(M)$ is abelian; see Wilking [19]. Using the lengths of Gromov's short generators $r_i \rightarrow \infty$ [9], we get an equivariant tangent cone $(\tilde{Y}, \tilde{y}, G)$ of $(\tilde{M}, \pi_1(M, x))$ at infinity, where the G -orbit at \tilde{y} is not connected (see Lemma 2.2). On the other hand, if \tilde{M} is k -Euclidean at infinity, then by Theorem 1.4 the orbit $G \cdot \tilde{y}$ is connected, a contradiction.

We illustrate our approach to Theorem 1.4. Put $\Gamma = \pi_1(M, x)$. Given two equivariant tangent cones of (\tilde{M}, Γ) at infinity $(\tilde{Y}_i, \tilde{y}_i, G_i)$ ($i = 1, 2$), assume that their projected actions $(\mathbb{R}^k, 0, p(G_i))$ are different. We consider the set of all equivariant tangent cones of (\tilde{M}, Γ) at infinity, denoted as $\Omega(\tilde{M}, \Gamma)$. It is known that $\Omega(\tilde{M}, \Gamma)$ is compact and connected in the equivariant Gromov-Hausdorff topology. Consequently, for any $\epsilon > 0$, there are finitely many spaces $(W_j, w_j, H_j) \in \Omega(\tilde{M}, \Gamma)$ ($j = 1, \dots, l$) such that $(W_1, w_1, H_1) = (\tilde{Y}_1, \tilde{y}_1, G_1)$, $(W_l, w_l, H_l) = (\tilde{Y}_2, \tilde{y}_2, G_2)$, and

$$d_{\text{GH}}((W_j, w_j, H_j), (W_{j+1}, w_{j+1}, H_{j+1})) \leq \epsilon$$

for all $j = 1, \dots, l-1$. When \tilde{M} is k -Euclidean at infinity, $W_i = \mathbb{R}^k \times C(Z_i)$ with $\text{diam}(Z_i) \leq \pi - \eta(\tilde{M})$. Under this control, the associated chain of \mathbb{R}^k -factors in W_j

with projected $p(H_j)$ -action $\{(\mathbb{R}^k, 0, p(H_j))\}_{j=1}^l$ form a $\psi(\epsilon)$ -chain, where $\psi(\epsilon)$ is a positive function with $\psi(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

To see a contradiction without involving the complexity in general situation, we restrict to the special case that all $p(H_j)$ -actions fix 0. Then this leads to the following stability of isometric actions on the unit sphere $S^{k-1} \subseteq \mathbb{R}^k$: given S^{k-1} with an effective isometric G -action, there is $\epsilon > 0$ such that if (S^{k-1}, H) is ϵ -close to (S^{k-1}, G) in the equivariant Gromov–Hausdorff topology, then either G and H are conjugate in $O(k)$, or $\dim(H) < \dim(G)$ (see [Theorem 4.1](#) for the precise statement). It turns out this stability result is enough for us to derive a contradiction. For instance, if $p(G)$ is not unique among $\Omega(\tilde{M}, \Gamma)$, we choose $(\tilde{Y}_1, \tilde{y}_1, G_1)$ such that $p(G_1)$ has the minimal dimension among all elements in $\Omega(\tilde{M}, \Gamma)$. Then for all $(\tilde{Y}_2, \tilde{y}_2, G_2)$ with $p(G_2)$ not conjugate to $p(G_1)$, we can form a $\psi(\epsilon)$ -chain $\{(\mathbb{R}^k, 0, p(H_j))\}_j$ as explained above. Using the stability of isometric actions, when $\psi(\epsilon)$ is sufficiently small, we see that all $p(H_j)$ ($j \geq 2$) are conjugate to $p(H_1)$, which leads to a desired contradiction. To deal with the general situation where these $p(H_i)$ -actions may not fix 0, we develop a key technical tool, referred as critical rescaling (see [Section 2](#) for details).

We start with some preliminaries in [Section 2](#). To illustrate the critical rescaling argument used in the proof of [Theorem 1.4](#), we consider a rudimentary version of [Theorem 1.4](#) in [Section 3](#) (see [Proposition 3.5](#)). In [Section 4](#), we prove a stability result on the isometric actions, which is another main ingredient in the proof of [Theorem 1.4](#) as mentioned above. Afterwards, we prove [Theorem 1.4](#) in [Section 5](#).

The author would like to thank Professor Xiaochun Rong for his suggestions and encouragement during the preparation of this paper. Part of the paper was written during the author's visit to Capital Normal University in Beijing. The author would like to thank Capital Normal University for its support and hospitality.

2 Preliminaries

In this section, we supply notions and results that will be used through the rest of this paper.

For a tuple (Y, y, G) , we always assume that (Y, y) is a pointed metric space, and G is a closed subgroup of $\text{Isom}(Y)$ acting effectively on Y . Depending on the context, we use d_{GH} to denote the (pointed) Gromov–Hausdorff distance between two metric

spaces, or the (pointed) equivariant Gromov–Hausdorff distance between two metric spaces with isometric actions.

2.1 Tangent cones at infinity

Let (M, x) be an open n –manifold of $\text{Ric} \geq 0$, and let $(\widetilde{M}, \widetilde{x})$ be its Riemannian universal cover. By path lifting, $\Gamma = \pi_1(M, x)$ acts on \widetilde{M} isometrically, freely and discretely. For any sequence $r_i \rightarrow \infty$, we can pass to a subsequence and obtain (pointed) equivariant Gromov–Hausdorff convergence [8]:

$$\begin{array}{ccc} (r_i^{-1}\widetilde{M}, \widetilde{x}, \Gamma) & \xrightarrow{\text{GH}} & (\widetilde{Y}, \widetilde{y}, G) \\ \downarrow \pi & & \downarrow \pi \\ (r_i^{-1}M, x) & \xrightarrow{\text{GH}} & (Y = \widetilde{Y}/G, y) \end{array}$$

where G is a closed subgroup of $\text{Isom}(\widetilde{Y})$, the isometry group of \widetilde{Y} . We call the limit space $(\widetilde{Y}, \widetilde{y}, G)$ an equivariant tangent cone of (\widetilde{M}, Γ) at infinity. $\text{Isom}(\widetilde{Y})$ is a Lie group [4; 6], thus G is a Lie group. For convenience, we introduce the set of all tangent cones of M at infinity,

$$\Omega(M) = \{(Y, y) \mid (Y, y) \text{ is a tangent cone of } M \text{ at infinity}\}.$$

The set $\Omega(M)$ has a natural topology, the pointed Gromov–Hausdorff topology. Similarly, we can consider the set of all equivariant tangent cones of (\widetilde{M}, Γ) at infinity,

$\Omega(\widetilde{M}, \Gamma) = \{(\widetilde{Y}, \widetilde{y}, G) \mid (\widetilde{Y}, \widetilde{y}, G) \text{ is an equivariant tangent cone of } (\widetilde{M}, \Gamma) \text{ at infinity}\}$
endowed with the pointed equivariant Gromov–Hausdorff topology. The following fact serves as an idea behind the proof of Theorem 1.4.

Proposition 2.1 *Let (M, x) be an open n –manifold with $\text{Ric} \geq 0$. Then the set $\Omega(\widetilde{M}, \Gamma)$ is compact and connected in the pointed equivariant Gromov–Hausdorff topology.*

Proposition 2.1 is well known among the experts. We include a proof below for the convenience of readers.

Proof For simplicity, we write $\Omega = \Omega(\widetilde{M}, \Gamma)$. The compactness of Ω follows directly from a standard diagonal argument. Let $\{(\widetilde{Y}_j, \widetilde{y}_j, G_j)\}_j$ be a sequence of elements in Ω , and for each j let $\{s_{i,j}\}_i$ be a sequence of scalars such that

$$(s_{i,j}^{-1}\widetilde{M}, \widetilde{x}, \Gamma) \xrightarrow{\text{GH}} (\widetilde{Y}_j, \widetilde{y}_j, G_j).$$

Passing to a subsequence if necessary, $(\tilde{Y}_j, \tilde{y}_j, G_j)$ converges to $(\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty)$. By a standard diagonal argument, there is a subsequence $i(j)$ such that $s_{i(j),j} \rightarrow \infty$ and

$$(s_{i(j),j}^{-1}\tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty).$$

Thus $(\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty) \in \Omega$.

Next we prove connectedness. Suppose that Ω is not connected; then we can write $\Omega = K_1 \cup K_2$, where K_1 and K_2 are nonempty compact subsets of Ω with $K_1 \cap K_2 = \emptyset$. We choose U_i ($i = 1, 2$) as an open neighborhood of K_i in the pointed equivariant Gromov–Hausdorff topology such that $U_1 \cap U_2 = \emptyset$. We regard

$$\sigma(s) = (s^{-1}\tilde{M}, \tilde{x}, \Gamma), \quad s \geq 1,$$

as a continuous curve in the pointed equivariant Gromov–Hausdorff topology. Note that any element in Ω is an accumulation point of this curve as $s \rightarrow \infty$, thus there is s_0 large such that $\sigma(s) \in U_1 \cup U_2$ for all $s \geq s_0$. Since open sets U_1 and U_2 have empty intersection, $\sigma|_{[s_0, \infty)}$ lies in exactly one of U_1 and U_2 . Consequently, one of K_1 and K_2 must be empty, a contradiction. □

2.2 Short generators

To study whether the fundamental group is finitely generated, we use Gromov’s short generators of $\Gamma = \pi_1(M, x)$ [9]. We say that $\{\gamma_1, \dots, \gamma_i, \dots\}$ is a set of short generators of Γ if

$$d(\gamma_1 \tilde{x}, \tilde{x}) \leq d(\gamma \tilde{x}, \tilde{x}) \quad \text{for all } \gamma \in \Gamma,$$

and for each $i \geq 2$, $\gamma_i \in \Gamma - \langle \gamma_1, \dots, \gamma_{i-1} \rangle$ satisfies that

$$d(\gamma_i \tilde{x}, \tilde{x}) \leq d(\gamma \tilde{x}, \tilde{x}) \quad \text{for all } \gamma \in \Gamma - \langle \gamma_1, \dots, \gamma_{i-1} \rangle,$$

where $\langle \gamma_1, \dots, \gamma_{i-1} \rangle$ is the subgroup generated by $\gamma_1, \dots, \gamma_{i-1}$.

If Γ has infinitely many short generators, then we can use the lengths of short generators to get an equivariant tangent cone of (\tilde{M}, Γ) at infinity. In this way, we find a special element in $\Omega(\tilde{M}, \Gamma)$ whose orbit at \tilde{y} is not connected.

Lemma 2.2 *Let (M, x) be an open n -manifold with $\text{Ric} \geq 0$. Suppose that Γ has infinitely many short generators $\{\gamma_1, \dots, \gamma_i, \dots\}$. Then, in the equivariant tangent cone of $(\tilde{M}, \tilde{x}, \Gamma)$ at infinity*

$$(r_i^{-1}\tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G),$$

the orbit $G \cdot \tilde{y}$ is not connected, where $r_i = d(\gamma_i \tilde{x}, \tilde{x}) \rightarrow \infty$.

Lemma 2.2 follows directly from the definition of short generators and equivariant Gromov–Hausdorff convergence (see [17] for the proof of **Lemma 2.2**).

Another theorem related to the Milnor conjecture is Wilking’s reduction.

Theorem 2.3 [19] *Let M be an open manifold with $\text{Ric} \geq 0$. If $\pi_1(M)$ is not finitely generated, then it contains a nonfinitely generated abelian subgroup.*

2.3 Structures of Ricci limit spaces

Next we recall some results on Ricci limit spaces. We denote $\mathcal{M}(n, 0)$ as the set of all the Gromov–Hausdorff limit spaces coming from some sequence (M_i, x_i) of n –manifolds with $\text{Ric} \geq 0$.

Theorem 2.4 [2] *Let $(X, x) \in \mathcal{M}(n, 0)$ be a Ricci limit space. If X contains a line, then X splits isometrically as $\mathbb{R} \times Y$.*

Theorem 2.5 [2] *Let (M, x) be an open n –manifold of $\text{Ric} \geq 0$. If M has Euclidean volume growth, then any tangent cone of M at infinity (Y, y) is a metric cone $C(Z)$ with vertex y of Hausdorff dimension n . Moreover, $\text{diam}(Z) \leq \pi$.*

When $(Y, y) \in \Omega(M)$ is a metric cone $Y = \mathbb{R}^k \times C(Z)$, we always put $0 \in \mathbb{R}^k$ as the projection of $y \in Y$ to the Euclidean factor.

The metric cone structure and **Theorem 2.4** immediately imply the following properties on its isometries:

Proposition 2.6 *Let $(\mathbb{R}^k \times C(Z), (0, z)) \in \mathcal{M}(n, 0)$ be a metric cone, where $C(Z)$ has vertex z and $\text{diam}(Z) < \pi$. Then, for any isometry g of $C(Z)$, we have $g \cdot (\mathbb{R}^k \times \{z\}) \subseteq \mathbb{R}^k \times \{z\}$.*

Proposition 2.7 *Let $Y = \mathbb{R}^k \times C(Z) \in \mathcal{M}(n, 0)$ be a metric cone with $\text{diam}(Z) < \pi$. Then its isometry group $\text{Isom}(Y)$ splits as $\text{Isom}(\mathbb{R}^k) \times \text{Isom}(Z)$.*

Due to this proposition, for a metric cone $Y = \mathbb{R}^k \times C(Z) \in \mathcal{M}(n, 0)$ with $\text{diam}(Z) < \pi$, there is a natural projection map

$$p: \text{Isom}(Y) \rightarrow \text{Isom}(\mathbb{R}^k).$$

Throughout this paper, we always use p to denote this projection map.

Remark 2.8 For any $v \in \mathbb{R}^k$, the orbit $G \cdot (v, z) = (p(G) \cdot v, z)$ can be naturally identified as $p(G) \cdot v \subseteq \mathbb{R}^k \times \{z\}$, where z is the vertex of $C(Z)$.

[Theorem 1.2](#) directly follows from these preparations and the equivariant stability of [Theorem 1.4](#).

Proof of Theorem 1.2 by assuming Theorem 1.4 Suppose that $\pi_1(M)$ is not finitely generated; then, by [Lemma 2.2](#), there is $(\mathbb{R}^k \times C(Z), (0, z), G) \in \Omega(\tilde{M}, \Gamma)$ such that the orbit $G \cdot (0, z) = (p(G) \cdot 0, z)$ is not connected. This contradicts [Theorem 1.4](#). \square

When $\text{diam}(Z) < \pi$, $C(Z)$ has no lines, and [Proposition 2.6](#) says that any isometry of $C(Z)$ must fix its vertex. This simple observation implies that if \tilde{M} is 0-Euclidean at infinity, then its fundamental group must be finite.

Proposition 2.9 *Let M be an open n -manifold of $\text{Ric} \geq 0$. If \tilde{M} is 0-Euclidean at infinity, then $\pi_1(M)$ is finite.*

Proof Suppose that $\Gamma = \pi_1(M, x)$ is an infinite group; then there are elements $\gamma_i \in \pi_1(M, x)$ with $r_i := d(\gamma_i \tilde{x}, \tilde{x}) \rightarrow \infty$. Passing to a subsequence if necessary, we consider an equivariant tangent cone of (\tilde{M}, Γ) at infinity,

$$(r_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G).$$

By our choice of r_i , there is $g \in G$ such that $d(g \cdot \tilde{y}, \tilde{y}) = 1$. On the other hand, since \tilde{M} is 0-Euclidean at infinity, (\tilde{Y}, \tilde{y}) is a metric cone with no lines and \tilde{y} is the unique vertex. Thus the orbit $G \cdot \tilde{y}$ must be a single point \tilde{y} by [Proposition 2.6](#), a contradiction. \square

Corollary 2.10 *Let (M, x) be an open n -manifold of $\text{Ric} \geq 0$. If its universal cover (\tilde{M}, \tilde{x}) has Euclidean volume growth and nonmaximal diameter growth*

$$\limsup_{R \rightarrow \infty} \frac{\text{diam}(\partial B_R(\tilde{x}))}{R} < 2,$$

then $\pi_1(M, x)$ is a finite group. Consequently, M itself has Euclidean volume growth.

Here we use extrinsic metric on $\text{diam}(\partial B_R(\tilde{x}))$, so we always have

$$\frac{\text{diam}(\partial B_R(\tilde{x}))}{R} \leq 2.$$

3 A critical rescaling argument

In this section, we develop the critical rescaling argument, a key technical tool as mentioned in the introduction, to prove a special case of [Theorem 1.4](#): if there is $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$ such that $p(G)$ is trivial, then for any $(\tilde{W}, \tilde{w}, H) \in \Omega(\tilde{M}, \Gamma)$, $p(H)$ is also trivial (see [Proposition 3.5](#)). The proof of [Theorem 1.4](#) is also modeled on the proofs in this section.

We first show that if M is k -Euclidean at infinity, then for any $Y = \mathbb{R}^k \times C(Z) \in \Omega(M)$, there is a uniform gap between Y and any Ricci limit space splitting off an \mathbb{R}^{k+1} -factor. This is indeed a direct consequence of being k -Euclidean at infinity.

Lemma 3.1 *Let M be an open n -manifold of $\text{Ric} \geq 0$. If M is k -Euclidean at infinity, then there is $\epsilon(M) > 0$ such that for any $(Y, y) \in \Omega(M)$ and any Ricci limit space $\mathbb{R}^l \times X \in \mathcal{M}(n, 0)$ with $l > k$, we have*

$$d_{\text{GH}}((Y, y), (\mathbb{R}^l \times X, (0, x))) \geq \epsilon(M).$$

Proof Suppose the contrary; then we would have a sequence $(Y_i, y_i) \in \Omega(M)$ such that as $i \rightarrow \infty$,

$$d_{\text{GH}}((Y_i, y_i), (\mathbb{R}^{l_i} \times X_i, (0, x_i))) \rightarrow 0,$$

where $\mathbb{R}^{l_i} \times X_i \in \mathcal{M}(n, 0)$ and $k < l_i \leq n$. By precompactness, we can pass to a subsequence and have convergence

$$(\mathbb{R}^{l_i} \times X_i, (0, x_i)) \xrightarrow{\text{GH}} (\mathbb{R}^{l_\infty} \times X_\infty, (0, x_\infty))$$

with an integer $l_\infty > k$. The corresponding subsequence of (Y_i, y_i) has the same limit. By a standard diagonal argument, $(\mathbb{R}^{l_\infty} \times X_\infty, (0, x_\infty))$ is also a tangent cone of M at infinity. This is a contradiction to the assumption that M is k -Euclidean at infinity. \square

Remark 3.2 For a metric cone $C(Z) \in \mathcal{M}(n, 0)$, $C(Z)$ splits off a line if and only if $\text{diam}(Z) = \pi$. From this perspective, [Lemma 3.1](#) implies that if M is k -Euclidean at infinity, then there exists $\eta(M) > 0$ such that for any $\mathbb{R}^k \times C(Z) \in \Omega(M)$, Z has diameter no more than $\pi - \eta(M)$.

Next we prove a gap phenomenon between two classes of group actions on spaces in $\Omega(M)$, which is a key property needed in the critical rescaling argument.

Lemma 3.3 *Let M be an open n -manifold of $\text{Ric} \geq 0$. Suppose that \widetilde{M} is k -Euclidean at infinity. Then there exists a constant $\epsilon(M) > 0$ such that the following holds:*

For two spaces $(\widetilde{Y}_j, \widetilde{y}_j, G_j) \in \Omega(\widetilde{M}, \Gamma)$ with $(\widetilde{Y}_j, \widetilde{y}_j) = (\mathbb{R}^k \times C(Z_j), (0, z_j))$ ($j = 1, 2$), if

- (1) $p(G_1)$ is trivial, and
- (2) there is $g \in G_2$ such that $p(g) \neq e$ and $d(g \cdot \widetilde{y}_2, \widetilde{y}_2) \leq 1$,

then

$$d_{\text{GH}}((\widetilde{Y}_1, \widetilde{y}_1, G_1), (\widetilde{Y}_2, \widetilde{y}_2, G_2)) \geq \epsilon(M).$$

For the Euclidean space \mathbb{R}^k with isometric G -action and a nonidentity element $g \in G$, if $d(g \cdot 0, 0) \leq 1$, then it is obvious that

$$d_{\text{GH}}((\mathbb{R}^k, 0, G), (\mathbb{R}^k, 0, \{e\})) \geq \frac{1}{2}.$$

Let $(\widetilde{Y}_1, \widetilde{y}_1, G_1)$ and $(\widetilde{Y}_2, \widetilde{y}_2, G_2)$ be two spaces in $\Omega(\widetilde{M}, \Gamma)$ as in Lemma 3.3. Roughly speaking, Lemma 3.1 assures that for ϵ sufficiently small, any ϵ -approximation from $(\widetilde{Y}_1, \widetilde{y}_1)$ to $(\widetilde{Y}_2, \widetilde{y}_2)$ cannot map the \mathbb{R}^k -factor to the non-Euclidean cone factor $C(Z_2)$. In other words, an ϵ -approximation map should map the \mathbb{R}^k -factor to the \mathbb{R}^k -factor. Together with the $p(G_j)$ -action on the \mathbb{R}^k -factor, we see that there should be a gap between $(\widetilde{Y}_1, \widetilde{y}_1, G_1)$ and $(\widetilde{Y}_2, \widetilde{y}_2, G_2)$.

Proof of Lemma 3.3 Suppose the contrary; then we have two sequences in $\Omega(M, \Gamma)$, $\{(\widetilde{Y}_{i1}, \widetilde{y}_{i1}, G_{i1})\}$ and $\{(\widetilde{Y}_{i2}, \widetilde{y}_{i2}, G_{i2})\}$, such that

- (1) $p(G_{i1})$ is trivial,
- (2) there is $g_i \in G_{i2}$ such that $p(g_i) \neq e$ and $d(g_i \cdot \widetilde{y}_{i2}, \widetilde{y}_{i2}) \leq 1$,
- (3) $d_{\text{GH}}((\widetilde{Y}_{i1}, \widetilde{y}_{i1}, G_{i1}), (\widetilde{Y}_{i2}, \widetilde{y}_{i2}, G_{i2})) \rightarrow 0$ as $i \rightarrow \infty$.

Passing to some subsequences if necessary, the above two sequences converge to the same limit:

$$(\widetilde{Y}_{i1}, \widetilde{y}_{i1}, G_{i1}) \xrightarrow{\text{GH}} (\widetilde{Y}_\infty, \widetilde{y}_\infty, G_\infty), \quad (\widetilde{Y}_{i2}, \widetilde{y}_{i2}, G_{i2}) \xrightarrow{\text{GH}} (\widetilde{Y}_\infty, \widetilde{y}_\infty, G_\infty),$$

with $(\widetilde{Y}_\infty, \widetilde{y}_\infty) = (\mathbb{R}^k \times C(Z_\infty), (0, z_\infty))$. By Lemma 3.1, $C(Z_\infty)$ does not split off any line, and thus

$$\begin{aligned} (\mathbb{R}^k \times \{z_{i1}\}, \widetilde{y}_{i1}, p(G_{i1})) &\xrightarrow{\text{GH}} (\mathbb{R}^k \times \{z_\infty\}, \widetilde{y}_\infty, p(G_\infty)), \\ (\mathbb{R}^k \times \{z_{i2}\}, \widetilde{y}_{i2}, p(G_{i2})) &\xrightarrow{\text{GH}} (\mathbb{R}^k \times \{z_\infty\}, \widetilde{y}_\infty, p(G_\infty)); \end{aligned}$$

see [Remark 2.8](#). From the first sequence, we see that $p(G_\infty)$ is trivial because $p(G_{i1}) = \{e\}$. On the other hand, $p(G_{i2})$ contains an element β_i with $d_{\mathbb{R}^k}(\beta_i \cdot 0, 0) \leq 1$. These β_i subconverge to some element $\beta_\infty \in G_\infty$ with $d(\beta_\infty \cdot 0, 0) \leq 1$. If $\beta_\infty \neq e$, then $p(G_\infty)$ is nontrivial. If $\beta_\infty = e$, then we consider the subgroup $H_i = \langle \beta_i \rangle$. The sequence of subgroups H_i subconverges to some nontrivial subgroup H_∞ of $p(G_\infty)$ because $D_1(H_i) \geq \frac{1}{20}$, where $D_1(H_i)$ is the displacement of H_i on $B_1(0) \subseteq \mathbb{R}^k$. In either case, $p(G_\infty)$ is nontrivial, a contradiction. \square

Remark 3.4 The gap in [Lemma 3.1](#) plays a key role in the above proof; it guarantees that symmetries on the non-Euclidean cone factor and on the Euclidean factor cannot interchange. If there is no gap between the non-Euclidean cone factor $C(Z)$ and spaces splitting off lines, then [Lemma 3.3](#) fails. As an example, we construct a continuous family of metric cones (Y_t, y_t, G_t) ($-\delta \leq t \leq \delta$) such that $Y_t = \mathbb{R}^2 \times C(Z_t)$, where $\text{diam}(Z_t) \leq \pi$. As $t \rightarrow 0$,

$$d_{\text{GH}}((\mathbb{R}^2 \times \{z_{-t}\}, (0, z_{-t}), p(G_{-t})), (\mathbb{R}^2 \times z_t, (0, z_t), p(G_t))) \rightarrow 0.$$

For $|t| < \delta$ small, we put $Y_t = \mathbb{R}^2 \times C(S_t^1)$, where S_t^1 is the round circle of diameter $\pi - |t|$. When $t = 0$, then $Y_t = \mathbb{R}^4$. Next we define G_t -action on Y_t . For $t > 0$, $G_t = S^1$ acts as rotations on the $C(S_t^1)$ -factor, while for $t \leq 0$, $G_t = S^1$ acts as rotations about the origin on the \mathbb{R}^2 -factor. It is clear that (Y_t, y_t, G_t) is a continuous path in the equivariant Gromov-Hausdorff topology. However, $p(G_t)$ is trivial for $t > 0$ while $p(G_t) = S^1$ for $t < 0$; they cannot be arbitrarily close as $t \rightarrow 0$.

We are ready to prove the following rudimentary version of [Theorem 1.4](#):

Proposition 3.5 *Let (M, x) be an open n -manifold of $\text{Ric} \geq 0$, whose universal cover is k -Euclidean at infinity. If there is $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$ such that $p(G)$ is trivial, then for any space $(\tilde{W}, \tilde{w}, H) \in \Omega(\tilde{M}, \Gamma)$, $p(H)$ is also trivial.*

Proof We argue by contradiction. Suppose that there are $r_i \rightarrow \infty$ and $s_i \rightarrow \infty$ such that

$$(r_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (s_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}_2, \tilde{y}_2, G_2),$$

where $p(G_1)$ is trivial but $p(G_2)$ is not. Scaling down the sequence s_i^{-1} by a constant if necessary, we assume that there is $g \in G_2$ such that $p(g_2) \neq e$ and $d(g \cdot \tilde{y}_2, \tilde{y}_2) \leq 1$. We pass to a subsequence and assume that $t_i := s_i^{-1}/r_i^{-1} \rightarrow \infty$. This enables us to regard the above first sequence as a rescaling of the second one. Put

$$(N_i, q_i, \Gamma_i) = (s_i^{-1} \tilde{M}, \tilde{x}, \Gamma).$$

In this way, we can rewrite these two convergent sequences as (with $t_i \rightarrow \infty$)

$$(N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (t_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}_2, \tilde{y}_2, G_2).$$

We look for a contradiction in some intermediate rescaling sequence. For each i , we define a set of scales

$$L_i := \left\{ 1 \leq l \leq t_i \mid d_{\text{GH}}((lN_i, q_i, \Gamma_i), (W, w, H)) \leq \frac{1}{3}\epsilon \text{ for some } (W, w, H) \in \Omega(\tilde{M}, \Gamma) \right. \\ \left. \text{such that } H \text{ has some } h \text{ with } p(h) \neq e \text{ and } d(h \cdot \tilde{w}, \tilde{w}) \leq 1 \right\},$$

where $\epsilon = \epsilon(M) > 0$ is the constant in [Lemma 3.3](#). It is clear that $t_i \in L_i$ for all i large, thus L_i is nonempty. We choose $l_i \in L_i$ such that $\inf L_i \leq l_i \leq \inf L_i + 1/i$. We regard this l_i as the critical rescaling sequence.

Claim 1 $l_i \rightarrow \infty$.

Suppose that l_i subconverges to $C < \infty$; then, for this subsequence, we can pass to a subsequence again and obtain the convergence

$$(l_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (C \cdot \tilde{Y}_1, \tilde{y}_1, G_1).$$

Since $l_i \in L_i$, by definition of L_i and the above convergence, we conclude that

$$d_{\text{GH}}((C \cdot \tilde{Y}_1, \tilde{y}_1, G_1), (W, w, H)) \leq \frac{1}{2}\epsilon$$

for some space (W, w, H) such that there is $h \in H$ with $p(h) \neq e$ and $d(h \cdot \tilde{w}, \tilde{w}) \leq 1$. On the other hand, on $(C \cdot \tilde{Y}_1, \tilde{y}_1, G_1)$, $p(G_1)$ is trivial. This is a contradiction to the choice of ϵ and [Lemma 3.3](#). Hence [Claim 1](#) is true.

Next we consider the convergence

$$(l_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}', \tilde{y}', G') \in \Omega(\tilde{M}, \Gamma).$$

We will derive a contradiction by ruling out all the possibilities of $p(G')$ -action on the \mathbb{R}^k -factor of \tilde{Y}' .

Claim 2 $p(G')$ is nontrivial.

For each i , because $l_i \in L_i$, we know that

$$d_{\text{GH}}((l_i N_i, q_i, \Gamma_i), (W_i, w_i, K_i)) \leq \frac{1}{3}\epsilon$$

for some $(W_i, w_i) \in \Omega(\widetilde{M})$ with K_i -action such that there is $k_i \in K_i$ with $p(k_i) \neq e$ and $d(k_i \cdot \tilde{w}_i, \tilde{w}_i) \leq 1$. Since $(l_i N_i, q_i, \Gamma_i)$ converges to $(\tilde{Y}', \tilde{y}', G')$, the limit space satisfies

$$d_{\text{GH}}((\tilde{Y}', \tilde{y}', G'), (W_i, w_i, K_i)) \leq \frac{1}{2}\epsilon$$

for i large. By Lemma 3.3, $p(G')$ is nontrivial.

By Claim 2, there is some $g' \in G'$ such that $p(g') \neq e$. We put $d := d(g' \cdot \tilde{y}', \tilde{y}')$. If $d \leq 1$, we consider the scaling sequence $\frac{1}{2}l_i$:

$$(\tfrac{1}{2}l_i \cdot N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tfrac{1}{2} \cdot \tilde{Y}', \tilde{y}', G').$$

Note that on $(\frac{1}{2} \cdot \tilde{Y}', \tilde{y}', G')$, there is some element $g' \in G'$ with $p(g') \neq e$ and $d(g' \cdot \tilde{y}', \tilde{y}') \leq \frac{1}{2}$. This shows that $\frac{1}{2}l_i \in L_i$ for i large, which is a contradiction to our choice of l_i with $\inf L_i \leq l_i \leq \inf L_i + 1/i$. If $d > 1$, then we consider the scaling sequence $l_i/(2d)$:

$$\left(\frac{l_i}{2d} \cdot N_i, q_i, \Gamma_i\right) \xrightarrow{\text{GH}} \left(\frac{1}{2d} \cdot \tilde{Y}, \tilde{y}, H'\right).$$

and a similar contradiction would arise because $l_i/(2d) \in L_i$. In any case, we see a contradiction. This completes the proof. □

Remark 3.6 In the above proof, when defining L_i , we include all the contradictory H -actions with $p(H) \neq \{e\}$ and a constraint on its displacement. In particular, this allows $p(H)$ to be different from $p(G_2)$. Doing so is necessary. For example, if $p(G_2) = \mathbb{Z}$ and we require $p(H) = \mathbb{Z}$ when defining L_i , then the intermediate sequence

$$(l_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}', \tilde{y}', G')$$

may have a limit group with $p(G') = \mathbb{Z}_p$, where p is a large integer. This \mathbb{Z}_p acts on \mathbb{R}^k as rotations on a plane about a point far away from 0, so that $(\mathbb{R}^k, 0, \mathbb{Z}_p)$ and $(\mathbb{R}^k, 0, \mathbb{Z})$ are very close. In this situation, one cannot derive a contradiction by dividing l_i by a constant.

To conclude this section, we use Proposition 3.5 to prove Corollary 1.3 in dimension 3.

Corollary 3.7 *Let M be an open 3-manifold with $\text{Ric} \geq 0$. Suppose that the universal cover \widetilde{M} has Euclidean volume growth and the unique tangent cone at infinity. Then $\pi_1(M)$ is finitely generated.*

Proof We argue by contradiction. Suppose that Γ has infinitely many short generators $\{\gamma_1, \dots, \gamma_i, \dots\}$. Let $r_i = d(\gamma_i \cdot \tilde{x}, \tilde{x}) \rightarrow \infty$. Passing to a subsequence, we consider an equivariant tangent cone of (\tilde{M}, Γ) at infinity,

$$(r_{i(j)}^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G).$$

We know that \tilde{Y} is a metric cone of Hausdorff dimension 3 (Theorem 2.5), and the orbit $G \cdot \tilde{y}$ is not connected (Lemma 2.2). By Theorem 2.3, we assume that Γ is abelian.

\tilde{Y} is isometric to $\mathbb{R}^k \times C(Z)$ with vertex $\tilde{y} = (0, z)$, where $k \in \{0, 1, 2, 3\}$, $C(Z)$ has vertex z and $\text{diam}(Z) < \pi$. If $k = 0$, then Γ is finite (Proposition 2.9). If $k = 3$, then $\tilde{Y} = \mathbb{R}^3$, and consequently \tilde{M} is isometric to \mathbb{R}^3 [5]. If $k = 2$, then \tilde{Y} is actually isometric to \mathbb{R}^3 according to the fact that the singular set of \tilde{Y} has codimension at least 2 [3]. It remains to handle the case $\tilde{Y} = \mathbb{R} \times C(Z)$. Since the nonconnected orbit $G \cdot \tilde{y}$ is contained in $\mathbb{R} \times \{z\}$, together with the assumption that G is abelian, we see that the orbit $G \cdot \tilde{y}$ is either a \mathbb{Z} -translation orbit, or a \mathbb{Z}_2 reflection orbit in $\mathbb{R} \times \{z\}$. In either case, $p(G)$ -action is free at 0. For the equivariant tangent cone of $(\tilde{Y}, \tilde{y}, G)$ at \tilde{y} ($j \rightarrow \infty$),

$$(j \tilde{Y}, \tilde{y}, G) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, H),$$

it is clear that $p(H) = \{e\}$. By a standard diagonal argument, we can find $s_j \rightarrow \infty$ such that

$$(s_j^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, H).$$

Right now there are two equivariant tangent cones of (\tilde{M}, Γ) at infinity, $(\tilde{Y}, \tilde{y}, H)$ and $(\tilde{Y}, \tilde{y}, G)$, with $p(H) = \{e\}$ but $p(G) \neq \{e\}$, a contradiction to Proposition 3.5. \square

4 Stability of isometric actions

In this section, for an isometric G -action on a Riemannian manifold M , we always assume that G is a closed subgroup of $\text{Isom}(M)$. The goal is the following stability result on isometric actions on any compact manifold M , which will be used in the proof of Theorem 1.4 with M being the unit sphere S^{k-1} .

Theorem 4.1 *Let (M, G) be a compact Riemannian manifold with isometric G -action. Then there exists a constant $\epsilon > 0$, depending on (M, G) , such that the following holds:*

For any isometric H –action on M , if

$$d_{\text{GH}}((M, G), (M, H)) \leq \epsilon,$$

then either H –action is conjugate to G –action by an isometry, or $\dim(H) < \dim(G)$.

One may compare [Theorem 4.1](#) with the result below by Grove and Karcher [\[11\]](#):

Theorem 4.2 *Let M a compact Riemannian manifold. Then there exists $\epsilon(M) > 0$ such that for any two isometric G –actions*

$$\mu_1, \mu_2\colon G \times M \rightarrow M$$

with $d_M(\mu_1(g, x), \mu_2(g, x)) \leq \epsilon(M)$ for all $g \in G$ and $x \in M$, these two actions are conjugate by an isometry.

We mention that the stability of group actions can be traced back to Palais [\[16\]](#). He shows that any two C^1 –close G –actions, as diffeomorphisms on M , can be conjugated by a diffeomorphism, where G is a compact Lie group. Grove and Karcher use the center-of-mass technique, and explicitly construct the conjugation map. They also interpret the C^1 –closeness in terms of curvature bounds of M , when one of the actions is by isometries. For our purpose, we restrict our attention to isometric actions only here.

[Theorem 4.1](#) is different from [Theorem 4.2](#) in the following aspects. [Theorem 4.1](#) considers two isometric actions with possibly different groups. For instance, $G = S^1$ and we can take $H = \mathbb{Z}_p \subseteq G$ with large integer p . Even if one assume $G = H$, the closeness of these two actions in the equivariant Gromov–Hausdorff topology is weaker than the pointwise closeness condition in [Theorem 4.2](#). For example, we know that there is a sequence of circle actions on the standard torus $T^2 = S^1 \times S^1$ converging to a T^2 –action,

$$(T^2, S_i^1) \xrightarrow{\text{GH}} (T^2, T^2).$$

Thus, for any $\epsilon > 0$, we can find two different circle actions in the tail of this sequence such that

$$d_{\text{GH}}((T^2, S_j^1), (T^2, S_k^1)) \leq \epsilon,$$

where j and k are sufficiently large. However, these circle actions are not pointwise close. This example also illustrates that the ϵ in [Theorem 4.1](#) has to depend on the G –action.

To prove [Theorem 4.1](#), we recall some facts on equivariant Gromov–Hausdorff convergence [\[8\]](#). Given $(M, H_i) \xrightarrow{\text{GH}} (M, G)$, one can always assume that the identity map on M gives equivariant ϵ_i -approximations for some $\epsilon_i \rightarrow 0$. We endow $\text{Isom}(M)$ with a natural bi-invariant metric d from its action on M ,

$$d_G(g_1, g_2) = \max_{x \in M} d_M(g_1 \cdot x, g_2 \cdot x).$$

Then H_i converges to the limit G with respect to the Hausdorff distance induced by $(\text{Isom}(M), d)$.

In our proof of [Theorem 4.1](#), we use the following results:

Proposition 4.3 [\[11\]](#) *Let $\mu_1, \mu_2: H \rightarrow G$ be two homomorphisms of compact Lie group H into the Lie group G with a bi-invariant Riemannian metric. There exists $\epsilon(G) > 0$ such that if $d(\mu_1(h), \mu_2(h)) \leq \epsilon(G)$ for all $h \in H$, then the subgroups $\mu_1(H)$ and $\mu_2(H)$ are conjugate in G .*

Proposition 4.4 [\[14\]](#) *Let G be a Lie group with left-invariant Riemannian metric. Then there exists a constant $\epsilon(G) > 0$ such that if $\phi: H \rightarrow G$ is a map from a Lie group H to G such that*

$$d(\phi(h_1 h_2), \phi(h_1) \phi(h_2)) \leq \epsilon < \epsilon(G)$$

for all $h_1, h_2 \in H$, then there is a Lie group homomorphism $\bar{\phi}: H \rightarrow G$ with $d(\bar{\phi}(h), \phi(h)) \leq 2\epsilon$ for all $h \in H$.

We call such a map $\phi: H \rightarrow G$ with

$$d(\phi(h_1 h_2), \phi(h_1) \phi(h_2)) \leq \epsilon$$

an ϵ -homomorphism. In practice, we may start with some bi-invariant distance function d on G . We can equip G with a bi-invariant Riemannian metric d_0 (we can do this because G is compact). Note that d and d_0 generate the same topology on the compact group G . Thus, for a sequence of ϵ_i -homomorphisms with respect to d , it is a sequence of ϵ'_i -homomorphisms with respect to d_0 for some $\epsilon'_i \rightarrow 0$. Therefore, we can still apply [Proposition 4.4](#) for i large.

Proof of Theorem 4.1 Let $\{H_i\}$ be a sequence of group actions on M such that $(M, H_i) \xrightarrow{\text{GH}} (M, G)$. We show that if $\dim(H_i) \geq \dim(G)$, then (M, H_i) is conjugate to (M, G) by an isometry for all i large.

As pointed out, for $(M, H_i) \xrightarrow{\text{GH}} (M, G)$, it is equivalent to consider the Hausdorff convergence $H_i \xrightarrow{\text{H}} G$ in $(\text{Isom}(M), d)$, where d is given by

$$d(g_1, g_2) = \max_{x \in M} d_M(g_1 \cdot x, g_2 \cdot x).$$

We know that there is $\epsilon_i \rightarrow 0$ such that $d_H(H_i, G) \leq \epsilon_i$. For each $h \in H_i$, we choose $\phi_i(h)$ as an element in G that is ϵ_i -close to h . This defines a map

$$\phi_i \colon H_i \rightarrow G.$$

It is straightforward to check that ϕ_i is a $3\epsilon_i$ -homomorphism with respect the metric $d|_G$:

$$\begin{aligned} d(\phi_i(h_1h_2), \phi_i(h_1)\phi_i(h_2)) \\ \leq d(\phi_i(h_1h_2), h_1h_2) + d(h_1h_2, h_1\phi_i(h_2)) + d(h_1\phi_i(h_2), \phi_i(h_1)\phi_i(h_2)) \\ \leq 3\epsilon_i \end{aligned}$$

for any $h_1, h_2 \in H_i$. Apply [Proposition 4.4](#); we obtain a sequence of Lie group homomorphisms

$$\bar{\phi}_i \colon H_i \rightarrow G.$$

Claim $\bar{\phi}_i$ is a Lie group isomorphism for all i large.

We first show that $\bar{\phi}_i$ is injective. Suppose that $\ker(\bar{\phi}_i) \neq \{e\}$; then we have a sequence of nontrivial subgroups converging to $\{e\}$,

$$\ker(\bar{\phi}_i) \xrightarrow{\text{H}} \{e\}.$$

However, there exists $\delta > 0$ such that any nontrivial subgroup of $\text{Isom}(M)$ has displacement at least δ on M . This is because $\text{Isom}(M)$ is a Lie group, which cannot have arbitrarily small nontrivial subgroups. Thus $\ker(\bar{\phi}_i) = \{e\}$ for all i large. Recall the assumption that $\dim(H_i) \geq \dim(G)$. Since $\bar{\phi}_i$ is injective, we must have $\dim(H_i) = \dim(G)$. Also note that the image $\bar{\phi}_i(H_i)$ is $\Psi(\epsilon_i)$ -dense in G , thus $\bar{\phi}_i$ must be surjective for i large.

Now H_i has two embeddings into $\text{Isom}(M)$,

$$\iota_i \colon H_i \rightarrow \text{Isom}(M), \quad \bar{\phi}_i \colon H_i \rightarrow G \subseteq \text{Isom}(M),$$

where ι_i is the inclusion map. Note that

$$d(h, \bar{\phi}_i(h)) \leq \Psi(\epsilon_i) \rightarrow 0$$

for all $h \in H_i$ and some constant C . By [Proposition 4.3](#), we conclude that for i large, $G = \bar{\phi}_i(H_i)$ is conjugate to H_i as subgroups in $\text{Isom}(M)$. In other words, there is some isometry $g_i \in \text{Isom}(M)$ such that $g_i^{-1}Gg_i = H_i$. This shows that (M, H_i) and (M, G) are conjugate through the isometry g_i for i large. \square

Remark 4.5 The proof of [Theorem 4.1](#) can be immediately extended to any compact metric space (X, d) whose isometry group $\text{Isom}(X)$ is a Lie group. In particular, for any metric cone $C(Z) \in \mathcal{M}(n, 0)$, since $\text{Isom}(Z)$ is a Lie group [[4](#); [6](#)], we can extend [Theorem 4.1](#) to such a space Z .

5 Proof of equivariant stability at infinity

Without mentioning, we always assume that groups in this section are abelian. We prove [Theorem 1.4](#) in two steps. First we show that for all

$$(\tilde{Y}, \tilde{y}, G) = (\mathbb{R}^k \times C(Z), (0, z), G) \in \Omega(\tilde{M}, \Gamma),$$

the isotropy subgroup of $p(G)$ at 0 is independent of $(\tilde{Y}, \tilde{y}, G)$, and $(\mathbb{R}^k, 0, p(G))$ satisfies property (P) (see [Lemma 5.2](#) below). Secondly, we prove the noncompact factor in $p(G)$ is also independent of $(\tilde{Y}, \tilde{y}, G)$. The proof of each step shares the same structure as [Proposition 3.5](#): we show that there exists a gap between two certain classes of group actions, then choose a critical rescaling to derive a desired contradiction in the corresponding limit space.

Recall that once we specify a point in \mathbb{R}^k as the origin 0, then every element in $\text{Isom}(\mathbb{R}^k) = \mathbb{R}^k \rtimes O(k)$ can be written as (A, v) , where $A \in O(k)$ fixes 0 and $v \in \mathbb{R}^k$. For convenience, we introduce a definition.

Definition 5.1 Let $(\mathbb{R}^k, 0, G)$ be the k -dimensional Euclidean space with an isometric abelian G -action. We say that $(\mathbb{R}^k, 0, G)$ satisfies property (P) if:

(P) For any element $(A, v) \in G$, $(A, 0)$ is also an element of G .

Property (P) has the following consequence:

Lemma 5.2 If $(\mathbb{R}^k, 0, G)$ satisfies property (P), then

- (1) any compact subgroup of G fixes 0;
- (2) G admits decomposition $G = \text{Iso}_0 G \times \mathbb{R}^l \times \mathbb{Z}^m$, and any element in the subgroup $\{e\} \times \mathbb{R}^l \times \mathbb{Z}^m$ is a translation, where $\text{Iso}_0 G$ is the isotropy subgroup of G at 0.

Proof (1) Let K be any compact subgroup of G . Suppose that K does not fix 0 . Then there is $g = (A, v) \in K$ such that

$$0 \neq g \cdot 0 = (A, v) \cdot 0 = v.$$

By assumption, $(A, 0) \in G$. Hence $(A, v) \cdot (A^{-1}, 0) = (I, v)$ is also an element of G . Because G is abelian, $(A, 0)$ and (I, v) commute. This implies that $A \cdot v = v$, and thus $(A, v)^k = (A^k, kv)$ for any integer k . We see that the subgroup generated by (A, v) cannot be contained in any compact group, a contradiction.

(2) This follows from (1) and the structure of abelian Lie groups. \square

It is clear that (2) in [Lemma 5.2](#) is equivalent to property (P).

Remark 5.3 Let M be an open n -manifold of $\text{Ric} \geq 0$. Suppose that

$$(\tilde{Y}, \tilde{y}, G) = (\mathbb{R}^k \times C(Z), (0, z), G) \in \Omega(\tilde{M}, \Gamma)$$

is a metric cone with isometric G -action, where $C(Z)$ has vertex z and $\text{diam}(Z) < \pi$. We do not know any example such that $(\mathbb{R}^k, 0, p(G))$ does not satisfy property (P). However, we can always find ones with property (P) in $\Omega(\tilde{M}, \Gamma)$ by passing to the equivariant tangent cone of $(\tilde{Y}, \tilde{y}, G)$ at \tilde{y} , or at infinity ($j \rightarrow \infty$):

$$(j\tilde{Y}, \tilde{y}, G) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G_{\tilde{y}}), \quad (j^{-1}\tilde{Y}, \tilde{y}, G) \xrightarrow{\text{GH}} (\tilde{Y}, \tilde{y}, G_{\infty}).$$

It is clear that both $(\mathbb{R}^k, 0, p(G_{\tilde{y}}))$ and $(\mathbb{R}^k, 0, p(G_{\infty}))$ satisfy property (P), because they satisfy (2) in [Lemma 5.2](#) (with no \mathbb{Z} -factors).

Remark 5.4 If $(\mathbb{R}^k, 0, G)$ does not satisfy property (P), then there is an element $(A, v) \in G$, but $(A, 0) \notin G$. After blowing down,

$$(j^{-1}\mathbb{R}^k, 0, G, (A, v)) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, G_{\infty}, (A, 0)).$$

Thus $(A, 0) \in G_{\infty}$. Note that $\text{Iso}_0 p(G)$ is preserved as a subgroup of $\text{Iso}_0 p(G_{\infty})$. Hence $\text{Iso}_0 p(G)$ is a proper subgroup of $\text{Iso}_0 p(G_{\infty})$.

We restate [Theorem 1.4](#) in terms of [Definition 5.1](#):

Theorem 5.5 Let M be an open n -manifold, with abelian fundamental group and $\text{Ric} \geq 0$, whose universal cover \tilde{M} is k -Euclidean at infinity. Then there exist a closed abelian subgroup K of $O(k)$ and an integer $l \in [0, k]$ such that for any space $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$, $(\mathbb{R}^k, 0, p(G))$ satisfies property (P) and $p(G) = K \times \mathbb{R}^l$.

For convenience, we introduce a definition:

Definition 5.6 Let (Y_j, y_j) be a metric space with isometric Lie group G_j -action ($j = 1, 2$). We say that (Y_1, y_1, G_1) is equivalent to (Y_2, y_2, G_2) if

$$d_{\text{GH}}((Y_1, y_1, G_1), (Y_2, y_2, G_2)) = 0,$$

or, equivalently, there is an isometry $F: Y_1 \rightarrow Y_2$ with $F(y_1) = y_2$, and a Lie group isomorphism $\psi: G_1 \rightarrow G_2$ such that $F(g_1 \cdot x_1) = \psi(g_1) \cdot F(x_1)$ for any $g_1 \in G$ and $x_1 \in Y_1$.

We first establish a gap phenomenon between two classes of actions with property (P) but different projected isotropy groups.

Lemma 5.7 Let M be an open n -manifold of $\text{Ric} \geq 0$ whose universal cover is k -Euclidean at infinity. Let K be an isometric action on \mathbb{R}^k fixing 0. Then there exists $\epsilon > 0$, depending on M and the K -action on \mathbb{R}^k , such that the following holds:

For any two spaces $(\tilde{Y}_j, \tilde{y}_j, G_j) \in \Omega(\tilde{M}, \Gamma)$ ($j = 1, 2$), if

- (1) $(\mathbb{R}^k, 0, p(G_j))$ satisfies property (P) ($j = 1, 2$),
- (2) $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_1))$ is equivalent to $(\mathbb{R}^k, 0, K)$,
- (3) $\dim(\text{Iso}_0 p(G_2)) \geq \dim(K)$, and
- (4) $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_2))$ is not equivalent to $(\mathbb{R}^k, 0, K)$,

then

$$d_{\text{GH}}((\tilde{Y}_1, \tilde{y}_1, G_1), (\tilde{Y}_2, \tilde{y}_2, G_2)) \geq \epsilon.$$

Proof Suppose that there are two sequences in $\Omega(M)$, $\{(\tilde{Y}_{ij}, \tilde{y}_{ij}, G_{ij})\}_i$ ($j = 1, 2$), such that for all i ,

- (1) $(\mathbb{R}^k, 0, p(G_{ij}))$ satisfies property (P) ($j = 1, 2$);
- (2) $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_{i1}))$ is equivalent to $(\mathbb{R}^k, 0, K)$;
- (3) $\dim(K_i) \geq \dim(K)$ and $(\mathbb{R}^k, 0, K_i)$ is not equivalent to $(\mathbb{R}^k, 0, K)$, where $K_i = \text{Iso}_0 p(G_{i2})$;
- (4) $d_{\text{GH}}((\tilde{Y}_{i1}, \tilde{y}_{i1}, G_{i1}), (\tilde{Y}_{i2}, \tilde{y}_{i2}, G_{i2})) \rightarrow 0$ as $i \rightarrow \infty$.

After passing to some subsequences, this gives convergences

$$(\tilde{Y}_{i1}, \tilde{y}_{i1}, G_{i1}) \xrightarrow{\text{GH}} (\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty), \quad (\tilde{Y}_{i2}, \tilde{y}_{i2}, G_{i2}) \xrightarrow{\text{GH}} (\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty),$$

with $(\tilde{Y}_\infty, \tilde{y}_\infty) = (\mathbb{R}^k \times C(Z_\infty), (0, z_\infty))$, where $\text{diam}(Z_\infty) < \pi$ and z_∞ is the vertex of $C(Z_\infty)$ (Lemma 3.1). Consequently,

$$(\mathbb{R}^k, 0, p(G_{i1})) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, p(G_\infty)), \quad (\mathbb{R}^k, 0, p(G_{i2})) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, p(G_\infty)).$$

Because each $(\mathbb{R}^k, 0, p(G_{ij}))$ satisfies property (P) for all i and j , we conclude that

$$(S^{k-1}, K) \xrightarrow{\text{GH}} (S^{k-1}, K_\infty), \quad (S^{k-1}, K_i) \xrightarrow{\text{GH}} (S^{k-1}, K_\infty),$$

where S^{k-1} is the unit sphere in \mathbb{R}^k and $K_\infty = \text{Iso}_0 p(G_\infty)$. It is obvious that $(S^{k-1}, 0, K_\infty)$ is equivalent to $(S^{k-1}, 0, K)$. By Theorem 4.1, for all i sufficiently large, either $\dim(K_i) < \dim(K)$ or $(S^{k-1}, 0, K_i)$ is equivalent to $(S^{k-1}, 0, K)$. This contradicts the hypotheses (3)–(4) on K_i . \square

Lemma 5.8 *Let M be an open n -manifold of $\text{Ric} \geq 0$ and abelian fundamental group. Suppose that \tilde{M} is k -Euclidean at infinity. Then, for any space $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$, the $p(G)$ -action on $(\mathbb{R}^k, 0, G)$ satisfies property (P). Moreover, $\text{Iso}_0 p(G)$ is independent of $(\tilde{Y}, \tilde{y}, G)$.*

The key to proving Lemma 5.8 is the following lemma:

Lemma 5.9 *Let M be an open n -manifold of $\text{Ric} \geq 0$ and abelian fundamental group. Suppose that \tilde{M} is k -Euclidean at infinity. Then, for any two spaces $(\tilde{Y}_j, \tilde{y}_j, G_j) \in \Omega(\tilde{M}, \Gamma)$ with $(\mathbb{R}^k, 0, p(G_j))$ satisfying property (P) ($j = 1, 2$), $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_1))$ must be equivalent to $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_2))$.*

To prove Lemma 5.9, we introduce an order on the set of all compact abelian Lie groups.

Definition 5.10 For a compact Lie group K , we define $D(K) = (\dim K, \#K/K_0)$. For two compact Lie groups K and H , with $D(K) = (l_1, l_2)$ and $D(H) = (m_1, m_2)$, we say that $D(K) < D(H)$ if $l_1 < m_1$ or if $l_1 = m_1$ and $l_2 < m_2$. We say that $D(K) \leq D(H)$ if $D(K) = D(H)$ or $D(K) < D(H)$.

Proof of Lemma 5.9 We argue by contradiction. Suppose that there are two spaces $(\tilde{Y}_j, \tilde{y}_j, G_j) \in \Omega(\tilde{M}, \Gamma)$ such that $(\mathbb{R}^k, 0, p(G_j))$ satisfies property (P) ($j = 1, 2$), and

$(\mathbb{R}^k, 0, \text{Iso}_0 p(G_1))$ is not equivalent to $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_2))$. Put $K_j = \text{Iso}_0 p(G_j)$. We choose $(\tilde{Y}_1, \tilde{y}_1, G_1)$ so that $D(K_1)$ is the minimal among all $D(K)$ for $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$ with property (P), where $K = \text{Iso}_0 p(G)$. We derive a contradiction by the critical rescaling argument and [Lemma 5.7](#).

Set $D(K_1) = (m_1, m_2)$. Let $r_i \rightarrow \infty$ and $s_i \rightarrow \infty$ be two sequences such that

$$(r_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (s_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \xrightarrow{\text{GH}} (\tilde{Y}_2, \tilde{y}_2, G_2),$$

and $t_i := (s_i^{-1})/(r_i^{-1}) \rightarrow \infty$. Put $(N_i, q_i, \Gamma_i) = (r_i^{-1} \tilde{M}, \tilde{x}, \Gamma)$; then we have

$$(N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (t_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\tilde{Y}_2, \tilde{y}_2, G_2).$$

We know that $(\mathbb{R}^k, 0, p(G_j))$ satisfies property (P) ($j = 1, 2$), $D(K_1) = (m_1, m_2) \leq D(K_2)$, and $(\mathbb{R}^k, 0, K_1)$ is not equivalent to $(\mathbb{R}^k, 0, K_2)$.

For each i , we define a set of scales

$$L_i := \{1 \leq l \leq t_i \mid d_{\text{GH}}((lN_i, q_i, \Gamma_i), (W, w, H)) \leq \frac{1}{3}\epsilon \text{ for some } (W, w, H) \in \Omega(\tilde{M}, \Gamma)$$

such that $(\mathbb{R}^k, 0, p(H))$ satisfies property (P), and, moreover,

$$D(\text{Iso}_0 p(H)) > (m_1, m_2), \text{ or } D(\text{Iso}_0 p(H)) = (m_1, m_2) \\ \text{but } (\mathbb{R}^k, 0, \text{Iso}_0 p(H)) \text{ is not equivalent to } (\mathbb{R}^k, 0, K_1)\}.$$

We choose the above $\epsilon > 0$ as follows: by [Lemma 5.7](#), there is $\epsilon > 0$, depending on M and $(\mathbb{R}^k, 0, K_1)$, such that, for any $(W_j, w_j, H_i) \in \Omega(\tilde{M}, \Gamma)$ ($j = 1, 2$) for which

- (1) $(\mathbb{R}^k, 0, p(H_j))$ satisfies property (P) ($j = 1, 2$),
- (2) $(\mathbb{R}^k, 0, \text{Iso}_0 p(H_1))$ is equivalent to $(\mathbb{R}^k, 0, K_1)$,

if $d_{\text{GH}}((W_1, w_1, H_1), (W_2, w_2, H_2)) \leq \epsilon$, then either $\dim(\text{Iso}_0 p(H_2)) < \dim(K_1)$, or $(\mathbb{R}^k, 0, \text{Iso}_0 p(H_2))$ is equivalent to $(\mathbb{R}^k, 0, K_1)$.

Since $t_i \in L_i$ for i large, we choose $l_i \in L_i$ with $\inf L_i \leq l_i \leq \inf L_i + 1/i$.

Claim 1 $l_i \rightarrow \infty$.

Suppose that $l_i \rightarrow C < \infty$ for some subsequence; then, for this subsequence,

$$(l_i N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (C \cdot \tilde{Y}_1, \tilde{y}_1, G_1).$$

Together with the fact that $l_i \in L_i$, we know that there is some space $(W, w, H) \in \Omega(\tilde{M}, \Gamma)$ with the properties below:

- (1) $(\mathbb{R}^k, 0, p(H))$ satisfies property (P) .
- (2) Either $D(\text{Iso}_0 p(H)) > (m_1, m_2)$, or $D(\text{Iso}_0 p(H)) = (m_1, m_2)$ but the space $(\mathbb{R}^k, 0, \text{Iso}_0 p(H))$ is not equivalent to $(\mathbb{R}^k, 0, K_1)$.
- (3) $d_{\text{GH}}((C \cdot \tilde{Y}_1, \tilde{y}_1, G_1), (W, w, H)) \leq \frac{1}{3}\epsilon$.

Since $(\mathbb{R}^k, 0, p(H))$ satisfies property (P) , we see that $(\mathbb{R}^k, 0, \text{Iso}_0 p(H))$ is equivalent to $(C \cdot \mathbb{R}^k, 0, \text{Iso}_0 p(H))$. By [Lemma 5.7](#) (the choice of ϵ), we conclude that either $\dim(\text{Iso}_0 p(H)) < \dim K_1$, or $(\mathbb{R}^k, 0, \text{Iso}_0 p(H))$ is equivalent to $(\mathbb{R}^k, 0, K_1)$, which is a contradiction to the condition (2) above. We have verified [Claim 1](#).

Passing to a subsequence if necessary, we have convergence

$$(l_i N_i, q_i, \Gamma_i) \overset{\text{GH}}{\longrightarrow} (\tilde{Y}', \tilde{y}', G').$$

To draw a contradiction, the goal is to rule out all the possibilities of $p(G')$ –action.

Claim 2 $D(K') \geq (m_1, m_2)$, where $K' = \text{Iso}_0 p(G')$.

If $D(K') < (m_1, m_2)$, we pass to the equivariant tangent cone of $(\tilde{Y}', \tilde{y}', G')$ at \tilde{y}' . In this way, we have $(\tilde{Y}', \tilde{y}', G'_{\tilde{y}'})$ with $(\mathbb{R}^k, 0, p(G'_{\tilde{y}'}))$ satisfying property (P) (see [Remark 5.3](#)). Note that $(\mathbb{R}^k, 0, \text{Iso}_0 p(G'_{\tilde{y}'}))$ is equivalent to $(\mathbb{R}^k, 0, K')$ and $D(K') < (m_1, m_2)$. We know that this cannot happen due to our choice of $(\tilde{Y}_1, \tilde{y}_1, G_1)$ with the minimal $D(K_1)$.

Claim 3 $(\mathbb{R}^k, 0, p(G'))$ satisfies property (P) , and $D(K') = (m_1, m_2)$.

In fact, we pass to the equivariant tangent cone of $(\tilde{Y}', \tilde{y}', G')$ at infinity:

$$(j^{-1} \tilde{Y}', \tilde{y}', G') \overset{\text{GH}}{\longrightarrow} (\tilde{Y}', \tilde{y}', G'_{\infty}).$$

For this space, $(\mathbb{R}^k, 0, p(G'_{\infty}))$ satisfies property (P) (see [Remark 5.3](#)). Suppose that [Claim 3](#) fails; then $D(\text{Iso}_0 p(G'_{\infty})) > (m_1, m_2)$ (see [Remark 5.4](#)). We choose a large integer J such that

$$d_{\text{GH}}((J^{-1} \tilde{Y}', \tilde{y}', G'), (\tilde{Y}', \tilde{y}', G'_{\infty})) \leq \frac{1}{4}\epsilon.$$

Hence, for all i large, we have

$$d_{\text{GH}}((J^{-1} l_i N_i, q_i, \Gamma_i), (\tilde{Y}', \tilde{y}', G'_{\infty})) \leq \frac{1}{3}\epsilon.$$

This implies that $l_i/J \in L_i$ for all i large, which is a contradiction to our choice of l_i .

Claim 4 $(\mathbb{R}^k, 0, K')$ is equivalent to $(\mathbb{R}^k, 0, K_1)$.

Suppose not; then we consider the sequence $\frac{1}{2}l_i$:

$$(\frac{1}{2}l_i \cdot N_i, q_i, \Gamma_i) \xrightarrow{\text{GH}} (\frac{1}{2} \cdot \tilde{Y}', \tilde{y}', G').$$

Since $(\mathbb{R}^k, 0, p(G'))$ satisfies property (P), $(\frac{1}{2} \cdot \mathbb{R}^k, 0, K')$ is equivalent to $(\mathbb{R}^k, 0, K')$, which is not equivalent to $(\mathbb{R}^k, 0, K_1)$. This means that $\frac{1}{2}l_i \in L_i$ for i large, a contradiction.

This leads to the ultimate contradiction: Because $l_i \in L_i$, there is some space (W, w, H) in $\Omega(\tilde{M}, \Gamma)$ satisfying the conditions (1)–(2) in the proof of [Claim 1](#), and

$$d_{\text{GH}}((\tilde{Y}', \tilde{y}', G'), (W, w, H)) \leq \frac{1}{2}\epsilon.$$

On the other hand, by [Claims 3](#) and [4](#), [Lemma 5.7](#) and the choice of ϵ , (W, w, H) cannot fulfill condition (2) (see the proof of [Claim 1](#)). \square

Proof of [Lemma 5.8](#) With [Lemma 5.9](#), it is enough to show that for any space $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$, $(\mathbb{R}^k, 0, p(G))$ always satisfies property (P). Suppose the contrary, that is, $(\mathbb{R}^k, 0, p(G))$ does not satisfy property (P) for some $(\tilde{Y}, \tilde{y}, G)$ in $\Omega(\tilde{M}, \Gamma)$. We pass to the equivariant tangent cone of $(\tilde{Y}, \tilde{y}, G)$ at \tilde{y} and at infinity, respectively (see [Remark 5.3](#)). We obtain $(\tilde{Y}, \tilde{y}, G_{\tilde{y}})$ and $(\tilde{Y}, \tilde{y}, G_{\infty})$. For these two spaces, $(\mathbb{R}^k, 0, p(G_{\tilde{y}}))$ and $(\mathbb{R}^k, 0, p(G_{\infty}))$ always satisfy property (P). By [Lemma 5.9](#), $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_{\tilde{y}}))$ is equivalent to $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_{\infty}))$.

On the other hand, because $(\mathbb{R}^k, 0, p(G))$ does not satisfy property (P), $\text{Iso}_0 p(G)$ is a proper subgroup of $\text{Iso}_0 p(G_{\infty})$ ([Remark 5.4](#)). Since $\text{Iso}_0 p(G) = \text{Iso}_0 p(G_{\tilde{y}})$, we conclude that $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_{\tilde{y}}))$ and $(\mathbb{R}^k, 0, \text{Iso}_0 p(G_{\infty}))$ cannot be equivalent, a contradiction to [Lemma 5.9](#). \square

[Lemmas 5.2](#) and [5.8](#) imply that there exists a closed subgroup K of $O(k)$ such that for any $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$, $(\mathbb{R}^k, 0, p(G))$ satisfies property (P), and $p(G) = K \times \mathbb{R}^l \times \mathbb{Z}^m$. To finish the proof of [Theorem 5.5](#), we need to show that l is independent of $(\tilde{Y}, \tilde{y}, G)$ and m is always 0.

We prove the following gap lemma on the noncompact factor of $p(G)$, which does not require [Lemma 5.7](#):

Lemma 5.11 *Let M be an open n -manifold of $\text{Ric} \geq 0$. Suppose that \tilde{M} is k -Euclidean at infinity. Then there exists $\epsilon(M) > 0$ such that the following holds:*

For any two spaces $(\tilde{Y}_j, \tilde{y}_j, G_j) \in \Omega(\tilde{M}, \Gamma)$ ($j = 1, 2$), suppose:

- (1) $(\mathbb{R}^k, 0, p(G_j))$ satisfies property (P) ($j = 1, 2$).
- (2) $p(G_1) = \text{Iso}_0 p(G_1) \times \mathbb{R}^l$ (see Lemma 5.2).
- (3) $p(G_2)$ contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup; for this extra \mathbb{Z} subgroup, it has generator γ with $d_{\mathbb{R}^k}(\gamma \cdot 0, 0) \leq 1$.

Then

$$d_{\text{GH}}((\tilde{Y}_1, \tilde{y}_1, G_1), (\tilde{Y}_2, \tilde{y}_2, G_2)) \geq \epsilon(M).$$

Proof We argue by contradiction. Suppose that there are two sequences of spaces in $\Omega(\tilde{M}, \Gamma)$: $\{(\tilde{Y}_{ij}, \tilde{y}_{ij}, G_{ij})\}_i$ ($j = 1, 2$) such that:

- (1) $(\mathbb{R}^k, 0, p(G_{ij}))$ satisfies property (P) ($j = 1, 2$).
- (2) $p(G_{i1}) = K_{i1} \times \mathbb{R}^l$, where $K_{i1} = \text{Iso}_0 p(G_{i1})$.
- (3) $p(G_{i2})$ contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup; for this extra \mathbb{Z} subgroup, it has generator γ_i with $d_{\mathbb{R}^k}(\gamma_i \cdot 0, 0) \leq 1$.
- (4) $d_{\text{GH}}((\tilde{Y}_{i1}, \tilde{y}_{i1}, G_{i1}), (\tilde{Y}_{i2}, \tilde{y}_{i2}, G_{i2})) \rightarrow 0$ as $i \rightarrow \infty$.

This gives the convergence

$$(\tilde{Y}_{i1}, \tilde{y}_{i1}, G_{i1}) \xrightarrow{\text{GH}} (\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty), \quad (\tilde{Y}_{i2}, \tilde{y}_{i2}, G_{i2}) \xrightarrow{\text{GH}} (\tilde{Y}_\infty, \tilde{y}_\infty, G_\infty);$$

thus,

$$(\mathbb{R}^k, 0, p(G_{i1})) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, p(G_\infty)), \quad (\mathbb{R}^k, 0, p(G_{i2})) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, p(G_\infty)).$$

Since $p(G_{i1}) = K_{i1} \times \mathbb{R}^l$ and $(\mathbb{R}^k, 0, p(G_{i1}))$ satisfies property (P), we conclude that $(\mathbb{R}^k, 0, p(G_\infty))$ also satisfies property (P), and $p(G_\infty) = K_\infty \times \mathbb{R}^l$ with K_∞ fixing 0. On the other hand, by hypothesis (3), $p(G_{i2})$ contains a proper closed subgroup $H_i = K_{i2} \times \mathbb{R}^l$ with $K_{i2} = \text{Iso}_0 p(G_{i2})$. Moreover, there is some element $\alpha_i \in p(G_{i2})$ outside H_i such that $d(H_i \cdot 0, \alpha_i \cdot 0) \in (1, 3)$. This yields

$$(\mathbb{R}^k, 0, H_i, \alpha_i) \xrightarrow{\text{GH}} (\mathbb{R}^k, 0, H_\infty, \alpha_\infty),$$

where α_∞ is outside H_∞ with $d(H_\infty \cdot 0, \alpha_\infty \cdot 0) \in (1, 3)$. Therefore, $p(G_\infty)$ also contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup, a contradiction. \square

Proof of Theorem 5.5 By Lemmas 5.2 and 5.8, for any space $(\tilde{Y}, \tilde{y}, G) \in \Omega(\tilde{M}, \Gamma)$, $(\mathbb{R}^k, 0, p(G))$ always satisfies property (P) and $p(G) = K \times \mathbb{R}^l \times \mathbb{Z}^m$, where K is a closed subgroup of $O(k)$ independent of $(\tilde{Y}, \tilde{y}, G)$. It remains to show that l is independent of $(\tilde{Y}, \tilde{y}, G)$ and m is always 0.

First note that if $\Omega(\tilde{M}, \Gamma)$ has a space $(\tilde{Y}, \tilde{y}, G)$ with $p(G) = K \times \mathbb{R}^l \times \mathbb{Z}^m$, then it also contains $(\tilde{Y}, \tilde{y}, G_{\tilde{y}})$ and $(\tilde{Y}, \tilde{y}, G_\infty)$ (see Remark 5.3) with $p(G_{\tilde{y}}) = K \times \mathbb{R}^l$ and $p(G_\infty) = K \times \mathbb{R}^{l+m}$, respectively. We choose $(\tilde{Y}_1, \tilde{y}_1, G_1) \in \Omega(\tilde{M}, \Gamma)$ with $p(G_1) = K \times \mathbb{R}^l$ having the minimal l among all spaces in $\Omega(\tilde{M}, \Gamma)$. Using the critical rescaling argument and Lemma 5.11, we will rule out the situation that $\Omega(\tilde{M}, \Gamma)$ has $(\tilde{Y}_2, \tilde{y}_2, G_2)$ such that $p(G_2)$ contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup.

Let $r_i \rightarrow \infty$ and $s_i \rightarrow \infty$ be two sequences such that

$$(r_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \overset{\text{GH}}{\longrightarrow} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (s_i^{-1} \tilde{M}, \tilde{x}, \Gamma) \overset{\text{GH}}{\longrightarrow} (\tilde{Y}_2, \tilde{y}_2, G_2),$$

and $t_i := (s_i^{-1})/(r_i^{-1}) \rightarrow \infty$. Rescaling s_i^{-1} down by a constant if necessary, we assume that the extra \mathbb{Z} subgroup in $p(G_2)$ has generator γ with $d_{\mathbb{R}^k}(\gamma \cdot 0, 0) \leq 1$. We put $(N_i, q_i, \Gamma_i) = (r_i^{-1} \tilde{M}, \tilde{x}, \Gamma)$; then

$$(N_i, q_i, \Gamma_i) \overset{\text{GH}}{\longrightarrow} (\tilde{Y}_1, \tilde{y}_1, G_1), \quad (t_i N_i, q_i, \Gamma_i) \overset{\text{GH}}{\longrightarrow} (\tilde{Y}_2, \tilde{y}_2, G_2).$$

For each i , we define

$$\begin{aligned} L_i := \{ 1 \leq l \leq t_i \mid d_{\text{GH}}((l N_i, q_i, \Gamma_i), (W, w, H)) \leq \tfrac{1}{3} \epsilon \text{ for some } (W, w, H) \in \Omega(\tilde{M}, \Gamma) \\ \text{such that } p(H) \text{ contains } \mathbb{R}^l \times \mathbb{Z} \text{ as a closed subgroup,} \\ \text{and, moreover, this extra } \mathbb{Z} \text{ subgroup has generator } h \\ \text{with } d_{\mathbb{R}^k}(h \cdot 0, 0) \leq 1. \} \end{aligned}$$

In the above definition, we choose $\epsilon = \epsilon(M) > 0$ as the constant in Lemma 5.11. $t_i \in L_i$ for i large. We choose $l_i \in L_i$ with $\inf L_i \leq l_i \leq \inf L_i + 1/i$.

Claim 1 $l_i \rightarrow \infty$.

If $l_i \rightarrow C$, then

$$(l_i N_i, q_i, \Gamma_i) \overset{\text{GH}}{\longrightarrow} (C \cdot \tilde{Y}_1, \tilde{y}_1, G_1).$$

The projection to the Euclidean factor $(C \cdot \mathbb{R}^k, 0, p(G_1))$ is equivalent to $(\mathbb{R}^k, 0, p(G'))$, because the latter one satisfies property (P) and $p(G') = K \times \mathbb{R}^l$. Since $l_i \in L_i$,

$$d_{\text{GH}}((C \cdot \tilde{Y}_1, \tilde{y}_1, G_1), (W, w, H)) \leq \tfrac{1}{2} \epsilon$$

for some $(W, w, H) \in \Omega(\widetilde{M}, \Gamma)$ with $p(H)$ containing $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup. Moreover, the extra \mathbb{Z} subgroup has generator h with $d_{\mathbb{R}^k}(h \cdot 0, 0) \leq 1$. This is a contradiction to our choice of ϵ and [Lemma 5.11](#).

Next we consider the convergence

$$(l_i N_i, q_i, \Gamma_i) \overset{\text{GH}}{\longrightarrow} (\widetilde{Y}', \widetilde{y}', G').$$

Claim 2 $p(G') = K \times \mathbb{R}^l$.

Indeed, since $(\mathbb{R}^k, 0, p(G'))$ satisfies property (P), we can write $p(G') = K \times \mathbb{R}^{l'} \times \mathbb{Z}^{m'}$. We can also assume that $l' \geq l$ due to the induction assumption. If $l' > l$ or $m' \neq 0$, then $p(G')$ contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup. Consequently, $l_i/d \in L_i$ for some constant $d \geq 2$, which contradicts our choice of l_i . Hence [Claim 2](#) holds.

We derive the desired contradiction: $l_i \in L_i$, so

$$d_{\text{GH}}((\widetilde{Y}', \widetilde{y}', G'), (W, w, H)) \leq \frac{1}{2}\epsilon$$

for some space $(W, w, H) \in \Omega(\widetilde{M}, \Gamma)$, where $p(H)$ contains $\mathbb{R}^l \times \mathbb{Z}$ as a closed subgroup, and the extra \mathbb{Z} subgroup has generator h with $d_{\mathbb{R}^k}(h \cdot 0, 0) \leq 1$, a contradiction to [Lemma 5.11](#). □

We conclude this paper with remarks on some extensions of [Theorems 1.2](#) and [1.4](#). As indicated in the introduction, we do not pursue these extensions because we do not use them in this paper.

Remark 5.12 With some mild additional arguments, one can generalize [Theorem 1.4](#) to any nilpotent fundamental group:

Let M be an open n -manifold, with nilpotent fundamental group $\text{Ric} \geq 0$, whose universal cover \widetilde{M} is k -Euclidean at infinity. Then there exist a closed nilpotent subgroup K of $O(k)$ and an integer $l \in [0, k]$ such that for any space $(\widetilde{Y}, \widetilde{y}, G) \in \Omega(\widetilde{M}, \Gamma)$, $(\mathbb{R}^k, 0, p(G))$ satisfies property (P) and $p(G) = K \times \mathbb{R}^l$.

Remark 5.13 The proof of [Theorem 1.4](#) does not rely the simple-connectedness of \widetilde{M} , nor the fact that $\pi_1(M, x)$ -action is free. Thus [Theorem 1.4](#) can be generalized to any isometric nilpotent G -action on M if M has $\text{Ric}_M \geq 0$ and is k -Euclidean at infinity.

Remark 5.14 Theorems 1.2 and 1.4 remain true if one replace the k -Euclidean at infinity condition by the following: there is k such that any tangent cone of \widetilde{M} at infinity splits as $(\mathbb{R}^k \times X, (0, x))$, where (X, x) satisfies

- (1) X has no lines,
- (2) any isometry of X fixes x .

With this assumption, tangent cones of \widetilde{M} at infinity may not be metric cones nor be polar spaces. Nevertheless, we still have the desired properties on $\text{Isom}(\widetilde{Y})$ for any $\widetilde{Y} = \mathbb{R}^k \times X \in \Omega(\widetilde{M})$ (see Propositions 2.6 and 2.7 and Remark 2.8):

- (1) $\text{Isom}(\mathbb{R}^k \times X) = \text{Isom}(\mathbb{R}^k) \times \text{Isom}(X)$,
- (2) $g \cdot (v, x) = (p(g) \cdot v, x)$ for any $g \in \text{Isom}(\widetilde{Y})$ and $v \in \mathbb{R}^k$, where $p: \text{Isom}(\widetilde{Y}) \rightarrow \text{Isom}(\mathbb{R}^k)$ is the natural projection.

These properties are all that we required in our proof of Theorems 1.2 and 1.4.

Remark 5.15 If M has the unique tangent cone at infinity as a metric cone $(C(Z), z)$, where $C(Z)$ may contain lines, and if G is a nilpotent group acting as isometries on M , then equivariant stability at infinity holds on the entire cone, that is, $\Omega(M, G)$ consists of a single element. More precisely, there exist a closed nilpotent subgroup K and an integer l such that $(C(Z), z, \mathbb{R}^l \times K)$ is the only element in $\Omega(M, G)$, where K fixes z and the subgroup $\{e\} \times \mathbb{R}^l$ acts as translations in the Euclidean factor of $C(Z)$.

This can be proved with Remark 4.5 and a similar argument as in Section 5.

References

- [1] **MT Anderson**, *On the topology of complete manifolds of nonnegative Ricci curvature*, Topology 29 (1990) 41–55 [MR](#)
- [2] **J Cheeger, TH Colding**, *Lower bounds on Ricci curvature and the almost rigidity of warped products*, Ann. of Math. 144 (1996) 189–237 [MR](#)
- [3] **J Cheeger, TH Colding**, *On the structure of spaces with Ricci curvature bounded below, I*, J. Differential Geom. 46 (1997) 406–480 [MR](#)
- [4] **J Cheeger, TH Colding**, *On the structure of spaces with Ricci curvature bounded below, II*, J. Differential Geom. 54 (2000) 13–35 [MR](#)
- [5] **TH Colding**, *Ricci curvature and volume convergence*, Ann. of Math. 145 (1997) 477–501 [MR](#)

- [6] **T H Colding, A Naber**, *Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications*, Ann. of Math. 176 (2012) 1173–1229 [MR](#)
- [7] **T H Colding, A Naber**, *Characterization of tangent cones of noncollapsed limits with lower Ricci bounds and applications*, Geom. Funct. Anal. 23 (2013) 134–148 [MR](#)
- [8] **K Fukaya, T Yamaguchi**, *The fundamental groups of almost non-negatively curved manifolds*, Ann. of Math. 136 (1992) 253–333 [MR](#)
- [9] **M Gromov**, *Almost flat manifolds*, J. Differential Geometry 13 (1978) 231–241 [MR](#)
- [10] **M Gromov**, *Metric structures for Riemannian and non-Riemannian spaces*, Birkhäuser, Boston, MA (2007) [MR](#)
- [11] **K Grove, H Karcher**, *How to conjugate C^1 -close group actions*, Math. Z. 132 (1973) 11–20 [MR](#)
- [12] **P Li**, *Large time behavior of the heat equation on complete manifolds with nonnegative Ricci curvature*, Ann. of Math. 124 (1986) 1–21 [MR](#)
- [13] **G Liu**, *3-Manifolds with nonnegative Ricci curvature*, Invent. Math. 193 (2013) 367–375 [MR](#)
- [14] **M Mazur, X Rong, Y Wang**, *Margulis lemma for compact Lie groups*, Math. Z. 258 (2008) 395–406 [MR](#)
- [15] **J Milnor**, *A note on curvature and fundamental group*, J. Differential Geometry 2 (1968) 1–7 [MR](#)
- [16] **RS Palais**, *Equivalence of nearby differentiable actions of a compact group*, Bull. Amer. Math. Soc. 67 (1961) 362–364 [MR](#)
- [17] **J Pan**, *A proof of Milnor conjecture in dimension 3*, J. Reine Angew. Math. (online publication March 2018)
- [18] **C Sormani**, *Nonnegative Ricci curvature, small linear diameter growth and finite generation of fundamental groups*, J. Differential Geom. 54 (2000) 547–559 [MR](#)
- [19] **B Wilking**, *On fundamental groups of manifolds of nonnegative curvature*, Differential Geom. Appl. 13 (2000) 129–165 [MR](#)

Department of Mathematics, University of California, Santa Barbara
 Santa Barbara, CA, United States

jypan10@gmail.com

Proposed: Tobias H Colding

Seconded: Bruce Kleiner, John Lott

Received: 8 October 2018

Revised: 2 March 2019

