

DR/DZ equivalence conjecture and tautological relations

ALEXANDR BURYAK

JÉRÉMY GUÉRE

PAOLO ROSSI

We present a family of conjectural relations in the tautological ring of the moduli spaces of stable curves which implies the strong double ramification/Dubrovin–Zhang equivalence conjecture introduced by the authors with Dubrovin (Comm. Math. Phys. 363 (2018) 191–260). Our tautological relations have the form of an equality between two different families of tautological classes, only one of which involves the double ramification cycle. We prove that both families behave the same way upon pullback and pushforward with respect to forgetting a marked point. We also prove that our conjectural relations are true in genus 0 and 1 and also when first pushed forward from $\overline{\mathcal{M}}_{g,n+m}$ to $\overline{\mathcal{M}}_{g,n}$ and then restricted to $\mathcal{M}_{g,n}$ for any $g, n, m \geq 0$. Finally we show that, for semisimple CohFTs, the DR/DZ equivalence only depends on a subset of our relations, finite in each genus, which we prove for $g \leq 2$. As an application we find a new formula for the class λ_g as a linear combination of dual trees intersected with kappa- and psi-classes, and we check it for $g \leq 3$.

14H10; 37K10

| | |
|--|------|
| 1. Introduction | 3537 |
| 2. Tautological relations | 3540 |
| 3. DR/DZ equivalence conjecture and the new tautological relations | 3552 |
| 4. Further structure of the relations | 3559 |
| 5. Restricted set of relations | 3581 |
| Appendix. Proof of the restricted genus 2 relations | 3584 |
| References | 3599 |

1 Introduction

A cohomological field theory (CohFT) $c_{g,n}$ is a family of cohomology classes on the moduli spaces $\overline{\mathcal{M}}_{g,n}$ of genus g stable curves with n marked points (parametrized by n tensor copies of a vector space) which satisfy certain compatibility axioms with respect

to the natural morphisms among different moduli spaces. They were introduced by Kontsevich and Manin [20] to axiomatize the properties of Gromov–Witten classes for a given smooth projective variety, but have since then also proved to be a powerful probe for the cohomology and Chow rings of $\overline{\mathcal{M}}_{g,n}$ itself, and their tautological subrings in particular; see Janda, Pandharipande, Pixton and Zvonkine [22; 18; 19].

Recall that the tautological rings $R^*(\overline{\mathcal{M}}_{g,n})$ for $g, n \geq 0$ satisfying $2g - 2 + n > 0$ are the smallest \mathbb{Q} –subalgebras of $H^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ closed under pushforward along the morphisms forgetting marked points and gluing two marked points together to form a node. $R^*(\overline{\mathcal{M}}_{g,n})$ is much smaller than the full cohomology ring, but still has a rich structure and contains most of the natural and geometrically interesting classes. The ring structure of $R^*(\overline{\mathcal{M}}_{g,n})$, however, is not yet completely under control. We know a system of additive generators, the so-called strata algebra, formed by basic classes which are represented by the closure of the loci of curves with fixed dual stable graph intersected with a given monomial in kappa- and psi-classes. The product of basic classes is explicitly described, but the full system of relations is still unknown, although Pixton has found a large set of relations that is conjectured to be complete; see [22].

In this paper we present a new family of conjectural relations in the form of an equality between two families of tautological classes. We denote these classes in $R^*(\overline{\mathcal{M}}_{g,n})$ by A_{d_1, \dots, d_n}^g and B_{d_1, \dots, d_n}^g , where the n integer nonnegative parameters d_1, \dots, d_n satisfy $2g - 1 \leq \sum d_i \leq 3g - 3 + n$. Their precise definition is given in Sections 2.2 and 2.3, respectively, but here we stress that they can be described as two different linear combinations of stable trees with psi-classes at the half-edges and, moreover, for the A –classes only, a double ramification cycle times the Hodge class λ_g is attached at each vertex.

The motivation for this conjecture comes from the study of the double ramification (DR) hierarchy, an integrable system of Hamiltonian PDEs associated to a CohFT and involving the geometry of the DR cycle, introduced by the first author in [1] and further studied by the authors and Dubrovin in [8; 7; 5; 3; 4] (see also Buryak [2] and Rossi [23] for a review). In [3], sharpening a conjecture from [1], it was conjectured that the (logarithm of the) tau-function of (a particular solution of) the DR hierarchy coincides with the reduced potential of the CohFT. The reduced potential is obtained from the full potential, ie the generating series of the intersection numbers of the CohFT with monomials in the psi-classes, by an explicit procedure, also described in [3], which only depends on the potential itself and which ultimately forgets part of the information.

If the CohFT is semisimple (a technical condition on its genus 0 part), the conjecture translates into a statement about the relation between the DR hierarchy and the Dubrovin–Zhang hierarchy, another, more classical, construction associating an integrable system to a semisimple CohFT for which we have the Witten-type result that the (logarithm of the) tau-function of (a special solution of) the DZ hierarchy coincides with the potential of the CohFT.

In this case the strong DR/DZ equivalence conjecture states that the two hierarchies are related by a normal Miura transformation, ie a change of coordinates preserving the tau-structure, and hence acting in particular on the tau-functions. This action on the tau-functions precisely corresponds to the reduction procedure described above for the potential of the CohFT.

As we have seen, the DR/DZ equivalence conjecture is about intersection numbers, not cohomology classes. However, in Section 3 we show how the coefficients of the two involved generating series, the (logarithm of the) DR tau-function and the reduced potential of the CohFT, are the intersection numbers of the CohFT with two different families of cohomology classes. These two families are precisely the A - and B -classes above. So the DR/DZ equivalence conjecture states that the intersection numbers of the A - and B -classes with any (possibly nontautological) CohFT are equal:

$$\int_{\overline{\mathcal{M}}_{g,n}} A_{d_1, \dots, d_n}^g c_{g,n} = \int_{\overline{\mathcal{M}}_{g,n}} B_{d_1, \dots, d_n}^g c_{g,n}.$$

This motivates us to conjecture that the A - and B -classes themselves are equal:

$$A_{d_1, \dots, d_n}^g = B_{d_1, \dots, d_n}^g.$$

In the rest of the paper we work towards the proof of this conjecture. In Section 4 we prove the string and dilaton equations for both A - and B -classes, establishing that their behaviour upon pullback and pushforward along the morphism $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ that forgets the last marked point is the same.

The string equation allows us to prove that the conjecture is true if and only if it is true when all the parameters d_1, \dots, d_n are strictly positive. This in turn yields a full proof of the conjecture in genus 0 and genus 1.

The dilaton equation is used to show that the relations in $R^*(\mathcal{M}_{g,n})$ obtained by pushing forward our conjectural relations from $R^*(\overline{\mathcal{M}}_{g,n+m})$ to $R^*(\overline{\mathcal{M}}_{g,n})$ and then restricting them to $R^*(\mathcal{M}_{g,n})$ are valid. This is what we mean by saying that the conjecture is valid on $\mathcal{M}_{g,n}$.

We then show that our relations imply in particular a new expression for the top Chern class of the Hodge bundle λ_g as a linear combination of basic classes whose dual graph is a tree (with psi- and kappa-classes). No expression of this type for λ_g was known before. We check its validity for $g \leq 3$.

Finally, in Section 5 we show that, for semisimple CohFTs, the DR/DZ equivalence conjecture actually depends on just a subset of our conjectural relations, namely the ones for which $\sum d_i = 2g$ and $d_i > 0$. This means that the number of relations one needs to check is finite in each genus, and equal to the number of partitions of $2g$.

In the appendix we check this finite subset of relations for $g = 2$, thereby proving the strong DR/DZ equivalence conjecture in genus $g \leq 2$ for any semisimple CohFT.

Acknowledgements

We would like to thank Boris Dubrovin, Rahul Pandharipande, Sergey Shadrin and Dimitri Zvonkine for useful discussions.

Buryak received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 797635, was supported by Grant ERC-2012-AdG-320368-MCSK in the group of Rahul Pandharipande at ETH Zürich and Grant RFFI-16-01-00409. Guéré was supported by the Einstein Foundation. Rossi was partially supported by a Chaire CNRS/Enseignement supérieur 2012-2017 grant.

Part of the work was completed during the visit of Guéré and Rossi to the Forschungsinstitut für Mathematik at ETH Zürich in 2017.

2 Tautological relations

In this section we present our conjectural tautological relations.

2.1 Tautological ring of $\overline{\mathcal{M}}_{g,n}$

Here we fix notation concerning tautological cohomology classes on $\overline{\mathcal{M}}_{g,n}$. We will use the notation from [22, Sections 0.2 and 0.3].

Recall that for any stable graph Γ we have the associated moduli space

$$\overline{\mathcal{M}}_{\Gamma} := \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}$$

and the canonical morphism

$$\xi_\Gamma: \overline{\mathcal{M}}_\Gamma \rightarrow \overline{\mathcal{M}}_{g(\Gamma), |L(\Gamma)|}.$$

Recall [22] that given numbers $x_i[v], y[h] \geq 0, i \geq 1, v \in V(\Gamma)$ and $h \in H(\Gamma)$, we can define a basic cohomology class on $\overline{\mathcal{M}}_\Gamma$ by

$$(1) \quad \gamma = \prod_{v \in V(\Gamma)} \prod_{i \geq 1} \kappa_i[v]^{x_i[v]} \cdot \prod_{h \in H(\Gamma)} \psi_h^{y[h]} \in H^*(\overline{\mathcal{M}}_\Gamma, \mathbb{Q}),$$

where $\kappa_i[v]$ is the i^{th} kappa-class on the space $\overline{\mathcal{M}}_{g(v), n(v)}$ and ψ_h is the psi-class on $\overline{\mathcal{M}}_{g(v(h)), n(v(h))}$. A cohomology class on $\overline{\mathcal{M}}_{g,n}$ of the form $\xi_{\Gamma*}(\gamma)$, where Γ is a stable graph of genus g with n legs and γ is a basic class on $\overline{\mathcal{M}}_\Gamma$, will be called a basic tautological class. Denote by $R^*(\overline{\mathcal{M}}_{g,n})$ the subspace of $H^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ spanned by all basic tautological classes. The subspace $R^*(\overline{\mathcal{M}}_{g,n})$ is closed under multiplication and is called the tautological ring of the moduli space of curves. Let

$$R^i(\overline{\mathcal{M}}_{g,n}) := R^*(\overline{\mathcal{M}}_{g,n}) \cap H^{2i}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q}).$$

Denote by $\mathcal{M}_{g,n}^{\text{ct}} \subset \overline{\mathcal{M}}_{g,n}$ the moduli space of curves of compact type and by $\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$ the moduli space of smooth curves. We will use the notation

$$R^i(\mathcal{M}_{g,n}^{\text{ct}}) := R^i(\overline{\mathcal{M}}_{g,n})|_{\mathcal{M}_{g,n}^{\text{ct}}}, \quad R^i(\mathcal{M}_{g,n}) := R^i(\overline{\mathcal{M}}_{g,n})|_{\mathcal{M}_{g,n}}.$$

Linear relations between basic tautological classes are called tautological relations. The class $\xi_{\Gamma*}(1) \in R^{|E(\Gamma)|}(\overline{\mathcal{M}}_{g(\Gamma), |L(\Gamma)|})$ will be called a boundary stratum.

We will represent a basic tautological class $\xi_{\Gamma*}(\gamma)$ on $\overline{\mathcal{M}}_{g(\Gamma), |L(\Gamma)|}$ by a picture of the graph Γ , where we put the monomial $\prod_{i \geq 1} \kappa_i[v]^{x_i[v]}$ next to each vertex v and the power of the psi-class $\psi_h^{y[h]}$ next to each half-edge h . For example, we have the well-known formulas

$$\psi_1 = \begin{array}{c} 1 \\ \diagdown \\ \textcircled{0} \\ \diagup \\ 2 \end{array} \text{---} \begin{array}{c} 3 \\ \diagdown \\ \textcircled{0} \\ \diagup \\ 4 \end{array} \in R^1(\overline{\mathcal{M}}_{0,4}), \quad \lambda_1 = \frac{1}{24} \text{---} \begin{array}{c} \textcircled{0} \\ \textcircled{0} \end{array} \in R^1(\overline{\mathcal{M}}_{1,1}),$$

where we denote by $\lambda_i \in H^{2i}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ the i^{th} Chern class of the Hodge vector bundle over $\overline{\mathcal{M}}_{g,n}$. It is well-known that the class λ_i is tautological (see eg [12]).

Suppose Γ_1 and Γ_2 are two stable graphs, both of genus g and with n legs. They are called isomorphic if there exists a pair $f = (f_1, f_2)$ of set isomorphisms $f_1: V(\Gamma_1) \rightarrow V(\Gamma_2)$ and $f_2: H(\Gamma_1) \rightarrow H(\Gamma_2)$ that preserve all the additional structure of the stable

graphs. Suppose γ_1 and γ_2 are two basic classes on the spaces $\overline{\mathcal{M}}_{\Gamma_1}$ and $\overline{\mathcal{M}}_{\Gamma_2}$, respectively,

$$\gamma_1 = \prod_{v \in V(\Gamma_1)} \prod_{i \geq 1} \kappa_i[v]^{x_{1,i}[v]} \cdot \prod_{h \in H(\Gamma_1)} \psi_h^{y_1[h]},$$

$$\gamma_2 = \prod_{v \in V(\Gamma_2)} \prod_{i \geq 1} \kappa_i[v]^{x_{2,i}[v]} \cdot \prod_{h \in H(\Gamma_2)} \psi_h^{y_2[h]}.$$


We will say that the pairs (Γ_1, γ_1) and (Γ_2, γ_2) are combinatorially equivalent if there exists a pair of maps $f = (f_1, f_2)$ with $f_1: V(\Gamma_1) \rightarrow V(\Gamma_2)$ and $f_2: H(\Gamma_1) \rightarrow H(\Gamma_2)$ that defines an isomorphism between the stable graphs Γ_1 and Γ_2 and also satisfies the properties

$$x_{1,i}[v] = x_{2,i}[f_1(v)] \quad \text{for any } i \geq 1 \text{ and } v \in V(\Gamma_1),$$

$$y_1[h] = y_2[f_2(h)] \quad \text{for any } h \in H(\Gamma_1).$$

Obviously, if the pairs (Γ_1, γ_1) and (Γ_2, γ_2) are combinatorially equivalent, then $\xi_{\Gamma_1*}(\gamma_1) = \xi_{\Gamma_2*}(\gamma_2)$.

Consider the set of stable graphs of genus g with n legs. Suppose I is a subset of $\{1, 2, \dots, n\}$. The symmetric group $S_{|I|}$ acts on our set of stable graphs by permutations of markings from the set I . This gives an $S_{|I|}$ -action on the set of pairs (Γ, γ) , where Γ is a stable graph and γ is a basic class on $\overline{\mathcal{M}}_{\Gamma}$. Let us fix some stable graph Γ and a basic class γ . The sum of the basic tautological classes corresponding to combinatorially nonequivalent pairs in the $S_{|I|}$ -orbit of the pair (Γ, γ) will be represented by the picture corresponding to the class $\xi_{\Gamma*}(\gamma)$, where we erase the labels from the set I . Let us give two examples in order to illustrate this rule:



As another useful example, we can write the topological recursion relations in genus 0 and 1:

$$(2) \quad \psi_1 = \sum_{\substack{i+j=n-3 \\ i \geq 1, j \geq 0}} 1 \cdot \underbrace{\begin{matrix} \text{---} 0 & \text{---} 0 \\ \vdots & \vdots \\ \text{---} & \text{---} \\ \underbrace{\hspace{1cm}}_{i \text{ legs}} & \underbrace{\hspace{1cm}}_{j \text{ legs}} \end{matrix}}_2 \in R^1(\overline{\mathcal{M}}_{0,n}), \quad n \geq 4,$$

$$(3) \quad \psi_1 = \frac{1}{24} \begin{array}{c} \textcircled{0} \\ \vdots \\ \underbrace{\quad\quad\quad}_{n-1 \text{ legs}} \end{array} + \sum_{\substack{i+j=n-1 \\ i \geq 1, j \geq 0}} \begin{array}{c} \textcircled{0} \text{---} \textcircled{1} \\ \vdots \quad \quad \quad \vdots \\ \underbrace{\quad\quad\quad}_{i \text{ legs}} \quad \underbrace{\quad\quad\quad}_{j \text{ legs}} \end{array} \in R^1(\overline{\mathcal{M}}_{1,n}).$$

By stable tree we mean a stable graph Γ with the first Betti number $b_1(\Gamma)$ equal to zero. Suppose Γ is a stable tree. Let

$$H^e(\Gamma) := H(\Gamma) \setminus L(\Gamma).$$

A path in Γ is a sequence of pairwise distinct vertices $v_1, v_2, \dots, v_k \in V(\Gamma)$ such that for any $1 \leq i \leq k - 1$ the vertices v_i and v_{i+1} are connected by an edge. For a vertex $v \in V(\Gamma)$ define a number $r(v)$ by

$$r(v) := 2g(v) - 2 + n(v).$$

Denote by $ST_{g,n}^m$ the set of stable trees of genus g with m vertices and with n legs marked by numbers $1, \dots, n$. For a stable tree $\Gamma \in ST_{g,n}^m$ denote by $l_i(\Gamma)$ the leg in Γ that is marked by i . For a leg $l \in L(\Gamma)$ denote by $1 \leq i(l) \leq n$ its marking.

A stable rooted tree is a pair (Γ, v_1) , where Γ is a stable tree and $v_1 \in V(\Gamma)$. The vertex v_1 is called the root. Denote by $H_+(\Gamma)$ the set of half-edges of Γ that are directed away from the root v_1 . Clearly, $L(\Gamma) \subset H_+(\Gamma)$. Let

$$H_+^e(\Gamma) := H_+(\Gamma) \setminus L(\Gamma).$$

A vertex w is called a descendant of a vertex v if v is on the unique path from the root v_1 to w . Note that according to our definition the vertex v is a descendant of itself. Denote by $\text{Desc}[v]$ the set of all descendants of v . A vertex w is called a direct descendant of v if $w \in \text{Desc}[v]$, $w \neq v$ and w and v are connected by an edge. In this case the vertex v is called the mother of w .

2.2 Double ramification cycle and the definition of the A-class

Consider integers a_1, \dots, a_n such that $a_1 + \dots + a_n = 0$. The double ramification cycle $\text{DR}_g(a_1, \dots, a_n)$ is a cohomology class in $H^{2g}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$. If not all of the a_i are equal to zero, then the restriction $\text{DR}_g(a_1, \dots, a_n)|_{\mathcal{M}_{g,n}}$ can be defined as the Poincaré dual to the locus of pointed smooth curves (C, p_1, \dots, p_n) satisfying $\mathcal{O}_C(\sum_{i=1}^n a_i p_i) \cong \mathcal{O}_C$, and we refer the reader, for example, to [9] for the definition of the double ramification cycle on the whole moduli space $\overline{\mathcal{M}}_{g,n}$. We will often consider the Poincaré dual to the double ramification cycle $\text{DR}_g(a_1, \dots, a_n)$. It is an

element of $H_{2(2g-3+n)}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ and, abusing our notation a little bit, it will also be denoted by $\text{DR}_g(a_1, \dots, a_n)$.

The double ramification cycle $\text{DR}_g(a_1, \dots, a_n)$ is a tautological class on $\overline{\mathcal{M}}_{g,n}$ [13]. A simple explicit formula for the restriction $\text{DR}_g(a_1, \dots, a_n)|_{\mathcal{M}_{g,n}^{\text{ct}}}$ was derived in [16; 21]:

$$(4) \quad \text{DR}_g(a_1, \dots, a_n)|_{\mathcal{M}_{g,n}^{\text{ct}}} = \frac{1}{g!} \left(\sum_{i=1}^n \frac{a_i^2 \psi_i}{2} - \frac{1}{2} \sum_{\substack{I \subset \{1, \dots, n\} \\ |I| \geq 2}} a_I^2 \delta_0^I - \frac{1}{4} \sum_{I \subset \{1, \dots, n\}} \sum_{h=1}^{g-1} a_I^2 \delta_h^I \right)^g,$$

where for a subset $I \subset \{1, 2, \dots, n\}$ and a number $0 \leq h \leq g$ we use the notation

$$a_I := \sum_{i \in I} a_i,$$

$$\delta_h^I := \left(\begin{array}{c} \textcircled{h} \text{---} \textcircled{h'} \\ \vdots \quad \quad \vdots \\ \underbrace{\quad \quad \quad}_I \quad \underbrace{\quad \quad \quad}_{I^c} \end{array} \right) \in R^1(\overline{\mathcal{M}}_{g,n}), \quad I^c := \{1, 2, \dots, n\} \setminus I, \quad h' := g - h.$$

From (4), which is known as Hain’s formula, we see that the class $\text{DR}_g(a_1, \dots, a_n)|_{\mathcal{M}_{g,n}^{\text{ct}}}$ is a polynomial in the variables a_1, \dots, a_n homogeneous of degree $2g$. We obtain from $\lambda_g|_{\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}^{\text{ct}}} = 0$ that the class $\lambda_g \text{DR}_g(a_1, \dots, a_n) \in R^{2g}(\overline{\mathcal{M}}_{g,n})$ is a polynomial in a_1, \dots, a_n homogeneous of degree $2g$. The full double ramification cycle is also polynomial, but not necessarily homogeneous [19].

The following properties of the double ramification cycle will be useful for us. Let $\pi_i: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the forgetful map that forgets the i^{th} marked point. Then

$$\text{DR}_g(a_1, \dots, a_n, 0) = \pi_{n+1}^* \text{DR}_g(a_1, \dots, a_n).$$

Let $\pi: \overline{\mathcal{M}}_{g,n+g} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the forgetful map that forgets the last g marked points. Then we have [9, Example 3.7]

$$(5) \quad \pi_* \text{DR}_g(a_1, \dots, a_{n+g}) = g! a_{n+1}^2 \cdots a_{n+g}^2 [\overline{\mathcal{M}}_{g,n}].$$

It is also useful to remember that (see eg [19])

$$\text{DR}_g(0, 0, \dots, 0) = (-1)^g \lambda_g \in R^g(\overline{\mathcal{M}}_{g,n}).$$

We denote by $\text{DR}_g(a_1, \dots, \tilde{a}_i, \dots, a_n)$ the class $\pi_{i*} \text{DR}_g(a_1, \dots, a_n) \in R^{g-1}(\overline{\mathcal{M}}_{g,n-1})$. Recall the following important divisibility property:

Lemma 2.1 [3] *Let $g, n \geq 1$. Then the polynomial class*

$$\text{DR}_g \left(-\sum a_i, a_1, a_2, \dots, \tilde{a}_n \right) \Big|_{\mathcal{M}_{g,n}^{\text{ct}}} \in R^{g-1}(\mathcal{M}_{g,n}^{\text{ct}})$$

is divisible by a_n^2 .

Consider a stable tree $\Gamma \in \text{ST}_{g,n}^m$ and integers a_1, \dots, a_n such that $a_1 + \dots + a_n = 0$. To each half-edge $h \in H(\Gamma)$ we assign an integer $a(h)$ in such a way that the following conditions hold:

- (a) If $h \in L(\Gamma)$, then $a(h) = a_i(h)$.
- (b) If $h \in H^e(\Gamma)$, then $a(h) + a(t(h)) = 0$.
- (c) For any vertex $v \in V(\Gamma)$, we have $\sum_{h \in H[v]} a(h) = 0$.

Clearly, such a function $a: H(\Gamma) \rightarrow \mathbb{Z}$ exists and is uniquely determined by the numbers a_1, \dots, a_n . For each moduli space $\bar{\mathcal{M}}_{g(v),n(v)}$, $v \in V(\Gamma)$, the numbers $a(h)$ for $h \in H[v]$ define the double ramification cycle

$$\text{DR}_{g(v)}(A_{H[v]}) \in R^{g(v)}(\bar{\mathcal{M}}_{g(v),n(v)}).$$

Here $A_{H[v]}$ denotes the list $a(h_1), \dots, a(h_{n(v)})$, where $\{h_1, \dots, h_{n(v)}\} = H[v]$. If we multiply all these cycles, we get the class

$$\prod_{v \in V(\Gamma)} \text{DR}_{g(v)}(A_{H[v]}) \in H^{2g}(\bar{\mathcal{M}}_\Gamma, \mathbb{Q}).$$

We define a class $\text{DR}_\Gamma(a_1, \dots, a_n) \in R^{g+m-1}(\bar{\mathcal{M}}_{g,n})$ by

$$\text{DR}_\Gamma(a_1, \dots, a_n) := \xi_{\Gamma*} \left(\prod_{v \in V(\Gamma)} \text{DR}_{g(v)}(A_{H[v]}) \right).$$

Clearly, the class

$$\lambda_g \text{DR}_\Gamma(a_1, \dots, a_n) \in R^{2g+m-1}(\bar{\mathcal{M}}_{g,n})$$

is a polynomial in a_1, \dots, a_n homogeneous of degree $2g$.

Suppose now that a_1, \dots, a_n are arbitrary integers and let $a_0 := -\sum_{i=1}^n a_i$. Consider the set of stable trees $\text{ST}_{g,n+1}^m$. It would be convenient for us to assume that the legs of stable trees from $\text{ST}_{g,n+1}^m$ are marked by $0, 1, \dots, n$. Let $\Gamma \in \text{ST}_{g,n+1}^m$ be an arbitrary stable tree. Consider it as a rooted tree with the root $v_1(\Gamma) := v(l_0(\Gamma))$. As above, the

numbers a_0, a_1, \dots, a_n define a function $a: H(\Gamma) \rightarrow \mathbb{Z}$. Define a coefficient $a(\Gamma)$ by

$$a(\Gamma) := \left(\prod_{h \in H_+^e(\Gamma)} a(h) \right) \left(\prod_{v \in V(\Gamma)} \frac{r(v)}{\sum_{\tilde{v} \in \text{Desc}[v]} r(\tilde{v})} \right).$$

Let $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the forgetful map that forgets the first marked point. Define a class $\tilde{A}^{g,m}(a_1, \dots, a_n) \in R^{2g+m-2}(\overline{\mathcal{M}}_{g,n})$ by

$$\tilde{A}^{g,m}(a_1, \dots, a_n) := \sum_{\Gamma \in \text{ST}_{g,n+1}^m} a(\Gamma) \lambda_g \pi_* \text{DR}_\Gamma(a_0, a_1, \dots, a_n).$$

We know that this class is a polynomial in a_1, \dots, a_n homogeneous of degree $2g+m-1$. Note that the expression for the class $\tilde{A}^{g,1}(a_1, \dots, a_n)$ is actually very simple:

$$\tilde{A}^{g,1}(a_1, \dots, a_n) = \lambda_g \text{DR}_g(\tilde{a}_0, a_1, \dots, a_n).$$

Lemma 2.2 *The polynomial class $\tilde{A}^{g,m}(a_1, \dots, a_n)$ is divisible by $\sum_{i=1}^n a_i$.*

Proof If $m = 1$, then the lemma follows from Lemma 2.1. Suppose $m \geq 2$ and $a_0 = -\sum_{i=1}^n a_i = 0$. We have to prove that

$$\tilde{A}^{g,m}(a_1, \dots, a_n) = 0.$$

Consider a stable tree $\Gamma \in \text{ST}_{g,n+1}^m$. If $g(v_1(\Gamma)) \geq 1$, then, again by Lemma 2.1,

$$\lambda_g \pi_* \text{DR}_\Gamma(0, a_1, \dots, a_n) = 0.$$

If $g(v_1(\Gamma)) = 0$, then $\pi_* \text{DR}_\Gamma(0, a_1, \dots, a_n)$, unless $n(v_1(\Gamma)) = 3$. We obtain

$$(6) \quad \tilde{A}^{g,m}(a_1, \dots, a_n) = \sum_{\substack{\Gamma \in \text{ST}_{g,n+1}^m \\ g(v_1(\Gamma))=0 \\ n(v_1(\Gamma))=3}} a(\Gamma) \lambda_g \pi_* \text{DR}_\Gamma(0, a_1, \dots, a_n).$$

Let us define certain maps

$$\text{ST}_{g,n}^{m-1} \rightarrow \{\Gamma \in \text{ST}_{g,n+1}^m \mid g(v_1(\Gamma)) = 0, n(v_1(\Gamma)) = 3\}.$$

Note that we mark the legs of stable trees from $\text{ST}_{g,n}^{m-1}$ by $1, \dots, n$ and the legs of stable trees from $\text{ST}_{g,n+1}^m$ by $0, 1, \dots, n$. Let $\Gamma \in \text{ST}_{g,n}^{m-1}$. Choose a leg $l \in L(\Gamma)$. Suppose that it is marked by the number $1 \leq i \leq n$. Let us attach to the leg l a new vertex of genus 0 with two legs marked by numbers 0 and i . Denote the resulting stable tree by $\Phi_l(\Gamma) \in \text{ST}_{g,n+1}^m$. Similarly, if we choose an edge $e \in E(\Gamma)$, then we can break this edge and insert a genus 0 vertex with one leg marked by 0. Denote

the resulting stable tree by $\Phi_e(\Gamma) \in \text{ST}_{g,n+1}^m$. Using these operations, we can rewrite formula (6) as

$$\tilde{A}^{g,m}(a_1, \dots, a_n) = \sum_{\Gamma \in \text{ST}_{g,n}^{m-1}} \left(\sum_{l \in L(\Gamma)} a(\Phi_l(\Gamma)) + \sum_{e \in E(\Gamma)} a(\Phi_e(\Gamma)) \right) \lambda_g \text{DR}_{\Gamma}(a_1, \dots, a_n).$$

We see that it is sufficient to prove that for any stable tree $\Gamma \in \text{ST}_{g,n}^{m-1}$ we have the identity

$$(7) \quad \sum_{l \in L(\Gamma)} a(\Phi_l(\Gamma)) + \sum_{e \in E(\Gamma)} a(\Phi_e(\Gamma)) = 0.$$

We prove (7) by induction on m . It will be convenient for us to assume that the genus $g(v)$ of a vertex $v \in V(\Gamma)$ can be a rational number such that $2g(v) - 2 + n(v) > 0$. So the total genus $g = \sum_{v \in V(\Gamma)} g(v)$ can also be rational. If $m = 2$, then

$$\sum_{l \in L(\Gamma)} a(\Phi_l(\Gamma)) = \sum_{i=1}^n \frac{-a_i}{2g - 1 + n} = 0.$$

Suppose $m \geq 3$. Choose a vertex $v \in V(\Gamma)$ such that $|H[v] \setminus L[v]| = 1$. Let h be the unique half-edge from the set $H[v] \setminus L[v]$. Write

$$h' := \iota(h), \quad v' := v(h'), \quad r := r(v), \quad r' := r(v'), \quad R := 2g - 2 + n.$$

Denote by e the edge of Γ corresponding the pair of half-edges (h, h') . Let us erase the vertex v together with all half-edges incident to it. Then the half-edge h' becomes a leg. Let us denote it by l' and mark by $n + 1$. Finally, let us increase the genus of the vertex v' by $\frac{r}{2}$. As a result, we get a stable tree from $\text{ST}_{g+r/2, n-|L[v]|+1}^{m-2}$, which we denote by Γ' . Note that the legs of Γ' are marked by the numbers $i(l)$ for $l \in L(\Gamma) \setminus L[v]$ and $n + 1$. We want to apply the induction assumption to the tree Γ' . Naturally, we assign to a leg $l \in L(\Gamma')$ the number $a_{i(l)}$ if $l \neq l'$, and the number $a(h') = \sum_{\tilde{l} \in L[v]} a(\tilde{l})$ if $l = l'$. It is not hard to see that

$$\sum_{l \in L[v]} a(\Phi_l(\Gamma)) = (-a(h')) \frac{rr'}{(R-r)(r+r')} a(\Phi_{l'}(\Gamma')),$$

$$a(\Phi_e(\Gamma)) = a(h') \frac{r'R}{(R-r)(r+r')} a(\Phi_{l'}(\Gamma')).$$

It is also easy to see that for any leg $l \in L(\Gamma')$ with $l \neq l'$ and for any edge $e' \in E(\Gamma')$ we have

$$a(\Phi_l(\Gamma)) = \frac{r'}{r+r'} a(h') a(\Phi_l(\Gamma')), \quad a(\Phi_{e'}(\Gamma)) = \frac{r'}{r+r'} a(h') a(\Phi_{e'}(\Gamma')).$$

Therefore, we obtain

$$\begin{aligned} \sum_{l \in L(\Gamma)} a(\Phi_l(\Gamma)) + \sum_{e' \in E(\Gamma)} a(\Phi_{e'}(\Gamma)) &= \frac{r'}{r+r'} a(h') \left(\sum_{l \in L(\Gamma')} a(\Phi_l(\Gamma')) + \sum_{e' \in E(\Gamma')} a(\Phi_{e'}(\Gamma')) \right) \\ &= 0, \end{aligned}$$

where the last equality follows from the induction assumption. The lemma is proved. \square

The lemma allows us to define a class $A^{g,m}(a_1, \dots, a_n)$ by

$$A^{g,m}(a_1, \dots, a_n) := \frac{1}{\sum a_i} \tilde{A}^{g,m}(a_1, \dots, a_n) \in R^{2g+m-2}(\overline{\mathcal{M}}_{g,n}).$$

It is a polynomial in a_1, \dots, a_n homogeneous of degree $2g + m - 2$.

Definition 2.3 For any $d_1, \dots, d_n \geq 0$ such that $\delta := \sum_{i=1}^n d_i \geq 2g - 1$ we define

$$A^{g}_{d_1, \dots, d_n} := \text{Coef}_{a_1^{d_1} \dots a_n^{d_n}} A^{g, \delta - 2g + 2}(a_1, \dots, a_n) \in R^{\delta}(\overline{\mathcal{M}}_{g,n}).$$

If $\sum d_i = 2g - 1$, then the formula for $A^{g}_{d_1, \dots, d_n}$ becomes particularly simple:

$$A^{g}_{d_1, \dots, d_n} = \text{Coef}_{a_1^{d_1} \dots a_n^{d_n}} \left(\frac{1}{\sum a_i} \lambda_g \text{DR}_g \left(\widetilde{-\sum a_i, a_1, \dots, a_n} \right) \right).$$

2.3 Definition of the B -class and the main conjecture

Let T be a stable rooted tree with at least n legs, where we split the set of legs in two subsets:

- the legs $\sigma_1, \dots, \sigma_n$ corresponding to the markings;
- some extra legs, whose set is denoted by $F(T)$, corresponding to additional marked points that we will eventually forget.

We will never call marking an element of $F(T)$ and let

$$H_+^{\text{em}}(T) := H_+(T) \setminus F(T).$$

There is a natural level function $l: V(T) \rightarrow \mathbb{N}^*$ such that the root is of level 1 and if a vertex v is the mother of a vertex v' , then $l(v') = l(v) + 1$. The total number of levels in T will be denoted by $\text{deg}(T)$ and called the degree of T . It is also convenient to extend the level function to $H_+^{\text{em}}(T)$ by taking $l(h) := k$ if the half-edge h is attached to a vertex of level k . We say that T is complete if the following conditions are satisfied:

- every vertex has at least one of its descendants with level $\text{deg}(T)$;
- all the markings are attached to the vertices of level $\text{deg}(T)$;
- each vertex of level $\text{deg}(T)$ is attached to at least one marking;
- there are no extra legs attached to the root;
- for every vertex except the root there is at least one extra leg attached to it.

For a complete tree T define a power function

$$q: H_+^e(T) \rightarrow \mathbb{N}$$

by requiring that for a half-edge $h \in H_+^e(T)$ there are exactly $q(h) + 1$ extra legs attached to the vertex v which is the direct descendant of h . We say that T is stable if

- for every $1 \leq k \leq \text{deg}(T)$, there is at least one vertex $v \in V(T)$ of level k such that v remains stable once we forget all the extra legs;
- every vertex of genus 0 with exactly one half-edge $h \in H_+^{\text{em}}(T)$ attached to it has exactly $q(h) + 1$ extra legs attached to it;
- every vertex of genus 0 with exactly two half-edges $h_1, h_2 \in H_+^{\text{em}}(T)$ attached to it has exactly $q(h_1) + q(h_2)$ extra legs attached to it.

We say that a stable complete tree T is admissible if for every $1 \leq k < \text{deg}(T)$ we have the condition

$$(8) \quad \sum_{\substack{h \in H_+^e(T) \\ l(h)=k}} q(h) \leq 2 \sum_{\substack{v \in V(T) \\ l(v) \leq k}} g(v) - 2.$$

We denote by $\Omega_{d_1, \dots, d_n}^{B, g}$ the set of pairs (T, q) , where T is an admissible stable complete tree with total genus g and n markings, and $q: H_+^{\text{em}}(T) \rightarrow \mathbb{N}$ is the extension of the

power function from above defined by $q(\sigma_i) := d_i$. We denote by

$$[T, q] := \xi_{T*} \left(\prod_{h \in H_+^{em}(T)} \psi_h^{q(h)} \right) \in R^*(\overline{\mathcal{M}}_{g,n+\#F(T)})$$

and by

$$e: \overline{\mathcal{M}}_{g,n+\#F(T)} \rightarrow \overline{\mathcal{M}}_{g,n}$$

the map forgetting all the extra legs.

Definition 2.4 For any $d_1, \dots, d_n \geq 0$ with $\delta := d_1 + \dots + d_n$, we define

$$(9) \quad B_{d_1, \dots, d_n}^g = \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_* [T, q] \in R^\delta(\overline{\mathcal{M}}_{g,n}).$$

Conjecture 2.5 Suppose $g \geq 0, n \geq 1$ and $2g - 2 + n > 0$. Then for any $d_1, \dots, d_n \geq 0$ such that $\sum d_i \geq 2g - 1$, we have

$$(10) \quad A_{d_1, \dots, d_n}^g = B_{d_1, \dots, d_n}^g.$$

Remark 2.6 Let us show how to express the B -class in terms of basic tautological classes. Let T be a stable complete tree with n markings. For a vertex $v \in V(T)$ denote by $F[v]$ the set of extra legs incident to v and by $H_+^{em}[v]$ the set of half-edges $h \in H_+^{em}(T)$ incident to v . The vertex v will be called strongly stable if it remains stable once we forget all the extra legs. Otherwise, we call it weakly stable. Clearly, the vertex v is weakly stable if and only if $g(v) = 0$ and $|H_+^{em}[v]| = 1$. The set of all strongly stable vertices of T will be denoted by $V^{ss}(T)$.

For a stable complete tree T denote by $st(T)$ the stable rooted tree obtained by forgetting all extra legs of T and then contracting all unstable vertices. Clearly, we can make the identification $V(st(T)) = V^{ss}(T)$ and we also identify the set $H(st(T))$ with the set of half-edges $h \in H(T)$ such that $v(h)$ is strongly stable.

Suppose $\pi: \overline{\mathcal{M}}_{g,n+m} \rightarrow \overline{\mathcal{M}}_{g,n}$ is the forgetful map that forgets the last m marked points. Then for any numbers $c_1, \dots, c_n \geq 0$ we have

$$\pi_*(\psi_1^{c_1} \dots \psi_n^{c_n}) = \sum_{\substack{b_1, \dots, b_n \geq 0 \\ b_i \leq c_i \\ \sum b_i + m = \sum c_i}} \frac{m!}{\prod (c_i - b_i)!} \prod_{i=1}^n \psi_i^{b_i}.$$

Using this formula, it is easy to see that (9) can be rewritten as

$$B_{d_1, \dots, d_n}^g = \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \times \xi_{\text{st}(T)*} \prod_{v \in V(\text{st}(T))} \sum_{\substack{p: H_+[v] \rightarrow \mathbb{Z}_{\geq 0} \\ p(h) \leq q(h) \\ \sum p(h) + |F[v]| = \sum q(h)}} \frac{|F[v]|!}{\prod (q(h) - p(h))!} \prod_{h \in H_+[v]} \psi_h^{p(h)}.$$

Let us immediately present some examples of relations (10). Consider genus 0. Then it is easy to see that for any $d_1, \dots, d_n \geq 0$ we have

$$B_{d_1, \dots, d_n}^0 = \psi_1^{d_1} \dots \psi_n^{d_n}.$$

On the other hand, let us compute, for example, $A_{1,0,0,0}^0$. We have

$$\begin{aligned} & \tilde{A}^{0,3}(a_1, a_2, a_3, a_4) \\ &= \pi_* \left(\sum_{\substack{\{i,j,k,l\}=\{1,2,3,4\} \\ i < j}} \frac{1}{6} (a_i + a_j)(a_i + a_j + a_k) \text{0} \text{---} \text{0} \text{---} \text{0} \begin{matrix} i \\ l \\ k \\ j \end{matrix} \right. \\ & \quad \left. + \sum_{\substack{\{i,j,k,l\}=\{1,2,3,4\} \\ i < j, k < l, i < k}} \frac{1}{3} (a_i + a_j)(a_k + a_l) \text{0} \text{---} \text{0} \begin{matrix} i \\ j \\ k \\ l \end{matrix} \right) \\ &= \frac{1}{3} a^2 \text{0} \text{---} \text{0} = a^2 \begin{matrix} 1 & & 3 \\ & \text{0} \text{---} \text{0} & \\ 2 & & 4 \end{matrix}, \end{aligned}$$

where $a := \sum a_i$. This gives

$$A_{1,0,0,0}^0 = \begin{matrix} 1 & & 3 \\ & \text{0} \text{---} \text{0} & \\ 2 & & 4 \end{matrix},$$

which is indeed equal to $\psi_1 = B_{1,0,0,0}^0$.

Consider genus 1 and the case $n = 1, d_1 = 1$. Then we have

$$A_1^1 = \text{Coef}_a \left(\frac{1}{a} \lambda_1 \pi_* \text{DR}_1(-a, a) \right) = \lambda_1 = \psi_1, \quad B_1^1 = \psi_1.$$

Let us give one more example with $g = 2, n = 1$ and $d_1 = 3$. We compute

$$(11) \quad A_3^2 = \text{Coef}_{a^3} \left(\frac{1}{a} \lambda_2 \text{DR}_2(\widetilde{-a}, a) \right), \quad B_3^2 = \psi_1^3 - \text{0} \text{---} \text{0} \text{---} \text{0} \frac{\psi^2}{a}.$$

Now the relation $A_3^2 = B_3^2$ is not so trivial, and we will prove it in the appendix.

Below we will check that the conjecture is true in genus 0 and 1 for arbitrary d_i , and also in genus 2 in the case $\sum d_i \leq 4$.

3 DR/DZ equivalence conjecture and the new tautological relations

In this section we explain the relation between the above Conjecture 2.5 and the strong double ramification/Dubrovin–Zhang hierarchies equivalence conjecture from [3]. After recalling the main notions, we prove in particular how the first implies the second.

3.1 Dubrovin–Zhang hierarchy

Consider an arbitrary cohomological field theory (CohFT; see [20]) $c_{g,n}: V^{\otimes n} \rightarrow H^{\text{even}}(\overline{\mathcal{M}}_{g,n}, \mathbb{C})$, with V its N –dimensional vector space, e_1, \dots, e_N a basis of V , e_1 the unit and η its symmetric nondegenerate bilinear form. Let $F = F(t_*, \varepsilon)$ denote its potential, ie the generating series of its intersection numbers with monomials in the psi-classes:

$$\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} c_{g,n} \left(\bigotimes_{i=1}^n e_{\alpha_i} \right) \prod_{i=1}^n \psi_i^{d_i}$$

for $2g - 2 + n > 0, 1 \leq \alpha_i \leq N$,

$$F(t_*, \varepsilon) := \sum_{g \geq 0} \varepsilon^{2g} F_g(t_*),$$

where

$$F_g(t_*) := \sum_{\substack{n \geq 0 \\ 2g - 2 + n > 0}} \frac{1}{n!} \sum_{d_1, \dots, d_n \geq 0} \left\langle \prod_{i=1}^n \tau_{d_i}(e_{\alpha_i}) \right\rangle_g \prod_{i=1}^n t_{d_i}^{\alpha_i}.$$

When the cohomological field theory is semisimple, in [10; 6] the authors associate to it an integrable hierarchy of Hamiltonian PDEs. Let $\widehat{\mathcal{A}}_w^{[d]}$ be the degree d part of $\mathbb{C}[[w_*, \varepsilon]]$, where w_k^α for $\alpha = 1, \dots, N$ and $k = 0, 1, 2, \dots$ are formal variables of degree $\deg w_k^\alpha = k$ and $\deg \varepsilon = -1$. Let $\widehat{\Lambda}_w^{[d]}$ be its quotient with respect to constants and the image of the operator $\partial_x = \sum_{k \geq 0} w_{k+1}^\alpha \partial / \partial w_k^\alpha$ (we perform sums over repeated Greek indices here and in what follows) and, if $f \in \widehat{\mathcal{A}}_w^{[d]}$, let \bar{f} denote its equivalence class in $\widehat{\Lambda}_w^{[d]}$. The Dubrovin–Zhang (DZ) hierarchy consists in Hamiltonian densities

$$h_{\alpha,p}^{\text{DZ}} \in \widehat{\Lambda}_w^{[0]}, \quad 1 \leq \alpha \leq N, p \geq -1,$$

with $h_{\alpha,-1}^{\text{DZ}} = \eta_{\alpha\mu} w^\mu$, and a Hamiltonian operator

$$(K^{\text{DZ}})^{\mu\nu} = \sum_{j \geq 0} (K^{\text{DZ}})_j^{\mu\nu} \partial_x^j, \quad (K^{\text{DZ}})_j^{\mu\nu} \in \widehat{\mathcal{A}}_w^{[-j+1]},$$

such that

$$\{\bar{h}_{\alpha,p}^{\text{DZ}}, \bar{h}_{\beta,q}^{\text{DZ}}\}_{K^{\text{DZ}}} := \int \frac{\delta \bar{h}_{\alpha,p}^{\text{DZ}}}{\delta w^\mu} (K^{\text{DZ}})^{\mu\nu} \left(\frac{\delta \bar{h}_{\beta,q}^{\text{DZ}}}{\delta w^\nu} \right) dx = 0, \quad 1 \leq \alpha, \beta \leq N, \quad p, q \geq -1,$$

where we have used the variational derivative $\delta \bar{f} / \delta w^\mu := \sum_{i \geq 0} (-\partial_x)^i \partial f / \partial w_i^\mu$. This guarantees that solutions $w_k^\alpha(x, t_*^*, \varepsilon) = \partial_x^k w^\alpha(x, t_*^*, \varepsilon) \in \mathbb{C}[[x, t_*^*, \varepsilon]]$ exist for the system of Hamiltonian PDEs

$$\frac{\partial}{\partial t_q^\beta} w^\alpha = (K^{\text{DZ}})^{\alpha\nu} \left(\frac{\delta \bar{h}_{\beta,q}^{\text{DZ}}}{\delta w^\nu} \right), \quad 1 \leq \alpha, \beta \leq N, \quad q \geq 0.$$

Notice how this Hamiltonian system in fact only depends on the Hamiltonian functionals $\bar{h}_{\alpha,p}^{\text{DZ}} \in \widehat{\Lambda}_w^{[0]}$ and not on the Hamiltonian densities $h_{\alpha,p}^{\text{DZ}} \in \widehat{\mathcal{A}}_w^{[0]}$. Nonetheless, Dubrovin and Zhang’s construction of specific Hamiltonian densities $h_{\alpha,p}^{\text{DZ}} \in \widehat{\mathcal{A}}_w^{[0]}$ is important because it is a tau-structure (see [3] for details), which implies in particular that, for any solution $w^\alpha(x, t_*^*, \varepsilon) \in \mathbb{C}[[x, t_*^*, \varepsilon]]$, there exists a formal series, called the (logarithm of the) tau-function, $\mathcal{F}(t_*^*, \varepsilon) \in \mathbb{C}[[t_*^*, \varepsilon]]$, such that

$$\left. \frac{\partial h_{\alpha,p-1}^{\text{DZ}}}{\partial t_q^\beta} \right|_{w_*^* = w_*^*(x, t_*^*, \varepsilon)|_{x=0}} = \frac{\partial^3 \mathcal{F}}{\partial t_0^1 \partial t_p^\alpha \partial t_q^\beta}, \quad 1 \leq \alpha, \beta \leq N, \quad p, q \geq 0.$$

An important property of the DZ hierarchy is that the so-called topological solution, ie the solution with the initial condition $(w^{\text{top}})^\alpha(x, t_*^*, \varepsilon)|_{t_*^*=0} = \delta^{\alpha,1} x$, has the potential $F(t_*^*, \varepsilon)$ of the underlying semisimple CohFT as the logarithm of its tau-function:

$$\left. \frac{\partial h_{\alpha,p-1}^{\text{DZ}}}{\partial t_q^\beta} \right|_{w_*^* = (w^{\text{top}})_*^*(x, t_*^*, \varepsilon)|_{x=0}} = \frac{\partial^3 F}{\partial t_0^1 \partial t_p^\alpha \partial t_q^\beta}, \quad 1 \leq \alpha, \beta \leq N, \quad p, q \geq 0.$$

3.2 Double ramification hierarchy

The double ramification (DR) hierarchy — see [1; 3] — is another tau-symmetric hierarchy of Hamiltonian PDEs associated to an arbitrary CohFT, this time even without the requirement of semisimplicity. This time it is the Hamiltonians that are constructed as generating series of certain intersection numbers of the CohFT with psi-classes, the λ_g -class and the double ramification cycle. Written in formal variables \tilde{u}_k^α for

$\alpha = 1, \dots, N$ and $k = 0, 1, 2, \dots$, it consists of differential polynomials

$$h^{\text{DR}} \in \widehat{\mathcal{A}}_{\tilde{u}}^{[0]}, \quad 1 \leq \alpha \leq N, \quad p \geq -1,$$

with $h_{\alpha,-1}^{\text{DR}} = \eta_{\alpha\mu} \tilde{u}^\mu$, and a Hamiltonian operator

$$(K^{\text{DR}})^{\mu\nu} = \sum_{j \geq 0} (K^{\text{DR}})_j^{\mu\nu} \partial_x^j, \quad (K^{\text{DR}})_j^{\mu\nu} \in \widehat{\mathcal{A}}_{\tilde{u}}^{[-j+1]},$$

such that

$$\{\bar{h}_{\alpha,p}^{\text{DR}}, \bar{h}_{\beta,q}^{\text{DR}}\}_{K^{\text{DR}}} := \int \frac{\delta \bar{h}_{\alpha,p}^{\text{DR}}}{\delta \tilde{u}^\mu} (K^{\text{DR}})^{\mu\nu} \left(\frac{\delta \bar{h}_{\beta,q}^{\text{DR}}}{\delta \tilde{u}^\nu} \right) dx = 0, \quad 1 \leq \alpha, \beta \leq N, \quad p, q \geq -1.$$

Like for the DZ hierarchy, the DR Hamiltonian densities $h_{\alpha,p}^{\text{DR}} \in \widehat{\mathcal{A}}_{\tilde{u}}^{[0]}$ form a tau-structure and we can define the DR potential as the (logarithm of the) tau-function of the topological solution, $(\tilde{u}^{\text{top}})^\alpha \in \mathbb{C}[[x, t_*^*, \varepsilon]]$ with $(\tilde{u}^{\text{top}})^\alpha(x, t_*^*, \varepsilon)|_{t_*^*=0} = \delta^{\alpha,1} x$, ie $F^{\text{DR}}(t_*^*, \varepsilon) \in \mathbb{C}[[t_*^*, \varepsilon]]$ satisfies

$$\left. \frac{\partial h_{\alpha,p-1}^{\text{DR}}}{\partial t_q^\beta} \right|_{\tilde{u}_*^* = (\tilde{u}^{\text{top}})_*^*(x, t_*^*, \varepsilon)|_{x=0}} = \frac{\partial^3 F^{\text{DR}}}{\partial t_0^1 \partial t_p^\alpha \partial t_q^\beta}, \quad 1 \leq \alpha, \beta \leq N, \quad p, q \geq 0.$$

Clearly, this equation doesn't determine the function F^{DR} uniquely, but we can additionally require that F^{DR} should satisfy the string and the dilaton equations. Then this fixes the potential F^{DR} completely. We define the DR correlators as the coefficients of the power series $F^{\text{DR}}(t_*^*, \varepsilon)$,

$$F^{\text{DR}}(t_*^*, \varepsilon) =: \sum_{g \geq 0} \varepsilon^{2g} F_g^{\text{DR}}(t_*^*),$$

where

$$F_g^{\text{DR}}(t_*^*) := \sum_{\substack{n \geq 0 \\ 2g-2+n > 0}} \frac{1}{n!} \sum_{d_1, \dots, d_n \geq 0} \left\langle \prod_{i=1}^n \tau_{d_i}(e_{\alpha_i}) \right\rangle_g^{\text{DR}} \prod_{i=1}^n t_{d_i}^{\alpha_i}.$$

3.3 Strong DR/DZ equivalence conjecture

In the effort toward understanding the relation between the DR and DZ hierarchies associated to the same semisimple CohFT, in [3] it was conjectured that a change of coordinates $w^\alpha \mapsto \tilde{u}^\alpha$ existed, transforming one hierarchy into the other and preserving the given tau-structures. A natural family of such changes of coordinates (called normal Miura transformations) has the form

$$(12) \quad \tilde{u}^\alpha(w) = w^\alpha + \eta^{\alpha\mu} \partial_x \{ \mathcal{P}, \bar{h}_{\mu,0}^{\text{DZ}} \}_{K^{\text{DZ}}},$$

where $\mathcal{P} \in \widehat{\mathcal{A}}_w^{[-2]}$ is an arbitrary differential polynomial and

$$\{\mathcal{P}, \bar{h}_{\mu,0}^{\text{DZ}}\}_{K^{\text{DZ}}} = \sum_{k \geq 0} \frac{\partial \mathcal{P}}{\partial w_k^\mu} \partial_x^k \left((K^{\text{DZ}})^{\mu\nu} \left(\frac{\delta \bar{h}_{\mu,0}^{\text{DZ}}}{\delta w^\nu} \right) \right).$$

The effect of such a transformation on the topological tau-function of the DZ hierarchy is

$$F \mapsto F + \mathcal{P}(w_*^*, \varepsilon)|_{w_*^*=(w^{\text{top}})_*(x,t_*^*,\varepsilon)|_{x=0}}.$$

In [3] the following results were proved:

Proposition 3.1 [3] *Let $g, m \geq 0$ be such that $2g - 2 + m > 0$. Then*

$$\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_m}(e_{\alpha_m}) \rangle_g^{\text{DR}} = 0 \quad \text{if } \sum_{i=1}^m d_i < 2g - 1.$$

Proposition 3.2 [3] *There exists a unique differential polynomial $\mathcal{P} \in \widehat{\mathcal{A}}_w^{[-2]}$ such that for the power series $F^{\text{red}} \in \mathbb{C}[[t_*^*, \varepsilon]]$ defined by*

$$(13) \quad F^{\text{red}} := F + \mathcal{P}(w_*^*, \varepsilon)|_{w_*^*=(w^{\text{top}})_*(x,t_*^*,\varepsilon)|_{x=0}},$$

the correlators

$$\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{\text{red}} := \text{Coef}_{\varepsilon^{2g}} \frac{\partial^n F^{\text{red}}}{\partial t_{d_1}^{\alpha_1} \cdots \partial t_{d_n}^{\alpha_n}} \Big|_{t_*^*=0}$$

satisfy the following vanishing property:

$$(14) \quad \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{\text{red}} = 0 \quad \text{if } \sum_{i=1}^n d_i < 2g - 1.$$

In light of these two results the following conjecture was formulated in [3]:

Conjecture 3.3 (strong DR/DZ equivalence) *Consider a semisimple cohomological field theory and the associated DZ and DR hierarchies. Then the normal Miura transformation (12) defined by the differential polynomial \mathcal{P} of Proposition 3.2 maps the DZ hierarchy to the DR hierarchy respecting their tau-structures.*

As proved in [3] this conjecture is equivalent to saying that $F^{\text{red}} = F^{\text{DR}}$. This last form of the conjecture can be generalized to arbitrary CohFTs, forgetting about the DZ hierarchy and concentrating on the reduced and DR potentials.

Conjecture 3.4 (generalized strong DR/DZ equivalence) *For an arbitrary cohomological field theory we have $F^{\text{DR}} = F^{\text{red}}$.*

3.4 From intersection numbers to cohomology classes

The following result makes the relation between Conjectures 2.5 and 3.4 explicit, showing in particular how the first implies the second.

Proposition 3.5 *Consider an arbitrary cohomological field theory $c_{g,n}: V^{\otimes n} \rightarrow H^{\text{even}}(\overline{\mathcal{M}}_{g,n}, \mathbb{C})$. Then for any $g, n \geq 0$ with $2g - 2 + n > 0$ and numbers $d_1, \dots, d_n \geq 0$ such that $\sum d_i \geq 2g - 1$, we have*

$$(15) \quad \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{\text{DR}} = \int_{\overline{\mathcal{M}}_{g,n}} A_{d_1, \dots, d_n}^g c_{g,n}(e_{\alpha_1} \otimes \cdots \otimes e_{\alpha_n}),$$

$$(16) \quad \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{\text{red}} = \int_{\overline{\mathcal{M}}_{g,n}} B_{d_1, \dots, d_n}^g c_{g,n}(e_{\alpha_1} \otimes \cdots \otimes e_{\alpha_n}).$$

Proof In [4] the authors proved that for any $d \geq 2g - 1$ we have

$$\begin{aligned} & \sum_{\substack{d_1, \dots, d_n \geq 0 \\ \sum d_i = d}} \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{\text{DR}} a_1^{d_1} \cdots a_n^{d_n} \\ &= \frac{1}{\sum a_i} \sum_{\Gamma \in \text{ST}_{g,n+1}^{d-2g+2}} a(\Gamma) \int_{\overline{\mathcal{M}}_{g,n+1}} \text{DR}_{\Gamma} \left(-\sum a_i, a_1, \dots, a_n \right) \lambda_g c_{g,n+1} \left(e_1 \otimes \bigotimes_{i=1}^n e_{\alpha_i} \right) \\ &= \int_{\overline{\mathcal{M}}_{g,n}} A^{g,d-2g+2}(a_1, \dots, a_n) c_{g,n} \left(\bigotimes_{i=1}^n e_{\alpha_i} \right). \end{aligned}$$

Equation (15) is proved.

Let us prove (16). The reduced potential F^{red} can be constructed in the following way. Let us recursively construct a sequence of power series

$$F^{(0,-2)} = F, F^{(1,0)}, F^{(2,0)}, F^{(2,1)}, F^{(2,2)}, \dots, F^{(j,0)}, F^{(j,1)}, \dots, F^{(j,2j-2)}, \dots \in \mathbb{C}[[t_*^*, \varepsilon]].$$

Suppose that a series $F^{(j,k)}$ is already defined. Introduce correlators

$$\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{(j,k)} := \text{Coef}_{\varepsilon^{2g}} \frac{\partial^n F^{(j,k)}}{\partial t_{d_1}^{\alpha_1} \cdots \partial t_{d_n}^{\alpha_n}} \Big|_{t_*^*=0}.$$

If $k < 2j - 2$, then we define the series $F^{(j,k+1)}$ by

$$(17) \quad F^{(j,k+1)} := F^{(j,k)} - \sum_{n \geq 0} \sum_{\substack{d_1, \dots, d_n \geq 0 \\ \sum d_i = k+1}} \frac{\varepsilon^{2j}}{n!} \left\langle \prod \tau_{d_i}(e_{\alpha_i}) \right\rangle_j^{(j,k)} \times \prod ((w^{\text{top}})_{d_i}^{\alpha_i} - \delta^{\alpha_i, 1} \delta_{d_i, 1})|_{x=0}.$$

If $k = 2j - 2$, then we define the series $F^{(j+1,0)}$ by an analogous formula

$$F^{(j+1,0)} := F^{(j,2j-2)} - \sum_{n \geq 0} \frac{\varepsilon^{2j+2}}{n!} \left\langle \prod_{i=1}^n \tau_0(e_{\alpha_i}) \right\rangle_{j+1}^{(j,2j-2)} \prod (w^{\text{top}})^{\alpha_i} |_{x=0}.$$

Recall that

$$(w^{\text{top}})^{\alpha} = \eta^{\alpha\mu} \frac{\partial^2 F}{\partial t_0^\mu \partial t_0^1} \Big|_{t_0^1 \mapsto t_0^1 + x}.$$

The string equation for the potential F ,

$$\frac{\partial F}{\partial t_0^1} = \sum_{n \geq 0} t_{n+1}^\alpha \frac{\partial F}{\partial t_n^\alpha} + \frac{1}{2} \eta_{\alpha\beta} t_0^\alpha t_0^\beta + \varepsilon^2 \langle \tau_0(e_1) \rangle_1,$$

implies that the function $(w^{\text{top}})_n^\alpha |_{x=0}$ has the form

$$(w^{\text{top}})_n^\alpha |_{x=0} = \delta^{\alpha,1} \delta_{n,1} + t_n^\alpha + r_n^\alpha + O(\varepsilon^2),$$

where the power series $r_n^\alpha \in \mathbb{C}[[t_*^*]]$ doesn't contain monomials $t_{b_1}^{\beta_1} \dots t_{b_m}^{\beta_m}$ with $\sum b_i \leq n$. Clearly, if $g \leq j$, then we have the property

$$\langle \tau_{d_1}(e_{\alpha_1}) \dots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{(j,k)} = 0 \quad \text{if} \quad \sum d_i \leq \begin{cases} 2g-2 & \text{if } g < j, \\ k & \text{if } g = j. \end{cases}$$

Define a series F' by $F' := \lim_{j \rightarrow \infty} F^{(j,2j-2)}$. The series F' has the form

$$F' = F + \mathcal{P}'(w^{\text{top}}, w_x^{\text{top}}, \dots, \varepsilon) |_{x=0}$$

for some nonhomogeneous differential polynomial $\mathcal{P}' = \sum_{i \leq -2} \mathcal{P}'_i$ with $\mathcal{P}'_i \in \hat{\mathcal{A}}_w^{[i]}$. Moreover, we have the property

$$\text{Coef}_{\varepsilon^{2g}} \frac{\partial^n F'}{\partial t_{d_1}^{\alpha_1} \dots \partial t_{d_n}^{\alpha_n}} \Big|_{t_*^*=0} = 0 \quad \text{if} \quad \sum d_i \leq 2g - 2.$$

One can notice that the recursive construction, described above, is slightly different from the recursive construction of the reduced potential F^{red} presented in the proof of Proposition 7.2 in [3]. However, using the uniqueness argument given there we can see that $F' = F^{\text{red}}$ and that actually $\mathcal{P}' \in \hat{\mathcal{A}}_w^{[-2]}$.

For a stable complete tree T and $1 \leq m \leq \text{deg}(T)$, let

$$g_m(T) := \sum_{\substack{v \in V(T) \\ l(v) \leq m}} g(v).$$

Before we proceed, let us prove the following simple lemma:

Lemma 3.6 Let $d_1, \dots, d_n \geq 0$, $(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}$ and $1 \leq m < \text{deg}(T)$. Suppose that $g_{m+1}(T) = g_m(T)$ and $e_*[T, q] \neq 0$. Then

$$\sum_{\substack{h \in H_+^e(T) \\ l(h)=m+1}} q(h) > \sum_{\substack{h \in H_+^e(T) \\ l(h)=m}} q(h).$$

Proof Consider a half-edge $h \in H_+^e(T)$ with $l(h) = m$ and let $v := v(\iota(h))$. We have $g(v) = 0$ and the map e forgets all $q(h) + 1$ extra legs incident to v . Therefore, if v is strongly stable, then $\sum_{h' \in H_+^e[v]} q(h') > q(h)$. If v is weakly stable, then $|H_+^e[v]| = 1$ and $q(h') = q(h)$, where $h' \in H_+^e[v]$. Since at least one vertex of level $m + 1$ is strongly stable, the lemma is true. \square

A stable complete tree T will be called (j, k) -admissible if for any $1 \leq m < \text{deg}(T)$ we have $g_m(T) \leq j$ and

$$\sum_{\substack{h \in H_+^e(T) \\ l(h)=m}} q(h) \leq \begin{cases} 2g_m(T) - 2 & \text{if } g_m(T) < j, \\ k & \text{if } g_m(T) = j. \end{cases}$$

Let $\Omega_{d_1, \dots, d_n}^{B, g, (j, k)} := \{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g} \mid T \text{ is } (j, k)\text{-admissible}\}$. Define the class

$$B_{d_1, \dots, d_n}^{g, (j, k)} := \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g, (j, k)}} (-1)^{\text{deg}(T)-1} e_*[T, q] \in R^{\sum d_i}(\overline{\mathcal{M}}_{g, n}).$$

Clearly, $B_{d_1, \dots, d_n}^{g, (j, k)} = B_{d_1, \dots, d_n}^g$ if $j > g$.

In order to prove (16), it is sufficient to prove that for any pair (j, k) from the sequence

$$(18) \quad (0, -2), (1, 0), (2, 0), (2, 1), (2, 2), \dots, (j, 0), (j, 1), \dots, (j, 2j - 2), \dots$$

we have

$$(19) \quad \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g^{(j, k)} = \int_{\overline{\mathcal{M}}_{g, n}} B_{d_1, \dots, d_n}^{g, (j, k)} c_{g, n}(e_{\alpha_1} \otimes \cdots \otimes e_{\alpha_n})$$

if either $g > j$, or $g \leq j$ and

$$\sum d_i > \begin{cases} 2g - 2 & \text{if } g < j, \\ k & \text{if } g = j. \end{cases}$$

We proceed by induction. Obviously, (19) is true for $(j, k) = (0, -2)$. Suppose that (19) is true for a pair (j, k) from the sequence (18). Let us check it for the next pair.

Suppose $k < 2j - 2$. For any $d_1, \dots, d_n \geq 0$ we have $\Omega_{d_1, \dots, d_n}^{B,g,(j,k)} \subset \Omega_{d_1, \dots, d_n}^{B,g,(j,k+1)}$. Using the induction assumption and formula (17), we see that it remains to check that

$$\begin{aligned}
 (20) \quad & \sum_{g,n \geq 0} \frac{\varepsilon^{2g}}{n!} \sum_{d_1, \dots, d_n \geq 0} \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g,(j,k+1)} \setminus \Omega_{d_1, \dots, d_n}^{B,g,(j,k)}} (-1)^{\deg(T)} \\
 & \times \left(\int_{\overline{\mathcal{M}}_{g,n}} e_*[T, q] c_{g,n} \left(\bigotimes_{i=1}^n e_{\alpha_i} \right) \right) \prod t_{d_i}^{\alpha_i} \\
 & = \sum_{n \geq 0} \sum_{\substack{d_1, \dots, d_n \geq 0 \\ \sum d_i = k+1}} \frac{\varepsilon^{2j}}{n!} \langle \prod \tau_{d_i}(e_{\alpha_i}) \rangle_j^{(j,k)} \left(\prod ((w^{\text{top}})_{d_i}^{\alpha_i} - \delta^{\alpha_i, 1} \delta_{d_i, 1})|_{x=0} - \prod t_{d_i}^{\alpha_i} \right).
 \end{aligned}$$

Consider a pair $(T, q) \in \Omega_{d_1, \dots, d_n}^{B,g,(j,k+1)} \setminus \Omega_{d_1, \dots, d_n}^{B,g,(j,k)}$ such that $e_*[T, q] \neq 0$. Then there exists $1 \leq m < \deg(T)$ such that $g_m(T) = j$ and $\sum_{h \in H_+^e(T), l(h)=m} q(h) = k + 1$. By Lemma 3.6, $m = \deg(T) - 1$. Denote by T' the stable rooted tree obtained by erasing all vertices in T of level $m + 1$ together with half-edges incident to them. Half-edges $h \in H_+^e(T)$ with $l(h) = m$ become marked legs of T' . Clearly, T' is a stable complete tree. By Lemma 3.6, the tree T' is (j, k) -admissible. Using the induction assumption, we conclude that (20) is true. This completes the induction step in the case $k < 2j - 2$. The case $k = 2j - 2$ is analogous. The proposition is proved. \square

4 Further structure of the relations

In this section we discuss the structure of the conjectural relations (10) in more details. In Section 4.1 we recall the formulas for the intersections of the double ramification cycle with a ψ -class and with a boundary divisor on $\overline{\mathcal{M}}_{g,n}$. In Section 4.2 we show that for a fixed $g \geq 1$ all relations $A_d^g = B_d^g$ with $d \geq 2g - 1$ follow from the relation $A_{2g-1}^g = B_{2g-1}^g$. In Section 4.3 we prove that the A - and the B -class behave in the same way upon the pullback along the forgetful map. We then use this result in Section 4.4 in order to show that Conjecture 2.5 is true if and only if it is true when all the d_i are positive. In Section 4.5 we prove that the classes $A_{d_1, \dots, d_n, 1}^g$ and $B_{d_1, \dots, d_n, 1}^g$ behave in the same way upon the pushforward along the map forgetting the last marked point. Using this result, in Section 4.6 we show that Conjecture 2.5 is valid on $\mathcal{M}_{g,n}$. In Section 4.7 we show that the conjectural relations (10) give a new formula for the class $\lambda_g \in R^g(\overline{\mathcal{M}}_g)$ and check the resulting formula for $g \leq 3$.

4.1 Formulas with the double ramification cycles

First of all, let us recall the formula from [9] for the product of the double ramification cycle with a ψ -class. Denote by

$$\text{gl}_k: \overline{\mathcal{M}}_{g_1, n_1+k} \times \overline{\mathcal{M}}_{g_2, n_2+k} \rightarrow \overline{\mathcal{M}}_{g_1+g_2+k-1, n_1+n_2}$$

the gluing map that glues a curve from $\overline{\mathcal{M}}_{g_1, n_1+k}$ to a curve from $\overline{\mathcal{M}}_{g_2, n_2+k}$ along the last k marked points on the first curve and the last k marked points on the second curve. Introduce the notation

$$\begin{aligned} \text{DR}_{g_1}(a_1, \dots, a_n) \boxtimes_k \text{DR}_{g_2}(b_1, \dots, b_m) &:= \text{gl}_{k*}(\text{DR}_{g_1}(a_1, \dots, a_n) \times \text{DR}_{g_2}(b_1, \dots, b_m)) \\ &\in R^{g_1+g_2+k}(\overline{\mathcal{M}}_{g_1+g_2+k-1, n+m-2k}). \end{aligned}$$

Let a_1, \dots, a_n be a list of integers with vanishing sum. For a subset $I = \{i_1, \dots, i_k\} \subset \{1, \dots, n\}$ with $i_1 < i_2 < \dots < i_k$, let us denote by A_I the list a_{i_1}, \dots, a_{i_k} and by a_I the sum $\sum_{i \in I} a_i$. Assume that $a_s \neq 0$ for some $1 \leq s \leq n$. Then we have [9, Theorem 4]

$$\begin{aligned} (21) \quad &a_s \psi_s \text{DR}_g(a_1, \dots, a_n) \\ &= \sum_{\substack{I \sqcup J = \{1, \dots, n\} \\ a_I > 0}} \sum_{p \geq 1} \sum_{\substack{g_1, g_2 \geq 0 \\ g_1+g_2+p-1=g}} \sum_{\substack{k_1, \dots, k_p \geq 1 \\ \sum k_j = a_I}} \frac{\rho \prod_{i=1}^p k_i}{r \cdot p!} \\ &\quad \times \text{DR}_{g_1}(A_I, -k_1, \dots, -k_p) \boxtimes_p \text{DR}_{g_2}(A_J, k_1, \dots, k_p), \end{aligned}$$

where $r := 2g - 2 + n$ and

$$\rho := \begin{cases} 2g_2 - 2 + |J| + p & \text{if } s \in I; \\ -(2g_1 - 2 + |I| + p) & \text{if } s \in J. \end{cases}$$

Let us also recall the formula for the intersection of the double ramification cycle with a boundary divisor on $\overline{\mathcal{M}}_{g,n}$. For $0 \leq h \leq g$ and a subset $I \subset \{1, \dots, n\}$ we have [9]

$$\delta_h^I \cdot \text{DR}_g(a_1, \dots, a_n) = \text{DR}_h(A_I, -a_I) \boxtimes_1 \text{DR}_{g-h}(A_{I^c}, a_I),$$

where $I^c := \{1, 2, \dots, n\} \setminus I$.

4.2 One-point case

Lemma 4.1 *Let $g \geq 1$. Then for any $k \geq 0$ we have $A_{2g-1+k}^g = \psi_1^k A_{2g-1}^g$.*

Proof Let $\pi: \overline{\mathcal{M}}_{g,2} \rightarrow \overline{\mathcal{M}}_{g,1}$ be the forgetful map that forgets the second marked point. We compute

$$\begin{aligned} a\psi_1 A^{g,1}(a) &= \psi_1 \lambda_g \text{DR}_g(a, \widetilde{-a}) = \pi_*(\pi^* \psi_1 \cdot \lambda_g \text{DR}_g(a, -a)) \\ &= \pi_*(\psi_1 \cdot \lambda_g \text{DR}_g(a, -a)) - \pi_*(\delta_0^{\{1,2\}} \cdot \lambda_g \text{DR}_g(a, -a)) \\ &= \sum_{\substack{g_1, g_2 \geq 1 \\ g_1 + g_2 = g}} \frac{g_2}{g} \pi_*(\lambda_g \text{DR}_{g_1}(a, -a) \boxtimes_1 \text{DR}_{g_2}(-a, a)) \\ &\quad - \pi_*(\lambda_g \text{DR}_g(0) \boxtimes_1 \text{DR}_0(a, -a, 0)) \\ &= A^{g,2}(a), \end{aligned}$$

where we used that $\lambda_g \text{DR}_g(0) = (-1)^g \lambda_g^2 = 0$. If $k = 1$, then the lemma is proved. If $k \geq 2$, then we write the equation $(a_1 \psi_1)^k A^{g,1}(a) = (a_1 \psi_1)^{k-1} A^{g,2}(a)$ and apply formula (21) to the right-hand side of it $k - 1$ times. The lemma is proved. \square

On the other hand, it is not hard to get an explicit expression for the class B_d^g . Let $g_1, g_2, \dots, g_k \geq 1$ and $d_1, \dots, d_k \geq 0$. Introduce a class

$$C_{d_1, \dots, d_k}^{g_1, \dots, g_k} := \textcircled{g_1}^{\psi^{d_1}} \textcircled{g_2}^{\psi^{d_2}} \dots \textcircled{g_k}^{\psi^{d_k}} \in R^{\sum d_i + k - 1}(\overline{\mathcal{M}}_{\sum g_i, 1}).$$

Then it is easy to see that for $g \geq 1$ and $d \geq 2g - 1$ we have

$$B_d^g = \sum_{k=1}^g \sum_{\substack{g_1, \dots, g_k \geq 1 \\ \sum g_i = g}} \sum_{d_1, \dots, d_k} (-1)^{k-1} C_{d_1, \dots, d_k}^{g_1, \dots, g_k},$$

where the last sum is taken over all nonnegative integers d_1, \dots, d_k satisfying

$$\sum_{i=1}^l d_i + l - 1 \leq 2 \sum_{i=1}^l g_i - 2 \quad \text{if } 1 \leq l \leq g - 1, \quad \sum_{i=1}^k d_i + k - 1 = d.$$

We see that $B_d^g = \psi_1^{d-2g+1} B_{2g-1}^g$. Thus, for $n = 1$, Conjecture 2.5 is equivalent to the sequence of relations

$$A_{2g-1}^g = B_{2g-1}^g, \quad g \geq 1.$$

4.3 String equation

In this section we prove that the A - and the B -class behave in the same way upon the pullback along the forgetful map $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$.

Proposition 4.2 Denote by $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map that forgets the last marked point. Then we have

$$(22) \quad A_{d_1, \dots, d_n, 0}^g = \begin{cases} \pi^* A_{d_1, \dots, d_n}^g & \text{if } \sum d_i = 2g - 1, \\ \pi^* A_{d_1, \dots, d_n}^g + \sum_{\substack{1 \leq i \leq n \\ d_i \geq 1}} \delta_0^{\{i, n+1\}} \pi^* A_{d_1, \dots, d_i-1, \dots, d_n}^g & \text{if } \sum d_i \geq 2g. \end{cases}$$

Proof Let $m := \sum d_i - 2g + 2$. The proposition is equivalent to the equation

$$(23) \quad \tilde{A}^{g,m}(a_1, \dots, a_n, 0) = \begin{cases} \pi^* \tilde{A}^{g,m}(a_1, \dots, a_n) & \text{if } m = 1, \\ \pi^* \tilde{A}^{g,m}(a_1, \dots, a_n) + \sum_{i=1}^n a_i \delta_0^{\{i, n+1\}} \pi^* \tilde{A}^{g,m-1}(a_1, \dots, a_n) & \text{if } m \geq 2, \end{cases}$$

where a_1, \dots, a_n are arbitrary integers. Let $a_0 := -\sum_{i=1}^n a_i$. Introduce a class $\hat{A}^{g,m}(a_0, a_1, \dots, a_n)$ by

$$\hat{A}^{g,m}(a_0, a_1, \dots, a_n) := \sum_{\Gamma \in \text{ST}_{g,n+1}^m} a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n).$$

Formula (23) follows from the equation

$$(24) \quad \hat{A}^{g,m}(a_0, \dots, a_n, 0) = \begin{cases} \pi^* \hat{A}^{g,m}(a_0, \dots, a_n) & \text{if } m = 1, \\ \pi^* \hat{A}^{g,m}(a_0, \dots, a_n) + \sum_{i=1}^n a_i \delta_0^{\{i, n+1\}} \pi^* \hat{A}^{g,m-1}(a_0, \dots, a_n) & \text{if } m \geq 2, \end{cases}$$

where the map $\pi: \overline{\mathcal{M}}_{g,n+2} \rightarrow \overline{\mathcal{M}}_{g,n+1}$ forgets the last marked point.

For $m = 1$, equation (24) is clear. Suppose that $m \geq 2$. Consider a stable tree $\Gamma \in \text{ST}_{g,n+2}^m$. Recall that we denote by $l_i(\Gamma)$ the leg of Γ marked by $0 \leq i \leq n + 1$. We will call a vertex $v \in V(\Gamma)$ exceptional if $g(v) = 0$, $n(v) = 3$ and the leg $l_{n+1}(\Gamma)$ is incident to v . An exceptional vertex $v \in V(\Gamma)$ will be called bad if it is not incident to any leg $l_i(\Gamma)$, where $1 \leq i \leq n$. We will call the tree Γ bad if it has a bad vertex. Otherwise, it will be called good. For a vertex $v \in V(\Gamma)$ let

$$r'(v) := \begin{cases} 2g(v) + n(v) - 2 & \text{if } l_{n+1}(\Gamma) \text{ is not incident to } v, \\ 2g(v) + n(v) - 3 & \text{if } l_{n+1}(\Gamma) \text{ is incident to } v. \end{cases}$$

For a good stable tree $\Gamma \in \text{ST}_{g,n+2}^m$ introduce a constant $a'(\Gamma)$ by

$$a'(\Gamma) := \left(\prod_{h \in H_+^e(\Gamma)} a(h) \right) \prod_{\substack{v \in V(\Gamma) \\ v \text{ is not exceptional}}} \frac{r'(v)}{\sum_{\tilde{v} \in \text{Desc}[v]} r'(\tilde{v})}.$$

Using this notation, we can rewrite the right-hand side of (24) as

$$\begin{aligned} \pi^* \widehat{A}^{g,m}(a_0, \dots, a_n) + \sum_{i=1}^n a_i \delta_0^{\{i,n+1\}} \pi^* \widehat{A}^{g,m-1}(a_0, \dots, a_n) \\ = \sum_{\substack{\Gamma \in \text{ST}_{g,n+2}^m \\ \Gamma \text{ is good}}} a'(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n, 0). \end{aligned}$$

On the other hand, by definition,

$$\widehat{A}^{g,m}(a_0, \dots, a_n, 0) = \sum_{\Gamma \in \text{ST}_{g,n+2}^m} a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n, 0).$$

We see that we have to prove the equation

$$\begin{aligned} (25) \quad \sum_{\Gamma \in \text{ST}_{g,n+2}^m} a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n, 0) \\ = \sum_{\substack{\Gamma \in \text{ST}_{g,n+2}^m \\ \Gamma \text{ is good}}} a'(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n, 0). \end{aligned}$$

Let us prove equation (25). Suppose Γ is a bad stable tree. Let us show how to express the class $a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, a_1, \dots, a_n, 0)$ as a linear combination of the classes $\lambda_g \text{DR}_{\widetilde{\Gamma}}(a_0, a_1, \dots, a_n, 0)$, where the stable trees $\widetilde{\Gamma}$ are good. Suppose that $s \geq 2$ and b_1, \dots, b_s are integers with vanishing sum. We have the following relation in the cohomology of $\overline{\mathcal{M}}_{g,s+2}$ (see eg [1, equation (5.2)]):

$$(26) \quad \lambda_g \sum_{\substack{I \sqcup J = \{1, \dots, s\} \\ I, J \neq \emptyset}} \sum_{g_1 + g_2 = g} b_I \text{DR}_{g_1}(0, B_I, -b_I) \boxtimes_1 \text{DR}_{g_2}(0, B_J, -b_J) = 0.$$

Suppose that the point with the zero multiplicity in the second double ramification cycle is marked by $s + 2$. Let us multiply relation (26) by ψ_{s+2} and push it forward to $\overline{\mathcal{M}}_{g,s+1}$ by forgetting the last marked point:

$$(27) \quad \lambda_g \sum_{\substack{I \sqcup J = \{1, \dots, s\} \\ I, J \neq \emptyset}} \sum_{\substack{g_1 + g_2 = g \\ 2g_2 + |J| - 1 > 0}} b_I (2g_2 + |J| - 1) \times \text{DR}_{g_1}(0, B_I, -b_I) \boxtimes_1 \text{DR}_{g_2}(B_J, -b_J) = 0.$$

Suppose that the level of the bad vertex in our bad stable tree Γ is equal to k . Then relation (27) allows us to express the class $a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, \dots, a_n, 0)$ in terms of the classes $\lambda_g \text{DR}_{\widetilde{\Gamma}}(a_0, \dots, a_n, 0)$, where the tree $\widetilde{\Gamma}$ is good or bad with the bad vertex of level $k + 1$. Therefore, applying relation (27) sufficiently many times, we come to a




Figure 1: Stable tree Γ .

decomposition

$$a(\Gamma)\lambda_g \text{DR}_\Gamma(a_0, \dots, a_n, 0) = \sum_{\substack{\tilde{\Gamma} \in \text{ST}_{g,n+2}^m \\ \tilde{\Gamma} \text{ is good}}} a(\Gamma, \tilde{\Gamma})\lambda_g \text{DR}_{\tilde{\Gamma}}(a_0, \dots, a_n, 0),$$

where $a(\Gamma, \tilde{\Gamma})$ are certain coefficients. We see that for any good graph Γ we have to prove the identity

$$(28) \quad a(\Gamma) + \sum_{\substack{\tilde{\Gamma} \in \text{ST}_{g,n+2}^m \\ \tilde{\Gamma} \text{ is bad}}} a(\tilde{\Gamma}, \Gamma) = a'(\Gamma).$$

Let us prove (28). Suppose that the leg $l_{n+1} = l_{n+1}(\Gamma)$ is incident to a vertex of level k . Denote it by v_k . Denote by v_1 the root of Γ . Let v_1, v_2, \dots, v_k be the unique path connecting v_1 and v_k . Denote by $v_{k+1}^1, \dots, v_{k+1}^l$, $l \geq 0$, the direct descendants of v_k . Let $L' := L[v_k] \setminus \{l_{n+1}\}$. In Figure 1 we draw our tree Γ . Note that each vertex v in the picture is decorated by the number $r(v)$, instead of the genus. This is more convenient for the computations. We use the notation $r_i := r(v_i)$ for $1 \leq i \leq k$ and $r_{k+1}^j := r(v_{k+1}^j)$ for $1 \leq j \leq l$. The symbols A_i and A_{k+1}^j indicate the pieces of the tree Γ that don't contain the vertices v_i and v_{k+1}^j . Let us also introduce the notation

$$R_i := \sum_{v \in \text{Desc}[v_i]} r(v), \quad 1 \leq i \leq k,$$

$$R_{k+1}^j := \sum_{v \in \text{Desc}[v_{k+1}^j]} r(v), \quad 1 \leq j \leq l,$$

$$\tilde{a} := a(\Gamma) / \left(\prod_{i=1}^k \frac{r_i}{R_i} \prod_{j=1}^l \frac{r_{k+1}^j}{R_{k+1}^j} \right).$$




Figure 2: Bad stable tree of the first type.

There are two cases: the vertex v_k can be exceptional or not.

Suppose that v_k is not exceptional. Then

$$a'(\Gamma) = \tilde{a} \frac{r_1 \cdots r_{k-1} (r_k - 1)}{(R_1 - 1) \cdots (R_k - 1)} \prod_{j=1}^l \frac{r_{k+1}^j}{R_{k+1}^j}.$$

It is not hard to understand the structure of bad stable trees $\tilde{\Gamma}$ such that $a(\tilde{\Gamma}, \Gamma) \neq 0$. These trees are of two types. A bad tree of the first type is shown in Figure 2, where $1 \leq i \leq k - 1$ and $\tilde{r} = r_{k-1} + r_k - 1$. A bad tree of the second type is shown in Figure 3, where $1 \leq i \leq k$, $1 \leq j \leq l$ and $\tilde{r} = r_k + r_{k+1}^j - 1$. It is not hard to see that

$$a(\tilde{\Gamma}, \Gamma) = \tilde{a} \frac{r_1 \cdots r_{k-1}}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)} \prod_{j=1}^l \frac{r_{k+1}^j}{R_{k+1}^j}$$

if $\tilde{\Gamma}$ is of the first type, and

$$a(\tilde{\Gamma}, \Gamma) = -\tilde{a} \frac{r_1 \cdots r_{k-1}}{R_1 \cdots R_i (R_i - 1) \cdots (R_k - 1)} R_{k+1}^j \prod_{m=1}^l \frac{r_{k+1}^m}{R_{k+1}^m}$$

if $\tilde{\Gamma}$ is of the second type.




Figure 3: Bad stable tree of the second type.

Therefore, (28) follows from the identity

$$\prod_{i=1}^k \frac{r_i}{R_i} + \sum_{i=1}^{k-1} \frac{r_1 \cdots r_{k-1}}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)} - \sum_{i=1}^k \sum_{j=1}^l \frac{r_1 \cdots r_{k-1}}{R_1 \cdots R_i (R_i - 1) \cdots (R_k - 1)} R_{k+1}^j = \frac{r_1 \cdots r_{k-1} (r_k - 1)}{(R_1 - 1) \cdots (R_k - 1)},$$

or, equivalently,

$$(29) \quad \frac{r_k}{R_1 \cdots R_k} + \sum_{i=1}^{k-1} \frac{1}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)} - \sum_{i=1}^k \frac{R_k - r_k}{R_1 \cdots R_i (R_i - 1) \cdots (R_k - 1)} = \frac{r_k - 1}{(R_1 - 1) \cdots (R_k - 1)}.$$

Note that

$$\frac{r_k}{R_1 \cdots R_k} - \frac{R_k - r_k}{R_1 \cdots R_k (R_k - 1)} = \frac{r_k - 1}{R_1 \cdots R_{k-1} (R_k - 1)},$$

$$\frac{1}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)} - \frac{R_k - r_k}{R_1 \cdots R_i (R_i - 1) \cdots (R_k - 1)} = \frac{r_k - 1}{R_1 \cdots R_i (R_i - 1) \cdots (R_k - 1)},$$

where $1 \leq i \leq k - 1$. Therefore, (29) is equivalent to the equation

$$(30) \quad \frac{1}{R_1 \cdots R_{k-1}} + \sum_{i=1}^{k-1} \frac{1}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)} = \frac{1}{(R_1 - 1) \cdots (R_{k-1} - 1)},$$

which can be easily proved by induction on k .

Suppose that v_k is exceptional. Then $l = 0$, the set L' consists of only one leg and $r_k = R_k = 1$. We have

$$a'(\Gamma) = \tilde{a} \frac{r_1 \cdots r_{k-1}}{(R_1 - 1) \cdots (R_{k-1} - 1)}.$$

A bad stable tree $\tilde{\Gamma}$ with $a(\tilde{\Gamma}, \Gamma) \neq 0$ should necessarily be of the first type (see Figure 2) and then we have

$$a(\tilde{\Gamma}, \Gamma) = \tilde{a} \frac{r_1 \cdots r_{k-1}}{R_1 \cdots R_i (R_i - 1) \cdots (R_{k-1} - 1)}.$$

We immediately see that again (28) follows from the elementary identity (30). The proposition is proved. \square

Proposition 4.3 Denote by $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map that forgets the last marked point. Then we have

$$(31) \quad B_{d_1, \dots, d_n, 0}^g = \begin{cases} \pi^* B_{d_1, \dots, d_n}^g & \text{if } \sum d_i = 2g - 1, \\ \pi^* B_{d_1, \dots, d_n}^g + \sum_{\substack{1 \leq i \leq n \\ d_i \geq 1}} \delta_0^{\{i, n+1\}} \pi^* B_{d_1, \dots, d_{i-1}, \dots, d_n}^g & \text{if } \sum d_i \geq 2g. \end{cases}$$

Proof Let $(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}$ be an admissible and stable complete tree with a power function

$$q: H_+^{\text{em}}(T) \rightarrow \mathbb{N},$$

as in Definition 2.4. We denote by $\text{deg}(T)$ its number of levels. In particular, there are extra legs at every vertex (except the root) that we will eventually forget when computing the B -class.

Choose a vertex $v \in V(T)$. Let $C = (e_1^C, v_1^C, \dots, e_{\text{deg}(T)-l(v)}^C, v_{\text{deg}(T)-l(v)}^C, \sigma_{n+1})$ be a chain of weakly stable vertices with a new marking σ_{n+1} . Precisely, the edge e_1^C is attached to the vertex v_1^C , the edge e_k^C links the vertex v_{k-1}^C to v_k^C , and the leg σ_{n+1} is attached to the vertex $v_{\text{deg}(T)-l(v)}^C$. Moreover, every vertex is of genus 0 and contains an extra leg. We construct a tree T_v , obtained from T by gluing the edge e_1^C (and thus the chain C) to the vertex v . We have $H_+^{\text{em}}(T) \subset H_+^{\text{em}}(T_v)$ and we extend the power function q into a function $q_v: H_+^{\text{em}}(T_v) \rightarrow \mathbb{N}$ by taking

$$q_v(h_k^C) := 0 \quad \text{and} \quad q_v(\sigma_{n+1}) := 0,$$

where h_k^C is the half-edge in $H_+^{\text{em}}(T_v)$ contained in the edge e_k^C . It is easy to see that we get

$$(T_v, q_v) \in \Omega_{d_1, \dots, d_n, 0}^{B, g}.$$

Choose a half-edge $h \in H_+^{\text{em}}(T)$ attached to the vertex v and such that $q(h) > 0$. We construct a tree $T_{(v, h)}$, obtained from T by adding an extra level between the levels $l(v)$ and $l(v) + 1$ of T as follows:

- denote by $h_0, \dots, h_m \in H_+^{\text{em}}(T)$ the half-edges of level $l(v)$, with $h_0 := h$;
- insert a pair (e_k, v_k) between the half-edge h_k and the vertex it is attached to, where $e_k = (h'_k, h''_k)$ is an edge and v_k is a vertex of genus 0;
- glue the half-edge h_1^C from the chain C to the vertex v_0 ;
- add $q(h)$ extra legs to the vertex v_0 and $q(h_k) + 1$ extra legs to the vertex v_k for $1 \leq k \leq m$.

Therefore, the number of levels of the tree $T_{(v,h)}$ is $\text{deg}(T) + 1$, the vertex $v_0 \in V(T_{(v,h)})$ is the only strongly stable vertex at its level, and we have a natural inclusion $H_+^{\text{em}}(T_v) \subset H_+^{\text{em}}(T_{(v,h)})$. Then, we extend the power function q_v into a function $q_{(v,h)}: H_+^{\text{em}}(T_{(v,h)}) \rightarrow \mathbb{N}$ by taking

$$q_{(v,h)}(h'_k) := \begin{cases} q(h_k) - 1 & \text{if } k = 0, \\ q(h_k) & \text{if } k \neq 0. \end{cases}$$

The complete tree $T_{(v,h)}$ is obviously stable, but not necessarily admissible. We get

$$(T_{(v,h)}, q_{(v,h)}) \in \Omega_{d_1, \dots, d_n, 0}^{B,g} \iff l(v) \neq \text{deg}(T) \text{ or } \sum_{i=1}^n d_i = 2g - 1.$$

Furthermore, observe that when $l(v) = \text{deg}(T)$, the half-edge h corresponds to a marking σ_i and we get

$$e_*[T_{(v,h)}, q_{(v,h)}] = \sigma_i^* e_*[T, q_i] = \delta_0^{\{i, n+1\}} \cdot \pi^* e_*[T, q_i] \in R^*(\overline{\mathcal{M}}_{g, n+1}),$$

where the morphism σ_i denotes here the section of the i^{th} marking in the universal curve $\mathcal{C}_{g,n} \simeq \overline{\mathcal{M}}_{g, n+1}$, and where $q_i: H_+^{\text{em}}(T) \rightarrow \mathbb{N}$ is defined by

$$q_i(h) := \begin{cases} d_i - 1 & \text{if } h = \sigma_i, \\ q(h) & \text{otherwise.} \end{cases}$$

Conversely, let $(T', q') \in \Omega_{d_1, \dots, d_n, 0}^{B,g}$ and denote by v the first strongly stable ancestor of the marking σ_{n+1} . In particular, the marking σ_{n+1} is attached to the vertex v via a chain C of weakly stable vertices and we denote by $h^{(n+1)} \in H_+^{\text{em}}(T')$ the half-edge from C attached to v . We have two possibilities:

- (1) v is a vertex of genus 0 with exactly two half-edges $h, h^{(n+1)} \in H_+^{\text{em}}(T')$ attached to it and v is the only strongly stable vertex of level $l(v)$;
- (2) v is another type of vertex.

Denote by T the tree obtained from T' by forgetting the chain C containing the marking σ_{n+1} , and contracting the level $l(v)$ in case (1). In particular, the power function q' restricts to a function q and we get

$$(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g} \quad \text{and} \quad (T', q') = \begin{cases} (T(v, h), q(v, h)) & \text{in case (1),} \\ (T_v, q_v) & \text{in case (2).} \end{cases}$$

Furthermore, from the formula

$$(32) \quad \pi^* \begin{array}{c} | \\ \circ \\ | \\ q_1 \cdots q_r \end{array} = \begin{array}{c} | \\ \circ \\ | \\ q_1 \cdots q_r \end{array} \begin{array}{c} | \\ \circ \\ | \\ n+1 \end{array} - \sum_{\substack{1 \leq i \leq r \\ q_i > 0}} \begin{array}{c} | \\ \circ \\ | \\ q_1 \cdots \hat{q}_i \cdots q_r \end{array} \begin{array}{c} | \\ \circ \\ | \\ q_i - 1 \end{array} \begin{array}{c} | \\ \circ \\ | \\ i \end{array} \begin{array}{c} | \\ \circ \\ | \\ n+1 \end{array}$$

expressing the pullback of ψ -classes via the map π , we obtain

$$e_* \pi^* ([T, q]) = \sum_{v \in T} e_* \left([T_v, q_v] - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v, q(h) > 0}} [T(v, h), q(v, h)] \right)$$

for every $(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}$ and where $h \rightarrow v$ means that the half-edge h is incident to the vertex v . As a consequence, when $d_1 + \dots + d_n \geq 2g$, we obtain

$$\begin{aligned} \pi^* (B_{d_1, \dots, d_n}^g) &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \pi^* e_* [T, q] \\ &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} e_* \pi^* [T, q], \end{aligned}$$

where the second equality comes from the general fact that $\overline{M}_{g, n+2}$ is birational to the fibre product $\overline{M}_{g, n+1} \times_{\overline{M}_{g, n}} \overline{M}_{g, n+1}$, and then

$$\begin{aligned} &\pi^* (B_{d_1, \dots, d_n}^g) \\ &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \sum_{v \in T} \left(e_* [T_v, q_v] - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v, q(h) > 0}} e_* [T(v, h), q(v, h)] \right) \\ &= \sum_{(T', q') \in \Omega_{d_1, \dots, d_n, 0}^{B, g}} (-1)^{\deg(T')-1} e_* [T', q'] \\ &\quad - \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \sum_{\substack{v \in T \\ l(v) = \deg(T)}} \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v, q(h) > 0}} e_* [T(v, h), q(v, h)] \end{aligned}$$

$$\begin{aligned}
 &= B_{d_1, \dots, d_n, 0}^g - \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \sum_{\substack{1 \leq i \leq n \\ \bar{d}_i > 0}} \delta_0^{\{i, n+1\}} \cdot \pi^* e_* [T, q_i] \\
 &= B_{d_1, \dots, d_n, 0}^g - \sum_{\substack{1 \leq i \leq n \\ \bar{d}_i > 0}} \delta_0^{\{i, n+1\}} \cdot \pi^* B_{d_1, \dots, d_i-1, \dots, d_n}^g.
 \end{aligned}$$

When $d_1 + \dots + d_n = 2g - 1$, we have seen that $(T_{(v,h)}, q_{(v,h)})$ is always admissible, so that the first three equalities are the same, but there is no second term in the last three equalities. Hence we get

$$\pi^* B_{d_1, \dots, d_n}^g = B_{d_1, \dots, d_n, 0}^g. \quad \square$$

4.4 Reduction of the conjecture

Proposition 4.4 *Conjecture 2.5 is true if and only if it is true when all the d_i are positive. Furthermore, Conjecture 2.5 is true in genus 0 and in genus 1.*

Proof The first statement follows immediately from Propositions 4.2 and 4.3.

Assume $g = 0$. Since $\dim \bar{\mathcal{M}}_{0,n} = n - 3$, the classes A_{d_1, \dots, d_n}^0 and B_{d_1, \dots, d_n}^0 are nontrivial only if $\sum d_i \leq n - 3$. Therefore, we can always apply formulas (22) and (31) to them, unless $n = 3$ and $d_1 = d_2 = d_3 = 0$, where we get

$$A_{0,0,0}^0 = B_{0,0,0}^0 = 1 \in H^0(\bar{\mathcal{M}}_{0,3}, \mathbb{Q}).$$

Assume $g = 1$. Since $\dim \bar{\mathcal{M}}_{1,n} = n$, the classes A_{d_1, \dots, d_n}^1 and B_{d_1, \dots, d_n}^1 are nontrivial only if $\sum d_i \leq n$. Therefore, we can always apply formulas (22) and (31) to them, unless $d_1 = d_2 = \dots = d_n = 1$. In order to prove that $A_{1,1, \dots, 1}^1 = B_{1,1, \dots, 1}^1$, it is sufficient to check that $\int_{\bar{\mathcal{M}}_{1,n}} A_{1,1, \dots, 1}^1 = \int_{\bar{\mathcal{M}}_{1,n}} B_{1,1, \dots, 1}^1$. Note that these two integrals are equal to $\langle \tau_1(e_1)^n \rangle_1^{\text{DR}}$ and $\langle \tau_1(e_1)^n \rangle_1^{\text{red}}$, respectively, for the trivial cohomological field theory. The equality $F^{\text{DR}} = F^{\text{red}}$ for the trivial cohomological field theory was checked in [3]. Therefore, Conjecture 2.5 is true in genus 1. □

4.5 Dilaton equation

Here we prove that the classes $A_{d_1, \dots, d_n, 1}^g$ and $B_{d_1, \dots, d_n, 1}^g$ behave in the same way upon the pushforward along the map forgetting the last marked point.

Proposition 4.5 Denote by $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map that forgets the last marked point. Then we have

$$(33) \quad \pi_*(A_{d_1, \dots, d_n, 1}^g) = \begin{cases} (2g - 2 + n)A_{d_1, \dots, d_n}^g & \text{if } \sum d_i > 2g - 2, \\ 0 & \text{if } \sum d_i = 2g - 2. \end{cases}$$

Before proving the proposition let us formulate three auxiliary statements. Recall that for a stable tree $\Gamma \in \text{ST}_{g,n+1}^m$ we denote by $v_1(\Gamma)$ the root of Γ and by $l_i(\Gamma)$ for $0 \leq i \leq n$ the leg of Γ marked by i .

Lemma 4.6 Let a_0, \dots, a_n with $n \geq 1$ be integers with vanishing sum and $m \geq 2$. Then we have

$$\begin{aligned} \widehat{A}^{g,m}(a_0, \dots, a_n) - a_1 \psi_1 \widehat{A}^{g,m-1}(a_0, \dots, a_n) \\ = \sum_{\substack{\Gamma \in \text{ST}_{g,n+1}^m \\ v(l_1(\Gamma)) = v_1(\Gamma)}} \frac{2g - 1 + n}{r(v_1(\Gamma))} a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, \dots, a_n). \end{aligned}$$

Proof Using formula (21), for an arbitrary stable tree $\Gamma \in \text{ST}_{g,n+1}^{m-1}$ we can write a decomposition

$$a_1 \psi_1 \cdot a(\Gamma) \lambda_g \text{DR}_\Gamma(a_0, \dots, a_n) = \sum_{\tilde{\Gamma} \in \text{ST}_{g,n+1}^m} a(\Gamma, \tilde{\Gamma}) \lambda_g \text{DR}_{\tilde{\Gamma}}(a_0, \dots, a_n),$$

where $a(\Gamma, \tilde{\Gamma})$ are certain coefficients. Let $\Gamma \in \text{ST}_{g,n+1}^m$. The statement of the lemma is equivalent to the equation

$$(34) \quad a(\Gamma) - \sum_{\tilde{\Gamma} \in \text{ST}_{g,n+1}^{m-1}} a(\tilde{\Gamma}, \Gamma) = \begin{cases} \frac{2g - 1 + n}{r(v_1(\Gamma))} & \text{if } l_1(\Gamma) \text{ is incident to } v_1(\Gamma), \\ 0 & \text{otherwise.} \end{cases}$$

Let $v \in V(\Gamma)$ be the vertex incident to $l_1 = l_1(\Gamma)$. Denote by v_1'', \dots, v_l'' with $l \geq 0$ the direct descendants of v . Let $L' := L[v] \setminus \{l_1\}$, $r := r(v)$, $r_i'' := r(v_i'')$, $R := \sum_{\tilde{v} \in \text{Desc}[v]} r(\tilde{v})$ and $R_i'' := \sum_{\tilde{v} \in \text{Desc}[v_i'']} r(\tilde{v})$.

Suppose that $v \neq v_1(\Gamma)$. Denote by $v' \in V(\Gamma)$ the mother of v and let $r' := r(v')$ and $R' := \sum_{\tilde{v} \in \text{Desc}[v']} r(\tilde{v})$. We draw the stable tree Γ in Figure 4. Similarly to the figures in the proof of Proposition 4.2, we decorate a vertex w of Γ by number $r(w)$. It is not hard to see that there are exactly $l + 1$ stable trees $\tilde{\Gamma} \in \text{ST}_{g,n+1}^{m-1}$ such that $a(\tilde{\Gamma}, \Gamma) \neq 0$. The first one is shown on the left-hand side of Figure 5, and the other l




Figure 4: Stable tree Γ .

trees are on the right-hand side, where $1 \leq j \leq l$. Let

$$\tilde{a} := a(\Gamma) / \left(\frac{r'}{R'} \frac{r}{R} \prod_{j=1}^l \frac{r''_j}{R''_j} \right).$$

The coefficient $a(\tilde{\Gamma}, \Gamma)$ for the left tree in Figure 5 is equal to $(\tilde{a}r'/R') \prod_{k=1}^l r''_k/R''_k$ and for the right tree in Figure 5 it is equal to $-(\tilde{a}r'R''_j/R'R) \prod_{k=1}^l r''_k/R''_k$. We compute

$$\sum_{\tilde{\Gamma} \in \text{ST}_{g,n+1}^{m-1}} a(\tilde{\Gamma}, \Gamma) = \tilde{a} \left(\frac{r'}{R'} - \sum_{j=1}^l \frac{r'R''_j}{R'R} \right) \prod_{k=1}^l \frac{r''_k}{R''_k} = \tilde{a} \frac{r'r}{R'R} \prod_{k=1}^l \frac{r''_k}{R''_k} = a(\Gamma).$$

Therefore, formula (34) is proved in the case when l_1 is not incident to $v_1(\Gamma)$.

Suppose that $v = v_1(\Gamma)$. The tree Γ and stable trees $\tilde{\Gamma}$ such that $a(\tilde{\Gamma}, \Gamma) \neq 0$ are shown in Figure 6.




Figure 5: Stable trees $\tilde{\Gamma}$ such that $a(\tilde{\Gamma}, \Gamma) \neq 0$.




Figure 6: Stable tree Γ and stable trees $\tilde{\Gamma}$ such that $a(\tilde{\Gamma}, \Gamma) \neq 0$.

Let

$$\tilde{a} := a(\Gamma) / \left(\frac{r}{R} \prod_{j=1}^l \frac{r''_j}{R''_j} \right).$$

The coefficient $a(\tilde{\Gamma}, \Gamma)$ for the right tree in Figure 6 equals $-(\tilde{a} R''_j / R) \prod_{k=1}^l r''_k / R''_k$. So we compute

$$a(\Gamma) - \sum_{\tilde{\Gamma} \in \text{ST}_{g,n+1}^{m-1}} a(\tilde{\Gamma}, \Gamma) = \tilde{a} \left(\frac{r}{R} + \sum_{j=1}^l \frac{R''_j}{R} \right) \prod_{k=1}^l \frac{r''_k}{R''_k} = \tilde{a} \prod_{k=1}^l \frac{r''_k}{R''_k} = \frac{R}{r} a(\Gamma).$$

The lemma is proved. □

Lemma 4.7 Let $a_0, \dots, a_n, n \geq 1$, be integers with vanishing sum and $m \geq 2$. Then we have

$$\begin{aligned} \hat{A}^{g,m}(a_0, \dots, a_n) - a_0 \psi_0 \hat{A}^{g,m-1}(a_0, \dots, a_n) \\ = \sum_{\Gamma \in \text{ST}_{g,n+1}^m} \frac{2g-1+n}{r(v_1(\Gamma))} a(\Gamma) \lambda_g \text{DR}_{\Gamma}(a_0, \dots, a_n). \end{aligned}$$

Proof The proof is analogous to the proof of the previous lemma. □

Corollary 4.8 Let a_1, \dots, a_n with $n \geq 1$ be arbitrary integers and $m \geq 2$. Denote by $\pi: \bar{\mathcal{M}}_{g,n+1} \rightarrow \bar{\mathcal{M}}_{g,n}$ the forgetful map that forgets the first marked point. Then we

have

$$\begin{aligned}
 (35) \quad & A^{g,m}(a_1, \dots, a_n) - a_1 \psi_1 A^{g,m-1}(a_1, \dots, a_n) \\
 &= \sum_{\substack{\Gamma \in \text{ST}_{g,n+1}^m \\ v(l_1(\Gamma))=v_1(\Gamma) \\ g(v_1(\Gamma)) \geq 1}} \frac{2g-1+n}{r(v_1(\Gamma))} \frac{a(\Gamma)}{\sum a_i} \lambda_g \pi_* \text{DR}_\Gamma \left(-\sum a_i, a_1, \dots, a_n \right) \\
 &\quad + \widehat{A}^{g,m-1} \left(-\sum_{i=2}^n a_i, a_2, \dots, a_n \right).
 \end{aligned}$$

Proof The corollary is an elementary exercise that uses the two previous lemmas and the fact that

$$A^{g,m}(a_1, \dots, a_n) = \frac{1}{\sum a_i} \pi_* \widehat{A}^{g,m} \left(-\sum a_i, a_1, \dots, a_n \right). \quad \square$$

Proof of Proposition 4.5 Let $m := \sum d_i - 2g + 3$. Let us prove that

$$\begin{aligned}
 (36) \quad & \left. \frac{\partial}{\partial a_{n+1}} \pi_* A^{g,m}(a_1, \dots, a_{n+1}) \right|_{a_{n+1}=0} \\
 &= \begin{cases} 0 & \text{if } m = 1, \\ \pi_*(\psi_{n+1} A^{g,m-1}(a_1, \dots, a_n, 0)) & \text{if } m \geq 2. \end{cases}
 \end{aligned}$$

For $m = 1$ this equation immediately follows from Lemma 2.1. Suppose $m \geq 2$. Let us rewrite (35) in the way that is more suitable for us:

$$\begin{aligned}
 (37) \quad & A^{g,m}(a_1, \dots, a_{n+1}) - a_{n+1} \psi_{n+1} A^{g,m-1}(a_1, \dots, a_{n+1}) \\
 &= \sum_{\substack{\Gamma \in \text{ST}_{g,n+2}^m \\ v(l_{n+1}(\Gamma))=v_1(\Gamma) \\ g(v_1(\Gamma)) \geq 1}} \frac{2g+n}{r(v_1(\Gamma))} \frac{a(\Gamma)}{\sum a_i} \lambda_g \pi_{0*} \text{DR}_\Gamma \left(-\sum a_i, a_1, \dots, a_{n+1} \right) \\
 &\quad + \widehat{A}^{g,m-1} \left(-\sum_{i=1}^n a_i, a_1, \dots, a_n \right),
 \end{aligned}$$

where the map $\pi_0: \overline{\mathcal{M}}_{g,n+2} \rightarrow \overline{\mathcal{M}}_{g,n+1}$ forgets the first marked point. The last term on the right-hand side of this equation doesn't depend on a_{n+1} . Note also that, by Lemma 2.1, after applying the pushforward π_* each term in the sum on the right-hand side of (37) becomes divisible by a_{n+1}^2 . This proves (36).

Equation (36) immediately implies the statement of the proposition for $m = 1$. In the case $m \geq 2$, (36) yields

$$\begin{aligned}
 \pi_* A_{d_1, \dots, d_n, 1}^g &= \pi_*(\psi_{n+1} A_{d_1, \dots, d_n, 0}^g) \quad (\text{by Proposition 4.2}) \\
 &= \pi_*(\psi_{n+1} \pi^* A_{d_1, \dots, d_n}^g) \\
 &= (2g - 2 + n) A_{d_1, \dots, d_n}^g. \quad \square
 \end{aligned}$$

Proposition 4.9 Denote by $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map that forgets the last marked point. Then we have

$$(38) \quad \pi_*(B_{d_1, \dots, d_n, 1}^g) = \begin{cases} (2g - 2 + n)B_{d_1, \dots, d_n}^g & \text{if } \sum d_i > 2g - 2, \\ 0 & \text{if } \sum d_i = 2g - 2. \end{cases}$$

Proof Let $(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}$ be an admissible and stable complete tree with a power function

$$q: H_+^{\text{em}}(T) \rightarrow \mathbb{N},$$

as in Definition 2.4. We denote by $\epsilon: \{1, \dots, \text{deg}(T) - 1\} \rightarrow \mathbb{N}$ the function

$$\epsilon(k) := 2 \sum_{\substack{v \in V(T) \\ l(v) \leq k}} g(v) - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ l(h) = k}} q(h) - 2$$

measuring the distance to nonadmissibility at the level k . As in the proof of Proposition 4.3, we have two possible ways to add a new marking labelled by $n + 1$.

First, choose a vertex $v \in V(T)$. Let $C = (e_1^C, v_1^C, \dots, e_{\text{deg}(T)-l(v)}^C, v_{\text{deg}(T)-l(v)}^C, \sigma_{n+1})$ be a chain of weakly stable vertices with a new marking σ_{n+1} . Precisely, the edge e_1^C is attached to the vertex v_1^C , the edge e_k^C links the vertex v_{k-1}^C to v_k^C , and the leg σ_{n+1} is attached to the vertex $v_{\text{deg}(T)-l(v)}^C$. Moreover, every vertex is of genus 0 and contains two extra legs. We construct a tree T_v , obtained from T by gluing the edge e_1^C (and thus the chain C) to the vertex v . We have $H_+^{\text{em}}(T) \subset H_+^{\text{em}}(T_v)$ and we extend the power function q into a function $q_v: H_+^{\text{em}}(T_v) \rightarrow \mathbb{N}$ by taking

$$q_v(h_k^C) := 1 \quad \text{and} \quad q_v(\sigma_{n+1}) := 1,$$

where h_k^C is the half-edge in $H_+^{\text{em}}(T_v)$ contained in the edge e_k^C . It is easy to see that we get

$$(T_v, q_v) \in \Omega_{d_1, \dots, d_n, 1}^{B, g} \iff \epsilon(k) \geq 1 \text{ for all } k \in [l(v), \text{deg}(T) - 1].$$

In particular, when the vertex v is at the maximal level $\text{deg}(T)$, then the tree T_v is always admissible.

Second, choose a half-edge $h \in H_+^{\text{em}}(T)$ attached to the vertex v . We construct a tree $T_{(v,h)}$, obtained from T by adding an extra level between the levels $l(v)$ and $l(v) + 1$ of T as follows:

- denote by $h_0, \dots, h_m \in H_+^{\text{em}}(T)$ the half-edges of level $l(v)$, with $h_0 := h$;

- insert a pair (e_k, v_k) between the half-edge h_k and the vertex it is attached to, where $e_k = (h'_k, h''_k)$ is an edge and v_k is a vertex of genus 0;
- glue the half-edge h_1^C from the chain C to the vertex v_0 ;
- add $q(h_k) + 1$ extra legs to the vertex v_k for $0 \leq k \leq m$.

Therefore, the number of levels of the tree $T_{(v,h)}$ is $\text{deg}(T) + 1$, the vertex $v_0 \in V(T_{(v,h)})$ is the only strongly stable vertex at its level, and we have a natural inclusion $H_+^{\text{em}}(T_v) \subset H_+^{\text{em}}(T_{(v,h)})$. Then, we extend the power function q_v into a function $q_{(v,h)}: H_+^{\text{em}}(T_{(v,h)}) \rightarrow \mathbb{N}$ by taking

$$q_{(v,h)}(h'_k) := q(h_k).$$

We obtain

$$(T_{(v,h)}, q_{(v,h)}) \in \Omega_{d_1, \dots, d_n, 1}^{B,g} \iff \begin{cases} \epsilon(k) \geq 1 \text{ for all } k \in [l(v), \text{deg}(T) - 1] \\ \text{and } (l(v) \neq \text{deg}(T) \text{ or } \sum_{i=1}^n d_i = 2g - 2). \end{cases}$$

In particular, when the vertex v is at the maximal level $\text{deg}(T)$, the tree $T_{(v,h)}$ is admissible if and only if $d_1 + \dots + d_n = 2g - 2$.

Let $l_T \in [1, \text{deg}(T)]$ be the smallest integer such that

$$\epsilon(k) \geq 1 \quad \text{for all } k \in [l_T, \text{deg}(T) - 1].$$

When $d_1 + \dots + d_n > 2g - 2$ (resp. when $d_1 + \dots + d_n = 2g - 2$), the two constructions

$$(T, q, v) \mapsto (T_v, q_v) \quad \text{and} \quad (T, q, v, h) \mapsto (T_{(v,h)}, q_{(v,h)})$$

give a bijection from the set

$$\bigsqcup_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} \{v \in V(T) \mid l(v) \geq l_T\} \quad \sqcup \{(v, h) \in V(T) \times H_+^{\text{em}}(T) \mid h \rightarrow v, l_T \leq l(v) < \text{deg}(T)\}$$

(resp. the same set with the inequality $l_T \leq l(v) \leq \text{deg}(T)$) to the set $\Omega_{d_1, \dots, d_n, 1}^{B,g}$. Furthermore, we get the contributions

$$(39) \quad e_* \pi_*([T_v, q_v]) = (2g(v) - 2 + n(v) + q(v) + 1)e_*[T, q],$$

$$(40) \quad e_* \pi_*([T_{(v,h)}, q_{(v,h)}]) = (q(h) + 1)e_*[T, q],$$

where $q(v)$ denotes the value of the power function $q: H_+^{\text{em}}(T) \rightarrow \mathbb{N}$ at the (half-)edge linking the mother of the vertex v to the vertex v and $n(v)$ denotes the number of half-edges attached to the vertex v , without counting the extra legs. Thus, the total number of half-edges attached to the vertex v is indeed $n(v) + q(v) + 1$.

Finally, when $d_1 + \dots + d_n > 2g - 2$, we get

$$\begin{aligned}
 & \pi_*(B_{d_1, \dots, d_n, 1}^g) \\
 &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n, 1}^{B, g}} (-1)^{\deg(T)-1} \pi_* e_* [T, q] \\
 &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \left(\sum_{\substack{v \in V(T) \\ l(v) = \deg(T)}} \pi_* e_* [T_v, q_v] \right. \\
 & \quad \left. + \sum_{\substack{v \in V(T) \\ l_T \leq l(v) < \deg(T)}} \left(\pi_* e_* [T_v, q_v] - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} \pi_* e_* [T_{(v, h)}, q_{(v, h)}] \right) \right) \\
 &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} \left(\sum_{\substack{v \in V(T) \\ l(v) = \deg(T)}} e_* \pi_* [T_v, q_v] \right. \\
 & \quad \left. + \sum_{\substack{v \in V(T) \\ l_T \leq l(v) < \deg(T)}} \left(e_* \pi_* [T_v, q_v] - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} e_* \pi_* [T_{(v, h)}, q_{(v, h)}] \right) \right),
 \end{aligned}$$

where the minus sign in the second line of the second equality comes from the fact that the number of levels in the tree $T_{(v, h)}$ is $\deg(T) + 1$; the third equality comes from the relation $e \circ \pi = \pi \circ e$ among the forgetful maps. Using equations (39) and (40), we get

$$\begin{aligned}
 & \pi_*(B_{d_1, \dots, d_n, 1}^g) \\
 &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} e_* [T, q] \cdot \left(\sum_{\substack{v \in V(T) \\ l(v) = \deg(T)}} (2g(v) - 2 + n(v) + q(v) + 1) \right. \\
 & \quad \left. + \sum_{\substack{v \in V(T) \\ l_T \leq l(v) < \deg(T)}} \left(2g(v) - 1 + n(v) + q(v) - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} (q(h) + 1) \right) \right) \\
 &= \sum_{(T, q) \in \Omega_{d_1, \dots, d_n}^{B, g}} (-1)^{\deg(T)-1} e_* [T, q] \cdot \left(\sum_{\substack{v \in V(T) \\ l(v) = \deg(T)}} (2g(v) + n(v) - 1 + q(v)) \right. \\
 & \quad \left. + \sum_{\substack{v \in V(T) \\ l_T \leq l(v) < \deg(T)}} \left(2g(v) + q(v) - \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} q(h) \right) \right)
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_*[T, q] \\
 &\quad \cdot \left(2 \sum_{\substack{v \in V(T) \\ l(v) \geq l_T}} g(v) + n + \left(\sum_{\substack{v \in V(T) \\ l(v) \geq l_T}} q(v) - \sum_{\substack{v \in V(T) \\ l_T \leq l(v) < \deg(T)}} \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} q(h) \right) \right) \\
 &= \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_*[T, q] \cdot \left(2 \sum_{\substack{v \in V(T) \\ l(v) \geq l_T}} g(v) + n + \sum_{\substack{v \in V(T) \\ l(v) = l_T}} q(v) \right).
 \end{aligned}$$

We conclude using the equality $\epsilon(l_T - 1) = 0$:

$$\begin{aligned}
 &\pi_*(B_{d_1, \dots, d_n, 1}^g) \\
 &= \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_*[T, q] \cdot \left(2 \sum_{\substack{v \in V(T) \\ l(v) \geq l_T}} g(v) + n + 2 \sum_{\substack{v \in V(T) \\ l(v) < l_T}} g(v) - 2 \right) \\
 &= (2g - 2 + n) \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_*[T, q] = (2g - 2 + n) B_{d_1, \dots, d_n}^g.
 \end{aligned}$$

When $d_1 + \dots + d_n = 2g - 2$, we have the same sequence of equalities with the additional term

$$\begin{aligned}
 &- \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} \sum_{\substack{v \in V(T) \\ l(v) = \deg(T)}} \sum_{\substack{h \in H_+^{\text{em}}(T) \\ h \rightarrow v}} (-1)^{\deg(T)-1} e_* \pi_*[T_{(v,h)}, q_{(v,h)}] \\
 &= - \sum_{(T,q) \in \Omega_{d_1, \dots, d_n}^{B,g}} (-1)^{\deg(T)-1} e_*[T, q] \cdot (n + d_1 + \dots + d_n) \\
 &= -(2g - 2 + n) B_{d_1, \dots, d_n}^g,
 \end{aligned}$$

coming from the fact that $(T_{(v,h)}, q_{(v,h)}) \in \Omega_{d_1, \dots, d_n, 1}^{B,g}$ when $l(v) = \deg(T)$. □

4.6 Validity of the conjecture on $\mathcal{M}_{g,n}$

Let $g, n, m \geq 0$ be such that $2g - 2 + n > 0$ and denote by $\pi^{(m)}: \overline{\mathcal{M}}_{g,n+m} \rightarrow \overline{\mathcal{M}}_{g,n}$ the map forgetting the last m markings. By definition, the restriction of Conjecture 2.5 to $\mathcal{M}_{g,n}$ is the following statement:

Proposition 4.10 *The restriction of Conjecture 2.5 to $\mathcal{M}_{g,n}$ is true. Precisely, for all integers $d_1, \dots, d_{n+m} \geq 1$ such that*

$$d_1 + \dots + d_{n+m} > 2g - 2,$$

we have

$$(\pi_*^{(m)}(A_{d_1, \dots, d_{n+m}}^g))|_{\mathcal{M}_{g,n}} = (\pi_*^{(m)}(B_{d_1, \dots, d_{n+m}}^g))|_{\mathcal{M}_{g,n}} \in R^*(\mathcal{M}_{g,n}).$$

Proof Using Propositions 4.5 and 4.9, we can assume that $d_{n+1}, \dots, d_{n+m} \geq 2$. Furthermore, the Chow degree of the two classes in the statement is

$$\delta := d_1 + \dots + d_n + (d_{n+1} - 1) + \dots + (d_{n+m} - 1).$$

We get

$$\delta > 2g - 2 - m \quad \text{and} \quad \delta \geq n + m.$$

Summing these two inequalities yields

$$\delta > g + \frac{1}{2}n - 1 \geq g - 1.$$

We conclude with the following result from [17]:

$$R^p(\mathcal{M}_{g,n}) = 0 \quad \text{for all } p > g - 1. \quad \square$$

4.7 New expression for λ_g

Let us show that our conjectural relations (10) give a new formula for the class $\lambda_g \in R^g(\overline{\mathcal{M}}_g)$.

Let $g \geq 2$ and consider the class

$$A^{g,1}(a_1, \dots, a_{g-1}) = \lambda_g \frac{1}{\sum a_i} \text{DR}_g\left(-\widetilde{\sum a_i}, a_1, \dots, a_{g-1}\right).$$

Let $\pi: \overline{\mathcal{M}}_{g,g-1} \rightarrow \overline{\mathcal{M}}_g$ be the forgetful map that forgets all marked points. Then, by (5),

$$\pi_* A^{g,1}(a_1, \dots, a_{g-1}) = g! \lambda_g a_1^2 \cdots a_{g-1}^2 \sum a_i.$$

Thus,

$$\lambda_g = \frac{1}{g!} \pi_* A_{3,2,\dots,2}^g \in R^g(\overline{\mathcal{M}}_g).$$

So, Conjecture 2.5 implies that

$$(41) \quad \lambda_g = \frac{1}{g!} \pi_* B_{3,2,\dots,2}^g \in R^g(\overline{\mathcal{M}}_g).$$

We can easily see that the expression on the right-hand side of this equation is a linear combination of basic tautological classes $\xi_{\Gamma^*}(\gamma)$, where Γ is a tree. No such expressions for the class λ_g were known before. Let us write explicitly and prove the resulting formulas in genus 2 and 3.

4.7.1 Genus 2 We already wrote the expression for B_3^2 in (11). Pushing it forward to $\overline{\mathcal{M}}_2$ and dividing by 2, we get that Conjecture 2.5 implies

$$(42) \quad \lambda_2 = \frac{1}{2}\kappa_2 - \frac{1}{2} \textcircled{1} - \textcircled{1}^{\kappa_1}.$$

The relation $A_3^2 = B_3^2$ is proved in the appendix, so formula (42) is true.

4.7.2 Genus 3 We compute

$$\begin{aligned}
 B_{3,2}^3 = & \Psi_1^3 \psi_2^2 - 10 \textcircled{3} \textcircled{\psi^4} \textcircled{0} \textcircled{1} - 2 \textcircled{2} \textcircled{\psi} \textcircled{1} \textcircled{\psi^2} \textcircled{1} - 2 \textcircled{2} \textcircled{\psi} \textcircled{1} \textcircled{\psi^2} \textcircled{1} - 2 \textcircled{2} \textcircled{\psi} \textcircled{1} \textcircled{\psi^3} \textcircled{1} - 2 \textcircled{2} \textcircled{\psi^2} \textcircled{1} \textcircled{\psi^2} \textcircled{1} \\
 & - 3 \textcircled{2} \textcircled{\psi^2} \textcircled{1} \textcircled{\psi} \textcircled{1} - 3 \textcircled{2} \textcircled{\psi^2} \textcircled{1} \textcircled{\psi^2} \textcircled{1} - 2 \textcircled{2} \textcircled{1} \textcircled{\psi^2} \textcircled{1} - 1 \textcircled{1} \textcircled{2} \textcircled{\psi^3} \textcircled{1} - 1 \textcircled{1} \textcircled{2} \textcircled{\psi^2} \textcircled{1} \\
 & + 10 \textcircled{2} \textcircled{\psi} \textcircled{1} \textcircled{\psi^2} \textcircled{0} \textcircled{1} + 10 \textcircled{2} \textcircled{\psi^2} \textcircled{1} \textcircled{\psi} \textcircled{0} \textcircled{1} + 2 \textcircled{2} \textcircled{\psi} \textcircled{0} \textcircled{1} \textcircled{\psi^2} \textcircled{1} \\
 & + 1 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi} \textcircled{1} + 2 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi^2} \textcircled{1} + 1 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi^3} \textcircled{1} + 1 \textcircled{1} - \textcircled{1} \textcircled{\psi} \textcircled{1} \textcircled{\psi^2} \textcircled{1} \\
 & + 3 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi} \textcircled{1} + 3 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi^2} \textcircled{1} + 1 \textcircled{1} - \textcircled{1} \textcircled{\psi} \textcircled{1} \textcircled{\psi^2} \textcircled{1} \\
 & + 10 \textcircled{1} - \textcircled{2} \textcircled{\psi^3} \textcircled{0} \textcircled{1} - 10 \textcircled{1} - \textcircled{1} - \textcircled{1} \textcircled{\psi^2} \textcircled{0} \textcircled{1} - 10 \textcircled{1} - \textcircled{1} \textcircled{\psi} \textcircled{1} \textcircled{\psi} \textcircled{0} \textcircled{1} \\
 & - \textcircled{1} - \textcircled{1} - \textcircled{0} \textcircled{1} \textcircled{\psi^2} \textcircled{1}.
 \end{aligned}$$

Pushing forward this expression to $\overline{\mathcal{M}}_3$ and dividing it by 6, we get that Conjecture 2.5 implies

$$\begin{aligned}
 (43) \quad \lambda_3 = & -\frac{3}{2}\kappa_3 + \frac{1}{6}\kappa_1\kappa_2 + \frac{2}{3} \textcircled{2} \textcircled{\psi} \textcircled{1}^{\kappa_1} - \frac{1}{6} \textcircled{2} \textcircled{1}^{\kappa_1} - \frac{\kappa_1}{6} \textcircled{1} + \frac{5}{6} \textcircled{2} \textcircled{1}^{\kappa_2} - \frac{1}{6} \textcircled{2} \textcircled{1}^{\kappa_1^2} \\
 & - \frac{1}{3} \textcircled{1} - \textcircled{1} - \textcircled{1}^{\kappa_1}.
 \end{aligned}$$

Let us prove this equation. In [11], C Faber proved that the whole cohomology ring of $\overline{\mathcal{M}}_3$ is tautological and that it is generated by the classes

$$\delta_0 := \textcircled{2}, \quad \delta_1 := \textcircled{2} - \textcircled{1}, \quad \lambda_1, \quad \kappa_2.$$

There are 13 monomials of cohomological degree 6 in these classes. Faber proved that $\dim R^3(\overline{\mathcal{M}}_3) = 10$ and found 3 relations between the 13 monomials (see [11, page 407]). These relations easily imply that the following 10 classes form a basis in $R^3(\overline{\mathcal{M}}_3)$:

$$\delta_0^2 \lambda_1, \quad \delta_0 \delta_1 \lambda_1, \quad \delta_0 \lambda_1^2, \quad \delta_0 \kappa_2, \quad \delta_1^3, \quad \delta_1^2 \lambda_1, \quad \delta_1 \lambda_1^2, \quad \lambda_1^3, \quad \delta_1 \kappa_2, \quad \lambda_1 \kappa_2.$$

It is not hard to check that each of these 10 classes has the same intersection numbers with both sides of equation (43). So, formula (43) is true.

5 Restricted set of relations

In this section we show that the strong DR/DZ equivalence conjecture for semisimple cohomological field theories follows from the restricted set of relations (10), where $\sum d_i = 2g$ and $d_i \geq 1$.

Consider an arbitrary cohomological field theory in genus 0,

$$c_{0,n}: V^{\otimes n} \rightarrow H^*(\overline{\mathcal{M}}_{0,n}, \mathbb{C}).$$

Let $F_0(t_*)$ be its potential. Suppose we have a deformation $F(t_*, \varepsilon)$ of F_0 the form

$$F = F_0 + \sum_{g \geq 1} \varepsilon^{2g} F_g, \quad F_g \in \mathbb{C}[[t_*^*]].$$

Introduce formal power series $(w^{\text{sol}})^\alpha(x, t_*, \varepsilon)$ by

$$(w^{\text{sol}})^\alpha := \eta^{\alpha\mu} \frac{\partial^2 F}{\partial t_0^\mu \partial t_0^1} \Big|_{t_0^1 \mapsto t_0^1 + x},$$

and let $(w^{\text{sol}})_n^\alpha := \partial_x^n (w^{\text{sol}})^\alpha$. We will use the notation

$$\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g := \frac{\partial^n F_g}{\partial t_{d_1}^{\alpha_1} \cdots \partial t_{d_n}^{\alpha_n}} \Big|_{t_*^* = 0}.$$

A correlator $\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g$ will be called admissible if $\sum d_i \leq 2g$.

Lemma 5.1 *Suppose that the following conditions are satisfied:*

- we have the vanishing property

$$(44) \quad \langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g = 0 \quad \text{if } \sum d_i \leq 2g - 2;$$

- the string and the dilaton equations hold:

$$\begin{aligned} \frac{\partial F}{\partial t_0^1} &= \sum_{n \geq 0} t_{n+1}^\alpha \frac{\partial F}{\partial t_n^\alpha} + \frac{1}{2} \eta_{\alpha\beta} t_0^\alpha t_0^\beta, \\ \frac{\partial F}{\partial t_1^1} &= \varepsilon \frac{\partial F}{\partial \varepsilon} + \sum_{n \geq 0} t_n^\alpha \frac{\partial F}{\partial t_n^\alpha} - 2F + \varepsilon^2 \frac{N}{24}; \end{aligned}$$

- for each μ there exists a differential polynomial $\Omega_{1,1;\mu,0} \in \hat{\mathcal{A}}_w^{[0]}$ such that

$$(45) \quad \Omega_{1,1;\mu,0} |_{w_n^\gamma = (w^{\text{sol}})^\gamma_n} = \frac{\partial^2 F}{\partial t_1^1 \partial t_0^\mu} \Big|_{t_0^1 \mapsto t_0^1 + x}.$$

Then all correlators $\langle \tau_{d_1}(e_{\alpha_1}) \cdots \tau_{d_n}(e_{\alpha_n}) \rangle_g$ are uniquely determined by the admissible correlators.

Proof The topological recursion relation in genus zero implies that the primary correlators $\langle \tau_0(e_{\alpha_1}) \cdots \tau_0(e_{\alpha_n}) \rangle_0$ determine all correlators in genus zero. Denote by \mathcal{R}_d the subspace of $\mathbb{C}[[t_*^*]]$ defined by

$$\mathcal{R}_d := \left\{ \sum c_{\alpha_1, \dots, \alpha_n}^{d_1, \dots, d_n} \prod t_{d_i}^{\alpha_i} \in \mathbb{C}[[t_*^*]] \mid c_{\alpha_1, \dots, \alpha_n}^{d_1, \dots, d_n} = 0 \text{ if } \sum d_i \leq d - 1 \right\}.$$

From the string equation and the vanishing property (44) it follows that the function $(w^{\text{sol}})_n^\alpha |_{x=0}$ has the form

$$(46) \quad (w^{\text{sol}})_n^\alpha |_{x=0} = \delta^{\alpha,1} \delta_{n,1} + t_n^\alpha + r_n^\alpha + \sum_{g \geq 1} q_{g,n}^\alpha \varepsilon^{2g}, \quad r_n^\alpha \in \mathcal{R}_{n+1}, \quad q_{g,n}^\alpha \in \mathcal{R}_{2g+n}.$$

Introduce a grading in the ring $\mathbb{C}[[t_*^*]]$ by $\deg t_d^\alpha := d$ and consider the expansion

$$q_{g,n}^\alpha = \sum_{k \geq 0} q_{g,n,k}^\alpha, \quad \deg q_{g,n,k}^\alpha = 2g + n + k.$$

Note that the functions $q_{g,n,0}^\alpha$ and $q_{g,n,1}^\alpha$ are determined by the admissible correlators.

Let us show that $\partial^2 \Omega_{1,1;\mu,0} / \partial (w_x^1)^2 = 0$. Consider a monomial f of the form

$$(47) \quad f = \varepsilon^{2h} (w_x^1)^l w_{d_1}^{\alpha_1} \cdots w_{d_n}^{\alpha_n}, \quad l + \sum d_i = 2h, \quad (\alpha_i, d_i) \neq (1, 1).$$

Then property (46) implies that

$$(48) \quad (f|_{w_n^\alpha=(w^{\text{sol}})_n^\alpha})|_{x=0} = \varepsilon^{2h} (t_{d_1}^{\alpha_1} \cdots t_{d_n}^{\alpha_n} + h_0) + \sum_{\substack{k \geq 1 \\ m \geq 0}} \varepsilon^{2h+2k} h_{k,m},$$

where $h_0 \in \mathcal{R}_{2h-l+1}$, $\deg h_{k,m} = 2h + 2k - l + m$ and the functions $h_{k,0}$ and $h_{k,1}$ are completely determined by the admissible correlators. Suppose now that $\partial^2 \Omega_{1,1;\mu,0} / \partial(w_x^1)^2 \neq 0$. Consider monomials f as in (47) with the minimal h such that $l \geq 2$ and such that the coefficient of f in the differential polynomial $\Omega_{1,1;\mu,0}$ is nonzero. Let us choose such a monomial with l as big as possible. Then using (45) we can see that

$$\langle \tau_0(e_\mu) \prod \tau_{d_i}(e_{\alpha_i}) \rangle_h = \frac{1}{2h-1+n} \langle \tau_1(e_1) \tau_0(e_\mu) \prod \tau_{d_i}(e_{\alpha_i}) \rangle_h \neq 0.$$

This contradicts the vanishing property (44), because $\sum d_i = 2h - l \leq 2h - 2$. We conclude that $\partial^2 \Omega_{1,1;\mu,0} / \partial(w_x^1)^2 = 0$.

Let us now prove that the differential polynomial $\Omega_{1,1;\mu,0}$ is completely determined by the admissible correlators. Let

$$c_{g;\alpha_1,\dots,\alpha_n}^{d_1,\dots,d_n} := \text{Coef}_{\varepsilon^{2g}} \frac{\partial^n \Omega_{1,1;\mu,0}}{\partial w_{d_1}^{\alpha_1} \cdots \partial w_{d_n}^{\alpha_n}}, \quad \sum d_i = 2g.$$

Let us prove by induction on g that all coefficients $c_{g;\alpha_1,\dots,\alpha_n}^{d_1,\dots,d_n}$ are uniquely determined by the admissible correlators. We already know it for $g = 0$. Suppose $g \geq 1$. Using property (48) we see that if $(\beta_i, q_i) \neq (1, 1)$ and $\sum q_i = 2g - 1$, then the difference

$$\left\langle \tau_1(e_1) \tau_0(e_\mu) \prod \tau_{q_i}(e_{\beta_i}) \right\rangle_g - c_{g;1,\beta_1,\dots,\beta_m}^{1,q_1,\dots,q_m}$$

can be expressed in terms of the admissible correlators and the coefficients $c_{h;\gamma_1,\dots,\gamma_i}^{r_1,\dots,r_i}$ with $h < g$. Similarly, if $(\beta_i, q_i) \neq (1, 1)$ and $\sum q_i = 2g$, then the difference

$$\left\langle \tau_1(e_1) \tau_0(e_\mu) \prod \tau_{q_i}(e_{\beta_i}) \right\rangle_g - c_{g;\beta_1,\dots,\beta_m}^{q_1,\dots,q_m}$$

can be expressed in terms of the admissible correlators, the coefficients $c_{h;\gamma_1,\dots,\gamma_i}^{r_1,\dots,r_i}$ with $h < g$ and the coefficients $c_{g;1,\rho_1,\dots,\rho_j}^{1,s_1,\dots,s_j}$. We conclude that the differential polynomial $\Omega_{1,1;\mu,0}$ is completely determined by the admissible correlators.

We see that the functions $(w^{\text{sol}})^\alpha$ are solutions of the system of partial differential equations

$$\frac{\partial w^\alpha}{\partial t^1} = \eta^{\alpha\mu} \partial_x \Omega_{1,1;\mu,0}, \quad 1 \leq \alpha \leq N.$$

The argument from the proof of Proposition 5.2 in [5] shows that using this system together with the string and the dilaton equations for F one can uniquely reconstruct the whole solution $(w^{\text{sol}})^\alpha$ starting from the dispersionless part $(w^{\text{sol}})^\alpha|_{\varepsilon=0}$. After that using the string and the dilaton equations it is easy to reconstruct the whole function F . The lemma is proved. \square

Proposition 5.2 *Suppose that all relations (10) with $\sum d_i = 2g$ and $d_i \geq 1$ are true. Then the strong DR/DZ equivalence conjecture is true for any semisimple cohomological field theory.*

Proof Consider an arbitrary semisimple cohomological field theory. Propositions 4.2, 4.3, 4.5 and 4.9 imply that all relations (10) with $\sum d_i \leq 2g$ are true. Therefore, $(\prod \tau_{d_i}(e_{\alpha_i}))_g^{\text{DR}} = (\prod \tau_{d_i}(e_{\alpha_i}))_g^{\text{red}}$ if $\sum d_i \leq 2g$. Both potentials F^{DR} and F^{red} satisfy the assumptions of Lemma 5.1 (see [3]). Therefore, the lemma implies that $F^{\text{DR}} = F^{\text{red}}$. So the strong DR/DZ equivalence conjecture is true. \square

In the appendix we will prove that relations (10) are true when $g = 2$, $d_i \geq 1$ and $\sum d_i \leq 4$. Therefore, the strong DR/DZ equivalence conjecture is true for all semisimple cohomological field theories at the approximation up to genus 2.

Appendix Proof of the restricted genus 2 relations

Here we prove relations (10) when $g = 2$, $d_i \geq 1$ and $\sum d_i \leq 4$.

A.1 Relation $A_d^2 = B_d^2$

As we know from Section 4.2, in order to prove that $A_d^2 = B_d^2$ for any $d \geq 3$, it is sufficient to prove that $A_3^2 = B_3^2$. We have

$$B_3^2 = \psi_1^3 - \textcircled{1} - \textcircled{1} \psi^2,$$

and $A_3^2 = \text{Coef}_{a^4}(\lambda_2 \text{DR}_2(\widetilde{-a}, a))$. The group $H^2(\overline{\mathcal{M}}_{2,1}, \mathbb{Q})$ has dimension 3 and a basis is given by (see eg [15])

$$\psi_1 \quad \delta_0 := \textcircled{1} - \quad \delta_1 := \textcircled{1} - \textcircled{1} -$$

So it is sufficient to check that the intersection of the difference $A_3^2 - B_3^2$ with these three classes is zero. We compute

$$\int_{\text{DR}_2(\widetilde{-a}, a)} \lambda_2 \psi_1 = \frac{1}{1152} a^4 \implies \int_{\overline{\mathcal{M}}_{2,1}} A_3^2 \psi_1 = \frac{1}{1152},$$

$$\int_{\text{DR}_2(\tilde{-a},a)} \lambda_2 \delta_0 = 0 \quad \Rightarrow \quad \int_{\overline{\mathcal{M}}_{2,1}} A_3^2 \delta_0 = 0,$$

$$\int_{\text{DR}_2(\tilde{-a},a)} \lambda_2 \delta_1 = \frac{1}{576} a^4 \quad \Rightarrow \quad \int_{\overline{\mathcal{M}}_{2,1}} A_3^2 \delta_1 = \frac{1}{576},$$

and

$$\int_{\overline{\mathcal{M}}_{2,1}} B_3^2 \psi_1 = \int_{\overline{\mathcal{M}}_{2,1}} \psi_1^4 = \frac{1}{1152},$$

$$\int_{\overline{\mathcal{M}}_{2,1}} B_3^2 \delta_0 = \text{diagram} - \text{diagram} = 0,$$

$$\int_{\overline{\mathcal{M}}_{2,1}} B_3^2 \delta_1 = \text{diagram} = \frac{1}{576}.$$

Thus, $A_3^2 = B_3^2$.

A.2 Relation $A_{2,1}^2 = B_{2,1}^2$

We have

$$(49) \quad B_{2,1}^2 = \psi_1^2 \psi_2 - 3 \text{diagram} - \text{diagram} - \text{diagram} + 3 \text{diagram}.$$

In [15], E Getzler proved that $H^*(\overline{\mathcal{M}}_{2,2}, \mathbb{Q}) = R^*(\overline{\mathcal{M}}_{2,2})$. Moreover, he proved that the group $R^2(\overline{\mathcal{M}}_{2,2})$ has dimension 14 with a basis given by

$$(50) \quad \begin{aligned} \delta_{22} &:= \text{diagram} & \delta_{11|1} &:= \text{diagram} & \delta_{111|1} &:= \text{diagram} \\ \delta_{11|2} &:= \text{diagram} & \delta_{111|2} &:= \text{diagram} & \delta_{01|1} &:= \text{diagram} \\ \delta_{01|1} &:= \text{diagram} & \delta_{01|2} &:= \text{diagram} & \delta_{01|2} &:= \text{diagram} \\ \delta_{0|1} &:= \text{diagram} & \delta_{0|1} &:= \text{diagram} & \delta_{0|2} &:= \text{diagram} \\ \delta_{0|12} &:= \text{diagram} & \delta_{00} &:= \text{diagram} \end{aligned}$$

We compute

$$\begin{aligned}
 & \int_{\text{DR}_2(\overline{-a_1-a_2, a_1, a_2})} \lambda_2 \delta_{22} \\
 &= \int_{\text{DR}_2(\overline{-a_1-a_2, a_1+a_2})} \lambda_2 \psi_1 = \frac{1}{1152} (a_1 + a_2)^4 \implies \int_{\overline{\mathcal{M}}_{2,2}} A_{2,1}^2 \delta_{22} = \frac{1}{384}, \\
 & \int_{\text{DR}_2(\overline{-a_1-a_2, a_1, a_2})} \lambda_2 \delta_{11|} = 0 \implies \int_{\overline{\mathcal{M}}_{2,2}} A_{2,1}^2 \delta_{11|} = 0, \\
 & \int_{\text{DR}_2(\overline{-a_1-a_2, a_1, a_2})} \lambda_2 \delta_{11|1} = \frac{1}{576} a_1^2 (a_1 + a_2)^2 \implies \int_{\overline{\mathcal{M}}_{2,2}} A_{2,1}^2 \delta_{11|1} = \frac{1}{576}, \\
 & \int_{\text{DR}_2(\overline{-a_1-a_2, a_1, a_2})} \lambda_2 \delta_{11|2} = \frac{1}{576} a_2^2 (a_1 + a_2)^2 \implies \int_{\overline{\mathcal{M}}_{2,2}} A_{2,1}^2 \delta_{11|2} = 0, \\
 & \int_{\text{DR}_2(\overline{-a_1-a_2, a_1, a_2})} \lambda_2 \delta_{11|12} = \frac{1}{576} (a_1 + a_2)^4 \implies \int_{\overline{\mathcal{M}}_{2,2}} A_{2,1}^2 \delta_{11|12} = \frac{1}{192}.
 \end{aligned}$$

Since $\lambda_g|_{\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}^{\text{ct}}} = 0$, the intersections of all remaining 9 classes from (50) with $A_{2,1}^2$ are equal to zero. It is not hard to compute the intersections of the class $B_{2,1}^2$ with the classes from (50) and see that they agree with what we have just computed for $A_{2,1}^2$. Thus, $A_{2,1}^2 = B_{2,1}^2$.


A.3 Relation $A_{1,1,1}^2 = B_{1,1,1}^2$

We have

$$\begin{aligned}
 (51) \quad B_{1,1,1}^2 &= \psi_1 \psi_2 \psi_3 - 2 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) - 6 \left(\begin{array}{c} \psi^2 \\ \circ \\ \psi \end{array} \right) - 2 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) - \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) \\
 &+ 4 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) + 2 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) + 2 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) \\
 &+ 6 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) + 2 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) - 4 \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right).
 \end{aligned}$$

Introduce the notation

$$\begin{aligned}
 \alpha_1 &:= \left(\begin{array}{c} \circ \\ \psi \end{array} \right) & \alpha_2 &:= \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) & \alpha_3 &:= \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) \\
 \alpha_4 &:= \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) & \alpha_5 &:= \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right) & \alpha_6 &:= \left(\begin{array}{c} \psi \\ \circ \\ \psi \end{array} \right)
 \end{aligned}$$



Denote by L' the subspace of $R^3(\overline{\mathcal{M}}_{2,3})$ spanned by boundary strata $\xi_{\Gamma^*}(1)$, where the first Betti number of a stable graph Γ is equal to 2. The symmetric group S_3 acts on $\overline{\mathcal{M}}_{2,3}$ by permutations of marked points. This action induces an action on $R^*(\overline{\mathcal{M}}_{2,3})$. Define a map $\text{Sym}: R^*(\overline{\mathcal{M}}_{2,3}) \rightarrow R^*(\overline{\mathcal{M}}_{2,3})$ by

$$\text{Sym}(\alpha) := \frac{1}{3!} \sum_{\sigma \in S_3} \sigma \alpha, \quad \alpha \in R^*(\overline{\mathcal{M}}_{2,3}).$$

Let $L := \text{Sym}(L') \subset R^3(\overline{\mathcal{M}}_{2,3})^{S_3}$. For two classes $\alpha, \beta \in R^3(\overline{\mathcal{M}}_{2,3})$ we will write $\alpha \stackrel{\text{mod } L}{=} \beta$ if $\alpha - \beta \in L$. Using the formulas for $\psi_1^2 \in R^2(\overline{\mathcal{M}}_{2,1})$ and $\psi_1 \psi_2 \in R^2(\overline{\mathcal{M}}_{2,2})$ from [15] and also the topological recursion relations in genus 0 and 1, after a long computation we obtain

$$\begin{aligned} B_{1,1,1}^2 \stackrel{\text{mod } L}{=} & -\frac{3}{5}\alpha_1 + \frac{3}{10}\alpha_2 + \frac{4}{5}\alpha_3 + \frac{4}{15}\alpha_4 + \frac{1}{10}\alpha_5 - \frac{4}{5}\alpha_6 + \frac{1}{360}a_{1,2,3}^{0,1,2} - \frac{1}{90}a_{1,2,3}^{0,2,1} \\ & - \frac{1}{24}a_{1,2,3}^{0,3,0} + \frac{1}{40}a_{1,2,3}^{1,1,1} + \frac{1}{180}a_{1,2,3}^{1,2,0} - \frac{1}{120}a_{1,2,3}^{2,1,0} - \frac{5}{48}a_{1,4,1}^{0,0,3} - \frac{1}{80}a_{1,4,1}^{0,1,2} \\ & - \frac{1}{144}a_{1,4,1}^{1,0,2} + \frac{1}{80}a_{2,3,1}^{0,0,3} - \frac{7}{240}a_{2,3,1}^{0,1,2} + \frac{1}{240}a_{2,3,1}^{1,0,2} + \frac{1}{90}a_{3,2,1}^{0,1,2} + \frac{7}{90}b_{1,3,2}^{0,1,2} \\ & + \frac{1}{45}b_{1,3,2}^{0,2,1} + \frac{1}{90}b_{1,3,2}^{1,1,1} - \frac{11}{30}b_{2,3,1}^{0,0,3} - \frac{1}{10}b_{2,3,1}^{0,1,2} - \frac{1}{90}b_{2,3,1}^{1,0,2} - \frac{1}{30}b_{3,2,1}^{0,1,2} \\ & + \frac{1}{5}c_{2,2,2}^{0,2,1} + \frac{2}{45}c_{2,2,2}^{1,1,1}. \end{aligned}$$

On the other hand, a direct computation using Hain's formula gives

$$\begin{aligned} \text{Coef}_{a_1 a_2 a_3} \left(\frac{1}{a_1 + a_2 + a_3} \text{DR}_2(\overline{-a_1 - a_2 - a_3, a_1, a_2, a_3}) \right) \Big|_{\mathcal{M}_{2,3}^{\text{ct}}} \\ = (\psi_1 + \psi_2 + \psi_3) - 6 \textcircled{2} - \textcircled{0} - 3 \textcircled{2} - \textcircled{0} - \frac{6}{5} \textcircled{1} - \textcircled{1} - \frac{1}{5} \textcircled{1} - \textcircled{1}. \end{aligned}$$

In this computation one should use that (see eg [15])

$$\kappa_1 = \frac{7}{10} \textcircled{1} - \textcircled{1} + \frac{1}{10} \textcircled{\textcircled{1}} \in R^1(\overline{\mathcal{M}}_2).$$

Using the formula [19]

$$\lambda_2 = \frac{1}{960} \textcircled{\textcircled{0}} + \frac{1}{240} \textcircled{1} - \textcircled{0} \in R^2(\overline{\mathcal{M}}_2),$$

we obtain the following formula for the class $A^2_{1,1,1}$:

$$A^2_{1,1,1} \stackrel{\text{mod } L}{=} \frac{1}{120} a_{1,2,3}^{0,1,2} + \frac{1}{60} a_{1,2,3}^{0,2,1} + \frac{1}{40} a_{1,2,3}^{0,3,0} + \frac{1}{120} a_{1,2,3}^{1,1,1} + \frac{1}{60} a_{1,2,3}^{1,2,0} + \frac{1}{120} a_{1,2,3}^{2,1,0} \\ - \frac{1}{80} a_{1,4,1}^{0,0,3} - \frac{1}{240} a_{1,4,1}^{0,1,2} - \frac{1}{240} a_{1,4,1}^{1,0,2} - \frac{1}{80} a_{2,3,1}^{0,0,3} - \frac{1}{240} a_{2,3,1}^{0,1,2} - \frac{1}{240} a_{2,3,1}^{1,0,2}.$$

Thus,

$$B^2_{1,1,1} - A^2_{1,1,1} \stackrel{\text{mod } L}{=} -\frac{3}{5} \alpha_1 + \frac{3}{10} \alpha_2 + \frac{4}{5} \alpha_3 + \frac{4}{15} \alpha_4 + \frac{1}{10} \alpha_5 - \frac{4}{5} \alpha_6 - \frac{1}{180} a_{1,2,3}^{0,1,2} \\ - \frac{1}{36} a_{1,2,3}^{0,2,1} - \frac{1}{15} a_{1,2,3}^{0,3,0} + \frac{1}{60} a_{1,2,3}^{1,1,1} - \frac{1}{90} a_{1,2,3}^{1,2,0} - \frac{1}{60} a_{1,2,3}^{2,1,0} \\ - \frac{11}{120} a_{1,4,1}^{0,0,3} - \frac{1}{120} a_{1,4,1}^{0,1,2} - \frac{1}{360} a_{1,4,1}^{1,0,2} + \frac{1}{40} a_{2,3,1}^{0,0,3} - \frac{1}{40} a_{2,3,1}^{0,1,2} \\ + \frac{1}{120} a_{2,3,1}^{1,0,2} + \frac{1}{90} a_{3,2,1}^{0,1,2} + \frac{7}{90} b_{1,3,2}^{0,1,2} + \frac{1}{45} b_{1,3,2}^{0,2,1} + \frac{1}{90} b_{1,3,2}^{1,1,1} \\ - \frac{11}{30} b_{2,3,1}^{0,0,3} - \frac{1}{10} b_{2,3,1}^{0,1,2} - \frac{1}{90} b_{2,3,1}^{1,0,2} - \frac{1}{30} b_{3,2,1}^{0,1,2} + \frac{1}{5} c_{2,2,2}^{0,2,1} + \frac{2}{45} c_{2,2,2}^{1,1,1}.$$

The famous Getzler relation [14] says that

$$(52) \quad \gamma := \textcircled{0} - \textcircled{1} - \textcircled{0} - \frac{1}{3} \textcircled{1} - \textcircled{0} - \textcircled{0} - \frac{1}{6} \textcircled{1} - \textcircled{0} - \textcircled{0} + \frac{1}{2} \textcircled{1} - \textcircled{0} - \textcircled{0} \\ + \frac{1}{24} \textcircled{\textcircled{0}} + \frac{1}{24} \textcircled{\textcircled{0}} - \frac{1}{12} \textcircled{\textcircled{0}} \\ = 0 \in R^2(\overline{\mathcal{M}}_{1,4}).$$

We will adopt the following notation. Suppose $g_1, g_2 \geq 0$ and let i_1, \dots, i_k and j_1, \dots, j_l be two lists of integers such that $\{i_1, \dots, i_k, j_1, \dots, j_l\} = \{1, 2, \dots, k+l\}$. Consider the moduli spaces $\overline{\mathcal{M}}_{g_1, k+1}$ and $\overline{\mathcal{M}}_{g_2, l+1}$, but let us label the marked points on curves from $\overline{\mathcal{M}}_{g_1, k+1}$ and $\overline{\mathcal{M}}_{g_2, l+1}$ by the numbers $i_1, \dots, i_k, k+l+1$ and $j_1, \dots, j_l, k+l+2$, respectively. Denote by

$$\mathfrak{gl}_{i_1, \dots, i_k | j_1, \dots, j_l}^{g_1 | g_2} : \overline{\mathcal{M}}_{g_1, k+1} \times \overline{\mathcal{M}}_{g_2, l+1} \rightarrow \overline{\mathcal{M}}_{g_1+g_2, k+l}$$

the gluing map that glues the marked points labelled by $k + l + 1$ and $k + l + 2$. From Getzler's relation we obtain

$$(53) \quad (\text{gl}_{|1,2,3,4}^{1,1})_*([\overline{\mathcal{M}}_{1,1}] \times \gamma) = \frac{1}{3}\alpha_1 - \frac{1}{6}\alpha_2 - \alpha_3 - \frac{1}{3}\alpha_4 + \frac{1}{2}\alpha_5 + \alpha_6 + \frac{1}{24}a_{1,2,3}^{0,2,1} \\ + \frac{1}{24}a_{1,2,3}^{0,3,0} + \frac{1}{24}a_{1,4,1}^{0,0,3} - \frac{1}{12}b_{1,3,2}^{0,1,2} = 0 \in R^3(\overline{\mathcal{M}}_{2,3}).$$

Notice that the WDVV relation on $\overline{\mathcal{M}}_{0,5}$ implies that $-\frac{1}{3}\alpha_1 + \frac{1}{6}\alpha_2 + \frac{1}{2}\alpha_5 = 0$. Using this observation and expressing the class α_6 via formula (53), we get

$$(54) \quad B_{1,1,1}^2 - A_{1,1,1}^2 \stackrel{\text{mod } L}{=} -\frac{1}{180}a_{1,2,3}^{0,1,2} + \frac{1}{180}a_{1,2,3}^{0,2,1} - \frac{1}{30}a_{1,2,3}^{0,3,0} + \frac{1}{60}a_{1,2,3}^{1,1,1} - \frac{1}{90}a_{1,2,3}^{1,2,0} \\ - \frac{1}{60}a_{1,2,3}^{2,1,0} - \frac{7}{120}a_{1,4,1}^{0,0,3} - \frac{1}{120}a_{1,4,1}^{0,1,2} - \frac{1}{360}a_{1,4,1}^{1,0,2} + \frac{1}{40}a_{2,3,1}^{0,0,3} \\ - \frac{1}{40}a_{2,3,1}^{0,1,2} + \frac{1}{120}a_{2,3,1}^{1,0,2} + \frac{1}{90}a_{3,2,1}^{0,1,2} + \frac{1}{90}b_{1,3,2}^{0,1,2} + \frac{1}{45}b_{1,3,2}^{0,2,1} \\ + \frac{1}{90}b_{1,3,2}^{1,1,1} - \frac{11}{30}b_{2,3,1}^{0,0,3} - \frac{1}{10}b_{2,3,1}^{0,1,2} - \frac{1}{90}b_{2,3,1}^{1,0,2} - \frac{1}{30}b_{3,2,1}^{0,1,2} \\ + \frac{1}{5}c_{2,2,2}^{0,2,1} + \frac{2}{45}c_{2,2,2}^{1,1,1}.$$

Let $\pi: \overline{\mathcal{M}}_{1,5} \rightarrow \overline{\mathcal{M}}_{1,4}$ be the forgetful map that forgets the fifth marked point and $\text{gl}_{1,5}: \overline{\mathcal{M}}_{1,5} \rightarrow \overline{\mathcal{M}}_{2,3}$ be the gluing map that glues the first and the fifth marked points. Then from Getzler's relation (52) we obtain

$$\text{gl}_{1,5}^*(\pi^* \gamma) \stackrel{\text{mod } L}{=} \frac{1}{2}a_{1,2,3}^{0,1,2} - \frac{1}{6}a_{1,2,3}^{0,2,1} - \frac{1}{3}a_{1,2,3}^{1,1,1} + \frac{1}{2}a_{1,4,1}^{0,0,3} - \frac{1}{6}a_{1,4,1}^{0,1,2} - \frac{1}{3}a_{1,4,1}^{1,0,2} + a_{2,3,1}^{0,1,2} \\ - \frac{1}{3}a_{3,2,1}^{0,1,2} + \frac{1}{2}b_{1,3,2}^{0,0,3} + \frac{1}{3}b_{1,3,2}^{0,1,2} - \frac{1}{6}b_{1,3,2}^{0,2,1} - \frac{1}{3}b_{1,3,2}^{1,0,2} - \frac{1}{3}b_{1,3,2}^{1,1,1} \\ + \frac{1}{2}b_{2,3,1}^{0,0,3} - \frac{1}{2}b_{2,3,1}^{0,1,2} - \frac{1}{3}b_{2,3,1}^{1,0,2} + b_{3,2,1}^{0,1,2} + c_{2,2,2}^{0,2,1} - \frac{2}{3}c_{2,2,2}^{1,1,1} \in L.$$

We can obtain another consequence from (52). Let $\text{gl}_{1,2}: \overline{\mathcal{M}}_{1,5} \rightarrow \overline{\mathcal{M}}_{2,3}$ be the gluing map that glues the first two marked points. Then Getzler's relation implies that

$$\text{Sym}(\text{gl}_{1,2}^*(\pi^* \gamma)) \stackrel{\text{mod } L}{=} \frac{1}{3}a_{1,2,3}^{0,1,2} + \frac{5}{18}a_{1,2,3}^{0,2,1} - \frac{1}{6}a_{1,2,3}^{0,3,0} + \frac{1}{18}a_{1,2,3}^{1,1,1} - \frac{5}{18}a_{1,2,3}^{1,2,0} - \frac{2}{9}a_{1,2,3}^{2,1,0} \\ - \frac{1}{6}a_{1,4,1}^{0,0,3} - \frac{1}{18}a_{1,4,1}^{0,1,2} - \frac{1}{18}a_{1,4,1}^{1,0,2} + a_{2,3,1}^{0,0,3} + \frac{1}{3}a_{2,3,1}^{0,1,2} + \frac{1}{3}a_{2,3,1}^{1,0,2} \\ + b_{1,3,2}^{0,0,3} + \frac{1}{9}b_{1,3,2}^{0,1,2} - \frac{2}{9}b_{1,3,2}^{0,2,1} - \frac{1}{9}b_{1,3,2}^{1,0,2} - \frac{1}{3}b_{1,3,2}^{1,1,1} - \frac{4}{9}b_{1,3,2}^{2,0,1} \\ - \frac{2}{3}b_{2,3,1}^{0,0,3} - \frac{2}{9}b_{2,3,1}^{0,1,2} - \frac{2}{9}b_{2,3,1}^{1,0,2} + \frac{4}{9}c_{2,2,2}^{0,2,1} + \frac{2}{9}c_{2,2,2}^{1,1,1} \in L.$$

Adding $\frac{1}{30} \text{gl}_{1,5^*}(\pi^* \gamma) - \frac{1}{40} \text{Sym}(\text{gl}_{1,2^*}(\pi^* \gamma))$ to the right-hand side of (54), we get


$$(55) \quad B_{1,1,1}^2 - A_{1,1,1}^2 \stackrel{\text{mod } L}{=} \frac{1}{360} a_{1,2,3}^{0,1,2} - \frac{1}{144} a_{1,2,3}^{0,2,1} - \frac{7}{240} a_{1,2,3}^{0,3,0} + \frac{1}{240} a_{1,2,3}^{1,1,1} - \frac{1}{240} a_{1,2,3}^{1,2,0} \\ - \frac{1}{90} a_{1,2,3}^{2,1,0} - \frac{3}{80} a_{1,4,1}^{0,0,3} - \frac{1}{80} a_{1,4,1}^{0,1,2} - \frac{1}{80} a_{1,4,1}^{1,0,2} - \frac{1}{120} b_{1,3,2}^{0,0,3} \\ + \frac{7}{360} b_{1,3,2}^{0,1,2} + \frac{1}{45} b_{1,3,2}^{0,2,1} - \frac{1}{120} b_{1,3,2}^{1,0,2} + \frac{1}{120} b_{1,3,2}^{1,1,1} + \frac{1}{90} b_{1,3,2}^{2,0,1} \\ - \frac{1}{3} b_{2,3,1}^{0,0,3} - \frac{1}{9} b_{2,3,1}^{0,1,2} - \frac{1}{60} b_{2,3,1}^{1,0,2} + \frac{2}{9} c_{2,2,2}^{0,2,1} + \frac{1}{60} c_{2,2,2}^{1,1,1}.$$

The WDVV relations on $\overline{\mathcal{M}}_{0,4}$, $\overline{\mathcal{M}}_{0,5}$ and $\overline{\mathcal{M}}_{0,6}$ imply that

$$c_{2,2,2}^{1,1,1} = b_{2,3,1}^{1,0,2}, \\ c_{2,2,2}^{0,2,1} = \frac{3}{2} b_{2,3,1}^{0,0,3} + \frac{1}{2} b_{2,3,1}^{0,1,2}, \\ a_{1,2,3}^{2,1,0} = b_{1,3,2}^{2,0,1}, \\ b_{1,3,2}^{1,0,2} = \frac{1}{2} a_{1,2,3}^{1,1,1} + a_{1,2,3}^{1,2,0} - \frac{1}{2} b_{1,3,2}^{1,1,1}, \\ b_{1,3,2}^{1,1,1} = a_{1,2,3}^{1,2,0} + a_{1,4,1}^{1,0,2}, \\ b_{1,3,2}^{0,0,3} = \frac{1}{3} a_{1,2,3}^{0,1,2} + \frac{2}{3} a_{1,2,3}^{0,2,1} + a_{1,2,3}^{0,3,0} - \frac{2}{3} b_{1,3,2}^{0,1,2} - \frac{1}{3} b_{1,3,2}^{0,2,1}, \\ b_{1,3,2}^{0,2,1} = \frac{1}{2} a_{1,2,3}^{0,2,1} + \frac{3}{2} a_{1,2,3}^{0,3,0} + \frac{3}{2} a_{1,4,1}^{0,0,3} + \frac{1}{2} a_{1,4,1}^{0,1,2} - b_{1,3,2}^{0,1,2}.$$

Using these relations, one can easily check that the right-hand side of (55) is zero. We conclude that $B_{1,1,1}^2 - A_{1,1,1}^2 \in L$.

It is easy to see that the space L is spanned by the classes




The WDVV relations on $\overline{\mathcal{M}}_{0,7}$ give the relations

$$\beta_2 - \beta_4 + \beta_7 - \beta_8 = 0, \\ \beta_1 + \beta_2 + 2\beta_3 + \frac{1}{3}\beta_5 + \frac{1}{3}\beta_7 - \frac{4}{3}\beta_8 - \frac{2}{3}\beta_9 = 0.$$

Therefore, $\dim L \leq 7$. On the other hand, in Table 1 we compute the intersection matrix of the classes β_1, \dots, β_9 with the seven classes $\psi_1^3, \psi_1^2\psi_2, \psi_1\psi_2\psi_3, \kappa_3,$

$\kappa_1\kappa_2$, $\psi_1\kappa_2$ and $\psi_1^2\delta$, where



This matrix is nondegenerate, so $\dim L = 7$. Thus, in order to prove that $A_{1,1,1}^2 = B_{1,1,1}^2$ it is sufficient to check that the intersections of $A_{1,1,1}^2 - B_{1,1,1}^2$ with the classes ψ_1^3 , $\psi_1^2\psi_2$, $\psi_1\psi_2\psi_3$, κ_3 , $\kappa_1\kappa_2$, $\psi_1\kappa_2$ and $\psi_1^2\delta$ are zero. This is a simple direct computation. The relation $A_{1,1,1}^2 = B_{1,1,1}^2$ is proved.

| | β_1 | β_2 | β_3 | β_4 | β_5 | β_6 | β_7 | β_8 | β_9 |
|----------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| ψ_1^3 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 2 |
| $\psi_1^2\psi_2$ | 0 | 3 | 0 | 3 | 0 | 1 | 1 | 1 | 3 |
| $\psi_1\psi_2\psi_3$ | 0 | 6 | 0 | 6 | 0 | 0 | 6 | 6 | 0 |
| κ_3 | 0 | 1 | 0 | 1 | 3 | 0 | 0 | 0 | 3 |
| $\kappa_1\kappa_2$ | 1 | 9 | 1 | 9 | 27 | 3 | 3 | 3 | 27 |
| $\psi_1\kappa_2$ | 1 | 4 | 0 | 4 | 4 | 2 | 1 | 1 | 8 |
| $\psi_1^2\delta$ | 0 | -2 | 1 | 0 | 2 | 0 | 2 | 0 | 2 |

Table 1: Intersection matrix of β_1, \dots, β_9 with $\psi_1^3, \psi_1^2\psi_2, \psi_1\psi_2\psi_3, \kappa_3, \kappa_1\kappa_2, \psi_1\kappa_2$ and $\psi_1^2\delta$.

A.4 Relations $A_{3,1}^2 = B_{3,1}^2$ and $A_{2,2}^2 = B_{2,2}^2$

Suppose $g, n \geq 1$ and $a_1, \dots, a_n \in \mathbb{Z}$. Let $a := \sum a_i$. The following formula is the particular case of Corollary 4.8 when $m = 2$:

$$\begin{aligned}
 (56) \quad & A^{g,2}(a_1, \dots, a_n) - a_1\psi_1 A^{g,1}(a_1, \dots, a_n) \\
 &= \lambda_g \text{DR}_g(a_1 - a, a_2, \dots, a_n) \\
 &+ \lambda_g \sum_{\substack{g_1 \geq 1, g_2 \geq 0 \\ g_1 + g_2 = g}} \sum_{\substack{I \sqcup J = \{1, \dots, n\} \\ 1 \in I \\ 2g_2 - 1 + |J| > 0}} \frac{a_J}{a} \text{DR}_{g_1}(\widetilde{-a}, A_I, a_J) \boxtimes_1 \text{DR}_{g_2}(A_J, -a_J).
 \end{aligned}$$

Let us prove now that $A_{d_1, d_2}^2 = B_{d_1, d_2}^2$, where $(d_1, d_2) = (3, 1)$ or $(d_1, d_2) = (2, 2)$. By (56), we have

$$\begin{aligned}
 & A_{d_1, d_2}^2 - \psi_1 A_{d_1-1, d_2}^2 \\
 &= \text{Coef}_{a_1^{d_1} a_2^{d_2}} \left(\frac{1}{a_1 + a_2} \lambda_2 \text{DR}_1(\widetilde{-a_1 - a_2}, a_1, a_2) \boxtimes_1 \text{DR}_1(a_2, -a_2) \right) \\
 &= \text{Coef}_{a_1^{d_1} a_2^{d_2}} \left(\frac{(a_1 + a_2)a_2^3}{576} \left(\text{Diagram} \right) \right) = 0.
 \end{aligned}$$

On the other hand, it is easy to compute that

$$\begin{aligned}
 B_{3,1}^2 &= \psi_1^3 \psi_2 - \textcircled{1} - \textcircled{1} \begin{array}{c} \nearrow \psi^2 1 \\ \searrow \psi 2 \end{array} - \textcircled{1} - \textcircled{1} \begin{array}{c} \nearrow \psi^3 1 \\ \searrow \psi 2 \end{array}, \\
 B_{2,2}^2 &= \psi_1^2 \psi_2^2 - \textcircled{1} - \textcircled{1} \begin{array}{c} \nearrow \psi^2 1 \\ \searrow \psi 2 \end{array} - \textcircled{1} - \textcircled{1} \begin{array}{c} \nearrow \psi 1 \\ \searrow \psi^2 2 \end{array}.
 \end{aligned}$$

Comparing these expressions with formula (49), we can easily see that $B_{3,1}^2 = \psi_1 B_{2,1}^2$ and $B_{2,2}^2 = \psi_1 B_{1,2}^2$. Since the relation $A_{2,1}^2 = B_{2,1}^2$ is already checked, the relations $A_{3,1}^2 = B_{3,1}^2$ and $A_{2,2}^2 = B_{2,2}^2$ are now also proved.

A.5 Relation $A_{2,1,1}^2 = B_{2,1,1}^2$

Using (56), we compute $A_{2,1,1}^2 - \psi_1 A_{1,1,1}^2$ as

$$\begin{aligned}
 (57) \quad & \sum_{\substack{I \sqcup J = \{1,2,3\} \\ 1 \in I, |J| \geq 1}} \text{Coef}_{a_1^2 a_2 a_3} \left(\frac{a_J}{\sum a_i} \lambda_2 \text{DR}_1 \left(-\widetilde{\sum a_i}, A_I, a_J \right) \boxtimes_1 \text{DR}_1(A_J, -a_J) \right) \\
 & + \text{Coef}_{a_1^2 a_2 a_3} \left(\frac{a_2 + a_3}{\sum a_i} \lambda_2 \text{DR}_2 \left(-\widetilde{\sum a_i}, a_1, a_2 + a_3 \right) \boxtimes_1 \text{DR}_0(a_2, a_3, -a_2 - a_3) \right).
 \end{aligned}$$

Let us look at a term in the sum in the first line of (57). The class $\lambda_1 \text{DR}_1(A_J, -a_J)$ is a polynomial in the variables a_j for $j \in J$, and it doesn't depend on a_1 . We have

$$\frac{1}{a_1 + a_2 + a_3} \lambda_1 \text{DR}_1(\widetilde{-a_1 - a_2 - a_3}, A_I, a_J) = (a_1 + a_2 + a_3) \lambda_1 \in R^1(\overline{\mathcal{M}}_{1,|I|+1}).$$

So, the polynomial class in the brackets in the first line of (57) depends on a_1 at most linearly. Therefore, the expression in the first line of (57) is equal to zero. Let us look at the expression in the second line of (57). We can easily see that it is equal to

$$\begin{aligned}
 (58) \quad & 2 \cdot \text{Coef}_{a^2 b^2} (\text{gl}_{1|2,3}^{2,0})_* \left(\frac{b}{a+b} \lambda_2 \text{DR}_2(\widetilde{-a-b}, a, b) \times [\overline{\mathcal{M}}_{0,3}] \right) \\
 & = 2 \cdot (\text{gl}_{1|2,3}^{2|0})_* (A_{2,1}^2 \times [\overline{\mathcal{M}}_{0,3}]).
 \end{aligned}$$

As a result, we obtain

$$A_{2,1,1}^2 = \psi_1 A_{1,1,1}^2 + 2 \cdot (\text{gl}_{1|2,3}^{2|0})_* (A_{2,1}^2 \times [\overline{\mathcal{M}}_{0,3}]).$$

On the other hand, we have

$$\begin{aligned}
 B_{2,1,1}^2 = & \psi_1^2 \psi_2 \psi_3 - 6 \text{ (diagram)} - 3 \text{ (diagram)} - \text{ (diagram)} - \text{ (diagram)} \\
 & + 6 \text{ (diagram)} + 3 \text{ (diagram)}.
 \end{aligned}$$

Using also formula (51), we compute

$$\begin{aligned}
 B_{2,1,1}^2 - \psi_1 B_{1,1,1}^2 = & -3 \text{ (diagram)} + 3 \text{ (diagram)} + 2 \text{ (diagram)} \\
 & - 2 \text{ (diagram)} - 2 \text{ (diagram)} \\
 = & 2 \text{ (diagram)} - 6 \text{ (diagram)} - 2 \text{ (diagram)} \\
 & - 2 \text{ (diagram)} + 6 \text{ (diagram)}.
 \end{aligned}$$

Using (49) we see that the last expression is equal to $2 \cdot (\text{gl}_{1|2,3}^{2|0})_* (B_{2,1}^2 \times [\overline{\mathcal{M}}_{0,3}])$ and we get

$$B_{2,1,1}^2 = \psi_1 B_{1,1,1}^2 + 2 \cdot (\text{gl}_{1|2,3}^{2|0})_* (B_{2,1}^2 \times [\overline{\mathcal{M}}_{0,3}]).$$

Since the relations $A_{2,1}^2 = B_{2,1}^2$ and $A_{1,1,1}^2 = B_{1,1,1}^2$ are proved, we conclude that relation $A_{2,1,1}^2 = B_{2,1,1}^2$ is true.

A.6 Relation $A_{1,1,1,1}^2 = B_{1,1,1,1}^2$

We follow the same strategy as in the previous section. Using (56), we compute $A_{1,1,1,1}^2 - \psi_1 A_{0,1,1,1}^2$ as

$$\begin{aligned}
 (59) \quad & \sum_{\substack{I \sqcup J = \{1,2,3,4\} \\ I \ni 1, |J| \geq 1}} \text{Coef}_{a_1 a_2 a_3 a_4} \left(\frac{a_J}{a} \lambda_2 \text{DR}_1(\widetilde{-a}, A_I, a_J) \boxtimes_1 \text{DR}_1(A_J, -a_J) \right) \\
 & + \sum_{\substack{I \sqcup J = \{1,2,3,4\} \\ I \ni 1, |J| \geq 2}} \text{Coef}_{a_1 a_2 a_3 a_4} \left(\frac{a_J}{a} \lambda_2 \text{DR}_2(\widetilde{-a}, A_I, a_J) \boxtimes_1 \text{DR}_0(A_J, -a_J) \right),
 \end{aligned}$$

where $a := \sum_{i=1}^4 a_i$. Let us look at a term in the sum in the first line of (59). We have $\frac{1}{a} \lambda_1 \text{DR}_1(\overline{a}, A_I, a_J) = a \lambda_1 \in R^1(\overline{\mathcal{M}}_{1,|I|+1})$ and the class $\lambda_1 \text{DR}_1(A_J, -a_J)$ doesn't depend on the variables a_i for $i \in I$. Therefore, the coefficient of $a_1 a_2 a_3 a_4$ can be nonzero only if $I = \{1\}$. So the expression in the first line of (59) is equal to

$$(60) \quad (\text{gl}_{1|2,3,4}^{1|1})_* (\lambda_1 \times \text{Coef}_{a_2 a_3 a_4} ((a_2 + a_3 + a_4) \lambda_1 \text{DR}_1(a_2, a_3, a_4, -a_2 - a_3 - a_4)))$$

$$= - \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} - 3 \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} + 2 \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} + 3 \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} + 3 \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} + 3 \begin{array}{c} \lambda_1 \\ \circ \\ | \\ 1 \end{array} - \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array} + 3 \begin{array}{c} \lambda_1 \\ \circ \\ / \backslash \end{array}.$$

The expression in the second line of (59) is equal to

$$(61) \quad 6(\text{gl}_{1|2,3,4}^{2|0})_*(A_{1,2}^2 \times [\overline{\mathcal{M}}_{0,4}]) + 2 \sum_{\substack{\{i,j,k\}=\{2,3,4\} \\ j < k}} (\text{gl}_{1,i|j,k}^{2|0})_*(A_{1,1,1}^2 \times [\overline{\mathcal{M}}_{0,3}]).$$

On the other hand, we have

$$B_{1,1,1,1}^2 = \psi_1 \psi_2 \psi_3 \psi_4 - 6 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} - 2 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} - 2 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} - 1 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array}$$

$$- 4 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 4 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 6 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 2 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array}$$

$$+ 2 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 2 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 8 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array}$$

$$- 4 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} + 4 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array} - 8 \begin{array}{c} \psi \\ \circ \\ / \backslash \end{array}.$$

After a long direct computation, which uses only the genus 0 topological recursion relation, we obtain that $B_{1,1,1,1}^2 - \psi_1 B_{0,1,1,1}^2$ equals

$$6(\text{gl}_{1|2,3,4}^{2|0})_*(B_{1,2}^2 \times [\overline{\mathcal{M}}_{0,4}]) + 2 \sum_{\substack{\{i,j,k\}=\{2,3,4\} \\ j < k}} (\text{gl}_{1,i|j,k}^{2|0})_*(B_{1,1,1}^2 \times [\overline{\mathcal{M}}_{0,3}]) + (\text{contd.})$$

$$\begin{aligned}
 &+ 6 \text{ (diagram)} + 2 \text{ (diagram)} - 2 \text{ (diagram)} \\
 &- \text{ (diagram)} + \left(\text{ (diagram)} - 2 \text{ (diagram)} - 2 \text{ (diagram)} \right. \\
 &\quad \left. - 6 \text{ (diagram)} + 2 \text{ (diagram)} \right).
 \end{aligned}$$

Using the formula

$$\psi_1 = \lambda_1 + \text{ (diagram)} \in R^1(\overline{\mathcal{M}}_{1,2}),$$

we can rewrite the expression in brackets above as

$$\begin{aligned}
 (62) \quad &\left(\lambda_1 \text{ (diagram)} - 2 \lambda_1 \text{ (diagram)} - 2 \lambda_1 \text{ (diagram)} - 6 \lambda_1 \text{ (diagram)} \right. \\
 &\quad \left. + 2 \lambda_1 \text{ (diagram)} \right) \\
 &+ \text{ (diagram)} - 2 \text{ (diagram)} - 2 \text{ (diagram)} - 6 \text{ (diagram)} \\
 &\quad + 2 \text{ (diagram)}.
 \end{aligned}$$

The expression on the right-hand side of (60) has the form $(g1_{1|2,3,4}^{1|1})_*(\lambda_1 \times \alpha)$, where

$$\alpha = -4 \text{ (diagram)} - 3 \text{ (diagram)} + 2 \text{ (diagram)} + 3 \text{ (diagram)} + 3 \text{ (diagram)}.$$

The part in brackets in expression (62) has the form $(\text{gl}_{1|2,3,4}^{1|1})_*(\lambda_1 \times \beta)$, where

$$\beta = 4 \begin{array}{c} \textcircled{1} \\ \psi \quad \psi \end{array} - 2 \begin{array}{c} 4 \\ \textcircled{1} \quad \textcircled{0} \\ \psi \quad \quad \quad \end{array} - 2 \begin{array}{c} 4 \\ \textcircled{1} \quad \textcircled{0} \\ \psi \quad \quad \quad \end{array} - 6 \begin{array}{c} 4 \\ \textcircled{1} \quad \textcircled{0} \\ \psi \quad \quad \quad \end{array} + 2 \begin{array}{c} 4 \\ \textcircled{1} \quad \textcircled{0} \quad \textcircled{0} \end{array} .$$

Expressing all psi-classes using the genus 1 topological recursion relation and also using the WDVV relation, it is easy to show that $\alpha = \beta$. Since $A_{2,1}^2 = B_{2,1}^2$ and $A_{1,1,1}^2 = B_{1,1,1}^2$, we obtain

$$(63) \quad B_{1,1,1,1}^2 - A_{1,1,1,1}^2 = 6 \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi^2 \quad \quad \quad \\ 1 \quad \quad \quad \end{array} + 2 \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - 2 \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ - \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} + \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \\ \psi \quad \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - 2 \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ - 2 \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - 6 \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ + 2 \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \textcircled{0} \\ \quad \quad \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} .$$

Define an operator $\text{Sym}: R^*(\overline{\mathcal{M}}_{2,4}) \rightarrow R^*(\overline{\mathcal{M}}_{2,4})$ by

$$\text{Sym}(\alpha) := \frac{1}{4!} \sum_{\sigma \in S_4} \sigma \alpha, \quad \alpha \in R^*(\overline{\mathcal{M}}_{2,4}),$$

where the symmetric group S_4 acts on $\overline{\mathcal{M}}_{2,4}$ by permutations of marked points. Applying the operator Sym to both sides of (63) we obtain that

$$B_{1,1,1,1}^2 - A_{1,1,1,1}^2 = \frac{3}{2} \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi^2 \quad \quad \quad \\ 1 \quad \quad \quad \end{array} + \frac{1}{2} \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - \frac{1}{2} \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ - \frac{1}{4} \begin{array}{c} \textcircled{1} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} + \frac{1}{4} \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \\ \psi \quad \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - \frac{1}{2} \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ - \frac{1}{2} \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \psi \quad \quad \quad \\ 1 \quad \quad \quad \end{array} - \frac{3}{2} \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \\ \psi \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} \\ + \frac{1}{2} \begin{array}{c} \textcircled{1} \textcircled{0} \textcircled{1} \textcircled{0} \textcircled{0} \\ \quad \quad \quad \quad \quad \quad \quad \\ 1 \quad \quad \quad \end{array} .$$

We see that the expression on the right-hand side has the form $(g_{|1,2,3,4}^{1|1})_*(\overline{\mathcal{M}}_{1,1} \times \rho)$, where

$$\begin{aligned} \rho = & \frac{3}{2} \text{diagram}_1 + \frac{1}{2} \text{diagram}_2 - \frac{1}{2} \text{diagram}_3 - \frac{1}{4} \text{diagram}_4 \\ & + \frac{1}{4} \text{diagram}_5 - \frac{1}{2} \text{diagram}_6 - \frac{1}{2} \text{diagram}_7 \\ & - \frac{3}{2} \text{diagram}_8 + \frac{1}{2} \text{diagram}_9. \end{aligned}$$

It is sufficient to prove that $\rho = 0$. For this we express all the psi-classes using the genus 1 topological recursion relation, and then prove that $\rho = 0$ using the WDVV relation and Getzler’s relation. This computation is straightforward, but quite long, so we present here only the most interesting parts of it. Expressing all the psi-classes, we obtain

$$\rho = \lambda_1 \theta_1 + \theta_2,$$

where $\theta_1 \in R^2(\overline{\mathcal{M}}_{1,5})$ and $\theta_2 \in R^3(\overline{\mathcal{M}}_{1,5})$ are sums of boundary strata. Using the WDVV relation it is not hard to prove that $\theta_2 = 0$. For the class $\lambda_1 \theta_1$ we get the expression

$$\begin{aligned} \lambda_1 \theta_1 = & a_1^{0,1,3} + \frac{1}{6} a_1^{0,2,2} - \frac{1}{6} a_1^{1,1,2} - \frac{1}{4} a_2^{0,0,4} + \frac{7}{8} a_2^{0,1,3} + \frac{1}{8} a_2^{0,2,2} + \frac{5}{4} a_2^{1,0,3} + \frac{1}{4} a_2^{1,1,2} \\ & + a_2^{2,0,2} - \frac{7}{16} a_3^{0,1,3} - \frac{3}{4} a_3^{0,2,2} + \frac{1}{16} a_3^{0,3,1} - \frac{7}{12} a_3^{1,1,2} + \frac{1}{12} a_3^{1,2,1} + b_1^{0,2,2} \\ & - \frac{3}{2} b_2^{0,1,3} - \frac{1}{2} b_2^{1,1,2}, \end{aligned}$$

where we use the notation

$$\begin{aligned} a_1^{i,j,k} & := \text{diagram}_1 & a_2^{i,j,k} & := \text{diagram}_2 \\ a_3^{i,j,k} & := \text{diagram}_3 & b_1^{0,2,2} & := \text{diagram}_4 \\ b_2^{i,j,k} & := \text{diagram}_5 \end{aligned}$$

Consider Getzler’s relation (52). Let $\pi: \overline{\mathcal{M}}_{1,5} \rightarrow \overline{\mathcal{M}}_{1,4}$ be the forgetful map that forgets the last marked point. We have

$$\begin{aligned} \lambda_1 \pi^* \gamma &= \frac{1}{2} a_1^{0,1,3} - \frac{1}{6} a_1^{0,2,2} - \frac{1}{3} a_1^{1,1,2} + \frac{1}{2} a_2^{0,1,3} - \frac{1}{6} a_2^{0,2,2} - \frac{1}{3} a_2^{1,1,2} + \frac{1}{2} a_3^{0,1,3} \\ &\quad - \frac{1}{6} a_3^{0,2,2} - \frac{1}{3} a_3^{1,1,2} + b_1^{0,2,2} + b_2^{0,2,2} \\ &= 0 \in R^3(\overline{\mathcal{M}}_{1,5}). \end{aligned}$$

Let $\pi': \overline{\mathcal{M}}_{1,5} \rightarrow \overline{\mathcal{M}}_{1,4}$ be the forgetful map that forgets the first marked point. We assume that, after forgetting the first marked point, a point labelled by i with $i \geq 2$ on a curve from $\overline{\mathcal{M}}_{1,5}$ becomes a point labelled by $i - 1$ on a curve in $\overline{\mathcal{M}}_{1,4}$. The symmetric group S_4 acts on $\overline{\mathcal{M}}_{1,5}$ by permutations of the first four marked points. Define a map $\text{Sym}' : R^*(\overline{\mathcal{M}}_{1,5}) \rightarrow R^*(\overline{\mathcal{M}}_{1,5})$ by

$$\text{Sym}'(\alpha) := \frac{1}{4!} \sum_{\sigma \in S_4} \sigma \alpha, \quad \alpha \in R^*(\overline{\mathcal{M}}_{1,5}).$$

We have

$$\begin{aligned} \lambda_1 \text{Sym}'((\pi')^* \gamma) &= -\frac{1}{4} a_1^{0,1,3} - \frac{1}{6} a_1^{0,2,2} - \frac{1}{12} a_1^{1,1,2} + \frac{1}{2} a_2^{0,0,4} - \frac{1}{12} a_2^{0,2,2} - \frac{1}{8} a_2^{1,0,3} \\ &\quad - \frac{1}{8} a_2^{1,1,2} - \frac{1}{6} a_2^{2,0,2} + \frac{3}{8} a_3^{0,1,3} + \frac{1}{6} a_3^{0,2,2} - \frac{1}{8} a_3^{0,3,1} - \frac{1}{24} a_3^{1,1,2} \\ &\quad - \frac{5}{24} a_3^{1,2,1} - \frac{1}{6} a_3^{2,1,1} + \frac{3}{4} b_2^{0,1,3} + \frac{1}{2} b_2^{0,2,2} + \frac{1}{4} b_2^{1,1,2} \\ &= 0 \in R^3(\overline{\mathcal{M}}_{1,5}). \end{aligned}$$

We compute

$$\begin{aligned} \lambda_1 \theta_1 &= \lambda_1 \theta_1 - \lambda_1 \pi^* \gamma + 2 \lambda_1 \text{Sym}'((\pi')^* \gamma) \\ &= \frac{3}{4} a_2^{0,0,4} + \frac{3}{8} a_2^{0,1,3} + \frac{1}{8} a_2^{0,2,2} + a_2^{1,0,3} + \frac{1}{3} a_2^{1,1,2} + \frac{2}{3} a_2^{2,0,2} - \frac{3}{16} a_3^{0,1,3} - \frac{1}{4} a_3^{0,2,2} \\ &\quad - \frac{3}{16} a_3^{0,3,1} - \frac{1}{3} a_3^{1,1,2} - \frac{1}{3} a_3^{1,2,1} - \frac{1}{3} a_3^{2,1,1}. \end{aligned}$$

Finally, applying the WDVV relations

$$\begin{aligned} a_3^{2,1,1} &= 2 a_2^{2,0,2}, \\ a_2^{1,0,3} &= -\frac{1}{3} a_2^{1,1,2} + \frac{1}{3} a_3^{1,1,2} + \frac{1}{3} a_3^{1,2,1}, \\ a_2^{0,0,4} &= -\frac{1}{2} a_2^{0,1,3} - \frac{1}{6} a_2^{0,2,2} + \frac{1}{4} a_3^{0,1,3} + \frac{1}{3} a_3^{0,2,2} + \frac{1}{4} a_3^{0,3,1}, \end{aligned}$$

it is easy to see that $\lambda_1 \theta_1 = 0$. The relation $A_{1,1,1,1}^2 = B_{1,1,1,1}^2$ is proved.

References

- [1] **A Buryak**, *Double ramification cycles and integrable hierarchies*, *Comm. Math. Phys.* 336 (2015) 1085–1107 MR
- [2] **A Y Buryak**, *New approaches to hierarchies of topological type*, *Uspekhi Mat. Nauk* 72 (2017) 63–112 MR In Russian; translated in *Russian Math. Surveys* 72 (2017) 841–887
- [3] **A Buryak, B Dubrovin, J Guéré, P Rossi**, *Tau-structure for the double ramification hierarchies*, *Comm. Math. Phys.* 363 (2018) 191–260 MR
- [4] **A Buryak, B Dubrovin, J Guéré, P Rossi**, *Integrable systems of double ramification type*, *Int. Math. Res. Notices* (online publication February 2019) art. id. rnz029
- [5] **A Buryak, J Guéré**, *Towards a description of the double ramification hierarchy for Witten’s r -spin class*, *J. Math. Pures Appl.* 106 (2016) 837–865 MR
- [6] **A Buryak, H Posthuma, S Shadrin**, *A polynomial bracket for the Dubrovin–Zhang hierarchies*, *J. Differential Geom.* 92 (2012) 153–185 MR
- [7] **A Buryak, P Rossi**, *Double ramification cycles and quantum integrable systems*, *Lett. Math. Phys.* 106 (2016) 289–317 MR
- [8] **A Buryak, P Rossi**, *Recursion relations for double ramification hierarchies*, *Comm. Math. Phys.* 342 (2016) 533–568 MR
- [9] **A Buryak, S Shadrin, L Spitz, D Zvonkine**, *Integrals of ψ -classes over double ramification cycles*, *Amer. J. Math.* 137 (2015) 699–737 MR
- [10] **B Dubrovin, Y Zhang**, *Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants*, preprint (2001) arXiv
- [11] **C Faber**, *Chow rings of moduli spaces of curves, I: The Chow ring of $\overline{\mathcal{M}}_3$* , *Ann. of Math.* 132 (1990) 331–419 MR
- [12] **C Faber, R Pandharipande**, *Hodge integrals and Gromov–Witten theory*, *Invent. Math.* 139 (2000) 173–199 MR
- [13] **C Faber, R Pandharipande**, *Relative maps and tautological classes*, *J. Eur. Math. Soc.* 7 (2005) 13–49 MR
- [14] **E Getzler**, *Intersection theory on $\overline{\mathcal{M}}_{1,4}$ and elliptic Gromov–Witten invariants*, *J. Amer. Math. Soc.* 10 (1997) 973–998 MR
- [15] **E Getzler**, *Topological recursion relations in genus 2*, from “Integrable systems and algebraic geometry” (M-H Saito, Y Shimizu, K Ueno, editors), World Sci., River Edge, NJ (1998) 73–106 MR
- [16] **R Hain**, *Normal functions and the geometry of moduli spaces of curves*, from “Handbook of moduli, I” (G Farkas, I Morrison, editors), Adv. Lect. Math. 24, International, Somerville, MA (2013) 527–578 MR

- [17] **E-N Ionel**, *Topological recursive relations in $H^{2g}(\mathcal{M}_{g,n})$* , *Invent. Math.* 148 (2002) 627–658 MR
- [18] **F Janda**, *Relations on $\overline{\mathcal{M}}_{g,n}$ via equivariant Gromov–Witten theory of \mathbb{P}^1* , preprint (2015) arXiv
- [19] **F Janda, R Pandharipande, A Pixton, D Zvonkine**, *Double ramification cycles on the moduli spaces of curves*, *Publ. Math. Inst. Hautes Études Sci.* 125 (2017) 221–266 MR
- [20] **M Kontsevich, Y Manin**, *Gromov–Witten classes, quantum cohomology, and enumerative geometry*, *Comm. Math. Phys.* 164 (1994) 525–562 MR
- [21] **S Marcus, J Wise**, *Stable maps to rational curves and the relative Jacobian*, preprint (2013) arXiv
- [22] **R Pandharipande, A Pixton, D Zvonkine**, *Relations on $\overline{\mathcal{M}}_{g,n}$ via 3–spin structures*, *J. Amer. Math. Soc.* 28 (2015) 279–309 MR
- [23] **P Rossi**, *Integrability, quantization and moduli spaces of curves*, *Symmetry Integrability Geom. Methods Appl.* 13 (2017) art. id. 60 MR

*School of Mathematics, University of Leeds
Leeds, United Kingdom*

*Université Grenoble Alpes, CNRS, Institut Fourier
Grenoble, France*

*Dipartimento di Matematica “Tullio Levi-Civita”, Università degli Studi di Padova
Padova, Italy*

a.buryak@leeds.ac.uk, jeremy.guere@univ-grenoble-alpes.fr,
paolo.rossi@math.unipd.it

<https://sites.google.com/site/alexandrburyakhomepage/home>,

<https://www-fourier.ujf-grenoble.fr/~guerej/index.html>,

<http://www.math.unipd.it/~rossip/>

Proposed: Yasha Eliashberg

Received: 5 May 2018

Seconded: Richard Thomas, Jim Bryan

Accepted: 4 November 2018