

Sutured manifolds and polynomial invariants from higher rank bundles

ALIAKBAR DAEMI

YI XIE

For each integer $N \geq 2$, Mariño and Moore defined generalized Donaldson invariants by the methods of quantum field theory, and made predictions about the values of these invariants. Subsequently, Kronheimer gave a rigorous definition of generalized Donaldson invariants using the moduli spaces of anti-self-dual connections on hermitian vector bundles of rank N . We confirm the predictions of Mariño and Moore for simply connected elliptic surfaces without multiple fibers and certain surfaces of general type in the case that $N = 3$. The primary motivation is to study 3–manifold instanton Floer homologies which are defined by higher rank bundles. In particular, the computation of the generalized Donaldson invariants is exploited to define a Floer homology theory for sutured 3–manifolds.

57M27, 57R57, 57R58

1. Introduction	49
2. Higher rank bundles and polynomial invariants	57
3. Floer homologies for closed 3–manifolds	74
4. Computing polynomial invariants	97
5. Sutured Floer homology	130
6. Gluing theory	142
7. Questions and conjectures	168
Appendix. Invariants of flat connections on $\Sigma(2, 3, 23)$	171
References	173

1 Introduction

Sutured manifolds were introduced by Gabai [34] to study foliations and the Thurston norm of 3–manifolds [82]. A sutured manifold is a pair of a 3–manifold M and an oriented 1–manifold $\alpha \subset M$ which decomposes the boundary of M in an appropriate

way. In [34], Gabai also defines an operation on sutured manifolds, which is called *surface decomposition*. Surface decompositions can be used to simplify sutured manifolds. Foliations of sutured manifolds are also well behaved with respect to surface decompositions. As a result, Gabai was able to construct *taut* foliations for certain families of 3-manifolds in an inductive way.

Floer homological invariants serve as another set of tools for studying topology and geometry of 3-dimensional manifolds. Such invariants were initially constructed for closed and oriented 3-manifolds: $U(N)$ -instanton Floer homology — see Floer [27; 28] and Kronheimer and Mrowka [58]; Heegaard Floer homology — see Ozsváth and Szabó [74]; monopole Floer homology — see Kronheimer and Mrowka [55]; embedded contact homology — see Hutchings [44; 45]. Later, Juhász [46] defined *sutured Floer homology*, a generalization of Heegaard Floer homology to *balanced* sutured 3-manifolds.¹ Subsequently, Kronheimer and Mrowka [56] constructed a sutured version of $U(2)$ -instanton Floer homology and monopole Floer homology, and embedded contact homology was constructed by Colin, Ghiggini, Honda and Hutchings [10; 11] and Kutluhan, Sivek and Taubes [60]. In particular, Kronheimer and Mrowka [57] used sutured $U(2)$ -instanton homology as the main ingredient to establish that Khovanov homology detects the unknot. This invariant was also used to reprove Property P for knots [56], and it lies at the core of a program that hopes to find a computer-free proof of the famous four-color theorem; see Kronheimer and Mrowka [59]. The primary motivation for this article is to extend $U(N)$ -instanton Floer homology to sutured manifolds for higher values of N .

1.1 Motivation

Fix an integer $N \geq 2$, and let K be a knot in an integral homology sphere Y . Also let μ denote an element of the knot group, $\pi_1(Y \setminus K)$, represented by a meridian of K :

Question 1.1 *Does there exist a representation $\varphi: \pi_1(Y \setminus K) \rightarrow \mathrm{SU}(N)$ with non-abelian image such that*

$$(1.2) \quad \varphi(\mu) = c \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & \zeta & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \zeta^{N-1} \end{bmatrix}$$

where $\zeta = e^{2\pi i/N}$, and $c = e^{\pi i/N}$ or $c = 1$ depending on whether N is even or odd?

¹For the definition of balanced sutured 3-manifolds, see Definition 5.20.

In the case that K is the unknot, the answer to the above question is clearly negative. Note also that if for a knot K , there is a representation to $SU(N)$ with the mentioned properties, then there is also a desired representation from $\pi_1(Y \setminus K)$ to $SU(lN)$ for any positive integer l .

Suppose Y is a homotopy sphere² and the answer to [Question 1.1](#) for any nontrivial knot K in Y is positive. A nonabelian representation φ satisfying (1.2) determines a nontrivial representation of $\pi_1(\Sigma_N(K))$ with $\Sigma_N(K)$ being the N -fold cyclic branched cover of Y , branched along K . This verifies the *covering conjecture*, which asserts that $\Sigma_N(K)$, for a nontrivial knot K , is not a homotopy sphere; see Kirby [48, Problem 3.38]. A consequence of the covering conjecture is the *Smith conjecture*, stating that a nontrivial knot is not the fixed-point set of an orientation-preserving homeomorphism $f: S^3 \rightarrow S^3$ of order N ; see Kirby [48, Problem 3.38]. The covering conjecture and the Smith conjecture have both been proved by *geometrization* techniques; see [67].

The sutured $U(2)$ -instanton homology group SHI_*^2 of Kronheimer and Mrowka [56] can be employed to answer [Question 1.1](#) affirmatively for $N = 2$ (and hence for any even N) and any nontrivial knot K .³ Associated to any knot K , there is a sutured manifold $(M(K), \alpha(K))$, where $M(K)$ is the knot complement and $\alpha(K)$ is the union of two oppositely oriented meridional curves. Kronheimer and Mrowka proved that if the dimension of $\text{SHI}_*^2(M(K), \alpha(K))$ is greater than 1, then there is a nonabelian representation of the knot group of K that satisfies (1.2). Similar to foliations, SHI_*^2 also behaves well with respect to surface decomposition, and one can inductively construct nontrivial elements of $\text{SHI}_*^2(M(K), \alpha(K))$ after simplifying $(M(K), \alpha(K))$ by a series of sutured decompositions. In particular, the dimension of $\text{SHI}_*^2(M(K), \alpha(K))$ is at least two for a nontrivial knot K . It is also shown by Boden and Friedl [5; 29] that if K is a knot with nontrivial Alexander polynomial, then the answer to [Question 1.1](#) is positive for infinitely many values of N . In light of the success of SHI_*^2 in addressing [Question 1.1](#), it is natural to look for the generalization of SHI_*^2 for higher values of N .

The essential device in the definition of the sutured Floer homology group SHI_*^2 is an excision theorem for $U(2)$ -instanton Floer homology; see Braam and Donaldson [8],

²By the Poincaré conjecture, this is equivalent to saying that $Y = S^3$. However, we are making this assumption to show that our proposed approach does not require the Poincaré conjecture.

³The original notation for sutured $U(2)$ -instanton homology is SHI_* . Here we use the superscript 2 to indicate that this invariant is the sutured version of $U(2)$ -instanton homology.

Floer [28] and Kronheimer and Mrowka [56]. The proof of the excision theorem is in turn based on Muñoz's characterization of the structure of a $U(2)$ -instanton Floer homology group associated to the 3-manifold $S^1 \times \Sigma$, where Σ is a Riemann surface [73]. Muñoz's work borrows some results about the cohomology ring of the moduli space of rank 2 stable bundles — see Baranovskii [3], King and Newstead [47], Siebert and Tian [78] and Zagier [86] — which are not available for higher values of the rank.

In the present paper, we establish an excision theorem for $N = 3$ using the relationship between instanton Floer homology and generalizations of Donaldson invariants from Kronheimer [52]. Roughly speaking, there is a $(3+1)$ -dimensional topological quantum field theory which associates $U(N)$ -instanton Floer homology to 3-manifolds and $U(N)$ analogues of Donaldson polynomial invariants to closed 4-manifolds. This relationship between $U(2)$ -instanton Floer homology and polynomial invariants has been extensively used to compute the invariants of 4-manifolds. In this paper, we first use the TQFT structure to compute the $U(3)$ -polynomial invariants of some families of smooth 4-manifolds. Next, we work in the other direction, and use our knowledge of $U(3)$ -polynomial invariants to obtain a better understanding of certain $U(3)$ -Floer homologies. This allows us to prove the excision theorem and define a Floer homology group SHI_*^3 for sutured manifolds in the case that $N = 3$. Computations of generalized polynomial invariants in the physics literature — see Mariño and Moore [64] — suggest that our approach can be also exploited for higher values of N .

1.2 Statement of results

In his groundbreaking work [17], Donaldson defined polynomial invariants for a smooth manifold X using the moduli space of *anti-self-dual* (ASD) connections on X . In his work, X is simply connected, $b^+(X)$ is an integer greater than 1, and the ASD connections are assumed to be defined on an $SU(2)$ -bundle E over X . Although the assumption on $b^+(X)$ is essential, the definition of polynomial invariants was subsequently generalized to the case that X is not simply connected — see Kronheimer and Mrowka [54] — and E is a $U(N)$ -bundle — see Culler [12] and Kronheimer [52]. Polynomial invariants have been extensively studied in the case that $N = 2$. However, there is not much known about these invariants for higher values of N .

For a smooth and connected 4-manifold X , suppose the algebra $\mathbb{A}(X)$ is defined as

$$\mathbb{A}(X) := \text{Sym}^*(H_0(X) \oplus H_2(X)) \otimes \Lambda^*(H_1(X)),$$

where $H_i(X)$ is computed with coefficients in \mathbb{C} . Form the tensor product algebra $\mathbb{A}(X)^{\otimes(N-1)}$, and for $\alpha \in H_i(X)$ and $2 \leq r \leq N$, let $\alpha_{(r)}$ be the corresponding element in the $(r-1)^{\text{st}}$ factor of $\mathbb{A}(X)^{\otimes(N-1)}$. In the case that α is the generator of $H_0(X)$, this element of $\mathbb{A}(X)^{\otimes(N-1)}$ is denoted by a_r . We also define a grading on $\mathbb{A}(X)^{\otimes(N-1)}$ such that for $\alpha \in H_i(X)$, the degree of $\alpha_{(r)}$ is equal to $2r - i$. A Hermitian vector bundle E of rank N on X is determined by its first and second Chern classes. Suppose $c_1(E)$ is represented by an embedded surface w in X and $c_2(E)[X] = k$. Then the $U(N)$ -polynomial invariants associated to the bundle E are given by a linear map,⁴

$$D_{X,w,k}^N : \mathbb{A}(X)^{\otimes(N-1)} \rightarrow \mathbb{C}.$$

For $z \in \mathbb{A}(X)^{\otimes(N-1)}$, the complex number $D_{X,w,k}^N(z)$ is nonzero only if

$$(1.3) \quad \deg(z) = 4Nk - 2(N-1)w \cdot w - (N^2 - 1)\frac{1}{2}(\chi(X) + \sigma(X)).$$

Therefore, we will not lose any information if we combine these invariants as

$$(1.4) \quad D_{X,w}^N := \sum_k D_{X,w,k}^N.$$

A substantial part of the present paper is devoted to computing $U(3)$ -polynomial invariants of some families of algebraic surfaces. Our first result in this direction is the following:

Theorem 1 *Suppose X is a K3 surface. Then for any embedded oriented surface w in X and any element $z \in \mathbb{A}(X)^{\otimes 2}$,*

$$(1.5) \quad D_{X,w}^3(a_2^3 z) = 27 D_{X,w}^3(z) \quad \text{and} \quad D_{X,w}^3(a_3 z) = 0.$$

Moreover, if Γ and Λ are two elements of $H_2(X)$, then

$$(1.6) \quad D_{X,w}^3\left(\left(1 + \frac{1}{3}a_2 + \frac{1}{9}a_2^2\right) \cdot e^{\Gamma(2)+\Lambda(3)}\right) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}.$$

In order to clarify the statement of the above theorem, the following remarks are in order. The left-hand side of (1.6) is defined as

$$D_{X,w}^3\left(\left(1 + \frac{1}{3}a_2 + \frac{1}{9}a_2^2\right) \cdot e^{\Gamma(2)+\Lambda(3)}\right) := \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{D_{X,w}^3\left(\left(1 + \frac{1}{3}a_2 + \frac{1}{9}a_2^2\right) \Gamma_{(2)}^i \Lambda_{(3)}^j\right)}{i!j!}.$$

⁴Our definition of $U(N)$ -polynomial invariants slightly differs from Culler's. See Section 2.1 for more details.

Theorem 1 asserts that the above series for a K3 surface is convergent, and the resulting number is equal to $e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}$. Here Q denotes the intersection form of X . That is to say, $Q(\Gamma)$ is the algebraic intersection number of Γ with itself. In general, the intersection number of two homology classes Γ and Γ' is denoted by $\Gamma \cdot \Gamma'$. Since (1.6) holds for all choices of Γ and Λ , the dimension formula (1.3) allows us to compute the following polynomial invariants for all choices of nonnegative integers i and j , an integer $k \in \{0, 1, 2\}$, and homology classes Γ and Λ :

$$D_{X,w}^3(a_2^k \Gamma_{(2)}^i \Lambda_{(3)}^j).$$

These invariants determine $D_{X,w}^3(z)$ for all $z \in \mathbb{A}(X)^{\otimes 2}$, because the K3 surface satisfies (1.5) and $b_1(X) = 0$.

Our computation of the invariants of K3 surfaces motivates the following definition: a smooth 4-manifold X with $b^+(X) \geq 2$ and $b^1(X) = 0$ has w -simple type with respect to an embedded surface w if

$$(1.7) \quad D_{X,w}^3(a_2^3 z) = 27 D_{X,w}^3(z) \quad \text{and} \quad D_{X,w}^3(a_3 z) = 0$$

for all $z \in \mathbb{A}(X)^{\otimes 2}$. The 4-manifold X has *simple type* if it has w -simple type with respect to any w in X . As in the case of the K3 surfaces, if X has simple type and the series

$$\widehat{D}_{X,w}(e^{\Gamma_{(2)}+\Lambda_{(3)}}) := D_{X,w}^3\left(\left(1 + \frac{1}{3}a_2 + \frac{1}{9}a_2^2\right) \cdot e^{\Gamma_{(2)}+\Lambda_{(3)}}\right)$$

is convergent for all choices of w and $\Gamma, \Lambda \in H_2(X)$, then these series determine all polynomial invariants of X .

We can extend our calculation for the K3 surfaces to a larger family of complex surfaces. Suppose that $W(m, n)$ is the blowup of $\mathbb{C}P^1 \times \mathbb{C}P^1$ at the $4mn$ singular points of the (complex) curve

$$B := \{p_1, \dots, p_{2m}\} \times \mathbb{C}P^1 \cup \mathbb{C}P^1 \times \{q_1, \dots, q_{2n}\}.$$

Let \widetilde{B} be the proper transform of B , and define $X(m, n)$ to be the branched double cover of $W(m, n)$, branched along the smooth curve \widetilde{B} . The horizontal and vertical fibrations of $W(m, n)$ by projective lines lift to two fibrations of $X(m, n)$ whose generic fibers are denoted by f_{m-1} and f_{n-1} . The Riemann surface f_i , for $i \in \{m-1, n-1\}$, has genus i . The complex surface $X(2, 2)$ is a K3 surface. More generally, $X(m, 2)$ is an elliptic surface without multiple fibers, which is usually denoted by $E(m)$; see Gompf and Stipsicz [40].

Theorem 2 *The elliptic surface $E(n)$ has simple type. Moreover, there are rational numbers \hbar_1 and \hbar_2 independent of n such that for any embedded surfaces w in $E(n)$ and $\Gamma, \Lambda \in H_2(E(n))$, the series $\widehat{D}_{E(n),w}(e^{\Gamma(2)+\Lambda(3)})$ is equal to*

$$e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}\left[\hbar_1 \cosh(\sqrt{3}f \cdot \Gamma) - 2\hbar_2 \cos\left(-\frac{2}{3}\pi w \cdot f + \sqrt{3}f \cdot \Lambda\right)\right]^{n-2},$$

where $f = f_1$ represents an elliptic fiber of $E(n)$. Furthermore, $\hbar_1 + \hbar_2 = \pm 1$ for an appropriate choice of the sign.

The constants \hbar_1 and \hbar_2 in [Theorem 2](#) are respectively equal to $\frac{2}{3}$ and $\frac{1}{3}$ (work in progress). The set of surfaces $X(m, n)$, as a set of smooth 4-manifolds, is closed with respect to taking *fiber sums*.⁵ For example, we can take the fiber sum of $X(m, n_1)$ and $X(m, n_2)$ along the fiber f_{m-1} , and the resulting 4-manifold is diffeomorphic to $X(m, n_1 + n_2)$. Given two embedded surfaces $\Sigma_1 \subset X(m, n_1)$ and $\Sigma_2 \subset X(m, n_2)$ which intersect a fiber in the same number of points, we can form a surface $\Sigma_1 \# \Sigma_2 \subset X(m, n_1 + n_2)$. Suppose $\mathcal{H}(m, n_1, n_2) \subset H_2(X(m, n_1 + n_2))$ is the space of homology classes generated by homology classes of the surfaces of the form $\Sigma \# \Sigma'$. The following theorem about $X(m, 4)$ is a consequence of [Theorem 4.93](#) about the polynomial invariants of fiber sums. In fact, [Theorem 4.93](#) can be used to obtain similar results about other surfaces in the family $X(m, n)$.

Theorem 3 *For $m \geq 3$, let $w \subset X(m, 4)$ be an embedded surface which has the form $w_1 \# w_2$ for $w_i \subset X(m, 2)$ and $w \cdot f_{m-1} \not\equiv 0 \pmod{3}$. Let K denote the canonical class of $X(m, 4)$. Then there are rational numbers \hbar_3 and \hbar_4 , independent of m , such that for $\Gamma, \Lambda \in \mathcal{H}(m, 2, 2)$ the series $\widehat{D}_{X(m,4),w}(e^{\Gamma(2)+\Lambda(3)})$ is convergent and is equal to*

$$e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}\left[\frac{1}{2}\hbar_1^2\hbar_3^{m-2} \cosh(\sqrt{3}K \cdot \Gamma) + 2\hbar_2^2\hbar_4^{m-2} \cos\left(-\frac{2}{3}\pi w \cdot K + \sqrt{3}K \cdot \Lambda\right)\right],$$

where \hbar_1 and \hbar_2 are the constants of [Theorem 2](#).

We do not attempt to find the undetermined constants \hbar_3 and \hbar_4 here. We also believe that $X(m, 4)$ has simple type, and the above theorem holds for any choice of $w \subset X(m, 4)$ and homology classes Γ and Λ . But the current version of the theorem is sufficient for our 3-dimensional applications.

The algebraic surfaces in [Theorems 1, 2 and 3](#) are representatives of surfaces with different possible finite Kodaira dimensions. K3 surfaces, elliptic surface $E(n)$ with $n \geq 3$ and $X(m, 4)$ for $m \geq 3$ have Kodaira dimensions 0, 1 and 2, respectively.

⁵See [Section 3.3](#) for a review of the definition of fiber sums.

Theorem 1 shows that the $U(3)$ -polynomial invariants of a K3 surface associated to homology classes Γ and Λ are determined by the self-intersection of these homology classes. On the other hand, for the $U(3)$ -polynomial invariants of $E(n)$ and $X(m, 4)$ we also need the pairing of Γ and Λ with the canonical class. Recall that the first Chern class of the canonical classes of $E(n)$ and $X(m, 4)$ are represented by $(n - 2)f$ and $(m - 2)f_3 + 2f_{m-1}$, respectively.

In **Section 3**, we introduce various Floer homology groups associated to the 3-manifolds $S^1 \times \Sigma$, and explain how these vector spaces admit ring structure. We also characterize the vector space structure on these Floer homology groups. **Theorems 2 and 3** allow us to obtain further information about the ring structure of these rings. In particular, in **Proposition 5.7**, which plays an essential role for us, we construct certain elements of Floer homology groups of $S^1 \times \Sigma$ that interact nicely with the ring structure. We use these elements to obtain an excision theorem for $U(3)$ -instanton Floer homology. With the aid of this excision theorem, we construct the promised sutured Floer homology SHI_*^3 , following the approach of Kronheimer and Mrowka [56]. This sutured Floer homology group has the following property:

Theorem 4 *For a knot K in a homology sphere Y , suppose that the dimension of $\text{SHI}_*^3(M(K), \alpha(K))$ is greater than 1. Then there is a nonabelian representation of $\pi_1(Y \setminus K)$ into $SU(3)$ that satisfies the holonomy condition (1.2) for $N = 3$.*

This is established by **Corollary 5.34**. We conjecture that $\dim(\text{SHI}_*^3(M(K), \alpha(K))) > 1$ for any nontrivial knot K in a homology sphere Y such that $Y \setminus K$ is irreducible. This answers **Question 1.1** affirmatively for $N = 3$ and any nontrivial knot K in an integral homology sphere Y (without the irreducibility assumption on $Y \setminus K$). We hope to come back to this conjecture elsewhere.

1.3 Outline of contents

Section 2 gives a review of the moduli spaces of anti-self-dual connections on 4-manifolds (possibly with boundary) and $U(N)$ -polynomial invariants. This section also contains a nonvanishing theorem for $U(N)$ -polynomial invariants of algebraic surfaces. The second half of **Section 2** discusses how the $U(3)$ -polynomial invariants behave in the presence of negative embedded spheres. In particular, we recall the results of Culler's thesis [12] about the blowup formula for $U(3)$ -polynomial invariants and discuss how this formula can be simplified for smooth 4-manifolds with simple

type. Section 3 deals with various Floer homology groups which appear in this paper. After giving an exposition of $U(N)$ -instanton Floer homology, we study various Floer homologies of $\Sigma \times S^1$ where Σ is an oriented surface. We also discuss a generalization of $U(N)$ -instanton Floer homology, which is known as Fukaya–Floer homology in the case that $N = 2$.

The Floer homology groups of Section 3 are our main tools in computing $U(3)$ -polynomial invariants of several complex surfaces in Section 4. In particular, the proofs of Theorems 1, 2 and 3 are given in this section. In Section 4, we also study the behavior of $U(3)$ -polynomial invariants with respect to fiber sum. In Section 5, we prove our excision theorem and define the sutured Floer homology group SHI_*^3 . To make the exposition of the paper more comprehensible, we postpone providing proofs for technical results in Sections 2 and 3 until Section 6. These results are proved by gluing theory of the moduli spaces of anti-self-dual connections. Section 7 concerns various questions and conjectures which naturally arise from our work on this paper.

All manifolds in this paper are smooth and oriented. Given such a manifold X , we will write $H_i(X)$ and $H^i(X)$ for the homology and cohomology groups of X with complex coefficients. If we need to work with another coefficient ring R , then we use the notation $H_i(X, R)$ and $H^i(X, R)$. Our main results for this paper concern $U(3)$ -polynomial invariants and $U(3)$ -instanton Floer homologies. However, we believe that our method for the construction of SHI_*^3 should work for arbitrary N . Therefore, we try to state our results for general N , when it is possible.

Acknowledgements We thank Lucas Culler, Simon Donaldson, Peter Kronheimer, Claude LeBrun and Tomasz Mrowka for helpful conversations. We are thankful to Victor Mikhaylov for verifying that our definition of the simple type condition matches the predictions from the physics literature. We also appreciate the referee's close reading of the paper and helpful comments about the exposition of the paper. We are very grateful to the Simons Center for Geometry and Physics for providing a stimulating environment while we were working on this project.

2 Higher rank bundles and polynomial invariants

2.1 $U(N)$ -polynomial invariants

In this section, we review the definition of $U(N)$ -polynomial invariants of 4-manifolds based on [52; 12]. For $N = 2$, there is an extensive literature on the subject (see, for

example, [17; 20; 68; 54]). For higher values of N , these invariants were first defined in [64] by the methods of quantum field theory. A rigorous definition of polynomial invariants for higher rank bundles is given in [52]. As we mentioned earlier, the polynomial invariants of a 4-manifold X are homomorphisms defined on the algebra $\mathbb{A}(X)^{\otimes(N-1)}$. In [52], the polynomial invariants are defined only on the subalgebra

$$\mathbb{A}(X) \otimes 1 \otimes \cdots \otimes 1.$$

Kronheimer's definition was subsequently generalized to the algebra $\mathbb{A}(X)^{\otimes(N-1)}$ in [12]. The construction of Fukaya–Floer homology in Section 6.3 is based on Culler's modification of $U(N)$ –polynomial invariants in [12]. Therefore, we attempt to give enough background on his treatment to motivate the construction of $U(N)$ –Fukaya–Floer homology.

Suppose X is a smooth, closed, oriented and connected 4-manifold, w is an oriented embedded surface in X , and k is an integer. Then there is a $U(N)$ –bundle P , unique up to isomorphism, over X such that $c_1(P) = \text{P.D.}[w]$ and $c_2(P)[X] = k$. An explicit construction of this $U(N)$ –bundle can be given as follows. Suppose $D(w)$ is a regular neighborhood of w in X whose boundary is denoted by $S(w)$. Then we can consider a Hermitian line bundle on $D(w)$ which is trivialized on $S(w)$ and whose relative first Chern class is given by the Thom class of the disc bundle $D(w)$. By extending the trivialization to the complement of $D(w)$, we obtain a Hermitian line bundle L_w where $c_1(L_w) = \text{P.D.}[w]$. The direct sum of L_w and the trivial bundle $\underline{\mathbb{C}}^{N-1}$ defines a $U(N)$ –bundle P_0 on X with $c_2(P_0)[X] = 0$ and $c_1(P_0) = c_1(L_w)$. Next, fix a $U(N)$ –bundle on the 4-dimensional ball D^4 which is trivialized on the boundary and whose relative second Chern class is given by $k \text{ P.D.}[pt]$. Removing a ball from $X \setminus D(w)$ and gluing the above ball gives rise to the same 4-manifold. We can also use the trivializations to glue the $U(N)$ –bundle on D^4 to P_0 and produce a $U(N)$ –bundle P such that $c_1(P) = \text{P.D.}[w]$, $c_2(P)[X] = k$ and the determinant bundle of P is identified with L_w .

A 2-cycle w in a closed 4-manifold is a union of embedded closed surfaces in X . We can apply the construction of the previous paragraph to obtain a Hermitian line bundle L_{w_i} for each connected component w_i of w . Then we can replace L_w in the previous paragraph with the tensor product of the line bundles L_{w_i} and produce a $U(N)$ –bundle P with $c_1(P) = \text{P.D.}[w]$ and $c_2(P)[X] = k$. The *topological energy* of P is defined to be

$$\kappa := k - \frac{N-1}{2N} w \cdot w.$$

Thus the bundle P is determined by the pair (κ, w) up to a canonical isomorphism. We say a closed 2-cycle w in X is coprime to N if there is an embedded oriented surface $\Sigma \subset X$ such that the intersection number $w \cdot \Sigma$ is coprime to N .

Suppose P is a $U(N)$ -bundle on a closed 4-manifold determined by a pair (κ, w) . Fix an integer $l \geq 3$ and an arbitrary smooth connection B_0 on L_w . Let $\mathcal{A}_\kappa(X, w)$ be the space of L^2_l connections on P whose induced connections on $\det(P) = L_w$ are each equal to B_0 . If $\mathfrak{su}(P)$ is the bundle associated to the conjugation action of $U(N)$ on the Lie algebra $\mathfrak{su}(N)$ of $SU(N)$, then $\mathcal{A}_\kappa(X, w)$ is an affine space modeled on the Banach space $L^2_l(X, \mathfrak{su}(P) \otimes \Lambda^1)$. We will also write $\mathcal{G}_\kappa(X, w)$ for the space of L^2_{l+1} automorphisms of P whose fiberwise determinant is equal to 1. Then $\mathcal{G}_\kappa(X, w)$ forms a Banach Lie group with Lie algebra $L^2_{l+1}(X, \mathfrak{su}(P))$. This Lie group acts on $\mathcal{A}_\kappa(X, w)$, and the quotient space is denoted by $\mathcal{B}_\kappa(X, w)$. We will write $[A]$ for an element of $\mathcal{B}_\kappa(X, w)$ represented by a connection A . The center of the Lie group $U(N)$ induces a finite subgroup of $\mathcal{G}_\kappa(X, w)$. If this subgroup is the stabilizer of a connection A , then A is an *irreducible* connection. Otherwise, the connection A is called *reducible*. The space of irreducible connections on P is denoted by $\mathcal{A}^*_\kappa(X, w)$, and we will write $\mathcal{B}^*_\kappa(X, w)$ for the quotient space.

Fix a Riemannian metric on X and let $*$ denote the associated Hodge operator on differential forms of X . Then $*$ defines an involution on the space of 2-forms on X , and 2-forms in the 1-eigenspace (respectively, (-1) -eigenspace) are called *self-dual* (respectively, *anti-self-dual*). A connection $A \in \mathcal{A}_\kappa(X, w)$ is *anti-self-dual* if it satisfies

$$(2.1) \quad F_0^+(A) = 0,$$

where $F_0(A)$ denotes the projection of the curvature of A to the space $\mathfrak{su}(P)$, and $F_0^+(A)$ is the self-dual part of $F_0(A)$. In other words, the connection induced by A on the associated $PU(N)$ -bundle to P has anti-self-dual curvature. Equation (2.1) is invariant with respect to the action of $\mathcal{G}_\kappa(X, w)$ and the quotient space of ASD connections is denoted by $\mathcal{M}_\kappa(X, w)$.

The local behavior of the moduli space $\mathcal{M}_\kappa(X, w)$ around an element $[A]$ is governed by the following elliptic complex, denoted by \mathcal{D}_A :

$$(2.2) \quad L^2_{l+1}(X, \mathfrak{su}(P)) \xrightarrow{d_A} L^2_l(X, \mathfrak{su}(P) \otimes \Lambda^1) \xrightarrow{d_A^+} L^2_{l-1}(X, \mathfrak{su}(P) \otimes \Lambda^+),$$

where Λ^+ denotes the bundle of self-dual forms on X . The i^{th} cohomology group of this complex is denoted by H^i_A . The connection A is irreducible if and only if H^0_A is

trivial. We say A is *regular* if $H^2(A)$ is trivial. If A is an irreducible and regular ASD connection, then in a neighborhood of $[A]$, the moduli space is a smooth manifold of the same dimension as H_A^1 . In this case, the dimension of H_A^1 is given explicitly by

$$(2.3) \quad 4N\kappa - (N^2 - 1)\frac{1}{2}(\chi(X) + \sigma(X)).$$

In general, the index of the elliptic complex \mathcal{D}_A is given by (2.3).

The ASD equation (2.1) can be perturbed by changing the metric on X . *Holonomy perturbations* determine another useful family of perturbations of the ASD equation [16; 80; 27; 19; 52; 58]. By abuse of notation, a solution of the perturbation of the ASD equation by a holonomy perturbation is still called an ASD connection, and the moduli space of the solutions of the perturbed equation is still denoted by $\mathcal{M}_\kappa(X, w)$. Suppose w is coprime to N and $b^+(X) \geq 1$. Then for a generic choice of the metric on X and a small holonomy perturbation the moduli space $\mathcal{M}_\kappa(X, w)$ consists of only irreducible and regular connections [52]. Therefore, the moduli space is a smooth manifold whose dimension is given by (2.3). This manifold is also orientable and in the case that N is odd [52], a canonical choice of orientation can be fixed. If N is even, to fix an orientation of the moduli space, we need an orientation of the determinant line of the elliptic operator

$$d^+ \oplus d^*: L_1^2(X, \Lambda^1) \rightarrow L_{l-1}^2(X, \Lambda^+) \oplus L_{l-1}^2(X).$$

Such an orientation of the determinant line is called a *homology orientation* of X . The $U(N)$ -polynomial invariants of X are given by integrating appropriate cohomology classes on $\mathcal{M}_\kappa(X, w)$.

The pullback of P to the product space $\mathcal{A}_\kappa(X, w) \times X$ admits an action of $\mathcal{G}_\kappa(X, w)$ which lifts the obvious action on the base. The quotient space defines a $PU(N)$ -bundle \mathbb{P} over $\mathcal{B}_\kappa^*(X, w) \times X$, called the *universal bundle* associated to P . In general, \mathbb{P} cannot be lifted to an $SU(N)$ -bundle. However, we can define Chern classes of \mathbb{P} as rational cohomology classes of $\mathcal{B}_\kappa^*(X, w) \times X$, because the rational cohomology groups of the classifying spaces $BPU(N)$ and $BSU(N)$ are isomorphic [7]. With a slight abuse of notation, we will denote the i^{th} Chern class of \mathbb{P} with $c_i(\mathbb{P})$ for $2 \leq i \leq N$.

The slant product of the Chern classes of the universal bundle and the homology classes of X gives rise to cohomology classes of $\mathcal{B}_\kappa^*(X, w)$. This construction can be used to define an algebra homomorphism,

$$(2.4) \quad \mu: \mathbb{A}(X)^{\otimes(N-1)} \rightarrow H^*(\mathcal{B}_\kappa^*(X, w)),$$

which is the unique algebra homomorphism that satisfies the property

$$(2.5) \quad \mu(\alpha_{(r)}) = (-1)^r c_r(\mathbb{P})/\alpha.$$

Our convention for the definition of μ slightly differs from that of [12] where $(-1)^r$ does not appear in the definition of $\mu(\alpha_{(r)})$.

Let the 2-cycle w be coprime to N , and arrange a metric and a small holonomy perturbation such that the resulting moduli space consists of irreducible and regular points. We will temporarily write $\mathcal{M}_\kappa(X, w, \pi_0)$ for the moduli space to emphasize the dependence on π_0 , which denotes the metric and the holonomy perturbation. If N is even, fix a homology orientation for X . Then $\mathcal{M}_\kappa(X, w, \pi_0)$ can be canonically oriented. Let $d = \dim(\mathcal{M}_\kappa(X, w, \pi_0))$, and fix $z \in \mathbb{A}(X)^{\otimes(N-1)}$ such that $\deg(z) = d$. If the moduli space $\mathcal{M}_\kappa(X, w, \pi_0)$ is compact, then we can evaluate $\mu(z)$ with respect to the fundamental class of $\mathcal{M}_\kappa(X, w, \pi_0)$ and obtain a number $D_{X,w,k}^N(z)$. (Recall that k is the second Chern number of P .) We wish to show that this number does not depend on π_0 , the metric and the holonomy perturbation. Suppose π_1 is another choice of a metric and a small holonomy perturbation avoiding reducible and irregular points. If $b^+(X) \geq 2$, then we can find a path $\{\pi_t\}_{0 \leq t \leq 1}$ of metrics and small holonomy perturbations such that the 1-parameter family of moduli spaces

$$(2.6) \quad \bigcup_t \mathcal{M}_\kappa(X, w, \pi_t)$$

is a smooth manifold of dimension $d + 1$. Because the class $\mu(z)$ can be pulled back to (2.6), Stokes' theorem implies that π_0 and π_1 give rise to the same number $D_{X,w,k}^N(z)$, assuming (2.6) is also compact. However, $\mathcal{M}_\kappa(X, w, \pi_0)$ and the 1-parameter family of moduli spaces are not compact in general and we need to pursue a geometric approach to define the evaluation of $\mu(z)$ on $\mathcal{M}_\kappa(X, w, \pi_0)$.

Uhlenbeck compactification of the moduli space $\mathcal{M}_\kappa(X, w)$ compensates for the noncompactness of this space. Suppose $\{[A_n]\}$ is a sequence of the elements of $\mathcal{M}_\kappa(X, w)$. Then there is a multiset \mathbf{x} of m points in X and a connection $A_\infty \in \mathcal{M}_{\kappa-m}(X, w)$ such that, after passing to a subsequence, $(h_n)_*(A_n)$ is L_1^p -convergent to A_∞ on $X \setminus \mathbf{x}$ for any given real number p [52, Proposition 11]. Here h_n is an isomorphism from the $U(N)$ -bundle carrying A_n to the $U(N)$ -bundle carrying A_∞ , which is defined only on $X \setminus \mathbf{x}$ and whose determinant does not depend on n .

Another key input is that \mathbb{P} can be replaced with a Hermitian vector bundle [12]. Consider the standard representation of $SU(N)$ on \mathbb{C}^N . The tensor product $SU(N)$ -space $(\mathbb{C}^N)^{\otimes N}$ induces a representation of the group $PU(N)$. Therefore, we can

associate a vector bundle \mathbb{E} of rank N^N to \mathbb{P} . We call \mathbb{E} the *universal complex vector bundle*. The Chern class $c_i(\mathbb{P})$ can be written as a polynomial in terms of Chern classes $c_j(\mathbb{E})$ for $2 \leq j \leq i$. For example, $c_2(\mathbb{P})$ is equal to $(1/N^N)c_2(\mathbb{E})$. Therefore, it suffices to define $D_{X,w,k}^N(z)$ for the elements $z \in \mathbb{A}(X)^{\otimes(N-1)}$ such that

$$(2.7) \quad \mu(z) = c_{i_1}(\mathbb{E})/\alpha_1 \cdots c_{i_m}(\mathbb{E})/\alpha_m \quad \text{for } 2 \leq i_j \leq N.$$

In order to define this polynomial invariant, let Σ_i be a submanifold of codimension at least 2 which represents the homology class α_i .

The Chern classes of a vector bundle V of rank r over a manifold M can be represented by stratified subspaces of M . The vector bundle $\text{Hom}(\mathbb{C}^{r-i+1}, V)$ is stratified by rank. If s is a generic section of V , then

$$(2.8) \quad N_i := \{x \in M \mid \text{rank}(s(x)) \leq r - i\}$$

is a stratified subspace of M with strata of even codimension. Therefore, the fundamental class of N_i determines a well-defined homology class which is the Poincaré dual of $c_i(V)$.

In order to define $D_{X,w,k}^N(z)$ for z in (2.7), fix an open neighborhood $\nu(\Sigma_j)$ of the submanifold Σ_j such that the inclusion of this open set in X induces a surjective map of fundamental groups. The unique continuation theorem implies that if the holonomy perturbation in the definition of $\mathcal{M}_\kappa(X, w)$ is small enough, then the restriction of any element of this moduli space to $\nu(\Sigma_j)$ is irreducible [52]. We also assume that $\nu(\Sigma_j)$ and $\nu(\Sigma_k)$ intersect only if Σ_j and Σ_k are embedded surfaces and each point of X lies on at most two such open neighborhoods. Analogous to \mathbb{E} , we can form a vector bundle \mathbb{E}_j of rank N^N on $\mathcal{B}^*(\nu(\Sigma_j)) \times \nu(\Sigma_j)$. In order to produce a representative for $c_{i_j}(\mathbb{E}_j)$, we form the bundle $\text{Hom}(\mathbb{C}^{N^N-i_j+1}, \mathbb{E}_j)$, and form a subspace $V_{i_j}(\Sigma_j)$ of $\mathcal{B}^*(\nu(\Sigma_j)) \times \nu(\Sigma_j)$ as in (2.8).

The vector bundles \mathbb{E} and \mathbb{E}_j are related to each other. The pullback of \mathbb{E}_j with respect to the restriction map $r_j: \mathcal{M}_\kappa(X, w) \times \Sigma_j \rightarrow \mathcal{B}^*(\nu(\Sigma_j)) \times \nu(\Sigma_j)$ is the restriction of the bundle \mathbb{E} . This suggests that $D_{X,w,k}^N(z)$ can be defined as the signed count of the points in the *cut-down moduli space*

$$\mathcal{N}_\kappa(X, w; z) := \{([A], x_1, \dots, x_m) \in \mathcal{M}_\kappa(X, w) \times \Sigma_1 \times \cdots \times \Sigma_m \mid r_j([A], x_j) \in V_{i_j}(\Sigma_j)\}.$$

For a generic choice of $V_{i_j}(\Sigma_j)$, the cut-down moduli space $\mathcal{N}_\kappa(X, w; z)$ is a compact 0-dimensional space, and the orientation of $\mathcal{M}_\kappa(X, w)$ fixes a sign on each point of this space [52; 12]. The compactness of $\mathcal{N}_\kappa(X, w; z)$ is a consequence of Uhlenbeck

compactification using a standard counting argument [17; 20; 52; 12]. Counting the points of the cut-down with respect to the associated signs defines a number which only depends on the homology class of $\Sigma_1, \dots, \Sigma_m$, and this dependence for each homology class is linear. If $b^+(X) \geq 2$, we can adapt the geometric counting argument to the moduli space (2.6), and show that the invariant $D_{X,w,k}^N$ does not depend on the choice of the metric on X and the holonomy perturbation of the ASD equation.

There is also a standard trick which allows us to define $D_{X,w,k}^N$ in the case that w is not coprime to N . Suppose \hat{X} denotes the blowup of X at an arbitrary point. Suppose also E is the exceptional sphere of \hat{X} . Then the cycle $w + E$ is coprime to N . In the non-coprime case, define

$$D_{X,w,k}^N(z) := D_{\hat{X},w+E,k}^N(E_{(2)}^{N-1}z).$$

In the case that w is already coprime, the above definition turns into an identity which is proved in [52].

The invariant $D_{X,w,k}^N(z)$ is defined to be zero if $\deg(z)$ is not equal to the dimension of $\mathcal{M}_\kappa(X, w)$. For a fixed 2-cycle w in X , the moduli spaces of ASD connections appear in different dimensions whose values mod $4N$ are constant. We use (1.4) to define $D_{X,w}^N(z)$, which is called the $U(N)$ -polynomial invariant of (X, w) evaluated at z . This number is nonzero only if

$$\deg(z) \equiv 2(N+1)w \cdot w - (N^2 - 1)\frac{1}{2}(\chi(X) + \sigma(X)) \pmod{4N}.$$

In particular, if N is an odd integer, the invariant $D_{X,w}^N(z)$ vanishes for the classes z such that $\deg(z)$ is not divisible by 4.

There are relationships among the polynomial invariants of X associated to different 2-cycles. If w and w' are two 2-cycles in X , then

$$(2.9) \quad D_{X,w+Nw'}^N(z) = (-1)^{cw' \cdot w'} D_{X,w}^N(z),$$

where c is zero if N is odd or divisible by 4, and is equal to 1 if N is 2 mod 4. The invariants associated to the 2-cycle w and the same 2-cycle with the reverse orientation, denoted by $-w$, are also related. As explained in [52], there is a diffeomorphism from $\mathcal{M}_\kappa(X, w)$ to $\mathcal{M}_\kappa(X, -w)$. This diffeomorphism is orientation-preserving if N is odd, and changes the orientation by the factor of $(-1)^{w \cdot w}$ if N is even. The diffeomorphism lifts to an antilinear isomorphism of the universal complex vector bundles. Therefore,

$$(2.10) \quad D_{X,-w}^N(z) = (-1)^{(N-1)w \cdot w} D_{X,w}^N(\tau(z)),$$

where $\tau: \mathbb{A}(X)^{\otimes(N-1)} \rightarrow \mathbb{A}(X)^{\otimes(N-1)}$ is the algebra homomorphism that maps $\alpha_{(r)}$ to $(-1)^r \alpha_{(r)}$.

Suppose $\Gamma^2, \dots, \Gamma^N$ are elements of $H_2(X)$ and $z \in \mathbb{A}(X)^{\otimes(N-1)}$. To avoid the convergence issue, we define $D_{X,w}^N(z e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N})$ in a slightly different way compared to the introduction:

$$D_{X,w}^N(z e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}) := \sum_{0 \leq i_2, \dots, i_N < \infty} \frac{D_{X,w}(z (\Gamma_{(2)}^2)^{i_2} \dots (\Gamma_{(N)}^N)^{i_N})}{i_2! \dots i_N!} t_2^{i_2} \dots t_N^{i_N},$$

where t_i is a formal variable. Therefore, the $U(N)$ -series $D_{X,w}^N(z e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N})$ is an element of the ring of formal power series $\mathbb{C}[[t_2, \dots, t_N]]$. We can also define an element of $\mathbb{C}[[t_2, \dots, t_N]]$ for linear functionals $f_2, \dots, f_N: H_2(X) \rightarrow \mathbb{C}$ and a power series $g(x) = \sum_{i=0}^{\infty} b_i x^i$:

$$(2.11) \quad g(f_2(\Gamma^2) + \dots + f_N(\Gamma^N)) := \sum_{i=0}^{\infty} b_i (f_2(\Gamma^2)t_2 + \dots + f_N(\Gamma^N)t_N)^i.$$

We use a similar convention in the case that f_i are homogeneous polynomials of higher degree (eg the intersection form Q). These conventions allow us to rephrase Theorems 1, 2 and 3 in terms of the identities of the elements of $\mathbb{C}[[t_2, t_3]]$.

Remark 2.12 Suppose X is a 4-manifold with $b^+(X) = 1$. For a generic metric and for small holonomy perturbations the moduli spaces $\mathcal{M}_\kappa(X, w)$ contains only irreducible connections. For each such choice of metric, we can apply the construction of this section to define a $U(N)$ -polynomial invariant $D_{X,w}^N$. However, this polynomial invariant depends on the choice of the metric, because a 1-parameter family of moduli spaces as in (2.6) might have reducible connections. As a result, the space of metrics on X can be divided into chambers such that $D_{X,w}^N$ is constant only inside the interior of each chamber. Such polynomial invariants have been studied for $N = 2$ in [50; 51].

2.2 Cylindrical ends and moduli spaces

One of the important themes of this article is the interplay between gauge theory on 3-manifolds and 4-manifolds. One can see the interaction by considering the analogues of the geometrical objects from the previous subsection on 4-manifolds with boundary. Suppose W is a 4-manifold with boundary Y , and fix a metric which is a product metric in a collar neighborhood of Y . A smooth 2-cycle in W is a properly embedded 2-dimensional submanifold of W . A 2-cycle w in W is a union of smooth 2-cycles in W whose boundary determines a smooth 1-manifold $\gamma \subset Y$. We will say that the

boundary of the pair (W, w) is the pair (Y, γ) . We also form a pair of noncompact manifolds (W^+, w^+) by adding the cylindrical ends $[0, \infty) \times Y$ and $[0, \infty) \times \gamma$ to W and w . As in the previous part, we can associate a $U(N)$ -bundle Q to the pair (Y, γ) . This $U(N)$ -bundle is trivialized on the complement of a regular neighborhood $D(\gamma)$ of γ , and its relative first Chern class on $D(\gamma)$ is given by the Thom class. We will also write L_γ for the determinant bundle of Q . Similar to the 4-dimensional case, let $\mathcal{B}(Y, \gamma)$ be the space of equivalence classes of connections on Q whose determinants are equal to a fixed connection on L_γ . Given $\alpha \in \mathcal{B}(Y, \gamma)$, the stabilizer of a connection representing α is denoted by Γ_α . The element α is irreducible if Γ_α is equal to the center of $SU(N)$. The bundle Q can be extended to a $U(N)$ -bundle P on W using the 2-cycle w . The bundle P also determines in an obvious way a $U(N)$ -bundle on W^+ , which will be also denoted by P .

Fix an element $\alpha \in \mathcal{B}(Y, \gamma)$, and let A_0 and A_1 be two connections on P whose restrictions to the end $[0, \infty) \times Y$ are the pullback of representatives of α . Then we say A_0 and A_1 represent the same *path* if there is an automorphism g of P with determinant 1 such that $g^*(A_1) - A_0$ is a compactly supported 1-form. The equivalence class of a connection under this equivalence relation is called a *path along (W, w) based at α* . The *topological energy* of a path p represented by a connection A is defined by the following Chern–Weil integral:

$$\kappa(p) := \frac{1}{16N\pi^2} \int_{W^+} \text{tr}(\text{ad}(F(A)) \wedge \text{ad}(F(A))).$$

Here $\text{ad}(F(A))$ is regarded as a 2-form with coefficients in $\text{End}(\mathfrak{su}(P))$. The product \wedge in the integrand is induced by the wedge product of differential forms and composition of the elements of $\text{End}(\mathfrak{su}(P))$. The above integral is independent of the chosen connection A and only depends on p . The constant in front of the integral is chosen such that if p is replaced by another path p' based at α , then the energy changes by an integer.

An important special case for us is the cylinder manifold $W = [0, 1] \times Y$. Fix connections $\alpha, \beta \in \mathcal{B}(Y, \gamma)$ and let p be a path along $([0, 1] \times Y, [0, 1] \times \gamma)$ based at α and β on $\{0\} \times Y$ and $\{1\} \times Y$. Then p induces a path, in the ordinary sense, from α to β in $\mathcal{B}(Y, \gamma)$. The topological energy of the path p defines a number whose value, up to an integer, depends only on α and β . Therefore, we can fix $\beta_0 \in \mathcal{B}(Y, \gamma)$ and define a functional $\text{CS}: \mathcal{B}(Y, \gamma) \rightarrow \mathbb{R}/\mathbb{Z}$, called the *Chern–Simons functional*, where $\text{CS}(\alpha)$ is equal to the topological energy of any path from α to β_0 . Since β_0 is chosen arbitrarily, CS is well defined only up to a constant. But in the case that γ is empty, the trivial

connection Θ gives a canonical choice of β_0 . Critical points of the Chern–Simons functional are represented by connections A on Q such that

$$F_0(A) = 0.$$

A critical point $\alpha \in \mathcal{B}(Y, \gamma)$ is called *nondegenerate* if the Hessian of the Chern–Simons functional is nondegenerate at α .

Suppose α is a nondegenerate critical point of the Chern–Simons functional. Suppose also A_0 is a connection on W^+ that represents a path p based at α . Then $\mathcal{A}_p(W, w; \alpha)$ is the space of connections

$$\mathcal{A}_p(W, w; \alpha) := \{A_0 + a \mid a \in L_{l,\delta}^2(W, \mathfrak{su}(P) \otimes \Lambda^1)\},$$

where $l \geq 3$, δ is a small positive number, and $L_{l,\delta}^2(W, \mathfrak{su}(P))$ is a weighted Sobolev space defined as follows. Let t be a function on W^+ that agrees with the cylindrical coordinate on the end of W^+ . For a vector bundle E on W and a positive constant δ , the Banach space $L_{l,\delta}^2(W, E)$ is defined as $e^{-\delta t} L_k^2(W^+, E)$. Suppose $\mathcal{G}_p(W, w; \alpha)$ is also defined as

$$\mathcal{G}_p(W, w; \alpha) := \{g \in \text{Aut}(P) \mid \det(g) = 1, \nabla_{A_0} g \in L_{l,\delta}^2(W, \mathfrak{su}(P) \otimes \Lambda^1)\}.$$

Then $\mathcal{G}_p(W, w; \alpha)$ is a Banach Lie group. This group acts on $\mathcal{A}_p(W, w; \alpha)$ and the quotient space is denoted by $\mathcal{B}_p(W, w; \alpha)$. The space of elements $[A] \in \mathcal{B}_p(W, w; \alpha)$ such that $F_0^+(A) = 0$ forms the *moduli space of ASD connections associated to the path p* . This moduli space is denoted by $\mathcal{M}_p(W, w; \alpha)$.

It is useful to form a framed version of the moduli space of ASD connections. Any gauge transformation in $\mathcal{G}_p(W, w; \alpha)$ is asymptotic to an element of Γ_α . Define the *framed gauge group* $\mathcal{G}_p^0(W, w; \alpha)$ to be the subspace of the elements of $\mathcal{G}_p(W, w; \alpha)$ which are asymptotic to the trivial element of Γ_α . The framed gauge group is also a Banach Lie group and its Lie algebra can be identified with $L_{l+1,\delta}^2(W, \mathfrak{su}(P))$. We write $\tilde{\mathcal{B}}_p(W, w; \alpha)$ for the quotient of $\mathcal{A}_p(W, w; \alpha)$ with respect to the action of $\mathcal{G}_p^0(W, w; \alpha)$. The set of the elements of $\tilde{\mathcal{B}}_p(W, w; \alpha)$ which satisfy the ASD equation is called the *framed (or based) moduli space of ASD connections associated to p* and is denoted by $\tilde{\mathcal{M}}_p(W, w; \alpha)$.

There is an important relationship between the ASD equation and the Chern–Simons functional. We can define an inner product on $\mathcal{B}(Y, \gamma)$ using

$$\langle a, b \rangle := -\frac{1}{16N\pi^2} \int_Y \text{tr}(\text{ad}(a) \wedge * \text{ad}(b)) \quad \text{for } a, b \in \Omega^1(Y, \mathfrak{su}(P)),$$

where tr is defined using the trace on $\text{End}(\mathfrak{su}(N))$. Suppose $\{\alpha(t)\}_{t \in [0,1]}$ is a path in $\mathcal{B}(Y, \gamma)$. This path defines a trajectory of the downward gradient flow of CS with respect to the above metric if it satisfies

$$(2.13) \quad \frac{d\alpha(t)}{dt} = - * F_0(\alpha(t)).$$

The path $\{\alpha(t)\}$ determines a connection A , in temporal gauge, on $[0, 1] \times Y$, and (2.13) is equivalent to the ASD equation $F_0^+(A) = 0$. This relationship between the ASD equation and the Chern–Simons functional allows us to conclude from nondegeneracy of the critical points of CS that the moduli spaces $\widetilde{\mathcal{M}}_p(W, w; \alpha)$ are analytically well behaved.

The local behavior of the framed moduli space $\widetilde{\mathcal{M}}_p(W, w; \alpha)$ around an element $[A]$ is modeled by the following elliptic complex:

$$(2.14) \quad L_{l+1,\delta}^2(W, \mathfrak{su}(P)) \xrightarrow{d_A} L_{l,\delta}^2(W, \Lambda^1 \otimes \mathfrak{su}(P)) \xrightarrow{d_A^+} L_{l-1,\delta}^2(W, \Lambda^+ \otimes \mathfrak{su}(P)).$$

This complex is Fredholm and its homology groups are denoted by H_A^0 , H_A^1 and H_A^2 . Then $H_A^0 = 0$, and the element $[A]$ is called *regular* if H_A^2 is also trivial. In this case, $\widetilde{\mathcal{M}}_p(W, w; \alpha)$ is smooth in a neighborhood of A , and H_A^1 gives a model for the tangent space of the framed moduli space at $[A]$. Therefore, the index of the above complex for a (not necessarily regular) ASD connection A is called the *expected dimension* of $\widetilde{\mathcal{M}}_p(W, w; \alpha)$ and is denoted by $\text{dim}_e(\widetilde{\mathcal{M}}_p(W, w; \alpha))$.

We slightly generalize the above discussion to include the case of 4–dimensional cobordisms. A cobordism $W: Y_0 \rightarrow Y_1$ is a 4–manifold with boundary $\bar{Y}_0 \amalg Y_1$. We also assume that a 2–cycle $w: \gamma_0 \rightarrow \gamma_1$ in W is given, and P is the associated $U(N)$ –bundle. Suppose also that α_0 and α_1 are flat connections on Y_0 and Y_1 and that p is a path along (W, w) from α_0 to α_1 . As before, we assume that α_0 and α_1 are nondegenerate. In this case, W^+ , $\mathcal{B}_p(W, w; \alpha_0, \alpha_1)$ and $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ are defined as in the previous case by regarding W as a 4–manifold with boundary. However, there is an alternative elliptic complex that one can associate to the elements of $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$. Suppose W^+ is the Riemannian 4–manifold given by adding cylindrical ends to W . In this case, we identify the cylindrical end corresponding to the incoming end with $(-\infty, 0] \times Y_0$. As in the previous case, suppose t is a function on W^+ that agrees with the cylindrical coordinates on the ends and define $L_{l,\delta}^2(W, E)$. Therefore, an element of $L_{l,\delta}^2(W, E)$ is allowed to have exponential growth on the incoming end and it is forced to have an exponential decay on the outgoing end.

For $[A] \in \mathcal{B}_p(W, w; \alpha_0, \alpha_1)$, the ASD operator \mathcal{D}_A is defined by

$$(2.15) \quad d_A^* \oplus d_A^+ : L_{l,\delta}^2(W, \Lambda^1 \otimes \mathfrak{su}(P)) \rightarrow L_{l-1,\delta}^2(W, (\Lambda^0 \oplus \Lambda^+) \otimes \mathfrak{su}(P)).$$

The operator \mathcal{D}_A is elliptic, and the excision property of elliptic operators shows that the index of \mathcal{D}_A is additive with respect to composition of cobordisms [19, Section 3.3]. It can be computed explicitly using the Atiyah–Patodi–Singer index theorem [81; 70]:

$$(2.16) \quad \text{index}(\mathcal{D}_A) = 4N\kappa(p) - (N^2 - 1) \frac{\chi(W) + \sigma(W)}{2} + \sum_{i=0,1} (-1)^i \frac{h^0(Y_i; \text{ad}_{\alpha_i}) - \rho_{\text{ad}_{\alpha_i}}(Y_i)}{2},$$

where $h^0(Y_i; \text{ad}_{\alpha_i})$ denotes the dimension of $H^0(Y_i; \text{ad}_{\alpha_i})$. Moreover, for a flat connection a on a vector bundle V over Y , ρ_a is the Atiyah–Patodi–Singer ρ -invariant of a [2]. As an example, let χ_j be a $U(1)$ -connection on $L(a, b)$ whose holonomy around the standard generator of $\pi_1(L(a, b))$ is equal to $e^{2\pi i j/a}$. It is shown in [2] that

$$(2.17) \quad \rho_{\chi_j}(L(a, b)) = -\frac{4}{a} \sum_{k=1}^{a-1} \cot\left(\frac{\pi k}{a}\right) \cot\left(\frac{\pi k b}{a}\right) \sin^2\left(\frac{\pi k j}{a}\right).$$

Suppose a 4-manifold W is regarded as a cobordism from the empty 3-manifold to the boundary of W . Then $\dim_e(\widetilde{\mathcal{M}}_p(W, w; \alpha))$ is equal to $\text{index}(\mathcal{D}_A) + h^0(\alpha)$ where A is a connection that represents the path p .

In the case of a cylinder $[0, 1] \times Y$, the ASD operator can be used to define a relative $\mathbb{Z}/4N\mathbb{Z}$ -grading on critical points of the Chern–Simons functional associated to a pair (Y, γ) . Fix an arbitrary critical point β_0 of CS, and let α be another critical point of CS. Let the connection A represent an arbitrary path p from α to β_0 . Then the Floer grading of α , denoted by $\text{deg}(\alpha)$, is defined to be $\text{index}(\mathcal{D}_A) \bmod 4N$. Since the choice of β_0 is arbitrary, this grading only gives rise to a relative $\mathbb{Z}/4N\mathbb{Z}$ -grading. In the case that γ is empty, we can make this grading absolute by requiring $\beta_0 = \Theta$. In this case, we can invoke the index formula (2.16) to compute $\text{deg}(\alpha)$ as follows:

$$(2.18) \quad \text{deg}(\alpha) \equiv 4N \cdot \text{CS}(\alpha) - \frac{1}{2}(N^2 - 1) + \frac{1}{2}(h^0(Y; \text{ad}_\alpha) - \rho_{\text{ad}_\alpha}(Y)) \pmod{4N}.$$

Analogous to the case of closed 4-manifolds, we can avoid nonregular points in $\mathcal{M}_p(W, w; \alpha)$ by perturbing the ASD equation. Suppose α is irreducible and nondegenerate. Then there are small holonomy perturbations of the ASD equation, supported in $[0, 1] \times Y \subset W^+$, such that the resulting moduli spaces consist of regular points [58]. Alternatively, if $b^+(X) \geq 2$ and w is coprime to N , then we can arrange for a small

perturbation of the ASD equation such that the moduli spaces $\mathcal{M}_p(W, w; \alpha)$, for arbitrary α and p with $\kappa(p)$ bounded, are regular [52]. In the case that $\mathcal{M}_p(W, w; \alpha)$ consists of only regular points, it is an orientable smooth manifold.

In general, the critical points of the Chern–Simons functional might not be nondegenerate. Suppose that all critical points of the Chern–Simons functional associated to a pair (Y, γ) are irreducible. Then, CS can be also perturbed by appropriate perturbations such that the critical points of the resulting functional are irreducible and nondegenerate [58, Proposition 3.10]. The perturbations used in [58] are also defined in terms of the holonomies of connections on Y . Suppose CS_π is such a perturbation of CS by a small holonomy perturbation and α is a critical point of CS_π . Suppose also (W, w) is a pair whose boundary is equal to (Y, γ) . The negative-gradient flow line of CS_π determines a perturbation of the ASD equation on the end $[0, \infty) \times Y$ of W^+ . This perturbation can be extended to W^+ so that the corresponding moduli space contains only regular points [58]. As in the previous case, the elliptic operator (2.15) can be used to study the local behavior of the moduli spaces. In particular, in the case of a cylinder, the index of \mathcal{D}_A can be used to define a relative $\mathbb{Z}/4N\mathbb{Z}$ -grading on the critical points of CS_π .

2.3 Nonvanishing theorem for algebraic surfaces

For a complex projective surface, the moduli spaces of ASD connections can be identified with the moduli spaces of *stable bundles* with fixed determinant [15]. Stable bundles have been studied extensively using various techniques in algebraic geometry. Thus one can use algebrogeometric methods to extract information about the polynomial invariants of a complex projective surface. For example, the following theorem about nonvanishing of $U(N)$ -polynomial invariants of algebraic surfaces is a generalization of Donaldson’s celebrated theorem about $U(2)$ -invariants [17; 20]:

Theorem 2.19 *Suppose X is a complex projective surface with positive geometric genus, h is a hyperplane class (or equivalently a very ample class), and w is a 2-cycle representing c_1 of a holomorphic line bundle L . Suppose also w is coprime to N . Then*

$$D_{X,w}^N(h_{(2)}^d) > 0$$

when d is large enough and

$$d \equiv (N + 1)w \cdot w - (N^2 - 1)\frac{1}{4}(\chi(X) + \sigma(X)) \pmod{2N}.$$

Proof Let $M_\kappa^N(X, L)$ be the moduli space of stable bundles with rank N , determinant L , and energy κ . We first review the proof of the nonvanishing theorem in the rank 2 case. In this case, the key idea is to find a projective embedding of $M_\kappa^2(X, L)$ and then interpret the polynomial invariant $D_{X,w}^2(h_{(2)}^d)$ as a multiple of the degree of the moduli space. The main steps of the proof can be summarized as follows:

- (1) Suppose C is an algebraic curve and $\mathcal{N}(C, d)$ is the moduli space of stable bundles of rank 2 and degree d on C . Then there is a projective embedding of $\mathcal{N}(C, d)$ into a projective space $\mathbb{P}(W)$ [36]. Moreover, this projective embedding is given by the sections of a large power \mathcal{L}^d of the *determinant line bundle* over $\mathcal{N}(C, d)$ [20].
- (2) For any stable bundle $\mathcal{E} \in M_\kappa^2(X, L)$, there is a p_0 such that if $C \subset X$ is a generic curve in the linear system $|\mathcal{O}(p)|$, for $p \geq p_0$, then the restriction of \mathcal{E} to C is also stable [66]. Using this result together with the fact that the moduli space $M_\kappa^2(X, L)$ has finite type, we can find a projective embedding $J: M_\kappa^2(X, L) \rightarrow \mathbb{P}(V)$. This embedding is given by restricting the elements of $M_\kappa^2(X, L)$ to finitely many curves in the linear system $|\mathcal{O}(p)|$ and applying the Gieseker embedding from the first step.
- (3) For κ large enough, the moduli space $M_\kappa^2(X, L)$ is not empty [79; 37].
- (4) The dimension of the irregular part of $M_\kappa^2(X, L)$ is strictly smaller than the virtual dimension of the moduli space, when κ is large enough [17].

By combining these facts, it can be shown that the map J can be chosen such that the degree of the quasiprojective variety $J(M_\kappa(X, L))$ is equal to

$$K^d D_{X,w}^N(h_{(2)}^d)$$

for an appropriate integer K , and for a large enough d . Therefore, the invariant $D_{X,w}^N(h_{(2)}^d)$ is positive. This proof can be generalized to the case that $N > 2$. Steps 1 and 2 are proved for an arbitrary rank in the references mentioned above. The generalization of the third step to the higher rank case is given in [62]. The higher rank version of the fourth fact is also proved in [38]. Then the same arguments as in [17; 20] can be used to realize $D_{X,w}^N(h_{(2)}^d)$ as a multiple of the degree of a quasiprojective variety, which verifies the claim in [Theorem 2.19](#). \square

Remark 2.20 In [Theorem 2.19](#), we assume that X is a projective surface. But a similar nonvanishing theorem can be formulated for any Kähler surface, because Kähler surfaces can be deformed into projective surfaces [49] and the $U(N)$ -polynomial invariants only depend on the smooth structure.

2.4 Negative embedded spheres

Motivated by [76], Fintushel and Stern used embedded 2–spheres with negative self-intersection to study polynomial invariants of a 4–manifold X [26; 25] (see also [83]). The same idea is exploited in [12] to obtain information about the properties of the polynomial invariants associated to higher rank bundles. Suppose τ is an embedded sphere in a 4–manifold X which has self-intersection -2 . Fix a 2–cycle w with $w \cdot \tau = 0$, and $z \in \mathbb{A}(\langle \tau \rangle^\perp)^{\otimes 2}$. In general, $\mathbb{A}(V)$ for a vector subspace $V \subseteq H_2(X)$ denotes the subalgebra $\text{Sym}^*(H_0(X) \oplus V) \otimes \Lambda^*(H_1(X))$, and $\langle \tau \rangle^\perp$ is the subspace of $H_2(X)$ consisting of the homology classes orthogonal to τ . The following formulas about the $U(3)$ –polynomial invariants of X are proved in [12, Section 3.2]:

- (C1) $D_{X,w}^3(\tau_{(2)}^2 z) = -2 D_{X,w+\tau}^3(z)$.
- (C2) $D_{X,w}^3(\tau_{(2)}^4 z) = -4 D_{X,w}^3(a_2 \tau_{(2)}^2 z) - 3 D_{X,w}^3(\tau_{(3)}^2 z)$.
- (C3) $D_{X,w}^3(\tau_{(2)}^3 \tau_{(3)} z) = -3 D_{X,w}^3(a_3 \tau_{(2)}^2 z) - D_{X,w}^3(a_2 \tau_{(2)} \tau_{(3)} z)$.

By a similar approach, we shall prove the following proposition in Section 6.2:

Proposition 2.21 *Suppose X is a smooth 4–manifold with $b^+(X) \geq 2$ and w is a 2–cycle. Suppose also σ is an embedded sphere with self-intersection -3 and $z \in \mathbb{A}(\langle \sigma \rangle^\perp)^{\otimes 2}$.*

- (i) *If $w \cdot \sigma \equiv 1 \pmod 3$, then there is a constant number c such that*

$$D_{X,w}^3\left(\left(-\frac{3}{2}\sigma_{(3)} - \frac{3}{2}\sigma_{(2)}^2 - a_2\right)z\right) = c D_{X,w-\sigma}^3(z).$$

- (ii) *If $w \cdot \sigma \equiv 2 \pmod 3$, then for the same constant c as above,*

$$D_{X,w}^3\left(\left(\frac{3}{2}\sigma_{(3)} - \frac{3}{2}\sigma_{(2)}^2 - a_2\right)z\right) = c D_{X,w+\sigma}^3(z).$$

- (iii) *If $w \cdot \sigma \equiv 0 \pmod 3$, then the following two formulas hold:*

$$D_{X,w}^3\left(\left(\sigma_{(2)}^4 + 4a_2\sigma_{(2)}^2 + 3\sigma_{(3)}^2\right)z\right) = 0,$$

$$D_{X,w}^3\left(\left(\sigma_{(2)}^3\sigma_{(3)} + 3a_3\sigma_{(2)}^2 + a_2\sigma_{(2)} \cdot \sigma_{(3)}\right)z\right) = 0.$$

In Section 4.2, we will show that the constant c in Proposition 2.21 is equal to -3 .

2.5 Blowup formula for 4–manifolds with simple type

In this subsection, we review the properties of $U(3)$ –polynomial invariants of blown-up 4–manifolds. This part has the same theme as the previous subsection, because the

exceptional divisor of a blown-up 4–manifold gives rise to a (-1) –sphere. We start with an exposition of the main result in Culler’s thesis [12]. Given a 4–manifold X , let \widehat{X} denote the blowup of X at one point, which is diffeomorphic to the 4–manifold $X \# \overline{\mathbb{C}\mathbb{P}^2}$. We will also denote the exceptional sphere in \widehat{X} with E . If w is a 2–cycle in X , then it induces a 2–cycle in \widehat{X} , which will be also denoted by w . Similarly, we can regard $\mathbb{A}(X)$ as a subalgebra of $\mathbb{A}(\widehat{X})$.

Proposition 2.22 *Suppose w is a 2–cycle in X and $z \in \mathbb{A}(X)^{\otimes 2}$. For $0 \leq i \leq 2$ and $0 \leq j \leq 1$, the invariant $D_{\widehat{X},w}^3(E_{(2)}^i E_{(3)}^j z)$ is equal to $D_{X,w}^3(z)$ if $i = j = 0$ and it is zero otherwise. The invariant $D_{\widehat{X},w+E}^3(E_{(3)} z)$ is equal to $D_{X,w}^3(z)$. Finally,*

$$D_{\widehat{X},w+E}^3(E_{(2)}^i z) = \begin{cases} 0 & \text{if } i = 0, 1, 3, \\ D_{X,w}^3(z) & \text{if } i = 2, \\ -D_{X,w}^3(a_2 z) & \text{if } i = 4, \\ D_{X,w}^3(a_3 z) & \text{if } i = 5. \end{cases}$$

Proof As we mentioned in Section 2.1, the identity $D_{\widehat{X},w+E}^3(E_{(2)}^2 z) = D_{X,w}^3(z)$ is proved in [52]. Proposition 62 in [12], together with the discussion succeeding it, essentially contains the proof of the remaining identities using a method similar to that in [52]. To be a bit more detailed, one stretches the neck region in \widehat{X} between X and $\overline{\mathbb{C}\mathbb{P}^2}$, and studies the moduli spaces of ASD connections over \widehat{X} in terms of the moduli spaces associated to X and $\overline{\mathbb{C}\mathbb{P}^2}$. □

According to this proposition, some of the polynomial invariants of \widehat{X} are determined by the invariants of X . The following theorem claims that a similar pattern holds for all invariants of \widehat{X} . This theorem is essentially proved in [12]. We just slightly expand the results of this thesis using the same methods:

Theorem 2.23 *Suppose w is a 2–cycle in X and $z \in \mathbb{A}(X)^{\otimes 2}$. For nonnegative integers i and j , there are polynomials $B_{i,j}, S_{i,j} \in \mathbb{Q}[a_2, a_3]$ independent of X, w and z which satisfy the following identities:*

$$D_{\widehat{X},w}^3(e^{E_{(2)}+E_{(3)}} z) = D_{X,w}^3\left(z \sum_{i,j} B_{i,j}(a_2, a_3) t_2^i t_3^j\right),$$

$$D_{\widehat{X},w+E}^3(e^{E_{(2)}+E_{(3)}} z) = D_{X,w}^3\left(z \sum_{i,j} S_{i,j}(a_2, a_3) t_2^i t_3^j\right).$$

Moreover, $B := \sum_{i,j} B_{i,j}(a_2, a_3)t_2^i t_3^j$ and $S := \sum_{i,j} S_{i,j}(a_2, a_3)t_2^i t_3^j$, as power series in the variables t_2 and t_3 with coefficients in the polynomial ring of a_2 and a_3 , are uniquely determined by the initial values given in Proposition 2.22 and the following four PDEs:

$$(2.24) \quad B_{22}B - B_2B_2 = -S \circ \tau \cdot S, \quad S_{22}S - S_2S_2 = -S \circ \tau \cdot B,$$

$$(2.25) \quad B_{2222}B - 4B_{222}B_2 + 3B_{22}B_{22} = -4a_2(B_{22}B - B_2B_2) - 3(B_{33}B - B_3B_3),$$

$$(2.26) \quad B_{2223}B - 3B_{223}B_2 - B_{222}B_3 + 3B_{23}B_{23} \\ = -3a_3(B_{22}B - B_2B_2) - a_2(B_{23}B - B_2B_3).$$

Here τ maps (t_2, t_3) to $(-t_2, -t_3)$. The subscript 2 means taking the partial derivative with respect to t_2 and the subscript 3 should be interpreted similarly.

Using (2.10), the power series S can be used to compute the invariant $D_{X,w-E}^3$.

Sketch of the proof The main tool is a trick due to Fintushel and Stern [26]. Suppose E and E' are the two exceptional spheres in $X \# 2\overline{\mathbb{C}P}^2$. Then the homology class $E - E'$ is represented by a (-2) -sphere. Suppose $\hat{C}_{k,l}$ (respectively, $\hat{C}'_{k,l}$) is the identity that is given by applying (C_1) from Section 2.4 to w (respectively, $w + E + E'$) and $\tilde{z} := (E + E')^k_{(2)}(E + E')^l_{(3)}z$. Similarly, we can derive identities $\bar{C}_{k,l}$ and $\check{C}_{k,l}$ by applying (C_2) and (C_3) to w and \tilde{z} as above. These identities can be used to prove the existence of $B_{i,j}$ and $S_{i,j}$ inductively. To be a bit more detailed, first one shows the existence of $B_{k,0}$ and $S_{k,0}$ inductively using the initial values in Proposition 2.22, $\hat{C}_{k,0}$ and $\hat{C}'_{k,0}$ [12, Proposition 69]. Then another inductive argument with the aid of the identities $\bar{C}_{k,l}$ and $\check{C}_{k,l}$ and Proposition 2.22 shows the existence of $B_{k,l}$ [12, Proposition 73]. Finally, $\hat{C}_{k,l}$, $\hat{C}'_{k,l}$, the initial values of Proposition 2.22 and the fact that $S_{0,1}$ is nonzero imply the existence of $S_{k,l}$ in an inductive way. After proving the existence of the polynomials $B_{k,l}$ and $S_{k,l}$, the identities $\hat{C}_{k,l}$, $\hat{C}'_{k,l}$, $\bar{C}_{k,l}$ and $\check{C}_{k,l}$ can be used to write four differential equations for B and S which are given in (2.24), (2.25) and (2.26). The existence proof shows that these PDEs and the initial values are enough to uniquely determine B and S . \square

In general, computing the exact form of the power series B and S (equivalently, solving the PDEs in the statement of Theorem 2.23) is not straightforward. In the following corollary, we show that for 4-manifolds with simple type, the blowup formula has a simple form:

Theorem 2.27 Suppose (X, w) is a pair of a 4–manifold and a 2–cycle such that X has w –simple type. Suppose also \hat{X} is the blowup of X at one point and E is the exceptional class. Then there are power series $b(t_2, t_3), s(t_2, t_3) \in \mathbb{Q}[[t_2, t_3]]$ such that

$$(2.28) \quad \hat{D}_{\hat{X},w}(e^{E(2)+E(3)}z) = \hat{D}_{X,w}(z)b(t_2, t_3),$$

$$(2.29) \quad \hat{D}_{\hat{X},w+E}(e^{E(2)+E(3)}z) = \hat{D}_{X,w}(z)s(t_2, t_3)$$

for $z \in \mathbb{A}(X)^{\otimes 2}$. The power series b and s are given by the formulas

$$(2.30) \quad b(t_2, t_3) = \frac{1}{3}e^{-\frac{1}{2}t_2^2+t_3^2}[\cosh(\sqrt{3}t_2) + 2 \cos(\sqrt{3}t_3)],$$

$$(2.31) \quad s(t_2, t_3) = \frac{1}{3}e^{-\frac{1}{2}t_2^2+t_3^2}[\cosh(\sqrt{3}t_2) - \cos(\sqrt{3}t_3) + \sqrt{3} \sin(\sqrt{3}t_3)].$$

Formula (2.30) for the power series $b(t_2, t_3)$ was previously found by means of quantum field theory arguments in [22, Formula 6.22]. When comparing the formulas here and in [22], the reader should note that t_3 needs to be replaced with it_3 because of different conventions in the definition of the μ map.

Proof Evaluating B and S of Theorem 2.23 at $a_2 = 3$ and $a_3 = 0$ produces b and s with the required property. These power series satisfy (2.24), (2.25) and (2.26) with a_2 and a_3 in the last two equations replaced by 3 and 0. In fact, the same proof as the existence proof in Theorem 2.23 shows that b and s are uniquely determined by these equations and the initial values in Proposition 2.22 (with a_2 and a_3 replaced by 3 and 0). The power series (2.30) and (2.31) satisfy the required conditions. Therefore, they are equal to b and s . □

Remark 2.32 Identity (2.10) implies that

$$(2.33) \quad \hat{D}_{\hat{X},w-E}(e^{E(2)+E(3)}z) = \hat{D}_{X,w}(z)s(t_2, -t_3).$$

3 Floer homologies for closed 3–manifolds

3.1 Admissible pairs

A pair (Y, γ) of a connected closed 3–manifold and an embedded oriented 1–manifold is N –admissible if there is an embedded oriented surface Σ in Y such that the integer $\Sigma \cdot \gamma$ is coprime to N . Suppose Q is the $U(N)$ –bundle associated to the pair (Y, γ) . The admissibility of (Y, γ) is what is called the *nonintegral* condition for the $U(N)$ –bundle Q in [58]. In particular, we can use the construction of [58] and

associate the *instanton Floer homology group* $I_*^N(Y, \gamma)$ to an N -admissible pair (Y, γ) . Instanton Floer homology can be lifted to a functor from a cobordism category $\text{COB}_{\mathbb{A}}$ to a certain category of vector spaces.

An object of the category $\text{COB}_{\mathbb{A}}$ is an N -admissible pair. A morphism from a pair (Y_0, γ_0) to another pair (Y_1, γ_1) is a triple (W, w, z) , where $W: Y_0 \rightarrow Y_1$ is a cobordism, $w: \gamma_0 \rightarrow \gamma_1$ is a 2-cycle, and $z \in \mathbb{A}(W)^{\otimes(N-1)}$. The composition of two morphisms

$$(W, w, z): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1) \quad \text{and} \quad (W', w', z'): (Y_1, \gamma_1) \rightarrow (Y_2, \gamma_2)$$

is equal to $(W' \circ W, w' \circ w, z' \cdot z)$.

Suppose VECTOR_n is the category of relatively $\mathbb{Z}/n\mathbb{Z}$ -graded vector spaces over \mathbb{C} . An object of this category is a vector space V with a direct sum decomposition

$$V = \bigoplus_{j \in J} V_j,$$

where $\mathbb{Z}/n\mathbb{Z}$ acts transitively and freely on J . A morphism in this category from $V = \bigoplus_{j \in J} V_j$ to $V' = \bigoplus_{j' \in J'} V'_{j'}$ is a complex linear map $f: V \rightarrow V'$ such that f maps each V_j to a summand $V'_{h(j)}$ of V' such that $h(j+k) = h(j) + k$. Let P-VECTOR_n be the category that has the same objects as VECTOR_n . A morphism in P-VECTOR_n is a vector space homomorphism as above which is well defined only up to a sign. Instanton Floer homology gives a functor $I_*^N: \text{COB}_{\mathbb{A}} \rightarrow \text{P-VECTOR}_{4N}$.

Remark 3.1 The invariant constructed in [58] is more general than the one we described here. In [58], a version of instanton Floer homology is constructed for a triple (Y, γ, K) , where K is a link in Y and γ determines a $U(N)$ -bundle on Y that satisfies a certain nonintegral condition. We need to consider only the case that K is the empty link. On the other hand, in [58], the cobordism maps $I_*^N(W, w, z)$ are defined only in the case that $z = 1$. The more general case is a straightforward generalization and is reviewed below.

For an N -admissible pair (Y, γ) , the vector space $I_*^N(Y, \gamma)$ is defined by applying Morse homological methods to the Chern–Simons functional $\text{CS}: \mathcal{B}(Y, \gamma) \rightarrow \mathbb{R}$. The admissibility condition implies that all critical points of CS are irreducible [58, Proposition 3.1]. Therefore, we can arrange for CS_{π} , a perturbation of the Chern–Simons functional with a small holonomy perturbation, such that all of its critical points are irreducible and nondegenerate [58, Proposition 3.10]. Then CS_{π} has finitely

many critical points. Suppose α and β are two critical points of CS_π and p is a path along $([0, 1] \times Y, [0, 1] \times \gamma)$ based at α and β on $\{0\} \times Y$ and $\{1\} \times Y$. We will write $\mathcal{M}_p(\alpha, \beta)$ for the moduli space of the solutions to the perturbed ASD equation on $\mathbb{R} \times Y$ associated to the path p . Here the perturbation of the ASD equation is induced by the perturbation of the Chern–Simons functional; ie the elements of $\mathcal{M}_p(\alpha, \beta)$ can be regarded as the downward gradient flow lines of CS_π . We can also assume that CS_π is chosen such that the elements of $\mathcal{M}_p(\alpha, \beta)$ for all choices of α, β and p are regular [58, Proposition 3.18]. There is an \mathbb{R} -action on $\mathcal{M}_p(\alpha, \beta)$ given by translation along the \mathbb{R} -factor. The quotient space is denoted by $\check{\mathcal{M}}_p(\alpha, \beta)$. The dimension formula of the previous section implies that

$$\dim(\check{\mathcal{M}}_p(\alpha, \beta)) = \deg(\alpha) - \deg(\beta) - 1 \pmod{4N}.$$

Instanton Floer homology of the pair (Y, γ) is given by the homology of a chain complex (\mathfrak{C}_*^π, d) associated to the functional CS_π . The vector space \mathfrak{C}_*^π is freely generated by the critical points of CS_π . The differential of a generator α of \mathfrak{C}_*^π is also defined as

$$(3.2) \quad d(\alpha) := \sum_{p: \alpha \rightarrow \beta} \# \check{\mathcal{M}}_p(\alpha, \beta) \beta,$$

where the sum is over all paths p such that $\check{\mathcal{M}}_p(\alpha, \beta)$ is 0-dimensional. These 0-dimensional moduli spaces are compact and we orient them as in [58, Section 3.6]. Then $\# \check{\mathcal{M}}_p(\alpha, \beta)$ denotes the signed count of the points in $\check{\mathcal{M}}_p(\alpha, \beta)$. We use the Floer grading \deg to define a relative $\mathbb{Z}/4N\mathbb{Z}$ -grading on \mathfrak{C}_*^π . Then the differential d decreases this grading by 1. The relatively $\mathbb{Z}/4N\mathbb{Z}$ -graded vector space $\mathbb{I}_*^N(Y, \gamma)$ is defined to be the homology of the chain complex (\mathfrak{C}_*^π, d) . This homology group is independent of the choice of the Riemannian metric on Y and the perturbation of the Chern–Simons functional.

Suppose the Chern–Simons functional associated to the pair (Y, γ) is *Morse–Bott*. That is to say, the set of critical points of CS is a smooth manifold, and the Hessian of CS is invertible only in the normal direction to the critical manifold. Following the above definition, we need to work with perturbations of the Chern–Simons functional to define $\mathbb{I}_*^N(Y, \gamma)$. However, one can still derive some information about $\mathbb{I}_*^N(Y, \gamma)$ using the unperturbed Chern–Simons functional:

Proposition 3.3 *If the Chern–Simons functional of the pair (Y, γ) is Morse–Bott, then $\dim(\mathbb{I}_*^N(Y, \gamma)) \leq \dim(H_*(\text{crit}(\text{CS})))$.*

Proof This claim can be verified using the standard spectral sequence that starts from the homology of the critical manifold of CS and abuts to the instanton Floer homology of (Y, γ) . \square

Suppose $(W, w): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1)$ is a cobordism of N -admissible pairs. After fixing Riemannian metrics on Y_0 and Y_1 , we form the noncompact manifold W^+ by adding cylindrical ends to W . Suppose also a perturbation of the Chern–Simons functional for (Y_i, γ_i) is fixed such that we can form the chain complex $(\mathfrak{C}_*^{\pi_i}, d)$ as above. We use a perturbation of the ASD equation on W^+ which is compatible with the chosen perturbations of the Chern–Simons functionals. Given a generator α_i of $\mathfrak{C}_*^{\pi_i}$ and a path $p: \alpha_0 \rightarrow \alpha_1$ along (W, w) , let $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ be the moduli space of the corresponding equation. We can pick a perturbation of the ASD equation on W such that this moduli space is a smooth manifold. By fixing a homology orientation of W , we can also assume that all such moduli spaces are oriented [58]. Moreover, if $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ is 0-dimensional, then it is compact.

Define a map $\mathfrak{C}(W, w): \mathfrak{C}_*^{\pi_0} \rightarrow \mathfrak{C}_*^{\pi_1}$ by

$$(3.4) \quad \mathfrak{C}(W, w)(\alpha_0) := \sum_{p: \alpha_0 \rightarrow \alpha_1} \# \mathcal{M}_p(W, w; \alpha_0, \alpha_1) \alpha_1,$$

where the sum is over all paths such that $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ is 0-dimensional. The term $\# \mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ is equal to the signed count of the points in $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$. As a matter of fact, (3.4) defines a chain map and the induced map at the level of homology, $\mathbf{I}_*^N(W, w, 1): \mathbf{I}_*^N(Y_0, \gamma_0) \rightarrow \mathbf{I}_*^N(Y_1, \gamma_1)$, determines a morphism of the category P-VECTOR_{4N} . More generally, given $z \in \mathbb{A}(W)^{\otimes(N-1)}$, we can cut down the moduli space $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ using z as in Section 2.1, and construct a smooth submanifold $\mathcal{M}_p(W, w; \alpha_0, \alpha_1, z)$ such that

$$(3.5) \quad \dim(\mathcal{M}_p(W, w; \alpha_0, \alpha_1, z)) = \dim(\mathcal{M}_p(W, w; \alpha_0, \alpha_1)) - \deg(z).$$

To be more precise, $\mathcal{M}_p(W, w; \alpha_0, \alpha_1, z)$ is a linear combination of the spaces whose dimensions are given by (3.5). Replacing $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ in the right-hand side of (3.4) with $\mathcal{M}_p(W, w; \alpha_0, \alpha_1, z)$ determines a new chain map and the associated map at the level of homology is $\mathbf{I}_*^N(W, w, z): \mathbf{I}_*^N(Y_0, \gamma_0) \rightarrow \mathbf{I}_*^N(Y_1, \gamma_1)$. This map is well defined; namely, it does not depend on the metric on W , the perturbation of the ASD equation and the choices involved in the construction of $\mathcal{M}_p(W, w; \alpha_0, \alpha_1, z)$. Because we fix the central part of connections associated to (W, w) , the cobordism maps only depend on the induced $\text{PU}(N)$ -bundles. In particular, if w is replaced with $w + nw'$, for

a closed 2–cycle w' , then the induced $\mathrm{PU}(N)$ –bundles are the same and they determine the same cobordism maps. This property is the analogue of (2.9) for closed 4–manifolds.

Remark 3.6 A priori it might seem that the cycles in 3–manifolds and 4–dimensional cobordisms only keep track of the first Chern classes of $\mathrm{U}(N)$ –bundles. However, they have strictly more information than these cohomology classes. For example, an element of $H^2(Y, \mathbb{Z})$, for a 3–manifold Y , determines a $\mathrm{U}(N)$ –bundle up to an isomorphism of $\mathrm{U}(N)$ –bundles. But an embedded 1–manifold γ in Y determines a $\mathrm{U}(N)$ –bundle up to a canonical isomorphism. (See [57, Section 4] for more details.) As a manifestation of this issue, suppose cohomology classes α and α' are given on cobordisms $W: Y_0 \rightarrow Y_1$ and $W': Y_1 \rightarrow Y_2$ such that the restrictions of these classes on Y_1 agree with each other. Then there might be an ambiguity in gluing these cohomology classes to construct an element of $H^2(W' \circ W; \mathbb{Z})$. On the other hand, there is no such ambiguity if α and α' are represented by embedded surfaces $w \subset W$ and $w' \subset W'$ such that $w|_{Y_1} = w'|_{Y_1}$.

Remark 3.7 There is not much difficulty in extending the definition of instanton Floer homology to the case of disconnected 3–manifolds. Suppose Y is a disconnected 3–manifold and $\gamma \subset Y$ is a 1–cycle. Then we say (Y, γ) is N –admissible if for each connected component Y_0 of Y , the pair $(Y_0, \gamma \cap Y_0)$ is N –admissible. Then we can repeat the definition analogous to the connected case and construct instanton Floer homology for the pair (Y, γ) . This instanton Floer homology can be computed in terms of the invariants for the connected components:

$$I_*^N(Y, \gamma) := I_*^N(Y_1, \gamma_1) \otimes \cdots \otimes I_*^N(Y_n, \gamma_n).$$

Here the Y_i are connected components of Y and $\gamma_i = \gamma \cap Y_i$. It is also possible to define cobordism maps for a cobordism of pairs (W, w) between two (not necessarily connected) N –admissible pairs and $z \in \mathbb{A}(W)^{\otimes(N-1)}$. However, these maps are not as well behaved with respect to composition as in the previous case. Consider two triples

$$(W, w, z): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1) \quad \text{and} \quad (W', w', z'): (Y_1, \gamma_1) \rightarrow (Y_2, \gamma_2).$$

If Y_1 is not connected, what we can say about the cobordism maps is that

$$I_*^N(W' \circ W, w' \circ w, z \cdot z') = c \cdot I_*^N(W', w', z') \circ I_*^N(W, w, z)$$

for some nonzero constant c . The simplest way to fix this issue about functoriality is to work with a variation of the category $\mathrm{P}\text{-VECTOR}_{4N}$ where the morphisms are well defined only up to a nonzero scalar. We follow this approach when it is necessary to work with disconnected 3–manifolds.

Remark 3.8 A slightly unsatisfying point about I_*^N is the sign ambiguity in the definition of the cobordism maps. This issue can be avoided in a straightforward way. In the case that N is an even number, we need to change the definition of the category $\text{COB}_{\mathbb{A}}$ slightly. Let $\widetilde{\text{COB}}_{\mathbb{A}}$ have the same objects as $\text{COB}_{\mathbb{A}}$. But a morphism of this new category is a quadruple (W, w, z, o_W) where W, w and z are as before and o_W is a homology orientation for W . Then I_*^N can be lifted to a functor $\widetilde{I}_*^N: \widetilde{\text{COB}}_{\mathbb{A}} \rightarrow \text{VECTOR}_{4N}$. The main point is that initially there is an ambiguity in the orientation of the moduli spaces $\mathcal{M}_p(W, w; \alpha_0, \alpha_1)$ that appear in the definition of the cobordism maps, and a homology orientation of W fixes this ambiguity. In the case that N is odd, there is not such ambiguity and one can readily lift I_*^N to $\widetilde{I}_*^N: \text{COB}_{\mathbb{A}} \rightarrow \text{VECTOR}_{4N}$.

The definition of cobordism maps can be extended to the case that one of the ends is the empty pair. Suppose X is a 4-manifold with boundary Y and w is a properly embedded surface in X such that $\gamma := \partial w = w \cap Y$. Assume that (Y, γ) is an N -admissible pair. Given any element $z \in \mathbb{A}(X)^{\otimes(N-1)}$, we can form an element $D_{X,w}^N(z)$ of $I_*^N(Y, \gamma)$. This construction is the extension of the $U(N)$ -polynomial invariants for closed 4-manifolds in the previous section. Alternatively, (X, w, z) can be regarded as a cobordism from the empty pair to the N -admissible pair (Y, γ) . Although the empty pair is not N -admissible, (3.4) can be used to define $D_{X,w}^N(z)$.

Remark 3.9 Given $z_i \in \mathbb{A}(X_i)^{\otimes(N-1)}$, we can consider the relative elements

$$D_{X_i, w_i}^N(z_i) \in I_*^N(Y, \gamma).$$

Each of these relative elements lies in a graded summand of $I_*^N(Y, \gamma)$. Therefore, the difference $\text{deg}(D_{X_2, w_2}^N(z_2)) - \text{deg}(D_{X_1, w_1}^N(z_1))$ of the relative $\mathbb{Z}/4N\mathbb{Z}$ -gradings is a well-defined number in $\mathbb{Z}/4N\mathbb{Z}$ and is equal to

$$2(N+1)(w_2^2 - w_1^2) - (N^2 - 1) \left(\frac{\chi(X_2) + \sigma(X_2)}{2} - \frac{\chi(X_1) + \sigma(X_1)}{2} \right) - (\text{deg}(z_2) - \text{deg}(z_1)).$$

Note that the term w_i^2 is not well defined and depends on a framing of the 1-cycle γ . However, the difference $w_2^2 - w_1^2$ is independent of the framing and the above expression is well defined. A similar formula can be written for the difference between the gradings of two cobordisms with the same ends.

Suppose (X, w) is a cobordism from an N -admissible pair (Y, γ) to the empty pair; namely, X is a 4-manifold whose boundary is identified with \bar{Y} , the 3-manifold Y with the reverse orientation. The boundary of the embedded surface w is also identified

with $\bar{\gamma}$. Suppose also $z \in \mathbb{A}(X)^{\otimes(N-1)}$. Similarly, we can construct a functional $D_N^{X,w}(z): I_*^N(Y, \gamma) \rightarrow \mathbb{C}$.

Similar to cobordism maps, $D_{X,w}^N(z)$ and $D_N^{X,w}(z)$ satisfy some functorial properties. For example, if (X, w, z) and (W, w', z') are chosen such that

$$\partial(X, w) = (Y_0, \gamma_0) \quad \text{and} \quad (W, w', z'): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1),$$

then

$$D_{W \circ X, w' \circ w}^N(z \cdot z') = I_*^N(W', w', z') \circ D_{X,w}^N(z).$$

A similar property holds for $D_N^{X,w}(z)$. There is also an important relation among these invariants and invariants of closed manifolds from the previous section:

Proposition 3.10 *Suppose Y is a closed and connected 3–manifold, (Y, γ) is an N –admissible pair, and X_1 and X_2 are two smooth 4–manifolds with $\partial X_1 = Y$ and $\partial X_2 = \bar{Y}$. Suppose also that oriented properly embedded surfaces $w_i \subset X_i$ are given such that $\partial w_1 = \gamma$ and $\partial w_2 = \bar{\gamma}$. If $b^+(X_2 \circ X_1) \geq 2$, then for $z_i \in \mathbb{A}(X_i)^{\otimes(N-1)}$,*

$$(3.11) \quad D_{X_2 \circ X_1, w_2 \circ w_1}^N(z_1 \cdot z_2) = D_N^{X_2, w_2}(z_2) \circ D_{X_1, w_1}^N(z_1).$$

If $b^+(X_2 \circ X_1) = 1$, then a similar formula holds where the left-hand side of (3.11) is interpreted as the invariant of the chamber associated to metrics with a long neck along Y .

There is another way that the invariants of a closed 4–manifold can be related to the cobordism maps:

Proposition 3.12 *Let (Y, γ) be an N –admissible pair and $(W, w): (Y, \gamma) \rightarrow (Y, \gamma)$ a cobordism of pairs. Let W and Y be connected. Let \tilde{W} be the closed 4–manifold given by gluing the incoming end of W to its outgoing end. Suppose also that $\tilde{w} \subset \tilde{W}$ is defined similarly. If $b^+(\tilde{W}) \geq 2$, then for $z \in \mathbb{A}(W)^{\otimes(N-1)}$,*

$$(3.13) \quad D_{\tilde{W}, \tilde{w}}^N(z) = N \cdot \text{str}(I_*^N(W, w, z)),$$

where str denotes the supertrace of $I_^N(W, w, z)$.⁶ If $b^+(\tilde{W}) = 1$, then a similar formula holds where the left-hand side of (3.13) is interpreted as the invariant of the chamber associated to metrics with a long neck along Y .*

⁶Since the $\mathbb{Z}/4N\mathbb{Z}$ –grading on instanton Floer homology is defined only relatively, there is a sign ambiguity on the right-hand side, even after fixing a homology orientation for W . There is also a sign ambiguity on the left-hand side because we did not fix any homology orientation for \tilde{W} . Therefore, (3.13) should be interpreted as an equality up to sign. Although we do not need to do so here, both sign ambiguities can be fixed after fixing a homology orientation for W .

3.2 Floer homology of $S^1 \times \Sigma$

Suppose Σ is the connected oriented Riemann surface of genus g . The 3-manifold $Y_g := \Sigma \times S^1$ and the embedded 1-manifold $\gamma_{g,d} := \{x_1, \dots, x_d\} \times S^1$ determine an N -admissible pair if $(d, N) = 1$. The $U(N)$ -bundle associated to the pair $(Y_g, \gamma_{g,d})$ is the pullback of a $U(N)$ -bundle Q_d of degree d on Σ . Let also L_d denote the determinant of Q_d . Recall that the space $\mathcal{A}(Y_g, \gamma_{g,d})$ is constructed using an auxiliary connection on $L_{\gamma_{g,d}}$. We assume that this connection is the pullback of a connection B_0 on L_d . Similar to the 3-dimensional case, $\mathcal{A}(\Sigma, Q_d)$ is defined to be the space of $U(N)$ -connections on Q_d whose determinants are equal to B_0 . The space \mathcal{G}_d is also defined to be the group of determinant 1 automorphisms of Q_d .

Let $\mathbb{V}_{g,d}^N$ be the vector space $I_*^N(Y_g, \gamma_{g,d})$. The critical points of the Chern–Simons functional for the pair $(Y_g, \gamma_{g,d})$ can be identified with N copies of the space

$$\mathcal{N}_{N,d} := \{A \in \mathcal{A}(\Sigma, Q_d) \mid F_0(A) = 0\} / \mathcal{G}_d.$$

In fact, we can pull back any element of $\mathcal{N}_{N,d}$ to $\Sigma \times [0, 1]$ and then identify the connections on $\Sigma \times \{0\}$ and $\Sigma \times \{1\}$ using an element of \mathcal{G}_d which is induced by a central element of $SU(N)$. For any connection A representing an element of $\mathcal{N}_{N,d}$,

$$(3.14) \quad h^0(\Sigma, \text{ad}_A) = 0, \quad h^1(\Sigma, \text{ad}_A) = (N^2 - 1)(2g - 2) \quad \text{and} \quad h^2(\Sigma, \text{ad}_A) = 0,$$

where $h^i(\Sigma, \text{ad}_A)$ again denotes the dimension of the cohomology group $H^i(\Sigma, \text{ad}_A)$. This implies the space $\mathcal{N}_{N,d}$ is a smooth manifold of dimension $(N^2 - 1)(2g - 2)$ [1]. If α is a critical point of the Chern–Simons functional of the pair $(Y_g, \gamma_{g,d})$ associated to the connection A and a central element of \mathcal{G}_d , then the Künneth formula and the identities in (3.14) imply that $H^1(Y_g, \text{ad}_\alpha)$ is isomorphic to $H^1(\Sigma, \text{ad}_A)$. Consequently, the Chern–Simons functional of $(Y_g, \gamma_{g,d})$ is Morse–Bott, and hence $\dim(\mathbb{V}_{g,d}^N) \leq N \dim(H^*(\mathcal{N}_{N,d}))$.

The space $\mathcal{N}_{N,d}$ has been extensively studied in the literature. This space is a Kähler manifold and can be identified with the moduli space of *stable bundles* of rank N and degree d on a Riemann surface of genus g with a fixed determinant. The Poincaré polynomial of this manifold can be computed inductively [42; 14; 1]. Furthermore, a set of generators for the cohomology ring of this space is given [1]. We review a slightly reformulated description of these generators which are more suitable for our purposes here.

Consider the 4-manifold $X_g := \Sigma \times S^2$ and the surface $\chi_g := \{x_1, \dots, x_d\} \times S^2$. The pullback of the elements of $\mathcal{N}_{N,d}$ to X_g are ASD connections associated to

the pair (X_g, χ_g) with $\kappa = 0$. In particular, $\mathcal{N}_{N,d}$ can be regarded as a subset of $\mathcal{B}_0(X_g, \chi_g)$. Consider the subalgebra $\mathbb{A}_g^N := \mathbb{A}(\Sigma)^{\otimes(N-1)}$ of $\mathbb{A}(X_g)^{\otimes(N-1)}$. The μ -map in (2.4) determines a graded algebra homomorphism $\Psi: \mathbb{A}_g^N \rightarrow H^*(\mathcal{N}_{N,d})$. Note that this map is equivariant with respect to $\text{Diff}(\Sigma)$, the group of diffeomorphisms of Σ .

Proposition 3.15 *The map Ψ is surjective. In particular, the cohomology ring of $\mathcal{N}_{N,d}$ is generated by the elements*

$$(3.16) \quad p_r := \Psi(a_r), \quad q_r^j := \Psi(l_{(r)}^j) \quad \text{and} \quad s_r := \Psi(\Sigma_{(r)}),$$

where $2 \leq r \leq N$ and $\{l^j\}_{1 \leq j \leq 2g}$ forms a set of generators for $H_1(\Sigma, \mathbb{Z})$.

The action of $\text{Diff}(\Sigma)$ on $H_1(\Sigma)$ factors through the action of the symplectic group $\text{Sp}(2g)$. Therefore, this proposition implies that the same holds for $H^*(\mathcal{N}_{N,d})$.

Proof In [1], a universal $U(N)$ -bundle \mathbb{F} is constructed over the product manifold $\mathcal{N}_{N,d} \times \Sigma$ and it is shown that the cohomology ring of $\mathcal{N}_{N,d}$ is generated by the classes

$$(3.17) \quad \tilde{p}_r := c_r(\mathbb{F})/[a], \quad \tilde{q}_r^j := c_r(\mathbb{F})/[l^j] \quad \text{and} \quad \tilde{s}_r := c_r(\mathbb{F})/[\Sigma].$$

The map Ψ is defined similarly using the universal $PU(N)$ -bundle \mathbb{P} over the space $\mathcal{B}_\kappa(X_g, \chi_g) \times X_g$. The restriction of \mathbb{P} to $\mathcal{N}_{N,d} \times \Sigma \subset \mathcal{B}_\kappa(X_g, \chi_g) \times X_g$ is isomorphic to the $PU(N)$ -bundle associated to \mathbb{F} . Thus the cohomology classes in (3.17) can be identified with the corresponding ones in (3.16) and this verifies the claim. \square

The vector space $\mathbb{V}_{g,d}^N$ admits a ring structure which is the analogue of the cup product on $H^*(\mathcal{N}_{N,d})$. Suppose P is the pair of pants cobordism from two copies of S^1 to one copy of S^1 . Then the triple $(\Sigma \times P, \{x_1, \dots, x_d\} \times P, 1)$ defines a map $m: \mathbb{V}_{g,d}^N \otimes \mathbb{V}_{g,d}^N \rightarrow \mathbb{V}_{g,d}^N$. To be more precise, we need to fix a homology orientation in the case that N is even. Suppose $\Delta_g := \Sigma \times D^2$ and $\delta_{g,d} := \{x_1, \dots, x_d\} \times D^2$, where D^2 is the 2-dimensional disc. We fix an arbitrary homology orientation on Δ_g and let $e := D_{\Delta_g, \delta_{g,d}}^N(1) \in \mathbb{V}_{g,d}^N$. We shall see in the proof of Proposition 3.23 that e is nonzero. We choose a homology orientation on $\Sigma \times P$ such that $m(e, e) = e$. Then functoriality of instanton Floer homology can be used to show that m defines a ring structure on $\mathbb{V}_{g,d}^N$ with unit e . We turn the relative $\mathbb{Z}/4N\mathbb{Z}$ -grading on $\mathbb{V}_{g,d}^N$ into an absolute grading by requiring that the unit element have degree 0. With this convention, the multiplication map is $\mathbb{Z}/4N\mathbb{Z}$ -graded; namely, the product of two elements of degree i_1 and i_2 has degree $i_1 + i_2$.

Suppose B is a cylinder, regarded as a cobordism with two circles as the incoming end and empty outgoing end. Then the pair

$$(3.18) \quad \Omega_g := \Sigma \times B \quad \text{and} \quad \omega_g := \{x_1, \dots, x_d\} \times B$$

determines a pairing $\langle \cdot, \cdot \rangle: \mathbb{V}_{g,d}^N \otimes \mathbb{V}_{g,d}^N \rightarrow \mathbb{C}$ which is defined in the following way after we choose an arbitrary homology orientation on Ω_g :

$$\langle \alpha, \alpha' \rangle = D_N^{\Omega_g, \omega_g}(\alpha \otimes \alpha') \quad \text{for } \alpha, \alpha' \in \mathbb{V}_{g,d}^N$$

Proposition 3.19 *The space $\mathbb{V}_{g,d}^N$ as a complex vector space with an action of $\text{Diff}(\Sigma)$ is isomorphic to $H^*(\mathcal{N}_{N,d}(\Sigma))[u]/(u^N - 1)$. In particular, the action of $\text{Diff}(\Sigma)$ on $\mathbb{V}_{g,d}^N$ factors through an action of $\text{Sp}(2g)$*

When $N = 2$, one can show that $\mathbb{V}_{g,d}^N$ is isomorphic to $H^*(\mathcal{N}_{N,d}(\Sigma))[u]/(u^N - 1)$ using the results of [21]. Muñoz [71] gave an alternative proof of this proposition for $N = 2$, and the proof of the general case is based on his approach.

Proof We can define a $\text{Diff}(\Sigma)$ -equivariant algebra homomorphism

$$\Phi: \mathbb{A}_g^N[u]/(u^N - 1) \rightarrow \mathbb{V}_{g,d}^N \quad \text{by} \quad \Phi(u^i z) := D_{\Delta_g, \delta_{g,d} + i\Sigma}^N(z),$$

where on the right-hand side of the defining equation z is regarded as an element of $\mathbb{A}(\Delta_g)^{\otimes(N-1)}$. Let $S: H^*(\mathcal{N}_{N,d}(\Sigma)) \rightarrow \mathbb{A}_g^N$ be a graded linear $\text{Sp}(2g)$ -equivariant right inverse for the map Ψ . Extend Ψ to an algebra homomorphism from $\mathbb{A}_g^N[u]/(u^N - 1)$ to $H^*(\mathcal{N}_{N,d}(\Sigma))[u]/(u^N - 1)$ by requiring that $\Psi(u) = u$. Similarly, we can extend S to $H^*(\mathcal{N}_{N,d}(\Sigma))[u]/(u^N - 1)$. We claim that the $\text{Diff}(\Sigma)$ -equivariant map $\Phi \circ S: H^*(\mathcal{N}_{N,d}(\Sigma))[u]/(u^N - 1) \rightarrow \mathbb{V}_{g,d}^N$ is injective. If the claim does not hold, then there is

$$p = \sum_{m=1}^M z_m u^{i_m} \in \mathbb{A}_g^N[u]/(u^N - 1) \quad \text{with } z_m \in \mathbb{A}_g^N \text{ and } 0 \leq i_m < N$$

such that $\Psi(p) \neq 0$ and $\Phi(p) = 0$. We assume that each z_m is nonzero and lies in one of the graded summands of \mathbb{A}_g^N . Furthermore, if $m < n$, then $\deg(z_m) \geq \deg(z_n)$ and equality holds only if $i_m \neq i_n$. Let $z' \in \mathbb{A}_g^N$ be such that

$$(3.20) \quad \deg(z') + \deg(z_1) = (N^2 - 1)(2g - 2)$$

and the cup product of $\Psi(z')$ and $\Psi(z_1)$ is nonzero. By Proposition 3.10, the pairing $\langle \Phi(p), \Phi(u^{N-i_1} z') \rangle$ is equal to

$$(3.21) \quad \sum_{m=1}^M D_{\Sigma \times S^2, w_{g,d} + (N+i_m-i_1)\Sigma}(z' z_m),$$

where the polynomial invariants are computed in the chamber such that the fiber Σ is small. The expression for the dimension in (2.3) shows that the dimension of each component of the moduli space associated to the pair $(\Sigma \times S^2, w_{g,d} + k\Sigma)$ is at least $(N^2 - 1)(2g - 2)$. Moreover, if k is not divisible by N , this dimension is strictly greater than $(N^2 - 1)(2g - 2)$. In the case that $k = 0$ (or divisible by N), the moduli space of dimension $(N^2 - 1)(2g - 2)$ is given by the pullback of the elements of $\mathcal{N}_{N,d}$ to X_g . Therefore, the only nonzero term in (3.21) is $D_{\Sigma \times S^2, w_{g,d}}(z' z_1)$, which is given by evaluating the cohomology class $\mu(z' z_1)$ on the pullback of the elements of $\mathcal{N}_{N,d}$ to X_g . Note that the moduli space is compact in this case and we do not need to use the geometric representatives to evaluate this invariant. Therefore, $D_{\Sigma \times S^2, w_{g,d}}(z' z_1)$ is equal to $\Psi(z') \cup \Psi(z)[\mathcal{N}_{N,d}]$, which is nonzero by assumption. This contradicts the assumption that $\Phi(p) = 0$. Therefore, the map $\Phi \circ S$ is injective. We already know that the dimension of $\mathbb{V}_{g,d}^N$ is not greater than $N \dim(H^*(\mathcal{N}_{N,d}))$. Therefore, $\Phi \circ S$ is a bijection. In particular, the action of $\text{Diff}(\Sigma)$ on $\mathbb{V}_{g,d}^N$ factors through an action of $\text{Sp}(2g)$. \square

Corollary 3.22 *The ring $\mathbb{V}_{g,d}^N$ is generated by the elements*

$$\begin{aligned} \epsilon &= D_{\Delta_g, \delta_{g,d} + \Sigma}(1), & \mathfrak{S}_r &= D_{\Delta_g, \delta_{g,d}}(a_r), \\ o_r^j &= D_{\Delta_g, \delta_{g,d}}(l_{(r)}^j), & \rho_r &= D_{\Delta_g, \delta_{g,d}}(\Sigma(r)), \end{aligned}$$

where $2 \leq r \leq N$ and $1 \leq j \leq 2g$. Furthermore, the $\mathbb{Z}/4N\mathbb{Z}$ -gradings of these elements are given by

$$\deg(\epsilon) = 4d, \quad \deg(\mathfrak{S}_r) = -2r, \quad \deg(o_r^j) = -2r + 1, \quad \deg(\rho_r) = -2r + 2.$$

Proof The first part is an immediate consequence of Proposition 3.19. The second part can be also verified easily using Remark 3.9. \square

Fix $S: H^*(\mathcal{N}_{N,d}) \rightarrow \mathbb{A}_g^N$ as in the proof of Proposition 3.19. Then we can use the isomorphism $\Phi \circ S$ to pull back the ring structure, the pairing and the $\mathbb{Z}/4N\mathbb{Z}$ -grading of $\mathbb{V}_{g,d}^N$ into $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$. Suppose the new multiplication map and the pairing on $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$ are also denoted by m and $\langle \cdot, \cdot \rangle$. We also fix the cohomological \mathbb{Z} -grading on $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$, where we set $\deg(u) = 0$. The

cohomological and the $\mathbb{Z}/4N\mathbb{Z}$ -gradings differ by a sign after collapsing into $\mathbb{Z}/4\mathbb{Z}$ -gradings. In the proof of Proposition 3.19, we show that for any element p of degree i in $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$, there is an element q of degree $(N^2 - 1)(2g - 2) - i$ with $\langle p, q \rangle \neq 0$. In particular, \langle, \rangle defines a nondegenerate pairing.

Suppose p_1 and p_2 are two elements of degrees i_1 and i_2 in $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$. Then the product $m(p_1, p_2)$ consists of terms in various gradings. However, there are some constraints on the degrees of these terms. Firstly, they all have the same $\mathbb{Z}/4\mathbb{Z}$ -grading because the multiplication map is graded with respect to the $\mathbb{Z}/4N\mathbb{Z}$ -grading on $\mathbb{V}_{g,d}^N$. Moreover, an argument similar to that of Proposition 3.19 shows that the pairing of $m(p_1, p_2) - p_1 \cup p_2$ and any element of $H^*(\mathcal{N}_{N,d})[u]/(u^N - 1)$ with cohomological degree less than or equal to $(N^2 - 1)(2g - 2) - i_1 - i_2$ is zero. Therefore, the degree of the terms in $m(p_1, p_2)$ are at most $i_1 + i_2$ and the term with the maximal degree is $p_1 \cup p_2$. Therefore, the product m is a deformation of the cup product. We summarize these properties of $\mathbb{V}_{g,d}^N$ in the following proposition:

Proposition 3.23 *The pairing \langle, \rangle on $\mathbb{V}_{g,d}^N$ is nondegenerate and the product*

$$m: \mathbb{V}_{g,d}^N \times \mathbb{V}_{g,d}^N \rightarrow \mathbb{V}_{g,d}^N$$

is a deformation of the cup product preserving the $\mathbb{Z}/4\mathbb{Z}$ -grading.

Multiplication with the elements of $\mathbb{V}_{g,d}^N$, constructed in Corollary 3.22, defines a series of operators on $\mathbb{V}_{g,d}^N$. We will use the same notation to denote these operators. The operator ϵ can be alternatively described as the cobordism map associated to the triple $([0, 1] \times Y_g, [0, 1] \times \gamma_{g,d} \cup \Sigma, 1)$. Similarly the remaining operators are cobordism maps associated to triples $([0, 1] \times Y_g, [0, 1] \times \gamma_{g,d}, z)$ for appropriate choices of z . The operators ϵ , \mathfrak{N}_r and ρ_r commute with each other and o_r^j . However, o_r^j and $o_{r'}^{j'}$ anticommute with each other.

In the special case that $g = 1$, the moduli space $\mathcal{N}_{N,d}(\Sigma)$ consists of only one point. Therefore, $\mathbb{V}_{1,d}^N$ has N generators with exactly one generator α_i in degree $4i$ with respect to $\mathbb{Z}/4N\mathbb{Z}$ -grading. In fact, the (nonperturbed) Chern–Simons functional associated to the N -admissible pair $(Y_1, \gamma_{1,d})$ has irreducible and nondegenerate critical points (see [52]). The operator ϵ maps α_i to α_{i+d} . The following proposition characterizes the action of some of the point classes in the case that $d = 1$:

Proposition 3.24 *The operators $\mathfrak{N}_i: \mathbb{V}_{1,1}^N \rightarrow \mathbb{V}_{1,1}^N$ satisfy the identities*

$$(3.25) \quad \mathfrak{N}_2 = N\epsilon^{-1} \quad \text{and} \quad \mathfrak{N}_{2i-1} = 0.$$

Proof The second identity can be verified easily, because $\deg(\mathfrak{N}_{2i-1})$ is not divisible by 4. The first claim is proved in [85]. Since \mathfrak{N}_2 and ϵ commute with each other and $\deg(\mathfrak{N}_2) = -4$, we can conclude that $\mathfrak{N}_2 = c\epsilon^{-1}$. Therefore, we just need to show that $\text{tr}(\mathfrak{N}_2 \circ \epsilon) = N^2$. Using Proposition 3.12, this can be reduced to showing that

$$D_{T^4, T^2 \times \{\text{pt}\} \cup \{\text{pt}\} \times T^2}^N(\mathfrak{N}_2) = N^3,$$

which is established in [85] using properties of stable bundles on abelian varieties. \square

In a work in progress, we will give another proof of this proposition which is independent of the results of [85].

3.3 Fukaya–Floer homology

Suppose X_1 and X_2 are 4–manifolds with $\partial X_1 = Y$ and $\partial X_2 = \bar{Y}$, and X is given by gluing these manifolds along their boundaries. Suppose also a 1–cycle $\gamma \subset Y$ and 2–cycles $w_i \subset X_i$ are chosen such that $\partial w_1 = \gamma$ and $\partial w_2 = \bar{\gamma}$. The cycles w_1 and w_2 can be glued to each other to form a 2–cycle $w \subset X$. We also assume that (Y, γ) is an N –admissible pair. Then Floer homology for this N –admissible pair provides a useful device to relate $U(N)$ –polynomial invariants of the form

$$(3.26) \quad D_{X,w}^N(z_1 \cdot z_2) \quad \text{with } z_i \in \mathbb{A}(X_i)^{\otimes(N-1)}$$

to the relative invariants associated to (X_1, w_1, z_1) and (X_2, w_2, z_2) (see Proposition 3.10). As we shall see in the next section, this decomposition theorem for polynomial invariants is a useful tool for computational purposes. However, not all polynomial invariants of (X, w) have the form in (3.26). There are homology classes $\Gamma \in H_2(X)$ such that Γ is not the sum of elements in $H_2(X_1)$ and $H_2(X_2)$. Then, for example, $D_{X,w}^N(\Gamma_{(2)}^i \Gamma_{(3)}^j)$ cannot be expressed in terms of the relative invariants. In this section, we introduce an extension of Floer homology which admits relative invariants for such polynomial invariants. This extension of Floer homology was already constructed in [33; 9] for $N = 2$ and is known as *Fukaya–Floer homology*.

Our extension of Floer homology is a module over a ring R_N . Let $R_{N,j}$ be the polynomial ring over the variables $t_{k,i}$ for $2 \leq k \leq N$ and $1 \leq i \leq j$, modulo the relations $t_{k,i}^2 = 0$:

$$R_{N,j} := \mathbb{C}[t_{k,i} : 2 \leq k \leq N, 1 \leq i \leq j] / (t_{k,i}^2).$$

For $j \geq l \geq 0$, there is an obvious map from $R_{N,j}$ to $R_{N,l}$ which maps $t_{k,i}$ to $t_{k,i}$ when $i \leq l$ and maps $t_{k,i}$ to 0 when $i > l$. The ring R_N is defined to be the inverse

limit of this system of rings. For example, for each $2 \leq k \leq N$, we have an element of R_N as follows:

$$s_k := \sum_{i=1}^{\infty} t_{k,i}.$$

The ring of polynomials $\mathbb{C}[[t_2, \dots, t_N]]$ can be regarded as a subring of R_N by mapping t_k to $s_k \in R_N$. Under this inclusion we have

$$\frac{t_k^l}{l!} \rightarrow \sum_{\substack{S \subset \mathbb{N} \\ |S|=l}} \prod_{i \in S} t_{k,i}.$$

The full version of Fukaya–Floer homology for $N = 2$, whose construction is sketched in [9], is expected to be a module over $\mathbb{C}[[t_2]]$. However, our construction is slightly different and we obtain a module over the ring R_N for general N . This is partly because the definition of polynomial invariants for higher rank bundles slightly differs from the classical definition of $U(2)$ –polynomial invariants. Another reason is that even for $N = 2$, the authors were not able to avoid some analytical difficulties related to the noncompactness of the moduli space of ASD connections and construct a ring over $\mathbb{C}[[t_2]]$.

Consider the N –admissible pair (Y, γ) , and let $L = (l_2, \dots, l_N)$ be an $(N-1)$ –tuple of the elements of $H_1(Y)$. Fukaya–Floer homology associates to (Y, γ, L) an $R_{N,j}$ –module $\mathbb{I}_*^{N,j}(Y, \gamma, L)$ for each nonnegative integer j and an $R_{N,j}$ –module homomorphism $f_j^k: \mathbb{I}_*^{N,j}(Y, \gamma, L) \rightarrow \mathbb{I}_*^{N,k}(Y, \gamma, L)$ for each pair (j, k) of nonnegative integers such that $j \geq k \geq 0$. If $j \geq k \geq l \geq 0$, we require $f_k^l \circ f_j^k = f_j^l$. Fukaya–Floer homology of (Y, γ, L) is the inverse limit of the inverse system $(\{\mathbb{I}_*^{N,j}(Y, \gamma, L)\}_j, \{f_j^k\}_{j \geq k})$. The $R_{N,j}$ –module $\mathbb{I}_*^{N,j}(Y, \gamma, L)$ is the homology of a chain complex $(\mathfrak{C}_*^{N,j}(Y, \gamma, L), d_{N,j})$. In fact, we can arrange for a perturbation CS_{π_j} of the Chern–Simons functional of the N –admissible pair (Y, γ) such that $\mathfrak{C}_*^{N,j}(Y, \gamma, L) = \mathfrak{C}_*^{\pi_j}(Y, \gamma) \otimes_{\mathbb{C}} R_{N,j}$.

The differential $d_{N,j}$ of the Fukaya–Floer chain complex has the form

$$(3.27) \quad d_{N,j}(\alpha) = \sum_{\substack{\bar{S}=(S_2, \dots, S_N) \\ p: \alpha \rightarrow \beta}} h_{\bar{S}}(\alpha, \beta) \left(\prod_{i \in S_k} t_{k,i} \right) \beta,$$

where $S_k \subset \{1, \dots, j\}$ and the path p is chosen such that the dimension of the moduli space $\mathcal{M}_p(\alpha, \beta)$ is equal to

$$2|S_2| + 4|S_3| + \dots + 2(N-1)|S_N| + 1.$$

The constant term of the differential is equal to the differential d of the Floer chain complex. That is to say, if we evaluate all variables $t_{k,i}$ at zero, then we recover d . The definitions of the other terms in (3.27) are discussed in Section 6.3. We extend the Floer grading to $\widetilde{\mathfrak{C}}_*^{N,j}(Y, \gamma, L)$ by requiring that $\deg(t_{k,i}) = 2(k - 1)$. Then the differential $d_{N,j}$ has degree -1 .

Suppose (X_1, w_1) is a pair of a 4-manifold and a 2-cycle which fills the N -admissible pair (Y, γ) . Suppose also $z_1 \in \mathbb{A}(X_1)^{\otimes(N-1)}$ and $\Gamma^2, \dots, \Gamma^N$ are properly embedded surfaces in X_1 where $[\partial\Gamma^i] = l_i \subset H_1(Y)$. Then one can associate an element of $\mathbb{I}_*^N(Y, \gamma, L)$ to $(X_1, w_1, z_1, \Gamma^2, \dots, \Gamma^N)$, which is denoted by

$$(3.28) \quad D_{X_1, w_1}^N(z_1 \cdot e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}).$$

The element in (3.28) is given by a system of cycles $D_{X_1, w_1}^{N,j}(z_1 \cdot e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}) \in \mathfrak{C}_*^{N,j}(Y, \gamma, L)$. We have

$$D_{X_1, w_1}^{N,j}(z_1 \cdot e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}) = \sum_{\bar{s}=(S_2, \dots, S_N), \alpha} m_{\bar{s}}(\alpha) \left(\prod_{i \in S_k} t_{k,i} \right) \alpha$$

for appropriate choices of complex numbers $m_{\bar{s}}(\alpha)$. Evaluating all the variables $t_{k,i}$ at zero produces a cycle in $\mathfrak{C}_*^{\pi j}(Y, \gamma)$ which represents the relative invariant $D_{X_1, w_1}^N(z_1)$.

Next, let (X_2, w_2) be a cobordism from (Y, γ) to the empty pair. Suppose that $z_2 \in \mathbb{A}(X_2)^{\otimes(N-1)}$ and $\Lambda_2, \dots, \Lambda_N$ are properly embedded surfaces in X_2 where $[\partial\Lambda^j] = -l_j$. In this case, there is an R_N -linear map from $\mathbb{I}_*^N(Y, \gamma, L)$ to R_N associated to $(X_2, w_2, z_2, \Lambda^2, \dots, \Lambda^N)$, which is denoted by

$$(3.29) \quad D_N^{X_2, w_2}(z_2 \cdot e^{\Lambda_{(2)}^2 + \dots + \Lambda_{(N)}^N}).$$

The construction of the element (3.28) and that of the functional (3.29) are given in Section 6.3. We can glue the 4-manifolds (X_1, w_1) and (X_2, w_2) to form a closed pair $(X_2 \circ X_1, w_2 \circ w_1)$. The embedded surfaces Γ^j and Λ^j can also be glued to each other to form a closed embedded surface $\Gamma^j \# \Lambda^j$. Then we have

$$(3.30) \quad D_{X_2 \circ X_1, w_2 \circ w_1}^N(z_1 \cdot z_2 \cdot e^{(\Gamma^2 \# \Lambda^2)_{(2)} + \dots + (\Gamma^N \# \Lambda^N)_{(N)}}) \\ = D_N^{X_2, w_2}(z_2 \cdot e^{\Lambda_{(2)}^2 + \dots + \Lambda_{(N)}^N}) \circ D_{X_1, w_1}^N(z_1 \cdot e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}).$$

This claim shall be proved as Proposition 6.51 in Section 6.3. A priori, the right-hand side of the above equality is an element of R_N . Part of the claim is that the right-hand side belongs to $\mathbb{C}[[t_2, \dots, t_N]] \subset R_N$ and is equal to the given power series on the left-hand side.

Fukaya–Floer homology is also functorial with respect to cobordisms of N –admissible pairs. Suppose $(W, w): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1)$ is such a cobordism. For $2 \leq i \leq N$, suppose also Γ^i is a properly embedded surface in W such that $\Gamma^i \cap Y_j$ represents the homology class $l_i^j \in H_1(Y_j)$. For any $z \in \mathbb{A}(W)$, there is a homomorphism

$$\mathbb{I}_*^N(W, w, ze^{\Gamma^2_2 + \dots + \Gamma^N_{(N)}}): \mathbb{I}_*^N(Y_0, \gamma_0, L_0) \rightarrow \mathbb{I}_*^N(Y_1, \gamma_1, L_1),$$

where $L_j = (l_2^j, \dots, l_N^j)$. This construction is functorial with respect to composition of cobordisms.

Suppose $(Y_g, \gamma_{g,d})$ is the N –admissible pair from Section 3.2 and $L_g = (l_2, \dots, l_N)$ is the $(N-1)$ –tuple of the elements of $H_1(Y_g)$ where l_i is an S^1 fiber. In this article, the main example of Fukaya–Floer homology for us is $\mathbb{I}_*^N(Y_g, \gamma_{g,d}, L_g)$, which is denoted by $\mathbb{I}_{g,d}^N$. Analogously to the previous subsection, we can define a ring homomorphism

$$\tilde{\Phi}: \mathbb{A}_g^N[u]/(u^N - 1) \rightarrow \mathbb{I}_{g,d}^N \quad \text{by} \quad \tilde{\Phi}(u^i z) := D_{\Delta_g, \delta_{g,d} + i\Sigma}^N(ze^{D^2_{(2)} + \dots + D^2_{(N)}}).$$

Recall that D^2 is a 2–dimensional disc, $\Delta_g = \Sigma \times D^2$ and $\delta_{g,d} = \{x_1, \dots, x_d\} \times D^2$. Similarly, we can define a multiplication and a pairing on $\mathbb{I}_{g,d}^N$ by repeating the construction of Section 3.2.

Proposition 3.31 *For any nonzero element $q \in \mathbb{I}_{g,d}^N$, there are $z \in \mathbb{A}_g^N$ and $1 \leq i \leq N$ such that*

$$\langle q, D_{\Delta_g, \delta_{g,d} + i\Sigma}^N(ze^{D^2_{(2)} + \dots + D^2_{(N)}}) \rangle \neq 0.$$

Proof Suppose q is given by the sequence $\{q_j\}_{j \geq 0}$, where $q_j \in \mathbb{I}_*^{N,j}(Y_g, \gamma_{g,d}, L_g)$. Since q is nonzero, there is a j such that q_j is nonzero. To verify the claim, it suffices to show that there are $z \in \mathbb{A}_g^N$ and $1 \leq i \leq N$ such that

$$\langle q_j, D_{\Delta_g, \delta_{g,d} + i\Sigma}^{N,j}(ze^{D^2_{(2)} + \dots + D^2_{(N)}}) \rangle \neq 0.$$

Suppose $(\mathfrak{C}_*^{\pi_j}(Y_g, \gamma_{g,d}), d)$ is a Floer chain complex for the pair $(Y_g, \gamma_{g,d})$ such that $\mathfrak{C}_*^{\pi_j}(Y_g, \gamma_{g,d}) \otimes R_{N,j}$ can be used to define $\mathbb{I}_*^{N,j}(Y_g, \gamma_{g,d}, L_g)$. Suppose q_j is represented by the element

$$\sum_{\bar{S} = (S_2, \dots, S_N) \in \mathcal{I}} b_{\bar{S}} \left(\prod_{i \in S_k} t_{k,i} \right) \alpha_{\bar{S}}$$

of this complex, where \mathcal{I} is a set consisting of $(N-1)$ –tuples of finite subsets of $\{1, \dots, j\}$, each $b_{\bar{S}}$ is a nonzero complex number and $\alpha_{\bar{S}} \in \mathfrak{C}_*^{\pi_j}(Y_g, \gamma_{g,d})$. Suppose $\bar{S}_0 = (S_2^0, \dots, S_N^0) \in \mathcal{I}$ is a minimal element with respect to the partial order on \mathcal{I}

induced by inclusion. Then $d\alpha_{\bar{S}_0} = 0$, and by changing the representative if necessary, we can assume that $\alpha_{\bar{S}_0}$ is not a boundary in $\mathfrak{C}_*^{\pi_j}(Y_g, \gamma_{g,d})$. Therefore, by Corollary 3.22 and Proposition 3.23, there are $z \in \mathbb{A}_g^N$ and $1 \leq i \leq N$ such that

$$\langle \alpha_{\bar{S}_0}, D_{\Delta_g, \delta_{g,d} + i\Sigma}^N(z) \rangle \neq 0.$$

This implies that the coefficient of $\prod_{2 \leq k \leq N} \prod_{i \in S_k^0} t_{k,i}$ in the pairing

$$\langle q_j, D_{\Delta_g, \delta_{g,d} + i\Sigma}^{N,j}(ze^{D_{(2)}^2 + \dots + D_{(N)}^2}) \rangle.$$

is nonzero. □

Example 3.32 In the case that $g = 1$, the Fukaya–Floer homology group of the triple $(Y_g, \gamma_{g,d}, L_g)$ is equal to $R_N^{\oplus N}$. This can be easily seen from the construction of Fukaya–Floer homology in Section 6.3. In fact, this R_N –module is freely generated by the critical points $\alpha_1, \dots, \alpha_N$ of the Chern–Simons functional of $(Y_1 = T \times S^1, \gamma_{1,d})$. Consider the operator

$$(3.33) \quad \tilde{\epsilon} := \mathbb{I}_*^N([0, 1] \times Y_1, [0, 1] \times \gamma_{1,d} + T, e^{([0,1] \times \gamma)_{(2)} + \dots + ([0,1] \times \gamma)_{(N)}}),$$

where γ is an S^1 –fiber of Y_1 . This operator is of order N and has degree $4d$. Moreover, $\alpha_i, \tilde{\epsilon}(\alpha_i), \dots, \tilde{\epsilon}^{N-1}(\alpha_i)$ form a basis for $\mathbb{I}_*^N((Y_g, \gamma_{g,d}, L_g))$ for any i . In particular, the kernel of $\tilde{\epsilon} - 1$ is equal to $R_N \cdot (1 + \tilde{\epsilon} + \dots + \tilde{\epsilon}^{N-1})(\alpha_i)$.

Next, we discuss a prototype for 4–manifolds with boundary Y_g which are of interest to us. Suppose (X_1, Σ) is a pair of a 4–manifold and an embedded surface of genus g with self-intersection 0 such that a regular neighborhood of Σ in X_1 is identified with Δ_g . Suppose (X_2, Σ) is another such pair. As the notation suggests, the embedded surfaces in X_1 and X_2 are identified with each other. Remove regular neighborhoods of Σ in X_1 and X_2 to produce 4–manifolds whose boundaries are Y_g , and then glue the resulting two 4–manifolds along their common boundaries by the orientation-reversing diffeomorphism that maps $(z, x) \in S^1 \times \Sigma$ to (\bar{z}, x) . This 4–manifold is denoted by $X_1 \#_{\Sigma} X_2$, and is called the *fiber sum* of X_1 and X_2 along Σ . We will also write X_i° for the complement of a neighborhood of Σ in X_i . Then X_i° can be also regarded as a subspace of $X_1 \#_{\Sigma} X_2$.

Elements of $H_2(X_1)$ and $H_2(X_2)$ can be glued to each other to construct elements of $H_2(X_1 \#_{\Sigma} X_2)$. Suppose $\iota_i: H_2(X_i) \rightarrow \mathbb{C}$ denotes the map that computes the intersection number of an element of $H_2(X_i)$ with Σ . Suppose also \mathcal{K} is the subspace

of the elements $(\Gamma, \Lambda) \in H_2(X_1) \oplus H_2(X_2)$ such that $\iota_1(\Gamma) = \iota_2(\Lambda)$. Then there is a homomorphism $\#: \mathcal{K} \rightarrow H_2(X_1 \#_{\Sigma} X_2)$ with the property that

$$(3.34) \quad j_1^{\#}(\Gamma \# \Lambda) = j_1^{\circ}(\Gamma) \quad \text{and} \quad j_2^{\#}(\Gamma \# \Lambda) = j_2^{\circ}(\Lambda).$$

Here $j_i^{\circ}: H_2(X_i) \rightarrow H_2(X_i^{\circ}, \partial X_i^{\circ})$ is the composition of the map from $H_2(X_i)$ to the relative homology $H_2(X_i, \Delta_g)$ and the excision isomorphism. To abbreviate our notation, from now on, we will write Γ° and Λ° for $j_1^{\circ}(\Gamma)$ and $j_2^{\circ}(\Lambda)$. The maps $j_i^{\#}: H_2(X_1 \#_{\Sigma} X_2) \rightarrow H_2(X_i^{\circ}, \partial X_i^{\circ})$ are also defined similarly.

The homomorphism $\#$ is not uniquely defined and we proceed as follows to fix one such homomorphism. Suppose Γ and Λ are integral homology classes. Then these homology classes can be represented by oriented embedded surfaces, which we denote with the same notation. We can assume that these surfaces are transversal to Σ , and intersect Σ in the same set of points with the same signs. Then there is an obvious way to glue Γ and Λ and to produce an oriented embedded surface in $X_1 \#_{\Sigma} X_2$. The homology class $\Gamma \# \Lambda$ is defined to be the homology of the glued-up surface. We apply this construction to an integral basis of \mathcal{K} and extend it linearly.

We use Poincaré duality to define $\mathcal{L} \subseteq H^2(X_1) \oplus H^2(X_2)$, the counterpart of \mathcal{K} , and the gluing map $\#: \mathcal{L} \rightarrow H^2(X_1 \# X_2)$. Suppose $(K, L) \in \mathcal{L}$ and $(\Gamma, \Lambda) \in \mathcal{K}$. Then we have the equality of the pairing of cohomology classes with homology classes

$$(K \# L)[\Gamma \# \Lambda] = K[\Gamma] + L[\Lambda].$$

Similarly, we can glue two cycles $w_1 \subset X_1$ and $w_2 \subset X_2$ that intersect Σ transversely in the same set of points with the same signs. The resulting 2-cycle in $X_1 \#_{\Sigma} X_2$ is denoted by $w_1 \# w_2$. We will also write w_i° for the intersection $w_i \cap X_i^{\circ}$.

Proposition 3.35 *For $1 \leq i \leq 4$, suppose X_i is a 4-manifold and T is an embedded surface of genus one in X_i . Suppose also $w_i \subset X_i$ is a 2-cycle such that $w_i \cdot T$ is coprime to N . For each $2 \leq l \leq N$, suppose also Γ_i^l is an element of $H_2(X_i)$ such that $\Gamma_i^j \cdot T = 1$. For $1 \leq i, k \leq 4$, let $D_{i,k}$ be the following element of $\mathbb{C}[[t_2, \dots, t_N]]$:*

$$\sum_{j=1}^N D_{X_i \# T X_k, w_i \# w_k + jT}^N (e^{(\Gamma_i^2 \# \Gamma_k^2)_{(2)} + \dots + (\Gamma_i^N \# \Gamma_k^N)_{(N)}}).$$

Then

$$D_{1,2} D_{3,4} = D_{1,4} D_{3,2}.$$

Proof The following elements lie in the kernel of the operator $\tilde{\epsilon}$ in (3.33):

$$D_i := \sum_{j=1}^N D_{X_i^\circ, w_i^\circ + jT}^N (e^{(\Gamma_i^2)^\circ_{(2)} + \dots + (\Gamma_i^N)^\circ_{(N)}}).$$

Moreover, (3.30) implies that

$$(3.36) \quad D_{i,k} = \frac{1}{N} \langle D_i, D_k \rangle.$$

Because $D_i \in \ker(\tilde{\epsilon} - 1)$, the claim is a consequence of the description of $\ker(\tilde{\epsilon} - 1)$ in Example 3.32. □

3.4 An SU(3)–instanton Floer homology for $\Sigma(2, 3, 23)$

In Section 3.1, Floer homology is defined for an N –admissible pair (Y, γ) . Then the computation of the polynomial invariants for a closed pair (X, w) which can be decomposed along a copy of (Y, γ) can be reduced to computing relative invariants for each component of $X \setminus Y$ (Proposition 3.10). One wishes to extend the definition of Floer homology so that it can be used in studying polynomial invariants of a pair (X, w) which is decomposed along a nonadmissible pair. However, there is little known in this direction even when $N = 2$. For $N = 2$, the most satisfactory answer is provided in the case that Y is an integral homology sphere and γ is empty [27], which will be denoted by $I_*^2(Y)$. The main reason that one can define $I_*^2(Y)$ for an integral homology sphere is that the only reducible flat connection on Y is the nondegenerate trivial connection. In order to extend $I_*^N(Y)$ to higher values of N , one would face more complicated reducible connections. Due to this complication, there are some difficulties in extending the definition of Floer homology of integral homology spheres to higher values of N . In this section, we make a modest progress in this direction and define $I_*^3(Y)$ for the Brieskorn homology sphere $\Sigma(2, 3, 23)$. Meanwhile, we compute some of the gauge-theoretical invariants for flat connections on $\Sigma(2, 3, 23)$.

Suppose the positive integers a_1, a_2 and a_3 are pairwise coprime, and $\Sigma(a_1, a_2, a_3)$ is the associated Brieskorn sphere:

$$\Sigma(a_1, a_2, a_3) := \{(z_1, z_2, z_3) \in \mathbb{C}^3 \mid z_1^{a_1} + z_2^{a_2} + z_3^{a_3} = 0, |z_1|^2 + |z_2|^2 + |z_3|^2 = 1\}.$$

This 3–manifold is an integral homology sphere. There is an S^1 –action on this 3–manifold,

$$e^{2\pi i \theta} \cdot (z_1, z_2, z_3) := (e^{2\pi i a_2 a_3 \theta} z_1, e^{2\pi i a_1 a_3 \theta} z_2, e^{2\pi i a_1 a_2 \theta} z_3).$$

This action turns $\Sigma(a_1, a_2, a_3)$ into a Seifert fiber space over S^2 with three exceptional orbits. Complex conjugation on \mathbb{C}^3 induces a diffeomorphism of $\Sigma(2, 3, 23)$, which will be also called complex conjugation. There is also a standard presentation of the fundamental group of this 3-manifold, given as

$$(3.37) \quad \pi_1(\Sigma(a_1, a_2, a_3)) = \langle x_1, x_2, x_3, h \mid [h, x_i] = 1, x_i^{a_i} h^{\beta_i} = 1, x_1 x_2 x_3 = 1 \rangle,$$

where β_i is given by the identity

$$\frac{\beta_1}{a_1} + \frac{\beta_2}{a_2} + \frac{\beta_3}{a_3} = \frac{1}{a}.$$

with $a = a_1 a_2 a_3$. The central element h in (3.37) is represented by a generic fiber of the Seifert fibration.

Suppose W is the space $(\Sigma(a_1, a_2, a_3) \times D^2)/S^1$, where the S^1 -action is the product of the Seifert action on $\Sigma(a_1, a_2, a_3)$ and the standard action on D^2 . Alternatively, W is the mapping cylinder of the fibration of $\Sigma(a_1, a_2, a_3)$ over S^2 . This space is an orbifold and has three singular points. Neighborhoods of these singular points are diffeomorphic to cones on the lens spaces $L(a_i, \beta_i)$. Therefore, removing neighborhoods of the orbifold points produces a cobordism W_0 from the union of three lens spaces $L(a_i, \beta_i)$ to $\Sigma(a_1, a_2, a_3)$. We will denote the union of the lens spaces by Y .

The fundamental group of W_0 is equal to $\pi_1(\Sigma(a_1, a_2, a_3))/\langle h \rangle$ and the inclusion of $\Sigma(a_1, a_2, a_3)$ in W_0 induces the quotient map at the level of fundamental groups. Moreover, the induced map from the fundamental group of $L(a_i, \beta_i)$ to the group $\pi_1(\Sigma(a_1, a_2, a_3))/\langle h \rangle$ maps the standard generator of $\pi_1(L(a_i, \beta_i))$ to x_i . The description of the fundamental group implies that the first homology of W_0 is trivial. We also have the short exact sequence

$$0 \rightarrow H^2(W_0, \partial W_0, \mathbb{Z}) \cong \mathbb{Z} \xrightarrow{\iota} H^2(W_0, \mathbb{Z}) \cong \mathbb{Z} \rightarrow H^2(\partial W_0, \mathbb{Z}) \cong \mathbb{Z}/a\mathbb{Z} \rightarrow 0,$$

where the map ι is multiplication by a . The self-intersection pairing, defined on the image of ι , maps a generator of $\text{im}(\iota)$ to $-a$. In particular, $b^+(W_0)$ is equal to 0.

The space $L := \Sigma(a_1, a_2, a_3) \times D^2$ defines an orbifold S^1 -bundle on W . In particular, the restriction of L to W_0 , denoted by L_0 , is a smooth S^1 -bundle. The first Chern class of L_0 is a generator of $H^2(W_0, \mathbb{Z})$. The restriction of this Chern class to $L(a_i, \beta_i)$ is equal to β_i times the standard generator of $H^2(L(a_i, \beta_i), \mathbb{Z})$. In particular, for any complex line bundle on ∂W_0 , there is a k such that the restriction of L_0^k to the boundary is isomorphic to the given line bundle.

The rational cohomology class induced by $c_1(L_0)$ can be lifted to $H^2(W_0, \partial W_0, \mathbb{Q})$. In particular, $c_1(L_0)^2$ is well defined and is equal to $-1/a$. For our purposes, we also fix a connection B_0 on L_0 whose restriction to a neighborhood of ∂W_0 is the pullback of a flat connection on ∂W_0 . In particular, we can assume that the restriction of this connection in a regular neighborhood of $\Sigma(a_1, a_2, a_3)$ is the trivial connection.

Suppose α is a flat $SU(N)$ -connection on $\Sigma(a_1, a_2, a_3)$ whose holonomy around the fiber is central. Then α can be extended as a flat $PU(N)$ -connection A to W_0 . The holonomy of α induces a conjugacy class r_i in $PU(N)$ corresponding to the loop x_i . The class r_i has order a_i and determines a flat $PU(N)$ -connection on $\pi_1(L(a_i, \beta_i))$ which matches the restriction of A to $L(a_i, \beta_i)$. This connection will be also denoted by r_i . As it was pointed in [24], the connection A can be used to compute some of the gauge-theoretical invariants of α :

Proposition 3.38 *Let α and A be given as above. Then $\rho_{\text{ad}(\alpha)}$ is equal to*

$$(3.39) \quad (T_\alpha + 1 - N^2) + \sum_{i=1}^3 \rho_{\text{ad}(r_i)}(L(a_i, \beta_i)),$$

where T_α is the number of trivial summands in the irreducible decomposition of the representation associated to ad_α .

Using the calculations of [2] for lens spaces, (3.39) allows us to compute the ρ -invariant of any connection as above.

Proof According to [2],

$$\rho_{\text{ad}(\alpha)} - \sum_{i=1}^3 \rho_{\text{ad}(r_i)}(L(a_i, \beta_i)) = (N^2 - 1)\sigma(W_0) - \sigma_A(W_0),$$

where $\sigma_A(W_0)$ denotes the signature of the twisted cohomology group $H^2(W_0; \text{ad}(A))$ determined by the flat $PU(N)$ -connection $\text{ad}(A)$. This twisted cohomology group can be decomposed according to the irreducible decomposition of A (or equivalently α). The argument of [24, Lemma 2.6] shows that the contribution of the nontrivial summands is equal to 0. On the other hand, each trivial summand contributes -1 to the sum, because $\sigma(W_0) = -1$. □

The underlying $PU(N)$ -bundle of the connection A on W_0 can be lifted to a $U(N)$ -bundle E . There are also nonnegative integers k_1, \dots, k_N such that the restriction of

the connection

$$(3.40) \quad B_0^{k_1} \oplus \cdots \oplus B_0^{k_N}$$

to Y , the union of the three lens spaces, is equal to the restriction of A to Y . In particular, $L_0^{k_1} \oplus \cdots \oplus L_0^{k_N}$, the underlying $U(N)$ -bundle of (3.40), has the same restriction as E on Y . Therefore, the determinant of $L_0^{k_1} \oplus \cdots \oplus L_0^{k_N}$ is equal to $\det(E) \otimes L_0^{ak}$ for an appropriate integer k . After replacing k_1 with $k_1 + ak$, the new bundle $L_0^{k_1} \oplus \cdots \oplus L_0^{k_N}$ has the same determinant as E and the connection in (3.40) and A give rise to the same connection after restriction to Y .

Since A and the connection in (3.40) agree on boundary components except on $\Sigma(a_1, a_2, a_3)$, where A gives the connection α and B_0 restricts to the trivial connection, we can conclude that

$$(3.41) \quad \begin{aligned} \text{CS}(\alpha) &= \mathcal{E}(B_0^{k_1} \oplus \cdots \oplus B_0^{k_N}) - \mathcal{E}(A) \\ &= -\frac{1}{2N} \sum_{i < j} (k_i - k_j)^2 c_1(L)^2 \\ &= \frac{1}{2N} \sum_{i < j} \frac{(k_i - k_j)^2}{a}. \end{aligned}$$

Therefore, this gives us a strategy to compute the Chern–Simons functional of the flat connection α .

Now we focus on $N = 3$ and the Brieskorn homology sphere $\Sigma(2, 3, 23)$. Suppose α is an irreducible connection on $\Sigma(2, 3, 23)$. The irreducibility assumption implies that the holonomy of α along the generic fiber of the Seifert fibration is central. In fact, this central element is equal to the identity [4]. It is also shown in [4] that there are 44 such irreducible representations which are nondegenerate. One can use the method of [4] to find the conjugacy classes of holonomies corresponding to the elements x_1 , x_2 and x_3 of $\pi_1(\Sigma(2, 3, 23))$. Any irreducible flat connection on $\Sigma(2, 3, 23)$ is characterized by its holonomies along x_3 . The possible conjugacy classes for x_3 are listed in Table 4. On the other hand, the holonomies along x_1 and x_2 are conjugate to the diagonal matrices $\text{Diag}(1, -1, -1)$ and $\text{Diag}(1, \zeta, \zeta^2)$, where $\zeta = e^{2\pi i/3}$. Knowledge of these conjugacy classes allows us to apply Proposition 3.38 and (3.41) to compute the ρ -invariants, the Chern–Simons functional and hence the degrees of irreducible flat connections on $\Sigma(2, 3, 23)$:

Proposition 3.42 *There are ten irreducible flat connections of degree 0, five irreducible flat connections of degree 2, nine irreducible flat connections of degree 4, five irreducible flat connections of degree 6, nine irreducible flat connections of degree 8 and six irreducible flat connections of degree 10. There are not any irreducible flat connections of odd degree. The Chern–Simons functional and the ρ -invariants of these irreducible flat connections can be found in Table 4.*

Any reducible flat connection on $\Sigma(2, 3, 23)$ is either trivial or $SU(2)$ -reducible, because this 3-manifold is an integral homology sphere. In particular, nontrivial reducible $SU(3)$ -connections on $\Sigma(2, 3, 23)$ can be regarded as irreducible $SU(2)$ -connections. The results and the methods of [24] can be utilized to study such connections. The holonomy of any nontrivial flat $SU(2)$ -connection along the fiber of the Seifert fibration is the central element $-\text{id}$. The holonomies of this connection along x_1 and x_2 are respectively conjugate to $\text{Diag}(\mathbf{i}, -\mathbf{i})$ and $\text{Diag}(e^{\pi\mathbf{i}/3}, e^{-\pi\mathbf{i}/3})$. As in the $SU(3)$ -case, an irreducible flat $SU(2)$ -connection on $\Sigma(2, 3, 23)$ is determined by the conjugacy class of its holonomy along x_3 . The eigenvalues of the holonomy term along x_3 are equal to $e^{2\pi\mathbf{i}k/23}$ and $e^{-2\pi\mathbf{i}k/23}$ for $2 \leq k \leq 9$:

Proposition 3.43 [24] *There are eight irreducible $SU(2)$ -connections on $\Sigma(2, 3, 23)$. For each degree $2i + 1 \in \mathbb{Z}/8\mathbb{Z}$, there are exactly two such irreducible connections and there is no irreducible connection of even degree.*

Proposition 3.38 and (3.41) give a strategy to compute the Chern–Simons functional and the ρ -invariants of irreducible $SU(2)$ -connections. These computations can be used to verify the second part of the above proposition.

We also need to compute the degrees of flat $SU(2)$ -connections when they are regarded as $SU(3)$ -connections. To distinguish between these connections, we will write $\tilde{\alpha}$ for the $SU(3)$ -connection associated to an $SU(2)$ -connection α . The values of the Chern–Simons functional of α and $\tilde{\alpha}$ are equal to each other. However, the ρ -invariants of these two connections are different because ad_α and $\text{ad}_{\tilde{\alpha}}$ define two different representations of the fundamental group. We cannot use Proposition 3.38 to compute the ρ -invariant of $\tilde{\alpha}$ because this connection does not extend to the cobordism W_0 . In [6], cut-and-paste methods have been utilized to compute the difference $\rho_{\text{ad}_{\tilde{\alpha}}} - \rho_{\text{ad}_\alpha}$ for flat $SU(2)$ -connections on a family of homology spheres which includes $\Sigma(2, 3, 23)$. In particular, the following proposition can be extracted from [6]. The claim in it about the nondegeneracy of reducible flat connections on $\Sigma(2, 3, 23)$ is also proved in [6]. For more details about reducible flat $SU(3)$ -connections on $\Sigma(2, 3, 23)$ see Table 5.

Proposition 3.44 *The eight nontrivial reducible flat connections on $\Sigma(2, 3, 23)$ are nondegenerate. The $SU(3)$ –degrees of these connections are given as follows: there is one connection of degree 1, one connection of degree 3, one connection of degree 5, two connections of degree 7, two connections of degree 9 and one connection of degree 11.*

Define $I_*^3(\Sigma(2, 3, 23))$ to be the complex vector space generated by the irreducible flat connections on $\Sigma(2, 3, 23)$. The significance of this vector space for us is a gluing theorem for the 4–manifolds which split along a copy of $\Sigma(2, 3, 23)$. The proof of the following theorem will be given in Section 6.1. This theorem is analogous to Proposition 3.10 and implies that $I_*^3(\Sigma(2, 3, 23))$ plays a role similar to instanton Floer homology of admissible pairs. Motivated by this, we call $I_*^3(\Sigma(2, 3, 23))$ instanton Floer homology of $\Sigma(2, 3, 23)$.

Proposition 3.45 *Let X_1 and X_2 be two 4–manifolds such that $b^+(X_1), b^+(X_2) \geq 1$, $\partial X_1 = \Sigma(2, 3, 23)$ and $\partial X_2 = \overline{\Sigma(2, 3, 23)}$. Let w_i be a closed 2–cycle in X_i and $z_i \in \mathbb{A}(X_i)^{\otimes 2}$. Assume that*

$$\deg(z_1) \equiv -4w_1^2 - 4(\chi(x_1) + \sigma(X_1)) + 4 \pmod{12}.$$

Then there are

$$D_{X_1, w_1}^3(z_1) \in I_4^3(\Sigma(2, 3, 23)) \quad \text{and} \quad D_3^{X_2, w_2}(z_2): I_*^3(\Sigma(2, 3, 23)) \rightarrow \mathbb{C}$$

such that

$$D_3^{X_2, w_2}(z_2) \circ D_{X_1, w_1}^3(z_1) = D_{X_1 \#_{\Sigma(2, 3, 23)} X_2, w_1 \cup w_2}^3(z_1 \cdot z_2).$$

Moreover, $D_3^{X_2, w_2}(z_2)$ is nonzero only on the terms of degree

$$4w_2^2 + 4(\chi(X_2) + \sigma(X_2)) - 4 + \deg(z_2).$$

4 Computing polynomial invariants

In this section, which is the heart of this paper, we first compute the $U(3)$ –polynomial invariants of $E(n)$. The rank 2 invariants of elliptic surfaces were partially computed in [30; 32; 31] using algebrogeometric techniques. Complete calculations of the $U(2)$ –polynomial invariants of elliptic surfaces are given in [53; 25; 63]. In [25] and [63], *Gompf decomposition* of elliptic surfaces plays a key role in computing the invariants.

In the introduction, we also recalled a construction of elliptic surfaces which give rise to a decomposition of $E(n)$ into fiber sum of n copies of $E(1)$. This decomposition of elliptic surfaces can be also exploited to compute some of the $U(2)$ -polynomial invariants [69; 18].

Our method for computing the $U(3)$ -invariants of elliptic surfaces uses the Gompf decomposition, the fiber-sum decomposition and the rich group of the symmetries of elliptic surfaces. In Section 4.1, we first review the Gompf decomposition of elliptic surfaces. Then we show how symmetries of elliptic surfaces give constraints on the structure of their $U(3)$ -invariants. Among the elliptic surfaces $E(n)$, the K3 surface $E(2)$ is special because it has a larger symmetry group, which gives more restrictions on the invariants of this 4-manifold. In particular, most of the $U(3)$ -invariants of this 4-manifold are computed in Section 4.2 using identities (C_1) , (C_2) and (C_3) of Section 2.4. In the same subsection, we extend this computation to all $U(3)$ -invariants of $E(2)$ using the blowup formula, and then we explain what can be said about the invariants of the other surfaces $E(n)$ using the same techniques. In Section 4.3, we complete the computation of the invariants of $E(3)$ using the universal formulas of Proposition 2.21 about (-3) -spheres.

Finally, we consider the general case of elliptic surfaces $E(n)$ with $n \geq 4$. The key ingredient here is Proposition 3.45 about the Floer homology of $\Sigma(2, 3, 23)$. This together with Gompf decomposition allows us to compute all invariants of $E(n)$ by induction on n . The calculations of the invariants of $E(2)$ and $E(3)$ in the previous subsections serve as the basis of the induction. In the last subsection of this section, we also give a general gluing theorem for fiber sums. Similar results for $U(2)$ -invariants are proved in [71]. The proof of the rank 3 case follows the same strategy as in [71].

In the next two sections of the paper, we mainly focus on polynomial invariants and instanton Floer homology in the case $N = 3$. Therefore, we shall drop 3 from our notation I_*^3 , $D_{X,w}^3$, et cetera, when it does not create any confusion. Because we are working with an odd value of N , there is not any sign ambiguity in the definition of $I_*^3(W, w, z)$, $D_{X,w}^3(z)$, et cetera and we do not need to fix a homology orientation for the underlying 4-manifold.

4.1 Structure of the invariants of $E(n)$

A construction of the elliptic surface $E(n)$ was reviewed in the introduction. The simplest 4-manifold in this family, $E(1)$, can be also constructed by blowing up the

projective plane $\mathbb{C}P^2$ at the nine intersection points of two generic cubics. The pencil of cubics generated by the two cubics determines an elliptic fibration of $E(1)$. The nine exceptional divisors give rise to sections of this elliptic fibration, which are embedded spheres with self-intersection -1 . The manifold $E(n)$ is the fiber sum of n copies of $E(1)$ along the fibers of the elliptic fibration. The fibration of $E(1)$ induces an elliptic fibration for $E(n)$, and we will write f for a regular fiber of this fibration. By taking the connected sums of the exceptional sections of $E(1)$, we can form nine disjoint embedded spheres in $E(n)$. These are sections of the elliptic fibration of $E(n)$ and have self-intersection $-n$. We fix one of these sections and we will denote it by σ . When it does not cause any confusion, we will use the same notation to denote the homology and the cohomology classes associated to f and σ .

We can assume that there is a cusp fiber in the elliptic fibration of $E(1)$ by choosing appropriate cubics. This gives a cusp fiber f_0 in the fibration of $E(n)$. A regular neighborhood of $\sigma \cup f_0$ is a 4-manifold with boundary $\Sigma(2, 3, 6n - 1)$, which is called the *Gompf nucleus* and is denoted by $G(n)$ [39]. The intersection form of $G(n)$ is given by

$$\begin{bmatrix} 0 & 1 \\ 1 & -n \end{bmatrix}.$$

The complement of $G(n)$ in $E(n)$, denoted by $B(2, 3, 6n - 1)$, is a *Milnor fiber* and its intersection form is given by

$$(4.1) \quad n(-E_8) \oplus 2(n-1) \begin{bmatrix} 0 & 1 \\ 1 & -2 \end{bmatrix}.$$

Here each $-E_8$ summand has a basis of embedded spheres with self-intersection -2 which intersect each other according to $-E_8$. The i^{th} summand of the second type in (4.1) has a basis of an embedded torus g_i with self-intersection 0 and an embedded (-2) -sphere τ_i , where g_i and τ_i intersect each other positively at one point [40].

The 4-manifold $E(2)$, which is a K3 surface, plays a special role in this family. For example, $E(2)$ enjoys a rich group of symmetries. As a manifestation of this fact, we have the following proposition:

Lemma 4.2 *Suppose that e and e' are two nonzero elements of $H^2(E(2), \mathbb{Z})$ with $e \cup e \equiv e' \cup e' \pmod{3}$. Then there is an orientation-preserving diffeomorphism Φ of $E(2)$ such that $\Phi^*(e) \equiv e' \pmod{3}$. In particular, the action of $\text{Diff}(E(2))$ on $H^2(E(2), \mathbb{Z}/3\mathbb{Z})$ has four orbits.*

Proof Let $e_1, e_2 \in H^2(E(2), \mathbb{Z})$ be chosen such that $e_i \cup e_j$ is zero when $i = j$, and is equal to 1 when $i \neq j$. Because the action of $\text{Diff}(E(2))$ on the primitive cohomology classes of a fixed self-intersection is transitive, there is an element Ψ of $\text{Diff}(E(2))$ such that $\Psi^*(e) = m(e_1 + n e_2)$ for $m, n \in \mathbb{Z}$. Therefore $\Psi^*(e) \bmod 3$ is equal to one of the following elements:

$$(4.3) \quad 0, \quad e_1, \quad e_1 - e_2, \quad e_1 + e_2.$$

The nonzero classes in (4.3) can be distinguished from each other by their self-intersection. Thus, there is a $\Psi' \in \text{Diff}(E(2))$ such that $\Psi^*(e) \equiv \Psi'^*(e') \bmod 3$. \square

For $1 \leq i \leq 3$, suppose w_i is the 2-cycle in $E(2)$ given by $\sigma - (i - 1)f$. Then $w_i^2 \equiv i \bmod 3$. Therefore, these 2-cycles and the empty cycle give representatives for the orbits of the action of $\text{Diff}(E(2))$ on $H^2(E(2), \mathbb{Z}/3\mathbb{Z})$. Alternatively, let w'_i be the union of i elements of the nine disjoint spheres of self-intersection -2 in $E(2)$ that were constructed above. If i is a positive integer and $i \equiv j \bmod 3$ with $1 \leq j \leq 3$, then define $w_i := w_j$ and $w'_i := w'_j$. We also define w_0 and w'_0 to be the empty cycles.

The group of diffeomorphisms of $E(n)$ for $n \geq 3$ is more constrained than that of $E(2)$. For example, any diffeomorphism of $E(n)$ maps the homology class f to $\pm f$. Therefore, we cannot expect that the analogue of Lemma 4.2 holds for an arbitrary n . However, $E(n)$ still has a big diffeomorphism group and we can prove the following weakened version of Lemma 4.2:

Lemma 4.4 *Suppose that the integers $n \geq 3$ and $1 \leq i \leq 3$ are given and that $u \in H^2(B(2, 3, 6n - 1), \mathbb{Z})$ satisfying $u \cup u \equiv i \bmod 3$ is fixed. Suppose also that $e \in H^2(E(n), \mathbb{Z})$ is such that $e \cup f \equiv 0 \bmod 3$ and $e \cup e \equiv i \bmod 3$. Then there is an element Φ of $\text{Diff}(E(n))$ such that*

$$(4.5) \quad \Phi^*(e) \equiv kf + u \bmod 3 \quad \text{or} \quad \Phi^*(e) \equiv kf \bmod 3,$$

where $k = 0$ or 1 . Moreover, the map on $H^2(G(n), \mathbb{Z})$ induced by Φ is $\pm \text{id}$. In particular, the action of $\text{Diff}(E(n))$ on $\langle f \rangle^\perp$ in $H^2(E(n), \mathbb{Z}/3\mathbb{Z})$ has eight orbits.

In the statement of Lemma 4.4 we regard u as an element of $H^2(E(n), \mathbb{Z})$ using the inclusion of $B(2, 3, 6n - 1)$ in $E(n)$. Note also that the second case in (4.5) holds only if $i = 3$.

Proof The element e can be written as the sum

$$rf + s\sigma + v,$$

where $v \in H^2(B(2, 3, 6n - 1), \mathbb{Z}) \subset H^2(E(n), \mathbb{Z})$. Because $e \cup f \equiv 0 \pmod 3$, we know s is divisible by 3. There is also a diffeomorphism of $E(n)$ that maps f to $-f$ and σ to $-\sigma$ [39, Lemma 3.7]. After applying this diffeomorphism if it is necessary, we can assume $e \equiv kf + v \pmod 3$ with $k = 0$ or 1. Suppose $\text{SO}(H^2(E(n), \mathbb{Z}))$ denotes an element of the special orthogonal group of the lattice $H^2(E(n), \mathbb{Z})$ with respect to the intersection bilinear form. According to [41, Proposition 3.3], there is a spinor norm one element of $\text{SO}(H^2(E(n), \mathbb{Z}))$ that fixes f and σ and maps v to an element of the form $m(\tau_1 + ng_1)$. This element can be realized by a diffeomorphism of $E(n)$ [32, Section 6.2]. This can be used to verify the claim as in Lemma 4.2. \square

The eight orbits in Lemma 4.4 can be represented by

$$w_{k,0} = kf \quad \text{and} \quad w_{k,l} = kf + \tau_1 - (l - 1)g_1$$

for $k = 0, 1$ and $l = 1, 2, 3$. Alternatively, we can use $w'_{k,l} = kf + \tau_1 + \dots + \tau_l$. If l is a positive integer and $l \equiv j \pmod 3$ with $1 \leq j \leq 3$, then let $w_{k,l} = w_{k,j}$ and $w'_{k,l} = w'_{k,j}$.

Consider the $\text{U}(3)$ -polynomial invariant $D^3_{E(n),w}(\Gamma^i_{(2)}\Lambda^j_{(3)})$ for the homology classes Γ and Λ . This polynomial is invariant with respect to the action of the diffeomorphism group of $E(n)$ on (w, Γ, Λ) . Therefore, we can use Lemmas 4.2 and 4.4 to focus on a smaller subset of possible values for w . Since changing $w \pmod 3$ would not change the polynomial invariant and $H^2(E(n), \mathbb{Z}/3\mathbb{Z})$ is finite, the polynomial $D^3_{E(n),w}(\Gamma^i_{(2)}\Lambda^j_{(3)})$ is invariant with respect to the action of a finite-index subgroup of $\text{Diff}(E(n))$ on (Γ, Λ) . This action of the diffeomorphism group factors through the action of the algebraic group $\text{O}(H_2(E(n)))$. Here the orthogonal group is defined using the intersection form on the complex vector space $H_2(E(n))$. Suppose also that $\text{SO}(H_2(E(n)); f)$ is the subgroup of $\text{O}(H_2(E(n)))$ consisting of the orthogonal transformations that map $f \in H_2(E(n))$ to itself and have determinant 1. As another manifestation of the big diffeomorphism group of $E(n)$, it is shown in [32, Section 6.2] that the image of any finite-index subgroup of $\text{Diff}(E(n))$ in $\text{O}(H_2(E(n)))$ contains an algebraically dense subgroup of $\text{SO}(H_2(E(n)); f)$. Therefore, the polynomial $D^3_{E(n),w}(\Gamma^i_{(2)}\Lambda^j_{(3)})$ is invariant with respect to the action of $\text{SO}(H_2(E(n)); f)$ on (Γ, Λ) . In the case that $n = 2$, one can even replace $\text{SO}(H_2(E(n)); f)$ with $\text{SO}(H_2(E(n)))$.

Lemma 4.6 Suppose (V, Q) is a pair of a complex vector space of dimension greater than 2 and a quadratic form. Suppose also τ_1 and τ_2 are two vectors orthogonal to each other such that $Q(\tau_1)$ and $Q(\tau_2)$ are nonzero. Suppose $P: V \oplus V \rightarrow \mathbb{C}$ is a bihomogeneous polynomial of bidegree (d_1, d_2) that is invariant with respect to the diagonal action of $SO(V)$ on $V \oplus V$. Then P is determined by its value on $W_0 \oplus W_1$, where W_0 is the span of τ_1 , and W_1 is the span of τ_1 and τ_2 . Moreover, P has the form

$$P(x, y) = \sum_{\substack{i,j,k \geq 0 \\ 2i+j=d_1 \\ j+2k=d_2}} c_{i,j,k} Q(x)^i Q(x, y)^j Q(y)^k$$

for appropriate constants $c_{i,j,k} \in \mathbb{C}$.

In our application, V will be $H_2(E(2))$, and τ_1 and τ_2 will be two disjoint embedded spheres in $E(2)$.

Proof Suppose $(\alpha, \beta) \in V \oplus V$ is such that $Q(\alpha) \neq 0$. Then the vector β can be written as the sum $\beta_0 + \beta_1$ where β_0 is a multiple of α , and β_1 is orthogonal to α . We also assume that $Q(\beta_1) \neq 0$. It is straightforward to find an element of $SO(V)$ which maps (α, β) to an element of the form $W_0 \oplus W_1$. The set of vectors (α, β) as above is also dense in $V \oplus V$. Therefore, P is determined by its values on $W_0 \oplus W_1$.

By evaluating P on elements of $W_0 \oplus W_1$, we have

$$P(\lambda\tau_1, \mu_1\tau_1 + \mu_2\tau_2) = \sum m_{a,b,c} \lambda^a \mu_1^b \mu_2^c.$$

There is an element of $SO(V)$ which maps τ_1 to itself (respectively, $-\tau_1$) and τ_2 to $-\tau_2$ (respectively, itself). Therefore, $m_{a,b,c}$ is nonzero only if $a + b$ and c are even. This implies that there are constants $c_{i,j,k}$ such that

$$P(x, y) = \sum_{\substack{2i+j=d_1 \\ j+2k=d_2}} c_{i,j,k} Q(x)^i Q(x, y)^j Q(y)^k$$

for $(x, y) \in W_0 \oplus W_1$, where j and k are nonnegative integers. This implies the second part of the lemma because each side of the above equality is invariant with respect to the action of $SO(V)$. Arguing as in [32, Chapter 6, Lemma 2.2], we can also assume that i only takes nonnegative integers. □

Lemma 4.7 Suppose that (V, Q) is a pair of a complex vector space of dimension greater than 3 and a quadratic form. Fix a vector $f \in V$ with $Q(f) = 0$, and

let the vectors k and τ be chosen such that $Q(k, f)$ and $Q(\tau)$ are nonzero, and $Q(\tau, k) = Q(\tau, f) = 0$. Suppose also $P: V \oplus V \rightarrow \mathbb{C}$ is a polynomial that is invariant with respect to the diagonal action of $SO(V; f)$. Then $P(x, y)$ is determined by its values on $W_0 \oplus W_1$, where W_0 is the span of the vectors f and k , and W_1 is the span of f, k and τ .

In our application, V, f, k and τ will be $H_2(E(n)), f, \sigma$ and an embedded sphere in $B(2, 3, 6n - 1)$, respectively.

Proof The proof is similar to that of the first part of Lemma 4.6. For a given $(\alpha, \beta) \in V \oplus V$, assume $Q(\alpha, f) \neq 0$. Then the vector β can be uniquely written as $\beta_0 + \beta_1$ where $\beta_0 \in \text{Span}(f, \alpha)$, and β_1 is orthogonal to $\text{Span}(f, \alpha)$. As another assumption, we require that $Q(\beta_1) \neq 0$. These assumptions hold for a dense subset of $V \oplus V$. It can be easily checked that there is an element of $SO(V; k)$ which maps (α, β) to $W_0 \oplus W_1$. Therefore, $P|_{W_0 \oplus W_1}$ determines P on $V \oplus V$. □

4.2 Invariants of $E(2)$

In this section, we study the $U(3)$ -polynomial invariants of $E(2)$ and a 2-cycle w . Lemma 4.4 shows that we can assume that the 2-cycle w is either empty or w_i for some $1 \leq i \leq 3$. Equivalently, we can replace w_i with w'_i . Throughout this subsection, we will follow the same notation as in the previous part to denote the surfaces σ, τ_i and g_i embedded in $E(2)$.

Proposition 4.8 *The invariant $D_{E(2), w_i}$ satisfies the identities*

$$(4.9) \quad D_{E(2), w_i} \left(\left(\frac{1}{3} a_2 \right)^j z \right) = D_{E(2), w_{i+j}}(z) \quad \text{and} \quad D_{E(2), w_i}(a_3 z) = 0$$

for $z \in \mathbb{A}(E(2))^{\otimes 2}$. In particular, for $1 \leq i \leq 3$, $E(2)$ has w_i -simple type and $\hat{D}_{E(2), w_i}(e^{\Gamma(2) + \Lambda(3)})$ is independent of the choice of i .

Proof Suppose $N(f)$ is a neighborhood of a regular fiber and X is the complement of $N(f)$. We also identify the boundary of X with $Y_1 = S^1 \times f$. The 4-manifold X contains a copy of $B(2, 3, 11)$. Therefore, we can find two disjoint embedded spheres τ_1 and τ_2 of self-intersection -2 in X . Let S be the subspace of $H_2(X)$ spanned by the vectors τ_1 and τ_2 , and $\mathbb{A}(S)$ be the subalgebra $\text{Sym}^*(H_0(X) \oplus S)$ of $\mathbb{A}(X)$. Then w_i can be decomposed as $w \# w'$, where w (respectively, w') is a 2-cycle in X (respectively, $N(f)$), and w and w' intersect Y_1 in $\gamma := S^1 \times \{\text{pt}\}$. For $z \in \mathbb{A}(S)^{\otimes 2}$,

Proposition 3.24, with the aid of the functoriality properties discussed in [Section 3.1](#), implies that

$$\begin{aligned} D_{E(2),w_i}(a_2^j a_3^k z) &= D^{N(f),w'}(1) \circ I_*(Y_1 \times [0, 1], \gamma \times [0, 1], a_2^j a_3^k) \circ D_{X,w}(z) \\ &= 3^j 0^k D^{N(f),w'}(1) \circ I_*(Y_1 \times [0, 1], \gamma \times [0, 1] - jf, 1) \circ D_{X,w}(z) \\ &= 3^j 0^k D_{E(2),w_i-jf}(z). \end{aligned}$$

This verifies [\(4.9\)](#) for the case $z \in \mathbb{A}(S)^{\otimes 2}$. Using [Lemma 4.6](#), the same claim holds for general z , and as a result $\widehat{D}_{E(2),w_i}(e^{\Gamma(2)+\Lambda(3)})$ is equal to

$$D_{E(2),w_1}(e^{\Gamma(2)+\Lambda(3)}) + D_{E(2),w_2}(e^{\Gamma(2)+\Lambda(3)}) + D_{E(2),w_3}(e^{\Gamma(2)+\Lambda(3)}).$$

In particular, $\widehat{D}_{E(2),w_i}(e^{\Gamma(2)+\Lambda(3)})$ does not depend on i . □

A similar argument, using [Lemma 4.7](#) instead of [Lemma 4.6](#), proves the following analogous statement for $E(n)$:

Proposition 4.10 *The polynomial invariants $D_{E(n),w_{k,l}}$ for $1 \leq l \leq 3$ satisfy the identities*

$$(4.11) \quad D_{E(n),w_{k,l}}\left(\left(\frac{1}{3}a_2\right)^j z\right) = D_{E(2),w_{k,l+j}}(z) \quad \text{and} \quad D_{E(n),w_{k,l}}(a_3 z) = 0$$

for $z \in \mathbb{A}(E(n))^{\otimes 2}$. In particular, for $1 \leq l \leq 3$, $E(n)$ has $w_{k,l}$ -simple type and $\widehat{D}_{E(n),w_{k,l}}(e^{\Gamma(2)+\Lambda(3)})$ is independent of the choice of l .

Proposition 4.12 *For $1 \leq i \leq 3$, we have*

$$(4.13) \quad \widehat{D}_{E(2),w_i}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}.$$

The two sides of [\(4.13\)](#) are power series in t_2 and t_3 where the coefficients of $t_2^i t_3^j$ are bihomogeneous polynomials on $H_2(E(2)) \oplus H_2(E(2))$ of degree (i, j) . Identity [\(4.18\)](#) means for each choice of (i, j) these coefficients are equal to each other.

Proof Using [Lemma 4.6](#), we can find a power series $g(r, s, t) \in \mathbb{C}[[r, s, t]]$ such that

$$\widehat{D}_{E(2),w_i}(e^{\Gamma(2)+\Lambda(3)}) \cdot e^{-\frac{1}{2}Q(\Gamma)+Q(\Lambda)} = g(Q(\Gamma), Q(\Gamma, \Lambda), Q(\Lambda)).$$

The constant term of g is equal to 1 [\[52\]](#). [Lemma 4.6](#) also implies that it suffices to show that $g(Q(\Gamma), Q(\Gamma, \Lambda), Q(\Lambda))$ is equal to 1 for homology classes Γ and Λ and

2–cycles w_i which are orthogonal to an embedded (-2) –sphere τ . For such choices of Γ and Λ , we have

$$\widehat{D}_{E(2),w_i}(e^{\Gamma(2)+\epsilon\tau(2)+\Lambda(3)}) \cdot e^{-\frac{1}{2}\mathcal{Q}(\Gamma)+\epsilon^2+\mathcal{Q}(\Lambda)} = g(\mathcal{Q}(\Gamma) - 2\epsilon^2, \mathcal{Q}(\Gamma, \Lambda), \mathcal{Q}(\Lambda)).$$

By taking the second derivatives of the above expression with respect to ϵ , evaluating ϵ at 0 and using (C_1) of Section 2.4, we obtain

$$\frac{\partial g}{\partial r}(r, s, t) = 0.$$

Similarly, identities (C_2) and (C_3) imply

$$4 \frac{\partial^2 g}{\partial r^2}(r, s, t) - \frac{\partial g}{\partial t}(r, s, t) = 0 \quad \text{and} \quad 2 \frac{\partial^2 g}{\partial s \partial r}(r, s, t) + \frac{\partial g}{\partial s}(r, s, t) = 0.$$

Therefore, g is equal to the constant power series 1. □

Suppose X is the blowup of $E(2)$ and w is the 2–cycle f in X . If E is the exceptional sphere in X , then Theorem 2.27 can be used to compute the invariants of (X, w) :

$$\widehat{D}_{X,w}(e^{\Gamma(2)+\Lambda(3)}) = \frac{1}{3}e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}(\cosh(\sqrt{3}E \cdot \Gamma) + 2 \cos(\sqrt{3}E \cdot \Lambda)).$$

The homology class $\sigma + E$ can be represented by an embedded (-3) –sphere σ' in X . Fix $\Gamma, \Lambda \in H_2(X)$ which are orthogonal to σ' . Then the above formula can be used to show

$$(4.14) \quad \widehat{D}_{X,w}\left(\left(-\frac{3}{2}\sigma'_{(3)} - \frac{3}{2}\sigma'_{(2)} - a_2\right)e^{\Gamma(2)+\Lambda(3)}\right) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}(-\sqrt{3} \sin(\sqrt{3}E \cdot \Lambda) + \cos(\sqrt{3}E \cdot \Lambda) - \cosh(\sqrt{3}E \cdot \Gamma)).$$

By another application of Theorem 2.27 and Remark 2.32, $\widehat{D}_{X,w-\sigma'}(e^{\Gamma(2)+\Lambda(3)})$ is equal to

$$(4.15) \quad \frac{1}{3}e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}(\cosh(\sqrt{3}E \cdot \Gamma) - \cos(\sqrt{3}E \cdot \Lambda) + \sqrt{3} \sin(\sqrt{3}E \cdot \Lambda)).$$

Using Proposition 2.21 and comparing these two identities, we can find the undetermined constant c in Proposition 2.21:

Proposition 4.16 *The constant c is equal to -3 .*

Now we are ready to complete the computation of the invariants of $E(2)$:

Theorem 4.17 *Suppose w is a 2–cycle in $E(2)$. Then $E(2)$ has w –simple type and the $U(3)$ –series of $E(2)$ is given by the formula*

$$(4.18) \quad \widehat{D}_{E(2),w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}.$$

Proof In light of Lemma 4.2 and Proposition 4.12, it suffices to consider only the empty 2-cycle w_0 . Let σ' be the embedded (-3) -sphere in $E(2) \# \overline{\mathbb{C}\mathbb{P}^2}$ constructed above. Consider the 2-cycle $w' := \sigma$ of $E(2) \# \overline{\mathbb{C}\mathbb{P}^2}$ and the element $z := (\sigma_{(2)} - 2E_{(2)})^2 z'$ of $\mathbb{A}(E(2) \# \overline{\mathbb{C}\mathbb{P}^2})^{\otimes 2}$, where $z' \in \mathbb{A}(\langle \sigma \rangle^\perp)^{\otimes 2} \cap \mathbb{A}(E(2))^{\otimes 2}$. By Proposition 2.22, the following invariant of $E(2) \# \overline{\mathbb{C}\mathbb{P}^2}$ is equal to $4D_{E(2),w_0}(z')$:

$$(4.19) \quad D_{E(2) \# \overline{\mathbb{C}\mathbb{P}^2}, w' - \sigma'}((\sigma_{(2)} - 2E_{(2)})^2 z').$$

Moreover, the first identity of Proposition 2.21 can be used to show (4.19) is equal to $D_{E(2) \# \overline{\mathbb{C}\mathbb{P}^2}, w'}(z'')$ for an appropriate choice of $z'' \in \mathbb{A}(E(2) \# \overline{\mathbb{C}\mathbb{P}^2})^{\otimes 2}$. Replacing w' with $w'' := \sigma + \tau_1 + g_1$ shows that

$$4D_{E(2), \tau_1 + g_1}(z') = D_{E(2) \# \overline{\mathbb{C}\mathbb{P}^2}, w''}(z'').$$

Since $w' \cdot w' \equiv w'' \cdot w'' \pmod 3$, the left-hand side of the above identity is equal to $D_{E(2) \# \overline{\mathbb{C}\mathbb{P}^2}, w'}(z'')$ by the blowup formula. Therefore, we can deduce that

$$(4.20) \quad D_{E(2), w_0}(z') = D_{E(2), \tau_1 + g_1}(z')$$

for $z' \in \mathbb{A}(\langle \sigma \rangle^\perp)^{\otimes 2}$. As a consequence of Lemma 4.6, Identity (4.20) holds for any choice of z' . In particular, $E(2)$ has simple type with respect to w_0 and (4.18) holds for this choice of w . □

Proposition 4.21 *Suppose that $\Gamma, \Lambda \in H_2(B(2, 3, 6n - 1)) \subset H_2(E(n))$ and that $\Gamma', \Lambda' \in H_2(G(n)) \subset H_2(E(n))$. Then*

$$(4.22) \quad \widehat{D}_{E(n), w_{k,l}}(e^{(\Gamma + \Gamma')_{(2)} + (\Lambda + \Lambda')_{(3)}}) = e^{\frac{1}{2}Q(\Gamma) - Q(\Lambda)} \widehat{D}_{E(n), w_{k,3}}(e^{(\Gamma')_{(2)} + (\Lambda')_{(3)}}).$$

Proof The group of orthogonal transformations $SO(H_2(B(2, 3, 6n - 1)), Q)$, regarded as a subgroup of $SO(H_2(E(n)))$, acts as the identity on the series $\widehat{D}_{E(n), w}(e^{\Gamma_{(2)} + \Lambda_{(3)}})$. This fact can be combined with the argument of Proposition 4.12 to verify (4.22) for $1 \leq l \leq 3$. To finish the proof, it suffices to show that $\widehat{D}_{E(n), w_{k,0}}(e^{\Gamma_{(2)} + \Lambda_{(3)}})$ is equal to $\widehat{D}_{E(n), w_{k,3}}(e^{\Gamma_{(2)} + \Lambda_{(3)}})$. This also can be achieved with the method of the proof of Theorem 4.17. □

By Proposition 4.10, we already know that $E(n)$ has $w_{k,l}$ -simple type for $1 \leq l \leq 3$. The above proof can be used to show that $E(n)$ has $w_{k,0}$ -simple type, too.

4.3 Invariants of $E(3)$

In this section, we study the $U(3)$ -polynomial invariants of $E(3)$ up to a constant and a sign ambiguity. The following is [Theorem 2](#) from the introduction:

Theorem 4.23 *The 4-manifold $E(3)$ has simple type. There are also real numbers \hbar_1 and \hbar_2 such that for any 2-cycle w in $E(3)$ and $\Gamma, \Lambda \in H_2(E(3))$, the series $\widehat{D}_{E(3),w}(e^{\Gamma(2)+\Lambda(3)})$ is equal to*

$$e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}(\hbar_1 \cosh(\sqrt{3}f \cdot \Gamma) - 2\hbar_2 \cos(-\frac{2}{3}\pi w \cdot f + \sqrt{3}f \cdot \Lambda)).$$

Furthermore, $\hbar_1 + \hbar_2 = \pm 1$ for an appropriate choice of the sign.

In a work in progress, we show that $\hbar_1 = \frac{2}{3}$ and $\hbar_2 = \frac{1}{3}$. However, we do not need the exact values of these constants in this paper. Later, we only use the fact that \hbar_1 and \hbar_2 are nonzero. To abbreviate our notation, define

$$(4.24) \quad G(\Gamma, \Lambda, j) := \hbar_1 \cosh(\sqrt{3}f \cdot \Gamma) - 2\hbar_2 \cos(-\frac{2}{3}\pi j + \sqrt{3}f \cdot \Lambda).$$

Proposition 4.25 *Suppose w is a 2-cycle in $E(3)$ with $w \cdot f \not\equiv 0 \pmod{3}$. Then*

$$(4.26) \quad D_{E(3),w}((\frac{1}{3}a_2)^j z) = D_{E(3),w-jf}(z) \quad \text{and} \quad D_{E(3),w}(a_3 z) = 0$$

for $z \in \mathbb{A}(E(3))^{\otimes 2}$. In particular, $E(3)$ has w -simple type. Furthermore, there is a power series $g \in \mathbb{Q}[[t_2, t_3]]$ such that for $\Gamma, \Lambda \in H_2(E(3))$,

$$\widehat{D}_{E(3),w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)} g(\Gamma \cdot f, \Lambda \cdot f)$$

when $w \cdot f \equiv 1 \pmod{3}$, and

$$\widehat{D}_{E(3),w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)} g(-\Gamma \cdot f, -\Lambda \cdot f)$$

when $w \cdot f \equiv 2 \pmod{3}$. Furthermore, g is even with respect to the variable t_2 and its constant term is equal to ± 1 .

Proof The 4-manifold $E(3)$ is given as the fiber sum $E(2) \#_f E(1)$. In this proof, let σ_n denote a section of the elliptic fibration of $E(n)$, which is a sphere of self-intersection $-n$. We can assume that $\sigma_3 = \sigma_2 \# \sigma_1$. Firstly, consider the case $w \cdot f \equiv 1 \pmod{3}$. Arguing as in [Lemma 4.4](#), we can assume that $w = w_1 \# \sigma_1$, where w_1 is a 2-cycle in $E(2)$ with $w_1 \cdot f = 1$. Suppose Γ_0 and Λ_0 are two elements of $H_2(E(2))$

such that $\Gamma_0 \cdot f = \Lambda_0 \cdot f = 1$. Then [Proposition 3.35](#) for $X_1 = E(2)$, $X_2 = E(1)$ and $X_3 = X_4 = S^2 \times f$ implies that

$$p(t_2, t_3) \sum_{0 \leq j \leq 2} D_{E(2)\#_f E(1), w+jf} (e^{(\Gamma_0\#\sigma_1)(2)+(\Lambda_0\#\sigma_1)(3)}) \\ = q(t_2, t_3) \sum_{0 \leq j \leq 2} D_{E(2)\#_f S^2 \times f, w_2+jf} (e^{(\Gamma_0)(2)+(\Lambda_0)(3)}),$$

where

$$p(t_2, t_3) = \sum_{0 \leq j \leq 2} D_{S^2 \times f, S^2 \times \{pt\}+jf} (e^{\Delta(2)+\Delta(3)})$$

and

$$q(t_2, t_3) = \sum_{0 \leq j \leq 2} D_{E(1), w_1+jf} (e^{(\sigma_1)(2)+(\sigma_1)(3)}).$$

Note that $b^+(S^2 \times f) = b^+(E(1)) = 1$ and we use the invariants with respect to the metrics that have long necks along f in the above identities. The power series $p(t_2, t_3)$ is invertible, because $p(0, 0) = 1$. Therefore, we can conclude there is a power series $g(t_2, t_3)$ such that

$$(4.27) \quad \sum_{0 \leq j \leq 2} D_{E(3), w+jf} (e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)} g(t_2, t_3)$$

with $\Gamma = \Gamma_0\#\sigma_1$ and $\Lambda = \Lambda_0\#\sigma_1$ for Γ_0 and Λ_0 given as above. Then [Lemma 4.7](#) implies that (4.27) holds for arbitrary Γ and Λ with $\Gamma \cdot f = \Lambda \cdot f = 1$. By [Proposition 4.8](#), we can use a similar argument as above to show

$$\sum_{0 \leq j \leq 2} D_{E(3), w+jf} (P(a_2, a_3)e^{\Gamma(2)+\Lambda(3)}) = P(3, 0)e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)} g(t_2, t_3).$$

In particular, this shows that

$$D_{E(3), w} \left(\left(\frac{1}{3}a_2\right)^j e^{s\Gamma(2)+t\Lambda(3)} \right) = D_{E(3), w-jf} (e^{\Gamma(2)+\Lambda(3)}), \\ D_{E(3), w} (a_3 e^{s\Gamma(2)+t\Lambda(3)}) = 0.$$

The power series g is even with respect to t_2 , because $D(e^{\Gamma(2)+\Lambda(3)})$ is even with respect to t_2 . A similar application of [Proposition 3.35](#) for $X_1 = X_2 = E(1)$ and $X_3 = X_4 = S^2 \times f$ shows that

$$e^{-t_2^2+2t_3^2} p(t_2, t_3) = q(t_2, t_3)^2.$$

The constant term of the above equality and the identity $p(0, 0) = 1$ shows that the constant term of g is equal to ± 1 . This fact completes the proof for the case

that $w \cdot f \equiv 1 \pmod 3$. Using a diffeomorphism of $E(3)$ which maps f to $-f$, we can also treat the case that $w \cdot f \equiv 2 \pmod 3$. \square

To determine the power series g , let σ and σ' be two disjoint sections of the elliptic fibration of $E(3)$. Let also w be chosen such that $w \cdot f = 1$ and $w \cdot \sigma = 2$. Then

$$\begin{aligned} \widehat{D}_{E(3),w}(e^{(s\sigma+s'\sigma')_{(2)}+(r\sigma+r'\sigma')_{(3)}}) \\ = e^{-3t_2^2(s^2+s'^2)/2+3t_3^2(r^2+r'^2)}g((s+s')t_2, (r+r')t_3). \end{aligned}$$

By differentiating with respect to s and r , we can conclude that

$$\widehat{D}_{E(3),w}(\sigma_{(2)}^i \sigma_{(3)}^j z) = h \frac{d^i}{ds^i} \frac{d^j}{dr^j} \Big|_{s=t=0} e^{-3t_2^2 s^2/2+3t_3^2 r^2} g((s+s')t_2, (r+r')t_3),$$

where $z = e^{s'\sigma'_{(2)}+r'\sigma'_{(3)}}$ and $h = e^{-3t_2^2 s'^2/2+3t_3^2 r'^2}$. By applying these identities to the second formula in Proposition 2.21, we can conclude

$$(4.28) \quad -\frac{1}{2}g_3 + \frac{1}{2}g_{22} - \frac{1}{2}g = g \circ \tau,$$

where g_{22} means the second derivative of the power series $g(t_2, t_3)$ with respect to t_2 , and so on. Moreover, τ maps (t_2, t_3) to $(-t_2, -t_3)$. We can use (4.28) to derive the identity

$$(4.29) \quad \frac{1}{2}(g \circ \tau)_3 + \frac{1}{2}(g \circ \tau)_{22} - \frac{1}{2}g \circ \tau = g.$$

Replacing $g \circ \tau$ in (4.29) with the left-hand side of (4.28) gives rise to the following PDE for g :

$$(4.30) \quad g_{2222} - g_{33} - 2g_{22} - 3g = 0.$$

Next, let w' be a 2-cycle with $w' \cdot f = 1$ and $w' \cdot \sigma = 0$ and consider

$$\widehat{D}_{E(3),w'}(e^{(s\sigma+s'\sigma')_{(2)}+(r\sigma+r'\sigma')_{(3)}})$$

instead. With the same argument, the last part of Proposition 2.21 implies that

$$(4.31) \quad g_{2222} - 6g_{22} + 3g_{33} + 9g = 0 \quad \text{and} \quad g_{2223} - 6g_{23} = 0.$$

We can combine (4.30) and the first equation in (4.31) to come up with the following simpler PDE:

$$(4.32) \quad g_{33} - g_{22} + 3g = 0.$$

The second PDE in (4.31) and the fact that g is even with respect to the variable t_2 imply that $g_{st} = p(t) \cosh(\sqrt{6}s)$. The equations (4.30) and (4.32) can be used to write two linear ordinary differential equations for p . It is straightforward to check that the only solution of these ODEs is $p(t) = 0$. Therefore, the power series g has the form $q_1(t) + q_2(s)$. Equation (4.32) can be used to find differential equations for q_1 and q_2 . By solving these ODEs and using the fact that g is even with respect to t_2 , we can conclude

$$(4.33) \quad g(t_2, t_3) = a \cosh(\sqrt{3}t_2) + b \cos(\sqrt{3}t_3) + c \sin(\sqrt{3}t_3).$$

If $g(0, 0) = 1$, then the initial value and (4.28) imply the following constraints on a , b and c , which can be used to prove Theorem 4.23 in the case $w \cdot f \not\equiv 0 \pmod 3$:

$$a + b = 1 \quad \text{and} \quad a - \frac{1}{2}b - \frac{\sqrt{3}}{2}c = 1.$$

A similar argument can be used in the case that $g(0, 0) = -1$.

Next, let $w \cdot f \equiv 0 \pmod 3$. Using Lemma 4.4 and Proposition 4.10, it suffices to consider the case that $w = w_{k,1}$ for $k = 0$ or 1 . The following proposition computes the invariants of $E(3)$ for this choice of 2-cycle. In its proof, we use the basis for the homology of $H_2(E(3), \mathbb{Z})$ which is introduced in Section 4.1:

Proposition 4.34 For any $(\Gamma, \Lambda) \in H_2(E(3)) \oplus H_2(E(3))$,

$$(4.35) \quad \widehat{D}_{E(3), w_{k,l}}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}G(\Gamma, \Lambda, 0).$$

Proof Let σ' be a section of the elliptic fibration of $E(3)$ which is disjoint from σ . Then the homology class of σ' is equal to $\sigma + 3f + u$, where $u \in H_2(B(2, 3, 17), \mathbb{Z})$ and $u \cdot u = -6$. Arguing as in Lemma 4.4, there is a diffeomorphism Φ of $E(3)$ that fixes $H_2(G(3))$ and maps u to a linear combination of g_1 and τ_1 . Furthermore, $\Phi_*(u) \equiv 2g_1 - \tau_1 \pmod 3$. In particular, $\alpha := \Phi(\sigma')$ is a (-3) -sphere with $\alpha \cdot w_{k,1} \equiv 1 \pmod 3$. Suppose W is the subspace of the elements of $H_2(E(3))$ whose intersection numbers with α are equal to 0. Using Proposition 2.21, the series $\widehat{D}_{E(3), w_{k,l}}(e^{\Gamma(2)+\Lambda(3)})$, for any $(\Gamma, \Lambda) \in W \oplus W$, is equal to

$$(4.36) \quad -\frac{1}{3}\widehat{D}_{E(3), w_{k,1}+\alpha}\left(\left(-\frac{3}{2}\alpha(3) - \frac{3}{2}\alpha(2) - a_2\right)e^{\Gamma(2)+\Lambda(3)}\right).$$

Since $(w_{k,1} + \alpha) \cdot f = 1$, we can evaluate (4.36) using our current knowledge of the invariants of $E(3)$. It is straightforward to check that the resulting series is equal to that in (4.35).

The homology classes f , $k := \sigma + \frac{3}{2}f$ and τ_2 satisfy the assumption of Lemma 4.7 for $V = H_2(E(3))$. Suppose W_0 and W_1 are given as in Lemma 4.7, and U is the subset of $W_0 \oplus W_1$ consisting of the pairs that satisfy (4.35). Then U is a Zariski closed subset of $W_0 \oplus W_1$. In order to complete the proof, we shall show that U contains a Euclidean open set in $W_0 \oplus W_1$ and hence $U = W_0 \oplus W_1$. Let $(\Gamma, \Lambda) \in W_0 \oplus W_1$ be given by

$$\Gamma := af + bk \quad \text{and} \quad \Lambda = a'f + b'k + c\tau_2.$$

Consider the homology classes

$$u_1 := \frac{1}{\sqrt{2}}(\tau_1 + \tau_3) \quad \text{and} \quad u_2 := \frac{1}{\sqrt{2}}(\tau_1 - \tau_3),$$

which have nonzero intersection with α . There is an element $A_{t,\theta} \in \text{SO}(H_2(E(n)); f)$ such that

$$\begin{aligned} A_{t,\theta}(\Gamma) &:= af + b(k + t^2f + tu_1), \\ A_{t,\theta}(\Lambda) &:= a'f + b'(k + t^2f + tu_1) + c'(\cos(\theta)\tau_2 + \sin(\theta)u_2). \end{aligned}$$

If a and a' are close enough to each other and b , b' and c' are close enough to 1, then t and θ can be chosen such that $A_{t,\theta}(\Gamma), A_{t,\theta}(\Lambda) \in W$. Therefore, U contains an open subset of $W_0 \oplus W_1$. □

4.4 Invariants of $E(n)$

In this section, we compute the invariants of the elliptic surface $E(n)$. We start with the simpler case that $w \cdot f \not\equiv 0 \pmod 3$:

Proposition 4.37 *Suppose w is a 2-cycle in $E(n)$ with $w \cdot f \not\equiv 0 \pmod 3$. Then $E(n)$ has w -simple type, and for $\Gamma, \Lambda \in H_2(E(n))$,*

$$\widehat{D}_{E(n),w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-2}.$$

Proof The proof of this proposition is similar to that of Proposition 4.25. Applying Proposition 3.35 for $X_1 = E(n - 2)$, $X_2 = E(2)$ and $X_3 = X_4 = E(1)$ gives us enough relations to verify the proposition by induction. □

In the blown-up elliptic surface $E(n) \# \overline{\mathbb{C}\mathbb{P}^2}$, there is an embedded sphere with self-intersection $-(n + 1)$, given by tubing a section of the elliptic fibration and the exceptional sphere in $\overline{\mathbb{C}\mathbb{P}^2}$. Therefore, there is a copy of the Gompf nucleus $G(n + 1)$ in $G(n) \# \overline{\mathbb{C}\mathbb{P}^2}$. The homology class $E + f$ can be realized by an embedded surface \bar{E}

in $M_n := G(n) \# \overline{\mathbb{C}\mathbb{P}^2} \setminus G(n+1)$. The surface \bar{E} determines a generator of $H_2(M_n)$. Similarly, the nucleus $G(n+2)$ can be embedded in $G(n) \# 2\overline{\mathbb{C}\mathbb{P}^2}$. Therefore, there are copies of $G(4)$ in the 4-manifolds $E(4)$, $E(3) \# \overline{\mathbb{C}\mathbb{P}^2}$ and $E(2) \# 2\overline{\mathbb{C}\mathbb{P}^2}$. Let $Z_0 \subset E(4)$, $Z_1 \subset E(3) \# \overline{\mathbb{C}\mathbb{P}^2}$ and $Z_2 \subset E(2) \# 2\overline{\mathbb{C}\mathbb{P}^2}$ be the complements of $G(4)$ in these manifolds. Then the boundary of Z_i is diffeomorphic to $\Sigma(2, 3, 23)$. It is clear from the inductive construction of $E(n)$ that there is an embedding of Z_i in $E(n-i) \# i\overline{\mathbb{C}\mathbb{P}^2}$ for $n \geq 4$. In fact, if $W(n)$ is the fiber sum $E(n-4) \#_f G(4)$, then $E(n-i) \# i\overline{\mathbb{C}\mathbb{P}^2}$ is diffeomorphic to $Z_i \#_{\Sigma(2,3,23)} W(n)$.

The 4-manifold Z_i gives rise to elements of $I_*(\Sigma(2, 3, 23))$ as it is explained in Proposition 3.45. Suppose $V_0 \subseteq I_*(\Sigma(2, 3, 23))$ is the vector space generated by the element $D_{Z_0, v_0}(1)$ where v_0 is a 2-cycle in Z_0 with $v_0^2 \equiv 0 \pmod 3$. Similarly, define V_1 to be the vector space generated by the three elements $D_{Z_1, w}(1)$ where w is one of the following elements, which satisfy $w^2 \equiv 1 \pmod 3$:

$$v_1 := \bar{E} + \tau_1 - g_1, \quad v_2 := -\bar{E} + \tau_1 - g_1, \quad v_3 := \tau_1.$$

Finally, let V_2 be the subspace of $I_*(\Sigma(2, 3, 23))$ which is generated by the elements of the form $D_{Z_2, w}(1)$ where $w^2 \equiv 2 \pmod 3$.

Proposition 4.38 *The space V_i is a subspace of $I_4(\Sigma(2, 3, 23))$. Furthermore, the dimension of V_i is at least $\frac{1}{2}(i+2)(i+1)$.*

Proof The first part of the proposition is an immediate consequence of Proposition 3.45. To show that $\dim(V_0) = 1$, let $D_{Z_0, v_0}(1) = 0$. By Proposition 3.45, $D_{E(4), v_0+w_0}(z)$ vanishes for a 2-cycle w_0 in $G(4)$ and $z \in \mathbb{A}(G(4))^{\otimes 2}$. If w_0 is chosen such that $w_0 \cdot f \not\equiv 0 \pmod 3$, then Proposition 4.37 asserts that there is a z such that $D_{E(4), v_0+w_0}(z) \neq 0$, which is a contradiction.

Next, we consider the case that $i = 1$. By Proposition 3.45, a linear relation among the vectors $D_{Z_1, v_l}(1)$ for $1 \leq l \leq 3$ implies that there are constant numbers c_l such that

$$(4.39) \quad \sum_{1 \leq l \leq 3} c_l \hat{D}_{E(3) \# \overline{\mathbb{C}\mathbb{P}^2}, v_l+w_0}(e^{\sigma(2)+\sigma(3)}) = 0.$$

Here w_0 is a 2-cycle in $G(4)$ and σ is the embedded (-4) -sphere in $G(4)$. We can use the blowup formula and the results of the previous subsection to evaluate the above power series and to conclude that

$$\sum_{1 \leq l \leq 3} c_l (\cosh(\sqrt{3}t_2) + \zeta^{-(v_l+w_0) \cdot E} e^{i\sqrt{3}t_3} + \zeta^{(v_l+w_0) \cdot E} e^{-i\sqrt{3}t_3}) = 0.$$

Since $v_l \cdot E \equiv -l \pmod 3$, the constants c_l are equal to zero. Therefore, $\dim(V_1) = 3$. Similar proofs can be applied to the case that $i = 2$. \square

Theorem 4.40 *Suppose w is a 2-cycle in $E(n)$ and $\Gamma, \Lambda \in H_2(E(n))$. Then*

$$(4.41) \quad \widehat{D}_{E(n),w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-2}.$$

Proof Since the dimension of $I_4(\Sigma(2, 3, 23))$ is equal to 9, Proposition 4.38 implies that there are vectors $\phi_i \in V_i$ such that

$$(4.42) \quad \phi_0 + \phi_1 + \phi_2 = 0.$$

We use this identity to verify (4.41) inductively for the remaining invariants of $E(n)$. Firstly, consider the case that $\phi_0 \neq 0$. Then we can assume that $\phi_0 = D_{Z_0,v_0}(1)$. There are also 2-cycles w_1, \dots, w_k in Z_2 and constants $c_1, c_2, c_3, c'_1, \dots, c'_k$ such that

$$(4.43) \quad \phi_1 = \sum_{l=1}^3 c_l D_{Z_1,v_l}(1) \quad \text{and} \quad \phi_2 = \sum_{j=1}^k c'_j D_{Z_2,w_j}(1).$$

Suppose $\Gamma, \Lambda \in H_2(W(n))$ and w is a 2-cycle in $W(n)$. Then Proposition 3.45 implies that the $U(3)$ -series $\widehat{D}_{E(n),v_0+w}(e^{\Gamma(2)+\Lambda(3)})$ is equal to

$$(4.44) \quad -\sum_l c_l \widehat{D}_{E(n-1)\#\overline{\mathbb{C}\mathbb{P}^2},v_l+w}(e^{\Gamma(2)+\Lambda(3)}) - \sum_j c'_j \widehat{D}_{E(n-2)\#2\overline{\mathbb{C}\mathbb{P}^2},w_j+w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-4}A(\Gamma, \Lambda),$$

where the term $A(\Gamma, \Lambda)$ is a linear combination of the six expressions

$$(4.45) \quad \begin{matrix} \cosh(\sqrt{3}f \cdot \Gamma)^2, & \cosh(\sqrt{3}f \cdot \Gamma)\zeta^{\mp w \cdot f} e^{\pm i\sqrt{3}f \cdot \Lambda}, \\ \zeta^{\mp 2w \cdot f} e^{\pm 2i\sqrt{3}f \cdot \Lambda}, & 1, \end{matrix}$$

and the coefficients of this linear combination do not depend on w . To derive (4.44) we use the facts that $\Gamma \cdot E = -\Gamma \cdot f$ and $\Lambda \cdot E = -\Lambda \cdot f$. In the case $w \cdot f \not\equiv 0 \pmod 3$, $A(\Gamma, \Lambda)$ is equal to $G(\Gamma, \Lambda, w \cdot f)^2$ using Proposition 4.37. This identity holds also in the case that $w \cdot f \equiv 0 \pmod 3$, because $A(\Gamma, \Lambda)$ is a linear combination of the terms in (4.45) with coefficients which are independent of w . Therefore, for a general 2-cycle w in $W(n)$ and $\Gamma, \Lambda \in H_2(W(n))$,

$$\widehat{D}_{E(n),v_0+w}(e^{\Gamma_2+\Lambda_3}) = e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-2}.$$

Then Proposition 4.21 implies $\widehat{D}_{E(n),w}(e^{\Gamma_2+\Lambda_3})$ equals $e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}G(\Gamma, \Lambda, w)^{n-2}$ for any 2-cycle w in $E(n)$ and $\Gamma, \Lambda \in H_2(E(n))$.

Next, assume that $\phi_0 = 0$. We assume that the nonzero vectors ϕ_1 and ϕ_2 are given as in (4.43). Fix an arbitrary 2-cycle w in $W(n + 1)$ and homology classes $\Gamma, \Lambda \in H_2(W(n + 1))$. Another application of Proposition 3.45 shows that

$$(4.46) \quad \sum_{1 \leq l \leq 3} c_l \widehat{D}_{E(n) \# \overline{\mathbb{C}\mathbb{P}^2}, v_l+w}(e^{\Gamma(2)+\Lambda(3)})$$

is equal to

$$\sum_j c'_j \widehat{D}_{E(n-1) \# 2\overline{\mathbb{C}\mathbb{P}^2}, w_j+w}(e^{\Gamma(2)+\Lambda(3)}).$$

By our inductive calculation of the invariants of $E(n)$, the latter expression is equal to

$$e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-3}B(\Gamma, \Lambda).$$

Here $B(\Gamma, \Lambda)$ is a linear combination of the six expressions

$$\cosh(\sqrt{3}f \cdot \Gamma)^2, \quad \cosh(\sqrt{3}f \cdot \Gamma)\zeta^{\mp w \cdot f} e^{\pm i\sqrt{3}f \cdot \Lambda}, \quad \zeta^{\mp 2w \cdot f} e^{\pm 2i\sqrt{3}f \cdot \Lambda}, \quad 1.$$

As in the previous case, the coefficients of this linear combination are determined by c'_j and w_j . In particular, they do not depend on w . Therefore, we can determine these coefficients by considering the case that $w \cdot f \not\equiv 0 \pmod 3$, for which we already computed the invariants. Therefore, $B(\Gamma, \Lambda)$ is equal to

$$G(\Gamma, \Lambda, w \cdot f) \sum_{1 \leq l \leq 3} \frac{1}{3}c_l(\cosh(\sqrt{3}E \cdot \Gamma) + \zeta^{-w \cdot E} e^{i\sqrt{3}E \cdot \Lambda} + \zeta^{w \cdot E} e^{-i\sqrt{3}E \cdot \Lambda}).$$

For an arbitrary 2-cycle $w \subset E(n) \# \overline{\mathbb{C}\mathbb{P}^2}$ and $\Gamma, \Lambda \in E(n) \# \overline{\mathbb{C}\mathbb{P}^2}$, let $\widehat{P}_w(\Gamma, \Lambda)$ be the power series given by subtracting $\widehat{D}_w(e^{\Gamma(2)+\Lambda(3)})$ from the power series

$$e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)}G(\Gamma, \Lambda, w \cdot f)^{n-2} \frac{1}{3}(\cosh(\sqrt{3}E \cdot \Gamma) + \zeta^{w \cdot E} e^{i\sqrt{3}E \cdot \Lambda} + \zeta^{-w \cdot E} e^{-i\sqrt{3}E \cdot \Lambda}).$$

Then we can rephrase our conclusion in the form of the identity

$$(4.47) \quad \sum_{1 \leq l \leq 3} c_l \widehat{P}_{v_l+w}(\Gamma, \Lambda) = 0$$

where w is a 2-cycle in $W(n + 1)$ and $\Gamma, \Lambda \in H_2(W(n + 1))$. Suppose $p_w^{i,j}$ is the polynomial on $H_2(E(n) \# \overline{\mathbb{C}\mathbb{P}^2}) \oplus H_2(E(n) \# \overline{\mathbb{C}\mathbb{P}^2})$ of bidegree (i, j) determined by the coefficient of $t_2^i t_3^j$ in \widehat{P}_w . Then $p_w^{i,j}$ can be evaluated at

$$(\Gamma_1, \dots, \Gamma_i; \Lambda_1, \dots, \Lambda_j)$$

for $\Gamma_k, \Lambda_k \in H_2(E(n) \# \overline{\mathbb{C}\mathbb{P}^2})$. By induction on $i + j$, we shall show that $p_w^{i,j}$ vanishes for all possible choices of w . By considering the constant terms of (4.47) for empty w , we have

$$c_1 p_{v_1}^{0,0} + c_2 p_{v_2}^{0,0} + c_3 p_{v_3}^{0,0} = 0.$$

The blowup formula asserts that $p_{v_1}^{0,0} = p_{v_2}^{0,0} = 0$. Therefore, if $c_3 \neq 0$, then $p_{v_3}^{0,0} = 0$. Thus Proposition 4.21 and the blowup formula show that $p_w^{0,0} = 0$ for all 2-cycles w in $E(n) \# \overline{\mathbb{C}\mathbb{P}^2}$. If $c_3 = 0$, then by (4.47),

$$c_1 p_{v_1}^{2,0}(\sigma + E, \sigma + E) + c_2 p_{v_2}^{2,0}(\sigma + E, \sigma + E) = 0$$

and

$$c_1 p_{v_1}^{0,1}(\sigma + E) + c_2 p_{v_2}^{0,1}(\sigma + E) = 0.$$

The blowup formula asserts that

$$c_1 p_{v_1-E}^{0,0} + c_2 p_{v_2+E}^{0,0} = 0 \quad \text{and} \quad c_1 p_{v_1-E}^{0,0} - c_2 p_{v_2+E}^{0,0} = 0.$$

Consequently, at least one of the numbers $p_{v_1-E}^{0,0}$ and $p_{v_2+E}^{0,0}$ is zero and we can derive the same conclusion as in the previous case.

Now assume that the polynomial $p_w^{i,j}$ vanishes for $i + j \leq k$ and any 2-cycle in $E(n) \# \overline{\mathbb{C}\mathbb{P}^2}$. Fix (i, j) , $\tilde{\Gamma}_1, \dots, \tilde{\Gamma}_i$ and $\tilde{\Lambda}_1, \dots, \tilde{\Lambda}_j$ such that $i + j = k + 1$ and $\tilde{\Gamma}_l$ and $\tilde{\Lambda}_{l'}$ are equal to either $\sigma + E$ or f . Then apply (4.47) to conclude that

$$(4.48) \quad \sum_{1 \leq l \leq 3} c_l p_{v_l}^{i,j}(\tilde{\Gamma}_1, \dots, \tilde{\Gamma}_i; \tilde{\Lambda}_1, \dots, \tilde{\Lambda}_j) = 0.$$

The blowup formula and the induction hypothesis imply that every term of the form $\sigma + E$ can be replaced with σ . Thus, if $c_3 \neq 0$,

$$(4.49) \quad p_{v_1}^{i,j}(\Gamma_1, \dots, \Gamma_i; \Lambda_1, \dots, \Lambda_j) = 0$$

for $\Gamma_k, \Lambda_k \in H_2(G(n))$. Therefore, Proposition 4.21 and the blowup formula allow us to complete the verification of the induction step. If $c_3 = 0$, we can use the analogue of (4.48) for

$$(\sigma + E, \sigma + E, \tilde{\Gamma}_1, \dots, \tilde{\Gamma}_i; \tilde{\Lambda}_1, \dots, \tilde{\Lambda}_j) \quad \text{and} \quad (\tilde{\Gamma}_1, \dots, \tilde{\Gamma}_i; \sigma + E, \tilde{\Lambda}_1, \dots, \tilde{\Lambda}_j)$$

and argue as in the basis of the induction. This completes the proof of the theorem. \square

4.5 Gluing 4-manifolds along surfaces of self-intersection zero

In this subsection, we use the calculation of $U(3)$ -polynomial invariants for elliptic surfaces to study invariants of another family of closed 4-manifolds:

Definition 4.50 Suppose that X is a smooth 4-manifold, Σ is an oriented surface of genus $g \geq 1$ embedded in X , w is a 2-cycle in X , and \mathcal{H} is a subspace of $H_2(X)$. Then (X, Σ, w) is *permissible with respect to the subspace \mathcal{H}* if the following hold:

- (i) $b^1(X) = 0$ and $b^+(X) > 1$.
- (ii) $\Sigma \cdot \Sigma = 0$.
- (iii) $w \cdot \Sigma \not\equiv 0 \pmod{3}$.
- (iv) Let $z \in \mathbb{A}(\mathcal{H})^{\otimes 2}$, and let u be the 2-cycle $w + l\Sigma$ for $l = 0, 1$ or 2 . Then

$$(4.51) \quad D_{X,u}((\frac{1}{3}a_2)^3 z) = D_{X,u}(z) \quad \text{and} \quad D_{X,u}(a_3 z) = 0.$$

Moreover, there are cohomology classes $K_i \in H^2(X, \mathbb{Z})$ such that K_i is an integral lift of $w_2(TX)$ and $|K_i \cdot \Sigma| \leq 2g - 2$, and for $\Gamma, \Lambda \in \mathcal{H}$, the power series $\widehat{D}_{X,u}(e^{\Gamma(2)+\Lambda(3)})$ is equal to

$$(4.52) \quad e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)} \sum_{i,j} c_{ij} \zeta^{-l\Sigma \cdot (\frac{1}{2}(K_i - K_j))} e^{\frac{\sqrt{3}}{2}(K_i + K_j) \cdot \Gamma + \frac{\sqrt{3}}{2}i(K_i - K_j) \cdot \Lambda}.$$

The cohomology class K_i is called a *basic class* of the triple (X, Σ, w) and $c_{i,j}$ is called the coefficient associated to the pair (K_i, K_j) . In the case that $\mathcal{H} = H_2(X)$, we say (X, Σ, w) is permissible.

Example 4.53 The results of Section 4.2 show that the triple $(E(2), f, w)$ forms a permissible triple where f is a fiber in an elliptic fibration of $E(2)$ and w is a 2-cycle such that $w \cdot f \not\equiv 0 \pmod{3}$. In this case, the only basic class in (4.52) is the zero cohomology class. More generally, we can embed a surface of genus g in $E(2)$ whose self-intersection is equal to $2g - 2$. For example, we can construct such an embedded surface by considering the union of g fibers and a section of the fibration, and then smoothing out the intersection points. Let Σ be the proper transform of this surface after blowing up $E(2)$ at $2g - 2$ points on the surface. Let also w be a 2-cycle in $E(2) \# (2g - 2)\overline{\mathbb{C}\mathbb{P}^2}$ such that $w \cdot \Sigma \not\equiv 0 \pmod{3}$. Then the blowup formula implies that $(E(2) \# (2g - 2)\overline{\mathbb{C}\mathbb{P}^2}, \Sigma, w)$ is a permissible triple. If E_1, \dots, E_{2g-2} are the exceptional classes, then a basic class of this triple has the form $\pm E_1 \pm \dots \pm E_{2g-2}$.

Example 4.54 One can further generalize the previous example by considering a surface Σ with self-intersection 0 embedded in the 4-manifold $E(n) \# k \overline{\mathbb{C}\mathbb{P}^2}$. Let w be a 2-cycle in X such that $w \cdot \Sigma \not\equiv 0 \pmod 3$. Then the blowup formula and [Theorem 4.40](#) can be utilized to verify most requirements of [Definition 4.50](#) for permissibility of (X, Σ, w) . The only missing part is to verify the inequality $|K \cdot \Sigma| \leq 2g - 2$ for basic classes K of X . To check this inequality, note that our basic classes for (X, Σ, w) are the same as $U(2)$ -basic classes for X [[53](#); [25](#); [63](#); [54](#); [26](#)]. Therefore, the desired inequality is a consequence of the adjunction inequality in [[54](#)].

Remark 4.55 Among the conditions in [Definition 4.50](#), (iv) might seem the most mysterious one. The first part of this condition is a weaker version of the simple type condition in (1.7). We expect that if X has simple type in the sense of [[54](#)] and (X, w, Σ) satisfies properties (i), (ii) and (iii) of [Definition 4.50](#), then condition (iv) automatically holds. In fact, we propose a more general conjecture as [Conjecture 7.2](#) in [Section 7](#). This conjecture is the $U(3)$ version of Kronheimer and Mrowka’s structure theorem [[54](#)].

Definition 4.56 For $g \geq 1$, the set of all integer pairs (a, b) which satisfy the following two properties is denoted by \mathcal{C}_g :

- (i) a and b have the same parity.
- (ii) $|a| + |b| \leq 2g - 2$.

We will write N_g for the number of the elements of \mathcal{C}_g , which is equal to $2g(g - 1) + 1$.

The following proposition can be verified using the permissible triples provided by [Example 4.53](#):

Proposition 4.57 For any $(a, b) \in \mathcal{C}_g$, there are a permissible triple (X, w, Σ) and basic classes K_i and K_j of the triple such that Σ has genus g and

$$(4.58) \quad a = \frac{1}{2}((K_i + K_j) \cdot \Sigma) \quad \text{and} \quad b = \frac{1}{2}((K_i - K_j) \cdot \Sigma).$$

For a permissible triple (X, Σ, w) , define

$$D_{X,w,\Sigma}(e^{\Gamma(2)+\Lambda(3)}) := \sum_{0 \leq l \leq 2} D_{X,w+l\Sigma}(e^{\Gamma(2)+\Lambda(3)}).$$

More generally, $D_{X,w,\Sigma}(z)$ is defined to be the sum $D_{X,w+l\Sigma}(z)$ for different values of $0 \leq l \leq 2$. The dimension formula shows that if all terms in z have a fixed degree, then only one of the 2-cycles $w + l\Sigma$ is involved in the definition of $D_{X,w,\Sigma}(z)$.

Lemma 4.59 Suppose (X, w, Σ) is a permissible triple as above and $d_w \in \mathbb{Z}/3\mathbb{Z}$ is defined to be $b^+ + 1 - w \cdot w$. Then $D_{X,w,\Sigma}((\frac{1}{3}a_2)^m e^{\Gamma(2)+\Lambda(3)})$ is equal to

$$\sum c_{ij} \zeta^{l_{i,j}(d_w-m)} e^{\zeta^{2l_{i,j}(\frac{1}{2}Q(\Gamma)+\frac{\sqrt{3}}{2}i(K_i-K_j)\cdot\Lambda)+\zeta^{l_{i,j}(-Q(\Lambda)+\frac{\sqrt{3}}{2}(K_i+K_j)\cdot\Gamma)},$$

where the inner sum is over all pairs of basic classes (K_i, K_j) . Moreover, $l_{i,j} \in \mathbb{Z}/3\mathbb{Z}$ is equal to $(w \cdot \Sigma)(\frac{1}{2}(K_i - K_j)) \cdot \Sigma$.

Proof For a 4-manifold W with simple type, the power series $D_{X,w}$ can be recovered from $\widehat{D}_{X,w}$ in the following way:

$$D_{X,w}((\frac{1}{3}a_2)^m e^{\Gamma(2)+\Lambda(3)}) = \frac{1}{3} \sum_{0 \leq k \leq 2} \zeta^{k(d_w-m)} \widehat{D}_{X,w}(e^{\zeta^k \Gamma(2)+\zeta^{2k} \Lambda(3)}).$$

Therefore

$$\begin{aligned} \sum_{0 \leq l \leq 2} D_{X,w+l\Sigma}((\frac{1}{3}a_2)^m e^{\Gamma(2)+\Lambda(3)}) \\ = \frac{1}{3} \sum_{0 \leq k,l \leq 2} \zeta^{k(d_w+l\Sigma-m)} \widehat{D}_{X,w+l\Sigma}(e^{\zeta^k \Gamma(2)+\zeta^{2k} \Lambda(3)}). \end{aligned}$$

Then, we can use the permissibility of (X, w, Σ) to rewrite the right-hand side in terms of basic classes. A straightforward simplification gives the desired result. \square

Suppose (X, w, Σ) and (X', w', Σ) are two permissible triples such that the embedded surfaces of genus g are identified with each other, and this identification is lifted to the normal bundles. Suppose also w and w' intersect Σ in the same number of points with the same signs. As explained in Section 3.3, we can form the triple $(X \#_{\Sigma} X', w \# w', \Sigma)$. There is also a subspace $\mathcal{K} \subseteq H_2(X) \oplus H_2(X')$ such that there is a map $\#: \mathcal{K} \rightarrow H_2(X \#_{\Sigma} X')$. The main goal of this section is to compute $\widehat{D}_{X \#_{\Sigma} X', w \# w'}(e^{\Gamma(2)+\Lambda(3)})$ for $\Gamma, \Lambda \in \text{im}(\#)$ in terms of the invariants of the pairs (X, w) and (X', w') .

The basic idea to achieve this goal is to use the gluing property in (3.30). Therefore, the R_N -module $\mathbb{I}_{g,d}^N$, introduced in Section 3.3, for $N = 3$ and $d = w \cdot \Sigma$, plays a key role in computing the invariants of $(X \#_{\Sigma} X', w \# w')$. In fact, we can replace $\mathbb{I}_{g,d}^3$ with a smaller module. Before giving the definition of this smaller module, we introduce some conventions. From now on, we drop 3 from our notation and denote this Fukaya-Floer homology module with $\mathbb{I}_{g,d}$. Moreover, for a permissible triple (X, w, Σ) the intersection $w \cdot \Sigma$ is denoted by d , unless otherwise is stated.

Let $\tilde{\mathbb{I}}_{g,d} \subset \mathbb{I}_{g,d}$ be the $\mathbb{C}[[t_2, t_3]]$ -module generated by the following relative elements in $\mathbb{I}_{g,d}$:

$$(4.60) \quad D_{X^\circ, w^\circ, \Sigma}(e^{\Gamma_{(2)}^\circ + \Lambda_{(3)}^\circ}) := \sum_{l \in \mathbb{Z}/3\mathbb{Z}} D_{X^\circ, w^\circ + l\Sigma}(e^{\Gamma_{(2)}^\circ + \Lambda_{(3)}^\circ}),$$

where (X, w, Σ) is a permissible triple, and $\Gamma, \Lambda \in H_2(X, \mathbb{Z})$ with $\Gamma \cdot \Sigma = \Lambda \cdot \Sigma = 1$. By (3.30), the pairing of this element with $D_{\Delta_g, \delta_g}(ze^{D_{(2)} + D_{(3)}})$ is equal to the following element of $\mathbb{C}[[t_2, t_3]]$:

$$(4.61) \quad D_{X, w, \Sigma}(ze^{\Gamma_{(2)} + \Lambda_{(3)}}).$$

Suppose $\mathbb{C}[[t_2, t_3]][x, y, z]$ is the ring of polynomials of three variables with coefficients in $\mathbb{C}[[t_2, t_3]]$ and $z = P(a_2, \Sigma_{(2)}, \Sigma_{(3)})$ for $P \in \mathbb{C}[[t_2, t_3]][x, y, z]$. Then Lemma 4.59 shows that the pairing of (4.60) and $D_{\Delta_g, \delta_g}(ze^{D_{(2)} + D_{(3)}})$ is equal to

$$\sum_{(a,b) \in C_g} \left(\zeta^{dbd_w} P(u(a, b)) \times \sum c_{i,j} e^{\zeta^{2db}(\frac{1}{2}Q(\Gamma) + \frac{\sqrt{3}}{2}i(K_i - K_j) \cdot \Lambda) + \zeta^{db}(-Q(\Lambda) + \frac{\sqrt{3}}{2}(K_i + K_j) \cdot \Gamma)} \right),$$

where the inner sum is over all pairs of basic classes (K_i, K_j) such that

$$(4.62) \quad \frac{1}{2}(K_i + K_j) \cdot \Sigma = a \quad \text{and} \quad \frac{1}{2}(K_i - K_j) \cdot \Sigma = b,$$

and

$$(4.63) \quad u(a, b) := (3\zeta^{2db}, \zeta^{db}\sqrt{3}a + \zeta^{2db}t_2, \zeta^{2db}\sqrt{3}ib - 2\zeta^{db}t_3).$$

For $\lambda = (\alpha, \beta) \in C_g$, fix a polynomial $P_\lambda \in \mathbb{C}[[t_2, t_3]][x, y, z]$ such that

$$(4.64) \quad P_\lambda(u(\alpha, \beta)) = 1 \quad \text{and} \quad P_\lambda(u(a, b)) = 0 \quad \text{for } (a, b) \neq (\alpha, \beta).$$

Define a map $\Phi: \tilde{\mathbb{I}}_{g,d} \rightarrow \mathbb{C}[[t_1, t_2]]^{N_g}$ by

$$\Phi(\eta) := \left\{ \langle \eta, D_{\Delta_g, \delta_g}(P_\lambda(a_2, \Sigma_{(2)}, \Sigma_{(3)})e^{D_{(2)} + D_{(3)}}) \rangle \right\}_{\lambda \in C_g}.$$

By (3.30), the homomorphism Φ maps the relative element in (4.60) to an element of $\mathbb{C}^{N_g}[[t_2, t_3]]$, whose component corresponding to $\lambda = (a, b)$, denoted by $c_{X, w, \Sigma}^\lambda(\Gamma, \Lambda)$, is equal to

$$(4.65) \quad \zeta^{dbd_w} \sum c_{i,j} e^{\zeta^{2db}(\frac{1}{2}Q(\Gamma) + \frac{\sqrt{3}}{2}i(K_i - K_j) \cdot \Lambda) + \zeta^{db}(-Q(\Lambda) + \frac{\sqrt{3}}{2}(K_i + K_j) \cdot \Gamma)},$$

where the sum is over the pairs of basic classes (K_i, K_j) that satisfy (4.62). Let $\mathbb{C}((t_1, t_2))$ denote the field of fractions of $\mathbb{C}[[t_2, t_3]]$. Then Φ induces a map

$$\bar{\Phi}: \tilde{\mathbb{I}}_{g,d} \otimes_{\mathbb{C}[[t_1,t_2]]} \mathbb{C}((t_1, t_2)) \rightarrow \mathbb{C}((t_1, t_2))^{N_g}.$$

Proposition 4.66 *The map Φ is injective. Moreover, $\bar{\Phi}$ is an isomorphism of vector spaces.*

Proof Suppose \mathcal{I} is the ideal in $\mathbb{A}_g^3 \otimes_{\mathbb{C}} \mathbb{C}[[t_2, t_3]]$ generated by $a_3, \gamma_{(2)}, \gamma_{(3)}$ and the elements $P(a_2, \Sigma_{(2)}, \Sigma_{(3)})$, where $P \in \mathbb{C}[[t_2, t_3]][x, y, z]$ is a polynomial evaluating to zero at the points in (4.63). Then the pairing of (4.60) and $D_{\Delta_g, \delta_g}(ze^{D_{(2)}+D_{(3)}})$ vanishes when $z \in \mathcal{I}$. Any element of $\tilde{\mathbb{I}}_{g,d}$ is also invariant with respect to the action of

$$(4.67) \quad \tilde{\epsilon} := \mathbb{I}_*^N([0, 1] \times Y_g, [0, 1] \times \gamma_{g,d} + \Sigma, e^{([0,1] \times \gamma)_{(2)} + \dots + ([0,1] \times \gamma)_{(N)}}).$$

Recall that $Y_g = \Sigma \times S^1$ and γ in (4.67) denotes an S^1 -fiber of Y_g . Any $z \in \mathbb{A}_g^3 \otimes_{\mathbb{C}} \mathbb{C}[[t_2, t_3]]$ can be written as a sum of an element of \mathcal{I} and a $\mathbb{C}[[t_2, t_3]]$ -linear combination of the polynomials $\{P_\lambda\}_{\lambda \in \mathcal{C}_g}$. Therefore, injectivity of Φ is a consequence of Proposition 3.31.

For a given $\lambda_0 \in \mathcal{C}_g$, Proposition 4.57 gives a permissible triple (X, w, Σ) such that the component of the relative element (4.60) corresponding to λ_0 is nonzero. Furthermore, we can change the relative class as in (4.60) by replacing Γ with $\Gamma + s\Sigma$ and Λ with $\Lambda + s\Sigma$. The component of this relative element corresponding to $\lambda = (a, b)$ picks the factor $e^{s(\xi^{2b}\sqrt{3}at_2 + \xi^b\sqrt{3}bt_3 + \xi^b t_2^2 - 2\xi^{2b}t_3^2)}$. Therefore, by taking $\mathbb{C}[[t_1, t_2]]$ -linear combinations of such expressions for different values of s , we can produce elements of $\tilde{\mathbb{I}}_{g,d}$ such that the component corresponding to λ_0 is the only nonzero element. This verifies the second part of the proposition. □

Consider the restriction of the pairing $\langle \cdot, \cdot \rangle$ on $\mathbb{I}_{g,d}$. The above proposition implies that this pairing induces a pairing on $\mathbb{C}((t_1, t_2))^{N_g}$ using the map $\bar{\Phi}$:

$$(4.68) \quad \langle \cdot, \cdot \rangle: \mathbb{C}((t_1, t_2))^{N_g} \times \mathbb{C}((t_1, t_2))^{N_g} \rightarrow \mathbb{C}((t_1, t_2)).$$

Suppose this pairing is given by $\{\xi^{2bd(g-1)}h_{\lambda, \lambda'}^{g,d}\}_{\lambda, \lambda' \in \mathcal{C}_g}$ with respect to the standard basis of $\mathbb{C}((t_1, t_2))^{N_g}$, where b is the second coordinate of λ . The constant $\xi^{2bd(g-1)}$ does not play an important role here. It will be used to obtain a slightly simpler form for our gluing formulas in Proposition 4.71.

Proposition 4.69 *The element $h_{\lambda, \lambda'}^{g,d}(t_2, t_3) \in \mathbb{C}((t_1, t_2))$ is nonzero only if $\lambda = \lambda'$. Furthermore, if $\lambda = (a, b)$ and $|\lambda|_1 := |a| + |b| < 2g - 2$, then $h_{\lambda, \lambda}^{g,d}$ is zero.*

Proof Suppose (X, w, Σ) and (X', w', Σ) are two permissible triples such that w and w' intersect Σ in the same set of points with the same signs. Suppose the homology classes $\Gamma, \Lambda \in H_2(X)$ and $\Gamma', \Lambda' \in H_2(X')$ are chosen such that their intersection with Σ is equal to 1. Then (3.30) asserts that

$$(4.70) \quad D_{X\#_{\Sigma}X', w\#w', \Sigma}(e^{(\Gamma\#\Gamma')_{(2)}+(\Lambda\#\Lambda')_{(3)}}) \\ = \sum_{\lambda, \lambda' \in C_g} \zeta^{2bd(g-1)} h_{\lambda, \lambda'}^{g,d} c_{X, w, \Sigma}^{\lambda}(\Gamma, \Lambda) c_{X', w', \Sigma}^{\lambda'}(\Gamma', \Lambda').$$

Replacing Γ, Γ', Λ and Λ' with $\Gamma + r\Sigma, \Gamma' - r\Sigma, \Lambda + s\Sigma$ and $\Lambda' - s\Sigma$ does not change the left-hand side of the above identity. On the right-hand side, the term corresponding to λ and λ' changes by a factor of the form $e^{r \cdot f(\lambda, \lambda') + s \cdot g(\lambda, \lambda')}$. Here $f(\lambda, \lambda')$ and $g(\lambda, \lambda')$, which can be computed explicitly, are zero if and only if $\lambda = \lambda'$. Therefore, $h_{\lambda, \lambda'}^{g,d}$ has to be nonzero when $\lambda \neq \lambda'$.

Let (X, w, Σ) be the permissible triple of Example 4.53 where Σ has genus $g - 1$. Taking the connected sum of Σ and a homologically trivial torus embedded in a 4-ball produces a permissible triple (X, w, Σ') such that Σ' has genus g . Then $X \#_{\Sigma} X$ can be decomposed as the connected sum of $S^2 \times S^2$ and another 4-manifold with $b^+ > 0$. Theorem 6.14 asserts that for Γ, Λ, Γ' and $\Lambda' \in H_2(X)$, the invariant

$$D_{X\#_{\Sigma'}X, w\#w, \Sigma}(e^{(\Gamma\#\Gamma')_{(2)}+(\Lambda\#\Lambda')_{(3)}})$$

vanishes. If $|\lambda|_1 < 2g - 2$, then we can find Γ and Λ such that $c_{X, w, \Sigma'}^{\lambda}$ is nonzero. Consequently, $h_{\lambda, \lambda}^{g,d}$ is zero for this choice of λ . □

In light of the above proposition, let $h_{a,b}^{g,d}$ be $h_{\lambda, \lambda}^{g,d}$ for $\lambda = (a, b) \in C_g \setminus C_{g-1}$. These are the only nonzero terms among the coefficients of the pairing.

Proposition 4.71 *Suppose (X, w, Σ) and (X', w', Σ) are two permissible triples such that w and w' intersect Σ in the same set of points with the same signs. Then*

$$(4.72) \quad D_{X\#_{\Sigma}X', w\#w'}\left(\left(\frac{1}{3}a_2\right)^3 z\right) = D_{X\#_{\Sigma}X', w\#w'}(z) \quad \text{and} \quad D_{X\#_{\Sigma}X', w\#w'}(a_3 z) = 0$$

for $z \in \text{Sym}^*(H_0(X \#_{\Sigma} X') \oplus \text{im}(\#))^{\otimes 2}$. Suppose also that the intersection number of the homology classes $\Gamma, \Lambda \in H_2(X)$ and $\Gamma', \Lambda' \in H_2(X')$ with Σ is 1. Then $\hat{D}_{X\#_{\Sigma}X', w\#w'}(e^{(\Gamma\#\Gamma')_{(2)}+(\Lambda\#\Lambda')_{(3)}})$ is equal to

$$e^{\frac{1}{2}Q(\Gamma\#\Gamma') - Q(\Gamma\#\Gamma')} \\ \times \sum_{(a,b) \in C_g \setminus C_{g-1}} h_{a,b}^{g,d}(t_2, t_3) \sum c_{ij} d_{i'j'} e^{(\frac{\sqrt{3}}{2}(M_{i,i'} + M_{j,j'}) \cdot \Gamma\#\Gamma' + \frac{\sqrt{3}}{2} \mathbf{i}(M_{i,i'} - M_{j,j'}) \cdot \Lambda\#\Lambda')}.$$

For each (a, b) , the second sum is over the pairs of basic classes (K_i, K_j) and $(L_{i'}, L_{j'})$ such that

$$(4.73) \quad \frac{1}{2}(K_i + K_j) \cdot \Sigma = \frac{1}{2}(L_{i'} + L_{j'}) \cdot \Sigma = a, \quad \frac{1}{2}(K_i - K_j) \cdot \Sigma = \frac{1}{2}(L_{i'} - L_{j'}) \cdot \Sigma = b,$$

and $M_{i,i'}$ and $M_{j,j'}$ are respectively equal to $K_i \# L_{i'}$ and $K_j \# L_{j'}$.

Proof The series $D_{X \#_{\Sigma} X', w \# w', \Sigma}(e^{(\Gamma \# \Gamma')_{(2)} + (\Lambda \# \Lambda')_{(3)}})$ can be computed in terms of the cohomology classes $M_{i,i'}$ by substituting (4.65) into (4.70) and applying Proposition 4.69. Then we argue as in Lemma 4.59 to obtain the desired formula for the $U(N)$ -series $\widehat{D}_{X \#_{\Sigma} X', w \# w', \Sigma}(e^{(\Gamma \# \Gamma')_{(2)} + (\Lambda \# \Lambda')_{(3)}})$. We can follow a similar strategy to compute $\widehat{D}_{X \#_{\Sigma} X', w \# w', \Sigma}(a_2^i a_3^j e^{(\Gamma \# \Gamma')_{(2)} + (\Lambda \# \Lambda')_{(3)}})$ in terms of the classes $M_{i,i'}$. The resulting formulas prove the identities in (4.72). \square

The goal of the remaining part of this section is to determine the power series $h_{a,b}^{g,d}(t_2, t_3)$ up to two constants. Firstly, one can obtain a constraint on this power series by changing the orientation of Σ :

$$(4.74) \quad h_{a,b}^{g,2}(t_2, t_3) = h_{-a,-b}^{g,1}(-t_2, -t_3).$$

We shall obtain more constraints by looking at some explicit 4-manifolds. In the case $g = 1$, we can in fact determine $h_{0,0}^{1,d}(t_2, t_3)$ completely using our calculation of the invariants of $E(n)$:

Corollary 4.75 For $g = 1$, the only nonzero term among the pairing coefficients is

$$h_{0,0}^{1,d}(t_2, t_3) = (\hbar_1 \cosh(\sqrt{3}t_2) - 2\hbar_2 \cos(-\frac{2}{3}\pi d + \sqrt{3}t_3))^2.$$

In particular, if (X, w, Σ) and (X', w', Σ) are permissible triples such that the genus of Σ is equal to 1, and w and w' intersect Σ in the same set of points with the same signs, then

$$(4.76) \quad \widehat{D}_{X \#_{\Sigma} X', w \# w', \Sigma}(e^{(\Gamma \# \Gamma')_{(2)} + (\Lambda \# \Lambda')_{(3)}}) = h_{0,0}^{1,d} \widehat{D}_{X,w}(e^{\Gamma_{(2)} + \Lambda_{(3)}}) \widehat{D}_{X',w'}(e^{\Gamma'_{(2)} + \Lambda'_{(3)}}).$$

Remark 4.77 Identity (4.76) is a consequence of Proposition 3.35, and it holds even if we only require (i), (ii) and (iii) of Definition 4.50 for the triples (X, w, Σ) and (X', w', Σ) .

The 4-manifold $B(1) := E(1) \# \overline{\mathbb{C}\mathbb{P}^2}$ has two elliptic fibrations with fibers f and f' such that $f \cdot f' = 1$. There are also disjoint embedded spheres E and E' in $B(1)$

such that E (respectively, E') intersects f (respectively, f') positively in one point and is disjoint from f' (respectively, f). The fiber sum of n copies of $B(1)$ along f' produces a 4-manifold $B(n)$ which is diffeomorphic to $E(n) \# n\overline{\mathbb{C}\mathbb{P}^2}$ (Figure 1). The exceptional spheres of $B(n)$ are denoted by E^1, \dots, E^n , where E^i is given by the exceptional sphere E in the i^{th} summand of $B(n)$.⁷ Furthermore, one can glue copies of f to produce a surface f_n of genus n with self-intersection zero. The homology class of f_n is equal to $[S] - [E^1] - \dots - [E^n]$ for an appropriate surface S of genus n in $E(n)$. Similarly, one can glue copies of E' to produce σ_n , a sphere of self-intersection $-n$ that is disjoint from f_n and the exceptional spheres. The intersection number of σ_n and f' is equal to 1. The following proposition shows that $(B(n), f_n, w)$ is permissible if $w \cdot f \not\equiv 0 \pmod 3$:

Proposition 4.78 *Suppose w is a 2-cycle in $B(n)$ such that $w \cdot f_n \not\equiv 0 \pmod 3$. For $n \geq 2$, the triple $(B(n), w, f_n)$ is permissible and $\widehat{D}_{B(n),w}(e^{\Gamma(2)+\Lambda(3)})$ is given by*

$$e^{\frac{1}{2}\mathcal{Q}(\Gamma)-\mathcal{Q}(\Lambda)} G'(\Gamma, \Lambda, w \cdot f')^{n-2} \prod_i \frac{1}{3} (\cosh(\sqrt{3}E^i \cdot \Gamma) + 2 \cos(-\frac{2}{3}\pi w \cdot E^i + \sqrt{3}E^i \cdot \Lambda))$$

for $\Gamma, \Lambda \in H_2(B(n))$. Here G' is given by (4.24) after replacing f with f' . In particular, a basic class of $B(n)$ has the form

$$(n - 2k)f' \pm E^1 \pm \dots \pm E^n \quad \text{for } 1 \leq k \leq n - 1.$$

Proof This follows from the blowup formula and Theorem 4.40. □

From the construction, it is clear that there is a diffeomorphism (see Figures 2 and 3)

$$(4.79) \quad \Phi_n: B(n) \#_{f_n} B(n) \rightarrow B(2) \#_{f_2} \dots \#_{f_2} B(2).$$

This diffeomorphism maps f_n and $f'_2 := f' \# f'$ in $B(n) \#_{f_n} B(n)$ to $f'_n := f' \# \dots \# f'$ and f_2 in $B(2) \#_{f_2} \dots \#_{f_2} B(2)$. The sphere $\sigma_n \subset B(n)$ determines two spheres of self-intersection $-n$ in $B(n) \#_{f_n} B(n)$, which are denoted by σ_n^1 and σ_n^2 . The diffeomorphism Φ_n maps σ_n^i to $E_n^i := E^i \# \dots \# E^i$. Therefore, the following elements of $\mathbb{C}[[r_2, s_2, r_3, s_3, t_2, t_3]]$ are equal to each other:

$$(4.80) \quad \widehat{D}_{B(n) \#_{f_n} B(n), kf'_2 + lf_n}(e^{(r_2 f_n + s_2 f'_2)(2) + (r_3 f_n + s_3 f'_2)(3)}) \\ = \widehat{D}_{B(2) \#_{f_2} \dots \#_{f_2} B(2), kf_2 + lf'_n}(e^{(r_2 f'_n + s_2 f_2)(2) + (r_3 f'_n + s_3 f_2)(3)}).$$

⁷For the special case of exceptional spheres in $B(n)$, we deviate from our previous notation and put i as a superscript.




Figure 1: A schematic of $B(n)$: each embedded curve represents an embedded surface and each intersection point of these curves represents an element in the intersection of the corresponding surfaces. Note that E^i and σ_n do not intersect.

Proposition 4.81 Suppose $(a, b) \in \mathcal{C}_2 \setminus \{(0, 0)\}$. Then

$$(4.82) \quad h_{a,b}^{2,d}(t_2, t_3) = \sum_{(\gamma, \eta) \in \mathcal{C}_2 \setminus \{(0,0)\}} h_{a,b,\gamma,\eta}^{2,d} e^{\sqrt{3}\gamma t_2 + \sqrt{3}i\eta t_3},$$

where $h_{a,b,\gamma,\eta}^{2,d}$ is a constant number.

Proof The triple $(B(2), df' + 2df_2, f_2)$ is permissible, and Proposition 4.71 can be utilized to compute the following element of $\mathbb{C}[[r_2, r_3, s_2, s_3, t_2, t_3]]$:

$$\widehat{D}_{B(2) \#_{f_2} B(2), df'_2 + df_2} (e^{(s_2 f'_2 + r_2 f_2)_{(2)} + (s_3 f'_2 + r_3 f_2)_{(3)}}).$$




Figure 2: A schematic of $B(n) \#_{f_n} B(n)$.




Figure 3: A schematic of $B(2) \#_{f_2} \dots \#_{f_2} B(2)$.

We can evaluate this series at $t_2 = t_3 = 1$ to obtain a well-defined element of $\mathbb{C}[[r_2, r_3, s_2, s_3]]$. This power series is equal to

$$(4.83) \quad e^{r_2 s_2 - 2r_3 s_3} \sum_{(a,b) \in \mathcal{C}_2 \setminus \{(0,0)\}} h_{a,b}^{2,d}(s_2, s_3) e^{\sqrt{3}ar_2 + \sqrt{3}ibr_3} \left(\sum_{(K_i, K_j)} c_{ij} \right)^2,$$

where the inner sum is over all pairs of basic classes (K_i, K_j) of $(B(2), df' + 2df_2, f_2)$ such that

$$\frac{1}{2}(K_i + K_j) \cdot f_2 = a \quad \text{and} \quad \frac{1}{2}(K_i - K_j) \cdot f_2 = b.$$

For each choice of (a, b) , the inner sum is nonzero. The identity (4.80) shows that the expression (4.83) is invariant with respect to the symmetry of $\mathbb{C}[[r_2, r_3, s_2, s_3]]$ that switches r_2 with s_2 and r_3 with s_3 . This can be used to show that $h_{(a,b)}^{2,d}(t_2, t_3)$ has the form in (4.82). \square




Figure 4: A schematic of $B(2) \#_{f_2} B(2)$: reflection with respect to the dashed line represents the diffeomorphism Φ_2 of this 4-manifold.

Proposition 4.84 The constant numbers $h_{a,b,\gamma,\eta}^{2,d}$ are zero except possibly

$$h_{2,0,2,0}^{2,d}, \quad h_{0,2,0,2}^{2,d}, \quad h_{0,-2,0,-2}^{2,d} \quad \text{and} \quad h_{-2,0,-2,0}^{2,d}.$$

Furthermore, there are real numbers \hbar_3 and \hbar_4 such that

$$(4.85) \quad h_{2,0,2,0}^{2,d} = h_{-2,0,-2,0}^{2,d} = \hbar_3, \quad h_{0,2,0,2}^{2,d} = \zeta^d \hbar_4 \quad \text{and} \quad h_{0,-2,0,-2}^{2,d} = \zeta^{-d} \hbar_4.$$

Proof Firstly, for the purpose of brevity, let

$$N_{X,w}(\Gamma, \Lambda) := \widehat{D}_{X,w}(e^{\Gamma(2)+\Lambda(3)})e^{\frac{1}{2}\mathcal{Q}(\Gamma)+\mathcal{Q}(\Lambda)}.$$

Proposition 4.81 implies that $N_{B(2)\#_{f_2}B(2),df'_2+d'f_2}(\Gamma, \Lambda)$, for $\Gamma, \Lambda \in \text{im}(\#)$, has the form

$$\sum M_{a,b,\gamma,\eta}^{d,d'}(i, j, i', j')e^{\left(\frac{\sqrt{3}}{2}((K_i\#K_{i'}+K_j\#K_{j'}+2\gamma f_2)\cdot\Gamma)+\frac{\sqrt{3}}{2}i((K_i\#K_{i'}-K_j\#K_{j'}+2\eta f_2)\cdot\Lambda)\right)},$$

where the sum is over the pairs $(a, b), (\gamma, \eta) \in \mathcal{C}_2 \setminus \{(0, 0)\}$ and the basic classes $K_i, K_j, K_{i'}$ and $K_{j'}$ of the permissible triple $(B(2), df' + 2d'f_2, f_2)$ such that

$$\frac{1}{2}(K_i + K_j) \cdot f_2 = \frac{1}{2}(K_{i'} + K_{j'}) \cdot f_2 = a, \quad \frac{1}{2}(K_i - K_j) \cdot f_2 = \frac{1}{2}(K_{i'} - K_{j'}) \cdot f_2 = b.$$

In the above expression, the constant $M_{a,b,\gamma,\eta}^{d,d'}(i, j, i', j')$ is equal to $h_{a,b,\gamma,\eta}^{2,d} c_{ij} c_{i'j'}$, where c_{ij} and $c_{i'j'}$ are the coefficients associated to the pairs of basic classes (K_i, K_j) and $(K_{i'}, K_{j'})$ for the permissible triple $(B(2), df' + 2d'f_2, f_2)$. We need the following elementary lemma:

Lemma 4.86 Suppose V is a vector space and $\{f_i\}_{1 \leq i \leq N}$ is a finite set of distinct complex-valued linear functionals on V . Then the functions $\{e^{f_i}\}_{1 \leq i \leq N}$ are linearly independent over \mathbb{C} .

This lemma is an immediate consequence of the existence of a line $l \subseteq V$ such that the restrictions $f_i|_l$ are distinct. We apply this lemma in the case that V is the following subspace of $H_2(B(2)\#_{f_2}B(2))^{\oplus 2}$:

$$(\text{im}(\#) \cap (\Phi_2)_*(\text{im}(\#))) \oplus (\text{im}(\#) \cap (\Phi_2)_*(\text{im}(\#))).$$

Since Φ maps $df'_2 + d'f_2$ to $df_2 + d'f'_2$, we have

$$N_{B(2)\#_{f_2}B(2),df'_2+d'f_2}(\Gamma, \Lambda) = N_{B(2)\#_{f_2}B(2),df_2+d'f'_2}((\Phi_2)_*(\Gamma), (\Phi_2)_*(\Lambda))$$

for $(\Gamma, \Lambda) \in V$. This identity implies that

$$(4.87) \quad \sum M_{a,b,\gamma,\eta}^{d,d'}(i, j, i', j')e^{f_{a,b,\gamma,\eta}^{i,j,i',j'}(\Gamma,\Lambda)} - M_{a,b,\gamma,\eta}^{d,d'}(i, j, i', j')e^{g_{a,b,\gamma,\eta}^{i,j,i',j'}(\Gamma,\Lambda)} = 0.$$

	f_2	f'_2	E_2^1	E_2^2	σ_2^1	σ_2^2
$L_1^\delta := E_2^1 + E_2^2 + \delta f_2$	2	δ	$\delta - 2$	$\delta - 2$	0	0
$L_2^\delta := -E_2^1 - E_2^2 + \delta f_2$	-2	δ	$\delta + 2$	$\delta + 2$	0	0
$L_3^\delta := -E_2^1 + E_2^2 + \delta f_2$	0	δ	$\delta + 2$	$\delta - 2$	0	0
$L_4^\delta := E_2^1 - E_2^2 + \delta f_2$	0	δ	$\delta - 2$	$\delta + 2$	0	0
$L_5^\delta := \zeta_1 - \zeta_2 + \delta f_2$	0	δ	δ	δ	0	0
$L_6^\delta := \zeta_2 - \zeta_1 + \delta f_2$	0	δ	δ	δ	0	0
$J_1^\delta := \sigma_2^1 + \sigma_2^2 + \delta f'_2$	δ	2	0	0	$\delta - 2$	$\delta - 2$
$J_2^\delta := -\sigma_2^1 - \sigma_2^2 + \delta f'_2$	δ	-2	0	0	$\delta + 2$	$\delta + 2$
$J_3^\delta := -\sigma_2^1 + \sigma_2^2 + \delta f'_2$	δ	0	0	0	$\delta + 2$	$\delta - 2$
$J_4^\delta := \sigma_2^1 - \sigma_2^2 + \delta f'_2$	δ	0	0	0	$\delta - 2$	$\delta + 2$
$J_5^\delta := \xi_1 - \xi_2 + \delta f'_2$	δ	0	0	0	δ	δ
$J_6^\delta := \xi_2 - \xi_1 + \delta f'_2$	δ	0	0	0	δ	δ

Table 1: Pairing of the cohomology classes L_m^δ and J_m^δ with some elements of $(\text{im}(\#) \cap (\Phi_2)_*(\text{im}(\#)))$.

Here $f_{a,b,\gamma,\eta}^{i,j,i',j'}$ is defined by the pair of cohomology classes

$$\left(\frac{\sqrt{3}}{2}(K_i \# K_{i'} + K_j \# K_{j'} + 2\gamma f_2), \frac{\sqrt{3}}{2}i(K_i \# K_{i'} - K_j \# K_{j'} + 2\eta f_2)\right),$$

and $g_{a,b,\gamma,\eta}^{i,j,i',j'} = \Phi_2^*(f_{a,b,\gamma,\eta}^{i,j,i',j'})$. In the case of $f_{a,b,\gamma,\eta}^{i,j,i',j'}$, this pair is equal to

$$\left(\frac{\sqrt{3}}{2}(L_m^\delta + L_{m'}^{\delta'}), \frac{\sqrt{3}}{2}i(L_m^\delta - L_{m'}^{\delta'})\right)$$

for an appropriate choice of m and δ , where L_m^δ is as defined in Table 1.

There are similar formulas for $g_{a,b,\gamma,\eta}^{i,j,i',j'}$, where L_m^δ and $L_{m'}^{\delta'}$ are replaced by J_m^δ and $J_{m'}^{\delta'}$. In Table 1, the cohomology classes ζ_1 and ζ_2 are equal to $E^1 \# E^2$ and $E^2 \# E^1$. Moreover, $\xi_i := \Phi_2^*(\zeta_i)$. The table contains evaluations of L_m^δ and J_m^δ at some elements of $\text{im}(\#) \cap (\Phi_2)_*(\text{im}(\#))$. The evaluations of Table 1 show that the only possible identities among the functionals

$$L_i^\delta \quad \text{and} \quad J_i^\delta \quad \text{for } 1 \leq i \leq 4$$

on V are the identities of the elements in the pairs (L_1^2, J_1^2) and (L_2^{-2}, J_2^{-2}) . Therefore, Lemma 4.86 can be used to prove the first part of the proposition. The second part is a

consequence of the remaining information in (4.87), the identity (4.74) and rationality of polynomial invariants. \square

Proposition 4.88 *The constant number \hbar_3 is nonzero.*

Proof Recall that the 4-manifold $X(m, n)$ is a branched double cover of $W(m, n)$ which is the blowup of $\mathbb{C}P^1 \times \mathbb{C}P^1$ at $4mn$ points. Suppose $\pi: X(m, n) \rightarrow W(m, n)$ is the covering map. Since π does not contract any curve, the pullback of any ample divisor is still ample by the Nakai–Moishezon criterion and projection formula [43]. In particular, the divisor

$$\pi^* \left(\{p\} \times \mathbb{C}P^1 + \mathbb{C}P^1 \times \{q\} - \sum_i E_i \right) = f_{n-1} + f_{m-1} - \sum_{i=1}^{4mn} \pi^*(E_i),$$

where $\{E_i\}_{1 \leq i \leq 4mn}$ is the set of exceptional classes, is ample.

Next we focus on the 4-manifold $X(3, 4)$. Suppose $w_1 = f_3$ and $w_2 = f_3 + f_2$. Then there is a holomorphic line bundle L_i on $X(3, 4)$ such that $c_1(L_i)$ is represented by w_i and w_i can be decomposed as $w_i^1 \# w_i^2$ with respect to the decomposition of $X(3, 4)$ as $E(3) \#_{f_2} E(3)$. Let S denote the ample class $f_2 + f_3 - \sum_{i=1}^{48} \pi^*(E_i)$. According to Theorem 2.19, the coefficient of t_2^k in the series $\widehat{D}_{X(3,4),w}(e^{S(2)}) \in \mathbb{C}[[t_2]]$ is positive for large values of k . Since $X(3, 4) = E(3) \#_{f_2} E(3)$ and the homology class S lies in the image of $\#$, we can use Proposition 4.84 to show that

$$\begin{aligned} &\widehat{D}_{X(3,4),w_i}(e^{S(2)}) \\ &= e^{\frac{1}{2}Q(S)} \left[\frac{1}{2} \hbar_1^2 \hbar_3 \cosh(\sqrt{3}(f_3 + 2f_2) \cdot S) + 2\hbar_2^2 \hbar_4 \cos\left(-\frac{2}{3}\pi w_i \cdot (f_3 + 2f_2)\right) \right]. \end{aligned}$$

Therefore

$$2\widehat{D}_{X(3,4),w_1}(e^{S(2)}) + \widehat{D}_{X(3,4),w_2}(e^{S(2)}) = \frac{3}{2} \hbar_1^2 \hbar_3 e^{\frac{1}{2}Q(S)} \cosh(\sqrt{3}(f_3 + 2f_2) \cdot S).$$

This implies that \hbar_3 is a positive number (and \hbar_1 is nonzero). \square

Proposition 4.89 *The power series $h_{a,b}^{g,d}(t_2, t_3)$ is zero, unless*

$$(a, b) \in \{(\pm(2g - 2), 0), (0, \pm(2g - 2))\}.$$

Furthermore, we have

$$\begin{aligned} h_{\pm(2g-2),0}^{g,d}(t_2, t_3) &= \hbar_3^{g-1} \left(\frac{2}{\hbar_1}\right)^{2g-4} e^{\pm 2\sqrt{3}t_2}, \\ h_{0,\pm(2g-2)}^{g,d}(t_2, t_3) &= \hbar_4^{g-1} \hbar_2^{4-2g} \zeta^{\pm \frac{2}{3}\pi d} e^{\pm 2\sqrt{3}it_3}. \end{aligned}$$

Proof This theorem can be proved by exploiting the diffeomorphism in (4.79) for $n=g$. Let the 2-cycle w in $B(g) \#_{f_g} B(g)$ be equal to $df'_2 + df_g$. Using Propositions 4.71 and 4.78, we can show that $N_{B(g) \#_{f_g} B(g), w}(\Gamma, \Lambda)$, for $\Gamma, \Lambda \in \mathcal{H} := \text{im}(\#)$, is equal to

$$(4.90) \quad \sum_{(a,b) \in \mathcal{C}_g \setminus \mathcal{C}_{g-1}} h_{a,b}^{g,d} (t_2, t_3) \sum c_{ij} c_{i'j'} e^{(\frac{\sqrt{3}}{2}(M_{i,i'} + M_{j,j'}) \cdot \Gamma + \frac{\sqrt{3}}{2} \mathbf{i} (M_{i,i'} - M_{j,j'}) \cdot \Lambda)},$$

where the second sum is over the pairs of basic classes (K_i, K_j) and $(K_{i'}, K_{j'})$ of the permissible triple $(B(g), df' + 2df_g, f_g)$ such that

$$(4.91) \quad \begin{aligned} \frac{1}{2}(K_i + K_j) \cdot f_g &= \frac{1}{2}(K_{i'} + K_{j'}) \cdot f_g = a, \\ \frac{1}{2}(K_i - K_j) \cdot f_g &= \frac{1}{2}(K_{i'} - K_{j'}) \cdot f_g = b, \end{aligned}$$

$$M_{i,i'} = K_i \# K_{i'} \text{ and } M_{j,j'} = K_j \# K_{j'}.$$

Recall that the 4-manifolds $B(g) \#_{f_g} B(g)$ and $B(2) \#_{f_2} \cdots \#_{f_2} B(2)$ are diffeomorphic to each other using the diffeomorphism Φ_g . Therefore, $N_{B(g) \#_{f_g} B(g), w}(\Gamma, \Lambda)$ can be also computed by regarding $B(g) \#_{f_g} B(g)$ as the fiber sum of g copies of $B(2)$ along surfaces of genus 2. In particular, Propositions 4.81 and 4.84 allow us to obtain the following explicit form for $N_{B(g) \#_{f_g} B(g), w}(\Gamma, \Lambda)$:

$$(4.92) \quad \begin{aligned} \hbar_3^{g-1} \left(\frac{1}{36}\right)^g e^{\sqrt{3}M_g \cdot \Gamma} + \hbar_3^{g-1} \left(\frac{1}{36}\right)^g e^{-\sqrt{3}M_g \cdot \Gamma} + \hbar_4^{g-1} \left(\frac{1}{9}\zeta^d\right)^g e^{\sqrt{3}\mathbf{i} M_g \cdot \Lambda} \\ + \hbar_4^{g-1} \left(\frac{1}{9}\zeta^{-d}\right)^g e^{-\sqrt{3}\mathbf{i} M_g \cdot \Lambda}, \end{aligned}$$

where $M_g = \sigma_g^1 + \sigma_g^2 + 2(g-1)f'_2$. However, this approach works for the homology classes $\Gamma, \Lambda \in \mathcal{H}'$ where \mathcal{H}' is the image of the iterated applications of $\#$ using the decomposition of $B(g) \#_{f_g} B(g)$ as the fiber sum of g copies of $B(2)$. Therefore, (4.90) and (4.92) are equal to each other for $\Gamma, \Lambda \in \mathcal{H} \cap \mathcal{H}'$. Fix $\Gamma, \Lambda \in \mathcal{H} \cap \mathcal{H}'$ and let

$$l := \{\Gamma + sf_g \mid s \in \mathbb{C}\} \quad \text{and} \quad l' := \{\Lambda + sf_g \mid s \in \mathbb{C}\}.$$

Applying Lemma 4.86 to the subspace $l \oplus l'$ of $(\mathcal{H} \cap \mathcal{H}')^{\oplus 2}$ shows that

$$\begin{aligned} h_{\pm(2g-2), 0}^{g,d} (t_2, t_3) \left(\frac{1}{36}\right)^g \left(\frac{1}{2}\hbar_1\right)^{2g-4} e^{\pm\sqrt{3}M'_g \cdot \Gamma} &= \hbar_3^{g-1} \left(\frac{1}{36}\right)^g e^{\pm\sqrt{3}M_g \cdot \Gamma}, \\ h_{0, \pm(2g-2)}^{g,d} (t_2, t_3) \left(\frac{1}{9}\right)^g \hbar_2^{2g-4} \zeta^{\mp(2g-2)d} e^{\pm\sqrt{3}\mathbf{i} M'_g \cdot \Lambda} &= \hbar_4^{g-1} \left(\frac{1}{9}\zeta^{\pm d}\right)^g e^{\pm\sqrt{3}\mathbf{i} M_g \cdot \Lambda}, \end{aligned}$$

where $M'_g := (g-2)f'_2 + E_2^1 + \cdots + E_2^g$. Since \hbar_1 and \hbar_2 are nonzero (work in progress), the above identities prove the second part of the proposition. If $(a, b) \notin \{(\pm(2g-2), 0), (0, \pm(2g-2))\}$, then the same argument shows that

$$h_{a,b}^{g,d} (t_2, t_3) \sum c_{ij} d_{i'j'} e^{(\frac{\sqrt{3}}{2}(M_{i,i'} + M_{j,j'}) \cdot \Gamma + \frac{\sqrt{3}}{2} \mathbf{i} (M_{i,i'} - M_{j,j'}) \cdot \Lambda)} = 0$$

for $\Gamma, \Lambda \in \mathcal{H} \cap \mathcal{H}'$. This sum is over the pairs that satisfy (4.91). Another application of Lemma 4.86 for the homology classes

$$\Gamma = s(E_2^1 \pm \dots \pm E_2^g) \quad \text{and} \quad \Lambda = s(E_2^1 \pm \dots \pm E_2^g)$$

in $\mathcal{H} \cap \mathcal{H}'$ shows that $h_{a,b}^{g,d}(t_2, t_3)$ has to be zero. □

The following theorem summarizes our results in this subsection:

Theorem 4.93 *Suppose (X, w, Σ) and (X', w', Σ) are two permissible triples and the genus of Σ is at least 2. Then the triple $(X \#_{\Sigma} X', w \# w', \Sigma)$ is permissible with respect to the image of the map $\#: \mathcal{K} \rightarrow H_2(X \#_{\Sigma} X')$. The basic classes for $(X \#_{\Sigma} X', w \# w', \Sigma)$ are*

$$(4.94) \quad M_{i,i'}^{\gamma} = K_i \# L_{i'} + 2\gamma\Sigma,$$

where K_i and $L_{i'}$ are basic classes of (X, w, Σ) and (X', w', Σ) , $K_i \cdot \Sigma = L_{i'} \cdot \Sigma$, and

$$(K_i \cdot \Sigma, \gamma) \in \{(2g - 2, 1), (-(2g - 2), -1)\}.$$

For a pair of basic classes $M_{i,i'}^{\gamma} = K_i \# L_{i'} + 2\gamma\Sigma$ and $M_{j,j'}^{\eta} = K_j \# L_{j'} + 2\eta\Sigma$, let

$$\begin{aligned} \frac{1}{2}(K_i + K_j) \cdot \Sigma &= \frac{1}{2}(L_{i'} + L_{j'}) \cdot \Sigma = a, \\ \frac{1}{2}(K_i - K_j) \cdot \Sigma &= \frac{1}{2}(L_{i'} - L_{j'}) \cdot \Sigma = b. \end{aligned}$$

Then the coefficient associated to this pair is equal to $c_{i,j} d_{i',j'} h_{a,b,\gamma,\eta}^{g,d}$, where $c_{i,j}$ is the coefficient associated to (K_i, K_j) for the triple (X, w, Σ) , $d_{i',j'}$ is the coefficient associated to $(L_{i'}, L_{j'})$ for the triple (X', w', Σ) , and $h_{a,b,\gamma,\eta}^{g,d}$ is nonzero in the cases

$$\begin{aligned} h_{(2g-2),0,1,0}^{g,d} &= h_{-(2g-2),0,-1,0}^{g,d} = \hbar_3^{g-1} \left(\frac{2}{\hbar_1} \right)^{2g-4}, \\ h_{0,(2g-2),0,1}^{g,d} &= \frac{\hbar_4^{g-1}}{\hbar_2^{2g-4}} \zeta^d \quad \text{and} \quad h_{0, -(2g-2),0,-1}^{g,d} = \frac{\hbar_4^{g-1}}{\hbar_2^{2g-4}} \zeta^{-d}. \end{aligned}$$

5 Sutured Floer homology

5.1 Eigenvectors

For arbitrary N , we introduce a set of generators of the algebra $\mathbb{V}_{g,d}^N$ in Corollary 3.22. In particular, in the special case that $N = 3$ and $d \equiv 1$ or $2 \pmod 3$, we have the

following generators of $\mathbb{V}_{g,d}^3$ (which will be denoted by $\mathbb{V}_{g,d}$ from now on):

$$(5.1) \quad \begin{aligned} \epsilon &= D_{\Delta_g, \delta_{g,d} + \Sigma}(1), & \mathfrak{N}_r &= D_{\Delta_g, \delta_{g,d}}(a_r), \\ o_r^j &= D_{\Delta_g, \delta_{g,d}}(l_{(r)}^j), & \rho_r &= D_{\Delta_g, \delta_{g,d}}(\Sigma(r)), \end{aligned}$$

where $r = 2, 3$ and $\{l^j\}_{1 \leq j \leq 2g}$ is a set of generators for $H_1(\Sigma, \mathbb{Z})$. If y is one of the above elements, then the product $m(\cdot, y)$ defines an operator on $\mathbb{V}_{g,d}$, which is still denoted by y . Recall that there is also a pairing on $\mathbb{V}_{g,d}$, which is denoted by $\langle \cdot, \cdot \rangle$.

The operator ϵ is equal to $I_*(Y_g \times [0, 1], \gamma_{g,d} \times [0, 1] + \Sigma, 1)$ and the remaining operators can be described as

$$(5.2) \quad I_*(Y_g \times [0, 1], \gamma_{g,d} \times [0, 1], q),$$

where $q = a_r, l_{(r)}^j$ or $\Sigma(r)$. This alternative description allows us to extend the definition of these operators to arbitrary admissible pairs. Suppose (Y, γ) is a 3–admissible pair, and Σ is an embedded surface of genus g in Y . We also assume that an integral basis $\{l^j\}_{1 \leq j \leq 2g}$ for $H_1(\Sigma)$ is fixed. By replacing $(Y_g, \gamma_{g,d})$ with (Y, γ) , we can define analogues of the operators $\epsilon, \mathfrak{N}_r, o_r^j$ and ρ_r on $I_*(Y, \gamma)$. These operators on $I_*(Y, \gamma)$ are denoted by $\epsilon(\Sigma), \mathfrak{N}_r, o_r^j(\Sigma)$ and $\rho_r(\Sigma)$. When the choice of Σ is clear from our discussion, we drop Σ from our notation for $\epsilon(\Sigma), o_r^j(\Sigma)$ and $\rho_r(\Sigma)$.

Definition 5.3 An element $v \in \mathbb{V}_{g,d}$ is called an *exhaustive eigenvector* if it is a simultaneous eigenvector of the action of the operators in (5.1). An *exhaustive eigenspace* is the set of all exhaustive eigenvectors which have the same eigenvalues with respect to these operators. An exhaustive eigenvector v is called *nondegenerate* if $\langle v, v \rangle \neq 0$.

Remark 5.4 Since $(o_r^j)^2 = 0$, the only eigenvalue of this operator is zero.

Suppose (X, w, Σ) is a permissible triple, Σ is a surface of genus g , and $w \cdot \Sigma = d$. For a pair of basic classes (K_i, K_j) of this triple, let $c_{i,j}$ denote the associated coefficient. Suppose also for a fixed $\lambda = (\alpha, \beta) \in \mathcal{C}_g$ that

$$(5.5) \quad \sum_{(K_i, K_j)} c_{i,j} \neq 0,$$

where the sum is over all pairs of basic classes (K_i, K_j) that satisfy the following equalities for $(a, b) = (\alpha, \beta)$:

$$(5.6) \quad \frac{1}{2}(K_i + K_j) \cdot \Sigma = a \quad \text{and} \quad \frac{1}{2}(K_i - K_j) \cdot \Sigma = b.$$

Recall that P_λ is the polynomial that satisfies (4.64). Suppose $Q_\lambda \in \mathbb{C}[x, y, z]$ is defined as the evaluation of $P_{(a,b)}$ at $t_2 = t_3 = 0$ and consider the following element of $\mathbb{V}_{g,d}$:

$$v_{(\alpha,\beta)} := D_{X^\circ, w^\circ, \Sigma}(Q_\lambda(a_2, \Sigma_{(2)}, \Sigma_{(3)})).$$

Proposition 5.7 *The element $v_{(\alpha,\beta)} \in \mathbb{V}_{g,d}$ is a nonzero exhaustive eigenvector. The eigenvalues of $v_{(\alpha,\beta)}$ with respect to the actions of ϵ , \mathfrak{N}_2 , \mathfrak{N}_3 , ρ_2 and ρ_3 are respectively equal to 1, $3\zeta^{2d\beta}$, 0, $\zeta^{d\beta} \sqrt{3}\alpha$ and $\zeta^{2d\beta} \sqrt{3}i\beta$. Furthermore, if $(\alpha, \beta) = (\pm(2g - 2), 0)$, then the eigenvector $v_{(\alpha,\beta)}$ is nondegenerate.*

Proof Lemma 4.59 can be used to show that for an arbitrary polynomial $P \in \mathbb{C}[x, y, z]$,

$$(5.8) \quad \begin{aligned} D_{X,w,\Sigma}(P(a_2, \Sigma_{(2)}, \Sigma_{(3)})) \\ = \sum_{(a,b) \in \mathcal{C}_g} \zeta^{dbd_w} P(3\zeta^{2db}, \zeta^{bd} \sqrt{3}a, \zeta^{2bd} \sqrt{3}ib) \sum_{(K_i, K_j)} c_{i,j}, \end{aligned}$$

where the inner sum is over all pairs of basic classes (K_i, K_j) that satisfy (5.6) and $c_{i,j}$ is the coefficient associated to the pair (K_i, K_j) . Functorial properties of Floer homology imply that the pairing of $v_{(\alpha,\beta)}$ and $D_{\Delta_g, \delta_{g,d}}(1)$ is equal to $D_{X,w,\Sigma}(Q_\lambda(a_2, \Sigma_{(2)}, \Sigma_{(3)}))$, which is nonzero. Therefore, $v_{(\alpha,\beta)}$ is a nonzero vector.

Suppose y is one of the elements of $\mathbb{V}_{g,d}^3$ in (5.1). Proposition 3.19 and the nondegeneracy of the pairing on $\mathbb{V}_{g,d}$ (Proposition 3.23) imply that $v_{(\alpha,\beta)}$ is an eigenvector of $m(\cdot, y)$ with eigenvalue c_y if and only if

$$(5.9) \quad \langle m(v_{(\alpha,\beta)}, y), D_{\Delta_g, \delta_{g,d} + i\Sigma}(z) \rangle = c_y \langle v_{(\alpha,\beta)}, D_{\Delta_g, \delta_{g,d} + i\Sigma}(z) \rangle$$

for any $z \in \mathbb{A}_g^3$ and $0 \leq i \leq 2$. Another application of the functorial properties of Floer homology shows that both of the expressions in (5.9) can be written as $U(3)$ -invariants of (X, w) . In particular, (5.8) shows that $v_{(\alpha,\beta)}$ is an eigenvector of ϵ , \mathfrak{N}_2 , ρ_2 and ρ_3 . The vector $v_{(\alpha,\beta)}$ is in the kernel of the operators \mathfrak{N}_3 and σ_r^j because X has w -simple type and $b_1(X) = 0$. The pairing $\langle v_{(\alpha,\beta)}, v_{(\alpha,\beta)} \rangle$ can be also computed by Theorem 4.93. Using Proposition 4.88, this number is nonzero for $(\alpha, \beta) = (\pm(2g - 2), 0)$. □

Example 4.53 gives a permissible triple such that the condition in (5.5) is satisfied for any $\lambda \in \mathcal{C}_g$. Therefore, for each λ there is an exhaustive eigenvector in $\mathbb{V}_{g,d}$. The condition in (5.5) is not very essential in constructing such an eigenvector. This

condition is used to show that $v_{(\alpha,\beta)}$ is a nonzero element of $\mathbb{V}_{g,d}$. It is possible to replace $v_{(\alpha,\beta)}$ with the element

$$v'_{(\alpha,\beta)} := D_{X^\circ, w^\circ, \Sigma}(Q_\lambda(a_2, \Sigma_{(2)}, \Sigma_{(3)})z),$$

where $z \in \mathbb{A}(X^\circ)^{\otimes 2}$. If the triple (X, w, Σ) has at least one pair of basic classes (K_i, K_j) which satisfy (5.6) for $(a, b) = (\alpha, \beta)$, then z can be chosen such that the above element of $\mathbb{V}_{g,d}$ is nonzero.

Proposition 5.10 *An exhaustive eigenspace is 1–dimensional.*

Proof Suppose $V \subset \mathbb{V}_{g,d}$ is an exhaustive eigenspace, and s_1, s_2, s_3, s_4 and s_5 are respectively the corresponding eigenvalues of $\epsilon, \mathfrak{N}_2, \mathfrak{N}_3, \rho_2$ and ρ_3 . Suppose also that $\mathfrak{J} \subset \mathbb{V}_{g,d}$ is the ideal generated by the elements of the set

$$(5.11) \quad G := \{\epsilon - s_1, \mathfrak{N}_2 - s_2, \mathfrak{N}_3 - s_3, \rho_2 - s_4, \rho_3 - s_5\} \cup \{o_r^j \mid 1 \leq j \leq 2g, 2 \leq r \leq 3\}.$$

Then an element of \mathfrak{J} is the sum of the elements of the form $m(x, y)$ with $x \in G$ and $y \in \mathbb{V}_{g,d}$. For any $v \in V$,

$$\langle v, m(x, y) \rangle = \langle m(v, x), y \rangle = 0.$$

Therefore, V is orthogonal to \mathfrak{J} . Since $\mathbb{V}_{g,d}/\mathfrak{J}$ is a 1–dimensional vector space and the pairing is nondegenerate, the dimension of the vector space V is at most 1. \square

Lemma 5.12 *Suppose that $v \in \mathbb{V}_{g,d}$ is a nondegenerate exhaustive eigenvector, and s_1, s_2, s_3, s_4 and s_5 are respectively the corresponding eigenvalues of $\epsilon, \mathfrak{N}_2, \mathfrak{N}_3, \rho_2$ and ρ_3 . Then the space*

$H := \ker_{\text{gen}}(\epsilon - s_1) \cap \ker_{\text{gen}}(\mathfrak{N}_2 - s_2) \cap \ker_{\text{gen}}(\mathfrak{N}_3 - s_3) \cap \ker_{\text{gen}}(\rho_2 - s_4) \cap \ker_{\text{gen}}(\rho_3 - s_5)$ *is 1–dimensional. Here $\ker_{\text{gen}}(T)$, for an operator T , is the union of the kernel of the operators T^k for all values of k .*

Proof Suppose the claim does not hold and $v' \in H$ is a vector which is linearly independent of v . Let G be defined as in (5.11). All the operators involved in the definition of H have even degree with respect to the $\mathbb{Z}/2\mathbb{Z}$ –grading of $\mathbb{V}_{g,d}$ induced by the $\mathbb{Z}/12\mathbb{Z}$ –grading. Therefore, H can be decomposed as $H_0 \oplus H_1$ with respect to the $\mathbb{Z}/2\mathbb{Z}$ –grading of $\mathbb{V}_{g,d}$, and we may assume that $v' \in H_i$ for $i = 0$ or 1 . By Proposition 5.10, there is an $x_0 \in G$ such that $m(v', x_0) \neq 0$. Since the restrictions of the elements of G on H are nilpotent, we can also assume, without loss of generality,

that the product of $m(v', x_0)$ and any element $x \in G$ is zero. Therefore, $m(v', x_0) = cv$ for a nonzero complex number c . This implies that

$$\langle m(v', x_0), m(v', x_0) \rangle = \langle cv, cv \rangle \neq 0.$$

On the other hand, we have

$$\langle m(v', x_0), m(v', x_0) \rangle = \pm \langle m(m(v', x_0), x_0), v' \rangle = 0,$$

which is a contradiction. □

Proposition 5.13 [56, Proposition 7.2] *Suppose (Y, γ) is a 3–admissible pair and Σ is an embedded surface in Y of genus g such that $\gamma \cdot \Sigma \equiv d \pmod 3$. If $(s_1, s_2, s_3, s_4, s_5)$ is a simultaneous eigenvalue of the operators $(\epsilon(\Sigma), \mathfrak{N}_2, \mathfrak{N}_3, \rho_2(\Sigma), \rho_3(\Sigma))$, then it is also a simultaneous eigenvalue of the operators $(\epsilon, \mathfrak{N}_2, \mathfrak{N}_3, \rho_2, \rho_3)$ acting on $\mathbb{V}_{g,d}$.*

By Proposition 5.7, $v_{2g-2,0}$ is a nondegenerate exhaustive eigenvector of $\mathbb{V}_{g,d}$. Suppose $s_1^g, s_2^g, s_3^g, s_4^g$ and s_5^g denote the corresponding eigenvalues of $\epsilon, \mathfrak{N}_2, \mathfrak{N}_3, \rho_2$ and ρ_3 . Then $s_1^g = 1, s_2^g = 3, s_3^g = 0, s_4^g = \sqrt{3}(2g-2)$ and $s_5^g = 0$. Following [56], we can define a variation of instanton Floer homology:

Definition 5.14 Suppose (Y, γ) is a 3–admissible pair and Σ is an embedded surface in Y of genus g such that $\gamma \cdot \Sigma \equiv d \pmod 3$. Then $I_*(Y, \gamma | \Sigma)$ is defined as

$$\ker_{\text{gen}}(\epsilon(\Sigma) - s_1^g) \cap \ker_{\text{gen}}(\mathfrak{N}_2 - s_2^g) \cap \ker_{\text{gen}}(\mathfrak{N}_3 - s_3^g) \cap \ker_{\text{gen}}(\rho_2(\Sigma) - s_4^g) \cap \ker_{\text{gen}}(\rho_3(\Sigma) - s_5^g).$$

In this definition, we allow Σ to have more than one connected component. In that case, each connected component Σ' of Σ is required to have genus g and $\gamma \cdot \Sigma' \equiv d \pmod 3$. In the definition of $I_*(Y, \gamma | \Sigma)$, we include the operators $\epsilon(\Sigma'), \rho_2(\Sigma') - s_4^g$ and $\rho_3(\Sigma') - s_5^g$ for each connected component Σ' of Σ .

Remark 5.15 In the case that $g = 1$, the actions of the operators \mathfrak{N}_3, ρ_2 and ρ_3 on $\mathbb{V}_{1,d}$ are trivial, the operator \mathfrak{N}_2 is equal to $3\epsilon^{-1}$ and $\epsilon^3 = 1$. Therefore, similar relationships hold among the operators $\epsilon(\Sigma), \mathfrak{N}_2, \mathfrak{N}_3, \rho_2(\Sigma), \rho_3(\Sigma)$ acting on $I_*(Y, \gamma)$, where Σ is a genus one surface in Y . This can be verified in a similar way as in Proposition 5.13. (See [56, Proposition 7.2].) Therefore, $I_*(Y, \gamma | \Sigma)$, for a genus one surface Σ , is simply equal to $\ker(\mathfrak{N}_2 - 3)$.

This variant of instanton Floer homology is also functorial with respect to cobordisms. Suppose $(W, w): (Y_0, \gamma_0) \rightarrow (Y_1, \gamma_1)$ is a cobordism of admissible pairs, $z \in \mathbb{A}(W)^{\otimes 2}$, Σ_i is an embedded oriented and connected surface in Y_i such that $\Sigma_i \cdot \gamma_i \equiv d \pmod 3$, and Σ_0 and Σ_1 induce the same homology classes of W . More generally, if Σ_1 is disconnected, then each connected component of Σ_1 is required to be homologous to a connected component of Σ_0 inside W . Properties of instanton Floer homology discussed in Section 3.1 imply that $I_*(W, w, z)$ maps $I_*(Y_0, \gamma_0 | \Sigma_0) \subseteq I_*(Y_0, \gamma_0)$ to $I_*(Y_1, \gamma_1 | \Sigma_1) \subseteq I_*(Y_1, \gamma_1)$. Moreover, suppose (X, w) is a cobordism from an admissible pair (Y, γ) to the empty pair and $z \in \mathbb{A}(X)^{\otimes 2}$. Then the restriction of the map $D^{X,w}(z)$ gives rise to a functional on $I_*(Y, \gamma | \Sigma)$, which is denoted with the same notation.

Lemma 5.12 asserts that $I_*(Y_g, \gamma_{g,d} | \Sigma)$ is 1-dimensional. The following pairs from Section 3.2 define cobordisms from two copies of $(Y_g, \gamma_{g,d})$ to the empty pair:

$$(\Omega_g, \omega_{g,d}) \quad \text{and} \quad (\Delta_g \amalg \Delta_g, \delta_{g,d} \amalg \delta_{g,d}).$$

Therefore, they determine two functionals $D^{\Omega_g, \omega_{g,d}}(1)$ and $D^{\Delta_g, \delta_{g,d}}(1) \otimes D^{\Delta_g, \delta_{g,d}}(1)$ on the 1-dimensional vector space $I_*(Y_g, \gamma_{g,d} | \Sigma) \otimes I_*(Y_g, \gamma_{g,d} | \Sigma)$. The nondegeneracy of the exhaustive eigenvector involved in the definition of $I_*(Y_g, \gamma_{g,d} | \Sigma)$ implies that the former functional is nonzero. Therefore, we have the following lemma which provides the distinguishing property of working with a nondegenerate exhaustive eigenspace for us:

Lemma 5.16 *The map*

$$D^{\Delta_g, \delta_{g,d}}(1) \otimes D^{\Delta_g, \delta_{g,d}}(1): I_*(Y_g, \gamma_{g,d} | \Sigma) \otimes I_*(Y_g, \gamma_{g,d} | \Sigma) \rightarrow \mathbb{C}$$

is a multiple of $D^{\Omega_g, \omega_{g,d}}(1): I_*(Y_g, \gamma_{g,d} | \Sigma) \otimes I_*(Y_g, \gamma_{g,d} | \Sigma) \rightarrow \mathbb{C}$.

5.2 Excision and sutured manifolds invariants

Suppose R_1 and R_2 are two embedded surfaces of genus $g \geq 1$ in a 3-manifold Y . Suppose also there is a 1-cycle γ in Y such that $\gamma \cdot R_1 = \gamma \cdot R_2$. We also assume γ intersects R_1 and R_2 transversally, and all the intersection points have the same signs. Fix a diffeomorphism $\phi: R_1 \rightarrow R_2$ such that ϕ maps $\gamma \cap R_1$ to $\gamma \cap R_2$. We cut Y along the surfaces R_1 and R_2 , and then identify the four boundary components of the resulting 3-manifold using ϕ so that the final 3-manifold, \tilde{Y} , is an oriented closed 3-manifold with embedded surfaces \tilde{R}_1 and \tilde{R}_2 . Our assumption on ϕ implies that γ




Figure 5: The saddle cobordism \mathcal{P} : the union of the two vertical boundary components on the left (respectively, right) is denoted by $\partial_l^v \mathcal{P}$ (respectively, $\partial_r^v \mathcal{P}$).

determines a 1-cycle $\tilde{\gamma}$ in \tilde{Y} . We will also write R (respectively, \tilde{R}) for the union $R_1 \cup R_2$ (respectively, $\tilde{R}_1 \cup \tilde{R}_2$). Now we are ready to state our excision theorem:

Theorem 5.17 *The following Floer homology groups are isomorphic:*

$$I_*(Y, \gamma|R) = I_*(\tilde{Y}, \tilde{\gamma}|\tilde{R}).$$

Proof This theorem is the analogue of excision theorem for $U(2)$ -instanton Floer homology. The $U(2)$ version of the excision theorem is proved in [28] for $g = 1$ (see also [8]) and in [56] for higher values of g . We follow the same strategy as in the $U(2)$ case to prove the theorem. In particular, the isomorphism between $I_*(Y, \gamma|R)$ and $I_*(\tilde{Y}, \tilde{\gamma}|\tilde{R})$ is induced by a cobordism of pairs $(W, w): (Y, \gamma) \rightarrow (\tilde{Y}, \tilde{\gamma})$. Let Y° be the complement of a regular neighborhood of R in Y . Then the cobordism W is the result of gluing $[0, 1] \times Y^\circ$ to $\mathcal{P} \times R_1$ where \mathcal{P} is the saddle cobordism in Figure 5. The boundary of the 3-manifold Y° is equal to $R_1 \cup \bar{R}_1 \cup R_2 \cup \bar{R}_2$. Then $[0, 1] \times (R_1 \cup \bar{R}_1) \subset [0, 1] \times Y^\circ$ is glued to $\partial_l^v \mathcal{P} \times R_1$ by the identity map and $[0, 1] \times (R_2 \cup \bar{R}_2) \subset [0, 1] \times Y^\circ$ is glued to $\partial_r^v \mathcal{P} \times R_1$ by the map ϕ . (For the definition of $\partial_l^v \mathcal{P}$ and $\partial_r^v \mathcal{P}$ see Figure 5.) The surface cobordism $w: \gamma \rightarrow \tilde{\gamma}$ is also constructed in a similar way. This surface is given by gluing one copy of \mathcal{P} for each intersection point in $\gamma \cap R_1$ to $[0, 1] \times (\gamma \cap Y^\circ)$. Reversing the cobordism (W, w) determines another cobordism $(\bar{W}, \bar{w}): (\tilde{Y}, \tilde{\gamma}) \rightarrow (Y, \gamma)$. To prove the excision theorem, we claim that

$$(5.18) \quad \begin{aligned} I_*(\bar{W}, \bar{w}) \circ I_*(W, w): I_*(Y, \gamma|R) &\rightarrow I_*(Y, \gamma|R), \\ I_*(\bar{W}, \bar{w}) \circ I_*(W, w): I_*(\tilde{Y}, \tilde{\gamma}|\tilde{R}) &\rightarrow I_*(\tilde{Y}, \tilde{\gamma}|\tilde{R}) \end{aligned}$$

are nonzero multiples of the identity map. In the composite cobordism $\bar{W} \circ W: Y \rightarrow Y$, a copy of $\mathcal{P} \times R_1$ is glued to another copy of $\mathcal{P} \times R_1$ with the reverse orientation.

In this part of $\overline{W} \circ W$, the union of the two copies of $\delta \times R_1$ gives rise to a copy of $Y_g := S^1 \times R_1$. The intersection of $\overline{w} \circ w$ with $S^1 \times R_1$ produces a copy of $\gamma_{g,d}$. According to Lemma 5.16 and functoriality of I_* , replacing a neighborhood of $S^1 \times R_1$ with $(\Delta_g \amalg \Delta_g, \delta_{g,d} \amalg \delta_{g,d})$ does not change the map $I_*(\overline{W} \circ W, \overline{w} \circ w)$, up to multiplication by a nonzero constant number. But the resulting cobordism of the pair is the product cobordism $([0, 1] \times Y, [0, 1] \times \gamma)$. Therefore, the first map in (5.18) is a nonzero multiple of the identity map. Replacing δ with δ' and using the same argument proves a similar result for the second map in (5.18). \square

The following proposition is the analogue of Corollary 4.8 in [56] and can be proved in a similar way using the excision theorem:

Proposition 5.19 *Let Y be a 3-manifold, γ be a 1-cycle in Y , and $R \subset Y$ be a connected surface of genus $g \geq 1$ such that $\gamma \cdot R \not\equiv 0 \pmod 3$ and the intersection points of $\gamma \cap R$ are transversal and have the same signs. Let \tilde{Y} be the 3-manifold obtained by cutting Y along R and regluing by an orientation-preserving diffeomorphism $\phi: R \rightarrow R$. Suppose ϕ maps $R \cap \gamma$ to $R \cap \gamma$, and $\tilde{\gamma}$ and \tilde{R} are the induced 1-cycle and the embedded surface in \tilde{Y} . Then*

$$I_*(Y, \gamma | R) \cong I_*(\tilde{Y}, \tilde{\gamma} | \tilde{R}).$$

Now we can define invariants for balanced sutured manifolds almost verbatim from [56]. Firstly we recall the definition of balanced sutured manifolds (see [35; 46; 56]):

Definition 5.20 *A sutured manifold (M, α) consists of an oriented 3-manifold M , an oriented closed 1-manifold α in ∂M and a decomposition of ∂M as*

$$(5.21) \quad \partial M = R_+ \cup A(\alpha) \cup R_-.$$

Each connected component of α is called a *suture* and $A(\alpha)$ is the closure of a tubular neighborhood of α . The spaces R_+ and R_- are disjoint and each of them is a union of some of the connected components of $\overline{\partial M \setminus A(\alpha)}$. In particular, each component of ∂R_+ and ∂R_- is a parallel copy of a suture. Suppose R_+ and R_- are oriented with the outward-normal-first convention. Similarly, each component of ∂R_+ (respectively, ∂R_-) inherits an orientation from R_+ (respectively, R_-). This orientation is required to agree (respectively, disagree) with the orientation of the corresponding suture. The sutured manifold (M, α) is *balanced* if neither M nor R_\pm has closed components, and $\chi(R_+) = \chi(R_-)$.

Note that the required conditions on R_+ , R_- and $A(\alpha)$ imply that the decomposition (5.21) is unique.

Example 5.22 Suppose $F_{g,k}$ is a surface of genus g with $k \geq 1$ boundary components which is not the 2-dimensional disc. Then $([-1, 1] \times F_{g,k}, \partial F_{g,k} \times \{0\})$ determines a balanced sutured manifold. The decomposition of the boundary of this product sutured manifold is given by

$$(5.23) \quad R_+ = \{1\} \times F_{g,k}, \quad A(\alpha) = [-1, 1] \times \partial F_{g,k}, \quad R_- = \{-1\} \times F_{g,k}.$$

Example 5.24 Suppose Y is a 3-manifold and $K \subset Y$ is a knot. Let $M(K)$ be the complement of a regular neighborhood of K in Y , and $\alpha(K)$ be the union of two oppositely oriented meridional curves on the boundary of $M(K)$. Then $(M(K), \alpha(K))$ determines a balanced sutured manifold. The manifolds R_+ and R_- are homeomorphic to $[0, 1] \times S^1$.

Example 5.25 Suppose that K is a null-homologous knot in Y and S is a Seifert surface for K . We can also associate a sutured manifold $(N(S), \alpha(S))$ to S . The 3-manifold $N(S)$ is defined to be $Y \setminus ((-1, 1) \times \text{int}(S))$, where $(-1, 1) \times S$ is a regular neighborhood of S in Y . The only suture $\alpha(S)$ of this sutured manifold is defined to be $\{0\} \times \partial S$.

Let (M, α) be a balanced sutured manifold such that the number of sutures is greater than one and neither R_+ nor R_- is a union of 2-dimensional discs. We attach the product sutured manifold $([-1, 1] \times F_{0,k}, \{0\} \times \partial F_{0,k})$ to (M, α) where k is the number of sutures of (M, α) . More precisely, we glue M to $[-1, 1] \times F_{0,k}$ by identifying $A(\alpha)$ with $[-1, 1] \times \partial F_{0,k}$ using an orientation-reversing map. The resulting manifold is oriented and has two boundary components,

$$\bar{R}_+ = R_+ \cup \{1\} \times F_{0,k} \quad \text{and} \quad \bar{R}_- = R_- \cup \{-1\} \times F_{0,k}.$$

Since (M, α) is balanced, the oriented surfaces \bar{R}_+ and \bar{R}_- have the same positive genus. We choose an orientation-reversing diffeomorphism $\phi: \bar{R}_+ \rightarrow \bar{R}_-$. Identifying \bar{R}_+ and \bar{R}_- using the map ϕ determines a closed 3-manifold Y_ϕ , which is called a closure of the sutured manifold (M, α) . The 3-manifold Y_ϕ only depends on (M, α) and the choice of the diffeomorphism ϕ . The surface \bar{R}_+ also induces an oriented surface in Y_ϕ , which is denoted by \bar{R} . We also fix a point y on $F_{0,k}$ and require that ϕ map $(1, y) \in \bar{R}_+$ to $(-1, y) \in \bar{R}_-$. Therefore, the path $[-1, 1] \times \{y\} \subset [-1, 1] \times F_{0,k}$ induces a 1-cycle γ_ϕ in Y_ϕ .

Definition 5.26 The *sutured instanton homology* of the sutured manifold (M, α) is

$$(5.27) \quad \text{SHI}_*(M, \alpha) := \text{I}_*(Y_\phi, \gamma_\phi | \bar{R}).$$

If (M, α) has one suture or R_+ (or equivalently R_-) is a union of discs, we replace $F_{0,k}$ with $F_{1,k}$ in the definition of (Y, γ, \bar{R}) and then extend sutured instanton homology to these sutured manifolds using (5.27).

Proposition 5.19 implies that sutured instanton homology SHI_* is well defined. Notice that our choices imply that the surface \bar{R} in (5.27) has genus at least one. In particular, the required assumption for **Proposition 5.19** is satisfied.

Remark 5.28 In the definition of the closure of a sutured manifold (M, α) , we can replace $F_{0,k}$ with $F_{g,k}$ for an arbitrary g . Then for each choice of g , we can define a sutured Floer homology group as above. Using our excision theorem and the method of [56], we can show that the rank of these sutured Floer homology groups is nondecreasing in g . We expect that these sutured Floer homology groups for various choices of g are isomorphic to each other. However, proving this seems to need a further study of the algebra $\mathbb{V}_{g,1}$. For example, if **Conjecture 7.3** holds, then all of the above versions of sutured Floer homology of (M, α) are isomorphic. We hope to come back to this issue elsewhere.

5.3 Instanton knot homology

Definition 5.29 Given a knot K in a 3-manifold Y , let $(M(K), \alpha(K))$ be the sutured manifold of **Example 5.24**. The *U(3)-knot homology* of K , denoted by $\text{KHI}_*(Y, K)$, is defined to be $\text{SHI}_*(M(K), \alpha(K))$.

As explained in [56, Lemma 5.2], a closure of $(M(K), \alpha(K))$ can be described as follows. Suppose F is a genus 1 surface with one boundary component, and $c, c' \subset F$ are two oriented nonseparating simple closed curves which intersect in exactly one point and $c \cdot c' = 1$. Let $Z(K)$ be the result of gluing the knot complement $M(K)$ to the product 3-manifold $F \times S^1$ such that the meridian of K is identified with $\{\text{point}\} \times S^1$ and the longitude of K is identified with $\partial F \times \{\text{point}\}$. Suppose also $\gamma(K) \subset Z(K)$ is the 1-cycle given by $c \times \{\text{point}\} \subset F \times S^1$. Then $Z(K)$ is a closure of the sutured manifold $(M(K), \alpha(K))$ and $\gamma(K)$ is the corresponding 1-cycle. The embedded surface \bar{R} is also given by the torus $T = c' \times S^1 \subset F \times S^1$. Consequently,

$$(5.30) \quad \text{KHI}_*(Y, K) = \text{I}_*(Z(K), \gamma(K) | T).$$

Next, we characterize the set of the critical points of the Chern–Simons functional associated to the pair $(Z(K), \gamma(K))$:

Proposition 5.31 *For a knot K in a 3–manifold Y , the set of the critical points of the Chern–Simons functional associated to the admissible pair $(Z(K), \gamma(K))$ is a 3–sheeted covering space of*

$$(5.32) \quad \mathcal{R} = \left\{ \rho: \pi_1(Y \setminus K) \rightarrow \text{SU}(3) \mid \rho(\mu) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \zeta & 0 \\ 0 & 0 & \zeta^2 \end{bmatrix} \right\}.$$

Recall that μ is a meridian of K and $\zeta = e^{2\pi i/3}$.

Proof The set of the critical points of the Chern–Simons functional is given by the conjugacy classes of representations $\rho: \pi_1(Z(K) \setminus \gamma(K)) \rightarrow \text{SU}(3)$ such that a meridian of the closed curve $\gamma(K)$ is mapped to ζ . We fix a base point for $Z(K)$ on the torus $c' \times S^1 \subset F \times S^1$. Suppose $J_1 = \rho(c' \times \{\text{point}\})$ and $J_2 = \{\text{point}\} \times S^1$. Since $c' \times S^1$ intersects c in one point, we can assume

$$(5.33) \quad [J_2, J_1] = \zeta \cdot \text{id}.$$

This implies that J_2 and $\zeta^{-1}J_2$ are conjugate to each other. Thus there is a unique representative for the conjugacy class of ρ such that

$$J_1 = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \quad \text{and} \quad J_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \zeta & 0 \\ 0 & 0 & \zeta^2 \end{bmatrix}.$$

Therefore, the conjugacy class of the representation $\rho|_{\pi_1(F \times S^1 \setminus \gamma(K))}$ is uniquely determined by $J_3 \in \text{SU}(3)$, which is equal to the image of ρ for a parallel copy of c . Since J_3 has to commute with J_2 it is a diagonal matrix. Therefore, the restriction of the above representative of ρ to the knot group $\pi_1(M(K))$ determines an element of \mathcal{R} . Furthermore, ρ maps the longitude of K to $[J_3, J_1]$. Now the claim can be easily verified, because the map that sends a diagonal matrix J_3 to $[J_3, J_1]$ is 3-to-1. \square

Corollary 5.34 *Suppose K is a knot in a homology sphere Y . If $\dim(\text{KHI}(Y, K)) > 1$, then there exists a nonabelian representation $\rho: \pi_1(Y \setminus K) \rightarrow \text{SU}(3)$ such that the image of the meridian is conjugate to*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \zeta & 0 \\ 0 & 0 & \zeta^2 \end{bmatrix}.$$

Proof Suppose there is not a representation with this property. Then the only critical points of the Chern–Simons functional of $(Z(K), \gamma(K))$ are the three flat connections induced by the abelian representation in (5.32). Since these critical points are non-degenerate,⁸ $I_*(Z(K), \gamma(K))$ is the homology of a chain complex which has three generators. The order-three map ϵ has degree 4 with respect to the $\mathbb{Z}/12\mathbb{Z}$ grading. Thus the three eigenspaces of this operator have the same dimensions. Therefore, $\ker(\epsilon - 1)$ has to be at most 1–dimensional, which is a contradiction. \square

Proposition 5.35 *Let K be a null-homologous knot in Y and S be a Seifert surface of genus $g \geq 1$ for K . Then $\dim(\text{KHI}(Y, K)) \geq 2 \dim(\text{SHI}_*(N(S), \alpha(S)))$, where $(N(S), \alpha(S))$ is the sutured manifold of Example 5.25.*

Proof According to (5.30), the $U(3)$ –knot homology is equal to $I_*(Z(K), \gamma(K)|T)$. In order to form a closure of $(N(S), \alpha(S))$ and to define $\text{SHI}_*(N(S), \alpha(S))$, we first glue $[-1, 1] \times F_{1,1}$ along the suture $\alpha(S)$. In this case, \bar{R}^\pm are two copies of $S \cup F_{1,1}$. If we identify \bar{R}^\pm in the obvious way, then the resulting space is again the 3–manifold $Z(K)$. Let $\gamma(S)$ and $\bar{R}(S)$ be the resulting 1–cycle and the surface in the closure $Z(K)$. The 1–cycle $\gamma(S)$ is a copy of $S^1 \times \{\text{point}\} \subset S^1 \times F_{1,1}$. The surface $\bar{R}(S)$ has genus $g + 1$ and is given by gluing the Seifert surface S to $F_{1,1}$. Arguing as in [56, Proposition 7.9] and using our excision theorem, we can show that

$$I_*(Z(K), \gamma(K)|T) = I_*(Z(K), \gamma(K) + \gamma(S)|T)$$

and

$$I_*(Z(S), \gamma(S)|\bar{R}(S)) = I_*(Z(S), \gamma(K) + \gamma(S)|\bar{R}(S)).$$

Remark 5.15 implies that the homology group $I_*(Z(K), \gamma(K) + \gamma(S)|T)$ is equal to

$$(5.36) \quad I_*(Z(K), \gamma(K) + \gamma(S)) \cap \ker(\mathfrak{S}_2 - 3).$$

We can further decompose the vector space in (5.36) using the eigenvalues of the operators $\rho_2(\bar{R}(S))$. In particular, the vector spaces

$$(5.37) \quad I_*(Z(K), \gamma(K) + \gamma(S)) \cap \ker(\mathfrak{S}_2 - 3) \cap \ker_{\text{gen}}(\rho_2(\bar{R}(S)) - 2\sqrt{3}g)$$

and

$$(5.38) \quad I_*(Z(K), \gamma(K) + \gamma(S)) \cap \ker(\mathfrak{S}_2 - 3) \cap \ker_{\text{gen}}(\rho_2(\bar{R}(S)) + 2\sqrt{3}g)$$

⁸This is a consequence of the fact that the Alexander polynomial of a knot K in an integral homology sphere does not have a root which is a third root of unity.

are two distinct summands of $I_*(Z(K), \gamma(K) + \gamma(S)|T)$. Since the operator $\bar{R}(S)$ has degree 2 with respect to the $\mathbb{Z}/12\mathbb{Z}$ -grading of $I_*(Z(K), \gamma(K) + \gamma(S)|T)$, the vector spaces in (5.37) and (5.38) have the same dimension. It is also clear from the definition that (5.37) contains $I_*(Z(S), \gamma(K) + \gamma(S)|\bar{R}(S))$, which verifies the claim of this proposition. \square

Conjecture 5.39 *If $Y \setminus K$ is irreducible, then $\dim(\text{SHI}_*(N(S), \alpha(S))) \geq 1$.*

If $Y \setminus K$ is irreducible, then the sutured manifold $(N(S), \alpha(S))$ is *taut*. Kronheimer and Mrowka proved a nonvanishing theorem for the $U(2)$ -sutured Floer homology SHI_*^2 of taut sutured manifolds [56]. As explained in Section 7, we expect that a similar nonvanishing theorem holds for our version of sutured Floer homology.

If Conjecture 5.39 holds, then the answer to Question 1.1 is positive for $N = 3$ and for any nontrivial knot in an integral homology sphere Y . Using the prime decomposition theorem for 3-manifolds, one can decompose Y as a connected sum $Y_1 \# Y_2$ such that $K \subset Y_1$ and $Y_1 \setminus K$ is irreducible. With the aid of Corollary 5.34 and Proposition 5.35, we can conclude from Conjecture 5.39 that the answer to Question 1.1 for $N = 3$ and the pair (Y_1, K) is positive. This clearly would imply the claim for (Y, K) .

6 Gluing theory

6.1 Moduli spaces on manifolds with long neck

Suppose that Y is a connected 3-manifold and $\gamma \subset Y$ is a cycle. We do not assume that (Y, γ) is N -admissible. However, we assume that the critical points of the (possibly perturbed) Chern–Simons functional of (Y, γ) are nondegenerate. Suppose also that (X, w) is a pair with boundary (Y, γ) . As explained in Section 2.2, we can form moduli spaces $\mathcal{M}_p(X, w)$ and their framed counterparts $\widetilde{\mathcal{M}}_p(X, w)$ by working with perturbations of the ASD equation. The Uhlenbeck compactness theorem of Section 2.1 has an analogue for 4-manifolds with cylindrical ends. A proof of this result for $N = 2$ is given in [19, Chapter 5], and it can be extended to higher rank by combining the arguments of [19] and [52]:

Theorem 6.1 *Suppose $\{[A_i]\}_{i \in \mathbb{N}}$ is a sequence of connections in the moduli space $\widetilde{\mathcal{M}}_p(X, w)$. Then there is an element $([B], [C_1], \dots, [C_k])$ of the space*

$$(6.2) \quad \widetilde{\mathcal{M}}_{p_0}(X, w; \alpha_0) \times_{\Gamma_{\alpha_0}} \widetilde{\mathcal{M}}_{p_1}(\alpha_0, \alpha_1) \times_{\Gamma_{\alpha_1}} \cdots \times_{\Gamma_{\alpha_{k-1}}} \widetilde{\mathcal{M}}_{p_k}(\alpha_{k-1}, \alpha_k)$$

and an element $(\mathbf{x}, \mathbf{y}_1, \dots, \mathbf{y}_k)$ of the space

$$(6.3) \quad (X^+)^{m_0} / S_{m_0} \times (Y \times \mathbb{R})^{m_1} / S_{m_1} \times \dots \times (Y \times \mathbb{R})^{m_k} / S_{m_k}$$

for appropriate nonnegative integers m_i such that $\{[A_i]\}_{i \in \mathbb{N}}$, after passing to a sequence, is **weakly chain convergent** to $(([B], \mathbf{x}), ([C_1], \mathbf{y}_1), \dots, ([C_k], \mathbf{y}_k))$. Furthermore,

$$(6.4) \quad \kappa(p) = \kappa(p_0) + \kappa(p_1) + \dots + \kappa(p_k) + m_0 + m_1 + \dots + m_k.$$

Note that $\tilde{\mathcal{M}}_q(\alpha, \beta)$ is the moduli space of framed connections associated to a path $q: \alpha \rightarrow \beta$ over $(Y \times \mathbb{R}, \gamma \times \mathbb{R})$. This moduli space is equipped with an action of $\Gamma_\alpha \times \Gamma_\beta$. The weak chain convergence of $\{[A_i]\}_{i \in \mathbb{N}}$ to $(([B], \mathbf{x}), ([C_1], \mathbf{y}_1), \dots, ([C_k], \mathbf{y}_k))$ means that the following holds [19]: the sequence $\{[A_i]\}_{i \in \mathbb{N}}$, after choosing appropriate gauge representatives, is L_1^P -convergent to B on compact sets of $X \setminus \mathbf{x}$. Moreover, there is a sequence of real numbers

$$t_i^1 := 0 < t_i^2 < \dots < t_i^k$$

with $\lim_i t_i^{j+1} - t_i^j = \infty$ such that the translation of $A_i|_{Y \times [0, \infty)}$ by the constant t_i^j is L_1^P -convergent to C_j on compact sets of $Y \times \mathbb{R} \setminus \mathbf{y}_j$. Identity (6.4) implies that

$$(6.5) \quad \text{index}(\mathcal{D}_{A_i}) \geq \text{index}(\mathcal{D}_{B_1}) + \text{index}(\mathcal{D}_{C_1}) + \dots + \text{index}(\mathcal{D}_{C_k})$$

and the two sides of the inequality are also equal to each other mod $4N$. In (6.5), equality holds if and only if the integers m_j are all zero, and in this case, L_1^P convergence on compact subspaces can be improved to C^∞ convergence on compact subspaces.

Remark 6.6 Theorem 6.1 can be extended to the case that W has more than one boundary component in an obvious way. In this case, we need to fix a chain of the elements of moduli spaces for each boundary component.

There is another compactness theorem we need to review, in which we stretch a 4-manifold along a neck and consider the associated moduli spaces. Suppose (X_1, w_1) and (X_2, w_2) are pairs whose boundaries are (Y, γ) and $(\bar{Y}, \bar{\gamma})$, respectively. Then we can glue these pairs to form (X, w) . We also fix Riemannian metrics on the X_i which are product metrics in neighborhoods of their boundaries associated to a fixed metric on Y . Let X^T be the Riemannian manifold diffeomorphic to X which has an isometric copy of $Y \times (-T, T)$ and is such that $X^T \setminus Y \times (-T, T)$ is isometric to the disjoint union of X_1 and X_2 . The proof of the following theorem is similar to that of Theorem 6.1.

Theorem 6.7 Suppose A_i is a connection on the moduli space $\mathcal{M}_\kappa(X^{T_i}, w)$ such that $\lim_{i \rightarrow \infty} T_i = \infty$. Then there is an element $([B_1], [C_1], \dots, [C_k], [B_2])$ of

$$(6.8) \quad \widetilde{\mathcal{M}}_p(X_1, w_1; \alpha_0) \times_{\Gamma_{\alpha_0}} \widetilde{\mathcal{M}}_{p_1}(\alpha_0, \alpha_1) \times_{\Gamma_{\alpha_1}} \cdots \times_{\Gamma_{\alpha_{k-1}}} \widetilde{\mathcal{M}}_{p_k}(\alpha_{k-1}, \alpha_k) \\ \times_{\Gamma_{\alpha_k}} \widetilde{\mathcal{M}}_{p'}(X_2, w_2; \alpha_k)$$

and an element $(\mathbf{x}_1, \mathbf{y}_1, \dots, \mathbf{y}_k, \mathbf{x}_2)$ of the space

$$(6.9) \quad (X_1^+)^{m_0}/S_{m_0} \times (Y \times \mathbb{R})^{m_1}/S_{m_1} \times \cdots \times (Y \times \mathbb{R})^{m_k}/S_{m_k} \times (X_2^+)^{m_{k+1}}/S_{m_{k+1}}$$

such that $\{[A_i]\}_{i \in \mathbb{N}}$ is weakly chain convergent to

$$(([B_1]\mathbf{x}_1), ([C_1], \mathbf{y}_1), \dots, ([C_k], \mathbf{y}_k), ([B_2], \mathbf{x}_2)).$$

Moreover, we have

$$(6.10) \quad \kappa = \kappa(p) + \kappa(p_1) + \cdots + \kappa(p_k) + \kappa(p') + m_0 + m_1 + \cdots + m_k + m_{k+1}.$$

The following gluing theorem can be regarded as an inverse to [Theorem 6.7](#). There are various places in the literature where similar gluing theorems are discussed [[81](#); [19](#); [55](#)].

[Theorem 6.11](#) can be proved with similar strategies (see eg [[19](#), Theorem 4.17 and Section 4.7.1]):

Theorem 6.11 Let (X_i, w_i) be given as above. For $1 \leq i \leq k$, let $p_i: \alpha_{i-1} \rightarrow \alpha_i$ be a path along $(Y \times \mathbb{R}, w \times \mathbb{R})$, and let \widetilde{N}_i be a compact $(\Gamma_{\alpha_{i-1}} \times \Gamma_{\alpha_i})$ -invariant subspace of $\widetilde{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i)$ which consists of regular points. Suppose we are also given two other compact spaces

$$\widetilde{N}_0 \subset \widetilde{\mathcal{M}}_p(X_1, w_1; \alpha_0) \quad \text{and} \quad \widetilde{N}_{k+1} \subset \widetilde{\mathcal{M}}_{p'}(X_2, w_2; \alpha_k)$$

which contain only regular points and are respectively invariant with respect to the action of Γ_{α_0} and Γ_{α_k} . Here we assume that the perturbations of the ASD equation on the ends of X_1 and X_2 are induced by a fixed perturbation of the Chern–Simons functional of (Y, γ) . Then there is a space \widetilde{U}_i containing \widetilde{N}_i which is an open subset of the relevant moduli space and is invariant with respect to the action of the relevant group. Moreover, for large enough values of T , there is a **gluing map**

$$\Phi_T: \widetilde{U}_0 \times_{\Gamma_{\alpha_0}} \widetilde{U}_1 \times_{\Gamma_{\alpha_1}} \cdots \times_{\Gamma_{\alpha_k}} \widetilde{U}_{k+1} \rightarrow \mathcal{M}_\kappa(X^T, w),$$

where

$$(6.12) \quad \kappa = \kappa(p) + \kappa(p_1) + \cdots + \kappa(p_k) + \kappa(p').$$

The gluing map is a diffeomorphism into its image and satisfies the following properties:

For any fixed element a in the domain of Φ_T , the sequence $\Phi_T(a)$ is chain convergent to a , as T goes to infinity. Moreover, if A_i is a connection on the moduli space $\mathcal{M}_\kappa(X^{T_i}, w)$ such that $\lim_{i \rightarrow \infty} T_i = \infty$ and the sequence $\{A_i\}$ is chain convergent to an element of

$$\tilde{N}_0 \times_{\Gamma_{\alpha_0}} \tilde{N}_1 \times_{\Gamma_{\alpha_1}} \cdots \times_{\Gamma_{\alpha_k}} \tilde{N}_{k+1}$$

then A_i lies in the image of the gluing map for large enough values of i .

Remark 6.13 Theorems 6.7 and 6.11 are strong enough to study the cut-down moduli spaces on a 4-manifold with a long neck. To demonstrate this in the context of an example, suppose (X_i, w_i) is as above and $b^+(X_i) \geq 1$. Let Σ be an embedded surface in X_2 . Let $\nu(\Sigma) \subset X_2$ be an open neighborhood of Σ such that the inclusion of $\nu(\Sigma)$ in X_2 induces a surjection of fundamental groups. Let κ be chosen such that $\mathcal{M}_\kappa(X^T, w)$ has expected dimension two. We make the simplifying assumption that all critical points of the Chern–Simons functional on (Y, γ) are irreducible and nondegenerate, and all the moduli spaces on $Y \times \mathbb{R}$ are regular. By choosing a generic metric and a small compactly supported perturbation, we can assume that the moduli spaces of the form $\mathcal{M}_p(X_1, w_1; \alpha)$ with expected dimension at most 0 contain only irreducible and regular elements [52, Lemma 24]. Similarly, we can arrange for a metric, a small compactly supported perturbation on X_2 , and a geometric representative $V_2(\Sigma) \subset \mathcal{B}^*(\nu(\Sigma)) \times \nu(\Sigma)$ such that:

- (i) The moduli spaces of the form $\mathcal{M}_p(X_2, w_2; \alpha)$ with dimension at most two consist of irreducible and regular solutions.
- (ii) The map $r: \mathcal{M}_p(X_2, w_2; \alpha) \times \Sigma \rightarrow \mathcal{B}^*(\nu(\Sigma)) \times \nu(\Sigma)$, for any moduli space $\mathcal{M}_p(X_2, w_2; \alpha)$ of dimension at most two, is transversal to $V_2(\Sigma)$. Suppose $\mathcal{N}_p(X_1, w_1; \alpha, \Sigma)$ denotes the cut-down moduli space.

The chosen holonomy perturbations on X_1 and X_2 induce a holonomy perturbation on X^T for large values of T . Theorem 6.1 implies that for large enough values of T , the space $\mathcal{M}_\kappa(X^T, w) \times \Sigma$ is also cut-down transversely by $V_2(\Sigma)$ and the resulting space is compact. Furthermore, the elements in the cut-down space $\mathcal{N}_\kappa(X^T, w; \Sigma)$ are in correspondence with the elements of the space

$$\bigcup_{\kappa(p_1)+\kappa(p_2)=\kappa} \mathcal{M}_{p_1}(X_1, w_1; \alpha, \Sigma) \times \mathcal{N}_{p_2}(X_2, w_2; \alpha).$$

In the following, we use a similar strategy to study the cut-down moduli spaces on 4-manifolds with long necks, without going into details.

The following vanishing theorem is a standard application of the above theorems [17; 20; 68]:

Theorem 6.14 *Suppose X_1 and X_2 are two 4-manifolds with $b^+(X_i) \geq 1$. Then for any 2-cycle w in the connected sum $X_1 \# X_2$ and any $z \in \mathbb{A}(X_1 \# X_2)^{\otimes 2}$, the number $D_{X_1 \# X_2, w}(z)$ is equal to zero.*

Proof We can assume that $w = w_1 \cup w_2$ and $z = z_1 \cdot z_2$, where w_i is a 2-cycle in X_i and $z_i \in \mathbb{A}(X_i)^{\otimes 2}$. By replacing X_i , w_i and z_i with $X_i \# \overline{\mathbb{C}\mathbb{P}^2}$, $w_i \cup E_i$ and $z_i \cdot (E_i)_{(2)}^2$, we can also assume that w_i is coprime to N . Here E_i is the exceptional class in $X_i \# \overline{\mathbb{C}\mathbb{P}^2}$. We fix a Riemannian metric on $X_1 \# X_2$ whose restriction to the connected sum region is isometric to the product metric $[-T, T] \times S^3$, where T is a large constant and S^3 has the standard metric. Using the standard metric on S^3 allows us to ensure that all the framed moduli spaces on the cylinder $\mathbb{R} \times S^3$ are regular. We fix a holonomy perturbation of the ASD equation on each X_i such that the perturbation is supported outside of a neighborhood of the connected sum region, and the cut-down moduli spaces $\mathcal{N}_p(X_i, w_i; \Theta, z_i)$ of expected dimension at most zero are regular. Here Θ is the trivial connection, which is the only flat connection on S^3 . Now we can use [Theorem 6.11](#) to conclude that the 0-dimensional moduli space $\mathcal{N}_k(X_1 \# X_2, w_1 \cup w_2; z_1 \cdot z_2)$ is empty when T is large enough. \square

Next, we utilize the gluing and the compactness theorems to prove [Proposition 3.45](#) on connected sums along $\Sigma(2, 3, 23)$. Recall that in addition to the trivial connections, there are 44 irreducible and eight $SU(2)$ -reducible connections on the trivial $SU(3)$ -bundle over $\Sigma(2, 3, 23)$. All these connections are nondegenerate. We choose a perturbation of the Chern–Simons functional of $\Sigma(2, 3, 23)$ (and the empty cycle) such that the critical points of the perturbed functional are the same as those of the Chern–Simons functional and the following assumption about the regularity of the elements of $[A] \in \mathcal{M}_p(\Sigma(2, 3, 23); \alpha, \beta)$ holds: if A is irreducible, then A is regular, and if A is reducible and induced by an $SU(2)$ -connection, then A is regular as a solution of the (perturbed) ASD equation for $SU(2)$ -connections [19; 58].

Suppose X_1 is a 4-manifold with $b^+(X_1) \geq 1$ and $\partial X_1 = \Sigma(2, 3, 23)$. Suppose also that w_1 is a closed 2-cycle in X_1 which is coprime to 3. Fix a metric with cylindrical ends on X_1 and a small holonomy perturbation of the ASD equation which is compatible with the perturbation of the Chern–Simons functional. Let $\mathcal{M}_p(X_1, w_1; \alpha)$ be the moduli space of solutions to a perturbation of the ASD equation associated to the pair (X_1, w_1) and the path p :

Proposition 6.15 *Suppose a positive integer n_0 is given. There is a metric and a holonomy perturbation of the ASD equation on X_1 such that the following holds: Let p be a path along (X_1, w_1) based at a flat connection α on $\Sigma(2, 3, 23)$ such that the index of the elements of $\mathcal{M}_p(X_1, w_1; \alpha)$ is at most n_0 . Then the moduli space $\mathcal{M}_p(X_1, w_1; \alpha)$ consists of regular solutions and does not have any reducible connections. Moreover, suppose $z_1 \in \mathbb{A}(X)^{\otimes 2}$ is such that*

$$d_{X_1, w_1} := -4w_1^2 - 4(\chi(X_1) + \sigma(X_1)) - 4 \equiv \deg(z_1) + 4 \pmod{12}.$$

Suppose also that the expected dimension of $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ is zero. Then there is a geometric representative for z_1 such that the cut-down moduli space $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ is compact.

Note that $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ might be a linear combination of different spaces. Then compactness of this space is defined to be the compactness of all the involved spaces in the linear combination. In what follows, we gloss over this point about the nature of the spaces $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$.

Proof The arguments of [52] can be used to show that the metric and the perturbation can be chosen such that if p is a path as in the statement of the proposition, then $\mathcal{M}_p(X_1, w_1; \alpha)$ is regular and does not contain any reducible solutions. We can also assume that $V(z_1)$ is chosen such that all moduli spaces $\mathcal{N}_p(X_1, w_1; \alpha)$ with expected dimension at most $n_0 - \deg(z_1)$ are cut down transversely. Next, let the dimension of $\mathcal{M}_p(X_1, w_1; \alpha)$ be equal to $\deg(z_1)$, and $\{A_i\}_i$ be a sequence of connections in the cut-down moduli space $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$. By Theorem 6.1, this sequence converges to an element $(([B], \mathbf{x}), ([C_1], \mathbf{x}), \dots, ([C_k], \mathbf{y}_k))$, where $[B] \in \widetilde{\mathcal{N}}_{p_0}(X_1, w_1; \alpha_0, z_1)$, $[C_i] \in \widetilde{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i)$, $\alpha_k = \alpha$, and

$$(6.16) \quad \deg(z_1) \geq \text{index}(\mathcal{D}_B) + \text{index}(\mathcal{D}_{C_1}) + \dots + \text{index}(\mathcal{D}_{C_k}),$$

with equality holding if and only if the multisets $\mathbf{x}, \mathbf{y}_1, \dots, \mathbf{y}_k$ are empty. The main point of the rest of the argument is to show that instantons C_i with nonpositive values of $\text{index}(\mathcal{D}_{C_i})$ do not appear in the limit points of $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$.

We first show that $\text{index}(\mathcal{D}_B)$ in (6.16) is bounded above. If C_i is irreducible, then $\text{index}(\mathcal{D}_{C_i})$ is positive. In the case that C_i is reducible, we cannot guarantee that $\text{index}(\mathcal{D}_{C_i})$ is positive. However, the index of \mathcal{D}_{C_i} as an $\text{SU}(2)$ -connection is positive. Using Table 5, it is straightforward to check that for a reducible connection $C_i \in \widetilde{\mathcal{M}}_{p_i}(\mathbb{R} \times \Sigma(2, 3, 23); \alpha_{i-1}, \alpha_i)$ with nonpositive $\text{index}(\mathcal{D}_{C_i})$, the flat connection α_i

has to be β_8 , and α_{i-1} is equal to either the trivial connection Θ or the $SU(2)$ -connection β_2 . In the first case, $\text{index}(\mathcal{D}_{C_i})$ is equal to -3 and in the latter case $\text{index}(\mathcal{D}_{C_i})$ is equal to -2 . Suppose

$$(6.17) \quad \{i_1, \dots, i_l\} \subseteq \{0, \dots, k-1\}$$

is the set of indices such that $\alpha_{i_j} = \beta_8$. We have

$$(6.18) \quad \text{index}(\mathcal{D}_{C_{i_{j+1}}}) + \text{index}(\mathcal{D}_{C_{i_{j+2}}}) + \dots + \text{index}(\mathcal{D}_{C_{i_{j+1}}}) \equiv 0 \pmod{12}$$

for $1 \leq j \leq k-1$. In (6.18), the last term is at least -3 and the other terms are positive. This shows that the sum in (6.18) is nonnegative. We can use this to conclude that the sum $\text{index}(\mathcal{D}_{C_1}) + \dots + \text{index}(\mathcal{D}_{C_k})$ in (6.16) is at least -3 . That is to say, $\text{index}(\mathcal{D}_B)$ is not greater than $\text{deg}(z_1) + 3$. By increasing the value of n_0 if necessary, we can assume that $\mathcal{N}_{p_0}(X_1, w_1; \alpha_0, z_1)$, which contains B , is cut down transversely. In particular, $\text{index}(\mathcal{D}_B) \geq \text{deg}(z_1)$.

If $l \geq 1$, then the index formula (2.16) shows

$$\text{index}(\mathcal{D}_B) + \text{index}(\mathcal{D}_{C_1}) + \dots + \text{index}(\mathcal{D}_{C_{i_1}}) \equiv d_{X_1, w_1} - 3 \equiv \text{deg}(z_1) + 1 \pmod{12}.$$

In the above expression, the first term on the left-hand side is not less than $\text{deg}(z_1)$, the last term is not less than -3 , and the remaining terms are positive. Therefore, the sum on the left-hand side is not less than $\text{deg}(z_1) + 1$. This also shows that the right-hand side of (6.16) is at least $\text{deg}(z_1) + 1$, which is a contradiction, and as a result, $l = 0$. Therefore, in (6.16), $\text{index}(\mathcal{D}_{C_i})$ is always positive. This also implies that $k = 0$ and x is empty. Consequently, $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ is compact. \square

Form the moduli spaces $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ with the perturbations from Proposition 6.15, and define the following element of $I_4(\Sigma(2, 3, 23))$:

$$(6.19) \quad D_{X_1, w_1}(z_1) := \sum \# \mathcal{N}_p(X_1, w_1; \alpha, z_1) \cdot \alpha,$$

where the sum is over all irreducible connections α in $\Sigma(2, 3, 23)$ and the paths p such that

$$(6.20) \quad \mathcal{N}_p(X_1, w_1; \alpha, z_1)$$

is 0-dimensional. Here we follow the standard conventions to orient (6.20) [58; 52]. Since $N = 3$, we do not need a homology orientation of (X_1, w_1) to fix a sign for $D_{X_1, w_1}(z_1)$.

Next, let X_2 be a 4-manifold with $b^+(X_2) \geq 1$ and $\partial X_2 = \overline{\Sigma(2, 3, 23)}$. Let also w_2 be a closed 2-cycle in X_2 which is coprime to 3. Fix a metric with cylindrical ends on X_2 and a small holonomy perturbation of the ASD equation which is compatible with the perturbation of the Chern–Simons functional:

Proposition 6.21 *Suppose a positive integer n_0 is given. There is a metric and a holonomy perturbation of the ASD equation on X_2 such that the following holds: Let p be a path along (X_2, w_2) based at a flat connection α on $\Sigma(2, 3, 23)$ such that the index of the elements of $\mathcal{M}_p(X_2, w_2; \alpha)$ is at most n_0 . Then the moduli space $\mathcal{M}_p(X_2, w_2; \alpha)$ consists of regular solutions and does not have any reducible connections. Moreover, suppose $z_2 \in \mathbb{A}(X) \otimes^2$ is chosen such that $\deg(z_2)$ is divisible by 4. Suppose also the expected dimension of $\mathcal{N}_p(X_2, w_2; \alpha, z_2)$ is zero. Then there is a geometric representative for z_2 such that the cut-down moduli space $\mathcal{N}_p(X_2, w_2; \alpha, z_2)$ is compact.*

The proof of this proposition is analogous to that of Proposition 6.15, and we leave it to the reader. The cut-down moduli spaces in Proposition 6.21 can be used to define the following functional on $I_*(\Sigma(2, 3, 23))$:

$$(6.22) \quad D^{X_2, w_2}(z_2)(\alpha) := \#\mathcal{N}_p(X_2, w_2; \alpha, z_2),$$

where α is an irreducible connection on $\Sigma(2, 3, 23)$ and the path p is chosen such that $\mathcal{N}_p(X_2, w_2; \alpha, z_2)$ is 0-dimensional. The index formula shows that this space is 0-dimensional only if

$$(6.23) \quad \deg(\alpha) \equiv 4w_2^2 + 4(\chi(X_2) + \sigma(X_2)) - 4 + \deg(z_2) \pmod{12}.$$

Thus, $D^{X_2, w_2}(z_2)$ is nonzero only on the elements of $I_*(\Sigma(2, 3, 23))$ that satisfy (6.23).

Proposition 6.24 *For $i = 1, 2$, suppose (X_i, w_i) and z_i are as in Propositions 6.15 and 6.21. Then*

$$(6.25) \quad D_{X_1 \circ X_2, w_1 \cup w_2}(z_1 \cdot z_2) = D^{X_2, w_2}(z_2) \circ D_{X_1, w_1}(z_1).$$

Proof Use Propositions 6.15 and 6.21 to fix metrics on X_1 and X_2 . Let also $X_1 \circ X_2$ be equipped with a Riemannian metric which is compatible with the metrics on X_1 and X_2 and has a long neck along $\Sigma(2, 3, 23)$. By slightly modifying the holonomy perturbations provided by Propositions 6.15 and 6.21, we can assume that the perturbation of X_i on the complement of a compact subset of X_i^+ is induced by the chosen perturbation of the Chern–Simons functional of $\Sigma(2, 3, 23)$ and the claims of these propositions about

the moduli spaces of the form $\mathcal{N}_p(X_1, w_1; \alpha, z_1)$ and $\mathcal{N}_p(X_2, w_2; \alpha, z_2)$ still hold. The perturbations of the ASD equations on X_1 and X_2 induce a perturbation of the ASD equation on $X_1 \circ X_2$. By applying [Theorem 6.11](#) and arguments similar to those in the proof of [Proposition 6.15](#), we can conclude that, for a long enough neck along $\Sigma(2, 3, 23)$, we have the following diffeomorphism of 0-dimensional moduli spaces:

$$\mathcal{N}_\kappa(X_1 \circ X_2, w_1 \cup w_2, z_1 \cdot z_2) \cong \bigcup_{\alpha} \mathcal{N}_{p_1}(X_1, w_1; \alpha, z_1) \times \mathcal{N}_{p_2}(X_2, w_2; \alpha, z_2).$$

Here α is an irreducible connection on $\Sigma(2, 3, 23)$, and κ, p_1 and p_2 are chosen such that all the above cut-down moduli spaces are 0-dimensional. Standard arguments show that the above diffeomorphism is compatible with respect to the orientation of the involved moduli spaces. This diffeomorphism implies the claim in [\(6.25\)](#). \square

This proposition essentially proves [Proposition 3.45](#) from [Section 3.4](#). We only need to extend the above proposition to the case that w_1 and w_2 are not necessarily coprime to 3 and $\deg(z_2)$ is not divisible by 4. The assumption on the w_i can be removed by the blowing-up trick. The dimension of the moduli space of rank 3 instantons on the closed 4-manifold $X_1 \circ X_2$ is always divisible by 4. Therefore, if we define $D^{X_2, w_2}(z_2) = 0$ in the case that $\deg(z_2) \not\equiv 0 \pmod 4$, then the above theorem still holds.

6.2 Gluing theory for negative embedded spheres

In this subsection, we give a proof of [Proposition 2.21](#) based on the techniques which are discussed in [Section 6.1](#). Suppose X is a smooth 4-manifold with $b^+(X) \geq 2$. Suppose also σ is an embedded sphere in X with $\sigma \cdot \sigma = -3$. A tubular neighborhood of σ , denoted by Z , is a disc bundle over σ with Euler class -3 and boundary $Y = L(3, -1)$. Thus X can be split as $Z \circ X_1$, where X_1 is the closure of the complement of Z . We will also write w_0 for a fiber of the disc bundle Z . Fix the orientation on w_0 such that w_0 intersects σ negatively in one point.

Fix a Riemannian metric with a cylindrical end on Z . Let also L be the complex line bundle over Z associated to the 2-cycle w_0 . Since $b^+(Z) = b^1(Z) = 0$, there is a unique ASD connection on L with finite energy. This connection, denoted by B , is asymptotic to a flat connection χ which maps the generator of $\pi_1(Y)$ to $\zeta = e^{2\pi i/3}$. Consider the ASD $U(N)$ -connection $A := B^{k_1} \oplus \dots \oplus B^{k_N}$ which is asymptotic to the flat connection $\alpha := \chi^{k_1} \oplus \dots \oplus \chi^{k_N}$. The first Chern class of the underlying $U(N)$ -connection is equal to $(k_1 + \dots + k_N) \text{P.D.}[w_0]$. The topological energy of A

is given by the formula

$$\kappa(A) = \frac{1}{12N} \sum_{1 \leq i, j \leq N} (k_i - k_j)^2.$$

Therefore, we can use (2.16) to compute the index of \mathcal{D}_A . In particular, if $|k_i - k_j| \leq 3$ for all i and j , then (2.16) shows [12] that

$$\text{index}(\mathcal{D}_A) = 1 - N^2 + \sum_{1 \leq i, j \leq N} |k_i - k_j|.$$

Table 2 consists of various choices of $U(3)$ -connections. For each connection A , the first Chern class of the underlying bundle of A is equal to k P.D. $[w_0]$, where k is also given in Table 2.

A	α	k	$\text{index } \mathcal{D}_A$	$\dim \tilde{\mathcal{M}}(A)$	Γ_α
$B^2 \oplus 1 \oplus 1$	$\chi^2 \oplus 1 \oplus 1$	2	0	4	$S(U(1) \times U(2))$
$B \oplus B \oplus 1$	$\chi \oplus \chi \oplus 1$	2	-4	0	$S(U(2) \times U(1))$
$B^{-1} \oplus B^2 \oplus B$	$\chi^2 \oplus \chi^2 \oplus \chi$	2	4	8	$S(U(2) \times U(1))$
$B \oplus B^{-1} \oplus B^{-1}$	$\chi \oplus \chi^2 \oplus \chi^2$	-1	0	4	$S(U(1) \times U(2))$
$B^{-1} \oplus 1 \oplus 1$	$\chi^2 \oplus 1 \oplus 1$	-1	-4	0	$S(U(1) \times U(2))$
$B^{-2} \oplus B \oplus 1$	$\chi \oplus \chi \oplus 1$	-1	4	8	$S(U(2) \times U(1))$

Table 2

We will write $\mathcal{M}(A)$ (respectively, $\tilde{\mathcal{M}}(A)$) for the moduli space (respectively, the framed moduli space) of connections on Z corresponding to the path which is represented by A . From now on, we assume that an anti-self-dual metric with positive scalar curvature is fixed on Z [61].⁹

Proposition 6.26 *For each A in Table 2, A has the minimal topological energy among all ASD connections with the same limiting flat connection as A and the same c_1 .*

Proof Let A' be an ASD connection with a smaller topological energy and the same limiting flat connection and c_1 as A . The difference $\dim(\tilde{\mathcal{M}}(A)) - \dim(\tilde{\mathcal{M}}(A'))$ is at least 12. On the other hand, any such connection is regular because of the choice of

⁹We use such metrics to ensure regularity of moduli spaces of ASD connections with respect to these metrics. The construction of [61] provides asymptotically cylindrical ASD metrics rather than cylindrical metrics. However, that is enough for our purposes because we can pick a cylindrical approximation to these metrics and then argue as in [13, Corollary 5.13].

the metric. Therefore, the dimension of $\widetilde{\mathcal{M}}(A')$ is at least $\dim(\Gamma_\alpha) - \dim(\Gamma_{A'})$. This can be used to rule out the existence of A' . \square

We study the moduli spaces $\mathcal{M}(A)$ for various choices of A . The same argument as in Proposition 6.26 shows that if $A = B \oplus B \oplus 1$, then the moduli space $\mathcal{M}(A)$ contains only the completely reducible connection A . Next, we turn to the case that $A = B^2 \oplus 1 \oplus 1$. An argument analogous to that in Proposition 6.26 shows that there are three types of connections in this space:

- finitely many irreducible connections;
- the complete reducible connection $A = B^2 \oplus 1 \oplus 1$;
- reducible connections of the form $R \oplus 1$, where R is an irreducible $U(2)$ -connection in the 1-dimensional space $\mathcal{M}(B^2 \oplus 1)$.

Proposition 6.27 *The unique component of $\mathcal{M}(B^2 \oplus 1)$ containing $B^2 \oplus 1$ is a half-line $[0, \infty)$. All the other components are either circles or copies of \mathbb{R} consisting of only irreducible connections. The corresponding components of $\widetilde{\mathcal{M}}(B^2 \oplus 1 \oplus 1)$ are \mathbb{C}^2 , $S^3 \times S^1$ and $S^3 \times \mathbb{R}$.*

Proof The proof of the first part is straightforward. For the second part, note that the orbit of the connection $R \oplus 1$ in the framed moduli space, for an irreducible connection R , is $\Gamma_\alpha / \Gamma_{R \oplus 1} = S(U(1) \times U(2)) / S(U(1)_2 \times U(1)) = S^3$, where $U(1)_2 \times U(1)$ denotes 3×3 diagonal matrices where the first two diagonal entries are equal to each other. \square

Now we are ready to prove the first part of Proposition 2.21:

Proposition 6.28 *Suppose X and σ are as above and $z \in \mathbb{A}(\langle \sigma \rangle^\perp)^{\otimes 2}$. Suppose also that w is a 2-cycle in X such that $w \cdot \sigma \equiv 1 \pmod{3}$. Then there is a constant c such that*

$$D_{X,w}^3 \left(\left(-\frac{3}{2}\sigma_{(3)} - \frac{3}{2}\sigma_{(2)}^2 - a_2 \right) z \right) = c D_{X,w-\sigma}^3(z).$$

Proof For simplicity of exposition, we assume that $z = 1$. A similar proof works in the more general case. Equip X with a Riemannian metric that has a neck of length T along the lens space Y , and denote the resulting Riemannian manifold by X^T . We first study the 4-dimensional moduli spaces of the form $\mathcal{M}_{\kappa_0}(X^T, w)$ for large values of T . Let α_1 and α_2 denote the flat connections $\chi^2 \oplus 1 \oplus 1$ and $\chi \oplus \chi \oplus 1$. We write G_i for the stabilizer group Γ_{α_i} . We also write \mathcal{R}_1 for $\widetilde{\mathcal{M}}(B^2 \oplus 1 \oplus 1)$ and \mathcal{R}_2 for $\widetilde{\mathcal{M}}(B \oplus B \oplus 1)$. Clearly \mathcal{R}_i is a G_i -manifold. By Theorem 6.11 and

the above description of the low-dimensional moduli spaces on Z , the moduli space $\mathcal{M}_{\kappa_0}(X^T, w)$ can be covered with two open sets \mathcal{U} and \mathcal{V} of the form

$$\mathcal{U} = \widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{R}_1 \quad \text{and} \quad \mathcal{V} = \widetilde{\mathcal{M}}_{p_2}(X_1, w \cap X_1; \alpha_2) \times_{G_2} \mathcal{R}_2,$$

where the framed moduli spaces $\widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1)$ and $\widetilde{\mathcal{M}}_{p_2}(X_1, w \cap X_1; \alpha_2)$ are respectively of dimensions 4 and 8. By choosing a generic metric on X_1 and a small holonomy perturbation of the ASD equation supported in X_1 , we can assume the action of Γ_{α_i} on $\widetilde{\mathcal{M}}_{p_i}(X_1, w \cap X_1; \alpha_i)$ is free. In particular, there are G_i equivariant maps

$$(6.29) \quad \tilde{f}_i: \widetilde{\mathcal{M}}_{p_i}(X_1, w \cap X_1; \alpha_i) \rightarrow EG_i,$$

where EG_i is the universal principal G_i -bundle over the classifying space BG_i . Every connected component of $\widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1)$ can be identified with G_1 and we can assume that the map \tilde{f}_1 is the same on different connected components of $\widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1)$.

Gluing theory also gives a description of the intersection $\mathcal{U} \cap \mathcal{V}$ for large values of T . Let \mathcal{N} be the $(G_1 \times G_2)$ -manifold $\widetilde{\mathcal{M}}_q(\alpha_1, \alpha_2)$, where the path q is chosen such that this space is 8-dimensional. Since the orbit of any irreducible connection with respect to the action of $G_1 \times G_2$ and translations is at least 9-dimensional, \mathcal{N} does not contain any irreducible connections. Thus \mathcal{N} consists of ASD connections of the form $R \oplus 1$ on $\mathbb{R} \times L(3, 1)$, where R is asymptotic to $\chi^2 \oplus 1$ and $\chi \oplus \chi$ on the two ends. After dividing by the action of translations, there are finitely many choices for the connection R . Thus, this manifold, as a $(G_1 \times G_2)$ -space, is the union of finitely many spaces of the form

$$(6.30) \quad \mathbb{R} \times \frac{G_1 \times G_2}{S(U(1)_2 \times U(1))}.$$

In particular, the action of G_i on this space is free. The G_1 -space $\mathcal{N} \times_{G_2} \mathcal{R}_2$ can be identified with the end of \mathcal{R}_1 . The G_2 -space $\widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{N}$ can be identified with the end of $\widetilde{\mathcal{M}}_{p_2}(X_1, w \cap X_1; \alpha_2)$. Using these identifications, the overlap $\mathcal{U} \cap \mathcal{V}$ is identified with the space

$$(6.31) \quad \widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2.$$

In summary, $\mathcal{M}_{\kappa_0}(X^T, w)$ is the pushout of the following diagram:

$$(6.32) \quad \begin{array}{ccc} \widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2 & \hookrightarrow & \widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{R}_1 \\ \downarrow & \lrcorner & \downarrow \\ \widetilde{\mathcal{M}}_{p_2}(X_1, w \cap X_1; \alpha_2) \times_{G_2} \mathcal{R}_2 & \longrightarrow & \mathcal{M}_{\kappa_0}(X^T, w) \end{array}$$

We can form a universal version of the above diagram as follows:

$$(6.33) \quad \begin{array}{ccc} EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2 & \xrightarrow{h_2} & EG_1 \times_{G_1} \mathcal{R}_1 \\ \downarrow h_1 & \lrcorner & \downarrow \\ EG_2 \times_{G_2} \mathcal{R}_2 & \longrightarrow & M \end{array}$$

The free action of G_2 on $EG_1 \times_{G_1} \mathcal{N}$ gives a G_2 -equivariant map

$$f: EG_1 \times_{G_1} \mathcal{N} \rightarrow EG_2.$$

The vertical map h_1 is induced by f . The horizontal map h_2 is induced by the inclusion of $\mathcal{N} \times_{G_2} \mathcal{R}_2$ into \mathcal{R}_1 . The space M is then the pushout of h_1 and h_2 . The maps \tilde{f}_1 and \tilde{f}_2 determine the maps

$$f_i: \tilde{\mathcal{M}}_{p_i}(X_1, w \cap X_1; \alpha_i) \times_{G_i} \mathcal{R}_i \rightarrow EG_i \times_{G_i} \mathcal{R}_i$$

and

$$g: \tilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2 \rightarrow EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2.$$

The maps \tilde{f}_1 and \tilde{f}_2 can be chosen such that f_1 , f_2 and g give rise to a map from diagram (6.32) to diagram (6.33). We will write F for the induced map from $\mathcal{M}_{\kappa_0}(X^T, w)$ to M .

There is a similar description of the universal bundle over the space $\mathcal{M}_{\kappa_0}(X^T, w) \times \sigma$. Suppose q_1 is the path over Z determined by the connection $B^2 \oplus 1 \oplus 1$. We can form an equivariant universal bundle $\tilde{\mathbb{P}}_1$ over $\tilde{\mathcal{B}}_{q_1}(Z, w \cap Z; \alpha) \times \sigma$ similar to the universal bundles in Section 2.1. There is an action of G_1 on $\tilde{\mathbb{P}}_1$ which lifts the obvious action of this group on $\tilde{\mathcal{B}}_{q_1}(Z, w \cap Z; \alpha) \times \sigma$. This bundle induces a PU(3)-bundle over $\mathcal{R}_1 \times \sigma$ with an action of G_1 , which we still denote by $\tilde{\mathbb{P}}_1$. Similarly, we can construct a PU(3)-bundle $\tilde{\mathbb{P}}_2$ over $\mathcal{R}_2 \times \sigma$ with an action of G_2 . Since \mathcal{R}_2 consists of only the class of the connection $B \oplus B \oplus 1$, the bundle $\tilde{\mathbb{P}}_2$ has the form [12, Proposition 46]

$$(6.34) \quad \tilde{\mathbb{P}}_2 = F \boxtimes L \oplus P,$$

where F and P are the standard U(2)- and U(1)-representations of G_2 , and L is the line bundle defined at the beginning of the subsection. To be more precise, (6.34) gives a lift of $\tilde{\mathbb{P}}_2$ to a U(3)-bundle with an action of G_2 . The restriction of $\tilde{\mathbb{P}}_1$, as a PU(3)-bundle with an action of G_1 , to the subset $\mathcal{N} \times_{G_2} \mathcal{R}_2 \times \sigma$ of $\mathcal{R}_1 \times \sigma$ can be identified

with $\mathcal{N} \times_{G_2} \widetilde{\mathbb{P}}_2$. Thus, $\widetilde{\mathbb{P}}_1$ and $\widetilde{\mathbb{P}}_2$ give rise to the following diagram of $\text{PU}(3)$ -bundles:

$$(6.35) \quad \begin{array}{ccc} \mathbb{P}_3 := EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \widetilde{\mathbb{P}}_2 & \xrightarrow{\bar{h}_2} & EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1 \\ \downarrow \bar{h}_1 & \lrcorner & \downarrow \\ EG_2 \times_{G_2} \widetilde{\mathbb{P}}_2 & \longrightarrow & \bar{\mathbb{P}} \end{array}$$

The $\text{PU}(3)$ -bundle $\bar{\mathbb{P}}$ over $M \times \sigma$ is the pushout of the above diagram. The pullback of $\bar{\mathbb{P}}$ with respect to the map $(F, \text{id}): \mathcal{M}_{\kappa_0}(X^T, w) \times \sigma \rightarrow M \times \sigma$ is equal to the universal bundle.

Suppose $\varphi \in H^4(M, \mathbb{Q})$ is defined to be

$$\frac{3}{2}c_3(\bar{\mathbb{P}})/\sigma - \frac{3}{2}(c_2(\bar{\mathbb{P}})/\sigma)^2 - c_2(\bar{\mathbb{P}})/x.$$

We claim the restriction of φ to the subspace $EG_2 \times_{G_2} \mathcal{R}_2$ vanishes. To see this, let

$$p := c_1(P) \quad \text{and} \quad f := c_2(F) \in H_{G_2}^*(\mathcal{R}_2) = H^*(BG_2)$$

be the equivariant Chern classes of the bundles P and F in the description of $\widetilde{\mathbb{P}}_2$ in (6.34). Then it is easy to verify our claim from the following identities:

$$\begin{aligned} c_2(EG_2 \times_{G_2} \widetilde{\mathbb{P}}_2)/\sigma &= -p, & c_3(EG_2 \times_{G_2} \widetilde{\mathbb{P}}_2)/\sigma &= \frac{1}{3}p^2 + \frac{2}{3}f, \\ c_2(EG_2 \times_{G_2} \widetilde{\mathbb{P}}_2)/x &= -p^2 + f. \end{aligned}$$

By (6.30), the space $EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2$ is homotopy equivalent to finitely many copies of

$$EG_1/S(U(1)_2 \times U(1)) \cong BS(U(1)_2 \times U(1)) \cong B(U(1) \times U(1))$$

whose odd-degree cohomology groups vanish. Therefore, we can apply the Mayer-Vietoris exact sequence for the pushout diagram in (6.33) to conclude that there is a unique choice of a relative cohomology class

$$\psi \in H^4(EG_1 \times_{G_1} \mathcal{R}_1, EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2)$$

such that φ is equal to the image of ψ in $H^4(M, \mathbb{Q})$. This discussion shows that the pairing of $F^*(\varphi)$ and the fundamental class of the 4-dimensional moduli space $\mathcal{M}_{\kappa_0}(X^T, w)$ is equal to the pairing of $f_1^*(\psi)$ and the relative fundamental class of $\widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{G_1} \mathcal{R}_1$. The description of the moduli spaces and the map \tilde{f}_1 shows that there is a constant c such that the latter pairing is equal to

$$c \cdot \#\mathcal{M}_{p_1}(X_1, w \cap X_1; \alpha_1).$$

Since $\widetilde{\mathcal{M}}(B^{-1} \oplus 1 \oplus 1)$ consists of a single point, we have

$$(6.36) \quad \begin{aligned} \mathcal{M}_{p_1}(X_1, w \cap X_1; \alpha_1) &= \widetilde{\mathcal{M}}_{p_1}(X_1, w \cap X_1; \alpha_1) \times_{\Gamma_{\alpha_1}} \widetilde{\mathcal{M}}(B^{-1} \oplus 1 \oplus 1) \\ &= \mathcal{M}_{\kappa_1}(X^T, w - \sigma). \end{aligned}$$

Here κ_1 is chosen such that the space $\mathcal{M}_{\kappa_1}(X^T, w - \sigma)$ is 0-dimensional. We use [Theorem 6.11](#) to conclude the second identity for large enough values of T . The identities in (6.36) allow us to verify the desired claim. □

The second part of [Proposition 2.21](#) can be proved by applying the first part to the (-3) -sphere σ with the reverse orientation. Next, we turn to the proof of the last part of [Proposition 2.21](#). As in the previous case, we need to study some low-dimensional moduli spaces over Z . [Table 3](#) consists of various choices of $U(3)$ -connections on Z with vanishing c_1 . The proof of [Proposition 6.26](#) shows that each connection A in this table has the minimal energy among all ASD connections with the same limiting flat connection and vanishing c_1 .

A	α	index \mathcal{D}_A	$\dim \widetilde{\mathcal{M}}(A)$	Γ_α
$1 \oplus 1 \oplus 1$	$1 \oplus 1 \oplus 1$	-8	0	$SU(3)$
$B \oplus B^{-1} \oplus 1$	$\chi \oplus \chi^2 \oplus 1$	0	2	$S(U(1) \times U(1) \times U(1))$
$B \oplus B \oplus B^{-2}$	$\chi \oplus \chi \oplus \chi$	4	12	$SU(3)$
$B^{-1} \oplus B^{-1} \oplus B^2$	$\chi^2 \oplus \chi^2 \oplus \chi^2$	4	12	$SU(3)$

Table 3

For $A = B \oplus B^{-1} \oplus 1$, the moduli space $\mathcal{M}(A)$ contains three types of connections:

- finitely many irreducible connections;
- the completely reducible connection $A = B \oplus B^{-1} \oplus 1$;
- reducible connections of the form $R \oplus 1$, where R is an irreducible $SU(2)$ -connection in the 1-dimensional space $\mathcal{M}(B \oplus B^{-1})$.

Proposition 6.37 *Each of the connected components of $\mathcal{M}(B \oplus B^{-1})$ which contains $B \oplus B^{-1}$ is a half-line $[0, \infty)$. All the other components are either circles or copies of \mathbb{R} consisting of only irreducible connections. The corresponding components of $\widetilde{\mathcal{M}}(A)$ are \mathbb{C} , $S^1 \times S^1$ and $S^1 \times \mathbb{R}$, and the action of $\Gamma_\alpha = S^1$ is standard.*

Proposition 6.38 *Suppose X and σ are as above and $z \in \mathbb{A}((\sigma)^\perp)^{\otimes 2}$. Suppose also w is a 2-cycle in X_1 . Then*

$$D_{X,w}^3((\sigma_{(2)}^4 + 4a_2\sigma_{(2)}^2 + 3\sigma_{(3)}^2)z) = 0$$

and

$$D_{X,w}^3((\sigma_{(2)}^3\sigma_{(3)} + 3a_3\sigma_{(2)}^2 + a_2\sigma_{(2)}\sigma_{(3)})z) = 0.$$

Proof Similar to the proof of Proposition 6.28, we may assume that $z = 1$. We need to study the 8-dimensional moduli space for the first identity and the 10-dimensional moduli space for the second identity. Suppose κ_0 is a constant number such that the expected dimension of the moduli space $\mathcal{M}_{\kappa_0}(X^T, w)$ is 8 or 10. This moduli space is compact. Moreover, Theorem 6.11 and the above description of the low-dimensional moduli spaces on Z show that $\mathcal{M}_{\kappa_0}(X^T, w)$ can be given as the pushout of the diagram

$$(6.39) \quad \begin{array}{ccc} \widetilde{\mathcal{M}}_{p_1}(X_1, w; \alpha_1) \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2 & \hookrightarrow & \widetilde{\mathcal{M}}_{p_1}(X_1, w; \alpha_1) \times_{G_1} \mathcal{R}_1 \\ \downarrow & \Gamma & \downarrow \\ \widetilde{\mathcal{M}}_{p_2}(X_1, w; \alpha_2) \times_{G_2} \mathcal{R}_2 & \longrightarrow & \mathcal{M}_{\kappa_0}(X^T, w) \end{array}$$

where $\alpha_1 = \chi \oplus \chi^2 \oplus 1$, $\alpha_2 = 1 \oplus 1 \oplus 1$, $G_i = \Gamma_{\alpha_i}$, $\mathcal{R}_1 = \widetilde{\mathcal{M}}(B \oplus B^{-1} \oplus 1)$, $\mathcal{R}_2 = \widetilde{\mathcal{M}}(1 \oplus 1 \oplus 1)$, $\mathcal{N} = \widetilde{\mathcal{M}}_q(\alpha_1, \alpha_2)$ and the path q is chosen such that the index of any element in \mathcal{N} is equal to 0. As in the previous case, we can form the universal version of the above diagram as

$$(6.40) \quad \begin{array}{ccc} EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2 & \xrightarrow{h_2} & EG_1 \times_{G_1} \mathcal{R}_1 \\ \downarrow h_1 & \Gamma & \downarrow \\ EG_2 \times_{G_2} \mathcal{R}_2 & \longrightarrow & M \end{array}$$

with a map $F: \mathcal{M}_{\kappa_0}(X^T, w) \rightarrow M$.

There are also $\text{PU}(3)$ -bundles $\widetilde{\mathbb{P}}_i$ on \mathcal{R}_i with an action of G_i . These bundles give rise to a pushout diagram of bundles

$$(6.41) \quad \begin{array}{ccc} \mathbb{P}_3 := EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \widetilde{\mathbb{P}}_2 & \xrightarrow{\bar{h}_2} & EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1 \\ \downarrow \bar{h}_1 & \Gamma & \downarrow \\ EG_2 \times_{G_2} \widetilde{\mathbb{P}}_2 & \longrightarrow & \widetilde{\mathbb{P}} \end{array}$$

such that $(F, \text{id})^*(\bar{\mathbb{P}})$ is equal to the universal bundle on $\mathcal{M}_{\kappa_0}(X^T, w) \times \sigma$. Let $\varphi_1 \in H^8(M, \mathbb{Q})$ and $\varphi_2 \in H^{10}(M, \mathbb{Q})$ be defined by

$$\begin{aligned} \varphi_1 &:= (c_2(\bar{\mathbb{P}})/\sigma)^4 + 4(c_2(\bar{\mathbb{P}})/x) \cdot (c_2(\bar{\mathbb{P}})/\sigma)^2 + 3(c_3(\bar{\mathbb{P}})/\sigma)^2, \\ \varphi_2 &:= (c_2(\bar{\mathbb{P}})/\sigma)^3 \cdot (c_3(\bar{\mathbb{P}})/\sigma) + 3(c_3(\bar{\mathbb{P}})/x) \cdot (c_2(\bar{\mathbb{P}})/\sigma)^2 \\ &\quad + (c_2(\bar{\mathbb{P}})/x) \cdot (c_2(\bar{\mathbb{P}})/\sigma) \cdot (c_3(\bar{\mathbb{P}})/\sigma). \end{aligned}$$

The restrictions of cohomology classes φ_1 and φ_2 to the open sets $EG_i \times_{G_i} \mathcal{R}_i$ vanish. We show this claim for the connected component $EG_1 \times_{G_1} \mathbb{C}$ of $EG_1 \times_{G_1} \mathcal{R}_1$. The restriction of the bundle $\widetilde{\mathbb{P}}_1$ to the G_1 -manifold \mathbb{C} is given by [12, Proposition 46]

$$P \boxtimes L \oplus Q \boxtimes L^{-1} \oplus R \boxtimes \underline{\mathbb{C}},$$

where P , Q and R are the G_1 -equivariant line bundles over \mathbb{C} associated to the three standard 1-dimensional representations of G_1 . Note that $P \otimes Q \otimes R$ is the trivial bundle with the trivial action of G_1 . Suppose $p, q \in H_{G_1}^*(\mathbb{C})$ denote the equivariant first Chern classes of the bundles P and Q . Then

$$\begin{aligned} c_2(EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1)/\sigma &= p - q, & c_3(EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1)/\sigma &= q^2 - p^2, \\ c_2(EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1)/x &= -p^2 - q^2 - pq, & c_3(EG_1 \times_{G_1} \widetilde{\mathbb{P}}_1)/x &= -pq(p + q). \end{aligned}$$

These identities can be used to show that the restrictions of φ_1 and φ_2 to $EG_1 \times_{G_1} \mathbb{C}$ are equal to zero. Using similar arguments, it is even easier to show that the restrictions of these two cohomology classes to $EG_2 \times_{G_2} \mathcal{R}_2$ and the remaining connected components of $EG_1 \times_{G_1} \mathcal{R}_1$ are equal to zero. Since the cohomology groups of $EG_1 \times_{G_1} \mathcal{N} \times_{G_2} \mathcal{R}_2$ in odd degrees vanish, the Mayer-Vietoris exact sequence for diagram (6.40) implies that φ_1 and φ_2 are both equal to zero. In particular, this verifies the claim of this proposition. □

6.3 Gluing theory for Fukaya–Floer homology

One of the primary goals of this subsection is to define the Fukaya–Floer homology for an N -admissible pair (Y, γ) and an $(N-1)$ -tuple $L = (l_2, \dots, l_N)$ of the elements of $H_1(Y)$. As it is mentioned in Section 3.3, we shall construct a chain complex $(\mathfrak{C}_*^{N,j}(Y, \gamma, L), d_{N,j})$ over $R_{N,j}$, for each nonnegative integer number j . We also define a chain map

$$F_j^k: \mathfrak{C}_*^{N,j}(Y, \gamma, L) \rightarrow \mathfrak{C}_*^{N,k}(Y, \gamma, L) \quad \text{for } j \geq k \geq 0$$

such that F_j^k is a homomorphism of $R_{N,j}$ -modules, and for a triple of integers satisfying $j \geq k \geq l \geq 0$, the map $F_k^l \circ F_j^k$ is chain homotopy equivalent to F_j^l .

Let $\mathbb{I}_*^{N,j}(Y, \gamma, L)$ be the homology of the chain complex $(\mathfrak{C}_*^{N,j}(Y, \gamma, L), d_{N,j})$ and $f_j^k: \mathbb{I}_*^{N,j}(Y, \gamma, L) \rightarrow \mathbb{I}_*^{N,k}(Y, \gamma, L)$ be the map induced by F_j^k . Then the Fukaya–Floer homology group $\mathbb{I}_*^N(Y, \gamma, L)$ is defined to be the inverse limit of the inverse system $(\{\mathbb{I}_*^{N,j}(Y, \gamma, L)\}_j, \{f_j^k\}_{j \geq k})$.

For simplicity of exposition, we assume that $l_3 = \dots = l_N = 0$ and l_2 is an integral homology class. Later we shall explain how the definition should be adapted to the arbitrary case. For each $i \in \mathbb{N}$, let η_i be an oriented closed curve that represents l_2 . Let also $\nu(\eta_i)$ be a regular neighborhood of η_i such that the inclusion of $\nu(\eta_i)$ into Y induces a surjective map at the level of fundamental groups. Furthermore, we assume that the open sets $\nu(\eta_i)$ are disjoint.

For a fixed integer $j \geq 0$, let CS_{π_j} be a perturbation of the Chern–Simons functional associated to (Y, γ) such that all critical points of CS_{π_j} are irreducible and nondegenerate, and all moduli spaces $\mathcal{M}_p(\alpha, \beta)$ with dimension at most $2j + 1$ consist of regular points. If i is a nonnegative integer number not greater than j and $\dim(\mathcal{M}_p(\alpha, \beta)) \leq 2j + 1$, then we may also assume that the restriction of any element in $\mathcal{M}_p(\alpha, \beta)$ to the open subspace $\nu(\eta_i) \times (0, 1)$ is irreducible.¹⁰ We define

$$(6.42) \quad \mathfrak{C}_*^{N,j}(Y, \gamma, L) := \mathfrak{C}_*^{\pi_j}(Y, \gamma) \otimes R_{N,j}.$$

Next, we need to define the differential $d_{N,j}$.

For any path p between two critical points α and β of CS_{π_j} and $S = \{i_1, \dots, i_k\} \subset \{1, \dots, j\}$, we have the diagonal \mathbb{R} –action $\{\sigma_t\}_{t \in \mathbb{R}}$ on the space

$$(6.43) \quad \mathcal{M}_p(\alpha, \beta) \times (\eta_{i_1} \times \mathbb{R}) \times \dots \times (\eta_{i_k} \times \mathbb{R}).$$

Here the action of $t \in \mathbb{R}$ maps $(x, r) \in \eta_i \times \mathbb{R}$ to $(x, r - t)$. If p is not the constant path or S is not the empty set, then this action is free and we will write $\check{\mathcal{M}}_p(\alpha, \beta; S)$ for the quotient space. Otherwise, $\check{\mathcal{M}}_p(\alpha, \beta; S)$ is defined to be empty. In the case that S is empty, the quotient space is the space $\check{\mathcal{M}}_p(\alpha, \beta)$. If $S' \subset S$, then there is an obvious projection map from $\check{\mathcal{M}}_p(\alpha, \beta; S)$ to $\check{\mathcal{M}}_p(\alpha, \beta; S')$, which is denoted by $\pi_{S \rightarrow S'}$.

We can form a partial compactification of $\check{\mathcal{M}}_p(\alpha, \beta; S)$ by defining

$$\check{\mathcal{M}}_p^+(\alpha, \beta; S) := \bigcup_k \bigcup_{\{(p_i, S_i)\}_{1 \leq i \leq k}} \prod_{i=1}^k \check{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i),$$

¹⁰This is a consequence of the unique continuation property of (nonperturbed) ASD connections and the Uhlenbeck compactness theorem. For more details see [52].

where $S_i \subset S$ and $p_i: \alpha_{i-1} \rightarrow \alpha_i$ is a path between two critical points of CS_{π_j} such that the sets S_i are disjoint, their union is equal to S , and the composition $p_1 \circ \dots \circ p_k$ is equal to p . Note that the set S_i may be empty. A sequence $u_l \in \check{\mathcal{M}}_p(\alpha, \beta; S)$ is chain convergent to $u_\infty = (u_\infty^1, \dots, u_\infty^k) \in \prod_{i=1}^k \check{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i)$ if there is a sequence of k -tuples of real numbers (t_l^1, \dots, t_l^k) such that

$$t_l^1 < \dots < t_l^k, \quad \lim_{l \rightarrow \infty} t_l^{m+1} - t_l^m = \infty \quad \text{for all } 1 \leq m \leq k-1$$

and

$$\sigma_{t_l^m} \circ \pi_{S \rightarrow S_m}(u_l) \xrightarrow{C_{\text{loc}}^\infty} u_\infty^m.$$

We use this notion of convergence to define a topology on $\check{\mathcal{M}}_p^+(\alpha, \beta; S)$.

Remark 6.44 Due to the possibility of bubbling off instantons, the space $\check{\mathcal{M}}_p^+(\alpha, \beta; S)$ is not necessarily compact.

Example 6.45 Suppose $A \in \mathcal{M}_p(\alpha, \beta)$, where $p: \alpha \rightarrow \beta$ is a nontrivial path between two critical points of the perturbed Chern–Simons functional. Suppose also $x \in \eta_1$ is fixed. Then $u_i = [A, (x, i)]$ defines a sequence of elements of $\check{\mathcal{M}}_p(\alpha, \beta; S)$ where $S = \{1\}$. This sequence is convergent to (u_∞^1, u_∞^2) :

$$u_\infty^1 := [A] \in \check{\mathcal{M}}_p(\alpha, \beta) \quad \text{and} \quad u_\infty^2 := [\pi^*(\beta), (x, 0)] \in \check{\mathcal{M}}_q(\beta, \beta; S).$$

Here q is the constant path from β to β , and $\pi^*(\beta)$ is the pullback of the connection β to $\mathbb{R} \times Y$.

For $S = \{i_1, \dots, i_k\}$, define the map

$$(6.46) \quad \Phi_S: \check{\mathcal{M}}_p(\alpha, \beta; S) \rightarrow \prod_{i \in S} \mathcal{B}^*(v(\eta_i) \times (0, 1)) \times \eta_i$$

by

$$\begin{aligned} \Phi_S([A, (x_{i_1}, t_{i_1}), \dots, (x_{i_k}, t_{i_k})]) \\ = ((T_{t_{i_1}}^* A|_{v(\eta_{i_1}) \times (0,1)}, x_{i_1}), \dots, (T_{t_{i_k}}^* A|_{v(\eta_{i_k}) \times (0,1)}, x_{i_k})), \end{aligned}$$

where $T_{t_i}: \mathbb{R} \times Y \rightarrow \mathbb{R} \times Y$ denotes the translation that maps (x, t) to $(x, t + t_i)$. This map extends to $\check{\mathcal{M}}_p^+(\alpha, \beta; S)$ in the obvious way and the extension is continuous. As in Section 2.1, we can form a universal $\text{PU}(N)$ -bundle \mathbb{P}_i and an associated $\text{SU}(N^N)$ vector bundle \mathbb{E}_i on $\mathcal{B}^*(v(\eta_i) \times (0, 1)) \times \eta_i$. We need to find a subspace of $\mathcal{B}^*(v(\eta_i) \times (0, 1)) \times \eta_i$ representing $c_2(\mathbb{P}_i)$ or equivalently $(1/N^N)c_2(\mathbb{E}_i)$, and use the

inverse image of this representative to cut down the moduli space $\check{\mathcal{M}}_p(\alpha, \beta; S)$. We can proceed as in Section 2.1. The rank stratification of the vector bundle $\text{Hom}(\mathbb{C}^{N^{N-1}}, \mathbb{E}_i)$ determines a codimension-four representative $V_2(\eta_i)$ for the second Chern class of the universal bundle. For all choices of α, β, S and p , we may assume that Φ_S in (6.46) is transversal to

$$V_2(\eta_{i_1}) \times \cdots \times V_2(\eta_{i_k}).$$

Let $\check{\mathcal{N}}_p(\alpha, \beta; S)$ be the inverse image of the above space by the map Φ_S . Then the closure of $\check{\mathcal{N}}_p(\alpha, \beta; S)$ in $\check{\mathcal{M}}_p^+(\alpha, \beta; S)$, denoted by $\check{\mathcal{N}}_p^+(\alpha, \beta; S)$, is also a stratified space with smooth strata of the form

$$\prod_{i=1}^k \check{\mathcal{N}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i).$$

Lemma 6.47 *If the dimension of $\check{\mathcal{N}}_p^+(\alpha, \beta; S)$ is at most 1, then this space is compact.*

Proof This is a consequence of Remark 6.6 and the standard dimension counting argument used for the definition of the polynomial invariants for closed 4-manifolds. \square

Now we are in a position to define the differential $d_{N,j}$. Firstly consider the operator

$$d_S: \mathfrak{C}_*^{\pi_j}(Y, \gamma) \rightarrow \mathfrak{C}_*^{\pi_j}(Y, \gamma), \quad d_S(\alpha) = \sum_{p: \alpha \rightarrow \beta} \#\check{\mathcal{N}}_p^+(\alpha, \beta; S) \cdot \beta,$$

where the sum is over all paths p from α to other critical points of CS_{π_j} such that the dimension of the moduli space $\mathcal{M}_p(\alpha, \beta)$ is equal to $2|S| + 1$. The cut-down moduli space $\check{\mathcal{N}}_p^+(\alpha, \beta; S)$ is 0-dimensional and hence it is compact by Lemma 6.47. Finally, the Fukaya–Floer differential $d_{N,j}$ is defined as

$$d_{N,j} := \sum_{S \subset \mathbb{N}^+} \frac{\prod_{i \in S} t_{2,i}}{N^{N|S|}} d_S.$$

The term $N^{N|S|}$ appears in the above expression due to the relationship between $c_2(\mathbb{P}_i)$ and $c_2(\mathbb{E}_i)$.

Proposition 6.48 *The map $d_{N,j}$ defines a differential; ie $d_{N,j}^2 = 0$.*

Proof For critical points α and β of CS_{π_j} and $S \subset \mathbb{N}$, let $h_S(\alpha, \beta)$ be a number that satisfies the identity

$$d_{N,j}^2 \alpha = \sum_{\beta, S} \frac{m_S(\alpha, \beta)}{N^{N|S|}} \left(\prod_{i \in S} t_{2,i} \right) \beta.$$

Fix α, β and $S \subseteq \{1, \dots, k\}$, and suppose $p: \alpha \rightarrow \beta$ is a path such that $\dim(\mathcal{M}_p(\alpha, \beta))$ is equal to $2|S| + 2$. Then the term $m_S(\alpha, \beta)$ is equal to

$$m_S(\alpha, \beta) = \sum_{\substack{p_1: \alpha \rightarrow \gamma \\ p_2: \gamma \rightarrow \beta}} \sum_{S_1, S_2} \#\check{\mathcal{N}}_{p_1}^+(\alpha, \gamma; S_1) \cdot \#\check{\mathcal{N}}_{p_2}^+(\gamma, \beta; S_2)$$

with $p_1 \circ p_2 = p$, $S_1 \cup S_2 = S$, S_1 and S_2 disjoint,

$$\dim(\mathcal{M}_{p_1}(\alpha, \gamma)) = 2|S_1| + 1 \quad \text{and} \quad \dim(\mathcal{M}_{p_2}(\gamma, \beta)) = 2|S_2| + 1.$$

Therefore, $m_S(\alpha, \beta)$ is equal to the signed count of the boundary points of the compact 1-manifold $\check{\mathcal{N}}_p^+(\alpha, \beta; S)$. This implies that $m_S(\alpha, \beta)$ is equal to zero. \square

The above discussion can be generalized in a straightforward way to the case of an arbitrary $(N-1)$ -tuple (l_2, \dots, l_N) . For each l_k , we choose a sequence of disjoint representatives $\{\eta_{k,i}\}_{i \in \mathbb{N}}$ and the open neighborhoods $\nu(\eta_{k,i})$. We keep assuming that $\eta_{k,i}$ is an oriented simple closed curve in Y , the open sets $\nu(\eta_{k,i})$ are disjoint and the inclusion of $\nu(\eta_{k,i})$ in Y induces a surjective map at the level of fundamental groups. In a more general case that l_i is a homology class with complex coefficients, we need to consider the straightforward generalization that $\eta_{k,i}$ is a linear combination of disjoint closed curves. Then for each $(N-1)$ -tuple $\bar{S} = (S_2, \dots, S_N)$ of subsets of $\{1, 2, \dots, j\}$, we can form the moduli space $\check{\mathcal{M}}_p(\alpha, \beta; \bar{S})$ and its partial compactification $\check{\mathcal{M}}_p^+(\alpha, \beta; \bar{S})$. As in the previous case, each stratum of the partial compactification is a product of moduli spaces of the form $\check{\mathcal{M}}_p(\alpha, \beta; \bar{S})$. The next step is to cut down $\check{\mathcal{M}}_p^+(\alpha, \beta; \bar{S})$ with divisors $V_k(\eta_{k,i})$ for $i \in S_k$. As in the case of closed 4-manifolds, we can use the auxiliary complex vector bundles of rank N^N associated to the universal $\text{PU}(N)$ -bundle and construct geometric representatives for $V_k(\eta_{k,i})$. The desired representatives are linear combinations of codimension $2k$ stratified spaces with even-dimensional strata. We will write $\check{\mathcal{N}}_p^+(\alpha, \beta; \bar{S})$ for the cut-down moduli space. A generic choice of these divisors allow us to obtain a cut-down moduli space. This space is compact when its dimension is less than or equal to 1.

The 0-dimensional cut-down moduli spaces $\check{\mathcal{N}}_p^+(\alpha, \beta; \bar{S})$ can be used to define an operator $d_{\bar{S}}$ acting on $\mathcal{C}_*^{\pi_j}(Y, \gamma)$:

$$d_{\bar{S}}\alpha = \sum_{p: \alpha \rightarrow \beta} \#\check{\mathcal{N}}_p^+(\alpha, \beta; \bar{S}) \cdot \beta.$$

We combine these operators to form

$$d_{N,j} := \sum_{\bar{S}=(S_2, \dots, S_N)} \left(\prod_{i \in S_k} t_{k,i} \right) d_{\bar{S}}.$$

As in the previous case, the 1–dimensional moduli spaces can be used to show that $d_{N,j}^2 = 0$.

Before giving the definition of the maps F_j^k , we study the functorial properties of the chain complex $(\mathcal{C}_*^{N,j}(Y, \gamma, L), d_{N,j})$. Let (X, w) be a pair whose boundary is (Y, γ) . Suppose that $\Gamma^2, \dots, \Gamma^N$ are properly embedded surfaces in X such that $[\partial\Gamma^i] = l_i \in H_1(Y)$. For $z \in \mathbb{A}(X)^{N-1}$, we want to define the relative invariant

$$D_{X,w}^{N,j}(ze^{\sum_i \Gamma^i}) \in \mathbb{I}_*^{N,j}(Y, \gamma, L),$$

where $L = (l_2, \dots, l_N)$. This is similar to the definition of the differential of the Fukaya–Floer chain complex. For simplicity of exposition, we assume that $\Gamma^3 = \dots = \Gamma^N = 0$, and $z = 1$. As in the case of the definition of the differential in Fukaya–Floer homology, the more general case is just slightly different. In the manifold X^+ with a cylindrical end, let $\{\Sigma_i\}_{i \in \mathbb{N}}$ be a sequence of surfaces which are given by perturbing the surface Γ^2 . We assume that these surfaces intersect generically and their intersections with $\mathbb{R}^{\geq 0} \times Y$ are equal to the disjoint product surfaces $\{\eta_i \times \mathbb{R}^{\geq 0}\}_{i \in \mathbb{N}}$. We choose a holonomy perturbation of the ASD equation on X^+ , compatible with the chosen perturbation CS_{π_j} of the Chern–Simons functional of Y , such that all moduli spaces $\mathcal{M}_p(X, w; \beta)$ consists of regular points. Given a subset $S = \{i_1, \dots, i_k\} \subset \{1, \dots, j\}$ and a path p along (X, w) based at the critical point β of CS_{π_j} , consider the space

(6.49)
$$\mathcal{M}_p(X, w; \beta, S) := \mathcal{M}_p(X, w; \beta) \times \Sigma_{i_1} \times \dots \times \Sigma_{i_k}.$$

For $S' \subset S$, the obvious projection map from $\mathcal{M}_p(X, w; \beta, S)$ to $\mathcal{M}_p(X, w; \beta, S')$ is denoted by $\pi_{S \rightarrow S'}$. There is also a partially defined translation map for $\mathcal{M}_p(X, w; \beta, S)$. Suppose $\mathcal{M}_p^{\text{cyl}}(X, w; \beta, S)$ denotes the following subset of $\mathcal{M}_p(X, w; \beta, S)$:

$$\mathcal{M}_p(X, w; \beta) \times (\eta_{j_1} \times \mathbb{R}^{\geq 0}) \times \dots \times (\eta_{j_k} \times \mathbb{R}^{\geq 0}).$$

For $u = ([A], (x_1, t_1), \dots, (x_k, t_k)) \in \mathcal{M}_p^{\text{cyl}}(X, w; \beta, S)$ define $\sigma_t(u)$ to be the element

$$([T_t^*(A|_{Y \times \mathbb{R}^{\geq 0}})], (x_1, t_1 - t), \dots, (x_k, t_k - t)).$$

Note that the connection $T_t^*(A|_{Y \times \mathbb{R}^{\geq 0}})$ is partially defined on the cylinder $Y \times \mathbb{R}$.

As in the cylinder case, we form a partial compactification of $\mathcal{M}_p(X, w; \beta, S)$ given by

$$\mathcal{M}_p^+(X, w; \beta, S) := \bigcup_k \bigcup_{\{(p_i, S_i)\}_{1 \leq i \leq k}} \mathcal{M}_{p_1}(X, w, \alpha_1, S_1) \times \prod_{i=2}^k \check{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i),$$

where $\alpha_1, \dots, \alpha_k$ are critical points of CS_{π_j} , $\alpha_k = \beta$, p_1 is a path along (X, w) , $p_i: \alpha_{i-1} \rightarrow \alpha_i$ is a path along the cylinder, $p = p_1 \circ \dots \circ p_k$, the sets S_i are disjoint and their union is equal to S . As before, S_i may be empty. In a little more detail, a sequence $u_l \in \mathcal{M}_p(X, w; \beta, S)$ is chain convergent to $u_\infty = (u_\infty^1, \dots, u_\infty^k) \in \mathcal{M}_{p_1}(X, w, \alpha_1, S_1) \times \prod_{i=2}^k \check{\mathcal{M}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i)$ if there is a sequence of $(k-1)$ -tuples of real numbers (t_l^2, \dots, t_l^k) such that

$$t_l^1 := 0 < t_l^2 < \dots < t_l^k, \quad \lim_{l \rightarrow \infty} t_l^{m+1} - t_l^m = \infty \quad \text{for all } 1 \leq m \leq k-1$$

and

$$\pi_{S \rightarrow S_1}(u_l) \xrightarrow{C_{loc}^\infty} u_\infty^1, \quad \sigma_{t_l^i} \circ \pi_{S \rightarrow S_i}(u_l) \xrightarrow{C_{loc}^\infty} u_\infty^i \quad \text{for } i \geq 2.$$

Here part of the assumption is that $\sigma_{t_l^i} \circ \pi_{S \rightarrow S_i}(u_l)$ is well defined for $i \geq 2$. That is to say, $\pi_{S \rightarrow S_i}(u_l) \in \mathcal{M}_p^{cyl}(X, w; \beta, S_i)$.

We momentarily assume that $S = \{1\}$. Suppose $\nu(\Sigma_1)$ is an open neighborhood of Σ_1 such that the inclusion map of $\nu(\Sigma_1)$ induces a surjective map. Suppose also that $\mathcal{B}^{**}(\nu(\Sigma_1))$ is the set of connections on $\nu(\Sigma_1)$ whose restrictions to the sets of the form $\nu(\eta_1) \times (t-1, t)$ are irreducible. We can assume that the perturbation π_j is small enough that the restriction map $r_1: \mathcal{M}_p(X, w; \beta, S) \rightarrow \mathcal{B}^{**}(\nu(\Sigma_1)) \times \Sigma_1$ is well defined.¹¹ We can form a universal $PU(N)$ -bundle \mathbb{P}_1 and the associated $SU(N^N)$ -bundle \mathbb{E}_1 on $\mathcal{B}^{**}(\nu(\Sigma_1)) \times \Sigma_1$. Our goal is to define a geometric representative for $c_2(\mathbb{P}_1)$ or equivalently $(1/N^N)c_2(\mathbb{E}_1)$ which is compatible with our choice of the geometric representative in the case of cylinders. Note that $\mathcal{M}_p(X, w; \beta, S)$ is the union of the two sets

$$B_1 = \mathcal{M}_p(X, w; \beta) \times (\eta_1 \times [1, \infty)) \quad \text{and} \quad B_2 = \mathcal{M}_p(X, w; \beta) \times (\Sigma_1 \setminus (\eta_1 \times (2, \infty))).$$

The map $r_1|_{B_1}$ can be composed with the map

$$F: \mathcal{B}^{**}(\nu(\Sigma_1)) \times (\eta_1 \times [1, \infty)) \rightarrow \mathcal{B}^*(\nu(\eta_1) \times (0, 1)) \times \eta_1, \\ F([A], (x, t)) = ([A|_{\nu(\Sigma) \times (t-1, t)}], x).$$

Therefore, we can choose the geometric representative $V_2(\Sigma_1) \subset \mathcal{B}^{**}(\nu(\Sigma_1)) \times \Sigma_1$ for $c_2(\mathbb{E}_1)$ such that

$$V_2(\Sigma_1) \cap (\mathcal{B}^{**}(\nu(\Sigma_1)) \times (\eta_1 \times [1, \infty))) = F^{-1}(V_2(\eta_1)).$$

Arguing as in [52], we can also assume that $V_2(\Sigma_1)$ is transversal to the map r_1 for all choices of the path p . In this process, first we slightly modify $V_2(\eta)$ to make

¹¹This is again a consequence of unique continuation and the Uhlenbeck compactness theorem [52].

the map $r_1|_{B_1}$ transversal. Then we extend $F^{-1}(V_2(\eta_1))$ to $\mathcal{B}^{**}(v(\Sigma_1)) \times \Sigma_1$ so that $r_1|_{B_2}$ is also transversal. We will write $\mathcal{N}_p(X, w; \beta, S)$ for the cut-down moduli space $r_1^{-1}(V_2(\Sigma_1))$. A similar construction can be used to define $\mathcal{N}_p(X, w; \beta, S)$ in the case that S has more than one element. The closure of $\mathcal{N}_p(X, w; \beta, S)$ in $\mathcal{M}_p^+(X, w; \beta, S)$, denoted by $\mathcal{N}_p^+(X, w; \beta, S)$, is also a stratified space with smooth strata of the form

$$\mathcal{N}_{p_1}(X, w, \alpha_1, S_1) \times \prod_{i=2}^k \check{\mathcal{N}}_{p_i}(\alpha_{i-1}, \alpha_i; S_i).$$

As in the cylinder case, the cut-down moduli space $\mathcal{N}_p^+(X, w; \beta, S)$ is compact when its dimension is at most one. The relative invariant $D_{X,w}^N(e^{\Gamma^{(2)}})$ is defined using 0-dimensional moduli spaces as below:

$$(6.50) \quad D_{X,w}^{N,j}(e^{\Gamma^{(2)}}) := \sum_{S \subset \{1, \dots, j\}} \frac{\prod_{i \in S} t_{2,i}}{N^{|S|}} (\#\mathcal{N}_p^+(X, w; \beta, S)) \cdot \beta \in \mathfrak{C}_*^{N,j}(Y, \gamma, L).$$

Following the proof of Proposition 6.48, we can show that (6.50) determines a cycle in $\mathfrak{C}_*^{N,j}(Y, \gamma, L)$. We will denote the corresponding element in $\mathbb{I}_*^{N,j}(Y, \gamma, L)$ with the same notation.

The definition of the relative element in (6.50) can be extended to similar situations. For example, suppose (Y_0, γ_0) and (Y_1, γ_1) are two N -admissible pairs and $L_i = (l_i^2, \dots, l_i^N)$ is an $(N-1)$ -tuple of the elements in $H_1(Y_i)$. Suppose also (W, w, z) is a morphism from (Y_0, γ_0) to (Y_1, γ_1) , and Γ^j is a properly embedded surface in W such that $[\partial_i \Gamma^j] = l_i^j$. Then there is a chain map,

$$\mathfrak{C}_*^{N,j}(W, w, ze^{\Gamma^{(2)} + \dots + \Gamma_{(N)}^N}): \mathfrak{C}_*^{N,j}(Y_0, \gamma_0, L_0) \rightarrow \mathfrak{C}_*^{N,j}(Y_1, \gamma_1, L_1),$$

which induces a map at the level of homology,

$$\mathbb{I}_*^{N,j}(W, w, ze^{\Gamma^{(2)} + \dots + \Gamma_{(N)}^N}): \mathbb{I}_*^{N,j}(Y_0, \gamma_0, L_0) \rightarrow \mathbb{I}_*^{N,j}(Y_1, \gamma_1, L_1).$$

Alternatively, if (Y, γ, L) is as above, (X, w) is a 4-manifold whose boundary is equal to $(\bar{Y}, \bar{\gamma})$, and Γ^j is an embedded surface such that $[\partial \Gamma^j] = -l_j$, then we have an $R_{N,j}$ -linear map

$$D_{N,j}^{X,w}(ze^{\Gamma^{(2)} + \dots + \Gamma_{(N)}^N}): \mathbb{I}_*^{N,j}(Y, \gamma, L) \rightarrow R_{N,j}.$$

Now we are ready to define the maps F_j^k for a triple (Y, γ, L) . For a pair of nonnegative integers j and k with $j \geq k$, we have chosen perturbations CS_{π_j} and CS_{π_k} of the Chern-Simons functional of the pair (Y, γ) . Since $j \geq k$, the functional CS_{π_j}

satisfies all the properties that we required for CS_{π_k} . In particular, the chain complex $(\mathfrak{C}_*^{N,j}(Y, \gamma, L) \otimes R_{N,k}, d_{N,j})$ gives an alternative chain complex to define $\mathbb{I}_*^{N,k}(Y, \gamma, L)$. The functoriality mentioned in the previous paragraph implies that there is a chain map

$$\mathfrak{C}^{N,j}([0, 1] \times Y, [0, 1] \times \gamma, e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}): \mathfrak{C}_*^{\pi_j}(Y, \gamma) \otimes R_{N,k} \rightarrow \mathfrak{C}_*^{\pi_k}(Y, \gamma) \otimes R_{N,k},$$

where Γ^i is the surface $[0, 1] \times \eta_{i,1}$. The map F_j^k is the composition of the above map and the obvious map from $\mathfrak{C}_*^{\pi_j}(Y, \gamma) \otimes R_{N,j}$ to $\mathfrak{C}_*^{\pi_j}(Y, \gamma) \otimes R_{N,k}$. A standard argument shows that $F_k^l \circ F_j^k$ is chain homotopy equivalent to F_j^l for any triple $j \geq k \geq l \geq 0$. Therefore, the maps f_j^k , induced by the maps F_j^k , have the required properties for an inverse system. This completes the definition of Fukaya–Floer homology $\mathbb{I}_*^N(Y, \gamma, L)$ for the triple. It is also standard to show that this R_N –module is independent of the choices that were made. The homomorphism f_j^k maps $D_{X,w}^{N,j}(ze^{\sum_i \Gamma^i}) \in \mathbb{I}_*^{N,j}(Y, \gamma, L)$ to $D_{X,w}^{N,k}(ze^{\sum_i \Gamma^i}) \in \mathbb{I}_*^{N,k}(Y, \gamma, L)$. Thus we have an induced element of $\mathbb{I}_*^N(Y, \gamma, L)$, which we denote by $D_{X,w}^N(ze^{\sum_i \Gamma^i})$. Similarly, we can use the construction of the previous paragraph to define $\mathbb{I}_*^N(W, w, ze^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N})$ and $D_N^{X,w}(ze^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N})$.

Now we are ready to give a proof of (3.30). For the convenience of the reader, we restate the claim as the following proposition:

Proposition 6.51 *Let (X_1, w_1) be a pair whose boundary is equal to an admissible pair (Y, γ) . Let (X_2, w_2) be another pair whose boundary is equal to $(\bar{Y}, \bar{\gamma})$. Let $z_1 \in \mathbb{A}(X_1)^{\otimes(N-1)}$ and $z_2 \in \mathbb{A}(X_2)^{\otimes(N-1)}$. Let Γ^j be a properly embedded surface in X_1 with boundary l_j . Let Λ^j be a properly embedded surface in X_2 whose boundary is equal to l_j with the reverse orientation. Then we can form the closed 4–manifold $X_2 \circ X_1$ and the 2–cycle $w_2 \circ w_1$. The embedded surfaces Γ^j and Λ^j can be glued to each other along their boundary to form a closed surface $\Gamma^j \# \Lambda^j$. Then the invariant*

$$(6.52) \quad D_{X_2 \circ X_1, w_2 \circ w_1}^N(z_1 \cdot z_2 \cdot e^{(\Gamma^2 \# \Lambda^2)_{(2)} + \dots + (\Gamma^N \# \Lambda^N)_{(N)}}) \in \mathbb{C}[[t_2, \dots, t_N]] \subset R_N$$

of the closed 4–manifold $X_2 \circ X_1$ is equal to

$$D_N^{X_2, w_2}(z_2 \cdot e^{\Lambda_{(2)}^2 + \dots + \Lambda_{(N)}^N}) \circ D_{X_1, w_1}^N(z_1 \cdot e^{\Gamma_{(2)}^2 + \dots + \Gamma_{(N)}^N}).$$

Proof We make simplifying assumptions as before: assume $z_1 = z_2 = 1$ and $\Gamma^3, \dots, \Gamma^N$ and $\Lambda^3, \dots, \Lambda^N$ are empty. Choose two series of properly embedded surfaces $\{\Sigma_i\}_{i \in \mathbb{N}} \subset X_1$ and $\{T_i\}_{i \in \mathbb{N}} \subset X_2$ such that Σ_i (respectively, T_i) is given by perturbing Γ^2 (respectively, Λ^2). We also assume that $\partial \Sigma_i$ (respectively, ∂T_i) is

equal to η_i (respectively, η_i with the reverse orientation), the curves η_i are disjoint and the embedded surfaces Σ_i and T_i intersect generically. We fix metrics on X_1 and X_2 which are product metrics in a neighborhood of their boundaries corresponding to a fixed metric on Y . For each nonnegative integer j , we prove that the image of the element in (6.52) in $R_{N,j}$ is equal to

$$D_{N,j}^{X_2,w_2}(e^{\Lambda_{(2)}^2}) \circ D_{X_1,w_1}^{N,j}(e^{\Gamma_{(2)}^2}).$$

To achieve this goal, we proceed as before to define open sets $v(\eta_i) \subset Y$, $v(\Sigma_i) \subset X_1^+$ and $v(T_i) \subset X_2^+$ and the geometric representatives

$$\begin{aligned} V_2(\eta_i) &\subset \mathcal{B}^*(v(\eta_i) \times (0, 1)) \times \eta_i, \\ V_2(\Sigma_i) &\subset \mathcal{B}^{**}(v(\Sigma_i)) \times \Sigma_i, \\ V_2(T_i) &\subset \mathcal{B}^{**}(v(T_i)) \times T_i \end{aligned}$$

suitable for the definition of the differential $d_{N,j}$ of the Fukaya–Floer chain complex and the relative elements $D_{X_1,w_1}^{N,j}(e^{\Gamma_{(2)}^2})$ and $D_{N,j}^{X_2,w_2}(e^{\Lambda_{(2)}^2})$.

Suppose X^T is the metric on $X_2 \circ X_1$ induced by the metrics on X_1 and X_2 with a neck of length T along Y . Suppose $w \subset X^T$ is the 2–cycle induced by w_1 and w_2 , and $R_i^T \subset X^T$ is the embedded surface induced by the surfaces Σ_i and T_i . The 4–manifold X can be decomposed into the three pieces

$$X_1, \quad X_2 \quad \text{and} \quad Y \times \left[-\frac{1}{2}T, \frac{1}{2}T\right].$$

We assume that X_1 and X_2 are disjoint and that they intersect $Y \times \left[-\frac{1}{2}T, \frac{1}{2}T\right]$ in $Y \times \left[-\frac{1}{2}T, -\frac{1}{2}T + 2\right]$ and $Y \times \left[\frac{1}{2}T - 2, \frac{1}{2}T\right]$, respectively. The Riemann surface R_i^T can be decomposed into a union of the three sets

$$\Sigma_i^c \subset X_1, \quad T_i^c \subset X_2 \quad \text{and} \quad \eta_i \times \left[-\frac{1}{2}T + 1, \frac{1}{2}T - 1\right] \subset Y \times \left[-\frac{1}{2}T + 1, \frac{1}{2}T - 1\right].$$

Suppose $\mathcal{B}_\kappa^{**}(X^T, w)$ is the subset of $\mathcal{B}_\kappa^*(X, w)$ which consists of connections whose restriction to any set of the following form is irreducible:

$$v(\eta_i) \times (t - 1, t), \quad -T + 1 < t < T.$$

Suppose κ is chosen such that the dimension of the moduli space $\mathcal{M}_\kappa(X^T, w)$ is at most $2j$. Then unique continuation and gluing theory show that for large enough values of T , the moduli space $\mathcal{M}_\kappa(X^T, w)$ is a subset of $\mathcal{B}_\kappa^{**}(X^T, w)$. Similar to the case of 4–manifolds with cylindrical ends, $V_2(\eta_i)$, $V_2(\Sigma_i)$ and $V_2(T_i)$ can be used to define a geometric representative $V(R_i^T) \subset \mathcal{B}_\kappa^{**}(X^T, w) \times v(R_i^T)$. Another application

of gluing theory shows that for large enough values of T , these divisors determine a transversal cut of $\mathcal{M}_\kappa(X^T, w) \times R_{i_1}^T \times \cdots \times R_{i_k}^T$ for any set $S = \{i_1, \dots, i_k\} \subset \{1, \dots, j\}$. In the case that the cut-down moduli space is zero-dimensional, it can be identified, for large values of T , with the set

$$\bigcup_{p_1, p_2, S_1, S_2} \mathcal{N}_{p_1}(X_1, w_1; \alpha, S_1) \times \mathcal{N}_{p_2}(X_2, w_2; \alpha, S_2),$$

where each p_i is a loop along (X_i, w_i) based at the connection α such that $\kappa = \kappa(p_1) + \kappa(p_2)$. Moreover, S_1 and S_2 are disjoint sets with $S = S_1 \cup S_2$. These geometric results for different choices of S can be translated to the algebraic identity

$$\begin{aligned} (6.53) \quad D_{N,j}^{X_2, w_2}(e^{\Lambda^2_{(2)}}) \circ D_{X_1, w_1}^{N, j}(e^{\Gamma^2_{(2)}}) &= \sum_{S \subset \{1, \dots, j\}} \left(\prod_{i \in S} t_{2,i} \right) D_{X^T, w}^N \left(\prod_{i \in S} (R_i^T)_{(2)} \right) \\ &= \sum_{S \subset \{1, \dots, j\}} \left(\prod_{i \in S} t_{2,i} \right) D_{X^T, w}^N \left((\Gamma^2 \# \Lambda^2)_{(2)}^{|S|} \right). \end{aligned}$$

In the second equality, we used the fact that R_j^T represents the homology class of $\Gamma^2 \# \Lambda^2$. The term in (6.53) is equal to the image of $D_{X^T, w}^N(e^{(\Gamma^2 \# \Lambda^2)_{(2)}})$ in $R_{N,j}$. \square

7 Questions and conjectures

In this section, we propose some questions and conjectures for future directions. This section is divided into two parts: the first subsection is concerned with the polynomial invariants of 4-manifolds. In the second part, we discuss some conjectures related to the algebra $\mathbb{V}_{g,d}^N$.

7.1 Structure of polynomial invariants and 4-manifolds with simple type

In Section 2.5, the definition of the simple type property of 4-manifolds uses $U(3)$ -polynomial invariants. As we pointed out earlier, the definition is motivated by Kronheimer and Mrowka’s simple type property, defined using $U(2)$ -polynomial invariants [54]. There is another version of simple type property defined by Seiberg–Witten invariants. It is unknown whether there are examples of smooth 4-manifolds with $b^+ \geq 2$ which do not have Kronheimer–Mrowka simple type or Seiberg–Witten simple type. It is also shown in [23] that many 4-manifolds with Seiberg–Witten simple type have Kronheimer–Mrowka simple type. Therefore, it is natural to ask whether

there is any relationship among $U(3)$ -simple type, Kronheimer–Mrowka simple type and Seiberg–Witten simple type. A more challenging question would be to investigate whether there is a 4-manifold with $b^+ \geq 2$ which does not have $U(3)$ -simple type. A more approachable question is the following:

Question 7.1 *What is the analogue of the simple type condition with respect to $U(N)$ -polynomial invariants?*

As in the $U(2)$ and the $U(3)$ case, the simple condition has to be formulated in terms of point classes. In light of [Proposition 3.24](#), it is plausible that one of the required conditions is

$$D_{X,w}^N(a_2^N z) = N^N D_{X,w}^N(z).$$

For $N = 2, 3$, the blowup formula for $U(N)$ simple type manifolds has a simpler form [\[26; 12\]](#). One might hope that the same holds for higher values of N and follow this direction to gain more insights into the correct definition for the simple type condition. The physics literature [\[64; 22\]](#) suggests that the blowup formula for an arbitrary N is related to function theory on a hyperelliptic curve with coefficients in $\mathbb{Q}[a_2, \dots, a_N]$. Evaluation of (a_2, \dots, a_N) determines a hyperelliptic curve on the complex numbers, and the simple type condition is related to the evaluations that produce a fully degenerate curve.

The relationship between $U(2)$ -polynomial invariants and Seiberg–Witten invariants goes beyond the simple type conditions. In [\[54\]](#), Kronheimer and Mrowka prove that $U(2)$ -polynomial invariants are completely determined by a finite set of cohomology classes (known as Kronheimer–Mrowka basic classes) and a set of rational numbers, one for each basic class. In [\[84\]](#), Witten argues that basic classes and corresponding rational numbers can be determined in terms of Seiberg–Witten invariants. To be more detailed, recall that for each spin^c structure \mathfrak{s} on a 4-manifold X (satisfying appropriate conditions, such as $b^+(X) \geq 2$) there is a Seiberg–Witten invariant $\text{SW}_X(\mathfrak{s})$. Then Witten’s conjecture states that any basic class of X is equal to $c_1(S_{\mathfrak{s}}^+)$, where \mathfrak{s} is a spin^c structure with nonzero $\text{SW}_X(\mathfrak{s})$ and $S_{\mathfrak{s}}^+$ is the half-spin bundle associated to \mathfrak{s} . Witten’s conjecture is generalized to higher values of N in [\[64\]](#). The calculations of $U(3)$ -polynomial invariants in this paper agree with the Moore–Mariño conjecture in [\[64\]](#) and can be exploited to fix the undetermined constants in this conjecture. In particular, a modified version of the Moore–Mariño conjecture runs as follows:

Conjecture 7.2 *Let X be a four-manifold which has $U(3)$ -simple type. Let $\{K_i\}$ be the set of Kronheimer–Mrowka basic classes. Then the $U(3)$ -series of X has the form*

$$\widehat{D}_{X,w}(e^{\Gamma(2)+\Lambda(3)}) = e^{\frac{1}{2}Q(\Gamma)-Q(\Lambda)} \sum_{i,j} c_{ij} \zeta^{-w \cdot (\frac{1}{2}(K_i-K_j))} e^{\frac{\sqrt{3}}{2}(K_i+K_j) \cdot \Gamma + \frac{\sqrt{3}}{2} \mathbf{i}(K_i-K_j) \cdot \Lambda},$$

where $c_{i,j}$ is given as

$$2^{\chi+\frac{3}{2}\sigma+\frac{1}{2}K_i \cdot K_j} 3^{2+\frac{7}{4}\chi+\frac{11}{4}\sigma} SW_X(\mathfrak{s}_i) SW_X(\mathfrak{s}_j).$$

Here \mathfrak{s}_i is chosen such that the associated basic class is equal to K_i .

7.2 The algebra $\mathbb{V}_{g,d}^N$

In Section 5.1, a list of simultaneous eigenvectors for the operators ϵ , \mathfrak{N}_2 , \mathfrak{N}_3 , $\rho_{(2)}$ and $\rho_{(3)}$, acting on $\mathbb{V}_{g,d}^3$, is constructed. We also showed that there are at least two nondegenerate simultaneous eigenvectors, which form the essential ingredient to prove the excision theorem in Section 5.2. However, we do not know whether our approach produces all simultaneous eigenvectors:

Conjecture 7.3 *Suppose $V_{g,d}^N \subset \mathbb{V}_{g,d}^N$ is the set of vectors which are invariant with respect to the action of ϵ . Then for $N = 3$, any simultaneous eigenvector of the operators acting on $V_{g,d}^N$ that are induced by \mathfrak{N}_2 , \mathfrak{N}_3 , $\rho_{(2)}$ and $\rho_{(3)}$ has the form $(3\zeta^{2d\beta}, 0, \zeta^{d\beta} \sqrt{3}\alpha, \zeta^{2d\beta} \sqrt{3}\mathbf{i}\beta)$ with $(\alpha, \beta) \in C_g$.*

For $N = 2$, there are three operators ϵ , \mathfrak{N}_2 and $\rho_{(2)}$. In this case, Muñoz obtains a complete understanding of the action of ϵ , \mathfrak{N}_2 and $\rho_{(2)}$ in [73]. In particular, his results show that the simultaneous eigenvectors of \mathfrak{N}_2 and $\rho_{(2)}$, acting on $V_{g,d}^2$, have the form

$$((-1)^r 2, \pm 2r \mathbf{i}^{r+1}), \quad 0 \leq r \leq g - 1.$$

All of these eigenvalues can be produced using the method of Proposition 5.7. Therefore, the analogue of Conjecture 7.3 holds for $N = 2$. Muñoz’s method of understanding the action of ϵ , \mathfrak{N}_2 and $\rho_{(2)}$ is based on the characterization of the ring structure of the cohomology ring $\mathcal{N}_{2,d}(\Sigma_g)$ in [86; 47; 78; 3], which is not available for higher values of N .

If Conjecture 7.3 holds, then we can use the method of [56] and show that $\text{SHI}_*(M, \alpha)$ is nonzero for a taut balanced sutured manifold [46; 56]. This nonvanishing result can be used to show that Conjecture 5.39 holds. We can also use this to show that

$\text{KHI}_*(K)$ detects the genus of K . Thus the answer to [Question 1.1](#) for a nontrivial knot and $N = 3$ is positive. In fact, in order to make this series of conclusions, we need the following weaker version of [Conjecture 7.3](#):

Conjecture 7.4 *If (x, y) is a pair of simultaneous eigenvalues for $(\rho_{(2)}, \rho_{(3)})$ in $\mathbb{V}_{g,d}$, then $|x| + |y| \leq \sqrt{3}(2g - 2)$.*

There is also a symplectic analogue of the algebra $\mathbb{V}_{g,d}^N$. The manifold $\mathcal{N}_{N,d}(\Sigma_g)$ is Kähler and the associated Gromov-Witten invariants can be used to define the *quantum cohomology* ring $\text{QH}^*(\mathcal{N}_{N,d})$ [75; 65]. The underlying vector space of $\text{QH}^*(\mathcal{N}_{N,d})$ is $H^*(\mathcal{N}_{N,d})$ and the ring structure is also a deformation of the cup product. Thus, it has similar structure to $V_{g,d}^N = \ker(\epsilon - 1)$, and it is natural to make the following conjecture:

Conjecture 7.5 *The ring $V_{g,d}^N$ is isomorphic to $\text{QH}^*(\mathcal{N}_{N,d})$.*

This conjecture for $N = 2$ is proved by Muñoz [72] using the characterization of the cohomology ring $H^*(\mathcal{N}_{N,d})$ in [86; 47; 78; 3]. The $N = 2$ special case of this conjecture was also proved using an adiabatic limit argument in [77].

Appendix Invariants of flat connections on $\Sigma(2, 3, 23)$

There are 44 irreducible flat $\text{SU}(3)$ -connections on $\Sigma(2, 3, 23)$ [4]. These flat connections are determined by their holonomies along the loop x_3 in the standard presentation of the fundamental group of $\Sigma(2, 3, 23)$ (see (3.37)). For each flat connection, the conjugacy class of this holonomy is determined by its eigenvalues which have the form $e^{2\pi i k/23}$, $e^{2\pi i l/23}$ and $e^{2\pi i m/23}$. The possible values of $\{k, l, m\}$ are given in [Table 4](#).

The complex conjugation diffeomorphism of $\Sigma(2, 3, 23)$ maps a flat connection with the associated triple $\{k, l, m\}$ to the flat connection with the complementary associated triple $\{23 - k, 23 - l, 23 - m\}$. If both these triples give the same flat connection, we denote this connection with α_j for an appropriate choice of the integer j . Otherwise, the resulting connections are denoted by α_j^1 and α_j^2 .

The gauge-theoretical invariants of these flat connections are also given in [Table 4](#).

There are eight nontrivial flat $\text{SU}(2)$ -connections on $\Sigma(2, 3, 23)$. As in the irreducible case, these connections are determined by the conjugacy class of their holonomies along x_3 . For each $2 \leq k \leq 9$, there is a unique flat $\text{SU}(2)$ -connection on $\Sigma(2, 3, 23)$ where the eigenvalues of holonomy along x_3 are equal to $e^{2\pi i k/23}$ and $e^{-2\pi i k/23}$. We

holonomy							holonomy						
α	k	l	m	$CS(\alpha)$	ρ_{ad_α}	deg	α	k	l	m	$CS(\alpha)$	ρ_{ad_α}	deg
α_1	0	4	19	$\frac{1}{138}$	$-\frac{364}{23}$	4	α_{16}^1	2	4	17	$\frac{19}{138}$	$-\frac{476}{23}$	8
α_2	0	5	18	$\frac{49}{138}$	$-\frac{540}{23}$	0	α_{16}^2	6	19	21	$\frac{19}{138}$	$-\frac{476}{23}$	8
α_3	0	6	17	$\frac{31}{138}$	$-\frac{520}{23}$	10	α_{17}^1	2	5	16	$\frac{1}{138}$	$-\frac{456}{23}$	6
α_4	0	7	16	$\frac{85}{138}$	$-\frac{488}{23}$	2	α_{17}^2	7	18	21	$\frac{1}{138}$	$-\frac{456}{23}$	6
α_5	0	8	15	$\frac{73}{138}$	$-\frac{444}{23}$	0	α_{18}^1	2	6	15	$\frac{55}{138}$	$-\frac{516}{23}$	0
α_6	0	9	14	$\frac{133}{138}$	$-\frac{572}{23}$	8	α_{18}^2	8	17	21	$\frac{55}{138}$	$-\frac{516}{23}$	0
α_7	0	10	13	$\frac{127}{138}$	$-\frac{504}{23}$	6	α_{19}^1	2	7	14	$\frac{43}{138}$	$-\frac{472}{23}$	10
α_8	0	11	12	$\frac{55}{138}$	$-\frac{424}{23}$	10	α_{19}^2	9	16	21	$\frac{43}{138}$	$-\frac{472}{23}$	10
α_9^1	1	4	18	$-\frac{43}{138}$	$-\frac{472}{23}$	10	α_{20}^1	2	8	13	$\frac{103}{138}$	$-\frac{508}{23}$	4
α_9^2	5	19	22	$-\frac{43}{138}$	$-\frac{472}{23}$	10	α_{20}^2	10	15	21	$\frac{103}{138}$	$-\frac{508}{23}$	4
α_{10}^1	1	5	17	$\frac{127}{138}$	$-\frac{412}{23}$	4	α_{21}^1	2	9	12	$\frac{97}{138}$	$-\frac{440}{23}$	2
α_{10}^2	6	18	22	$\frac{127}{138}$	$-\frac{412}{23}$	4	α_{21}^2	11	14	21	$\frac{97}{138}$	$-\frac{440}{23}$	2
α_{11}^1	1	6	16	$\frac{7}{138}$	$-\frac{524}{23}$	8	α_{22}^1	3	4	16	$\frac{67}{138}$	$-\frac{468}{23}$	0
α_{11}^2	7	17	22	$\frac{7}{138}$	$-\frac{524}{23}$	8	α_{22}^2	7	19	20	$\frac{67}{138}$	$-\frac{468}{23}$	0
α_{12}^1	1	7	15	$\frac{97}{138}$	$-\frac{532}{23}$	4	α_{23}^1	3	5	15	$\frac{85}{138}$	$-\frac{488}{23}$	2
α_{12}^2	8	16	22	$\frac{97}{138}$	$-\frac{532}{23}$	4	α_{23}^2	8	18	20	$\frac{85}{138}$	$-\frac{488}{23}$	2
α_{13}^1	1	8	14	$\frac{121}{138}$	$-\frac{528}{23}$	6	α_{24}^1	3	6	14	$\frac{37}{138}$	$-\frac{404}{23}$	8
α_{13}^2	9	15	22	$\frac{121}{138}$	$-\frac{528}{23}$	6	α_{24}^2	9	17	20	$\frac{37}{138}$	$-\frac{404}{23}$	8
α_{14}^1	1	9	13	$\frac{79}{138}$	$-\frac{420}{23}$	0	α_{25}^1	3	7	13	$\frac{61}{138}$	$-\frac{492}{23}$	0
α_{14}^2	10	14	22	$\frac{79}{138}$	$-\frac{420}{23}$	0	α_{25}^2	10	16	20	$\frac{61}{138}$	$-\frac{492}{23}$	0
α_{15}^1	1	10	12	$\frac{109}{138}$	$-\frac{484}{23}$	4	α_{26}^1	3	8	12	$\frac{19}{138}$	$-\frac{476}{23}$	8
α_{15}^2	11	13	22	$\frac{109}{138}$	$-\frac{484}{23}$	4	α_{26}^2	11	15	20	$\frac{19}{138}$	$-\frac{476}{23}$	8

Table 4: Holonomies and gauge-theoretical invariants of irreducible flat SU(3)-connections on $\Sigma(2, 3, 23)$.

will write β_k for this connection. The gauge-theoretical invariants of these connections are given in Table 5. In this table, $\tilde{\alpha}$ denotes the reducible SU(3)-connection associated to an SU(2)-connection α .

α	$\text{CS}(\alpha)$	ρ_{ad_α}	$\rho_{\text{ad}_{\tilde{\alpha}}}$	$\text{deg } \alpha$	$\text{deg } \tilde{\alpha}$
β_2	$\frac{1}{552}$	$-\frac{343}{69}$	$-\frac{206}{23}$	1	1
β_3	$\frac{169}{552}$	$-\frac{559}{69}$	$-\frac{406}{23}$	5	9
β_4	$\frac{73}{552}$	$-\frac{475}{69}$	$-\frac{410}{23}$	3	7
β_5	$\frac{265}{552}$	$-\frac{643}{69}$	$-\frac{402}{23}$	7	11

α	$\text{CS}(\alpha)$	ρ_{ad_α}	$\rho_{\text{ad}_{\tilde{\alpha}}}$	$\text{deg } \alpha$	$\text{deg } \tilde{\alpha}$
β_6	$\frac{193}{552}$	$-\frac{511}{69}$	$-\frac{382}{23}$	5	9
β_7	$\frac{409}{552}$	$-\frac{631}{69}$	$-\frac{534}{23}$	1	5
β_8	$\frac{361}{552}$	$-\frac{451}{69}$	$-\frac{490}{23}$	7	3
β_9	$\frac{49}{552}$	$-\frac{523}{69}$	$-\frac{434}{23}$	3	7

Table 5: Gauge-theoretical invariants of reducible flat $\text{SU}(3)$ -connections on $\Sigma(2, 3, 23)$.

References

- [1] **MF Atiyah, R Bott**, *The Yang–Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A 308 (1983) 523–615 [MR](#)
- [2] **MF Atiyah, V K Patodi, IM Singer**, *Spectral asymmetry and Riemannian geometry, II*, Math. Proc. Cambridge Philos. Soc. 78 (1975) 405–432 [MR](#)
- [3] **V Y Baranovskii**, *The cohomology ring of the moduli space of stable bundles with odd determinant*, Izv. Ross. Akad. Nauk Ser. Mat. 58 (1994) 204–210 [MR](#) In Russian; translated in *Russ. Acad. Sci. Izv. Math.* 45 (1995) 207–213
- [4] **H U Boden**, *Unitary representations of Brieskorn spheres*, Duke Math. J. 75 (1994) 193–220 [MR](#)
- [5] **H U Boden, S Friedl**, *Metabelian $\text{SL}(n, \mathbb{C})$ representations of knot groups*, Pacific J. Math. 238 (2008) 7–25 [MR](#)
- [6] **H U Boden, C M Herald, P A Kirk, E P Klassen**, *Gauge theoretic invariants of Dehn surgeries on knots*, Geom. Topol. 5 (2001) 143–226 [MR](#)
- [7] **A Borel**, *Topology of Lie groups and characteristic classes*, Bull. Amer. Math. Soc. 61 (1955) 397–432 [MR](#)
- [8] **P J Braam, S K Donaldson**, *Floer’s work on instanton homology, knots and surgery*, from “The Floer memorial volume” (H Hofer, C H Taubes, A Weinstein, E Zehnder, editors), Progr. Math. 133, Birkhäuser, Basel (1995) 195–256 [MR](#)
- [9] **P J Braam, S K Donaldson**, *Fukaya–Floer homology and gluing formulae for polynomial invariants*, from “The Floer memorial volume” (H Hofer, C H Taubes, A Weinstein, E Zehnder, editors), Progr. Math. 133, Birkhäuser, Basel (1995) 257–281 [MR](#)
- [10] **V Colin, P Ghiggini, K Honda**, *Embedded contact homology and open book decompositions*, preprint (2010) [arXiv](#)
- [11] **V Colin, P Ghiggini, K Honda, M Hutchings**, *Sutures and contact homology, I*, Geom. Topol. 15 (2011) 1749–1842 [MR](#)

- [12] **LH Culler**, *The blowup formula for higher rank Donaldson invariants*, PhD thesis, Massachusetts Institute of Technology (2014) [MR](#) Available at <https://search.proquest.com/docview/1566428730>
- [13] **L Culler, A Daemi, Y Xie**, *Surgery, polygons and $SU(N)$ -Floer Homology*, preprint (2017) [arXiv](#)
- [14] **U V Desale, S Ramanan**, *Poincaré polynomials of the variety of stable bundles*, Math. Ann. 216 (1975) 233–244 [MR](#)
- [15] **SK Donaldson**, *Anti self-dual Yang–Mills connections over complex algebraic surfaces and stable vector bundles*, Proc. London Math. Soc. 50 (1985) 1–26 [MR](#)
- [16] **SK Donaldson**, *The orientation of Yang–Mills moduli spaces and 4-manifold topology*, J. Differential Geom. 26 (1987) 397–428 [MR](#)
- [17] **SK Donaldson**, *Polynomial invariants for smooth four-manifolds*, Topology 29 (1990) 257–315 [MR](#)
- [18] **SK Donaldson**, *Floer homology and algebraic geometry*, from “Vector bundles in algebraic geometry” (NJ Hitchin, PE Newstead, WM Oxbury, editors), London Math. Soc. Lecture Note Ser. 208, Cambridge Univ. Press (1995) 119–138 [MR](#)
- [19] **SK Donaldson**, *Floer homology groups in Yang–Mills theory*, Cambridge Tracts in Mathematics 147, Cambridge Univ. Press (2002) [MR](#)
- [20] **SK Donaldson, PB Kronheimer**, *The geometry of four-manifolds*, Clarendon, New York (1990) [MR](#)
- [21] **S Dostoglou, DA Salamon**, *Self-dual instantons and holomorphic curves*, Ann. of Math. 139 (1994) 581–640 [MR](#)
- [22] **JD Edelstein, M Gómez-Reino, M Mariño**, *Blowup formulae in Donaldson–Witten theory and integrable hierarchies*, Adv. Theor. Math. Phys. 4 (2000) 503–543 [MR](#)
- [23] **PMN Feehan, TG Leness**, *Witten’s conjecture for many four-manifolds of simple type*, J. Eur. Math. Soc. 17 (2015) 899–923 [MR](#)
- [24] **R Fintushel, R J Stern**, *Instanton homology of Seifert fibred homology three spheres*, Proc. London Math. Soc. 61 (1990) 109–137 [MR](#)
- [25] **R Fintushel, R J Stern**, *Donaldson invariants of 4-manifolds with simple type*, J. Differential Geom. 42 (1995) 577–633 [MR](#)
- [26] **R Fintushel, R J Stern**, *The blowup formula for Donaldson invariants*, Ann. of Math. 143 (1996) 529–546 [MR](#)
- [27] **A Floer**, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. 118 (1988) 215–240 [MR](#)
- [28] **A Floer**, *Instanton homology and Dehn surgery*, from “The Floer memorial volume” (H Hofer, CH Taubes, A Weinstein, E Zehnder, editors), Progr. Math. 133, Birkhäuser, Basel (1995) 77–97 [MR](#)

- [29] **S Friedl**, *Eta invariants as sliceness obstructions and their relation to Casson–Gordon invariants*, *Algebr. Geom. Topol.* 4 (2004) 893–934 [MR](#)
- [30] **R Friedman**, *Rank two vector bundles over regular elliptic surfaces*, *Invent. Math.* 96 (1989) 283–332 [MR](#)
- [31] **R Friedman**, *Vector bundles and $SO(3)$ –invariants for elliptic surfaces*, *J. Amer. Math. Soc.* 8 (1995) 29–139 [MR](#)
- [32] **R Friedman, J W Morgan**, *Smooth four-manifolds and complex surfaces*, *Ergeb. Math. Grenzgeb.* 27, Springer (1994) [MR](#)
- [33] **K Fukaya**, *Floer homology for oriented 3–manifolds*, from “Aspects of low-dimensional manifolds” (Y Matsumoto, S Morita, editors), *Adv. Stud. Pure Math.* 20, Kinokuniya, Tokyo (1992) 1–92 [MR](#)
- [34] **D Gabai**, *Foliations and the topology of 3–manifolds*, *J. Differential Geom.* 18 (1983) 445–503 [MR](#)
- [35] **D Gabai**, *Foliations and the topology of 3–manifolds*, *Bull. Amer. Math. Soc.* 8 (1983) 77–80 [MR](#)
- [36] **D Gieseker**, *Geometric invariant theory and applications to moduli problems*, from “Invariant theory” (F Gherardelli, editor), *Lecture Notes in Math.* 996, Springer (1983) 45–73 [MR](#)
- [37] **D Gieseker**, *A construction of stable bundles on an algebraic surface*, *J. Differential Geom.* 27 (1988) 137–154 [MR](#)
- [38] **D Gieseker, J Li**, *Moduli of high rank vector bundles over surfaces*, *J. Amer. Math. Soc.* 9 (1996) 107–151 [MR](#)
- [39] **R E Gompf**, *Nuclei of elliptic surfaces*, *Topology* 30 (1991) 479–511 [MR](#)
- [40] **R E Gompf, A I Stipsicz**, *4–manifolds and Kirby calculus*, *Graduate Studies in Mathematics* 20, Amer. Math. Soc., Providence, RI (1999) [MR](#)
- [41] **V Gritsenko, K Hulek, G K Sankaran**, *Abelianisation of orthogonal groups and the fundamental group of modular varieties*, *J. Algebra* 322 (2009) 463–478 [MR](#)
- [42] **G Harder, M S Narasimhan**, *On the cohomology groups of moduli spaces of vector bundles on curves*, *Math. Ann.* 212 (1974/75) 215–248 [MR](#)
- [43] **R Hartshorne**, *Algebraic geometry*, *Graduate Texts in Mathematics* 52, Springer (1977) [MR](#)
- [44] **M Hutchings**, *Embedded contact homology and its applications*, from “Proceedings of the International Congress of Mathematicians, II” (R Bhatia, A Pal, G Rangarajan, V Srinivas, M Vanninathan, P Gastesi, editors), Hindustan Book Agency, New Delhi (2010) 1022–1041 [MR](#)

- [45] **M Hutchings**, *Lecture notes on embedded contact homology*, from “Contact and symplectic topology” (F Bourgeois, V Colin, A Stipsicz, editors), Bolyai Soc. Math. Stud. 26, János Bolyai Math. Soc., Budapest (2014) 389–484 [MR](#)
- [46] **A Juhász**, *Holomorphic discs and sutured manifolds*, *Algebr. Geom. Topol.* 6 (2006) 1429–1457 [MR](#)
- [47] **A D King, P E Newstead**, *On the cohomology ring of the moduli space of rank 2 vector bundles on a curve*, *Topology* 37 (1998) 407–418 [MR](#)
- [48] **R C Kirby, editor**, *Problems in low-dimensional topology*, from “Geometric topology, II” (W H Kazez, editor), AMS/IP Stud. Adv. Math. 2, Amer. Math. Soc., Providence, RI (1997) 35–473 [MR](#)
- [49] **K Kodaira**, *On compact complex analytic surfaces, I*, *Ann. of Math.* 71 (1960) 111–152 [MR](#)
- [50] **D Kotschick**, *SO(3)-invariants for 4-manifolds with $b_2^+ = 1$* , *Proc. London Math. Soc.* 63 (1991) 426–448 [MR](#)
- [51] **D Kotschick, J W Morgan**, *SO(3)-invariants for 4-manifolds with $b_2^+ = 1$, II*, *J. Differential Geom.* 39 (1994) 433–456 [MR](#)
- [52] **P B Kronheimer**, *Four-manifold invariants from higher-rank bundles*, *J. Differential Geom.* 70 (2005) 59–112 [MR](#)
- [53] **P B Kronheimer, T S Mrowka**, *Recurrence relations and asymptotics for four-manifold invariants*, *Bull. Amer. Math. Soc.* 30 (1994) 215–221 [MR](#)
- [54] **P B Kronheimer, T S Mrowka**, *Embedded surfaces and the structure of Donaldson’s polynomial invariants*, *J. Differential Geom.* 41 (1995) 573–734 [MR](#)
- [55] **P Kronheimer, T Mrowka**, *Monopoles and three-manifolds*, New Mathematical Monographs 10, Cambridge Univ. Press (2007) [MR](#)
- [56] **P Kronheimer, T Mrowka**, *Knots, sutures, and excision*, *J. Differential Geom.* 84 (2010) 301–364 [MR](#)
- [57] **P B Kronheimer, T S Mrowka**, *Khovanov homology is an unknot-detector*, *Publ. Math. Inst. Hautes Études Sci.* 113 (2011) 97–208 [MR](#)
- [58] **P B Kronheimer, T S Mrowka**, *Knot homology groups from instantons*, *J. Topol.* 4 (2011) 835–918 [MR](#)
- [59] **P B Kronheimer, T S Mrowka**, *Tait colorings, and an instanton homology for webs and foams*, *J. Eur. Math. Soc.* 21 (2019) 55–119 [MR](#)
- [60] **C Kutluhan, S Sivek, C H Taubes**, *Sutured ECH is a natural invariant*, preprint (2013) [arXiv](#)
- [61] **C LeBrun**, *Explicit self-dual metrics on $\mathbb{C}P_2 \# \cdots \# \mathbb{C}P_2$* , *J. Differential Geom.* 34 (1991) 223–253 [MR](#)

- [62] **W-P Li, Z Qin**, *Stable vector bundles on algebraic surfaces*, Trans. Amer. Math. Soc. 345 (1994) 833–852 [MR](#)
- [63] **P Lisca**, *Computations of instanton invariants using Donaldson–Floer theory*, Invent. Math. 119 (1995) 347–359 [MR](#)
- [64] **M Mariño, G Moore**, *The Donaldson–Witten function for gauge groups of rank larger than one*, Comm. Math. Phys. 199 (1998) 25–69 [MR](#)
- [65] **D McDuff, D Salamon**, *J–holomorphic curves and symplectic topology*, 2nd edition, American Mathematical Society Colloquium Publications 52, Amer. Math. Soc., Providence, RI (2012) [MR](#)
- [66] **V B Mehta, A Ramanathan**, *Restriction of stable sheaves and representations of the fundamental group*, Invent. Math. 77 (1984) 163–172 [MR](#)
- [67] **J W Morgan, H Bass** (editors), *The Smith conjecture*, Pure Appl. Math. 112, Academic, Orlando (1984) [MR](#)
- [68] **J W Morgan, T S Mrowka**, *A note on Donaldson’s polynomial invariants*, Int. Math. Res. Not. 1992 (1992) 223–230 [MR](#)
- [69] **J W Morgan, T S Mrowka**, *The smooth classification of elliptic surfaces*, from “Geometry, topology, & physics” (S-T Yau, editor), Conf. Proc. Lecture Notes Geom. Topology 4, International, Cambridge, MA (1995) 246–292 [MR](#)
- [70] **J W Morgan, T Mrowka, D Ruberman**, *The L^2 -moduli space and a vanishing theorem for Donaldson polynomial invariants*, Monographs in Geometry and Topology 2, International, Cambridge, MA (1994) [MR](#)
- [71] **V Muñoz**, *Gluing formulae for Donaldson invariants for connected sums along surfaces*, Asian J. Math. 1 (1997) 785–800 [MR](#)
- [72] **V Muñoz**, *Quantum cohomology of the moduli space of stable bundles over a Riemann surface*, Duke Math. J. 98 (1999) 525–540 [MR](#)
- [73] **V Muñoz**, *Ring structure of the Floer cohomology of $\Sigma \times \mathbb{S}^1$* , Topology 38 (1999) 517–528 [MR](#)
- [74] **P Ozsváth, Z Szabó**, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. 159 (2004) 1027–1158 [MR](#)
- [75] **Y Ruan, G Tian**, *A mathematical theory of quantum cohomology*, J. Differential Geom. 42 (1995) 259–367 [MR](#)
- [76] **D Ruberman**, *Relations among Donaldson invariants arising from negative 2–spheres and tori*, Duke Math. J. 83 (1996) 645–660 [MR](#)
- [77] **D A Salamon**, *Quantum products for mapping tori and the Atiyah–Floer conjecture*, from “Northern California Symplectic Geometry Seminar” (Y Eliashberg, D Fuchs, T Ratiu, A Weinstein, editors), Amer. Math. Soc. Transl. Ser. 2 196, Amer. Math. Soc., Providence, RI (1999) 199–235 [MR](#)

- [78] **B Siebert, G Tian**, *Recursive relations for the cohomology ring of moduli spaces of stable bundles*, Turkish J. Math. 19 (1995) 131–144 [MR](#)
- [79] **CH Taubes**, *Self-dual connections on 4–manifolds with indefinite intersection matrix*, J. Differential Geom. 19 (1984) 517–560 [MR](#)
- [80] **CH Taubes**, *Casson’s invariant and gauge theory*, J. Differential Geom. 31 (1990) 547–599 [MR](#)
- [81] **CH Taubes**, *L^2 moduli spaces on 4–manifolds with cylindrical ends*, Monographs in Geometry and Topology 1, International, Cambridge, MA (1993) [MR](#)
- [82] **WP Thurston**, *A norm for the homology of 3–manifolds*, Mem. Amer. Math. Soc. 339, Amer. Math. Soc., Providence, RI (1986) 99–130 [MR](#)
- [83] **W Wieczorek**, *The Donaldson invariant and embedded 2–spheres*, J. Reine Angew. Math. 489 (1997) 15–51 [MR](#)
- [84] **E Witten**, *Monopoles and four-manifolds*, Math. Res. Lett. 1 (1994) 769–796 [MR](#)
- [85] **Y Xie**, *On the framed singular instanton Floer homology from higher rank bundles*, PhD thesis, Harvard University (2016) [MR](#) Available at <https://tinyurl.com/Xie-thesis>
- [86] **D Zagier**, *On the cohomology of moduli spaces of rank two vector bundles over curves*, from “The moduli space of curves” (R Dijkgraaf, C Faber, G van der Geer, editors), Progr. Math. 129, Birkhäuser, Boston (1995) 533–563 [MR](#)

*Simons Center for Geometry and Physics, State University of New York
Stony Brook, NY, United States*

*Beijing International Center for Mathematical Research, Peking University
Beijing, China*

adaemi@scgp.stonybrook.edu, yixie@pku.edu.cn

Proposed: Simon Donaldson

Received: 4 March 2018

Seconded: Ciprian Manolescu, Peter Ozsváth

Revised: 25 March 2019