

# Recognition of being fibered for compact 3–manifolds

ANDREI JAIKIN-ZAPIRAIN

Let  $M$  be a compact orientable aspherical 3–manifold. We show that if the profinite completion of  $\pi_1(M)$  is isomorphic to the profinite completion of a free-by-cyclic group or to the profinite completion of a surface-by-cyclic group, then  $M$  fibers over the circle with compact fiber.

57M27; 20E18, 20J05, 57M05

## 1 Introduction

Let  $\Gamma$  be a finitely generated residually finite group. It is natural to ask what properties of  $\Gamma$  are determined by the set of its finite quotients. Recall that two finitely generated groups have the same collection of finite quotients if and only if their profinite completions are isomorphic (see the discussion around Theorem 2.2 of A Reid [15]). Thus, our previous question can be reformulated as follows:

**Question 1** *Let  $\Gamma_1$  and  $\Gamma_2$  be two finitely generated groups. Assume that they have isomorphic profinite completions. What group-theoretic properties are shared by  $\Gamma_1$  and  $\Gamma_2$ ?*

There are many recent developments around this question. We recommend the reader to consult the survey of Reid [15], where all these new results are described.

In this paper we consider a particular case of Question 1, where the groups  $\Gamma_1$  and  $\Gamma_2$  are the fundamental groups of 3–manifolds. This case has been also studied actively in recent years (for more details, see [15, Section 4]).

Throughout this paper, all manifolds are connected and aspherical. We allow 3–manifolds to have nonempty boundary, but we assume that there are no spherical components.

By a free-by-cyclic group we mean a group  $F \rtimes \mathbb{Z}$ , where  $F$  is a finitely generated free group, and a surface-by-cyclic group is a group  $S \rtimes \mathbb{Z}$ , where  $S$  is the fundamental group of a compact closed orientable surface. Our result shows that the profinite

completion of the fundamental group of a 3–manifold determines whether the manifold fibers over the circle with compact fiber.

**Theorem 1.1** *Let  $M$  be a compact orientable 3–manifold. Assume that the profinite completion of  $\pi_1(M)$  is isomorphic to the profinite completion of a free-by-cyclic group or to the profinite completion of a surface-by-cyclic group. Then  $M$  fibers over the circle with compact fiber.*

Under some additional conditions, this result has been proved by Bridson and Reid [5], Boileau and Friedl [4] and Bridson, Reid and Wilton [6]. Our proof and all the previous proofs use in an essential way results of I Agol [1; 2] and P Przytycki and D Wise [14; 18] on separability of 3–manifold groups.

**Corollary 1.2** *Let  $M$  and  $N$  be two compact orientable 3–manifolds such that  $\widehat{\pi_1(M)} \cong \widehat{\pi_1(N)}$ . Then  $M$  fibers over the circle if and only if  $N$  does.*

Let us explain briefly the strategy of the proof of Theorem 1.1. If  $\Gamma$  is a finitely generated group, then we say that a nontrivial class  $\phi \in H^1(\Gamma, \mathbb{Z}) = \text{Hom}(\Gamma, \mathbb{Z})$  is *fibred* if  $\ker \phi$  is finitely generated. If  $\Gamma = \pi_1(M)$  is the fundamental group of a compact orientable 3–manifold  $M$ , then, by a well-known result of J Stallings [17], fibred classes in  $H^1(\pi_1(M), \mathbb{Z})$  are in one-to-one correspondence with fibrations of  $M$  over the circle with compact fiber.

Now, let us assume that  $M$  satisfies the conditions of Theorem 1.1. First we have to find a useful criterion for a class  $\phi \in H^1(\pi_1(M), \mathbb{Z})$  to be fibred. We start with a criterion proved by S Friedl and S Vidussi (Proposition 3.3). In Section 3.1 we explain the notion of cohomological goodness introduced by J-P Serre [16]. Using that the fundamental group of a 3–manifold is cohomologically good we reformulate Proposition 3.3 in terms of the profinite completion of  $\pi_1(M)$  and obtain Corollary 3.4. It tells us that  $\phi \in H^1(\pi_1(M), \mathbb{Z})$  is fibred if certain homology groups of some subgroups of  $\widehat{\pi_1(M)}$  are finite. The latter property can be understood in terms of certain rank functions, which we study in Section 2. In Section 4 we prove a proposition that together with Corollary 2.2 allows us to find a desired class. All this is explained again in more detail in Section 5.

## Acknowledgements

This paper is partially supported by the grants MTM2017-82690-P and MTM2014-53810-C2-01 of the Spanish MINECO, the grant PRX16/00179 of the mobility program

“Salvador de Madariaga” of the Spanish MECD and by the ICMAT Severo Ochoa project SEV-2015-0554.

This paper was written while the author was visiting the Mathematical Institute of the University of Oxford. I would like to thank everyone involved for their fine hospitality.

I would like to thank Pavel Zalesskii for bringing this problem to my attention and to Martin Bridson, Alan Reid and Henry Wilton for very useful discussions. I would also like to thank Diego López-Álvarez and an anonymous referee whose remarks and suggestions substantially improved the article.

## 2 Rank functions

Let  $R$  be a commutative domain and let  $Q(R)$  denote its field of fractions. Given a matrix  $A$  over  $R$  we denote by  $\text{rk}_R(A)$  the  $Q(R)$ -rank of  $A$ . Similarly, if  $\phi: M_1 \rightarrow M_2$  is a homomorphism of finitely generated free left  $R$ -modules, we denote by  $\text{rk}_R(\phi)$  the  $Q(R)$ -rank of the induced map  $Q(R) \otimes_R M_1 \rightarrow Q(R) \otimes_R M_2$ .

We may extend the previous notation in the following way. Let  $S$  be a ring and assume that  $S_0$  is a subring of  $S$  such that  $S \cong (S_0)^k$  for some  $k \geq 1$  as a left  $S_0$ -module. If  $\alpha: S_0 \rightarrow R$  is a homomorphism of rings, then  $\alpha$  induces a structure of right  $S_0$ -module on  $R$ . Take  $A \in \text{Mat}_{n \times m}(S)$  and consider the homomorphism of finitely generated free left  $R$ -modules

$$\phi_\alpha^A: (R \otimes_{S_0} S)^n \rightarrow (R \otimes_{S_0} S)^m, \quad (x_1, \dots, x_n) \mapsto (x_1, \dots, x_n)A.$$

If  $R$  is a commutative domain, then we put  $\text{rk}_\alpha(A) = \text{rk}_R(\phi_\alpha^A)$ .

Let us describe two concrete examples of rank functions that will appear in this paper. Let  $\Gamma$  be a residually finite group and  $H$  a subgroup of finite index. Put  $S = \mathbb{Z}[\Gamma]$  and  $S_0 = \mathbb{Z}[H]$ . It is clear that  $S$  is a left free  $S_0$ -module. We fix a prime  $p$  once and for all throughout the paper.

(A) Let  $K$  be a normal subgroup of  $\Gamma$  such that  $\Gamma/K$  is abelian and torsion-free. Let  $\alpha_1: \mathbb{Z}[H] \rightarrow \mathbb{F}_p[H/H \cap K]$  be the canonical map. We define

$$\text{rk}_{H,K}(A) = \frac{\text{rk}_{\alpha_1}(A)}{|\Gamma : H|}, \quad \text{where } A \text{ is a matrix over } \mathbb{Z}[\Gamma].$$

(B) For a profinite group  $G$  and a profinite ring  $T$  we denote by  $T[[G]]$  the completed group algebra

$$T[[G]] = \varprojlim_{U \trianglelefteq_o G} T[[G/U]].$$

Now let  $G$  be a profinite completion of  $\Gamma$ . We identify  $\Gamma$  with its canonical image in  $G$ . Let  $U$  be the closure of  $H$  in  $G$  and let  $N$  be a closed normal subgroup of  $G$  such that  $G/N$  is an abelian and torsion-free pro- $p$  group. Let  $\alpha_2: \mathbb{Z}[H] \rightarrow \mathbb{F}_p[[U/U \cap N]]$  be the canonical map. In this context we define

$$\text{rk}_{H,N}(A) = \frac{\text{rk}_{\alpha_2}(A)}{|G : U|}, \quad \text{where } A \text{ is a matrix over } \mathbb{Z}[\Gamma].$$

This rank function coincides with the previous one in the case  $K = \Gamma \cap N$ . Indeed, for every matrix  $A$  over  $\mathbb{Z}[\Gamma]$  we have the equality

$$\text{rk}_{\alpha_1}(A) = \text{rk}_{\alpha_2}(A)$$

because  $\alpha_2$  is the composition of  $\alpha_1$  and the canonical embedding of  $\mathbb{F}_p[H/H \cap K]$  into  $\mathbb{F}_p[[U/U \cap N]]$ . Thus, we obtain that

$$(1) \quad \text{rk}_{H,\Gamma \cap N}(A) = \text{rk}_{H,N}(A).$$

The rank functions  $\text{rk}_{H,K}$  and  $\text{rk}_{H,N}$  are examples of Sylvester matrix rank functions on  $\mathbb{Z}[\Gamma]$ . The interested reader can find more information about them in [11]. We need the following elementary lemma about rank functions:

**Lemma 2.1** *Let  $S$  be a ring and assume that  $S_0$  is a subring of  $S$  such that  $S$  is a finitely generated free left  $S_0$ -module. Let  $\pi: R_1 \rightarrow R_2$  be a homomorphism between two commutative domains and  $\alpha_1: S_0 \rightarrow R_1$  a homomorphism. We put  $\alpha_2 = \pi \circ \alpha_1$ . Then for any matrix  $A$  over  $S$  we have*

$$\text{rk}_{\alpha_1}(A) \geq \text{rk}_{\alpha_2}(A).$$

**Proof** For simplicity of exposition, let us assume that  $A \in S$  (that is,  $A$  is a 1-by-1 matrix).

Let  $\{v_1, \dots, v_k\}$  be a basis of  $S$  as a left  $S_0$ -module. Consider the matrix associated to  $\phi_{\text{id}}^A$  with respect to  $\{v_1, \dots, v_k\}$ . It is a  $k$ -by- $k$  matrix  $B = (b_{ij})$  over  $S_0$  defined by means of

$$v_i A = \sum_{j=1}^k b_{ij} v_j.$$

Let  $B_1 = \alpha_1(B)$  and  $B_2 = \alpha_2(B)$ , and observe that  $B_2 = \pi(B_1)$ . Thus,  $\text{rk}_{R_1}(B_1) \geq \text{rk}_{R_2}(B_2)$ . Therefore, we obtain that

$$\text{rk}_{\alpha_1}(A) = \text{rk}_{R_1}(B_1) \geq \text{rk}_{R_2}(B_2) = \text{rk}_{\alpha_2}(A). \quad \square$$

We will apply the lemma in the following situation:

**Corollary 2.2** *Let  $\Gamma$  be a residually finite group and let  $G = \widehat{\Gamma}$ . Let  $N$  be a closed normal subgroup of  $G$  such that  $G/N \cong \mathbb{Z}_p$  and let  $K$  be a normal subgroup of  $\Gamma$  such that  $\Gamma/K \cong \mathbb{Z}$ . Assume that  $\Gamma \cap N \leq K$ . Then, for every matrix  $A$  over  $\mathbb{Z}[\Gamma]$  and every subgroup  $H$  of  $\Gamma$  of finite index,*

$$\text{rk}_{H,N}(A) \geq \text{rk}_{H,K}(A).$$

**Proof** Using (1), we obtain that  $\text{rk}_{H,\Gamma \cap N}(A) = \text{rk}_{H,N}(A)$ . Since  $\Gamma \cap N \leq K$ , the previous lemma implies that  $\text{rk}_{H,\Gamma \cap N}(A) \geq \text{rk}_{H,K}(A)$ . Hence we are done.  $\square$

### 3 Preliminaries on 3–manifolds

#### 3.1 Cohomologically good groups

Let  $\Gamma$  be a group and  $\widehat{\Gamma}$  its profinite completion. The group  $\Gamma$  is called *cohomologically good* if the homomorphism of cohomology groups

$$i^n(M): H^n(\widehat{\Gamma}, M) \rightarrow H^n(\Gamma, M)$$

induced by the natural homomorphism  $i: \Gamma \rightarrow \widehat{\Gamma}$  is an isomorphism for every  $\mathbb{Z}[\Gamma]$ –module  $M$  having a finite number of elements. This notion was introduced by Serre (see [16, Section I.2.6]) and has been studied in several papers (see for example [10; 9; 13]).

It is known that free, surface, free-by-cyclic, surface-by-cyclic and 3–manifold groups are cohomologically good. The result on 3–manifold groups (see for example [7; 9, Remark 5.14; 3]) uses the recent advances in the theory of 3–manifolds, including the solution of the virtual Haken conjecture by Agol [2]. We want to point out that under the hypothesis of Theorem 1.1 that the profinite completion of  $\pi_1(M)$  is isomorphic to the profinite completion of a free-by-cyclic group or to the profinite completion of a surface-by-cyclic group, the cohomological goodness of  $\pi_1(M)$  can be proved directly as in [9, Corollary 5.4].

In the case where  $\Gamma$  is an  $\text{FP}_\infty$ –group there is an alternative way to define the cohomological goodness.

**Proposition 3.1** *Let  $\Gamma$  be an  $\text{FP}_\infty$ –group and let*

$$\dots \rightarrow \mathbb{Z}[\Gamma]^{d_i} \xrightarrow{\phi_i} \mathbb{Z}[\Gamma]^{d_{i-1}} \rightarrow \dots \rightarrow \mathbb{Z}[\Gamma]^{d_1} \xrightarrow{\phi_1} \mathbb{Z}[\Gamma] \rightarrow \mathbb{Z} \rightarrow 0$$

be a resolution of the trivial  $\mathbb{Z}[\Gamma]$ -module  $\mathbb{Z}$ . Then  $\Gamma$  is cohomologically good if and only if the induced sequence

$$\dots \rightarrow \widehat{\mathbb{Z}}[\widehat{\Gamma}]^{d_i} \xrightarrow{\widehat{\phi}_i} \widehat{\mathbb{Z}}[\widehat{\Gamma}]^{d_{i-1}} \rightarrow \dots \rightarrow \widehat{\mathbb{Z}}[\widehat{\Gamma}]^{d_1} \xrightarrow{\widehat{\phi}_1} \widehat{\mathbb{Z}}[\widehat{\Gamma}] \rightarrow \widehat{\mathbb{Z}} \rightarrow 0$$

is exact.

**Proof** The “if” part is clear. Let us show the “only if” part.

By [16, page 15], since  $\Gamma$  is cohomologically good we have that for every  $n \geq 1$ , every prime  $p$ , every subgroup  $H$  of  $\Gamma$  of finite index and every  $\alpha \in H^k(H, \mathbb{F}_p)$ , there exists another subgroup  $T$  of finite index in  $H$  such that the restriction of  $\alpha$  to  $T$  vanishes in  $H^k(T, \mathbb{F}_p)$ . Hence the direct limit of  $H^k(H, \mathbb{F}_p)$  (when  $H$  runs over all subgroups of finite index of  $\Gamma$ ) is equal to zero, and so the inverse limit of  $H_k(H, \mathbb{F}_p)$  is also equal to zero. Now we can apply [12, Theorem 2.5] and conclude that  $H_k(\Gamma, \mathbb{Z}_p[\widehat{\Gamma}]) = 0$  for each  $p$ , and therefore  $H_k(\Gamma, \widehat{\mathbb{Z}}[\widehat{\Gamma}]) = 0$ . This implies the exactness of the second sequence in the statement of the proposition.  $\square$

### 3.2 Profinite completions of 3-manifold groups

In the following proposition we show that the hypotheses of Theorem 1.1 impose several restrictions on the boundary of  $M$ .

**Proposition 3.2** *Let  $M$  be a compact orientable 3-manifold and let  $\Gamma$  be a free-by-cyclic group or a surface-by-cyclic group. Assume that  $\widehat{\pi_1(M)} \cong \widehat{\Gamma}$ . Then the boundary of  $M$  is a union of incompressible tori or it is empty.*

**Proof** If  $\Gamma$  is a free-by-cyclic group, the proposition follows from [5, Corollary 4.3]. If  $\Gamma$  is a surface-by-cyclic group, then  $\widehat{\pi_1(M)}$  is of cohomological dimension 3. Since  $\pi_1(M)$  is cohomologically good,  $\pi_1(M)$  is of cohomological dimension 3 as well, and so  $M$  does not have boundary.  $\square$

### 3.3 A criterion for fibering

Let  $\phi: \Gamma \rightarrow \mathbb{Z}$  be a nontrivial homomorphism and let  $H$  be a subgroup of finite index in  $\Gamma$ . We denote by  $H_\phi$  the intersection  $H \cap \ker \phi$ . We have the following criterion of fibering, stated in a slightly different way in a paper of Friedl and Vidussi [8]:

**Proposition 3.3** *Let  $M$  be a compact orientable 3-manifold with toroidal or empty boundary and let  $\phi: \pi_1(M) \rightarrow \mathbb{Z}$  be a nontrivial map. Then the class  $\phi$  is fibered if and only if for every normal subgroup  $H$  of  $\pi_1(M)$  of finite index the group  $H_1(H_\phi, \mathbb{F}_p)$  is finite.*

**Proof** The “only if” part is clear, since if  $\phi$  fibers,  $\ker \phi$  is finitely generated.

Let  $H$  be a normal subgroup of  $\pi_1(M)$  of finite index and assume that  $H_1(H_\phi, \mathbb{F}_p)$  is finite. Hence  $H_1(\pi_1(M), \mathbb{F}_p[\pi_1(M)/H_\phi]) \cong H_1(H_\phi, \mathbb{F}_p)$  is also finite. Put  $L = \mathbb{F}_p[\pi_1(M)/H]$ . The tensor product

$$\tilde{L} = \mathbb{F}_p[\pi_1(M)/\ker \phi] \otimes_{\mathbb{F}_p} L$$

is a finitely generated right  $\pi_1(M)$ -module (the elements of  $\pi_1(M)$  act diagonally on  $L_\phi$ ). The  $i^{\text{th}}$  twisted Alexander polynomial  $\Delta_{M,\phi,i}^L$  is the order of the  $\mathbb{F}_p[\pi_1(M)/\ker \phi]$ -module  $H_i(\pi_1(M), \tilde{L})$  (see [4] for details). Observe that  $\tilde{L}$  is a direct sum of a finite number of copies of  $\mathbb{F}_p[\pi_1(M)/H_\phi]$ . Therefore,  $H_1(\pi_1(M), \tilde{L})$  is also finite. Hence

$$\Delta_{M,\phi,1}^{\mathbb{F}_p[\pi_1(M)/H]} \neq 0.$$

Thus, we can apply [8, Theorem 1.1] and obtain the “if” part of the proposition.  $\square$

Let  $\Gamma$  be a finitely generated group. A map  $\phi: \Gamma \rightarrow \mathbb{Z}$  induces a homomorphism  $\phi_k: \hat{\Gamma} \rightarrow \mathbb{Z}_p/p^k\mathbb{Z}_p$ . We put

$$\hat{\phi} = \lim_k \phi_k: \hat{\Gamma} \rightarrow \mathbb{Z}_p$$

and say that  $\hat{\phi}$  is associated to  $\phi$ .

Let  $G$  be a profinite group,  $U$  an open subgroup and  $\varphi: G \rightarrow \mathbb{Z}_p$  a nontrivial homomorphism. We denote by  $U_\varphi$  the intersection  $U \cap \ker \varphi$ . Note that  $U/U_\varphi \cong \mathbb{Z}_p$ . Now we present a profinite analogue of Proposition 3.3.

**Corollary 3.4** *Let  $M$  be a compact orientable 3-manifold with toroidal or empty boundary and let  $\phi: \pi_1(M) \rightarrow \mathbb{Z}$  be a nontrivial map. Denote by  $\hat{\phi}: \widehat{\pi_1(M)} \rightarrow \mathbb{Z}_p$  the homomorphism associated to  $\phi$ . Then the class  $\phi$  is fibered if and only if for every normal open subgroup  $U$  of  $\widehat{\pi_1(M)}$  the group  $H_1(U_{\hat{\phi}}, \mathbb{F}_p)$  is finite.*

**Proof** The “only if” part is clear. Let us prove the “if” part.

We write a projective resolution of  $\mathbb{Z}$  as a left  $\mathbb{Z}[\pi_1(M)]$ -module

$$\mathbb{Z}[\pi_1(M)]^r \xrightarrow{m_B} \mathbb{Z}[\pi_1(M)]^d \rightarrow \mathbb{Z}[\pi_1(M)] \rightarrow \mathbb{Z} \rightarrow 0,$$

where  $m_B$  is multiplication by a matrix  $B$ . Since,  $\pi_1(M)$  is cohomologically good, using Proposition 3.1, we obtain an exact sequence

$$\widehat{\mathbb{Z}}[\widehat{\pi_1(M)}]^r \xrightarrow{m_B} \widehat{\mathbb{Z}}[\widehat{\pi_1(M)}]^d \rightarrow \widehat{\mathbb{Z}}[\widehat{\pi_1(M)}] \rightarrow \widehat{\mathbb{Z}} \rightarrow 0.$$

Let  $H$  be a normal subgroup of  $\pi_1(M)$  of finite index and let  $U$  be the closure of  $H$  in  $\widehat{\pi_1(M)}$ . Then  $U$  is a normal open subgroup of  $\widehat{\pi_1(M)}$ . Note that

$$H_1(U_{\widehat{\phi}}, \mathbb{F}_p) \cong H_1(\widehat{\pi_1(M)}, \mathbb{F}_p[\widehat{\pi_1(M)}/U_{\widehat{\phi}}]).$$

Thus, since  $H_1(U_{\widehat{\phi}}, \mathbb{F}_p)$  is finite, we obtain that  $\text{rk}_{H, \ker \widehat{\phi}}(B) = d - 1$ . Now, observe that, by (1),

$$\text{rk}_{H, \ker \phi}(B) = \text{rk}_{H, \ker \widehat{\phi}}(B) = d - 1.$$

Therefore,

$$H_1(H_{\phi}, \mathbb{F}_p) \cong H_1(\pi_1(M), \mathbb{F}_p[\pi_1(M)/H_{\phi}])$$

is also finite. Hence we can apply Proposition 3.3 and obtain the ‘‘if’’ part of the corollary. □

### 4 Finding a fibered class

The following proposition will be a key result that will help us to find a fibered class in Theorem 1.1.

**Proposition 4.1** *Let  $V = \mathbb{Z}_p^k$  and let  $I \cong J \cong \mathbb{Z}^k$  be two dense subgroups of  $V$ . Put  $V^* = \text{Hom}(V, \mathbb{Z}_p)$ . For  $\varphi \in V^*$  we denote by  $\varphi|_I$  the restriction of  $\varphi$  to  $I$ . Then, for every  $\psi \in V^*$  such that  $\psi(I) \leq \mathbb{Z}$ , there exists an element  $\varphi \in V^*$  such that  $\varphi(J) \leq \mathbb{Z}$  and  $\ker \varphi|_I \leq \ker \psi|_I$ .*

**Proof** Let  $\{v_1, \dots, v_k\}$  and  $\{u_1, \dots, u_k\}$  be  $\mathbb{Z}$ -bases of  $I$  and  $J$ , respectively. Define elements  $v_1^*, \dots, v_k^*, u_1^*, \dots, u_k^*$  of  $V^*$  by means of

$$v_i^*(v_j) = u_i^*(u_j) = \delta_{ij}.$$

Since  $\{v_1, \dots, v_k\}$  and  $\{u_1, \dots, u_k\}$  are  $\mathbb{Z}_p$ -bases of  $V$ ,  $\{v_1^*, \dots, v_k^*\}$  and  $\{u_1^*, \dots, u_k^*\}$  are  $\mathbb{Z}_p$ -bases of  $V^*$ . Hence there exists a matrix  $A \in \text{GL}_k(\mathbb{Z}_p)$  such that

$$(2) \quad (u_1^*, \dots, u_k^*) = (v_1^*, \dots, v_k^*)A.$$

Write  $A$  as

$$(3) \quad A = \sum_{i=1}^s A_i z_i,$$

where  $A_i \in \text{Mat}_k(\mathbb{Z})$  and  $\{z_1, \dots, z_s\} \subset \mathbb{Z}_p$  are  $\mathbb{Z}$ -linearly independent.

Let  $R = \mathbb{Q}[t_1, \dots, t_s]$  be a polynomial ring. We put  $B = \sum_{i=1}^s A_i t_i \in \text{Mat}_k(R)$ . Since  $A = \sum_{i=1}^s A_i z_i$  is invertible over  $\mathbb{Z}_p$ ,  $\det A \neq 0$ . Hence  $\det B \neq 0$  as well and, as a consequence, there are  $q_1, \dots, q_s \in \mathbb{Z}$  such that

$$\det\left(\sum_{i=1}^s A_i q_i\right) \neq 0.$$

Let us put

$$(4) \quad C = \sum_{i=1}^s A_i q_i.$$

We can express  $\psi$  as

$$(5) \quad \psi = \alpha_1 v_1^* + \dots + \alpha_k v_k^*, \quad \text{where } \alpha_i \in \mathbb{Z}.$$

Since  $C$  is invertible over  $\mathbb{Q}$ , there are  $\beta_1, \dots, \beta_k \in \mathbb{Z}$  and  $0 \neq n \in \mathbb{Z}$  such that

$$(6) \quad C \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix} = n \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_k \end{pmatrix}.$$

Now we set

$$\begin{aligned} \varphi &= \sum_{i=1}^k \beta_i u_i^* = (v_1^*, \dots, v_k^*) A \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix} && \text{(by (2))} \\ &= \sum_{j=1}^s z_j (v_1^*, \dots, v_k^*) A_j \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix} && \text{(by (3)),} \end{aligned}$$

and define

$$(7) \quad \varphi_j = (v_1^*, \dots, v_k^*) A_j \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix} \in \mathbb{Z} v_1^* + \dots + \mathbb{Z} v_k^*.$$

Then, since  $\{z_1, \dots, z_s\}$  are  $\mathbb{Z}$ -linearly independent,

$$\ker \varphi|_I = \ker(z_1(\varphi_1)|_I + \dots + z_s(\varphi_s)|_I) = \bigcap_{i=1}^s \ker(\varphi_j)|_I.$$

Thus, we obtain

$$\begin{aligned} \ker \varphi|_I &\leq \ker \left( \sum_{j=1}^s q_j(\varphi_j)|_I \right) = \ker \left( (v_1^*, \dots, v_k^*) C \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix} |_I \right) && \text{(by (7) and (4))} \\ &= \ker \left( (v_1^*, \dots, v_k^*) \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_k \end{pmatrix} |_I \right) && \text{(by (6))} \\ &= \ker \psi|_I && \text{(by (5)).} \quad \square \end{aligned}$$

### 5 Proof of Theorem 1.1

In this section we prove Theorem 1.1. Let  $\Gamma = S \rtimes \mathbb{Z}$  be a free-by-cyclic or a surface-by-cyclic group. Then  $\widehat{\Gamma} \cong \widehat{S} \rtimes \widehat{\mathbb{Z}}$ . We write a projective resolution of  $\mathbb{Z}$  as a left  $\mathbb{Z}[\Gamma]$ -module,

$$\mathbb{Z}[\Gamma]^r \xrightarrow{m_A} \mathbb{Z}[\Gamma]^d \rightarrow \mathbb{Z}[\Gamma] \rightarrow \mathbb{Z} \rightarrow 0,$$

where  $m_A$  is multiplication by a matrix  $A$ . We will need the following lemma:

**Lemma 5.1** *Let  $\varphi: \widehat{\Gamma} \rightarrow \mathbb{Z}_p$  be a nontrivial homomorphism and let  $H$  be a subgroup of  $\Gamma$  of finite index. Denote by  $U$  the closure of  $H$  in  $\widehat{\Gamma}$ . Then  $H_1(U_\varphi, \mathbb{F}_p)$  is finite if and only if  $\text{rk}_{H, \ker \varphi}(A) = d - 1$ .*

**Proof** We may use the same argument as in the proof of Corollary 3.4, because  $\Gamma$  is cohomologically good. □

Let  $M$  be a compact orientable 3-manifold and assume that  $\widehat{\pi_1(M)}$  and  $\widehat{\Gamma}$  are isomorphic. By Proposition 3.2,  $M$  has toroidal or empty boundary.

Denote by  $\alpha: \widehat{\pi_1(M)} \rightarrow \widehat{\Gamma}$  an isomorphism between  $\widehat{\pi_1(M)}$  and  $\widehat{\Gamma}$ . In order to show that  $\pi_1(M)$  is fibered we want to use Corollary 3.4. To do so we are going to work in  $\widehat{\Gamma}$  and then transport the obtained information from  $\widehat{\Gamma}$  to  $\widehat{\pi_1(M)}$  using  $\alpha^{-1}$ .

Let  $f: \Gamma \rightarrow \mathbb{Z}$  be such that  $\ker f = S$  and let  $\widehat{f}: \widehat{\Gamma} \rightarrow \mathbb{Z}_p$  be the map associated to  $f$ . Denote by  $V$  the maximal torsion-free abelian pro- $p$  quotient of  $\widehat{\Gamma}$ . Then  $V \cong \mathbb{Z}_p^k$  for some  $k \geq 1$ . Let  $I$  and  $J$  be the images in  $V$  of  $\Gamma$  and  $\alpha(\pi_1(M))$ , respectively.

Applying Proposition 4.1 for  $\psi = \widehat{f}$ , we obtain that there exists  $\varphi: \widehat{\Gamma} \rightarrow \mathbb{Z}_p$  such that

$$\varphi(\alpha(\pi_1(M))) \leq \mathbb{Z} \quad \text{and} \quad \ker \varphi \cap \Gamma \leq \ker f.$$

Let  $U$  be an open normal subgroup of  $\widehat{\Gamma}$ . Put  $H = \Gamma \cap U$ . Observe that  $U$  is equal to the closure of  $H$  in  $\widehat{\Gamma}$ . By Corollary 2.2,

$$\mathrm{rk}_{H, \ker \varphi}(A) \geq \mathrm{rk}_{H, \ker f}(A) = d - 1.$$

Hence, by Lemma 5.1,  $H_1(U_\varphi, \mathbb{F}_p)$  is finite.

Now, we put

$$\phi = \varphi \circ \alpha|_{\pi_1(M)} \in H^1(\pi_1(M), \mathbb{Z}).$$

Let  $\widehat{\phi}: \widehat{\pi_1(M)} \rightarrow \mathbb{Z}_p$  be associated to  $\phi$ . Then we have that  $\widehat{\phi} = \varphi \circ \alpha$ .

Thus, we obtain that

$$H_1(U_{\widehat{\phi}}, \mathbb{F}_p) \cong H_1(\alpha(U_{\widehat{\phi}}), \mathbb{F}_p) = H_1(\alpha(U)_\varphi, \mathbb{F}_p)$$

is finite for every normal open subgroup  $U$  of  $\widehat{\pi_1(M)}$ . Therefore, applying Corollary 3.4, we obtain that  $\phi$  is fibered. This finishes the proof of Theorem 1.1.

## References

- [1] **I Agol**, *Criteria for virtual fibering*, J. Topol. 1 (2008) 269–284 MR
- [2] **I Agol**, *The virtual Haken conjecture*, Doc. Math. 18 (2013) 1045–1087 MR
- [3] **M Aschenbrenner, S Friedl, H Wilton**, *3-Manifold groups*, Eur. Math. Soc., Zürich (2015) MR
- [4] **M Boileau, S Friedl**, *The profinite completion of 3-manifold groups, fiberedness and the Thurston norm*, preprint (2015) arXiv
- [5] **M Bridson, A Reid**, *Profinite rigidity, fibering, and the figure-eight knot*, preprint (2015) arXiv
- [6] **M R Bridson, A W Reid, H Wilton**, *Profinite rigidity and surface bundles over the circle*, Bull. Lond. Math. Soc. 49 (2017) 831–841 MR
- [7] **W Cavendish**, *Finite-sheeted covering spaces and solenoids over 3-manifolds*, PhD thesis, Princeton University (2012) MR <https://search.proquest.com/docview/1040698746>
- [8] **S Friedl, S Vidussi**, *A vanishing theorem for twisted Alexander polynomials with applications to symplectic 4-manifolds*, J. Eur. Math. Soc. 15 (2013) 2027–2041 MR
- [9] **F Grunewald, A Jaikin-Zapirain, A G S Pinto, P A Zalesskii**, *Normal subgroups of profinite groups of non-negative deficiency*, J. Pure Appl. Algebra 218 (2014) 804–828 MR

- [10] **F Grunewald, A Jaikin-Zapirain, P A Zalesskii**, *Cohomological goodness and the profinite completion of Bianchi groups*, *Duke Math. J.* 144 (2008) 53–72 MR
- [11] **A Jaikin-Zapirain**,  *$L^2$ -Betti numbers and their analogues in positive characteristic*, from “Groups St Andrews 2017 in Birmingham” (C M Campbell, C W Parker, M R Quick, E F Robertson, C M Roney-Dougal, editors), *London Math. Soc. Lecture Note Ser.* 455, Cambridge Univ. Press (2019) 346–405 MR
- [12] **D H Kochloukova, P A Zalesskii**, *Profinite and pro- $p$  completions of Poincaré duality groups of dimension 3*, *Trans. Amer. Math. Soc.* 360 (2008) 1927–1949 MR
- [13] **K Lorensen**, *Groups with the same cohomology as their profinite completions*, *J. Algebra* 320 (2008) 1704–1722 MR
- [14] **P Przytycki, D T Wise**, *Separability of embedded surfaces in 3-manifolds*, *Compos. Math.* 150 (2014) 1623–1630 MR
- [15] **A W Reid**, *Profinite rigidity*, from “Proceedings of the International Congress of Mathematicians” (B Sirakov, P N de Souza, M Viana, editors), volume 2, World Sci., Hackensack, NJ (2018) 1211–1234 MR
- [16] **J-P Serre**, *Galois cohomology*, Springer (2002) MR
- [17] **J Stallings**, *On fibering certain 3-manifolds*, from “Topology of 3-manifolds and related topics” (J M K Fort, editor), Prentice-Hall, Englewood Cliffs, NJ (1962) 95–100 MR
- [18] **D T Wise**, *Research announcement: the structure of groups with a quasiconvex hierarchy*, *Electron. Res. Announc. Math. Sci.* 16 (2009) 44–55 MR

*Departamento de Matemáticas, University Autónoma de Madrid and Instituto de Ciencias Matemáticas  
Madrid, Spain*

andrei.jaikin@uam.es

<http://matematicas.uam.es/~andrei.jaikin>

Proposed: Ian Agol

Seconded: Martin Bridson, Cameron Gordon

Received: 27 September 2018

Revised: 11 March 2019