

Anomalous partially hyperbolic diffeomorphisms III: Abundance and incoherence

CHRISTIAN BONATTI
ANDREY GOGOLEV
ANDY HAMMERLINDL
RAFAEL POTRIE

Let M be a closed 3–manifold which admits an Anosov flow. We develop a technique for constructing partially hyperbolic representatives in many mapping classes of M . We apply this technique both in the setting of geodesic flows on closed hyperbolic surfaces and for Anosov flows which admit transverse tori. We emphasize the similarity of both constructions through the concept of h –transversality, a tool which allows us to compose different mapping classes while retaining partial hyperbolicity.

In the case of the geodesic flow of a closed hyperbolic surface S we build stably ergodic, partially hyperbolic diffeomorphisms whose mapping classes form a subgroup of the mapping class group $\mathcal{M}(T^1S)$ which is isomorphic to $\mathcal{M}(S)$. At the same time we show that the totality of mapping classes which can be realized by partially hyperbolic diffeomorphisms does not form a subgroup of $\mathcal{M}(T^1S)$.

Finally, some of the examples on T^1S are absolutely partially hyperbolic, stably ergodic and robustly nondynamically coherent, disproving a conjecture of Rodriguez Hertz, Rodriguez Hertz and Ures (Ann. Inst. H Poincaré Anal. Non Linéaire 33 (2016) 1023–1032).

37C15, 37D30

1 Introduction

A diffeomorphism $f: M \rightarrow M$ of a closed Riemannian manifold is *partially hyperbolic* if the tangent bundle TM splits into three (nontrivial) Df –invariant continuous subbundles $TM = E^{ss} \oplus E^c \oplus E^{uu}$ such that for some $\ell > 0$, for all $x \in M$ and all unit vectors $v^\sigma \in E^\sigma(x)$ ($\sigma = ss, c, uu$) one has

$$\|Df^\ell v^{ss}\| < \min\{1, \|Df^\ell v^c\|\} \leq \max\{1, \|Df^\ell v^c\|\} < \|Df^\ell v^{uu}\|.$$

Sometimes, one uses the stronger notion of *absolute partial hyperbolicity*. This means that f is partially hyperbolic and there exist constants $\lambda_1 < 1 < \lambda_2$ such that

$$\|Df^\ell v^{ss}\| < \lambda_1 < \|Df^\ell v^c\| < \lambda_2 < \|Df^\ell v^{uu}\|.$$

The subbundles E^{ss} , E^c and E^{uu} depend on f and we will indicate this, when needed, by using a subscript, eg E_f^{ss} .

Partially hyperbolic diffeomorphisms appear naturally in many contexts and provide natural classes to study phenomena such as robust transitivity and stable ergodicity (see eg Bonatti, Díaz and Viana [6], Wilkinson [42], Carrasco, Rodriguez Hertz, Rodriguez Hertz and Ures [22] and Hammerlindl and Potrie [26]). For a long time, the known examples in dimension three were rather restrictive, and efforts were made to try to show that the behavior seen in that restricted family of examples was indeed general: such results hold, for instance, on manifolds with solvable fundamental group (see Hammerlindl and Potrie [25]). Recently new examples have started to appear; see Rodriguez Hertz, Rodriguez Hertz and Ures [38] and Bonatti, Gogolev, Parwani and Potrie [12; 8].

In this paper, we consider a closed 3–manifold and look for mapping classes (ie the diffeomorphisms up to homotopy) which contain partially hyperbolic representatives. In fact, the constructions we present here all start with an Anosov flow, and therefore we consider 3–manifolds which support Anosov flows. For some of the new examples we study specific properties of the dynamics and geometry that makes them different from the previously known behaviors (see Theorem 1.2 below).

A general fact about a 3–manifold M is that its mapping class group $\mathcal{M}(M) := \pi_0(\text{Diff}_+^1(M))$ is strongly related to the group generated by the Dehn twists associated to incompressible tori (see Johannson [29]). We will consider two cases, which correspond to different positions of the tori with respect to the Anosov flow.

In Theorem 1.3, the tori will be transverse to the flow. In Theorem 1.1, the Anosov flow will be a geodesic flow and we will perform Dehn twist along tori over closed geodesics: these tori are not transverse to the geodesic flow, but they are *Birkhoff sections* for the flow.

These two cases already appear in [8]. The novelty here is that we are able to compose an arbitrary number of such Dehn twists, and that this involves a conceptual understanding of the mechanism underlying the examples in [12; 8]. Note that a priori the mapping classes which admit partially hyperbolic representatives do not form a

subgroup of $\mathcal{M}(M)$ (eg when $M = \mathbb{T}^3$, it is not a subgroup). In both settings described above, this paper exhibits infinite subgroups of $\mathcal{M}(M)$ such that each element admits a partially hyperbolic representative.

In recent years, much of the study of partially hyperbolic systems has focused on questions of robust transitivity and stable ergodicity. When analyzing such systems, an important first property to establish is whether or not there exists an invariant foliation tangent to the center direction. Indeed, a longstanding open question, recently answered by Rodriguez Hertz, Rodriguez Hertz and Ures, was whether a partially hyperbolic system with one-dimensional center necessarily had a center foliation [38]. They showed this was not the case by constructing a counterexample on the 3–torus. This surprising and important result introduced new techniques of constructing partially hyperbolic example in the nontransitive setting. Their construction fundamentally relies on having an embedded 2–torus tangent to the center-stable direction along which the dynamical incoherence occurs. As this torus is normally attracting, it cannot be used in the construction of a stably ergodic or robustly transitive example. Further, they conjectured [38; 22] that for transitive, partially hyperbolic systems in dimension three, invariant center foliation must always exist.

In fact, this conjecture is false as a consequence of some of the new examples we present in this paper. Our results imply the following:

There is a C^1 –open set of partially hyperbolic diffeomorphisms which are both transitive and dynamically incoherent.

These examples can also be made conservative and stably ergodic. See Theorem 1.2 for a precise statement. These systems do not have center-stable or center-unstable foliations. However, they do possess unique invariant branching foliations as defined by Burago and Ivanov [18]. Because of this, we may discuss the branching center foliation defined by intersecting the center-stable and center-unstable branching foliations.

As a consequence of transitivity, this “branching” occurs everywhere. In particular, for any nonempty open subset U of M , one can find distinct center leaves L_1 and L_2 such that $L_1 \cap L_2 \cap U$ is nonempty.

Despite the branching, these center leaves are comparable to the orbits of the Anosov geodesic flow defined on M ; there is a bijective correspondence between the leaves of the branching center foliation and the orbits of the flow. In particular, compact center



Figure 1: Possible dynamics on invariant center curves. Previous examples, such as on the 3–torus, have large-scale dynamics as in the left and center figures. Examples in the current paper also have dynamics as shown on the right. Notice that there may be more than one fixed point in the leaves, but they should all lie in a compact interval.

leaves are associated to periodic orbits of the flow. However, the partially hyperbolic dynamics on these leaves behaves very differently from the flow:

There is a C^1 open set of partially hyperbolic diffeomorphisms such that the union of circles tangent to E^c is dense in M , but none of these circles is periodic.

In light of this result, any center leaf which has a periodic point must be a line. In previously known examples, the dynamics on such periodic center lines is either contracting on large scales or expanding on large scales as depicted in Figure 1, left and center. The new systems studied in the current paper have periodic center lines which are contracting on one end and expanding on the other, as shown in Figure 1, right. This large-scale behavior of the leaves is a key tool in proving dynamical incoherence.

We also point out that, while in “small manifolds” a stronger notion called *absolute* partial hyperbolicity is enough to guarantee dynamical coherence (see Brin, Burago and Ivanov [16] and Hammerlindl and Potrie [25]), our new examples are absolutely partially hyperbolic, thus producing the first such examples of dynamical incoherence in the three-dimensional setting.

We begin by stating our result in the geometric setting.

1.1 Partially hyperbolic diffeomorphisms on the unit tangent bundle of a hyperbolic surface

Let S denote the surface of genus $g \geq 2$, and let T^1S denote its unit tangent bundle (equivalently the circle bundle over S whose Euler class is $2-2g$, the Euler characteristic of S). Let $\mathcal{M}(S)$ denote the group of homotopy (or equivalently isotopy) classes of orientation-preserving diffeomorphisms of S ; in other words, $\mathcal{M}(S) = \pi_0(\text{Diff}_+(S))$.

Given a diffeomorphism $\varphi: S \rightarrow S$, its normalized differential $T\varphi(v) = D\varphi v / \|D\varphi v\|$ is a diffeomorphism of T^1S and so $\text{Diff}_+(S)$ acts on the unit tangent bundle T^1S , yielding an injective homomorphism $\iota: \mathcal{M}(S) \hookrightarrow \mathcal{M}(T^1S)$ (see [29, Proposition 25.3]).

The main theorem in this setting is the following:

Theorem 1.1 *For any $\varphi \in \mathcal{M}(S)$ there exists a diffeomorphism $f: T^1S \rightarrow T^1S$ such that*

- *f belongs to $\iota(\varphi)$,*
- *f is absolutely partially hyperbolic,*
- *f preserves the volume and is stably ergodic.*

In Section 3.1 we use the results from Hammerlindl, Potrie and Shannon [27] to see that not every mapping class of T^1S can be realized and, moreover, that the set of mapping classes which are realizable do not form a subgroup of $\mathcal{M}(T^1S)$.

1.2 Dynamical incoherence

For the examples given by Theorem 1.1 applied to a *pseudo-Anosov* mapping class φ (see eg Farb and Margalit [24] for a definition of pseudo-Anosov homeomorphism) we are able to obtain properties for f which were unexpected. Recall that a partially hyperbolic diffeomorphism is called *dynamically coherent* if there exist f -invariant foliations \mathcal{F}^{cs} and \mathcal{F}^{cu} tangent to $E^{cs} := E^{ss} \oplus E^c$ and $E^{cu} := E^c \oplus E^{uu}$, respectively (and, therefore, there is also an invariant center foliation obtained by taking intersection). Otherwise, we say that f is *dynamically incoherent*.

Recall that the strong stable and unstable bundles are always uniquely integrable for dynamical reasons (see eg [26]). In higher dimensions, when the center bundle has dimension at least two, there are examples where the center bundle is smooth but does not satisfy the Frobenius integrability condition (see Burns and Wilkinson [19]). When the center bundle is one-dimensional the problem of integrability only comes from the lack of regularity. Assuming absolute partial hyperbolicity, geometric conditions on the strong foliations (which are always satisfied on the torus \mathbb{T}^3) are known to imply dynamical coherence (see Brin, Burago and Ivanov [15; 16]). For certain families of 3-manifolds, exact conditions for dynamical coherence are known (see [22; 26]). However, there is no known general criterion for deciding whether the center bundle is integrable.

This motivated Burago and Ivanov [18] to construct objects called *branching foliations* which exist for any 3-dimensional partially hyperbolic system and serve as substitutes for the true invariant foliations. Originally, these branching foliations were used only

as a tool to establish dynamical coherence. Later, Rodriguez Hertz, Rodriguez Hertz and Ures [38] constructed a concrete example which has branching foliations, but not true foliations. For this example the branching occurs exactly on a finite collection of attracting or repelling 2-tori (which is incompatible with transitivity and absolute partial hyperbolicity). They further conjectured [38] that the existence of such tori is the unique obstruction for dynamical coherence (see also [22]).

Theorem 1.2 provides counterexamples to this conjecture:

Theorem 1.2 *For any pseudo-Anosov mapping class $\varphi \in \mathcal{M}(S)$ there exists a diffeomorphism $f: T^1S \rightarrow T^1S$ such that*

- *f belongs to $\iota(\varphi)$,*
- *f is absolutely partially hyperbolic,*
- *f preserves the volume and is stably ergodic,*
- *f is robustly transitive,*
- *f is robustly dynamically incoherent.*

For these examples we also establish minimality of the strong stable and unstable foliations. This implies that the branching sets of the branching foliations are dense in the whole manifold. Distributions with this kind of behavior were already known to exist but not associated to partially hyperbolic dynamics; see Bonatti and Franks [7].

It is natural to expect that given a maximal curve η tangent to center distribution E^c the stable saturation $W^s(\eta)$ is a center-stable leaf. However, it was realized early on — see Burns and Wilkinson [13] — that theoretically the stable saturation $W^s(\eta)$ may fail to coincide with the full center-stable leaf. This may happen if $W^s(\eta)$ is not *complete*. The example in [38] also showed that such hypothetical behavior indeed may occur: $W^s(\eta)$ are not complete and their nonempty boundary is contained in the repelling tori. One could believe that the minimality of the strong foliations could imply completeness, but our examples show that this is not so.

Another important tool for the study of the dynamics of a partially hyperbolic diffeomorphism is the understanding of the dynamics in the center leaves. Our examples exhibit a new type of behavior: one can show that there are many compact center curves but none of them is periodic, and that there exist simultaneously contracting, repelling and saddle-node center leaves (see eg [6, Section 7.3.4]). In particular, the examples give a complete answer to [6, Problem 7.26] (see also [8, Section 3.8]). See Remark 5.13 for a detailed explanation of these phenomena.

1.3 Partially hyperbolic diffeomorphisms associated to a family of tori transverse to an Anosov flow

Given a torus T embedded in a 3-manifold M , we associate an element $\tau(T, \gamma) \in \mathcal{M}(M)$, called *the Dehn twist along T directed by γ* , to each homotopy class of closed curves $\gamma \in \pi_1(T)$ (see Section 4.1 for a precise definition).

Given a torus $(T, \mathcal{F}, \mathcal{G})$ endowed with a pair of transverse C^1 -foliations \mathcal{F} and \mathcal{G} , we consider the subset $G(T, \mathcal{F}, \mathcal{G})$ of homotopy classes $\gamma \in \pi_1(T)$ such that there exists a C^1 -continuous loop $\{\psi_t\}_{t \in \mathbb{R}/\mathbb{Z}}$ of diffeomorphisms of T such that

- $\psi_0 = \text{id}$,
- $\psi_t(\mathcal{F})$ is transverse to \mathcal{G} for all $t \in \mathbb{R}/\mathbb{Z}$,
- the homotopy class of the loop $\{\psi_t(x)\}_{t \in \mathbb{R}/\mathbb{Z}}$, for $x \in T$ is γ .

It was shown by Bonatti and Zhang [14] that either $G(T, \mathcal{F}, \mathcal{G}) = \pi_1(T) \simeq \mathbb{Z}^2$ or both foliations contain circle leaves in the same homotopy class and $G(T, \mathcal{F}, \mathcal{G})$ is the cyclic group \mathbb{Z} generated by this homotopy class.

Now, if X is a C^2 -Anosov flow on a 3-manifold M and if $T \subset M$ is a torus transverse to X then the center-stable and unstable foliations of X induce a pair of transverse C^1 -foliation (F_T^s, F_T^u) on T . Then we denote by $G(T, X)$ the group $G(T, F_T^s, F_T^u)$.

Given an Anosov flow X on a 3-manifold M we denote by $\mathcal{T}(X) \subset \mathcal{M}(M)$ the subgroup generated by the Dehn twists $\tau(T, \gamma)$ where T is a torus transverse to X and γ belongs to $G(T, X)$.

Theorem 1.3 *Let X be an Anosov flow on a closed 3-manifold M . Then every element of $\mathcal{T}(X)$ contains an absolutely partially hyperbolic representative.*

In certain cases, this theorem produces partially hyperbolic representatives in virtually all mapping classes.

Corollary 1.4 *For each $n \geq 1$ there exists a closed 3-dimensional graph manifold M_n and a finite-index abelian subgroup $G \simeq \mathbb{Z}^{2n} \oplus \mathbb{Z}/n\mathbb{Z}$ of the mapping class group $\mathcal{M}(M_n)$ such that each mapping class in G can be represented by an absolutely partially hyperbolic, volume-preserving diffeomorphism.*

Manifolds M_n above are n -fold cyclic covers of the ambient manifold of the Bonatti–Langevin Anosov flow [10].

1.4 Questions

It is our opinion that this paper has opened even more questions than the previous one [8]. In particular, it remains a challenge to better understand these new examples, which have several new features which contrast our previous beliefs and should be now taken into account in the study of the dynamics of partially hyperbolic diffeomorphisms.

We state here some questions which we believe arise naturally from our work, but the reader is of course invited to add more.

We have constructed many mapping classes which admit partially hyperbolic representatives on manifolds which admit Anosov flows. It is natural to ask for characterization of all such mapping classes.

Problem 1.5 *Given a closed 3–manifold M admitting an Anosov flow, determine which mapping classes admit partially hyperbolic representatives.*

In this direction, the following question is natural:

Question 1.6 *For a manifold M , let \mathcal{P} be those elements of $\mathcal{M}(M)$ admitting a partially hyperbolic representative. When is \mathcal{P} a subgroup of $\mathcal{M}(M)$? When is \mathcal{P} a finite-index group? When does it equal $\mathcal{M}(M)$?*

In this direction, Corollary 1.4 shows that on some 3–manifolds, the classes which can be realized by partially hyperbolic diffeomorphisms contain a finite-index subgroup. See also Section 3.1 for results in this direction.

This paper reduces the realization of mapping classes admitting partially hyperbolic diffeomorphisms to understanding geometry of the subbundles and how can they be made h –transverse (see Definition 2.1). Under certain assumptions on the Anosov flow, we have tools to check that certain Dehn twists along certain tori respect transversality between the subbundles of the Anosov flow. In particular, our understanding seems to be complete in the case when tori are transverse to the Anosov flow. However, there are other configurations where h –transversality is not completely understood. In particular, we address the case of Birkhoff tori only in the geometric setting — we perform Dehn twists using the geometric considerations of [8]. The general case of Birkhoff tori remains to be understood.

Problem 1.7 *Determine which Dehn twists respect transversalities between the bundles of an Anosov flow.*

Understanding the above problem may yield even more mapping classes with partially hyperbolic representatives and provide a way to attack Problem 1.5. Notice that there are some general results about Anosov flows which indicate that the study of Birkhoff tori might be sufficient (see Barbot [1]).

Further questions arise about the geometry and dynamics of these new examples. Recall that for certain examples which realize pseudo-Anosov mapping class we are able to establish dynamical incoherence as well as noncompleteness of stable manifolds of center leaves (see Corollary 5.12). Also these examples possess saddle-node center leaves. However, for the general examples, many questions arise:

Question 1.8 *Which of the new examples are dynamically coherent (compare [12])? Are the dynamically coherent examples plaque-expansive? Can the transitive examples transverse to tori be always made volume-preserving? Robustly transitive?*

The examples given by Theorem 1.2 come from the pseudo-Anosov mapping class and, hence, have some strong and better-understood dynamical and geometric properties. A natural question that one can pose is:

Question 1.9 *Is every absolutely partially hyperbolic diffeomorphism of T^1S homotopic to the lift of a pseudo-Anosov diffeomorphism of S transitive (or at least chain-recurrent)?*

Compare with Bonatti and Guelman [9], Shi [39] and Potrie [34].

Outline of the paper

The paper is organized as follows. In Section 2 we present a flexible criterion, which conceptualizes the results of [12; 8] and allows us to compose maps while retaining the partially hyperbolic property.

We apply this criterion in Sections 3 and 4 to prove Theorems 1.1 and 1.3, respectively. To be able to apply the criterion we need specific constructions for each case, using the ideas in [8] to prove Theorem 1.1 and the ones of [14] for Theorem 1.3. In addition, we show in Section 3.1 that not every mapping class can be realized as a partially hyperbolic diffeomorphism and in Section 4.8 that there are manifolds with infinite mapping class for which virtually every mapping class can be realized as a partially hyperbolic diffeomorphism.

Finally, in Section 5 we show dynamical incoherence of some of the examples presented here (proving Theorem 1.2), and in particular we show some relevant properties about the structure of the center curves of such examples.

2 A criterion for partial hyperbolicity

We provide a general criterion for partial hyperbolicity based on cone fields. This criterion is obtained by extracting and formalizing the key point of the argument in the previous paper [8] (see also [26]). The criterion has no restrictions on the dimensions of subbundles.

Definition 2.1 Let $f: M \rightarrow M$ and $g: N \rightarrow N$ be partially hyperbolic diffeomorphisms and let $h: M \rightarrow N$ be a diffeomorphism. We say that f is h -transverse to g and use the notation

$$f \rightsquigarrow_h g$$

if $Dh(E_f^{uu})$ is transverse to E_g^{cs} and $Dh^{-1}(E_g^{ss})$ is transverse to E_f^{cu} .

Remark 2.2 It is easy to see that $f \rightsquigarrow_h g$ holds if and only if $g^{-1} \rightsquigarrow_{h^{-1}} f^{-1}$. Further, if $f \rightsquigarrow_h g$ then $f \rightsquigarrow_{g^l h f^k} g$ for any nonnegative integers k and l .

The h -transversality property is transitive in the following sense:

Lemma 2.3 Let $f_i: M_i \rightarrow M_i$ for $i \in \{1, \dots, \ell\}$ be a family of partially hyperbolic diffeomorphisms and let $h_j: M_j \rightarrow M_{j+1}$ for $j \in \{1, \dots, \ell - 1\}$ be a family of diffeomorphisms such that f_i is h_i -transverse to f_{i+1} . Then there exist positive $m_2, \dots, m_{\ell-1}$ such that, for every $n_i \geq m_i$,

$$f_1 \rightsquigarrow_{h_{\bar{n}}} f_{\ell},$$

where $h_{\bar{n}} = h_{\ell} \circ f_{\ell-1}^{n_{\ell-1}} \circ h_{\ell-1} \circ \dots \circ f_2^{n_2} \circ h_1$.

Proof We will show that

$$f_1 \rightsquigarrow_{h_1} f_2 \rightsquigarrow_{h_2} f_3$$

implies that for large n one has that $f_1 \rightsquigarrow_{\hat{h}} f_3$ with $\hat{h} = h_2 \circ f_2^n \circ h_1$, as the statement follows by several applications of this.

As $Dh_1(E_{f_1}^{uu})$ is transverse to $E_{f_2}^{cs}$ it follows that $Df_2^n(Dh_1(E_{f_1}^{uu}))$ converges uniformly to $E_{f_2}^{uu}$. The fact that $Dh_2(E_{f_2}^{uu})$ is transverse to $E_{f_3}^{cs}$ allows to conclude that $D(h_2 \circ f_2^n \circ h_1)(E_{f_1}^{uu})$ is transverse to $E_{f_3}^{cs}$.

A symmetric argument gives the other condition. □

Now, we can formulate the criteria for partial hyperbolicity:

Proposition 2.4 *Let $f: M \rightarrow M$ be an absolutely partially hyperbolic diffeomorphism and $h: M \rightarrow M$ a diffeomorphism of M such that $f \rightsquigarrow_h f$. Then there exists $N > 0$ such that $f^n \circ h$ is absolutely partially hyperbolic for any $n \geq N$.*

Proof We prove that $f^n \circ h$ has a weak absolutely partially hyperbolic splitting of the form $E^{cs} \oplus E^{uu}$. A similar proof shows that $f^n \circ h$ has a splitting of the form $E^{ss} \oplus E^{cu}$, and together these show that $f^n \circ h$ is absolutely partially hyperbolic. We will use the classical cone criteria; see eg [6, Appendix B].

Up to replacing f by an iterate, there are $1 < \gamma < \lambda$ such that $\|Df v^c\| < \gamma < \lambda < \|Df v^u\|$ for all unit vectors $v^c \in E_f^{cs}$ and $v^u \in E_f^{uu}$. By properties of partial hyperbolicity, there is a cone family $\mathcal{C}^{uu} \subset TM$ such that $Df(\mathcal{C}^{uu}) \subset \mathcal{C}^{uu}$,

$$\bigcap_{n \geq 0} Df^n(\mathcal{C}^{uu}) = E_f^{uu}$$

and $\|Df v\| \geq \lambda \|v\|$ for all $v \in \mathcal{C}^{uu}$. Since

$$\bigcap_{n \geq 0} D(h \circ f^n)(\mathcal{C}^{uu}) = Dh(E_f^{uu})$$

is transverse to E_f^{cs} , it follows that $D(h \circ f^k)(\mathcal{C}^{uu}) \cap E_f^{cs} = \{0\}$ for some integer k . By replacing \mathcal{C}^{uu} with $Df^k(\mathcal{C}^{uu})$, we freely assume that $Dh(\mathcal{C}^{uu}) \cap E^{cs} = \{0\}$.

For any nonzero vector $v \in \mathcal{C}^{uu}$, the angle between $D(f^n \circ h)(v)$ and E_f^{uu} tends to zero as $n \rightarrow +\infty$ and so there is N such that $D(f^n \circ h)(v) \in \mathcal{C}^{uu}$ for all $n > N$. By a compactness argument, this N may be chosen independently of v . Further, there is a constant $a > 0$ such that $\|D(f^n \circ h)v\| \geq a\lambda^n \|v\|$ for all $n > N$ and $v \in \mathcal{C}^{uu}$. If n is sufficiently large, then $a\lambda^n > 1$.

By a similar argument (and possibly adjusting a and N), there is a cone family \mathcal{C}^{cs} such that, if $v \in \mathcal{C}^{cs}$, then

$$D(f^n \circ h)^{-1}v \in \mathcal{C}^{cs} \quad \text{and} \quad \|D(f^n \circ h)^{-1}v\| \geq \frac{a}{\gamma^n} \|v\|$$

for all $n > N$. Note also that $\gamma^n/a < a\lambda^n$ for all large n . These conditions together imply that $f^n \circ h$ has a weak absolutely partially hyperbolic splitting of the form $E^{cs} \oplus E^{uu}$. \square

Indeed, it follows from the proof of the proposition that we have:

Scholium 2.5 *In the setting of Proposition 2.4, given any sufficiently narrow cone field \mathcal{C}^{uu} (resp. \mathcal{C}^{ss}) in M around E^{uu} (resp. around $Dh^{-1}(E^{ss})$), there exists N such that $f^n \circ h$ preserves the cone fields for $n > N$.*

Remark 2.6 If one writes $F_{n,m} = f^n \circ h \circ f^m$ it follows that $F_{n,m}$ is smoothly conjugate to $f^{n+m} \circ h$ via f^{-m} . It follows that for $n+m$ sufficiently large, $F_{n,m}$ is absolutely partially hyperbolic. Further, if both n and m are large, one gets that the bundles of $F_{n,m}$ are as close to those of f as one wants.

Using Lemma 2.3 one gets:

Proposition 2.7 *Let f and g be two partially hyperbolic diffeomorphisms on M such that there is a C^1 -continuous path $\{f_t\}_{t \in [0,1]}$ of partially hyperbolic diffeomorphisms such that $f_0 = f$ and $f_1 = g$. Then there exists a diffeomorphism h of M such that f is h -transverse to g .*

Moreover:

- h is isotopic to f^N for some large N . In particular, if f (and therefore g) is isotopic to the identity, then h can be chosen to be isotopic to the identity.
- If all the diffeomorphisms in the path $\{f_t\}$ preserve a smooth volume vol_t which varies smoothly with t , then one can choose h so that it maps vol_0 to vol_1 .

Proof First notice that, by continuity of the bundles with respect to t , and by compactness of the parameters set $[0, 1]$, there is $\varepsilon > 0$ such that, for any $s, t \in [0, 1]$ with $|t - s| \leq \varepsilon$, one has

$$f_t \rightsquigarrow_{\text{id}} f_s.$$

Then, one applies the above transitivity lemma to obtain h -transversality. The fact that h is isotopic to a power of f is immediate from the lemma and the fact that all f_t are isotopic to f .

To see that one can choose h to preserve volume, notice that one can find, using Moser's trick (see [30, Theorem 5.1.27]), a path φ_t of diffeomorphisms sending vol_t into vol_0 . It follows that the diffeomorphisms $\{\varphi_t \circ f_t \circ \varphi_t^{-1}\}$ form a smooth path of diffeomorphisms which preserve vol_0 . So, applying the previous reasoning to the path $\varphi_t \circ f_t \circ \varphi_t^{-1}$ which connects f and $\hat{g} = \varphi_1 \circ g \circ \varphi_1^{-1}$, one obtains that f is \hat{h} -transverse to \hat{g} with \hat{h} a vol_0 -preserving diffeomorphism. Because φ_1 conjugates g and \hat{g} we see that $h = \varphi_1^{-1} \circ \hat{h}$ is the posited diffeomorphism. \square

Remark 2.8 We have worked in the specific setting that is needed for this paper. It is not hard to see that the concept of h -transversality can be generalized to give general statements about *dominated splittings*, *normal hyperbolicity*, etc. Moreover, in Proposition 2.4 one just need to ensure transversality of certain flags and domination between bundles; then, to obtain partial (or absolute partial, or r -normal) hyperbolicity, it is enough to have the property for just one of the diffeomorphisms involved. We leave these statements to the reader as they will not be used here.

3 Geodesic flows on higher-genus surfaces

In this section we prove Theorem 1.1. Let us first rewrite and reinterpret in our language a key proposition of the proof of [8, Theorems 2.8 and 2.10]. Recall that if γ is a simple closed curve on a surface S then τ_γ denotes the element of $\mathcal{M}(S)$ containing the Dehn twist along γ .

Proposition 3.1 *Consider a closed oriented surface S and let γ be the free homotopy class of a simple closed curve in S . Then there exist*

- a hyperbolic metric μ on S , and
- a volume-preserving diffeomorphism $h: T^1S \rightarrow T^1S$ in the mapping class $\iota(\tau_\gamma)$

such that the time-one map f of the geodesic flow of (S, μ) is h -transverse to itself.

Proof This follows from [8, Proposition 2.9]. In that proposition it is shown that one can choose a sequence of metrics μ_n in S and a sequence of maps $h_n = D\rho_n$ in the class $\iota(\tau_\gamma)$ such that the bundles of the geodesic flow on (S, μ_n) are mapped by h_n making arbitrarily small angle as $n \rightarrow \infty$ with themselves, which implies h_n -transversality for a sufficiently large n (this is called *ph-respectful* in [8]).

In [8, Theorem 2.10] it is explained how to modify the map to make it volume-preserving using local coordinates around γ . \square

Proof of Theorem 1.1 Let $\varphi \in \mathcal{M}(S)$. According to the Dehn–Lickorish theorem (see for instance [24, Theorem 4.1]) one can write φ as a composition $\varphi = \tau_{\gamma_\ell}^{\varepsilon_\ell} \circ \dots \circ \tau_{\gamma_1}^{\varepsilon_1}$ where γ_i are simple closed curves (not necessary disjoint nor distinct) and $\varepsilon_i \in \{-1, 1\}$.

Proposition 3.1 yields hyperbolic metrics μ_1, \dots, μ_ℓ on S associated to γ_i for $i = 1, \dots, \ell$ and volume-preserving diffeomorphisms $h_i: T^1 S \rightarrow T^1 S$ for $i = 1, \dots, \ell$ such that each time-one map f_i of the geodesic flow for μ_i is h_i –transverse to itself and h_i is in the mapping class of $\iota(\tau_{\gamma_i}^{\varepsilon_i})$.

Recall that the Teichmüller space of hyperbolic metrics on S is diffeomorphic to an open ball and, hence, is path-connected (see eg [24, Chapter 10]). Thus f_i is isotopic to f_{i+1} through a path of time-one maps of geodesic flows which correspond to a path of hyperbolic metrics connecting μ_i to μ_{i+1} . Clearly the path connecting f_i to f_{i+1} consists of volume-preserving partially hyperbolic diffeomorphisms (the preserved volume varying smoothly with the parameter).

By Proposition 2.7, we obtain diffeomorphisms $g_i: T^1 S \rightarrow T^1 S$ such that

- f_i is g_i –transverse to f_{i+1} ,
- g_i sends the Liouville volume of μ_i to that of μ_{i+1} for $i \in \mathbb{Z}/\ell\mathbb{Z}$,
- g_i is isotopic to the identity.

Therefore we have

$$f_1 \rightsquigarrow_{h_1} f_1 \rightsquigarrow_{g_1} f_2 \rightsquigarrow_{h_2} f_2 \rightsquigarrow_{g_2} \dots \rightsquigarrow_{g_{\ell-1}} f_\ell \rightsquigarrow_{h_\ell} f_\ell \rightsquigarrow_{g_\ell} f_1$$

Now, according to Proposition 2.4 (and Lemma 2.3), for every large m the diffeomorphism

$$F = f_1^m \circ g_\ell \circ f_\ell^m \circ h_\ell \circ \dots \circ h_2 \circ f_2^m \circ g_1 \circ f_1^m \circ h_1$$

is absolutely partially hyperbolic. By construction it is volume-preserving. Finally, as the g_i and the f_i are isotopic to the identity, the isotopy class of F is the same as that of $h_\ell \circ \dots \circ h_1$. Hence the mapping class of F is given by $\iota(\tau_{\gamma_\ell}^{\varepsilon_\ell}) \circ \dots \circ \iota(\tau_{\gamma_1}^{\varepsilon_1}) = \iota(\varphi)$ as ι is a group homomorphism.

Finally, according to [11; 37], there exists a small perturbation of F which is stably ergodic, which completes the proof. □

Remark 3.2 The proof can be adapted to give that for any given hyperbolic metric μ with geodesic flow \mathcal{G}_t and $\varphi \in \mathcal{M}(S)$ there exists a volume-preserving diffeomorphism h in $\iota(\varphi)$ such that \mathcal{G}_t is h transverse to itself.

Remark 3.3 If we choose a reducible mapping class, ie one that leaves invariant a simple closed geodesic, then we can apply the same reasoning as in [8, Proposition 2.12] to make a robustly transitive and stably ergodic perturbation of the example. Robust transitivity can also be obtained in other contexts, as we will explain in Section 5.6.

Remark 3.4 Every closed surface S admits a metric of constant negative curvature with an orientation-reversing isometry, and this lifts to an h -transversality for the geodesic flow of the metric with itself. This allows us to extend Theorem 1.1 to the case of an orientation-reversing mapping classes τ of $\mathcal{M}(S)$. Note that in this case $\iota(\tau)$ is orientation-preserving because the derivative reverses orientation on the fibers.

Remark 3.5 It was pointed out to us by Livio Flamino that the above proof also admits the following variant. Equip S with hyperbolic metrics μ_1 and μ_2 such that $\phi: (S, \mu_1) \rightarrow (S, \mu_2)$ is an isometry. Clearly ϕ defines an h -transversality for corresponding geodesic flows. Then connect μ_1 and μ_2 by a path of hyperbolic metrics. Then, similarly to the proof above, by subdividing the path into small intervals and using $\text{id}: S \rightarrow S$ as h -transversalities, one realizes the mapping class of ϕ by a partially hyperbolic diffeomorphism.

3.1 Some obstructions

Recall (see [29, Chapter 25]) that the mapping class group of T^1S fits into the short exact sequence

$$1 \rightarrow H_1(S, \mathbb{Z}) \rightarrow \mathcal{M}(T^1S) \rightarrow \mathcal{M}(S) \rightarrow 1.$$

Theorem 1.1 implies that the set of mapping classes of T^1S which are realized by partially hyperbolic diffeomorphisms surjects onto the whole $\mathcal{M}(S)$. Here we show the following:

Theorem 3.6 *If a diffeomorphism $f: T^1S \rightarrow T^1S$ is partially hyperbolic and its mapping class projects to the identity in $\mathcal{M}(S)$, then f is isotopic to the identity.*

The proof assumes familiarity with branching foliations [18] and uses a result from [27].

Proof Let z be the element of $\pi_1(T^1S)$ given by a path around a fiber of the circle fibering. Equivalently, z is a generator for the cyclic subgroup which is the center of $\pi_1(T^1S)$. Every element of $\pi_1(T^1S)$ may then be written in the form $[\gamma'] \cdot z^k$ where k is an integer and the curve γ' in T^1S is the derivative of a closed geodesic γ

in S . Up to replacing f with f^2 , we may assume $f_*(z) = z$ and so, if f_* is not the identity, there is a closed geodesic γ such that $f_*([\gamma']) = [\gamma'] \cdot z^k$ for some nonzero k .

By [27, Section 5.2] we know that the center-stable branching foliation $\mathcal{F}_{\text{bran}}^{cs}$ for f is horizontal. To be precise, we may first approximate $\mathcal{F}_{\text{bran}}^{cs}$ by a true foliation \mathcal{F}_ϵ where every leaf of the true foliation is isotopic to a leaf of the branching foliation and vice versa. Then there exists a smooth conjugacy which is isotopic to the identity and which puts \mathcal{F}_ϵ transverse to the fibers of the circle fibering. Using a semiconjugacy result of Matsumoto [31], one may further show the following: for every leaf \mathcal{L} of the weak stable foliation of the geodesic flow on T^1S , there is a leaf L of $\mathcal{F}_{\text{bran}}^{cs}$ isotopic to \mathcal{L} and vice versa. In particular, one of the leaves L of $\mathcal{F}_{\text{bran}}^{cs}$ is isotopic to the weak stable leaf containing γ' and this implies that L itself contains a closed curve isotopic to γ' .

Both the leaf L and its image $f(L)$ are cylinders and $\pi_1(f(L))$ is the cyclic subgroup of $\pi_1(T^1S)$ generated by $f_*([\gamma']) = [\gamma'] \cdot z^k$. However, the semiconjugacy implies that $f(L)$ is isotopic to a leaf of the weak stable foliation of the geodesic flow. Since no such leaf has such a fundamental group, we arrive at a contradiction. \square

Corollary 3.7 *The set of mapping classes realized by partially hyperbolic diffeomorphisms does not form a subgroup of $\mathcal{M}(T^1S)$.*

Proof It is enough to show that one can create different mapping classes projecting to the same mapping class in $\mathcal{M}(S)$. Then, by composing one with the inverse of the other, one obtains a nontrivial mapping class projecting to the identity, which would yield a contradiction. To see that one has examples in such mapping classes one just has to observe that Theorem 1.1 produces mapping classes projecting into all of $\mathcal{M}(S)$ and that the mapping classes which can be realized by partially hyperbolic diffeomorphisms are closed under conjugacy. For example, one can take two homologically nontrivial curves α and β in S whose intersection number is one. Then denote by $f_\alpha: T^1S \rightarrow T^1S$ a partially hyperbolic diffeomorphism in the mapping class of the (differential of) the Dehn twist along α and denote by $v_\beta: T^1S \rightarrow T^1S$ a diffeomorphism whose mapping class projects to the identity in $\mathcal{M}(S)$ but for which β' is not homotopic to $v_\beta(\beta')$. Then f_α and $v_\beta \circ f_\alpha \circ v_\beta^{-1}$ are partially hyperbolic diffeomorphisms from different mapping classes which project to the same mapping class in $\mathcal{M}(S)$. \square

4 Anosov flows transverse to tori

In this section we prove Theorem 1.3.

4.1 Dehn twists in three dimensions

Here, we introduce a three-dimensional analogue of two-dimensional Dehn twists. Consider a two-dimensional torus T embedded in a three-dimensional manifold M . Let U be a tubular neighborhood of this torus. Then we may realize U as the image of an embedding $i: \mathbb{T}^2 \times (-1, +1) \rightarrow M$, where \mathbb{T}^2 is the standard 2-torus given by the quotient $\mathbb{R}^2/\mathbb{Z}^2$. Further assume that i maps $\mathbb{T}^2 \times \{0\}$ to T . Consider now a curve γ embedded in T . After a homotopy, we can reduce to the case where γ is the image of a linear curve in \mathbb{T}^2 . That is, there is a vector $(p, q) \in \mathbb{Z}^2$ such that γ is the image under i of the set

$$\{(tp, tq, 0) : t \in [0, 1]\} \subset \mathbb{T}^2 \times (-1, +1).$$

We now define *the Dehn twist along T directed by γ* . This will be an element of the mapping class group of M and so it is enough to define a diffeomorphism $\tau: M \rightarrow M$ representing this element. In fact, this map will be the identity outside of U and so we first define a Dehn twist on $\mathbb{T}^2 \times (-1, +1)$. Let $h: (-1, +1) \rightarrow [0, 1]$ be a smooth bump function such that $h(z) = 0$ for $z < -\frac{1}{2}$ and $h(z) = 1$ for $z > \frac{1}{2}$. Then define $\phi: \mathbb{T}^2 \times (-1, +1) \rightarrow \mathbb{T}^2 \times (-1, +1)$ by

$$\phi(x, y, z) = (x + ph(z), y + qh(z), z).$$

Finally, define $\tau: M \rightarrow M$ by $\tau(v) = i \circ \phi \circ i^{-1}(v)$ if $v \in U$, and $\tau(v) = v$ otherwise.

4.2 Anosov flows transverse to tori

Let X be an Anosov vector field on a 3-manifold M and let $T \subset M$ be a torus transverse to X . It is well known that T must be incompressible [17]. A systematic study of Anosov flows transverse to tori has been recently carried out in [2], yet some questions still remain open.

4.3 Connecting different vector fields

The following proposition is immediate. We state it to emphasize the parallel between this case and the proof of Theorem 1.1, where an analogous statement was obtained using the connectedness of the Teichmüller space.

Proposition 4.1 *Let X be a smooth Anosov vector field on M and let $\rho_1, \rho_2: M \rightarrow (0, 1]$ be smooth functions. Then the time-one maps of the flows generated by the vector fields $\rho_1 X$ and $\rho_2 X$ are isotopic through partially hyperbolic diffeomorphisms.*

As an immediate consequence of this proposition and Proposition 2.7 we have the following:

Corollary 4.2 *Let X be a smooth Anosov vector field on M and let $\rho_1, \rho_2: M \rightarrow (0, 1]$ be smooth functions. Then there exists a diffeomorphism $g: M \rightarrow M$ such that the time-one map of the flow of $\rho_1 X$ is g -transverse to the time-one map of the flow of $\rho_2 X$. Moreover, g is isotopic to the identity.*

Remark 4.3 If X is volume-preserving, then, again by Proposition 2.7, one can choose g so that it sends the volume preserved by $\rho_1 X$ to the volume preserved by $\rho_2 X$.

4.4 Coordinates in flow boxes

We consider an Anosov vector field X transverse to a torus T . Without loss of generality we can assume that we have chosen a metric in M such that X is everywhere orthogonal to T and such that $\|X(x)\| = 1$ for all $x \in M$.

Since X is transverse to T we obtain that the foliations W_X^{cs} and W_X^{cu} induce (transverse) foliations F_X^s and F_X^u on T by taking the intersections.

We can consider coordinates $\theta_T: T \rightarrow \mathbb{T}^2$ where $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ is equipped with the usual (x, y) -coordinates (mod 1). We denote by F^s and F^u the foliations $\theta_T(F_X^s)$ and $\theta_T(F_X^u)$, respectively.

For each $\eta > 0$ consider a manifold \widehat{M}_η diffeomorphic to M with an Anosov vector field \widehat{X}^η obtained as follows:

- first cut M along T to obtain a 3-manifold M' with two boundary components T_0 and T_1 such that the vector field X points outwards on T_0 and inwards on T_1 ,
- glue in a flow box $\mathcal{U}_\eta = [0, \eta] \times T$ to M' so that $\{0\} \times T$ glues to T_0 and $\{\eta\} \times T$ glues to T_1 ,
- define \widehat{X}^η to be the unit horizontal vector field on $[0, \eta] \times T$ which glues well with X .

Denote by \widehat{T} the torus $T_0 \cong \{0\} \times T$ in \widehat{M}_η which can be identified with T in the obvious way. We will simply write \widehat{X} for \widehat{X}^η and we will write \widehat{X}_t for the time- t map of the flow generated by \widehat{X} . It is straightforward to check that the flow generated by \widehat{X} is Anosov (even if the constants worsen as η increases). The fact that

\widehat{M}_η is diffeomorphic to M is also immediate because T has a neighborhood in M diffeomorphic to $(0, 1) \times T$.

The following proposition will be helpful to choose nice deformations of the flow in order to apply Proposition 2.4 and Corollary 4.2:

Proposition 4.4 *There exists a smooth function $\rho_\eta: M \rightarrow (0, 1]$ such that the vector field $\rho_\eta X$ is smoothly conjugate to \widehat{X}^η .*

Proof Pick a small $\varepsilon > 0$ and an increasing diffeomorphism $\psi: [-\varepsilon, \varepsilon + \eta] \rightarrow [-\varepsilon, \varepsilon]$ such that $\psi' \leq 1$ and $\psi - x$ is C^∞ -flat at the endpoints. Let $T_\varepsilon = \bigcup_{|t| \leq \varepsilon} X_t(T) \subset M$. In a similar way, let $\widehat{T}_\varepsilon^\eta = \bigcup_{-\varepsilon \leq t \leq \eta + \varepsilon} \widehat{X}_t(T_0) \subset \widehat{M}_\eta$. Clearly these are smooth embeddings of $[-\varepsilon, \varepsilon] \times T$ and $[-\varepsilon, \eta + \varepsilon] \times T$ in M and \widehat{M}_η , respectively, so that orbits of the flows correspond to horizontal lines. Now, choose a diffeomorphism from \widehat{M}_η to M which is the identity outside $\widehat{T}_\varepsilon^\eta$ and identifies $\widehat{T}_\varepsilon^\eta$ with T_ε via $\psi \times \text{id}$. This diffeomorphism yields the conjugacy and the posited time change $\rho_\eta X$. \square

Recall that the inserted flow box $\mathcal{U}_\eta = [0, \eta] \times T \subset \widehat{M}_\eta$ can be viewed a sweep-out of T up to time η :

$$\mathcal{U}_\eta = \bigcup_{0 \leq t \leq \eta} \widehat{X}_t(\widehat{T}).$$

Consider the “straightening diffeomorphism” $H_\eta: \mathcal{U}_\eta \rightarrow [0, 1] \times \mathbb{T}^2$ given by

$$(4-1) \quad H_\eta(\widehat{X}_t(p)) = \left(\frac{t}{\eta}, \theta_T(p) \right), \quad p \in \widehat{T}.$$

For fixed η we will denote by \widehat{W}_η^σ and \widehat{E}_η^σ the corresponding foliations and invariant bundles for \widehat{X} ($\sigma = cs, cu, ss, uu$).

We also denote by \mathcal{F}^{uu} and \mathcal{F}^{ss} the one-dimensional foliations of $[0, 1] \times \mathbb{T}^2$ which in $\{t\} \times \mathbb{T}^2$ coincide with the foliations $\{t\} \times F^u$ and $\{t\} \times F^s$, respectively.

Lemma 4.5 *The diffeomorphism H_η has the following properties:*

- $H_\eta(\widehat{W}_\eta^{cs} \cap \mathcal{U}_\eta) = [0, 1] \times F^s := \mathcal{F}^{cs}$.
- $H_\eta(\widehat{W}_\eta^{cu} \cap \mathcal{U}_\eta) = [0, 1] \times F^u := \mathcal{F}^{cu}$.
- $DH_\eta(\widehat{E}_\eta^{ss})$ converges to the tangent bundle of the foliation \mathcal{F}^{ss} as $\eta \rightarrow \infty$.
- $DH_\eta(\widehat{E}_\eta^{uu})$ converges to the tangent bundle of the foliation \mathcal{F}^{uu} as $\eta \rightarrow \infty$.

It is important to note that the above convergence is with respect to the standard metric in $[0, 1] \times \mathbb{T}^2$ and not with respect to the pushforward metric from the manifold via H_η .

Proof Because the differential of H_η maps the vector field \widehat{X} to the vector field $\frac{1}{\eta} \frac{\partial}{\partial t}$, the first two properties follow. Note that the component of \widehat{E}_η^σ along \widehat{X} is uniformly bounded (for $\sigma = ss, uu$). Therefore, contraction by a factor $\frac{1}{\eta}$ implies the posited limit behavior in the latter properties. \square

4.5 A diffeomorphism in a flow box which preserves transversalities

Assume that there exists a smooth path $\{\varphi_s\}_{s \in [0,1]}$ of diffeomorphisms of \mathbb{T}^2 such that

- $\varphi_s = \text{id}$ for s in neighborhoods of 0 and 1,
- the closed path $s \mapsto \varphi_s$ is not homotopically trivial in $\text{Diff}(\mathbb{T}^2)$,
- for every $s \in [0, 1]$,

$$\varphi_s(F^u) \pitchfork F^s.$$

We use the coordinate chart $H_\eta: \mathcal{U}_\eta \rightarrow [0, 1] \times \mathbb{T}^2$ to define a diffeomorphism $\mathcal{G}_\eta: \mathcal{U}_\eta \rightarrow \mathcal{U}_\eta$ by

$$(s, x, y) \mapsto (s, \varphi_s(x, y)).$$

The following lemma is immediate from our choice of $\{\varphi_s\}_{s \in [0,1]}$:

Lemma 4.6 *The diffeomorphism \mathcal{G}_η has the following properties:*

- $\mathcal{G}_\eta(H_\eta^{-1}(\mathcal{F}^{uu}))$ is transverse to $\widehat{W}_{\widehat{X}}^{cs} = H_\eta^{-1}(\mathcal{F}^{cs})$.
- $\mathcal{G}_\eta(\widehat{W}_{\widehat{X}}^{cu}) = \mathcal{G}_\eta(H_\eta^{-1}(\mathcal{F}^{cu}))$ is transverse to $H_\eta^{-1}(\mathcal{F}^{ss})$.

Let $h_\eta: \widehat{M}_\eta \rightarrow \widehat{M}_\eta$ be the diffeomorphism defined as

$$(4-2) \quad h_\eta(x) = \begin{cases} x & \text{if } x \notin \mathcal{U}_\eta, \\ \mathcal{G}_\eta(x) & \text{if } x \in \mathcal{U}_\eta. \end{cases}$$

If γ denotes the element of $\pi_1(T)$ associated to the loop $\{\theta_T^{-1} \circ \varphi_t \circ \theta_T(x)\}_t$, then, by construction, h_η is a diffeomorphism that belongs to $\tau(T, \gamma)$, ie a Dehn twist along γ (see Section 4.1).

Proposition 4.7 *There exists $\eta_0 > 0$ such that, for all $\eta \geq \eta_0$,*

- $Dh_\eta(E_{\widehat{X}}^{uu})$ is transverse to $E_{\widehat{X}}^{cs}$,
- $Dh_\eta(E_{\widehat{X}}^{cu})$ is transverse to $E_{\widehat{X}}^{ss}$.

Proof This follows by combining Lemmas 4.6 4.5 and the fact that outside \mathcal{U}_η the diffeomorphism h_η is the identity. \square

As a consequence, using Proposition 4.4, we have the following:

Corollary 4.8 *Given X an Anosov flow on M transverse to an embedded torus T and $\gamma \in G(T, X)$, there exists a smooth function $\rho_T: M \rightarrow (0, 1]$ and a diffeomorphism $h_\gamma: M \rightarrow M$ in the mapping class $\tau(T, \gamma)$ such that if f_T is the time-one map of the flow generated by $\rho_T X$, then f_T is h_γ -transverse to itself.*

Remark 4.9 If there is a volume form ω on T such that φ_s preserves the form $(\theta_T)_*(\omega)$ for every s , then h_η preserves the volume form $\omega \wedge dX$.

4.6 Proof of Theorem 1.3

Let X be an Anosov flow on M and let T_1, \dots, T_ℓ be tori (not necessarily disjoint or distinct) transverse to X . Choose $\gamma_i \in G(T_i, X)$.

Using Corollaries 4.2 and 4.8, one obtains diffeomorphisms h_i and g_i of M such that if f_i denotes the time-one map of the flow generated by $\rho_{T_i} X$, then we have

$$f_1 \rightsquigarrow_{h_1} f_1 \rightsquigarrow_{g_1} f_2 \rightsquigarrow_{h_2} f_2 \rightsquigarrow_{g_2} \cdots \rightsquigarrow_{g_{\ell-1}} f_\ell \rightsquigarrow_{h_\ell} f_\ell \rightsquigarrow_{g_\ell} f_1$$

with g_i isotopic to the identity and $h_i \in \tau(T_i, \gamma_i)$.

Now, according to Proposition 2.4 (and Lemma 2.3), for every large m the diffeomorphism

$$F = f_1^m \circ g_\ell \circ f_\ell^m \circ h_\ell \circ \cdots \circ h_2 \circ f_2^m \circ g_1 \circ f_1^m \circ h_1$$

is absolutely partially hyperbolic. Clearly, the homotopy class of F is the composition of $\tau(T_\ell, \gamma_\ell) \circ \cdots \circ \tau(T_1, \gamma_1)$. □

4.7 Mapping class groups of 3-manifolds

In order to proceed with the proof of Corollary 1.4, we need to briefly recall the description of the mapping class group $\mathcal{M}(M)$ of a Haken manifold M . This description relies on Johannson’s characteristic submanifold theory, which we briefly summarize. For full generality and full details we refer to [29, Chapters VI and IX]; see also [32, Section 4; 21, Chapter 4].

Let M be a closed ($\partial M = \emptyset$) Haken 3-manifold. Then the *characteristic submanifold* $\Sigma \subset M$ is a codimension-0 submanifold of M which consists of all Seifert-fibered pieces of the JSJ decomposition of M . A torus which separates two Seifert-fibered

pieces of the JSJ decomposition yields two torus boundary components of Σ such that the number of connected components of Σ is precisely the number of Seifert-fibered pieces in the JSJ decomposition. Johannson's classification theorem asserts that every diffeomorphism of M is homotopic to a homeomorphism which preserves the characteristic submanifold Σ . Consider the subgroup $\mathcal{M}(M, \Sigma)$ of $\mathcal{M}(M)$ given by diffeomorphisms whose restriction to the closure of the complement of Σ is isotopic to the identity. By work of Johannson, $\mathcal{M}(M, \Sigma)$ has finite index in $\mathcal{M}(M)$. (In the proof of Corollary 1.4 below, the fact that $\mathcal{M}(M, \Sigma)$ has finite index will be immediate from Johannson's classification theorem.)

Restricting to connected components Σ_i for $i = 1, \dots, n$ of Σ yields an exact sequence (see [32, Section 4])

$$(4-3) \quad 1 \rightarrow K \rightarrow \mathcal{M}(M, \Sigma) \rightarrow \prod_{i=1}^n \overline{\mathcal{M}}(\Sigma_i, \partial\Sigma_i) \rightarrow 1,$$

where K consists of mapping classes which are represented by homeomorphisms $h: M \rightarrow M$ which are the identity outside a small product neighborhood of $\partial\Sigma$ and $\overline{\mathcal{M}}(\Sigma_i, \partial\Sigma_i)$ is the subgroup of the full mapping class group $\mathcal{M}(\Sigma_i, \partial\Sigma_i)$ given by homeomorphisms $h: \Sigma_i \rightarrow \Sigma_i$ whose restriction to $\partial\Sigma_i$ is isotopic to the identity.

Proposition 4.10 *Let M be a Haken manifold admitting an Anosov flow X which is transverse to the boundary $\partial\Sigma$ of the characteristic submanifold $\Sigma = \bigcup_{i=1}^n \Sigma_i$. Assume that for each torus boundary component T of Σ the group of permitted Dehn twists $G(T, X)$ is maximal, ie isomorphic to \mathbb{Z}^2 . Also assume that the mapping class groups of the Seifert-fibered pieces $\overline{\mathcal{M}}(\Sigma_i, \partial\Sigma_i)$ for $i = 1, \dots, n$ are finite. Then the mapping class group $\mathcal{M}(M)$ has a finite-index subgroup K such that each mapping class in K can be represented by an absolutely partially hyperbolic diffeomorphism.*

Proof By the above discussion and the assumption on $\overline{\mathcal{M}}(\Sigma_i, \partial\Sigma_i)$ for $i = 1, \dots, n$, the subgroup K from (4-3) has finite index in $\mathcal{M}(M)$. Each element of K is represented by a Dehn twist along $\partial\Sigma$. Hence, by the assumption on $G(T, X)$, we have $K \subset \mathcal{T}(X)$ (recall that $\mathcal{T}(X)$ is the group generated by permitted Dehn twists). Hence each mapping class from K is represented by a partially hyperbolic diffeomorphism according to Theorem 1.3. \square

In order to demonstrate examples to which Proposition 4.10 applies, we need to have Seifert-fibered manifolds Σ_i with finite $\overline{\mathcal{M}}(\Sigma_i, \partial\Sigma_i)$. Thus we need to recall

Johannson’s description of the mapping class group of Seifert-fibered manifolds with boundary [29, Section 25].

Let Σ be a Seifert-fibered manifold with a nonempty boundary $\partial\Sigma$. Consider the mapping class group $\mathcal{M}(\Sigma, \partial\Sigma)$ of self-homeomorphisms $h: (\Sigma, \partial\Sigma) \rightarrow (\Sigma, \partial\Sigma)$ modulo boundary-preserving isotopies (h is allowed to permute boundary components). If one excludes six exceptional manifolds [29, 5.1.1–5.1.6] (which we will not encounter here) then the homeomorphism h is isotopic to a fiber-preserving homeomorphism [29, Corollary 5.9]. It follows [29, Proposition 25.3] that the mapping class group fits into a split short exact sequence

$$(4-4) \quad 1 \rightarrow \mathcal{M}^0(\Sigma, \partial\Sigma) \rightarrow \mathcal{M}(\Sigma, \partial\Sigma) \rightarrow \mathcal{M}(B, \partial B) \rightarrow 1,$$

where $\mathcal{M}^0(\Sigma, \partial\Sigma)$ is the subgroup of the mapping class group given by homeomorphisms which preserve each fiber (ie homeomorphisms which fiber over the identity) and $\mathcal{M}(B, \partial B)$ is the mapping class group of the underlying orbifold B . (In the setup to which we are about to specify in order to obtain Corollary 1.4, the manifold Σ will be circle-fibered and, hence, $\mathcal{M}(B, \partial B)$ will simply be the mapping class group of a surface B .) Further, $\mathcal{M}^0(\Sigma, \partial\Sigma)$ is isomorphic to $H_1(B, \partial B)$ (relative to first homology) and is generated by certain vertical Dehn twists [29, Lemma 25.2].


Recall that, by Johannson’s work, elements of $\mathcal{M}(\Sigma, \partial\Sigma)$ are represented by fiber-preserving homeomorphisms. Further if $h_1, h_2: \Sigma \rightarrow \Sigma$ are isotopic fiber-preserving homeomorphisms then, by work of Waldhausen [41] (see discussion on page 85), h_1 is isotopic to h_2 via a path of fiber-preserving homeomorphisms. It follows that the short exact sequence (4-4) restricts to a short exact sequence

$$(4-5) \quad 1 \rightarrow \overline{\mathcal{M}}^0(\Sigma, \partial\Sigma) \rightarrow \overline{\mathcal{M}}(\Sigma, \partial\Sigma) \rightarrow \overline{\mathcal{M}}(B, \partial B) \rightarrow 1,$$

where a bar indicates the subgroup (of the corresponding mapping class group) induced by homeomorphisms whose restriction to the boundary is isotopic to the identity.

4.8 Proof of Corollary 1.4

Denote by Σ the total space of the (unique) nonoriented circle bundle $S^1 \rightarrow \Sigma \xrightarrow{P} B$, where B is the projective plane $\mathbb{R}P^2$ with two disks removed (see Figure 2). Then the manifold Σ is orientable and has two torus boundary components. Bonatti and Langevin exhibited a self-gluing of Σ (via an orientation-reversing diffeomorphism of the 2–torus) which yields a closed manifold M which supports an Anosov flow X transverse to the gluing torus [10]. The closed manifold M is not Seifert-fibered, hence

Figure 2: The surface B .

the characteristic submanifold of M is the complement of a product neighborhood of the gluing torus and, therefore, can be identified with Σ . Note that, for M , the subgroup K appearing in (4-3) is the subgroup of Dehn twists along the gluing torus and, hence, is isomorphic to \mathbb{Z}^2 . Further, by combining the description of invariant foliations near the gluing torus provided by [10] with the description of permitted Dehn twists [14] we have that all Dehn twists are permitted. Hence Proposition 4.10 applies and, according to (4-5), in order to complete the proof of Corollary 1.4 in the case when $n = 1$ it only remains to show that the groups $\overline{\mathcal{M}}^0(\Sigma, \partial\Sigma)$ and $\overline{\mathcal{M}}(B, \partial B)$ are finite. In fact, the form of the intersection of the foliations with the tori allow to see that one can make the Dehn twists in a volume-preserving way (see [8, Lemma 3.8]).

The mapping class group $\overline{\mathcal{M}}(B, \partial B)$ can be easily seen to be $\mathbb{Z}/2\mathbb{Z}$. By Lemma 25.2 of [29], we have the isomorphism $\mathcal{M}^0(\Sigma, \partial\Sigma) \simeq H_1(B, \partial B)$. The group $H_1(B, \partial B)$ is the first homology of $\mathbb{R}P^2$ with two points identified. Hence $\mathcal{M}^0(\Sigma, \partial\Sigma) \simeq H_1(B, \partial B) \simeq \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$. Further, the proof of Case 2 of Lemma 25.2 of [29] gives an explicit description of generators of $\mathcal{M}^0(\Sigma, \partial\Sigma)$ as follows. Let α be a simple curve which connects the boundary components of B and let h be a simple curve which connects a boundary component to itself as indicated in Figure 2. Let $A_\alpha = p^{-1}\alpha$ and $A_h = p^{-1}h$ be the corresponding annuli. Then the Dehn twists along A_α and A_h are the generators of $\mathcal{M}^0(\Sigma, \partial\Sigma)$, the first one being the infinite-order generator and the second one being the generator of order two. Clearly the Dehn twist along A_α restricts to a Dehn twist on each torus boundary component and, hence, is not isotopic to the identity on the boundary. However, the restriction of Dehn twist along A_h to the boundary $\partial\Sigma$ can be seen to be isotopic to the identity (two Dehn twists on the boundary component given by h are inverses of each other). We conclude that $\overline{\mathcal{M}}^0(\Sigma, \partial\Sigma)$ is $\mathbb{Z}/2\mathbb{Z}$ and that $K \simeq \mathbb{Z}^2$ indeed has finite index in $\mathcal{M}(M)$.

To address the case $n > 1$, consider finite cyclic covers $M_n \rightarrow M$ obtained by gluing n copies of Σ using the Bonatti–Langevin gluing. Then the lifted Anosov flow on M_n is transverse to each of the gluing tori. The characteristic submanifold of M_n is the disjoint union of n copies of Σ . Then the same considerations as above apply to show that $K \simeq \mathbb{Z}^{2n}$ has finite index in $\mathcal{M}(M_n)$. Further, because the group of deck transformations $\mathbb{Z}/n\mathbb{Z}$ of the covering $M_n \rightarrow M$ commutes with the Anosov flow, one can, in fact, realize a larger subgroup $K \oplus \mathbb{Z}/n\mathbb{Z}$ by partially hyperbolic diffeomorphisms by following the arguments for the proof of Theorem 1.3. More specifically, one has to utilize the fact that if h denotes a deck transformation of the covering $M_n \rightarrow M$ then the time-one map of the Anosov flow is h -transverse to itself. As this is only needed to go from K to $K \oplus \mathbb{Z}/n\mathbb{Z}$, we leave the details to the reader.

5 Dynamical incoherence

In this section, we consider the examples built in Theorem 1.1 on the unit tangent bundle T^1S of an oriented higher-genus surface S which are isotopic to the projectivizations of pseudo-Anosov diffeomorphisms of S .

Consider a hyperbolic surface (S, μ) and an element φ of the mapping class group $\mathcal{M}(S)$. We denote by $\{\mathcal{G}^t\}_{t \in \mathbb{R}}$ the geodesic flow for the hyperbolic metric μ and by Z the vector field which generates it. Let $E_Z^{ss} \oplus \mathbb{R}Z \oplus E_Z^{uu}$ be the hyperbolic splitting for the geodesic flow.

According to Theorem 1.1 and Remarks 2.6 and 3.2, there exists a diffeomorphism $f: T^1S \rightarrow T^1S$ such that

- f belongs to the mapping class $\iota(\varphi)$;
- for any $t \geq 0$ the diffeomorphism $f_t = \mathcal{G}^t \circ f \circ \mathcal{G}^t$ is absolutely partially hyperbolic and volume-preserving; we denote by $E_t^{ss} \oplus E_t^c \oplus E_t^{uu}$ the corresponding partially hyperbolic splitting;
- the subbundles E_t^{ss} , E_t^c and E_t^{uu} converge uniformly to E_Z^{ss} , $\mathbb{R}Z$ and E_Z^{uu} , respectively, as $t \rightarrow +\infty$.

The goal of this section is to prove the following result (which is nonvacuous according to the above discussion):

Theorem 5.1 *Consider a hyperbolic surface (S, μ) and a pseudo-Anosov mapping class $\varphi \in \mathcal{M}(S)$. Let Z be the vector field which generates the geodesic flow on T^1S .*

Then there exists $\eta > 0$ such that, if $f: T^1S \rightarrow T^1S$ is a diffeomorphism satisfying that

- f is partially hyperbolic,
- f belongs to the mapping class $\iota(\varphi)$,
- the subbundles of the partially hyperbolic splitting $E^{ss} \oplus E^c \oplus E^{uu}$ of f are η -close to the bundles of the hyperbolic splitting $E_Z^{ss} \oplus \mathbb{R}Z \oplus E_Z^{uu}$ of the geodesic flow,

then f is dynamically incoherent.

In fact, we will prove a slightly more general result. To state the result we need to recall the Nielsen–Thurston classification of mapping classes of S [40; 28]. According to this classification, each homeomorphism of a compact oriented surface S of negative Euler characteristic with (possibly empty) boundary ∂S is isotopic to one of the following “normal forms” $h: S \rightarrow S$:

- h is *periodic*, that is, $h^n = \text{id}_S$ for some $n \geq 1$.
- h is *pseudo-Anosov*.
- h is *reducible*. Namely, there exists a finite collection of simple closed curves whose union is invariant under h . This collection partitions S into a finite collection of subsurfaces each of which is invariant under h^n for some $n \geq 1$. Moreover, the restriction of h^n to each of the subsurfaces is either periodic or pseudo-Anosov.

Addendum 5.2 *Theorem 5.1 also holds if instead of assuming that mapping class φ is pseudo-Anosov we assume that φ is a reducible mapping class with at least one pseudo-Anosov component.*

From now on we fix the hyperbolic surface (S, μ) , its geodesic flow \mathcal{G}^t and the generating vector field Z .

5.1 Shadowing property of the geodesic flow \mathcal{G}^t

Let us first state the standard shadowing lemma for Anosov flows (see Theorem 18.1.6 of [30]). Recall that a δ -*pseudorbit* for a flow generated by a vector field X is a curve $c: \mathbb{R} \rightarrow M$ such that $\|c'(t) - X(c(t))\| \leq \delta$ for all $t \in \mathbb{R}$. We say that a curve $c: \mathbb{R} \rightarrow M$ is ε -*shadowed* by an orbit of X if there exists $x \in M$ and a reparametrization $s: \mathbb{R} \rightarrow \mathbb{R}$ such that $|s'(t) - 1| \leq \varepsilon$ for all t and such that $d(c(s(t)), X_t(x)) \leq \varepsilon$ for all $t \in \mathbb{R}$.

Theorem 5.3 *Let X be an Anosov flow on a closed manifold M . Then, for every $\varepsilon > 0$, there exists $\delta > 0$ such that every δ -pseudorbit is ε -shadowed by an orbit of X . Moreover, if ε is sufficiently small, then the shadowing orbit is unique.*

In our setting, we also obtain shadowing injectivity as follows. We can parametrize center leaves as $(c, \vec{c}): \mathbb{R} \rightarrow T^1S$ and the notation is to indicate that the projection into S is the curve $c: \mathbb{R} \rightarrow S$.

Theorem 5.4 *Let Z be the vector field which generates the geodesic flow \mathcal{G}^t of a hyperbolic surface S . Then there is $\varepsilon_0 > 0$ such that for every $\varepsilon \in (0, \varepsilon_0)$ there is $\delta > 0$ with the following property:*

- *for any dynamically coherent partially hyperbolic diffeomorphism $f: T^1S \rightarrow T^1S$ whose subbundles E_f^s, E_f^c and E_f^u of its partially hyperbolic splitting are δ -close to the corresponding ones of Z , and let X_f^c be a vector field orienting E_f^c such that X_f^c is δ -close to Z ,*
- *and any pair of different center leaves $(c_1, \vec{c}_1), (c_2, \vec{c}_2): \mathbb{R} \rightarrow T^1S$ tangent to X_f^c ,*

the orbits of the geodesic flow \mathcal{G}^t that ε -shadow (c_1, \vec{c}_1) and (c_2, \vec{c}_2) are different.

The argument will be carried out on the cover $T^1\mathbb{D}^2$ and we need to introduce some notation first. The Poincaré disk \mathbb{D}^2 covers the hyperbolic surface (S, μ) so that the covering map is a local isometry. Let $d_{\mathbb{D}^2}$ denote the metric on \mathbb{D}^2 induced by the hyperbolic Riemannian metric. We equip $T^1\mathbb{D}^2$ with the product Riemannian metric (which is π_1S -equivariant) and denote by $d_{T^1\mathbb{D}^2}$ the induced metric.

Proof To prove the theorem, we lift all the structures to $T^1\mathbb{D}^2$. We denote the lift of f by $\tilde{f}: T^1\mathbb{D}^2 \rightarrow T^1\mathbb{D}^2$. First note that it is sufficient to establish the “shadowing injectivity property” on $T^1\mathbb{D}^2$. (Indeed, if a geodesic $(\gamma, \vec{\gamma}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$ shadows a center leaf $(c, \vec{c}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$, and its image under a deck transformation $g(\gamma, \vec{\gamma})$ shadows another center leaf (c', \vec{c}') , then $(\gamma, \vec{\gamma})$ shadows both (c, \vec{c}) and $g^{-1}(c', \vec{c}')$. Hence, by shadowing injectivity on the cover we would have $(c', \vec{c}') = g(c, \vec{c})$ and, hence, the posited injectivity on T^1S .)

Note that according to our assumptions on subbundles of f the size of local product structure neighborhoods is uniform in ε ; that is, the size stays fixed when we let $\varepsilon \searrow 0$

(and $\delta \searrow 0$). Hence we assume that ε is much smaller than the size of local product structure neighborhoods.

Each center leaf of \tilde{f} has the form $(c, \vec{c}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$, where $c: \mathbb{R} \rightarrow \mathbb{D}^2$. Further, by dynamical coherence, each center leaf (c, \vec{c}) is contained in a unique center-unstable leaf $W^{cu}(c, \vec{c})$ which is uniformly transverse (ie with angle bounded away from 0) to the circle fibers of π . It follows that $\pi: W^{cu}(c, \vec{c}) \rightarrow \mathbb{D}^2$ is a covering map and, hence, is a diffeomorphism. Further, because π is a Riemannian submersion and the angle between $E^s \oplus E^{uu}$ and the circle fibers is uniformly bounded away from 0 (uniformly in ε), we also have that $\|D((\pi|_{W^{cu}(x)})^{-1})\|$ is uniformly bounded and, hence,

$$(5-1) \quad d_{W^{cu}(x)}(x_1, x_2) \leq C d_{\mathbb{D}^2}(\pi(x_1), \pi(x_2)),$$

where $x_2 \in W^{cu}(x_1)$, $d_{W^{cu}(x)}$ is the metric induced by the restriction of the Riemannian metric to $W^{cu}(x)$ and C is a constant which does not depend on ε .

We will argue by contradiction. Assume that a geodesic $(\gamma, \vec{\gamma}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$ ε -shadows two distinct center leaves $(c_1, \vec{c}_1): \mathbb{R} \rightarrow T^1\mathbb{D}^2$ and $(c_2, \vec{c}_2): \mathbb{R} \rightarrow T^1\mathbb{D}^2$. Then, because π does not increase distance and using the triangle inequality, we have

$$(5-2) \quad d_{\mathbb{D}^2}^{\mathcal{H}}(c_1, c_2) \leq d_{T^1\mathbb{D}^2}^{\mathcal{H}}((c_1, \vec{c}_1), (c_2, \vec{c}_2)) \leq 2\varepsilon,$$

where superscript \mathcal{H} indicates the Hausdorff distance with respect to the corresponding metric.

Without loss of generality we can assume that (c_1, \vec{c}_1) and (c_2, \vec{c}_2) belong to the same center-unstable leaf. To see this consider the “heteroclinic center leaf” $(c_3, \vec{c}_3) = W^{u,loc}(c_1, \vec{c}_1) \cap W^{s,loc}(c_2, \vec{c}_2)$. Clearly this leaf is also $K\varepsilon$ -shadowed by γ , where K is a constant associated with the local product structure and is independent of ε . Also c_3 is different either from c_1 or from c_2 (or both). We assume that c_1 and c_3 are distinct (otherwise the arguments are the same with the time direction reversed). Hence, we can conclude that, by choosing ε smaller and by replacing c_2 with c_3 , we indeed can assume that the center leaves (c_1, \vec{c}_1) and (c_2, \vec{c}_2) are contained in the same center-unstable leaf.

For any $n \geq 1$ consider center leaves $\tilde{f}^n(c_1, \vec{c}_1)$ and $\tilde{f}^n(c_2, \vec{c}_2)$. Let γ_1 and γ_2 be the geodesics which shadow them, respectively. By (5-2) and because $\|D\tilde{f}\|$ is uniformly bounded, we see that γ_1 and γ_2 are a finite distance apart (as oriented geodesics). Recall that in \mathbb{D}^2 geodesics either coincide or diverge exponentially in positive or

negative time. Hence $\gamma_1 = \gamma_2$. By the same token as before (5-2), we also have

$$(5-3) \quad d_{\mathbb{D}^2}^{\mathcal{H}}(\pi \tilde{f}^n(c_1, \vec{c}_1), \pi \tilde{f}^n(c_2, \vec{c}_2)) \leq d_{T^1\mathbb{D}^2}^{\mathcal{H}}(\tilde{f}^n(c_1, \vec{c}_1), \tilde{f}^n(c_2, \vec{c}_2)) \leq 2\varepsilon$$

Now, for each center leaf, consider its neighborhood in the center-unstable leaf,

$$W^{cu,loc}(c, \vec{c}) = \bigcup_{x \in (c, \vec{c})} W^{u,loc}(x) \subset W^{cu}(c, \vec{c}),$$

where the size of $W^{u,loc}$ is the size of local product structure neighborhoods and hence is much bigger than ε . For all $n \geq 0$, using (5-1) and (5-3), we have

$$\begin{aligned} d_{W^{cu}(\tilde{f}^n(c_1, \vec{c}_1))}^{\mathcal{H}}(\tilde{f}^n(c_1, \vec{c}_1), \tilde{f}^n(c_2, \vec{c}_2)) &\leq C d_{\mathbb{D}^2}^{\mathcal{H}}(\pi \tilde{f}^n(c_1, \vec{c}_1), \pi \tilde{f}^n(c_2, \vec{c}_2)) \\ &\leq C d_{T^1\mathbb{D}^2}^{\mathcal{H}}(\tilde{f}^n(c_1, \vec{c}_1), \tilde{f}^n(c_2, \vec{c}_2)) \leq 2C\varepsilon. \end{aligned}$$

Recall that C is independent of ε . Hence, by the above inequality, we can pick ε sufficiently small that $\tilde{f}^n(c_2, \vec{c}_2)$ belongs to $W^{u,loc}(\tilde{f}^n(c_1, \vec{c}_1))$ for all $n \geq 0$. Now pick an $x \in (c_1, \vec{c}_1)$ and let $y \in W^{u,loc}(x) \cap (c_2, \vec{c}_2)$. Then $\tilde{f}^n(x)$ and $\tilde{f}^n(y)$ diverge exponentially quickly along the unstable leaf. Pick the smallest $n \geq 1$ such that $\tilde{f}^n(y) \notin W^{u,loc}(\tilde{f}^n(x))$. On the other hand, $\tilde{f}^n(y) \in \tilde{f}^n(c_2, \vec{c}_2) \subset W^{u,loc}(c_1, \vec{c}_1)$. Therefore there exists a point $z \in (c_1, \vec{c}_1)$ such that $\tilde{f}^n(y) \in W^{u,loc}(\tilde{f}^n(z))$. We conclude that the unstable leaf $W^u(x)$ intersects the center leaf (c_1, \vec{c}_1) in two distinct points x and z . But such a configuration is impossible inside $W^{cu}(x)$ (which is diffeomorphic to \mathbb{D}^2) by the standard Poincaré–Bendixson argument (see eg [30, Section 14.1]). \square

The following global shadowing property of (general) Anosov flows is a straightforward consequence of [30, Theorem 18.1.7]:

Theorem 5.5 *Let X be an Anosov flow on a compact manifold M . Then, for any $\varepsilon > 0$, there is $\delta > 0$ with the following property: consider any C^0 -foliation \mathcal{F} directed by a vector field Y such that, for every $x \in M$,*

$$\|Y(x) - X(x)\| < \delta.$$

Then there is $h: M \rightarrow M$ with the following properties:

- *For any $x \in M$ one has $d(h(x), x) < \varepsilon$.*
- *For any leaf L of \mathcal{F} , the image $h(L)$ is exactly an orbit of X .*
- *h is continuous and onto; in particular, for any orbit γ of X there is a leaf L with $h(L) = \gamma$.*

Putting together Theorems 5.5 and 5.4 we immediately obtain the following:

Corollary 5.6 *For every $\varepsilon > 0$ there exists $\delta > 0$ such that if $f: T^1S \rightarrow T^1S$ is a dynamically coherent partially hyperbolic diffeomorphism whose splitting is δ -close to the one of Z , then every center leaf $c: \mathbb{R} \rightarrow T^1S$ tangent to X_f^c parametrized by arc-length is ε -shadowed by a unique orbit of Z . Moreover, for every orbit γ of Z , there is a unique center curve which is ε -shadowed by γ .*

5.2 Dynamics at infinity

We compactify the Poincaré disk \mathbb{D}^2 by adding the “the circle at infinity” $\partial\mathbb{D}^2$ with the topology induced by the Euclidean topology of $\mathbb{R}^2 \supset \mathbb{D}^2 \cup \partial\mathbb{D}^2$. Recall that $\partial\mathbb{D}^2$ can be viewed as the space of asymptotic classes of geodesics in \mathbb{D}^2 . Further, each unit speed geodesic γ in \mathbb{D}^2 is uniquely determined by an ordered pair of *boundary points* $(\gamma_-, \gamma_+) \in \partial\mathbb{D}^2 \times \partial\mathbb{D}^2 \setminus \Delta$, where

$$\gamma_- = \lim_{t \rightarrow -\infty} \gamma(t), \quad \gamma_+ = \lim_{t \rightarrow +\infty} \gamma(t).$$

In the same way, any curve $c: \mathbb{R} \rightarrow \mathbb{D}^2$ which stays a bounded distance away from a geodesic determines a pair of boundary points $(c_-, c_+) \in \partial\mathbb{D}^2 \times \partial\mathbb{D}^2 \setminus \Delta$.

Now let $\Psi: S \rightarrow S$ be a homeomorphism and let $\tilde{\Psi}: \mathbb{D}^2 \rightarrow \mathbb{D}^2$ be a lift. Then the action of $\tilde{\Psi}$ on geodesic rays induces a homeomorphism $\partial\tilde{\Psi}: \partial\mathbb{D}^2 \rightarrow \partial\mathbb{D}^2$ which depends only on the isotopy class of Ψ (but does depend on the choice of the lift). Namely, if $\Psi' \simeq \Psi$ and $\tilde{\Psi}' \simeq \tilde{\Psi}$ is the lift of the isotopy, then $\partial\tilde{\Psi}' = \partial\tilde{\Psi}$. Further, if Ψ is pseudo-Anosov, then, for any lift $\tilde{\Psi}$, the boundary homeomorphism $\partial\tilde{\Psi}$ has finitely many fixed points on $\partial\mathbb{D}^2$ alternatively expanding and contracting (see eg [23, Theorem 5.5]). Moreover, the lift of the unstable geodesic lamination \tilde{L}^u of the mapping class of Ψ contains the geodesics joining consecutive expanding fixed points.

In the case when Ψ is reducible, the structure of the fixed points at infinity can be more complicated, but the following proposition is still true:

Proposition 5.7 [23, Sections 5 and 6; 33, Section 9] *Let $\Psi: S \rightarrow S$ be a pseudo-Anosov homeomorphism or a reducible homeomorphism with a pseudo-Anosov component. Then there is a positive iterate $\partial\tilde{\Psi}^n$ which has two distinct expanding fixed points γ_- and γ_+ , and the geodesic γ which is determined by these points belongs to the lift of the unstable geodesic lamination \tilde{L}^u . Further, Ψ^n can be isotoped to Φ so that the corresponding lift $\tilde{\Phi} \simeq \tilde{\Psi}^n$ preserves the unstable geodesic lamination and hence preserves γ .*

Remark 5.8 There is a discrepancy between the terminology in hyperbolic dynamics and the terminology for laminations (and singular foliations) of pseudo-Anosov diffeomorphisms. Namely, the *stable geodesic lamination* is the one which *expands* sufficiently large segments of its geodesics. We refer to [23; 20] for further background on geodesic laminations and pseudo-Anosov theory.

5.3 Dynamics at infinity and the center leaves

Consider a dynamically coherent partially hyperbolic diffeomorphism $f: T^1S \rightarrow T^1S$ which satisfies the assumptions of Theorem 5.4. Assume that there is a diffeomorphism $\Phi: S \rightarrow S$ such that f is isotopic to $D\Phi: T^1S \rightarrow T^1S$. Consider the π_1S -covers $\mathbb{D}^2 \rightarrow S$ and $T^1\mathbb{D}^2 \rightarrow T^1S$ and lifts $\tilde{\Phi}: \mathbb{D}^2 \rightarrow \mathbb{D}^2$ and $\tilde{f}: T^1\mathbb{D}^2 \rightarrow T^1\mathbb{D}^2$ such that the isotopy $f \simeq D\Phi$ lifts to an isotopy $\tilde{f} \simeq D\tilde{\Phi}$. As $\tilde{f} \simeq D\tilde{\Phi}$ is an equivariant isotopy, we have

$$(5-4) \quad d_{T^1\mathbb{D}^2}(\tilde{f}, D\tilde{\Phi}) < C$$

and, because the bundle map $\pi: T^1\mathbb{D}^2 \rightarrow \mathbb{D}^2$ is a Riemannian submersion, we also have

$$(5-5) \quad d_{\mathbb{D}^2}(\pi \circ \tilde{f}, \tilde{\Phi}) < C.$$

Consider the center leaves for \tilde{f} , ie connected components of preimages of center leaves for f under the covering map. Recall that each center leaf has the form $(c, \bar{c}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$, where $c: \mathbb{R} \rightarrow \mathbb{D}^2$. We slightly abuse terminology by referring to the underlying curve $c: \mathbb{R} \rightarrow \mathbb{D}^2$ as a center leaf as well. This is harmless because $c: \mathbb{R} \rightarrow \mathbb{D}^2$ uniquely determines the center leaf. By Theorem 5.4, each center leaf c is shadowed by a unique geodesic γ , and hence each center leaf c is uniquely determined by a pair of points on the boundary and vice versa. According to the following claim the dynamics of \tilde{f} on the space of center leaves is uniquely determined by the dynamics on the boundary.

Claim 5.9 For a center leaf (c, \bar{c}) of \tilde{f} , let $(c_1, \bar{c}_1) = \tilde{f}(c, \bar{c})$. Then

$$(c_{1-}, c_{1+}) = (\partial\tilde{\Phi}(c_-), \partial\tilde{\Phi}(c_+)),$$

where (c_-, c_+) and (c_{1-}, c_{1+}) are the boundary points of c and c_1 , respectively.

Proof Let γ be the geodesic which shadows c . Then γ and c have the same boundary points (c_-, c_+) . Applying \tilde{f} , we have that (c_1, \bar{c}_1) is of bounded distance away

from $\tilde{f}(\gamma, \gamma')$. By (5-4), $\tilde{f}(\gamma, \gamma')$ is of bounded distance away from $D\tilde{\Phi}(\gamma, \gamma') = (\tilde{\Phi}(\gamma), \tilde{\Phi}(\gamma'))$. But $\tilde{\Phi}(\gamma)$ has boundary points $(\partial\tilde{\Phi}(c_-), \partial\tilde{\Phi}(c_+))$ by the definition of $\partial\tilde{\Phi}$. □

5.4 Proof of Theorem 5.1 and the addendum

Let f be a partially hyperbolic diffeomorphism and let $\varphi \in \mathcal{M}(S)$ be the mapping class as in Theorem 5.1 or the addendum with $f \in \iota(\varphi)$. First we pass to positive iterates of f and φ , so that Proposition 5.7 applies and yields a representative Φ of φ such that the lift $\tilde{\Phi}$ preserves a geodesic γ . Further, γ belongs to the unstable lamination and, hence, is *coarsely contracting* (see Remark 5.8), ie for all sufficiently long segments $[\gamma(a), \gamma(b)] \subset \gamma(\mathbb{R})$ we have

$$d_{\mathbb{D}^2}(\tilde{\Phi}(\gamma(a)), \tilde{\Phi}(\gamma(b))) \leq \lambda d_{\mathbb{D}^2}(\gamma(a), \gamma(b)),$$

where $\lambda < 1$ and depends only on the mapping class; see eg [20, Section 1.7.4].

Next assume that f is dynamically coherent. By choosing sufficiently small $\eta > 0$ we have that the rest of the assumptions of Theorem 5.4 are also satisfied. By Corollary 5.6, there exists a unique center leaf $(c, \vec{c}): \mathbb{R} \rightarrow T^1\mathbb{D}^2$ which is shadowed by γ . Claim 5.9 implies that (c, \vec{c}) is preserved by \tilde{f} . Using (5-5) and the fact that γ is coarsely contracting under $\tilde{\Phi}$, it is easy to show that (c, \vec{c}) is coarsely contracting under \tilde{f} . Hence, \tilde{f} has a fixed point $x \in (c, \vec{c})$.

Recall that the center foliation is obtained by taking the intersection of the center-stable and the center-unstable foliations. Consider the center-unstable leaf L which contains (c, \vec{c}) . Because $E^c \oplus E^{ss}$ is transverse to the circle fibers of T^1S , the restriction of the bundle map $\pi|_L: L \rightarrow \mathbb{D}^2$ is a diffeomorphism. Note that π sends a center leaves to its underlying curve. Pick a point $y \in L$ on the local stable manifold of x . Consider the sequence $\{\tilde{f}^n(y) : n \geq 0\}$ and corresponding sequence of center leaves which contain these points $\{(c_n, \vec{c}_n) : n \geq 0\} \subset L$. Because $\pi|_L$ is a diffeomorphism, $c_n: \mathbb{R} \rightarrow \mathbb{D}^2$ are all distinct and therefore have distinct pairs of boundary points (c_{n-}, c_{n+}) for $n \geq 0$. Each c_n partitions \mathbb{D}^2 into two connected components: one containing $\pi(x)$ and the other one containing $\pi(\tilde{f}^m(y))$ for $m < n$. It follows that one of the sequences $\{c_{n-}, n \geq 0\}$ and $\{c_{n+}, n \geq 0\}$ is nonincreasing and the other one is nondecreasing (we orient $\partial\mathbb{D}^2$ counterclockwise). For the sake of concreteness, we can assume that $\{c_{n-} : n \geq 0\}$ is nonincreasing and that $\{c_{n+} : n \geq 0\}$ is nondecreasing (see Figure 3). Let

$$\ell_- = \lim_{n \rightarrow \infty} c_{n-}, \quad \ell_+ = \lim_{n \rightarrow \infty} c_{n+}.$$




Figure 3: Depiction of the elements in the proof.

Also let c_{∞} be the center leaf determined by (ℓ_{-}, ℓ_{+}) .

The pairs (c_{n-}, c_{n+}) are all distinct and, hence, at least one of these sequences does not stabilize at the limit point, say $\{c_{n-} : n \geq 0\}$ does not stabilize. Recall that, by Claim 5.9, we have $c_{n-} = \partial \tilde{\Phi}^n(c_{0-})$. Hence ℓ_{-} is a contracting fixed point for $\partial \tilde{\Phi}$ (at least on one side). By Proposition 5.7, c_{-} and c_{+} are expanding fixed points. Hence, $\ell_{-} \neq c_{-}$ and consequently c_{∞} and c are disjoint. By construction, c_{∞} separates c from the sequence $\{c_n : n \geq 0\}$. In particular, we have, for all $n \geq 0$,

$$d_{\mathbb{D}^2}(\pi(x), \pi(\tilde{f}^n y)) \geq d_{\mathbb{D}^2}(\pi(x), c_{\infty}) = \rho > 0,$$

which yields a contradiction as $\tilde{f}^n y$ lies on the stable manifold of x . □

Remark 5.10 The proof can be extended to rule out a weaker version of dynamical coherence. Namely, we show that there does not exist an invariant foliation \mathcal{F}^{cu} tangent to $E^c \oplus E^{uu}$. Indeed, using a leaf of the center-stable branching foliation (see [18])¹ and, assuming that there is an invariant foliation tangent to $E^c \oplus E^{uu}$, one can use the same argument because the argument takes place inside a center-stable leaf.

¹It is always possible to consider a “lowermost” center-stable leaf which will be fixed by \tilde{f} .

5.5 Further remarks and extensions

Using the branching foliations of Burago and Ivanov [18], some of the intermediate results in the proof above may be established in the dynamically incoherent setting. In particular, we obtain information about complete curves tangent to the center distribution as well as their coarse dynamics.

Putting together Theorem 5.5, the ideas in Theorem 5.4 and a key result from [18], we obtain the following:

Proposition 5.11 *For every $\varepsilon > 0$ there exists $\delta > 0$ such that if $f: T^1S \rightarrow T^1S$ is a partially hyperbolic diffeomorphism whose splitting is δ -close to that of \mathcal{G}^t , then every $c: \mathbb{R} \rightarrow T^1S$ tangent to E_f^c and parametrized by arc-length is ε -shadowed by a unique orbit of \mathcal{G}^t . Moreover, for every orbit γ of \mathcal{G}^t , there is a unique center curve which is ε -shadowed by γ .*

Proof If the center bundle E_f^c integrates to an invariant foliation then the result is a direct consequence of Theorems 5.5 and 5.4 (which in particular give that the center bundle should be uniquely integrable).

Now consider the case when E_f^c does not integrate to a foliation. Then the existence of a unique orbit of \mathcal{G}^t shadowing a given center curve is given by Theorem 5.5 and the fact that no two center curves are shadowed by the same orbit of \mathcal{G}^t is given by Theorem 5.4. Indeed, in Theorem 5.4 one does not actually need that the diffeomorphism f is dynamically coherent: the one important point where dynamical coherence was used was to reduce to the case when the two center curves which were shadowed by the same orbit of the flow were in the same center-unstable leaf (or center-stable leaf). This can be done in the absence of dynamical coherence by appealing to the fundamental results of [18]. In particular, Proposition 3.1 of [18] implies that saturating by strong stable (resp. unstable) manifolds gives a surface tangent to the center-stable bundle (resp. center-unstable). Then the main technical result of [18] provides branching foliations which can be used to replace the center-stable and center-unstable leaves.

To show that every orbit γ of \mathcal{G}^t shadows some center curve, we apply [18, Key Lemma] to approximate $E_f^s \oplus E_f^c$ and $E_f^c \oplus E_f^u$ by integrable distributions E_n and F_n . Moreover, Theorem 7.2 of [18] implies that the integral leaves of E_n are pairwise at bounded distance in $T^1\mathbb{D}$ (and similarly for F_n). The intersection gives a one-dimensional foliation \mathcal{F}_n^c and as $n \rightarrow \infty$ the leaves converge uniformly to center curves. By applying Theorem 5.5, we have that for all sufficiently large n there exists




Figure 4: Plausible behavior on a periodic cs -leaf associated to a regular periodic point of the pseudo-Anosov. The dotted lines correspond to the strong stable manifolds.

a unique leaf ℓ_n in \mathcal{F}_n^c which is shadowed by γ . As $n \rightarrow \infty$ the curves ℓ_n converge to the posited center leaf. □

Using this improvement, one can obtain the following:

Corollary 5.12 *Let f be a diffeomorphism that satisfies the assumptions of Theorem 5.1. Then there exists a center leaf c such that the union of the stable manifolds through c does not form a complete surface.*

Proof Let c be the center leaf constructed in the proof of Theorem 5.1, ie the center leaf which is shadowed by the geodesic which corresponds to two repelling fixed points on the boundary. Let $W^s(c) = \bigcup_{y \in c} W^{ss}(y)$. According to [18] one knows that $W^s(c)$ is a surface tangent to $E_f^{ss} \oplus E_f^c$. Hence, its lift $\widetilde{W}^s(c)$ to $T^1\mathbb{D}^2$ is transverse to the circle fibers. By a similar argument as in Theorem 5.1, we see that $\widetilde{W}^s(c)$ cannot be a covering of \mathbb{D}^2 . Indeed, one can choose a center leaf in $\widetilde{W}^s(c)$ whose endpoints leave the attracting fixed point at infinity in between; the fact that this center curve belongs to $\widetilde{W}^s(c)$ gives a contradiction with the dynamics induced in the boundary. In particular, it cannot be complete as a surface in T^1S . □

See Figure 4 for a possible behavior inside certain periodic center-stable leaf. Finally, we make some comments on the dynamics of the center curves related to [6, Problem 7.26].

Remark 5.13 The considerations about coarse dynamics of center leaves made in Section 5.3 all apply to the center curves even when the diffeomorphism is dynamically incoherent. In particular, using the dynamics at infinity of the mapping class one can simultaneously create center curves which are contracting and expanding at infinity (noncompact center leaves of stable and unstable type in the nomenclature of [6, Problem 7.26]). Further, one can also show the following facts:

- If the mapping class Φ is pseudo-Anosov, then there are no closed periodic center curves.
- When two center curves merge as shown in Figure 4, one can see that there is an invariant center curve with coarse saddle-node behavior and fixed points.

5.6 Minimality of the strong foliations

In [5] it was shown that amongst robustly partially hyperbolic diffeomorphisms on 3-manifolds, those for which one of the two strong foliations is minimal form an open and dense subset. The proof of [5] is done by performing certain C^1 perturbations and robust transitivity is used to avoid checking that these perturbations are still transitive. Because all the perturbations made in [5] (see also [6, Section 7.3.3]) can be made conservative,² the same proof immediately applies to yield the following statement: there exists an open set \mathcal{U} of conservative partially hyperbolic diffeomorphisms which are stably ergodic, satisfy the conclusions of Theorem 5.1 and for which either the strong stable foliation or the strong unstable foliation is minimal.

In our setting we can use the specific knowledge on the dynamics on center leaves (see Remark 5.13) together with the arguments of [5] to show the following stronger result:

Proposition 5.14 *Let f be a diffeomorphism which satisfies the assumptions of Theorem 5.1. Then there exists a small conservative perturbation \hat{f} of f and a C^1 -neighborhood \mathcal{U} of \hat{f} such that for every $g \in \mathcal{U}$ the strong stable and unstable foliations are both minimal. In particular, \hat{f} is stably ergodic and robustly transitive.*

Proof Without loss of generality we can assume that f has dense periodic points all of which are hyperbolic (see eg [3]).

From stable ergodicity of f we know that there is a neighborhood \mathcal{U}_0 of f such that for every volume-preserving $g \in \mathcal{U}_0$ and every periodic point p of g of stable index 2, the

²The perturbations needed are the closing lemma, C^1 -connecting lemma [3] and creation of blenders [4; 36].

stable manifold $W^s(p)$ is dense in T^1S and symmetrically for the unstable manifolds of a periodic point q of g of stable index 1 (see [5, Proposition 2.1]; notice that the proof there does not use any perturbation, just that f is transitive and partially hyperbolic).

Take p and q to be periodic points of f which belongs to a saddle-node center leaf (see Section 5.3) and are respectively the first and the last periodic points in that leaf. For the sake of concreteness assume that p has stable index 2 and q has stable index 1. Using the fact that the strong stable and strong unstable foliations are close to those of the geodesic flow, which are minimal, we can assume that every strong stable leaf of some length $L > 0$ intersects the unstable manifold of q while every strong unstable manifold of length L intersects the stable manifold of p . In the terminology of [5], this means that the stable manifold of p is a u -section and the unstable manifold of q is an s -section (see [5, Section 3]).

One can create blenders associated to p and q by conservative perturbation of f (see [36]). This implies that in a neighborhood $\mathcal{U}_1 \subset \mathcal{U}_0$ of f we have that, for every $g \in \mathcal{U}_1$, the strong unstable manifold of p_g activates the blender of q_g and vice versa. In particular, we obtain for every $g \in \mathcal{U}_1$ which preserves volume that the strong stable manifold of q_g and the strong unstable of p_g are dense in T^1S (see [5, Proposition 2.6]). The fact that every strong stable manifold of length L intersects the unstable manifold of p_g implies minimality of the strong unstable foliation for each conservative $g \in \mathcal{U}_1$ and the same argument gives minimality of the strong stable foliation (for more details, see [5, Proposition 4.1]).

It follows from minimality of the strong unstable foliation that there exists $L > 0$ such that every strong unstable manifold intersects a blender. This implies that each conservative $g \in \mathcal{U}_1$ is in the hypothesis of the main theorem of [35]. As the other strong stable foliation is minimal, this implies that it is robustly minimal even among perturbations which may not be conservative. The same argument applies to the strong unstable foliation and completes the proof. \square

Remark 5.15 This implies that the set of branching points for the center curves is dense in T^1S .

Remark 5.16 This argument applies to the other examples we have constructed in Theorem 1.1 thanks to Remark 3.3 (see also the discussions in [5; 6, Section 7.3.3]).

Acknowledgements Potrie would like to thank Thomas Barthelémé, Sergio Fenley and Steven Frankel for very helpful discussions.

Bonatti was partially supported by IFUM. Gogolev was partially supported by NSF grant DMS-1266282 and Simons grant 427063. Potrie was partially supported by CSIC group 618. This research was partially supported by the Australian Research Council.

References

- [1] **T Barbot**, *Mise en position optimale de tores par rapport à un flot d'Anosov*, Comment. Math. Helv. 70 (1995) 113–160 MR
- [2] **F Béguin, C Bonatti, B Yu**, *Building Anosov flows on 3–manifolds*, Geom. Topol. 21 (2017) 1837–1930 MR
- [3] **C Bonatti, S Crovisier**, *Réurrence et généricité*, Invent. Math. 158 (2004) 33–104 MR
- [4] **C Bonatti, L J Díaz**, *Persistent nonhyperbolic transitive diffeomorphisms*, Ann. of Math. 143 (1996) 357–396 MR
- [5] **C Bonatti, L J Díaz, R Ures**, *Minimality of strong stable and unstable foliations for partially hyperbolic diffeomorphisms*, J. Inst. Math. Jussieu 1 (2002) 513–541 MR
- [6] **C Bonatti, L J Díaz, M Viana**, *Dynamics beyond uniform hyperbolicity*, Encycl. Math. Sci. 102, Springer (2005) MR
- [7] **C Bonatti, J Franks**, *A Hölder continuous vector field tangent to many foliations*, from “Modern dynamical systems and applications” (M Brin, B Hasselblatt, Y Pesin, editors), Cambridge Univ. Press (2004) 299–306 MR
- [8] **C Bonatti, A Gogolev, R Potrie**, *Anomalous partially hyperbolic diffeomorphisms, II: Stably ergodic examples*, Invent. Math. 206 (2016) 801–836 MR
- [9] **C Bonatti, N Guelman**, *Axiom A diffeomorphisms derived from Anosov flows*, J. Mod. Dyn. 4 (2010) 1–63 MR
- [10] **C Bonatti, R Langevin**, *Un exemple de flot d'Anosov transitif transverse à un tore et non conjugué à une suspension*, Ergodic Theory Dynam. Systems 14 (1994) 633–643 MR
- [11] **C Bonatti, C Matheus, M Viana, A Wilkinson**, *Abundance of stable ergodicity*, Comment. Math. Helv. 79 (2004) 753–757 MR
- [12] **C Bonatti, K Parwani, R Potrie**, *Anomalous partially hyperbolic diffeomorphisms, I: Dynamically coherent examples*, Ann. Sci. École Norm. Sup. 49 (2016) 1387–1402 MR
- [13] **C Bonatti, A Wilkinson**, *Transitive partially hyperbolic diffeomorphisms on 3–manifolds*, Topology 44 (2005) 475–508 MR
- [14] **C Bonatti, J Zhang**, *Transverse foliations on the torus \mathbb{T}^2 and partially hyperbolic diffeomorphisms on 3–manifolds*, Comment. Math. Helv. 92 (2017) 513–550 MR

- [15] **M Brin**, *On dynamical coherence*, Ergodic Theory Dynam. Systems 23 (2003) 395–401 MR
- [16] **M Brin, D Burago, S Ivanov**, *Dynamical coherence of partially hyperbolic diffeomorphisms of the 3–torus*, J. Mod. Dyn. 3 (2009) 1–11 MR
- [17] **M Brunella**, *Separating the basic sets of a nontransitive Anosov flow*, Bull. Lond. Math. Soc. 25 (1993) 487–490 MR
- [18] **D Burago, S Ivanov**, *Partially hyperbolic diffeomorphisms of 3–manifolds with abelian fundamental groups*, J. Mod. Dyn. 2 (2008) 541–580 MR
- [19] **K Burns, A Wilkinson**, *Dynamical coherence and center bunching*, Discrete Contin. Dyn. Syst. 22 (2008) 89–100 MR
- [20] **D Calegari**, *Foliations and the geometry of 3–manifolds*, Oxford Univ. Press (2007) MR
- [21] **R D Canary, D McCullough**, *Homotopy equivalences of 3–manifolds and deformation theory of Kleinian groups*, Mem. Amer. Math. Soc. 812, Amer. Math. Soc., Providence, RI (2004) MR
- [22] **P D Carrasco, F Rodriguez-Hertz, J Rodriguez-Hertz, R Ures**, *Partially hyperbolic dynamics in dimension three*, Ergodic Theory Dynam. Systems 38 (2018) 2801–2837 MR
- [23] **A J Casson, S A Bleiler**, *Automorphisms of surfaces after Nielsen and Thurston*, Lond. Math. Soc. Stud. Texts 9, Cambridge Univ. Press (1988) MR
- [24] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Math. Ser. 49, Princeton Univ. Press (2012) MR
- [25] **A Hammerlindl, R Potrie**, *Classification of partially hyperbolic diffeomorphisms in 3–manifolds with solvable fundamental group*, J. Topol. 8 (2015) 842–870 MR
- [26] **A Hammerlindl, R Potrie**, *Partial hyperbolicity and classification: a survey*, Ergodic Theory Dynam. Systems 38 (2018) 401–443 MR
- [27] **A Hammerlindl, R Potrie, M Shannon**, *Seifert manifolds admitting partially hyperbolic diffeomorphisms*, J. Mod. Dyn. 12 (2018) 193–222 MR
- [28] **M Handel, W P Thurston**, *New proofs of some results of Nielsen*, Adv. Math. 56 (1985) 173–191 MR
- [29] **K Johannson**, *Homotopy equivalences of 3–manifolds with boundaries*, Lecture Notes in Math. 761, Springer (1979) MR
- [30] **A Katok, B Hasselblatt**, *Introduction to the modern theory of dynamical systems*, Encycl. Math. Appl. 54, Cambridge Univ. Press (1995) MR
- [31] **S Matsumoto**, *Some remarks on foliated S^1 bundles*, Invent. Math. 90 (1987) 343–358 MR
- [32] **D McCullough**, *Virtually geometrically finite mapping class groups of 3–manifolds*, J. Differential Geom. 33 (1991) 1–65 MR

- [33] **R T Miller**, *Geodesic laminations from Nielsen’s viewpoint*, Adv. Math. 45 (1982) 189–212 MR
- [34] **R Potrie**, *A few remarks on partially hyperbolic diffeomorphisms of \mathbb{T}^3 isotopic to Anosov*, J. Dynam. Differential Equations 26 (2014) 805–815 MR
- [35] **E R Pujals**, **M Sambarino**, *A sufficient condition for robustly minimal foliations*, Ergodic Theory Dynam. Systems 26 (2006) 281–289 MR
- [36] **F Rodriguez Hertz**, **MA Rodriguez Hertz**, **A Tahzibi**, **R Ures**, *Creation of blenders in the conservative setting*, Nonlinearity 23 (2010) 211–223 MR
- [37] **F Rodriguez Hertz**, **MA Rodriguez Hertz**, **R Ures**, *Accessibility and stable ergodicity for partially hyperbolic diffeomorphisms with 1D-center bundle*, Invent. Math. 172 (2008) 353–381 MR
- [38] **F Rodriguez Hertz**, **MA Rodriguez Hertz**, **R Ures**, *A non-dynamically coherent example on \mathbb{T}^3* , Ann. Inst. H Poincaré Anal. Non Linéaire 33 (2016) 1023–1032 MR
- [39] **Y Shi**, *Perturbations of partially hyperbolic automorphisms on Heisenberg nilmanifold*, PhD thesis, Peking University (2014) Available at <http://www.theses.fr/2014DIJ0S048>
- [40] **W P Thurston**, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. 19 (1988) 417–431 MR
- [41] **F Waldhausen**, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. 87 (1968) 56–88 MR
- [42] **A Wilkinson**, *Conservative partially hyperbolic dynamics*, from “Proceedings of the International Congress of Mathematicians, III” (R Bhatia, A Pal, G Rangarajan, V Srinivas, M Vanninathan, editors), Hindustan, New Delhi (2010) 1816–1836 MR

*Institut de Mathématiques de Bourgogne, CNRS - URM 5584, Université de Bourgogne
Dijon, France*

*Department of Mathematics, The Ohio State University
Columbus, OH, United States*

*School of Mathematical Sciences, Monash University
Clayton VIC, Australia*

*Facultad de Ciencias - Centro de Matemática, Universidad de la República
Montevideo, Uruguay*

*Current address for RP: Institute for Advanced Study
Princeton, NJ, United States*

*bonatti@u-bourgogne.fr, gogolyev.1@osu.edu,
andy.hammerlindl@monash.edu, rpotrie@cmat.edu.uy*

Proposed: Dmitri Burago

Received: 17 August 2018

Seconded: Leonid Polterovich, David M Fisher

Revised: 9 October 2019