



# *Geometry & Topology*

Volume 28 (2024)

**The desingularization of the theta divisor of a cubic threefold  
as a moduli space**

AREND BAYER

SJOERD VIKTOR BEENTJES

SOHEYLA FEYZBAKHS

GEORG HEIN

DILETTA MARTINELLI

FATEMEH REZAAE

BENJAMIN SCHMIDT

# The desingularization of the theta divisor of a cubic threefold as a moduli space

AREND BAYER  
SJOERD VIKTOR BEENTJES  
SOHEYLA FEYZBAKHS  
GEORG HEIN  
DILETTA MARTINELLI  
FATEMEH REZAAE  
BENJAMIN SCHMIDT

We show that the moduli space  $\overline{M}_X(v)$  of Gieseker stable sheaves on a smooth cubic threefold  $X$  with Chern character  $v = (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$  is smooth and of dimension four. Moreover, the Abel–Jacobi map to the intermediate Jacobian of  $X$  maps it birationally onto the theta divisor  $\Theta$ , contracting only a copy of  $X \subset \overline{M}_X(v)$  to the singular point  $0 \in \Theta$ .

We use this result to give a new proof of a categorical version of the Torelli theorem for cubic threefolds, which says that  $X$  can be recovered from its Kuznetsov component  $\text{Ku}(X) \subset \text{D}^b(X)$ . Similarly, this leads to a new proof of the description of the singularity of the theta divisor, and thus of the classical Torelli theorem for cubic threefolds, ie that  $X$  can be recovered from its intermediate Jacobian.

14D20; 14F08, 14J30, 14J45

1. Introduction	128
2. Cubic threefolds and intermediate Jacobians	131
3. Divisors on hyperplane sections	133
4. Notions of stability	134
5. Construction of sheaves	142
6. Variation of stability	143
7. Proof of the main theorem	150
8. Kuznetsov component	153
List of symbols	158
References	158

# 1 Introduction

Moduli spaces of sheaves provide examples of algebraic varieties with an interesting and rich geometry and they have been widely studied in the past few decades. In particular, there are many strong results regarding moduli spaces on surfaces, while the situation on threefolds is less understood. We refer to Huybrechts and Lehn [23] for a more detailed account of the theory, which has been revolutionized by the introduction of stability conditions on triangulated categories by Bridgeland [12].

Perhaps the main player of the seminal paper by Clemens and Griffiths [14] on the geometry of cubic threefolds is the theta divisor  $\Theta$  of its intermediate Jacobian  $J(X)$ . Various authors have studied parametrizations of the theta divisor by moduli spaces of sheaves; see Artebani, Kloosterman and Pacini [3], Beauville [9] and Iliev [24].

In this paper, we find a new, and in a sense most efficient, parametrization of this type: a smooth four-dimensional moduli space of stable sheaves isomorphic to the desingularization of the theta divisor.

Let  $X \subset \mathbb{P}^4$  be a smooth cubic threefold over  $\mathbb{C}$  and  $H$  the hyperplane section. Let  $\overline{M}_X(v)$  be the moduli space of Gieseker-semistable sheaves on  $X$  with Chern character  $v := (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$ .

**Theorem 7.1** *The moduli space  $\overline{M}_X(v)$  is smooth and irreducible of dimension 4. More precisely, it is the blowup of  $\Theta$  in its unique singular point. The exceptional divisor is isomorphic to the cubic threefold  $X$  itself, and parametrizes non-locally-free sheaves in  $\overline{M}_X(v)$ .*

## Moduli space in the Kuznetsov component

The original motivation for our analysis of the moduli space  $\overline{M}_X(v)$  comes from the study of moduli spaces of stable objects in a full triangulated subcategory  $\text{Ku}(X) \subset \text{D}^b(X)$  called the *Kuznetsov component*. It is defined through the semiorthogonal decomposition

$$\text{D}^b(X) = \langle \text{Ku}(X), \mathcal{O}_X, \mathcal{O}_X(H) \rangle.$$

See Kuznetsov [25] for details on the decomposition and on the Kuznetsov component.

Stability conditions on  $\text{Ku}(X)$  have been constructed in Bernardara, Macrì, Mehrotra and Stellari [11] and Bayer, Lahoz, Macrì and Stellari [5]. These stability conditions are Serre-invariant, which roughly means that stability of an object is preserved by the action of the Serre functor of  $\text{Ku}(X)$ ; see Section 8 for the precise definition. This property allows us to study stability of objects irrespective of the specific construction of stability conditions.

The class  $v$  in Theorem 7.1 is chosen as the class of the projection  $K_P$  of a skyscraper sheaf  $\mathcal{O}_P$  for a point  $P \in X$ , which is defined by the short exact sequence

$$0 \rightarrow K_P \rightarrow \mathcal{O}^{\oplus 4} \rightarrow I_P(1) \rightarrow 0.$$

These are the non-locally-free torsion-free slope-stable sheaves appearing in [Theorem 7.1](#), and we show that they are also stable as objects of  $\mathrm{Ku}(X)$  with respect to any Serre-invariant stability condition. Hence, the moduli space  $M_\sigma(v)$  of  $\sigma$ -stable objects in  $\mathrm{Ku}(X)$  of Chern character  $v$  contains  $X$ , yet its expected dimension is four. This was our first clue that this moduli space is of interest. Indeed, our next result says that the moduli spaces  $M_\sigma(v)$  and  $\overline{M}_X(v)$  agree entirely.

**Theorem 1.1** ([Theorem 8.7](#) and [Proposition 8.10](#)) *Let  $\sigma$  be an arbitrary Serre-invariant stability condition on  $\mathrm{Ku}(X)$ . Then the moduli space  $M_\sigma(v)$  is isomorphic to the moduli space  $\overline{M}_X(v)$ .*

To summarize, we project the structure sheaf of a point into the Kuznetsov component and take its moduli space. It obviously contains  $X$  but is bigger. It is the resolution of the theta divisor, with  $X$  as the exceptional divisor. Thus, we recover  $X$  from  $\mathrm{Ku}(X)$  or from the intermediate Jacobian, ie we obtain new proofs of both the categorical and classical Torelli theorem for cubic threefolds:

**Theorem 1.2** ([Corollary 7.6](#) and [Theorem 8.1](#)) *Let  $X_1$  and  $X_2$  be smooth cubic threefolds. The following are equivalent:*

- (i)  $X_1$  and  $X_2$  are isomorphic.
- (ii)  $\mathrm{Ku}(X_1)$  and  $\mathrm{Ku}(X_2)$  are equivalent as triangulated categories.
- (iii)  $J(X_1)$  and  $J(X_2)$  are isomorphic as principally polarized abelian varieties.

## Proof ideas

The proof of [Theorem 7.1](#) relies on two classical ingredients. Firstly, we use the fact that any irreducible theta divisor is normal, due to Ein and Lazarsfeld [\[16\]](#). Secondly, we use a characterization of the theta divisor of the intermediate Jacobian in terms of twisted cubics; see [Proposition 2.2](#). This was proved by Beauville in [\[9\]](#), but it can also be deduced from the description of  $\Theta$  as differences of lines in Clemens and Griffiths [\[14\]](#); see [Remark 2.3](#).

The strategy to prove [Theorem 7.1](#) is to vary the notion of stability and reach a detailed description of the objects that belong to the moduli space  $\overline{M}_X(v)$  through wall-crossing. Since  $X$  has Picard rank one, Gieseker stability cannot be varied. This is where the derived category comes into play in the form of tilt-stability introduced in Bridgeland [\[13\]](#) for K3 surfaces, and then further generalized to other surfaces and threefolds in Arcara and Bertram [\[2\]](#) and Bayer, Macrì and Toda [\[7\]](#). In fact, we give a complete description of the wall and chamber structure; see [Section 6](#). Once a set-theoretic description of  $\overline{M}_X(v)$  has been reached, we use standard deformation theory arguments in [Corollary 6.9](#) to deduce that it is smooth and of dimension four.

To prove [Theorem 8.7](#), we first show the claim for the specific stability condition constructed in Bayer, Lahoz, Macrì and Stellari [\[5\]](#) which are Serre-invariant by Pertusi and Yang [\[35\]](#). We then prove in a completely separate argument that our moduli space is independent of the choice of Serre-invariant stability conditions  $\sigma$ . The essential ingredient in this last argument is the weak Mukai lemma from [\[35\]](#).

## Related work

In the recent paper [1], Altavilla, Petković and Rota studied moduli spaces of some torsion sheaves in the Kuznetsov components of Fano threefolds with Picard rank one and index two. In the case of cubic threefolds they study  $M_\sigma([S^2(K_P)])$  ( $S$  is the Serre functor on  $\text{Ku}(X)$ ), but do not obtain our detailed geometric description. A key difference is that in their case the moduli space in the Kuznetsov component is different from the moduli space of Gieseker-semistable sheaves.

Classical Torelli is the implication (iii)  $\implies$  (i) in [Theorem 1.2](#), which was first proved in Clemens and Griffiths [14]. The implication (ii)  $\implies$  (iii) was first established in Bernardara, Macrì, Mehrotra and Stellari [11, Theorem 1.1], where it was shown that the Fano variety of lines  $F(X)$  can be recovered from  $\text{Ku}(X)$  as a moduli space of stable objects. Thus, one obtains the intermediate Jacobian  $J(X)$  as the Albanese variety of  $F(X)$ . A more recent argument for (ii)  $\implies$  (iii) can be deduced from Perry’s categorical construction of intermediate Jacobians [34, Section 5.3], when the equivalence is given by a Fourier–Mukai kernel on  $X_1 \times X_2$ . Instead, our paper gives a very direct geometric argument for (ii)  $\implies$  (i), as well as a variant of the proof of classical Torelli via the description of the singularity of theta divisor implied by [Theorem 7.1](#).

Since this article originally appeared on the arXiv, Feyzbakhsh and Pertusi [17] and Zhang [40] proved uniqueness of Serre-invariant stability conditions on  $\text{Ku}(X)$ . [Proposition 8.10](#) in the last section could now be obtained as an immediate corollary.

## Acknowledgements

This work originated from a working group at the workshop *Semiorthogonal decompositions, stability conditions and sheaves of categories* in Toulouse in May 2018, a meeting of the CIMI network “Semiorthogonal decompositions and stability condition”. We are deeply grateful to the organizers of the workshop for the nice and productive atmosphere. In particular, we would like to thank Aaron Bertram, Naoki Koseki, Marin Petković, Maxim Smirnov and Benjamin Sung for very useful discussions we had in Toulouse at the beginning of the project. We would also like to express our gratitude to Marcello Bernardara, Emanuele Macrì, Alex Perry, Laura Pertusi and Richard Thomas for many useful comments on the subject of this paper. In addition, we are very grateful to the referee for many detailed comments, making the exposition more accurate and readable.

Bayer, Beentjes, Feyzbakhsh, Martinelli and Rezaee were in part supported by ERC starting grant 337039. Bayer and Rezaee were in part supported by ERC consolidator grant 819864 WallCrossAG. Rezaee was partially supported by EPSRC grant EP/T015896/1 and Maxwell Institute Research Fellowship EP/W522648/1. Feyzbakhsh was partially supported by EPSRC postdoctoral fellowship EP/T018658/1. This material is also partially based upon work supported by the NSF under grant DMS-1440140 while Bayer, Martinelli and Rezaee were in residence at the MSRI in Berkeley, California, during the Spring 2019 semester.

## 2 Cubic threefolds and intermediate Jacobians

Let  $X \subset \mathbb{P}^4$  be a smooth cubic threefold. In their celebrated article [14], Clemens and Griffiths introduced the *intermediate Jacobian* of  $X$ . It is the complex torus defined as

$$J(X) := H^{2,1}(X)^\vee / H_3(X, \mathbb{Z}) = H^1(\Omega_X^2)^\vee / H_3(X, \mathbb{Z}).$$

It turns out that  $J(X)$  is a principally polarized abelian variety of dimension five.

Let  $\{Z_b\}_{b \in \mathcal{B}}$  be a family of 1-cycles over a variety  $\mathcal{B}$ . The choice of a basepoint  $b_0 \in \mathcal{B}$  leads to an Abel–Jacobi map  $\Psi_{\mathcal{B}}: \mathcal{B} \rightarrow J(X)$  as follows. For any  $b \in \mathcal{B}$  the cycle  $Z_b - Z_{b_0}$  has degree 0, ie it is homologically trivial, and can be written as the boundary  $\partial\Gamma$  for a 3-chain  $\Gamma$ . The integral  $\int_{\Gamma}$  is an element in  $H^{1,2}(X)^\vee$  whose class in  $J(X)$  is the image of the Abel–Jacobi map. By [19, Theorem 2.20] the map  $\Psi_{\mathcal{B}}$  is algebraic along the smooth locus of  $\mathcal{B}$ .

If  $Z_b = C$  is a smooth curve, then the induced morphism on tangent spaces has been described by Welters; see [39, Section 2]. Recall that the tangent space of the Hilbert scheme at  $C$  is naturally given by  $H^0(\mathcal{N}_{C/X})$ , where  $\mathcal{N}_{C/X}$  is the normal bundle. The tangent space of  $J(X)$  at any point is given by  $H^{1,2}(X)^\vee = H^1(\Omega_X^2)^\vee$ . By definition, the *infinitesimal Abel–Jacobi map*

$$\psi_C: H^0(\mathcal{N}_{C/X}) \rightarrow H^1(\Omega_X^2)^\vee$$

is the map of tangent spaces induced by  $\Psi_{\mathcal{B}}$ . We get a dual morphism

$$\psi_C^\vee: H^1(\Omega_X^2) \rightarrow H^0(\mathcal{N}_{C/X})^\vee.$$

**Lemma 2.1** *The following diagram is commutative and has exact rows and columns:*

$$\begin{array}{ccccc} & & 0 & & \\ & & \downarrow & & \\ & & H^0(\mathcal{I}_C(H)) & & \\ & & \downarrow & & \\ & & H^0(\mathcal{O}_X(H)) & \xrightarrow{\cong} & H^1(\Omega_X^2) \\ & & \downarrow & & \downarrow \psi_C^\vee \\ H^0(\mathcal{N}_{C/\mathbb{P}^4}(-2H)) & \longrightarrow & H^0(\mathcal{O}_C(H)) & \longrightarrow & H^0(\mathcal{N}_{C/X})^\vee \end{array}$$

**Proof** This is mostly [39, Lemma 2.8] and the preceding construction of the morphisms. The map  $H^0(\mathcal{O}_X(H)) \rightarrow H^1(\Omega_X^2)$  is the connecting morphism in a long exact sequence

$$H^0(\Omega_{\mathbb{P}^4}^3 \otimes \mathcal{O}_X(3H)) \rightarrow H^0(\mathcal{O}_X(H)) \rightarrow H^1(\Omega_X^2) \rightarrow H^1(\Omega_{\mathbb{P}^4}^3 \otimes \mathcal{O}_X(3H)).$$

The wedge product induces a perfect pairing  $\Omega_{\mathbb{P}^4}^3 \otimes \Omega_{\mathbb{P}^4} \rightarrow \mathcal{O}_{\mathbb{P}^4}(-5)$ . Therefore,  $\Omega_{\mathbb{P}^4}^3 = T_{\mathbb{P}^4}(-5)$ . For  $i = 0, 1$  we have

$$H^i(T_{\mathbb{P}^4} \otimes \mathcal{O}_X(-2H)) = 0. \quad \square$$

Recall that the Lefschetz hyperplane theorem says that the hyperplane section  $H \in \text{Pic}(X)$  generates the Picard group. One can use twisted cubics to characterize the theta divisor of  $J(X)$ . A proof of the following result can be found in [9, Proposition 5.2]. Let  $\mathcal{T}$  be the open locus of smooth twisted cubics in the Hilbert scheme of  $X$ , and let  $\overline{\mathcal{T}}$  be its closure.

**Proposition 2.2** *The Abel–Jacobi map  $\varphi: \overline{\mathcal{T}} \rightarrow J(X)$  with basepoint of class  $H^2$  is algebraic. Its image is a theta divisor  $\Theta \subset J(X)$  and its generic fiber is isomorphic to  $\mathbb{P}^2$ .*

**Remark 2.3** Proposition 2.2 can be deduced from the description of  $\Theta$  as differences of lines as well. We give a rough sketch of the argument here.

Let  $F$  be the Fano variety of lines on  $X$ . According to [14] the morphism  $F \times F \rightarrow J(X)$  that maps  $(L, L') \mapsto [L] - [L']$  is generically a 6-to-1 cover of  $\Theta$ .

Since a twisted cubic  $C \subset X$  lies in a unique cubic surface  $Y \subset X$ , the morphism  $\mathcal{T} \rightarrow J(X)$  factors via the moduli space  $\mathcal{F}$  of pairs  $(D, Y)$ , where  $Y$  is a cubic surface and  $D$  is the divisor class of a twisted cubic. The generic fiber of the morphism  $\mathcal{T} \rightarrow \mathcal{F}$  is given by  $\mathbb{P}(H^0(\mathcal{O}_Y(D))) = \mathbb{P}^2$ . Indeed,  $\mathcal{O}_Y(D)$  is the pullback of  $\mathcal{O}_{\mathbb{P}^2}(1)$  if  $Y$  is written as the blowup of six general points in  $\mathbb{P}^2$ .

If  $D$  is the class of a twisted cubic on a smooth cubic surface, then  $D - H^2$  can be written as the difference of two lines on a cubic surface. Therefore, the Abel–Jacobi morphism maps onto  $\Theta$ . Moreover, there are precisely six ways to write  $D - H^2$  as the difference of two lines. Together with the fact that  $F \times F \rightarrow J(X)$  is generically a 6-to-1 cover of  $\Theta$ , we get that  $\mathcal{F} \rightarrow \Theta$  has degree 1.

**Lemma 2.4** *Let  $\mathbb{P}^1 \cong C \subset X \subset \mathbb{P}^4$  be a twisted cubic. Then*

$$\mathcal{N}_{C/\mathbb{P}^4} = \mathcal{O}_{\mathbb{P}^1}(5)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(3), \quad h^0(\mathcal{N}_{C/X}) = 6 \quad \text{and} \quad h^1(\mathcal{N}_{C/X}) = 0.$$

*In particular, the Hilbert scheme  $\mathcal{T}$  is smooth of dimension six.*

**Proof** We have a short exact sequence

$$0 \rightarrow \mathcal{N}_{C/\mathbb{P}^3} = \mathcal{O}_{\mathbb{P}^1}(5)^{\oplus 2} \rightarrow \mathcal{N}_{C/\mathbb{P}^4} \rightarrow \mathcal{N}_{\mathbb{P}^3/\mathbb{P}^4} \otimes \mathcal{O}_C = \mathcal{O}_{\mathbb{P}^1}(3) \rightarrow 0.$$

Since  $\text{Ext}^1(\mathcal{O}_{\mathbb{P}^1}(3), \mathcal{O}_{\mathbb{P}^1}(5)) = 0$ , we get  $\mathcal{N}_{C/\mathbb{P}^4} = \mathcal{O}_{\mathbb{P}^1}(5)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(3)$ . Next, we have a short exact sequence

$$0 \rightarrow \mathcal{N}_{C/X} \rightarrow \mathcal{N}_{C/\mathbb{P}^4} = \mathcal{O}_{\mathbb{P}^1}(5)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(3) \rightarrow \mathcal{N}_{X/\mathbb{P}^4} \otimes \mathcal{O}_C = \mathcal{O}_{\mathbb{P}^1}(9) \rightarrow 0.$$

Thus,  $\mathcal{N}_{C/X}$  has degree 4 and can only be  $\mathcal{O}_{\mathbb{P}^1}(m) \oplus \mathcal{O}_{\mathbb{P}^1}(4-m)$  for some  $-1 \leq m \leq 5$ . The claim about the cohomology of  $\mathcal{N}_{C/X}$  holds for each of them.  $\square$

**Lemma 2.5** *Along the locus of smooth curves  $\mathcal{T} \subset \overline{\mathcal{T}}$ , the Abel–Jacobi morphism  $\varphi$  has differential of rank four.*

**Proof** Let  $C \subset X$  be a smooth twisted cubic. Clearly, restriction maps  $H^0(\mathcal{O}_X(H)) \cong \mathbb{C}^5$  surjectively onto  $H^0(\mathcal{O}_C(H)) \cong \mathbb{C}^4$ . By Lemma 2.4, we have  $h^0(\mathcal{N}_{C/\mathbb{P}^4}(-2H)) = h^0(\mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(-3)) = 0$ .

By Lemma 2.1, we get a commutative diagram

$$\begin{CD} \mathbb{C}^5 @>\cong>> H^1(\Omega_X^2) \\ @VVV @VV\psi_C^\vee V \\ \mathbb{C}^4 @<< \hookrightarrow H^0(\mathcal{N}_{C/X})^\vee \end{CD}$$

Therefore,  $\psi_C^\vee$  has rank four. □

The singularities of the theta divisor were computed in [33, page 348]. Another proof was given in [8, Main Theorem and Proposition 2]. We will not need this full description and instead rely only on normality.

**Theorem 2.6** [16, Theorem 1] *Any irreducible theta divisor of an abelian variety is normal.*

**Lemma 2.7** *Up to numerical equivalence, the Todd class of  $X$  is  $\text{td}(X) = (1, H, \frac{2}{3}H^2, \frac{1}{3}H^3)$ . In particular, for any  $E \in D^b(X)$ ,*

$$\chi(E) = \text{ch}_3(E) + H \cdot \text{ch}_2(E) + \frac{2}{3}H^2 \cdot \text{ch}_1(E) + \frac{1}{3}H^3 \cdot \text{ch}_0(E).$$

**Proof** By Kodaira vanishing  $H^i(\mathcal{O}_X) = 0$  for  $i \neq 0$ , and therefore,  $\chi(\mathcal{O}_X) = 1$ . By the Hirzebruch–Riemann–Roch Theorem we get  $\text{td}_3(X) = \chi(\mathcal{O}_X) = \frac{1}{3}H^3$ . Similarly, Kodaira vanishing implies  $H^i(\mathcal{O}_X(-H)) = 0$  for  $i \neq 0$ . Again by Hirzebruch–Riemann–Roch,

$$0 = \chi(\mathcal{O}_X(-H)) = -\frac{1}{6}H^3 + H \cdot \frac{1}{2}H^2 - \text{td}_2(X) \cdot H + \frac{1}{3}H^3.$$

Since  $X$  has Picard rank one, this is only possible if  $\text{td}_2(X) = \frac{2}{3}H^2$ . □

**Lemma 2.8** *The numerical Chow ring  $\text{CH}_n^*(X)$  has a basis given by  $1, H, \frac{1}{3}H^2$  and  $\frac{1}{3}H^3$ . In particular, if  $E \in D^b(X)$ , then  $\text{ch}_2(E) \in \frac{1}{6}H^2 \cdot \mathbb{Z}$ , and  $\text{ch}_3(E) \in \frac{1}{6}H^3 \cdot \mathbb{Z}$ .*

**Proof** Since  $\text{Pic}(X)$  is generated by  $H$ , the group  $\text{CH}_n^2(X)$  is generated by a rational multiple of  $H^2$ . A general hyperplane section of  $X$  is a smooth cubic surface, which contains lines. The class of such a line is  $\frac{1}{3}H^2$ . Since  $H^3 = 3$ , the class has to be indivisible. Since  $\frac{1}{3}H^3$  is the class of a point, the group  $\text{CH}_n^3(X)$  must be generated by it.

The claim about second Chern characters follows directly from  $\text{ch}_2(E) = \frac{1}{2}c_1^2(E) - c_2(E)$ . The claim about  $\text{ch}_3(E)$  follows from Lemma 2.7 and the fact that  $\chi(E) \in \mathbb{Z}$ . □

### 3 Divisors on hyperplane sections

We need to understand the singularities that can occur on hyperplane sections of  $X$ .

**Proposition 3.1** *Any cubic hyperplane section  $Y = V \cap X \subset \mathbb{P}^4$  is normal and integral.*



**Proof** Since hypersurfaces satisfy condition S2, by Serre's condition [10, Section 031S], it is enough to show that  $Y$  has isolated singularities. Assume for contradiction that  $Y$  contains a curve  $C$  of singular points. Let  $F$  and  $x$  be the defining equations of  $X$  and  $V$ , respectively. Then  $\partial F/\partial x$  is a homogeneous degree 2 polynomial and hence vanishes somewhere along  $C$ . At such a point, all partial derivatives of  $F$  vanish, hence it is a singular point of  $X$ , a contradiction.  $\square$

In order to deal with singular hyperplane sections, we need to recall the relation between Weil divisors and rank-one reflexive sheaves on integral normal varieties. This is very similar to the standard relation between line bundles and Cartier divisors. We refer to [10, Tag 0EBK] or [36] for proofs of the following facts. They can also be found in [22] in more generality.

Let  $Y$  be a normal integral projective variety. By  $\text{Cl}(Y)$  we denote the *group of Weil divisors modulo rational equivalence*. For two rank-one reflexive sheaves  $L_1, L_2 \in \text{Coh}(Y)$  we can define a new rank-one reflexive sheaf by  $(L_1 \otimes L_2)^{\vee\vee}$ . This defines a group law for rank-one reflexive sheaves on  $Y$ , where inverses are given by  $L \mapsto L^\vee$ . For any effective prime divisor  $D$  one can define a rank-one reflexive sheaf  $\mathcal{O}_Y(D) := \mathcal{I}_D^\vee$ . This can be linearly extended to any divisor.

- Proposition 3.2** (i) *The group of isomorphism classes of rank-one reflexive sheaves is isomorphic to  $\text{Cl}(Y)$  under the homomorphism  $D \mapsto \mathcal{O}_Y(D)$ .*
- (ii) *To every nonzero section  $s \in H^0(L)$  of a rank-one reflexive sheaf  $L$ , one can associate an effective divisor  $D$  on  $Y$ .*
- (iii) *For any effective Weil divisor  $D$  on  $Y$ , there is a section  $s \in H^0(\mathcal{O}_Y(D))$  such that the associated divisor is given by  $D$ .*
- (iv) *Two sections  $s_1, s_2 \in H^0(L)$  define the same divisor if they satisfy  $s_1 = \lambda s_2$  for some  $\lambda \in \mathbb{C}^*$ .*

## 4 Notions of stability

In this section, we recall a number of notions of stability for sheaves. Let  $X$  be a smooth projective threefold, and let  $H$  be an ample divisor on  $X$ .

**Definition 4.1** [32; 38] (i) For any  $E \in \text{Coh}(X)$ , the *Mumford–Takemoto slope* is defined as

$$\mu(E) := \begin{cases} \frac{H^2 \cdot \text{ch}_1(E)}{H^3 \cdot \text{ch}_0(E)} & \text{for } \text{ch}_0(E) \neq 0, \\ +\infty & \text{for } \text{ch}_0(E) = 0. \end{cases}$$

- (ii) A sheaf  $E \in \text{Coh}(X)$  is *slope-(semi)stable* if for any nontrivial proper subsheaf  $F \hookrightarrow E$  the inequality  $\mu(F) < (\leq) \mu(E/F)$  holds.

From the definition it follows immediately that if  $\text{Pic}(X) = \mathbb{Z} \cdot H$  and  $E$  is slope-semistable with  $\text{gcd}(\text{ch}_0(E), H^2 \text{ch}_1(E)/H^3) = 1$ , then  $E$  is slope-stable.

While slope-stability suffices to construct moduli spaces of vector bundles on curves, a refinement is necessary in higher dimensions.

**Definition 4.2** We define a preorder on the polynomial ring  $\mathbb{R}[m]$  as follows.

- (i) For all nonzero  $f \in \mathbb{R}[m]$ , we have  $f < 0$ .
- (ii) If  $\text{deg}(f) > \text{deg}(g)$  for nonzero  $f, g \in \mathbb{R}[m]$ , then  $f < g$ .
- (iii) Let  $\text{deg}(f) = \text{deg}(g)$  for nonzero  $f, g \in \mathbb{R}[m]$ , and let  $a_f$  and  $a_g$  be the leading coefficients of  $f$  and  $g$ , respectively. Then  $f \leq g$  if and only if  $f(m)/a_f \leq g(m)/a_g$  for all  $m \gg 0$ .
- (iv) If  $f, g \in \mathbb{R}[m]$  with  $f \leq g$  and  $g \leq f$ , we write  $f \asymp g$ .


For any  $E \in \text{Coh}(X)$ , we denote its *Hilbert polynomial* and the terms  $\alpha_i(E)$  by

$$P(E, m) := \chi(E(mH)) = \sum_{i=0}^3 \alpha_i(E)m^i.$$

Moreover, let  $P_2(E, m) = \sum_{i=1}^3 \alpha_i(E)m^i$ .

- Definition 4.3**
- (i) The sheaf  $E$  is Gieseker-(semi)stable if for all nontrivial proper subsheaves  $F \subset E$ , the inequality  $P(F, m) < (\leq) P(E, m)$  holds.
  - (ii) The sheaf  $E$  is 2-Gieseker-(semi)stable if for all nontrivial proper subsheaves  $F \subset E$ , the inequality  $P_2(F, m) < (\leq) P_2(E/F, m)$  holds.

Note that for 2-Gieseker-semistability we could have equivalently asked  $P_2(F, m) \leq P_2(E, m)$ , but for 2-Gieseker-stability,  $P_2(F, m) < P_2(E, m)$  is a stronger condition that is almost never fulfilled for all such subsheaves. These notions imply each other as follows:



The intermediate notion of 2-Gieseker stability is not classical and will just appear in the technical parts of our arguments.

Due to [18; 30; 31; 37] there exists a projective moduli space  $\overline{M}_X(v)$  parametrizing  $S$ -equivalence classes of Gieseker-semistable sheaves with Chern character  $v$ . Here two semistable sheaves are called  $S$ -equivalent if they have the same stable factors, up to order and isomorphism, in their Jordan-Hölder filtrations:

**Proposition 4.4** [23, Proposition 1.5.2] *Any Gieseker-semistable sheaf  $E \in \text{Coh}(X)$  has a filtration*

$$0 = E_0 \hookrightarrow E_1 \hookrightarrow \dots \hookrightarrow E_n = E$$

*such that the factors  $A_i := E_i/E_{i-1}$  are Gieseker-stable with  $P(A_i, m) \asymp P(E, m)$  for  $i = 1, \dots, n$ . The sheaf*

$$\bigoplus_{i=1}^n A_i$$

*is uniquely determined (up to isomorphism) by  $E$ .*

Moreover, any sheaf  $E$  has a *Harder–Narasimhan filtration* into Gieseker-semistable factors.

**Proposition 4.5** [23, Theorem 1.3.4] *Let  $E \in \text{Coh}(X)$ . There is a unique filtration*

$$0 = E_0 \hookrightarrow E_1 \hookrightarrow \dots \hookrightarrow E_n = E$$

*such that the factors  $A_i := E_i/E_{i-1}$  are Gieseker-semistable with*

$$P(A_1, m) \succ P(A_2, m) \succ \dots \succ P(A_n, m).$$

Based on Bridgeland stability on surfaces, the notion of tilt stability was introduced in [7]. It is not quite a Bridgeland stability condition, but it turns out to suffice for our purposes. The basic idea is to change the category in which subobjects are taken when defining stability. This is done via the theory of tilting introduced in [20]. As before, let  $X$  be a smooth projective threefold with an ample divisor  $H$ .

**Definition 4.6** For any  $\beta \in \mathbb{R}$ , we define two full additive subcategories of  $\text{Coh}(X)$ :

$$\mathcal{F}_\beta(X) := \{E \in \text{Coh}(X) : \text{any slope-semistable factor } F \text{ of } E \text{ satisfies } \mu(F) \leq \beta\},$$

$$\mathcal{T}_\beta(X) := \{E \in \text{Coh}(X) : \text{any slope-semistable factor } F \text{ of } E \text{ satisfies } \mu(F) > \beta\}.$$

The category

$$\text{Coh}^\beta(X) := \langle \mathcal{T}_\beta(X), \mathcal{F}_\beta(X)[1] \rangle$$

is the full additive subcategory of those  $E \in \text{D}^b(X)$  for which  $\mathcal{H}^0(E) \in \mathcal{T}_\beta(X)$ ,  $\mathcal{H}^{-1}(E) \in \mathcal{F}_\beta(X)$  and  $\mathcal{H}^i(E) = 0$  for all  $i \neq -1, 0$ .

Note that  $\text{Hom}(T, F) = 0$  for all  $T \in \mathcal{T}_\beta(X)$  and  $F \in \mathcal{F}_\beta(X)$ , by semistability. It is well known that the category  $\text{Coh}^\beta(X)$  is abelian. A sequence of morphisms

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

in  $\text{Coh}^\beta(X)$  is a short exact sequence if and only if the induced sequence

$$A \rightarrow B \rightarrow C \rightarrow A[1]$$

is a distinguished triangle in  $\text{D}^b(X)$ .

To simplify notation, we define for any  $E \in \mathbf{D}^b(X)$  its twisted Chern character  $\text{ch}^\beta(E) := \text{ch}(E) \cdot e^{-\beta H}$ . Note that when  $\beta \in \mathbb{Z}$ , this is nothing but  $\text{ch}(E \otimes \mathcal{O}_X(-\beta H))$ .

**Definition 4.7** For  $\alpha > 0$ ,  $\beta \in \mathbb{R}$  and  $E \in \text{Coh}^\beta(X)$ , we define a slope function

$$v_{\alpha,\beta}(E) := \frac{H \cdot \text{ch}_2^\beta(E) - \frac{1}{2}\alpha^2 H^3 \cdot \text{ch}_0^\beta(E)}{H^2 \cdot \text{ch}_1^\beta(E)},$$

where again division by zero needs to be interpreted as  $+\infty$ . Analogously to slope-stability, an object  $E \in \text{Coh}^\beta(X)$  is called  $v_{\alpha,\beta}$ -(semi)stable if for all nontrivial proper subobjects  $F \hookrightarrow E$  in  $\text{Coh}^\beta(X)$  the inequality  $v_{\alpha,\beta}(F) < (\leq) v_{\alpha,\beta}(E/F)$  holds.

If it is clear from context, we will sometimes abuse notation and write *tilt*-(semi)stable instead of  $v_{\alpha,\beta}$ -(semi)stable. Note that by definition, any  $E \in \text{Coh}^\beta(X)$  satisfies  $H^2 \cdot \text{ch}_1^\beta(E) \geq 0$ . Therefore, this function plays the same role in  $\text{Coh}^\beta(X)$  as the rank does in  $\text{Coh}(X)$ .

As previously, Harder–Narasimhan filtrations exist. However, note that a version of Jordan–Hölder filtrations exists, but the stable factors are not unique up to order.

The notion of 2–Gieseker stability occurs as a limit of tilt stability as follows.

**Proposition 4.8** [13, Proposition 14.2] *Let  $E \in \mathbf{D}^b(X)$  and  $\beta < \mu(E)$ . Then  $E \in \text{Coh}^\beta(X)$  and  $E$  is  $v_{\alpha,\beta}$ -(semi)stable for  $\alpha \gg 0$  if and only if  $E \in \text{Coh}(X)$  and  $E$  is 2–Gieseker-(semi)stable.*

The statement in [13] is for K3 surfaces, but the same proof works in our setting. If  $\beta > \mu(E)$  the situation is slightly more complicated. The following proposition is a combination of [6, Lemma 2.7] and [27, Proposition 3.1].

**Proposition 4.9** *Take a  $v_{\alpha,\beta}$ -semistable object  $E \in \text{Coh}^\beta(X)$ . If  $\beta \neq \mu(E)$ , then  $\mathcal{H}^{-1}(E)$  is a reflexive sheaf, and if  $\beta \geq \mu(E)$  and  $\alpha \gg 0$ , then  $\mathcal{H}^{-1}(E)$  is a torsion-free slope-semistable sheaf and  $\mathcal{H}^0(E)$  is supported in dimension less than or equal to one.*

Semistable sheaves satisfy the Bogomolov inequality; see [23, Theorem 3.4.1]. A version for tilt stability was proved in [7, Corollary 7.3.2].

**Theorem 4.10** (Bogomolov inequality) *Let  $E \in \text{Coh}^\beta(X)$  be  $v_{\alpha,\beta}$ -semistable. Then*

$$\Delta_H(E) := (H^2 \cdot \text{ch}_1(E))^2 - 2(H^3 \cdot \text{ch}_0(E))(H \cdot \text{ch}_2(E)) \geq 0.$$

Most applications of tilt stability come from varying  $(\alpha, \beta)$  and determining what that means for the stability of a given set of objects. We visualize the parameter space of tilt stability,  $(\alpha, \beta) \in \mathbb{R}^2$  with  $\alpha > 0$ , as the upper half-plane via  $i\alpha + \beta$ . For a given class  $v \in K_0(X)$ , it turns out that there is a locally finite wall and chamber structure such that stability only changes as we cross a wall. These walls are




Figure 1: Walls are nested semicircles or a unique vertical wall (Theorem 4.12(ii)).

either semicircles with center on the  $\beta$ -axis or vertical lines; see Figures 1 and 2. In the following, we recall what this means formally.

For  $v \in K_0(X)$  we write  $\text{ch}(v)$ ,  $\mu(v)$ ,  $v_{\alpha,\beta}(v)$  and  $\Delta(v)$  to mean the appropriate versions where  $E$  is replaced by  $v$ .

**Definition 4.11** For  $v, w \in K_0(X)$ , we define

$$W(v, w) := \{(\alpha, \beta) \in \mathbb{R}_{>0} \times \mathbb{R} : v_{\alpha,\beta}(v) = v_{\alpha,\beta}(w)\}.$$

The set  $W(v, w)$  is a *numerical wall* if  $W(v, w) \neq \emptyset$  and  $W(v, w) \neq \mathbb{R}_{>0} \times \mathbb{R}$ , ie if it is a proper nontrivial subset of the upper half-plane.

Numerical walls in tilt stability have a rather simple structure, as shown in [28]:

**Theorem 4.12** (nested wall theorem) *Let  $v \in K_0(X)$  with  $\Delta(v) \geq 0$ .*

- (i) *A numerical wall for  $v$  is either a semicircle centered along the  $\beta$ -axis, or a vertical line parallel to the  $\alpha$ -axis in the upper half-plane.*




Figure 2: Walls are nested semicircles (Theorem 4.12(iii)).

- (ii) If  $\text{ch}_0(v) \neq 0$ , then there is a unique numerical vertical wall for  $v$  given by  $\beta = \mu(v)$ . The remaining numerical walls for  $v$  are split into two sets of nested semicircles, whose apexes lie on the hyperbola  $v_{\alpha,\beta}(v) = 0$ . In particular, no two distinct walls intersect.
- (iii) If  $\text{ch}_0(v) = 0$  and  $H^2 \cdot \text{ch}_1(v) \neq 0$ , then every numerical wall for  $v$  is a semicircle, whose apex lies on the ray  $\beta = (H \cdot \text{ch}_2(v))/(H^2 \cdot \text{ch}_1(v))$ .

The following is a well-known consequence of the fact that walls do not intersect.

**Corollary 4.13** *Let*

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

*be a short exact sequence of  $v_{\alpha,\beta}$ -semistable objects in  $\text{Coh}^{\beta_0}(X)$  for some  $(\alpha_0, \beta_0) \in W(F, E)$ . Then this is a short exact sequence of  $v_{\alpha,\beta}$ -semistable objects in  $\text{Coh}^{\beta}(X)$  for any  $(\alpha, \beta) \in W(E, F)$ .*

**Definition 4.14** *Let  $v \in K_0(X)$ . A numerical wall  $W$  for  $v$  is called an *actual wall* for  $v$  if there is a short exact sequence*

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

*of  $v_{\alpha,\beta}$ -semistable objects in  $\text{Coh}^{\beta}(X)$  for one  $(\alpha, \beta) \in W(F, E)$  such that  $W = W(F, E)$  and  $\text{ch}(E) = v$ .*

The above corollary implies that this is a short exact sequence in  $\text{Coh}^{\beta}(X)$  for all  $(\alpha, \beta) \in W(F, E)$ . Determining walls is the key technique in this paper. It will allow us to classify sheaves with certain Chern characters in terms of short exact sequences; see [Theorem 6.1](#). Note that the condition  $W(F, E) \neq \mathbb{R}_{>0} \times \mathbb{R}$  implies  $v_{\alpha,\beta}(F) > v_{\alpha,\beta}(E)$  on one side of such a wall. We say that the short exact sequence

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0,$$

or sometimes the wall  $W(F, E)$ , *destabilizes*  $E$ .

**Proposition 4.15** [\[6, Appendix A\]](#) *If an actual wall is induced by a short exact sequence of tilt-semistable objects  $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ , then*

$$\Delta_H(F) + \Delta_H(G) \leq \Delta_H(E),$$

*and equality can only occur if either  $F$  or  $G$  is a sheaf supported in dimension zero.*

It turns out that walls of large radius can only be induced by subobjects of small rank. The following precise statement is close to [\[15, Proposition 8.3\]](#). A proof of this version can be found in [\[29, Lemma 2.4\]](#) for the case of nonnegative ranks. The case of nonpositive ranks has the exact same proof, with reversed signs.

**Proposition 4.16** *Assume that an object  $E$  is destabilized by a semicircular wall induced by a subobject  $F \hookrightarrow E$  or quotient  $E \twoheadrightarrow F$  with  $\text{ch}_0(F) > \text{ch}_0(E) \geq 0$  or  $\text{ch}_0(F) < \text{ch}_0(E) \leq 0$ . Then the radius  $\rho$  of  $W(F, E)$  satisfies*

$$\rho^2 \leq \frac{\Delta_H(E)}{4(H^3 \cdot \text{ch}_0(F))(H^3 \cdot \text{ch}_0(F) - H^3 \cdot \text{ch}_0(E))}.$$

Tilt stability interacts nicely with the derived dual  $\mathbb{D}(\cdot) := \mathbf{R}\text{Hom}(\cdot, \mathcal{O}_X)[1]$ .

**Proposition 4.17** [7, Proposition 5.1.3] *Suppose that  $E \in \text{Coh}^\beta(X)$  is a  $v_{\alpha, \beta}$ -semistable object with  $v_{\alpha, \beta}(E) \neq \infty$ . Then there is a  $v_{\alpha, -\beta}$ -semistable object  $\tilde{E} \in \text{Coh}^{-\beta}(X)$ , a torsion sheaf  $T$  supported in dimension zero, and a distinguished triangle*

$$\tilde{E} \rightarrow \mathbb{D}(E) \rightarrow T[-1] \rightarrow \tilde{E}[1].$$

The following proposition seems to be well known to experts, but we could find no proof in the literature.

**Proposition 4.18** *Let  $E \in \text{Coh}(X)$  be torsion-free. Then  $E[1]$  is tilt-stable along the vertical wall  $\beta = \mu(E)$  if and only if  $E$  is slope-stable and reflexive. In particular, slope-stable reflexive sheaves do not get destabilized along the vertical wall.*

**Proof** If  $E$  is slope-unstable, then  $E \notin \text{Coh}^{\mu(E)}(X)$ . Assume that  $E$  is strictly slope-semistable. Then there is a short exact sequence of slope-semistable sheaves

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

such that  $\mu(F) = \mu(G)$ . Taking a shift by one, this becomes a short exact sequence in  $\text{Coh}^{\mu(E)}(X)$  with  $v_{\alpha, \mu(E)}(F[1]) = v_{\alpha, \mu(E)}(G[1])$ .

Assume that  $E$  is not reflexive, but slope-stable. Then we have a short exact sequence in  $\text{Coh}^{\mu(E)}(X)$  given by

$$0 \rightarrow T \rightarrow E[1] \rightarrow E^{\vee\vee}[1] \rightarrow 0,$$

where  $T$  is a nontrivial sheaf supported in dimension less than or equal to one. However, this sequence makes  $E[1]$  strictly tilt-semistable along  $\beta = \mu(E)$ .

Assume conversely that  $E$  is a slope-semistable reflexive sheaf. Then it is an object in  $\text{Coh}^{\mu(E)}(X)$  of maximal phase, and in particular tilt-semistable. If it is strictly semistable, then it admits a short exact sequence

$$0 \rightarrow F \rightarrow E[1] \rightarrow G[1] \rightarrow 0,$$

where  $F, G[1], \mathcal{H}^{-1}(F)[1]$  and  $\mathcal{H}^0(F)$  are also of maximal phase. In particular,  $\mathcal{H}^{-1}(F)$  and  $G$  are torsion-free and slope-semistable of slope  $\mu(E)$ , and  $\mathcal{H}^0(F)$  has support of dimension at most one.

Consider the long exact sequence

$$0 \rightarrow \mathcal{H}^{-1}(F) \rightarrow E \rightarrow G \rightarrow \mathcal{H}^0(F) \rightarrow 0.$$

Since we assume that  $E$  is strictly stable, this is a contradiction unless  $\mathcal{H}^{-1}(F) = 0$ . Taking duals we get an exact sequence

$$0 \rightarrow G^\vee \rightarrow E^\vee \rightarrow \mathcal{E}xt^1(F, \mathcal{O}_X).$$

Since  $F$  is supported in dimension less than or equal to one, this implies  $\mathcal{E}xt^1(F, \mathcal{O}_X) = 0$  and  $G^\vee \cong E^\vee$ . Hence,  $E \subsetneq G = G^{\vee\vee} = E^{\vee\vee}$ , a contradiction to  $E$  being reflexive.  $\square$

From now on, we assume  $X \subset \mathbb{P}^4$  is a smooth cubic threefold. In the later sections, we need the following result of [26, Proposition 3.2], which improves the Bogomolov inequality in the case of a Fano threefold of Picard rank one. Be aware that our notation differs from Li's.

**Theorem 4.19** *Let  $E$  be a tilt-stable with  $\text{ch}_0(E) \neq 0$  for some  $\alpha > 0$ ,  $\beta \in \mathbb{R}$ . If  $-\frac{1}{2} \leq \mu_H(E) \leq \frac{1}{2}$ , then*

$$\frac{H \cdot \text{ch}_2(E)}{H^3 \cdot \text{ch}_0(E)} \leq 0.$$

In the case of cubic threefolds, direct sums of line bundles can be detected among semistable sheaves or objects by their Chern characters, as follows.

**Proposition 4.20** (i) *If  $E$  is slope-semistable, or  $\nu_{\alpha,\beta}$ -semistable for some  $\alpha > 0$  and  $\beta < 0$ , with  $\text{ch}(E) = (r, 0, 0, eH^3)$  where  $r > 0$ , then  $e \leq 0$ . If, additionally,  $e = 0$ , then  $E \cong \mathcal{O}_X^{\oplus r}$ .*

(ii) *If  $E$  is  $\nu_{\alpha,\beta}$ -semistable for some  $\alpha > 0$  and  $\beta > 0$ , with  $\text{ch}(E) = (-r, 0, 0, eH^3)$  where  $r > 0$ , then  $e = 0$  and  $E \cong \mathcal{O}_X^{\oplus r}[1]$ .*

**Proof** In either case, Proposition 4.15 and  $\Delta(E) = 0$  imply that  $E$  has no semicircular walls.

We first claim that the only slope-stable reflexive sheaf of class  $(r, 0, 0, eH^3)$  is  $\mathcal{O}_X$ . Assume otherwise. By Proposition 4.18, such an  $E$  is also stable at the vertical wall  $\beta = 0$ , and thus, it is  $\nu_{\alpha,\beta}$ -stable for all  $\alpha > 0$  and  $\beta \in \mathbb{R}$ . Since  $\nu_{0,\beta}(E) = -\frac{1}{2}\beta > -\frac{1}{2}\beta - 1 = \nu_{0,\beta}(\mathcal{O}_X(-2H)[1])$  and both objects are stable for  $\alpha \ll 1$  and  $\beta \in (-2, 0)$ , we have  $\text{Ext}^2(\mathcal{O}_X, E) = \text{Hom}(E, \mathcal{O}_X(-2H)[1]) = 0$ . Similarly, from  $\nu_{\alpha,\beta}$ -stability for  $\alpha \ll 1$  and  $\beta \in (0, 2)$  we obtain  $\text{Ext}^2(E, \mathcal{O}_X) = \text{Hom}(\mathcal{O}_X(2H), E[1]) = 0$ . However, at least one of  $\chi(\mathcal{O}_X, E) = r + 3e$  or  $\chi(E, \mathcal{O}_X) = r - 3e$  is positive, and so  $E$  admits a morphism from  $\mathcal{O}_X$  or a morphism to  $\mathcal{O}_X$ . As both are reflexive and slope-stable of slope 0, this shows  $E \cong \mathcal{O}_X$ .

Now consider an object  $E$  as in case (i). Then  $E[1]$  is  $\nu_{\alpha,0}$ -semistable. By Proposition 4.18, its Jordan–Hölder factors are either of the form  $F[1]$  for a slope-stable reflexive sheaf  $F$  with  $\text{ch}(F) = (r_F, 0, d_F H^2, e_F H^3)$ , or a torsion sheaf supported in dimension  $\leq 1$ . In fact, Proposition 4.15 shows  $d_F = 0$  in the former case, and thus,  $F = \mathcal{O}_X$  by the previous case, and that the torsion sheaves are supported in dimension zero. As  $-3e$  is the total length of the torsion sheaves, we get  $e \leq 0$ . If  $e = 0$ , all factors are isomorphic to  $\mathcal{O}_X[1]$  and the claim follows from  $\text{Ext}^1(\mathcal{O}_X, \mathcal{O}_X) = 0$ .



In case (ii), we again consider a Jordan–Hölder filtration with respect to  $v_{\alpha,0}$ -stability. Let  $E_i \hookrightarrow E_{i+1}$  be the first filtration step where the quotient  $E_{i+1}/E_i$  is a zero-dimensional torsion sheaf  $T$ , should one exist. Then  $E_i = \mathcal{O}_X[1]^{\oplus k}$  for some  $k > 0$ . Since  $\text{Ext}^1(T, \mathcal{O}_X[1]) = H^1(T)^\vee = 0$ , we have  $E_{i+1} = E_i \oplus T$ , and so  $T$  is a subobject of  $E$ . This contradicts stability of  $E$  for  $\beta > 0$ . Thus,  $E = \mathcal{O}_X[1]^{\oplus r}$ , as claimed.  $\square$

## 5 Construction of sheaves

In this section, we introduce the sheaves that make up our moduli space  $\overline{M}_X(v)$ . It turns out that all of them are at least reflexive, and the generic one is a vector bundle. From now on  $X \subset \mathbb{P}^4$  is an arbitrary smooth cubic threefold.

Let  $Y \subset X$  be an arbitrary hyperplane section,  $D$  be an effective Weil divisor on  $Y$ , and  $V \subset H^0(\mathcal{O}_Y(D))$  be a nontrivial subspace. Then we define  $\mathcal{E}_{D,V} \in \text{D}^b(X)$  to be the cone of the induced morphism  $\mathcal{O}_X \otimes V \rightarrow \mathcal{O}_Y(D)$ . Moreover, let  $E_{D,V} := \mathcal{H}^{-1}(\mathcal{E}_{D,V})$ . Hence, we have a long exact sequence

$$0 \rightarrow E_{D,V} \rightarrow \mathcal{O}_X \otimes V \rightarrow \mathcal{O}_Y(D) \rightarrow \mathcal{H}^0(\mathcal{E}_{D,V}) \rightarrow 0.$$

If  $V = H^0(\mathcal{O}_Y(D))$ , we will drop  $V$ , and just write  $\mathcal{E}_D$  and  $E_D$ .

**Lemma 5.1** *The sheaf  $E_{D,V}$  is slope-stable and reflexive. If, additionally,  $\mathcal{H}^0(\mathcal{E}_{D,V}) = 0$ , then  $E_{D,V}$  is a vector bundle.*

**Proof** The quotient  $(\mathcal{O}_X \otimes V)/E_{D,V}$  embeds into  $\mathcal{O}_Y(D)$ . Since  $Y$  is integral by Proposition 3.1, the sheaf  $(\mathcal{O}_X \otimes V)/E_{D,V}$  must be supported on  $Y$ . Therefore,  $\text{ch}_{\leq 1}(E_{D,V}) = (\dim V, -H)$  is primitive and it is enough to show that  $E_{D,V}$  is slope-semistable. If not, let  $F \subset E_{D,V}$  be the slope-semistable subsheaf in the Harder–Narasimhan filtration of  $E_{D,V}$ . Then  $\mu(F) > \mu(E_{D,V})$  and the quotient  $E_{D,V}/F$  is torsion-free. Since  $F$  is also a subsheaf of  $\mathcal{O}_X \otimes V$ , we must have  $\mu(F) = 0$ . Let  $\text{ch}(F) = (r, 0, dH^2, eH^3)$ . The quotient  $(\mathcal{O}_X \otimes V)/F$  satisfies  $\text{ch}((\mathcal{O}_X \otimes V)/F) = (\dim V - r, 0, -dH^2, -eH^3)$ . By the snake lemma this quotient is either torsion-free or has a torsion subsheaf purely supported on  $Y$ . However, if it is not torsion-free, then its torsion-free quotient would destabilize  $\mathcal{O}_X \otimes V$ , a contradiction. As a torsion-free quotient of  $\mathcal{O}_X \otimes V$  with slope zero,  $(\mathcal{O}_X \otimes V)/F$  has to be slope-semistable as well.

The classical Bogomolov inequalities  $\Delta_H(F) \geq 0$  and  $\Delta_H((\mathcal{O}_X \otimes V)/F) \geq 0$  imply  $d = 0$ . Applying Proposition 4.20 to both  $F$  and  $(\mathcal{O}_X \otimes V)/F$  implies  $e = 0$ , and finally,  $F = \mathcal{O}_X^{\oplus r}$ . However, by construction,  $E_{D,V}$  has no global sections, a contradiction.

To see that  $E_{D,V}$  is reflexive it suffices to show that  $\mathcal{E}xt^q(E_{D,V}, \mathcal{O}_X) = 0$  for  $q \geq 2$  and  $\mathcal{E}xt^1(E_{D,V}, \mathcal{O}_X)$  is supported in dimension zero. If additionally  $\mathcal{E}xt^1(E_{D,V}, \mathcal{O}_X) = 0$ , then  $E_{D,V}$  is a vector bundle.

Clearly,  $\mathcal{E}xt^q(\mathcal{O}_X \otimes V, \mathcal{O}_X) = 0$  for  $q \neq 0$ . Because  $\mathcal{O}_Y(D)$  is a rank-one reflexive sheaf on the codimension one subvariety  $Y$ , the quotient  $(\mathcal{O}_X \otimes V)/E_{D,V} \subset \mathcal{O}_Y(D)$  is purely supported on  $Y$ . We

can use [23, Proposition 1.1.10] to see that  $\mathcal{E}xt^q((\mathcal{O}_X \otimes V)/E_{D,V}, \mathcal{O}_X) = 0$  for all  $q \neq 1, 2$ , and  $\mathcal{E}xt^2((\mathcal{O}_X \otimes V)/E_{D,V}, \mathcal{O}_X)$  is supported in dimension zero. The long exact sequence obtained from dualizing the short exact sequence

$$(1) \quad 0 \rightarrow E_{D,V} \rightarrow \mathcal{O}_X \otimes V \rightarrow (\mathcal{O}_X \otimes V)/E_{D,V} \rightarrow 0$$

implies the required vanishings.

If additionally  $\mathcal{H}^0(\mathcal{E}_{D,V}) = 0$ , then  $(\mathcal{O}_X \otimes V)/E_{D,V} = \mathcal{O}_Y(D)$  is a reflexive sheaf on the codimension-one subvariety  $Y$ , and we can use [23, Proposition 1.1.10] again to see that  $\mathcal{E}xt^2(\mathcal{O}_Y(D), \mathcal{O}_X) = 0$ . The same long exact sequence as above now implies  $\mathcal{E}xt^1(E_{D,V}, \mathcal{O}_X) = 0$ .  $\square$

Note that we will use this lemma for the case  $\text{ch}(\mathcal{O}_Y(D)) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ . It will turn out that in this case  $h^0(\mathcal{O}_Y(D)) = 3$  for any such  $D$  (see Theorem 6.1) and we will choose  $V = H^0(\mathcal{O}_Y(D))$ . Moreover, we will show that in that case  $\mathcal{H}^0(\mathcal{E}_D) = 0$ , ie  $\mathcal{O}_Y(D)$  is globally generated; see Theorem 6.1. A straightforward computation shows that in this example  $\text{ch}(E_D) = (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$ .

**Corollary 5.2** *Let  $P \in X$ . Then  $h^0(\mathcal{I}_P(H)) = 4$  and the sheaf  $K_P$  defined through the exact sequence*

$$(2) \quad 0 \rightarrow K_P \rightarrow \mathcal{O}_X^{\oplus 4} \rightarrow \mathcal{I}_P(H) \rightarrow 0$$

*satisfies  $\text{ch}(K_P) = (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$ . Moreover,  $K_P$  is reflexive and slope-stable, and locally free except at  $P$ .*

**Proof** By choosing an embedding  $K_P \hookrightarrow \mathcal{O}_X^{\oplus 3}$  we get a short exact sequence

$$0 \rightarrow K_P \rightarrow \mathcal{O}_X^{\oplus 3} \rightarrow \mathcal{I}_{P/Y}(H) \rightarrow 0$$

for some hyperplane section  $Y$ . The statement then follows from Lemma 5.1 by choosing  $D = H$  and  $V = H^0(\mathcal{I}_{P/Y}(H)) \subset H^0(\mathcal{O}_Y(H))$ .

From the defining short exact sequence (2) one immediately sees that  $K_P$  is locally free away from  $P$  (as it is the kernel of a surjective map of vector bundles), and not locally free at  $P$  (as  $\text{Ext}^2(\mathcal{O}_P, K_P) = \text{Ext}^1(\mathcal{O}_P, \mathcal{I}_P(H)) \neq 0$ ).  $\square$

## 6 Variation of stability

In this section, we investigate semistable sheaves with Chern character

$$v := (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3).$$

The main goal is to use wall-crossing to prove the following theorem, which gives a set-theoretic description of the moduli space  $\overline{M}_X(v)$ .

- Theorem 6.1** (i) Suppose that  $D$  is a Weil divisor on a (possibly singular) hyperplane section  $Y$  with  $\text{ch}(\mathcal{O}_Y(D)) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ . Then  $\mathcal{O}_Y(D)$  is globally generated, and  $h^0(\mathcal{O}_Y(D)) = 3$ . In particular, there exists a smooth twisted cubic  $C$  in  $Y$  of class  $D$ .
- (ii) A sheaf  $E$  with Chern character  $v$  is Gieseker-semistable if and only if it is either equal to the reflexive sheaf  $K_P$  for a point  $P \in X$  as in (2), or the vector bundle  $E_D$  for a Weil divisor  $D$  on a hyperplane section  $Y \subset X$  as in (1) with  $\text{ch}(\mathcal{O}_Y(D)) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ .

Note that since  $\text{ch}_1(E) = -H$ , any Gieseker-semistable sheaf of class  $v$  is slope-stable. The argument will essentially boil down to a detailed analysis of the numerical wall  $W$  defined by

$$(3) \quad \alpha^2 + \left(\beta - \frac{1}{2}\right)^2 = \frac{1}{4}.$$

At this wall, the short exact sequences (2) and (1) become destabilizing short exact sequences in  $\text{Coh}^\beta(X)$  in the form

$$0 \rightarrow \mathcal{O}_Y(D) \rightarrow E_D[1] \rightarrow \mathcal{O}_X[1]^{\oplus 3} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow I_P(H) \rightarrow K_P[1] \rightarrow \mathcal{O}_X[1]^{\oplus 4} \rightarrow 0.$$

Moreover, we can show that every object gets destabilized, and the destabilizing short exact sequence must be of one of these types; see Lemma 6.8.

### 6.1 Classification of some torsion sheaves

In this section, we prove the following proposition.

**Proposition 6.2** The wall  $W$  of equation (3) is the unique actual wall in tilt stability for objects  $G$  with Chern character  $\text{ch}(G) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ .

- (i) Above  $W$  the moduli space of tilt-semistable objects is the moduli space of Gieseker-semistable sheaves, and contains precisely the following two types of sheaves  $G$ :
- (a)  $G = \mathcal{I}_{P/Y}(H)$  for  $Y \in |H|$  and  $P \in Y$ , and
  - (b)  $G = \mathcal{O}_Y(D)$ , where  $D$  is a Weil divisor on some  $Y \in |H|$ .
- (ii) Below  $W$  the moduli space of tilt-semistable objects contains precisely the following two types of objects  $G$ :
- (a) the unique nontrivial extensions
- (4) 
$$0 \rightarrow \mathcal{O}_X[1] \rightarrow G_P \rightarrow \mathcal{I}_P(H) \rightarrow 0$$
- for points  $P \in X$ , and
- (b)  $G = \mathcal{O}_Y(D)$ , where  $D$  is a Weil divisor on some  $Y \in |H|$ .

We start by dealing with slightly more general objects without fixing  $\text{ch}_3$ .

**Lemma 6.3** *The wall  $W$  of equation (3) is the unique actual wall in tilt stability for objects  $G$  with Chern character  $\text{ch}_{\leq 2}(G) = (0, H, \frac{1}{2}H^2)$ . If  $G$  is strictly semistable along  $W$ , then any Jordan–Hölder filtration of  $G$  is given by either*

$$0 \rightarrow \mathcal{I}_Z(H) \rightarrow G \rightarrow \mathcal{O}_X[1] \rightarrow 0 \quad \text{or} \quad 0 \rightarrow \mathcal{O}_X[1] \rightarrow G \rightarrow \mathcal{I}_Z(H) \rightarrow 0,$$

where  $Z \subset X$  is a zero-dimensional subscheme of length  $\frac{1}{6}H^3 - \text{ch}_3(G)$ .

**Proof** All walls for  $(0, H, \frac{1}{2}H^2)$  intersect the vertical ray  $\beta = \frac{1}{2}$ . If  $G$  is strictly semistable along some numerical wall intersecting  $\beta = \frac{1}{2}$ , then there is a short exact sequence in  $\text{Coh}^{1/2}(X)$  of tilt-semistable objects

$$0 \rightarrow A \rightarrow G \rightarrow B \rightarrow 0$$

with equal tilt-slope. Let  $\text{ch}_{\leq 2}(A) = (r, cH, dH^2)$ . By definition of  $\text{Coh}^{1/2}(X)$  and the fact that neither  $A$  nor  $B$  can have infinite tilt-slope, we get

$$0 < H^2 \cdot \text{ch}_1^{1/2}(A) = H^3(c - \frac{1}{2}r) < H^2 \cdot \text{ch}_1^{1/2}(G) = H^3.$$

Therefore,  $c = \frac{1}{2}r + \frac{1}{2}$ , and in particular,  $r$  is odd. We will deal with the case  $r < 0$ . If  $r > 0$ , then  $B$  has negative rank and one simply has to exchange the roles of  $A$  and  $B$  in the following argument.

For  $(\alpha, \frac{1}{2}) \in W(A, G)$  we have

$$-\alpha^2 r + 2d - \frac{1}{4}r - \frac{1}{2} = v_{\alpha, 1/2}(A) = v_{\alpha, 1/2}(G) = 0.$$

Since  $\alpha^2 > 0$ , this implies  $d < \frac{1}{8}r + \frac{1}{4}$ . The fact

$$0 \leq \frac{\Delta_H(A)}{(H^3)^2} = -2dr + \frac{1}{4}r^2 + \frac{1}{2}r + \frac{1}{4}$$

implies  $d \geq \frac{1}{8}r + (1/8r) + \frac{1}{4}$ . Since  $d \in \frac{1}{6}\mathbb{Z}$ , these restrictions on  $d$  are only possible for  $r \in \{-1, -3\}$ .

If  $r = -3$ , then  $\text{ch}_{\leq 2}(A) = (-3, -H, -\frac{1}{6}H^2)$ . This case is immediately ruled out by [Theorem 4.19](#).

If  $r = -1$ , then  $\text{ch}_{\leq 2}(A) = (-1, 0, 0)$ , and by [Proposition 4.20](#), we know  $A = \mathcal{O}_X[1]$ . Then  $\text{ch}(B) = (1, H, \frac{1}{2}H^2, \text{ch}_3(G))$ . By [Proposition 4.15](#), there is no semicircular wall for  $B$ , and by [Proposition 4.8](#), the object  $B$  has to be a 2–Gieseker-stable sheaf. Since  $\text{ch}(B(-H)) = (1, 0, 0, \text{ch}_3(G) - \frac{1}{6}H^3)$ , the remaining statement follows by applying [Proposition 4.20](#) to  $B(-H)$ .  $\square$

The next step is to gain further control over the third Chern character.

**Lemma 6.4** *Let  $G$  be a  $v_{\alpha, \beta}$ –semistable object with  $\text{ch}_{\leq 2}(G) = (0, H, \frac{1}{2}H^2)$ . Then  $\text{ch}_3(G) \leq \frac{1}{6}H^3$ . If  $\text{ch}_3(G) = \frac{1}{6}H^3$  and  $(\alpha, \beta)$  is above  $W$ , then  $G \cong \mathcal{O}_Y(H)$  for some  $Y \in |H|$ .*

**Proof** We may assume  $\text{ch}_3(G) \geq \frac{1}{6}H^3$ . By [Lemma 6.3](#), the only possible wall is given by  $W$ . Therefore,  $G$  has to be tilt-semistable along  $W$ . Since  $W$  lies below the numerical wall  $W(G, \mathcal{O}_X(-H)[1])$ , we get  $\text{ext}^2(\mathcal{O}_X(H), G) = \text{hom}(G, \mathcal{O}_X(-H)[1]) = 0$ . Thus,

$$\text{hom}(\mathcal{O}_X(H), G) \geq \chi(\mathcal{O}_X(H), G) = \text{ch}_3(G) + \frac{1}{6}H^3 > 0.$$

Therefore,  $W$  is a wall for  $G$  and by Lemma 6.3, the destabilizing sequence is

$$0 \rightarrow \mathcal{O}_X(H) \rightarrow G \rightarrow \mathcal{O}_X[1] \rightarrow 0.$$

This implies  $G = \mathcal{O}_Y(H)$  for some  $Y \in |H|$  and  $\text{ch}_3(G) = \frac{1}{6}H^3$ . □

**Proof of Proposition 6.2** Assume that  $G$  is strictly tilt-semistable along  $W$ . Then Lemma 6.3 splits our problem into two cases.

Firstly, assume that  $G$  fits into a nonsplitting short exact sequence

$$0 \rightarrow \mathcal{I}_P(H) \rightarrow G \rightarrow \mathcal{O}_X[1] \rightarrow 0$$

for a point  $P \in X$ . Then clearly  $G = \mathcal{I}_{P/Y}(H)$  for some  $Y \in |H|$ . This object is tilt-stable above  $W$ , and tilt-unstable below  $W$  by precisely this sequence.

Secondly, assume that  $G$  fits into a nonsplitting short exact sequence

$$(5) \quad 0 \rightarrow \mathcal{O}_X[1] \rightarrow G \rightarrow \mathcal{I}_P(H) \rightarrow 0$$

for some  $P \in X$ . By Serre duality,  $\text{Ext}^1(\mathcal{I}_P(H), \mathcal{O}_X[1]) = h^1(\mathcal{I}_P(-H)) = 1$  and hence, there is a unique  $G$  for each  $P \in X$ . Clearly, this object is tilt-unstable above  $W$ . Assume it is also tilt-unstable below  $W$ . Then there is a short exact sequence  $0 \rightarrow A \rightarrow G \rightarrow B \rightarrow 0$  destabilizing  $G$  below the wall. However,  $G$  is strictly semistable at  $W$ , and by Lemma 6.3, this implies  $B = \mathcal{O}_X[1]$ . However, that means the short exact sequence (5) splits, a contradiction.

Lastly, assume that  $G$  is  $v_{\alpha,\beta}$ -stable for all  $(\alpha, \beta)$ . By Proposition 4.17,  $\mathbb{D}(G)$  lies in a distinguished triangle

$$(6) \quad \tilde{G} \rightarrow \mathbb{D}(G) \rightarrow T[-1] \rightarrow \tilde{G}[1],$$

where  $T$  is a torsion sheaf supported in dimension zero and  $\tilde{G} \in \text{Coh}^{-\beta}(X)$  is  $v_{\alpha,-\beta}$ -semistable. If  $\text{ch}_3(T) = t$ , then  $\text{ch}(\tilde{G}) = (0, H, -\frac{1}{2}H^2, -\frac{1}{6}H^3 + t)$ . Thus,  $\tilde{G}$  is a pure sheaf supported on a hyperplane section  $Y \in |H|$ . We can compute

$$\text{ch}(\tilde{G} \otimes \mathcal{O}_X(H)) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3 + t).$$

Thus, Lemma 6.4 gives  $t = 0$  or  $t = 1$ , and if  $t = 1$ , then  $\tilde{G} \otimes \mathcal{O}_X(H) \cong \mathcal{O}_Y(H)$ , ie  $\tilde{G} \cong \mathcal{O}_Y(H)$ . Hence there is a nontrivial morphism  $\mathcal{O}_X \rightarrow \tilde{G}$ . Since  $\text{hom}(\mathcal{O}_X, T[-i]) = 0$  for  $i > 0$ , The triangle (6) shows that there is a nontrivial morphism  $\mathcal{O}_X \rightarrow \mathbb{D}(G)$ . Dualizing this morphism leads to a nontrivial morphism  $G \rightarrow \mathcal{O}_X[1]$ . However, this is in contradiction to the assumption that  $G$  is stable along  $W$ .

If  $t = 0$ , then  $\mathbb{D}(G) = \tilde{G}$  is a sheaf, so  $\text{Ext}^q(G, \mathcal{O}_X) = 0$  for  $q > 1$ . Thus, [23, Proposition 1.1.10] implies that  $G$  is reflexive and supported on a hyperplane section  $Y \in |H|$ . This means  $G = \mathcal{O}_Y(D)$  for some Weil divisor  $D$  on  $Y$ . □

## 6.2 Set-theoretic description of the moduli space

We now prepare the proof of [Theorem 6.1](#).

**Lemma 6.5** *There are no walls along  $\beta = -1$  for tilt-semistable objects  $E$  with Chern character  $\text{ch}_{\leq 2}(E) = (3, -H, -\frac{1}{2}H^2)$ .*

**Proof** Assume there is such a wall induced by a short exact sequence

$$0 \rightarrow A \rightarrow E \rightarrow B \rightarrow 0$$

with  $\text{ch}_{\leq 2}^{-1}(A) = (r, xH, yH^2)$ . Then

$$0 < H \cdot \text{ch}_1^{-1}(A) = xH^3 < H \cdot \text{ch}_1^{-1}(E) = 2H^3$$

implies  $x = 1$ . By exchanging the roles of  $A$  and  $B$  if necessary, we may assume  $r \geq 2$ .

Using  $\Delta_H(A) \geq 0$  we get  $y \leq 1/2r$ . A straightforward computation shows that there exists  $\alpha > 0$  with  $v_{\alpha, -1}(A) = v_{\alpha, -1}(E)$  if and only if  $y > 0$ . Since  $y \in \frac{1}{6}\mathbb{Z}$ , this is only possible if  $y = \frac{1}{6}$  and  $r \in \{2, 3\}$ . Both cases  $\text{ch}_{\leq 2}^{-1}(A) = (3, H, \frac{1}{6}H^2)$  and  $\text{ch}_{\leq 2}^{-1}(A) = (2, H, \frac{1}{6}H^2)$  are directly ruled out by [Theorem 4.19](#).  $\square$

**Proposition 6.6** *Take a slope-stable sheaf  $E$  of Chern character  $(3, -H, \text{ch}_2, \text{ch}_3)$ . Then  $H \cdot \text{ch}_2 \leq -\frac{1}{2}H^3$ , and if  $\text{ch}_2 \cdot H = -\frac{1}{2}H^3$ , then  $\text{ch}_3 \leq \frac{1}{6}H^3$ . In particular, this implies that any slope-stable sheaf of Chern character  $v$  is a reflexive sheaf.*

**Proof** Since  $E$  is slope-stable, the classical Bogomolov inequality gives

$$\Delta_H(E) = (H^3)^2 - 2(3H^3)(H \cdot \text{ch}_2(E)) \geq 0,$$

which implies  $H \cdot \text{ch}_2(E) \leq \frac{1}{6}H^3$ . The case  $H \cdot \text{ch}_2(E) = \frac{1}{6}H^3$  is immediately ruled out by [Theorem 4.19](#). Since  $c_2(E) = \frac{1}{2}H^2 - \text{ch}_2(E)$  has to be an integral class, we are left to rule out  $H \cdot \text{ch}_2(E) = -\frac{1}{6}H^3$ . Assume  $H \cdot \text{ch}_2(E) = -\frac{1}{6}H^3$ . We may assume that  $E$  is a reflexive sheaf. If not, we replace it by the double dual  $E^{\vee\vee}$ , which satisfies  $H \cdot \text{ch}_2(E) \leq H \cdot \text{ch}_2(E^{\vee\vee})$ . By the first part of the argument  $H \cdot \text{ch}_2(E^{\vee\vee}) = -\frac{1}{6}H^3$  holds as well.

We first show that  $\text{ext}^2(E, E) = 0$ . Since  $H^3 \cdot \text{ch}_1^{-1/2}(E) = \frac{1}{2}H$ , any destabilizing subobject  $F \subset E$  along  $\beta = -\frac{1}{2}$  must satisfy  $H^3 \cdot \text{ch}_1^{-1/2}(F) = \frac{1}{2}H$  or  $H^3 \cdot \text{ch}_1^{-1/2}(F) = 0$ . Thus, either  $F$  or the quotient  $E/F$  have infinite tilt-slope, a contradiction. This means  $E$  is  $v_{\alpha, -1/2}$ -stable for all  $\alpha > 0$ .

By [Proposition 4.18](#), the object  $E[1]$  is tilt-stable for  $\beta = 0$  and  $\alpha \gg 0$ . Since  $H^3 \cdot \text{ch}_1(E[1]) = H^3$ , the same type of argument as above shows that there cannot be any wall along  $\beta = 0$ . Hence,  $E(-2H)[1]$  is  $v_{\alpha, \beta}$ -stable for  $\beta = -2$  and any  $\alpha > 0$ .

A straightforward computation shows that  $W(E, E(-2H)[1])$  intersects both the vertical lines  $\beta = -2$  and  $\beta = -\frac{1}{2}$ . Therefore,  $E$  and  $E(-2H)[1]$  are tilt-stable for any  $(\alpha, \beta) \in W(E, E(-2H)[1])$  and have the same phase, and thus,  $\text{ext}^2(E, E) = \text{hom}(E, E(-2H)[1]) = 0$ . Since  $E$  is stable, we know  $\text{hom}(E, E) = 1$  and hence,  $3 = \chi(E, E) = 1 - \text{ext}^1(E, E) - \text{ext}^3(E, E) \leq 1$ , a contradiction.

Now assume  $H \cdot \text{ch}_2 = -\frac{1}{2}H^3$ . We know  $E \in \text{Coh}^\beta(X)$  is  $\nu_{\alpha, \beta}$ -stable for  $\alpha \gg 0$  and  $\beta < -\frac{1}{3}$ . By Lemma 6.5, we have that  $E$  is  $\nu_{\alpha, -1}$ -stable for any  $\alpha > 0$ . One can easily compute

$$\nu_{0, -1}(\mathcal{O}_X(-2H)[1]) < \nu_{0, -1}(E),$$

which implies  $h^2(E) = \text{hom}(E, \mathcal{O}_X(-2H)[1]) = 0$ . Moreover, since  $\mu(E) = -\frac{1}{3} < \mu(\mathcal{O}_X)$ , we get  $\text{hom}(\mathcal{O}_X, E) = 0$ . Therefore,  $\chi(E) = \text{ch}_3(E) - \frac{1}{6}H^3 \leq 0$ , as claimed.

Lastly, assume that a slope-stable sheaf  $E$  of Chern character  $v$  is not reflexive. We have a short exact sequence

$$0 \rightarrow E \rightarrow E^{\vee\vee} \rightarrow T \rightarrow 0.$$

Since  $E^{\vee\vee}$  is also slope-stable, and both  $H \cdot \text{ch}_2(E)$  and  $H \cdot \text{ch}_3(E)$  are maximal, one gets  $\text{ch}(E) = \text{ch}(E^{\vee\vee})$ . This is only possible if  $T = 0$ . □

To prove Theorem 6.1, we start in the large volume limit.

**Lemma 6.7** *Take  $\beta > -\frac{1}{3}$ . An object  $\tilde{E} \in \text{Coh}^\beta(X)$  of Chern character  $-v$  is  $\nu_{\alpha, \beta}$ -semistable for  $\alpha \gg 0$  if and only if  $\tilde{E} \cong E[1]$  for a slope-stable reflexive sheaf  $E$ .*

**Proof** Take a  $\nu_{\alpha, \beta}$ -semistable object  $\tilde{E}$  of class  $-v$ . Proposition 4.9 implies that  $\mathcal{H}^{-1}(\tilde{E})$  is a slope-stable reflexive sheaf and  $\mathcal{H}^0(\tilde{E})$  is a torsion sheaf supported in dimension  $\leq 1$ . Therefore,

$$\text{ch}(\mathcal{H}^{-1}(\tilde{E})) = (3, -H, -\frac{1}{2}H^2 + \text{ch}_2(\mathcal{H}^0(\tilde{E})), \frac{1}{6}H^3 + \text{ch}_3(\mathcal{H}^0(\tilde{E}))).$$

By Proposition 6.6, this is only possible if  $\text{ch}_2(\mathcal{H}^0(\tilde{E})) = \text{ch}_3(\mathcal{H}^0(\tilde{E})) = 0$ , ie  $\mathcal{H}^0(\tilde{E}) = 0$ .

Conversely, any slope-stable reflexive sheaf  $E$  of class  $v$  is  $\nu_{\alpha, \beta}$ -stable for  $\alpha \gg 0$  and  $\beta < \mu(E) = -\frac{1}{3}$ . Proposition 4.18 implies that  $E[1]$  is  $\nu_{\alpha, \beta}$ -stable for  $\alpha \gg 0$  and  $\beta > \mu(E) = -\frac{1}{3}$ . □

Next, we move down from the large volume limit and investigate walls for objects of class  $-v$ . Note that all walls to the right of the vertical wall must intersect  $\beta = -\frac{1}{3}$ .

**Lemma 6.8** *The wall  $W$  of equation (3) is the unique actual wall for objects with Chern character  $-v$  to the right of the vertical wall. There are no tilt-semistable objects below  $W$ . Any tilt-semistable  $\tilde{E}$  with Chern character  $-v$  fits into one of the following two cases:*

- (i)  $\tilde{E}$  fits into a short exact sequence

$$0 \rightarrow \mathcal{O}_Y(D) \rightarrow \tilde{E} \rightarrow \mathcal{O}_X^{\oplus 3}[1] \rightarrow 0,$$

where  $D$  is a Weil divisor on hyperplane section  $Y \in |H|$ .

(ii)  $\tilde{E}$  fits into a short exact sequence

$$0 \rightarrow \mathcal{I}_P(H) \rightarrow \tilde{E} \rightarrow \mathcal{O}_X^{\oplus 4}[1] \rightarrow 0,$$

where  $P \in X$ .

**Proof** Let  $\tilde{E}$  be a tilt-semistable object with Chern character  $-v$ . Let  $W'$  be a wall strictly above  $W$  induced by a short exact sequence  $0 \rightarrow F \rightarrow \tilde{E} \rightarrow G \rightarrow 0$ . Then the wall  $W'$  contains points  $(\alpha, 0)$  with  $\alpha > 0$ . In particular,  $0 < H \cdot \text{ch}_1(F) < H \cdot \text{ch}_1(\tilde{E}) = H^3$ , a contradiction.

Since the wall  $W(\mathcal{O}_X(2H), \tilde{E})$  is larger than  $W$ , we get  $\text{hom}(\tilde{E}, \mathcal{O}_X[3]) = \text{hom}(\mathcal{O}_X(2H), \tilde{E}) = 0$  and

$$\text{hom}(\tilde{E}, \mathcal{O}_X[1]) = \text{hom}(\tilde{E}, \mathcal{O}_X) + \text{ext}^2(\tilde{E}, \mathcal{O}_X) - \chi(\tilde{E}, \mathcal{O}_X) \geq -\chi(\tilde{E}, \mathcal{O}_X) = 3.$$

Clearly, any morphism  $\tilde{E} \rightarrow \mathcal{O}_X[1]$  destabilizes  $\tilde{E}$  below  $W$ .

Let  $r := \text{hom}(\tilde{E}, \mathcal{O}_X[1]) \geq 3$ . We get a short exact sequence of tilt-semistable objects along  $W$  given by

$$0 \rightarrow G \rightarrow \tilde{E} \rightarrow \mathcal{O}_X^{\oplus r}[1] \rightarrow 0.$$

If  $r \geq 4$ , then [Proposition 4.16](#) says

$$\frac{1}{4} \leq \frac{1}{r(r-3)},$$

ie  $r \leq 4$ . For  $r = 4$ , we get  $\text{ch}(G(-H)) = (1, 0, 0, -\frac{1}{3}H^3)$  and so  $G = \mathcal{I}_P(H)$  for some  $P \in X$ .

If  $r = 3$ , then  $\text{ch}(G) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ . Assume  $G$  is not of the form  $\mathcal{O}_Y(D)$  for some Weil divisor  $D$  on a hyperplane section  $Y \in |H|$ . Then [Proposition 6.2](#) implies that  $G$  has to be strictly semistable along our wall  $W$ . Since  $\tilde{E}$  is tilt-semistable above the wall, we know  $\text{Hom}(\mathcal{O}_X[1], E) = 0$ .

Therefore, [Lemma 6.3](#) shows that there is a short exact sequence

$$0 \rightarrow \mathcal{I}_P(H) \rightarrow G \rightarrow \mathcal{O}_X[1] \rightarrow 0$$

for a point  $P \in X$ . But then there is an inclusion  $\mathcal{I}_P(H) \hookrightarrow \tilde{E}$  and we are in the second case. □

**Proof of Theorem 6.1** Let  $D$  be a Weil divisor on a hyperplane section  $Y \in |H|$  with  $\text{ch}(\mathcal{O}_Y(D)) = (0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ . By [Proposition 6.2](#), the sheaf  $\mathcal{O}_Y(D)$  is tilt-stable for all  $\alpha > 0$  and  $\beta \in \mathbb{R}$ . A straightforward computation shows that the numerical wall  $W(\mathcal{O}_Y(D), \mathcal{O}_X(-2H)[1])$  is nonempty, and therefore,  $h^2(\mathcal{O}_Y(D)) = \text{hom}(\mathcal{O}_Y(D), \mathcal{O}_X(-2H)[1]) = 0$ . We conclude

$$h^0(\mathcal{O}_Y(D)) = \chi(\mathcal{O}_Y(D)) + h^1(\mathcal{O}_Y(D)) + h^3(\mathcal{O}_Y(D)) \geq \chi(\mathcal{O}_Y(D)) = 3.$$

We pick a three-dimensional subspace  $V \subset h^0(\mathcal{O}_Y(D))$  to get an object  $\mathcal{E}_{D,V} \in \text{D}^b(X)$  as in [Section 5](#). By [Lemma 5.1](#), the sheaf  $E_{D,V} = \mathcal{H}^{-1}(\mathcal{E}_{D,V})$  is slope-stable and reflexive. If  $\mathcal{H}^0(\mathcal{E}_{D,V}) \neq 0$ , then  $E_{D,V}$  has a Chern character in contradiction to [Proposition 6.6](#). This shows that  $\mathcal{O}_Y(D)$  is globally generated.

Since  $E_{D,V}$  is slope-stable, we know  $h^0(E_{D,V}) = 0$  and  $h^3(E_{D,V}) = \text{hom}(E_{D,V}, \mathcal{O}_X(-2H)) = 0$ . Moreover, as in the proof of [Proposition 6.6](#) we get  $h^2(E_{D,V}) = 0$ . This implies  $h^1(E_{D,V}) = -\chi(E_{D,V}) = 0$ .



The long exact sequence obtained from taking sheaf cohomology of

$$0 \rightarrow E_{D,V} \rightarrow \mathcal{O}_X \otimes V \rightarrow \mathcal{O}_Y(D) \rightarrow 0$$

implies  $H^i(\mathcal{O}_Y(D)) = 0$  for  $i > 0$  and  $h^0(\mathcal{O}_Y(D)) = 3$ . Therefore,  $V = H^0(\mathcal{O}_Y(D))$  and for each  $D$  there is a unique slope-stable sheaf  $E_D = E_{D,V}$ .

Let  $U \subset Y$  be the smooth locus of  $Y$ . By Proposition 3.1, we know that  $Y$  is normal, and therefore,  $Y \setminus U$  has dimension zero. In particular, a general section of  $\mathcal{O}_Y(D)$  leads to a curve completely contained in  $U$ . Since we work in characteristic 0, we can use a version of Bertini’s theorem [21, Corollary III.10.9, Remark III.10.9.1, Remark III.10.9.2] on the open subset  $U$  to see that a general section cuts out a smooth curve  $C$ . By adjunction,

$$\begin{aligned} \text{ch}(K_C) &= \text{ch}(\mathcal{O}_Y(-H + D)|_D) = \text{ch}(\mathcal{O}_Y(-H + D)) - \text{ch}(\mathcal{O}_Y(-H)) \\ &= (0, H, -\frac{1}{2}H^2, -\frac{1}{6}H^3) - (0, H, -\frac{3}{2}H^2, \frac{7}{6}H^3) = (0, 0, H^2, -\frac{4}{3}H^3), \end{aligned}$$

which shows that  $C$  is of degree 3 with  $\chi(K_C) = -1$ , ie a twisted cubic. This completes the proof of part (i).

For part (ii), we already showed in Corollary 5.2 that  $K_P$  is slope-stable for any  $P \in X$ . Conversely, if  $E$  is slope-stable, we can immediately conclude by Lemma 6.8. □

As a consequence we can already infer that our moduli space  $\overline{M}_X(v)$  is smooth.

**Corollary 6.9** *Every Gieseker-semistable sheaf  $E$  with  $\text{ch}(E) = (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$  satisfies*

$$\text{Ext}^i(E, E) = \begin{cases} \mathbb{C} & \text{if } i = 0, \\ \mathbb{C}^4 & \text{if } i = 1, \\ 0 & \text{otherwise.} \end{cases}$$

*In particular, the moduli space  $\overline{M}_X(v)$  is smooth and 4-dimensional.*

**Proof** Since  $(3, -H)$  is primitive, we know that  $E$  is slope-stable. Therefore,  $\text{hom}(E, E) = 1$ . Moreover, we must have  $\text{Ext}^3(E, E) = \text{Hom}(E, E(-2H))^\vee = 0$ . By Lemma 6.5, the sheaf  $E$  is  $\nu_{\alpha,-1}$ -stable for any  $\alpha > 0$ . Proposition 6.6 shows that  $E(-2H)$  is reflexive, so its shift  $E(-2H)[1]$  lies in the heart  $\text{Coh}^{\beta=-1}(X)$  and it is  $\nu_{\alpha,-1}$ -stable for any  $\alpha > 0$  by Lemma 6.8. Since

$$\nu_{0,-1}(E) = 0 > -\frac{1}{2} = \nu_{0,-1}(E(-2H)[1]),$$

we get  $\text{Ext}^2(E, E) = \text{Hom}(E, E(-2H)[1]) = 0$ . We can conclude that

$$\text{ext}^1(E, E) = \text{hom}(E, E) - \chi(E, E) = 4. \quad \square$$

## 7 Proof of the main theorem

Recall that  $\overline{M}_X(v)$  is the moduli space of Gieseker-semistable sheaves with Chern character

$$v := (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$$

and  $M_X(v) \subset \overline{M}_X(v)$  is the open locus of Gieseker-semistable vector bundles. The aim of this section is to prove the following theorem.

**Theorem 7.1** *The moduli space  $\overline{M}_X(v)$  is smooth and irreducible of dimension 4. Moreover, there is an Abel–Jacobi morphism  $\Psi: \overline{M}_X(v) \rightarrow J(X)$  sending  $E \mapsto \tilde{c}_2(E) - H^2$ , whose image is a theta divisor  $\Theta$  in the intermediate Jacobian  $J(X)$ . The theta divisor has a unique singular point, and  $\overline{M}_X(v)$  is the blowup of  $\Theta$  in this point. The exceptional divisor is isomorphic to the cubic threefold  $X$  itself.*

We have already shown that  $\overline{M}_X(v)$  is smooth of dimension 4 in Corollary 6.9. By Proposition 2.2, the image of  $\varphi: \overline{\mathcal{T}} \rightarrow J(X)$  is  $\Theta \subset J(X)$ , where  $\mathcal{T}$  is the open locus of smooth twisted cubics in the Hilbert scheme of  $X$ , and  $\overline{\mathcal{T}}$  is its closure. By Theorem 2.6, we know that  $\Theta$  is normal.

**Proposition 7.2** *There is a surjective map  $\varphi': \mathcal{T} \rightarrow M_X(v)$  that sends a twisted cubic  $C$  to the vector bundle  $E_C$ . The map  $\varphi|_{\mathcal{T}}: \mathcal{T} \rightarrow J(X)$  factors through  $\varphi'$ :*

$$\begin{array}{ccc}
 & \mathcal{T} & \\
 \varphi' \swarrow & & \searrow \varphi|_{\mathcal{T}} \\
 M_X(v) & \xrightarrow{\Psi|_{M_X(v)}} & J(X)
 \end{array}$$

Therefore, the image of  $\Psi: \overline{M}_X(v) \rightarrow J(X)$  is  $\Theta \subset J(X)$ .

**Proof** Let  $C$  be a twisted cubic in  $X$ . Then it lies in a unique hyperplane section  $Y$ . There is a short exact sequence

$$0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Y(C) \rightarrow T \rightarrow 0,$$

where  $T$  is a sheaf supported on  $C$  with rank one. Therefore,  $\widetilde{\text{ch}}_{\leq 2}(\mathcal{O}_Y(C)) = (0, H, C - \frac{1}{2}H^2)$  and we get  $\widetilde{\text{ch}}_{\leq 2}(E_C) = (3, -H, \frac{1}{2}H^2 - C)$ . It follows that  $\tilde{c}_2(E_C) = C$ . Thus, the composition  $\Psi|_{M_X(v)} \circ \varphi': \mathcal{T} \rightarrow M_X(v) \rightarrow J(X)$  is the Abel–Jacobi map  $\varphi: \overline{\mathcal{T}} \rightarrow J(X)$  restricted to  $\mathcal{T}$ . Surjectivity of  $\varphi'$  is a direct consequence of Theorem 6.1.  $\square$

**Lemma 7.3** *The morphism  $i: X \rightarrow \overline{M}_X(v)$  that maps  $P \mapsto K_P$  is an embedding with normal bundle  $\mathcal{O}_X(-H)$ .*

**Proof** We interpret  $X$  as the moduli spaces of twisted ideal sheaves  $\mathcal{I}_P(H)$  for all  $P \in X$ . By definition of  $K_P$ , we have a canonical short exact sequence

$$(7) \quad 0 \rightarrow K_P \rightarrow \mathcal{O}_X^{\oplus 4} \rightarrow \mathcal{I}_P(H) \rightarrow 0.$$

The appropriate version in families, considered below, induces the morphism  $i$ . It is injective, as  $P$  is the unique point where  $K_P$  is not locally free by Corollary 5.2.

Applying  $\text{Hom}(\cdot, K_P)$  to (7), we get an isomorphism  $\text{Ext}^1(K_P, K_P) \cong \text{Ext}^2(\mathcal{I}_P(H), K_P)$ . Next, we apply the functor  $\text{Hom}(\mathcal{I}_P(H), \cdot)$  to (7) to show that the induced morphism on tangent spaces  $\text{Ext}^1(\mathcal{I}_P(H), \mathcal{I}_P(H)) \hookrightarrow \text{Ext}^2(\mathcal{I}_P(H), K_P) = \text{Ext}^1(K_P, K_P)$  is an embedding. Since both  $X$  and  $\overline{M}_X(v)$  are smooth, the morphism is an embedding.

To determine the normal bundle, we need a relative version of the previous arguments to determine the cokernel of this embedding as a line bundle on  $X$ . The universal family inducing  $i$  is given by the sheaf  $\mathcal{K}$  on  $X \times X$  fitting into the short exact sequence

$$0 \rightarrow \mathcal{K} \rightarrow p^* \Omega_{\mathbb{P}^4}|_X(H) \rightarrow \mathcal{I}_\Delta(0, H) \rightarrow 0,$$

where  $p: X \times X \rightarrow X$  is the projection to the first factor. The pullback of the tangent bundle via  $i$  is  $i^* T_{\overline{M}_X(v)} = \mathcal{H}^1(p_* \mathcal{H}om(\mathcal{K}, \mathcal{K}))$ . Since  $p_* \mathcal{H}om(p^* \Omega_{\mathbb{P}^4}|_X(H), \mathcal{K}) = 0$ , we have an isomorphism

$$\mathcal{H}^1(p_* \mathcal{H}om(\mathcal{K}, \mathcal{K})) = \mathcal{H}^2(p_* \mathcal{H}om(\mathcal{I}_\Delta(0, H), \mathcal{K})).$$

The differential  $d_i$  of  $i$  fits into the four-term long exact sequence

$$\begin{aligned} 0 \rightarrow T_X = \mathcal{H}^1(p_* \mathcal{H}om(\mathcal{I}_\Delta(0, H), \mathcal{I}_\Delta(0, H))) &\xrightarrow{d_i} \mathcal{H}^2(p_* \mathcal{H}om(\mathcal{I}_\Delta(0, H), \mathcal{K})) \\ &\rightarrow \mathcal{H}^2(p_* \mathcal{H}om(\mathcal{I}_\Delta(0, H), p^* \Omega_{\mathbb{P}^4}|_X(H))) \rightarrow \mathcal{H}^2(p_* \text{Hom}(\mathcal{I}_\Delta(0, H), \mathcal{I}_\Delta(0, H))) \rightarrow 0. \end{aligned}$$

Using Grothendieck duality and the projection formula, the third term becomes

$$\Omega_{\mathbb{P}^4}|_X(H) \otimes \mathcal{H}^1(p_* \mathcal{I}_\Delta(0, -H))^\vee = \Omega_{\mathbb{P}^4}|_X(H) \otimes \mathcal{H}^0(p_* \mathcal{O}_\Delta(0, -H))^\vee = \Omega_{\mathbb{P}^4}|_X(2H).$$

A similar computation using the short exact sequence  $\mathcal{I}_\Delta \hookrightarrow \mathcal{O}_X \boxtimes \mathcal{O}_X \twoheadrightarrow \mathcal{O}_\Delta$  gives

$$\mathcal{H}^2(p_* \text{Hom}(\mathcal{I}_\Delta, \mathcal{I}_\Delta)) = \Omega_X(2H)$$

for the fourth term. Thus, the cokernel of  $d_i$  is isomorphic to  $\mathcal{N}_{X/\mathbb{P}^4}^\vee(2H) = \mathcal{O}_X(-H)$ , as claimed.  $\square$

**Lemma 7.4** *The morphism  $\Psi$  induces an isomorphism  $M_X(v) \rightarrow \Theta \setminus \{0\}$ . Moreover,  $\Psi$  contracts the irreducible divisor  $\overline{M}_X(v) \setminus M_X(v)$  to the zero point. In particular,  $\Theta$  is smooth away from 0.*

**Proof** By Lemma 5.1 and Corollary 5.2, the locus  $\overline{M}_X(v) \setminus M_X(v)$  coincides with vector bundles  $E_C$  associated to a twisted cubic  $C$ . By Lemma 2.5, the map  $\varphi|_{\mathcal{T}}$  has full rank four on tangent spaces. Thus, the commutative diagram in Proposition 7.2 implies that  $\Psi|_{M_X(v)}$  has full rank four on tangent spaces. Since  $M_X(v)$  is smooth of dimension four,  $\Psi|_{M_X(v)}$  must be injective on tangent spaces. In particular, the morphism  $\Psi|_{M_X(v)}$  must have finite fibers. Since  $\varphi|_{\mathcal{T}}$  has generically connected fibers by Proposition 2.2, the same holds for  $\Psi|_{M_X(v)}$ . Since  $\Theta$  is normal, Zariski’s main theorem implies that  $\Psi|_{M_X(v)}$  is an open embedding. Since  $\Theta$  is singular at the origin, we must have  $\Psi(M_X(v)) \subset \Theta - \{0\}$ .

By definition,  $\tilde{c}_2(K_P) = H^2$  and we get  $\Psi(K_P) = 0$ . Thus  $\Psi^{-1}(0) = \overline{M}_X(v) \setminus M_X(v)$ , and the image of  $M_X(v)$  is indeed  $\Theta \setminus \{0\}$  by Proposition 2.2.  $\square$

We can finish the proof of [Theorem 7.1](#) with the following lemma.

**Lemma 7.5** *The formal neighborhood of  $0 \in \Theta$  is isomorphic to the vertex of the affine cone over  $X \subset \mathbb{P}^4$ . Moreover, we have an isomorphism  $\overline{M}_X(v) = \text{Bl}_0(\Theta)$ . Thus,  $X$  is the union of all rational curves on  $\overline{M}_X(v)$ , and the unique divisor contracted by any morphism to a complex abelian variety.*

**Proof** The first two claims are scheme-theoretic enhancements of the set-theoretic statements in the previous lemma, which hold for any contraction of a divisor with ample conormal bundle to a point. We will only sketch the arguments.

Since the normal bundle of  $X \subset \overline{M}_X(v)$  is antiample, by Artin’s contractibility criterion [4, Corollary 6.12] there is a contraction  $\Psi': \overline{M}_X(v) \rightarrow N$  to an algebraic space  $N$  of finite type over  $\mathbb{C}$  that is an isomorphism away from  $X$ , and contracts  $X$  to a point  $0 \in N$ . Moreover, by Artin’s construction in [4, Theorem 6.2], the formal neighborhood of  $0 \in N$  is given by the affinization of the formal neighborhood of  $X \subset \overline{M}_X(v)$ . More precisely, if  $\mathcal{I}$  is the ideal of  $X$ , then it is given by

$$\text{Spec} \varprojlim_n H^0(X, \mathcal{O}_{\overline{M}_X(v)}/\mathcal{I}^{n+1}) = \text{Spec} \varprojlim_n \bigoplus_{0 \leq k \leq n} H^0(X, \mathcal{O}_X(k)),$$

ie the completion of the vertex of the affine cone over  $X$ . Since the image of every infinitesimal neighborhood of  $X$  under  $\Psi$  is affine, it factors via its affinization. Taking the limit, we see that  $\Psi$  factors via  $\Psi'$  both in the formal neighborhood of  $X$ , and in its complement. Hence (eg by [4, Theorem 3.1]) we get an induced morphism  $j: N \rightarrow \Theta$  factoring  $\Psi$ . As  $j$  is bijective on points and has normal target, it is an isomorphism.

For the last claim, note that  $X$  is uniruled, hence the union  $U$  of all rational curves in  $\overline{M}_X(v)$  contains  $X$ . If there was any other rational curve  $C$  not contained in  $X$ , then  $\Psi: C \rightarrow \Theta$  is a nonconstant map from a rational to an abelian variety, a contradiction. □

**Corollary 7.6** *If  $X_1$  and  $X_2$  are smooth projective threefolds with  $J(X_1) = J(X_2)$  as principally polarized abelian varieties, then  $X_1 = X_2$ .*

**Proof** As in the classical argument, this is an immediate consequence of the description of the singularity of the theta divisor in [Lemma 7.5](#). □

## 8 Kuznetsov component

The bounded derived category of a cubic threefold  $X$  admits a semiorthogonal decomposition

$$D^b(X) = \langle \text{Ku}(X), \mathcal{O}_X, \mathcal{O}_X(1) \rangle,$$

whose nontrivial part  $\text{Ku}(X)$  is called the Kuznetsov component. The goal of this section is to give a new proof of the following theorem.

**Theorem 8.1** *Let  $X_1$  and  $X_2$  be smooth cubic threefolds. Then  $\text{Ku}(X_1)$  and  $\text{Ku}(X_2)$  are equivalent as triangulated categories if and only if  $X_1$  and  $X_2$  are isomorphic.*

Let  $S$  be the Serre functor of  $\text{Ku}(X)$ . By [25, Lemmas 4.1 and 4.2], for any object  $F \in \text{Ku}(X)$ , we have

$$(8) \quad S(F) = L_{\mathcal{O}_X}(F \otimes \mathcal{O}_X(H))[1],$$

where  $L_{\mathcal{O}_X}$  is the left mutation functor with respect to  $\mathcal{O}_X$ . By [11, Proposition 2.7], the numerical Grothendieck group  $\mathcal{N}(\text{Ku}(X))$  is a two-dimensional lattice

$$\mathcal{N}(\text{Ku}(X)) \cong \mathbb{Z}^2 \cong \mathbb{Z}[\mathcal{I}_\ell] \oplus \mathbb{Z}[S(\mathcal{I}_\ell)],$$

where  $\mathcal{I}_\ell$  is the ideal sheaf of a line  $\ell$  in  $X$ . With respect to this basis, the Euler characteristic  $\chi(-, -)$  on  $\mathcal{N}(\text{Ku}(X))$  has the form

$$\begin{bmatrix} -1 & -1 \\ 0 & -1 \end{bmatrix}.$$

For any line  $\ell$  in  $X$ , we know  $\text{ch}(\mathcal{I}_\ell) = (1, 0, -\frac{1}{3}H^2, 0)$ . The Chern character of our second basis vector of  $\text{ch}(\text{Ku}(X))$ , and the action of the Serre functor  $S$  on our chosen basis are given as follows.

**Lemma 8.2** *We have  $\text{ch}(S(\mathcal{I}_\ell)) = (2, -H, -\frac{1}{6}H^2, \frac{1}{6}H^3)$  and  $\text{ch}(S^2(\mathcal{I}_\ell)) = (1, -H, \frac{1}{6}H^2, \frac{1}{6}H^3)$ . Thus, the class  $[S^2(\mathcal{I}_\ell)]$  in  $\mathcal{N}(\text{Ku}(X))$  is equal to  $[S(\mathcal{I}_\ell)] - [\mathcal{I}_\ell]$ .*

**Proof** By (8) we have  $[S(E)] = -[E(H)] + \chi(E(H))[\mathcal{O}_X]$  for  $E \in \text{Ku}(X)$ . Hence  $\text{ch}(I_\ell(H)) = (1, H, \frac{1}{6}H^2, -\frac{1}{6}H^3)$  and  $\chi(I_\ell(H)) = 3$  imply the formula for  $\text{ch}(S(\mathcal{I}_\ell))$ . The formula for  $\text{ch}(S^2(\mathcal{I}_\ell))$  follows from the last claim, which in turn follows from the Euler characteristic form above with

$$\begin{aligned} \chi(I_\ell, S^2(\mathcal{I}_\ell)) &= \chi(S^2(\mathcal{I}_\ell), S(\mathcal{I}_\ell)) = \chi(S(\mathcal{I}_\ell), I_\ell) = 0 = \chi([\mathcal{I}_\ell], [S(\mathcal{I}_\ell)] - [\mathcal{I}_\ell]), \\ \chi(S(\mathcal{I}_\ell), S^2(\mathcal{I}_\ell)) &= \chi(I_\ell, S(\mathcal{I}_\ell)) = -1 = \chi([S(\mathcal{I}_\ell)], [S(\mathcal{I}_\ell)] - [\mathcal{I}_\ell]). \end{aligned} \quad \square$$

For a point  $P \in X$ , the sheaf  $K_P$ , which is defined through the sequence (2), lies in the Kuznetsov component  $\text{Ku}(X)$ .

**Lemma 8.3** *Let  $[A]$  be a class in  $\mathcal{N}(\text{Ku}(X))$  such that  $\chi([A], [A]) = -3$ . Then, up to a sign,  $[A]$  is either  $[K_P] = [\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]$ , or  $[S(K_P)] = -[\mathcal{I}_\ell] + 2[S(\mathcal{I}_\ell)]$ , or  $[S^2(K_P)] = -2[\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]$ .*

Let  $\sigma_{\alpha, -1/2}^0 = (\text{Coh}_{\alpha, -1/2}^0(X), Z_{\alpha, -1/2}^0)$  be the weak stability condition on  $\text{D}^b(X)$  constructed in [5, Proposition 2.14]. Here  $\text{Coh}_{\alpha, -1/2}^0(X)$  is the usual double tilt and

$$(9) \quad Z_{\alpha, -1/2}^0(E) = H^2 \cdot \text{ch}_1^{-1/2}(E) + i(H \cdot \text{ch}_2^{-1/2}(E) - \frac{1}{2}\alpha^2 H^3 \cdot \text{ch}_0(E)).$$

As proven in [5, Theorem 6.8], for  $0 < \alpha \ll 1$  it induces the stability condition  $\sigma(\alpha) = (\mathcal{A}(\alpha), Z(\alpha))$  on  $\text{Ku}(X)$ , where

$$\mathcal{A}(\alpha) := \text{Coh}_{\alpha, -1/2}^0(X) \cap \text{Ku}(X) \quad \text{and} \quad Z(\alpha) := Z_{\alpha, -1/2}^0|_{\text{Ku}(X)}.$$

**Lemma 8.4** *There is an embedding  $M_X(v) \hookrightarrow M_{\sigma(\alpha)}([\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)])$  from the moduli space  $M_X(v)$  for  $v = \text{ch}(\mathcal{I}_\ell) + \text{ch}(S(\mathcal{I}_\ell)) = (3, -H, -\frac{1}{2}H^2, \frac{1}{6}H^3)$  to  $M_{\sigma(\alpha)}([\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)])$ , which parametrizes  $\sigma(\alpha)$ -semistable objects in  $\text{Ku}(X)$  of class  $[\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)] \in \mathcal{N}(\text{Ku}(X))$ .*

**Proof** According to Lemma 6.5 there is no wall for objects of Chern character  $v$  to the left of the vertical wall. Thus,  $E$  is  $\nu_{\alpha,-1/2}$ -stable for any  $\alpha > 0$ . Since  $\sigma_{\alpha,-1/2}^0$  is just a rotation of  $\nu_{\alpha,-1/2}$ , we obtain that  $E$  is  $\sigma_{\alpha,-1/2}^0$ -stable. By Theorem 6.1(ii), the sheaf  $E \in \text{Ku}(X)$  lies in the Kuznetsov component. Thus,  $E$  is  $\sigma(\alpha)$ -stable. Note that the object  $E$  could be destabilized by objects with  $Z_{\alpha,-1/2}^0 = 0$  after rotation. But we know that these are all sheaves supported in dimension zero and would not be in  $\text{Ku}(X)$  and therefore,  $E$  is stable after restriction to  $\text{Ku}(X)$ .  $\square$

Corollary 5.6 of [35] implies that the stability condition  $\sigma(\alpha)$  is  $S$ -invariant, ie  $S \cdot \sigma(\alpha) = \sigma(\alpha) \cdot \tilde{g}$  for  $\tilde{g} \in \widetilde{\text{GL}}^+(2, \mathbb{R})$ . Thus, there is an isomorphism

$$(10) \quad S : M_{\sigma(\alpha)}(2[\mathcal{I}_\ell] - [S(\mathcal{I}_\ell)]) \rightarrow M_{\sigma(\alpha)}([\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]), \quad E \mapsto S(E).$$

The following proposition is a slight strengthening of [1, Theorem 1.2], which describes all elements of the moduli space. The idea of the proof is the same as [1, Lemma 2.2].

**Proposition 8.5** *Any  $\sigma(\alpha)$ -semistable object in  $\text{Ku}(X)$  of class  $2[\mathcal{I}_\ell] - [S(\mathcal{I}_\ell)]$  is of the form  $G[2k]$  for  $k \in \mathbb{Z}$ , where  $G$  is either equal to  $G_P(-H)$  described in (4) for a point  $P \in X$ , or  $\mathcal{O}_Y(D - H)$ , where  $D$  is a Weil divisor on some  $Y \in |H|$ .*

**Proof** Lemma 8.2 implies  $\text{ch}(G) = (0, H, -\frac{1}{2}H^2, -\frac{1}{6}H^3)$ . Since  $G$  is  $\sigma(\alpha)$ -semistable, its shift  $G[2k]$  lies in the heart  $\mathcal{A}(\alpha)$  for some  $k \in \mathbb{Z}$ . We know its image under the stability function  $Z(\alpha)$  is equal to  $-H^3$ , so it has maximum phase in the heart  $\mathcal{A}(\alpha)$ , which immediately implies  $G[2k]$  is  $\sigma_{\alpha,-1/2}^0$ -semistable. We claim that  $G[2k]$  has no subobject  $Q \in \text{Coh}_{\alpha,-1/2}^0$  with  $Z_{\alpha,-1/2}^0(Q) = 0$ , so it is  $\nu_{\alpha,-1/2}$ -semistable. Assume for a contradiction that there is such a subobject  $Q$ . By the definition of  $\text{Coh}_{\alpha,-1/2}^0(X)$ , it is a sheaf supported in dimension zero. Thus,  $\text{hom}(\mathcal{O}_X, Q) \neq 0$ . Since  $\mathcal{O}_X \in \text{Coh}_{\alpha,-1/2}^0(X)$ , we have  $\text{hom}(\mathcal{O}_X, (G[2k]/Q)[-1]) = 0$ . Therefore,  $\text{hom}(\mathcal{O}_X, G[2k]) \neq 0$ , which is not possible because  $G[2k] \in \text{Ku}(X)$ . Finally, since  $G[2k]$  is  $\nu_{\alpha,-1/2}$ -semistable for  $0 < \alpha \ll 1$ , the claim follows by Proposition 6.2(ii).  $\square$

**Remark 8.6** Since the class  $2[\mathcal{I}_\ell] - [S(\mathcal{I}_\ell)]$  is primitive in  $\mathcal{N}(\text{Ku}(X))$ , any  $\sigma(\alpha)$ -semistable object of this class is  $\sigma(\alpha)$ -stable if we choose  $\alpha$  sufficiently small.

We now describe the image of the semistable objects  $G \in M_{\sigma(\alpha)}(2[\mathcal{I}_\ell] - [S(\mathcal{I}_\ell)])$  under the Serre functor  $S$ . If  $G = G_P(-H)$ , then by (4), we know there is a distinguished triangle

$$\mathcal{O}_X[1] \rightarrow G_P \rightarrow \mathcal{I}_P(H) \rightarrow \mathcal{O}_X[2],$$

which gives  $L_{\mathcal{O}_X}(G_P) = L_{\mathcal{O}_X}(\mathcal{I}_P(H)) = K_P[1]$ , so

$$(11) \quad S(G_P) = K_P[2].$$

If  $G = \mathcal{O}_Y(D - H)$ , then  $G(H) = \mathcal{O}_Y(D)$  is of class  $(0, H, \frac{1}{2}H^2, -\frac{1}{6}H^3)$ , and lies in a distinguished triangle

$$\mathcal{O}_X^{\oplus 3} \rightarrow \mathcal{O}_Y(D) \rightarrow E_D[1] \rightarrow \mathcal{O}_X^{\oplus 3}[1].$$

Thus,

$$(12) \quad S(G) = L_{\mathcal{O}_X}(\mathcal{O}_Y(D))[1] = L_{\mathcal{O}_X}(E_D[1])[1] = E_D[2].$$

Combining (11) and (12) with Lemma 8.4 implies the next result.

**Theorem 8.7** *The moduli space  $M_{\sigma(\alpha)}([\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)])$  is isomorphic to the moduli space  $\overline{M}_X(v)$  parametrizing Gieseker-stable sheaves of class  $v$ .*

The next step is to show that we can replace  $\sigma(\alpha)$  by any  $S$ -invariant stability condition on  $\text{Ku}(X)$ .

**Lemma 8.8** [35, Lemmas 5.8 and 5.10] *Let  $\sigma$  be an  $S$ -invariant stability condition on  $\text{Ku}(X)$  and  $F \in \text{Ku}(X)$  be  $\sigma$ -semistable of phase  $\varphi(F)$ . Then*

- (i)  $\varphi(F) < \varphi(S(F)) < \varphi(F) + 2$ ,
- (ii)  $\dim \text{Ext}^1(F, F) \geq 2$ .

For cubic threefolds, we also have a weak version of the Mukai lemma for K3 surfaces.

**Lemma 8.9** (weak Mukai lemma [35, Lemma 5.11]) *Let  $\sigma$  be an  $S$ -invariant stability condition. Let  $A \rightarrow E \rightarrow B$  be a triangle in  $\text{Ku}(X)$  such that  $\text{hom}(A, B) = 0$  and the  $\sigma$ -semistable factors of  $A$  have phase greater than or equal to the phase of the  $\sigma$ -semistable factors of  $B$ . Then*

$$\dim_{\mathbb{C}} \text{Ext}^1(A, A) + \dim_{\mathbb{C}} \text{Ext}^1(B, B) \leq \dim_{\mathbb{C}} \text{Ext}^1(E, E).$$

**Proposition 8.10** *Let  $\sigma_1$  and  $\sigma_2$  be two  $S$ -invariant stability conditions on  $\text{Ku}(X)$ . An object  $E \in \text{Ku}(X)$  of class  $[\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]$  is  $\sigma_1$ -stable if and only if it is  $\sigma_2$ -stable.*

**Proof** By [35, Proposition 4.6],  $\mathcal{I}_\ell$  and  $S(\mathcal{I}_\ell)$  are  $\sigma$ -stable with respect to any  $S$ -invariant stability condition. Thus, Lemma 8.8 implies that

$$(13) \quad \varphi_{\sigma}(\mathcal{I}_\ell) < \varphi_{\sigma}(S(\mathcal{I}_\ell)) < \varphi_{\sigma}(\mathcal{I}_\ell) + 2.$$

Take a  $\sigma_1$ -stable object  $E \in \text{Ku}(X)$  of class  $[\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]$ . Since  $\sigma_1$  is  $S$ -invariant, Lemma 8.8 gives

$$\varphi_{\sigma_1}(E) < \varphi_{\sigma_1}(S(E)) < \varphi_{\sigma_1}(E) + 2.$$

Thus, for  $i < 0$  or  $i \geq 2$ , we get

$$\text{hom}(E, E[i]) = \text{hom}(E[i], S(E)) = 0.$$

Since  $E$  is  $\sigma_1$ -stable, we get  $\text{hom}(E, E) = 1$ , which gives

$$\text{hom}(E, E[1]) = -\chi(E, E) + 1 = 4.$$

Suppose now for a contradiction that  $E$  is  $\sigma_2$ -unstable. There is a distinguished triangle of destabilizing objects  $F_1 \rightarrow E \rightarrow F_2 \rightarrow F_1[1]$  with respect to  $\sigma_2$ . We may assume  $F_1$  is  $\sigma_2$ -semistable. Thus, [Lemma 8.8](#) implies that

$$(14) \quad \text{hom}(F_1, F_1[1]) \geq 2.$$

Since the phase of  $F_1$  is bigger than the phase of  $\sigma_2$ -semistable factors of  $F_2$ , we have

$$(15) \quad \text{hom}(F_1, F_2) = 0.$$

Thus, the weak Mukai lemma ([Lemma 8.9](#)) implies

$$\text{hom}(F_1, F_1[1]) + \text{hom}(F_2, F_2[1]) \leq \text{hom}(E, E[1]) = 4.$$

By (14), we get  $\text{hom}(F_2, F_2[1]) \leq 2$ . If  $\text{hom}(F_2, F_2[1]) = 0$  or  $1$ , then all its  $\sigma_2$ -semistable factors would satisfy the same property by the weak Mukai lemma ([Lemma 8.9](#)), which is not possible by [Lemma 8.8](#). Therefore,

$$\text{hom}(F_1, F_1[2]) = \text{hom}(F_2, F_2[1]) = 2,$$

and [\[35, Lemma 5.12\]](#) implies that  $F_1$  and  $F_2$  are  $\sigma_2$ -stable. This gives  $\chi(F_i, F_i) = -1$  for  $i = 1, 2$ , so  $[F_i]$  is either  $\pm[\mathcal{I}_\ell]$ , or  $\pm[S(\mathcal{I}_\ell)]$ , or  $\pm([S(\mathcal{I}_\ell)] - [\mathcal{I}_\ell])$ . Since there are only 2 stable factors and the object  $E$  is of class  $[\mathcal{I}_\ell] + [S(\mathcal{I}_\ell)]$ , the destabilizing objects must be of class  $[\mathcal{I}_\ell]$  and  $[S(\mathcal{I}_\ell)]$ . Thus, [\[35, Proposition 4.6\]](#) implies that the destabilizing objects are  $\mathcal{I}_\ell[2k]$  and  $S(\mathcal{I}_{\ell'})[2k']$  for two lines  $\ell, \ell'$  and integers  $k, k' \in \mathbb{Z}$ .

Let  $F_1 = \mathcal{I}_\ell[2k]$  and  $F_2 = S(\mathcal{I}_{\ell'})[2k']$ . Since  $E$  is  $\sigma_1$ -stable, we have  $\varphi_{\sigma_1}(F_1) < \varphi_{\sigma_1}(F_2)$ , thus (13) gives  $k \leq k'$ . But  $F_1$  and  $F_2$  are the destabilizing objects with respect to  $\sigma_2$ , hence  $\varphi_{\sigma_2}(F_1) > \varphi_{\sigma_2}(F_2)$  and (13) gives  $k' + 1 \leq k$ , which is not possible. By a similar argument, we reach a contradiction if  $F_1 = S(\mathcal{I}_{\ell'})[2k']$  and  $F_2 = \mathcal{I}_\ell[2k]$ . Finally, note that  $E$  cannot be strictly  $\sigma_2$ -semistable because the phases of  $\mathcal{I}_\ell[2k]$  and  $S(\mathcal{I}_{\ell'})[2k']$  cannot be equal, by (13).  $\square$

**Proof of Theorem 8.1** As a cubic threefold has free Picard group of rank one, the first implication is obvious. As for the second implication, assume there is an exact equivalence  $\Phi: \text{Ku}(X_1) \rightarrow \text{Ku}(X_2)$ . [Lemma 8.3](#) implies that, up to composing with a power of the Serre functor of  $\text{Ku}(X_1)$  and shift functor, we may assume  $[\Phi_*(K_P)] = [K_{P'}]$  for points  $P$  and  $P'$  in  $X_1$  and  $X_2$ , respectively. Take an  $S$ -invariant stability condition  $\sigma$  on  $\text{Ku}(X_1)$ . [Theorem 8.7](#) and [Proposition 8.10](#) imply that

$$(16) \quad M_{X_1}(v) \cong M_\sigma(\text{Ku}(X_1), [K_P]) \cong M_{\varphi \cdot \sigma}(\text{Ku}(X_2), [K_{P'}]).$$

Since the Serre functor commutes with autoequivalences,  $\varphi \cdot \sigma$  is an  $S$ -invariant stability condition on  $\text{Ku}(X_2)$ . Thus, [Theorem 8.7](#) gives

$$M_{\varphi \cdot \sigma}(\text{Ku}(X_2), [K_{P'}]) \cong M_{X_2}(v).$$



Combining this with (16) gives  $M_{X_1}(v) \cong M_{X_2}(v)$ . By Lemma 7.5, we know  $X_1$  and  $X_2$  are the unique exceptional divisors of  $M_{X_1}(v)$  and  $M_{X_2}(v)$  which get contracted by any map to a complex abelian variety. Thus,  $X_1 \cong X_2$ .  $\square$

## List of symbols

$X$	smooth cubic threefold in $\mathbb{P}^4$ over $\mathbb{C}$
$H$	the ample generator of $\text{Pic}(X)$
$Y$	a hyperplane section of $X$
$D^b(X)$	bounded derived category of coherent sheaves on $X$
$\text{Ku}(X)$	the Kuznetsov component inside $D^b(X)$
$\text{CH}^*(X)$	the Chow ring of $X$
$\text{CH}_n^*(X)$	the numerical Chow ring of $X$ , obtained as $\text{CH}^*(X)$ modulo numerical equivalence
$\mathcal{H}^i(E)$	the $i^{\text{th}}$ cohomology sheaf of a complex $E \in D^b(X)$
$H^i(E)$	the $i^{\text{th}}$ sheaf cohomology group of a complex $E \in D^b(X)$
$\text{ch}(E)$	total Chern character of an object $E \in D^b(X)$ up to numerical equivalence
$\widetilde{c}(E)$	total Chern class of an object $E \in D^b(X)$ up to numerical equivalence
$\widetilde{\text{ch}}(E)$	total Chern character of an object $E \in D^b(X)$ up to rational equivalence
$\widetilde{c}(E)$	total Chern class of an object $E \in D^b(X)$ up to rational equivalence
$\text{ch}_{\leq l}(E)$	$(\text{ch}_0(E), \dots, \text{ch}_l(E))$
$\widetilde{\text{ch}}_{\leq l}(E)$	$(\widetilde{\text{ch}}_0(E), \dots, \widetilde{\text{ch}}_l(E))$

## References

- [1] M Altavilla, M Petković, F Rota, *Moduli spaces on the Kuznetsov component of Fano threefolds of index 2*, *Épjournal Géom. Algébrique* 6 (2022) art. id. 13 MR Zbl
- [2] D Arcara, A Bertram, *Bridgeland-stable moduli spaces for  $K$ -trivial surfaces*, *J. Eur. Math. Soc.* 15 (2013) 1–38 MR Zbl
- [3] M Artebani, R Kloosterman, M Pacini, *A new model for the theta divisor of the cubic threefold*, *Matematiche (Catania)* 58 (2003) 201–236 MR Zbl
- [4] M Artin, *Algebraization of formal moduli, II: Existence of modifications*, *Ann. of Math.* 91 (1970) 88–135 MR Zbl
- [5] A Bayer, M Lahoz, E Macrì, P Stellari, *Stability conditions on Kuznetsov components*, *Ann. Sci. Éc. Norm. Supér.* 56 (2023) 517–570 MR Zbl
- [6] A Bayer, E Macrì, P Stellari, *The space of stability conditions on abelian threefolds, and on some Calabi–Yau threefolds*, *Invent. Math.* 206 (2016) 869–933 MR Zbl
- [7] A Bayer, E Macrì, Y Toda, *Bridgeland stability conditions on threefolds, I: Bogomolov–Gieseker type inequalities*, *J. Algebraic Geom.* 23 (2014) 117–163 MR Zbl
- [8] A Beauville, *Les singularités du diviseur  $\Theta$  de la Jacobienne intermédiaire de l’hypersurface cubique dans  $\mathbb{P}^4$* , from “Algebraic threefolds” (A Conte, editor), *Lecture Notes in Math.* 947, Springer (1982) 190–208 MR Zbl

- [9] **A Beauville**, *Vector bundles on the cubic threefold*, from “Symposium in honor of CH Clemens” (A Bertram, J A Carlson, H Kley, editors), Contemp. Math. 312, Amer. Math. Soc., Providence, RI (2002) 71–86 [MR](#) [Zbl](#)
- [10] **P Belmans, A J de Jong**, et al., *The Stacks project*, electronic reference (2005–) Available at <http://stacks.math.columbia.edu>
- [11] **M Bernardara, E Macrì, S Mehrotra, P Stellari**, *A categorical invariant for cubic threefolds*, Adv. Math. 229 (2012) 770–803 [MR](#) [Zbl](#)
- [12] **T Bridgeland**, *Stability conditions on triangulated categories*, Ann. of Math. 166 (2007) 317–345 [MR](#) [Zbl](#)
- [13] **T Bridgeland**, *Stability conditions on K3 surfaces*, Duke Math. J. 141 (2008) 241–291 [MR](#) [Zbl](#)
- [14] **CH Clemens, PA Griffiths**, *The intermediate Jacobian of the cubic threefold*, Ann. of Math. 95 (1972) 281–356 [MR](#) [Zbl](#)
- [15] **I Coskun, J Huizenga**, *The ample cone of moduli spaces of sheaves on the plane*, Algebr. Geom. 3 (2016) 106–136 [MR](#) [Zbl](#)
- [16] **L Ein, R Lazarsfeld**, *Singularities of theta divisors and the birational geometry of irregular varieties*, J. Amer. Math. Soc. 10 (1997) 243–258 [MR](#) [Zbl](#)
- [17] **S Feyzbakhsh, L Pertusi**, *Serre-invariant stability conditions and Ulrich bundles on cubic threefolds*, Épijournal Géom. Algébrique 7 (2023) art. id. 1 [MR](#) [Zbl](#)
- [18] **D Gieseker**, *On the moduli of vector bundles on an algebraic surface*, Ann. of Math. 106 (1977) 45–60 [MR](#) [Zbl](#)
- [19] **PA Griffiths**, *Periods of integrals on algebraic manifolds, II: Local study of the period mapping*, Amer. J. Math. 90 (1968) 805–865 [MR](#) [Zbl](#)
- [20] **D Happel, I Reiten, SO Smalø**, *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. 575, Amer. Math. Soc., Providence, RI (1996) [MR](#) [Zbl](#)
- [21] **R Hartshorne**, *Algebraic geometry*, Graduate Texts in Math. 52, Springer (1977) [MR](#) [Zbl](#)
- [22] **R Hartshorne**, *Generalized divisors on Gorenstein schemes, K–Theory* 8 (1994) 287–339 [MR](#) [Zbl](#)
- [23] **D Huybrechts, M Lehn**, *The geometry of moduli spaces of sheaves*, 2nd edition, Cambridge Univ. Press (2010) [MR](#) [Zbl](#)
- [24] **A Iliev**, *Minimal sections of conic bundles*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. 2 (1999) 401–428 [MR](#) [Zbl](#)
- [25] **AG Kuznetsov**, *Derived category of a cubic threefold and the variety  $V_{14}$* , Tr. Mat. Inst. Steklova 246 (2004) 183–207 [MR](#) In Russian: translated in Proc. Steklov Inst. Math. 246 (2004) 171–194
- [26] **C Li**, *Stability conditions on Fano threefolds of Picard number 1*, J. Eur. Math. Soc. 21 (2019) 709–726 [MR](#) [Zbl](#)
- [27] **J Lo, Y More**, *Some examples of tilt-stable objects on threefolds*, Comm. Algebra 44 (2016) 1280–1301 [MR](#) [Zbl](#)
- [28] **A Maciocia**, *Computing the walls associated to Bridgeland stability conditions on projective surfaces*, Asian J. Math. 18 (2014) 263–279 [MR](#) [Zbl](#)
- [29] **E Macrì, B Schmidt**, *Derived categories and the genus of space curves*, Algebr. Geom. 7 (2020) 153–191 [MR](#) [Zbl](#)

- [30] **M Maruyama**, *Moduli of stable sheaves. I*, J. Math. Kyoto Univ. 17 (1977) 91–126 [MR](#) [Zbl](#)
- [31] **M Maruyama**, *Moduli of stable sheaves. II*, J. Math. Kyoto Univ. 18 (1978) 557–614 [MR](#) [Zbl](#)
- [32] **D Mumford**, *Projective invariants of projective structures and applications*, from “Proceedings of the International Congress of Mathematicians” (V Stenström, editor), Institut Mittag-Leffler, Djursholm (1963) 526–530 [MR](#) [Zbl](#)
- [33] **D Mumford**, *Prym varieties, I*, from “Contributions to analysis: a collection of papers dedicated to Lipman Bers” (L V Ahlfors, I Kra, B Maskit, L Nirenberg, editors), Academic (1974) 325–350 [MR](#) [Zbl](#)
- [34] **A Perry**, *The integral Hodge conjecture for two-dimensional Calabi–Yau categories*, Compos. Math. 158 (2022) 287–333 [MR](#) [Zbl](#)
- [35] **L Pertusi, S Yang**, *Some remarks on Fano threefolds of index two and stability conditions*, Int. Math. Res. Not. 2022 (2022) 12940–12983 [MR](#) [Zbl](#)
- [36] **K Schwede**, *Generalized divisors and reflexive sheaves*, lecture notes (2007) Available at <https://www.math.utah.edu/~schwede/Notes/GeneralizedDivisors.pdf>
- [37] **C T Simpson**, *Moduli of representations of the fundamental group of a smooth projective variety, I*, Inst. Hautes Études Sci. Publ. Math. 79 (1994) 47–129 [MR](#) [Zbl](#)
- [38] **F Takemoto**, *Stable vector bundles on algebraic surfaces*, Nagoya Math. J. 47 (1972) 29–48 [MR](#) [Zbl](#)
- [39] **G E Welters**, *Abel–Jacobi isogenies for certain types of Fano threefolds*, Mathematical Centre Tracts 141, Mathematisch Centrum, Amsterdam (1981) [MR](#) [Zbl](#)
- [40] **S Zhang**, *Bridgeland moduli spaces for Gushel–Mukai threefolds and Kuznetsov’s Fano threefold conjecture*, preprint (2021) [arXiv 2012.12193v2](https://arxiv.org/abs/2012.12193v2)

AB, SVB: *School of Mathematics and Maxwell Institute, University of Edinburgh  
Edinburgh, United Kingdom*

SF: *Department of Mathematics, Imperial College London  
London, United Kingdom*

GH: *Fakultät für Mathematik, Universität Duisburg-Essen  
Essen, Germany*

DM: *Korteweg-de Vries Institute for Mathematics, University of Amsterdam  
Amsterdam, Netherlands*

FR: *Centre for Mathematical Sciences, University of Cambridge  
Cambridge, United Kingdom*

FR: *ETH Zürich  
Zürich, Switzerland*

BS: *Institut für Algebraische Geometrie, Gottfried Wilhelm Leibniz Universität Hannover  
Hannover, Germany*

[arend.bayer@ed.ac.uk](mailto:arend.bayer@ed.ac.uk), [sjoerd.beentjes@ed.ac.uk](mailto:sjoerd.beentjes@ed.ac.uk), [s.feyzbakhsh@imperial.ac.uk](mailto:s.feyzbakhsh@imperial.ac.uk),  
[georg.hein@uni-due.de](mailto:georg.hein@uni-due.de), [d.martinelli@uva.nl](mailto:d.martinelli@uva.nl), [fr414@cam.ac.uk](mailto:fr414@cam.ac.uk),  
[bschmidt@math.uni-hannover.de](mailto:bschmidt@math.uni-hannover.de)

Proposed: Mark Gross

Seconded: Richard P Thomas, Jim Bryan

Received: 21 January 2021

Revised: 19 February 2022

# GEOMETRY & TOPOLOGY

[msp.org/gt](http://msp.org/gt)

## MANAGING EDITOR

András I Stipsicz    Alfréd Rényi Institute of Mathematics  
[stipsicz@renyi.hu](mailto:stipsicz@renyi.hu)

## BOARD OF EDITORS

Mohammed Abouzaid	Stanford University <a href="mailto:abouzaid@stanford.edu">abouzaid@stanford.edu</a>	Mark Gross	University of Cambridge <a href="mailto:mgross@dpms.cam.ac.uk">mgross@dpms.cam.ac.uk</a>
Dan Abramovich	Brown University <a href="mailto:dan_abramovich@brown.edu">dan_abramovich@brown.edu</a>	Rob Kirby	University of California, Berkeley <a href="mailto:kirby@math.berkeley.edu">kirby@math.berkeley.edu</a>
Ian Agol	University of California, Berkeley <a href="mailto:ianagol@math.berkeley.edu">ianagol@math.berkeley.edu</a>	Bruce Kleiner	NYU, Courant Institute <a href="mailto:bkleiner@cims.nyu.edu">bkleiner@cims.nyu.edu</a>
Arend Bayer	University of Edinburgh <a href="mailto:arend.bayer@ed.ac.uk">arend.bayer@ed.ac.uk</a>	Sándor Kovács	University of Washington <a href="mailto:skovacs@uw.edu">skovacs@uw.edu</a>
Mark Behrens	University of Notre Dame <a href="mailto:mbehrens1@nd.edu">mbehrens1@nd.edu</a>	Urs Lang	ETH Zürich <a href="mailto:urs.lang@math.ethz.ch">urs.lang@math.ethz.ch</a>
Mladen Bestvina	University of Utah <a href="mailto:bestvina@math.utah.edu">bestvina@math.utah.edu</a>	Marc Levine	Universität Duisburg-Essen <a href="mailto:marc.levine@uni-due.de">marc.levine@uni-due.de</a>
Martin R Bridson	University of Oxford <a href="mailto:bridson@maths.ox.ac.uk">bridson@maths.ox.ac.uk</a>	Ciprian Manolescu	University of California, Los Angeles <a href="mailto:cm@math.ucla.edu">cm@math.ucla.edu</a>
Jim Bryan	University of British Columbia <a href="mailto:jbryan@math.ubc.ca">jbryan@math.ubc.ca</a>	Haynes Miller	Massachusetts Institute of Technology <a href="mailto:hmr@math.mit.edu">hmr@math.mit.edu</a>
Dmitri Burago	Pennsylvania State University <a href="mailto:burago@math.psu.edu">burago@math.psu.edu</a>	Tomasz Mrowka	Massachusetts Institute of Technology <a href="mailto:mrowka@math.mit.edu">mrowka@math.mit.edu</a>
Tobias H Colding	Massachusetts Institute of Technology <a href="mailto:colding@math.mit.edu">colding@math.mit.edu</a>	Aaron Naber	Northwestern University <a href="mailto:anaber@math.northwestern.edu">anaber@math.northwestern.edu</a>
Simon Donaldson	Imperial College, London <a href="mailto:s.donaldson@ic.ac.uk">s.donaldson@ic.ac.uk</a>	Peter Ozsváth	Princeton University <a href="mailto:petero@math.princeton.edu">petero@math.princeton.edu</a>
Yasha Eliashberg	Stanford University <a href="mailto:eliash-gt@math.stanford.edu">eliash-gt@math.stanford.edu</a>	Leonid Polterovich	Tel Aviv University <a href="mailto:polterov@post.tau.ac.il">polterov@post.tau.ac.il</a>
Benson Farb	University of Chicago <a href="mailto:farb@math.uchicago.edu">farb@math.uchicago.edu</a>	Colin Rourke	University of Warwick <a href="mailto:gt@maths.warwick.ac.uk">gt@maths.warwick.ac.uk</a>
David M Fisher	Rice University <a href="mailto:davidfisher@rice.edu">davidfisher@rice.edu</a>	Roman Sauer	Karlsruhe Institute of Technology <a href="mailto:roman.sauer@kit.edu">roman.sauer@kit.edu</a>
Mike Freedman	Microsoft Research <a href="mailto:michaelf@microsoft.com">michaelf@microsoft.com</a>	Stefan Schwede	Universität Bonn <a href="mailto:schwede@math.uni-bonn.de">schwede@math.uni-bonn.de</a>
David Gabai	Princeton University <a href="mailto:gabai@princeton.edu">gabai@princeton.edu</a>	Natasa Sesum	Rutgers University <a href="mailto:natasas@math.rutgers.edu">natasas@math.rutgers.edu</a>
Stavros Garoufalidis	Southern U. of Sci. and Tech., China <a href="mailto:stavros@mpim-bonn.mpg.de">stavros@mpim-bonn.mpg.de</a>	Gang Tian	Massachusetts Institute of Technology <a href="mailto:tian@math.mit.edu">tian@math.mit.edu</a>
Cameron Gordon	University of Texas <a href="mailto:gordon@math.utexas.edu">gordon@math.utexas.edu</a>	Ulrike Tillmann	Oxford University <a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>
Jesper Grodal	University of Copenhagen <a href="mailto:jg@math.ku.dk">jg@math.ku.dk</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Misha Gromov	IHÉS and NYU, Courant Institute <a href="mailto:gromov@ihes.fr">gromov@ihes.fr</a>	Anna Wienhard	Universität Heidelberg <a href="mailto:wienhard@mathi.uni-heidelberg.de">wienhard@mathi.uni-heidelberg.de</a>

---

See inside back cover or [msp.org/gt](http://msp.org/gt) for submission instructions.

---

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

---

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# GEOMETRY & TOPOLOGY

Volume 28

Issue 1 (pages 1–496)

2024

---

Homological invariants of codimension 2 contact submanifolds	1
LAURENT CÔTÉ and FRANÇOIS-SIMON FAUTEUX-CHAPLEAU	
The desingularization of the theta divisor of a cubic threefold as a moduli space	127
AREND BAYER, SJOERD VIKTOR BEENTJES, SOHEYLA FEYZBAKSH, GEORG HEIN, DILETTA MARTINELLI, FATEMEH REZAAE and BENJAMIN SCHMIDT	
Coarse-median preserving automorphisms	161
ELIA FIORAVANTI	
Classification results for expanding and shrinking gradient Kähler–Ricci solitons	267
RONAN J CONLON, ALIX DERUELLE and SONG SUN	
Embedding calculus and smooth structures	353
BEN KNUDSEN and ALEXANDER KUPERS	
Stable maps to Looijenga pairs	393
PIERRICK BOUSSEAU, ANDREA BRINI and MICHEL VAN GARREL	