



# *Geometry & Topology*

Volume 28 (2024)

**Moduli spaces of Ricci positive metrics in dimension five**

MCFEELY JACKSON GOODMAN

# Moduli spaces of Ricci positive metrics in dimension five

MCFEELY JACKSON GOODMAN

We use the  $\eta$  invariants of  $\text{spin}^c$  Dirac operators to distinguish connected components of moduli spaces of Riemannian metrics with positive Ricci curvature. We then find infinitely many nondiffeomorphic five-dimensional manifolds for which these moduli spaces each have infinitely many components. The manifolds are total spaces of principal  $S^1$  bundles over  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  and the metrics are lifted from Ricci positive metrics on the bases. Along the way we classify 5-manifolds with fundamental group  $\mathbb{Z}_2$  admitting free  $S^1$  actions with simply connected quotients.

53C20, 53C27, 58D19, 58D27, 58J28; 19K56

Many closed manifolds are known to admit Riemannian metrics of positive Ricci curvature — for example, all compact, simply connected homogeneous spaces, biquotients, and cohomogeneity-one manifolds; see Berestovskii [5], Grove and Ziller [23], Schwachhöfer and Tuschmann [35]. Systematic methods for constructing such metrics on certain connected sums and bundles have been explored in Corro and Galaz-García [11], Gilkey, Park and Tuschmann [20], Nash [33], Searle and Wilhelm [36], Sha and Yang [37] and Wraith [44].

Once we know that a manifold admits positive Ricci curvature we ask how many such metrics it admits. The space of geometrically distinct metrics of positive Ricci curvature on a manifold  $M$  is the moduli space  $\mathfrak{M}_{\text{Ric}>0}(M) = \mathfrak{R}_{\text{Ric}>0}(M)/\text{Diff}(M)$ , where  $\mathfrak{R}_{\text{Ric}>0}(M)$  is the set of positive Ricci curvature metrics on  $M$  and  $\text{Diff}(M)$  is the diffeomorphism group, acting by pullbacks. The number of path components of  $\mathfrak{M}_{\text{Ric}>0}$  serves as a coarse measure of distinct positive Ricci curvature metrics on  $M$ .

We identify an infinite family of 5-manifolds  $M$  with  $\pi_1(M) = \mathbb{Z}_2$  such that  $\mathfrak{M}_{\text{Ric}>0}(M)$  has infinitely many path components.

**Theorem A** *Let  $B^4 = \#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  with  $a+b \geq 2$ , and let  $S^1 \rightarrow M^5 \rightarrow B^4$  be a principal bundle with first Chern class  $2d$ , where  $d \in H^2(B^4, \mathbb{Z})$  is primitive and  $w_2(TB^4) = d \pmod{2}$ . Then  $\mathfrak{M}_{\text{Ric}>0}(M^5)$  has infinitely many path components.*

Here  $w_2$  is the second Stiefel–Whitney class, and a primitive class is one that is not a positive integer multiple of any other. We will see that for each 4-manifold  $B$  there are 2, 3 or 4 diffeomorphism types of such total spaces  $M$ , depending on the value of  $|a-b| \pmod{4}$ , each of which admits infinitely many inequivalent free  $S^1$  actions with quotient  $B$ . The only other five-dimensional manifolds for which  $\mathfrak{M}_{\text{Ric}>0}$  is known to have infinitely many components are the four homotopy real projective spaces

recently described by Dessai and González-Álvaro [14] and five quotients of  $S^2 \times S^3$  recently described by Wermelinger [42].

The conditions on the first Chern class in [Theorem A](#) are equivalent to the statement that  $\pi_1(M^5) = \mathbb{Z}_2$ ,  $M^5$  is nonspin, and the universal cover of  $M^5$  is spin.  $M^5$  can be constructed by taking five-dimensional homotopy real projective spaces, removing tubular neighborhoods of generators of the fundamental group, and gluing along the boundaries of the tubular neighborhoods. By the classification of Smale [39] and Barden [4], the universal cover  $\tilde{M}^5$  is diffeomorphic to  $\#^{a+b-1}S^3 \times S^2$ . But we do not know an explicit description of the deck group action by  $\mathbb{Z}_2$  on  $\tilde{M}^5$ .

Our second theorem identifies conditions under which  $M^5$  admits one, and infinitely many, free  $S^1$  actions. As an application, we will show that the manifolds in [Theorem A](#) admit infinitely many free  $S^1$  actions. We construct the metrics used in [Theorem A](#) by lifting metrics from the quotients of  $M^5$  by those actions. Here  $b_2(M)$  is the second Betti number of  $M$ .

**Theorem B** *Let  $M^5$  be a 5–manifold with  $\pi_1 = \mathbb{Z}_2$ . Then  $M$  admits a free  $S^1$  action with a simply connected quotient if and only if  $M$  is orientable,  $H_2(M, \mathbb{Z})$  is torsion-free and  $\pi_1(M)$  acts trivially on  $\pi_2(M)$ . Furthermore, if  $b_2(M) = 0$ , then  $M$  is diffeomorphic to  $\mathbb{R}P^5$ . If  $b_2(M) > 0$  and  $M$  admits a free  $S^1$  action with simply connected quotient  $B^4$ , then  $M$  admits infinitely many inequivalent free  $S^1$  actions with quotients diffeomorphic to  $B^4$ .*

Note that here  $B^4$  can be any simply connected 4–manifold, and need not be one of the manifolds of [Theorem A](#). [Theorem 1.11](#) provides greater detail about the correspondence between a 5–manifold  $M^5$  and the set  $Q(M)$  of possible quotients  $B^4 = M^5/S^1$ . Given  $M^5$  satisfying the hypotheses of [Theorem B](#), we give conditions on the cohomology ring of a 4–manifold  $B^4$  which are necessary and sufficient for  $B$  to be in  $Q(M)$ . In particular, any smooth manifold homeomorphic to a manifold in  $Q(M)$  is in  $Q(M)$ . In [Corollary 1.12](#) we see that for any such  $M$ ,  $Q(M)$  contains either  $\#^c S^2 \times S^2$  or  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  for some  $a, b, c \in \mathbb{Z}$ . Those manifolds admit metrics with positive Ricci curvature, which can be lifted to  $M$ . Thus we have:

**Corollary** *Let  $M$  be a 5–manifold with  $\pi_1(M) = \mathbb{Z}_2$  admitting a free  $S^1$  action with a simply connected quotient. Then  $M$  admits a metric with positive Ricci curvature.*

Furthermore, it follows from [Theorem 1.11](#) that given a simply connected 4–manifold  $B^4$ , the set of diffeomorphism types of total spaces  $M^5$  with  $\pi_1(M^5) = \mathbb{Z}_2$  of  $S^1$  bundles over  $B^4$  depends only on the cohomology ring of  $B^4$ . In particular, [Theorem A](#) would describe the same set of 5–manifolds if we replaced  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  with one of the manifolds homeomorphic to it.

We first review previous work with methods and results relevant to [Theorem A](#). In [30] Kreck and Stolz invented a moduli space invariant  $s(M, g) \in \mathbb{Q}$  for a metric  $g$  of positive scalar curvature on a closed spin manifold  $M$ . The metric is based on the  $\eta$  spectral invariant of the Dirac operator defined in Atiyah, Patodi and Singer [1]. If  $s(M, g_1) \neq s(M, g_2)$  then  $g_1$  and  $g_2$  represent elements in different path components

of  $\mathfrak{M}_{\text{scal}>0}$ . Kreck and Stolz use the invariant to prove that for  $M^{4k+3}$  with a unique spin structure and vanishing rational Pontryagin classes  $\mathfrak{M}_{\text{scal}>0}(M)$  is either empty or has infinitely many components.

Since a path of Riemannian metrics which maintains positive Ricci curvature maintains positive scalar curvature as well, the  $s$  invariant can detect connected components of  $\mathfrak{M}_{\text{Ric}>0}$ . Kreck and Stolz calculated  $s$  for the Einstein metrics on  $S^1$  bundles  $N_{k,l}^7$  over  $\mathbb{C}P^1 \times \mathbb{C}P^2$  described by Wang and Ziller [41]. Kreck and Stolz showed, using the diffeomorphism classification in [28], that when  $k$  is even and  $\gcd(k, l) = 1$ ,  $N_{k,l}$  is diffeomorphic to infinitely many manifolds in the same family. As the  $s$  invariant takes infinitely many values on those metrics, the authors concluded that  $\mathfrak{M}_{\text{Ric}>0}(N_{k,l})$  has infinitely many components. Similar results have since been proved for  $S^1$  bundles over  $\mathbb{C}P^1 \times \mathbb{C}P^{2n}$  with  $n \geq 1$ ; see Dessai, Klaus and Tuschmann [15].

Wraith showed that for a homotopy sphere  $\sigma^{4k-1}$  bounding a parallelizable manifold,  $\mathfrak{M}_{\text{Ric}>0}(\sigma)$  has infinitely many components. The procedure known as plumbing with disc bundles over spheres produces infinitely many parallelizable manifolds with boundaries diffeomorphic to  $\sigma$ . Wraith [43] constructed metrics of positive Ricci curvature on each boundary, and calculated the  $s$  invariant of each metric in [45].

Dessai [13] and the author [21] used the  $s$  invariant to find several infinite families of seven-dimensional sphere bundles  $M^7$  such that  $\mathfrak{M}_{\text{Ric}>0}(M)$  and  $\mathfrak{M}_{\text{sec}\geq 0}(M)$  have infinitely many path components. Grove and Ziller [22; 24] constructed metrics of nonnegative sectional curvature on the manifolds in those families, and the diffeomorphism classifications in Crowley and Escher [12] and Escher and Ziller [18] show that each manifold is diffeomorphic to infinitely many other members of the family.

More recently, Dessai and González-Álvaro [14] showed that if  $M^5$  is one of the four closed manifolds homotopy equivalent to  $\mathbb{R}P^5$  then  $\mathfrak{M}_{\text{sec}\geq 0}(M)$  and  $\mathfrak{M}_{\text{Ric}>0}(M)$  have infinitely many path components. López de Medrano [32] showed that each such  $M^5$  admits infinitely many descriptions as a quotient of a Brieskorn variety, and Grove and Ziller [23] showed the each quotient admits a metric of nonnegative sectional curvature. Dessai and González-Álvaro calculated the relative  $\eta$  invariant for those metrics to distinguish the path components. Wermelinger [42] extended their method to prove the same conclusion for five  $\mathbb{Z}_2$  quotients of  $S^2 \times S^3$ .

We now outline the proof of [Theorem A](#). We use [Theorem B](#) to show that each manifold  $M^5$  in [Theorem A](#) admits infinitely many inequivalent free  $S^1$  actions with quotient  $B^4 = \#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ . We modify a result of Perelman [34] to show that  $B$  admits a metric of positive Ricci curvature. That metric can be lifted to a metric of positive Ricci curvature on  $M$  by Gilkey, Park and Tuschmann [20]. The lifted metrics depend on the  $S^1$  action, and we get infinitely many distinct metrics on  $M$ .

We show that in dimensions  $4k + 1$ , the  $\eta$  invariant of a certain spin<sup>c</sup> Dirac operator constructed for a positive Ricci curvature metric  $g$  depends only on the connected component of the class of  $g$  in  $\mathfrak{M}_{\text{Ric}>0}$ . To complete the proof we calculate  $\eta$  for each metric on  $M$  and show that it obtains infinitely many values. This is the most intricate part of our proof.

The standard method for calculating the  $\eta$  invariant of a spin Dirac operator on a manifold  $M$  with positive scalar curvature is to extend the metric over a manifold  $W$  with  $\partial W = M$  so that the extension has positive scalar curvature as well. When  $M$  is not spin but  $\text{spin}^c$ , both the metric and a unitary connection on the complex line bundle associated to the  $\text{spin}^c$  structure must be extended. The desired condition then involves the curvatures of both metric and connection. In their work, Dessai and González-Álvaro passed to the universal cover to find a suitable  $W$  over which the connection could be extended to a flat connection. They use equivariant  $\eta$  invariants on the cover to compute the  $\eta$  invariant on the quotient.

In this paper we work directly on  $M$  and use a manifold with boundary  $W$  over which the connection cannot be extended to a flat connection, but the curvature of the extension can be explicitly controlled. To be specific, we extend the metric and connection on  $M$  to a metric  $h$  and connection  $\nabla$  on the disc bundle  $W = M \times_{S^1} D^2$  associated to the  $S^1$  bundle. We then use the Atiyah–Patodi–Singer index theorem [1] to obtain a formula for  $\eta$  in terms of the index of the  $\text{spin}^c$  Dirac operator on  $W$  and topological data on  $W$ . The index will vanish as long as

$$\text{scal}(h) > 2|F^\nabla|_h,$$

where  $F^\nabla$  is the curvature form of the connection  $\nabla$ . We accomplish the extension for a general class of  $S^1$ -invariant metrics of positive scalar curvature. This is more general than we need but may be of independent interest. In fact we construct  $h$  and  $\nabla$  such that

$$\text{scal}(h) > \ell|F^\nabla|_h,$$

where  $\ell$  is a positive integer such that the first Chern class of the  $S^1$  bundle is  $\ell$  times the canonical class of a  $\text{spin}^c$  structure on the quotient.

Sha and Yang [38] constructed metrics of positive Ricci curvature on the 4-manifolds  $\#^{a-b} \mathbb{C}P^2 \#^b S^2 \times S^2$  with  $a > b$ . Those manifolds are diffeomorphic to  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ , so a manifold  $M$  satisfying the hypotheses of [Theorem A](#) also admits a free  $S^1$  action with quotient  $\#^{a-b} \mathbb{C}P^2 \#^b S^2 \times S^2$ . One can lift the Sha–Yang metric to  $M$ , and there is no reason to expect that the resulting metric lies in the same component as the metric lifted from  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  in the proof of [Theorem A](#). We will see, however, that the computation of the  $\eta$  invariant involves only the cohomology ring of the quotient, and we cannot distinguish any new components in this way.

In [37] Sha and Yang also found metrics of positive Ricci curvature on  $\#^b S^2 \times S^2$ . One might expect our methods to yield a similar result in this case. The 5-manifolds, however, would be spin, and the  $\eta$  invariant of the spin Dirac operator in dimension  $4k + 1$  vanishes, even when twisted with certain complex line bundles; see Botvinnik and Gilkey [7].

We now discuss [Theorem B](#). In [26], Hambleton and Su find a complete diffeomorphism classification of 5-manifolds  $M$  with  $\pi_1(M) = \mathbb{Z}_2$  when  $M$  is orientable,  $H_2(M, \mathbb{Z})$  is torsion-free, and  $\pi_1(M)$  acts trivially on  $\pi_2(M)$ . They apply the classification to investigate the diffeomorphism type of the total space of an  $S^1$  bundle over a simply connected 4-manifold. When the total space is nonspin but has a

spin universal cover, as is the case in [Theorem A](#), they can only restrict the diffeomorphism type to two possibilities. Furthermore, an error is present in that calculation, which we correct in [Lemma 1.7](#).

To prove [Theorem B](#), we use the data of a principal  $S^1$  bundle, namely the base and the first Chern class, to compute the diffeomorphism invariants used by Hambleton and Su for the total space. One, the second Betti number, is calculated easily. When the total space is nonspin but has a spin universal cover, we show how the other invariant can be computed by applying a map from  $\Omega_4^{\text{Spin}^c} \rightarrow \Omega_4^{\text{Pin}^+}$  to the base. While a two-fold ambiguity remains in determining which diffeomorphism type corresponds to a specific first Chern class, we are nonetheless able to determine which pairs of invariants are achieved, and achieved infinitely many times, by bundles over a given 4-manifold.

The paper is organized as follows. In [Section 1](#) we examine  $S^1$  actions on 5-manifolds with  $\pi_1 = \mathbb{Z}_2$  and prove [Theorem B](#). In [Section 2](#) we discuss the  $\eta$  invariant of a  $\text{spin}^c$  Dirac operator and show that it can be used to detect connected components of the moduli space in the context of [Theorem A](#). In [Section 3](#) we compute  $\eta$  in the case of certain  $(4n+1)$ -manifolds admitting free  $S^1$  actions and prove [Theorem A](#). In [Section 4](#) we construct the metrics and connections used in the computations of [Section 3](#).

**Acknowledgements** I would like to acknowledge my PhD advisor Wolfgang Ziller for all his help, as well as Anand Dessai, David González-Álvarez, Fernando Galaz-García and Diego Corro for helpful discussions. I am further grateful to Yang Su for pointing out how to work around an error in [\[26\]](#) and for other useful insights. This research was partially supported by National Science Foundation grant DMS-2001985.

## 1 $S^1$ actions on 5-manifolds with $\pi_1 = \mathbb{Z}_2$

Our methods for constructing metrics with positive Ricci curvature and for calculating  $\eta$  use the structure of a principal  $S^1$  bundle. In this section we prove [Theorem 1.11](#), which classifies 5-manifolds with  $\pi_1 = \mathbb{Z}_2$  admitting one, or infinitely many, free  $S^1$  actions with simply connected quotients. [Theorem 1.11](#) also identifies those quotients. In particular, we prove [Theorem B](#) and show that a manifold  $M^5$  satisfying the hypotheses of [Theorem A](#) admits infinitely many inequivalent  $S^1$  actions with the same quotient. Our proof relies on a diffeomorphism classification of 5-manifolds with fundamental group  $\mathbb{Z}_2$  carried out by Hambleton and Su [\[26\]](#).

Given a manifold  $M$  with  $\pi_1(M) = \mathbb{Z}_2$ , a characteristic submanifold  $P \subset M$  is defined as follows. For  $N$  sufficiently large let  $f: M \rightarrow \mathbb{R}P^N$  be a classifying map of the universal covering  $\tilde{M} \rightarrow M$ . We can choose  $f$  to be transverse to  $\mathbb{R}P^{N-1}$ , and hence  $P = f^{-1}(\mathbb{R}P^{N-1})$  is a smooth manifold. One checks that any two manifolds defined in this way are cobordant.

Alternatively, assume that  $P \subset M$  is a submanifold such that the inverse image  $\tilde{P} \subset \tilde{M}$  under the universal covering splits  $\tilde{M}$  into two components  $\tilde{M}_1$  and  $\tilde{M}_2$ . Furthermore  $\partial\tilde{M}_1 = \partial\tilde{M}_2 = \tilde{P}$  and the covering transformation acting on  $\tilde{M}$  switches  $\tilde{M}_1$  and  $\tilde{M}_2$ . One can then construct a map  $f: M \rightarrow \mathbb{R}P^N$  such that  $P = f^{-1}(\mathbb{R}P^{N-1})$ . For details see [\[19; 32\]](#).

The key invariant of the classification in [26] is the class of  $P$  in an appropriate cobordism group. The appropriate structure on  $P$  depends on the second Stiefel–Whitney classes  $w_2$  of  $M$  and  $\tilde{M}$ . Hambleton and Su use the following labels for a manifold  $M$  with  $\pi_1(M) = \mathbb{Z}_2$  and universal cover  $\tilde{M}$ :

- **Type I**  $w_2(T\tilde{M}) \neq 0$ .
- **Type II**  $w_2(TM) = 0$ .
- **Type III**  $w_2(TM) \neq 0$  and  $w_2(T\tilde{M}) = 0$ .

A characteristic submanifold  $P$  of a Type III manifold admits a  $\text{pin}^+$  structure, and all such  $P$  are  $\text{pin}^+$  cobordant. Here  $\text{Pin}^\pm(n)$  is the extension of  $O(n)$  by  $\mathbb{Z}_2$  such that a preimage of a reflection squares to  $\pm 1$  and  $\Omega_n^{\text{Pin}^\pm}$  is the cobordism group of  $n$ -manifolds with  $\text{pin}^\pm$  structures. For details, see [26] and [19].

We review the construction of a  $\text{pin}^+$  structure on  $P$  as we will use it later. Let  $\mu = \tilde{M} \times_{\mathbb{Z}_2} \mathbb{R}$  be the unique nontrivial real line bundle over  $M$ . Recall that  $\tilde{M} = \tilde{M}_1 \cup_{\tilde{f}} \tilde{M}_2$ , and the covering transformation exchanges the components. Thus the normal bundle  $N\tilde{P}$  of  $\tilde{P}$  is trivial and the covering transformation reverses the orientation of the fibers. The normal bundle  $NP$  of  $P$  satisfies

$$NP = N\tilde{P}/\mathbb{Z}_2 \cong \tilde{P} \times_{\mathbb{Z}_2} \mathbb{R} = \mu|_P.$$

Since  $M$  is orientable,

$$w_1(NP) = w_1(TP) = w_1(\det(TP)),$$

so  $NP \cong \det(TP)$ . Thus

$$(1.1) \quad (TM \oplus 2\mu)|_P = TP \oplus 3NP = TP \oplus 3\det(TP).$$

Using [19, Lemma 9; 26, Lemma 2.3], one checks that  $w_2(TM \oplus 2\mu) = 0$ . We can apply [27, Lemma 1.7] to see that a spin structure on  $TP \oplus 3\det(TP)$  induces a  $\text{pin}^+$  structure on  $TP$ . A similar argument on a cobordism shows that any two characteristic submanifolds are  $\text{pin}^+$  cobordant.

Let  $b_2(M)$  denote the second Betti number of a manifold  $M$ . The main theorem for Type III manifolds is [26, Theorem 3.1]:

**Theorem 1.2** [26] *Let  $M_1$  and  $M_2$  be Type III 5-manifolds such that  $\pi_1(M_i) \cong \mathbb{Z}_2$  acts trivially on  $\pi_2(M_i)$  and  $H_2(M_i, \mathbb{Z})$  is torsion-free for  $i = 1, 2$ . Then  $M_1$  is diffeomorphic to  $M_2$  if and only if*

$$b_2(M_1) = b_2(M_2) \quad \text{and} \quad [P_1] = \pm[P_2] \in \Omega_4^{\text{Pin}^+},$$

where  $P_i$  is a characteristic submanifold of  $M_i$ .

We will take the data of a principal  $S^1$  bundle, namely the base and the first Chern class, and identify the diffeomorphism type of the total space. In particular, we will identify when the total space satisfies the hypotheses of Theorem 1.2, and then compute  $b_2$  and  $[P]$ . That computation combined with the classification of Type I and II total spaces in [26, Theorems 6.5 and 6.8] finishes the proof of Theorem 1.11, which in turn implies Theorem B.

A straightforward computation using the long exact homotopy and Gysin sequences proves the following; see for instance [26, Proposition 6.1].



**Lemma 1.3** *Let  $B^n$  be a simply connected manifold and let  $M^{n+1} \rightarrow B^n$  be a nontrivial principal  $S^1$  bundle with first Chern class  $kd$ , where  $d$  is a primitive element of  $H^2(B, \mathbb{Z})$  and  $k \neq 0$  is an integer. Then  $M$  is orientable,  $H_2(M, \mathbb{Z})$  is torsion-free and  $b_2(M) = b_2(B) - 1$ . The fundamental group  $\pi_1(M) \cong \mathbb{Z}_k$  is generated by any  $S^1$  fiber and acts trivially on  $\pi_2(M)$ . The universal cover of  $M$  is the total space of an  $S^1$  bundle over  $B$  with first Chern class  $d$ . If  $k = 2$ ,  $M$  is Type III if and only if and  $w_2(TB) = d \pmod 2$ .*

The condition  $w_2(TB) = d \pmod 2$  implies the existence of a  $\text{spin}^c$  structure on  $B$ . We call  $d$  the canonical class of that  $\text{spin}^c$  structure. On a simply connected manifold a  $\text{spin}^c$  structure is uniquely determined by its canonical class. Thus in the Type III case, given a simply connected  $\text{spin}^c$  4-manifold  $B^4$  with primitive canonical class  $d$ , we want to know the diffeomorphism type of the total space  $M^5$  of the  $S^1$  bundle over  $B^4$  with first Chern class  $2d$ . Since  $b_2(M)$  is determined by Lemma 1.3, it remains to find the  $\text{pin}^+$  cobordism class of a characteristic submanifold  $P^4 \subset M^5$ . In fact, the  $\text{spin}^c$  structure on  $B^4$  will naturally induce a  $\text{pin}^+$  structure on  $P^4$ .

To see this let  $\rho: M \rightarrow B$  be the bundle map and let  $\lambda \rightarrow B$  be a complex line bundle with first Chern class  $d$ ; then  $\rho^*d$  is the unique nontrivial torsion element of  $H^2(M, \mathbb{Z})$ . Let  $\mu \rightarrow M$  be the unique nontrivial real line bundle over  $M$ . As in the proof that a characteristic submanifold of  $M$  will admit a  $\text{pin}^+$  structure — see [19, Lemma 9; 26, Lemma 2.3] —  $w_2(\mu \oplus \mu) = w_1(\mu)^2 \neq 0$ . So  $\mu \oplus \mu$  with its natural orientation is a nontrivial complex line bundle. Since  $\mu \otimes \mu$  is trivial,  $c_1(\mu \oplus \mu)$  is torsion, and we conclude that  $\rho^*\lambda \cong \mu \oplus \mu$ .

The  $S^1$  action on  $M$  splits  $TM$  into a horizontal bundle isomorphic to  $\rho^*TB$  and a vertical bundle, trivialized by an action field, which we call  $TS^1$ . The  $\text{spin}^c$  structure on  $B$  is equivalent to a spin structure on  $TB \oplus \lambda$ . That spin structure induces a spin structure on

$$(1.4) \quad \rho^*(TB \oplus \lambda) \oplus TS^1 \cong TM \oplus \mu \oplus \mu$$

and in turn a  $\text{pin}^+$  structure on  $P \subset M$  using (1.1). Denote by  $\beta(B, d) \in \Omega_4^{\text{pin}^+}$  the cobordism class of  $P$  with this  $\text{pin}^+$  structure. We synthesize the construction with the results of Lemma 1.3 as follows:

**Lemma 1.5** *Let  $B^4$  be a simply connected 4-manifold and let  $M^5$  be the total space of a principal  $S^1$  bundle over  $B$  with first Chern class  $2d \in H^2(B, \mathbb{Z})$  where  $d$  is a primitive element such that  $w_2(TB) = d \pmod 2$ . Then  $M$  satisfies the conditions of Theorem 1.2 with  $b_2(M) = b_2(B) - 1$  and  $[P] = \beta(B, d)$ .*

In the next lemma, we will see that  $\beta$  is a  $\text{spin}^c$  cobordism invariant whenever it is defined.

**Lemma 1.6** *Let  $B_1$  and  $B_2$  be  $\text{spin}^c$  manifolds with primitive canonical classes  $d_1$  and  $d_2$ , respectively. Then:*

- (a)  $\beta(B_1 \amalg B_2, d_1 + d_2) = \beta(B_1, d_1) + \beta(B_2, d_2)$ .
- (b) If  $B_1$  is  $\text{spin}^c$  cobordant to  $B_2$  then  $\beta(B_1, d_1) = \beta(B_2, d_2)$ .



**Proof** Part (a) follows immediately since the total space of the relevant bundle and the characteristic submanifold of that total space will be disjoint unions.

To prove part (b), let  $W$  be a simply connected  $\text{spin}^c$  cobordism between  $B_1$  and  $B_2$  with canonical class  $d$ . Then  $d|_{B_i} = d_i$  for each  $i = 1, 2$ , and  $d$  must be a primitive class. Let  $\pi: N \rightarrow B$  be the principal  $S^1$  bundle over  $W$  with first Chern class  $2d$ . By Lemma 1.3,  $\pi_1(N) = \mathbb{Z}_2$ . We have that  $\partial N = \pi^{-1}(B_1) \amalg \pi^{-1}(B_2)$  and  $M_i = \pi^{-1}(B_i) \rightarrow B_i$  is the principal  $S^1$  bundle with first Chern class  $2d_i$ .

Let  $f: N \rightarrow \mathbb{R}P^N$  be a classifying map for the universal cover of  $N$  which is transverse to  $\mathbb{R}P^{N-1}$ . By Lemma 1.3,  $\pi_1(N)$  is generated by any  $S^1$  orbit, so  $\pi_1(M_i) \rightarrow \pi_1(N)$  is an isomorphism, and  $f|_{M_i}$  is a classifying map for the universal cover of  $M_i$ . Thus  $P_i = f^{-1}(\mathbb{R}P^{N-1}) \cap M_i$  is a characteristic submanifold of  $M_i$  and  $f^{-1}(\mathbb{R}P^{N-1})$  is a cobordism between  $P_1$  and  $P_2$ . The argument before Lemma 1.5 proves that the  $\text{spin}^c$  structure on  $W$  induces a  $\text{pin}^+$  structure on  $f^{-1}(\mathbb{R}P^{N-1})$ . That  $\text{pin}^+$  structure restricts to the  $\text{pin}^+$  structures induced on  $P_i$  by the  $\text{spin}^c$  structures on  $B_i$ . To see this one must simply note that the nontrivial real line bundle over  $N$  restricts to the nontrivial real line bundle over  $M_i$ . We conclude that

$$\beta(B_1, d_1) = [P_1] = [P_2] = \beta(B_2, d_2). \quad \square$$

We now see that  $\beta$  defines a map between the  $\text{spin}^c$  and  $\text{pin}^+$  cobordism groups. The four-dimensional  $\text{spin}^c$  cobordism group  $\Omega_4^{\text{Spin}^c}$  is isomorphic to  $\mathbb{Z}^2$ . The isomorphism takes a  $\text{spin}^c$  manifold  $B$  with canonical class  $d$  to the characteristic numbers

$$\langle d^2, [B] \rangle \quad \text{and} \quad \frac{1}{8}(\langle d^2, [B] \rangle - \text{sign } B).$$

Here  $\text{sign}(B)$  is the signature, and the second integer is the index of the  $\text{spin}^c$  Dirac operator, which we denote by  $\text{ind}(B, d)$ . See [3; 40] for details. To construct generators of  $\Omega_4^{\text{Spin}^c}$  let  $x \in H^*(\mathbb{C}P^2, \mathbb{Z})$  be the generator which is the first Chern class of the Hopf bundle. Give  $X = \mathbb{C}P^2$  the  $\text{spin}^c$  structure with canonical class  $x$  and  $Y = \mathbb{C}P^2 \# \mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$  the  $\text{spin}^c$  structure with canonical class  $d_Y = (3x, x, x) \in H^2(Y, \mathbb{Z}) \cong \oplus^3 H^2(\mathbb{C}P^2, \mathbb{Z})$ . Then  $[X], [Y] \in \Omega_4^{\text{Spin}^c}$  represent  $(1, 0)$  and  $(9, 1)$  under the isomorphism with  $\mathbb{Z}^2$  and form a minimal generating set of  $\Omega_4^{\text{Spin}^c}$ . Since  $X$  and  $Y$  have primitive canonical classes, and their inverses in the cobordism group are given by reversing orientation, we conclude that every class in  $\Omega_4^{\text{Spin}^c}$  can be represented by a simply connected manifold  $B$  with primitive canonical class  $d$ . Lemma 1.6 implies that by mapping the cobordism class of such a pair to  $\beta(B, d)$  we can define a homomorphism  $\beta: \Omega_4^{\text{Spin}^c} \rightarrow \Omega_4^{\text{Pin}^+}$ .

Using the isomorphism  $\Omega_4^{\text{Pin}^+} \cong \mathbb{Z}_{16}$  generated by a  $\text{pin}^+$  structure on  $\mathbb{R}P^4$  we prove the following:

**Lemma 1.7** *We have that*

$$\beta(B, d) = \langle d^2, [B] \rangle + 4\epsilon \text{ind}(B, d) \pmod{16}$$

for an unknown  $\text{sign } \epsilon = \pm 1$ .

This lemma corrects a mistake in the statement of [26, Theorem 6.7]. Our argument uses ideas from the proof in [26] as well as corrections suggested to the author by Yang Su.

**Proof** We will see that  $\beta(X, x) = 1$  and  $\beta(Y, d_Y) = 5$  or  $13$ . The lemma then follows since  $\beta$  is a homomorphism and  $\Omega_4^{\text{Spin}^c} \cong \mathbb{Z}^2$ .

The principal  $S^1$  bundle  $\mathbb{R}P^5 \rightarrow \mathbb{C}P^2$  which is a  $\mathbb{Z}_2$  quotient of the Hopf bundle has first Chern class  $2x$ . Since  $\mathbb{R}P^4$  is a characteristic submanifold of  $\mathbb{R}P^5$ , it follows that

$$\beta(X, x) = [\mathbb{R}P^4] = 1 \in \Omega_4^{\text{Pin}^+}.$$

The second calculation is more involved. We use the notation  $[z_0, z_1, z_2] \in \mathbb{C}P^2$  and  $[z_0, z_1, z_2]_{\pm} \in \mathbb{R}P^5$  for the respective images of the point  $(z_0, z_1, z_2) \in S^5 \subset \mathbb{C}^3$ . Let  $\rho: M \rightarrow Y$  be the principal  $S^1$  bundle with first Chern class  $2d_Y \in H^2(Y, \mathbb{Z})$  as defined above. By Lemma 1.3, the double cover  $\tilde{M}$  of  $M$  is the total space of a principal  $S^1$  bundle  $\tilde{\rho}: \tilde{M} \rightarrow Y$  with first Chern class  $d_Y$ . Let  $g: Y \rightarrow \mathbb{C}P^2$  be a classifying map for  $\tilde{\rho}$  which is transverse to  $\mathbb{C}P^1 \subset \mathbb{C}P^2$  and has a regular value  $[1, 0, 0] \in \mathbb{C}P^1$ . Then  $g^*x = d_Y$  and the pullback of  $\pi: \mathbb{R}P^5 \rightarrow \mathbb{C}P^2$  by  $f$  has first Chern class  $2d_Y$ . There is a map of principal  $S^1$  bundles  $f: M \rightarrow \mathbb{R}P^5$  covering  $g$ , that is, an  $S^1$  equivariant map making the following diagram commute:

$$\begin{array}{ccc} M & \xrightarrow{f} & \mathbb{R}P^5 \\ \downarrow \rho & & \downarrow \pi \\ Y & \xrightarrow{g} & \mathbb{C}P^2 \end{array}$$

Since the fundamental groups of  $M$  and  $\mathbb{R}P^5$  are generated by  $S^1$  orbits (see Lemma 1.3), the homomorphism  $f_*: \pi_1(M) \rightarrow \pi_1(\mathbb{R}P^5)$  is an isomorphism and  $f$  is a classifying map for the double cover  $\tilde{M} \rightarrow M$ . Thus if we show that  $f$  is transverse to  $\mathbb{R}P^4 \subset \mathbb{R}P^5$ , we can conclude that  $P = f^{-1}(\mathbb{R}P^4)$  is a characteristic submanifold of  $M$ . Then given the correct  $\text{pin}^+$  structure on  $P$ ,  $\beta(Y, d_Y) = [P] \in \Omega_4^{\text{Pin}^+}$ .

To see that  $f$  is transverse to  $\mathbb{R}P^4 = \{[z_0, z_1, r]_{\pm} \in \mathbb{R}P^5 \mid r \in \mathbb{R}\}$  note that at points in  $\pi^{-1}(\mathbb{C}P^2 \setminus \mathbb{C}P^1)$ ,  $\mathbb{R}P^4$  is transverse to the  $S^1$  orbits, which are contained in the image of the equivariant map  $f$ . At points in  $\pi^{-1}(\mathbb{C}P^1)$ , we associate the horizontal space of the  $S^1$  action with  $T\mathbb{C}P^2$ . By assumption on  $g$ ,  $f$  is transverse to  $T\mathbb{C}P^1$ , and  $T\mathbb{C}P^1 \subset T\mathbb{R}P^4$ .

For later, we also note that  $f$  is transverse to  $\mathbb{R}P^2 = \{[z_0, r, 0] \in \mathbb{R}P^5 \mid r \in \mathbb{R}\}$  since  $T\mathbb{C}P^1 \subset T\mathbb{R}P^2$  except at  $[1, 0, 0]$ , which is a regular value of  $f$  by assumption on  $g$ .

There is a short exact sequence

$$(1.8) \quad 0 \rightarrow \mathbb{Z}_2 \rightarrow \Omega_4^{\text{Pin}^+} \xrightarrow{\phi} \Omega_2^{\text{Pin}^-} \rightarrow 0,$$

where  $\phi$  is given by taking the cobordism class of a submanifold dual to  $w_1^2$ ; see [26, page 172] and [27, page 217] for details. Thus  $\Omega_2^{\text{Pin}^-}$  is isomorphic to  $\mathbb{Z}_8$  with generator  $[\mathbb{R}P^2]$ . We now compute  $\phi([P]) = 5$ , which restricts the possible values of  $\beta(Y, d_Y) = 5$  or  $13$ , as desired.

We need to find a submanifold of  $P$  dual to  $w_1^2(TP)$ . Denote by  $N\mathbb{R}P^4$  the normal bundle of  $\mathbb{R}P^4$  in  $\mathbb{R}P^5$  and by  $NP$  the normal bundle of  $P$  in  $M$ . Then  $f^*N\mathbb{R}P^4 = NP$ . Since  $\mathbb{R}P^5$  and  $M$  are orientable,

$$w_1(TP) = w_1(NP) = f^*w_1(N\mathbb{R}P^4) = f^*w_1(T\mathbb{R}P^4).$$

Since  $w_1(T\mathbb{R}P^4)^2$  is dual to  $\mathbb{R}P^2 \subset \mathbb{R}P^4$ , as long as the mod 2 degree of  $f: f^{-1}(\mathbb{R}P^2) \rightarrow \mathbb{R}P^2$  is 1, it follows that  $f^{-1}(\mathbb{R}P^2)$  is dual to  $w_1(TP)^2$ . For convenience let  $\Sigma = f^{-1}(\mathbb{R}P^2)$ . Since  $[1, 0, 0]$  is a regular point of  $g$ ,  $[1, 0, 0]_{\pm}$  is a regular point of  $f$ , and the degree of  $f$  is the same as the degree of  $f|_{\Sigma}$ . The degree of  $f$  is the same as the degree of  $g$ . The degree of  $g$  is given by

$$\langle g^*x^2, Y \rangle = \langle d_Y^2, [Y] \rangle = 9.$$

Thus the mod 2 degree of  $f|_{\Sigma}$  is 1 and  $\phi([P]) = [\Sigma] \in \Omega_2^{\text{Pin}^-}$ .

Let  $U$  be a tubular neighborhood of the  $S^1$  orbit of  $[1, 0, 0]_{\pm}$  and  $V = \mathbb{R}P^2 \setminus U$ . Since  $[1, 0, 0]$  is a regular value of  $g$  we can choose  $U$  to be made up of regular values of  $f$ . Then  $f|_{f^{-1}(U)}$  is a covering map. Since  $f$  maps  $S^1$  fibers to  $S^1$  fibers,  $f_*: \pi_1(f^{-1}(U)) \rightarrow \pi_1(U)$  is surjective and the covering is trivial. Thus  $f^{-1}(U)$  is the disjoint union of  $\text{deg}(f) = 9$  copies of  $U$  and  $f^{-1}(U \cap \mathbb{R}P^2)$  is 9 copies of  $U \cap \mathbb{R}P^2$ . The  $S^1$  orbit of  $[1, 0, 0]_{\pm}$  is a nontrivial loop in  $\mathbb{R}P^2$ , and  $U \cap \mathbb{R}P^2$  is a tubular neighborhood of that loop, diffeomorphic to  $\mathbb{R}P^2 \setminus D^2$  (the Möbius band). The local inverses to  $f|_{f^{-1}(U)}$  are equivariant embeddings of the oriented tubular neighborhood  $U$  and are all isotopic. It follows that the 9 embedding of  $\mathbb{R}P^2 \setminus D^2$  making up  $f^{-1}(U \cap \mathbb{R}P^2)$  are all isotopic. Thus the process by which  $TM$  induces a  $\text{pin}^+$  structure on  $P$ , which in turn induces a  $\text{pin}^-$  structure on  $\Sigma$ , will induce the same  $\text{pin}^-$  structure on each of the 9 copies of  $\mathbb{R}P^2 \setminus D^2$ .

Since  $\pi(\mathbb{R}P^2) = \mathbb{C}P^1$  and  $\pi(U) \cap \mathbb{C}P^1$  is diffeomorphic to a disc  $D^2$  around  $[1, 0, 0]$  made up of regular values of  $g$ ,  $g^{-1}(\pi(U) \cap \mathbb{C}P^1)$  is 9 copies of  $D^2$  and  $\pi(V) = \mathbb{C}P^1 \setminus D^2$ .  $\pi|_{\mathbb{R}P^2}$  is injective away from the orbit of  $[1, 0, 0]_{\pm}$ , and thus is injective on  $V$ . It follows that  $\rho$  maps  $f^{-1}(V)$  injectively onto  $g^{-1}(\pi(V))$ . Thus  $f^{-1}(V)$  is diffeomorphic to  $g^{-1}(\mathbb{C}P^2)$  with 9 discs removed while  $f^{-1}(U \cap \mathbb{R}P^2)$  is 9 copies of  $\mathbb{R}P^2 \setminus D^2$ . In other words,

$$(1.9) \quad \Sigma \cong g^{-1}(\mathbb{C}P^1) \# \mathbb{R}P^2 \# \dots \# \mathbb{R}P^2,$$

and the nine summands of  $\mathbb{R}P^2$  all have the same  $\text{pin}^-$  structure.  $\Omega_2^{\text{Pin}^-}$  is generated by  $[\mathbb{R}P^2]$ , and so it remains to compute the value of  $[g^{-1}(\mathbb{C}P^1)]$ .

Let  $\chi = g^{-1}(\mathbb{C}P^2)$ . We will use a general method to define a  $\text{pin}^-$  structure called  $r_{\chi}$  on  $\chi$  and compute  $[\chi] \in \Omega_2^{\text{Pin}^-}$  with this structure. We will then show that  $r_{\chi}$  is the correct  $\text{pin}^-$  structure to use, that is,  $r_{\chi}$  is compatible under (1.9) with the  $\text{pin}^-$  structure used to identify  $[\Sigma]$  with  $\phi([P])$ , which we will call  $r$ .

Consider a simply connected  $\text{spin}^c$  4-manifold  $B$  with canonical class  $d$  and  $\nu$  the complex line bundle with  $c_1(\nu) = d$ . Let  $N \subset B$  be a smooth submanifold dual to  $d$ . Then  $\nu|_N$  is isomorphic to the normal bundle of  $N$ . The  $\text{spin}^c$  structure on  $B$  is equivalent to a  $\text{spin}$  structure, called  $s$ , on  $TB \oplus \nu$ . Restricted

to  $N$ , this is a spin structure on  $TN \oplus 2\nu$ . The transition functions for  $2\nu$  admit a canonical lift from  $SO(4)$  to  $Spin(4)$ ; simply multiply two copies of any lift for the transition functions of  $\nu$ , and the sign ambiguities cancel. Note that the identity lifts to the identity in this way. Using this lift,  $s$  induces a spin structure  $s_N$  on  $N$ .

The spin cobordism class of  $N$  depends only on the  $spin^c$  cobordism class of  $B$ . To see this, note that the dual to the canonical class of a  $spin^c$  cobordism will be a spin cobordism between the two relevant submanifolds. Thus we have a homomorphism

$$\psi : \Omega_4^{Spin^c} \rightarrow \Omega_2^{Spin} \cong \mathbb{Z}_2$$

defined by  $\psi([B]) = [N]$ . Indeed, there is a long exact sequence

$$\rightarrow \Omega_4^{Spin} \rightarrow \Omega_4^{Spin^c} \rightarrow \Omega_2^{Spin}(BU(1)) \rightarrow \Omega_3^{Spin} = 0$$

as in [26, page 154; 25, page 654]. We see that  $\psi$  is surjective by noting that  $\psi$  is the composition of  $\Omega_4^{Spin^c} \rightarrow \Omega_2^{Spin}(BU(1))$  with the surjective map  $\Omega_2^{Spin}(BU(1)) \rightarrow \Omega_2^{Spin}$ , which ignores the map to  $BU(1)$ .

Recall that  $X, Y$  generate  $\Omega_4^{Spin^c}$ . The canonical class of  $X$  is dual to  $\mathbb{C}P^1 \subset \mathbb{C}P^2$ , which is nullcobordant, so  $\psi([X]) = 0$ . Since  $\psi$  is surjective,  $\psi([Y])$  generates  $\Omega_2^{Spin}$ . Since  $\mathbb{C}P^1$  contains a regular value of  $g$ , the degree of  $g|_\chi$  equals the degree of  $g$  and  $\chi$  is dual to  $g^*x = d_Y$ . Giving  $\chi$  the spin structure  $s_\chi$  used to define  $\psi$ , we have  $\psi([Y]) = [\chi] \neq 0$ .

$Spin(n)$  embeds naturally into both  $Pin^\pm(n)$ , so a spin structure induces a natural  $pin^-$  structure. Kirby and Taylor show that in dimension 2, the corresponding map

$$\Omega_2^{Spin} \cong \mathbb{Z}_2 \rightarrow \Omega_2^{Pin^-} \cong \mathbb{Z}_8$$

is injective; see [27, Proposition 3.8]. Let  $r_\chi$  be the  $Pin^-$  structure on  $\chi$  induced by  $s_\chi$ . Using that structure, we have  $[\chi] = 4 \in \Omega_2^{Pin^-}$ . Once we confirm that  $r_\chi$  is the correct structure, we conclude with (1.9) that  $\phi([P]) = 5$ , completing the proof of Lemma 1.7.

Let  $r$  be the  $pin^-$  structure on  $\Sigma$  used to define  $\phi([P])$ . Recall that  $\rho$  is a diffeomorphism between the open set  $O = f^{-1}(V) \subset \Sigma$  and  $\rho(O)$ , which is  $\chi$  with 9 discs removed. It remains only to check that  $r = \rho^*r_\chi$  on  $O$ .

We first recall the definition of  $r$ . Let  $\mu$  be the nontrivial real line bundle over  $M$  and let  $E = TM \oplus 2\mu$ . Let  $\lambda$  be the complex line bundle over  $Y$  with  $c_1(\lambda) = d_Y$  and let  $s$  be spin structure on  $TY \oplus \lambda$  used in the definition of  $\psi$ . With the isomorphism (1.4),  $s$  induces a spin structure on  $E$  called  $s_E$ . Then (1.1) shows

$$E|_P = TP \oplus 3 \det(TP)$$

and we induce a  $pin^+$  structure on  $TP$  using a canonical lift of the transition functions of  $3 \det(TP)$  from  $O(3)$  to  $Pin^-(3)$ . In turn,

$$TP|_\Sigma = T\Sigma \oplus 2 \det(T\Sigma),$$

and using a canonical lift of the transition functions of  $2 \det(T\Sigma)$  from  $O(2)$  to  $\text{Pin}^+(2)$  we induce the  $\text{pin}^-$  structure  $r$  on  $\Sigma$ . Note that the normal bundle of  $\Sigma$  in  $P$  is orientable and thus

$$w_1(\det(T\Sigma)) = w_1(\det(TP)|_\Sigma).$$

In this way we can combine the two steps and see that  $s_E$  induces  $r$  on  $T\Sigma$  using the isomorphism

$$(1.10) \quad E|_\Sigma = T\Sigma \oplus 5 \det(T\Sigma)$$

and a canonical lift of the transition functions of  $5 \det(T\Sigma)$  from  $O(5)$  to  $\text{Pin}^+(5)$ . The details of the canonical lifts involved can be found in [27, Lemma 1.7]; the salient fact is that each lifts the identity to the identity.

Next, we note that  $\det(T\Sigma)$  and  $\rho^*\lambda$  are trivial over  $O$ . The former follows because  $O$  is an open set in  $\Sigma$ , but is orientable since it is diffeomorphic to an open set in  $\chi$ . As for the latter, we have seen that  $\rho^*\lambda \cong 2\mu$ ,  $\mu|_P = \det(TP)$ , and  $\det(TP)|_\Sigma = \det(T\Sigma)$ . Since  $\rho$  is a diffeomorphism on  $O$  and  $\rho^*\lambda$  is trivial,  $\lambda$  is trivial on  $\rho(O)$ .

Let  $t_{ij}$  be transition functions with values in  $\text{SO}(2)$  for  $T\chi$ . As we saw in the definition of  $\psi$ , for points in  $\chi$ ,

$$TY \oplus \lambda \cong T\chi \oplus 2\lambda.$$

Thus on  $\rho(O)$  the transition functions for  $\lambda$  can be chosen to be the identity and the transition functions for  $(TY \oplus \lambda)|_\chi$  can be chosen to be  $t_{ij}$ . The spin structure  $s$  gives a lift of  $t_{ij}$  to  $\tilde{t}_{ij}$  in  $\text{Spin}(2)$ . Since the canonical lift of the transition functions for  $2\lambda$  will also be the identity,  $\tilde{t}_{ij}$  is also the lift given by  $s_\chi$  and  $r_\chi$ .

Furthermore, using (1.4),  $t_{ij} \circ \rho$  are transition functions for  $E$  on  $O$ . By definition,  $s_E$  gives the lift  $\tilde{t}_{ij} \circ \rho$ . Using (1.10),  $t_{ij} \circ \rho$  are transition functions for both  $E|_O$  and  $T\Sigma$ , compatible by picking trivial transition functions for  $5 \det(T\Sigma)$ . The canonical lift of the transition functions for  $5 \det(T\Sigma)$  will also be trivial, and the lift given by  $r$  will simply be the inclusion of  $\tilde{t}_{ij} \circ \rho$  into  $\text{Pin}^-(2)$ . Thus  $r = \rho^*r_\chi$  on  $O$ . This completes the proof of Lemma 1.7.  $\square$

We can now prove Theorem B. In fact, we prove the following more detailed theorem, which includes the statement of Theorem B. Here we use the notation of Hambleton and Su, where  $\#_{S^1}$  is gluing along the boundary of a tubular neighborhood of a generator of  $\pi_1$ . The  $X(q)$  for  $q = 1, 3, 5, 7$  are the four closed manifolds homotopy equivalent to  $\mathbb{R}P^5$ , with  $X(1) = \mathbb{R}P^5$ , and the  $X(q)$  for  $q = 0, 2, 4, 6, 8$  are constructed from pairs of homotopy  $\mathbb{R}P^5$ 's using the operation  $\#_{S^1}$ . The labeling is such that a characteristic submanifold  $P \subset X(q)$  has class  $q \in \Omega_4^{\text{Pin}^+}/\pm = \{0, \dots, 8\}$ . See the discussion before [26, Theorem 3.7] for details.

**Theorem 1.11** *Let  $M$  be a 5-manifold with  $\pi_1 = \mathbb{Z}_2$ . Let  $P \subset M$  be a characteristic submanifold.*

- (1)  *$M$  admits a free  $S^1$  action with a simply connected quotient if and only if  $M$  is orientable,  $H_2(M, \mathbb{Z})$  is torsion-free, and  $\pi_1(M)$  acts trivially on  $\pi_2(M)$ . Furthermore if  $b_2(M) = 0$  then  $M$  is diffeomorphic to  $\mathbb{R}P^5$ .*

$M^5$	$Q(M^5) =$ simply connected 4–manifolds $B^4$ such that:
Type II	$B$ is spin and $b_2(B) = b_2(M) + 1$ .
Type III	$B$ is nonspin, $b_2(B) = b_2(M) + 1$ and $\text{sign}(B) = \pm[P] \pmod{4}$ .
Type I and $M \notin S$	$B$ is nonspin and $b_2 = b_2(M) + 1$ .
$X(q) \#_{S^1} (\mathbb{C}P^2 \times S^1)$ with $q = 0, 4$	$B$ is nonspin, $b_2 = 3$ and $ \text{sign } B  = 1$ .
$X(q) \#_{S^1} (S^2 \times \mathbb{R}P^3)$ with $q = 0, 4$	$B$ is nonspin, $b_2 = 4$ and $ \text{sign } B  < 4$ .

Table 1

(2) Suppose  $M^5$  satisfies the conditions in (1). Let  $Q(M)$  be the set of quotients of  $M$  by free  $S^1$  actions. Table 1 gives necessary and sufficient conditions for a 4–manifold to be in  $Q(M)$ .  $S$  is a set of four exceptional 5–manifolds of Type I described in the final two rows. If  $b_2(M) > 0$  then for each  $B \in Q(M)$ ,  $M$  admits infinitely many inequivalent  $S^1$  actions with quotients diffeomorphic to  $B$ .

Thus given  $M^5$  satisfying the hypotheses of (1) and matching the description of one of the rows in the left column, a 4–manifold  $B^4$  is diffeomorphic to a quotient of  $M^5$  by a free  $S^1$  action if and only if it satisfies the conditions given in the corresponding row of the right column.

**Proof** We prove (2) first. Let  $M$  be an orientable 5–manifold with  $\pi_1(M) = \mathbb{Z}_2$  acting trivially on  $\pi_2(M)$ ,  $H_2(M, \mathbb{Z})$  torsion-free, and  $b_2(M) > 0$  unless  $M \cong \mathbb{R}P^2$ . Let  $P \subset M$  be a characteristic submanifold.

If  $M \rightarrow B$  is a principal  $S^1$  bundle, the long exact homotopy sequence implies that  $\pi_1(M) \rightarrow \pi_1(B)$  is surjective. If  $\pi_1(B) = \mathbb{Z}_2$ , then the Gysin sequence implies that  $H^3(B) \rightarrow H^3(M)$  is injective. Since  $M$ , and thus  $B$ , is orientable,  $H^3(B) = \mathbb{Z}_2$  and  $H_2(M)$  would not be torsion-free. Thus any quotient of  $M$  by a free  $S^1$  action is simply connected.

**$M$  is Type II** First, suppose  $M \rightarrow B$  is a principal  $S^1$  bundle. By Lemma 1.3,  $b_2(B) = b_2(M) + 1$  and by [26, Proposition 6.1],  $B$  is spin.

Conversely, let  $B$  be a simply connected spin 4–manifold with  $b_2(B) = b_2(M) + 1$ . It follows from [26, Proposition 6.1] that all of the total spaces of principal  $S^1$  bundle over  $B$  with  $\pi_1 = \mathbb{Z}_2$  are Type II and have second Betti number  $b_2(B) - 1$ . By [26, Theorem 3.1] all such total spaces are diffeomorphic to  $M$ . If  $b_2(M) \geq 1$  there are infinitely many primitive elements of  $H^2(B, \mathbb{Z}) = \mathbb{Z}^{b_2(M)+1}$  and thus infinitely many nonisomorphic such bundles.

**$M$  is Type III** Suppose  $M \rightarrow B$  is a principal  $S^1$  bundle. By Lemma 1.3,  $b_1(B) = b_1(M) + 1$  and the first Chern class of the bundle is  $2d$ , where  $d$  is a primitive element of  $H^2(B, \mathbb{Z})$  such that  $w_2(TB) = d \pmod{2}$ . It follows that  $B$  is nonspin, and by [31, Corollary II.2.12] the intersection form of  $B$  is odd. By the classification of integral forms and Donaldson’s theorem [16, page 5 and Theorem 1.3.1],

the intersection form of  $B$  is diagonal, and so  $H^*(B, \mathbb{Z}) = H^*(\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}, \mathbb{Z})$  for some integers  $a$  and  $b$ . Then using [31, Corollary II.2.12] again we see that

$$w_2(B) = (1, 1, \dots, 1) \in H^2(B, \mathbb{Z}_2) \cong \mathbb{Z}_2^{a+b}.$$

Thus  $d = (d_1, \dots, d_{a+b}) \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}^{a+b}$ , where each  $d_i$  is an odd integer. This completes the proof of one direction of (2) since

$$[P] = \beta(B, d) = \langle d^2, [B] \rangle = \sum_{i=1}^a d_i^2 - \sum_{j=a+1}^b d_j^2 = \text{sign } B \pmod{4}.$$

Conversely, Let  $B$  be a nonspin simply connected 4–manifold with  $b_2(B) = b_2(M) + 1$ . Assume further that  $\text{sign}(B) = [P] \in \mathbb{Z}_4/\pm$ . Again,  $H^*(B, \mathbb{Z}) = H^*(\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}, \mathbb{Z})$ , where  $b_2(B) = a + b$  and  $\text{sign}(B) = a - b$ . Choose  $c \in \{0, 1, 2, 3\}$  such that  $\pm[P] = a - b + 4c \pmod{16}$ . If  $b_2(M) > 0$ , choose  $k$  such that

$$(4 + 2\epsilon)k(k + 1) = 4c \pmod{16},$$

where  $\epsilon = \pm 1$  is the sign from Lemma 1.7. If  $b_2(M) = 0$  then choose  $k = 0$ . Set

$$d_k = (1 + 2k, 1, \dots, 1) \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}^{a+b}.$$

Then  $d$  is primitive and as above, we see that  $w_2(TB) = d \pmod{2}$ . Using Lemma 1.7 we have

$$\beta(B, d_k) = \text{sign } B + (4 + 2\epsilon)k(k + 1) = \pm[P] \pmod{16}.$$

Hence, by Lemma 1.5 and Theorem 1.2,  $M$  is diffeomorphic to the total space of an  $S^1$  bundle over  $B$  with first Chern class  $2d_k$ . In the case where  $b_2(M) > 1$ , there are infinitely many choices of  $k$  yielding distinct classes  $d_k$ , and  $M$  is diffeomorphic to infinitely many total spaces of nonisomorphic  $S^1$  bundles over  $B$ .

**$M$  is Type I** Suppose  $M \rightarrow B$  is a principal  $S^1$  bundle. By Lemma 1.3,  $b_1(B) = b_1(M) + 1$  and by [26, Proposition 6.1]  $B$  is nonspin and the first Chern class of the bundle is  $2d$ , where  $d$  is a primitive element of  $H^2(B, \mathbb{Z})$  such that  $w_2(TB) \neq d \pmod{2}$ .

If  $M = X(q) \#_{S^1} (\mathbb{C}P^2 \times S^1)$  with  $q = 0, 4$ , then  $b_2(B) = 3$  and by [26, Theorem 6.8]  $\langle d^2, [B] \rangle = \pm q \pmod{8}$ . If  $\text{sign}(B) = \pm 3$ , then up to orientation as above  $H^*(B, \mathbb{Z}) = H^*(\#^3 \mathbb{C}P^2, \mathbb{Z})$  and  $w_2(TB) = (1, 1, 1)$ . Thus

$$d = (d_1, d_2, d_3) \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}^3,$$

and some  $d_i$  must be even. Since  $d$  is primitive, some  $d_i$  must be odd. One easily checks that under these conditions,  $\langle d^2, [B] \rangle \neq 0, 4 \pmod{8}$ . So  $\text{sign}(B) = \pm 1$ .

If  $M = X(q) \#_{S^1} (S^2 \times \mathbb{R}P^3)$  with  $q = 0, 4$ , then  $b_2(B) = 4$  and  $\langle d^2, [B] \rangle = \pm q \pmod{8}$ . If  $\text{sign}(B) = \pm 4$ , then up to orientation by the argument in the Type III case,  $H^*(B, \mathbb{Z}) = H^*(\#^4 \mathbb{C}P^2, \mathbb{Z})$  and

$$d = (d_1, d_2, d_3, d_4) \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}^3$$

with at least one  $d_i$  even and at least one  $d_i$  odd. Again  $\langle d^2, [B] \rangle \neq 0, 4 \pmod{8}$ , so  $|\text{sign}(B)| < 4$ .



Conversely, Let  $B$  be a simply connected nonspin 4–manifold satisfying the conditions given by the table for  $Q(M)$ . Then  $H^*(B, \mathbb{Z}) = H^*(\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}, \mathbb{Z})$  for some integers  $a, b$  such that  $a + b = b_1(M) + 1$ . Let  $(q, s) \in \mathbb{Z}_8 \oplus \mathbb{Z}_2$  represent the cobordism class of  $P \subset M$  in the pin<sup>c</sup> cobordism group  $\Omega_4^{\text{Pin}^c} \cong \mathbb{Z}_8 \oplus \mathbb{Z}_2$ ; see [26, page 154]. By [26, Theorem 3.6],  $q + s = b_2(M) + 1 \pmod 2$ .

If  $q = 0, 4$  then [26, Theorem 3.7] implies that  $a + b \geq 3$ , so we can assume that up to orientation  $a \geq 2$  and using Table 1 either  $a + b \geq 5$  or  $|\text{sign}(B)| < b_2(B)$ , which implies  $b > 0$ . Define the following elements  $d_k \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}^a \oplus \mathbb{Z}^b$  for each  $k \in \mathbb{Z}$ :

$$\begin{aligned}
 q = 0: \quad d_k &= \begin{cases} (1 + 8k, 0, \dots, 0, 1) & \text{if } b > 0, \\ (2 + 8k, 1, 1, 1, 0, \dots, 0) & \text{if } b = 0, \end{cases} \\
 q = 4: \quad d_k &= \begin{cases} (2 + 8k, 1, 0, \dots, 0, 1) & \text{if } b > 0, \\ (1 + 8k, 1, 1, 1, 0, \dots, 0) & \text{if } b = 0. \end{cases}
 \end{aligned}$$

If  $q = 2$ , [26, Theorem 3.7] implies that  $a + b \geq 3$  and we can assume  $a \geq 2$  and define

$$q = 2: \quad d_k = (1 + 8k, 1, 0, \dots, 0).$$

If  $q$  is odd, by [26, Theorem 3.7]  $a + b \geq 2$ , and we can assume  $a \geq 1$ . Define

$$\begin{aligned}
 q = 1: \quad d_k &= (1 + 8k, 4, 0, \dots, 0), \\
 q = 3: \quad d_k &= (1 + 8k, 2, 0, \dots, 0).
 \end{aligned}$$

In each case  $d_k$  is primitive,  $w_2(TB) \neq d_k \pmod 2$ , and  $q = \pm \langle d_k^2, [B] \rangle \pmod 8$ . By [26, Theorem 6.8] the  $S^1$  bundle over  $B$  with first Chern class  $2d_k$  is diffeomorphic to  $M$ . Again, infinitely many  $k$  yield distinct classes  $d_k$  and thus nonisomorphic bundles.

To prove (1), first assume  $M$  is a 5–manifold with  $\pi_1(M) = \mathbb{Z}_2$  admitting a free  $S^1$  action with simply connected quotient  $B$ . By Lemma 1.3,  $M$  is orientable,  $\pi_1(M)$  acts trivially on  $\pi_2(M)$  and  $H_2(M, \mathbb{Z})$  is torsion-free. If  $b_2(M) = 0$ , then  $b_2(B) = 1$  and up to orientation  $H^*(B, \mathbb{Z}) \cong H^*(\mathbb{C}P^2, \mathbb{Z})$  and  $w_2(TB)$  is nonzero. There are only two primitive classes  $\pm d \in H^2(B, \mathbb{Z}) \cong \mathbb{Z}$ , each restricting to  $w_2(B) \pmod 2$ . Thus  $B$  is of Type III and  $\beta([B, d]) = \pm 1$ . By Theorem 1.2,  $M$  is diffeomorphic to  $\mathbb{R}P^5$ .

To prove the converse, suppose  $M$  is an orientable 5–manifold with  $\pi_1(M) = \mathbb{Z}_2$  acting trivially on  $\pi_2(M)$  and  $H_2(M, \mathbb{Z})$  torsion-free. Let  $P \subset M$  be a characteristic submanifold. Since  $\mathbb{R}P^5$  admits a free  $S^1$  action induced by the Hopf action we assume  $b_2(M) > 0$ . We must show the set  $Q(M)$  described in Table 1 is nonempty.

If  $M$  is Type II, by [26, Theorem 3.6]  $b_2(M)$  is odd. Then  $B = \#^{(b_2(M)+1)/2} S^2 \times S^2 \in Q(M)$ . If  $M$  is Type I then  $B = \#^{b_2(M)} \mathbb{C}P^2 \# \overline{\mathbb{C}P^2} \in Q(M)$ . If  $M$  is Type III, let  $0 \leq c < 16$  be an integer such that  $[P] = c \pmod{16}$ . By [26, Theorem 3.6] we see that  $c = b_2(M) + 1 \pmod 2$ . Choose  $l$  such that

$$0 \leq c - 4l < 4.$$

Then

$$a = \frac{1}{2}(b_2(M) + 1 + c - 4l) \quad \text{and} \quad b = \frac{1}{2}(b_2(M) + 1 - c + 4l)$$

are nonnegative integers. Let  $B = \#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ . Then  $b_2(B) = b_2(M) + 1$  and  $\text{sign}(B) = [P] \in \mathbb{Z}_4/\pm$ . So  $B \in Q(M)$ . □

We note that the final paragraph of the proof above in fact shows the following, which we will make use of later.

**Corollary 1.12** *Let  $M$  be a 5–manifold with  $\pi_1 = \mathbb{Z}_2$  admitting a free  $S^1$  action with a simply connected quotient. Then  $M$  admits a free  $S^1$  action with quotient diffeomorphic to either  $\#^c S^2 \times S^2$  or  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  for some  $a, b, c \in \mathbb{Z}$ .*

Combining [Theorem 1.11](#) with [[26](#), [Theorem 3.7](#)], we can characterize the manifolds satisfying [Theorem A](#).

**Corollary 1.13** *Let  $M^5$  be a 5–manifold. The following are equivalent:*

- (1)  $M^5$  is Type III and admits a free  $S^1$  action with a simply connected quotient.
- (2) There exists  $B^4 = \#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  with  $a, b \in \mathbb{Z}_{\geq 0}$  such that  $M^5$  is the total space of a principal bundle  $S^1 \rightarrow M^5 \rightarrow B^4$  with first Chern class  $2d$ , where  $d \in H^2(B^4, \mathbb{Z})$  is primitive and  $w_2(TB^4) = d \pmod 2$ .
- (3) There exist  $k \in \mathbb{Z}_{\geq 0}$  and  $q \in \{0, 1, \dots, 8\}$ , with  $k > 0$  if  $q$  is 3, 5 or 7, such that  $M^5$  is diffeomorphic to

$$X(q) \#_{S^1} (\#^k (S^2 \times S^2) \times S^1).$$

If those conditions are satisfied then  $\mathfrak{M}_{\text{Ric}>0}(M^5)$  has infinitely many path components.

**Proof** (1) implies (2) by [Lemma 1.3](#) and [Corollary 1.12](#). If we assume (2), [Lemma 1.3](#) implies that  $M$  is a Type III manifold with  $\pi_1$  acting trivially on  $\pi_2$  and  $H_2(M, \mathbb{Z})$  torsion-free. [[26](#), [Theorem 3.7](#)] shows that every such manifold is diffeomorphic to  $X(q) \#_{S^1} (\#^k (S^2 \times S^2) \times S^1)$  for some  $q \in \{0, \dots, 8\}$  and some  $k \in \mathbb{Z}_{\geq 0}$ . If  $k = 0$  and  $q$  is odd,  $b_2(X(q)) = 0$  and using [Theorem 1.11](#),  $M$  must be diffeomorphic to  $\mathbb{R}P^5 = X(1)$ .

By [[26](#), [Theorem 3.7](#)],  $M = X(q) \#_{S^1} (\#^k (S^2 \times S^2) \times S^1)$  is an orientable Type III manifold with  $\pi_1(M)$  acting trivially on  $\pi_2(M)$ ,  $H_2(M, \mathbb{Z})$  torsion-free, and  $b_2(M) = 2k + (1 + (-1)^q)/2$ . Thus by [Theorem 1.11](#), (3) implies (1).

Now assume  $M$  satisfies the conditions. If  $b_2(M) > 0$ , then by [Lemma 1.3](#) the integers  $a, b$  in (2) must satisfy  $a + b \geq 2$ . Then [Theorem A](#) implies that  $\mathfrak{M}_{\text{Ric}>0}(M)$  has infinitely many path components. By [Theorem 1.11](#), if  $b_2(M) = 0$ , then (1) implies that  $M \cong \mathbb{R}P^5$ .  $\mathfrak{M}_{\text{Ric}>0}(\mathbb{R}P^5)$  is shown to have infinitely many path components in [[14](#)]. □

**Remark 1.14** • By the discussion preceding [[26](#), [Theorem 3.7](#)] the manifolds described in [Corollary 1.13](#) can also be constructed by applying  $\#_{S^1}$  to the homotopy  $\mathbb{R}P^5$ 's. For instance,

$$X(q) \#_{S^1} (\#^k (S^2 \times S^2) \times S^1) \cong X(q) \#_{S^1} X(0) \#_{S^1} \dots \#_{S^1} X(0).$$

- It is shown in [14] that  $\mathfrak{M}_{\text{Ric}>0}$  also has infinitely many components for the homotopy  $\mathbb{R}P^5$ 's  $X(3)$ ,  $X(5)$  and  $X(7)$ .
- A characteristic submanifold  $P \subset X(q)\#_{S^1}(\#^k(S^2 \times S^2) \times S^1)$  has class  $q \in \Omega_4^{\text{Pin}^+}/\pm = \{0, \dots, 8\}$ . If we fix a nonspin simply connected 4-manifold  $B^4$ , then a Type III total space of a principal  $S^1$  bundle over  $B$  will be diffeomorphic to  $X(q)\#_{S^1}(\#^k(S^2 \times S^2) \times S^1)$ . Using Table 1 we see that  $q$  must satisfy  $q = \pm \text{sign } B \pmod 4$ . It follows that there are 2, 3 or 4 choices of  $q$ , and the same number of diffeomorphism types of Type III total spaces, if  $\text{sign}(B)$  is 2, 0 or  $\pm 1 \pmod 4$ , respectively. The value of  $q$  can be determined, up to two possibilities, using Lemma 1.7. The set of diffeomorphism types of Type I total spaces is more complicated, but can be computed using Theorem 1.11 and [26, Theorem 3.7]. If  $B^4$  is a simply connected spin 4-manifold, there exists a unique diffeomorphism type of total spaces with  $\pi_1 = \mathbb{Z}_2$ , represented by  $(S^2 \times \mathbb{R}P^3)\#_{S^1}(\#^{(b_2(B)-2)/2}(S^2 \times S^2) \times S^1)$ .

Using a result of Gilkey, Park and Tuschmann, we can lift metrics from the quotients described by Corollary 1.12 to prove the following:

**Corollary 1.15** *Let  $M$  be a 5-manifold with  $\pi_1(M) = \mathbb{Z}_2$  admitting a free  $S^1$  action with a simply connected quotient. Then  $M$  admits a metric with positive Ricci curvature.*

**Proof** In [37] Sha and Yang put a metric of positive Ricci curvature on  $\#^c S^2 \times S^2$ . A modification of Perelman's construction in [34] puts such a metric on  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ ; see Lemma 3.10. Corollary 1.12 shows that  $M^5$  admits a free  $S^1$  action with quotient  $B^4$  diffeomorphic to one of those manifolds. Gilkey, Park and Tuschmann [20] showed that if  $B^4$  admits  $\text{Ric} > 0$ ,  $M^5$  is the total space of a principal bundle over  $B^4$  with compact connected structure group  $G$ , and  $\pi_1(M^5)$  is finite, then  $M$  admits a  $G$ -invariant metric with  $\text{Ric} > 0$ . In this case  $G = S^1$ ,  $\pi_1(M) = \mathbb{Z}_2$  and the corollary follows.  $\square$

The corresponding result in the simply connected case was proved by Corro and Galaz-Garcia in [11]. By Lichnerowicz's theorem, many simply connected 4-manifolds, such as a K3 surface, do not admit even positive scalar curvature. It is interesting to note that Corollary 1.15 and the results of [11] imply that total spaces with  $\pi_1 = 0$  or  $\mathbb{Z}_2$  of principal  $S^1$  bundles over such manifolds nonetheless admit metrics of positive Ricci curvature.

## 2 The $\eta$ invariant

We use the  $\eta$  invariant of the  $\text{spin}^c$  Dirac operator, which we define in this section, to distinguish components of geometric moduli spaces. A manifold  $M$  is  $\text{spin}^c$  if there exists a complex line bundle  $\lambda$  over  $M$  such that the frame bundle of  $TM \oplus \lambda$ , a principal  $\text{SO}(n) \times U(1)$  bundle, lifts to a principal  $\text{Spin}^c(n) = \text{Spin}(n) \times_{\mathbb{Z}_2} U(1)$  bundle. A manifold is  $\text{spin}^c$  if and only if the second Stiefel–Whitney class  $w_2(TM)$  is the image of an integral class  $c \in H^2(M, \mathbb{Z})$  under the map  $H^2(M, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Z}_2)$ . In

this case  $c$ , which we call the canonical class of the  $\text{spin}^c$  structure, is the first Chern class of  $\lambda$ , which we call the canonical bundle.

Using complex representations of  $\text{Spin}^c(n)$  we form  $\text{spin}^c$  spinor bundles and equip them with actions of the complex Clifford algebra bundle  $\mathbb{C}\ell(TM)$ . When the dimension of  $M$  is even there is a unique irreducible such bundle  $S$  with a natural grading  $S = S^+ \oplus S^-$ . Given a metric  $g$  on  $M$  and a unitary connection  $\nabla$  on  $\lambda$ , we can construct a spinor connection  $\nabla^S$  on  $S$ , compatible with Clifford multiplication, and a  $\text{spin}^c$  Dirac operator  $D_{g,\nabla}^c$  acting on sections of  $S$ . See [31, Appendix D] for details. The Bochner–Lichnerowicz identity for this operator is

$$(2.1) \quad (D_{g,\lambda}^c)^2 = (\nabla^S)^* \nabla^S + \frac{1}{4} \text{scal}(g) + \frac{i}{2} F^\nabla,$$

where the complex two-form  $F^\nabla$  is the curvature of  $\nabla$ . This form acts on the spinor bundle  $S$  by way of the vector bundle isomorphism  $\Lambda T^*M \rightarrow \Lambda TM \rightarrow \mathbb{C}\ell(TM)$  given by  $g$ . The operator  $(\nabla^S)^* \nabla^S$  is nonnegative definite with respect to the  $L^2$  inner product on a closed manifold or a compact manifold with boundary on which the Atiyah–Patodi–Singer boundary conditions have been applied. See [2, Theorem 3.9] for details. The remaining term  $\frac{1}{4} \text{scal}(g) + \frac{i}{2} F^\nabla$  is positive definite if

$$(2.2) \quad \text{scal}(g) > 2|F^\nabla|_g,$$

where the norm  $|\cdot|_g$  is the operator norm on  $\mathbb{C}\ell(TM)$  acting on  $S$ . In particular,  $\ker(D_{g,\nabla}^c) = 0$  if (2.2) is satisfied. For a later purpose we note that for  $\omega \in \Omega^2(M, \mathbb{C})$  and an orthonormal basis  $\{e_i\}$  of  $TM$  with respect to  $g$ , we have

$$(2.3) \quad |\omega|_g \leq \sum_{i < j} |\omega(e_i, e_j)|.$$

Suppose  $W$  is a  $\text{spin}^c$  manifold with boundary  $\partial W = M$ , with  $\lambda$  and  $c$  defined on  $W$  as above.  $W$  induces a  $\text{spin}^c$  structure on  $M$  with canonical class  $c|_{\partial W}$  and canonical bundle  $\lambda|_{\partial W}$ . Choose a metric  $h$  on  $W$  and a connection  $\nabla$  on  $\lambda$  which are product-like near  $\partial W$ , ie

$$h = h|_{\partial W} + dr^2 \quad \text{and} \quad \nabla = \text{proj}_M^*(\nabla|_{\partial W})$$

on a collar neighborhood  $U \cong M \times I$ , where  $I$  is an interval with coordinate  $r$ . Applying the Atiyah–Patodi–Singer boundary conditions, the Atiyah–Patodi–Singer index theorem [1] states that

$$(2.4) \quad \text{ind}(D_{h,\nabla}^c|_{S^+}) = \int_W e^{c_1(\nabla)/2} \hat{A}(p(g)) - \frac{1}{2}(\dim(\ker(D_{h|_{\partial W}, \nabla|_{\partial W}}^c))) + \eta(D_{h|_{\partial W}, \nabla|_{\partial W}}^c).$$

Here  $c_1(\nabla)$  and  $p(g)$  are the Chern–Weil Chern and Pontryagin forms constructed from the curvature tensors of the connection and metric, respectively.  $\hat{A}$  is the polynomial in the Pontryagin forms and  $D_{h|_{\partial W}, \nabla|_{\partial W}}^c$  is the  $\text{spin}^c$  Dirac operator on  $M$  constructed using the induced metric and connection.

The  $\eta$  invariant is an analytic invariant of the spectrum of an elliptic operator defined in [1]. Given an elliptic differential operator  $D$  with spectrum  $\{\lambda_i\}$ , we define a complex function

$$\eta(D, s) = \sum_{\lambda_i \neq 0} \text{sign}(\lambda_i) |\lambda_i|^{-s}.$$

One shows that the function is analytic when the real part of  $s$  is large, and Atiyah, Patodi and Singer showed that it can be analytically continued to a meromorphic function which is analytic at 0. Thus we define  $\eta(D) = \eta(D, 0)$ . If a diffeomorphism  $\phi$  preserves the  $\text{spin}^c$  structure, then  $D_{\phi^*g, \phi^*\nabla}^c$  is conjugate to  $D_{g, \nabla}^c$ , and hence they have the same spectrum and the same values of  $\eta$ . We will use (2.4) to calculate  $\eta$  for an operator  $D_{g, \bar{\nabla}}$  on a manifold  $M$  by finding a suitable  $W$  with  $\partial W = M$  and extending  $g$  and  $\bar{\nabla}$  to product-like  $h$  and  $\nabla$  on  $W$ .

Kreck and Stolz combined the  $\eta$  invariant with information about the Chern–Weil forms of the metric to get an invariant for metrics on  $(4n+3)$ -dimensional spin manifolds. We prove that the  $\eta$  invariant alone provides the desired invariant for certain  $(4n+1)$ -dimensional  $\text{spin}^c$  manifolds.

**Theorem 2.5** *Let  $M^{4n+1}$  be a closed  $\text{spin}^c$  manifold with canonical class  $c \in H^2(M, \mathbb{Z})$  and canonical bundle  $\lambda$ . Suppose  $c$  and the Pontryagin classes  $p_i(TM)$  are torsion and  $g_t$ , where  $t \in [0, 1]$ , is a smooth path of metrics on  $M$  with  $\text{scal}(g_t) > 0$ . If  $\nabla_0$  and  $\nabla_1$  are flat unitary connections on  $\lambda$ , then*

$$\eta(D_{g_0, \nabla_0}^c) = \eta(D_{g_1, \nabla_1}^c).$$

**Proof** Modifying  $g_t$  if necessary, we assume it is a constant path for  $t$  near 0 and 1. Given  $L \in \mathbb{R}_{>0}$ , define a smooth metric  $g$  on  $M \times [0, 1]$  by

$$g = g_t + L^2 dt^2.$$

Then  $g$  is product-like near  $M \times \{0, 1\}$ . One sees that  $\text{scal}(g)$  differs from  $\text{scal}(g_t)$  by terms depending on the second fundamental form of each slice  $M \times \{t\}$ , but the second fundamental form tends to 0 as  $L \rightarrow \infty$ , so for large  $L$  we have  $\text{scal}(g) > 0$ .

The difference of unitary connections on a complex line bundle is an imaginary one-form. Define  $\alpha \in \Omega(M)$  such that

$$i\alpha = \nabla_1 - \nabla_0.$$

Since both connections are flat,  $d\alpha = 0$ . Let  $\pi: M \times [0, 1] \rightarrow M$  be the projection and let  $f: M \times [0, 1] \rightarrow [0, 1]$  be the projection onto  $[0, 1]$  followed by a smooth function which is 0 in a neighborhood of 0 and 1 in a neighborhood of 1. Define a connection on  $\pi^*\lambda$  by

$$\nabla = \pi^*\nabla_0 + if\pi^*\alpha.$$

Then, since  $\nabla_0$  is flat,

$$F^\nabla = idf \wedge \pi^*\alpha.$$

Let  $e_i$  be an orthonormal frame for  $g$  at a point  $(p, t)$ , such that  $e_1 = (1/L)\partial_t$ . Then

$$2 \sum_{i < j} |(df \wedge \alpha)(e_i, e_j)| = \frac{2\partial_t f}{L} \sum_{i > 1} \alpha(e_i).$$

Since  $e_i$  for  $i > 2$  is tangent to  $M \times \{t\}$ , it does not depend on  $L$ . Using (2.3), for large  $L$  we have

$$\text{scal}(g) > 2|F^\nabla|_g.$$

The definition of  $f$  ensures that  $\nabla$  is product-like near  $\partial(M \times I)$ . Then by (2.1),  $D_{g,\nabla}^c$  has trivial kernel and  $\text{ind}(D_{g,\nabla}^c|_{S^+}) = 0$ .

Since  $F^{\nabla_i} = 0$  for  $i = 1, 2$ ,

$$\text{scal}(g_i) > 0 = 2|F^{\nabla_i}|_{g_i}$$

and hence (2.1) implies  $\ker D_{g_i,\nabla_i}^c = \{0\}$ . We now apply the Atiyah–Patodi–Singer index theorem (2.4). The boundary of  $M \times I$  is two copies of  $M$  with opposite orientations. The spectrum of the Dirac operator on  $M \times \{0, 1\}$  is the union of the spectra on  $M \times \{0\}$  and  $M \times \{1\}$ , and the  $\eta$  invariant is the sum of the two  $\eta$  invariants. When we change the orientation of an odd-dimensional manifold, the Dirac operator changes by a sign. Thus the Atiyah–Patodi–Singer theorem yields

$$\begin{aligned} \text{ind}(D_{g,\nabla}^c|_{S^+}) \\ = \int_{M \times [0,1]} e^{c_1(\nabla)/2} \widehat{A}(p(g)) - \frac{1}{2} (\dim(\ker(D_{g_0,\nabla_0}^c)) + \dim(\ker(D_{g_1,\nabla_1}^c)) + \eta(D_{g_0,\nabla_0}^c) - \eta(D_{g_1,\nabla_1}^c)), \end{aligned}$$

and hence

$$\eta(D_{g_1,\nabla_1}^c) - \eta(D_{g_0,\nabla_0}^c) = 2 \int_{M \times [0,1]} e^{c_1(\nabla)} \widehat{A}(p(g)).$$

Since  $\pi_1^*c$  is torsion,  $c_1(\nabla)$  is exact. Because  $\nabla$  is flat near the boundary,  $c_1(\nabla)|_{\partial(M \times I)} = 0$ . Furthermore,  $g$  is product-like near the boundary so  $p(g)|_{M \times \{i\}} = p(g_i)$ . Since the real Pontryagin classes of  $M$  vanish,  $p_j(g_i)$  is exact for  $j > 0$ . By Stokes' theorem, and since the dimension of  $M$  is  $4n + 1$ , the integral vanishes.  $\square$

As a corollary we show how to use the  $\eta$  invariant to detect path components of moduli spaces of metrics with curvature conditions no weaker than positive scalar curvature.

**Corollary 2.6** *Let  $M$  be as in Theorem 2.5. Let  $(g_i, \nabla_i)$  be a sequence of Riemannian metrics  $g_i$  with  $\text{Ric}(g_i) > 0$ , and flat connections  $\nabla_i$  on  $\lambda$  such that  $\{\eta(D_{g_i,\nabla_i}^c)\}_i$  is infinite. Then  $\mathfrak{M}_{\text{Ric}>0}(M)$  and  $\mathfrak{M}_{\text{scal}>0}(M)$  have infinitely many path components.*

**Proof** Let  $\text{Diff}^c(M)$  be the set of diffeomorphisms of  $M$  which fix the  $\text{spin}^c$  structure. For  $g \in \mathfrak{R}_{\text{scal}>0}$  let  $[g]$  represent the image in  $\mathfrak{M}_{\text{scal}>0}$  and  $[g]^c$  the image in  $\mathfrak{R}_{\text{scal}>0}/\text{Diff}^c(M)$ . It follows from Ebin's slice theorem [17; 9] that if  $[g_i]$  and  $[g_j]$  are in the same connected component of  $\mathfrak{R}_{\text{scal}>0}/\text{Diff}^c(M)$  then  $g_i$  and  $\phi^*g_j$  are in the same path component of  $\mathfrak{R}_{\text{scal}>0}$  for some  $\phi \in \text{Diff}^c(M)$ . Then there is a path between them maintaining positive scalar curvature, and by Theorem 2.5 and the  $\text{spin}^c$  diffeomorphism invariance of  $\eta$  we have  $\eta(D_{g_i,\nabla_i}^c) = \eta(D_{\phi^*g_j,\phi^*\nabla_j}^c) = \eta(D_{g_j,\nabla_j}^c)$ . Since  $\{\eta(D_{g_i,\nabla_i}^c)\}$  is infinite,  $\mathfrak{R}_{\text{scal}>0}/\text{Diff}^c(M)$  has infinitely many components.

Any diffeomorphism  $\phi$  pulls back the  $\text{spin}^c$  structure to another one with canonical class  $\phi^*c$ , a torsion class in  $H^2(M, \mathbb{Z})$ . There are finitely many such classes. The finite group  $H^1(M, \mathbb{Z}_2)$  indexes the  $\text{spin}^c$  structures associated to each class. Thus the orbit of the  $\text{spin}^c$  structure under  $\text{Diff}(M)$  and

the set  $\text{Diff}(M)/\text{Diff}^c(M)$  are finite. The fibers of  $\mathfrak{M}_{\text{scal}>0}/\text{Diff}^c(M) \rightarrow \mathfrak{M}_{\text{scal}>0}$  are no larger than  $\text{Diff}(M)/\text{Diff}^c(M)$ , implying that  $\mathfrak{M}_{\text{scal}>0}$  has infinitely many components.

The proof is identical for  $\mathfrak{M}_{\text{Ric}>0}$  since  $\text{Ric} > 0$  implies  $\text{scal} > 0$ . □

### 3 The $\eta$ invariant in dimension $4n + 1$ with free $S^1$ actions

In this section we prove [Theorem A](#). We want to use the Atiyah–Patodi–Singer index theorem to calculate the  $\eta$  invariant of a metric on  $M$ . Many authors have computed  $\eta$  and related invariants on spin manifolds  $M$  by extending metrics to manifolds  $W$  with boundary diffeomorphic to  $M$ . If the extension has positive scalar curvature, the index of the Dirac operator will vanish. In the  $\text{spin}^c$  case, we must also extend an auxiliary connection. A difficulty arises when the extended connection cannot be flat because the canonical class of the  $\text{spin}^c$  structure on  $W$  is not torsion. Then the metric and connection must satisfy (2.2). The following theorem, which we prove in [Section 4](#), illustrates how to use certain free  $S^1$  actions on  $M$  to accomplish this.

**Theorem 3.1** *Let  $S^1$  act freely on  $M$  by isometries of a Riemannian metric  $g_M$  with  $\text{scal}(g_M) > 0$  and assume  $\pi_1(M)$  is finite. Let  $B = M/S^1$  be the quotient and  $\rho: W = M \times_{S^1} D^2 \rightarrow B$  the associated disc bundle. Suppose the first Chern class of the principal  $S^1$  bundle  $\pi: M \rightarrow B$  is  $\ell d$  for  $d \in H^2(B, \mathbb{Z})$  and  $\ell \in \mathbb{Z}$ . If  $\lambda$  is the complex line bundle over  $W$  with first Chern class  $\rho^*d$ , then there exists a metric  $g_W$  on  $W$  and a connection  $\nabla$  on  $\lambda$  such that*

$$(3.2) \quad \text{scal}(g_W) > \ell |F^\nabla|_{g_W}.$$

Furthermore there is a collar neighborhood  $V \cong M \times [0, N]$  of  $\partial W \cong M$  such that for  $t \in [0, N]$  near 0,  $g_W$  is a product metric

$$(3.3) \quad g_W \cong g_M + dt^2,$$

and

$$(3.4) \quad \nabla \cong \text{proj}_{V, M}^* \bar{\nabla},$$

where  $\bar{\nabla}$  is any flat unitary connection on  $\lambda|_{\partial W}$ .

Notice that here there are no restrictions on the dimension or Pontryagin classes of  $M$ ,  $d$  need not be primitive, and no  $\text{spin}^c$  structure is required. We next use [Theorem 3.1](#) and (2.4) to calculate  $\eta$  for  $S^1$ -invariant metrics on certain  $\text{spin}^c$  manifolds in dimensions  $4n + 1$ .

**Theorem 3.5** *Let  $S^1$  act freely on a  $4n + 1$  manifold  $M$  by isometries of a Riemannian metric  $g$  with  $\text{scal}(g) > 0$ . Assume  $\pi_1(M)$  is finite and let  $B = M/S^1$  be the quotient. Suppose the first Chern class of the principal bundle  $S^1 \rightarrow M \xrightarrow{\pi} B$  is  $\ell d$ , where  $\ell$  is a positive even integer and  $w_2(TB) = d \pmod 2$ .*



Finally assume the real Pontryagin classes of  $M$  vanish. Then  $M$  admits a  $\text{spin}^c$  structure with canonical class  $\pi^*d$ . If  $\bar{\nabla}$  is a flat connection on the canonical bundle of this  $\text{spin}^c$  structure and  $D_{g,\bar{\nabla}}^c$  is the  $\text{spin}^c$  Dirac operator, then

$$\eta(D_{g,\bar{\nabla}}^c) = \left\langle \frac{\sinh(\frac{1}{2}d)\widehat{A}(TB)}{\sinh(\frac{1}{2}\ell d)}, [B] \right\rangle.$$

When  $n = 1$ ,

$$(3.6) \quad \eta(D_{g,\bar{\nabla}}^c) = \left\langle -\frac{(\ell^2 - 1)d^2 + p_1(TB)}{24\ell}, [B] \right\rangle.$$

**Proof** Since  $TM$  is the direct sum of  $\pi^*TB$  and a trivial bundle generated by the action field of the  $S^1$  action,

$$w_2(TM) = \pi^*w_2(TB) = \pi^*d \pmod{2}.$$

Let  $\mu$  be the complex line bundle over  $B$  associated to  $\pi: M \rightarrow B$ . Let  $W = M \times_{S^1} D^2$  and let  $\rho: W \rightarrow B$  be the disc bundle associated to  $\pi: M \rightarrow B$ . Then  $TW = \rho^*(TB \oplus \mu)$  and, since  $\ell$  is even,

$$w_2(TW) = \rho^*(d + \ell d) \pmod{2} = \rho^*d \pmod{2}.$$

It follows that  $W$  admits a  $\text{spin}^c$  structure with canonical class  $\rho^*d$ . We call the canonical bundle  $\lambda$ . The  $\text{spin}^c$  structure on  $W$  induces one on  $M$  with canonical class  $\pi^*d$ .

Then  $M$ ,  $W$  and  $\lambda$  satisfy the hypotheses of [Theorem 3.1](#). We construct the metric  $g_W$  on  $W$  and connection  $\nabla$  on  $\lambda$  as in the theorem such that  $g_W|_M = g_M$  and  $\nabla|_M = \bar{\nabla}$ . Define the  $\text{spin}^c$  Dirac operator  $D_{g_W,\nabla}^c$  on  $W$  and  $D_{g_M,\bar{\nabla}}^c$  as in [Section 2](#). Given that  $g_W$  and  $\nabla$  are product-like near  $\partial W$ , we can apply [\(2.4\)](#). Since  $\bar{\nabla}$  is flat,

$$\text{scal}(g_M) > 2|F^{\bar{\nabla}}|_{g_M} = 0,$$

and, by [\(3.2\)](#),

$$\text{scal}(g_W) > \ell|F^\nabla|_{g_W} \geq 2|F^\nabla|_{g_W}.$$

Then [\(2.1\)](#) implies that  $\text{ind}(D_{g_W,\nabla}^c) = 0$  and  $\ker(D_{g_M,\bar{\nabla}}^c) = \{0\}$ . It follows from [\(2.4\)](#) that

$$(3.7) \quad \eta(D_{g_M,\bar{\nabla}}^c) = 2 \int_W e^{c_1(\nabla)/2} \widehat{A}(p(g_W)).$$

To evaluate that integral, we use [\[29, Lemma 2.7\]](#):

**Lemma 3.8** [\[30\]](#) *Let  $W$  be a manifold with boundary, and let  $\alpha$  and  $\beta$  be closed forms on  $W$  such that  $\alpha|_{\partial W} = d\hat{\alpha}$  and  $\beta|_{\partial W} = d\hat{\beta}$ . Then*

$$\int_W \alpha \wedge \beta = \int_{\partial W} \hat{\alpha} \wedge \hat{\beta} + \langle j^{-1}(\alpha) \cup j^{-1}(\beta), [W, \partial W] \rangle,$$

where  $j^{-1}$  represents any preimage under the long exact sequence map

$$j: H^*(W, \partial W; \mathbb{Q}) \rightarrow H^*(W, \mathbb{Q}).$$

To apply Lemma 3.8 to (3.7), let  $\alpha = e^{c_1(\nabla)/2}$  and  $\beta = \widehat{A}(p(g_W))$ . Since  $g_W$  is product-like near the boundary,  $p_i(g_W)|_{\partial W} = p_i(g_M)$ . For  $i > 0$ ,  $p_i(g_M)$  is exact by the assumption on the Pontryagin classes of  $M$ . Since  $c_1(\nabla)|_{\partial W} = c_1(\bar{\nabla})$  and  $\bar{\nabla}$  is flat, we can choose  $\widehat{\alpha} = 0$ . The form  $c_1(\nabla)$  represents the cohomology class  $c_1(\lambda) = \rho^*d$ . Thus

$$\eta(D_{g,\bar{\nabla}}^c) = 2\langle j^{-1}[e^{\rho^*d/2}] \cup j^{-1}[\widehat{A}(TW)], [W, \partial W] \rangle.$$

The following cup product diagram commutes:

$$\begin{CD} H^s(W, \partial W) \oplus H^t(W, \partial W) @>\cup>> H^{s+t}(W, \partial W) \\ @VV(\text{Id}, j)V @VVV \\ H^s(W, \partial W) \oplus H^t(W) @>\cup>> H^{s+t}(W, \partial W) \end{CD}$$

Thus,

$$(3.9) \quad \eta(D_{g,\bar{\nabla}}^c) = 2\langle j^{-1}[e^{\rho^*d/2}] \cup [\widehat{A}(TW)], [W, \partial W] \rangle.$$

Since the terms of  $\widehat{A}(TW)$  have degree  $4k$ , with  $k \in \mathbb{Z}$ , and the dimension of  $W$  is  $4n + 2$ , only terms of degree  $4k + 2$  in  $e^{\rho^*d/2}$  will contribute. In those degrees,  $e^{\rho^*d/2} = \sinh(\rho^*d/2)$  as power series.

Since  $TW = \rho^*(TB \oplus \mu)$ , we have  $\widehat{A}(TW) = \rho^*(\widehat{A}(TB)\widehat{A}(\mu))$ . For the complex line bundle  $\mu$ , we have

$$\widehat{A}(\mu) = \frac{\frac{1}{2}c_1(\mu)}{\sinh(\frac{1}{2}c_1(\mu))} = \frac{\ell d}{2 \sinh(\frac{1}{2}\ell d)}$$

as a formal power series. The series  $\sinh(\frac{1}{2}d)$  is divisible by  $d$ , so

$$\rho^*\left(\frac{\sinh(\frac{1}{2}d)}{\ell d}\right) \in H^*(W, \mathbb{Q}).$$

Let  $\Phi \in H^2(W, \partial W, \mathbb{Z})$  be the Thom class of  $\rho: W \rightarrow B$ . Then  $j(\Phi) = \rho^*c_1(\mu) = \rho^*(\ell d)$ . By means of another commutative diagram

$$\begin{CD} H^*(W, \partial W) \oplus H^*(W) @>\cup>> H^*(W, \partial W) \\ @VV(j, \text{Id})V @VVjV \\ H^*(W) \oplus H^*(W) @>\cup>> H^*(W) \end{CD}$$

we see

$$j\left(\Phi \cup \rho^*\left(\frac{\sinh(\frac{1}{2}d)}{\ell d}\right)\right) = \rho^*\left(\ell d \cup \frac{\sinh(\frac{1}{2}d)}{\ell d}\right) = \rho^* \sinh(\frac{1}{2}d).$$

Substituting into (3.9),

$$\begin{aligned} \eta(D_{g,\bar{\nabla}}^c) &= 2\left\langle \Phi \cup \rho^*\left(\frac{\sinh(\frac{1}{2}d)}{\ell d}\right) \cup \rho^*\left(\frac{\widehat{A}(TB) \cdot \ell d}{2 \sinh(\frac{1}{2}\ell d)}\right), [W, \partial W] \right\rangle \\ &= \left\langle \Phi \cup \rho^*\left(\frac{\sinh(\frac{1}{2}d)\widehat{A}(TB)}{\sinh(\frac{1}{2}\ell d)}\right), [W, \partial W] \right\rangle. \end{aligned}$$

The Thom isomorphism yields

$$\eta(D_{g, \bar{\nu}}^c) = \left\langle \frac{\sinh(\frac{1}{2}d)\widehat{A}(TB)}{\sinh(\frac{1}{2}\ell d)}, [B] \right\rangle.$$

When  $n = 1$  the dimension of  $B$  is four and we have, as series in  $H^*(B, \mathbb{Z})$ ,

$$\widehat{A}(TB) = 1 - \frac{1}{24}p_1(TB) \quad \text{and} \quad \frac{\sinh(\frac{1}{2}d)}{\sinh(\frac{1}{2}\ell d)} = \frac{1}{\ell} \left( 1 - \frac{1}{24}(\ell^2 - 1)d^2 \right).$$

Multiplying and isolating terms of degree four yields (3.6). □

We are now ready to prove [Theorem A](#). We first construct metrics of  $\text{Ric} > 0$  on  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ . Perelman [34] constructed a metric with  $\text{Ric} > 0$  on arbitrary connected sums of  $\mathbb{C}P^2$  with its standard orientation. More details on Perelman’s proof can be found in [10; 8]. With a slight adjustment to the construction one can reverse the orientation on some of the copies of  $\mathbb{C}P^2$ , proving the following.

**Lemma 3.10**  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$  admits a metric with positive Ricci curvature for all  $a$  and  $b$ .

**Proof** In [34], Perelman puts a metric on  $\#^c \mathbb{C}P^2$  for all values of  $c$ . The construction involves  $c$  copies of  $\mathbb{C}P^2$  attached to a central  $S^4$  by “necks”  $S^3 \times I$ . The metric on the necks is of the form

$$ds^2 = dt^2 + A^2(t, x) dx^2 + B^2(t, x) d\sigma^2,$$

where  $t$  is the coordinate on the interval  $I$ ; see [34, page 159]. Furthermore,  $S^3$  is represented as the product of  $S^2$  and an interval with the top and bottom each identified to a point, and  $x$  is the coordinate on that interval, while  $d\sigma^2$  is the standard metric on  $S^2$ .

An orientation-reversing isometry of  $d\sigma^2$ , such as the antipodal map, extends naturally to a diffeomorphism of  $\phi: S^3 \rightarrow S^3$ , which induces an isometry of  $ds^2$ . Let  $c = a + b$ , and take Perelman’s metric on  $\#^c \mathbb{C}P^2$ . For  $b$  of the necks, we cut along a copy of  $S^3$  and reglue with  $\phi$  rather than the identity. Because  $\phi$  reverses orientation, the resulting manifold is  $\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ . Because  $\phi$  induces an isometry on  $S^3 \times I$ , the same metrics on the pieces extend smoothly over the gluing, completing the proof of the lemma. □

Let  $M^5$  satisfy the hypotheses of [Theorem A](#). By [Lemma 1.3](#),  $M$  is Type III and by [Theorem 1.11](#)  $M$  is the total space of infinitely many nonisomorphic principal  $S^1$  bundles  $\pi_k: M^5 \rightarrow B^4 = \#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}$ . From the proof of [Theorem 1.11](#) we see that the first Chern class of  $\pi_k$  is  $2d_k$ , where

$$d_k = (1 + 2k, 1, \dots, 1) \in H^2(\#^a \mathbb{C}P^2 \#^b \overline{\mathbb{C}P^2}, \mathbb{Z}) \cong \mathbb{Z}^{a+b}$$

for a certain infinite set of integers  $k$ .

Using the result of [20] (see [Corollary 1.15](#)) we see that since  $B$  admits a metric of positive Ricci curvature by [Lemma 3.10](#),  $\pi_k: M \rightarrow B$  is a principal  $S^1$  bundle, and  $\pi_1(M)$  is finite, then for each  $k$   $M$  admits a metric  $g_k$  with  $\text{Ric}(g_k) > 0$  such that the  $S^1$  action corresponding to the principal bundle  $\pi_k: M \rightarrow B$  acts by isometries of  $g_k$ .

Using the Gysin sequence it follows that  $H^4(M, \mathbb{R}) = 0$  and  $M$ ,  $g_k$  and  $B$  satisfy the hypotheses of [Theorem 3.5](#) with  $g_M = g_k$ ,  $d = d_k$ ,  $\ell = 2$  and  $\bar{\nabla}$  any flat connection on the canonical bundle of the  $\text{spin}^c$  structure. By [\(3.6\)](#) we have

$$\eta(D_{g_k, \bar{\nabla}}^c) = -\frac{1}{16}(\langle c_k^2, [B] \rangle + \text{sign } B) = -\frac{1}{16}(\pm 4k^2 \pm 4k + 2 \text{sign } B),$$

using the fact that  $\langle \frac{1}{3}p_1(TB), [B] \rangle = \langle L(TB), [B] \rangle$  is equal to the signature of  $B$ .

Thus  $\eta(D_{g_k, \bar{\nabla}}^c)$  is a nontrivial polynomial in  $k$  and takes on infinitely many values for the infinite set of integers  $k$ . [Corollary 2.6](#) implies that  $\mathfrak{M}_{\text{Ric}>0}(M)$  has infinitely many components, completing the proof of [Theorem A](#). □

Note that [Corollary 2.6](#) also implies that  $\mathfrak{M}_{\text{scal}>0}(M)$  has infinitely many components.

### 4 Metric and connection

In this section we prove [Theorem 3.1](#). We first set up notation for the tangent space to  $W$ . We consider  $D^2$  to be the unit disc in  $\mathbb{C}$ . Let  $\sigma: M \times D^2 \rightarrow W$  be the quotient map so  $\sigma(p, x) = [p, x]$ . Then  $\rho([p, x]) = \pi(p)$ . The metric  $g_M$  and the  $S^1$  action induce an orthogonal splitting  $T_p M = \bar{H}_p \oplus \bar{V}_p$  into horizontal space  $\bar{H}_p$  and vertical space  $\bar{V}_p$  of  $\pi$ . Define horizontal and vertical spaces of  $\rho$  to be

$$H_{[p,x]} = \sigma_*(\bar{H}_p \oplus \{0\}) \quad \text{and} \quad V_{[p,x]} = \sigma_*(\{0\} \oplus T_x D^2)$$

for  $p \in M$  and  $x \in D^2$ .

These are well defined since for  $z \in S^1$ ,  $\bar{H}_{z p} = z_* \bar{H}_p$  and  $T_{z x} D^2 = z_* T_x D^2$ . One can use a local section of  $\sigma$  to see that  $H_{[p,x]}$  and  $V_{[p,x]}$  are smooth distributions on  $W$ . Note that  $V_{[p,x]}$  is the tangent space to the fiber  $\rho^{-1}(\pi(p)) = \sigma(\{p\} \times D^2)$  and  $T_{[p,x]} W = H_{[p,x]} \oplus V_{[p,x]}$ . Away from the zero section of  $\rho$ ,  $V_{[p,x]}$  is spanned by

$$W_r = \sigma_*(0, \partial_r) \quad \text{and} \quad W_\theta = \sigma_*(0, \partial_\theta).$$

These are well-defined smooth vector fields since  $\partial_\theta$  and  $\partial_r$  are  $S^1$ -invariant vector fields on  $D^2$ .

Fix  $0 < L < 1$  and define a diffeomorphism

$$\tau: M \times [L, 1] \hookrightarrow M \times D^2 \xrightarrow{\sigma} W$$

of  $M \times [L, 1]$  to a collar neighborhood  $U$  of  $\partial W$ . Let  $t$  be the coordinate on  $[L, 1]$  and, in a slight abuse of notation, let  $\text{proj}_{U, M}: M \times [L, 1] \rightarrow M$  be the projection. Thus,

$$\rho \circ \tau = \pi \circ \text{proj}_{U, M}, \quad \tau_*(\bar{H}_p \oplus \{0\}) = H_{[p,x]}, \quad \tau_*(0, \partial_t) = W_r.$$

Let

$$X^*(p) = \left. \frac{d}{dt} \right|_{t=0} e^{it} \cdot p$$

be the action field of the  $S^1$  action on  $M$ , which spans  $\bar{V}_p$ . Then, since  $\sigma_*(X^*, \partial_\theta) = 0$ ,

$$\tau_*(X^*, 0) = -W_\theta.$$

Furthermore,  $\tau|_{M \times \{1\}}$  identifies  $M$  and  $\partial W$ , sending  $\bar{H}_p$  to  $H_{[p,1]}$  and  $X^*$  to  $-W_\theta$ .

We keep track of the maps in the following diagram:

$$\begin{array}{ccc} & & M \times D^2 \\ & \nearrow & \downarrow \sigma \\ M \times I & \xrightarrow{\tau} & W \\ \text{proj}_{U,M} \downarrow & & \downarrow \rho \\ M & \xrightarrow{\pi} & B \end{array}$$

To construct  $g_W$  and  $\nabla$  we will use two smooth functions on the interval  $[0, 1]$ . Let  $f_1 : [0, 1] \rightarrow [0, 1]$  be a smooth monotone function which is 0 in a neighborhood of 0, and 1 in a neighborhood of  $[L, 1]$ .

For a constant  $\epsilon > 0$ , let

$$f_2(r) = -\frac{1}{2} \int_0^r f_1(t) dt - \epsilon r^3 + r.$$

One easily sees that  $f_2 > 0$  on  $(0, 1]$  for small  $\epsilon$ .

### 4.1 Metric

We define a Riemannian metric at a point  $(p, (r, \theta)) \in M \times D^2$ , where  $r$  and  $\theta$  are polar coordinates on  $D^2$ , by

$$g_{M \times D^2}(p, (r, \theta)) = g_M(p) + \epsilon^2 |X^*(p)|_{g_M}^2 \left( dr^2 + \frac{f_2(r)^2}{1 - \epsilon^2 f_2(r)^2} d\theta^2 \right).$$

By converting to Cartesian coordinates on  $D^2$ , one sees that  $g_{M \times D^2}$  is smooth as long as

$$\frac{1}{r^4} \left( \frac{f_2^2}{1 - \epsilon^2 f_2^2} - r^2 \right)$$

is a smooth function of  $r \in [0, 1]$ . This is easily seen to hold since for  $r$  near 0,  $f_2(r) = r - \epsilon r^3$ . Since  $g_{M \times D^2}$  is invariant under the diagonal action of  $S^1$  on  $M \times D^2$ , it induces a metric  $g_W$  on  $W$  such that  $g_{M \times D^2}$  and  $g_W$  make  $\sigma$  into a Riemannian submersion. Similarly, let  $g_B$  be the metric on  $B$  such that  $g_M$  and  $g_B$  make  $\pi$  into a Riemannian submersion.

**Lemma 4.1** *The metrics  $g_W$  and  $g_B$  make  $\rho$  into a Riemannian submersion.*

**Proof** With respect to  $g_{M \times D^2}$ ,  $\bar{H}_p \oplus \{0\}$  is orthogonal to  $X^*$  and  $TD^2$ . Thus  $\bar{H}_p \oplus \{0\}$  is orthogonal to the vertical space of  $\sigma$ , which is spanned by  $(X^*, \partial_\theta)$ , and to the horizontal projection of  $TD^2$  as well. It follows that with respect to  $g_W$ ,  $H_{[p,x]}$  is orthogonal to  $V_{[p,x]}$  and is the horizontal space of  $\rho$ . Finally, we have

$$g_W|_{H_{[p,x]}} \cong g_{M \times D^2}|_{\bar{H}_p \oplus \{0\}} \cong g_M|_{\bar{H}_p} \cong g_B|_{T_{\pi(p)}B}.$$

□

We first describe the induced metric on the  $D^2$  fibers of  $\rho$ .

**Lemma 4.2**  $g_W|_{\rho^{-1}(\pi(p))} \cong \epsilon^2 |X^*(p)|_{g_M} (dr^2 + f_2(r)^2 d\theta^2)$ .

**Proof** The restriction  $\sigma|_{\{p\} \times D^2}: D^2 \rightarrow \rho^{-1}(\pi(p))$  is a diffeomorphism such that  $\partial_r$  and  $\partial_\theta$  are mapped to  $W_r$  and  $W_\theta$ , respectively. Since  $\sigma$  is a Riemannian submersion with vertical space generated by  $(X^*, \partial_\theta)$ , we calculate:

$$\begin{aligned} |W_r|_{g_W}^2 &= |(0, \partial_r)|_{g_{M \times D^2}}^2 = \epsilon^2 |X^*|_{g_M}^2, \\ |W_\theta|_{g_W}^2 &= |(0, \partial_\theta)|_{g_{M \times D^2}}^2 - \frac{\langle (0, \partial_\theta), (X^*, \partial_\theta) \rangle_{g_{M \times D^2}}^2}{\langle (X^*, \partial_\theta), (X^*, \partial_\theta) \rangle_{g_{M \times D^2}}} \\ &= \epsilon^2 |X^*|_{g_M}^2 \left( \frac{f_2(r)^2}{1 - \epsilon^2 f_2(r)^2} \right) \\ &\quad - \epsilon^4 |X^*|_{g_M}^4 \left( \frac{f_2(r)^2}{1 - \epsilon^2 f_2(r)^2} \right)^2 \left( \frac{1}{|X^*|_{g_M}^2 + \epsilon^2 |X^*|_{g_M}^2 (f_2(r)^2 / (1 - \epsilon^2 f_2(r)^2))} \right) \\ &= \epsilon^2 |X^*|_{g_M}^2 f_2(r)^2, \\ \langle W_r, W_\theta \rangle_{g_W} &= \langle (0, \partial_r), (0, \partial_\theta) \rangle_{g_{M \times D^2}} = 0. \end{aligned} \quad \square$$

We next modify  $g_W$  to have the desired product structure near  $\partial W$ . We use a technique of Wraith, which allows deformations of metrics with positive mean curvature at the boundary.

**Lemma 4.3**  $\partial W$  has positive mean curvature with respect to an inward normal vector.

**Proof** Let  $\bar{X}_i$  be local  $S^1$ -invariant vector fields extending an orthonormal frame of  $\bar{H}_p$ , and define  $X_i = \sigma_*(\bar{X}_i, 0)$ . At a point  $[p, 1]$ ,

$$\left\{ X_i, \frac{1}{\epsilon |X^*|_{g_M} f_2} W_\theta \right\}$$

is an orthonormal basis of  $T\partial W$  and

$$-\frac{1}{\epsilon |X^*|_{g_M}} W_r$$

is an inward-pointing unit normal vector. Since

$$[X_i, W_r] = [\sigma_*(\bar{X}_i, 0), \sigma_*(0, \partial_r)] = \sigma_*[(\bar{X}_i, 0), (0, \partial_r)] = 0$$

and  $|X_i| = 1$ ,

$$\frac{1}{\epsilon |X^*|_{g_M}} \langle \nabla_{X_i} X_i, -W_r \rangle = \frac{1}{\epsilon |X^*|_{g_M}} \langle X_i, \nabla_{X_i} W_r \rangle = \frac{1}{\epsilon |X^*|_{g_M}} \langle X_i, \nabla_{W_r} X_i \rangle = 0.$$

Thus

$$\frac{1}{\epsilon^3 |X^*|_{g_M}^3 f_2(1)^2} \langle \nabla_{W_\theta} W_\theta, -W_r \rangle = \frac{1}{2\epsilon^3 |X^*|_{g_M}^3 f_2(1)^2} W_r(|W_\theta|^2) = \frac{f_2'(1)}{\epsilon |X^*|_{g_M} f_2(1)}.$$

Evaluating that quantity at  $r = 1$  we see that the mean curvature is

$$\frac{\frac{1}{2} - 3\epsilon}{\epsilon |X^*|_{g_M} f_2(1)} > 0$$

for sufficiently small  $\epsilon$ . □

We see that  $g_W|_{\partial W}$  is obtained from  $g_M$  by shrinking the  $S^1$  fibers of  $\pi$ , a process which preserves positive scalar curvature.

**Lemma 4.4** *There exists a smooth path of metrics  $g_M(s)$  on  $M$ , with  $s \in [\epsilon^2 f_2(1)^2, 1]$ , such that  $g_M(\epsilon^2 f_2(1)^2) = g_W|_{\partial W}$ ,  $g_M(1) = g_M$  and  $\text{scal}(g_M(s)) > 0$  for all  $s$ .*

**Proof** We recall that  $\tau|_{M \times \{1\}}: M \rightarrow \partial W$  is a diffeomorphism. We see that

$$((\tau|_{M \times \{1\}})^* g_W)|_{\bar{H}_p} = g_W|_{H_{[p,1]}} = g_M|_{\bar{H}_p}$$

and

$$|X^*(p)|_{(\tau|_{M \times \{1\}})^* g_W}^2 = |W_\theta([p, 1])|_{g_W}^2 = \epsilon^2 f_2(1)^2 |X^*(p)|_{g_M}^2.$$

Thus, defining

$$g_M(s) = g_M|_{\bar{H}_p} + s g_M|_{\bar{V}_p},$$

we have, for  $\epsilon$  small enough, that

$$\epsilon^2 f_2(1)^2 < 1, \quad g_M(\epsilon^2 f_2(1)^2) = (\tau|_{M \times \{1\}})^* g_W \quad \text{and} \quad g_M(1) = g_M.$$

Since the metric is not changing on the horizontal space of  $\pi$ , each  $g_M(s)$  makes  $\pi$  into a Riemannian submersion with  $g_B$ . The O’Neil formula [6] then implies

$$\text{scal}(g_M(s)) = \text{scal}(g_B) - s |A_\pi|^2 - |T_\pi|^2 - |N_\pi|^2 - 2\delta N_\pi \geq \text{scal}(g_M) > 0,$$

where  $A_\pi$ ,  $T_\pi$  and  $N_\pi$  are the tensors defined for the Riemannian submersion  $\pi$  with respect to  $g_M$ . □

Use the normal exponential map from  $\partial W$  to define a collar neighborhood  $V \cong M \times [0, N]$ , where  $t \in [0, N]$  is the distance to  $\partial W$ . We choose  $N$  small such that  $V \subset U$ . Using this identification,  $g_W$  has the form

$$g_W = g(t) + dt^2,$$

where  $g(t) = g_W|_{M \times t}$  is a smooth path of metrics on  $M$ . Since  $g(0) = g_W|_{\partial W}$  has positive scalar curvature, we can choose  $N$  small such that  $\text{scal}(g(t)) > 0$  for all  $t \in [0, N]$ .

**Lemma 4.5** *We can alter  $g_W$  inside of  $V$  such that it is product-like near  $\partial W$  with  $g_W|_{\partial W} = g_M$  and  $\text{scal}(g_W|_V) > 0$ .*



**Proof** We use the paths  $g_M(s)$  and  $g(s)$  and the following lemma from [45] to replace  $g_W$  near the boundary with a product metric restricting to  $g_M$  at the boundary.

**Lemma 4.6** [45] *Let  $g(t) + dt^2$  be a metric of positive scalar curvature on  $M \times [0, N]$  such that  $\text{scal}(g(t)) > 0$  and  $M \times \{0\}$  has positive mean curvature with respect to the inward normal vector  $\partial_t$ . Let  $\bar{g}(t)$  be a smooth path of metrics on  $M$  such that  $\text{scal}(\bar{g}(t)) > 0$  for  $t \in [0, N]$  and  $\bar{g}(t) = g(t)$  for  $t$  in a neighborhood of  $N$ . Then there exists a function  $\beta: [0, N] \rightarrow \mathbb{R}_+$  such that  $\beta = 1$  for  $t$  in a neighborhood of  $N$ ,  $\beta = \beta(0)$  is constant for  $t$  in a neighborhood of  $0$ , and  $\bar{g}(t) + \beta(t) dt^2$  has positive scalar curvature.*

To define our replacement path  $\bar{g}$ , we define two smooth functions:

$$\begin{aligned} \chi_1: [0, \frac{1}{2}N] &\rightarrow [\epsilon^2 f_2(1)^2, 1] \text{ such that } \chi_1(t) = 1 \text{ for } t \text{ near } 0 \text{ and } \chi_1(t) = \epsilon^2 f_2(1)^2 \text{ for } t \text{ near } \frac{1}{2}N, \\ \chi_2: [\frac{1}{2}N, N] &\rightarrow [0, 1] \text{ such that } \chi_2(t) = 0 \text{ for } t \text{ near } \frac{1}{2}N \text{ and } \chi_2(t) = t \text{ for } t \text{ near } N. \end{aligned}$$

We then define a smooth path of metrics

$$\bar{g}(t) = \begin{cases} g_M \circ \chi_1(t) & \text{if } t \in [0, \frac{1}{2}N], \\ g \circ \chi_2(t) & \text{if } t \in [\frac{1}{2}N, N]. \end{cases}$$

By Lemma 4.4 and the definition of  $g$ ,  $\text{scal}(\bar{g}(t)) > 0$  for all  $t$ . Then Lemmas 4.3 and 4.6 imply that  $\bar{g}(t) + \beta(t) dt^2$  has positive scalar curvature for the function  $\beta(t)$  given by Lemma 4.6. For  $t$  near  $N$ ,  $\bar{g}(t) = g(t)$  and  $\beta(t) = 1$ , so  $\bar{g}(t) + \beta(t) dt^2 = g_W$ . Thus replacing  $g_W|_V$  with this metric results in a new smooth metric, for which we reuse the notation  $g_W$ . Since  $\bar{g}(t) = g$  and  $\beta(t)$  is constant for  $t$  near  $0$ ,  $\bar{g}(t) + \beta(t) dt^2$  has the desired product structure (3.3). This proves Lemma 4.5. □

## 4.2 Connection

Let  $\beta \in \Omega^2(B)$  represent the image of  $\ell d$  in  $H^2(B, \mathbb{R})$ . The Gysin sequence for an  $S^1$  bundle shows that  $\pi^* \ell d = 0$ , so we can choose  $\alpha \in \Omega^1(M)$  such that  $\pi^* \beta = d\alpha$ . Since  $\pi^* \beta$  is  $S^1$ -invariant, we can choose  $\alpha$  to be  $S^1$ -invariant by averaging.

**Lemma 4.7** 
$$\alpha(X^*) = -\frac{1}{2\pi}.$$

**Proof** Let  $\Phi \in \Omega^2(W)$  be a Thom form of the disc bundle  $\rho: W \rightarrow B$ . Since

$$[\Phi] \mapsto \rho^* \ell d$$

under the long exact sequence map  $H^2(W, \partial W) \rightarrow H^2(W)$ , we have

$$\rho^* \beta - \Phi = d\bar{\alpha}$$

for some  $\bar{\alpha} \in \Omega^1(W)$ . Since  $\Phi$  vanishes near  $\partial W$ ,

$$d\bar{\alpha}|_M = \rho^* \beta|_M = \pi^* \beta = d\alpha.$$

Since  $\pi_1(M)$  is finite,  $\bar{\alpha}|_M - \alpha$  is exact. By the defining property of the Thom form, for any point  $q \in B$ ,  $\int_{\rho^{-1}(q)} \Phi = 1$ . We use Stokes' theorem to compute

$$-1 = \int_{\rho^{-1}(q)} \rho^* \beta - \Phi = \int_{\rho^{-1}(q)} d\bar{\alpha} = \int_{\pi^{-1}(q)} \bar{\alpha} = \int_{\pi^{-1}(q)} \alpha = 2\pi\alpha(X^*). \quad \square$$

We next construct a form  $\gamma \in \Omega^1(W)$  extending  $2\pi\alpha/\ell$ . We first define a form  $\bar{\gamma} \in \Omega(M \times D^2)$ .

At  $(p, x) \in M \times D^2$ ,  $x \neq 0$ , set

$$\bar{\gamma}|_{\bar{H}_p \times \{0\}} = \frac{2\pi}{\ell} \alpha_{\bar{H}_p}, \quad \bar{\gamma}(X^*, 0) = -\frac{f_1(r)}{\ell}, \quad \bar{\gamma}(0, \partial_r) = 0, \quad \bar{\gamma}(0, \partial_\theta) = \frac{f_1(r)}{\ell},$$

where  $r$  is the radial coordinate on  $D^2$ . This form extends smoothly to the origin of  $D^2$  since  $f_1$  is zero in a neighborhood of  $r = 0$ . Since  $r, \bar{H}_p \oplus \{0\}, \alpha, \partial_r, \partial_\theta$  and  $X^*$  are all preserved by the  $S^1$  action,  $\bar{\gamma}$  is  $S^1$ -invariant. The vertical space of  $\sigma$  is generated by  $(X^*, \partial_\theta)$ , and so  $\bar{\gamma}$  vanishes on the vertical space. It follows that there is a unique form  $\gamma \in \Omega(W)$  such that  $\sigma^*\gamma = \bar{\gamma}$ .

**Lemma 4.8**

$$\tau^*\gamma = \frac{2\pi}{\ell} \text{proj}_{U,M}^* \alpha.$$

**Proof** Recall that  $f_1(r) = 1$  for  $r$  in the image of  $\tau$  and note that  $\tau^*\gamma = (\sigma^*\gamma)|_{M \times [L,1]} = \bar{\gamma}|_{M \times [L,1]}$ . Thus:

$$\begin{aligned} \tau^*\gamma|_{\bar{H}_p \oplus \{0\}} &= \bar{\gamma}|_{\bar{H}_p \oplus \{0\}} = \frac{2\pi}{\ell} \alpha_{\bar{H}_p}, \\ \tau^*\gamma(X^*, 0) &= \bar{\gamma}(X^*, 0) = -\frac{f_1(r)}{\ell} = \frac{2\pi}{\ell} \alpha(X^*), \\ \tau^*\gamma(0, \partial_t) &= \bar{\gamma}(0, \partial_r) = 0 = \frac{2\pi}{\ell} \alpha(\text{proj}_{M^*}(0, \partial_t)). \end{aligned} \quad \square$$

Let  $\lambda_B$  be the complex line bundle with  $c_1(\lambda_B) = d$ . Given a differential form in the de Rham cohomology class of  $2\pi i$  times the first Chern class of a complex line bundle, there is a unitary connection on the line bundle whose curvature is that differential form. Thus, since  $\beta$  represents  $\ell d$ , let  $\nabla_B$  be a unitary connection on  $\lambda_B$  with curvature

$$F^{\nabla_B} = \frac{2\pi i}{\ell} \beta.$$

We now define a connection on  $\lambda$ :

$$\nabla = \rho^*\nabla_B - i\gamma.$$

**Lemma 4.9**  $\nabla$  is flat on  $U$ .

**Proof** We need to show that  $F^{\tau^*\nabla} = 0$ . Using Lemma 4.8 it follows that

$$\tau^*\nabla = \tau^*\rho^*\nabla_B - i\tau^*\gamma = \text{proj}_{U,M}^* \left( \pi^*\nabla_B - \frac{2\pi i}{\ell} \alpha \right),$$

and hence the curvature of the term in the parentheses is

$$\frac{2\pi i}{\ell} \pi^* \beta - \frac{2\pi i}{\ell} d\alpha = 0. \quad \square$$

We finish the construction of  $\nabla$  by modifying it so that it is product-like near  $\partial W$  and restricts to  $\bar{\nabla}$  at  $\partial W$ . Let  $\text{proj}_{V,M}^* : V \rightarrow M$  be the projection defined by the identification  $V \cong M \times [0, N]$  from Section 4.1. Note that while  $V \subset U$ ,  $\text{proj}_{V,M}$  and  $\text{proj}_{U,M}$  will not in general agree (the latter was defined independently of  $h$ , and the former using  $h$ ). Since  $V \subset U$ ,  $\nabla$  is flat on  $V$ . Since  $\text{proj}_{V,M}$  and the inclusion of  $\partial W \cong M \times \{0\}$  are homotopy inverses,  $\text{proj}_{M,V}^*(\lambda|_M) = \lambda|_V$ . Thus  $\nabla|_V$  and  $\text{proj}_{V,M}^* \bar{\nabla}$  are both flat unitary connections on  $\lambda|_V$  and

$$\text{proj}_{V,M}^*(\bar{\nabla}) - \nabla|_V = i\delta$$

for some closed form  $\delta \in \Omega^1(V)$ . Since  $\pi_1(V) = \pi_1(M)$  is finite,  $\delta = df$  for a smooth function  $f$  on  $V$ . We modify  $f$  to a function  $\bar{f}$  which is equal to  $f$  near  $\partial W \cong M \times \{0\}$  and equal to 0 near  $M \times \{N\}$ . We then replace  $\nabla$  with  $\nabla + i d\bar{f}$  on  $V$ . We see that

$\nabla$  is still smooth, flat on  $V$ , and near  $\partial W$ ,  $\nabla = \text{proj}_{V,M}^* \bar{\nabla}$ , satisfying (3.4).

### 4.3 Curvature

We complete the proof of Theorem 3.1 by showing that (3.2) holds. On  $V$ ,  $\nabla$  is flat and by Lemma 4.5  $\text{scal}(g_W) > 0$ , so the inequality is satisfied. For the remainder of the proof we consider  $W \setminus V$ . Then  $\text{scal}(g_W)$  is given by Lemma 4.1 and the O’Neil formula for the scalar curvature of Riemannian submersion

$$\text{scal}(g_W) = \text{scal}(g_W|_{\rho^{-1}(\pi(p))}) + \text{scal}(g_B) - |A_\rho|^2 - |T_\rho|^2 - |N_\rho|^2 - 2\delta N_\rho.$$

As  $\epsilon \rightarrow 0$ ,  $|A_\rho| \rightarrow 0$ , while the final three terms remain constant. By Lemma 4.2,

$$\text{scal}(g_W|_{\rho^{-1}(\pi(p))}) = -\frac{2}{\epsilon^2 |X^*|_{g_M}^2} \left( \frac{f_2''}{f_2} \right).$$

Therefore, as  $\epsilon \rightarrow 0$ ,

$$\text{scal}(g_W) = -\frac{2}{\epsilon^2 |X^*|_{g_M}^2} \left( \frac{f_2''}{f_2} \right) + O(1).$$

Let  $\bar{X}_i$  be an orthonormal basis of  $\bar{H}_p$  with respect to  $g_M$ . Let  $X_i = \sigma_*(\bar{X}_i, 0)$ . Then  $\{X_i\}$  is an orthonormal basis of  $H_{[p,x]}$  with respect to  $g_W$  outside of  $V$ . Away from the zero section,

$$\left\{ \frac{1}{\epsilon |X^*|_{g_M}} W_r, \frac{1}{\epsilon |X^*|_{g_M} f_2} W_\theta \right\}$$

is an orthonormal basis of  $V_{[p,x]}$ . Neither the  $\bar{X}_i$  nor  $\nabla$  depend on  $\epsilon$ . Then as  $\epsilon \rightarrow 0$ , using (2.3),

$$\begin{aligned} & |F^\nabla|_{g_M} \\ & \leq \frac{1}{\epsilon^2 |X^*|_{g_M}^2 f_2} |F^\nabla(W_r, W_\theta)| + \sum_i \frac{1}{\epsilon |X^*|_{g_M}} |F^\nabla(W_r, X_i)| + \frac{1}{\epsilon |X^*|_{g_M} f_2} |F^\nabla(W_\theta, X_i)| + O(1). \end{aligned}$$

**Lemma 4.10**  $F^\nabla(W_r, W_\theta) = -if_1'(r)/\ell$  and  $F^\nabla(W_r, X_i) = F^\nabla(W_\theta, X_i) = 0$ .

**Proof** Since  $\rho_* W_r = \rho_* W_\theta = 0$ ,

$$\begin{aligned} F^\nabla(W_r, W_\theta) &= -i d\gamma(W_r, W_\theta) = -i d\gamma(\sigma_*(0, \partial_r), \sigma_*(0, \partial_\theta)) \\ &= -i \sigma^* d\gamma((0, \partial_r), (0, \partial_\theta)) = -i d\bar{\gamma}((0, \partial_r), (0, \partial_\theta)) = -i \partial_r \bar{\gamma}(0, \partial_\theta) = -i \frac{f_1'(r)}{\ell}. \end{aligned}$$

Similarly,

$$F^\nabla(W_r, X_i) = -i d\bar{\gamma}((0, \partial_r), (\bar{X}_i, 0)) = -i \left( \partial_r \left( \frac{2\pi}{\ell} \alpha(\bar{X}_i) \right) - \bar{X}_i \left( \frac{f_1(r)}{\ell} \right) \right) = 0,$$

$$F^\nabla(W_\theta, X_i) = -i d\bar{\gamma}((0, \partial_\theta), (\bar{X}_i, 0)) = -i \left( \partial_\theta \left( \frac{2\pi}{\ell} \alpha(\bar{X}_i) \right) \right) = 0. \quad \square$$

**Lemma 4.10** implies that as  $\epsilon \rightarrow 0$ ,

$$\text{scal}(g_W) - \ell |F^\nabla|_{g_M} = \frac{1}{\epsilon^2 |X^*|_{g_M}^2} \left( \frac{-2f_2'' - f_1'}{f_2} \right) + O(1) = \frac{12}{\epsilon |X^*|_{g_M}^2} \left( \frac{r}{f_2} \right) + O(1).$$

From the definition of  $f_2$  one sees that  $r/f_2 \rightarrow 1$  as  $r \rightarrow 0$ . It follows that we can choose  $\epsilon$  small enough that (3.2) holds, completing the proof of **Theorem 3.1**.  $\square$

In [30, Lemma 4.2], Kreck and Stolz constructed positive scalar curvature metrics on associated disc bundles in order to calculate their invariant for spin manifolds with free  $S^1$  actions. In their proof, they needed to assume that the  $S^1$  orbits were geodesics. The metric  $g_W$  constructed in **Theorem 3.1** generalizes their method to a free isometric  $S^1$  action without the geodesic condition.

## References

- [1] **MF Atiyah, V K Patodi, IM Singer**, *Spectral asymmetry and Riemannian geometry, I*, Math. Proc. Cambridge Philos. Soc. 77 (1975) 43–69 [MR](#) [Zbl](#)
- [2] **MF Atiyah, V K Patodi, IM Singer**, *Spectral asymmetry and Riemannian geometry, II*, Math. Proc. Cambridge Philos. Soc. 78 (1975) 405–432 [MR](#) [Zbl](#)
- [3] **A Bahri, P Gilkey**, *The eta invariant, Pin<sup>c</sup> bordism, and equivariant Spin<sup>c</sup> bordism for cyclic 2–groups*, Pacific J. Math. 128 (1987) 1–24 [MR](#) [Zbl](#)
- [4] **D Barden**, *Simply connected five-manifolds*, Ann. of Math. 82 (1965) 365–385 [MR](#) [Zbl](#)
- [5] **V N Berestovskii**, *Homogeneous Riemannian manifolds of positive Ricci curvature*, Mat. Zametki 58 (1995) 334–340 [MR](#) [Zbl](#) In Russian; translated in *Math. Notes* 58 (1995) 905–909
- [6] **A L Besse**, *Einstein manifolds*, Ergebnisse der Math. (3) 10, Springer (1987) [MR](#) [Zbl](#)
- [7] **B Botvinnik, P B Gilkey**, *The eta invariant and metrics of positive scalar curvature*, Math. Ann. 302 (1995) 507–517 [MR](#) [Zbl](#)
- [8] **B Botvinnik, M G Walsh, D J Wraith**, *Homotopy groups of the observer moduli space of Ricci positive metrics*, Geom. Topol. 23 (2019) 3003–3040 [MR](#) [Zbl](#)
- [9] **J-P Bourguignon**, *Une stratification de l'espace des structures riemanniennes*, Compositio Math. 30 (1975) 1–41 [MR](#) [Zbl](#)

- [10] **B L Burdick**, *Ricci-positive metrics on connected sums of projective spaces*, Differential Geom. Appl. 62 (2019) 212–233 [MR](#) [Zbl](#)
- [11] **D Corro, F Galaz-García**, *Positive Ricci curvature on simply-connected manifolds with cohomogeneity-two torus actions*, Proc. Amer. Math. Soc. 148 (2020) 3087–3097 [MR](#) [Zbl](#)
- [12] **D Crowley, C M Escher**, *A classification of  $S^3$ -bundles over  $S^4$* , Differential Geom. Appl. 18 (2003) 363–380 [MR](#) [Zbl](#)
- [13] **A Dessai**, *On the moduli space of nonnegatively curved metrics on Milnor spheres*, preprint (2018) [arXiv 1712.08821v2](#)
- [14] **A Dessai, D González-Álvaro**, *Moduli space of metrics of nonnegative sectional or positive Ricci curvature on homotopy real projective spaces*, Trans. Amer. Math. Soc. 374 (2021) 1–33 [MR](#) [Zbl](#)
- [15] **A Dessai, S Klaus, W Tuschmann**, *Nonconnected moduli spaces of nonnegative sectional curvature metrics on simply connected manifolds*, Bull. Lond. Math. Soc. 50 (2018) 96–107 [MR](#) [Zbl](#)
- [16] **S K Donaldson, P B Kronheimer**, *The geometry of four-manifolds*, Oxford Univ. Press (1990) [MR](#) [Zbl](#)
- [17] **D G Ebin**, *The manifold of Riemannian metrics*, from “Global analysis” (S-S Chern, S Smale, editors), Proc. Sympos. Pure Math. XIV–XVI, Amer. Math. Soc., Providence, RI (1970) 11–40 [MR](#) [Zbl](#)
- [18] **C Escher, W Ziller**, *Topology of non-negatively curved manifolds*, Ann. Global Anal. Geom. 46 (2014) 23–55 [MR](#) [Zbl](#)
- [19] **H Geiges, C B Thomas**, *Contact topology and the structure of 5-manifolds with  $\pi_1 = \mathbb{Z}_2$* , Ann. Inst. Fourier (Grenoble) 48 (1998) 1167–1188 [MR](#) [Zbl](#)
- [20] **P B Gilkey, J Park, W Tuschmann**, *Invariant metrics of positive Ricci curvature on principal bundles*, Math. Z. 227 (1998) 455–463 [MR](#) [Zbl](#)
- [21] **M J Goodman**, *On the moduli spaces of metrics with nonnegative sectional curvature*, Ann. Global Anal. Geom. 57 (2020) 305–320 [MR](#) [Zbl](#)
- [22] **K Grove, W Ziller**, *Curvature and symmetry of Milnor spheres*, Ann. of Math. 152 (2000) 331–367 [MR](#) [Zbl](#)
- [23] **K Grove, W Ziller**, *Cohomogeneity one manifolds with positive Ricci curvature*, Invent. Math. 149 (2002) 619–646 [MR](#) [Zbl](#)
- [24] **K Grove, W Ziller**, *Lifting group actions and nonnegative curvature*, Trans. Amer. Math. Soc. 363 (2011) 2865–2890 [MR](#) [Zbl](#)
- [25] **I Hambleton, M Kreck, P Teichner**, *Nonorientable 4-manifolds with fundamental group of order 2*, Trans. Amer. Math. Soc. 344 (1994) 649–665 [MR](#) [Zbl](#)
- [26] **I Hambleton, Y Su**, *On certain 5-manifolds with fundamental group of order 2*, Q. J. Math. 64 (2013) 149–175 [MR](#) [Zbl](#)
- [27] **R C Kirby, L R Taylor**, *Pin structures on low-dimensional manifolds*, from “Geometry of low-dimensional manifolds, II” (S K Donaldson, C B Thomas, editors), London Math. Soc. Lecture Note Ser. 151, Cambridge Univ. Press (1990) 177–242 [MR](#) [Zbl](#)
- [28] **M Kreck, S Stolz**, *A diffeomorphism classification of 7-dimensional homogeneous Einstein manifolds with  $SU(3) \times SU(2) \times U(1)$ -symmetry*, Ann. of Math. 127 (1988) 373–388 [MR](#) [Zbl](#)
- [29] **M Kreck, S Stolz**, *Some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature*, J. Differential Geom. 33 (1991) 465–486 [MR](#) [Zbl](#)

- [30] **M Kreck, S Stolz**, *Nonconnected moduli spaces of positive sectional curvature metrics*, J. Amer. Math. Soc. 6 (1993) 825–850 [MR](#) [Zbl](#)
- [31] **H B Lawson, Jr, M-L Michelsohn**, *Spin geometry*, Princeton Mathematical Series 38, Princeton Univ. Press (1989) [MR](#) [Zbl](#)
- [32] **S López de Medrano**, *Involutions on manifolds*, Ergebnisse der Math. (3) 59, Springer (1971) [MR](#) [Zbl](#)
- [33] **J C Nash**, *Positive Ricci curvature on fibre bundles*, J. Differential Geometry 14 (1979) 241–254 [MR](#) [Zbl](#)
- [34] **G Perelman**, *Construction of manifolds of positive Ricci curvature with big volume and large Betti numbers*, from “Comparison geometry” (K Grove, P Petersen, editors), Math. Sci. Res. Inst. Publ. 30, Cambridge Univ. Press (1997) 157–163 [MR](#) [Zbl](#)
- [35] **L J Schwachhöfer, W Tuschmann**, *Metrics of positive Ricci curvature on quotient spaces*, Math. Ann. 330 (2004) 59–91 [MR](#) [Zbl](#)
- [36] **C Searle, F Wilhelm**, *How to lift positive Ricci curvature*, Geom. Topol. 19 (2015) 1409–1475 [MR](#) [Zbl](#)
- [37] **J-P Sha, D Yang**, *Positive Ricci curvature on the connected sums of  $S^n \times S^m$* , J. Differential Geom. 33 (1991) 127–137 [MR](#) [Zbl](#)
- [38] **J-P Sha, D Yang**, *Positive Ricci curvature on compact simply connected 4-manifolds*, from “Differential geometry: Riemannian geometry, III” (R Greene, S T Yau, editors), Proc. Sympos. Pure Math. 54, Amer. Math. Soc., Providence, RI (1993) 529–538 [MR](#) [Zbl](#)
- [39] **S Smale**, *On the structure of 5-manifolds*, Ann. of Math. 75 (1962) 38–46 [MR](#) [Zbl](#)
- [40] **R E Stong**, *Relations among characteristic numbers, I*, Topology 4 (1965) 267–281 [MR](#) [Zbl](#)
- [41] **M Y Wang, W Ziller**, *Einstein metrics on principal torus bundles*, J. Differential Geom. 31 (1990) 215–248 [MR](#) [Zbl](#)
- [42] **J Wermelinger**, *Moduli space of metrics of nonnegative sectional or positive Ricci curvature on Brieskorn quotients in dimension 5*, preprint (2020) [arXiv 2006.13697](#)
- [43] **D Wraith**, *Exotic spheres with positive Ricci curvature*, J. Differential Geom. 45 (1997) 638–649 [MR](#) [Zbl](#)
- [44] **D J Wraith**, *New connected sums with positive Ricci curvature*, Ann. Global Anal. Geom. 32 (2007) 343–360 [MR](#) [Zbl](#)
- [45] **D J Wraith**, *On the moduli space of positive Ricci curvature metrics on homotopy spheres*, Geom. Topol. 15 (2011) 1983–2015 [MR](#) [Zbl](#)

Department of Mathematics, Colby College  
Waterville, ME, United States

[mjgoodma@colby.edu](mailto:mjgoodma@colby.edu)

Proposed: Tomasz S Mrowka

Seconded: Tobias H Colding, David Fisher

Received: 4 December 2020

Revised: 1 April 2022

# GEOMETRY & TOPOLOGY

[msp.org/gt](http://msp.org/gt)

## MANAGING EDITOR

András I Stipsicz    Alfréd Rényi Institute of Mathematics  
[stipsicz@renyi.hu](mailto:stipsicz@renyi.hu)

## BOARD OF EDITORS

Mohammed Abouzaid	Stanford University <a href="mailto:abouzaid@stanford.edu">abouzaid@stanford.edu</a>	Mark Gross	University of Cambridge <a href="mailto:mgross@dpms.cam.ac.uk">mgross@dpms.cam.ac.uk</a>
Dan Abramovich	Brown University <a href="mailto:dan_abramovich@brown.edu">dan_abramovich@brown.edu</a>	Rob Kirby	University of California, Berkeley <a href="mailto:kirby@math.berkeley.edu">kirby@math.berkeley.edu</a>
Ian Agol	University of California, Berkeley <a href="mailto:ianagol@math.berkeley.edu">ianagol@math.berkeley.edu</a>	Bruce Kleiner	NYU, Courant Institute <a href="mailto:bkleiner@cims.nyu.edu">bkleiner@cims.nyu.edu</a>
Arend Bayer	University of Edinburgh <a href="mailto:arend.bayer@ed.ac.uk">arend.bayer@ed.ac.uk</a>	Sándor Kovács	University of Washington <a href="mailto:skovacs@uw.edu">skovacs@uw.edu</a>
Mark Behrens	University of Notre Dame <a href="mailto:mbehrens1@nd.edu">mbehrens1@nd.edu</a>	Urs Lang	ETH Zürich <a href="mailto:urs.lang@math.ethz.ch">urs.lang@math.ethz.ch</a>
Mladen Bestvina	University of Utah <a href="mailto:bestvina@math.utah.edu">bestvina@math.utah.edu</a>	Marc Levine	Universität Duisburg-Essen <a href="mailto:marc.levine@uni-due.de">marc.levine@uni-due.de</a>
Martin R Bridson	University of Oxford <a href="mailto:bridson@maths.ox.ac.uk">bridson@maths.ox.ac.uk</a>	Ciprian Manolescu	University of California, Los Angeles <a href="mailto:cm@math.ucla.edu">cm@math.ucla.edu</a>
Jim Bryan	University of British Columbia <a href="mailto:jbryan@math.ubc.ca">jbryan@math.ubc.ca</a>	Haynes Miller	Massachusetts Institute of Technology <a href="mailto:hmr@math.mit.edu">hmr@math.mit.edu</a>
Dmitri Burago	Pennsylvania State University <a href="mailto:burago@math.psu.edu">burago@math.psu.edu</a>	Tomasz Mrowka	Massachusetts Institute of Technology <a href="mailto:mrowka@math.mit.edu">mrowka@math.mit.edu</a>
Tobias H Colding	Massachusetts Institute of Technology <a href="mailto:colding@math.mit.edu">colding@math.mit.edu</a>	Aaron Naber	Northwestern University <a href="mailto:anaber@math.northwestern.edu">anaber@math.northwestern.edu</a>
Simon Donaldson	Imperial College, London <a href="mailto:s.donaldson@ic.ac.uk">s.donaldson@ic.ac.uk</a>	Peter Ozsváth	Princeton University <a href="mailto:petero@math.princeton.edu">petero@math.princeton.edu</a>
Yasha Eliashberg	Stanford University <a href="mailto:eliash-gt@math.stanford.edu">eliash-gt@math.stanford.edu</a>	Leonid Polterovich	Tel Aviv University <a href="mailto:polterov@post.tau.ac.il">polterov@post.tau.ac.il</a>
Benson Farb	University of Chicago <a href="mailto:farb@math.uchicago.edu">farb@math.uchicago.edu</a>	Colin Rourke	University of Warwick <a href="mailto:gt@maths.warwick.ac.uk">gt@maths.warwick.ac.uk</a>
David M Fisher	Rice University <a href="mailto:davidfisher@rice.edu">davidfisher@rice.edu</a>	Roman Sauer	Karlsruhe Institute of Technology <a href="mailto:roman.sauer@kit.edu">roman.sauer@kit.edu</a>
Mike Freedman	Microsoft Research <a href="mailto:michaelf@microsoft.com">michaelf@microsoft.com</a>	Stefan Schwede	Universität Bonn <a href="mailto:schwede@math.uni-bonn.de">schwede@math.uni-bonn.de</a>
David Gabai	Princeton University <a href="mailto:gabai@princeton.edu">gabai@princeton.edu</a>	Natasa Sesum	Rutgers University <a href="mailto:natasas@math.rutgers.edu">natasas@math.rutgers.edu</a>
Stavros Garoufalidis	Southern U. of Sci. and Tech., China <a href="mailto:stavros@mpim-bonn.mpg.de">stavros@mpim-bonn.mpg.de</a>	Gang Tian	Massachusetts Institute of Technology <a href="mailto:tian@math.mit.edu">tian@math.mit.edu</a>
Cameron Gordon	University of Texas <a href="mailto:gordon@math.utexas.edu">gordon@math.utexas.edu</a>	Ulrike Tillmann	Oxford University <a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>
Jesper Grodal	University of Copenhagen <a href="mailto:jg@math.ku.dk">jg@math.ku.dk</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Misha Gromov	IHÉS and NYU, Courant Institute <a href="mailto:gromov@ihes.fr">gromov@ihes.fr</a>	Anna Wienhard	Universität Heidelberg <a href="mailto:wienhard@mathi.uni-heidelberg.de">wienhard@mathi.uni-heidelberg.de</a>

---

See inside back cover or [msp.org/gt](http://msp.org/gt) for submission instructions.

---

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

---

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2024 Mathematical Sciences Publishers



# GEOMETRY & TOPOLOGY

Volume 28 Issue 3 (pages 1005–1499) 2024

---

Homological mirror symmetry for hypertoric varieties, I: Conic equivariant sheaves	1005
MICHAEL MCBREEN and BEN WEBSTER	
Moduli spaces of Ricci positive metrics in dimension five	1065
MCFEELY JACKSON GOODMAN	
Riemannian manifolds with entire Grauert tube are rationally elliptic	1099
XIAOYANG CHEN	
On certain quantifications of Gromov’s nonsqueezing theorem	1113
KEVIN SACKEL, ANTOINE SONG, U MUT VAROLGUNES and JONATHAN J ZHU	
Zariski dense surface groups in $SL(2k + 1, \mathbb{Z})$	1153
D DARREN LONG and MORWEN B THISTLETHWAITE	
Scalar and mean curvature comparison via the Dirac operator	1167
SIMONE CECCHINI and RUDOLF ZEIDLER	
Symplectic capacities, unperturbed curves and convex toric domains	1213
DUSA MCDUFF and KYLER SIEGEL	
Quadric bundles and hyperbolic equivalence	1287
ALEXANDER KUZNETSOV	
Categorical wall-crossing formula for Donaldson–Thomas theory on the resolved conifold	1341
YUKINOBU TODA	
Nonnegative Ricci curvature, metric cones and virtual abelianness	1409
JIAYIN PAN	
The homology of the Temperley–Lieb algebras	1437
RACHAEL BOYD and RICHARD HEPWORTH	