



# *Geometry & Topology*

Volume 28 (2024)

**A nonexistence result for wing-like mean curvature flows in  $\mathbb{R}^4$**

KYEONGSU CHOI  
ROBERT HASLHOFER  
OR HERSHKOVITS

# A nonexistence result for wing-like mean curvature flows in $\mathbb{R}^4$

KYEONGSU CHOI  
ROBERT HASLHOFFER  
OR HERSHKOVITS

Some of the most worrisome potential singularity models for the mean curvature flow of three-dimensional hypersurfaces in  $\mathbb{R}^4$  are noncollapsed wing-like flows, ie noncollapsed flows that are asymptotic to a wedge. We rule out this potential scenario, not just among self-similarly translating singularity models, but in fact among all ancient noncollapsed flows in  $\mathbb{R}^4$ . Specifically, we prove that for any ancient noncollapsed mean curvature flow  $M_t = \partial K_t$  in  $\mathbb{R}^4$  the blowdown  $\lim_{\lambda \rightarrow 0} \lambda \cdot K_{t_0}$  is always a point, halfline, line, halfplane, plane or hyperplane, but never a wedge. In our proof we introduce a fine bubble-sheet analysis, which generalizes the fine neck analysis that has played a major role in many recent papers. Our result is also a key first step towards the classification of ancient noncollapsed flows in  $\mathbb{R}^4$ , which we will address in a series of subsequent papers.

53E10

1. Introduction	3096
2. Coarse properties of ancient noncollapsed flows	3102
3. Foliations and barriers	3105
4. Setting up the fine bubble-sheet analysis	3107
5. Bubble-sheet analysis in the neutral mode	3115
6. Bubble-sheet analysis in the unstable mode	3120
7. Conclusion in the unstable mode case	3128
Appendix A. Local $L^\infty$ -estimate	3130
Appendix B. Merle–Zaag ODE lemma	3132
References	3132

## 1 Introduction

The mean curvature flow of mean-convex surfaces in  $\mathbb{R}^3$ , and also of 2-convex hypersurfaces in  $\mathbb{R}^{n+1}$ , is by now well understood. In particular, White [30; 31] proved that the evolving surfaces are smooth away from a small set of times, and Huisken and Sinestrari [22], Brendle and Huisken [8] and Haslhofer and Kleiner [18] constructed a flow with surgery. More recently, in significant work by Angenent, Daskalopoulos and Sesum [3; 4] and Brendle and Choi [6; 7] a complete classification of all possible singularity models for such flows has been obtained (these classification results have in turn be generalized in our recent proof of the mean-convex neighborhood conjecture [10; 11]).

Some important results are of course valid in arbitrary dimensions. In particular, tangent flows, ie blowup limits centered at some fixed space-time point, are always self-similarly shrinking thanks to Huisken's monotonicity formula [20]. By fundamental results of Huisken [21] and Colding and Minicozzi [13] the only mean-convex shrinkers (and also the only stable shrinkers) are round spheres and cylinders. This of course is an important ingredient in the above quoted regularity theory for mean-convex flows due to White. More recently, Colding and Minicozzi [14] proved uniqueness of cylindrical tangent flows, and applied this in [15] to obtain refined information about the structure of the singular set in flows that only encounter stable singularities. On the other hand, general blowup limits, ie blowup limits along arbitrary sequences of space-time points, are much less understood already for the flow of three-dimensional hypersurfaces in  $\mathbb{R}^4$ . In particular, one of the most worrisome potential scenarios is that one encounters so-called wing-like flows as blowup limits, ie flows that are asymptotic to a wedge.

To explain the background, let us first discuss the situation of two-dimensional flows in  $\mathbb{R}^3$ , and in fact let us restrict the discussion even further to the case of strictly convex, graphical, self-similarly translating solutions. The most well known such solution is the translating bowl soliton of Altschuler and Wu [1], which is given by a rotationally symmetric entire graph. However, in addition to the bowl there is also a one-parameter family of solutions defined over strip regions  $(-b, b) \times \mathbb{R}$  for every  $b > \pi/2$ . These solutions have been constructed by Ilmanen (unpublished) and Wang [29], and have been classified in recent work by Hoffman, Ilmanen, Martin and White [19], building also on important prior work by Spruck and Xiao [28]. Ilmanen called them  $\Delta$ -wings, capturing their shape. They are asymptotic to a wedge with its sides modeled on the grim-reaper times  $\mathbb{R}$ .

The two-dimensional  $\Delta$ -wings do not cause any deep concerns for the singularity analysis of mean curvature flow. This is because they are collapsed. In fact, it is known that collapsed solutions cannot arise as blowup limit of any mean-convex flow [30; 31] (see also Sheng and Wang [27], Andrews [2] and Haslhofer and Kleiner [17]), and conjectured that they cannot arise as blowup limit of any embedded flow; see Ilmanen [23]. We recall that a mean-convex flow is called noncollapsed, if there is some  $\alpha > 0$  such that every point admits interior and exterior balls of radius at least  $\alpha/H(p)$ .

However, the situation changes dramatically when one moves one dimension higher. This is because the grim-reaper is collapsed, but the bowl is noncollapsed. Hence, one has to worry about the potential




Figure 1: One of the most worrisome potential singularity models would be a three-dimensional noncollapsed wing-like flow in  $\mathbb{R}^4$  with its sides modeled on the two-dimensional bowl times  $\mathbb{R}$ .

scenario that a blowup limit could be a wing-like translator with its sides modeled on the two-dimensional bowl times  $\mathbb{R}$ . This is illustrated in Figure 1. More generally, this leads to the question:

**Question 1.1** (noncollapsed wing-like translators) *Do there exist any noncollapsed wing-like translators in  $\mathbb{R}^4$ ?*

Here, by “wing-like” we mean any solution that is asymptotic to a wedge, ie a two-dimensional convex cone with angle strictly less than  $\pi$ . Such solutions would cause a major complication for understanding the flow through singularities in  $\mathbb{R}^4$ , and would also be a major obstacle for the construction of a flow with surgery. More generally, one also has to worry about blowup limits that are not necessarily self-similar:

**Question 1.2** (noncollapsed wing-like ancient flows) *Do there exist any noncollapsed wing-like ancient flows in  $\mathbb{R}^4$ ?*

### 1.1 Main results

To address Questions 1.1 and 1.2 (and even to properly define the notion of wing-like), we study the blowdown of the time slices. To this end, let  $M_t = \partial K_t \subset \mathbb{R}^4$  be a noncollapsed ancient solution of the mean curvature flow. We recall that such flows are always convex thanks to [17, Theorem 1.10].

**Definition 1.3** (blowdown) Given any time  $t_0$ , the *blowdown* of  $K_{t_0}$  is defined by

$$(1) \quad \check{K}_{t_0} := \lim_{\lambda \rightarrow 0} \lambda \cdot K_{t_0}.$$

For example, if the flow is a three-dimensional bowl soliton, then its blowdown is a halfline. On the other hand, if the flow is a two-dimensional bowl soliton times  $\mathbb{R}$ , then its blowdown is a two-dimensional halfplane. The scenario of a wing-like flow would correspond to the case where the blowdown is a wedge, ie a two-dimensional convex cone with angle strictly less than  $\pi$ .

Our main theorem completely classifies all possible blowdowns of noncollapsed flows in  $\mathbb{R}^4$ :

**Theorem 1.4** (blowdown of noncollapsed flows in  $\mathbb{R}^4$ ) *Let  $M_t = \partial K_t$  be an ancient noncollapsed mean curvature flow in  $\mathbb{R}^4$ . Then its blowdown is time-independent and is*

- *either a point (which only happens if the solution is compact),*
- *or a halfline,*
- *or a line (which only happens for  $S^2 \times \mathbb{R}$  or oval times  $\mathbb{R}$ ),*
- *or a two-dimensional halfplane (which only happens if the solution is a two-dimensional bowl soliton times  $\mathbb{R}$ ),*
- *or a two-dimensional plane (which only happens for  $S^1 \times \mathbb{R}^2$ ),*
- *or a three-dimensional hyperplane (which only happens for flat  $\mathbb{R}^3$ ).*

*In particular, the blowdown can never be a wedge.*

**Theorem 1.4** shows that the asymptotic structure of noncollapsed flows, ie the flows that are actually relevant for singularity analysis, is much more rigid than the asymptotic structure of arbitrary convex flows. In fact, the example of the  $\Delta$ -wings illustrates that even for two-dimensional translators in  $\mathbb{R}^3$  one can get the wedge of any angle less than  $\pi$  as blowdown. In  $\mathbb{R}^4$  there is a zoo of collapsed ancient convex solutions, see eg Wang [29] and Hoffman, Ilmanen, Martín and White [19], whose blowdown can be much more arbitrary than the ones in **Theorem 1.4**.

As an immediate corollary, we can answer **Question 1.2**. Namely, since the blowdown can never be a wedge, we can rule out the potential scenario of noncollapsed wing-like ancient flows in  $\mathbb{R}^4$ :

**Corollary 1.5** (nonexistence of wing-like noncollapsed flows) *Wing-like noncollapsed ancient flows in  $\mathbb{R}^4$  do not exist.*

In particular, this also answers **Question 1.1**:

**Corollary 1.6** (nonexistence of wing-like noncollapsed translators) *Wing-like noncollapsed translators in  $\mathbb{R}^4$  do not exist.*

Corollaries **1.5** and **1.6** rule out some of the most worrisome potential singularity models for the mean curvature flow of three-dimensional mean-convex hypersurfaces in  $\mathbb{R}^4$  (and more generally also for mean-convex flows in 4-manifolds, since after blowup the ambient manifold always becomes Euclidean).

## 1.2 Outline and further results

Let us now outline our approach. Let  $M_t = \partial K_t$  be an ancient noncollapsed mean curvature flow in  $\mathbb{R}^4$ . We can assume that the solution is noncompact and nonflat, as otherwise the blowdown would simply be a point or a three-dimensional hyperplane, respectively.

As we explain in Section 2, it follows from the general theory of noncollapsed flows (see Haslhofer and Kleiner [17]) that the blowdown

$$(2) \quad \check{K} := \lim_{\lambda \rightarrow 0} \lambda \cdot K_{t_0}$$

always is a convex cone of dimension at most 2 (also, this limit is in fact independent of  $t_0$ ). If  $\check{K}$  splits off a line, then so does the flow  $M_t = \partial K_t$ , and we are done by the classification of two-dimensional noncollapsed flows in  $\mathbb{R}^3$  by Brendle and Choi [6] and Angenent, Daskalopoulos and Sesum [4]. After these reductions, the task is thus to rule out the scenario where  $\check{K}$  is wedge.

We next consider the tangent flow at  $-\infty$ . Flows with a neck-tangent flow at  $-\infty$ , ie with

$$(3) \quad \lim_{\lambda \rightarrow 0} \lambda M_{\lambda^{-2}t} = \mathbb{R} \times S^2(\sqrt{-4t}),$$

have already been classified by Brendle and Choi [7]. We can thus assume that we have a bubble-sheet tangent flow at  $-\infty$ , ie that

$$(4) \quad \lim_{\lambda \rightarrow 0} \lambda M_{\lambda^{-2}t} = \mathbb{R}^2 \times S^1(\sqrt{-2t}).$$

In Section 3, to facilitate various calibration and barrier arguments, we build a foliation for flows close to a bubble sheet in  $\mathbb{R}^4$ . We do this by shifting and rotating the two-dimensional shrinker foliation in  $\mathbb{R}^3$  from Angenent, Daskalopoulos and Sesum [3].

In Section 4, we set up the fine bubble-sheet analysis, which generalizes the fine neck-analysis that played a major role in Angenent, Daskalopoulos and Sesum [3; 4], Brendle and Choi [6; 7], Choi, Haslhofer and Hershkovits [10] and Choi, Haslhofer, Hershkovits and White [11]. Given any space-time point  $X_0$ , we consider the renormalized flow

$$(5) \quad \bar{M}_\tau = e^{\tau/2}(M_{t_0-e^{-\tau}} - x_0).$$

Then, the hypersurfaces  $\bar{M}_\tau$  converge for  $\tau \rightarrow -\infty$  to the cylinder

$$(6) \quad \Gamma = \mathbb{R}^2 \times S^1(\sqrt{2}).$$

In particular,  $\bar{M}_\tau$  can be written as the graph of a function  $u = u_{X_0}(\cdot, \tau)$  with small norm over  $\Gamma \cap B_{2\rho(\tau)}$ , where  $\rho(\tau) \rightarrow \infty$  as  $\tau \rightarrow -\infty$ . The goal is then to derive a sharp asymptotic expansion for the graph function  $u$ .

The analysis over  $\Gamma$  is governed by the Ornstein–Uhlenbeck operator

$$(7) \quad \mathcal{L} = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{1}{2} \frac{\partial^2}{\partial \theta^2} - \frac{1}{2} x_1 \frac{\partial}{\partial x_1} - \frac{1}{2} x_2 \frac{\partial}{\partial x_2} + 1.$$

The operator  $\mathcal{L}$  has 5 unstable eigenfunctions, namely

$$(8) \quad 1, \cos \theta, \sin \theta, x_1, x_2,$$

and 7 neutral eigenfunctions, namely

$$(9) \quad x_1 \cos \theta, x_1 \sin \theta, x_2 \cos \theta, x_2 \sin \theta, x_1^2 - 2, x_2^2 - 2, x_1 x_2.$$

Using the ODE lemma from Merle and Zaag [26], we show that for  $\tau \rightarrow -\infty$  either the unstable mode is dominant or the neutral mode is dominant. Furthermore, considering instead an expansion for

$$(10) \quad \tilde{M}_\tau = S(\tau) \bar{M}_\tau,$$

where the fine-tuning rotation  $S(\tau) \in \text{SO}(4)$  is obtained via the implicit function theorem, we can assume that in the neutral mode case the truncated function  $\hat{u}$  is orthogonal to the  $\theta$ -dependent functions from (9).

In Section 5, we consider the case where the neutral mode is dominant. By the reductions from above we can assume that the blowdown  $\check{K}$  is a convex cone that does not contain any line. Furthermore, rotating coordinates, we can arrange that  $\check{K}$  contains the positive  $x_1$ -axis, namely

$$(11) \quad \{\lambda e_1 \mid \lambda \geq 0\} \subseteq \check{K}.$$

In this setting, we have:

**Theorem 1.7** (blowdown in neutral mode) *If the neutral mode is dominant, then its blowdown is a halfline, namely*

$$(12) \quad \check{K} = \{\lambda e_1 \mid \lambda \geq 0\}.$$

Moreover,

$$(13) \quad \lim_{\tau \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|} = -c(x_2^2 - 2),$$

where  $c > 0$ .

To prove this, remembering (9) and the fine-tuning, we first show that along a subsequence we have

$$(14) \quad \lim_{\tau_{i_m} \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau_{i_m})}{\|\hat{u}(\cdot, \tau_{i_m})\|} = q_{11}(x_1^2 - 2) + q_{22}(x_2^2 - 2) + 2q_{12}x_1x_2.$$

Using Brunn's concavity principle, we show that  $Q = \{q_{ij}\}$  is a nontrivial seminegative definite  $2 \times 2$ -matrix.

The crucial step is then to relate the algebra of the quadratic form  $Q$  with the geometry of the blowdown  $\check{K}$ . Specifically, we show that

$$(15) \quad \check{K} \subseteq \ker Q.$$

The basic idea for this is that in directions  $v \notin \ker Q$  we have an inwards quadratic bending, which implies that  $v$  is a "short" direction, ie  $v \notin \check{K}$ . This can be made precise using again Brunn's concavity principle. Once (15) is established, Theorem 1.7 follows easily. In fact, our argument also gives a quantitative estimate for the diameter of the short directions:

**Corollary 1.8** (diameter of level sets) *If  $M_t = \partial K_t$  is as above, then for any  $\delta > 0$  we have*

$$\bar{M}_\tau \cap \{x_1 = 0\} \subseteq B_{e^{\delta|\tau|}}(0) \quad \text{for } \tau \ll 0.$$

In [Section 6](#), we consider the case where the unstable mode is dominant. Our main technical result for such flows is the following:

**Theorem 1.9** (fine bubble-sheet theorem) *Let  $\{M_t\}$  be an ancient noncollapsed flow in  $\mathbb{R}^4$ , with a bubble sheet tangent flow at  $-\infty$ , whose unstable mode is dominant. Then there exist universal constants  $a_1$  and  $a_2$  such that, for every  $X$ , after suitable recentering in the  $x_3x_4$ -plane, the truncated graph function  $\check{u}^X(\cdot, \tau)$  of the renormalized flow  $\bar{M}_\tau^X$  satisfies*

$$(16) \quad \check{u}^X = e^{\tau/2} (a_1x_1 + a_2x_2) + o(e^{\tau/2})$$

for  $\tau \ll 0$  depending only on an upper bound on the bubble-sheet scale  $Z(X)$ . Moreover, the expansion parameters satisfy

$$(17) \quad |a_1| + |a_2| > 0.$$

[Theorem 1.9](#) shows that the bubble-sheet increases its bubble-size slightly, in a precisely described way, as one moves in direction of the vector  $(a_1, a_2)$ . The constants  $a_1$  and  $a_2$  are genuine constants of the flow. For example, if  $M_t$  is  $\mathbb{R}$  times a two-dimensional bowl translating in  $x_2$ -direction, then  $a_1 = 0$  and  $a_2$  is proportional to the reciprocal of the speed of translation.

The fine-bubble sheet theorem ([Theorem 1.9](#)) generalizes the fine-neck theorem from our prior work [[10](#); [11](#)]. To prove it, we follow the scheme from our prior work with the necessary modifications. In particular, we now expand in terms of the unstable eigenfunctions from ([8](#)), and we use the new foliation from [Section 3](#). Another new step is to show that the unstable mode is dominant even after removing the fine-tuning rotation.

In [Section 7](#), we play the following end game. Let  $M_t = \partial K_t$  be a noncompact ancient noncollapsed flow in  $\mathbb{R}^4$ , with bubble-sheet tangent flow at  $-\infty$ , whose unstable mode is dominant, and suppose towards a contradiction that its blowdown is a wedge. Then taking suitable limits along the sides, we see  $\mathbb{R}$  times a two-dimensional bowl translating soliton. As observed above, the vector  $(a_1, a_2)$  points in the translation direction. However, since the limits obtained along the two different sides of the wedge translate in different directions this gives the desired contradiction. In fact, the argument also works for a degenerate wedge (ie a halfline) and thus proves:

**Theorem 1.10** (classification in unstable mode) *The only noncompact ancient noncollapsed flow in  $\mathbb{R}^4$ , with bubble-sheet tangent flow at  $-\infty$ , whose unstable mode is dominant, is  $\mathbb{R} \times 2d$ -bowl.*

[Theorem 1.10](#), in addition to ruling out the wedge blowdown, in fact completes the classification of noncompact ancient noncollapsed flows in  $\mathbb{R}^4$  in case the unstable mode is dominant. We will address the neutral mode case in subsequent work, based on [Theorem 1.7](#) and [Corollary 1.8](#).



**Acknowledgements** Choi has been supported by the KIAS Individual grant MG078902, an Asian Young Scientist Fellowship, and the National Research Foundation of Korea (NRF) grants RS-2023-00219980 and RS-2024-00345403 funded by the Korean government (MSIT). Haslhofer has been supported by the NSERC Discovery grants RGPIN-2016-04331 and RGPIN-2023-04419, and a Sloan Research Fellowship. Hershkovits has been supported by the ISF grant 437/20. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program, grant agreement 101116390. We are very grateful to Wenkui Du and the referees for very detailed and helpful comments.

## 2 Coarse properties of ancient noncollapsed flows

Recall that by [17, Theorem 1.14] and [14], if  $M_t = \partial K_t$  is an ancient noncollapsed<sup>1</sup> mean curvature flow in  $\mathbb{R}^4$ , that is noncompact and nonflat, then, up to rotation, either

$$(18) \quad \lim_{\lambda \rightarrow 0} \lambda K_{\lambda^{-2}t} = \mathbb{R} \times D^3(\sqrt{-4t})$$

or

$$(19) \quad \lim_{\lambda \rightarrow 0} \lambda K_{\lambda^{-2}t} = \mathbb{R}^2 \times D^2(\sqrt{-2t}).$$

In the case (18) we say that  $M_t$  has a *neck tangent flow at  $-\infty$* , whereas in case (19) we say that  $M_t$  has a *bubble-sheet tangent flow at  $-\infty$* .

In the neck tangent case, it follows from Brendle and Choi [7] (or alternatively from Choi, Haslhofer, Hershkovits and White [11]) that  $M_t$  is either a round shrinking  $S^2 \times \mathbb{R}$  or a three-dimensional bowl. Hence, we can focus on the bubble-sheet case.

### 2.1 The blowdown of time slices

In this section, we establish several elementary properties of the blowdown of time slices (Definition 1.3).

**Proposition 2.1** (basic properties of blowdown) *For any ancient noncollapsed mean curvature flow  $M_t = \partial K_t$  in  $\mathbb{R}^4$ , that is noncompact and nonflat, the blowdown  $\check{K}_{t_0}$  is a convex cone of dimension at most 2.*

**Proof** Assume  $t_0 = 0$  and  $0 \in M_0$  for ease of notation. By convexity (see [17, Theorem 1.10]), we have  $\lambda \cdot K_0 \subseteq \mu \cdot K_0$  whenever  $\lambda \leq \mu$ . Hence,

$$(20) \quad \check{K}_0 := \lim_{\lambda \rightarrow 0} \lambda \cdot K_0 = \bigcap_{\lambda > 0} \lambda \cdot K_0$$

exists and is convex. Since  $K_0$  is noncompact,  $\check{K}_0$  contains a halfline. Clearly, we have  $\lambda \cdot \check{K}_0 = \check{K}_0$  for any  $\lambda > 0$ , ie  $\check{K}_0$  is a cone.

<sup>1</sup>By [5] and [16] we can always take  $\alpha = 1$  as noncollapsing constant.

As the flow is moving inwards, for every  $t < 0$  we have

$$(21) \quad \lambda K_0 \subseteq \lambda K_{\lambda^{-2}t}.$$

Thus,

$$(22) \quad \check{K}_0 = \lim_{\lambda \rightarrow 0} \lambda K_0 \subseteq \lim_{\lambda \rightarrow 0} \lambda K_{\lambda^{-2}t},$$

where the limit on the right-hand side is described (after suitable rotation) either by (18) or by (19). In either case, we infer that

$$(23) \quad \check{K}_0 \subseteq \mathbb{R}^2 \times \{0\},$$

so  $\check{K}_0$  is at most two-dimensional. This proves the proposition. □

**Proposition 2.2** (dimension reduction) *Let  $M_t = \partial K_t$  be an ancient noncollapsed mean curvature flow in  $\mathbb{R}^4$ . If  $\check{K}_{t_0}$  contains a line, then  $M_t$  is either a static hyperplane, a round shrinking  $\mathbb{R} \times S^2$ , a round shrinking  $\mathbb{R}^2 \times S^1$ , a two-dimensional bowl times  $\mathbb{R}$ , or a two-dimensional ancient oval times  $\mathbb{R}$ .*

**Proof** If  $\check{K}_{t_0}$  contains a line, then so does  $K_{t_0}$ . Hence, the flow  $M_t = \partial K_t$  splits off an  $\mathbb{R}$ -factor. By the classification of two-dimensional noncollapsed flows in  $\mathbb{R}^3$  by Angenent, Daskalopoulos and Sesum [3] and Brendle and Choi [6], this implies the assertion. □

We will now show that the blowdown  $\check{K}_{t_0}$  is independent of  $t_0 < T_{\text{ext}}(\mathcal{M})$ , where  $T_{\text{ext}}(\mathcal{M})$  denotes the extinction time of our flow. To show this, we need the following lemma (we write  $\nu$  for the outward unit normal).

**Lemma 2.3** (interior balls) *For every  $\theta_0 > 0$  and  $H_0 < \infty$  there exists some  $\delta = \delta(H_0, \theta_0) > 0$  with the following significance: If  $p \in M_{t_0}$  is such that  $H(p) \leq H_0$  and  $\langle -\nu(p), \nu \rangle \geq \theta_0$  for every  $v \in \check{K}_{t_0}$  with  $|v| = 1$ , then we have  $B(p + v, \delta) \subseteq K_{t_0}$  for every  $v \in \check{K}_{t_0}$  with  $|v| \geq 1$ .*

**Proof** First, note that as  $\lim_{\lambda \rightarrow 0} \lambda K_{t_0} = \lim_{\lambda \rightarrow 0} \lambda(K_{t_0} - p)$ , we have  $p + sv \in K_{t_0}$  for every  $v \in \check{K}_{t_0}$  with  $|v| = 1$  and every  $s \geq 0$ . The convexity of  $K_{t_0}$  implies that the function  $(0, \infty) \ni s \mapsto \text{dist}(p + sv, M_{t_0})$  is concave and positive on  $K_{t_0}$ , and thus, nondecreasing. Taking an interior tangent ball at  $p$  of radius  $\alpha/H_0$  yields the desired result. □

**Proposition 2.4** (time-independence of blowdown) *The blowdown  $\check{K}_{t_0}$  is independent of  $t_0 < T_{\text{ext}}(\mathcal{M})$ .*

**Proof** It suffices to prove this in case  $M_t = \partial K_t$  in  $\mathbb{R}^4$  is noncompact and strictly convex. Fix some  $t_0$  and consider the set  $I$  of all  $t \in [t_0, T_{\text{ext}}(\mathcal{M})$  such that  $\check{K}_{t_0} = \check{K}_t$ . Since  $K_{t_2} \subseteq K_{t_1}$  for  $t_2 \geq t_1$  we clearly have that  $\check{K}_{t_2} \subseteq \check{K}_{t_1}$  for  $t_2 \geq t_1$ , so  $I$  is a (potentially degenerate) subinterval of  $[t_0, T_{\text{ext}}(\mathcal{M})$  that contains  $t_0$ . Letting  $t_1 \in I$ , and taking any  $p \in \partial K_{t_1}$ , strict convexity implies that

$$(24) \quad \inf_{v \in \check{K}_{t_1} - \{0\}} \left\langle -\nu(p), \frac{v}{|v|} \right\rangle > 0.$$

By the lemma above, there exists some  $\delta > 0$  such that for every  $|v| \geq 1$ ,  $B(p + v, \delta) \subseteq K_{t_1}$ . Therefore, by avoidance,  $p + v \in K_t$  for all  $t$  with  $|t - t_1| \leq \delta^2/6$ , so all such  $t$  are also in  $I$ .

Thus, if  $I \neq [t_0, T_{\text{ext}}(\mathcal{M}))$  then  $I = [t_0, t_1)$  for some  $t_1 < T_{\text{ext}}(\mathcal{M})$ . We want to show that this is impossible. Suppose, for the sake of contradiction, that there exists some  $v \in \check{K}_{t_0} \setminus \check{K}_{t_1}$ . Given any  $p \in M_{t_1}$ , by smoothness, we can find  $t \in I$  arbitrarily close to  $t_1$  and  $p_t \in M_t$  such  $p_t \rightarrow p$  and such that  $H(p_t) \leq H(p) + 1$ . As above, Lemma 2.3 and avoidance imply that

$$(25) \quad \langle v(p_t), v \rangle \rightarrow 0,$$

so  $v \in T_p M_{t_1}$ . But since  $p \in M_{t_1}$  was arbitrary, this implies that  $M_{t_1}$  splits a line in the direction  $v$ , contradicting the strict convexity of  $M_{t_1}$ .  $\square$

We end this section by giving a (well-known) description of the blowdown in terms of graphical directions:

**Proposition 2.5** (alternative description of blowdown) *Let  $K_t = \partial M_t$  be a noncollapsed strictly convex flow. Then*

$$(26) \quad \check{K}_{t_0} = \{\omega \in \mathbb{R}^{n+1} \text{ such that } \langle v(p), \omega \rangle < 0 \text{ for every } p \in M_{t_0}\} \cup \{0\}.$$

**Proof** Let  $\Omega = \{\omega \in \mathbb{R}^{n+1} \text{ such that } \langle v(p), \omega \rangle < 0 \text{ for every } p \in M_{t_0}\}$ . The containment  $\check{K}_{t_0} \subseteq \Omega \cup \{0\}$  is clear, as for every  $p \in M_{t_0}$  and  $v \in \check{K}_{t_0}$  we have  $p + sv \in K_{t_0}$  for every positive  $s$ . Conversely, if  $v \in \Omega$ , for every  $p \in M_{t_0}$  we have that  $p + sv \in K_{t_0}$  for small values of  $s$ . Thus, if  $p + s_0 v \in M_{t_0}$  for some  $s_0 > 0$  then for  $s > s_0$  with  $s - s_0$  sufficiently small,  $p + sv \in K_{t_0}$ , which contradicts the concavity of the distance to the boundary. Thus, the entire line  $\{p + sv \mid s \in [0, \infty)\}$  is contained in  $K_{t_0}$ , so  $v \in \check{K}_{t_0}$ .  $\square$

### 2.2 The bubble-sheet scale

Let  $\mathcal{M}$  be an ancient mean curvature flow, with a bubble-sheet tangent flow at  $-\infty$ . Given a point  $X = (x, t) \in \mathcal{M}$  and a scale  $r > 0$ , we consider the flow

$$(27) \quad \mathcal{M}_{X,r} = \mathcal{D}_{1/r}(\mathcal{M} - X)$$

which is obtained from  $\mathcal{M}$  by translating  $X$  to the space-time origin and parabolically rescaling by  $1/r$ . Here,  $\mathcal{D}_\lambda(x, t) = (\lambda x, \lambda^2 t)$ .

**Definition 2.6** ( $\varepsilon$ -cylindrical) We say that  $\mathcal{M}$  is  $\varepsilon$ -cylindrical around  $X$  at scale  $r$  if  $\mathcal{M}_{X,r}$  is  $\varepsilon$ -close in  $C^{[1/\varepsilon]}$  in  $B(0, 1/\varepsilon) \times [-2, -1]$  to the evolution of a round shrinking bubble-sheet cylinder with radius  $r(t) = \sqrt{-2t}$  and axis through the origin.

Note that in [10] and [11], being  $\varepsilon$ -cylindrical is defined by closeness to the evolution of a neck, rather than the evolution of a bubble-sheet. We hope that the benefits of the analogies coming from overloading the term outweigh the potential confusion. In any case, throughout the present paper  $\varepsilon$ -cylindrical is always meant in the sense of Definition 2.6.

Given any point  $X = (x, t) \in \mathcal{M}$ , we analyze the solution around  $X$  at the dyadic scales  $r_j = 2^j$ , where  $j \in \mathbb{Z}$ .

**Theorem 2.7** For any small enough  $\varepsilon > 0$ , there is a positive integer  $N = N(\varepsilon) < \infty$  with the following significance. If  $\mathcal{M}$  is an ancient flow asymptotic to a bubble-sheet, which is not the round shrinking bubble-sheet cylinder, then for every  $X \in \mathcal{M}$  there exists an integer  $J(X) \in \mathbb{Z}$  such that

$$(28) \quad \mathcal{M} \text{ is not } \varepsilon\text{-cylindrical around } X \text{ at scale } r_j \text{ for all } j < J(X)$$

and

$$(29) \quad \mathcal{M} \text{ is } \frac{1}{2}\varepsilon\text{-cylindrical around } X \text{ at scale } r_j \text{ for all } j \geq J(X) + N.$$

**Proof** The proof, based on quantitative differentiation (cf [9]), is identical to that of [10, Theorem 2.7].  $\square$

We fix a small enough parameter  $\varepsilon > 0$  quantifying the quality of the bubble-sheet for the rest of the paper.

**Definition 2.8** (bubble-sheet scale) The *bubble-sheet scale* of  $X \in \mathcal{M}$  is defined by

$$(30) \quad Z(X) = 2^{J(X)}.$$

### 3 Foliations and barriers

The proof of [Theorem 1.4](#) relies on the spectral analysis of the equation describing the evolution by mean curvature flow of a graph over the cylinder. Importantly, the flow is not an entire graph over a cylinder, but rather, a graph over a large portion of it. Thus, one needs to introduce some truncation, and control the error introduced by it. Moreover, one needs quantitative bounds on the graphical radius — the radius in which the flow is a small graph over the cylinder. In the neck-singularity setting, both of these goals are achieved in [3] by using a foliation whose leaves are fixed points of the rescaled MCF. In this section, we construct analogues of that foliation which are suited for the analysis over the bubble-sheet cylinder  $\mathbb{R}^2 \times S^1$ .

We recall from Angenent, Daskalopoulos and Sesum that there exists some  $L_0 > 1$  such that for every  $a \geq L_0$  and  $b > 0$  there are self-shrinkers

$$(31) \quad \begin{aligned} \Sigma_a &= \{\text{surface of revolution with profile } r = u_a(x_1), \text{ where } 0 \leq x_1 \leq a\}, \\ \tilde{\Sigma}_b &= \{\text{surface of revolution with profile } r = \tilde{u}_b(x_1), \text{ where } 0 \leq x_1 < \infty\}, \end{aligned}$$

as illustrated in [3, Figure 1]; see also [24]. We will refer to these shrinkers as ADS-shrinkers and KM-shrinkers, respectively. Here, the parameter  $a$  captures where the concave functions  $u_a$  meet the  $x_1$ -axis, namely  $u_a(a) = 0$ , and the parameter  $b$  is the asymptotic slope of the convex functions  $\tilde{u}_b$ , namely  $\lim_{x_1 \rightarrow \infty} \tilde{u}'_b(x_1) = b$ . A detailed description of these shrinkers can be found in [3, Section 8]. We recall:

**Lemma 3.1** [3, Lemmas 4.9 and 4.10] *There exists some  $b_0 > 0$  such that the self-shrinkers  $\Sigma_a$ ,  $\tilde{\Sigma}_b$  and the cylinder  $\Sigma := \{x_2^2 + x_3^2 = 2\} \subset \mathbb{R}^3$  foliate the region*

$$(32) \quad \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_2^2 + x_3^2 \leq b_0^2 x_1^2 \text{ and } x_1 \geq L_0\}.$$

Moreover, denoting by  $\nu_{\text{fol}}$  the outward unit normal of this family, we have that

$$(33) \quad \text{div}(e^{-x^2/4} \nu_{\text{fol}}) = 0,$$

ie the shrinker family forms a calibration for the Gaussian area.

We will now shift and rotate this foliation to construct a suitable foliation in  $\mathbb{R}^4$ :

**Definition 3.2** (bubble-sheet foliation) For every  $a \geq L_0$ , we denote by  $\Gamma_a$  the doubly-rotationally symmetric hypersurface in  $\mathbb{R}^4$  given by

$$(34) \quad \Gamma_a = \{(r \cos \theta, r \sin \theta, x_3, x_4) \in \mathbb{R}^4 \mid \theta \in [0, 2\pi), (r - 1, x_3, x_4) \in \Sigma_a\}.$$

Similarly, for each  $b > 0$  we denote by  $\tilde{\Gamma}_b$  the doubly-rotationally symmetric hypersurface in  $\mathbb{R}^4$  given by

$$(35) \quad \tilde{\Gamma}_b = \{(r \cos \theta, r \sin \theta, x_3, x_4) \in \mathbb{R}^4 \mid \theta \in [0, 2\pi), (r - 1, x_3, x_4) \in \tilde{\Sigma}_b\}.$$

**Lemma 3.3** (foliation lemma) *There exist  $b_0 > 0$  and  $L_1 \geq 3$  such that the hypersurfaces  $\Gamma_a$ ,  $\tilde{\Gamma}_b$  and the cylinder  $\Gamma := \mathbb{R}^2 \times S^1(\sqrt{2})$  foliate the domain*

$$(36) \quad \Omega := \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid x_3^2 + x_4^2 \leq b_0^2(x_1^2 + x_2^2) \text{ and } x_1^2 + x_2^2 \geq L_1^2\}.$$

Moreover, denoting by  $\nu_{\text{fol}}$  the outward pointing unit normal of this foliation, we have that

$$(37) \quad \text{div}(\nu_{\text{fol}} e^{-|x|^2/4}) \leq 0 \quad \text{inside the cylinder,}$$

$$(38) \quad \text{div}(\nu_{\text{fol}} e^{-|x|^2/4}) \geq 0 \quad \text{outside the cylinder.}$$

**Proof** Let  $b_0$  be as in Lemma 3.1, and set  $L_1 = L_0 + 1$ . The fact that  $\Gamma_a$ ,  $\tilde{\Gamma}_b$  and  $\Gamma$  foliate  $\Omega$  follows from Lemma 3.1 and Definition 3.2.

Now, observe that for every element  $\Gamma_*$  in the foliation of  $\Omega$  we have

$$(39) \quad \text{div}(\nu_{\text{fol}} e^{-|x|^2/4}) = (H_{\Gamma_*} - \frac{1}{2} \langle x, \nu_{\text{fol}} \rangle) e^{-|x|^2/4}.$$

Hence, (37) is equivalent to the condition

$$(40) \quad H_{\Gamma_*} \leq \frac{1}{2} \langle x, \nu_{\text{fol}} \rangle.$$

To show (40), note that by symmetry, it suffices to compute the curvatures  $H_{\Gamma_*}$  of  $\Gamma_*$  in the region  $\{x_2 = 0, x_1 > 0\}$ , where we can identify points and unit normals in  $\Gamma_*$  with the corresponding ones in  $\Sigma_*$ ,

by disregarding the  $x_2$ -component. The relation between the mean curvature of a surface  $\Sigma_* \subset \mathbb{R}^3$  and its (unshifted) rotation  $\Gamma_* \subset \mathbb{R}^4$  on points with  $x_2 = 0$  and  $x_1 > 0$  is given by

$$(41) \quad H_{\Gamma_*} = H_{\Sigma_*} + \frac{1}{x_1} \langle e_1, \nu \rangle, \quad \text{where } e_1 = (1, 0, 0).$$

For the ADS-shrinkers, the concavity of  $u_a$  implies  $\langle e_1, \nu_{\text{fol}} \rangle \geq 0$ , so using (41) and the shrinker equation we infer that

$$(42) \quad H_{\Gamma_a} = \frac{1}{2} \langle x - e_1, \nu_{\text{fol}} \rangle + \frac{1}{x_1 - 1} \langle e_1, \nu_{\text{fol}} \rangle \leq \frac{1}{2} \langle x, \nu_{\text{fol}} \rangle,$$

as in  $\Omega \cap \{x_2 = 0, x_1 > 0\}$ , we have  $x_1 \geq L_1 \geq 3$ .

For the KM-shrinkers, the convexity of  $\tilde{u}_b$  implies that  $\langle e_1, \nu_{\text{fol}} \rangle \leq 0$ , so by a similar calculation as in (42),

$$(43) \quad H_{\tilde{\Gamma}_b} \geq \frac{1}{2} \langle x, \nu_{\text{fol}} \rangle. \quad \square$$

**Corollary 3.4** (inner barriers) *The hypersurfaces  $\Gamma_a$  are inner barriers for the renormalized mean curvature flow, in the following sense: Assume  $\{K_\tau\}_{\tau \in [\tau_1, \tau_2]}$  are compact domains, the boundary of which evolve by renormalized mean curvature flow. Assume further that  $K_\tau$  for every  $\tau \in [\tau_1, \tau_2]$  contains the region bounded by  $\Gamma_a$  and  $x_1^2 + x_2^2 = L_1^2$ , and that  $\partial K_\tau \cap \Gamma_a = \emptyset$  for all  $\tau < \tau_2$ . Then*

$$(44) \quad \partial K_{\tau_2} \cap \Gamma_a \subseteq \partial \Gamma_a.$$

**Proof** Lemma 3.3 implies that the vector  $\vec{H} + \frac{1}{2}x^\perp$  points outwards of  $\Gamma_a$ . The result now follows from the maximum principle.  $\square$

**Remark 3.5** (outer barriers) Although this is not needed in the convex case, the hypersurfaces  $\tilde{\Gamma}_b$  are evidently outer barriers for the renormalized mean curvature flow.

## 4 Setting up the fine bubble-sheet analysis

Throughout this section, let  $\mathcal{M}$  be a noncollapsed ancient flow in  $\mathbb{R}^4$  whose tangent flow for  $t \rightarrow -\infty$  is a bubble-sheet. Assume further that  $\mathcal{M}$  is not self-similarly shrinking. Given any space-time point  $X_0$ , we consider the renormalized flow

$$(45) \quad \bar{M}_\tau^{X_0} = e^{\tau/2} (M_{t_0 - e^{-\tau}} - x_0).$$

Then  $\bar{M}_\tau^{X_0}$  converges to the cylinder

$$(46) \quad \Gamma = \mathbb{R}^2 \times S^1(\sqrt{2}) \quad \text{as } \tau \rightarrow -\infty.$$

Since the renormalized MCF is invariant under rotations, the corresponding rotation vector fields appear as Jacobi fields in its linearization. Therefore, to obtain a useful geometric information from the spectral

analysis in the case where the neutral mode is dominant, one needs to make sure that those rotation vector fields are not the dominant ones. There are two ways that have been successfully employed in doing that:

- (i) using a neck-improvement theorem to show that the rotations are nondominant (cf [10, Section 4.3]), or
- (ii) rotating the evolution in such a way that the graphs are orthogonal to all rotations (cf [6, Section 2]).

The difference between the two methods, in terms of the argument, is where those rotations are dealt with: in the latter approach, the labor lies in showing that the modes of the linearization dominate the evolution even after one rotates the surfaces. In the former approach, the analysis in the neutral mode case is harder.

In our current setting, we have found it easier to use the second alternative, as in [6].

After normalizing we can assume that  $Z(X_0) \leq 1$ , and we can find a universal function  $\rho(\tau) > 0$  with

$$(47) \quad \lim_{\tau \rightarrow -\infty} \rho(\tau) = \infty \quad \text{and} \quad -\rho(\tau) \leq \rho'(\tau) \leq 0,$$

such that for every  $S \in \text{SO}(4)$  with  $\angle(S(\Gamma), \Gamma) < \frac{1}{100}\rho(\tau)^{-3}$ , the rotated surface  $S(\bar{M}_\tau^{X_0})$  is the graph of a function  $u = u_S(\cdot, \tau)$  over  $\Gamma \cap B_{2\rho(\tau)}(0)$ , namely

$$(48) \quad \{x + u(x, \tau)v_\Gamma(x) \mid x \in \Gamma \cap B_{2\rho(\tau)}(0)\} \subset S(\bar{M}_\tau^{X_0}),$$

where  $v_\Gamma$  denotes the outward pointing unit normal to  $\Gamma$ , such that

$$(49) \quad \|u(\cdot, \tau)\|_{C^4(\Gamma \cap B_{2\rho(\tau)}(0))} \leq \rho(\tau)^{-2}.$$

For later use, let us also arrange that in the special case where the rotation matrix is the identity, we have the better decay

$$(50) \quad \|u_I(\cdot, \tau)\|_{C^4(\Gamma \cap B_{2\rho(\tau)}(0))} \leq \rho(\tau)^{-4}.$$

We fix a nonnegative smooth cutoff function  $\chi$  satisfying  $\chi(s) = 1$  for  $|s| \leq \frac{1}{2}$  and  $\chi(s) = 0$  for  $|s| \geq 1$ . We consider the truncated function

$$(51) \quad \hat{u}(x, \tau) := u(x, \tau)\chi\left(\frac{r}{\rho(\tau)}\right),$$

where

$$(52) \quad r(x) := \sqrt{x_1^2 + x_2^2}.$$

**Proposition 4.1** (orthogonality) *There exists a differentiable function  $S(\tau) := S^{X_0}(\tau) \in \text{SO}(4)$ , defined for  $\tau$  sufficiently negative, such that, setting  $u := u_{S^{X_0}(\tau)}$ , we have*

$$(53) \quad \int_{\Gamma \cap B_{\rho(\tau)}(0)} \langle Ax, v_\Gamma \rangle \hat{u}(x, \tau) e^{-|x|^2/4} = 0,$$

for every  $A \in o(4)$ . Moreover, the matrix  $A(\tau) = S'(\tau)S(\tau)^{-1} \in o(4)$  satisfies  $A_{12}(\tau) = 0$  and  $A_{34}(\tau) = 0$  for all  $\tau \ll 0$ .

**Proof** Let  $\text{Gr}(2, \mathbb{R}^4)$  be the space of two-dimensional planes through the origin in  $\mathbb{R}^4$ . The rotation group  $O(4)$  acts transitively on  $\text{Gr}(2, \mathbb{R}^4)$  with stabilizer  $O(2) \times O(2)$ , and hence the Grassmannian can be expressed as homogeneous space

$$(54) \quad \text{Gr}(2, \mathbb{R}^4) = \frac{O(4)}{O(2) \times O(2)}.$$

In particular,  $\dim \text{Gr}(2, \mathbb{R}^4) = 6 - 2 = 4$ . Let us select an explicit choice of fibration

$$(55) \quad p: O(4) \rightarrow \text{Gr}(2, \mathbb{R}^4),$$

by declaring that  $p$  maps each rotation matrix to the span of its first two column vectors.

Denote by  $U_\tau$  the open set of all  $R \in O(4)$  satisfying  $\angle(R(\Gamma), \Gamma) < \frac{1}{100}\rho(\tau)^{-3}$ . Now, given any  $\tau \ll 0$  and  $R \in U_\tau$  we can write  $R(\overline{M}_\tau)$  as a graph of a function  $u_R(x, \tau)$  over  $\Gamma \cap B_{2\rho(\tau)}(0)$ . Observe that the expression

$$(56) \quad \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} u_R^2(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right),$$

as well as the relation

$$(57) \quad \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} \langle Ax, \nu_\Gamma \rangle u_R(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right) = 0 \quad \text{for all } A \in o(4),$$

is constant along the fibers of  $p$ . Set  $V_\tau := p(U_\tau)$ , and observe that this is an open neighborhood of  $[x_1 x_2] \in \text{Gr}(2, \mathbb{R}^4)$ .

**Claim 4.2** *Possibly after decreasing  $U_\tau$ , for every  $\tau \ll 0$  there exists a unique  $\pi(\tau) \in V_\tau$  such that*

$$(58) \quad \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} \langle Ax, \nu_\Gamma \rangle u_{\tilde{\pi}(\tau)}(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right) = 0 \quad \text{for all } A \in o(4),$$

where  $\tilde{\pi}(\tau)$  denotes some lift of  $\pi(\tau)$  to  $O(4)$ . Moreover, the function  $\tau \mapsto \pi(\tau)$  is smooth.

**Proof of the claim** We will use the quantitative version of the implicit function theorem.

In order to first construct the required approximate solution, we consider the functional

$$(59) \quad H(\tau, \pi) = \frac{1}{2} \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} u_\pi^2(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right).$$

$\overline{M}_\tau$  is a small graph over the cylinder with axis given by the 2-plane  $[x_1 x_2] \in \text{Gr}(2, \mathbb{R}^4)$ , so for every  $\tau \ll 0$  there is a minimizer  $\pi(\tau)$  for  $H$  in a small neighborhood of  $[x_1 x_2]$ , and we have  $H(\tau, \pi(\tau)) \leq C/\rho(\tau)^4$ . Here, using (50) it is not hard to see that the minimum is indeed attained (well inside) the open set  $V_\tau$ .

Fix some  $\tau \ll 0$  and abbreviate  $\tilde{\pi} := \tilde{\pi}(\tau)$ . Now, let  $R(\eta) \in \text{SO}(4)$  be a one-parameter family of rotations with  $R(0) = \tilde{\pi}$  and  $R'(0) = A\tilde{\pi}$ , where  $A \in o(4)$ . Note that

$$(60) \quad R(\eta)^{-1}(x + u_{R(\eta)}(x, \tau)\nu_\Gamma(x)) \in \overline{M}_\tau.$$



Taking  $(d/d\eta)|_{\eta=0}$  of this expression, observing the resulting vector is of course orthogonal to  $v(x + u_{\tilde{\pi}}v_{\Gamma})$ , and using the condition  $\langle \tilde{\pi}(\Gamma), \Gamma \rangle < \frac{1}{100}\rho(\tau)^{-3}$ , we infer that

$$(61) \quad \frac{d}{d\eta} \Big|_{\eta=0} u_{R(\eta)}(x, \tau) = \langle Ax, v_{\Gamma} \rangle + O\left(\frac{1 + |x|}{\rho(\tau)^2}\right).$$

Using this, we obtain that the Euler–Lagrange equation for  $H$  reads

$$(62) \quad \nabla_A H = \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} u_{\pi}(x, \tau) \langle Ax, v_{\Gamma} \rangle \chi\left(\frac{r}{\rho(\tau)}\right) + O\left(\frac{1}{\rho(\tau)^4}\right) = 0 \quad \text{for all } A \in o(4).$$

Thus, our minimizer  $\pi(\tau)$  is an approximate solution of (58).

Now, to show existence and uniqueness of solutions to (58), as well as smooth dependence, let  $A^1, \dots, A^4$  be the standard basis of

$$(63) \quad \mathcal{A} := \{A \in \text{so}(4) \mid A_{12} = A_{34} = 0\},$$

and define a map  $F : \{(\tau, \pi) \mid \tau \leq \tau_0, \pi \in V_{\tau}\} \rightarrow \mathbb{R}^4$  by

$$(64) \quad F(\tau, \pi)_i = \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} \langle A^i x, v_{\Gamma} \rangle u_{\pi}(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right) \quad \text{for } i = 1, \dots, 4.$$

Then, computing similarly as above we obtain

$$(65) \quad \nabla_{A^i} F_j = \int_{\Gamma \cap B_{\rho(\tau)}(0)} e^{-|x|^2/4} \langle A^i x, v_{\Gamma} \rangle \langle A^j x, v_{\Gamma} \rangle \chi\left(\frac{r}{\rho(\tau)}\right) + O\left(\frac{1}{\rho(\tau)^2}\right).$$

Since the only antisymmetric matrix in  $A \in \mathcal{A}$  such that  $\langle Ax, v_{\Gamma} \rangle \equiv 0$  on  $\Gamma$  is 0, we see that the form  $\nabla_{A^i} F_j$  is positive definite for  $\tau \ll 0$ , and so is in particular invertible.

Note that equation (65) holds in an  $\rho(\tau)^{-3}/100$  neighborhood of  $[x_1 x_2]$  with uniform constants. Thus by the quantitative version of the inverse function theorem, in light of (62) and (65), there is a neighborhood of our fixed time  $\tau$  such that the function  $\pi(\cdot)$  can be chosen in it to satisfy (58). Finally, since by the above,  $F$  is uniformly bounded in spatial  $C^1$ , the size of that neighborhood can be taken to be independent of  $\tau$ . This finishes the proof of the claim. □

Now, we can take a lift  $S(\tau)$  of  $\pi(\tau)$  to  $\text{SO}(4)$ . Moreover, by post-composing with two  $\text{SO}(2)$  rotations, of the first two variables, and of the last two variables, we can further arrange that  $A = S'S^{-1}$  satisfies  $A_{12} = A_{34} = 0$ . This finishes the proof of the proposition. □

We now set

$$(66) \quad \tilde{M}_{\tau}^{X_0} = S^{X_0}(\tau) \bar{M}_{\tau}^{X_0},$$

where  $S^{X_0}(\tau) \in \text{SO}(4)$  if from Proposition 4.1 (orthogonality), and set

$$(67) \quad u := u_{S^{X_0}(\tau)},$$

so  $u$  describes  $\tilde{M}_\tau^{X_0}$  as a graph over the cylinder  $\Gamma$ . Recall that we defined

$$(68) \quad \hat{u}(x, \tau) = u(x, \tau) \chi\left(\frac{r}{\rho(\tau)}\right).$$

Our next task it to show that, despite of the truncation and the rotation, the function  $\hat{u}$  evolves by the linearization of the rescaled MCF equation over the cylinder, up to a small error. Set

$$(69) \quad e_r := \nabla r = \frac{(x_1, x_2, 0, 0)}{r}.$$

Let  $\Delta_\tau = \Delta_\tau^+ \cup \Delta_\tau^-$  be the region bounded by  $\tilde{M}_\tau$  and  $\Gamma$ . Here,  $\Delta_\tau^+$  denotes the region that is outside of  $\Gamma$  and inside of  $\tilde{M}_\tau$ , and  $\Delta_\tau^-$  denotes the region that is inside of  $\Gamma$  outside of  $\tilde{M}_\tau$ .

**Proposition 4.3** (Gaussian area) *For all  $L \in [L_1, \rho(\tau)]$  and  $\tau \ll 0$  we have the Gaussian area estimate*

$$(70) \quad \int_{\tilde{M}_\tau \cap \{r \geq L\}} e^{-|x|^2/4} - \int_{\Gamma \cap \{r \geq L\}} e^{-|x|^2/4} \geq - \int_{\Delta_\tau \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle|.$$

**Proof** Let  $\tilde{M}_\tau^+$  (respectively  $\tilde{M}_\tau^-$ ) be the part of  $\tilde{M}_\tau$  that lies outside (respectively inside) the cylinder. As above, let  $\Delta_\tau^\pm$  be the corresponding parts of  $\Delta_\tau$ , and let  $\Gamma^\pm = \Delta_\tau^\pm \cap \Gamma$ .

Considering the foliation of  $\Omega$  from Lemma 3.3, since  $\text{div}(e^{-|x|^2/4} \nu_{\text{fol}}) \geq 0$  in  $\Delta_\tau^+ \cap \{r \geq L\}$ , the divergence theorem yields that for every  $R > L$  we have

$$(71) \quad \int_{\tilde{M}_\tau^+ \cap \{L \leq r \leq R\}} e^{-|x|^2/4} \langle \nu, \nu_{\text{fol}} \rangle - \int_{\Gamma^+ \cap \{L \leq r \leq R\}} e^{-|x|^2/4} \\ \geq - \int_{\Delta_\tau^+ \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle| - \int_{\Delta_\tau^+ \cap \{r=R\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle|.$$

Similarly, since  $\text{div}(e^{-|x|^2/4} \nu_{\text{fol}}) \leq 0$  in  $\Delta_\tau^- \cap \{r \geq L\}$ , the divergence theorem yields

$$(72) \quad \int_{\Gamma^- \cap \{L \leq r \leq R\}} e^{-|x|^2/4} - \int_{\tilde{M}_\tau^- \cap \{L \leq r \leq R\}} e^{-|x|^2/4} \langle \nu, \nu_{\text{fol}} \rangle \\ \leq \int_{\Delta_\tau^- \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle| + \int_{\Delta_\tau^- \cap \{r=R\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle|.$$

Putting these together, we get

$$(73) \quad \int_{\tilde{M}_\tau \cap \{L \leq r \leq R\}} e^{-|x|^2/4} \langle \nu, \nu_{\text{fol}} \rangle - \int_{\Gamma \cap \{L \leq r \leq R\}} e^{-|x|^2/4} \\ \geq - \int_{\Delta_\tau \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle| - \int_{\Delta_\tau \cap \{r=R\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle|.$$

Using  $|\langle \nu, \nu_{\text{fol}} \rangle| \leq 1$  and  $|\Delta_\tau \cap \{r = R\}| \leq CR^3$  and passing  $R \rightarrow \infty$ , we conclude that

$$(74) \quad \int_{\tilde{M}_\tau \cap \{r \geq L\}} e^{-|x|^2/4} - \int_{\Gamma \cap \{r \geq L\}} e^{-|x|^2/4} \geq - \int_{\Delta_\tau \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle|. \quad \square$$

Next, we have an inverse Poincaré inequality:

**Proposition 4.4** (inverse Poincaré inequality) *The graph function  $u$  satisfies the integral estimates*

$$(75) \quad \int_{\Gamma \cap \{|r| \leq L\}} e^{-|x|^2/4} |\nabla u(x, \tau)|^2 \leq C \int_{\Gamma \cap \{|r| \leq L/2\}} e^{-|x|^2/4} u(x, \tau)^2,$$

$$(76) \quad \int_{\Gamma \cap \{L/2 \leq |r| \leq L\}} e^{-|x|^2/4} u(x, \tau)^2 \leq \frac{C}{L^2} \int_{\Gamma \cap \{|r| \leq L/2\}} e^{-|x|^2/4} u(x, \tau)^2,$$

for all  $L \in [L_1, \rho(\tau)]$  and  $\tau \ll 0$ , where  $C < \infty$  is a numerical constant.

**Proof** The proof is quite similar to that of [6, Proposition 2.3]. Since  $|\langle e_r, \nu_{\text{fol}} \rangle| \leq CL^{-1} |x_3^2 + x_4^2 - 2|$  by [3, Lemma 4.11], we infer that

$$(77) \quad \int_{\Delta_\tau \cap \{r=L\}} e^{-|x|^2/4} |\langle e_r, \nu_{\text{fol}} \rangle| \leq CL^{-1} \int_{\Gamma \cap \{r=L\}} e^{-|x|^2/4} u^2.$$

Thus, Proposition 4.3 (Gaussian area) implies

$$(78) \quad \int_{\tilde{M}_\tau \cap \{r \geq L\}} e^{-|x|^2/4} - \int_{\Gamma \cap \{r \geq L\}} e^{-|x|^2/4} \geq -CL^{-1} \int_{\Gamma \cap \{r=L\}} e^{-|x|^2/4} u^2.$$

On the other hand, we have

$$(79) \quad \int_{\tilde{M}_\tau \cap \{r \leq L\}} e^{-|x|^2/4} - \int_{\Gamma \cap \{r \leq L\}} e^{-|x|^2/4} \geq \int_{\{r \leq L\}} \int_0^{2\pi} e^{-r^2/4} \left[ -Cu^2 + \frac{1}{C} |\nabla^\Gamma u|^2 \right] d\theta dA,$$

where  $C < \infty$  is a numerical constant. Hence, we can do the same computation as in the proof of [6, Proposition 2.3] to obtain the desired result. □

Recall that  $\tilde{M}_\tau$  is expressed as graph of a function  $u(x, \tau)$  over  $\Gamma \cap B_{2\rho(\tau)}(0)$  satisfying the estimate (49). Using that  $\bar{M}_\tau = S(\tau)^{-1} \tilde{M}_\tau$  moves by rescaled mean curvature flow, one obtains:

**Lemma 4.5** (evolution of graph function) *The function  $u(x, \tau)$  satisfies*

$$(80) \quad \partial_\tau u = \mathcal{L}u + E + \langle A(\tau)x, \nu_\Gamma \rangle,$$

where  $A = S'S^{-1}$  and  $\mathcal{L}$  is the linear operator on  $\Gamma$  defined by

$$(81) \quad \mathcal{L}f = \Delta f - \frac{1}{2} \langle x^{\text{tan}}, \nabla f \rangle + f,$$

and where the error term can be estimated by

$$(82) \quad |E| \leq C\rho(\tau)^{-1} (|u| + |\nabla u| + |A(\tau)|) \quad \text{for } \tau \ll 0.$$

**Proof** The proof is identical to the proof of [6, Lemma 2.4]. □

Denote by  $\mathcal{H}$  the Hilbert space of all functions  $f$  on  $\Gamma$  such that

$$(83) \quad \|f\|_{\mathcal{H}}^2 = \int_{\Gamma} \frac{1}{(4\pi)^{3/2}} e^{-|x|^2/4} f^2 < \infty.$$

**Proposition 4.6** (evolution of the truncated graph function) *The truncated graph function  $\hat{u}(x, \tau) = u(x, \tau)\chi(r/\rho(\tau))$  satisfies*

$$(84) \quad \partial_{\tau}\hat{u} = \mathcal{L}\hat{u} + \hat{E} + \langle A(\tau)x, \nu_{\Gamma} \rangle \chi\left(\frac{r}{\rho(\tau)}\right),$$

where

$$(85) \quad \|\hat{E}\|_{\mathcal{H}} \leq C\rho^{-1}(\|\hat{u}\|_{\mathcal{H}} + |A(\tau)|) \quad \text{for } \tau \ll 0.$$

**Proof** We compute

$$(86) \quad \partial_{\tau}\hat{u} = \mathcal{L}\hat{u} + \hat{E} + \langle A(\tau)x, \nu_{\Gamma} \rangle \chi\left(\frac{r}{\rho(\tau)}\right),$$

where

$$(87) \quad \hat{E} = E\chi\left(\frac{r}{\rho(\tau)}\right) - \frac{2}{\rho(\tau)} \frac{\partial u}{\partial r} \chi'\left(\frac{r}{\rho(\tau)}\right) - \frac{1}{\rho(\tau)^2} u \chi''\left(\frac{r}{\rho(\tau)}\right) + \frac{r}{2\rho(\tau)} u \chi'\left(\frac{r}{\rho(\tau)}\right) - \frac{r\rho'(\tau)}{\rho(\tau)^2} u \chi'\left(\frac{r}{\rho(\tau)}\right).$$

Using [Lemma 4.5](#) (evolution of graph function), we deduce that

$$|\hat{E}| \leq C\rho(\tau)^{-1}(|u| + |\nabla^{\Gamma}u| + |A(\tau)|) \quad \text{for } r \leq \frac{1}{2}\rho(\tau).$$

Moreover, since  $|\rho'(\tau)| \leq \rho(\tau)$  and  $\rho(\tau) \rightarrow \infty$ , we obtain

$$|\hat{E}| \leq C|u| + C\rho^{-1}(|\nabla^{\Gamma}u| + |A(\tau)|) \quad \text{for } \frac{1}{2}\rho(\tau) \leq r \leq \rho(\tau).$$

Thus, we can obtain the desired result as in the proof of [\[6, Lemma 2.5\]](#). □

**Lemma 4.7** (estimate for rotation function) *The rotation can be estimated by*

$$(88) \quad |A(\tau)| \leq C\rho^{-1}\|u\|_{\mathcal{H}}.$$

In particular, we have

$$(89) \quad \|\hat{u}_{\tau} - \mathcal{L}\hat{u}\|_{\mathcal{H}} \leq C\rho^{-1}\|\hat{u}\|_{\mathcal{H}}.$$

**Proof** The proof is similar to the one of [\[6, Lemma 2.6\]](#). The conditions  $A_{12}(\tau) = A_{34}(\tau) = 0$  imply

$$(90) \quad |A(\tau)|^2 \leq C \int_{\Gamma} e^{-|x|^2/4} \langle A(\tau)x, \nu_{\Gamma} \rangle^2 \chi\left(\frac{r}{\rho(\tau)}\right).$$

Indeed, since nonzero antisymmetric matrices  $A$  with  $A_{12} = A_{34} = 0$  are the velocity fields of rotations which change the axis of the cylinder, the bilinear form

$$(91) \quad (A, B) := \int_{\Gamma} e^{-|x|^2/4} \langle Ax, v_{\Gamma} \rangle \langle Bx, v_{\Gamma} \rangle \chi\left(\frac{r}{\rho(\tau)}\right)$$

is uniformly positive definite on such matrices.

**Proposition 4.1** (orthogonality) implies that  $\hat{u}$ , hence also  $\partial_{\tau}\hat{u}$ , is orthogonal to all  $\langle Ax, v \rangle$  for each antisymmetric  $A$ . Also  $\mathcal{L}\hat{u}$  is orthogonal to  $\langle Ax, v \rangle$ , as the latter is in the kernel of  $\mathcal{L}$ . Thus,  $\langle Ax, v \rangle$  is orthogonal to  $\partial_{\tau}\hat{u} - \mathcal{L}\hat{u} = \hat{E} + \langle A(\tau)x, v_{\Gamma} \rangle \chi(r/\rho(\tau))$ . From this and (90), we get

$$|A(\tau)|^2 \leq C \int_{\Gamma} e^{-|x|^2/4} |\hat{E}| |\langle A(\tau)x, v_{\Gamma} \rangle| \leq C \|\hat{E}\|_{\mathcal{H}} |A(\tau)| \leq C\rho^{-1} \|\hat{u}\|_{\mathcal{H}} |A(\tau)| + C\rho^{-1} |A(\tau)|^2,$$

where in the last step we have used Proposition 4.6. Consequently,

$$(92) \quad |A(\tau)| \leq C\rho^{-1} \|\hat{u}\|_{\mathcal{H}}.$$

Using Proposition 4.6 once more, we get (89) as well. □

The operator  $\mathcal{L}$  is explicitly given by

$$(93) \quad \mathcal{L} = \frac{\partial^2}{\partial x_1^2} f + \frac{\partial^2}{\partial x_2^2} f + \frac{1}{2} \frac{\partial^2}{\partial \theta^2} f - \frac{1}{2} x_1 \frac{\partial}{\partial x_1} f - \frac{1}{2} x_2 \frac{\partial}{\partial x_2} f + f.$$

Analyzing the spectrum of  $\mathcal{L}$ , the Hilbert space  $\mathcal{H}$  from (83) can be decomposed as

$$(94) \quad \mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_0 \oplus \mathcal{H}_-,$$

where

$$(95) \quad \mathcal{H}_+ = \text{span}\{1, \cos \theta, \sin \theta, x_1, x_2\},$$

$$(96) \quad \mathcal{H}_0 = \text{span}\{x_1 \cos \theta, x_1 \sin \theta, x_2 \cos \theta, x_2 \sin \theta, x_1^2 - 2, x_2^2 - 2, x_1 x_2\}.$$

We have

$$(97) \quad \begin{aligned} \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &\geq \frac{1}{2} \|f\|_{\mathcal{H}}^2 && \text{for } f \in \mathcal{H}_+, \\ \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &= 0 && \text{for } f \in \mathcal{H}_0, \\ \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &\leq -\frac{1}{2} \|f\|_{\mathcal{H}}^2 && \text{for } f \in \mathcal{H}_-. \end{aligned}$$

Consider the functions

$$(98) \quad U_+(\tau) := \|P_+\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \quad U_0(\tau) := \|P_0\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \quad U_-(\tau) := \|P_-\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2,$$

where  $P_+$ ,  $P_0$  and  $P_-$  denote the orthogonal projections to  $\mathcal{H}_+$ ,  $\mathcal{H}_0$  and  $\mathcal{H}_-$ , respectively.

**Theorem 4.8** (Merle–Zaag alternative) For  $\tau \rightarrow -\infty$  either the neutral mode is dominant, ie

$$(99) \quad U_- + U_+ = o(U_0),$$

or the unstable mode is dominant, ie

$$(100) \quad U_- + U_0 \leq C\rho^{-1}U_+.$$

**Proof** Using Proposition 4.6 (evolution of the truncated graph function) and Lemma 4.7 (estimate for the rotation function) we obtain

$$(101) \quad \begin{aligned} \frac{d}{d\tau}U_+(\tau) &\geq U_+(\tau) - C\rho^{-1}(U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \left| \frac{d}{d\tau}U_0(\tau) \right| &\leq C\rho^{-1}(U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \frac{d}{d\tau}U_-(\tau) &\leq -U_-(\tau) + C\rho^{-1}(U_+(\tau) + U_0(\tau) + U_-(\tau)). \end{aligned}$$

Hence, the Merle–Zaag ODE lemma (Lemma B.1) implies the assertion. □

## 5 Bubble-sheet analysis in the neutral mode

In this section, we prove Theorem 1.4 in the case where the neutral mode is dominant. Namely, throughout this section we consider a noncompact ancient noncollapsed flow in  $\mathbb{R}^4$  whose truncated graph function  $\hat{u}(\cdot, \tau)$  satisfies

$$(102) \quad U_- + U_+ = o(U_0).$$

The following lemma gives a rough bound, showing that for  $\tau \rightarrow -\infty$  the function  $U_0$  decays slower than any exponential.

**Lemma 5.1** (rough decay estimate) For any  $\delta > 0$  we have

$$(103) \quad U_0(\tau) \geq e^{\delta\tau}$$

for sufficiently large  $-\tau$ .

**Proof** Given any  $\delta > 0$ , the inequality (101) together with the assumption  $U_- + U_+ = o(U_0)$  implies

$$(104) \quad \left| \frac{d}{d\tau}U_0(\tau) \right| \leq \frac{1}{2}\delta U_0$$

for sufficiently large  $-\tau$ . Rewriting this as

$$(105) \quad \left| \frac{d}{d\tau} \log U_0(\tau) \right| \leq \frac{1}{2}\delta,$$

integration gives  $\log U_0(\tau) \geq -C + \frac{1}{2}\delta\tau$ . This yields the desired result. □

**Proposition 5.2** Every sequence  $\{\tau_i\}$  converging to  $-\infty$  has a subsequence  $\{\tau_{i_m}\}$  such that

$$(106) \quad \lim_{\tau_{i_m} \rightarrow -\infty} \frac{\widehat{u}(\cdot, \tau_{i_m})}{\|\widehat{u}(\cdot, \tau_{i_m})\|_{\mathcal{H}}} = q_{11}(x_1^2 - 2) + q_{22}(x_2^2 - 2) + 2q_{12}x_1x_2$$

in  $\mathcal{H}$ -norm, where  $\{q_{ij}\}$  is a nontrivial seminegative definite  $2 \times 2$ -matrix.

**Proof** By Proposition 4.1 (orthogonality), we have

$$(107) \quad P_0\widehat{u} \in \text{span}\{x_1^2 - 2, x_2^2 - 2, x_1x_2\} \subset \mathcal{H}_0.$$

Therefore, every sequence  $\tau_i \rightarrow -\infty$  has a subsequence  $\{\tau_{i_m}\}$  such that

$$(108) \quad \lim_{\tau_{i_m} \rightarrow -\infty} \frac{\widehat{u}(\cdot, \tau_{i_m})}{\|\widehat{u}(\cdot, \tau_{i_m})\|_{\mathcal{H}}} = \mathcal{Q}(x_1, x_2),$$

with respect to the  $\mathcal{H}$ -norm, where

$$(109) \quad \mathcal{Q}(x_1, x_2) = q_{11}(x_1^2 - 2) + q_{22}(x_2^2 - 2) + 2q_{12}x_1x_2$$

for some nontrivial matrix  $Q = \{q_{ij}\}$ . After an orthogonal change of coordinates in the  $x_1x_2$ -plane we can assume that  $q_{12} = 0$ . Let us show that  $q_{11} \leq 0$  (the same argument yields  $q_{22} \leq 0$ ).

We denote by  $\widetilde{K}_\tau$  the region enclosed by  $\widetilde{M}_\tau$  and denote by  $\mathcal{A}(x'_1, x'_2, \tau)$  the area of the cross-section  $\widetilde{K}_\tau \cap \{(x_1, x_2) = (x'_1, x'_2)\}$ . Explicitly, for  $x_1^2 + x_2^2 \leq \rho(\tau)^2$  we have

$$(110) \quad \begin{aligned} \mathcal{A}(x_1, x_2, \tau) &= \frac{1}{2} \int_0^{2\pi} (\sqrt{2} + u(\theta, x_1, x_2, \tau))^2 d\theta \\ &= 2\pi + \sqrt{2} \int_0^{2\pi} u(\theta, x_1, x_2, \tau) d\theta + \frac{1}{2} \int_0^{2\pi} u(\theta, x_1, x_2, \tau)^2 d\theta. \end{aligned}$$

By Brunn’s concavity principle, the function

$$(111) \quad (x_1, x_2) \mapsto \sqrt{\mathcal{A}(x_1, x_2, \tau)}$$

is concave. In particular, we have

$$(112) \quad \sqrt{\mathcal{A}(x_1, x_2, \tau)} \geq \frac{1}{2} \sqrt{\mathcal{A}(x_1 - 2, x_2, \tau)} + \frac{1}{2} \sqrt{\mathcal{A}(x_1 + 2, x_2, \tau)}.$$

This implies

$$(113) \quad 3 \int_{-1}^1 \int_{-1}^1 \sqrt{\mathcal{A}} dx_2 dx_1 \geq \int_{-3}^3 \int_{-1}^1 \sqrt{\mathcal{A}} dx_2 dx_1.$$

Hence, for sufficiently large  $-\tau_{i_m}$ , combining (108), (110) and (113) yields

$$(114) \quad (3 + o(1))\|\widehat{u}\|_{\mathcal{H}} \int_{-1}^1 \int_{-1}^1 \mathcal{Q} dx_2 dx_1 \geq (1 - o(1))\|\widehat{u}\|_{\mathcal{H}} \int_{-3}^3 \int_{-1}^1 \mathcal{Q} dx_2 dx_1 - O(\|\widehat{u}\|_{\mathcal{H}}^2).$$

Indeed, given any compact intervals  $I_1, I_2 \subset \mathbb{R}$  setting  $u_m = \hat{u}(\cdot, \tau_{i_m})$ , we can compute

$$(115) \quad \int_{I_1} \int_{I_2} (\sqrt{d} - \sqrt{2\pi}) dx_2 dx_1 = \frac{1 \pm o(1)}{2\sqrt{\pi}} \int_{I_1} \int_{I_2} \int_0^{2\pi} u_m d\theta dx_2 dx_1$$

$$= (1 \pm o(1))\sqrt{\pi}\|u_m\|_{\mathcal{H}} \int_{I_1} \int_{I_2} \mathcal{Q} dx_2 dx_1 \pm o(1)\|u_m\|_{\mathcal{H}},$$

which readily implies (114). Now, since  $\|\hat{u}\|_{\mathcal{H}} > 0$  and  $\|\hat{u}\|_{\mathcal{H}} \rightarrow 0$ , taking the limit as  $m \rightarrow \infty$ , we obtain

$$(116) \quad 3 \int_{-1}^1 \int_{-1}^1 \mathcal{Q} dx_2 dx_1 \geq \int_{-3}^3 \int_{-1}^1 \mathcal{Q} dx_2 dx_1.$$

This implies

$$(117) \quad 3q_{11} \int_{-1}^1 \int_{-1}^1 (x_1^2 - 2) dx_2 dx_1 \geq q_{11} \int_{-3}^3 \int_{-1}^1 (x_1^2 - 2) dx_2 dx_1.$$

Since the integral of the left-hand side is negative, while the integral on the right-hand side is positive, we conclude that  $q_{11} \leq 0$ . This proves the proposition.  $\square$

Recall that in this section  $M_t = \partial K_t$  denotes an ancient noncollapsed flow in  $\mathbb{R}^4$  with dominant neutral mode, ie (102) holds. Now, by the reduction from Section 2, we can assume that its blowdown  $\check{K} \equiv \check{K}_{t_0}$  is a convex cone that does not contain any line. Furthermore, rotating coordinates, we can arrange that  $\check{K}$  contains the positive  $x_1$ -axis, namely

$$(118) \quad \{\lambda e_1 \mid \lambda \geq 0\} \subseteq \check{K}.$$

**Theorem 5.3** (blowdown in neutral mode) *If  $M_t = \partial K_t$  is as above, then its blowdown is a halfline, namely*

$$(119) \quad \check{K} = \{\lambda e_1 \mid \lambda \geq 0\}.$$

Moreover,

$$(120) \quad \lim_{\tau \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}} = -c(x_2^2 - 2)$$

in  $\mathcal{H}$ -norm, where  $c = |x_2^2 - 2|_{\mathcal{H}}^{-1}$ .

**Proof** By Proposition 5.2 given any sequence converging to  $-\infty$  we can find a subsequence  $\tau_m$  such that

$$(121) \quad \lim_{\tau_m \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau_{i_m})}{\|\hat{u}(\cdot, \tau_m)\|_{\mathcal{H}}} = q_{11}(x_1^2 - 2) + q_{22}(x_2^2 - 2) + 2q_{12}x_1x_2$$

in  $\mathcal{H}$ -norm, where  $Q = \{q_{ij}\}$  is a nontrivial seminegative definite  $2 \times 2$ -matrix. We will show that

$$(122) \quad \check{K} \subseteq \ker Q.$$



To this end, observe that if  $v$  is a unit vector in the  $x_1x_2$ -plane, denoting by  $w$  the unit vector that is obtained from  $v$  by a  $\pi/2$ -rotation, then

$$(123) \quad \int_0^1 \int_0^1 \mathfrak{Q}(rv + sw) \, dr \, ds = -\frac{5}{3}\text{tr}(Q) + \frac{1}{2}w^T Qv \geq -\text{tr}(Q) > 0.$$

On the other hand, if  $v \notin \ker Q$ , we see that for sufficiently large  $d = d(\langle v, \ker Q \rangle)$ , we have

$$(124) \quad \int_0^1 \int_d^{d+1} \mathfrak{Q}(rv + sw) \, dr \, ds \leq \frac{1}{2}d^2 v^T Qv < 0.$$

Let  $\tilde{K}_\tau$  be, as before, the region enclosed by  $\tilde{M}_\tau$ . Defining  $\mathcal{A}$  as in (110), similarly as in the previous proof we have

$$(125) \quad \int_0^1 \int_a^b \frac{\sqrt{\mathcal{A}(rv + sw, \tau_m)} - \sqrt{2\pi}}{|\hat{u}|_{\mathfrak{H}}} \, dr \, ds \rightarrow \sqrt{\pi} \int_0^1 \int_a^b \mathfrak{Q}(rv + sw) \, dr \, ds$$

as  $m \rightarrow \infty$ . Combining (123), (124) and (125) we see that for every  $m$  sufficiently large, there exists  $r_m, s_m \in [0, 1]$  such that

$$(126) \quad \mathcal{A}(r_m v + s_m w, \tau_m) > \mathcal{A}((r_m + d)v + s_m w, \tau_m).$$

Now, suppose towards a contradiction there is some  $\omega \in \tilde{K} \setminus \ker Q$ . Since  $S(\tau) \rightarrow I$  as  $\tau \rightarrow -\infty$ , for all  $-\tau$  sufficiently large we have

$$(127) \quad \langle P_{12}(S(\tau)\omega), \ker Q \rangle > \frac{1}{2}\langle \omega, \ker Q \rangle,$$

where  $P_{12}$  denotes the projection to the  $\text{span}\{e_1, e_2\}$ . Set

$$(128) \quad v_m := \frac{P_{12}(S(\tau_m)\omega)}{|P_{12}(S(\tau_m)\omega)|}.$$

Take  $v = v_m$  in the previous discussion (and let  $w_m$  be its  $\pi/2$ -rotation). It follows from Proposition 2.5 (alternative description of blowdown) that for any  $x \in \tilde{K}_\tau$  and  $\omega \in \tilde{K}$  we have  $x + \lambda S(\tau)\omega \in \tilde{K}_\tau$  for every  $\lambda \in [0, \infty)$ . Therefore, since  $r_m v_m + s_m w_m \in \tilde{K}_{\tau_m}$ , we see that

$$(129) \quad r_m v_m + s_m w_m + \lambda S(\tau_m)\omega \in \tilde{K}_{\tau_m} \quad \text{for every } \lambda \in [0, \infty).$$

On the other hand, thanks to Brunn’s concavity principle, the function

$$(130) \quad r \mapsto \sqrt{\mathcal{A}(r v_m + s_m w_m, \tau)}$$

is concave, for as long as it does not vanish. Together with (126) this implies that for all  $m$  sufficiently large, the area of the cross-sections is decreasing for  $r > r_m$ , and vanishes at some finite  $r_*$ . This contradicts (129), as the ray would have nowhere to go. This proves (122).

Using the inclusion (122), since  $\ker Q$  is one-dimensional, we infer that  $\tilde{K}$  is one-dimensional. Hence

$$(131) \quad \tilde{K} = \{\lambda e_1 \mid \lambda \geq 0\},$$

$$(132) \quad \ker Q = \{x_1\text{-axis}\}.$$

Finally, since a normalized seminegative  $2 \times 2$ -matrix is uniquely characterized by its one-dimensional kernel, we see that subsequential convergence in (121) in fact entails full convergence, and

$$(133) \quad \lim_{\tau \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}} = -c(x_2^2 - 2),$$

where  $c = |x_2^2 - 2|_{\mathcal{H}}^{-1}$ . This proves the theorem.  $\square$

**Corollary 5.4** (diameter of level sets) *If  $M_t = \partial K_t$  is as above, then given any  $X$  and  $\delta > 0$ , assuming that the neutral mode dominates, we have*

$$\overline{M}_\tau^X \cap \{x_1 = 0\} \subset B_{e^{-\delta\tau}}(0)$$

for sufficiently large  $-\tau$ .

**Proof** By Theorem 5.3 we have

$$(134) \quad \lim_{\tau \rightarrow -\infty} \frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}} = -\frac{(x_2^2 - 2)}{|x_2^2 - 2|_{\mathcal{H}}}.$$

Hence, arguing similarly as above, for sufficiently large  $-\tau$  we obtain

$$(135) \quad \int_1^2 \int_{-1}^1 \sqrt{\mathcal{A}} \, dx_2 \, dx_1 - \sup_{s \in [-1, 1]} \int_{1+s}^{2+s} \int_9^{11} \sqrt{\mathcal{A}} \, dx_2 \, dx_1 \geq (c + o(1)) \|\hat{u}\|_{\mathcal{H}},$$

where  $c > 0$  is a numerical constant. Moreover, by Lemma 5.1 (rough decay estimate) we know that

$$(136) \quad \|\hat{u}\|_{\mathcal{H}} \geq e^{\delta\tau}.$$

Therefore, there is some  $a_+(\tau) \in (1, 2) \times (-1, 1)$  such that

$$(137) \quad \sqrt{\mathcal{A}(a_+(\tau), \tau)} - \sup_{s \in [-1, 1]} \sqrt{\mathcal{A}(a_+(\tau) + se_1 + 10e_2, \tau)} \geq ce^{\delta\tau}.$$

In particular, for all  $\tau$  sufficiently negative this implies that

$$(138) \quad \sqrt{\mathcal{A}(a_+(\tau), \tau)} - \sqrt{\mathcal{A}(a_+(\tau) + 10\langle e_2, S(\tau)e_2 \rangle^{-1} S(\tau)e_2, \tau)} \geq ce^{\delta\tau}.$$

Hence, the concavity of  $(x_1, x_2) \mapsto \sqrt{\mathcal{A}(x_1, x_2, \tau)}$  yields

$$(139) \quad \mathcal{A}(a_+(\tau) + Ce^{-\delta\tau} S(\tau)e_2, \tau) = 0$$

for some constant  $C < \infty$ . Finally, recall that  $\tilde{M}_\tau^X$  is a convex graph with respect to the height function  $S(\tau)e_1$ , and that the level sets of a convex graph monotonically increase. Observing also that  $\langle a_+(\tau), S(\tau)e_1 \rangle \geq 0$  we infer that

$$(140) \quad \mathcal{A}(Ce^{-\delta\tau} S(\tau)e_2, \tau) = 0.$$

Similarly, we can show  $\mathcal{A}(-Ce^{-\delta\tau} S(\tau)e_2, \tau) = 0$ . This proves the corollary.  $\square$

## 6 Bubble-sheet analysis in the unstable mode

In this section, we consider ancient noncollapsed flows with a bubble sheet tangent flow at  $-\infty$ , whose unstable mode is dominant. We will show that each such flow has some nonvanishing expansion parameters associated to it. This is the content of the fine bubble-sheet theorem (Theorem 6.7) and the nonvanishing expansion theorem (Theorem 6.9).

More precisely, throughout this section we assume the tilted rescaled flow  $\tilde{M}_\tau^{X_0}$  around some point  $X_0$  has a dominant unstable mode, ie

$$(141) \quad U_0 + U_- \leq C\rho^{-1}U_+.$$

### 6.1 Analysis of the untilted flow

When dealing with the unstable mode case, it is convenient to work with the renormalized flow  $\bar{M}_\tau^{X_0}$  itself, and not with its tilted version  $\tilde{M}_\tau^{X_0}$ . Our first task is to show that when dealing with the *untilted* renormalized flow around *any* point  $X$ , the unstable mode is still the dominant one.

Given any point  $X$ , for  $\tau \ll 0$  (depending only on  $Z(X)$ ) we can write  $\bar{M}_\tau^X$  as a graph of a function  $\bar{u}(\cdot, \tau)$  over  $\Gamma \cap B_{2\rho(\tau)}(0)$ , where  $\rho = \rho^X(\cdot, \tau)$ , which satisfies (47) and (49). We will work with the truncated function

$$(142) \quad \check{u} = \bar{u}\chi\left(\frac{r}{\rho(\tau)}\right).$$

**Proposition 6.1** (evolution of truncated graph function) *There exists some universal constant  $C < \infty$  such that*

$$(143) \quad \|(\partial_\tau - \mathcal{L})\check{u}\|_{\mathcal{H}} \leq C\rho^{-1}\|\check{u}\|_{\mathcal{H}} \quad \text{for } \tau \ll 0.$$

**Proof** This follows from repeating the argument in Lemma 4.5 and Proposition 4.6 with  $A(\tau) = 0$ .  $\square$

Now, the functions

$$\check{U}_+(\tau) := \|P_+\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \quad \check{U}_0(\tau) := \|P_0\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \quad \check{U}_-(\tau) := \|P_-\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2,$$

satisfy the evolution inequalities (101). Applying the Merle–Zaag ODE lemma (Lemma B.1), we infer that there are universal constants  $C_0, R < \infty$  (where we can assume that  $R > 10^3$ ) such that for every  $X$ , either

$$(144) \quad \check{U}_+ + \check{U}_- = o(\check{U}_0),$$

or

$$(145) \quad \check{U}_0 + \check{U}_- \leq C_0\rho^{-1}\check{U}_+ \quad \text{whenever } \rho \geq R.$$

**Proposition 6.2** (dominant mode) *The unstable mode is dominant for the untilted flow. More precisely, the inequality (145) holds for every  $X$ .*

**Proof** Let us first show that the statement hold for  $X = X_0$ . By assumption, the tilted flow with center  $X_0$  has dominant unstable mode, ie

$$(146) \quad U_0 + U_- \leq C\rho^{-1}U_+.$$

Together with the evolution inequalities (101) this also implies

$$(147) \quad \frac{d}{d\tau}U_+(\tau) \geq (1 - C\rho^{-1})U_+.$$

Note that (147) implies

$$(148) \quad \frac{d}{d\tau}e^{-\frac{9}{10}\tau}U_+(\tau) \geq 0$$

for sufficiently large  $-\tau$ . Hence,

$$(149) \quad \|u(\cdot, \tau)\|_{\mathcal{H}}^2 = (1 + o(1))U_+ \leq Ce^{\frac{9}{10}\tau}.$$

Therefore, Lemma 4.7 yields that

$$(150) \quad |S^{X_0}(\tau) - I| \leq Ce^{\frac{9}{20}\tau}.$$

Recall that

$$(151) \quad \|u(\cdot, \tau)\|_{\mathcal{H}}^2 = \frac{1}{(4\pi)^{3/2}} \int_{\Gamma} \hat{u}_{S^{X_0}(\tau)}(\cdot, \tau)^2 e^{-|x|^2/4},$$

and note that for  $X = X_0$  we have  $\check{u} = \hat{u}_I$ , hence

$$(152) \quad \|\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2 = \frac{1}{(4\pi)^{3/2}} \int_{\Gamma} \hat{u}_I(\cdot, \tau)^2 e^{-|x|^2/4}.$$

Combining (149) and (150) we thus infer that the untilted flow satisfies

$$(153) \quad \|\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2 \leq Ce^{\frac{9}{20}\tau}.$$

On the other hand, if we had  $\check{U}_+ + \check{U}_- = o(\check{U}_0)$ , then arguing as in Lemma 5.1 we would see that  $\|\check{u}(\cdot, \tau)\|_{\mathcal{H}}^2 \geq e^{\delta\tau}$  for every  $\delta > 0$  and  $-\tau$  sufficiently large, which is inconsistent with (153). Thus, for  $X = X_0$ , we indeed get

$$(154) \quad \check{U}_0 + \check{U}_- \leq C\rho^{-1}\check{U}_+.$$

Finally, any neck centered at a general point  $X$  merges with the neck centered at  $X_0$  as  $\tau \rightarrow -\infty$ . Thus, (145) holds for every  $X$ . □

Recapping, for any center  $X$  we therefore have

$$(155) \quad \check{U}_0 + \check{U}_- \leq C_0 \rho^{-1} \check{U}_+$$

and, thanks to the evolution inequalities (101), also

$$(156) \quad \frac{d}{d\tau} \check{U}_+ \geq (1 - C_0 \rho^{-1}) \check{U}_+$$

whenever  $\rho \geq R$ , where  $R \geq 10^3$  and  $C_0$  are some universal constants. Increasing  $R$  further, we can also assume that  $R \geq 10C_0$ .

### 6.2 Graphical radius

The goal of this section is to prove Proposition 6.4, which gives a lower bound for the optimal graphical radius.

We now fix some  $\varepsilon < 1/R$  in the definition of the bubble-sheet scale (Definition 2.6). In what follows,  $C < \infty$  and  $\mathcal{T} > -\infty$  will denote constants that are allowed to change from line to line, but depend *only* on an upper bound of  $Z(X)$ . In particular, our initial choice of graphical radius satisfies  $\rho^X(\tau) \geq R$  for  $\tau \in (-\infty, \mathcal{T}]$  and (155) and (156) hold for all  $\tau \in (-\infty, \mathcal{T}]$ .

**Lemma 6.3** (decay estimate) *There exists some constant  $C < \infty$ , depending only on an upper bound for  $Z(X)$ , such that for  $\tau \leq \mathcal{T}$ ,*

$$(157) \quad \|\check{u}(\cdot, \tau)\|_{\mathcal{H}} \leq C e^{\frac{9}{20}\tau},$$

$$(158) \quad \|\check{u}(\cdot, \tau)\|_{C^{10}(\{r \leq 100\})} \leq C e^{\frac{9}{20}\tau}.$$

**Proof** Since  $\bar{M}_{\mathcal{T}}^X$  is an  $\varepsilon$ -graph over  $\Gamma \cap B_{2\rho^X}(0)$ , we get  $\|\check{u}(\cdot, \mathcal{T})\|_{\mathcal{H}} \leq 1$ . Note that (156) implies  $(d/d\tau)(e^{-\frac{9}{10}\tau} \check{U}_+) \geq 0$  for  $\tau \leq \mathcal{T}$ . Hence,

$$(159) \quad e^{-\frac{9}{10}\tau} \check{U}_+ \leq e^{-\frac{9}{10}\mathcal{T}} \|\check{u}(\cdot, \mathcal{T})\|_{\mathcal{H}}^2 \leq C.$$

This gives the first inequality (157). The second inequality (158) follows from (157) by parabolic estimates. Indeed, one can first establish an  $L^\infty$ -bound via De Giorgi–Nash–Moser iteration (more precisely, one can apply Theorem A.1 with  $k = 0$ ), and then upgrade this to a  $C^{10}$ -bound via standard Schauder theory. □

**Proposition 6.4** (improved graphical radius) *There exists some  $\mathcal{T} > -\infty$ , depending only on an upper bound for  $Z(X)$ , such that*

$$(160) \quad \bar{\rho}(\tau) = e^{-\frac{1}{9}\tau}$$

*is a graphical radius function satisfying (47) and (49) for  $\tau \leq \mathcal{T}$ .*

**Proof** We recall from [3, Theorem 8.2] that the profile function  $u_a$  of the ADS-barriers satisfies

$$(161) \quad u_a(x) \geq \sqrt{2} \left( 1 - \frac{x^2}{a^2} \right)$$

on the interval  $0 \leq x \leq a$ , and

$$(162) \quad u_a(x) \leq \sqrt{2} \left( 1 - \frac{x^2 - 10}{1000a^2} \right)$$

on the interval  $8 \leq x \leq 100$ , provided that  $a$  is large enough.

The upper bound for  $u_a$  and Lemma 6.3 (decay estimate) imply that  $\{r = 10\} \cap \Gamma_a$  is enclosed by  $\bar{M}_\tau^X$  for  $a^2 \leq (1/C)e^{-\frac{9}{20}\tau}$ . Since  $\Gamma_a$  is enclosed by  $\bar{M}_\tau^X$  for  $\tau \ll 0$ , the maximum principle guarantees that

$$(163) \quad \bar{M}_\tau^X \text{ encloses } \{r \geq 10\} \cap \Gamma_a \text{ for } a^2 \leq Ce^{-\frac{9}{20}\tau}.$$

Hence, by the estimate (158), the convexity of  $\bar{M}_\tau^X$  and the lower bound for  $u_a$ , it follows that  $\bar{M}_\tau^X$  is graphical over the cylinder up to a radius  $(1/C)e^{-\frac{9}{60}\tau}$ , with  $C^0$ -norm less than  $Ce^{\frac{9}{60}\tau}$ .

We will now upgrade this  $C^0$ -estimate to a bound for the regularity scale by arguing similarly as in the proof of [10, Proposition 4.10]. Specifically, for any  $X' = (x', t') \in \mathcal{M}$  denote by  $R(X')$  the maximal radius  $r$  such that  $|A| \leq 1/r$  in the parabolic pall  $P(X', r)$ . Suppose towards a contradiction that  $R(X') \ll \sqrt{-t'}$  for some  $X'$  in the unrenormalized flow in the region under consideration. Consider Huisken's monotone quantity  $\Theta_{X'}(r_j)$  from [20] centered at  $X'$  at scales  $r_j = 2^j R(X')$ . Given any  $\delta > 0$  by quantitative differentiation [9] there exists some  $j \in \{1, \dots, \lceil 2/\delta \rceil\}$  such that  $\Theta_{X'}(r_{j+1}) - \Theta_{X'}(r_{j-1}) < \delta$ , and consequently the flow must be  $\varepsilon$ -close to a self-shrinker at scale  $r_j$  centered at  $X'$ , where  $\varepsilon(\delta) \rightarrow 0$  for  $\delta \rightarrow 0$ . Fixing  $\delta > 0$  small enough we can arrange that  $\varepsilon \ll 1$  and  $r_j \ll \sqrt{-t'}$ . However, since in our noncollapsed setting the only self-shrinkers are round spheres, necks and bubble-sheets, this contradicts our graphical  $C^0$ -estimate.

Finally, interpolating the regularity scale estimate and the  $C^0$ -estimate, we see that up to a radius of  $2e^{-\frac{1}{9}\tau}$ , the hypersurface  $\bar{M}_\tau^X$  is a  $C^1$ -graph over the cylinder with norm less than  $\varepsilon_0 e^{\frac{9}{60}\tau}$ . Hence, using standard parabolic Schauder estimates for the renormalized flow around any point within that radius  $e^{-\frac{1}{9}\tau}$  of the point  $X$  yields the desired result.  $\square$

From now on, we work with  $\rho = \bar{\rho}$  from Proposition 6.4 (improved graphical radius), and in particular define  $\check{u}, \check{U}_0, \check{U}_\pm, \dots$  with respect to this improved graphical radius. Note that the unstable mode is still dominant.

**Corollary 6.5** (sharp decay estimate) *There exist constants  $C < \infty$  and  $\mathcal{T} > -\infty$ , depending only an upper bound for  $Z(X)$ , such that for  $\tau \leq \mathcal{T}$ ,*

$$(164) \quad \|\check{u}\|_{\mathcal{H}} \leq Ce^{\tau/2},$$

$$(165) \quad \|\check{u}(\cdot, \tau)\|_{C^{10}(\{r \leq 100\})} \leq Ce^{\tau/2}.$$

**Proof** Combining the evolution inequality (156), Lemma 6.3 (decay estimate) and Proposition 6.4 (improved graphical radius) yields

$$(166) \quad \frac{d}{d\tau}(e^{-\tau}\check{U}_+) \geq -Ce^{-\tau+\frac{1}{9}\tau+\frac{9}{10}\tau} = -Ce^{\frac{1}{90}\tau}$$

for all  $\tau \leq \mathcal{T}$ . Thus,  $\check{U}_+ \leq Ce^\tau$ . Since the unstable mode is dominant, this proves that  $\|\check{u}\|_{\mathcal{H}} \leq Ce^{\tau/2}$ .

Moreover, Proposition 6.4 (improved graphical radius), inequality (155) and Proposition 6.1 (evolution of truncated graph function) then give

$$(167) \quad \check{U}_0 + \check{U}_- + \|(\partial_\tau - \mathcal{L})\check{u}\|_{\mathcal{H}}^2 \leq Ce^{\frac{10}{9}\tau}.$$

Using this, (165) follows from parabolic estimates; see Theorem A.1 for details. □

### 6.3 The fine bubble-sheet theorem

The goal of this section is to prove Theorem 6.7. Recalling the basis of  $\mathcal{H}_+$  from (95) we can write

$$(168) \quad P_+\check{u}^X = a_0^X(\tau) + a_1^X(\tau)x_1 + a_2^X(\tau)x_2 + a_3^X(\tau)\cos\theta + a_4^X(\tau)\sin\theta,$$

where the superscript is to remind us that all these quantities can (a priori) depend on the center point  $X$ .

**Proposition 6.6** (estimate for coefficients) *The coefficients defined in (168) satisfy, for  $\tau \leq \mathcal{T}$ , the estimates*

$$(169) \quad |a_0^X(\tau)| \leq Ce^{\frac{11}{18}\tau},$$

$$(170) \quad \sum_{i=1}^4 |e^{-\tau/2}a_i^X(\tau) - \bar{a}_i^X| \leq Ce^{\frac{1}{9}\tau},$$

where  $\bar{a}_i^X$  are numbers that might depend on  $X$ .

**Proof** Letting  $\check{E} := (\partial_\tau - \mathcal{L})\check{u}$ , and using  $\mathcal{L}1 = 1$ , we compute

$$(171) \quad \frac{d}{d\tau}a_0^X(\tau) = \left(\frac{e}{2\pi}\right)^{\frac{1}{4}} \int (\mathcal{L}\check{u} + \check{E}) \frac{1}{4\pi} e^{-|x|^2/4} = a_0^X(\tau) + \left(\frac{e}{2\pi}\right)^{\frac{1}{4}} \int \frac{\check{E}}{4\pi} e^{-|x|^2/4}.$$

Hence, using Proposition 6.1, Proposition 6.4 and Corollary 6.5, we obtain

$$(172) \quad \left| \frac{d}{d\tau}(e^{-\tau}a_0^X(\tau)) \right| \leq Ce^{-\tau}\|\check{E}\|_{\mathcal{H}} \leq Ce^{-\tau+\tau/2+\tau/9} \leq Ce^{-\frac{7}{18}\tau}.$$

Integrating this from  $\tau$  to  $\mathcal{T}$  implies (169).

In a similar manner, using  $\mathcal{L}x_i = \frac{1}{2}x_i$  for  $i = 1, \dots, 4$ , we get

$$(173) \quad \left| \frac{d}{d\tau}(e^{-\tau/2}a_i^X(\tau)) \right| \leq Ce^{-\tau/2}\|\check{E}\|_{\mathcal{H}} \leq Ce^{\tau/9},$$

so integrating from  $-\infty$  to  $\tau$  yields (170) with

$$(174) \quad \bar{a}_i^X = \lim_{\tau \rightarrow -\infty} e^{-\tau/2}a_i^X(\tau). \quad \square$$

**Theorem 6.7** (the fine bubble-sheet theorem) *Let  $\{M_t\}$  be an ancient noncollapsed flow in  $\mathbb{R}^4$ , with a bubble sheet tangent flow at  $-\infty$ , whose unstable mode is dominant. Then there exists some constants  $a_1, \dots, a_4$ , independent of the center point  $X = (x_0^1, x_0^2, x_0^3, x_0^4, t_0)$ , such that*

$$(175) \quad \bar{a}_1^X = a_1, \quad \bar{a}_2^X = a_2, \quad \bar{a}_3^X = a_3 - x_0^3, \quad \bar{a}_4^X = a_4 - x_0^4.$$

Moreover, for every center point  $X$ , the truncated graph function  $\check{u}^X(\cdot, \tau)$  of the renormalized flow  $\bar{M}_\tau^X$  satisfies, for  $\tau \leq \mathcal{T}$ , the estimates

$$(176) \quad \|\check{u}^X - e^{\tau/2}(a_1x_1 + a_2x_2 + \bar{a}_3^X \cos \theta + \bar{a}_4^X \sin \theta)\|_{\mathcal{H}} \leq Ce^{\frac{5}{9}\tau},$$

$$(177) \quad \|\check{u}^X - e^{\tau/2}(a_1x_1 + a_2x_2 + \bar{a}_3^X \cos \theta + \bar{a}_4^X \sin \theta)\|_{L^\infty(\{r \leq 100\})} \leq Ce^{\frac{19}{36}\tau},$$

where  $C < \infty$  and  $\mathcal{T} > -\infty$  only depend on an upper bound on the bubble sheet scale  $Z(X)$ .

**Proof** Consider the difference

$$(178) \quad D^X := \check{u}^X - e^{\tau/2}(\bar{a}_1^X x_1 + \bar{a}_2^X x_2 + \bar{a}_3^X \cos \theta + \bar{a}_4^X \sin \theta).$$

Using Proposition 6.6 (estimate for coefficients) we see that

$$(179) \quad |D^X| \leq |\check{u}^X - P_+ \check{u}^X| + C(|x| + 1)e^{\frac{11}{18}\tau}.$$

Since by (167) we have  $\check{U}_0 + \check{U}_- \leq Ce^{\frac{10}{9}\tau}$ , it follows that

$$(180) \quad \|D^X\|_{\mathcal{H}} \leq Ce^{\frac{5}{9}\tau},$$

which proves (176) modulo the claim about the coefficients.

Next, we observe that

$$(181) \quad \|D^X\|_{L^2(\{r \leq 100\})} \leq C \|D^X\|_{\mathcal{H}} \leq Ce^{\frac{5}{9}\tau}$$

and, using Corollary 6.5 (sharp decay estimate), that

$$(182) \quad \|\nabla^3 D^X\|_{L^2(\{r \leq 100\})} \leq C \|\hat{u}^X\|_{C^3(\{r \leq 100\})} + Ce^{\tau/2} \leq Ce^{\tau/2}.$$

Applying Agmon's inequality, this yields

$$(183) \quad \|D^X\|_{L^\infty(\{r \leq 100\})} \leq C \|D^X\|_{L^2(\{r \leq 100\})}^{\frac{1}{2}} \|D^X\|_{H^3(\{r \leq 100\})}^{\frac{1}{2}} \leq Ce^{\frac{19}{36}\tau}.$$

Finally, let us show that the parameters  $a_1, \dots, a_4$ , defined via (175), are independent of  $X$ .

First, let us show that they are independent of time translation. Denote by  $\check{u}^{X'}$  the function obtained by considering the renormalized mean curvature flow with center  $X' = (x_0^1, x_0^2, x_0^3, x_0^4, 0)$ . A direct calculation shows that for  $\theta \in (0, 2\pi]$  and  $x_1^2 + x_2^2 \leq 100$ , we have

$$(184) \quad \check{u}^X \left( \frac{x_1}{\sqrt{1+t_0e^\tau}}, \frac{x_2}{\sqrt{1+t_0e^\tau}}, \theta, \tau - \log(1+t_0e^\tau) \right) = \frac{1}{\sqrt{1+t_0e^\tau}} (\sqrt{2} + \check{u}^{X'}(x_1, x_2, \theta, \tau)) - \sqrt{2}.$$



This implies

$$(185) \quad \|\check{u}^X - \check{u}^{X'}\|_{L^\infty(\{r \leq 100\})} = o(e^{\tau/2}),$$

and thus together with (183) yields  $\bar{a}_i^X = \bar{a}_i^{X'}$  for every  $i = 1, \dots, 4$ .

Comparing the renormalized flows with center  $X' = (x_0^1, x_0^2, x_0^3, x_0^4, 0)$  and center  $X'' = ((0, 0, 0, 0), 0)$ , we need to relate both the parameters of the functions  $\check{u}^{X'}$  and  $\check{u}^{X''}$  which describe the same point in the original flow, and the distance of such a point from the respective axii. This leads to

$$(186) \quad \begin{aligned} \sqrt{2} + \check{u}^{X'}(x_1 - x_0^1 e^{\tau/2}, x_2 e - x_0^2 e^{\tau/2}, \theta + O(e^{\tau/2}), \tau) \\ = \text{dist}((\sqrt{2} + \check{u}^{X''}(x_1, x_2, \theta, \tau))(\cos \theta, \sin \theta), e^{\tau/2}(x_0^3, x_0^4)). \end{aligned}$$

By Taylor expansion and Corollary 6.5 we have

$$(187) \quad \begin{aligned} \text{dist}((\sqrt{2} + \check{u}^{X''}(x_1, x_2, \theta, \tau))(\cos \theta, \sin \theta), e^{\tau/2}(x_0^3, x_0^4)) \\ = \sqrt{2} + \check{u}^{X''}(x_1, x_2, \theta, \tau) - x_0^3 \cos \theta e^{\tau/2} - x_0^4 \sin \theta e^{\tau/2} + o(e^{\tau/2}). \end{aligned}$$

Together with (183), the above formulas imply that

$$(188) \quad \bar{a}_1^{X'} = \bar{a}_1^{X''}, \quad \bar{a}_2^{X'} = \bar{a}_2^{X''}, \quad \bar{a}_3^{X'} = \bar{a}_3^{X''} - x_0^3, \quad \bar{a}_4^{X'} = \bar{a}_4^{X''} - x_0^4.$$

This finishes the proof of the theorem. □

### 6.4 The nonvanishing expansion theorem

Our next goal is to show that  $a_1$  and  $a_2$  cannot simultaneously vanish.

We decompose  $\mathcal{H}_+$  into  $\mathcal{H}_{1/2} = \text{span}\{x_1, x_2, \cos \theta, \sin \theta\}$  and  $\mathcal{H}_1 = \text{span}\{1\}$ . Also, we define  $P_{1/2}$  and  $P_1$  as the projections to  $\mathcal{H}_{1/2}$  and  $\mathcal{H}_1$ , respectively. In addition, we denote  $\check{U}_{1/2} = \|P_{1/2}\check{u}\|_{\mathcal{H}}^2$  and  $\check{U}_1 = \|P_1\check{u}\|_{\mathcal{H}}^2$ .

**Lemma 6.8** (decay if coefficients vanished) *If the center  $X$  was such that  $\bar{a}_1^X = \dots = \bar{a}_4^X = 0$ , then there would be a constant  $K_0 > 0$  such that*

$$(189) \quad \check{U}_1 = K_0 e^{2\tau} (1 + O(e^{\frac{1}{9}\tau})),$$

and moreover we would have

$$(190) \quad \check{U}_{1/2} + \check{U}_0 + \check{U}_- \leq C e^{\frac{1}{9}\tau} \check{U}_1.$$

**Proof** Assuming  $\bar{a}_1^X = \dots = \bar{a}_4^X = 0$ , Proposition 6.6 implies

$$(191) \quad a_0^X(\tau)^2 + \dots + a_4^X(\tau)^2 \leq C e^{\frac{11}{9}\tau},$$

hence

$$(192) \quad \check{U}_1 + \check{U}_{1/2} \leq e^{\frac{11}{9}\tau}.$$

Moreover, since the unstable mode is dominant, we have

$$(193) \quad \check{U}_0 + \check{U}_- \leq C e^{\frac{1}{9}\tau} (\check{U}_1 + \check{U}_{1/2}).$$

Now, using Proposition 6.1 we get the evolution inequalities

$$(194) \quad \left| \frac{d}{d\tau} \check{U}_{1/2} - \check{U}_{1/2} \right| \leq C e^{\frac{1}{9}\tau} (\check{U}_{1/2} + \check{U}_1),$$

$$(195) \quad \left| \frac{d}{d\tau} \check{U}_1 - 2\check{U}_1 \right| \leq C e^{\frac{1}{9}\tau} (\check{U}_{1/2} + \check{U}_1).$$

Applying the Merle–Zaag ODE lemma (Lemma B.1) with  $U_0 = e^{-\tau} \check{U}_{1/2}$ ,  $U_- = 0$ , and  $U_+ = e^{-\tau} \check{U}_1$ , we get either  $\check{U}_1 = o(\check{U}_{1/2})$  or  $\check{U}_{1/2} \leq C e^{\frac{1}{9}\tau} \check{U}_1$ . In the former case, arguing as in Lemma 5.1 we could infer that  $e^{-\tau} \hat{U}_{1/2} \geq e^{\frac{1}{9}\tau}$  for every  $-\tau$  sufficiently large, contradicting (192). Hence,

$$(196) \quad \check{U}_{1/2} \leq C e^{\frac{1}{9}\tau} \check{U}_1.$$

Together with (193) this proves the estimate (190).

Moreover, using (196), our differential inequality takes the form

$$(197) \quad \left| \frac{d}{d\tau} \check{U}_1 - 2\check{U}_1 \right| \leq C e^{\frac{1}{9}\tau} \check{U}_1,$$

which can be rewritten as

$$(198) \quad \left| \frac{d}{d\tau} (\log(e^{-2\tau} \check{U}_1)) \right| \leq C e^{\frac{1}{9}\tau}.$$

Integrating this from  $-\infty$  to  $\tau$  gives that there exists some constant  $\tilde{K}_0$  such

$$(199) \quad |\log(e^{-2\tau} \check{U}_1) - \tilde{K}_0| \leq C e^{\frac{1}{9}\tau} \quad \text{for all } \tau \leq \mathcal{F}.$$

Exponentiating both sides and using the approximation  $e^x \approx 1 + x$  for small  $x \in \mathbb{R}$  give (189). In particular,  $K_0 = e^{\tilde{K}_0} > 0$ . □

**Theorem 6.9** (the nonvanishing expansion theorem) *The coefficients from the fine-bubble sheet theorem satisfy  $|a_1| + |a_2| > 0$ .*

**Proof** Suppose towards a contradiction that  $a_1 = a_2 = 0$ . Then, we can choose a point  $X$  such that  $\bar{a}_1^X = \dots = \bar{a}_4^X = 0$ . By Lemma 6.8 and parabolic estimates (see Theorem A.1) we get

$$(200) \quad \|u(\cdot, \tau)\|_{C^4(\{r \leq 100\})} \leq C e^\tau,$$

$$(201) \quad x_3^2 + x_4^2 = 2(1 + K e^\tau) + o(e^\tau) \quad \text{on } \{r \leq 100\},$$

where  $K = (2e/\pi)^{1/4} K_0^{1/2}$ . Here, we have determined the constant  $K$  using the identity

$$(202) \quad \|2^{-1/2} K e^\tau\|_{\mathcal{H}^2}^2 = K_0 e^{2\tau} = \check{U}_1 + O(e^{\frac{10}{9}\tau}),$$

which holds since

$$(203) \quad \int_{\Sigma} e^{-|x|^2/4} = 2^{\frac{3}{2}} \pi e^{-1/2} \left( \int_{-\infty}^{\infty} e^{-z^2/4} dz \right)^2 = \left( \frac{2\pi}{e} \right)^{\frac{1}{2}} (4\pi)^{\frac{3}{2}}.$$

Hence, the rescaled flow with the center  $X' = X + (0, K)$  satisfies

$$(204) \quad x_3^2 + x_4^2 = 2 + o(e^\tau),$$

uniformly on  $\{r \leq 100\}$ ; see also [3, Lemma 5.11] for a more detailed explanation. Since we re-centered by shifting only in time direction, the new point  $X'$  still satisfies  $\bar{a}_1^{X'} = \dots = \bar{a}_4^{X'} = 0$ , so Lemma 6.8 gives some  $K'_0 > 0$  such that

$$(205) \quad e^{-2\tau} \check{U}_1 = K'_0 + O(e^{\frac{1}{9}\tau}).$$

Since  $K'_0 \neq 0$ , this contradicts (204). This proves the theorem. □

## 7 Conclusion in the unstable mode case

The goal of this section is to prove the following theorem.

**Theorem 7.1** (unstable mode) *The only noncompact ancient noncollapsed flow in  $\mathbb{R}^4$ , with bubble-sheet tangent flow at  $-\infty$ , whose unstable mode is dominant, is  $\mathbb{R} \times 2d$ -bowl.*

**Proof** By the reduction from Section 2 it is enough to prove that if the unstable mode is dominant, then its blowdown contains a line.

So let  $M_t = \partial K_t$  be a noncompact ancient noncollapsed flow in  $\mathbb{R}^4$ , with bubble-sheet tangent flow at  $-\infty$ , whose unstable mode is dominant, and suppose towards a contradiction that its blowdown  $\check{K}$  does not contain a line. Then the flow is strictly convex, and  $\check{K}$  is a halfline or a wedge of angle less than  $\pi$  in  $\mathbb{R}^2 \times \{0\}$ . Choosing suitable coordinates we can assume that  $\check{K}$  is symmetric across the  $x_1$ -axis, and is contained in the half space  $\{x_1 \geq 0\}$ . By translating, we may also assume that  $0 \in M_0$  is the point in  $M_0$  with smallest  $x_1$ -value. This implies that for every  $h > 0$  there exist a unique point  $x_h^\pm \in M_0 \cap \{x_1 = h\}$  at which  $x_2$  is maximized/minimized.

By the fine bubble-sheet theorem (Theorem 6.7) and the nonvanishing expansion theorem (Theorem 6.9) there exists expansion parameters  $a_1, a_2$  associated to our flow such that  $|a_1| + |a_2| > 0$ .

**Claim 7.2** (bubble-sheet scale) *There exists some constant  $C < \infty$  such that*

$$(206) \quad \sup_h Z(x_h^\pm) \leq C.$$

**Proof of the claim** We will argue as in the proofs of [10, Proposition 5.8] and [11, Proposition 6.2].

Suppose towards a contradiction that  $Z(x_{h_i}^\pm) \rightarrow \infty$  for some sequence  $\{h_i\}$  with  $\lim_{i \rightarrow \infty} h_i = \infty$ . Let  $\mathcal{M}^i$  be the sequence of flows obtained by shifting  $x_{h_i}^\pm$  to the origin, and parabolically rescaling by  $Z(x_{h_i}^\pm)^{-1}$ .

By [17, Theorem 1.14] we can pass to a subsequential limit  $\mathcal{M}^\infty$ , which is an ancient noncollapsed flow that is weakly convex and smooth until it becomes extinct. Note also that  $\mathcal{M}^\infty$  has bubble-sheet tangent flow at  $-\infty$ .

We next observe that,  $\mathcal{M}^\infty$  cannot be a round shrinking  $\mathbb{R}^2 \times S^1$ . Indeed, if such a cylinder became extinct at time 0 that would contradict the definition of the bubble-sheet scale, and if it became extinct at some later time that would contradict the fact that  $M_0^\infty \cap (\mathbb{R}^2 \times \{0\})$  is a strict subset of  $\mathbb{R}^2 \times \{0\}$  by construction.

Thus, by Theorem 4.8 (Merle–Zaag alternative) for the flow  $\mathcal{M}^\infty$  either the neutral mode is dominant or the unstable mode is dominant. If the neutral mode is dominant, then for large  $i$ , this contradicts the fact that  $\mathcal{M}^i$  has dominant unstable mode. Indeed, on the one hand by Lemma 5.1 (rough decay estimate) and Theorem 5.3 (blowdown in neutral mode) the hypersurfaces  $\overline{M}_\tau^{\infty,0}$  have some definite inwards quadratic bending, but on the other hand by the fine-bubble sheet theorem (Theorem 6.7) the hypersurfaces  $\overline{M}_\tau^{i,0}$  converge exponentially fast to the bubble sheet  $\Gamma$ . Since  $\overline{M}_\tau^{i,0}$  converges locally smoothly to  $\overline{M}_\tau^{\infty,0}$  this gives the desired contradiction for  $i$  large enough. If the unstable mode is dominant, then by the fine-bubble sheet theorem (Theorem 6.7) and the nonvanishing expansion theorem (Theorem 6.9) the limit  $\mathcal{M}^\infty$  has some expansion parameters  $a_1^\infty, a_2^\infty$  that do not vanish simultaneously. However, this contradicts the fact that the expansion parameters of  $\mathcal{M}^i$  are obtained from the expansion parameters  $(a_1, a_2)$  of  $\mathcal{M}$  by scaling by  $Z(x_{h_i}^+)^{-1} \rightarrow 0$ . This concludes the proof of the claim.  $\square$

Continuing the proof of Theorem 7.1, let  $h_i \rightarrow \infty$  and consider the sequence  $\mathcal{M}^i := \mathcal{M} - (x_{h_i}^+, 0)$ , which is obtained by translating in space time without rescaling. Taking a subsequential limit, Claim 7.2 implies that this limit  $\mathcal{M}^\infty$  is an ancient noncollapsed flow with a bubble-sheet tangent at  $-\infty$ . Moreover, arguing as in the proof of Claim 7.2 we see that  $\mathcal{M}^\infty$  has a dominant unstable mode, with the same expansion parameters  $a_1, a_2$  as  $\mathcal{M}$ .

On the other hand, by the choice of  $x_{h_i}^+$ , we have

$$(207) \quad \frac{x_{h_i}^+}{\|x_{h_i}^+\|} \rightarrow w^+ \in \partial \check{K},$$

with  $\langle w^+, e_2 \rangle \geq 0$ . Thus,  $\mathcal{M}^\infty$  splits off a line in the direction  $w^+$ . Therefore, by [6] the limit  $\mathcal{M}^\infty$  is  $\mathbb{R}$  times a two-dimensional bowl, where the  $\mathbb{R}$ -factor is in the direction  $w^+$ , and where the translation direction  $v^+$  is the orthogonal complement of  $w^+$  in  $\mathbb{R}^2 \times \{0\}$  with

$$(208) \quad \langle v^+, e_2 \rangle < 0.$$

As the expansion parameters of  $\mathcal{M}^\infty$  are also  $a_1, a_2$  we see by observation (or by the fine-neck theorem from [10]) that

$$(209) \quad \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \gamma v^+ \quad \text{for some } \gamma > 0.$$

Combining this with (208), we get that  $a_2 < 0$ .

Arguing similarly using  $x_{h_i}^-$  gives that  $a_2 > 0$ , a contradiction. This completes the proof of Theorem 7.1.  $\square$

### Appendix A Local $L^\infty$ -estimate

In this appendix, we consider the renormalized mean curvature flow given (in some ball) as a graph over the cylinder  $\Gamma = \mathbb{R}^{n-d} \times S^d(\sqrt{2d})$ , namely our variables are  $(y, \sqrt{2d} \omega) \in \Gamma$  where  $y \in \mathbb{R}^{n-d}$  and  $\omega \in S^d$ . Let

$$(210) \quad \mathcal{L}u = \Delta_\Gamma u - \frac{1}{2}x^{\text{tau}} \cdot \nabla_\Gamma u + u = \rho^{-1} \operatorname{div}(\rho \nabla u) + \frac{1}{2d} \Delta_{S^d} u + u,$$

where  $\rho(y)$  is the Gaussian density given by

$$(211) \quad \rho(y) = (4\pi)^{-(d+1)/2} e^{-d/2} e^{-|y|^2/4}.$$

Given  $R \gg 1$ , we let  $\Gamma_R = \{(y, \omega) \in \Gamma \mid |y| \leq R\}$  and  $Q(R) = \Gamma_R \times [-R^2, 0]$ . We consider smooth solutions  $u: Q(2R) \rightarrow \mathbb{R}$  to the equation

$$(212) \quad u_\tau = \mathcal{L}u + E.$$

**Theorem A.1** (cf [25, Theorem 6.17]) *Suppose that for some constants  $C_0, k < \infty$ , the error  $E$  satisfies*

$$(213) \quad |E| \leq C_0(|u| + |\nabla u|) + k.$$

Then

$$(214) \quad \sup_{Q(R)} |u| \leq C \left[ k + \left( \int_{Q(2R)} u^2 \, d\operatorname{vol}_\Gamma \, d\tau \right)^{\frac{1}{2}} \right], \quad \text{where } C = C(C_0, R, n) < \infty.$$

**Proof** Instead of  $u$ , we consider the function  $\bar{u} = \rho u$  which solves

$$(215) \quad \bar{u}_\tau = \operatorname{div}(\rho \nabla_\Gamma(\rho^{-1} \bar{u})) + \rho E.$$

Since this equation satisfies the conditions of [25, Theorem 6.17], given  $q > 2$  we can obtain the following inequality as the proof in Lieberman<sup>2</sup>

$$(216) \quad \int_\Omega \bar{u}^{q-2} |\nabla \bar{u}|^2 v^{\alpha q - n - 2} \xi^2 \, d\operatorname{vol}_\Gamma \, d\tau \leq \frac{C q^2}{R^2} \int_\Omega \bar{u}^q v^{\alpha q - n - 2} \, d\operatorname{vol}_\Gamma \, d\tau,$$

where

- $\Omega = Q(2R) \cap \{\bar{u} \geq kR\}$ ,
- $\xi = (1 - |x|^2/4R^2)_+ (1 + t/4R^2)_+$  is a cut-off function,

<sup>2</sup>The book states a stronger inequality, which is wrong, but easily correctable. Here, we provide the necessary modification of the argument for convenience of the reader.

- $v = (1 - kR/\bar{u})_+$  is in  $[0, 1)$ , and
- $\alpha = (n + 2)/2$ .

Thus,  $h = \bar{u}v^\alpha$  satisfies

$$(217) \quad \int_{\Omega} |\nabla h|^2 \leq Cq^4 R^{-2} \int_{\Omega} h^2 v^{-2}.$$

Hence, the Sobolev type inequality from [Lemma A.2](#) below yields

$$(218) \quad \begin{aligned} Cq^4 \int_{\Omega} h^2 v^{-2} &\geq \left( \int_{\Omega} |\nabla h|^2 + \int_{\Omega} |h|^2 \right) + \int_{\Omega} |h|^2 \\ &\geq \frac{1}{C} \left( \int_{\Omega} |\nabla h|^2 + |h|^2 \right)^{n/(n+2)} \left( \int_{\Omega} h^2 \right)^{2/(n+2)} \\ &\geq \frac{1}{C} \left( \int_{\Omega} h^{2(n+2)/n} \right)^{2/(n+2)}. \end{aligned}$$

Thus, setting

$$\kappa = (n+2)/n, \quad w = \bar{u}v^\alpha \quad \text{and} \quad d\mu = R\xi(1 - kR/\bar{u})_+^{n-2} d\text{vol}_{\Gamma} d\tau$$

gives

$$(219) \quad \left( \int_{\Omega} w^{\kappa q} d\mu \right)^{1/\kappa q} \leq C^{1/q} q^{4/q} \left( \int_{\Omega} w^q d\mu \right)^{1/q}.$$

Hence, iterating this process with  $q = 2\kappa^j$  for  $j \in \mathbb{N}$  yields the result. □

The following lemma has been used in lieu of [\[25, Theorem 6.9\]](#):

**Lemma A.2** *Suppose that  $u \in C^\infty(Q(R))$  is a nonnegative function satisfying  $u = 0$  on  $\partial\Gamma_R \times [-R^2, 0]$ . Then, there exists some constant  $C = C(n, R) < \infty$  such that*

$$(220) \quad \int_{Q(R)} u^{2(n+2)/n} \leq C \left( \int_{Q(R)} u^2 \right)^{2/n} \left( \int_{Q(R)} |\nabla u|^2 + u^2 \right).$$

**Proof** The Hölder inequality yields

$$(221) \quad \int_Q u^{2(n+2)/n} \leq \left( \int_Q u^2 \right)^{1/n} \left( \int_Q u^{2(n+1)/(n-1)} \right)^{(n-1)/n}.$$

Applying the Michael–Simon inequality this implies

$$(222) \quad \int_Q u^{2(n+2)/n} \leq C \left( \int_Q u^2 \right)^{1/n} \left( \int_Q u^{(n+2)/n} |Du| + u^{2(n+1)/n} \right)^{(n-1)/n}.$$

Using again Hölder’s inequality we obtain the desired result. □

## Appendix B Merle–Zaag ODE lemma

We recall the following variant of the Merle–Zaag ODE lemma [26, Lemma A.1], which has been proved in [12, Lemma B.1]:

**Lemma B.1**  $U_0, U_+, U_- : (-\infty, 0] \rightarrow \mathbb{R}$  are absolutely continuous nonnegative functions satisfying  $U_0 + U_+ + U_- > 0$  and

$$(223) \quad \liminf_{s \rightarrow -\infty} U_-(s) = 0.$$

Suppose that there exist some constant  $c_0 > 0$  and positive increasing function  $\sigma : (-\infty, 0] \rightarrow \mathbb{R}$  such that  $\lim_{s \rightarrow -\infty} \sigma(s) = 0$  and the following hold

$$(224) \quad |U'_0| \leq \sigma(U_0 + U_- + U_+),$$

$$(225) \quad U'_- \leq -c_0 U_- + \sigma(U_0 + U_+),$$

$$(226) \quad U'_+ \geq c_0 U_+ - \sigma(U_0 + U_-).$$

Then there exists  $c = c(c_0) > 0$  and  $\delta = \delta(c_0) > 0$  such that if  $\sigma(s_0) < \delta$  then

$$(227) \quad U_- \leq c\sigma(U_0 + U_+) \text{ on } (-\infty, s_0],$$

and either

$$(228) \quad U_+ \leq o(U_0),$$

or

$$(229) \quad U_0 \leq c\sigma U_+$$

for all  $s \leq s_0$ .

## References

- [1] S J Altschuler, L F Wu, *Translating surfaces of the non-parametric mean curvature flow with prescribed contact angle*, Calc. Var. Partial Differential Equations 2 (1994) 101–111 [MR](#) [Zbl](#)
- [2] B Andrews, *Noncollapsing in mean-convex mean curvature flow*, Geom. Topol. 16 (2012) 1413–1418 [MR](#) [Zbl](#)
- [3] S Angenent, P Daskalopoulos, N Sesum, *Unique asymptotics of ancient convex mean curvature flow solutions*, J. Differential Geom. 111 (2019) 381–455 [MR](#) [Zbl](#)
- [4] S Angenent, P Daskalopoulos, N Sesum, *Uniqueness of two-convex closed ancient solutions to the mean curvature flow*, Ann. of Math. 192 (2020) 353–436 [MR](#) [Zbl](#)
- [5] S Brendle, *A sharp bound for the inscribed radius under mean curvature flow*, Invent. Math. 202 (2015) 217–237 [MR](#) [Zbl](#)

- [6] **S Brendle, K Choi**, *Uniqueness of convex ancient solutions to mean curvature flow in  $\mathbb{R}^3$* , *Invent. Math.* 217 (2019) 35–76 [MR](#) [Zbl](#)
- [7] **S Brendle, K Choi**, *Uniqueness of convex ancient solutions to mean curvature flow in higher dimensions*, *Geom. Topol.* 25 (2021) 2195–2234 [MR](#) [Zbl](#)
- [8] **S Brendle, G Huisken**, *Mean curvature flow with surgery of mean convex surfaces in  $\mathbb{R}^3$* , *Invent. Math.* 203 (2016) 615–654 [MR](#) [Zbl](#)
- [9] **J Cheeger, R Haslhofer, A Naber**, *Quantitative stratification and the regularity of mean curvature flow*, *Geom. Funct. Anal.* 23 (2013) 828–847 [MR](#) [Zbl](#)
- [10] **K Choi, R Haslhofer, O Hershkovits**, *Ancient low-entropy flows, mean-convex neighborhoods, and uniqueness*, *Acta Math.* 228 (2022) 217–301 [MR](#) [Zbl](#)
- [11] **K Choi, R Haslhofer, O Hershkovits, B White**, *Ancient asymptotically cylindrical flows and applications*, *Invent. Math.* 229 (2022) 139–241 [MR](#) [Zbl](#)
- [12] **K Choi, C Mantoulidis**, *Ancient gradient flows of elliptic functionals and Morse index*, *Amer. J. Math.* 144 (2022) 541–573 [MR](#) [Zbl](#)
- [13] **T H Colding, W P Minicozzi, II**, *Generic mean curvature flow, I: Generic singularities*, *Ann. of Math.* 175 (2012) 755–833 [MR](#) [Zbl](#)
- [14] **T H Colding, W P Minicozzi, II**, *Uniqueness of blowups and Łojasiewicz inequalities*, *Ann. of Math.* 182 (2015) 221–285 [MR](#) [Zbl](#)
- [15] **T H Colding, W P Minicozzi, II**, *The singular set of mean curvature flow with generic singularities*, *Invent. Math.* 204 (2016) 443–471 [MR](#) [Zbl](#)
- [16] **R Haslhofer, B Kleiner**, *On Brendle’s estimate for the inscribed radius under mean curvature flow*, *Int. Math. Res. Not.* 2015 (2015) 6558–6561 [MR](#) [Zbl](#)
- [17] **R Haslhofer, B Kleiner**, *Mean curvature flow of mean convex hypersurfaces*, *Comm. Pure Appl. Math.* 70 (2017) 511–546 [MR](#) [Zbl](#)
- [18] **R Haslhofer, B Kleiner**, *Mean curvature flow with surgery*, *Duke Math. J.* 166 (2017) 1591–1626 [MR](#) [Zbl](#)
- [19] **D Hoffman, T Ilmanen, F Martín, B White**, *Graphical translators for mean curvature flow*, *Calc. Var. Partial Differential Equations* 58 (2019) art. id. 117 [MR](#) [Zbl](#)
- [20] **G Huisken**, *Asymptotic behavior for singularities of the mean curvature flow*, *J. Differential Geom.* 31 (1990) 285–299 [MR](#) [Zbl](#)
- [21] **G Huisken**, *Local and global behaviour of hypersurfaces moving by mean curvature*, from “Differential geometry: partial differential equations on manifolds” (R Greene, S-T Yau, editors), *Proc. Sympos. Pure Math.* 54.1, Amer. Math. Soc., Providence, RI (1993) 175–191 [MR](#) [Zbl](#)
- [22] **G Huisken, C Sinestrari**, *Mean curvature flow with surgeries of two-convex hypersurfaces*, *Invent. Math.* 175 (2009) 137–221 [MR](#) [Zbl](#)
- [23] **T Ilmanen**, *Singularities of mean curvature flow of surfaces*, preprint (1995) Available at <https://people.math.ethz.ch/~ilmanen/papers/sing.ps>
- [24] **S Kleene, NM Møller**, *Self-shrinkers with a rotational symmetry*, *Trans. Amer. Math. Soc.* 366 (2014) 3943–3963 [MR](#) [Zbl](#)
- [25] **G M Lieberman**, *Second order parabolic differential equations*, World Sci., River Edge, NJ (1996) [MR](#) [Zbl](#)



- [26] **F Merle, H Zaag**, *Optimal estimates for blowup rate and behavior for nonlinear heat equations*, Comm. Pure Appl. Math. 51 (1998) 139–196 [MR](#) [Zbl](#)
- [27] **W Sheng, X-J Wang**, *Singularity profile in the mean curvature flow*, Methods Appl. Anal. 16 (2009) 139–155 [MR](#) [Zbl](#)
- [28] **J Spruck, L Xiao**, *Complete translating solitons to the mean curvature flow in  $\mathbb{R}^3$  with nonnegative mean curvature*, Amer. J. Math. 142 (2020) 993–1015 [MR](#) [Zbl](#)
- [29] **X-J Wang**, *Convex solutions to the mean curvature flow*, Ann. of Math. 173 (2011) 1185–1239 [MR](#) [Zbl](#)
- [30] **B White**, *The size of the singular set in mean curvature flow of mean-convex sets*, J. Amer. Math. Soc. 13 (2000) 665–695 [MR](#) [Zbl](#)
- [31] **B White**, *The nature of singularities in mean curvature flow of mean-convex sets*, J. Amer. Math. Soc. 16 (2003) 123–138 [MR](#) [Zbl](#)

*School of Mathematics, Korea Institute for Advanced Study  
Seoul, South Korea*

*Department of Mathematics, University of Toronto  
Toronto, ON, Canada*

*Institute of Mathematics, Hebrew University  
Jerusalem, Israel*

[choiks@kias.re.kr](mailto:choiks@kias.re.kr), [roberth@math.toronto.edu](mailto:roberth@math.toronto.edu), [or.hershkovits@mail.huji.ac.il](mailto:or.hershkovits@mail.huji.ac.il)

Proposed: Tobias H Colding

Seconded: David Fisher, Bruce Kleiner

Received: 11 August 2021

Revised: 2 May 2023

# GEOMETRY & TOPOLOGY

[msp.org/gt](http://msp.org/gt)

## MANAGING EDITOR

András I Stipsicz    Alfréd Rényi Institute of Mathematics  
[stipsicz@renyi.hu](mailto:stipsicz@renyi.hu)

## BOARD OF EDITORS

Mohammed Abouzaid	Stanford University <a href="mailto:abouzaid@stanford.edu">abouzaid@stanford.edu</a>	Mark Gross	University of Cambridge <a href="mailto:mgross@dpms.cam.ac.uk">mgross@dpms.cam.ac.uk</a>
Dan Abramovich	Brown University <a href="mailto:dan_abramovich@brown.edu">dan_abramovich@brown.edu</a>	Rob Kirby	University of California, Berkeley <a href="mailto:kirby@math.berkeley.edu">kirby@math.berkeley.edu</a>
Ian Agol	University of California, Berkeley <a href="mailto:ianagol@math.berkeley.edu">ianagol@math.berkeley.edu</a>	Bruce Kleiner	NYU, Courant Institute <a href="mailto:bkleiner@cims.nyu.edu">bkleiner@cims.nyu.edu</a>
Arend Bayer	University of Edinburgh <a href="mailto:arend.bayer@ed.ac.uk">arend.bayer@ed.ac.uk</a>	Sándor Kovács	University of Washington <a href="mailto:skovacs@uw.edu">skovacs@uw.edu</a>
Mark Behrens	University of Notre Dame <a href="mailto:mbehrens1@nd.edu">mbehrens1@nd.edu</a>	Urs Lang	ETH Zürich <a href="mailto:urs.lang@math.ethz.ch">urs.lang@math.ethz.ch</a>
Mladen Bestvina	University of Utah <a href="mailto:bestvina@math.utah.edu">bestvina@math.utah.edu</a>	Marc Levine	Universität Duisburg-Essen <a href="mailto:marc.levine@uni-due.de">marc.levine@uni-due.de</a>
Martin R Bridson	University of Oxford <a href="mailto:bridson@maths.ox.ac.uk">bridson@maths.ox.ac.uk</a>	Ciprian Manolescu	University of California, Los Angeles <a href="mailto:cm@math.ucla.edu">cm@math.ucla.edu</a>
Jim Bryan	University of British Columbia <a href="mailto:jbryan@math.ubc.ca">jbryan@math.ubc.ca</a>	Haynes Miller	Massachusetts Institute of Technology <a href="mailto:hmr@math.mit.edu">hmr@math.mit.edu</a>
Dmitri Burago	Pennsylvania State University <a href="mailto:burago@math.psu.edu">burago@math.psu.edu</a>	Tomasz Mrowka	Massachusetts Institute of Technology <a href="mailto:mrowka@math.mit.edu">mrowka@math.mit.edu</a>
Tobias H Colding	Massachusetts Institute of Technology <a href="mailto:colding@math.mit.edu">colding@math.mit.edu</a>	Aaron Naber	Northwestern University <a href="mailto:anaber@math.northwestern.edu">anaber@math.northwestern.edu</a>
Simon Donaldson	Imperial College, London <a href="mailto:s.donaldson@ic.ac.uk">s.donaldson@ic.ac.uk</a>	Peter Ozsváth	Princeton University <a href="mailto:petero@math.princeton.edu">petero@math.princeton.edu</a>
Yasha Eliashberg	Stanford University <a href="mailto:eliash-gt@math.stanford.edu">eliash-gt@math.stanford.edu</a>	Leonid Polterovich	Tel Aviv University <a href="mailto:polterov@post.tau.ac.il">polterov@post.tau.ac.il</a>
Benson Farb	University of Chicago <a href="mailto:farb@math.uchicago.edu">farb@math.uchicago.edu</a>	Colin Rourke	University of Warwick <a href="mailto:gt@maths.warwick.ac.uk">gt@maths.warwick.ac.uk</a>
David M Fisher	Rice University <a href="mailto:davidfisher@rice.edu">davidfisher@rice.edu</a>	Roman Sauer	Karlsruhe Institute of Technology <a href="mailto:roman.sauer@kit.edu">roman.sauer@kit.edu</a>
Mike Freedman	Microsoft Research <a href="mailto:michaelf@microsoft.com">michaelf@microsoft.com</a>	Stefan Schwede	Universität Bonn <a href="mailto:schwede@math.uni-bonn.de">schwede@math.uni-bonn.de</a>
David Gabai	Princeton University <a href="mailto:gabai@princeton.edu">gabai@princeton.edu</a>	Natasa Sesum	Rutgers University <a href="mailto:natasas@math.rutgers.edu">natasas@math.rutgers.edu</a>
Stavros Garoufalidis	Southern U. of Sci. and Tech., China <a href="mailto:stavros@mpim-bonn.mpg.de">stavros@mpim-bonn.mpg.de</a>	Gang Tian	Massachusetts Institute of Technology <a href="mailto:tian@math.mit.edu">tian@math.mit.edu</a>
Cameron Gordon	University of Texas <a href="mailto:gordon@math.utexas.edu">gordon@math.utexas.edu</a>	Ulrike Tillmann	Oxford University <a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>
Jesper Grodal	University of Copenhagen <a href="mailto:jg@math.ku.dk">jg@math.ku.dk</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Misha Gromov	IHÉS and NYU, Courant Institute <a href="mailto:gromov@ihes.fr">gromov@ihes.fr</a>	Anna Wienhard	Universität Heidelberg <a href="mailto:wienhard@mathi.uni-heidelberg.de">wienhard@mathi.uni-heidelberg.de</a>

---

See inside back cover or [msp.org/gt](http://msp.org/gt) for submission instructions.

---

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

---

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# GEOMETRY & TOPOLOGY

Volume 28 Issue 7 (pages 3001–3510) 2024

---

Holomorphic 1-forms on the moduli space of curves	3001
FILIPPO FRANCESCO FAVALE, GIAN PIETRO PIROLA and SARA TORELLI	
Two-dimensional metric spaces with curvature bounded above, I	3023
KOICHI NAGANO, TAKASHI SHIOYA and TAKAO YAMAGUCHI	
A nonexistence result for wing-like mean curvature flows in $\mathbb{R}^4$	3095
KYEONGSU CHOI, ROBERT HASLHOFER and OR HERSHKOVITS	
Higgs bundles, harmonic maps and pleated surfaces	3135
ANDREAS OTT, JAN SWOBODA, RICHARD WENTWORTH and MICHAEL WOLF	
Multiple cover formulas for K3 geometries, wall-crossing, and Quot schemes	3221
GEORG OBERDIECK	
Ancient solutions to the Kähler Ricci flow	3257
YU LI	
CAT(0) 4-manifolds are Euclidean	3285
ALEXANDER LYTCHAK, KOICHI NAGANO and STEPHAN STADLER	
Gromov–Witten theory via roots and logarithms	3309
LUCA BATTISTELLA, NAVID NABIJOU and DHARUV RANGANATHAN	
3-Manifolds without any embedding in symplectic 4-manifolds	3357
ALIAKBAR DAEMI, TYE LIDMAN and MIKE MILLER EISMEIER	
Orbit closures of unipotent flows for hyperbolic manifolds with Fuchsian ends	3373
MINJU LEE and HEE OH	
When does the zero fiber of the moment map have rational singularities?	3475
HANS-CHRISTIAN HERBIG, GERALD W SCHWARZ and CHRISTOPHER SEATON	