



Geometry & Topology

Volume 28 (2024)

CAT(0) 4-manifolds are Euclidean

ALEXANDER LYTCHAK

KOICHI NAGANO

STEPHAN STADLER



CAT(0) 4–manifolds are Euclidean

ALEXANDER LYTCHAK
KOICHI NAGANO
STEPHAN STADLER

We prove that a topological 4–manifold of globally nonpositive curvature is homeomorphic to Euclidean space.

51H20, 51K10, 53C23, 54F65, 57N16

1 Introduction

1.1 Main result

This paper concerns the topology of CAT(0) *manifolds*. These are synthetic generalizations of complete simply connected Riemannian manifolds of nonpositive sectional curvature. By the classical theorem of Cartan and Hadamard, any such Riemannian manifold is diffeomorphic to the Euclidean space \mathbb{R}^n . In his seminal paper, Gromov [1981], asked if there exist simply connected topological manifolds other than Euclidean space which admit a metric of nonpositive curvature in a synthetic sense. (See Section 1.6 for further discussion.) The most important synthetic notion of nonpositive curvature is the one due to Alexandrov. The corresponding spaces were named CAT(0) by Gromov. Any CAT(0) space is contractible, thus any CAT(0) 2–manifold is homeomorphic to \mathbb{R}^2 by the classification of surfaces. In dimensions strictly greater than two, contractible manifolds are Euclidean precisely when they are simply connected at infinity, thanks to classical topological results [Freedman 1982; Husch and Price 1970; Stallings 1962]. In dimension three, CAT(0) manifolds are indeed Euclidean [Brown 1961; Rolfsen 1968; Thurston 1996b]. In dimensions strictly greater than four, Davis and Januszkiewicz [1991] constructed examples of non-Euclidean CAT(0) manifolds. We deal with the remaining open case:

Theorem 1.1 *Let X be a CAT(0) space which is topologically a 4–dimensional manifold. Then X is homeomorphic to \mathbb{R}^4 .*

This theorem also answers the first question in [Davis et al. 2012, Section 6].

1.2 Related statements: the simplicial case

Examples of Davis and Januszkiewicz [1991] mentioned above are simplicial complexes with piecewise Euclidean metrics. On the other hand, Stone [1976] verified that a *PL* n –manifold which is CAT(0) with

respect to a piecewise Euclidean metric is homeomorphic to the Euclidean space \mathbb{R}^n . By the resolution of the Poincaré conjecture, any simplicial complex homeomorphic to a 4–manifold is a PL–manifold. Consequently, any CAT(0) 4–manifold with a piecewise Euclidean metric is homeomorphic to \mathbb{R}^4 .

In higher dimensions there exist a large supply of exotic CAT(0) manifolds. Ancel and Guilbault [1997] have shown that the interior of every compact contractible PL n –manifold (for $n \geq 5$) supports a complete geodesic metric of strictly negative curvature. They also point out that their result continues to hold without the PL assumption in dimensions strictly larger than 5. The question as to which manifolds carry a piecewise Euclidean CAT(0) metric has been further investigated in [Adiprasito and Benedetti 2020; Adiprasito and Funar 2015]. Motivated by Gromov’s question, it is natural to ask the following; compare [Adiprasito and Funar 2015, Question 3]:

Question 1.2 Are there CAT(0) topological manifolds which do not carry a piecewise Euclidean CAT(0) metric?

Question 1.3 What are necessary and sufficient conditions for the existence of a CAT(0) metric on a contractible manifold?

1.3 Related statements: the cocompact case

Gromov’s question has been thoroughly studied in the *cocompact* setting. (Recall that universal coverings of locally CAT(0) spaces are CAT(0) [Alexander and Bishop 1990]). By [Davis and Januszkiewicz 1991, Theorem 5b.1], in all dimensions strictly greater than four, there exist compact locally CAT(0) manifolds whose universal coverings are not Euclidean; see also [Ancel et al. 1997]. Moreover, in such dimensions there exist compact locally CAT(0) manifolds which have no PL structure at all [Davis and Januszkiewicz 1991, Section 5a]. We refer to [Davis et al. 2012, Section 3] for an overview of further similar results.

In dimension four, several classes of smoothable compact topological manifolds carrying a locally CAT(0) metric, yet not admitting a smooth metric of nonpositive curvature, have been constructed in [Davis et al. 2012; Sathaye 2017; Stadler 2015]. In these examples, the universal covering is homeomorphic to \mathbb{R}^4 (as follows from our main theorem), and identifying the obstructions to the existence of smooth metrics relies on an intricate analysis of the group actions involved.

1.4 Distance spheres

Our proof depends upon an important contribution by Thurston [1996b]. He showed that if all distance spheres to some fixed point $o \in X$ of a 4–dimensional CAT(0) manifold X are topological 3–manifolds, then X is homeomorphic to \mathbb{R}^4 . Using a finer analysis of the metric structure of the space, we verify this condition in the more general setting of *homology 4–manifolds*; see Sections 1.5 and 3.4 for the relevant definition and properties.

Theorem 1.4 *Let X be a CAT(0) space which is a homology 4–manifold. Let $o \in X$ and $R > 0$ be arbitrary. Then the distance sphere $S_R(o)$ of radius R around o is a topological 3–manifold.*

We remark that this result does not hold true in dimensions $n \geq 5$, even for piecewise Euclidean topological n -manifolds. Indeed, this can be seen in the aforementioned examples of Davis and Januszkiewicz; compare [Davis and Januszkiewicz 1991, Proposition 3d.3].

If X is a CAT(0) 4-manifold (and not just a homology 4-manifold), then the resolution of the Poincaré conjecture together with [Thurston 1996b] implies that all distance spheres are homeomorphic to \mathbb{S}^3 . Moreover, the homeomorphism in Theorem 1.1 is not completely abstract, but rather has the following geometric feature:

Corollary 1.5 *Let X be a 4-dimensional CAT(0) manifold and let $o \in X$ be an arbitrary point. Then the distance function $d_o: X \setminus \{o\} \rightarrow (0, \infty)$ is a trivial fiber bundle with fiber \mathbb{S}^3 .*

On the other hand, for a general CAT(0) homology 4-manifold, the topology of the distance spheres may depend on the radius, despite the fact that all of the spheres involved are manifolds. This can already be seen in the Euclidean cone $X = C(\Sigma)$ over the Poincaré homology sphere Σ . The fine topological analysis of [Thurston 1996b], using the fact that the ambient space is a manifold, is therefore indispensable for the conclusion of our main theorem.

Corollary 1.5 extends to the *ideal boundary* $\partial_\infty X$ and the *natural compactification* $\bar{X} := X \cup \partial_\infty X$ of X ; see [Bridson and Haefliger 1999, Section II.8] for the definition and properties of ideal boundaries. In dimensions $n \geq 5$, there are CAT(0) spaces homeomorphic to \mathbb{R}^n with ideal boundary different from \mathbb{S}^{n-1} [Davis and Januszkiewicz 1991, Theorem 5c.1]. In dimension four we show:

Corollary 1.6 *Let X be a 4-dimensional topological manifold with a CAT(0) metric. Then the ideal boundary $\partial_\infty X$ of X is homeomorphic to \mathbb{S}^3 and the canonical compactification $\bar{X} = X \cup \partial_\infty X$ is homeomorphic to the closed unit ball in \mathbb{R}^4 .*

1.5 Related statements: homology manifolds

A homology n -manifold (without boundary) is a locally compact metric space X of finite topological dimension such that, for all $x \in X$, the local homology $H_*(X, X \setminus \{x\})$ equals $H_*(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\})$. The structure theory of homology manifolds has been a central topic in geometric topology for many decades and is of fundamental importance in topological *manifold recognition* [Cannon 1978; Cavicchioli et al. 2016; Repovš 1994].

A homology n -manifold for $n \leq 2$ is always a topological n -manifold by a theorem of Moore [Wilder 1949, Chapter IX]. On the other hand, in dimensions $n \geq 3$, a homology n -manifold may not have any manifold points at all [Daverman and Walsh 1981]. While there are deep and rather robust results allowing one to recognize when a homology manifold is a manifold in dimensions five and up, much less is known in dimensions three and four [Repovš 1994]. Even though the tools of algebraic topology allow us to recognize homology manifolds in many instances, in particular, in the situation of Theorem 1.4 (in all dimensions), passing from homology manifolds to topological manifolds is difficult and requires some geometric insight.

In the situation of Theorem 1.4, we achieve the needed local control of the topology of large spheres by slicing them and verifying that slices are 2-dimensional spheres. Subsequently, these slices can be controlled uniformly with the help of Jordan's curve theorem. The control of the slices allows us to recover the local topology.

In contrast to the situation for general homology n -manifolds, CAT(0) homology n -manifolds are not too far from being manifolds. More precisely, a CAT(0) homology n -manifold is a topological n -manifold on the complement of a discrete subset [Lytchak and Nagano 2022, Theorem 1.2]; see [Wu 1997] for corresponding statements on spaces with lower curvature bounds.

We mention in passing a question of Busemann [1955; Berestovsky et al. 2011], which is related in spirit to the origins of this paper:

Question 1.7 Let X be a locally compact geodesic metric space. Assume that X is geodesically complete and that there are no branching geodesics. Does X have finite dimension? Is any such finite-dimensional X a topological manifold?

If such a space X has finite dimension n then X is a homology n -manifold, and if $n \leq 4$ then X is a manifold [Busemann 1955; Krakus 1968; Thurston 1996a; Berestovsky et al. 2011]. For $n \geq 5$, the question remains open. Finally, we mention that an answer to Busemann's question would follow from a purely topological conjecture of Bing and Borsuk [Halverson and Repovš 2008].

1.6 Minor generalizations

An application of [Lytchak and Stadler 2020, Theorem 1.1] extends Theorem 1.1 and Corollary 1.5 to other curvature bounds:

Corollary 1.8 Let X be a 4-dimensional topological manifold which is a CAT(κ) space. Let $R > 0$ be a real number with $R < \pi/(2\sqrt{\kappa})$ if $\kappa > 0$. Then, for any $o \in X$, the open ball $B_R(o)$, the closed ball $\bar{B}_R(o)$ and the distance sphere $S_R(o)$ are homeomorphic to the open unit ball in \mathbb{R}^4 , to the closed unit ball in \mathbb{R}^4 and to \mathbb{S}^3 , respectively.

There are several notions of nonpositive curvature for metric spaces. In all works cited above, Gromov's question has been studied for CAT(0) spaces, even though the original question was posed for Busemann convex spaces, that is, geodesic spaces with a convex distance function. Any CAT(0) space is Busemann convex. Conversely, examples of Busemann convex spaces that are neither CAT(0) nor normed spaces are extremely rare; compare [Ivanov and Lytchak 2019]. Our ideas will apply to this more general setting, once some structural results developed in [Lytchak and Nagano 2019] for CAT(0) spaces are generalized to Busemann convex spaces.




Figure 1

1.7 Comment on strategy and technique

Our proof relies on the structural theory of geodesically complete spaces with upper curvature bounds developed in [Lytchak and Nagano 2019; 2022]. Since any CAT(0) manifold is geodesically complete, this theory applies to the present situation. So-called “strainer maps”, first appearing in [Burago et al. 1992] and defined by distance functions to points, are particularly useful. It has been verified [Lytchak and Nagano 2022] that for any point $o \in X$ as in Theorem 1.4, all sufficiently small distance spheres around o are (pairwise homeomorphic) 3-manifolds. In order to get sufficient control of remote spheres, an extension of the theory of strainer maps by one additional “orthogonal but nonstraining” coordinate is required. This extension may be useful beyond the present work. We refer the reader to Sections 5 and 7 for more details. Here, we only formulate a special case of Proposition 5.5, essential for the proof of Theorem 1.4:

Theorem 1.9 *Let X be a locally compact geodesically complete CAT(0) space, let $o \in X$ and let $p \in X$ be at distance $R > 0$ from o . Then there exists $\epsilon > 0$ such that for all $0 < s < \epsilon$, the intersection of the distance sphere $S_s(p)$ with the closed ball $\bar{B}_R(o)$ is contractible.*

Acknowledgments Lytchak and Stadler were supported by DFG grant SPP 2026. Nagano is supported by JSPS KAKENHI grant 20K03603. The authors are grateful to Ronan Conlon for useful comments and to Tadeusz Januskiewicz for helpful suggestions including Corollary 1.6. The authors would also like to thank the referee for valuable comments.

2 Preliminaries

2.1 Metric spaces

We refer the reader to [Alexander et al. 2024; Burago et al. 2001; Bridson and Haefliger 1999] for general background. We denote by d the distance in a metric space X . For $x \in X$, we denote by d_x the distance function $d_x(\cdot) = d(x, \cdot)$. For $x \in X$ and $r > 0$, we denote by $B_r(x)$ and $\bar{B}_r(x)$ the open and closed r -balls around x , respectively. Similarly, $B_r(A)$ denotes the open r -neighborhood of a subset $A \subset X$.

Moreover, $S_r(x)$ denotes the r -sphere around x and by $\dot{B}_r(x)$ we denote the punctured r -ball $B_r(x) \setminus \{x\}$. For $\lambda > 0$, we denote by $\lambda \cdot X$ the metric space resulting from X by rescaling the metric by λ . A *geodesic* is an isometric embedding of an interval. A *triangle* is a union of three geodesics connecting three points. X is a *geodesic metric space* if any pair of points of X is connected by a geodesic. It is *geodesically complete* if every isometric embedding of an interval extends to a locally isometric embedding of \mathbb{R} .

A map $F: X \rightarrow Y$ between metric spaces is called *L -Lipschitz* if $d(F(x), F(\bar{x})) \leq Ld(x, \bar{x})$, for all $x, \bar{x} \in X$.

The map F is called *L -open* if, for any $x \in X$ and any $r > 0$ such that the closed ball $\bar{B}_{Lr}(x)$ is complete, $B_r(F(x)) \subset F(B_{Lr}(x))$.

An ANR will denote an *absolute neighborhood retract*. For finite-dimensional metric spaces, the only case relevant here, being an ANR is equivalent to being locally contractible [Hu 1965].

2.2 Spaces with an upper curvature bound

For $\kappa \in \mathbb{R}$, let $R_\kappa \in (0, \infty]$ be the diameter of the complete simply connected surface M_κ^2 of constant curvature κ . A complete metric space is called a *CAT(κ) space* if any pair of its points with distance $< R_\kappa$ is connected by a geodesic and if all triangles with perimeter $< 2R_\kappa$ are not thicker than the *comparison triangle* in M_κ^2 . A metric space is called a *space with curvature bounded above by κ* if any point has a CAT(κ) neighborhood. We refer to [Alexander et al. 2024; Burago et al. 2001; Bridson and Haefliger 1999] for basic facts about such spaces.

For any CAT(κ) space X , the angle between each pair of geodesics starting at the same point is well defined. The *space of directions* $\Sigma_x X$ at $x \in X$, equipped with the angle metric, is a CAT(1) space. The Euclidean cone over Σ_x is a CAT(0) space. It is denoted by $T_x X$ and called the *tangent space* at x of X . Its tip will be denoted by o_x .

Let x, y and z be three points at pairwise distance $< R_\kappa$ in a CAT(κ) space X . Whenever $x \neq y$, the geodesic between x and y is unique and will be denoted by xy . For $y, z \neq x$, the angle at x between xy and xz will be denoted by $\angle yxz$.

In a CAT(κ) space X , all balls of radius smaller $\frac{1}{2}R_\kappa$ are convex, and hence X is locally contractible. In fact, X is an ANR [Kramer 2011, Theorem 3.2].

3 Geometric topology

3.1 Homology manifolds

Denote by D^n the closed unit ball in \mathbb{R}^n .

Let M be a locally compact separable metric space of finite topological dimension. We say that M is a *homology n -manifold with boundary* if for any $p \in M$ we have a point $x \in D^n$ such that the local homology $H_*(M, M \setminus \{p\})$ at p is isomorphic to $H_*(D^n, D^n \setminus \{x\})$. The *boundary* ∂M of M is defined

as the set of all points at which the n^{th} local homologies are trivial. In the case where the boundary of M is empty, we simply say that M is a *homology n -manifold*.

If M is a homology n -manifold with boundary then ∂M is a closed subset of M and it is a homology $(n-1)$ -manifold by [Mitchell 1990].

Any homology n -manifold (with boundary) has dimension n . For $n \leq 2$, we have the following theorem of Moore [Wilder 1949, Chapter IX]:

Theorem 3.1 *Any homology n -manifold with $n \leq 2$ is a topological manifold.*

A *homology n -sphere* is a homology n -manifold X with the homology of the n -sphere: $H_*(X) = H_*(S^n)$.

3.2 Uniform local contractibility

A function $\rho: [0, r_0) \rightarrow [0, \infty)$ is called a *contractibility function* if it is continuous at 0 with $\rho(0) = 0$ and $\rho(t) \geq t$ holds for all $t \in [0, r_0)$ [Petersen 1993].

Definition 3.2 We say that a family \mathcal{F} of metric spaces is *uniformly locally contractible* if there exists a contractibility function $\rho: [0, r_0) \rightarrow [0, \infty)$ such that, for any space X in the family \mathcal{F} , any point $x \in X$ and any $0 < r < r_0$, the ball $B_r(x)$ is contractible within the ball $B_{\rho(r)}(x)$.

For example, the family of all CAT(κ) spaces is uniformly locally contractible with $\rho: [0, \pi/\sqrt{\kappa}) \rightarrow [0, \infty)$ being the identity map.

Here is a special case of [Petersen 1990, Theorem A; 1993, Theorem 9]:

Theorem 3.3 *For any natural number n and any family \mathcal{F} of uniformly locally contractible metric spaces of dimension at most n , there exists some $\delta > 0$ such that any pair of spaces $X, Y \in \mathcal{F}$ with Gromov–Hausdorff distance at most δ is homotopy equivalent.*

The homotopy equivalences and the corresponding homotopies in Theorem 3.3 can be chosen arbitrarily close to the identity [Petersen 1993].

When dealing with a family of fibers of some map, we will use the following more convenient variant of Definition 3.2 [Ungar 1969]:

Definition 3.4 Let $F: X \rightarrow Y$ be a map between metric spaces. We say that F has *uniformly locally contractible fibers* if the following condition holds true for any point $x \in X$ and every neighborhood U of x in X : there exists a neighborhood $V \subset U$ of x in X such that for any fiber $F^{-1}(y)$ with nonempty intersection $F^{-1}(y) \cap V$, this intersection is contractible in $F^{-1}(y) \cap U$.

For X compact, a map $F: X \rightarrow Y$ has uniformly locally contractible fibers in the sense of Definition 3.4 if and only if the family of the fibers is uniformly locally contractible in the sense of Definition 3.2.

3.3 Fibrations and fiber bundles

A map $F: X \rightarrow Y$ between metric spaces is called a *Hurewicz fibration* if it satisfies the homotopy lifting property with respect to all spaces [Hatcher 2002, Section 4.2; Ungar 1969].

The map F is called open if the images of open sets are open.

We will use the following result to recognize Hurewicz fibrations:

Theorem 3.5 [Ferry 1978, Theorem 2; Ungar 1969, Theorem 1] *Let X and Y be finite-dimensional, compact metric spaces and let Y be an ANR. Let $F: X \rightarrow Y$ be an open surjective map with uniformly locally contractible fibers. Then X is an ANR and F is a Hurewicz fibration.*

In some situations, Hurewicz fibrations turn out to be fiber bundles. We will rely on the following:

Theorem 3.6 [Ferry 1991, Theorems 1.1–1.4; Raymond 1965, Theorem 2] *Let X and Y be finite-dimensional locally compact ANRs. Let $F: X \rightarrow Y$ be a Hurewicz fibration. If all fibers of F are closed n -manifolds then F is a locally trivial fiber bundle.*

3.4 CAT(0) (homology) manifolds

Following [Lytchak and Nagano 2019], we will abbreviate a locally compact locally geodesically complete separable space with an upper curvature bound as GCBA. Here we are concerned with CAT(0) spaces which are homeomorphic to (homology) manifolds. We will call such spaces CAT(0) *homology manifolds* and CAT(0) *manifolds*, respectively. Every CAT(0) homology manifold is geodesically complete [Lytchak and Schroeder 2007, Theorem 1.5] and therefore GCBA. Hence, we can rely on the results from [Lytchak and Nagano 2019; 2022]. For the local arguments of [Lytchak and Nagano 2019; 2022], the notion of a *tiny ball* played a role. We point out that in a CAT(0) homology manifold, a tiny ball is any ball of radius at most 1. After rescaling, the bound of 1 becomes irrelevant.

From [Lytchak and Nagano 2022, Lemma 3.1, Corollary 3.4 and Theorem 6.4] we infer:

Proposition 3.7 *Let X be a CAT(0) homology n -manifold. Then any space of directions $\Sigma_x X$ is a homology $(n-1)$ -sphere. If $n \leq 4$, then $\Sigma_x X$ is a topological manifold.*

Any CAT(0) homology n -manifold is a topological n -manifold on the complement of a discrete subset [Lytchak and Nagano 2022, Theorem 1.2]. For $n \leq 3$, a CAT(0) homology n -manifold is a manifold homeomorphic to \mathbb{R}^n [Lytchak and Nagano 2022, Theorem 6.4; Thurston 1996b]. The Euclidean cone over the Poincaré sphere is a CAT(0) homology 4-manifold which is not a manifold.

Any CAT(0) homology n -manifold is locally bi-Lipschitz equivalent to \mathbb{R}^n away from a closed set of Hausdorff dimension at most $n - 2$, as follows from [Lytchak and Nagano 2019, Theorem 1.2 and Section 10.2].

4 Strainer maps

We recall the definition and basic properties of strainer maps in the framework of CAT(0) spaces from [Lytchak and Nagano 2019; 2022]. Originally, strainer maps were introduced in [Burago et al. 1992] to study Alexandrov spaces with curvature bounded below.

Throughout this section X will denote a locally compact and geodesically complete CAT(0) space.

4.1 Almost spherical directions

Let v be a direction at a point $x \in X$. An *antipode* of v is a direction $\hat{v} \in \Sigma_x X$ at distance at least π from v . If v has a unique antipode \hat{v} , then $\Sigma_x X$ splits isometrically as a spherical join $\Sigma_x X \cong \{v, \hat{v}\} * Z$. More generally, the subset Σ^0 of points with unique antipodes in $\Sigma_x X$ is isometric to \mathbb{S}^k , for some k , and Σ^0 is a spherical join factor of $\Sigma_x X$ [Lytchak 2005, Corollary 4.4].

A quantitative version is provided by the notion of δ -spherical points and tuples. The direction $v \in \Sigma_x X$ is called δ -spherical if there exists some $\bar{v} \in \Sigma_x X$ such that for any $w \in \Sigma_x X$,

$$d(v, w) + d(w, \bar{v}) < \pi + \delta.$$

Moreover, we say that v and \bar{v} are *opposite* δ -spherical points. A δ -spherical direction v has a set of antipodes of diameter at most 2δ [Lytchak and Nagano 2019, Lemma 6.3]. Therefore, if δ is small, this “almost leads to a splitting” of $\Sigma_x X$ [loc. cit., Proposition 6.6].

A k -tuple (v_1, \dots, v_k) of points in $\Sigma_x X$ is called δ -spherical if there exists another k -tuple (\bar{v}_i) in $\Sigma_x X$ with the following two properties:

- For $1 \leq i \leq k$, the directions v_i and \bar{v}_i are opposite δ -spherical.
- For $1 \leq i \neq j \leq k$, the distances $d(v_i, \bar{v}_j)$, $d(v_i, v_j)$ and $d(\bar{v}_i, \bar{v}_j)$ are less than $\frac{1}{2}\pi + \delta$.

Moreover, (\bar{v}_i) and (v_i) are called *opposite* δ -spherical k -tuples.

4.2 Strainers and strainer maps

A k -tuple (p_1, \dots, p_k) is called a (k, δ) -strainer at a point $x \in X \setminus \{p_1, \dots, p_k\}$ if the starting directions $v_i \in \Sigma_x X$ of the geodesics $x p_i$ constitute a δ -spherical k -tuple in $\Sigma_x X$.

Two (k, δ) -strainers (p_i) and (q_i) at x are *opposite* if the corresponding δ -spherical k -tuples (v_i) and (w_i) are opposite in $\Sigma_x X$.

A k -tuple (p_i) in X is a (k, δ) -strainer in $A \subset X \setminus \{p_1, \dots, p_k\}$ if (p_i) is a (k, δ) -strainer at every point $x \in A$.

The set of all points $U \subset X \setminus \{p_1, \dots, p_k\}$ at which (p_i) is a (k, δ) -strainer is open in X [loc. cit., Corollary 7.9].

Each k -tuple (p_i) yields a *distance map* $F: X \rightarrow \mathbb{R}^k$ via $F = (d_{p_1}, \dots, d_{p_k})$. If (p_i) is a (k, δ) -strainer on a subset $A \subset X$, then the associated distance map F is called a (k, δ) -strainer map on A .

4.3 Properties of strainer maps

For $\delta \leq 1/(4k)$ and $L = 2\sqrt{k}$, every (k, δ) -strainer map $F: U \rightarrow \mathbb{R}^k$ on an open subset $U \subset \mathbb{R}^k$ is L -open and L -Lipschitz [loc. cit., Lemma 8.2].

Two observations form the building blocks for straining maps. Recall the definition of a punctured distance ball $\dot{B}_r(x) := B_r(x) \setminus \{x\}$. First, for any $\delta > 0$ and any $x \in X$, the function $d_x: \dot{B}_r(x) \rightarrow (0, r)$ is a $(1, \delta)$ -strainer map if r is chosen small enough [loc. cit., Proposition 7.3]. Second, let $F: U \rightarrow \mathbb{R}^k$ be a (k, δ) -strainer map and let p be a point in a fiber Π of F . Then there exists $r > 0$ and a neighborhood W of $\dot{B}_r(p) \cap \Pi$ in U such that the map $\hat{F} = (F, d_p): W \rightarrow \mathbb{R}^{k+1}$ is a $(k+1, 4\delta)$ -strainer map [loc. cit., Proposition 9.4].

Any (k, δ) -strainer map on an open subset of a k -dimensional CAT(0) space X provides a bi-Lipschitz chart [loc. cit., Corollary 11.2]. In general, we have the following topological structure:

Theorem 4.1 [Lytchak and Nagano 2022, Theorem 5.1 and Corollary 5.2] *Let U be an open subset of a GCBA space X . Let $F: U \rightarrow \mathbb{R}^k$ be a (k, δ) -strainer map for some k and some $\delta < 1/(20k)$. Then any $x \in U$ has arbitrarily small open contractible neighborhoods V such that the restriction $F: V \rightarrow F(V)$ is a Hurewicz fibration with contractible fibers.*

If a fiber $F^{-1}(b)$ is compact, then there exists an open neighborhood V of $F^{-1}(b)$ in U such that $F: V \rightarrow F(V)$ is a Hurewicz fibration.

If U is a homology n -manifold, then any fiber $F^{-1}(b)$ is a homology $(n-k)$ -manifold.

5 Extended strainer maps

5.1 Definition and basic properties

Throughout this section, X will denote a locally compact and geodesically complete CAT(0) space.

Let (p_1, \dots, p_k) be a k -tuple in X and let $q \in X$ be an additional point. We say that (p_1, \dots, p_k, q) is an *extended (k, δ) -strainer* in a subset $A \subset X \setminus \{p_1, \dots, p_k, q\}$ if the following holds true for all $x \in A$:

The k -tuple (p_i) is a (k, δ) -strainer at x and any continuation qq' of the geodesic qx beyond x is such that, for all $1 \leq i \leq k$,

$$\angle qx p_i < \frac{1}{2}\pi + \delta \quad \text{and} \quad \angle q' x p_i < \frac{1}{2}\pi + \delta.$$

By the semicontinuity of angles, the set U of all points at which (p_1, \dots, p_k, q) is an extended (k, δ) -strainer is open in $X \setminus \{p_1, \dots, p_k, q\}$ [Lytchak and Nagano 2019, Section 3.3 and Corollary 7.9].

Let (p_1, \dots, p_k, q) be an extended (k, δ) -strainer in an open set $U \subset X$. Then we call the map

$$\hat{F} = (d_{p_1}, \dots, d_{p_k}, d_q) = (F, d_q): U \rightarrow (0, \infty)^{k+1}$$

an *extended (k, δ) -strainer map*.

By definition, an extended (k, δ) -strainer map $\widehat{F}: U \rightarrow \mathbb{R}^{k+1}$ is also an extended (k, δ') -strainer map for any $0 < \delta' < \delta$.

5.2 Basic properties

Let (p_1, \dots, p_k, q) be an extended (k, δ) -strainer at a point $x \in X$ and let qq' be an extension of the geodesic qx . Since $\angle qxq' = \pi$, for $1 \leq i \leq k$, we have

$$\angle p_i x q > \frac{1}{2}\pi - \delta \quad \text{and} \quad \angle p_i x q' > \frac{1}{2}\pi - \delta.$$

We fix an opposite (k, δ) -strainer (p'_1, \dots, p'_k) to (p_i) at the point x . The definition of opposite strainers implies

$$\angle p'_i x q < \frac{1}{2}\pi + 2\delta \quad \text{and} \quad \angle p'_i x q' < \frac{1}{2}\pi + 2\delta.$$

Therefore

$$\angle p'_i x q > \frac{1}{2}\pi - 2\delta \quad \text{and} \quad \angle p'_i x q' > \frac{1}{2}\pi - 2\delta.$$

Applying [loc. cit., Lemma 8.1] (compare [loc. cit., Lemma 8.2]) we get:

Lemma 5.1 For $\delta \leq 1/(20k)$ and $L = 2\sqrt{k+1}$, any extended (k, δ) -strainer map $\widehat{F}: U \rightarrow \mathbb{R}^{k+1}$ is L -Lipschitz and L -open.

Remark 5.2 The argument in [loc. cit., Lemma 8.3] allows us to choose the constant L above arbitrarily close to 1 if δ is sufficiently small.

By definition, any point q is an extended $(0, \delta)$ -strainer in $X \setminus \{q\}$ for any $\delta > 0$ and the distance function $d_q: X \setminus \{q\} \rightarrow (0, \infty)$ is an extended $(0, \delta)$ -strainer map. We are interested in distance spheres, and thus in fibers of such $(0, \delta)$ -strainer maps. As in [Burago et al. 1992; Lytchak and Nagano 2019], we approach the structure of these fibers by finding more strainers:

Lemma 5.3 Let $\widehat{F} = (d_{p_1}, \dots, d_{p_k}, d_q): U \rightarrow \mathbb{R}^{k+1}$ be an extended (k, δ) -strainer map for some $k \geq 0$ and $\delta < 1/(20k)$. Let $p \in U$ be arbitrary and let $\widehat{\Pi}_p := \widehat{F}^{-1}(\widehat{F}(p))$ be the fiber of \widehat{F} through p .

Then there exists $r > 0$ such that (p, p_1, \dots, p_k, q) is an extended $(k+1, 4\delta)$ -strainer in the intersection of $\widehat{\Pi}_p$ and the punctured ball $\dot{B}_r(p)$.

Proof We apply [Lytchak and Nagano 2019, Proposition 9.4] and find some $r > 0$ such that (p, p_1, \dots, p_k) is a $(k+1, 4\delta)$ -strainer in $\widehat{\Pi}_p \cap \dot{B}_r(p)$.

For any $x \in \widehat{\Pi}_p \cap \dot{B}_r(p)$, the points x and p are at equal distance to q , and hence $\angle pxq < \frac{1}{2}\pi$. It remains to prove that, for sufficiently small $r > 0$, $\angle pxq' < \frac{1}{2}\pi + \delta$, for all $x \in \widehat{\Pi}_p \cap \dot{B}_r(p)$ and all points q' with $\angle qxq' = \pi$. Suppose for the sake of contradiction that we find $x_i \in \widehat{\Pi}_p \setminus \{p\}$ converging to p and points q'_i lying on extensions of the geodesics qx_i such that $\angle px_i q'_i \geq \frac{1}{2}\pi + \delta$. We may assume that $d(x_i, q'_i) = 1$ and that the q'_i converge to a point q'_∞ . Using geodesic completeness, we extend $x_i p$ to a geodesic $x_i p_i$ of length 1 such that

$$\angle p_i p q'_\infty = \pi - \angle q'_\infty p x_i \leq \frac{1}{2}\pi.$$




Figure 2

Taking a subsequence, we may assume that p_i converges to a point p_∞ . Hence, $\angle p_\infty p q'_\infty \leq \frac{1}{2}\pi$. But, by semicontinuity of angles,

$$\angle p_\infty p q'_\infty \geq \limsup_{i \rightarrow \infty} \angle p_i x_i q'_i \geq \frac{1}{2}\pi + \delta.$$

This contradiction finishes the proof. □

5.3 Halfspaces

Let $\widehat{F} = (F, d_o)$ be an extended strainer map on an open set U . We denote the F -fiber and \widehat{F} -fiber through a point x by Π_x and $\widehat{\Pi}_x$, respectively. We define the \widehat{F} -halfspace through x by

$$\widehat{\Pi}_x^+ = \{y \in \Pi_x \mid d_o(y) \leq d_o(x)\}.$$

The proof of our main results will rely on the following structural results about the fibers and halfspaces of extended strainer maps. The proofs of these results are postponed to Section 7.

The first result generalizes [Thurston 1996b, Proposition 2.7; Lytchak and Nagano 2022, Corollary 5.2]:

Proposition 5.4 *Let U be an open subset of X . Then for any extended (k, δ) -strainer map $\widehat{F}: U \rightarrow \mathbb{R}^{k+1}$ with $\delta < 1/(64k)$, the following holds true for any $x \in U$:*

- (1) *The halfspace $\widehat{\Pi}_x^+$ is an ANR.*
- (2) *If U is a homology n -manifold, then $\widehat{\Pi}_x^+$ is a homology $(n-k)$ -manifold with boundary $\widehat{\Pi}_x$. The fiber $\widehat{\Pi}_x$ is a homology $(n-k-1)$ -manifold without boundary.*

The second statement is an extension of Theorem 1.9 which constitutes the special case where $k = 0$ and $y = x$. In the proof of our main results we will only need the cases $k = 0$ and $k = 1$.

Proposition 5.5 *For every relatively compact set $V \subset X$ there exists $\delta_0 > 0$ with the following property. Let $\widehat{F}: X \rightarrow \mathbb{R}^{k+1}$ be a distance map which is an extended (k, δ_0) -strainer map at a point $x \in V$. Then there exist $\epsilon_0, s_0 > 0$ such that, for any $0 < \epsilon < \epsilon_0$ and any point y with $d(x, y) < s_0\epsilon$, the “hemisphere” $S_\epsilon(x) \cap \widehat{\Pi}_y^+$ is contractible and locally contractible.*

6 Proof of the main theorem

6.1 Topology of intersecting spheres

Fix a CAT(0) homology 4-manifold X , a point $o \in X$ and some radius $R > 0$. We denote by S the distance sphere $S = S_R(o)$, and we are going to verify that S is a topological 3-manifold.

We fix an arbitrary $p \in S$ for the rest of the proof. We need to find a neighborhood of p in S which is homeomorphic to \mathbb{R}^3 . For this we aim to show that the restriction of d_p to S is a fiber bundle on a punctured neighborhood of p in S . The proof boils down to understanding how distance spheres intersect in our CAT(0) homology 4-manifold X .

We apply Proposition 5.5 to the relatively compact set $V := B_{2R}(o)$. Note that $\hat{F} := d_o: V \rightarrow \mathbb{R}$ is an extended $(0, \delta)$ -strainer map for any $\delta > 0$. The halfspace $\hat{\Pi}_p^+$ is exactly the ball $\bar{B}_R(o)$ and $\hat{\Pi}_p = S$.

Corollary 6.1 *There exists a radius $r_p > 0$ such that $S_r(p) \cap S$ is homeomorphic to \mathbb{S}^2 for every $0 < r \leq r_p$.*

Proof By [Lytchak and Nagano 2019, Proposition 9.4] and Lemma 5.3, we can choose r_p such that p is a $(1, \delta)$ -strainer in $\dot{B}_{r_0}(p)$ and (p, o) is an extended $(1, \delta)$ -strainer on $\dot{B}_{r_0}(p) \cap S$. In addition, we choose r_p smaller than the constant $\epsilon_0 = \epsilon_0(p)$ from Proposition 5.5. Then by Propositions 5.4 and 5.5, $S_r(p) \cap \bar{B}_R(o)$ is a contractible homology 3-manifold with boundary $S_r(p) \cap S$, for all $r < r_p$. Thus, $S_r(p) \cap S$ is a homology 2-manifold and therefore a 2-manifold by Theorem 3.1. By Poincaré duality, $S_r(p) \cap S$ is a homology 2-sphere; see [Thurston 1996b, Proposition 2.8]. Due to the classification of surfaces, $S_r(p) \cap S$ is homeomorphic to \mathbb{S}^2 . \square

Lemma 6.2 *There exists $r_0 = r_0(p) > 0$ such that the distance function $d_p: \dot{B}_{r_0}(p) \cap S \rightarrow (0, r_0)$ has uniformly locally contractible fibers.*

Proof Let δ_0 be the constant from Proposition 5.5 and set $\delta = \frac{1}{4}\delta_0$. Let r_p be as in Corollary 6.1. By [Lytchak and Nagano 2019, Proposition 9.4] and Lemma 5.3, we can choose $r_0 < r_p$ such that p is a $(1, \delta)$ -strainer in $\dot{B}_{r_0}(p)$ and (p, o) is an extended $(1, \delta)$ -strainer on $\dot{B}_{r_0}(p) \cap S$.

We fix an arbitrary $x \in \dot{B}_{r_0}(p) \cap S$ and set $t_0 := d_p(x)$. In addition, we fix a positive number $\rho_0 < r_0 - t_0$.

Sublemma *There exists a positive $\epsilon_0 < \rho_0$ and a positive $s_0 < 1$ such that for all t with $|t - t_0| < s_0\epsilon_0$ the intersection of spheres $S_{\epsilon_0}(x) \cap S_t(p) \cap S$ is homeomorphic to \mathbb{S}^1 .*

Proof We apply [Lytchak and Nagano 2019, Proposition 9.4] and Lemma 5.3 and find $\epsilon_0 < \rho_0$ small enough that (p, x) is a $(2, \delta_0)$ -strainer in $\dot{B}_{2\epsilon_0}(x) \cap S_{t_0}(p)$ and (p, x, o) is an extended $(2, \delta_0)$ -strainer in $\dot{B}_{2\epsilon_0}(x) \cap S_{t_0}(p) \cap S$.

Using the openness of the strainer property, we find some small $s_0 > 0$ such that for all t with $|t - t_0| < s_0\epsilon_0$, the pair (p, x) is a $(2, \delta_0)$ -strainer in $\dot{B}_{2\epsilon_0}(x) \cap S_t(p)$ and the triple (p, x, o) is an extended $(2, \delta_0)$ -strainer in $\dot{B}_{2\epsilon_0}(x) \cap S_t(p) \cap S$.

We apply Proposition 5.4 and deduce that for all such t the intersection $S_{\epsilon_0}(x) \cap S_t(p) \cap \bar{B}_R(o)$ is a homology 2–manifold with boundary $S_{\epsilon_0}(x) \cap S_t(p) \cap S$. By Theorem 3.1, these intersections $S_{\epsilon_0}(x) \cap S_t(p) \cap \bar{B}_R(o)$ are 2–manifolds with boundary $S_{\epsilon_0}(x) \cap S_t(p) \cap S$.

By our choice of $\delta_0 = 4\delta$, we may apply Proposition 5.5. By possibly making ϵ_0 and s_0 even smaller, we deduce that all intersections $S_{\epsilon_0}(x) \cap S_t(p) \cap \bar{B}_R(o)$ are contractible and therefore homeomorphic to closed discs. Hence their boundaries $S_{\epsilon_0}(x) \cap S_t(p) \cap S$ are circles. \square

Now we can easily finish the proof of the lemma. By the choice of r_0 and ρ_0 , and Corollary 6.1, any fiber $S_t(p) \cap S$ is homeomorphic to \mathbb{S}^2 .

In order to verify the uniform local contractibility of the fibers of the restriction of d_p , we will argue that for every t with $|t - t_0| < s_0\epsilon_0$, the set $B_{\epsilon_0}(x) \cap S_t(p) \cap S$ is contractible inside $B_{\rho_0}(x) \cap S_t(p) \cap S$.

In the same parameter range as above, $B_{\epsilon_0}(x) \cap S_t(p) \cap S$ is an open subset of the 2–sphere $S_t(p) \cap S$ whose topological boundary inside $S_t(p) \cap S$ is contained in the circle $S_{\epsilon_0}(x) \cap S_t(p) \cap S$. Therefore, by the Jordan curve theorem, $\bar{B}_{\epsilon_0}(x) \cap S_t(p) \cap S$ is either a topological disc or all of $S_t(p) \cap S$. In both cases, $B_{\epsilon_0}(x) \cap S_t(p) \cap S$ is contractible inside $\bar{B}_{\epsilon_0}(x) \cap S_t(p) \cap S$. Therefore $B_{\epsilon_0}(x) \cap S_t(p) \cap S$ is contractible inside the larger set $B_{\rho_0}(x) \cap S_t(p) \cap S$. \square

6.2 The main results

Proof of Theorem 1.4 Let $p \in S = S_R(o)$ be arbitrary. Choose r_p as in Corollary 6.1 and $r_0 < r_p$ as in Lemma 6.2. The distance function $d_p: \dot{B}_{r_0}(p) \cap S \rightarrow (0, r_0)$ has uniformly locally contractible fibers homeomorphic to \mathbb{S}^2 by Corollary 6.1 and Lemma 6.2. By Lemma 5.1 and Theorem 3.5, d_p is a fiber bundle. Hence $\dot{B}_{r_0}(p) \cap S$ is homeomorphic to $\mathbb{S}^2 \times (0, r_0)$. Therefore $B_{r_0}(p) \cap S$ is homeomorphic to a 3–ball. Since p was arbitrary, $S = S_R(o)$ is a 3–manifold, as required. \square

Now the main result of [Thurston 1996b] implies Theorem 1.1.

Before turning to Corollaries 1.5 and 1.6, we recall some notation. We fix a CAT(0) 4–manifold X and a point $o \in X$.

For $R > r > 0$ we have a canonical *geodesic contraction map*

$$c_{R,r}: S_R(o) \rightarrow S_r(o)$$

defined by sending $y \in S_R(o)$ to the point of intersection of the geodesic oy with $S_r(o)$.

The ideal boundary $\partial_\infty X$ is canonically identified with the *inverse limit for the bonding maps* $c_{R,r}$ [Bridson and Haefliger 1999, Section II.8.5; Fujiwara et al. 2004, page 872]:

$$\partial_\infty X = \varprojlim S_r(o).$$

The canonical map $c_{\infty,r}: \partial_\infty X \rightarrow S_r(o)$, the inverse limit of the bonding maps $c_{R,r}$, sends a point $\xi \in \partial_\infty X$ to the intersection with $S_r(o)$ of the ray starting in o and determined by ξ [Bridson and Haefliger 1999, Section II.8].

We recall in a special case the notions of cell-like mappings, referring to [Mitchell and Repovš 1988; Thurston 1996b] for details. A compact subset K of a 3-manifold M is called *cell-like* if K is contractible in any neighborhood of K in M . A map $f: M \rightarrow N$ between 3-manifolds is called a *cell-like map* if the preimage of any point is cell-like.

The following result is a combination of [Mitchell and Repovš 1988, Theorem 1.2 and Corollary 1.4; McMillan 1967, Corollary 2.2]:

Theorem 6.3 *Let M be a compact 3-manifold and $f: M \rightarrow M$ be a surjective cell-like map. Then f is a uniform limit of homeomorphisms. For any open subset $U \subset M$, the restriction $f: f^{-1}(U) \rightarrow U$ is a homotopy equivalence.*

Now we turn to the proof of Corollary 1.5, which is essentially contained in the proof of [Thurston 1996b, Theorem 4.3].

Proof of Corollary 1.5 By Theorems 1.1 and 1.4 and [Thurston 1996b, Theorem 4.3] all distance spheres $S_R(o)$ are homotopy equivalent to $\mathbb{R}^4 \setminus \{o\}$, and hence to \mathbb{S}^3 . By Theorem 1.4 and the resolution of the Poincaré conjecture, any sphere $S_R(o)$ is homeomorphic to \mathbb{S}^3 .

Hence, for any geodesic contraction $c_{R,r}: S_R(o) \rightarrow S_r(o)$ the preimage of any point is contained in a subset homeomorphic to \mathbb{R}^3 . Combining [Thurston 1996b, Corollary 2.10 and Theorem 2.13] and the subsequent remark, we deduce that $c_{R,r}$ are cell-like maps.

By Theorem 6.3, for any open contractible set $W \subset S_r(o)$, the preimages $c_{R,r}^{-1}(W)$ are contractible for all $R > r$. This implies that the map $d_o: X \setminus \{o\} \rightarrow (0, \infty)$ has uniformly locally contractible fibers. Indeed, for any $t > \epsilon > 0$ and any $x \in S_t(o)$, consider the point $z = c_{t,t-\epsilon}(x)$. Find an open contractible neighborhood W of diameter less than ϵ around x in the manifold $S_t(o)$. Then $\widehat{W} := \bigcup_{t-\epsilon < s < t+\epsilon} c_{s,t+\epsilon}^{-1}(W)$ is an open neighborhood of x in X of diameter at most 4ϵ . As we have seen above, the intersection of \widehat{W} with any fiber of the function d_o is contractible. Hence, d_o has uniformly locally contractible fibers.

Therefore an application of Theorem 3.6 completes the proof. □

In the proof of Corollary 1.6 below we assume some knowledge of the ideal boundary, its cone topology and the canonical compactification $\bar{X} = X \cup \partial_\infty X$ of a CAT(0) space X .

Proof of Corollary 1.6 We fix a point o in the CAT(0) 4-manifold X . By Corollary 1.5, every distance sphere $S_R(o)$ is homeomorphic to \mathbb{S}^3 .

As we have seen in the proof of Corollary 1.5 above, all geodesic contractions $c_{R,r}: S_R(o) \rightarrow S_r(o)$ are cell-like maps. Hence, due to Theorem 6.3, the map $c_{R,r}$ is a uniform limit of homeomorphisms. Thus, an application of the main result of [Brown 1960] implies that the ideal boundary $\partial_\infty X$ is homeomorphic to \mathbb{S}^3 . Moreover, the proof in [Brown 1960] shows that the canonical projection

$$c_{\infty,r}: \partial_\infty X \rightarrow S_r(o)$$

is a uniform limit of homeomorphisms.

Consider the map $f: \bar{X} \rightarrow (0, 1]$ which sends $\partial_\infty X$ to 1 and is defined as

$$f(x) := \frac{d_o(x)}{1 + d_o(x)}$$

on X . The map f is continuous on \bar{X} and coincides on X with d_o up to a homeomorphism of the image interval. Hence f is a fiber bundle on $X \setminus \{o\}$. We claim that f is also a fiber bundle on all of $\bar{X} \setminus \{o\}$.

All fibers of f are 3–spheres. Moreover, for any open contractible W in any sphere $S_r(o)$ the preimages $c_{R,r}^{-1}(W)$ are contractible, for all $R \in [r, \infty]$, since all geodesic contractions including $c_{\infty,r}$ are cell-like. This implies that f has uniformly locally contractible fibers.

Theorem 3.6 implies the claim. Therefore, $\bar{X} \setminus \{o\}$ is homeomorphic to $\mathbb{S}^3 \times (0, 1]$. Thus \bar{X} , the one-point compactification of this space, is homeomorphic to the 4–ball. \square

7 Structure of fibers of extended strainer maps

7.1 Generalized distance functions

In this final section, we want to use information on limits of distance maps to conclude topological properties of their fibers. This requires a slight generalization of the notion of distance functions and strainer maps. For this purpose we make the following definitions; see also [Nagano 2022, Section 5]. Recall that a convex function on a CAT(0) space attains its minimal value on a closed convex set or doesn't attain a minimum at all. A *generalized distance function* on a CAT(0) space X is a convex function $b: X \rightarrow \mathbb{R}$ whose (negative) gradient has unit norm on the complement of its minimal set:

$$\|\nabla_x(-b)\| := \max\left\{0, \limsup_{y \rightarrow x} \frac{b(x) - b(y)}{d(x, y)}\right\} \equiv 1.$$

This definition unifies the concept of distance functions to convex subsets and Busemann functions. Adding a constant to a generalized distance function results in a generalized distance function. On every bounded open set, a generalized distance function equals the distance function to a convex set up to a constant. In particular, the integral curves of the “negative gradient” of a generalized distance function are geodesics and the *negative gradient* $\nabla_x(-b) \in \Sigma_x X$ is well defined.

A map $F: X \rightarrow \mathbb{R}^k$ will be called a *generalized distance map* if all coordinates f_i of F are generalized distance functions.

A generalized distance map $F: X \rightarrow \mathbb{R}^k$ with components f_i will be called a *generalized (k, δ) –strainer map in a subset $A \subset X$* if the minimum sets of any f_i are disjoint from A and, for any $x \in A$, the negative gradients $\nabla_x(-f_i) \in \Sigma_x X$ form a δ –spherical k –tuple of directions in $\Sigma_x X$.

Two generalized distance maps $F, \bar{F}: X \rightarrow \mathbb{R}^k$ with coordinates f_i and \bar{f}_i are *opposite generalized (k, δ) –strainer maps* on $A \subset X$ if, for all $x \in A$, the corresponding k –tuples of negative gradients are opposite (k, δ) –strainers.

As for (nongeneralized) distance maps, the set of points $x \in X$ at which a generalized distance map $F: X \rightarrow \mathbb{R}^k$ is a generalized (k, δ) -strainer map is open. Similarly, the set of points at which a pair of distance maps F and \bar{F} are opposite generalized (k, δ) -strainers is open.

Let $F: X \rightarrow \mathbb{R}^k$ be a generalized distance map with coordinates f_i and denote by b another generalized distance function. Suppose that F is a generalized (k, δ) -strainer map on a subset $A \subset X$ and b does not attain its minimum on A . Then the map $\hat{F} = (F, b): X \rightarrow \mathbb{R}^{k+1}$ is called a *generalized extended (k, δ) -strainer map* on A if, at all points $x \in A$ and for every antipode $w_x \in \Sigma_x X$ of $\nabla_x(-b)$, the following holds for all $1 \leq i \leq k$:

$$\angle_x(\nabla_x(-b), \nabla_x(-f_i)) < \frac{1}{2}\pi + \delta \quad \text{and} \quad \angle_x(w_x, \nabla_x(-f_i)) < \frac{1}{2}\pi + \delta.$$

The set of points where a given generalized distance map is a generalized extended (k, δ) -strainer map is open, again due to the semicontinuity of angles.

All statements about (extended) strainer maps transfer to the generalized setting. For instance, the concept of “straining radius” introduced in [Lytchak and Nagano 2019, Section 7.5] generalizes as follows. Let $F: X \rightarrow \mathbb{R}^k$ be a generalized distance map with coordinates f_i . Suppose that F is a generalized (k, δ) -strainer map at a point x . Then the *straining radius* is the largest radius σ_x with the following property: For every $y \in B_{\sigma_x}(x)$ and $1 \leq i \leq k$ let \bar{p}_i be any point with $d(x, \bar{p}_i) = 1$ and such that the direction at x of the geodesic $x\bar{p}_i$ is antipodal to $\nabla(-f_i)$. Then F and $\bar{F} = (d_{\bar{p}_1}, \dots, d_{\bar{p}_k})$ are opposite generalized $(k, 2\delta)$ -strainer maps on $B_{\sigma_x}(y)$. The proof of positivity of σ_x is identical to [loc. cit., Lemma 7.10]. Similarly, we define an “extended straining radius”. If $\hat{F} = (F, b)$ is an extended generalized (k, δ) -strainer map at x , then the *extended straining radius* is the largest radius $\hat{\sigma}_x \leq \sigma_x$ such that for all $y \in B_{\hat{\sigma}_x}(x)$ the map \hat{F} is an extended generalized $(k, 2\delta)$ -strainer map on $B_{\hat{\sigma}_x}(y)$.

Let (X_n, x_n) be a sequence of pointed locally compact CAT(0) spaces converging in the pointed Gromov–Hausdorff topology to a space (X, x) . Then, for any sequence of generalized distance functions $f_n: X_n \rightarrow \mathbb{R}$ with uniformly bounded $f_n(x_n)$, we find a subsequence converging to a generalized distance function $f: X \rightarrow \mathbb{R}$.

Let $F_n: X_n \rightarrow \mathbb{R}^k$ be a sequence of generalized distance maps converging to a generalized distance map $F: X \rightarrow \mathbb{R}^k$. The semicontinuity of angles under convergence implies the following, as in [loc. cit., Lemma 7.8]: if F is a generalized (extended) (k, δ) -strainer at x then F_n is a generalized (extended) (k, δ) -strainer at x_n , for all n large enough. Moreover, for all n large enough, the (extended) straining radius σ_{x_n} of F_n at x_n is bounded from below by half of the (extended) straining radius σ_x .

7.2 Local topology of halfspaces

The following result on strainer maps translates to the generalized setting as well, but since we only apply it in the nongeneralized setting, and since this allows us to directly rely on [loc. cit., Theorem 9.1], we refrained from formulating a generalized version even though proofs extend literally.

Proposition 7.1 Let $\widehat{F} = (F, d_o): X \rightarrow \mathbb{R}^k$ be a distance map and $\delta \leq 1/(64k)$. Suppose that \widehat{F} is an extended (k, δ) -strainer map at a point x with extended strainer radius $\widehat{\sigma}_x$. Denote by W a ball $B_r(x)$ with radius $r \leq \widehat{\sigma}_x$. Then there exists a deformation retraction of W onto the halfspace $\widehat{\Pi}_x^+ \cap W$.

Proof The proof is an adaption of the proof of [loc. cit., Theorem 9.1]. For convenience of the reader, we stick to the notation used there. Hence (p_i) denotes the strainer defining F , and (q_i) is a k -tuple in $X \setminus W$ such that x lies on the geodesic $p_i q_i$ and the tuples (p_i) and (q_i) are opposite $(k, 2\delta)$ -strainers in W .

Set $s_0 := d(o, x)$ and define the function $M: W \rightarrow \mathbb{R}$ by

$$M(z) := \max_{1 \leq i \leq k} |d(p_i, z) - d(p_i, x)|.$$

Denote by $\Phi: W \times [0, 1] \rightarrow W$ the homotopy which retracts W onto $\Pi_x \cap W$, provided by [loc. cit., Theorem 9.1]. Recall that the length of the path $\gamma_z(t) := \Phi(z, t)$ is at most $8kM(z)$. Moreover, $\gamma_z(t)$ is an infinite piecewise geodesic all of whose segments are directed towards one of the points p_i or q_i . By the first variation formula, the value of d_o changes along $\gamma_z(t)$ with velocity at most 4δ . Hence for all $t \in [0, 1]$,

$$(7-1) \quad |d_o(z) - d_o(\gamma_z(t))| \leq 4\delta \cdot 8k \cdot M(z) \leq \frac{1}{2}M(z).$$

Denote by $\varphi: W \times [0, 1] \rightarrow W$ the flow which deformation retracts W onto $\overline{B}_{s_0}(o) \cap W$. More precisely, φ moves a point $z \in W \setminus \overline{B}_{s_0}(o)$ towards o at unit speed until it reaches $S_{s_0}(o)$ and then stops. Note that φ does indeed preserve W , by the CAT(0) property of X . We define a concatenated homotopy Ψ by setting $\Psi(z, t) = \Phi(z, 2t)$ for $t \leq \frac{1}{2}$ and $\Psi(z, t) = \varphi(\Phi(z, 1), 2t - 1)$ for $t \geq \frac{1}{2}$.

By definition, $d(o, \Psi(z, 1)) \leq s_0$ for all $z \in W$ and Ψ fixes $\widehat{\Pi}_x^+ \cap W$.

The length of the Ψ -flow line of a general point $z \in W$ is bounded above by $8kM(z) + 2r$. However, if $d(o, z) \leq s_0$ holds, then the length of the Ψ -flow line starting at z is at most $(1 + 4\delta)8kM(z)$ by (7-1).

Along the homotopy φ , the value of M changes at most with velocity 4δ , due to the first variation formula. Hence, for any z with $d(o, z) \leq s_0$, we deduce, using $M(\Phi(z, 1)) = 0$ and (7-1),

$$M(\Psi(z, 1)) \leq 2\delta M(z).$$

To obtain the required deformation retraction, we take a limit of iterated concatenations of Ψ . More precisely, for $m \geq 1$ we define homotopies $\Psi_m: W \times [0, 1] \rightarrow W$ as follows. The homotopy Ψ_m is the identity on the interval $[1 - 2^{-m}, 1]$ and it equals a rescaling of Ψ on any of the intervals $[1 - 2^{-l}, 1 - 2^{-l-1}]$, for $l = 0, \dots, m - 1$.

The above inequalities imply $M(\Psi_m(z, 1)) \leq (2\delta)^m M(z)$, by induction. Moreover, the flow line of Ψ_m starting at $z \in W$ has length uniformly bounded above by $2r + 24kM(z)$. Therefore (Ψ_m) converges uniformly to a homotopy $\Psi_\infty: W \times [0, 1] \rightarrow W$, as required. \square

Recall that a closed subset Π of a topological space Y is called *homotopy negligible in Y* if for each open set U of Y the inclusion $U \setminus \Pi \rightarrow U$ is a homotopy equivalence. If Y is an ANR, this condition is satisfied

if any point $z \in \Pi$ has a neighborhood basis \mathcal{U}_z of contractible neighborhoods U_z with contractible complements $U_z \setminus \Pi$ [Eells and Kuiper 1969, Theorem 1]. In our setting, we have:

Corollary 7.2 *Let $\hat{F} = (F, d_o): U \rightarrow \mathbb{R}^k$ be an extended (k, δ) -strainer map on an open set $U \subset X$ with $\delta \leq 1/(64k)$. Then, for every $x \in U$, the halfspace $\hat{\Pi}_x^+$ is an ANR and the fiber $\hat{\Pi}_x$ is homotopy negligible in $\hat{\Pi}_x^+$.*

Proof By [Lytchak and Nagano 2019, Theorem 9.1], for all $y \in \hat{\Pi}_x^+ \setminus \hat{\Pi}_x$ the set $\hat{\Pi}_x^+ \cap B_r(y)$ is contractible as a retract of $B_r(y)$, as long as the radius r is less than the straining radius σ_y and the difference of levels $d_o(x) - d_o(y)$. Similarly, by Proposition 7.1, for all $z \in \hat{\Pi}_x$, the set $\hat{\Pi}_x^+ \cap B_r(z)$ is contractible as a retract of $B_r(z)$ for some radius $r < \hat{\sigma}_z$. Hence $\hat{\Pi}_x^+$ is an ANR. Now for $z \in \hat{\Pi}_x$, set $W = B_{\hat{\sigma}_z}(z)$ as above. It remains to show that $(\hat{\Pi}_x^+ \setminus \hat{\Pi}_x) \cap W$ is contractible. Since it is an ANR as an open subset of $\hat{\Pi}_x^+$, it suffices to verify that all of its homotopy groups vanish [Hu 1965, Corollary VII.8.5]. This will follow once we have shown that, for any compact subset $K \subset (\hat{\Pi}_x^+ \setminus \hat{\Pi}_x) \cap W$, the inclusion map $K \hookrightarrow (\hat{\Pi}_x^+ \setminus \hat{\Pi}_x) \cap W$ is nullhomotopic. By continuity of the straining radius $\hat{\sigma}_x$, given such a set K , we find a point $w \in (\hat{\Pi}_x^+ \setminus \hat{\Pi}_x) \cap W$ and $s \leq \hat{\sigma}_w$ with $K \subset B_s(w) \subset W$. By Proposition 7.1, $\hat{\Pi}_w^+ \cap B_s(w) \subset (\hat{\Pi}_x^+ \setminus \hat{\Pi}_x) \cap W$ is contractible and the proof is complete. \square

Proof of Proposition 5.4 We have already seen in Corollary 7.2 that the halfspace $\hat{\Pi}_x^+$ is an ANR.

Assume now that U is a homology n -manifold. Since $\delta < 1/(20k)$, Theorem 4.1 implies that the fibers of F are homology $(n-k)$ -manifolds. The complement of $\hat{\Pi}_x$ in $\hat{\Pi}_x^+$ is open in Π_x and therefore a homology $(n-k)$ -manifold. By Corollary 7.2, $\hat{\Pi}_x$ is homotopy negligible in $\hat{\Pi}_x^+$. In particular, every singleton $\{y\} \subset \hat{\Pi}_x$ is homotopy negligible in $\hat{\Pi}_x^+$ [Toruńczyk 1978, Corollary 2.6]. We conclude that the local homology groups $H_*(\hat{\Pi}_x^+, \hat{\Pi}_x^+ \setminus \{y\})$ vanish at all points $y \in \hat{\Pi}_x$. By [Mitchell 1990], $\hat{\Pi}_x$ is the boundary of a homology manifold and therefore is itself a homology manifold without boundary. \square

7.3 Contractibility of hemispheres

Proof of Proposition 5.5 Suppose for the sake of contradiction that there is a sequence $\delta_l \rightarrow 0$ and distance maps $\hat{F}_l = (F_l, d_{q_l}): X \rightarrow \mathbb{R}^{k+1}$ with $F_l = (d_{p_1^l}, \dots, d_{p_k^l})$ which are extended (k, δ_l) -strainer maps at points $x_l \in V$ where the statement fails. Thus we find arbitrarily small “hemispheres” around x_l which are either not contractible or not locally contractible. More precisely, we find sequences $\epsilon_l \rightarrow 0$, $s_l \rightarrow 0$ and a sequence of points $y_l \in X$ with $d(x_l, y_l) < s_l \epsilon_l$ and the following additional properties:

(1) **Gromov–Hausdorff close to tangent space**

$$|\bar{B}_{\epsilon_l}(x_l), \bar{B}_{\epsilon_l}(o_{x_l})|_{\text{GH}} < \frac{\epsilon_l}{l}.$$

(2) **Improved strainer** The map (d_{x_l}, F_l) is a $(k+1, 4\delta_l)$ -strainer map on $\hat{B}_{\epsilon_l}(x_l)$.

(3) **Improved extended strainer** The map (d_{x_l}, \hat{F}_l) is an extended $(k+1, 4\delta_l)$ -strainer map on an open neighborhood V_l of $\hat{\Pi}_{x_l} \cap \hat{B}_{\epsilon_l}(x_l)$.

(4) **Large levels** $\min\{d_{p_1^l}(x_l), \dots, d_{p_k^l}(x_l), d_{q^l}(x_l)\} \geq l\epsilon_l$.

(5) **Fiber lies in extended domain** $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l} \subset V_l$.

(6) **Noncontractible** $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+$ is either not contractible or not locally contractible.

The first item can be arranged because, in our setting, tangent spaces are Gromov–Hausdorff limits of rescaled balls around a particular point [Lytchak and Nagano 2019, Corollary 5.7]. The second and third items follow from [loc. cit., Proposition 9.4] and Lemma 5.3, respectively, by choosing ϵ_l small enough. Similarly, the fourth item can be achieved by choosing ϵ_l small enough. Finally, the fifth item can be guaranteed by choosing s_l small enough.

We define the shifted strainer maps $\Phi_l = (d_{p_1^l} - d_{p_1^l}(x_l), \dots, d_{p_k^l} - d_{p_k^l}(x_l))$, as well as the shifted distance function $b_l = d_{q^l} - d_{q^l}(x_l)$. In particular, $\Phi_l(x_l) = 0$. Now we rescale space and functions by $1/\epsilon_l$. Since V is relatively compact, up to passing to subsequences, we can take a pointed Gromov–Hausdorff limit $(X_\infty, x_\infty) = \lim_{l \rightarrow \infty} ((1/\epsilon_l) \cdot X, x_l)$ [loc. cit., Proposition 5.10]. We also pass to corresponding limits of functions $\Phi_\infty = \lim_{l \rightarrow \infty} (1/\epsilon_l)\Phi_l$ and $b_\infty = \lim_{l \rightarrow \infty} (1/\epsilon_l)b_l$. Item (4) ensures that all coordinates of Φ_∞ , as well as the function b_∞ , are Busemann functions on X_∞ [Kapovich and Leeb 1997, Lemma 2.3].

By (1), (X_∞, x_∞) is isometric to a pointed Gromov–Hausdorff limit of the sequence of tangent spaces $(T_{x_l} X, o_{x_l})$. In particular, (X_∞, x_∞) is isometric to a Euclidean cone with tip x_∞ . Therefore $S_1(x_\infty)$ is a CAT(1) space [Berestovsky 1983]. Moreover, the spaces of directions $\Sigma_{x_l} X$ converge to $S_1(x_\infty)$ [Lytchak and Nagano 2019, Theorem 13.1]. By assumption, the negative gradients of the components of Φ_l provide a δ_l –spherical k –tuple of directions at x_l . Then [loc. cit., Proposition 6.6] implies that $S_1(x_\infty)$ splits isometrically as a spherical join $S_1(x_\infty) \cong \mathbb{S}^{k-1} * \Sigma'$; equivalently, X_∞ splits isometrically as a direct product $X_\infty \cong \mathbb{R}^k \times X'_\infty$. Moreover, the negative gradients of the components of Φ_∞ form a spherical k –tuple inside the \mathbb{S}^{k-1} –factor of $S_1(x_\infty)$.

From the L –openness of (d_{x_l}, \widehat{F}_l) , given by Lemma 5.1, we conclude the Gromov–Hausdorff convergence: $\lim_{l \rightarrow \infty} (1/\epsilon_l) \cdot (S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+) = S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$.

Sublemma *The sequence $(1/\epsilon_l) \cdot (S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+)$ is uniformly locally contractible.*

Proof By compactness of $S_1(x_\infty)$, we find $r > 0$ such that the straining radius of Φ_∞ satisfies $\sigma_z > 2r$ at all $z \in S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}$ and the extended straining radius of (Φ_∞, b_∞) satisfies $\hat{\sigma}_z > 2r$ at all $z \in S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$. Then, for l large enough, the straining radius of F_l and the extended straining radius of (F_l, b_l) are larger than $r\epsilon_l$ on $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}$ and $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+$, respectively.

Let $z_l \in S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+$ be a point. If the distance from z_l to the fiber $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}$ is at least $\frac{1}{2}r\epsilon_l$, then $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+ \cap B_{(r/2)\epsilon_l}(z_l)$ is contractible by [loc. cit., Theorem 9.1]. On the other hand, if the distance from z_l to the fiber $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}$ is smaller than $\frac{1}{2}r\epsilon_l$, then $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+ \cap B_{(r/2)\epsilon_l}(z_l)$ is contained in an ball $B_{r\epsilon_l}(w_l)$ with $w_l \in S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}$. By Proposition 7.1, the set $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+ \cap B_{r\epsilon_l}(w_l)$ is contractible. It follows that any $\frac{1}{2}r$ –ball in $(1/\epsilon_l) \cdot (S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+)$ is contractible inside its concentric $2r$ –ball. \square

As one consequence, the same local contractibility holds for the hemispheres $S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$ [Petersen 1993, Theorem 9]. Moreover, $S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$ is homotopy equivalent to $S_{\epsilon_l}(x_l) \cap \widehat{\Pi}_{y_l}^+$, for large enough l . Hence, to arrive at a contradiction, it remains to show that $S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$ is contractible.

Let $v \in S_1(x_\infty)$ denote the point corresponding to the negative gradient of b_∞ . By semicontinuity of angles and the splitting $S_1(x_\infty) \cong \mathbb{S}^{k-1} * \Sigma'$, we see $v \in \Sigma'$; see Section 5.2. In particular, $b_\infty = b'_\infty \circ \pi'$, where π' denotes the projection $X_\infty \cong \mathbb{R}^k \times X'_\infty \rightarrow X'_\infty$, and b'_∞ is a Busemann function on X'_∞ . We infer $\widehat{\Pi}_{x_\infty}^+ \cong \{0\} \times \{b'_\infty \leq 0\}$ since $\Phi_\infty(x_\infty) = 0$. Hence,

$$S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+ \cong \Sigma_{x_\infty} X'_\infty \cap \{b'_\infty \leq 0\}.$$

But $\Sigma_{x_\infty} X'_\infty \cap \{b'_\infty \leq 0\} = \bar{B}_{\pi/2}(v) \subset \Sigma_{x_\infty} X'_\infty$ and therefore $\Sigma_{x_\infty} X'_\infty \cap \{b'_\infty \leq 0\}$ is contractible, since $\Sigma_{x_\infty} X'_\infty$ is CAT(1). Consequently, the hemisphere $S_1(x_\infty) \cap \widehat{\Pi}_{x_\infty}^+$ is contractible. \square

References

- [Adiprasito and Benedetti 2020] **K Adiprasito, B Benedetti**, *Collapsibility of CAT(0) spaces*, *Geom. Dedicata* 206 (2020) 181–199 MR Zbl
- [Adiprasito and Funar 2015] **K A Adiprasito, L Funar**, *CAT(0) metrics on contractible manifolds*, preprint (2015) arXiv 1512.06403
- [Alexander and Bishop 1990] **S B Alexander, R L Bishop**, *The Hadamard–Cartan theorem in locally convex metric spaces*, *Enseign. Math.* 36 (1990) 309–320 MR Zbl
- [Alexander et al. 2024] **S Alexander, V Kapovitch, A Petrunin**, *Alexandrov geometry: foundations*, *Graduate Studies in Math.* 236, Amer. Math. Soc., Providence, RI (2024) MR Zbl
- [Ancel and Guilbault 1997] **F D Ancel, C R Guilbault**, *Interiors of compact contractible n -manifolds are hyperbolic ($n \geq 5$)*, *J. Differential Geom.* 45 (1997) 1–32 MR Zbl
- [Ancel et al. 1997] **F D Ancel, M W Davis, C R Guilbault**, *CAT(0) reflection manifolds*, from “Geometric topology, I” (W H Kazez, editor), *AMS/IP Stud. Adv. Math.* 2, Amer. Math. Soc., Providence, RI (1997) 441–445 MR Zbl
- [Berestovsky 1983] **V N Berestovsky**, *Borsuk’s problem on metrization of a polyhedron*, *Dokl. Akad. Nauk SSSR* 268 (1983) 273–277 MR Zbl In Russian; translated in *Soviet Math. Dokl.* 27 (1983) 56–59
- [Berestovsky et al. 2011] **V N Berestovsky, D M Halverson, D Repovš**, *Locally G -homogeneous Busemann G -spaces*, *Differential Geom. Appl.* 29 (2011) 299–318 MR Zbl
- [Bridson and Haefliger 1999] **M R Bridson, A Haefliger**, *Metric spaces of non-positive curvature*, *Grundle. Math. Wissen.* 319, Springer (1999) MR Zbl
- [Brown 1960] **M Brown**, *Some applications of an approximation theorem for inverse limits*, *Proc. Amer. Math. Soc.* 11 (1960) 478–483 MR Zbl
- [Brown 1961] **M Brown**, *The monotone union of open n -cells is an open n -cell*, *Proc. Amer. Math. Soc.* 12 (1961) 812–814 MR Zbl
- [Burago et al. 1992] **Y Burago, M Gromov, G Perelman**, *A D Aleksandrov spaces with curvatures bounded below*, *Uspekhi Mat. Nauk* 47 (1992) 3–51 MR Zbl In Russian; translated in *Russian Math. Surv.* 47 (1992) 1–58

- [Burago et al. 2001] **D Burago, Y Burago, S Ivanov**, *A course in metric geometry*, Graduate Studies in Math. 33, Amer. Math. Soc., Providence, RI (2001) MR Zbl
- [Busemann 1955] **H Busemann**, *The geometry of geodesics*, Academic, New York (1955) MR Zbl
- [Cannon 1978] **J W Cannon**, *The recognition problem: what is a topological manifold?*, Bull. Amer. Math. Soc. 84 (1978) 832–866 MR Zbl
- [Cavicchioli et al. 2016] **A Cavicchioli, F Hegenbarth, D Repovš**, *Higher-dimensional generalized manifolds: surgery and constructions*, Eur. Math. Soc., Zürich (2016) MR Zbl
- [Daverman and Walsh 1981] **R J Daverman, J J Walsh**, *A ghastly generalized n -manifold*, Illinois J. Math. 25 (1981) 555–576 MR Zbl
- [Davis and Januszkiewicz 1991] **M W Davis, T Januszkiewicz**, *Hyperbolization of polyhedra*, J. Differential Geom. 34 (1991) 347–388 MR Zbl
- [Davis et al. 2012] **M Davis, T Januszkiewicz, J-F Lafont**, *4-Dimensional locally CAT(0)-manifolds with no Riemannian smoothings*, Duke Math. J. 161 (2012) 1–28 MR Zbl
- [Eells and Kuiper 1969] **J Eells, Jr, N H Kuiper**, *Homotopy negligible subsets*, Compos. Math. 21 (1969) 155–161 MR Zbl
- [Ferry 1978] **S Ferry**, *Strongly regular mappings with compact ANR fibers are Hurewicz fiberings*, Pacific J. Math. 75 (1978) 373–382 MR Zbl
- [Ferry 1991] **S C Ferry**, *Alexander duality and Hurewicz fibrations*, Trans. Amer. Math. Soc. 327 (1991) 201–219 MR Zbl
- [Freedman 1982] **M H Freedman**, *The topology of four-dimensional manifolds*, J. Differential Geom. 17 (1982) 357–453 MR Zbl
- [Fujiwara et al. 2004] **K Fujiwara, T Shioya, S Yamagata**, *Parabolic isometries of CAT(0) spaces and CAT(0) dimensions*, Algebr. Geom. Topol. 4 (2004) 861–892 MR Zbl
- [Gromov 1981] **M Gromov**, *Hyperbolic manifolds, groups and actions*, from “Riemann surfaces and related topics” (I Kra, B Maskit, editors), Ann. of Math. Stud. 97, Princeton Univ. Press (1981) 183–213 MR Zbl
- [Halverson and Repovš 2008] **D M Halverson, D Repovš**, *The Bing–Borsuk and the Busemann conjectures*, Math. Commun. 13 (2008) 163–184 MR Zbl
- [Hatcher 2002] **A Hatcher**, *Algebraic topology*, Cambridge Univ. Press (2002) MR Zbl
- [Hu 1965] **S-T Hu**, *Theory of retracts*, Wayne State Univ. Press, Detroit, MI (1965) MR Zbl
- [Husch and Price 1970] **L S Husch, T M Price**, *Finding a boundary for a 3-manifold*, Ann. of Math. 91 (1970) 223–235 MR Zbl
- [Ivanov and Lytchak 2019] **S Ivanov, A Lytchak**, *Rigidity of Busemann convex Finsler metrics*, Comment. Math. Helv. 94 (2019) 855–868 MR Zbl
- [Kapovich and Leeb 1997] **M Kapovich, B Leeb**, *Quasi-isometries preserve the geometric decomposition of Haken manifolds*, Invent. Math. 128 (1997) 393–416 MR Zbl
- [Krakus 1968] **B Krakus**, *Any 3-dimensional G -space is a manifold*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 16 (1968) 737–740 MR Zbl
- [Kramer 2011] **L Kramer**, *On the local structure and the homology of CAT(κ) spaces and Euclidean buildings*, Adv. Geom. 11 (2011) 347–369 MR Zbl

- [Lytchak 2005] **A Lytchak**, *Rigidity of spherical buildings and joins*, *Geom. Funct. Anal.* 15 (2005) 720–752 MR Zbl
- [Lytchak and Nagano 2019] **A Lytchak, K Nagano**, *Geodesically complete spaces with an upper curvature bound*, *Geom. Funct. Anal.* 29 (2019) 295–342 MR Zbl
- [Lytchak and Nagano 2022] **A Lytchak, K Nagano**, *Topological regularity of spaces with an upper curvature bound*, *J. Eur. Math. Soc.* 24 (2022) 137–165 MR Zbl
- [Lytchak and Schroeder 2007] **A Lytchak, V Schroeder**, *Affine functions on CAT(κ)-spaces*, *Math. Z.* 255 (2007) 231–244 MR Zbl
- [Lytchak and Stadler 2020] **A Lytchak, S Stadler**, *Improvements of upper curvature bounds*, *Trans. Amer. Math. Soc.* 373 (2020) 7153–7166 MR Zbl
- [McMillan 1967] **D R McMillan, Jr**, *Strong homotopy equivalence of 3-manifolds*, *Bull. Amer. Math. Soc.* 73 (1967) 718–722 MR Zbl
- [Mitchell 1990] **W J R Mitchell**, *Defining the boundary of a homology manifold*, *Proc. Amer. Math. Soc.* 110 (1990) 509–513 MR Zbl
- [Mitchell and Repovš 1988] **W J R Mitchell, D Repovš**, *The topology of cell-like mappings*, *Rend. Sem. Fac. Sci. Univ. Cagliari* 58 (1988) 265–300 MR Zbl
- [Nagano 2022] **K Nagano**, *Asymptotic topological regularity of CAT(0) spaces*, *Ann. Global Anal. Geom.* 61 (2022) 427–457 MR Zbl
- [Petersen 1990] **P Petersen, V**, *A finiteness theorem for metric spaces*, *J. Differential Geom.* 31 (1990) 387–395 MR Zbl
- [Petersen 1993] **P Petersen, V**, *Gromov–Hausdorff convergence of metric spaces*, from “Differential geometry: Riemannian geometry” (R Greene, S-T Yau, editors), *Proc. Sympos. Pure Math.* 54, Amer. Math. Soc., Providence, RI (1993) 489–504 MR Zbl
- [Raymond 1965] **F Raymond**, *Local triviality for Hurewicz fiberings of manifolds*, *Topology* 3 (1965) 43–57 MR Zbl
- [Repovš 1994] **D Repovš**, *The recognition problem for topological manifolds: a survey*, *Kodai Math. J.* 17 (1994) 538–548 MR Zbl
- [Rolfsen 1968] **D Rolfsen**, *Strongly convex metrics in cells*, *Bull. Amer. Math. Soc.* 74 (1968) 171–175 MR Zbl
- [Sathaye 2017] **B Sathaye**, *Link obstruction to Riemannian smoothings of locally CAT(0) 4-manifolds*, preprint (2017) arXiv 1707.03433
- [Stadler 2015] **S Stadler**, *An obstruction to the smoothability of singular nonpositively curved metrics on 4-manifolds by patterns of incompressible tori*, *Geom. Funct. Anal.* 25 (2015) 1575–1587 MR Zbl
- [Stallings 1962] **J Stallings**, *The piecewise-linear structure of Euclidean space*, *Proc. Cambridge Philos. Soc.* 58 (1962) 481–488 MR Zbl
- [Stone 1976] **D A Stone**, *Geodesics in piecewise linear manifolds*, *Trans. Amer. Math. Soc.* 215 (1976) 1–44 MR Zbl
- [Thurston 1996a] **P Thurston**, *4-Dimensional Busemann G -spaces are 4-manifolds*, *Differential Geom. Appl.* 6 (1996) 245–270 MR Zbl
- [Thurston 1996b] **P Thurston**, *CAT(0) 4-manifolds possessing a single tame point are Euclidean*, *J. Geom. Anal.* 6 (1996) 475–494 MR Zbl

- [Toruńczyk 1978] **H Toruńczyk**, *Concerning locally homotopy negligible sets and characterization of l_2 -manifolds*, *Fund. Math.* 101 (1978) 93–110 MR Zbl
- [Ungar 1969] **G S Ungar**, *Conditions for a mapping to have the slicing structure property*, *Pacific J. Math.* 30 (1969) 549–553 MR Zbl
- [Wilder 1949] **R L Wilder**, *Topology of manifolds*, *Amer. Math. Soc. Colloq. Publ.* 32, Amer. Math. Soc., Providence, RI (1949) MR Zbl
- [Wu 1997] **J-Y Wu**, *Topological regularity theorems for Alexandrov spaces*, *J. Math. Soc. Japan* 49 (1997) 741–757 MR Zbl

*Institut für Algebra und Geometrie, Karlsruher Institut für Technologie
Karlsruhe, Germany*

*Department of Mathematics, University of Tsukuba
Tsukuba, Japan*

*Max Planck Institute for Mathematics
Bonn, Germany*

alexander.lytchak@kit.edu, nagano@math.tsukuba.ac.jp, stadler@mpim-bonn.mpg.de

Proposed: Urs Lang

Seconded: John Lott, Bruce Kleiner

Received: 27 May 2022

Revised: 29 March 2023

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

| | | | |
|----------------------|--|--------------------|---|
| Mohammed Abouzaid | Stanford University abouzaid@stanford.edu | Mark Gross | University of Cambridge mgross@dpmms.cam.ac.uk |
| Dan Abramovich | Brown University dan_abramovich@brown.edu | Rob Kirby | University of California, Berkeley kirby@math.berkeley.edu |
| Ian Agol | University of California, Berkeley ianagol@math.berkeley.edu | Bruce Kleiner | NYU, Courant Institute bkleiner@cims.nyu.edu |
| Arend Bayer | University of Edinburgh arend.bayer@ed.ac.uk | Sándor Kovács | University of Washington skovacs@uw.edu |
| Mark Behrens | University of Notre Dame mbehren1@nd.edu | Urs Lang | ETH Zürich urs.lang@math.ethz.ch |
| Mladen Bestvina | University of Utah bestvina@math.utah.edu | Marc Levine | Universität Duisburg-Essen marc.levine@uni-due.de |
| Martin R Bridson | University of Oxford bridson@maths.ox.ac.uk | Ciprian Manolescu | University of California, Los Angeles cm@math.ucla.edu |
| Jim Bryan | University of British Columbia jbryan@math.ubc.ca | Haynes Miller | Massachusetts Institute of Technology hrm@math.mit.edu |
| Dmitri Burago | Pennsylvania State University burago@math.psu.edu | Tomasz Mrowka | Massachusetts Institute of Technology mrowka@math.mit.edu |
| Tobias H Colding | Massachusetts Institute of Technology colding@math.mit.edu | Aaron Naber | Northwestern University anaber@math.northwestern.edu |
| Simon Donaldson | Imperial College, London s.donaldson@ic.ac.uk | Peter Ozsváth | Princeton University petero@math.princeton.edu |
| Yasha Eliashberg | Stanford University eliash-gt@math.stanford.edu | Leonid Polterovich | Tel Aviv University polterov@post.tau.ac.il |
| Benson Farb | University of Chicago farb@math.uchicago.edu | Colin Rourke | University of Warwick gt@maths.warwick.ac.uk |
| David M Fisher | Rice University davidfisher@rice.edu | Roman Sauer | Karlsruhe Institute of Technology roman.sauer@kit.edu |
| Mike Freedman | Microsoft Research michaelf@microsoft.com | Stefan Schwede | Universität Bonn schwede@math.uni-bonn.de |
| David Gabai | Princeton University gabai@princeton.edu | Natasa Sesum | Rutgers University natasas@math.rutgers.edu |
| Stavros Garoufalidis | Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de | Gang Tian | Massachusetts Institute of Technology tian@math.mit.edu |
| Cameron Gordon | University of Texas gordon@math.utexas.edu | Ulrike Tillmann | Oxford University tillmann@maths.ox.ac.uk |
| Jesper Grodal | University of Copenhagen jg@math.ku.dk | Nathalie Wahl | University of Copenhagen wahl@math.ku.dk |
| Misha Gromov | IHÉS and NYU, Courant Institute gromov@ihes.fr | Anna Wienhard | Universität Heidelberg wienhard@mathi.uni-heidelberg.de |

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2024 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 28 Issue 7 (pages 3001–3510) 2024

| | |
|---|------|
| Holomorphic 1-forms on the moduli space of curves | 3001 |
| FILIPPO FRANCESCO FAVALE, GIAN PIETRO PIROLA and SARA TORELLI | |
| Two-dimensional metric spaces with curvature bounded above, I | 3023 |
| KOICHI NAGANO, TAKASHI SHIOYA and TAKAO YAMAGUCHI | |
| A nonexistence result for wing-like mean curvature flows in \mathbb{R}^4 | 3095 |
| KYEONGSU CHOI, ROBERT HASLHOFER and OR HERSHKOVITS | |
| Higgs bundles, harmonic maps and pleated surfaces | 3135 |
| ANDREAS OTT, JAN SWOBODA, RICHARD WENTWORTH and MICHAEL WOLF | |
| Multiple cover formulas for K3 geometries, wall-crossing, and Quot schemes | 3221 |
| GEORG OBERDIECK | |
| Ancient solutions to the Kähler Ricci flow | 3257 |
| YU LI | |
| CAT(0) 4-manifolds are Euclidean | 3285 |
| ALEXANDER LYTCHAK, KOICHI NAGANO and STEPHAN STADLER | |
| Gromov–Witten theory via roots and logarithms | 3309 |
| LUCA BATTISTELLA, NAVID NABIJOU and DHARUV RANGANATHAN | |
| 3-Manifolds without any embedding in symplectic 4-manifolds | 3357 |
| ALIAKBAR DAEMI, TYE LIDMAN and MIKE MILLER EISMEIER | |
| Orbit closures of unipotent flows for hyperbolic manifolds with Fuchsian ends | 3373 |
| MINJU LEE and HEE OH | |
| When does the zero fiber of the moment map have rational singularities? | 3475 |
| HANS-CHRISTIAN HERBIG, GERALD W SCHWARZ and CHRISTOPHER SEATON | |