

Geometry & Topology Volume 29 (2025)

Lower Ricci curvature and nonexistence of manifold structure

ERIK HUPP Aaron Naber Kai-Hsiang Wang





Lower Ricci curvature and nonexistence of manifold structure

Erik Hupp

AARON NABER KAI-HSIANG WANG

We build limit spaces $(M_j^n, g_j) \to (X^k, d)$ of manifolds M_j with uniform lower bounds on Ricci curvature such that X^k is nowhere a topological manifold, and in fact every open set $U \subseteq X$ has infinitely generated homology.

More completely, it is known that any such X^k must be k-rectifiable for some unique dim $X := k \le n = \dim M_j$. It is also known that if k = n, then X^n is a topological manifold on an open dense subset, and it has been an open question as to whether this holds for k < n. Consider now any smooth complete 4-manifold (X^4, h) with Ric > λ and $\lambda \in \mathbb{R}$. Then for each $\epsilon > 0$ we construct a complete 4-rectifiable metric space $(X_{\epsilon}^4, d_{\epsilon})$ with $d_{\text{GH}}(X_{\epsilon}^4, X^4) < \epsilon$ such that the following hold. First, X_{ϵ}^4 is a limit space $(M_j^6, g_j) \to X_{\epsilon}^4$, where M_j^6 are smooth manifolds satisfying the same lower Ricci bound Ric_j > λ . Additionally, X_{ϵ}^4 has no open subset which is topologically a manifold. Indeed, for any open $U \subseteq X_{\epsilon}^4$ we have that the second homology $H_2(U)$ is infinitely generated. Topologically, X_{ϵ}^4 is the connect sum of X^4 with an infinite number of densely spaced copies of $\mathbb{C} P^2$.

In this way we see that every 4-manifold X^4 may be approximated arbitrarily closely by 4-dimensional limit spaces X_{ϵ}^4 which are nowhere manifolds. We will see that there is a sense, as yet imprecise, in which generically one should expect manifold structures to not exist on spaces with higher-dimensional Ricci curvature lower bounds.

53B20, 53C21; 51F30, 53C23, 58C07

1.	Introduction	444
2.	Geometric outline of construction	446
3.	Preliminaries	454
4.	Step 1: the gluing block	456
5.	Step 2: adding conical singularities	464
References		476

^{© 2025} The Authors, under license to MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via Subscribe to Open.

1 Introduction

Let us begin with a historical discussion of measured Gromov Hausdorff limit spaces

(1)
$$(M_j^n, g_j, v_j, p_j) \to (X^k, d, \nu, p), \quad \operatorname{Ric}_j > \lambda, \quad \nu_j := \frac{dv_{g_j}}{\operatorname{Vol}(B_1(p_j))},$$

and their structure. That metric space limits even exist in this context was the result of Gromov [1999, Theorem 5.3]. Structural results for the limits X began in earnest with the almost rigidities of Cheeger and Colding [1996, Theorem 6.62]. Their almost splitting theorem allowed them to show that X was the union of rectifiable pieces of various dimensions [Cheeger and Colding 1997], and they conjectured that the dimension is locally constant and hence unique. Colding and Naber [2012] resolved this conjecture and showed that the limit X^k is k-rectifiable for a unique k. More recently, an example of Pan and Wei [2022] has shown that while X^k is k-rectifiable, its Hausdorff dimension might be larger than k. More specifically, it is possible for the singular part of X to have larger dimension with respect to the Hausdorff measure than it does with respect to the limit ν -measure.

In the context where (1) is noncollapsed, which is to say $Vol(B_1(p_j)) > v > 0$, one can say quite a bit more. In this case one has that k = n, and by volume convergence [Cheeger and Colding 1997, Theorem 5.9] the limit measure v is the *n*-dimensional Hausdorff measure on X. The starting point for a more refined analysis of X in the noncollapsed context is another almost rigidity of Cheeger and Colding [1996, Theorem 4.85]. This time one considers the monotone quantity

$$\theta_r(x) := \frac{\operatorname{Vol}(B_r(x))}{\omega_n r^n}$$

and shows that in the limit X it is a constant in r if and only if $B_r(x)$ is a metric cone. This opens the door to the techniques of Federer [1970], which have been applied to many nonlinear equations. In particular, one can decompose $X = \operatorname{Reg}(X) \cup \operatorname{Sing}(X)$ into a regular and singular part and stratify the singular part $\mathscr{P}^0(X) \subseteq \cdots \subseteq \mathscr{P}^{n-1}(X) = \operatorname{Sing}(X)$. Cheeger-Colding were able to then use the Federer dimension reduction to prove the dimensional estimates dim $\mathscr{P}^{\ell} \leq \ell$. More recently, the work of Cheeger, Jiang and Naber [Cheeger et al. 2021] was able to prove that $\mathscr{P}^{\ell}(X)$ is ℓ -rectifiable. This result is sharp for $\ell \leq n-2$ by an example of Li and Naber [2020], who built examples whose singular strata $\mathscr{P}^{\ell}(X)$ are ℓ -rectifiable, ℓ -cantor sets.

For the regular set Reg(X) of a noncollapsed limit one can say even more. A Reifenberg-type result of Cheeger and Colding [1997, Theorem 5.14] allows one to show that there is an open dense subset on which X is a topological manifold. In the case where the M_j are boundary free it was shown in [Cheeger and Colding 1997, Theorem 6.1] that X is a manifold away from a codimension 2 set. More recently, it was shown by Brué, Naber and Semola [2022] that even in the boundary case one has a manifold structure away from a codimension 2 set. That is, the top stratum of the singular set $\mathcal{G}^{n-1}(X)$ is itself a manifold away from a codimension 2 set, and thus X is a topological manifold with boundary away from a codimension 2 set.

1.1 Main result on topological structure

It has remained an open question as to whether in the collapsed case X needs to have a topological manifold structure on some open dense subset. The main result of this paper is to answer this question in the negative:

Theorem 1.1 Let (X^4, h) be a smooth complete manifold with $\operatorname{Ric}_X > \lambda$, where $\lambda \in \mathbb{R}$. Then for every $\epsilon > 0$ there exists a metric space $(X^4_{\epsilon}, d_{\epsilon})$ such that

- (i) $d_{\mathrm{GH}}(X^4, X_{\epsilon}) < \epsilon$,
- (ii) X_{ϵ}^4 is 4-rectifiable,
- (iii) there exist $(M_i^6, g_j) \xrightarrow{\text{GH}} (X_{\epsilon}^4, d)$ with $\operatorname{Ric}_{g_j} > \lambda$,
- (iv) for all open sets $U \subseteq X_{\epsilon}$, we have that the homology group $H_2(U)$ is infinitely generated. Consequently, every open set $U \subseteq X$ is noncontractible and therefore not homeomorphic to Euclidean space.

We will see that it is possible to build many such X_{ϵ}^4 . In short, for each countable dense subset $\mathscr{C} = \{x_j\} \subseteq X$ and each collection of sufficiently decaying constants $\epsilon \ge \epsilon_j \to 0$ we will build X_{ϵ} by connect-summing X^4 with a $\mathbb{C}P^2$ of size $\approx \epsilon_j$ at the collection of points \mathscr{C} . The topological picture will be similar to a complex algebraic blow up, where we replace a point x_j with a 2-sphere $S_{\epsilon_j}^2$, though it is important to note that this is purely topological and there is no complex structure being preserved in this process. In particular, the geometric properties of the *blow down* map that sends the newly added $S_{\epsilon_j}^2$ to the chosen x_j will be important to the construction. These blow down maps will explain how the rectifiable charts of the space X_{ϵ}^4 collapse each of our added 2-spheres to points, which form a set of measure zero. Most examples of rectifiable structures which are not manifolds arise by allowing for holes in the space. The blow down picture here explains the ability to build a rectifiable space which is nowhere a manifold, but also has no holes. By choosing these points $\{x_j\}$ and scales ϵ_j fairly freely we see that a smooth structure is actually quite hard to obtain under higher-dimensional lower Ricci curvature bounds, and in a certain generic sense we should expect limit spaces to not have manifold structures.

The above raises the question about whether if we assume bounds on topology we might obtain more. There are two versions of such a question:

Question 1.1 Let $(M_j^n, g_j, v_j, p_j) \xrightarrow{\text{mGH}} (X^k, d, v, p)$, where $\text{Ric}_j \ge \lambda$, $v_j := dv_{g_j}/\text{Vol}(B_1(p_j))$ and $|H_*(M_j, \mathbb{Z})| < A < \infty$. Then is X^k a topological manifold on an open dense subset?

Here, $|H_*(M_j, \mathbb{Z})|$ refers to the number of generators for the abelian group in question. We can phrase the question directly on the metric-measure space itself in a very similar form:

Question 1.2 If (X^k, d, ν) is an RCD (n, λ) space with bounded homology $|H_*(X, \mathbb{Z})| < \infty$, then is X a topological manifold on an open dense subset?

Rephrasing the above: is the nonmanifold structure in Theorem 1.1 only possible in the presence of infinitely generated homology?

Acknowledgements Naber was funded by National Science Foundation grant DMS-1809011 for much of this work.

2 Geometric outline of construction

Let us turn our attention to the construction of the smooth manifolds $(M_j^6, g_j) := (X_j^4 \times S^2, h_j + f_j^2 g_{S^2})$. We will often write $X_j \times_{f_j} S^2$ to represent that we are geometrically considering the product of two spaces with a warping factor f_j . We can view X_j^4 as the blow up of X^4 at an increasingly dense sequence of points. That is, to construct X_j^4 we will effectively take a collection of points $\{x_j^a\} \subseteq X^4$ and replace these points with 2-spheres. Each time we blow up a point, we are introducing a new noncontractible S^2 into the space. Geometrically, each S^2 being introduced will be of size at most ϵ , but their sizes will decrease quickly as the sequence continues. We will additionally alter the warping factor on the S^2 factor of M_j^6 in order to preserve the strict Ricci curvature lower bound. As our collection of blow up points becomes dense, we will arrive at our limit X_{ϵ}^4 .

Our construction will be inductive. That is, given M_j^6 which satisfies a handful of inductive properties, we will explicitly construct from it M_{j+1}^6 with similar inductive properties. Our goals in this section will first be to prove Theorem 1.1 given our inductive sequence, which we will do in Section 2.0.1. We will then focus on the inductive construction itself, which will be broken down into steps. Each step will consist of a main inductive lemma. The inductive lemmas will be proved later in the paper, but in the meantime we will finish the inductive construction in Section 2.3 based on these lemmas. As such, our goal for this section is to complete the proof of Theorem 1.1 modulo the proof of the inductive lemmas.

In order to state the conditions of our inductive construction, let us introduce the correct notion of regularity scale for this paper.

Definition 2.1 (regularity scale) Let $(M^n, g) = (X^{n-2} \times S^2, h + f^2 g_{S^2})$ be a smooth manifold with $x \in X$, and $0 < \eta < 1$ a constant. Then we define the regularity scale

(2)
$$r_x = r_x^{\eta} := \max_{0 < r \le 1} \left\{ \inf_{X(x) \ge r} \text{ and } \sup_{B_r(x)} \sum_{0 \le k \le 2} (r^{2+k} |\nabla^k \operatorname{Rm}_h| + r^{1+k} |\nabla^{1+k} \ln f|) \le \eta \right\}.$$

Remark 2.1 The definition depends on a constant $0 < \eta < 1$, though this constant may be fixed somewhat arbitrarily. Pictorially, when η is small we are increasingly close to $\mathbb{R}^{n-2} \times S^2$ with a product metric.

Remark 2.2 It follows that if $r_x > 2r$, then topologically $B_r(x) \subseteq X$ is contained in a Euclidean ball.

Remark 2.3 If $r_x > 2r$, then we can write the metric *h* in exponential coordinates on $B_r(x)$ so that h_{ab} is C^2 close to the identity and $\ln f$ is C^3 close to a constant. See Lemma 3.2.

Now let (X^4, h) be our chosen manifold from Theorem 1.1 with $\operatorname{Ric}_X > \lambda$. We will go ahead and assume X^4 is compact, but it will be clear that this is not needed as all constructions are purely local. Primarily, this allows us to discuss the construction in terms of some global parameters instead of choosing them locally. Let us define

(3)
$$\lambda^+ := \max\{\lambda : \operatorname{Ric}_X[v, v] \ge \lambda |v|^2\},$$

and observe that as X is compact we have $\lambda^+ > \lambda$. There is no loss in generality in us then assuming that the constant $\epsilon > 0$ from Theorem 1.1 satisfies

(4)
$$\epsilon < \frac{1}{2}(\lambda^+ - \lambda)$$

Our inductive construction of (M_j^6, g_j) will produce a choice of parameters

(5)
$$r_j := \frac{1}{2^j}, \quad \delta_j \le \delta^{1+j}, \quad \epsilon_j := \frac{\epsilon}{2^j} < \epsilon, \quad \lambda_j := \lambda^+ - \sum_{k=1}^J \epsilon_k > \lambda,$$

where $0 < \delta \ll 1$ will later be chosen sufficiently small. We will let our base step of the induction be represented by the space $M_0^6 := X^4 \times S^2$ with product metric $g_0 = h + \delta^2 g_{S^2}$. Our inductive assumptions will be the following:

- (I1) We can write $M_j^6 = X_j^4 \times S^2$, with metric $g_j = h_j + f_j^2 g_{S^2}$ satisfying $\operatorname{Ric}_{g_j} \ge \lambda_j$. The space (X_j^4, h_j) and warping function $f_j : X_j \to \mathbb{R}^+$ are smooth with $|f_j| < \delta_j r_j$.
- (I2) We have a smooth mapping $\phi_j : X_j \to X_{j-1}$ and a maximal disjoint collection $\{B_{r_i}(x_i^a) \subseteq X_j\}_a$ subject to the conditions
 - (I2.a) $\{x_i^a\}_a \subseteq \{x \in X_j : r_x \ge 4r_j\},\$

(I2.b)
$$\phi_{ji}(B_{4r_j}(x_i^a)) \cap \{x_i^b\} = \emptyset$$
, where $\phi_{ji} := \phi_{i+1} \circ \cdots \circ \phi_j : X_j \to X_i$ for $i < j$.

- (I3) The mapping $\phi_j : X_j \to X_{j-1}$ is a diffeomorphism away from $\phi_j^{-1}(x_{j-1}^a) \cong S^2$, an isometry away from $\phi_j^{-1}(B_{r_{j-1}}(x_{j-1}^a))$, and $\phi_j^{-1}(B_{r_{j-1}}(x_{j-1}^a))$ is diffeomorphic to the generating line bundle $E \to S^2$.
- (I4) $\phi_j^{-1}(x_{j-1}^a) \cong S^2$ are totally geodesic, round 2-spheres of radius $\leq \delta_j r_j$ in X_j . We have $|D\phi_j| < C(6)$, and if $x, y \in X_j$ with $d(x, y) > \delta_j r_j$ then $(1-\delta_j)d(x, y) < d(\phi_j(x), \phi_j(y)) < (1+\delta_j)d(x, y)$.

The notation C(6) above tells us that $|D\phi|$ is uniformly bounded by a dimensional n = 6 constant, which in this case is just a uniform constant.

Let us discuss some of the above properties and their implications. Condition (11) tells us that from a geometric standpoint M_i^6 is globally a warped product over X_i^4 , and that geometrically the S^2 factor is disappearing in the limit. Condition (I2) is an enumeration of the points we will be doing surgery around to move from X_i to X_{i+1} . The important point to observe is that necessarily this set is becoming increasingly dense by condition (I2.a), as the points are maximal subsets inside the set whose regularity scale is too large.

To move from X_j to X_{j+1} we will be performing surgeries on the balls $B_{r_j}(x_j^a)$. Condition (13) is telling us that the surgery is topologically a connect sum with $\mathbb{C}P^2$, where we are replacing each x_j^a with a 2-sphere. Near x_j^a this has the effect of replacing the diffeomorphic ball $B_{r_j}(x_j^a)$ with the total space $E \to S^2$ of the generating line bundle over S^2 . Note that the unit sphere bundle in E is S^3 , and hence this is the right object for gluing. From a topological perspective, moving from X_j to X_{j+1} adds a second homology generator for each x_j^a . Condition (12.b) is telling us that our surgeries never intersect the previously added 2-spheres.

Condition (I4) is explaining the geometric properties of our blow down maps $\phi_j : X_j \to X_{j-1}$. These will each be smooth mappings, indeed uniformly Lipschitz, and will be Riemannian isometries away from some small neighborhoods of the blow up points. We will see that the limit map $\Phi := \lim_{j\to\infty} \phi_{j1} : X_{\epsilon}^4 \to X^4$ is uniformly Hölder and locally bi-Lipschitz away from a set of measure zero. In particular we will have a single rectifiable chart of X_{ϵ}^4 over X^4 , that is an a.e. defined locally bi-Lipschitz map onto a full-measure subset of a smooth manifold. The blow up 2-spheres will all be collapsed to single points under this mapping.

2.0.1 Proving Theorem 1.1 given the inductive spaces M_j^6 Before focusing on the inductive construction itself, let us see how to use (I1)–(I4) in order to finish the proof of Theorem 1.1.

Let us begin by studying properties of the spaces X_j^4 . For each i < j let us write $\{S_{jia}^2\}_a := \{\phi_{ji}^{-1}(x_i^a)\}_a$. Notice by conditions (I4) and (I2.b) that these are disjoint totally geodesic round 2-spheres inside of X_j^4 . Additionally, by (I3) and a Mayer–Vietoris sequence we have that

(6)
$$\pi_1(X_j^4) = \pi_1(X^4) \quad \text{with } H_2(X_j^4) = H_2(X^4) \oplus \bigoplus_{i < j, a} \langle [S_{jia}^2] \rangle.$$

In particular, the rank of the second homology is growing in step with these two spheres.

Let us now flesh out the geometric properties implied by (I4) more completely. From (I4), we expect the Lipschitz constant to be uniformly bounded, but not necessarily close to 1. On the other hand, (I4) also tells us that the Lipschitz constant is close to 1 away from a small neighborhood of the diagonal in $X_i^4 \times X_i^4$. Consequently, we have the following bounds:

(7) If
$$x, y \in X_j$$
 with
$$\begin{cases} d(x, y) > \delta_j r_j, & \text{then } (1 - \delta_j) d(x, y) < d(\phi_j(x), \phi_j(y)) < (1 + \delta_j) d(x, y), \\ d(x, y) \le \delta_j r_j, & \text{then } d(\phi_j(x), \phi_j(y)) < Cd(x, y) \text{ for } C = C(6). \end{cases}$$

Thus we can take the Lipschitz constant to be small if the distance between two points is also not too small. This in particular implies the much weaker estimate

(8)
$$\phi_j$$
 is a $C\delta_j \leq C2^{-j}\delta$ -Gromov Hausdorff map.

By composing, we see that X_j^4 is always $C\delta < \epsilon$ -GH close to X^4 for small enough δ .

The importance of these estimates is that our real goal is to obtain uniform continuity for the maps $\phi_{ji}: X_j \to X_i$ from (I2.b). It is too much to ask that they be uniformly Lipschitz. However, for

449

 $\delta < \delta(C) = \delta(6)$ in the construction, condition (I4) will now allow us to show that the ϕ_{ji} are uniformly C^{α} maps. Indeed, for any fixed $x, y \in X_j$, write r := d(x, y), and then using (7) we can estimate

(9)
$$d(\phi_{ji}(x),\phi_{ji}(y)) \leq \prod_{k \leq j: r \leq \delta_k r_k} C \cdot \prod_{k \leq j: r > \delta_k r_k} (1+\delta_k) r.$$

To estimate the above we will use that $\delta_j \leq \delta^{1+j}$, which gives us that

(10)
$$d(\phi_{ji}(x), \phi_{ji}(y)) \le (1+\delta)C^{\ln(r)/\ln(\delta/2)}r = (1+\delta)r^{\alpha(\delta)},$$

where $\alpha(\delta) = 1 + \ln(C) / \ln(\delta/2) \to 1$ as $\delta \to 0$.

Now recall by (8) that ϕ_j is a $C\delta_j \leq C2^{-j}\delta$ -Gromov Hausdorff map. Consequently we have that $\phi_{ji}: X_j \to X_i$ is a $C\sum_{j\geq k\geq i+1} \delta_k \leq C2^{-i}\delta$ -Gromov Hausdorff map. This tells us that $\{X_j\}$ is a Cauchy sequence, and so

(11)
$$(X_j, d_j) \xrightarrow{\operatorname{GH}} (X_{\epsilon}, d)$$

Note that this is not a subsequential convergence but an actual convergence by the Cauchy condition. It follows from (I1) that $M_j \xrightarrow{\text{GH}} X_{\epsilon}^4$ as well, since $|f_j| \leq \delta_j r_j \to 0$. As the maps ϕ_{ji} witness the GH-Cauchy condition, we can take limits

(12)
$$\lim_{j \to \infty} \phi_{ji} := \Phi_i \colon X_{\epsilon}^4 \to X_i$$

where the Φ_i are also $C2^{-i}\delta$ -Gromov Hausdorff maps. It follows from (10) that the Φ_i are C^{α} -Hölder maps,¹ or more precisely that

(13)
$$d(\Phi_i(x), \Phi_i(y)) \le (1+\delta)d(x, y)^{\alpha}.$$

Importantly, we have that the Φ_i are continuous maps.

Now recall from (I3) and (I4) that $\{S_{jia}^2\} \subseteq X_j^4$ are totally geodesic round 2-spheres in X_j^4 . Note also that for i < j < k we have by (I2.b) and (I3) that $\phi_{kj}|_{S_{kia}^2}: S_{kia}^2 \to S_{jia}^2$ is an isometry. Consequently, we can limit our sequences of 2-spheres to get round 2-spheres

(14)
$$\lim_{j \to \infty} \{S_{jia}^2 \subseteq X_j\} = \{S_{ia}^2 \subseteq X_{\epsilon}^4\}.$$

Note that $\Phi_j|_{S_{ia}^2}: S_{ia}^2 \to S_{jia}^2$ is an isometry for every j > i; in particular, the radius of each 2-sphere S_{ia}^2 is at most $r_i \delta_i$. Now we claim that each $S_{ia}^2 \subseteq X_{\epsilon}^4$ is a nontrivial generator in the second homology group as well. Indeed, assuming this is not the case, there must be a continuous 3-chain $\psi: \Delta^3 \to X_{\epsilon}^4$ with boundary $\partial \psi = S_{ia}^2$. If we compose with Φ_j then this gives us a continuous chain $\Phi_j \circ \psi: \Delta^3 \to X_{\epsilon}^4$, whose boundary is the 2-sphere S_{jia}^2 . However, as we know that S_{jia}^2 is a nontrivial generator in the second homology group, this is not possible. A similar argument shows that each $\{S_{ia}^2\}$ generates an independent factor in the second homology group.

¹In fact, the Φ_i are rectifiable charts. Indeed, the ϕ_j are $(1 + \delta_j)$ -bi-Lipschitz away from the bubbles $\phi_j^{-1}(B_{\delta_{j-1}r_{j-1}}(x_{j-1}^a))$, and uniformly locally bi-Lipschitz away from the added 2-spheres. Composing these estimates as in (10) shows that for any j > 0, the Φ_i are C(i, j)-bi-Lipschitz away from $\bigcup_{k>i,a} B_{r_k\delta_j \wedge k}(S_{ka}^2)$.

Finally, let us show more carefully that the 2-spheres $\{S_{ia}^2\}_{i,a}$ are dense in X_{ϵ} . Fix any $x \in X_{\epsilon}$ and $\epsilon' > 0$. Consider $\Phi_i(x) \in X_i$ for some large *i* such that Φ_i is an ϵ' -GH map. We *claim* that we can find $x_j^a \in X_j$ for some $j \ge i$ such that $d(\phi_{ji}(x_j^a), \Phi_i(x)) < \epsilon'$. Since $\Phi_i = \phi_{ji} \circ \Phi_j$, it is then clear that the 2-sphere $S_{ja}^2 = \Phi_j^{-1}(x_j^a)$ is contained in the ball $B_{2\epsilon'}(x)$. To prove the claim, suppose it is not true. Since ϕ_{ji} is a $C2^{-i}\delta$ -Gromov Hausdorff map, we then have $x_j^a \notin B_{\epsilon'-C2^{-i}\delta}(\Phi_j(x))$ for any $j \ge i$ and any blow up point x_j^a . Thus the ball $B_{\epsilon'-C2^{-i}\delta-r_i}(\Phi_j(x))$ has the same Riemannian metric for all such *j*. In particular, $\Phi_j(x)$ will have the same regularity scale for all such *j* — note that the regularity scale is invariant under scaling of the warping function *f*. Now for *j* large enough, a point in $B_{\epsilon'-C2^{-i}\delta-r_i}(\Phi_j(x))$ has to be blown up since the collection of blow up points is maximal. This is a contradiction.

Thus we have our limit space X_{ϵ}^4 and a dense collection of two spheres $\{S_{ia}^2\}$ which are all generators in the second homology group, as claimed. This finishes the proof of Theorem 1.1 under the assumption that we have built our inductive sequence M_i^6 satisfying (I1)–(I4).

2.1 Step 1: the gluing block $\mathfrak{B}(\epsilon, \alpha, \delta)$

In order to prove Theorem 1.1 we are therefore left with showing how to build $M_{j+1} = X_{j+1} \times_{f_{j+1}} S^2$ from $M_j = X_j \times_{f_j} S^2$ in the inductive construction. The first step of the construction will build what is our main gluing block. When we move from X_j to X_{j+1} we will take our appropriately dense collection of points $\{x_j^a\}$ and replace a small neighborhood of each with our gluing block \mathcal{B} . From a topological perspective, it will be a connect sum with a copy of $\mathbb{C} P^2$ near each x_j^a , so that we are blowing up the points $\{x_i^a\}$ and replacing them with 2-spheres.

Our main constructive lemma in this step of the construction is the following:

Lemma 2.2 (inductive step 1) For every $0 < \epsilon < \frac{1}{10}$, $0 < \alpha < \alpha(\epsilon)$ and $0 < \delta < \delta(\epsilon, \alpha)$, there exists a smooth Riemannian manifold $\Re(\epsilon, \alpha, \delta)$ with $\operatorname{Rie}_{\Re} > 0$ and such that:

- (i) \mathfrak{B} is diffeomorphic to $E \times S^2$, where *E* is the total space of the generating line bundle $E \to S^2$. Further, (\mathfrak{B}, g) has a warped product structure $g := g_E + f^2 g_{S^2}$.
- (ii) There exists $U = U_E \times S^2 \subseteq \mathfrak{B}$ such that $\mathfrak{B} \setminus U$ is isometric to a neighborhood of infinity in $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ with the metric $dr^2 + (1-\epsilon)^2 r^2 g_{S^3} + \delta^2 r^{2\alpha} g_{S^2}$.

Further, there exists a $\phi: (E, g_E) \to C(S^3_{1-\epsilon})$ such that:

- (i) ϕ is a diffeomorphism away from the cone point $0 \in C(S^3)$, with $\phi^{-1}(0) \cong S^2$ an isometric sphere.
- (ii) $|D\phi| \leq C$ is uniformly bounded with ϕ an isometry away from U_E .

Note that $\mathfrak{B} \setminus U$ looks very pGH-close to $\mathbb{R}^4 \times S^2$, where the degree of closeness is being measured by ϵ , α and δ in a quantitative manner.

2.2 Step 2: adding cone singularities

The second step of the construction involves changing the geometry near each x_j^a so that the gluing blocks \mathcal{B} , which have a very specific geometry at infinity, may be isometrically glued along an annulus into X_i . Let us begin with a broader discussion before stating the main constructive lemma.

Let us start with a discussion of the density of singularities. It is known—see for instance [Otsu and Shioya 1994, Example 2]—that one can build examples (X, h) with $\operatorname{Ric}_h > 0$ for which the singular set is dense. In fact, the example in [loc. cit.] has positive sectional $\sec_h > 0$ and even full positive curvature operator $\operatorname{Rm}_h > 0$. What is at first counterintuitive is that the constructions of [loc. cit.] not only produce but essentially rely on these stronger curvature conditions. That is, the construction of singularities with $\operatorname{Rm}_h > 0$ in [loc. cit.] is very analogous to the construction of convex functions with nonsmooth points.

Let us now consider what is almost the reverse direction. Begin with a smooth space (X, h) with some form of lower bound on the curvature and ask about adding cone singularities near any point $x \in X$ without destroying the lower curvature bound. If for instance $\sec_h > \lambda > 0$, then one can accomplish this by performing a C^0 gluing in the spirit of [Perelman 1997, Section 4]. Namely, one can remove a sufficiently small neighborhood of $x \in X$ and isometrically glue in a rescaled spherical suspension of the boundary. There will be what is essentially distributional curvature added along the gluing, but the $\sec_h > \lambda$ assumption will allow us to guarantee that these distributional curvatures all have the right sign. In particular, one can smooth near the gluing region and preserve the $\sec_h > \lambda > 0$ condition.

The procedure described above of adding cone singularities near any point does not work if we are only assuming $\operatorname{Ric}_h > \lambda$. In short, the distributional curvature added from the C^0 gluing is due to the difference in second fundamental forms of the boundary on the two sides of the gluing. This second fundamental form in turn is closely related to sectional curvature, and Ricci curvature control is not sufficient. It turns out that we need to exploit better the local geometry near $x \in X$ in order to control the Ricci curvature.

In Step 2.1 of Section 5, we will see how to resolve this problem and add such cone singularities to arbitrary spaces satisfying $\operatorname{Ric}_h > \lambda$ without destroying the lower Ricci curvature bound.² The inductive lemma of this step of the construction is a generalization of this discussion and will allow us to also add conical singularities with a fixed warping structure near every point. This extra warping control is necessary in order to use the inductive lemma of Step 1 to add our desired topology. We will focus on the 6-dimensional case $M = X^4 \times S^2$ of interest, though it is clear that dimension is not a relevant constraint. Precisely, we have the following local gluing lemma, which focuses on a ball with controlled regularity scale:

Lemma 2.3 (inductive step 2) Consider a warped product space $(B_2^4(p) \times S^2, g)$, where the metric $g = g_B + f^2 g_{S^2}$ satisfies $inj_{g_B}(p) > 2$, $Ric_g > \lambda$ and

(15) $|\operatorname{Rm}_{g_B}|, |\nabla \operatorname{Rm}_{g_B}|, |\nabla^2 \operatorname{Rm}_{g_B}|, |\nabla \ln f|, |\nabla^2 \ln f|, |\nabla^3 \ln f| < \eta < 1.$

²The construction in Section 5.1 is a bit more general; one can drop the warping factor to obtain the claimed result.

Write $r := \text{dist}_{g_B}(\cdot, p)$. Then for all choices of parameters $0 < \epsilon < \epsilon(|\lambda|), 0 < \alpha < \alpha(\epsilon), 0 < \hat{r} < \hat{r}(\alpha, \lambda, \epsilon)$ and $0 < \hat{\delta} < \hat{\delta}(\lambda, \alpha, \epsilon, ||f||_{L^{\infty}}, \hat{r})$, there exists $0 < \delta = \delta(\hat{\delta}||f||_{\infty} | \alpha, \lambda, \epsilon)$ and a warped product metric $\hat{g} = \hat{g}_B + \hat{f}^2 g_{S^2}$ such that:

- (i) The Ricci lower bound $\operatorname{Ric}_{\widehat{g}} > \lambda C(4)\epsilon$ holds for $\widehat{r}/2 \leq r \leq 2$.
- (ii) $\hat{g} = g_B + \hat{\delta}^2 f^2 g_{S^2}$ is unchanged up to scaling the warping factor f by $\hat{\delta}$ for $1 \le r \le 2$.
- (iii) $\hat{g} = dr^2 + (1-\epsilon)^2 r^2 g_{S^3} + \delta^2 r^{2\alpha} g_{S^2}$ has the cone warping structure $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ for $r \le \hat{r}$.
- (iv) The identity map Id: $(B_2(p), \hat{g}_B) \rightarrow (B_2(p), g_B)$ is $(1 + 2\epsilon)$ -bi-Lipschitz.

Remark 2.4 Our notation of the constant dependence $\delta(\hat{\delta} \| f \|_{\infty} | \alpha, \lambda, \epsilon)$ means $\delta \to 0$ as $\hat{\delta} \| f \|_{\infty} \to 0$ with the other constants α , λ and ϵ fixed.

Remark 2.5 The caveat in (i) that $\operatorname{Ric}_{\widehat{g}} > \lambda - C(4)\epsilon$ only away from a small neighborhood of the cone point, eg $B_{\widehat{r}/2}(p) \times S^2$, cannot be improved. This is simply due to the fact that the fixed warping structure $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ has infinite negative curvature at the pole, in particular it has $\operatorname{Ric}|_{TS^2} \to -\infty$ as $r \to 0$. In practice, $B_{\widehat{r}/2}(p) \times S^2$ will be replaced by a rescaled bubble \mathfrak{B} obtained from Lemma 2.2, which does have the appropriate Ricci lower bound.

In practice the above works as follows. If $M = X \times_f S^2$ is a smooth manifold with a lower Ricci curvature bound and $x \in X$, then the above tells us we can find a potentially very small neighborhood $B_{2\rho}(x) \times S^2$ in which we can change the geometry of M. Specifically, after shrinking the warping factor by $\hat{\delta}$, we can alter the metric so that the ball $B_{\hat{r}\rho}(x) \times S^2$ will be isometric to that of the warped cone $C(S_{1-\epsilon}^3) \times_{\delta\rho(r/\rho)^{\alpha}} S^2$.

In the case of no warping factor, as per the discussion before the statement of the above lemma, we can repeat this process indefinitely in order to produce a dense set of singularities. In the case of a warping factor we can similarly repeat this process indefinitely, but also combine with the inductive lemma (Lemma 2.2) in order to glue topology in at each step. We will discuss this construction more carefully in the next step.

2.3 Step 3: constructing M_{j+1}

Let us now see how to use the inductive steps of Lemmas 2.2 and 2.3 in order to complete the inductive step of the construction and build M_{j+1} from M_j . Thus let us assume we have built $M_j = X_j \times S^2$ with $g_j = h_j + f_j^2 g_{S^2}$ and $\operatorname{Ric}_{g_j} > \lambda_j$. Let us also choose a maximal disjoint collection $\{B_{r_j}(x_j^a)\}$ such that $\{x_j^a\} \subseteq \{x \in X_j^4 : r_x \ge 4r_j\}$ and $\phi_{ji}(B_{4r_j}(x_j^a)) \cap \{x_i^b\} = \emptyset$.

Due to the unfortunate number of constants being accounted for in the construction, it is helpful to briefly remark on what will happen. The goal is to construct M_{j+1} in two steps. First we will take each ball $B_{r_j}(x_j^a) \subseteq X_j$ and apply the inductive step of Lemma 2.3 in order to add a warped cone singularity on $B_{\hat{r}r_j}(x_j^a)$. This space is not smooth at the cone point x_j^a of course, but will have appropriately positive Ricci at least outside of $B_{\hat{r}r_j/2}(x_j^a)$. We will then apply the inductive step of Lemma 2.2 to replace each warped

cone $B_{\hat{r}r_j}(x_j^a)$ with the smooth bubble metric $\hat{\mathcal{B}}$ of positive Ricci. This will produce X_{j+1} , and if we choose the various constants sufficiently small at each stage, we can do this while keeping control of both the space and its relationship to X_j . That is, we can show the inductive hypotheses (I1)–(I4) are satisfied.

Let us now describe the modifications to each disjoint ball $B_{2r_j}(x_j^a) \times S^2$ in more detail. It will be more convenient to describe the process on a single rescaled ball $B_2(p) \times S^2 := r_j^{-1}(B_{2r_j}(x_j^a) \times S^2)$ with metric $g = g_B + f^2 g_{S^2} := r_j^{-2}(h_j + f_j^2 g_{S^2}) = r_j^{-2} g_j$. Observe that the ball $B_2(p) \times S^2$ satisfies the criteria of Lemma 2.3, with Ricci lower bound $r_j^2 \lambda_j$. Therefore let us apply Lemma 2.3 with input parameters ϵ' , α , \hat{r} , $\hat{\delta}$, and output parameter δ_I . We choose ϵ' , α , \hat{r} and $\hat{\delta}$ to satisfy the conditions of Lemmas 2.3 and 2.2, but we may further shrink these constants later.

The previous application of Lemma 2.3 to $B_2(p) \times S^2$ produced a metric $\hat{g} = \hat{g}_B + \hat{f}^2 g_{S^2}$ such that $\hat{f} = \hat{\delta} f$ and $\hat{g}_B = g_B$ on the open set $A_{1,2}(p)$. Note additionally that by choosing $\epsilon' < \epsilon'(\epsilon_{j+1}, r_j)$ small enough, we can ensure that $\operatorname{Ric}_{\hat{g}} \ge r_j^2 \lambda_{j+1}$ holds away from $B_{\hat{r}/2}(p) \times S^2$. Inside the inner radius, $(B_{\hat{r}}(p) \times S^2, \hat{g})$ is isometric to the cone warping structure $B_{\hat{r}}(0) \times S^2 \subseteq C(S_{1-\epsilon'}^3) \times_{\delta_I r^\alpha} S^2$. We wish to replace this inner ball with a rescaled copy of the gluing model $\mathfrak{B}(\epsilon', \alpha, \delta(\epsilon', \alpha))$ from Lemma 2.2. Recall from Lemma 2.2 that $U \subseteq \mathfrak{B}$ is an open set such that $\mathfrak{B} \setminus U$ is isometric to $dr^2 + (1-\epsilon')^2 r^2 g_{S^3} + (\delta(\epsilon', \alpha))^2 r^{2\alpha} g_{S^2}$. Let $\hat{\mathfrak{B}}$ be a rescaling of \mathfrak{B} so that the rescaled core $\hat{U} \subseteq \hat{\mathfrak{B}}$ is contained in a bounded region with a collar neighborhood \hat{V} of its boundary isometric to $A_{\hat{r}/2,\hat{r}}(0) \times S^2 \subseteq C(S_{1-\epsilon'}^3) \times_{\delta_{II}r^\alpha} S^2$, for some $\delta_{II} = \delta_{II}(\hat{r}, \alpha, \epsilon')$. We additionally require that $\hat{r} < \hat{r}(|\lambda_{j+1}|, r_j)$ is small enough so that $\operatorname{Ric}_{\hat{\mathfrak{B}}} > r_j^2 \lambda_{j+1}$ in this region. In particular, if we multiply \hat{f} by $\min(\delta_I, \delta_{II})/\delta_I$ and multiply the warping factor of $\hat{\mathfrak{B}}$ by $\min(\delta_I, \delta_{II})/\delta_{II}$, then we can define a warped product $(\overline{\mathfrak{B}, \overline{g})$ from $(B_2(p) \times S^2, \widehat{g})$ by removing $B_{\hat{r}/2}(p) \times S^2$ and isometrically gluing the rescaled model $\hat{\mathfrak{B}}$ into $B_{\hat{r}}(p) \times S^2$ along \hat{V} . Note that for warped product geometries, multiplying the warping factor by a constant ≤ 1 only increases the Ricci curvature — see Remark 3.2 — so $\operatorname{Ric}_{\overline{g}} > r_j^2 \lambda_{j+1}$.

We now seek to replace the original ball $B_{2r_j}(x_j^a) \times S^2 \subseteq M_j$ with the rescaled $r_j\overline{\mathfrak{B}}$. By construction, $r_j\overline{\mathfrak{B}}$ is a warped product over a compact manifold with boundary, with a collar neighborhood of its boundary isometric to $(A_{r_j,2r_j}(p) \times S^2, h_j + (\overline{\delta}f_j)^2 g_{S^2})$, where $\overline{\delta} := \widehat{\delta} \min(\delta_I, \delta_{II})/\delta_I$. If we multiply the warping factor of M_j by $\overline{\delta}$ — which is independent of the choice of point in $\{x_j^a\}!$ — then we see that $B_{r_j}(x_j^a) \times S^2$ can be replaced by $r_j\overline{\mathfrak{B}}$, glued isometrically along $A_{r_j,2r_j}(p) \times S^2$. We similarly replace $B_{2r_j}(x_j^a) \times S^2$ for each other a to form (M_{j+1}, g_{j+1}) .

We have almost completed our construction. For any choices of $\epsilon' > 0$ and $\hat{r} > 0$ appropriately small our construction above holds, and we need only make choices. Observe that we can define $\phi_{j+1}: X_{j+1} \to X_j$ to be the identity outside of $\bigcup_a B_{r_j}(x_j^a)$. On the union of annuli $\bigcup_a (B_{r_j}(x_j^a) \setminus B_{\hat{r}r_j}(x_j^a))$ we have by Lemma 2.3 that $|D\phi_{j+1} - I| < C\epsilon'$ if we set $\phi_{j+1} :=$ Id setwise on this region. We can use the second part of Lemma 2.2 in order to extend ϕ_{j+1} to each glued bubble $r_j \overline{\mathfrak{B}}$, so that $|D\phi_{j+1}| < C(6)$ on $\{B_{\hat{r}r_j}(x_j^a)\}$. If we now choose $\epsilon' < \epsilon'(\delta_{j+1})$ and $\hat{r} < \delta_{j+1}/2$ sufficiently small, (7) holds. This finishes the construction of M_{j+1} .

3 Preliminaries

3.1 Ricci curvature of warping geometry

The underlying ansatz for all constructions going forward is the warped product with S^2 . We have several formulas for the curvature of such spaces which will be used in this paper. In this section we collect together some elementary remarks and formulas about such constructions. These will be the starting point for many of the other formulas computed in this paper.

Let us now be more precise: consider the data of a smooth Riemannian manifold (X, h) with a positive function $f: X \to (0, \infty)$. We can form the warped product of X with S^2 as follows:

(16)
$$X \times_f S^2 := (X \times S^2, h + f^2 g_{S^2}).$$

That is, $X \times_f S^2$ is a Riemannian manifold that topologically has the structure of a (trivial) sphere bundle $S^2 \to X \times_f S^2 \xrightarrow{\pi} X$. The fibers $\pi^{-1}(x) = S_{f(x)}^2$, for $x \in X$, of the projection map π are metrically round spheres of radius f, and are orthogonal to the natural sections $X \times \{\omega\}$ for $\omega \in S^2$. Denoting $M := X \times_f S^2$, we will therefore use the orthogonal splitting to make the identification $TM \cong TX \oplus TS^2$. We obtain the following concise formulas for the Ricci curvature of M in the complementary directions:

(17)
$$\operatorname{Ric}_{M}|_{TX} = \operatorname{Ric}_{h} - 2\frac{(\nabla_{h}^{2} f)}{f},$$

(18)
$$\operatorname{Ric}_{M}|_{TS^{2}} = \left(\frac{1}{f^{2}} - \frac{|\nabla_{h}f|_{h}^{2}}{f^{2}} - \frac{\Delta_{h}f}{f}\right) f^{2}g_{S^{2}}.$$

Let us make a couple of remarks about the general form of these identities, which we will use without further comment throughout the rest of the paper:

Remark 3.1 There is no cross-term in Ric_M , ie $\operatorname{Ric}_M(v, w) = 0$ for $v \in TX$ and $w \in TS^2$. In practice, this splitting of the Ricci curvature into orthogonal blocks will allow us to subdivide the problem of lower bounding Ric_M into two distinct steps.

Remark 3.2 The scaling action $f \mapsto \lambda f$ for $\lambda > 0$ leaves the *TX* directions $\operatorname{Ric}_M |_{TX}$ invariant, and acts on the TS^2 directions by $\operatorname{Ric}_M |_{TS^2} \mapsto \lambda^2 \operatorname{Ric}_M |_{TS^2} + (\lambda^{-2} - 1)\lambda^2 g_{S^2}$. Thus, Ric_M is nondecreasing under the scaling $f \mapsto \lambda f$ when $\lambda \leq 1$. In practice, this will mean that $\operatorname{Ric}_M |_{TS^2}$ can be made as large as desired, without disrupting $\operatorname{Ric}_M |_{TM}$, by multiplying f by a suitably small positive number.

3.2 C^1 gluing lemma for warping geometry

We state in this subsection a C^1 gluing lemma. It is a slight generalization of a result of Menguy [2000, Lemma 1.170], and its proof is essentially verbatim. We will make use of it in both steps of the construction.

Lemma 3.1 (C^1 warped geometry gluing) Let (M^n, h) be a smooth manifold with $N^{n-1} \subseteq M$ a complete smooth submanifold. Consider $(M^n \times S^k, g)$, where $g = h + f^2 g_{S^k}$ is a C^1 warped product metric. Assume further that

- (i) *h* and *f* are smooth on $M \setminus N$ and have smooth limits from each side of *N*,
- (ii) $\inf \operatorname{Ric}_g > \lambda \text{ on } (M \setminus N) \times S^k$.

Then for every open set $N \subseteq U$ and number $\epsilon > 0$, there exists a smooth metric $g_U = h_U + f_U^2 g_{S^k}$ on $M \times S^k$ such that $\operatorname{Ric}_{g_U} > \lambda$ and $g_U = g$ outside of $U \times S^k$. Moreover, one can arrange that $\|g - g_U\|_{C^1(U \times S^k)} < \epsilon$.

Remark 3.3 It is enough to assume that g is C^4 on $M \setminus N$.

Remark 3.4 The verbatim result is true for more general warping factors other than spheres.

3.3 Regularity under the exponential map

The following is relatively standard, and the proof goes through a series of Jacobi field estimates; however it is surprisingly difficult to find a precise reference for it. For the convenience of the reader we state the result precisely below:

Lemma 3.2 (regularity of exponential map) Let $0 < \eta < 1$ be a number and $(B_1(p), g)$ a metric ball which satisfies the regularity scale estimates

(19)
$$\operatorname{inj}(p) > 1 \quad \text{with } \sum_{\ell=0}^{k} \|\nabla^{\ell} \operatorname{Rm}\|_{L^{\infty}} < \eta.$$

Given an orthonormal basis $\{\partial_a\}$ of T_pM , let $g_{ab} = \exp^* g$ be the metric in exponential coordinates. Then we can estimate

(20)
$$\sum_{a,b} \left(r^{-2} \| g_{ab} - \delta_{ab} \|_{L^{\infty}} + \sum_{c} r^{-1} \| \partial_{c} g_{ab} \|_{L^{\infty}} + \sum_{\ell=2}^{k} \sum_{c_{1},...,c_{\ell}} \| \partial_{c_{1},...,c_{\ell}}^{\ell} g_{ab} \|_{L^{\infty}} \right) \leq C(n,k) \eta.$$

Remark 3.5 Let us say a few words about how these estimates can be proved. One rewrites the metric derivatives in terms of the Jacobi vector fields $J_a := r\partial_a$ along radial geodesics passing through p, eg $\partial_c g_{ab} = r^{-3}(g(\nabla_{J_c} J_a, J_b) + g(J_a, \nabla_{J_c} J_b))$ for $\partial_c \perp \nabla r$. To obtain estimates on the iterated covariant derivatives $J_{a_1,...,a_k,b} := \nabla_{J_{a_1}} \cdots \nabla_{J_{a_k}} J_b$, one uses the equation they solve:

$$0 = \nabla_{J_{a_1}} \cdots \nabla_{J_{a_k}} (J_b'' + R(J_b, \partial_r)\partial_r) = J_{a_1, \dots, a_k, b}'' + R(J_{a_1, \dots, a_k, b}, \partial_r)\partial_r + E_{a_1, \dots, a_k, b}.$$

The inhomogeneous term $E_{a_1,...,a_k,b}$ depends only on lower-order covariant derivatives $J_{c_1,...,c_\ell}J_d$ for $\ell < k$, which inductively have already been estimated, the base case $\ell = 0$ being the standard estimates for Jacobi vector fields as in [Jost 2017, Chapter 6.5]. One can then proceed to estimate solutions of this inhomogeneous ODE, eg by Duhamel's principle.

Remark 3.6 Let us briefly compare this regularity estimate and proof sketch with another possible approach. First, one switches to harmonic coordinates and uses the assumed injectivity radius and curvature bounds (19) to obtain $C^{k+1,\alpha}$ control on the metric in these coordinates for any $0 < \alpha < 1$, following [Anderson 1990]. Then, one converts this into $C^{k-1,\alpha}$ control on the metric in exponential coordinates by [DeTurck and Kazdan 1981, Theorem 2.1]. The loss of $(1 - \alpha)$ derivatives compared to Lemma 3.2 is immaterial in our context, where we are only ever working on the regularity scale in a smooth manifold.

Our primary use of the above will be to view the metric g as a form of twisted cone. Namely, in the context where inj(p) > 1 as above we can use exponential coordinates to write the metric g on $B_1(0) \subseteq C(S^{n-1}) = \mathbb{R}^n$ as

$$g = dr^2 + r^2 g_r$$

where g_r is a smooth family of metrics on S^{n-1} . The estimates above can then be understood as estimates on this family g_r :

Corollary 3.3 (cone regularity of exponential map) Let $(B_1(p), g)$ be a metric ball which satisfies the regularity scale estimates (19) with $0 < \eta < 1$. Let us use exponential coordinates to write $g = dr^2 + r^2g_r$ as in (21), where g_r is a family of metrics on S^{n-1} . Then we have the estimates

(22)
$$\sum_{\ell=0}^{k} \|\nabla_{g_{S^3}}^{\ell}(g_r - g_{S^3})\|_{L^{\infty}(S^3, g_{S^3})} \le C(n, k) \eta r^2,$$
$$\sum_{\ell=0}^{k-1} \|\nabla_{g_{S^3}}^{\ell}g_r'\|_{L^{\infty}(S^3, g_{S^3})} \le C(n, k) \eta r, \qquad \sum_{\ell=0}^{k-2} \|\nabla_{g_{S^3}}^{\ell}g_r''\|_{L^{\infty}(S^3, g_{S^3})} \le C(n, k) \eta.$$

4 Step 1: the gluing block

In this section we build the gluing block of Step 1 for our construction. Our 4-manifold of interest for this gluing block is the generating line bundle $E \to S^2$, which we can topologically also view as \mathbb{C}^2 blown-up at the origin. We will build a metric of positive Ricci curvature on $E \times S^2$ which has the property that at infinity it looks roughly like $\mathbb{R}^4 \times S^2$. More precisely, it will be isometric to the warped product $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ near infinity. The precise lemma is the following:

Lemma 4.1 (inductive step 1) For every $0 < \epsilon < \frac{1}{10}$, $0 < \alpha < \alpha(\epsilon)$ and $0 < \delta < \delta(\epsilon, \alpha)$, there exists a smooth Riemannian manifold $\Re(\epsilon, \alpha, \delta)$ with $\operatorname{Ric}_{\Re} > 0$ and such that:

- (i) \mathfrak{B} is diffeomorphic to $E \times S^2$, where *E* is the total space of the generating line bundle $E \to S^2$. Further, (\mathfrak{B}, g) has a warped product structure $g := g_E + f^2 g_{S^2}$ for some smooth $f : E \to \mathbb{R}^+$.
- (ii) There exists a $U = U_E \times S^2 \subseteq \Re$ such that $\Re \setminus U$ is isometric to a neighborhood of ∞ in $C(S^3_{1-\epsilon}) \times_{\delta r^{\alpha}} S^2$ with the metric $dr^2 + (1-\epsilon)^2 r^2 g_{S^3} + \delta^2 r^{2\alpha} g_{S^2}$.

Further, there exists a $\phi: (E, g_E) \to C(S^3_{1-\epsilon})$ such that:

- (i) ϕ is a diffeomorphism away from the cone point $0 \in C(S_{1-\epsilon}^3)$, with $\phi^{-1}(0) \cong S^2$ an isometric sphere.
- (ii) $|D\phi| \leq C$ is uniformly bounded with ϕ an isometry away from U_E .

Remark 4.1 As usual our use of the notation $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ means we are looking at the warped product metric on $C(S^3) \times S^2$ given by $g := dr^2 + (1-\epsilon)^2 r^2 g_{S^3} + \delta^2 r^{2\alpha} g_{S^2}$.

The proof of the above lemma is broken down over the remainder of this Section. In Step 1.1 of Section 4.1 we begin by writing down a metric on $(E, g_{E,1})$ with nonnegative Ricci curvature, and which looks like a cone at infinity. This cone however may not be close to \mathbb{R}^4 at this stage.

In Step 1.2 of Section 4.2 we will write down a metric on $E \times S^2$ of the form $g_2 = g_{E,2} + f_2^2 g_{S^2}$. The base metric $g_{E,2} := g_{E,1}$ will simply be the metric from the first step, however we will now equip the metric with a warping S^2 factor $f_2(r) := \delta_2(1 + r^2)^{\alpha_2/2}$. The polynomial growth of the warping factor will add extra curvature which will be useful in flattening out the cone in the third step.

In Step 1.3 of Section 4.3 we will use the extra curvature provided by the warping factor to slowly increase the cone angle of g_E until it is close to Euclidean. In Step 1.4 we will fix the warping factor so that our space becomes isometric to the warped cone $C(S_{1-\epsilon}^3) \times_{\delta r^{\alpha}} S^2$ near infinity. At several steps we will only build geometries which are globally C^1 , but we will end the construction of Lemma 4.1 by applying the C^1 smoothing Lemma 3.1 in order to fix this issue.

4.1 Step 1.1: bubble metric with positive Ricci

Consider the generating line bundle $E \to S^2$, and let us observe that the unit sphere bundle is diffeomorphic to S^3 . In particular, if we remove the zero section then $E \setminus S^2$ is diffeomorphic to $\mathbb{R}^+ \times S^3$. We will begin by writing our metric in this degenerate coordinate system. To do so let us choose the canonical left-invariant vector fields X, Y, Z on S^3 so that they satisfy the commutator relations [X, Y] = 2Z, [Y, Z] = 2X and [Z, X] = 2Y. Let dX, dY and dZ denote the dual frames. We first consider a metric on $\mathbb{R}^+ \times S^3$ of the form

(23)
$$g_{E,1} := dr^2 + A(r)^2 dX^2 + B(r)^2 (dY^2 + dZ^2).$$

In order for this to define a smooth metric on E it is required that A(0) = 0 with $A^{(\text{even})}(0) = 0$ and B(0) > 0 with $B^{(\text{odd})} = 0$. Our construction for this step is the following:

Lemma 4.2 Let $g_{E,1}$ be as in (23), and let $0 < m < \frac{1}{100}$ with $r_1 := 2$. Then there exist A(r) and B(r) such that

- (i) A(r) = B(r) with A'(r) = B'(r) = m for $r \ge r_1$,
- (ii) $g_{E,1}$ defines a C^1 metric on E, smooth away from $r = r_1$, with Ric ≥ 0 on the smooth part, and Ric $\ge \frac{1}{2}k^2 > 0$ on $U_{r_1} := \{r < r_1\}$ for some constant k = k(m).

Remark 4.2 For $r \ge r_1$, we have that $g_{E,1} = dr^2 + A(r)^2 g_{S_3}$ is exactly the cone metric on $C(S_m^3)$.

Proof of Lemma 4.2 Let k > 0 be the smallest number satisfying the relation

$$(24) m = \cos(kr_1).$$

Clearly we then have $\frac{1}{3}\pi \le kr_1 < \frac{1}{2}\pi$. Let us define A(r) by the formula

(25)
$$A(r) = \begin{cases} \frac{1}{k} \sin(kr) & \text{if } r \le r_1, \\ \frac{\sqrt{1-m^2}}{k} + m(r-r_1) & \text{if } r \ge r_1. \end{cases}$$

To define B(r) let $b = b(m) < \sqrt{1 - m^2}/k$ be taken so that we can find a smooth function with

(26)
$$B(r) = \begin{cases} b & \text{if } r \leq \frac{1}{2}r_1, \\ 0 \leq B'' \leq \frac{4m}{r_1} & \text{if } \frac{1}{2}r_1 \leq r \leq r_1, \\ \frac{\sqrt{1-m^2}}{k} + m(r-r_1) & \text{if } r \geq r_1. \end{cases}$$

Note that in this case we necessarily have $0 \le B' \le m \le A'$ and $B \ge A$. Together with $kr_1 < \frac{1}{2}\pi$ and $m < \frac{1}{100}$, we can estimate

(27)
$$b = B\left(\frac{1}{2}r_1\right) \ge B(r_1) - m\left(r_1 - \frac{1}{2}r_1\right) \ge \frac{1}{k}\left(\sqrt{1 - m^2} - \frac{1}{4}m\pi\right) > \frac{1}{2k}.$$

Observe from the behavior of A and B as $r \to 0^+$ that $g_{E,1}$ indeed defines a smooth metric on E near the zero section; see eg [Perelman 1997].

Let us now estimate the Ricci curvatures. For $r < \frac{1}{2}r_1$, we have (note that B' = 0 = B'' here)

(28)

$$\operatorname{Ric}(\partial_{r}, \partial_{r}) = -\frac{A''}{A} - 2\frac{B''}{B} = k^{2},$$

$$\operatorname{Ric}\left(\frac{X}{|X|}, \frac{X}{|X|}\right) = -\frac{A''}{A} - 2\frac{A'B'}{AB} + 2\frac{A^{2}}{B^{4}} \ge k^{2},$$

$$\operatorname{Ric}\left(\frac{Y}{|Y|}, \frac{Y}{|Y|}\right) = \operatorname{Ric}\left(\frac{Z}{|Z|}, \frac{Z}{|Z|}\right) = -\frac{B''}{B} - \frac{A'B'}{AB} - \left(\frac{B'}{B}\right)^{2} + \frac{2}{B^{2}} + \frac{2B^{2} - 2A^{2}}{B^{4}}$$

$$\ge \frac{2}{B(r_{1})^{2}} = \frac{2k^{2}}{1 - m^{2}},$$

which is clearly appropriately positive. For $\frac{1}{2}r_1 \le r < r_1$, we can estimate

(29)
$$\operatorname{Ric}(\partial_{r}, \partial_{r}) = -\frac{A''}{A} - 2\frac{B''}{B} \ge k^{2} - 2\frac{4m/r_{1}}{1/2k} \ge k^{2}\left(1 - \frac{48m}{\pi}\right),$$
$$\operatorname{Ric}\left(\frac{X}{|X|}, \frac{X}{|X|}\right) = -\frac{A''}{A} - 2\frac{A'B'}{AB} + 2\frac{A^{2}}{B^{4}} \ge k^{2} - 2k\operatorname{cot}(kr_{1}/2)\frac{m}{1/2k} + \frac{2k^{2}\sin^{2}(kr_{1}/2)}{(1 - m^{2})^{2}} \ge k^{2}(1 - 4\sqrt{3}m + \frac{1}{2(1 - m^{2})^{2}}),$$

Lower Ricci curvature and nonexistence of manifold structure

$$\operatorname{Ric}\left(\frac{Y}{|Y|}, \frac{Y}{|Y|}\right) = \operatorname{Ric}\left(\frac{Z}{|Z|}, \frac{Z}{|Z|}\right) = -\frac{B''}{B} - \frac{A'B'}{AB} - \left(\frac{B'}{B}\right)^2 + \frac{2}{B^2} + \frac{2B^2 - 2A^2}{B^4}$$
$$\geq -\frac{4m/r_1}{1/2k} - k \cot(kr_1/2)\frac{m}{1/2k} - \left(\frac{m}{1/2k}\right)^2 + \frac{2k^2}{1 - m^2}$$
$$\geq k^2 \left(\frac{2}{1 - m^2} - \frac{24m}{\pi} - 2\sqrt{3}m - 4m^2\right).$$

Again observe that for $m < \frac{1}{100}$ we have the appropriate positivity. Finally for $r > r_1$, we have (note that A = B is affine here)

(30)
$$\operatorname{Ric}(\partial_r, \partial_r) = 0, \quad \operatorname{Ric}\left(\frac{X}{|X|}, \frac{X}{|X|}\right) = 2\frac{1 - (A')^2}{A^2} = 2\frac{1 - m^2}{A^2}.$$

Since $m < \frac{1}{100}$ we have Ric ≥ 0 for $r > r_1$. This completes the construction.

4.2 Step 1.2: bubble metric with S^2 warping factor

Recall that we ended the last step by constructing a metric $g_{E,1}$ on $E \to S^2$ which has nonnegative Ricci curvature and is a cone $C(S_m^3)$ outside a compact subset. The sphere S_m^3 in this cone is, however, potentially quite small, and we will want to take the radius of this sphere closer to 1 in order to geometrically flatten out the space.

In this next step of the construction, we want to add to $g_{E,1}$ a warped S^2 factor. This factor will add additional curvature to the radial directions, which will be used in subsequent sections to flatten out our cone structure. In this step we will look for a metric of the form

(31)
$$g_2 := g_{E,2} + f_2(r)^2 g_{S^2} = g_{E,1} + f_2(r)^2 g_{S^2}.$$

In particular, we will not change the metric on our base E in this step. The warping factor $f_2(r)$ will be given explicitly by

(32)
$$f_2(r) = \delta_2 (1 + r^2)_2^{\alpha} / 2$$

Our main purpose in this step is to see that g_2 always has positive Ricci curvature:

Lemma 4.3 Let g_2 be as in (31) and (32). Then for any $0 < \alpha_2 \le \alpha_2(m) \le \frac{1}{2}$ and $0 < \delta_2 < 1$,

(33)
$$g_2 = g_{E,2} + f_2^2 g_{S^2} = dr^2 + A(r)^2 dX^2 + B(r)^2 (dY^2 + dZ^2) + f_2(r)^2 g_{S^2}$$

defines a C^1 metric on $E \times S^2$, smooth away from $r = r_1$ with Ric > 0 on the smooth part.

Proof Note first that as $f_2^{\text{(odd)}}(0) = 0$, the metric is smooth near r = 0. Calculations together with the results from the previous subsection give that for $r < r_1$,

$$(34) \quad \frac{f_2'}{f_2} = \frac{\alpha_2 r}{1+r^2} \le \frac{\alpha_2}{2}, \qquad \qquad \frac{f_2''}{f_2} = \frac{\alpha_2}{1+r^2} (1 + \frac{(\alpha_2 - 2)r^2}{1+r^2}) \le \alpha_2, \\ \frac{A'f_2'}{Af_2} = \frac{k\cos(kr)\alpha_2 r}{\sin(kr)(1+r^2)} \le \frac{\alpha_2\cos(kr)}{1+r^2} \frac{\pi}{2} \le \frac{\pi}{2}\alpha_2, \qquad \frac{B'f_2'}{Bf_2} \le \frac{m}{B(\frac{1}{2}r_1)} \frac{\alpha_2}{2} \le mk\alpha_2.$$

Geometry & Topology, Volume 29 (2025)

 \Box

We use the subscript index α for coordinate directions on S^2 that are orthonormal for g_{S^2} (and diagonalize the corresponding block of Ric) at the point of calculation. We now estimate the Ricci curvature of the whole space for $r < r_1$ (with Lemma 4.2):

(35)

$$\operatorname{Ric}(\partial_{r},\partial_{r}) = -\frac{A''}{A} - 2\frac{B''}{B} - 2\frac{f_{2}''}{f_{2}} \ge \frac{1}{2}k^{2} - 2\alpha_{2},$$

$$\operatorname{Ric}\left(\frac{X}{|X|},\frac{X}{|X|}\right) = -\frac{A''}{A} - 2\frac{A'B'}{AB} + 2\frac{A^{2}}{B^{4}} - 2\frac{A'f_{2}'}{Af_{2}} \ge \frac{1}{2}k^{2} - \alpha_{2}\pi,$$

$$\operatorname{Ric}\left(\frac{Y}{|Y|},\frac{Y}{|Y|}\right) = \operatorname{Ric}\left(\frac{Z}{|Z|},\frac{Z}{|Z|}\right) = -\frac{B''}{B} - \frac{A'B'}{AB} - \left(\frac{B'}{B}\right)^{2} + 2\frac{2B^{2} - A^{2}}{B^{4}} - 2\frac{B'f_{2}'}{Bf_{2}}$$

$$\ge \frac{1}{2}k^{2} - 2mk\alpha_{2},$$

$$f_{2}^{-2}\operatorname{Ric}(\partial_{\alpha},\partial_{\alpha}) = \frac{1}{f_{2}^{2}} - \frac{f_{2}''}{f_{2}} - \frac{(f_{2}')^{2}}{f_{2}^{2}} - \frac{A'f_{2}'}{Af_{2}} - 2\frac{B'f_{2}'}{Bf_{2}}$$

$$\ge \frac{1}{\delta_{2}^{2}(1+r_{1}^{2})^{\alpha_{2}}} - \alpha_{2} - \frac{1}{4}\alpha_{2}^{2} - \frac{1}{2}\pi\alpha_{2} - 2mk\alpha_{2}.$$

Observe for any $0 < \alpha_2 \le \alpha_2(m)$ and any $0 < \delta_2 < 1$ that we have Ric > 0. For $r > r_1 = 2$ we use different estimates (recall that A = B and they are affine here, and we impose $\alpha_2 \le \frac{1}{2}$):

(36)
$$\frac{f_2'}{f_2} = \frac{\alpha_2 r}{1+r^2},$$
$$\frac{f_2''}{f_2} = \frac{\alpha_2}{1+r^2} \left(1 + \frac{(\alpha_2 - 2)r^2}{1+r^2}\right) \le \frac{\alpha_2}{1+r^2} \left(1 + \frac{(\alpha_2 - 2)r_1^2}{1+r_1^2}\right) \le \frac{-\alpha_2}{5(1+r^2)},$$
$$\frac{A'f_2'}{Af_2} = \frac{m\alpha_2 r}{A(1+r^2)}.$$

Thus we can estimate the Ricci curvatures for $r > r_1$:

(37)
$$\operatorname{Ric}(\partial_{r},\partial_{r}) = -3\frac{A''}{A} - 2\frac{f_{2}''}{f_{2}} > 0,$$
$$\operatorname{Ric}\left(\frac{X}{|X|},\frac{X}{|X|}\right) = -\frac{A''}{A} + 2\frac{1-(A')^{2}}{A^{2}} - 2\frac{A'f_{2}'}{Af_{2}} = 2\frac{1-m^{2}}{A^{2}} - 2\frac{m\alpha_{2}r}{A(1+r^{2})}$$
$$\geq \frac{2}{Ar}(kr_{1}\sqrt{1-m^{2}}-m\alpha_{2}),$$
$$f_{2}^{-2}\operatorname{Ric}(\partial_{\alpha},\partial_{\alpha}) = \frac{1}{f_{2}^{2}} - \frac{f_{2}''}{f_{2}} - \frac{(f_{2}')^{2}}{f_{2}^{2}} - 3\frac{A'f_{2}'}{Af_{2}}$$
$$\geq \frac{1}{\delta_{2}^{2}(1+r^{2})^{\alpha_{2}}} + \frac{\alpha_{2}}{5(1+r^{2})} - \left(\frac{\alpha_{2}r}{1+r^{2}}\right)^{2} - 3\frac{m\alpha_{2}r}{A(1+r^{2})}$$
$$\geq \frac{1}{1+r^{2}}\left(\frac{1}{\delta_{2}^{2}} + \frac{\alpha_{2}}{5} - \alpha_{2}^{2} - 3\alpha_{2}\right).$$

Thus we again have, for any $0 < \alpha_2 \le \alpha_2(m)$ and $0 < \delta_2 < 1$, that Ric > 0.

Geometry & Topology, Volume 29 (2025)

4.3 Step 1.3: flattening the cone

Our goal in this step of the construction is to look for a metric on $E \times S^2$ of the form

(38)
$$g_3 := g_{E,3} + f_3(r)^2 g_{S^2}$$

where the warping factor $f_3(r) := f_2(r) = \delta_2(1+r^2)^{\alpha_2/2}$ will remain unchanged, up to further restrictions on the parameters δ_2 and α_2 . The base metric $g_{E,3}$ should look like a flat cone $C(S_{1-\epsilon}^3)$ outside some large radius, and will more generally satisfy

(39)
$$g_{E,3} = \begin{cases} g_{E,2} & \text{if } r \le r_1, \\ dr^2 + h_3(r)^2 g_{S^3} & \text{if } r \ge r_1, \end{cases}$$

where the warping function h_3 will be smooth on $[r_1, \infty)$ and satisfy

(40)
$$h_3(r_1) = A(r_1), \quad h'_3(r_1) = m, \quad \begin{cases} 0 \le r h''_3 \le 10/\ln r_3 & \text{if } r \in [r_1, r_3], \\ h'_3(r) = 1 - \epsilon & \text{if } r \ge r_3. \end{cases}$$

The following lemma will tell us that for r_3 sufficiently large, the resulting metric will have positive Ricci curvature:

Lemma 4.4 Let g_3 satisfy (38)–(40) with $\alpha_2 < \alpha_2(\epsilon, m) \le \frac{1}{2}$, $r_3 \ge r_3(m, \alpha_2, \epsilon)$ and $\delta_2 < \delta_2(m)$. Then g_3 is smooth away from $r = r_1$, globally C^1 , and satisfies Ric > 0 on the smooth region.

Proof We will focus our computations in the range $r \in [r_1, r_3]$, as in the range $r \in [r_3, \infty)$ the metric $g_{E,3}$ is again conic and the estimate will be similar as in the previous subsection. We also introduce the subscript index *i* for coordinate directions on S^3 that are orthonormal for g_{S^3} (and diagonalize the corresponding block of Ric) at the point of calculation. Let us begin by computing the Ricci curvature of the ansatz $g_3 = dr^2 + h_3(r)^2 g_{S^3} + f_3(r)^2 g_{S^2}$ as:

(41)

$$\operatorname{Ric}_{rr} = -3\frac{h_3''}{h_3} - 2\frac{f_3''}{f_3},$$

$$h_3^{-2}\operatorname{Ric}_{ii} = 2\frac{1 - (h_3')^2}{h_3^2} - \frac{h_3''}{h_3} - 2\frac{h_3'}{h_3}\frac{f_3'}{f_3},$$

$$f_3^{-2}\operatorname{Ric}_{\alpha\alpha} = \frac{1 - (f_3')^2}{f_3^2} - \frac{f_3''}{f_3} - 3\frac{f_3'}{f_3}\frac{h_3'}{h_3}.$$

Let us impose the restriction $\alpha_2 \leq \frac{1}{2}$ and calculate $mr_1 \leq A(r_1) \leq \frac{9}{10}r_1 \leq (1-\epsilon)r_1$. Then in the range $r \in [r_1, r_3]$, let us observe the estimates

$$(42) \qquad \frac{f_3'}{f_3}\frac{h_3'}{h_3} \le \frac{\alpha_2 r}{1+r^2}\frac{1-\epsilon}{m(r-r_1)+A(r_1)} \le \frac{\alpha_2}{m}\frac{1}{r^2},\\ \frac{1-(h_3')^2}{(h_3)^2} \ge \frac{1-(1-\epsilon)^2}{((1-\epsilon)(r-r_1)+A(r_1))^2} \ge \frac{\epsilon}{r^2},\\ \frac{1-(f_3')^2}{(f_3)^2} = \frac{1}{\delta_2^2(1+r^2)^{\alpha_2}} - \frac{\alpha_2^2 r^2}{(1+r^2)^2} \ge \frac{1}{\delta_2^2 r^{2\alpha_2}} \left(\frac{r^2}{1+r^2}\right)^{\alpha_2} - \frac{1}{4r^2} \ge \frac{1}{2\delta_2^2 r^{2\alpha_2}} \ge \frac{1}{2\delta_2^2 r^2},$$

Erik Hupp, Aaron Naber and Kai-Hsiang Wang

$$\left|\frac{h_3''}{h_3}\right| \le \frac{10}{m\ln r_3} \frac{1}{r^2},$$

$$\frac{f_3''}{f_3} = \frac{\alpha_2}{1+r^2} \left(1 + \frac{(\alpha_2 - 2)r^2}{1+r^2}\right) \le \frac{-\alpha_2}{5(1+r^2)} \le -\frac{4\alpha_2}{25} \frac{1}{r^2}.$$

If we plug these estimates into (41) then we arrive at

(43)

$$\operatorname{Ric}_{rr} \geq \left(\frac{8\alpha_{2}}{25} - \frac{30}{m\ln r_{3}}\right)\frac{1}{r^{2}},$$

$$h_{3}^{-2}\operatorname{Ric}_{ii} \geq \left(2\epsilon - \frac{10}{m\ln r_{3}} - \frac{2\alpha_{2}}{m}\right)\frac{1}{r^{2}},$$

$$f_{3}^{-2}\operatorname{Ric}_{\alpha\alpha} \geq \frac{1}{2\delta_{2}^{2}r} + \left(\frac{4}{25} - \frac{3}{m}\right)\frac{\alpha_{2}}{r^{2}} \geq \frac{1}{2\delta_{2}^{2}r} + \frac{1}{2}\left(\frac{4}{25} - \frac{3}{m}\right)\frac{1}{r}.$$

It follows from the first inequality that if $r_3 \ge r_3(m, \alpha_2)$ then $\operatorname{Ric}_{rr} > 0$. It follows from the second inequality that if $\alpha_2 \le \alpha_2(m, \epsilon)$ and $r_3 \ge r_3(m, \epsilon)$ then $\operatorname{Ric}_{ii} > 0$. Finally we see from the last equation that if $\delta_2 \le \delta_2(m)$ that $\operatorname{Ric}_{\alpha\alpha} > 0$.

4.4 Step 1.4: the warped cone metric

In the last step of the construction we have built a global metric g_3 on $E \times S^2$ such that outside the compact set $U_3 := \{r \le r_3\}$, the metric g_3 can be written as

(44)
$$g_3 = dr^2 + (1-\epsilon)^2 (r-R_3)^2 g_{S^3} + \delta_2^2 (1+r^2)^{\alpha_2} g_{S^2},$$

where R_3 solves $h_3(r_3) = (1-\epsilon)(r_3 - R_3)$. Observe that $R_3 > 0$ under our assumptions of the parameters. Our goal in this step of the construction is to build a metric g_4 which agrees with g_3 for $r \le r_3$, but for $r \ge r_3$ should take the form

(45)
$$g_4 = dr^2 + (1-\epsilon)^2 (r-R_3)^2 g_{S^3} + f_4(r)^2 g_{S^2},$$

where our warping factor satisfies

(46)
$$f_4(r) = \begin{cases} f_3(r) & \text{if } r \le r_3, \\ \delta(r - R_3)^{\alpha} & \text{if } r \ge r_3. \end{cases}$$

We want our warping function to be globally C^1 , and thus we will choose the constants

(47)
$$\alpha := \alpha_2 \frac{r_3 \frac{n_3(r_3)}{1 - \epsilon}}{1 + r_3^2} < \alpha_2 \quad \text{and} \quad \delta := \delta_2 \frac{(1 + r_3^2)^{\frac{1}{2}\alpha_2}}{\left(\frac{h_3(r_3)}{1 - \epsilon}\right)^{\alpha}}.$$

Our final lemma is that for α_2 and δ_2 sufficiently small, our new metric g_4 has positive Ricci curvature:

Lemma 4.5 Let g_4 satisfy (45) and (46). If we further choose $\alpha_2 \le \alpha_2(\epsilon)$ and $\delta_2 \le \delta_2(\alpha_2)$, then for $r > r_3$ we have that Ric > 0.

Geometry & Topology, Volume 29 (2025)

462

Remark 4.3 After the change of coordinates $t := r - R_3$, the metric g_4 above becomes the desired format

(48)
$$dt^{2} + ((1-\epsilon)t)^{2}g_{S^{3}} + (\delta t^{\alpha})^{2}g_{S^{2}} =: dt^{2} + h(t)^{2}g_{S^{3}} + f(t)^{2}g_{S^{2}},$$

as in Lemma 4.1.

Proof The range $r > r_3$ corresponds exactly to $t > h_3(r_3)/(1-\epsilon)$, and in these new coordinates we can compute the Ricci curvature of g_4 as

(49)
$$\operatorname{Ric}_{tt} = -2\frac{\alpha(\alpha-1)}{t^2} > 0,$$
$$h^{-2}\operatorname{Ric}_{ii} = \left(\frac{1}{(1-\epsilon)^2} - 1 - \alpha\right)\frac{2}{t^2},$$
$$f^{-2}\operatorname{Ric}_{\alpha\alpha} = \left(\alpha(1-\alpha) + \frac{t^{2-2\alpha}}{\delta^2} - \alpha^2 - 3\alpha\right)\frac{1}{t^2} \ge \left(\alpha(1-\alpha) + \frac{\left(\frac{h_3(r_3)}{1-\epsilon}\right)^{2-2\alpha}}{\delta^2} - \alpha^2 - 3\alpha\right)\frac{1}{t^2}.$$

Notice that as $\alpha_2 \to 0$ we have that $\alpha \to 0$, and similarly (after fixing α_2, α, r_3) we have that $\delta_2 \to 0$ as $\delta \to 0$. Thus for $\alpha_2 \le \alpha_2(\epsilon)$ the second term is uniformly positive. Finally for $\delta_2 \le \delta_2(\alpha_2)$ we have that the third term is uniformly positive.

4.5 Finishing the proof of Lemma 4.1

For $\epsilon > 0$ and $m = \frac{1}{10^3}$ fixed we can now choose $\alpha < \alpha(\epsilon)$ and $\delta < \delta(\epsilon, \alpha)$. Let us now equip $E \times S^2$ with the metric g_4 . Recall that this metric is smooth away from $r \in \{r_1, r_3\}$, globally C^1 and satisfies Ric > 0 on the smooth part. We can now apply the C^1 smoothing Lemma 3.1 in order to build a smooth metric $g = g_E + f^2 g_{S^2}$ on $E \times S^2$ with Ric > 0 such that $g = g_4$ for $r \ge 2r_3$. This completes the construction of $\mathfrak{B} = \mathfrak{B}(\epsilon, \alpha, \delta)$. What remains is to define and study the projection map $\phi: (E, g_E) \to C(S_{1-\epsilon}^3)$.

Recall that we have coordinates (r, ω) on $\mathbb{R}^+ \times S^3$, and we have identified $E \setminus S^2$ and $C(S^3) \setminus \{0\}$ with $\mathbb{R}^+ \times S^3$. Our mapping $\phi: E \to C(S^3)$ will then take the form

(50)
$$\phi(r,\omega) = (\lambda(r),\omega),$$

where $\lambda: [0, \infty) \to [0, \infty)$ is a smooth function with

(51)
$$\begin{cases} \lambda(0) = 0, \\ \lambda', \lambda'' > 0, \\ \lambda(r_3) = r_3 - R_3 < r_3, \\ \lambda'(r) = 1 \quad \text{for } r \ge r_3. \end{cases}$$

Note that since $\lambda' \leq 1$ and $\lambda(0) = 0$, we have $\lambda(r) \leq r$ for all r > 0. Also observe that ϕ sends the zero section S^2 of *E* to the cone point 0 of $C(S_{1-\epsilon}^3)$. It then suffices to estimate $|D\phi|$ in terms of the metrics on *E* and $C(S_{1-\epsilon}^3)$. Since the C^1 gluing lemma produces a smooth metric on *E* which is an arbitrarily small C^1 perturbation, it is enough to estimate in the metric g_4 on *E*.

For $r < r_1$, using the notation and results in Lemma 4.2 we have

(52)
$$|D\phi(\partial_r)| = \lambda'(r) \le 1,$$
$$\left| D\phi\left(\frac{X}{|X|}\right) \right| = \frac{(1-\epsilon)\lambda(r)}{A(r)} \le \frac{(1-\epsilon)r}{(1/k)\sin(kr)} \le \frac{(1-\epsilon)kr_1}{\sin(kr_1)} \le \frac{\pi}{2}(1-\epsilon),$$
$$\left| D\phi\left(\frac{Y}{|Y|}\right) \right| = \left| D\phi\left(\frac{Z}{|Z|}\right) \right| = \frac{(1-\epsilon)\lambda(r)}{B(r)} \le \frac{(1-\epsilon)r_1}{B(\frac{1}{2}r_1)} \le \pi(1-\epsilon).$$

For $r_1 < r < r_3$, using the notation and results in Lemma 4.4 we have

(53)
$$|D\phi(\partial_r)| = \lambda'(r) \le 1,$$
$$\left| D\phi\left(\frac{\partial\omega}{|\partial\omega|}\right) \right| = \frac{(1-\epsilon)\lambda(r)}{h_3(r)} \le \frac{(1-\epsilon)r}{A(r_1) + (r-r_1)m} \le \frac{1-\epsilon}{m}$$

Since $m = 10^{-3}$ is chosen universally, the bounds above do not depend on any other parameters.

For $r > r_3$, $D\phi$ is an isometry. This concludes the proof of Lemma 4.1.

5 Step 2: adding conical singularities

We complete Step 2 of the construction in this section. Namely, we want to see how to take a manifold $M = X^4 \times_f S^2$ and add a cone point in any arbitrarily small neighborhood of X^4 while (almost) preserving a Ricci curvature lower bound. Our primary setup, essentially after rescaling on the regularity scale of M, is to assume we are faced with a warped product space $(B_2(p) \times S^2, g)$ with metric $g = g_B + f^2 g_{S^2}$, under the assumptions

(54)

$$\operatorname{Ric}_{g} > \lambda g, \quad \operatorname{inj}_{g_{B}}(p) > 2,$$

$$|\operatorname{Rm}_{g_{B}}|_{g_{B}}, |\nabla \operatorname{Rm}_{g_{B}}|_{g_{B}}, |\nabla^{2} \operatorname{Rm}_{g_{B}}|_{g_{B}} < \eta < 1,$$

$$|\nabla_{g_{B}} \ln f|_{g_{B}}, |\nabla^{2}_{g_{B}} \ln f|_{g_{B}}, |\nabla^{3}_{g_{B}} \ln f|_{g_{B}} < \eta.$$

Note that there are no assumptions about the sign of $\lambda \in \mathbb{R}$. Observe that the above hold for any warped product $M^4 \times_f S^2$ so long as we work on the regularity scale. Our main result in this section is:

Lemma 5.1 (inductive Step 2) Consider a warped product space $(B_2^4(p) \times S^2, g)$ with metric $g = g_B + f^2 g_{S^2}$ satisfying (54), and write $r := \text{dist}_{g_B}(\cdot, p)$. Then for all choices of parameters $0 < \epsilon < \epsilon(|\lambda|)$, $0 < \alpha < \alpha(\epsilon)$, $0 < \hat{r} < \hat{r}(\alpha, \lambda, \epsilon)$ and $0 < \hat{\delta} < \hat{\delta}(\lambda, \alpha, \epsilon, ||f||_{L^{\infty}}, \hat{r})$, there exists $0 < \delta = \delta(\hat{\delta} ||f||_{\infty} | \alpha, \lambda, \epsilon)$ and a warped product metric $\hat{g} = \hat{g}_B + \hat{f}^2 g_{S^2}$ such that:

- (i) The Ricci lower bound $\operatorname{Ric}_{\widehat{g}} > \lambda C(4)\epsilon$ holds for $\widehat{r}/2 \leq r \leq 2$.
- (ii) $\hat{g} = g_B + \hat{\delta}^2 f^2 g_{S^2}$ is unchanged up to scaling the warping factor f by $\hat{\delta}$ for $1 \le r \le 2$.
- (iii) $\hat{g} = dr^2 + (1-\epsilon)^2 r^2 g_{S^3} + \delta^2 r^{2\alpha} g_{S^2}$ has the cone warping structure $C(S^3_{1-\epsilon}) \times_{\delta r^{\alpha}} S^2$ for $r \le \hat{r}$.
- (iv) The identity map Id: $(B_2(p), \hat{g}_B) \rightarrow (B_2(p), g_B)$ is $(1 + 2\epsilon)$ -bi-Lipschitz.

Remark 5.1 Most of the results of this section hold in general dimensions, where the constants should then include a dimensional dependence. Our notation C(4) denotes that the constant only depends on the dimension n = 4.

Remark 5.2 Recall that our notation of the constant dependence $\delta = \delta(\hat{\delta} || f ||_{\infty} | \alpha, \lambda, \epsilon)$ means that $\delta \to 0$ as $\hat{\delta} || f ||_{\infty} \to 0$ with the other constants $\alpha, \lambda, \epsilon$ fixed.

The construction will be broken down into three steps. In Step 2.1 of Section 5.1 we begin by writing the metric $g_B = dr^2 + r^2g_r$ in exponential polar coordinates, where g_r is a smooth family of metrics on S^3 which naturally converges to the standard metric as $r \to 0$. Our primary goal in Step 2.1 is to alter the base metric g_B to a metric $g_{B,1}$. The metric $g_{B,1}$ will agree with g_B for $1 \le r \le 2$, but will take the form $g_{B,1} = dr^2 + (1-\epsilon)^2 r^2 g_r$ for $r \le r_1$. This will give $g_{B,1}$ a large amount of additional positive Ricci curvature in the nonradial directions, which we will exploit in future steps. Additionally, we are able to ensure that Ric of the total space drops by at worst an ϵ -small amount when passing from g to g_1 .

In Step 2.2 of Section 5.2 we focus on the S^2 warping factor and leave the base $g_{B,2} := g_{B,1}$ fixed. Our goal will be to construct a warping factor f_2 so that $f_2 = \hat{\delta} f_1$ for $r \ge r_1$, while $f_2 = \delta r^{\alpha}$ for $r \le r_2$. The effect of this will be to add a large amount of Ricci curvature to the radial direction of $g_2 := g_{B,2} + f_2^2 g_{S^2}$.

In the final Step 2.3 of Section 5.3 we will use the additional positive Ricci curvature introduced in the first two steps to once again alter the base metric $g_{B,3}$, while fixing $f_3 := f_2$. We will preserve $g_{B,3} = g_{B,2}$ for $r \ge r_2$, however for $r \le r_3$ we will ensure that $g_{B,3} = dr^2 + (1-\epsilon)^2 r^2 g_{S^3}$ is the standard cone $C(S_{1-\epsilon}^3)$. This will complete the construction of \hat{g} , and in Section 5.4 we will check the final bi-Lipschitz property of the construction.

5.1 Step 2.1: decreasing the cone angle

Consider the metric g_B on $B_2(p)$, and by using the radial function $r := d(\cdot, p)$ and exponential coordinates let us write g_B as

where g_r is a smooth family of metrics on S^3 . It follows from Corollary 3.3, recalling that $\eta < 1$, that

(56)
$$(2 - C(4)\eta r^2)g_r \le \operatorname{Ric}_{g_r} \text{ and } \|g_r'\|_{L^{\infty}(S^3, g_r)} \le C(4)\eta r.$$

Let us remark that the first inequality of (56) follows either from the C^2 estimate Corollary 3.3 or directly from the second inequality of (56) by applying the Gauss equations and the identity $\frac{1}{2}(r^2g_r)' = II^{\partial B_r(p)}$.

In this subsection we will look for a metric

(57)
$$g_1 := g_{B,1} + f_1^2 g_{S^2}$$

under the ansatz

(58)
$$g_{B,1} = dr^2 + h(r)^2 g_r$$
 and $f_1 := f$.

Observe that g_r is the original family of metrics on S^3 and that f_1 is a function on $B_2(p)$. For $r_1 := \frac{1}{2}$, the function h(r) will be chosen as any smooth function with the properties that

(59)
$$\begin{cases} h(r) = r & \text{if } r \ge 1, \\ |h - r| + |h' - 1| + |h''| \le 10\epsilon & \text{if } r_1 \le r \le 1, \\ h(r) = (1 - \epsilon)r & \text{if } r \le r_1. \end{cases}$$

In particular, this construction implies that $g_{B,1} = g_B$ for $1 \le r \le 2$, while $g_{B,1} = dr^2 + (1-\epsilon)^2 r^2 g_r$ for $r \le r_1$. Note that on $B_{r_1}(p)$ we have introduced a cone singularity at p, and we will see that this introduces a scale invariant (positive) blow up of the Ricci curvature near p. Our main result in this subsection is that for ϵ sufficiently small, the metric g_1 has Ricci curvature that drops by an arbitrarily small amount from that of g:

Lemma 5.2 Let g satisfy the assumptions of Lemma 5.1 with $g_{B,1}$ defined as in (58) and (59). Then for any $0 < \epsilon < \epsilon(|\lambda|)$, we have that

- (i) $\operatorname{Ric}_{g_1}|_{TB_2(p)} > \lambda C(4)\epsilon$ on $(B_2(p) \setminus \{p\}) \times S^2$, and
- (ii) $\operatorname{Ric}_{g_1}|_{TS^2} > f^{-2} C(4)\eta r^{-1}$ on $(B_2(p) \setminus \{p\}) \times S^2$.

Remark 5.3 The reader may wonder at the disagreeable $\operatorname{Ric}_{g_1}|_{TS^2}$ estimate. This is simply an artifact of the division of the proof of Lemma 5.1 into distinct steps. As soon as the radius r_2 is chosen in Step 2.2 of Section 5.2, we will multiply f by the small but positive number $0 < \hat{\delta} = \hat{\delta}(r_2, \ldots)$ so that the first term of the above $\operatorname{Ric}_{g_1}|_{TS^2}$ lower bound dominates the second in the range $r_2 \leq r \leq 2$.

Proof Our standard notation for the rest of this section will be to use i, j, ... to represent coordinate directions on S^3 and $\alpha, \beta, ...$ to represent coordinate directions on S^2 . The index r is reserved for the radial direction, and we will sometimes write $\partial_r h = h'$ to represent radial derivatives. Let us compute the Ricci curvature of the ansatz (58) as follows:

$$(60) \quad (\operatorname{Ric}_{g_1})_{rr} = (\operatorname{Ric}_g)_{rr} - 3\frac{h''}{h} + \left(\frac{1}{r} - \frac{h'}{h}\right) \operatorname{tr}_{g_r}(g'_r), (\operatorname{Ric}_{g_1})_{ir} = (\operatorname{Ric}_g)_{ir} + \frac{2}{h} \left(\frac{h'}{h} - \frac{1}{r}\right) \frac{h\partial_i f_1}{f_1}, (\operatorname{Ric}_{g_1})_{ij} = (\operatorname{Ric}_g)_{ij} + \frac{h^2 - r^2}{r^2} (\operatorname{Ric}_g)_{ij} + \left(\frac{1}{h^2} - \frac{1}{r^2}\right) (h^2 \operatorname{Ric}_{g_r})_{ij} - \frac{h''}{h} (g_1)_{ij} + 2 \left(\frac{1}{r^2} - \frac{(h')^2}{h^2}\right) (g_1)_{ij} + \left(\frac{1}{r} - \frac{h'}{h}\right) \left(\frac{1}{2} \operatorname{tr}_{g_r}(g'_r)(g_1)_{ij} + \frac{3}{2} (h^2 g'_r)_{ij} + 2\frac{f'_1}{f_1} (g_1)_{ij}\right) + 2 \frac{h^2 - r^2}{r^2} \left(\frac{(\nabla^2_{g_B} f_1)_{ij}}{f_1} - \frac{1}{2} \frac{f'_1}{f_1} (g'_r)_{ij}\right),$$

Lower Ricci curvature and nonexistence of manifold structure

$$\begin{aligned} (\operatorname{Ric}_{g_1})_{\alpha\beta} &= \left[\frac{1}{f_1^2} - \frac{r^2}{h^2} \left(\frac{\Delta_{g_B} f_1}{f_1} + \frac{|\nabla^{g_B} f_1|_{g_B}^2}{f_1^2} \right) + \frac{r^2 - h^2}{h^2} \left(\frac{f_1''}{f_1} + \frac{(f_1')^2}{f_1^2} + \frac{1}{2} \frac{f_1'}{f_1} \operatorname{tr}_{g_r}(g_r') \right) \\ &+ 3 \left(\frac{r^2 - h^2}{rh^2} + \frac{1}{r} - \frac{h'}{h} \right) \frac{f_1'}{f_1} \right] (g_1)_{\alpha\beta}. \end{aligned}$$

As in (56) we have that

(61)
$$(2 - C\eta r^2)g_r \le \operatorname{Ric}_{g_r} \quad \text{and} \quad \|g_r'\|_{L^{\infty}(S^3, g_r)} \le C\eta r.$$

Additionally, we recall the assumptions

(62)
$$|\nabla_{g_B} \ln f|_{g_B} < \eta$$
 and $\left| \frac{(\nabla^2_{g_B} f)}{f} \right|_{g_B} \le |\nabla^2_{g_B} \ln f|_{g_B} + |\nabla_{g_B} \ln f|^2_{g_B} < C\eta.$

Let us focus first on the region $r \in [r_1, 1]$, which is the worst of the two regions. If we plug the above estimates and (59) into (60), we arrive at

(63)

$$(\operatorname{Ric}_{g_{1}})_{rr} > \lambda - C(1 + C\eta)\epsilon,$$

$$|(\operatorname{Ric}_{g_{1}} - \operatorname{Ric}_{g})_{ir}|_{(g_{1})_{ij}} < C\eta\epsilon,$$

$$(\operatorname{Ric}_{g_{1}})_{ij} > (\lambda - C(1 + C\eta)\epsilon)(g_{1})_{ij},$$

$$(\operatorname{Ric}_{g_{1}})_{\alpha\beta} > \left(\frac{1}{f^{2}} - C\eta\right)(g_{1})_{\alpha\beta}.$$

We see from the first three inequalities that if $\epsilon < \epsilon(|\lambda|)$, then $\operatorname{Ric}_{g_1}|_{TB_2(p)} > \lambda - C(4)\epsilon$. If we focus now on $r \le r_1$, where $h(r) = (1 - \epsilon)r$, then most of the error terms of (60) vanish. What remains are the estimates

(64)

$$(\operatorname{Ric}_{g_{1}})_{rr} > \lambda,$$

$$(\operatorname{Ric}_{g_{1}})_{ri} = (\operatorname{Ric}_{g})_{ri},$$

$$(\operatorname{Ric}_{g_{1}})_{ij} > \left(\lambda + \frac{(2 - C\eta r^{2})\epsilon}{r^{2}} - C\frac{\eta\epsilon}{r} - C\eta\epsilon\right)(g_{1})_{ij},$$

$$(\operatorname{Ric}_{g_{1}})_{\alpha\beta} > \left(\frac{1}{f^{2}} - C\frac{\eta\epsilon}{r} - C\eta\right)(g_{1})_{\alpha\beta}.$$

We again see that for $\epsilon < \epsilon(|\lambda|)$, the Ricci curvature satisfies $\operatorname{Ric}_{g_1} \ge \lambda - C\epsilon$ in the $TB_2(p)$ directions. We also see that $\operatorname{Ric}_{g_1}|_{TS^2} > f^{-2} - C\eta r^{-1}$ holds for both the $r_1 \le r \le 1$ region and the $r \le r_1$ region. \Box

5.2 Step 2.2: the S^2 warping factor

We now want to alter the metric g_1 in the range $r_2 \le r \le r_1 = \frac{1}{2}$. We are looking for a metric g_2 of the form

(65)
$$g_2 := g_{B,2} + f_2(x)^2 g_{S^2} := g_{B,1} + f_2(x)^2 g_{S^2}.$$

In particular, we will not alter the base metric $g_{B,1}$ in this step. Our warping function $f_2: B_2 \to \mathbb{R}^+$ is not a radial function everywhere, though one of our goals will be to make it radial on small radii. We will want f_2 to satisfy the properties

(66)
$$\begin{cases} f_2 = \hat{\delta} f_1 & \text{if } r \ge r_1 \\ f_2 = \delta r^{\alpha} & \text{if } r \le r_2 \end{cases}$$

Due to the nonradial nature of f_1 , there is some subtlety which makes a naive interpolation between f_1 and δr^{α} insufficient. Morally, this is due to the uncontrolled positivity of Ric, which can contribute very negative terms if altered in a careless manner. The following will be the main constructive lemma for f_2 in this subsection:

Lemma 5.3 Let $(B_2(p), g_B)$ and f_1 be as in Lemma 5.2. Then for each $0 < \alpha < 1$, $0 < \hat{\delta} < 1$ and $0 < r_2 < r_2(\alpha)$, there exists $f_2: B_2(p) \to \mathbb{R}^+$ with $r_2 < C(4)r_2 = r_2^+ < r_1$ and $\delta = \delta(\hat{\delta} || f ||_{\infty} || r_2)$ such that

- (i) $f_2 = \hat{\delta} f_1$ on the region $r_2^+ \le r \le 2$,
- (ii) $f_2 = \delta r^{\alpha}$ on the region $r \leq r_2$,
- (iii) $-f_2''/f_2 \ge \alpha r^{-2}$ on the region $r_2 \le r \le r_2^+$,

(iv)
$$\alpha^{-1} |\nabla_{g_B} \ln f_2|_{g_B}^2$$
, $|\nabla_{g_B}^2 \ln f_2|_{g_B} < C(4) \alpha r^{-2}$ on the region $r_2 \le r \le r_2^+$,

- (v) $||f_2||_{\infty} \le C(4)(\hat{\delta}||f||_{\infty})^{1/2}$ on the region $r_2 \le r \le r_2^+$, and
- (vi) f_2 is smooth away from $r \in \{0, r_2, r_2^+\}$, and C^1 everywhere except $\{p\}$.

Remark 5.4 The C^1 nature of f_2 is due to (iii), where we force a definite amount of radial concavity throughout the interpolation region. This will later be smoothed with a C^1 gluing lemma.

We will wait until the end of the subsection to prove the above, which will require a bit of work. Let us begin instead with how to use it in order to build our desired g_2 .

Lemma 5.4 Let $(B_2(p) \times S^2, g_1)$ be as in Lemma 5.2, and for $0 < \alpha < \alpha(\epsilon)$, $r_2 = r_2(\lambda, \alpha, \epsilon)$ and $0 < \hat{\delta} < \hat{\delta}(\lambda, \alpha, \epsilon, \|f\|_{\infty}, \hat{r})$, let f_2 be as in Lemma 5.3. Then $g_2 := g_{B,1} + f_2^2 g_{S^2}$ satisfies $\operatorname{Ric}_{g_2} > \lambda - C(4)\epsilon$ away from $r \in [0, \hat{r}/2) \cup \{r_2, r_2^+\}$.

Proof of Lemma 5.4 given Lemma 5.3 Let us apply Lemma 5.3, to obtain f_2 . We will choose our constants r_2 , α and $\hat{\delta}$ later in the proof. We begin with the Ricci curvature computation, where g_B is the base metric for the *original* metric from Lemma 5.1 :

(67)
$$(\operatorname{Ric}_{g_2})_{rr} = (\operatorname{Ric}_{g_B})_{rr} - 2\frac{f_2''}{f_2},$$

 $(\operatorname{Ric}_{g_2})_{ir} = (\operatorname{Ric}_{g_B})_{ir} - 2\frac{(\nabla_{g_B}^2 f_2)_{ir}}{f_2},$

Lower Ricci curvature and nonexistence of manifold structure

$$\begin{split} (\operatorname{Ric}_{g_2})_{ij} &= (1-\epsilon)^2 (\operatorname{Ric}_{g_B})_{ij} + (1-(1-\epsilon)^2) (\operatorname{Ric}_{g_r})_{ij} - 2 \frac{(\nabla_{g_B}^2 f_2)_{ij}}{f_2} \\ &+ (1-(1-\epsilon)^2) \left(\frac{2}{r} (g_2)_{ij} + (r^2 g_r')_{ij}\right) \frac{f_2'}{f_2}, \\ (\operatorname{Ric}_{g_2})_{\alpha\beta} &= \left[\frac{1}{f_2^2} - \frac{1}{(1-\epsilon)^2} \left(\frac{\Delta_{g_B} f_2}{f_2} + \frac{|\nabla^{g_B} f_2|_{g_B}^2}{f_2^2}\right) \\ &+ \frac{1-(1-\epsilon)^2}{(1-\epsilon)^2} \left(\frac{f_2''}{f_2} + \frac{(f_2')^2}{f_2^2} + \left(\frac{1}{2}\operatorname{tr}_{g_r}(g_r') + \frac{3}{r}\right) \frac{f_2'}{f_2}\right) \right] (g_2)_{\alpha\beta}. \end{split}$$

To estimate the Ricci curvature, we have three regions to study, namely $r \in (0, r_2) \cup (r_2, r_2^+) \cup (r_2^+, r_1)$. The region $r \in (r_1, 2)$ is covered by Remark 3.2 and the previous steps of the construction. Let us begin with the region $r \in (r_2^+, r_1)$, and let us use Lemma 5.2. If we allow $\hat{\delta} < \hat{\delta}(||f||_{\infty}, r_2^+, |\lambda|)$ then we can use Lemma 5.2 to estimate

(68)
$$\operatorname{Ric}_{g_2}|_{TB_2(p)} > (\lambda - C(4)\epsilon) \text{ and } \operatorname{Ric}_{g_2}|_{TS^2} > f_2^{-2} - C(4)\eta r^{-1} \ge \frac{1}{\hat{\delta}^2 f^2} - \frac{C\eta}{r_2^+} > \lambda.$$

In particular, once r_2^+ has been fixed we may fix $\hat{\delta}$ sufficiently small to control the above. Let us therefore focus on the more challenging situations: when $r \in (0, r_2) \cup (r_2, r_2^+)$.

To begin, let us record some basic estimates which will be used. Recalling (56), we have that

(69)
$$(2-C\eta r^2)g_r \leq \operatorname{Ric}_{g_r} \quad \text{and} \quad \|g_r'\|_{L^{\infty}(S^3,g_r)} \leq C\eta r.$$

In view of Lemma 5.3, we have in the region $r_2 < r < r_2^+$ that

(70)
$$\alpha^{-1} |\nabla_{g_B} \ln f_2|_{g_B}^2, |\nabla_{g_B}^2 \ln f_2|_{g_B} < C \frac{\alpha}{r^2} \quad \text{and} \quad \frac{\alpha}{2r^2} \le -\frac{f_2''}{f_2}.$$

Moreover, the same estimates hold when $0 < r < r_2$, because $f_2 = \delta r^{\alpha}$ in this region. Let us analyze each $B_2(p) \times S^2$ block of Ric_{g_2} according to (67). For the r, i, j, \ldots block corresponding to the base $B_2(p)$ we can use the above estimates to get

(71)

$$(\operatorname{Ric}_{g_{2}})_{rr} > \frac{\alpha}{2r^{2}} - C\eta,$$

$$|(\operatorname{Ric}_{g_{2}})_{ir}| < C(\eta)\frac{\alpha}{r^{2}} + C\eta,$$

$$(\operatorname{Ric}_{g_{2}})_{ij} > \left(\frac{(2 - C\eta r^{2})\epsilon}{r^{2}} - C\left(\frac{\alpha}{r^{2}} + \eta\right)\right)(g_{2})_{ij}.$$

If we require that $\alpha < \alpha(\epsilon)$ and $r_2^+ < r_2^+(\alpha, \epsilon)$ are small enough, then

(72)
$$(\operatorname{Ric}_{g_2})_{rr} > \frac{\alpha}{4r^2}, \quad (\operatorname{Ric}_{g_2})_{ij} > \frac{\epsilon}{r^2}(g_2)_{ij}, \quad |(\operatorname{Ric}_{g_2})_{ir}| < C\frac{\alpha}{r^2}.$$

If $\alpha < \alpha(\epsilon)$ is again small enough, then this gives the estimate $\operatorname{Ric}_{g_2}|_{TB_2(p)} > \epsilon/2r^2$. If we finally require that $r_2^+ < r_2^+(\epsilon, |\lambda|)$, then this gives the required estimate for the B_2 block.

For the S^2 block of Ric_{g_2} , we argue separately for $r \in (\hat{r}/2, r_2)$ and $r \in (r_2, r_2^+)$. When $r \in (r_2, r_2^+)$ let us use (67) and (70) in order to estimate

(73)
$$(\operatorname{Ric}_{g_2})_{\alpha\beta} > \left(\frac{1}{\|f_2\|_{\infty}^2} - C\left(\frac{\alpha}{r_2^2} + \epsilon\eta\alpha\right)\right)(g_2)_{\alpha\beta}.$$

Recalling the bound $||f_2||_{\infty} \leq C(\hat{\delta}||f||_{\infty})^{1/2}$ from Lemma 5.3, it is clear that a small enough choice of $\hat{\delta} < \hat{\delta}(||f||_{\infty}, \lambda, r_2)$ ensures that $(\operatorname{Ric}_{g_2})_{\alpha\beta} > \lambda(g_2)_{\alpha\beta}$ for $r \in (r_2, r_2^+)$. We can now focus on the region $r \in (\hat{r}/2, r_2)$, where

(74)
$$(\operatorname{Ric}_{g_2})_{\alpha\beta} > \left(\frac{1}{(\delta r_2^{\alpha})^2} - C\frac{\alpha}{\hat{r}^2}\right)(g_2)_{\alpha\beta}$$

We see that for $\delta < \delta(\hat{r}, |\lambda|)$, we have the desired $(\operatorname{Ric}_{g_2})_{\alpha\beta} > \lambda(g_2)_{\alpha\beta}$ in this region, which would complete the proof if δ were sufficiently small. However, we have from Lemma 5.3 that $\delta = \delta(\hat{\delta} || f ||_{\infty} | r_2)$, so it suffices to require that $\hat{\delta} < \hat{\delta}(|| f ||_{\infty}, r_2, \hat{r}, |\lambda|)$ to ensure that δ is sufficiently small.

Proof of Lemma 5.3 Before diving into the proof, we establish some new notation for the sake of legibility. We label

(75)
$$f_+(r,\omega) := \widehat{\delta} f_1(r,\omega) \quad \text{and} \quad f_-(r,\omega) = f_-(r) := \delta r^{\alpha},$$

where δ will be specified later in the proof. Also, instead of referring to the radii r_2 and r_2^+ directly, it will be convenient to write the interval of interpolation (r_2, r_2^+) in terms of a midpoint $r_m := (r_2 + r_2^+)/2$ and a radius $\rho r_m := (r_2^+ - r_2)/2$. In particular, the choice of radii r_2, r_2^+ from the statement of Lemma 5.3 will instead take the form of a choice of a constant ρ and radius $r_m < r_m(\alpha)$.

The remaining proof has multiple steps, which we will break down into pieces:

Locating the intersection set of f_- and f_+ We first seek to set it up so that the intersection set $\{f_+ = f_-\} = \{(r, \omega) \mid f_+(r, \omega) = f_-(r)\}$ is approximately at our radius r_m , which requires selecting δ depending on r_m . We will estimate the deviation of the intersection set from this radius. These estimates will be used in the next steps of the proof.

Begin by observing that the $\nabla_{g_B} \ln f$ bound gives the estimate

(76)
$$|\ln f_+(r,\omega) - \ln f_+(p)| \le \eta r \implies f_+(r,\omega) \in [f_+(p)e^{-2\eta r_m}, f_+(p)e^{2\eta r_m}]$$
 for all $r \le 2r_m$.

One then checks under what conditions f_{-} takes values in this same interval:

(77)
$$f_{-}(r) = \delta r^{\alpha} \in [f_{+}(p)e^{-2\eta r_{m}}, f_{+}(p)e^{2\eta r_{m}}] \qquad \Longleftrightarrow r \in [(\delta^{-1}f_{+}(p)e^{-2\eta r_{m}})^{1/\alpha}, (\delta^{-1}f_{+}(p)e^{2\eta r_{m}})^{1/\alpha}].$$

It is at this point that we make the choice

$$\delta := f_+(p) r_m^{-\alpha} e^{2\eta r_m}.$$

For this choice of δ , we have

$$(\delta^{-1} f_+(p) e^{2\eta r_m})^{1/\alpha} = r_m$$

and thus when (77) holds we have $r \leq 2r_m$. Observe that we now have the relationships

(78)
$$\begin{cases} f_{-}(r) < f_{+}(p)e^{-2\eta r_{m}} \le f_{+}(r,\omega) & \text{if } r < r_{m}e^{-4\eta r_{m}/\alpha}, \\ f_{-}(r) \in [f_{+}(p)e^{-2\eta r_{m}}, f_{+}(p)e^{2\eta r_{m}}] & \text{if } r_{m}e^{-4\eta r_{m}/\alpha} \le r \le r_{m}, \\ f_{-}(r) > f_{+}(p)e^{2\eta r_{m}} \ge f_{+}(r,\omega) & \text{if } r_{m} < r \le 2r_{m}. \end{cases}$$

In particular, we have

(79)
$$\{f_{-}=f_{+}\}\cap B_{2r_{m}}(p)\subset A_{r_{m}e^{-4\eta r_{m}/\alpha},r_{m}}(p).$$

Let us additionally remark that $\{f_- = f_+\} \cap B_{2r_m}(p) \cap (\mathbb{R}_+ \times \{\omega\})$ is nonempty for each $\omega \in S^3$ by the intermediate value theorem. We may therefore define $g(\omega) \in [r_m e^{-4\eta r_m/\alpha}, r_m]$ for each $\omega \in S^3$ to be a radius such that $f_+(g(\omega), \omega) = f_-(g(\omega))$. We can ensure that $(g(\omega), \omega)$ always lies in the interpolation region $r \in ((1 - \rho)r_m, (1 + \rho)r_m)$ by requiring that $r_m < r_m(\alpha)$ is small enough that $e^{-4\eta r_m/\alpha} > 1 - \rho$. It is not necessary that g be continuously defined.

Definition of f_2 and C^1 cubic interpolation Our construction of f_2 will make use of the general notion of a C^1 cubic interpolation. Namely, we will ask that f_2 be the uniquely defined C^1 function satisfying

(80)
$$\ln f_2(r,\omega) = \begin{cases} \ln f_-(r) & \text{if } r \le (1-\rho)r_m \\ Q(r,\omega) = \text{cubic polynomial in } r & \text{if } r \in ((1-\rho)r_m, (1+\rho)r_m) \\ \ln f_+(r,\omega) & \text{if } r \ge (1+\rho)r_m. \end{cases}$$

We see from the above that $Q(r, \omega)$ is therefore well defined by the values of f_{\pm} and f'_{\pm} at the end points of the interval $[(1 - \rho)r_m, (1 + \rho)r_m]$. It will be helpful to write the form of $Q(r, \omega)$ explicitly by

$$(81) \quad Q(r,\omega) := \left[\frac{r - (1 - \rho)r_m}{2\rho r_m}\right]^2 \left[\ln f_+((1 + \rho)r_m, \omega) - ((1 + \rho)r_m - r)\left((\ln f_+)'((1 + \rho)r_m, \omega) - \frac{1}{\rho r_m}\ln f_+((1 + \rho)r_m, \omega)\right)\right] + \left[\frac{(1 + \rho)r_m - r}{2\rho r_m}\right]^2 \left[\ln f_-((1 - \rho)r_m) + (r - (1 - \rho)r_m)\left((\ln f_-)'((1 - \rho)r_m) + \frac{1}{\rho r_m}\ln f_-((1 - \rho)r_m)\right)\right]$$

Concavity estimates for ln f_2 We can now estimate $(\ln f_2)''$ in the interpolation region

$$r \in ((1-\rho)r_m, (1+\rho)r_m).$$

Next, we use the mean value theorem to estimate the difference, at $g(\omega)$, between $r \mapsto \ln f_{\pm}(r, \omega)$ and its linearization centered at $r = (1 \pm \rho)r_m$:

$$(82) \quad (\ln f_{2})'' = \frac{1}{2\rho r_{m}} ((\ln f_{+})'((1+\rho)r_{m},\omega) - (\ln f_{-})'((1-\rho)r_{m})) \\ + \frac{3(r-r_{m})}{2\rho^{3}r_{m}^{3}} (g(\omega) - r_{m}) ((\ln f_{+})'((1+\rho)r_{m},\omega) - (\ln f_{-})'((1-\rho)r_{m})) \\ + \underbrace{\frac{3(r-r_{m})}{2\rho^{3}r_{m}^{3}} [((1+\rho)r_{m} - g(\omega))(\ln f_{+})'((1+\rho)r_{m},\omega) - (\ln f_{+}((1+\rho)r_{m},\omega) - \ln f_{+}(g(\omega),\omega))]}_{=O(\| \ln f_{+}''\|_{\infty})} \\ + \underbrace{\frac{3(r-r_{m})}{2\rho^{3}r_{m}^{3}} [(g(\omega) - (1-\rho)r_{m})(\ln f_{-})'((1-\rho)r_{m}) - (\ln f_{-}(g(\omega)) - \ln f_{-}((1-\rho)r_{m}))]}_{=O(\| \ln f_{-}''\|_{\infty})} \\ \leq \left(-\frac{1}{2\rho} + \frac{3}{2} \frac{|g(\omega) - r_{m}|}{\rho^{2}r_{m}} + \frac{C}{1-\rho} \right) \frac{\alpha}{(1-\rho)r_{m}^{2}} + \left(\frac{1}{2\rho} + \frac{3}{2} \frac{|g(\omega) - r_{m}|}{\rho^{2}r_{m}} + Cr_{m} \right) \frac{\eta}{r_{m}}.$$

The desired concavity should arise from the first term on the right-hand side of (82). We therefore pick the universal constant $\rho > 0$ small enough so that

$$-\frac{1}{2\rho} + \frac{C}{1-\rho} < -\frac{4}{1-\rho},$$

and then $r_m \leq r_m(\alpha)$ small enough so that

$$|1 - e^{-4\eta r_m/\alpha}| \le \frac{2\rho^2}{3(1-\rho)}$$

Since $g(\omega) \in (r_m e^{-4\eta r_m/\alpha}, r_m)$, the constraint on r_m implies that

$$|g(\omega) - r_m| \le \frac{2\rho^2}{3(1-\rho)}r_m$$

Thus,

(83)
$$\left(-\frac{1}{2\rho} + \frac{3}{2}\frac{|g(\omega) - r_m|}{\rho^2 r_m} + \frac{C}{1-\rho}\right) \le -\frac{3}{1-\rho}$$

We also require that $r_m \leq r_m(\alpha)$ is small enough so that

$$\frac{\eta}{r_m} \le \left(\frac{1}{2\rho} + \frac{1}{1-\rho} + Cr_m\right)^{-1} \frac{\alpha}{(1-\rho)^2 r_m^2}$$

in order to absorb the lower-order second term on the right-hand side of (82). This yields the claimed concavity

(84)
$$(\ln f_2)'' \le -2\frac{\alpha}{[(1-\rho)r_m]^2} \le -2\frac{\alpha}{r^2} \text{ for all } r \in ((1-\rho)r_m, (1+\rho)r_m).$$

Radial zeroth- and first-order estimates on f_2 We begin this step by obtaining a bound on $(\ln f_2)'$. This allows us to turn the concavity estimates for $\ln f_2$ into concavity estimates for f_2 . Additionally, we will use the estimate on $(\ln f_2)'$ to obtain L^{∞} control for f_2 in the interpolation region.

Geometry & Topology, Volume 29 (2025)

472

The previous step implies that for fixed $\omega \in S^3$, the map $r \mapsto \ln f'_2(r, \omega)$ is decreasing from the value $\ln f'_2((1-\rho)r_m, \omega) = \alpha[(1-\rho)r_m]^{-1}$ at $r = (1-\rho)r_m$. Thus,

(85)
$$|(\ln f_2)'| \le \frac{\alpha}{(1-\rho)r_m} \le \alpha \frac{1+\rho}{1-\rho}r^{-1}$$

Combining this with our concavity estimate for $\ln f_2$ gives

(86)
$$\frac{f_2''}{f_2} = (\ln f_2)'' + (\ln f_2')^2 \le -2\frac{\alpha}{r^2} + \left(\frac{1+\rho}{1-\rho}\right)^2 \frac{\alpha^2}{r^2} \le -\alpha r^{-2} \text{ as long as } \alpha \le \left(\frac{1-\rho}{1+\rho}\right)^2.$$

This proves Lemma 5.3.(iii). To obtain L^{∞} control on f_2 , let us estimate f_2 on $((1 - \rho)r_m, (1 + \rho)r_m)$ by using the explicit formula (81) for Q:

(87)
$$Q(r,\omega) \leq \left[\frac{r - (1 - \rho)r_m}{2\rho r_m}\right]^2 \left[\ln f_+((1 + \rho)r_m, \omega) + \eta((1 + \rho)r_m - r)\right] \\ + \left[\frac{(1 + \rho)r_m - r}{2\rho r_m}\right]^2 \left[\ln f_-((1 - \rho)r_m) + \alpha \frac{(r - (1 - \rho)r_m)}{(1 - \rho)r_m}\right] \\ \leq \left(\left[\frac{r - (1 - \rho)r_m}{2\rho r_m}\right]^2 + \left[\frac{(1 + \rho)r_m - r}{2\rho r_m}\right]^2\right) \ln(\|f_+\|_{\infty}) + \frac{2\rho}{1 - \rho}\alpha + 2\rho r_m \eta \\ \leq \frac{1}{2}\ln\|f_+\|_{\infty} + C\alpha + Cr_m \eta.$$

Note that for the second inequality above, we used that $f_{-}((1-\rho)r_m) \leq f_{-}(g(\omega)) = f_{+}(g(\omega))$. Since f_{-} is increasing, this holds if $(1-\rho)r_m \leq g(\omega)$, which itself will hold if $r_m < r_m(\alpha)$ is small enough because $g(\omega) \geq r_m e^{-4\eta r_m/\alpha}$. Taking exp on both sides proves Lemma 5.3(v).

Remaining derivative estimates The last remaining item is Lemma 5.3(iv), which we now turn to. The constraint $r_m \leq r_m(\alpha)$ is also finalized in this last step.

For the remaining computations, we pick coordinates ∂_i on S^3 that are normal at $\omega \in S^3$ for the metric g_r , and assume that we are working in the region $r \in ((1 - \rho)r_m, (1 + \rho)r_m)$. Before launching into the remaining derivative estimates, we recall two basic bounds that we will need, both from Corollary 3.3:

(88)
$$\|g'_r\|_{L^{\infty}(S^3, g_r)} \le C(4)\eta r \text{ and } \|g''_r\|_{L^{\infty}(S^3, g_r)} \le C(4)\eta$$

We begin with the tangential derivatives of f_2 . One notices the identity, using the notation (81),

(89)
$$\partial_i \ln f_2 = \left[\frac{r - (1 - \rho)r_m}{2\rho r_m}\right]^2 \left[\partial_i \ln f_+((1 + \rho)r_m, \omega) - ((1 + \rho)r_m - r)((\partial_i \ln f_+)'((1 + \rho)r_m, \omega) - \frac{1}{\rho r_m}\partial_i \ln f_+((1 + \rho)r_m, \omega)\right].$$

What is important is that the right-hand side is a linear combination of $\partial_i \ln f_+((1 + \rho)r_m, \omega)$ and $O(r)\partial_i \ln f'_+((1 + \rho)r_m, \omega)$ with uniformly bounded coefficients. The first term $\partial_i \ln f_+((1 + \rho)r_m, \omega)$ is bounded by η , while the second equals

$$O(r)(\partial_i \ln f'_+) = O(r)(\nabla_{i,r}^2 \ln f_+ + r^{-1}(\delta_i^k + rg_r^{kj}(g'_r)_{ji}/2)\partial_k \ln f_+),$$

which can be bounded by $C\eta$. Thus, for $r_m \leq r_m(\alpha)$, we can estimate

(90)
$$|\nabla_i \ln f_2|_{r^2(g_r)_{ij}} \le C(4)\eta \le \frac{\alpha}{r}.$$

Similarly, $\partial_i \partial_j \ln f_2$ is a linear combination of $\partial_i \partial_j \ln f_+((1+\rho)r_m, \omega)$ and $O(r)\partial_i \partial_j \ln f'_+((1+\rho)r_m, \omega)$ with bounded coefficients, where

$$(91) \quad \partial_{i}\partial_{j}\ln f_{+} = \nabla_{i,j}^{2}f_{+} - \frac{1}{2}(2r(g_{r})_{ij} + r^{2}(g_{r}')_{ij})\partial_{r}f_{+}, \\ \partial_{i}\partial_{j}\ln f_{+}' = \nabla_{r,i,j}^{3}f_{+} - \frac{1}{2}(2(g_{r})_{ij} + 4r(g_{r}')_{ij} + r^{2}(g_{r}'')_{ij})\partial_{r}f_{+} - \frac{1}{2}(2r(g_{r})_{ij} + r^{2}(g_{r}')_{ij})\partial_{r}^{2}f_{+} \\ + \frac{1}{2r}(2\delta_{i}^{k} + rg_{r}^{k\ell}(g_{r}')_{\ell i})\nabla_{k,j}^{2}f_{+} + \frac{1}{2r}(2\delta_{j}^{k} + rg_{r}^{k\ell}(g_{r}')_{\ell j})\nabla_{k,j}^{2}f_{+} + \frac{1}{2r}(2\delta_{j}^{k$$

The highest-order term is $-(g_r)_{ij}\partial_r f_+$ from the second equation, and can be bounded by ηr^{-2} . The other terms can be estimated by $C\eta r^{-1}$. Using this and the previously obtained bound on $\ln f'_2$, and again requiring that $r_m \leq r_m(\alpha)$, we get

(92)
$$|\nabla_{ij}^2 \ln f_2|_{r^2(g_r)_{ij}} \le |\partial_i \partial_j \ln f_2|_{r^2(g_r)_{ij}} + \frac{1}{2} |(2r(g_r)_{ij} + r^2(g_r')_{ij}) \ln f_2'|_{r^2(g_r)_{ij}} \\ \le C \frac{\eta}{r} + C \frac{\alpha}{r^2} \le C \frac{\alpha}{r^2}.$$

Finally we repeat this strategy once more and observe as in (89) that $\partial_i \partial_r \ln f_2$ is a linear combination of $\partial_i \ln f_+((1+\rho)r_m, \omega)$ and $O(r)\partial_i \ln f'_+((1+\rho)r_m, \omega)$, with coefficients that are bounded times a factor $\leq r_m^{-1}$. We have already bounded these two expressions by $C\eta$, so for $r_m \leq r_m(\alpha)$ we get

(93)
$$|\nabla_{i,r}^{2} \ln f_{2}|_{r^{2}(g_{r})_{ij}} \leq |\partial_{i}\partial_{r} \ln f_{2}|_{r^{2}(g_{r})_{ij}} + \frac{1}{2r} |(2\delta_{i}^{k} + rg_{r}^{k\ell}(g_{r}')_{\ell i})\partial_{k} \ln f_{2}|_{r^{2}(g_{r})_{ij}} \\ \leq C(4)\eta r_{m}^{-1} + C(4)\eta r^{-1} \leq C(4)\eta \frac{\alpha}{r^{2}},$$

which finishes the proof of Lemma 5.3.

5.3 Step 2.3: interpolating the warped cone to a homogeneous cone

We ended the last subsection having constructed a metric

(94)
$$g_2 = g_{B,2} + f_2^2 g_{S^2}.$$

This metric has the property that for $r \leq r_2$ we can write

(95)
$$g_{B,2} = dr^2 + (1-\epsilon)^2 r^2 g_r$$
 and $f_2 = f_2(r) = \delta r^{\alpha}$,

where recall that $g_B = dr^2 + r^2g_r$ was the original smooth metric with regularity scale $r_x > 2$; see (54). Our final goal in this section is to interpolate g_r with the unit sphere metric g_{S^3} . In this way we will build a metric

(96)
$$g_3 := g_{B,3} + f_3^2 g_{S^2},$$

Geometry & Topology, Volume 29 (2025)

where $f_3 := f_2$ and $g_{B,3} = dr^2 + (1 - \epsilon)^2 r^2 g_{r,3}$, with

(97)
$$g_{r,3} := \begin{cases} g_r & \text{if } r \ge 2r_3, \\ \xi(r)g_r + (1-\xi(r))g_{S^3} & \text{if } r \in [r_3, 2r_3], \\ g_{S^3} & \text{if } r \le r_3. \end{cases}$$

Here, $\xi(r)$ is a smooth cutoff on $[r_3, 2r_3]$ satisfying the r_3 -independent bounds

(98)
$$|\xi'(r)| \le Cr^{-1}$$
 and $|\xi''(r)| \le Cr^{-2}$.

We will show that for r_3 sufficiently small, the metric g_3 will have the appropriate Ricci curvature bounds:

Lemma 5.5 Let g_3 be defined as in (96) and (97) under the assumptions of Lemma 5.1. Then for $r_3 \leq r_3(\alpha, \lambda, \epsilon)$, we have that $\operatorname{Ric}_{g_3} > \lambda - C(4)\epsilon$ on $(B_2(p) \setminus B_{\hat{r}/2}(p)) \times S^2$.

Proof For radii $r > 2r_3$, the Ricci lower bound follows from Lemma 5.4 and the fact that $g_2 = g_3$ in this region. Thus, we assume for the rest of the proof that $r \le 2r_3$. The main basic estimates we will need are from Corollary 3.3, namely

(99)
$$||g_r - g_{S^3}||_{C^2(S^3, g_r)} \le C\eta r^2, \quad ||g_r'||_{C^1(S^3, g_r)} \le C\eta r, \quad ||g_r''||_{L^{\infty}(S^3, g_r)} \le C\eta.$$

Let us first apply the above in order to see the estimates

(100)
$$\|g_{r,3} - g_{S^3}\|_{C^2(S^3, g_{r,3})} \le C\eta r^2,$$
$$\|g_{r,3}'\|_{C^1(S^3, g_{r,3})} = \|\xi'(g_r - g_{S^3}) + \xi g_r'\|_{C^1(S^3, g_{r,3})} \le C\eta r,$$
$$\|g_{r,3}''\|_{L^{\infty}(S^3, g_{r,3})} = \|\xi''(g_r - g_{S^3}) + 2\xi'g_r' + \xi g_r''\|_{L^{\infty}(S^3, g_{r,3})} \le C\eta.$$

We are now in a position to estimate Ric_{g_3} , which we compute as follows:

$$(101) \quad (\operatorname{Ric}_{g_{3}})_{rr} = -\frac{1}{2} \operatorname{tr}_{g_{r,3}}(g_{r,3}'') - \frac{1}{r} \operatorname{tr}_{g_{r,3}}(g_{r,3}') + \frac{1}{4} |g_{r,3}'|_{g_{r,3}}^{2} - 2\frac{\alpha(\alpha-1)}{r^{2}},$$

$$(\operatorname{Ric}_{g_{3}})_{ir} = \frac{1}{2} (\partial_{i} \operatorname{tr}_{g_{r,3}}(g_{r,3}') - (\operatorname{tr}_{g_{r,3}}^{1/2}(\nabla g_{r,3}'))_{i}),$$

$$(\operatorname{Ric}_{g_{3}})_{ij} = (1 - (1 - \epsilon)^{2})(\operatorname{Ric}_{g_{S^{3}}})_{ij} + (\operatorname{Ric}_{g_{r,3}} - \operatorname{Ric}_{g_{S^{3}}})_{ij} - \frac{1}{2}(1 - \epsilon)^{2}(r^{2}g_{r,3}')_{ij}$$

$$-(1 - \epsilon)^{2} \left(\frac{3}{2r} + \frac{1}{4} \operatorname{tr}_{g_{r,3}}(g_{r,3}')\right)(r^{2}g_{r,3}')_{ij} + \frac{1}{2}(1 - \epsilon)^{2}(rg_{r,3}')_{ik}g_{r,3}^{k\ell}(rg_{r,3}')_{\ell j}$$

$$+ \frac{1}{2r} \operatorname{tr}_{g_{r,3}}(g_{r,3}')(g_{3})_{ij} - \alpha \left(\frac{2}{r^{2}}(g_{3})_{ij} + (1 - \epsilon)^{2}(rg_{r,3}')_{ij}\right) + 2((g_{S^{3}})_{ij} - (g_{r,3})_{ij}),$$

$$(\operatorname{Ric}_{g_{3}})_{\alpha\beta} = \left[\frac{1}{\delta^{2}r^{2\alpha}} - \frac{\alpha}{r^{2}}(2(1 + \alpha) + \frac{1}{2}\operatorname{tr}_{g_{r,3}}(rg_{r,3}'))\right](g_{3})_{\alpha\beta}.$$

We may now use the bounds (100) above to estimate the components of Ric_{g_3} by

(102)
$$(\operatorname{Ric}_{g_3})_{rr} > 2\frac{\alpha(1-\alpha)}{r^2} - C(4)\eta, \quad (\operatorname{Ric}_{g_3})_{ij} > \left(2\frac{\epsilon}{r^2} - C\left(\frac{\alpha}{r^2} + C(4)\eta\right)\right)(g_3)_{ij},$$
$$|(\operatorname{Ric}_{g_3})_{ir}|_{(g_3)_{ij}} < C(4)\eta, \qquad (\operatorname{Ric}_{g_3})_{\alpha\beta} > \left(\frac{1}{\delta^2 r^{2\alpha}} - C\left(\frac{\alpha}{r^2} + C(4)\eta\right)\right)(g_3)_{\alpha\beta}.$$

We see from these estimates that requiring $\alpha \leq C\epsilon$, $r_3 \leq r_3(\alpha, \epsilon, \lambda)$ and $\delta < \delta(\hat{r}, |\lambda|)$ small enough guarantees that $\operatorname{Ric}_{g_3} > \lambda$ for the range $\hat{r}/2 \leq r \leq 2r_3$ under consideration. Note that without loss of generality these requirements on α and δ (by way of $\hat{\delta}$) held when they were chosen in Lemma 5.4. \Box

5.4 Proof of Lemma 5.1

By construction, the metric g_3 with inner radius \hat{r} subject to $\hat{r} < r_3 = r_3(\alpha, \lambda, \epsilon)$ from Lemma 5.5 satisfies $\operatorname{Ric}_{\widehat{g}} > \lambda - C(4)\epsilon$ on $(B_2(p) \setminus B_{\widehat{r}/2}(p)) \times S^2$ and all the conditions of Lemma 5.1, except that it is not smooth. This metric is globally C^1 away from p, but fails to be smooth at $r \in \{0, r_2, r_2^+\}$. The singularity at 0 is as described in the statement of Lemma 5.1, and the other two radii can be smoothed within the class of warped product metrics to a metric $\hat{g} = \hat{g}_B + \hat{f}^2 g_{S^2}$ while preserving the Ric bound by the C^1 gluing of Lemma 3.1.

The last remaining statement of Lemma 5.1 to be proved is that the identity map Id: $(B_2(p), \hat{g}_B) \rightarrow$ $(B_2(p), g_B)$ is $(1+2\epsilon)$ -bi-Lipschitz. As the smooth metric \hat{g} can be made C^1 close to g_3 , we can prove a slightly stronger estimate directly for g_3 , and the result will follow for our final smooth metric as in Lemma 3.1.

The bi-Lipschitz condition will follow from the following metric comparisons for different ranges of r, which themselves follow by construction and the estimate $||g_{r,3} - g_r||_{C^0(S^3, g_r)} \le C \eta r^2$:

(103)
$$\begin{array}{c} g_B \\ (1-\epsilon)^2 g_B \\ (1-\epsilon)^2 (1-C(4)\eta r^2) g_B \end{array} \right\} \leq g_{B,3} \leq \begin{cases} g_B & \text{if } r \in (1,2], \\ g_B & \text{if } r \in (2r_3,1], \\ (1+C(4)\eta r^2) g_B & \text{if } r \in (0,2r_3]. \end{cases}$$

By again requiring that $r_3 \leq r_3(\epsilon)$ in Lemma 5.5, we can simplify the above by replacing all lower and upper bounds by $(1 \pm \frac{3}{2}\epsilon)^2 g_B$, respectively. Estimating the bi-Lipschitz constant of ϕ then becomes a matter of unraveling definitions as

(104)
$$(1-\frac{3}{2}\epsilon)|D\phi(v)|_{g_B} = (1-\frac{3}{2}\epsilon)|v|_{g_B} \le |v|_{g_{B,3}} \le (1+\frac{3}{2}\epsilon)|D\phi(v)|_{g_B}$$
 for all $v \in (TB_2(p), g_{B,3})$,
which completes the proof.

which completes the proof.

References

- [Anderson 1990] MT Anderson, Convergence and rigidity of manifolds under Ricci curvature bounds, Invent. Math. 102 (1990) 429-445 MR Zbl
- [Bruè et al. 2022] E Bruè, A Naber, D Semola, Boundary regularity and stability for spaces with Ricci bounded below, Invent. Math. 228 (2022) 777-891 MR Zbl
- [Cheeger and Colding 1996] J Cheeger, TH Colding, Lower bounds on Ricci curvature and the almost rigidity of warped products, Ann. of Math. 144 (1996) 189-237 MR Zbl
- [Cheeger and Colding 1997] J Cheeger, TH Colding, On the structure of spaces with Ricci curvature bounded below, I, J. Differential Geom. 46 (1997) 406-480 MR Zbl

- [Cheeger et al. 2021] J Cheeger, W Jiang, A Naber, *Rectifiability of singular sets of noncollapsed limit spaces with Ricci curvature bounded below*, Ann. of Math. 193 (2021) 407–538 MR Zbl
- [Colding and Naber 2012] **TH Colding**, **A Naber**, *Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications*, Ann. of Math. 176 (2012) 1173–1229 MR Zbl
- [DeTurck and Kazdan 1981] **DM DeTurck**, **JL Kazdan**, *Some regularity theorems in Riemannian geometry*, Ann. Sci. École Norm. Sup. 14 (1981) 249–260 MR Zbl
- [Federer 1970] H Federer, The singular sets of area minimizing rectifiable currents with codimension one and of area minimizing flat chains modulo two with arbitrary codimension, Bull. Amer. Math. Soc. 76 (1970) 767–771 MR Zbl
- [Gromov 1999] **M Gromov**, *Metric structures for Riemannian and non-Riemannian spaces*, Progr. Math. 152, Birkhäuser, Boston, MA (1999) MR Zbl
- [Jost 2017] J Jost, Riemannian geometry and geometric analysis, 7th edition, Springer (2017) MR Zbl
- [Li and Naber 2020] N Li, A Naber, *Quantitative estimates on the singular sets of Alexandrov spaces*, Peking Math. J. 3 (2020) 203–234 MR Zbl
- [Menguy 2000] **X Menguy**, *Noncollapsing examples with positive Ricci curvature and infinite topological type*, Geom. Funct. Anal. 10 (2000) 600–627 MR Zbl
- [Otsu and Shioya 1994] Y Otsu, T Shioya, *The Riemannian structure of Alexandrov spaces*, J. Differential Geom. 39 (1994) 629–658 MR Zbl
- [Pan and Wei 2022] **J Pan**, **G Wei**, *Examples of Ricci limit spaces with non-integer Hausdorff dimension*, Geom. Funct. Anal. 32 (2022) 676–685 MR Zbl
- [Perelman 1997] G Perelman, Construction of manifolds of positive Ricci curvature with big volume and large Betti numbers, from "Comparison geometry" (K Grove, P Petersen, editors), Math. Sci. Res. Inst. Publ. 30, Cambridge Univ. Press (1997) 157–163 MR Zbl

Department of Mathematics, Northwestern University Evanston, IL, United States Institute for Advanced Study Princeton, NJ, United States Department of Mathematics, Northwestern University Evanston, IL, United States

hupp@iam.uni-bonn.de, anaber@ias.edu, khwang2025@u.northwestern.edu

Proposed: Tobias H Colding Seconded: Dmitri Burago, Bruce Kleiner Received: 3 September 2023 Revised: 2 February 2024



GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITORS

Robert Lipshitz

University of Oregon lipshitz@uoregon.edu

András I Stipsicz

Alfréd Rényi Institute of Mathematics

stipsicz@renyi.hu

BOARD OF EDITORS

Mohammed Abouzaid	Stanford University abouzaid@stanford.edu	Mark Gross	University of Cambridge mgross@dpmms.cam.ac.uk
Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Arend Bayer	University of Edinburgh arend.bayer@ed.ac.uk	Sándor Kovács	University of Washington skovacs@uw.edu
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	Haynes Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Aaron Naber	Northwestern University anaber@math.northwestern.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Peter Ozsváth	Princeton University petero@math.princeton.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Benson Farb	University of Chicago farb@math.uchicago.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David M Fisher	Rice University davidfisher@rice.edu	Roman Sauer	Karlsruhe Institute of Technology roman.sauer@kit.edu
Mike Freedman	Microsoft Research michaelf@microsoft.com	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
David Gabai	Princeton University gabai@princeton.edu	Natasa Sesum	Rutgers University natasas@math.rutgers.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Cameron Gordon	University of Texas gordon@math.utexas.edu	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2025 is US \$865/year for the electronic version, and \$1210/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY



© 2025 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 29 Issue 1 (pages 1–548) 2025	
Helly groups	1
Jérémie Chalopin, Victor Chepoi, Anthony Genevois, Hiroshi Hirai and Damian Osajda	
Topologically trivial proper 2-knots	71
Robert E Gompf	
The stable Adams operations on Hermitian K-theory	127
JEAN FASEL and OLIVIER HAUTION	
On Borel Anosov subgroups of $SL(d, \mathbb{R})$	171
SUBHADIP DEY	
Global Brill–Noether theory over the Hurwitz space	193
ERIC LARSON, HANNAH LARSON and ISABEL VOGT	
Hyperbolic hyperbolic-by-cyclic groups are cubulable	259
FRANÇOIS DAHMANI, SURAJ KRISHNA MEDA SATISH and JEAN PIERRE MUTANGUHA	
The smooth classification of 4-dimensional complete intersections	269
DIARMUID CROWLEY and CSABA NAGY	
An embedding of skein algebras of surfaces into localized quantum tori from Dehn–Thurston coordinates	313
RENAUD DETCHERRY and RAMANUJAN SANTHAROUBANE	
Virtual classes via vanishing cycles	349
TASUKI KINJO	
On termination of flips and exceptionally noncanonical singularities	399
JINGJUN HAN and JIHAO LIU	
Lower Ricci curvature and nonexistence of manifold structure	443
ERIK HUPP, AARON NABER and KAI-HSIANG WANG	
Independence of singularity type for numerically effective Kähler–Ricci flows	479
HOSEA WONDO and ZHOU ZHANG	
Subgroups of genus-2 quasi-Fuchsian groups and cocompact Kleinian groups	495
ZHENGHAO RAO	