



Geometry & Topology

Volume 29 (2025)

Asymptotically Calabi metrics and weak Fano manifolds

HANS-JOACHIM HEIN

SONG SUN

JEFFREY VIACLOVSKY

RUOBING ZHANG

Asymptotically Calabi metrics and weak Fano manifolds

HANS-JOACHIM HEIN

SONG SUN

JEFFREY VIACLOVSKY

RUOBING ZHANG

We show that any asymptotically Calabi manifold which is Calabi–Yau can be compactified complex analytically to a weak Fano manifold. Furthermore, the Calabi–Yau structure arises from a generalized Tian–Yau construction on the compactification, and we prove a strong uniqueness theorem. We also give an application of this result to the surface case.

53C21, 53C25, 53C55; 14J32, 14J45

1 Introduction

This paper is concerned with the Tian–Yau construction [35] of complete Ricci-flat Kähler metrics on the complement of a smooth anticanonical divisor in a smooth Fano manifold. We begin by describing their geometry at infinity.

1.1 Asymptotically Calabi metrics

We begin by defining the asymptotic models for the metrics constructed in [35]. Let D be an $(n-1)$ -dimensional compact Kähler manifold with trivial canonical bundle and let $L \rightarrow D$ be an ample line bundle. Define

$$(1-1) \quad \deg(L) = \int_D c_1(L)^{n-1},$$

and fix a nowhere vanishing holomorphic $(n-1)$ -form Ω_D on D satisfying

$$(1-2) \quad \frac{1}{2} \int_D (\sqrt{-1})^{(n-1)^2} \Omega_D \wedge \bar{\Omega}_D = (2\pi c_1(L))^{n-1}.$$

Using Yau’s resolution [37] of the Calabi conjecture, there exists a unique Ricci-flat Kähler metric $\omega_D \in 2\pi c_1(L)$ with

$$(1-3) \quad \omega_D^{n-1} = \frac{1}{2} (\sqrt{-1})^{(n-1)^2} \Omega_D \wedge \bar{\Omega}_D.$$

There exists a unique hermitian metric h on L whose curvature form is $-\sqrt{-1}\omega_D$, up to scaling. Fixing a choice of h , the *Calabi model space* is the subset \mathcal{C} of L consisting of all elements ξ with $0 < |\xi|_h < 1$. We next define a nowhere vanishing holomorphic volume form $\Omega_{\mathcal{C}}$ and a Ricci-flat Kähler metric $\omega_{\mathcal{C}}$ which is incomplete as $|\xi|_h \rightarrow 1$ and complete as $|\xi|_h \rightarrow 0$. Let $p: \mathcal{C} \rightarrow D$ denote the bundle projection and Z be the holomorphic vector field generating the natural \mathbb{C}^* -action on the fibers of p . The holomorphic volume form $\Omega_{\mathcal{C}}$ is uniquely determined by the equation

$$(1-4) \quad Z \lrcorner \Omega_{\mathcal{C}} = p^* \Omega_D,$$

and the metric $\omega_{\mathcal{C}}$ is given by the *Calabi ansatz*

$$(1-5) \quad \omega_{\mathcal{C}} = \frac{n}{n+1} \sqrt{-1} \partial \bar{\partial} (-\log |\xi|_h^2)^{(n+1)/n},$$

which satisfies the complex Monge–Ampère equation

$$\omega_{\mathcal{C}}^n = \frac{1}{2} (\sqrt{-1})^{n^2} \Omega_{\mathcal{C}} \wedge \bar{\Omega}_{\mathcal{C}},$$

hence is Ricci-flat. The function $z = (-\log |\xi|_h^2)^{1/n}$ is the $\omega_{\mathcal{C}}$ -moment map for the natural S^1 -action on L . It is easily verified that the $\omega_{\mathcal{C}}$ -distance function r to a fixed point in \mathcal{C} satisfies

$$(1-6) \quad r^{-1} z^{(n+1)/2} = C + o(1) \quad \text{as } z \rightarrow \infty.$$

We will consider Kähler manifolds with trivial canonical bundle which will be denoted by (X, I, ω, Ω) , where X is a smooth manifold of real dimension $2n$, I is a complex structure, ω is a Kähler form, and Ω is a holomorphic volume form. We say that (X, I, ω, Ω) is a *Calabi–Yau structure* if it satisfies the complex Monge–Ampère equation

$$(1-7) \quad \omega_X^n = \tau^n \cdot \frac{1}{2} (\sqrt{-1})^{n^2} \cdot \Omega_X \wedge \bar{\Omega}_X$$

for some constant $\tau > 0$. We are interested in Calabi–Yau structures which are asymptotic to the above Calabi model spaces. More precisely, we make the following definition.

Definition 1.1 A Calabi–Yau structure $(X, I_X, \omega_X, \Omega_X)$ is *asymptotically Calabi* if there exists $\underline{\delta} > 0$, a Calabi model space \mathcal{C} , and a diffeomorphism $\Phi: \mathcal{C} \setminus K' \rightarrow X \setminus K$, where $K \subset X$ is compact and $K' = \{|\xi|_h \geq \frac{1}{2}\}$, such that for all $k \in \mathbb{N}_0$, the following hold uniformly as $z \rightarrow +\infty$:

$$(1-8) \quad |\nabla_{g_{\mathcal{C}}}^k (\Phi^* I_X - I_{\mathcal{C}})|_{g_{\mathcal{C}}} = O(e^{-\underline{\delta} z^{n/2}}),$$

$$(1-9) \quad |\nabla_{g_{\mathcal{C}}}^k (\Phi^* \Omega_X - \Omega_{\mathcal{C}})|_{g_{\mathcal{C}}} = O(e^{-\underline{\delta} z^{n/2}}),$$

$$(1-10) \quad |\nabla_{g_{\mathcal{C}}}^k (\Phi^* \omega_X - \omega_{\mathcal{C}})|_{g_{\mathcal{C}}} = O(e^{-\underline{\delta} z^{n/2}}).$$

Remark 1.2 The decay assumption on the error terms is the natural one since harmonic functions on the Calabi model space have this type of decay; see [19; 32].

Our main theorem is the following.

Theorem 1.3 Any Calabi–Yau structure (X, I, ω, Ω) which is asymptotically Calabi can be compactified complex analytically to a weak Fano manifold \bar{X} . Furthermore, the Calabi–Yau structure arises from a generalized Tian–Yau construction on \bar{X} and ω is the unique Calabi–Yau metric with respect to (I, Ω) satisfying (1-10) and representing $[\omega] \in H^2(X)$.

The above theorem solves a particular case of Yau’s conjecture in [38]. We give a precise description of this generalized Tian–Yau construction in Section 2. The main step in the proof is to show that for a certain choice of a parallel complex structure I , the underlying complex manifold of an Calabi–Yau asymptotically Calabi metric can be compactified to a weak Fano manifold. This involves producing holomorphic functions with controlled growth at infinity, which is typically done using weighted Fredholm theory. This strategy runs into trouble due to the fact that, in the Tian–Yau construction, the decay rate of the metric is much slower than the decay rate of the complex structure. To overcome this difficulty, our strategy is based on the L^2 -estimates in several complex variables pioneered by Hörmander [21]. The proof of Theorem 1.3 can be found in Section 2. We also have the following corollary. Recall the index of \bar{X} is the largest integer k such that $K_{\bar{X}}^{-1} = H^k$ for some line bundle H .

Corollary 1.4 Let (X^n, I, ω, Ω) be a Calabi–Yau structure which is asymptotically Calabi. Then $\pi_1(X)$ is a cyclic group with order the index of \bar{X} . Furthermore, there exists a constant $C(n)$, depending only upon n , such that $\deg(L) \leq C(n)$.

This degree bound is remarkable because in Definition 1.1, $\deg(L)$ could a priori be any integer, but if \mathcal{C} occurs at infinity for a Calabi–Yau asymptotically Calabi metric, then $\deg(L)$ must be bounded. The proof of this degree bound uses Theorem 1.3 and deep results in birational geometry.

1.2 The surface case

A gravitational instanton is by definition a complete noncompact hyperkähler 4-manifold (X, g, ω) with square-integrable curvature. By the results of the recent paper by Sun and Zhang [33], a gravitational instanton is always asymptotic to a model end. Accordingly, gravitational instantons can be classified into 6 families: ALE, ALF, ALG, ALH, ALG*, ALH*. There has been extensive work, much of it quite recent, on classifying the 6 families completely; see Chen and Chen [3; 2; 4], Chen and Viaclovsky [5], Kronheimer [25] and Minerbe [27]. The results in this paper are relevant to the ALH* family. This has the unique intriguing feature that its members have fractional asymptotic volume growth; indeed, the volume growth exponent of an ALH* model end is $\frac{4}{3}$. Gravitational instantons of type ALH* also appear as singularity models in polarized degenerations of K3 surfaces; see Hein, Sun, Viaclovsky and Zhang [19] and Sun and Zhang [31]. Their precise definition can be found in Section 3.

There are two known mathematical constructions of ALH* gravitational instantons. Both of them come with a preferred choice of complex structure, and are based on solving a complex Monge–Ampère equation on a quasiprojective surface with trivial canonical bundle. First, we have the Tian–Yau construction [35], which involves the complement of a smooth anticanonical divisor in a del Pezzo surface. Second, we

have the construction of Hein [18], which involves the complement of a singular fiber of Kodaira type I_b in a rational elliptic surface. Our next theorem relates the complex structures involved in these two constructions.

Theorem 1.5 *Let (X, g, ω) be an ALH* gravitational instanton.*

- (i) *Letting I denote the complex structure corresponding to ω_1 , then (X, I) is biholomorphic to weak del Pezzo surface \bar{X} minus a smooth anticanonical elliptic curve. Furthermore, the hyperkähler structure arises from a generalized Tian–Yau construction on this compactification and is the unique Calabi–Yau metric with respect to $(I, \Omega = \omega_2 + \sqrt{-1}\omega_3)$ satisfying (1-10) and representing $[\omega_1] \in H^2(X)$.*
- (ii) *Letting J denote the complex structure corresponding to I_2 , then (X, J) compactifies to a rational elliptic surface S with a global section by adding F , a Kodaira type I_b fiber of multiplicity 1. The 2-form $\Omega = \omega_2 + \sqrt{-1}\omega_3$ is a rational 2-form on S with a simple pole along F .*

The proof of Theorem 1.5 will be given in Section 3 and is our originally intended proof of some claims made in [19, Remark 2.4]. The main ingredients in the proof are Theorem 1.3, the decay estimates of [19, Section 3], and the analysis of harmonic functions on asymptotically Calabi spaces of [19, Section 4]. We also note that in the meantime Collins, Jacob and Lin [8, Theorem 1.3] have proved, using an entirely different method, that a Tian–Yau space can be compactified to a rational elliptic surface.

Acknowledgements The authors thank Gao Chen for valuable discussions on gravitational instantons. Hein was supported by NSF grant DMS-1745517 and by the DFG under Germany’s Excellence Strategy EXC 2044-390685587 *Mathematics Münster: Dynamics-Geometry-Structure* as well as by the CRC 1442 *Geometry: deformations and rigidity* of the DFG. Sun was supported by the Simons Collaboration on *Special holonomy in geometry, analysis and physics* (#488633), and NSF grant DMS-2004261. Viaclovsky was partially supported by NSF grants DMS-1811096 and DMS-2105478. Zhang was partially supported by NSF grants DMS-1906265 and DMS-2304818. Finally, the authors thank the referee for their careful reading of the paper, and for their questions and comments which improved the exposition of the paper.

2 Compactification to weak Fano manifold

In this section, we give the proof of Theorem 1.3. Let (X, I, ω, Ω) be a complete Calabi–Yau manifold that is asymptotically Calabi. We identify $X \setminus K$ smoothly with a Calabi model space $\mathcal{C} \setminus K'$, where $K' = \{z \leq z_0\}$ for some $z_0 \gg 0$, and assume that (1-8)–(1-10) are satisfied. Let $\phi_0 \equiv z^n - \delta z^{n/2}$, for some $\delta \in (0, \frac{1}{2})$ to be chosen later. Then

$$\begin{aligned}
 (2-1) \quad dd_{\mathbb{C}}^c \phi_0 &= -dJ_{\mathbb{C}} d\phi_0 = -d\left(\left(nz^{n-1} - \frac{1}{2}n\delta z^{n/2-1}\right)J_{\mathbb{C}} dz\right) \\
 &= d\left(\left(n - \frac{1}{2}n\delta z^{-n/2}\right)\theta\right) \\
 &= \left(n - \frac{1}{2}n\delta z^{-n/2}\right) d\theta + \delta\left(\frac{1}{2}n\right)^2 z^{-n/2-1} dz \wedge \theta,
 \end{aligned}$$

where $\theta \equiv -z^{n-1} J_\epsilon dz$. Note also that

$$(2-2) \quad \omega_D = \sqrt{-1} \partial \bar{\partial} (-\log \|\xi\|_h^2) = \sqrt{-1} \partial \bar{\partial} (z^n) = \frac{1}{2} n d\theta,$$

$$(2-3) \quad \begin{aligned} \omega_\epsilon &= \frac{n}{n+1} \frac{1}{2} d d_\epsilon^c z^{n+1} = -\frac{n}{2(n+1)} d J_\epsilon dz^{n+1} \\ &= -\frac{1}{2} n d(z^n J dz) = \frac{1}{2} n d(z\theta) \\ &= \frac{1}{2} n (z d\theta + dz \wedge \theta). \end{aligned}$$

Then for $z_1 \gg z_0$,

$$(2-4) \quad \begin{aligned} d d_\epsilon^c \phi_0 &= z^{-n/2-1} \left((nz^{n/2+1} - \frac{1}{2} n \delta z) d\theta + \delta (\frac{1}{2} n)^2 dz \wedge \theta \right) \\ &= z^{-n/2-1} \left((nz^{n/2+1} - \frac{1}{2} n \delta z - (n/2)^2 \delta z) d\theta + \delta (\frac{1}{2} n)^2 (z d\theta + dz \wedge \theta) \right) \\ &\geq z^{-n/2-1} \frac{1}{2} n \delta \omega_\epsilon, \end{aligned}$$

since $\frac{1}{2} n d\theta = \omega_D$ is positive definite on D . Let $A = z_1^n - \delta z_1^{n/2}$, and choose a smooth increasing and convex function $u: \mathbb{R} \rightarrow \mathbb{R}$ such that $u(t) = 2A/3$ for $t \leq A/2$ and $u(t) = t$ for $t \geq A$. Write $\phi_1 \equiv u \circ \phi_0$. Then

$$(2-5) \quad \begin{aligned} d d_\epsilon^c \phi_1 &= -d J_\epsilon d(u \circ \phi_0) = -d J_\epsilon u'(\phi_0) d\phi_0 \\ &= -u''(\phi_0) d\phi_0 \wedge J_\epsilon d\phi_0 + u'(\phi_0) d d_\epsilon^c \phi_0. \end{aligned}$$

Since the form $-d\phi_0 \wedge J_\epsilon d\phi_0$ is positive semidefinite, we see that $d d_\epsilon^c \phi_1 \geq 0$ for all $z \geq z_0$ and $d d_\epsilon^c \phi_1 \geq C \delta z^{-n/2-1} \omega_\epsilon$ for $z > z_1$.

Notice that ϕ_1 can be naturally viewed as a smooth function on X , and it satisfies $d d_I^c \phi_1 \geq P(z)\omega$ for a nonnegative function $P(z)$ with $P(z) \geq C \delta z^{-n/2-1}$ when z is large. In particular, we know that (X, I) is 1-convex. So, by [13, Section 2], there is a Remmert reduction $\pi: X \rightarrow \tilde{X}$, where \tilde{X} is Stein and $\text{Sing}(\tilde{X})$ is a finite set contained in the region $\{z \leq z_1\}$. Then \tilde{X} admits an exhaustion function $\psi_{\tilde{X}}$ which is smooth on $\tilde{X} \setminus \text{Sing}(\tilde{X})$ and satisfies $d d_I^c \psi_{\tilde{X}} > 0$. Denote $\psi_X \equiv \pi^* \psi_{\tilde{X}}$. Then $d d_I^c \psi_X > 0$ on $X \setminus E$, where $E \equiv \pi^{-1}(\text{Sing}(\tilde{X}))$. Choose a cutoff function χ on X supported in $\{z \leq z_1 + 1\}$ with $\chi \equiv 1$ on $\{z \leq z_1\}$. Then let $\phi \equiv \epsilon \chi \psi_X + \phi_1$ for a fixed $0 < \epsilon \ll 1$. The above calculation shows that ϕ also satisfies $d d_I^c \phi \geq P(z)\omega$, where $P(z) \geq 0$ on X , $P(z) > 0$ on $X \setminus E$, and $P(z) \geq C \delta z^{-n/2-1}$ outside a compact set.

Any holomorphic section $s \in H^0(D, L^k)$ gives rise to a holomorphic function f_s on $L \setminus \mathbf{0}_L$ by defining

$$(2-6) \quad f_s(\xi) = s(\pi(\xi))/\xi^{\otimes k},$$

where $\xi \in L \setminus \mathbf{0}_L$, and $\pi: L \rightarrow D$ is the bundle projection. In particular, f_s restricts to a holomorphic function on \mathcal{C} . Taking the logarithm of (2-6), and using that $z^n = -2 \log \|\xi\|_h$, we see that $f_s = \tilde{f}_s e^{(k/2)z^n}$, where \tilde{f}_s is a function on the unit circle bundle of L .

Lemma 2.1 For all $l \geq 0$ and any $\varepsilon > 0$, we have $|\nabla_{g_\varepsilon}^l f_s|_{g_\varepsilon} = O(e^{(k/2)z^n + \varepsilon z^{n/2}})$ as $z \rightarrow \infty$.

Proof The estimate for $l = 0$ holds with $\varepsilon = 0$, and follows from the previous remarks. Since f_s is holomorphic, it is harmonic. The curvature of the Calabi metric is in particular bounded at infinity; see [35, Lemma 4.3] and [19, Remark 3.2]. Given $p \in \mathcal{C}$, let $B_1(p)$ be a unit ball around p , and $B_1(\hat{p})$ be a unit ball in its universal cover. Then standard elliptic estimates for harmonic functions yield that

$$(2-7) \quad |\nabla^l f_s|(p) \leq \|\nabla^l f_s\|_{C^0(B_{1/2}(\hat{p}))} \leq C_l \|f_s\|_{C^0(B_1(\hat{p}))}.$$

Using (1-6), one can easily check that if $|r(q) - r(p)| < 1$, then

$$(2-8) \quad |z(q)^n - z(p)^n| \leq C z(p)^{(n-1)/2}.$$

Since $(n - 1)/2 < n/2$, the claim follows. □

Denote by $\mathcal{O}(X)$ the space of I -holomorphic functions on X .

Proposition 2.2 There is an injective linear map $\mathcal{L}: \bigoplus_{k=0}^\infty H^0(D, L^k) \rightarrow \mathcal{O}(X)$ such that for any nonzero section $s \in H^0(D, L^k)$, we have that

$$(2-9) \quad |\nabla_{\Phi^*g}^l (\mathcal{L}(s) \circ \Phi - f_s)|_{\Phi^*g} = O(e^{(k/2)z^n - (\delta_k/2)z^{n/2}})$$

for all $l \geq 0$ and for some $\delta_k > 0$ as $z \rightarrow \infty$.

Proof In the following proof, for simplicity of notation we will omit the diffeomorphism Φ . Fix a cutoff function χ on \mathcal{C} which is equal to 1 when $z \geq z_1 + 1$ and vanishes when $z \leq z_1$. Then for any $s \in H^0(D, L^k)$, the function χf_s naturally extends to a smooth function on X . Notice that for $z > z_0 + 1$ we have that

$$(2-10) \quad |\bar{\partial}_I(\chi f_s)|_g = |\bar{\partial}_I f_s|_g = |(\bar{\partial}_I - \bar{\partial}_{\mathcal{C}})f_s|_g \leq e^{(k/2)z^n - \delta z^{n/2}}.$$

Set $\delta \equiv \min(\delta/(4k), \frac{1}{4})$ in the definition of ϕ_0 above. Then we have that

$$(2-11) \quad \int_{X \setminus E} \frac{1}{P(z)} |\bar{\partial}_I(\chi f_s)|_g^2 e^{-k\phi} d\text{Vol}_g < \infty.$$

Notice that $X \setminus E \cong \tilde{X} \setminus \text{Sing}(\tilde{X})$ admits a complete Kähler metric; see [29, Proposition 4.1]. Also, by assumption K_X is trivial, so we can apply the standard L^2 -estimates for the $\bar{\partial}$ -operator on $X \setminus E$ — see for example [11, Chapter VIII.6, Theorem 6.1] — to find a unique solution u to the equation $\bar{\partial}_I u = \bar{\partial}_I(\chi f_s)$ with

$$(2-12) \quad \int_{X \setminus E} |u|^2 e^{-k\phi} d\text{Vol}_g \leq \int_{X \setminus E} \frac{1}{P(z)} |\bar{\partial}_I(\chi f_s)|_g^2 e^{-k\phi} d\text{Vol}_g$$

such that u is L^2 orthogonal to $\ker(\bar{\partial}_I)$. Notice that

$$(2-13) \quad \Delta_g u = \bar{\partial}_{I,g}^* \bar{\partial}_I(\chi f_s) = O(e^{(k/2)z^n - (\delta/2)z^{n/2}}).$$

Similar to the proof of Lemma 2.1, it follows from local L^2 elliptic estimates that for all $l \geq 0$,

$$(2-14) \quad |\nabla_g^l u|_g = O(e^{(k/2)z^n - \delta z^{n/2}}) \quad \text{as } z \rightarrow \infty.$$

Indeed, by straightforward computations, one can find a $C > 0$ such that $\text{Vol}_g(B_1(p)) \geq C \cdot z^{-(n-1)/2}(p)$. Therefore using (2-8) and (2-12), there exists a uniform constant $C > 0$ such that for any $z(p) \gg 1$,

$$(2-15) \quad \int_{B_1(p)} |u|^2 \, d\text{vol}_g \leq C e^{k \cdot z(p)^n - \delta' \cdot z(p)^{n/2}},$$

where $\delta' > 0$ is any smaller number than $\underline{\delta}$. By an obvious volume comparison argument, we obtain a similar estimate for the lifting of u to $B_{1/2}(\widehat{p})$ in the local universal cover $\widehat{B_1(p)}$ of $B_1(p)$. Since $\widehat{B_1(p)}$ is noncollapsed with bounded curvature, standard elliptic theory applied to the lifting of (2-13) yields (2-14).

To finish the proof, we let $\mathcal{L}(s) \equiv \chi f_s - u$. This function is holomorphic away from E , so by Hartogs's theorem (applied to \widetilde{X}) one can see that it is globally holomorphic on X . The conclusion then follows. \square

Fix k such that $L^l|_D$ is very ample for all $l \geq k$. Then we have a holomorphic embedding

$$(2-16) \quad F_k: L \rightarrow \mathbb{P}(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*), \quad (x, \xi) \mapsto (\text{ev}_{x,k} / \xi^{\otimes k}, \text{ev}_{x,k+1} / \xi^{\otimes(k+1)}),$$

where $x \in D$, $\xi \in L_x$ and $\text{ev}_{x,l}: H^0(D, L^l) \rightarrow L^l|_x$ is the evaluation map. Alternatively, we can describe F_k as follows. By assumption, we have the embeddings

$$(2-17) \quad i_{L^k}: D \rightarrow \mathbb{P}(H^0(D, L^k)^*), \quad i_{L^{k+1}}: D \rightarrow \mathbb{P}(H^0(D, L^{k+1})^*).$$

We can view i_{L^k} and $i_{L^{k+1}}$ as mapping into $\mathbb{P}(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*)$. Then for $x \in D$, F_k maps the fiber $\pi^{-1}(x) \subset L$ linearly to the line between $i_{L^k}(\pi(p))$ and $i_{L^{k+1}}(\pi(p))$, so it is clearly an embedding. Obviously, $F_k(\mathbf{0}_L)$ is isomorphic to D and is contained in the linear subspace $\mathbb{P}(H^0(D, L^{k+1})^*) \subset \mathbb{P}(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*)$.

Now we define a holomorphic map $G_k: X \rightarrow \mathbb{P}(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*)$ via

$$(2-18) \quad p \in X \mapsto (\widetilde{\text{ev}}_{p,k}, \widetilde{\text{ev}}_{p,k+1}),$$

where $\widetilde{\text{ev}}_{p,k}: H^0(D, L^k) \rightarrow \mathbb{C}$ is given by $\widetilde{\text{ev}}_{p,k}(s) = \mathcal{L}(s)(p)$.

We denote by \overline{X} the topological compactification of X by adding $F_k(D)$ to the end of $G_k(X)$. This is justified by the following.

Proposition 2.3 *There exists a compact set $K \subset X$ such that G_k is a holomorphic embedding on $X \setminus K$ with $G_k(X \setminus K) \cap F_k(D) = \emptyset$. Furthermore there is a neighborhood of $F_k(D)$ in \overline{X} which is homeomorphic to a neighborhood of $F_k(\mathbf{0}_L)$ in $F_k(L)$.*

Proof The key point is that we can compare G_k with F_k via the fixed embedding of the end of X into $\mathcal{C} \subset L$. Given any point $q_0 \in \mathbf{0}_L \cong D$, we can find sections $s_0, \dots, s_{n-1} \in H^0(D, L^k)$ and $s_n \in H^0(D, L^{k+1})$ such that $s_0(q_0) \neq 0$, $s_1(q_0) = \dots = s_{n-1}(q_0) = 0$, $ds_1(q_0), \dots, ds_{n-1}(q_0)$ are linearly independent, and $s_n(q_0) \neq 0$. Then $w_k \equiv s_k/s_0$ for $1 \leq k \leq n-1$ and $w_n \equiv s_0/s_n$ form local holomorphic coordinates in a neighborhood U of q_0 in L . To clarify this definition, note that w_n is

a local section of L^{-1} on D , but by duality we can view such a section as a local function on the total space of L which is linear on fibers. We therefore can think of w_1, \dots, w_{n-1} as coordinates on the divisor, and w_n as a fiber coordinate. Denote this coordinate system by $w: U \rightarrow \mathbb{C}^n$. Notice that $|\xi|_{h_L}^2 = |w_n|^2 e^{-\varphi(w_1, \dots, w_{n-1})}$ for a smooth function φ satisfying $\sqrt{-1} \partial \bar{\partial} \varphi = \omega_D$. The Calabi metric in the (w_1, \dots, w_n) coordinates is given by

$$(2-19) \quad \omega_\xi = (-\log |\xi|_{h_L}^2)^{1/n} \omega_D + 1/n (-\log |\xi|_{h_L}^2)^{(1/n)-1} \cdot \sqrt{-1} \cdot \left(\frac{dw_n}{w_n} - \partial \varphi \right) \wedge \left(\frac{d\bar{w}_n}{\bar{w}_n} - \bar{\partial} \varphi \right).$$

We define a mapping Ψ from an open subset $V \subset P(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*)$ containing $F_k(q_0)$ to \mathbb{C}^n by

$$(2-20) \quad [(\alpha, \beta)] \mapsto \left(\frac{\alpha(s_1)}{\alpha(s_0)}, \dots, \frac{\alpha(s_{n-1})}{\alpha(s_0)}, \frac{\alpha(s_0)}{\beta(s_n)} \right).$$

Then the restriction of Ψ to $\text{Image}(F_k)$ is a coordinate chart for $\text{Image}(F_k)$ near $F_k(q_0)$, and $\Psi \circ F_k = w$ in a neighborhood of $q_0 \in \mathbf{0}_L$. Next, let $\eta_j = \mathcal{L}(s_j)/\mathcal{L}(s_0)$ for $1 \leq j \leq n-1$, and $\eta_n = \mathcal{L}(s_0)/\mathcal{L}(s_n)$, and we denote this mapping $\eta: \Phi(U \setminus \mathbf{0}_L) \rightarrow \mathbb{C}^n$ (after possibly shrinking U). Note that for any p in the domain of η , we have

$$(2-21) \quad \begin{aligned} \Psi(G_k(p)) &= \Psi([\tilde{e}_{v_{p,k}}, \tilde{e}_{v_{p,k+1}}]) = \left(\frac{\tilde{e}_{v_{p,k}}(s_1)}{\tilde{e}_{v_{p,k}}(s_0)}, \dots, \frac{\tilde{e}_{v_{p,k}}(s_0)}{\tilde{e}_{v_{p,k+1}}(s_n)} \right) \\ &= \left(\frac{\mathcal{L}(s_1)}{\mathcal{L}(s_0)}(p), \dots, \frac{\mathcal{L}(s_0)}{\mathcal{L}(s_n)}(p) \right) = \eta(p). \end{aligned}$$

By Proposition 2.2, on $U \setminus \mathbf{0}_L$ we have that $\eta \circ \Phi = w(1 + \zeta)$ with $|\nabla_g^l \zeta|_g = O(e^{-\delta' z^{n/2}})$ for all $l \geq 0$ for some $\delta' > 0$. In the following, the constant δ' is allowed to change from line to line. Using (2-19), we have that

$$(2-22) \quad \left(\frac{\partial(\eta \circ \Phi)_\alpha}{\partial w_\beta} \right) = \begin{pmatrix} \mathbb{I}_{n-1} + O(e^{-\delta' z^{n/2}}) & O(e^{-\delta' z^{n/2}} |w_n|^{-1}) \\ \dots & O(e^{-\delta' z^{n/2}} |w_n|) & \dots & 1 + O(e^{-\delta' z^{n/2}}) \end{pmatrix}.$$

Thus, the Jacobian matrix $(\partial(\eta \circ \Phi)_\alpha / \partial w_\beta)$ is nondegenerate for $z \gg 1$, which implies that G_k is an immersion outside a compact set.

Next, we show that G_k is injective onto its image for $z \gg 1$. Suppose we have two points $p_1, p_2 \in X$ with $G_k(p_1) = G_k(p_2)$, and $z(q_2) \geq z(q_1) \gg 1$, where $\Phi(q_j) = p_j$. Let d_{FS} denote the Fubini–Study distance on $\mathbb{P}(H^0(D, L^k)^* \oplus H^0(D, L^{k+1})^*)$. We claim that there exists $\varepsilon > 0$ such that $d_{\text{FS}}(F_k(q), G_k(\Phi(q))) < \varepsilon$ if $q \in L$ is sufficiently near D . To see this, using the formula

$$(2-23) \quad d_{\text{FS}}(Z, W) = \arccos \sqrt{\frac{\langle Z, W \rangle^2}{\|Z\|^2 \|W\|^2}}$$

and Proposition 2.2, we obtain that

$$(2-24) \quad d_{\text{FS}}(F_k(q), G_k(\Phi(q))) = \arccos(1 + O(e^{-\delta'z(q)^{n/2}})) \quad \text{as } z \rightarrow \infty,$$

since L^{k+1} is very ample. Then

$$(2-25) \quad d_{\text{FS}}(F_k(q_1), F_k(q_2)) \leq d_{\text{FS}}(F_k(q_1), G_k(p_1)) + d_{\text{FS}}(G_k(p_2), F_k(q_2)) < 2\varepsilon.$$

This implies that q_1 and q_2 must be contained in the same w coordinate patch above, and therefore $\eta_\alpha(p_1) = \eta_\alpha(p_2)$ for $\alpha = 1, \dots, n$. Denote $\tau_\alpha \equiv |w_\alpha(q_2) - w_\alpha(q_1)|$. Then $\tau_\alpha = w_\alpha(q_1)O(e^{-\delta'z(q_1)^{n/2}})$. So we know in particular that $|w_n(q_1)| \leq C|w_n(q_2)|$. Notice that

$$(2-26) \quad |\tau_\alpha| = |w_\alpha(q_2)\zeta_\alpha(q_2) - w_\alpha(q_1)\zeta_\alpha(q_1)| \leq Ce^{-\delta'z(q_1)^{n/2}}|\tau_\alpha| + |w_\alpha(q_1)| \cdot |\zeta_\alpha(q_2) - \zeta_\alpha(q_1)|.$$

Let q_t with $t \in [0, 1]$ be the straight line connecting q_1 and q_2 in the w coordinates. Then

$$(2-27) \quad |\zeta_\alpha(q_2) - \zeta_\alpha(q_1)| \leq \sum_{\beta=1}^n \sup_{t \in [0,1]} \left| \frac{\partial \zeta_\alpha}{\partial w_\beta}(q_t) \right| \cdot |\tau_\beta| \leq Ce^{-\delta'z(q_1)^{n/2}} \left(\sum_{j=1}^{n-1} |\tau_j| + \frac{1}{|w_n(q_1)|} |\tau_n| \right).$$

Combining (2-26) and (2-27) with $\alpha = n$, we get that when $z_1 \gg 1$,

$$|\tau_n| \leq Ce^{-\delta'z(q_1)^{n/2}}|w_n(q_1)| \cdot \sum_{j=1}^{n-1} |\tau_j|.$$

If we now sum (2-26) for $\alpha = 1, \dots, n - 1$ and use (2-27), then we obtain that $\tau_\alpha = 0$ for $1 \leq \alpha \leq n - 1$ and hence that $\tau_n = 0$ as well.

Next, we claim that away from a sufficiently large compact subset, G_k maps no point into $F_k(\mathbf{0}_L)$. To see this, assume that there exists a point $p \in X$ with $G_k(p) = F_k(x)$, where $x \in \mathbf{0}_L$. Letting $q = \Phi^{-1}(p)$, if $z(q)$ is sufficiently large then q and x must be contained in a same w -coordinate chart as above. Taking the n^{th} component of $\Psi(G_k(p)) = \eta(p)$, yields

$$(2-28) \quad 0 = w_n(x) = \Psi_n(F_k(x)) = \eta_n(p) = \eta_n(\Phi(q)) = w_n(q)(1 + \zeta_n(q)).$$

But if $z(q) \gg 1$, we obtain a contradiction since $w_n(q) \neq 0$. The proposition then follows from the above. □

Remark 2.4 Recall the coordinate $(\eta \circ \Phi)_n = w_n(1 + \eta_n)$, and since $e^{-\delta z^{n/2}} \sim e^{-\delta' \sqrt{-\log |w_n|}}$, the complex structure I is not even Hölder continuous at the divisor. Thus, one cannot invoke any known weak version of the Newlander–Nirenberg theorem to construct a complex-analytic compactification; see for example [20]. This is the reason why we had to deviate from the known approaches to constructing such compactifications; compare for example the proofs of [16, Theorem 3.1] and [26, Theorem 1.6].

Denote by Z the image of \bar{G}_k . It follows from the Remmert–Stein theorem [11, Chapter II.8, Theorem 8.7] that Z is a complex analytic variety in a neighborhood of $F_k(\mathbf{0}_L) \cong D$. Since Z is topologically a manifold by Proposition 2.3, it follows that Z is locally irreducible. Thus, by [11, Chapter II.7, Corollary 7.13], the normalization map $Z' \rightarrow Z$ is a homeomorphism.

Denote by D' the copy of D in Z' . Notice that, in the proof of Proposition 2.3, the functions η_α for $1 \leq \alpha \leq n$ can be viewed as local holomorphic functions on a neighborhood of a point $q \in D'$ in Z' . Moreover, they define a local topological embedding of Z' into \mathbb{C}^n . So the inverse map is holomorphic, which implies that Z' must be smooth near D' . Now we may glue Z' to X and obtain a smooth complex-analytic compactification of X , which we denote by \bar{X}' . By construction, D' is a smooth divisor in \bar{X}' .

Write $\omega' \equiv \sqrt{-1}(\sum_{j=1}^{n-1} d\eta_j \wedge d\bar{\eta}_j + |\eta_n|^{-2} d\eta_n \wedge d\bar{\eta}_n)$. This is a locally defined Kähler metric in a punctured neighborhood of D' , with cylindrical behavior normal to D' . It is easy to check using the calculations in the proof of Proposition 2.3 that for all $l \geq 0$,

$$(2-29) \quad \sum_{j=1}^{n-1} |\nabla_{\omega'}^l(w_j)|_{\omega'} + |w_n^{-1} \nabla_{\omega'}^l(w_n)|_{\omega'} \leq C_l.$$

This is a crucial fact for us. One can also reinterpret this as saying that the metric ω' is C^∞ uniformly equivalent to the corresponding cylindrical Kähler metric defined using (w_1, \dots, w_n) . Note that $|\partial_{w_j} z| \leq Cz^{1-n}$ for $1 \leq j \leq n-1$ and $|\partial_{w_n} z| \leq Cz^{1-n}|w_n|^{-1}$. In particular, we have that

$$(2-30) \quad |\nabla_{\omega'}^l z|_{\omega'} \leq Clz^{1-nl}.$$

Recall that Ω is the holomorphic volume form on (X, g) with respect to I , and $\Omega_{\mathcal{C}}$ is the corresponding holomorphic 2-form on the Calabi model space \mathcal{C} . Notice that $\Omega_{\mathcal{C}}$ is a meromorphic 2-form on L with a simple pole along the zero section $\mathbf{0}_L$, and Ω can be viewed as a holomorphic 2-form on $\bar{X}' \setminus D'$.

Lemma 2.5 Ω is a meromorphic volume form on \bar{X}' with a simple pole along D' . In particular, D' is an anticanonical divisor.

Proof We may locally, near a point $p \in D$, write $\Omega_{\mathcal{C}} = f(w_1, \dots, w_n)w_n^{-1}dw_1 \wedge \dots \wedge dw_n$, where f is a nowhere vanishing local holomorphic function. Then $\Omega = f(w_1, \dots, w_n)w_n^{-1}dw_1 \wedge \dots \wedge dw_n + \Gamma$ with $|\Gamma \wedge \bar{\Gamma}/\omega_{\mathcal{C}}^n| = O(e^{-\delta z^{n/2}})$. But $\omega_{\mathcal{C}}^n = \frac{1}{2}(\sqrt{-1})^{n^2} \Omega_{\mathcal{C}} \wedge \bar{\Omega}_{\mathcal{C}} \sim -|w_n|^{-2}dw_1 \wedge d\bar{w}_1 \wedge \dots \wedge dw_n \wedge d\bar{w}_n$ near D . It follows that $C^{-1} \leq |\Omega|_{\omega'} \leq C$, so the form $w_n\Omega$ extends holomorphically to D' locally. \square

Now choose a finite open cover $\{V_j\}$ of D such that each V_j has holomorphic coordinates $w_{1,j}, \dots, w_{n-1,j}$ given by the quotients $s_{k,j}/s_{0,j}$ for some sections $s_{0,j}, \dots, s_{n-1,j} \in H^0(D, L^k)$ for $1 \leq k \leq n-1$, and $L|_{V_j}$ has a trivializing section $e_j = s_{n,j}/s_{0,j}$ for some $s_{n,j} \in H^0(D, L^{k+1})$. Consequently we obtain an open cover $\{U_j\}$ of a neighborhood of $\mathbf{0}_L \simeq D$ in L , with holomorphic coordinates $\{w_{1,j}, \dots, w_{n-1,j}\}$, such that a point $\xi \in L$ is represented as $\xi = w_{n,j} \cdot e_j(w_{1,j}, \dots, w_{n-1,j})$. Write $e^{-\varphi_j(w_{1,j}, \dots, w_{n-1,j})} = |e_j(w_{1,j}, \dots, w_{n-1,j})|_h^2$.

Accordingly, we get an open cover $\{U'_j\}$ of a neighborhood of D' in \bar{X}' . In each U'_j we then have holomorphic coordinates $\eta_{k,j} = \mathcal{L}(s_{k,j})/\mathcal{L}(s_{0,j})$ and $\eta_{n,j} = \mathcal{L}(s_{0,j})/\mathcal{L}(s_{n,j})$. From the proof of Proposition 2.3 we know that $\eta_{k,j} = w_{k,j}$ on $U'_j \cap D'$ for $1 \leq k \leq n-1$, and the transition functions

satisfy $\partial\eta_{n,j}/\partial\eta_{n,i} = \partial w_{n,j}/\partial w_{n,i}$ on $U'_i \cap U'_j \cap D'$. This implies that the conormal bundle $N_{D'}^{-1}$ is isomorphic to L^{-1} , so in particular D' has ample normal bundle in \bar{X}' . Moreover, the local functions $\eta_{n,j}$ together give rise to a global section $S_{D'}$ of the line bundle $[D'] \simeq K_{\bar{X}'}^{-1}$ with a simple zero along D' .

Now the local functions φ_j define a hermitian metric on $N_{D'} \simeq [D']$. Fix an arbitrary extension of this to a hermitian metric h' on $[D']$ on a neighborhood of D' . On each chart U_j , this extension h' has a local representation given by $e^{-\phi_j}$, with $\phi_j = \varphi_j$ along D' . We consider a smooth closed $(1, 1)$ -form

$$(2-31) \quad \omega_m \equiv \frac{n}{n+1} \sqrt{-1} \partial\bar{\partial} (-\log |S_{D'}|_{h'}^2)^{(n+1)/n}.$$

It is well-defined and positive definite outside a compact subset of $\bar{X}' \setminus D'$.

One can check that ω_m and ω' satisfy

$$(2-32) \quad C^{-1} z^{1-n} \omega' \leq \omega_m \leq C z \omega'$$

for some constant C , and for each $k \geq 0$,

$$(2-33) \quad |\nabla_{\omega'}^k \omega_m|_{\omega'} = O(z^{mk})$$

for some $m_k \in \mathbb{N}_0$, as $z \rightarrow \infty$. Therefore covariant derivative with respect to ω' and ω_m differ by at most polynomial error terms.

Proposition 2.6 For all $l > 0$ and all $\delta < \min(\delta_k, \underline{\delta})$, we have that

$$(2-34) \quad |\nabla_{\omega_m}^l (\omega - \omega_m)|_{\omega_m} = O(e^{-\delta z^{n/2}}).$$

Consequently, $[\omega] \in \text{im}(H_c^2(X) \rightarrow H^2(X))$.

Proof By assumption, $\omega = \omega_\epsilon + \beta_1$, where $|\nabla_{\omega_\epsilon}^l \beta_1|_{\omega_\epsilon} = O(e^{-\delta z^{n/2}})$ for all $l \geq 0$. Let

$$\tilde{\omega} \equiv \frac{n}{n+1} \sqrt{-1} \partial_I \bar{\partial}_I z^{n+1}.$$

Writing $\omega_\epsilon = \tilde{\omega} + \beta_2$, using (2-30), (2-32) and (2-33), we have that $|\nabla_{\tilde{\omega}}^l \beta_2|_{\tilde{\omega}} = O(e^{-\delta z^{n/2}})$ for all $l \geq 0$ and $\delta < \underline{\delta}$. In U_j , we have that

$$z^n = -\log |\xi|_{h_L}^2 = -\log |w_{n,j}|^2 + \varphi_j(w_{1,j}, \dots, w_{n-1,j}).$$

On the other hand, we have that

$$|S_{D'}|_{h'}^2 = -\log |\eta_{n,j}|^2 + \phi_j(\eta_{1,j}, \dots, \eta_{n,j}).$$

Notice that $\eta_{n,j} = w_{n,j}(1 + \beta_3)$, with $|\nabla_{\omega}^l \beta_3|_{\omega} = O(e^{-\delta_k z^{n/2}})$ for all $l \geq 0$, and $\phi_j(\eta_{1,j}, \dots, \eta_{n,j}) = \varphi_j(w_{1,j}, \dots, w_{n-1,j}) + \eta_{n,j} P(\eta_{1,j}, \dots, \eta_{n,j})$ for some smooth function P . Estimate (2-34) now follows from a straightforward computation, using the key property (2-29), (2-32) and (2-33).

By integrating from infinity as in [19, Lemma 3.7], away from a compact set we may write $\omega = \omega_m + d\sigma$, where σ is a smooth real-valued 1-form such that $|\nabla_{\omega_m}^l \sigma|_{\omega_m} = O(e^{-\delta z^{n/2}})$ for all $l \geq 0$. Then the smooth real-valued 2-form

$$(2-35) \quad \omega - d \left(\chi \left(\sigma + \frac{n}{2(n+1)} d_I^c (-\log |S_{D'}|_{\tilde{h}'}^2)^{(n+1)/n} \right) \right)$$

is cohomologous to ω and has compact support, where χ is a cutoff function which is 1 in a neighborhood of infinity and 0 on a compact set. \square

Now we claim that \bar{X}' is a *weak Fano manifold*, ie \bar{X}' is projective and $K_{\bar{X}'}^{-1}$ is big and nef. Assuming projectivity, the nef property is obvious because $K_{\bar{X}'}^{-1} = [D']$ for some smooth divisor D' with ample normal bundle, and it is also elementary to deduce the big property from this fact; see for example [36, Lemma 2.3]. Projectivity follows from a similar reasoning as in the proof of [10, Theorem 2.1]; see in particular [10, page 4, proof in the smooth case]. In short, one first proves, using the theory of the Remmert reduction (see the beginning of Section 2), that $K_{\bar{X}'}^{-1}$ is semiample. This also strongly relies on the fact that $K_{\bar{X}'}^{-1} = [D']$, where D' has ample normal bundle. It then follows that \bar{X}' is Moishezon, which implies that Hodge theory holds on \bar{X}' . Also, $h^{0,2}(\bar{X}') = 0$ thanks to the Grauert–Riemenschneider vanishing theorem [14, Satz 2.1]. Using the latter two properties, the fact that \bar{X}' admits compactly supported Kähler classes by Proposition 2.6, and another vanishing theorem on the open manifold X due to Grauert–Riemenschneider [14, page 278, Korollar], one can explicitly write down a Kähler form on \bar{X}' . The Kodaira embedding theorem now implies that \bar{X}' is projective because we already know that $h^{0,2}(\bar{X}') = 0$. We note that a similar gluing argument of Kähler forms will appear again in the proof of Lemma 2.7 below. The argument works here even though, unlike in [10], D' is not Fano, so that $H^{1,1}(\bar{X}')$ may not surject onto $H^2(X)$. The key point is that we are restricting ourselves to the case of compactly supported Kähler classes on X .

To finish the proof of Theorem 1.3, it remains to identify the Ricci-flat Kähler metric ω on X with a slight generalization of the construction given by Tian and Yau [35, Theorem 4.2]. Let M be an n -dimensional projective manifold containing a smooth divisor D with ample normal bundle such that $K_M^{-1} = [D]$. Then M is necessarily weakly Fano. Fix a defining section $S \in H^0(M, K_M^{-1})$ of the divisor D and denote $X \equiv M \setminus D$. Following the beginning of the argument from [10] sketched above, one first proves that K_M^{-1} is semiample. Here this also follows from the Kawamata basepoint-free theorem [22, Theorem 6.1] because we already know that M is projective. Let E be the nonample locus of K_M^{-1} . This is a union of some subvarieties of M that are disjoint from D . We fix a smooth hermitian metric \tilde{h} on K_M^{-1} such that its curvature form is nonnegative everywhere and is positive away from E . Write $\tilde{v} \equiv -\log |S|_{\tilde{h}}^2$.

We may view S^{-1} as a holomorphic volume form Ω_X on X with a simple pole along D . Let Ω_D be the holomorphic volume form on D given by the residue of Ω_X along D . Let h_D be a hermitian metric on $K_M^{-1}|_D$ such that its curvature form is a Calabi–Yau metric ω_D on D . Rescaling S if necessary, we may assume that $\omega_D^{n-1} = \frac{1}{2}(\sqrt{-1})^{(n-1)^2} \Omega_D \wedge \bar{\Omega}_D$.

One can extend h_D to a smooth hermitian metric h on K_M^{-1} such that its curvature form is positive definite in a neighborhood of D . Fix such an extension, denoted by h_M . For any $A \in \mathbb{R}$, we write $h_A \equiv h_M e^{-A}$. Then it is easy to see that outside a compact subset of X , $(-\log |S|_{h_A}^2)^{(n+1)/n}$ is strictly plurisubharmonic. As in the beginning of this section, by composing with a convex function we may find a global smooth function v_A on X with $dd^c v_A \geq 0$ such that

$$v_A = \frac{n}{n+1} (-\log |S|_{h_A}^2)^{(n+1)/n}$$

outside a compact set and v_A is constant in a neighborhood of E .

Abusing notation, we denote by $H_{c,+}^2(X)$ the subset of $\text{im}(H_c^2(X) \rightarrow H^2(X))$ consisting of classes \mathfrak{k} such that $\int_Y \mathfrak{k}^p > 0$ for any compact analytic subset Y of X of pure complex dimension $p > 0$. Notice that any such Y must be contained in E .

Lemma 2.7 *For all $\mathfrak{k} \in H_{c,+}^2(X)$ and all numbers $A \in \mathbb{R}$, $\tau > 0$, there exists a smooth Kähler form $\omega_{A,\tau}$ on X such that $[\omega_{A,\tau}] = \mathfrak{k}$ and $|\nabla_{\omega_{A,\tau}}^l (\omega_{A,\tau} - \tau dd^c v_A)|_{\omega_{A,\tau}} = O(|S|_{h_M}^{\delta_0})$ for some $\delta_0 > 0$ and all $l \geq 0$.*

Proof By hypothesis, \mathfrak{k} is represented by a closed 2-form β on X such that $\beta = 0$ away from some compact subset of X . In particular, β trivially extends to a smooth closed 2-form on M . By the Grauert–Riemenschneider vanishing theorem [14, Satz 2.1], or (because we are assuming that M is projective) by the Kawamata–Viehweg vanishing theorem (see for example [24, Theorem 2.64]), we have that $h^{0,2}(M) = 0$. Thus, there exist a smooth closed $(1, 1)$ -form η and a smooth 1-form γ on M such that $\beta = \eta + d\gamma$. Because β has compact support in $X \subset M$, it follows that η is d -exact on some open neighborhood of D in M . In particular, $\eta|_D = dd^c f$ for some smooth function f on D . We choose an arbitrary smooth extension of f to M and replace β by $\eta - dd^c f$. Thus, we are now assuming without loss of generality that β is a smooth closed $(1, 1)$ -form on M such that $\beta|_D = 0$ and $[\beta|_M] = \mathfrak{k}$. It is straightforward to check that such a form β satisfies the exponential decay estimate from the statement of the lemma with respect to any reference metric of the form $\tau dd^c v_A$ outside a sufficiently large compact subset of X . (If $n \geq 3$, then instead of applying the dd^c -lemma on D it is possible to apply a stronger dd^c -lemma on the complement of a large compact subset of X to arrange directly that β still has compact support in X ; see for example [7, Lemma 4.3].)

Because $\mathfrak{k} \in H_{c,+}^2(X)$, by the generalized Demailly–Păun criterion of [9, Theorem 1.1] there exists a smooth function u_0 on X such that $\beta + dd^c u_0$ is positive in a neighborhood U of E . Choose a cutoff function χ of compact support contained in U which equals 1 on a neighborhood of E , and let $\rho = \rho(t)$ be a cutoff function on $[0, \infty)$ which equals 1 when $t \leq 1$ and vanishes when $t \geq 2$. For fixed A and τ , we define

$$(2-36) \quad \omega_{A,\tau} \equiv \beta + dd^c \left(\chi u_0 + C_1 \left(\rho \circ \left(\frac{1}{C_2} \tilde{v} \right) \right) \cdot \tilde{v} + \tau v_A \right).$$

It is straightforward to verify that if we first choose C_1 large and then C_2 large (depending on C_1), then $\omega_{A,\tau}$ is globally positive on X . \square

One can also see that $\omega_{A,\tau}^n = e^{-f_{A,\tau}} \tau^n \cdot \frac{1}{2}(\sqrt{-1})^{n^2} \Omega_X \wedge \bar{\Omega}_X$ for some function $f_{A,\tau}$ which tends to zero at infinity.

Lemma 2.8 *Given any $\tau > 0$, there is a unique $A = A(\tau)$ such that*

$$(2-37) \quad \int_X (\omega_{A,\tau}^n - \tau^n \frac{1}{2}(\sqrt{-1})^{n^2} \Omega_X \wedge \bar{\Omega}_X) = 0.$$

Proof It is easy to check that the integral is finite for any A . For a given A , add and subtract $\omega_{0,\tau}^n$ under the integral sign and split up the integral accordingly. For $\varepsilon > 0$ sufficiently small, we have

$$\int_{|S|_h \leq \varepsilon} (\omega_{A,\tau}^n - \omega_{0,\tau}^n) = \tau \int_{|S|_h = \varepsilon} (d^c v_A \wedge \omega_{A,\tau}^{n-1} - d^c v_0 \wedge \omega_{0,\tau}^{n-1}).$$

By computing the boundary term explicitly and letting $\varepsilon \rightarrow 0$, one sees that

$$\int_X (\omega_{A,\tau}^n - \omega_{0,\tau}^n) = \lambda \cdot A \tau^n \int_D \omega_D^{n-1}$$

for some $\lambda = \lambda(n) \neq 0$. The key point is that we only care the limit as $\varepsilon \rightarrow 0$, and thus we only need to understand the leading term and ignore all the lower order terms. This argument goes back to [35, Lemma 4.6]. For a more detailed exposition, see also the very recent paper [6, page 14]. Then the desired condition becomes a linear equation for A . □

Using Lemmas 2.7 and 2.8 as ingredients, we now get the following existence theorem, which generalizes the classical existence result of Tian and Yau [35, Theorem 4.2].

Theorem 2.9 *Given any $\mathfrak{k} \in H_{c,+}^2(M)$ and $\tau > 0$, there is a complete Kähler metric $\omega_\tau = \omega_{A(\tau),\tau} + dd^c \phi$ on X such that $[\omega_\tau] = \mathfrak{k}$ and $\omega_\tau^n = \tau^n \frac{1}{2}(\sqrt{-1})^{n^2} \Omega_X \wedge \bar{\Omega}_X$, and such that for some $\delta_0 > 0$ and for all $l \geq 0$ we have that $|\nabla_{\omega_{A(\tau),\tau}}^l \phi|_{\omega_{A(\tau),\tau}} = O(e^{-\delta_0(-\log |S|_{h_M}^2)^{1/2}})$.*

This follows from [35, Theorem 1.1] and [18, Proposition 2.4, Proposition 2.9(ia) and (ii)]. The correct choice of $A = A(\tau)$ is crucial because otherwise the relevant Monge–Ampère equation cannot be solved by direct methods. Once the equation has been solved for $A = A(\tau)$, it follows that a solution exists for all A by adding $v_{A(\tau),\tau} - v_{A,\tau}$ to the potential, but $v_{A(\tau),\tau} - v_{A,\tau}$ is comparable to $(A(\tau) - A)\tau(-\log |S|_{h_M}^2)^{1/n}$ at infinity, which is not even uniformly bounded for $A \neq A(\tau)$.

Remark 2.10 The decay rate of the complex structure is of order $O(e^{-(\frac{1}{2}-\varepsilon)z^n})$ for all $\varepsilon > 0$. This is much faster than the decay rate of the metric, which is in general only of order $O(e^{-\delta z^{n/2}})$.

Theorem 2.11 *Suppose we have a Kähler metric ω on X such that $\omega^n = C \Omega_X \wedge \bar{\Omega}_X$ for some $C > 0$ and such that there exist $\tau > 0$ and $A \in \mathbb{R}$ such that $|\nabla_\omega^l (\omega - \tau dd^c v_A)|_\omega = O(e^{-\delta(-\log |S|_h^2)^{1/2}})$ for some $\delta > 0$ and for all $l \geq 0$. Then $\omega = \omega_\tau$, where ω_τ is the metric constructed in Theorem 2.9 in the class $\mathfrak{k} = [\omega] \in H_{c,+}^2(X)$.*

Proof By rescaling ω if necessary, we may assume that $C = 1$. As in the proof of Proposition 2.6, we can see that $\omega = \tau dd^c v_A + d\sigma$ outside a compact set, where σ is a smooth real-valued 1-form

such that $|\nabla_{\omega_m}^l \sigma|_{\omega_m} = O(e^{-\delta(-\log |S|_h^2)^{1/2}})$ for all $l \geq 0$. In particular, $[\omega] \in \text{im}(H_c^2(X) \rightarrow H^2(X))$. Thus, if we let \mathfrak{k} denote the de Rham cohomology class of ω in $H^2(X)$, then we have that $\mathfrak{k} \in H_{c,+}^2(X)$. Consequently, we can apply [Theorem 2.9](#) to obtain a complete Calabi–Yau metric ω_τ on X with $[\omega_\tau] = \mathfrak{k}$. By construction, we have that $\omega^n = \omega_\tau^n$. Then, as in the proof of [Lemma 2.8](#), we can see that $A = A(\tau)$, so we know that $\omega = \omega_\tau + d\tilde{\sigma}$, with $|\nabla_{\omega_\tau}^l \tilde{\sigma}|_{\omega_\tau} = O(e^{-\delta_1(-\log |S|_h^2)^{1/2}})$ for some $\delta_1 > 0$.

Next, we solve the Poisson equation $\bar{\partial}^* \bar{\partial} u = \bar{\partial}^* \tilde{\sigma}^{0,1}$ with respect to ω_τ . Note that $\int_X (\bar{\partial}^* \tilde{\sigma}^{0,1}) \omega_\tau^n = 0$ by definition, so we can apply [\[17, Theorem 1.5\]](#) to conclude the existence of a C^∞ solution u which is uniformly bounded and satisfies $\int_X |du|_{\omega_\tau}^2 \omega_\tau^n < \infty$. Moreover, by the integration argument in the proof of [\[18, Proposition 2.9\(ia\)\]](#), after subtracting a constant from u one actually has the asymptotics $|\nabla_{\omega_\tau}^l u|_{\omega_\tau} = O(e^{-\delta_2(-\log |S|_h^2)^{1/2}})$ for some $\delta_2 > 0$ and all $l \geq 0$. Let $\gamma \equiv \tilde{\sigma}^{0,1} - \bar{\partial} u$. By construction, $\bar{\partial} \gamma = \bar{\partial}^* \gamma = 0$ with respect to ω_τ . Thus, by the Bochner formula, $\Delta |\gamma|^2 \geq 0$. Since γ tends to zero at infinity, it must vanish identically. It follows that $\omega - \omega_\tau = dd^c(\sqrt{-1}(\bar{u} - u))$. Writing

$$(2-38) \quad 0 = \omega^n - \omega_\tau^n = (\omega - \omega_\tau)(\omega^{n-1} + \dots + \omega_\tau^{n-1}),$$

then multiplying [\(2-38\)](#) by $\bar{u} - u$ and integrating by parts, since $\bar{u} - u$ is decaying, we conclude that $u = \bar{u}$, hence $\omega = \omega_\tau$. □

The second statement in [Theorem 1.3](#) follows from [Theorem 2.11](#) (applied to $M = (\bar{X}', \bar{I})$) and [Proposition 2.6](#).

Proof of Corollary 1.4 By [\[34, Theorem 1.1\]](#), the weak Fano manifold $\bar{X} = X \cup D$ is simply connected. Since $-K_{\bar{X}}$ is big, D does not consist of a pencil, so by [\[28, Corollary 2.10\]](#), $\pi_1(X) \simeq \pi_1(-K_{\bar{X}}^\times)$, where $-K_{\bar{X}}^\times = -K_{\bar{X}} \setminus \mathbf{0}_{-K_{\bar{X}}}$. Letting N denote the unit circle bundle of $-K_{\bar{X}}$, the homotopy sequence of the fibration $S^1 \rightarrow N \rightarrow \bar{X}$ yields

$$(2-39) \quad \pi_2(\bar{X}) \rightarrow \mathbb{Z} \rightarrow \pi_1(X) \rightarrow 0.$$

Since \bar{X} is simply connected and $-K_{\bar{X}}^\times$ deformation retracts onto N , noticing that $K_{\bar{X}}$ is trivial over X , we have $\pi_1(N) \cong \pi_1(-K_{\bar{X}}^\times) \cong \pi_1(X)$. By [\[30, Proposition 2\]](#), the first mapping factors as the Hurewicz projection from $\pi_2(\bar{X}) \rightarrow H_2(\bar{X}, \mathbb{Z})$ composed with the mapping $\delta: H_2(\bar{X}, \mathbb{Z}) \rightarrow \mathbb{Z}$ given by the intersection pairing with $[D] \in H_{n-2}(\bar{X}, \mathbb{Z})$, that is,

$$(2-40) \quad \delta(\Sigma) = \int_{\Sigma} c_1([D]).$$

By Poincaré duality, this mapping is nontrivial, so the image is an infinite subgroup $k\mathbb{Z} \subset \mathbb{Z}$, and consequently $\pi_1(X) = \mathbb{Z}/k\mathbb{Z}$ is cyclic, with k equal to the index of \bar{X} .

The bound on the degree is a consequence of the following results. It is a classical result that the degree of a weak del Pezzo surface satisfies $1 \leq d \leq 9$; see for example [\[12, Chapter 8\]](#). A bound on the degree of a weak Fano threefold is proved in [\[23, Corollary 1.3\]](#). Finally, Birkar [\[1, Theorem 1.1\]](#) proved that in any dimension there is an a priori bound on the degree of a weak Fano manifold. □

3 ALH* gravitational instantons

The 3-dimensional Heisenberg group is

$$(3-1) \quad H(1, \mathbb{R}) \equiv \left\{ \begin{bmatrix} 1 & x & t \\ 0 & 1 & y \\ 0 & 0 & 1 \end{bmatrix} : x, y, t \in \mathbb{R} \right\}.$$

Let $(\mathbb{T}^2, g_{c,\tau})$ be the flat 2-torus corresponding to the lattice $c(\mathbb{Z} \oplus \mathbb{Z}\tau) \subset \mathbb{C}$ with $\tau \in \mathbb{H}$ and $c > 0$, where \mathbb{H} is the upper half-plane in \mathbb{C} . Write $\tau_1 = \text{Re}(\tau)$ and $\tau_2 = \text{Im}(\tau)$. For $b \in \mathbb{Z}_+$, let $\text{Nil}_{b,c,\tau}^3$ be the nilmanifold corresponding to the quotient of the Heisenberg group by the subgroup generated by

$$(3-2) \quad (x, y, t) \mapsto (x + c, y, t + cy),$$

$$(3-3) \quad (x, y, t) \mapsto (x + c\tau_1, y + c\tau_2, t + c\tau_1 y),$$

$$(3-4) \quad (x, y, t) \mapsto (x, y, t + \tau_2 b^{-1} c^2).$$

Definition 3.1 For $b \in \mathbb{Z}_+, c > 0, \tau \in \mathbb{H}$, and $R > 0$, the ALH* model space is

$$(3-5) \quad \mathfrak{M}_{b,c,\tau}(R) \equiv (R, \infty) \times \text{Nil}_{b,\tau,c}^3$$

together with the hyperkähler Riemannian metric

$$(3-6) \quad g^{\mathfrak{M}} \equiv V(dx^2 + dy^2 + dz^2) + V^{-1}\theta^2,$$

where z is the coordinate on (R, ∞) , $V \equiv 2\pi bc^{-2}\tau_2^{-1}z$ and $\theta \equiv 2\pi bc^{-2}\tau_2^{-1}(dt - x dy)$.

Choose the following orthonormal frame for $T^*\mathfrak{M}$:

$$(3-7) \quad \{e_1, e_2, e_3, e_4\} = \{V^{\frac{1}{2}} dx, V^{\frac{1}{2}} dy, V^{\frac{1}{2}} dz, V^{-\frac{1}{2}} \theta\}.$$

Define three almost-complex structures on $T^*\mathfrak{M}$ by

$$(3-8) \quad I_{\mathfrak{M}}^*(e_1) = e_2, \quad I_{\mathfrak{M}}^*(e_3) = e_4,$$

$$(3-9) \quad J_{\mathfrak{M}}^*(e_1) = e_4, \quad J_{\mathfrak{M}}^*(e_2) = e_3,$$

$$(3-10) \quad K_{\mathfrak{M}}^*(e_2) = e_4, \quad K_{\mathfrak{M}}^*(e_3) = e_1,$$

which are dual to almost-complex structures $I_{\mathfrak{M}}, J_{\mathfrak{M}}, K_{\mathfrak{M}}$ on $T\mathfrak{M}$, respectively. Denote the Kähler forms associated to $I_{\mathfrak{M}}^*, J_{\mathfrak{M}}^*, K_{\mathfrak{M}}^*$ by $\omega_1^{\mathfrak{M}}, \omega_2^{\mathfrak{M}}, \omega_3^{\mathfrak{M}}$, respectively. These are explicitly given by

$$(3-11) \quad \omega_1^{\mathfrak{M}} = dz \wedge \theta + V dx \wedge dy,$$

$$(3-12) \quad \omega_2^{\mathfrak{M}} = dx \wedge \theta + V dy \wedge dz,$$

$$(3-13) \quad \omega_3^{\mathfrak{M}} = dy \wedge \theta + V dz \wedge dx.$$

Taken together, this data defines a hyperkähler structure on \mathfrak{M} , which we denote as $(\mathfrak{M}, g^{\mathfrak{M}}, \omega^{\mathfrak{M}})$. This structure can equivalently also be specified as $(\mathfrak{M}, g^{\mathfrak{M}}, I_{\mathfrak{M}}, J_{\mathfrak{M}}, K_{\mathfrak{M}})$.

Definition 3.2 (ALH* gravitational instanton) A hyperkähler structure (X, g, ω) on a 4-manifold X is called an ALH* gravitational instanton with parameters $b \in \mathbb{Z}_+$, $c > 0$ and $\tau \in \mathbb{H}$ if there exist $\delta, R > 0$, a compact subset $X_R \subset X$, an ALH* model space $(\mathfrak{M}_{b,c,\tau}(R), g^{\mathfrak{M}})$ and a diffeomorphism

$$(3-14) \quad \Phi: \mathfrak{M}_{b,c,\tau}(R) \rightarrow X \setminus X_R$$

such that for all $k \in \mathbb{N}_0$,

$$(3-15) \quad |\nabla_{g^{\mathfrak{M}}}^k (\Phi^* g - g^{\mathfrak{M}})|_{g^{\mathfrak{M}}} = O(e^{-\delta z}),$$

$$(3-16) \quad |\nabla_{g^{\mathfrak{M}}}^k (\Phi^* \omega_i - \omega_i^{\mathfrak{M}})|_{g^{\mathfrak{M}}} = O(e^{-\delta z}) \quad \text{for } i = 1, 2, 3,$$

as $z \rightarrow \infty$.

According to [19, Proposition 3.1], any ALH* model space \mathfrak{M} together with its complex structure $I_{\mathfrak{M}}$ is holomorphically isometric to a Calabi model space up to rescaling. This means the following. We can view the flat torus $(\mathbb{T}^2, g_{c,\tau})$ as an elliptic curve D . Then there exists an ample line bundle $L \rightarrow D$ of degree b , together with a hermitian metric h_L whose curvature form defines the flat metric

$$(3-17) \quad 2\pi b c^{-2} \tau_2^{-1} g_{c,\tau}$$

on D , such that the underlying complex manifold $(\mathfrak{M}_{b,c,\tau}(R), I_{\mathfrak{M}})$ can be identified with the open set

$$(3-18) \quad \mathcal{C} \equiv \{\xi \in L : 0 < |\xi|_{h_L} < e^{-\frac{1}{2}z_0^2}\}$$

for some $z_0 > 0$. Moreover, the Kähler form and the holomorphic 2-form on \mathfrak{M} are respectively given by $\omega_1 = \mu \omega_{\mathcal{C}}$ for some $\mu > 0$ and $\omega_2 + \sqrt{-1}\omega_3 = \nu \Omega_{\mathcal{C}}$ for some $\nu \in \mathbb{C}^*$, where

$$(3-19) \quad \omega_{\mathcal{C}} \equiv \frac{2}{3} \sqrt{-1} \partial_{\mathcal{C}} \bar{\partial}_{\mathcal{C}} (-\log |\xi|_{h_L}^2)^{\frac{3}{2}}$$

and $\Omega_{\mathcal{C}}$ is a holomorphic volume form which has a simple pole along the zero section $\mathbf{0}_L$ and is invariant under the natural \mathbb{C}^* -action on L . In addition, we have that

$$(3-20) \quad z = (-\log |\xi|_{h_L}^2)^{\frac{1}{2}},$$

and this is the $\omega_{\mathcal{C}}$ -moment map for the natural S^1 -action on L . We note here that h_L is unique only up to scaling, and the scale of h_L is an important free parameter; see Lemma 2.8.

Thanks to this identification, an ALH* gravitational instanton is the same as a complete hyperkähler 4-manifold which is asymptotically Calabi in the sense of [19, Definition 4.1]. The proof of part (i) of Theorem 1.5 follows from this equivalence and Theorem 1.3.

3.1 Compactification to rational elliptic surfaces

In this subsection, we prove part (ii) of Theorem 1.5. First, we recall some basic facts about the lowest nontrivial eigenvalue of the Laplacian on $(\mathbb{T}^2, g_{c,\tau})$. The \mathbb{Z}^2 -action on \mathbb{C} is generated by

$$(3-21) \quad (x, y) \mapsto (x + c, y), \quad (x, y) \mapsto (x + c\tau_1, y + c\tau_2).$$

The eigenfunctions of the Laplacian on $(\mathbb{T}^2, g_{c,\tau})$ are given by

$$(3-22) \quad \phi_{m,n}(x, y) = e^{2\pi imc^{-1}(x-\tau_1\tau_2^{-1}y)} e^{2\pi inc^{-1}\tau_2^{-1}y}$$

for $(m, n) \in \mathbb{Z}^2$. The eigenvalues are

$$(3-23) \quad \lambda_{m,n} = 4\pi^2 c^{-2} \{m^2 + \tau_2^{-2}(n - m\tau_1)^2\}.$$

Using the $\text{PSL}(2, \mathbb{Z})$ -action on \mathbb{H} , we can assume without loss of generality that $|\tau| \geq 1$. Then the lowest nontrivial eigenvalue is given by $\lambda_1 = \lambda_{0,1} = 4\pi^2 c^{-2} \tau_2^{-2}$, with eigenfunction $\phi_{0,1}$.

Next, we establish an almost optimal decay rate for any ALH* gravitational instanton.

Proposition 3.3 *Let (X, g, ω) be an ALH* gravitational instanton with parameters $b \in \mathbb{Z}_+, c > 0, \tau \in \mathbb{H}$. Then, in suitable ALH* coordinates, (3-15)–(3-16) hold with $\delta = \sqrt{\lambda_1} - \varepsilon$ for any $\varepsilon > 0$.*

Proof By Theorem 1.3, (X, g, I) arises as a Tian–Yau metric on a weak del Pezzo surface M minus an anticanonical divisor D . Thus, we can use the coordinate system Φ from [19, Proposition 3.4]. The Tian–Yau metric is of the form $\omega = \omega_0 + \sqrt{-1} \partial\bar{\partial}u$ with $|\nabla_{g_\varepsilon}^k u|_{g_\varepsilon} = O(e^{-\varepsilon z})$ as $z \rightarrow \infty$ for some $\varepsilon > 0$ and all $k \geq 0$. By [19, Proposition 3.4], the background Kähler form satisfies the asymptotics

$$(3-24) \quad |\nabla_{g_\varepsilon}^k (\omega_0 - \omega_\varepsilon)|_{g_\varepsilon} = O(e^{-\varepsilon z^2})$$

as $z \rightarrow \infty$, for some $\varepsilon > 0$ and all $k \geq 0$. Expanding the Calabi–Yau equation

$$(3-25) \quad (\omega_0 + \sqrt{-1} \partial\bar{\partial}u)^2 = \frac{1}{2} \Omega \wedge \bar{\Omega}$$

yields that $\Delta_{g_\varepsilon} u = O(e^{-2\varepsilon z})$ as $z \rightarrow \infty$. Assume that $2\varepsilon < \sqrt{\lambda_1}$. Then the arguments of [19, Section 4] allow us to conclude that $u = O(e^{-2\varepsilon z})$ for any $\underline{\varepsilon} \in (0, \varepsilon)$. Indeed, notice that [19, Proposition 4.12] holds with $\underline{\delta} = \sqrt{\lambda_1}$, as is clear from the proof. Using this, we can find a function $u_{-2\underline{\varepsilon}}$ defined on $\{z > R_1\}$ for some $R_1 > R$ such that $u_{-2\underline{\varepsilon}} = O(e^{-2\underline{\varepsilon} z})$ for any $\underline{\varepsilon} \in (0, \varepsilon)$ and $\Delta_{g_\varepsilon} u_{-2\underline{\varepsilon}} = \Delta_{g_\varepsilon} u$. The function $u - u_{-2\underline{\varepsilon}}$ is then harmonic with respect to g_ε and is also $o(1)$ as $z \rightarrow \infty$, so from [19, Proposition 4.10] (part (2) of which is easily seen to hold with $\underline{\delta}/2$ replaced by any number greater than $-\sqrt{\lambda_1}$), we have

$$(3-26) \quad u - u_{-2\underline{\varepsilon}} = O(e^{(-\sqrt{\lambda_1} + \varepsilon')z})$$

for any $\varepsilon' > 0$, which implies that $u = O(e^{-2\varepsilon z})$ for any $\underline{\varepsilon} \in (0, \varepsilon)$. We can iterate this argument together with standard local derivative estimates for the equation (3-25) to get

$$(3-27) \quad |\nabla_{g_\varepsilon}^k u|_{g_\varepsilon} = O(e^{(-\sqrt{\lambda_1} + \varepsilon)z})$$

as $z \rightarrow \infty$, for all $k \geq 0$ and $\varepsilon > 0$. This implies (3-16) for $i = 1$ with $\delta = \sqrt{\lambda_1} - \varepsilon$ for any $\varepsilon > 0$.

Also, from Theorem 1.3 and [19, Proposition 3.4(a)], we have that

$$(3-28) \quad |\nabla_{g_{\mathfrak{M}}}^k (\Phi^* I - I_{\mathfrak{M}})|_{g_{\mathfrak{M}}} = O(e^{-\varepsilon' z^2})$$

for some $\varepsilon' > 0$ and any $k \geq 0$ as $z \rightarrow \infty$. Since g is Kähler with respect to I , the metric condition (3-15) with $\delta = \sqrt{\lambda_1} - \varepsilon$ then follows from (3-28) and the above estimates on $\Phi^* \omega_1 - \omega_1^{\mathfrak{M}}$.

Finally, from [19, Proposition 3.4(b)], we have that

$$(3-29) \quad |\nabla_{g_{\mathfrak{M}}}^k (\Phi^* \Omega_X - \Omega_{\mathfrak{M}})|_{g_{\mathfrak{M}}} = O(e^{-\varepsilon' z^2})$$

for some $\varepsilon' > 0$ and any $k \geq 0$ as $z \rightarrow \infty$. Because $\Omega_X = \omega_2 + \sqrt{-1}\omega_3$ and $\Omega_{\mathfrak{M}} = \omega_2^{\mathfrak{M}} + \sqrt{-1}\omega_3^{\mathfrak{M}}$, the conditions (3-16) for $i = 2$ and $i = 3$ actually hold for any $\delta > 0$. \square

Remark 3.4 Proposition 3.3 could also be proved by following the arguments in [33, Section 6.5]. This way of reasoning is analytically more involved, but it does not require Theorem 1.3.

We now produce a J -holomorphic function on X asymptotic to a nice $J_{\mathfrak{M}}$ -holomorphic function on \mathfrak{M} that defines an elliptic fibration of \mathfrak{M} with the desired behavior at infinity.

Proposition 3.5 *Let (X, g, I, J, K) be an ALH* gravitational instanton. Then there exists a function $u: X \rightarrow \mathbb{C}$ which is holomorphic with respect to J and satisfies*

$$(3-30) \quad u = e^{\sqrt{\lambda_1}(z+iy)} + f,$$

where $|\nabla^k f|_g = O(e^{\varepsilon z})$ for all $k \geq 0$, as $z \rightarrow \infty$, for any $\varepsilon > 0$.

Proof The function $z + \sqrt{-1}y$ is a locally defined $J_{\mathfrak{M}}$ -holomorphic function on \mathfrak{M} because

$$(3-31) \quad J_{\mathfrak{M}}(dz + i dy) = -dy + i dz = i(dz + i dy).$$

Consequently, $h = e^{\sqrt{\lambda_1}(z+iy)}$ is a globally defined $J_{\mathfrak{M}}$ -holomorphic function on \mathfrak{M} . Identify X and \mathfrak{M} at infinity using the ALH* coordinate system of Proposition 3.3. Fix a cutoff function χ such that $\chi = 1$ for $z > 4R$ and $\chi = 0$ for $z < 2R$. Using the fact that $\Delta_{g_{\mathfrak{M}}} h = 0$ and Proposition 3.3, we have that

$$(3-32) \quad \Delta_g(\chi h) = O(e^{\varepsilon z}) \quad \text{as } z \rightarrow \infty, \text{ for all } \varepsilon > 0.$$

Claim 1 *There exists a function $f \in C^\infty(X)$ such that $\Delta_g f = -\Delta_g(\chi h)$ and*

$$(3-33) \quad f = O(e^{\varepsilon z}) \quad \text{as } z \rightarrow \infty, \text{ for all } \varepsilon > 0.$$

Proof of claim Using [19, Proposition 4.12], we can find a smooth function $f_{\mathfrak{M}}$ defined on \mathfrak{M} such that $f_{\mathfrak{M}} = O(e^{\varepsilon z})$ as $z \rightarrow \infty$ for any $\varepsilon > 0$ and such that $\Delta_{g_{\mathfrak{M}}} f_{\mathfrak{M}} = -\Delta_g h$. Thus, aiming to set $f = \chi f_{\mathfrak{M}} + f'$, we can reduce to solving the equation $\Delta_g f' = h'$ for $f' = O(e^{\varepsilon z})$ (for any $\varepsilon > 0$), where

$$(3-34) \quad h' \equiv -\Delta_g(\chi h) - \Delta_g(\chi f_{\mathfrak{M}}) = O(e^{(-\sqrt{\lambda_1} + \varepsilon)z})$$

for any $\varepsilon > 0$ thanks to Proposition 3.3. In fact, we will now show that for any $\delta > 0$ there exists a $\underline{\delta} > 0$ such that the equation $\Delta_g f' = h'$ with $h' = O(e^{-\delta z})$ is solvable with $f' = \alpha z + O(e^{-\delta z})$ for some $\alpha \in \mathbb{R}$. This implies what we want. To prove this new claim, we first observe that the function z is $g_{\mathfrak{M}}$ -harmonic. Thus, by Proposition 3.3, $\Delta_g(\chi z) = O(e^{(-\sqrt{\lambda_1} + \varepsilon)z})$ for any $\varepsilon > 0$, so in particular $\Delta_g(\chi z) \in L^1(X, d\text{Vol}_g)$. The divergence theorem now allows us to conclude that $\int_X \Delta_g(\chi z) d\text{Vol}_g > 0$.

Indeed, the corresponding boundary integrals in (X, g) and in $(\mathfrak{M}, g_{\mathfrak{M}})$ are asymptotic to each other because z grows more slowly than any exponential in z , and the boundary integral in $(\mathfrak{M}, g_{\mathfrak{M}})$ approaches a positive constant as $z \rightarrow \infty$ by direct computation. Thus, by replacing h' by $h' - \alpha \Delta_g(\chi z)$ for some suitable $\alpha \in \mathbb{R}$, we can assume without loss of generality that h' has mean value zero over X with respect to $d\text{Vol}_g$. It now follows from [17, Theorem 1.5] that the equation $\Delta_g f' = h'$ is solvable for some smooth and uniformly bounded f' with $\int_X |df'|_g^2 d\text{Vol}_g < \infty$. It then also follows from the integration argument in the proof of [18, Proposition 2.9(ia)] that any such f' satisfies the estimate $|\nabla^k(f' - C)|_g = O(e^{-\delta z})$ for some constant C and for any $k \geq 0$ as $z \rightarrow \infty$. After subtracting this constant, the (new) claim follows. \square

Since the function $u \equiv \chi h + f$ is harmonic, the 1-form $\alpha \equiv \bar{\partial}_J u$ satisfies

$$(3-35) \quad \bar{\partial}_J \alpha = 0, \quad \bar{\partial}_{J,g}^* \alpha = 0.$$

Since g is Kähler with respect to J , from Proposition 3.3, we have that

$$(3-36) \quad |J - J_{\mathfrak{M}}|_{g_{\mathfrak{M}}} = O(e^{(-\sqrt{\lambda_1} + \varepsilon)z}) \quad \text{as } z \rightarrow \infty.$$

Using the fact that $h = O(e^{\sqrt{\lambda_1}z})$ and $\bar{\partial}_{J_{\mathfrak{M}}} h = 0$, we can deduce from (3-36) that $\bar{\partial}_J h = O(e^{\varepsilon z})$ for any $\varepsilon > 0$ with respect to $g_{\mathfrak{M}}$. By (3-33) and standard local gradient estimates, the same bound holds for $\bar{\partial}_J f$. Thus, it holds for α as well. Since α has J -type $(0, 1)$, $\text{Re}(\alpha)$ is half-harmonic with respect to g thanks to (3-35). By [19, Theorem 5.1], we conclude that $\alpha \equiv 0$, which implies that u is J -holomorphic. \square

We now complete the proof of part (ii) of Theorem 1.5. Given our work so far, the proof is similar to the proof in [3, Section 4.7]. Let u be the holomorphic function from Proposition 3.5. From (3-30), it follows that all fibers of u near infinity are regular and are diffeomorphic to tori, hence all the regular fibers of u are diffeomorphic to tori, and so $u: X \rightarrow \mathbb{C}$ is an elliptic fibration. The ALH* coordinates define a submanifold near infinity by

$$(3-37) \quad \Sigma_0 = \{(x, y, t, z) : x = 0, t = 0\}.$$

This is clearly a smooth section of the model elliptic fibration over $U \equiv \mathbb{C} \setminus B_R(0)$ for some $R \gg 1$. It is moreover $J_{\mathfrak{M}}$ -holomorphic because for all $p \in \Sigma_0$,

$$(3-38) \quad T_p \Sigma_0 = \text{span}_p \left\{ \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\} = \{Y \in T_p \mathfrak{M} : e_1(Y) = e_4(Y) = 0\},$$

and $\text{span}\{e_1, e_4\}$ is invariant under $J_{\mathfrak{M}}$ by (3-9).

We view the section Σ_0 as a mapping $\sigma_0 : U \rightarrow X \setminus X_R$. We next want to perturb σ_0 to a section σ which is holomorphic with respect to J . Given any smooth (not necessarily holomorphic) section σ over U , we can define $\bar{\partial} \sigma \in \Gamma(\Lambda^{0,1}(U) \otimes \sigma^* T^{1,0}(F))$, where $T^{1,0}(F)$ is the $(1, 0)$ part of the vertical tangent bundle, by restricting the differential $\sigma_* \otimes \mathbb{C}$ to $T^{0,1}(U)$, and then projecting to the $(1, 0)$ part. Next, we use the 2-form $\Omega = \omega_1 + \sqrt{-1}\omega_3$, which is holomorphic with respect to J . If we insert the $T^{1,0}(F)$

component of $\bar{\partial}\sigma$ into Ω , we can define $\Omega \odot \bar{\partial}\sigma \in C^\infty(U, \mathbb{C})$, since $\Lambda^{1,0}(U) \otimes \Lambda^{0,1}(U) \cong \Lambda^{1,1}(U)$ is a trivial bundle. Denoting $h(u, \bar{u}) = \Omega \odot \bar{\partial}\sigma_0$, from basic theory of the $\bar{\partial}$ -operator in U , we can solve the equation $(\partial/\partial\bar{u})H = h$ on U . Choose an arbitrary point $p \in U$ and an affine holomorphic fiber coordinate w over a small neighborhood U_p of p in U . Then $\{w = 0\}$ is a local holomorphic section over U_p , and the holomorphic 2-form can be written as $\Omega = f(u) du \wedge dw$, where $f(u)$ is nowhere vanishing. It is easy to see that the smooth local sections

$$(3-39) \quad \sigma_{h,p} \equiv \left(\frac{H(u, \bar{u})}{f(u)} \right) \frac{\partial}{\partial w}$$

over U_p patch up to a well-defined smooth section σ_h over U , independent of the choice of local w coordinate. Consequently, the section $\sigma \equiv \sigma_0 - \sigma_h$ is a holomorphic section defined over all of U .

After identifying U with a punctured disc Δ^* using $z = u^{-1}$ as a holomorphic coordinate, we can then identify the elliptic surface with $(\Delta_z^* \times \mathbb{C}_w)/(\mathbb{Z} \oplus \mathbb{Z})$, with the action given by

$$(3-40) \quad (m, n) \cdot (z, w) = (z, w + mt_1(z) + nt_2(z)), \quad \text{where } (m, n) \in \mathbb{Z} \oplus \mathbb{Z}$$

(t_1, t_2 are the periods), such that $\{w = 0\}$ defines σ ; see for example [18, pages 369–370]. Consequently, there exists a compactification of X to an elliptic surface S such that $X = S \setminus F$, where F is the fiber at infinity. Since the cross-sections are diffeomorphic to nilmanifolds of degree b , the only possibility is that the monodromy is of type I_b . It is easy to see that the form Ω is then a meromorphic 2-form on S with a pole of order 1 along F , which implies that F has multiplicity 1. Since $\text{div}(\Omega) = -F$, we have that $K_S = -[F]$. From Corollary 1.4, $b^1(X) = 0$. A Mayer–Vietoris argument similar to that in the proof of Corollary 1.4 above shows that $b^1(S) = 0$. Arguing exactly as in [4, Theorem 3.3], we see that S is a rational elliptic surface, with projection $u: S \rightarrow \mathbb{P}^1$, so is the blowup of \mathbb{P}^2 in 9 points. Consequently, there exists a (-1) -curve E (the exceptional divisor of the last blowup). The adjunction formula then implies that $K_S \cdot E = -1$, so the condition that $K_S = -[F]$ implies that there are no multiple fibers, and E is a global section; see [15, Proposition 4.1].

References

- [1] **C Birkar**, *Singularities of linear systems and boundedness of Fano varieties*, Ann. of Math. 193 (2021) 347–405 [MR](#) [Zbl](#)
- [2] **G Chen, X Chen**, *Gravitational instantons with faster than quadratic curvature decay, II*, J. Reine Angew. Math. 756 (2019) 259–284 [MR](#) [Zbl](#)
- [3] **G Chen, X Chen**, *Gravitational instantons with faster than quadratic curvature decay, I*, Acta Math. 227 (2021) 263–307 [MR](#) [Zbl](#)
- [4] **G Chen, X Chen**, *Gravitational instantons with faster than quadratic curvature decay, III*, Math. Ann. 380 (2021) 687–717 [MR](#) [Zbl](#)
- [5] **G Chen, J Viaclovsky**, *Gravitational instantons with quadratic volume growth*, J. Lond. Math. Soc. 109 (2024) art. id. e12886 [MR](#) [Zbl](#)

- [6] **Y Chen**, *Calabi–Yau metrics of Calabi type with polynomial rate of convergence*, preprint (2024) [arXiv 2404.18070](#)
- [7] **C van Coevering**, *A construction of complete Ricci-flat Kähler manifolds*, preprint (2008) [arXiv 0803.0112](#)
- [8] **T C Collins, A Jacob, Y-S Lin**, *Special Lagrangian submanifolds of log Calabi–Yau manifolds*, *Duke Math. J.* 170 (2021) 1291–1375 [MR](#) [Zbl](#)
- [9] **T C Collins, V Tosatti**, *A singular Demailly–Păun theorem*, *C. R. Math. Acad. Sci. Paris* 354 (2016) 91–95 [MR](#) [Zbl](#)
- [10] **R J Conlon, H-J Hein**, *Classification of asymptotically conical Calabi–Yau manifolds*, *Duke Math. J.* 173 (2024) 947–1015 [MR](#) [Zbl](#)
- [11] **J-P Demailly**, *Complex analytic and differential geometry*, book manuscript (2012) Available at <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>
- [12] **I V Dolgachev**, *Classical algebraic geometry: a modern view*, Cambridge Univ. Press (2012) [MR](#) [Zbl](#)
- [13] **H Grauert**, *Über Modifikationen und exzeptionelle analytische Mengen*, *Math. Ann.* 146 (1962) 331–368 [MR](#) [Zbl](#)
- [14] **H Grauert, O Riemenschneider**, *Verschwindungssätze für analytische Kohomologiegruppen auf komplexen Räumen*, *Invent. Math.* 11 (1970) 263–292 [MR](#) [Zbl](#)
- [15] **B Harbourne, W E Lang**, *Multiple fibers on rational elliptic surfaces*, *Trans. Amer. Math. Soc.* 307 (1988) 205–223 [MR](#) [Zbl](#)
- [16] **M Haskins, H-J Hein, J Nordström**, *Asymptotically cylindrical Calabi–Yau manifolds*, *J. Differential Geom.* 101 (2015) 213–265 [MR](#) [Zbl](#)
- [17] **H-J Hein**, *Weighted Sobolev inequalities under lower Ricci curvature bounds*, *Proc. Amer. Math. Soc.* 139 (2011) 2943–2955 [MR](#) [Zbl](#)
- [18] **H-J Hein**, *Gravitational instantons from rational elliptic surfaces*, *J. Amer. Math. Soc.* 25 (2012) 355–393 [MR](#) [Zbl](#)
- [19] **H-J Hein, S Sun, J Viaclovsky, R Zhang**, *Nilpotent structures and collapsing Ricci-flat metrics on the $K3$ surface*, *J. Amer. Math. Soc.* 35 (2022) 123–209 [MR](#) [Zbl](#)
- [20] **C D Hill, M Taylor**, *Integrability of rough almost complex structures*, *J. Geom. Anal.* 13 (2003) 163–172 [MR](#) [Zbl](#)
- [21] **L Hörmander**, *An introduction to complex analysis in several variables*, 3rd edition, North-Holland Math. Libr. 7, North-Holland, Amsterdam (1990) [MR](#) [Zbl](#)
- [22] **Y Kawamata**, *Pluricanonical systems on minimal algebraic varieties*, *Invent. Math.* 79 (1985) 567–588 [MR](#) [Zbl](#)
- [23] **J Kollár, Y Miyaoka, S Mori, H Takagi**, *Boundedness of canonical \mathbb{Q} -Fano 3-folds*, *Proc. Japan Acad. Ser. A Math. Sci.* 76 (2000) 73–77 [MR](#) [Zbl](#)
- [24] **J Kollár, S Mori**, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math. 134, Cambridge Univ. Press (1998) [MR](#) [Zbl](#)
- [25] **P B Kronheimer**, *A Torelli-type theorem for gravitational instantons*, *J. Differential Geom.* 29 (1989) 685–697 [MR](#) [Zbl](#)
- [26] **C Li**, *On sharp rates and analytic compactifications of asymptotically conical Kähler metrics*, *Duke Math. J.* 169 (2020) 1397–1483 [MR](#) [Zbl](#)

- [27] **V Minerbe**, *Rigidity for multi-Taub-NUT metrics*, J. Reine Angew. Math. 656 (2011) 47–58 [MR](#) [Zbl](#)
- [28] **M V Nori**, *Zariski's conjecture and related problems*, Ann. Sci. École Norm. Sup. 16 (1983) 305–344 [MR](#) [Zbl](#)
- [29] **T Ohsawa**, *Vanishing theorems on complete Kähler manifolds*, Publ. Res. Inst. Math. Sci. 20 (1984) 21–38 [MR](#) [Zbl](#)
- [30] **I Shimada**, *Remarks on fundamental groups of complements of divisors on algebraic varieties*, Kodai Math. J. 17 (1994) 311–319 [MR](#) [Zbl](#)
- [31] **S Sun, R Zhang**, *Complex structure degenerations and collapsing of Calabi–Yau metrics*, preprint (2019) [arXiv 1906.03368](#)
- [32] **S Sun, R Zhang**, *A Liouville theorem on asymptotically Calabi spaces*, Calc. Var. Partial Differential Equations 60 (2021) art. id. 103 [MR](#) [Zbl](#)
- [33] **S Sun, R Zhang**, *Collapsing geometry of hyperkähler 4-manifolds and applications*, Acta Math. 232 (2024) 325–424 [MR](#) [Zbl](#)
- [34] **S Takayama**, *Simple connectedness of weak Fano varieties*, J. Algebraic Geom. 9 (2000) 403–407 [MR](#) [Zbl](#)
- [35] **G Tian, S-T Yau**, *Complete Kähler manifolds with zero Ricci curvature, I*, J. Amer. Math. Soc. 3 (1990) 579–609 [MR](#) [Zbl](#)
- [36] **C Voisin**, *Coniveau 2 complete intersections and effective cones*, Geom. Funct. Anal. 19 (2010) 1494–1513 [MR](#) [Zbl](#)
- [37] **S T Yau**, *On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I*, Comm. Pure Appl. Math. 31 (1978) 339–411 [MR](#) [Zbl](#)
- [38] **S T Yau**, *The role of partial differential equations in differential geometry*, from “Proceedings of the International Congress of Mathematicians, I” (O Lehto, editor), Acad. Sci. Fennica, Helsinki (1980) 237–250 [MR](#) [Zbl](#)

H-JH: *Mathematisches Institut, Universität Münster
Münster, Germany*

SS: *Institute for Advanced Study in Mathematics, Zhejiang University
Hangzhou, China*

JV: *Department of Mathematics, University of California, Irvine
Irvine, CA, United States*

RZ: *Department of Mathematics, University of California
San Diego, CA, United States*

hhein@uni-muenster.de, songsun@zju.edu.cn, jviaclov@uci.edu, rzhang573@wisc.edu

Proposed: Simon Donaldson

Seconded: Tobias H Colding, Mark Gross

Received: 12 September 2023

Revised: 19 July 2024

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITORS

Robert Lipshitz University of Oregon
lipshitz@uoregon.edu
András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Mohammed Abouzaid	Stanford University abouzaid@stanford.edu	Mark Gross	University of Cambridge mgross@dpms.cam.ac.uk
Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Arend Bayer	University of Edinburgh arend.bayer@ed.ac.uk	Sándor Kovács	University of Washington skovacs@uw.edu
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	Haynes Miller	Massachusetts Institute of Technology hmr@math.mit.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Aaron Naber	Northwestern University anaber@math.northwestern.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Peter Ozsváth	Princeton University petero@math.princeton.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Benson Farb	University of Chicago farb@math.uchicago.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David M Fisher	Rice University davidfisher@rice.edu	Roman Sauer	Karlsruhe Institute of Technology roman.sauer@kit.edu
Mike Freedman	Microsoft Research michaelf@microsoft.com	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
David Gabai	Princeton University gabai@princeton.edu	Natasa Sesum	Rutgers University natasas@math.rutgers.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Cameron Gordon	University of Texas gordon@math.utexas.edu	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2025 is US \$865/year for the electronic version, and \$1210/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2025 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 29 Issue 5 (pages 2251–2782) 2025

Quantitative Thomas–Yau uniqueness	2251
YANG LI	
Knot surgery formulae for instanton Floer homology, I: The main theorem	2269
ZHENKUN LI and FAN YE	
Morse actions of discrete groups on symmetric spaces: local-to-global principle	2343
MICHAEL KAPOVICH, BERNHARD LEEB and JOAN PORTI	
Nearly geodesic immersions and domains of discontinuity	2391
COLIN DAVALO	
Graded character sheaves, HOMFLY-PT homology, and Hilbert schemes of points on \mathbb{C}^2	2463
QUOC P HO and PENGHUI LI	
Asymptotically Calabi metrics and weak Fano manifolds	2547
HANS-JOACHIM HEIN, SONG SUN, JEFFREY VIACLOVSKY and RUOBING ZHANG	
The distribution of critical graphs of Jenkins–Strebel differentials	2571
FRANCISCO ARANA-HERRERA and AARON CALDERON	
Tian’s stabilization problem for toric Fanos	2609
CHENZI JIN and YANIR A RUBINSTEIN	
On the logarithmic slice filtration	2653
FEDERICO BINDA, DOOSUNG PARK and PAUL ARNE ØSTVÆR	
Joyce structures on spaces of quadratic differentials	2695
TOM BRIDGELAND	
Big monodromy for higher Prym representations	2733
AARON LANDESMAN, DANIEL LITT and WILL SAWIN	