



# *Geometry & Topology*

Volume 30 (2026)

**Prym representations and twisted cohomology  
of the mapping class group with level structures**

XIYAN ZHONG

# Prym representations and twisted cohomology of the mapping class group with level structures

XIYAN ZHONG

We compute the twisted cohomology of the mapping class group with level structures, with coefficients in the  $r$ -tensor powers of the Prym representations for any positive integer  $r$ . When  $r \geq 2$ , we show that the cohomology exhibits instability for large genus, whereas it remains stable for  $r = 0$  or  $r = 1$ . As a corollary, we prove that the symplectic Prym representation associated with any finite abelian regular cover of a nonclosed finite-type surface is infinitesimally rigid.

1. Introduction	155
2. Preliminaries	161
3. Stable cohomology of $\text{Mod}(\Sigma_{g,p}^b)$ with coefficients in $H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}$	168
4. Moduli spaces of Riemann surfaces with level structures	176
5. Twisted cohomology of the level- $l$ mapping class groups	184
6. Infinitesimal rigidity of symplectic Prym representations	196
Acknowledgements	200
References	201

## 1 Introduction

The (pure) mapping class group  $\text{Mod}(S)$  of an oriented surface  $S$  is the group of isotopy classes of orientation-preserving homeomorphisms that fix the boundary pointwise, including punctures and boundary components. For each integer  $\ell \geq 2$ , the level- $\ell$  mapping class group of  $S$  is the finite-index subgroup of  $\text{Mod}(S)$  consisting of elements that act trivially on  $H_1(S; \mathbb{Z}/\ell)$ . An important family of representations defined on these level subgroups are the Prym representations, first studied by Looijenga [1997].

The main result of this paper is the computation of the cohomology of the level- $\ell$  mapping class group with twisted coefficients given by the  $r$ -tensor powers of the Prym representation, for arbitrary  $r \geq 1$ , in a range where the genus of  $S$  is sufficiently large. Our calculations uncover a new phenomenon: when  $r \geq 2$ , this twisted cohomology exhibits instability with respect to the genus of  $S$ , in contrast to the known cohomological stability of mapping class groups and the stable behavior in the case  $r = 1$ .

As another consequence of our work, we prove that the Prym representation is infinitesimally rigid when the genus of  $S$  is at least 41.

The central technical innovation of this paper is the construction of a refined bordification of the moduli space of Riemann surfaces with level structures in Section 4. The boundary strata of this space are described using the theory of orbit configuration spaces, which allows us to compute its cohomology.

MSC2020: 20F05, 20J06, 57K20, 57S05.

© 2026 MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

By analyzing a specific Leray spectral sequence associated with this bordified moduli space and applying the theory of mixed Hodge structures, we derive our main results.

Since the statement of infinitesimal rigidity for the Prym representations is more straightforward, we begin by presenting this result.

### 1.1 Rigidity theorem

Let  $\tilde{S} \rightarrow S$  be a regular cover whose deck transformation group is a finite abelian group  $A$ . Let  $\text{Mod}(S, A)$  denote the subgroup of  $\text{Mod}(S)$  consisting of mapping classes that lift to homeomorphisms of  $\tilde{S}$ . The induced action

$$\text{Mod}(S, A) \rightarrow \text{Aut}(H^1(\tilde{S}; \mathbb{Q}))$$

defines the associated *Prym representation*. When  $S$  is closed, Looijenga [1997, Theorem 2.5] determined the image of the Prym representation up to finite index.

When  $S$  is not closed, let  $\hat{S}$  denote the closed surface obtained by gluing disks to all boundary components and filling in all punctures of  $\tilde{S}$ . Any lift of a homeomorphism of  $S$  to  $\tilde{S}$  preserves homology classes of loops around boundary components of  $\tilde{S}$ , and commutes with the deck group  $A$ . It is therefore natural to consider a refined version of the Prym representation:

$$\Phi : \text{Mod}(S, A) \rightarrow \text{Aut}(H^1(\hat{S}; \mathbb{R}))^A.$$

We refer to this as the *symplectic Prym representation*. We work over  $\mathbb{R}$  rather than  $\mathbb{Q}$  because, when the image of  $\Phi$  is a Lie group, this allows us to study its rigidity in the following sense.

Let  $\Gamma$  be a finitely generated group and  $G$  a Lie group. A homomorphism  $\Phi : \Gamma \rightarrow G$  is called *infinitesimally rigid* if

$$H^1(\Gamma; \mathfrak{g}) = 0,$$

where  $\mathfrak{g}$  is the Lie algebra of  $G$ , and  $\Gamma$  acts on  $\mathfrak{g}$  via the composition of  $\Phi$  and the adjoint representation  $\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$ . Intuitively, infinitesimal rigidity means that  $\Phi$  admits no nontrivial first-order deformations. We establish this kind of rigidity for the symplectic Prym representation:

**Theorem A** *Let  $S$  be a nonclosed surface of genus  $g \geq 41$ . For any finite abelian cover  $\tilde{S} \rightarrow S$  with deck transformation group  $A$ , the associated symplectic Prym representation*

$$\Phi : \text{Mod}(S, A) \rightarrow \text{Aut}(H^1(\hat{S}; \mathbb{R}))^A$$

*is infinitesimally rigid.*

**Remark 1.1** Infinitesimal rigidity implies other kinds of rigidity as follows:

- (1) Weil [1964, page 152] proved that an infinitesimally rigid representation is *locally rigid*. A representation  $\Phi$  is locally rigid if  $[\Phi]$  is an isolated point in  $\text{Hom}(\Gamma, G)/G$ , i.e., any  $\Phi' \in \text{Hom}(\Gamma, G)$  sufficiently close to  $\Phi$  is conjugate to  $\Phi$ . The converse is false: locally rigidity does not imply infinitesimal rigidity; see for instance [Lubotzky and Magid 1985, (2.10.4)].

- (2) For local systems on quasiprojective varieties, infinitesimal rigidity also implies *cohomological rigidity* when the Lie group  $G$  is semisimple. See the definition of cohomological rigidity in [Esnault and Groechenig 2018] and [Klevdal and Patrikis 2020].

## 1.2 Computations of twisted cohomology

To state our results on the twisted cohomology of mapping class groups with level structures, we introduce the following setup. Let  $\Sigma_{g,p}^b$  denote a genus- $g$  surface with  $p$  (ordered) punctures and  $b$  boundary components. We omit  $p$  or  $b$  when it is 0. Let  $\text{Mod}_{g,p}^b$  denote the mapping class group of  $\Sigma_{g,p}^b$ , and define the level- $\ell$  mapping class group of  $\Sigma_{g,p}^b$  by

$$\text{Mod}_{g,p}^b(\ell) = \text{Ker}(\text{Mod}_{g,p}^b \rightarrow \text{Aut}(H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell))).$$

We remark that when  $p + b \geq 2$ , the kernel of the action of  $\text{Mod}_{g,p}^b$  on  $H_1(\Sigma_g; \mathbb{Z}/\ell)$  strictly contains  $\text{Mod}_{g,p}^b(\ell)$ .

Let  $\mathcal{D} = H_1(\Sigma_g; \mathbb{Z}/\ell)$ . Consider the composition of group homomorphisms

$$\pi_1(\Sigma_{g,p}^b) \rightarrow H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_g; \mathbb{Z}/\ell),$$

where the second map is induced by the embedding  $\Sigma_{g,p}^b \rightarrow \Sigma_g$  obtained by gluing disks to all boundary components and filling in all punctures of  $\Sigma_{g,p}^b$ . This group homomorphism induces a regular cover of  $\Sigma_{g,p}^b$  with deck transformation group  $\mathcal{D}$ , which we denote by  $S_{\mathcal{D}} \rightarrow \Sigma_{g,p}^b$ . By covering-space theory, elements of  $\text{Mod}_{g,p}^b(\ell)$  lift to homeomorphisms of  $S_{\mathcal{D}}$  that fix all punctures and boundary components pointwise. Define

$$\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) := H^1(S_{\mathcal{D}}; \mathbb{Q}),$$

which is a  $\text{Mod}_{g,p}^b(\ell)$ -module. We refer to  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})$  as the Prym representation of  $\text{Mod}_{g,p}^b(\ell)$ . Our central result is the computation of the cohomology

$$H^k(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$$

in a range where the genus  $g$  is sufficiently large relative to  $k$ , for arbitrary  $r \geq 1$ :

**Theorem B** *Let  $A'_r(\ell)^\bullet$  be the graded  $\mathbb{Q}$ -vector space defined in (5-1) of Section 5. The symmetric group  $S_r$  acts naturally on  $A'_r(\ell)^\bullet$  by definition, and also on  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}$  by permuting tensor factors. There is a graded  $S_r$ -equivariant map of  $H^\bullet(\text{Mod}_{g,p}^b(\ell); \mathbb{Q})$ -modules*

$$H^\bullet(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}) \otimes A'_r(\ell)^\bullet \rightarrow H^{\bullet-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

Here  $A'_r(\ell)^\bullet$  is a subspace of the rational cohomology ring of a moduli space denoted by  $\mathcal{C}_{g,r}(\ell)$ . This moduli space contains the moduli space  $\mathcal{M}_{g,r}(\ell)$  of Riemann surfaces homeomorphic to  $\Sigma_{g,r}$  with a level- $\ell$  structure as an open subvariety. In Section 4, we construct  $\mathcal{C}_{g,r}(\ell)$  explicitly, which is a bordification of  $\mathcal{M}_{g,r}(\ell)$  with boundary strata indexed by orbit configuration spaces. We compute in

**Theorem 4.8** the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$  for sufficiently large  $g$ . As defining  $A'_r(\ell)^\bullet$  requires additional notation from [Section 4](#), we postpone its formal definition to [Section 5](#).

**Theorem B** enables us to compute the rational dimension of

$$H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$$

explicitly, thanks to Putman’s theorem [[2022](#), Theorem A], which states that

$$H^k(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}) \cong H^k(\text{Mod}_{g,p}^b; \mathbb{Q}) \quad \text{for } g \geq 2k^2 + 7k + 2.$$

By the Madsen–Weiss theorem [[2007](#), Theorem 1.1.1] and Looijenga’s extension to punctured surfaces [[1996](#), Proposition 2.2], this implies that

$$H^\bullet(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}) \cong \mathbb{Q}[e_1, \dots, e_p] \otimes \mathbb{Q}[\kappa_1, \kappa_2, \dots],$$

where  $\kappa_i \in H^{2i}(\text{Mod}_{g,p}^b; \mathbb{Q})$  ( $i \geq 1$ ) are the Miller–Morita–Mumford classes [[Miller 1986](#); [Morita 1987](#)], and  $e_j \in H^2(\text{Mod}_{g,p}^b; \mathbb{Q})$  is the Euler class associated with the central extension

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_{g,p-1}^{b+1} \rightarrow \text{Mod}_{g,p}^b \rightarrow 1$$

obtained by blowing up the  $j$ -th puncture of  $\Sigma_{g,p}^b$  to a boundary component. The central  $\mathbb{Z}$  is generated by the Dehn twist around this boundary component. See also [[Morita 1987](#), Section 7] for an alternative definition of  $e_j$ .

**Remark 1.2** From the definition of  $(A'_{r,\ell})^\bullet$  (see (5-1)), it follows that all its elements are of even degree. Therefore, **Theorem B** implies that for odd  $k$ ,

$$H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) = 0 \quad \text{for } g \geq 2k^2 + 7k + 2.$$

In particular, **Theorem A** follows as a corollary of the special case  $k = 3, r = 2$ .

### 1.3 Instability and historical remarks

Computations via **Theorem B** show that for  $r \geq 2$  and an even number  $k$ ,

$$H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$$

is independent of  $b$ , but depends on  $g, p$ , and  $\ell$  for  $r \geq 2$ . This instability arises from the fact that our calculations involve orbit configuration spaces, which depend on the deck transformation group  $\mathcal{D} = H_1(\Sigma_g; \mathbb{Z}/\ell)$ .

This instability also helps us identify relationship with some other twisted cohomology groups related to mapping class groups, which we now recall.

The mapping class group  $\text{Mod}_{g,p}^b$  acts nontrivially on  $H^1(\Sigma_{g,p}^b; \mathbb{Q})$ . For any positive integer  $r$ , the twisted cohomology

$$(1-1) \quad H^{k-r}(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r})$$

has been studied by Looijenga [[1996](#)], Kawazumi [[2008](#)], Randal-Williams [[2018](#)] and Putman [[2022](#)].

In the case  $p = b = 0$ , this twisted cohomology (1-1) was computed by Looijenga [1996, Corollary 3.3] for  $g \geq \frac{3}{2}k + 1$ . He constructed an  $S_r$ -equivariant map analogous to the one in Theorem B, which allowed him to further compute the cohomology of  $\text{Mod}_g$  with coefficients in any irreducible representation of  $\text{Sp}_{2g}(\mathbb{C})$ .

Kawazumi [2008, Theorem 1.A] computed the twisted cohomology (1-1) for  $\Sigma_g^b$  with  $b \geq 1$ , also in the same stable range, but over  $\mathbb{Z}$ , using a different inductive approach. Kawazumi’s results exhibit stability with respect to both  $g$  and  $b$ .

Randal-Williams [2018, Appendix B] identified (1-1) as a specific  $S_r$ -module in the cases of  $\Sigma_g$  and  $\Sigma_g^1$ . Later, Kupers and Randal-Williams [2020, Theorem 3.15] incorporated additional structure into the coefficients, allowing them to compute the cohomology of  $\text{Mod}_g^1$  with coefficients in any algebraic representation of  $\text{Sp}_{2g}$ .

**Remark 1.3** Randal-Williams suggested that it might be possible to generalize the results of Theorem B from coefficients in tensor powers to coefficients in Schur functors. We plan to explore this direction in future work.

For completeness, we derive from the work of Looijenga and Kawazumi the remaining case of (1-1) where  $b = 0$  and  $p \geq 1$ , and state the general result for any nonclosed surface  $\Sigma_{g,p}^b$  in Theorem 3.9.

The twisted cohomology (1-1) is related to the cohomology

$$H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$$

in the following way. Recall that  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) = H^1(S_{\mathfrak{D}}; \mathbb{Q})$ , where  $S_{\mathfrak{D}} \rightarrow \Sigma_{g,p}^b$  is the regular cover with deck group  $\mathfrak{D} = H_1(\Sigma_g; \mathbb{Z}/\ell)$ . The covering map induces a natural map between tensor powers

$$H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r} \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r},$$

which in turn induces a map on twisted cohomology:

$$(1-2) \quad H^{k-r}(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}) \rightarrow H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}).$$

Putman [2022, Theorem C] studied the case  $r = 1$  and showed that the map (1-2) is an isomorphism when  $g \geq 2k^2 + 7k + 1$ . This result implies that for  $r = 1$ , the right-hand side of (1-2) is independent of  $g$ ,  $b$ , and  $\ell$ , since the left-hand side is.

For general  $r \geq 2$ , Putman [2022, Remark 1.8] conjectured the following:

**Conjecture 1.4** For  $r \geq 2$ , the map (1-2) is not an isomorphism.

Since Theorem B shows that the right-hand side of (1-2) is not stable with respect to the genus  $g$ , in contrast with the known stability of the left-hand side in (1-1), we show that Putman’s conjecture is true:

**Corollary 1.5** For  $r \geq 2$  and  $k$  even, the map (1-2) is not an isomorphism when

$$g \geq \max(\frac{3}{2}k + 1, 2k^2 + 7k + 2) = 2k^2 + 7k + 2.$$

**Remark 1.6** For  $k$  odd, both sides of the map (1-2) vanish when  $g \geq 2k^2 + 7k + 2$ .

### 1.4 An example

To gain more insight into how these twisted cohomology groups in the map (1-2) differ when  $r \geq 2$ , we present an example for the case  $r = 2$ . By taking  $r = 2$  in Theorem 3.9, we obtain

$$H^{\bullet-2}(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes 2}) \cong H^{\bullet}(\text{Mod}_{g,p}^b; \mathbb{Q}) \otimes (u_1 u_2 \mathbb{Q}[u_1, u_2] \oplus \mathbb{Q}[u_{\{1,2\}}] a_{\{1,2\}}),$$

in degrees  $k$  such that  $g \geq \frac{3}{2}k + 1$ , with  $\deg(u_1) = \deg(u_2) = \deg(a_{\{1,2\}}) = 2$ . Notably, this twisted cohomology is independent of the genus  $g$ .

In contrast, taking  $r = 2$  in Theorem B yields

$$\begin{aligned} H^{\bullet-2}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2}) \\ \cong H^{\bullet}(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}) \otimes \left( v_1 v_2 \mathbb{Q}[v_1, v_2] \oplus \bigoplus_{d \in \mathcal{D}} \mathbb{Q}[v_{\{1<2\},d}] a_{\{1<2\},d} \right) \end{aligned}$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ , with  $\deg(v_1) = \deg(v_2) = \deg(a_{\{1<2\},d}) = 2$ . Since there is a nontrivial summand corresponding to each  $d \in \mathcal{D} = (\mathbb{Z}/\ell)^{2g}$ , the twisted cohomology group  $H^{\bullet-2}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2})$  exhibits a dependence on the genus  $g$ .

The key distinction between these two twisted cohomology groups lies in the nature of the cohomology classes  $a_{\{1,2\}}$  and  $a_{\{1<2\},d}$ . The class  $a_{\{1,2\}}$  is associated with the moduli space  $\mathcal{C}_{g,2}$  of closed genus- $g$  Riemann surfaces with two (not necessarily distinct) marked points. More precisely,  $a_{\{1,2\}}$  represents the Poincaré dual of the subvariety of  $\mathcal{C}_{g,2}$  where the two marked points coincide (i.e.,  $x_2 = x_1$ ).

In contrast, for each  $d \in \mathcal{D} = (\mathbb{Z}/\ell)^{2g}$ , the class  $a_{\{1<2\},d}$  relates to the moduli space  $\mathcal{C}_{g,2}(\ell)$  that we construct in Section 4. This moduli space is a bordification of  $\mathcal{M}_{g,2}(\ell)$ , the moduli space of Riemann surfaces homeomorphic to  $\Sigma_{g,2}$  with level- $\ell$  structures. One can roughly view  $\mathcal{C}_{g,2}(\ell)$  as the moduli space of two (not necessarily distinct) marked points on the regular  $\mathcal{D}$ -cover of a closed genus- $g$  Riemann surface. The class  $a_{\{1<2\},d}$  is defined as the Poincaré dual of the subvariety of  $\mathcal{C}_{g,2}(\ell)$  where the marked points  $y_1, y_2$  on the  $\mathcal{D}$ -cover satisfy the relation  $y_2 = d \cdot y_1$ .

This distinction illustrates that the additional classes in  $H^{\bullet-2}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2})$  arise from the richer geometric structure of the moduli space  $\mathcal{C}_{g,2}(\ell)$  compared to  $\mathcal{C}_{g,2}$ .

### 1.5 Proof ideas

To conclude the introduction, we outline our approach to proving Theorem B.

Our strategy is inspired by Looijenga’s computation [1996, Corollary 3.3] of  $H^{\bullet}(\text{Mod}_g; H^1(\Sigma_g; \mathbb{Q})^{\otimes r})$ . Looijenga studied the Leray spectral sequence associated with the  $r$ -fold fiber product of the universal curve  $\mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ . Since we work with the level- $\ell$  mapping class group, we shift our focus from  $\mathcal{M}_g$  to  $\mathcal{M}_{g,r}(\ell)$ , the moduli space of Riemann surfaces homeomorphic to  $\Sigma_{g,r}$  equipped with a level- $\ell$  structure. Our approach proceeds as follows:

- (1) We reinterpret  $\mathcal{M}_{g,r}(\ell)$  as the moduli space of one marked point on  $\Sigma_{g,r}$  and  $(r - 1)$  marked points on the regular  $H_1(\Sigma_g; \mathbb{Z}/\ell)$ -cover of  $\Sigma_{g,r}$ , subject to certain conditions.

- (2) We construct a partial bordification of  $\mathcal{M}_{g,r}(\ell)$ , denoted by  $\mathcal{C}_{g,r}(\ell)$ , whose boundary strata are described in terms of orbit configuration spaces. We remark that, unlike Looijenga’s bordification, this space  $\mathcal{C}_{g,r}(\ell)$  is not the total space of the  $r$ -fold fiber product of  $\mathcal{M}_{g,1}(\ell) \rightarrow \mathcal{M}_g(\ell)$ .
- (3) We compute the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$  for  $g$  sufficiently large, stated in [Theorem 4.8](#).
- (4) We construct a map  $\mathcal{C}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$ , where the cohomology of the fiber contains a summand isomorphic to  $\mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}$ , the  $r$ -fold tensor power of the Prym representation.
- (5) By a theorem of Deligne [[1974, Section 8.1](#)] (see [Theorem 2.17](#)), the Leray spectral sequence associated with the map  $\mathcal{C}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  degenerates at the  $E_2$ -page.
- (6) We compute the  $E_2$ -page of the Leray spectral sequence, and using mixed Hodge theory, we identify  $H^\bullet(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r})$  as a subspace of  $H^\bullet(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q})$ .
- (7) Finally, to extend our results from  $\text{Mod}_{g,1}(\ell)$  to  $\text{Mod}_{g,p}^b(\ell)$  with  $p + b \geq 1$ , we apply Putman’s theory [[2022](#)] of partial level- $\ell$  representations.

**Outline** In [Section 2](#), we introduce key preliminaries, including background on mapping class groups, group cohomology, and mixed Hodge theory. In [Section 3](#), we synthesize the results of Looijenga and Kawazumi to give a complete account of the cohomology groups (1-1). In [Section 4](#), we construct the space  $\mathcal{C}_{g,r}(\ell)$  and compute its rational cohomology for  $g$  sufficiently large. In [Section 5](#), we present the proof of [Theorem B](#). In [Section 6](#), we prove [Theorem A](#).

## 2 Preliminaries

### 2.1 Stable cohomology of the mapping class group

We summarize key stability results for the cohomology of the mapping class group.

**Theorem 2.1** (Harer stability [[1985](#)])  $H^k(\text{Mod}_{g,p}^b; \mathbb{Z})$  is independent of  $g$  and  $b$  in degrees  $\leq N(g)$ .

**Remark 2.2** Here the number  $N(g)$  is the maximal degree  $N$  such that the two homomorphisms

$$H^N(\text{Mod}_{g+1,p}^b; \mathbb{Z}) \rightarrow H^N(\text{Mod}_{g,p}^{b+1}; \mathbb{Z}) \quad \text{and} \quad H^N(\text{Mod}_{g,p}^b; \mathbb{Z}) \rightarrow H^N(\text{Mod}_{g,p}^{b+1}; \mathbb{Z})$$

are isomorphisms. The original bound given by Harer [[1985](#)] was  $N(g; \mathbb{Z}) \geq \frac{1}{3}g$ , which was later improved by Ivanov [[1989](#)], Boldsen [[2012](#)] and Randal-Williams [[2016](#)] to  $N(g) \geq \frac{2}{3}(g - 1)$ .

Unlike the genus  $g$  and boundary components  $b$ , the stable cohomology of  $\text{Mod}_{g,p}^b$  does depend on the number of punctures  $p$ . For  $1 \leq i \leq p$ , we have an Euler class  $e_i \in H^2(\Sigma_{g,p}^b; \mathbb{Z})$ , as we mentioned in the introduction. Below, we provide an alternate definition following Looijenga [[1996, Section 2](#)] for future purposes:

- Let  $\mathcal{M}_{g,p}$  denote the moduli space of closed Riemann surfaces of genus  $g$  with  $p$  distinct marked points. Since  $\mathcal{M}_{g,p}$  is a rational classifying space for the mapping class group, we have

$$H^\bullet(\mathcal{M}_{g,p}; \mathbb{Q}) \cong H^\bullet(\text{Mod}_{g,p}; \mathbb{Q}).$$

- Consider the universal curve  $\pi : \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ . Let  $\theta$  be its relative tangent sheaf.
- For each  $i$ , define  $f_i : \mathcal{M}_{g,p} \rightarrow \mathcal{M}_{g,1}$  as the map that forgets all but the  $i$ -th marked point. Then, we define the Euler class  $e_i$  as the first Chern class:

$$e_i = c_1(f_i^*(\theta)) \in H^2(\text{Mod}_{g,p}; \mathbb{Q}).$$

**Theorem 2.3** [Looijenga 1996, Proposition 2.2] *The ring homomorphism*

$$H^*(\text{Mod}_g; \mathbb{Z})[e_1, \dots, e_p] \rightarrow H^*(\text{Mod}_{g,p}^b; \mathbb{Z})$$

*is an isomorphism in degrees  $\leq N(g)$ .*

The stable integral cohomology of the mapping class group is complicated, but the stable rational cohomology has a beautiful form. Conjectured by Mumford [1983], proved by Madsen and Weiss [2007, Theorem 1.1.1], the stable rational cohomology of  $\text{Mod}_g$  is isomorphic to polynomial ring generated by these Miller–Morita–Mumford classes [Miller 1986; Morita 1987]  $\kappa_i \in H^{2i}(\text{Mod}_g; \mathbb{Q})$  for  $i \geq 1$ . See also [Hatcher 2011], [Galatius 2013] and [Wahl 2013] for alternate proofs and expositions.

**Theorem 2.4** [Madsen and Weiss 2007, Theorem 1.1.1] *In degrees  $\leq N(g)$ , we have*

$$H^*(\text{Mod}_g; \mathbb{Q}) \cong \mathbb{Q}[\kappa_1, \kappa_2, \kappa_3, \dots].$$

Combining these two theorems above, in degrees  $\leq N(g)$ , we have

$$H^*(\text{Mod}_{g,p}^b; \mathbb{Q}) \cong \mathbb{Q}[e_1, \dots, e_p] \otimes \mathbb{Q}[\kappa_1, \kappa_2, \kappa_3, \dots].$$

## 2.2 Level- $l$ mapping class groups

Recall that the level- $\ell$  mapping class group  $\text{Mod}_{g,p}^b(\ell)$  is the subgroup of  $\text{Mod}_{g,p}^b$  which acts trivially on  $H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$ . It has many similar properties to  $\text{Mod}_{g,p}^b$ . For example:

**Proposition 2.5** [Putman 2022, Proposition 2.10] *Fix some  $g, p, b \geq 0$  such that  $\pi_1(\Sigma_{g,p}^{b+1})$  is nonabelian, and let  $\partial$  be a boundary component of  $\Sigma_{g,p}^{b+1}$ . Let  $\ell \geq 2$ . Then there is a central extension*

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_{g,p}^{b+1}(\ell) \rightarrow \text{Mod}_{g,p+1}^b(\ell) \rightarrow 1,$$

where the central  $\mathbb{Z}$  is generated by the Dehn twist  $T_\partial$ .

We also have the mod- $\ell$  version of the Birman exact sequence:

**Theorem 2.6** (mod- $l$  Birman exact sequence [Putman 2022, Theorem 2.8]) *Fix  $g, p, b \geq 0, \ell \geq 2$  such that  $\pi_1(\Sigma_{g,p}^b)$  is nonabelian. Let  $x_0$  be a puncture of  $\Sigma_{g,p+1}^b$ . There is a short exact sequence obtained by forgetting  $x_0$ :*

$$1 \rightarrow \text{PP}_{x_0}(\Sigma_{g,p}^b; \ell) \rightarrow \text{Mod}_{g,p+1}^b(\ell) \rightarrow \text{Mod}_{g,p}^b(\ell) \rightarrow 1,$$

where the level- $\ell$  point pushing group  $\text{PP}_{x_0}(\Sigma_{g,p}^b; \ell)$  is as follows:

- If  $p = b = 0$ , then  $\text{PP}_{x_0}(\Sigma_{g,p}^b; \ell) = \pi_1(\Sigma_{g,p}^b, x_0)$ .
- If  $p + b \geq 1$ , then  $\text{PP}_{x_0}(\Sigma_{g,p}^b; \ell) = \text{Ker}(\pi_1(\Sigma_{g,p}^b, x_0) \rightarrow H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_g; \mathbb{Z}/\ell))$ .

A natural question is whether the finite-index subgroup  $\text{Mod}_{g,p}^b(\ell)$  of  $\text{Mod}_{g,p}^b$  has the same stable cohomology as  $\text{Mod}_{g,p}^b$ . The answer is negative for integral cohomology, as exotic torsion elements in  $H^1(\text{Mod}_{g,p}^b(\ell); \mathbb{Z})$  have been discovered by Perron [2008], Sato [2010], and Putman [2012]. These elements do not arise from  $H^1(\text{Mod}_{g,p}^b; \mathbb{Z})$ , and they are not stable.

However, Putman established that the result does hold for rational cohomology:

**Theorem 2.7** [Putman 2022, Theorem A] *We have*

$$H^k(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}) \cong H^k(\text{Mod}_{g,p}^b; \mathbb{Q})$$

if  $g \geq 2k^2 + 7k + 2$ .

Putman also studied the twisted cohomology of  $\text{Mod}_{g,p}^b(\ell)$  with coefficients in tensor powers of  $H^1(\Sigma_{g,p}^b; \mathbb{Q})$  and showed this cohomology is stable by the following theorem:

**Theorem 2.8** [Putman 2022, Theorem B] *The map*

$$H^k(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}) \rightarrow H^k(\text{Mod}_{g,p}^b(\ell); H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r})$$

is an isomorphism when  $g \geq 2(k+r)^2 + 7k + 6r + 2$ .

### 2.3 Group cohomology

Here we review some useful theorems about group cohomology. We will apply these theorems in later sections.

Consider a group  $G$  and a finite-index subgroup  $H$ , along with a  $\mathbb{Z}[G]$ -module  $M$ . There is a natural map

$$\text{Res}_H^G : H^k(G; M) \rightarrow H^k(H; M)$$

in group cohomology obtained from the inclusion  $\mathbb{Z}[H] \hookrightarrow \mathbb{Z}[G]$ . We call it the restriction map. There is a “wrong-way” map, called the transfer map,

$$\text{cor}_H^G : H^k(H; M) \rightarrow H^k(G; M),$$

which satisfies:

**Proposition 2.9** [Brown 1982, Proposition 9.5] *If  $H$  is a finite-index subgroup of  $G$  with index  $[G : H]$ , then the composition of transfer maps and restriction maps is the multiplication map by  $[G : H]$ , in other words,  $\text{cor}_H^G \cdot \text{Res}_H^G = [G : H] \text{id}$ .*

**Remark 2.10** In particular, supposing  $H$  is a finite-index subgroup of  $G$ , if  $M$  is a  $\mathbb{Q}$ (or  $\mathbb{R}$ )-vector space, we see that  $\text{cor}_H^G$  is surjective and  $\text{Res}_H^G$  is injective.

**Proposition 2.11** [Putman 2014, Proposition 1.1] *Let  $X$  be a simplicial complex and  $G$  be a finite group acting simplicially on  $X$ . Then for all  $k \geq 0$ , we have*

$$H^k(X/G; \mathbb{Q}) \cong (H^k(X; \mathbb{Q}))^G.$$

The next technique we will use a lot is the Gysin sequence, which is a long exact sequence obtained from the second page of the Hochschild–Serre spectral sequence [1953, Proposition 7] associated to a central extension of groups:

**Proposition 2.12** (Gysin sequence [Hochschild and Serre 1953, Proposition 7]) *Consider a central extension*

$$1 \rightarrow \mathbb{Z} \rightarrow G \rightarrow K \rightarrow 1$$

and a  $\mathbb{Z}[K]$  module  $M$  (thus  $M$  is also a  $\mathbb{Z}[G]$  module through the map  $G \rightarrow K$ ). We have the long exact sequence

$$\dots \rightarrow H^{k-2}(K; M) \rightarrow H^k(K; M) \rightarrow H^k(G; M) \rightarrow H^{k-1}(K; M) \rightarrow H^{k+1}(K; M) \rightarrow \dots,$$

where  $H^{k-2}(K; M) \rightarrow H^k(K; M)$  is the differential on the  $E_2$ -page of the Hochschild–Serre spectral sequence.

**Remark 2.13** The geometric version of the Gysin sequence is that, for an oriented sphere bundle  $S^d \hookrightarrow E \rightarrow M$ , we have the long exact sequence

$$\dots \rightarrow H^{k-d-1}(M) \rightarrow H^k(M) \rightarrow H^k(E) \rightarrow H^{k-d}(M) \rightarrow H^{k+1}(M) \rightarrow \dots,$$

where the map  $H^{k-d-1}(M) \rightarrow H^k(M)$  is the wedge product with the Euler class, and the map  $H^k(E) \rightarrow H^{k-d}(M)$  is fiberwise integration.

There is a generalization of the Gysin sequence, called the Thom–Gysin sequence. This sequence is widely known, appearing in [Atiyah and Bott 1983, equation 1.16] and [Mumford et al. 1994, page 167] for example.

**Proposition 2.14** (Thom–Gysin sequence [Atiyah and Bott 1983, equation 1.16]) *Let  $X$  be a smooth complex variety, and let  $Y$  be an open smooth subvariety of  $X$  whose complement  $X \setminus Y$  has constant (real) codimension  $d$ . We then have the following long exact sequence:*

$$\dots \rightarrow H^{k-d}(X \setminus Y; \mathbb{Q}) \rightarrow H^k(X; \mathbb{Q}) \rightarrow H^k(Y; \mathbb{Q}) \rightarrow H^{k-d+1}(X \setminus Y; \mathbb{Q}) \rightarrow H^{k+1}(X; \mathbb{Q}) \rightarrow \dots$$

**Remark 2.15** When  $X$  is the total space of a sphere bundle and  $X \setminus Y$  is the zero section, the associated Thom–Gysin sequence is exactly the Gysin sequence. In general, one can derive the Thom–Gysin sequence by applying the Thom isomorphism theorem and the excision theorem to the tubular neighborhood of  $X \setminus Y$  in  $X$ .

**Remark 2.16** In most of our cases in Section 4, we apply the Thom–Gysin sequence to quasiprojective varieties, such as  $\mathcal{M}_{g,r}(\ell)$  for  $\ell \geq 3$ . The case of  $\mathcal{M}_{g,r}(2)$  is the only instance where we use the Thom–Gysin sequence on a quasiprojective orbifold, which is a quotient of a quasiprojective variety by a finite group. In this case, by Proposition 2.11, the Thom–Gysin sequence retains the same form, so we proceed without additional markings.

## 2.4 Deligne’s degeneration theorem

Our computations will be based on an important theorem due to Deligne [1974, Section 8.1], which was proved using mixed Hodge theory. Here we quote the version in Griffiths and Schmid’s survey [1975, page 42]:

**Theorem 2.17** (Deligne’s degeneration theorem) *Let  $E$  be a Kähler manifold,  $X$  a complex manifold, and  $f : E \rightarrow X$  a smooth, proper holomorphic mapping, which implies  $f$  is a differential fiber bundle whose fibers  $X_b$ , for  $b \in B$ , are compact Kähler manifolds. The corresponding Leray spectral sequence*

$$E_2^{p,q} = H^p(B, R^q f_*(\mathbb{Q})) \Rightarrow H^{p+q}(E; \mathbb{Q})$$

*degenerates at the second page, i.e.,  $E_2 = E_g$ . Here*

$$R^q f_*(\mathbb{Q}) \text{ comes from the presheaf } U \mapsto H^q(f^{-1}(U); \mathbb{Q}).$$

**Remark 2.18** Smooth quasiprojective varieties are Kähler manifolds, so we can apply Deligne’s degeneration theorem to  $\mathcal{C}_{g,r}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  for  $\ell \geq 3$ . This covers nearly all cases in Section 5, though there are two cases where we apply Deligne’s degeneration theorem to quasiprojective orbifolds: for  $\mathcal{C}_{g,r} \rightarrow \mathcal{M}_{g,1}$  in Section 3 and for  $\mathcal{C}_{g,r}(2) \rightarrow \mathcal{M}_{g,1}(2)$  in Section 5. Deligne’s degeneration theorem still applies in these cases, as demonstrated by the following argument:

Let  $E$  be a quasiprojective orbifold and  $X$  a complex orbifold. Suppose  $f : E \rightarrow X$  is a map of orbifolds that lifts to a map  $\tilde{f} : E' \rightarrow X'$  with the following properties:

- $E'$  and  $B'$  are quasiprojective varieties.
- $E' \rightarrow E$  and  $B' \rightarrow B$  are finite branched  $G$ -covers for a finite group  $G$ .
- $\tilde{f} : E' \rightarrow X'$  is a smooth, proper holomorphic mapping.

By applying Deligne’s degeneration Theorem 2.17 to  $\tilde{f} : E' \rightarrow X'$ , we obtain the Leray spectral sequence

$$(2-1) \quad E_2^{p,q} = H^p(B', R^q \tilde{f}_*(\mathbb{Q})) \Rightarrow H^{p+q}(E'; \mathbb{Q}),$$

which degenerates at the second page.

By Proposition 2.11, we have

$$H^{p+q}(E; \mathbb{Q}) \cong H^{p+q}(E'; \mathbb{Q})^G,$$

and by a similar argument, we obtain

$$H^p(B, R^q f_*(\mathbb{Q})) := H^p(B', R^q \tilde{f}_*(\mathbb{Q}))^G.$$

Restricting the spectral sequence (2-1) to  $G$ -invariants, we conclude that the Leray spectral sequence

$$E_2^{p,q} = H^p(B, R^q f_*(\mathbb{Q})) \Rightarrow H^{p+q}(E; \mathbb{Q})$$

also degenerates at the second page.

### 2.5 Mixed Hodge theory

Mixed Hodge theory is used in the proof of Deligne’s degeneration [Theorem 2.17](#), and is also a powerful tool for determining terms in spectral sequences. We will introduce some basic properties according to the survey by Griffiths and Schmid [\[1975\]](#). First, we start with definitions of pure Hodge structures.

**Definition 2.19** [\[Griffiths and Schmid 1975, Definition 1.1, 1.2\]](#) Let  $H_{\mathbb{R}}$  be a finite-dimensional real vector space, and  $H_{\mathbb{Z}}$  be a lattice in  $H_{\mathbb{R}}$ . Let  $H = H_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$  be its complexification.

(1) A Hodge structure of weight  $m$  on  $H$  consists of a direct sum decomposition

$$H = \bigoplus_{p+q=m} H^{p,q}, \quad \text{with } H^{q,p} = \bar{H}^{p,q},$$

where  $\bar{H}^{p,q}$  denotes the complex conjugate of  $H^{p,q}$ .

(2) A morphism of Hodge structures of type  $(r, r)$  is a linear map (defined over  $\mathbb{Q}$  relative to the lattices  $H_{\mathbb{Z}}$  and  $H'_{\mathbb{Z}}$ )

$$\varphi : H \rightarrow H', \quad \text{with } \varphi(H^{p,q}) \subset (H')^{p+r,q+r}.$$

(3) A Hodge structure  $H$  of weight  $m$  is polarized by a nondegenerate integer bilinear form  $Q$  on  $H_{\mathbb{Z}}$  if the extended bilinear form  $Q$  on  $H$  satisfies the following conditions

$$\begin{aligned} Q(v, w) &= (-1)^m Q(w, v) && \text{for all } v, w \in H, \\ Q(H^{p,q}, H^{p',q'}) &= 0 && \text{unless } p = q', q = p', \\ \sqrt{-1}^{p-q} Q(v, \bar{v}) &> 0 && \text{for } v \in H^{p,q}, v \neq 0. \end{aligned}$$

**Remark 2.20** Let  $H$  be a Hodge structure of weight  $m$  and  $H'$  be a Hodge structure of weight  $m'$ . The tensor product  $H \otimes H'$  inherits a Hodge structure of weight  $m + m'$ :

$$H \otimes H' = \sum_{p+q=m+m'} H^{p,m-p} \otimes (H')^{q,m'-q}.$$

Moreover, if  $H$  is polarized by  $Q$  and  $H'$  is polarized by  $Q'$ , then  $H \otimes H'$  is polarized by the induced bilinear form  $Q \otimes Q'$ .

A Hodge structure can also be given as follows:

**Proposition 2.21** [\[Griffiths and Schmid 1975, page 35\]](#) Let  $H, H_{\mathbb{R}}, H_{\mathbb{Z}}$  be the same as above.

(1) There is a Hodge structure of weight  $m$  on  $H$  if and only if  $H$  has a **Hodge filtration**

$$H \supset \dots \supset F^{p-1} \supset F^p \supset F^{p+1} \supset \dots \supset 0, \quad \text{with } F^p \oplus \bar{F}^{m-p+1} \xrightarrow{\cong} H \text{ for all } p.$$

(2) A map  $\varphi : H \rightarrow H'$  is a morphism of Hodge structures of type  $(r, r)$  if and only if  $\varphi$  preserves the Hodge filtration with a shift by  $r$ , i.e.,

$$\varphi(F^p) \subset (F')^{p+r} \quad \text{for all } p.$$

In particular, a morphism of Hodge structures of type  $(r, r)$  preserves the Hodge filtration strictly:

$$\varphi(F^p) = (F')^{p+r} \cap \text{Im}(\varphi) \quad \text{for all } p.$$

A mixed Hodge structure is a generalization of a Hodge structure.

**Definition 2.22** [Griffiths and Schmid 1975, Definition 1.11] Let  $H_{\mathbb{Z}}$  be a finitely generated free abelian group.

(1) A mixed Hodge structure is a triple  $(H_{\mathbb{Z}}, W_{\bullet}, F^{\bullet})$  such that:

(a) The weight filtration  $W_{\bullet}$  is

$$0 \subset \cdots \subset W_{m-1} \subset W_m \subset W_{m+1} \subset \cdots \subset H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q} = H_{\mathbb{Q}}.$$

(b) The Hodge filtration  $F^{\bullet}$  is

$$H = H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C} \supset \cdots \supset F^{p-1} \supset F^p \supset F^{p+1} \supset \cdots \supset 0.$$

(c) For each  $m \in \mathbb{Z}$ , on the graded piece  $\text{Gr}_m(W_{\bullet}) = W_m/W_{m-1}$ , the induced filtration by  $F^{\bullet}$  defines a Hodge structure of weight  $m$ .

(2) A morphism of mixed Hodge structures of type  $(r, r)$  consists of a linear map

$$\varphi : H_{\mathbb{Q}} \rightarrow (H')_{\mathbb{Q}} \quad \text{with } \varphi(W_m) \subset (W')_{m+2r} \text{ and } \varphi(F^p) \subset (F')^{p+r}.$$

The morphisms of mixed Hodge structures are also strict in the following sense.

**Lemma 2.23** [Griffiths and Schmid 1975, Lemma 1.13] A morphism of type  $(r, r)$  between mixed Hodge structures is strict with respect to both the weight and Hodge filtrations. More precisely,

$$\varphi(W_m) = (W')_{m+2r} \cap \text{Im}(\varphi), \quad \varphi(F^p) = (F')^{p+r} \cap \text{Im}(\varphi).$$

**Remark 2.24** Let  $(H_{\mathbb{Z}}, W_{\bullet}, F^{\bullet})$  and  $(\tilde{H}_{\mathbb{Z}}, \tilde{W}_{\bullet}, \tilde{F}^{\bullet})$  be two mixed Hodge structures. Their tensor product  $H \otimes \tilde{H}$  inherits a mixed Hodge structure with the weight filtration

$$0 \subset \cdots \subset \sum_{a+b \leq m-1} W_a \otimes \tilde{W}_b \subset \sum_{a+b \leq m} W_a \otimes \tilde{W}_b \subset \sum_{a+b \leq m+1} W_a \otimes \tilde{W}_b \subset \cdots \subset H_{\mathbb{Q}} \otimes \tilde{H}_{\mathbb{Q}},$$

and the Hodge filtration

$$H \otimes \tilde{H} \supset \cdots \supset \sum_{a+b \geq p-1} F^a \otimes \tilde{F}^b \supset \sum_{a+b \geq p} F^a \otimes \tilde{F}^b \supset \sum_{a+b \geq p+1} F^a \otimes \tilde{F}^b \supset \cdots \supset 0.$$

We are interested in the cohomology of quasiprojective complex varieties, which has a canonical polarizable mixed Hodge structure by the following theorem of Deligne:

**Theorem 2.25** [Deligne 1971, Theorem 3.2.5] Let  $X$  be a quasiprojective complex variety. Then  $H^{\bullet}(X; \mathbb{Q})$  carries a canonical polarizable mixed Hodge structure.

Here a polarizable mixed Hodge structure means all graded pieces  $\text{Gr}_m(W_\bullet)$  are polarizable Hodge structures. We can decompose a polarized Hodge structure into a direct sum of simple objects by the following theorem:

**Theorem 2.26** [Peters and Steenbrink 2008, Corollary 2.12] *The category of polarizable Hodge structures of weight  $m$  is semisimple.*

### 3 Stable cohomology of $\text{Mod}(\Sigma_{g,p}^b)$ with coefficients in $H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}$

In this section, we will review results of Looijenga [1996] and Kawazumi [2008] and give an explicit expression of  $H^\bullet(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r})$  for  $p + b \geq 1$ .

#### 3.1 Looijenga’s results

Let  $\mathcal{C}_{g,r}$  be the moduli space of closed genus- $g$  Riemann surfaces with  $r$  (not necessarily distinct) ordered marked points. The map  $\mathcal{C}_{g,r} \rightarrow \mathcal{M}_g$  obtained by forgetting all the marked points can also be viewed as the  $r$ -fold fiber product of the universal curve  $\pi : \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ .

Looijenga computed the rational stable cohomology of  $\mathcal{C}_{g,r}$  in a stable range, in terms of the following cohomology classes:

- (1) For  $1 \leq i \leq r$ , let

$$u_i \in H^2(\mathcal{C}_{g,r}; \mathbb{Q})$$

be the first Chern class of  $\theta_i = f_i^*(\theta)$ , where  $f_i : \mathcal{C}_{g,r} \rightarrow \mathcal{M}_{g,1}$  is the map forgetting all but the  $i$ -th marked point, and  $\theta$  is the relative tangent sheaf of  $\pi : \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ .

- (2) For a subset  $I$  of  $[r] = \{1, 2, \dots, r\}$  with  $|I| \geq 2$ , let

$$a_I \in H^{2|I|-2}(\mathcal{C}_{g,r}; \mathbb{Q})$$

be the Poincaré dual of the subvariety of  $\mathcal{C}_{g,r}$  where the marked points  $\{x_i\}_{i \in I}$  coincide:

$$x_i = x_j \quad \text{for all } i, j \in I.$$

Looijenga [1996, Lemma 2.4] showed that these cohomology classes satisfy the relations

$$(3-1) \quad u_i a_I = u_j a_I \quad \text{if } i, j \in I,$$

$$(3-2) \quad a_I a_J = u_i^{|I \cap J| - 1} a_{I \cup J} \quad \text{if } si \in I \cap J \neq \emptyset.$$

We now state Looijenga’s theorem on the rational cohomology of  $\mathcal{C}_{g,r}$ .

**Theorem 3.1** [Looijenga 1996, Theorem 2.3] *Let  $A_r^\bullet$  denote the graded commutative  $\mathbb{Q}$ -algebra generated by all  $u_i$  and  $a_I$  subject to the relations (3-1) and (3-2). There is an algebra homomorphism*

$$H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes A_r^\bullet \rightarrow H^\bullet(\mathcal{C}_{g,r}; \mathbb{Q}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq \frac{3}{2}k + 1$ .

**Remark 3.2** Looijenga provided an explicit description of  $A_r^\bullet$  as a vector space:

$$A_r^\bullet = \bigoplus_{P|[r]} \mathbb{Q}[u_I : I \in P] a_P, \quad \text{where } a_P = \prod_{I \in P, |I| \geq 2} a_I.$$

Here, the notation is as follows:

- $P|[r]$  denotes a *partition* of the set  $[r]$ , i.e.,  $P = \{I_1, I_2, \dots, I_m\}$ , where  $I_i$  are disjoint nonempty subsets of  $[r]$  and  $I_1 \cup I_2 \cup \dots \cup I_m = [r]$ . The class  $a_P = \prod_{I \in P, |I| \geq 2} a_I$  represents the Poincaré dual of the subvariety of  $\mathcal{C}_{g,r}$  where the marked points indexed by  $I_a \in P$  coincide for each  $1 \leq a \leq m$ .
- For a given partition  $P$  of  $[r]$ , we define an equivalence relation on  $\{u_1, \dots, u_r\}$ :

$$u_i \sim u_j \quad \text{if and only if } i, j \in I \text{ for some } I \in P.$$

The equivalence class of  $u_i$  for  $i \in I$  is denoted by  $u_I$ , with the convention that if  $I$  is a singleton  $\{i\}$ , then  $u_I = u_i$ .

By observing the two relations (3-1) and (3-2) that  $u_i, a_I$  satisfy, it is not hard to derive the expression of  $A_r^\bullet$  above. There is a simple mixed Hodge structure on  $A_r^\bullet$  where the degree- $2m$  part has Hodge type  $(m, m)$ . Since  $H^\bullet(\text{Mod}_g; \mathbb{Q})$  carries a canonical mixed Hodge structure (see for instance [Looijenga 1996]), this induces a mixed Hodge structure on  $H^\bullet(\mathcal{C}_{g,r}; \mathbb{Q})$ .

To study the twisted cohomology of  $\text{Mod}_g$ , Looijenga [1996, Corollary 2.7] introduced a key subspace of  $A_r^\bullet$ :

$$(3-3) \quad A_r^{\bullet} = \bigoplus_{P|[r]} \left( \prod_{\{i\} \in P} u_i^2 \right) \mathbb{Q}[u_I : I \in P] a_P.$$

By analyzing the Leray spectral sequence associated with the forgetful map  $\mathcal{C}_{g,r} \rightarrow \mathcal{M}_g$ , Looijenga embedded the twisted cohomology groups of  $\text{Mod}_g$  with coefficients  $H^1(\Sigma_g; \mathbb{Q})^{\otimes r}$  into  $H^\bullet(\mathcal{C}_{g,r}; \mathbb{Q})$ , leading to the following theorem:

**Theorem 3.3** [Looijenga 1996, Corollary 3.3] *Let  $A_r^{\bullet}$  be as defined in (3-3). There is a graded  $S_r$ -equivariant map of  $H^\bullet(\text{Mod}_g; \mathbb{Q})$ -modules which is also a morphism of mixed Hodge structures*

$$H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes A_r^{\bullet} \rightarrow H^{\bullet-r}(\text{Mod}_g; H^1(\Sigma_g; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq \frac{3}{2}k + 1$ .

Here, the symmetric group  $S_r$  acts on the left-hand side by permuting the indices in  $[r]$ , and on the right-hand side by permuting the tensor factors of  $H^1(\Sigma_g; \mathbb{Q})^{\otimes r}$ . This equivariance allows Looijenga to compute the cohomology of  $\text{Mod}_g$  with coefficients in any Schur functors of  $\text{Sp}_{2g}(\mathbb{C})$ .

### 3.2 Kawazumi’s results

Instead of studying Leray spectral sequences, Kawazumi [2008] computed the twisted cohomology of  $\text{Mod}_{g,p}^b$  with  $b \geq 1$  in a different way. He first introduced a notation called weighted partitions:

**Definition 3.4** [Kawazumi 2008, Definition 1.1] A set  $\widehat{P} = \{(S_1, i_1), (S_2, i_2), \dots, (S_\nu, i_\nu)\}$  is a *weighted partition* of the index set  $\{1, 2, \dots, r\}$  if

- (1) the set  $\{S_1, S_2, \dots, S_\nu\}$  is a partition of the set  $\{1, 2, \dots, r\}$ ,
- (2)  $i_1, i_2, \dots, i_\nu \geq 0$  are nonnegative integers,
- (3) each  $(S_a, i_a)$  for  $1 \leq a \leq \nu$  satisfies  $i_a + |S_a| \geq 2$ .

We denote by  $\mathcal{P}_r$  the set of all weighted partitions of  $\{1, 2, \dots, r\}$ .

Given a weighted partition  $\widehat{P}$ , Kawazumi [2008, equation 1.10] defined the twisted Morita–Mumford class

$$m_{\widehat{P}} \in H^{2(\sum_{a=1}^{\nu} i_a) + r - 2\nu}(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Z})^{\otimes r}),$$

and then computed the twisted cohomology of  $\text{Mod}_{g,1}$  with coefficients in  $H^1(\Sigma_g^1; \mathbb{Z})^{\otimes r}$ :

**Theorem 3.5** [Kawazumi 2008, Theorem 1.B] For degrees  $\leq N(g) - r$ ,

$$H^\bullet(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Z})^{\otimes r}) \cong \bigoplus_{\widehat{P} \in \mathcal{P}_r} H^\bullet(\text{Mod}_g^1; \mathbb{Z}) m_{\widehat{P}}.$$

Kawazumi also computed the twisted cohomology of  $\text{Mod}_{g,p}^b$  with coefficients  $H^1(\Sigma_{g,p}^b; \mathbb{Z})^{\otimes r}$  for general  $b \geq 1$ . His proof is based on the Gysin sequence (Proposition 2.12) and induction.

**Theorem 3.6** [Kawazumi 2008, Theorem 1.A] For  $b \geq 1, p \geq 0$ , we have

$$H^\bullet(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Z})^{\otimes r}) \cong H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Z}) \otimes_{H^\bullet(\text{Mod}_g^1; \mathbb{Z})} H^\bullet(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Z})^{\otimes r})$$

in degrees  $\leq N(g) - r$ .

### 3.3 Reproving the case of $\Sigma_g^1$

While Kawazumi’s theorem, Theorem 3.5, allows us to obtain the cohomology of  $\text{Mod}_g^1$  with coefficients in  $H^1(\Sigma_g^1; \mathbb{Q})^{\otimes r}$ , we provide an alternative derivation using Looijenga’s framework.

We define a subspace of  $A_r^\bullet$  from Theorem 3.1 with mixed Hodge substructures:

$$(3-4) \quad A_r^{\bullet\bullet} = \bigoplus_{P|[r]} \left( \prod_{\{i\} \in P} u_i \right) \mathbb{Q}[u_I : I \in P] a_P.$$

A reinterpretation of Theorem 3.5 using Looijenga’s setup is:

**Proposition 3.7** Let  $A_r^{\bullet\bullet}$  be as defined in (3-4). There is a graded map of  $H^\bullet(\text{Mod}_g^1; \mathbb{Q})$ -modules, which is also a morphism of mixed Hodge structures

$$H^\bullet(\text{Mod}_g^1; \mathbb{Q}) \otimes A_r^{\bullet\bullet} \rightarrow H^{\bullet-r}(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq \frac{3}{2}k + 1$ .

A nonrigid proof is identifying the twisted Morita–Mumford class  $m_{\hat{P}}$  with  $\prod_{S_a \in P} u_{S_a}^{i_a} \cdot a_P$ , where  $\hat{P} = \{(S_1, i_1), (S_2, i_2), \dots, (S_\nu, i_\nu)\}$  and  $P = \{S_1, S_2, \dots, S_\nu\}$ . It can be observed that their degrees differ by  $r$ , which is because Proposition 3.7 is stated with a degree shift by  $r$ . To rigorously prove Proposition 3.7, we first need the following lemma:

**Lemma 3.8**  $H^0(\text{Mod}_{g,1}; H^2(\Sigma_g; \mathbb{Q}))$  is isomorphic to  $\mathbb{Q}$  generated by  $a_{\{1,2\}}$ .

**Proof** Recall that  $\mathcal{C}_{g,2}(\ell)$  is the moduli space of closed genus- $g$  Riemann surfaces with two ordered marked points that are not necessarily distinct. By forgetting the first marked point in  $\mathcal{C}_{g,2}(\ell)$ , we obtain a map  $\mathcal{C}_{g,2} \rightarrow \mathcal{M}_{g,1}$ . The associated Leray spectral sequence is

$$E_2^{p,q} = H^p(\mathcal{M}_{g,1}; H^q(\Sigma_g; \mathbb{Q})) \Rightarrow H^{p+q}(\mathcal{C}_{g,2}; \mathbb{Q}).$$

We then identify  $H^0(\text{Mod}_{g,1}; H^2(\Sigma_g; \mathbb{Q}))$  with  $E_2^{0,2}$ . This Leray spectral sequence degenerates at  $E_2$  by Deligne’s theorem, Theorem 2.17, so  $E_g = E_2$ . Therefore, the map

$$H^k(\mathcal{C}_{g,2}; \mathbb{Q}) \rightarrow E_g^{k-2,2} \rightarrow E_2^{k-2,2} = H^{k-2}(\mathcal{M}_{g,1}; H^2(\Sigma_g; \mathbb{Q}))$$

is integration along fibers. Recall that  $a_{\{1,2\}}$  denotes the Poincaré dual of  $\mathcal{M}_{g,1}$  in  $\mathcal{C}_{g,2}$ , where  $\mathcal{M}_{g,1}$  embeds into  $\mathcal{C}_{g,2}$  via the trivial section,

$$\mathcal{M}_{g,1} \rightarrow \mathcal{C}_{g,2}, \quad (C, x_1) \mapsto (C, x_1, x_2 = x_1).$$

Then for any  $\omega \in H^{k-2}(\mathcal{M}_{g,1}; H^2(\Sigma_g; \mathbb{Q}))$ , letting  $\tilde{\omega}$  be its preimage in  $H^k(\mathcal{C}_{g,2}; \mathbb{Q})$ , we have

$$\int_{\mathcal{C}_{g,2}} \tilde{\omega} \wedge a_{\{1,2\}} = \int_{\mathcal{M}_{g,1}} \omega.$$

Thus  $E_2^{0,2} = H^0(\text{Mod}_{g,1}; H^2(\Sigma_g; \mathbb{Q}))$  is isomorphic to  $\mathbb{Q}$  generated by  $a_{\{1,2\}}$ . □

**Proof of Proposition 3.7** We first prove by induction on  $r$  that there is an isomorphism

$$H^\bullet(\text{Mod}_{g,1}; \mathbb{Q}) \otimes A_r^{\bullet} \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$$

in degrees  $\leq \frac{2}{3}(g-1)$ . When  $r = 0$ , the statement is vacuously true.

For  $r \geq 1$ , we suppose it is true for any  $s \leq r-1$  there is an isomorphism in degrees  $\leq \frac{2}{3}(g-1)$ :

$$H^\bullet(\text{Mod}_{g,1}; \mathbb{Q}) \otimes A_s^{\bullet} \rightarrow H^{\bullet-s}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes s}).$$

Recall that  $\mathcal{C}_{g,r+1}$  is the moduli space of closed genus- $g$  Riemann surfaces with  $(r+1)$  ordered marked points. Looijenga’s theorem, Theorem 3.1, gives us the isomorphism of mixed Hodge structures

$$H^\bullet(\mathcal{C}_{g,r+1}; \mathbb{Q}) \cong H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes A_{r+1}^\bullet$$

in degrees  $\leq \frac{2}{3}(g-1)$ .

By forgetting all but the last marked point in  $\mathcal{C}_{g,r+1}$ , we obtain a map  $f : \mathcal{C}_{g,r+1} \rightarrow \mathcal{M}_{g,1}$ , where  $\mathcal{C}_{g,r+1}$  is a complex orbifold. The associated Leray spectral sequence with  $\mathbb{Q}$ -coefficients is

$$E_2^{p,q} = H^p(\mathcal{M}_{g,1}; R^q f_*(\mathbb{Q})) \Rightarrow H^{p+q}(\mathcal{C}_{g,r+1}; \mathbb{Q}).$$

Since we are working over  $\mathbb{Q}$ , we have  $R^q f_*(\mathbb{Q}) = H^q((\Sigma_g)^{\times r}; \mathbb{Q})$ . Notice that each fiber of  $f$  is projective. Thus we can apply the orbifold version of Deligne’s theorem, [Theorem 2.17](#), which means the above spectral sequence degenerates at page 2. This implies

$$H^k(\mathcal{C}_{g,r+1}; \mathbb{Q}) \cong \bigoplus_{p+q=k} H^p(\mathcal{M}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q})).$$

The moduli space  $\mathcal{M}_{g,1}$  of closed genus- $g$  Riemann surfaces with one marked point has the same rational cohomology as  $\text{Mod}_{g,1}$ . Thus we can rewrite the above as

$$H^k(\mathcal{C}_{g,r+1}; \mathbb{Q}) \cong \bigoplus_{p+q=k} H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q})).$$

The Leray filtration preserves the mixed Hodge structure of  $H^\bullet(\mathcal{C}_{g,r+1}; \mathbb{Q})$ , therefore the  $E_2$ -page terms  $E_2^{p,q} = H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q}))$  inherit mixed Hodge structures.

We can expand the coefficients  $H^q((\Sigma_g)^{\times r}; \mathbb{Q})$  by the Künneth formula:

$$H^q((\Sigma_g)^{\times r}; \mathbb{Q}) \cong \bigoplus_{i_1+i_2+\dots+i_r=q} H^{i_1}(\Sigma_g; \mathbb{Q}) \otimes H^{i_2}(\Sigma_g; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\Sigma_g; \mathbb{Q}).$$

Identifying  $H^\bullet(\Sigma_g; \mathbb{Q})$  with  $H^\bullet(\Sigma_{g,1}; \mathbb{Q})$  as  $\text{Mod}_{g,1}$  modules, we have

$$\begin{aligned} H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q})) \\ \cong \bigoplus_{i_1+i_2+\dots+i_r=q} H^p(\text{Mod}_{g,1}; H^{i_1}(\Sigma_{g,1}; \mathbb{Q}) \otimes H^{i_2}(\Sigma_{g,1}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\Sigma_{g,1}; \mathbb{Q})). \end{aligned}$$

Observe that  $H^{k-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$  is the component of  $H^{k-r}(\text{Mod}_{g,1}; H^r((\Sigma_g)^{\times r}; \mathbb{Q}))$  with  $i_1 = i_2 = \dots = i_r = 1$ . Let’s consider the other components of  $H^{k-r}(\text{Mod}_{g,1}; H^r((\Sigma_g)^{\times r}; \mathbb{Q}))$ :

- (1) When some  $i_j = 0$ , the cohomology

$$H^p(\text{Mod}_{g,1}; H^{i_1}(\Sigma_{g,1}; \mathbb{Q}) \otimes \dots \otimes H^{i_j}(\Sigma_{g,1}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\Sigma_{g,1}; \mathbb{Q}))$$

is a component of  $H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q}))$  where  $q \leq 2r - 2$ . For each  $2 \leq i \leq r + 1$ , define  $\psi_i : \mathcal{C}_{g,r+1} \rightarrow \mathcal{C}_{g,r}$  to be the map that forgets the  $i$ -th marked point. Notice the map  $f : \mathcal{C}_{g,r+1} \rightarrow \mathcal{M}_{g,1}$  factors through  $\psi_i$ , so we have the diagram

$$\begin{array}{ccccc} (\Sigma_g)^{\times r} & \longrightarrow & \mathcal{C}_{g,r+1} & \xrightarrow{f} & \mathcal{M}_{g,1} \\ \downarrow & & \downarrow \psi_i & & \downarrow \text{id} \\ (\Sigma_g)^{\times(r-1)} & \longrightarrow & \mathcal{C}_{g,r} & \longrightarrow & \mathcal{M}_{g,1} \end{array}$$

The map  $\psi_i^* : H^\bullet(\mathcal{C}_{g,r}; \mathbb{Q}) \rightarrow H^\bullet(\mathcal{C}_{g,r+1}; \mathbb{Q})$  induces maps between items in the two Leray spectral sequences. That is,

$$H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times(r-1)}; \mathbb{Q})) \rightarrow H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q})).$$

When  $q \leq 2r - 2$ , the image of the above map is clear by the Künneth formula and induction on  $r$ .

(2) When some  $i_j = 2$ , the cup product

$$\begin{aligned} H^p(\text{Mod}_{g,1}; H^{i_1}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes \widehat{H^{i_j}}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes H^{i_r}(\Sigma_{g,1}; \mathbb{Q})) \otimes H^0(\text{Mod}_{g,1}; H^2(\Sigma_{g,1}; \mathbb{Q})) \\ \rightarrow H^p(\text{Mod}_{g,1}; H^{i_1}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes H^{i_j}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes H^{i_r}(\Sigma_{g,1}; \mathbb{Q})) \end{aligned}$$

turns out to be an isomorphism by direct computations. The term

$$H^p(\text{Mod}_{g,1}; H^{i_1}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes \widehat{H^{i_j}}(\Sigma_{g,1}; \mathbb{Q}) \otimes \cdots \otimes H^{i_r}(\Sigma_{g,1}; \mathbb{Q}))$$

is a component of  $H^p(\text{Mod}_{g,1}; H^\bullet((\Sigma_g)^{\times(r-1)}; \mathbb{Q}))$ , so it is known by induction. By Lemma 3.8, we know  $H^0(\text{Mod}_{g,1}; H^2(\Sigma_{g,1}; \mathbb{Q}))$  is  $\mathbb{Q}$  generated by  $a_{\{1,j+1\}}$ . We also make use of the relations (3-1) and (3-2) satisfied by  $u_i, a_I$  to simplify the notation (e.g., write  $a_{\{1,2\}}a_{\{1,3\}} = a_{\{1,2,3\}}$ ).

The maps in (1) and (2) are morphisms of mixed Hodge structures. All the Hodge structures involved are polarizable, hence semisimple by Theorem 2.26. Therefore, after carefully writing terms of the above two types in terms of partitions  $P$ , we can exclude them to get  $H^{k-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$  in Table 1 as follows. (For the polynomials in the table, we mean the degree  $k$  parts of them. For the  $j_1, j_2, \dots$  indices in the table, they should be distinct and between 2 and  $r + 1$ . The order listed is by increasing  $q$ . As a shorthand, we write  $K = H^\bullet(\text{Mod}_g; \mathbb{Q})$ . We always have degree  $k \leq \frac{2}{3}(g - 1)$ .)

From Table 1, we conclude that the map

$$H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes \mathbb{Q}[u_{r+1}] \otimes \left( \bigoplus_{P|[r]} \left( \prod_{\{i\} \in P} u_i \right) \mathbb{Q}[u_I : I \in P] a_P \right) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$$

is an isomorphism of mixed Hodge structures in degrees  $\leq \frac{2}{3}(g - 1)$ .

Notice that via the map  $f : \mathcal{C}_{g,r+1} \rightarrow \mathcal{C}_{g,1}$ , the image of  $H^\bullet(\mathcal{C}_{g,1}; \mathbb{Q})$  in  $H^\bullet(\mathcal{C}_{g,r+1})$  is exactly

$$H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes \mathbb{Q}[u_{r+1}].$$

Thus we can rewrite the result as

$$(3-5) \quad H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \cong H^\bullet(\text{Mod}_{g,1}; \mathbb{Q}) \otimes \left( \bigoplus_{P|[r]} \left( \prod_{\{i\} \in P} u_i \right) \mathbb{Q}[u_I : I \in P] a_P \right)$$

in degrees  $\leq \frac{2}{3}(g - 1)$ . We finish the induction.

The last step is getting the twisted cohomology of  $\text{Mod}_g^1$ . By gluing a punctured disk to  $\Sigma_g^1$ , we obtain a map

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_g^1 \rightarrow \text{Mod}_{g,1} \rightarrow 1.$$

$P [r+1]$	$H^k(\mathcal{C}_{g,r+1}; \mathbb{Q})$	$\bigoplus_{p+q=k} H^p(\text{Mod}_{g,1}; H^q((\Sigma_g)^{\times r}; \mathbb{Q}))$
$\{1\}, \dots, \{r+1\}$	$K \otimes \mathbb{Q}[u_1, u_2, \dots, u_{r+1}]$	$K \otimes \mathbb{Q}[u_{r+1}]$ $K \otimes u_{j_1} \mathbb{Q}[u_{r+1}, u_{j_1}]$ $K \otimes u_{j_1} \cdot u_{j_2} \mathbb{Q}[u_{r+1}, u_{j_1}, u_{j_2}]$ $\vdots$ $K \otimes u_{j_1} \cdots u_{j_{r-1}} \mathbb{Q}[u_{r+1}, u_{j_1}, \dots, u_{j_{r-1}}]$ $? \subset H^{k-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$
$\{r+1\}, I_2 = \{1, \dots, r\}$	$K \otimes \mathbb{Q}[u_{r+1}, u_{I_2}] a_{I_2}$	$? \subset H^{k-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$
$\{r+1\},$ $P \setminus \{r+1\}$ not as above	$K \otimes \mathbb{Q}[u_I : I \in P] a_P$	$K \otimes \mathbb{Q}[u_{r+1}, u_I : I \in P,  I  \geq 2] a_P$ $K \otimes u_{j_1} \mathbb{Q}[u_{r+1}, u_{j_1}, u_I : \{j_1\} \in P, I \in P,  I  \geq 2] a_P$ <div style="text-align: right;"><math>\text{if } \sum_{ I  \geq 2}  I  &lt; r-1</math></div> $\vdots$ $K \otimes u_{j_1} \cdots u_{j_m} \mathbb{Q}[u_{r+1}, u_{j_1}, \dots, u_{j_m}, u_I :$ <div style="text-align: right;"><math>\{j_1\}, \dots, \{j_m\}, I \in P,  I  \geq 2] a_P</math></div> <div style="text-align: right;"><math>\text{if } m + \sum_{I \in P,  I  \geq 2}  I  = r-1</math></div> $? \subset H^{k-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$
$r+1 \in I_1,  I_1  \geq 2$	$K \otimes \mathbb{Q}[u_I : I \in P] a_P$	<p>(*Take the degree <math>k-2( I_1 -1)</math> part of polynomials)</p> $K \otimes \mathbb{Q}[u_I : I \in P,  I  \geq 2] \prod_{I \in P, I \neq I_1} a_I$ $K \otimes u_{j_1} \mathbb{Q}[u_{j_1}, u_I : \{j_1\} \in P, I \in P,  I  \geq 2]$ <div style="text-align: right;"><math>\prod_{I \in P, I \neq I_1} a_I</math></div> $K \otimes u_{j_1} \cdots u_{j_n} \mathbb{Q}[u_{j_1}, \dots, u_{j_n}, u_I : \{j_1\}, \dots, \{j_n\},$ <div style="text-align: right;"><math>I \in P,  I  \geq 2] \prod_{I \in P, I \neq I_1} a_I</math></div> <div style="text-align: right;"><math>\text{if } n + \sum_{I \in P,  I  \geq 2}  I  = r+1</math></div> <p>(*The above is equivalent to the degree <math>k</math> part of the polynomial multiplied by <math>u_{I_1}^{ I_1 -1}</math>, thus equivalent to polynomials whose last term is <math>a_P</math>.)</p>

Table 1: Rational cohomology of  $\mathcal{C}_{g,r+1}$  written in two ways.

The associated Gysin sequence (Proposition 2.12) with coefficients  $H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}$  is

$$\begin{aligned} \dots \rightarrow H^{\bullet-r-2}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) &\rightarrow H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \\ &\rightarrow H^{\bullet-r}(\text{Mod}_g^1; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r-1}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \dots \end{aligned}$$

Here the map

$$H^{\bullet-r-2}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r})$$

is the multiplication by the Euler class  $e_1 \in H^2(\text{Mod}_{g,1}; \mathbb{Q})$ , so it is injective. Thus the above long exact sequence splits into short exact sequences:

$$0 \rightarrow H^{\bullet-r-2}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \xrightarrow{\cdot \ell_1} H^{\bullet-r}(\text{Mod}_{g,1}; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r}(\text{Mod}_g^1; H^1(\Sigma_{g,1}; \mathbb{Q})^{\otimes r}) \rightarrow 0.$$

The first two nonzero terms can be obtained from (3-5). Taking cokernel in the short exact sequence gives

$$H^{\bullet-r}(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Q})^{\otimes r}) \cong H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes \left( \bigoplus_{P[[r]]} \left( \prod_{\{i\} \in P} u_i \right) \mathbb{Q}[u_I : I \in P]_{a_P} \right)$$

in degrees  $\leq \frac{2}{3}(g-1)$ . □

### 3.4 Extending the results to general $\Sigma_{g,p}^b$

By combining the results of Looijenga and Kawazumi, we establish the following theorem on the twisted cohomology of  $\text{Mod}_{g,p}^b$  for  $p+b \geq 1$ :

**Theorem 3.9** *Let  $g, p, b$  be positive integers such that  $p+b \geq 1$ . The  $\mathbb{Q}$ -vector space  $A_r''^\bullet$  is defined in (3-4). There is a graded  $S_r$ -equivariant map of  $H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Q})$ -modules, which is also a morphism of mixed Hodge structures*

$$H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Q}) \otimes A_r''^\bullet \rightarrow H^{\bullet-r}(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $\leq \frac{2}{3}(g-1)$ .

**Remark 3.10** One could derive this result by repeating the method used to prove Proposition 3.7. However, that approach requires lots of repetitive computations. Instead, we present a simplified proof using Kawazumi’s results.

**Proof** Applying Kawazumi’s theorem, Theorem 3.6, over  $\mathbb{Q}$ , we obtain for  $b \geq 1$ :

$$H^\bullet(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}) \cong H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Z}) \otimes_{H^\bullet(\text{Mod}_g^1; \mathbb{Z})} H^\bullet(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Z})^{\otimes r}).$$

Using the expression of  $H^\bullet(\text{Mod}_g^1; H^1(\Sigma_g^1; \mathbb{Q})^{\otimes r})$  in Proposition 3.7, we have

$$(3-6) \quad \begin{aligned} H^{\bullet-r}(\text{Mod}_{g,p}^b; H^1(\Sigma_{g,p}^b; \mathbb{Q})^{\otimes r}) &\cong H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Q}) \otimes_{H^\bullet(\text{Mod}_g^1; \mathbb{Q})} (H^\bullet(\text{Mod}_g^1; \mathbb{Q}) \otimes A_r''^\bullet) \\ &\cong H^\bullet(\text{Mod}_{g,p}^b; \mathbb{Q}) \otimes A_r''^\bullet \end{aligned}$$

in degrees  $\leq \frac{2}{3}(g-1)$ .

The remaining case to consider is when  $b=0$  and  $p \geq 1$ . We proceed by gluing a punctured disk to the boundary component of  $\Sigma_{g,p-1}^1$ , which gives rise to the exact sequence

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_{g,p-1}^1 \rightarrow \text{Mod}_{g,p} \rightarrow 1.$$

Taking the Gysin sequence (Proposition 2.12) with coefficients  $H^1(\Sigma_{g,p-1}^1; \mathbb{Q})^{\otimes r} \cong H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}$ , we obtain

$$\begin{aligned} \dots \rightarrow H^{\bullet-r-2}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) &\rightarrow H^{\bullet-r}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \\ &\rightarrow H^{\bullet-r}(\text{Mod}_{g,p-1}^1; H^1(\Sigma_{g,p-1}^1; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r-1}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \rightarrow \dots \end{aligned}$$

Here, the map

$$H^{\bullet-r-2}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r})$$

is left multiplication by the Euler class  $e_p \in H^2(\text{Mod}_{g,p}; \mathbb{Q})$ , which is injective in degrees  $\leq \frac{2}{3}(g-1)$ . This allows us to split the above sequence into short exact sequences:

$$0 \rightarrow H^{\bullet-r-2}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \xrightarrow{\cdot e_p} H^{\bullet-r}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r}(\text{Mod}_{g,p-1}^1; H^1(\Sigma_{g,p-1}^1; \mathbb{Q})^{\otimes r}) \rightarrow 0.$$

From this, we conclude

$$H^{\bullet-r}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \cong \mathbb{Q}[e_p] \otimes H^{\bullet-r}(\text{Mod}_{g,p-1}^1; H^1(\Sigma_{g,p-1}^1; \mathbb{Q})^{\otimes r}).$$

Using equation (3-6), we obtain

$$H^{\bullet-r}(\text{Mod}_{g,p}; H^1(\Sigma_{g,p}; \mathbb{Q})^{\otimes r}) \cong \mathbb{Q}[e_p] \otimes H^{\bullet}(\text{Mod}_{g,p-1}^1; \mathbb{Q}) \otimes A_r'' \cong H^{\bullet}(\text{Mod}_{g,p}^b; \mathbb{Q}) \otimes A_r''$$

in degrees  $\leq \frac{2}{3}(g-1)$ , where the last isomorphism follows from [Theorem 2.3](#). This completes the proof. □

## 4 Moduli spaces of Riemann surfaces with level structures

In this section, we study a moduli space denoted by  $\mathcal{C}_{g,r}(\ell)$  and compute its rational cohomology for sufficiently large genus  $g$ .

### 4.1 The moduli space $\mathcal{M}_{g,r}(\ell)$

Recall that the moduli space  $\mathcal{M}_{g,r}(\ell)$  of Riemann surfaces homeomorphic to  $\Sigma_{g,r}$  with a level- $\ell$  structure is defined as

$$\mathcal{M}_{g,r}(\ell) := \text{Teich}(\Sigma_{g,r}) / \text{Mod}_{g,r}(\ell),$$

where

$$\text{Teich}(\Sigma_{g,p}) = \{\text{complex structures on } \Sigma_{g,p}\} / \text{homotopy}$$

is the Teichmüller space of  $\Sigma_{g,p}$ . When  $\ell \geq 3$ , the action of  $\text{Mod}_{g,r}(\ell)$  on  $\text{Teich}(\Sigma_{g,r})$  is free; see for example [\[Putman 2012, Section 7.1\]](#). Consequently  $\mathcal{M}_{g,r}(\ell)$  is a quasiprojective variety whose cohomology coincides with the group cohomology of  $\text{Mod}_{g,r}(\ell)$ .

For  $\ell = 2$ , the moduli space  $\mathcal{M}_{g,r}(2)$  is an orbifold, but it admits a finite branched cover (e.g.,  $\mathcal{M}_{g,r}(4)$ ) that is a quasiprojective variety.

A more explicit description of  $\mathcal{M}_{g,r}(\ell)$  is given by

$$\left\{ (C, x_1, x_2, \dots, x_r, h) \left| \begin{array}{l} C \text{ is a closed Riemann surface of genus } g, \\ x_i \text{ are distinct points on } C, \\ h : H_1(C \setminus \{x_1, \dots, x_r\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,r}; \mathbb{Z}/\ell) \text{ is an isomorphism} \end{array} \right. \right\} / \sim,$$

where

$$(C, x_1, x_2, \dots, x_r, h) \sim (C', x'_1, x'_2, \dots, x'_r, h')$$

if there exists a biholomorphism  $f : C \rightarrow C'$  such that  $f(x_i) = x'_i$  for  $1 \leq i \leq r$  and the induced map  $f_*$  on  $H_1(-; \mathbb{Z}/\ell)$  satisfies  $h' \circ f_* = h$ .

To define  $\mathcal{C}_{g,r}(\ell)$  as a moduli space containing  $\mathcal{M}_{g,r}(\ell)$ , we introduce an alternative perspective on  $\mathcal{M}_{g,r}(\ell)$ . Consider the map  $\pi^{(r+1)} : \mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  defined by

$$\pi^{(r+1)} : (C, x_1, \dots, x_{r+1}, h) \mapsto (C, x_1, \hat{h}),$$

where  $\hat{h} : H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,1}; \mathbb{Z}/\ell)$  is the map  $p_2 \circ h \circ s$  as shown in the diagram

$$\begin{array}{ccc} H_1(C \setminus \{x_1, \dots, x_{r+1}\}; \mathbb{Z}/\ell) & \xrightarrow{h} & H_1(\Sigma_{g,r+1}; \mathbb{Z}/\ell) \\ \begin{array}{c} \nearrow \\ s \downarrow \\ \searrow \end{array} p_1 & & \downarrow p_2 \\ H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) & & H_1(\Sigma_{g,1}; \mathbb{Z}/\ell) \end{array}$$

where the vertical maps  $p_1, p_2$  are obtained by forgetting all but the first marked points, while  $s$  is an arbitrary section of  $p_1$ . The definition of  $\hat{h}$  is independent of the choice of the section  $s$ , since the homology class of a loop around the  $i$ -th puncture ( $2 \leq i \leq r + 1$ ) maps to 0 in  $H_1(\Sigma_{g,1}; \mathbb{Z}/\ell)$ . It can be verified directly that  $\pi^{(r+1)} : \mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  is well-defined.

Let  $\mathcal{D} = H_1(\Sigma_{g,1}; \mathbb{Z}/\ell)$ . Denote by  $\tilde{\Sigma}_{g,1}^{(\mathcal{D})}$  the regular  $\mathcal{D}$ -cover of  $\Sigma_{g,1}$  associated with the homomorphism

$$\pi_1(\Sigma_{g,1}) \rightarrow H_1(\Sigma_{g,1}; \mathbb{Z}/\ell).$$

The fiber of the map  $\pi^{(r+1)}$  can be interpreted as follows:

**Proposition 4.1** *The fiber of the map  $\pi^{(r+1)} : \mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  is isomorphic to the orbit configuration space*

$$\text{Conf}_r^{\mathcal{D}}(\tilde{\Sigma}_{g,1}^{(\mathcal{D})}) = \{(y_2, y_3, \dots, y_{r+1}) \mid y_i \in \tilde{\Sigma}_{g,1}^{(\mathcal{D})}; y_i \neq d \cdot y_j \text{ for all } d \in \mathcal{D}, i \neq j\},$$

which is the space of  $r$  ordered points in  $\tilde{\Sigma}_{g,1}^{(\mathcal{D})}$  in different  $\mathcal{D}$ -orbits.

**Proof** We prove this by induction on  $r$ .

For  $r = 1$ , the orbit configuration space  $\text{Conf}_1^{\mathcal{D}}(\tilde{\Sigma}_{g,1}^{(\mathcal{D})})$  is  $\tilde{\Sigma}_{g,1}^{(\mathcal{D})}$ . For a given element in  $\mathcal{M}_{g,1}(\ell)$ ,

$$(C, x_1, h : H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,1}; \mathbb{Z}/\ell)),$$

its preimage under  $\pi^{(2)} : \mathcal{M}_{g,2}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  consists of elements

$$(C, x_1, x_2, \tilde{h} : H_1(C \setminus \{x_1, x_2\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,2}; \mathbb{Z}/\ell)) \in \mathcal{M}_{g,2}(\ell),$$

where  $x_2 \in C \setminus \{x_1\}$  and  $p_2 \circ \tilde{h} \circ s = h$ . For two such elements  $(C, x_1, x_2, \tilde{h})$  and  $(C, x_1, x_2, \tilde{h}')$ , they differ by

$$\tilde{h}^{-1} \circ \tilde{h}' : H_1(C \setminus \{x_1, x_2\}; \mathbb{Z}/\ell) \rightarrow H_1(C \setminus \{x_1, x_2\}; \mathbb{Z}/\ell).$$

Fix a symplectic basis  $\alpha_1, \beta_1, \dots, \alpha_g, \beta_g$  of  $H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell)$ , and let  $\delta \in H_1(C \setminus \{x_1, x_2\}; \mathbb{Z}/\ell)$  denotes the homology class of the loop around the  $x_2$ . Then the matrix corresponding to  $\tilde{h}^{-1} \circ \tilde{h}'$  under the basis  $\alpha_1, \beta_1, \dots, \alpha_g, \beta_g, \delta$  is

$$\begin{bmatrix} I_{2g \times 2g} & 0_{2g \times 1} \\ \vec{v} & 0_{1 \times 1} \end{bmatrix},$$

where  $\vec{v} \in H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \cong \mathcal{D}$ . A point  $x_2 \in C \setminus \{x_1\}$  and a deck transformation  $\vec{v} \in \mathcal{D}$  together determine a point  $y_2$  in the regular  $\mathcal{D}$ -cover of  $C \setminus \{x_1\}$ . Thus the fiber of  $\pi^{(2)}$  is homeomorphic to  $\tilde{\Sigma}_{g,1}(\mathcal{D})$ . We remark that the long exact sequence of homotopy groups associated to the fibration  $\tilde{\Sigma}_{g,1}(\mathcal{D}) \hookrightarrow \mathcal{M}_{g,2}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  when  $\ell \geq 3$  is

$$0 \rightarrow \pi_1(\tilde{\Sigma}_{g,1}(\mathcal{D})) \rightarrow \text{Mod}_{g,2}(\ell) \rightarrow \text{Mod}_{g,1}(\ell) \rightarrow 0,$$

which is exactly the mod- $\ell$  Birman exact sequence.

For  $r \geq 2$ , assume that the fiber of  $\pi^{(r)} : \mathcal{M}_{g,r}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  is

$$\text{Conf}_{r-1}^{\mathcal{D}}(\tilde{\Sigma}_{g,1}(\mathcal{D})) = \{(y_2, y_3, \dots, y_r) \mid y_i \in \tilde{\Sigma}_{g,1}(\mathcal{D}); \text{ if } i \neq j, \text{ then } y_i \neq d \cdot y_j \text{ for all } d \in \mathcal{D}\}.$$

Our goal is to find the fiber of  $\pi^{(r+1)} : \mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$ . Notice that by definition  $\pi^{(r+1)}$  is the composition map  $\pi^{(r)} \circ \tilde{\pi}$ , where  $\tilde{\pi} : \mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,r}(\ell)$  is defined in a similar way to  $\pi^{(r+1)}$  by forgetting the marked point  $x_{r+1}$  and projecting  $h : H_1(C \setminus \{x_1, \dots, x_{r+1}\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,r+1}; \mathbb{Z}/\ell)$  to  $\hat{h} : H_1(C \setminus \{x_1, \dots, x_r\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,r}; \mathbb{Z}/\ell)$ . By the same argument as  $r = 1$ , the fiber of  $\tilde{\pi}$  is the regular  $\mathcal{D}$ -cover of  $\Sigma_{g,r}$ , which we denote by  $\tilde{\Sigma}_{g,r}(\mathcal{D})$ . Therefore the fiber of the composition  $\pi^{(r)} \circ \tilde{\pi}$  is

$$\{(y_1, y_2, \dots, y_{r-1}) \in \text{Conf}_{r-1}^{\mathcal{D}}(\tilde{\Sigma}_{g,1}(\mathcal{D})), y_r \in \tilde{\Sigma}_{g,r}(\mathcal{D})\},$$

which is exactly  $\text{Conf}_r^{\mathcal{D}}(\tilde{\Sigma}_{g,1}(\mathcal{D}))$ . This completes the induction. □

Since the fiber of  $\mathcal{M}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  can be described as an orbit configuration space of  $r$  points on the regular  $\mathcal{D}$ -cover of  $\Sigma_{g,1}$ , we can express  $\mathcal{M}_{g,r+1}(\ell)$  as follows:

**Corollary 4.2** *The moduli space  $\mathcal{M}_{g,r+1}(\ell)$  can be described as*

$$\left\{ (C, x_1, y_2, \dots, y_{r+1}, h) \left| \begin{array}{l} C \text{ closed genus-}g \text{ Riemann surface, } x_1 \in C, y_2, \dots, y_{r+1} \in \tilde{C}(\mathcal{D}), \\ y_2, \dots, y_{r+1} \text{ project to distinct points in } C \setminus \{x_1\}, \\ h : H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,1}; \mathbb{Z}/\ell) \text{ is an isomorphism} \end{array} \right. \right\} / \sim,$$

where  $\tilde{C}(\mathcal{D})$  is the regular  $\mathcal{D}$ -cover of  $C$  associated with the homomorphism

$$\pi_1(C, x_1) \rightarrow H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \xrightarrow{h} H_1(\Sigma_{g,1}; \mathbb{Z}/\ell) = \mathcal{D}.$$

Two tuples  $(C, x_1, y_2, \dots, y_{r+1}, h)$  and  $(C', x'_1, y'_2, \dots, y'_{r+1}, h')$  are equivalent if there exists a biholomorphism  $f : C \rightarrow C'$  satisfying

- $f(x_1) = x'_1$ ,
- the induced map  $f_*$  on  $H_1(-; \mathbb{Z}/\ell)$  satisfies  $f_* \circ h = h'$ ,
- the unique lifting of  $f$  with  $\tilde{f}(y_1) = y'_1$  satisfies  $\tilde{f}(y_i) = y'_i$  for  $2 \leq i \leq r + 1$ .

### 4.2 The moduli space $\mathcal{C}_{g,r}(\ell)$

In Corollary 4.2, if we relax the condition on the marked points  $y_2, \dots, y_{r+1}$  and allow these points to be located anywhere in the regular  $\mathcal{D}$ -cover of  $\Sigma_g$ , we obtain a larger moduli space that contains  $\mathcal{M}_{g,r+1}(\ell)$  as an open subvariety provided  $\ell \geq 3$ . Let  $\mathcal{C}_{g,r+1}(\ell)$  denote this moduli space, defined as

$$\left\{ (C, x_1, y_2, \dots, y_{r+1}, h) \mid \begin{array}{l} C \text{ closed genus-}g \text{ Riemann surface, } x_1 \in C, y_2, \dots, y_{r+1} \in \tilde{C}_{(\mathcal{D})}, \\ h : H_1(C \setminus \{x_1\}; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,1}; \mathbb{Z}/\ell) \text{ is an isomorphism} \end{array} \right\} / \sim,$$

where the equivalence relation is the same as that in  $\mathcal{M}_{g,r+1}(\ell)$ .

**Remark 4.3** The moduli space  $\mathcal{C}_{g,r+1}(\ell)$  is strictly larger than the  $r$ -fold fiber product of  $\pi^{(2)} : \mathcal{M}_{g,2}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$ . This is because the definition of  $\mathcal{C}_{g,r+1}(\ell)$  permits the marked points  $y_2, \dots, y_{r+1}$  on  $\tilde{C}_{(\mathcal{D})}$  to project to the same point as  $x_1$ , which is not allowed in the fiber product.

To describe subvarieties of  $\mathcal{C}_{g,r}(\ell)$  with different conditions on the marked points  $y_2, \dots, y_r$  on the regular  $\mathcal{D}$ -cover of  $\Sigma_g$ , we introduce the following notion:

**Definition 4.4** Fix a group  $\mathcal{D}$ . A set  $\tilde{P} = \{(S_1, \vec{d}_1), (S_2, \vec{d}_2), \dots, (S_\nu, \vec{d}_\nu)\}$  is called a  $\mathcal{D}$ -weighted partition of the index set  $[r] = \{1, 2, \dots, r\}$ , if:

- (1) The set  $\{S_1, S_2, \dots, S_\nu\}$  is a partition of the set  $\{1, 2, \dots, r\}$ .
- (2) For each  $1 \leq a \leq \nu$ , there is an order in  $S_a = \{i_1 < i_2 < \dots < i_{|S_a|}\}$ .
- (3) For each  $1 \leq a \leq \nu$ , the element  $\vec{d}_a$  is a tuple  $(d_a^{(1)}, d_a^{(2)}, \dots, d_a^{(|S_a|-1)})$ , with  $d_a^{(i)} \in \mathcal{D}$ . By convention,  $\vec{d}_a$  is empty if  $|S_a| = 1$ .

We denote by  $\mathcal{P}_r^{\mathcal{D}}$  the set of all  $\mathcal{D}$ -weighted partitions of the index set  $\{1, 2, \dots, r\}$ . For  $\tilde{I} = (S, \vec{d}) \in \tilde{P}$ , define  $|\tilde{I}|$  to be  $|S|$ .

We then apply this notion to denote subvarieties of  $\mathcal{C}_{g,r}(\ell)$  in the following way:

**Notation 4.5** For a  $\mathcal{D}$ -weighted partition  $\tilde{P} = \{(S_1, \vec{d}_1), (S_2, \vec{d}_2), \dots, (S_\nu, \vec{d}_\nu)\}$  of the index set  $[r] = \{1, 2, \dots, r\}$ , we denote by  $\mathcal{M}(\tilde{P})$  the subvariety of  $\mathcal{C}_{g,r}(\ell)$  where the marked points  $y_1$  (lifting of  $x_1$ ),  $y_2, \dots, y_r$  satisfy:

- Points indexed by elements in the same  $S_a$  lie in the same  $\mathcal{D}$ -orbit, while points indexed by different  $S_a$ 's do not.
- For each  $1 \leq a \leq \nu$ , writing  $S_a = \{i_1^{(a)} < \dots < i_{|S_a|}^{(a)}\}$  and  $\vec{d}_a = (d_a^{(1)}, \dots, d_a^{(|S_a|-1)})$ , the points satisfy

$$y_{i_j^{(a)}} = d_a^{(j-1)} \cdot y_{i_1^{(a)}} \quad \text{for } 2 \leq j \leq |S_a|.$$

A  $\mathcal{D}$ -weighted partition encodes two pieces of information about the positions of the marked points on the regular  $\mathcal{D}$ -cover of  $\Sigma_g$ . First, it specifies whether any of the points project to the same point on  $C$ . Second, if they do, it indicates how these points differ under the  $\mathcal{D}$ -action.

By definition, each  $\mathcal{M}(\tilde{P})$  can be identified with  $\mathcal{M}_{g,|\tilde{P}|}(\ell)$ , where  $|\tilde{P}| = \nu$ . Considering all possible configurations of the marked points, we get a natural stratification of  $\mathcal{C}_{g,r}(\ell)$  as a disjoint union of subvarieties

$$(4-1) \quad \mathcal{C}_{g,r}(\ell) = \coprod_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}} \mathcal{M}_{g,|\tilde{P}|}(\ell),$$

where  $\mathcal{P}_r^{\mathfrak{D}}$  denotes the set of all  $\mathfrak{D}$ -weighted partitions of  $[r]$ .

### 4.3 Rational cohomology of $\mathcal{C}_{g,r}(\ell)$

The stratification (4-1) of  $\mathcal{C}_{g,r}(\ell)$  allows us to compute the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$ .

First, let us introduce the following cohomology classes in  $H^\bullet(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$ :

(1) For  $1 \leq i \leq r$ , let

$$v_i \in H^2(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$$

be the first Chern class of  $\theta_i = g_i^*(\theta)$ , where  $\theta$  is the relative tangent sheaf of the universal curve  $\pi : \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ , and

$$g_i : \mathcal{C}_{g,r}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell) \rightarrow \mathcal{M}_{g,1}$$

is the composition map, where the first map is

$$\mathcal{C}_{g,r}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell), \quad (C, x_1, y_2, \dots, y_r, h) \mapsto \begin{cases} (C, x_1, h) & \text{if } i = 1, \\ (C, \bar{y}_i, h) & \text{if } i \geq 2. \end{cases}$$

Here  $\bar{y}_i$  is the image of  $y_i$  under the covering map  $\tilde{C}_{(\mathfrak{D})} \rightarrow C$ . One can also view  $v_i$  as the pullback of the Euler class  $e_1 \in H^2(\mathcal{M}_{g,1}; \mathbb{Q})$  via  $g_i$ . The first Chern class of  $\theta_i$  restricted to the open subvariety  $\mathcal{M}_{g,r+1}(\ell)$  is exactly the Euler class

$$e_i \in H^2(\text{Mod}_{g,r+1}(\ell); \mathbb{Q}) \cong H^2(\text{Mod}_{g,r+1}; \mathbb{Q}).$$

(2) For a  $\mathfrak{D}$ -weighted partition  $\tilde{P}$  of the index set  $[r]$ , and  $\tilde{I} = (S, \vec{d}) \in \tilde{P}$  with  $|S| \geq 2$ , write

$$S = \{i_1 < i_2 < \dots < i_{|S|}\}, \quad \vec{d} = (d^{(1)}, d^{(2)}, \dots, d^{(|S|-1)}).$$

Let

$$a_{\tilde{I}} \in H^{2|\tilde{I}|-2}(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$$

be the Poincaré dual of the subvariety of  $\mathcal{C}_{g,r}(\ell)$  whose points  $y_1, y_2, \dots, y_r$  satisfy

$$(4-2) \quad y_{i_{j+1}} = d^{(j)} \cdot y_{i_1} \quad \text{for } 1 \leq j \leq |S| - 1.$$

Denote this subvariety by  $\mathcal{C}_{g,r}(\ell)[\tilde{I}]$ .

**Lemma 4.6** *The cohomology classes  $v_i$  and  $a_{\tilde{I}}$  satisfy the relations*

$$(4-3) \quad v_i a_{\tilde{I}} = v_j a_{\tilde{I}} \quad \text{if } i, j \in S \text{ with } \tilde{I} = (S, \vec{d}),$$

$$(4-4) \quad a_{\tilde{I}} a_{\tilde{J}} = v_i^{|\tilde{I} \cap \tilde{J}| - 1} a_{\tilde{I} \cup \tilde{J}} \quad \text{if } i \in S_0 \text{ with } \tilde{I} \cap \tilde{J} = (S_0, \vec{d}_0) \neq \emptyset,$$

$$(4-5) \quad a_{\tilde{I}} a_{\tilde{J}} = 0 \quad \text{if conditions (4-2) for } \tilde{I} \text{ and } \tilde{J} \text{ contradict.}$$

**Proof** First, for any  $\tilde{I} = (S, \vec{d}) \in \tilde{P}$ , if  $i, j \in S$ , then since  $\theta_i = g_i^*(\theta)$  and  $\theta_j = g_j^*(\theta)$  have isomorphic restrictions to  $\mathcal{C}_{g,r}(\ell)[\tilde{I}]$ , we obtain the relation (4-3).

Next, consider two elements  $\tilde{I} = (S, \vec{d})$  and  $\tilde{J} = (T, \vec{p})$ , the intersection of their corresponding subvarieties,  $\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}]$ , falls into one of the following cases:

- If  $S \cap T = \emptyset$ , the intersection  $\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}]$  is a nonempty closed subvariety of  $\mathcal{C}_{g,r+1}(\ell)$  that combines the conditions (4-2) imposed by  $\tilde{I}$  and  $\tilde{J}$ . In this case, no additional relations hold between  $a_{\tilde{I}}$  and  $a_{\tilde{J}}$ .
- If  $S \cap T \neq \emptyset$  and the conditions (4-2) imposed by  $\tilde{I}$  and  $\tilde{J}$  are compatible, we define the union of  $\tilde{I}$  and  $\tilde{J}$  as

$$\tilde{I} \cup \tilde{J} := (S \cup T, \vec{d} \cup \vec{p}).$$

In this case, we have

$$\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}] = \mathcal{C}_{g,r+1}(\ell)[\tilde{I} \cup \tilde{J}].$$

Applying [Looijenga 1996, Lemma 2.4], we obtain the relation (4-4).

- If  $S \cap T \neq \emptyset$  but the conditions (4-2) imposed by  $\tilde{I}$  and  $\tilde{J}$  are inconsistent, then

$$\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}] = \emptyset.$$

Consequently, we obtain the vanishing relation (4-5).

This completes the proof. □

**Example 4.7** Let us look at some examples of these relations.

- (1) If  $\tilde{I} = (\{1 < 2\}, d)$ , then  $1, 2 \in \tilde{I}$  and the first relation in the above lemma applies:

$$v_1 a_{\{1 < 2\}, d} = v_2 a_{\{1 < 2\}, d}.$$

- (2) If  $\tilde{I} = (\{1 < 2\}, d)$  and  $\tilde{J} = (\{1 < 3\}, d')$ , then  $\tilde{I} \cup \tilde{J} = (\{1 < 2 < 3\}, (d, d'))$ , because

$$\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}] = \mathcal{C}_{g,r+1}(\ell)[\tilde{I} \cup \tilde{J}].$$

The intersection is  $\tilde{I} \cap \tilde{J} = (\{1\})$ . The second relation above applies:

$$a_{\{1 < 2\}, d} \cdot a_{\{1 < 3\}, d'} = 1 \cdot a_{\{1 < 2 < 3\}, (d, d')}.$$

- (3) If  $\tilde{I} = (\{1 < 2\}, d)$  and  $\tilde{J} = (\{1 < 2\}, d')$  with  $d \neq d'$ , then we can see

$$\mathcal{C}_{g,r+1}(\ell)[\tilde{I}] \cap \mathcal{C}_{g,r+1}(\ell)[\tilde{J}] = \emptyset,$$

since you cannot simultaneously require that the second marked point is  $d \cdot x_1$  and  $d' \cdot x_1$ . In this case we should have  $\tilde{I} \cap \tilde{J} = \emptyset$  and we do not have the second relation above for  $a_{\tilde{I}}$  and  $a_{\tilde{J}}$ . Instead, since  $\mathcal{C}_{g,r+1}(\ell)[\tilde{I}]$  and  $\mathcal{C}_{g,r+1}(\ell)[\tilde{J}]$  are disjoint, we have

$$a_{\{1 < 2\}, d} a_{\{1 < 2\}, d'} = 0 \quad \text{if } d \neq d'.$$

Now we state the result of the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$ .

**Theorem 4.8** *Let  $v_i, a_{\tilde{\gamma}}$  be as defined above. Define  $A_r(\ell)^\bullet$  as the graded commutative  $\mathbb{Q}$ -module generated by all  $v_i$  and  $a_{\tilde{\gamma}}$ , subject to the relations (4-3), (4-4) and (4-5). Then, there exists an algebra homomorphism*

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes A_r(\ell)^\bullet \rightarrow H^\bullet(\mathcal{C}_{g,r}(\ell); \mathbb{Q}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

**Remark 4.9** To express  $A_r(\ell)^\bullet$  explicitly as a vector space, following Looijenga’s approach in Remark 3.2, we introduce the following notation:

- Given a  $\mathbb{D}$ -weighted partition  $\tilde{P}$  of the index set  $[r]$ , define

$$a_{\tilde{P}} = \prod_{\tilde{I} \in \tilde{P}, |\tilde{I}| \geq 2} a_{\tilde{I}}.$$

By relations (4-4) and (4-5), the module  $A_r(\ell)^\bullet$  can be viewed as a  $\mathbb{Q}[v_i : 1 \leq i \leq r]$ -module generated by all  $a_{\tilde{P}}$ , subject to the relation (4-3).

- Given a  $\mathbb{D}$ -weighted partition  $\tilde{P}$  of  $[r]$ , define an equivalence relation on the set  $\{v_1, \dots, v_r\}$  by

$$v_i \sim v_j \quad \text{if and only if} \quad i, j \in S \text{ for some } \tilde{I} = (S, \vec{d}) \in \tilde{P}.$$

Let  $v_{\tilde{I}}$  denote the equivalence class of  $v_i$  for any  $i \in S$  with  $\tilde{I} = (S, \vec{d}) \in \tilde{P}$ . In particular, when  $\tilde{I} = (\{i\}, \emptyset)$ , we have  $v_{\tilde{I}} = v_i$ .

Using this notation, the vector space  $A_r(\ell)^\bullet$  admits the decomposition

$$(4-6) \quad A_r(\ell)^\bullet = \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathbb{D}}} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}}.$$

**Remark 4.10** Fix the grading of the generators so that

- $v_i$  has degree 2,
- $a_{\tilde{I}}$  has degree  $2|\tilde{I}| - 2$ .

Then  $A_r(\ell)^\bullet$  decomposes as

$$A_r(\ell)^\bullet = \bigoplus_{m=0}^g A_r(\ell)^{2m},$$

where  $A_r(\ell)^{2m}$  denotes the degree  $2m$  part. This grading induces a mixed Hodge structure on  $A_r(\ell)^\bullet$ , in which  $A_r(\ell)^{2m}$  has Hodge type  $(m, m)$ .

By Putman’s theorem, Theorem 2.7, we have an isomorphism

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \cong H^\bullet(\text{Mod}_g; \mathbb{Q})$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . Consequently,  $H^\bullet(\text{Mod}_g(\ell); \mathbb{Q})$  inherits a canonical mixed Hodge structure as described in [Looijenga 1996, Section 2.5].

Moreover, by [Theorem 2.25](#), the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$  carries a canonical polarizable mixed Hodge structure. In [Theorem 4.8](#), the map

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes A_r(\ell)^\bullet \rightarrow H^\bullet(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$$

is a morphism of mixed Hodge structures.

**Proof of Theorem 4.8** We compute  $H^\bullet(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$  based on the stratification (4-1).

For integers  $k \geq 0$ , let  $U_k$  be the union of the strata  $\mathcal{M}_{g,|\tilde{P}|}$  of codimension  $\leq 2k$ , and let  $S_k$  be the union of the strata  $\mathcal{M}_{g,|\tilde{P}|}$  of codimension  $2k$ . We prove by induction on  $k$  that

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \bigoplus_{\tilde{P} \in \mathfrak{P}_r^{\oplus}, r-|\tilde{P}| \leq k} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \rightarrow H^\bullet(U_k; \mathbb{Q})$$

is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

The final case  $k = r$  is what the theorem statement is since  $U_r = \mathcal{C}_{g,r}(\ell)$ .

In the base case  $k = 0$ , the space  $U_0$  is exactly  $\mathcal{M}_{g,r}(\ell)$  with the weighted partition being  $\tilde{P} = \{\{1\}, \{2\}, \dots, \{r\}\}$ . In degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ , by [Theorem 2.7](#) we have

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \mathbb{Q}[v_1, \dots, v_r] \xrightarrow{\cong} H^\bullet(U_0; \mathbb{Q}) \cong H^\bullet(\text{Mod}_g; \mathbb{Q}) \otimes \mathbb{Q}[e_1, \dots, e_r].$$

For general  $k$ , suppose the map

$$(4-7) \quad H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \bigoplus_{\tilde{P} \in \mathfrak{P}_r^{\oplus}, r-|\tilde{P}| \leq k-1} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \rightarrow H^\bullet(U_{k-1}; \mathbb{Q})$$

is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . Notice that  $U_{k-1}$  is an open subvariety of  $U_k$ , whose complement  $S_k$  has codimension  $2k$ . The corresponding Thom–Gysin sequence ([Proposition 2.14](#)) is

$$\dots \rightarrow H^{\bullet-2k}(S_k; \mathbb{Q}) \rightarrow H^\bullet(U_k; \mathbb{Q}) \rightarrow H^\bullet(U_{k-1}; \mathbb{Q}) \rightarrow H^{\bullet+1}(U_k; \mathbb{Q}) \rightarrow \dots$$

Here the map  $H^\bullet(U_k; \mathbb{Q}) \rightarrow H^\bullet(U_{k-1}; \mathbb{Q})$  is surjective since the map (4-7) factorizes over  $H^\bullet(U_k; \mathbb{Q})$ . Thus we have a splitting short exact sequence

$$0 \rightarrow H^{\bullet-2k}(S_k; \mathbb{Q}) \rightarrow H^\bullet(U_k; \mathbb{Q}) \rightarrow H^\bullet(U_{k-1}; \mathbb{Q}) \rightarrow 0.$$

Recall

$$S_k = \coprod_{\tilde{P} \in \mathfrak{P}_{r+1}^{\oplus}, |\tilde{P}|=r-k} \mathcal{M}_{g,|\tilde{P}|}(\ell).$$

We know  $H^\bullet(\mathcal{M}_{g,|\tilde{P}|}(\ell); \mathbb{Q})$  from [Theorem 2.7](#), and the Gysin map  $H^{\bullet-2k}(S_k; \mathbb{Q}) \rightarrow H^\bullet(U_k; \mathbb{Q})$  restricted to the component  $H^\bullet(\mathcal{M}_{g,|\tilde{P}|}(\ell); \mathbb{Q})$  is the multiplication by  $a_{\tilde{P}} \in H^{2k}(U_k; \mathbb{Q})$ . Thus we conclude

that in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$  we have

$$\begin{aligned}
 H^\bullet(U_k; \mathbb{Q}) &\cong H^\bullet(U_{k-1}; \mathbb{Q}) \oplus H^{\bullet-2k}(S_k; \mathbb{Q}) \\
 &\cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}, r-|\tilde{P}| \leq k-1} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \oplus H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \\
 &\qquad \qquad \qquad \otimes \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}, r-|\tilde{P}|=k} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \\
 &\cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}, r-|\tilde{P}| \leq k} \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}}.
 \end{aligned}$$

This finishes the induction. □

### 5 Twisted cohomology of the level- $l$ mapping class groups

In this section, we prove [Theorem B](#) by embedding the twisted cohomology of the level- $l$  mapping class group, with coefficients in the  $r$ -tensor power of Prym representation, into  $H^\bullet(\mathcal{C}_{g,r}(\ell); \mathbb{Q})$ .

From [Theorem 4.8](#), we recall that the rational cohomology of  $\mathcal{C}_{g,r}(\ell)$  is described using an algebra  $A_r(\ell)^\bullet$ , explicitly given by (4-6). To proceed, we define a  $\mathbb{Q}$ -subspace of  $A_r(\ell)^\bullet$  as

$$(5-1) \quad A'_r(\ell)^\bullet = \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}},$$

where  $\mathcal{P}_r^{\mathfrak{D}}$  is the set of all  $\mathfrak{D}$ -weighted partition of the index set  $[r]$ ; see [Definition 4.4](#).

The symmetric group  $S_r$  naturally acts on  $A'_r(\ell)^\bullet$  by permuting the indices of  $[r]$ . Furthermore,  $A'_r(\ell)^\bullet$  inherits a mixed Hodge substructure from the mixed Hodge structure of  $A_r(\ell)^\bullet$  in [Remark 4.10](#).

#### 5.1 Case of $\text{Mod}_{g,1}(\ell)$

We first analyze the case of  $\text{Mod}_{g,1}(\ell)$ .

**Theorem 5.1** *There is a graded  $S_r$ -equivariant map of  $H^\bullet(\text{Mod}_{g,1}(\ell); \mathbb{Q})$ -modules which is also a morphism of mixed Hodge structures*

$$H^\bullet(\text{Mod}_{g,1}(\ell); \mathbb{Q}) \otimes A'_r(\ell)^\bullet \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

Recall that  $\mathfrak{D} = H_1(\Sigma_g; \mathbb{Z}/\ell)$  and  $\mathfrak{H}_{g,1}(\ell; \mathbb{Q}) = H^1(\tilde{\Sigma}_{g,1}^{(\mathfrak{D})}; \mathbb{Q})$ , where  $\tilde{\Sigma}_{g,1}^{(\mathfrak{D})}$  denotes the regular  $\mathfrak{D}$ -cover of  $\Sigma_{g,1}$ . Let  $\tilde{\Sigma}_g^{(\mathfrak{D})}$  denote the regular  $\mathfrak{D}$ -cover of  $\Sigma_g$ .

To compute  $H^\bullet(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r})$ , we also need to consider twisted cohomology with coefficients in

$$H^1(\tilde{\Sigma}_g^{(\mathfrak{D})}; \mathbb{Q})^{\otimes r} \quad \text{and} \quad H^1(\tilde{\Sigma}_g^{(\mathfrak{D})}; \mathbb{Q})^{\otimes r-1} \otimes \mathfrak{H}_{g,1}(\ell; \mathbb{Q}),$$

and similar mixed tensor products. To handle these cases systematically, we introduce the following indexing notation:

**Definition 5.2** Set  $f(1) = \mathfrak{H}_{g,1}(\ell; \mathbb{Q})$  and  $f(0) = H^1(\tilde{\Sigma}_{g(\mathfrak{D})}; \mathbb{Q})$ . For any  $r \geq 1$ , let  $J = (J_1, J_2, \dots, J_r)$  be a sequence where  $J_i \in \{0, 1\}$  for all  $i$ . Then, we define the corresponding  $r$ -tensor product

$$\mathfrak{H}^r(J) := f(J_1) \otimes f(J_2) \otimes \dots \otimes f(J_r).$$

This notation allows us to track different tensor products involving  $H^1(\tilde{\Sigma}_{g(\mathfrak{D})}; \mathbb{Q})$  and  $\mathfrak{H}_{g,1}(\ell; \mathbb{Q})$  in a structured manner.

To describe the cohomology of  $\text{Mod}_{g,1}(\ell)$  with coefficients in  $\mathfrak{H}^r(J)$ , we define when a  $\mathfrak{D}$ -weighted partition is compatible with  $J$ :

**Definition 5.3** Given  $\mathfrak{D}, r, J, \mathfrak{H}^r(J)$  as above, a  $\mathfrak{D}$ -weighted partition

$$\tilde{P} = \{(S_1, \vec{d}_1), (S_2, \vec{d}_2), \dots, (S_\nu, \vec{d}_\nu)\}$$

indexed by  $[r + 1]$  is said to be *compatible with  $J$*  if:

- (1) By convention, we assume  $1 \in S_1$ .
- (2) For  $\vec{d}_1 = (d_1^{(1)}, d_1^{(2)}, \dots, d_1^{(|S_1|-1)})$ , each  $d_1^{(i)}$  ( $1 \leq i \leq |S_1| - 1$ ) is not the unit 1 in  $\mathfrak{D}$ .
- (3)  $S_1$  does not contain any  $2 \leq a \leq r + 1$  such that  $J_{a-1} = 1$ .

Let  $\mathcal{P}_{r+1}^{\mathfrak{D}}(J)$  denote the set of all  $\mathfrak{D}$ -weighted partitions indexed by  $[r + 1]$  which are compatible with  $J$ .

Using this notation, the twisted cohomology  $H^\bullet(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J))$  is described as follows:

**Theorem 5.4** *There is a graded map of  $H^\bullet(\text{Mod}_g(\ell); \mathbb{Q})$ -modules*

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_{r+1}^{\mathfrak{D}}(J)} \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J)),$$

which is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

**Remark 5.5** *Theorem 5.1* follows as a special case of *Theorem 5.4* when  $J = (1, 1, \dots, 1)$ . By *Definition 5.3*, a  $\mathfrak{D}$ -weighted partition indexed by  $[r + 1]$  that is compatible with  $(1, 1, \dots, 1)$  takes the form

$$\tilde{P} = \{(S_1, \vec{d}_1) = (\{1\}, \emptyset), (S_2, \vec{d}_2), \dots, (S_\nu, \vec{d}_\nu)\}.$$

This is in one-to-one correspondence with a  $\mathfrak{D}$ -weighted partition indexed by  $[r]$ :

$$\tilde{P}' = \{(S_2, \vec{d}_2), \dots, (S_\nu, \vec{d}_\nu)\}.$$

Thus, when  $J = (1, 1, \dots, 1)$ , we obtain

$$\begin{aligned} H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_{r+1}^{\mathfrak{D}}(J)} \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right) \\ \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P}' \in \mathcal{P}_r^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}'} v_i \right) \mathbb{Q}[v_1, v_{\tilde{I}} : \tilde{I} \in \tilde{P}'] a_{\tilde{P}'} \right) \\ \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \mathbb{Q}[v_1] \otimes \left( \bigoplus_{\tilde{P}' \in \mathcal{P}_r^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}'} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}'] a_{\tilde{P}'} \right) \\ \cong H^\bullet(\text{Mod}_{g,1}(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P}' \in \mathcal{P}_r^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}'} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}'] a_{\tilde{P}'} \right). \end{aligned}$$

Now it suffices to prove [Theorem 5.4](#). We begin with proving the case  $J = (0, 0, \dots, 0)$ , where the coefficients  $\mathfrak{H}^r(J) = H^1(\tilde{\Sigma}_{g(\mathfrak{D})}; \mathbb{Q})^{\otimes r}$ :

**Proof of [Theorem 5.4](#) when  $J = (0, 0, \dots, 0)$**  We proceed by induction on  $r$ . For  $r = 0$ , the statement is

$$H^\bullet(\text{Mod}_{g,1}(\ell); \mathbb{Q}) \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \mathbb{Q}[v_1],$$

which follows from Putman’s theorem ([Theorem 2.7](#)) and Looijenga’s theorem ([Theorem 2.3](#)).

Let  $r \geq 1$ . In the previous section, we studied the moduli space  $\mathcal{C}_{g,r+1}(\ell)$  and computed its stable rational cohomology. Define the map

$$\pi : \mathcal{C}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell), \quad (C, x_1, y_2, \dots, y_{r+1}, h) \mapsto (C, x_1, h),$$

which is a holomorphic map of quasiprojective varieties when  $\ell \geq 3$  and a map of orbifolds when  $\ell = 2$ .

The fiber of  $\pi$  consists of  $r$  ordered (not necessarily distinct) points in the regular  $\mathfrak{D}$ -cover of the closed genus- $g$  Riemann surface  $C$ , so the fiber is homeomorphic to  $(\tilde{\Sigma}_{g(\mathfrak{D})})^{\times r}$ . Since the fiber is projective, we can apply Deligne’s degeneration theorem ([Theorem 2.17](#)) to the map  $f$ , implying that the associated Leray spectral sequence degenerates at the second page:

$$E_2^{p,q} = H^p(\mathcal{M}_{g,1}(\ell); H^q((\tilde{\Sigma}_{g(\mathfrak{D})})^{\times r}; \mathbb{Q})) \Rightarrow H^{p+q}(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q}).$$

Thus,

$$H^k(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q}) \cong \bigoplus_{p+q=k} H^p(\mathcal{M}_{g,1}(\ell); H^q((\tilde{\Sigma}_{g(\mathfrak{D})})^{\times r}; \mathbb{Q})).$$

Since the Leray filtration respects the mixed Hodge structure of  $H^*(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q})$ , terms on the second page inherit mixed Hodge structures.

On the one hand, by [Theorem 4.8](#), the rational cohomology of  $\mathcal{C}_{g,r+1}(\ell)$  is given by

$$H^\bullet(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q}) \cong \bigoplus_{\tilde{P} \in \mathcal{P}_{r+1}^{\mathfrak{D}}} H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}}$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . This is also an isomorphism of mixed Hodge structures, as discussed in [Remark 4.10](#).

On the other hand, we can expand the terms on the second page of the Leray spectral sequence using the Künneth formula:

$$H^p(\text{Mod}_{g,1}(\ell); H^q((\tilde{\Sigma}_g^{\text{g}})^{\times r}; \mathbb{Q})) \cong \bigoplus_{i_1+i_2+\dots+i_r=q} H^p(\text{Mod}_{g,1}(\ell); H^{i_1}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes H^{i_2}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q})).$$

Our target  $H^{k-r}(\text{Mod}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q})^{\otimes r})$  is the component where  $i_1 = \dots = i_r = 1$ . Other components can be determined as follows:

(1) If some  $i_j = 0$ , the component

$$H^p(\text{Mod}_{g,1}(\ell); H^{i_1}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_j}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}))$$

appears in  $H^p(\text{Mod}_{g,1}; H^q((\tilde{\Sigma}_g^{\text{g}})^{\times r}; \mathbb{Q}))$ , where  $q \leq 2r - 2$ . For  $2 \leq i \leq r + 1$ , let the map  $\pi^{(i)} : \mathcal{C}_{g,r+1}(\ell) \rightarrow \mathcal{C}_{g,r}(\ell)$  be defined by

$$(C, x_1, y_2, \dots, y_{r+1}, h) \mapsto (C, x_1, y_2, \dots, \hat{y}_i, \dots, y_{r+1}, h).$$

By definition, the map  $\pi : \mathcal{C}_{g,r+1}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$  factors through  $\pi^{(i)}$ . We then have the commuting diagram

$$\begin{array}{ccccc} (\tilde{\Sigma}_g^{\text{g}})^{\times r} & \longrightarrow & \mathcal{C}_{g,r+1}(\ell) & \longrightarrow & \mathcal{M}_{g,1}(\ell) \\ \downarrow & & \downarrow \pi^{(i)} & & \downarrow \text{id} \\ (\tilde{\Sigma}_g^{\text{g}})^{\times(r-1)} & \longrightarrow & \mathcal{C}_{g,r}(\ell) & \longrightarrow & \mathcal{M}_{g,1}(\ell) \end{array}$$

which induces maps between the  $E_2$  terms of the two Leray spectral sequences:

$$H^p(\mathcal{M}_{g,1}(\ell); H^q((\tilde{\Sigma}_g^{\text{g}})^{\times(r-1)}; \mathbb{Q})) \rightarrow H^p(\mathcal{M}_{g,1}(\ell); H^q((\tilde{\Sigma}_g^{\text{g}})^{\times r}; \mathbb{Q})) \quad \text{for } q \leq 2r - 2.$$

After we expand the left-hand side by the Künneth formula, these terms can be identified by induction.

(2) If some  $i_j = 2$ , such a component appears as the image of the cup product

$$\begin{aligned} &H^p(\text{Mod}_{g,1}(\ell); H^{i_1}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{\hat{i}_j}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q})) \\ &\quad \otimes H^0(\text{Mod}_{g,1}(\ell); H^2(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q})) \\ &\rightarrow H^p(\text{Mod}_{g,1}(\ell); H^{i_1}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_j}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q})). \end{aligned}$$

Here the term

$$H^p(\text{Mod}_{g,1}(\ell); H^{i_1}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{\hat{i}_j}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}) \otimes \dots \otimes H^{i_r}(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}))$$

is known by induction. Additionally, by following essentially the same argument as in [Lemma 3.8](#), but applied to the Leray spectral sequence associated with the map  $\mathcal{C}_{g,2}(\ell) \rightarrow \mathcal{M}_{g,1}(\ell)$ , we deduce that  $H^0(\text{Mod}_{g,1}(\ell); H^2(\tilde{\Sigma}_g^{\text{g}}; \mathbb{Q}))$  is isomorphic to  $\mathbb{Q}$  generated by  $a_{(\{1 < j+1\}, 1 \in \mathbb{g})}$ . Using this, we can compute the image and simplify the result by employing the relations (4-3)–(4-5) satisfied by  $v_i$  and  $a_{\tilde{\gamma}}$ .

$\tilde{P} \in \mathcal{P}_{r+1}^{\otimes}$	$H^k(\mathcal{C}_{g,r+1}(\ell); \mathbb{Q})$	$\bigoplus_{p+q=k} H^p(\mathcal{M}_{g,1}(\ell); H^q(\tilde{\Sigma}_g^{\otimes} \times \tilde{\Sigma}_g^{\otimes}; \mathbb{Q}))$
$\{1\}, \{2\}, \dots, \{r+1\}$	$K \otimes \mathbb{Q}[v_1, v_2, \dots, v_{r+1}]$	$K \otimes \mathbb{Q}[u_1]$ $K \otimes v_a \mathbb{Q}[v_1, v_a]$ for all $a \geq 2$ $K \otimes v_b v_c \mathbb{Q}[v_1, v_b, v_c], 2 \leq b < c$ $\vdots$ $v_{a_1} v_{a_2} \cdots v_{a_{r-1}} [v_1, v_{a_1}, v_{a_2}, \dots, v_{a_{r-1}}],$ $2 \leq a_1 < a_2 < \cdots < a_{r-1}$ $? \subset H^{k-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r})$
$\{1\}, I_2 = (S_2, \vec{d}_2),$ $\dots, I_v = (S_v, \vec{d}_v),$ $ S_j  \geq 2$ for all $j \geq 2$	$K \otimes \mathbb{Q}[v_1, v_{I_j} : j \geq 2] a \tilde{p}$	$? \subset H^{k-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r})$
$\{1\}, S_2 = \{s_2\},$ $\dots, S_{m-1} = \{s_{m-1}\},$ $I_m = (S_m, \vec{d}_m),$ $\dots, I_v = (S_v, \vec{d}_v),$ $m > 2,  S_j  \geq 2$ for all $j \geq m$	$K \otimes \mathbb{Q}[v_1, v_{s_2}, \dots, v_{s_m},$ $v_{I_j} : j \geq m] a \tilde{p}$	$K \otimes \mathbb{Q}[v_1, v_{I_j} : j \geq m] a \tilde{p}$ $K \otimes v_a \mathbb{Q}[v_1, v_{s_a}, v_{I_j} : j \geq m] a \tilde{p}, 2 \leq a \leq m-1, m > 3$ $\vdots$ $v_{s_{a_1}} \cdots v_{s_{a_{m-3}}} \mathbb{Q}[v_1, v_{s_{a_1}}, \dots, v_{s_{a_{m-3}}}, v_{I_j} : j \geq m] a \tilde{p},$ $2 \leq a_1 < \cdots < a_{m-3} \leq m-1$ $? \subset H^{k-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r})$
$(S_1, \vec{d}_1), \dots, (S_v, \vec{d}_v),$ $1 \in S_1,  S_1  \geq 2$	$K \otimes \mathbb{Q}[u_{I_j} : 1 \leq j \leq v] a \tilde{p}$	For $\vec{d}_1 = (d_1^{(1)}, d_1^{(2)}, \dots, d_1^{( S_1 -1)})$ , if there exists an $i$ such that $d_1^{(i)} = 1$ , then all can be realized outside $H^{k-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r})$ ; if $d_1^{(i)} \neq 1$ for all $i$ , then all except $(\prod_{\{i\} \in \tilde{P}, i \neq 1} v_i) \mathbb{Q}[v_I : I \in \tilde{P}] a \tilde{p}$ can be realized outside $H^{k-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r})$ .

Table 2: Rational cohomology of  $\mathcal{C}_{g,r+1}(\ell)$  written in two ways.

We carefully list components of these two types in Table 2, where polynomials in the table refer to the degree  $k$  terms and  $K$  denotes  $H^\bullet(\text{Mod}_g(\ell); \mathbb{Q})$ . Since all isomorphisms in the table are isomorphisms of (polarized) mixed Hodge structures, by semisimplicity, we have, when  $g \geq 2k^2 + 7k + 2$ ,

$$H^{\bullet-r}(\mathcal{M}_{g,1}(\ell); H^1(\tilde{\Sigma}_g^{\otimes}; \mathbb{Q})^{\otimes r}) \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_{r+1}^{\otimes}(J)} \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a \tilde{p} \right)$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . Here  $J = (0, 0, \dots, 0)$ . □

**Proof of Theorem 5.4 for general  $J$**  Let  $J = (J_1, J_2, \dots, J_r)$ . We have just proved the case where  $J = (0, 0, \dots, 0)$ . We now prove the theorem for a general  $J$  by induction on  $r$  and on  $\sum_{i=1}^r J_i$ .

When  $r = 0$ , the theorem holds by the same reasoning as in the beginning of the previous proof.

For  $r \geq 1$ , assume by induction that the theorem holds for cases  $\leq (r - 1)$ . When  $\sum_{i=1}^r J_i = 0$ , the proof has already been provided. Now, let  $\sum_{i=1}^r J_i = m > 0$ , and assume by induction that the theorem holds for  $\sum_{i=1}^r J_i = m - 1$ .

For  $J = (J_1, J_2, \dots, J_r)$ , since  $\sum_{i=1}^r J_i > 0$ , there exists at least one  $J_t$  that is 1. Replacing the  $t$ -th entry in  $J$  by 0, we obtain a new array  $\tilde{J}$  with  $\sum_{i=1}^r \tilde{J}_i = m - 1$ . We can then write

$$\begin{aligned} \mathfrak{H}^r(J) &= f(J_1) \otimes \dots \otimes f(J_t = 1) \otimes \dots \otimes f(J_r), \\ \mathfrak{H}^r(\tilde{J}) &= f(J_1) \otimes \dots \otimes f(\tilde{J}_t = 0) \otimes \dots \otimes f(J_r). \end{aligned}$$

Here  $f(0) = H^1(\tilde{\Sigma}_{g,1}^{\mathfrak{D}(\mathfrak{Q})}; \mathbb{Q})$  and  $f(1) = \mathfrak{H}_{g,1}(\ell; \mathbb{Q})$ . Recall that the Prym representation  $\mathfrak{H}_{g,1}(\ell; \mathbb{Q})$  is  $H^1(\tilde{\Sigma}_{g,1}^{\mathfrak{D}(\mathfrak{Q})}; \mathbb{Q})$ , where  $\tilde{\Sigma}_{g,1}^{\mathfrak{D}(\mathfrak{Q})}$  is the regular  $\mathfrak{D}$ -cover of  $\Sigma_{g,1}$ . By filling in all punctures of  $\tilde{\Sigma}_{g,1}^{\mathfrak{D}(\mathfrak{Q})}$ , we obtain the following short exact sequence of  $\text{Mod}_{g,1}(\ell)$ -modules:

$$0 \rightarrow H^1(\tilde{\Sigma}_{g,1}^{\mathfrak{D}(\mathfrak{Q})}; \mathbb{Q}) \rightarrow \mathfrak{H}_{g,1}(\ell; \mathbb{Q}) \rightarrow \mathbb{Q}^{|\mathfrak{Q}|-1} \rightarrow 0.$$

Tensoring this short exact sequence with  $f(J_1) \otimes \dots \otimes f(J_{t-1})$  on the left and with  $f(J_{t+1}) \otimes \dots \otimes f(J_r)$  on the right, we obtain

$$0 \rightarrow \mathfrak{H}^r(\tilde{J}) \rightarrow \mathfrak{H}^r(J) \rightarrow f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes \mathbb{Q}^{|\mathfrak{Q}|-1} \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r) \rightarrow 0.$$

This short exact sequence of  $\text{Mod}_{g,1}(\ell)$ -modules induces the following long exact sequence:

$$\begin{aligned} \dots \rightarrow H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes \mathbb{Q}^{|\mathfrak{Q}|-1} \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r)) \\ \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(\tilde{J})) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J)) \\ \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes \mathbb{Q}^{|\mathfrak{Q}|-1} \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r)) \rightarrow \dots \end{aligned}$$

Our goal is to compute  $H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J))$ . Let  $\phi_\bullet$  denote the map

$$(5-2) \quad H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes \mathbb{Q}^{|\mathfrak{Q}|-1} \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r)) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(\tilde{J})).$$

Then we have

$$(5-3) \quad 0 \rightarrow \text{Coker}(\phi_\bullet) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J)) \rightarrow \text{Ker}(\phi_{\bullet+1}).$$

On the one hand, the source term of the map (5-2) is

$$\begin{aligned} H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes \mathbb{Q}^{|\mathfrak{Q}|-1} \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r)) \\ \cong \bigoplus_{|\mathfrak{Q}|-1} H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r)). \end{aligned}$$

Since the coefficients are  $(r-1)$ -tensor powers, this cohomology is known by the induction hypothesis on  $r$ . For each  $1 \neq d \in \mathfrak{D}$ , the component

$$(5-4) \quad H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); f(J_1) \otimes \dots \otimes f(J_{t-1}) \otimes f(J_{t+1}) \otimes \dots \otimes f(J_r))$$

is isomorphic to the direct sum of

$$\left( H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right)^{\bullet-2},$$

where  $\tilde{P}$  ranges over all  $\mathcal{D}$ -weighted partitions of the index set  $\{1, \dots, t, \widehat{t+1}, t+2, \dots, r+1\}$  that are compatible with  $(J_1, \dots, J_{t-1}, J_{t+1}, \dots, J_r)$ . Here  $[2]$  indicates a degree shift by 2. Inserting  $(t+1)$  into the index set, we can rewrite this as a direct sum of

$$(5-5) \quad \left( H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{I_1 \setminus \{t+1\}} \prod_{m \geq 2} a_{I_m} \right)^{\bullet-2},$$

where  $\tilde{P} = \{I_1 = (S_1, \vec{d}_1), \dots, I_\nu = (S_\nu, \vec{d}_\nu)\}$  ranges over all  $\mathcal{D}$ -weighted partitions of the index set  $[r+1]$  such that:

- $1 \in S_1$  and  $t+1 \in S_1$ .
- Each entry in  $\vec{d}_1$  is not  $1 \in \mathcal{D}$ . In particular, the entry corresponding to  $t+1$  is  $d$ .
- $S_1$  does not contain any  $2 \leq a \leq r+1$  such that  $J_{a-1} = 1$ .

On the other hand, since  $\sum_{i=1}^{r+1} \tilde{J}_i = m-1 < m$ , the target term  $H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(\tilde{J}))$  of the map (5-2) is known by the induction hypothesis on  $\sum_{i=1}^{r+1} \tilde{J}_i$ . We have

$$H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(\tilde{J})) \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{D}_{r+1}^{\mathcal{D}}(\tilde{J})} \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right)$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

The map (5-2) restricted to the component (5-4) indexed by  $1 \neq d \in \mathcal{D}$  is multiplication by the Poincaré dual  $a_{(\{1 < t+1\}, d)}$ . Hence, the image of the summand (5-5) is

$$(5-6) \quad H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{I_1 \setminus \{t+1\}} \prod_{m \geq 2} a_{I_m} \cdot a_{(\{1 < t+1\}, d)}.$$

The relation (4-4) for  $I_1 = (S_1, \vec{d}_1) = (\{1 < t+1\}, d) \cup (I_1 \setminus \{t+1\})$  tells us

$$a_{(\{1 < t+1\}, d)} \cdot a_{I_1 \setminus \{t+1\}} = a_{I_1}.$$

Thus (5-6) can be rewritten as

$$\begin{aligned} H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{I_1 \setminus \{t+1\}} \prod_{m \geq 2} a_{I_m} \cdot a_{(\{1 < t+1\}, d)} \\ = H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{I_1} \prod_{m \geq 2} a_{I_m} \\ = H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}}. \end{aligned}$$

Thus the image of the map (5-2) is the direct sum of (5-6) as  $\tilde{P}$  ranges over  $\mathcal{D}$ -weighted partitions of the index set  $[r+1]$  that satisfy the conditions specified above. By excluding these particular  $\mathcal{D}$ -weighted

partitions from  $\mathcal{P}_{r+1}^{\mathfrak{D}}(\tilde{J})$ , the cokernel of the map (5-2) is the direct sum of

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}},$$

as  $\tilde{P}$  ranges over  $\mathfrak{D}$ -weighted partitions of the index set  $[r + 1]$  that are compatible with  $\tilde{J}$  and  $S_1$  does not contain  $t + 1$ . According to the definition of compatibility (Definition 5.3), these are precisely the  $\mathfrak{D}$ -weighted partitions of the index set  $[r + 1]$  that are compatible with  $J$ , since  $J_t = 1$ .

Since the map (5-2) is injective, the kernel is zero. Thus by (5-3) we have

$$H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}^r(J)) \cong H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_{r+1}^{\mathfrak{D}}(J)} \left( \prod_{\{i\} \in \tilde{P}, i \neq 1} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right).$$

This completes the induction. □

Our next goal is to generalize the result from  $\Sigma_{g,1}$  to any nonclosed surface  $\Sigma_{g,p}^b$ . In order to do that, we first need the following result for  $\Sigma_g^1$ :

**Corollary 5.6** *There is a graded map of  $H^\bullet(\text{Mod}_g^1(\ell); \mathbb{Q})$ -modules*

$$H^\bullet(\text{Mod}_g^1(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_g^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right) \rightarrow H^{\bullet-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{Q})^{\otimes r}),$$

which is an isomorphism in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ .

**Proof** By Proposition 2.5, we have the short exact sequence

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_g^1(\ell) \rightarrow \text{Mod}_{g,1}(\ell) \rightarrow 1$$

obtained by gluing a punctured disk to the boundary of  $\Sigma_g^1$ . This sequence induces a Gysin sequence (Proposition 2.12) for cohomology with coefficients in  $\mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r} \cong \mathfrak{H}_g^1(\ell; \mathbb{Q})^{\otimes r}$ :

$$\begin{aligned} \dots \rightarrow H^{\bullet-r-2}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}) &\rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}) \\ &\rightarrow H^{\bullet-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r-1}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}) \\ &\rightarrow H^{\bullet-r+1}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}) \rightarrow \dots \end{aligned}$$

Let  $\phi_\bullet$  denote the boundary map

$$\phi_\bullet : H^{\bullet-r-2}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r}(\text{Mod}_{g,1}(\ell); \mathfrak{H}_{g,1}(\ell; \mathbb{Q})^{\otimes r}).$$

We then have the short exact sequence

$$1 \rightarrow \text{Coker}(\phi_\bullet) \rightarrow H^{\bullet-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{Q})^{\otimes r}) \rightarrow \text{Ker}(\phi_{\bullet+1}) \rightarrow 1.$$

Since  $\phi_\bullet$  is multiplication by the Euler class  $e_1$ , it is injective. The target term of  $\phi_\bullet$  is known from [Theorem 5.1](#), so we can determine the cokernel of  $\phi_\bullet$ :

$$H^\bullet(\text{Mod}_g(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathfrak{D}}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right)$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . This is isomorphic to  $H^{\bullet-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{Q})^{\otimes r})$ . □

To generalize our results to any nonclosed surface  $\Sigma_{g,p}^b$ , we introduce Putman’s theory [\[2022\]](#) on partial level- $\ell$  representations as follows.

### 5.2 Putman’s theory on partial level- $\ell$ representations

Let

$$\mathfrak{H}_{g,p}^b(\ell; \mathbb{C}) = \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C}.$$

The finite group  $\mathfrak{D} \cong (\mathbb{Z}/\ell)^{2g}$  acts on  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{C})$  via deck transformations. Irreducible representations of  $\mathfrak{D}$  are characterized by characters. A character  $\chi : \mathfrak{D} \rightarrow \mathbb{C} \setminus \{0\}$  gives an irreducible representation  $\mathbb{C}_\chi$ , where  $d \cdot \vec{v} = \chi(d)\vec{v}$  for  $d \in \mathfrak{D}$  and  $\vec{v} \in \mathbb{C}_\chi$ .

We denote by  $\widehat{\mathfrak{D}}$  the group of all characters of  $\mathfrak{D}$ , and let  $\mathfrak{H}_{g,p}^b(\chi)$  present the  $\mathbb{C}_\chi$ -isotypic component of  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{C})$ . Thus we can decompose  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{C})$  as

$$\mathfrak{H}_{g,p}^b(\ell; \mathbb{C}) = \bigoplus_{\chi \in \widehat{\mathfrak{D}}} \mathfrak{H}_{g,p}^b(\chi).$$

Since the action of  $\text{Mod}_{g,p}^b(\ell)$  on  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{C})$  commutes with the action of  $\mathfrak{D}$ , each isotypic component is preserved, giving a decomposition into  $\text{Mod}_{g,p}^b(\ell)$ -modules. Taking the  $r$ -tensor power of  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{C})$ , we get

$$(5-7) \quad \mathfrak{H}_{g,p}^b(\ell; \mathbb{C})^{\otimes r} = \bigoplus_{\chi_1, \dots, \chi_r \in \widehat{\mathfrak{D}}} \mathfrak{H}_{g,p}^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_{g,p}^b(\chi_r).$$

A subgroup  $H < H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$  is called *symplectic* if the algebraic intersection pairing on  $H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$  restricts to a nondegenerate pairing on  $H$ . Here “nondegenerate” means that it identifies  $H$  with its dual  $\text{Hom}(H, \mathbb{Z}/\ell)$ . We can write  $H \cong (\mathbb{Z}/\ell)^{2h}$ , where  $h$  is called the genus of  $H$ .

For a symplectic subgroup  $H$ , there is a surjective homomorphism

$$(5-8) \quad \pi_1(\Sigma_{g,p}^b) \rightarrow H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell) = H \oplus H^\perp \xrightarrow{\text{proj}} H.$$

This map factors through  $\mathfrak{D} = H_1(\Sigma_g; \mathbb{Z}/\ell)$  since the homology classes of loops around boundary components and punctures lie in  $H^\perp$ . Thus we have a surjective homomorphism  $\mathfrak{D} \rightarrow H$ . A character  $\chi \in \widehat{\mathfrak{D}}$  is said to be *compatible* with  $H$  if it factors through the map  $\mathfrak{D} \rightarrow H$ . The following lemma is a special case of Lemma 6.9 in [\[Putman 2022\]](#):

**Lemma 5.7** For  $g > r$ , given  $r$  characters  $\chi_1, \dots, \chi_r \in \widehat{\mathcal{D}}$ , there exists a symplectic subgroup  $H$  of genus  $r$  such that  $\chi_1, \dots, \chi_r$  are all compatible with  $H$ .

**Proof** We define the group homomorphism

$$\mu_r : H_1(\Sigma_g; \mathbb{Z}) \rightarrow H_1(\Sigma_g; \mathbb{Z}/\ell) \rightarrow (\mathbb{C} \setminus \{0\})^r, \quad x \mapsto \bar{x} \mapsto (\chi_1(\bar{x}), \dots, \chi_r(\bar{x})).$$

Since elements in  $H_1(\Sigma_g; \mathbb{Z}/\ell) \cong (\mathbb{Z}/\ell)^{2g}$  have order dividing  $\ell$ , the image of each  $\chi_i$  lies in the cyclic group of  $\ell$ -th roots of unity. Thus we can rewrite  $\mu_r$  as

$$\mu_r : H_1(\Sigma_g; \mathbb{Z}) \rightarrow (\mathbb{Z}/\ell)^r.$$

By Lemma 3.5 in [Putman 2023], there exists a symplectic subspace  $V$  of  $H_1(\Sigma_g; \mathbb{Z})$  of genus  $(g - r)$  such that  $\mu_r|_V = 0$ . Letting  $H$  be the orthogonal complement of the image of  $V$  under the map  $H_1(\Sigma_g; \mathbb{Z}) \rightarrow H_1(\Sigma_g; \mathbb{Z}/\ell)$ , we have the required symplectic subgroup with genus  $r$ .  $\square$

Let  $\chi_1, \dots, \chi_r \in \widehat{\mathcal{D}}$  be  $r$  characters that are compatible with  $H$ . Define

$$\mathfrak{H}_{g,p}^b(\underline{\chi}) = \mathfrak{H}_{g,p}^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_{g,p}^b(\chi_r).$$

The *partial level- $\ell$  subgroup* of  $\text{Mod}_{g,p}^b$  is

$$\text{Mod}_{g,p}^b(H) = \{f \in \text{Mod}_{g,p}^b \mid f_* : H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell) \rightarrow H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell) \text{ fixes } H \text{ pointwise}\}.$$

Let  $S_H$  denotes the regular  $H$ -cover of  $\Sigma_{g,p}^b$  corresponding to the map (5-8). By definition, the partial level- $\ell$  subgroup  $\text{Mod}_{g,p}^b(H)$  acts on  $\mathfrak{H}_{g,p}^b(H; \mathbb{C}) := H^1(S_H; \mathbb{C})$ . In particular, this induces an action of  $\text{Mod}_{g,p}^b(H)$  on  $\mathfrak{H}_{g,p}^b(\underline{\chi})$  due to the decomposition of  $\mathfrak{H}_{g,p}^b(H; \mathbb{C})$  by Putman [2022, Lemma 6.5]

$$(5-9) \quad \mathfrak{H}_{g,p}^b(H; \mathbb{C}) \cong \bigoplus_{\chi \in \widehat{H}} \mathfrak{H}_{g,p}^b(\chi),$$

where  $\widehat{H}$  denotes the set of all characters of  $\mathcal{D}$  that are compatible with  $H$ .

By definition,  $\text{Mod}_{g,p}^b(\ell) < \text{Mod}_{g,p}^b(H)$  since  $H < H^1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$ . The following theorem of Putman allows us to identify the twisted cohomology of  $\text{Mod}_{g,p}^b(\ell)$  with the twisted cohomology of  $\text{Mod}_{g,p}^b(H)$ :

**Theorem 5.8** [Putman 2022, Theorem D] Let  $g, p, b \geq 0$  and  $\ell \geq 2$  be such that  $p + b \geq 1$ . Let  $H$  be a symplectic subgroup of  $H^1(\Sigma_g^b; \mathbb{Z}/\ell)$  and let  $\chi_1, \dots, \chi_r \in \mathcal{D}$  be  $r$  characters that are compatible with  $H$ . Assume that  $g \geq 2(k + r)^2 + 7k + 6r + 2$ . Then the map

$$H^k(\text{Mod}_{g,p}^b(H); \mathfrak{H}_{g,p}^b(\underline{\chi})) \rightarrow H^k(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\underline{\chi}))$$

induced by the inclusion  $\text{Mod}_{g,p}^b(\ell) \hookrightarrow \text{Mod}_{g,p}^b(H)$  is an isomorphism.

The following theorem of Putman implies a phenomenon where, when  $p = 0$ , the twisted cohomology of  $\text{Mod}_g^b(H)$  is independent of  $b$ , the number of boundary components. This phenomenon is important for our later proof.

**Theorem 5.9** [Putman 2022, Theorem 8.1] *Let  $\iota : \Sigma_g^b \rightarrow \Sigma_{g'}^{b'}$  be an orientation-preserving embedding between surfaces with nonempty boundary. For some  $l \geq 2$ , let  $H$  be a genus- $h$  symplectic subgroup of  $H_1(\Sigma_g^b; \mathbb{Z}/\ell)$ . Fix some  $k, r \geq 0$ , and assume that  $g \geq (2h+2)(k+r) + (4h+2)$ . Then the induced map*

$$H^k(\text{Mod}_{g'}^{b'}(H); \mathfrak{H}_{g'}^{b'}(H; \mathbb{C})^{\otimes r}) \rightarrow H^k(\text{Mod}_g^b(H); \mathfrak{H}_g^b(H; \mathbb{C})^{\otimes r})$$

is an isomorphism.

**Remark 5.10** If we take  $H$  to be a symplectic subgroup of genus  $g$ , the partial level- $\ell$  subgroup  $\text{Mod}_{g,p}^b(H)$  is exactly the level- $\ell$  subgroup  $\text{Mod}_{g,p}^b(\ell)$ , since  $\text{Mod}_{g,p}^b$  acts trivially on the subgroup of  $H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$  generated by loops around the boundary components and punctures of  $\Sigma_{g,p}^b$ . However, one cannot apply Theorem 5.9 directly to  $\text{Mod}_{g,p}^b(\ell)$ , because the condition  $g \geq (2h+2)(k+r) + (4h+2)$  cannot hold if we let  $h = g$ .

### 5.3 Proof of Theorem B

We now prove the case of  $\Sigma_{g,p}^b$  with  $p + b \geq 1$  by induction on  $p$ . Recall that we aim to show

$$H^{\bullet-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) \cong H^{\bullet}(\text{Mod}_{g,p}(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\otimes}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right)$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . We will prove the statement over  $\mathbb{C}$ .

For  $p = 0$ , the case of  $b = 1$  follows from Corollary 5.6. It then suffices to show that this twisted cohomology is independent of  $b$  for  $b \geq 1$ . Fix  $b \geq 1$ . Observe that there is an orientation-preserving embedding  $\Sigma_g^1 \hookrightarrow \Sigma_g^b$  by gluing a surface homeomorphic to  $\Sigma_0^{b+1}$  to the boundary of  $\Sigma_g^1$ . Then, by Theorem 5.9, we have the following isomorphism for any symplectic subgroup  $H$  of genus  $h$ , when  $g \geq (2h+2)k + (4h+2)$ :

$$(5-10) \quad H^{k-r}(\text{Mod}_g^b(H); \mathfrak{H}_g^b(H; \mathbb{C})^{\otimes r}) \cong H^{k-r}(\text{Mod}_g^1(H); \mathfrak{H}_g^1(H; \mathbb{C})^{\otimes r}).$$

These coefficients can be decomposed into direct sums of  $H$ -isotypic components by (5-9):

$$\begin{aligned} \mathfrak{H}_g^1(H; \mathbb{C})^{\otimes r} &\cong \left( \bigoplus_{\chi \in \hat{H}} \mathfrak{H}_g^1(\chi) \right)^{\otimes r} = \bigoplus_{\chi_1, \dots, \chi_r \in \hat{H}} \mathfrak{H}_g^1(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^1(\chi_r), \\ \mathfrak{H}_g^b(H; \mathbb{C})^{\otimes r} &\cong \left( \bigoplus_{\chi \in \hat{H}} \mathfrak{H}_g^b(\chi) \right)^{\otimes r} = \bigoplus_{\chi_1, \dots, \chi_r \in \hat{H}} \mathfrak{H}_g^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^b(\chi_r). \end{aligned}$$

We can expand both sides of (5-10) using the Künneth formula. Since the actions of  $\text{Mod}_g^b$  and  $\text{Mod}_g^1$  commute with the action of  $H$ , when  $g \geq (2h+2)k + (4h+2)$  we have an isomorphism in each direct-sum component:

$$(5-11) \quad H^{k-r}(\text{Mod}_g^b(H); \mathfrak{H}_g^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^b(\chi_r)) \cong H^{k-r}(\text{Mod}_g^1(H); \mathfrak{H}_g^1(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^1(\chi_r)).$$

By Lemma 5.7, for each tuple  $\underline{\chi} = (\chi_1, \dots, \chi_r)$ , where  $\chi_1, \dots, \chi_r \in \widehat{\mathcal{D}}$ , there exists a genus- $r$  symplectic subgroup, which we denote by  $H_{\underline{\chi}}$ , such that  $\chi_1, \dots, \chi_r$  are compatible with  $H_{\underline{\chi}}$ . By Theorem 5.8, we can identify the twisted cohomology of  $\text{Mod}_g^b(H_{\underline{\chi}})$  with the twisted cohomology of  $\text{Mod}_g^b(\ell)$ , with coefficients in  $\mathfrak{H}_g^b(\underline{\chi}) = \mathfrak{H}_g^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^b(\chi_r)$ :

$$(5-12) \quad H^{k-r}(\text{Mod}_g^b(H_{\underline{\chi}}); \mathfrak{H}_g^b(\underline{\chi})) \cong H^{k-r}(\text{Mod}_g^b(\ell); \mathfrak{H}_g^b(\underline{\chi})) \quad \text{if } g \geq 2k^2 + 7k - r + 2.$$

Combining these facts, we have

$$\begin{aligned} H^{k-r}(\text{Mod}_g^b(\ell); \mathfrak{H}_g^b(\ell; \mathbb{C})^{\otimes r}) &\cong H^{k-r}\left(\text{Mod}_g^b(\ell); \bigoplus_{\chi_1, \dots, \chi_r \in \widehat{\mathcal{D}}} \mathfrak{H}_g^b(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^b(\chi_r)\right) && \text{(by (5-7))} \\ &\cong \bigoplus_{\underline{\chi} \in (\widehat{\mathcal{D}})^{\times r}} H^{k-r}(\text{Mod}_g^b(\ell); \mathfrak{H}_g^b(\underline{\chi})) && \text{(K\"unneth)} \\ &\cong \bigoplus_{\underline{\chi} \in (\widehat{\mathcal{D}})^{\times r}} H^{k-r}(\text{Mod}_g^b(H_{\underline{\chi}}); \mathfrak{H}_g^b(\underline{\chi})) && \text{(by (5-12))} \\ &\cong \bigoplus_{\underline{\chi} \in (\widehat{\mathcal{D}})^{\times r}} H^{k-r}(\text{Mod}_g^1(H_{\underline{\chi}}); \mathfrak{H}_g^1(\underline{\chi})) && \text{(by (5-11))} \\ &\cong \bigoplus_{\underline{\chi} \in (\widehat{\mathcal{D}})^{\times r}} H^{k-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\underline{\chi})) && \text{(by (5-12))} \\ &\cong H^{k-r}\left(\text{Mod}_g^1(\ell); \bigoplus_{\chi_1, \dots, \chi_r \in \widehat{\mathcal{D}}} \mathfrak{H}_g^1(\chi_1) \otimes \dots \otimes \mathfrak{H}_g^1(\chi_r)\right) && \text{(K\"unneth)} \\ &\cong H^{k-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{C})^{\otimes r}) && \text{(by (5-7)),} \end{aligned}$$

where we need to be a little bit careful with the range. In the places where we apply (5-12), the range is  $g \geq 2k^2 + 7k - r + 2$ . In the fourth isomorphism where we apply (5-11), we let  $h = r$ , so the range is  $g \geq (2h + 2)k + (4h + 2) = (2r + 2)k + (4r + 2)$ . Since  $k \geq r$ , we have

$$2k^2 + 7k - r + 2 \geq (2r + 2)k + (4r + 2).$$

Thus the above statement holds in the range  $g \geq 2k^2 + 7k - r + 2$ . Since the isomorphism in Corollary 5.6 for  $H^{k-r}(\text{Mod}_g^1(\ell); \mathfrak{H}_g^1(\ell; \mathbb{C})^{\otimes r})$  holds when  $g \geq 2k^2 + 7k + 2$ , and  $2k^2 + 7k + 2 \geq 2k^2 + 7k - r + 2$ , the theorem for  $H^{k-r}(\text{Mod}_g^b(\ell); \mathfrak{H}_g^b(\ell; \mathbb{C})^{\otimes r})$  holds when  $g \geq 2k^2 + 7k + 2$  as well.

Now, let  $p \geq 1$ . By Proposition 2.5, we have the following short exact sequence obtained by gluing a punctured disk to  $\Sigma_{g,p-1}^{b+1}$ :

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Mod}_{g,p-1}^{b+1}(\ell) \rightarrow \text{Mod}_{g,p}^b(\ell) \rightarrow 1.$$

This sequence induces a Gysin sequence (Proposition 2.12) with coefficients in  $\mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r} \cong \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}$ :

$$\begin{aligned} \dots \rightarrow H^{\bullet-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) &\rightarrow H^{\bullet-r}(\text{Mod}_{g,p-1}^{b+1}(\ell); \mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r}) \\ &\rightarrow H^{\bullet-r-1}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r+1}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) \rightarrow \dots \end{aligned}$$

Here the map

$$\phi_{\bullet-r-1} : H^{\bullet-r-1}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) \rightarrow H^{\bullet-r+1}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$$

is multiplication by the Euler class  $e_p \in H^2(\text{Mod}_{g,p}^b(\ell); \mathbb{Q})$ , so it is injective. Similar to what we did in [Corollary 5.6](#), by observing the short exact sequence

$$1 \rightarrow \text{Coker}(\phi_{\bullet-r-2}) \rightarrow H^{\bullet-r}(\text{Mod}_{g,p-1}^{b+1}(\ell); \mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r}) \rightarrow \text{Ker}(\phi_{\bullet-r-1}) \rightarrow 1,$$

we get

$$H^{\bullet}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) \cong H^{\bullet}(\text{Mod}_{g,p-1}^{b+1}(\ell); \mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r}) \otimes_{\mathbb{Q}} \mathbb{Q}[e_p].$$

We know  $H^{\bullet}(\text{Mod}_{g,p-1}^{b+1}(\ell); \mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r})$  by the induction hypothesis. Thus we conclude

$$\begin{aligned} H^{\bullet-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r}) &\cong H^{\bullet}(\text{Mod}_{g,p-1}^{b+1}(\ell); \mathfrak{H}_{g,p-1}^{b+1}(\ell; \mathbb{Q})^{\otimes r}) \otimes_{\mathbb{Q}} \mathbb{Q}[e_p] \\ &\cong H^{\bullet}(\text{Mod}_{g,p-1}(\ell); \mathbb{Q}) \otimes \mathbb{Q}[e_p] \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathbb{Q}}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right) \\ &\cong H^{\bullet}(\text{Mod}_{g,p}(\ell); \mathbb{Q}) \otimes \left( \bigoplus_{\tilde{P} \in \mathcal{P}_r^{\mathbb{Q}}} \left( \prod_{\{i\} \in \tilde{P}} v_i \right) \mathbb{Q}[v_{\tilde{I}} : \tilde{I} \in \tilde{P}] a_{\tilde{P}} \right) \end{aligned}$$

in degrees  $k$  such that  $g \geq 2k^2 + 7k + 2$ . Note that this twisted cohomology is independent of  $b$  (the number of boundary components) but depends on  $p$  (the number of punctures) and  $g$  (the genus).  $\square$

## 6 Infinitesimal rigidity of symplectic Prym representations

In this section, we aim to prove [Theorem A](#) about infinitesimal rigidity of symplectic Prym representations.

Recall that the symplectic Prym representation is defined in the following way. Let  $\tilde{S} \rightarrow S$  be a finite-abelian cover with deck group  $A$ . Let  $\text{Mod}(S, A)$  be the subgroup of  $\text{Mod}(S)$  fixing  $A$  pointwise. Denote by  $\hat{S}$  the closed surface obtained by gluing disks to all boundary components and filling in all punctures of  $\tilde{S}$ . Then there is a symplectic action of  $\text{Mod}(S, A)$  on  $H^1(\hat{S}; \mathbb{R})$  which commutes with the deck group  $A$ :

$$\Phi : \text{Mod}(S, A) \rightarrow \text{Aut}(H^1(\hat{S}; \mathbb{R}))^A.$$

Denote the target Lie group by  $G_A$ . Let  $h$  be the genus of  $\hat{S}$ . Then we have

$$G_A = \text{Sp}(2h; \mathbb{R})^A = \{P \in \text{Sp}(2h; \mathbb{R}) \mid Pa = aP \text{ for all } a \in A\}.$$

First of all, we compute the Lie algebra  $\mathfrak{g}_A$  of  $G_A$ :

**Lemma 6.1** *The Lie algebra  $\mathfrak{g}_A$  of  $G_A = \text{Sp}(2h; \mathbb{R})^A$  is*

$$\mathfrak{g}_A = \mathfrak{sp}(2h; \mathbb{R})^A = \{X \in \mathfrak{sp}(2h; \mathbb{R}) \mid Xa = aX \text{ for all } a \in A\}.$$

**Proof** The Lie group  $G_A$  can be described as

$$G_A = \{P \in \text{GL}(2h; \mathbb{R}) \mid P^T J P = J, Pa = aP \text{ for all } a \in A\}, \quad \text{where } J = \begin{pmatrix} 0 & I_h \\ -I_h & 0 \end{pmatrix}.$$

The Lie algebra  $\mathfrak{g}_A$  of the matrix Lie group  $G_A$  is given by

$$\mathfrak{g}_A = \{X \in \text{Mat}(2h; \mathbb{R}) \mid e^{tX} \in G_A \text{ for all } t \in \mathbb{R}\}.$$

The Lie algebra of  $\text{Sp}(2h; \mathbb{R})$ , denoted by  $\mathfrak{sp}(2h; \mathbb{R})$ , satisfies  $X^T J + JX = 0$ . It remains to determine how the condition  $Pa = aP, \forall a \in A$  descends to  $\mathfrak{g}_A$ .

For  $X \in \mathfrak{g}_A$ , we require that  $e^{tX}a = ae^{tX}$  for any  $a \in A$ . Now expanding  $e^{tX}$  as a power series  $e^{tX} = \sum_{j=0}^{\infty} (tX)^j / j!$  and disregarding terms of order  $t^2$  or higher, we obtain  $tXa = atX$  for all  $t \in \mathbb{R}$ . Thus we have  $Xa = aX$  for any  $a \in A$ , so we conclude that  $\mathfrak{g}_A = \mathfrak{sp}(2h; \mathbb{R})^{\mathfrak{Q}}$ . □

Notice that by composing  $\Phi : \text{Mod}(S, A) \rightarrow G_A$  with the adjoint representation  $\text{Ad} : G_A \rightarrow \text{GL}(\mathfrak{g}_A)$ , we can view the Lie algebra  $\mathfrak{g}_A$  is a  $\text{Mod}(S, A)$ -module. We observe:

**Lemma 6.2** *The Lie algebra  $\mathfrak{g}_A$  is a  $\text{Mod}(S, A)$ -submodule of  $H^1(\widehat{S}; \mathbb{R})^{\otimes 2} \cong (\mathbb{R}^{2h})^{\otimes 2}$ .*

**Proof** We first embed  $\mathfrak{g}_A$  into  $H^1(\widehat{S}; \mathbb{R})^{\otimes 2} \cong (\mathbb{R}^{2h})^{\otimes 2}$  as

$$\mathfrak{g}_A = \mathfrak{sp}(2h; \mathbb{R})^A \subset \text{Mat}(2h; \mathbb{R}) \cong (\mathbb{R}^{2h})^* \otimes \mathbb{R}^{2h} \cong (\mathbb{R}^{2h})^{\otimes 2},$$

where the isomorphism  $(\mathbb{R}^{2h})^* \cong \mathbb{R}^{2h}$  is induced by the nondegenerate algebraic intersection form  $i : H_1(\widehat{S}; \mathbb{R}) \times H_1(\widehat{S}; \mathbb{R}) \rightarrow \mathbb{R}$ .

It remains to show that the action of  $\text{Mod}(S, A)$  on  $\mathfrak{g}_A$  is compatible with its action on  $H^1(\widehat{S}; \mathbb{R})^{\otimes 2}$ .

Take a symplectic basis  $\{\alpha_1, \beta_1, \dots, \alpha_h, \beta_h\}$  of  $H^1(\widehat{S}; \mathbb{R})$ . For  $X \in \mathfrak{g}_A$ , we can write

$$\begin{aligned} X &= \sum_{j=1}^h (\alpha_j)^* \otimes X\alpha_j + \sum_{j=1}^h (\beta_j)^* \otimes X\beta_j \in (\mathbb{R}^{2h})^* \otimes \mathbb{R}^{2h} \\ &= \sum_{j=1}^h \beta_j \otimes X\alpha_j + \sum_{j=1}^h (-\alpha_j) \otimes X\beta_j \in (\mathbb{R}^{2h})^{\otimes 2}. \end{aligned}$$

The action of  $f \in \text{Mod}(S, A)$  on  $X \in \mathfrak{g}_A$  is defined by  $f \cdot X = FXF^{-1}$ , where  $F = \Phi(f) \in \text{Sp}(2h; \mathbb{R})$ .

Thus we can express  $f \cdot X$  as

$$\begin{aligned} f \cdot X &= \sum_{j=1}^h (\alpha_j)^* \otimes FXF^{-1}\alpha_j + \sum_{j=1}^h (\beta_j)^* \otimes FXF^{-1}\beta_j \in (\mathbb{R}^{2h})^* \otimes \mathbb{R}^{2h} \\ &= \sum_{j=1}^h \beta_j \otimes FXF^{-1}\alpha_j + \sum_{j=1}^h (-\alpha_j) \otimes FXF^{-1}\beta_j \in (\mathbb{R}^{2h})^{\otimes 2} \\ &= \sum_{j=1}^h F(F^{-1}\beta_j) \otimes FX(F^{-1}\alpha_j) + \sum_{j=1}^h F(-F^{-1}\alpha_j) \otimes FX(F^{-1}\beta_j) \in (\mathbb{R}^{2h})^{\otimes 2}. \end{aligned}$$

Since  $\{F^{-1}(\alpha_1), F^{-1}(\beta_1), \dots, F^{-1}(\alpha_h), F^{-1}(\beta_h)\}$  is also a symplectic basis of  $H^1(\widehat{S}; \mathbb{R})$ , we can rewrite  $f \cdot X$  as

$$\sum_{j=1}^h F(\beta_j) \otimes F(X\alpha_j) + \sum_{j=1}^h F(-\alpha_j) \otimes F(X\beta_j).$$

This expression confirms that the action of  $\text{Mod}(S, A)$  on  $\mathfrak{g}_A$  coincides with its action on  $H^1(\widehat{S}; \mathbb{R})^{\otimes 2}$ .  $\square$

Next we prove that the degree-2 twisted cohomology group of  $\text{Mod}(S, A)$  with coefficients in  $H^1(\widehat{S}; \mathbb{R})^{\otimes 2}$  is 0 for sufficiently large  $g$ :

**Lemma 6.3** *Let  $g, p, b$  be integers such that  $p + b \geq 1$ . When  $g \geq 41$ , we have*

$$H^1(\text{Mod}(S, A); H^1(\widehat{S}; \mathbb{R})^{\otimes 2}) = 0.$$

**Proof Step 1** We first prove the result for  $A = H_1(\Sigma_g; \mathbb{Z}/\ell)$ . Note that by definition  $\text{Mod}(S, A)$  is usually larger than the level- $\ell$  mapping class group  $\text{Mod}_{g,p}^b(\ell)$  which acts trivially on  $H_1(\Sigma_{g,p}^b; \mathbb{Z}/\ell)$ .

In this case, the Prym representation  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})$  is  $H^1(\widetilde{S}; \mathbb{Q})$ . By filling in all punctures and gluing disks to all boundary components of  $\widetilde{S}$ , we obtain the short exact sequence

$$(6-1) \quad 0 \rightarrow H^1(\widehat{S}; \mathbb{Q}) \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \rightarrow \mathbb{Q}^{(p+b) \cdot |A| - 1} \rightarrow 0.$$

We then compute the following twisted cohomology groups one by one:

(1) We computed  $H^{k-r}(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes r})$  in [Theorem B](#) in the range  $g \geq 2k^2 + 7k + 2$ , and one feature of this twisted cohomology is that it is 0 when  $k$  is odd, since the left-hand side of the isomorphism in [Theorem B](#) consists of even-degree terms. In particular, setting  $k = 3$  and  $r = 2$ , we get

$$H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2}) = 0 \quad \text{if } g \geq 41.$$

Similarly, for  $k = 3$  and  $r = 1$ , we have

$$H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})) = 0 \quad \text{if } g \geq 41.$$

(2) The short exact sequence (6-1) of  $\text{Mod}_{g,p}^b(\ell)$ -modules induces a long exact sequence of twisted cohomology groups

$$\rightarrow H^3(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}^{(p+b) \cdot |A| - 1}) \rightarrow H^2(\text{Mod}_{g,p}^b(\ell); H^1(\widehat{S}; \mathbb{Q})) \rightarrow H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})) \rightarrow \dots$$

When  $g \geq 41$ , the term  $H^3(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}^{(p+b) \cdot |A| - 1})$  is 0 by [\[Putman 2022, Theorem A\]](#) ([Theorem 2.7](#)), and  $H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}))$  is also 0 as discussed in (1). Thus we have

$$H^2(\text{Mod}_{g,p}^b(\ell); H^1(\widehat{S}; \mathbb{Q})) = 0 \quad \text{if } g \geq 41.$$

(3) By tensoring the above short exact sequence (6-1) with  $\mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})$  on the left, we obtain another short exact sequence:

$$0 \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes H^1(\widehat{S}; \mathbb{Q}) \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2} \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes \mathbb{Q}^{(p+b) \cdot |A| - 1} \rightarrow 0.$$

This short exact sequence of  $\text{Mod}_{g,p}^b(\ell)$ -modules induces a long exact sequence of twisted cohomology groups:

$$\begin{aligned} \rightarrow H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes \mathbb{Q}^{(p+b)\cdot|A|-1}) &\rightarrow H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes H^1(\widehat{S}; \mathbb{Q})) \\ &\rightarrow H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2}) \rightarrow \dots \end{aligned}$$

Here  $H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes \mathbb{Q}^{(p+b)\cdot|A|-1}) \cong \bigoplus_{(p+b)\cdot|A|-1} H^2(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}))$  and  $H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q})^{\otimes 2})$  are both 0 when  $g \geq 41$  as discussed in (1), so

$$H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes H^1(\widehat{S}; \mathbb{Q})) = 0 \quad \text{if } g \geq 41.$$

(4) By tensoring the above short exact sequence (6-1) with  $H^1(\widehat{S}; \mathbb{Q})$  on the right, we obtain the following short exact sequence:

$$0 \rightarrow H^1(\widehat{S}; \mathbb{Q})^{\otimes 2} \rightarrow \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes H^1(\widehat{S}; \mathbb{Q}) \rightarrow \mathbb{Q}^{(p+b)\cdot|A|-1} \otimes H^1(\widehat{S}; \mathbb{Q}) \rightarrow 0.$$

This short exact sequence of  $\text{Mod}_{g,p}^b(\ell)$ -modules induces a long exact sequence of twisted cohomology groups:

$$\begin{aligned} \rightarrow H^2(\text{Mod}_{g,p}^b(\ell); \mathbb{Q}^{(p+b)\cdot|A|-1} \otimes H^1(\widehat{S}; \mathbb{Q})) &\rightarrow H^1(\text{Mod}_{g,p}^b(\ell); H^1(\widehat{S}; \mathbb{Q})^{\otimes 2}) \\ &\rightarrow H^1(\text{Mod}_{g,p}^b(\ell); \mathfrak{H}_{g,p}^b(\ell; \mathbb{Q}) \otimes H^1(\widehat{S}; \mathbb{Q})) \rightarrow \dots \end{aligned}$$

Here the first term is 0 when  $g \geq 41$  as discussed in (2), and the third term is 0 when  $g \geq 41$  as discussed in (3). Thus we have

$$H^1(\text{Mod}_{g,p}^b(\ell); H^1(\widehat{S}; \mathbb{Q})^{\otimes 2}) = 0 \quad \text{if } g \geq 41.$$

(5) Consider the finite-index subgroup  $\text{Mod}_{g,p}^b(\ell)$  of  $\text{Mod}(S, A)$ . By Proposition 2.9, the associated transfer map

$$H^1(\text{Mod}_{g,p}^b(\ell); H^1(\widehat{S}; \mathbb{R})^{\otimes 2}) \rightarrow H^1(\text{Mod}(S, A); H^1(\widehat{S}; \mathbb{R})^{\otimes 2})$$

is surjective. Thus from (4) we have

$$H^1(\text{Mod}(S, A); H^1(\widehat{S}; \mathbb{Q})^{\otimes 2}) = 0 \quad \text{if } g \geq 41.$$

Tensoring the result with  $\mathbb{R}$ , we finish the proof for  $A = H_1(\Sigma_g; \mathbb{Z}/\ell)$ .

**Step 2** We now extend the proof to a general finite abelian group  $A$ .

Recall that  $\widetilde{S}$  denotes the regular cover of  $S$  corresponding to the homomorphism

$$\pi_1(S) \rightarrow A.$$

Since  $A$  is abelian, this map factors through  $\pi_1(S) \rightarrow H_1(S; \mathbb{Z})$ . Letting  $\omega = |A|$ , this map furthermore factors through  $\pi_1(S) \rightarrow H_1(S; \mathbb{Z}/\omega)$ . Therefore  $\text{Mod}_{g,p}^b(\omega)$  is a finite-index subgroup of  $\text{Mod}(S, A)$ .

Denote by  $\tilde{S}_\omega$  the regular cover of  $\Sigma_g$  corresponding to the map  $\pi_1(\Sigma_g) \rightarrow H_1(\Sigma_g; \mathbb{Z}/\omega)$ . By part (4) in Step 1, we have

$$H^1(\text{Mod}_{g,p}^b(\omega); H^1(\tilde{S}_\omega; \mathbb{R})^{\otimes 2}) = 0 \quad \text{if } g \geq 41.$$

By filling in all punctures and gluing disks to all boundary components in the cover  $\tilde{S} \rightarrow S$ , we obtain a cover  $\hat{S} \rightarrow \Sigma_g$ , where the deck group is a quotient of  $H_1(\Sigma_g; \mathbb{Z}/\omega)$ . Therefore  $\tilde{S}_\omega$  is a finite cover of  $\hat{S}$ .

Since the action of  $\text{Mod}_{g,p}^b(\omega)$  on  $\hat{S}_\omega$  commutes with the deck group of  $\tilde{S}_\omega \rightarrow \hat{S}$ , Maschke's theorem implies that  $H^1(\hat{S}; \mathbb{R})^{\otimes 2}$  is a direct summand of  $H^1(\tilde{S}_\omega; \mathbb{R})^{\otimes 2}$  as an  $\text{Mod}_{g,p}^b(\omega)$ -module. Consequently,

$$H^1(\text{Mod}_{g,p}^b(\omega); H^1(\hat{S}; \mathbb{R})^{\otimes 2}) = 0 \quad \text{if } g \geq 41.$$

Finally, the transfer map associated with the finite-index subgroup  $\text{Mod}_{g,p}^b(\omega) < \text{Mod}(S, A)$  with coefficients in  $H^1(\hat{S}; \mathbb{Q})^{\otimes 2}$ ,

$$H^1(\text{Mod}_{g,p}^b(\omega); H^1(\hat{S}; \mathbb{R})^{\otimes 2}) \rightarrow H^1(\text{Mod}(S, A); H^1(\hat{S}; \mathbb{R})^{\otimes 2}),$$

is surjective by [Proposition 2.9](#). Thus we conclude that

$$H^1(\text{Mod}(S, A); H^1(\hat{S}; \mathbb{R})^{\otimes 2}) = 0 \quad \text{if } g \geq 41. \quad \square$$

Now we combine the above lemmas to prove [Theorem A](#):

**Proof of Theorem A** To prove the infinitesimal rigidity of the symplectic Prym representation

$$\Phi : \text{Mod}(S, A) \rightarrow G_A,$$

we need to show that

$$H^1(\text{Mod}(S, A); \mathfrak{g}_A) = 0 \quad \text{if } g \geq 41,$$

where  $\mathfrak{g}_A$  is the Lie algebra of the Lie group  $G_A$ .

From [Lemma 6.1](#), we have  $\mathfrak{g}_A = \mathfrak{sp}(2h; \mathbb{R})^A$ . By [Lemma 6.2](#), we see that  $\mathfrak{sp}(2h; \mathbb{R})^A$  is a  $\text{Mod}(S, A)$ -submodule of  $H^1(\hat{S}; \mathbb{R})^{\otimes 2}$ . Therefore, it suffices to prove that

$$H^1(\text{Mod}(S, A); H^1(\hat{S}; \mathbb{R})^{\otimes 2}) = 0 \quad \text{if } g \geq 41,$$

which is precisely the result of [Lemma 6.3](#). This completes the proof. □

### Acknowledgements

I sincerely thank my advisor Andrew Putman for suggesting the problem, providing many helpful comments, and offering continuous encouragement throughout the entire project, as well as for numerous suggestions on earlier drafts of the paper. I would also like to thank Eduard Looijenga, Oscar Randal-Williams and Eric Riedl for helpful conversations. Finally, I thank the referees for their many useful comments.

## References

- [Atiyah and Bott 1983] **M F Atiyah, R Bott**, *The Yang–Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A 308:1505 (1983) 523–615 [MR](#)
- [Boldsen 2012] **S K Boldsen**, *Improved homological stability for the mapping class group with integral or twisted coefficients*, Math. Z. 270:1-2 (2012) 297–329 [MR](#)
- [Brown 1982] **K S Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer (1982) [MR](#)
- [Deligne 1971] **P Deligne**, *Théorie de Hodge, II*, Inst. Hautes Études Sci. Publ. Math. 40 (1971) 5–57 [MR](#)
- [Deligne 1974] **P Deligne**, *Théorie de Hodge, III*, Inst. Hautes Études Sci. Publ. Math. 44 (1974) 5–77 [MR](#)
- [Esnault and Groechenig 2018] **H Esnault, M Groechenig**, *Cohomologically rigid local systems and integrality*, Selecta Math. 24:5 (2018) 4279–4292 [MR](#)
- [Galatius 2013] **S Galatius**, *Lectures on the Madsen–Weiss theorem*, from “Moduli spaces of Riemann surfaces” (B Farb, R Hain, E Looijenga, editors), IAS/Park City Math. Ser. 20, Amer. Math. Soc., Providence, RI (2013) 139–167 [MR](#)
- [Griffiths and Schmid 1975] **P Griffiths, W Schmid**, *Recent developments in Hodge theory: a discussion of techniques and results*, from “Discrete subgroups of Lie groups and applications to moduli” (Bombay, 1973), Tata Inst. Fundam. Res. Stud. Math. 7, Oxford Univ. Press, Bombay (1975) 31–127 [MR](#)
- [Harer 1985] **J L Harer**, *Stability of the homology of the mapping class groups of orientable surfaces*, Ann. of Math. 121:2 (1985) 215–249 [MR](#)
- [Hatcher 2011] **A Hatcher**, *A short exposition of the Madsen–Weiss theorem*, preprint (2011) [arXiv 1103.5223](#)
- [Hochschild and Serre 1953] **G Hochschild, J-P Serre**, *Cohomology of group extensions*, Trans. Amer. Math. Soc. 74 (1953) 110–134 [MR](#)
- [Ivanov 1989] **N V Ivanov**, *Stabilization of the homology of Teichmüller modular groups*, Algebra i Analiz 1:3 (1989) 110–126 [MR](#) In Russian; translated in Leningrad Math. J. 1 (1990) 675–691
- [Kawazumi 2008] **N Kawazumi**, *On the stable cohomology algebra of extended mapping class groups for surfaces*, from “Groups of diffeomorphisms” (R Penner, D Kotschick, T Tsuboi, N Kawazumi, T Kitano, Y Mitsumatsu, editors), Adv. Stud. Pure Math. 52, Math. Soc. Japan, Tokyo (2008) 383–400 [MR](#)
- [Klewdal and Patrikis 2020] **C Klewdal, S Patrikis**, *G-rigid local systems are integral*, preprint (2020) [arXiv 2009.07350](#)
- [Kupers and Randal-Williams 2020] **A Kupers, O Randal-Williams**, *On the cohomology of Torelli groups*, Forum Math. Pi 8 (2020) e7, 83 [MR](#)
- [Looijenga 1996] **E Looijenga**, *Stable cohomology of the mapping class group with symplectic coefficients and of the universal Abel–Jacobi map*, J. Algebraic Geom. 5:1 (1996) 135–150 [MR](#)
- [Looijenga 1997] **E Looijenga**, *Prym representations of mapping class groups*, Geom. Dedicata 64:1 (1997) 69–83 [MR](#)
- [Lubotzky and Magid 1985] **A Lubotzky, A R Magid**, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. 336, Amer. Math. Soc., Providence, RI (1985) [MR](#)
- [Madsen and Weiss 2007] **I Madsen, M Weiss**, *The stable moduli space of Riemann surfaces: Mumford’s conjecture*, Ann. of Math. 165:3 (2007) 843–941 [MR](#)
- [Miller 1986] **E Y Miller**, *The homology of the mapping class group*, J. Differential Geom. 24:1 (1986) 1–14 [MR](#)
- [Morita 1987] **S Morita**, *Characteristic classes of surface bundles*, Invent. Math. 90:3 (1987) 551–577 [MR](#)
- [Mumford 1983] **D Mumford**, *Towards an enumerative geometry of the moduli space of curves*, from “Arithmetic and geometry, II” (M Artin, J Tate, editors), Progr. Math. 36, Birkhäuser, Boston, MA (1983) 271–328 [MR](#)
- [Mumford et al. 1994] **D Mumford, J Fogarty, F Kirwan**, *Geometric invariant theory*, 3rd edition, Ergebnisse der Math. 34, Springer (1994) [MR](#)
- [Perron 2008] **B Perron**, *Filtration de Johnson et groupe de Torelli modulo  $p$ ,  $p$  premier*, C. R. Math. Acad. Sci. Paris 346:11-12 (2008) 667–670 [MR](#)
- [Peters and Steenbrink 2008] **C A M Peters, J H M Steenbrink**, *Mixed Hodge structures*, Ergebnisse der Math. (3) 52, Springer (2008) [MR](#)

- [Putman 2012] **A Putman**, *The Picard group of the moduli space of curves with level structures*, Duke Math. J. 161:4 (2012) 623–674 [MR](#)
- [Putman 2014] **A Putman**, *The action on homology of finite groups of automorphisms of surfaces and graphs*, notes (2014) Available at <https://academicweb.nd.edu/~andyp/notes/FiniteOrderHomology.pdf>
- [Putman 2022] **A Putman**, *The stable cohomology of the moduli space of curves with level structures*, preprint (2022) [arXiv 2209.06183](https://arxiv.org/abs/2209.06183)
- [Putman 2023] **A Putman**, *Partial Torelli groups and homological stability*, Algebr. Geom. Topol. 23:8 (2023) 3417–3496 [MR](#)
- [Randal-Williams 2016] **O Randal-Williams**, *Resolutions of moduli spaces and homological stability*, J. Eur. Math. Soc. (JEMS) 18:1 (2016) 1–81 [MR](#)
- [Randal-Williams 2018] **O Randal-Williams**, *Cohomology of automorphism groups of free groups with twisted coefficients*, Selecta Math. 24:2 (2018) 1453–1478 [MR](#)
- [Sato 2010] **M Sato**, *The abelianization of the level  $d$  mapping class group*, J. Topol. 3:4 (2010) 847–882 [MR](#)
- [Wahl 2013] **N Wahl**, *The Mumford conjecture, Madsen–Weiss and homological stability for mapping class groups of surfaces*, from “Moduli spaces of Riemann surfaces” (B Farb, R Hain, E Looijenga, editors), IAS/Park City Math. Ser. 20, Amer. Math. Soc., Providence, RI (2013) 109–138 [MR](#)
- [Weil 1964] **A Weil**, *Remarks on the cohomology of groups*, Ann. of Math. 80 (1964) 149–157 [MR](#)

XIYAN ZHONG [xiyanmath@gmail.com](mailto:xiyanmath@gmail.com)

*Department of Mathematics, University of Notre Dame, South Bend, IN, United States*

Proposed: Dan Abramovich

Received: April 3, 2024

Seconded: Benson Farb, Nathalie Wahl

Revised: April 17, 2025

# GEOMETRY & TOPOLOGY

[msp.org/gt](http://msp.org/gt)

## MANAGING EDITORS

Robert Lipshitz University of Oregon  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)  
András I Stipsicz Alfréd Rényi Institute of Mathematics  
[stipsicz@renyi.hu](mailto:stipsicz@renyi.hu)

## BOARD OF EDITORS

Mohammed Abouzaid	Stanford University <a href="mailto:abouzaid@stanford.edu">abouzaid@stanford.edu</a>	Rob Kirby	University of California, Berkeley <a href="mailto:kirby@math.berkeley.edu">kirby@math.berkeley.edu</a>
Dan Abramovich	Brown University <a href="mailto:dan_abramovich@brown.edu">dan_abramovich@brown.edu</a>	Bruce Kleiner	NYU, Courant Institute <a href="mailto:bkleiner@cims.nyu.edu">bkleiner@cims.nyu.edu</a>
Ian Agol	University of California, Berkeley <a href="mailto:ianagol@math.berkeley.edu">ianagol@math.berkeley.edu</a>	Sándor Kovács	University of Washington <a href="mailto:skovacs@uw.edu">skovacs@uw.edu</a>
Arend Bayer	University of Edinburgh <a href="mailto:arend.bayer@ed.ac.uk">arend.bayer@ed.ac.uk</a>	Urs Lang	ETH Zürich <a href="mailto:urs.lang@math.ethz.ch">urs.lang@math.ethz.ch</a>
Agnès Beaudry	University of Colorado Boulder <a href="mailto:agnes.beaudry@colorado.edu">agnes.beaudry@colorado.edu</a>	Marc Levine	Universität Duisburg-Essen <a href="mailto:marc.levine@uni-due.de">marc.levine@uni-due.de</a>
Mark Behrens	University of Notre Dame <a href="mailto:mbehren1@nd.edu">mbehren1@nd.edu</a>	Jianfeng Lin	Tsinghua University <a href="mailto:linjian5477@mail.tsinghua.edu.cn">linjian5477@mail.tsinghua.edu.cn</a>
Mladen Bestvina	University of Utah <a href="mailto:bestvina@math.utah.edu">bestvina@math.utah.edu</a>	Ciprian Manolescu	University of California, Los Angeles <a href="mailto:cm@math.ucla.edu">cm@math.ucla.edu</a>
Martin R Bridson	University of Oxford <a href="mailto:bridson@maths.ox.ac.uk">bridson@maths.ox.ac.uk</a>	Haynes Miller	Massachusetts Institute of Technology <a href="mailto:hmm@math.mit.edu">hmm@math.mit.edu</a>
Tobias H Colding	Massachusetts Institute of Technology <a href="mailto:colding@math.mit.edu">colding@math.mit.edu</a>	Aaron Naber	Institute for Advanced Studies <a href="mailto:anaber@ias.edu">anaber@ias.edu</a>
Simon Donaldson	Imperial College, London <a href="mailto:s.donaldson@ic.ac.uk">s.donaldson@ic.ac.uk</a>	Peter Ozsváth	Princeton University <a href="mailto:petero@math.princeton.edu">petero@math.princeton.edu</a>
Yasha Eliashberg	Stanford University <a href="mailto:eliash-gt@math.stanford.edu">eliash-gt@math.stanford.edu</a>	Leonid Polterovich	Tel Aviv University <a href="mailto:polterov@post.tau.ac.il">polterov@post.tau.ac.il</a>
Benson Farb	University of Chicago <a href="mailto:farb@math.uchicago.edu">farb@math.uchicago.edu</a>	Colin Rourke	University of Warwick <a href="mailto:gt@maths.warwick.ac.uk">gt@maths.warwick.ac.uk</a>
David M Fisher	Rice University <a href="mailto:davidfisher@rice.edu">davidfisher@rice.edu</a>	Roman Sauer	Karlsruhe Institute of Technology <a href="mailto:roman.sauer@kit.edu">roman.sauer@kit.edu</a>
Mike Freedman	Microsoft Research <a href="mailto:michaelf@microsoft.com">michaelf@microsoft.com</a>	Stefan Schwede	Universität Bonn <a href="mailto:schwede@math.uni-bonn.de">schwede@math.uni-bonn.de</a>
David Gabai	Princeton University <a href="mailto:gabai@princeton.edu">gabai@princeton.edu</a>	Natasa Sesum	Rutgers University <a href="mailto:natasas@math.rutgers.edu">natasas@math.rutgers.edu</a>
Stavros Garoufalidis	Southern U. of Sci. and Tech., China <a href="mailto:stavros@mpim-bonn.mpg.de">stavros@mpim-bonn.mpg.de</a>	Gang Tian	Massachusetts Institute of Technology <a href="mailto:tian@math.mit.edu">tian@math.mit.edu</a>
Cameron Gordon	University of Texas <a href="mailto:gordon@math.utexas.edu">gordon@math.utexas.edu</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Jesper Grodal	University of Copenhagen <a href="mailto:jg@math.ku.dk">jg@math.ku.dk</a>	Kirsten Wickelgren	Duke University <a href="mailto:kirsten.wickelgren@duke.edu">kirsten.wickelgren@duke.edu</a>
Misha Gromov	IHES and NYU, Courant Institute <a href="mailto:gromov@ihes.fr">gromov@ihes.fr</a>	Anna Wienhard	Universität Heidelberg <a href="mailto:wienhard@mathi.uni-heidelberg.de">wienhard@mathi.uni-heidelberg.de</a>
Mark Gross	University of Cambridge <a href="mailto:mgross@dpmms.cam.ac.uk">mgross@dpmms.cam.ac.uk</a>		

---

See inside back cover or [msp.org/gt](http://msp.org/gt) for submission instructions.


The subscription price for 2026 is US \$905/year for the electronic version, and \$1275/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

---

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

# GEOMETRY & TOPOLOGY

Volume 30

Issue 1 (pages 1–388)

2026

---

Free boundary flow with surgery ROBERT HASLHOFER	1
Limits of asymptotically Fuchsian surfaces in a closed hyperbolic 3-manifold FERNANDO AL ASSAL	23
The 3-fold K-theoretic DT/PT vertex correspondence holds NIKOLAS KUHN, HENRY LIU and FELIX THIMM	71
Prym representations and twisted cohomology of the mapping class group with level structures XIYAN ZHONG	155
Modular sheaves with many moduli KIERAN G O'GRADY	203
Degenerating conic Kähler–Einstein metrics to the normal cone OLIVIER BIQUARD and HENRI GUENANCIA	247
Degree two Gopakumar–Vafa invariants of local curves BEN DAVISON and NAOKI KOSEKI	307
On pro-cdh descent on derived schemes SHANE KELLY, SHUJI SAITO and GEORG TAMME	337
On the cohomological dimension of kernels of maps to $\mathbb{Z}$ SAM P. FISHER	373