



Geometry & Topology

Volume 30 (2026)

K3 surfaces associated with varieties of generalized Kummer type

SALVATORE FLOCCARI

K3 surfaces associated with varieties of generalized Kummer type

SALVATORE FLOCCARI

Any hyper-Kähler variety K of generalized Kummer type is associated via Hodge theory with a K3 surface S_K . We show how they are related geometrically through a moduli space of sheaves on S_K . As a consequence, building fundamentally on the works of O’Grady, Markman, Voisin and Varesco, we establish the Hodge conjecture for all powers of any of these K3 surfaces as well as for all abelian fourfolds of Weil type with discriminant 1 and their powers, strengthening a result of Markman.

1 Introduction

A manifold of generalized Kummer type is, by definition, a compact Kähler manifold which is deformation equivalent to the generalized Kummer variety $K^n(A)$ on an abelian surface A constructed by Beauville [1983]. Manifolds of generalized Kummer type give examples of hyper-Kähler manifolds [Beauville 1983; Huybrechts 1999] in each even dimension $2n$, commonly referred to as manifolds of Kum ^{n} -type; by a variety of Kum ^{n} -type we shall indicate one such manifold which is projective. For any $n > 1$, these projective varieties form countably many 4-dimensional families. Besides Beauville’s construction of generalized Kummer varieties, other examples can be obtained by considering moduli of stable sheaves on abelian surfaces [Yoshioka 2001]; however, it is known that the general projective variety of generalized Kummer type cannot be realized from such a construction.

In this article we consider some K3 surfaces naturally attached via Hodge theory to varieties of generalized Kummer type. Let K be a variety of Kum ^{n} -type. By the Torelli theorem there exists a unique K3 surface S_K such that there exists a Hodge isometry $H_{\text{tr}}^2(S_K, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Z})(2)$ of transcendental lattices, where on the right-hand side the Beauville–Bogomolov form is multiplied by a factor 2. If $K = K^n(A)$ is the generalized Kummer variety on an abelian surface A , then S_K is the Kummer K3 surface $\text{Km}(A)$, which motivates our definition.

We will explain how the K3 surface S_K is related geometrically to K , for any variety K of Kum ^{n} -type. Indeed, there exists a natural subvariety W_K of K which is a hyper-Kähler variety of K3^[m]-type where either $2m = n$ or $2m = n + 1$, with the key property of deforming everywhere together with K ; i.e., W_K is a trianalytic submanifold of K in the sense of Verbitsky [1995]. It is obtained as the component of maximal dimension of the fixed locus of an involution of K which acts trivially on the second cohomology group; such automorphisms were studied already in [Hassett and Tschinkel 2013; Boissière et al. 2011; Oguiso 2020; Kamenova et al. 2022; Floccari 2024] and used to produce trianalytic cycles in [Kaledin and Verbitsky 1998]. Our contribution lies in the determination of the period of W_K :

MSC2020: 14C15, 14C25, 14C30, 14J28, 14J42.

© 2026 The Author, under license to MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

we show that the pullback along $\iota : W_K \hookrightarrow K$ induces a primitive embedding of lattices and Hodge structures $\iota^* : H_{\text{tr}}^2(K, \mathbb{Z})(2) \xrightarrow{\sim} H_{\text{tr}}^2(W_K, \mathbb{Z})$, where on the left-hand side the form is multiplied by 2. It follows that $W_K \subset K$ is a smooth and projective moduli space of stable sheaves on the K3 surface S_K .

As a consequence of the above construction, we obtain the following result.

Theorem 1.1 *There exists an algebraic cycle on $S_K \times K$ inducing a Hodge isometry*

$$H_{\text{tr}}^2(S_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(2).$$

This algebraic cycle gives in fact an isomorphism between the transcendental parts of the homological motives of S_K and K in degree 2; see Theorem 5.9.

Building on results of O’Grady [2021], Markman [2023], Voisin [2022] and Varesco [2025], we use Theorem 1.1 to prove some cases of the Hodge conjecture.

Theorem 1.2 *Let K be any projective variety of Kumⁿ-type, and let S_K be the K3 surface with transcendental lattice $H_{\text{tr}}^2(S_K, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Z})(2)$. Then:*

- (i) *The Kuga–Satake correspondence is algebraic for S_K .*
- (ii) *The Hodge conjecture holds for any power of S_K .*

The theorem gives countably many new families of K3 surfaces of general Picard rank 16 for which it is possible to prove (i) and (ii), generalizing previous results from [Paranjape 1988; Ingalls et al. 2022; Floccari 2024]. Using that isogenies of K3 surfaces are algebraic by [Buskin 2019; Huybrechts 2019], we show that the conclusions of Theorem 1.2 hold for any K3 surface S such that there exists an isometric embedding of rational quadratic spaces $H_{\text{tr}}^2(S, \mathbb{Q}) \hookrightarrow (U^{\oplus 3} \oplus \langle -m \rangle_{\mathbb{Q}})$ for some positive integer m , where U denotes a hyperbolic plane; see Theorem 5.11.

Let us summarize the main points of the proof. Recall that the Kuga–Satake construction [1967] (see also [Deligne 1972]) produces an abelian variety $\text{KS}(X)$ from a polarized hyper-Kähler variety X , with an embedding of Hodge structures $H^2(X, \mathbb{Q})_{\text{prim}} \hookrightarrow H^2(\text{KS}(S)^2, \mathbb{Q})$ called the Kuga–Satake correspondence; according to the Hodge conjecture, this embedding should be induced by an algebraic cycle.

It was shown by O’Grady [2021] and Markman [2023] that the Kuga–Satake variety of a Kumⁿ-variety is isogenous to a power of its third intermediate Jacobian $J^3(K)$, which is an abelian fourfold of Weil type with discriminant 1. Moreover, varying K in polarized families, one obtains all complete up to isogeny families of such abelian fourfolds; see also [van Geemen 2023]. Abelian fourfolds of Weil type are characterized by the presence of exceptional Hodge classes: there is a 2-dimensional space of Hodge classes in their middle cohomology which, for the very general such fourfold, cannot be represented as intersection of divisors; see [van Geemen 2000; Moonen and Zarhin 1995]. Markman [2023] has proven that these Hodge–Weil classes are algebraic for any fourfold of Weil type of discriminant 1; his results further imply that the Kuga–Satake correspondence is algebraic for any variety of Kumⁿ-type, as shown by Voisin [2022].

Theorem 1.2 is then obtained as follows. By definition, the Kuga–Satake variety of S_K is isogenous to a power of the Kuga–Satake variety of K , and, hence, to a power of a Weil fourfold with discriminant 1. We deduce Theorem 1.2(i) from Theorem 1.1 and the algebraicity of the Kuga–Satake correspondence for the Kumⁿ-variety K . Statement (ii) is a consequence of (i) thanks to the work of Varesco [2025].

The Hodge conjecture for a very general fourfold A of Weil type with discriminant 1 is proven by Markman [2023], which implies the Hodge conjecture also for powers of such A by [Varesco 2025; Milne 2021]. However, the results of [Markman 2023] do not directly imply the Hodge conjecture for an arbitrary Weil fourfold with discriminant 1. We complete the proof of the Hodge conjecture for all Weil fourfolds with discriminant 1, and their powers.

Theorem 1.3 *Let A be an abelian fourfold of Weil type with discriminant 1. Then the Hodge conjecture holds for A and all of its powers.*

To prove this statement, we show in Theorem 3.5 that if S is any K3 surface such that the Kuga–Satake correspondence is algebraic and the Hodge conjecture holds for any power of S , then the Hodge conjecture holds for any power of the Kuga–Satake variety of S as well. We then obtain Theorem 1.3 from Theorem 1.2 using that any Weil fourfold with discriminant 1 appears as isogeny factor of the Kuga–Satake variety of some K3 surface S_K as in Theorem 1.2.

2 Motives

2.1 A neutral Tannakian category \mathcal{C} is a \mathbb{Q} -linear abelian rigid tensor category in which $\text{End}(1) = \mathbb{Q}$ and which admits an exact faithful tensor functor $\omega : \mathcal{C} \rightarrow \text{Vect}_{\mathbb{Q}}$. We refer the reader to [Deligne and Milne 1982] for more details. The group of tensor automorphisms of ω is a (pro-)algebraic group $\mathcal{G}(\mathcal{C})$ with the property that \mathcal{C} is \otimes -equivalent to the category of finite-dimensional representations of $\mathcal{G}(\mathcal{C})$. Thus, neutral Tannakian subcategories of \mathcal{C} correspond bijectively to quotients of $\mathcal{G}(\mathcal{C})$. Given $M \in \mathcal{C}$, we denote by $\langle M \rangle_{\mathcal{C}}$ the neutral Tannakian subcategory of \mathcal{C} generated by M ; more directly, it is defined as the smallest thick and full subcategory containing M and closed under direct sums, duals, tensor products and subobjects. Then $\langle M \rangle_{\mathcal{C}}$ is equivalent to the category of finite-dimensional representations of a linear algebraic group $G_{\mathcal{C}}(M) \subset \text{GL}(\omega(M))$, called the Tannakian fundamental group of M .

2.2 The category HS of polarizable pure \mathbb{Q} -Hodge structures is an abelian and semisimple rigid tensor category, which is neutral Tannakian via the forgetful functor to \mathbb{Q} -vector spaces. The Tannakian fundamental group of a polarizable Hodge structure V is a reductive and connected algebraic group over \mathbb{Q} ([Moonen 1999]) denoted by $\text{MT}(V) \subset \text{GL}(V)$ and called the Mumford–Tate group of V . It is equivalently defined as the subgroup of $\text{GL}(V)$ which fixes all Hodge classes in any tensor construction

$$V^{\otimes i} \otimes V^{\vee, \otimes j} \otimes \mathbb{Q}(k),$$

where $\mathbb{Q}(k)$ is the Tate Hodge structure of weight $-2k$. Provided that V is not entirely of weight 0, there is a central torus $\mathbb{G}_{m, \mathbb{Q}} \subset \text{MT}(V)$ whose action induces the weight decomposition of the objects in $\langle V \rangle_{\text{HS}}$.

Remark 2.1 Let $V \in \text{HS}$ be a Hodge structure of odd weight. Denote by $\langle V \rangle_{\text{HS}}^{\text{ev}}$ the full subcategory of $\langle V \rangle_{\text{HS}}$ of Hodge structures of even weight, which is a neutral Tannakian subcategory of HS . If $w : \mathbb{G}_{m, \mathbb{Q}} \rightarrow \text{MT}(V)$ denotes the inclusion of the weight torus, the objects of $\langle V \rangle_{\text{HS}}^{\text{ev}}$ are precisely the Hodge structures in $\langle V \rangle_{\text{HS}}$ on which $w(-1)$ acts trivially. Hence, we have $\text{MT}(\langle V \rangle_{\text{HS}}^{\text{ev}}) = \text{MT}(V)/w(-1)$. It follows that if W is a Hodge structure of even weight in $\langle V \rangle_{\text{HS}}$ such that the induced surjective morphism $\phi : \text{MT}(V) \rightarrow \text{MT}(W)$ of Mumford–Tate groups is an isogeny of degree 2, we then have the equality $\langle V \rangle_{\text{HS}}^{\text{ev}} = \langle W \rangle_{\text{HS}}$ of subcategories of HS . Indeed, since W has even weight, it belongs to $\langle V \rangle_{\text{HS}}^{\text{ev}}$, and, therefore, ϕ factors as the composition of the projection $\psi : \text{MT}(V) \rightarrow \text{MT}(\langle V \rangle_{\text{HS}}^{\text{ev}})$ followed by a surjective morphism $\phi' : \text{MT}(\langle V \rangle_{\text{HS}}^{\text{ev}}) \rightarrow \text{MT}(W)$ of algebraic groups. But both ϕ and ψ are isogenies of degree 2, and, hence, ϕ' is an isomorphism; it follows that $\langle W \rangle_{\text{HS}} = \langle V \rangle_{\text{HS}}^{\text{ev}}$.

2.3 An algebraic class on a smooth and projective complex variety X is a cohomology class α in $H^{2k}(X, \mathbb{Q})$ which is \mathbb{Q} -linear combination of fundamental classes of subvarieties of X . Any algebraic class is a Hodge class, and the Hodge conjecture predicts that, conversely, all Hodge classes are algebraic. A correspondence from X to Y of degree k is by definition an algebraic class in $H^{2k+2 \dim X}(X \times Y, \mathbb{Q})$; correspondences may be composed [Fulton 1998]. Grothendieck’s theory of motives is very useful in the study of algebraic cycles; in brief, the main insight is to use correspondences as morphisms. We denote by Mot the category of homological motives over \mathbb{C} with rational coefficients; see [André 2004]. Its objects are triples (X, p, n) where $X \in \text{SmProj}_{\mathbb{C}}$, $p \in H^{2 \dim X}(X \times X, \mathbb{Q})$ is an idempotent correspondence, and n is an integer. Morphisms from (X, p, n) to (Y, q, m) are defined to be the correspondences $f \in H^{2m-2n+2 \dim X}(X \times Y, \mathbb{Q})$ of degree $m - n$ such that $f \circ p = q \circ f$. There is a realization functor $R : \text{Mot} \rightarrow \text{HS}$ which sends (X, p, n) to $p_*(H^\bullet(X, \mathbb{Q})(n))$. Moreover, there is a contravariant functor $h : \text{SmProj}_{\mathbb{C}} \rightarrow \text{Mot}$ which attaches to the smooth and projective variety X its motive $h(X) := (X, \Delta_X, 0)$, where $\Delta_X \in H^{2 \dim X}(X \times X, \mathbb{Q})$ is the class of the diagonal. The composition $R \circ h$ is the functor $H^\bullet : \text{SmProj}_{\mathbb{C}} \rightarrow \text{HS}$ which attaches to the smooth and projective complex variety X its rational Hodge structure $H^\bullet(X, \mathbb{Q})$.

2.4 The category Mot of homological motives is a pseudoabelian tensor category via the product of varieties. However, it is not known to be abelian nor Tannakian: these properties are conditional to the validity of Grothendieck’s standard conjectures [1969]. By the work of Jannsen [1992] and André [1996b], we have the following. Given $m \in \text{Mot}$, we let $\langle m \rangle_{\text{Mot}}$ be the smallest thick and full subcategory of Mot containing m and closed under direct sums, duals, tensor products and subquotients.

Theorem 2.2 [Arapura 2006, Theorem 4.1] *Let X be a smooth and projective variety. The standard conjectures hold for X if and only if $\langle h(X) \rangle_{\text{Mot}}$ is a semisimple abelian neutral Tannakian category.*

Thus, provided that the standard conjectures hold for X , we can attach to it a reductive algebraic group $G_{\text{Mot}}(h(X))$ over \mathbb{Q} , whose representation category is equivalent to $\langle h(X) \rangle_{\text{Mot}}$. The algebraic group $G_{\text{Mot}}(h(X))$ is characterized as the subgroup of $\prod_j \text{GL}(H^j(X, \mathbb{Q}))$ fixing exactly the algebraic classes in $H^{2j}(X^k, \mathbb{Q}(j))$ for any j and k . There is a canonical inclusion $i_X : \text{MT}(H^\bullet(X, \mathbb{Q})) \hookrightarrow G_{\text{Mot}}(h(X))$,

which is an isomorphism if and only if the Hodge conjecture holds for X and all of its powers. Moreover, any object of $\langle h(X) \rangle_{\text{Mot}}$ admits a weight decomposition induced by the action of the weight torus $\mathbb{G}_m \subset \text{MT}(H^\bullet(X, \mathbb{Q})) \subset G_{\text{Mot}}(X)$. In particular, we have the Künneth decomposition $h(X) = \bigoplus_i h^i(X)$ in Mot .

Remark 2.3 Let X be a smooth and projective variety and assume that the Künneth component $h^2(X)$ of $h(X)$ is well-defined. Thanks to the Lefschetz theorem on divisors, this motive splits as $h^2(X) = h_{\text{tr}}^2(X) \oplus h_{\text{alg}}^2(X)$: the realization of $h_{\text{alg}}^2(X)$ is the Néron-Severi group $\text{NS}(X)_{\mathbb{Q}}$ while that of $h_{\text{tr}}^2(X)$ is the transcendental part $H_{\text{tr}}^2(X, \mathbb{Q})$ of $H^2(X, \mathbb{Q})$, the smallest sub-Hodge structure whose complexification contains $H^{2,0}(X)$.

Remark 2.4 In light of Theorem 2.2 we shall say that the standard conjectures hold for a motive $m \in \text{Mot}$ if the category $\langle m \rangle_{\text{Mot}}$ is an abelian and semisimple neutral Tannakian category, thus equivalent to the category of representations of a reductive algebraic group $G_{\text{Mot}}(m)$. When this happens, the realization functor $R : \langle m \rangle_{\text{Mot}} \rightarrow \text{HS}$ is conservative [André 2004, Corollaire 5.1.3.3], i.e., a morphism in $\langle m \rangle_{\text{Mot}}$ is an isomorphism if and only if its realization is an isomorphism of \mathbb{Q} -Hodge structures.

Remark 2.5 Let $m \in \text{Mot}$ be a motive for which the standard conjectures hold. The torus $w : \mathbb{G}_{m, \mathbb{Q}} \hookrightarrow G_{\text{Mot}}(m)$ inducing the weight decomposition for the objects in $\langle m \rangle_{\text{Mot}}$ coincides with the weight torus $\mathbb{G}_{m, \mathbb{Q}} \subset \text{MT}(R(m))$ from Hodge theory. Assume that m is a motive of odd weight and let $\langle m \rangle_{\text{Mot}}^{\text{ev}}$ be the full subcategory of $\langle m \rangle_{\text{Mot}}$ of objects of even weight, which is again abelian, semisimple and neutral Tannakian. Analogously to Remark 2.1, the motives in $\langle m \rangle_{\text{Mot}}^{\text{ev}}$ correspond to the representations of $G_{\text{Mot}}(m)$ on which $w(-1)$ acts trivially. Therefore, the natural morphism of algebraic groups $G_{\text{Mot}}(m) \rightarrow G_{\text{Mot}}(\langle m \rangle_{\text{Mot}}^{\text{ev}})$ is an isogeny of degree 2. Moreover, by the same argument given in Remark 2.1, if $n \in \langle m \rangle_{\text{Mot}}$ is a motive of even weight such that the corresponding morphism $G_{\text{Mot}}(m) \rightarrow G_{\text{Mot}}(n)$ is an isogeny of degree 2, we then have the equality $\langle n \rangle_{\text{Mot}} = \langle m \rangle_{\text{Mot}}^{\text{ev}}$ of subcategories of Mot .

2.5 The standard conjectures are known to hold for curves, surfaces and abelian varieties; they hold for varieties X and Y if and only if they hold for $X \times Y$; see [Kleiman 1968]. The standard conjectures are not known to hold in general for hyper-Kähler varieties. They hold for the varieties of $\text{K3}^{[n]}$, OG10 and OG6-type built via moduli spaces of sheaves on K3 or abelian surfaces by [Bülles 2020; Floccari et al. 2021; Floccari 2023b], and for some hyper-Kähler varieties admitting a Lagrangian fibration [Ancona et al. 2025]. The standard conjectures have been fully proven for varieties of $\text{K3}^{[n]}$ -type by Charles and Markman [2013]. Recently, Foster [2024] proved that they hold in certain degrees for varieties of Kum^n -type; for later reference we state the following consequence of his results; see also [Floccari and Varesco 2025, Remark 3.2].

Theorem 2.6 [Foster 2024] *Let K be a variety of Kum^n -type. The Künneth projector $H^\bullet(K, \mathbb{Q}) \rightarrow H^2(K, \mathbb{Q})$ is algebraic. Moreover, the standard conjectures hold for the Künneth component $h^2(K) \in \text{Mot}$ of the motive of K .*

3 The Kuga–Satake construction

3.1 Let V be a \mathbb{Q} -vector space equipped with a nondegenerate symmetric bilinear form q . The Clifford algebra

$$C(V) := \bigoplus_{j \geq 0} V^{\otimes j} / \langle v \cdot v - q(v, v) \rangle_{v \in V}$$

is naturally $\mathbb{Z}/2\mathbb{Z}$ -graded: $C(V) = C^+(V) \oplus C^-(V)$. The Clifford algebra acts on itself via left and right multiplication. We fix a vector $v_0 \in V$ such that $q(v_0, v_0) \neq 0$. Setting $\mu(v)(a) := v \cdot a \cdot v_0$ we obtain an embedding

$$(1) \quad \mu : V \hookrightarrow \text{End}(C^+(v)).$$

Remark 3.1 We denote by $\text{End}_{C^+}(C^+(V))$ the algebra of linear endomorphisms of $C^+(V)$ commuting with the action of $C^+(V)$ on itself by right multiplication. There is a canonical isomorphism $C^+(V) = \text{End}_{C^+}(C^+(V))$, i.e., any $f \in \text{End}_{C^+}(C^+(V))$ is induced by left multiplication by some $\alpha_f \in C^+(V)$; see [Deligne 1972, §3] or [André 1996a, §4.2].

3.2 Assume now that (V, q) is a polarized \mathbb{Q} -Hodge structure of K3-type, by which we mean that V is an effective Hodge structure of weight 2 with Hodge numbers $(1, k, 1)$. Following [Kuga and Satake 1967] and [Deligne 1972], the Hodge structure on V determines an effective polarizable Hodge structure of weight 1 on $C^+(V)$. This thus defines an abelian variety $\text{KS}(V)$ up to isogeny, such that $H^1(\text{KS}(V), \mathbb{Q}) \cong C^+(V)$. With a suitable Tate twist to make V of weight 0, the embedding $\mu : V(1) \rightarrow \text{End}(H^1(\text{KS}(V), \mathbb{Q}))$ of (1) is a morphism of Hodge structures (for any choice of the vector $v_0 \in V$).

Remark 3.2 It is explained by Deligne [1972, §4] how the Hodge structure on $C^+(V)$ is induced through the action of the Clifford group $\text{CSpin}(V)$, which is a subgroup of $C^+(V)^*$ acting on $C^+(V)$ via left multiplication. This means that the Mumford–Tate group of $H^1(\text{KS}(V), \mathbb{Q})$ is contained in $\text{CSpin}(V) \subset \text{GL}(C^+(V))$. Moreover the morphism $\text{MT}(H^1(\text{KS}(V)), \mathbb{Q}) \rightarrow \text{MT}(V)$ induced by the embedding μ is an isogeny of degree 2, being the restriction of the double cover $\text{CSpin}(V) \rightarrow \text{GO}(V)$, where $\text{GO}(V)$ is the group of linear automorphisms of V preserving the form up to scalar.

3.3 Let S be a polarized K3 surface or hyper-Kähler variety, and consider the polarized Hodge structure of K3-type $V = H^2(S, \mathbb{Q})_{\text{prim}}$, the orthogonal to the given polarization, equipped with the restriction q of the form. We may also take for q the restriction of the Beauville–Bogomolov form, a natural pairing on the second cohomology of a hyper-Kähler manifold [Beauville 1983]. We define the Kuga–Satake variety $\text{KS}(S)$ of S as the abelian variety obtained from (V, q) via the Kuga–Satake construction. As above, we have an embedding of Hodge structures

$$\mu : H^2(S, \mathbb{Q})_{\text{prim}}(1) \hookrightarrow \text{End}(H^1(\text{KS}(S), \mathbb{Q}));$$

identifying $H^1(\text{KS}(S), \mathbb{Q})^\vee$ with $H^1(\text{KS}(S), \mathbb{Q})(1)$ by means of a polarization, we get the embedding of Hodge structures

$$(2) \quad \mu' : H^2(S, \mathbb{Q})_{\text{prim}} \hookrightarrow H^1(\text{KS}(S), \mathbb{Q})^{\otimes 2} \subset H^2(\text{KS}(S)^2, \mathbb{Q}).$$

Conjecture 3.3 *The morphism μ' is induced by an algebraic cycle on $S \times \text{KS}(S)^2$.*

This is the Kuga–Satake Hodge conjecture, which is of course a special case of the Hodge conjecture. It has been proven for many K3 surfaces of Picard rank at least 17 (see [Morrison 1985]), but it is widely open otherwise. In [Paranjape 1988] and [Ingalls et al. 2022] the conjecture is proven for two 4-dimensional families of K3 surfaces of general Picard rank 16; countably many more such families of K3 surfaces for which the conjecture holds are found in [Floccari 2024]. The paper [Bolognesi and Laterveer 2024] proves Conjecture 3.3 for a 9-dimensional family of K3 surfaces related to certain cubic fourfolds. The Kuga–Satake Hodge conjecture has been proven for all hyper-Kähler varieties of generalized Kummer type by Markman [2023] and Voisin [2022].

Remark 3.4 As a variant, we may define $\text{KS}'(S)$ as the Kuga–Satake variety built from the smaller Hodge structure of K3-type $H_{\text{tr}}^2(S, \mathbb{Q})$; then the Kuga–Satake variety $\text{KS}(S)$ built from $H^2(S, \mathbb{Q})_{\text{prim}}$ is isogenous to a power of $\text{KS}'(S)$; see [Huybrechts 2016, Chapter 4, Example 2.4]. Moreover, Conjecture 3.3 is equivalent to the statement that the embedding $\mu' : H_{\text{tr}}^2(S, \mathbb{Q}) \hookrightarrow H^1(\text{KS}'(S), \mathbb{Q})^{\otimes 2} \subset H^2(\text{KS}'(S)^2, \mathbb{Q})$ analogous to (2) is induced by an algebraic cycle.

3.4 We are now ready to prove the main result of this section.

Theorem 3.5 *Let S be a projective K3 surface. Assume that the Kuga–Satake Hodge conjecture holds for S . Then the Hodge conjecture holds for all powers of S if and only if it holds for all powers of its Kuga–Satake variety $\text{KS}(S)$.*

We will deduce this result from the following proposition. Recall (Section 2.5) that the standard conjectures hold for S as well as for $\text{KS}(S)$.

Proposition 3.6 *Let S be a projective K3 surface. Assume that the Kuga–Satake Hodge conjecture holds for S . Then, the motive $h^2(S)$ belongs to the subcategory $\langle h^1(\text{KS}(S)) \rangle_{\text{Mot}}$ of the category Mot of homological motives and the induced morphism of algebraic groups $G_{\text{Mot}}(h^1(\text{KS}(S))) \rightarrow G_{\text{Mot}}(h^2(S))$ is an isogeny of degree 2.*

Proof By Theorem 2.2, the subcategory $\langle h(\text{KS}(S) \times S) \rangle_{\text{Mot}}$ of Mot is an abelian and semisimple neutral Tannakian category. Hence, if Conjecture 3.3 holds for S , then the motive $h^2(S)$ is a direct summand of $h^2(\text{KS}(S)^2)$, and so it belongs to $\langle h^1(\text{KS}(S)) \rangle_{\text{Mot}}$. We obtain the commutative diagram

$$\begin{array}{ccc} G_{\text{Mot}}(h^1(\text{KS}(S))) & \xrightarrow{\psi} & G_{\text{Mot}}(h^2(S)) \\ i_{\text{KS}} \uparrow & & \uparrow i_S \\ \text{MT}(H^1(\text{KS}(S), \mathbb{Q})) & \xrightarrow{\phi} & \text{MT}(H^2(S, \mathbb{Q})) \end{array}$$

We know from Remark 3.2 that the morphism ϕ is an isogeny of degree 2; we will show that $\ker(\psi) = \iota_{\text{KS}}(\ker(\phi))$, which implies that ψ is also an isogeny of degree 2.

To ease notation, we let $A := \text{KS}(S)$ and $V := H^2(S, \mathbb{Q})_{\text{prim}}$; we denote by q the polarization form on V . By Remark 3.2, the action of $C^+(V)$ on $H^1(A, \mathbb{Q})$ via right multiplication commutes with $\text{MT}(H^1(A, \mathbb{Q}))$, i.e., $C^+(V)$ acts via Hodge endomorphisms. By Lefschetz (1,1)-theorem, any such endomorphism is algebraic. Therefore, $G_{\text{Mot}}(h^1(A)) \subset \text{GL}(H^1(A, \mathbb{Q}))$ commutes with the $C^+(V)$ -action given by right multiplication. By Remark 3.1, we can thus identify $G_{\text{Mot}}(h^1(A))$ with a subgroup of $C^+(V)^*$, the group of units in the Clifford algebra, acting on $C^+(V)$ via left multiplication.

Let $g \in \ker(\psi : G_{\text{Mot}}(h^1(A)) \rightarrow G_{\text{Mot}}(h^2(S)))$; then $g : H^1(A, \mathbb{Q}) \rightarrow H^1(A, \mathbb{Q})$ is left multiplication by some $\alpha_g \in C^+(V)^*$. We claim that α_g is a central element of $C^+(V)$.

By assumption, the embedding $\mu : V(1) \hookrightarrow \text{End}(H^1(A, \mathbb{Q}))$ is induced by an algebraic cycle. As $g \in \ker(\psi)$, it follows that g commutes with the image of μ . Recall (Section 3.1) that the definition of μ depends on a vector $v_0 \in V$ with $q(v_0, v_0) \neq 0$ such that $\mu(v)(\beta) := v \cdot \beta \cdot v_0$. In the Clifford algebra we have $v_0^{2k} = q(v_0, v_0)^k \cdot e$, where e is the identity element, and any $\beta \in C^+(V)$ is a linear combination of products $v_1 \cdots v_{2k}$ of an even number of vectors in V . As g commutes with $\mu(v)$ for any $v \in V$, we find

$$\begin{aligned} \alpha_g(v_1 v_2 \cdots v_{2k}) &= \frac{1}{q(v_0, v_0)^k} \cdot \alpha_g \cdot (v_1 v_2 \cdots v_{2k}) \cdot v_0^{2k} \\ &= \frac{1}{q(v_0, v_0)^k} \cdot (g \circ \mu(v_1) \circ \mu(v_2) \circ \cdots \circ \mu(v_{2k}))(e) \\ &= \frac{1}{q(v_0, v_0)^k} \cdot (\mu(v_1) \circ \mu(v_2) \circ \cdots \circ \mu(v_{2k}) \circ g)(e) = (v_1 v_2 \cdots v_{2k}) \alpha_g. \end{aligned}$$

Therefore, $\ker(\psi)$ is contained in the center of $C^+(V)$. But $V = H^2(S, \mathbb{Q})_{\text{prim}}$ has odd dimension 21, and in such case $C^+(V)$ is a central simple algebra; see [Deligne 1972, page 216]. The units in the center of $C^+(V)$ thus form a torus $\mathbb{G}_{m, \mathbb{Q}}$ acting on $C^+(V)$ via scalar multiplication; this torus induces the weight filtration on objects of $\langle H^1(A, \mathbb{Q}) \rangle_{\text{HS}}$ and so it is contained in $\text{MT}(H^1(A, \mathbb{Q}))$. Hence, $\ker(\psi)$ is contained in $i_{\text{KS}}(\ker(\phi))$. □

Proof of Theorem 3.5 One direction is immediate: as Conjecture 3.3 holds for S by assumption, we have $h^2(S) \in \langle h^1(\text{KS}(S)) \rangle_{\text{Mot}}$, and the Hodge conjecture for all powers of $\text{KS}(S)$ clearly implies the Hodge conjecture for all powers of S .

Let us prove the converse implication. The Hodge conjecture for all powers of $\text{KS}(S)$ predicts that the realization functor $R : \text{Mot} \rightarrow \text{HS}$ induces a surjection

$$(3) \quad \text{Hom}_{\text{Mot}}(\mathbb{Q}(-j), h^{2j}(\text{KS}(S)^k)) \rightarrow \text{Hom}_{\text{HS}}(\mathbb{Q}(-j), H^{2j}(\text{KS}(S)^k, \mathbb{Q}))$$

for any k and j . Only motives and Hodge structures of even weight are involved in (3), and so these Hom-spaces are actually taken in the categories $\langle h^1(\text{KS}(S)) \rangle_{\text{Mot}}^{\text{ev}}$ and $\langle H^1(\text{KS}(S), \mathbb{Q}) \rangle_{\text{HS}}^{\text{ev}}$, respectively. Proposition 3.6 and Remark 2.5 yield the equality $\langle h^1(\text{KS}(S)) \rangle_{\text{Mot}}^{\text{ev}} = \langle h^2(S) \rangle_{\text{Mot}}$ of subcategories of Mot ;

in particular, $h^{2j}(\text{KS}(S)^k)$ belongs to $\langle h^2(S) \rangle_{\text{Mot}}$ for any j and k . Therefore, the map in (3) is the map

$$\text{Hom}_{\langle h^2(S) \rangle_{\text{Mot}}}(\mathbb{Q}(-j), h^{2j}(\text{KS}(S)^k)) \rightarrow \text{Hom}_{\langle H^2(S, \mathbb{Q}) \rangle_{\text{HS}}}(\mathbb{Q}(-j), H^{2j}(\text{KS}(S)^k, \mathbb{Q}))$$

given by the realization functor R . If the Hodge conjecture holds for all powers of S , then $R : \langle h^2(S) \rangle_{\text{Mot}} \rightarrow \langle H^2(S, \mathbb{Q}) \rangle_{\text{HS}}$ is full, and the maps in (3) are surjective. \square

3.5 For later use, we recall the following functoriality property of the Kuga–Satake construction, following Varesco [2023]. Let X and Y be hyper-Kähler varieties, not necessarily deformation equivalent and possibly of different dimensions. Assume that $\phi : H_{\text{tr}}^2(X, \mathbb{Q}) \rightarrow H_{\text{tr}}^2(Y, \mathbb{Q})$ is a rational Hodge similitude, i.e., ϕ is an isomorphism of \mathbb{Q} -Hodge structures which multiplies the Beauville–Bogomolov form on the left-hand side by some nonzero $k \in \mathbb{Q}$. As in Remark 3.4, let $\text{KS}'(X)$ and $\text{KS}'(Y)$ be the Kuga–Satake varieties constructed from $H_{\text{tr}}^2(X, \mathbb{Q})$ and $H_{\text{tr}}^2(Y, \mathbb{Q})$, respectively. Varesco [2023, Proposition 3.1] then proved that there exists an isogeny $\Psi : \text{KS}'(X) \rightarrow \text{KS}'(Y)$ of abelian varieties such that the diagram

$$\begin{array}{ccc} H_{\text{tr}}^2(X, \mathbb{Q}) & \xrightarrow{\phi} & H_{\text{tr}}^2(Y, \mathbb{Q}) \\ \downarrow \mu'_X & & \downarrow \mu'_Y \\ H^1(\text{KS}'(X), \mathbb{Q})^{\otimes 2} & \xrightarrow{\Psi_*} & H^1(\text{KS}'(Y), \mathbb{Q})^{\otimes 2} \end{array}$$

commutes, where μ'_X and μ'_Y are the respective Kuga–Satake correspondences and the bottom arrow is the isomorphism induced by the isogeny Ψ .

Remark 3.7 With notation as above, assume that $\phi : H_{\text{tr}}^2(X, \mathbb{Q}) \rightarrow H_{\text{tr}}^2(Y, \mathbb{Q})$ is induced by an algebraic cycle on $X \times Y$. Then, Conjecture 3.3 for Y implies Conjecture 3.3 for X . Indeed, the above diagram yields $\mu'_X = (\Psi_*)^{-1} \circ \mu'_Y \circ \phi$; note that $(\Psi_*)^{-1} : H^1(\text{KS}'(Y), \mathbb{Q})^{\otimes 2} \rightarrow H^1(\text{KS}'(X), \mathbb{Q})^{\otimes 2}$ is induced by an algebraic cycle, so that the Kuga–Satake correspondence μ'_X is algebraic if μ'_Y is so.

4 Some $\text{K3}^{[m]}$ -type submanifolds of generalized Kummer varieties

4.1 Let A be an abelian surface. We denote by $K^n(A)$ the generalized Kummer variety on A of dimension $2n$ introduced in [Beauville 1983]. It is a hyper-Kähler variety constructed as follows: consider the Hilbert scheme $A^{[n+1]}$ of 0-dimensional subschemes of A of length $n + 1$, which is a crepant resolution $\nu : A^{[n+1]} \rightarrow A^{(n+1)}$ of the symmetric power. Denoting by $\Sigma : A^{(n+1)} \rightarrow A$ the summation map $\Sigma(a_1, \dots, a_{n+1}) = \sum_{i=1}^{n+1} a_i$, the variety $K^n(A)$ is defined as the fiber of $\Sigma \circ \nu$ over $0 \in A$. We let $A_0^{(n+1)} \subset A^{(n+1)}$ be the fiber of Σ over 0 ; the Hilbert–Chow morphism ν restricts to a crepant resolution $\nu : K^n(A) \rightarrow A_0^{(n+1)}$. For $n = 1$, the construction yields nothing but the Kummer K3 surface $\text{Km}(A)$ associated to A , that is, the minimal resolution of the quotient $A/\pm 1$.

We will study certain natural subvarieties of $K^n(A)$, $n \geq 2$. Consider the morphisms

$$(4) \quad \begin{aligned} f : A^m &\rightarrow A_0^{(2m)}, & (a_1, \dots, a_m) &\mapsto (a_1, -a_1, \dots, a_m, -a_m), \\ f' : A^m &\rightarrow A_0^{(2m+1)}, & (a_1, \dots, a_m) &\mapsto (a_1, -a_1, \dots, a_m, -a_m, 0). \end{aligned}$$

- Definition 4.1** (i) If n is odd, $n = 2m - 1$, we let $\iota : W \hookrightarrow K^n(A)$ be the strict transform of the image of $f : A^m \rightarrow A_0^{(2m)}$ under $\nu : K^n(A) \rightarrow A_0^{(2m)}$.
- (ii) If n is even, $n = 2m$, we let $\iota : W \hookrightarrow K^n(A)$ be the strict transform of the image of $f' : A^m \rightarrow A_0^{(2m+1)}$ under $\nu : K^n(A) \rightarrow A_0^{(2m+1)}$.

Notice that the automorphism -1 of A naturally acts on $K^n(A)$ for all n , as the natural action of -1 on $A^{[n+1]}$ stabilizes $K^n(A)$. It is known that -1 acts trivially on $H^2(K^n(A), \mathbb{Z})$; see [Boissière et al. 2011]. The map f (resp. f') factors through an embedding of $(A/\pm 1)^{(m)}$ into $A_0^{(2m)}$ (resp. into $A_0^{(2m+1)}$); a natural crepant resolution of $(A/\pm 1)^{(m)}$ is given by $\text{Km}(A)^{[m]}$, the Hilbert scheme of points on the Kummer K3 surface $\text{Km}(A)$.

Lemma 4.2 For any $n \geq 2$, the subvariety $W \subset K^n(A)$ is the unique component of maximal dimension of the fixed locus of -1 acting on $K^n(A)$. Moreover, if $n = 2m$ (resp. $n = 2m - 1$), then f' (resp. f) induces an embedding $\iota : \text{Km}(A)^{[m]} \hookrightarrow K^{2m}(A)$ (resp. $\iota : \text{Km}(A)^{[m]} \hookrightarrow K^{2m-1}(A)$) with image W .

Proof The lemma is proven by Kamenova, Mongardi and Oblomkov [Kamenova et al. 2022]. They show in [loc. cit., Theorem 1.3] that the fixed locus of -1 on $K^n(A)$ is the union of components which are varieties of $\text{K3}^{[k]}$ -type or points, and that there exists a unique component of maximal dimension $2m$ for $n = 2m - 1$ or $n = 2m$. As our $W \subset K^n(A)$ is evidently fixed by -1 , it must be the component of maximal dimension of the fixed locus. The local study done in [loc. cit., Appendix B] shows that the birational map $\text{Km}(A)^{[m]} \dashrightarrow W$ induced by f' if $n = 2m$ or f if $n = 2m - 1$ extends to an isomorphism. \square

The main result of this section is the following.

Proposition 4.3 Let $\iota : W \hookrightarrow K^n(A)$ denote the embedding of Definition 4.1. Then the pullback along ι induces a primitive embedding

$$\iota^* : H^2(K^n(A), \mathbb{Z})(2) \hookrightarrow H^2(W, \mathbb{Z})$$

of lattices and Hodge structures, where on the left-hand side the form is multiplied by a factor 2.

The cases $n = 2$ and $n = 3$ may be deduced from [Hassett and Tschinkel 2013] and [Floccari 2023a], respectively.

4.2 Let us fix some notation for the cohomology groups of the varieties we shall consider. By [Beauville 1983], for $n \geq 2$ there is a canonical primitive embedding of $H^2(A, \mathbb{Z})$ into the second cohomology of $K^n(A)$. We have

$$(5) \quad H^2(K^n(A), \mathbb{Z}) = H^2(A, \mathbb{Z}) \oplus \mathbb{Z} \cdot \xi,$$

where ξ is half the class of the divisor $E \subset K^n(A)$ parametrizing nonreduced subschemes, which is the exceptional divisor of the Hilbert–Chow resolution. The class ξ has square $-2(n + 1)$ with respect to the Beauville–Bogomolov form; as a lattice, $H^2(K^n(A), \mathbb{Z})$ is thus isometric to $U^{\oplus 3} \oplus \langle -2(n + 1) \rangle$.

The Kummer surface $\text{Km}(A)$ associated to A contains 16 exceptional curves C_τ , parametrized by the 2-torsion points $\tau \in A_2$. The pushforward along the rational map $\pi : A \dashrightarrow \text{Km}(A)$ induces a primitive

embedding of $H^2(A, \mathbb{Z})(2)$ into $H^2(\text{Km}(A), \mathbb{Z})$ whose orthogonal is the Kummer lattice (see [Nikulin 1975]), which is the saturation of the sublattice generated by the 16 pairwise orthogonal (-2) -classes $[C_\tau]$. Hence, $H^2(\text{Km}(A), \mathbb{Z})$ contains with finite index the sublattice $H^2(A, \mathbb{Z})(2) \oplus \langle [C_\tau] \rangle_{\tau \in A_2}$.

For the cohomology of the Hilbert scheme $\text{Km}(A)^{[m]}$, $m \geq 2$, we have by [Beauville 1983] a canonical primitive embedding of $H^2(\text{Km}(A), \mathbb{Z})$ into $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$, and

$$(6) \quad H^2(\text{Km}(A)^{[m]}, \mathbb{Z}) = H^2(\text{Km}(A), \mathbb{Z}) \oplus \mathbb{Z} \cdot \delta,$$

where δ is half the class of the divisor $D \subset \text{Km}(A)^{[m]}$ of nonreduced subschemes; the Beauville–Bogomolov square of δ is $-2(m - 1)$. Under the embedding of $H^2(\text{Km}(A), \mathbb{Z})$ into $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$, the class $[C_\tau]$ of the exceptional curve over the node given by $\tau \in A_2$ corresponds to the class $[R_\tau]$ of the divisor R_τ of subschemes whose support intersects C_τ . Therefore, $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$ contains with finite index a sublattice $H^2(A, \mathbb{Z})(2) \oplus \langle [R_\tau] \rangle_{\tau \in A_2} \oplus \langle \delta \rangle$ and the saturation of $\langle [R_\tau] \rangle_{\tau \in A_2}$ is the Kummer lattice.

4.3 The next lemma is the first step towards Proposition 4.3.

Lemma 4.4 *Consider the embedding $\iota : W \hookrightarrow K^n(A)$ for $n \geq 2$.*

(i) *For $n = 2$, the pullback along $\iota : \text{Km}(A) \hookrightarrow K^2(A)$ yields*

$$\iota^*(\xi) = [C_0] + \frac{1}{2} \sum_{\tau \in A_2} [C_\tau] \quad \text{in } H^2(\text{Km}(A), \mathbb{Z}).$$

(ii) *For $n = 2m - 1$, the pullback along $\iota : \text{Km}(A)^{[m]} \hookrightarrow K^{2m-1}(A)$ yields*

$$\iota^*(\xi) = 2\delta + \frac{1}{2} \sum_{\tau \in A_2} [R_\tau] \quad \text{in } H^2(\text{Km}(A)^{[m]}, \mathbb{Z}).$$

(iii) *For $n = 2m \geq 4$, the pullback along $\iota : \text{Km}(A)^{[m]} \hookrightarrow K^{2m}(A)$ yields*

$$\iota^*(\xi) = 2\delta + [R_0] + \frac{1}{2} \sum_{\tau \in A_2} [R_\tau] \quad \text{in } H^2(\text{Km}(A)^{[m]}, \mathbb{Z}).$$

Notice that $\frac{1}{2} \sum_{\tau} [C_\tau]$ is indeed an integral class; see [Huybrechts 2016, Chapter 14, §3.3]. The proof of the lemma will be reduced to some local computations.

Example 4.5 Consider the involution $-1 : \mathbb{C}^2 \rightarrow \mathbb{C}^2$. The quotient $\mathbb{C}^2/\pm 1$ has a nodal singularity at the image $\bar{0}$ of the origin, which is resolved by the blowup $\text{Bl}_{\bar{0}}(\mathbb{C}^2/\pm 1) \rightarrow \mathbb{C}^2$. We denote by $C \subset \text{Bl}_{\bar{0}}(\mathbb{C}^2/\pm 1)$ the exceptional curve.

(a) The graph map $\hat{\phi} : \mathbb{C}^2 \rightarrow (\mathbb{C}^2)^2$ given by $y \mapsto (y, -y)$ yields a Cartesian diagram

$$(7) \quad \begin{array}{ccc} \text{Bl}_{\bar{0}}(\mathbb{C}^2/\pm 1) & \xleftarrow{\phi} & \mathbb{C}^{[2]} \\ \downarrow \nu & & \downarrow \nu \\ (\mathbb{C}^2/\pm 1) & \xleftarrow{\bar{\phi}} & (\mathbb{C}^2)^{(2)} \end{array}$$

By [Kamenova et al. 2022, Lemma 3.13], the image of ϕ is the fixed locus of -1 acting on $\mathbb{C}^{[2]}$. Let $F \subset \mathbb{C}^{[2]}$ be the divisor of nonreduced subschemes. Then, as divisors, $\phi^*(F) = C$; indeed, the image of ϕ and F intersect transversely in the curve $\phi(C)$.

(b) Consider now the map $\hat{\phi} : \mathbb{C}^2 \rightarrow (\mathbb{C}^2)^3$ given by $y \mapsto (y, -y, 0)$. It induces the Cartesian commutative diagram

$$(8) \quad \begin{array}{ccc} \text{Bl}_0(\mathbb{C}^2/\pm 1) & \xleftarrow{\phi} & \mathbb{C}^{[3]} \\ \downarrow \nu & & \downarrow \nu \\ (\mathbb{C}^2/\pm 1) & \xleftarrow{\bar{\phi}} & (\mathbb{C}^2)^{(3)} \end{array}$$

By [Kamenova et al. 2022, Lemma 3.13], the fixed locus of -1 acting on $\mathbb{C}^{[3]}$ consists of the image of ϕ and an isolated fixed point. Let $F \subset \mathbb{C}^{[3]}$ be the divisor of nonreduced subschemes. Then we claim that, as divisors, we have $\phi^*(F) = 3C$.

To see this, let $B \subset (\mathbb{C}^2)^{[3]}$ be the Briançon variety at 0, that is, the subvariety $B := \nu^{-1}((0, 0, 0))$ that parametrizes subschemes fully supported at $0 \in \mathbb{C}^2$. Let $j : B \hookrightarrow \mathbb{C}^{[3]}$ denote the embedding. The set-theoretic intersection of F and the image of ϕ is $\phi(C)$, which is contained in B . In our case, by [Briançon 1977, IV.2], the Briançon variety B is isomorphic to the weighted projective plane $\mathbb{P}(1, 1, 3)$; following [Hassett and Tschinkel 2013, §4], there exists a closed embedding $\mathbb{P}(1, 1, 3) \hookrightarrow \mathbb{P}^4$ realizing B as the cone over the twisted cubic curve and the restriction $j^*(\mathcal{O}_{(\mathbb{C}^2)^{[3]}}(F))$ equals $\mathcal{O}_B(-2H)$, where we denote by $\mathcal{O}_B(H)$ the line bundle which induces the embedding $B \hookrightarrow \mathbb{P}^4$. Moreover, $-1 : \mathbb{C}^{[3]} \rightarrow \mathbb{C}^{[3]}$ acts on B with fixed locus the vertex of the cone and the twisted cubic curve at the base, which is thus identified with $\phi(C)$. The embedding $\phi|_C : C \hookrightarrow (\mathbb{C}^2)^{[3]}$ factors as the composition of embeddings $C \xrightarrow{\phi'} B \xrightarrow{j} (\mathbb{C}^2)^{[3]}$. Therefore,

$$(9) \quad \deg(\mathcal{O}_{(\mathbb{C}^2)^{[3]}}(F)|_{\phi(C)}) = \deg(\mathcal{O}_B(-2H)|_{\phi'(C)}) = -6,$$

as H is a hyperplane divisor in \mathbb{P}^4 and the twisted cubic curve has degree 3. If k denotes the multiplicity of $\phi^*(F)$ at C , we have $-2k = \deg(\mathcal{O}_{(\mathbb{C}^2)^{[3]}}(F)|_{\phi(C)}) = -6$, and we conclude that $k = 3$.

To justify the last claim, consider a smooth and projective surface Y with an involution θ with isolated fixed points. Each fixed point gives a node on the quotient Y/θ ; the blowup of these nodes is a smooth and projective surface X . We choose a fixed point $p \in Y$, and let $C_p \subset X$ be the exceptional curve over the corresponding node. We then have the Cartesian diagram of projective varieties

$$\begin{array}{ccc} X & \xleftarrow{\phi} & Y^{[3]} \\ \downarrow \nu & & \downarrow \nu \\ Y/\theta & \xleftarrow{\bar{\phi}} & Y^{(3)} \end{array}$$

induced by the map $Y \rightarrow Y^3$ sending y to $(y, \theta(y), p)$. We can find an analytic neighborhood $U \subset Y$ of p stable under θ which is identified with a neighborhood V of the origin in \mathbb{C}^2 in such a way that the

action of θ on U corresponds to the action of -1 on V . Thus, denoting by F the exceptional divisor of $Y^{[3]} \rightarrow Y$ and by k the multiplicity of $\phi^*(F)$ at C_p , we have to show that $-2k = \deg(\mathcal{O}_{Y^{[3]}}(F)|_{\phi(C_p)})$. But $\phi^*(F)$ is supported over the exceptional curves C_q of $X \rightarrow Y/\theta$, which are pairwise orthogonal (-2) -curves. Therefore, by the projection formula, we have

$$-2k = \int_X [C_p] \cdot [\phi^*(F)] = \int_{Y^{[3]}} [\phi(C_p)] \cdot [F] = \deg(\mathcal{O}_{Y^{[3]}}(F)|_{\phi(C_p)}).$$

Proof of Lemma 4.4 The exceptional divisor $E \subset K^n(A)$ is the restriction to $K^n(A) \subset A^{[n+1]}$ of the exceptional divisor of the Hilbert–Chow resolution $A^{[n+1]} \rightarrow A^{(n+1)}$, which we still denote by E . We consider the Cartesian diagram

$$(10) \quad \begin{array}{ccc} \mathrm{Km}(A)^{[m]} & \xleftarrow{\iota} & A^{[n+1]} \\ \downarrow & & \downarrow \nu \\ (A/\pm 1)^{(m)} & \xleftarrow{\bar{\iota}} & A^{(n+1)} \end{array}$$

where $n = 2m$ or $n = 2m - 1$: in the first case, $\bar{\iota}(\bar{a}_1, \dots, \bar{a}_m) = (a_1, -a_1, \dots, a_m, -a_m, 0)$, while in the second $\bar{\iota}(\bar{a}_1, \dots, \bar{a}_m) = (a_1, -a_1, \dots, a_m, -a_m)$. It is immediate to check that, if $m \geq 2$, the restriction $\iota^*(E)$ is supported over the union of the Hilbert–Chow divisor $D \subset \mathrm{Km}(A)^{[m]}$ and the 16 divisors $R_\tau \subset \mathrm{Km}(A)^{[m]}$, for $\tau \in A_2$, while if $m = 1$ then $\iota^*(E)$ is supported over the 16 exceptional curves $C_\tau \subset \mathrm{Km}(A)$, $\tau \in A_2$, which we will also denote by R_τ to ease notation. We now have the following cases to consider.

Case 1: Let $\tau \in A_2$ be a 2-torsion point; if n is even, assume further that $\tau \neq 0$. We claim that then $\iota^*(E)$ has multiplicity 1 along the divisor R_τ .

Indeed, the image in $(A/\pm 1)^{(m)}$ of a general subscheme $\zeta \in R_\tau$ is a point of the form $(\bar{\tau}, \bar{a}_2, \dots, \bar{a}_m)$ with the \bar{a}_i pairwise distinct smooth points of $A/\pm 1$, while $\iota(\zeta)$ is a subscheme in $A^{[n+1]}$ which is supported over the point $(\tau, \tau, a_2, -a_2, \dots, a_m, -a_m)$ of $A^{(n+1)}$ if $n = 2m - 1$ is odd, or over $(\tau, \tau, a_2, -a_2, \dots, a_m, -a_m, 0)$ if $n = 2m$ is even. Let \mathcal{U} be an analytic neighborhood of $\iota(\zeta)$ in $A^{[n+1]}$ which is stable under -1 , and let \mathcal{V} be the neighborhood $\iota^{-1}(\mathcal{U})$ of ζ in $\mathrm{Km}(A)^{[m]}$. Assume first that $n = 2m - 1$, for some $m \geq 2$. Choosing \mathcal{U} small enough, diagram (10) restricts to

$$\begin{array}{ccc} \mathcal{V} \cong (\mathrm{Bl}_{\bar{\tau}}(U_\tau/\pm 1) \times \prod_{j=2}^m U_j) & \xleftarrow{\iota} & \mathcal{U} \cong (U_\tau^{[2]} \times \prod_{j=2}^m (U_j \times U_j^-)) \\ \downarrow & & \downarrow \nu \\ (U_\tau/\pm 1) \times \prod_{j=2}^m U_j & \xleftarrow{\bar{\iota}} & U_\tau^{(2)} \times \prod_{j=2}^m (U_j \times U_j^-) \end{array}$$

where: $U_\tau \subset A$ is a neighborhood of τ which is stable under -1 ; $U_j \subset A$ is a neighborhood of a_j and $U_j^- := -1(U_j)$ for $j = 2, \dots, m$; the morphism ν is the product of the Hilbert–Chow resolution $U_\tau^{[2]} \rightarrow U_\tau^{(2)}$ with the identity of each factor $(U_j \times U_j^-)$; the morphism $\bar{\iota}$ is given by $\bar{\iota}(\bar{\alpha}, \alpha_2, \dots, \alpha_m) = ((\alpha, -\alpha), \alpha_2, -\alpha_2, \dots, \alpha_m, -\alpha_m)$. If instead $n = 2m$ is even, choosing \mathcal{U} small enough, diagram (10)

restricts to

$$\begin{CD} \mathcal{V} \cong (\mathrm{Bl}_{\bar{\tau}}(U_{\tau}/\pm 1) \times \prod_{j=2}^m U_j) @<\iota<< \mathcal{U} \cong (U_{\tau}^{[2]} \times \prod_{j=2}^m (U_j \times U_j^-) \times U_0) \\ @VVV @VVV \\ (U_{\tau}/\pm 1) \times \prod_{j=2}^m U_j @<\bar{\iota}<< U_{\tau}^{(2)} \times \prod_{j=2}^m (U_j \times U_j^-) \times U_0 \end{CD}$$

where: $U_{\tau} \subset A$ is a neighborhood of τ stable under -1 ; $U_j \subset A$ is a neighborhood of a_j and $U_j^- := -1(U_j)$ for $j = 2, \dots, m$; $U_0 \subset A$ is a neighborhood of 0 ; the morphism ν is the product of the Hilbert–Chow resolution $U_{\tau}^{[2]} \rightarrow U_{\tau}^{(2)}$ with the identity of each factor $(U_j \times U_j^-)$ and of U_0 ; we have $\bar{\iota}(\bar{\alpha}, \alpha_2, \dots, \alpha_m) = ((\alpha, -\alpha), \alpha_2, -\alpha_2, \dots, \alpha_m, -\alpha_m, 0)$.

In both cases, we have $E|_{\mathcal{U}} = \mathrm{pr}_1^*(F)$, where $\mathrm{pr}_1 : \mathcal{U} \rightarrow U_{\tau}^{[2]}$ is the projection onto the first factor and $F \subset U_{\tau}^{[2]}$ is the divisor of nonreduced subschemes. Moreover, the divisor $R_{\tau|_{\mathcal{V}}}$ equals $\tilde{\mathrm{pr}}_1^*(C)$, where $\tilde{\mathrm{pr}}_1 : \mathcal{V} \rightarrow \mathrm{Bl}_{\bar{\tau}}(U_{\tau}/\pm 1)$ is the projection and $C \subset \mathrm{Bl}_{\bar{\tau}}(U_{\tau}/\pm 1)$ is the exceptional curve. We thus have the commutative diagram

$$\begin{CD} \mathcal{V} @<\iota<< \mathcal{U} \\ @V\tilde{\mathrm{pr}}_1VV @VV\mathrm{pr}_1V \\ \mathrm{Bl}_{\bar{\tau}}(U_{\tau}/\pm 1) @<\phi<< U_{\tau}^{[2]} \end{CD}$$

where ϕ coincides with the restriction of the map $\mathrm{Bl}_0(\mathbb{C}^2/\pm 1) \rightarrow (\mathbb{C}^2)^{[2]}$ of Example 4.5(a) in a neighborhood of the exceptional curve $C \subset \mathrm{Bl}_0(\mathbb{C}^2 \pm 1)$. We deduce that, as divisors, we have

$$\iota^*(E|_{\mathcal{U}}) = (\iota \circ \mathrm{pr}_1)^*(F) = (\tilde{\mathrm{pr}}_1^*(\phi^*(F))) = \tilde{\mathrm{pr}}_1^*(C) = R_{\tau|_{\mathcal{V}}}.$$

Hence, $\iota^*(E)$ has multiplicity 1 along the divisor $R_{\tau} \subset \mathrm{Km}(A)^{[m]}$.

Case 2: Let $n = 2m$ be even and consider the divisor R_0 of $\mathrm{Km}(A)^{[m]}$. We claim that $\iota^*(E)$ has multiplicity 3 along the divisor R_0 .

Indeed, the image in $(A/\pm 1)^{(m)}$ of a general subscheme $\zeta \in R_0$ is a point of the form $(\bar{0}, \bar{a}_2, \dots, \bar{a}_m)$ with the \bar{a}_i pairwise distinct smooth points of $A/\pm 1$, while $\iota(\zeta)$ is a subscheme in $A^{[n+1]}$ supported over the point $(0, 0, a_2, -a_2, \dots, a_m, -a_m, 0)$ of $A^{(n+1)}$. Let \mathcal{U} be an analytic neighborhood of $\iota(\zeta)$ in $A^{[n+1]}$ such that $-1(\mathcal{U}) = \mathcal{U}$, and let \mathcal{V} be the analytic neighborhood $\iota^{-1}(\mathcal{U})$ of ζ in $\mathrm{Km}(A)^{[m]}$. Choosing \mathcal{U} small enough, diagram (10) restricts to

$$\begin{CD} \mathcal{V} \cong (\mathrm{Bl}_0(U_0/\pm 1) \times \prod_{j=2}^m U_j) @<\iota<< \mathcal{U} \cong (U_0^{[3]} \times \prod_{j=2}^m (U_j \times U_j^-)) \\ @VVV @VVV \\ (U_0/\pm 1) \times \prod_{j=2}^m U_j @<\bar{\iota}<< U_0^{(3)} \times \prod_{j=2}^m (U_j \times U_j^-) \end{CD}$$

where: $U_0 \subset A$ is a neighborhood of 0 stable under -1 ; $U_j \subset A$ is a neighborhood of a_j and $U_j^- := -1(U_j)$, for $j = 2, \dots, m$; the morphism ν is the product of the Hilbert–Chow resolution $U_0^{[3]} \rightarrow U_0^{(3)}$ with the identity of each factor $(U_j \times U_j^-)$; we have $\bar{\iota}(\bar{\alpha}, \alpha_2, \dots, \alpha_m) = ((\alpha, -\alpha, 0), \alpha_2, -\alpha_2, \dots, \alpha_m, -\alpha_m)$.

Then $E|_{\mathcal{U}} = \text{pr}_1^*(F)$, where $\text{pr}_1 : \mathcal{U} \rightarrow U_0^{[3]}$ is the projection and $F \subset U_0^{[3]}$ is the divisor of nonreduced subschemes. Moreover, $R_0|_{\mathcal{V}} = \widetilde{\text{pr}}_1^*(C)$, where $\widetilde{\text{pr}}_1 : \mathcal{V} \rightarrow \text{Bl}_0(U_0/\pm 1)$ is the projection and $C \subset \text{Bl}_0(U_0/\pm 1)$ is the exceptional curve. We have the commutative diagram

$$\begin{array}{ccc} \mathcal{V} & \xrightarrow{\iota} & \mathcal{U} \\ \downarrow \widetilde{\text{pr}}_1 & & \downarrow \text{pr}_1 \\ \text{Bl}_0(U_0/\pm 1) & \xrightarrow{\phi} & U_0^{[3]} \end{array}$$

where ϕ coincides with the restriction of the map $\text{Bl}_0(\mathbb{C}^2/\pm 1) \rightarrow (\mathbb{C}^2)^{[3]}$ of Example 4.5(b) in a neighborhood of $C \subset \text{Bl}_0(\mathbb{C}^2/\pm 1)$, and we deduce that

$$\iota^*(E|_{\mathcal{U}}) = (\iota \circ \text{pr}_1)^*(F) = (\widetilde{\text{pr}}_1^*(\phi^*(F))) = \widetilde{\text{pr}}_1^*(3C) = 3R_0|_{\mathcal{V}}$$

as divisors. Hence, $\iota^*(E)$ has multiplicity 3 along the divisor $R_0 \subset \text{Km}(A)^{[m]}$.

Case 3: Assume now that $m \geq 2$ and consider the Hilbert–Chow divisor $D \subset \text{Km}(A)^{[m]}$. We claim that $\iota^*(E)$ has multiplicity 2 along D .

In fact, the image in $(A/\pm 1)^{(m)}$ of a general subscheme $\zeta \in D$ is a point $(\bar{x}, \bar{x}, \bar{a}_3, \dots, \bar{a}_m)$, for $m - 1$ distinct smooth points $\bar{x}, \bar{a}_3, \dots, \bar{a}_m$ of $(A/\pm 1)$. The subscheme $\iota(\zeta)$ in $A^{[n+1]}$ is supported over $(x, x, -x, -x, a_3, -a_3, \dots, a_m, -a_m)$ if $n = 2m - 1$ is odd, or $(x, x, -x, -x, a_3, -a_3, \dots, a_m, -a_m, 0)$ if $n = 2m$ is even. Let $\mathcal{U} \subset A^{[n+1]}$ be an analytic neighborhood of $\iota(\zeta)$, and let $\mathcal{V} \subset \text{Km}(A)^{[m]}$ be its preimage $\iota^{-1}(\mathcal{U})$. If $n = 2m - 1$ is odd, shrinking \mathcal{U} if necessary, there exist neighborhoods U_x of x and U_j of a_j in A for $j = 3, \dots, m$, such that diagram (10) restricts to

$$\begin{array}{ccc} \mathcal{V} \cong (U_x^{[2]} \times \prod_{j=3}^m U_j) & \xrightarrow{\iota} & \mathcal{U} \cong (U_x^{[2]} \times (U_x^-)^{[2]} \times \prod_{j=3}^m (U_j \times U_j^-)) \\ \downarrow & & \downarrow \nu \\ U_x^{(2)} \times \prod_{j=3}^m U_j & \xrightarrow{\bar{\iota}} & U_x^{(2)} \times (U_x^-)^{(2)} \times \prod_{j=3}^m (U_j \times U_j^-) \end{array}$$

where: U_x^- and U_j^- denote the image via -1 of U_x and U_j respectively, the map ν is the product of the Hilbert–Chow morphisms $U_x^{[2]} \rightarrow U_x^{(2)}$ and $(U_x^-)^{[2]} \rightarrow (U_x^-)^{(2)}$ with the identity of each factor $(U_j \times U_j^-)$ for $j = 3, \dots, m$, and $\bar{\iota}$ is given by

$$\bar{\iota}((x_1, x_2), \alpha_3, \dots, \alpha_m) = ((x_1, x_2), (-x_1, -x_2), \alpha_3, -\alpha_3, \dots, \alpha_m, -\alpha_m).$$

If instead $n = 2m$ is even, there exist neighborhoods $U_x \subset A$ of x and $U_j \subset A$ of a_j for $j = 3, \dots, m$, and a neighborhood $U_0 \subset A$ of 0 such that diagram (10) restricts to

$$\begin{array}{ccc} \mathcal{V} \cong (U_x^{[2]} \times \prod_{j=3}^m U_j) & \xrightarrow{\iota} & \mathcal{U} \cong (U_x^{[2]} \times (U_x^-)^{[2]} \times \prod_{j=3}^m (U_j \times U_j^-) \times U_0) \\ \downarrow & & \downarrow \nu \\ U_x^{(2)} \times \prod_{j=3}^m U_j & \xrightarrow{\bar{\iota}} & U_x^{(2)} \times (U_x^-)^{(2)} \times \prod_{j=3}^m (U_j \times U_j^-) \times U_0 \end{array}$$

where: U_x^- and U_j^- denote the image via -1 of U_x and U_j respectively, the map ν is the product of the Hilbert–Chow morphisms $U_x^{[2]} \rightarrow U_x^{(2)}$ and $(U_x^-)^{[2]} \rightarrow (U_x^-)^{(2)}$ with the identity of U_0 and of each

factor $(U_j \times U_j^-)$ for $j = 3, \dots, m$, and we have

$$\bar{\iota}((x_1, x_2), \alpha_3, \dots, \alpha_m) = ((x_1, x_2), (-x_1, -x_2), \alpha_3, -\alpha_3, \dots, \alpha_m, -\alpha_m, 0).$$

In both cases, the divisor $E|_{\mathcal{U}}$ breaks into two components E_1 and E_2 such that $E_1 = \text{pr}_1^*(F)$ and $E_2 = \text{pr}_2^*(F^-)$, where $\text{pr}_1 : \mathcal{U} \rightarrow U_x^{[2]}$ and $\text{pr}_2 : \mathcal{U} \rightarrow (U_x^-)^{[2]}$ denote the projections and $F \subset U_x^{[2]}$ (resp. $F^- \subset (U_x^-)^{[2]}$) is the divisor of nonreduced subschemes. Moreover, denoting the projection by $\tilde{\text{pr}}_1 : \mathcal{V} \rightarrow U_x^{[2]}$, we have $D|_{\mathcal{V}} = \tilde{\text{pr}}_1^*(F)$. Notice that the element $-1 \in \text{Aut}(A)$ induces an isomorphism $-1 : U_x^{[2]} \rightarrow (U_x^-)^{[2]}$ which identifies F with F^- . From the commutative diagram

$$\begin{CD} \mathcal{V} @<\iota<< \mathcal{U} \\ @V{\tilde{\text{pr}}_1}VV @VV{\text{pr}_1 \times \text{pr}_2}V \\ U_x^{[2]} @<{\text{id} \times (-1)}<< U_x^{[2]} \times (U_x^-)^{[2]} \end{CD}$$

we deduce that $\iota^*(E_1) = D|_{\mathcal{V}} = \iota^*(E_2)$, and thus $\iota^*(E)$ has multiplicity 2 at the divisor $D \subset \text{Km}(A)^{[m]}$.

Conclusion: Recalling that we have $[E] = 2\xi$ in $H^2(K^n(A), \mathbb{Z})$ and $[D] = 2\delta$ in $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$, the three cases discussed above yield (i), (ii) and (iii). □

4.4 The key property of the embedding $\iota : \text{Km}(A)^{[m]} \rightarrow K^n(A)$ of Definition 4.1 is that it deforms together with $K^n(A)$. More precisely, we have the following.

Proposition 4.6 *Let $\mathcal{K} \rightarrow B$ be a smooth proper family of complex manifolds of Kumⁿ-type over a connected manifold B , such that for some $0 \in B$ we have $\mathcal{K}_0 = K^n(A)$ for some abelian surface A . Let $\iota_0 : \text{Km}(A)^{[m]} \rightarrow K^n(A)$ be the embedding discussed above, where $n = 2m - 1$ or $n = 2m$. Then, up to a finite étale base-change, there exist a smooth and proper family $\mathcal{W} \rightarrow B$ of manifolds of K3^[m]-type with $\mathcal{W}_0 = \text{Km}(A)^{[m]}$, and a closed embedding $\iota : \mathcal{W} \hookrightarrow \mathcal{K}$ over B extending ι_0 .*

Proof By [Hassett and Tschinkel 2013, Theorem 2.1], the group $\text{Aut}_0(X)$ of automorphisms inducing the identity on $H^2(X, \mathbb{Z})$ is deformation invariant for any hyper-Kähler manifold X . It was computed in [Boissière et al. 2011] that for $K^n(A)$ we have $\text{Aut}_0(K^n(A)) = A_{n+1} \rtimes \langle -1 \rangle$; thus, any K of Kumⁿ-type admits an action of $\text{Aut}_0(K) \cong (\mathbb{Z}/(n+1)\mathbb{Z})^4 \rtimes \mathbb{Z}/2\mathbb{Z}$. In particular, if $\mathcal{K} \rightarrow B$ is a family as in the statement, the action of $-1 \in \text{Aut}(A)$ on the fiber \mathcal{K}_0 extends to a fiberwise automorphism of $\mathcal{K} \rightarrow B$, up to a finite étale base-change possibly needed in order to trivialize the monodromy action on $\text{Aut}_0(\mathcal{K}_0)$. For each $b \in B$, the action of -1 on \mathcal{K}_b is a symplectic involution. By [Camere 2012], each component \mathcal{Y} of the fixed locus $\mathcal{K}^{(-1)}$ is a smooth and proper family $\mathcal{Y} \rightarrow B$ of symplectic manifolds, or just the image of a section of $\mathcal{K} \rightarrow B$. By Lemma 4.2, for $n = 2m$ or $n = 2m - 1$ there exists a unique such component \mathcal{W} of maximal relative dimension $2m$, such that $\mathcal{W}_0 = \text{Km}(A)^{[m]}$ is embedded into \mathcal{K}_0 via $\iota_0 : \text{Km}(A)^{[m]} \hookrightarrow K^n(A)$. We thus have obtained the desired subfamily $\mathcal{W} \rightarrow B$ of manifolds of K3^[m]-type. □

As a consequence, the cohomology class $[\iota(\text{Km}(A)^{[m]})]$ remains algebraic, and in particular a Hodge class, on any deformation of $K^n(A)$.

Definition 4.7 A canonical Hodge class on a hyper-Kähler manifold X is a Hodge class $\eta \in H^{2k}(X, \mathbb{Q})$ which remains Hodge on any deformation of X .

The most obvious examples are given by the Chern classes of X . Any canonical Hodge class $\alpha \in H^{4k}(X, \mathbb{Q})$ on X satisfies the generalized Fujiki relation [1987]: there exists a constant $c(\alpha) \in \mathbb{Q}$ such that for any $\gamma \in H^2(X, \mathbb{Q})$ we have

$$(11) \quad \int_X \alpha \cdot (\gamma)^{2n-2k} = \frac{(2n-2k)!}{(n-k)! 2^{n-k}} \cdot c(\alpha) \cdot q_X(\gamma, \gamma)^{n-k},$$

where $\dim X = 2n$ and q_X is the Beauville–Bogomolov form on $H^2(X, \mathbb{Q})$. The constant $c(1)$ is called the reduced Fujiki constant of X .

Let now $\mathcal{K} \rightarrow B$ be a family of manifolds of Kumⁿ-type as in Proposition 4.6. Up to a finite étale base-change, that proposition yields a subfamily $\mathcal{W} \rightarrow B$ of manifolds of K3^[m]-type, where $n = 2m$ or $n = 2m - 1$. We let $\iota : \mathcal{W} \rightarrow \mathcal{K}$ denote the embedding.

Lemma 4.8 For any $b \in B$, the pullback $\iota_b^* : H^2(\mathcal{K}_b, \mathbb{Z}) \rightarrow H^2(\mathcal{W}_b, \mathbb{Z})$ is injective and multiplies the form by a factor 2.

Proof Extending our family $\mathcal{K} \rightarrow B$ if necessary, we may assume that the very general fiber has Picard rank 0, so that $H^2(\mathcal{K}_b, \mathbb{Z})$ is an irreducible Hodge structure for very general b . A symplectic form on \mathcal{K}_b restricts to a symplectic form on \mathcal{W}_b (see [Camere 2012, Proposition 3]), and hence ι_b^* is not zero. But then $\iota_b^* : H^2(\mathcal{K}_b, \mathbb{Q}) \rightarrow H^2(\mathcal{W}_b, \mathbb{Q})$ must be an embedding of rational Hodge structures for very general $b \in B$. Since $H^2(\mathcal{K}_b, \mathbb{Z})$ is torsion-free, we conclude that $\iota_b^* : H^2(\mathcal{K}_b, \mathbb{Z}) \rightarrow H^2(\mathcal{W}_b, \mathbb{Z})$ is injective for all $b \in B$.

For the second assertion, it suffices to consider $\iota : \text{Km}(A)^{[m]} \rightarrow K^n(A)$. Assume first that $n = 2$ and $m = 1$. The cohomology class $[\iota(\text{Km}(A))] \in H^4(K^2(A), \mathbb{Z})$ is a canonical Hodge class and, hence, there exists a constant c such that

$$q_{\text{Km}(A)}(\iota^*(\gamma), \iota^*(\gamma)) = \int_{\text{Km}(A)} \iota^*(\gamma)^2 = \int_{K^2(A)} [\text{Km}(A)] \cdot \gamma^2 = c \cdot q_{K^2(A)}(\gamma, \gamma)$$

for any $\gamma \in H^2(K^2(A), \mathbb{Z})$. For the class $\xi \in H^2(K^2(A), \mathbb{Z})$, we have $q_{K^2(A)}(\xi, \xi) = -6$ while $q_{\text{Km}(A)}(\iota^*(\xi), \iota^*(\xi)) = -12$ by Lemma 4.4(i), and we get $c = 2$.

Consider next the case when $n = 2m - 1$ is odd, with $m \geq 2$. The cohomology class $[\iota(\text{Km}(A)^{[m]})]$ in $H^{4m-4}(K^{2m-1}(A), \mathbb{Z})$ is a canonical Hodge class. Using the generalized Fujiki relation (11) for $[\iota(\text{Km}(A)^{[m]})]$ and Fujiki relation for $\text{Km}(A)^{[m]}$, there exists a constant c such that

$$(12) \quad \frac{(2m)!}{m! 2^m} \cdot c_{\text{K3}^{[m]}} \cdot q_{\text{Km}(A)^{[m]}}(\iota^*(\gamma), \iota^*(\gamma))^m = \int_{\text{Km}(A)^{[m]}} \iota^*(\gamma)^{2m} = \frac{(2m)!}{m! 2^m} \cdot c \cdot q_{K^{2m-1}(A)}(\gamma, \gamma)^m$$

for any $\gamma \in H^2(K^{2m-1}(A), \mathbb{Z})$, where $c_{\text{K3}^{[m]}} = 1$ is the reduced Fujiki constant of manifolds of K3^[m]-type (computed in [Beauville 1983]). For the class $\xi \in H^2(K^{2m-1}(A), \mathbb{Z})$ which is half the class of the

divisor of nonreduced subschemes, we have $q_{K^{2m-1}(A)}(\xi, \xi) = -4m$ and, by Lemma 4.4(ii), we have $q_{\text{Km}(A)^{[m]}}(\iota^*(\xi), \iota^*(\xi)) = -8m$. We thus obtain

$$(-8m)^m = c \cdot (-4m)^m,$$

which gives $c = 2^m$. Therefore, for any $\gamma \in H^2(K^{2m-1}(A), \mathbb{Z})$ equation (12) yields

$$q_{\text{Km}(A)^{[m]}}(\iota^*(\gamma), \iota^*(\gamma))^m = 2^m \cdot q_{K^{2m-1}(A)}(\gamma, \gamma)^m,$$

and hence ι^* multiplies the form by 2 (for m even, the sign is determined as ι^* sends Kähler classes to Kähler classes).

Finally, assume $n = 2m \geq 4$ is even. As above, there exists a constant c such that

$$(13) \quad c_{K3^{[m]}} \cdot q_{\text{Km}(A)^{[m]}}(\iota^*(\gamma), \iota^*(\gamma))^m = \int_{\text{Km}(A)^{[m]}} \iota^*(\gamma)^{2m} = c \cdot q_{K^{2m}(A)}(\gamma, \gamma)^m$$

for all $\gamma \in H^2(K^{2m}(A), \mathbb{Z})$. For the class $\xi \in H^2(K^{2m}(A), \mathbb{Z})$ which is half the class of nonreduced subschemes, we have $q_{K^{2m}(A)}(\xi, \xi) = -4m - 2$, while we have $q_{\text{Km}(A)^{[m]}}(\iota^*(\xi), \iota^*(\xi)) = -8m - 4$ by Lemma 4.4(iii). As $c_{K3^{[m]}} = 1$, we obtain

$$(-8m - 4)^m = c \cdot (-4m - 2)^m,$$

and hence $c = 2^m$; by (13), we have $q_{\text{Km}(A)^{[m]}}(\iota^*(\gamma), \iota^*(\gamma))^m = 2^m \cdot q_{K^{2m}(A)}(\gamma, \gamma)^m$ for any γ in $H^2(K^{2m}(A), \mathbb{Z})$, and again we deduce that ι^* multiplies the form by 2. □

4.5 We can finally conclude our study of the pullback ι^* .

Proof of Proposition 4.3 Thanks to Lemma 4.8, it remains to show that the image of

$$\iota^* : H^2(K^n(A), \mathbb{Z}) \hookrightarrow H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$$

is a saturated sublattice. As in (5), we have $H^2(K^n(A), \mathbb{Z}) = H^2(A, \mathbb{Z}) \oplus \langle \xi \rangle$, and the Hodge structure on $H^2(K^n(A), \mathbb{Z})$ is determined by that on $H^2(A, \mathbb{Z})$ via this equality. Moreover, the Hodge structure on $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$ is determined by that on its primitive sublattice $H^2(A, \mathbb{Z})(2)$ (Section 4.2). Deforming to a very general complex torus A such that $H^2(A, \mathbb{Z})$ is an irreducible Hodge structure, we see that the restriction ι^* necessarily maps the summand $H^2(A, \mathbb{Z})$ of $H^2(K^n(A), \mathbb{Z})$ to the sublattice $H^2(A, \mathbb{Z})(2)$ of $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$. In fact, the restriction of ι^* to $H^2(A, \mathbb{Z}) \subset H^2(K^n(A), \mathbb{Z})$ is surjective onto $H^2(A, \mathbb{Z})(2) \subset H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$, as follows from the discriminant-index formula (see [Huybrechts 2016, Chapter 14]) since ι^* is injective and multiplies the form by 2 by Lemma 4.8.

Assume first that $n = 2$ and $m = 1$. Then, by the above and Lemma 4.4, the image of ι^* is the sublattice of $H^2(\text{Km}(A), \mathbb{Z})$ generated by $H^2(A, \mathbb{Z})(2)$ and the class $[C_0] + \frac{1}{2} \sum_{\tau \in A_2} [C_\tau]$. This sublattice is saturated: if $w := \alpha u + \beta \cdot ([C_0] + \frac{1}{2} \sum_{\tau \in A_2} [C_\tau])$ is an integral class in $H^2(\text{Km}(A), \mathbb{Z})$ for some primitive $u \in H^2(A, \mathbb{Z})(2)$ and some rational numbers α and β , then from $q_{\text{Km}(A)}(w, [C_\tau]) = -\beta$ for any $\tau \neq 0$ we deduce that α and β are integers, and therefore w belongs to $\text{im}(\iota^*)$.

For $m \geq 2$, the lattice $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$ contains with finite index the sublattice

$$H^2(A, \mathbb{Z})(2) \oplus \langle [R_\tau] \rangle_{\tau \in A_2} \oplus \langle \delta \rangle.$$

By Lemma 4.4, if $n = 2m - 1$, the image of ι^* is the sublattice of $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$ generated by $H^2(A, \mathbb{Z})(2)$ and the class $2\delta + \frac{1}{2} \sum_{\tau \in A_2} [R_\tau]$, while if $n = 2m$ the image of ι^* is the sublattice of $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$ generated by $H^2(A, \mathbb{Z})(2)$ and the class $2\delta + [R_0] + \frac{1}{2} \sum_{\tau \in A_2} [R_\tau]$. In both cases, the same argument used above shows that the image of ι^* is a saturated sublattice of $H^2(\text{Km}(A)^{[m]}, \mathbb{Z})$. \square

Remark 4.9 A manifold of Kum^n -type K admits at least $(n + 1)^4$ involutions acting trivially on the second cohomology, whose fixed loci are union of $\text{K3}^{[j]}$ -manifolds by [Kamenova et al. 2022]. Our results adapt immediately to all components of maximal dimension. For a lower-dimensional component $\iota : Z \hookrightarrow K$, the argument of Lemma 4.8 gives that $\iota^* : H^2(K, \mathbb{Z}) \rightarrow H^2(Z, \mathbb{Z})$ is injective and multiplies the form by some positive constant k ; however, determining this constant and the saturation of the image of ι^* becomes more difficult.

5 K3 surfaces associated with varieties of generalized Kummer type

5.1 In this section we shall conclude the proof of our main results.

Definition 5.1 Let K be a variety of Kum^n -type. The associated K3 surface S_K is the K3 surface with transcendental lattice Hodge isometric to $H_{\text{tr}}^2(K, \mathbb{Z})(2)$.

In the special case of the generalized Kummer variety $K^n(A)$ on an abelian surface A , we have $H_{\text{tr}}^2(K^n(A), \mathbb{Z}) = H_{\text{tr}}^2(A, \mathbb{Z})$, and the K3 surface $S_{K^n(A)}$ is isomorphic to the Kummer K3 surface $\text{Km}(A)$. This motivates our Definition 5.1. Note that for K very general S_K is projective of Picard rank 16, and thus it is not a Kummer K3 surface.

Lemma 5.2 *The K3 surface S_K exists and it is unique up to isomorphism.*

Proof The lattice $H_{\text{tr}}^2(K, \mathbb{Z})(2)$ has rank at most 6 and signature $(2, k)$. Any such lattice can be embedded primitively in the K3 lattice and, by the Torelli theorem, there exists a K3 surface S_K whose transcendental lattice $H_{\text{tr}}^2(S_K, \mathbb{Z})$ is Hodge isometric to $H_{\text{tr}}^2(K, \mathbb{Z})(2)$; see [Morrison 1984, Corollary 2.10]. The K3 surface S_K is necessarily projective, of Picard rank at least 16, and it is determined up to isomorphism because projective K3 surfaces with Hodge isometric transcendental lattices and Picard rank at least 12 are isomorphic; see [Huybrechts 2016, Chapter 16, Corollary 3.8]. \square

The K3 surfaces associated with varieties of generalized Kummer type can be easily characterized via the Torelli theorem.

Lemma 5.3 *Let S be a projective K3 surface. Then S is isomorphic to the K3 surface S_K associated to some variety K of Kum^n -type if and only if there exists a primitive embedding of lattices*

$$H_{\text{tr}}^2(S, \mathbb{Z}) \hookrightarrow U(2)^{\oplus 3} \oplus \langle -4n - 4 \rangle.$$

Proof The lattice $U(2)^{\oplus 3} \oplus \langle -4n-4 \rangle$ is the Beauville–Bogomolov lattice $\Lambda_{\text{Kum}^n}(2)$ of Kumⁿ-manifolds with the form multiplied by 2. Thus, by definition, for the K3 surface S_K associated to a variety K of Kumⁿ-type we have a primitive embedding of $H_{\text{tr}}^2(S_K, \mathbb{Z})$ into the lattice $U(2)^{\oplus 3} \oplus \langle -4n-4 \rangle$. Conversely, assume that S is a projective K3 surface and $l : H_{\text{tr}}^2(S, \mathbb{Z}) \hookrightarrow \Lambda_{\text{Kum}^n}(2)$ is a primitive embedding. Let $T \subset \Lambda_{\text{Kum}^n}$ be the primitive sublattice such that $\text{im}(l) = T(2)$, and equip T with the Hodge structure induced by that on $H_{\text{tr}}^2(S, \mathbb{Z})$. By the surjectivity of the period map for hyper-Kähler manifolds [Huybrechts 1999, Theorem 8.1], there exists a manifold K of Kumⁿ-type with $H_{\text{tr}}^2(K, \mathbb{Z})$ Hodge isometric to T . By construction, we have a Hodge isometry

$$H_{\text{tr}}^2(S, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Z})(2);$$

K is projective by [Huybrechts 1999, Theorem 3.11], since its transcendental lattice has signature $(2, k)$. Moreover, the K3 surface S is isomorphic to S_K , as they have Hodge isometric transcendental lattices and Picard rank at least 16. □

Passing to rational coefficients, a similar argument yields the following.

Lemma 5.4 *Let S be a projective K3 surface. The following are equivalent:*

- (i) *There exists a rational Hodge isometry $H_{\text{tr}}^2(S, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(S_K, \mathbb{Q})$, where S_K is the K3 surface associated to a variety K of Kumⁿ-type for some $n \geq 2$.*
- (ii) *There exists an isometric embedding of $H_{\text{tr}}^2(S, \mathbb{Q})$ into $(U^{\oplus 3} \oplus \langle -m \rangle) \otimes_{\mathbb{Z}} \mathbb{Q}$ for some positive integer m .*

Proof For any nonzero rational number k , we have $U(k) \otimes_{\mathbb{Z}} \mathbb{Q} \cong U \otimes_{\mathbb{Z}} \mathbb{Q}$. If (i) holds, then $H_{\text{tr}}^2(S, \mathbb{Q})$ is Hodge isometric to $H_{\text{tr}}^2(K, \mathbb{Q})(2)$, and, hence, it embeds isometrically into $(U(2)^{\oplus 3} \oplus \langle -4n-4 \rangle) \otimes_{\mathbb{Z}} \mathbb{Q} \cong (U^{\oplus 3} \oplus \langle -n-1 \rangle) \otimes_{\mathbb{Z}} \mathbb{Q}$. Conversely, if S is a projective K3 surface with an embedding $H_{\text{tr}}^2(S, \mathbb{Q}) \hookrightarrow (U^{\oplus 3} \oplus \langle -m \rangle) \otimes_{\mathbb{Z}} \mathbb{Q}$ for some positive $m \in \mathbb{Z}$, we find as well an isometric embedding

$$j : H_{\text{tr}}^2(S, \mathbb{Q}) \hookrightarrow (U(2)^{\oplus 3} \oplus \langle -4n-4 \rangle) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for some $n \geq 2$. The quadratic space on the right-hand side is $\Lambda_{\text{Kum}^n}(2) \otimes_{\mathbb{Z}} \mathbb{Q}$. We choose a primitive sublattice $T \subset \Lambda_{\text{Kum}^n}$ such that $\text{im}(j)$ coincides with $T(2) \otimes_{\mathbb{Z}} \mathbb{Q}$; under this identification, the Hodge structure on $H_{\text{tr}}^2(S, \mathbb{Q})$ induces a Hodge structure of K3-type on T . By the surjectivity of the period map, there exists a projective variety K of Kumⁿ-type such that $H_{\text{tr}}^2(K, \mathbb{Z})$ is Hodge isometric to T , equipped with this Hodge structure. Hence, by construction, $H_{\text{tr}}^2(S, \mathbb{Q})$ is Hodge isometric to $H_{\text{tr}}^2(S_K, \mathbb{Q})$. □

5.2 A priori, the K3 surface S_K and the variety K of Kumⁿ-type are only related at the level of lattices and Hodge structures. But in fact they are related geometrically, as follows. Choose a family $\mathcal{H} \rightarrow B$ of manifolds of Kumⁿ-type, with one fiber $\mathcal{H}_{b_0} = K$ and another fiber $\mathcal{H}_{b_1} = K^n(A)$ isomorphic to the generalized Kummer variety on an abelian surface A . By Proposition 4.6, up to a finite étale base-change, we obtain a family $\mathcal{W} \rightarrow B$ of K3^[m]-manifolds with an embedding $\iota : \mathcal{W} \rightarrow \mathcal{H}$ over B , where $n = 2m - 1$

or $n = 2m$, which extends the embedding $\iota_{b_1} : \text{Km}(A)^{[m]} \hookrightarrow K^n(A)$ of Lemma 4.2. Taking the fiber over b_0 , we get a submanifold $\iota : W_K \hookrightarrow K$ of $\text{K3}^{[m]}$ -type.

Lemma 5.5 *The pullback along $\iota : W_K \hookrightarrow K$ induces a primitive embedding of lattices and Hodge structures*

$$\iota^* : H^2(K, \mathbb{Z})(2) \hookrightarrow H^2(W_K, \mathbb{Z}).$$

In particular, ι^ induces a Hodge isometry $H_{\text{tr}}^2(K, \mathbb{Z})(2) \xrightarrow{\sim} H_{\text{tr}}^2(W_K, \mathbb{Z})$ of transcendental lattices.*

Proof The lemma follows immediately from Proposition 4.3 as $\iota : W_K \hookrightarrow K$ deforms to the embedding $\text{Km}(A)^{[m]} \hookrightarrow K^n(A)$ there considered. \square

We refer to [Huybrechts and Lehn 2010; Bridgeland 2008; Bayer and Macrì 2014b] for background on moduli spaces of sheaves on K3 surfaces.

Proposition 5.6 *Let S_K be the K3 surface associated to the Kum^n -variety K . Then:*

- (i) *If $n = 2$, W_K is isomorphic to the K3 surface S_K .*
- (ii) *If $n \geq 3$, $W_K \subset K$ is birational to a smooth and projective moduli space $M_{S_K, H}(v)$ for some primitive Mukai vector v and a v -generic polarization H .*

Proof A Hodge isometry $H_{\text{tr}}^2(W_K, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(S_K, \mathbb{Z})$ of transcendental lattices exists by Lemma 5.5 and the definition of S_K . If $n = 2$, W_K and S_K are isomorphic as they are K3 surfaces with Hodge isometric transcendental lattice and large Picard rank. If $n \geq 3$, the proposition follows applying [Addington 2016, Proposition 4]. \square

Remark 5.7 In fact, for $n \geq 3$, W_K is isomorphic to a moduli space of Bridgeland stable objects on S_K , by [Bayer and Macrì 2014a, Theorem 1.2(c)]; see also [Mongardi and Wandel 2017, Proposition 2.3].

5.3 Let S be a projective K3 surface, and let $M = M_{S, H}(v)$ be a smooth and projective moduli space of stable sheaves of dimension ≥ 4 , for a primitive Mukai vector v and a v -generic polarization H . Then, following [Mukai 1987; O’Grady 1997], there exists a quasiuniversal sheaf \mathcal{U} over $S \times M$ of similitude ρ , and the map

$$\alpha \mapsto \frac{1}{\rho} \cdot \text{pr}_{M, *}[\text{ch}(\mathcal{U})^\vee \cdot \sqrt{\text{td}(S)} \cdot \text{pr}_S^*(\alpha)]_3$$

induces a Hodge isometry $\theta : H_{\text{tr}}^2(S, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(M, \mathbb{Z})$ of transcendental lattices. Here, the notation $[-]_3$ indicates that we take the Künneth component in $H^6(S \times M, \mathbb{Z})$ and pr_M, pr_S denote the projections from $S \times M$ onto the two factors. The standard conjectures hold for both the surface S and the $\text{K3}^{[n]}$ -variety M , as we recalled in Section 2.5. Hence, they hold for $S \times M$, and it follows that θ is the correspondence induced by the algebraic cycle $[(1/\rho)\text{ch}(\mathcal{U})^\vee \cdot \sqrt{\text{td}(S)}]_2$ on $S \times M$.

Remark 5.8 In fact, the Künneth components $h^2(S)$ and $h^2(M)$ as well as their transcendental parts $h_{\text{tr}}^2(S)$ and $h_{\text{tr}}^2(M)$ are well-defined homological motives; the Hodge isometry θ is the realization of an isomorphism of motives $\Theta : h_{\text{tr}}^2(S) \rightarrow h_{\text{tr}}^2(M)$ in Mot . This again uses the standard conjectures for

$S \times M$ and in particular the conservativity of the realization functor when restricted to $\langle h(S \times M) \rangle_{\text{Mot}}$ (Remark 2.4).

5.4 Let K be a variety of Kum^{*n*}-type. By the results of Foster [2024] recalled in Theorem 2.6, the Künneth component $h^2(K)$ of the motive of K is well-defined in Mot and $\langle h^2(K) \rangle_{\text{Mot}}$ is a semisimple and abelian neutral Tannakian category; the transcendental part $h_{\text{tr}}^2(K)$ of $h^2(K)$ is also a well-defined homological motive. The next result implies Theorem 1.1 from the introduction.

Theorem 5.9 *Let K be a variety of Kum^{*n*}-type and let S_K be the K3 surface associated with K . There exists an isomorphism*

$$h_{\text{tr}}^2(S_K) \xrightarrow{\sim} h_{\text{tr}}^2(K)$$

of motives, whose realization is a Hodge isometry $H_{\text{tr}}^2(S_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(2)$.

Proof For $n = 2m$ or $n = 2m - 1$, we find as in Section 5.2 a $\text{K3}^{[m]}$ -variety W_K which is isomorphic to S_K for $n = 2$ and birational to a moduli space of stable sheaves on S_K if $n \geq 3$, and that admits an embedding $\iota : W_K \hookrightarrow K$ such that the restriction induces a Hodge isometry $\iota^* : H_{\text{tr}}^2(K, \mathbb{Z})(2) \xrightarrow{\sim} H_{\text{tr}}^2(W_K, \mathbb{Z})$.

As ι^* is clearly induced by an algebraic cycle, it is the realization of a morphism $t : h_{\text{tr}}^2(K) \rightarrow h_{\text{tr}}^2(W_K)$ of motives. But in fact t is a morphism in the subcategory $\langle h_{\text{tr}}^2(K) \oplus h_{\text{tr}}^2(W_K) \rangle_{\text{Mot}}$, which is an abelian and semisimple neutral Tannakian category as the standard conjectures hold for W_K and $h^2(K)$. Since the realization of t is an isomorphism of rational Hodge structures, it follows that t is an isomorphism of motives; the realization of its inverse t^{-1} is a Hodge isometry $H_{\text{tr}}^2(W_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(2)$. This concludes the proof for $n = 2$, since S_K is isomorphic to W_K in this case.

If $n \geq 3$, let $\phi : M_{S_K, H}(v) \dashrightarrow W_K$ be a birational map, where $M_{S_K, H}(v)$ is a smooth and projective moduli space of stable sheaves on S_K . Then ϕ induces a morphism $\Phi_* : h^2(M_{S_K, H}(v)) \rightarrow h^2(W_K)$ of motives whose realization is a Hodge isometry $\phi_* : H^2(M_{S_K, H}(v), \mathbb{Z}) \xrightarrow{\sim} H^2(W_K, \mathbb{Z})$, by [Huybrechts 1999, Corollary 5.2]. As the standard conjectures hold for $\text{K3}^{[m]}$ -varieties, Φ_* is an isomorphism of motives. From Remark 5.8 we obtain an isomorphism of motives $\Theta : h_{\text{tr}}^2(S_K) \xrightarrow{\sim} h_{\text{tr}}^2(M_{S_K, H}(v))$, whose realization is a Hodge isometry $H_{\text{tr}}^2(S_K, \mathbb{Z}) \xrightarrow{\sim} H_{\text{tr}}^2(M_{S_K, H}(v), \mathbb{Z})$. We conclude that the composition $t^{-1} \circ \Phi_* \circ \Theta : h_{\text{tr}}^2(S_K) \rightarrow h_{\text{tr}}^2(K)$ is an isomorphism of motives, whose realization is the desired Hodge isometry $H_{\text{tr}}^2(S_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(2)$. □

Thanks to the work of O’Grady, Markman, Voisin and Varesco, recalled in the introduction, Theorem 5.9 yields the following result, which was stated as Theorem 1.2.

Theorem 5.10 *Let K be a variety of Kum^{*n*}-type with associated K3 surface S_K . Then:*

- (i) *The Kuga–Satake correspondence for S_K is algebraic.*
- (ii) *The Hodge conjecture holds for any power of S_K .*

Proof By Theorem 5.9 there exists an algebraic cycle inducing a Hodge isometry

$$H_{\text{tr}}^2(S_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(2).$$

The Kuga–Satake Hodge conjecture (Conjecture 3.3) holds for K by [Voisin 2022, Theorem 0.5]. By Remark 3.7, the same conclusion holds for S_K , and we obtain (i).

The K3 surfaces S_K come in countably many 4-dimensional families of general Picard rank 16, corresponding to the polarized families of varieties of generalized Kummer type. By construction (see Section 3.5) the Kuga–Satake variety of S_K is isogenous to a power of $\text{KS}(K)$, and, hence, to a power of a Weil fourfold with discriminant 1 by [Markman 2023; O’Grady 2021]. As Conjecture 3.3 holds for any of the K3 surfaces S_K , we can apply [Varesco 2025, Theorem 0.2] to deduce that the Hodge conjecture holds for all powers of the K3 surfaces S_K . \square

5.5 We finally complete the proof of our results.

Theorem 5.11 *Let S be a projective K3 surface such that there exists an isometric embedding*

$$H_{\text{tr}}^2(S, \mathbb{Q}) \hookrightarrow (U^{\oplus 3} \oplus \langle -m \rangle) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for some positive integer m . Then:

- (i) *The Kuga–Satake correspondence for S is algebraic.*
- (ii) *The Hodge conjecture holds for any power of S .*

Proof Let S be as in the statement. By Lemma 5.4, for some $n \geq 2$ there exist a variety K of Kum^n -type and a rational Hodge isometry $\phi : H_{\text{tr}}^2(S, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(S_K, \mathbb{Q})$. By [Buskin 2019; Huybrechts 2019], ϕ is algebraic and the motives of S and S_K are isomorphic. Therefore, by Remark 3.7 and Theorem 5.10(i), the Kuga–Satake correspondence for S is algebraic, and, by Theorem 5.10(ii), the Hodge conjecture holds for any power of S . \square

We next prove Theorem 1.3, which we state again below.

Theorem 5.12 *Let A be an abelian fourfold of Weil type with discriminant 1. Then the Hodge conjecture holds for A and any of its powers.*

Proof By [O’Grady 2021] and [Markman 2023], there exists a Kum^n -variety K such that A is isogenous to the intermediate Jacobian $J^3(K)$; moreover, the Kuga–Satake variety of K is isogenous to a power of A . The Kuga–Satake variety of the K3 surface S_K associated to K is thus also isogenous to a power of A . By Theorem 5.10, the Kuga–Satake Hodge conjecture holds for S_K and the Hodge conjecture holds for any power of S_K . Therefore, Theorem 3.5 implies that the Hodge conjecture holds for A and all of its powers. \square

We conclude with the following corollary, which follows from a result of Varesco [2023].

Corollary 5.13 *Let S be a projective K3 surface and let K be a projective variety of Kum^n -type. Assume that $\phi : H_{\text{tr}}^2(S_K, \mathbb{Q}) \xrightarrow{\sim} H_{\text{tr}}^2(K, \mathbb{Q})(k)$ is a Hodge isometry, where on the right-hand side the form is multiplied by some positive $k \in \mathbb{Q}$. Then ϕ is induced by an algebraic cycle on $S \times K$.*

Proof As $\Lambda_{\text{Kum}^n} = U^{\oplus 3} \oplus \langle -2n - 2 \rangle$ and $U(k) \otimes_{\mathbb{Z}} \mathbb{Q} \cong U \otimes_{\mathbb{Z}} \mathbb{Q}$ for any $0 \neq k \in \mathbb{Q}$, there exists by hypothesis an isometric embedding

$$H_{\text{tr}}^2(S_K, \mathbb{Q}) \hookrightarrow U_{\mathbb{Q}}^{\oplus 3} \oplus \langle -m \rangle_{\mathbb{Q}}$$

for some positive integer m . Hence, by Theorem 1.2, the Kuga–Satake Hodge conjecture holds for S . Moreover, this conjecture holds for K by [Voisin 2022, Theorem 0.5], and the Lefschetz standard conjecture in degree 2 for K is proven in [Foster 2024, Theorem 1.1]. We can therefore apply [Varesco 2023, Corollary 4.6] to obtain that ϕ is algebraic. \square

Acknowledgements

I am grateful to Lie Fu, Bert van Geemen, Ben Moonen, Ángel Ríos Ortiz and Charles Vial for useful conversations and to the referee for helpful comments. I thank Stefan Schreieder for his support. Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 491392403 - TRR 358.

References

- [Addington 2016] **N Addington**, *On two rationality conjectures for cubic fourfolds*, *Math. Res. Lett.* 23:1 (2016) 1–13 MR
- [Ancona et al. 2025] **G Ancona, M Cavicchi, R Laterveer, G Saccà**, *Relative and absolute Lefschetz standard conjectures for some Lagrangian fibrations*, *J. Lond. Math. Soc. (2)* 111:4 (2025) art. id. e70133 MR
- [André 1996a] **Y André**, *On the Shafarevich and Tate conjectures for hyper-Kähler varieties*, *Math. Ann.* 305:2 (1996) 205–248 MR
- [André 1996b] **Y André**, *Pour une théorie inconditionnelle des motifs*, *Inst. Hautes Études Sci. Publ. Math.* :83 (1996) 5–49 MR
- [André 2004] **Y André**, *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*, *Panoramas et Synthèses* 17, Soc. Math. France, Paris (2004) MR
- [Arapura 2006] **D Arapura**, *Motivation for Hodge cycles*, *Adv. Math.* 207:2 (2006) 762–781 MR
- [Bayer and Macrì 2014a] **A Bayer, E Macrì**, *MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations*, *Invent. Math.* 198:3 (2014) 505–590 MR
- [Bayer and Macrì 2014b] **A Bayer, E Macrì**, *Projectivity and birational geometry of Bridgeland moduli spaces*, *J. Amer. Math. Soc.* 27:3 (2014) 707–752 MR
- [Beauville 1983] **A Beauville**, *Variétés Kähleriennes dont la première classe de Chern est nulle*, *J. Differential Geom.* 18:4 (1983) 755–782 MR
- [Boissière et al. 2011] **S Boissière, M Nieper-Wißkirchen, A Sarti**, *Higher dimensional Enriques varieties and automorphisms of generalized Kummer varieties*, *J. Math. Pures Appl. (9)* 95:5 (2011) 553–563 MR
- [Bolognesi and Laterveer 2024] **M Bolognesi, R Laterveer**, *A 9-dimensional family of K3 surfaces with finite-dimensional motive*, *Rend. Circ. Mat. Palermo (2)* 73:7 (2024) 2313–2331 MR
- [Brianchon 1977] **J Brianchon**, *Description de $\text{Hilb}^n C\{x, y\}$* , *Invent. Math.* 41:1 (1977) 45–89 MR
- [Bridgeland 2008] **T Bridgeland**, *Stability conditions on K3 surfaces*, *Duke Math. J.* 141:2 (2008) 241–291 MR
- [Bülles 2020] **T-H Bülles**, *Motives of moduli spaces on K3 surfaces and of special cubic fourfolds*, *Manuscripta Math.* 161:1-2 (2020) 109–124 MR
- [Buskin 2019] **N Buskin**, *Every rational Hodge isometry between two K3 surfaces is algebraic*, *J. Reine Angew. Math.* 755 (2019) 127–150 MR

- [Camere 2012] **C Camere**, *Symplectic involutions of holomorphic symplectic four-folds*, Bull. Lond. Math. Soc. 44:4 (2012) 687–702 MR
- [Charles and Markman 2013] **F Charles, E Markman**, *The standard conjectures for holomorphic symplectic varieties deformation equivalent to Hilbert schemes of K3 surfaces*, Compos. Math. 149:3 (2013) 481–494 MR
- [Deligne 1972] **P Deligne**, *La conjecture de Weil pour les surfaces K3*, Invent. Math. 15 (1972) 206–226 MR
- [Deligne and Milne 1982] **P Deligne, JS Milne**, *Tannakian categories*, from “Hodge cycles, motives, and Shimura varieties” (A Dold, B Eckmann, editors), Lecture Notes in Mathematics 900, Springer (1982) 101–228 MR
- [Floccari 2023a] **S Floccari**, *The Hodge and Tate conjectures for hyper-Kähler sixfolds of generalized Kummer type*, preprint (2023) arXiv 2308.02267
- [Floccari 2023b] **S Floccari**, *On the motive of O’Grady’s six-dimensional hyper-Kähler varieties*, Épijournal Géom. Algébrique 7 (2023) art. id. 4 MR
- [Floccari 2024] **S Floccari**, *Sixfolds of generalized Kummer type and K3 surfaces*, Compos. Math. 160:2 (2024) 388–410 MR
- [Floccari and Varesco 2025] **S Floccari, M Varesco**, *Algebraic cycles on hyper-Kähler varieties of generalized Kummer type*, Math. Ann. 391:3 (2025) 4443–4453 MR
- [Floccari et al. 2021] **S Floccari, L Fu, Z Zhang**, *On the motive of O’Grady’s ten-dimensional hyper-Kähler varieties*, Commun. Contemp. Math. 23:4 (2021) art. id. 2050034 MR
- [Foster 2024] **J Foster**, *The Lefschetz standard conjectures for IHSMs of generalized Kummer deformation type in certain degrees*, Eur. J. Math. 10:2 (2024) art. id. 34 MR
- [Fujiki 1987] **A Fujiki**, *On the de Rham cohomology group of a compact Kähler symplectic manifold*, from “Algebraic geometry” (Sendai, 1985) (T Oda, editor), Adv. Stud. Pure Math. 10, North-Holland, Amsterdam (1987) 105–165 MR
- [Fulton 1998] **W Fulton**, *Intersection theory*, 2nd edition, Ergebnisse der Math. (3) 2, Springer (1998) MR
- [van Geemen 2000] **B van Geemen**, *Kuga–Satake varieties and the Hodge conjecture*, from “The arithmetic and geometry of algebraic cycles” (Banff, 1998) (B B Gordon, J D Lewis, S Müller-Stach, S Saito, N Yui, editors), NATO Sci. Ser. C Math. Phys. Sci. 548, Kluwer, Dordrecht (2000) 51–82 MR
- [van Geemen 2023] **B van Geemen**, *Fourfolds of Weil type and the spinor map*, Expo. Math. 41:2 (2023) 418–447 MR
- [Grothendieck 1969] **A Grothendieck**, *Standard conjectures on algebraic cycles*, from “Algebraic geometry” (Bombay, 1968) (S S Abhyankar, editor), Tata Inst. Fundam. Res. Stud. Math. 4, Tata Inst. Fund. Res., Bombay (1969) 193–199 MR
- [Hassett and Tschinkel 2013] **B Hassett, Y Tschinkel**, *Hodge theory and Lagrangian planes on generalized Kummer fourfolds*, Mosc. Math. J. 13:1 (2013) 33–56, 189 MR
- [Huybrechts 1999] **D Huybrechts**, *Compact hyper-Kähler manifolds: basic results*, Invent. Math. 135:1 (1999) 63–113 MR
- [Huybrechts 2016] **D Huybrechts**, *Lectures on K3 surfaces*, Cambridge Studies in Advanced Mathematics 158, Cambridge Univ. Press (2016) MR
- [Huybrechts 2019] **D Huybrechts**, *Motives of isogenous K3 surfaces*, Comment. Math. Helv. 94:3 (2019) 445–458 MR
- [Huybrechts and Lehn 2010] **D Huybrechts, M Lehn**, *The geometry of moduli spaces of sheaves*, 2nd edition, Cambridge Univ. Press (2010) MR
- [Ingalls et al. 2022] **C Ingalls, A Logan, O Patashnick**, *Explicit coverings of families of elliptic surfaces by squares of curves*, Math. Z. 302:2 (2022) 1191–1238 MR
- [Jannsen 1992] **U Jannsen**, *Motives, numerical equivalence, and semi-simplicity*, Invent. Math. 107:3 (1992) 447–452 MR
- [Kaledin and Verbitsky 1998] **D Kaledin, M Verbitsky**, *Partial resolutions of Hilbert type, Dynkin diagrams, and generalized Kummer varieties*, preprint (1998) arXiv math/9812078
- [Kamenova et al. 2022] **L Kamenova, G Mongardi, A Oblomkov**, *Symplectic involutions of $K3^{[n]}$ type and Kummer n type manifolds*, Bull. Lond. Math. Soc. 54:3 (2022) 894–909 MR
- [Kleiman 1968] **SL Kleiman**, *Algebraic cycles and the Weil conjectures*, from “Dix exposés sur la cohomologie des schémas” (A Grothendieck, NH Kuiper, editors), Adv. Stud. Pure Math. 3, North-Holland, Amsterdam (1968) 359–386 MR
- [Kuga and Satake 1967] **M Kuga, I Satake**, *Abelian varieties attached to polarized K_3 -surfaces*, Math. Ann. 169 (1967) 239–242 MR

- [Markman 2023] **E Markman**, *The monodromy of generalized Kummer varieties and algebraic cycles on their intermediate Jacobians*, J. Eur. Math. Soc. (JEMS) 25:1 (2023) 231–321 MR
- [Milne 2021] **J S Milne**, *The Tate and standard conjectures for certain abelian varieties*, preprint (2021) arXiv 2112.12815
- [Mongardi and Wandel 2017] **G Mongardi, M Wandel**, *Automorphisms of O’Grady’s manifolds acting trivially on cohomology*, Algebr. Geom. 4:1 (2017) 104–119 MR
- [Moonen 1999] **B Moonen**, *Notes on Mumford–Tate groups*, unpublished lecture notes, Centre Emile Borel (1999) Available at <https://www.math.ru.nl/~bmoonen/Lecturenotes/CEBnotesMT.pdf>
- [Moonen and Zarhin 1995] **B J J Moonen, Y G Zarhin**, *Hodge classes and Tate classes on simple abelian fourfolds*, Duke Math. J. 77:3 (1995) 553–581 MR
- [Morrison 1984] **D R Morrison**, *On K3 surfaces with large Picard number*, Invent. Math. 75:1 (1984) 105–121 MR
- [Morrison 1985] **D R Morrison**, *The Kuga–Satake variety of an abelian surface*, J. Algebra 92:2 (1985) 454–476 MR
- [Mukai 1987] **S Mukai**, *On the moduli space of bundles on K3 surfaces, I*, from “Vector bundles on algebraic varieties” (Bombay, 1984) (MF Atiyah, editor), Tata Inst. Fund. Res. Stud. Math. 11, Tata Inst. Fund. Res., Bombay (1987) 341–413 MR
- [Nikulin 1975] **V V Nikulin**, *Kummer surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. 39:2 (1975) 278–293, 471 MR In Russian; translated in Math. USSR-Izv. 9:2 (1976), 261–275
- [O’Grady 1997] **K G O’Grady**, *The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface*, J. Algebraic Geom. 6:4 (1997) 599–644 MR
- [O’Grady 2021] **K G O’Grady**, *Compact tori associated to hyperkähler manifolds of Kummer type*, Int. Math. Res. Not. 2021:16 (2021) 12356–12419 MR
- [Oguiso 2020] **K Oguiso**, *No cohomologically trivial nontrivial automorphism of generalized Kummer manifolds*, Nagoya Math. J. 239 (2020) 110–122 MR
- [Paranjape 1988] **K Paranjape**, *Abelian varieties associated to certain K3 surfaces*, Compositio Math. 68:1 (1988) 11–22 MR
- [Varesco 2023] **M Varesco**, *Hodge similarities, algebraic classes, and Kuga–Satake varieties*, Math. Z. 305:4 (2023) art. id. 69 MR
- [Varesco 2025] **M Varesco**, *The Hodge conjecture for powers of K3 surfaces of Picard number 16*, Michigan Math. J. 75:5 (2025) 1085–1116 MR
- [Verbitsky 1995] **M Verbitsky**, *Tri-analytic subvarieties of hyper-Kähler manifolds*, Geom. Funct. Anal. 5:1 (1995) 92–104 MR
- [Voisin 2022] **C Voisin**, *Footnotes to papers of O’Grady and Markman*, Math. Z. 300:4 (2022) 3405–3416 MR
- [Yoshioka 2001] **K Yoshioka**, *Moduli spaces of stable sheaves on abelian surfaces*, Math. Ann. 321:4 (2001) 817–884 MR

SALVATORE FLOCCARI salvatore.floccari@hu-berlin.de

Fakültat für Mathematik, Universität Bielefeld, Bielefeld, Germany

Current address: *Institut für Mathematik, Humboldt-Universität zu Berlin, Berlin, Germany*

Proposed: Sándor J Kovács

Received: January 18, 2025

Seconded: Marc Levine, Dan Abramovich

Revised: July 12, 2025

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITORS

Robert Lipshitz University of Oregon
lipshitz@uoregon.edu
András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Mohammed Abouzaid	Stanford University abouzaid@stanford.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Dan Abramovich	Brown University dan_abramovich@brown.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Sándor Kovács	University of Washington skovacs@uw.edu
Arend Bayer	University of Edinburgh arend.bayer@ed.ac.uk	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Agnès Beaudry	University of Colorado Boulder agnes.beaudry@colorado.edu	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Jianfeng Lin	Tsinghua University linjian5477@mail.tsinghua.edu.cn
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Haynes R Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Aaron Naber	Institute for Advanced Studies anaber@ias.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Peter Ozsváth	Princeton University petero@math.princeton.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
David M Fisher	Rice University davidfisher@rice.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
Mike Freedman	Microsoft Research michaelf@microsoft.com	Roman Sauer	Karlsruhe Institute of Technology roman.sauer@kit.edu
David Gabai	Princeton University gabai@princeton.edu	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Natasa Sesum	Rutgers University natasas@math.rutgers.edu
Cameron Gordon	University of Texas gordon@math.utexas.edu	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHES and NYU, Courant Institute gromov@ihes.fr	Kirsten Wickelgren	Duke University kirsten.wickelgren@duke.edu
Mark Gross	University of Cambridge mgross@dpms.cam.ac.uk	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2026 is US \$905/year for the electronic version, and \$1275/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 30 Issue 3 (pages 835–1201) 2026

The level structure in quantum K -theory and mock theta functions	835
YONGBIN RUAN, MING ZHANG and YAOXIONG WEN	
Springer theory for symplectic Galois groups	883
KEVIN MCGERTY and THOMAS NEVINS	
Unipotent morphisms	929
DANIEL BRAGG, JACK HALL and SIDDHARTH MATHUR	
On the Donaldson–Scaduto conjecture	959
SAMAN HABIBI ESFAHANI and YANG LI	
Milnor’s invariants for knots and links in closed orientable 3-manifolds	983
RYAN STEES	
Chromatic defect, Wood’s theorem and higher real K -theories	1051
CHRISTIAN CARRICK	
Compact holonomy G_2 manifolds need not be formal	1109
LUCÍA MARTÍN-MERCHÁN	
$K3$ surfaces associated with varieties of generalized Kummer type	1129
SALVATORE FLOCCARI	
Ulrich sheaves, the arithmetic writhe and algebraic isotopies of space curves	1155
DANIELE AGOSTINI and MARIO KUMMER	