

On braid groups and homotopy groups

FR COHEN

J WU

This article is an exposition of certain connections between the braid groups, classical homotopy groups of the 2–sphere, as well as Lie algebras attached to the descending central series of pure braid groups arising as Vassiliev invariants of pure braids. Natural related questions are posed at the end of this article.

20F36, 55Q40; 20F40, 55U10, 18G30

1 Introduction

The purpose of this article is to give an exposition of certain connections between the braid groups (Artin [1], see Birman [4]) and classical homotopy groups which arises in joint work of Jon Berrick, Yan-Loi Wong and the authors [9; 10; 3; 38]. These connections emerge through several other natural contexts such as Lie algebras attached to the descending central series of pure braid groups arising as Vassiliev invariants of the pure braid groups as developed by T Kohno [23; 24].

The main feature of this article is to identify certain “non-standard” free subgroups of the braid groups via Vassiliev invariants of pure braids. A second feature is to indicate natural ways in which these subjects fit together with classical homotopy theory. This article is an attempt to draw together these connections.

Since this paper was submitted, other connections to principal congruence subgroups in natural matrix groups as well as other extensions have developed. The authors have taken the liberty of adding an additional Section 7 with some of these new connections.

Natural related questions are posed at the end of this article.

Although not yet useful for direct computations, there is a strong connection between braid groups and homotopy groups. The braids which naturally arise in this setting also give a large class of special knots and links arising from Brunnian braids as described below as well as by Stanford [37]. It is natural to wonder whether and how these fit together.

The authors take this opportunity to thank Toshitake Kohno, Shigeyuki Morita, Dai Tamaki as well as other friends for this very enjoyable opportunity to participate in this conference. The first author is especially grateful for this mathematical opportunity to learn and to work on mathematics with friends. The authors also thank an excellent referee who gave elegant, useful suggestions regarding the exposition and organization of this article.

2 Braid groups, Vassiliev invariants of pure braids and certain free subgroups of braid groups

The section addresses a naive construction with the braid groups arising as a “cabling” construction. This construction is interpreted in later sections in terms of the structure of braid groups, Vassiliev invariants of pure braids as developed by Toshitake Kohno [23; 24], associated Lie algebras and the homotopy groups of the 2–sphere (Berrick, Wong and the authors [9; 10; 3; 38]).

Let B_k denote Artin’s k –stranded braid group while P_k denotes the pure k –stranded braid group, the subgroup of B_k which corresponds to the trivial permutation of the endpoints of the strands. The group P_k is the fundamental group of the configuration space of ordered k –tuples of distinct points in the plane

$$\text{Conf}(\mathbb{R}^2, k)$$

for which $\text{Conf}(M, k) = \{(m_1, m_2, \dots, m_k) \in M^k \mid m_i \neq m_j \text{ for all } i \neq j\}$ for any space M .

The group B_k is the fundamental group of the orbit space

$$\text{Conf}(\mathbb{R}^2, k) / \Sigma_k$$

obtained from the natural, free (left)-action of the symmetric group on k letters Σ_k . The k –stranded braid group of an arbitrary connected surface S , $B_k(S)$, is defined to be the fundamental group of the configuration space of unordered k –tuples of distinct points in S , $\pi_1 \text{Conf}(S, k) / \Sigma_k$. The pure braid group $P_k(S)$ is defined to be the fundamental group

$$\pi_1 \text{Conf}(S, k).$$

The pure braid groups P_k and $P_k(S^2)$ are intimately related to the loop space of the 2–sphere as elucidated below in the Section 4. Similar properties are satisfied for any sphere as described in Section 7.

To start to address this last point, first consider the free group on N letters $F_N = F[y_1, \dots, y_N]$ together with elements x_i for $1 \leq i \leq N$ in P_{N+1} given by the naive

“cabling” pictured in Figure 1 below. The braid x_1 with $N = 1 = i$ in Figure 1 is Artin’s generator $A_{1,2}$ of P_2 . The braids x_i for $1 \leq i \leq N$ in Figure 1 yield homomorphisms from a free group on N letters $F_N = F[y_1, \dots, y_N]$ to P_{N+1}

$$\Theta_N: F[y_1, \dots, y_N] \rightarrow P_{N+1}$$

defined on generators y_i in F_N by the formula

$$\Theta_N(y_i) = x_i.$$

The maps Θ_N are the subject of their authors’ papers [9; 10] where it is shown $\Theta_N: F_N \rightarrow P_{N+1}$ is faithful for every N . Three natural questions arise: (1) Why would one want to know whether Θ_n is faithful, (2) are there sensible applications and (3) why is Θ_n faithful? The answers to these three questions provide the main content of this expository article.




Figure 1: The braid x_i in P_{N+1}

3 On Θ_n

This section addresses one reason why the map Θ_n is faithful [9; 10]. The method of proof is to appeal to the structure of the Lie algebras obtained from the descending central series for both the source and the target of Θ_n . The structure of these Lie algebras is reviewed below.

Recall the descending central series of a discrete group π given by

$$\pi = \Gamma^1(\pi) \geq \Gamma^2(\pi) \geq \dots$$

where $\Gamma^i(\pi)$ is the subgroup of π generated by all commutators

$$[\dots[x_1, x_2]x_3] \dots x_t]$$

for $t \geq i$ with $x_i \in \pi$. The group $\Gamma^i(\pi)$ is a normal subgroup of π with the successive sub-quotients

$$gr_i(\pi) = \Gamma^i(\pi) / \Gamma^{i+1}(\pi)$$

which are abelian groups having additional structure as follows (cf Magnus, Karrass and Solitar [25]).

Consider the direct sum of all of the $gr_i(\pi) = \Gamma^i(\pi) / \Gamma^{i+1}(\pi)$ denoted

$$gr_*(\pi) = \bigoplus_{i \geq 1} \Gamma^i(\pi) / \Gamma^{i+1}(\pi).$$

The commutator function

$$[-, -]: \pi \times \pi \rightarrow \pi$$

given by

$$[x, y] = xyx^{-1}y^{-1},$$

passes to quotients to give a bilinear map

$$[-, -]: gr_s(\pi) \otimes_{\mathbb{Z}} gr_t(\pi) \rightarrow gr_{s+t}(\pi)$$

which satisfies both the antisymmetry law and Jacobi identity for a Lie algebra. (One remark about definitions: The abelian group $gr_*(\pi)$ is both a graded abelian group and a Lie algebra, but not a graded Lie algebra as the sign conventions do not work properly in this context. This situation can be remedied by doubling all degrees of elements in $gr_*(\pi)$.)

The associated graded Lie algebra obtained from the descending central series for the target yields Vassiliev invariants of pure braids by work of Kohno [23; 24]. This Lie algebra has been used by both Kohno and Drinfel'd [13] in their work on the KZ equations. The Lie algebra obtained from the descending central series of the free group F_N is a free Lie algebra by a classical result due to P Hall [18]; see also Serre [35].

The proof described next yields more information than just the fact that Θ_N is faithful. The method of proof gives a natural connection of Vassiliev invariants of braids to a classical spectral sequence abutting to the homotopy groups of the 2-sphere. Sections 5, 6 and 7 below provide an elucidation of this interconnection.

A discrete group π is said to be residually nilpotent provided

$$\bigcap_{i \geq 1} \Gamma^i(\pi) = \{\text{identity}\}$$

where $\Gamma^i(\pi)$ denotes the i -th stage of the descending central series for π . Examples of residually nilpotent groups are free groups, and P_n .

Lemma 3.1

(1) Assume that π is a residually nilpotent group. Let

$$\alpha: \pi \rightarrow G$$

be a homomorphism of discrete groups such that the morphism of associated graded Lie algebras

$$\text{gr}_*(\alpha): \text{gr}_*(\pi) \rightarrow \text{gr}_*(G)$$

is a monomorphism. Then α is a monomorphism.

(2) If π is a free group, and $\text{gr}_*(\alpha)$ is a monomorphism, then α is a monomorphism.

Thus one step in the proof of Theorem 3.2 below is to describe the map

$$\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$$

on the level of associated graded Lie algebras

$$\text{gr}_*(\Theta_n): \text{gr}_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{gr}_*(P_{n+1}).$$

Recall Artin's generators $A_{i,j}$ for P_{n+1} together with the projections of the $A_{i,j}$ to $\text{gr}_*(P_{n+1})$ labeled $B_{i,j}$ [9; 10].

Theorem 3.2 *The induced morphism of Lie algebras*

$$\text{gr}_*(\Theta_n): \text{gr}_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{gr}_*(P_{n+1})$$

satisfies the formula

$$\text{gr}_*(\Theta_n)(y_q) = \sum_{1 \leq i \leq n-q+1 < j \leq n+1} B_{i,j}.$$

Examples of this theorem are listed next.

Example 3.3

(1) If $q = 1$, then

$$\mathrm{gr}_*(\Theta_n)(y_1) = B_{1,n+1} + B_{2,n+1} + \cdots + B_{n,n+1}.$$

Thus if $q = 1$, and $n = 3$,

$$\mathrm{gr}_*(\Theta_3)(y_1) = B_{1,4} + B_{2,4} + B_{3,4}.$$

(2) If $q = 2$, then

$$\mathrm{gr}_*(\Theta_n)(y_2) = (B_{1,n+1} + B_{2,n+1} + \cdots + B_{n-1,n+1}) + (B_{1,n} + B_{2,n} + \cdots + B_{n-1,n}).$$

Thus if $q = 2$, and $n = 3$,

$$\mathrm{gr}_*(\Theta_3)(y_2) = (B_{1,4} + B_{2,4}) + (B_{1,3} + B_{2,3}).$$

(3) In general,

$$\mathrm{gr}_*(\Theta_n)(y_q) = V_{n-q+2} + V_{n-q+3} + \cdots + V_{n+1}$$

$$\text{where } V_r = B_{1,r} + B_{2,r} + \cdots + B_{r-1,r}.$$

Thus if $q = 3$, and $n = 3$,

$$\mathrm{gr}_*(\Theta_3)(y_3) = B_{1,2} + B_{1,3} + B_{1,4}.$$

To determine the map of Lie algebras with a more global view, the structure of the Lie algebra $\mathrm{gr}_*(P_n)$ is useful, and is given as follows. Let $L[S]$ denote the free Lie algebra over \mathbb{Z} generated by a set S . The next theorem was given in work of Kohno [23; 24], and Falk and Randell [17].

Theorem 3.4 *The Lie algebra $\mathrm{gr}_*(P_n)$ is the quotient of the free Lie algebra generated by $B_{i,j}$ for $1 \leq i < j \leq n$ modulo the infinitesimal braid relations (also called the horizontal 4T relations or Yang–Baxter–Lie relations)*

$$\mathrm{gr}_*(P_n) = L[B_{i,j} \mid 1 \leq i < j \leq n] / I$$

where I denotes the 2-sided (Lie) ideal generated by the infinitesimal braid relations as listed next:

- (1) $[B_{i,j}, B_{s,t}] = 0$, if $\{i, j\} \cap \{s, t\} = \emptyset$
- (2) $[B_{i,j}, B_{i,s} + B_{s,j}] = 0$
- (3) $[B_{i,j}, B_{i,t} + B_{j,t}] = 0$

A computation with these maps gives the following result of [9; 10] for which further connections are elucidated in sections 5 and 7.

Theorem 3.5 *The maps $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ on the level of associated graded Lie algebras*

$$\text{gr}_*(\Theta_n): \text{gr}_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{gr}_*(P_{n+1})$$

are monomorphisms. Thus the maps Θ_n are monomorphisms.

Remark 3.6 Two remarks concerning Θ_n are given next.

- (1) That Θ_n is a monomorphism identifies $F[y_1, y_2, \dots, y_n]$ as a free subgroup of rank n in P_{n+1} . However, there are other, natural free groups of rank n in P_{n+1} . These arise from the fibrations of Fadell and Neuwirth given by projection maps $p_i: \text{Conf}(\mathbb{R}^2, n+1) \rightarrow \text{Conf}(\mathbb{R}^2, n)$ which delete the i -th coordinate and have fibre of the homotopy type of an n -fold wedge of circles $\vee_n S^1$ (Fadell and Neuwirth [16]).

Let $q_i: P_{n+1} \rightarrow P_n$ denote the map induced by p_i on the level of fundamental groups. The kernel of q_i is a free group of rank n .

The image of Θ_n has a contrasting feature: Any composite of the natural projection maps $d_I: P_{n+1} \rightarrow P_2$ precomposed with Θ_n ,

$$F[y_1, y_2, \dots, y_n] \xrightarrow{\Theta_n} P_{n+1} \xrightarrow{d_I} P_2,$$

is a surjection.

- (2) The combinatorial behavior of the map $\text{gr}_*(\Theta_n)$ is intricate even though the definition is elementary as well as natural. For example, various powers of 2 arise in the computation of the map

$$\text{gr}_*(\Theta_n): \text{gr}_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{gr}_*(P_{n+1})$$

for $n > 2$. One example is listed next.

Example 3.7 $\Theta_3([[[y_1, y_2]y_3]y_2]) = -[[[\gamma_1, \gamma_2]\gamma_3]\gamma_2] + 2[[[\gamma_1, \gamma_3]\gamma_2]\gamma_2] + \delta$ where δ is independent of the other terms with $\gamma_1 = B_{1,4} + B_{2,4} + B_{3,4}$, $\gamma_2 = B_{3,4}$ and $\gamma_3 = B_{2,4} + B_{3,4}$. At first glance, these elements may appear to be "random". However, this formula represents a systematic behavior which arises naturally from kernels of certain morphisms of Lie algebras.

The crucial feature which makes the computations effective is the "infinitesimal braid relations". In addition, the behavior of the map $\text{gr}_*(\Theta_n)$ is more regular after restricting

to certain sub-Lie algebras arising in the third stage of the descending central series [9; 10]. Finally, the maps Θ_n also induce monomorphisms of restricted Lie algebras on passage to the Lie algebras obtained from the mod- p descending central series [9; 10].

4 Simplicial objects, and Δ -objects

Some basic constructions which are part of an algebraic topologist's toolkit are described in this section (Moore [31], Curtis [11], Bousfield and Kan [5], May [27], and Rourke and Sanderson [32]).

One of the great insights in classical homotopy theory, due to Moore and then 'extended' by Kan as well as Rourke and Sanderson is that not only are homology groups a combinatorial invariant, but so are homotopy groups. The basic combinatorial framework is that of a simplicial set developed in [31; 11; 5; 27] and a Δ -set developed in [32] both of which model the combinatorics of a simplicial complex.

Definition 4.1 A Δ -set is a collection of sets

$$K_\bullet = \{K_0, K_1, \dots\}$$

with functions, *face operations*,

$$d_i: K_t \rightarrow K_{t-1} \text{ for } 0 \leq i \leq t$$

which satisfy the identities

$$d_i d_j = d_{j-1} d_i$$

if $i < j$. A Δ -object in a small category \mathcal{C} is a Δ -set $K_\bullet = \{K_0, K_1, \dots\}$ with the K_i given by objects in \mathcal{C} and the maps $d_i: K_t \rightarrow K_{t-1}$ given by morphisms in \mathcal{C} . Thus, a Δ -group is a Δ -set for which all $d_i: K_t \rightarrow K_{t-1}$ are group homomorphisms.

Let $\Delta[n]$ denote the n -simplex with $\delta_i: \Delta[n] \rightarrow \Delta[n+1]$ the inclusion of the i -th face. Assume that each set S_i is given the discrete topology unless otherwise stated. The geometric realization of a Δ -set K_\bullet is given by

$$|K_\bullet| = (\coprod K_n \times \Delta[n]) / \sim$$

where \sim is the equivalence relation generated by requiring that if $x \in K_{n+1}$ and $\alpha \in \Delta[n]$, then

$$(d_i(x), \alpha) \sim (x, \delta_i(\alpha)).$$

Example 4.2 Examples of Δ -sets are given next.

- (1) A choice of Δ -set with exactly one 0-simplex and one 1-simplex has geometric realization given by the circle.
- (2) A natural example of a Δ -group arises from the pure braid groups $P_n(S) = \pi_1 \text{Conf}(S, n)$ for a path-connected surface S [3]. Define

$$\Delta_\bullet(S) \quad \text{by} \quad \Delta_n(S) = P_{n+1}(S)$$

the $(n + 1)$ -st pure braid group for the surface S .

There are $n + 1$ homomorphisms

$$d_i: P_{n+1}(S) \rightarrow P_n(S)$$

for $0 \leq i \leq n$ obtained by deleting the $(i + 1)$ -st strand in $P_{n+1}(S)$. The homomorphisms d_i are induced on the level of fundamental groups of configuration spaces by the projection maps

$$p_{i+1}: \text{Conf}(S, n + 1) \rightarrow \text{Conf}(S, n)$$

given by deleting the $(i + 1)$ -st coordinate and satisfy the identities required for a Δ -group.

In case $S = \mathbb{C}P^1 = S^2$, the associated Δ -group gives basic information about the homotopy groups of the 2-sphere [3]. In case S_g is a closed orientated surface, the Δ -group $\Delta_\bullet(S_g)$ does not admit the structure of a simplicial group as given in the next Definition 4.3.

Simplicial sets are defined next.

Definition 4.3 A *simplicial set* is

- (1) a Δ -set $K_\bullet = \{K_0, K_1, \dots\}$ together with
- (2) functions, *degeneracy operations*,

$$s_j: K_t \rightarrow K_{t+1} \quad \text{for} \quad 0 \leq j \leq t$$

which satisfy the *simplicial identities*

$$d_i d_j = d_{j-1} d_i \quad \text{if} \quad i < j,$$

$$s_i s_j = s_{j+1} s_i \quad \text{if} \quad i \leq j \quad \text{and}$$

$$d_i s_j = \begin{cases} s_{j-1} d_i & \text{if } i < j, \\ \text{identity} & \text{if } i = j \text{ or } i = j + 1, \\ s_j d_{i-1} & \text{if } i > j + 1. \end{cases}$$

A simplicial-object in a small category \mathcal{C} is a simplicial-set $K_\bullet = \{K_0, K_1, \dots\}$ with the K_i given by objects in \mathcal{C} for which both face maps $d_i: K_t \rightarrow K_{t-1}$ and degeneracies $s_j: K_t \rightarrow K_{t+1}$ are given by morphisms in \mathcal{C} .

Thus, a *simplicial-group*

$$G_\bullet = \{G_0, G_1, \dots\}$$

is a simplicial-set for which all of the G_i are groups with face and degeneracies given by group homomorphisms.

Example 4.4 Two examples of simplicial sets are given next.

- (1) The simplicial 1-simplex $\Delta[1]$ has two 0-simplices $\langle 0 \rangle$ and $\langle 1 \rangle$. The n -simplices of $\Delta[1]$ are sequences $\langle 0^i, 1^{n+1-i} \rangle$ for $0 \leq i \leq n+1$. All of the non-degenerate simplices are $\langle 0 \rangle$, $\langle 1 \rangle$, and $\langle 0, 1 \rangle$.
- (2) The simplicial circle S^1 is a quotient of the simplicial 1-simplex $\Delta[1]$ obtained by identifying $\langle 0 \rangle$ and $\langle 1 \rangle$. There are exactly two equivalence classes of non-degenerate simplices given by $\langle 0 \rangle$, and $\langle 0, 1 \rangle$. Furthermore, the simplicial circle S^1 is given in degree k by
 - (a) a single point $\langle 0 \rangle$ in case $k = 0$, and
 - (b) $n+1$ points $\langle 0^i, 1^{n+1-i} \rangle$ for $0 \leq i < n+1$ in case $k = n$ for which $\langle 0^{n+1} \rangle$ and $\langle 1^{n+1} \rangle$ are identified.

In what follows below, it is useful to label these simplices by

$$y_{n+1-i} = \langle 0^i, 1^{n+1-i} \rangle$$

for $0 < i \leq n+1$ with $y_0 = s_0^n(\langle 0 \rangle)$.

In addition, classical, elegant constructions for the standard simplicial n -simplex $\Delta[n]$ as well as the n -sphere are given in [5; 11; 27].

Homotopy groups are defined for any Δ -set, but the definition admits a simple description for simplicial sets which satisfy an additional condition.

Definition 4.5 A simplicial set K_\bullet is said to satisfy the *extension condition* if for every set of n -simplices $x_0, x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_{n+1} \in K_n$ which satisfy the compatibility condition

$$d_i x_j = d_{j-1} x_i, \quad i < j, \quad i \neq k, \quad j \neq k,$$

there exists an $(n+1)$ -simplex $x \in K_{n+1}$ such that $d_i x = x_i$ for $i \neq k$. A simplicial set which satisfies the extension condition is also called a *Kan complex*.

For each Kan complex K_\bullet with base-point $* \in K_0$ and $s_0^n(*) \in K_n$, the n -th homotopy set $\pi_n(K_\bullet, *)$ is defined for $n \geq 1$ as the equivalence classes of simplices $a \in K_n$, denoted $[a]$, such that

- (1) $d_i(a) = s_0^{n-1}(*)$ for all $0 \leq i \leq n$, and
- (2) two such simplices $a, a' \in K_n$ are equivalent provided there exists a simplex $y \in K_{n+1}$ such that $d_{n+1}y = a'$, $d_ny = a$, and $d_iy = s_0^n(*)$ for all $0 \leq i < n$.

If K_\bullet satisfies the extension condition, and $n \geq 1$, then $\pi_n(K_\bullet, *)$ is a group. In case $n \geq 2$, $\pi_n(K_\bullet, *)$ is also abelian.

Basic examples of (i) Kan complexes as well as (ii) Δ -groups which do not admit the structure of a Kan complex are given next.

Example 4.6 A simplicial group $G_\bullet = \{G_0, G_1, \dots\}$ always satisfies the extension condition as shown in [31].

- (1) If $G_\bullet = \{G_0, G_1, \dots\}$ is a simplicial group, then the n -th homotopy group of G_\bullet is the quotient group

$$\pi_n(G_\bullet) = Z_n/B_n$$

for which

$$Z_n = \bigcap_{0 \leq i \leq n} \text{Ker}(d_i: G_n \rightarrow G_{n-1}),$$

and
$$B_n = d_0 \left(\bigcap_{1 \leq i \leq n+1} \text{Ker}(d_i: G_{n+1} \rightarrow G_n) \right).$$

- (2) Additional, related information is stated next for certain Δ -groups $\Delta_\bullet = \{\Delta_0, \Delta_1, \dots\}$ which are not necessarily simplicial groups.
 - (a) In this case, the set of left cosets $\pi_n(\Delta_\bullet) = Z_n/B_n$ is still well-defined.
 - (b) The left cosets Z_n/B_n may not admit a natural structure as a group. One case occurs for S_g a closed, oriented surface with $\Delta_\bullet = \Delta_\bullet(S_g)$ as given in Example 4.2 and in [3].
 - (c) The special case of $S = S^2$ has further properties given in in [3] and below.

An example of a simplicial group obtained naturally from Artin's pure braid groups is described next.

Example 4.7 Consider Δ -groups with $\Delta_n(S) = P_{n+1}(S)$ as given in Example 4.2 for surfaces S . Specialize to the surface

$$S = \mathbb{R}^2.$$

In this case, there are also $n + 1$ homomorphisms

$$s_i: P_{n+1} \rightarrow P_{n+2}$$

obtained by “doubling” the $(i + 1)$ -st strand. The homomorphisms s_i are induced on the level of fundamental groups by the maps for configuration spaces

$$\mathbb{S}_i: \text{Conf}(\mathbb{R}^2, n + 1) \rightarrow \text{Conf}(\mathbb{R}^2, n + 2)$$

defined by the formula

$$\mathbb{S}_i(x_1, \dots, x_{n+1}) = (x_1, \dots, x_{i+1}, \lambda(x_{i+1}), x_{i+2}, \dots, x_{n+1})$$

where $\lambda(x_{i+1}) = x_{i+1} + (\epsilon, 0)$ for $(\epsilon, 0)$ a point in \mathbb{R}^2 with

$$\epsilon = (1/2) \cdot \min_{t \neq i+1} \|x_{i+1} - x_t\|.$$

The homomorphisms d_i and s_j satisfy the simplicial identities [9; 10; 3].

Thus the pure groups in case $S = \mathbb{R}^2$ provide an example of a simplicial group denoted

$$\text{AP}_\bullet \quad \text{with} \quad \text{AP}_n = P_{n+1}$$

for $n = 0, 1, 2, 3, \dots$.

Consider a pointed topological space $(X, *)$. The pointed loop space of X , $\Omega(X)$, has a natural multiplication coming from “loop sum” which is not associative, but homotopy associative. Milnor proved that the loop space of a connected simplicial complex is homotopy equivalent to a topological group [28]. James [22] proved that the loop space of the suspension of a connected CW-complex is naturally homotopy equivalent to a free monoid as explained by Hatcher [19, page 282]. Milnor realized that the James construction could be translated directly into the the language of simplicial groups as described next [30].

Definition 4.8 Let K_\bullet denote a pointed simplicial set (with base-point $* \in K_0$ and $s_0^n(*) \in K_n$). Define *Milnor’s free simplicial group* $F[K]_\bullet$ in degree n by

$$F[K]_n = F[K_n]/s_0^n(*) = 1.$$

Then $F[K]_\bullet$ is a simplicial group with face and degeneracy operations given by the natural multiplicative extension of those for K_\bullet . In addition, the face and degeneracy operations applied to a generator is either another generator or the identity element.

Example 4.9 An example of $F[K]_\bullet$ is given by $K_\bullet = S^1_\bullet$ the simplicial circle. Notice that $F[S^1]_\bullet$ in degree n is isomorphic to the free group $F[S^1]_n = F[y_1, \dots, y_n]$ by Example 4.4.

Milnor defined the geometric realization of a simplicial set $K_\bullet = \{K_0, K_1, \dots\}$ for which the underlying topology of K_\bullet is discrete [29]. Recall the inclusion of the i -th face $\delta_i: \Delta[n-1] \rightarrow \Delta[n]$ together with the projection maps to the j -th face $\sigma_j: \Delta[n+1] \rightarrow \Delta[n]$ [5; 11; 27].

Definition 4.10 The *geometric realization* of K_\bullet is

$$|K_\bullet| = (\coprod K_n \times \Delta[n]) / \sim$$

where \sim denotes the equivalence relation generated by requiring

- (1) if $x \in K_{n+1}$ and $\alpha \in \Delta[n]$, then $(d_i(x), \alpha) \sim (x, \delta_i(\alpha))$, and
- (2) if $y \in K_n$ and $\beta \in \Delta[n+1]$, then $(x, \sigma_j(\beta)) \sim (s_j(x), \beta)$.

Theorem 4.11 If K_\bullet is a reduced simplicial set (that is K_0 is equal to a single point $\{*\}$), then the geometric realization $|F[K_\bullet]|$ is homotopy equivalent to $\Omega\Sigma|K_\bullet|$. Thus the homotopy groups of $F[K_\bullet]$ (as given in [31] and Example 4.6) are isomorphic to the homotopy groups of the space $\Omega\Sigma|K_\bullet|$.

Example 4.12 Consider the special case of $K_\bullet = S^1_\bullet$. Then the geometric realization $|F[S^1_\bullet]|$ is homotopy equivalent to ΩS^2 , and there are isomorphisms

$$\pi_n F[S^1_\bullet] \rightarrow \pi_n \Omega S^2 \cong \pi_{n+1} S^2.$$

A partial synthesis of this information is given sections 5, 6 and 7.

5 Pure braid groups of surfaces as simplicial groups and Δ -groups

The homomorphism $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ which arises from the cabling operation described in Figure 1 satisfies the following properties.

- (1) The homomorphisms $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ give a morphism of simplicial groups

$$\Theta: F[S^1_\bullet] \rightarrow AP_\bullet$$

for which the homomorphism Θ_n is the restriction of Θ to $F[S^1_n]$.

- (2) By Theorem 3.5, the homomorphisms $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ are monomorphisms and so the morphism $\Theta: F[S^1_\bullet] \rightarrow AP_\bullet$ is a monomorphism of simplicial groups.

- (3) There is exactly one morphism of simplicial groups Θ with the property that $\Theta_1(y_1) = A_{1,2}$.

Thus, the picture given in Figure 1 is a description for generators of $F[S^1]_n$ in the simplicial group $F[S^1]_\bullet$. These features are summarized next.

Theorem 5.1 *The homomorphisms $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ (“pictured” in Figure 1) give the unique morphism of simplicial groups*

$$\Theta: F[S^1]_\bullet \rightarrow AP_\bullet$$

with $\Theta_1(y_1) = A_{1,2}$. The map Θ is an embedding. Hence the n -th homotopy group of $F[S^1]$, isomorphic to $\pi_{n+1}(S^2)$, is a natural sub-quotient of AP_\bullet . Furthermore, the smallest sub-simplicial group of AP_\bullet which contains the element $\Theta_1(y_1) = A_{1,2}$ is isomorphic to $F[S^1]_\bullet$.

On the other-hand, the homotopy sets for the Δ -group $\Delta_\bullet(S^2)$ are also giving the homotopy groups of the 2-sphere via a different occurrence of $F[S^1]_\bullet$. The homeomorphism of spaces

$$\text{Conf}(S^2, k) \rightarrow PGL(2, \mathbb{C}) \times \text{Conf}(S^2 - Q_3, k - 3)$$

for $k \geq 3$ and where Q_3 denotes a set of three distinct points in S^2 is basic for the next Theorem [3].

Theorem 5.2 *If $S = S^2$ and $n \geq 4$, then there are isomorphisms*

$$\pi_n(\Delta_\bullet(S^2)) \rightarrow \pi_n(S^2).$$

The descriptions of homotopy groups implied by these Theorems admit interpretations in terms of classical, well-studied features of the braid groups as given in the next section. An extension to all spheres is given in [9; 10] as pointed out in Section 7.

6 Brunnian braids, “almost Brunnian” braids, and homotopy groups

The homotopy groups of a simplicial group, or the homotopy sets of a Δ -group admit a combinatorial description as pointed out in Example 4.6. These homotopy sets are the set of left cosets Z_n/B_n where Z_n is the group of n -cycles and B_n is the group of n -boundaries for the Δ -group.

Recall Example 4.2 in which the Δ -group $\Delta_\bullet(S)$ is specified by $\Delta_n(S) = P_{n+1}(S)$ the $(n + 1)$ -stranded pure braid group for a connected surface S . The main point of this section is that the n -cycles Z_n are given by the "Brunnian braids" while the n -boundaries B_n are given by the "almost Brunnian braids", subgroups considered next which are also important in other applications (Magnum and Stanford [26]).

Definition 6.1 Consider the n -stranded pure braid group for any (connected) surface S , the fundamental group of $\text{Conf}(S, n)$. The group of Brunnian braids $\text{Brun}_n(S)$ is the subgroup of $P_n(S)$ given by those braids which become trivial after deleting any single strand. That is,

$$\text{Brun}_n(S) = \bigcap_{0 \leq i \leq n-1} \text{Ker}(d_i: P_n(S) \rightarrow P_{n-1}(S)).$$

The "almost Brunnian" $(n + 1)$ -stranded braid group is

$$\text{QBrun}_{n+1}(S) = \bigcap_{1 \leq i \leq n} \text{Ker}(d_i: P_{n+1}(S) \rightarrow P_n(S)).$$

The subgroup $\text{QBrun}_{n+1}(S)$ of $P_{n+1}(S)$ consists of those braids which are trivial after deleting any one of the strands $2, 3, \dots, n + 1$, but not necessarily the first.

Example 6.2 Consider the simplicial group AP_\bullet with

$$\text{AP}_n = P_{n+1}$$

for $n = 0, 1, 2, 3, \dots$ as given in Example 4.7.

In this case, notice that the map $d_0: \text{QBrun}_{k+2} \rightarrow \text{Brun}_{k+1}$ is a split epimorphism. Thus the homotopy groups of the simplicial group AP_\bullet are all trivial.

An inspection of definitions gives that

- (1) the group of n -cycles of $\Delta_\bullet(S)$, $Z_n(S)$, is precisely $\text{Brun}_{n+1}(S)$ while
- (2) the group of n -boundaries, $B_n(S)$, is exactly $d_0(\text{QBrun}_{n+2}(S))$.

This feature is recorded next as a lemma.

Lemma 6.3 Let S denote a connected surface with associated Δ -group $\Delta_\bullet(S)$ (as given in Example 4.2). Then the following hold.

- (1) The group of n -cycles $Z_n(S)$ is $\text{Brun}_{n+1}(S)$.
- (2) The group of boundaries $B_n(S)$ is $d_0(\text{QBrun}_{n+2}(S))$.

(3) *There is an isomorphism*

$$\pi_k(\text{AP}\bullet) \rightarrow \text{Brun}_{k+1}/d_0(\text{QBrun}_{k+2}).$$

Furthermore, $\pi_k(\text{AP}\bullet)$ is the trivial group.

(4) *There is an isomorphism of left cosets which is natural for pointed embeddings of connected surfaces S*

$$\pi_k(\Delta_*(S)) \rightarrow \text{Brun}_{k+1}(S)/d_0(\text{QBrun}_{k+2}(S)).$$

Properties of the Δ -group for the 2-sphere $S = \mathbb{C}\mathbb{P}^1 = S^2$ is the main subject of [3] where the next result is proven.

Theorem 6.4 *If $S = S^2$ and $k \geq 4$, then*

$$\pi_k(\Delta_\bullet(S^2)) = \text{Brun}_{k+1}(S^2)/d_0(\text{QBrun}_{k+2}(S^2))$$

is a group which is isomorphic to the classical homotopy group $\pi_k(S^2)$.

Furthermore, there is an exact sequence of groups

$$1 \rightarrow \text{Brun}_{k+2}(S^2) \rightarrow \text{Brun}_{k+1}(\mathbb{R}^2) \rightarrow \text{Brun}_{k+1}(S^2) \rightarrow \pi_k(S^2) \rightarrow 1.$$

The next lemma follows by a direct check of the long exact homotopy sequence obtained from the Fadell–Neuwirth fibrations for configuration spaces [16; 15].

Lemma 6.5 *If S is a surface not homeomorphic to either S^2 or $\mathbb{R}\mathbb{P}^2$, and $k \geq 3$, then $\text{Brun}_k(S)$ and $\text{QBrun}_k(S)$ are free groups. If S is any surface, and $k \geq 4$, then $\text{Brun}_k(S)$ and $\text{QBrun}_k(S)$ are free groups.*

Example 6.6 One classical example of a Brunnian braid group is $\text{Brun}_4(S^2)$ which is isomorphic to the principle congruence subgroup of level 4 in $PSL(2, \mathbb{Z})$ as given in Section 7 below.

One question below in Section 8 is to consider properties of the free groups obtained from the intersections $\Theta_k(F_k) \cap \text{Brun}_{k+1}$ as well as $\Theta_k(F_k) \cap d_0(\text{QBrun}_{k+2})$ where $\Theta_k: F[y_1, y_2, \dots, y_k] \rightarrow P_{k+1}$ is the homomorphism in Section 3. These groups are precisely the cycles and boundaries for $F[S^1]_\bullet$.

Lemma 6.7 *If $k \geq 3$, then $\Theta_k(F_k) \cap \text{Brun}_{k+1}$ as well as $\Theta_k(F_k) \cap d_0(\text{QBrun}_{k+2})$ are countably infinitely generated free groups.*

The standard Hall collection process or natural variations can be used to give inductive recipes rather than closed forms for generators. T Stanford has given a related elegant exposition of the Hall collection process [37]. The analogous process was applied by Cohen and Levi [7] to give group theoretic models for iterated loop spaces.

The connection of the homotopy groups of S^2 as well as the Lie algebra attached to the descending central series of the pure braid groups is discussed next.

Theorem 6.8 *The group $\Theta_k(F_k) \cap d_0(\text{QBrun}_{k+2})$ is a normal subgroup of $\Theta_k(F_k) \cap \text{Brun}_{k+1}$. There are isomorphisms*

$$\Theta_k(F_k) \cap \text{Brun}_{k+1} / \Theta_k(F_k) \cap d_0(\text{QBrun}_{k+2}) \rightarrow \pi_{k+1} S^2.$$

The method of proving that the maps $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ are monomorphisms via Lie algebras admits an interpretation in terms of classical homotopy theory. The method is to filter both simplicial groups $F[S^1]_\bullet$ and AP_\bullet via the descending central series, and then to analyze the natural map on the level of associated graded Lie algebras.

On the other-hand, the Lie algebra arising from filtering any simplicial group by its' descending central series gives the E^0 -term of the Bousfield–Kan spectral sequence for the simplicial group in question [5]. Similarly, filtering via the mod- p descending central series gives the classical unstable Adams spectral sequence [5; 11; 38].

Thus the method of proof of Theorem 3.5 is precisely an analysis of the behavior of the natural map $\Theta: F[S^1]_\bullet \rightarrow \text{AP}_\bullet$ on the level of the E^0 -term of the Bousfield–Kan spectral sequence. This method exhibits a close connection between Vassiliev invariants of pure braids and these natural spectral sequences. The next result is restatement of Theorem 3.5 proven in [9; 10].

Corollary 6.9 *The maps $\Theta_n: F[y_1, y_2, \dots, y_n] \rightarrow P_{n+1}$ on the level of associated graded Lie algebras*

$$\text{gr}_*(\Theta_n): \text{gr}_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{gr}_*(P_{n+1})$$

are monomorphisms. Thus the maps Θ_n induce embeddings on the level of the E^0 -term of the Bousfield–Kan spectral sequences for $E^0(\Theta): E^0(F[S^1]_\bullet) \rightarrow E^0(\text{AP}_\bullet)$.

7 Other connections

Several further connections, outlined next, emerged after this article was submitted.

Connection to principal congruence subgroups

One basic construction above is the Brunnian braid groups $\text{Brun}_k(S)$. Recently, the authors have proven (unpublished) that the Brunnian braid group $\text{Brun}_4(S^2)$ is isomorphic to the principal congruence subgroup of level 4 in $PSL(2, \mathbb{Z})$ [3].

This identification may admit an extension by considering the Brunnian braid groups $\text{Brun}_{2g}(S^2)$ as natural subgroups of mapping class groups for genus g surfaces. The subgroups $\text{Brun}_{2g}(S^2)$ may embed naturally in $Sp(2g, \mathbb{Z})$ via classical surface topology using branched covers of the 2-sphere (work in progress).

Connections to other spheres

The work above has been extended to all spheres as well as other connected CW-complexes [10]. One way in which other spheres arise is via the induced embedding of free products of simplicial groups

$$\Theta \amalg \Theta: F[S^1]_{\bullet} \amalg F[S^1]_{\bullet} \rightarrow \text{AP}_{\bullet} \amalg \text{AP}_{\bullet}.$$

The geometric realization of $F[S^1]_{\bullet} \amalg F[S^1]_{\bullet}$ is homotopy equivalent to $\Omega(S^2 \vee S^2)$ by Milnor's theorem stated above as 4.11.

Furthermore, $\Omega(S^2 \vee S^2)$ is homotopy equivalent to a weak infinite product of spaces $\Omega(S^n)$ for all $n > 1$.

Connection to certain Galois groups

Consider automorphism groups $\text{Aut}(H)$ where H is F_n or a completion of F_n given by either the pro-finite completion \widehat{F}_n or the pro- ℓ completion $(F_n)_{\ell}$. Certain Galois groups G are identified as natural subgroups of these automorphism groups by Belyi [2], Deligne [12], Drinfel'd [13; 14], Ihara [20; 21] and Schneps [33].

One example is Drinfel'd's Grothendieck–Teichmüller Galois group $G = \widehat{GT}$, a subgroup of $\text{Aut}(\widehat{F}_2)$.

Let $\text{Der}(L^R[V_n])$ denote the Lie algebra of derivations of the free Lie algebra $L^R[V_n]$ where V_n denotes a free module of rank n over R a commutative ring with identity. Two natural morphisms of Lie algebras which take values in $\text{Der}(L^R[V_n])$ occur in this context as follows.

Properties of the infinitesimal braid relations as stated in Theorem 3.4 above give a natural second map

$$\text{Ad}: gr_*(P_{n+1}) \rightarrow \text{Der}(L^{\mathbb{Z}}[V_n])$$

for which the kernel of Ad is precisely the center of $gr_*(P_{n+1})$ (Cohen and Prassidis [8]). Combining this last fact with Theorem 3.2 gives properties of the composite morphism of Lie algebras

$$gr_*(F_n) \xrightarrow{gr_*(\Theta_n)} gr_*(P_{n+1}) \xrightarrow{Ad} \text{Der}(L^{\widehat{\mathbb{Z}}}[V_n]).$$

Proposition 7.1 *If $n \geq 2$, the induced morphism of Lie algebras*

$$Ad \circ gr_*(\Theta_n): gr_*(F[y_1, y_2, \dots, y_n]) \rightarrow \text{Der}(L^{\widehat{\mathbb{Z}}}[V_n])$$

is a monomorphism.

In addition, certain Galois groups G above are filtered with induced morphisms of Lie algebras

$$gr_*(G) \rightarrow \text{Der}(L^{\widehat{\mathbb{Z}}}[V_n])$$

where $\widehat{\mathbb{Z}}$ denotes the pro-finite completion of the integers. One example is $G = \widehat{GT}$ with

$$gr_*(\widehat{GT}) \rightarrow \text{Der}(L^{\widehat{\mathbb{Z}}}[V_2])$$

as given in in [12; 20; 21; 34].

This raises the question of (i) whether the images of $Ad \circ gr_*(\Theta_2)$ and $gr_*(\widehat{GT})$ in $\text{Der}(L^{\widehat{\mathbb{Z}}}[V_2])$ have a non-trivial intersection and (ii) whether these intersections are meaningful.

8 Questions

The point of this section is to consider whether the connections between the braid groups and homotopy groups above are useful. Some natural as well as speculative problems are listed next.

- (1) The combinatorial problem of distinguishing elements in the pure braid groups has been well-studied. For example, the Lie algebra associated to the descending central series of the pure braid group P_n has been connected with Vassiliev theory and has been shown to be a complete set of invariants which distinguish all elements in P_n [24]. Furthermore, these Lie algebras have been applied to other questions arising from the classical KZ-equations [23; 13] as well as the structure of certain Galois groups [20; 12; 13; 14].

Find invariants of braids up to the coarser equivalence relation given in Theorem 6.8. In particular, can one identify the subset of Vassiliev invariants of braids which are "homotopy invariant"?

- (2) Give combinatorial properties of the natural map $\text{Brun}_{k+1}(\mathbb{R}^2) \rightarrow \text{Brun}_{k+1}(S^2)$ which provide information about the cokernel. Two concrete problems are stated next.
- (a) Give group theoretic reasons why the order of the 2-torsion in $\pi_*(S^2)$ is bounded above by 4 and why the p -torsion for an odd prime p is bounded above by p .
- (b) If $k + 1 \geq 5$, the image of $\text{Brun}_{k+1}(\mathbb{R}^2) \rightarrow \text{Brun}_{k+1}(S^2)$ is a normal subgroup of finite index.

This fact follows from Serre's classical theorem that $\pi_k(S^2)$ is finite for $k > 3$ and Theorems 5.2 and 6.4 proven in [3].

Do natural features of the braid groups imply this result?

- (3) Let F_n denote the image of $\Theta_n(F_n)$. Observe that the groups $\text{QBrun}_{n+2} \cap F_{n+1}$, and $\text{Brun}_{n+1} \cap F_n$ are free. Furthermore, there is a short exact sequence of groups

$$1 \rightarrow F_n \cap d_0(\text{QBrun}_{n+2}) \rightarrow F_n \cap \text{Brun}_{n+1} \rightarrow \pi_{n+1}S^2 \rightarrow 1$$

as well as isomorphisms

$$F_n \cap \text{Brun}_{n+1} / (F_n \cap d_0(\text{QBrun}_{n+2})) \rightarrow \pi_{n+1}S^2$$

by Theorem 6.8.

Consider the Serre 5-term exact sequence for the group extension directly above to obtain information about the induced surjection

$$H_1(F_n \cap \text{Brun}_{n+1}) \rightarrow \pi_{n+1}(S^2).$$

This 5-term exact sequence specializes to

$$\begin{aligned} H_2(\pi_{n+1}(S^2)) &\rightarrow H_1(F_n \cap d_0(\text{QBrun}_{n+2}))_{\pi_{n+1}(S^2)} \\ &\rightarrow H_1(F_n \cap \text{Brun}_{n+1}) \rightarrow \pi_{n+1}(S^2) \end{aligned}$$

where A_π denotes the group of co-invariants of a π -module A . Thus $\pi_{n+1}(S^2)$ is a quotient of the free abelian group $H_1(F_n \cap \text{Brun}_{n+1})$ with relations given by the image of the coinvariants $H_1(F_n \cap d_0(\text{QBrun}_{n+2}))_{\pi_{n+1}(S^2)}$.

Give combinatorial descriptions of the induced map on the level of the first homology groups

$$H_1(F_n \cap d_0(\text{QBrun}_{n+2})) \rightarrow H_1(F_n \cap \text{Brun}_{n+1}).$$

A similar problem arises with the the epimorphism $\text{Brun}_{n+1}(S^2) \rightarrow \pi_n S^2$ with kernel in the image of $\text{Brun}_{k+1}(\mathbb{R}^2)$ for $n + 1 \geq 5$.

- (4) This problem addresses a similarity between Vassiliev invariants and modular forms in the sense that both can be regarded as functions defined on certain braid groups. The initial ingredient here is given by the natural epimorphism

$$B_3 \rightarrow SL(2, \mathbb{Z})$$

with kernel given by the integers \mathbb{Z} . A generator for this kernel is equal to twice a generator of the center of B_3 .

Recall that Shimura [36] gives isomorphisms

$$H^1(SL(2, \mathbb{Z}); \mathbb{R}[x_1, x_2]) \rightarrow \mathcal{M},$$

and
$$H^1(SL(2, \mathbb{Z}); \text{Sym}^k(x_1, x_2)) \rightarrow \mathcal{M}_{2k+2}$$

with the following properties.

- (a) The action of $SL(2, \mathbb{Z})$ on $\mathbb{R}[x_1, x_2]$ is specified by the tautological representation on the two dimensional vector space with basis x_1, x_2 extended multiplicatively to the polynomial ring $\mathbb{R}[x_1, x_2]$.
- (b) The module $\text{Sym}^k(x_1, x_2)$ denotes the $SL(2, \mathbb{Z})$ -submodule of $\mathbb{R}[x_1, x_2]$ given by polynomials of classical degree k .
- (c) The vector space \mathcal{M}_{2k+2} denotes the summand of the ring of modular forms of weight $2k + 2$ (with Shimura's weight convention).

Also recall that the first cohomology group $H^1(SL(2, \mathbb{Z}); \mathbb{R}[x_1, x_2])$ is given by the quotient of the module of crossed homomorphisms $SL(2, \mathbb{Z}) \rightarrow \mathbb{R}[x_1, x_2]$ modulo principal crossed homomorphisms. Thus elements in the classical ring of modular forms can be regarded as equivalence classes of certain functions defined on $SL(2, \mathbb{Z})$.

Furthermore, there are isomorphisms

$$H^1(B_3; \text{Sym}^k(x_1, x_2)) \rightarrow E[u] \otimes \mathcal{M}_{2k+2}$$

where $E[u]$ denotes an exterior algebra with u of degree 1. It is natural to ask whether and how crossed homomorphisms representing $H^1(B_3; \text{Sym}^k(x_1, x_2))$ distinguish braids in B_3 .

There is also a natural analogue for subgroups of B_{2g+2} as follows. Determine $H^1(B_{2g+2}; \mathbb{R}[x_1, \dots, x_{2g}])$ where B_{2g+2} acts via a natural symplectic representation on a vector space with basis $\{x_1, \dots, x_{2g}\}$ the generating module for the polynomial ring and $\mathbb{R}[x_1, \dots, x_{2g}]$. How do the natural crossed homomorphisms distinguish braids?

- (5) Consider Brunnian braids Brun_k . Fix a braid γ with image in the k -th symmetric group Σ_k given by a k -cycle. For any braid α in Brun_k , the braid closure

of $\alpha \circ \gamma$ is a knot. Describe features of these knots or those obtained from the analogous constructions for $\Theta_k(F_{k-1}) \cap \text{Brun}_k$. Where do these fit in Budney’s description of the space of long knots [6]?

- (6) Let $L[V]$ denote the free Lie algebra over the integers generated by the free abelian group V . Let $\text{Der}(L[V])$ denote the Lie algebra of derivations of $L[V]$ and consider the classical adjoint representation

$$\text{Ad}: L[V] \rightarrow \text{Der}(L[V]).$$

Recall that the map $\Theta_k: F_k \rightarrow P_{k+1}$ induces a monomorphism of Lie algebras $\text{gr}_*(\Theta_k): \text{gr}_*(F_k) \rightarrow \text{gr}_*(P_{k+1})$ where $\text{gr}_*(F_k)$ is isomorphic to the free Lie algebra $L[V_k]$ with V_k a free abelian group of rank k . In addition, properties of the "infinitesimal braid relations" give a representation

$$\rho_k: \text{gr}_*(P_{k+1}) \rightarrow \text{Der}(L[V_k])$$

appearing in work on certain Galois groups [20] (with the integers \mathbb{Z} replaced by the profinite completion of \mathbb{Z}) and, when restricted to the integers \mathbb{Z} , also addressed in [8].

Identify F_k with $\Theta_k(F_k)$ in what follows below.

Give methods to describe combinatorial properties of the composite

$$\text{gr}_*(F_k \cap \text{Brun}_{k+1}) \xrightarrow{\text{gr}_*(i_k)} \text{gr}_*(F_k) \xrightarrow{\text{gr}_*(\Theta_k)} \text{gr}_*(P_{k+1}) \xrightarrow{\rho_k} \text{Der}(L[V_k])$$

where $i_k: F_k \cap \text{Brun}_{k+1} \rightarrow F_k$ is the natural inclusion. Let Φ_{k+1} denote this composite. When restricted to $\text{gr}_*(F_k) = L[V_k]$, this map is a monomorphism. Give methods to describe the sub-quotient

$$\Phi_{k+1}(\text{gr}_*(F_k \cap \text{Brun}_{k+1})) / \Phi_{k+1} \text{gr}_*(F_k \cap d_0(\text{QBrun}_{k+2})).$$

- (7) Assume that the pure braid groups $P_n(S)$ are replaced by either their pro-finite completion $\widehat{P_n(S)}$ or their pro- ℓ completions. Describe the associated changes for the homotopy groups arising in Theorems 5.2, 6.4, or 5.1. For example, is the torsion in these homotopy groups left unchanged by replacing $P_n(S)$ by $\widehat{P_n(S)}$?

9 Acknowledgements

Some of the work on this project was accomplished at the University of Tokyo during 2004. The first author was partially supported by the National Science Foundation under Grant No. 9704410 as well as Darpa Grant No. 2006-06918-01, and the Institute

for Advanced Study. The second author was partially supported by the Academic Research Fund of the National University of Singapore No. R-146-000-048-112.

References

- [1] **E Artin**, *Theorie die Zöpfe*, Hamburg Abhandlung (1924)
- [2] **G V Belyĭ**, *Galois extensions of a maximal cyclotomic field*, *Izv. Akad. Nauk SSSR Ser. Mat.* 43 (1979) 267–276, 479 MR534593 Russian; English translation: *Math. USSR Izv.* 14 (1980) 247–256
- [3] **A J Berrick, F R Cohen, Y L Wong, J Wu**, *Configurations, braids, and homotopy groups*, *J. Amer. Math. Soc.* 19 (2006) 265–326 MR2188127
- [4] **J S Birman**, *Braids, links, and mapping class groups*, *Ann of Math Studies* 82, Princeton University Press (1974) MR0375281
- [5] **A K Bousfield, D M Kan**, *Homotopy limits, completions and localizations*, *Lecture Notes in Mathematics* 304, Springer, Berlin (1972) MR0365573
- [6] **R Budney**, *The topology of knotspaces in dimension 3* arXiv:math.GT/0506523
- [7] **F R Cohen, R Levi**, *Combinatorial models for iterated loop spaces* Available at <http://www.maths.abdn.ac.uk/~ran/papers.html>
- [8] **F R Cohen, S Prassidis**, *On injective homomorphisms for pure braid groups, and associated Lie algebras*, *J. Algebra* 298 (2006) 363–370 MR2217620
- [9] **F R Cohen, J Wu**, *On braid groups, free groups, and the loop space of the 2-sphere*, from: “Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001)”, *Progr. Math.* 215, Birkhäuser, Basel (2004) 93–105 MR2039761
- [10] **F R Cohen, J Wu**, *Braid groups, free groups, and the loop space of the 2-sphere* arXiv:math.AT/0409307
- [11] **E B Curtis**, *Simplicial homotopy theory*, *Advances in Math.* 6 (1971) 107–209 (1971) MR0279808
- [12] **P Deligne**, *Le groupe fondamental de la droite projective moins trois points*, from: “Galois groups over \mathbb{Q} (Berkeley, CA, 1987)”, *Math. Sci. Res. Inst. Publ.* 16, Springer, New York (1989) 79–297 MR1012168
- [13] **V G Drinfel’d**, *Quasi-Hopf algebras and Knizhnik-Zamolodchikov equations*, from: “Problems of modern quantum field theory (Alushta, 1989)”, (A A Belavin, A U Klimyk, A B Zamolodchikov, editors), *Res. Rep. Phys.*, Springer, Berlin (1989) 1–13 MR1091757
- [14] **V G Drinfel’d**, *On quasitriangular quasi-Hopf algebras and on a group that is closely connected with $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$* , *Algebra i Analiz* 2 (1990) 149–181 MR1080203 Russian; English translation: *Leningrad Math. J.* 2 (1991) 829–860

- [15] **E R Fadell, S Y Hussein**, *Geometry and topology of configuration spaces*, Springer Monographs in Mathematics, Springer, Berlin (2001) MR1802644
- [16] **E Fadell, L Neuwirth**, *Configuration spaces*, Math. Scand. 10 (1962) 111–118 MR0141126
- [17] **M Falk, R Randell**, *The lower central series of a fiber-type arrangement*, Invent. Math. 82 (1985) 77–88 MR808110
- [18] **P Hall**, *A contribution to the theory of groups of prime power order*, Proc. London Math. Soc. 36 (1933) 29–95
- [19] **A Hatcher**, *Algebraic topology*, Cambridge University Press, Cambridge (2002) MR1867354
- [20] **Y Ihara**, *On the stable derivation algebra associated with some braid groups*, Israel J. Math. 80 (1992) 135–153 MR1248930
- [21] **Y Ihara**, *On the embedding of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ into $\widehat{\text{GT}}$* , from: “The Grothendieck theory of dessins d’enfants (Luminy, 1993)”, LMS Lecture Note Ser. 200, Cambridge Univ. Press, Cambridge (1994) 289–321 MR1305402 With an appendix: the action of the absolute Galois group on the moduli space of spheres with four marked points by M Emsalem and P Lochak
- [22] **IM James**, *Reduced product spaces*, Ann. of Math. (2) 62 (1955) 170–197 MR0073181
- [23] **T Kohno**, *Linear representations of braid groups and classical Yang-Baxter equations*, from: “Braids (Santa Cruz, CA, 1986)”, Contemp. Math. 78, Amer. Math. Soc., Providence, RI (1988) 339–363 MR975088
- [24] **T Kohno**, *Vassiliev invariants and de Rham complex on the space of knots*, from: “Symplectic geometry and quantization (Sanda and Yokohama, 1993)”, Contemp. Math. 179, Amer. Math. Soc., Providence, RI (1994) 123–138 MR1319605
- [25] **W Magnus, A Karrass, D Solitar**, *Combinatorial group theory: Presentations of groups in terms of generators and relations*, Interscience Publishers [John Wiley & Sons] (1966) MR0207802
- [26] **B Mangum, T Stanford**, *Brunnian links are determined by their complements*, Algebr. Geom. Topol. 1 (2001) 143–152 MR1823496
- [27] **J P May**, *Simplicial objects in algebraic topology*, Van Nostrand Mathematical Studies 11 (1967) MR0222892
- [28] **J Milnor**, *Construction of universal bundles. I*, Ann. of Math. (2) 63 (1956) 272–284 MR0077122
- [29] **J Milnor**, *The geometric realization of a semi-simplicial complex*, Ann. of Math. (2) 65 (1957) 357–362 MR0084138

- [30] **J Milnor**, *On the construction $F[K]$* , from: “Algebraic topology—a student’s guide”, (JF Adams, editor), LMS Lecture Note Ser. 4, Cambridge University Press (1972) 119–136 MR0445484
- [31] **J C Moore**, *Homotopie des complexes monoïdaux*, Séminaire H Cartan (1954/55)
- [32] **C P Rourke, B J Sanderson**, Δ -sets. I. *Homotopy theory*, Quart. J. Math. Oxford Ser. (2) 22 (1971) 321–338 MR0300281
- [33] **L Schneps**, *Groupe de Grothendieck-Teichmüller et automorphismes de groupes de tresses*, C. R. Acad. Sci. Paris Sér. I Math. 317 (1993) 729–734 MR1244420
- [34] **L Schneps**, *On the Poisson bracket on the free Lie algebra in two generators*, J. Lie Theory 16 (2006) 19–37 MR2196410
- [35] **J-P Serre**, *Lie algebras and Lie groups*, Lectures given at Harvard University 1964, Benjamin, New York–Amsterdam (1965) MR0218496
- [36] **G Shimura**, *Introduction to the arithmetic theory of automorphic functions*, Publ. Math. Soc. Japan 11, Princeton University Press (1994) MR1291394 Reprint of the 1971 original, Kanô Memorial Lectures, 1
- [37] **T Stanford**, *Brunnian braids and some of their generalizations*, to appear in Bull. London Math. Soc. arXiv:math.GT/9907072
- [38] **J Wu**, *Combinatorial descriptions of homotopy groups of certain spaces*, Math. Proc. Cambridge Philos. Soc. 130 (2001) 489–513 MR1816806

Department of Mathematics, University of Rochester
Rochester, NY 14627, USA

Department of Mathematics, National University of Singapore
Singapore 117543, Republic of Singapore

cohf@math.rochester.edu, matwuj@nus.edu.sg

<http://www.math.rochester.edu/people/faculty/cohf/>, <http://www.math.nus.edu.sg/~matwujie>

Received: 11 July 2006 Revised: 29 May 2007

