

Kernel(J) warns of false vacua

MICHAEL FREEDMAN

JHC Whitehead defined a map $J_r: \pi_r(\mathrm{SO}) \rightarrow \pi_r^s$ from the homotopy of the special orthogonal group to the stable homotopy of spheres. Within a toy model we show how the known computation for $\mathrm{kernel}(J)$ leads to nonlinear σ -models with spherical source (space) and spherical target which admit false vacua separated from the true vacuum by an energy barrier. In this construction, the dimension of space must be at least 8 and the dimension of the σ -model target at least 5.

55Q50; 19L20, 81P99

1 Introduction

Homotopy theory and, in particular, K -theory have recently played a prominent role in both condensed matter (CM) and high energy physics (Kane and Mele [7], Kitaev [8], Ryu, Schnyder, Furusaki and Ludwig [12], Horava [6], and Minasian and Moore [10]). In CM applications, the discussion can often be reduced to the study of nonlinear σ -models with symmetric space targets (Ludwig, Moore and Ryu [9]). In physical applications (Freedman et al [2]), there may be a potential function V on the target symmetric space which can lead to distinct phases separated by an energy barrier. The purpose of this paper is to raise the possibility that even when there is *no* potential on the target symmetric space X , and the physical space B is as simple as possible—say a sphere or Euclidean space—the energy landscape on (smooth) maps $\mathcal{M}(B, X)$, maps f from the physical space B to X , may have local minima, effectively creating sectors of false vacua. Similarly, the space-time action of any field $F: B \times [0, 1] \rightarrow X$ interpolating between distinct vacua may be large compared to fields F' remaining within a single vacuum sector. Such an F is a novel kind of instanton protected by the intrinsic inefficiency of *any* homotopy between distinct vacua. This may be contrasted with conventional instantons, for example in Yang–Mills theory, where the instanton results from a *willfully* inefficient null homotopy:

$$F: S^3 \times [0, 1] \rightarrow S^4 \cong HP^1 \subset HP^\infty,$$

where $F(S^3 \times 0) = F(S^3 \times 1) = *$, is *chosen* to have nonzero degree.

I will explain why no such minima are expected to arise when $\dim B, \dim X \leq 2$. After this, as the dimension increases, little is known until we meet the example, f_0 , defined after equation (3). The focus of this paper is the surprisingly complex geometry necessarily associated to *any* null homotopy of $f_0: B \rightarrow X$, where $B = S^8$ and $X = S^5$, and where f_0 is a specific field on an 8–dimensional spherical space taking values in the 5–dimensional sphere. Of course, there is the question of what energy functional to use for a general $f: S^8 \rightarrow S^5$. Something like

$$E(f) = m^{(n-9)\hbar} \int_{S^3} d^8x \sqrt{g} |\nabla f|^n, \quad 8 \leq n < \infty,$$

where we have chosen units with $c = 1$ and m a constant with units of length, is a candidate (note that for $n < 8$, rescaling f toward a point $x \in S^8$ by precomposing with a conformal transformation with x a repelling fixed point makes E approach zero). Another possibility related to the choice $n = \infty$ is to let E be the Lipschitz constant of f (times a unit of energy). These choices entail serious analytical difficulties which are circumvented here by choosing a surrogate energy

$$(1) \quad E_{\text{top}} = \max_{x \in S^5} \sum_{i=0}^7 b_i(f^{-1}(x)),$$

f is presumed to be a smooth (C^∞) function and b_i the i^{th} Betti number, $b_i = \text{rank } H^*(X; R)$ (I have now dropped coupling constants and reference to units). We use the Čech cohomology \check{H}^i because it is appropriate for the nonregular point inverse images, which may be arbitrary closed subsets of B .

Similarly, define the action S_{top} of a field F on space-time $F := f_t: S^8 \times [0, 1] \rightarrow S^5$ by

$$(2) \quad S_{\text{top}}(F) = \max_{x \in S^5} \sum_{i=0}^7 b_i(F^{-1}(x)).$$

Both E_{top} and S_{top} are nonlocal—that is not integrated up from a locally defined quantity—but nevertheless seem to capture an essential feature of more physical Hamiltonians and Lagrangians, respectively: maps of high energy (action) are generally those with complicated point preimages (see [Figure 1](#)). With this definition, f_0 has $E_{\text{top}}(f_0) = 2$ and can be deformed to the constant map f_1 , that is the true vacuum, with $E_{\text{top}}(f_1) = 1$, but during the deformation we find that $E_{\text{top}}(f_{t_0}) > 2$ for some $t_0 \in (0, 1)$. Thus $E_{\text{top}}(f_{t_0})$ is an energy barrier between a false vacuum f_0 and the true vacuum. Similarly, the action must be large for any F interpolating between f_0 and f_1 : $S_{\text{top}}(F) \geq 23$ whereas for paths staying within a vacua, for example $S_{\text{top}}((F(y, t)) := f_0(y)) = 2$ and $S_{\text{top}}((F(y, t)) := f_1(y)) = 1$, the action can be

much smaller. I should emphasize that although topology is used to define E_{top} and S_{top} , these quantities are *not* homotopy invariants: as f_0 is deformed, there will be times when $E_{\text{top}}(f_t)$ jumps; similarly for $S_{\text{top}}(F)$ if F were also deformed. What E_{top} and S_{top} *do* depend on is the topological complexity of the inverse images of points $x \in S^5$, and these can jump whenever x is (transiently) a singular value.

So one may understand (Figure 1) how preimage complexity roughly encodes energy. The surprise is that it may be necessary to increase the energy en route to decreasing it. It would be as if in passing from Figure 1(a) to Figure 1(c) we had to go through Figure 1(b).




Figure 1

If we distance ourselves from energy functionals, the basic idea of needing to “pass over a saddle” on the way to lowering complexity can easily be illustrated using immersions. The path components of immersions $\mathcal{I}(S^1, \mathbb{R}^2)$ are known to be indexed by the winding number in \mathbb{Z} . It is easy (Figure 2) to find two immersions α_0 and α_1 with equal winding where the “complexity” as measured by the number of multiple points, must increase during any regular homotopy $\alpha_t, t \in [0, 1]$, beyond the initial or final values.




Figure 2

Unlike E_{top} , the complexity in this example seems, in spirit, opposite to conventional energy functionals: for example, a constant map, though not an immersion, would have infinite multiplicity and hence high, not low, energy.

It is hoped that some insight can be gained even from the “toy functionals” E_{top} and S_{top} . The key to the high dimensional example is the noninjectivity of Whitehead’s J -homomorphism from homotopy theory.

In summary, this paper is a message from topology to physics that fields *might* become trapped in local minima for the energy of a nonlinear σ -model, even when the space-time and the target space are as homogeneous as possible: both spheres. In our example, the dimensions are a bit high (except for string theorists) and the action is a toy. The challenge for the reader is to determine if this same phenomenon can occur, or contrarywise is excluded, in more conventional and perhaps lower dimensional, physical situations.

2 The Example

In homotopy theory, the J -homomorphism is the map $J_r: \pi_r(\text{SO}) \rightarrow \pi_r^s$ from the homotopy of the special orthogonal group to the r^{th} stable stem π_r^s ($:= \pi_{r+d}(S^d)$ for $d > r + 1$). We can interpret $\alpha \in \pi_r(\text{SO})$ as a (stable) normal framing of the r -sphere S^r and $\beta \in \pi_r^s$ can, by the Pontrjagin–Thom construction, be encoded as a normally framed r -manifold M^r . The Pontrjagin–Thom construction associates to $\beta: S^{r+d} \rightarrow S^d$, the preimage of a regular value $* \in S^d$, $M^r := \beta^{-1}(*)$. The normal bundle of M^r in S^{r+d} is framed by pulling back the normal framing of $*$ in S^d . For $d > r + 1$, the stable normal cobordism class of M^r is equivalent to the homotopy class of β . In these terms, the J -homomorphism simply includes spheres (with framed normal bundle) into manifolds (with framed normal bundle). It is known that J_r is injective unless $r = 4n - 1$, $n \geq 1$ and is the epimorphism $J_r: \mathbb{Z} \rightarrow \mathbb{Z}_{4n/B_{2n}}$, where B_{2n} is the $2n^{\text{th}}$ Bernoulli denominator. The first relevant images of J are $\text{Im} J_3 = \mathbb{Z}_{24}$, $\text{Im} J_7 = \mathbb{Z}_{240}$ and $\text{Im} J_9 = \mathbb{Z}_{504}$. The discussion requires only that J_r have a kernel, so there are examples of exotic local minima for the obvious dimensional extension¹ of E_{top} (and S_{top} after crossing the source with $[0, 1]$) to maps $S^{2(4n-1)+2+i} \rightarrow S^{(4n-1)+2+i}$ for all $n > 0$ and $i \geq 0$. Increasing n means going to a higher stable stem and increasing i merely means suspending maps within that stable stem. I analyze the case $n = 1$ and $i = 0$. Nothing changes as i increases, however with increasing n the characteristic class computations are slightly different and more significantly a refinement coming from Seiberg–Witten theory is not present. In all cases, however, there is a similar energy barrier.

¹Extend the definition of E_{top} and S_{top} given in equations (1) and (2) by extending the range of the summations to (spatial dimension) $- 1$.

The homotopy group $\pi_3(\text{SO}) \cong \mathbb{Z}$ is generated by the composition

$$(3) \quad g: S^3 \cong SU(2) \xrightarrow{\text{double cover}} \text{SO}(3) \hookrightarrow \text{SO}.$$

$24g$ is obtained by precomposing g with $(\phi, \theta_1; \theta_2) \mapsto (24\phi, \theta_1; \theta_2)$ in spherical coordinates. The map we study $f_0: S^8 \rightarrow S^5$ may be defined by writing $S^8 = S^3 \wedge S^4$, where the join symbol \wedge indicates the space of line segments “joining” the two factors: $P \wedge Q := P \times Q \times [0, 1] / (p, q, 0) = (p, q', 0)$ and $(p, q, 1) \equiv (p', q, 1)$, for all $p, p' \in P$ and all $q, q' \in Q$. In these coordinates, $f_0(S^3) = +$ and $f_0(S^4) = -$, $+(-)$ the north (south) pole of S^5 . The normal direction to S^3 in the join consists of a unit 5-disk D_s^5 at every point $s \in S^3$ and f_0 wraps this D_s^5 degree = one over S^5 by exponentiation. The only ambiguity is that we have not yet said how D_s^5 is identified with the unit disk in the target space $T_+(S^5)$ as a function of s . When $s = * \in S^3$, the base point, say the identity of $SU(2)$, the identification is an arbitrary isometry. For general s , precompose this arbitrary identification with $24g$ as defined above, where SO acts on \mathbb{R}^5 through the intermediate $\text{SO}(5)$, $\text{SO}(3) \subset \text{SO}(5) \subset \text{SO}$.

By inspection, every point preimage of f_0 is a 3-sphere except the preimage of the south pole, which is a 4-sphere. Since

$$\text{rank } \check{H}^i(S^n; R) = \begin{cases} 1 & \text{for } i = 0 \text{ or } n, \\ 0 & \text{otherwise,} \end{cases}$$

we see that $S_{\text{top}}(f_0) = 2$.

Since it is known that $J(24g) \simeq f_0$ is homotopically trivial, f_0 is homotopic to f_1 , the constant map $S^8 \rightarrow S^5$, taking each point of S^8 to the south pole $-$ of S^5 . The only nonempty preimage of f_1 is $f_1^{-1}(-) = S^8$, so $E_{\text{top}}(f_1) = \sum_{i=0}^7 b_i = 1$. We now state:

Theorem 1 *If f_t is a smooth family, $F := f_t: S^8 \times [0, 1] \rightarrow S^5$ with f_1 constant, then $S_{\text{top}}(F) \geq 23$.*

Proof By Sard’s theorem, the regular values on S^5 are an open dense set of full Lebesgue measure; let $* \in S^5 \setminus (-)$ be one. By the Inverse Function Theorem, $F^{-1}(x)$ is a smooth, normally framed 4-dimensional manifold $M \subset S^8 \times [0, 1]$, with $\partial M = S^3 \subset S^8 \times 0$. Cap M off with a 4-disk to obtain a smooth, closed 4-manifold \hat{M} . Since \hat{M} has a normal framing away from a point, the normal characteristic classes $w_1(\nu\hat{M}) = w_2(\nu\hat{M}) = 0$. By the Whitney sum formula those tangential classes also vanish $w_1(\tau\hat{M}) = w_2(\tau\hat{M}) = 0$, so \hat{M} is spin. The first obstruction to trivializing the normal frame bundle over \hat{M} is $24\gamma \in H^4(\hat{M}; \pi_3(\text{SO})) \cong \mathbb{Z}$, where γ is a generator, $\gamma = \pm 1 \in \mathbb{Z}$. The first Pontrjagin class $p_1(\tau\hat{M})$ is twice this obstruction,

$p_1(\tau_{\hat{M}}) = \pm 48\gamma$. On the other hand, in dimension 4 the Hirzebruch L -genus reduces to:

$$(4) \quad \text{signature of cup-product on } H_2(\hat{M}; \mathbb{Z})/\text{torsion} =: \sigma(\hat{M}) = \frac{p_1(\hat{M})}{3},$$

so $\sigma(\hat{M}) = \pm 16$. The $|\text{signature}|$ is a lower bound on the second Betti number $b_2(\hat{M})$. But, by excision, $b_2(M) = b_2(\hat{M})$. Putting this together we find

$$(5) \quad b_2(M) \geq 16$$

since $b_0(M) \geq 1$. We may conclude that

$$(6) \quad \sum_{i=0}^7 b_i(M) \geq 17.$$

Since S_{top} is defined as the maximal such sum over point preimages, $S_{\text{top}}(F) \geq 17$.

Because \hat{M} is smooth, spin and 4-dimensional (since we are considering kernel J_3), there is more refined information on the lower bound to b_2 available from the Seiberg–Witten equations. It is known that \hat{M} must have b_2 as large as that of the $K3$ surface, $b_2(\hat{M}) \geq 22$. This follows from Furuta’s “10/8–theorem” [5] and also from previous unpublished work of Peter Kronheimer. The estimate $b_2(\hat{M}) \geq 22$ is Donaldson’s “Theorem C” in the special case $\pi_1(\hat{M}) = 0$. This estimate implies $S_{\text{top}}(F) \geq 23$. \square

This is the Lagrangian result. The corresponding energy barrier is identified by the next theorem.

Theorem 2 *If f_t is a smooth family $F := f_t: S^8 \times [0, 1] \rightarrow S^5$ with f_0 as defined and f_1 constant, then there exists a $t \in (0, 1)$ where $E_{\text{top}}(f_t) \geq 3$. Recall $E_{\text{top}}(f_0) = 2$ and $E_{\text{top}}(f_1) = 1$.*

Proof Begin with the same $(M, \partial M)$, $\partial M = S^3$, as in the proof of [Theorem 1](#), except now restrict M to be only the connected component containing ∂M . Let $h: M \rightarrow [0, 1]$ be the inclusion $M \subset S^8 \times [0, 1]$ followed by projection to the second factor. Intuitively, we may think of h as a Morse function, but this cannot in fact be assumed. All we really know is $h^{-1}(0) = \partial M \cong S^3$, $h^{-1}(1) = \emptyset$ and h is smooth. A lower bound to

$$b := \max_{t \in [0, 1]} \sum_{i=1}^7 b_i(h^{-1}(t))$$

is the lower bound to the barrier $\max_{t \in [0, 1]} E_{\text{top}}(f_t)$ obtained by restricting attention to preimages of $* \in S^5$. We show $b \geq 3$ by assuming for a contradiction that $b = 2$.

Let $x_0 \in [0, 1]$ be the maximum value for which $h^{-1}(x) \neq \emptyset$. Since we assumed M connected, for all $x \in [0, x_0]$ we have $h^{-1}(x) \neq \emptyset$. Let $x_1 \in (0, x_0)$ be any regular value for h . $h^{-1}(x_1)$ is a smooth, closed 3–manifold so by the assumption $b = 2$, $h^{-1}(x_1)$ must be a real (equivalent rational) homology 3–sphere (otherwise its first and second Betti numbers would contribute to the sum).

A key property of Čech cohomology is that it commutes with inverse limits. This will now be used twice.

Lemma 1 *For all x_1, x_2 h –regular values in $[0, x_0]$, $h^{-1}[x_1, x_2]$ must have the real Čech cohomology of S^3 .*

Proof Suppose this fails for some $[x_1, x_2]$ pick an h –regular $x_3 \in (x_1, x_2)$ near $(x_1 + x_2)/2$ and write

$$h^{-1}[x_1, x_2] = h^{-1}[x_1, x_3] \bigcup_{h^{-1}(x_3)} h^{-1}[x_3, x_2].$$

Since $h^{-1}(x_3)$ is normally collared (x_3 is an h –regular value) we may use the usual Mayer–Vietoris sequence to conclude that either $h^{-1}[x_1, x_3]$ or $h^{-1}[x_3, x_2]$ fails to have the real Čech cohomology of S^3 and the additional homology $\check{H}^i(h^{-1}[x_1, x_3]; \mathbb{R})$, $i = 1, 2$ restricts from $\check{H}^i(h^{-1}[x_1, x_2]; \mathbb{R})$ (for notational simplicity, write $h^{-1}[x_1, x_3]$ where $h^{-1}[x_3, x_2]$ could instead occur).

Picking a regular $x_4 \in (x_1, x_3)$ and near $(x_1 + x_3)/2$, we may conclude similarly that one of the pieces, say $h^{-1}[x_1, x_4]$, is not a real Čech cohomology S^3 . Proceeding in this way, find a sequence of nested intervals I_j each no more than, say, 60% the length of the previous with the cohomology of $h^{-1}(I_j)$ containing a nontrivial subspace which restricts from $\check{H}^i(h^{-1}[x_1, x_2]; \mathbb{R})$, $i = 1, 2$. Taking inverse limits by letting $z = \bigcap_{j=1}^{\infty} I_j$, we find $\check{H}^i(h^{-1}(z); \mathbb{R})$, $i = 1, 2$, also contains a nontrivial subspace restricting from $\check{H}^i(h^{-1}[x_1, x_2])$, contradicting $b = 2$. □

From the lemma, and once again applying the Mayer–Vietoris sequence to decompositions inverse to h –regular values, we conclude:

Lemma 2 *For every regular value $x_1 \in [0, x_0]$, the natural map*

$$\check{H}^*(M; \mathbb{R}) \rightarrow \check{H}^*(h^{-1}[x_1, x_0]; \mathbb{R})$$

is an isomorphism.

Proof Write

$$M = h^{-1}[x_1, x_0] \cup_{h^{-1}(x_1)} h^{-1}[0, x_1]$$

and use Lemma 3 to recognize the second piece as a cohomology collar. Apply Mayer-Vietoris along $h^{-1}(x_1)$. □

The proof of Theorem 2 is now completed by a second passage to inverse limits. Write

$$h^{-1}(x_0) = \bigcap_{h\text{-regular } x_{1_i}} h^{-1}[x_{1_i}, x_0].$$

By Lemma 4, the cohomology of each $h^{-1}[x_{1_i}, x_0]$ is given isomorphically by restriction from $\check{H}^*(M; \mathbb{R})$, so passing to limits, $\check{H}^*(h^{-1}(x_0); \mathbb{R}) \cong \check{H}^*(M; \mathbb{R})$ via inc^* . But since $b_2(M) \geq 22$, this implies $b_2(h^{-1}(x_0)) \geq 16$. Thus $2 = b \geq b_2(h^{-1}(x_0)) + b_0(h^{-1}(x_0)) \geq 22 + 1 = 23$, a contradiction. □

3 Low Dimensions

The spaces of maps $\mathcal{M}(S^1, S^1)$, $\mathcal{M}(S^2, S^1)$ and $\mathcal{M}(S^2, S^2)$ probably do not hold any surprises though in the last case there are some subtleties.

When the target is S^1 , “circular” Morse theory can be used to steadily simplify maps until they are constant or in the $S^1 \rightarrow S^1$ case, possibly a covering map. If we try to proceed to the case of maps $S^3 \rightarrow S^1$, $\mathcal{M}(S^3, S^1)$, unknown issues are encountered. We may lift each $f: S^3 \rightarrow S^1$ to $\tilde{f}: S^3 \rightarrow \mathbb{R}$. If \tilde{f} happens to be a “self-indexing” (critical values of higher index are larger) a theorem of Waldhausen [14] shows that \tilde{f} can be monotonely simplified to the standard height function. But the existence of such monotone simplifications appears to be open if \tilde{f} is not self-indexing. If no simplifying path from \tilde{f} is monotone, then the passage between a local minimum \tilde{f} and global minimum would be analogous to the instanton we produced between the vacua on S^8 .

$\mathcal{M}(S^2, S^2)$ is quite interesting. I would conjecture that there are no energy barriers: that for reasonable choices of energy topological or otherwise, energy may be monotonically reduced until a cyclic branch cover is reached. This appears to be true for E_{top} , unsigned area and harmonic map energy,

$$\int_{S^2} d^2x |\nabla f|^2.$$

In fact, work of Topping [13] on harmonic map flow might be used in verifying such conjectures. The essential point is that because the target S^2 is *positively curved*,

the local energy density $|du|^2$ is not bounded by the usual Eells–Sampson maximal principle but instead may exhibit a specific singularity called “bubbling” under harmonic map flow. Intuitively, according to Topping, one should be able to run harmonic map flow until just before a singularity and then pause to do a π -twist to prevent the flow from being caught on the singularity and then resume the flow. The phenomenon of bubbling and the flow catching on a singularity are certainly intrinsic since the Hopf map $S^3 \rightarrow S^2$ can be regarded as an “essential loop of degree zero maps $S^2 \rightarrow S^2$ ”; only the flow catching the singularity prevents the loop from being contracted.

Proceeding by dimension, the techniques of Freedman and Berger [3], and DeTurk, Gluck and Storm [1] might be useful in bounding the height of energy barriers within $\mathcal{M}(S^3, S^2)$ if they in fact exist, but so far nothing is known.

For $\mathcal{M}(S^1, S^3)$ (and, more generally, $\mathcal{M}(S^m, S^{m+2})$ where $m \geq 1$), the phenomenon of knotting implies the existence of E_{top} energy barriers. Furthermore, there is a physically interesting conformally invariant energy associated to $\mathcal{M}(S^1, S^3)$ (Freedman, He and Wang [4]) but it is unknown if it has local (but not global) minima within a given knot type. The gradient flow of this functional has been studied as a potential “unknotting” algorithm.

4 Energy Barriers from Mathematical Logic

One surprising geometric output of mathematical logic is the existence of enormous energy barriers related to recognition problems. It was shown in the 1950s by Boone and Novikov that the triviality problem for finitely presented groups is undecidable. However the Tietze theorem tells us that any two finite group presentations of the same group are joined by a finite string of four simple moves on presentations. The only way these two statements can be compatible is:

Fact 1 *The minimal number of Tietze moves required to reduce a presentation of the trivial group of total size n to the empty presentation must be a function $f(n)$ which grows more rapidly than any recursive function.*

This fact finds an even more geometric echo in work of Nabutovsky and Weinberger [11]. An example is:

Fact 2 *If one considers flat piecewise linear imbedding $e: S^5 \hookrightarrow \mathbb{R}^6$ (or one may take $S^d \subset \mathbb{R}^{d+1}$, for and $d \geq 5$) consisting of n top dimensional simplices, then the degree of refinement $f(n)$, which might be required before there is a piecewise linear isotopy to a convex imbedding $e_{\text{convex}}: S^5 \hookrightarrow \mathbb{R}^6$, also grows faster than any recursive function.*

The point is that any recursive growth rate would lead to an algorithm for recognizing 5–spheres, which is impossible. In attempting to recognize S^5 , the chief difficulty is determining the triviality of the fundamental group.

5 Summary

We have shown in a toy model that even for nonlinear σ –models whose range and domains are round spheres, false vacua and energy barriers can arise in the space of maps. The tool is JHC Whitehead’s J –homomorphism, $J_r: \pi_r(\mathrm{SO}) \rightarrow \pi_r^s$, from the homotopy of the special orthogonal group to the stable homotopy of spheres. Energy barriers and the instantons which cross them have diverse mathematical origins. Here the kernel of J has been used to describe a new class of barrier and instanton.

Acknowledgements This work was done partially at the Aspen Center for Physics and I would like to thank Greg Moore for valuable comments regarding an earlier draft of this paper.

References

- [1] **D DeTurck, H Gluck, P A Storm**, *Lipschitz minimality of Hopf fibrations and Hopf vector fields* [arXiv:1009:5439](#)
- [2] **MH Freedman, et al**, *Projective ribbon permutation statistics: a remnant of non-Abelian braiding in higher dimensions*, Phys. Rev. B 83 (2011) 115132 See section 3C for a discussion of charge density wave vs. superconducting order
- [3] **MH Freedman, MA Berger**, *Combinatorial relaxation of magnetic fields*, Geophys. Astrophys. Fluid Dynam. 73 (1993) 91–96 [MR1289021](#)
- [4] **MH Freedman, Z-X He, Z Wang**, *Möbius energy of knots and unknots*, Ann. of Math. 139 (1994) 1–50 [MR1259363](#)
- [5] **M Furuta**, *Monopole equation and the $\frac{11}{8}$ –conjecture*, Math. Res. Lett. 8 (2001) 279–291 [MR1839478](#)
- [6] **P Horava**, *Stability of Fermi surfaces and K –theory*, Phys. Rev. Lett. 95 (2005) 016405
- [7] **CL Kane, E J Mele**, *\mathbb{Z}_2 topological order and the quantum spin Hall effect*, Phys. Rev. Lett. 95 (2005) 146802
- [8] **A Kitaev**, *Periodic table for topological insulators and superconductors* [arXiv:0901.2686](#)
- [9] **A Ludwig, JE Moore, S Ryu**, *Electromagnetic and gravitational responses and anomalies in topological insulators and superconductors*, Phys. Rev. B, to appear

- [10] **R Minasian, G Moore**, *K-theory and Ramond–Ramond charge*, J. High Energy Phys. (1997) Paper 2, 7 pp. [MR1606278](#)
- [11] **A Nabutovsky, S Weinberger**, *Algorithmic aspects of homeomorphism problems*, from: “Tel Aviv Topology Conference: Rothenberg Festschrift”, Contemp. Math. 231, Amer. Math. Soc. (1999) 245–250 [MR1707346](#)
- [12] **S Ryu, A Schnyder, A Furusaki, A Ludwig**, *Topological insulators and superconductors: ten-fold way and dimensional hierarchy*, New J. Phys. 12 (2010) 065010
- [13] **P Topping**, *Bubbling of almost-harmonic maps between 2-spheres at points of zero energy density*, from: “Variational problems in Riemannian geometry”, Progr. Nonlinear Differential Equations Appl. 59, Birkhäuser, Basel (2004) 33–42 [MR2076265](#) Also private correspondence
- [14] **F Waldhausen**, *Heegaard-Zerlegungen der 3-Sphäre*, Topology 7 (1968) 195–203 [MR0227992](#)

Microsoft Station Q, University of California
Santa Barbara, CA 93106-6105, USA

michaelf@microsoft.com

<http://stationq.cnsi.ucsb.edu/~freedman/>

Received: 3 January 2012 Revised: 25 January 2012