



Base subsets of the Hilbert Grassmannian

Mark Pankov

Abstract

Let H be a separable Hilbert space. We consider the Hilbert Grassmannian $\mathcal{G}_\infty(H)$ consisting of closed subspaces having infinite dimension and codimension and show that every bijective transformation of $\mathcal{G}_\infty(H)$ preserving the class of base subsets is induced by an element of $\text{GL}(H)$ or it is the composition of the transformation induced by an element of $\text{GL}(H)$ and the bijection sending a subspace to its orthogonal complement.

Keywords: Hilbert Grassmannian, base subset, infinite-dimensional topological projective space

MSC 2000: 46C05, 51E24

1 Introduction

We start from the classical Grassmannian $\mathcal{G}_k(V)$ consisting of all k -dimensional subspaces of an n -dimensional vector space V . A base subset of $\mathcal{G}_k(V)$ is the set of $\binom{n}{k}$ distinct k -dimensional subspaces spanned by vectors of a certain base of V . This construction is closely related with Tits buildings [5]: the Grassmannians $\mathcal{G}_k(V)$ ($k = 1, \dots, n - 1$) are the shadow spaces of the building associated with the group $\text{GL}(V)$ and their base subsets are the shadows of the corresponding apartments. It was proved in [2] that transformations of $\mathcal{G}_k(V)$ preserving the class of base subsets are induced by a semilinear isomorphism of V to itself or to V^* (the second possibility can be realized only for the case when $n = 2k$); a more general result can be found in [3].

Suppose that V is an infinite-dimensional vector space and $\dim V = \aleph$. The group $\text{GL}(V)$ acts on the set \mathcal{S} of all subspaces of V and the orbits of this action are called *Grassmannians*. There are the following three types of Grassmannians:

$$\mathcal{G}_\alpha(V) := \{ S \in \mathcal{S} \mid \dim S = \alpha, \text{ codim } S = \aleph \}$$

$$\mathcal{G}^\alpha(V) := \{ S \in \mathcal{S} \mid \dim S = \aleph, \text{ codim } S = \alpha \}$$

for any cardinality $\alpha < \aleph$ and

$$\mathcal{G}_\aleph(V) := \{ S \in \mathcal{S} \mid \dim S = \text{codim } S = \aleph \}.$$

As in the finite-dimensional case, we define base subsets of Grassmannians. It was shown in [4] that every bijective transformation of $\mathcal{G}_\alpha(V)$, $\alpha < \aleph$ is induced by a semilinear automorphism of V . The methods of [4] can not be applied to other Grassmannians; this is related with the fact that V and V^* have different dimensions and the duality principles do not hold for infinite-dimensional vector spaces.

Now suppose that V is a topological vector space. Denote by Π_V the projective space associated with V . The topology of V induces a topology on Π_V and we talk about an *infinity-dimensional topological projective space*. A collineation of Π_V to itself will be called *closed* if it preserves the class of closed subspaces. It follows from Mackey's results [1] that every closed collineation of Π_V to itself is induced by an invertible bounded linear operator if V is a real normed space. In the general case, closed collineations are not determined.

In the present note we give a geometrical characterization of closed collineations of the projective space over a separable Hilbert space.

2 Result

Let H be a separable Hilbert space (real or complex). We write $\mathcal{G}_\infty(H)$ for the Hilbert Grassmannian consisting of all closed subspaces with infinite dimension and codimension. Let $B = \{x_i\}_{i \in \mathbb{N}}$ be a base of H (possibly non-orthogonal). The set of all elements of $\mathcal{G}_\infty(H)$ spanned by subsets of B is called the *base subset* of $\mathcal{G}_\infty(H)$ associated with (or defined by) the base B .

Every closed collineation of Π_H to itself induces a bijective transformation of $\mathcal{G}_\infty(H)$ preserving the class of base subsets. The bijection

$$\circ : \mathcal{G}_\infty(H) \rightarrow \mathcal{G}_\infty(H)$$

sending subspaces to their orthogonal complements also preserves the class of base subsets.

Theorem 2.1. *If f is a bijective transformation of $\mathcal{G}_\infty(H)$ preserving the class of base subsets, then f is induced by a closed collineation of Π_H to itself or it is the composition of the transformation \circ and the transformation induced by a closed collineation of Π_H to itself.*

3 Preliminaries

Let $B = \{x_i\}_{i \in \mathbb{N}}$ be base of H , and let \mathcal{B} be the associated base subset of $\mathcal{G}_\infty(H)$. We denote by P_i the 1-dimensional subspace containing x_i and write \mathcal{B}_{+i} and \mathcal{B}_{-i} for the sets of all elements of \mathcal{B} which contain P_i or do not contain P_i , respectively. Then \mathcal{B}_{-i} consists of all elements of \mathcal{B} contained in the hyperplane

$$S_i := \overline{B \setminus \{x_i\}}.$$

We say that $\mathcal{X} \subset \mathcal{B}$ is an *exact* subset if there is only one base subset of $\mathcal{G}_\infty(H)$ containing \mathcal{X} ; otherwise, \mathcal{X} is said to be *inexact*.

Remark 3.1. It is trivial that \mathcal{X} is exact if and only if for each $i \in \mathbb{N}$ the intersection of all $U \in \mathcal{X}$ containing P_i coincides with P_i . For example,

$$\mathcal{R}_{ij} := (\mathcal{B}_{+i} \cap \mathcal{B}_{+j}) \cup \mathcal{B}_{-i} \quad i \neq j$$

is an inexact subset. Indeed, the intersection of all elements containing P_k coincides with P_k if $k \neq i$, however for $k = i$ this intersection is $P_i + P_j$. Every element of $\mathcal{B} \setminus \mathcal{R}_{ij}$ intersects $P_i + P_j$ by P_i . This means that

$$\mathcal{R}_{ij} \cup \{U\}$$

is exact and the inexact subset \mathcal{R}_{ij} is maximal. Conversely, every maximal inexact subset of \mathcal{B} coincides with certain \mathcal{R}_{ij} (the proof of this fact is similar to the proof of Lemma 1 in [4]).

Let \mathcal{B}' be the base subset of $\mathcal{G}_\infty(H)$ defined by a base $B' = \{x'_i\}_{i \in \mathbb{N}}$. We write \mathcal{B}'_{+i} and \mathcal{B}'_{-i} for the sets of all elements of \mathcal{B}' which contain P'_i or do not contain P'_i (respectively); here P'_i is the 1-dimensional subspace containing x'_i . We also define

$$S'_i := \overline{B' \setminus \{x'_i\}}.$$

A bijection $g : \mathcal{B} \rightarrow \mathcal{B}'$ is called *special* if g and g^{-1} map inexact subsets to inexact subsets.

Lemma 3.2. *If $g : \mathcal{B} \rightarrow \mathcal{B}'$ is a special bijection then there exists a bijective transformation $\delta : \mathbb{N} \rightarrow \mathbb{N}$ such that*

$$g(\mathcal{B}_{+i}) = \mathcal{B}'_{+\delta(i)}, \quad g(\mathcal{B}_{-i}) = \mathcal{B}'_{-\delta(i)} \quad \forall i \in \mathbb{N}$$

or

$$g(\mathcal{B}_{+i}) = \mathcal{B}'_{-\delta(i)}, \quad g(\mathcal{B}_{-i}) = \mathcal{B}'_{+\delta(i)} \quad \forall i \in \mathbb{N}.$$

Proof. This is similar to the proof of Lemma 3 in [4]. □

We say that a special bijection $g : \mathcal{B} \rightarrow \mathcal{B}'$ is of *first type* if the first equality of Lemma 3.2 holds; otherwise, g is said to be of *second type*.

Lemma 3.3. *Let g and δ be as in the previous lemma. Let also $S, U \in \mathcal{B}$. If g is of first type then*

$$S \cap U = P_i \Leftrightarrow g(S) \cap g(U) = P'_{\delta(i)} \quad \text{and} \quad \overline{S+U} = S_i \Leftrightarrow \overline{g(S)+g(U)} = S'_{\delta(i)}.$$

If g is of second type then

$$S \cap U = P_i \Leftrightarrow \overline{g(S)+g(U)} = S'_{\delta(i)} \quad \text{and} \quad \overline{S+U} = S_i \Leftrightarrow g(S) \cap g(U) = P'_{\delta(i)}.$$

Proof. The equality $S \cap U = P_i$ holds if and only if S, U belong to \mathcal{B}_{+i} and there is no $j \neq i$ such that \mathcal{B}_{+j} contains both S, U . Similarly, we have $\overline{S+U} = S_i$ if and only if S, U belong to \mathcal{B}_{-i} and there is no $j \neq i$ such that \mathcal{B}_{-j} contains both S, U . Lemma 3.2 gives the claim. \square

For every 1-dimensional subspace P we denote by $[P]$ the set of all elements of $\mathcal{G}_\infty(H)$ containing P . If S is a closed hyperplane then we write $[S]$ for the set of all elements of $\mathcal{G}_\infty(H)$ contained in S .

Lemma 3.4. *For any $i \in \mathbb{N}$ and any $U \in [P_i] \setminus \mathcal{B}$ there exist $M, N \in \mathcal{B}_{+i}$ such that*

$$M \cap N = P_i, \quad \text{codim } \overline{M+N} > 1,$$

and M, N, U are contained in a certain base subset of $\mathcal{G}_\infty(H)$.

Proof. Let $\{U_j\}_{j \in X \subset \mathbb{N}}$ be a countable collection of elements of \mathcal{B} such that

$$U_k \cap U_m = 0 \quad \text{if } k \neq m.$$

For every $j \in X$ we choose $P_j \in U_j \setminus U$ (this is possible, since $U \neq U_j$) and denote by T the element of \mathcal{B} spanned by P_i and all $P_j, j \in X$. Then $U \cap T = P_i$. We take any $M, N \in \mathcal{B}_{+i}$ contained in T and such that $M \cap N = P_i$. These subspaces are as required. \square

Since $\circ|_{\mathcal{B}} : \mathcal{B} \rightarrow \circ(\mathcal{B})$ is a special bijection of second type, we have the following dual version of Lemma 3.4.

Lemma 3.5. *For any $i \in \mathbb{N}$ and any $U \in [S_i] \setminus \mathcal{B}$ there exist $M, N \in \mathcal{B}_{-i}$ such that*

$$\overline{M+N} = S_i, \quad \dim M \cap N > 1,$$

and M, N, U are contained in a certain base subset of $\mathcal{G}_\infty(H)$.

4 Proof of the theorem

Let f be a bijective transformation of $\mathcal{G}_\infty(H)$ preserving the class of base subsets. We consider an arbitrary base subset $\mathcal{B} \subset \mathcal{G}_\infty(H)$. Then $\mathcal{B}' := f(\mathcal{B})$ is a base subset and $f|_{\mathcal{B}} : \mathcal{B} \rightarrow \mathcal{B}'$ is a special bijection. Suppose that $B = \{x_i\}_{i \in \mathbb{N}}$ and $B' = \{x'_i\}_{i \in \mathbb{N}}$ are bases associated with \mathcal{B} and \mathcal{B}' , respectively. Let P_i, P'_i, S_i, S'_i and $\delta : \mathbb{N} \rightarrow \mathbb{N}$ be as in the previous section. It is clear that we can assume that δ is identical. We have to consider the following possibilities:

- (A) $f|_{\mathcal{B}}$ is a special bijection of first type,
- (B) $f|_{\mathcal{B}}$ is a special bijection of second type.

Case (A). First we claim that

$$f([P_i]) = [P'_i] \quad \forall i \in \mathbb{N}.$$

Let U, M, N be as in Lemma 3.4. By Lemma 3.3,

$$f(M) \cap f(N) = P'_i, \quad \text{codim } \overline{f(M) + f(N)} > 1. \quad (1)$$

If $\tilde{\mathcal{B}}$ is a base subset containing U, M, N then (1) together with Lemma 3.3 show that the restriction of f to $\tilde{\mathcal{B}}$ is a special bijection of first type. Bases associated with $\tilde{\mathcal{B}}$ and its f -image contain vectors lying in P_i and P'_i (respectively), and Lemma 3.2 implies that $f(U)$ belongs to $[P'_i]$. Thus $f([P_i]) \subset [P'_i]$. Since f^{-1} preserves the class of base subsets, we can also prove the inverse inclusion. The claim follows.

Using Lemma 3.5 we show that

$$f([S_i]) = [S'_i] \quad \forall i \in \mathbb{N}.$$

For any 1-dimensional subspace P there exists S_i which do not contain P . We consider a base $\tilde{\mathcal{B}}$ of H such that each vector of $\tilde{\mathcal{B}}$ is contained in S_i or P . The latter equality guarantees that the restriction of f to the associated base subset is a special bijection of first type, and arguments given above imply the existence of a 1-dimensional subspace P' such that $f([P]) = [P']$.

Similarly, for every closed hyperplane S we choose $P_i \notin S$ and establish the existence of a closed hyperplane S' such that $f([S]) = [S']$.

Therefore, f gives a closed collineation of Π_H . This collineation induces f .

Case (B). Since $\circ f|_{\mathcal{B}}$ is a special bijection of first type, $\circ f$ is induced by a closed collineation.

Acknowledgment

The result of this paper was presented on 32. *Arbeitstagung über Geometrie und Algebra* (9 – 11 February, 2006, Hamburg). The author thanks the organizers (A. Blunck, H. Kiechle, A. Kreuzer, H.-J. Samaga, H. Wefelscheid) for the invitation and financial support.

References

- [1] **G. W. Mackey**, Isomorphisms of normed linear spaces, *Ann. of Math.* **43** (1942), 244-260.
- [2] **M. Pankov**, Transformations of Grassmannians and automorphisms of classical groups, *J. Geom.* **75** (2002), 132-150.
- [3] _____, A characterization of geometrical mappings of Grassmann spaces, *Results Math.* **45** (2004), 319-327.
- [4] _____, Base subsets of Grassmannians: infinite-dimensional case, *European J. Combin.* **28** (2007), 26-32.
- [5] **J. Tits**, *Buildings of spherical type and finite BN-pairs*, Lect. Notes in Math. **386**, Springer-Verlag, 1974.

Mark Pankov

DEPARTMENT OF MATHEMATICS AND INFORMATION TECHNOLOGY,, UNIVERSITY OF WARMIA AND MAZURY,, ŻOLNIERSKA 14A,, 10-561 OLSZTYN,, POLAND

e-mail: pankov@matman.uwm.edu.pl