

page 1 / 13

go back

full screen

close

quit

Directions in $AG(2, p^2)$

András Gács*

László Lovász

Tamás Szőnyi

Dedicated to Gábor Korchmáros on his 60th birthday.

Abstract

In this paper we prove that if q is the square of a prime and U is a set of q points determining at least $\frac{q+3}{2}$ directions, then either U is affinely equivalent to the graph of the function $x^{\frac{q+1}{2}}$ or it determines at least $\frac{q+p}{2} + 1$ directions. This is sharp, the example is due to Polverino, Szőnyi and Weiner [10]. Our method combines the lacunary polynomial and the double power sum approach.

Keywords: affine planes, directions, blocking sets of Rédei type

MSC 2000: 51E21

1. Introduction

Throughout this paper $U = \{(a_i, b_i) : i = 1, \dots, q\}$ will denote a q -element point set in $AG(2, q)$, the Desarguesian affine plane of order q . We write

$$D = \left\{ \frac{b_i - b_j}{a_i - a_j} \mid i \neq j \right\}$$

and call elements of this set *the directions determined by U* . This is a subset of $GF(q) \cup \{\infty\}$ and consists of slopes of lines joining two points of U . Finally, let $N = |D|$, the number of determined directions.

The problem of determining the possible values of N and characterizing the corresponding point sets has received a lot of attention in recent years. For motivation and the history of the problem we refer to [3] and [4]. Here we summarize some known results.

*The authors were supported by Magyary and Bolyai grants, TÉT grants E16/04 and SI-2/2007, OTKA grants F043772, T043758, T049662, T067867.

ACADEMIA
PRESS





page 2 / 13

go back

full screen

close

quit

Result 1.1 (Rédei [11]; Blokhuis, Ball, Brouwer, Storme, Szőnyi [4] and Ball [1]). Let $q = p^h$ and let $s = p^e$ be maximal with the property that any line containing at least two points of U meets U in a multiple of s points. Then one of the following holds:

- (i) $s = 1$ and $\frac{q+3}{2} \leq N \leq q + 1$;
- (ii) $\text{GF}(s)$ is a subfield of $\text{GF}(q)$ and $1 + q/s \leq N \leq (q - 1)/(s - 1)$;
- (iii) $s = q$ and $N = 1$.

Moreover, if $s > 2$, then U can be regarded as a coset of a vector space over $\text{GF}(s)$.

This result solves the problem entirely for the case $N < \frac{q+3}{2}$. It was first proved in [4] with some exceptions for the characteristic 2 and 3 cases. Recently S. Ball [1] found an easier proof which also handles the missing cases.

For the case $N \geq \frac{q+3}{2}$ there have been results only when q is a prime:

Result 1.2. (i) (Lovász and Schrijver [9]) If q is a prime, then the only sets determining $\frac{q+3}{2}$ directions are the affine equivalents of the graph of the function $x^{\frac{q+1}{2}}$.

(ii) (Gács [7]) If q is prime and $N > \frac{q+3}{2}$, then $N \geq [2 \frac{q-1}{3} + 1]$.

Note that for the case q is a prime Result 1.1 gives that N is at least $\frac{q+3}{2}$, unless $N = 1$ (that is U is a line). This was already observed by Rédei and Megyesi, see [11]. The graph of $x^{\frac{q+1}{2}}$ determines $\frac{q+3}{2}$ directions for any odd prime power q , showing that the bound in Result 1.1(i) is sharp.

In this paper we consider the next case, that is, when q is the square of a prime. We prove an analogous result to the two statements of Result 1.2:

Theorem 1.3. Suppose that $q = p^2$, where p is prime and U is a set of q points in $\text{AG}(2, q)$ determining at least $\frac{q+3}{2}$ directions. Then either U is affinely equivalent to the graph of $x^{\frac{q+1}{2}}$, or the number of directions is at least $\frac{q+p}{2} + 1$.


The bound is sharp: Polverino, Szőnyi and Weiner [10] constructed an example determining $\frac{q+\sqrt{q}}{2} + 1$ directions when q is a square. We conjecture that for any square prime power, this should be the next possible value for N , that is, $\frac{q+3}{2} < N < \frac{q+\sqrt{q}}{2} + 1$ cannot happen for q square.

We continue with some preliminary remarks on polynomials over finite fields and Lucas' theorem. For proofs, see [8].

When U does not determine all directions, that is, when $N < q + 1$, one can find a suitable transformation (not affecting N) to achieve that $U = \{(x, f(x)) \mid x \in \text{GF}(q)\}$ for some function f . After this we have another form for D ; namely,

ACADEMIA
PRESS





page 3 / 13

go back

full screen

close

quit

it is easy to see that D consists of those $c \in \text{GF}(q)$ for which $f(x) - cx$ is not bijective.

Over the finite field $\text{GF}(q)$ any function can be written as a polynomial of degree at most $q - 1$. This is called the *reduced form* of f . For any $f(x) = c_{q-1}x^{q-1} + \dots + c_1x + c_0$, we have $\sum_{x \in \text{GF}(q)} f(x) = -c_{q-1}$. Such an f is called a *permutation polynomial* if it is bijective as a function. For such polynomials we have $\sum_{x \in \text{GF}(q)} f(x)^k = 0$ for any $1 \leq k \leq q - 2$; this is equivalent to saying that in the reduced form of $f(x)^k$ the coefficient of x^{q-1} is zero.

Lucas' theorem tells how binomial coefficients behave modulo a prime p . Let the p -adic expansion of n and k be $n = \sum_{i=1}^r n_i p^{i-1}$ and $k = \sum_{i=1}^r k_i p^{i-1}$, respectively. Then $\binom{n}{k} \equiv \binom{n_1}{k_1} \dots \binom{n_r}{k_r}$ modulo p .

Finally, we show how the direction problem is connected to blocking sets in $\text{PG}(2, q)$. A *blocking set* B in the projective plane $\text{PG}(2, q)$ is a set of points meeting every line. A blocking set B is called *non-trivial* if it contains no line, and *minimal* if it does not properly contain a blocking set.

If U is a set of q points in $\text{AG}(2, q)$ and D is the set of determined directions, then embedding $\text{AG}(2, q)$ into $\text{PG}(2, q)$ and adding to U the infinite points corresponding to elements in D , we get a blocking set B of the projective plane. It contains a line if and only if either U is an affine line or U determines every direction. Then B has the property that there is a line (the line at infinity) missing exactly q points of B . It is easy to see that this property characterizes minimal blocking sets arising from the above construction; they are called *blocking sets of Rédei type*.

In the next section we deal with blocking sets in general. After some easy observations we will end up in a result about Rédei type blocking sets (Proposition 2.4), which will be used in the proof of Theorem 1.3.

We will consider $\text{PG}(2, q)$ as $\text{AG}(2, q)$ extended by the line at infinity, l_∞ . The infinite point of lines with slope c will be denoted by (c) , the infinite point of the vertical lines will be denoted by (∞) .


2. Blocking sets

Suppose that B is a blocking set in $\text{PG}(2, q)$ with $|B| < 2q - 1$ and $(\infty) \notin B$. Write $U = B \setminus l_\infty = \{(a_i, b_i) \mid i = 1, \dots, k\}$ and let $D = B \cap l_\infty = \{(y_i) \mid i = 1, \dots, N\}$. The *Rédei polynomial* of U is defined as

$$H(X, Y) = \prod_{i=1}^k (X + a_i Y + b_i).$$

ACADEMIA
PRESS





We will often use the Rédei polynomial of B also, which is

$$H^*(X, Y) = \prod_{i=1}^N (Y + y_i) H(X, Y).$$

Finally, the *homogeneous Rédei polynomial* of B is defined as

$$R(X, Y, Z) = \prod_{i=1}^N (Y + y_i Z) \prod_{i=1}^k (X + a_i Y + b_i Z).$$

Note that R is the homogenization of H^* .

The partial derivatives of R with respect to X , Y and Z will be denoted by R'_X , R'_Y and R'_Z , respectively.

Lemma 2.1. (i) *There exist polynomials f_1 and f_2 of degree at most $|B| - q$ such that*

$$H^*(X, Y) = (X^q - X)f_1(X, Y) + (Y^q - Y)f_2(X, Y);$$

(ii) *there exist homogeneous polynomials f, g, h of degree $|B| - q$ such that*

$$R(X, Y, Z) = X^q f + Y^q g + Z^q h;$$

(iii) *$Xf + Yg + Zh = 0$ for the polynomials found in (ii);*

(iv) *for any $(x, y, z) \in \text{GF}(q)^3 \setminus (0, 0, 0)$, we have*

$$\begin{aligned} f(x, y, z) &= -R'_X(x, y, z), \\ g(x, y, z) &= -R'_Y(x, y, z), \\ h(x, y, z) &= -R'_Z(x, y, z), \end{aligned}$$

for the polynomials found in (ii).

Proof. (i) is well-known, see Blokhuis [2]. For the homogenization of H^* it gives that it is of the form $R = X^q f_1^* + Y^q f_2^* - Z^{q-1}(X f_1^* + Y f_2^*)$. Hence the polynomials $f = f_1^*$, $g = f_2^*$ and $h = -\frac{1}{Z}(X f_1^* + Y f_2^*)$ will be appropriate for (ii) and (iii), provided that we can prove that $Z \mid X f_1^* + Y f_2^*$. Consider the terms of R not containing Z . These are $Y^N \prod_i (X + a_i Y)$. Since $(\infty) \notin B$, each element of $\text{GF}(q)$ occurs at least once as an a_i . Hence the terms we are looking for can be written as $X^M Y^N (X^{q-1} - Y^{q-1})s(X, Y)$ ($M, N \geq 1$). They all come from $X^q f_1^* + Y^q f_2^*$, so we have to find out the terms containing X^q and Y^q , which are $X^q X^{M-1} Y^N s(X, Y)$ and $-Y^q X^M Y^{N-1} s(X, Y)$. Hence the terms in $X f_1^* + Y f_2^*$ not containing Z are $X X^{M-1} Y^N s(X, Y) - Y X^M Y^{N-1} s(X, Y) = 0$.

Now (iv) easily follows from (iii) and the derivative of (ii). \square



page 4 / 13

go back


full screen

close

quit

ACADEMIA
PRESS





page 5 / 13

go back

full screen

close

quit

The following lemma gives an easy consequence of Lemma 2.1.

Lemma 2.2. *Suppose that the line $l : aX + bY + cZ = 0$ is a 1-secant to B . Then the unique intersection point of l and B is $(f(a, b, c), g(a, b, c), h(a, b, c))$.*

Proof. The point $(f(a, b, c), g(a, b, c), h(a, b, c))$ is on the line in question because of Lemma 2.1(iii). At this stage it is more convenient not to distinguish between the affine and infinite points of B , so write $B = \{(u_i, v_i, w_i) \mid i = 1, \dots, k + N\}$, hence $R(X, Y, Z) = \prod_{i=1}^{k+N} (u_i X + v_i Y + w_i Z)$. Differentiate R with respect to X to find $R'_X(X, Y, Z) = \sum_i u_i \prod_{j \neq i} (u_j X + v_j Y + w_j Z)$. If we substitute $X = a$, $Y = b$ and $Z = c$, then all products in the sum will be zero, except for the case when (u_i, v_i, w_i) is the intersection point, hence $R'_X(a, b, c) = u_i \prod_{j \neq i} (u_j a + v_j b + w_j c)$ (for this i). Similarly we have $R'_Y(a, b, c) = v_i \prod_{j \neq i} (u_j a + v_j b + w_j c)$ and $R'_Z(a, b, c) = w_i \prod_{j \neq i} (u_j a + v_j b + w_j c)$. Since the line $aX + bY + cZ = 0$ is a 1-secant to B , this product is non-zero. By Lemma 2.1(iv), we are done. \square

Note that when the line $aX + bY + cZ = 0$ meets B in more than one point, then $f(a, b, c) = g(a, b, c) = h(a, b, c) = 0$.

Now we suppose that B is of Rédei type. Hence we have $k = q$ and $D = \{(y_i) \mid i = 1, \dots, N\}$ is the set of determined directions of the affine set $U = \{(a_i, b_i) \mid i = 1, \dots, q\}$.

Note that in this case we have $f(X, Y, Z) = \prod_{i=1}^N (Y + y_i Z)$ in Lemma 2.1.

Definition 2.3. The index I of B is defined so that the X -degree of the polynomial $Y^q g + Z^q h$ is $q - I$.

From Lemma 2.1(iii) we see that $q - I$ is also the X -degree of g and h (unless one of them is 1 and the other is 0). Note that considering H as a polynomial in X (with coefficient polynomials in Y), X^{q-I} is the first term after X^q with non-vanishing coefficient polynomial.


Proposition 2.4. (i) *If the infinite point (y) does not belong to B (that is, (y) is not a determined direction), then the affine part of B is equivalent to $\{(\frac{1}{c}g(t, y, -1), t) \mid t \in \text{GF}(q)\}$, where $c \neq 0$ depends on y ;*

(ii) *If $q - N > N + I - q$, then by a suitable linear transformation we can suppose that the affine part of B is the graph of a polynomial of degree $q - I$.*

Proof. If (y) is not determined, then all affine lines through it are 1-secants, hence by calculating the intersections of B and these lines, we can determine the q affine points.

ACADEMIA
PRESS





page 6 / 13

go back

full screen

close

quit

The lines in question have equation $tX_0 + yX_1 - X_2 = 0$, so by Lemma 2.1, we find that the affine part of B is

$$\{(f(t, y, -1), g(t, y, -1), h(t, y, -1)) \mid t \in \text{GF}(q)\}.$$

We know that $f(t, y, -1) = \prod(y - y_i)$, and from Lemma 2.1(iii) we have $yg(t, y, -1) - h(t, y, -1) = -\prod(y - y_i)t$. Note that $c := \prod(y - y_i)$ is a constant, hence after the transformation $X'_2 = X_2 - yX_1$ we find the form claimed in (i).

By the definition of I , after (i) we only need that there is a suitable non-determined direction (y) , for which the degree of $g(X, y, -1)$ is the same as the X -degree of g . We have $q - N$ choices for y . The coefficient of X^{q-I} in $g(X, Y, Z)$ is a homogeneous polynomial $g_0(Y, Z)$ of degree $|B| - q - (q - I) = N + I - q$. If this is smaller than $q - N$, then from the fact that g_0 is not identically zero, we should have an appropriate y . \square

Remark 2.5. Similar ideas and some of the results were used by Sziklai to prove results about small blocking sets, see [12].

3. Results about directions for general q

In the spirit of the introduction, from now on $U = \{(x, f(x)) \mid x \in \text{GF}(q)\}$,

$$D = \left\{ \frac{f(x) - f(y)}{x - y} \mid x \neq y \right\} = \{c \in \text{GF}(q) \mid f(x) - cx \text{ is not a perm. pol.}\},$$

and $N(f) = |D|$. Here $f(x) = c_n x^n + \dots + c_0$ with $\deg(f) = n \leq q - 1$. In this section we introduce two more parameters depending on f and relate them to $N(f)$.

By the remarks at the end of the introduction, $U \cup D$ is a blocking set of Rédei type. In this case the Rédei polynomial is

$$H(X, Y) = \prod_{t \in \text{GF}(q)} (X + tY + f(t)).$$


Expanding H in powers of X , we have

$$H(X, Y) = X^q + h_1(Y)X^{q-1} + \dots + h_q(Y). \quad (1)$$

Write $h_i(Y) = \sum_j \sigma_{i-j, j} Y^j$. Note that h_i is the i -th elementary symmetric polynomial of the multiset $\{Yt + f(t) \mid t \in \text{GF}(q)\}$. It is easy to see that $\sigma_{0, i}$, that is, the coefficient of Y^i in h_i , is the i -th elementary symmetric polynomial of the set $\text{GF}(q)$. This is zero for $1 \leq i \leq q - 2$, so for these i 's, $\deg(h_i) \leq i - 1$.

ACADEMIA
PRESS





page 7 / 13

go back

full screen

close

quit

In general for $(a, b) \neq (0, 0)$ we have the following for $\sigma_{a,b}$ (the coefficient of Y^a in h_{a+b} , that is, the coefficient of $X^{q-a-b}Y^a$ in $H(X, Y)$):

$$\sigma_{a,b} = \sum_{t_1, \dots, t_a, u_1, \dots, u_b} t_1 \cdots t_a \cdot f(u_1) \cdots f(u_b),$$

where the sum is over all choices of $t_1, \dots, t_a, u_1, \dots, u_b$ all different. For $a = b = 0$, we have $\sigma_{0,0} = 1$.

The use of $H(X, Y)$ is that it translates intersection properties of U to algebraic ones. This was used in all proofs mentioned in the Introduction and in Section 2.

Lemma 3.1. *Fixing $Y = y_0$ and considering $H(X, y_0)$ as a polynomial in X , the multiplicities of its roots are the same as a multiset as the intersection sizes of lines through the infinite point $(-y_0)$ with U .*

Proof. See Rédei [11]. □

We introduce another series of polynomials:

$$g_k(c) := \sum_{t \in \text{GF}(q)} (f(t) + ct)^k = \sum_{i=0}^k \binom{k}{i} \pi_{i, k-i} c^i, \quad (2)$$

where $\pi_{a,b} = \sum_{t \in \text{GF}(q)} t^a f(t)^b$. Define $\pi_{0,0}$ to be 1. Note that the g_k 's are the power sums of the multiset $\{f(t) + ct \mid t \in \text{GF}(q)\}$.

The two parameters (depending on the reduced polynomial f) to be introduced are the following.

Definition 3.2. The first index $I_1(f)$ of f is defined to be the smallest positive integer k for which the polynomial h_k defined by (1) is not identically zero.

The second index $I_2(f)$ of f is defined to be the smallest positive integer k for which the polynomial g_k defined by (2) is not identically zero.


Note that $I_1(f)$ coincides with the index (of the blocking set) defined in the previous section. The reason for not using the same notation is that we want to stress that I is a parameter of the blocking set B , while I_1 (and I_2) are parameters of the affine part U of the blocking set.

The proofs in [1] and in [4] make use of lacunary polynomials arising from $H(X, Y)$, this means the consideration of the parameter $I_1(f)$. On the other hand, [7] and [9] use double power sums in the proof, this is the consideration of the parameter $I_2(f)$.

In this paper we use both parameters, it seems that this might be the way of attacking the $N \geq \frac{q+3}{2}$ case. Next we relate I_1, I_2 and N to each other. Most of

ACADEMIA
PRESS





page 8 / 13

go back

full screen

close

quit

these observations are at least implicitly stated in one of the above mentioned papers. The first part was also observed by Evans, Greene and Niederreiter [6].

Lemma 3.3. (i) If $I_1(f) \geq \frac{q+1}{2}$, then $N(f) \leq \frac{q-1}{p-1}$;

(ii) if $N(f) > 1$, then $q + 1 - N(f) \leq I_1(f) \leq I_2(f)$, with $I_1(f) = I_2(f)$ if and only if p does not divide $I_1(f)$.

Proof. For (i) we refer to [4]. This is the first easy step of the proof which was already found by Rédei [11].

The fact that $I_1(f) \leq I_2(f)$ and the characterization of the case of equality is a consequence of the Newton formulas relating power sums and elementary symmetric polynomials.

For $q+1-N(f) \leq I_1(f)$ note that fixing any $-y \notin D$ we have $H(X, y) = X^q - X$, hence for these y 's we have $h_1(y) = \dots = h_{q-2}(y) = 0$. For $h_1, \dots, h_{q-N(f)}$ these are more roots than their degrees, so these h_i 's are identically zero. \square

Corollary 3.4. Suppose q is odd. If $\frac{q+3}{2} \leq N(f) \leq \frac{q+p}{2}$, then $I_1(f) = I_2(f)$.

Proof. From Lemma 3.3 we deduce $\frac{q-p}{2} + 1 \leq I_1(f) \leq \frac{q-1}{2}$, so I_1 cannot be divisible by p . The same lemma gives $I_1 = I_2$. \square

Lemma 3.5. Suppose $N(f) < 3q/4$ and $I_1(f) \leq \frac{q-1}{2}$. Then one can make a linear transformation for the graph of f to find the graph of another polynomial f_1 with $\deg(f_1) = q - I_1(f_1) = q - I_1(f)$.

Proof. This is a consequence of Proposition 2.4(ii). The conditions are easily seen to be satisfied, so after transformation, we can find the desired f_1 . Since this is in fact a transformation of the blocking set that fixes the Rédei line, $N(f_1) = N(f)$ and by the original definition of I , we see that $I_1(f) = I_1(f_1)$. \square


The following lemma gives a relation between the above defined $\sigma_{k,l}$'s and $\pi_{k,l}$'s. It can be considered as a generalization of the Newton-Girard formulas relating elementary symmetric polynomials to power sums.

Lemma 3.6. Fix two positive integers k and l . The following formula holds:

$$\sum_{r=0}^k \sum_{s=0}^l w(r, s) \pi_{r,s} \sigma_{k-r, l-s} = 0,$$

ACADEMIA
PRESS





page 9 / 13

go back

full screen

close

quit

where the function $w(r, s)$ is defined as follows: fix two field-elements a and b , then $w(0, 0) := ak + bl$; while for $(r, s) \neq (0, 0)$

$$w(r, s) := (-1)^{r+s} \left(\binom{r+s-1}{s} a + \binom{r+s-1}{r} b \right).$$

(That is, we get a formula for every choice of a and b .)

Proof. It is easy to see that after multiplication on the left hand side we have monomials of the form $t^r f(t)^s t_1 \cdots t_{k-r} f(u_1) \cdots f(u_{l-s})$, where t, t_1, \dots, u_{l-s} are different field elements and $0 \leq r \leq k, 0 \leq s \leq l$.

If r and s are both positive, then we can get such a term from three summands: $\pi_{r,s} \sigma_{k-r,l-s}$, $\pi_{r-1,s} \sigma_{k+1-r,l-s}$ and $\pi_{r,s-1} \sigma_{k-r,l+1-s}$. Hence the coefficient of such a monomial is $w(r, s) + w(r, s-1) + w(r-1, s) = 0$.

If $r = 0, s > 1$, then there are two summands giving the monomial in question: $\pi_{0,s} \sigma_{k,l-s}$ and $\pi_{0,s-1} \sigma_{k,l+1-s}$, so the coefficient of such a monomial is $w(0, s) + w(0, s-1) = 0$. The $s = 0, r > 1$ case is similar.

The $\{r, s\} = \{0\}$ and $\{r, s\} = \{0, 1\}$ cases are the same, so what is left is to show that monomials of the form $t_1 \cdots t_k f(u_1) \cdots f(u_k)$ also have zero coefficient. There are three summands giving them: $\pi_{0,0} \sigma_{k,l}$ (one time), $\pi_{1,0} \sigma_{k-1,l}$ (k times) and $\pi_{0,1} \sigma_{k,l-1}$ (l times). Hence the coefficient in question is $w(0, 0) + lw(0, 1) + kw(1, 0) = 0$. \square

We will use two corollaries of this lemma. The first one was noticed by Chou [5].

Corollary 3.7. (i) $\deg(f) \leq q - I_1(f)$;

(ii) in the reduced form of f^2 the only non-zero terms of degree higher than $q + 1 - I_1(f)$ can be those of degree divisible by p .

Proof. (i) By the definition of I_1 , we know that $\sigma_{k,l} = 0$ for every $0 < k + l < I_1(f)$. We use the formula of Lemma 3.6 with $l = 1, a = 0$ and $b = 1$. It gives $(-1)^{k+1} \binom{k}{k} \pi_{k,1} = 0$ for all $k \leq I_1(f) - 2$, which means that the coefficients of $x^{q-1}, x^{q-2}, \dots, x^{q-I_1(f)+1}$ are zero in f .

(ii) Similarly to (i), we now use the formula with $l = 2, a = 0, b = 1$ to find $(-1)^{k+2} \binom{k+1}{k} \pi_{k,2} = 0$ for all $k \leq I_1(f) - 3$. This gives that in the reduced form of $f^2 x^k$, the coefficient of x^{q-1} is zero, unless $p \mid k + 1$; this is exactly what we wanted. \square

In the next theorem, we summarize what we have so far.

ACADEMIA
PRESS





page 10 / 13

go back

full screen

close

quit

Theorem 3.8. Suppose U is a set of q points in $AG(2, q)$ determining $N \leq \frac{q+p}{2}$ directions, where q is a proper power of the odd prime p . Then either $N \leq \frac{q+1}{2}$ and we know all such examples from the classification [1, 4], or U is affinely equivalent to the graph of a polynomial f with $I_1(f) = I_2(f) = \frac{q-1}{2}$ and $\deg(f) = \frac{q+1}{2}$.

Proof. The previous lemmas together yield that after transformation, U is the graph of a polynomial $f(x) = x^n + \dots + c_2x^2 + c_1x + c_0$ with

$$\frac{q-p}{2} + 1 \leq I_1(f) = I_2(f) = q + 1 - n \leq \frac{q+1}{2}.$$

All we need is that $n = \frac{q+1}{2}$.

Write $n = \frac{q-1}{2} + r$ with $1 \leq r \leq \frac{p-1}{2}$. Consider the reduced form of f^p . The term x^n will give $x^{(q-1)/2+rp}$ (after reduction). It follows that

$$\sum x^{(q-1)/2-rp} f(x)^p \neq 0.$$

Since $\binom{(q-1)/2-rp+p}{p} \neq 0$ by Lucas' theorem, this gives that $g_{(q-1)/2-(r-1)p} \neq 0$ identically, hence by the definition of I_2 , we have $(q-1)/2 - (r-1)p \geq I_2 > (q-p)/2$. This is only possible for $r = 1$. \square

The above theorem implies in particular, that to prove Theorem 1.3 or even its generalization to an arbitrary odd prime power q (which is not a prime), one can suppose that the set U is the graph of a polynomial of degree $\frac{q+1}{2}$.

4. Proof of Theorem 1.3

From now on suppose $q = p^2$ for a prime p . For $p = 2$ and $p = 3$ there is nothing to prove, so suppose $p \geq 5$. By Theorem 3.8 let $f(x) = x^{\frac{q+1}{2}} + \dots + c_0$ be a polynomial with $\frac{q+3}{2} \leq N(f) \leq \frac{q+p}{2}$ and $I_1 = I_2 = \frac{q-1}{2}$. We make a transformation to achieve $c_{\frac{q-1}{2}} = c_1 = c_0 = 0$. It is not difficult to see that this does not affect I_1 or I_2 . We have to prove that f is equivalent to $x^{\frac{q+1}{2}}$. The proof will be carried out in several steps.

Claim 1. Consider the intervals

$$A_i = \left(\frac{q-1}{2} - (i+1)p, \frac{q-1}{2} - ip \right] \text{ for } i = 0, 1, \dots, \frac{p-3}{2}.$$


The only possible indices $j \in A_i$ with $c_j \neq 0$ are

$$\frac{q-1}{2} - ip, \frac{q-1}{2} - ip - 1, \dots, \frac{q-1}{2} - ip - (i-1)$$

(for $i = 0$ this means that for all $j \in (\frac{q-1}{2} - p, \frac{q-1}{2}]$, $c_j = 0$).

ACADEMIA
PRESS





page 11 / 13

go back

full screen

close

quit

Proof. We use Corollary 3.7(ii). Here $q + 1 - I_1 = \frac{q+3}{2}$.

Using that in f^2 the coefficients of $x^{q-1}, \dots, x^{q-p+1}$ are zero, we find that $c_{\frac{q-3}{2}} = \dots = c_{(q-1)/2-p+1} = 0$. Then using the fact that in f^2 there is no $x^{q-p-1}, \dots, x^{q-2p+1}$, we find that $c_{(q-1)/2-p-1} = \dots = c_{(q-1)/2-2p+1} = 0$. Again we cannot cancel $c_{(q-1)/2-2p}$, but also $c_{(q-1)/2-2p-1}$. The reason for the latter is that in f^2 the terms $x^{\frac{q+1}{2}} x^{(q-1)/2-2p-1}$ and $(x^{(q-1)/2-p})^2$ give terms of the same degree, so they might cancel each other.

In general we use induction on i . Suppose we have proved the statement for $0, \dots, i-1$ but $c_{(q-1)/2-ip-j} \neq 0$ for a $j \geq i$. Considering f^2 again, we find a term of degree $q - ip - j$. This is a contradiction unless we can have this term from the product of two terms of the form $(q-1)/2 - i_1p - j_1$ and $(q-1)/2 - i_2p - j_2$ for some $i_1, i_2 \leq i-1$ and $j_1 \leq i_1 - 1, j_2 \leq i_2 - 1$ (by the induction hypothesis). An easy calculation shows that this is not possible. \square

Note that what we have proved implies in particular that all terms below $\frac{q+1}{2}$ have degree between 0 and $\frac{p-1}{2}$ modulo p .

Claim 2. If $f(x) \neq x^{\frac{q+1}{2}}$, then $f(x) = x^{\frac{q+1}{2}} + c_{(q-1)/2-jp} x^{(q-1)/2-jp} + \dots$ with $c_{(q-1)/2-jp} \neq 0$. (That is, the first term after $\frac{q+1}{2}$ with nonzero coefficient has to be congruent to $\frac{p-1}{2}$ modulo p .)

Proof. Let $f(x) = x^{\frac{q+1}{2}} + c_s x^s + \dots$. Then in the reduced form of $f(x)^2$ the coefficient of the term $x^{(q+1)/2+s}$ is not zero, hence by Corollary 3.7(ii), $\frac{q+1}{2} + s$ is divisible by p . \square


Claim 3. If $f(x) \neq x^{\frac{q+1}{2}}$, then $f(x) = x^{\frac{q+1}{2}} + c_{(q-1)/2-p} x^{(q-1)/2-p} + \dots$ with $c_{(q-1)/2-p} \neq 0$ (hence $j = 1$ in Claim 2) and $c_{(q-1)/2-p}^p + c_{(q-1)/2-p} = 0$.

Proof. By the remark after Claim 1, one can show that after reduction $f(x)^p = x^{(q-1)/2+sp} + c_{(q-1)/2-jp}^p x^{(q-1)/2-jp} + \dots$. Multiplying f and the reduced form of f^p we find that in the reduced form of f^{p+1} there are two big terms, namely $c_{(q-1)/2-jp}^p x^{q-j}$ and $c_{(q-1)/2-jp} x^{q-1-(j-1)p}$. These can cancel each other if $j = 1$ and $c_{(q-1)/2-jp}^p + c_{(q-1)/2-jp} = 0$, in all other cases f^{p+1} has a non-zero term of degree $q - j$. But this cannot happen, since this would mean that g_{p+j} is not zero identically (note that $\binom{p+j}{p+1}$ is not zero for $j \geq 1$). \square

Claim 4. $f(x) = x^{\frac{q+1}{2}}$.

ACADEMIA
PRESS





page 12 / 13

go back

full screen

close

quit

Proof. Suppose $f(x) \neq x^{\frac{q+1}{2}}$. We summarize what we have from the previous three claims:

$$\begin{aligned} f(x) &= x^{(q+1)/2} + c_{(q-1)/2-p} x^{(q-1)/2-p} + \dots ; \\ f(x)^p &= x^{(q+1)/2+p-1} + c_{(q-1)/2-p}^p x^{(q-3)/2} + \dots ; \\ 0 &= c_{(q-1)/2-p}^p + c_{(q-1)/2-p} . \end{aligned} \quad (3)$$

Now consider $x^2 (x^{p-1} f(x) - f(x)^p)^2$. This polynomial is the linear combination of polynomials of the form $x^k f(x)^l$ with $\binom{k+l}{k} \neq 0$, hence by the definition of I_2 , after reduction it cannot have a term of degree $q-1$ (unless $2p+2 \geq I_1$). It is easy to see that this gives $c_{(q-1)/2-p}^p - c_{(q-1)/2-p} = 0$, which together with (3) implies $c_{(q-1)/2-p} = 0$, a contradiction.


So the proof is finished except for the case $2p+2 \geq I_1 = \frac{q-1}{2}$; this can only happen for $p=5$. This case can be ruled out by computer. \square

References

- [1] **S. Ball**, The number of directions determined by a function over a finite field, *J. Combin. Theory Ser. A* **104** (2003), 341–350.
- [2] **A. Blokhuis**, On the size of a blocking set in $\text{PG}(2, p)$, *Combinatorica* **14** (1994), 273–275.
- [3] **A. Blokhuis**, **A. E. Brouwer** and **T. Szőnyi**, The number of directions determined by a function f on a finite field, *J. Combin. Theory Ser. A* **70** (1995), 349–353.
- [4] **A. Blokhuis**, **S. Ball**, **A. Brouwer**, **L. Storme** and **T. Szőnyi**, On the number of slopes of the graph of a function defined on a finite field, *J. Combin. Theory Ser. A* **86** (1999), 187–196.
- [5] **W. S. Chou**, *Permutation polynomials on finite fields and combinatorial applications*, Ph.D. thesis, Penn. State Univ., 1990.
- [6] **R. J. Evans**, **J. Green** and **H. Niederreiter**, Linearized polynomials and permutation polynomials of finite fields, *Michigan Math. J.* **39** (1992) 405–413.
- [7] **A. Gács**, On a generalization of Rédei's theorem, *Combinatorica* **23** (2003), 585–598.

ACADEMIA
PRESS





page 13 / 13

go back

full screen

close

quit

- [8] **R. Lidl** and **H. Niederreiter**, *Finite fields*, Cambridge University Press, 1997.
- [9] **L. Lovász** and **A. Schrijver**, Remarks on a theorem of Rédei, *Studia Scient. Math. Hungar.* **16** (1981), 449–454.
- [10] **O. Polverino**, **T. Szőnyi** and **Zs. Weiner**, Blocking sets in Galois planes of square order, *Acta Sci. Math. (Szeged)* **65** (1999), 737–748.
- [11] **L. Rédei**, *Lückenhafte Polynome über endlichen Körpern*, Birkhäuser Verlag, Basel, 1970. (English translation: *Lacunary Polynomials over Finite Fields*, North Holland, Amsterdam, 1973.)
- [12] **P. Sziklai**, On small blocking sets and their linearity, *J. Combin. Theory Ser. A* **115** (2008), 1167–1182.

András Gács

DEPARTMENT OF COMPUTER SCIENCE, EÖTVÖS LORÁND UNIVERSITY, H-1117 BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY

e-mail: gacs@cs.elte.hu

László Lovász

DEPARTMENT OF COMPUTER SCIENCE, EÖTVÖS LORÁND UNIVERSITY, H-1117 BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY

e-mail: lovasz@cs.elte.hu

Tamás Szőnyi

DEPARTMENT OF COMPUTER SCIENCE, EÖTVÖS LORÁND UNIVERSITY, H-1117 BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY

COMPUTER AND AUTOMATION RESEARCH INSTITUTE OF THE HUNGARIAN ACADEMY OF SCIENCES, H-1111 BUDAPEST, LÁGYMÁNYOSI U. 11, HUNGARY

e-mail: szonyi@cs.elte.hu, szonyi@sztaki.hu

ACADEMIA
PRESS

