



Innovations in Incidence Geometry

Algebraic, Topological and Combinatorial



On triples of ideal chambers in A_2 -buildings

Anne Parreau



On triples of ideal chambers in A_2 -buildings

Anne Parreau

We investigate the geometry in a real Euclidean building X of type A_2 of some simple configurations in the associated projective plane at infinity \mathbb{P} , seen as ideal configurations in X , and relate it with the projective invariants (from the cross ratio on \mathbb{P}). In particular we establish a geometric classification of generic triples of ideal chambers of X and relate it with the triple ratio of triples of flags.

Introduction

The triples of objects in the boundaries of geometric spaces X are basic tools, for example in the study of surface group representations. For instance, in the case where $X = \mathbb{H}^2$, the hyperbolic plane, ideal triples of points may be used to define the notion of Euler class [Goldman 1980], and Penner–Thurston shear coordinates on the Teichmüller space. In the case where $X = \mathbb{H}_{\mathbb{C}}^2$, the ideal triples are classified by Cartan’s angular invariant, see for example [Goldman 1999, §7.1], and they may be for instance used to define Toledo’s invariant and maximal representations [Toledo 1989]. See for instance [Clerc and Neeb 2006; Burger et al. 2010] for generalization to higher rank Hermitian symmetric spaces X , and triples in their Shilov boundary.

For higher rank symmetric spaces X of type A_{N-1} , corresponding to the group $\mathrm{PGL}_N(\mathbb{R})$, ideal configurations in X may be seen as configurations in the projective space $\mathbb{P} = \mathbb{P}(\mathbb{R}^N)$. In particular, ideal chambers of X correspond to complete flags in \mathbb{P} , and opposite pairs of flags (or generic N -tuples of points) in \mathbb{P} correspond to maximal flats in X . This is still true in the non-Archimedean setting, i.e., replacing \mathbb{R} by an ultrametric valued field \mathbb{K} , in which case X is a Euclidean building of type A_{N-1} .

MSC2010: primary 51E24; secondary 20E42.

Keywords: Euclidean buildings, projective planes, ideal configurations, triple ratio.

Configurations in projective spaces $\mathbb{P}(\mathbb{R}^N)$ have been widely studied and used. In particular, triples of flags in $\mathbb{P}(\mathbb{R}^N)$ and their classical invariants (the triple ratio for $N = 3$), are the basic building blocks used by Fock and Goncharov [2006] to define generalized shearing coordinates for higher Teichmüller space, parametrizing positive representations of punctured surface groups in $G = \mathrm{SL}_N(\mathbb{R})$. But the geometric properties in the symmetric space or Euclidean building X of these configurations remain mysterious.

In this article, we investigate the geometry of some simple ideal configurations in a (not necessarily discrete) Euclidean building X of type A_2 , mainly the generic triples of ideal chambers, and the relationship with their projective geometry in the projective plane \mathbb{P} . Our first motivation is to use it to study actions of surface groups on Euclidean buildings of type A_2 , and degenerations of Hitchin representations in $\mathrm{SL}_3(\mathbb{R})$ (see [Parreau 2015]).

The main result is a classification of ideal triples of chambers by the geometry of the five naturally associated flats in X , in relation with their triple ratio as triples of flags in \mathbb{P} . In the case where X is a real tree (e.g., a Euclidean building of type A_1), any generic ideal triple bounds a *tripod* in X , that is a convex subset consisting of union of three rays from a point $x \in X$ (the *center* of the tripod). This is no longer the case in general in higher rank buildings like A_2 -buildings, and many types of configurations are possible. A special case was studied by A. Balsler [2008], who established a characterization of triples of points in $\partial_\infty X$ bounding a tripod in X , and used it to study convex rank 1 subsets in A_2 -buildings. We give here a complete and precise description.

We now get into more details. Let X be a real Euclidean building of (vectorial) type A_2 , i.e., with model flat the Euclidean plane

$$\mathbb{A} = \left\{ \lambda = (\lambda_1, \lambda_2, \lambda_3) \in \mathbb{R}^3 / \sum_i \lambda_i = 0 \right\}$$

endowed with the finite reflection group $W = \mathfrak{S}_3$ acting by permutation of the coordinates. Note that X is not necessarily discrete (simplicial) nor locally compact, and possibly exotic.

The boundary at infinity of X may be identified with the incidence graph of an associated projective plane $\mathbb{P} = \mathbb{P}_\infty(X)$, equipped with an \mathbb{R} -valued additive cross ratio β (called a projective valuation in [Tits 1986]) defined on quadruples of pairwise distinct collinear points in \mathbb{P} [Tits 1986]. In the algebraic case, i.e., when X is the Bruhat–Tits building $X(\mathbb{K}^3)$ associated with the group $\mathrm{PGL}(\mathbb{K}^3)$ for some ultrametric field \mathbb{K} , the projective plane \mathbb{P} is $\mathbb{P}(\mathbb{K}^3)$ and β is the logarithm

$$\beta = \log|b|$$

of the absolute value of the usual \mathbb{K} -valued cross ratio \mathbf{b} on $\mathbb{P}(\mathbb{K}^3)$, where conventions on cross ratios are taken such that

$$\mathbf{b}(\infty, -1, 0, Z) = Z$$

in $\mathbb{P}^1\mathbb{K} = \mathbb{K} \cup \{\infty\}$ (following [Fock and Goncharov 2006]). We will then call $\boldsymbol{\beta}$ the *geometric* cross ratio and \mathbf{b} the *algebraic* cross ratio to distinguish between them.

We now turn to ideal triples of chambers. Let $T = (F_1, F_2, F_3)$ be a triple of chambers at infinity of X . We denote by $F_i = (p_i, D_i)$ the corresponding flag of \mathbb{P} , with p_i the point and D_i the line. The set $\{1, 2, 3\}$ of indices will be canonically identified with $\mathbb{Z}/3\mathbb{Z}$. A triple $T = (F_1, F_2, F_3)$ will be called *generic* if the flags $(F_i)_i$ are pairwise opposite, the points $(p_i)_i$ are not collinear and the lines $(D_i)_i$ are not concurrent.

In the algebraic case $\mathbb{P} = \mathbb{P}(\mathbb{K}^3)$ generic triples of flags $T = (F_1, F_2, F_3)$ are classified by one \mathbb{K} -valued invariant, the (*algebraic*) triple ratio (see for example [Fock and Goncharov 2006, §9.4]), that may be defined by:

$$\text{Tri}(F_1, F_2, F_3) = \mathbf{b}(D_1, p_1p_2, p_1p_{23}, p_1p_3) \tag{0-1}$$

where $p_{ij} = D_i \cap D_j$. We recall that it is invariant under cyclic permutations of T , and that reversing the order inverses the algebraic triple ratio: $\text{Tri}(\bar{T}) = \text{Tri}(T)^{-1}$, where $\bar{T} = (F_3, F_2, F_1)$.

In the general case, we introduce an invariant for generic triples of flags in \mathbb{P} , analogous to the algebraic triple ratio: the *geometric triple ratio*, which still make sense when the building X is exotic (nonalgebraic), whereas the algebraic triple ratio is not defined anymore. We define it as the triple

$$\text{tri}(T) = (\text{tri}_m(T))_{m=1,2,3}$$

of the following cross ratios in \mathbb{P} , which are the cross ratios obtained from the four lines $D_1, p_1p_2, p_1p_{23}, p_1p_3$ by cyclic permutation of the three last one:

$$\begin{aligned} \text{tri}_1(F_1, F_2, F_3) &= \boldsymbol{\beta}(D_1, p_1p_2, p_1p_{23}, p_1p_3) \\ \text{tri}_2(F_1, F_2, F_3) &= \boldsymbol{\beta}(D_1, p_1p_3, p_1p_2, p_1p_{23}) \\ \text{tri}_3(F_1, F_2, F_3) &= \boldsymbol{\beta}(D_1, p_1p_{23}, p_1p_3, p_1p_2). \end{aligned}$$

To simplify notations, we denote from now on

$$z_m = \text{tri}_m(T) \quad \text{and} \quad z = (z_1, z_2, z_3) = \text{tri}(T)$$

In the algebraic case, we have $\mathbb{P} = \mathbb{P}(\mathbb{K}^3)$ and the geometric triple ratio is obtained from the algebraic cross ratio $Z = \text{Tri}(T) \in \mathbb{K}$ by:

$$z_1 = \log|Z|, \quad z_2 = \log\left|\frac{1}{1+Z}\right| = -\log|1+Z|, \quad z_3 = \log|1+Z^{-1}|.$$

The geometric triple ratio z enjoys the following properties. It is invariant by cyclic permutations of the flags, and changed to $(-z_1, -z_3, -z_2)$ by permutations reversing the cyclic order. We also have $z_1 + z_2 + z_3 = 0$, and the stronger following property: for all $m \in \mathbb{Z}/3\mathbb{Z}$, if $z_m > 0$ then $z_{m-1} = 0$ and $z_{m+1} = -z_m < 0$. Note that the three natural cases: $z \in \mathbb{R}_+(0, 1, -1)$, $z \in \mathbb{R}_+(-1, 0, 1)$, and $z \in \mathbb{R}_+(1, -1, 0)$ subdivide in two types, as the case $z_1 = 0$ is invariant under reversing the order of T , whereas the two other cases are exchanged.

We now turn to the geometry inside the Euclidean building X . A generic triple $T = (F_1, F_2, F_3)$ of ideal chambers defines five natural flats in X : the three flats $A_{ij} = A(F_i, F_j)$ containing the opposite chambers F_i and F_j in their boundaries, the flat $A_p = A(p_1, p_2, p_3)$ containing the triple of ideal singular points (p_1, p_2, p_3) in its boundary, and the similarly defined flat $A_D = A(D_1, D_2, D_3)$. We will show that there are also six particular points in X naturally associated with the triple T , that may be defined as the orthogonal projections y_i and y_i^* (which happen to be unique) of p_i and D_i on the flat A_{jk} where $j = i + 1$ and $k = i + 2$.

We say that (F_1, F_2, F_3) is of type “tripod” if there exists a tripod in X joining the three (middle points of the) ideal chambers (F_1, F_2, F_3) . The set of centers of such tripods is the intersection I of the three flats A_{ij} .

We show that either the three flats A_{ij} have a nonempty intersection, that is (F_1, F_2, F_3) is of type “tripod”, or the two flats A_p and A_D have non empty intersection Δ , which is then a *flat singular triangle* (that is, a triangle in \mathbb{A} with singular sides) (we then say that (F_1, F_2, F_3) is of type “flat”). The two following results describe more precisely the two possible types, and relate them with the points y_i, y_i^* and the geometric triple ratio z . We denote by $\mathfrak{C} = \{\lambda \in \mathbb{A} / \lambda_1 > \lambda_2 > \lambda_3\}$ the model Weyl chamber of \mathbb{A} and we use the corresponding *simple roots coordinates* on \mathbb{A} , that is $\lambda = (\lambda_1 - \lambda_2, \lambda_2 - \lambda_3)$.

Theorem 0.1 (type “tripod”). *The intersection $I = A_{12} \cap A_{23} \cap A_{31}$ is nonempty if and only if $z_1 = 0$. Then $z_2 \geq 0$ and there exist a unique pair (x, x^*) in X such that:*

- (i) $y_1 = y_2 = y_3 = x$ and $y_1^* = y_2^* = y_3^* = x^*$.
- (ii) I is the segment $[x, x^*]$.
- (iii) $[x, x^*]$ is the unique shortest segment joining A_p to A_D .
- (iv) Identifying A_{ij} with \mathbb{A} by a marked flat $f : \mathbb{A} \mapsto A_{ij}$ sending \mathfrak{C} to F_j , in simple roots coordinates, we have $\overrightarrow{x x^*} = (-z_2, z_2)$. In particular x^* is on the ray $[x, p_{ij})$ from x to p_{ij} .

Theorem 0.2 (type “flat”). *The intersection $A_p \cap A_D$ is nonempty if and only if $(z_2 = 0$ or $z_3 = 0)$, or, equivalently, if and only if $z_2 \leq 0$. Then there exists a unique flat singular triangle $\Delta \subset X$ with vertices x_1, x_2, x_3 such that:*




Figure 1. Type “tripod”. Bottom row: projections in the flat A_{ij} (left), in the flat A_p (middle) and in the flat A_D .

- (i) $A_p \cap A_D = \Delta$.
- (ii) $A_{ij} \cap A_{ik}$ is the Weyl chamber from x_i to F_i .
- (iii) Let $i \in \{1, 2, 3\}$ and $j = i + 1$. In a marked flat $f : \mathbb{A} \mapsto A_{ij}$ sending \mathfrak{C} to F_j , in simple roots coordinates, we have $\overrightarrow{x_i x_j} = (z_1^+, z_1^-)$ where $z_1^+ = \max(z_1, 0)$ and $z_1^- = \max(-z_1, 0)$. In particular x_j is on the ray from x_i to p_j (if $z_1 \geq 0$) or D_j (if $z_1 < 0$).
- (iv) The germs of Weyl chambers at x_i respectively defined by Δ and F_i are opposite (in the spherical building of directions at x_i). In particular there exists a flat containing Δ , and containing F_i in its boundary.

Furthermore if $z_1 \geq 0$ we have $x_i = y_{i-1} = y_{i+1}^*$ for all i , and if $z_1 \leq 0$ we have $x_i = y_{i+1} = y_{i-1}^*$ for all i .

The intersections of each flat with the four other flats form a partition (i.e., a covering with disjoint interiors), which is described in Figure 1 for the type “tripod”, and in Figure 2 for the type “flat” (see Proposition 4.2, Corollary 4.3 and Proposition 4.5).

The special case where the hypotheses of both Theorems 0.1 and 0.2 are satisfied corresponds to the case where $z_1 = z_2 = z_3 = 0$. Then the five flats intersect in




Figure 2. Type “flat”, in the case where $z_1 \geq 0$. Bottom row: projections in A_{ij} , with $j = i + 1$. (left), in A_p (middle) and in A_D . The case $z_1 \leq 0$ is obtained from the case $z_1 \geq 0$ by reversing the order of the flags F_i , that is, by exchanging 1 and 3 and i and j in the diagrams.

a unique point x , and, in the spherical building of directions at x , the triple of chambers induced by $T = (F_1, F_2, F_3)$ is generic.

In particular we recover the characterization of [Balser 2008] for triples of points in $\partial_\infty X$ bounding a tripod in X . Note that M. Talbi [2006] established some analogous geometric classification for interior triangles in discrete Euclidean buildings of type A_2 .

Theorem 0.2 will be used in [Parreau 2015] to study actions of punctured surface groups on Euclidean buildings of type A_2 . It allows us to give a metric interpretation, in the building, of Fock–Goncharov parameters associated with ideal triangulations. We are then able to construct in X an invariant weakly convex cocompact 2-complex for large families of actions. Theorem 0.2 enables us to associate to each triangle of the triangulation a flat singular triangle in X , the complex is then obtained by connecting them gluing flat strips. This allows to describe length spectra for large families of degenerations of convex projective structures on surfaces.

We also show that generic quadruples of points in \mathbb{P} (which will be called *projective frames*) define a nice center in X , with various characterizations, see Proposition 2.4 (this result generalizes to higher rank \mathbb{R} -buildings of type A_{N-1}).

1. Preliminaries

1A. The model flat (\mathbb{A}, W) of type A_{N-1} . Let $N \geq 2$ be an integer. The *model flat* of type A_{N-1} is the vector space $\mathbb{A} = \mathbb{R}^N / \mathbb{R}(1, \dots, 1)$, endowed with the action of the *Weyl group* $W = \mathfrak{S}_N$ acting on \mathbb{A} by permutation of coordinates (finite reflection group). We denote by $[\lambda]$ the projection in \mathbb{A} of a vector λ in \mathbb{R}^N . The vector space \mathbb{A} may be identified with the hyperplane $\{\lambda = (\lambda_1, \dots, \lambda_N) \in \mathbb{R}^N / \sum_i \lambda_i = 0\}$ of \mathbb{R}^N . Recall that a vector in \mathbb{A} is called *singular* if it belongs to one the hyperplanes $\lambda_i = \lambda_j$, and *regular* otherwise. A (*open*) (*vectorial*) *Weyl chamber* of \mathbb{A} is a connected component of regular vectors. We will call a *sector* a more general convex cone in \mathbb{A} , in particular the closed convex cone formed by the union of the closed Weyl chambers containing a given singular ray. The *model Weyl chamber* is the simplicial cone

$$\mathfrak{C} = \{\lambda \in \mathbb{A} / \lambda_1 > \dots > \lambda_N\}.$$

Its closure $\bar{\mathfrak{C}}$ is a strict fundamental domain for the action of W on \mathbb{A} . Recall that two nonzero vectors λ and λ' of \mathbb{A} are called *opposite* if $\lambda' = -\lambda$. Similarly, two Weyl chambers C and C' of \mathbb{A} are *opposite* if $C' = -C$. The *type* of a vector $\lambda \in \mathbb{A}$ is its projection (modulo W) in $\bar{\mathfrak{C}}$.

We denote by $\partial\mathbb{A}$ the sphere of unitary vectors in \mathbb{A} , identified with the set $\mathbb{P}^+(\mathbb{A}) = (\mathbb{A} - \{0\}) / \mathbb{R}_{>0}$ of rays issued from 0, and by $\partial : \mathbb{A} - \{0\} \rightarrow \partial\mathbb{A}$ the corresponding projection. The *type (of direction)* of a nonzero vector $\lambda \in \mathbb{A}$ is its canonical projection in $\partial\bar{\mathfrak{C}}$.

We denote by $(\varepsilon_1, \dots, \varepsilon_N)$ the canonical basis of \mathbb{R}^N . For $d = 1, \dots, N - 1$, we will say that a nonzero vector in \mathbb{A} (or a point in the sphere $\partial\mathbb{A}$) is *singular of type d* if its canonical projection in $\partial\bar{\mathfrak{C}}$ is $[\varepsilon_1 + \dots + \varepsilon_d]$.

The *simple roots* (associated with \mathfrak{C}) are the following linear forms on \mathbb{A}

$$\varphi_i : \lambda \mapsto \lambda_i - \lambda_{i+1}$$

for $i = 1, \dots, N - 1$. The set of simple roots is denoted by Λ . We will also use the root $\varphi_N : \lambda \mapsto \lambda_N - \lambda_1$ satisfying

$$\varphi_1 + \dots + \varphi_N = 0.$$

The vector space \mathbb{A} is endowed with the unique W -invariant Euclidean scalar product, which is well defined up to homothety (induced by the standard Euclidean scalar product of \mathbb{R}^N). We will normalize it by requiring that the simple roots have unit norm, i.e., the distance between the two hyperplanes with equation $\varphi_i = 0$ and $\varphi_i = 1$ is 1 for one (all) i . When $\dim \mathbb{A} = 1$, we will identify \mathbb{A} with \mathbb{R} by the basis $\{[\varepsilon_1]\}$, i.e., by the map from $s \mapsto [(s, 0)]$ from \mathbb{R} to \mathbb{A} , which is an isometry in the above normalization.




Figure 3. The model flat \mathbb{A} of type A_2 (for $N = 3$), and simple roots coordinates. The arrows denote the singular directions of type 1.

1B. Projective spaces. We here collect the notations and vocabulary for projective spaces, which will be used throughout this article. We refer to [Tits 1974, §6.2]. Let \mathbb{P} be a projective space of dimension $N - 1$, with $N \geq 2$. We denote by $\text{Flags}(\mathbb{P})$ the set of flags of \mathbb{P} , that is increasing sequences (V_1, \dots, V_M) of proper linear subspaces of \mathbb{P} . We denote by \mathbb{P}^* the dual projective space, whose set of points is the set of hyperplanes of \mathbb{P} .

Two maximal flags (V_1, \dots, V_{N-1}) and (V'_1, \dots, V'_{N-1}) are *opposite* if they are in generic position, that is if $V_i \oplus V'_{n-i} = \mathbb{P}$ for all i . A finite subset p_1, \dots, p_M in \mathbb{P} , with $2 \leq M \leq N$, is called *independent* if it is not contained in any linear subspace of dimension $M - 2$ of \mathbb{P} . Then it is contained in a unique $(M - 1)$ -dimensional linear subspace of \mathbb{P} , which will be denoted by $p_1 \oplus \dots \oplus p_M$. When $M = 2$, we will also denote the line $p \oplus q$ by pq .

A *frame* of \mathbb{P} is an independent N -tuple. A *projective frame* in \mathbb{P} is a $(N + 1)$ -tuple (p_0, p_1, \dots, p_N) of points in \mathbb{P} in generic position, i.e., such that the induced N -tuple $(p_0, \dots, \hat{p}_i, \dots, p_N)$ is a frame in \mathbb{P} for all i .

If p is a point in \mathbb{P} , we denote by \mathbb{P}/p the set of lines through p , which is a projective space of dimension $N - 2$ whose linear subspaces are the linear subspaces of \mathbb{P} containing p . The *projection at p* is the corresponding projection $\text{proj}_p : q \mapsto pq$ from $\mathbb{P} - \{p\}$ to \mathbb{P}/p . If p is a point of \mathbb{P} and $H \subset \mathbb{P}$ an hyperplane with $p \notin H$, then the projection proj_p induces a canonical isomorphism $\text{proj}_{H_p} : H \xrightarrow{\sim} \mathbb{P}/p$ (called *perspectivity*).

Note that if $\mathcal{F} = (p_1, \dots, p_M)$ is independent in \mathbb{P} , then its projection $\text{proj}_{p_1}(\mathcal{F}) = (p_1 p_2, \dots, p_1 p_M)$ at p_1 is independent in \mathbb{P}/p_1 . In particular the projection of a (projective) frame at one of its points is a (projective) frame.

1C. Spherical buildings of type A_{N-1} and associated projective spaces. See [Tits 1974, §6]. A spherical building \mathcal{B} of type A_{N-1} is the building of flags of an associated projective space $\mathbb{P} = \mathbb{P}(\mathcal{B})$ of dimension $N - 1$. For $d = 0, 1, \dots, N - 1$, the set of linear subspaces of dimension d of \mathbb{P} identifies with the subset of vertices of type $d + 1$ of \mathcal{B} . In particular, the projective space \mathbb{P} itself is identified with the set of vertices of type 1 of \mathcal{B} , and the dual projective space \mathbb{P}^* is identified with the set of vertices of type $N - 1$.

In the algebraic case, that is when \mathcal{B} is the spherical building of flags of some vector space V of dimension N over a field \mathbb{K} , then $\mathbb{P} = \mathbb{P}(V)$.

A basic fact is that frames in \mathbb{P} correspond to apartments of \mathcal{B} .

Recall that, in (the geometric realization modeled on $(\partial\mathbb{A}, W)$ of) a spherical building, any two points (resp. chambers) are contained in a common apartment, and that they are *opposite* if they are opposite in that apartment, that is, for two points ξ and ξ' , if and only if $\angle(\xi, \xi') = \pi$ for the canonical metric \angle on \mathcal{B} . Note that $p \in \mathbb{P}$ and $H \in \mathbb{P}^*$ are opposite if and only if $\angle(p, H) = \pi$, if and only if $p \notin H$. Two chambers are opposite if and only if they are opposite as maximal flags in \mathbb{P} . In particular, in the type A_2 case, two chambers $F_1 = (p_1, D_1)$, $F_2 = (p_2, D_2)$ are opposite if and only if $p_1 \notin D_2$ and $p_2 \notin D_1$.

For any simplex σ of \mathcal{B} the *residue* $\text{St}(\sigma)$ of σ is the spherical building formed by the simplices of \mathcal{B} containing σ . If H is a hyperplane of \mathbb{P} , the residue $\text{St}(H)$ of H in \mathcal{B} is the subset of flags of \mathbb{P} containing H . It canonically identifies with the spherical building $\text{Flags}(H)$ of flags of H by the map $(V_1, \dots, V_M, H) \mapsto (V_1, \dots, V_M)$. The residue $\text{St}(p)$ of a point p in \mathbb{P} identifies canonically with the flag building $\text{Flags}(\mathbb{P}/p)$ of \mathbb{P}/p by the map $(V_1 = p, \dots, V_M) \mapsto (V_2/p, \dots, V_M/p)$. If $p \notin H$ then the projection proj_p induces a canonical isomorphism $\text{proj}_{Hp} : \text{St}(H) \xrightarrow{\sim} \text{St}(p)$ of spherical buildings (perspectivity).

1D. Euclidean buildings. Euclidean buildings considered in this article are (not necessarily discrete) Euclidean buildings of type A_{N-1} . We refer for example to [Parreau 2000] for the definition and properties of Euclidean buildings we use below (see also [Tits 1986; Kleiner and Leeb 1997; Rousseau 2009]). Recall that a *Euclidean building of type A_{N-1}* is a CAT(0) metric space X endowed with a (maximal) collection \mathcal{A} of isometric embeddings $f : \mathbb{A} \rightarrow X$ called *marked apartments*, or *marked flats* by analogy with Riemannian symmetric spaces, satisfying the following properties:

- (A1) \mathcal{A} is invariant by precomposition by W_{aff} .
- (A2) If f and f' are two marked flats, then the transition map $f^{-1} \circ f'$ is the restriction of an element of W_{aff} .
- (A3') Any two rays of X are initially contained in a common marked flat.

Where W_{aff} denotes the subgroup of all affine isomorphisms of \mathbb{A} with linear part in W . The *flats* and the *Weyl chambers* of X are the images of \mathbb{A} and \mathcal{C} by the marked flats, respectively.

Algebraic case. Let \mathbb{K} be an ultrametric field, i.e., a field endowed with an ultrametric absolute value $|\cdot|$ (not necessarily discrete). When V is a finite N -dimensional vector space over \mathbb{K} , we denote by $X = X(V)$ the Euclidean building associated with $G = \text{PGL}(V)$. We refer for example to [Parreau 2000] for the model of norms for X (see [Goldman and Iwahori 1963; Bruhat and Tits 1984]). To each basis \mathbf{v} of V is then associated a marked flat $f_{\mathbf{v}} : \mathbb{A} \rightarrow A_{\mathbf{v}} \subset X$, such that, if a is an element of G with diagonal matrix $\text{diag}(a_1, \dots, a_N)$ in the basis \mathbf{v} , then a translates the flat $A_{\mathbf{v}}$ by the vector

$$\nu(a) = [(\log|a_i|)_i]$$

in \mathbb{A} (identifying the flat $A_{\mathbf{v}}$ with the model flat \mathbb{A} through the marking $f_{\mathbf{v}}$).

From now to Section 1H, X will denote a Euclidean building of type A_{N-1} .

1E. Spherical building and projective space at infinity. The $\text{CAT}(0)$ boundary $\partial_{\infty}X$ of X is the geometric realization modeled on $(\partial\mathbb{A}, W)$ of a spherical building of type A_{N-1} whose chambers are the boundaries of the Weyl chambers of X , and whose apartments are the boundaries of the flats of X . It will be identified with the building of flags on the associated projective space $\mathbb{P} = \mathbb{P}_{\infty}(X)$, whose points are the vertices of type 1 of $\partial_{\infty}X$. If c_+ and c_- are opposite ideal chambers, then we denote by $A(c_-, c_+)$ the unique flat *joining* c_- to c_+ in X , that is, containing c_- and c_+ in its boundary. If \mathcal{F} is a frame of \mathbb{P} or \mathbb{P}^* , then there is a unique flat $A(\mathcal{F})$ of X containing \mathcal{F} in its boundary.

1F. Local spherical building and projective space at a point. Recall that, in Euclidean buildings, two (unit speed) geodesic segments issued from a common point x have zero angle if and only if they have same germ at x (i.e., coincide in a neighborhood of x). A *direction* at $x \in X$ is a germ of nontrivial geodesic segment from x . A direction, geodesic segment, ray or line has a well-defined *type (of direction)* in $\partial\bar{\mathcal{C}}$, which is its canonical projection (through a marked flat) in $\partial\bar{\mathcal{C}}$. It is called *singular* or *regular* accordingly.

The *space of directions* at x of X is the quotient space of non trivial geodesic segments from x for this relation, with the induced angular metric, and is denoted by $\Sigma_x X$. We denote by $\Sigma_x : X - \{x\} \rightarrow \Sigma_x X, y \rightarrow \Sigma_x y$, the associated projection. Its extension to the boundary at infinity will also be denoted by $\Sigma_x : \partial_{\infty}X \rightarrow \Sigma_x X, \xi \rightarrow \Sigma_x \xi$ and called the *canonical projection*.

The space of directions $\Sigma_x X$ inherits the structure of a spherical A_{N-1} -building, whose apartments are the germs $\Sigma_x A$ at x of the flats A of X passing through x , and whose chambers are the germs $\Sigma_x C$ at x of the Weyl chambers C of X with vertex

x (see for example [Parreau 2000]). The canonical projection $\Sigma_x : \partial_\infty X \rightarrow \Sigma_x X$ sends chambers to chambers (and, more generally, simplices to simplices) and preserves the type of points.

The *local projective space* $\mathbb{P}_x = \mathbb{P}_x(X)$ at x is the projective space of dimension $N - 1$ associated with the spherical building $\Sigma_x X$ of type A_{N-1} (see Section 1C). Its underlying set is the set of vertices of type 1 of $\Sigma_x X$.

The canonical projection $\Sigma_x : \partial_\infty X \rightarrow \Sigma_x X$ induces (by restriction to vertices) a surjective morphism (of projective spaces) $\Sigma_x : \mathbb{P} \rightarrow \mathbb{P}_x$ from the projective space at infinity \mathbb{P} to the local projective space \mathbb{P}_x at x . Note that, in particular, if \mathcal{F} is a frame of \mathbb{P} , then x belongs to the associated flat $A(\mathcal{F})$ if and only if $\Sigma_x(\mathcal{F})$ is a frame of \mathbb{P}_x .

1G. Transverse spaces at infinity. See for example [Tits 1986, §8; Leeb 2000, 1.2.3; Mühlherr et al. 2014, §4]. Let ξ be a vertex of $\partial_\infty X$ of type 1 or $N - 1$, i.e., either a point p in the projective plane at infinity \mathbb{P} or a hyperplane H of \mathbb{P} .

The *transverse space* X_ξ at ξ may be defined, from the metric viewpoint (as in [Leeb 2000, 1.2.3]), as the quotient space of the set of all rays to ξ by the pseudodistance d_ξ given by

$$d_\xi(r_1, r_2) = \inf_{t_1, t_2} d(r_1(t_1), r_2(t_2)).$$

We denote by $\pi_\xi : X \rightarrow X_\xi$ the canonical projection (which maps x to the class of the unique ray from x to ξ). The space X_ξ is a Euclidean building of type A_{N-2} , whose flats are the projections to X_ξ of the flats of X containing a ray to ξ . In particular, when X is of type A_2 , the transverse space X_ξ is an \mathbb{R} -tree, and we will call it the *transverse tree* at ξ .

In the algebraic case, i.e., when $X = X(V)$, the transverse space X_H canonically identifies with the building $X(H)$ of H , where H is seen as an hyperplane of V , and X_p identifies with $X(V/p)$, where p is seen as a 1-dimensional subspace of V .

The spherical building $\partial_\infty X_\xi$ at infinity of X_ξ identifies canonically with the residue $\text{St}(\xi)$ of ξ . In particular, if p is a point in \mathbb{P} , the projective space at infinity of X_p identifies with \mathbb{P}/p , and if H is an hyperplane of \mathbb{P} , the projective space at infinity of X_H identifies with H .

If $\mathcal{F} = (p_1, \dots, p_N)$ is a frame in $\mathbb{P} \subset \partial_\infty X$, then the projection on X_{p_1} of the flat $A(p_1, \dots, p_N)$ is the flat defined by the projection $\text{proj}_{p_1}(\mathcal{F}) = (p_1 p_2, \dots, p_1 p_N)$ of the frame \mathcal{F} , i.e., $\pi_{p_1}(A(\mathcal{F})) = A(\text{proj}_{p_1}(\mathcal{F}))$.

We now describe the canonical isomorphism $\pi_{\xi^-\xi^+} : X_{\xi^-} \xrightarrow{\sim} X_{\xi^+}$ for opposite points ξ^- and ξ^+ of $\partial_\infty X$. The union $F_{\xi^-\xi^+}$ of all geodesics joining ξ^- to ξ^+ is a convex closed subspace and a subbuilding, whose flats are the flats of X containing a geodesic joining ξ^- to ξ^+ (see [Kleiner and Leeb 1997, Proposition 4.8.1] and [Parreau 2012, 2.2.1]). We denote by $F_{\xi^-\xi^+} = X^{\xi^-\xi^+} \times \mathbb{R}$ the

canonical decomposition (see [Parreau 2011, 1.2.10]). The restriction of the projection π_{ξ^+} to F_{ξ^+} is surjective and factorizes through the projection on the first factor, inducing a canonical isomorphism of Euclidean buildings $X^{\xi^+} \xrightarrow{\sim} X_{\xi^+}$. We similarly have a isomorphism $X^{\xi^-} \xrightarrow{\sim} X_{\xi^-}$, so it induces a canonical isomorphism $\pi_{\xi^-} : X_{\xi^-} \xrightarrow{\sim} X_{\xi^+}$. It is easy to see that the map π_{ξ^-} extends to the boundaries at infinity of X_{ξ^-} and X_{ξ^+} by the canonical isomorphism of spherical buildings $\text{proj}_{\xi^-} : \text{St}(\xi^-) \xrightarrow{\sim} \text{St}(\xi^+)$ (perspectivity).

1H. The \mathbb{A} -valued Busemann cocycle. Let c be a chamber at infinity of X . We now define the \mathbb{A} -valued *Busemann cocycle*

$$B_c : X \times X \rightarrow \mathbb{A}$$

associated to c . It can be simply defined from canonical retractions as

$$B_c(x, y) := r(y) - r(x)$$

where $r : X \rightarrow \mathbb{A}$ is any canonical retraction centered at c , sending c to $\partial\mathcal{C}$ (see [Parreau 2000, Proposition 1.19]). More precisely, the Busemann cocycle at c is characterized by the property:

$$B_c(f(\lambda), f'(\lambda')) = \lambda' - \lambda$$

for any two marked flats $f, f' : \mathbb{A} \rightarrow X$ sending $\partial\mathcal{C}$ to c and such that $f = f'$ on some subchamber of \mathcal{C} .

We clearly have

$$B_c(x, z) = B_c(x, y) + B_c(y, z).$$

When $\dim \mathbb{A} = 1$, it coincides with the usual Busemann cocycle, which is defined for $\xi \in \partial_\infty X$ by

$$B_\xi(x, y) = \lim_{z \rightarrow \xi} d(x, z) - d(y, z).$$

In the type A_2 case, the simple root coordinates of \mathbb{A} -valued Busemann cocycles may be determined by projecting in transverse trees at infinity, using the following relations (using the normalization of the metric).

$$\begin{aligned} \varphi_1(B_{(p,D)}(x, y)) &= B_p(\pi_D(x), \pi_D(y)), \\ \varphi_2(B_{(p,D)}(x, y)) &= B_D(\pi_p(x), \pi_p(y)). \end{aligned} \tag{1-1}$$

We now turn to cross ratios.


1I. Cross ratio on the boundary of a tree. See [Tits 1986, §7], and in a more general setting [Otal 1992; Bourdon 1996]. In this section, we suppose that X is an \mathbb{R} -tree. Given three distinct ideal points ξ_1, ξ_2, ξ_3 in $\partial_\infty X$, we denote by $c(\xi_1, \xi_2, \xi_3)$ the *center* of the ideal triple (ξ_1, ξ_2, ξ_3) , that is the unique common

intersection point of the three geodesic lines joining two of the three points. Note that $c(\xi_1, \xi_2, \xi_3)$ is the (orthogonal) projection of ξ_3 on the geodesic joining ξ_1 to ξ_2 . We denote by $B_{\xi}(x, y)$ the Busemann cocycle (see Section 1H).

Define the *cross ratio* of four pairwise distinct points $\xi_1, \xi_2, \xi_3, \xi_4$ in $\partial_{\infty} X$ by

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) = \frac{1}{2}(\ell_{12} - \ell_{23} + \ell_{34} - \ell_{41})$$

where ℓ_{ij} is the length of the geodesic in X from ξ_i to ξ_j after removing disjoint fixed horoballs centered at each ξ_k . It does not depend on the choice of the horoballs since the horoballs centered at a given point are equidistant along the rays to that point.



The cross ratio naturally extends to *nondegenerate* quadruples, that are quadruples $(\xi_1, \xi_2, \xi_3, \xi_4)$ *without triple point* (i.e., any three of the points are not equal), which is equivalent to the following condition:

$$(\xi_1 \neq \xi_4 \text{ and } \xi_2 \neq \xi_3) \quad \text{or} \quad (\xi_1 \neq \xi_2 \text{ and } \xi_3 \neq \xi_4). \tag{1-2}$$


We then set

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) = \begin{cases} 0 & \text{when } \xi_1 = \xi_3 \text{ or } \xi_2 = \xi_4, \\ -\infty & \text{when } \xi_1 = \xi_2 \text{ or } \xi_3 = \xi_4, \\ +\infty & \text{when } \xi_1 = \xi_4 \text{ or } \xi_2 = \xi_3. \end{cases}$$

We now recall some basic properties that we will use.

The cross ratio may be read inside the tree on the oriented geodesic from ξ_3 to ξ_1 , as the oriented distance \overrightarrow{xy} from the center x of the ideal triple (ξ_3, ξ_1, ξ_2) to the center y of the ideal triple (ξ_3, ξ_1, ξ_4) :

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) = \overrightarrow{xy} = B_{\xi_1}(x, y). \tag{1-3}$$



The cocycle identity is

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) + \beta(\xi_1, \xi_4, \xi_3, \xi_5) = \beta(\xi_1, \xi_2, \xi_3, \xi_5).$$

The cross ratio β is left unchanged by the double transpositions and changed to $-\beta$ by the transpositions (13) and (24). We now consider the behavior under cyclic permutations of the three last terms. We have

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) + \beta(\xi_1, \xi_4, \xi_2, \xi_3) + \beta(\xi_1, \xi_3, \xi_4, \xi_2) = 0. \tag{1-4}$$

Moreover, the following *ultrametricity* property (specific to the case of trees) is easy to prove using (1-3) (see [Tits 1986, §7, Proposition 3]):

If $\beta(\xi_1, \xi_2, \xi_3, \xi_4) > 0$,
 then $\beta(\xi_1, \xi_3, \xi_4, \xi_2) = 0$ and $\beta(\xi_1, \xi_4, \xi_2, \xi_3) = -\beta(\xi_1, \xi_2, \xi_3, \xi_4)$. (1-5)

Note that (1-5) is equivalent (under (1-4)) to

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) \leq \max(0, -\beta(\xi_1, \xi_4, \xi_2, \xi_3)). \quad (1-6)$$

which in the algebraic case follows from the symmetry properties of the cross ratio under 3-cyclic permutations (1-9).

1J. Algebraic case: link with usual cross ratio. Suppose that X is the tree $X(V)$ associated with a 2-dimensional vector space V over an ultrametric field \mathbb{K} (see Section 1D). Then $\partial_\infty X$ identifies with the projective line $\mathbb{P}(V)$.

The usual cross ratio \mathbf{b} on $\mathbb{P}(V)$ of a nondegenerate quadruple of points (see (1-2)) is defined by (following the convention of [Fock and Goncharov 2007], and taking values in $\mathbb{K} \cup \{\infty\}$)

$$\mathbf{b}(a_1, a_2, a_3, a_4) = \frac{(a_1 - a_2)(a_3 - a_4)}{(a_1 - a_4)(a_2 - a_3)} \quad (1-7)$$

in any affine chart $\mathbb{P}(V) \xrightarrow{\sim} \mathbb{K} \cup \{\infty\}$, so that $\mathbf{b}(\infty, -1, 0, a) = a$.

The cross ratio β defined in Section II will then be called the *geometric* cross ratio, to distinguish it from \mathbf{b} , which will be called the *algebraic* cross ratio. They are then related as follows:

$$\beta(\xi_1, \xi_2, \xi_3, \xi_4) = \log|\mathbf{b}(\xi_1, \xi_2, \xi_3, \xi_4)|. \quad (1-8)$$

Proof. Let $x_4 = c(\xi_3, \xi_1, \xi_2)$ and $x_2 = c(\xi_3, \xi_1, \xi_4)$. In a suitable basis $\mathbf{v} = (v_1, v_2)$ of V , we have in homogeneous coordinates $\xi_1 = [1 : 0]$, $\xi_3 = [0 : 1]$, $\xi_2 = [-1 : 1]$ and $\xi_4 = [b : 1]$, where $b = \mathbf{b}(\xi_1, \xi_2, \xi_3, \xi_4)$. Then $g = \begin{pmatrix} -b & 0 \\ 0 & 1 \end{pmatrix}$ fixes ξ_1 and ξ_3 and sends ξ_2 to ξ_4 . Hence $g(x_4) = x_2$. In the flat $A(\xi_3, \xi_1)$ identified with $\mathbb{A} = \mathbb{R}^2/\mathbb{R}(1, 1)$ by the marked flat f_v , we have $\overrightarrow{x_4 x_2} = v(g) = [(\log|b|, 0)]$, hence $\overrightarrow{x_4 x_2} = \log|b|$ as needed. \square

We recall that the algebraic cross ratio \mathbf{b} satisfies the following symmetry properties: It is left unchanged by the double transpositions and changed to \mathbf{b}^{-1} by the transpositions (13) and (24). Furthermore we have an additional symmetry under 3-cycles not satisfied by the geometric cross ratio:

$$\begin{aligned} \mathbf{b}(a_1, a_3, a_4, a_2) &= -1 - \mathbf{b}(a_1, a_2, a_3, a_4)^{-1}, \\ \mathbf{b}(a_1, a_4, a_2, a_3) &= -(1 + \mathbf{b}(a_1, a_2, a_3, a_4))^{-1}. \end{aligned} \quad (1-9)$$

1K. Cross ratio on the boundary of an A_2 -Euclidean building. See [Tits 1986]. Let X be a Euclidean building of type A_2 , and \mathbb{P} the associated projective plane at infinity.

Let (p_1, p_2, p_3, p_4) be a nondegenerate quadruple of points of \mathbb{P} on a common line D . Then their *cross ratio* $\beta(p_1, p_2, p_3, p_4)$ (i.e., *projective valuation* in [Tits 1986]) is by definition their cross ratio as ideal points of the transverse tree X_D . The cross ratio of a nondegenerate quadruple of lines in \mathbb{P} passing through a common point p is similarly defined as their cross ratio as ideal points of the transverse tree X_p .

The main additional property is that perspectivities preserve cross ratio, which follows from the fact that perspectivities extend isometries between the transverse trees (see Section 1G):

Proposition 1.1. *Let p be a point of \mathbb{P} and D a line of \mathbb{P} with $p \notin D$. The canonical isomorphisms (perspectivities) $\text{proj}_{pD} : \text{St}(D) \xrightarrow{\sim} \text{St}(p)$, $q \mapsto pq$ and $\text{proj}_{Dp} : \text{St}(p) \xrightarrow{\sim} \text{St}(D)$, $L \mapsto D \cap L$, preserve the cross ratio β , i.e.,*

$$\begin{aligned} \beta(p_1, p_2, p_3, p_4) &= \beta(pp_1, pp_2, pp_3, pp_4), \\ \beta(D_1, D_2, D_3, D_4) &= \beta(D \cap D_1, D \cap D_2, D \cap D_3, D \cap D_4) \quad \square \end{aligned}$$

2. Some basic ideal configurations

2A. Extension of orthogonal projection to the boundary in CAT(0) spaces. In this section X is a general CAT(0) metric space, and we prove the following basic property: the usual orthogonal projection onto a proper convex subset $Y \subset X$ extends to the boundary outside the closed $\frac{\pi}{2}$ -neighborhood of $\partial_\infty Y$ for the Tits metric (note that the projection is no longer unique). This property is quite elementary but we did not see it in the classical literature, so we include the proof. We refer to the book [Bridson and Haefliger 1999] for CAT(0) spaces.

We denote by $\partial_\infty X$ the CAT(0) boundary of X , and by $\angle_{Tits}(\xi, \eta)$ the Tits angle between two ideal points $\xi, \eta \in \partial_\infty X$. For a subset A of $\partial_\infty X$, we define $\angle_{Tits}(\xi, A) = \inf_{\eta \in A} \angle_{Tits}(\xi, \eta)$.

Definition 2.1. Let Y be a subspace of X and $\xi \in \partial_\infty X$ an ideal point. We say that a point $x \in Y$ is an *orthogonal projection of ξ on Y* if $\angle_x(\xi, y) \geq \frac{\pi}{2}$ for all $y \in Y - \{x\}$.

Proposition 2.2. *Let Y be a convex subspace of a CAT(0) space X which is proper for the induced metric, and ξ in $\partial_\infty X$. Suppose that $\angle_{Tits}(\xi, \partial_\infty Y) > \frac{\pi}{2}$. Then there exists an orthogonal projection x of ξ on Y .*

Proof. Consider a sequence (x_n) converging to ξ in X , and let y_n be the orthogonal projection of x_n on Y . If $(y_n)_{n \in \mathbb{N}}$ is not bounded then, up to passing to a subsequence, y_n converges to η in $\partial_\infty Y$. Then for any fixed y in Y we have $\angle_y(\xi, y_n) \leq \frac{\pi}{2}$

for all n , hence $\angle_y(\xi, \eta) \leq \frac{\pi}{2}$. Therefore $\angle_{Tits}(\xi, \eta) \leq \frac{\pi}{2}$. Thus $(y_n)_{n \in \mathbb{N}}$ is bounded, hence, since Y is proper, it has a converging subsequence, and the limit point x is then an orthogonal projection of ξ on Y . \square

2B. Centers of generic $(N + 1)$ -tuples. In this section, we show that the notion of center of ideal triples in trees extends to Euclidean buildings of type A_{N-1} , for generic $(N + 1)$ -tuples of points (or hyperplanes) in the associated projective space at infinity (Proposition 2.4).

Let X be a Euclidean building of type A_{N-1} , and \mathbb{P} be its projective space at infinity (i.e., the set of singular points of type 1 in $\partial_\infty X$, see Section 1). Recall from Section 1B that a *projective frame* in a projective space of dimension $N - 1$ is a generic $(N + 1)$ -tuple of points.

We first observe that the orthogonal projection of a point of \mathbb{P} on a flat of X exists under a simple necessary and sufficient condition.

Proposition 2.3. *Let A be a flat of X and $p \in \mathbb{P}$. Let $(p_1, \dots, p_N) = (\partial_\infty A) \cap \mathbb{P}$ be the points of type 1 in $\partial_\infty A$. Then p admits an orthogonal projection on A if and only if (p, p_1, \dots, p_N) is a projective frame.*

The analogous property is also valid for points $H \in \mathbb{P}^*$. Note that these properties also hold in symmetric spaces of type A_{N-1} .

Proof. If $p \in H$ for some hyperplane H in $\mathbb{P}^* \cap \partial_\infty A$, then p and H are in a common chamber of the spherical building $\partial_\infty X$, and, as the diameter d of the model spherical Weyl chamber $\partial \bar{\mathcal{C}}$ is strictly less than $\pi/2$ (for the angle metric), we have $\angle_{Tits}(p, H) < \pi/2$, hence the orthogonal projection does not exist. Else, for every hyperplane H in $\mathbb{P}^* \cap \partial_\infty A$, we have $p \notin H$, hence $\angle_{Tits}(p, H) = \pi$, which implies that since $\angle_{Tits}(p, \eta) \geq \pi - d > \pi/2$ for all $\eta \in \partial_\infty A$, and the orthogonal projection exists by Proposition 2.2. \square

We now turn to the main result of this section.

Proposition 2.4. *Let $\mathcal{F} = (p_0, p_1, \dots, p_N)$ be a projective frame in $\mathbb{P} \subset \partial_\infty X$. For each $i \in \{0, \dots, N\}$ let A_i be the unique flat of X through $(p_0, \dots, \hat{p}_i, \dots, p_N)$. There exists a unique point $x \in X$ satisfying the following equivalent conditions:*

- (i) $x \in \cap_i A_i$.
- (ii) For all i and for all H in $\partial_\infty A_i \cap \mathbb{P}^*$ the angle $\angle_x(p_i, H)$ is π .
- (iii) The $(N + 1)$ -tuple $\Sigma_x \mathcal{F} = (\Sigma_x p_i)_{i=0, \dots, N}$ of directions at x form a projective frame in \mathbb{P}_x .
- (iv) For all i , the point x is an orthogonal projection of p_i on the flat A_i .
- (v) There exists i such that x is an orthogonal projection of p_i on A_i .




Figure 4. The center $x \in X$ of a projective frame (p_1, p_2, p_3, p_4) (for $N = 3$).

We will call x the center of the projective frame $\mathcal{F} = (p_0, p_1, \dots, p_N)$ and denote it by $c(p_0, p_1, \dots, p_N)$ or $c(\mathcal{F})$.

Proof. The existence of x , as an orthogonal projection of p_0 on A_0 , is ensured by Proposition 2.3.

For $i \neq j$, denote by H_{ij} the hyperplane $\bigoplus_{k \neq i, j} p_k$ in the projective space \mathbb{P} . Let $x \in X$. Conditions (iii) and (i) are equivalent (see Section 1F).

We first show (i) \Rightarrow (ii): Fix i and $H \in \mathbb{P}^*$ in $\partial_\infty A_i$. The opposite of H in $\partial_\infty A_i$ is some p_j . Then $H = H_{ij}$, so H is also the opposite of p_i in the apartment $\partial_\infty A_j$. As $x \in A_j$, we then have $\angle_x(p_i, H) = \pi$. We now prove (ii) \Rightarrow (iii): First recall that for $p \in \mathbb{P}$ and $H \in \mathbb{P}^*$, we have $\angle_x(p, H) = \pi$ if and only if $\Sigma_x p \notin \Sigma_x H$ in the projective space \mathbb{P}_x . So (ii) means that $\Sigma_x p_i \notin \Sigma_x H_{ij}$ for all $i \neq j$. Let U_i be the minimal linear subspace of the projective space \mathbb{P}_x containing $\Sigma_x p_0, \dots, \Sigma_x p_i$. Then, for $i \leq N - 1$, we have that $\Sigma_x p_i$ is not in U_{i-1} , else $\Sigma_x p_i$ would belong to $\Sigma_x H_{i, i+1}$. Hence $(\Sigma_x p_0, \dots, \Sigma_x p_i)$ is independent in \mathbb{P}_x by induction on i . Therefore $(\Sigma_x p_0, \dots, \Sigma_x p_{N-1})$ is a frame, and (iii) follows by permuting the p_i .

We now prove (ii) \Rightarrow (iv). Let $i \in \{0, \dots, N\}$. Let $v \in \Sigma_x A_i$. Let $C \subset A_i$ be a closed Weyl chamber with vertex x containing v . Let $H \in \mathbb{P}^*$ be the singular point of type $N - 1$ in $\partial_\infty C$. Then $\angle_x(p_i, H) = \pi$, hence $\angle_x(p_i, v) \geq \pi - d > \frac{\pi}{2}$, as the diameter d of $\partial \bar{C}$ is strictly less than $\pi/2$.

(iv) \Rightarrow (v) is clear. Assume now that (v) holds. For $j \neq i$ in $\{0, \dots, N\}$, as $\angle_x(p_i, H_{ij}) \geq \frac{\pi}{2}$, the direction $\Sigma_x p_i$ is not in a closed chamber of $\Sigma_x X$ containing $\Sigma_x H_{ij}$. Hence by type considerations we must have $\angle_x(p_i, H_{ij}) = \pi$. So (ii) holds.

So the equivalence of all assertions is proven. We now prove the uniqueness of x . Suppose that x' is another point of X with the same properties, and $x' \neq x$. We proved above that we have then $\angle_x(p_i, x') > \frac{\pi}{2}$ and $\angle_{x'}(p_i, x) > \frac{\pi}{2}$, which is impossible. \square

We now state some properties of centers of projective frames. Consider a projective frame $\mathcal{F} = (p_0, p_1, \dots, p_N)$ in \mathbb{P} , and let $x \in X$ be its center. Consider

the $N + 1$ associated flats $A_i = A(p_0, \dots, \hat{p}_i, \dots, p_N)$ in X . We first describe the intersections of the flats A_i with A_0 .

Proposition 2.5. *For $i = 1, \dots, N$, let S_i be the convex hull of the rays from x to the points $p_1, \dots, \hat{p}_i, \dots, p_N$ — in other words the sector on those points, with basepoint x . Let $H_i = p_1 \oplus \dots \oplus \hat{p}_i \oplus \dots \oplus p_N$ denote the point in $\partial_\infty A_0$ opposite to p_i . For $i \in \{1, \dots, N\}$, we have:*

- (i) *Let y be an interior point of S_i . Then $\Sigma_y p_0 = \Sigma_y p_i$.*
- (ii) *For $y \in A_0$, we have $y \in A_0 \cap A_i$ if and only if $\Sigma_y p_0$ is opposite to $\Sigma_y H_i$.*
- (iii) $A_0 \cap A_i = S_i$.

In particular, the intersections $A_0 \cap A_i$, $i = 1, \dots, N$, form a partition (i.e., a covering with disjoint interiors) of A_i .

Note that the sector S_i is the union of the Weyl chambers of the flat A_0 based at x and containing the singular ray to H_i .

Proof. The inclusion $S_i \subset A_0 \cap A_i$ is clear since $x \in A_0 \cap A_i$ and p_j is in both $\partial_\infty A_0$ and $\partial_\infty A_i$ for $j \neq i$ in $\{1, \dots, N\}$.

If y is an interior point of S_i , then in the local spherical building $\Sigma_y X$ at y , we have that $\Sigma_y p_0 \in \Sigma_y A_0$. Moreover, $y \in A_i$ as previously observed, so $\Sigma_y p_0$ is opposite to $\Sigma_y H_i$ (in $\Sigma_y A_i$). Hence $\Sigma_y p_0$ is equal to the opposite of $\Sigma_y H_i$ in $\Sigma_y A_0$, which is $\Sigma_y p_i$, proving (i).

We now prove (ii): In \mathbb{P}_y , the points $(\Sigma_y p_1, \dots, \Sigma_y p_N)$ form a frame (since $y \in A_0$). Hence the $N - 1$ points $(\Sigma_y p_1, \dots, \widehat{\Sigma_y p_i}, \dots, \Sigma_y p_N)$ are independent. Therefore $(\Sigma_y p_0, \dots, \widehat{\Sigma_y p_i}, \dots, \Sigma_y p_N)$ is a frame in \mathbb{P}_y (i.e., $y \in A_i$) if and only if $\Sigma_y p_0 \notin \Sigma_y H_i$.

We finish by proving the remaining inclusion $A_0 \cap A_i \subset S_i$: The S_i clearly form a partition of A_0 . So it is enough to prove that $A_0 \cap A_i$ does not meet the interior of S_j for $j \neq i$. Else, at such a point y , by (i), we would have $\Sigma_y p_0 = \Sigma_y p_j$, which is not opposite to $\Sigma_y H_i$, providing a contradiction. \square

The following proposition shows that the notion of center of projective frames behaves well with respect to projections to transverse spaces at infinity.

Proposition 2.6. *For each i , the projection of x in the transverse building at infinity X_{p_i} is the center of the projective frame of $\partial_\infty X_{p_i}$ formed by the projections $\text{proj}_{p_i}(p_j) = p_i p_j$ of the p_j , $j \neq i$, that is:*

$$\pi_{p_i}(c(p_0, p_1, \dots, p_N)) = c(p_i p_0, p_i p_1, \dots, \widehat{p_i p_i}, \dots, p_i p_N).$$

Proof. For all $j \neq i$, the ray from x to p_j is in the flat A_j hence its projection $\pi_{p_i}(x)$ in the transverse building X_{p_i} is in $\pi_{p_i}(A_j)$, which is the flat defined by the frame $\text{proj}_{p_i}(p_k) = p_i p_k$, $k \neq i, j$. \square

In the algebraic case, i.e., when X is the Euclidean building $X(V)$ associated with some vector space V of dimension N over an ultrametric field \mathbb{K} , we have the following characterization of the center as a norm on V .

Proposition 2.7. *Let $\mathcal{F} = (p_0, p_1, \dots, p_N)$ be a projective frame in $\mathbb{P} = \mathbb{P}(V)$. The center of \mathcal{F} is the norm η on V canonically associated to any basis $\mathbf{v} = (v_i)_{i=1, \dots, N}$ of V such that $p_i = [v_i]$ for $1 \leq i \leq N$ and $p_0 = [v_1 + \dots + v_N]$ in $\mathbb{P}(V)$, i.e., the norm defined by*

$$\eta \left(\sum_{i=1}^N a_i v_i \right) = \max_{1 \leq i \leq N} |a_i|.$$

Proof. Let $\mathbf{v} = (v_1, \dots, v_N)$ be a basis of V such that $p_0 = [v_1 + \dots + v_N]$ in $\mathbb{P}(V)$ and $p_i = [v_i]$. Let η be the associated canonical norm on V . We clearly have $\eta \in A_0$ by the definition of marked flats in the model of norms. Let g be the element of $\text{GL}(V)$ sending the basis \mathbf{v} to the basis $(v_1, \dots, v_{N-1}, v_1 + \dots + v_N)$. Then g preserves the norm η and sends A_0 to A_N and hence η is in the flat A_N . Permuting the basis \mathbf{v} , we similarly get that η is in the flat A_i for all $i \neq 0$. \square

Remark 2.8. By duality, the similar properties hold for generic $(N + 1)$ -tuples (projective frames) in $\mathbb{P}^* \subset \partial_\infty X$.

2C. Projecting two ideal points onto a flat. From now on we return to the case where $N = 3$ (type A_2).

Proposition 2.9. *Let (p_1, p_2, p_3) be a independent triple in \mathbb{P} . Let p, q be two points in \mathbb{P} , in generic position relatively to the p_i (i.e., not on any of the lines $p_i p_j$). Denote by x and y the respective orthogonal projections of p and q on the flat $A = A(p_1, p_2, p_3)$. Identify A with \mathbb{A} by a marked flat sending $\partial \mathcal{C}$ to $(p_1, p_1 p_2)$. Then the roots coordinates of \overrightarrow{xy} are given by the three natural cross ratios at the vertices of the triangle:*

$$\begin{aligned} \varphi_1(\overrightarrow{xy}) &= \boldsymbol{\beta}(p_3 p_1, p_3 p, p_3 p_2, p_3 q), \\ \varphi_2(\overrightarrow{xy}) &= \boldsymbol{\beta}(p_1 p_2, p_1 p, p_1 p_3, p_1 q), \\ \varphi_3(\overrightarrow{xy}) &= \boldsymbol{\beta}(p_2 p_3, p_2 p, p_2 p_1, p_2 q). \end{aligned}$$

The analogous dual result holds for projections of two lines of \mathbb{P} on a flat (exchanging the roles of points and lines in \mathbb{P}).

Proof. Projecting on the transverse tree X_{p_1} in direction p_1 , we have

$$\varphi_2(\overrightarrow{xy}) = \varphi_2(B_{(p_1, p_1 p_2)}(x, y)) = B_{p_1 p_2}(\pi_{p_1}(x), \pi_{p_1}(y))$$

by (1-1). Since the projections of x and y on the tree X_{p_1} are the respective centers of the ideal triples $(p_1 p_2, p_1 p_3, p_1 p)$ and $(p_1 p_2, p_1 p_3, p_1 q)$ (Proposition 2.6), we

have

$$B_{p_1 p_2}(\pi_{p_1}(x), \pi_{p_1}(y)) = \beta(p_1 p_2, p_1 p, p_1 p_3, p_1 q)$$

by (1-3), hence $\varphi_2(\vec{x}y) = \beta(p_1 p_2, p_1 p, p_1 p_3, p_1 q)$. The remaining assertions follow by applying cyclic permutation, since

$$\varphi_1(B_{(p_1, p_1 p_2)}(x, y)) = \varphi_2(B_{(p_3, p_3 p_1)}(x, y)),$$

$$\varphi_3(B_{(p_1, p_1 p_2)}(x, y)) = \varphi_2(B_{(p_2, p_2 p_3)}(x, y)). \quad \square$$

For the projections of a point and a line, we have the following result.

Proposition 2.10. *Let $F_- = (p_-, D_-)$ and $F_+ = (p_+, D_+)$ be two opposite flags in \mathbb{P} and A the flat in X joining them, identified with \mathbb{A} by a marked flat sending $\partial\mathcal{C}$ to F_+ . Let p be a point and D a line in \mathbb{P} in generic position with respect to F_- and F_+ , (i.e., p does not belong to any of the lines $p_- p_+$, D_- , D_+ , and D does not contain any of the points $D_- \cap D_+$, p_-, p_+).*

Denote by x and x^ the respective orthogonal projections of p and D on A . Then in simple roots coordinates we have*

$$\vec{x}x^* = (z_-, z_+),$$

with

$$z_- = \beta(p_+, D_+ \cap (p_- p), D_+ \cap D_-, D_+ \cap D)$$

$$= \beta(D_-, p_- \oplus (D_+ \cap D), p_- p_+, p_- p),$$

$$z_+ = \beta(p_-, D_- \cap D, D_- \cap D_+, D_- \cap (p_+ p))$$

$$= \beta(D_+, p_+ p, p_+ p_-, p_+ \oplus (D_- \cap D))$$

Proof. See Figure 5. The projection of x on the transverse tree X_{p_-} is the center of the ideal triple $(p_- p_+, p_-(D_- \cap D_+), p_- p)$, and the projection of x^* on the tree X_{D_+} is the center of the ideal triple $(p_+, D_+ \cap D_-, D_+ \cap D)$ (Proposition 2.6). As x lies on a geodesic from p_- to D_+ , we have

$$\pi_{D_+}(x) = \pi_{D_+, p_-}(\pi_{p_-}(x))$$

$$= \pi_{D_+, p_-}(c(p_- p_+, p_-(D_- \cap D_+), p_- p))$$

$$= c(p_+, D_- \cap D_+, D_+ \cap (p_- p)).$$

Then projecting on the transverse tree X_{D_+} we have

$$\varphi_1(\vec{x}x^*) = B_{p_+}(\pi_{D_+}(x), \pi_{D_+}(x^*)) = \beta(p_+, D_+ \cap (p_- p), D_+ \cap D_-, D_+ \cap D)$$

as needed. The remaining assertions have identical proofs. \square




Figure 5. Projecting a point and a line on a flat. The left part of the diagram represents the situation in X , and the right part the situation in X_{D_+} .

3. Triple ratio of a triple of ideal chambers

In this section, we introduce the (*geometric*) *triple ratio* of a nondegenerate triple of ideal chambers in a real Euclidean building X of type A_2 , establish its basic properties, and the links with the usual \mathbb{K} -valued (algebraic) triple ratio of triples of flags (see e.g., [Fock and Goncharov 2007]) in the algebraic case $\mathbb{P} = \mathbb{P}(\mathbb{K}^3)$.

We first give a precise definition of *nondegenerate* and *generic* triples of flags in an arbitrary projective plane \mathbb{P} .

3A. Nondegenerate and generic triples of flags. Let \mathbb{P} be a projective plane and $T = (F_1, F_2, F_3)$ be a triple of flags $F_i = (p_i, D_i)$ in \mathbb{P} . We will denote by p_{ij} the point $D_i \cap D_j$ (resp. D_{ij} the line $p_i p_j$), when defined.

The natural nondegeneracy condition on the triple (F_1, F_2, F_3) for the triple ratios to be well defined is the following:

$$\text{either } \forall i, p_i \notin D_{i+1} \text{ or } \forall i, p_i \notin D_{i-1}. \tag{ND}$$

This condition is clearly equivalent to: the points are pairwise distinct, the lines are pairwise distinct, none of the points is on the three lines (i.e., $D_i \cap D_j \neq p_k$ for all $\{i, j, k\} = \{1, 2, 3\}$) and none of the lines contains the three points (i.e., $p_i p_j \neq D_k$ for all i, j, k). We will then say that the triple (F_1, F_2, F_3) is *nondegenerate*.

It is easy to check that the triple T defines then a nondegenerate quadruple $(D_i, p_i p_j, p_i p_{jk}, p_i p_k)$ of lines through each point p_i , and a nondegenerate quadruple $(p_i, D_i \cap D_j, D_i \cap D_{jk}, D_i \cap D_k)$ of points on each line D_i .

The triple of flags $T = (F_1, F_2, F_3)$ is *generic* if the flags $F_i = (p_i, D_i)$ are pairwise opposite, the points $(p_i)_i$ are not collinear and the lines $(D_i)_i$ are not concurrent. In particular, T is then nondegenerate, and the induced quadruples of

points on each line (resp. of lines through each point) are generic (i.e., pairwise distinct).

3B. Algebraic triple ratio. When $\mathbb{P} = \mathbb{P}(\mathbb{K}^3)$ is the projective plane associated with an arbitrary field \mathbb{K} , the algebraic triple ratio of a nondegenerate triple of flags $T = (F_1, F_2, F_3)$ (see Section 3A), with values in $\mathbb{K} \cup \{\infty\}$, is defined by (see [Fock and Goncharov 2006, §9.4])

$$\text{Tri}(F_1, F_2, F_3) = \frac{\tilde{D}_1(\tilde{p}_2)\tilde{D}_2(\tilde{p}_3)\tilde{D}_3(\tilde{p}_1)}{\tilde{D}_1(\tilde{p}_3)\tilde{D}_2(\tilde{p}_1)\tilde{D}_3(\tilde{p}_2)},$$

where \tilde{p}_i is any vector in \mathbb{K}^3 representing p_i and \tilde{D}_i is any linear form in $(\mathbb{K}^3)^*$ representing D_i , and $F_i = (p_i, D_i)$. It is invariant under cyclic permutation of the flags and inverted by reversing the order

$$\text{Tri}(F_3, F_2, F_1) = \text{Tri}(F_1, F_2, F_3)^{-1}.$$

It may be expressed as the following cross ratio:

$$\text{Tri}(F_1, F_2, F_3) = \mathbf{b}(D_1, p_1 p_2, p_1 p_{23}, p_1 p_3). \quad (3-1)$$

3C. Geometric triple ratio. We suppose now that the projective plane \mathbb{P} is the projective plane at infinity of some a real Euclidean building X of type A_2 , possibly exotic. Let β be the associated geometric cross ratio on \mathbb{P} (see Section 1K). Let $T = (F_1, F_2, F_3)$ be a nondegenerate triple of ideal chambers of X , i.e., a nondegenerate triple of flags $F_i = (p_i, D_i)$ in \mathbb{P} .

The idea is to define the geometric triple ratio of T by analogy with the expression of the algebraic triple ratio as a cross ratio (3-1), replacing \mathbf{b} by β , in such a way that, in the algebraic case, the geometric triple ratio of a triple T with algebraic triple ratio Z should be $\log|Z|$. But for the purpose of geometric classification, this geometric cross ratio $\beta(D_1, p_1 p_2, p_1 p_{23}, p_1 p_3)$ alone will not retain enough information. In particular, in contrast to the algebraic cross ratio, it does not determine the geometric cross ratios obtained from the original 4-tuple by cyclic permutations of the three last arguments, which in the algebraic case are $\log|1 + Z^{-1}|$ and $-\log|1 + Z|$, see (1-9), and have geometric significance. For example, in the algebraic case, it will not distinguish between two triples T and T' with respective algebraic triple ratios $Z = -1$ and $Z' = -1 + a$ with $|a| < 1$.

In order to retain this information we define the *geometric triple ratio* of T as the triple

$$\text{tri}(T) = (\text{tri}_m(T))_{m=1,2,3}$$

where

$$\text{tri}_1(F_1, F_2, F_3) = \beta(D_1, p_1 p_2, p_1 p_{23}, p_1 p_3),$$

$$\text{tri}_2(F_1, F_2, F_3) = \beta(D_1, p_1 p_3, p_1 p_2, p_1 p_{23}),$$

$$\text{tri}_3(F_1, F_2, F_3) = \beta(D_1, p_1 p_{23}, p_1 p_3, p_1 p_2),$$

are the geometric cross ratios obtained from $(D_1, p_1 p_2, p_1 p_{23}, p_1 p_3)$ by cyclic permutation of the three last lines. Note these cross ratios are well defined, since the four lines $D_1, p_1 p_2, p_1 p_{23}, p_1 p_3$ are well defined and form a nondegenerate quadruple of lines through p_1 (see Section 3A above).

The next proposition gathers the properties of the geometric triple ratio, and show in particular that this invariant is in fact 1-dimensional, as it takes values in one of the three rays $\mathbb{R}_+(0, 1, -1)$, $\mathbb{R}_+(-1, 0, 1)$, and $\mathbb{R}_+(1, -1, 0)$.

Proposition 3.1. (i) *The geometric triple ratio is invariant by cyclic permutations of the flags; i.e., for $m = 1, 2, 3$,*

$$\text{tri}_m(F_2, F_3, F_1) = \text{tri}_m(F_1, F_2, F_3).$$

(ii) *Exchanging two flags, we have*

$$\text{tri}_1(F_1, F_3, F_2) = -\text{tri}_1(F_1, F_2, F_3), \quad \text{tri}_2(F_1, F_3, F_2) = -\text{tri}_3(F_1, F_2, F_3).$$

(iii) *We have $\text{tri}_1(T) + \text{tri}_2(T) + \text{tri}_3(T) = 0$.*

(iv) *For all $m \in \mathbb{Z}/3\mathbb{Z}$, if $\text{tri}_m(T) > 0$, then we have $\text{tri}_{m-1}(T) = 0$ and $\text{tri}_{m+1}(T) = -\text{tri}_m(T) < 0$.*

In order to prove this proposition, in particular, the invariance of the triple ratio by cyclic permutation of the flags, we first introduce the natural dual invariants given by the cross ratios of the natural induced quadruple of points on the line D_1 (that is, exchanging the role of points and lines):

$$\text{tri}_1^*(F_1, F_2, F_3) = \beta(p_1, D_2 \cap D_1, D_{23} \cap D_1, D_3 \cap D_1),$$

$$\text{tri}_2^*(F_1, F_2, F_3) = \beta(p_1, D_3 \cap D_1, D_2 \cap D_1, D_{23} \cap D_1),$$

$$\text{tri}_3^*(F_1, F_2, F_3) = \beta(p_1, D_{23} \cap D_1, D_3 \cap D_1, D_2 \cap D_1).$$

The following property is straightforward.

$$\begin{aligned} \text{tri}_1^*(F_1, F_3, F_2) &= -\text{tri}_1^*(F_1, F_2, F_3), \\ \text{tri}_2^*(F_1, F_3, F_2) &= -\text{tri}_3^*(F_1, F_2, F_3). \end{aligned} \tag{3-2}$$

We will need the following property showing that the invariants behave nicely under duality.

Lemma 3.2. *For $m = 1, 2, 3$, we have $\text{tri}_m^*(F_1, F_2, F_3) = \text{tri}_m(F_3, F_2, F_1)$.*

Proof of Lemma 3.2. By invariance under perspectivities and double transpositions, we have

$$\begin{aligned} \text{tri}_1^*(F_1, F_2, F_3) &= \beta(p_1, D_2 \cap D_1, D_{23} \cap D_1, D_3 \cap D_1) \\ &= \beta(p_1 p_3, p_{12} p_3, D_{23}, D_3) \\ &= \beta(D_3, p_2 p_3, p_{12} p_3, p_1 p_3) \\ &= \text{tri}_1(F_3, F_2, F_1). \end{aligned}$$

The proof of $\text{tri}_m^*(F_1, F_2, F_3) = \text{tri}_m(F_3, F_2, F_1)$ for $m = 2, 3$ is similar. \square

We now turn to the proof of Proposition 3.1.

Proof of Proposition 3.1. Assertions (iii) and (iv) follow immediately from the properties of the cross ratio β under cyclic permutation of the three last points (see (1-4) and (1-5)).

Assertion (ii) follows immediately from the definition and from the symmetries of the cross ratio.

We finally prove Proposition 3.1(i). Using (ii), Lemma 3.2 and (3-2), we have

$$\begin{aligned} \text{tri}_1(F_2, F_3, F_1) &= -\text{tri}_1(F_2, F_1, F_3) \\ &= -\text{tri}_1^*(F_3, F_1, F_2) = \text{tri}_1^*(F_3, F_2, F_1) = \text{tri}_1(F_1, F_2, F_3), \\ \text{tri}_2(F_2, F_3, F_1) &= -\text{tri}_3(F_2, F_1, F_3) \\ &= -\text{tri}_3^*(F_3, F_1, F_2) = \text{tri}_2^*(F_3, F_2, F_1) = \text{tri}_2(F_1, F_2, F_3). \end{aligned}$$

The case where $m = 3$ is similar to the case $m = 2$. \square

3D. Geometric triple ratio from algebraic triple ratio. When \mathbb{P} is the projective plane on some field \mathbb{K} endowed with some ultrametric absolute value, and $\beta = \log|\mathbf{b}|$ where \mathbf{b} is the usual \mathbb{K} -valued cross ratio on \mathbb{P} , the three geometric triple ratios $\text{tri}_m(T)$, $m = 1, 2, 3$ of T are obtained from the single algebraic triple ratio $Z = \text{Tri}(T)$ of T by the relations

$$\begin{aligned} \text{tri}_1(T) &= \log|Z|, \\ \text{tri}_2(T) &= \log\left|\frac{1}{1+Z}\right| = -\log|1+Z|, \\ \text{tri}_3(T) &= \log|1+Z^{-1}|, \end{aligned} \tag{3-3}$$

which are easily derived from the expression of algebraic triple ratio as a cross ratio (3-1) and from the symmetry properties of the algebraic cross ratio (1-9).

Remark 3.3. Note that the geometric invariants do not determine the triple of flags up to automorphisms of \mathbb{P} (unlike the usual (algebraic) triple ratio): for example in the algebraic case $\mathbb{P} = \mathbb{P}(\mathbb{K}^3)$, take T with triple ratio $Z \in \mathbb{K}$ with $|Z| > 1$ and T' with triple ratio $Z' = Za$ where $a \in \mathbb{K}$ with $|a| = 1$ and $a \neq 1$. Then T and T' are

not in the same $\mathrm{PGL}(\mathbb{K}^3)$ -orbit, but have the same three geometric invariants, as $\mathrm{tri}_1(T) = \log|Z| = \mathrm{tri}_1(T')$, $\mathrm{tri}_2(T) = -\log|Z| = \mathrm{tri}_2(T')$, $\mathrm{tri}_3(T) = 0 = \mathrm{tri}_3(T')$.

4. Proof of the main result

In this section we prove Theorems 0.1 and 0.2. Let X be a Euclidean building of type A_2 and $T = (F_1, F_2, F_3)$ be a generic triple of flags in the projective plane \mathbb{P} at infinity of X . We denote by $z_m = \mathrm{tri}_m(F_1, F_2, F_3)$, $m = 1, 2, 3$, its geometric triple ratio, and by $A_{ij} = A(F_i, F_j)$, $A_p = A(p_1, p_2, p_3)$ and $A_D = A(D_1, D_2, D_3)$ the five associated flats.

We first define the six associated points in X .

4A. Associated points in the building. For $\{i, j, k\} = \{1, 2, 3\}$, denote by y_k the center in X of the projective frame (p_1, p_2, p_3, p_{ij}) , where $p_{ij} = D_i \cap D_j$, and by y_k^* the center of the projective frame (D_1, D_2, D_3, D_{ij}) , where $D_{ij} = p_i p_j$, as defined in Proposition 2.4. In particular the point y_k is the orthogonal projection of p_{ij} on A_p , the point y_k^* is the orthogonal projection of D_{ij} on A_D , the point y_k is the orthogonal projection of p_k on $A_{ij} = A(p_i, p_j, p_{ij})$, and the point y_k^* is the orthogonal projection of D_k on $A_{ij} = A(D_i, D_j, D_{ij})$.

4B. In the flat A_{ij} . We now link the respective position of the points y_k and y_k^* in the flat A_{ij} to the geometric triple ratio of T . Suppose that the indices i, j, k respect the cyclic order, i.e., that $(i, j, k) = (123)$ as cyclic permutations. We identify A_{ij} with the model flat \mathbb{A} by a marked flat $f_{ij} : \mathbb{A} \rightarrow A_{ij}$ sending $\partial\mathfrak{C}$ to F_j . For x, y in $A_{ij} \simeq \mathbb{A}$, we define then $\overrightarrow{xy} = y - x = B_{F_j}(x, y)$. Recall that $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$ denotes the canonical basis of \mathbb{R}^3 . In particular, the directions of p_i, p_{ij} and p_j are respectively identified with the directions of $[\varepsilon_1], [\varepsilon_2]$, and $[\varepsilon_3]$ in \mathbb{A} .

Proposition 4.1. (i) *In simple roots coordinates, we have $\overrightarrow{y_k^* y_k} = (z_2, z_3)$.*

(ii) *For $m = 1, 2, 3$, if $z_m > 0$ then $\overrightarrow{y_k y_k^*} = z_m[\varepsilon_m]$. In particular y_k^* is on one of the three singular rays of type 1 issued from y_k (i.e the rays to p_i, p_j and p_{ij}).*

Proof. As y_k and y_k^* are the respective orthogonal projections on the flat A_{ij} of p_k and D_k , by Proposition 2.10 and cyclic invariance of the geometric triple ratio, we have

$$\begin{aligned} \overrightarrow{y_k^* y_k} &= \boldsymbol{\beta}(D_i, p_i p_k, p_i p_j, p_i p_{jk}) = \mathrm{tri}_2(F_i, F_j, F_k) = z_2 \quad \text{and} \\ \overrightarrow{y_k y_k^*} &= \boldsymbol{\beta}(D_j, p_j p_{ki}, p_j p_i, p_j p_k) = \mathrm{tri}_3(F_j, F_k, F_i) = z_3. \end{aligned}$$

Assertion (ii) follows, since we have then $z_{m-1} = 0$ and $z_{m+1} = -z_m$ by ultrametricity of the geometric triple ratio (Proposition 3.1(iv)). \square

We now describe the intersections of A_{ij} with the four other flats (see Figures 1 and 2 in the introduction). These intersections happen to be sectors in \mathbb{A} bounded

by two singular rays of same type, equivalently the union of two adjacent Weyl chambers.

Proposition 4.2. *Let $x \in A_{ij}$. Then:*

- (i) *The intersection $A_{ij} \cap A_p$ is the sector at y_k bounded by the rays to p_i and p_j . That is,*

$$x \in A_p \text{ if and only if } \begin{cases} \varphi_1(x) \geq \varphi_1(y_k), \\ \varphi_2(x) \leq \varphi_2(y_k). \end{cases}$$

- (ii) *The intersection $A_{ij} \cap A_D$ is the sector at y_k^* bounded by the rays to D_i and D_j . That is,*

$$x \in A_D \text{ if and only if } \begin{cases} \varphi_1(x) \leq \varphi_1(y_k^*), \\ \varphi_2(x) \geq \varphi_2(y_k^*). \end{cases}$$

- (iii) *The intersection $A_{ij} \cap A_{jk}$ is the intersection of the sector at y_k bounded by the rays to p_j and $D_i \cap D_j$, and the sector at y_k^* bounded by the rays to D_j and $p_i p_j$. That is,*

$$x \in A_{jk} \text{ if and only if } \begin{cases} \varphi_1(x) \geq \varphi_1(y_k^*), \\ \varphi_2(x) \geq \varphi_2(y_k), \\ \varphi_3(x) \leq \min(\varphi_3(y_k), \varphi_3(y_k^*)). \end{cases}$$

- (iv) *The intersection $A_{ij} \cap A_{ki}$ is the intersection of the sector at y_k bounded by the rays to p_i and $D_i \cap D_j$, and the sector at y_k^* bounded by the rays to D_i and $p_i p_j$. That is,*

$$x \in A_{ki} \text{ if and only if } \begin{cases} \varphi_1(x) \leq \varphi_1(y_k), \\ \varphi_2(x) \leq \varphi_2(y_k^*), \\ \varphi_3(x) \geq \max(\varphi_3(y_k), \varphi_3(y_k^*)). \end{cases}$$

Proof. Since y_k is the center of the projective frame (p_i, p_j, p_{ij}, p_k) , assertion (i) comes from Proposition 2.5, as $A_{ij} = A(p_i, p_j, p_{ij})$ and $A_p = A(p_i, p_j, p_k)$. Assertion (ii) is similar. Assertion (iii): A point $x \in A_{ij}$ lies in A_{jk} if and only if, in the spherical building of directions at $\Sigma_x X$, the direction $\Sigma_x D_j$ is opposite to $\Sigma_x p_k$ and $\Sigma_x p_j$ is opposite to $\Sigma_x D_k$. Moreover, $\Sigma_x D_j$ is opposite to $\Sigma_x p_k$ if and only if $x \in A(p_k, p_j, p_{ij})$. As y_k is the center of the projective frame (p_i, p_j, p_{ij}, p_k) and $A_{ij} = A(p_i, p_j, p_{ij})$, the set of such x is the sector at y_k bounded by the rays to p_j and $D_i \cap D_j$ (by Proposition 2.5). This is the subset of $x \in A_{ij}$ satisfying: $\varphi_2(x) \geq \varphi_2(y_k)$ and $\varphi_3(x) \leq \varphi_3(y_k)$. Similarly, as y_k^* is the center of the projective frame (D_i, D_j, D_{ij}, D_k) and $A_{ij} = A(D_i, D_j, D_{ij})$, the direction $\Sigma_x p_j$ is opposite to $\Sigma_x D_k$ if and only if x is in the sector at y_k^* bounded by the rays to D_j and $D_{ij} = p_i p_j$. That is, if and only if $\varphi_1(x) \geq \varphi_1(y_k^*)$ and $\varphi_3(x) \leq \varphi_3(y_k^*)$, and we are done. Assertion (iv) is similar. \square

In particular, as y_k^* is on one of the three singular rays of type 1 issued from y_k by Propositions 4.1, from Proposition 4.2 we easily get the following result.

Corollary 4.3. *The intersections with A_{ij} of A_{jk}, A_{ki}, A_p and A_D form a partition of A_{ij} . \square*

4C. In the flat A_p . We now consider the flat $A_p = A(p_1, p_2, p_3)$. The following proposition describes the respective positions in A_p of the points y_1, y_2, y_3 . We identify A_p with \mathbb{A} by a marked flat $f_p : \mathbb{A} \rightarrow A_p$ sending $\partial\mathcal{C}$ to $(p_1, p_1 p_2)$ (hence direction $[\varepsilon_i]$ to p_i for $i = 1, 2, 3$). Recall that we then have $\overrightarrow{xx'} = x' - x = B_{(p_1, p_1 p_2)}(x, x')$ for $x, x' \in A_p$.

Proposition 4.4. *In the flat A_p we have:*

- (i) *In simple roots coordinates, we have $\overrightarrow{y_2 y_3} = (z_1, 0)$.*
- (ii) *If $z_1 \geq 0$, the point y_{i+1} is in the ray $[y_i, p_{i+2})$ (for all i), and if $z_1 \leq 0$, the point y_i is in the ray $[y_{i+1}, p_{i+2})$ for all i .*

In particular the triangle $\Delta \subset A_p$ with vertices y_1, y_2, y_3 is singular, i.e., the sides have singular type in $\bar{\mathcal{C}}$.

Proof. Recall that the point y_k is the orthogonal projection on the flat A_p of the singular boundary point $p_{ij} = D_i \cap D_j$. Then, by Proposition 2.6 the points y_2 and y_3 have the same projection in the transverse tree X_{p_1} , that is the center of the ideal triple $(p_1 p_{13}, p_1 p_2, p_1 p_3) = (D_1, p_1 p_2, p_1 p_3) = (p_1 p_{23}, p_1 p_2, p_1 p_3)$, proving that $\varphi_2(\overrightarrow{y_2 y_3}) = 0$. Furthermore, by Proposition 2.9 we have

$$\begin{aligned} \varphi_2(\overrightarrow{y_3 y_1}) &= \beta(p_1 p_2, p_1 p_{12}, p_1 p_3, p_1 p_{23}) \\ &= \beta(p_1 p_2, D_1, p_1 p_3, p_1 p_{23}) \\ &= \beta(D_1, p_1 p_2, p_1 p_{23}, p_1 p_3) \\ &= z_1, \end{aligned}$$

proving that $\varphi_2(\overrightarrow{y_3 y_1}) = z_1$. Applying this to the permuted triple (F_3, F_1, F_2) , we obtain $\varphi_1(\overrightarrow{y_2 y_3}) = z_1$ (by invariance of the geometric triple ratio z_1 by cyclic permutation). Assertion (ii) follows from (i), applying cyclic permutations. \square

We now describe the intersections of A_p with the other flats; see Figure 6.

Proposition 4.5. *Let $S_i = A_p \cap A_{i, i+1}$ and let Δ be the triangle with vertices y_1, y_2, y_3 . Then:*

- (i) *S_i is the sector of A_p bounded by the rays from y_{i+2} to p_i and p_{i+1} .*
- (ii) *S_1, S_2, S_3 and Δ form a partition of A_p .*

Proof. Assertion (i) follows from Proposition 4.2(i). In the case where $z_1 \geq 0$, assertion (ii) then comes from the fact that for all i , y_{i+1} is in the ray $[y_i, p_{i+2})$ (Proposition 4.4). The case where $z_1 \leq 0$ is similar. \square




Figure 6. Situation in the flat A_p : when $z_1 \geq 0$ (left) and when $z_1 \leq 0$ (right).




Figure 7. Situation in the flat A_D : when $z_1 \geq 0$ (left) and when $z_1 \leq 0$ (right).

4D. In the flat A_D . We now state the similar properties in the dual flat $A_D = A(D_1, D_2, D_3)$, which have same proofs, exchanging the role of points and lines.

Proposition 4.6. *In the flat A_D identified with \mathbb{A} by a marked flat sending $\partial\mathcal{C}$ to $(D_1 \cap D_2, D_1)$, we have:*

- (i) $\overrightarrow{y_2^* y_3^*} = (0, -z_1)$ in simple roots coordinates. In particular y_2^* and y_3^* are on a common singular geodesic to D_1 .
- (ii) The points y_1^*, y_2^*, y_3^* form a singular triangle Δ^* in A_D .
- (iii) For all $i \in \mathbb{Z}/3\mathbb{Z}$, $S_i^* = A_D \cap A_{i,i+1}$ is the sector of A_D bounded by the rays from y_{i+2}^* to D_i and D_{i+1} .
- (iv) S_1^*, S_2^*, S_3^* and Δ^* form a partition of A_D . □

4E. The classification. We now combine the previous results to establish the classification in two geometric types, finishing to prove Theorems 0.1 and 0.2.

Proof of Theorem 0.1. Let $x = y_3$ and $x^* = y_3^*$. We identify the flat A_{12} with the model flat \mathbb{A} by a marked flat sending $\partial\mathcal{C}$ to F_2 , and 0 to y_3^* . By Proposition 4.2

applied to the flat A_{12} , we have $\varphi_1(y_3) = z_2$, $\varphi_2(y_3) = z_3$, and $\varphi_3(y_3) = z_1$. By Proposition 4.2 applied to the flat A_{12} , the intersection $I = A_{12} \cap A_{23} \cap A_{31}$ is the subset of $y \in A_{12}$ such that

$$\begin{aligned} 0 &\leq \varphi_1(y) \leq \varphi_1(y_3) = z_2, \\ 0 &\geq \varphi_2(y) \geq \varphi_2(y_3) = z_3, \\ \max(\varphi_3(y_3), 0) &\leq \varphi_3(y) \leq \min(\varphi_3(y_3), 0). \end{aligned}$$

In particular, if I is not empty, then $z_1 = \varphi_3(y_3) = 0$.

Suppose from now on that $z_1 = 0$. Then $z_2 \geq 0$ and $z_3 = -z_2$ by the ultrametricity of the geometric triple ratio (Proposition 3.1(iv)). By the description above, I is then the subset of the line $\varphi_3 = 0$ (which contains $y_3^* = 0$ and y_3) consisting of the y such that $0 \leq \varphi_1(y) \leq \varphi_1(y_3)$ (since $\varphi_2(y) = -\varphi_1(y)$ when $\varphi_3(y) = 0$). Hence I is not empty and is the segment from $0 = y_3^*$ to y_3 i.e., $[x, x^*]$. Furthermore, as $z_1 = 0$, Proposition 4.4 implies that $y_1 = y_2 = y_3$. Similarly, we have $y_1^* = y_2^* = y_3^*$ by Proposition 4.6. Suppose now $x \neq x^*$. Since the segment $[x, x^*]$ lies in the ray $[x, p_{ij})$, and $x = y_k$ is the orthogonal projection of p_{ij} on A_p , we have $\angle_x(x^*, D) = \pi$ for all lines D in $\partial_\infty A_p$ (Proposition 2.4). Therefore we have $\angle_x(x^*, y) \geq \frac{2\pi}{3}$ for all $y \neq x$ in A_p . Similarly, we have that $\angle_{x^*}(x, y) \geq \frac{2\pi}{3}$ for all $y \neq x$ in A_p . Hence $[x, x^*]$ is the unique segment of minimal length joining A_p to A_D . Assertion (iv) follows from Proposition 4.1. \square

Proof of Theorem 0.2. If $z_2 > 0$, then $z_1 = 0$ by the ultrametricity of the geometric triple ratio (Proposition 3.1(iv)), and $A_p \cap A_D$ is empty by Theorem 0.1. Suppose now that $z_2 \leq 0$. Since the case $z_1 \leq 0$ reduces to the case $z_1 \geq 0$ by exchanging F_2 and F_3 , it is enough to handle the case $z_1 \geq 0$. Then $z_3 = 0$ and $z_2 = -z_1$. Let $x_i = y_{i+2}$ for $i \in \mathbb{Z}/3\mathbb{Z}$. In A_{ij} identified with \mathbb{A} in such a way that $y_k^* = 0$, by Proposition 4.1 we have $\varphi_1(y_k) = z_2 = -z_1 \leq 0$, $\varphi_2(y_k) = z_3 = 0$, hence $\varphi_3(y_k) = z_1 \geq 0$. By Proposition 4.2(iv), $A_{ij} \cap A_{ik}$ is the set of $x \in A_{ij} \simeq \mathbb{A}$ such that $\varphi_1(x) \leq \varphi_1(y_k)$, $\varphi_2(x) \leq 0 = \varphi_2(y_k)$ and $\varphi_3(x) \geq \max(\varphi_3(y_k), 0) = \varphi_3(y_k)$. This is the Weyl chamber $y_k - \bar{\mathcal{C}}$, i.e., the Weyl chamber from $y_k = x_i$ to F_i . Similarly, $A_{ij} \cap A_{jk}$ is the Weyl chamber from y_k^* to F_j . Applying a cyclic permutation (ijk) , i.e., working in the flat A_{jk} , we also similarly get that $A_{ij} \cap A_{jk}$ is the Weyl chamber from y_i to F_j . Therefore $y_k^* = y_i$.

By Proposition 4.2 $A_p \cap A_D \cap A_{ij}$ is the intersection of the sector at y_k^* bounded by the rays to D_i and D_j , with the sector at y_k bounded by the rays to p_i and p_j . As the point y_k is on the ray from y_k to D_i , this is equal to the segment $[y_k, y_k^*]$. In particular $A_p \cap A_D$ contains y_k . Then $A_p \cap A_D$ contains y_1, y_2 and y_3 , hence the triangle Δ with vertices y_1, y_2 and y_3 , and since $A_p \cap A_D \cap A_{ij} = [y_k, y_i] \subset \Delta$, Proposition 4.5(ii) provides the reverse inclusion. Assertion (iii) comes from Proposition 4.1.

We finally prove (iv). Let $(i, j, k) = (123)$. Looking in the flat A_p , we see that the singular triangle Δ is contained in the Weyl chamber of X with tip x_i and that at x_i , we have $\Sigma_{x_i} x_j = \Sigma_{x_i} p_j$. Looking in the flat A_D we get $\Sigma_{x_i} x_k = \Sigma_{x_i} D_k$. Hence $\Sigma_{x_i} \Delta = (\Sigma_{x_i} p_j, \Sigma_{x_i} D_k)$. Since x_i belongs to the flats $A(F_i, F_j)$ and $A(F_i, F_k)$, we have that $\Sigma_{x_i} p_j$ is opposite to $\Sigma_{x_i} D_i$ and that $\Sigma_{x_i} D_k$ is opposite to $\Sigma_{x_i} p_i$. Therefore the Weyl chambers $\Sigma_{x_i} \Delta$ and $\Sigma_{x_i} F_i$ are opposite. It implies that Δ and the Weyl chamber from x_i to F_i are contained in a common flat of X by basic properties of real Euclidean buildings (see property (CO) of [Parreau 2000]). \square

In the algebraic case the following remark provides an alternative proof of some of the assertions of Theorem 0.2.

Remark 4.7. Let \tilde{p}_i in $V = \mathbb{K}^3$ be a vector representing p_i and \tilde{D}_i in V^* be a linear form representing D_i . Let $\mathbf{v} = (v_1, v_2, v_3)$ be the basis of V dual to the basis $(\tilde{D}_1, \tilde{D}_2, \tilde{D}_3)$ of V^* . Then in the projective plane $[v_i] = D_j \cap D_k$. We may suppose that $\tilde{p}_1 = (0, 1, 1)$, $\tilde{p}_2 = (Z, 0, 1)$, $\tilde{p}_3 = (1, 1, 0)$ in the basis \mathbf{v} , with $Z = \text{Tri}(F_1, F_2, F_3)$. Then the element $g \in \text{GL}(V)$ whose matrix in the basis \mathbf{v} is

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1/Z & 0 & 1 \end{pmatrix}$$

sends $[v_i]$ to p_{i+1} , hence A_D to A_p . If $|1 + Z| \geq 1$ and $z = \log|Z| \geq 0$, then the fixed point set of g in A_D is the image by the marked flat $f_{\mathbf{v}}$ of the singular triangle $\{\lambda \in \bar{\mathcal{C}} \mid \lambda_1 - \lambda_3 \leq \log|Z|\}$ (that is, Δ).

4F. Complements. We add here for future use a simple description of the vertices x_i, x_j, x_k of the singular triangle Δ in Theorem 0.2 by the projections on transverse trees at infinity.

Lemma 4.8. *We keep the hypotheses and notation of Theorem 0.2.*

- (i) *The projection $\pi_{p_i}(x_i)$ of x_i on the tree X_{p_i} is the center of the ideal tripod $(D_i, p_i p_j, p_i p_k)$.*
- (ii) *The projection $\pi_{D_i}(x_i)$ of x_i on the tree X_{D_i} is the center of the ideal tripod $(p_i, D_i \cap D_j, D_i \cap D_k)$.*
- (iii) *The projection $\pi_{p_i}(x_j)$ is the center of the ideal tripod $(D_i, p_i p_j, p_i p_{jk})$.*
- (iv) *The projection $\pi_{D_i}(x_j)$ is the center of the ideal tripod $(p_i, D_i \cap D_j, D_i \cap D_{jk})$.*

Proof. As the point x_i belongs to the three flats $A(F_k, F_i)$ and $A(F_j, F_i)$ and $A(p_i, p_j, p_k)$, its projection in the tree X_{p_i} belongs to the projection of $A(F_j, F_i)$, which is the line from D_i to $p_i p_j$, to the projection of $A(F_k, F_i)$, which is the line from D_i to $p_i p_k$, and to the projection of $A(p_i, p_j, p_k)$, which is the line from $p_i p_j$ to $p_i p_k$. Hence (i) is proven. Assertion (ii) is proven in the same way.

We now prove (iii). By (ii) applied to x_j , we have that $\pi_{D_j}(x_j)$ is the center of the ideal tripod p_j , $p_{jk} = D_j \cap D_k$, $D_j \cap D_i$. As x_j is on a geodesic from D_j to p_i , we may deduce that $\pi_{p_i}(x_j)$ is the center of the ideal tripod $p_i p_j$, $p_i p_{jk}$, D_i (using the canonical isomorphism $X_{D_j} \xrightarrow{\sim} X_{p_i}$). The last assertion (iv) has identical proof. \square

Acknowledgments

I thank Frédéric Paulin for useful discussions and comments, and the members of the Institut Fourier for their support.

References

- [Balsler 2008] A. Balsler, “Convex rank 1 subsets of Euclidean buildings (of type A_2)”, *Geom. Dedicata* **131** (2008), 123–158. MR Zbl
- [Bourdon 1996] M. Bourdon, “Sur le birapport au bord des CAT(−1)-espaces”, *Inst. Hautes Études Sci. Publ. Math.* 83 (1996), 95–104. MR Zbl
- [Bridson and Haefliger 1999] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] **319**, Springer, 1999. MR Zbl
- [Bruhat and Tits 1984] F. Bruhat and J. Tits, “Schémas en groupes et immeubles des groupes classiques sur un corps local”, *Bull. Soc. Math. France* **112**:2 (1984), 259–301. MR Zbl
- [Burger et al. 2010] M. Burger, A. Iozzi, and A. Wienhard, “Surface group representations with maximal Toledo invariant”, *Ann. of Math. (2)* **172**:1 (2010), 517–566. MR Zbl
- [Clerc and Neeb 2006] J.-L. Clerc and K.-H. Neeb, “Orbits of triples in the Shilov boundary of a bounded symmetric domain”, *Transform. Groups* **11**:3 (2006), 387–426. MR Zbl
- [Fock and Goncharov 2006] V. Fock and A. Goncharov, “Moduli spaces of local systems and higher Teichmüller theory”, *Publ. Math. Inst. Hautes Études Sci.* 103 (2006), 1–211. MR Zbl
- [Fock and Goncharov 2007] V. Fock and A. B. Goncharov, “Moduli spaces of convex projective structures on surfaces”, *Adv. Math.* **208**:1 (2007), 249–273. MR Zbl
- [Goldman 1980] W. M. Goldman, *Discontinuous groups and the Euler class*, Ph.D. thesis, University of California, Berkeley, 1980, Available at <https://search.proquest.com/docview/302998874>. MR
- [Goldman 1999] W. M. Goldman, *Complex hyperbolic geometry*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1999. Oxford Science Publications. MR Zbl
- [Goldman and Iwahori 1963] O. Goldman and N. Iwahori, “The space of p -adic norms”, *Acta Math.* **109** (1963), 137–177. MR Zbl
- [Kleiner and Leeb 1997] B. Kleiner and B. Leeb, “Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings”, *Inst. Hautes Études Sci. Publ. Math.* 86 (1997), 115–197. MR Zbl
- [Leeb 2000] B. Leeb, *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, Bonner Mathematische Schriften [Bonn Mathematical Publications] **326**, Universität Bonn, Mathematisches Institut, 2000. MR Zbl
- [Mühlherr et al. 2014] B. Mühlherr, K. Struyve, and H. Van Maldeghem, “Descent of affine buildings—I, Large minimal angles”, *Trans. Amer. Math. Soc.* **366**:8 (2014), 4345–4366. MR Zbl

- [Otal 1992] J.-P. Otal, “Sur la géométrie symplectique de l’espace des géodésiques d’une variété à courbure négative”, *Rev. Mat. Iberoamericana* **8**:3 (1992), 441–456. MR Zbl
- [Parreau 2000] A. Parreau, “Immeubles affines: construction par les normes et étude des isométries”, pp. 263–302 in *Crystallographic groups and their generalizations* (Kortrijk, 1999), edited by P. Igodt et al., Contemp. Math. **262**, Amer. Math. Soc., Providence, RI, 2000. MR Zbl
- [Parreau 2011] A. Parreau, “Espaces de représentations complètement réductibles”, *J. Lond. Math. Soc.* (2) **83**:3 (2011), 545–562. MR Zbl
- [Parreau 2012] A. Parreau, “Compactification d’espaces de représentations de groupes de type fini”, *Math. Z.* **272**:1-2 (2012), 51–86. MR Zbl
- [Parreau 2015] A. Parreau, “Invariant subspaces for some surface groups acting on A_2 -euclidean buildings”, preprint, 2015. arXiv
- [Rousseau 2009] G. Rousseau, “Euclidean buildings”, pp. 77–116 in *Géométries à courbure négative ou nulle, groupes discrets et rigidités*, edited by L. Bessières et al., Sémin. Congr. **18**, Soc. Math. France, Paris, 2009. MR Zbl
- [Talbi 2006] M. Talbi, “A Haagerup inequality, deformation of triangles and affine buildings”, *J. Inst. Math. Jussieu* **5**:2 (2006), 187–227. MR Zbl
- [Tits 1974] J. Tits, *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Mathematics **386**, Springer, Berlin, 1974. MR Zbl
- [Tits 1986] J. Tits, “Immeubles de type affine”, pp. 159–190 in *Buildings and the geometry of diagrams* (Como, 1984), edited by L. A. Rosati, Lecture Notes in Math. **1181**, Springer, 1986. MR Zbl
- [Toledo 1989] D. Toledo, “Representations of surface groups in complex hyperbolic space”, *J. Differential Geom.* **29**:1 (1989), 125–133. MR Zbl

Received 21 Jul 2016. Revised 10 Dec 2018.

ANNE PARREAU:

anne.parreau@univ-grenoble-alpes.fr

Université Grenoble Alpes, CNRS, IF, Grenoble, France

Innovations in Incidence Geometry

msp.org/iig

MANAGING EDITOR

Tom De Medts Ghent University
tom.demedts@ugent.be

Linus Kramer Universität Münster
linus.kramer@wwu.de

Klaus Metsch Justus-Liebig Universität Gießen
klaus.metsch@math.uni-giessen.de

Bernhard Mühlherr Justus-Liebig Universität Gießen
bernhard.m.muehlherr@math.uni-giessen.de

Joseph A. Thas Ghent University
thas.joseph@gmail.com

Koen Thas Ghent University
koen.thas@gmail.com

Hendrik Van Maldeghem Ghent University
hendrik.vanmaldeghem@ugent.be

HONORARY EDITORS

Jacques Tits
Ernest E. Shult †

EDITORS

Peter Abramenko University of Virginia

Francis Buekenhout Université Libre de Bruxelles

Philippe Cara Vrije Universiteit Brussel

Antonio Cossidente Università della Basilicata

Hans Cuypers Eindhoven University of Technology

Bart De Bruyn University of Ghent

Alice Devillers University of Western Australia

Massimo Giulietti Università degli Studi di Perugia

James Hirschfeld University of Sussex

Dimitri Leemans Université Libre de Bruxelles

Oliver Lorscheid Instituto Nacional de Matemática Pura e Aplicada (IMPA)

Guglielmo Lunardon Università di Napoli “Federico II”

Alessandro Montinaro Università di Salento

James Parkinson University of Sydney

Antonio Pasini Università di Siena (emeritus)

Valentina Pepe Università di Roma “La Sapienza”

Bertrand Rémy École Polytechnique

Tamás Szonyi ELTE Eötvös Loránd University, Budapest

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

See inside back cover or msp.org/iig for submission instructions.

The subscription price for 2019 is US \$275/year for the electronic version, and \$325/year (+\$15, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Innovations in Incidence Geometry: Algebraic, Topological and Combinatorial (ISSN 2640-7345 electronic, 2640-7337 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

IIG peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>
© 2019 Mathematical Sciences Publishers

Innovation in Incidence Geometry

Vol. 17 No. 2

2019

- Regular pseudo-hyperovals and regular pseudo-ovals in even characteristic 77
JOSEPH A. THAS
- Conics in Baer subplanes 85
SUSAN G. BARWICK, WEN-AI JACKSON and PETER WILD
- On triples of ideal chambers in A_2 -buildings 109
ANNE PARREAU
- Opposition diagrams for automorphisms of small spherical buildings 141
JAMES PARKINSON and HENDRIK VAN MALDEGHEM

