

# involve

a journal of mathematics

Numerical semigroups from open intervals

Vadim Ponomarenko and Ryan Rosenbaum

 mathematical sciences publishers

2010

vol. 3, no. 3

# Numerical semigroups from open intervals

Vadim Ponomarenko and Ryan Rosenbaum

(Communicated by Scott Chapman)

We consider numerical semigroups  $\mathbb{N} \cap \mathbb{N}I$ , for intervals  $I$ . We compute the Frobenius number and multiplicity of such semigroups, and show that we may freely restrict  $I$  to be open, closed, or half-open, as we prefer.

Given an interval  $I \subseteq \mathbb{Q}^{>0}$  in the positive rationals, consider the set

$$S(I) = \mathbb{N} \cap \mathbb{N}I = \{m \in \mathbb{N} : \exists n \in \mathbb{N}, m/n \in I\}.$$

This turns out to be a numerical semigroup, and has been the subject of considerable recent investigation (see [Rosales and García-Sánchez 2009, Chapter 4] for an introduction). Special cases include modular numerical semigroups [Rosales et al. 2005], where  $I = [m/n, m/(n-1)]$  ( $m, n \in \mathbb{N}$ ); proportionally modular numerical semigroups [Rosales et al. 2003], where  $I = [m/n, m/(n-s)]$  ( $m, n, s \in \mathbb{N}$ ); and opened modular numerical semigroups [Rosales and Urbano-Blanco 2006] where  $I = (m/n, m/(n-1))$  ( $m, n \in \mathbb{N}$ ).

We consider instead arbitrary open intervals  $I = (a, b)$ . We show that this set of semigroups coincides with the set of semigroups generated by closed and half-open intervals. Consequently, this class of semigroups contains modular numerical semigroups, proportionally modular numerical semigroups, as well as opened modular numerical semigroups. We also compute two important invariants of these numerical semigroups: the Frobenius number  $g(S(I))$  and multiplicity  $m(S(I))$ .

## 1. Preliminaries

We begin by defining a helpful function  $\kappa(a, b)$ . For  $a, b \in \mathbb{R}$  with  $a < b$  we define  $\kappa(a, b) = \lfloor b/(b-a) \rfloor$ . The function  $\kappa$  has various nice properties, for example  $\kappa(a, b) = \kappa(ac, bc)$  for  $c > 0$ . In the special case of  $a = m/n, b = m/(n-s)$ , we have  $\kappa(a, b) = \lfloor n/s \rfloor$ . The following properties of  $\kappa(a, b)$  are needed in the sequel.

**Lemma 1.1.** *Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $b \neq 0$ . Set  $\kappa = \kappa(a, b)$ . If  $\kappa \neq 0$ , then  $(\kappa - 1)/\kappa \leq a/b$ . If  $\kappa \neq -1$ , then  $a/b < \kappa/(\kappa + 1)$ .*

*MSC2000:* 20M10, 20M14.

*Keywords:* numerical semigroup, modular numerical semigroup.

*Proof.* We have  $\kappa \leq b/(b-a) < \kappa + 1$ . Assume that  $\kappa \notin \{-1, 0\}$ . Then

$$\kappa, \frac{b}{b-a}, \kappa + 1$$

all have the same sign, and we have  $1/\kappa \geq (b-a)/b > 1/(\kappa + 1)$ , hence

$$\frac{1-\kappa}{\kappa} \geq -\frac{a}{b} > \frac{-\kappa}{\kappa+1},$$

and the results follow. If  $\kappa = 0$ , then  $0 \leq b/(b-a) < 1$ , so  $b > 0$  and  $b < b-a$  so  $a < 0$  hence  $a/b < 0 = \kappa/(\kappa + 1)$ . If  $\kappa = -1$ , then  $-1 \leq b/(b-a) < 0$ , so  $b < 0$  and  $a-b \leq b$  so  $a \leq 2b$  and  $a/b \geq 2 = (\kappa - 1)/\kappa$ .  $\square$

**Lemma 1.2.** *Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $b > 0$ . Then*

$$\mathbb{N} \setminus S((a, b)) = \mathbb{N} \cap \bigcup_{n=1}^{\kappa(a,b)} [b(n-1), an].$$

*Proof.* Because  $S((a, b)) = \mathbb{N} \cap \bigcup_{n=1}^{\infty} (an, bn)$ , we have

$$\mathbb{N} \setminus S((a, b)) = \mathbb{N} \cap \bigcup_{n=1}^{\infty} [b(n-1), an].$$

Since  $b > 0$ ,  $\kappa(a, b) \neq -1$  and hence by Lemma 1.1,  $b\kappa(a, b) > a(\kappa(a, b) + 1)$ . Hence for  $n > \kappa(a, b)$ , the intervals are empty and may be excluded.  $\square$

Lemma 1.2 yields an upper bound for  $g$ . This bound will later be improved in Theorem 3.1, but for the purposes of Theorem 2.3 the following weaker bound suffices.

**Corollary 1.3.** *Suppose  $0 < a < b$ . Then  $g(S((a, b))) \leq \lfloor a\kappa(a, b) \rfloor$ .*

## 2. Intervals

We now prove that restricting  $I$  to be open is harmless, as this class of semigroups coincides with ones generated by closed or half-open intervals. To reduce the number of cases to consider, we introduce the symbols  $\{, \}$  to denote endpoints of an interval that are either open or closed. For example,  $(a, b\}$  indicates an interval that is open on the left. The meaning of these symbols is determined when first used, and then remains consistent; that is, if  $(a, b\}$  is open then  $[a', b\}$  means  $[a', b)$ , and if  $(a, b\}$  is half-open then  $[a', b\}$  means  $[a', b]$ .

The following lemma is the cornerstone of the interval equivalence results. Let  $d(x)$  denote the denominator of reduced rational  $x$ .

**Lemma 2.1.** *Let  $a \in \mathbb{Q}^{>0}$ ,  $n \in \mathbb{N}$ . Then all rationals in the interval*

$$\left( a - \frac{a}{nd(a)+1}, a + \frac{a}{nd(a)+1} \right),$$

other than possibly  $a$ , have numerator greater than  $n$ .

*Proof.* Suppose  $a = p/q$ , so  $d(a) = q$ . Consider any rational  $x/y$  with

$$0 < \left| \frac{x}{y} - a \right| < \frac{a}{nq + 1} = \frac{p}{q(nq + 1)}.$$

Also, we have

$$\left| \frac{x}{y} - \frac{p}{q} \right| = \left| \frac{xq - yp}{yq} \right| \geq \frac{1}{yq},$$

because  $xq - yp \neq 0$  since  $x/y \neq a$ . Combining, we get

$$\frac{p}{q(nq + 1)} > \frac{1}{yq},$$

hence

$$\frac{px}{nq + 1} > \frac{x}{y} > a - \frac{a}{nq + 1} = \frac{pnq}{q(nq + 1)} = \frac{pn}{nq + 1},$$

and thus  $x > n$ . □

**Lemma 2.2.** *Suppose that  $I, J, I \cup J$  are all intervals. Then  $S(I \cup J) = S(I) \cup S(J)$ . Also, if  $I \subseteq J$ , then  $S(I) \subseteq S(J)$  and  $g(S(I)) \geq g(S(J))$ .*

*Proof.* An integer  $m \in S(I \cup J)$  if and only if  $m/n \in I \cup J$  for some  $n$ . This is true if and only if  $m/n \in I$  or  $m/n \in J$ . Hence  $m \in S(I \cup J)$  if and only if  $m \in S(I) \cup S(J)$ . If  $I \subseteq J$ , then  $S(J) = S(I \cup J) = S(I) \cup S(J) \supseteq S(I)$ . □

The following theorem lets us replace a closed endpoint with an open one nearby, leaving the semigroup unchanged. Given a modular or proportionally modular numerical semigroup  $S$ , it explicitly gives an open interval  $I$  with  $S(I) = S$ .

**Theorem 2.3.** *Let  $0 < a < b$ . Set*

$$a' = \begin{cases} a - \frac{a}{[\text{a}\kappa(a, b)]d(a) + 1} & \text{if } a \in \mathbb{Q}, \\ a & \text{if } a \notin \mathbb{Q}, \end{cases}$$

$$b' = \begin{cases} b + \frac{b}{[\text{a}\kappa(a, b)]d(b) + 1} & \text{if } b \in \mathbb{Q}, \\ b & \text{if } b \notin \mathbb{Q}. \end{cases}$$

Then

$$S([a, b]) = S((a', b)) \quad \text{and} \quad S(\{a, b\}) = S(\{a, b'\}).$$

*Proof.* We consider only  $[a, b]$ ;  $\{a, b\}$  is symmetric. Suppose first that  $a \notin \mathbb{Q}$ . By Lemma 2.2,  $S([a', b]) = S((a', b)) \cup S([a', a']) = S((a', b))$  since  $S([a', a']) = \emptyset$ . We now assume  $a \in \mathbb{Q}$ . Since  $a < 2a - a'$ , Lemma 2.2 implies that  $S((a', b)) = S((a', 2a - a')) \cup S([a, b])$ . We will show  $S((a', 2a - a')) \subseteq S([a, b])$ , implying  $S((a', b)) \subseteq S([a, b])$  (and  $S((a', b)) \supseteq S([a, b])$  by Lemma 2.2).

Let  $c \in S((a', 2a - a'))$ . Hence there is some  $d \in \mathbb{N}$  so that  $c/d \in (a', 2a - a')$ . By Lemma 2.1, either  $c/d = a$  (in which case  $c \in S([a, b])$ ), or  $c > [\text{a}\kappa(a, b)]$ .

In the latter case, we apply Corollary 1.3 and  $c > \lfloor a\kappa(a, b) \rfloor \geq g(S((a, b))) \geq g(S(\lfloor a, b \rfloor))$ , so  $c \in S(\lfloor a, b \rfloor)$ . □

Theorem 2.4 is a counterpoint to Theorem 2.3, allowing us to replace an open endpoint with a closed one nearby. Proposition 5 in [Rosales and Urbano-Blanco 2006] tells us more: that every  $S(I)$  is proportionally modular; that is, there are  $m, n, s \in \mathbb{N}$  where  $S(I) = S(\lfloor m/n, m/(n - s) \rfloor)$ . Unfortunately neither of these results give an explicit formula such as in Theorem 2.3.

**Theorem 2.4.** *Let  $0 < a < b$ . Then there are  $a', b'$  with  $S((a, b)) = S(\lfloor a', b \rfloor)$  and  $S(\{a, b\}) = S(\{a', b\})$ . Further,  $a/a', b/b' \in \mathbb{Q}$ .*

*Proof.* We consider only  $(a, b)$ ;  $\{a, b\}$  is symmetric. Suppose first that  $a \notin \mathbb{Q}$ . By Lemma 2.2,  $S(\lfloor a', b \rfloor) = S((a', b)) \cup S(\lfloor a', a' \rfloor) = S((a', b))$  since  $S(\lfloor a', a' \rfloor) = \emptyset$ . We now assume  $a \in \mathbb{Q}$ . Let  $a_0$  be any rational in  $(a, b)$ , and consider the sequence given by  $a_i = \frac{1}{2}(a + a_{i-1})$ , for  $i \geq 1$ . By Lemma 2.2, we have  $S(\lfloor a_1, b \rfloor) \subseteq S(\lfloor a_2, b \rfloor) \subseteq \dots \subseteq S((a, b))$ . Set

$$X = S((a, b)) \setminus S(\lfloor a_1, b \rfloor),$$

a finite set. Set  $Z = \{x/y : x \in X, x/y \in (a, b)\}$ . Since  $a_i \rightarrow a$  and  $\min Z > a$ , there is some  $j > 0$  with  $Z \subseteq \lfloor a_j, b \rfloor$ , and hence  $X \subseteq S(\lfloor a_j, b \rfloor)$ . We take  $a' = a_j$ ; note that  $a' \in \mathbb{Q}$  by construction. □

### 3. Calculating $g(S((a, b)))$ and $m(S((a, b)))$

We now improve on Corollary 1.3 by calculating  $g(S((a, b)))$  exactly. Various other results are known in related contexts. For example, if  $S(\lfloor a, b \rfloor)$  is not a half-line, in [Rosales and Urbano-Blanco 2006] it was shown that

$$\frac{g(S(\lfloor a, b \rfloor))}{g(S(\lfloor a, b \rfloor)) - 1} < a < b < g(S(\lfloor a, b \rfloor)).$$

Also, if  $2 \leq a < b$  with  $a, b \in \mathbb{N}$ , in [Rosales and Vasco 2009] it was shown that  $g(S((a, b))) = b$ .

**Theorem 3.1.** *Suppose  $0 < a < b$ . Set  $\kappa = \kappa(a, b), \kappa' = \max(\kappa(a - 1, b - 1), 0)$ . Then  $g(S((a, b))) = \lfloor a\alpha \rfloor$ , where  $\alpha \in \mathbb{Z}$  satisfies  $\kappa' \leq \alpha \leq \kappa$ . Specifically,*

$$\alpha = \kappa - \sum_{i=\kappa'+1}^{\kappa} \prod_{j=i}^{\kappa} (1 + \lfloor aj \rfloor + \lfloor b(1 - j) \rfloor).$$

*Proof.* By Lemma 1.2,  $g(S((a, b))) = \lfloor a\alpha \rfloor$ , for the greatest integer  $\alpha$  where

$$\mathbb{N} \cap \lfloor b(a - 1), a\alpha \rfloor$$

is nonempty; in particular  $\alpha \leq \kappa$ . The lower bound  $\alpha \geq \kappa'$  is trivial when  $\kappa' = 0$ ; if  $b \leq 1$  then

$$\kappa(a - 1, b - 1) = \left\lfloor \frac{b - 1}{b - a} \right\rfloor \leq 0,$$

and hence  $\kappa' = 0$ . Otherwise,  $b > 1$  and so by Lemma 1.1,

$$\frac{\kappa' - 1}{\kappa'} \leq \frac{a - 1}{b - 1};$$

rearranging we get  $b(\kappa' - 1) \leq a\kappa' - 1$ . Hence the interval  $[b(\kappa' - 1), a\kappa']$  has length at least 1. It must therefore contain an integer, so  $\alpha \geq \kappa'$ .

To prove the formula for  $\alpha$ , for  $i \leq \kappa$  we define the function

$$f(i) = \begin{cases} 1 & \text{if } \alpha \leq i, \\ 0 & \text{if } \alpha > i; \end{cases}$$

this gives us  $\alpha = \kappa - \sum_{i=0}^{\kappa} f(i) = \kappa - \sum_{i=\kappa'+1}^{\kappa} f(i)$ . We define  $f$  via  $f(i) = \prod_{j=i}^{\kappa} \chi(j)$ , for

$$\chi(j) = \begin{cases} 1 & \text{if } [b(j - 1), aj] \cap \mathbb{N} \neq \emptyset, \\ 0 & \text{if } [b(j - 1), aj] \cap \mathbb{N} = \emptyset. \end{cases}$$

We now have  $\alpha = \kappa - \sum_{i=\kappa'+1}^{\kappa} \prod_{j=i}^{\kappa} \chi(j)$ .

We now calculate  $\chi(j)$  explicitly by showing that for  $j \geq \kappa' + 1$ , the interval  $[b(j - 1), aj]$  contains at most one integer. For  $b \leq 1$ , we have  $bj > aj \geq aj + (b - 1)$  so  $b(j - 1) > aj - 1$ . For  $b > 1$ , by Lemma 1.1 we have

$$\frac{a - 1}{b - 1} < \frac{\kappa'}{\kappa' + 1} \leq \frac{j - 1}{(j - 1) + 1}, \quad \text{for any } j - 1 \geq \kappa'.$$

Rearranging, we get  $b(j - 1) > aj - 1$ . Hence  $|[b(j - 1), aj] \cap \mathbb{N}| \leq 1$  and in fact  $\chi(j)$  equals the number of integers in  $[b(j - 1), aj]$ , that is,

$$\chi(j) = 1 + \lfloor aj \rfloor + \lfloor b(1 - j) \rfloor. \quad \square$$

We have  $\alpha \in [\kappa', \kappa]$ ; in general, neither bound can be improved. The size of this interval,  $\kappa - \kappa'$ , can be arbitrarily large, when  $b/a$  is small. On the other hand, the following shows that  $\kappa - \kappa'$  is small if  $b/a > 2$ . This is desirable, as it shortens the calculation for  $g(S(I))$ .

**Proposition 3.2.** *Let  $0 < 2a < b$ . Let  $\kappa, \kappa'$  be as in Theorem 3.1. Then*

$$\kappa - \kappa' = \begin{cases} 1 & \text{if } a < 1, \\ 0 & \text{if } a \geq 1. \end{cases}$$

*Proof.* For convenience, set

$$I = \left( \frac{b-1}{b-a}, \frac{b}{b-a} \right);$$

$\kappa - \kappa'$  counts the number of integers in  $I$ . Suppose first that  $b \leq 1$ . Then

$$\kappa(a-1, b-1) \leq 0,$$

so  $\kappa' = 0$ . Note that  $b > 2a$  implies  $b - a > a$ , and hence  $1/(b - a) < 1/a$ , so  $1 + a/(b - a) < 1 + a/a = 2$ , and hence  $\kappa = \lfloor 1 + a/(b - a) \rfloor = 1$ . Suppose now that  $a < 1 < b$ . If  $a \leq \frac{1}{2}$ , then  $b > 1 = \frac{1}{2} + a$ . Alternatively, if  $a > \frac{1}{2}$ , then  $b > 2a > \frac{1}{2} + a$ . Hence  $b > \frac{1}{2} + a$ ; rearranging we get  $1/(b - a) < 2$ . Hence  $I$  is of length less than 2, and can contain at most one integer. Therefore  $\kappa - \kappa' \leq 1$ . But  $I$  contains the integer  $1 = (b - a)/(b - a)$ , so  $\kappa - \kappa' = 1$ . Lastly, we consider the case  $a \geq 1$ . We have  $b - 1 \geq b - a$ , hence  $(b - 1)/(b - a) \geq 1$  and  $I$  does not contain 0 or 1. Suppose  $I$  contains integer  $n \geq 2$ . Then  $2 \leq b/(b - a)$ ; rearranging we get  $b \leq 2a$ , a contradiction. Hence  $I$  contains no integers, and  $\kappa - \kappa' = 0$ .  $\square$

Computing  $m(S((a, b)))$  is similar to computing  $g(S((a, b)))$ , in that we must count integers in an interval, only this time the intervals are open. We first prove a technical lemma, for which we recall Farey sequences (for an introduction see [Graham et al. 1994]). The  $n$ th Farey sequence  $F_n$  consists of all reduced fractions in  $[0, 1]$  whose denominator is at most  $n$ , arranged in increasing order. The key property we require is that if  $a/b$  and  $c/d$  are consecutive terms in a Farey sequence, then  $bc - ad = 1$ .

**Lemma 3.3.** *Let  $0 < a < b$ . Let  $n \in \mathbb{N}$  be minimal such that  $(an, bn)$  contains an integer. Suppose  $n > 1$ . Then  $(an, bn)$  contains exactly one integer.*

*Proof.* Suppose by way of contradiction that  $(an, bn)$  contains at least two integers. Then there is some  $m \in \mathbb{N}$  such that  $m, m + 1 \in (an, bn)$ . Set  $d = \gcd(m, n)$ . If  $d > 1$  then  $m/d \in (an/d, bn/d)$  violates the minimality of  $n$ . Similarly,  $\gcd(m + 1, n) = 1$ . Let  $m' \in (0, n - 1)$  with  $m = m' + kn$  for some integer  $k$ . We now consider the  $n$ th Farey sequence  $F_n$ . Both  $m'/n$  and  $(m' + 1)/n$  are elements of  $F_n$ ; however  $(m' + 1)n - m'n = n > 1$ , so they are not consecutive terms and there must be some  $p/q$  in  $F_n$  with  $m'/n < p/q < (m' + 1)/n$ , with  $q < n$ . But then

$$\frac{m' + kn}{n} < \frac{p + qk}{q} < \frac{m' + 1 + kn}{n},$$

so  $p + qk \in (aq, bq)$ , violating the minimality of  $n$ .  $\square$

We now compute the multiplicity  $m(S((a, b)))$ . The reverse problem of finding an open interval whose semigroup possesses a given multiplicity, is solved in [Rosales and Vasco 2009]. A nondiscrete version is proved as Proposition 5 in [Rosales et al. 2003].

**Theorem 3.4.** *Suppose  $0 < a < b$ . Set  $\kappa'' = \kappa(1, b - a + 1)$ . Then*

$$m(S((a, b))) = \lceil a\alpha \rceil,$$

where  $\alpha \in \mathbb{N}$  satisfies  $1 \leq \alpha \leq \kappa''$ . Specifically,

$$\alpha = \sum_{i=0}^{\kappa''} \prod_{j=1}^i (2 + \lfloor aj \rfloor + \lfloor -bj \rfloor).$$

*Proof.* Set  $m = m(S((a, b)))$ , and let  $\alpha \in \mathbb{N}$  be minimal such that  $m/\alpha \in (a, b)$ ; then  $m(S((a, b))) = \lceil a\alpha \rceil$ . By Lemma 1.1,  $1/(b - a + 1) < \kappa''/(\kappa'' + 1)$ . Rearranging, we find  $\kappa''b - \kappa''a > 1$ , so there is an integer  $t \in (\kappa''a, \kappa''b)$ . Suppose that  $a > \kappa''$ . We then have  $m/\alpha < m/\kappa'' \leq t/\kappa''$ ; since  $m/\alpha$  and  $t/\kappa''$  are in  $(a, b)$ , we conclude that  $m/\kappa'' \in (a, b)$ , which contradicts the minimality of  $\alpha$ . Hence  $\alpha \leq \kappa''$ .

We now prove the  $\alpha$  formula. We proceed in a manner similar to Theorem 3.1, by defining

$$f(i) = \begin{cases} 1 & i \leq \alpha, \\ 0 & i > \alpha, \end{cases}$$

via  $f(i) = \prod_{j=1}^i (1 - \chi(j))$ , where  $\chi(j)$  is the number of integers in  $(aj, bj)$ . For  $i < \alpha$ ,  $\chi(i) = 0$ . By Lemma 3.3,  $\chi(\alpha) = 1$ , so  $f(i) = 0$  for  $i \geq \alpha$ . Hence

$$\alpha = \sum_{i=0}^{\kappa''} f(i) = \sum_{i=0}^{\kappa''} \prod_{j=1}^i (1 - \chi(j)),$$

but  $1 - \chi(j) = 2 + \lfloor aj \rfloor + \lfloor -bj \rfloor$ . □

We have  $\alpha \in [1, \kappa'']$ ; in general, neither bound can be improved. The upper bound  $\kappa''$  can be arbitrarily large, when  $b - a$  is small. On the other hand, the following shows that  $\kappa''$  is small if  $b - a$  is large, thus simplifying computation of  $m$ .

**Proposition 3.5.** *Let  $0 < a < b$ . Let  $n \in \mathbb{N}$  be minimal with  $b - a > 1/n$ . Then  $\kappa'' = n$ , in the notation of Theorem 3.4.*

*Proof.* We have  $1/n < b - a \leq 1/(n - 1)$ , hence  $n > 1/(b - a) \geq n - 1$ , so  $\lfloor 1/(b - a) \rfloor = n - 1$ , and

$$\kappa'' = \lfloor \frac{b - a + 1}{b - a} \rfloor = \lfloor 1 + \frac{1}{b - a} \rfloor = 1 + (n - 1) = n. \quad \square$$

### Acknowledgments

The authors would like to thank the anonymous referee for extensive and helpful suggestions.



## References

- [Graham et al. 1994] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete mathematics*, 2nd ed., Addison-Wesley Publishing Company, Reading, MA, 1994. MR 97d:68003 Zbl 0836.00001
- [Rosales and García-Sánchez 2009] J. C. Rosales and P. A. García-Sánchez, *Numerical semigroups*, Developments in Mathematics **20**, Springer, New York, 2009. MR 2010j:20091 Zbl 05623301
- [Rosales and Urbano-Blanco 2006] J. C. Rosales and J. M. Urbano-Blanco, “Opened modular numerical semigroups”, *J. Algebra* **306**:2 (2006), 368–377. MR 2007h:20063 Zbl 1109.20052
- [Rosales and Vasco 2009] J. C. Rosales and P. Vasco, “Opened modular numerical semigroups with a given multiplicity”, *Internat. J. Algebra Comput.* **19**:2 (2009), 235–246. MR 2010i:20082 Zbl 1187.20067
- [Rosales et al. 2003] J. C. Rosales, P. A. García-Sánchez, J. I. García-García, and J. M. Urbano-Blanco, “Proportionally modular Diophantine inequalities”, *J. Number Theory* **103**:2 (2003), 281–294. MR 2004k:20127 Zbl 1039.20036
- [Rosales et al. 2005] J. C. Rosales, P. A. García-Sánchez, and J. M. Urbano-Blanco, “Modular Diophantine inequalities and numerical semigroups”, *Pacific J. Math.* **218**:2 (2005), 379–398. MR 2007a:20056 Zbl 1184.20052

Received: 2010-06-30    Revised: 2010-09-29    Accepted: 2010-09-29

vadim@sciences.sdsu.edu

*Department of Mathematics and Statistics,  
San Diego State University, 5500 Campanile Dr.,  
San Diego, CA 92182-7720, United States  
<http://www-rohan.sdsu.edu/~vadim/>*

bhappening@gmail.com

*San Diego State University, San Diego, CA 92182-7720,  
United States*

# involve

pjm.math.berkeley.edu/involve

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

### BOARD OF EDITORS

John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Pietro Cerone	Victoria University, Australia pietro.cerone@vu.edu.au	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Ken Ono	University of Wisconsin, USA ono@math.wisc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	Robert J. Plemmons	Wake Forest University, USA plemmons@wfu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Sat Gupta	U of North Carolina, Greensboro, USA sgupta@uncg.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Filip Saidak	U of North Carolina, Greensboro, USA f.saidak@uncg.edu
Karen Kafadar	University of Colorado, USA karen.kafadar@cudenver.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
David Larson	Texas A&M University, USA larson@math.tamu.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu

## PRODUCTION

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor

Cover design: ©2008 Alex Scorpan

See inside back cover or <http://pjm.math.berkeley.edu/involve> for submission instructions.

The subscription price for 2010 is US \$100/year for the electronic version, and \$120/year (+\$20 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94704-3840, USA.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY  
 mathematical sciences publishers  
<http://www.mathscipub.org>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2010 by Mathematical Sciences Publishers

# involve

2010

vol. 3

no. 3

Gracefulness of families of spiders	241
PATRICK BAHLS, SARA LAKE AND ANDREW WERTHEIM	
Rational residuacity of primes	249
MARK BUDDEN, ALEX COLLINS, KRISTIN ELLIS LEA AND STEPHEN SAVIOLI	
Coexistence of stable ECM solutions in the Lang–Kobayashi system	259
ERICKA MOCHAN, C. DAVIS BUENGER AND TAMAS WIANDT	
A complex finite calculus	273
JOSEPH SEABORN AND PHILIP MUMMERT	
$\zeta(n)$ via hyperbolic functions	289
JOSEPH D'AVANZO AND NIKOLAI A. KRYLOV	
Infinite family of elliptic curves of rank at least 4	297
BARTOSZ NASKRECKI	
Curvature measures for nonlinear regression models using continuous designs with applications to optimal experimental design	317
TIMOTHY O'BRIEN, SOMSRI JAMROENPINYO AND CHINNAPHONG BUMRUNGSUP	
Numerical semigroups from open intervals	333
VADIM PONOMARENKO AND RYAN ROSENBAUM	
Distinct solution to a linear congruence	341
DONALD ADAMS AND VADIM PONOMARENKO	
A note on nonresidually solvable hyperlinear one-relator groups	345
JON P. BANNON AND NICOLAS NOBLETT	



1944-4176(2010)3:3;1-E