

involve

a journal of mathematics

Diameter, girth and cut vertices of the graph of equivalence
classes of zero-divisors

Blake Allen, Erin Martin, Eric New and Dane Skabelund

 mathematical sciences publishers

Diameter, girth and cut vertices of the graph of equivalence classes of zero-divisors

Blake Allen, Erin Martin, Eric New and Dane Skabelund

(Communicated by Scott Chapman)

We explore the properties of $\Gamma_E(R)$, the graph of equivalence classes of zero-divisors of a commutative Noetherian ring R . We determine the possible combinations of diameter and girth for the zero-divisor graph $\Gamma(R)$ and the equivalence class graph $\Gamma_E(R)$, and examine properties of cut-vertices of $\Gamma_E(R)$.

Introduction

The zero-divisor graph of a commutative ring R , was first introduced in [Beck 1988] and has since been investigated in various forms. It was shown in [Anderson and Livingston 1999] that the zero-divisor graph of any ring is connected with diameter less than or equal to 3. Mulay [2002] proved many interesting results about cycles in the zero-divisor graph.

In 2009, Spiroff and Wickham [2011] introduced $\Gamma_E(R)$, the graph of equivalence classes of zero-divisors, which is a simplification of the zero-divisor graph $\Gamma(R)$. The vertices of $\Gamma_E(R)$ are, instead of individual zero-divisors of R , equivalence classes of zero-divisors determined by annihilator ideals. The graph $\Gamma_E(R)$ provides a more succinct view of the zero-divisor activity of the ring. In many cases, the equivalence class graph is finite even though the zero-divisor graph is infinite. For example, for $S = \mathbb{Z}[X, Y]/(X^4, XY)$, the graph $\Gamma(S)$ is infinite, while the graph $\Gamma_E(S)$ has only 6 vertices. Specifically, the vertices corresponding to $X^3, 2X^3, 3X^3, \dots$ are all distinct in $\Gamma(S)$. However, since they all have the same annihilator, they all belong to the same equivalence class, and so are represented by a single vertex $[X^3]$ in $\Gamma_E(S)$.

The equivalence class graph also lets us view the interplay between the annihilator ideals of R and helps to easily identify the associated primes of the ring. The vertices of $\Gamma_E(R)$ which correspond to associated primes have special properties which will help us to prove several interesting results related to $\Gamma_E(R)$. In

MSC2010: 13A99.

Keywords: zero-divisor graph, diameter, girth, cut vertices.

Section 1, we provide basic definitions and background. In Section 2, we determine all possible diameter combinations of $\Gamma(R)$ and $\Gamma_E(R)$, and do the same for the girth of the two graphs in Section 3. In Section 4, we look at properties of the cut-vertices of $\Gamma_E(R)$. Throughout, R will denote a commutative Noetherian ring.

1. Background and basic results

Graph theory. We briefly review basic graph theory terms that we will use throughout the paper. All graphs we deal with will be *simple* graphs in the sense that they contain no loops or double edges. We will denote the set of vertices of a graph Γ by $V(\Gamma)$. If two vertices x and y are joined by an edge, we say x and y are *adjacent*, and write $x - y$. A *path* is defined as an alternating sequence of distinct vertices and edges, and the *length of a path* is the number of edges in the path. If x and y are two vertices, then the *distance between x and y* , denoted $d(x, y)$, is the length of the shortest path from x to y . If there is no path connecting x to y , we say that $d(x, y) = \infty$, and we define $d(x, x) = 0$. The *diameter of a graph* is the maximum distance between any two vertices of the graph. We will denote the diameter of a graph Γ by $\text{diam } \Gamma$. A *cycle* is a closed path, or a path that starts and ends on the same vertex. The *girth* of a graph is the length of its smallest cycle. We denote the girth of a graph Γ by $g(\Gamma)$ and say that $g(\Gamma) = \infty$ if the graph Γ contains no cycle. Note that the smallest possible cycle length is 3, so if Γ contains a cycle, $g(\Gamma) \geq 3$.

A graph is said to be *connected* if every pair of vertices is joined by a path and *complete* if every pair of vertices is joined by an edge. A *connected component* of a graph Γ is a maximal connected subgraph of Γ . If removing a vertex v from a graph along with all its incident edges increases the number of connected components in the graph, then v is called a *cut vertex*. A graph is *complete bipartite* if its vertices can be partitioned into two subsets, V_1 and V_2 , such that every vertex of V_1 is adjacent to every vertex of V_2 , but no two vertices of V_1 are adjacent and no two vertices of V_2 are adjacent. Such a graph will be denoted $K_{n,m}$, where $n = |V_1|$ and $m = |V_2|$. If the vertices of a graph can be partitioned into r subsets in a similar fashion, then the graph is said to be *r -partite*.


Zero-divisor graphs. Let $Z(R)$ denote the set of zero-divisors of R and $Z^*(R)$ denote the set $Z(R) \setminus \{0\}$. We define the *zero-divisor graph* of R as the simple graph $\Gamma(R)$ where the vertices of $\Gamma(R)$ are the elements of $Z^*(R)$, and there is an edge between $x, y \in \Gamma(R)$ whenever $xy = 0$.

Recall that the annihilator ideal associated to an element $x \in R$ is the set $\text{ann } x = \{r \in R : xr = 0\}$. We define an equivalence relation \sim on R such that for all $x, y \in R$, we say $x \sim y$ if $\text{ann } x = \text{ann } y$. Let $[x]$ denote the equivalence class of x . Notice

that $[0] = \{0\}$, $[1] = R \setminus Z(R)$ and the relation \sim partitions the remaining zero-divisors into distinct classes. Furthermore, it follows that the multiplication of these equivalence classes $[x] \cdot [y] = [xy]$ is well-defined.

The graph of equivalence classes of zero-divisors of R , $\Gamma_E(R)$, is the graph whose vertices are the classes of nonzero zero-divisors of R determined by the relation \sim , where there is an edge between two vertices $[x]$ and $[y]$ if $[x] \cdot [y] = [0]$.

Here, as an example, are the zero-divisor graph of \mathbb{Z}_{12} and the graph of its equivalence classes:



We see that since $\text{ann } 2 = \text{ann } 10$, the elements 2 and 10 are in the same equivalence class, and therefore collapse to the single vertex $[2]$ in $\Gamma_E(R)$.

Previous results. Spiroff and Wickham [2011] have several interesting results linking the associated primes of R with the structure of $\Gamma_E(R)$. These will be useful in furthering our investigation of $\Gamma_E(R)$. Remember that a prime ideal \mathfrak{p} of R is an *associated prime* if it is the annihilator of some element of R . The set of associated primes is denoted $\text{ass } R$. It is well known that if R is a Noetherian ring, then $\text{ass } R$ is nonempty and finite and that any maximal element of the family of annihilator ideals $\mathfrak{F} = \{\text{ann } x : 0 \neq x \in R\}$ is an associated prime. Note also that since every zero divisor is contained in an annihilator ideal and maximal annihilators are associated primes, the set of zero-divisors of R equals the union of all associated primes of R . Since there is exactly one vertex of $\Gamma_E(R)$ for each distinct annihilator ideal of R , we have a natural injection of $\text{ass } R$ into the vertex set of $\Gamma_E(R)$ given by $\mathfrak{p} \mapsto [y]$ where $\mathfrak{p} = \text{ann } y$. We adopt the conventions of Spiroff and Wickham and by a slight abuse of terminology will refer to the vertex $[y]$ as an associated prime if $\text{ann } y \in \text{ass } R$. It will be clear from context whether $[y]$ refers to an equivalence class, a vertex, or a specific annihilator.

Lemma 1.1 [Spiroff and Wickham 2011, Lemma 1.2]. *Any two distinct elements of $\text{ass } R$ are connected by an edge. Furthermore, every vertex $[v]$ of $\Gamma_E(R)$ is either an associated prime or adjacent to an associated prime maximal in \mathfrak{F} .*

Lemma 1.2 [Spiroff and Wickham 2011, Proposition 1.7]. *Let R be a ring such that $\Gamma_E(R)$ is complete r -partite. Then $r = 2$ and $\Gamma_E(R) = K_{n,1}$ for some $n \geq 1$.*

2. Diameter

In this section, we explore the relationship between the diameters of the graphs $\Gamma(R)$ and $\Gamma_E(R)$. It is shown in [Anderson and Livingston 1999] that $\Gamma(R)$ has diameter at most 3 for any commutative ring R . In [Spiroff and Wickham 2011] it is shown that $\text{diam } \Gamma_E(R) \leq 3$ for R commutative and Noetherian. The following results further demonstrate the relationship between the diameters of the two graphs.

Proposition 2.1. *If R is a commutative ring, then $\text{diam } \Gamma_E(R) \leq \text{diam } \Gamma(R)$.*

Proof. Let $[a], [b] \in \Gamma_E(R)$ with $d([a], [b]) = n$, and let $[a] = [x_1] - [x_2] - \cdots - [x_{n+1}] = [b]$ be a path of minimal length from $[a]$ to $[b]$. From each $[x_i]$, choose one $y_i \in [x_i]$. Then $y_1 - y_2 - \cdots - y_{n+1}$ is a path in $\Gamma(R)$ of length n . We claim that this path is minimal, and thus $d(y_1, y_{n+1}) = n$. If this path is not minimal, there is some shorter path $y_1 = z_1 - z_2 - \cdots - z_{m+1} = y_{n+1}$, with $m < n$. Since either $[z_i] = [z_{i+1}]$ or $[z_i] - [z_{i+1}]$, the path $[y_1] = [z_1] - [z_2] - \cdots - [z_{m+1}] = [y_{n+1}]$ has length less than or equal to m , a contradiction. \square

Theorem 2.2. *If $\text{diam } \Gamma_E(R) = 0$, then $\text{diam } \Gamma(R) = 0$ or 1.*

Proof. Let $\Gamma_E(R)$ have diameter 0. Since $\Gamma_E(R)$ has only one vertex, $[x] = [y]$ for every $x, y \in Z^*(R)$. Since the graph $\Gamma(R)$ is connected and every element in $\Gamma(R)$ has the same annihilator, $xy = 0$ for every $x, y \in Z^*(R)$. Thus the graph $\Gamma(R)$ is complete and $\text{diam } \Gamma(R) = 0$ or 1. \square

Theorem 2.3. *If $\text{diam } \Gamma(R) = 3$, then $\text{diam } \Gamma_E(R) = 3$.*

Proof. Let $\Gamma(R)$ have diameter 3. Then for some elements $x, w \in \Gamma(R)$, $d(x, w) = 3$ in $\Gamma(R)$. Let $x - y - z - w$ be a path from x to w of minimal length. Since this path is minimal, $xz \neq 0$, but $zw = 0$, so $\text{ann } x \neq \text{ann } w$. By similar reasoning we see that each of $\text{ann } x$, $\text{ann } y$, $\text{ann } z$, and $\text{ann } w$ are distinct. Hence $[x]$, $[y]$, $[z]$, and $[w]$ are distinct equivalence classes in $\Gamma_E(R)$. Thus $[x]$ is not adjacent to $[w]$ and there exist no paths $[x] - [y] - [w]$ or $[x] - [z] - [w]$ in $\Gamma_E(R)$. Now suppose there is some other $[v]$ such that $[x] - [v] - [w]$. This is impossible because it implies that there is a path $x - v - w$ in $\Gamma(R)$, contradicting the supposition that $x - y - z - w$ is a minimal path. Therefore $d([x], [w]) = 3$ and since $\text{diam } \Gamma_E(R) \leq 3$, $\text{diam } \Gamma_E(R) = 3$. \square

We summarize with Table 1, which shows all possible combinations of diameter for $\Gamma(R)$ and $\Gamma_E(R)$.

We see from our examples that it is possible for the diameter of the zero-divisor graph to shrink under the equivalence relation. We consider the situations where this happens.

If $\text{diam } \Gamma(R) = 1$ and $\text{diam } \Gamma_E(R) = 0$, then R has a unique annihilator ideal $\text{ann } x$. This annihilator is maximal in \mathfrak{F} and an associated prime of the ring. Since $Z(R) = \bigcup_{\mathfrak{p} \in \text{ass } R} \mathfrak{p} = \text{ann } x$, $Z(R)$ forms an ideal of R .


| diam $\Gamma(R)$ | diam $\Gamma_E(R) =$ | | | |
|------------------|---------------------------------------|--|-------------------|-------------------|
| | 0 | 1 | 2 | 3 |
| 0 | $\mathbb{Z}_4, \mathbb{Z}_2[x]/(x^2)$ | – | – | – |
| 1 | $\mathbb{Z}_9, \mathbb{Z}_{25}$ | $\mathbb{Z}_2 \times \mathbb{Z}_2$ | – | – |
| 2 | impossible (Theorem 2.2) | $\mathbb{Z}_6, \mathbb{Z}_{21}, \mathbb{Z}_2[x]/(x^3)$ | \mathbb{Z}_{16} | – |
| 3 | impossible (Theorem 2.3) | | | \mathbb{Z}_{12} |

Table 1. Possibilities for $\text{diam } \Gamma(R)$ and $\text{diam } \Gamma_E(R)$, with examples.

Next we consider the situation in which the diameter reduces from 2 to 1. Since there are no complete equivalence class graphs on 3 or more vertices, by [Spiroff and Wickham 2011, Proposition 1.5], $\Gamma_E(R)$ must have exactly two vertices, and R must have exactly 2 distinct annihilator ideals, $\text{ann } x$ and $\text{ann } y$. Let $\text{ann } x$ be maximal in \mathfrak{F} . If $\text{ann } y \subseteq \text{ann } x$, then $Z(R) = \bigcup_{\mathfrak{p} \in \text{ass } R} \mathfrak{p} = \text{ann } x$ forms an ideal of R . Otherwise, both $\text{ann } x$ and $\text{ann } y$ are maximal in \mathfrak{F} and $\text{ann } x \cap \text{ann } y = \{0\}$. If we have nonzero a, b with $a \in \text{ann } x$ and $b \in \text{ann } y$ such that $a + b \in \text{ann } x$, then $b \in \text{ann } x$, a contradiction. So in this case $Z(R) = \bigcup_{\mathfrak{p} \in \text{ass } R} \mathfrak{p} = \text{ann } x \cup \text{ann } y$ does not form an ideal of R .

Therefore we see that if the diameter shrinks in the equivalence class graph, R has 1 or 2 associated primes. If R is a finite ring, this corresponds to R being the direct product of 1 or 2 local rings, since every finite ring R is expressible as the product of finite local rings, with the number of factors equal to the number of associated primes of R .

We show below examples of graphs of rings with shrinking diameter, one from each of the situations considered above. Note that \mathbb{Z}_{25} has a unique annihilator,



$\text{ann } 5 = (5)$, $\mathbb{Z}_4[x]/(2x, x^2 - 2)$ has two annihilators, $\text{ann } x = (2) \subseteq (2, x) = \text{ann } 2$, and \mathbb{Z}_{15} has two annihilators, $\text{ann } 3 = (5)$ and $\text{ann } 5 = (3)$, which intersect trivially.

3. Girth

Mulay [2002] proved that if the zero-divisor graph, $\Gamma(R)$, contains a cycle then $g(\Gamma(R)) \leq 4$. In this section we will demonstrate an even stronger restriction on the girth of the equivalence class graph, and find all possible combinations of girth for $\Gamma(R)$ and $\Gamma_E(R)$. The following result gives a girth restriction for $\Gamma_E(R)$ similar to that shown by Mulay for $\Gamma(R)$.

Theorem 3.1. *If R is a commutative Noetherian ring, and if $\Gamma_E(R)$ contains a cycle, then $g(\Gamma_E(R)) \leq 4$.*

Proof.

Case 1: If R has at least 3 distinct associated primes, say $\text{ann } x$, $\text{ann } y$, and $\text{ann } z$, then the vertices $[x]$, $[y]$, and $[z]$ in $\Gamma_E(R)$ are all adjacent to each other by Lemma 1.1, and therefore span a complete subgraph of $\Gamma_E(R)$. Hence $\Gamma_E(R)$ contains a 3-cycle, so $g(\Gamma_E(R)) = 3$.

Case 2: If R has exactly one associated prime, $\text{ann } y$, then every other vertex in $\Gamma_E(R)$ is adjacent to $[y]$ by Lemma 1.1. If there is any cycle in $\Gamma_E(R)$, then there are some vertices $[x_1], [x_2]$ distinct from $[y]$ with $[x_1] - [x_2]$. But these are both adjacent to $[y]$, creating the 3-cycle $[y] - [x_1] - [x_2] - [y]$. So $g(\Gamma_E(R)) = 3$.

Case 3: Now assume that R has exactly 2 associated primes, and let $\text{ass } R = \{\text{ann } x, \text{ann } y\}$. Let $[x_1]$ and $[x_2]$ be two vertices distinct from $[x]$ and $[y]$ such that $[x_1] - [x_2]$. By Lemma 1.1, $[x_1]$ is adjacent to an associated prime. Without loss of generality, let $[x_1] - [x]$. Also, $[x_2]$ is adjacent to either $[x]$ or $[y]$. In the first case, we have a 3-cycle $[x] - [x_1] - [x_2] - [x]$ and in the second case, we have a 4-cycle $[x] - [x_1] - [x_2] - [y]$. Now assume that given any two vertices of $\Gamma_E(R)$, at least one is an associated prime. Let $[x_1] - [x_2] - \cdots - [x_n] - [x_1]$ be a cycle in $\Gamma_E(R)$ of minimal length, and let $n \geq 4$. Since at least one of $[x_1]$ and $[x_2]$ is an associated prime, without loss of generality let $[x_1]$ be an associated prime. Also, at least one of $[x_3]$ and $[x_4]$ is an associated prime. If $[x_3]$ is an associated prime, we have the 3-cycle $[x_1] - [x_2] - [x_3] - [x_1]$, and if $[x_4]$ is an associated prime, we have the 4-cycle $[x_1] - [x_2] - [x_3] - [x_4] - [x_1]$. \square

The following corollary is a direct result of the proof of Theorem 3.1.

Corollary 3.2. *If $\Gamma_E(R)$ has girth 4, then R must have exactly 2 associated primes.*

The following proposition gives a relationship between the girths of the two graphs. Note that the inequality is opposite that of the diameter relationship stated in the previous section.

Proposition 3.3. *If $\Gamma_E(R)$ contains a cycle, then $g(\Gamma_E(R)) \geq g(\Gamma(R))$.*

Proof. Let $[x_1] - [x_2] - \cdots - [x_n] - [x_1]$ be a cycle in $\Gamma_E(R)$. For each $[x_i]$, choose one $y_i \in [x_i]$. Then by the definition of multiplication of our equivalence classes, $y_1 - y_2 - \cdots - y_n - y_1$ is a cycle in $\Gamma(R)$ of equal length. So $g(\Gamma_E(R)) \geq g(\Gamma(R))$. \square

Corollary 3.4. *If $g(\Gamma_E) = 3$, then $g(\Gamma) = 3$.*

We now examine the situation in which $\Gamma_E(R)$ has girth 4 and conclude that it is impossible.

Theorem 3.5. *For R a commutative Noetherian ring, $g(\Gamma_E(R)) \neq 4$.*

Proof.

Suppose that $\Gamma_E(R)$ has girth 4. By Corollary 3.2, R has exactly two associated primes, so let $\text{ass } R = \{\text{ann } x, \text{ann } y\}$.

Since $\text{ann } x$ and $\text{ann } y$ are associated primes, $[x] - [y]$ by Lemma 1.1. Let $[z]$ be some other vertex of $\Gamma_E(R)$. Then $[z]$ must be adjacent to at least one of $[x]$ or $[y]$. But if it is adjacent to both $[x]$ and $[y]$ we have a 3-cycle, so $[z]$ is adjacent to exactly one of $[x]$ or $[y]$. Thus the vertex set of $\Gamma_E(R)$ minus $\{[x], [y]\}$ can be partitioned into two disjoint subsets, one adjacent to $[x]$ and one adjacent to $[y]$. We refer to these subsets as X and Y , respectively.

As mentioned earlier, since R is Noetherian, there is at least one maximal element of \mathfrak{F} , and this annihilator is an associated prime. Without loss of generality, let $\text{ann } x$ be maximal in \mathfrak{F} . We claim that $\text{ann } y$ is also maximal in \mathfrak{F} . Now if $\text{ann } y \subseteq \text{ann } w$ for some w , then $\text{ann } w \subseteq \text{ann } m$ for some maximal element $\text{ann } m \in \mathfrak{F}$, but since $\text{ann } m$ is an associated prime, $\text{ann } m = \text{ann } y$ or $\text{ann } m = \text{ann } x$. In the latter case, $\text{ann } y \subseteq \text{ann } x$, so $[x]$ and $[y]$ are both adjacent to a common vertex. This creates a 3-cycle, contradicting that $g(\Gamma_E(R)) = 4$. So both $\text{ann } y$ and $\text{ann } x$ are maximal in \mathfrak{F} .

Suppose that $[x]^2 = [0]$ and $[y]^2 = [0]$, and consider the class $[x + y]$. This class is annihilated by both $[x]$ and $[y]$, so either $[x + y] = [0]$ or $[x + y]$ is in the vertex set of $\Gamma_E(R)$. If $[x + y] = [0]$, then $[x] = [y]$, contrary to our assumption. So $[x + y]$ is in the vertex set of $\Gamma_E(R)$. Since $[y]$ is adjacent to no vertex of X , $[x + y] \neq [x]$. Similarly, since $[x]$ is adjacent to no vertex of Y , $[x + y] \neq [y]$. So $\Gamma_E(R)$ contains the 3-cycle $[x + y] - [x] - [y] - [x + y]$, a contradiction.

Now suppose that $[x]^2 \neq [0]$ and $[y]^2 \neq [0]$. Then $\text{ann } x \cap \text{ann } y = \{0\}$. Now multiplying any $[x_j] \in X$ and $[y_i] \in Y$, we see that since $[x_j] \in \text{ann } x$ and $[y_i] \in \text{ann } y$, $[x_j y_i] \in \text{ann } x \cap \text{ann } y = \{0\}$. If we break up the vertex set of $\Gamma_E(R)$ into $X \cup \{[y]\}$ and $Y \cup \{[x]\}$, we see that $\Gamma_E(R)$ is complete bipartite, and $\Gamma_E(R) = K_{n,m}$ with $n, m \neq 1$, which contradicts Lemma 1.2.


Without loss of generality, let $[x]^2 = [0]$ and $[y]^2 \neq [0]$. Let $[x] - [y] - [z] - [w] - [x]$ be a 4-cycle in $\Gamma_E(R)$, with $[w] \in X$, $[z] \in Y$. Then there is a 4-cycle $x - y - z - w - x$ in $\Gamma(R)$. By the previous discussion, $x^2 = 0$ and $y^2 \neq 0$.

| diam $\gamma(R)$ | diam $\gamma_E(R) =$ | | |
|------------------|----------------------|------------------------------|--------------------------|
| | ∞ | 3 | 4 |
| ∞ | \mathbb{Z}_4 | impossible (Proposition 3.3) | |
| 3 | \mathbb{Z}_{12} | \mathbb{Z}_{24} | impossible (Theorem 3.5) |
| 4 | \mathbb{Z}_{15} | impossible (Corollary 3.2) | impossible (Theorem 3.5) |

Table 2. Possibilities for $g(\Gamma(R))$ and $g(\Gamma_E(R))$, with examples.

Since $\text{ann } y$ is maximal in the set of annihilators of R , there is some m in $\text{ann } y$ but not in $\text{ann } w$. Note that $mw \neq 0$, but $\text{ann } mw \supseteq \{x, z, y\}$. Since $mw - y$ but $y^2 \neq 0$, $\text{ann } mw \neq \text{ann } y$. Also since mw is adjacent to both x and z , and x and z are not adjacent, $\text{ann } mw \neq \text{ann } x$ and $\text{ann } mw \neq \text{ann } z$. So we have the 3-cycles $x - y - mw - x$ and $z - y - mw - z$ that do not reduce under the equivalence relation. So $\Gamma_E(R)$ contains a 3-cycle and $g(\Gamma_E(R)) \neq 4$. \square

We summarize with Table 2, which shows all possible combinations of girths for $\Gamma(R)$ and $\Gamma_E(R)$. We illustrate the case (3, 3) with the graphs of the ring \mathbb{Z}_{24} , which does not have shrinking girth:



4. Cut-vertices

In this section, we examine the properties of cut-vertices of $\Gamma_E(R)$. Since $\Gamma_E(R)$ is connected, the vertex $[a]$ is a cut-vertex of $\Gamma_E(R)$ exactly when removing the vertex $[a]$ and its incident edges causes $\Gamma_E(R)$ to no longer be connected.

We begin with an interesting result concerning cut-vertices and ideals of the ring. The following theorem is very similar to [Axtell et al. 2009, Theorem 4.4], which deals with cut-vertices of the original zero-divisor graph $\Gamma(R)$.

Theorem 4.1. *If $[a]$ is a cut-vertex of $\Gamma_E(R)$, then $[a] \cup \{0\}$ forms an ideal of R .*

Proof. Let $[a]$ be a cut-vertex of $\Gamma_E(R)$ and let $[a]$ partition $\Gamma_E(R)$ into Γ_b and Γ_c . Let $[b] \in \Gamma_b$ with $[a] - [b]$ and $[c] \in \Gamma_c$ with $[a] - [c]$. Let $a_1, a_2 \in [a] \cup \{0\}$. Since $a_1 + a_2 \in \text{ann } b \cap \text{ann } c$, $a_1 + a_2 \in [a] \cup \{0\}$. If $r \in R$, then $c(ra) = r(ca) = 0$, so $ra \in \text{ann } c$. Similarly, $ra \in \text{ann } b$. So $ra \in \text{ann } b \cap \text{ann } c = [a] \cup \{0\}$. This shows that $[a] \cup \{0\}$ is an ideal of R . \square



Theorem 4.2. *If $[a]$ is a cut-vertex of $\Gamma_E(R)$, then $\text{ann } a$ is maximal in \mathfrak{F} .*

Proof. Let $[a]$ be a cut-vertex of $\Gamma_E(R)$, and let X and Y be mutually separated subgraphs of $\Gamma_E(R)$ with $V(X \cup Y) = V(\Gamma_E(R)) \setminus [a]$. Let $[x] \in X$ and $[y] \in Y$. Then for any $[x_1] \in X$ we have $y \in \text{ann } a \setminus \text{ann } x_1$, and for any $[y_1] \in Y$ we have $x \in \text{ann } a \setminus \text{ann } y_1$. Thus $\text{ann } a \not\subseteq \text{ann } x_1$ and $\text{ann } a \not\subseteq \text{ann } y_1$, and so $\text{ann } a$ is maximal in \mathfrak{F} . \square

The converse of this theorem does not hold. We may have $\text{ann } x$ maximal in \mathfrak{F} , yet not have $[x]$ be a cut-vertex. For example, here are two equivalence graphs, one on 6 vertices and one on 8, each with no cut vertex:

$$\Gamma_E(\mathbb{Z}_2[x, y, z] / (x^3, y^2, z^2, xy, xz))$$

$$\Gamma_E(\mathbb{Z}_2[x, y] / (x^4, xy, x^3 + y^2))$$



Both of these rings contains an annihilator ideal which maximal in \mathfrak{F} , and therefore an associated prime.

The next corollary follows immediately from Theorem 4.2:

Corollary 4.3. *If $[a]$ is a cut-vertex of $\Gamma_E(R)$, then $\text{ann } a$ is an associated prime.*

Theorem 4.4. *If $[a]$ is a cut-vertex of $\Gamma_E(R)$, then all other associated primes of $\Gamma_E(R)$ are contained in only one connected component of $\Gamma_E(R) \setminus [a]$.*

Proof. Suppose that X and Y are two mutually separated connected components of $\Gamma_E(R) \setminus [a]$, and that each contains an associated prime. By Lemma 1.1, these associated primes are adjacent, and so X and Y are connected, a contradiction. \square

Theorem 4.5. *If $\Gamma_E(R)$ has at least 2 cut-vertices, then it has diameter 3.*

Proof. Let $[a]$ and $[b]$ be cut-vertices of $\Gamma_E(R)$. Since $[a]$ is a cut-vertex, there is some $[x_a]$ such that any path connecting $[x_a]$ and $[b]$ must include $[a]$. Similarly, since $[b]$ is a cut-vertex, there is some $[x_b]$ such that any path connecting $[x_b]$ and $[a]$ must include $[b]$. Therefore any path from $[x_a]$ to $[x_b]$ must include both $[a]$ and $[b]$ and so $d([a], [b]) \geq 3$. Since $\Gamma_E(R)$ is connected, $\text{diam } \Gamma_E(R) = 3$. \square

Acknowledgements

This paper was written during the Research Experience for Undergraduates conducted at Brigham Young University in the summer of 2010. This research was funded by the National Science Foundation (DMS-0453421) and Brigham Young University. We would like to especially acknowledge the help of fellow REU students Cathryn Holm and Kaylee Kooiman, as well as TAs Chelsea Johnson and Donald Sampson.

References

- [Anderson and Livingston 1999] D. F. Anderson and P. S. Livingston, “The zero-divisor graph of a commutative ring”, *J. Algebra* **217**:2 (1999), 434–447. MR 2000e:13007 Zbl 0941.05062
- [Axtell et al. 2009] M. Axtell, J. Stickles, and W. Trampbachls, “Zero-divisor ideals and realizable zero-divisor graphs”, *Involve* **2**:1 (2009), 17–27. MR 2010b:13011 Zbl 1169.13301
- [Beck 1988] I. Beck, “Coloring of commutative rings”, *J. Algebra* **116**:1 (1988), 208–226. MR 89i:13006 Zbl 0654.13001
- [Mulay 2002] S. B. Mulay, “Cycles and symmetries of zero-divisors”, *Comm. Algebra* **30**:7 (2002), 3533–3558. MR 2003j:13007a Zbl 1087.13500
- [Spiroff and Wickham 2011] S. Spiroff and C. Wickham, “A zero divisor graph determined by equivalence classes of zero divisors”, *Comm. Algebra* **39**:7 (2011), 2338–2348. MR 2821714 Zbl 1225.13007

Received: 2011-01-19 Revised: 2011-08-02 Accepted: 2011-08-18

blakej2@hotmail.com

*Department of Mathematics, Utah Valley University,
Orem, UT 84058, United States*

martine@william.jewell.edu

*Department of Physics and Mathematics,
William Jewell College, Liberty, MO 64068, United States*

new4@tcnj.edu

*Department of Mathematics and Statistics,
The College of New Jersey, Ewing, NJ 08628, United States*

dane.skabelund@gmail.com

*Department of Mathematics, Brigham Young University,
Provo, UT 84602, United States*

involve

msp.berkeley.edu/involve

EDITORS

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

| | | | |
|----------------------|---|------------------------|---|
| John V. Baxley | Wake Forest University, NC, USA baxley@wfu.edu | Chi-Kwong Li | College of William and Mary, USA ckli@math.wm.edu |
| Arthur T. Benjamin | Harvey Mudd College, USA benjamin@hmc.edu | Robert B. Lund | Clemson University, USA lund@clemson.edu |
| Martin Bohner | Missouri U of Science and Technology, USA bohner@mst.edu | Gaven J. Martin | Massey University, New Zealand g.j.martin@massey.ac.nz |
| Nigel Boston | University of Wisconsin, USA boston@math.wisc.edu | Mary Meyer | Colorado State University, USA meyer@stat.colostate.edu |
| Amarjit S. Budhiraja | U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu | Emil Minchev | Ruse, Bulgaria eminchev@hotmail.com |
| Pietro Cerone | Victoria University, Australia pietro.cerone@vu.edu.au | Frank Morgan | Williams College, USA frank.morgan@williams.edu |
| Scott Chapman | Sam Houston State University, USA scott.chapman@shsu.edu | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir |
| Jem N. Corcoran | University of Colorado, USA corcoran@colorado.edu | Zuhair Nashed | University of Central Florida, USA znashed@mail.ucf.edu |
| Toka Diagana | Howard University, USA tdiagana@howard.edu | Ken Ono | Emory University, USA ono@mathcs.emory.edu |
| Michael Dorff | Brigham Young University, USA mdorff@math.byu.edu | Timothy E. O'Brien | Loyola University Chicago, USA tdorrie1@luc.edu |
| Sever S. Dragomir | Victoria University, Australia sever@matilda.vu.edu.au | Joseph O'Rourke | Smith College, USA orourke@cs.smith.edu |
| Behrouz Emamizadeh | The Petroleum Institute, UAE bemamizadeh@pi.ac.ae | Yuval Peres | Microsoft Research, USA peres@microsoft.com |
| Errin W. Fulp | Wake Forest University, USA fulp@wfu.edu | Y.-F. S. Pétermann | Université de Genève, Switzerland petermann@math.unige.ch |
| Joseph Gallian | University of Minnesota Duluth, USA jgallian@d.umn.edu | Robert J. Plemmons | Wake Forest University, USA rplemmons@wfu.edu |
| Stephan R. Garcia | Pomona College, USA stephan.garcia@pomona.edu | Carl B. Pomerance | Dartmouth College, USA carl.pomerance@dartmouth.edu |
| Ron Gould | Emory University, USA rg@mathcs.emory.edu | Vadim Ponomarenko | San Diego State University, USA vadim@sciences.sdsu.edu |
| Andrew Granville | Université Montréal, Canada andrew@dms.umontreal.ca | Bjorn Poonen | UC Berkeley, USA poonen@math.berkeley.edu |
| Jerrold Griggs | University of South Carolina, USA griggs@math.sc.edu | James Propp | U Mass Lowell, USA jpropp@cs.uml.edu |
| Ron Gould | Emory University, USA rg@mathcs.emory.edu | József H. Przytycki | George Washington University, USA przytyck@gwu.edu |
| Sat Gupta | U of North Carolina, Greensboro, USA sngupta@uncg.edu | Richard Rebarber | University of Nebraska, USA rrebarbe@math.unl.edu |
| Jim Haglund | University of Pennsylvania, USA jhaglund@math.upenn.edu | Robert W. Robinson | University of Georgia, USA rwr@cs.uga.edu |
| Johnny Henderson | Baylor University, USA johnny_henderson@baylor.edu | Filip Saidak | U of North Carolina, Greensboro, USA f_saidak@uncg.edu |
| Natalia Hritonenko | Prairie View A&M University, USA nahritonenko@pvamu.edu | James A. Sellers | Penn State University, USA sellersj@math.psu.edu |
| Charles R. Johnson | College of William and Mary, USA crjohnso@math.wm.edu | Andrew J. Sterge | Honorary Editor andy@ajsterge.com |
| Karen Kafadar | University of Colorado, USA karen.kafadar@cudenver.edu | Ann Trenk | Wellesley College, USA atrenk@wellesley.edu |
| K. B. Kulasekera | Clemson University, USA kk@ces.clemson.edu | Ravi Vakil | Stanford University, USA vakil@math.stanford.edu |
| Gerry Ladas | University of Rhode Island, USA gladas@math.uri.edu | Ram U. Verma | University of Toledo, USA verma99@msn.com |
| David Larson | Texas A&M University, USA larson@math.tamu.edu | John C. Wierman | Johns Hopkins University, USA wierman@jhu.edu |
| Suzanne Lenhart | University of Tennessee, USA lenhart@math.utk.edu | Michael E. Zieve | University of Michigan, USA zieve@umich.edu |

PRODUCTION

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor

Cover design: ©2008 Alex Scorpan

See inside back cover or <http://msp.berkeley.edu/involve> for submission instructions.

The subscription price for 2012 is US \$105/year for the electronic version, and \$145/year (+\$35 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94704-3840, USA.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Mathematical Sciences Publishers

involve

2012

vol. 5

no. 1

| | |
|---|----|
| Elliptic curves, eta-quotients and hypergeometric functions DAVID PATHAKJEE, ZEF ROSNBRICK AND EUGENE YOONG | 1 |
| Trapping light rays aperiodically with mirrors ZACHARY MITCHELL, GREGORY SIMON AND XUEYING ZHAO | 9 |
| A generalization of modular forms ADAM HAQUE | 15 |
| Induced subgraphs of Johnson graphs RAMIN NAIMI AND JEFFREY SHAW | 25 |
| Multiscale adaptively weighted least squares finite element methods for convection-dominated PDEs BRIDGET KRAYNIK, YIFEI SUN AND CHAD R. WESTPHAL | 39 |
| Diameter, girth and cut vertices of the graph of equivalence classes of zero-divisors BLAKE ALLEN, ERIN MARTIN, ERIC NEW AND DANE SKABELUND | 51 |
| Total positivity of a shuffle matrix AUDRA MCMILLAN | 61 |
| Betti numbers of order-preserving graph homomorphisms LAUREN GUERRA AND STEVEN KLEE | 67 |
| Permutation notations for the exceptional Weyl group F_4 PATRICIA CAHN, RUTH HAAS, ALOYSIUS G. HELMINCK, JUAN LI AND JEREMY SCHWARTZ | 81 |
| Progress towards counting D_5 quintic fields ERIC LARSON AND LARRY ROLEN | 91 |
| On supersingular elliptic curves and hypergeometric functions KEENAN MONKS | 99 |



1944-4176(2012)5:1;1-C