

# involve

a journal of mathematics

Ecological systems, nonlinear boundary conditions, and  
 $\Sigma$ -shaped bifurcation curves

Kathryn Ashley, Victoria Sincavage and Jerome Goddard II



# Ecological systems, nonlinear boundary conditions, and $\Sigma$ -shaped bifurcation curves

Kathryn Ashley, Victoria Sincavage and Jerome Goddard II

(Communicated by John Baxley)

We examine a one-dimensional reaction diffusion model with a weak Allee growth rate that appears in population dynamics. We combine grazing with a certain nonlinear boundary condition that models negative density dependent dispersal on the boundary and analyze the effects on the steady states. In particular, we study the bifurcation curve of positive steady states as the grazing parameter is varied. Our results are acquired through the adaptation of a quadrature method and Mathematica computations. Specifically, we computationally ascertain the existence of  $\Sigma$ -shaped bifurcation curves with several positive steady states for a certain range of the grazing parameter.

## 1. Introduction

Within population dynamics, the most accepted exemplar for modeling a designated population is the logistic equation

$$f(u) = u(a - bu), \quad (1-1)$$

which illustrates the inference that as a population burgeons, the per capita growth rate

$$\tilde{f}(u) = a - bu \quad (1-2)$$

of that population declines linearly. Yet empirically several authors have witnessed that at lower population densities, the per capita growth rate initially increases (see [Allee 1938; Dennis 1989; Lewis and Kareiva 1993; Shi and Shivaji 2006]). This phenomenon is known in the literature as the Allee effect [1938]. Since the logistic growth model does not compensate for the initial increase, a model of the Allee effect must be implemented to account for this phenomenon.

---

*MSC2010:* 34B08, 34B18.

*Keywords:* nonlinear boundary conditions, weak Allee effect, positive solutions.

Research supported by National Science Foundation grant DMS 0852032.

The Allee effect can be either strong, in which the per capita growth rate is initially negative, or weak, in which the per capita growth rate is initially positive. The Allee effect is generally modeled in the literature via quadratic per capita growth rate functions of the population density. In this case, the analysis is more difficult since the per capita growth rate is not linear or even nonincreasing. As a contrast with (1-1), a weak Allee effect has been modeled as

$$f(u) = u(u + 1)(b - u), \quad (1-3)$$

where  $b > 1$ .

By analyzing additional factors that can influence a population, such as grazing or harvesting, a better understanding can be had of the dynamics of the population. Therefore, through the inclusion of an extra term to account for these factors, specifically grazing, a more precise model can be obtained. Grazing can be considered as a category of natural predation, for example, when an owl preys upon the surrounding rodent population. The term  $cu^2/(1 + u^2)$  is commonly employed to model grazing of a population (see [Causey et al. 2010; Lee et al. 2011; Poole et al. 2012; van Nes and Scheffer 2005]).

Density dependent dispersal, or more specifically density dependent emigration, describes a situation in which the dispersal/emigration of individuals living within a patch is based on the population density, in our case, on the habitat border. A positive density dependent emigration characterizes a case where individuals have a tendency to leave if the population density is large and a tendency to stay if the population density is small. On the contrary, a negative density dependent emigration represents a case where individuals have a tendency to stay if the population density is large and a tendency to leave if the population density is small.

Initially and intuitively it was believed that the majority of animals exhibit positive density dependent dispersal. However, recent studies of several animals, including the bighorn sheep, roe deer, house mouse, prairie vole, European badger, and the Glanville fritillary butterfly *Melitaea cinxia*, have proven otherwise (see [Kuussaari et al. 1996; Matthysen 2005]). In the literature, several factors have been suggested as a cause of negative density dependent dispersal, including: niche breadth, increased predator abundance, and, in particular, conspecific attraction (see [Kuussaari et al. 1996; Matthysen 2005]). Conspecific attraction most simply means that there is a predisposition of individuals within a population to become enticed to areas where there are more conspecifics.

Cantrel and Cosner proposed the following nonlinear boundary condition to model conspecific attraction on the boundary of a patch (see [Cantrell and Cosner 2003; 2007; Goddard et al. 2010a; 2010b; 2011a; 2011b; 2012]):

$$d(\nabla u \cdot \eta)\alpha(x, u) + [1 - \alpha(x, u)]u = 0; \quad \partial\Omega, \quad (1-4)$$

where  $\alpha : \bar{\Omega} \times [0, \infty) \rightarrow [0, 1]$  is  $C^1$  and nondecreasing,  $d > 0$  is the diffusion parameter,  $\nabla u \cdot \eta$  is the outward normal derivative, and  $\Omega \subset \mathbb{R}^n$  ( $n \geq 1$ ) is a smooth bounded domain. The  $\alpha(x, u)$ 's of biological importance are of the form

$$\alpha(x, u) = \alpha(u) = \frac{u}{u + g(u)}, \tag{1-5}$$

where  $g : [0, \infty) \rightarrow [\delta, \infty)$  is a  $C^1$  function,  $\delta > 0$ , and  $g(u)/u \rightarrow 0$  as  $u \rightarrow \infty$ . Here,  $\alpha(u)$  represents the fraction of the population that stays on the boundary when reached. Notice that if  $\alpha(u) \equiv 0$  then (1-4) becomes the Dirichlet boundary condition ( $u = 0; \partial\Omega$ ), and if  $\alpha(u) \equiv 1$  then (1-4) becomes the Neumann boundary condition ( $\nabla u \cdot \eta = 0; \partial\Omega$ ). In terms of this paper, we consider the case when  $g(u) \equiv d$ , where  $d > 0$  is the diffusion parameter.

Our purpose is to analyze the effects of grazing in combination with a weak Allee effect and the nonlinear boundary conditions (1-4) on the steady state solutions of a reaction diffusion model. In particular, we study the one-dimensional case when  $n = 1$  and  $\Omega = (0, 1)$ :

$$u_t = \frac{1}{\lambda} u_{xx} + u \tilde{f}(u) - \frac{cu^2}{1 + u^2}; \quad (0, 1), \tag{1-6}$$

with nonlinear boundary conditions, namely

$$\begin{aligned} -u'' &= \lambda \left[ u \tilde{f}(u) - \frac{cu^2}{1 + u^2} \right] = \lambda f(u); \quad (0, 1), \\ u(0) \left[ -\frac{1}{\lambda} u'(0) + \frac{1}{\lambda} \right] &= 0, \\ u(1) \left[ \frac{1}{\lambda} u'(1) + \frac{1}{\lambda} \right] &= 0, \end{aligned} \tag{1-7}$$

where  $u$  represents the population density,  $\tilde{f}(u)$  represents the per capita growth rate,  $\lambda = 1/d$  and  $d > 0$  represents the diffusion coefficient, and  $c \geq 0$  represents the maximum grazing rate. Notice that the boundary conditions found in (1-7) can be separated into the following four cases:


$$-u'' = \lambda f(u); (0, 1), \quad u(0) = 0, \quad u(1) = 0, \tag{1-8}$$

$$-u'' = \lambda f(u); (0, 1), \quad u(0) = 0, \quad u'(1) = -1, \tag{1-9}$$

$$-u'' = \lambda f(u); (0, 1), \quad u'(0) = 1, \quad u(1) = 0, \tag{1-10}$$

$$-u'' = \lambda f(u); (0, 1), \quad u'(0) = 1, \quad u'(1) = -1. \tag{1-11}$$


Thus, the positive solutions of (1-8)–(1-11) are the positive solutions of (1-7). Further, it is clear that if  $u(x)$  is a positive solution of (1-9), then  $v(x) = u(1 - x)$  also satisfies (1-10). Thus, it suffices to only consider (1-8), (1-9), and (1-11).



**Figure 1.** S-shaped bifurcation curve.

Prior studies have gathered information and analyzed the positive solutions to both strong and weak Allee problems. Additionally, the analysis of the positive solutions to the combination of grazing and the Allee effect has also been made; however, to the best of our understanding no analysis has been made in regards to the Allee effect with grazing and nonlinear boundary conditions. In the case when  $\alpha(u) \equiv 0$ , (1-8) has a rich history. For the logistic case with Dirichlet boundary conditions, Lee, Sasi, and Shivaji proved the existence of an S-shaped bifurcation curve in one dimension, as well as higher dimensions for a certain range of the grazing parameter [Lee et al. 2011]. Regarding the one-dimensional weak Allee effect model with Dirichlet boundary conditions, Poole, Roberson, and Stephenson showed the existence of an S-shaped bifurcation curve, resembling Figure 1, both computationally and analytically for certain parameter ranges [Poole et al. 2012]. In particular, our focus is to further examine the structure of positive solutions of (1-7) when the nonlinear boundary conditions (1-4) are satisfied for the range of the parameters where Poole et al. [2012] showed the existence of an S-shaped bifurcation curve of positive solutions. Computationally, we show the existence of  $\Sigma$ -shaped bifurcation curves as exemplified in Figure 2.

We employ and adapt the quadrature method first developed by Laetsch [1970] to study the structure of positive solutions of (1-7). First, some important preliminaries will be presented in Section 2, followed by a discussion of applying and adapting the quadrature method for the specific cases (1-8), (1-9), and (1-11). In Section 6, we provide the complete evolution of the bifurcation curve of positive solutions of (1-7), followed by analytical results confirming some of our observations in Section 7.



**Figure 2.**  $\Sigma$ -shaped bifurcation curves.

### 2. Preliminaries


We examine the combination of the weak Allee effect and grazing in the subsequent reaction term:

$$\begin{aligned}
 f(u) &= u(u + 1)(b - u) - \frac{cu^2}{1 + u^2} \quad \text{for } b > 1, c \geq 0 \\
 &= \frac{u(u + 1)(b - u)(1 + u^2) - cu^2}{1 + u^2}.
 \end{aligned}$$


Through observation it is apparent that the numerator of  $f(u)$  can be written as a fifth-degree polynomial. Regardless of any specific values for  $b$  and  $c$ , by analyzing the roots of  $f(u)$  the existence of three roots — a negative root, a positive root, and a root at  $u = 0$  — can be determined. As  $c$  is varied the remaining three roots alternate between imaginary and real values. For the purpose of this paper, denote  $\sigma = \sigma(b, c)$  as the smallest positive root of  $f(u)$ . Also, allow  $\sigma_0 = \sigma_0(b, c)$  and  $\sigma_1 = \sigma_1(b, c)$  to represent the remaining roots. Regardless of the value of  $c > 0$ , for certain values of  $b$ , specifically  $b \in (1, b_0)$  (some  $b_0 > 0$ ), there exists only one positive real root of  $f(u)$  represented by  $\sigma$ .

**Remark 1.** Through calculation and the use of Mathematica, it is estimated that  $b_0 \approx 2.852$ .

Specifically, when  $b \in (b_0, \infty)$ , it has been determined that the shape of  $f(u)$  changes when  $c$  is varied. Note when  $c \in [0, c_0)$  (some  $c_0 = c_0(b) > 0$ ), there exists exactly one positive real root denoted by  $\sigma(b, c)$ . Figure 3 depicts this case. The shape of  $f(u)$  is modified as  $c$  becomes larger. Specifically, when  $c \in [c_0, c_1)$  (some  $c_1 = c_1(b) \in (c_0, \infty)$ ),  $f(u)$  has 3 positive real roots, namely  $\sigma(b, c)$ ,  $\sigma_0(b, c)$ , and  $\sigma_1(b, c)$ , as depicted in Figure 4. For  $c > c_1$ ,  $f(u)$  is shifted downward resulting in



**Figure 3.** Graph of  $f(u)$  for  $b > b_0$  and  $c \in [0, c_0)$ .



**Figure 4.** Graph of  $f(u)$  for  $b > b_0$  and  $c \in [c_0, c_1)$ .

exactly one positive real root  $\sigma(b, c)$ , meaning  $\sigma_0(b, c)$  and  $\sigma_1(b, c)$  are imaginary roots. This particular case is illustrated in Figure 5.


In the preceding cases the structure of the positive solutions of (1-7) varies. As our primary interest is the structure of positive solutions for the range of parameters where Poole et al. [2012] showed the existence of S-shaped bifurcation curves, we focus on the case when  $c \in [0, c_0)$ .

### 3. Quadrature method for (1-8)


For completeness, we reestablish the results obtained through the quadrature method actualized by Laetsch [1970] and Brown, Ibrahim, and Shivaji [Brown et al. 1981]. Additionally we recapitulate the subsequent boundary value problem analyzed by Poole et al. [2012] for positive solutions:

$$-u''(x) = \lambda f(u(x)); x \in (0, 1), \quad u(0) = 0, \quad u(1) = 0, \quad (3-1)$$

where  $f : [0, \infty) \rightarrow (0, \infty)$  is a  $C^1$  function. Clearly, a positive solution of (3-1) must resemble Figure 6.



**Figure 5.** Graph of  $f(u)$  for  $b > b_0$  and  $c > c_1$ .



**Figure 6.** Graph of a typical positive solution of (3-1).

**Theorem 3.1** [Brown et al. 1981; Laetsch 1970]. *Suppose  $u(x)$  is a positive solution to (3-1) with  $\|u\|_\infty = \rho = u(\frac{1}{2})$ , where  $\rho > 0$ . Such a solution to (3-1) exists if and only if*

$$G_1(\rho) = \sqrt{2} \int_0^\rho \frac{ds}{\sqrt{F(\rho) - F(s)}} = \sqrt{\lambda}, \tag{3-2}$$

where  $F(x) = \int_0^x f(s) ds$ .

*Proof.* ( $\Rightarrow$ ) Recognizing that (3-1) is an autonomous differential equation, we see that if  $u$  is a positive solution to (3-1) with  $u'(x_0) = 0$  for a particular  $x_0 \in (0, 1)$ , then  $m(x) = u(x_0 + x)$  and  $n(x) = u(x_0 - x)$  both satiate the initial value problem

$$-k''(x) = \lambda f(k(x)), \quad k(0) = u(x_0), \quad k'(0) = 0, \tag{3-3}$$



where  $x \in [0, l)$  and  $l = \min\{x_0, 1 - x_0\}$ . Using Picard's existence and uniqueness theorem, we have that  $u(x_0 + x) \equiv u(x_0 - x)$  for all  $x \in [0, l)$  and thus  $u(x)$  is symmetric about  $x = \frac{1}{2}$ , which is notably where  $u(x)$  achieves its maximum.

By multiplying the differential equation in (3-1) by  $u'(x)$ , we have

$$-\left[\frac{[u'(x)]^2}{2}\right]' = [\lambda F(u(x))]' \quad (3-4)$$

Integration of both sides of (3-4) gives

$$\frac{u'(x)}{\sqrt{F(\rho) - F(u(x))}} = \sqrt{2\lambda}; \quad x \in [0, \frac{1}{2}). \quad (3-5)$$

By integrating a second time and using the fact that  $u(0) = 0$ , we have

$$\int_0^{u(x)} \frac{dt}{\sqrt{F(\rho) - F(t)}} = \sqrt{2\lambda}x; \quad x \in [0, \frac{1}{2}]. \quad (3-6)$$

Substituting  $x = \frac{1}{2}$  and utilizing  $u(\frac{1}{2}) = \rho$ , (3-6) can be written as

$$G_1(\rho) = \sqrt{2} \int_0^\rho \frac{dt}{\sqrt{F(\rho) - F(t)}} = \sqrt{\lambda}. \quad (3-7)$$

Therefore, if  $u(x)$  is a positive solution to (3-1) where  $\|u\|_\infty = \rho$ , then  $\rho$  must fulfill  $G_1(\rho) = \sqrt{\lambda}$ .

( $\Leftrightarrow$ ) Assume  $G_1(\rho) = \sqrt{\lambda}$  for  $\rho > 0$ . Now define a function  $u : [0, \frac{1}{2}] \rightarrow [0, \infty)$  by

$$\int_0^{u(x)} \frac{dt}{\sqrt{F(\rho) - F(t)}} = \sqrt{2\lambda}x; \quad x \in [0, \frac{1}{2}]. \quad (3-8)$$

We now show that  $u(x)$  satisfies (3-1). Notice that  $u(x)$  is well defined and via the implicit function theorem also twice differentiable. Hence, differentiating (3-8) yields

$$u'(x) = \sqrt{2\lambda[F(\rho) - F(u(x))]}.$$


By differentiating a second time we obtain

$$-u''(x) = \lambda f(u(x)).$$

In addition, it is clear that  $u(0) = 0$ . By defining  $u(x)$  as a symmetric solution on  $[0, 1]$ , it is apparent that  $u(x)$  is a positive solution to (3-1) with  $\|u\|_\infty = \rho$ .  $\square$

It is important to discern that  $G_1(\rho)$  is well defined and the improper integral is convergent. To that end, we state an important remark.

**Remark 2.** The improper integral in (3-7) is both well defined and convergent for  $\rho$ -values that fulfill:



**Figure 7.** Graph of a typical positive solution of (4-1).

- (1)  $f(\rho) > 0$ ;
- (2)  $F(\rho) > F(s)$  for all  $s \in [0, \rho)$ .

Notice that from Figure 3 if  $c \in [0, c_0)$ , then both (1) and (2) will be satisfied for all  $\rho \in (0, \sigma(b, c))$ . We close this section by recalling an important result from Brown et al.

**Theorem 3.2** [Brown et al. 1981].  $G_1(\rho)$  is both differentiable and continuous on the defined set  $T = \{\rho > 0 \mid f(\rho) > 0 \text{ and } F(\rho) - F(s) > 0 \text{ for all } s \in [0, \rho)\}$  where

$$G'_1(\rho) = \sqrt{2} \int_0^1 \frac{H(\rho) - H(\rho v)}{[F(\rho) - F(\rho v)]^{3/2}} dv,$$

in which

$$H(s) = F(s) - \frac{s}{2} f(s).$$

#### 4. Quadrature method for (1-9)

In this section, we adapt the quadrature method to analyze the structure of positive solutions of (1-9):

$$-u'' = \lambda \left[ u(u+1)(b-u) - \frac{cu^2}{(u^2+1)} \right]; \quad (0, 1), \quad u(0) = 0, \quad u'(1) = -1. \quad (4-1)$$

Define

$$f(u) = \left[ u(u+1)(b-u) - \frac{cu^2}{(u^2+1)} \right] \quad \text{and} \quad F(x) = \int_0^x f(s) ds.$$

It is apparent that a positive solution of (4-1) must resemble Figure 7.

Assume  $u(x)$  is a positive solution to (4-1) with  $\|u\|_\infty = \rho$  and  $u(1) = q$  for  $q \in [0, \rho)$ . By multiplying the differential equation in (4-1) by  $u'(x)$  we obtain

$$-\left[\frac{[u'(x)]^2}{2}\right]' = [\lambda F(u(x))]' \quad (4-2)$$

Integrating both sides of (4-2) yields

$$\frac{-(u'(x))^2}{2} = \lambda F(u(x)) + C. \quad (4-3)$$

Recalling that  $u(x_0) = \rho$  and  $u'(x_0) = 0$ , (4-3) becomes

$$C = -\lambda F(\rho). \quad (4-4)$$

Similarly, using  $u(1) = q$  and  $u'(1) = -1$ , (4-3) is utilized to determine a second value for  $C$ ,

$$C = -\frac{1}{2} - \lambda F(q). \quad (4-5)$$

Combining (4-4) and (4-5) gives

$$\sqrt{2\lambda} = \frac{1}{\sqrt{F(\rho) - F(q)}}. \quad (4-6)$$

In utilizing the  $C$ -value from (4-4) while solving for  $u'(x)$ , (4-3) becomes

$$u'(x) = \sqrt{2\lambda[F(\rho) - F(u(x))]}; \quad x \in [0, x_0], \quad (4-7)$$

$$u'(x) = -\sqrt{2\lambda[F(\rho) - F(u(x))]}; \quad x \in [x_0, 1]. \quad (4-8)$$

Rearranging (4-7) and (4-8) gives

$$\frac{u'(x)}{\sqrt{F(\rho) - F(u(x))}} = \sqrt{2\lambda}; \quad x \in [0, x_0], \quad (4-9)$$

$$\frac{u'(x)}{\sqrt{F(\rho) - F(u(x))}} = -\sqrt{2\lambda}; \quad x \in (x_0, 1]. \quad (4-10)$$

Integration of (4-9) from 0 to  $x$  and (4-10) from  $x_0$  to  $x$  yields

$$\int_0^x \frac{u'(x)}{\sqrt{F(\rho) - F(u(x))}} = \int_0^x \sqrt{2\lambda}; \quad x \in [0, x_0], \quad (4-11)$$

$$\int_{x_0}^x \frac{u'(x)}{\sqrt{F(\rho) - F(u(x))}} = \int_{x_0}^x -\sqrt{2\lambda}; \quad x \in (x_0, 1]. \quad (4-12)$$

Using a change of variables and recalling  $u(0) = 0$  and  $u(x_0) = \rho$  we obtain

$$\int_0^{u(x)} \frac{dw}{\sqrt{F(\rho) - F(w)}} = \sqrt{2\lambda}x; \quad x \in [0, x_0], \quad (4-13)$$

$$\int_\rho^{u(x)} \frac{dw}{\sqrt{F(\rho) - F(w)}} = -\sqrt{2\lambda}(x - x_0); \quad x \in [x_0, 1]. \quad (4-14)$$

By substituting  $x = x_0$  into (4-13) and  $x = 1$  into (4-14) we obtain

$$\int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} = \sqrt{2\lambda}x_0, \quad (4-15)$$

$$\int_\rho^q \frac{dw}{\sqrt{F(\rho) - F(w)}} = -\sqrt{2\lambda}(1 - x_0). \quad (4-16)$$

Subtracting (4-16) from (4-15) we have

$$\sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}} = \sqrt{\lambda}. \quad (4-17)$$

By synthesizing (4-6) with (4-17) we denote

$$\begin{aligned} \tilde{G}_2(\rho, q) = \\ \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}}. \end{aligned} \quad (4-18)$$

By Remark 2, the improper integral in  $\tilde{G}_2(\rho, q)$  exists and is convergent for  $\rho$  in  $(0, \sigma(b, c))$ . Also, for a given  $\rho \in (0, \sigma(b, c))$  Picard's existence and uniqueness theorem guarantees that the corresponding  $q = u(1) \in [0, \rho]$  must be unique. If for each  $\rho \in (0, \sigma(b, c))$  there exists a unique  $q(\rho) \in [0, \rho]$  where  $\tilde{G}_2(\rho, q(\rho)) = 0$ , then there exists a unique  $\lambda \in (0, \infty)$  such that

$$\begin{aligned} \sqrt{2} \int_0^\rho \frac{ds}{\sqrt{F(\rho) - F(s)}} - \frac{1}{\sqrt{2}} \int_0^{q(\rho)} \frac{ds}{\sqrt{F(\rho) - F(s)}} \\ = \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q(\rho))}} = \sqrt{\lambda} \end{aligned} \quad (4-19)$$

will be satisfied. Therefore it is imperative to examine the existence and uniqueness of such a  $q = q(\rho)$ . Hence, we recall and prove Lemma 1, adapted from [Goddard et al. 2010a], which outlines necessary properties of  $\tilde{G}_2(\rho, q)$ .

**Lemma 1** [Goddard et al. 2010a]. *If  $\rho \in (0, \sigma(b, c))$  then:*

- (1)  $\tilde{G}_2(\rho, q) \rightarrow -\infty$  as  $q \rightarrow \rho^-$  for fixed  $\rho \in (0, \sigma(b, c))$ .
- (2)  $[\tilde{G}_2]_q < 0$  for every  $q \in [0, \rho]$  and fixed  $\rho \in (0, \sigma(b, c))$ .
- (3)  $\tilde{G}_2(\rho, 0) \rightarrow \infty$  when  $\rho \rightarrow \sigma(b, c)^-$ .

(4)  $\tilde{G}_2(\rho, 0) \rightarrow -\infty$  when  $\rho \rightarrow 0^+$ .

*Proof.* (1) Accomplished via the mean value theorem and the fact that  $F(u)$  is an increasing function on  $(0, \sigma(b, c))$ .

(2) Let  $\rho \in (0, \sigma(b, c))$ . Thus

$$[\tilde{G}_2(\rho, q)]_q = -\frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} - \frac{f(q)}{2\sqrt{2}[F(\rho) - F(q)]^{\frac{3}{2}}} < 0$$

for all  $q \in [0, \rho)$ , since  $f(s) > 0$  for  $s \in (0, \sigma(b, c))$ .

(3) For every  $\rho \in (0, \sigma(b, c))$ , we have


$$\tilde{G}_2(\rho, 0) = \sqrt{2} \int_0^\rho \frac{ds}{\sqrt{F(\rho) - F(s)}} - \frac{1}{\sqrt{2}\sqrt{F(\rho)}} = G_1(\rho) - \frac{1}{\sqrt{2}\sqrt{F(\rho)}}. \tag{4-20}$$

Laetsch [1970] showed that  $G_1(\rho) \rightarrow \infty$  as  $\rho \rightarrow \sigma(b, c)^-$ . This implies that  $\tilde{G}_2(\rho, 0) \rightarrow \infty$  when  $\rho \rightarrow \sigma(b, c)^-$ .


(4) Ascertained via the mean value theorem and the monotonicity of  $F(u)$  on  $(0, \sigma(b, c))$ . □

According to Lemma 1,  $\tilde{G}_2(\rho, q)$  must resemble Figure 8, whereas Figures 9 and 10 illustrate  $\tilde{G}_2(\rho, 0)$ . Noteworthy from Lemma 1, if  $\tilde{G}_2(\rho, 0) \geq 0$  then there exists a unique  $q(\rho) \in [0, \rho)$  wherefore  $\tilde{G}_2(\rho, q(\rho)) = 0$ . We conjecture as a result of our computations that there is a unique  $\rho^* = \rho^*(b, c) > 0$  wherefore if  $\rho \geq \rho^*$ , then  $\tilde{G}_2(\rho, 0) \geq 0$ . Also if  $\rho < \rho^*$  then  $\tilde{G}_2(\rho, 0) < 0$ . So, for all  $\rho \in [\rho^*, \infty)$  there exists a unique  $q = q(\rho) \in [0, \rho)$  where  $\tilde{G}_2(\rho, q(\rho)) = 0$ . In this case, we have


$$G_2(\rho, q(\rho)) = \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} = \sqrt{\lambda}. \tag{4-21}$$



**Figure 8.** Graph of  $\tilde{G}_2(\rho, q)$ .



**Figure 9.** Graph of  $\tilde{G}_2(\rho, 0)$  when  $b = 10$  and  $c = 0$ .



**Figure 10.** Graph of  $\tilde{G}_2(\rho, 0)$  when  $b = 10$  and  $c = 33$ .

We now state and prove the main theorem of the section.

**Theorem 4.1.** *The function  $u(x)$  is a positive solution to (4-1) with*

$$\|u\|_\infty = \rho \in S(b, c) := [\rho^*(b, c), \sigma(b, c))$$

*if and only if*

$$G_2(\rho, q(\rho)) = \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} = \sqrt{\lambda}$$

*for a positive  $\lambda$  for which  $q = q(\rho) \in [0, \rho)$  is the unique solution of*

$$\begin{aligned} & \tilde{G}_2(\rho, q(\rho)) \\ &= \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} = 0. \end{aligned}$$

*Proof.* ( $\Rightarrow$ ) Accomplished in the above analysis.

( $\Leftarrow$ ) Assume that there exist  $\lambda \in (0, \infty)$  and  $\rho \in S(b, c)$  wherefore  $G_2(\rho, q(\rho)) = \sqrt{\lambda}$  in which the unique solution of  $\tilde{G}_2(\rho, q(\rho)) = 0$  is  $q(\rho) \in [0, \rho]$ . Define

$$u(x) : [0, 1] \rightarrow \mathbb{R}$$

via

$$\int_0^{u(x)} \frac{ds}{\sqrt{F(\rho) - F(s)}} = \sqrt{2\lambda}x; \quad x \in [0, x_0], \quad (4-22)$$

$$\int_\rho^{u(x)} \frac{ds}{\sqrt{F(\rho) - F(s)}} = -\sqrt{2\lambda}(x - x_0); \quad x \in [x_0, 1]. \quad (4-23)$$

Now, we will exhibit  $u(x)$  as a positive solution to (4-1). Note that  $u(x)$  has a turning point at  $x_0$  denoted by

$$x_0 = \frac{1}{\sqrt{2\lambda}} \int_0^\rho \frac{ds}{\sqrt{F(\rho) - F(s)}}. \quad (4-24)$$

For the given  $\lambda > 0$ , it is apparent that

$$\frac{1}{\sqrt{2\lambda}} \int_0^{u(x)} \frac{ds}{\sqrt{F(\rho) - F(s)}} \quad (4-25)$$

is both a differentiable function of  $u$  and an increasing function ranging from 0 to  $x_0$  when  $u$  takes on the values from 0 to  $\rho$ . Therefore, for each  $x \in [0, x_0]$  there is a unique  $u(x)$  wherefore

$$\int_0^{u(x)} \frac{ds}{\sqrt{F(\rho) - F(s)}} = \sqrt{2\lambda}x. \quad (4-26)$$

The implicit function theorem gives that  $u(x)$  is a twice-differentiable function with respect to  $x$ . Differentiating (4-26) with respect to  $x$  gives

$$u'(x) = \sqrt{2\lambda[F(\rho) - F(u(x))]}; \quad x \in [0, x_0]. \quad (4-27)$$

Through a similar argument,

$$u'(x) = -\sqrt{2\lambda[F(\rho) - F(u(x))]}; \quad x \in [x_0, 1]. \quad (4-28)$$

By utilizing (4-27) and (4-28) we obtain

$$\frac{[u'(x)]^2}{2} = \lambda[F(\rho) - F(u(x))]; \quad x \in [0, 1]. \quad (4-29)$$

Through differentiation of (4-29) we have

$$-u''u' = \lambda f(u)u'; \quad x \in (0, 1), \quad (4-30)$$

which can be rewritten as

$$-u'' = \lambda f(u); \quad x \in (0, 1). \tag{4-31}$$

Thus, we have proved that  $u(x)$  satisfies the differential equation in (4-1). Now, we show that  $u(x)$  satisfies the boundary value conditions in (4-1); however, it is apparent that  $u(0) = 0$ . Additionally, using  $G_2(\rho, q(\rho)) = \sqrt{\lambda}$ , we ascertain

$$\sqrt{F(\rho) - F(q)} = \frac{1}{\sqrt{2\lambda}}. \tag{4-32}$$

Substitution of  $x = 1$  in (4-28) yields

$$u'(1) = -\sqrt{2\lambda}\sqrt{F(\rho) - F(q)}. \tag{4-33}$$

When (4-32) and (4-33) are synthesized we obtain

$$u'(1) = -1. \tag{4-34}$$

Therefore, the boundary conditions in (4-1) are satisfied by  $u(x)$ . □

### 5. Quadrature method for (1-11)

Further extension of the quadrature method is performed in this section to analyze the structure of positive solutions of (1-11):

$$-u'' = \lambda \left[ u(u+1)(b-u) - \frac{cu^2}{(u^2+1)} \right]; \quad (0, 1), \quad u'(0) = 1, \quad u'(1) = -1. \tag{5-1}$$

Define

$$f(u) = \left[ u(u+1)(b-u) - \frac{cu^2}{(u^2+1)} \right] \quad \text{and} \quad F(x) = \int_0^x f(s) ds.$$

Clearly, a positive solution of (5-1) must resemble Figure 11, where  $\|u\|_\infty = \rho$ ,  $\rho \in (0, \infty)$ ,  $q = u(0) = u(1)$ , and  $q \in [0, \rho)$ . Through a similar argument as in Section 4, we articulate the main theorem of this section.


**Theorem 5.1.** *The function  $u(x)$  is a positive solution of (5-1) with*

$$\|u\|_\infty = \rho \in S(b, c) = [\rho^*(b, c), \sigma(b, c)]$$

*if and only if*

$$G_3(\rho, q(\rho)) = \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} = \sqrt{\lambda}, \tag{5-2}$$





**Figure 11.** Graph of a typical positive solution of (5-1).

for which  $q = q(\rho) \in [0, \rho)$  is the unique solution of

$$\begin{aligned} &\tilde{G}_3(\rho, q(\rho)) \\ &= \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \sqrt{2} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} = 0. \end{aligned}$$

### 6. Computational results


Within this section we exhibit the complete evolution of the bifurcation curve of positive solutions of (1-7) for  $c \in [0, c_0(b))$ . The results for (1-8) are reestablished via Mathematica computations and by recalling Theorem 3.1. For (1-9) and (1-11), we recall Theorems 4.1 and 5.1 and utilize a standard root-finding algorithm to find the unique  $\rho^*(b, c) > 0$ . Then for  $\rho \in [\rho^*(b, c), \sigma(b, c))$  we employ a root-finding algorithm to find the corresponding unique  $q(\rho)$ , which is delineated in Theorems 4.1 and 5.1. These diagrams were acquired via Mathematica for a single  $b$ -value as  $c$ -values are varied. If  $b \in (b_0, \infty)$  then there exist

$$0 < c_0^* < c_1^* < c_2^* < c_3^* < c_4^* < c_5^* < c_6^* < c_7^* < c_0(b)$$

such that we have the following cases. In the subsequent figures, (1-8) is represented in black, cases (1-9) and (1-10) in red, and (1-11) in blue.

**Case 1.** If  $c \in [0, c_0^*)$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, 3, 4$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in [\lambda_0, \infty)$ , then (1-7) has exactly 4 positive solutions;



**Figure 12.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 0$  (Case 1).

- $\lambda \in (\lambda_1, \lambda_0)$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_1]$ , then (1-7) has exactly 8 positive solutions.

Figure 12 illustrates Case 1.


**Case 2.** If  $c \in [c_0, c_1)$  (some  $c_1(b) > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 6$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in [\lambda_0, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda \in (\lambda_1, \lambda_5)$  and  $\lambda \in (\lambda_6, \lambda_0)$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$ ,  $\lambda = \lambda_5$ , and  $\lambda = \lambda_6$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$  and  $\lambda \in (\lambda_5, \lambda_6)$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_1]$ , then (1-7) has exactly 8 positive solutions.

Figure 13 illustrates Case 2.

**Case 3.** If  $c = c_1$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 5$  such that if


- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;



**Figure 13.**  $\rho$  versus  $\lambda$  (top) and cross-section (bottom) for  $b = 10$  and  $c = 8.97$  (Case 2).

- $\lambda = \lambda_3$  and  $\lambda \in [\lambda_0, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda \in (\lambda_1, \lambda_5)$  and  $\lambda = \lambda_0$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$ ,  $\lambda = \lambda_5$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$  and  $\lambda \in (\lambda_5, \lambda_0)$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_1]$ , then (1-7) has exactly 8 positive solutions.


Figure 14 illustrates Case 3.



**Figure 14.**  $\rho$  versus  $\lambda$  (top) and cross-section (bottom) for  $b = 10$  and  $c = 8.972$  (Case 3).

**Case 4.** If  $c \in (c_1, c_2)$  (some  $c_2(b) > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 6$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_6, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda \in (\lambda_1, \lambda_5)$  and  $\lambda = \lambda_6$ , then (1-7) has exactly 5 positive solutions;




**Figure 15.**  $\rho$  versus  $\lambda$  (top) and cross-section (bottom) for  $b = 10$  and  $c = 8.99$  (Case 4).

- $\lambda \in (\lambda_3, \lambda_4)$ ,  $\lambda \in [\lambda_0, \lambda_6)$ , and  $\lambda = \lambda_5$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$  and  $\lambda \in (\lambda_5, \lambda_0)$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_1]$ , then (1-7) has exactly 8 positive solutions.

Figure 15 illustrates Case 4.

**Case 5.** If  $c = c_2$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 5$  such that if


- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;



**Figure 16.**  $\rho$  versus  $\lambda$  (top) and cross-section (bottom) for  $b = 10$  and  $c = 9$  (Case 5).

- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_5, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda \in (\lambda_1, \lambda_0]$  and  $\lambda = \lambda_5$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$  and  $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_1]$ , then (1-7) has exactly 8 positive solutions.

Figure 16 illustrates Case 5.



**Figure 17.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 18$  (Case 6).


**Case 6.** If  $c \in (c_2, c_3]$  (some  $c_3(b) > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 5$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_5, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_5$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_1, \lambda_5)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0]$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_1]$ , then (1-7) has exactly 9 positive solutions.


Figure 17 illustrates Case 6.

**Case 7.** If  $c \in (c_3, c_4)$  (some  $c_4 > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 7$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_7, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_7$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_1, \lambda_7)$ , then (1-7) has exactly 6 positive solutions;



**Figure 18.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 26$  (Case 7).



**Figure 19.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 27.3$  (Case 8).

- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0]$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5), (\lambda_6, \lambda_1]$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_5, \lambda = \lambda_6$ , then (1-7) has exactly 11 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6)$ , then (1-7) has exactly 13 positive solutions.

Figure 18 illustrates Case 7. Notice that the red curve has become  $\Sigma$ -shaped and this shape persists through  $c \leq c_0(b)$ .



**Case 8.** If  $c = c_4$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 6$  such that if


- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_6, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_6$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$ ,  $(\lambda_1, \lambda_6)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0]$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_1$ ,  $\lambda = \lambda_5$ , then (1-7) has exactly 11 positive solutions;
- $\lambda \in (\lambda_5, \lambda_1)$ , then (1-7) has exactly 13 positive solutions.

Figure 19 illustrates Case 8.

**Case 9.** If  $c \in (c_4, c_5]$  (some  $c_5(b) > 0$ ), then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 7$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_7, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_7$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4)$ ,  $(\lambda_6, \lambda_7)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0]$ ,  $\lambda = \lambda_6$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda \in (\lambda_1, \lambda_6)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda = \lambda_5$ , then (1-7) has exactly 11 positive solutions;
- $\lambda \in (\lambda_5, \lambda_1]$ , then (1-7) has exactly 13 positive solutions.


Figure 20 illustrates Case 9.



**Figure 20.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 28$  (Case 9).

**Case 10.** If  $c \in (c_5, c_6)$  (some  $c_6(b) > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 9$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_9, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_9$ , then (1-7) has exactly 5 positive solutions;




**Figure 21.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 29$  (Case 10).

- $\lambda \in (\lambda_3, \lambda_4), (\lambda_8, \lambda_9)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0], \lambda = \lambda_8$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5), (\lambda_6, \lambda_7)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_5, \lambda = \lambda_6, \lambda \in (\lambda_1, \lambda_8)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6), \lambda = \lambda_7$ , then (1-7) has exactly 11 positive solutions;
- $\lambda \in (\lambda_7, \lambda_1]$ , then (1-7) has exactly 13 positive solutions.

Figure 21 illustrates Case 10. Notice that the blue curve has now also become  $\Sigma$ -shaped and its shape persists through  $c \leq c_0(b)$ .

**Case 11.** If  $c = c_6$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 8$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_8, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_8$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_7, \lambda_8)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0], \lambda = \lambda_7$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;




**Figure 22.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 30$  (Case 11).

- $\lambda = \lambda_5, \lambda \in (\lambda_1, \lambda_7)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6)$ , then (1-7) has exactly 11 positive solutions;
- $\lambda = \lambda_6$ , then (1-7) has exactly 12 positive solutions;
- $\lambda \in (\lambda_6, \lambda_1]$ , then (1-7) has exactly 13 positive solutions.

Figure 22 illustrates Case 11.

**Case 12.** If  $c \in (c_6, c_7)$  (some  $c_7(b) > 0$ ) then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 9$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_9, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_9$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_8, \lambda_9)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0], \lambda = \lambda_8$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_5, \lambda \in (\lambda_1, \lambda_8)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6)$ , then (1-7) has exactly 11 positive solutions;
- $\lambda = \lambda_6, \lambda \in (\lambda_7, \lambda_1]$ , then (1-7) has exactly 13 positive solutions;



**Figure 23.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 30.1$  (Case 12).


- $\lambda = \lambda_7$ , then (1-7) has exactly 14 positive solutions;
- $\lambda \in (\lambda_6, \lambda_7)$ , then (1-7) has exactly 15 positive solutions.

Figure 23 illustrates Case 12.


**Case 13.** If  $c = c_7$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 8$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_8, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_8$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_7, \lambda_8)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0], \lambda = \lambda_7$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_5, \lambda \in (\lambda_1, \lambda_7)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6)$ , then (1-7) has exactly 11 positive solutions;
- $\lambda = \lambda_6$ , then (1-7) has exactly 13 positive solutions;
- $\lambda = \lambda_1$ , then (1-7) has exactly 14 positive solutions;
- $\lambda \in (\lambda_6, \lambda_1)$ , then (1-7) has exactly 15 positive solutions.

Figure 24 illustrates Case 13.



**Figure 24.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 30.3$  (Case 13).



**Figure 25.**  $\rho$  versus  $\lambda$  when  $b = 10$  and  $c = 32$  (Case 14).

**Case 14.** If  $c \in (c_7, c_0(b))$  then there exist  $\lambda_i > 0$  for  $i = 1, 2, \dots, 9$  such that if

- $\lambda \in [0, \lambda_2)$ , then (1-7) has no positive solution;
- $\lambda = \lambda_2$ , then (1-7) has a unique positive solution;
- $\lambda \in (\lambda_2, \lambda_3)$ , then (1-7) has exactly 2 positive solutions;
- $\lambda = \lambda_3$  and  $\lambda \in (\lambda_9, \infty)$ , then (1-7) has exactly 4 positive solutions;
- $\lambda = \lambda_9$ , then (1-7) has exactly 5 positive solutions;
- $\lambda \in (\lambda_3, \lambda_4), (\lambda_8, \lambda_9)$ , then (1-7) has exactly 6 positive solutions;
- $\lambda = \lambda_4$ , then (1-7) has exactly 7 positive solutions;
- $\lambda \in (\lambda_4, \lambda_0], \lambda = \lambda_8$ , then (1-7) has exactly 8 positive solutions;
- $\lambda \in (\lambda_0, \lambda_5)$ , then (1-7) has exactly 9 positive solutions;
- $\lambda = \lambda_5, \lambda \in (\lambda_7, \lambda_8)$ , then (1-7) has exactly 10 positive solutions;
- $\lambda \in (\lambda_5, \lambda_6), \lambda = \lambda_7$ , then (1-7) has exactly 11 positive solutions;
- $\lambda \in (\lambda_1, \lambda_7)$ , then (1-7) has exactly 12 positive solutions;
- $\lambda = \lambda_6$ , then (1-7) has exactly 13 positive solutions;
- $\lambda \in (\lambda_6, \lambda_1]$ , then (1-7) has exactly 15 positive solutions;

Figure 25 illustrates Case 14.

## 7. Analytical results

In order to bolster our computational results as well as elaborate on the behavior of the bifurcation curves, we procure some analytical results. First, we recall some results from [Laetsch 1970] detailing the behavior of  $G_1(\rho)$  when  $\rho \rightarrow \sigma(b, c)^-$

and when  $\rho \rightarrow 0^+$  in the following lemmas, where  $\sigma(b, c)$  represents the smallest positive root of  $f(u)$ .

**Lemma 2** [Laetsch 1970].  $\lim_{\rho \rightarrow \sigma(b,c)^-} G_1(\rho) = \infty$ .

**Lemma 3** [Laetsch 1970].  $\lim_{\rho \rightarrow 0^+} G_1(\rho) = \pi/(2\sqrt{b})$ .

Our main goal for this section is to establish the following analytical results for (1-9) and (1-11). Recall that  $\lambda = [G_2(\rho, q)]^2$  and  $\lambda = [G_3(\rho, q)]^2$  from Theorems 4.1 and 5.1, respectively. Thus, we can obtain some global behavior of the  $\rho$  versus  $\lambda$  bifurcation curve via study of  $G_2(\rho, q)$  and  $G_3(\rho, q)$ .

**Theorem 7.1.**

$$(1) \left( \sqrt{2} - \frac{1}{\sqrt{2}} \right) \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} \leq G_2(\rho, q) \leq \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}};$$

$$(2) \quad G_3(\rho, q) \leq \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}.$$

*Proof.* To prove (1), recall

$$\begin{aligned} G_2(\rho, q) &= \frac{1}{\sqrt{2}\sqrt{F(\rho) - F(q)}} \\ &= \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \frac{1}{\sqrt{2}} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}}. \end{aligned} \quad (7-1)$$

We ascertain an upper bound by substituting  $q = 0$  into (7-1) yielding

$$G_2(\rho, q) \leq \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}.$$

Also, recalling  $q \in [0, \rho)$  and allowing  $q \rightarrow \rho^-$  in (7-1) we obtain

$$G_2(\rho, q) \geq \left( \sqrt{2} - \frac{1}{\sqrt{2}} \right) \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}$$

as the lower bound. Hence,

$$\left( \sqrt{2} - \frac{1}{\sqrt{2}} \right) \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} \leq G_2(\rho, q) \leq \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}.$$

Now to prove (2). Recall

$$G_3(\rho, q) = \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}} - \sqrt{2} \int_0^q \frac{dw}{\sqrt{F(\rho) - F(w)}}. \quad (7-2)$$

Similarly, by substituting  $q = 0$  into (7-2) we obtain

$$G_3(\rho, q) \leq \sqrt{2} \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}$$

as the upper bound. □

**Theorem 7.2.**  $\lim_{\rho \rightarrow \sigma(b,c)^-} G_2(\rho, q) = \infty.$

*Proof.* By Theorem 7.1, we have

$$G_2(\rho, q) \geq \left( \sqrt{2} - \frac{1}{\sqrt{2}} \right) \int_0^\rho \frac{dw}{\sqrt{F(\rho) - F(w)}}. \quad (7-3)$$

From Lemma 2, it is clear that the right side of (7-3) approaches infinity as

$$\rho \rightarrow \sigma(b, c)^-.$$

Therefore, it is apparent that

$$\lim_{\rho \rightarrow \sigma(b,c)^-} G_2(\rho, q) = \infty. \quad \square$$

### Acknowledgement

A significant portion of this research work was completed by Kathryn Ashley and Victoria Sincavage while attending the 2011 NSF REU Program at the Center for Computational Sciences at Mississippi State University. We would like to extend a special thanks to our advisor and mentor Professor Ratnasingham Shivaji for his guidance and support throughout this research project.

### References

- [Allee 1938] W. C. Allee, *The social life of animals*, Norton, New York, 1938.
- [Brown et al. 1981] K. J. Brown, M. M. A. Ibrahim, and R. Shivaji, "S-shaped bifurcation curves", *Nonlinear Anal.* **5**:5 (1981), 475–486. MR 82h:35007 Zbl 0458.35036
- [Cantrell and Cosner 2003] R. S. Cantrell and C. Cosner, *Spatial ecology via reaction-diffusion equations*, Wiley, Chichester, 2003. MR 2007a:92069 Zbl 1059.92051
- [Cantrell and Cosner 2007] R. S. Cantrell and C. Cosner, "Density dependent behavior at habitat boundaries and the Allee effect", *Bull. Math. Biol.* **69**:7 (2007), 2339–2360. MR 2341874 Zbl 05265737
- [Causey et al. 2010] R. Causey, S. Sasi, and R. Shivaji, "An ecological model with grazing and constant yield harvesting", *Bull. Belg. Math. Soc. Simon Stevin* **17**:5 (2010), 833–839. MR 2012a:35103 Zbl 1208.35153
- [Dennis 1989] B. Dennis, "Allee effects: population growth, critical density, and the chance of extinction", *Natur. Resource Modeling* **3**:4 (1989), 481–538. MR 91h:92032 Zbl 0850.92062
- [Goddard and Shivaji 2012] J. Goddard, II and R. Shivaji, "A population model with nonlinear boundary conditions and constant yield harvesting", *Proceedings of Dynamic Systems and Applications* **6** (2012), 150–157.
- [Goddard et al. 2010a] J. Goddard, II, E. K. Lee, and R. Shivaji, "A double S-shaped bifurcation curve for a reaction-diffusion model with nonlinear boundary conditions", *Bound. Value Probl.* (2010), Art. ID 357542. MR 2665818 Zbl 1211.35037
- [Goddard et al. 2010b] J. Goddard, II, E. K. Lee, and R. Shivaji, "Population models with nonlinear boundary conditions", *Electron. J. Differ. Equ. Conf.* **19** (2010), 135–149. MR 2012f:35306 Zbl 1204.34030



- [Goddard et al. 2011a] J. Goddard, II, E. K. Lee, and R. Shivaji, “Population models with diffusion, strong Allee effect, and nonlinear boundary conditions”, *Nonlinear Anal.* **74**:17 (2011), 6202–6208. MR 2012i:35192 Zbl 1227.35172
- [Goddard et al. 2011b] J. Goddard, II, R. Shivaji, and E. K. Lee, “Diffusive logistic equation with non-linear boundary conditions”, *J. Math. Anal. Appl.* **375**:1 (2011), 365–370. MR 2011i:35123 Zbl 1208.35082
- [Kuussaari et al. 1996] M. Kuussaari, M. Nieminen, and I. Hanski, “An experimental study of migration in the Glanville fritillary butterfly *Melitaea cinxia*”, *Journal of Animal Ecology* **65**:6 (1996), 791–801.
- [Laetsch 1970] T. Laetsch, “The number of solutions of a nonlinear two point boundary value problem”, *Indiana Univ. Math. J.* **20** (1970), 1–13. MR 42 #4815 Zbl 0215.14602
- [Lee et al. 2011] E. K. Lee, S. Sasi, and R. Shivaji, “S-shaped bifurcation curves in ecosystems”, *J. Math. Anal. Appl.* **381**:2 (2011), 732–741. MR 2012e:92080 Zbl 1221.35421
- [Lewis and Kareiva 1993] M. A. Lewis and P. Kareiva, “Allee dynamics and the spread of invading organisms”, *Theoretical Population Biology* **43**:2 (1993), 141–158. Zbl 0769.92025
- [Matthysen 2005] E. Matthysen, “Density-dependent dispersal in birds and mammals”, *Ecography* **28** (2005), 403–416.
- [van Nes and Scheffer 2005] E. H. van Nes and M. Scheffer, “Implications of spatial heterogeneity for catastrophic regime shifts in ecosystems”, *Ecology* **86**:7 (2005), 1797–1807.
- [Poole et al. 2012] E. Poole, B. Roberson, and B. Stephenson, “Weak Allee effect, grazing, and S-shaped bifurcation curves”, *Involve* **5**:2 (2012), 133–158.
- [Shi and Shivaji 2006] J. Shi and R. Shivaji, “Persistence in reaction diffusion models with weak Allee effect”, *J. Math. Biol.* **52**:6 (2006), 807–829. MR 2007g:92070 Zbl 1110.92055

Received: 2012-07-23    Revised: 2013-04-04    Accepted: 2013-04-10

klashley01@email.wm.edu    *Department of Mathematical Sciences, Clemson University,  
Clemson, SC 29634, United States*

vsincav@clemson.edu    *Department of Mathematical Sciences, Clemson University,  
Clemson, SC 29634, United States*

jgoddard@aum.edu    *Department of Mathematics, Auburn University Montgomery,  
Montgomery, AL 36124-4023, United States*

# involve

msp.org/involve

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

### BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	Victoria University, Australia pietro.cerone@vu.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moselehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsgdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA jgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sngupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

## PRODUCTION


Silvio Levy, Scientific Editor

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2013 is US \$105/year for the electronic version, and \$145/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# involve

2013

vol. 6

no. 4

Embeddedness for singly periodic Scherk surfaces with higher dihedral symmetry VALMIR BUCAJ, SARAH CANNON, MICHAEL DORFF, JAMAL LAWSON AND RYAN VIERTEL	383
An elementary inequality about the Mahler measure KONSTANTIN STULOV AND RONGWEI YANG	393
Ecological systems, nonlinear boundary conditions, and $\Sigma$ -shaped bifurcation curves KATHRYN ASHLEY, VICTORIA SINCAVAGE AND JEROME GODDARD II	399
The probability of randomly generating finite abelian groups TYLER CARRICO	431
Free and very free morphisms into a Fermat hypersurface TABES BRIDGES, RANKEYA DATTA, JOSEPH EDDY, MICHAEL NEWMAN AND JOHN YU	437
Irreducible divisor simplicial complexes NICHOLAS R. BAETH AND JOHN J. HOBSON	447
Smallest numbers beginning sequences of 14 and 15 consecutive happy numbers DANIEL E. LYONS	461
An orbit Cartan type decomposition of the inertia space of $SO(2m)$ acting on $\mathbb{R}^{2m}$ CHRISTOPHER SEATON AND JOHN WELLS	467
Optional unrelated-question randomized response models SAT GUPTA, ANNA TUCK, TRACY SPEARS GILL AND MARY CROWE	483
On the difference between an integer and the sum of its proper divisors NICHOLE DAVIS, DOMINIC KLYVE AND NICOLE KRAGHT	493
A Pexider difference associated to a Pexider quartic functional equation in topological vector spaces SAEID OSTADBASHI, ABBAS NAJATI, MAHSA SOLAIMANINIA AND THEMISTOCLES M. RASSIAS	505



1944-4176(2013)6:4;1-7