

# involve

a journal of mathematics

The number of convex topologies  
on a finite totally ordered set

Tyler Clark and Tom Richmond



# The number of convex topologies on a finite totally ordered set

Tyler Clark and Tom Richmond

(Communicated by Kenneth S. Berenhaut)

We give an algorithm to find the number  $T_{\text{cvx}}(n)$  of convex topologies on a totally ordered set  $X$  with  $n$  elements, and present these numbers for  $n \leq 10$ .

## 1. Introduction

A subset  $B$  of poset  $(X, \leq)$  is *increasing* if  $x \in B$  and  $y \geq x$  imply  $y \in B$ , and is *convex* if  $x, z \in B$  and  $x \leq y \leq z$  imply  $y \in B$ . An  $n$ -point totally ordered set  $X$  may be labeled  $X = \{1, 2, \dots, n\}$ , where  $1 < 2 < \dots < n$ . This set will be denoted  $[1, n]$ , and in general,  $[a, b]$  will denote  $\{a, a+1, \dots, b\} \subset \mathbb{N}$  with the natural order from  $\mathbb{N}$ . A topology on  $(X, \leq)$  is *convex* if it has a base of convex sets, or equivalently, if each point has a neighborhood base of convex sets. Because of these equivalent characterizations, convex topologies are often called *locally convex* topologies. (See [Nachbin 1965]). For finite sets, every point  $j$  has a minimal neighborhood  $\text{MN}(j)$ , which is the intersection of all neighborhoods of  $j$ . It is convenient to identify a topology on  $[1, n]$  with its base  $\{\text{MN}(j) : j \in [1, n]\}$  of minimal neighborhoods of each point. Finite topological spaces are used in computer graphics, where the Euclidean plane is modeled by a topology on a finite set of pixels. If  $a < b < c$  in a finite poset with a topology, if  $c$  is “near”  $a$  and there is any compatibility between the topology and order, we would expect  $b$  to also be near  $a$ . This is the convexity condition, which is a natural, weak compatibility condition between a topology and order assumed in most applications. We will consider the number of convex topologies on a finite totally ordered set  $[1, n]$ .

An excellent reference on finding the number  $T(n)$  of topologies on an  $n$ -element set is [Erné and Stege 1991]. Currently,  $T(n)$  is known for  $n \leq 18$ . A standard approach to counting topologies on a finite set  $X$  is to employ the one-to-one correspondence between a topology  $\tau$  on  $X$  and the associated specialization quasiorder defined by  $x \leq y$  if and only if  $x$  is in the closure of  $y$ . This correspondence

---

MSC2010: 05A15, 06F30, 54A10, 54F05.

Keywords: convex topology, totally ordered set, number of topologies.

dates back to [Alexandroff 1937]. (See [Richmond 1998] for a survey of this connection.) One approach to counting the convex topologies would be to find a (bordered) characterization of convex topologies using some compatibility between the specialization order and the given total order. Fruitful results in this direction have not been found.

For  $j \in [1, n]$ , a convex subset  $N(j)$  of  $[1, n]$  containing  $j$  has the form  $[a, b]$ , where  $1 \leq a \leq j \leq b \leq n$ . There are  $j$  choices for  $a$  and  $n - j + 1$  choices for  $b$ , and thus  $j(n + j - 1)$  choices for  $N(j)$ . Since a base of minimal neighborhoods for a locally convex topology on  $[1, n]$  consists of one convex subset  $N(j)$  for each  $j \in [1, n]$ , we see that

$$\prod_{j=1}^n (j)(n + j - 1) = (n!)^2$$

gives an upper bound on  $T_{\text{cvx}}(n)$ . Of course, arbitrarily selecting a convex set  $N(j)$  containing  $j$  for each  $j \in [1, n]$  is unlikely to give a base for a topology, so this upper bound is not sharp.

## 2. Nested convex topologies

Stephen [1968] gave a recursive formula for the number of nested topologies (or equivalently, ordered partitions) on an  $n$ -point set  $X$ , generating the sequence 1, 3, 13, 75, 541, 4683, 47293,  $\dots$ , which is A000670 in *The On-Line Encyclopedia of Integer Sequences* (OEIS); see [Sloane 2014]. If  $X = [1, n]$  is a totally ordered set with  $n$  elements, let  $T_{\text{Nest}}(n)$  be the number of nested convex topologies on  $X$ , and let  $T_{\text{Nest}}(n, k)$  be the number of those convex topologies consisting of  $k$  nested nonempty open sets  $U_1, U_2, \dots, U_k$ , where  $X = U_1 \supset U_2 \supset \dots \supset U_k \neq \emptyset$ . Since the indiscrete topology is the only nested topology with one nonempty open set,  $T_{\text{Nest}}(n, 1) = 1$ . Suppose we have found  $T_{\text{Nest}}(m, j)$  for all  $m \leq n$  and  $j \leq k$ . To find  $T_{\text{Nest}}(n, k + 1)$ , note that  $X = U_1 \supset U_2 \supset U_3 \supset \dots \supset U_{k+1} \neq \emptyset$  implies that  $U_2$  must contain at least  $k$  elements and at most  $n - 1$  elements. If  $|U_2| = j$ , there are  $n - j + 1$  ways to choose  $U_2$  as a convex subset of  $X$ , and  $T_{\text{Nest}}(j, k)$  ways to complete the nested convex topology  $\{U_2, \dots, U_{k+1}\}$  on the  $j$ -point totally ordered set  $U_2$ . Thus, we have

$$T_{\text{Nest}}(n, k + 1) = \sum_{j=k}^{n-1} (n - j + 1) T_{\text{Nest}}(j, k) = \sum_{m=2}^{n-k+1} m \cdot T_{\text{Nest}}(n - m + 1, k),$$

where the second equality follows from the substitution  $m = n - j + 1$ . In Table 1, we tabulate the values of  $T_{\text{Nest}}(n, k)$  for  $n, k \leq 10$ .

$n \backslash k$	1	2	3	4	5	6	7	8	9	10
1	1									
2	1	2								
3	1	5	4							
4	1	9	16	8						
5	1	14	41	44	16					
6	1	20	85	146	112	32				
7	1	27	155	377	456	272	64			
8	1	35	259	833	1,408	1,312	640	128		
9	1	44	406	1,652	3,649	4,712	3,568	1,472	256	
10	1	54	606	3,024	8,361	14,002	14,608	9,312	3,328	512

**Table 1.**  $T_{\text{Nest}}(n, k)$ , the number of topologies on a totally ordered  $n$ -point set consisting of  $k$  nested convex sets.

This table (sequence A056242 in the OEIS [Mallows 2014]) is also used by Hwang and Mallow [1995] to count the number of *order-consecutive partitions* of  $X = \{1, 2, \dots, n\}$ , which they define as follows: An ordered list  $S_1, S_2, \dots, S_m$  of subsets of  $X$  is an order-consecutive partition of  $X$  if  $\{S_1, \dots, S_m\}$  is a partition of  $X$  and each of the sets  $\bigcup_{j=1}^k S_j$  ( $1 \leq k \leq m$ ) is a consecutive set of integers. If  $\{S_1, \dots, S_m\}$  is an order-consecutive partition, clearly  $\{S_1, S_1 \cup S_2, S_1 \cup S_2 \cup S_3, \dots, X\}$  is a nested convex topology on  $X$ . Conversely, any nested convex topology  $\tau = \{U_1, U_2, \dots, U_k\}$  on  $X = \{1, 2, \dots, n\}$  generates the order-consecutive partition  $U_1, U_2 \setminus U_1, U_3 \setminus U_2, \dots, U_k \setminus U_{k-1}$ .

It is easy to confirm from our formula for  $T_{\text{Nest}}(n, k)$  that  $T_{\text{Nest}}(n, n) = 2^{n-1}$  and  $T_{\text{Nest}}(n, 2) = \Delta_n - 1$ , where  $\Delta_n$  is the  $n$ -th triangular number.

Now, we note that

$$T_{\text{Nest}}(n) = \sum_{k=1}^n T_{\text{Nest}}(n, k).$$

This sequence, whose first few elements are

$$(T_{\text{Nest}}(n))_{n=1}^{10} = (1, 3, 10, 34, 116, 396, 1352, 4616, 15760, 53808),$$

appears as A007052 in the OEIS [Mallows et al. 2014], where it is noted that

$$T_{\text{Nest}}(n) = 4 T_{\text{Nest}}(n - 1) - 2 T_{\text{Nest}}(n - 2) \quad \text{for } n > 2.$$

Solving this recurrence relation by standard techniques gives

$$T_{\text{Nest}}(n) = \frac{(2 + \sqrt{2})^n + (2 - \sqrt{2})^n}{4}.$$

Nested convex topologies have as much inclusion as possible. Not only are they totally ordered by inclusion, but they maximize “overlap”. The other extreme would be to have as little inclusion and overlap as possible. This suggests considering mutually disjoint collections. A collection  $\mathcal{D}$  of mutually disjoint convex subsets of  $X$  is not a basis for a topology if  $\bigcup \mathcal{D} \neq X$ , but  $\mathcal{D} \cup \{X\}$  is always a basis for a convex topology on  $X$ . The authors have shown that the number of topologies on an  $n$ -element totally ordered set having a base consisting of a mutually disjoint collection  $\mathcal{D}$  of convex sets, or such a collection  $\mathcal{D}$  together with  $X$ , is  $F_{2n+1} - 1$ , where  $F_k$  is the  $k$ -th Fibonacci number [Clark and Richmond 2010].


### 3. An algorithm for $T_{\text{cvx}}(n)$

We now present a recursive algorithm to find the number  $T_{\text{cvx}}(n)$  of convex topologies on a totally ordered set  $[1, n]$ . It is easy to check that  $T_{\text{cvx}}(1) = 1 = T(1)$  and  $T_{\text{cvx}}(2) = 4 = T(2)$ . That is, the only topology on a 1-point set is convex, as are all four topologies on a 2-point set.


Suppose  $T_{\text{cvx}}(n)$  is known. To find  $T_{\text{cvx}}(n+1)$ , note that each convex topology on  $[1, n+1]$ , when restricted to  $[1, n]$ , gives a unique convex topology on  $[1, n]$ . Thus, we may count  $T_{\text{cvx}}(n)$  by looping through each topology  $\tau$  counted in  $T_{\text{cvx}}(n)$ , adding  $n+1$  as the greatest point, adjusting the minimal neighborhoods of  $j \in [1, n]$ , and defining the minimal neighborhood of  $n+1$  so that the subspace topology on  $[1, n]$  is still  $\tau$ . That is, considering how each topology on  $[1, n]$  may be appropriately expanded to  $[1, n+1]$  gives a complete, unduplicated count of the convex topologies on  $[1, n+1]$ .

**Step 1: Redefining minimal neighborhoods of  $j \in [1, n]$ .** We loop through all convex topologies  $\tau$  on  $[1, n]$ . The simplest way to extend  $\tau$  to  $[1, n+1]$  so that the restriction of the extension is still  $\tau$  would be to keep the minimum neighborhoods of each  $j \in [1, n]$  unchanged. However, we may also expand some of the minimal neighborhoods of points  $j \in [1, n]$  to include  $n+1$ . To maintain convexity and to guarantee a topology on  $[1, n+1]$  whose restriction to  $[1, n]$  agrees with  $\tau$ , the minimal neighborhood  $\text{MN}(j)$  of  $j$  can be expanded to include  $n+1$  if and only if  $\text{MN}(j)$  already includes  $n$ . If  $n \in \text{MN}(j) \subseteq \text{MN}(k)$  and  $\text{MN}(j)$  is expanded to include  $n+1$ , then  $\text{MN}(k)$  must also be expanded to include  $n+1$ , for otherwise  $\text{MN}(k)$  would be a neighborhood of  $j$  not including  $n+1$ , contrary to the hypothesis that the minimal neighborhood of  $j$  was to include  $n+1$ .

As an immediate consequence, if  $n \in \text{MN}(j) = \text{MN}(k)$ , then  $\text{MN}(j)$  is expanded to include  $n+1$  if and only if  $\text{MN}(k)$  is. That is, a single basis element which happens to be the minimal neighborhood of distinct points  $j$  and  $k$  is still treated as a single entity in the expansion process.



**Figure 1.** A sample topology on  $[1, 8]$ .




**Figure 2.** Possible expansions of minimal neighborhoods containing previous right endpoint: none, outermost one, outermost two, outermost three.

Thus, if  $\mathcal{B} = \{MN(1), MN(2), \dots, MN(n)\}$  has  $m$  distinct sets containing  $n$ , we expand the outermost  $k$  of these to include  $n + 1$ , looping as  $k$  goes from 1 to  $m$ .


For example, consider the convex topology  $\tau$  on  $[1, 8]$  having a base of minimal neighborhoods  $\mathcal{B} = \{\{1\}, [2, 8], [3, 4], \{5\}, [5, 8], \{8\}\}$ , as shown in Figure 1.

We may add 9 to this topology without changing any of the minimal neighborhoods of  $j$  for  $j \in [1, 8]$ , or since  $MN(2), MN(6) = MN(7)$ , and  $MN(8)$  include the right endpoint 8, they may be extended to include the added point 9. Since  $8 \in MN(8) \subset MN(7) = MN(6) \subset MN(2)$ , we note that  $MN(6)$  is expanded if and only if  $MN(7)$  is expanded, so we do not need to treat  $MN(6)$  and  $MN(7)$  as distinct basis elements and we may effectively ignore the duplicate  $MN(7)$ . Also, if  $MN(6)$  is expanded, then  $MN(6) \subset MN(2)$  implies that  $MN(2)$  would also have to be expanded. Repeating this idea, we may expand nothing except the outermost (i.e., longest) minimal neighborhood containing 8, namely  $MN(2)$ , the outermost two minimal neighborhoods containing 8, namely  $MN(2)$  and  $MN(6)$ , or the outermost three,  $MN(2), MN(6)$ , and  $MN(8)$ . See Figure 2.

**Step 2: Defining the minimal neighborhood of the added point.** Having determined the expansion of minimal neighborhoods of  $j \in [1, n]$ , it remains to define the minimal neighborhood  $MN(n + 1)$  of  $n + 1$ . Clearly we must have  $n + 1 \in MN(n + 1)$ . The convexity condition and our need to retain the original topology  $\tau$  on  $[1, n]$  as a subspace imply that  $MN(n + 1)$  must be of form  $\{n + 1\} \cup I$ , where  $I$  is increasing and open in  $\tau$ . The final condition is the minimality of the neighborhood  $MN(n + 1)$ . In Step 1, we may have expanded some neighborhoods of  $n$  to contain  $n + 1$  and, if so, the minimal neighborhood of  $n + 1$  must be contained in each of these previously defined neighborhoods of  $n + 1$ . Thus,  $MN(n + 1)$  must be of the form  $\{n + 1\} \cup I$ ,



**Figure 3.** Possible choices for  $MN(9)$  if no minimal neighborhoods  $MN(j)$  are expanded for  $j \in [1, 8]$ .



**Figure 4.** Possible choices for  $MN(9)$  if  $MN(2)$  and  $MN(6)$  are expanded to include 9.

where  $I$  is increasing and  $\tau$ -open, and  $I$  is contained in the innermost (shortest) neighborhood  $MN(j)$  which was expanded in [Step 1](#).

Continuing the example presented above, we may expand none of the original minimal neighborhoods of  $j \in [1, 8]$  to include 9, and then the minimal neighborhood  $MN(9)$  of 9 may be defined as  $\{9\} \cup I$ , where  $I$  is an increasing  $\tau$ -open set in any of the six ways suggested in [Figure 3](#).

[Figure 4](#) shows the three possible choices for the minimal neighborhood  $MN(9)$  if the outermost two minimal neighborhoods containing 8, namely  $MN(2)$  and  $MN(6)$ , have been expanded to include 9.

A computer implementation of this algorithm yields the values for  $T_{\text{cvx}}(n)$  shown in [Table 2](#) below. With the  $T_{\text{cvx}}(2) = 4$  convex topologies on  $[1, 2]$  as input, the computer implementation loops through all the topologies  $\tau$  on  $[1, n]$ , adds  $n + 1$ , determines the number  $m$  of distinct minimal neighborhoods of  $j \in [1, n]$  containing  $n$ , expands the outermost  $k$  of these to contain  $n + 1$  (as  $k$  goes from 0 to  $m$ ), determines the increasing  $\tau$ -open sets, defines the minimal neighborhood  $MN(n + 1)$  of  $n + 1$  as  $\{n + 1\} \cup I$ , where  $I$  is one of the increasing  $\tau$ -open sets contained in the smallest  $MN(j)$  previously expanded to include  $n + 1$ , and, at each selection of an option above, increments the  $T_{\text{cvx}}(n + 1)$  counter and records the data for this new topology on  $[1, n + 1]$  required for the next iteration.

The efficiency of this algorithm can be improved by eliminating duplication of computations. For example, if  $p$  is the largest integer with  $MN(p) = X$  for two topologies  $s$  and  $t$  which agree to the right of  $p$ , then the computation for  $s$  duplicates that for  $t$ , as noted by a helpful referee.

$n$	$T_{\text{Nest}}(n)$	$T_{\text{cvx}}(n)$	$T(n)$
1	1	1	1
2	3	4	4
3	10	21	29
4	34	129	355
5	116	876	6,942
6	396	6,376	209,527
7	1,352	48,829	9,535,241
8	4,616	388,771	642,779,354
9	15,760	3,191,849	63,260,289,423
10	53,808	26,864,936	8,977,053,873,043

**Table 2.** The numbers  $T_{\text{Nest}}(n)$  and  $T_{\text{cvx}}(n)$  of nested convex topologies and convex topologies on an  $n$ -point totally ordered set, and the number  $T(n)$  of topologies on an  $n$ -point set.

The numbers  $T_{\text{cvx}}(n)$  in Table 2 were also verified for  $n \leq 8$  without the algorithm using an exhaustive generation scheme. For comparison, we also include the number  $T_{\text{Nest}}(n)$  of nested convex topologies and the number  $T(n)$  of topologies on  $n$  points in the table.

## References

- [Alexandroff 1937] P. Alexandroff, “Diskrete Räume”, *Mat. Sb. (N.S.)* **2**:3 (1937), 501–518.
- [Clark and Richmond 2010] T. Clark and T. Richmond, “Collections of mutually disjoint convex subsets of a totally ordered set”, *Fibonacci Quart.* **48**:1 (2010), 77–79. Currently also at <http://people.wku.edu/tom.richmond/Papers/FibQuarterly.pdf>. MR 2011e:11029 Zbl 1211.05017
- [Erné and Stege 1991] M. Erné and K. Stege, “Counting finite posets and topologies”, *Order* **8**:3 (1991), 247–265. MR 93b:06004 Zbl 0752.05002
- [Hwang and Mallows 1995] F. K. Hwang and C. L. Mallows, “Enumerating nested and consecutive partitions”, *J. Combin. Theory Ser. A* **70**:2 (1995), 323–333. MR 96e:05014 Zbl 0819.05005
- [Mallows 2014] C. Mallows, “A056242: Triangle read by rows”, entry A056242 in *The on-line encyclopedia of integer sequences* (<http://oeis.org>), 2014.
- [Mallows et al. 2014] C. Mallows, N. J. A. Sloane, and S. Plouffe, “A007052: Number of order-consecutive partitions of  $n$ ”, entry A007052 in *The on-line encyclopedia of integer sequences* (<http://oeis.org>), 2014.
- [Nachbin 1965] L. Nachbin, *Topology and order*, Van Nostrand Mathematical Studies **4**, D. Van Nostrand, Princeton, N.J., 1965. MR 36 #2125 Zbl 0131.37903
- [Richmond 1998] T. A. Richmond, “Quasiorders, principal topologies, and partially ordered partitions”, *Internat. J. Math. Math. Sci.* **21**:2 (1998), 221–234. MR 99h:06003 Zbl 0898.54005
- [Sloane 2014] N. J. A. Sloane, “A000670: Fubini numbers”, entry A000670 in *The on-line encyclopedia of integer sequences* (<http://oeis.org>), 2014.

[Stephen 1968] D. Stephen, “Topology on finite sets”, *Amer. Math. Monthly* **75** (1968), 739–741.  
[MR 38 #2725](#) [Zbl 0191.20701](#)

Received: 2011-10-19    Revised: 2013-06-14    Accepted: 2013-08-07

[thomas.clark973@topper.wku.edu](mailto:thomas.clark973@topper.wku.edu) *Department of Mathematics, Western Kentucky University,  
1906 College Heights Boulevard,  
Bowling Green, KY 42101, United States*

[tom.richmond@wku.edu](mailto:tom.richmond@wku.edu) *Department of Mathematics, Western Kentucky University,  
1906 College Heights Boulevard,  
Bowling Green, KY 42101, United States*

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, [berenhks@wfu.edu](mailto:berenhks@wfu.edu)

### BOARD OF EDITORS

Colin Adams	Williams College, USA <a href="mailto:colin.c.adams@williams.edu">colin.c.adams@williams.edu</a>	David Larson	Texas A&M University, USA <a href="mailto:larson@math.tamu.edu">larson@math.tamu.edu</a>
John V. Baxley	Wake Forest University, NC, USA <a href="mailto:baxley@wfu.edu">baxley@wfu.edu</a>	Suzanne Lenhart	University of Tennessee, USA <a href="mailto:lenhart@math.utk.edu">lenhart@math.utk.edu</a>
Arthur T. Benjamin	Harvey Mudd College, USA <a href="mailto:benjamin@hmc.edu">benjamin@hmc.edu</a>	Chi-Kwong Li	College of William and Mary, USA <a href="mailto:ckli@math.wm.edu">ckli@math.wm.edu</a>
Martin Bohner	Missouri U of Science and Technology, USA <a href="mailto:bohner@mst.edu">bohner@mst.edu</a>	Robert B. Lund	Clemson University, USA <a href="mailto:lund@clemson.edu">lund@clemson.edu</a>
Nigel Boston	University of Wisconsin, USA <a href="mailto:boston@math.wisc.edu">boston@math.wisc.edu</a>	Gaven J. Martin	Massey University, New Zealand <a href="mailto:g.j.martin@massey.ac.nz">g.j.martin@massey.ac.nz</a>
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA <a href="mailto:budhiraj@email.unc.edu">budhiraj@email.unc.edu</a>	Mary Meyer	Colorado State University, USA <a href="mailto:meyer@stat.colostate.edu">meyer@stat.colostate.edu</a>
Pietro Cerone	La Trobe University, Australia <a href="mailto:P.Cerone@latrobe.edu.au">P.Cerone@latrobe.edu.au</a>	Emil Minchev	Ruse, Bulgaria <a href="mailto:eminchev@hotmail.com">eminchev@hotmail.com</a>
Scott Chapman	Sam Houston State University, USA <a href="mailto:scott.chapman@shsu.edu">scott.chapman@shsu.edu</a>	Frank Morgan	Williams College, USA <a href="mailto:frank.morgan@williams.edu">frank.morgan@williams.edu</a>
Joshua N. Cooper	University of South Carolina, USA <a href="mailto:cooper@math.sc.edu">cooper@math.sc.edu</a>	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran <a href="mailto:moslehian@ferdowsi.um.ac.ir">moslehian@ferdowsi.um.ac.ir</a>
Jem N. Corcoran	University of Colorado, USA <a href="mailto:corcoran@colorado.edu">corcoran@colorado.edu</a>	Zuhair Nashed	University of Central Florida, USA <a href="mailto:znashed@mail.ucf.edu">znashed@mail.ucf.edu</a>
Toka Diagana	Howard University, USA <a href="mailto:tdiagana@howard.edu">tdiagana@howard.edu</a>	Ken Ono	Emory University, USA <a href="mailto:ono@mathcs.emory.edu">ono@mathcs.emory.edu</a>
Michael Dorff	Brigham Young University, USA <a href="mailto:mdorff@math.byu.edu">mdorff@math.byu.edu</a>	Timothy E. O'Brien	Loyola University Chicago, USA <a href="mailto:tbriell@luc.edu">tbriell@luc.edu</a>
Sever S. Dragomir	Victoria University, Australia <a href="mailto:sever@matilda.vu.edu.au">sever@matilda.vu.edu.au</a>	Joseph O'Rourke	Smith College, USA <a href="mailto:orourke@cs.smith.edu">orourke@cs.smith.edu</a>
Behrouz Emamizadeh	The Petroleum Institute, UAE <a href="mailto:bemamizadeh@pi.ac.ae">bemamizadeh@pi.ac.ae</a>	Yuval Peres	Microsoft Research, USA <a href="mailto:peres@microsoft.com">peres@microsoft.com</a>
Joel Foisy	SUNY Potsdam <a href="mailto:foisys@potsdam.edu">foisys@potsdam.edu</a>	Y.-F. S. Pétermann	Université de Genève, Switzerland <a href="mailto:petermann@math.unige.ch">petermann@math.unige.ch</a>
Errin W. Fulp	Wake Forest University, USA <a href="mailto:fulp@wfu.edu">fulp@wfu.edu</a>	Robert J. Plemmons	Wake Forest University, USA <a href="mailto:rplemmons@wfu.edu">rplemmons@wfu.edu</a>
Joseph Gallian	University of Minnesota Duluth, USA <a href="mailto:kgallian@d.umn.edu">kgallian@d.umn.edu</a>	Carl B. Pomerance	Dartmouth College, USA <a href="mailto:carl.pomerance@dartmouth.edu">carl.pomerance@dartmouth.edu</a>
Stephan R. Garcia	Pomona College, USA <a href="mailto:stephan.garcia@pomona.edu">stephan.garcia@pomona.edu</a>	Vadim Ponomarenko	San Diego State University, USA <a href="mailto:vadim@sciences.sdsu.edu">vadim@sciences.sdsu.edu</a>
Anant Godbole	East Tennessee State University, USA <a href="mailto:godbole@etsu.edu">godbole@etsu.edu</a>	Bjorn Poonen	UC Berkeley, USA <a href="mailto:poonen@math.berkeley.edu">poonen@math.berkeley.edu</a>
Ron Gould	Emory University, USA <a href="mailto:rg@mathcs.emory.edu">rg@mathcs.emory.edu</a>	James Propp	U Mass Lowell, USA <a href="mailto:jpropp@cs.uml.edu">jpropp@cs.uml.edu</a>
Andrew Granville	Université Montréal, Canada <a href="mailto:andrew.andrew@dms.umontreal.ca">andrew.andrew@dms.umontreal.ca</a>	József H. Przytycki	George Washington University, USA <a href="mailto:przytyck@gwu.edu">przytyck@gwu.edu</a>
Jerrold Griggs	University of South Carolina, USA <a href="mailto:griggs@math.sc.edu">griggs@math.sc.edu</a>	Richard Rebarber	University of Nebraska, USA <a href="mailto:rrebarbe@math.unl.edu">rrebarbe@math.unl.edu</a>
Sat Gupta	U of North Carolina, Greensboro, USA <a href="mailto:sgupta@uncg.edu">sgupta@uncg.edu</a>	Robert W. Robinson	University of Georgia, USA <a href="mailto:rwr@cs.uga.edu">rwr@cs.uga.edu</a>
Jim Haglund	University of Pennsylvania, USA <a href="mailto:jhaglund@math.upenn.edu">jhaglund@math.upenn.edu</a>	Filip Saidak	U of North Carolina, Greensboro, USA <a href="mailto:f_saidak@uncg.edu">f_saidak@uncg.edu</a>
Johnny Henderson	Baylor University, USA <a href="mailto:johnny_henderson@baylor.edu">johnny_henderson@baylor.edu</a>	James A. Sellers	Penn State University, USA <a href="mailto:sellersj@math.psu.edu">sellersj@math.psu.edu</a>
Jim Hoste	Pitzer College <a href="mailto:jhoste@pitzer.edu">jhoste@pitzer.edu</a>	Andrew J. Sterge	Honorary Editor <a href="mailto:andy@ajsterge.com">andy@ajsterge.com</a>
Natalia Hritonenko	Prairie View A&M University, USA <a href="mailto:nahritonenko@pvamu.edu">nahritonenko@pvamu.edu</a>	Ann Trenk	Wellesley College, USA <a href="mailto:atrenk@wellesley.edu">atrenk@wellesley.edu</a>
Glenn H. Hurlbert	Arizona State University, USA <a href="mailto:hurlbert@asu.edu">hurlbert@asu.edu</a>	Ravi Vakil	Stanford University, USA <a href="mailto:vakil@math.stanford.edu">vakil@math.stanford.edu</a>
Charles R. Johnson	College of William and Mary, USA <a href="mailto:crjohnso@math.wm.edu">crjohnso@math.wm.edu</a>	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy <a href="mailto:antonia.vecchio@cnr.it">antonia.vecchio@cnr.it</a>
K. B. Kulasekera	Clemson University, USA <a href="mailto:kk@ces.clemson.edu">kk@ces.clemson.edu</a>	Ram U. Verma	University of Toledo, USA <a href="mailto:verma99@msn.com">verma99@msn.com</a>
Gerry Ladas	University of Rhode Island, USA <a href="mailto:gladas@math.uri.edu">gladas@math.uri.edu</a>	John C. Wierman	Johns Hopkins University, USA <a href="mailto:wierman@jhu.edu">wierman@jhu.edu</a>
		Michael E. Zieve	University of Michigan, USA <a href="mailto:zieve@umich.edu">zieve@umich.edu</a>

## PRODUCTION

Silvio Levy, Scientific Editor

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# involve

2015

vol. 8

no. 1

<a href="#">Efficient realization of nonzero spectra by polynomial matrices</a>	1
NATHAN MCNEW AND NICHOLAS ORMES	
<a href="#">The number of convex topologies on a finite totally ordered set</a>	25
TYLER CLARK AND TOM RICHMOND	
<a href="#">Nonultrametric triangles in diametral additive metric spaces</a>	33
TIMOTHY FAVER, KATELYNN KOCHALSKI, MATHAV KISHORE MURUGAN, HEIDI VERHEGGEN, ELIZABETH WESSON AND ANTHONY WESTON	
<a href="#">An elementary approach to characterizing Sheffer A-type 0 orthogonal polynomial sequences</a>	39
DANIEL J. GALIFFA AND TANYA N. RISTON	
<a href="#">Average reductions between random tree pairs</a>	63
SEAN CLEARY, JOHN PASSARO AND YASSER TORUNO	
<a href="#">Growth functions of finitely generated algebras</a>	71
ERIC FREDETTE, DAN KUBALA, ERIC NELSON, KELSEY WELLS AND HAROLD W. ELLINGSEN, JR.	
<a href="#">A note on triangulations of sumsets</a>	75
KÁROLY J. BÖRÖCZKY AND BENJAMIN HOFFMAN	
<a href="#">An exploration of ideal-divisor graphs</a>	87
MICHAEL AXTELL, JOE STICKLES, LANE BLOOME, ROB DONOVAN, PAUL MILNER, HAILEE PECK, ABIGAIL RICHARD AND TRISTAN WILLIAMS	
<a href="#">The failed zero forcing number of a graph</a>	99
KATHERINE FETCIE, BONNIE JACOB AND DANIEL SAAVEDRA	
<a href="#">An Erdős–Ko–Rado theorem for subset partitions</a>	119
ADAM DYCK AND KAREN MEAGHER	
<a href="#">Nonreal zero decreasing operators related to orthogonal polynomials</a>	129
ANDRE BUNTON, NICOLE JACOBS, SAMANTHA JENKINS, CHARLES MCKENRY JR., ANDRZEJ PIOTROWSKI AND LOUIS SCOTT	
<a href="#">Path cover number, maximum nullity, and zero forcing number of oriented graphs and other simple digraphs</a>	147
ADAM BERLINER, CORA BROWN, JOSHUA CARLSON, NATHANAEL COX, LESLIE HOGBEN, JASON HU, KATRINA JACOBS, KATHRYN MANTERNACH, TRAVIS PETERS, NATHAN WARNBERG AND MICHAEL YOUNG	
<a href="#">Braid computations for the crossing number of Klein links</a>	169
MICHAEL BUSH, DANIELLE SHEPHERD, JOSEPH SMITH, SARAH SMITH-POLDERMAN, JENNIFER BOWEN AND JOHN RAMSAY	