

involve

a journal of mathematics

On attractors and their basins

Alexander Arbieto and Davi Obata



On attractors and their basins

Alexander Arbieto and Davi Obata

(Communicated by Kenneth S. Berenhaut)

We prove that the map assigning to a given vector field the Lebesgue measure of the union of the basins of its attractors is lower semicontinuous in a residual subset of vector fields. Moreover, we prove that the Lebesgue measure of the union of the basins of attractors of a generic sectional axiom A vector field is total. For this, we also improve a result of Morales about sectional-hyperbolic sets. We also remark that homoclinic classes are topologically ergodic and that for a generic tame diffeomorphism, the union of the stable manifolds of the hyperbolic periodic orbits is dense in the manifold.

1. Introduction

One of the key notions in the theory of dynamical systems is that of attractors. By definition, an attractor captures the asymptotic information of a large set of orbits, called its basin, which always contains an open set. As an example, if an attractor is hyperbolic, then the asymptotic behavior of an orbit in its basin is governed by the dynamics of one orbit inside it (a shadowing property).

Moreover, that essentially every orbit is attracted by one attractor and that the set of attractors is finite (and possibly hyperbolic) implies that the dynamics of the system are nicely described by the attractors. For instance, this led Palis [2005] to conjecture that “there is a dense set D of dynamical systems such that any element of D has finitely many attractors whose union of basins of attraction has total probability”.

Mathematicians have made many efforts to understand attractors and their basins, not only for finite-dimensional dynamics, but also for PDEs (infinite-dimensional dynamical systems). See, for instance, [Constantin et al. 1985] or [Hale 2000].

On the other hand, to understand properties of the entire set of dynamical systems is a difficult task, and it is more reasonable to try to understand a large part of the set of dynamical systems. This reasoning leads to the theory of generic dynamics. Since

MSC2010: 37C10, 37C20.

Keywords: attractors, basin, sectional axiom A, sectional hyperbolic, basin of attraction.

Partially supported by CNPq, FAPERJ and PRONEX/DS from Brazil.

the C^r -topology turns the space of diffeomorphisms (or vector fields) into a Baire space, it is natural to show that some properties holds for a residual subset of the space of dynamical systems, i.e., a countable intersection of open and dense subsets, since this will show the presence of this property for a dense subset and this property could be used to show another property in another residual subset. Indeed, the intersection of two residual subsets is also a residual subset. Usually, we say that a property holds for a generic system if it holds in a residual subset of dynamical systems.

The purpose of this article is to give some remarks about attractors, and their basins, of certain classes of dynamical systems, both diffeomorphisms and vector fields. These remarks are the results obtained by Obata [2010], guided by Arbieto, in his undergraduate monograph. We will state the results and refer the reader to the next section for the precise definitions of the more technical objects used in the statements.

Let M be a Riemannian closed manifold. We denote by $\text{Diff}^1(M)$ the space of diffeomorphisms and by $\mathfrak{X}^1(M)$ the space of vector fields, both endowed with the C^1 -topology. We denote by m the Lebesgue measure and by d the geodesic distance, both induced by the Riemannian metric. If $X \in \mathfrak{X}^1(M)$, we denote by X_t the flow generated by X .

Results for flows. An attractor is an invariant compact subset Λ of M such that there exists a neighborhood U of Λ with

$$X_t(\bar{U}) \subset U \text{ for } t > 0 \quad \text{and} \quad \bigcap_{t \geq 0} X_t(U) = \Lambda.$$

The set U is called the *local basin* of Λ and $B(\Lambda) := \bigcup_{t \leq 0} X_t(U)$ is the *basin* of Λ . We also define a set R to be a repeller if R is an attractor for $-X$.

Let X be a vector field, and denote by $m(B(X))$ the Lebesgue measure of the union of the basins of the attractors of X . This generates a map $\Phi : \mathfrak{X}^1(M) \rightarrow [0, +\infty]$, defined as $\Phi(X) := m(B(X))$ if there exists an attractor and $\Phi(X) := 0$ if not.

Theorem 1. *There exists a residual subset \mathcal{R} such that $\Phi|_{\mathcal{R}}$ is lower semicontinuous.*

The analogous statement holds for diffeomorphisms using the same proof.

Metzger and Morales [2008] extended the notion of axiom A vector fields for flows with singularities, called *sectional axiom A* vector fields. As an intermediate step to studying sectional axiom A vector fields, we have the following result:

Theorem 2. *There exists a residual subset \mathcal{R} such that if X is in \mathcal{R} and $\Gamma = \Lambda_1 \cup \dots \cup \Lambda_k$, with $\Gamma \subset \Omega(X)$, is a disjoint union of homogeneous sectional-hyperbolic sets for X or $-X$, and Γ is a proper subset of M , then $m(\Gamma) = 0$.*

We remark that it is well known that if M is a closed manifold which is a sectional-hyperbolic set for X , then X has no singularities and X is Anosov [Bautista and Morales 2011].

To prove this theorem, we extend a result of [Morales 2007]; see Theorem 13.

As a corollary, we obtain the following result, which improves Corollary D of [Alves et al. 2007] in two ways. We do not require that the vector field be $C^{1+\varepsilon}$ or that the dimension of the manifold be 3. Indeed, in [Alves et al. 2007] it is proved that if a sectional axiom A vector field X over M^3 is $C^{1+\varepsilon}$, then the Lebesgue measure of the union of the basins of its hyperbolic or sectional-hyperbolic attractors is total. We remark also that the union of the sets of $C^{1+\varepsilon}$ vector fields, over any $\varepsilon > 0$, is a meager subset of vector fields.

Theorem 3. *Let X be a generic sectional axiom A vector field. Then either X is Anosov, or the Lebesgue measure of the nonwandering set of X is zero and the Lebesgue measure of the union of the basins of its attractors is total.*

A difficulty in proving this theorem is that it is not known whether the set of sectional axiom A vector fields (without cycles) is open. This is an interesting question. Even so, there are open sets of vector fields formed by sectional axiom A sets [Bautista and Morales 2011]. Moreover, [Morales and Pacifico 2003] shows that in dimension 3, generically, either a vector field has infinitely many sinks or sources or it is sectional axiom A. So, we obtain the following corollary:

Corollary 4. *If $\dim(M) = 3$, a generic vector field either has infinitely many sinks or sources, or the Lebesgue measure of the union of the basins of its attractors is total.*

Results for diffeomorphisms. Abdenur [2003] proved that attractors for generic diffeomorphisms are homoclinic classes. These classes are always transitive. However it can be proved that they have another property called topological ergodicity.¹

Proposition 5. *Any homoclinic class of a periodic point p , with period k , of a diffeomorphism f is topologically ergodic. Moreover, for any two open sets U and V , the density of $N(U, V) = \{i \geq 1 : f^i(U) \cap V \neq \emptyset\}$ is bounded by below $1/k$.*

Finally, the techniques used in the proof of the results above can be used to prove a folklore result. Since, as far as the authors know, it was never written, we include here a proof of this result:

Proposition 6. *If f is a C^1 -generic tame diffeomorphism, then the union of the stable manifolds of the hyperbolic periodic orbits is dense in M .*

We observe that this result was proved in a more general setting (partially hyperbolic diffeomorphisms with one-dimensional central bundle) by Bonatti, Gan and Wen [Bonatti et al. 2007]. In particular, they obtain this corollary using stronger

¹Recently Abdenur and Crovisier [2012] investigated the mixing property for isolated sets.

methods. However, the short proof given here only uses the connecting lemma. This is a particular case of Bonatti's conjecture; see [Bonatti et al. 2007].

Conjecture 7. *There exists a residual subset $\mathcal{R} \subset \text{Diff}^1(M)$ such that for any $f \in \mathcal{R}$, the union of the stable manifolds of the hyperbolic periodic orbits is dense in M .*

This paper is organized as follows. In Section 2, we give precise definitions of terms used in the introduction. In Section 3, we prove Theorem 1. In Section 4, we prove Theorems 2 and 3 and also prove an extension of a theorem by Morales. In Section 5, we give a proof of Proposition 5. Finally, in Section 6, we give a proof of Proposition 6.

2. Preliminaries

In this section, we give precise definitions of terms used in the introduction and collect some useful results.

2.1. Topology. As remarked before, both $\text{Diff}^1(M)$ and $\mathcal{X}^1(M)$ are Baire spaces. We will say that a property P is generic if it holds for a residual subset of these spaces. If the residual subset is fixed, we also say that an element of it is generic.

Let $F(M)$ denote the space of compact subsets of M ; it is a metric space under the *Hausdorff metric*, given by

$$d_H(A, B) = \max\{d_A(B), d_B(A)\} \quad \text{for all } A, B \in F(M),$$

where $d_A(B) = \max_{b \in B} \{\min_{a \in A} (d(a, b))\}$.

Let (N, d) be a metric space. A map $\varphi : N \rightarrow F(M)$ is *lower semicontinuous* at $y \in N$ if $y_n \rightarrow y$ implies $d_H(\varphi(y_n), \varphi(y)) \rightarrow 0$. Analogously, a map $\varphi : N \rightarrow \mathbb{R}$ is lower semicontinuous at $x_0 \in X$ if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0).$$

It is well known that if (N, d) is a Baire space, then the set of continuity points of a lower semicontinuous map, in either definition above, is a residual subset of its domain; see [Kelley 1955].

2.2. Flows. Let $X \in \mathcal{X}^1(M)$. The orbit of a point p is the set $\{X_t(p)\}_{t \in \mathbb{R}}$. A *periodic orbit* of X is an orbit $\{X_t(p) : t \in \mathbb{R}\}$ of a point $p \in M$ satisfying $X_T(p) = p$ for some minimal $T > 0$. A singularity σ is a zero of X . By a *closed orbit* we mean a periodic orbit or a singularity. The nonwandering set of X is the set $\Omega(X)$ of points x such that for every neighborhood U of x and $N > 0$, there exists some $T > N$ such that $X_T(U) \cap U \neq \emptyset$.

A subset $\Lambda \subset M$ is *invariant* if $X_t(\Lambda) = \Lambda$ for all $t \in \mathbb{R}$; *transitive* if there exists $p \in \Lambda$ such that its orbit is dense in Λ ; *isolated* if there exists a neighborhood U

of Λ such that $\bigcap_{t \in \mathbb{R}} X_t(U) = \Lambda$; and Ω -isolated if there exists a neighborhood V of Λ such that $\Omega(X) \cap V = \Lambda$. We remark that any attractor is Ω -isolated.

We say that a subset $\Lambda \subset M$ is sectional-hyperbolic if every singularity in Λ is hyperbolic and it has a nontrivial partially hyperbolic splitting $T_\Lambda M = E \oplus F$ such that E is uniformly contracting and F is sectionally expanding; i.e.,

$$\dim(E_x^c) \geq 2 \quad \text{and} \quad |\det(DX_t(x)/L_x)| \geq K^{-1}e^{\lambda t}$$

for all $x \in \Lambda, t \geq 0$, and L_x a two-dimensional subspace of E_x^c .

We say that Λ is *hyperbolic* if there is a continuous invariant tangent bundle decomposition

$$T_\Lambda M = \hat{E}_\Lambda^s \oplus \hat{E}_\Lambda^X \oplus \hat{E}_\Lambda^u,$$

and positive constants K, λ , where \hat{E}_Λ^X is the subbundle generated by X and

$$\|DX_t(x)/\hat{E}_x^s\| \leq Ke^{-\lambda t} \quad \text{and} \quad \|DX_{-t}(x)/\hat{E}_{X_t(x)}^u\| \leq Ke^{-\lambda t}$$

for all $x \in \Lambda$ and $t \geq 0$.

A closed orbit is hyperbolic if it is a hyperbolic compact invariant set. A hyperbolic set is a basic set if it is isolated and transitive. Similar notions hold for diffeomorphisms.

Given an invariant splitting $T_\Lambda M = E_\Lambda \oplus F_\Lambda$ over an invariant set Λ of a vector field X , we say that the subbundle E_Λ *dominates* F_Λ if there are positive constants K, λ such that

$$\|DX_t(x)/E_x\| \|DX_{-t}(x)/F_{X_t(x)}\| \leq Ke^{-\lambda t} \quad \text{for all } x \in \Lambda \text{ and } t \geq 0.$$

In such a case we say that $T_\Lambda M = E_\Lambda \oplus F_\Lambda$ is a *dominated splitting*.

We say that Λ is *partially hyperbolic* if it has a dominated splitting $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$ whose dominating subbundle E_Λ^s is *contracting*, that is,

$$\|DX_t(x)/E_x^s\| \leq Ke^{-\lambda t} \quad \text{for all } x \in \Lambda \text{ and } t \geq 0.$$

Moreover, we call the central subbundle E_Λ^c *sectionally expanding* if

$$\dim(E_x^c) \geq 2 \quad \text{and} \quad |\det(DX_t(x)/L_x)| \geq K^{-1}e^{\lambda t}$$

for all $x \in \Lambda, t \geq 0$, and L_x a two-dimensional subspace of E_x^c .

Definition 8. We say that a compact and invariant set Λ of X is *sectional-hyperbolic* if every singularity contained in Λ is hyperbolic and it has a nontrivial partially hyperbolic set with a sectionally expanding central subbundle.

Now, we recall the notion of sectional axiom A vector field, given in [Metzger and Morales 2008]; see also [Morales et al. 1999].

Definition 9. A vector field X is sectional axiom A if there is a finite disjoint decomposition

$$\Omega(X) = \Lambda_1 \cup \dots \cup \Lambda_k,$$

where each Λ_i is a hyperbolic basic set or a sectional-hyperbolic attractor up to time reversion.

2.3. Diffeomorphisms. If p is a hyperbolic periodic point, of period k , of a diffeomorphism f , then its stable manifold is the set

$$W^s(p) = \{y \in M : d(f^{kn}(y), p) \rightarrow 0 \text{ as } n \rightarrow \infty\}.$$

This set is in fact an immersed manifold. The stable manifold of the orbit of p is the union of the stable manifolds of $f^i(p)$ for $i = 0, \dots, k-1$, and it is denoted by $W^s(O(p))$. Analogously, we define the unstable manifold of p and the orbit of p .

Definition 10. The homoclinic class of p is the set

$$H(p, f) = \overline{W^s(O(p)) \cap W^u(O(p))}.$$

A diffeomorphism is *tame* if its nonwandering set decomposes as a finite number of homoclinic classes and finitely many sinks or sources. Analogous definitions hold for vector fields.

Given two nonempty open sets U and V , we define the set of times that the orbit of U visits V as

$$N(U, V) = \{i \geq 1 : f^i(U) \cap V \neq \emptyset\}.$$

The following definition can be found in [Abdenur and Crovisier 2012]:

Definition 11. An invariant and compact subset Λ of f is topologically ergodic if for every two nonempty open sets $U, V \subset \Lambda$, we have

$$\limsup_{n \rightarrow \infty} \frac{\#N(U, V) \cap \{1, \dots, n\}}{n} > 0.$$

3. Proof of Theorem 1

First we observe that since attractors are isolated, there are at most countably many of them. Let $\Lambda_1, \Lambda_2, \dots$ be the attractors of a generic vector field X . Denote by $B(\Lambda_1), B(\Lambda_2), \dots$ its basins.

We select $\Lambda_1, \dots, \Lambda_r$ such that

$$\sum_{i=1}^r m(B(\Lambda_i)) \geq m(B(X)) - \varepsilon.$$

There exist compact sets K_1, \dots, K_r such that $\Lambda_i \subset K_i \subset B(\Lambda_i)$ for $i = 1, \dots, r$ and such that

$$m(B(\Lambda_i) - K_i) < \frac{\varepsilon}{r}.$$

Now, we recall a result of Abdenur. Actually, he works with diffeomorphisms, but his proof holds for vector fields with the necessary adaptations. Also, he states his theorem for Ω -isolated transitive sets, but we will only state it in the case of attractors, which is the context here.

Theorem 12 [Abdenur 2003]. *There exists a residual subset $\mathfrak{R} \subset \mathfrak{X}^1(M)$ such that if $X \in \mathfrak{R}$ and Λ is an attractor of X with local basin U which does not reduce to a singularity, then there exists a neighborhood \mathfrak{U} of X such that for any $Y \in \mathfrak{U} \cap \mathfrak{R}$, $\Lambda(Y) = \bigcap_{t \geq 0} Y_t(U)$ is an attractor. Moreover, there exists a periodic orbit $O(p)$ such that $\Lambda(Y) = H(O(p), Y)$.*

Thus, there are local basins U_i of Λ_i such that these local basins persist in a C^1 -generic neighborhood of X . Since $B(\Lambda_i) = \bigcup_{t \geq 0} X_{-t}(U_i)$ and $K_i \subset B(\Lambda_i)$ is a compact set, there is $T > 0$ such that

$$K_i \subset \bigcup_{t \in [0, T]} X_{-t}(U_i).$$

The set on the right is open. So, if Y is C^1 -close to X , we obtain that

$$K_i \subset \bigcup_{t \in [0, T]} Y_{-t}(U_i).$$

Thus, if Y is generic and C^1 -close to X , we have $m(B(\Lambda_i(Y))) \geq m(K_i)$.

Hence,

$$m(B(Y)) \geq \sum_{i=1}^r m(K_i) \geq m(B(X)) - \varepsilon.$$

This proves lower semicontinuity.

4. Proof of Theorems 2 and 3

Let Λ be a sectional-hyperbolic set for X . We recall that its strong stable manifold is the set

$$W^{ss}(x) = \left\{ y \in M : \lim_{t \rightarrow \infty} d(X_t(x), X_t(y)) = 0 \right\}.$$

Its local strong stable manifold is an ε -ball $W_\varepsilon^{ss}(x)$ in $W^{ss}(x)$ centered at x for some $\varepsilon > 0$.

Given $A \subset M$, we define $\alpha(A)$ as the set of points $y = \lim_{n \rightarrow \infty} X_{t_n}(z_n)$ for some sequences $t_n \rightarrow -\infty$ and $z_n \in A$. We say that a sectional-hyperbolic set is homogeneous if the splitting $E^s \oplus E^c$ given by the definition is such that $\dim E^s$ is constant.

The following result improves the main theorem in [Morales 2007] since we do not require transitivity.

Theorem 13. *Let $\Lambda \subset \Omega(X)$ be a homogeneous sectional-hyperbolic set for X . Denote by R the union of the hyperbolic repellers contained in Λ . Then $\Lambda - R$ does not contain any local strong stable manifold.*

Proof. By hypothesis, the map $x \in \Lambda \mapsto W_\varepsilon^{\text{ss}}(x)$ is continuous if $\varepsilon > 0$ is small, but fixed. Assume that $\Lambda - R$ contains some $W_\varepsilon^{\text{ss}}(x)$. Let $\delta < \varepsilon$ and take $H = \alpha(W_\delta^{\text{ss}}(x)) \subset \Lambda - R$, which is compact and invariant. Observe also that the set $\Lambda - R$ is compact and invariant, since $\Lambda \subset \Omega(X)$.

If H has a singularity σ then, by definition, $\sigma = \lim X_{t_n}(z_n)$ for some sequences $t_n \rightarrow -\infty$ and $z_n \in W_\delta^{\text{ss}}(x)$. Moreover, $W_\delta^{\text{ss}}(X_{t_n}(z_n)) \subset \Lambda$ for any natural number n . Taking the limit as $n \rightarrow \infty$, we obtain that $W_\delta^{\text{ss}}(\sigma) \subset \Lambda$.

However, by [Bautista and Morales 2011], since Λ is sectional-hyperbolic, we have that $\Lambda \cap W^{\text{ss}}(\sigma) = \{\sigma\}$, and this is a contradiction.

If H does not have a singularity, then by the hyperbolic lemma [Bautista and Morales 2011], H is a hyperbolic set. Now, let y be a cluster point of $X_{t_n}(x)$, with $t_n \rightarrow -\infty$. We will show that $W^{\text{ss}}(y) \subset H$. Indeed, let $z \in W^{\text{ss}}(y)$ and let $\varepsilon > 0$ be small enough. There exists $T > 0$ such that

$$d(X_T(z), X_T(y)) < \varepsilon.$$

Also, there exists n_0 such that for any $n \geq n_0$, we have

$$d(X_t(y), X_{t_n+T}(x)) < \varepsilon.$$

Finally, for any n large, there exists $z_n \in W_\delta^{\text{ss}}(x)$ such that

$$d(X_{t_n+T}(x), X_{t_n+T}(z_n)) < \varepsilon.$$

This implies that if n is large enough then

$$d(X_T(z), X_{t_n+T}(z_n)) < \varepsilon.$$

In particular, we can assume that $t_n + T \rightarrow -\infty$. Thus, $X_T(z) \in H$, and by invariance, $z \in H$. Thus H is a repeller inside $\Lambda - R$, a contradiction. \square

Remark 14. We could remove the homogeneity assumption. Indeed, the sets $\{x \in \Lambda : \dim(E^s(x)) = i\}$ for $1 \leq i \leq d - 1$ are compact. Hence, we could use the argument restricting ourselves to each of these sets.

Now, we observe that X_1 is a partially hyperbolic diffeomorphism over Λ since the dominated splitting $T_\Lambda M = E^s \oplus E^c$ has a contracting subbundle E^s . A strong stable disk of X_1 is a disk which is tangent to the subbundle E^s over Λ . Obviously, a strong stable disk of X_1 is a local strong stable manifold for some point $x \in \Lambda$. However, the following result was proved in [Alves et al. 2007, Theorem 2.2]:

Theorem 15. *Let $f : M \rightarrow M$ be a C^2 diffeomorphism and $\Lambda \subset M$ a partially hyperbolic set with positive volume. Then Λ contains a strong stable disk.*

Together with Theorem 13, we obtain the following:

Corollary 16. *Let Λ be a proper subset of M . If Λ is a homogeneous sectional-hyperbolic set of a C^2 vector field X and $\Lambda \subset \Omega(X)$, then $m(\Lambda) = 0$.*

Proof. First we remark that there are only countably many repellers in Λ , since they are isolated. Moreover, by [Bowen 1975], the measure of any hyperbolic repeller (or attractor) is zero if X is C^2 .

On the other hand, if R denotes the union of the hyperbolic repellers of Λ and $m(\Lambda - R) > 0$, then by Theorem 15, there exists a strong stable disk on $\Lambda - R$, and this contradicts Theorem 13. □

For any open set U , let $\Lambda_Y(U) = \bigcap_{t \in \mathbb{R}} Y_t(\bar{U})$. These sets have an upper semi-continuity property: $\limsup \Lambda_{X_n}(U) \subset \Lambda_X(U)$. Indeed, let $x \in \limsup \Lambda_{X_n}(U)$. So, there exists $x_n \in \Lambda_{X_n}(U)$ such that $x_n \rightarrow x$. Fix $t \in \mathbb{R}$. We have $(X_n)_t(x_n) \in \bar{U}$. Thus, $X_t(x) \in \bar{U}$. Since this holds for every $t \in \mathbb{R}$, this implies that $x \in \Lambda_X(U)$.

Now, let $\{U_k\}$ be a countable basis of the topology and $\{O_k\}$ the set of finite unions of the U_k . For every $n, k \in \mathbb{N}$, we define $\mathcal{U}_{n,k}$ as the set of vector fields Y such that $m(\Lambda_Y(O_k)) < 1/n$.

Lemma 17. *$\mathcal{U}_{n,k}$ is an open set.*

Proof. Let $Y \in \mathcal{U}_{n,k}$, and suppose that $m(\Lambda_Y(O_k)) = 1/n - \varepsilon$. There exists T large enough that

$$m\left(\bigcap_{t=-T}^T Y_t(\bar{O}_k)\right) < m(\Lambda_Y(O_k)) + \frac{\varepsilon}{2}.$$

Let W be a neighborhood of $\bigcap_{t=-T}^T Y_t(\bar{O}_k)$ such that

$$m(W) < m\left(\bigcap_{t=-T}^T Y_t(\bar{O}_k)\right) + \frac{\varepsilon}{2}.$$

If Z is close enough to Y , we have that $\bigcap_{t=-T}^T Z_t(\bar{O}_k) \subset W$. Thus

$$\begin{aligned} m(\Lambda_Z(O_k)) &\leq m\left(\bigcap_{t=-T}^T Z_t(\bar{O}_k)\right) < m\left(\bigcap_{t=-T}^T Y_t(\bar{O}_k)\right) + \frac{\varepsilon}{2} \\ &\leq m(\Lambda_Y(O_k)) + \varepsilon = \frac{1}{n}. \end{aligned} \quad \square$$

Now, we prove Theorem 2.

Proof of Theorem 2. By the previous lemma, $\mathcal{U}_{n,k}$ is an open set. Now, we define $\mathcal{N}_{n,k} = \mathcal{X}^1(M) - \overline{\mathcal{U}_{n,k}}$. Consider the residual subset

$$\mathcal{R} = \bigcap_n \bigcap_k (\mathcal{U}_{n,k} \cup \mathcal{N}_{n,k}).$$

Let $X \in \mathcal{R}$ and let $\Gamma = \Lambda_1 \cup \dots \cup \Lambda_k$, as in the statement of Theorem 2. Suppose that Λ_i is a homogeneous sectional-hyperbolic set for X . Since Λ_i is invariant, there exists $k(i)$ such that $\Lambda_i \subset \Lambda_X(\mathcal{O}_{k(i)})$ and $\Lambda_X(\mathcal{O}_{k(i)})$ is a homogeneous sectional-hyperbolic set. A similar argument holds when Λ_i is a homogeneous sectional-hyperbolic set for $-X$.

Now, suppose that $m(\Lambda_X(\mathcal{O}_{k(i)})) > 0$ for some i . Thus, there exists n such that $m(\Lambda_X(\mathcal{O}_{k(i)})) \geq 1/n$. So, $X \in \mathcal{N}_{n,k(i)}$. Since $\mathcal{N}_{n,k(i)}$ is an open set, there exists a neighborhood \mathcal{V} of X such that $m(\Lambda_Y(\mathcal{O}_{k(i)})) \geq 1/n$ for every $Y \in \mathcal{V}$.

Using the semicontinuity property, mentioned above, and the sectional hyperbolicity of $\Lambda_X(\mathcal{O}_{k(i)})$, we can assume, shrinking \mathcal{V} if necessary, that $\Lambda_Y(\mathcal{O}_{k(i)})$ is a homogeneous sectional-hyperbolic set for every $Y \in \mathcal{V}$.

Now, we can choose a C^2 vector field $Y \in \mathcal{V}$ and by Corollary 16, we have that $m(\Lambda_Y(\mathcal{O}_k)) = 0$, a contradiction. □

Proof of Theorem 3. The arguments given above show that there exists a residual subset \mathcal{S} such that if $X \in \mathcal{S}$ and Λ is a proper saddle-type isolated transitive sectional-hyperbolic set, then $m(\Lambda) = 0$.

Indeed, let U be an open set, and define $\mathcal{U}(U)$ as the (open) set formed by vector fields Y such that $\Lambda_Y(U)$ is hyperbolic of saddle type. Let $\mathcal{U}_n(U) = \{Y \in \mathcal{U}(U) : m(B(\Lambda_Y(U))) < 1/n\}$. Using the same argument as in the proof of Lemma 17, we obtain that $\mathcal{U}_n(U)$ is an open set.

Moreover, if $Y \in \mathcal{U}(U)$ is C^2 , we have that $m(B(\Lambda_Y(U))) = 0$ [Bowen 1975, p. 68]. So, $\mathcal{U}_n(U)$ is dense in $\mathcal{U}(U)$.

Defining \mathcal{O}_k as above, we set $\mathcal{S} = \bigcap_{k,n} \mathcal{U}_n(\mathcal{O}_k)$.

Let $X \in \mathcal{R} \cap \mathcal{S}$ be a sectional axiom A vector field. By definition, we have a spectral decomposition $\Omega(X) = \Lambda_1 \cup \dots \cup \Lambda_k$, formed by sectional-hyperbolic attractors, repellers and basic saddle-type hyperbolic sets. Moreover, since these sets have a dense orbit, they are homogeneous.

If $m(\Omega(X)) > 0$, then there exists $1 \leq i \leq k$ such that $m(\Lambda_i) > 0$. By the previous argument and Theorem 2, we have that $\Lambda_i = M$. If Λ_i is a saddle-type hyperbolic set, then X is Anosov. If Λ_i is a sectional-hyperbolic attractor then it cannot have any singularity. Indeed, if σ is a singularity, we must have $W^{ss}(\sigma) \cap \Lambda_i = \{\sigma\}$, but if $\Lambda_i = M$ this cannot be true. Hence, by the hyperbolic lemma, M would be hyperbolic again and X would be Anosov. If Λ_i is a sectional-hyperbolic attractor for $-X$, the same holds.

So, assuming that X is not Anosov, $m(\Omega(X)) = 0$. Using Lemma 2.2 of [Shub 1978], we have that

$$M = W^s(\Lambda_1) \cup \dots \cup W^s(\Lambda_k).$$

Since $X \in \mathcal{R}$, if Λ_i is a repeller then $W^s(\Lambda_i) = \Lambda_i$ and $m(\Lambda_i) = 0$. Since $X \in \mathcal{S}$, if Λ_i is a hyperbolic basic set then $m(W^s(\Lambda_i)) = 0$. Thus, the measure of the union of the basins of the attractors is total. \square

5. Proof of Proposition 5

In the following, we will work with the topology relative to the homoclinic class. First, we will show that any homoclinic class is topologically ergodic.²

Let $H(p, f)$ be a homoclinic class. Denote by k the period of p . We recall that the local stable manifold of p is the set $W_\varepsilon^s(p) = \{y \in M : d(f^n(y), f^n(p)) \leq \varepsilon\}$.

Fix two nonempty open subsets U and V of $H(p, f)$. Since the stable manifold of its orbit is dense, there exist $\varepsilon > 0$ and $N > 0$ such that

$$f^{-N}(W_\varepsilon^s(p)) \cap U \neq \emptyset.$$

In particular, there exists a disk $D \subset f^N(U)$ transversal to $W_\varepsilon^s(p)$. Moreover, since $W^u(O(p))$ is dense, there exists $K > 0$ such that $f^K(W_\varepsilon^u(p)) \cap V \neq \emptyset$. Using the λ -lemma [Palis and de Melo 1982], there exist m_0 and $0 \leq i < k$ such that for every $m \geq m_0$, we have that $f^{km+i}(D) \cap V \neq \emptyset$. Let $l \in \mathbb{N}$ such that $A = lk + (N + i)$ is the largest integer less than or equal to n ; in particular, $n < A + k$. By the previous remark, $\#N(U, V) \cap \{1, \dots, n\} \geq l - m_0$. So,

$$\limsup_{n \rightarrow \infty} \frac{\#N(U, V) \cap \{1, \dots, n\}}{n} \geq \limsup_{l \rightarrow \infty} \frac{l - m_0}{(l + 1)k + N + i} = \frac{1}{k}.$$

This shows that the homoclinic class is topologically ergodic.

6. Proof of Proposition 6

We recall that an invariant and compact subset $A \subset M$ is called Lyapunov stable if given U , an open neighborhood of A , there exists another neighborhood V of A such that $f^n(V) \subset U$ for every $n \in \mathbb{N}$.

Lemma 18 [Carballo et al. 2003, Lemma 3.4]. *If f is a C^1 -generic diffeomorphism, then $\overline{W^u(O(p))}$ is Lyapunov stable for f .*

Another source of Lyapunov stable sets is the following, which is [Morales and Pacifico 2002, Theorem A]:

Theorem 19. *There exists a residual subset $R^* \subset \text{Diff}^1(M)$ such that if $g \in R^*$, then the set $S = \{x \in M : \omega(x) \text{ is Lyapunov stable}\}$ is a residual subset of M .*

²We want to thank Professor Abdenur for pointing out this short argument to us.

We also recall Hayashi’s connecting lemma [1997], one of the most useful techniques in the C^1 -generic theory of dynamical systems. The formulation that we give here is taken from [Wen and Xia 2000].

Theorem 20 (connecting lemma). *Let $f \in \text{Diff}^1(M)$, and let z be a nonperiodic point of f . Given a neighborhood \mathcal{U} of f , there exist $\rho > 1$, $L \in \mathbb{N}$ and $\delta_0 > 0$ with the following property. Let $0 < \delta < \delta_0$ and*

$$p, q \notin \Delta(\delta) := \bigcup_{n=1}^L (f^{-n}(B(z, \delta))).$$

If there exist $a > L$ such that $f^a(p) \in B(z, \delta/\rho)$ and $b \geq 0$ such that $f^{-b}(q) \in B(z, \delta/\rho)$, then there exists $g \in \mathcal{U}$ such that q is a future g -iterate of p and $g \equiv f$ outside $\Delta(\delta)$.

We remark that the method used in the proof of Theorem 1 could be used to prove the topological semicontinuity of the basins of generic attractors. However, in the C^1 -topology a stronger property can be obtained, which, together with the continuity given by the stable manifold theorem, quickly implies this semicontinuity in this topology.

Proposition 21. *C^1 -generically, if a diffeomorphism has an attractor, then there exists a periodic point inside the attractor such that its stable manifold is dense in the basin of the attractor.*

Proof. Let U be an open set. We define the set

$$\mathcal{U}(U, m) = \left\{ f \in \text{Diff}^1(M) : \exists p \in \bigcap_{n \geq 0} f^n(U) \cap \text{Per}_h(f) \text{ with } W^s(p, f) \text{ } 1/m\text{-dense in } U \right\}.$$

If $f \in \mathcal{U}(U, m)$, then it has a hyperbolic periodic point in U such that its stable manifold is $1/m$ -dense in U . Since this point is hyperbolic, there exists V , a C^1 -neighborhood of f such that if $g \in V$ then $p(g) \in U$. Take $y \in U$ and $B = B(y, 1/m)$, so for f , we have that $W^s(p, f) \cap B \neq \emptyset$. By the stable manifold theorem, we have that $W^s(p(g)) \cap B \neq \emptyset$, so $W^s(p(g))$ is $1/m$ -dense in U and $g \in \mathcal{U}(U, m)$. This proves that the set $\mathcal{U}(U, m)$ is open.

Let $\{U_k\}$ be a countable basis of open sets of M , and let $\{O_n\}$ be the set of all possible unions of the elements U_k . Define

$$A(O_n, m) = \mathcal{U}(O_n, m) \cup \overline{\mathcal{U}(O_n, m)^c}.$$

Now, by the previous remark, and by construction, this set is open and dense in $\text{Diff}^1(M)$. So $R_1 = \bigcap_{n,m} A(O_n, m)$ is a residual subset. Let R_2 be the residual subset given in [Abdenur 2003].

Let $R = R_1 \cap R_2$. If $f \in R$ and Λ is an attractor of f , then there exists $p \in \text{Per}_h(f)$ such that $\Lambda = H(p, f)$. Fix n, m such that O_n is a local basin of Λ . Now, we must prove that $f \in \mathcal{U}(O_n, m)$. Suppose that $f \in \overline{\mathcal{U}(O_n, m)^c}$. Since this set is open, there is $W \subset \overline{\mathcal{U}(O_n, m)^c}$, a small open C^1 -neighborhood of f . The next step is to prove that we can find $g \in W$ such that $g \in \mathcal{U}(O_n, m)$, which will be a contradiction. To prove this we will use the C^1 -connecting lemma, and we will also need the following lemmas. From now on we will fix f and W as above.

Lemma 22. *The function $\Phi(g) = \overline{W^s(p(g), g)}$ for $g \in W$ is continuous in a residual subset of W .*

Proof. The map Φ is lower semicontinuous in W by the stable manifold theorem. Then, it is continuous in a residual subset $W^* \subset W$. □

Thus we have that the map Φ is continuous in $W^* \cap R$. Now, since $f \in \overline{\mathcal{U}(O_n, m)^c}$, there exists an $x \in O_n$ such that $B(x, 1/m) \cap \Phi(f) = \emptyset$.

This, together with Theorem 19, implies the following corollary:

Corollary 23. *There exists a residual subset $R_W \subset W$ such that if $g \in R_W$, then there exists a residual subset $P \subset O_m$ such that if $x \in P$ then $\omega(x) = \Lambda(g)$.*

Proof. Let R^* and S be given by Theorem 19. Define $R_W := R \cap R^* \cap W$ and $P = O_n \cap S$. Hence, if $x \in P$, then $x \in O_n$ and $\omega(x) \subset \Lambda$. However, since $x \in S$ as well, we know that $\omega(x)$ is Lyapunov stable. By the previous remark, since Λ is transitive, we have that $\Lambda \subset \omega(x)$. Thus $\omega(x) = \Lambda$. □

Now, we study the consequences of the continuity of Φ .

Lemma 24. *If Φ is continuous in $g \in R_W$ and S is the set given by Theorem 19, then $O_n \cap S \subset \Phi(g)$.*

Proof. If the lemma does not hold, then there exists $x \in (O_n \cap S) - \Phi(g)$. Let U be a neighborhood of $\Phi(g)$ such that $x \notin U$. By continuity there exists a neighborhood \mathcal{V} of g such that if $h \in \mathcal{V}$ then $\Phi(h) \subset U$.

Since $x \in O_n \cap S$, we have $\omega(x) = \Lambda(g)$. Thus, there exists a sequence $(l_n) \subset \mathbb{N}$ such that $g^{l_n}(x) \rightarrow p(g)$. By the Hartman–Grobman theorem [Palis and de Melo 1982], there exists another sequence $(t_n) \subset \mathbb{N}$ such that

$$g^{t_n}(x) \rightarrow q \in W_\varepsilon^s(p(g), g) - \{p(g)\}.$$

Let $\rho > 1$, $L \in \mathbb{N}$ and $\delta_0 > 0$, as given by the C^1 -connecting lemma applied to q and \mathcal{U} . Choose δ with $0 < \delta < \delta_0$, and let V be a neighborhood of the orbit of $p(g)$ such that

$$p(g), x \notin \Delta(\delta) = \bigcup_{n=1}^L (g^{-n}(B(q, \delta)))$$

and

$$\bigcup_{n=1}^L (g^{-n}(B(q, \delta))) \cap V = \emptyset.$$

Pick $y \in B(q, \delta/\rho) \cap W^s(p(g), g)$ such that, defining $z = g^k(y)$, we have $z \in (W^s(p(g), g) - \{p(g)\}) \cap V$. By definition, we have that $g^{-k}(z) = y \in B(q, \delta/\rho)$. Using that $g^{t_n}(x) \rightarrow q$, we obtain some $n_0 > L$ such that

$$g^{t_{n_0}}(x) \in B(q, \delta/\rho).$$

Applying the C^1 -connecting lemma, we obtain $h \in \mathcal{V}$ such that $h = g$ outside of $\Delta(\delta)$ and x belongs to the h -negative orbit of z . However, since $z \in (W_g^s(p(g)) - \{p(g)\}) \cap V$, we obtain that the h -positive orbit of z belongs to V .

Thus

$$z \in W_h^s(p(h)) \quad \text{and thus} \quad x \in W_h^s(p(h)).$$

This leads to a contradiction, since $h \in \mathcal{V}$ and $x \notin U$. \square

By the previous lemma, since $f \in W$, there is $g \in R_W$ such that Φ is continuous in g . So $O_n \cap S \subset \Phi(g)$, and there exists $y \in B(x, 1/m) \cap S$. Then $y \in \Phi(g)$, which is a contradiction since $f \in W \subset \overline{\mathcal{U}(O_n, m)^c}$. Then $f \in \mathcal{U}(O_n, m)$, which proves the proposition. \square

Now, to prove Proposition 6, it is enough to combine Proposition 21 with:

Theorem 25 [Carballo and Morales 2003]. *If f is a C^1 -generic tame diffeomorphism then the union of the basins of its attractors is an open and dense subset of M .*

References

- [Abdenur 2003] F. Abdenur, “Attractors of generic diffeomorphisms are persistent”, *Nonlinearity* **16**:1 (2003), 301–311. MR 2003k:37040 Zbl 1023.37007
- [Abdenur and Crovisier 2012] F. Abdenur and S. Crovisier, “Transitivity and topological mixing for C^1 diffeomorphisms”, pp. 1–16 in *Essays in mathematics and its applications*, edited by P. M. Pardalos and T. M. Rassias, Springer, Heidelberg, 2012. MR 2975581
- [Alves et al. 2007] J. F. Alves, V. Araújo, M. J. Pacifico, and V. Pinheiro, “On the volume of singular-hyperbolic sets”, *Dyn. Syst.* **22**:3 (2007), 249–267. MR 2008k:37069 Zbl 1226.37014
- [Bautista and Morales 2011] S. Bautista and C. A. Morales, “Lectures on sectional-Anosov flows”, IMPA preprint series D 86/2011, Instituto Nacional de Matemática Pura e Aplicada, Rio de Janeiro, 2011, available at http://preprint.impa.br/Shadows/SERIE_D/2011/86.html.
- [Bonatti et al. 2007] C. Bonatti, S. Gan, and L. Wen, “On the existence of non-trivial homoclinic classes”, *Ergodic Theory Dynam. Systems* **27**:5 (2007), 1473–1508. MR 2009d:37036 Zbl 1128.37021
- [Bowen 1975] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Math. **470**, Springer, Berlin, 1975. MR 56 #1364 Zbl 0308.28010

- [Carballo and Morales 2003] C. M. Carballo and C. A. Morales, “Homoclinic classes and finitude of attractors for vector fields on n -manifolds”, *Bull. London Math. Soc.* **35**:1 (2003), 85–91. MR 2003h:37021 Zbl 1035.37007
- [Carballo et al. 2003] C. M. Carballo, C. A. Morales, and M. J. Pacifico, “Homoclinic classes for generic C^1 vector fields”, *Ergodic Theory Dynam. Systems* **23**:2 (2003), 403–415. MR 2004e:37031 Zbl 1047.37009
- [Constantin et al. 1985] P. Constantin, C. Foias, and R. Temam, *Attractors representing turbulent flows*, Mem. Amer. Math. Soc. **53**:314, American Mathematical Society, Providence, RI, 1985. MR 86m:35137 Zbl 0567.35070
- [Hale 2000] J. K. Hale, “Dissipation and attractors”, pp. 622–637 in *International Conference on Differential Equations* (Berlin, 1999), vol. 1, edited by B. Fiedler et al., World Scientific, River Edge, NJ, 2000. MR 1870207 Zbl 0971.37037
- [Hayashi 1997] S. Hayashi, “Connecting invariant manifolds and the solution of the C^1 stability and Ω -stability conjectures for flows”, *Ann. of Math. (2)* **145**:1 (1997), 81–137. MR 98b:58096 Zbl 0871.58067
- [Kelley 1955] J. L. Kelley, *General topology*, Van Nostrand, Toronto, ON, 1955. Reprinted in Grad. Texts in Math. **27**, Springer, New York, 1995. MR 16,1136c Zbl 0066.16604
- [Metzger and Morales 2008] R. Metzger and C. A. Morales, “Sectional-hyperbolic systems”, *Ergodic Theory Dynam. Systems* **28**:5 (2008), 1587–1597. MR 2010g:37045 Zbl 1165.37010
- [Morales 2007] C. A. Morales, “Strong stable manifolds for sectional-hyperbolic sets”, *Discrete Contin. Dyn. Syst.* **17**:3 (2007), 553–560. MR 2008a:37036 Zbl 1137.37015
- [Morales and Pacifico 2002] C. A. Morales and M. J. Pacifico, “Lyapunov stability of ω -limit sets”, *Discrete Contin. Dyn. Syst.* **8**:3 (2002), 671–674. MR 2003b:37024 Zbl 1162.37302
- [Morales and Pacifico 2003] C. A. Morales and M. J. Pacifico, “A dichotomy for three-dimensional vector fields”, *Ergodic Theory Dynam. Systems* **23**:5 (2003), 1575–1600. MR 2005a:37030 Zbl 1040.37014
- [Morales et al. 1999] C. A. Morales, M. J. Pacifico, and E. R. Pujals, “Singular hyperbolic systems”, *Proc. Amer. Math. Soc.* **127**:11 (1999), 3393–3401. MR 2000c:37034 Zbl 0924.58068
- [Obata 2010] D. J. Obata, “Resultados na teoria de dinâmica genérica”, undergraduate monograph, Federal University of Rio de Janeiro, 2010, available at <http://im.ufrj.br/~arbiето/davimono.pdf>.
- [Palis 2005] J. Palis, Jr., “A global perspective for non-conservative dynamics”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**:4 (2005), 485–507. MR 2006b:37037 Zbl 1143.37016
- [Palis and de Melo 1982] J. Palis, Jr. and W. de Melo, *Geometric theory of dynamical systems: an introduction*, Springer, New York, 1982. MR 84a:58004 Zbl 0491.58001
- [Shub 1978] M. Shub, *Stabilité globale des systèmes dynamiques*, Astérisque **56**, Société Mathématique de France, Paris, 1978. Translated as *Global stability of dynamical systems*, Springer, New York, 1987. MR 80c:58015 Zbl 0396.58014
- [Wen and Xia 2000] L. Wen and Z. Xia, “ C^1 connecting lemmas”, *Trans. Amer. Math. Soc.* **352**:11 (2000), 5213–5230. MR 2001b:37024 Zbl 0947.37018

Received: 2011-12-06 Revised: 2013-12-12 Accepted: 2014-01-08

arbiето@im.ufrj.br

Instituto de Matemática, Universidade Federal do Rio de Janeiro, P. O. Box 68530, 21945-970 Rio de Janeiro, Brazil

davi.obata@gmail.com

Instituto de Matemática, Universidade Federal do Rio de Janeiro, P. O. Box 68530, 21945-970 Rio de Janeiro, Brazil

involve

msp.org/involve

EDITORS

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moselehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsgdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA jgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sngupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

PRODUCTION


Silvio Levy, Scientific Editor

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

involve

2015

vol. 8

no. 2

Enhancing multiple testing: two applications of the probability of correct selection statistic	181
ERIN IRWIN AND JASON WILSON	
On attractors and their basins	195
ALEXANDER ARBIETO AND DAVI OBATA	
Convergence of the maximum zeros of a class of Fibonacci-type polynomials	211
REBECCA GRIDER AND KRISTI KARBER	
Iteration digraphs of a linear function	221
HANNAH ROBERTS	
Numerical integration of rational bubble functions with multiple singularities	233
MICHAEL SCHNEIER	
Finite groups with some weakly s -permutably embedded and weakly s -supplemented subgroups	253
GUO ZHONG, XUANLONG MA, SHIXUN LIN, JIAYI XIA AND JIANXING JIN	
Ordering graphs in a normalized singular value measure	263
CHARLES R. JOHNSON, BRIAN LINS, VICTOR LUO AND SEAN MEEHAN	
More explicit formulas for Bernoulli and Euler numbers	275
FRANCESCA ROMANO	
Crossings of complex line segments	285
SAMULI LEPPÄNEN	
On the ε -ascent chromatic index of complete graphs	295
JEAN A. BREYTENBACH AND C. M. (KIEKA) MYNHARDT	
Bisection envelopes	307
NOAH FECHTOR-PRADINES	
Degree 14 2-adic fields	329
CHAD AWTRY, NICOLE MILES, JONATHAN MILSTEAD, CHRISTOPHER SHILL AND ERIN STROSNIDER	
Counting set classes with Burnside's lemma	337
JOSHUA CASE, LORI KOBAN AND JORDAN LEGRAND	
Border rank of ternary trilinear forms and the j -invariant	345
DEREK ALLUMS AND JOSEPH M. LANDSBERG	
On the least prime congruent to 1 modulo n	357
JACKSON S. MORROW	



1944-4176(2015)8:2;1-5