

involve

a journal of mathematics

Linear symplectomorphisms
as R -Lagrangian subspaces

Chris Hellmann, Brennan Langenbach and Michael VanValkenburgh



Linear symplectomorphisms as R -Lagrangian subspaces

Chris Hellmann, Brennan Langenbach and Michael VanValkenburgh

(Communicated by Ravi Vakil)

The graph of a real linear symplectomorphism is an R -Lagrangian subspace of a complex symplectic vector space. The restriction of the complex symplectic form is thus purely imaginary and may be expressed in terms of the generating function of the transformation. We provide explicit formulas; moreover, as an application, we give an explicit general formula for the metaplectic representation of the real symplectic group.

1. Introduction

1.1. Overview. As part of our symplectic upbringing, our ancestors impressed upon us the Symplectic Creed:

Everything is a Lagrangian submanifold [Weinstein 1981].

Obviously false if taken literally, rather than a “creed” it might be called the Maslow–Weinstein hammer, or, in French, *la déformation professionnelle symplectique*, saying that “if all you have is a [symplectic form], everything looks like a [Lagrangian submanifold],” or, in other words, to a symplectic geometer, everything should be expressed in terms of Lagrangian submanifolds. In this paper we consider a vector space endowed with *two* symplectic forms, namely the real and imaginary parts $\operatorname{Re} \omega^{\mathbb{C}}$ and $\operatorname{Im} \omega^{\mathbb{C}}$ of a complex symplectic form $\omega^{\mathbb{C}}$, and begin with the simple observation that

Not every Lagrangian submanifold [with respect to $\operatorname{Re} \omega^{\mathbb{C}}$] *is a Lagrangian submanifold* [with respect to $\operatorname{Im} \omega^{\mathbb{C}}$].

We study its implications for the classification of real linear symplectomorphisms \mathcal{H} , as the graph of \mathcal{H} is essentially by definition a Lagrangian subspace with respect to $\operatorname{Re} \omega^{\mathbb{C}}$; we ask, with some abuse of language:

MSC2010: 37J10, 51A50, 70H15, 81S10.

Keywords: complex symplectic linear algebra, linear symplectomorphisms, Lagrangian submanifolds, the metaplectic representation.

Open problem. Is every $2n \times 2n$ skew-symmetric matrix of the form $\text{Im } \omega^{\mathbb{C}}|_{\text{graph } \mathcal{H}}$ for some \mathcal{H} ?

We believe that an answer would shed some light on the structure of linear symplectomorphisms. While our primary reason for writing this article is to precisely formulate the above open problem, which we do in [Section 1.2](#), our primary technical result is to rewrite it in terms of generating functions; after all, if one guiding principle is the Symplectic Creed, another is that “symplectic topology is the geometry of generating functions” [[Viterbo 1992](#)]. Or, to go further back, while Sir William Rowan Hamilton first conceived of generating functions (or as he called them, *characteristic functions*) as mathematical tools in his symplectic formulation of optics, he later found, in his symplectic formulation of classical mechanics, that the generating function for a physical system is the least action function, in a sense that we will not make precise [[Abraham and Marsden 1978](#); [Hamilton 1834](#)]; this gives a striking connection with the calculus of variations. Moreover, in Fresnel optics and quantum mechanics, the generating function is used as the phase function of an oscillatory integral operator; the integral operator is said to “quantize” the corresponding symplectomorphism [[Grigis and Sjöstrand 1994](#); [Guillemin and Sternberg 1984](#)]. (Loosely speaking, when differentiating the integral, one finds that the phase function must satisfy the Hamilton–Jacobi equation.) This topic will be touched upon in [Section 3](#). For us, the generating function corresponding to the linear symplectomorphism \mathcal{H} is the scalar-valued function Φ in our main theorem:

Theorem 1. *For each $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$ there exists a quadratic form $\Phi : \mathbb{C}^n \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$ such that*

$$\text{graph}_{\mathbb{C}} \mathcal{H} = \left\{ \left(z, -2 \frac{\partial \Phi}{\partial z}(z, \theta) \right) : \frac{\partial \Phi}{\partial \theta}(z, \theta) = 0 \right\},$$

and the restriction of $\omega^{\mathbb{C}}$ to $\text{graph}_{\mathbb{C}} \mathcal{H}$ is given by

$$\begin{aligned} & \omega^{\mathbb{C}} \left(\left(z, -2 \frac{\partial \Phi}{\partial z}(z, \theta) \right), \left(w, -2 \frac{\partial \Phi}{\partial z}(w, \eta) \right) \right) \\ &= 2 \sum_{j=1}^n \sum_{\ell=1}^{2n} \frac{\partial^2 \Phi}{\partial z_j \partial \theta_{\ell}} (z_j \eta_{\ell} - w_j \theta_{\ell}) + 2 \sum_{j,m=1}^n \frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_m} (z_j \bar{w}_m - w_j \bar{z}_m). \quad (1) \end{aligned}$$

Moreover, our construction provides an explicit general formula for Φ .

Our notation will be explained in the following subsection, along with the necessary background and a restatement of the open problem. We prove the theorem in [Section 2](#), and in [Section 3](#) we show how our construction seems to adequately answer a question of Folland [[1989](#)] regarding the metaplectic representation. We

conclude with a broad indication of future work. In the [Appendix](#) we give additional linear-algebraic background and some new elementary results relevant to our problem, and also give an additional restatement of our open problem.

1.2. Background and restatement of the problem. In a real symplectic vector space there is already a natural complex structure; the model example is \mathbb{R}^{2n} with the $2n \times 2n$ matrix $\mathcal{J} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$, where of course $\mathcal{J}^2 = -I$. What we mean by “complex symplectic linear algebra” is something else; we instead consider \mathbb{C}^{2n} with the above matrix \mathcal{J} , that is, we consider

$$\omega^{\mathbb{C}} = \sum_{j=1}^n d\zeta_j \wedge dz_j \quad \text{on } \mathbb{C}_z^n \times \mathbb{C}_\zeta^n$$

(a nondegenerate alternating bilinear form over \mathbb{C}). The basic formalism of complex symplectic linear algebra is not new; indeed, complex symplectic structures naturally appear in the theory of differential equations and have been studied through that lens (see, for example, [Schapira 1981] and [Sjöstrand 1982], or [Everitt and Markus 2004] for another perspective). The point of view of this paper is that elementary linear-algebraic aspects remain unexplored in the complex case and may help us better understand the real case.

A *symplectic vector space* over a field¹ K is by definition a pair (V, ω) , where V is a finite-dimensional vector space over K and ω is a nondegenerate alternating bilinear form on V . The basic example is $\mathbb{R}_x^n \times \mathbb{R}_\xi^n$ with the symplectic form $\omega = \sum_{j=1}^n d\xi_j \wedge dx_j$:

$$\omega((x, \xi), (x', \xi')) = \sum_{j=1}^n (\xi_j x'_j - x_j \xi'_j). \tag{2}$$

In fact, this is essentially the only example: for a general symplectic vector space (V, ω) over a field K one can find a basis $\{e_1, \dots, e_n, f_1, \dots, f_n\}$ for V such that

$$\omega(e_j, e_k) = 0, \quad \omega(f_j, f_k) = 0, \quad \omega(f_j, e_k) = \delta_{jk} \quad \text{for all } j, k.$$

Such a basis is called a *symplectic basis*, and ω is of the form (2) in these coordinates. (In particular, a symplectic vector space is necessarily even-dimensional.) Note that ω vanishes on the span of the e_j , and it vanishes on the span of the f_j ; such a subspace is called a *Lagrangian subspace*: a maximal subspace on which ω vanishes. (A Lagrangian subspace of V is necessarily of dimension n .)

The symplectic formalism is fundamental in Hamiltonian mechanics: the symplectic form provides an isomorphism between tangent space and cotangent space,

¹Duistermaat’s book [1996] on Fourier integral operators contains a brief treatment of symplectic vector spaces over a general field.

mapping the Hamiltonian vector field of a function f to the differential of f :

$$df = \omega(\cdot, H_f).$$

A *linear symplectomorphism* T on (V, ω) is a linear isomorphism on V such that $T^*\omega = \omega$, that is,

$$\omega(Tv, Tv') = \omega(v, v') \quad \text{for all } v, v' \in V.$$

This is equivalent to the property that a symplectic basis is mapped to a symplectic basis.

We now let (V, ω) be a real symplectic vector space. Then

$$(V \times V, \omega \oplus -\omega)$$

is a real symplectic vector space. We write $\omega_0 = \omega \oplus -\omega$ so that, by definition,

$$\omega_0((v, w), (v', w')) = \omega(v, v') - \omega(w, w').$$

The following classical result (see [Tao 2012] for a broad perspective) justifies this choice of the symplectic form:

A map $\mathcal{H} : V \rightarrow V$ is a linear symplectomorphism if and only if its graph $\{(v, \mathcal{H}(v)) : v \in V\}$ is a Lagrangian subspace of $(V \times V, \omega_0)$.

For a basic example, let

$$\mathcal{H} : \mathbb{R}_x^n \times \mathbb{R}_\xi^n \rightarrow \mathbb{R}_y^n \times \mathbb{R}_\eta^n, \quad (x, \xi) \mapsto (y, \eta),$$

be a linear symplectomorphism. Then graph \mathcal{H} is a Lagrangian subspace for

$$\omega^{\mathbb{R}} = \sum_{j=1}^n d\xi_j \wedge dx_j - d\eta_j \wedge dy_j.$$

The point of view of this paper is to consider graph \mathcal{H} as an \mathbb{R} -linear subspace of a complex symplectic vector space. After all, with $z_j = x_j + iy_j$ and $\zeta_j = \xi_j + i\eta_j$, we have the complex symplectic form

$$\omega^{\mathbb{C}} = \sum_{j=1}^n d\zeta_j \wedge dz_j \quad \text{on } \mathbb{C}_z^n \times \mathbb{C}_\zeta^n,$$

which induces the two *real* symplectic forms

$$\operatorname{Re} \omega^{\mathbb{C}} = \sum_{j=1}^n d\xi_j \wedge dx_j - d\eta_j \wedge dy_j, \quad \operatorname{Im} \omega^{\mathbb{C}} = \sum_{j=1}^n d\xi_j \wedge dy_j + d\eta_j \wedge dx_j$$

on $\mathbb{R}_{x,\xi}^{2n} \times \mathbb{R}_{y,\eta}^{2n}$. We then say that an \mathbb{R} -linear $2n$ -dimensional subspace of $\mathbb{R}_{x,\xi}^{2n} \times \mathbb{R}_{y,\eta}^{2n}$ is an *R-Lagrangian subspace* if it is Lagrangian with respect to $\operatorname{Re} \omega^{\mathbb{C}}$, and an

I -Lagrangian subspace if it is Lagrangian with respect to $\text{Im } \omega^{\mathbb{C}}$. Thus the graph of $\mathcal{H} : \mathbb{R}_{x,\xi}^{2n} \rightarrow \mathbb{R}_{y,\eta}^{2n}$ may be considered as an R -Lagrangian subspace of $(\mathbb{C}_z^n \times \mathbb{C}_\zeta^n, \omega^{\mathbb{C}})$.

Writing a symplectic matrix $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$ as $\mathcal{H} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, we have

$$\text{graph } \mathcal{H} = \{((x, \xi), (Ax + B\xi, Cx + D\xi)) : (x, \xi) \in \mathbb{R}^{2n}\};$$

or, in terms of (z, ζ) , we have

$$\text{graph}_{\mathbb{C}} \mathcal{H} = \{(x + i(Ax + B\xi), \xi + i(Cx + D\xi)) : (x, \xi) \in \mathbb{R}^{2n}\}.$$

Thus

$$\omega^{\mathbb{C}}|_{\text{graph}_{\mathbb{C}} \mathcal{H}} = i \text{Im } \omega^{\mathbb{C}}|_{\text{graph } \mathcal{H}}$$

is given by

$$\begin{aligned} \omega^{\mathbb{C}}((x + i(Ax + B\xi), \xi + i(Cx + D\xi)), (x' + i(Ax' + B\xi'), \xi' + i(Cx' + D\xi'))) \\ = i \begin{pmatrix} x^T & \xi^T \end{pmatrix} \begin{pmatrix} C^T - C & -A^T - D \\ A + D^T & B - B^T \end{pmatrix} \begin{pmatrix} x' \\ \xi' \end{pmatrix}. \end{aligned}$$

The symplectic form $\text{Re } \omega^{\mathbb{C}}$ vanishes, but the symplectic form $\text{Im } \omega^{\mathbb{C}}$ might *not* vanish; that is, $\text{graph}_{\mathbb{C}} \mathcal{H}$ is R -Lagrangian but not necessarily I -Lagrangian.

We have thus defined a map from the group of symplectic matrices to the space of skew-symmetric matrices

$$\mathfrak{X} : \text{Sp}(2n, \mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R}), \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \mapsto \begin{pmatrix} C^T - C & -A^T - D \\ A + D^T & B - B^T \end{pmatrix}.$$

We can thus restate our open problem:

Open problem. Is the map $\mathfrak{X} : \text{Sp}(2n, \mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R})$ a surjection?

While we do not solve this problem, the main result of the paper is [Theorem 1](#); we can explicitly construct a generating function Φ for \mathcal{H} and thus give an alternate characterization of $\omega^{\mathbb{C}}|_{\text{graph}_{\mathbb{C}} \mathcal{H}}$ and hence of \mathfrak{X} .

2. In terms of generating functions: the proof of the theorem

Generating functions (in the sense of symplectic geometry) were discovered by Sir William Rowan Hamilton in his extensive work on optics. In modern language (and in the linear case), light rays are specified by the following data: \mathbb{R}_x^2 is a plane of initial positions perpendicular to the optical axis of the system, $\xi \in \mathbb{R}^2$ are the initial “directions” (multiplied by the index of refraction), \mathbb{R}_y^2 is a plane of terminal positions, and $\eta \in \mathbb{R}^2$ are the terminal directions. The spaces $\mathbb{R}_{x,\xi}^4$ and $\mathbb{R}_{y,\eta}^4$ are given the standard symplectic structures. Taken piece by piece, the optical system consists of a sequence of reflections and refractions for each light ray, the laws of which were long known; Hamilton’s discovery was that, taken as a whole, the optical

system is determined by a single function, the *generating function*, or, as Hamilton called it, the *characteristic function*, of the optical system. The transformation from initial conditions to terminal conditions is a symplectomorphism expressible in terms of a single scalar-valued function, “by which means optics acquires, as it seems to me, an uniformity and simplicity in which it has been hitherto deficient” [Hamilton 1828, Section IV, Paragraph 20].²

The optical framework gives an intuitive reason why, in the symplectic matrix $\mathcal{H} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, the rank of B plays a special role in characterizing \mathcal{H} and thus its generating function. Again, \mathcal{H} maps the initial (position, direction)-pair (x, ξ) to the terminal (position, direction)-pair

$$\begin{pmatrix} y \\ \eta \end{pmatrix} = \begin{pmatrix} Ax + B\xi \\ Cx + D\xi \end{pmatrix}.$$

The case $B = 0$ corresponds to perfect focusing: all the rays from a given position x arrive at the same position y , resulting in a perfect image. And the case $\det B \neq 0$ corresponds to *no* such focusing: two rays with initial position x but different initial directions must arrive at different positions y . (See [Guillemin and Sternberg 1984] for an exposition of symplectic techniques in optics.)

2.1. When B is invertible. We recall that

$$\text{graph}_{\mathbb{C}} \mathcal{H} = \{(x + i(Ax + B\xi), \xi + i(Cx + D\xi)) : (x, \xi) \in \mathbb{R}^{2n}\},$$

taken over the reals, is an R -Lagrangian subspace of $(\mathbb{C}_z^n \times \mathbb{C}_\zeta^n, \omega^{\mathbb{C}})$, and we note that

$$\pi : \text{graph}_{\mathbb{C}} \mathcal{H} \rightarrow \mathbb{C}^n, \quad (z, \zeta) \mapsto z,$$

is an \mathbb{R} -linear transformation whose kernel is given by $(x, \xi) \in \{0\} \times \ker B$. Thus it is an \mathbb{R} -linear isomorphism if and only if B is invertible. In this case, the general theory of symplectic geometry gives the existence of a real C^∞ function Φ defined on $\text{graph}_{\mathbb{C}} \mathcal{H}$ such that

$$\text{graph}_{\mathbb{C}} \mathcal{H} = \left\{ \left(z, -2 \frac{\partial \Phi}{\partial z}(z) \right) : z \in \mathbb{C}^n \right\}.$$

²There are different types of generating functions in symplectic geometry, and, as Arnold writes, “[the apparatus of generating functions] is unfortunately noninvariant and it uses, in an essential way, the coordinate structure in phase space” [Arnold 1978, Section 47]. For our purposes, we may take the term “generating function” to broadly mean a scalar-valued function which generates a symplectomorphism (or, more generally, a Lagrangian submanifold) in the same sense that a potential function generates a conservative vector field. Our generating functions are denoted by the symbol Φ below.

Hence if $\det B \neq 0$, then

$$\begin{aligned} \text{graph}_{\mathbb{C}} \mathcal{H} &= \{(x + i(Ax + B\xi), \xi + i(Cx + D\xi)) : (x, \xi) \in \mathbb{R}^{2n}\} \\ &= \{(z, -2(\partial\Phi/\partial z)(z)) : z \in \mathbb{C}^n\} \\ &= \{(p + iq, B^{-1}(q - Ap) + i(Cp + DB^{-1}(q - Ap))) : p + iq \in \mathbb{C}^n\}, \end{aligned}$$

where we write $z = p + iq$, so that

$$\Phi(p, q) = \frac{1}{2}p^T B^{-1}Ap - p^T B^{-1}q + \frac{1}{2}q^T DB^{-1}q. \quad (3)$$

This function appears in [Folland 1989, Equation (4.54)] and in [Guillemin and Sternberg 1984, Section 11]. (Note that $B^{-1}A$ and DB^{-1} are symmetric since \mathcal{H} is symplectic.) Substituting $p = x$ and $q = Ax + B\xi$, we arrive at the following expression, with the obvious abuse of notation:

$$\Phi(x, \xi) = \frac{1}{2}x^T A^T Cx + x^T C^T B\xi + \frac{1}{2}\xi^T B^T D\xi. \quad (4)$$

Or, writing Φ with respect to z and \bar{z} , we have

$$\begin{aligned} \Phi(z) &= \frac{1}{8}z^T (B^{-1}A + 2iB^{-1} - DB^{-1})z \\ &\quad + \frac{1}{4}\bar{z}^T (B^{-1}A - i(B^T)^{-1} + iB^{-1} + DB^{-1})z \\ &\quad + \frac{1}{8}\bar{z}^T (B^{-1}A - 2iB^{-1} - DB^{-1})\bar{z}. \end{aligned}$$

Thus

$$\left(\frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_k} \right) = \frac{1}{4}(B^{-1}A - iB^{-1} + i(B^T)^{-1} + DB^{-1}).$$

We can directly compute $\omega^{\mathbb{C}}$ restricted to $\text{graph}_{\mathbb{C}} \mathcal{H}$ in terms of z and \bar{z} :

$$\begin{aligned} \omega^{\mathbb{C}} \left(\left(z, -2 \frac{\partial \Phi}{\partial z}(z) \right), \left(z', -2 \frac{\partial \Phi}{\partial z}(z') \right) \right) &= 4i \operatorname{Im} \left(\sum_{j,k} z_j \left(\frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_k} \right) \bar{z}'_k \right) \\ &= 2 \sum_{j,k} \frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_k} (z_j \bar{z}'_k - z'_j \bar{z}_k). \end{aligned}$$

If we substitute

$$\begin{aligned} z &= x + i(Ax + B\xi), \\ z' &= x' + i(Ax' + B\xi'), \end{aligned}$$

then after a lengthy mechanical calculation we recover the expression

$$\begin{aligned} \omega^{\mathbb{C}} \left(\left(z, -2 \frac{\partial \Phi}{\partial z}(z) \right), \left(z', -2 \frac{\partial \Phi}{\partial z}(z') \right) \right) \\ = 2 \sum_{j,k} \frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_k} (z_j \bar{z}'_k - z'_j \bar{z}_k) = i \begin{pmatrix} x^T & \xi^T \end{pmatrix} \mathfrak{X}(\mathcal{H}) \begin{pmatrix} x' \\ \xi' \end{pmatrix}. \end{aligned}$$

2.2. When B is not invertible. When B is *not* invertible, we seek

$$\Phi = \Phi(z, \theta) \in C^\infty(\mathbb{C}^n \times \mathbb{R}^N)$$

such that

$$\text{graph}_{\mathbb{C}} \mathcal{H} = \left\{ \left(z, -2 \frac{\partial \Phi}{\partial z}(z, \theta) \right) : \frac{\partial \Phi}{\partial \theta}(z, \theta) = 0 \right\}. \quad (5)$$

We follow the general method outlined by Guillemin and Sternberg [1977].

Let

$$W = \text{graph}_{\mathbb{C}} \mathcal{H}, \quad X = \{(z, 0); z \in \mathbb{C}^n\}, \quad Y = \{(0, \zeta); \zeta \in \mathbb{C}^n\}.$$

Since W is an R -Lagrangian subspace, we know that $W \cap Y$ and $PW \subset X$ are orthogonal with respect to $\text{Re } \omega^{\mathbb{C}}$, where P is the projection onto X along Y . Indeed,

$$W \cap Y = \{(0, \xi + iD\xi) : \xi \in \ker B\}, \quad PW = \{(x + i(Ax + B\xi), 0) : (x, \xi) \in \mathbb{R}^{2n}\},$$

and we can check directly that, with $\xi \in \ker B$,

$$\omega^{\mathbb{C}}((0, \xi + iD\xi), (x' + i(Ax' + B\xi'), 0)) = i[\xi^T(A + D^T)x' + \xi^T B\xi'].$$

Since $\text{graph}_{\mathbb{C}} \mathcal{H}$ is not a \mathbb{C} -linear subspace but an \mathbb{R} -linear subspace, for now we prefer to write

$$\begin{aligned} W \cap Y &= \{(0, \xi; 0, D\xi) : \xi \in \ker B\}, \\ PW &= \{(x, 0; Ax + B\xi, 0) : (x, \xi) \in \mathbb{R}^{2n}\}. \end{aligned}$$

We note that $PW \oplus (W \cap Y)$ has real dimension $2n$, hence is a Lagrangian subspace of $(\mathbb{R}^{4n}, \text{Re } \omega^{\mathbb{C}})$.

We seek to write $\text{graph } \mathcal{H}$ as the graph of a function from $PW \oplus (W \cap Y)$ to a complementary Lagrangian subspace; as a first step, we choose a convenient symplectic basis. We let $\{b_1, \dots, b_k\}$ be an orthonormal basis for $\ker B$ and extend to an orthonormal basis $\{b_1, \dots, b_n\}$ for \mathbb{R}^n , so that

$$\{(0, b_j; 0, Db_j) : j = 1, \dots, k\}$$

is a basis for $W \cap Y$, and

$$\{(0, 0; Bb_j, 0) : j = k + 1, \dots, n\} \cup \{(b_j, 0; Ab_j, 0) : j = 1, \dots, n\}$$

is a basis for PW . We then extend to the following symplectic basis for $(\mathbb{R}^{4n}, \text{Re } \omega^{\mathbb{C}})$:

$$\begin{aligned} \{(0, 0; Ab_j, 0) : j = 1, \dots, k\} &\leftrightarrow \{(0, b_j; 0, Db_j) : j = 1, \dots, k\}, \\ \{(0, 0; Bb_j, 0) : j = k + 1, \dots, n\} &\leftrightarrow \{(0, A^T \beta_j; 0, \beta_j) : j = k + 1, \dots, n\}, \quad (6) \\ \{(b_j, 0; Ab_j, 0) : j = 1, \dots, n\} &\leftrightarrow \{(0, -b_j; 0, 0) : j = 1, \dots, n\}, \end{aligned}$$

where the $\{\beta_j\}_{j=k+1}^n$ satisfy

$$\begin{cases} A^T \beta_j \in (\ker B)^\perp = \text{Im } B^T, \\ b_J \cdot B^T \beta_j = \delta_{Jj} \quad \text{for all } J \in \{k+1, \dots, n\}. \end{cases} \tag{7}$$

One advantage of using the particular symplectic basis (6) is that the vectors on the left are all ‘‘horizontal,’’ and the vectors on the right are all ‘‘vertical’’. (The arrows signify the symplectically dual pairs.)

The following proposition implies the existence of $\{\beta_j\}_{j=k+1}^n$.

Proposition 2. *The set $\{Ab_1, \dots, Ab_k, Bb_{k+1}, \dots, Bb_n\}$ is a basis for \mathbb{R}^n .*

Proof. Suppose

$$\sum_{j=1}^k \alpha_j Ab_j + \sum_{j=k+1}^n \alpha_j Bb_j = 0.$$

We take the dot product with Db_J , $J \in \{1, \dots, k\}$, to get $\alpha_1 = \dots = \alpha_k = 0$, and the rest are zero by the linear independence of $\{Bb_{k+1}, \dots, Bb_n\}$. \square

Thus for $J \in \{k+1, \dots, n\}$ we can take β_J to be the unique vector orthogonal to the set

$$\{Ab_1, \dots, Ab_k, Bb_{k+1}, \dots, \widehat{Bb_J}, \dots, Bb_n\}$$

(where the wide hat denotes omission) and satisfying

$$\beta_J \cdot Bb_J = 1.$$

We will now describe graph \mathcal{H} in terms of the above symplectic coordinate system: we write a general linear combination of the $4n$ vectors and find necessary and sufficient conditions on the coefficients to make the vector in graph \mathcal{H} . Explicitly, we write the general vector in \mathbb{R}^{4n} as

$$\begin{aligned} & \sum_{j=1}^k t'_j(0, 0; Ab_j, 0) + \sum_{j=k+1}^n t''_j(0, 0; Bb_j, 0) + \sum_{j=1}^n t''_{n+j}(b_j, 0; Ab_j, 0) \\ & + \sum_{j=1}^k \theta'_j(0, b_j; 0, Db_j) + \sum_{j=k+1}^n \theta''_j(0, A^T \beta_j; 0, \beta_j) + \sum_{j=1}^n \theta''_{n+j}(0, -b_j; 0, 0) \end{aligned} \tag{8}$$

(the primes are not necessary but are useful for bookkeeping), and we will describe graph \mathcal{H} as (t', θ') as a function of (t'', θ') .

We have the following necessary and sufficient conditions for the vector (8) to be in graph \mathcal{H} :

Thus, defining the functions

$$f_i''(t'') = \sum_{j=k+1}^n [Bb_i \cdot Db_j]t_j'' + \sum_{j=1}^n [Bb_i \cdot Cb_j]t_{n+j}'' \quad \text{for } i = k+1, \dots, n,$$

$$f_{n+i}''(t'') = \sum_{j=k+1}^n [Cb_i \cdot Bb_j]t_j'' + \sum_{j=1}^n [Ab_i \cdot Cb_j]t_{n+j}'' \quad \text{for } i = 1, \dots, n,$$

we see that (9) is equivalent to the conditions $t' = 0$, $\theta'' = f''(t'')$. Noting that

$$\frac{\partial f_i''}{\partial t_j''} = \frac{\partial f_j''}{\partial t_i''} \quad \text{for all } i, j \in k+1, \dots, n,$$

and defining

$$F(t'') = \frac{1}{2} \sum_{i=k+1}^n \sum_{j=k+1}^n t_i'' [Bb_i \cdot Db_j] t_j''$$

$$+ \sum_{i=k+1}^n \sum_{j=1}^n t_i'' [Bb_i \cdot Cb_j] t_{n+j}'' + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n t_{n+i}'' [Ab_i \cdot Cb_j] t_{n+j}'',$$

we conclude that the vector is in graph \mathcal{H} if and only if

$$t' = 0, \quad \frac{\partial F}{\partial t''}(t'') = \theta''.$$

We now define

$$\varphi(t', t''; \theta', \theta'') = \theta' \cdot t' + F(t'') + (\theta'' - f''(t''))^2.$$

Then in $(t', t''; \theta', \theta'')$ -coordinates, graph \mathcal{H} is given as

$$\left\{ \left(t', t''; \frac{\partial \varphi}{\partial t'}, \frac{\partial \varphi}{\partial t''} \right) : \frac{\partial \varphi}{\partial \theta'} = 0, \frac{\partial \varphi}{\partial \theta''} = 0 \right\}.$$

Or, written in terms of the standard basis, graph \mathcal{H} is the set of values of

$$\sum_{j=1}^k t_j'(0, 0; Ab_j, 0) + \sum_{j=k+1}^n t_j''(0, 0; Bb_j, 0) + \sum_{j=1}^n t_{n+j}''(b_j, 0; Ab_j, 0)$$

$$+ \sum_{j=1}^k \frac{\partial \varphi}{\partial t_j'}(t, \theta)(0, b_j; 0, Db_j) + \sum_{j=k+1}^n \frac{\partial \varphi}{\partial t_j''}(t, \theta)(0, A^T \beta_j; 0, \beta_j)$$

$$+ \sum_{j=1}^n \frac{\partial \varphi}{\partial t_{n+j}''}(t, \theta)(0, -b_j; 0, 0) \quad (10)$$

subject to the condition that $\frac{\partial \varphi}{\partial \theta}(t, \theta) = 0$.

We return to complex coordinates, in the standard basis; for that purpose we write the “horizontal” parts of (10) as

$$z := \sum_{j=1}^k it'_j Ab_j + \sum_{j=k+1}^n it''_j Bb_j + \sum_{j=1}^n t''_{n+j} (I + iA)b_j.$$

That is,

$$\begin{aligned} \operatorname{Re} z &= \sum_{j=1}^n t''_{n+j} b_j, \\ \operatorname{Im} z &= \sum_{j=1}^k t'_j Ab_j + \sum_{j=k+1}^n t''_j Bb_j + \sum_{j=1}^n t''_{n+j} Ab_j. \end{aligned}$$

With the same notation as before, the inverse transformation is given by

$$\begin{aligned} t'_j &= -b_j \cdot \operatorname{Re} z + Db_j \cdot \operatorname{Im} z && \text{for } j \in \{1, \dots, k\}, \\ t''_j &= -A^T \beta_j \cdot \operatorname{Re} z + \beta_j \cdot \operatorname{Im} z && \text{for } j \in \{k+1, \dots, n\}, \\ t''_{n+j} &= b_j \cdot \operatorname{Re} z && \text{for } j \in \{1, \dots, n\}. \end{aligned} \quad (11)$$

We write the “vertical” part of (10) as:

$$\begin{aligned} \operatorname{Re} \zeta &= \sum_{j=1}^k \frac{\partial \varphi}{\partial t'_j}(t, \theta) b_j + \sum_{j=k+1}^n \frac{\partial \varphi}{\partial t''_j}(t, \theta) A^T \beta_j - \sum_{j=1}^n \frac{\partial \varphi}{\partial t''_{n+j}}(t, \theta) b_j, \\ \operatorname{Im} \zeta &= \sum_{j=1}^k \frac{\partial \varphi}{\partial t'_j}(t, \theta) Db_j + \sum_{j=k+1}^n \frac{\partial \varphi}{\partial t''_j}(t, \theta) \beta_j. \end{aligned} \quad (12)$$

Using $t = t(z)$ to denote the transformation (11), we define

$$\Phi(z, \theta) := \varphi(t(z), \theta),$$

so that (12) says

$$\zeta = -2 \frac{\partial \Phi}{\partial z}(z, \theta).$$

In summary, we now have the following expression for $\operatorname{graph}_{\mathbb{C}} \mathcal{H}$:

$$\operatorname{graph}_{\mathbb{C}} \mathcal{H} = \left\{ \left(z, -2 \frac{\partial \Phi}{\partial z}(z, \theta) \right) : \frac{\partial \Phi}{\partial \theta}(z, \theta) = 0 \right\}, \quad (13)$$

where the $\theta \in \mathbb{R}^{2n}$ are considered as auxiliary parameters, as in (5).

As for $\omega^{\mathbb{C}}|_{\operatorname{graph}_{\mathbb{C}} \mathcal{H}}$, we use the expression

$$\frac{\partial \Phi}{\partial z}(z, \theta) = \frac{\partial^2 \Phi}{\partial z \partial \theta} \cdot \theta + \frac{\partial^2 \Phi}{\partial z^2} \cdot z + \frac{\partial^2 \Phi}{\partial z \partial \bar{z}} \cdot \bar{z}$$

to compute

$$\begin{aligned} \omega^{\mathbb{C}} & \left(\left(z, -2 \frac{\partial \Phi}{\partial z}(z, \theta) \right), \left(w, -2 \frac{\partial \Phi}{\partial z}(w, \eta) \right) \right) \\ & = 2z \cdot \frac{\partial \Phi}{\partial z}(w, \eta) - 2w \cdot \frac{\partial \Phi}{\partial z}(z, \theta) \\ & = 2 \sum_{j=1}^n \sum_{\ell=1}^{2n} \frac{\partial^2 \Phi}{\partial z_j \partial \theta_\ell} (z_j \eta_\ell - w_j \theta_\ell) + 2 \sum_{j,m=1}^n \frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_m} (z_j \bar{w}_m - w_j \bar{z}_m), \end{aligned} \tag{14}$$

where the variables are related by the conditions

$$\frac{\partial \Phi}{\partial \theta}(z, \theta) = 0 \quad \text{and} \quad \frac{\partial \Phi}{\partial \theta}(w, \eta) = 0.$$

Of course, from Section 1, we know that (14) is equal to

$$i \begin{pmatrix} x^T & \xi^T \end{pmatrix} \mathfrak{X}(\mathcal{H}) \begin{pmatrix} x' \\ \xi' \end{pmatrix}, \tag{15}$$

where

$$\begin{aligned} z & = x + i(Ax + B\xi), & w & = x' + i(Ax' + B\xi'), \\ -2 \frac{\partial \Phi}{\partial z}(z, \theta) & = \xi + i(Cx + D\xi), & -2 \frac{\partial \Phi}{\partial z}(w, \eta) & = \xi' + i(Cx' + D\xi'). \end{aligned}$$

This completes the proof of the theorem.

We leave it as an illustrative exercise for the reader to compute Φ and its derivatives in the special cases when $B = 0$ and when B is invertible (to be compared with the generating function (3) in Section 2.1).

3. Application: the metaplectic representation

In the previous section, we showed how to associate to a linear symplectomorphism \mathcal{H} a (real-valued) generating function Φ . For the purposes of Fresnel optics and quantum mechanics one then associates to the generating function Φ an oscillatory integral operator

$$\mu(\mathcal{H}) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n), \quad u \mapsto a h^{-3n/2} \iint e^{i\Phi(x+iy,\theta)/h} u(x) dx d\theta. \tag{16}$$

The map $\mu : \mathcal{H} \rightarrow \mu(\mathcal{H})$ is called the *metaplectic representation* of the symplectic group, and $\mu(\mathcal{H})$ is said to be the “quantization” of the classical object \mathcal{H} . As defined, the operator $\mu(\mathcal{H})$ maps Schwartz functions to tempered distributions, but in fact it extends to a bounded operator on $L^2(\mathbb{R}^n)$; we choose a so that $\mu(\mathcal{H})$ is unitary on $L^2(\mathbb{R}^n)$, and here $h > 0$ is a small parameter. These are the operators

of “Fresnel optics,” a relatively simple model theory for optics which accounts for interference and diffraction, describing the propagation of light of wavelength h [Guillemin and Sternberg 1984]. For the analytic details we refer the reader to a text in semiclassical analysis [Dimassi and Sjöstrand 1999]; here we only show that the standard conditions are indeed satisfied.

The above (real-valued) generating function Φ , for an arbitrary $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$, has the property that the 1-forms $d(\partial\Phi/\partial\theta_1), \dots, d(\partial\Phi/\partial\theta_{2n})$ are linearly independent. Equivalently, with the notation from the previous section, the matrix

$$\begin{pmatrix} \frac{\partial^2\Phi}{\partial(\text{Re } z)\partial\theta'} & \frac{\partial^2\Phi}{\partial(\text{Re } z)\partial\theta''} \\ \frac{\partial^2\Phi}{\partial(\text{Im } z)\partial\theta'} & \frac{\partial^2\Phi}{\partial(\text{Im } z)\partial\theta''} \\ \frac{\partial^2\Phi}{\partial\theta'^2} & \frac{\partial^2\Phi}{\partial\theta'\partial\theta''} \\ \frac{\partial^2\Phi}{\partial\theta''\partial\theta'} & \frac{\partial^2\Phi}{\partial\theta''^2} \end{pmatrix} = \begin{pmatrix} \begin{array}{c} | \\ (-b_1) \end{array} & \cdots & \begin{array}{c} | \\ (-b_k) \end{array} & * \\ \begin{array}{c} | \\ | \\ Db_1 \end{array} & \cdots & \begin{array}{c} | \\ | \\ Db_k \end{array} & * \\ \begin{array}{c} | \\ | \\ 0_{k,k} \end{array} & & \begin{array}{c} | \\ | \\ 0_{k,(2n-k)} \end{array} & \\ \begin{array}{c} | \\ | \\ 0_{(2n-k),k} \end{array} & & \begin{array}{c} | \\ | \\ 2I_{(2n-k),(2n-k)} \end{array} & \end{pmatrix}$$

has linearly independent columns. (The asterisks denote irrelevant components.) This condition says that quadratic form $\Phi = \Phi(z, \theta)$ is a *nondegenerate phase function* in the sense of semiclassical analysis [Dimassi and Sjöstrand 1999].

Folland writes: “it seems to be a fact of life that there is no simple description of the operator $\mu(\mathcal{A})$ that is valid for all $\mathcal{A} \in \text{Sp}$ ” [Folland 1989, p. 193]; however, we believe that (16), combined with our construction of Φ in the proof of Theorem 1, is such a description.

4. Conclusion

The open problem and results presented in this paper were motivated by the basic question of the relationship between real and complex symplectic linear algebra. Our approach to this question was to consider a real symplectomorphism as a Lagrangian submanifold with regard to the real part of a complex symplectic form. We believe the resulting problem of the nature of the restriction of the imaginary part of the complex symplectic form to this submanifold (formally, $\mathfrak{X}(\mathcal{H})$ for a symplectomorphism \mathcal{H}) is relevant to the structure of the real symplectic group. (We direct the reader to the Appendix for a list of properties of \mathfrak{X} and reformulations of our open problem which lend credence to this belief.) Accordingly, we view the main result of this paper as primarily a means for further investigation of the open problem of the image of \mathfrak{X} . In addition to solving our open problem, we believe that, in line with our generating function formulation, it would be interesting to have a “complexified” theory of the calculus of variations. At present we only have trivial extensions of the real theory.

Appendix

A. Elementary properties of \mathfrak{X} . We first note some standard facts about symplectic matrices that are used throughout the paper; for further information, see, for example, [Cannas da Silva 2001] or [Folland 1989]. We write

$$\mathcal{J} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$$

for the matrix representing the standard symplectic form.

Proposition 3 [Folland 1989]. *Let $\mathcal{H} \in GL(2n, \mathbb{R})$. The following are equivalent:*

- (1) $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$.
- (2) $\mathcal{H}^T \mathcal{J} \mathcal{H} = \mathcal{J}$.
- (3) $\mathcal{H}^{-1} = \mathcal{J} \mathcal{H}^T \mathcal{J}^{-1} = \begin{pmatrix} D^T & -B^T \\ -C^T & A^T \end{pmatrix}$.
- (4) $\mathcal{H}^T \in \text{Sp}(2n, \mathbb{R})$.
- (5) $A^T D - C^T B = I$, $A^T C = C^T A$ and $B^T D = D^T B$.
- (6) $AD^T - BC^T = I$, $AB^T = BA^T$ and $CD^T = DC^T$.

While \mathfrak{X} may be extended to all of $\mathbb{M}^{2n}(\mathbb{R})$,

$$\mathfrak{X} : \mathbb{M}^{2n}(\mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R}), \quad M \mapsto \mathcal{J}M + M^T \mathcal{J}, \tag{1}$$

for purposes of our open problem the resulting linearity of \mathfrak{X} does not seem to help when \mathfrak{X} is restricted to $\text{Sp}(2n, \mathbb{R})$.

The following proposition presents some of the most interesting elementary linear algebraic properties of \mathfrak{X} , which follow immediately from the definition.

Proposition 4. *Let $\mathfrak{X} : \mathbb{M}^{2n}(\mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R})$ be defined as above. Then:*

- (1) $\ker(\mathfrak{X}) = \mathfrak{sp}(2n, \mathbb{R})$, the symplectic Lie algebra.
- (2) For any $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$, $\mathfrak{X}(\mathcal{H}) = \mathcal{J}(\mathcal{H} + \mathcal{H}^{-1})$.
 In particular, for $\mathcal{U} \in U(n) = \left\{ \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \in \text{Sp}(2n, \mathbb{R}) \right\}$ we have $\mathcal{U}^{-1} = \mathcal{U}^T$, so $\mathfrak{X}(\mathcal{U}) = \mathcal{J}(\mathcal{U} + \mathcal{U}^T)$.
- (3) For any $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$, $\mathfrak{X}(\mathcal{H})$ is invertible (equivalently, $\text{Im } \omega^{\mathbb{C}}|_{\text{graph } \mathcal{H}}$ is nondegenerate) if and only if -1 is not a member of the spectrum of \mathcal{H}^2 .
- (4) For $\mathcal{H}, \mathcal{R} \in \text{Sp}(2n, \mathbb{R})$, we have $\mathcal{H}^T \mathfrak{X}(\mathcal{R}) \mathcal{H} = \mathfrak{X}(\mathcal{H}^{-1} \mathcal{R} \mathcal{H})$.

We now take some examples.

Examples of symplectic matrices and their images under \mathfrak{X} .

(1)
$$\begin{pmatrix} A & 0 \\ 0 & (A^T)^{-1} \end{pmatrix} \mapsto \begin{pmatrix} 0 & -A^T - (A^T)^{-1} \\ A + A^{-1} & 0 \end{pmatrix}.$$

In particular,

$$\begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \mapsto \begin{pmatrix} 0 & -2I \\ 2I & 0 \end{pmatrix} = 2\mathcal{J}.$$

(2) For $B = B^T$,

$$\begin{pmatrix} I & B \\ 0 & I \end{pmatrix} \mapsto \begin{pmatrix} 0 & -2I \\ 2I & 0 \end{pmatrix}.$$

(3) For $C = C^T$,

$$\begin{pmatrix} I & 0 \\ C & I \end{pmatrix} \mapsto \begin{pmatrix} 0 & -2I \\ 2I & 0 \end{pmatrix}.$$

(4)
$$\mathcal{J} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \mapsto \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

(5) For $t \in \mathbb{R}$,

$$\begin{pmatrix} (\cos t)I & (-\sin t)I \\ (\sin t)I & (\cos t)I \end{pmatrix} \mapsto \begin{pmatrix} 0 & -2(\cos t)I \\ 2(\cos t)I & 0 \end{pmatrix}.$$

(6) For any $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$, we have $\mathfrak{X}(\mathcal{H}) = \mathfrak{X}(\mathcal{H}^{-1})$.

Thus in Examples (2) and (3), $\text{graph}_{\mathbb{C}} \mathcal{H}$ is an RI -subspace (R -Lagrangian and I -symplectic). And in Example (4), $\text{graph}_{\mathbb{C}} \mathcal{H}$ is a C -Lagrangian subspace (R -Lagrangian and I -Lagrangian).

The exact nature of the image of \mathfrak{X} is an open question. The following is a partial result

Proposition 5. *For each $k \in \{0, 1, \dots, n\}$, there exists $\mathcal{H}_k \in \text{Sp}(2n, \mathbb{R})$ such that $\text{rank}(\mathfrak{X}(\mathcal{H}_k)) = 2k$. Moreover, for any $\mathcal{H} \in \text{Sp}(2n, \mathbb{R})$, we have $\ker \mathfrak{X}(\mathcal{H}) = \ker(\mathcal{H}^2 + I)$.*

Proof. We fix $k \in \{0, 1, \dots, n\}$ and write

$$(x, \xi) = (x', x'', \xi', \xi''), \quad x', \xi' \in \mathbb{R}^k, \quad x'', \xi'' \in \mathbb{R}^{n-k}.$$

Let

$$\mathcal{H}_k(x', x'', \xi', \xi'') = (x', -\xi'', \xi', x'').$$

The matrix representation of \mathcal{H}_k is

$$\begin{pmatrix} I_k & 0_k & & \\ & 0_{n-k} & -I_{n-k} & \\ 0_k & I_k & & \\ & I_{n-k} & 0_{n-k} & \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{R}).$$

Then

$$\mathfrak{X}(\mathcal{H}_k) = \begin{pmatrix} & & -2I_k & \\ & & & 0_{n-k} \\ 2I_k & & & \\ & 0_{n-k} & & \end{pmatrix},$$

so that

$$\mathrm{rank}(\mathfrak{X}(\mathcal{H}_k)) = 2k.$$

The last statement of the proposition follows from (1). □

B. Restatement of the problem. It is sometimes convenient to work with the extension of \mathfrak{X} to all of $\mathbb{M}^{2n}(\mathbb{R})$:

$$\mathfrak{X}(M) = \mathcal{J}M + M^T \mathcal{J}.$$

Then $\mathfrak{X} : \mathbb{M}(2n, \mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R})$ is a linear epimorphism with kernel $\mathfrak{sp}(2n, \mathbb{R})$, the symplectic Lie algebra (see, for example, [Folland 1989, Proposition 4.2]). Thus the map $\mathfrak{X}|_{\mathrm{Sp}(2n, \mathbb{R})}$ is surjective if and only if every element of the quotient space $\mathbb{M}(2n, \mathbb{R})/\mathfrak{sp}(2n, \mathbb{R})$ contains a symplectic matrix. So our question is:

Question. Can every $M \in \mathbb{M}(2n, \mathbb{R})$ be written as $M = \mathcal{H} + \mathcal{A}$ for some $\mathcal{H} \in \mathrm{Sp}(2n, \mathbb{R})$ and some $\mathcal{A} \in \mathfrak{sp}(2n, \mathbb{R})$?

Proposition 6. *Every $M \in \mathbb{M}(2n, \mathbb{R})/\mathfrak{sp}(2n, \mathbb{R})$ has a unique representative of the form*

$$\begin{pmatrix} 0 & S_2 \\ S_3 & \mathcal{D} \end{pmatrix},$$

where S_2 and S_3 are skew-symmetric.

Proof. Existence: let

$$M = \begin{pmatrix} M_1 & M_2 \\ M_3 & M_4 \end{pmatrix} \in \mathbb{M}(2n, \mathbb{R}).$$

Since $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathfrak{sp}(2n, \mathbb{R})$ if and only if $\delta = -\alpha^T$, $\beta = \beta^T$, $\gamma = \gamma^T$, we may replace M by

$$\tilde{M} = M - \begin{pmatrix} M_1 & \frac{1}{2}(M_2 + M_2^T) \\ \frac{1}{2}(M_3 + M_3^T) & -M_1^T \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{2}(M_2 - M_2^T) \\ \frac{1}{2}(M_3 - M_3^T) & M_4 + M_1^T \end{pmatrix}.$$

Uniqueness: suppose

$$\begin{pmatrix} 0 & S_2 \\ S_3 & D \end{pmatrix} = \begin{pmatrix} 0 & S'_2 \\ S'_3 & D' \end{pmatrix} \in \mathbb{M}(2n, \mathbb{R}) / \mathfrak{sp}(2n, \mathbb{R}),$$

with the S_j and S'_j skew-symmetric. Thus

$$\begin{pmatrix} 0 & S_2 - S'_2 \\ S_3 - S'_3 & D - D' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha^T \end{pmatrix} \in \mathfrak{sp}(2n, \mathbb{R}).$$

This shows that $S_j - S'_j$ is symmetric and skew-symmetric, hence zero, and it is clear that $D = D'$. \square

Thinking geometrically, we are to find the projection of $\mathrm{Sp}(2n, \mathbb{R})$ onto

$$\left\{ \begin{pmatrix} 0 & S_2 \\ S_3 & D \end{pmatrix} : S_2, S_3 \text{ skew-symmetric} \right\}$$

along $\mathfrak{sp}(2n, \mathbb{R})$. That is, let $\mathcal{H} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{R})$. Then

$$\pi(\mathcal{H}) = \begin{pmatrix} 0 & \frac{1}{2}(B - B^T) \\ \frac{1}{2}(C - C^T) & A^T + D \end{pmatrix}.$$

Is every

$$\begin{pmatrix} 0 & S_2 \\ S_3 & D \end{pmatrix}$$

of this form?

For a possible simplification, the map

$$\mathcal{Y} : \mathrm{Sp}(2n, \mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R}), \quad \mathcal{H} \mapsto \mathfrak{X}(-\mathcal{J}\mathcal{H}) = \mathcal{H} - \mathcal{H}^T,$$

has the same image as $\mathfrak{X} : \mathrm{Sp}(2n, \mathbb{R}) \rightarrow \mathfrak{so}(2n, \mathbb{R})$ and may be easier to understand.

Acknowledgements

Funding for this project was generously provided by Grinnell College, as part of its summer directed research program. VanValkenburgh is currently a faculty member at California State University, Sacramento, but was a visiting faculty member at Grinnell College while the first draft of the paper was written; he thanks both institutions for their support. The authors are also grateful to the anonymous referee for a thoughtful reading of the paper and suggestions for its improvement.

References

[Abraham and Marsden 1978] R. Abraham and J. E. Marsden, *Foundations of mechanics*, 2nd ed., Benjamin/Cummings, Reading, MA, 1978. [MR 81e:58025](#) [Zbl 0393.70001](#)

- [Arnold 1978] V. I. Arnold, *Mathematical methods of classical mechanics*, Graduate Texts in Mathematics **60**, Springer, New York, 1978. [MR 57 #14033b](#) [Zbl 0386.70001](#)
- [Cannas da Silva 2001] A. Cannas da Silva, *Lectures on symplectic geometry*, Lecture Notes in Mathematics **1764**, Springer, Berlin, 2001. [MR 2002i:53105](#) [Zbl 1016.53001](#)
- [Dimassi and Sjöstrand 1999] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series **268**, Cambridge University Press, 1999. [MR 2001b:35237](#) [Zbl 0926.35002](#)
- [Duistermaat 1996] J. J. Duistermaat, *Fourier integral operators*, Progress in Mathematics **130**, Birkhäuser, Boston, MA, 1996. [MR 96m:58245](#) [Zbl 0841.35137](#)
- [Everitt and Markus 2004] W. N. Everitt and L. Markus, *Infinite dimensional complex symplectic spaces*, Mem. Amer. Math. Soc. **810**, Amer. Math. Soc., Providence, RI, 2004. [MR 2005d:46048](#) [Zbl 1054.46019](#)
- [Folland 1989] G. B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies **122**, Princeton University Press, 1989. [MR 92k:22017](#) [Zbl 0682.43001](#)
- [Grigis and Sjöstrand 1994] A. Grigis and J. Sjöstrand, *Microlocal analysis for differential operators: An introduction*, London Mathematical Society Lecture Note Series **196**, Cambridge University Press, 1994. [MR 95d:35009](#) [Zbl 0804.35001](#)
- [Guillemin and Sternberg 1977] V. Guillemin and S. Sternberg, *Geometric asymptotics*, Mathematical Surveys **14**, Amer. Math. Soc., Providence, RI, 1977. [MR 58 #24404](#) [Zbl 0364.53011](#)
- [Guillemin and Sternberg 1984] V. Guillemin and S. Sternberg, *Symplectic techniques in physics*, Cambridge University Press, 1984. [MR 86f:58054](#) [Zbl 0576.58012](#)
- [Hamilton 1828] Sir W. R. Hamilton, “Theory of systems of rays: Part first”, *Transactions of the Royal Irish Academy* **15** (1828), 69–174.
- [Hamilton 1834] Sir W. R. Hamilton, “On a general method in dynamics”, *Philosophical Transactions of the Royal Society* **124** (1834), 247–308.
- [Schapira 1981] P. Schapira, “Conditions de positivité dans une variété symplectique complexe: Application à l’étude des microfonctions”, *Ann. Sci. École Norm. Sup.* (4) **14**:1 (1981), 121–139. [MR 82i:58067](#) [Zbl 0473.58022](#)
- [Sjöstrand 1982] J. Sjöstrand, *Singularités analytiques microlocales*, Astérisque **95**, Société Mathématique de France, Paris, 1982. [MR 84m:58151](#) [Zbl 0524.35007](#)
- [Tao 2012] T. Tao, “The closed graph theorem in various categories”, blog post, 20 November 2012, <http://terrytao.wordpress.com/2012/11/20/the-closed-graph-theorem-in-various-categories/>.
- [Viterbo 1992] C. Viterbo, “Symplectic topology as the geometry of generating functions”, *Math. Ann.* **292**:1 (1992), 685–710. [MR 93b:58058](#) [Zbl 0735.58019](#)
- [Weinstein 1981] A. Weinstein, “Symplectic geometry”, *Bull. Amer. Math. Soc. (N.S.)* **5**:1 (1981), 1–13. [MR 83a:58044](#) [Zbl 0465.58013](#)

Received: 2013-09-20

Revised: 2014-08-24

Accepted: 2014-10-31

hellmann@grinnell.edu*Department of Mathematics and Statistics, Grinnell College, Grinnell, IA 50112, United States*langenba@grinnell.edu*Department of Mathematics and Statistics, Grinnell College, Grinnell, IA 50112, United States*mjv@csus.edu*Department of Mathematics and Statistics, California State University, Sacramento, CA 95819, United States*

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moselehian	Ferdowsi University of Mashhad, Iran ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA kgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Pomomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sgupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nhritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

involve

2015

vol. 8

no. 4

The Δ^2 conjecture holds for graphs of small order	541
COLE FRANKS	
Linear symplectomorphisms as R -Lagrangian subspaces	551
CHRIS HELLMANN, BRENNAN LANGENBACH AND MICHAEL VANVALKENBURGH	
Maximization of the size of monic orthogonal polynomials on the unit circle corresponding to the measures in the Steklov class	571
JOHN HOFFMAN, MCKINLEY MEYER, MARIYA SARDARLI AND ALEX SHERMAN	
A type of multiple integral with log-gamma function	593
DUOKUI YAN, RONGCHANG LIU AND GENG-ZHE CHANG	
Knight's tours on boards with odd dimensions	615
BAOYUE BI, STEVE BUTLER, STEPHANIE DEGRAAF AND ELIZABETH DOEBEL	
Differentiation with respect to parameters of solutions of nonlocal boundary value problems for difference equations	629
JOHNNY HENDERSON AND XUEWEI JIANG	
Outer billiards and tilings of the hyperbolic plane	637
FILIZ DOGRU, EMILY M. FISCHER AND CRISTIAN MIHAI MUNTEANU	
Sophie Germain primes and involutions of \mathbb{Z}_n^\times	653
KARENNA GENZLINGER AND KEIR LOCKRIDGE	
On symplectic capacities of toric domains	665
MICHAEL LANDRY, MATTHEW MCMILLAN AND EMMANUEL TSUKERMAN	
When the catenary degree agrees with the tame degree in numerical semigroups of embedding dimension three	677
PEDRO A. GARCÍA-SÁNCHEZ AND CATERINA VIOLA	
Cylindrical liquid bridges	695
LAMONT COLTER AND RAY TREINEN	
Some projective distance inequalities for simplices in complex projective space	707
MARK FINCHER, HEATHER OLNEY AND WILLIAM CHERRY	