

# involve

a journal of mathematics

Maximization of the size of  
monic orthogonal polynomials on the unit circle  
corresponding to the measures in the Steklov class

John Hoffman, McKinley Meyer, Mariya Sardarli and Alex Sherman





# Maximization of the size of monic orthogonal polynomials on the unit circle corresponding to the measures in the Steklov class

John Hoffman, McKinley Meyer, Mariya Sardarli and Alex Sherman

(Communicated by Sever S. Dragomir)

We investigate the size of monic, orthogonal polynomials defined on the unit circle corresponding to a finite positive measure. We find an upper bound for the  $L_\infty$  growth of these polynomials. Then we show, by example, that this upper bound can be achieved. Throughout these proofs, we use a method developed by Rahmanov to compute the polynomials in question. Finally, we find an explicit formula for a subsequence of the Verblunsky coefficients of the polynomials.

## 1. Introduction

Let  $V = C(\mathbb{T}; \mathbb{C})$ , where  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ . We define an inner product on  $V$  by

$$\langle f, g \rangle_{d\mu} = \int_{\mathbb{T}} f(z) \overline{g(z)} d\mu,$$

where  $d\mu$  is of the form

$$d\mu = p(\theta) d\theta + \sum_{j=1}^n m_j \delta(\theta - \theta_j),$$

where  $p(\theta)$  is a continuous function,  $\delta$  is the Dirac delta function, and the  $m_j$  are masses placed at the  $\theta_j$  satisfying  $m_j \geq 0$ . We will confine our analysis to measures in the restricted Steklov class of order  $\delta$ , denoted  $S_\delta$ , which consists of measures with the properties

$$p(\theta) > \delta, \quad m_j \geq 0, \quad \langle 1, 1 \rangle_{d\mu} = 2. \tag{1-1}$$

---

*MSC2010:* 42C05.

*Keywords:* OPUC, classical analysis, approximation theory, orthogonal polynomials on the unit circle.

At the time of writing, Hoffman, Sherman and Meyer were undergraduates at the University of Wisconsin–Madison, and Sardarli was an undergraduate at Princeton University. Hoffman is the corresponding author.

This inner product gives a norm  $\|\cdot\|$  defined as

$$\|f(z)\|_{d\mu} = \sqrt{\langle f, f \rangle_{d\mu}}.$$

Given a measure  $d\mu \in S_\delta$ , there exists a unique set of monic orthogonal polynomials  $\{\phi_n(z; d\mu)\}$  [Simon 2005]. We will adopt the convention that  $\phi_n(z; d\mu)$  is the polynomial of degree  $n$  in this set. When there is no ambiguity about what measure is being used, we will simply write these polynomials as  $\phi_n(z)$ . Corresponding to the set  $\{\phi_n(z; d\mu)\}$  is the set  $\{\varphi_n(z)\}$  of orthonormal polynomials, defined by

$$\varphi_n(z) = \frac{\phi_n(z)}{\|\phi_n(z)\|}.$$

These polynomials form an orthonormal set. Uniqueness of this set follows the from uniqueness of  $\{\phi_n(z)\}$ .<sup>1</sup>

A conjecture of Steklov stated that the sequence

$$M_{n,\delta} = \sup_{d\mu \in S_\delta} \max_{z \in \mathbb{T}} |\varphi_n(z; d\mu)|$$

is bounded in  $n$ . This was disproven by Rahmanov [1979]. In particular, Rahmanov proved the existence of a probability measure  $d\eta = \phi(\theta) d\theta + \sum_{j=1}^n m_j \delta(\theta - \theta_j)$  such that

$$|\varphi_n(1, d\eta)| \geq C \ln(n) + B$$

for some constants  $B, C$ . The hard part in making such estimates is that, in general, there are few tools available to compute  $\varphi_n(z)$  other than the Gram–Schmidt process. To establish his result, Rahmanov found a formula for computing the  $\phi_n(z; d\eta)$ , where  $d\eta = d\mu + \sum_{k=0}^n m_k \delta(\theta - \theta_k)$  in terms of  $\phi_n(z; d\mu)$ , meaning that  $d\eta$  differs from  $d\mu$  only in its masses. This formula uses the Christoffel–Darboux kernel

$$K_n(z, \xi) = \sum_{j=0}^n \overline{\varphi_j(z)} \varphi_j(\xi). \tag{1-2}$$

The roots of the Christoffel–Darboux kernel are those  $\xi_j$  satisfying

$$K_n(\xi_j, \xi_i) \begin{cases} = 0 & \text{for } i \neq j, \\ \neq 0 & \text{for } i = j. \end{cases} \tag{1-3}$$

Rahmanov’s formula, in light of these definitions, is

$$\phi_n(z; d\eta) = \phi_n(z; d\mu) - \sum_{j=1}^n \frac{m_j \phi_n(\xi_j; d\mu)}{1 + m_j K_{n-1}(\xi_j, \xi_j)} K_{n-1}(z, \xi_j). \tag{1-4}$$

---

<sup>1</sup>Originally, the last condition given for the Steklov class is stated as  $\langle 1, 1 \rangle_{d\mu} = 1$ . This is a minor modification though, because  $\varphi_n(z)/\sqrt{2}$  is in the Steklov class  $S_{\delta/\sqrt{2}}$ , given the original definition.

We now outline our results. In Theorem 2.3, we use Rahmanov’s method for computing  $\phi_n(\theta)$  for the measure  $d\eta = d\theta/2\pi + \sum_{j=1}^{\lfloor n/4 \rfloor} m_j \delta(\theta - \theta_j)$ , with  $\theta_j = (2\pi j - \pi)/n$ ,  $m_j = 4/n$  and show that the corresponding polynomials  $\phi_n(z; d\mu)$  are uniformly bounded above by  $8/(5\pi^2) \log(\lfloor n/4 \rfloor - 1) + C$ , where  $C$  is a constant. Our next main result is Theorem 4.1, where we construct a family of measures  $d\mu_n$  such that  $\phi_n(1; d\mu_n) > 1/\pi \log n + c$ ,  $c$  a bounded constant. Finally, in Theorem 5.1, we show that, given the measure  $d\mu = d\theta/2\pi + \sum_{j=1}^n m_j \delta(\theta - \theta_j)$ , with  $\theta_j = 2\pi j/n - \theta_0$ , the subsequence  $\{\alpha_{nk-1}\}_{k=1}^\infty$  of the Verblunsky coefficients  $\alpha_j(d\mu)$  satisfies

$$\alpha_{nk-1} = e^{ink\theta_0} \sum_{j=1}^n \frac{m_j}{1 + m_j nk}.$$

The reader may notice that all of our results are stated in terms of  $\phi_n(z)$ , while Steklov’s conjecture is stated in terms of  $\varphi_n(z)$ . We will end this introduction with a lemma, proven by Rahmanov [1979], that shows why bounds on  $\phi_n(z)$  imply bounds on  $\varphi_n(z)$ , and thus why it is sufficient to evaluate  $\{\phi_n(z)\}$ .

**Lemma 1.1.** *Given a measure  $d\mu \in S_\delta$ ,  $\delta > 0$ , there exists a constant  $C$  such that*

$$\frac{1}{C} \|\phi_n(z, d\mu)\| \leq \|\varphi_n(z, d\mu)\| \leq C \|\phi_n(z, d\mu)\|$$

for all  $n \geq 0$ .

*Proof.* Since

$$|\varphi_n(z)| = \frac{|\phi_n(z)|}{\|\phi_n(z)\|_{d\mu}},$$

it suffices to find constant upper and lower bounds on  $\|\phi_n(z)\|_{d\mu}$ .

To find an upper bound, we first claim that  $\phi_n(z)$  minimizes the integral

$$\int_{\mathbb{T}} |P(z)|^2 d\mu,$$

where  $P(z)$  is any monic polynomial of degree  $n$ . Let  $q(z)$  be an arbitrary polynomial of degree less than  $n$ . Then, since  $\phi_n(z)$  is orthogonal to all polynomials of degree less than  $n$  under the measure  $d\mu$  and the inner product of a polynomial with itself is nonnegative, we have

$$\begin{aligned} &\langle \phi_n(z) + q(z), \phi_n(z) + q(z) \rangle_{d\mu} \\ &= \langle \phi_n(z), \phi_n(z) \rangle_{d\mu} + \langle \phi_n(z), q(z) \rangle_{d\mu} + \langle q(z), \phi_n(z) \rangle_{d\mu} + \langle q(z), q(z) \rangle_{d\mu} \\ &= \langle \phi_n(z), \phi_n(z) \rangle_{d\mu} + \langle q(z), q(z) \rangle_{d\mu} \\ &\geq \langle \phi_n(z), \phi_n(z) \rangle_{d\mu}. \end{aligned}$$

Hence  $\phi_n(z)$  minimizes the integral  $\int_{\mathbb{T}} |P(z)|^2 d\mu$ . In particular this gives us

$$\|\phi_n(z)\|_{d\mu}^2 = \int_0^{2\pi} |\phi_n(z)|^2 d\mu \leq \int_0^{2\pi} |z^n|^2 d\mu = \int_0^{2\pi} 1 d\mu = 2. \tag{1-5}$$

We can derive a lower bound using the fact that  $d\mu \in S_\delta$  and, in particular, that  $d\mu$  satisfies (1-1), which gives

$$\begin{aligned} \|\phi_n(z)\|_{d\mu}^2 &= \frac{1}{2\pi} \int_0^{2\pi} |\phi_n(e^{i\theta})|^2 p(\theta) d\theta + \sum_{j=1}^l m_j |\phi_n(e^{i\theta_j})|^2 \\ &\geq \frac{\delta}{2\pi} \int_0^{2\pi} |\phi_n(e^{i\theta})|^2 d\theta. \end{aligned}$$

Let the coefficient of the  $z^k$  term of  $\phi_n(z)$  be  $a_k$ . In particular,  $a_n = 1$ . Using that the integral of  $e^{ik\theta}$  over the unit circle is 0 for a nonzero integer  $k$ , we get

$$\int_0^{2\pi} |\phi_n(e^{i\theta})|^2 d\theta = \int_0^{2\pi} \sum_{k=0}^n a_k^2 d\theta \geq \int_0^{2\pi} a_n^2 d\theta = 2\pi.$$

Hence,  $\|\phi_n(z)\|_{d\mu}^2 \geq \delta$ .

Combining this with the upper bound on  $\|\phi_n(z)\|_{d\mu}^2$  from (1-5) gives

$$\delta \leq \|\phi_n(z)\|_{d\mu}^2 \leq 2,$$

and as a result

$$\frac{|\phi_n(z)|}{\sqrt{2}} \leq |\phi_n(z)| \leq \frac{|\phi_n(z)|}{\sqrt{\delta}}. \quad \square$$

## 2. Review of Rahmanov’s result

We begin by reviewing Rahmanov’s argument [1979] to show that the growth of the monic polynomials under Rahmanov’s scheme is bounded below by  $c \log n$ , where  $c$  is a constant. Before doing that, though, we need to prove two lemmas that simplify our future calculations.

We now characterize the roots of the Christoffel–Darboux kernel (which we defined on page 2) for a certain measure:

**Lemma 2.1.** *For  $d\mu = d\theta/2\pi$ , the roots of the Christoffel–Darboux kernel are the  $n$ -th roots of unity times a constant of modulus one.*

*Proof.* Recall the definition of the Christoffel–Darboux kernel and its roots from (1-2) and (1-3). For our  $d\mu$ ,  $\varphi_j = z^j$ , so assuming  $\xi_j$  is of modulus one for all  $j$ ,

$$\begin{aligned} K_{n-1}(\xi_i, \xi_j) &= \sum_{j=0}^{n-1} \xi_i^j / \xi_j^j \\ &= \frac{\xi_i^n / \xi_j^n - 1}{\xi_i / \xi_j - 1} \quad (\text{since this is a geometric series}) \\ &= \frac{\xi_i^n - \xi_j^n}{(\xi_i - \xi_j)\xi_j^{n-1}}. \end{aligned} \tag{2-1}$$

Therefore, by (2-1),  $\xi_j = e^{2i\pi j/n} \xi_0$ , where  $1 \leq j \leq n$  and  $\xi_0$  is an arbitrary point on the unit circle, and so  $\xi_j$  is an  $n$ -th root of unity times a constant.  $\square$

**Lemma 2.2.** *We need only assess  $\phi_n(z; d\mu)$  at  $z = 1$ , since*

$$\sup_{\mu \in S_\delta} \max_{z \in \mathbb{T}} |\phi_n(z; d\mu)| = \sup_{\mu \in S_\delta} |\phi_n(1; d\mu)|.$$

*Proof.* Let

$$d\mu_1 = p(\theta) d\theta + \sum_{j=1}^m m_j \delta(\theta - \theta_j),$$

where  $d\mu_1 \in S_\delta$ . Then  $d\mu_2 \in S_\delta$ , where

$$d\mu_2 = p(\theta - \theta^*) d\theta + \sum_{j=1}^m m_j \delta(\theta - \theta^* - \theta_j), \quad \theta^* \in [0, 2\pi).$$

In particular,  $\phi_n(e^{i\theta}; d\mu_1) = \phi_n(e^{i(\theta+\theta^*); d\mu_2})$ .

Hence,

$$\begin{aligned} \max_{z \in \mathbb{T}} |\phi_n(z; d\mu_1)| &= \max_{z \in \mathbb{T}} |\phi_n(z; d\mu_2)|, \\ \sup_{\mu \in S_\delta} \max_{z \in \mathbb{T}} |\phi_n(z; d\mu)| &= \sup_{\mu \in S_\delta} |\phi_n(1; d\mu)|. \end{aligned}$$

Henceforth, we will only look at  $\phi_n(z; d\mu)$  evaluated at  $z = 1$ .  $\square$

**Theorem 2.3.** *Under a finite measure  $d\eta = d\mu + \sum_{j=1}^{\lfloor n/4 \rfloor} m_j \delta(\theta - \theta_j)$ , the monic polynomials are not uniformly bounded from above; specifically, there exists a  $d\eta$  such that the maximums are greater than or equal to  $8/(5\pi^2) \log(\lfloor n/4 \rfloor - 1)$ .*

**Remark 2.4.** This is Rahmanov’s result [1979], whose proof we have included for the reader’s convenience.

*Proof.* First, we will deal generally with some  $d\eta$  without specifying the added masses.

In light of Lemma 2.1, let  $\theta_j = (2\pi j - \pi)/n$  for  $1 \leq j \leq \lfloor n/4 \rfloor$ . Then, using Rahmanov’s formula in (1-4), we have

$$\phi_n(z; d\eta) = \phi_n(z; d\mu) - \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j \phi_n(\xi_j; d\mu)}{1 + m_j K_{n-1}(\xi_j, \xi_j)} K_{n-1}(z, \xi_j), \tag{2-2}$$

which, by noting that  $K_{n-1}(\xi_j, \xi_j) = \sum_{j=1}^{n-1} 1 = n$  and substituting  $z$  and  $\xi_j$  into (2-2), becomes

$$\begin{aligned} \phi_n(z; d\eta) &= z^n - \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j \xi_j^n}{1 + m_j n} \frac{-z^n - 1}{ze^{-i\theta_j} - 1} \\ &= z^n + \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{z^n + 1}{1 - ze^{-i\theta_j}}. \end{aligned}$$

Now we want to find a lower bound for  $|\phi_n|$ :

$$\max_{z \in \mathbb{T}} |\phi_n(z; d\eta)| \geq \max_{z \in \mathbb{T}} \left| \operatorname{Im} \left( z^n + \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{z^n + 1}{1 - ze^{-i\theta_j}} \right) \right|.$$

We take  $z = 1$ , in line with Lemma 2.2, to get

$$\begin{aligned} \max_{z \in \mathbb{T}} |\phi_n(z; d\eta)| &\geq \left| \operatorname{Im} \left( 1 + 2 \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{1}{1 - e^{-i\theta_j}} \right) \right| \\ &= \left| \operatorname{Im} \left( 1 - 2 \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{e^{i\theta_j} - 1}{|1 - e^{-i\theta_j}|^2} \right) \right|. \end{aligned}$$

Note that  $0 < \theta_j < \pi/2$ ,  $|1 - e^{-i\theta_j}| \leq \theta_j$ , and

$$\operatorname{Im}(e^{i\theta_j} - 1) = \sin \theta_j \geq \frac{2\theta_j}{\pi} \quad \text{for } \theta \in \left(0, \frac{\pi}{2}\right),$$

which gives

$$\left| \operatorname{Im} \left( 1 - 2 \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{e^{i\theta_j} - 1}{|1 - e^{-i\theta_j}|^2} \right) \right| \geq 2 \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{m_j}{1 + m_j n} \frac{2}{\pi \theta_j}. \tag{2-3}$$

Now, we specify the masses of  $d\eta$  to get a precise bound. Let  $m_j = 4/n$  for all  $j$ . This simplifies (2-3) to

$$\max_{z \in \mathbb{T}} |\phi_n(z; d\eta)| \geq \frac{16}{5\pi n} \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{1}{(2j - 1)\frac{\pi}{n}} \geq \frac{8}{5\pi^2} \sum_{j=1}^{\lfloor n/4 \rfloor} \frac{1}{j}.$$



Note that  $\log a = \int_1^a 1/x \, dx \geq \sum_{j=1}^{a-1} 1/j$  since  $1/x$  is decreasing.

Therefore,

$$\max_{z \in \mathbb{T}} |\phi_n(z; d\eta)| \geq \frac{8}{5\pi^2} \log\left(\left\lfloor \frac{n}{4} \right\rfloor - 1\right). \tag{2-4}$$

Since  $4/(5\pi^2) \log(\lfloor n/4 \rfloor - 1)$  is strictly increasing in  $n$ ,  $\max_{z \in \mathbb{T}} |\phi_n(z, d\eta)|$  is not uniformly bounded from above.  $\square$

### 3. Finding a general upper bound

In this section, we find a general upper bound for the growth of the monic orthogonal polynomials under a  $d\eta$  which differs from  $d\theta/2\pi$  only in the discrete portion. We prove the following theorem by making a sequence of overestimates of  $|\phi_n(1, d\eta)|$ .

**Theorem 3.1.** *Let  $d\eta$  be a measure such that*

$$d\eta = \frac{1}{2\pi} d\theta + \sum_{j=1}^n m_j \delta(\theta - \theta_j),$$

where  $m_j \geq 0$ ,  $\theta_j = 2\pi j/n + \theta_0$  for  $1 \leq j \leq n$ , and  $\theta_0 \in [0, 2\pi)$ .

Then

$$|\phi_n(1, d\eta)| \leq \frac{1}{\pi} \log n + C,$$

where  $C$  is a constant uniformly bounded in  $n$ .

**Remark 3.2.** Note the generalized offset  $\theta_0$  in the theorem. In Section 2, we used the specific offset of  $\theta_0 = -\pi/n$ , but here, we find a general upper bound under any offset.

We prove the theorem using two lemmas. The first, Lemma 3.3, finds an overestimate for  $|\phi_n(1, d\eta)|$  using Rahmanov’s formula [1979]. The second, Lemma 3.4, makes another overestimate using Taylor series.

**Lemma 3.3.** *Let  $d\eta$  be a measure such that*

$$d\eta = \frac{1}{2\pi} d\theta + \sum_{j=1}^n m_j \delta(\theta - \theta_j),$$

where  $m_j \geq 0$  and  $\theta_j = 2\pi j/n + \theta_0$ ,  $1 \leq j \leq n$ .

Then

$$|\phi_n(1)| = \begin{cases} \frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| + c_n & \text{if } \theta_0 \neq 0, \\ c_n & \text{if } \theta_0 = 0, \end{cases}$$

where  $|c_n| < 2$  for all  $n$ .

*Proof.* We first consider the case where  $\theta_0 = 0$ . If  $\theta_0 = 0$ , then  $K_{n-1}(1, e^{i\theta_j}) = 0$  for  $1 \leq j < n$  and  $K_{n-1}(1, e^{i\theta_n}) = n$ . From Rahmanov's formula (1-4),

$$|\phi_n(1)| = \left| 1 - \frac{m_n}{1 + m_n n} n \right| < 1.$$

Therefore,  $\phi_n(1)$  is not increasing in  $n$  for  $\theta_0 = 0$ . Henceforth, we restrict ourselves to working with  $\theta_0 \neq 0$ .

From Rahmanov's formula in (1-4) and applying Lemma 2.1, we derive

$$\phi_n(1) = 1 - \sum_{j=1}^n \frac{m_j}{1 + m_j n} e^{in\theta_j} \frac{\overline{e^{in\theta_j}} - 1}{e^{i\theta_j} - 1}.$$

Then, using algebra, we find that

$$\begin{aligned} \phi_n(1) &= 1 + \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{1 - e^{in\theta_j}}{1 - e^{i\theta_j}} \\ &= 1 + \frac{1 - e^{in\theta_0}}{2} \sum_{j=1}^n \frac{m_j}{1 + m_j n} \left( 1 - i \frac{\sin \theta_j}{1 - \cos \theta_j} \right), \end{aligned} \tag{3-1}$$

which implies by the triangle inequality that

$$\left| \phi_n(1) - \frac{1 - e^{in\theta_0}}{2} i \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| \leq 1 + \frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \right|.$$

Note that

$$|1 - e^{in\theta_0}| \leq 2 \quad \text{and} \quad 0 \leq \frac{m_j}{1 + m_j n} \leq \frac{1}{n}, \quad \text{so} \quad \frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \right| \leq 1.$$

Hence,

$$\left| \phi_n(1) - \frac{1 - e^{in\theta_0}}{2} i \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| \leq 2. \quad \square$$

Thus, it is sufficient to consider the growth of

$$\frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right|. \tag{3-2}$$

We want to eliminate the magnitude around the sum in (3-2).

Since  $m_j \geq 0$ , and  $\sin \theta_j / (1 - \cos \theta_j)$  is positive on  $(0, \pi)$  and negative on  $(\pi, 2\pi)$ , we have

$$\left| \sum_{j=1}^n \frac{m_j}{1+m_j n} \frac{\sin \theta_j}{1-\cos \theta_j} \right| \leq \max \left\{ \left| \sum_{\theta_j \in (0, \pi)} \frac{m_j}{1+m_j n} \frac{\sin \theta_j}{1-\cos \theta_j} \right|, \left| \sum_{\theta_j \in (\pi, 2\pi)} \frac{m_j}{1+m_j n} \frac{\sin \theta_j}{1-\cos \theta_j} \right| \right\}.$$

Now, if we alter  $d\eta$  so that the masses are instead located at  $\theta_{j*} = -2\pi j/n - \theta_0$ , essentially reflecting the discrete portion of the measure over the real axis, we see that (3-2) does not change.

Hence, since we are looking to find an upper bound of (3-1) that is independent of  $m_j$  and  $\theta_0$ , we can assume without loss of generality that we are only looking at  $\theta_j \in (0, \pi)$ , and thus take

$$\frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1+m_j n} \frac{\sin \theta_j}{1-\cos \theta_j} \right| \leq \frac{|1 - e^{in\theta_0}|}{2} \sum_{\theta_j \in (0, \pi)} \frac{m_j}{1+m_j n} \frac{\sin \theta_j}{1-\cos \theta_j}.$$

Since replacing  $\theta_0$  with  $\theta_0 + 2\pi/n$  and then shifting the index of the  $m_j$  does not affect the value of (3-2), assume  $\theta_0 \in (-2\pi/n, 0)$ . Having made these simplifications, we can now move on to the main lemma, which finds an upper bound as described in the theorem.

**Lemma 3.4.**

$$\frac{|1 - e^{in\theta_0}|}{2} \sum_{\theta_j \in (0, \pi)} \frac{\sin \theta_j}{1-\cos \theta_j} \leq n \left( 1 + \frac{1}{\pi} + \frac{1}{\pi} \log \left\lfloor \frac{n}{2} \right\rfloor \right),$$

where  $\theta_j = 2\pi j/n + \theta_0$ ,  $\theta_0 \in (-2\pi/n, 0)$ .

*Proof.* We separate the first term from the sum, since that term contributes the most to the magnitude. Recall that  $\theta_1 = 2\pi/n + \theta_0$ . Thus,

$$\frac{|1 - e^{in\theta_0}|}{2} \sum_{\theta_j \in (0, \pi)} \frac{\sin \theta_j}{1-\cos \theta_j} \leq \frac{|1 - e^{in\theta_0}|}{2} \frac{\sin \theta_1}{1-\cos \theta_1} + \sum_{\theta_j \in (2\pi/n, \pi)} \frac{\sin \theta_j}{1-\cos \theta_j}$$

since  $|1 - e^{in\theta_0}| \leq 2$ . We now bound these two terms of the sum separately.

We claim that

$$\frac{|1 - e^{in\theta_0}|}{2} \frac{\sin \theta_1}{1-\cos \theta_1} \leq n$$

for  $\theta_0 \in (-2\pi/n, 0)$ . Recall  $\theta_1 = \theta_0 + 2\pi/n$ , so hence  $|1 - e^{in\theta_0}| = |1 - e^{in\theta_1}|$ . Denote  $\theta_1$  by  $t$ , where  $t \in (0, 2\pi/n)$ . We do the calculation

$$\begin{aligned} \frac{|1 - e^{int}|}{2} \frac{\sin t}{1 - \cos t} &= \frac{\sqrt{(1 - \cos nt)^2 + (\sin nt)^2}}{2} \frac{\sin t}{2(\sin \frac{t}{2})^2} = \frac{\sqrt{2 - 2 \cos nt}}{2} \frac{\sin t}{2(\sin \frac{t}{2})^2}. \end{aligned}$$

Because  $\sin(nt/2)$  is nonnegative for  $t \in (0, 2\pi/n)$ , we have

$$\sin \frac{nt}{2} \frac{\sin t}{2(\sin \frac{t}{2})^2} = \frac{\sin \frac{nt}{2} (2 \sin \frac{t}{2} \cos \frac{t}{2})}{2(\sin \frac{t}{2})^2} = \frac{\sin \frac{nt}{2} \cos \frac{t}{2}}{\sin \frac{t}{2}},$$

$\sin(nt/2)/\sin(t/2)$  is nonnegative for  $t \in (0, 2\pi/n)$  and  $\cos t/2$  is bounded above by 1. Hence, the expression is bounded above by  $\sin(nt/2)/\sin(t/2)$ . It remains to show this is bounded above by  $n$ .

This is clearly true for  $n = 1$ . Let  $n > 1$ . Recall that  $nt \in (0, 2\pi)$ . Consider an  $(n + 1)$ -gon inscribed in a unit circle in which  $n$  of the sides of the polygon form a central angle of  $t$ . The last side of the polygon forms a central angle of  $nt$  (this angle may be reflexive.) Recall that the length of a chord of a unit circle which forms a central angle of  $t$  is  $2 \sin(t/2)$ . Similarly the length of the chord which forms a central angle of  $nt$  is  $2 \sin(nt/2)$ . As the polygon is not degenerate, the sum of the lengths of the  $n$  equal side lengths is greater than the length of the remaining side length. Namely  $2n \sin(t/2) \geq 2 \sin(nt/2)$  as desired.

We now handle the second term. To bound

$$\sum_{\theta_j \in (2\pi/n, \pi)} \frac{\sin \theta_j}{1 - \cos \theta_j},$$

note that  $\sin \theta_j / (1 - \cos \theta_j)$  is decreasing on  $(0, 2\pi)$ , so

$$\sum_{\theta_j \in (2\pi/n, \pi)} \frac{\sin \theta_j}{1 - \cos \theta_j} \leq \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{\sin(\frac{2\pi}{n} j)}{1 - \cos(\frac{2\pi}{n} j)}.$$

Recall from the Taylor expansion that we can approximate  $\sin x / (1 - \cos x)$  near 0 by  $2/x$ . In fact, since

$$\lim_{x \rightarrow 0} \frac{\sin x}{1 - \cos x} - \frac{2}{x} = 0$$

and for  $x \in (0, \pi]$ , we have

$$\frac{d}{dx} \left( \frac{\sin x}{1 - \cos x} - \frac{2}{x} \right) < 0,$$

we arrive at the inequality

$$\frac{\sin x}{1 - \cos x} \leq \frac{2}{x} \quad \text{for } x \in (0, \pi].$$

Therefore,

$$\sum_{j=1}^{\lfloor n/2 \rfloor} \frac{\sin(\frac{2\pi}{n} j)}{1 - \cos(\frac{2\pi}{n} j)} \leq \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{2}{\frac{2\pi}{n} j} = \frac{n}{\pi} \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{1}{j}.$$

Recall that  $\log a = \int_1^a 1/x \, dx \geq \sum_{j=2}^a 1/j$  since  $1/x$  is decreasing, so that

$$\frac{n}{\pi} \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{1}{j} = \frac{n}{\pi} + \frac{n}{\pi} \sum_{j=2}^{\lfloor n/2 \rfloor} \frac{1}{j} \leq \frac{n}{\pi} + \frac{n}{\pi} \log \left\lfloor \frac{n}{2} \right\rfloor,$$

and thus

$$\frac{|1 - e^{in\theta_0}|}{2} \sum_{\theta_j \in (0, \pi)} \frac{\sin \theta_j}{1 - \cos \theta_j} \leq n \left( 1 + \frac{1}{\pi} + \frac{1}{\pi} \log \left\lfloor \frac{n}{2} \right\rfloor \right). \quad \square$$

Returning to the statement of the theorem,

$$\begin{aligned} |\phi_n(1)| &\leq 2 + \frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| \\ &\leq 2 + \frac{|1 - e^{in\theta_0}|}{2} \left| \sum_{j=1}^n \frac{1}{n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| \\ &\leq 3 + \frac{1}{\pi} + \frac{1}{\pi} \log \left\lfloor \frac{n}{2} \right\rfloor. \end{aligned}$$

Since  $\log \lfloor n/2 \rfloor$  is equal to  $\log n$  plus some uniformly bounded term, we can conclude that

$$|\phi_n(1)| \leq \frac{1}{\pi} \log n + C,$$

where  $C$  is constant in  $n$ , which completes the proof of the theorem.

**Remark 3.5.** Note that here we used that

$$\frac{m_j}{1 + m_j n} \leq \frac{1}{n}.$$

If we were to use the Rahmanov scheme of distributing masses and set all  $m_j = 1/n$  then

$$\frac{m_j}{1 + m_j n} = \frac{1}{2n},$$

and the monic orthogonal polynomials given by the Rahmanov type of measure would have growth bounded from above by  $1/2\pi \log n + b$ , where  $b$  is a bounded constant.

### 4. Proving the lower bound

In this section, we construct a measure that achieves the upper bound of  $1/\pi \log n$  plus a bounded term, as described in Theorem 3.1. We accomplish this primarily by applying the technique of Lagrange multipliers to find an optimal measure.

**Theorem 4.1.** *For all  $n \in \mathbb{N}$ , there exists a measure*

$$d\eta = \frac{1}{2\pi} d\theta + \sum_{j=1}^n m_j \delta(\theta - \theta_j),$$

where  $m_j \geq 0$  and  $\sum_{j=1}^n m_j = 1$  such that

$$|\phi_n(1, d\eta)| \geq \frac{1}{\pi} \log n + c,$$

where  $c$  is a bounded constant.

We will prove this theorem as a sequence of lemmas.

The first lemma, Lemma 4.2, finds a lower bound for the expression from Lemma 3.3 which is simpler to manipulate. In the second lemma, Lemma 4.4, we apply the technique of Lagrange multipliers to that lower bound to find a critical “point”, in our case a scheme of  $m_j$ s. Finally, in the third lemma, Lemma 4.6, we insert those derived  $m_j$  into the approximation and find that we achieve the growth stated in the theorem.

Set  $\theta_j = (2\pi j - \pi)/n$ . Inserting those  $\theta_j$  into (3-1), we have that

$$|\phi_n(1)| = \frac{|1 - e^{-\pi i}|}{2} \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| + c_n = \left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| + c_n$$

for some constant  $|c_n| < 2$ . We know that  $\sin \theta_j / (1 - \cos \theta_j)$  is positive for  $\theta_j \in (0, \pi)$  and negative for  $\theta_j \in (\pi, 2\pi)$ . Thus, in order to maximize  $|\phi_n(1)|$ , we set  $m_j = 0$  for all  $j$  such that  $\theta_j \in (\pi, 2\pi)$ , which prevents destructive interference from the other side of the circle.

Under this setting, we can say that

$$\left| \sum_{j=1}^n \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \right| = \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j}.$$

We next bound this equation from below with a simpler expression.

**Lemma 4.2.** *For  $\theta_j = (2\pi j - \pi)/n$  and  $m_j \geq 0$ ,*

$$\sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} \geq \frac{1}{\pi} \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{n m_j}{1 + m_j n} \frac{1}{j} + d,$$

where  $d$  is some constant.

**Remark 4.3.** It may appear contradictory that we first find a lower bound when we want the  $n$ -th degree monic polynomial to be as large as possible. However, this lower bound is easier to manipulate, and we show in the subsequent lemmas that it actually achieves the growth stated in the theorem.

*Proof.* We prove this lemma using two approximations. We first approximate  $\sin \theta_j / (1 - \cos \theta_j)$  by  $2/\theta_j$ , from the Taylor series; we then approximate  $2/\theta_j$  by  $1/(\pi j)$ .

First, we show that  $2/\theta_j$  is a good approximation of  $\sin \theta_j / (1 - \cos \theta_j)$ . Let

$$M = \max_{\theta_j \in [0, \pi]} \left| \frac{\sin \theta_j}{1 - \cos \theta_j} - \frac{2}{\theta_j} \right|.$$

This maximum,  $M$ , is achieved because

$$\left| \frac{\sin \theta_j}{1 - \cos \theta_j} - \frac{2}{\theta_j} \right|$$

is continuous in an open neighborhood containing  $[0, \pi]$ . Thus,  $2/\theta_j$  is a good approximation and we can bound the following difference by a constant:

$$\begin{aligned} & \left| \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{\sin \theta_j}{1 - \cos \theta_j} - \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{2}{\theta_j} \right| \\ & \leq \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \left| \frac{\sin \theta_j}{1 - \cos \theta_j} - \frac{2}{\theta_j} \right| \leq \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{1}{n} M \leq M. \end{aligned}$$

Having established this, we can now replace  $2/\theta_j$  with  $2n/((2j - 1)\pi)$  and attain the inequality

$$\sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{2}{\theta_j} \geq \frac{n}{\pi} \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{m_j}{1 + m_j n} \frac{1}{j}.$$

Combining this and the previous inequality proves the lemma. □

Now that we have the simplified lower bound

$$\frac{1}{\pi} \sum_{j=1}^{\lfloor n/2 \rfloor} \frac{nm_j}{1 + m_j n} \frac{1}{j},$$

we can apply the method of Lagrange multipliers to it in order to construct the  $m_j$  that prove the theorem.

**Lemma 4.4.** *Let  $n \in \mathbb{N}$ . Consider any  $l \in \mathbb{N}$  with  $l \leq n$ . Under the constraints  $m_j \geq 0$  and  $\sum_{j=1}^l m_j = 1$ , we achieve the maximum of*

$$\sum_{j=1}^l \frac{m_j n}{(1 + nm_j)j}$$

by setting

$$m_j = \frac{m_1}{\sqrt{j}} + \frac{1}{n} \left( \frac{1}{\sqrt{j}} - 1 \right)$$

for all  $1 \leq j \leq l$ , where

$$m_1 = \left( 1 + \frac{l}{n} \right) \frac{1}{\sum_{j=1}^{\lfloor n/2 \rfloor} 1/\sqrt{j}} - \frac{1}{n}.$$

*Proof.* Set up  $f$ , the function to be maximized, and the constraint  $g$ , where  $\mathbf{m}$  is the vector listing all  $m_j$ :

$$\begin{aligned} f(\mathbf{m}) &= \sum_{j=1}^l \frac{m_j n}{(1 + nm_j)j}, \\ g(\mathbf{m}) &= \sum_{j=1}^l m_j - 1 = 0. \end{aligned} \tag{4-1}$$

If  $f$  under the constraint  $g$  has a local extremum at  $\mathbf{m}'$  and  $\mathbf{m}'$  is not on the boundary, for example  $m'_j > 0$  for all  $1 \leq j \leq l$ , then there is a  $\lambda \in \mathbb{R}$  such that

$$\nabla f(\mathbf{m}') = \lambda \nabla g(\mathbf{m}'). \tag{4-2}$$

To simplify the following expressions, denote

$$\sum_{j=1}^l \frac{1}{\sqrt{j}} = \alpha(l).$$

Calculations yield that, for all  $j$ ,

$$\frac{n}{(1 + nm'_1)^2} = \frac{n}{j(1 + nm'_j)^2},$$

which, substituting  $m'_1$  and  $m'_j$ , gives  $m'_j$  in terms of  $m'_1$ , that is,

$$m'_j = \frac{m'_1}{\sqrt{j}} + \frac{1}{n} \left( \frac{1}{\sqrt{j}} - 1 \right). \tag{4-3}$$

Inserting that expression for  $m'_j$  into  $g(\mathbf{m}') = \sum_{j=1}^l m'_j - 1 = 0$  yields

$$m'_1 = \left( 1 + \frac{l}{n} \right) \frac{1}{\alpha(l)} - \frac{1}{n}.$$



**Remark 4.5.** For all  $1 \leq j \leq l$ , we have  $m'_j > 0$  since  $\alpha(l) < \int_0^l 1/\sqrt{l} dl = 2\sqrt{l}$ . Thus, we satisfy the condition that  $m'_j \geq 0$ .

To insure that, in the computation for  $\mathbf{m}'$ , the Lagrange multipliers method did in fact give us the  $\mathbf{m}$  that maximized  $f(\mathbf{m})$  under the constraint  $m_j \geq 0$  and  $\sum_{j=1}^l m_j = 1$ , we must check the boundary. We next provide a quick proof that the maximum is not achieved at the boundary.

Consider the Lagrangian  $L(\mathbf{m}) = f(\mathbf{m}) - \lambda g(\mathbf{m})$  defined on  $(-1/n, \infty)^l$ , where  $\lambda$  is the constant in (4-2). Note that  $\mathbf{m}'$  is a critical point of  $L$  since  $\mathbf{m}'$  satisfies  $\nabla L = \nabla f - \lambda \nabla g = 0$ . It suffices to show that  $L$  is concave on  $(-1/n, \infty)^l$ .

We first calculate the entries of the Lagrangian  $L$ :

$$\frac{\partial^2 L}{\partial m_j^2} = -\frac{2n^2}{j} \frac{1}{(1 + nm_j)^3} < 0,$$

$$\frac{\partial^2 L}{\partial m_j \partial m_k} = 0 \quad \text{for } j \neq k.$$

The Hessian of  $L$  is then negative definite and hence  $L$  is concave on  $(-1/n, \infty)^l$ . Therefore,  $\mathbf{m}'$  as computed in (4-3) is a point where  $L$  achieves a global maximum on the open neighborhood  $(-1/n, \infty)^l$ . In particular,  $L(\mathbf{m}')$  is the maximum of  $L$  on the region defined by  $m_j \geq 0$  and  $\sum_{j=1}^l m_j = 1$ , a subset of  $(-1/n, \infty)^l$ . On this region,  $g = 0$ , so  $L = f$ . Hence  $f$ , constrained to the aforementioned region, achieves a global maximum at  $\mathbf{m}'$ . □

We conclude the proof by calculating the value of

$$\sum_{j=1}^l \frac{m_j n}{(1 + nm_j)j}$$

for  $m_j$  as described in Lemma 4.4. Since this function evaluated at  $l = \lfloor n/2 \rfloor$  is a lower bound of  $|\phi_n(1, d\eta)|$ , as proved in Lemma 4.2, this final lemma concludes the proof of the theorem.

**Lemma 4.6.** For the  $m_j$  described in Lemma 4.4 in (4-3),

$$\sum_{j=1}^l \frac{m_j n}{(1 + nm_j)j} = \frac{1}{\pi} \log l + c,$$

where  $c$  is uniformly bounded.

*Proof.* We simply evaluate  $f$  from (4-1) at the  $\mathbf{m}'$  given by (4-3):

$$\begin{aligned} f(\mathbf{m}') &= \sum_{j=1}^l \frac{m'_j}{1 + nm'_j} \frac{n}{j} = \sum_{j=1}^l \frac{\frac{1}{n} \left( \frac{1}{\sqrt{j}} (1 + nm'_1) - 1 \right)}{1 + n \frac{1}{n} \left( \frac{1}{\sqrt{j}} (1 + nm'_1) - 1 \right)} \frac{n}{j} \\ &= \sum_{j=1}^l \frac{\frac{1}{\sqrt{j}} (1 + nm'_1) - 1}{\sqrt{j} (1 + nm'_1)} = \sum_{j=1}^l \frac{1}{j} - \frac{1}{1 + nm'_1} \alpha(l) \\ &= \sum_{j=1}^l \frac{1}{j} - \frac{\alpha(l)^2}{\left(1 + \frac{l}{n}\right)n} \\ &= \sum_{j=1}^l \frac{1}{j} - \frac{\alpha(l)^2}{n + l}. \end{aligned}$$

Now  $\sum_{j=1}^l 1/j$  differs from  $\log l$  by at most 1, and  $\alpha(l)^2/(n + l)$  is bounded in  $n$  and  $l$  since

$$0 \leq \frac{\alpha(l)^2}{n + l} < \frac{(2\sqrt{l})^2}{n + l} = \frac{4l}{n + l} \leq \frac{4n}{n} = 4.$$

Therefore, for the  $\mathbf{m}'$  given by (4-3),  $f(\mathbf{m}') = \log l + d_l$ , where  $d_l$  is a constant bounded uniformly in  $l$ . In light of Lemma 4.2, we have constructed a  $d\eta$  such that  $|\phi_n(1, d\eta)| \geq 1/\pi \log n + c$ , where  $c$  is a bounded constant, completing the proof of Theorem 4.1. □

### 5. Investigating higher degree polynomials

In the previous sections, we described the magnitude of monic polynomials of degree less than or equal to  $n$ , where  $n$  is the number of discrete masses in the measure, using Rahmanov’s formula in (1-4). However, we also want to describe the higher degree monic polynomials, i.e.,  $\phi_{n'}(z; d\eta)$ , where  $n' > n$ . Unfortunately, we are not able to do this for all  $n' > n$ , but we can partially describe  $\phi_{n'}(z; d\eta)$ , where  $n' = kn$ ,  $k \in \mathbb{N}$ .

Recall the definition of Verblunsky coefficients [Simon 2005]:

$$\phi_{n+1}(z) = z\phi_n(z) - \bar{\alpha}_n \phi_n^*(z), \tag{5-1}$$

where

$$\begin{aligned} \phi_n(z) &= \beta_n z^n + \dots + \beta_0, \quad 0 \leq j \leq n, \quad \beta_j \in \mathbb{C}, \\ \phi_n^*(z) &= \bar{\beta}_0 z^n + \dots + \bar{\beta}_n. \end{aligned}$$

In the  $n' = kn$  case, we are able to derive the corresponding Verblunsky coefficients, and do so explicitly for a  $d\rho$  similar to that of Rahmanov’s in Section 2.

**Theorem 5.1.** For a measure  $d\eta = d\theta/2\pi + \sum_{j=1}^n m_j \delta(\theta - \theta_j)$ , with masses located at  $\xi_j = e^{i\theta_j}$  and  $\theta_j = 2\pi j/n + \theta_0$  (cf. Lemma 2.1),

$$\phi_{nk}(z, d\eta) = z^{nk} - \xi_0^{nk} \sum_{j=1}^n \frac{m_j}{1 + m_j nk} K_{nk-1}(z, \xi_j),$$

and

$$\alpha_{nk-1} = \frac{1}{\xi_0^{nk}} \sum_{j=1}^n \frac{m_j}{1 + m_j nk},$$

where  $\alpha_{nk-1}$  is a Verblunsky coefficient. Furthermore, under Rahmanov's scheme, where  $\theta_j = 2\pi j/n$  and

$$d\rho = \frac{d\theta}{2\pi} + \sum_{j=1}^n \frac{\delta(\theta - \theta_j)}{n},$$

the Verblunsky coefficients are

$$\alpha_{nk-1} = \frac{1}{1 + k}.$$

*Proof.* Note that, since  $\phi_n(z; d\eta)$  is a monic polynomial,  $\beta_n$  from the above definition of the Verblunsky coefficients is 1, so

$$\phi_n^*(0; d\eta) = 1,$$

which by (5-1) implies

$$\phi_{n+1}(0; d\eta) = -\bar{\alpha}_n. \tag{5-2}$$

Having set out these preliminaries, we can simply apply Rahmanov's formula [1979] from (1-4) to find a formula for  $\phi_{nk}(z; d\eta)$  under a measure  $d\eta$  as described in the statement of Theorem 5.1:

$$\phi_{nk}(z; d\eta) = z^{nk} - \sum_{j=1}^n \frac{m_j \phi_{nk}(\xi_j; d\mu)}{1 + m_j K_{nk-1}(\xi_j, \xi_j)} K_{nk-1}(z, \xi_j) \tag{5-3}$$

$$= z^{nk} - \xi_0^{nk} \sum_{j=1}^n \frac{m_j}{1 + m_j nk} K_{nk-1}(z, \xi_j). \tag{5-4}$$

**Remark 5.2.** The simplification of the numerator from (5-3) to (5-4) depends upon the  $\xi_j$  being roots of unity times a constant (as in Lemma 2.1). Such a simplification is only possible in the  $\phi_{nk}$  case, which is why the description of other higher-degree monic polynomials is considerably more complicated.

Now consider  $z = 0$  to find the Verblunsky coefficients:

$$\phi_{nk}(0, d\eta) = -\xi_0^{nk} \sum_{j=1}^n \frac{m_j}{1 + m_j nk} K_{nk-1}(0, \xi_j) = -\xi_0^{nk} \sum_{j=1}^n \frac{m_j}{1 + m_j nk},$$

and, applying (5-2), we obtain

$$\begin{aligned} -\bar{\alpha}_{nk-1} &= \phi_{nk}(0, d\eta) = -\xi_0^{nk} \sum_{j=1}^n \frac{m_j}{1 + m_j nk}, \\ \alpha_{nk-1} &= \overline{\xi_0^{nk}} \sum_{j=1}^n \frac{m_j}{1 + m_j nk}. \end{aligned} \tag{5-5}$$

If we now take  $\theta_0 = 0$ , as Rahmanov does, and

$$d\rho = \frac{d\theta}{2\pi} + \sum_{j=1}^n \frac{\delta(\theta - \theta_j)}{n},$$

then (5-5) simplifies to

$$\alpha_{nk-1} = \frac{1}{1 + k}. \tag{5-6}$$


□

**Remark 5.3.** It is noteworthy that, as  $k$  grows, the  $\alpha_{nk-1}$  decay at the rate of  $1/(1 + k)$ . In light of the fact that  $\sum_{j=1}^{\infty} \alpha_j^2 < \infty$  [Simon 2005], this suggests that the  $\alpha_j$  are small for  $j \in (n(k - 1), nk)$ , where  $k \in \mathbb{N}$ , and increase rapidly near  $j = kn$ . However, as mentioned above, describing  $\phi_j(z; d\eta)$  for  $j \neq kn$  is much more complicated.

### Appendix: Numerical appendix

In order to help visualize the results of this paper, the graphs of the magnitudes of four orthogonal monic polynomials induced by four respective measures have been included at the end of this section. Each measure has a continuous portion of  $d\theta/2\pi$  as well as masses placed at  $\theta_j = \pi/n(2j - 1)$ , where  $1 \leq j \leq n/2$  (cf. Lemma 2.1). For simplicity, throughout this section, we will consider only even  $n$ . For the first two polynomials (displayed in Figure 1), masses of uniform size  $2/n$  are used as suggested by Rahmanov (see Section 2). For the second two (Figure 2), the masses are given their weights according to (4-3).

These graphs have several key features in common, including the presence of two peaks that grow in  $n$ : one at  $\theta = 0$  and another at  $\theta = \pi$ . Also, both have much lower minimums in the range  $0 \leq \theta \leq \pi$  than in  $-\pi \leq \theta \leq 0$ . Upon closer inspection, it can be seen that the two peaks in Figure 1 are equal; in contrast, in




**Figure 1.** Left:  $|\phi_{10}(\theta)|$  for  $\theta_j = \frac{\pi}{10}(2j - 1)$  and  $m_j = \frac{1}{5}$ , where  $1 \leq j \leq 5$ . Right:  $|\phi_{100}(\theta)|$  for  $\theta_j = \frac{\pi}{100}(2j - 1)$  and  $m_j = \frac{1}{50}$ , where  $1 \leq j \leq 50$ .

Figure 2, the peak at  $\theta = 0$  is larger than the peak at  $\theta = \pi$ . Additionally, the peak at  $\theta = 0$  in the latter case is higher than in the former, as predicted by Theorem 4.1.

To explain some of these features, first note that with the above choice of placement of the masses, Rahmanov’s formula (1-4) [1979] reduces to

$$\begin{aligned} \operatorname{Re}(\phi_n(e^{i\theta})) &= (1 + \cos(n\theta)) \left( 1 + \frac{1}{2} \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \right) - \frac{1}{2} \sin(n\theta) \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \frac{\sin(\theta - \theta_j)}{1 - \cos(\theta - \theta_j)}, \\ \operatorname{Im}(\phi_n(e^{i\theta})) &= \sin(n\theta) \left( 1 + \frac{1}{2} \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \right) + \frac{1}{2} (1 + \cos(n\theta)) \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \frac{\sin(\theta - \theta_j)}{1 - \cos(\theta - \theta_j)}. \end{aligned}$$



**Figure 2.** Left:  $|\phi_{10}(\theta)|$  for  $\theta_j = \frac{\pi}{10}(2j - 1)$  and  $m_j$  chosen optimally, where  $1 \leq j \leq 5$ . Right:  $|\phi_{100}(\theta)|$  for  $\theta_j = \frac{\pi}{100}(2j - 1)$  and  $m_j$  chosen optimally, where  $1 \leq j \leq 50$ .

**Analysis of the minima.** Due to its prominent role in each term, let us evaluate both the real and imaginary parts at the extrema of  $1 + \cos(n\theta)$ , that is,  $\theta = \theta_k = (\pi/n)(2k-1)$  and  $\theta = \theta_k^* = 2\pi k/n$ . For  $\theta = \theta_k$ ,  $\sin(n\theta_k)$  and  $1 + \cos(n\theta_k)$  are each zero. However, we must be careful, because for  $1 \leq k \leq n/2$ , one of the terms in the sum will have a denominator of zero. Thus, using L'Hôpital's rule, we take the limits

$$\begin{aligned} \lim_{\theta \rightarrow \theta_k} \frac{\sin(n\theta) \sin(\theta - \theta_k)}{1 - \cos(\theta - \theta_k)} &= -2n, \\ \lim_{\theta \rightarrow \theta_k} \frac{(1 + \cos(n\theta)) \sin(\theta - \theta_k)}{1 - \cos(\theta - \theta_k)} &= 0. \end{aligned}$$

Substituting these values into our formulae, we then have that

$$|\phi_n(e^{i\theta_k})| = \begin{cases} 1 - nm_k/(1 + nm_k) & \text{if } 1 \leq k \leq n/2, \\ 1 & \text{otherwise.} \end{cases}$$

Thus, the minima will be lower in the region where the masses are placed than outside that region. Also, we can now see the reason for the minima increasing as  $\theta$  increases in the cases where the choice of  $m_j$  is optimal, as in Figure 2.

**Analysis of peaks at  $\theta = 0, \pi$ .** Now, let us examine the values of the polynomials at  $\theta = \theta_k^*$ . In this case,  $\sin(n\theta_k^*)$  is still zero, but  $1 + \cos(n\theta_k^*)$  is instead 2, so we need not worry about zero denominators. Immediately, we have that our previous formulae reduce to

$$\begin{aligned} \operatorname{Re}(\phi_n(e^{i\theta_k^*})) &= 1 + \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j}, \\ \operatorname{Im}(\phi_n(e^{i\theta_k^*})) &= \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \frac{\sin(\theta_k^* - \theta_j)}{1 - \cos(\theta_k^* - \theta_j)}. \end{aligned} \tag{A-1}$$

The real part is constant in  $\theta_k^*$  and can be ignored. For  $k = 0$ , we have precisely the sum that was analyzed in Section 4. For  $k = n/2$ , we obtain the sum

$$\begin{aligned} \operatorname{Im}(\phi_n(e^{i\theta_{n/2}^*})) &= \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \frac{\sin(\pi - \theta_j)}{1 - \cos(\pi - \theta_j)} \\ &= \sum_{j=1}^{n/2} \frac{m_j}{1 + nm_j} \frac{\sin \theta_j}{1 + \cos \theta_j}. \end{aligned}$$

It can easily be seen that, if  $m_j$  is constant, this sum will be identical to the sum for  $k = 0$ , and so the result will be two peaks of equal amplitude as we observed before in Figure 1. If  $m_j$  decreases proportionally to  $1/\sqrt{j}$ , however, this sum

will be very different from the sum for  $k = 0$ , since the largest terms of the sum will now be those  $\theta_j$  close to  $\pi$  rather than zero. The  $m_j$  with corresponding  $\theta_j$  close to  $\pi$  will all be of the order  $1/n$ , and so we would expect that the value of the polynomial here will behave something more similarly to the peaks of the uniform mass case than to those of the optimal  $m$  case.

**Analysis of peaks away from  $\theta = 0, \pi$ .** However, we have not yet explained why the peaks away from  $\theta = 0$  and  $\theta = \pi$  are all smaller, so now we consider the case where  $\theta = \theta_k^*$  for  $0 < k < n/2$ . First, note that

$$\theta_k^* - \theta_j = \frac{\pi}{n}(2(k - j) + 1),$$

and consider the terms in the sum (A-1), where  $j = k$  and  $j = k + 1$ . These terms will be

$$\frac{m_k}{1 + nm_k} \frac{\sin \frac{\pi}{n}}{1 - \cos \frac{\pi}{n}}$$

and

$$-\frac{m_{k+1}}{1 + nm_{k+1}} \frac{\sin \frac{\pi}{n}}{1 - \cos \frac{\pi}{n}}.$$

In the case that all the masses have equal weight, these terms will cancel out completely, and, even in the case of the optimal choice of  $m_j$ , they still mostly cancel out since the difference of  $m_{k+1}$  and  $m_k$  will be small. In general, for the  $j = k - l$  and  $j = k + l + 1$  terms, as long as  $k - l \geq 1$  and  $k + l + 1 \leq n/2$  are satisfied, similar cancellations will occur. Thus, the values at these peaks will be less than those at  $\theta = 0$  and  $\theta = \pi$ .

### Acknowledgements

Our research was done during the 2013 University of Wisconsin-Madison REU, sponsored by NSF grants DMS-1056327 and DMS-1147523. We would like to thank Serguei Denissov for introducing us to this problem and advising us as we wrote this paper.

### References

[Rahmanov 1979] E. A. Rahmanov, “Steklov’s conjecture in the theory of orthogonal polynomials”, *Mat. Sb. (N.S.)* **108(150)**:4 (1979), 581–608. In Russian; translated in *USSR-Sb* **36**:4 (1980), 549–575. MR 81j:42042 Zbl 0452.33012

[Simon 2005] B. Simon, *Orthogonal polynomials on the unit circle, I: Classical theory*, American Mathematical Society Colloquium Publications **54**, Amer. Math. Soc., Providence, RI, 2005. MR 2006a:42002a Zbl 1082.42020

hoffman.locke@gmail.com	<i>University of Wisconsin-Madison, Madison, WI 53706, United States</i>
mtmeyer3@wisc.edu	<i>Department of Applied and Natural Sciences, University of Wisconsin-Green Bay, 2420 Nicolet Drive, Green Bay, WI 54311, United States</i>
sardarli@princeton.edu	<i>Princeton University, Princeton, NJ 08544, United States</i>
ajsherman2@wisc.edu	<i>University of Wisconsin-Madison, Madison, WI 53706, United States</i>



## MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

## BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moselehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA jgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sngupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

## PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# involve

2015

vol. 8

no. 4

The $\Delta^2$ conjecture holds for graphs of small order	541
COLE FRANKS	
Linear symplectomorphisms as $R$ -Lagrangian subspaces	551
CHRIS HELLMANN, BRENNAN LANGENBACH AND MICHAEL VANVALKENBURGH	
Maximization of the size of monic orthogonal polynomials on the unit circle corresponding to the measures in the Steklov class	571
JOHN HOFFMAN, MCKINLEY MEYER, MARIYA SARDARLI AND ALEX SHERMAN	
A type of multiple integral with log-gamma function	593
DUOKUI YAN, RONGCHANG LIU AND GENG-ZHE CHANG	
Knight's tours on boards with odd dimensions	615
BAOYUE BI, STEVE BUTLER, STEPHANIE DEGRAAF AND ELIZABETH DOEBEL	
Differentiation with respect to parameters of solutions of nonlocal boundary value problems for difference equations	629
JOHNNY HENDERSON AND XUEWEI JIANG	
Outer billiards and tilings of the hyperbolic plane	637
FILIZ DOGRU, EMILY M. FISCHER AND CRISTIAN MIHAI MUNTEANU	
Sophie Germain primes and involutions of $\mathbb{Z}_n^\times$	653
KARENNA GENZLINGER AND KEIR LOCKRIDGE	
On symplectic capacities of toric domains	665
MICHAEL LANDRY, MATTHEW MCMILLAN AND EMMANUEL TSUKERMAN	
When the catenary degree agrees with the tame degree in numerical semigroups of embedding dimension three	677
PEDRO A. GARCÍA-SÁNCHEZ AND CATERINA VIOLA	
Cylindrical liquid bridges	695
LAMONT COLTER AND RAY TREINEN	
Some projective distance inequalities for simplices in complex projective space	707
MARK FINCHER, HEATHER OLNEY AND WILLIAM CHERRY	



1944-4176(2015)8:4;1-3